

Hardy Spaces Associated with Ball Quasi-Banach Function Spaces on Spaces of Homogeneous Type: Characterizations of Maximal Functions, Decompositions, and Dual Spaces

Xianjie Yan, Ziyi He, Dachun Yang* and Wen Yuan

Abstract Let (X, ρ, μ) be a space of homogeneous type in the sense of Coifman and Weiss, and $Y(X)$ a ball quasi-Banach function space on X , which supports a Fefferman–Stein vector-valued maximal inequality, and the boundedness of the powered Hardy–Littlewood maximal operator on its associate space. The authors first introduce the Hardy space $H_Y^*(X)$, associated with $Y(X)$, via the grand maximal function, and then establish its various real-variable characterizations, respectively, in terms of radial or non-tangential maximal functions, atoms or finite atoms, and molecules. As an application, the authors give the dual space of $H_Y^*(X)$, which proves to be a ball Campanato-type function space associated with $Y(X)$. All these results have a wide range of generality and, particularly, even when they are applied to variable Hardy spaces, the obtained results are also new. The major novelties of this article exist in that, to escape the reverse doubling condition of μ and the triangle inequality of ρ , the authors cleverly construct admissible sequences of balls, and fully use the geometrical properties of X expressed by dyadic reference points or dyadic cubes and, to overcome the difficulty caused by the lack of the good dense subset of $H_Y^*(X)$, the authors further prove that $Y(X)$ can be embedded into the weighted Lebesgue space with certain special weight, and then can fully use the known results of the weighted Lebesgue space.

Contents

1	Introduction	2
2	Ball quasi-Banach function spaces on spaces of homogeneous type	6
2.1	Spaces of homogeneous type	6
2.2	Ball quasi-Banach function spaces	7
2.3	Boundedness of the Hardy–Littlewood maximal operator	11
2.4	Embedding theorem for ball quasi-Banach function spaces	13

2020 *Mathematics Subject Classification*. Primary 42B30; Secondary 42B25, 42B35, 46E36, 30L99.

Key words and phrases. space of homogeneous type, ball quasi-Banach function space, Hardy space, maximal function, atom, molecule, ball Campanato-type function space.

This project is partially supported by the National Key Research and Development Program of China (Grant No. 2020YFA0712900), the National Natural Science Foundation of China (Grant Nos. 11971058, 12071197 and 11871100) and the Fundamental Research Funds for the Central Universities (Grant Nos. 500421359 and 500421126).

*Corresponding author, E-mail: dcyang@bnu.edu.cn/August 10, 2021/Final version.

3	Maximal function characterizations of $H_Y^*(X)$	19
3.1	Relations between $Y(X)$ and $H_Y^*(X)$	19
3.2	Maximal function characterizations	26
4	Atomic and finite atomic characterizations of $H_Y^*(X)$	29
4.1	Atomic characterizations	29
4.2	Finite atomic characterizations	45
5	Molecular characterizations of $H_Y^*(X)$	50
6	Dual space of $H_Y^*(X)$	56

1 Introduction

It is well known that the real-variable theory of Hardy-type spaces on \mathbb{R}^n including various real-variable equivalent characterizations and the boundedness of Calderón–Zygmund operators plays a key role in harmonic analysis and partial differential equations (see, for instance, [52, 63]). Recall that the classical Hardy space $H^p(\mathbb{R}^n)$ with $p \in (0, 1]$ was originally introduced by Stein and Weiss [64] and further studied by Fefferman and Stein [18]. From then on, various variants of classical Hardy spaces have been introduced and their real-variable theories have been well developed (see, for instance, [1, 35, 36, 37, 47, 49, 54, 73]). Recently, Sawano et al. [59] originally introduced the ball quasi-Banach function space X on \mathbb{R}^n , which further generalizes the Banach function space in [5] in order to include weighted Lebesgue spaces, Morrey spaces, mixed-norm Lebesgue spaces, Orlicz-slice spaces, and Musielak–Orlicz spaces. Observe that the aforementioned several function spaces are not quasi-Banach function spaces (see, for instance, [59, 61, 69, 76]). In the same article [59], Sawano et al. also introduced the Hardy space $H_X(\mathbb{R}^n)$ associated with X , and established its various maximal function characterizations by assuming that the Hardy–Littlewood maximal operator is bounded on the p -convexification of X , and several other real-variable characterizations, respectively, in terms of atoms, molecules, and Lusin-area functions by assuming that the Hardy–Littlewood maximal operator satisfies a Fefferman–Stein vector-valued inequality on X , and is bounded on the associate space of X . Later, Wang et al. [68] further established the Littlewood–Paley g -function and g_λ^* -function characterizations of both $H_X(\mathbb{R}^n)$ and its localized version $h_X(\mathbb{R}^n)$, and obtained the boundedness of Calderón–Zygmund operators and pseudo-differential operators, respectively, on $H_X(\mathbb{R}^n)$ and $h_X(\mathbb{R}^n)$. Under much weaker assumptions on the Littlewood–Paley functions than the corresponding ones in [59, 68], Chang et al. [11] obtained various Littlewood–Paley function characterizations of $H_X(\mathbb{R}^n)$. Recently, Zhang et al. [76] and Wang et al. [69] introduced weak Hardy-type spaces $WH_X(\mathbb{R}^n)$ associated with X and developed a complete real-variable theory of these spaces; Yan et al. [71] established the dual theory and obtained the intrinsic square function characterizations of $H_X(\mathbb{R}^n)$; Zhang et al. [75] introduced some new ball Campanato-type function space which proves the dual space of $H_X(\mathbb{R}^n)$, and established its Carleson measure characterization. For more studies about ball quasi-Banach function spaces on \mathbb{R}^n , we refer the reader to [41, 42, 58, 72].

On the other hand, Coifman and Weiss [13, 14] originally introduced the notion of spaces of homogeneous type in order to establish a uniform framework for both the theory of Calderón–Zygmund operators and the real-variable theory of Hardy spaces over different underlying spaces.

From then on, spaces of homogeneous type have become the most natural and the most general underlying space to study the real-variable theory of function spaces and the boundedness of operators (see, for instance, [15, 43, 46, 51, 53]). However, many existing results of both function spaces and boundedness of operators on spaces of homogeneous type require some additional geometrical assumptions on the underlying spaces such as the reverse doubling property of the equipped measure μ (see, for instance, [23, 60, 62, 78]). Recently, a breakthrough on the analysis over a space X of homogeneous type without any additional geometrical assumptions was made by Auscher and Hytönen [4] who constructed an orthonormal wavelet basis, with exponential decay, of $L^2(X)$ by using the system of random dyadic cubes. From then on, the real-variable theory of function spaces on a given space X of homogeneous type has rapidly been developed. To be precise, Han et al. [26] introduced the Hardy space via wavelets on X and established the criteria of the boundedness of Calderón–Zygmund operators on it and its dual space. Fu and Yang [20] obtained an unconditional basis and several equivalent characterizations of the atomic Hardy space on X in terms of wavelets. He et al. [31] introduced a kind of approximation of the identity with exponential decay and obtained new Calderón reproducing formulae on X , which proved necessary to establish various real-variable characterizations of Hardy spaces. Motivated by this, He et al. [30] developed a complete real-variable theory of Hardy spaces on X including various real-variable equivalent characterizations and the boundedness of sublinear operators. Fu et al. [19] and Zhou et al. [77] generalized the corresponding results in [30], respectively, to Musielak–Orlicz Hardy spaces and Hardy–Lorentz spaces. We refer the reader to [9, 10, 12, 21, 27, 32, 33, 44, 67] for more studies on the real-variable theory of function spaces over X , and [17, 45, 48, 66] for their applications in the boundedness of operators; particularly, we refer the reader to [6, 7, 8] for some recent progress on the real-variable theory of function spaces on X associated with operators.

Throughout this article, we always let (X, ρ, μ) be a space of homogeneous type in the sense of Coifman and Weiss, and $Y(X)$ a ball quasi-Banach function space on X , which supports a Fefferman–Stein vector-valued maximal inequality, and the boundedness of the powered Hardy–Littlewood maximal operator on its associate space. Motivated by [30] and [59], in this article, using the grand maximal function, we first introduce the Hardy space $H_Y^*(X)$ associated with $Y(X)$, and then establish its various real-variable characterizations, respectively, in terms of radial or non-tangential maximal functions, atoms or finite atoms, and molecules. As an application, we also give the dual space of $H_Y^*(X)$, which proves to be a ball Campanato-type function space associated with $Y(X)$. These results can further be used to establish the Littlewood–Paley function characterizations of $H_Y^*(X)$ and the boundedness of Calderón–Zygmund operators on $H_Y^*(X)$; to limit the length of this article, they are presented in [70]. The major novelties of this article exist in that, to escape the reverse doubling condition of μ and the triangle inequality of ρ , we cleverly construct admissible sequences of balls, and fully use the geometrical properties of X expressed by dyadic reference points and dyadic cubes and, to overcome the difficulty caused by the lack of the good dense subset of $Y(X)$, we further prove that $Y(X)$ can be embedded into the weighted Lebesgue space with certain special weight, and then can fully use the known results of the weighted Lebesgue space.

We point out that the results obtained in this article and [70] have a wide range of generality because $H_Y^*(X)$ includes various known Hardy spaces on X , for instance, Hardy spaces in [31], Hardy–Lorentz spaces in [77], weighted Hardy spaces in [19], Orlicz–Hardy spaces in [19], and variable Hardy spaces on RD-spaces in [78]. Moreover, even when these results are applied to these concrete cases, some results are also new; obviously, more applications are expectable.

The remainder of this article is organized as follows.

In Section 2, we recall some notation and notions used throughout this article. More precisely, in Subsection 2.1, we recall the definition of a space \mathcal{X} of homogeneous type, and some basic properties of \mathcal{X} . In Subsection 2.2, we introduce the ball quasi-Banach function space $Y(\mathcal{X})$ on \mathcal{X} and give some examples. Some mild assumptions on the boundedness of the Hardy–Littlewood maximal operator on $Y(\mathcal{X})$ are stated in Subsection 2.3, which are needed throughout this article. In Subsection 2.4, we first creatively introduce the notion of admissible sequences of balls to get rid of the dependence on the reverse doubling property of μ because an admissible sequence of balls are either doubling and also reverse doubling (see Remark 2.25 below for more details). Also, it is well known that a ball quasi-Banach function space may not have a good dense subset and hence the usual dense argument does not work anymore in this setting. To overcome this difficulty, using admissible sequences of balls, we further show that a ball quasi-Banach function space can be embedded to the weighted Lebesgue space with certain special $A_1(\mathcal{X})$ weight (see Theorem 2.17 below), which plays a crucial role in clarifying the relations between $Y(\mathcal{X})$ and the corresponding Hardy space (see the proof of Theorem 3.12 below) and also in establishing the atomic characterization of the corresponding Hardy space (see the proof of Proposition 4.5 below).

The aim of Section 3 is to introduce the Hardy space $H_Y^*(\mathcal{X})$ associated with $Y(\mathcal{X})$, discuss the relations between $Y(\mathcal{X})$ and $H_Y^*(\mathcal{X})$, and establish various maximal function characterizations of $H_Y^*(\mathcal{X})$. To be precise, in Subsection 3.1, we first introduce three Hardy spaces $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$ with $a \in (0, \infty)$, and $H_Y^*(\mathcal{X})$ associated with $Y(\mathcal{X})$, via using, respectively, the radial maximal function, the non-tangential maximal function, and the grand maximal function in Definition 3.9, and then clarify the relations between $Y(\mathcal{X})$ and $H_Y^*(\mathcal{X})$ in Theorem 3.12 below, whose proof strongly depends on the embedding of $Y(\mathcal{X})$ into the weighted Lebesgue space with certain special $A_1(\mathcal{X})$ weight obtained in Theorem 2.17. In Subsection 3.2, we first prove that, as subspaces of the space of distributions, the Hardy spaces mentioned above coincide with each other in the sense of equivalent quasi-norms (see Theorem 3.14 below). Indeed, by [30, (3.1)], we know that these maximal functions are pointwisely comparable with each other. From this, the boundedness of the Hardy–Littlewood maximal operator on $Y(\mathcal{X})$, and an argument similar to that used in the proof of [30, (3.4)], we deduce Theorem 3.14. Second, by borrowing some ideas from the proof of [30, Proposition 3.8], we prove that $H_Y^*(\mathcal{X})$ are independent of the choices of distributions (see Theorem 3.15 below). Finally, combining Theorems 3.14 and 3.15, we further obtain various maximal function characterizations of $H_Y^*(\mathcal{X})$.

Section 4 is devoted to establishing the atomic and the finite atomic characterizations of $H_Y^*(\mathcal{X})$. In Subsection 4.1, using a variant on \mathcal{X} of the key lemma obtained in [76, Lemma 4.8], the boundedness of the powered Hardy–Littlewood maximal operator on the associate space of $Y(\mathcal{X})$, and the Fefferman–Stein vector-valued maximal inequality on $Y(\mathcal{X})$, we prove a “reconstruction” result of $H_Y^*(\mathcal{X})$ (see Proposition 4.2 below). On the other hand, $Y(\mathcal{X})$ does not have an absolutely continuous quasi-norm, which makes the good dense subspace of $H_Y^*(\mathcal{X})$ be still unknown. Thus, the proof of the “decomposition” result (see Proposition 4.5 below) is quite different from the methods used in the proofs of [19, Theorem 5.4] and [30, Theorem 4.2]. Indeed, we borrow some ideas from the proof of [77, Theorem 4.6] to directly obtain the atomic decomposition of distributions in $H_Y^*(\mathcal{X})$ instead of using some good dense function subspace. However, due to the lack of an explicit norm expression of $Y(\mathcal{X})$, the methods used in the proof of the convergence of atomic decompositions (see, for instance, [77, (4.27) and (4.41)]) are inapplicable anymore. To overcome this difficulty, in Lemma 4.9 below, we establish a continuous embedding of $Y(\mathcal{X})$ into

the weighted Lebesgue space $L_w^p(X)$ with certain special $A_1(X)$ weight w , which is an application of Theorem 2.17. From this and the Fefferman–Stein vector-valued maximal inequality for $L_w^p(X)$, we prove the convergence of Calderón–Zygmund decompositions for a distribution in $H_Y^*(X)$ (see Lemma 4.11). By using Lemma 4.9 again, and the known atomic characterization of the weighted Hardy space which is a special case of the known atomic characterization of the Musielak–Orlicz Hardy space on X obtained in [19, Theorem 5.4], we further prove the convergence of atomic decompositions in (4.22) and hence complete the proof of Proposition 4.5. In Subsection 4.2, we further obtain the finite atomic characterization of $H_Y^*(X)$.

The aim of Section 5 is to characterize $H_Y^*(X)$ by molecules (see Theorem 5.3 below). We first introduce the notions of both the molecule and the molecular Hardy space $H_{\text{mol}}^{Y,q,d,\epsilon}(X)$ in Definition 5.1. By the fact that each $(Y(X), q)$ -atom is also a $(Y(X), q, \epsilon)$ -molecule, and Proposition 4.5, we prove $H_Y^*(X) \subset H_{\text{mol}}^{Y,q,d,\epsilon}(X)$. On the other hand, we show that any $(Y(X), q, \epsilon)$ -molecule can be divided into an infinite linear combination of $(Y(X), q)$ -atoms in Lemma 5.2, which, combined with an argument similar to that used in the proof of Proposition 4.2, further implies that $H_{\text{mol}}^{Y,q,d,\epsilon}(X) \subset H_Y^*(X)$ and hence completes the proof of the molecular characterization of $H_Y^*(X)$.

Throughout Section 6, we always assume that $Y(X)$ has an absolutely continuous quasi-norm. Under this additional assumption, we first establish a dominated convergence theorem on $Y(X)$, and then prove that the finite atomic Hardy is dense in $H_Y^*(X)$ (see Lemma 6.4 below). From this, Theorem 4.19, and Theorem 4.23, we deduce that the dual space of $H_Y^*(X)$ is the ball Campanato-type function space $\mathcal{L}_{Y,q,d}(X)$ associated with $Y(X)$. Remarkably, similarly to [39, 75], to overcome the difficulty caused by the fact that the quasi-norm of $Y(X)$ may not be concave, we consider the role as a whole of finite linear combinations of atoms instead of the role of a single atom in the construction of the ball Campanato-type function space; this makes perfect sense because the finite atomic Hardy space is dense in the corresponding Hardy space, and the most important thing is that both of them have equivalent quasi-norms.

At the end of this section, we make some conventions on notation. Let $\mathbb{N} := \{1, 2, \dots\}$ and $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$. We denote by C a *positive constant* which is independent of the main parameters, but may vary from line to line. We use $C_{(\alpha, \dots)}$ to denote a positive constant depending on the indicated parameters α, \dots . The symbol $f \lesssim g$ means $f \leq Cg$ and, if $f \lesssim g \lesssim f$, then we write $f \sim g$. If $f \leq Cg$ and $g = h$ or $g \leq h$, we then write $f \lesssim g \sim h$ or $f \lesssim g \lesssim h$, rather than $f \lesssim g = h$ or $f \lesssim g \leq h$. If E is a subset of X , we denote by $\mathbf{1}_E$ its *characteristic function* and by E^c the set $X \setminus E$. For any $r \in (0, \infty)$ and $x \in X$, we denote by $B(x, r)$ the ball centered at x with the radius r , namely, $B(x, r) := \{y \in X : \rho(x, y) < r\}$. For any ball B , we use x_B to denote its center and r_B its radius, and denote by λB for any $\lambda \in (0, \infty)$ the ball concentric with B having the radius λr_B . For any index $q \in [1, \infty]$, we denote by q' its *conjugate index*, namely, $1/q + 1/q' = 1$. For any $x, x_0, y \in X$ and $r, \gamma \in (0, \infty)$, let $V_r(x) := \mu(B(x, r))$,

$$V(x, y) := \begin{cases} \mu(B(x, \rho(x, y))) & \text{if } x \neq y, \\ 0 & \text{if } x = y, \end{cases}$$

and

$$P_\gamma(x_0, x; r) := \frac{1}{V_r(x_0) + V(x_0, x)} \left[\frac{r}{r + \rho(x_0, x)} \right]^\gamma. \quad (1.1)$$

2 Ball quasi-Banach function spaces on spaces of homogeneous type

In this section, we first recall some basic notions about spaces of homogeneous type and ball quasi-Banach function spaces, respectively, in Subsections 2.1 and 2.2, which are used throughout this article. Then we make some assumptions about the boundedness of the Hardy–Littlewood maximal operator in Subsection 2.3, and establish an embedding theorem from the ball quasi-Banach function space to the weighted Lebesgue space on space of homogeneous type in Subsection 2.4, which plays an important role in the remainder of this article.

2.1 Spaces of homogeneous type

In this subsection, we recall the notion of spaces of homogeneous type, and some related basic estimates.

Definition 2.1. A *quasi-metric space* (X, ρ) is a non-empty set X equipped with a *quasi-metric* ρ , namely, a non-negative function defined on $X \times X$ satisfying that, for any $x, y, z \in X$,

- (i) $\rho(x, y) = 0$ if and only if $x = y$;
- (ii) $\rho(x, y) = \rho(y, x)$;
- (iii) there exists a constant $A_0 \in [1, \infty)$, independent of x, y , and z , such that

$$\rho(x, z) \leq A_0[\rho(x, y) + \rho(y, z)]. \quad (2.1)$$

The *ball* B of X , centered at $x_0 \in X$ with radius $r \in (0, \infty)$, is defined by setting

$$B := B(x_0, r) := \{x \in X : \rho(x, x_0) < r\}.$$

For any ball B and any $\tau \in (0, \infty)$, we denote $B(x_0, \tau r)$ by τB if $B := B(x_0, r)$ for some $x_0 \in X$ and $r \in (0, \infty)$.

Definition 2.2. Let (X, ρ) be a quasi-metric space and μ a non-negative measure on X . The triple (X, ρ, μ) is called a *space of homogeneous type* if μ satisfies the following doubling condition: there exists a constant $C_{(\mu)} \in [1, \infty)$ such that, for any ball $B \subset X$,

$$\mu(2B) \leq C_{(\mu)}\mu(B). \quad (2.2)$$

The above doubling condition implies that, for any ball $B \subset X$ and any $\lambda \in [1, \infty)$,

$$\mu(\lambda B) \leq C_{(\mu)}\lambda^\omega \mu(B), \quad (2.3)$$

where $\omega := \log_2 C_{(\mu)}$ is called the *upper dimension* of X . If $A_0 = 1$, then (X, ρ, μ) is called a *metric measure space of homogeneous type* or, simply, a *doubling metric measure space*.

Both spaces of homogeneous type, with some additional assumptions, and function spaces on them have been extensively investigated in many articles. One special case of spaces of homogeneous type is the *RD-space*, originally introduced in [29] (see also [28, 74]), which is a doubling metric measure space satisfying the following additional *reverse doubling condition*:

there exist constants $\widetilde{C}_{(\mu)} \in (0, 1]$ and $\kappa \in (0, \omega]$ such that, for any ball $B(x, r)$ with $x \in X$ and $r \in (0, \text{diam } X/2)$, and any $\lambda \in [1, \text{diam } X/(2r))$,

$$\widetilde{C}_{(\mu)} \lambda^\kappa \mu(B(x, r)) \leq \mu(B(x, \lambda r)),$$

here and thereafter, $\text{diam } X := \sup_{x, y \in X} \rho(x, y)$.

Throughout this article, according to [14, pp. 587-588], we always make the following assumptions on (X, ρ, μ) :

- (i) for any point $x \in X$, the balls $\{B(x, r)\}_{r \in (0, \infty)}$ form a basis of open neighborhoods of x ;
- (ii) μ is *Borel regular* which means that all open sets are μ -measurable and every set $A \subset X$ is contained in a Borel set E such that $\mu(A) = \mu(E)$;
- (iii) for any $x \in X$ and $r \in (0, \infty)$, $\mu(B(x, r)) \in (0, \infty)$;
- (iv) $\text{diam } X = \infty$, and (X, ρ, μ) is *non-atomic*, which means $\mu(\{x\}) = 0$ for any $x \in X$.

Notice that $\text{diam } X = \infty$ implies that $\mu(X) = \infty$ (see [4, p. 284] or [55, Lemma 5.1]). From this, it follows that, under the above assumptions, $\mu(X) = \infty$ if and only if $\text{diam } X = \infty$.

The following basic estimates are from [28, Lemma 2.1], which can be proved by using (2.3).

Lemma 2.3. *Let $x, y \in X$ and $r \in (0, \infty)$. Then $V(x, y) \sim V(y, x)$ and*

$$V_r(x) + V_r(y) + V(x, y) \sim V_r(x) + V(x, y) \sim V_r(y) + V(x, y) \sim \mu(B(x, r + \rho(x, y))).$$

Moreover, if $\rho(x, y) \leq r$, then $V_r(x) \sim V_r(y)$. Here the positive equivalence constants are independent of x, y , and r .

2.2 Ball quasi-Banach function spaces

Throughout this article, let (X, ρ, μ) be a space of homogeneous type with $\mu(X) = \infty$. In this subsection, we introduce the notion of (ball) quasi-Banach function spaces on X (see [59] for the corresponding Euclidean case). Moreover, we give some examples of (ball) quasi-Banach function spaces.

Definition 2.4. Let $Y(X)$ be a quasi-Banach space consisting of μ -measurable functions on X . Then $Y(X)$ is called a *quasi-Banach function space* if it satisfies:

- (i) $\|f\|_{Y(X)} = 0$ implies that $f = 0$ almost everywhere in X ;
- (ii) $|g| \leq |f|$ almost everywhere in X implies that $\|g\|_{Y(X)} \leq \|f\|_{Y(X)}$;
- (iii) $0 \leq f_m \uparrow f$ almost everywhere in X implies that $\|f_m\|_{Y(X)} \uparrow \|f\|_{Y(X)}$;
- (iv) $\mathbf{1}_E \in Y(X)$ for any μ -measurable set $E \subset X$ with finite measure.

Moreover, a quasi-Banach function space $Y(X)$ is called a *Banach function space* if it is a Banach space and

- (v) for any given μ -measurable set $E \subset \mathcal{X}$ with finite measure, there exists a positive constant $C_{(E)}$ such that, for any $f \in Y(\mathcal{X})$,

$$\int_E |f(x)| d\mu(x) \leq C_{(E)} \|f\|_{Y(\mathcal{X})}.$$

Definition 2.5. Let $Y(\mathcal{X})$ be a quasi-Banach space consisting of μ -measurable functions on \mathcal{X} . Then $Y(\mathcal{X})$ is called a *ball quasi-Banach function space* if it satisfies (i), (ii), and (iii) of Definition 2.4, and

- (iv)' $\mathbf{1}_B \in Y(\mathcal{X})$ for any ball $B \subset \mathcal{X}$.

Moreover, a ball quasi-Banach function space $Y(\mathcal{X})$ is called a *ball Banach function space* if it is a Banach space and Definition 2.4(v) holds true with any μ -measurable set E of finite measure replaced by any ball B .

Obviously, a quasi-Banach function space is also a ball quasi-Banach function space and the converse is not necessary to be true.

Remark 2.6. (i) Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . By an elementary calculation, we know that $\|f\|_{Y(\mathcal{X})} = 0$ if and only if $f = 0$ almost everywhere in \mathcal{X} .

- (ii) Observe that, in Definition 2.5, if we replace any ball B by any bounded μ -measurable set E , we obtain its another equivalent formulation.

Next, we present several concrete examples of ball quasi-Banach function spaces on \mathcal{X} as follows.

Remark 2.7. (i) For any given μ -measurable set $E \subset \mathcal{X}$ and any given $p \in (0, \infty]$, the *Lebesgue space* $L^p(E)$ is defined by setting, when $p \in (0, \infty)$,

$$L^p(E) := \left\{ f \text{ is } \mu\text{-measurable on } E : \|f\|_{L^p(E)} := \left[\int_E |f(x)|^p d\mu(x) \right]^{1/p} < \infty \right\},$$

and

$$L^\infty(E) := \left\{ f \text{ is } \mu\text{-measurable on } E : \|f\|_{L^\infty(E)} := \operatorname{ess\,sup}_{x \in E} |f(x)| < \infty \right\}.$$

It is easy to show that, for any given μ -measurable set E and any given $p \in (0, \infty]$, $L^p(E)$ is a quasi-Banach function space and hence a ball quasi-Banach function space.

- (ii) Let $p \in (0, \infty]$ and $r \in (0, \infty]$. Recall that the *Lorentz space* $L^{p,r}(\mathcal{X})$ is defined to be the set of all μ -measurable functions f on \mathcal{X} such that

$$\|f\|_{L^{p,r}(\mathcal{X})} := \begin{cases} \left\{ \int_0^\infty \left[t^{1/p} R(f)(t) \right]^r \frac{dt}{t} \right\}^{1/r} < \infty & \text{when } r \in (0, \infty), \\ \sup_{t \in (0, \infty)} \left[t^{1/p} R(f)(t) \right] < \infty & \text{when } r = \infty, \end{cases}$$

with the usual modification made when $p = \infty$, where $R(f)$ denotes the *decreasing rearrangement function* of f defined by setting, for any $t \in (0, \infty)$,

$$R(f)(t) := \inf \{ \alpha \in (0, \infty) : d_f(\alpha) \leq t \}$$

with $d_f(\alpha) := \mu(\{x \in X : |f(x)| > \alpha\})$ for any $\alpha \in (0, \infty)$. By [22, Theorem 1.4.11], we know that, for any given $p \in (0, \infty]$ and $r \in (0, \infty]$, $L^{p,r}(X)$ is a quasi-Banach function space and hence a ball quasi-Banach function space.

- (iii) Recall that a function $\Phi : [0, \infty) \rightarrow [0, \infty)$ is called an *Orlicz function* if it is non-decreasing, $\Phi(0) = 0$, $\Phi(t) > 0$ for any $t \in (0, \infty)$, and $\lim_{t \rightarrow \infty} \Phi(t) = \infty$. The function Φ is said to be of *upper* (resp., *lower*) *type* p for some $p \in [0, \infty)$ if there exists a positive constant C such that, for any $s \in [1, \infty)$ (resp., $s \in [0, 1]$) and $t \in [0, \infty)$, $\Phi(st) \leq C s^p \Phi(t)$.

For a given function $\varphi : X \times [0, \infty) \rightarrow [0, \infty)$ such that, for almost every $x \in X$, $\varphi(x, \cdot)$ is an Orlicz function, φ is said to be of *uniformly upper* (resp., *lower*) *type* p for some $p \in [0, \infty)$ if there exists a positive constant C such that, for any $x \in X$, $s \in [1, \infty)$ (resp., $s \in [0, 1]$), and $t \in [0, \infty)$, $\varphi(x, st) \leq C s^p \varphi(x, t)$.

Let $\varphi : X \times [0, \infty) \rightarrow [0, \infty)$ satisfy that $x \mapsto \varphi(x, t)$ is μ -measurable for any $t \in [0, \infty)$. The function $\varphi(\cdot, t)$ is said to satisfy the *uniformly Muckenhoupt condition* for some $q \in [1, \infty)$, denoted by $\varphi \in \mathbb{A}_q(X)$, if, when $q \in (1, \infty)$,

$$[\varphi]_{\mathbb{A}_q(X)} := \sup_{t \in (0, \infty)} \sup_{B \subset X} \frac{1}{[\mu(B)]^q} \int_B \varphi(x, t) d\mu(x) \left\{ \int_B [\varphi(y, t)]^{-q'/q} d\mu(y) \right\}^{q/q'} < \infty,$$

or, when $q = 1$,

$$[\varphi]_{\mathbb{A}_1(X)} := \sup_{t \in (0, \infty)} \sup_{B \subset X} \frac{1}{\mu(B)} \int_B \varphi(x, t) d\mu(x) \left\{ \operatorname{ess\,sup}_{y \in B} [\varphi(y, t)]^{-1} \right\} < \infty,$$

where the second suprema are taken over all balls $B \subset X$ (see, for instance, [38]). Let

$$\mathbb{A}_\infty(X) := \bigcup_{q \in [1, \infty)} \mathbb{A}_q(X).$$

The *critical weight index* $q(\varphi)$ of $\varphi \in \mathbb{A}_\infty(X)$ is defined by setting

$$q(\varphi) := \inf \{ q \in [1, \infty) : \varphi \in \mathbb{A}_q(X) \}. \quad (2.4)$$

Recall that a function $\varphi : X \times [0, \infty) \rightarrow [0, \infty)$ is called a *growth function* if φ has the following properties:

- (a) φ is a *Musielak–Orlicz function*, namely,
 - (a)₁ the function $\varphi(x, \cdot)$ is an Orlicz function for almost every given $x \in X$;
 - (a)₂ the function $\varphi(\cdot, t)$ is a μ -measurable function on X for any given $t \in [0, \infty)$.
- (b) $\varphi \in \mathbb{A}_\infty(X)$.
- (c) φ is of uniformly lower type p for some $p \in (0, 1]$, and of uniformly upper type 1.

Let φ be a growth function. The *Musielak–Orlicz space* $L^\varphi(X)$ is defined to be the set of all μ -measurable functions f on X such that $\int_X \varphi(x, |f(x)|) d\mu(x) < \infty$ equipped with the *quasi-norm*

$$\|f\|_{L^\varphi(X)} := \inf \left\{ \lambda \in (0, \infty) : \int_X \varphi\left(x, \frac{|f(x)|}{\lambda}\right) d\mu(x) \leq 1 \right\}.$$

By the property of $\mathbb{A}_\infty(X)$, we know that, for any ball $B \subset X$, $\|\mathbf{1}_B\|_{L^\varphi(X)} < \infty$, which further implies that $\mathbf{1}_B \in L^\varphi(X)$ and hence $L^\varphi(X)$ is a ball quasi-Banach function space. However, by an argument similar to that used in [59, p. 86], we know that, if we define the growth function φ_0 by setting, for any $x \in \mathbb{R}$ and $t \in [0, \infty)$, $\varphi_0(x, t) := (1 + |x|)t$, then $L^{\varphi_0}(\mathbb{R})$ is *not* a quasi-Banach function space.

(iv) Let $p(\cdot) : X \rightarrow (0, \infty)$ be a μ -measurable function satisfying

$$0 < \widetilde{p}_- := \operatorname{ess\,inf}_{x \in X} p(x) \leq \operatorname{ess\,sup}_{x \in X} p(x) =: \widetilde{p}_+ < \infty. \quad (2.5)$$

The *variable Lebesgue space* $L^{p(\cdot)}(X)$ is defined to be the set of all μ -measurable functions f on X such that $\int_X |f(x)|^{p(x)} d\mu(x) < \infty$ and equipped with the *quasi-norm*

$$\|f\|_{L^{p(\cdot)}(X)} := \inf \left\{ \lambda \in (0, \infty) : \int_X \left[\frac{|f(x)|}{\lambda} \right]^{p(x)} d\mu(x) \leq 1 \right\}.$$

If $\widetilde{p}_- \in [1, \infty)$, then $L^{p(\cdot)}(X)$ is a Banach function space (see, for instance, [16]). From this, we deduce that $L^{p(\cdot)/\widetilde{p}_-}(X)$ is a Banach function space and hence, for any given $p(\cdot) : X \rightarrow (0, \infty)$ and any given μ -measurable set $E \subset X$ with finite measure,

$$\|\mathbf{1}_E\|_{L^{p(\cdot)}(X)} = \|\mathbf{1}_E\|_{L^{p(\cdot)/\widetilde{p}_-}(X)}^{1/\widetilde{p}_-} < \infty.$$

Thus, for any given $p(\cdot) : X \rightarrow (0, \infty)$ satisfying (2.5), $L^{p(\cdot)}(X)$ is a quasi-Banach function space and hence a ball quasi-Banach function space.

Remark 2.8. Let $Y(X)$ be a ball quasi-Banach function space on X . Then, by the Aoki–Rolewicz theorem (see [22, Exercise 1.4.6] and also [2, 56]), we find that there exists a constant $\nu \in (0, 1)$ such that, for any $N \in \mathbb{N}$ and $\{f_k\}_{k=1}^N \subset Y(X)$,

$$\left\| \sum_{k=1}^N |f_k| \right\|_{Y(X)}^\nu \leq 4 \left[\sum_{k=1}^N \|f_k\|_{Y(X)}^\nu \right].$$

The following Fatou lemma of $Y(X)$ when $X := \mathbb{R}^n$ is just [5, Chapter 1, Lemma 1.5(ii)], whose proof is a slight modification of [5, Chapter 1, Lemma 1.5(ii)]; we omit the details here.

Lemma 2.9. *Let $Y(X)$ be a ball quasi-Banach function space on X and $\{f_k\}_{k \in \mathbb{N}} \subset Y(X)$. If $f_k \rightarrow f$ almost everywhere in X as $k \rightarrow \infty$, and $\liminf_{k \rightarrow \infty} \|f_k\|_{Y(X)}$ is finite, then $f \in Y(X)$ and*

$$\|f\|_{Y(X)} \leq \liminf_{k \rightarrow \infty} \|f_k\|_{Y(X)}.$$

Next, we introduce the notion of the convexification of $Y(\mathcal{X})$.

Definition 2.10. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} , and $p \in (0, \infty)$. The p -convexification $Y^p(\mathcal{X})$ of $Y(\mathcal{X})$ is defined by setting

$$Y^p(\mathcal{X}) := \{f \text{ is } \mu\text{-measurable on } \mathcal{X} : |f|^p \in Y(\mathcal{X})\}$$

equipped with the *quasi-norm* $\|f\|_{Y^p(\mathcal{X})} := \| |f|^p \|_{Y(\mathcal{X})}^{\frac{1}{p}}$.

Remark 2.11. Let $p \in (0, \infty)$ and $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . By Definitions 2.5 and 2.10, it is easy to show that $Y^p(\mathcal{X})$ is also a ball quasi-Banach function space on \mathcal{X} ; see [11, Lemma 2.6] for the corresponding Euclidean case.

2.3 Boundedness of the Hardy–Littlewood maximal operator

Throughout this article, we always let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . In this subsection, we make two mild assumptions on the boundedness of the Hardy–Littlewood maximal operator on $Y(\mathcal{X})$ and its associate space. We first recall that the *Hardy–Littlewood maximal operator* \mathcal{M} is defined by setting, for any μ -measurable function f and any $x \in \mathcal{X}$,

$$\mathcal{M}(f)(x) := \sup_{B \ni x} \frac{1}{\mu(B)} \int_B |f(y)| d\mu(y), \quad (2.6)$$

where the supremum is taken over all balls B of \mathcal{X} containing x . For any given $\theta \in (0, \infty)$, the *powered Hardy–Littlewood maximal operator* $\mathcal{M}^{(\theta)}$ is defined by setting, for any μ -measurable function f and any $x \in \mathcal{X}$,

$$\mathcal{M}^{(\theta)}(f)(x) := \left\{ \mathcal{M}(|f|^\theta)(x) \right\}^{\frac{1}{\theta}}.$$

Throughout this article, we denote by $M(\mathcal{X})$ the *set of all μ -measurable functions* on \mathcal{X} . For any ball Banach function space $Y(\mathcal{X})$, its *associate space (Köthe dual)* $Y'(\mathcal{X})$ is defined by setting

$$Y'(\mathcal{X}) := \{f \in M(\mathcal{X}) : \|f\|_{Y'(\mathcal{X})} < \infty\}, \quad (2.7)$$

where, for any $f \in M(\mathcal{X})$,

$$\|f\|_{Y'(\mathcal{X})} := \sup \left\{ \|fg\|_{L^1(\mathcal{X})} : g \in Y(\mathcal{X}), \|g\|_{Y(\mathcal{X})} = 1 \right\}.$$

In this article, we need the following mild assumptions about the boundedness of \mathcal{M} on the ball quasi-Banach function space under consideration, and its associate space.

Assumption 2.12. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . Assume that there exists a $p_- \in (0, \infty)$ such that, for any given $p \in (0, p_-)$ and $r \in (1, \infty)$, there exists a positive constant C such that, for any $\{f_k\}_{k=1}^\infty \subset M(\mathcal{X})$,

$$\left\| \left\{ \sum_{k=1}^\infty [\mathcal{M}(f_k)]^r \right\}^{\frac{1}{r}} \right\|_{Y^{\frac{1}{p}}(\mathcal{X})} \leq C \left\| \left\{ \sum_{k=1}^\infty |f_k|^r \right\}^{\frac{1}{r}} \right\|_{Y^{\frac{1}{p}}(\mathcal{X})}.$$

In what follows, for any given $p_- \in (0, \infty)$, let

$$\underline{p} := \min\{p_-, 1\}. \quad (2.8)$$

The following lemma is used throughout this article, which is a trivial corollary of (2.6); we omit the details here.

Lemma 2.13. *For any $h \in (0, \infty)$, any $\alpha \in [1, \infty)$, any ball $B \subset X$, and any $x \in X$, it holds true that*

$$\mathbf{1}_{\alpha B}(x) \leq \left[\frac{\mu(\alpha B)}{\mu(B)} \right]^h [\mathcal{M}(\mathbf{1}_B)(x)]^h.$$

Remark 2.14. Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12, $\alpha \in [1, \infty)$, and $h \in (0, \underline{p})$ with \underline{p} as in (2.8). Then, by Lemma 2.13, (2.3), and Assumption 2.12, we conclude that there exists a positive constant C , independent of α , such that, for any sequence $\{B_k\}_{k \in \mathbb{N}}$ of balls of X ,

$$\begin{aligned} \left\| \sum_{k \in \mathbb{N}} \mathbf{1}_{\alpha B_k} \right\|_{Y(X)} &\leq C \alpha^{\frac{\omega}{h}} \left\| \sum_{k \in \mathbb{N}} [\mathcal{M}(\mathbf{1}_{B_k})]^{\frac{1}{h}} \right\|_{Y(X)} \\ &= C \alpha^{\frac{\omega}{h}} \left\| \left\{ \sum_{k \in \mathbb{N}} [\mathcal{M}(\mathbf{1}_{B_k})]^{\frac{1}{h}} \right\}^h \right\|_{Y^{\frac{1}{h}}(X)}^{\frac{1}{h}} \\ &\leq C \alpha^{\frac{\omega}{h}} \left\| \sum_{k \in \mathbb{N}} \mathbf{1}_{B_k} \right\|_{Y(X)}. \end{aligned}$$

Assumption 2.15. Let $p_- \in (0, \infty)$ and $Y(X)$ be a ball quasi-Banach function space on X . Assume that there exist a $\theta_0 \in (0, \underline{p})$ with \underline{p} as in (2.8), and a $p_0 \in (\theta_0, \infty)$ such that $Y^{1/\theta_0}(X)$ is a ball Banach function space and, for any $f \in (Y^{1/\theta_0})'(X)$,

$$\left\| \mathcal{M}^{((p_0/\theta_0)')}(f) \right\|_{(Y^{1/\theta_0})'(X)} \leq C \|f\|_{(Y^{1/\theta_0})'(X)}, \quad (2.9)$$

where C is a positive constant independent of f , and $\frac{1}{p_0/\theta_0} + \frac{1}{(p_0/\theta_0)'} = 1$.

Remark 2.16. Let $\theta_1, \theta_2 \in (0, \infty)$ satisfy $\theta_1 < \theta_2$. By the Hölder inequality, we know that, for any μ -measurable function f and any $x \in X$,

$$\begin{aligned} \mathcal{M}^{(\theta_1)}(f)(x) &= \sup_{B \ni x} \left\{ \frac{1}{\mu(B)} \int_B |f(y)|^{\theta_1} d\mu(y) \right\}^{\frac{1}{\theta_1}} \\ &\leq \sup_{B \ni x} \left\{ \frac{1}{\mu(B)} \left[\int_B |f(y)|^{\theta_2} d\mu(y) \right]^{\frac{\theta_1}{\theta_2}} [\mu(B)]^{1 - \frac{\theta_1}{\theta_2}} \right\}^{\frac{1}{\theta_1}} \\ &= \sup_{B \ni x} \left\{ \frac{1}{\mu(B)} \int_B |f(y)|^{\theta_2} d\mu(y) \right\}^{\frac{1}{\theta_2}} = \mathcal{M}^{(\theta_2)}(f)(x). \end{aligned} \quad (2.10)$$

Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} satisfying Assumption 2.15 with some $\theta_0 \in (0, \underline{p})$ and $p_0 \in (\theta_0, \infty)$. Then, by (2.10) and Definition 2.4(ii), we conclude that there exists a positive constant C such that, for any given $p_1 \in (p_0, \infty]$ and any $f \in (Y^{1/\theta_0})'(\mathcal{X})$,

$$\|\mathcal{M}^{((p_1/\theta_0)')} (f)\|_{(Y^{1/\theta_0})'(\mathcal{X})} \leq \|\mathcal{M}^{((p_0/\theta_0)')} (f)\|_{(Y^{1/\theta_0})'(\mathcal{X})} \leq C\|f\|_{(Y^{1/\theta_0})'(\mathcal{X})}.$$

2.4 Embedding theorem for ball quasi-Banach function spaces

In this subsection, we aim to embed $Y(\mathcal{X})$ continuously into the weighted Lebesgue space on \mathcal{X} . To this end, we first recall some notions on weighted Lebesgue spaces, and then state some basic properties of $Y(\mathcal{X})$.

Throughout this article, for any given $p \in (0, \infty)$, a function f is said to be *locally p -integrable* if, for any $x \in \mathcal{X}$, there exists an $r \in (0, \infty)$ such that

$$\int_{B(x,r)} |f(y)|^p d\mu(y) < \infty.$$

Denote by $L_{\text{loc}}^p(\mathcal{X})$ the set of all *locally p -integrable functions* on \mathcal{X} .

Let $q \in [1, \infty)$ and $w \in L_{\text{loc}}^1(\mathcal{X})$ be a nonnegative function. Then w is said to be a *Muckenhoupt A_q -weight*, denoted by $w \in A_q(\mathcal{X})$, if

$$[w]_{A_q(\mathcal{X})} := \sup_{B \subset \mathcal{X}} \frac{1}{[\mu(B)]^q} \int_B w(x) d\mu(x) \left\{ \int_B [w(y)]^{-q'/q} d\mu(y) \right\}^{q/q'} < \infty \quad (2.11)$$

when $q \in (1, \infty)$, or

$$[w]_{A_1(\mathcal{X})} := \sup_{B \subset \mathcal{X}} \frac{1}{\mu(B)} \int_B w(x) d\mu(x) \left(\text{ess sup}_{y \in B} [w(y)]^{-1} \right) < \infty,$$

where the suprema are taken over all balls $B \subset \mathcal{X}$ (see, for instance, [65]). Let

$$A_\infty(\mathcal{X}) := \bigcup_{q \in [1, \infty)} A_q(\mathcal{X}).$$

Recall that, for any given $p \in (0, \infty)$ and $w \in A_\infty(\mathcal{X})$, the *weighted Lebesgue space* $L_w^p(\mathcal{X})$ is defined to be the set of all μ -measurable functions f on \mathcal{X} such that

$$\|f\|_{L_w^p(\mathcal{X})} := \left\{ \int_{\mathcal{X}} |f(x)|^p w(x) d\mu(x) \right\}^{1/p} < \infty.$$

Obviously, $L_w^p(\mathcal{X})$ is a ball quasi-Banach function space, which even when $\mathcal{X} := \mathbb{R}^n$ may not be a quasi-Banach function space (see, for instance, [59, p. 86]).

Next, we state an embedding theorem from $Y(\mathcal{X})$ to the weighted Lebesgue space (see [11, Lemma 4.7] for the corresponding Euclidean case).

Theorem 2.17. *Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . Assume that there exists an $r \in (0, \infty)$ such that $Y^{1/r}(\mathcal{X})$ is a ball Banach function space and that \mathcal{M} is bounded on $(Y^{1/r})'(\mathcal{X})$. Fix $x_0 \in \mathcal{X}$. Then there exists an $\epsilon \in (0, 1)$ such that $Y(\mathcal{X})$ is continuously embedded into $L_w^r(\mathcal{X})$ with $w := [\mathcal{M}(\mathbf{1}_{B(x_0,1)})]^\epsilon \in A_1(\mathcal{X})$. Moreover, there exists a positive constant C , depending on x_0 , such that, for any $f \in Y(\mathcal{X})$,*

$$\|f\|_{L_w^r(\mathcal{X})} \leq C\|f\|_{Y(\mathcal{X})}. \quad (2.12)$$

Theorem 2.17 is the main result of this subsection. To prove it, we borrow some ideas from the proof of [11, Lemma 4.7] and establish several technical lemmas, some of which are used again in later sections. By Definition 2.4(i) and (2.7), we easily obtain the following Hölder inequality on $Y(X)$ (see [5, Chapter 1, Theorem 2.4] for the corresponding Euclidean case), whose proof is a slight modification of [5, Chapter 1, Theorem 2.4]; we omit the details here.

Lemma 2.18. *Let $Y(X)$ be a ball Banach function space on X , and $Y'(X)$ the associate space of $Y(X)$. If $f \in Y(X)$ and $g \in Y'(X)$, then fg is integrable and*

$$\int_X |f(x)g(x)| d\mu(x) \leq \|f\|_{Y(X)} \|g\|_{Y'(X)}.$$

For the corresponding Euclidean case, the following lemma is just [59, Proposition 2.3], whose proof is a slight modification of [59, Proposition 2.3]; we omit the details here.

Lemma 2.19. *Let $Y(X)$ be a ball Banach function space on X . Then its associate space $Y'(X)$ is also a ball Banach function space.*

By a method similar to that used in [5, Chapter 1, Theorem 2.7], and Lemmas 2.18 and 2.19, we have the following conclusion, which is a generalization of [76, Lemma 2.6] on \mathbb{R}^n to X ; we omit the details here.

Lemma 2.20. *Every ball Banach function space $Y(X)$ coincides with its second associate space $Y''(X)$. More precisely, a function f belongs to $Y(X)$ if and only if it belongs to $Y''(X)$. Moreover, for any $f \in Y(X)$ or $f \in Y''(X)$,*

$$\|f\|_{Y(X)} = \|f\|_{Y''(X)}.$$

The following lemma is a generalization of [59, Lemma 2.15(ii)] on \mathbb{R}^n to X , whose proof is also valid in this setting; we omit the details here.

Lemma 2.21. *Assume that $Y(X)$ is a ball quasi-Banach function space on X , and \mathcal{M} bounded on $Y(X)$. Then there exists an $\eta \in (1, \infty)$ such that $\mathcal{M}^{(\eta)}$ is bounded on $Y(X)$.*

The following lemma establishes the pointwise estimates of $\mathcal{M}(\mathbf{1}_{B(x_0,1)})$ with $x_0 \in X$ and \mathcal{M} as in (2.6), which can be regarded as a generalization of [22, (2.1.6)] on \mathbb{R}^n to X .

Lemma 2.22. *Let $x_0 \in X$ and \mathcal{M} be as in (2.6). Then there exists a constant $C \in [1, \infty)$, independent of x_0 , such that, for any $x \in X$,*

$$C^{-1} \min \left\{ 1, \frac{V_1(x_0)}{V(x_0, x)} \right\} \leq \mathcal{M}(\mathbf{1}_{B(x_0,1)})(x) \leq C \min \left\{ 1, \frac{V_1(x_0)}{V(x_0, x)} \right\}. \quad (2.13)$$

Proof. Let all the symbols be as in the present lemma. To obtain (2.13), we consider two cases on $\rho(x_0, x)$.

Case 1) $\rho(x_0, x) \in [0, 2A_0)$. In this case, it suffices to show that $\mathcal{M}(\mathbf{1}_{B(x_0,1)})(x) \sim 1$. On one hand, by (2.6), we have

$$\mathcal{M}(\mathbf{1}_{B(x_0,1)})(x) \leq 1.$$

On the other hand, by $\rho(x_0, x) < 2A_0$, (2.6), and (2.3), we conclude that

$$\mathcal{M}(\mathbf{1}_{B(x_0,1)})(x) \geq \frac{1}{\mu(B(x_0, 2A_0))} \int_{B(x_0, 2A_0)} \mathbf{1}_{B(x_0,1)}(y) d\mu(y) = \frac{\mu(B(x_0, 1))}{\mu(B(x_0, 2A_0))} \gtrsim 1.$$

Combining the above two inequalities, we obtain the desired estimates in this case.

Case 2) $\rho(x_0, x) \in [2A_0, \infty)$. In this case, we only need to show that

$$\mathcal{M}(\mathbf{1}_{B(x_0, 1)})(x) \sim \frac{V_1(x_0)}{V(x_0, x)}. \quad (2.14)$$

To this end, we first claim that

$$B(x_0, 1) \cap B(x, \rho(x_0, x)/[2A_0]) = \emptyset. \quad (2.15)$$

Indeed, if there exists a $y \in B(x_0, 1) \cap B(x, \rho(x_0, x)/[2A_0])$, then, by the quasi-triangle inequality and $\rho(x_0, x) \geq 2A_0$, we conclude that

$$\rho(x_0, x) \leq A_0[\rho(x_0, y) + \rho(y, x)] < A_0 \left[1 + \frac{\rho(x_0, x)}{2A_0} \right] \leq \rho(x_0, x),$$

which makes a contradiction. Thus, (2.15) holds true. On one hand, by (2.15), (2.3), and Lemma 2.3, we obtain

$$\begin{aligned} \mathcal{M}(\mathbf{1}_{B(x_0, 1)})(x) &\sim \sup_{r \in (0, \infty)} \frac{1}{\mu(B(x, r))} \int_{B(x, r)} \mathbf{1}_{B(x_0, 1)}(y) d\mu(y) \\ &\sim \sup_{r \in (\rho(x_0, x)/[2A_0], \infty)} \frac{\mu(B(x, r) \cap B(x_0, 1))}{\mu(B(x, r))} \\ &\lesssim \frac{\mu(B(x_0, 1))}{\mu(B(x, \rho(x_0, x)/[2A_0]))} \sim \frac{\mu(B(x_0, 1))}{\mu(B(x_0, \rho(x_0, x)))} \sim \frac{V_1(x_0)}{V(x_0, x)}. \end{aligned} \quad (2.16)$$

On the other hand, observing that $x \in B(x_0, 2\rho(x_0, x))$, from this, (2.6), and (2.3), it follows that

$$\begin{aligned} \mathcal{M}(\mathbf{1}_{B(x_0, 1)})(x) &\geq \frac{1}{\mu(B(x_0, 2\rho(x_0, x)))} \int_{B(x_0, 2\rho(x_0, x))} \mathbf{1}_{B(x_0, 1)}(y) d\mu(y) \\ &= \frac{\mu(B(x_0, 1))}{\mu(B(x_0, 2\rho(x_0, x)))} \sim \frac{V_1(x_0)}{V(x_0, x)}, \end{aligned}$$

which, combined with (2.16), further implies (2.14). Thus, we find that (2.13) also holds true in this case. This finishes the proof of Lemma 2.22. \square

By borrowing some ideas from [25], we establish the following lemma, which is regarded as a substitute of the reverse doubling condition and can be used to deal with integrals only in terms of measures of balls.

Lemma 2.23. *Let X be a space of homogeneous type with $\mu(X) = \infty$, $c \in (1, \infty)$, $x_0 \in X$, and $r_0 \in (0, \infty)$. Then there exists a sequence $\{r_k\}_{k=1}^\infty \subset (0, \infty)$ such that, for any $k \in \mathbb{N}$,*

$$(i) \quad r_k \geq 2r_{k-1};$$

$$(ii) \quad c\mu(B(x_0, r_{k-1})) < \mu(B(x_0, r_k)) \leq cC_{(\mu)}\mu(B(x_0, r_{k-1})), \text{ where } C_{(\mu)} \in (1, \infty) \text{ is as in (2.2).}$$

Proof. Let all the symbols be as in the present lemma. We use an inductive method to construct $\{r_k\}_{k=1}^\infty$. For any fixed $k \in \mathbb{N}$, if $r_{k-1} \in (0, \infty)$ has been constructed, we then let

$$A := \left\{ r \in (0, \infty) : \mu(B(x_0, r)) \leq cC_{(\mu)}\mu(B(x_0, r_{k-1})) \right\} \quad \text{and} \quad r_k := \sup A.$$

First, by $r_{k-1} \in A$, we know that $r_k \geq r_{k-1}$ and hence $r_k > 0$. Second, if $r_k = \infty$, then, from the definition of r_k , $\mu(X) = \infty$, and $r_{k-1} < \infty$, we infer that there exists an increasing sequence $\{\tau_n\}_{n=1}^\infty \subset A$ such that $\lim_{n \rightarrow \infty} \tau_n = \infty$ and

$$\infty = \lim_{n \rightarrow \infty} \mu(B(x_0, \tau_n)) \leq cC_{(\mu)}\mu(B(x_0, r_{k-1})) < \infty,$$

which makes a contradiction. Thus, $r_k < \infty$. Finally, by (2.2), we find that $2r_{k-1} \in A$ which implies that $r_k \geq 2r_{k-1}$. This proves (i).

Now, we prove (ii). Indeed, by the definition of r_k , we know that there exists an increasing sequence $\{\tau_n\}_{n=1}^\infty \subset A$ such that $\lim_{n \rightarrow \infty} \tau_n = r_k$, which further implies that

$$\mu(B(x_0, r_k)) = \lim_{n \rightarrow \infty} \mu(B(x_0, \tau_n)) \leq cC_{(\mu)}\mu(B(x_0, r_{k-1})).$$

On the other hand, note that $2r_k > r_k$. Thus, $2r_k \notin A$. From this and (2.2), we deduce that

$$\mu(B(x_0, r_k)) \geq C_{(\mu)}^{-1}\mu(B(x_0, 2r_k)) > c\mu(B(x_0, r_{k-1})).$$

These two inequalities then complete the proof of (ii) and hence of Lemma 2.23. \square

Definition 2.24. Let X be a space of homogeneous type with $\mu(X) = \infty$, $c \in (1, \infty)$, $x_0 \in X$, and $r_0 \in (0, \infty)$. A sequence $\{B_k\}_{k \in \mathbb{Z}_+} := \{B(x_0, r_k)\}_{k \in \mathbb{Z}_+}$ of balls centered at x_0 , associated with μ , c , and r_0 , is said to be *admissible* if $\{r_k\}_{k \in \mathbb{Z}_+}$ satisfy (i) and (ii) of Lemma 2.23.

Remark 2.25. (i) Obviously, for any given μ and c as in Definition 2.24, and any $x_0 \in X$ and $r_0 \in (0, \infty)$, Lemma 2.23 guarantees that there exists an admissible sequence $\{B_k\}_{k \in \mathbb{Z}_+}$ of balls centered at x_0 , associated with μ , c , and r_0 .

(ii) Observe that any admissible sequence $\{B_k\}_{k \in \mathbb{Z}_+}$ of balls satisfies both the doubling and the reverse doubling conditions, which is the key tool used in the proof of Theorem 2.17 to escape the dependence on the reverse doubling assumption of the measure μ under consideration. We believe that more applications are expectable.

Now, we give the proof of Theorem 2.17.

Proof of Theorem 2.17. Let all the symbols be as in the present theorem. By Lemma 2.19, we know that $(Y^{1/r})'(X)$ is a ball Banach function space. From this, the fact that \mathcal{M} is bounded on $(Y^{1/r})'(X)$, and Lemma 2.21, it follows that there exists an $\eta \in (1, \infty)$ such that $\mathcal{M}^{(\eta)}$ is bounded on $(Y^{1/r})'(X)$. Let $\epsilon \in (1/\eta, 1)$ and $w := [\mathcal{M}(\mathbf{1}_{B(x_0, 1)})]^\epsilon$. We first prove that $w \in A_1(X)$. To this end, from the definition of $A_1(X)$, it suffices to show that there exists a positive constant C such that, for any $B := B(x_B, r_B) \subset X$ with $x_B \in X$ and $r_B \in (0, \infty)$, and almost every $x \in B$,

$$\frac{1}{\mu(B)} \int_B w(y) d\mu(y) \leq Cw(x). \quad (2.17)$$

Decompose $\mathbf{1}_{B(x_0,1)}$ as

$$\mathbf{1}_{B(x_0,1)} = \mathbf{1}_{B(x_0,1) \cap 2A_0B} + \mathbf{1}_{B(x_0,1) \cap (2A_0B)^c} =: f_1 + f_2. \quad (2.18)$$

On one hand, by the boundedness of \mathcal{M} from $L^1(X)$ to $L^{1,\infty}(X)$ (see, for instance, [14, Theorem (3.5)]), (2.2), $\epsilon < 1$, and (2.6), we know that, for any $x \in B$,

$$\begin{aligned} & \frac{1}{\mu(B)} \int_B [\mathcal{M}(f_1)(y)]^\epsilon d\mu(y) \\ &= \frac{\epsilon}{\mu(B)} \int_0^\infty \lambda^{\epsilon-1} \mu(\{y \in B : \mathcal{M}(f_1)(y) > \lambda\}) d\lambda \\ &\lesssim \frac{\epsilon}{\mu(B)} \int_0^\infty \lambda^{\epsilon-1} \min\left\{\mu(B), \frac{\|f_1\|_{L^1(X)}}{\lambda}\right\} d\lambda \\ &\lesssim \frac{\epsilon}{\mu(B)} \left[\int_0^{\|f_1\|_{L^1(X)}/\mu(B)} \lambda^{\epsilon-1} \mu(B) d\lambda + \int_{\|f_1\|_{L^1(X)}/\mu(B)}^\infty \lambda^{\epsilon-2} \|f_1\|_{L^1(X)} d\lambda \right] \\ &\lesssim \|f_1\|_{L^1(X)}^\epsilon [\mu(B)]^{-\epsilon} \lesssim \left[\frac{1}{\mu(2A_0B)} \int_{2A_0B} \mathbf{1}_{B(x_0,1)}(y) d\mu(y) \right]^\epsilon \lesssim w(x), \end{aligned} \quad (2.19)$$

where the implicit positive constants are independent of B and x .

On the other hand, let $z \in B$ and $\tilde{B} := B(x_{\tilde{B}}, r_{\tilde{B}}) \subset X$ be such that $z \in \tilde{B}$ and $\int_{\tilde{B}} f_2(x) d\mu(x) > 0$. Then

$$r_B < 2A_0 r_{\tilde{B}}. \quad (2.20)$$

Otherwise, by the fact that $\int_{\tilde{B}} f_2(x) d\mu(x) > 0$, we know that there exists a $u \in \tilde{B} \cap (2A_0B)^c$, and hence

$$2A_0 r_B \leq \rho(u, x_B) \leq A_0[\rho(u, z) + \rho(z, x_B)] < A_0 \rho(u, z) + A_0 r_B,$$

which further implies that

$$r_B < \rho(u, z) \leq A_0[\rho(u, x_{\tilde{B}}) + \rho(x_{\tilde{B}}, z)] < 2A_0 r_{\tilde{B}} \leq r_B.$$

This is a contradiction and hence (2.20) holds true. By (2.20), we conclude that, for any $v \in B$,

$$\begin{aligned} \rho(v, x_{\tilde{B}}) &\leq A_0[\rho(v, x_B) + \rho(x_B, x_{\tilde{B}})] \leq A_0\{\rho(v, x_B) + A_0[\rho(x_B, z) + \rho(z, x_{\tilde{B}})]\} \\ &< (A_0 + A_0^2)r_B + A_0^2 r_{\tilde{B}} < (2A_0 + 3)A_0^2 \tilde{B}. \end{aligned}$$

This further implies that $v \in (2A_0 + 3)A_0^2 \tilde{B}$, and hence $B \subset (2A_0 + 3)A_0^2 \tilde{B}$ by the arbitrariness of $v \in B$. From this and (2.3), we deduce that, for any $x \in B$,

$$\frac{1}{\mu(\tilde{B})} \int_{\tilde{B}} f_2(y) d\mu(y) \lesssim \frac{1}{\mu((2A_0 + 3)A_0^2 \tilde{B})} \int_{(2A_0 + 3)A_0^2 \tilde{B}} f_2(y) d\mu(y) \lesssim \mathcal{M}(\mathbf{1}_{B(x_0,1)})(x).$$

Thus, by the arbitrariness of \tilde{B} , we find that, for any $x \in B$,

$$\mathcal{M}(f_2)(z) \lesssim \mathcal{M}(\mathbf{1}_{B(x_0,1)})(x),$$

which further implies that

$$\frac{1}{\mu(B)} \int_B [\mathcal{M}(f_2)(z)]^\epsilon d\mu(z) \lesssim w(x), \quad (2.21)$$

where the implicit positive constant is independent of B and x .

Combining (2.18), (2.19), and (2.21), we conclude that, for any $B \subset \mathcal{X}$ and $x \in B$,

$$\frac{1}{\mu(B)} \int_B w(y) d\mu(y) \lesssim \frac{1}{\mu(B)} \int_B \{[\mathcal{M}(f_1)(y)]^\epsilon + [\mathcal{M}(f_2)(y)]^\epsilon\} d\mu(y) \lesssim w(x).$$

This proves (2.17) and hence $w \in A_1(\mathcal{X})$.

Next, we prove (2.12). Let $r_0 = 1$, $c \in (1, \infty)$, x_0 be as in the present theorem, and $\{B_k\}_{k \in \mathbb{Z}_+}$ an admissible sequence of balls as in Definition 2.24. From Lemma 2.23(i), we deduce that, for any $f \in Y(\mathcal{X})$,

$$\begin{aligned} \|f\|_{L_w^r(\mathcal{X})}^r &= \int_{\mathcal{X}} |f(x)|^r w(x) d\mu(x) \\ &= \int_{B_0} |f(x)|^r w(x) d\mu(x) + \sum_{k=1}^{\infty} \int_{B_k \setminus B_{k-1}} \cdots \\ &=: I_1 + I_2. \end{aligned} \quad (2.22)$$

We first estimate I_1 . Indeed, by Lemmas 2.22, 2.18, and 2.19, and Definitions 2.10 and 2.4(iv), we know that

$$I_1 \lesssim \int_{B_0} |f(x)|^r d\mu(x) \lesssim \| |f|^r \|_{(Y^{\frac{1}{r}})'(\mathcal{X})} \| \mathbf{1}_{B_0} \|_{(Y^{\frac{1}{r}})'(\mathcal{X})} \sim \|f\|_{Y(\mathcal{X})}^r. \quad (2.23)$$

Next, we estimate I_2 . By Lemmas 2.22 and 2.23(ii), we know that, for any $k \in \mathbb{N}$ and $x \in B_k \setminus B_{k-1}$,

$$w(x) \lesssim \left[\frac{\mu(B_0)}{\mu(B_{k-1})} \right]^\epsilon \sim \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^\epsilon. \quad (2.24)$$

On the other hand, from the definition of $\mathcal{M}^{(\eta)}$, we deduce that, for any $k \in \mathbb{N}$ and $x \in B_k$,

$$\mathcal{M}^{(\eta)}(\mathbf{1}_{B_0})(x) \geq \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^{1/\eta},$$

which further implies that, for any $x \in \mathcal{X}$,

$$\mathbf{1}_{B_k}(x) \leq \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^{-1/\eta} \mathcal{M}^{(\eta)}(\mathbf{1}_{B_0})(x).$$

By this, (2.24), Lemma 2.18, Definition 2.10, the boundedness of $\mathcal{M}^{(\eta)}$ on $(Y^{1/r})'(\mathcal{X})$, Lemmas 2.19 and 2.23(ii), Definition 2.4(iv), and $\epsilon > 1/\eta$, we conclude that

$$I_2 \lesssim \sum_{k=1}^{\infty} \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^\epsilon \int_{B_k} |f(x)|^r d\mu(x) \lesssim \sum_{k=1}^{\infty} \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^\epsilon \| \mathbf{1}_{B_k} \|_{(Y^{\frac{1}{r}})'(\mathcal{X})} \|f\|_{Y(\mathcal{X})}^r$$

$$\lesssim \sum_{k=1}^{\infty} \left[\frac{\mu(B_0)}{\mu(B_k)} \right]^{\epsilon-1/\eta} \|\mathbf{1}_{B_0}\|_{(Y^\perp)^\gamma(X)} \|f\|_{Y(X)}^r \lesssim \|f\|_{Y(X)}^r \sum_{k=1}^{\infty} c^{-(k-1)(\epsilon-1/\eta)} \lesssim \|f\|_{Y(X)}^r.$$

This, combined with (2.22) and (2.23), implies (2.12) and hence finishes the proof of Theorem 2.17. \square

Remark 2.26. We point out that the proof of $w \in A_1(X)$ in Theorem 2.17 is a slight modification of the proof of [3, Lemma 3.3]. We still give some details here for the sake of completeness.

3 Maximal function characterizations of $H_Y^*(X)$

In this section, we first introduce the Hardy space $H_Y^*(X)$ associated with ball quasi-Banach function space $Y(X)$ via using the grand maximal function, then discuss the relations between $Y(X)$ and $H_Y^*(X)$, and finally establish the radial and the non-tangential maximal function characterizations of $H_Y^*(X)$.

3.1 Relations between $Y(X)$ and $H_Y^*(X)$

The aim of this subsection is to clarify the relations between $Y(X)$ and $H_Y^*(X)$. To this end, we first recall the notions of the spaces of test functions and distributions on X , which were originally introduced by Han et al. [28, Definition 2.2] (see also [29, Definition 2.8]).

Definition 3.1 (test functions). Let $x_0 \in X$, $r \in (0, \infty)$, $\beta \in (0, 1]$, and $\gamma \in (0, \infty)$. A function f on X is called a *test function of type* (x_0, r, β, γ) , denoted by $f \in \mathcal{G}(x_0, r, \beta, \gamma)$, if there exists a positive constant C such that

(i) (the *size condition*) for any $x \in X$,

$$|f(x)| \leq CP_\gamma(x_0, x; r), \quad (3.1)$$

here and thereafter, $P_\gamma(x_0, x; r)$ is as in (1.1);

(ii) (the *regularity condition*) for any $x, y \in X$ satisfying $\rho(x, y) \leq (2A_0)^{-1}[r + \rho(x_0, x)]$,

$$|f(x) - f(y)| \leq C \left[\frac{\rho(x, y)}{r + \rho(x_0, x)} \right]^\beta P_\gamma(x_0, x; r). \quad (3.2)$$

For any $f \in \mathcal{G}(x_0, r, \beta, \gamma)$, its *norm* $\|f\|_{\mathcal{G}(x_0, r, \beta, \gamma)}$ in $\mathcal{G}(x_0, r, \beta, \gamma)$ is defined by setting

$$\|f\|_{\mathcal{G}(x_0, r, \beta, \gamma)} := \inf \{C \in (0, \infty) : C \text{ satisfies (3.1) and (3.2)}\}.$$

Fix $x_0 \in X$. We denote $\mathcal{G}(x_0, 1, \beta, \gamma)$ simply by $\mathcal{G}(\beta, \gamma)$. Obviously, $\mathcal{G}(\beta, \gamma)$ is a Banach space. Note that, for any fixed $x \in X$ and $r \in (0, \infty)$, $\mathcal{G}(x, r, \beta, \gamma) = \mathcal{G}(\beta, \gamma)$ with equivalent norms, but the positive equivalence constants may depend on x and r .

Fix $\epsilon \in (0, 1]$ and $\beta, \gamma \in (0, \epsilon]$. Let $\mathcal{G}_0^\epsilon(\beta, \gamma)$ be the completion of the set $\mathcal{G}(\epsilon, \epsilon)$ in $\mathcal{G}(\beta, \gamma)$. Furthermore, the norm of $\mathcal{G}_0^\epsilon(\beta, \gamma)$ is defined by setting $\|\cdot\|_{\mathcal{G}_0^\epsilon(\beta, \gamma)} := \|\cdot\|_{\mathcal{G}(\beta, \gamma)}$. The space $\mathcal{G}_0^\epsilon(\beta, \gamma)$ is called the *space of test functions*. The *dual space* $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ is defined to be the set of all

continuous linear functionals from $\mathcal{G}_0^\epsilon(\beta, \gamma)$ to \mathbb{C} , equipped with the weak-* topology. The space $(\mathcal{G}_0^\epsilon(\beta, \gamma))'$ is called the *space of distributions*.

The following system of dyadic cubes of (X, ρ, μ) was established by Hytönen and Kairema in [40, Theorem 2.2].

Lemma 3.2. *Suppose that constants $0 < c_0 \leq C_0 < \infty$ and $\delta \in (0, 1)$ satisfy $12A_0^3 C_0 \delta \leq c_0$. Assume that a set of points, $\{z_\alpha^k : k \in \mathbb{Z}, \alpha \in \mathcal{A}_k\} \subset X$ with \mathcal{A}_k , for any $k \in \mathbb{Z}$, being a set of indices, has the following properties: for any $k \in \mathbb{Z}$,*

$$\rho(z_\alpha^k, z_\beta^k) \geq c_0 \delta^k \quad \text{if } \alpha \neq \beta, \quad \text{and} \quad \min_{\alpha \in \mathcal{A}_k} \rho(x, z_\alpha^k) < C_0 \delta^k \quad \text{for any } x \in X.$$

Then there exists a family of sets, $\{Q_\alpha^k : k \in \mathbb{Z}, \alpha \in \mathcal{A}_k\}$, satisfying

- (i) for any $k \in \mathbb{Z}$, $\bigcup_{\alpha \in \mathcal{A}_k} Q_\alpha^k = X$ and $\{Q_\alpha^k : \alpha \in \mathcal{A}_k\}$ is disjoint for any given $k \in \mathbb{Z}$;
- (ii) if $k, l \in \mathbb{Z}$ and $k \leq l$, then, for any $\alpha \in \mathcal{A}_k$ and $\beta \in \mathcal{A}_l$, either $Q_\beta^l \subset Q_\alpha^k$ or $Q_\beta^l \cap Q_\alpha^k = \emptyset$;
- (iii) for any $k \in \mathbb{Z}$ and $\alpha \in \mathcal{A}_k$, $B(z_\alpha^k, (3A_0^2)^{-1} c_0 \delta^k) \subset Q_\alpha^k \subset B(z_\alpha^k, 2A_0 C_0 \delta^k)$.

Throughout this article, for any $k \in \mathbb{Z}$, define

$$\mathcal{G}_k := \mathcal{A}_{k+1} \setminus \mathcal{A}_k \quad \text{and} \quad \mathcal{Y}^k := \{z_\alpha^{k+1}\}_{\alpha \in \mathcal{G}_k} =: \{y_\alpha^k\}_{\alpha \in \mathcal{G}_k} \quad (3.3)$$

and, for any $x \in X$, define

$$\rho(x, \mathcal{Y}^k) := \inf_{y \in \mathcal{Y}^k} \rho(x, y). \quad (3.4)$$

Next, we recall the notion of approximations of the identity with exponential decay which was introduced in [31, Definition 2.7].

Definition 3.3. Let δ be as in Lemma 3.2. A sequence $\{Q_k\}_{k \in \mathbb{Z}}$ of bounded linear integral operators on $L^2(X)$ is called an *approximation of the identity with exponential decay* (for short, exp-ATI) if there exist constants $C, \nu \in (0, \infty)$, $a \in (0, 1]$, and $\eta \in (0, 1)$ such that, for any $k \in \mathbb{Z}$, the kernel of the operator Q_k , a function on $X \times X$, which is still denoted by Q_k , has the following properties:

- (i) (the *identity condition*) $\sum_{k \in \mathbb{Z}} Q_k = I$ in $L^2(X)$, where I denotes the identity operator on $L^2(X)$;
- (ii) (the *size condition*) for any $x, y \in X$,

$$|Q_k(x, y)| \leq C E_k(x, y),$$

here and thereafter,

$$E_k(x, y) := \frac{1}{\sqrt{V_{\delta^k}(x) V_{\delta^k}(y)}} \exp \left\{ -\nu \left[\frac{\rho(x, y)}{\delta^k} \right]^a \right\} \exp \left\{ -\nu \left[\frac{\max\{\rho(x, \mathcal{Y}^k), \rho(y, \mathcal{Y}^k)\}}{\delta^k} \right]^a \right\};$$

- (iii) (the *regularity condition*) for any $x, x', y \in X$ with $\rho(x, x') \leq \delta^k$,

$$|Q_k(x, y) - Q_k(x', y)| + |Q_k(y, x) - Q_k(y, x')| \leq C \left[\frac{\rho(x, x')}{\delta^k} \right]^\eta E_k(x, y);$$

- (iv) (the *second difference regularity condition*) for any $x, x', y, y' \in X$ with $\rho(x, x') \leq \delta^k$ and $\rho(y, y') \leq \delta^k$,

$$|[Q_k(x, y) - Q_k(x', y)] - [Q_k(x, y') - Q_k(x', y')]| \leq C \left[\frac{\rho(x, x')}{\delta^k} \right]^\eta \left[\frac{\rho(y, y')}{\delta^k} \right]^\eta E_k(x, y);$$

- (v) (the *cancellation condition*) for any $x, y \in X$,

$$\int_X Q_k(x, y') d\mu(y') = 0 = \int_X Q_k(x', y) d\mu(x').$$

The following notion of 1-exp-ATIs is just [30, Definition 2.8].

Definition 3.4. Let $\eta \in (0, 1)$ be as in Definition 3.3. A sequence $\{P_k\}_{k \in \mathbb{Z}}$ of bounded linear integral operators on $L^2(X)$ is called an *approximation of the identity with exponential decay and integration 1* (for short, 1-exp-ATI) if $\{P_k\}_{k \in \mathbb{Z}}$ has the following properties:

- (i) for any $k \in \mathbb{Z}$, P_k satisfies (ii), (iii), and (iv) of Definition 3.3 but without the term $E_k(x, y)$;
- (ii) for any $k \in \mathbb{Z}$ and $x \in X$,

$$\int_X P_k(x, y) d\mu(y) = 1 = \int_X P_k(y, x) d\mu(y);$$

- (iii) for any $k \in \mathbb{Z}$, letting $Q_k := P_k - P_{k-1}$, then $\{Q_k\}_{k \in \mathbb{Z}}$ is an exp-ATI.

The following lemma is just [77, Lemma 3.3].

Lemma 3.5. Let $\{P_k\}_{k \in \mathbb{Z}}$ be a 1-exp-ATI and η as in Definition 3.3. Then the following statements hold true:

- (i) for any given $\beta, \gamma \in (0, \eta]$, there exists a positive constant C such that, for any $k \in \mathbb{Z}_+$ and $\varphi \in \mathcal{G}(\beta, \gamma)$,

$$\|P_k \varphi\|_{\mathcal{G}(\beta, \gamma)} \leq C \|\varphi\|_{\mathcal{G}(\beta, \gamma)};$$

- (ii) for any $\beta, \gamma \in (0, \eta]$, $\tilde{\beta} \in (0, \beta)$, and $f \in \mathcal{G}(\beta, \gamma)$,

$$\lim_{k \rightarrow \infty} P_k f = f$$

in $\mathcal{G}(\tilde{\beta}, \gamma)$;

- (iii) for any $\beta \in (0, \eta)$ and $\gamma \in (0, \eta]$, if $f \in \mathcal{G}_0^\eta(\beta, \gamma)$ [resp., $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$], then

$$\lim_{k \rightarrow \infty} P_k f = f$$

in $\mathcal{G}_0^\eta(\beta, \gamma)$ [resp., $(\mathcal{G}_0^\eta(\beta, \gamma))'$].

The following properties for 1-exp-ATIs as in Definition 3.4 are just [77, Lemmas 4.16 and 4.17], respectively.

Lemma 3.6. *Let $\beta, \gamma \in (0, \eta)$ with η as in Definition 3.3, $\{P_k\}_{k \in \mathbb{Z}}$ be a 1-exp-ATI as in Definition 3.4, and $f \in \mathcal{G}(x_0, r_0, \beta, \gamma)$ for some $x_0 \in X$ and $r_0 \in (0, \infty)$. Then there exists a positive constant C , independent of x_0, r_0 , and f , such that, for any $k \in \mathbb{Z}$ satisfying $r_0 \geq \delta^k$,*

$$\|P_k f\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)} \leq C \|f\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)}.$$

Lemma 3.7. *Let $\beta, \gamma \in (0, \eta)$ with η as in Definition 3.3, $\{P_k\}_{k \in \mathbb{Z}}$ be a 1-exp-ATI as in Definition 3.4, and $f \in \mathcal{G}(x_0, r_0, \beta, \gamma)$ for some $x_0 \in X$ and $r_0 \in (0, \infty)$. Then there exists a positive constant C , independent of x_0, r_0 , and f , such that, for any $k \in \mathbb{Z}$ satisfying $r_0 < \delta^k$,*

$$\|P_k f\|_{\mathcal{G}(x_0, \delta^k, \beta, \gamma)} \leq C \|f\|_{\mathcal{G}(x_0, r_0, \beta, \gamma)}.$$

Remark 3.8. Compared with the corresponding ones on RD-spaces (see [29]), these exp-ATIs on X have some essential differences presented via some terms such as

$$\exp \left\{ -\nu \left[\frac{\max\{\rho(x, \mathcal{Y}^k), \rho(y, \mathcal{Y}^k)\}}{\delta^k} \right]^a \right\}.$$

Observe that, here $x, y \in X$, \mathcal{Y}^k is the set of dyadic reference points as in (3.3), and $\rho(x, \mathcal{Y}^k)$ [resp. $\rho(y, \mathcal{Y}^k)$] the quasi-distance between x (resp. y) and \mathcal{Y}^k as in (3.4). Thus, such terms are closely connecting with the geometrical property of X and, therefore, help us to get rid of the dependence on the reverse doubling assumption of the equipped measure μ of X .

Next, we introduce several Hardy spaces, associated with $Y(X)$, via different maximal functions.

Definition 3.9. Let $\beta, \gamma \in (0, \eta)$ and $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$ with η as in Definition 3.3. Let $\{P_k\}_{k \in \mathbb{Z}}$ be a 1-exp-ATI as in Definition 3.4. The *radial maximal function* $\mathcal{M}^+(f)$ of f is defined by setting, for any $x \in X$,

$$\mathcal{M}^+(f)(x) := \sup_{k \in \mathbb{Z}} |P_k f(x)|$$

and, for any given $a \in (0, \infty)$, the *non-tangential maximal function* $\mathcal{M}_a(f)$, with aperture a , of f is defined by setting, for any $x \in X$,

$$\mathcal{M}_a(f)(x) := \sup_{k \in \mathbb{Z}} \sup_{y \in B(x, a\delta^k)} |P_k f(y)|,$$

where δ is as in Lemma 3.2. Moreover, the *grand maximal function* f^* of f is defined by setting, for any $x \in X$,

$$f^*(x) := \sup \left\{ |\langle f, h \rangle| : h \in \mathcal{G}_0^\eta(\beta, \gamma), \|h\|_{\mathcal{G}(x, r, \beta, \gamma)} \leq 1 \text{ for some } r \in (0, \infty) \right\}.$$

Let $Y(X)$ be a ball quasi-Banach function space on X . Then the *Hardy spaces* $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$ [with $a \in (0, \infty)$], and $H_Y^*(\mathcal{X})$, associated with $Y(X)$, are defined, respectively, by setting

$$H_Y^+(\mathcal{X}) := \left\{ f \in (\mathcal{G}_0^\eta(\beta, \gamma))' : \|f\|_{H_Y^+(\mathcal{X})} := \|\mathcal{M}^+(f)\|_{Y(X)} < \infty \right\}, \quad (3.5)$$

$$H_Y^a(X) := \left\{ f \in (\mathcal{G}_0^\eta(\beta, \gamma))' : \|f\|_{H_Y^a(X)} := \|\mathcal{M}_a(f)\|_{Y(X)} < \infty \right\}, \quad (3.6)$$

and

$$H_Y^*(X) := \left\{ f \in (\mathcal{G}_0^\eta(\beta, \gamma))' : \|f\|_{H_Y^*(X)} := \|f^*\|_{Y(X)} < \infty \right\}. \quad (3.7)$$

The following lemma shows that $H_Y^*(X)$ can be embedded into $(\mathcal{G}_0^\eta(\beta, \gamma))'$.

Lemma 3.10. *Assume that $Y(X)$ is a ball quasi-Banach function space on X . Let β , γ , and η be as in Definition 3.9. Then $H_Y^*(X)$ is continuously embedded into $(\mathcal{G}_0^\eta(\beta, \gamma))'$, namely, there exists a positive constant C such that, for any $f \in H_Y^*(X)$ and $h \in \mathcal{G}_0^\eta(\beta, \gamma)$,*

$$|\langle f, h \rangle| \leq C \|f\|_{H_Y^*(X)} \|h\|_{\mathcal{G}_0^\eta(\beta, \gamma)}. \quad (3.8)$$

Proof. Let all the symbols be as in the present lemma. Without loss of generality, we may assume that $h \in \mathcal{G}_0^\eta(\beta, \gamma)$ with $\|h\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \leq 1$. Let $x_0 \in X$ be as in Definition 3.1. By Definition 3.1, we know that, for any $x \in B(x_0, 1)$,

$$\|h\|_{\mathcal{G}(x, 1, \beta, \gamma)} \sim \|h\|_{\mathcal{G}(\beta, \gamma)} \sim \|h\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \lesssim 1,$$

where the implicit positive constants are independent of x and h . From this and Definition 3.9, we deduce that, for any $f \in H_Y^*(X)$ and $x \in B(x_0, 1)$, $|\langle f, h \rangle| \lesssim f^*(x)$, which, combined with (ii) and (iv) of Definition 2.4, further implies that

$$|\langle f, h \rangle| \lesssim \inf_{x \in B(x_0, 1)} f^*(x) \lesssim \|f^*\|_{Y(X)} \|\mathbf{1}_{B(x_0, 1)}\|_{Y(X)}^{-1} \sim \|f\|_{H_Y^*(X)}. \quad (3.9)$$

This shows (3.8) and hence finishes the proof of Lemma 3.10. \square

Remark 3.11. (i) Let $a \in (0, \infty)$ and η be as in Definition 3.3. By [23, Remark 2.9(ii)], we know that there exists a positive constant C , depending on a , such that, for any $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$ with $\beta, \gamma \in (0, \eta)$, and any $x \in X$,

$$\mathcal{M}^+(f)(x) \leq \mathcal{M}_a(f)(x) \leq C f^*(x).$$

(ii) Let $Y(X)$ be a ball quasi-Banach function space on X . By Lemmas 2.9 and 3.10, together with an argument similar to that used in the proof of [30, Proposition 3.1] (see also [77, Proposition 3.9]), we conclude that $H_Y^*(X)$ is complete.

Next, we state the main theorem of this subsection as follows.

Theorem 3.12. *Let $\beta, \gamma \in (0, \eta)$ with η as in Definition 3.3, and $Y(X)$ be a ball quasi-Banach function space on X . Assume that there exist an $r_1 \in (1, \infty)$ and an $r_2 \in [1, \infty)$ such that $Y^{1/r_1}(X)$ is a ball Banach function space and that \mathcal{M} is bounded, respectively, on both $(Y^{1/r_1})'(X)$ and $Y^{1/r_2}(X)$. Then the following statements hold true:*

- (i) $Y(X) \hookrightarrow (\mathcal{G}_0^\eta(\beta, \gamma))'$;
- (ii) if $f \in Y(X)$, then $f \in H_Y^*(X)$ and there exists a positive constant C , independent of f , such that $\|f\|_{H_Y^*(X)} \leq C \|f\|_{Y(X)}$;

(iii) if $f \in H_Y^+(X)$, then there exists an $\tilde{f} \in Y(X)$ such that $\tilde{f} = f$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ and $\|\tilde{f}\|_{Y(X)} \leq \|f\|_{H_Y^+(X)}$.

Consequently, for any fixed $a \in (0, \infty)$, $H_Y^+(X) = H_Y^a(X) = H_Y^*(X) = Y(X)$ in the sense of both representing the same distributions and having the equivalent norms.

Proof. Let all the symbols be as in the present theorem. First, we prove (i). Let $f \in Y(X)$, $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$, and r_2 be as in the present theorem. By (2.10), we know that $\mathcal{M}(f) \leq \mathcal{M}^{(r_2)}(f)$, which further implies that $[\mathcal{M}(f)]^{r_2} \leq \mathcal{M}(|f|^{r_2})$. From this, Definition 2.10, and the boundedness of \mathcal{M} on $Y^{1/r_2}(X)$, we deduce that

$$\|\mathcal{M}(f)\|_{Y(X)} = \|[\mathcal{M}(f)]^{r_2}\|_{Y^{\frac{1}{r_2}}(X)}^{\frac{1}{r_2}} \leq \|\mathcal{M}(|f|^{r_2})\|_{Y^{\frac{1}{r_2}}(X)}^{\frac{1}{r_2}} \lesssim \| |f|^{r_2} \|_{Y^{\frac{1}{r_2}}(X)}^{\frac{1}{r_2}} \sim \|f\|_{Y(X)}. \quad (3.10)$$

By this, Definition 3.1(i), (2.3), and Definition 2.4(iv), we conclude that

$$\begin{aligned} & \int_X |f(x)\varphi(x)| d\mu(x) \\ & \leq \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \int_X |f(x)| P_\gamma(x_0, x; 1) d\mu(x) \\ & = \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \left\{ \sum_{k \in \mathbb{N}} \int_{B(x_0, 2^{k+1}) \setminus B(x_0, 2^k)} + \int_{B(x_0, 2)} \right\} |f(x)| P_\gamma(x_0, x; 1) d\mu(x) \\ & \lesssim \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \sum_{k \in \mathbb{Z}_+} 2^{-k\gamma} \frac{1}{\mu(B(x_0, 2^k))} \int_{B(x_0, 2^{k+1})} |f(x)| d\mu(x) \\ & \lesssim \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \sum_{k \in \mathbb{Z}_+} 2^{-k\gamma} \inf_{y \in B(x_0, 1)} \mathcal{M}(f)(y) \\ & \lesssim \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \|\mathcal{M}(f)\|_{Y(X)} \|\mathbf{1}_{B(x_0, 1)}\|_{Y(X)}^{-1} \lesssim \|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \|f\|_{Y(X)}, \end{aligned}$$

where $P_\gamma(x_0, x; 1)$ is as in (1.1) with r replaced by 1. This further implies that $Y(X) \hookrightarrow (\mathcal{G}_0^\eta(\beta, \gamma))'$ and hence (i) holds true.

Next, we prove (ii). Let $f \in Y(X)$. By (i) and [23, Proposition 3.9], we know that $f^* \lesssim \mathcal{M}(f)$. From this and (3.10), we deduce that

$$\|f\|_{H_Y^+(X)} \lesssim \|\mathcal{M}(f)\|_{Y(X)} \lesssim \|f\|_{Y(X)},$$

which completes the proof of (ii).

Finally, we prove (iii). Let $f \in H_Y^+(X)$, $\{P_k\}_{k \in \mathbb{Z}}$ be a 1-exp-ATI as in Definition 3.4, and r_1 as in the present theorem. By Definition 3.9 and Theorem 2.17, we know that

$$\left\| \sup_{k \in \mathbb{Z}} |P_k f| \right\|_{L_w^{r_1}(X)} \lesssim \left\| \sup_{k \in \mathbb{Z}} |P_k f| \right\|_{Y(X)} \sim \|f\|_{H_Y^+(X)} < \infty,$$

where w is as in Theorem 2.17. From this and the Banach–Alaoglu theorem (see, for instance, [57, Theorem 3.17]), we deduce that there exist an $\tilde{f} \in L_w^{r_1}(X)$ and a sequence $\{k_j\}_{j \in \mathbb{N}} \subset \mathbb{Z}$ such that, for any $g \in (L_w^{r_1}(X))^*$ [the dual space of $L_w^{r_1}(X)$],

$$\int_X \tilde{f}(x)g(x) d\mu(x) = \lim_{j \rightarrow \infty} \int_X P_{k_j} f(x)g(x) d\mu(x). \quad (3.11)$$

Using this and $\mathcal{G}_0^\eta(\beta, \gamma) \subset (L_w^{r_1}(\mathcal{X}))^*$, we conclude that, for any $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$,

$$\begin{aligned} \left| \int_{\mathcal{X}} \widetilde{f}(x) \varphi(x) d\mu(x) \right| &= \lim_{j \rightarrow \infty} \left| \int_{\mathcal{X}} P_{k_j} f(x) \varphi(x) d\mu(x) \right| \\ &\leq \int_{\mathcal{X}} \sup_{k \in \mathbb{Z}} |P_k f(x)| |\varphi(x)| d\mu(x), \end{aligned}$$

which, combined with the arbitrariness of $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$ and the Lebesgue differentiation theorem (see, for instance, [34, Theorem 1.8]), further implies that, for almost every $x \in \mathcal{X}$,

$$|\widetilde{f}(x)| \leq \sup_{k \in \mathbb{Z}} |P_k f(x)|.$$

Thus, $\|\widetilde{f}\|_{Y(\mathcal{X})} \leq \|f\|_{H_Y^+(\mathcal{X})}$. Moreover, letting $w_1 := w^{1-r_1'}$, then, by the Hölder inequality and $\mathcal{G}_0^\eta(\beta, \gamma) \subset L_{w_1}^{r_1'}(\mathcal{X})$, we know that, for any $j \in \mathbb{N}$ and $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$,

$$\begin{aligned} &\int_{\mathcal{X}} |P_{k_j} f(x) \varphi(x)| d\mu(x) \\ &\leq \left[\int_{\mathcal{X}} |P_{k_j} f(x)|^{r_1} w(x) d\mu(x) \right]^{1/r_1} \left[\int_{\mathcal{X}} |\varphi(x)|^{r_1'} [w(x)]^{-r_1'/r_1} d\mu(x) \right]^{1/r_1'} \\ &= \|P_{k_j} f\|_{L_w^{r_1}(\mathcal{X})} \|\varphi\|_{L_{w_1}^{r_1'}(\mathcal{X})} < \infty. \end{aligned}$$

From this, Lemma 3.5(iii), $L_w^{r_1}(\mathcal{X}) \subset (\mathcal{G}_0^\eta(\beta, \gamma))'$, [29, (3.98)], and (3.11), we deduce that, for any $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$,

$$\langle f, \varphi \rangle = \lim_{j \rightarrow \infty} \langle P_{k_j} f, \varphi \rangle = \lim_{j \rightarrow \infty} \int_{\mathcal{X}} P_{k_j} f(x) \varphi(x) d\mu(x) = \int_{\mathcal{X}} \widetilde{f}(x) \varphi(x) d\mu(x).$$

This finishes the proof of (iii).

By (ii), (iii), and Remark 3.11(i), we obtain $H_Y^+(\mathcal{X}) = H_Y^a(\mathcal{X}) = H_Y^*(\mathcal{X}) = Y(\mathcal{X})$, which completes the proof of Theorem 3.12. \square

Remark 3.13. (i) Let $a \in (0, \infty)$, $p \in (1, \infty]$, and $Y(\mathcal{X}) := L^p(\mathcal{X})$. Then $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$, and $H_Y^*(\mathcal{X})$ are, respectively, the classical *Hardy spaces* $H^{+,p}(\mathcal{X})$, $H_a^p(\mathcal{X})$, and $H^{*,p}(\mathcal{X})$, which were introduced by He et al. in [30]. Choose $r_1, r_2 \in (1, p)$. By this, Remark 2.7(i), and [14, (3.6)], we conclude that $L^p(\mathcal{X})$ satisfies all the assumptions of Theorem 3.12. In this case, Theorem 3.12 covers [30, Theorem 3.4].

(ii) We point out that, in the proof of Theorem 3.12(iii), we can not find a desired \widetilde{f} directly since $Y(\mathcal{X})$ does not have an explicit norm expression. To overcome the difficulty caused by this, via Theorem 2.17, we embed $Y(\mathcal{X})$ to the weighted Lebesgue space $L_w^{r_1}(\mathcal{X})$ with w as in Theorem 2.17, and then apply the completeness of $L_w^{r_1}(\mathcal{X})$ together with the Banach–Alaoglu theorem to complete the proof of 3.12(iii).

3.2 Maximal function characterizations

In this subsection, we characterize $H_Y^*(X)$ by the radial and the non-tangential maximal functions, respectively. To this end, we first prove the following conclusion.

Theorem 3.14. *Let $Y(X)$ be a ball quasi-Banach function space on X , $a \in (0, \infty)$, and $p_- \in (\omega/(\omega + \eta), \infty)$ with ω and η , respectively, as in (2.3) and Definition 3.3. Assume that, for any given $\theta \in (0, \underline{p})$ with \underline{p} as in (2.8), the Hardy–Littlewood maximal operator \mathcal{M} in (2.6) is bounded on $Y^{1/\theta}(X)$. Let $\beta, \gamma \in (\omega(1/\underline{p} - 1), \eta)$. Then*

$$\left[H_Y^+(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))' \right] = \left[H_Y^a(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))' \right] = \left[H_Y^*(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))' \right]$$

with equivalent quasi-norms.

Proof. Let all the symbols be as in the present theorem. Assume that $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$. By Remark 3.11(i) and Definition 2.4(ii), we know that

$$\|\mathcal{M}^+(f)\|_{Y(X)} \leq \|\mathcal{M}_a(f)\|_{Y(X)} \lesssim \|f^*\|_{Y(X)}$$

and hence $H_Y^*(X) \subset H_Y^a(X) \subset H_Y^+(X)$. Thus, to complete the proof of this theorem, it suffices to show that $H_Y^+(X) \subset H_Y^*(X)$ and

$$\|f^*\|_{Y(X)} \lesssim \|\mathcal{M}^+(f)\|_{Y(X)}. \quad (3.12)$$

Choose $\tilde{\beta} \in (0, \min\{\beta, \gamma\})$ such that $\frac{\omega}{\omega + \tilde{\beta}} < \underline{p}$. Let $\theta \in (\frac{\omega}{\omega + \tilde{\beta}}, \underline{p})$. Then, similarly to the proof of [30, (3.5)], we conclude that, for any $x \in X$,

$$f^*(x) \lesssim \mathcal{M}^+(f)(x) + \left\{ \mathcal{M}([\mathcal{M}^+(f)]^\theta)(x) \right\}^{\frac{1}{\theta}},$$

which, combined with Definitions 2.4(ii) and 2.10, and the boundedness of \mathcal{M} on $Y^{1/\theta}(X)$, further implies that

$$\begin{aligned} \|f^*\|_{Y(X)} &\lesssim \|\mathcal{M}^+(f)\|_{Y(X)} + \left\| \left\{ \mathcal{M}([\mathcal{M}^+(f)]^\theta) \right\}^{\frac{1}{\theta}} \right\|_{Y(X)} \\ &\sim \|\mathcal{M}^+(f)\|_{Y(X)} + \left\| \mathcal{M}([\mathcal{M}^+(f)]^\theta) \right\|_{Y^{1/\theta}(X)}^{\frac{1}{\theta}} \\ &\lesssim \|\mathcal{M}^+(f)\|_{Y(X)}. \end{aligned}$$

This finishes the proof of (3.12) and hence of Theorem 3.14. \square

Next, we give an important property of $H_Y^*(X)$ as follows.

Theorem 3.15. *Let $Y(X)$ be a ball quasi-Banach function space on X , and $p_- \in (\omega/(\omega + \eta), \infty)$ with ω and η , respectively, as in (2.3) and Definition 3.3. Assume that, for any given $\theta \in (0, \underline{p})$ with \underline{p} as in (2.8), the Hardy–Littlewood maximal operator \mathcal{M} in (2.6) is bounded on $Y^{1/\theta}(X)$. Let $\beta_1, \beta_2, \gamma_1, \gamma_2 \in (\omega(1/\underline{p} - 1), \eta)$ and $H_Y^*(X)$ be as in Definition 3.9 with β and γ replaced, respectively, by β_1 and γ_1 . If $f \in H_Y^*(X)$, then $f \in (\mathcal{G}_0^\eta(\beta_2, \gamma_2))'$.*

Proof. Let β_1 and γ_1 be as in the present theorem, $f \in H_Y^*(X)$, and x_0 be as in Definition 3.1. Then, by an argument similar to that used in the proof of [30, (3.9)], we know that there exist an $a \in (0, \infty)$ and a $\theta \in (\omega/(\omega + \eta), \underline{p})$ such that, for any $\varphi \in \mathcal{G}(\eta, \eta)$ with $\|\varphi\|_{\mathcal{G}(\beta_2, \gamma_2)} \leq 1$, and any $x \in B(x_0, 1)$,

$$|\langle f, \varphi \rangle| \lesssim \left\{ \mathcal{M}([\mathcal{M}_a(f)]^\theta)(x) \right\}^{\frac{1}{\theta}},$$

which, combined with (ii) and (iv) of Definition 2.4, Definition 2.10, the boundedness of \mathcal{M} on $Y^{1/\theta}(X)$, and Theorem 3.14, further implies that

$$\begin{aligned} |\langle f, \varphi \rangle| &\lesssim \inf_{x \in B(x_0, 1)} \left\{ \mathcal{M}([\mathcal{M}_a(f)]^\theta)(x) \right\}^{\frac{1}{\theta}} \lesssim \left\| \left\{ \mathcal{M}([\mathcal{M}_a(f)]^\theta) \right\} \right\|_{Y(X)}^{\frac{1}{\theta}} \left\| \mathbf{1}_{B(x_0, 1)} \right\|_{Y(X)}^{-1} \\ &\lesssim \left\| \mathcal{M}([\mathcal{M}_a(f)]^\theta) \right\|_{Y^{1/\theta}(X)}^{\frac{1}{\theta}} \lesssim \|\mathcal{M}_a(f)\|_{Y(X)} \sim \|f\|_{H_Y^*(X)}. \end{aligned}$$

From this and the homogeneity of $\|\cdot\|_{\mathcal{G}(\beta_2, \gamma_2)}$, we deduce that, for any $\varphi \in \mathcal{G}(\eta, \eta)$,

$$|\langle f, \varphi \rangle| \lesssim \|f\|_{H_Y^*(X)} \|\varphi\|_{\mathcal{G}(\beta_2, \gamma_2)}. \quad (3.13)$$

Now, let $g \in \mathcal{G}_0^\eta(\beta_2, \gamma_2)$. By the definition of $\mathcal{G}_0^\eta(\beta_2, \gamma_2)$, we know that there exist $\{g_j\}_{j \in \mathbb{N}} \subset \mathcal{G}(\eta, \eta)$ such that

$$\|g - g_j\|_{\mathcal{G}(\beta_2, \gamma_2)} \rightarrow 0 \quad (3.14)$$

as $j \rightarrow \infty$. Moreover, from (3.13), we deduce that, for any $j, k \in \mathbb{N}$,

$$\left| \langle f, g_j \rangle - \langle f, g_k \rangle \right| = \left| \langle f, g_j - g_k \rangle \right| \lesssim \|f\|_{H_Y^*(X)} \|g_j - g_k\|_{\mathcal{G}(\beta_2, \gamma_2)}.$$

This, combined with (3.14), implies that $\lim_{j \rightarrow \infty} \langle f, g_j \rangle$ exists and the limit is independent of the choice of $\{g_j\}_{j \in \mathbb{N}}$. Define $\langle f, g \rangle := \lim_{j \rightarrow \infty} \langle f, g_j \rangle$. By this, (3.13), and (3.14), we conclude that

$$|\langle f, g \rangle| = \lim_{j \rightarrow \infty} \left| \langle f, g_j \rangle \right| \lesssim \|f\|_{H_Y^*(X)} \lim_{j \rightarrow \infty} \|g_j\|_{\mathcal{G}(\beta_2, \gamma_2)} \sim \|f\|_{H_Y^*(X)} \|g\|_{\mathcal{G}_0^\eta(\beta_2, \gamma_2)},$$

which further implies that $f \in (\mathcal{G}_0^\eta(\beta_2, \gamma_2))'$. This finishes the proof of Theorem 3.15. \square

Combining Theorems 3.14 and 3.15, we obtain the following main theorem of this subsection, which shows that the Hardy spaces in Definition 3.9 are all independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$ and coincide with equivalent quasi-norms.

Theorem 3.16. *Let $Y(X)$ be a ball quasi-Banach function space on X , $a \in (0, \infty)$, and $p_- \in (\omega/(\omega + \eta), \infty)$ with ω and η , respectively, as in (2.3) and Definition 3.3. Assume that, for any given $\theta \in (0, \underline{p})$ with \underline{p} as in (2.8), the Hardy–Littlewood maximal operator \mathcal{M} in (2.6) is bounded on $Y^{1/\theta}(X)$. Then $H_Y^+(X)$, $H_Y^a(X)$, and $H_Y^*(X)$ are all independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$ whenever $\beta, \gamma \in (\omega(1/\underline{p} - 1), \eta)$, and coincide with equivalent quasi-norms.*

Proof. Let $\beta_1, \beta_2, \gamma_1, \gamma_2 \in (\omega(1/\underline{p} - 1), \eta)$. For any $i \in \{1, 2\}$, define $H_Y^{+,i}(X)$, $H_Y^{a,i}(X)$, and $H_Y^{*,i}(X)$, respectively, as in (3.5), (3.6), and (3.7) with $(\mathcal{G}_0^\eta(\beta, \gamma))'$ replaced by $(\mathcal{G}_0^\eta(\beta_i, \gamma_i))'$. To prove this theorem, it suffices to show that

$$H_Y^{+,1}(X) = H_Y^{+,2}(X). \quad (3.15)$$

Indeed, using (3.15) and Theorem 3.14 twice, we obtain

$$H_Y^{a,1}(\mathcal{X}) = H_Y^{+,1}(\mathcal{X}) = H_Y^{+,2}(\mathcal{X}) = H_Y^{a,2}(\mathcal{X})$$

and

$$H_Y^{*,1}(\mathcal{X}) = H_Y^{*,1}(\mathcal{X}) = H_Y^{*,2}(\mathcal{X}) = H_Y^{*,2}(\mathcal{X}).$$

Next, we prove (3.15). By symmetry, we only need to show $H_Y^{+,1}(\mathcal{X}) \subset H_Y^{+,2}(\mathcal{X})$. To this end, let $f \in H_Y^{+,1}(\mathcal{X}) \subset (\mathcal{G}_0^\eta(\beta_1, \gamma_1))'$. Then, by Theorems 3.14 and 3.15, we know that $f \in H_Y^{*,1}(\mathcal{X}) \subset (\mathcal{G}_0^\eta(\beta_2, \gamma_2))'$, which, combined with $\mathcal{M}^+(f) \in Y(\mathcal{X})$, further implies that $f \in H_Y^{+,2}(\mathcal{X})$. Thus, $H_Y^{+,1}(\mathcal{X}) \subset H_Y^{+,2}(\mathcal{X})$. This finishes the proof of (3.15) and hence of Theorem 3.16. \square

Remark 3.17. Next, we apply Theorem 3.16 to several concrete examples of ball quasi-Banach function spaces on \mathcal{X} , namely, classical Lebesgue spaces, Lorentz spaces, Musielak–Orlicz spaces, and variable Lebesgue spaces.

- (i) Assume that $a \in (0, \infty)$, $p \in (\omega/(\omega + \eta), 1]$, and $\beta, \gamma \in (\omega(1/p - 1), \eta)$ with ω as in (2.3) and η as in Definition 3.3. Let $Y(\mathcal{X}) := L^p(\mathcal{X})$. Then $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$, and $H_Y^*(\mathcal{X})$ are, respectively, the classical *Hardy spaces* $H^{+,p}(\mathcal{X})$, $H_a^p(\mathcal{X})$, and $H^{*,p}(\mathcal{X})$, which were introduced by He et al. in [30]. Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p]$. By this, Remark 2.7(i), the fact that $Y^{1/\theta}(\mathcal{X}) = L^{p/\theta}(\mathcal{X})$ for any $\theta \in (0, \infty)$, and [14, (3.6)], we conclude that $L^p(\mathcal{X})$ satisfies all the assumptions of Theorem 3.16. In this case, Theorem 3.16 improves the corresponding results in [30, Theorem 3.5] by removing the assumption that Hardy spaces are restricted to the same space of distributions.
- (ii) Assume that $a \in (0, \infty)$, $r \in (0, \infty)$, $p \in (\omega/(\omega + \eta), \infty)$, and $\beta, \gamma \in (\omega[1/p - 1]_+, \eta)$ with ω as in (2.3), η as in Definition 3.3, and $[1/p - 1]_+ := \max\{1/p - 1, 0\}$. Let $Y(\mathcal{X}) := L^{p,r}(\mathcal{X})$. Then $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$, and $H_Y^*(\mathcal{X})$ are, respectively, the *Hardy–Lorentz spaces* $H_+^{p,r}(\mathcal{X})$, $H_a^{p,r}(\mathcal{X})$, and $H_\star^{p,r}(\mathcal{X})$, which were introduced by Zhou et al. in [77]. Choose

$$p_- \in \begin{cases} (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p] & \text{if } p \in (\omega/(\omega + \eta), 1), \\ [1, p] & \text{if } p \in [1, \infty). \end{cases}$$

By this, Remark 2.7(ii), the fact that $Y^{1/\theta}(\mathcal{X}) = L^{p/\theta, r/\theta}(\mathcal{X})$ for any $\theta \in (0, \infty)$, and [77, Lemma 3.5], we conclude that $L^{p,r}(\mathcal{X})$ satisfies all the assumptions of Theorem 3.16. In this case, Theorem 3.16 improves the corresponding results in [77, Theorem 3.6] by removing the assumption that Hardy spaces are restricted to the same space of distributions.

- (iii) Assume that $a \in (0, \infty)$, φ is a growth function as in Remark 2.7(iii) satisfying $p/q(\varphi) \in (\omega/(\omega + \eta), 1]$, and $\beta, \gamma \in (\omega[q(\varphi)/p - 1], \eta)$ with ω as in (2.3) and η as in Definition 3.3. Let $Y(\mathcal{X}) := L^\varphi(\mathcal{X})$. Then $H_Y^+(\mathcal{X})$, $H_Y^a(\mathcal{X})$, and $H_Y^*(\mathcal{X})$ are, respectively, the *Musielak–Orlicz Hardy spaces* $H^{+, \varphi}(\mathcal{X})$, $H_a^\varphi(\mathcal{X})$, and $H^{*, \varphi}(\mathcal{X})$, which were introduced by Fu et al. in [19]. Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p/q(\varphi)]$. By this, Remark 2.7(iii), the fact that $Y^{1/\theta}(\mathcal{X}) = L^{\tilde{\varphi}}(\mathcal{X})$ with $\tilde{\varphi}(\cdot, t) = \varphi(\cdot, t^{1/\theta})$ for any $t \in (0, \infty)$ and $\theta \in (0, \infty)$, and [19, Theorem 4.11], we conclude that $L^\varphi(\mathcal{X})$ satisfies all the assumptions of Theorem 3.16. In this setting, Theorem 3.16 improves the corresponding results in [19, Theorem 4.12] by removing the assumption that Hardy spaces are restricted to the same space of distributions.

- (iv) Recall that a μ -measurable function $p(\cdot) : \mathcal{X} \rightarrow (0, \infty)$ is said to be *globally log-Hölder continuous*, denoted by $p(\cdot) \in C^{\log}(\mathcal{X})$, if there exist positive constants $C_{\log}(p)$ and C_{∞} , $x_p \in \mathcal{X}$, and $p_{\infty} \in \mathbb{R}$ such that, for any $x, y \in \mathcal{X}$,

$$|p(x) - p(y)| \leq \frac{C_{\log}(p)}{\log(e + 1/\rho(x, y))}$$

and

$$|p(x) - p_{\infty}| \leq \frac{C_{\infty}}{\log(e + \rho(x, x_p))}.$$

Assume that $a \in (0, \infty)$, $p(\cdot) \in C^{\log}(\mathcal{X})$ satisfies $\widetilde{p}_{-} \in (\omega/(\omega + \eta), \infty)$, and $\beta, \gamma \in (\omega[1/\widetilde{p}_{-} - 1]_{+}, \eta)$ with ω as in (2.3), η as in Definition 3.3, and $[1/\widetilde{p}_{-} - 1]_{+} := \max\{1/\widetilde{p}_{-} - 1, 0\}$. Let $Y(\mathcal{X}) := L^{p(\cdot)}(\mathcal{X})$. Then $H_Y^*(\mathcal{X})$ is the *variable Hardy space* $H^{*,p(\cdot)}(\mathcal{X})$ which was introduced by Zhuo et al. in [78]. Choose

$$p_{-} \in \begin{cases} (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, \widetilde{p}_{-}] & \text{if } \widetilde{p}_{-} \in (\omega/(\omega + \eta), 1), \\ [1, \widetilde{p}_{-}] & \text{if } \widetilde{p}_{-} \in [1, \infty). \end{cases}$$

By this, Remark 2.7(iv), the fact that $Y^{1/\theta}(\mathcal{X}) = L^{p(\cdot)/\theta}(\mathcal{X})$ for any $\theta \in (0, \infty)$, and [78, Lemma 2.5], we conclude that $L^{p(\cdot)}(\mathcal{X})$ satisfies all the assumptions of Theorem 3.16. In this case, a corresponding conclusion of Theorem 3.16 was obtained in [78, Theorem 3.11]. However, we point out that, in [78, Theorem 3.11], (\mathcal{X}, ρ, μ) is assumed to be an RD-space. Thus, Theorem 3.16 improves the corresponding results in [78] by removing the assumptions that μ satisfies the reverse doubling condition and that Hardy spaces are restricted to the same space of distributions.

4 Atomic and finite atomic characterizations of $H_Y^*(\mathcal{X})$

In this section, we characterize $H_Y^*(\mathcal{X})$ by atoms and finite atoms. Indeed, using the boundedness of the powered Hardy–Littlewood maximal operator on the associate space of $Y(\mathcal{X})$, a key lemma (see Lemma 4.3 below), the continuous embedding of $Y(\mathcal{X})$ to $L_w^p(\mathcal{X})$, and the Fefferman–Stein vector-valued maximal inequality on $Y(\mathcal{X})$, we first establish the atomic characterization of $H_Y^*(\mathcal{X})$. By this, we further obtain the finite atomic characterization of $H_Y^*(\mathcal{X})$.

4.1 Atomic characterizations

In this subsection, we establish the atomic characterization of $H_Y^*(\mathcal{X})$. To this end, we first introduce the notion of atoms associated with $Y(\mathcal{X})$.

Definition 4.1. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} , and $q \in [1, \infty]$. A μ -measurable function a on \mathcal{X} is called a $(Y(\mathcal{X}), q)$ -atom if there exists a ball $B \subset \mathcal{X}$ such that

- (i) $\text{supp } a := \{x \in \mathcal{X} : a(x) \neq 0\} \subset B$;
- (ii) $\|a\|_{L^q(\mathcal{X})} \leq [\mu(B)]^{1/q} \|\mathbf{1}_B\|_{Y(\mathcal{X})}^{-1}$;
- (iii) $\int_{\mathcal{X}} a(x) d\mu(x) = 0$.

We first give a reconstruction result of $H_Y^*(X)$ as follows (see [59, Theorem 3.6] for the corresponding Euclidean case).

Proposition 4.2. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), p)$, and $p_0 \in (\theta_0, \infty)$, where p is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$ and $d \in (0, \theta_0]$. Suppose that $\{a_j\}_{j \in \mathbb{N}}$ is a sequence of $(Y(\bar{X}), q)$ -atoms supported, respectively, in balls $\{B_j\}_{j \in \mathbb{N}}$ of X , $\{\lambda_j\}_{j \in \mathbb{N}} \subset [0, \infty)$, and*

$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} < \infty.$$

Then $f := \sum_{j \in \mathbb{N}} \lambda_j a_j$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ for any given $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. Moreover, $f \in H_Y^(X)$ and there exists a positive constant C , independent of $\{a_j\}_{j \in \mathbb{N}}$, $\{B_j\}_{j \in \mathbb{N}}$, and $\{\lambda_j\}_{j \in \mathbb{N}}$, such that*

$$\|f\|_{H_Y^*(X)} \leq C \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)}. \quad (4.1)$$

The following technical lemma, which is a generalization of [76, Lemma 4.8] on \mathbb{R}^n to X , plays a key role in the proof of Proposition 4.2. It can be proved by an argument similar to that used in the proofs of [76, Lemma 4.8] and Lemmas 2.18 and 2.20; we omit the details here.

Lemma 4.3. *Let $r \in (0, \infty)$, $s \in (r, \infty]$, and $Y(X)$ be a ball quasi-Banach function space on X . Assume that $Y^{1/r}(X)$ is a ball Banach function space and that there exists a positive constant C_0 such that, for any $f \in (Y^{1/r})'(X)$,*

$$\|\mathcal{M}^{((s/r)')} (f)\|_{(Y^{1/r})'(X)} \leq C_0 \|f\|_{(Y^{1/r})'(X)}.$$

Then there exists a positive constant C such that, for any sequence $\{B_j\}_{j \in \mathbb{N}}$ of balls, $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$, and μ -measurable functions $\{a_j\}_{j \in \mathbb{N}}$ satisfying that, for any $j \in \mathbb{N}$, $\text{supp } a_j \subset B_j$ and $\|a_j\|_{L^s(X)} \leq [\mu(B_j)]^{1/s}$, it holds true that

$$\left\| \left(\sum_{j \in \mathbb{N}} |\lambda_j a_j|^r \right)^{\frac{1}{r}} \right\|_{Y(X)} \leq C \left\| \left(\sum_{j \in \mathbb{N}} |\lambda_j \mathbf{1}_{B_j}|^r \right)^{\frac{1}{r}} \right\|_{Y(X)}.$$

We also need the following basic inequality.

Lemma 4.4. *Let $\theta \in (0, 1]$. Then, for any $\{a_j\}_{j \in \mathbb{N}} \subset [0, \infty)$,*

$$\left(\sum_{j \in \mathbb{N}} a_j \right)^\theta \leq \sum_{j \in \mathbb{N}} a_j^\theta.$$

Now, we show Proposition 4.2.

Proof of Proposition 4.2. Let all the symbols be as in the present proposition, a a $(Y(\mathcal{X}), q)$ -atom supported in a ball $B := B(x_B, r_B)$ with $x_B \in \mathcal{X}$ and $r_B \in (0, \infty)$, and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. Let $\widetilde{B} := 2A_0B$. From the proof of [30, (4.1)], we deduce that, for any $x \in B$,

$$a^*(x) \lesssim \mathcal{M}(a)(x). \quad (4.2)$$

Moreover, by the cancellation of a , (3.2), Lemma 2.3, the Hölder inequality, (2.3), and the definition of \mathcal{M} , we know that, for any $x \in \widetilde{B}^c$ and $h \in \mathcal{G}_0^\eta(\beta, \gamma)$ satisfying that $\|h\|_{\mathcal{G}(x, r, \beta, \gamma)} \leq 1$ for some $r \in (0, \infty)$,

$$\begin{aligned} |\langle a, h \rangle| &= \left| \int_B a(y) [h(y) - h(x_B)] d\mu(y) \right| \leq \int_B |a(y)| |h(y) - h(x_B)| d\mu(y) \\ &\lesssim \int_B |a(y)| \left[\frac{\rho(x_B, y)}{r + \rho(x, x_B)} \right]^\beta \frac{1}{V_r(x) + V(x, x_B)} \left[\frac{r}{r + \rho(x, x_B)} \right]^\gamma d\mu(y) \\ &\lesssim \int_B |a(y)| d\mu(y) \left[\frac{r_B}{\rho(x_B, x)} \right]^\beta \frac{1}{V(x_B, x)} \lesssim \frac{\mu(B)}{\|\mathbf{1}_B\|_{Y(\mathcal{X})}} \left[\frac{r_B}{\rho(x_B, x)} \right]^\beta \frac{1}{V(x_B, x)} \\ &\lesssim \frac{1}{\|\mathbf{1}_B\|_{Y(\mathcal{X})}} \left[\frac{\mu(B)}{V(x_B, x)} \right]^{\frac{\beta+\omega}{\omega}} \lesssim \frac{1}{\|\mathbf{1}_B\|_{Y(\mathcal{X})}} [\mathcal{M}(\mathbf{1}_B)(x)]^{\frac{\beta+\omega}{\omega}}, \end{aligned}$$

which further implies that, for any $x \in \widetilde{B}^c$,

$$a^*(x) \lesssim \frac{1}{\|\mathbf{1}_B\|_{Y(\mathcal{X})}} [\mathcal{M}(\mathbf{1}_B)(x)]^{\frac{\beta+\omega}{\omega}}. \quad (4.3)$$

Let $\{a_j\}_{j \in \mathbb{N}}$ be a sequence of $(Y(\mathcal{X}), q)$ -atoms supported, respectively, in balls $\{B_j\}_{j \in \mathbb{N}}$ of \mathcal{X} , and $\{\lambda_j\}_{j \in \mathbb{N}} \subset [0, \infty)$. We first prove that $\sum_{j \in \mathbb{N}} \lambda_j a_j$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. Without loss of generality, we may assume that $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$ with $\|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \leq 1$. Let x_0 be as in Definition 3.1. Then, by an argument similar to that used in the proof of (3.9), (ii) and (iv) of Definition 2.4, (4.2), and (4.3), we know that

$$\begin{aligned} \sum_{j \in \mathbb{N}} \lambda_j |\langle a_j, \varphi \rangle| &\lesssim \sum_{j \in \mathbb{N}} \lambda_j \inf_{x \in B(x_0, 1)} a_j^*(x) \lesssim \inf_{x \in B(x_0, 1)} \sum_{j \in \mathbb{N}} \lambda_j a_j^*(x) \\ &\lesssim \left\| \sum_{j \in \mathbb{N}} \lambda_j a_j^* \right\|_{Y(\mathcal{X})} \|\mathbf{1}_{B(x_0, 1)}\|_{Y(\mathcal{X})}^{-1} \lesssim \left\| \sum_{j \in \mathbb{N}} \lambda_j a_j^* \mathbf{1}_{\widetilde{B}_j} \right\|_{Y(\mathcal{X})} + \left\| \sum_{j \in \mathbb{N}} \lambda_j a_j^* \mathbf{1}_{\widetilde{B}_j^c} \right\|_{Y(\mathcal{X})} \\ &\lesssim \mathbf{I}_1 + \mathbf{I}_2, \end{aligned} \quad (4.4)$$

where, for any $j \in \mathbb{N}$, $\widetilde{B}_j := 2A_0B_j$, the implicit positive constants depend on x_0 ,

$$\mathbf{I}_1 := \left\| \sum_{j \in \mathbb{N}} \lambda_j \mathcal{M}(a_j) \mathbf{1}_{\widetilde{B}_j} \right\|_{Y(\mathcal{X})},$$

and

$$\mathbf{I}_2 := \left\| \sum_{j \in \mathbb{N}} \frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} [\mathcal{M}(\mathbf{1}_{B_j})]^{\frac{\beta+\omega}{\omega}} \right\|_{Y(\mathcal{X})}.$$

For I_1 , by $q \in (\max\{p_0, 1\}, \infty]$, the Hölder inequality, and the boundedness of \mathcal{M} on $L^q(\mathcal{X})$ (see, for instance, [14, (3.6)]), we know that, for any $j \in \mathbb{N}$,

$$\text{supp} \left(\mathcal{M}(a_j) \mathbf{1}_{\widetilde{B}_j} \|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})} \right) \subset \widetilde{B}_j$$

and

$$\left\| \mathcal{M}(a_j) \mathbf{1}_{\widetilde{B}_j} \|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})} \right\|_{L^{p_0}(\mathcal{X})} \lesssim \|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})} \|\mathcal{M}(a_j)\|_{L^q(\mathcal{X})} [\mu(\widetilde{B}_j)]^{\frac{1}{p_0} - \frac{1}{q}} \lesssim [\mu(\widetilde{B}_j)]^{\frac{1}{p_0}}.$$

From this, Lemma 4.4, Definition 2.4(ii), $p_0 > \theta_0$, Assumption 2.15, Lemma 4.3, Remark 2.14, and $d \in (0, \theta_0]$, we deduce that

$$\begin{aligned} I_1 &\leq \left\| \left\{ \sum_{j \in \mathbb{N}} [\lambda_j \mathcal{M}(a_j) \mathbf{1}_{\widetilde{B}_j}]^{\theta_0} \right\}^{\frac{1}{\theta_0}} \right\|_{Y(\mathcal{X})} \lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^{\theta_0} \mathbf{1}_{\widetilde{B}_j} \right\}^{\frac{1}{\theta_0}} \right\|_{Y(\mathcal{X})} \\ &\lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^{\theta_0} \mathbf{1}_{B_j} \right\}^{\frac{1}{\theta_0}} \right\|_{Y(\mathcal{X})} \lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})} < \infty. \end{aligned} \quad (4.5)$$

For I_2 , by Definition 2.10, $\beta > \omega(1/\theta_0 - 1)$, $\theta_0 < \underline{p}$, Assumption 2.12, Lemma 4.4, and $d \in (0, \theta_0] \subset (0, 1)$, we conclude that

$$\begin{aligned} I_2 &= \left\| \left\{ \sum_{j \in \mathbb{N}} \frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} [\mathcal{M}(\mathbf{1}_{B_j})]^{\frac{\beta+\omega}{\omega}} \right\}^{\frac{\omega}{\beta+\omega}} \right\|_{Y^{\frac{\beta+\omega}{\omega}}(\mathcal{X})} \\ &\lesssim \left\| \sum_{j \in \mathbb{N}} \frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \mathbf{1}_{B_j} \right\|_{Y(\mathcal{X})} \lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})} < \infty, \end{aligned} \quad (4.6)$$

which, combined with (4.4) and (4.5), further implies that $\sum_{j \in \mathbb{N}} \lambda_j a_j$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$.

Next, we prove (4.1). Define $f := \sum_{j \in \mathbb{N}} \lambda_j a_j$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. Then, from (4.4), (4.5), and (4.6), it follows that

$$\begin{aligned} \|f\|_{H_Y^*(\mathcal{X})} &= \|f^*\|_{Y(\mathcal{X})} \lesssim \left\| \sum_{j \in \mathbb{N}} \lambda_j a_j^* \mathbf{1}_{\widetilde{B}_j} \right\|_{Y(\mathcal{X})} + \left\| \sum_{j \in \mathbb{N}} \lambda_j a_j^* \mathbf{1}_{\widetilde{B}_j^c} \right\|_{Y(\mathcal{X})} \\ &\lesssim \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})}. \end{aligned}$$

This shows (4.1) and hence finishes the proof of Proposition 4.2. \square

Now, we formulate a decomposition result of $H_Y^*(\mathcal{X})$ (see [59, Theorem 3.7] for the corresponding Euclidean case).

Proposition 4.5. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $d \in (0, \theta_0]$, $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$, and $f \in H_Y^*(X)$. Then there exist a sequence $\{a_j\}_{j \in \mathbb{N}}$ of $(Y(X), \infty)$ -atoms supported, respectively, in the balls $\{B_j\}_{j \in \mathbb{N}}$, and a sequence $\{\lambda_j\}_{j \in \mathbb{N}} \subset [0, \infty)$ such that*

$$f = \sum_{j \in \mathbb{N}} \lambda_j a_j$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and there exists a positive constant C , independent of f , such that

$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \leq C \|f\|_{H_Y^*(X)}.$$

To prove Proposition 4.5, we need several technical lemmas. The following two lemmas are, respectively, generalizations of [30, Propositions 4.4 and 4.5] from the measure μ under consideration to any given doubling measure ν , whose proofs are slight modifications of [30, Propositions 4.4 and 4.5]; we omit the details here.

Lemma 4.6. *Suppose that ν is a doubling measure, $\Omega \subset X$ an open set with $\nu(\Omega) \in (0, \infty)$, and $A \in [1, \infty)$. For any $x \in \Omega$, let*

$$r(x) := \frac{\rho(x, \Omega^c)}{2AA_0} \in (0, \infty)$$

with A_0 as in (2.1). Then there exist an $L_0 \in \mathbb{N}$ and a sequence $\{x_k\}_{k \in I} \subset \Omega$, where I is a countable index set, such that

- (i) $\{B(x_k, r_k/(5A_0^3))\}_{k \in I}$ is disjoint. Here and thereafter, $r_k := r(x_k)$ for any $k \in I$;
- (ii) $\bigcup_{k \in I} B(x_k, r_k) = \Omega$ and $B(x_k, Ar_k) \subset \Omega$ for any $k \in I$;
- (iii) for any $k \in I$ and $x \in B(x_k, r_k)$, $Ar_k \leq \rho(x, \Omega^c) \leq 3AA_0^2 r_k$;
- (iv) for any $k \in I$, there exists a $y_k \notin \Omega$ such that $\rho(x_k, y_k) < 3AA_0 r_k$;
- (v) for any given $k \in I$, the number of balls $\{B(x_j, Ar_j)\}_{j \in I}$ that intersect $B(x_k, Ar_k)$ is at most L_0 ;
- (vi) if, in addition, Ω is bounded, then, for any $\sigma \in (0, \infty)$, the set $\{k \in I : r_k > \sigma\}$ is finite.

Lemma 4.7. *Let ν be a doubling measure, $\Omega \subset X$ an open set with $\nu(\Omega) \in (0, \infty)$, and A_0 as in (2.1). Suppose that sequences $\{x_k\}_{k \in I}$ and $\{r_k\}_{k \in I}$ are as in Lemma 4.6 with $A := 16A_0^4$ therein. Then there exist non-negative functions $\{\phi_k\}_{k \in I}$ such that*

- (i) for any $k \in I$, $0 \leq \phi_k \leq 1$ and $\text{supp } \phi_k \subset B(x_k, 2A_0 r_k)$;
- (ii) $\sum_{k \in I} \phi_k = \mathbf{1}_\Omega$;

- (iii) for any $k \in I$, $\phi_k \geq L_0^{-1}$ in $B(x_k, r_k)$, where L_0 is as in Lemma 4.6;
- (iv) there exists a positive constant C such that, for any $k \in I$, $\|\phi_k\|_{\mathcal{G}(x_k, r_k, \eta, \eta)} \leq CV_{r_k}(x_k)$ with η as in Definition 3.3.

Let $Y(X)$, β , γ , and η be as in Proposition 4.5. Assume that $f \in H_Y^*(X)$. From Lemma 3.10, it follows that $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$. Next, we aim to obtain the Calderón–Zygmund decomposition of f by applying Lemmas 4.6 and 4.7 to the level set $\{x \in X : f^*(x) > \lambda\}$ with $\lambda \in (0, \infty)$. However, this level set may not be open even in the case that ρ is a metric. To solve this problem, we borrow some ideas from [23, Section 2] and [50, Theorem 2] (see also [30, Section 4.2]).

For the quasi-metric ρ , by the proof of [50, Theorem 2], we know that there exist a $\theta \in (0, 1)$ and a metric $\tilde{\rho}$ such that $\tilde{\rho} \sim \rho^\theta$. For any $x \in X$ and $r \in (0, \infty)$, define the $\tilde{\rho}$ -ball $\tilde{B}(x, r)$ by setting

$$\tilde{B}(x, r) := \{y \in X : \tilde{\rho}(x, y) < r\}.$$

Then $(X, \tilde{\rho}, \mu)$ is a doubling metric measure space and, for any $x, y \in X$ and $r \in (0, \infty)$,

$$\mu(B(x, r + \rho(x, y))) \sim \mu(\tilde{B}(x, [r + \rho(x, y)]^\theta)) \sim \mu(\tilde{B}(x, r^\theta + \tilde{\rho}(x, y))), \quad (4.7)$$

where the implicit positive constants are independent of x, y , and r . Based on the metric $\tilde{\rho}$, a variant of the space of test functions was introduced in [30, Definition 4.6].

Definition 4.8. For any $x_1 \in X$, $R \in (0, \infty)$, and $\tilde{\beta}, \tilde{\gamma} \in (0, \infty)$, the space $G(x_1, R, \tilde{\beta}, \tilde{\gamma})$ is defined to be the set of all functions f on X satisfying that there exists a positive constant C such that

- (i) (the *size condition*) for any $x \in X$,

$$|f(x)| \leq C \frac{1}{\mu(\tilde{B}(x, R + \tilde{\rho}(x_1, x)))} \left[\frac{R}{R + \tilde{\rho}(x_1, x)} \right]^{\tilde{\gamma}};$$

- (ii) (the *regularity condition*) for any $x, y \in X$ satisfying $\tilde{\rho}(x, y) \leq [R + \tilde{\rho}(x_1, x)]/2$,

$$|f(x) - f(y)| \leq C \left[\frac{\tilde{\rho}(x, y)}{R + \tilde{\rho}(x_1, x)} \right]^{\tilde{\beta}} \frac{1}{\mu(\tilde{B}(x, R + \tilde{\rho}(x_1, x)))} \left[\frac{R}{R + \tilde{\rho}(x_1, x)} \right]^{\tilde{\gamma}}.$$

For any $f \in G(x_1, R, \tilde{\beta}, \tilde{\gamma})$, its norm $\|f\|_{G(x_1, R, \tilde{\beta}, \tilde{\gamma})}$ in $G(x_1, R, \tilde{\beta}, \tilde{\gamma})$ is defined by setting

$$\|f\|_{G(x_1, R, \tilde{\beta}, \tilde{\gamma})} := \inf \{C \in (0, \infty) : \text{(i) and (ii) hold true}\}.$$

By Lemma 2.3 and (4.7), we know that there exists a $\theta \in (0, 1)$ such that

$$\mathcal{G}(x_1, r, \beta, \gamma) = G(x_1, r^\theta, \beta/\theta, \gamma/\theta)$$

with equivalent norms, and the positive equivalence constants are independent of x_1 and r . For any $\beta, \gamma \in (0, \eta)$ and $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$, define the *modified grand maximal function* f^* of f by setting, for any $x \in X$,

$$f^*(x) := \sup \left\{ |\langle f, \varphi \rangle| : \varphi \in \mathcal{G}_0^\eta(\beta, \gamma) \text{ with } \|\varphi\|_{G(x, r^\theta, \beta/\theta, \gamma/\theta)} \leq 1 \text{ for some } r \in (0, \infty) \right\}.$$

It is easy to see that $f^* \sim f^*$ pointwisely on X .

Next, we establish an embedding lemma for $Y(X)$ as in Proposition 4.5, which is actually an application of Theorem 2.17.

Lemma 4.9. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $x_0 \in X$. Then there exists an $\epsilon \in (0, 1)$ such that $\bar{Y}(X)$ is continuously embedded into $L_w^{\theta_0}(X)$ with $w := [\mathcal{M}(\mathbf{1}_{B(x_0, 1)})]^\epsilon \in A_1(X)$.*

Proof. By Assumption 2.15 and (2.10) with θ_1 and θ_2 replaced, respectively, by 1 and $(p_0/\theta_0)'$, we know that $Y^{1/\theta_0}(X)$ is a ball Banach function space and \mathcal{M} bounded on $(Y^{1/\theta_0})'(X)$. From this and Theorem 2.17, we further deduce that there exists an $\epsilon \in (0, 1)$ such that $Y(X)$ is continuously embedded into $L_w^{\theta_0}(X)$ with $w := [\mathcal{M}(\mathbf{1}_{B(x_0, 1)})]^\epsilon \in A_1(X)$. This finishes the proof of Lemma 4.9. \square

The following lemma shows that the set $\{x \in X : f^\star(x) > \lambda\}$ with $\lambda \in (0, \infty)$ is an open and μ_w -measurable set with finite measure.

Lemma 4.10. *Let $f \in H_Y^*(X)$. For any $\lambda \in (0, \infty)$ and any μ -measurable $E \subset X$, define*

$$\Omega_\lambda := \{x \in X : f^\star(x) > \lambda\}$$

and

$$\mu_w(E) := w(E) := \int_E w(x) d\mu(x),$$

where w is as in Lemma 4.9. Then the following two statements hold true:

- (i) μ_w is a doubling measure;
- (ii) for any given λ , Ω_λ is open under the topology induced by ρ , and $\mu_w(\Omega_\lambda) < \infty$.

Proof. First, by (2.3), the fact that $w \in A_1(X)$, and [65, Chapter 1, Lemma 12], we know that μ_w is a doubling measure, namely, (i) holds true. Next, we prove (ii). By an argument similar to that used in [23, Remark 2.9(iii)], we conclude that Ω_λ is open under the topology induced by ρ . Moreover, from the fact that $f^\star \sim f^*$, and Lemma 4.9, we deduce that, for any given $\lambda \in (0, \infty)$,

$$\begin{aligned} \mu_w(\Omega_\lambda) &= \int_{\Omega_\lambda} w(x) d\mu(x) \leq \int_X \left[\frac{f^\star(x)}{\lambda} \right]^{\theta_0} w(x) d\mu(x) \\ &\sim \lambda^{-\theta_0} \|f^*\|_{L_w^{\theta_0}(X)}^{\theta_0} \lesssim \lambda^{-\theta_0} \|f^*\|_{Y(X)}^{\theta_0} \sim \lambda^{-\theta_0} \|f\|_{H_Y^*(X)}^{\theta_0} < \infty, \end{aligned}$$

where θ_0 is as in Lemma 4.9 and the implicit positive constants are independent of λ and f . This proves (ii) and hence finishes the proof of Lemma 4.10. \square

Let $Y(X)$, β , γ , and η be as in Proposition 4.5. Assume that $f \in H_Y^*(X)$. To decompose f into a sum of $(Y(X), \infty)$ -atoms, we let $\{P_m\}_{m \in \mathbb{Z}}$ be a 1-exp-ATI and, for any $m \in \mathbb{Z}$, let

$$f^{(m)} := P_m f. \quad (4.8)$$

In the remainder of this section, we always assume that $m \in \mathbb{N}$. By the fact that $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$, Lemmas 3.5(iii), 3.6, and 3.7, and an argument similar to that used in [77, p. 210], we know that

$$f^{(m)} \rightarrow f \quad (4.9)$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ as $m \rightarrow \infty$ and, for any $m \in \mathbb{N}$ and $x \in X$, $(f^{(m)})^*(x) \lesssim f^*(x)$ with the implicit positive constant independent of m , f , and x . Next, we deal with $f^{(m)}$. To this end, for any $j \in \mathbb{Z}$, let

$$\Omega_j := \{x \in X : f^*(x) > 2^j\}. \quad (4.10)$$

By Lemma 4.10(ii), we know that, for any $j \in \mathbb{Z}$, Ω_j is open and $\mu_w(\Omega_j) < \infty$. From this, and Lemmas 4.6 and 4.7 with $\nu := \mu_w$ therein, we further deduce that, for any $j \in \mathbb{Z}$, there exist $\{x_{j,k}\}_{k \in I_j} \subset X$ with I_j being a countable index set, $\{r_{j,k}\}_{k \in I_j} \subset (0, \infty)$, $L_0 \in \mathbb{N}$, and a sequence $\{\phi_{j,k}\}_{k \in I_j}$ of non-negative functions satisfying all the conclusions of Lemmas 4.6 and 4.7. For any $m \in \mathbb{N}$, $j \in \mathbb{Z}$, and $k \in I_j$, let

$$P_{j,k}^{(m)} := \frac{1}{\|\phi_{j,k}\|_{L^1(X)}} \int_X f^{(m)}(\xi) \phi_{j,k}(\xi) d\mu(\xi) \quad \text{and} \quad b_{j,k}^{(m)} := (f^{(m)} - P_{j,k}^{(m)}) \phi_{j,k}. \quad (4.11)$$

We then have the following technical lemma.

Lemma 4.11. *For any $m \in \mathbb{N}$, $j \in \mathbb{Z}$, and $k \in I_j$, let $b_{j,k}^{(m)}$ be as in (4.11). Then the following statements hold true:*

(i) *there exists a positive constant C such that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,*

$$\left\| \sum_{k \in I_j} (b_{j,k}^{(m)})^* \right\|_{Y(X)} \leq C \|f^* \mathbf{1}_{\Omega_j}\|_{Y(X)}; \quad (4.12)$$

moreover, $b_j^{(m)} := \sum_{k \in I_j} b_{j,k}^{(m)}$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ satisfying that $b_j^{(m)} \in H_Y^(X)$ and, for any $x \in X$,*

$$(b_j^{(m)})^*(x) \leq C 2^j \sum_{k \in I_j} \frac{V_{r_{j,k}}(x_{j,k})}{V_{r_{j,k}}(x_{j,k}) + V(x_{j,k}, x)} \left[\frac{r_{j,k}}{r_{j,k} + \rho(x_{j,k}, x)} \right]^\beta + C f^*(x) \mathbf{1}_{\Omega_j}(x); \quad (4.13)$$

(ii) *for any $m \in \mathbb{N}$, $b_j^{(m)} \rightarrow 0$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ as $j \rightarrow \infty$;*

(iii) *for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, let $g_j^{(m)} := f^{(m)} - b_j^{(m)}$. Then, for any $m \in \mathbb{N}$, $g_j^{(m)} \rightarrow 0$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ as $j \rightarrow -\infty$.*

Proof. Let all the symbols be as in the present lemma and

$$J := \frac{V_{r_{j,k}}(x_{j,k})}{V_{r_{j,k}}(x_{j,k}) + V(x_{j,k}, x)} \left[\frac{r_{j,k}}{r_{j,k} + \rho(x_{j,k}, x)} \right]^\beta.$$

We first prove (i). By [77, (4.21)] and Definition 2.4(ii), we conclude that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,

$$\left\| \sum_{k \in I_j} (b_{j,k}^{(m)})^* \right\|_{Y(X)} \lesssim \left\| \sum_{k \in I_j} f^* \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{Y(X)} + \left\| \sum_{k \in I_j} 2^j J \mathbf{1}_{[B(x_{j,k}, Ar_{j,k})]^c} \right\|_{Y(X)}$$

$$=: I_1 + I_2, \quad (4.14)$$

where $A := 16A_0^4$. By (ii) and (v) of Lemma 4.6, we obtain

$$I_1 \lesssim \|f^* \mathbf{1}_{\Omega_j}\|_{Y(\mathcal{X})}. \quad (4.15)$$

To estimate I_2 , since $\beta \in (\omega(1/\theta_0 - 1), \eta)$, it follows that we can choose $h \in (\frac{\omega}{\omega+\beta}, \theta_0)$. By Definition 2.4(ii), Lemma 2.13, (2.3), Definition 2.10, $\omega(1/h - 1) < \beta$, Assumption 2.12, (4.10) with the fact that $f^\star \sim f^*$, and (ii) and (v) of Lemma 4.6, we conclude that

$$\begin{aligned} I_2 &\sim \left\| \sum_{k \in I_j} \sum_{s \in \mathbb{Z}_+} 2^j J \mathbf{1}_{B(x_{j,k}, 2^{s+1} Ar_{j,k}) \setminus B(x_{j,k}, 2^s Ar_{j,k})} \right\|_{Y(\mathcal{X})} \\ &\lesssim \left\| \sum_{k \in I_j} 2^j \sum_{s \in \mathbb{Z}_+} 2^{-s\beta} \frac{V_{r_{j,k}}(x_{j,k})}{V_{2^s Ar_{j,k}}(x_{j,k})} \mathbf{1}_{B(x_{j,k}, 2^{s+1} Ar_{j,k})} \right\|_{Y(\mathcal{X})} \\ &\sim \left\| \sum_{k \in I_j} 2^j \sum_{s \in \mathbb{Z}_+} 2^{-s\beta} \left[\frac{V_{2^s Ar_{j,k}}(x_{j,k})}{V_{Ar_{j,k}}(x_{j,k})} \right]^{\frac{1}{h}-1} \left[\frac{V_{Ar_{j,k}}(x_{j,k})}{V_{2^s Ar_{j,k}}(x_{j,k})} \right]^{\frac{1}{h}} \mathbf{1}_{B(x_{j,k}, 2^{s+1} Ar_{j,k})} \right\|_{Y(\mathcal{X})} \\ &\lesssim \left\| \sum_{k \in I_j} 2^j \sum_{s \in \mathbb{Z}_+} 2^{[\omega(\frac{1}{h}-1)-\beta]s} \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}) \right]^{\frac{1}{h}} \right\|_{Y(\mathcal{X})} \lesssim \left\| \left\{ \sum_{k \in I_j} 2^j \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}) \right]^{\frac{1}{h}} \right\}^h \right\|_{Y^{\frac{1}{h}}(\mathcal{X})}^{\frac{1}{h}} \\ &\lesssim \left\| \sum_{k \in I_j} 2^j \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{Y(\mathcal{X})} \lesssim \left\| \sum_{k \in I_j} f^* \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{Y(\mathcal{X})} \lesssim \|f^* \mathbf{1}_{\Omega_j}\|_{Y(\mathcal{X})}, \end{aligned}$$

which, combined with (4.14) and (4.15), further implies (4.12).

Next, we show that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, $\sum_{k \in I_j} b_{j,k}^{(m)}$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. Without loss of generality, we may assume that $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$ with $\|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \leq 1$. Let $x_0 \in \mathcal{X}$ be as in Definition 3.1. From an argument similar to that used in the proof of (3.9), (ii) and (iv) of Definition 2.4, and (4.12), we deduce that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,

$$\begin{aligned} \sum_{k \in I_j} |\langle b_{j,k}^{(m)}, \varphi \rangle| &\lesssim \sum_{k \in I_j} \inf_{x \in B(x_0, 1)} (b_{j,k}^{(m)})^*(x) \lesssim \inf_{x \in B(x_0, 1)} \sum_{k \in I_j} (b_{j,k}^{(m)})^*(x) \\ &\lesssim \left\| \sum_{k \in I_j} (b_{j,k}^{(m)})^* \right\|_{Y(\mathcal{X})} \|\mathbf{1}_{B(x_0, 1)}\|_{Y(\mathcal{X})}^{-1} \sim \left\| \sum_{k \in I_j} (b_{j,k}^{(m)})^* \right\|_{Y(\mathcal{X})} \\ &\lesssim \|f^* \mathbf{1}_{\Omega_j}\|_{Y(\mathcal{X})} \lesssim \|f\|_{H_Y^*(\mathcal{X})} < \infty, \end{aligned}$$

which further implies that $\sum_{k \in I_j} b_{j,k}^{(m)}$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, where the implicit positive constants depend on x_0 . Let $b_j^{(m)} := \sum_{k \in I_j} b_{j,k}^{(m)}$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. Then, by (4.12) and [77, (4.21)], we obtain $b_j^{(m)} \in H_Y^*(\mathcal{X})$ and (4.13). This finishes the proof of (i).

Next, we prove (ii). Without loss of generality, we may assume that $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$ with $\|\varphi\|_{\mathcal{G}_0^\eta(\beta, \gamma)} \leq 1$. Let $x_0 \in \mathcal{X}$ be as in Definition 3.1. By an argument similar to that used in the

proof of (3.9), we conclude that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,

$$\left| \langle b_j^{(m)}, \varphi \rangle \right| \lesssim \inf_{x \in B(x_0, 1)} \left(b_j^{(m)} \right)^*(x) \lesssim \left\| \left(b_j^{(m)} \right)^* \right\|_{L_w^{\theta_0}(\mathcal{X})} \left\| \mathbf{1}_{B(x_0, 1)} \right\|_{L_w^{\theta_0}(\mathcal{X})}^{-1} \lesssim \left\| \left(b_j^{(m)} \right)^* \right\|_{L_w^{\theta_0}(\mathcal{X})}, \quad (4.16)$$

where w is as in Lemma 4.9 and the implicit positive constants depend on x_0 . Moreover, on one hand, by (2.6), we know that, for any $j \in \mathbb{Z}$, $k \in I_j$, and $x \in B(x_{j,k}, 2A_0Ar_{j,k})$,

$$J \leq 1 \lesssim \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})})(x) \right]^{\frac{\beta+\omega}{\omega}};$$

on the other hand, by (2.3), we conclude that, for any $j \in \mathbb{Z}$, $k \in I_j$, and $x \in (B(x_{j,k}, 2A_0Ar_{j,k}))^c$,

$$J \leq \frac{V_{r_{j,k}}(x_{j,k})}{V(x_{j,k}, x)} \left[\frac{r_{j,k}}{\rho(x_{j,k}, x)} \right]^\beta \lesssim \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})})(x) \right]^{\frac{\beta+\omega}{\omega}}.$$

To summarize, for any $j \in \mathbb{Z}$, $k \in I_j$, and $x \in \mathcal{X}$, it holds true that

$$J \lesssim \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})})(x) \right]^{\frac{\beta+\omega}{\omega}}.$$

From this, (4.16), (4.13), the facts that $\beta > \omega(1/\theta_0 - 1)$ and $f^\star \sim f^*$, the Fefferman–Stein vector-valued maximal inequality of $L_w^{[\theta_0(\beta+\omega)]/\omega}(\mathcal{X})$ (see [19, Theorem 6.5(ii)]), and (ii) and (v) of Lemma 4.6, we deduce that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$,

$$\begin{aligned} \left| \langle b_j^{(m)}, \varphi \rangle \right| &\lesssim \left\| 2^j \sum_{k \in I_j} \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}) \right]^{\frac{\beta+\omega}{\omega}} \right\|_{L_w^{\theta_0}(\mathcal{X})} + \|f^* \mathbf{1}_{\Omega_j}\|_{L_w^{\theta_0}(\mathcal{X})} \\ &\sim \left\| \left\{ 2^j \sum_{k \in I_j} \left[\mathcal{M}(\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}) \right]^{\frac{\beta+\omega}{\omega}} \right\}^{\frac{\omega}{\beta+\omega}} \right\|_{L_w^{\frac{\theta_0(\beta+\omega)}{\omega}}(\mathcal{X})}^{\frac{\beta+\omega}{\omega}} + \|f^* \mathbf{1}_{\Omega_j}\|_{L_w^{\theta_0}(\mathcal{X})} \\ &\lesssim \left\| 2^j \sum_{k \in I_j} \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{L_w^{\theta_0}(\mathcal{X})} + \|f^* \mathbf{1}_{\Omega_j}\|_{L_w^{\theta_0}(\mathcal{X})} \lesssim \|f^* \mathbf{1}_{\Omega_j}\|_{L_w^{\theta_0}(\mathcal{X})}, \end{aligned}$$

which, combined with the dominated convergence theorem and the fact that $\Omega_j \downarrow \emptyset$ as $j \rightarrow \infty$, then completes the proof of (ii).

Finally, we prove (iii). By Lemmas 3.6, 3.7, 4.6, and 4.7, and an argument similar to that used in the proof of [77, (4.26)], we conclude that, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, $\|g_j^{(m)}\|_{L^\infty(\mathcal{X})} \lesssim 2^j$, which further implies (iii) and hence completes the proof of Lemma 4.11. \square

Next, we recall the atomic characterization of Musielak–Orlicz Hardy spaces on \mathcal{X} , which plays a key role in the proof of Proposition 4.5 and was originally established in [19]. To this end, we first recall some notions on Musielak–Orlicz Hardy spaces.

Definition 4.12. Let $\beta, \gamma \in (0, \eta)$ with η as in Definition 3.3, and φ be a growth function as in Remark 2.7(iii). Then the Musielak–Orlicz Hardy space $H^{*,\varphi}(\mathcal{X})$ is defined by setting

$$H^{*,\varphi}(\mathcal{X}) := \left\{ f \in \left(\mathcal{G}_0^\eta(\beta, \gamma) \right)' : \|f\|_{H^{*,\varphi}(\mathcal{X})} := \|f^*\|_{L^\varphi(\mathcal{X})} < \infty \right\}.$$

Definition 4.13. Let E be a μ -measurable subset of \mathcal{X} and $q \in [1, \infty]$, the space $L_\varphi^q(E)$ is defined to be the set of all μ -measurable functions f on E such that

$$\|f\|_{L_\varphi^q(E)} := \begin{cases} \sup_{t \in (0, \infty)} \left[\frac{1}{\varphi(E, t)} \int_E |f(x)|^q \varphi(x, t) d\mu(x) \right]^{1/q} & \text{when } q \in [1, \infty), \\ \|f\|_{L^\infty(E)} & \text{when } q = \infty \end{cases}$$

is finite, here and thereafter, $\varphi(E, t) := \int_E \varphi(x, t) d\mu(x)$.

Definition 4.14. Let ω be as in (2.3), φ a growth function as in Remark 2.7(iii), $q \in (q(\varphi), \infty]$, and

$$m(\varphi) := \left\lfloor \omega \left[\frac{q(\varphi)}{i(\varphi)} - 1 \right] \right\rfloor \leq 0,$$

where $q(\varphi)$ is as in (2.4) and

$$i(\varphi) := \sup \{ p \in (0, \infty) : \varphi \text{ is of uniformly lower type } p \}.$$

A μ -measurable function a is called a (φ, q) -atom supported in a ball $B \subset \mathcal{X}$ if the following three conditions hold true:

- (i) $\text{supp } a := \{x \in \mathcal{X} : a(x) \neq 0\} \subset B$;
- (ii) $a \in L_\varphi^q(B)$ and $\|a\|_{L_\varphi^q(B)} \leq \|\mathbf{1}_B\|_{L^\varphi(\mathcal{X})}^{-1}$;
- (iii) $\int_{\mathcal{X}} a(x) d\mu(x) = 0$.

The following lemma is just [19, Lemma 4.9].

Lemma 4.15. Let $\beta, \gamma \in (0, \eta)$ with η as in Definition 3.3, and φ be as in Definition 4.14. Then $H^{*,\varphi}(\mathcal{X})$ continuously embeds into $(\mathcal{G}_0^\eta(\beta, \gamma))'$.

The following lemma is just [19, Lemma 2.8(ii)] (see [73, Lemma 1.1.10(i)] for the corresponding Euclidean case).

Lemma 4.16. Let φ be as in Definition 4.14. Then, for any $f \in L^\varphi(\mathcal{X}) \setminus \{0\}$,

$$\int_{\mathcal{X}} \varphi \left(x, \frac{|f(x)|}{\|f\|_{L^\varphi(\mathcal{X})}} \right) d\mu(x) = 1.$$

To prove Proposition 4.5, we also need the following lemma.

Lemma 4.17. Let φ and q be as in Definition 4.14. Then there exists a positive constant C such that, for any (φ, q) -atom a , $\|a\|_{H^{*,\varphi}(\mathcal{X})} \leq C$.

Proof. Let a be a (φ, q) -atom supported in B . Then, by [19, (5.2)], the fact that φ is a growth function, and Lemma 4.16, we know that

$$\int_{\mathcal{X}} \varphi(x, a^*(x)) d\mu(x) \lesssim \varphi(B, \|a\|_{L_\varphi^q(B)}) \lesssim \varphi(B, \|\mathbf{1}_B\|_{L^\varphi(\mathcal{X})}^{-1}) \sim 1,$$

which further implies that

$$\|a\|_{H^{*,\varphi}(\mathcal{X})} = \|a^*\|_{L^\varphi(\mathcal{X})} \lesssim 1.$$

This finishes the proof of Lemma 4.17. \square

Now, we give the proof of Proposition 4.5.

Proof of Proposition 4.5. Let all the symbols be as in the present proposition. For any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, let $f^{(m)}$ be as in (4.8), and $b_j^{(m)}$ and $g_j^{(m)}$ as in Lemma 4.11. Then, by Lemma 4.11, we know that, for any $m \in \mathbb{N}$, and any $J_1, J_2 \in \mathbb{N}$,

$$\begin{aligned} f^{(m)} - \sum_{j=-J_1}^{J_2} [g_{j+1}^{(m)} - g_j^{(m)}] &= f^{(m)} - [g_{J_2+1}^{(m)} - g_{-J_1}^{(m)}] \\ &= b_{J_2+1}^{(m)} + g_{-J_1}^{(m)} \rightarrow 0 \end{aligned}$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, as $J_1, J_2 \rightarrow \infty$, which further implies that

$$f^{(m)} = \sum_{j \in \mathbb{Z}} [g_{j+1}^{(m)} - g_j^{(m)}] \quad (4.17)$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. For any $m \in \mathbb{N}$, $j \in \mathbb{Z}$, $k \in I_j$, and $\ell \in I_{j+1}$, let $\phi_{j,k}$, $P_{j,k}^{(m)}$, and $b_{j,k}^{(m)}$ be as in (4.11),

$$L_{j+1,k}^{(m),\ell} := \frac{1}{\|\phi_{j+1,\ell}\|_{L^1(X)}} \int_X [f^{(m)}(\xi) - P_{j+1,\ell}^{(m)}(\xi)] \phi_{j,k}(\xi) \phi_{j+1,\ell}(\xi) d\mu(\xi),$$

and

$$h_{j,k}^{(m)} := b_{j,k}^{(m)} - \sum_{\ell \in I_{j+1}} [b_{j+1,\ell}^{(m)} \phi_{j,k} - L_{j+1,k}^{(m),\ell} \phi_{j+1,\ell}].$$

Then, by (4.17) and an argument similar to that used in the proof of [77, p. 214], we conclude that, for any $m \in \mathbb{N}$, $j \in \mathbb{Z}$, and $k \in I_j$,

$$\|h_{j,k}^{(m)}\|_{L^\infty(X)} \lesssim 2^j, \quad \text{supp } h_{j,k}^{(m)} \subset B(x_{j,k}, Ar_{j,k}), \quad \int_X h_{j,k}^{(m)}(x) d\mu(x) = 0, \quad (4.18)$$

and

$$g_{j+1}^{(m)} - g_j^{(m)} = \sum_{k \in I_j} h_{j,k}^{(m)} \quad (4.19)$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, where $A := 16A_0^4$. Moreover, by (4.18) and the Banach–Alaoglu theorem (see, for instance, [57, Theorem 3.17]), together with the well-known diagonal rule for series two times, we conclude that there exist a subsequence $\{m_q\}_{q=1}^\infty \subset \mathbb{N}$ and a sequence $\{h_{j,k}\}_{k \in I_j} \subset L^\infty(X)$ with any $j \in \mathbb{Z}$ such that, for any $j \in \mathbb{Z}$ and $k \in I_j$,

$$\|h_{j,k}\|_{L^\infty(X)} \lesssim 2^j, \quad \text{supp } h_{j,k} \subset B(x_{j,k}, Ar_{j,k}), \quad \int_X h_{j,k}(x) d\mu(x) = 0, \quad (4.20)$$

and, for any $g \in L^1(X)$,

$$\langle h_{j,k}^{(m_q)}, g \rangle \rightarrow \langle h_{j,k}, g \rangle$$

as $q \rightarrow \infty$. Now, we claim that, for any given $j \in \mathbb{Z}$, $\sum_{k \in I_j} h_{j,k}$ converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. Indeed, by (4.20), (ii) and (v) of Lemma 4.6, and Lemma 4.10(ii), we know that, for any given $j \in \mathbb{Z}$ and $r \in (1, \infty)$,

$$\left\| \sum_{k \in I_j} h_{j,k} \right\|_{L_w^r(X)} \lesssim 2^j \left\| \sum_{k \in I_j} \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{L_w^r(X)} \lesssim 2^j \|\mathbf{1}_{\Omega_j}\|_{L_w^r(X)} \sim 2^j [\mu_w(\Omega_j)]^{\frac{1}{r}} < \infty,$$

where w and μ_w are as in Lemma 4.10. This further implies the above claim.

For any $j \in \mathbb{Z}$ and $k \in I_j$, let

$$\lambda_{j,k} := 2^j \|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{Y(X)} \quad \text{and} \quad a_{j,k} := \frac{h_{j,k}}{\lambda_{j,k}}. \quad (4.21)$$

Then, by (4.20), we know that $a_{j,k}$ is a $(Y(X), \infty)$ -atom supported in $B(x_{j,k}, Ar_{j,k})$. Next, we prove

$$f = \sum_{j \in \mathbb{Z}} \sum_{k \in I_j} \lambda_{j,k} a_{j,k} \quad (4.22)$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and

$$K := \left\| \left\{ \sum_{j \in \mathbb{Z}} \sum_{k \in I_j} \left[\frac{\lambda_{j,k}}{\|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{Y(X)}} \right]^d \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\} \right\|_{Y(X)}^{\frac{1}{d}} \lesssim \|f\|_{H_Y^*(X)}. \quad (4.23)$$

To this end, for any $m \in \mathbb{N}$ and $j \in \mathbb{Z}$, let $f_j^{(m)} := g_{j+1}^{(m)} - g_j^{(m)}$ and $f_j := \sum_{k \in I_j} h_{j,k}$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. It suffices to show that

$$\lim_{J_3 \rightarrow -\infty} \sup_{m \in \mathbb{N}} \left\| \sum_{j \leq J_3} f_j^{(m)} \right\|_{H_w^{p_1}(X)} = 0 \quad \text{and} \quad \lim_{J_4 \rightarrow \infty} \sup_{m \in \mathbb{N}} \left\| \sum_{j \geq J_4} f_j^{(m)} \right\|_{H_w^{p_1}(X)} = 0 \quad (4.24)$$

for some $p_1 \in (\frac{\omega}{\omega+\eta}, \theta_0)$, where w is as in Lemma 4.9. Assuming this for the moment, by (4.24), Lemma 4.15, and an argument similar to that used in [77, pp. 215-217], we obtain

$$\lim_{q \rightarrow \infty} \sum_{j \in \mathbb{Z}} f_j^{(m_q)} = \sum_{j \in \mathbb{Z}} f_j$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, which, combined with (4.9), (4.17), and (4.21), further implies that

$$\begin{aligned} f &= \lim_{m \rightarrow \infty} f^{(m)} = \lim_{m \rightarrow \infty} \sum_{j \in \mathbb{Z}} [g_{j+1}^{(m)} - g_j^{(m)}] = \lim_{q \rightarrow \infty} \sum_{j \in \mathbb{Z}} f_j^{(m_q)} \\ &= \sum_{j \in \mathbb{Z}} f_j = \sum_{j \in \mathbb{Z}} \sum_{k \in I_j} h_{j,k} = \sum_{j \in \mathbb{Z}} \sum_{k \in I_j} \lambda_{j,k} a_{j,k} \end{aligned}$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and hence (4.22) holds true. Moreover, from Definition 2.4(ii), (ii) and (v) of Lemma 4.6, and the fact that $f^* \sim f^*$, we deduce that

$$K \lesssim \left\| \left\{ \sum_{j \in \mathbb{Z}} 2^{jd} \mathbf{1}_{\Omega_j} \right\} \right\|_{Y(X)}^{\frac{1}{d}} \sim \left\| \left\{ \sum_{j \in \mathbb{Z}} \sum_{r=j}^{\infty} 2^{jd} \mathbf{1}_{\Omega_r \setminus \Omega_{r+1}} \right\} \right\|_{Y(X)}^{\frac{1}{d}}$$

$$\begin{aligned}
& \sim \left\| \left\{ \sum_{r \in \mathbb{Z}} 2^{rd} \sum_{j=-\infty}^r 2^{(j-r)d} \mathbf{1}_{\Omega_r \setminus \Omega_{r+1}} \right\} \right\|_{Y(\mathcal{X})}^{\frac{1}{d}} \sim \left\| \left\{ \sum_{r \in \mathbb{Z}} 2^{rd} \mathbf{1}_{\Omega_r \setminus \Omega_{r+1}} \right\} \right\|_{Y(\mathcal{X})}^{\frac{1}{d}} \\
& \lesssim \left\| f^\star \left\{ \sum_{r \in \mathbb{Z}} \mathbf{1}_{\Omega_r \setminus \Omega_{r+1}} \right\} \right\|_{Y(\mathcal{X})}^{\frac{1}{d}} \sim \|f^\star\|_{Y(\mathcal{X})} \sim \|f\|_{H_Y^*(\mathcal{X})}.
\end{aligned} \tag{4.25}$$

Thus, (4.23) holds true, which then completes the proof of this proposition.

Now, we show (4.24). On one hand, for any $x \in \mathcal{X}$ and $t \in [0, \infty)$, let $\varphi_1(x, t) := w(x)t$ with w as in Lemma 4.9. Obviously, φ_1 satisfies all the assumptions of Lemma 4.17. From this, (4.19), (4.18), Lemma 4.17, (ii) and (v) of Lemma 4.6, the fact that $f^\star \sim f^*$, and Lemma 4.9, we deduce that, for any $m \in \mathbb{N}$ and $J_3 \in \mathbb{Z}$,

$$\begin{aligned}
\left\| \sum_{j \leq J_3} f_j^{(m)} \right\|_{H_w^1(\mathcal{X})} & \leq \sum_{j \leq J_3} \sum_{k \in I_j} \|h_{j,k}^{(m)}\|_{H_w^1(\mathcal{X})} \\
& = \sum_{j \leq J_3} \sum_{k \in I_j} 2^j \|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{L_w^1(\mathcal{X})} \left\| \frac{h_{j,k}^{(m)}}{2^j \|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{L_w^1(\mathcal{X})}} \right\|_{H_w^1(\mathcal{X})} \\
& \lesssim \sum_{j \leq J_3} \sum_{k \in I_j} 2^j w(B(x_{j,k}, Ar_{j,k})) \lesssim \sum_{j \leq J_3} 2^j w(\Omega_j) \\
& \lesssim \sum_{j \leq J_3} 2^j \int_{\Omega_j} \left[\frac{f^\star(x)}{2^j} \right]^{\theta_0} w(x) d\mu(x) \lesssim \sum_{j \leq J_3} 2^{j(1-\theta_0)} \|f^\star\|_{L_w^{\theta_0}(\mathcal{X})}^{\theta_0} \\
& \lesssim \sum_{j \leq J_3} 2^{j(1-\theta_0)} \|f^\star\|_{Y(\mathcal{X})}^{\theta_0} \sim 2^{J_3(1-\theta_0)} \|f\|_{H_Y^*(\mathcal{X})}^{\theta_0}
\end{aligned}$$

with the implicit positive constants independent of m and J_3 , which, combined with $\theta_0 \in (0, 1)$ and $J_3 \rightarrow -\infty$, further implies the first estimate of (4.24).

On the other hand, choose $p_1 \in (\frac{\omega}{\omega+\eta}, \theta_0)$ and let $\varphi_2(x, t) := w(x)t^{p_1}$ for any $x \in \mathcal{X}$ and $t \in [0, \infty)$ with w as in Lemma 4.9. Obviously, φ_2 also satisfies all the assumptions of Lemma 4.17. Using this, (4.18), Lemma 4.17, (ii) and (v) of Lemma 4.6, the fact that $f^\star \sim f^*$, and Lemma 4.9, we know that, for any $m \in \mathbb{N}$ and $J_4 \in \mathbb{Z}$,

$$\begin{aligned}
\left\| \sum_{j \geq J_4} f_j^{(m)} \right\|_{H_w^{p_1}(\mathcal{X})}^{p_1} & \leq \sum_{j \geq J_4} \sum_{k \in I_j} \|h_{j,k}^{(m)}\|_{H_w^{p_1}(\mathcal{X})}^{p_1} \\
& = \sum_{j \geq J_4} \sum_{k \in I_j} 2^{jp_1} \|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{L_w^{p_1}(\mathcal{X})}^{p_1} \left\| \frac{h_{j,k}^{(m)}}{2^{jp_1} \|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{L_w^{p_1}(\mathcal{X})}} \right\|_{H_w^{p_1}(\mathcal{X})}^{p_1} \\
& \lesssim \sum_{j \geq J_4} \sum_{k \in I_j} 2^{jp_1} w(B(x_{j,k}, Ar_{j,k})) \lesssim \sum_{j \geq J_4} 2^{jp_1} w(\Omega_j) \\
& \lesssim \sum_{j \geq J_4} 2^{jp_1} \int_{\Omega_j} \left[\frac{f^\star(x)}{2^j} \right]^{\theta_0} w(x) d\mu(x) \lesssim \sum_{j \geq J_4} 2^{j(p_1-\theta_0)} \|f^\star\|_{L_w^{\theta_0}(\mathcal{X})}^{\theta_0}
\end{aligned}$$

$$\lesssim \sum_{j \geq J_4} 2^{j(p_1 - \theta_0)} \|f^\star\|_{Y(X)}^{\theta_0} \sim 2^{J_4(p_1 - \theta_0)} \|f\|_{H_Y^*(X)}^{\theta_0}$$

with the implicit positive constants independent of m and J_4 , which, combined with $p_1 \in (\frac{\omega}{\omega + \eta}, \theta_0)$ and $J_4 \rightarrow \infty$, further implies the second estimate of (4.24) and hence completes the proof of Proposition 4.5. \square

Next, we introduce the atomic Hardy spaces associated with $Y(X)$.

Definition 4.18. Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$, $d \in (0, \theta_0]$, and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. The *atomic Hardy space* $H_{\text{atom}}^{Y,q,d}(X)$ is defined to be the set of all $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$ satisfying that there exist a sequence $\{\lambda_j\}_{j \in \mathbb{N}}$ of non-negative numbers and a sequence $\{a_j\}_{j \in \mathbb{N}}$ of $(Y(X), q)$ -atoms supported, respectively, in balls $\{B_j\}_{j \in \mathbb{N}}$ of X such that

$$f = \sum_{j \in \mathbb{N}} \lambda_j a_j$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and

$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\} \right\|_{Y(X)}^{\frac{1}{d}} < \infty.$$

Moreover, for any $f \in H_{\text{atom}}^{Y,q,d}(X)$, let

$$\|f\|_{H_{\text{atom}}^{Y,q,d}(X)} := \inf \left\{ \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\} \right\|_{Y(X)}^{\frac{1}{d}} \right\},$$

where the infimum is taken over all decompositions of f as above.

Combining Propositions 4.2 and 4.5, we obtain the following conclusion.

Theorem 4.19. Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$, $d \in (0, \theta_0]$, and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. Then

$$[H_Y^*(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))'] = [H_{\text{atom}}^{Y,q,d}(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))']$$

with equivalent quasi-norms.

As a corollary of Theorems 3.16 and 4.19, the following conclusion shows that the atomic Hardy space in Definition 4.18 is independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$.

Corollary 4.20. Let $Y(X)$, q , d , η , ω , and θ_0 be as in Theorem 4.19. Then $H_{\text{atom}}^{Y,q,d}(X)$ is independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$ whenever $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$.

Remark 4.21. By the definitions of $Y^{1/\theta_0}(\mathcal{X})$ and $(Y^{1/\theta_0})'(\mathcal{X})$, we know that (2.9) is equivalent to

$$\|\mathcal{M}(f)\|_{[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(\mathcal{X})} \lesssim \|f\|_{[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(\mathcal{X})},$$

where the implicit positive constant is independent of f . Next, we apply Theorem 4.19 to several concrete examples of ball quasi-Banach function spaces on \mathcal{X} , namely, classical Lebesgue spaces, Lorentz spaces, weighted Lebesgue spaces, Orlicz spaces, and variable Lebesgue spaces.

- (i) Let $p \in (\omega/(\omega + \eta), 1]$ and $\beta, \gamma \in (\omega(1/p - 1), \eta)$ with ω as in (2.3) and η as in Definition 3.3. Assume that $Y(\mathcal{X}) := L^p(\mathcal{X})$. Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p]$, $\theta_0 \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_-)$, and $p_0 \in (p, \infty)$. By this and an argument similar to that used in [68, Remark 2.7(a)], we know that

$$[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(\mathcal{X}) = L^{(p/\theta_0)'/(p_0/\theta_0)'}(\mathcal{X})$$

and $(p/\theta_0)'/(p_0/\theta_0)' > 1$. From this, Remark 2.7(i), [14, (3.6)], and [24, Theorem 1.2], we deduce that $L^p(\mathcal{X})$ satisfies all the assumptions of Theorem 4.19. In this case, a different variant of Theorem 4.19 was obtained in [30, Theorem 4.2] which uses the same atoms, but equips a different quasi-norm.

- (ii) Let $r \in (0, \infty)$, $p \in (\omega/(\omega + \eta), 1]$, and $\beta, \gamma \in (\omega(1/p - 1), \eta)$ with ω as in (2.3) and η as in Definition 3.3. Assume that $Y(\mathcal{X}) := L^{p,r}(\mathcal{X})$ is defined as in Remark 2.7(ii). Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p]$, $\theta_0 \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_-)$, and $p_0 \in (p, \infty)$. By this and an argument similar to that used in [68, Remark 2.7(f)], we know that

$$[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(\mathcal{X}) = L^{(p/\theta_0)'/(p_0/\theta_0)', (r/\theta_0)'/(p_0/\theta_0)'}(\mathcal{X})$$

and $(p/\theta_0)'/(p_0/\theta_0)' > 1$. From this, Remark 2.7(ii), [77, Lemma 3.5], and [77, Theorem 7.3], we deduce that $L^{p,r}(\mathcal{X})$ satisfies all the assumptions of Theorem 4.19. Note that the definition of atomic Hardy spaces here is different from [77, Definition 4.2]. Thus, Theorem 4.19 is a different variant of [77, Theorem 4.4] which uses the same atoms, but equips a different quasi-norm.

- (iii) Let $p \in (\omega/(\omega + \eta), 1]$, $r \in (1, \infty)$, $w \in A_r(\mathcal{X})$ satisfy

$$r_w := \inf \{r \in [1, \infty) : w \in A_r(\mathcal{X})\} \in (1, p(\omega + \eta)/\omega),$$

and $\beta, \gamma \in (\omega(r_w/p - 1), \eta)$ with ω as in (2.3) and η as in Definition 3.3. Assume that $Y(\mathcal{X}) := L_w^p(\mathcal{X})$ is defined as in Section 2.4. Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p/r_w]$ and $\theta_0 \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_-)$. By this and (2.11), we obtain $w^{1-(p/\theta_0)'} \in A_{(p/\theta_0)'}(\mathcal{X})$. On one hand, choose $p_0 \in (p, \infty)$ large enough such that

$$w^{1-(p/\theta_0)'} \in A_{(p/\theta_0)'/(p_0/\theta_0)'}(\mathcal{X}). \quad (4.26)$$

On the other hand, by an argument similar to that used in [68, Remark 2.7(b)], we conclude that

$$[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(\mathcal{X}) = L_{w^{1-(p/\theta_0)'}}^{(p/\theta_0)'/(p_0/\theta_0)'}(\mathcal{X}).$$

From this, Remark 2.7(iii), (4.26), [19, Theorem 4.11], and [19, Theorem 6.5(ii)], we deduce that $L_w^p(\mathcal{X})$ satisfies all the assumptions of Theorem 4.19. In this case, a different variant of Theorem 4.19 was obtained in [19, Theorem 5.4] which uses different atoms and equips a different quasi-norm.

- (iv) Let Φ be an Orlicz function as in Remark 2.7(iii) with positive lower type p_Φ^- and positive upper type p_Φ^+ . Recall that the Orlicz space $L^\Phi(X)$ is defined to be the set of all μ -measurable functions f on X such that

$$\|f\|_{L^\Phi(X)} := \inf \left\{ \lambda \in (0, \infty) : \int_X \Phi \left(\frac{|f(x)|}{\lambda} \right) d\mu(x) \leq 1 \right\} < \infty.$$

Let $p_\Phi^- \in (\omega/(\omega + \eta), 1]$, $p_\Phi^+ = 1$, and $\beta, \gamma \in (\omega/(p_\Phi^- - 1), \eta)$ with ω as in (2.3) and η as in Definition 3.3. Assume that $Y(X) := L^\Phi(X)$. Choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_\Phi^-]$, $\theta_0 \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_-)$, and $p_0 \in (1, \infty)$. By this and an argument similar to that used in [68, Remark 2.7(g)], we know that

$$[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(X) = L^{\tilde{\Psi}}(X)$$

and $p_{\tilde{\Psi}}^- = (p_\Phi^+/\theta_0)'/(p_0/\theta_0)' > 1$, where, for any $t \in [0, \infty)$,

$$\tilde{\Psi}(t) := \sup_{r \in [0, \infty)} [rt^{1/(p_0/\theta_0)'} - \Phi(r^{1/s})].$$

By this, Remark 2.7(iii), [19, Theorem 4.11], and [19, Theorem 6.6], we know that $L^\Phi(X)$ satisfies all the assumptions of Theorem 4.19. In this case, a different variant of Theorem 4.19 was obtained in [19, Theorem 5.4] which uses the same atoms, but equips a different quasi-norm.

- (v) Let (X, ρ, μ) be an RD-space, $p(\cdot) \in C^{\log}(X)$ satisfy $\widetilde{p}_- \in (\omega/(\omega + \eta), \infty)$, and $\beta, \gamma \in (\omega[1/\widetilde{p}_- - 1]_+, \eta)$ with ω as in (2.3), η as in Definition 3.3, and $[1/\widetilde{p}_- - 1]_+ := \max\{1/\widetilde{p}_- - 1, 0\}$. Assume that $Y(X) := L^{p(\cdot)}(X)$. If $\widetilde{p}_- \in (\omega/(\omega + \eta), 1)$, we choose $p_- \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, \widetilde{p}_-]$, $\theta_0 \in (\max\{\omega/(\omega + \beta), \omega/(\omega + \gamma)\}, p_-)$, and $p_0 \in (\widetilde{p}_+, \infty)$; if $\widetilde{p}_- \in [1, \infty)$, we choose $p_- \in [1, \widetilde{p}_-]$, $\theta_0 \in (0, 1)$, and $p_0 \in (\widetilde{p}_+, \infty)$. By this and an argument similar to that used in [68, Remark 2.7(f)], we know that

$$[(Y^{1/\theta_0})']^{1/(p_0/\theta_0)'}(X) = L^{(p(\cdot)/\theta_0)'/(p_0/\theta_0)'}(X)$$

and $(p(\cdot)/\theta_0)'/(p_0/\theta_0)' \geq (\widetilde{p}_+/\theta_0)'/(p_0/\theta_0)' > 1$. From this, Remark 2.7(iv), [78, Lemma 2.5], and [78, Theorem 2.7], we deduce that $L^{p(\cdot)}(X)$ satisfies all the assumptions of Theorem 4.19. In this case, Theorem 4.19 improves [78, Theorem 4.3] by removing the reverse doubling assumption of μ .

Let φ be a growth function as in Remark 2.7(iii). Although the Musielak–Orlicz space $L^\varphi(X)$ is also a ball quasi-Banach function space [see Remark 2.7(iii) above], Theorem 4.19 is not applicable to the Musielak–Orlicz Hardy space $H_Y^{*,\varphi}(X)$. Indeed, the space variable x and the growth variable t in $\varphi(x, t)$ are in general not separable, which makes Assumption 2.15 fail on $L^\varphi(X)$. Recently, Fu et al. [19] obtained a corresponding conclusion of Theorem 4.19 in case that $Y(X) := L^\varphi(X)$ independently.

4.2 Finite atomic characterizations

In this subsection, we establish finite atomic characterizations of $H_Y^*(X)$. To this end, we first introduce the notion of finite atomic Hardy spaces associated with $Y(X)$.

Definition 4.22. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(\mathcal{X})$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$ and $d \in (0, \theta_0]$. The *finite atomic Hardy space* $H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ associated with $Y(\mathcal{X})$ is defined to be the set of all finite linear combinations of $(Y(\mathcal{X}), q)$ -atoms. The quasi-norm $\|\cdot\|_{H_{\text{fin}}^{Y,q,d}(\mathcal{X})}$ in $H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ is defined by setting, for any $f \in H_{\text{fin}}^{Y,q,d}(\mathcal{X})$,

$$\|f\|_{H_{\text{fin}}^{Y,q,d}(\mathcal{X})} := \inf \left\{ \left\| \left\{ \sum_{j=1}^m \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})} : m \in \mathbb{N}, \right. \\ \left. f = \sum_{j=1}^m \lambda_j a_j, \{\lambda_j\}_{j=1}^m \subset [0, \infty) \right\},$$

where the infimum is taken over all finite linear combinations of f via $(Y(\mathcal{X}), q)$ -atoms as above.

Next, we establish the finite atomic characterization of $H_Y^*(\mathcal{X})$ (see [72, Theorem 5.12] for the corresponding Euclidean case). In what follows, denote by the symbol $\mathcal{UC}(\mathcal{X})$ the *set of all uniformly continuous functions on \mathcal{X}* , that is, a function $f \in \mathcal{UC}(\mathcal{X})$ if and only if, for any given $\epsilon \in (0, \infty)$, there exists a $\delta \in (0, \infty)$ such that $|f(x) - f(y)| < \epsilon$ whenever $\rho(x, y) < \delta$.

Theorem 4.23. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(\mathcal{X})$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$ and $d \in (0, \theta_0]$.

- (i) If $q \in (\max\{p_0, 1\}, \infty)$, then $\|\cdot\|_{H_{\text{fin}}^{Y,q,d}(\mathcal{X})}$ and $\|\cdot\|_{H_Y^*(\mathcal{X})}$ are equivalent quasi-norms on the space $H_{\text{fin}}^{Y,q,d}(\mathcal{X})$.
- (ii) If $q = \infty$, then $\|\cdot\|_{H_{\text{fin}}^{Y,\infty,d}(\mathcal{X})}$ and $\|\cdot\|_{H_Y^*(\mathcal{X})}$ are equivalent quasi-norms on the space $H_{\text{fin}}^{Y,\infty,d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$.

Proof. Let all the symbols be as in the present theorem. From Theorem 4.19, we deduce that, for any $q \in (\max\{p_0, 1\}, \infty]$,

$$H_{\text{fin}}^{Y,q,d}(\mathcal{X}) \subset H_{\text{atom}}^{Y,q,d}(\mathcal{X}) \subset H_Y^*(\mathcal{X})$$

and, for any $f \in H_{\text{fin}}^{Y,q,d}(\mathcal{X})$,

$$\|f\|_{H_Y^*(\mathcal{X})} \lesssim \|f\|_{H_{\text{atom}}^{Y,q,d}(\mathcal{X})} \lesssim \|f\|_{H_{\text{fin}}^{Y,q,d}(\mathcal{X})}.$$

Thus, to complete the proof of this theorem, it suffices to show that

$$\|f\|_{H_{\text{fin}}^{Y,q,d}(\mathcal{X})} \lesssim \|f\|_{H_Y^*(\mathcal{X})}$$

for any $f \in H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ when $q \in (\max\{p_0, 1\}, \infty)$, or any $f \in H_{\text{fin}}^{Y,\infty,d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$.

Assume that $q \in (\max\{p_0, 1\}, \infty]$. By the homogeneity of both $\|\cdot\|_{H_Y^*(X)}$ and $\|\cdot\|_{H_{\text{fin}}^{Y,q,d}(X)}$, without loss of generality, we may assume that $f \in H_{\text{fin}}^{Y,q,d}(X)$ and $\|f\|_{H_Y^*(X)} = 1$. Since f is a finite linear combination of $(Y(X), q)$ -atoms, it follows that there exist an $x_1 \in X$ and a $K \in (0, \infty)$ such that $\text{supp } f \subset B(x_1, K)$. In the remainder of this proof, let $B_0 := B(x_1, AK)$ with $A := 16A_0^4$ and all the notation be as in the proof of Proposition 4.5. Let $x \notin B_0$ and $\varphi \in \mathcal{G}_0^\eta(\beta, \gamma)$ with $\|\varphi\|_{\mathcal{G}(x,r,\beta,\gamma)} \leq 1$ for some $r \in (0, \infty)$. We consider the following two cases on r .

If $r \geq 4A_0^2\rho(x_1, x)/3$, then, by an argument similar to that used in the proof of [30, (7.1)], we conclude that, for any $y \in B(x, \rho(x_1, x))$,

$$|\langle f, \varphi \rangle| \lesssim f^*(y). \quad (4.27)$$

Let $h \in (0, p)$. From the fact that $\mathbf{1}_{B_0} \leq [C_{(\mu)} 2^{-\omega} \mathcal{M}(\mathbf{1}_{B(x, \rho(x_1, x))})]^{1/h}$ with $C_{(\mu)}$ and ω as in (2.3), Definition 2.4(ii), and Assumption 2.12, we deduce that

$$\|\mathbf{1}_{B_0}\|_{Y(X)} \leq [C_{(\mu)} 2^{-\omega}]^{1/h} \left\| \left[\mathcal{M}(\mathbf{1}_{B(x, \rho(x_1, x))}) \right]^{1/h} \right\|_{Y(X)} \lesssim \|\mathbf{1}_{B(x, \rho(x_1, x))}\|_{Y(X)}.$$

By this and (4.27), we conclude that

$$|\langle f, \varphi \rangle| \lesssim \inf_{y \in B(x, \rho(x, x_1))} f^*(y) \lesssim \|f^*\|_{Y(X)} \|\mathbf{1}_{B(x, \rho(x_1, x))}\|_{Y(X)}^{-1} \lesssim \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1}. \quad (4.28)$$

This is the desired estimate.

If $r < 4A_0^2\rho(x_1, x)/3$, then, by an argument similar to that used in the proof of [30, (7.1)], we conclude that, for any $y \in B(x_1, \rho(x_1, x))$, $|\langle f, \varphi \rangle| \lesssim f^*(y)$. From this, we deduce that

$$|\langle f, \varphi \rangle| \lesssim \inf_{y \in B(x_1, \rho(x, x_1))} f^*(y) \lesssim \inf_{y \in B_0} f^*(y) \lesssim \|f^*\|_{Y(X)} \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1} \sim \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1}. \quad (4.29)$$

This is also the desired estimate.

Combining (4.28), (4.29), the arbitrariness of φ , and the fact that $f^\star \sim f^*$, we know that, for any $x \in B_0^c$,

$$f^\star(x) \leq C f^*(x) \leq c_0 \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1},$$

where c_0 and C are positive constants independent of f and x . Denote by j' the largest integer j such that $2^j < c_0 \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1}$. Then, for any $j \in (j', \infty) \cap \mathbb{Z}$,

$$\Omega_j \subset B_0. \quad (4.30)$$

Let

$$h := \sum_{j=-\infty}^{j'} \sum_{k \in I_j} \lambda_{j,k} a_{j,k} \quad \text{and} \quad \ell := \sum_{j=j'+1}^{\infty} \sum_{k \in I_j} \lambda_{j,k} a_{j,k}, \quad (4.31)$$

where the series converge both in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ and almost everywhere in X . By (4.30), (4.31), and Lemma 4.6(ii), we know that $\text{supp } \ell \subset B_0$, and hence $\text{supp } h \subset B_0$. Moreover, from (4.21) and Lemma 4.6(v), we deduce that

$$\|h\|_{L^\infty(X)} \lesssim \sum_{j=-\infty}^{j'} \left\| \sum_{k \in I_j} \lambda_{j,k} a_{j,k} \right\|_{L^\infty(X)} \lesssim \sum_{j=-\infty}^{j'} 2^j \lesssim 2^{j'} \lesssim \|\mathbf{1}_{B_0}\|_{Y(X)}^{-1}.$$

Notice that $f \in L^{\tilde{q}}(X)$ with $\tilde{q} := q$ if $q < \infty$, and $\tilde{q} := 2$ if $q = \infty$. From this and [30, Theorem 3.4], we deduce that $f^* \in L^{\tilde{q}}(X)$, which, combined with the Hölder inequality, (4.30), (4.21), and an argument similar to that used in the proof of (4.25), further implies that

$$\begin{aligned} \left\| \sum_{j=j'+1}^{\infty} \sum_{k \in I_j} |\lambda_{j,k} a_{j,k}| \right\|_{L^1(X)} &\lesssim \left\| \sum_{j=j'+1}^{\infty} \sum_{k \in I_j} |\lambda_{j,k} a_{j,k}| \right\|_{L^{\tilde{q}}(X)} \lesssim \left\| \sum_{j=j'+1}^{\infty} \sum_{k \in I_j} 2^j \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\|_{L^{\tilde{q}}(X)} \\ &\lesssim \left\| \sum_{j=j'+1}^{\infty} 2^j \mathbf{1}_{\Omega_j} \right\|_{L^{\tilde{q}}(X)} \lesssim \|f^*\|_{L^{\tilde{q}}(X)} < \infty. \end{aligned} \quad (4.32)$$

By this, the Lebesgue dominated convergence theorem, and the cancellation of $a_{j,k}$, we conclude that

$$\int_X \ell(x) d\mu(x) = \sum_{j=j'+1}^{\infty} \sum_{k \in I_j} \int_X \lambda_{j,k}(x) a_{j,k}(x) d\mu(x) = 0, \quad (4.33)$$

which further implies that

$$\int_X h(x) d\mu(x) = 0.$$

Thus, h is a harmless constant multiple of a $(Y(X), \infty)$ -atom supported in B_0 . For ℓ , we consider two cases on q .

Case 1) $q \in (\max\{p_0, 1\}, \infty)$. In this case, $f \in H_{\text{fin}}^{Y,q,d}(X)$. For any $i \in \mathbb{N}$, let

$$F_i := \{(j, k) \in \mathbb{Z} \times \mathbb{N} : j > j', k \in I_j, |j| + k \leq i\},$$

and $\ell_i := \sum_{(j,k) \in F_i} \lambda_{j,k} a_{j,k}$. By (4.32), we know that ℓ converges in $L^q(X)$, which further implies that there exists a positive integer i_0 such that

$$\|\ell - \ell_{i_0}\|_{L^q(X)} \leq \frac{[\mu(B_0)]^{1/q}}{\|\mathbf{1}_{B_0}\|_{Y(X)}}. \quad (4.34)$$

Moreover, from (4.33), we deduce that

$$\int_X [\ell(x) - \ell_{i_0}(x)] d\mu(x) = \int_X \ell(x) d\mu(x) - \int_X \ell_{i_0}(x) d\mu(x) = 0.$$

By this, the fact that $\text{supp}(\ell - \ell_{i_0}) \subset B_0$, and (4.34), we conclude that $\ell - \ell_{i_0}$ is a $(Y(X), q)$ -atom supported in B_0 . Therefore,

$$f = h + \ell = h + (\ell - \ell_{i_0}) + \ell_{i_0}$$

is a finite decomposition of f in terms of $(Y(X), q)$ -atoms. From this, Remark 2.8, (4.21), and (4.25), we deduce that there exists a $\nu \in (0, 1)$ such that

$$\|f\|_{H_{\text{fin}}^{Y,q,d}(X)}^\nu \lesssim \left\| \left\{ \left[\frac{1}{\|\mathbf{1}_{B_0}\|_{Y(X)}} \right]^d \mathbf{1}_{B_0} + \sum_{(j,k) \in F_{i_0}} \left[\frac{\lambda_{j,k}}{\|\mathbf{1}_{B(x_{j,k}, Ar_{j,k})}\|_{Y(X)}} \right]^d \mathbf{1}_{B(x_{j,k}, Ar_{j,k})} \right\}^{\frac{1}{d}} \right\|_{Y(X)}^\nu$$

$$\lesssim 1 + \left\| \left\{ \sum_{j \in \mathbb{Z}} \sum_{k \in I_j} 2^{jd} \mathbf{1}_{B(x_{jk}, Ar_{jk})} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})}^v \lesssim 1 + \|f\|_{H_Y^*(\mathcal{X})}^v \lesssim \|f\|_{H_Y^*(\mathcal{X})}^v,$$

which completes the proof of (i).

Case 2) $q = \infty$. In this case, $f \in H_{\text{fin}}^{Y, \infty, d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$. Assume that $\|f\|_{H_Y^*(\mathcal{X})} = 1$. Since f is bounded, it follows from the fact that $f^\star \sim f^*$, [23, Proposition 3.9], and the boundedness of \mathcal{M} on $L^\infty(\mathcal{X})$ (see, for instance, [14, (3.6)]) that

$$\|f^\star\|_{L^\infty(\mathcal{X})} \sim \|f^*\|_{L^\infty(\mathcal{X})} \lesssim \|\mathcal{M}(f)\|_{L^\infty(\mathcal{X})} \lesssim \|f\|_{L^\infty(\mathcal{X})} < \infty.$$

Thus, there exists a positive integer $j'' > j'$ such that, for any $j \in \{j'' + 1, j'' + 2, \dots\}$, $\mu(\Omega_j) = 0$. Consequently, in this case, we have

$$\ell = \sum_{j=j'+1}^{j''} \sum_{k \in I_j} \lambda_{j,k} a_{j,k}$$

almost everywhere in \mathcal{X} . Let $\epsilon \in (0, \infty)$. By the fact that $f \in \mathcal{UC}(\mathcal{X})$, we know that there exists a $\delta \in (0, \infty)$ such that, for any $x, y \in \mathcal{X}$ with $\rho(x, y) < \delta$, $|f(x) - f(y)| < \epsilon$. Write $\ell = \ell_1^\epsilon + \ell_2^\epsilon$ with $\ell_1^\epsilon := \sum_{(j,k) \in J_1} \lambda_{j,k} a_{j,k}$ and $\ell_2^\epsilon := \sum_{(j,k) \in J_2} \lambda_{j,k} a_{j,k}$, where

$$J_1 := \{(j, k) \in \mathbb{Z} \times \mathbb{N} : j' < j \leq j'', k \in I_j, 12A_0^3 r_{j,k} \geq \delta\}$$

and

$$J_2 := \{(j, k) \in \mathbb{Z} \times \mathbb{N} : j' < j \leq j'', k \in I_j, 12A_0^3 r_{j,k} < \delta\}.$$

On one hand, by (4.30), we know that, for any $j \in (j', j''] \cap \mathbb{Z}$, Ω_j is bounded, which, combined with (i) and Lemma 4.6(vi), further implies that ℓ_1^ϵ is a finite linear combination of $(Y(\mathcal{X}), \infty)$ -atoms, and $\|\ell_1^\epsilon\|_{H_{\text{fin}}^{Y, q, d}(\mathcal{X})} \lesssim \|f\|_{H_Y^*(\mathcal{X})}$. On the other hand, it is easy to see $\text{supp } \ell_2^\epsilon \subset B_0$ and $\int_{\mathcal{X}} \ell_2^\epsilon(x) d\mu(x) = 0$. By an argument similar to that used in the proof of [77, Theorem 5.7(ii)], we obtain $\|\ell_2^\epsilon\|_{L^\infty(\mathcal{X})} \lesssim \epsilon$. Thus, ℓ_2^ϵ is a small constant multiple of a $(Y(\mathcal{X}), \infty)$ -atom. Therefore, $f = h + \ell_1^\epsilon + \ell_2^\epsilon$ is a finite decomposition of f in terms of $(Y(\mathcal{X}), \infty)$ -atoms. Furthermore, using this fact and repeating the proof of (i), we obtain

$$\|f\|_{H_{\text{fin}}^{Y, \infty, d}(\mathcal{X})} \lesssim \|f\|_{H_Y^*(\mathcal{X})}.$$

This finishes the proof of (ii) and hence of Theorem 4.23. \square

Remark 4.24. We now give several applications of Theorem 4.23 as follows.

- (i) Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. If $Y(\mathcal{X}) := L^p(\mathcal{X})$, then, by Remark 4.21(i), we know that $L^p(\mathcal{X})$ satisfies all the assumptions of Theorem 4.23. In this case, Theorem 4.23 is a different variant of [30, Theorem 7.1] which uses the same atoms, but equips a different quasi-norm.
- (ii) Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3, and $r \in (0, \infty)$. If $Y(\mathcal{X}) := L^{p, r}(\mathcal{X})$, then, by Remark 4.21(ii), we know that $L^{p, r}(\mathcal{X})$ satisfies all the assumptions of Theorem 4.23. In this case, Theorem 4.23 is a different variant of [77, Theorem 5.7] which uses the same atoms, but equips a different quasi-norm.

(iii) Let $p \in (\omega/(\omega + \eta), 1]$, $r \in (1, \infty)$, $w \in A_r(X)$ satisfy

$$r_w := \inf \{r \in [1, \infty) : \varphi \in A_r(X)\} \in (1, p(\omega + \eta)/\omega)$$

with ω as in (2.3) and η as in Definition 3.3. If $Y(X) := L_w^p(X)$, then, by Remark 4.21(iii), we know that $L_w^p(X)$ satisfies all the assumptions of Theorem 4.23. In this case, Theorem 4.23 is a different variant of [19, Theorem 7.2] which uses different atoms and equips a different quasi-norm.

(iv) Let Φ be an Orlicz function, $p_\Phi^+ = 1$, and $p_\Phi^- \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. If $Y(X) := L^\Phi(X)$, then, by Remark 4.21(iv), we know that $L^\Phi(X)$ satisfies all the assumptions of Theorem 4.23. In this case, Theorem 4.23 is a different variant of [19, Theorem 7.2] which uses the same atoms, but equips a different quasi-norm.

(v) Let (X, ρ, μ) be an RD-space, and $p(\cdot) \in C^{\log}(X)$ satisfy $\widetilde{p_-} \in (\omega/(\omega + \eta), \infty)$ with ω as in (2.3) and η as in Definition 3.3. If $Y(X) := L^{p(\cdot)}(X)$, then, by Remark 4.21(v), we know that $L^{p(\cdot)}(X)$ satisfies all the assumptions of Theorem 4.23. In this case, Theorem 4.23 improves [78, Theorem 4.24] by removing the reverse doubling assumption of μ .

Let φ be a growth function as in Remark 2.7(iii). As was mentioned in Remark 4.21, Theorem 4.23 is not applicable to $H^{*,\varphi}(X)$. Fu et al. [19] obtained a similar conclusion of Theorem 4.23 in case that $Y(X) := L^\varphi(X)$ independently.

5 Molecular characterizations of $H_Y^*(X)$

In this section, we establish the molecular characterization of $H_Y^*(X)$. To this end, we first introduce the notion of molecules.

Definition 5.1. Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$, $\epsilon \in (0, \infty)$, and $\delta \in (0, 1)$ be as in Lemma 3.2.

(i) A μ -measurable function m on X is called a $(Y(X), q, \epsilon)$ -molecule centered at a ball $B \subset X$ if

(i)₁ for any $j \in \mathbb{Z}_+$,

$$\|m\|_{L^q(U_j(B))} \leq \delta^{j\epsilon} \left[\mu(\delta^{-j}B) \right]^{\frac{1}{q}} \|\mathbf{1}_B\|_{Y(X)}^{-1};$$

here and thereafter, $U_0(B) := B$ and, for any $j \in \mathbb{N}$, $U_j(B) := (\delta^{-j}B) \setminus (\delta^{-j+1}B)$;

(i)₂ $\int_X m(x) d\mu(x) = 0$.

(ii) Let $d \in (0, \theta_0]$ and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. The molecular Hardy space $H_{\text{mol}}^{Y,q,d,\epsilon}(X)$ is defined to be the set of all $f \in (\mathcal{G}_0^\eta(\beta, \gamma))'$ satisfying that there exist a sequence $\{\lambda_j\}_{j \in \mathbb{N}} \subset [0, \infty)$

and a sequence $\{m_j\}_{j \in \mathbb{N}}$ of $(Y(X), q, \epsilon)$ -molecules centered, respectively, at balls $\{B_j\}_{j \in \mathbb{N}}$ of X such that $f = \sum_{j \in \mathbb{N}} \lambda_j m_j$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and

$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} < \infty.$$

Moreover, for any $f \in H_{\text{mol}}^{Y,q,d,\epsilon}(X)$, let

$$\|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)} := \inf \left\{ \left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \right\},$$

where the infimum is taken over all decompositions of f as above.

The following lemma shows that a $(Y(X), q, \epsilon)$ -molecule can be decomposed into a sum of a sequence of $(Y(X), q)$ -atoms.

Lemma 5.2. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$, $\delta \in (0, 1)$ be as in Lemma 3.2, ω as in (2.3), and $\epsilon \in (\omega, \infty)$. If m is a $(Y(X), q, \epsilon)$ -molecule centered at a ball $B := B(x_0, r_0)$ for some $x_0 \in X$ and $r_0 \in (0, \infty)$, then it holds true that*

$$m = \sum_{j=0}^{\infty} u_j b_j + \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} v_{j,k} c_{j,k}$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ with $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$, where, for any $j \in \mathbb{Z}_+$ and $k \in \mathbb{Z} \cap [j, \infty)$,

$$u_j := \frac{2\delta^{\epsilon j} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}} \quad \text{and} \quad v_{j,k} := \frac{2C_{(\mu)} \delta^{(\epsilon-\omega)k + \omega j - \omega} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}}$$

with $C_{(\mu)}$ as in (2.3), and b_j and $c_{j,k}$ are both $(Y(X), q)$ -atoms supported in $\delta^{-j}B$.

Proof. Let m be a $(Y(X), q, \epsilon)$ -molecule centered at some ball $B \subset X$ with $Y(X)$, q , and ϵ as in the present lemma. For any $j \in \mathbb{Z}_+$, let

$$m_j := \frac{\mathbf{1}_{\delta^{-j}B}}{\mu(\delta^{-j}B)} \int_X m(y) \mathbf{1}_{U_j(B)}(y) d\mu(y)$$

and $M_j := m \mathbf{1}_{U_j(B)} - m_j$. Obviously,

$$m = \sum_{j=0}^{\infty} M_j + \sum_{j=0}^{\infty} m_j \tag{5.1}$$

pointwisely.

Now, we consider the first sum of (5.1). Fix $j \in \mathbb{Z}_+$. We claim that M_j is a multiple of a $(Y(X), q)$ -atom. Indeed, it is easy to see that

$$\text{supp } M_j \subset \delta^{-j}B \quad \text{and} \quad \int_X M_j(x) d\mu(x) = 0. \quad (5.2)$$

Moreover, by the Minkowski and the Hölder inequalities, we obtain

$$\begin{aligned} \|M_j\|_{L^q(X)} &\leq \|m\mathbf{1}_{U_j(B)}\|_{L^q(X)} + [\mu(\delta^{-j}B)]^{-\frac{1}{q'}} \|m\mathbf{1}_{U_j(B)}\|_{L^1(X)} \\ &\leq 2 \|m\mathbf{1}_{U_j(B)}\|_{L^q(X)} \leq 2\delta^{\epsilon j} [\mu(\delta^{-j}B)]^{\frac{1}{q}} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &= \frac{2\delta^{\epsilon j} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}} [\mu(\delta^{-j}B)]^{\frac{1}{q}} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}^{-1}. \end{aligned} \quad (5.3)$$

Now, for any $j \in \mathbb{Z}_+$, let

$$u_j := \frac{2\delta^{\epsilon j} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}} \quad \text{and} \quad b_j := \frac{M_j}{u_j}.$$

Then, by (5.2) and (5.3), we know that b_j is a $(Y(X), q)$ -atom supported in $\delta^{-j}B$. From the Minkowski inequality, (5.3), (2.3), and the fact that $\epsilon > \omega$, we deduce that

$$\begin{aligned} \sum_{j=0}^{\infty} \|M_j\|_{L^q(X)} &\lesssim \sum_{j=0}^{\infty} \delta^{\epsilon j} [\mu(\delta^{-j}B)]^{\frac{1}{q}} \|\mathbf{1}_B\|_{Y(X)}^{-1} \lesssim \sum_{j=0}^{\infty} \delta^{(\epsilon-\omega/q)j} [\mu(B)]^{\frac{1}{q}} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &\lesssim [\mu(B)]^{\frac{1}{q}} \|\mathbf{1}_B\|_{Y(X)}^{-1} < \infty. \end{aligned}$$

Thus, $\sum_{j=0}^{\infty} M_j$ converges in $L^q(X)$ and hence in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ with β and γ as in the present lemma.

Next, we consider the second sum of (5.1). For any $j \in \mathbb{Z}_+$, let

$$\mathbf{1}_j := \frac{\mathbf{1}_{\delta^{-j}B}}{\mu(\delta^{-j}B)}, \quad \tilde{m}_j := \int_X m(y) \mathbf{1}_{U_j(B)}(y) d\mu(y), \quad \text{and} \quad N_j := \sum_{k=j}^{\infty} \tilde{m}_k.$$

Then, by the cancellation of m , we obtain

$$N_0 = \sum_{k=0}^{\infty} \tilde{m}_k = \int_X m(y) d\mu(y) = 0,$$

which further implies that

$$\begin{aligned} \sum_{j=0}^{\infty} m_j &= \sum_{j=0}^{\infty} \mathbf{1}_j \tilde{m}_j = \sum_{j=0}^{\infty} \mathbf{1}_j (N_j - N_{j+1}) = \sum_{j=1}^{\infty} (\mathbf{1}_j - \mathbf{1}_{j-1}) N_j \\ &= \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} (\mathbf{1}_j - \mathbf{1}_{j-1}) \tilde{m}_k =: \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} d_{j,k}. \end{aligned} \quad (5.4)$$

Fix $j \in \mathbb{N}$ and $k \in [j, \infty) \cap \mathbb{N}$. We claim that $d_{j,k}$ is a multiple of a $(Y(X), q)$ -atom. Indeed, it is easy to see that

$$\text{supp } d_{j,k} \subset \delta^{-j}B \quad \text{and} \quad \int_X d_{j,k}(x) d\mu(x) = 0. \quad (5.5)$$

Moreover, by the Minkowski inequality, the Hölder inequality, $k \geq j$, and (2.3), we know that

$$\begin{aligned} \|d_{j,k}\|_{L^q(X)} &\leq \frac{2}{[\mu(\delta^{-j+1}B)]^{1/q'}} |\widetilde{m}_k| \leq \frac{2}{[\mu(\delta^{-j+1}B)]^{1/q'}} \|m\mathbf{1}_{U_k(B)}\|_{L^1(X)} \\ &\leq \frac{2[\mu(U_k(B))]^{1/q'}}{[\mu(\delta^{-j+1}B)]^{1/q'}} \|m\mathbf{1}_{U_k(B)}\|_{L^q(X)} \leq \frac{2\delta^{\epsilon k}\mu(\delta^{-k}B)}{[\mu(\delta^{-j+1}B)]^{1/q'}} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &\leq 2C_{(\mu)}\delta^{(\epsilon-\omega)k+\omega j-\omega} [\mu(\delta^{-j+1}B)]^{1/q} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &\leq \frac{2C_{(\mu)}\delta^{(\epsilon-\omega)k+\omega j-\omega} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}} [\mu(\delta^{-j}B)]^{1/q} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}^{-1}, \end{aligned} \quad (5.6)$$

where $C_{(\mu)}$ is as in (2.3). Thus, we find that $d_{j,k}$ is a multiple of a $(Y(X), q)$ -atom. Now, for any $j \in \mathbb{N}$ and $k \in [j, \infty) \cap \mathbb{N}$, define

$$v_{j,k} := \frac{2C_{(\mu)}\delta^{(\epsilon-\omega)k+\omega j-\omega} \|\mathbf{1}_{\delta^{-j}B}\|_{Y(X)}}{\|\mathbf{1}_B\|_{Y(X)}} \quad \text{and} \quad c_{j,k} := \frac{d_{j,k}}{v_{j,k}}.$$

Then, by (5.5) and (5.6), we know that $c_{j,k}$ is a $(Y(X), q)$ -atom supported in $\delta^{-j}B$. From (5.6), (2.3), and the fact that $\epsilon > \omega$, we deduce that

$$\begin{aligned} \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} \|d_{j,k}\|_{L^q(X)} &\lesssim \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} \delta^{(\epsilon-\omega)k+\omega j-\omega} [\mu(\delta^{-j+1}B)]^{1/q} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &\lesssim \sum_{j=1}^{\infty} \delta^{\epsilon j} [\mu(\delta^{-j+1}B)]^{1/q} \|\mathbf{1}_B\|_{Y(X)}^{-1} \lesssim \sum_{j=1}^{\infty} \delta^{(\epsilon-\omega/q)j} [\mu(B)]^{1/q} \|\mathbf{1}_B\|_{Y(X)}^{-1} \\ &\lesssim [\mu(B)]^{1/q} \|\mathbf{1}_B\|_{Y(X)}^{-1} < \infty. \end{aligned}$$

Thus, $\sum_{j=1}^{\infty} \sum_{k=j}^{\infty} d_{j,k}$ converges in $L^q(X)$, and hence in $(\mathcal{G}_0^\eta(\beta, \gamma))'$. By the convergences of both $\sum_{j=0}^{\infty} M_j$ and $\sum_{j=1}^{\infty} \sum_{k=j}^{\infty} d_{j,k}$, (5.1), and (5.4), we conclude that

$$m = \sum_{j=0}^{\infty} M_j + \sum_{j=0}^{\infty} m_j = \sum_{j=0}^{\infty} u_j b_j + \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} v_{j,k} c_{j,k}$$

converges in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, which completes the proof of Lemma 5.2. \square

Now, we establish the molecular characterization of $H_Y^*(X)$ as follows (see [59, Theorem 3.9] for the corresponding Euclidean case).

Theorem 5.3. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that*

$Y(X)$ satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$, $d \in (0, \theta_0]$, $\epsilon \in (\omega/d, \infty)$, and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. Then

$$\left[H_Y^*(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))' \right] = \left[H_{\text{mol}}^{Y,q,d,\epsilon}(X) \cap (\mathcal{G}_0^\eta(\beta, \gamma))' \right]$$

with equivalent quasi-norms.

Proof. Let $Y(X)$, q , and d be as in the present theorem. It is easy to see that, for any given $\epsilon \in (0, \infty)$, any $(Y(X), \infty)$ -atom is a $(Y(X), q, \epsilon)$ -molecule. Thus, by Theorem 4.19, we have

$$H_Y^*(X) \subset H_{\text{atom}}^{Y,\infty,d}(X) \subset H_{\text{mol}}^{Y,q,d,\epsilon}(X)$$

and, for any $f \in H_Y^*(X)$,

$$\|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)} \lesssim \|f\|_{H_{\text{atom}}^{Y,\infty,d}(X)} \lesssim \|f\|_{H_Y^*(X)}.$$

Next, we show that $H_{\text{mol}}^{Y,q,d,\epsilon}(X) \subset H_Y^*(X)$ with $\epsilon \in (\omega/d, \infty)$. By Theorem 4.19, it suffices to prove $H_{\text{mol}}^{Y,q,d,\epsilon}(X) \subset H_{\text{atom}}^{Y,q,d}(X)$. Let $f \in H_{\text{mol}}^{Y,q,d,\epsilon}(X)$. Then, by Definition 5.1, we know that there exist sequences $\{\lambda_i\}_{i \in \mathbb{N}} \subset [0, \infty)$ and $\{m_i\}_{i \in \mathbb{N}}$ of $(Y(X), q, \epsilon)$ -molecules centered, respectively, at balls $\{B_i\}_{i \in \mathbb{N}}$ of X such that $f = \sum_{i \in \mathbb{N}} \lambda_i m_i$ in $(\mathcal{G}_0^\eta(\beta, \gamma))'$ and

$$\left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \lesssim \|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)}. \quad (5.7)$$

From this and Lemma 5.2, we deduce that

$$f = \sum_{i \in \mathbb{N}} \sum_{j=0}^{\infty} \lambda_i u_{i,j} b_{i,j} + \sum_{i \in \mathbb{N}} \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} \lambda_i v_{i,j,k} c_{i,j,k}$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, where, for any $i \in \mathbb{N}$, $j \in \mathbb{Z}_+$, and $k \in \mathbb{Z}_+ \cap [j, \infty)$,

$$u_{i,j} := \frac{2\delta^{\epsilon j} \|\mathbf{1}_{\delta^{-j}B_i}\|_{Y(X)}}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \quad \text{and} \quad v_{i,j,k} := \frac{2C(\mu)\delta^{(\epsilon-\omega)k+\omega j-\omega} \|\mathbf{1}_{\delta^{-j}B_i}\|_{Y(X)}}{\|\mathbf{1}_{B_i}\|_{Y(X)}}$$

with $C(\mu)$ as in (2.3), and both $b_{i,j}$ and $c_{i,j,k}$ are $(Y(X), q)$ -atoms supported in $\delta^{-j}B_i$. Moreover,

$$\begin{aligned} \|f\|_{H_{\text{atom}}^{Y,q,d}(X)} &\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=0}^{\infty} \left[\frac{\lambda_i u_{i,j}}{\|\mathbf{1}_{\delta^{-j}B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{\delta^{-j}B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\ &\quad + \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} \left[\frac{\lambda_i v_{i,j,k}}{\|\mathbf{1}_{\delta^{-j}B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{\delta^{-j}B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\ &=: \mathbf{I}_1 + \mathbf{I}_2. \end{aligned} \quad (5.8)$$

We first estimate I_1 . Choose $h \in (\frac{\omega}{\epsilon d}, 1)$. By Lemma 2.13, (2.3), Assumption 2.12, and (5.7), we know that

$$\begin{aligned}
I_1 &\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=0}^{\infty} \left[\frac{\lambda_i \delta^{\epsilon j}}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{\delta^{-j}B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\
&\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=0}^{\infty} \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \delta^{(\epsilon d - \omega/h)j} [\mathcal{M}(\mathbf{1}_{B_i})]^{\frac{1}{h}} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\
&\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d [\mathcal{M}(\mathbf{1}_{B_i})]^{\frac{1}{h}} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\
&\sim \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d [\mathcal{M}(\mathbf{1}_{B_i})]^{\frac{1}{h}} \right\}^h \right\|_{Y^{\frac{1}{hd}}(X)}^{\frac{1}{hd}} \\
&\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \lesssim \|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)}. \tag{5.9}
\end{aligned}$$

For I_2 , from (5.9), we deduce that

$$\begin{aligned}
I_2 &\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=1}^{\infty} \sum_{k=j}^{\infty} \left[\frac{\lambda_i \delta^{(\epsilon - \omega)k + \omega j - \omega}}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{\delta^{-j}B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \\
&\lesssim \left\| \left\{ \sum_{i \in \mathbb{N}} \sum_{j=1}^{\infty} \left[\frac{\lambda_i \delta^{\epsilon j}}{\|\mathbf{1}_{B_i}\|_{Y(X)}} \right]^d \mathbf{1}_{\delta^{-j}B_i} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \lesssim \|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)}. \tag{5.10}
\end{aligned}$$

Combining (5.8), (5.9), and (5.10), we obtain $\|f\|_{H_{\text{atom}}^{Y,q,d,\epsilon}(X)} \lesssim \|f\|_{H_{\text{mol}}^{Y,q,d,\epsilon}(X)}$, which completes the proof of $H_{\text{mol}}^{Y,q,d,\epsilon}(X) \subset H_Y^*(X)$ and hence of Theorem 5.3. \square

As a corollary of Theorems 3.16 and 5.3, the following conclusion shows that the molecular Hardy space in Definition 5.1(ii) is independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$.

Corollary 5.4. *Let $Y(X)$, q , d , ϵ , η , ω , and θ_0 be as in Theorem 5.3. Then $H_{\text{mol}}^{Y,q,d,\epsilon}(X)$ is independent of the choices of $(\mathcal{G}_0^\eta(\beta, \gamma))'$ whenever $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$.*

Remark 5.5. We give several applications of Theorem 5.3 as follows.

- (i) Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. If $Y(X) := L^p(X)$, then, by Remark 4.21(i), we know that $L^p(X)$ satisfies all the assumptions of Theorem 5.3. In this case, Theorem 5.3 is a different variant of [31, Proposition 5.5] which uses the different molecules and equips a different quasi-norm.

- (ii) Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3, and $r \in (0, \infty)$. If $Y(\mathcal{X}) := L^{p,r}(\mathcal{X})$, then, by Remark 4.21(ii), we know that $L^{p,r}(\mathcal{X})$ satisfies all the assumptions of Theorem 5.3. In this case, Theorem 5.3 is a different variant of [77, Theorem 6.4] which uses the same molecules, but equips a different quasi-norm.
- (iii) Let $p \in (\omega/(\omega + \eta), 1]$, $r \in (1, \infty)$, $w \in A_r(\mathcal{X})$ satisfy

$$r_w := \inf \{r \in [1, \infty) : \varphi \in A_r(\mathcal{X})\} \in (1, p(\omega + \eta)/\omega)$$

with ω as in (2.3) and η as in Definition 3.3. If $Y(\mathcal{X}) := L_w^p(\mathcal{X})$, then, by Remark 4.21(iii), we know that $L_w^p(\mathcal{X})$ satisfies all the assumptions of Theorem 5.3. In this case, Theorem 5.3 is a different variant of [19, Theorem 6.8] which uses same molecules, but equips a different quasi-norm.

- (iv) Let Φ be an Orlicz function, $p_\Phi^+ = 1$, and $p_\Phi^- \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. If $Y(\mathcal{X}) := L^\Phi(\mathcal{X})$, then, by Remark 4.21(iv), we know that $L^\Phi(\mathcal{X})$ satisfies all the assumptions of Theorem 5.3. In this case, Theorem 5.3 is a different variant of [19, Theorem 6.8] which uses the same molecules, but equips a different quasi-norm.
- (v) Let (\mathcal{X}, ρ, μ) be an RD-space, $p(\cdot) \in C^{\log}(\mathcal{X})$ satisfy $\widetilde{p}^- \in (\omega/(\omega + \eta), \infty)$ with ω as in (2.3) and η as in Definition 3.3. If $Y(\mathcal{X}) := L^{p(\cdot)}(\mathcal{X})$, then, by Remark 4.21(v), we know that $L^{p(\cdot)}(\mathcal{X})$ satisfies all the assumptions of Theorem 5.3. In this case, Theorem 5.3 is new.

Let φ be a growth function as in Remark 2.7(iii). As was mentioned in Remark 4.21, Theorem 5.3 is not applicable to $H^{*,\varphi}(\mathcal{X})$. Fu et al. [19] obtained a similar conclusion of Theorem 5.3 in case that $Y(\mathcal{X}) := L^\varphi(\mathcal{X})$ independently.

6 Dual space of $H_Y^*(\mathcal{X})$

In this section, we give the dual space of $H_Y^*(\mathcal{X})$. To this end, we first introduce the notion of absolutely continuous quasi-norms as follows (see, for instance, [5, Chapter 1, Definition 3.1] and [68, Definition 3.2] for the corresponding Euclidean case).

Definition 6.1. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} . A function $f \in Y(\mathcal{X})$ is said to have an *absolutely continuous quasi-norm* in $Y(\mathcal{X})$ if $\|f\mathbf{1}_{E_j}\|_{Y(\mathcal{X})} \downarrow 0$ whenever $\{E_j\}_{j \in \mathbb{N}}$ is a sequence of μ -measurable sets satisfying $E_j \supset E_{j+1}$ for any $j \in \mathbb{N}$, and $\bigcap_{j \in \mathbb{N}} E_j = \emptyset$. Moreover, $Y(\mathcal{X})$ is said to have an *absolutely continuous quasi-norm* if, for any $f \in Y(\mathcal{X})$, f has an absolutely continuous quasi-norm in $Y(\mathcal{X})$.

Remark 6.2. (i) Assume that $Y(\mathcal{X})$ is a ball quasi-Banach function space on \mathcal{X} and has an absolutely continuous quasi-norm. Let $p \in (0, \infty)$. Then, by Definition 2.10 and Remark 2.11, we know that $Y^p(\mathcal{X})$ also has an absolutely continuous quasi-norm.

- (ii) Observe that, in Definition 6.1, if we replace $\bigcap_{j \in \mathbb{N}} E_j = \emptyset$ by $\mu(\bigcap_{j \in \mathbb{N}} E_j) = 0$, we obtain its another equivalent formulation.

We prove the following dominated convergence theorem on $Y(\mathcal{X})$ based on the assumption that $Y(\mathcal{X})$ has an absolutely continuous quasi-norm. For the case of Banach function spaces on \mathbb{R}^n , this conclusion is a simple corollary of [5, Chapter 1, Proposition 3.6].

Lemma 6.3. *Assume that $Y(X)$ is a ball quasi-Banach function space on X and has an absolutely continuous quasi-norm. Let $g \in Y(X)$ and $\{f_m\}_{m \in \mathbb{N}}$ be a sequence of μ -measurable functions satisfying that $|f_m| \leq |g|$ for any $m \in \mathbb{N}$, and $\lim_{m \rightarrow \infty} f_m = f$ almost everywhere in X . Then*

$$\lim_{m \rightarrow \infty} \|f_m - f\|_{Y(X)} = 0.$$

Proof. Without loss of generality, we may assume that $\lim_{m \rightarrow \infty} f_m = f$ pointwisely. First, it is easy to see that, for any $x \in X$,

$$|f(x)| \leq |g(x)|. \quad (6.1)$$

Now, we prove this lemma in case that $\text{supp } f \subset B$, where B is a ball of X . For any $n, j \in \mathbb{N}$, define

$$E_{n,j} := \{x \in B : |f_m(x) - f(x)| < 2^{-j}, \forall m \geq n\}.$$

It is easy to see that, for any given $j \in \mathbb{N}$,

$$\bigcup_{n \in \mathbb{N}} E_{n,j} = B,$$

$E_{n,j} \subset E_{n+1,j}$ for any $n \in \mathbb{N}$, and

$$\lim_{n \rightarrow \infty} \mu(E_{n,j}) = \mu(B).$$

Therefore, for any $j \in \mathbb{N}$, there exists an $n_j \in \mathbb{N}$ such that

$$\mu(B \setminus E_{n_j,j}) < 2^{-j}. \quad (6.2)$$

For any $J \in \mathbb{N}$, define

$$A_J := \bigcap_{j=J}^{\infty} E_{n_j,j} \quad \text{and} \quad B_J := B \setminus A_J.$$

Then, for any $J \in \mathbb{N}$,

$$B_{J+1} = B \setminus A_{J+1} = \bigcup_{j=J+1}^{\infty} (B \setminus E_{n_j,j}) \subset \bigcup_{j=J}^{\infty} (B \setminus E_{n_j,j}) = B \setminus A_J = B_J; \quad (6.3)$$

moreover, by (6.2), we know that, for any $J \in \mathbb{N}$,

$$\mu(B_J) = \mu\left(\bigcup_{j=J}^{\infty} (B \setminus E_{n_j,j})\right) \leq \sum_{j=J}^{\infty} \mu(B \setminus E_{n_j,j}) < \sum_{j=J}^{\infty} 2^{-j} \lesssim 2^{-J},$$

which, combined with (6.3), further implies that

$$\mu\left(\bigcap_{J \in \mathbb{N}} B_J\right) = 0. \quad (6.4)$$

On one hand, from (6.3), (6.4), and Remark 6.2(ii), we deduce that, for any $\epsilon \in (0, \infty)$, there exists a $J_1 \in \mathbb{N}$ such that

$$\|g \mathbf{1}_{B_{J_1}}\|_{Y(X)} < \frac{\epsilon}{2}. \quad (6.5)$$

On the other hand, it is easy to see that, for any $\epsilon \in (0, \infty)$, there exists a $J_2 \in (J_1, \infty) \cap \mathbb{N}$ such that

$$2^{-J_2} < \frac{\epsilon}{2\|\mathbf{1}_B\|_{Y(X)}}, \quad (6.6)$$

which further implies that there exists an $n_{J_2} \in \mathbb{N}$ such that, for any $x \in A_{J_2} \subset E_{n_{J_2}, J_2}$ and any $m > n_{J_2}$,

$$|f_m(x) - f(x)| < 2^{-J_2}. \quad (6.7)$$

Combining Definition 2.4(ii), (6.7), (6.1), (6.6), (6.3), and (6.5), we conclude that, for any $m > n_{J_2}$,

$$\begin{aligned} \|f_m - f\|_{Y(X)} &\lesssim \|(f_m - f)\mathbf{1}_{A_{J_2}}\|_{Y(X)} + \|(f_m - f)\mathbf{1}_{B_{J_2}}\|_{Y(X)} \\ &\lesssim 2^{-J_2} \|\mathbf{1}_B\|_{Y(X)} + \|g\mathbf{1}_{B_{J_2}}\|_{Y(X)} \\ &\lesssim \frac{\epsilon}{2} + \|g\mathbf{1}_{B_{J_1}}\|_{Y(X)} \lesssim \epsilon. \end{aligned}$$

Thus,

$$\lim_{m \rightarrow \infty} \|f_m - f\|_{Y(X)} = 0. \quad (6.8)$$

Next, we prove this lemma for any $f \in Y(X)$. Let $\{B_n\}_{n \in \mathbb{N}}$ be a sequence of balls satisfying that $X = \bigcup_{n \in \mathbb{N}} B_n$ and, for any $n \in \mathbb{N}$, $B_n \subset B_{n+1}$. On one hand, by Definition 6.1, we know that, for any $\epsilon \in (0, \infty)$, there exists an $N_1 \in \mathbb{N}$ such that

$$\left\| g(1 - \mathbf{1}_{B_{N_1}}) \right\|_{Y(X)} < \frac{\epsilon}{2}. \quad (6.9)$$

On the other hand, from (6.8), we deduce that there exists an $N_2 \in \mathbb{N}$ such that, for any $m > N_2$,

$$\|(f_m - f)\mathbf{1}_{B_{N_1}}\|_{Y(X)} < \frac{\epsilon}{2},$$

which, combined with Definition 2.4(ii), (6.1), and (6.9), further implies that, for any $m > N_2$,

$$\begin{aligned} \|f_m - f\|_{Y(X)} &\lesssim \|f_m(1 - \mathbf{1}_{B_{N_1}})\|_{Y(X)} + \|(f_m - f)\mathbf{1}_{B_{N_1}}\|_{Y(X)} + \|f(\mathbf{1}_{B_{N_1}} - 1)\|_{Y(X)} \\ &\lesssim \|g(1 - \mathbf{1}_{B_{N_1}})\|_{Y(X)} + \|(f_m - f)\mathbf{1}_{B_{N_1}}\|_{Y(X)} \lesssim \frac{\epsilon}{2} + \frac{\epsilon}{2} \sim \epsilon. \end{aligned}$$

This shows that $\lim_{m \rightarrow \infty} \|f_m - f\|_{Y(X)} = 0$ and hence finishes the proof of Lemma 6.3. \square

If we assume that $Y(X)$ has an absolutely continuous quasi-norm, then we have the following density conclusions of $H_Y^*(X)$.

Lemma 6.4. *Let $Y(X)$ be a ball quasi-Banach function space on X satisfying Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. Further assume that $Y(X)$ has an absolutely continuous quasi-norm and satisfies Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$ and $d \in (0, \theta_0]$. Then*

- (i) $H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ is dense in $H_Y^*(\mathcal{X})$;
- (ii) $H_{\text{fin}}^{Y,\infty,d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$ is dense in $H_Y^*(\mathcal{X})$.

Proof. Let all the symbols be as in the present lemma. We first prove (i). Let $f \in H_Y^*(\mathcal{X})$ and $\beta, \gamma \in (\omega(1/\theta_0 - 1), \eta)$. Then, by Theorem 4.19, we know that there exist a sequence $\{\lambda_j\}_{j \in \mathbb{N}}$ of non-negative numbers and a sequence $\{a_j\}_{j \in \mathbb{N}}$ of $(Y(\mathcal{X}), q)$ -atoms supported, respectively, in balls $\{B_j\}_{j \in \mathbb{N}}$ of \mathcal{X} such that

$$f = \sum_{j \in \mathbb{N}} \lambda_j a_j$$

in $(\mathcal{G}_0^\eta(\beta, \gamma))'$, and

$$\left\| \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\} \right\|_{Y(\mathcal{X})}^{\frac{1}{d}} \sim \|f\|_{H_Y^*(\mathcal{X})} < \infty.$$

Let

$$F := \left\{ \sum_{j \in \mathbb{N}} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}}$$

and, for any $N \in \mathbb{N}$, let

$$f_N := \sum_{j=1}^N \lambda_j a_j \quad \text{and} \quad F_N := \left\{ \sum_{j=1}^N \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}}.$$

Then $F^d \in Y^{\frac{1}{d}}(\mathcal{X})$, $f_N \in H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ and $F_N^d \leq F^d$ for any $N \in \mathbb{N}$, and $\lim_{N \rightarrow \infty} F_N^d = F^d$ almost everywhere in \mathcal{X} . From this, Theorem 4.19, Remark 6.2(i), and Lemma 6.3, we deduce that

$$\begin{aligned} \|f - f_N\|_{H_Y^*(\mathcal{X})} &\sim \|f - f_N\|_{H_{\text{atom}}^{Y,q,d}(\mathcal{X})} \lesssim \left\| \left\{ \sum_{j=N+1}^{\infty} \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_j} \right\} \right\|_{Y(\mathcal{X})}^{\frac{1}{d}} \\ &\sim \left\| (F^d - F_N^d)^{\frac{1}{d}} \right\|_{Y(\mathcal{X})} \sim \|F^d - F_N^d\|_{Y^{\frac{1}{d}}(\mathcal{X})}^{\frac{1}{d}} \rightarrow 0 \end{aligned}$$

as $N \rightarrow \infty$, which further implies (i). Thus, to prove the present lemma, it remains to show (ii). By (i), we only need to show that $H_{\text{fin}}^{Y,\infty,d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$ is dense in $H_Y^*(\mathcal{X})$ with the quasi-norm $\|\cdot\|_{H_Y^*(\mathcal{X})}$. To this end, let a be a $(Y(\mathcal{X}), \infty)$ -atom supported in a ball $B := B(x_B, r_B) \subset \mathcal{X}$ with $x_B \in \mathcal{X}$ and $r_B \in (0, \infty)$, and $k_0 \in \mathbb{Z}$ sufficiently large. Then, by the proof of [77, Lemma 5.3(ii)] (with the same notation as therein), we know that, for any $k > k_0$, $S_k a \in \mathcal{UC}(\mathcal{X})$, $S_k a$ is a $(Y(\mathcal{X}), \infty)$ -atom supported in $B(x_B, 2A_0 r_B)$, and

$$\lim_{k \rightarrow \infty} \|S_k a - a\|_{L^2(\mathcal{X})} = 0. \quad (6.10)$$

Thus,

$$\widetilde{a} := \frac{(S_k a - a)[\mu(B(x_B, 2A_0 r_B))]^{1/2}}{\|S_k a - a\|_{L^2(\mathcal{X})} \|\mathbf{1}_{B(x_B, 2A_0 r_B)}\|_{Y(\mathcal{X})}}$$

is a $(Y(\mathcal{X}), 2)$ -atom. From this, Theorem 4.19, and (6.10), we deduce that

$$\begin{aligned} \lim_{k \rightarrow \infty} \|S_k a - a\|_{H_Y^*(\mathcal{X})} &\sim \lim_{k \rightarrow \infty} \|S_k a - a\|_{H_{\text{atom}}^{Y,2,d}(\mathcal{X})} \\ &\sim \frac{\|\mathbf{1}_{B(x_B, 2A_0 r_B)}\|_{Y(\mathcal{X})}}{[\mu(B(x_B, 2A_0 r_B))]^{1/2}} \lim_{k \rightarrow \infty} \|S_k a - a\|_{L^2(\mathcal{X})} = 0. \end{aligned}$$

This finishes the proof of (iii) and hence of Lemma 6.4. \square

Let $q \in (0, \infty)$. Recall that the space $L_{\mathbb{B}}^q(\mathcal{X})$ is defined to be the set of all μ -measurable functions f on \mathcal{X} such that, for any ball $B \subset \mathcal{X}$,

$$\int_B |f(x)|^q d\mu(x) < \infty.$$

Next, we introduce the notion of ball Campanato-type function spaces associated with $Y(\mathcal{X})$ (see [39, Definition 2.4] and [75, Definition 3.2], respectively, for the cases of anisotropic mixed-norm Lebesgue spaces and ball quasi-Banach function spaces on \mathbb{R}^n).

Definition 6.5. Let $Y(\mathcal{X})$ be a ball quasi-Banach function space on \mathcal{X} , $q \in [1, \infty)$, and $d \in (0, \infty)$. Then the ball Campanato-type function space $\mathcal{L}_{Y,q,d}(\mathcal{X})$, associated with $Y(\mathcal{X})$, is defined to be the set of all $f \in L_{\mathbb{B}}^q(\mathcal{X})$ such that

$$\begin{aligned} \|f\|_{\mathcal{L}_{Y,q,d}(\mathcal{X})} &:= \sup \left\| \left\{ \sum_{i=1}^m \left[\frac{\lambda_i}{\|\mathbf{1}_{B_i}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_i} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})}^{-1} \\ &\quad \times \sum_{j=1}^m \left\{ \frac{\lambda_j \mu(B_j)}{\|\mathbf{1}_{B_j}\|_{Y(\mathcal{X})}} \left[\frac{1}{\mu(B_j)} \int_{B_j} |f(x) - m_{B_j}(f)|^q d\mu(x) \right]^{\frac{1}{q}} \right\} < \infty, \end{aligned}$$

where, for any ball $B \subset \mathcal{X}$,

$$m_B(f) := \frac{1}{\mu(B)} \int_B f(x) d\mu(x),$$

and the supremum is taken over all $m \in \mathbb{N}$, balls $\{B_j\}_{j=1}^m$ of \mathcal{X} , and $\{\lambda_j\}_{j=1}^m \subset [0, \infty)$ with $\sum_{j=1}^m \lambda_j \neq 0$.

Now, we give the dual space of $H_Y^*(\mathcal{X})$, which is the main result of this section.

Theorem 6.6. Assume that $Y(\mathcal{X})$ is a ball quasi-Banach function space on \mathcal{X} having an absolutely continuous quasi-norm. Further assume that $Y(\mathcal{X})$ satisfies Assumption 2.12 with $p_- \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3, and Assumption 2.15 with the same p_- as in Assumption 2.12, $\theta_0 \in (\omega/(\omega + \eta), \underline{p})$, and $p_0 \in (\theta_0, \infty)$, where \underline{p} is as in (2.8). Let $q \in (\max\{p_0, 1\}, \infty]$ and $d \in (0, \theta_0]$. Then the dual space of $H_Y^*(\mathcal{X})$, denoted by $(H_Y^*(\mathcal{X}))^*$, is $\mathcal{L}_{Y,q',d}(\mathcal{X})$ with $1/q + 1/q' = 1$ in the following sense:

(i) if $f \in \mathcal{L}_{Y,q',d}(\mathcal{X})$, then the linear functional

$$T_f : g \mapsto T_f(g) := \int_{\mathcal{X}} f(x)g(x) d\mu(x), \quad (6.11)$$

initially defined for any $g \in H_{\text{fin}}^{Y,q,d}(\mathcal{X})$, has a bounded extension to $H_Y^*(\mathcal{X})$;

- (ii) conversely, any continuous linear functional on $H_Y^*(X)$ arises as in (6.11) with a unique $f \in \mathcal{L}_{Y,q',d}(X)$.

Moreover, $\|f\|_{\mathcal{L}_{Y,q',d}(X)} \sim \|T_f\|_{(H_Y^*(X))^*}$, where the positive equivalence constants are independent of f .

Proof. Let all the symbols be as in the present theorem. We first prove (i) in the case $q \in (\max\{p_0, 1\}, \infty)$. Let $f \in \mathcal{L}_{Y,q',d}(X)$ and $g \in H_{\text{fin}}^{Y,q,d}(X)$. By Definition 4.22, we know that there exist an $N \in \mathbb{N}$, a sequence $\{\lambda_j\}_{j=1}^N \subset [0, \infty)$, and a sequence $\{a_j\}_{j=1}^N$ of $(Y(X), q)$ -atoms supported, respectively, in balls $\{B_j\}_{j=1}^N$ such that $g = \sum_{j=1}^N \lambda_j a_j$ almost everywhere in X and

$$\left\| \left\{ \sum_{j=1}^N \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \lesssim \|g\|_{H_{\text{fin}}^{Y,q,d}(X)}.$$

By this, Definition 4.1, the Hölder inequality, and Theorem 4.23(i), we obtain

$$\begin{aligned} |T_f(g)| &= \left| \int_X f(x)g(x) d\mu(x) \right| \leq \sum_{j=1}^N \lambda_j \left| \int_{B_j} f(x)a_j(x) d\mu(x) \right| \\ &\leq \sum_{j=1}^N \lambda_j \int_{B_j} |f(x) - m_{B_j}(f)| |a_j(x)| d\mu(x) \\ &\leq \sum_{j=1}^N \lambda_j \|a_j\|_{L^q(X)} \left\{ \int_{B_j} |f(x) - m_{B_j}(f)|^{q'} d\mu(x) \right\}^{1/q'} \\ &\leq \sum_{j=1}^N \frac{\lambda_j \mu(B_j)}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \left[\frac{1}{\mu(B_j)} \int_{B_j} |f(x) - m_{B_j}(f)|^{q'} d\mu(x) \right]^{1/q'} \\ &\leq \left\| \left\{ \sum_{j=1}^N \left[\frac{\lambda_j}{\|\mathbf{1}_{B_j}\|_{Y(X)}} \right]^d \mathbf{1}_{B_j} \right\}^{\frac{1}{d}} \right\|_{Y(X)} \|f\|_{\mathcal{L}_{Y,q',d}(X)} \\ &\lesssim \|g\|_{H_{\text{fin}}^{Y,q,d}(X)} \|f\|_{\mathcal{L}_{Y,q',d}(X)} \sim \|g\|_{H_Y^*(X)} \|f\|_{\mathcal{L}_{Y,q',d}(X)}. \end{aligned}$$

From this, Lemma 6.4(ii), and a standard density argument, we deduce that T_f can uniquely be extended to a bounded linear functional on $H_Y^*(X)$ and

$$\|T_f\|_{(H_Y^*(X))^*} \lesssim \|f\|_{\mathcal{L}_{Y,q',d}(X)},$$

which then completes the proof of (i) in the case $q \in (\max\{p_0, 1\}, \infty)$.

Next, we prove (i) in the case $q = \infty$. Indeed, using Theorem 4.23(ii) and Lemma 6.4(iii), and repeating the previous proof, we conclude that any $f \in \mathcal{L}_{Y,1,d}(X)$ induces a bounded linear functional on $H_Y^*(X)$, which is initially defined on $H_{\text{fin}}^{Y,\infty,d}(X) \cap \mathcal{UC}(X)$ given by setting, for any $g \in H_{\text{fin}}^{Y,\infty,d}(X) \cap \mathcal{UC}(X)$,

$$T_f : g \mapsto T_f(g) := \int_X f(x)g(x) d\mu(x), \quad (6.12)$$

and then has a bounded linear extension to $H_Y^*(\mathcal{X})$. Let $f \in \mathcal{L}_{Y,1,d}(\mathcal{X})$. Thus, it remains to prove that, for any $g \in H_{\text{fin}}^{Y,\infty,d}(\mathcal{X})$,

$$T_f(g) = \int_{\mathcal{X}} f(x)g(x) d\mu(x). \quad (6.13)$$

Indeed, suppose $g \in H_{\text{fin}}^{Y,\infty,d}(\mathcal{X})$ and $\text{supp } g \subset B(x_0, R)$ for some $x_0 \in \mathcal{X}$ and $R \in (0, \infty)$. By the proof of Lemma 6.4(iii), we know that there exists a sequence $\{S_k\}_{k \in \mathbb{N}}$ of bounded operators on $L^2(\mathcal{X})$ such that $S_k g \in H_{\text{fin}}^{Y,\infty,d}(\mathcal{X}) \cap \mathcal{UC}(\mathcal{X})$, $\lim_{k \rightarrow \infty} S_k g(x) = g(x)$ for almost every $x \in \mathcal{X}$, and

$$\lim_{k \rightarrow \infty} \|S_k g - g\|_{H_Y^*(\mathcal{X})} = 0.$$

From this, (6.12), the fact that $|f S_k g| \lesssim \|g\|_{L^\infty(\mathcal{X})} \mathbf{1}_{B(x_0, 2A_0 R)} |f|$ for any $k \in \mathbb{N}$, and the Lebesgue dominated convergence theorem, we deduce that

$$T_f(g) = \lim_{k \rightarrow \infty} T_f(S_k g) = \lim_{k \rightarrow \infty} \int_{\mathcal{X}} f(x) S_k g(x) d\mu(x) = \int_{\mathcal{X}} f(x) g(x) d\mu(x),$$

which shows that (6.13) holds true and hence completes the proof of (i).

Next, we prove (ii). Let $T \in (H_Y^*(\mathcal{X}))^* = (H_{\text{atom}}^{Y,q,d}(\mathcal{X}))^*$. For any given $q \in (\max\{p_0, 1\}, \infty]$ and any ball $B \subset \mathcal{X}$, let $L^q(B)$ be the set of all $L^q(\mathcal{X})$ functions vanishing outside B , and

$$L_0^q(B) := \left\{ g \in L^q(B) : \int_{\mathcal{X}} g(x) d\mu(x) = 0 \right\}.$$

Then $L_0^q(B) \subset H_{\text{atom}}^{Y,q,d}(\mathcal{X})$ and, for any $g \in L_0^q(B)$ with $\|g\|_{L^q(B)} \neq 0$, $a := \frac{[\mu(B)]^{1/q}}{\|g\|_{L^q(\mathcal{X})} \|\mathbf{1}_B\|_{Y(\mathcal{X})}} g$ is a $(Y(\mathcal{X}), q)$ -atom; moreover,

$$|T(g)| = \frac{\|g\|_{L^q(\mathcal{X})} \|\mathbf{1}_B\|_{Y(\mathcal{X})}}{[\mu(B)]^{1/q}} |T(a)| \leq \frac{\|g\|_{L^q(\mathcal{X})} \|\mathbf{1}_B\|_{Y(\mathcal{X})}}{[\mu(B)]^{1/q}} \|T\|_{(H_{\text{atom}}^{Y,q,d}(\mathcal{X}))^*},$$

which implies that

$$\|T\|_{(L_0^q(B))^*} \leq \frac{\|\mathbf{1}_B\|_{Y(\mathcal{X})} \|T\|_{(H_{\text{atom}}^{Y,q,d}(\mathcal{X}))^*}}{[\mu(B)]^{1/q}}$$

and hence T is a bounded linear functional on $L_0^q(B)$. From this and the Hahn–Banach theorem, we deduce that T can be extended to the whole space $L^q(B)$ without increasing its norm.

If $q \in (\max\{p_0, 1\}, \infty)$, by the duality $[L^q(B)]^* = L^{q'}(B)$ with $1/q + 1/q' = 1$, we know that there exists an $F \in L^{q'}(B)$ such that, for any $g \in L_0^q(B)$,

$$T(g) = \int_{\mathcal{X}} F(x)g(x) d\mu(x)$$

and

$$\|F\|_{L^{q'}(B)} = \|T\|_{(L^q(B))^*} \leq \frac{\|\mathbf{1}_B\|_{Y(\mathcal{X})} \|T\|_{(H_{\text{atom}}^{Y,q,d}(\mathcal{X}))^*}}{[\mu(B)]^{1/q}}.$$

For the case $q = \infty$, let $\tilde{q} \in (\max\{p_0, 1\}, \infty)$. By Theorem 4.19, we know that $T \in (H_{\text{atom}}^{Y, \infty, d}(\mathcal{X}))^*$ implies $T \in (H_{\text{atom}}^{Y, \tilde{q}, d}(\mathcal{X}))^*$ with equivalent norms. Thus, there exists an $F \in L^{\tilde{q}}(B) \subset L^1(B)$ such that, for any $g \in L_0^\infty(B)$, $T(g) = \int_{\mathcal{X}} F(x)g(x) d\mu(x)$ and

$$\|F\|_{L^1(B)} \leq \|F\|_{L^{\tilde{q}}(B)} [\mu(B)]^{1/\tilde{q}} \leq \frac{\|\mathbf{1}_B\|_{Y(\mathcal{X})} \|T\|_{(H_{\text{atom}}^{Y, \tilde{q}, d}(\mathcal{X}))^*}}{[\mu(B)]^{1/\tilde{q}}} [\mu(B)]^{1/\tilde{q}} \lesssim \|\mathbf{1}_B\|_{Y(\mathcal{X})} \|T\|_{(H_{\text{atom}}^{Y, \infty, d}(\mathcal{X}))^*}.$$

Altogether, we find that, for any given $q \in (\max\{1, p_0\}, \infty]$, there exists an $F \in L^{q'}(B)$ such that, for any $g \in L_0^q(B)$,

$$T(g) = \int_{\mathcal{X}} F(x)g(x) d\mu(x) \quad (6.14)$$

and

$$\|F\|_{L^{q'}(B)} \lesssim \frac{\|\mathbf{1}_B\|_{Y(\mathcal{X})} \|T\|_{(H_{\text{atom}}^{Y, q, d}(\mathcal{X}))^*}}{[\mu(B)]^{1/q}}.$$

Next, we claim that, if there exists another function $\tilde{F} \in L^{q'}(B)$ such that, for any $g \in L_0^q(B)$, $T(g) = \int_B \tilde{F}(x)g(x) d\mu(x)$, then $\tilde{F} - F = m_B(\tilde{F} - F)$ almost everywhere in B . Indeed, for any $g \in L_0^q(B)$,

$$\int_B [\tilde{F}(x) - F(x)] g(x) d\mu(x) = 0,$$

which, together with the fact that $h - m_B(h) \in L_0^q(B)$ for any $h \in L^q(B)$, further implies that, for any $h \in L^q(B)$,

$$\int_B [\tilde{F}(x) - F(x)] [h(x) - m_B(h)] d\mu(x) = 0. \quad (6.15)$$

Moreover, for any $H \in L^q(\mathcal{X})$ and $G \in L^{q'}(\mathcal{X})$, by the definitions of $m_B(H)$ and $m_B(G)$, we obtain

$$\begin{aligned} & \int_B [m_B(H)G(x) - m_B(G)H(x)] d\mu(x) \\ &= \int_B \{m_B(H)[G(x) - m_B(G)] + m_B(G)[m_B(H) - H(x)]\} d\mu(x) = 0, \end{aligned} \quad (6.16)$$

which, together with $h \in L^q(B)$ and $F, \tilde{F} \in L^{q'}(B)$, further implies that

$$\int_B m_B(h) [\tilde{F}(x) - F(x)] d\mu(x) = \int_B h(x) m_B(\tilde{F} - F) d\mu(x). \quad (6.17)$$

Combining (6.15) and (6.17), we conclude that, for any $h \in L^q(B)$,

$$\int_B h(x) [\tilde{F}(x) - F(x) - m_B(\tilde{F} - F)] d\mu(x) = 0,$$

which implies that, for almost every $x \in B$,

$$\tilde{F}(x) - F(x) = m_B(\tilde{F} - F)(x).$$

Now, we take a sequence $\{B_j\}_{j \in \mathbb{N}}$ of balls such that $B_1 \subset B_2 \subset \cdots \subset B_j \subset \cdots$ and $\bigcup_{j \in \mathbb{N}} B_j = \mathcal{X}$. Then, by (6.14), we know that there exists a sequence $\{F_j\}_{j \in \mathbb{N}}$ of measurable functions such that, for any $j \in \mathbb{N}$, $F_j \in L^{q'}(B_j)$ and, for any $g \in L^q_0(B_j)$,

$$T(g) = \int_{\mathcal{X}} F_j(x)g(x) d\mu(x). \quad (6.18)$$

Let $\widetilde{F}_1 := F_1$ and $\widetilde{F}_{j+1} := F_{j+1} + m_{B_j}(\widetilde{F}_j - F_{j+1})$ for any $j \in \mathbb{N}$. Then, by the above claim, we find that, for any $j \in \mathbb{N}$ and almost every $x \in B_j$, $\widetilde{F}_{j+1}(x) = \widetilde{F}_j(x)$ and $\widetilde{F}_j \in L^{q'}(B_j)$. Let $g \in H^{Y,q,d}_{\text{fin}}(\mathcal{X})$ and f be a measurable function satisfying that, for any $j \in \mathbb{N}$ and $x \in B_j$, $f(x) = \widetilde{F}_j(x)$. Then there exists some $j_0 \in \mathbb{N}$ such that $\text{supp } g \subset B_{j_0}$. By this, the cancellation of g , and (6.18), we know that

$$T(g) = \int_{B_{j_0}} F_{j_0}(x)g(x) d\mu(x) = \int_{B_{j_0}} \widetilde{F}_{j_0}(x)g(x) d\mu(x) = \int_{\mathcal{X}} f(x)g(x) d\mu(x). \quad (6.19)$$

It remains to prove that $f \in \mathcal{L}_{Y,q',d}(\mathcal{X})$. To this end, for any given $m \in \mathbb{N}$, any given balls $\{B_k\}_{k=1}^m$ of \mathcal{X} , and any given $\{\lambda_k\}_{k=1}^m \subset [0, \infty)$ with $\sum_{k=1}^m \lambda_k \neq 0$, and for any $k \in \{1, \dots, m\}$, let $h_k \in L^q(B_k)$ with $\|h_k\|_{L^q(B_k)} = 1$ be such that

$$\left[\int_{B_k} |f(x) - m_{B_k}(f)|^{q'} d\mu(x) \right]^{\frac{1}{q'}} = \int_{B_k} [f(x) - m_{B_k}(f)] h_k(x) d\mu(x) \quad (6.20)$$

and, for any $x \in \mathcal{X}$, let

$$a_k(x) := \begin{cases} 0 & \text{if } h_k \text{ is a constant function,} \\ \frac{[\mu(B_k)]^{\frac{1}{q}} [h_k(x) - m_{B_k}(h_k)] \mathbf{1}_{B_k}}{\|\mathbf{1}_{B_k}\|_{Y(\mathcal{X})} \|h_k - m_{B_k}(h_k)\|_{L^q(B_k)}} & \text{otherwise.} \end{cases}$$

Then it is easy to see that, for any $k \in \{1, \dots, m\}$, a_k is a $(Y(\mathcal{X}), q)$ -atom. From this and Theorem 4.19, it follows that $\sum_{k=1}^m \lambda_k a_k \in H_Y^*(\mathcal{X})$. This, together with (6.20), (6.16), (6.19), Theorem 4.23, $T \in (H_Y^*(\mathcal{X}))^*$, and $\|h_k - m_{B_k}(h_k)\|_{L^q(B_k)} \lesssim 1$ for any $k \in \{1, \dots, m\}$, further implies that

$$\begin{aligned} & \sum_{k=1}^m \frac{\lambda_k \mu(B_k)}{\|\mathbf{1}_{B_k}\|_{Y(\mathcal{X})}} \left[\frac{1}{\mu(B_k)} \int_{B_k} |f(x) - m_{B_k}(f)|^{q'} d\mu(x) \right]^{\frac{1}{q'}} \\ &= \sum_{k=1}^m \frac{\lambda_k [\mu(B_k)]^{\frac{1}{q}}}{\|\mathbf{1}_{B_k}\|_{Y(\mathcal{X})}} \int_{B_k} [f(x) - m_{B_k}(f)] h_k(x) d\mu(x) \\ &= \sum_{k=1}^m \frac{\lambda_k [\mu(B_k)]^{\frac{1}{q}}}{\|\mathbf{1}_{B_k}\|_{Y(\mathcal{X})}} \int_{B_k} [h_k(x) - m_{B_k}(h_k)] f(x) \mathbf{1}_{B_k}(x) d\mu(x) \\ &\lesssim \sum_{k=1}^m \lambda_k \int_{B_k} a_k(x) f(x) d\mu(x) \sim \sum_{k=1}^m \lambda_k T(a_k) \sim T\left(\sum_{k=1}^m \lambda_k a_k\right) \\ &\lesssim \|T\|_{(H_Y^*(\mathcal{X}))^*} \left\| \sum_{k=1}^m \lambda_k a_k \right\|_{H_Y^*(\mathcal{X})} \lesssim \|T\|_{(H_Y^*(\mathcal{X}))^*} \left\| \left\{ \sum_{k=1}^m \left[\frac{\lambda_k}{\|\mathbf{1}_{B_k}\|_{Y(\mathcal{X})}} \right]^d \mathbf{1}_{B_k} \right\}^{\frac{1}{d}} \right\|_{Y(\mathcal{X})}. \end{aligned}$$

This implies that $f \in \mathcal{L}_{Y,q',d}(\mathcal{X})$ and hence finishes the proof of (ii) and also of Theorem 6.6. \square

As a consequence of Theorem 6.6, we have the following equivalence of ball Campanato-type function spaces; we omit the details.

Corollary 6.7. *Let $Y(\mathcal{X})$, d , and p_0 be as in Theorem 6.6. Suppose $q \in [1, \infty)$ when $p_0 \in (0, 1)$, or $q \in [1, p'_0)$ when $p_0 \in [1, \infty)$. Then*

$$\mathcal{L}_{Y,q,d}(\mathcal{X}) = \mathcal{L}_{Y,1,d_0}(\mathcal{X})$$

with equivalent quasi-norms, where $d_0 \in (0, \theta_0)$ with θ_0 as in Theorem 6.6.

Remark 6.8. (i) We point out that the proof of Theorem 6.6 strongly depends on the fact that $H_{\text{fin}}^{Y,q,d}(\mathcal{X})$ is dense in $H_Y^*(\mathcal{X})$, which is guaranteed by $Y(\mathcal{X})$ having absolutely continuous quasi-norm (see Lemma 6.4). Therefore, the assumption that $Y(\mathcal{X})$ has an absolutely continuous quasi-norm in Theorem 6.6 is necessary.

(ii) Observe that we establish a continuous embedding of $Y(\mathcal{X})$ into the weighted Lebesgue space to overcome the difficulty caused by the lack of the good dense subset of $H_Y^*(\mathcal{X})$ in Section 4. However, this method does not work anymore in the proof of Theorem 6.6 because, in Section 4, we only need the convergence in the distributional sense of the atomic decomposition under consideration, but now we need its convergence in the quasi-norm $\|\cdot\|_{H_Y^*(\mathcal{X})}$.

(iii) Recall that, for any $\alpha \in (0, \infty)$, the *Lipschitz space* $\mathcal{L}_\alpha(\mathcal{X})$ is defined to be the set of all measurable functions f on \mathcal{X} such that

$$\|f\|_{\mathcal{L}_\alpha(\mathcal{X})} := \sup_{x \neq y} \frac{|f(x) - f(y)|}{[V(x, y)]^\alpha} < \infty.$$

Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. On one hand, by [14, Theorem B] and [31, Theorems 4.2 and 4.16], we know that the dual space of $H^{*,p}(\mathcal{X})$ is the space $\mathcal{L}_{1/p-1}(\mathcal{X})$ when $p < 1$, or $\text{BMO}(\mathcal{X})$ when $p = 1$ (see [14, p. 593] for the definition of $\text{BMO}(\mathcal{X})$). On the other hand, if $Y(\mathcal{X}) := L^p(\mathcal{X})$, then, by Remark 4.21(i) and Theorem 6.6, we conclude that the dual space of $H^{*,p}(\mathcal{X})$ is $\mathcal{L}_{Y,q',d}(\mathcal{X})$ with $Y(\mathcal{X})$ replaced by $L^p(\mathcal{X})$. Thus, Theorem 6.6 give an equivalent characterization of $\mathcal{L}_{1/p-1}(\mathcal{X})$ when $p < 1$, or $\text{BMO}(\mathcal{X})$ when $p = 1$.

(iv) Let $p \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3, and $r \in (0, \infty)$. If $Y(\mathcal{X}) := L^{p,r}(\mathcal{X})$, then, by Remark 4.21(ii), we know that $L^{p,r}(\mathcal{X})$ satisfies all the assumptions of Theorem 6.6. In this case, Theorem 6.6 is new.

(v) Let $p \in (\omega/(\omega + \eta), 1]$, $r \in (1, \infty)$, $w \in A_r(\mathcal{X})$ satisfy

$$r_w := \inf \{r \in [1, \infty) : \varphi \in A_r(\mathcal{X})\} \in (1, p(\omega + \eta)/\omega)$$

with ω as in (2.3) and η as in Definition 3.3. If $Y(\mathcal{X}) := L_w^p(\mathcal{X})$, then, by Remark 4.21(iii), we know that $L_w^p(\mathcal{X})$ satisfies all the assumptions of Theorem 6.6. From this and a generalization of [75, Proposition 3.7] on \mathbb{R}^n to \mathcal{X} , we deduce that [19, Theorem 8.1] is a special case of Theorem 6.6.

- (vi) Let Φ be an Orlicz function, $p_{\Phi}^{+} = 1$, and $p_{\Phi}^{-} \in (\omega/(\omega + \eta), 1]$ with ω as in (2.3) and η as in Definition 3.3. If $Y(X) := L^{\Phi}(X)$, then, by Remark 4.21(iv), we know that $L^{\Phi}(X)$ satisfies all the assumptions of Theorem 6.6. From this and a generalization of [75, Proposition 3.7] on \mathbb{R}^n to X , we deduce that [19, Theorem 8.1] is a special case of Theorem 6.6.
- (vii) Let (X, ρ, μ) be an RD-space, and $p(\cdot) \in C^{\log}(X)$ satisfy $p_{+} \in (0, 1]$ and $\widetilde{p}_{-} \in (\omega/(\omega + \eta), \infty)$, where ω is as in (2.3) and η as in Definition 3.3. If $Y(X) := L^{p(\cdot)}(X)$, then, by Remark 4.21(v), we know that $L^{p(\cdot)}(X)$ satisfies all the assumptions of Theorem 6.6. In this case, Theorem 6.6 improves the corresponding results in [78, Theorem 7.2] by removing the additional restriction $p_{+} \in (0, 1]$ and the reverse doubling assumption of μ .
- (viii) Let φ be a growth function as in Remark 2.7(iii). As was mentioned in Remark 4.21, Theorem 6.6 is not applicable to $H^{*,\varphi}(X)$. Fu et al. [19] obtained a similar conclusion of Theorem 6.6 in case that $Y(X) := L^{\varphi}(X)$ independently.

Acknowledgements The authors would like to thank Fan Wang for some helpful discussions on the subject of this article.

References

- [1] W. Abu-Shammala and A. Torchinsky, The Hardy–Lorentz spaces $H^{p,q}(\mathbb{R}^n)$, *Studia Math.* 182 (2007), 283–294.
- [2] T. Aoki, Locally bounded linear topological spaces, *Proc. Imp. Acad. Tokyo* 18 (1942), 588–594.
- [3] H. Arai and T. Mizuhara, Morrey spaces on spaces of homogeneous type and estimates for \square_b and the Cauchy–Szegő projection, *Math. Nachr.* 185 (1997), 5–20.
- [4] P. Auscher and T. Hytönen, Orthonormal bases of regular wavelets in spaces of homogeneous type, *Appl. Comput. Harmon. Anal.* 34 (2013), 266–296.
- [5] C. Bennett and R. Sharpley, *Interpolation of Operators*, Pure Appl. Math. 129, Academic Press, Boston, MA, 1988.
- [6] T. A. Bui, P. D’Ancona, X.-T. Duong and D. Müller, On the flows associated to selfadjoint operators on metric measure spaces, *Math. Ann.* 375 (2019), 1393–1426.
- [7] T. A. Bui, P. D’Ancona and F. Nicola, Sharp L^p estimates for Schrödinger groups on spaces of homogeneous type, *Rev. Mat. Iberoam.* 36 (2020), 455–484.
- [8] T. A. Bui and X.-T. Duong, Sharp weighted estimates for square functions associated to operators on spaces of homogeneous type, *J. Geom. Anal.* 30 (2020), 874–900.
- [9] T. A. Bui, X.-T. Duong and F. K. Ly, Maximal function characterizations for new local Hardy-type spaces on spaces of homogeneous type, *Trans. Amer. Math. Soc.* 370 (2018), 7229–7292.
- [10] T. A. Bui, X.-T. Duong and F. K. Ly, Maximal function characterizations for Hardy spaces on spaces of homogeneous type with finite measure and applications, *J. Funct. Anal.* 278 (2020), 108423, 55 pp.
- [11] D.-C. Chang, S. Wang, D. Yang and Y. Zhang, Littlewood–Paley characterizations of Hardy-type spaces associated with ball quasi-Banach function spaces, *Complex Anal. Oper. Theory* 14 (2020), Paper No. 40, 33 pp.

- [12] C. Chen, J. Li and F. Liao, Some function spaces via orthonormal bases on spaces of homogeneous type, *Abstr. Appl. Anal.* 2014, Art. ID 265378, 13 pp.
- [13] R. R. Coifman and G. Weiss, *Analyse Harmonique Non-Commutative sur Certains Espaces Homogènes. (French) Étude de Certaines Intégrales Singulières*, Lecture Notes in Mathematics 242, Springer-Verlag, Berlin–New York, 1971.
- [14] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* 83 (1977), 569–645.
- [15] D. Deng and Y. Han, *Harmonic Analysis on Spaces of Homogeneous Type*, Lecture Notes in Math. 1966, Springer-Verlag, Berlin, 2009.
- [16] L. Diening, P. Harjulehto, P. Hästö and M. Růžička, *Lebesgue and Sobolev Spaces with Variable Exponents*, Lecture Notes in Math. 2017, Springer, Heidelberg, 2011.
- [17] X.-T. Duong, R. Gong, M. S. Kuffner, J. Li, B. D. Wick and D. Yang, Two weight commutators on spaces of homogeneous type and applications, *J. Geom. Anal.* 31 (2021), 980–1038.
- [18] C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.* 129 (1972), 137–193.
- [19] X. Fu, T. Ma and D. Yang, Real-variable characterizations of Musielak–Orlicz Hardy spaces on spaces of homogeneous type, *Ann. Acad. Sci. Fenn. Math.* 45 (2020), 343–410.
- [20] X. Fu and D. Yang, Wavelet characterizations of the atomic Hardy space H^1 on spaces of homogeneous type, *Appl. Comput. Harmon. Anal.* 44 (2018), 1–37.
- [21] X. Fu, D. Yang and Y. Liang, Products of functions in $BMO(X)$ and $H_{at}^1(X)$ via wavelets over spaces of homogeneous type, *J. Fourier Anal. Appl.* 23 (2017), 919–990.
- [22] L. Grafakos, *Classical Fourier Analysis*, Third edition, Graduate Texts in Mathematics 249, Springer, New York, 2014.
- [23] L. Grafakos, L. Liu and D. Yang, Maximal function characterizations of Hardy spaces on RD-spaces and their applications, *Sci. China Ser. A* 51 (2008), 2253–2284.
- [24] L. Grafakos, L. Liu and D. Yang, Vector-valued singular integrals and maximal functions on spaces of homogeneous type, *Math. Scand.* 104 (2009), 296–310.
- [25] R. Gong, J. Li, E. Pozzi and M. N. Vempati, Commutators on weighted Morrey spaces on spaces of homogeneous type, *Anal. Geom. Metr. Spaces* 8 (2020), 305–334.
- [26] Ya. Han, Yo. Han and J. Li, Criterion of the boundedness of singular integrals on spaces of homogeneous type, *J. Funct. Anal.* 271 (2016), 3423–3464.
- [27] Y. Han, J. Li and L. A. Ward, Hardy space theory on spaces of homogeneous type via orthonormal wavelet bases, *Appl. Comput. Harmon. Anal.* 45 (2018), 120–169.
- [28] Y. Han, D. Müller and D. Yang, Littlewood–Paley characterizations for Hardy spaces on spaces of homogeneous type, *Math. Nachr.* 279 (2006), 1505–1537.
- [29] Y. Han, D. Müller and D. Yang, A theory of Besov and Triebel–Lizorkin spaces on metric measure spaces modeled on Carnot–Carathéodory spaces, *Abstr. Appl. Anal.* 2008, Art. ID 893409, 1–250.
- [30] Z. He, Y. Han, J. Li, L. Liu, D. Yang and W. Yuan, A complete real-variable theory of Hardy spaces on spaces of homogeneous type, *J. Fourier Anal. Appl.* 25 (2019), 2197–2267.
- [31] Z. He, L. Liu, D. Yang and W. Yuan, New Calderón reproducing formulae with exponential decay on spaces of homogeneous type, *Sci. China Math.* 62 (2019), 283–350.
- [32] Z. He, F. Wang, D. Yang and W. Yuan, Wavelet characterization of Besov and Triebel–Lizorkin spaces on spaces of homogeneous type and its applications, *Appl. Comput. Harmon. Anal.* 54 (2021), 176–226.

- [33] Z. He, D. Yang and W. Yuan, Real-variable characterizations of local Hardy spaces on spaces of homogeneous type, *Math. Nachr.* 294 (2021), 900–955.
- [34] J. Heinonen, *Lectures on Analysis on Metric Spaces*, Springer-Verlag, New York, 2001.
- [35] K.-P. Ho, Atomic decomposition of Hardy–Morrey spaces with variable exponents, *Ann. Acad. Sci. Fenn. Math.* 40 (2015), 31–62.
- [36] K.-P. Ho, Atomic decompositions and Hardy’s inequality on weak Hardy–Morrey spaces, *Sci. China Math.* 60 (2017), 449–468.
- [37] S. Hou, D. Yang and S. Yang, Lusin area function and molecular characterizations of Musielak–Orlicz Hardy spaces and their applications, *Commun. Contemp. Math.* 15 (2013), 1350029, 37 pp.
- [38] S. Hou, D. Yang and S. Yang, Musielak–Orlicz BMO-type spaces associated with generalized approximations to the identity, *Acta Math. Sin. (Engl. Ser.)* 30 (2014), 1917–1962.
- [39] L. Huang, D. Yang and W. Yuan, Anisotropic mixed-norm Campanato-type spaces with applications to duals of anisotropic mixed-norm Hardy spaces, *Banach J. Math. Anal.* 15 (2021), no. 4, Paper No. 62, 36 pp.
- [40] T. Hytönen and A. Kairema, Systems of dyadic cubes in a doubling metric space, *Colloq. Math.* 126 (2012), 1–33.
- [41] M. Izuki, T. Noi and Y. Sawano, The John–Nirenberg inequality in ball Banach function spaces and application to characterization of BMO, *J. Inequal. Appl.* 2019, Paper No. 268, 11 pp.
- [42] M. Izuki and Y. Sawano, Characterization of BMO via ball Banach function spaces, *Vestn. St.-Peterbg. Univ. Mat. Mekh. Astron.* 4(62) (2017), 78–86.
- [43] M. Kronz, Some function spaces on spaces of homogeneous type, *Manuscripta Math.* 106 (2001), 219–248.
- [44] J. Li, Atomic decomposition of weighted Triebel–Lizorkin spaces on spaces of homogeneous type, *J. Aust. Math. Soc.* 89 (2010), 255–275.
- [45] J. Li and L. A. Ward, Singular integrals on Carleson measure spaces CMO^p on product spaces of homogeneous type, *Proc. Amer. Math. Soc.* 141 (2013), 2767–2782.
- [46] W. Li, A maximal function characterization of Hardy spaces on spaces of homogeneous type, *Approx. Theory Appl. (N. S.)* 14 (1998), 12–27.
- [47] J. Liu, D. Yang and W. Yuan, Littlewood–Paley characterizations of anisotropic Hardy–Lorentz spaces, *Acta Math. Sci. Ser. B (Engl. Ed.)* 38 (2018), 1–33.
- [48] L. Liu, D.-C. Chang, X. Fu and D. Yang, Endpoint estimates of linear commutators on Hardy spaces over spaces of homogeneous type, *Math. Methods Appl. Sci.* 41 (2018), 5951–5984.
- [49] S. Lu and D. Yang, The weighted Herz-type Hardy spaces and its applications, *Sci. China Ser. A* 38 (1995), 662–673.
- [50] R. A. Macías and C. Segovia, Lipschitz functions on spaces of homogeneous type, *Adv. Math.* 33 (1979), 257–270.
- [51] R. A. Macías and C. Segovia, A decomposition into atoms of distributions on spaces of homogeneous type, *Adv. Math.* 33 (1979), 271–309.
- [52] S. Müller, Hardy space methods for nonlinear partial differential equations, *Tatra Mt. Math. Publ.* 4 (1994), 159–168.
- [53] E. Nakai, The Campanato, Morrey and Hölder spaces on spaces of homogeneous type, *Studia Math.* 176 (2006), 1–19.

- [54] E. Nakai and Y. Sawano, Hardy spaces with variable exponents and generalized Campanato spaces, *J. Funct. Anal.* 262 (2012), 3665–3748.
- [55] E. Nakai and K. Yabuta, Pointwise multipliers for functions of weighted bounded mean oscillation on spaces of homogeneous type, *Math. Japon.* 46 (1997), 15–28.
- [56] S. Rolewicz, On a certain class of linear metric spaces, *Bull. Acad. Polon. Sci. Cl. III.* 5 (1957), 471–473.
- [57] W. Rudin, *Functional Analysis*, Second edition, International Series in Pure and Applied Mathematics, McGraw-Hill, Inc., New York, 1991.
- [58] Y. Sawano, *Theory of Besov Spaces*, Developments in Mathematics 56, Springer, Singapore, 2018.
- [59] Y. Sawano, K.-P. Ho, D. Yang and S. Yang, Hardy spaces for ball quasi-Banach function spaces, *Dissertationes Math.* 525 (2017), 1–102.
- [60] Y. Sawano, T. Shimomura and H. Tanaka, A remark on modified Morrey spaces on metric measure spaces, *Hokkaido Math. J.* 47 (2018), 1–15.
- [61] Y. Sawano and H. Tanaka, The Fatou property of block spaces, *J. Math. Sci. Univ. Tokyo* 22 (2015), 663–683.
- [62] I. Sihwaningrum, H. Gunawan and E. Nakai, Maximal and fractional integral operators on generalized Morrey spaces over metric measure spaces, *Math. Nachr.* 291 (2018), 1400–1417.
- [63] E. M. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*, Princeton Mathematical Series 43, Princeton University Press, Princeton, NJ, 1993.
- [64] E. M. Stein and G. Weiss, On the theory of harmonic functions of several variables. I. The theory of H^p -spaces, *Acta Math.* 103 (1960), 25–62.
- [65] J.-O. Strömberg and A. Torchinsky, *Weighted Hardy Spaces*, Lecture Notes in Math. 1381, Springer-Verlag, Berlin, 1989.
- [66] F. Wang, Y. Han, Z. He and D. Yang, Besov and Triebel–Lizorkin spaces on spaces of homogeneous type with applications to boundedness of Calderón–Zygmund operators, *Dissertationes Math.* (2021), doi: 10.4064/dm821-4-2021.
- [67] F. Wang, Z. He, D. Yang and W. Yuan, Difference characterization of Besov and Triebel–Lizorkin spaces on spaces of homogeneous type, *Commun. Math. Stat.* (2021), <https://doi.org/10.1007/s40304-021-00243-w>.
- [68] F. Wang, D. Yang and S. Yang, Applications of Hardy spaces associated with ball quasi-Banach function spaces, *Results Math.* 75 (2020), Paper No. 26, 58 pp.
- [69] S. Wang, D. Yang, W. Yuan and Y. Zhang, Weak Hardy-type spaces associated with ball quasi-Banach function spaces II: Littlewood–Paley characterizations and real interpolation, *J. Geom. Anal.* 31 (2021), 631–696.
- [70] X. Yan, Z. He, D. Yang and W. Yuan, Hardy spaces associated with ball quasi-Banach function spaces on spaces of homogeneous type: Littlewood–Paley characterizations with applications to boundedness of Calderón–Zygmund operators, Submitted.
- [71] X. Yan, D. Yang and W. Yuan, Intrinsic square function characterizations of Hardy spaces associated with ball quasi-Banach function spaces, *Front. Math. China* 15 (2020), 769–806.
- [72] X. Yan, D. Yang and W. Yuan, Intrinsic square function characterizations of several Hardy-type spaces – a survey, *Anal. Theory Appl.* 37 (2021), 426–464.
- [73] D. Yang, Y. Liang and L. D. Ky, *Real-Variable Theory of Musielak–Orlicz Hardy Spaces*, Lecture Notes in Mathematics 2182, Springer, Cham, 2017.

- [74] D. Yang and Y. Zhou, New properties of Besov and Triebel–Lizorkin spaces on RD-spaces, *Manuscripta Math.* 134 (2011), 59–90.
- [75] Y. Zhang, L. Huang, D. Yang and W. Yuan, New ball Campanato-type function spaces and their applications, *J. Geom. Anal.* (to appear).
- [76] Y. Zhang, S. Wang, D. Yang and W. Yuan, Weak Hardy-type spaces associated with ball quasi-Banach function spaces I: decompositions with applications to boundedness of Calderón–Zygmund operators, *Sci. China Math.* 64 (2021), 2007–2064.
- [77] X. Zhou, Z. He and D. Yang, Real-variable characterizations of Hardy–Lorentz spaces on spaces of homogeneous type with applications to real interpolation and boundedness of Calderón–Zygmund operators, *Anal. Geom. Metr. Spaces* 8 (2020), 182–260.
- [78] C. Zhuo, Y. Sawano and D. Yang, Hardy spaces with variable exponents on RD-spaces and applications, *Dissertationes Math.* 520 (2016), 1–74.

Xianjie Yan, Dachun Yang (Corresponding author) and Wen Yuan

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, People’s Republic of China

E-mails: xianjieyan@mail.bnu.edu.cn (X. Yan)

dcyang@bnu.edu.cn (D. Yang)

wenyuan@bnu.edu.cn (W. Yuan)

Ziyi He

School of Science, Beijing University of Posts and Telecommunications, Beijing, 100876, People’s Republic of China

E-mail: ziyihe@bupt.edu.cn (Z. He)