TWO-TERM SILTING AND τ -CLUSTER MORPHISM CATEGORIES

ERLEND D. BØRVE

ABSTRACT. We generalise τ -cluster morphism categories to non-positive proper dg algebras. The compatibility of silting reduction with support τ -tilting reduction will be an essential ingredient when linking our definition to that of Buan–Marsh. We also define two-term presilting sequences in the bounded derived category in such a way that they correspond to signed τ -exceptional sequences in the module category.

CONTENTS

0. Introduction	1
0.1. Organisation	2
1. Notation and preliminaries	4
1.1. t-structures and wide subcategories of the heart	5
1.2. Silting and τ -tilting	9
2. Silting t-structures and reduction	12
3. Compatibility of the Iyama–Yang and Buan–Marsh bijections	18
4. τ -cluster morphism categories	22
5. Two-term (pre)silting sequences	25
References	27

0. INTRODUCTION

In tilting theory, it is valuable to find all tilting modules over a fixed finite-dimensional algebra A, or better still, all tilting objects in the bounded homotopy category $\mathcal{K}^{\mathrm{b}}(\mathrm{proj}\,A)$, where $\mathrm{proj}\,A$ is the category of finitely generated projective A-modules. If all tilting objects in $\mathcal{K}^{\mathrm{b}}(\mathrm{proj}\,A)$ could be identified, one could construct all finite-dimensional algebras B that are derived Morita equivalent to A [Ric89], which is to say that their derived categories are equivalent as triangulated categories. Many invariants of finite-dimensional algebras are preserved by derived Morita equivalence, and it might be more efficient to compute these for some B rather than for A directly.

Finding all tilting objects turns out to be an unrealistic project in general. However, one could instead study the larger class of silting objects, for which there is a mutation process [AI12], generating new examples in a reliable manner. In addition, one has a reduction procedure for silting objects. Iyama– Yang show that if P is a presilting object of $\mathcal{K}^{\mathrm{b}}(\mathrm{proj} A)$, then the (pre)silting objects in $\mathcal{K}^{\mathrm{b}}(\mathrm{proj} A)$ having P as a direct summand correspond to the (pre)silting objects in the Verdier quotient $\mathcal{K}^{\mathrm{b}}(\mathrm{proj} A)/\mathrm{thick}(P)$ [IY18, Theorem 3.7]. The class of two-term silting objects closely resembles that of tilting modules. Indeed, a generalisation of the Brenner–Butler Theorem applies [BZ16, Theorem 1.1] [BZ21, Theorem 2.1]. Also, when performing silting reduction with respect to a two-term presilting object, one can restrict Iyama–Yang's bijection to a bijection of two-term (pre)silting objects [Jas15, Proposition 4.11].

The mutation of two-term silting objects is closely tied to cluster combinatorics. For path algebras of a quiver of simply laced Dynkin type, the two-term silting objects of the bounded homotopy category are in bijection with the cluster-tilting objects in the cluster category [AIR14, Theorem 4.1]. Moreover, this bijection respects the mutation processes for cluster-tilting objects [AIR14, Corollary 4.8]. Cluster-tilting objects are in turn in bijection with the clusters of the corresponding cluster algebra [BMR⁺06, Corollary 4.4].

One may also model cluster combinatorics using τ -tilting theory [AIR14]. Adachi–Iyama–Reiten define support τ -tilting modules and support τ -rigid pairs, and show that they correspond bijectively with silting objects and presilting objects, respectively [AIR14, Theorem 3.2]. One would thus suspect that the reduction procedure of Iyama–Yang is transferable to τ -tilting theory. Given any finite-dimensional algebra, Jasso provides a suitable bijection between classes of support τ -tilting modules [Jas15, Theorem 3.15], and shows that it is compatible with Iyama–Yang's silting reduction [Jas15, Theorem 4.12(b)]. Buan–Marsh later extended Jasso's bijection to support τ -rigid pairs [BM21a, Theorem 3.6].

The bijection of Buan–Marsh was constructed in order to define the τ -cluster morphism category of a τ tilting finite algebra. This builds on Igusa–Todorov's cluster morphism category defined for representation finite hereditary algebras [IT17, Section 1], as well as a categorification of non-crossing partitions [Igu14]. The definition of τ -cluster morphism categories has recently been extended to all finite-dimensional algebras [BH21]. Although Buan–Marsh define the τ -cluster morphism category in terms of τ -tilting theory, they occasionally translate the setting to two-term silting in order to prove necessary results.

 τ -cluster morphism categories are useful in the study of picture groups and picture spaces, defined by Igusa–Todorov–Weyman [ITW16]. In the case of representation finite hereditary algebras, we have that the geometric realisation of the τ -cluster morphism category is a $K(\pi, 1)$ for the corresponding picture group [IT17, Theorem 3.1]. This result was later extended to include Nakayama algebras [HI21, Theorem 4.16].

A path in a τ -cluster morphism category can be interpreted as a signed τ -exceptional sequence, a notion defined in a parallel paper of Buan–Marsh [BM21b]. In the hereditary representation finite case, these are signed exceptional sequences [IT17, Section 2], which in informal terms can be seen as exceptional sequences where relative projective objects may be suspended in the derived category. A complete exceptional sequence need not exist for a finite-dimensional algebra, whereas one may always find a complete signed τ -exceptional sequence. One proves this fact using the correspondence between signed τ -exceptional sequences and ordered support τ -rigid modules [BM21b, Corollary 5.5]. If a τ -exceptional sequence consists only of modules (rather than suspended relative projectives) it can be obtained from the stratifying system of a TF-admissible decomposition of a τ -rigid module [MHT20, Theorem 5.1].

0.1. **Organisation.** Section 1 sets up the notation and covers the necessary preliminaries. We review how t-exact subcategories (see Definition 1.5) of a triangulated category with a bounded t-structure

correspond with wide subcategories in the heart (Theorem 1.10), the interaction between two-term silting

and τ -tilting theory, and τ -tilting reduction. In particular, we recall in Proposition-Definition 1.17 that a support τ -rigid pair (M, Q) in mod(A) gives rise to a wide subcategory $J(M, Q) \stackrel{\text{def}}{=} M^{\perp} \cap^{\perp}(\tau M) \cap Q^{\perp}$ of mod(A).

In Section 2, we work over a non-positive proper dg algebra A. Let $\mathcal{D}_{\mathrm{fd}}(A)$ be the subcategory of the derived category $\mathcal{D}(A)$ spanned by complexes of finite-dimensional total cohomology. In Corollary 2.9, we show that the perpendicular category $P^{\perp_{\mathbb{Z}}}$ of a two-term presilting object $P \in \mathrm{per}(A)$ is a t-exact subcategory of $\mathcal{D}_{\mathrm{fd}}(A)$. By Proposition 2.10, the zeroth cohomology of $P^{\perp_{\mathbb{Z}}}$ is the wide subcategory J(M, Q) of $\mathrm{mod}(A)$, where (M, Q) is the support τ -rigid pair corresponding to P.

Section 3 presents a proof that Iyama–Yang's silting reduction is compatible with Buan–Marsh' support τ -tilting reduction, generalising Jasso's compatibility result [Jas15, Theorem 4.12(b)].

Theorem 3.5. Let \mathcal{C} be a triangulated category and S be a silting object. Certain mild technical conditions are imposed. Consider a 2_S-term presilting object P, let $A = \operatorname{End}_{\mathcal{C}}(S)$, and let $C = \operatorname{End}_{\mathcal{Z}_P/[P]}(T_P^+)$ be the τ -tilting reduction of A with respect to the support τ -rigid pair corresponding to P. We have a commutative diagram of bijections

where φ_P is the Iyama–Yang bijection and $\psi_{H_S(P)}$ is the Buan–Marsh bijection. The horizontal maps are the correspondences between two-term presilting objects and support τ -rigid pairs (see Theorem 1.16).

Our main result appears in Section 4. Given a non-positive proper dg algebra A, we construct a category \mathfrak{W}_A , which we call the τ -cluster morphism category of A. Developing the theory purely in terms of silting, as opposed to τ -tilting, has a major advantage; since silting reduction is induced by a functor, it is easier to prove that the composition law in \mathfrak{W}_A is associative, as we do in Theorem 4.3. When we claim that the work of Buan–Marsh and Buan–Hanson has been generalised, it is in the following specific sense.

Theorem 4.4. Let A be a non-positive proper dg algebra, and let $\mathfrak{W}_{H^0A}^{BM}$ be τ -cluster morphism category (as defined by Buan–Hanson [BH21]) of the zeroth cohomology of A. Then the cohomological functor

$$\mathcal{D}_{\mathrm{fd}}(A, -) \colon \mathcal{D}_{\mathrm{fd}}(\mathrm{mod}\,A) \longrightarrow \mathrm{mod}\,H^0A$$

induces an equivalence of categories

$$\mathfrak{W}_A \longrightarrow \mathfrak{W}_{H^0A}^{\mathrm{BM}}$$

In particular, if A is a finite-dimensional algebra, then \mathfrak{W}_A is equivalent to \mathfrak{W}_A^{BM} .

Finally, Section 5 explores how signed τ -exceptional sequences can be lifted to our framework. We define two-term presilting sequences in a recursive manner, similarly to Buan–Marsh [BM21b, Definition 1.3]. A correspondence between our notion and theirs is provided: **Theorem 5.2.** Let A be a non-positive proper dg algebra, and let t be a non-negative integer. If H_A is as defined in Theorem 1.16, then the map

$$(X_1,\ldots,X_{t-1},X_t)\longmapsto (H_AX_1,\ldots,H_AX_{t-1},H_AX_t),$$

is a bijection from the first to the second of the following sets :

- (1) signed two-term presilting sequences of length t in $\mathcal{D}_{\rm fd}(A)$,
- (2) signed τ -exceptional sequences of length t in mod (H^0A) .

Acknowledgements. I would like to express my gratitude to my PhD supervisor, Aslak B. Buan, for introducing me to the topic, countless helpful discussions, and meticulous proofreading. I am also grateful to Eric J. Hanson and Yann Palu for helpful comments on this manuscript, and to Gustavo Jasso for useful conversations.

1. NOTATION AND PRELIMINARIES

Throughout, all subcategories will be full and closed under isomorphism. We fix a field k, and declare that all triangulated categories and their (not necessarily triangulated) subcategories will be k-linear.

Recall that a triangulated subcategory is *thick* if it is closed under direct summands. For a triangulated category \mathcal{T} , we write thick_{\mathcal{T}}(\mathcal{X}) for the smallest thick subcategory containing the subcategory \mathcal{X} , or simply thick(\mathcal{X}) when there is no risk of confusion.

All modules are right modules, unless otherwise specified. For a k-algebra R, the category of finitely generated right R-modules will be denoted by mod R, and the subcategory of projective R-modules by proj R. The triangulated category $\mathcal{D}^{\mathrm{b}}(\mathrm{mod}\,R)$ is the bounded derived category of the former, whereas $\mathcal{K}^{\mathrm{b}}(\mathrm{proj}\,R)$ is the bounded homotopy category of the latter. They are both subcategories of the full derived category $\mathcal{D}(R)$. More generally, if A is a differential graded (henceforth dg) k-algebra A, we denote its full derived category by $\mathcal{D}(A)$, and define $\mathrm{per}(A)$ to the thick closure of A in $\mathcal{D}(A)$, and $\mathcal{D}_{\mathrm{fd}}$ to be the (thick) subcategory spanned by complexes of finite-dimensional total cohomology.

Let \mathcal{C} be a k-linear category and let $\mathcal{X} \subseteq \mathcal{C}$ be a subcategory. The *additive closure* of \mathcal{X} is the full subcategory of \mathcal{C} containing all finite direct sums of direct summands of objects in \mathcal{X} . We denote this category by $\operatorname{add}(\mathcal{X})$, or by $\operatorname{add}(\mathcal{X})$ if \mathcal{X} contains a single object X.

Let \mathcal{P} be an additive subcategory of \mathcal{C} . A morphism $Q \xrightarrow{\beta} Y$ (resp. $X \xrightarrow{\alpha} Q$) is \mathcal{P} -epic (resp. \mathcal{P} monic) if the induced morphism $\mathcal{C}(P,\beta)$ (resp. $\mathcal{C}(\alpha, P)$) is surjective for all objects $P \in \mathcal{P}$. It is a right \mathcal{P} -approximation of Y (resp. left \mathcal{P} -approximation of X) if, in addition, we have that $Q \in \mathcal{P}$. We say that \mathcal{P} is a contravariantly finite subcategory (resp. covariantly finite subcategory) of \mathcal{C} if every object in \mathcal{C} has a right (resp. left) \mathcal{P} -approximation. A functorially finite subcategory of \mathcal{C} is one that is both contravariantly and covariantly finite.

The *perpendicular subcategories* of \mathcal{P} in \mathcal{C} are the full subcategories

$${}^{\perp}\mathcal{P} \stackrel{\text{def}}{=} \{ X \in \mathcal{C} \mid \mathcal{C}(X, \mathcal{P}) = 0 \},\$$
$$\mathcal{P}^{\perp} \stackrel{\text{def}}{=} \{ Y \in \mathcal{C} \mid \mathcal{C}(\mathcal{P}, Y) = 0 \}.$$

If \mathcal{T} is a triangulated category, we denote its suspension functor by Σ , unless otherwise specified. For each subset $I \subseteq \mathbb{Z}$, we define the *perpendicular subcategories*

$$\overset{\perp_{I}}{\stackrel{\mathcal{P}}{=}} \stackrel{\text{def}}{=} \{ X \in \mathcal{T} \mid \mathcal{T}(X, \Sigma^{i}\mathcal{P}) = 0 \quad \forall i \in I \},$$
$$\mathcal{P}^{\perp_{I}} \stackrel{\text{def}}{=} \{ Y \in \mathcal{T} \mid \mathcal{T}(\mathcal{P}, \Sigma^{i}Y) = 0 \quad \forall i \in I \}.$$

Suppose that \mathcal{T} is triangulated and let \mathcal{P}_1 and \mathcal{P}_2 be additive subcategories of \mathcal{T} . The category of extensions is the full subcategory $\mathcal{P}_1 * \mathcal{P}_2$ of \mathcal{T} containing the objects E that fit in a triangle

 $X_1 \longrightarrow E \longrightarrow X_2 \longrightarrow \Sigma X_1,$

where $X_1 \in \mathcal{P}_1$ and $X_2 \in \mathcal{P}_2$. It can be shown using the octahedral axiom that * is an associative operation on subcategories [BBD82, Lemme 1.3.10]. A full subcategory $\mathcal{P} \subseteq \mathcal{T}$ is closed under extensions if $\mathcal{P} = \mathcal{P} * \mathcal{P}$.

An additive subcategory $\mathcal{P} \subseteq \mathcal{T}$ is *suspended* (resp. *co-suspended*) if it is closed under extensions and the suspension functor Σ (resp. the desuspension functor Σ^{-1}).

The ideal $[\mathcal{P}]$ in \mathcal{C} contains precisely the morphisms that factor through an object in \mathcal{P} . The ideal quotient will be denoted by $\frac{\mathcal{C}}{|\mathcal{P}|}$. If $\mathcal{P} = \operatorname{add}(P)$ for some object P, we denote this quotient by $\frac{\mathcal{C}}{|\mathcal{P}|}$.

We use Deligne's convention, writing F = G when we mean that these functors are naturally isomorphic.

1.1. t-structures and wide subcategories of the heart. Truncation structures (t-structures for short) were introduced by Beilinson–Bernstein–Deligne [BBD82] (and also Gabber [BBDG18]). We give a quick survey of their elementary properties. A triangulated category \mathcal{D} is fixed.

Definition 1.1. A pair of full additive subcategories $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$ of \mathcal{D} constitutes a truncation structure (henceforth *t*-structure) if the conditions (t.1) and (t.2) below are met.

- (t.1) $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$ is a torsion pair in \mathcal{D} , i.e.
 - (t.1.1) The Hom-spaces $\mathcal{D}(X, Y)$ are trivial for all $X \in \mathcal{D}^{\leq 0}$ and $Y \in \mathcal{D}^{>0}$,

ar

(t.1.2) every object in \mathcal{D} is an extension of an object in $\mathcal{D}^{>0}$ with an object in $\mathcal{D}^{\leq 0}$. This is to say that $\mathcal{D}^{\leq 0} * \mathcal{D}^{>0} = \mathcal{D}$.

(t.2) The subcategory $\mathcal{D}^{\leq 0}$ is suspended (equivalently, the subcategory $\mathcal{D}^{>0}$ is co-suspended).

We let $\mathcal{D}^{\leq -1} \stackrel{\text{def}}{=} \Sigma \mathcal{D}^{\leq 0}$. More generally, we set

$$\mathcal{D}^{\leq n} \stackrel{\text{def}}{=} \Sigma^{-n} \mathcal{D}^{\leq 0},$$
$$\mathcal{D}^{>m} \stackrel{\text{def}}{=} \Sigma^{-m} \mathcal{D}^{>0},$$
ad
$$\mathcal{D}^{[m,n]} \stackrel{\text{def}}{=} \mathcal{D}^{>m-1} \cap \mathcal{D}^{\leq n}$$

for integers n and m. The t-structure is bounded if $\bigcup_{m,n\in\mathbb{Z}} \mathcal{D}^{[m,n]} = \mathcal{D}$. The subcategory $\mathcal{D}^0 \stackrel{\text{def}}{=} \mathcal{D}^{[0,0]}$ of \mathcal{D} is called the *heart* of the t-structure.

Remark 1.2. The subcategories $\mathcal{D}^{\leq 0}$ and $\mathcal{D}^{>0}$ determine each other; we have that $(\mathcal{D}^{\leq 0})^{\perp_0} = \mathcal{D}^{>0}$ and that $^{\perp_0}(\mathcal{D}^{>0}) = \mathcal{D}^{\leq 0}$. As a result of this fact, the subcategories $\mathcal{D}^{\leq 0}$ and $\mathcal{D}^{>0}$ are closed under extensions and direct summands.

Let \mathcal{A} be an abelian category. Then the bounded derived category $\mathcal{D}^{b}(\mathcal{A})$ can be equipped with the structure

$$\mathcal{D}^{\leq 0}(\mathcal{A}) = \{ X \in \mathcal{D}^{\mathbf{b}}(\mathcal{A}) \mid H^{i}(X) = 0 \text{ for all } i > 0 \},\$$
$$\mathcal{D}^{>0}(\mathcal{A}) = \{ X \in \mathcal{D}^{\mathbf{b}}(\mathcal{A}) \mid H^{i}(X) = 0 \text{ for all } i \leq 0 \}.$$

This is the *standard t-structure* on $\mathcal{D}^{\mathrm{b}}(\mathcal{A})$. The heart is equivalent to the abelian category \mathcal{A} . In general, we have that the heart of any t-structure is an abelian category [BBD82, Théorème 1.3.6].

For a suspended subcategory $\mathcal{U} \subseteq \mathcal{D}$, the pair $(\mathcal{U}, \mathcal{U}^{\perp})$ forms a t-structure on \mathcal{D} if and only if \mathcal{U} is co-reflective [KV88, 1.1 Proposition], i.e. the inclusion admits a right adjoint. A subcategory of \mathcal{D} is called an *aisle* (resp. a *co-aisle*) if it is suspended (resp. co-suspended) and co-reflective (resp. reflective, i.e. the inclusion admits a left adjoint). Since the pair $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$ is a t-structure on \mathcal{D} precisely when $\mathcal{D}^{\leq 0}$ is an aisle in \mathcal{D} (or, equivalently $\mathcal{D}^{>0}$ is a co-aisle), we say that $\mathcal{D}^{\leq 0}$ is the *aisle of the t-structure* (and that $\mathcal{D}^{>0}$ is the *co-aisle*).

We may hence define a truncation functor $\sigma^{\leq 0} \colon \mathcal{D} \to \mathcal{D}^{\leq 0}$ (resp. $\sigma^{>0} \colon \mathcal{D} \to \mathcal{D}^{>0}$) as the right adjoint of the inclusion functor $\iota^{\leq 0} \colon \mathcal{D}^{\leq 0} \to \mathcal{D}$ (resp. as the left adjoint of the inclusion functor $\iota^{>0} \colon \mathcal{D}^{>0} \to \mathcal{D}$).

Remark 1.3. It is conventional to denote the truncation functors by $\tau^{\leq 0}$ and $\tau^{>0}$. We have chosen to use $\sigma^{\leq 0}$ and $\sigma^{>0}$ instead, so that they will not be confused with the Auslander–Reiten translation.

For all objects $X \in \mathcal{D}$, there is a unique morphism $\sigma^{>0}X \xrightarrow{\partial_X} \Sigma \sigma^{\leq 0}X$ such that we have a triangle $\sigma^{\leq 0}X \xrightarrow{\varepsilon_X} X \xrightarrow{\eta_X} \sigma^{>0}X \xrightarrow{\partial_X} \Sigma \sigma^{\leq 0}X$,

where ε_X (resp. η_X) is induced by the co-unit of the adjunction $(\iota^{\leq 0}, \sigma^{\leq 0})$ (resp. by the unit of the adjunction $(\sigma^{>0}, \iota^{>0})$). Such triangles will be referred to as *truncation triangles*.

For any integer *i*, we define the cohomology functor in degree *i* by $H^i \stackrel{\text{def}}{=} \sigma^{\leq i} \sigma^{>i-1}$.

Theorem 1.4 ([BBD82, Théorème 1.3.6 and Remarque 3.1.17(ii)]). Let \mathcal{D} be a triangulated category with a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$.

- (1) The functor $H^0 = \sigma^{\leq 0} \sigma^{>-1} \colon \mathcal{D} \longrightarrow \mathcal{D}^0$ is cohomologial, i.e. it sends distinguished triangles to long exact sequences.
- (2) A complex

 $0 \longrightarrow U \stackrel{i}{\longrightarrow} V \stackrel{p}{\longrightarrow} W \longrightarrow 0$

in \mathcal{D}^0 is an exact sequence if and only if there exists a morphism $W \xrightarrow{\partial} \Sigma U$ in \mathcal{D} such that

 $U \xrightarrow{i} V \xrightarrow{p} W \xrightarrow{\partial} \Sigma U$

is a distinguished triangle in \mathcal{D} .

Applying Theorem 1.4(1) to the t-structure $(\mathcal{D}^{\leq i}, \mathcal{D}^{>i})$ shows that the cohomology functors

$$H^{i} \stackrel{\text{def}}{=} \sigma^{\leq i} \sigma^{>i-1} \colon \mathcal{D} \longrightarrow \mathcal{D}^{i}$$

are cohomological.

T-exactness, which we define presently, is a term used for triangle functors that preserve t-structures.

Definition 1.5 ([BBD82, Definition 1.3.16]). Let \mathcal{D} be a triangulated category with a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$, and let \mathcal{U} be a triangulated category with a t-structure $(\mathcal{U}^{\leq 0}, \mathcal{U}^{>0})$. A triangle functor $F: \mathcal{U} \longrightarrow \mathcal{D}$ is

- (1) left t-exact if $F(\mathcal{U}^{>0}) \subseteq \mathcal{D}^{>0}$,
- (2) right t-exact if $F(\mathcal{U}^{\leq 0}) \subseteq \mathcal{D}^{\leq 0}$,
- (3) *t-exact* if it is both left t-exact and right t-exact.

A *t-exact subcategory* of \mathcal{D} with respect to the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$, is a triangulated subcategory \mathcal{S} of \mathcal{D} such that $\sigma^{\leq 0}X \in \mathcal{S}$ for any $X \in \mathcal{S}$.

Remark 1.6. We might as well have defined a t-exact subcategory to be a triangulated subcategory which is closed under any truncation functor of the form $\sigma^{\leq i}$ or $\sigma^{>i}$, not just $\sigma^{\leq 0}$. First of all, a t-exact subcategory $S \subseteq D$ is closed under truncation by $\sigma^{\leq m}$, for all $m \in \mathbb{Z}$, since we have assumed that S is closed under suspension and desuspension. It is also preserved by $\sigma^{>0}$ (and thus $\sigma^{>m}$, for all $m \in \mathbb{Z}$), which one proves by forming a truncation triangle

$$\sigma^{\leq 0}X \xrightarrow{\varepsilon_X} X \longrightarrow \sigma^{>0}X \longrightarrow \Sigma \sigma^{\leq 0}X,$$

where $X \in \mathcal{S}$, and then pointing out that $\sigma^{>0}X$ is a mapping cone of ε_X , a morphism in \mathcal{S} .

The t-exact subcategories of \mathcal{D} are precisely those on which we can induce a t-structure $(\mathcal{D}^{\leq 0} \cap \mathcal{S}, \mathcal{D}^{>0} \cap \mathcal{S})$ such that the inclusion functor $\mathcal{S} \longrightarrow \mathcal{D}$ is t-exact [BBD82, p. 38]. The heart of the induced t-structure is $\mathcal{D}^{0} \cap \mathcal{S}$, and the cohomological functor $\mathcal{S} \longrightarrow \mathcal{S}^{0}$ is naturally isomorphic to the restriction of $H^{0}: \mathcal{D} \longrightarrow \mathcal{D}^{0}$.

We include the following well-known result to emphasise that t-exactness induces exactness.

Proposition 1.7 ([BBD82, Proposition 1.3.17]). Let $F: \mathcal{U} \longrightarrow \mathcal{D}$ be a t-exact functor. Then the composite functor

$$\mathcal{U}^{0} \longrightarrow \mathcal{U} \xrightarrow{F} \mathcal{D} \xrightarrow{H^{0}} \mathcal{D}^{0}$$

is an exact functor between abelian categories. In particular, if S is a t-exact subcategory of D, then S^0 is an exact abelian subcategory of D^0 .

Exact abelian subcategories are often called *wide subcategories* (or *weak Serre subcategories*). Equivalently, it is a subcategory which is closed under kernels, cokernels and extensions. In particular, every idempotent endomorphism admits a kernel, which is to say that the wide subcategory in question is idempotent complete. It follows that it is closed under direct summands.

Lemma 1.8. Let \mathcal{D} be a triangulated category equipped with a bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$, and let S be a t-exact subcategory of \mathcal{D} . Then S is thick.

Proof. It is to be shown that S is closed under direct summands. Let $X, Y \in D$ be such that $X \oplus Y \in S$. Since $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$ is bounded, only finitely many integers i are such that $H^i(X \oplus Y)$ is non-zero. We proceed by induction on the number d of such integers i.

The case d = 0 concerns only zero objects, and is therefore trivial to establish. If d = 1, we have that $X \oplus Y \in S^i$ for some *i*. As S^i is equivalent to S^0 , we may assume without loss of generality that i = 0. Since S^0 is a wide subcategory of \mathcal{D}^0 , it follows that the direct summand X of $X \oplus Y$ is in S^0 , as desired. Assume that the statement is true whenever $d < \ell$ for some $\ell \geq 2$. If $d = \ell$, we have a triangle

$$X_1 \longrightarrow X \longrightarrow X_2 \longrightarrow \Sigma X_1,$$

where X_1 has non-zero cohomology in at most $\ell - 1$ positions, and X_2 has non-zero cohomology in at most one position. In the same vein, one constructs a triangle

$$Y_1 \longrightarrow Y \longrightarrow Y_2 \longrightarrow \Sigma Y_1$$

where Y_1 and Y_2 have the same properties. Then $X_1 \oplus Y_1$ and $X_2 \oplus Y_2$ are objects in S with cohomology in fewer than ℓ positions, whence $X_1, X_2 \in S$ by the induction hypothesis. As S is a triangulated subcategory of D, it follows that $X \in S$, since it is an extension of X_2 with X_1 .

Following Zhang–Cai [ZC17, Definition 2.3], we say that a thick subcategory S of \mathcal{D} is H^0 -stable if $H^0(X) \in S$ for all $X \in S$.

Lemma 1.9. Let \mathcal{D} be a triangulated category with a t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$. A thick subcategory of $\mathcal{S} \subseteq \mathcal{D}$ is H^0 -stable if it is t-exact. If the t-structure is bounded, the converse also holds.

Proof. We explained in Remark 1.6 that a t-exact subcategory S is closed under truncation by $\sigma^{\leq 0}$ and $\sigma^{>-1}$. Hence, it is closed under $H^0 = \sigma^{\leq 0} \sigma^{>-1}$, which is to be H^0 -stable.

Conversely, suppose that the t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$ is bounded and let \mathcal{S} be an H^0 -stable subcategory of \mathcal{D} . For any $X \in \mathcal{S}$ and any integer n we can form a truncation triangle

$$\sigma^{\leq n-1}X \longrightarrow \sigma^{\leq n}X \longrightarrow H^nX \longrightarrow \Sigma\sigma^{\leq n-1}X.$$

The object $H^n X = H^0(\Sigma^{-n}X)$ lies in S, and hence $\sigma^{\leq n-1}X$ belongs to S if and only if $\sigma^{\leq n}X$ does. By induction, it follows that $\sigma^{\leq 0}X \in S$ if and only if $\sigma^{\leq n}X \in S$, where n is an arbitrary integer. The boundedness of the t-structure ensures that $\sigma^{\leq n}X = X$ for sufficiently large n. We conclude that S is t-exact, as we have shown that $\sigma^{\leq 0}X \in S$ for any $X \in S$,

The set of t-exact subcategories of \mathcal{D} is closed under arbitrary intersections. It thus has the structure of a complete lattice. The join of a family $\{S_i\}_{i \in I}$ is the smallest t-exact subcategory containing all S_i , whereas the meet is computed by intersection. We denote this lattice by t-exact(\mathcal{D}), and we let wide(\mathcal{D}^0) be the lattice of wide subcategories in \mathcal{D}^0 . We conclude this subsection with a result providing an explicit isomorphism of these lattices.

Theorem 1.10 ([ZC17, Theorem 2.5]). Let \mathcal{D} be a triangulated category with a bounded t-structure $(\mathcal{D}^{\leq 0}, \mathcal{D}^{>0})$. The cohomology functor $H^0: \mathcal{D} \longrightarrow \mathcal{D}^0$ induces an isomorphism of lattices

$$H^0: \operatorname{t-exact}(\mathcal{D}) \longrightarrow \operatorname{wide}(\mathcal{D}^0)$$

with inverse $\mathcal{D}_{(-)}$: wide $(\mathcal{D}^0) \rightarrow \text{t-exact}(\mathcal{D})$ sending a wide subcategory $\mathcal{W} \subseteq \mathcal{D}^0$ to the t-exact subcategory

$$\mathcal{D}_{\mathcal{W}} \coloneqq \{ X \in \mathcal{D} \mid H^n X \in \mathcal{W} \text{ for all } n \in \mathbb{Z} \} = \text{thick}_{\mathcal{D}}(\mathcal{W})$$

of \mathcal{D} .

Note that $\mathcal{D}_{\mathcal{W}}$ clearly is H^0 -stable. By Lemma 1.9, it is indeed a t-exact subcategory of \mathcal{D} . It is also possible to express the map H^0 : t-exact $(\mathcal{D}) \longrightarrow \text{wide}(\mathcal{D}^0)$ by intersecting t-exact subcategories with \mathcal{D}^0 .

1.2. Silting and τ -tilting.

Definition 1.11. Let C be a triangulated category. An object $X \in C$ is *presilting* if $C(X, \Sigma^n X) = 0$ for all positive integers n. A presilting object X is a *silting object* if, in addition, thick(X) = C.

We identify two presilting objects X and Y if $\operatorname{add}(X) = \operatorname{add}(Y)$. The sets of presilting and silting objects in \mathcal{C} are denoted by $\operatorname{presilt}(\mathcal{C})$ and $\operatorname{silt}(\mathcal{C})$, respectively. The subsets containing the objects having a fixed presilting object P as a direct summand are denoted by $\operatorname{presilt}_P(\mathcal{C})$ and $\operatorname{silt}_P(\mathcal{C})$, respectively.

Definition 1.12. For a fixed silting object $S \in C$, we say that an object $P \in C$ is 2_S -term it is belongs to $\operatorname{add}(S) * \Sigma \operatorname{add}(S)$, i.e if there exists a triangle

$$S^1 \longrightarrow S^0 \longrightarrow P \longrightarrow \Sigma S^1$$

where $S^1, S^0 \in \text{add}(S)$. The set of 2_S -term presilting objects is denoted by 2_S -presilt(\mathcal{C}), and the subset of 2_S -term silting objects by 2_S -silt(\mathcal{C}). The subsets of those having P is a direct summand are denoted by 2_S -presilt_P(\mathcal{C}) and 2_S -silt_P(\mathcal{C}), respectively.

If $\mathcal{C} = \mathcal{K}^{\mathrm{b}}(\mathrm{proj} A)$, the bounded homotopy category of a k-algebra A, then A is a silting object in \mathcal{C} . More generally, if A is a dg k-algebra which is non-positive (i.e. with trivial cohomology in positive degrees), then A is silting in per(A). In these cases, we write 2-silt(A) for the set of 2_A-term silting objects (and so on), omitting the subscript beneath the symbol 2.

In fact, if \mathcal{C} is a k-linear algebraic (i.e. triangle equivalent to the stable category of a Frobenius category) Hom-finite Krull–Schmidt triangulated category (which is to say that all Hom-spaces of the form $\mathcal{T}(X, X')$ are finite-dimensional for all $X, X' \in \mathcal{C}$, and any object $X \in \mathcal{C}$ is isomorphic to a finite direct sum $\bigoplus_{i=1}^{n} X_i$, where the endomorphism ring of each X_i is a local finite-dimensional k-algebra) with a silting object S, there exists a non-positive dg k-algebra A and a triangle equivalence $\mathcal{C} \longrightarrow \text{per}(A)$, sending S to A [KY14, Lemma 4.1(b)].

If C is a Krull–Schmidt triangulated k-linear category, an object $X \in C$ is *basic* if no fixed decomposition of X has a pair of distinct indecomposable direct summands that are isomorphic.

The set $\operatorname{silt}(\mathcal{C})$ admits a partial order \geq , where $P \geq Q$ provided that $\mathcal{C}(P, \Sigma^n Q) = 0$ whenever n > 0[AI12, Theorem 2.11 and Proposition 2.14]. Of particular interest is the subposet 2_S -silt $(\mathcal{C}) \subseteq \operatorname{silt}(\mathcal{C})$.

If \mathcal{C} is Hom-finite and Krull–Schmidt, it turns out that any 2_S -term presilting object $P \in \mathcal{C}$ can be completed to a 2_S -term silting object. One such silting object is the *Bongartz completion* $T_P^+ \stackrel{\text{def}}{=} P \oplus Q$ of P, where Q is defined by the triangle

$$S \longrightarrow Q \longrightarrow P_0 \xrightarrow{\beta_{\Sigma S}} \Sigma S,$$
 (1.A)

in which $\beta_{\Sigma S}$ is a minimal right add(P)-approximation of ΣS [IJY14]. Consequently, a 2_S-term presilting object is precisely the same thing as a direct summand of a 2_S-term silting object. If |P| denotes the number of non-isomorphic indecomposable direct summands of a 2_S-term presilting object P, we have that |P| is silting if and only if |P| = |S| [IJY14, Lemma 4.3]. The Bongartz completion is the maximal silting object (with respect to the partial order \geq defined above) in which P is a direct summand. There is a close connection between the 2_S -term silting theory of \mathcal{C} and the support τ -tilting theory of the finite-dimensional algebra $\operatorname{End}_{\mathcal{C}}(S)$. We recall the basic definitions and results of τ -tilting theory [AIR14].

Definition 1.13. For a finite-dimensional algebra A, a finitely generated right A-module M is τ -rigid if $\operatorname{Hom}_A(M, \tau M) = 0$, where τ is the Auslander–Reiten translation. A pair (M, Q) is support τ -rigid if M is a τ -rigid module and Q is a finitely generated projective right A-module such that $\operatorname{Hom}_A(Q, M) = 0$. A support τ -rigid pair (M, Q) is support τ -tilting |M| + |Q| = |A|, where |X| is the number of non-isomorphic indecomposable direct summands of X. An A-module M is support τ -tilting if there exists a projective A-module P such that (M, Q) is a τ -tilting pair.

A support τ -rigid pair (N, R) is a direct summand of the support τ -rigid pair (M, Q) if N is a direct summand of M and R is a direct summand of Q. We denote the set of support τ -rigid pairs in mod A by $s\tau$ -rigid pair(A), and those having a support τ -rigid pair (M, Q) as a direct summand by $s\tau$ -rigid pair $_{(M,Q)}(A)$. The subsets of support τ -tilting pairs are denoted by $s\tau$ -tilt pair(A) and $s\tau$ -tilt pair $_{(M,Q)}(A)$, respectively. We say that an algebra is τ -tilting finite if it has finitely many basic τ -tilting modules up to isomorphism.

If (M, Q) and (M, R) are support τ -tilting pairs, then $\operatorname{add}(Q) = \operatorname{add}(R)$ [AIR14, Proposition 2.3(b)]. In other words, we have a correspondence between basic support τ -tilting modules and basic support τ -tilting pairs, which truncates a pair (M, Q) to M.

The set of support τ -tilting A-modules is inextricably linked with functorially finite torsion classes of mod A. Recall that a *torsion class* (resp. *torsion-free class*) of mod A is a subcategory which is closed under factor modules (resp. submodules) and extensions. If \mathcal{G} is a torsion class, the right perpendicular category \mathcal{G}^{\perp} is a torsion-free class. Dually, the left perpendicular category of a torsion-free class is a torsion class. A *torsion pair* is a pair of subcategories $(\mathcal{G}, \mathcal{F})$ where $\mathcal{G}^{\perp} = \mathcal{F}$ and $^{\perp}\mathcal{F} = \mathcal{G}$. It is indeed the case that \mathcal{G} is a torsion class and that \mathcal{F} is torsion-free.

In the next theorem, and throughout, let gen(M) be the full subcategory of mod A containing the A-modules X such that there exists an epimorphism $M^{\oplus n} \twoheadrightarrow X$ for some $n \ge 1$.

Theorem 1.14 ([AIR14, Theorem 2.7]). Let M be a support τ -tilting right A-module. Then gen(M) is a functorially finite torsion class and we have a bijection

gen:
$$s\tau$$
-tilt $(A) \longrightarrow f$ -tors (A)

from the set of support τ -tilting A-modules to the set of functorially finite torsion classes in mod A.

Since the set of torsion classes is partially ordered under inclusion, this bijection gives a partial order on $s\tau$ -tilt(A). More explicitly, we impose that $M \ge N$ if $gen(M) \supseteq gen(N)$. If (M, Q) and (N, R) are support τ -tilting pairs, we say that $(M, Q) \ge (N, R)$ if $M \ge N$ as support τ -tilting modules.

Theorem 1.15 ([Jas15, Proposition 4.5], [IY08, Proposition 6.2(3)]). Let C be Hom-finite Krull–Schmidt triangulated category and $S \in C$ be a silting object. Let $\operatorname{add}(S) * \Sigma \operatorname{add}(S)$ be the subcategory of 2_S -term objects in C. The functor

$$\mathcal{C}(S,-)\colon \mathcal{C} \longrightarrow \operatorname{mod} \operatorname{End}_{\mathcal{C}}(S)$$

induces an equivalence of categories

$$\mathcal{C}(S,-) \colon \frac{\mathrm{add}(S) * \Sigma \mathrm{add}(S)}{[\Sigma S]} \longrightarrow \mathrm{mod} \, \mathrm{End}_{\mathcal{C}}(S),$$

where $[\Sigma S]$ is the ideal of morphisms factoring though ΣS .

The close connection between silting and support τ -tilting is expressed by the following theorem.

Theorem 1.16 ([IJY14, Theorem 4.5]). Let C and S be as in Theorem 1.15 and let $A = \text{End}_{C}(S)$. We have a bijection

where ΣX_1 is the maximal direct summand of X in $add(\Sigma S)$. It restricts to bijections

$$2_S$$
-presilt_P(\mathcal{C}) $\xrightarrow{H_S}$ s τ -rigid pair_{H_S(P)}(A),

and

$$2_{S}\text{-silt}_{P}(\mathcal{C}) \xrightarrow{H_{S}} s\tau\text{-tilt}\operatorname{pair}_{H_{S}(P)}(A), \qquad (1.B)$$

for each $P \in 2_S$ -presilt(C). The last bijection is an isomorphism of partially ordered sets.

Recall that $\mathcal{C}(S, -)$ is naturally isomorphic to the cohomology functor H^0 if $\mathcal{C} = \mathcal{K}^{\mathrm{b}}(\mathrm{proj}(A))$ (resp. $\mathrm{per}(A)$) and S = A for some k-algebra (resp. dg k-algebra) A.

The isomorphism (1.B) sends the Bongartz completion of P to a maximal object of $s\tau$ -tilt pair_{$H_S(P)$}(A). We refer to this as the *Bongartz completion* of the support τ -rigid pair $H_S(P)$. A module-theoretic construction of the Bongartz completion of support τ -rigid pairs is available [AIR14, Theorem 2.10], [DIR⁺17, Theorem 4.4]. We often denote the Bongartz completion of (M, Q) by (M^+, Q) , noting that the second component Q remains unaltered.

Proposition-Definition 1.17 ([Jas15, Proposition 3.6], [DIR⁺17, Theorem 4.12(a)]). Let A be a finitedimensional algebra, and let (M, Q) be a support τ -rigid pair in mod A. The pair (M, Q) determines a wide subcategory

$$J(M,Q) \stackrel{def}{=} M^{\perp} \cap {}^{\perp}(\tau M) \cap Q^{\perp} \subseteq \operatorname{mod} A$$

called the τ -perpendicular category of (M, Q).

Definition 1.18. Let J(M, Q) be the τ -perpendicular category of a support τ -rigid pair (M, Q) with Bongartz completion (M^+, Q) , and let $C = \text{End}_A(M^+)/[M]$, where [M] is the ideal of morphisms factoring through add(M). A support τ -rigid pair in J(M, Q) is a pair (U, R) in J(M, Q) such that $(\text{Hom}(M^+, U), \text{Hom}(M^+, R))$ is a support τ -rigid pair in mod C.

Theorem 1.19 ([DIR⁺17, Theorem 4.18], [Jas15, Theorem 3.8]). If A is τ -tilting finite, all wide subcategories of mod A are τ -perpendicular categories

Furthermore, τ -perpendicular subcategories are module categories, as the next result shows.

Theorem 1.20 ([Jas15, Theorem 3.8],[DIR⁺17, Theorem 4.12(b)]). Let (M, Q) be a support τ -rigid pair in mod A, let (M^+, Q) be the Bongartz completion of (M, Q), and let $C = \text{End}_A(M^+)/[M]$. We then have an exact equivalence

$$F_{(M,Q)} \stackrel{def}{=} \operatorname{Hom}_A(M^+, -) \colon J(M,Q) \longrightarrow \operatorname{mod} C.$$

Letting $B = \text{End}_A(M^+)$, we may express a quasi-inverse as the restriction of the tensor functor

$$G_{(M,Q)} \stackrel{def}{=} - \otimes_B M^+ \colon \operatorname{mod} B \longrightarrow \operatorname{mod} A$$

 $to \mod C$.

We will refer to the algebra C in the last theorem as the τ -tilting reduction of A with respect to (M, Q).

2. Silting t-structures and reduction

Silting reduction is a well-developed reduction technique, particularly for Hom-finite Krull–Schmidt triangulated categories [IY18]. The bijection in question restricts to two-term objects and is compatible with τ -tilting reduction [Jas15, Theorem 4.12]. In this section, we interpret silting reduction in terms of the t-structures they generate. Iyama–Yang prove the following theorem, though in greater generality.

Theorem 2.1 ([IY18, Theorems 3.1, 3.6, and 3.7], [Jas15, Proposition 4.11]). Let C be a Hom-finite Krull–Schmidt triangulated category containing a silting object S. Let P be a presilting object.

(1) Let $\mathcal{Z}_P = {}^{\perp_{>0}}P \cap P^{\perp_{>0}}$. Then the composite

$$\mathcal{Z}_P \longrightarrow \mathcal{C} \xrightarrow{\operatorname{loc}} \mathcal{C}/\operatorname{thick}(P)$$

induces a triangle equivalence $\mathcal{Z}_P/[P] \longrightarrow \mathcal{C}/\operatorname{thick}(P)$, where the triangulation on $\mathcal{Z}_P/[P]$ is described by Iyama-Yoshino [IY08, Theorem 4.2].

(2) The ideal quotient $\mathcal{Z}_P \longrightarrow \mathcal{Z}_P/[P]$ (and also the Verdier localisation $\mathcal{C} \longrightarrow \mathcal{C}/\operatorname{thick}(P)$) induces a bijection

$$\varphi_P : \operatorname{presilt}_P(\mathcal{C}) \longrightarrow \operatorname{presilt}(\mathcal{Z}_P/[P]) \cong \operatorname{presilt}(\mathcal{C}/\operatorname{thick}(P)).$$
 (2.A)

If P is 2_S -term with Bongartz completion T_P^+ , it restricts to a bijection

$$\varphi_P \colon 2_S \operatorname{-presilt}_P(\mathcal{C}) \longrightarrow 2_{T_P^+} \operatorname{-presilt}(\mathcal{Z}_P/[P]) \cong 2_{T_P^+} \operatorname{-presilt}(\mathcal{C}/\operatorname{thick}(P)).$$
 (2.B)

(3) The bijection (2.A) restricts to an isomorphism of posets

$$\varphi_P : \operatorname{silt}_P(\mathcal{C}) \longrightarrow \operatorname{silt}(\mathcal{Z}_P/[P]) \cong \operatorname{silt}(\mathcal{C}/\operatorname{thick}(P))$$

which, if P is 2_S -term with Bongartz completion T_P^+ , further restricts to an isomorphism

$$\varphi_P \colon 2_S \operatorname{-silt}_S(\mathcal{C}) \longrightarrow 2_{T_P^+} \operatorname{-silt}(\mathcal{Z}_P/[P]) \cong 2_{T_P^+} \operatorname{-silt}(\mathcal{C}/\operatorname{thick}(P)).$$
(2.C)

Since the bijection (2.A) is induced by the ideal quotient $Z_P \rightarrow Z_P/[P]$, our next two lemmas are immediate consequences.

Lemma 2.2. Let C is as in Theorem 2.1, and let P be a 2_S -term presilting object in C. We define ind_P 2_S -presilt_P(C) to be the objects in 2_S -presilt_P(C) of the form $P \oplus X$, where X is an indecomposable object which is not contained in add(P). Then the bijection (2.A) restricts to a bijection

$$\varphi_P \colon \operatorname{ind}_P 2_S \operatorname{-presilt}_P(\mathcal{C}) \longrightarrow \operatorname{ind} 2_{T_P^+} \operatorname{-presilt}(\mathcal{Z}_P/[P])$$

where the codomain is the set of indecomposable objects in $2_{T_P^+}$ -presilt $(\mathcal{Z}_P/[P])$.

Lemma 2.3. Let Q be a 2_S -term presilting object in C such that $P \oplus Q$ also is 2_S -term presilting. The bijection

$$\varphi_P: 2_S$$
-presilt $_P(\mathcal{C}) \longrightarrow 2_{T_P^+}$ -presilt $(\mathcal{Z}_P/[P])$

restricts to a bijection

$$\varphi_P \colon 2_S \operatorname{-presilt}_{P \oplus Q}(\mathcal{C}) \longrightarrow 2_{T_P^+} \operatorname{-presilt}_{\varphi_P(Q)}(\mathcal{Z}_P/[P])$$

Lemma 2.4. Let $P \oplus Q$ be a 2_S -term presilting object in C. Then the diagram

commutes.

Proof. By the Third Isomorphism Theorem, we have additive equivalences


$$\frac{\mathcal{Z}_{P\oplus Q}/[P]}{[Q]} \cong \frac{\mathcal{Z}_{P\oplus Q}/[P]}{[P\oplus Q]/[P]} \cong \frac{\mathcal{Z}_{P\oplus Q}}{[P\oplus Q]},\tag{2.E}$$

and similarly

$$\frac{\mathcal{Z}_{P\oplus Q}/[Q]}{[P]} \cong \frac{\mathcal{Z}_{P\oplus Q}}{[P\oplus Q]}.$$
(2.F)

To give an argument as to why $\varphi_{\varphi_Q(P)}$ (and similarly $\varphi_{\varphi_P(Q)}$) has the claimed codomain, it now suffices to show that $T^+_{P\oplus Q}$ is the Bongartz completion of $\varphi_Q(P)$ in $\mathcal{Z}_Q/[Q]$. This is a consequence of the formula $\varphi_Q T^+_X \cong T^+_{\varphi_Q(X)}$, due to Jasso [Jas15, Proposition 4.10(b)]; set $X = P \oplus Q$, and recall that φ_Q is induced by the ideal quotient $\mathcal{Z} \longrightarrow \mathcal{Z}/[P]$, which leaves the object X unaltered.

Further, the isomorphisms (2.E) and (2.F) enable the construction of an essentially commutative diagram of ideal quotient functors



Since these functors induce the bijections in (2.D), the proof is complete.

If C is as in Theorem 2.1, and is also assumed to be algebraic, we recall (from the discussion following Definition 1.12) that there exists a non-positive proper dg k-algebra A and a triangle equivalence $C \rightarrow \text{per}(A)$, sending S to A [KY14, Lemma 4.1(b)]. From this point and to the end of the article, this is the most general case we will consider. We state our setup below for future reference. Recall that a dg *k*-algebra is *non-positive* if it has no cohomology in positive degrees, and *proper* if its total cohomology is finite-dimensional.

Setup 2.5. Let A be a non-positive proper dg k-algebra, and let P be a two-term presilting object in per(A), i.e. an object in 2-presilt(A).

Note that finite-dimensional algebras are included in this setup. It is also important to emphasise that the regular module A is an object in $\mathcal{D}_{\mathrm{fd}}(A)$ (since A is proper).

It would be possible to generalise from Setup 2.5 to the setting of ST-triples $(\mathcal{C}, \mathcal{D}, S)$, where $\mathcal{C} \subseteq \mathcal{D}$. We do not give the definition of ST-triples here [AMY19, §4], but we note that if \mathcal{C} is algebraic, it is equivalent to a triple of the form $(\text{per}(A), \mathcal{D}_{\text{fd}}(A), A)$, where A is as in Setup 2.5 [AMY19, Corollary 6.13]. The assumption of algebraic does not reduce the scope of applications at time of writing; ST-triples seem not to have occurred outside an algebraic context.

In this section, we describe Iyama–Yang silting reduction and its interaction with silting t-structures. Silting reduction with respect to P occurs in the Verdier quotient per(A)/thick(P), but we will show that perpendicular category $P^{\perp_{\mathbb{Z}}}$ also plays a role; if per(A)/thick(P) is triangle equivalent to per(C) for some dg algebra C, then $P^{\perp_{\mathbb{Z}}}$ is triangle equivalent to $\mathcal{D}_{fd}(C)$ (see Corollary 2.12). Note that $P^{\perp_{\mathbb{Z}}}$ is a thick subcategory of $\mathcal{D}_{fd}(A)$. The t-structures defined by these silting objects will be restricted to $P^{\perp_{\mathbb{Z}}}$, which is achievable since $P^{\perp_{\mathbb{Z}}}$ is t-exact (see Corollary 2.9).

We first specify what it means for a t-structure to be generated by a silting object.

Lemma 2.6 ([ATJLSS03, Lemma 5.3]). Let T be a silting object in per(A). Then the pair $(T^{\perp_{>0}}, T^{\perp_{\leq 0}})$ is a bounded t-structure on $\mathcal{D}_{fd}(A)$.

A t-structure of the form $(T^{\perp>0}, T^{\perp\leq 0})$, where T is silting object, is said to be a *silting t-structure* and it is *generated* by T. The silting object A in per(A), where A is a non-positive dg algebra, generates the standard t-structure on $\mathcal{D}_{\mathrm{fd}}(A)$.

Lemma 2.7 ([AMY19, Propostion 4.6(a)]). Let T be a silting object in per(A), and let H_T^0 be the cohomology functor associated to the t-structure $(T^{\perp_{>0}}, T^{\perp_{\leq 0}})$. Then H_T^0 sends T to a projective generator of the heart $T^{\perp_{\neq 0}}$, and the Hom-functor $\mathcal{D}_{\mathrm{fd}}(A)(T, -)$ restricts to an exact equivalence of abelian categories

$$\mathcal{D}_{\mathrm{fd}}(A)(T,-)\colon T^{\perp_{\neq 0}} \longrightarrow \mathrm{mod}\,\mathrm{End}_{\mathcal{D}_{\mathrm{fd}}(A)}(T).$$

We include a useful lemma.

Lemma 2.8. Let A and P be as in Setup 2.5 and let $(\mathcal{U}, \mathcal{U}^{\perp})$ be a t-structure on $\mathcal{D}_{\mathrm{fd}}(A)$ such that $P \in \mathcal{U}$ and $\mathcal{U} \subseteq P^{\perp_{>1}}$. Then the subcategory $P^{\perp_{\mathbb{Z}}}$ is t-exact with respect to $(\mathcal{U}, \mathcal{U}^{\perp})$. In particular, the pair $(\mathcal{U} \cap P^{\perp_{\mathbb{Z}}}, \mathcal{U}^{\perp} \cap P^{\perp_{\mathbb{Z}}})$ is a t-structure on $P^{\perp_{\mathbb{Z}}}$.

Proof. Let $\sigma_{\mathcal{U}}$ and $\sigma_{\mathcal{U}^{\perp}}$ be the truncation functors for $(\mathcal{U}, \mathcal{U}^{\perp})$. Fixing an arbitrary object $X \in P^{\perp_{\mathbb{Z}}}$, it suffices to show that $\sigma_{\mathcal{U}}X \in P^{\perp_{\mathbb{Z}}}$. We have a truncation triangle

$$\sigma_{\mathcal{U}}X \longrightarrow X \longrightarrow \sigma_{\mathcal{U}^{\perp}}X \longrightarrow \Sigma \sigma_{\mathcal{U}}X.$$

Since $\sigma_{\mathcal{U}^{\perp}} X \in \mathcal{U}^{\perp} \subseteq P^{\perp_{\leq 0}}$ (the last inclusion holding because $P \in \mathcal{U}$ and \mathcal{U} being closed under suspension), the desuspension $\Sigma^{-1}\sigma_{\mathcal{U}^{\perp}} X$ is in $P^{\perp_{\leq 1}}$. Using that $P^{\perp_{\leq 1}}$ is closed under extensions, we deduce that $\sigma_{\mathcal{U}} X \in P^{\perp_{\leq 1}}$. As $\sigma_{\mathcal{U}} X$ also belongs to \mathcal{U} , and thus to $P^{\perp_{>1}}$, it follows that $\sigma_{\mathcal{U}} X \in P^{\perp_{\leq 1}} \cap P^{\perp_{>1}} = P^{\perp_{\mathbb{Z}}}$, as desired.

Lemma 2.8 applies in two very important cases pertaining to our setup.

Corollary 2.9. Let A be as in Setup 2.5.

- (1) Let T be a silting object in $\mathcal{D}_{\mathrm{fd}}(A)$ of which Q is a direct summand. Then $Q^{\perp_{\mathbb{Z}}}$ is t-exact with respect to the t-structure $(T^{\perp_{>0}}, T^{\perp_{\leq 0}})$ on $\mathcal{D}_{\mathrm{fd}}(A)$.
- (2) Let $P \in 2$ -presilt(A) (as in Setup 2.5). Then $P^{\perp_{\mathbb{Z}}}$ is t-exact with respect to the standard t-structure $(A^{\perp_{>0}}, A^{\perp_{\leq 0}})$ on $\mathcal{D}_{\mathrm{fd}}(A)$.

Proof. We will prove that both assertions are consequences of Lemma 2.8.

We first address (1). It is indeed the case that

$$Q \in T^{\perp_{>0}} \subseteq Q^{\perp_{>0}} \subseteq Q^{\perp_{>1}}$$

The first and penultimate inclusions hold whenever Q is a direct summand of T.

To prove (2), we simply have to point out that $P \in A^{\perp_{>0}}$ and $A^{\perp_{>0}} \subseteq P^{\perp_{>1}}$ both hold since P is two-term.

We claimed in the introduction that the perpendicular category $P^{\perp_{\mathbb{Z}}}$ plays that role of the τ -perpendicular category. This assertion will now be made explicit.

Proposition 2.10. Let A and P be as in Setup 2.5. Then the isomorphism of lattices

$$\mathcal{D}_{\rm fd}(A)(A,-) = H^0: \text{ t-exact}(\mathcal{D}_{\rm fd}(A)) \longrightarrow \text{ wide}(\text{mod } H^0A), \tag{2.G}$$

provided by Theorem 1.10, sends $P^{\perp_{\mathbb{Z}}}$ to J(M,Q).

Proof. The bijection (2.G) can be expressed as $- \cap A^{\perp \neq 0}$, and its inverse by thick_{$\mathcal{D}_{fd}(A)$}(-). Since these are mutually inverse isomorphism of posets, it suffices to prove that

$$P^{\perp_{\mathbb{Z}}} \cap A^{\perp_{\neq 0}} \subseteq J(M,Q) \tag{2.H}$$

$$\operatorname{thick}_{\mathcal{D}_{\mathrm{fd}}(A)}(J(M,Q)) \subseteq P^{\perp_{\mathbb{Z}}}$$

$$(2.1)$$

Write P as $P_M \oplus \Sigma Q$, where P_M is a minimal projective presentation of M. The perpendicular category $P^{\perp_{\mathbb{Z}}}$ can then be expressed as the intersection $P_M^{\perp_{\mathbb{Z}}} \cap Q^{\perp_{\mathbb{Z}}}$.

The inclusion (2.H) will be addressed first. Let $X \in P^{\perp_{\mathbb{Z}}} \cap A^{\perp_{\neq 0}}$. It is to be deduced that $X \in M^{\perp_{0}} \cap {}^{\perp_{0}}(\tau M) \cap Q^{\perp_{0}}$. The assertion that $X \in Q^{\perp_{0}}$ is immediate, having assumed that $X \in P^{\perp_{\mathbb{Z}}} \subseteq Q^{\perp_{\mathbb{Z}}}$. Consider the following truncation triangle in the standard t-structure

$$\sigma^{<0}P_M \longrightarrow P_M \longrightarrow M \longrightarrow \Sigma \sigma^{<0}P_M. \tag{2.J}$$

ERLEND D. BØRVE

As both $\sigma^{<0}P_M$ and $\Sigma\sigma^{<0}P_M$ are in $\mathcal{D}_{\mathrm{fd}}^{<0}(A)$, there are no maps from these objects to $X \in A^{\perp_{\neq 0}} \subseteq \mathcal{D}_{\mathrm{fd}}^{\geq 0}(A)$. Having assumed that $\mathcal{D}_{\mathrm{fd}}(A)(P_M, X) = 0$, it follows from a long exact sequence argument that $X \in M^{\perp_0}$. Lastly, consider the triangle

$$X_{<1} \longrightarrow X \longrightarrow X_{\geq 1} \longrightarrow \Sigma X_{<1}$$

$$(2.K)$$

where $X_{\geq 1}$ is the projective presentation of X, and $X_{<1}$ is what remains of its projective resolution. Since the Hom-spaces $\mathcal{D}_{\rm fd}(A)(P_M, \Sigma X) = 0$ and $\mathcal{D}_{\rm fd}(A)(P_M, \Sigma^2 X_{<1}) = 0$ vanish (the former by assumption, the latter since P_M is two-term), so does $\mathcal{D}_{\rm fd}(A)(P, \Sigma X_{\geq 1}) = 0$. Now, since the vanishing of $\mathcal{D}_{\rm fd}(A)(P, \Sigma X_{\geq 1})$ is equivalent to the assertion that $\operatorname{Hom}_A(X, \tau M) = 0$ [AIR14, Lemma 3.4], we are in a position to conclude that the inclusion (2.H) holds.

To prove (2.I), it suffices to show that $J(M,Q) \subseteq P^{\perp_{\mathbb{Z}}}$, since $P^{\perp_{\mathbb{Z}}}$ is thick. As $J(M,Q) \subseteq A^{\perp_{\neq 0}}$ and P is two-term, we can immediately assert that $J(M,Q) \subseteq P^{\perp_{\neq 0,1}}$. Fix an X in J(M,Q). We will proceed by mostly reversing the arguments in the last paragraph. To prove that $X \in P_M^{\perp_0}$, apply the contravariant Hom-functor $\mathcal{D}_{\mathrm{fd}}(A)(-,X)$ to the triangle (2.J) and use a long exact sequence argument. To show that $J(M,Q) \subseteq P_M^{\perp_1}$, apply the covariant Hom-functor $\mathcal{D}_{\mathrm{fd}}(A)(\Sigma^{-1}P_M,-)$ to (2.K). Finally, we have assumed that $X \in Q^{\perp_0}$, and the assertion that $X \in Q^{\perp_{\neq 0}}$ follows from the fact that X has no cohomology in non-zero degrees (see Lemma 2.7). We have shown that $X \in P^{\perp_Z}$, completing the proof.

Section 4 is devoted to the construction of the τ -cluster morphism category in terms of two-term silting. We conclude this section with results that will turn out useful for this purpose. To prove them, we will apply some basic results from the localisation theory of compactly generated triangulated categories, especially for the full derived category $\mathcal{D}(A)$, where the objects are possibly unbounded complexes of possibly infinitely generated modules. Proposition 2.11 provides references to the localisation theory we need, whereas the subsequent corollaries will be applied in later sections.

Proposition 2.11. Let A and P be as in Setup 2.5, let $P^{\perp_{\mathbb{Z}}}$ be the perpendicular category of P in the full derived category $\mathcal{D}(A)$, and let Loc(P) be the smallest thick subcategory of $\mathcal{D}(A)$ that is closed under set-indexed coproducts.

(1) [Kra08, Theorem 5.6.1] There is a recollement of triangulated categories

 $P^{\perp_{\mathbb{Z}}} \xrightarrow{\iota_{P}} \mathcal{D}(A) \xrightarrow{\rho_{P}} \operatorname{Loc}(P)$ (2.L)

i.e. the diagram displays four adjoint pair of triangle functors, the composite $\pi_P \lambda_P$ is zero, all functors into $\mathcal{D}(A)$ are fully faithful, and for all $X \in \mathcal{D}(A)$ we have triangles

$$\lambda_P \rho_P X \xrightarrow{\varepsilon} X \xrightarrow{\eta} \iota_P \pi_P X \longrightarrow \Sigma(\rho_P \lambda_P X),$$

 $\iota_P \kappa_P X \xrightarrow{\varepsilon'} X \xrightarrow{\eta'} \mu_P \rho_P X \longrightarrow \Sigma(\iota_P \kappa_P X),$

where η and η' (resp. ε and ε') are units (resp. co-units) of the adjunctions.

16

- (2) [Kel94, §4.3] We have a triangle equivalence $P^{\perp_{\mathbb{Z}}} \simeq \mathcal{D}(C_P)$, where C_P is the derived endomorphism algebra of $\pi_P A$ in $P^{\perp_{\mathbb{Z}}}$.
- (3) The functor π_P restricts to a functor $\operatorname{per}(A) \longrightarrow \operatorname{per}(C_P)$, which is a Verdier localisation functor with respect to thick(P). In particular, we have a triangle equivalence

$$\operatorname{per}(A)/\operatorname{thick}(P) \simeq \operatorname{per}(C_P).$$

(4) The functor $\mathcal{D}(C_P) \simeq P^{\perp_{\mathbb{Z}}} \longrightarrow \mathcal{D}(A)$ preserves boundedness, i.e. it restricts to an embedding $\mathcal{D}_{\mathrm{fd}}(C_P) \hookrightarrow \mathcal{D}_{\mathrm{fd}}(A).$

Proof. The assertions in (1) and (2) are taken directly from the cited references.

The assertion in (3) can be deduced from a result of Neeman [Nee92, Theorem 2.1], noting that $P^{\perp_{\mathbb{Z}}}$ is idempotent complete.

Since π_P restricts to a functor

$$\pi_P \colon \operatorname{per}(A) \longrightarrow \operatorname{per}(C_P)$$

the claim in (4) follows from a result of Gao–Psaroudakis [GP18, Lemma 4.2].

Having temporarily used the notation $P^{\perp_{\mathbb{Z}}}$ to denote the perpendicular category of P inside $\mathcal{D}(A)$, we now revert to the convention where $P^{\perp_{\mathbb{Z}}}$ is a subcategory of $\mathcal{D}_{\mathrm{fd}}(A)$.

Corollary 2.12. Let A and P be as in Setup 2.5, and let $(M,Q) = H_A(P)$ be the support τ -rigid Amodule corresponding to P. Let C_P as in Proposition 2.11(2). Then the perpendicular category $P^{\perp_{\mathbb{Z}}}$ in $\mathcal{D}_{\mathrm{fd}}(A)$ is triangle equivalent to $\mathcal{D}_{\mathrm{fd}}(C_P)$.

Proof. This follows directly from Proposition 2.11(4).

Corollary 2.13. Let A and P be as in Setup 2.5. The restriction of the functor π_P (from Proposition 2.11(1)) to per(A) fits into a commutative diagram

$$per(A) \xrightarrow{\pi_P} P^{\perp_{\mathbb{Z}}} \subseteq \mathcal{D}_{fd}(A)$$

$$(2.M)$$

$$per(A)/\operatorname{thick}(P)$$

where q is the Verdier localisation and i is fully faithful.

Proof. By Corollary 2.12, we have that $P^{\perp_{\mathbb{Z}}} \simeq \mathcal{D}_{\mathrm{fd}}(C_P)$ for some non-positive proper dg algebra C_P such that $\mathrm{per}(C_P) \simeq \mathrm{per}(A)/\mathrm{thick}(P)$. As C_P is proper, the perfect derived category $\mathrm{per}(C_P)$ is contained in $\mathcal{D}_{\mathrm{fd}}(C_P)$. Hence, the restriction of π_P to $\mathrm{per}(A)$ factors as follows:

$$\operatorname{per}(A) \xrightarrow{q} \operatorname{per}(A) / \operatorname{thick}(P) \simeq \operatorname{per}(C_P) \hookrightarrow \mathcal{D}_{\mathrm{fd}}(C_P) \simeq P^{\perp_{\mathbb{Z}}}$$

as desired.

Corollary 2.14. Let A and P be as in Setup 2.5. For any $X \in per(A)$, there is a triangle

 $X_P \longrightarrow X \longrightarrow \pi_P X \longrightarrow \Sigma X_P$

where $X_P \in \text{Loc}(P)$.

Proof. By Proposition 2.11(1), we have a triangle

$$\lambda_P \rho_P X \xrightarrow{\varepsilon} X \xrightarrow{\eta} \iota_P \pi_P X \longrightarrow \Sigma(\lambda_P \rho_P X),$$

where η is the unit of the adjunction $\pi_P \dashv \iota_P$ and ε is the co-unit of $\lambda_P \dashv \rho_P$. Since λ_P and ι_P are the obvious inclusions, we can put $X_P = \rho_P X$ and conclude.

3. Compatibility of the Iyama-Yang and Buan-Marsh bijections

Reduction techniques have been developed for both silting objects (known as *siling reduction* [IY18]) and support τ -tilting modules (τ -tilting reduction [Jas15]). They are compatible [Jas15, Theorem 4.12(b)]. In this section we generalise Jasso's compatibility theorem, by showing that the support τ -tilting reduction of Buan–Marsh [BM21a, Section 3] is compatible with silting reduction.

Buan–Marsh prove the following.

Theorem 3.1 ([BM21a, Theorem 3.6]). Let A be a finite-dimensional algebra, and let (M, Q) be a support τ -rigid pair in mod A. There is a bijection

$$\mathcal{E}_{(M,Q)}: \text{ } \text{s}\tau\text{-rigid } \text{pair}_{(M,Q)}(A) \longrightarrow \text{s}\tau\text{-rigid } \text{pair}(J(M,Q)),$$

where J(M,Q) is the τ -perpendicular category of (M,Q).

We will now recall how the map $\mathcal{E}_{(M,Q)}$ is constructed. Buan–Marsh first address the cases where (M,Q) is either of the form (M,0) or (0,Q). In each of the five cases below, they define a map of indecomposable objects

$$\mathcal{E}_{(M,Q)}: \text{ ind } s\tau\text{-rigid pair}_{(M,Q)}(A) \longrightarrow \text{ ind } s\tau\text{-rigid pair}(J(M,Q)),$$

which is extended to the bijection in Theorem 3.1 in the obvious way.

For a τ -rigid A-module X, let P_X denote its minimal projective presentation, considered as a two-term presilting object in $\mathcal{K}^{\mathrm{b}}(\mathrm{proj}(A))$. In abstract terms, one may set $P_X \stackrel{\mathrm{def}}{=} H_A^{-1}(X,0)$, where H_A was defined in Theorem 1.16.

Case I: Suppose that Q = 0.

Case I(a): If X is an indecomposable A-module such that $M \oplus X$ is τ -rigid and $X \notin \text{gen}(M)$, define $\mathcal{E}_{(M,0)}(X,0) \stackrel{\text{def}}{=} (f_M(X),0)$, where $f_M: \mod A \longrightarrow M^{\perp}$ is the torsion-free functor for the torsion pair $(\text{gen}(M), M^{\perp})$, that is, the natural functor $\mod A \longrightarrow M^{\perp}$.

Case I(b): If X is an indecomposable module such that $M \oplus X$ is τ -rigid and $X \in \text{gen}(M)$, define $\mathcal{E}_{(M,0)}(X,0) \stackrel{\text{def}}{=} (0, f_M(H^0 R_X))$, where

$$R_X \longrightarrow (P_M)_X \xrightarrow{\beta_{P_X}} P_X \longrightarrow \Sigma R_X$$

is a distinguished triangle and β_{P_X} is a minimal right $\operatorname{add}(P_M)$ -approximation.

Case I(c): If R is an indecomposable projective such that $\text{Hom}_A(R, M) = 0$, define

$$\mathcal{E}_{(M,0)}(0,R) \stackrel{\text{def}}{=} (0, f_M(H^0 C_{\Sigma R})),$$

where

$$C_{\Sigma R} \longrightarrow (P_M)_{\Sigma R} \xrightarrow{\beta_{\Sigma R}} \Sigma R \longrightarrow \Sigma C_{\Sigma R}$$

is a distinguished triangle and $\beta_{\Sigma R}$ is a minimal right $\operatorname{add}(P_M)$ -approximation.

Case II: Suppose that M = 0.

Case II(a): If X is an indecomposable τ -rigid module such that $\operatorname{Hom}_A(Q, X) = 0$, define

$$\mathcal{E}_{(0,Q)}(X,0) \stackrel{\text{def}}{=} (X,0)$$

Case II(b): If R is an indecomposable projective module such that $\operatorname{add}(Q) \cap \operatorname{add}(R) = \{0\}$, define $\mathcal{E}_{(0,Q)}(0,R) \stackrel{\text{def}}{=} (0, f_Q(R)).$

In general: Let (M, Q) be a support τ -rigid pair in mod A. Let $(M^+, 0)$ be the Bongartz completion of (M, 0). By Theorem 1.20 we have an exact equivalence

$$F_{(M,0)} = \operatorname{Hom}_A(M^+, -) \colon J(M,0) \longrightarrow \operatorname{mod} C = \operatorname{mod} \operatorname{End}_A(M^+) / [M]$$

which induces a bijection

$$s\tau$$
-rigid pair_(0,Q) $(J(M,0)) \longrightarrow s\tau$ -rigid pair_(0,Q') (C)

where Q' is the projective C-module $\operatorname{Hom}_A(M^+, Q)$. We define a map $\psi_{(M,0)}$ as the composite of this bijection with $\mathcal{E}_{(M,0)}$ (as seen in the upper triangle of (3.A) below). Then $\psi_{(M,0)}(0,Q) = (0,Q')$, which is a support τ -rigid pair in mod C. Case II above gives a bijection

$$\mathcal{E}^{C}_{(0,Q)}$$
: $s\tau$ -rigid pair $(C) \longrightarrow s\tau$ -rigid pair $J_{C}(0,Q')$,

where $J_C(0, Q')$ is the τ -perpendicular category of (0, Q') in mod C. Theorem 1.20 now gives another exact equivalence

$$J_C(0,Q') \xrightarrow{u} (Q')^{\perp} \xrightarrow{F_{(M,0)}^{-1}} M^{\perp} \cap {}^{\perp}(\tau M) \cap Q^{\perp} = J(M,Q).$$

$$i \cap mod \ C$$

We define $\mathcal{E}_{(M,Q)}(X,R) \stackrel{\text{def}}{=} u \circ \mathcal{E}^{C}_{(0,Q')} \circ \psi_{(M,0)}(X,R).$

Our aim is to link $\mathcal{E}_{(M,P)}$ to Iyama–Yang silting reduction. This is achievable in the following setup.

Setup 3.2. A Hom-finite Krull–Schmidt triangulated category C is fixed, as well as a silting object $S \in C$. Let P be a 2_S-term presilting object, and A the endomorphism algebra $\text{End}_{\mathcal{C}}(S)$.

In order to establish the compatibility of the two reduction techniques described above, we prove two important lemmas.

Lemma 3.3. Let C, S, P, and A be as in Setup 3.2. For all $X \in \operatorname{add}(S) * \Sigma \operatorname{add}(S)$, we have a natural isomorphism $\frac{\mathcal{C}}{[P]}(S,X) \cong f_{\mathcal{C}(S,P)}\mathcal{C}(S,X)$, as right A-modules, where $f_{\mathcal{C}(S,P)} \colon \operatorname{mod} A \longrightarrow \mathcal{C}(S,P)^{\perp}$ is the torsion-free functor for $(\operatorname{gen} \mathcal{C}(S,P), \mathcal{C}(S,P)^{\perp})$.

Proof. We have an exact sequence

$$[P](S,X) \xrightarrow{i} \mathcal{C}(S,X) \longrightarrow \frac{\mathcal{C}}{[P]}(S,X)$$
(3.B)

of A-modules, where [P](S, X) is the ideal of morphisms factoring through $\operatorname{add}(P)$. It suffices to show that $[P](S, X) \in \operatorname{gen}(\mathcal{C}(S, P))$ and that $\frac{\mathcal{C}}{|P|}(S, X) \in \mathcal{C}(S, P)^{\perp}$.

Let $P' \xrightarrow{\beta_X} X$ be a right $\operatorname{add}(P)$ -approximation of X. Using the definition of right approximations, one argues that the map

$$\mathcal{C}(S, P') \xrightarrow{\beta_X \circ -} [P](S, X)$$

surjects, whence $[P](S, X) \in \text{gen}(\mathcal{C}(S, P)).$

We now show that $\operatorname{Hom}_A(\mathcal{C}(S, P), \frac{\mathcal{C}}{[P]}(S, X)) = 0$. This k-vector space appears as the third term in the following long exact sequence of k-vector spaces:

$$0 \longrightarrow \operatorname{Hom}_{A}(\mathcal{C}(S, P), [P](S, X)) \xrightarrow{i \circ -} \operatorname{Hom}_{A}(\mathcal{C}(S, P), \mathcal{C}(S, X)) \longrightarrow \operatorname{Hom}_{A}(\mathcal{C}(S, P), \frac{\mathcal{C}}{[P]}(S, X))$$

 $\operatorname{Ext}_{A}^{1}(\mathcal{C}(S, P), [P](S, X)).$

Having already shown that $[P](S, X) \in \text{gen}(\mathcal{C}(S, P))$, a result of Auslander–Smalø [AS81, Proposition 5.8] asserts that the τ -rigidity of $\mathcal{C}(S, P)$ is equivalent to the vanishing of $\text{Ext}_A^1(\mathcal{C}(S, P), [P](S, X))$. Thus it suffices to show that $i \circ -$ surjects.

We will conclude the proof by showing that

$$\operatorname{Hom}_{A}(\mathcal{C}(S, P), \mathcal{C}(S, X)) = \operatorname{Hom}_{A}(\mathcal{C}(S, P), [P](S, X)),$$

which would make $i \circ -$ an injective endomorphism of a finite-dimensional vector space (thus also surjective). It follows from Theorem 1.15 that every A-homomorphisms from $\mathcal{C}(S, P)$ to $\mathcal{C}(S, X)$ is determined by an equivalence class of morphisms $P \longrightarrow X$. Consequently, all homomorphisms in $\text{Hom}_A(\mathcal{C}(S, P), \mathcal{C}(S, X))$ have image in [P](S, X), and $i \circ -$ surjects. \Box

Lemma 3.4. If C, S, and P are as in Setup 3.2, we have a bijection


$$\begin{array}{ccc} 2_{T_{P}^{+}}\text{-}\text{presilt}(\mathcal{Z}_{P}/[P]) & \xrightarrow{H_{S}} & \text{s}\tau\text{-}\text{rigid pair } J(H_{S}(P)) \\ & & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & Y & \longmapsto & \left(\begin{array}{c} \mathcal{C} \\ \overline{[P]}(S,Y), \begin{array}{c} \mathcal{C} \\ \overline{[P]}(S,Y_{1}) \end{array} \right) & \begin{array}{c} 3.3 \\ \cong \end{array} & \left(f_{\mathcal{C}(S,P)}\mathcal{C}(S,Y), f_{\mathcal{C}(S,P)}\mathcal{C}(S,Y_{1}) \right) \end{array} \right)$$

where $Y_1\langle 1 \rangle$ is the largest direct summand of Y in $\operatorname{add}(T_P^+)\langle 1 \rangle$. Here, the functor $\langle 1 \rangle$ is the suspension functor on the triangulation given to $\mathcal{Z}_P/[P]$ (one can compute it as the mapping cone of a left $\operatorname{add}(P)$ approximation of X). Moreover, if $F_{H_S(P)}$ is as in Theorem 1.20 and $H_{T_P^+}$ as in Theorem 1.16, then $F_{H_S(P)} \circ H'_S = H_{T_P^+}$ as maps from $2_{T_P^+}$ -presilt($\mathcal{Z}_P/[P]$) to s τ -rigid pair(C).

Proof. All maps are induced by functors. We can thus claim that $F_{H_S(P)} \circ H'_S = H_{T_P^+}$, since Jasso shows that the functors inducing them obey the same relation [Jas15, Proposition 4.15]. We just have to show that H'_S has the claimed codomain, and that it is a bijection. Indeed, since we defined the support τ -rigid pairs of $J(H_S(P))$ to be those that $F_{H_S(P)}$ sends to support τ -rigid pairs of C, it follows that H'_S maps to the support τ -rigid pairs of $J(H_S(P))$. Since $F_{H_S(P)}$ and $H_{T_P^+}$ are bijections, so is H'_S . Let $(M, Q) = H_S(P)$ (where H_S is as defined in Theorem 1.16). Theorem 1.20 gives a k-algebra $C = \operatorname{End}_A(M^+)/[M^+]$, where (M^+, Q) is the Bongartz completion of the support τ -rigid pair (M, Q). We have an isomorphism $C \cong \operatorname{End}_{\mathcal{Z}_P/[P]}(T_P^+)$, where T_P^+ is the Bongartz completion of P [Jas15, Theorem 4.12(a)]. Henceforth, we abuse notation by letting C denote the endomorphism algebra of T_P^+ in $\mathcal{Z}_P/[P]$.

We now have all ingredients to prove the main theorem of this section.

Theorem 3.5. Let C, S, P, and A be as in Setup 3.2 let $C = \operatorname{End}_{\mathcal{Z}_P/[P]}(T_P^+)$. We have a commutative diagram of bijections



where $\psi_{H_S(P)} \stackrel{def}{=} F_{H_S(P)} \circ \mathcal{E}_{H_S(P)}$.

Proof. The commutativity of the lower triangle was shown in Lemma 3.4, and the right triangle (with $\psi_{H_S(P)}$ along the diagonal) commutes by definition. What remains is proving that

$$H'_{S} \circ \varphi_{P}(X) = \mathcal{E}_{H_{S}(P)} \circ H_{S}(X), \qquad (3.C)$$

which should hold for all $X \in 2_S$ -presilt_P(\mathcal{C}). All maps in question are constructed to distribute over direct sums, whence it suffices to consider the case where X is indecomposable. We treat five cases, corresponding to the definition of $\mathcal{E}_{H_S(P)}$ we reviewed in the discussion following Theorem 3.1.

Throughout, the object $R_X \in \mathcal{Z}_P$ is defined by the triangle

$$R_X \longrightarrow P_X \xrightarrow{\beta_X} X \longrightarrow \Sigma R_X$$

where β_X is a minimal right add(P)-approximation. Note that $R_X \langle 1 \rangle$ and X are isomorphic in $\mathbb{Z}_P/[P]$. When addressing the cases I(b), I(c), and II(b), we will use the fact that $H'_S \circ \varphi_P(X) = (0, \frac{\mathcal{C}}{[P]}(S, R_X))$ provided that $\frac{\mathcal{C}}{[P]}(S, X) = 0$.

Case I: Suppose that *P* has no direct summand in $\Sigma \operatorname{add}(S)$.

Case I(a), where $X \in 2_S$ -presilt_P(\mathcal{C}) has no direct summand in Σ add(S) and $\mathcal{C}(S, X) \notin \text{gen}(\mathcal{C}(S, P))$, has already been treated by Jasso [Jas15, Theorem 4.12(b)].

Case I(b): Suppose that $X \in 2_S$ -presilt_P(\mathcal{C}) has no direct summand in Σ add(S), but $\mathcal{C}(S, X) \in \text{gen}(\mathcal{C}(S, P))$. We then have that $\mathcal{E}_{H_S(P)}H_S(X) = (0, f_{\mathcal{C}(S,P)}\mathcal{C}(S, R_X))$. Since $\mathcal{C}(S, X) \in \text{gen}(\mathcal{C}(S, P))$, it follows from Lemma 3.3 that $\frac{\mathcal{C}}{[P]}(S, X) = 0$, whence $H'_S \circ \varphi_P(X) = (0, \frac{\mathcal{C}}{[P]}(X, R_X))$, so (3.C) holds.

Case I(c): Suppose that $X = \Sigma Q$, where $Q \in \text{ind} \operatorname{add}(S)$. Then

 $\mathcal{E}_{H_S(P)}H_S(X) = (0, f_{\mathcal{C}(S,P)}\mathcal{C}(S,R_X)).$ Since $\mathcal{C}(S,X) = 0$, we have that $\frac{\mathcal{C}}{[P]}(S,X) = 0$, whence (3.C) holds, for the same reason as in Case I(b).

Case II: Suppose that $P \in \Sigma \operatorname{add}(S)$.

Case II(a): If X does not have direct summands in $\Sigma \operatorname{add}(S)$, we have that $\mathcal{E}_{H_S(P)}H_S(X) = (\mathcal{C}(S,X), 0)$. On the other hand $\frac{\mathcal{C}}{[P]}(S,X) = \mathcal{C}(S,X)$, which is sufficient for (3.C) to hold.

Case II(b): If $X = \Sigma Q$, where $Q \in \text{ind} \text{add}(S)$, then $\mathcal{E}_{H_S(P)}H_S(X) = (0, f_{\mathcal{C}(S,Q)}(\mathcal{C}(S,R_X)))$. Then $\frac{\mathcal{C}}{[P]}(S,X) = 0$ since $\mathcal{C}(S,X) = 0$ and $\frac{\mathcal{C}}{[P]}(S,R_X) = f_{\mathcal{C}(S,Q)}(S,R_X)$. Thus (3.C) holds, for the same reason as in Case I(b) and Case I(c).

We have accomplished the task of linking the Buan–Marsh bijection to that of Iyama–Yang.

4. τ -cluster morphism categories

We are now ready to give a generalisation of τ -cluster morphism categories. We keep Setup 2.5, which is repeated here for the convenience of the reader.

Setup 2.5. Let A be a non-positive proper dg k-algebra, and let P be a two-term presilting object in per(A), i.e. an object in 2-presilt(A).

Given a two-term presilting object $U \in \text{per}(A)$, we proved in Lemma 2.8 that the perpendicular category $U^{\perp_{\mathbb{Z}}}$ is t-exact in $\mathcal{D}_{\text{fd}}(A)$. The τ -cluster morphism category should keep track of this information. More specifically, we want there to be a morphism

$$(\mathcal{D}_{\mathrm{fd}}(A), A) \xrightarrow{U} (U^{\perp_{\mathbb{Z}}}, \pi_U A),$$


where π_U : per(A) $\longrightarrow U^{\perp_{\mathbb{Z}}} \subseteq \mathcal{D}_{\mathrm{fd}}(A)$ is the functor constructed in Corollary 2.13. If C_U is the derived endomorphism algebra of $\pi_U A$ in $U^{\perp_{\mathbb{Z}}}$, the functor π_U induces the silting reduction map

$$\pi_{U,P}: 2\text{-presilt}_U(C_U) \longrightarrow 2\text{-presilt}(C_P)$$

a bijection previously denoted by φ_P (we often omit the U in the subscript when it is clear what it should be). Moreover, there is a triangle equivalence $U^{\perp_{\mathbb{Z}}} \simeq \mathcal{D}_{\mathrm{fd}}(C_U)$ sending $\pi_U A$ to C_U (this follows from Corollary 2.12 and Theorem 1.20). For a 2_{C_U} -term silting object P in $\mathrm{per}(C_U)$, the τ -cluster morphism category will be constructed in such a way that it contains a morphism

$$(U^{\perp_{\mathbb{Z}}}, \pi_U A) \xrightarrow{P} (U^{\perp_{\mathbb{Z}}} \cap P^{\perp_{\mathbb{Z}}}, \pi_{U,P} A)$$

It can be shown (as shall be done in Lemma 4.2) that the perpendicular category $\pi_U^{-1}(P)^{\perp_{\mathbb{Z}}}$ coincides with $U^{\perp_{\mathbb{Z}}} \cap P^{\perp_{\mathbb{Z}}}$, and that $\pi_{U,P}A$ is coincides with the composite $\pi_P \circ \pi_U$. The τ -cluster morphism category will contain a commutative diagram of the form



Definition 4.1. Let A be as in Setup 2.5. The objects of \mathfrak{W}_A are pairs (\mathcal{S}, X) , where

- S is a (thick, by Lemma 1.8) t-exact subcategory of $\mathcal{D}_{\mathrm{fd}}(A)$ such that $S = U^{\perp_{\mathbb{Z}}}$ for some two-term presilting object U in per(A),
- $X = \pi_U(A)$, where π_U : per(A) $\longrightarrow U^{\perp_{\mathbb{Z}}} \subseteq \mathcal{D}_{\mathrm{fd}}(A)$ is the functor constructed in Corollary 2.13.

Let (\mathcal{S}_1, X) such a pair. For each 2_X -term presilting object P in \mathcal{S}_1 , we add a morphism

$$P \colon (\mathcal{S}_1, X) \longrightarrow (\mathcal{S}_2, Y)$$

provided that $S_2 = S_1 \cap P^{\perp_{\mathbb{Z}}}$ and $Y = \pi_P(X)$. Given two consecutive morphisms

$$(\mathcal{S}_1, X) \xrightarrow{P} (\mathcal{S}_2, Y) \xrightarrow{Q} (\mathcal{S}_3, Z)$$
 (4.A)

we define their composition to be $\pi_P^{-1}(Q)$.

Lemma 4.2. The composition rule proposed in Definition 4.1 is well-defined. That is, if P and Q are as in (4.A), then the perpendicular category $\pi_P^{-1}(Q)^{\perp_{\mathbb{Z}}}$ coincides with $P^{\perp_{\mathbb{Z}}} \cap Q^{\perp_{\mathbb{Z}}}$,

Proof. Let $\pi_P^{-1}(Q) = P \oplus Q'$. Then we clearly have that $\pi_P^{-1}(Q)^{\perp_{\mathbb{Z}}} = P^{\perp_{\mathbb{Z}}} \cap (Q')^{\perp_{\mathbb{Z}}}$. Corollary 2.14 gives a triangle

$$(Q')_P \longrightarrow Q' \longrightarrow \pi_P Q' \longrightarrow \Sigma(Q')_P$$

where $(Q')_P \in \text{Loc}(P) \subseteq \mathcal{D}(A)$. It follows from a long exact sequence argument that $P^{\perp_{\mathbb{Z}}} \cap (Q')^{\perp_{\mathbb{Z}}} = P^{\perp_{\mathbb{Z}}} \cap Q^{\perp_{\mathbb{Z}}}$, completing the proof.

Having defined its objects, morphisms, and composition rule, we now check that \mathfrak{W}_A is a category. It is clear that the 2_X -term presilting object P = 0 in 2_X -presilt(\mathcal{S}) gives rise to the identity morphism on (\mathcal{S}, X) . However, it is not obvious that the composition rule is associative. Buan–Marsh show that their τ -cluster morphism category has an associative composition rule [BM21a, Corollary 1.8] by considering several cases. Igusa–Todorov's treatment of the representation finite hereditary case makes use of the corresponding root systems [IT17, Section 1]. Our proof takes advantage of the functoriality of silting reduction.

Theorem 4.3. The composition rule for \mathfrak{W}_A is associative. Thus \mathfrak{W}_A is a category.

Proof. Given three composable morphisms

$$(\mathcal{S}_1, X) \xrightarrow{P} (\mathcal{S}_2, Y) \xrightarrow{Q} (\mathcal{S}_3, Z) \xrightarrow{R} (\mathcal{S}_4, W)$$

we verify the identity

$$R \circ (Q \circ P) = (R \circ Q) \circ P.$$

As the bijection

$$\pi_P: 2_X$$
-presilt $_P(\mathcal{S}_1) \longrightarrow 2_Y$ -presilt (\mathcal{S}_2)

is induced by the functor π_P (and similarly for π_Q), we may apply Lemma 2.4 when expanding the right hand side.

$$\begin{split} (R \circ Q) \circ P &= (\pi_Q^{-1}(R)) \circ P \\ &= \pi_P^{-1}(\pi_Q^{-1}(R)) \\ &= \pi_{\pi_P^{-1}(Q)}^{-1}(R) \\ &= \pi_{Q \circ P}^{-1}(R) \\ &= R \circ (Q \circ P). \end{split}$$

Our main aim of this section is to show that the category defined above generalises the τ -cluster morphism category of Buan–Marsh (and Buan–Hanson), which we denote by \mathfrak{W}_B^{BM} , where B is a finitedimensional algebra. For a τ -tilting finite algebra B, recall that the objects of \mathfrak{W}_B^{BM} are the wide subcategories of mod B. For general finite-dimensional algebras, its objects are the τ -perpendicular wide subcategories, namely those that are of the form J(M,Q) for some support τ -tilting pair (M,Q) in mod A [BH21]. There is a morphism $W_1 \xrightarrow{(M,Q)} W_2$ if (M,Q) is a support τ -rigid pair in W_1 such that $J_{W_1}(M,Q) = W_2$, where $J_{W_1}(M,Q)$ is the τ -perpendicular category of (M,Q) in W_1 . The composite of (M,Q) with (M',Q') is defined by $\mathcal{E}_{(M,Q)}^{-1}(M',Q')$, where $\mathcal{E}_{(M,Q)}^{-1}$ is the inverse of the bijection in Theorem 3.1.

Note that our approach is independent of that of Buan–Marsh and Buan–Hanson. Hence, it gives an alternative approach to defining τ -cluster morphism categories for all finite-dimensional algebras, and also for non-positive proper dg k-algebras.

Theorem 4.4. Let A be as in Setup 2.5. Then the cohomological functor

$$\mathcal{D}_{\mathrm{fd}}(A, -) \colon \mathcal{D}_{\mathrm{fd}}(\mathrm{mod}\,A) \longrightarrow \mathrm{mod}\,H^0A$$

induces an equivalence of categories

$$\mathfrak{W}_A \longrightarrow \mathfrak{W}_{H^0 A}^{\mathrm{BM}}.$$
(4.B)

Explicitly, this equivalence sends the object (\mathcal{S}, X) to \mathcal{S}^0 and the map $(\mathcal{S}_1, X) \xrightarrow{P} (\mathcal{S}_2, Y)$ to $H_X(P)$ (see Theorem 1.16).

In particular, if A is a finite-dimensional algebra, then \mathfrak{W}_A is equivalent to \mathfrak{W}_A^{BM} .

Proof. Let $(U^{\perp_{\mathbb{Z}}}, \pi_U A)$ be an object in \mathfrak{W}_A . By Proposition 2.10, it is indeed sent to $J(H_A(U))$ by $\mathcal{D}_{\mathrm{fd}}(A, -)$, which is a τ -perpendicular wide subcategory of mod H^0A . This provides a suitable map of objects. Since Theorem 1.10 gives a correspondence between t-exact subcategories of $\mathcal{D}_{\mathrm{fd}}(A)$ and wide subcategories of mod H^0A , this map of objects is surjective. Differently put, if we can show that $\mathcal{D}_{\mathrm{fd}}(A)(A, -)$ induces a functor, it will be essentially surjective.

The Hom-set $\mathfrak{W}_A((\mathcal{S}_1, X), (\mathcal{S}_2, Y))$ is a subset of 2_X -presilt (\mathcal{S}_1) , and $\mathfrak{W}_A^{BM}(W_1, W_2)$ is a subset of $s\tau$ -rigid (W_1) , where W_i is the wide subcategory $\mathcal{D}_{\mathrm{fd}}(A)(A, \mathcal{S}_i)$ of mod H^0A , for $i \in \{1, 2\}$. It follows from Proposition 2.10 that the map H_A induces

$$\mathcal{D}_{\mathrm{fd}}(A)(A,-)\colon \mathfrak{W}_A((\mathcal{S}_1,X),(\mathcal{S}_2,Y)) \longrightarrow \mathfrak{W}_{H^0A}^{\mathrm{BM}}(W_1,W_2). \tag{4.C}$$

Indeed, given morphism $P \in \mathfrak{W}_A((\mathcal{S}_1, X), (\mathcal{S}_2, Y))$, we have that the τ -perpendicular category of $\mathcal{D}_{\mathrm{fd}}(A)(A, P)$ in W_1 is W_2 . A map of Hom-sets has thus been constructed.

It is easy to check that identity maps are sent to identity maps; they are parameterised by trivial presilting objects and support τ -rigid pairs. It should also be shown that composition is respected. Let P and Q be a pair of composable morphisms, as shown in (4.A). Their composition is then given by

 $\pi_P^{-1}(Q)$. Theorem 3.5 produces a commutative diagram of maps

$$2_{X} \operatorname{-presilt}_{P}(\mathcal{S}_{1}) \xrightarrow{H_{X}} \operatorname{s\tau-rigid} \operatorname{pair}_{H_{X}(P)}(\operatorname{End}_{\mathcal{D}_{\mathrm{fd}}(A)}(X))$$

$$\downarrow^{\pi_{P}} \qquad \qquad \qquad \downarrow^{\varepsilon_{H_{X}(P)}}$$

$$2_{Y} \operatorname{-presilt}(\mathcal{S}_{2}) \xrightarrow{H_{Y}} \operatorname{s\tau-rigid} \operatorname{pair}(J(H_{X}(P)))$$

$$(4.D)$$

The morphisms in $\mathfrak{W}_{H^0A}^{BM}$ corresponding to P and Q, namely $H_X(P)$ and $H_Y(Q)$, compose to $\mathcal{E}_{H_X(P)}^{-1}(H_Y(Q))$. The commutativity of (4.D) now shows that composition is respected.

Since the functor (4.B) is a bijection as a map of objects, it is essentially surjective. The maps (4.C) are the restrictions of the map shown on top in the diagram presented in Theorem 3.5. The assertion that (4.C) bijects is equivalent to the following: for two 2_X -term partial silting objects P_1 and P_2 in S_1 , we have that $P_1^{\perp_z} = P_2^{\perp_z}$ if and only if $\mathcal{D}_{\rm fd}(A)(A, P_1^{\perp_z}) = \mathcal{D}_{\rm fd}(A)(A, P_2^{\perp_z})$. This is a simple consequence of Theorem 1.10 and Lemma 2.7. This shows that the functor (4.B) is fully faithful. Having already dealt with essential surjectivity in the first paragraph of this proof, the proof is complete.

5. Two-term (pre)silting sequences

The notion of signed τ -exceptional sequences has recently been defined by Buan–Marsh [BM21b] as a generalisation of the signed exceptional sequences for representation finite hereditary algebras [IT17]. They correspond to paths in the τ -cluster morphism category [BM21a, Theorem 11.8] and ordered support τ -rigid objects [BM21b, Theorem 5.4]. We devote this section to an interpretation in terms of two-term silting.

Definition 5.1. Let A be as in Setup 2.5, and let t be a non-negative integer. A two-term presilting sequence for A is a sequence of objects

$$(X_1,\ldots,X_{t-1},X_t)$$

in $\mathcal{D}_{\rm fd}(A)$ subject to the following (recursive) conditions:

- (1) The object X_t is an indecomposable two-term presilting object in per(A),
- (2) the truncated sequence (X_1, \ldots, X_{t-1}) is a $2_{\pi_{X_t}A}$ -term presilting sequence in $X_t^{\perp_{\mathbb{Z}}}$.

We will regard the empty sequence as a two-term presilting sequence, so that the above definition makes sense when t = 0. A two-term presilting sequence of length 1 is simply an indecomposable twoterm presilting object in per(A).

Although it is not directly stated in the definition, all the objects in a two-term presilting sequence $(X_1, \ldots, X_{t-1}, X_t)$ are in per(A). The definition does require that X_t lies in per(A). One shows that X_{t-1} is in per(A) by appealing to the fact that it is two-term with respect to $\pi_{X_t}(A)$, which by Corollary 2.13 is in per(A). One then continues recursively to show that all X_i are in per(A).

Buan–Marsh define a τ -exceptional sequence for A as a sequence of objects in the category

1.0

$$\mathcal{C}(A) \stackrel{\text{der}}{=} \mod A \sqcup \Sigma \mod A \subseteq \mathcal{D}^{\mathrm{b}}(\mod A),$$

such that

- (1) the last term X_t is an indecomposable τ -rigid pair,
- (2) the removal of X_t yields a τ -exceptional sequence for the τ -tilting reduction $J(X_t)$.

It is not true that Definition 5.1 specialises to Buan–Marsh' definition in this case; the last object X_t is two-term, whereas it is a module or a suspended projective in the original definition. We do, however, have a suitable bijection.

Theorem 5.2. Let A be as in Setup 2.5, and let t be a non-negative integer. Consider the map

where X_1 is the largest direct summand of X with no cohomology in degree 0 or greater. It induces a bijection

$$(X_1, \dots, X_{t-1}, X_t) \longmapsto (H_A X_1, \dots, H_A X_{t-1}, H_A X_t),$$
(5.A)

from the first to the second of the following sets:

- (1) two-term presilting sequences of length t in $\mathcal{D}_{\mathrm{fd}}(A)$,
- (2) signed τ -exceptional sequences of length t in $\mathcal{C}(H^0A)$.

Proof. We proceed by induction on t. The statement is vacuously true for t = 0, and the case t = 1 is proved by Theorem 1.16, treating a support τ -rigid pair (M, P) as the object $M \oplus \Sigma P \in \mathcal{C}(H^0A)$.

Suppose that the statement holds for $t = \ell - 1$, for some $\ell \ge 1$. To prove that the statement holds for $t = \ell$, it suffices to fix a two-term presilting object X_{ℓ} and prove that the map H_A provides a bijection between the sets

- (1) two-term presilting sequences of length ℓ in $\mathcal{D}_{\mathrm{fd}}(A)$ ending in X_{ℓ} ,
- (2) signed τ -exceptional sequences of length ℓ in $\mathcal{C}(H^0A)$ ending in $H_A X_{\ell}$.

Indeed, since H_A is a bijection, a two-term presilting sequence in $\mathcal{D}_{\mathrm{fd}}(A)$ ends in X_{ℓ} precisely when the corresponding sequence in $\mathcal{C}(H^0A)$ ends in $H_A X_{\ell}$.

Let $(X_1, \ldots, X_{\ell-1}, X_\ell)$ be such a 2_S -term presilting sequence of length ℓ . Then the truncated sequence $(X_1, \ldots, X_{\ell-1})$ is a signed $2_{\pi_P A}$ -term presilting sequence in $X_{\ell}^{\perp z}$. The induction hypothesis gives that the cohomology functor $H_{\pi_P A}$ provides a bijection between the sets

- (1) 2_{π_PA} -term presilting sequences of length $\ell 1$ in $X_{\ell}^{\perp_{\mathbb{Z}}}$,
- (2) signed τ -exceptional sequences of length $\ell 1$ in $(X_{\ell}^{\perp_{\mathbb{Z}}})^0 \cong J(H_A(X_{\ell}))$ (see Proposition 2.10).

The maps H_A and $H_{\pi_P A}$ are determined by the cohomological functors $\mathcal{D}_{\mathrm{fd}}(A)(A, -)$ and $\mathcal{D}_{\mathrm{fd}}(A)(\pi_P A, -)$. These coincide when restricted to $X_{\ell}^{\perp_{\mathbb{Z}}}$. Indeed, an object $Y \in X_{\ell}^{\perp_{\mathbb{Z}}}$ is expressed as $\pi_{X_{\ell}}A$, and by Corollary 2.14 there is a triangle

$$A_{X_{\ell}} \longrightarrow A \longrightarrow Y \longrightarrow \Sigma A_{X_{\ell}}$$

with $A_{X_{\ell}} \in \operatorname{Loc}(X_{\ell}) \subseteq \mathcal{D}(A)$, whence the claim follows from a long exact sequence argument.

It follows that the map (5.A) is a bijection. This completes the inductive step, and the proof. \Box

Buan-Marsh define a complete signed τ -exceptional sequence to be one of length |A|, the rank of the algebra. These sequences correspond to ordered support τ -tilting objects [BM21b, Theorem 5.4]. If we define a two-term presilting sequence to be a two-term silting sequence provided that its length is the same as the number of indecomposable direct summands of A, the bijection in Theorem 5.2 restricts to a bijection between two-term silting sequences and complete signed τ -exceptional sequences.

References

- [AI12] T. Aihara and O. Iyama, "Silting mutation in triangulated categories," J. Lond. Math. Soc. (2), vol. 85, no. 3, pp. 633–668, 2012. [Online]. Available: https://doi.org/10.1112/jlms/jdr055
- [AIR14] T. Adachi, O. Iyama, and I. Reiten, "τ-tilting theory," Compos. Math., vol. 150, no. 3, pp. 415–452, 2014.
 [Online]. Available: https://doi.org/10.1112/S0010437X13007422
- [AMY19] T. Adachi, Y. Mizuno, and D. Yang, "Discreteness of silting objects and t-structures in triangulated categories," Proc. Lond. Math. Soc. (3), vol. 118, no. 1, pp. 1–42, 2019. [Online]. Available: https://doi.org/10.1112/plms.12176
- [AS81] M. Auslander and S. O. Smalø, "Almost split sequences in subcategories," J. Algebra, vol. 69, no. 2, pp. 426–454, 1981. [Online]. Available: https://doi.org/10.1016/0021-8693(81)90214-3
- [ATJLSS03] L. Alonso Tarrío, A. Jeremías López, and M. J. Sauto Salorio, "Construction of t-structures and equivalences of derived categories," *Transactions of the American Mathematical Society*, vol. 355, no. 6, pp. 2523–2543, 2003.
- [BBD82] A. A. Beïlinson, J. Bernstein, and P. Deligne, "Faisceaux pervers," in Analysis and topology on singular spaces, I (Luminy, 1981), ser. Astérisque. Soc. Math. France, Paris, 1982, vol. 100, pp. 5–171.
- [BBDG18] A. Beilinson, J. Bernstein, P. Deligne, and O. Gabber, *Faisceaux pervers*. Société mathématique de France, 2018.
- [BH21] A. Bakke Buan and E. J. Hanson, "τ-perpendicular wide subcategories," arXiv e-prints, p. arXiv:2107.01141, Jul. 2021.
- [BM21a] A. B. Buan and B. R. Marsh, "A category of wide subcategories," Int. Math. Res. Not. IMRN, no. 13, pp. 10278–10338, 2021. [Online]. Available: https://doi.org/10.1093/imrn/rnz082
- $[BM21b] \qquad ---, \quad ``\tau\text{-exceptional sequences,''} \quad J. \quad Algebra, \quad \text{vol. 585, pp. 36-68, 2021. [Online]. Available: https://doi.org/10.1016/j.jalgebra.2021.04.038$
- [BMR⁺06] A. B. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov, "Tilting theory and cluster combinatorics," Adv. Math., vol. 204, no. 2, pp. 572–618, 2006. [Online]. Available: https://doi.org/10.1016/j.aim.2005.06.003
- [BZ16] A. B. Buan and Y. Zhou, "A silting theorem," J. Pure Appl. Algebra, vol. 220, no. 7, pp. 2748–2770, 2016.
 [Online]. Available: https://doi.org/10.1016/j.jpaa.2015.12.009
- [BZ21] A. Bakke Buan and Y. Zhou, "Weak cotorsion, τ -tilting and two-term categories," arXiv e-prints, p. arXiv:2111.10995, Nov. 2021.
- [DIR⁺17] L. Demonet, O. Iyama, N. Reading, I. Reiten, and H. Thomas, "Lattice theory of torsion classes," arXiv e-prints, p. arXiv:1711.01785, Nov. 2017.
- [GP18] N. Gao and C. Psaroudakis, "Ladders of compactly generated triangulated categories and preprojective algebras," Applied Categorical Structures, vol. 26, no. 4, pp. 657–679, 2018.
- [HI21] E. J. Hanson and K. Igusa, "τ-Cluster morphism categories and picture groups," Comm. Algebra, vol. 49, no. 10, pp. 4376–4415, 2021. [Online]. Available: https://doi.org/10.1080/00927872.2021.1921184
- [Igu14] K. Igusa, "The category of noncrossing partitions," *arXiv e-prints*, p. arXiv:1411.0196, Nov. 2014.
- [IJY14] O. Iyama, P. Jørgensen, and D. Yang, "Intermediate co-t-structures, two-term silting objects, τ-tilting modules, and torsion classes," Algebra Number Theory, vol. 8, no. 10, pp. 2413–2431, 2014. [Online]. Available: https://doi.org/10.2140/ant.2014.8.2413

ERLEND D. BØRVE

- [IT17] K. Igusa and G. Todorov, "Signed exceptional sequences and the cluster morphism category," arXiv e-prints, p. arXiv:1706.02041, Jun. 2017.
- [ITW16] K. Igusa, G. Todorov, and J. Weyman, "Picture groups of finite type and cohomology in type A_n ," arXiv *e-prints*, p. arXiv:1609.02636, Sep. 2016.
- [IY08] O. Iyama and Y. Yoshino, "Mutation in triangulated categories and rigid Cohen-Macaulay modules," Invent. Math., vol. 172, no. 1, pp. 117–168, 2008. [Online]. Available: https://doi.org/10.1007/s00222-007-0096-4
- [IY18] O. Iyama and D. Yang, "Silting reduction and Calabi–Yau reduction of triangulated categories," Trans. Amer. Math. Soc., vol. 370, no. 11, pp. 7861–7898, 2018. [Online]. Available: https://doi.org/10.1090/tran/7213
- [Jas15] G. Jasso, "Reduction of τ-tilting modules and torsion pairs," Int. Math. Res. Not. IMRN, no. 16, pp. 7190–7237, 2015. [Online]. Available: https://doi.org/10.1093/imrn/rnu163
- [Kel94] B. Keller, "Deriving dg categories," in Annales scientifiques de l'Ecole normale supérieure, vol. 27, no. 1, 1994, pp. 63–102.
- [Kra08] H. Krause, "Localization theory for triangulated categories," arXiv preprint arXiv:0806.1324, 2008.
- [KV88] B. Keller and D. Vossieck, "Aisles in derived categories," 1988, vol. 40, no. 2, pp. 239–253, deuxième Contact Franco-Belge en Algèbre (Faulx-les-Tombes, 1987).
- [KY14] S. Koenig and D. Yang, "Silting objects, simple-minded collections, t-structures and co-t-structures for finitedimensional algebras," Doc. Math., vol. 19, pp. 403–438, 2014.
- [MHT20] O. Mendoza Hernández and H. Treffinger, "Stratifying systems through τ-tilting theory," Documenta Mathematica, vol. 25, pp. 701–720, 2020.
- [Nee92] A. Neeman, "The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel," in Annales scientifiques de l'Ecole normale supérieure, vol. 25, no. 5, 1992, pp. 547–566.
- [Ric89] J. Rickard, "Morita theory for derived categories," J. Lond. Math. Soc., II. Ser., vol. 39, no. 3, pp. 436–456, 1989.
- [ZC17] C. Zhang and H. Cai, "A note on thick subcategories and wide subcategories," Homology Homotopy Appl., vol. 19, no. 2, pp. 131–139, 2017. [Online]. Available: https://doi.org/10.4310/HHA.2017.v19.n2.a8

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU, NO-7491 TRONDHEIM, NORWAY

Current address: Institut Fourier, Université Grenoble Alpes, 100 rue des mathématiques, 38610 Gières, France *Email address*: erlend.borve@univ-grenoble-alpes.fr

28