Nonlocal Fully Nonlinear Parabolic Differential Equations Arising in Time-Inconsistent Problems¹

Qian Lei^a, Chi Seng Pun^{a,*}

^aSchool of Physical and Mathematical Sciences, Nanyang Technological University, Singapore

Abstract

We prove the well-posedness results, i.e. existence, uniqueness, and stability, of the solutions to a class of nonlocal fully nonlinear parabolic partial differential equations (PDEs), where there is an external time parameter t on top of the temporal and spatial variables (s, y) and thus the problem could be considered as a flow of equations. The nonlocality comes from the dependence on the unknown function and its first- and second-order derivatives evaluated at not only the local point (t, s, y) but also at the diagonal line of the time domain (s, s, y). Such equations arise from time-inconsistent problems in game theory or behavioural economics, where the observations and preferences are (reference-)time-dependent. To address the open problem of the well-posedness of the corresponding nonlocal PDEs (or the time-inconsistent problems), we first study the linearized version of the nonlocal PDEs with an innovative construction of appropriate norms and Banach spaces and contraction mappings over which. With fixed-point arguments, we obtain the wellposedness of nonlocal linear PDEs and establish a Schauder-type prior estimate for the solutions. Then, by the linearization method, we analogously establish the well-posedness under the fully nonlinear case. Moreover, we reveal that the solution of a nonlocal fully nonlinear parabolic PDE is an adapted solution to a flow of second-order forward-backward stochastic differential equations.

Keywords: Existence and Uniqueness, Nonlocal Nonlinear Parabolic PDEs, Parametric PDEs, Time inconsistency, FBSDE, Mathematics of Behavioral Economics

¹Chi Seng Pun gratefully acknowledges Ministry of Education (MOE), AcRF Tier 2 grant (Reference No: MOE2017-T2-1-044) for the funding of this research.

^{*}Corresponding author

Email addresses: leiq0002@e.ntu.edu.sg(Qian Lei), cspun@ntu.edu.sg(Chi Seng Pun)

1. Introduction

We study the existence and uniqueness problems for a class of nonlocal fully nonlinear parabolic partial differential equations (PDEs) of the form

$$\begin{cases} u_s(t, s, y) = F(t, s, y, u(t, s, y), u_y(t, s, y), u_{yy}(t, s, y), \\ u(s, s, y), u_y(s, s, y), u_{yy}(s, s, y)), \\ u(t, 0, y) = g(t, y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^d. \end{cases}$$
(1)

where the mapping F could be nonlinear with respect to all its arguments, s and y are temporal and spatial variables, respectively, while t could be considered as an external temporal parameter, and the temporal variables (s,t) are defined in a triangular region $\Delta[0,T] := \{(t,s) \in [0,T]^2 : 0 \le s \le t \le T\}$. The nonlocality of (1) stems from the "diagonal dependence," where u, u_y , and u_{yy} in F are evaluated not only at (t, s, y) but also at (s, s, y). The PDEs (1) originate from various problems in theories of stochastic control and stochastic differential equation with their applications in financial economics when behavioural factors are considered.

Organization of this paper. Subsection 1.1 briefly introduces the nonlocal PDEs as the driving force of a Nobel-prize winning theory, namely prospect theory, while Subsection 1.2 reviews the literature on this research direction, followed by a succinct account of our insights and main contributions in Subsections 1.3 and 1.4. Our study begins with a linearized case (linear F) in Section 2, where we first introduce appropriate norms and Banach spaces for our study in Subsection 2.1 and then the well-posedness results in Subsection 2.2, which acquires a Schauder-type prior estimates revealing the relationship among solutions of the nonlocal equations, the nonhomogeneous terms, and the initial conditions. Under mild conditions, Section 3 is devoted to prove the existence and uniqueness of nonlocal fully nonlinear parabolic PDEs by fixed-point arguments. Section 4 presents the close connection between nonlocal parabolic PDEs and forward-backward stochastic differential equations (FBSDEs). Moreover Subsection 4.2 shows that the solution to a nonlocal fully nolinear parabolic PDE solves a corresponding flow of second-order FBSDEs, which implies a new Feynman–Kac formula. Finally, Section 5 concludes.

1.1. Nonlocality, Time-Inconsistency, and the Lack of Flow Property

Pontryagin's maximum principle and Bellman's principle of optimality (or dynamic programming, DP) are two main pillars of solving a stochastic control problem; see [1]. However, these two principles could be violated in a common situation in economics when a decision-maker's preferences change over time, i.e. dynamic inconsistency (in the context of game theory) or time inconsistency (in the context of behavioral economics). Some well-known examples are the financial mean-variance portfolio selection [2], hyperbolic discounting in behavioural economics [3], and endogenous habit formulation. As evidenced in the prospect theory ([4]) and [5], the psychological principles of reference dependence and state-dependent objectives should be taken into account during the decision making, but they cause time-inconsistent (TIC) issues. In what follows, we briefly introduce the relevance of solving (1) and its connection with stochastic differential equation (SDE) theory.

Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a completed filtered probability space on which a k-dimensional Brownian motion $W(\cdot)$ with the natural filtration $\mathbb{F} = \{\mathcal{F}_s\}_{s\geq 0}$ augmented by all the \mathbb{P} -null sets in \mathcal{F} is well-defined. By choosing a suitable control process $\alpha(\cdot) : [s, T] \times \Omega \to U$ with $U \subseteq \mathbb{R}^m$ being a non-empty set, we aim to optimize the following cost functional:

$$\inf_{\alpha} J(s, y; \alpha(\cdot)) \tag{2}$$

with
$$J(s, y; \alpha(\cdot)) := \mathbb{E}\left[\int_{s}^{T} h(s, \tau, X(\tau), \alpha(\tau)) d\tau + g(s, X(T)) \middle| \mathcal{F}_{s}\right]$$
 (3)

and the controlled state process $\{X(\tau)\}_{\tau \in [s,T]}$ driven by the following SDE:

$$\begin{cases} dX(\tau) = b(\tau, X(\tau), \alpha(\tau))d\tau + \sigma(\tau, X(\tau), \alpha(\tau))dW(\tau), & \tau \in [s, T], \\ X(s) = y, \quad y \in \mathbb{R}^d. \end{cases}$$
(4)

For illustration, we assume a Markovian framework and that all the coefficient and objective functions, $b : [0,T] \times \mathbb{R}^d \times U \mapsto \mathbb{R}^d$, $\sigma : [0,T] \times \mathbb{R}^d \times U \mapsto \mathbb{R}^{d \times k}$, $h : \nabla[0,T] \times \mathbb{R}^d \times U \mapsto \mathbb{R}$ and $g : [0,T] \times \mathbb{R}^d \mapsto \mathbb{R}$ are deterministic, where $\nabla[0,T] := \{(s,\tau) \in [0,T]^2 : 0 \le s \le \tau \le T\}$. A distinct feature of problem (2) is its running and terminal costs, h and g in (3), varying in the time variable s. Hyperbolic or delay discounting (see [5, 6]) is a typical example of (3), which induces time (dynamic) inconsistency and incents the decision-makers deviating their pre-committed plan for some future period when it arrives. This is also a common feature of decision-making under the prospect theory [4]. The time inconsistency (TIC) of problem (2) also manifests itself through a violation of Bellman's principle of optimality and thus standard DP arguments are not directly applicable in this case. In the context of game theory, a pre-commitment plan to problem (2) is subgame imperfect, where players are the incarnations of the agent parametrized by $s \in [0, T]$. In this context, finding a Nash subgame perfect equilibrium is essentially a consistent planning that is economically meaningful and appealing; see [7, 8].

Recent studies have contributed to convert the problem (2) to solving Hamilton–Jacobi–Bellman (HJB) equation(s), which is essentially a (nonlocal) nonlinear PDE. There are two derivations of the PDE based on a modified recursive equation of the value function V in (2) and a discretization method, respectively, with game-theoretic and DP arguments. [9] characterize the subgame perfect equilibrium via the solution (v(s, y), u(t, s, y)) of a system of HJB equations: for any $(t, s) \in \nabla[0, T]$ and $y \in \mathbb{R}^d$, v(s, y) = u(s, s, y) and

$$\begin{cases} v_s(s,y) + \inf_{a \in U} \left\{ \mathcal{H}(s,s,y,a,v_y(s,y),v_{yy}(s,y)) \right\} = 0, \\ u_s(t,s,y) + \mathcal{H}(t,s,y,e(s,y),u_y(t,s,y),u_{yy}(t,s,y)) = 0, \end{cases}$$
(5)

with boundary conditions v(T, y) = g(T, y) and u(t, T, y) = g(t, y) for $t \in [0, T]$ and $y \in \mathbb{R}^d$, where the Hamiltonian $\mathcal{H} : \nabla[0, T] \times \mathbb{R}^d \times U \times \mathbb{R}^d \times \mathbb{S}^d \to \mathbb{R}$ with $\mathbb{S}^d \subseteq \mathbb{R}^{d \times d}$ being the set of all $d \times d$ symmetric matrices is defined as:

$$\mathcal{H}(t,s,y,a,p,q) = \frac{1}{2} \operatorname{tr} \left[q \cdot \left(\sigma \sigma^{\top} \right)(s,y,a) \right] + p^{\top} b(s,y,a) + h\left(t,s,y,a\right)$$
(6)

and the e(s, y) in the second equation of (5) realizes the supremum in the first equation (if it always exists) and constitutes an equilibrium control policy for $s \in [0, T]$. We first provide two important observations about the system (5): first, we need to determine simultaneously u and v(or equivalently e that depends on v); second, from the first equation and the relation between vand u, we know that the e(s, y) depends on $s, y, v_y(s, y) = u_y(s, s, y)$, and $v_{yy}(s, y) = u_{yy}(s, s, y)$, i.e.

$$e(s,y) = \phi\left(s,s,y,u_y(s,s,y),u_{yy}(s,s,y)\right) \quad \text{for } \phi(t,s,y,p,q) \in \inf_a \mathcal{H}(t,s,y,a,p,q) \tag{7}$$

as we allow the control variate $\alpha(\cdot)$ entering both of the drift b and volatility σ of the controlled state process $X(\cdot)$. Hence, it suffices to solve for the second equation of (5) with e(s, y) of the form (7): for $(t, s) \in \nabla[0, T]$ and $y \in \mathbb{R}^d$,

$$u_{s}(t,s,y) + \mathcal{H}\Big(t,s,y,\phi\big(s,s,y,u_{y}(s,s,y),u_{yy}(s,s,y)\big), u_{y}(t,s,y),u_{yy}(t,s,y)\Big) = 0$$
(8)

with boundary condition u(t, T, y) = g(t, y) for $t \in [0, T]$ and $y \in \mathbb{R}^d$, while the verification theorem in [9] validates that the classical solution to (5) or equivalently (8) is the subgame perfect equilibrium solution. By a discretization method and taking the continuous-time limit, [10, 11, 12] also yield the same equation of (8) for the problem (2) and the authors call it as an *equilibrium HJB equation*. In contrast with the conventional HJB equation, which is a **local** fully nonlinear PDE, the equilibrium HJB equation (8) is a backward **nonlocal** fully nonlinear PDE as a special case of (1). Heuristically speaking, the decision-makers/players in TIC stochastic control problems are supposed to follow the principle of "globally thinking, locally acting." "Thinking globally" induces the *u* terms evaluated at (t, s, y), while "acting locally" implies them evaluated at (s, s, y)in the equation (8).

In addition to the originations and inspirations from TIC stochastic control problems, nonlocal PDEs are in close connection with the SDE theory, especially for a flow of FBSDEs or backward stochastic Volterra integral equation (BSVIE). As the well-known Feynman–Kac formula (see [13]) indicates, an adapted solution $(X(\cdot), Y(\cdot), Z(\cdot))$ of a system of FBSDE gives a stochastic representation of the solution u(s, y) of local parabolic PDEs. In the classical setting, the Feynman–Kac formula connects two sides of the adapted solution $Y(\cdot)$ and $Z(\cdot)$ of a BSDE and the solution u(s, y) of the corresponding parabolic PDE in the fashion that $Y(\cdot) = u(\cdot, X(\cdot))$ and $Z(\cdot) = \sigma^{\top}(\cdot, X(\cdot))u_y(\cdot, X(\cdot))$, where $X(\cdot)$ solves a forward SDE. Depending on whether the FBSDE is coupled or not, the parabolic PDEs are either semi-linear or quasi-linear. By introducing the dependence of the current time t into the generator and the terminal condition of BSDEs, a family of FBSDEs parameterized by t forms a flow of FBSDEs, which is linked to the system of HJB equations. Moreover, a parallel extension of BSDE leads to a BSVIE, which is linked to the equilibrium HJB equation. In the context of SDE theory, BSVIE is **lack of flow property** that a normal BSDE possesses. Analogous to the FBSDE theory, we can also anticipate a Feynman–Kactype representations of adapted solutions of FBSVIEs, which are characterized by the solution of a nonlocal parabolic PDE. As Section 4 elaborates, all the nonlocal parabolic PDEs in the existing Feynman–Kac-type formulas of FBSVIEs are special cases of our general nonlocal fully nonlinear PDE (1). It is noteworthy that a nonlocal fully nonlinear parabolic PDEs are corresponded to a flow of second-order FBSDEs (2BSDEs) or second-order FBSVIEs (2FBSVIEs), which creates new knowledge to the literature. A further exploration between nonlocal parabolic PDEs and a flow of FBSDEs (or 2FBSDEs) will be detailed in Section 4.

1.2. Limitation of the Existing Studies

In general, the existence and uniqueness problems of the equilibrium HJB equation (8) or the system of HJB equations (5) still remain open, although there are several attempts on their existence and uniqueness in the past decade. To the best of our knowledge, the best result in this regard so far is obtained in [11], where the authors presented a existence and uniqueness result for the case that the nonlocal PDEs are restricted to the linear dependence on the second-order derivative at local point $u_{yy}(t, s, y)$ and the removal of the second-order derivative at diagonal $u_{yy}(s, s, y)$. The removal was justified by prohibiting the control from entering the diffusion of the state process (4), i.e. $\sigma(s, y, a) = \sigma(s, y)$, which would lead to the $u_{yy}(s, s, y)$ -independence of the equilibrium control policy e(s, y) (7) and the equilibrium HJB equation (8). The similar restriction is inherited to the subsequent works, e.g., [14, 15, 16]. However, without controls in the diffusion of the state, we can hardly control the risk (noises) from the stochastic systems, which are crucial in many problems such as portfolio management, inventory control, etc. In fact, only when the controls could or would take effect on the magnitude of uncertainty, the stochastic problems differ from the deterministic ones. In the related works of [10, 11, 17, 14, 15, 18], all the authors admit that the existence and uniqueness problems of nonlocal fully nonlinear parabolic PDEs with the term $u_{yy}(s, s, y)$ is a complicated open problem. Hence, if we can establish the well-posedness of nonlocal fully nonlinear parabolic PDEs (1), the well-posedness of the subgame perfect equilibrium is resolved, so are some open problems listed in the discussion of [9].

From the perspective of FBSDEs, hindered by the limitation of the nonlocal PDE theory, there is lack of unified and general treatment for the PDE-side in Feynman–Kac formulas of BSVIEs.

In fact, as the Feynman–Kac formula indicates, advances of the well-posedness results of nonlocal PDEs allow us to solve directly the corresponding FBSVIE $(X(\cdot), Y(\cdot, \cdot), Z(\cdot, \cdot))$ by combining solutions of nonlocal PDEs $u(\cdot, \cdot, \cdot)$ and of the coupled forward SDE $X(\cdot)$. The well-posedness of nonlocal PDEs can be utilized to study the solvability of FBSVIEs in a similar manner of the classical four-step numerical scheme for studying the solvability of FBSDEs; see [19]. Likewise, limited by the development of nonlocal PDEs, the FBSVIEs in the existing literature only correspond to some special cases of (1), such as nonlocal *semi-linear* or *quasi-linear* PDEs as in [11, 20, 17]. Therefore, the study of nonlocal *fully nonlinear* PDEs (1) provides a solid foundation of studying a more general form of FBSVIEs.

1.3. Insights into Analysing Nonlocal Fully Nonlinear PDEs

Given any suitable pair (F, g) of nonlinearity and initial condition, the well-posedness of the nonlocal differential equations (1) requires

- 1. a solution exists in some sense;
- 2. the solution is unique in some space;
- 3. the map from (F, g) to solutions is continuous in some topology.

First of all, this paper adopts the concept of classical solutions with sufficient regularities, i.e. the smoothness of first- and second-order derivatives of u with respect to s and y. Within a classical solution framework, we can take advantage of methodologies for local parabolic PDEs in [21, 22, 23], such as various regularity results and prior estimates of solutions, as well as represent the equilibrium controls in a practical closed form (7). Moreover, for such a new class of nonlocal PDEs (1), both dependence on diagonal terms and the order relation between t and s distinguish themselves from the local PDEs. Hence, we need to identify suitable norms and Banach spaces for the nonlocal PDEs and even for all TIC problems for practicality. Under a well-defined norm and the induced Banach space, the non-linearity F is required to be closed in some sense.

In this paper, we make use of fixed-point arguments to show the well-posedness of (1). The key steps of which are to construct suitable mappings and to prove their contractions under the

norms introduced. However, the presence of the second-order diagonal term $u_{yy}(s, s, y)$ causes an essential difficulty for constructing a desired contraction. As discussed in the previous section, this difficulty has been the major obstacle for many studies on TIC problems.

To understand the essence of our problem, we illustrate the difficulty with a simplified problem of nonlocal linear PDE without the low-order terms. For the existence and uniqueness problems of such an equation, it is natural to consider a mapping from u to w that satisfies

$$w_s(t,s,y) = a(t,s,y)w_{yy}(t,s,y) + \overline{a}(t,s,y)u_{yy}(s,s,y)$$
(9)

It is easy to see that the mapping is well-defined, thanks to the classical PDE theory. By replacing the unmanageable diagonal term with a known function u, the well-posedness of the PDEs (9) parameterized by t promises the existence and uniqueness of the solution w. Moreover, it is also clear that the fixed point solves the original nonlocal linear PDE. However, since the input u is of the same order of the output w by simple analysis, it is not feasible to prove that they are contractive.

We overcome the major difficulties by developing a range of techniques and results, three essential ingredients of which are listed as follows:

- (a) studying the well-posedness of a linearized version of (1), i.e. (10), where the mapping F is linear with respect to all its arguments related to u. Giving up the kind of mappings directly from u to w like (9), we propose a coupled nonlocal PDE system of (u, u_t) , which consists of the unknown function u and its derivative u_t with respective to t, and the system is mathematically equivalent to the linearized nonlocal PDE via the linkage between their solutions (u). In this way, the smoothness of u in t can help us achieve a desired contraction such that the system of (u, u_t) admits a unique solution. Consequently, the well-posedness of original linearized nonlocal PDEs is established. The technical details are discussed in Theorem 2.1;
- (b) establishing a Schauder-type prior estimate for solutions of linear nonlocal PDEs studied in (a). In general, a successful treatment of nonlinear PDEs depends on a prior estimate. Based on the norms defined by us for nonlocal PDEs (and TIC problems), the prior estimate

not only controls the behavior of the solutions and provides quantitative information on the regularity of solutions, but also gives a certain compactness to the class of possible solutions. Such a compactness is necessary for Picard's iterative method and Banach's fixed-point theorem in the study of nonlocal fully nonlinear PDEs (1). The Schauder prior estimate is presented in Corollary 2.1;

(c) linearizing nonlocal fully nonlinear PDEs into nonlocal linear equations. The established results in (a) and (b) are utilized to show the well-posedness of nonlocal fully nonlinear PDEs of the form (1) with the linearization. We first construct a mapping from u to w

$$w_s = \mathcal{L}w + F(t, s, y, u, u_y, u_{yy}, u_{t=s}, u_y|_{t=s}, u_{yy}|_{t=s}) - \mathcal{L}u$$

where \mathcal{L} is a nonlocal linear differential operator defined in (34). Given u in a suitable space, the solvability of nonlocal linear PDEs proven in (a) ensures the operator well-defined. Furthermore, the prior estimate obtained in (b) provides a bound of the solution w with the nonhomogeneous term $F - \mathcal{L}u$ and a given initial condition g. Subsequently, we can prove that the mapping is a contraction in a suitable Banach space. Consequently, the unique fixed point solves the nonlocal fully nonlinear PDEs (1).

Finally, the stability of solutions with respect to the data (F, g) is also important for theoretical analysis and practical applications. The map from data to solutions is continuous in the topology induced by our norms. In summary, the essential difficulties of studying the nonlocal fully non-linear PDEs (or TIC problems) is not only to establish suitable Banach spaces and well-defined mappings over them, but also to ensure that the mappings are contractive over the norm-induced spaces.

1.4. Main Contributions

This paper aims to address some open problems in the theories of PDE, SDE, and stochastic controls for the cases where there is nonlocality (for PDEs), lack of flow property (for FBSDEs), or time-inconsistency (for optimal stochastic controls). By noting the connections among these three fields, the aforementioned three scenarios are haunted by the similar mathematical difficulty.

Hence, our study of nonlocality in PDEs is beneficial to understanding time-inconsistency in FB-SDE and control problems.

The main contributions of the this paper is threefold. First, we propose suitable norms and Banach spaces for studying nonlocal PDEs and TIC problems. Under them, the well-posedness results of nonlocal linear PDEs are first proven. After establishing a Schauder-type prior estimate, the method of linearization is adopted to show the existence and uniqueness of nonlocal fully nonlinear PDEs. Second, thanks to these regularity results of nonlocal PDEs (1), the well-posedness of solutions of the equilibrium HJB equation (8) and the system of HJB equations (5), both of which identify the equilibrium policy and the equilibrium value functions in TIC problems, can be also established even when the control $\alpha(\cdot)$ enters the diffusion term σ of state processes (the bottleneck of the existing studies). Third, a unified and general treatment of PDE representations of adapted solutions of a flow of second-order FBSDEs is initiated, which enhances the feasibility of studying FBSDEs or FBSVIEs from the modelling perspective of PDEs.

2. Nonlocal Linear Parabolic PDEs

In this section, we first propose suitable norms and Banach spaces for nonlocal PDEs. Based on them, we will prove the solvability of nonlocal linear PDEs and estimate their solutions.

2.1. Preliminaries: Norms and Banach Spaces

For the conventional local parabolic PDEs, the corresponding differential operator works well over Hölder spaces of sufficiently smooth functions. It is expected that some revision of the Hölder spaces are desired and effective for our nonlocal parabolic equations. Hence, we review some definitions of spaces of Hölder continuous functions here.

Given $0 \le a \le b \le T$, we denote by $C([a, b] \times \mathbb{R}^d)$ the set of all the continuous and bounded functions in $[a, b] \times \mathbb{R}^d$ endowed with the sup norm $|\cdot|_{[a,b] \times \mathbb{R}^d}^{\infty}$. Wherever no confusion arises, we write $|\cdot|^{\infty}$ instead of $|\cdot|_{[a,b]\times\mathbb{R}^d}^{\infty}$. Next, for $\alpha \in (0,1)$, we introduce Hölder spaces:

$$C^{\alpha,0}([a,b] \times \mathbb{R}^d) = \left\{ \varphi \in C([a,b] \times \mathbb{R}^d) : \langle \varphi \rangle_s^{(\alpha)} := \sup_{\substack{a \le s < s' \le b \\ y \in \mathbb{R}^d}} \frac{|\varphi(s,y) - \varphi(s',y)|}{|s-s'|^{\alpha}} < \infty, \\ |\varphi|_{C^{\alpha,0}([a,b] \times \mathbb{R}^d)} := |\varphi|^{\infty} + \langle \varphi \rangle_s^{(\alpha)} < \infty \right\}$$

and similarly,

$$C^{0,\alpha}([a,b] \times \mathbb{R}^d) = \left\{ \varphi \in C([a,b] \times \mathbb{R}^d) : \langle \varphi \rangle_y^{(\alpha)} := \sup_{\substack{a \le s \le b \\ y,y' \in \mathbb{R}^d}} \frac{|\varphi(s,y) - \varphi(s,y')|}{|y - y'|^{\alpha}} < \infty, \\ |\varphi|_{C^{0,\alpha}([a,b] \times \mathbb{R}^d)} := |\varphi|^{\infty} + \langle \varphi \rangle_y^{(\alpha)} < \infty \right\}.$$

Moreover, the classical set $C^{1,2}([a,b]\times \mathbb{R}^d)$ is defined as

$$C^{1,2}([a,b] \times \mathbb{R}^d) = \Big\{ \varphi \in C([a,b] \times \mathbb{R}^d) : \exists \varphi_s, \ \varphi_{y_i y_j} \in C([a,b] \times \mathbb{R}^d), \ i,j = 1, \dots, d \Big\}.$$

which is endowed with

$$|\varphi|_{C^{1,2}([a,b]\times\mathbb{R}^d)} = |\varphi|^\infty + \sum_{i=1}^d |\varphi_{y_i}|^\infty + |\varphi_s|^\infty + \sum_{i,j=1}^d |\varphi_{y_iy_j}|^\infty.$$

Now, we define "parabolic" Hölder spaces, which are very common in the study of local parabolic equations. We set

$$C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d) = C^{\frac{\alpha}{2},0}([a,b]\times\mathbb{R}^d)\bigcap C^{0,\alpha}([a,b]\times\mathbb{R}^d),$$

the norm of which is given by

$$|\varphi|_{C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d)} = |\varphi|_{C^{\frac{\alpha}{2},0}([a,b]\times\mathbb{R}^d)} + |\varphi|_{C^{0,\alpha}([a,b]\times\mathbb{R}^d)}.$$

Furthermore, we have

$$C^{1+\frac{\alpha}{2},2+\alpha}([a,b]\times\mathbb{R}^d) = \left\{\varphi \in C^{1,2}([a,b]\times\mathbb{R}^d): \varphi_s, \varphi_{y_iy_j} \in C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d), \\ i,j=1,\ldots,d\right\}$$

with the following norm

$$\begin{aligned} |\varphi|_{C^{1+\frac{\alpha}{2},2+\alpha}([a,b]\times\mathbb{R}^d)} &= |\varphi|^{\infty} + |\varphi_s|_{C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d)} \\ &+ \sum_{i=1}^d |\varphi_{y_i}|^{\infty} + \sum_{i,j=1}^d |\varphi_{y_iy_j}|_{C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d)} \end{aligned}$$

For convenience, $|\cdot|_{C^{\frac{\alpha}{2},\alpha}([a,b]\times\mathbb{R}^d)}$ and $|\cdot|_{C^{1+\frac{\alpha}{2},2+\alpha}([a,b]\times\mathbb{R}^d)}$ are usually denoted as $|\cdot|_{[a,b]\times\mathbb{R}^d}^{(\alpha)}$ and $|\cdot|_{[a,b]\times\mathbb{R}^d}^{(2+\alpha)}$. In addition, wherever no confusion arises, we do not distinguish between $|\varphi(\cdot)|_{[a,b]\times\mathbb{R}^d}^{(\alpha)}$ and $|\varphi(\cdot)|_{\mathbb{R}^d}^{(\alpha)}$ and between $|\varphi(\cdot)|_{[a,b]\times\mathbb{R}^d}^{(2+\alpha)}$ and $|\varphi(\cdot)|_{\mathbb{R}^d}^{(2+\alpha)}$ for functions $\varphi(y)$ independent of s.

After reviewing the classical Hölder spaces, we revise on top of them to fulfil requirements of nonlocal PDEs and TIC problems. Before that, we first investigate the following two essential features of nonlocal PDEs and TIC problems.

(The order relation between t and s) Since both TIC control problems and FBSVIEs are backward problems, t and s usually represent the initial time and the running time, respectively, where 0 ≤ t ≤ s ≤ T. Considering a symmetry between forward and backward problems, we can formulate our nonlocal PDEs as forward problems and consider a reverse order relation, i.e. 0 ≤ s ≤ t ≤ T. Given the relation between t and s, we are concerned only with functions defined over the triangle Δ[0, δ] as illustrated in Figure 1 instead of a rectangle [0, δ]²;




Figure 1: Time domain $\Delta[0, \delta]$ of solutions of nonlocal PDEs

2. (Regularity and smoothness) Inspired by parabolic Hölder spaces for local PDEs, it is natural to preserve sufficient smoothness of the functions with respect to the variables s and y. Specifically, we will study the functions in space C^{1+α/2,2+α}. Next, it is essential to determine regularity of the external parameter t of u(t, ·, ·). In this paper, we require that u(t, s, y) is C¹ continuous with respect to t. Two main reasons behind the requirement are as follows: (a) when constructing the contractions and estimating solutions of nonlocal PDEs, we often encounter the evaluation of the difference |φ(s, s, y) – φ(s', s', y)|. The differentiability of φ in t makes some theorems like mean-value theorem available; (b) More importantly, the regularity of u with respect to t can help us convert nonlocal linear PDEs (10) into a system of coupled nonlocal PDEs (15) for (u, u_t). Consequently, the difficulty of studying the original nonlocal linear PDEs by fixed-point arguments can be resolved by investigating the induced system. Informally speaking, our successful settlement depends substantially on the contribution of the differentiability in t to the establishment of contractions.

After the preliminary analysis, we begin to define the norms and Banach spaces for nonlocal PDEs and TIC problems. For $0 \le \delta \le T$, we introduce norms

$$\begin{split} & [\phi]_{[0,\delta]}^{(k+\alpha)} & := \sup_{t \in [0,\delta]} \left\{ |\phi(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(k+\alpha)} \right\}, \\ & \|\phi\|_{[0,\delta]}^{(k+\alpha)} & := \sup_{t \in [0,\delta]} \left\{ |\phi(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(k+\alpha)} + |\phi_t(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(k+\alpha)} \right\}, \end{split}$$

where k = 0, 1, 2. Then they induce the following normed spaces, respectively,

$$\begin{split} \Theta_{[0,\delta]}^{(k+\alpha)} &:= \left\{ \theta(\cdot,\cdot,\cdot) \in C(\Delta[0,\delta] \times \mathbb{R}^d) : [\theta]_{[0,\delta]}^{(k+\alpha)} < \infty \right\}, \\ \Omega_{[0,\delta]}^{(k+\alpha)} &:= \left\{ \omega(\cdot,\cdot,\cdot) \in C(\Delta[0,\delta] \times \mathbb{R}^d) : \|\omega\|_{[0,\delta]}^{(k+\alpha)} < \infty \right\} \end{split}$$

It is easy to see that both $[\phi]_{[0,\delta]}^{(k+\alpha)}$ and $\|\phi\|_{[0,\delta]}^{(k+\alpha)}$ are norms under which $\Theta_{[0,\delta]}^{(k+\alpha)}$ and $\Omega_{[0,\delta]}^{(k+\alpha)}$ are Banach spaces, respectively. The definitions above leverage not only the order relation between t and s but also the sufficient regularities in all arguments.

2.2. Wellposedness of Nonlocal Linear Parabolic PDEs

Now, we are ready to study the solvability of nonlocal linear parabolic PDEs with the form

$$\begin{cases} u_{s}(t,s,y) = \sum_{i,j=1}^{d} a_{ij}(t,s,y)u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} b_{i}(t,s,y)u_{y_{i}}(t,s,y) + c(t,s,y)u(t,s,y) \\ + \sum_{i,j=1}^{d} \overline{a}_{ij}(t,s,y)u_{y_{i}y_{j}}(s,s,y) + \sum_{i=1}^{d} \overline{b}_{i}(t,s,y)u_{y_{i}}(s,s,y) + \overline{c}(t,s,y)u(s,s,y) \\ + f(t,s,y), \\ u(t,0,y) = g(t,y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(10)$$

where all coefficients $a, \overline{a}, b, \overline{b}, c$, and \overline{c} belong to $\Omega_{[0,T]}^{(\alpha)}$ and satisfy the uniform ellipticity condition, i.e., there exists some $\lambda > 0$ such that

$$\sum_{i,j=1}^{d} a_{ij}(t,s,y)\xi_i\xi_j \geq \lambda |\xi|^2,$$
(11)

$$\sum_{i,j=1}^{d} \left(a_{ij}(t,s,y) + \overline{a}_{ij}(t,s,y) \right) \xi_i \xi_j \geq \lambda |\xi|^2,$$
(12)

for any $(t,s) \in \Delta[0,T]$ and $y, \xi \in \mathbb{R}^d$. Moreover, the nonhomogeneous term $f \in \Omega_{[0,T]}^{(\alpha)}$ and the initial condition $g \in \Omega_{[0,T]}^{(2+\alpha)}$.

Suppose that u is a solution of (10), then its first-order derivative $\frac{\partial u}{\partial t}$ with respect to t should satisfy the following differential equation:

$$\begin{cases} \left(\frac{\partial u}{\partial t}\right)_{s}\left(t,s,y\right) = \sum_{i,j=1}^{d} a_{ij}(\cdot) \left(\frac{\partial u}{\partial t}\right)_{y_{i}y_{j}}\left(t,s,y\right) + \sum_{i=1}^{d} b_{i}(\cdot) \left(\frac{\partial u}{\partial t}\right)_{y_{i}}\left(t,s,y\right) + c(\cdot) \left(\frac{\partial u}{\partial t}\right)\left(t,s,y\right) \\ + \sum_{i,j=1}^{d} \frac{\partial a_{ij}(\cdot)}{\partial t} u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} \frac{\partial b_{i}(\cdot)}{\partial t} u_{y_{i}}(t,s,y) + c_{t}(\cdot)u(t,s,y) \\ + \sum_{i,j=1}^{d} \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} u_{y_{i}y_{j}}(s,s,y) + \sum_{i=1}^{d} \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} u_{y_{i}}(s,s,y) + \overline{c}_{t}(\cdot)u(s,s,y) + f_{t}(\cdot), \\ u_{t}(t,0,y) = g_{t}(t,y), \quad 0 \leq s \leq t \leq T, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(13)$$

where we use a convention that $\phi(\cdot) = \phi(t, s, y)$. Next, for any $i, j = 1, \ldots, d$, we notice the

following integral representations

$$\begin{cases} u(t,s,y) - u(s,s,y) = \int_{s}^{t} \frac{\partial u}{\partial t}(\theta,s,y)d\theta, \\ u_{y_{i}}(t,s,y) - u_{y_{i}}(s,s,y) = \int_{s}^{t} \left(\frac{\partial u}{\partial t}\right)_{y_{i}}(\theta,s,y)d\theta, \\ u_{y_{i}y_{j}}(t,s,y) - u_{y_{i}y_{j}}(s,s,y) = \int_{s}^{t} \left(\frac{\partial u}{\partial t}\right)_{y_{i}y_{j}}(\theta,s,y)d\theta. \end{cases}$$
(14)

(14) allows us to rewrite the diagonal terms in (10) and (13) as integrals of $\frac{\partial u}{\partial t}$ or its derivatives. Hence, by coupling equations (10) and (13) with (14), we obtain a nonlocal linear PDE system for $(u, v := \frac{\partial u}{\partial t})$:

$$\begin{cases} u_{s}(t,s,y) = \sum_{i,j=1}^{d} \left(a_{ij}(\cdot) + \overline{a}_{ij}(\cdot)\right) u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} \left(b_{i}(\cdot) + \overline{b}_{i}(\cdot)\right) u_{y_{i}}(t,s,y) \\ - \sum_{i,j=1}^{d} \overline{a}_{ij}(\cdot) \int_{s}^{t} v_{y_{i}y_{j}}(\theta,s,y) d\theta - \sum_{i=1}^{d} \overline{b}_{i}(\cdot) \int_{s}^{t} v_{y_{i}}(\theta,s,y) d\theta \\ + \left(c(\cdot) + \overline{c}(\cdot)\right) u(t,s,y) - \overline{c}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f(\cdot), \\ v_{s}(t,s,y) = \sum_{i,j=1}^{d} a_{ij}(\cdot) v_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} b_{i}(\cdot) v_{y_{i}}(t,s,y) + c(\cdot) v(t,s,y) \\ + \sum_{i,j=1}^{d} \left(\frac{\partial a_{ij}(\cdot)}{\partial t} + \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t}\right) u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} \left(\frac{\partial b_{i}(\cdot)}{\partial t} + \frac{\partial \overline{b}_{i}(\cdot)}{\partial t}\right) u_{y_{i}}(t,s,y) \\ - \sum_{i,j=1}^{d} \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}y_{j}}(\theta,s,y) d\theta - \sum_{i=1}^{d} \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}}(\theta,s,y) d\theta \\ + \left(c_{t}(\cdot) + \overline{c}_{t}(\cdot)\right) u(t,s,y) - \overline{c}_{t}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f_{t}(\cdot), \\ (u,v) (t,0,y) = (g,g_{t}) (t,y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(15)$$

Before studying the well-posedness of the system (15) for (u, v), we first prove the equivalence between (10) and (15) in the following lemma.

Lemma 2.1. The nonlocal equation (10) is equivalent to the system (15), i.e.,

- 1. If u is a solution of (10), then (u, u_t) solves (15).
- 2. Conversely, if (15) admits a solution pair (u, v), then u solves (10) and $v = u_t$.

Proof. The first claim is straightforward as it follows how we derived the equations (15) before. Next, we are focused on proving the second claim. Let (u, v) be a solution of (15). Differentiating the first equation of (15) with respect to t gives

$$\begin{pmatrix} \frac{\partial u}{\partial t} \end{pmatrix}_{s}(t,s,y) = \sum_{i,j=1}^{d} \left(a_{ij}(\cdot) + \overline{a}_{ij}(\cdot) \right) \left(\frac{\partial u}{\partial t} \right)_{y_{i}y_{j}}(t,s,y) + \sum_{i,j=1}^{d} \left(\frac{\partial a_{ij}(\cdot)}{\partial t} + \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \right) u_{y_{i}y_{j}}(t,s,y)$$

$$+ \sum_{i=1}^{d} \left(b_{i}(\cdot) + \overline{b}_{i}(\cdot) \right) \left(\frac{\partial u}{\partial t} \right)_{y_{i}}(t,s,y) + \sum_{i=1}^{d} \left(\frac{\partial b_{i}(\cdot)}{\partial t} + \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \right) u_{y_{i}}(t,s,y)$$

$$+ \left(c(\cdot) + \overline{c}(\cdot) \right) \left(\frac{\partial u}{\partial t} \right) (t,s,y) + \left(\frac{\partial c(\cdot)}{\partial t} + \frac{\partial \overline{c}(\cdot)}{\partial t} \right) u(t,s,y)$$

$$- \sum_{i,j=1}^{d} \overline{a}_{ij}(\cdot) v_{y_{i}y_{j}}(t,s,y) - \sum_{i,j=1}^{d} \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}y_{j}}(\theta,s,y) d\theta$$

$$- \sum_{i=1}^{d} \overline{b}_{i}(\cdot) v_{y_{i}}(t,s,y) - \sum_{i=1}^{d} \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}}(\theta,s,y) d\theta$$

$$- \overline{c}(\cdot) v(t,s,y) - \frac{\partial \overline{c}(\cdot)}{\partial t} \int_{s}^{t} v(\theta,s,y) d\theta + f_{t}(\cdot),$$

$$\left(\frac{\partial u}{\partial t} \right) (t,0,y) = g_{t}(t,y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^{d}.$$

$$(16)$$

From (16) and (15), we can find that $\left(\frac{\partial u}{\partial t} - v\right)(t, s, y)$ satisfies:

$$\begin{cases} \left(\frac{\partial u}{\partial t} - v\right)_{s}\left(t, s, y\right) = \sum_{i,j=1}^{d} \left(a_{ij}(\cdot) + \overline{a}_{ij}(\cdot)\right) \left(\frac{\partial u}{\partial t} - v\right)_{y_{i}y_{j}}\left(t, s, y\right) \\ + \sum_{i=1}^{d} \left(b_{i}(\cdot) + \overline{b}_{i}(\cdot)\right) \left(\frac{\partial u}{\partial t} - v\right)_{y_{i}}\left(t, s, y\right) \\ + \left(c(\cdot) + \overline{c}(\cdot)\right) \left(\frac{\partial u}{\partial t} - v\right)\left(t, s, y\right) \\ \left(\frac{\partial u}{\partial t} - v\right)\left(t, 0, y\right) = 0, \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(17)$$

which implies that $\left(\frac{\partial u}{\partial t} - v\right)(t, s, y) \equiv 0$, thanks to the classical PDE theory.

Next, we replace v in the first equation of (15) with $\frac{\partial u}{\partial t}$. Then we have

$$\begin{aligned} u_s(t,s,y) &= \sum_{i,j=1}^d \left(a_{ij}(\cdot) + \overline{a}_{ij}(\cdot) \right) u_{y_i y_j}(t,s,y) + \sum_{i=1}^d \left(b_i(\cdot) + \overline{b}_i(\cdot) \right) u_{y_i}(t,s,y) \\ &- \sum_{i,j=1}^d \overline{a}_{ij}(\cdot) \int_s^t \left(\frac{\partial u}{\partial t} \right)_{y_i y_j} (\theta, s, y) d\theta - \sum_{i=1}^d \overline{b}_i(\cdot) \int_s^t \left(\frac{\partial u}{\partial t} \right)_{y_i} (\theta, s, y) d\theta \\ &+ \left(c(\cdot) + \overline{c}(\cdot) \right) u(t,s,y) - \overline{c}(\cdot) \int_s^t \left(\frac{\partial u}{\partial t} \right) (\theta, s, y) d\theta + f(\cdot) \\ &= \sum_{i,j=1}^d a_{ij}(\cdot) u_{y_i y_j}(t,s,y) + \sum_{i=1}^d b_i(\cdot) u_{y_i}(t,s,y) + c(\cdot) u(t,s,y) \\ &+ \sum_{i,j=1}^d \overline{a}_{ij}(\cdot) u_{y_i y_j}(s,s,y) + \sum_{i=1}^d \overline{b}_i(\cdot) u_{y_i}(s,s,y) + \overline{c}(\cdot) u(s,s,y) + f(\cdot) \end{aligned}$$

Therefore, the result follows.

With the equivalence between (10) and (15), the well-posedness of (15) guarantees the solvability of (10). Next, for the coupled system for (u, v) induced from (10), we have the following conclusions.

Theorem 2.1. Suppose that all coefficient functions and f of (10) belong to $\Omega_{[0,T]}^{(\alpha)}$ and assume that $g \in \Omega_{[0,T]}^{(2+\alpha)}$. Then there exist $\delta > 0$ and a unique solution pair $(u, v) \in \Theta_{[0,\delta]}^{(2+\alpha)} \times \Theta_{[0,\delta]}^{(2+\alpha)}$ satisfying (15) in $\Delta[0, \delta] \times \mathbb{R}^d$.

Proof. The theorem is proven with fixed-point arguments and the overall idea is as follows. We first establish a contraction Γ defined over a closed subset \mathcal{V} with a radius R in $\Theta_{[0,\delta]}^{(2+\alpha)}$ and then show that it admits a unique fixed point in \mathcal{V} . Finally, a contradiction is constructed to argue for the uniqueness of the solution to the system (15) in $\Theta_{[0,\delta]}^{(2+\alpha)} \times \Theta_{[0,\delta]}^{(2+\alpha)}$.

(Existence) We make use of (15) to construct a mapping from a conservative vector field v to a conservative vector field V by $\Gamma(v) = V$, where V is part of the solution pair (u, V) to a system of equations:

$$\begin{cases} u_{s}(t,s,y) = \sum_{i,j=1}^{d} \left(a_{ij}(\cdot) + \overline{a}_{ij}(\cdot)\right) u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} \left(b_{i}(\cdot) + \overline{b}_{i}(\cdot)\right) u_{y_{i}}(t,s,y) \\ - \sum_{i,j=1}^{d} \overline{a}_{ij}(\cdot) \int_{s}^{t} v_{y_{i}y_{j}}(\theta,s,y) d\theta - \sum_{i=1}^{d} \overline{b}_{i}(\cdot) \int_{s}^{t} v_{y_{i}}(\theta,s,y) d\theta \\ + \left(c(\cdot) + \overline{c}(\cdot)\right) u(t,s,y) - \overline{c}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f(\cdot), \end{cases}$$

$$V_{s}(t,s,y) = \sum_{i,j=1}^{d} a_{ij}(\cdot) V_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} b_{i}(\cdot) V_{y_{i}}(t,s,y) + c(\cdot) V(t,s,y) \\ + \sum_{i,j=1}^{d} \left(\frac{\partial a_{ij}(\cdot)}{\partial t} + \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t}\right) u_{y_{i}y_{j}}(t,s,y) + \sum_{i=1}^{d} \left(\frac{\partial b_{i}(\cdot)}{\partial t} + \frac{\partial \overline{b}_{i}(\cdot)}{\partial t}\right) u_{y_{i}}(t,s,y) \\ - \sum_{i,j=1}^{d} \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}y_{j}}(\theta,s,y) d\theta - \sum_{i=1}^{d} \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \int_{s}^{t} v_{y_{i}}(\theta,s,y) d\theta \\ + \left(c_{t}(\cdot) + \overline{c}_{t}(\cdot)\right) u(t,s,y) - \overline{c}_{t}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f_{t}(\cdot), \\ (u,V)(t,0,y) = (g,g_{t})(t,y), \quad 0 \le s \le t \le \delta, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(18)$$

The operator $\Gamma(v) = V$ is defined in the set

$$\mathcal{V} = \left\{ v \in \Theta_{[0,\delta]}^{(2+\alpha)} : v(t,0,y) = g_t(t,y), \ [v-g]_{[0,\delta]}^{(2+\alpha)} \le R \right\}.$$
17

It is clear that given a conservative vector field v satisfying the initial condition $g_t(t, y)$, the system (18) admits a unique solution V and thus $\Gamma(\cdot)$ is well-defined.

Next, we shall show that the mapping Γ has a unique fixed point. Let $v, \hat{v} \in \mathcal{V}$. Then, $w := \Gamma(v) - \Gamma(\hat{v})$ is part of the solution pair $(u - \hat{u}, w)$ to a system of equations:

$$\begin{cases} (u - \widehat{u})_{s}(t, s, y) = \sum_{i,j=1}^{d} (a_{ij}(\cdot) + \overline{a}_{ij}(\cdot)) (u - \widehat{u})_{y_{i}y_{j}}(t, s, y) \\ + \sum_{i=1}^{d} (b_{i}(\cdot) + \overline{b}_{i}(\cdot)) (u - \widehat{u})_{y_{i}}(t, s, y) \\ - \sum_{i,j=1}^{d} \overline{a}_{ij}(\cdot) \int_{s}^{t} (v - \widehat{v})_{y_{i}y_{j}}(\theta, s, y) d\theta - \sum_{i=1}^{d} \overline{b}_{i}(\cdot) \int_{s}^{t} (v - \widehat{v})_{y_{i}}(\theta, s, y) d\theta \\ + (c(\cdot) + \overline{c}(\cdot)) (u - \widehat{u}) (t, s, y) - \overline{c}(\cdot) \int_{s}^{t} (v - \widehat{v}) (\theta, s, y) d\theta, \\ w_{s}(t, s, y) = \sum_{i,j=1}^{d} a_{ij}(\cdot) w_{y_{i}y_{j}}(t, s, y) + \sum_{i=1}^{d} b_{i}(\cdot) w_{y_{i}}(t, s, y) + c(\cdot) w(t, s, y) \\ + \sum_{i,j=1}^{d} \left(\frac{\partial a_{ij}(\cdot)}{\partial t} + \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \right) (u - \widehat{u})_{y_{i}y_{j}}(t, s, y) \\ + \sum_{i=1}^{d} \left(\frac{\partial b_{i}(\cdot)}{\partial t} + \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \right) (u - \widehat{u})_{y_{i}}(t, s, y) \\ - \sum_{i,j=1}^{d} \frac{\partial \overline{a}_{ij}(\cdot)}{\partial t} \int_{s}^{t} (v - \widehat{v})_{y_{i}y_{j}}(\theta, s, y) d\theta - \sum_{i=1}^{d} \frac{\partial \overline{b}_{i}(\cdot)}{\partial t} \int_{s}^{t} (v - \widehat{v})_{y_{i}}(\theta, s, y) d\theta \\ + (c_{t}(\cdot) + \overline{c}_{t}(\cdot)) (u - \widehat{u}) (t, s, y) - \overline{c}_{t}(\cdot) \int_{s}^{t} (v - \widehat{v}) (\theta, s, y) d\theta, \\ (u - \widehat{u}, w) (t, 0, y) = (0, 0), \quad 0 \le s \le t \le \delta, \quad y \in \mathbb{R}^{d}. \end{cases}$$

$$(19)$$

By the classical theory of parabolic PDEs [21, 22, 23], for a fix $t \in [0, \delta]$, we have the following estimates for the solution to the first equation of (19):

$$\begin{aligned} \left| (u - \widehat{u}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} \\ &\leq c \left(\sum_{i,j=1}^d \left| \int_{\cdot}^t (v - \widehat{v})_{y_i y_j} (\theta, \cdot, \cdot) d\theta \right|_{[0,t] \times \mathbb{R}^d}^{(\alpha)} + \sum_{i=1}^d \left| \int_{\cdot}^t (v - \widehat{v})_{y_i} (\theta, \cdot, \cdot) d\theta \right|_{[0,t] \times \mathbb{R}^d}^{(\alpha)} \\ &+ \left| \int_{\cdot}^t (v - \widehat{v}) (\theta, \cdot, \cdot) d\theta \right|_{[0,t] \times \mathbb{R}^d}^{(\alpha)} \right), \end{aligned}$$
(20)

where the constant c depends on $||a||_{[0,T]}^{(\alpha)}$, $||\overline{a}||_{[0,T]}^{(\alpha)}$, $||b||_{[0,T]}^{(\alpha)}$, $||\overline{b}||_{[0,T]}^{(\alpha)}$, $||c||_{[0,T]}^{(\alpha)}$ and $||\overline{c}||_{[0,T]}^{(\alpha)}$. To further estimate the upper bound of (20), we investigate the Hölder continuity of $\int_{s}^{t} (v - \hat{v})_{y_i y_j} (\theta, s, y) d\theta$ with respect to s and y for any y_i and y_j , while the similar analyses can be conducted for other two integral terms in (20).

Let $\delta \leq 1$ and $0 \leq s \leq s' \leq t \leq \delta \leq T$, we have

$$\begin{aligned} \left| \int_{s}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) d\theta - \int_{s'}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s', y) d\theta \right| \\ \leq \left| \int_{s}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) d\theta - \int_{s'}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) d\theta \right| \\ + \left| \int_{s'}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) d\theta - \int_{s'}^{t} (v - \hat{v})_{y_{i}y_{j}} (\theta, s', y) d\theta \right| \\ \leq \int_{s}^{s'} \left| (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) \right| d\theta + \int_{s'}^{t} \left| (v - \hat{v})_{y_{i}y_{j}} (\theta, s, y) - (v - \hat{v})_{y_{i}y_{j}} (\theta, s', y) \right| d\theta \\ \leq \left(\sup_{t \in [0,\delta]} \left\{ \left| (v - \hat{v}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} \right\} (s' - s)^{1-\frac{\alpha}{2}} \right. \\ \left. + \sup_{t \in [0,\delta]} \left\{ \left| (v - \hat{v}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} \right\} \delta^{1-\frac{\alpha}{2}} \cdot |s' - s|^{\frac{\alpha}{2}} \end{aligned} \right.$$

which implies also, by noting $(v - \hat{v}) (\cdot, 0, \cdot) \equiv 0$, that

$$\left|\int_{\cdot}^{t} (v-\widehat{v})_{y_i y_j} (\theta, \cdot, \cdot) d\theta\right|_{[0,t] \times \mathbb{R}^d}^{\infty} \leq \sup_{t \in [0,\delta]} \left\{ \left| (v-\widehat{v}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} \right\} \delta$$
(22)

Moreover, let $y, y' \in \mathbb{R}^d$, then we have

$$\left| \int_{s}^{t} (v - \widehat{v})_{y_{i}y_{j}} (\theta, s, y') d\theta - \int_{s}^{t} (v - \widehat{v})_{y_{i}y_{j}} (\theta, s, y) d\theta \right| \\
\leq \int_{s}^{t} \left| (v - \widehat{v})_{y_{i}y_{j}} (\theta, s, y') - (v - \widehat{v})_{y_{i}y_{j}} (\theta, s, y) \right| d\theta \\
\leq \sup_{t \in [0,\delta]} \left\{ \left| (v - \widehat{v}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} \right\} (t - s) \cdot |y' - y|^{\alpha} \\
\leq \sup_{t \in [0,\delta]} \left\{ \left| (v - \widehat{v}) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} \right\} \delta \cdot |y' - y|^{\alpha}.$$
(23)

Consequently, from (21), (22), and (23), we have

$$\left|\int_{\cdot}^{t} \left(v-\widehat{v}\right)_{y_{i}y_{j}}(\theta,\cdot,\cdot)d\theta\right|_{\left[0,t\right]\times\mathbb{R}^{d}}^{(\alpha)} \leq \delta^{1-\frac{\alpha}{2}} \cdot \sup_{t\in\left[0,\delta\right]}\left\{\left|\left(v-\widehat{v}\right)\left(t,\cdot,\cdot\right)\right|_{\left[0,t\right]\times\mathbb{R}^{d}}^{(2+\alpha)}\right\}\right\}$$

Similarly, we can also estimate other lower-order terms of (20). Then,

$$\left|\left(u-\widehat{u}\right)\left(t,\cdot,\cdot\right)\right|_{\left[0,t\right]\times\mathbb{R}^{d}}^{\left(2+\alpha\right)} \leq C_{1}\delta^{1-\frac{\alpha}{2}} \cdot \sup_{t\in\left[0,\delta\right]}\left\{\left|\left(v-\widehat{v}\right)\left(t,\cdot,\cdot\right)\right|_{\left[0,t\right]\times\mathbb{R}^{d}}^{\left(2+\alpha\right)}\right\}.$$
(24)

By the classical theory of parabolic PDEs and (24), we can also acquire the inequality from the second equation of (19):

$$|w(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} \le C_2 \delta^{1-\frac{\alpha}{2}} \cdot \sup_{t\in[0,\delta]} \left\{ |(v-\widehat{v})(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} \right\}.$$

Consequently, under the norm $[\phi]_{[0,\delta]}^{(2+\alpha)} := \sup_{t \in [0,\delta]} \left\{ |\phi(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} \right\}$, choosing a suitably small δ yields that

$$[\Gamma(v) - \Gamma(\widehat{v})]_{[0,\delta]}^{(2+\alpha)} = \sup_{t \in [0,\delta]} \left\{ |w(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} \right\} \le \frac{1}{2} \left[v - \widehat{v} \right]_{[0,\delta]}^{(2+\alpha)}.$$

(A contraction Γ mapping \mathcal{V} into itself.) On the other hand, we also have

$$[\Gamma(v) - g_t]_{[0,\delta]}^{(2+\alpha)} \le [\Gamma(v) - \Gamma(g_t)]_{[0,\delta]}^{(2+\alpha)} + [\Gamma(g_t) - g_t]_{[0,\delta]}^{(2+\alpha)}$$
$$\le \frac{1}{2} [v - g_t]_{[0,\delta]}^{(2+\alpha)} + C' \le \frac{R}{2} + C'$$

Therefore, for a suitably large R, Γ is a contraction mapping \mathcal{V} into itself and thus it has a unique fixed point v in \mathcal{V} such that $\Gamma(v) = v$.

(Uniqueness) To complete the proof, we ought to show that the conservative vector field v is the unique fixed point of (18) in $\Theta_{[0,\delta]}^{(2+\alpha)}$. It can be done with some standard arguments. If (18) admits two fixed points v^1 and v^2 , let

$$t_0 = \sup \left\{ t \in [0, \delta] : v^1(t, s, y) = v^2(t, s, y), (t, s, y) \in \Delta[0, t] \times \mathbb{R}^d \right\}.$$

We shall focus only on the case when $t_0 < \delta$ because if $t_0 = \delta$, then $v^1 = v^2$ in the whole $\Delta[0, \delta] \times \mathbb{R}^d$ and the proof is completed.

According to the definition of $t_0(<\delta)$, we know that $v^1(t,s,y) = v^2(t,s,y)$ in $\Delta_1 \times \mathbb{R}^d$ in Figure 2. Furthermore, their corresponding $u^1(t,s,y)$ and $u^2(t,s,y)$ are also equal in $\Delta_1 \times \mathbb{R}^d$ according to the first equation of (18). Hence, we obtain diagonal conditions, namely $u^1(s,s,y) =$ $u^2(s,s,y), u^1_y(s,s,y) = u^2_y(s,s,y)$ and $u^1_{yy}(s,s,y) = u^2_{yy}(s,s,y)$ for any $s \in [0, t_0]$ and $y \in \mathbb{R}^d$. By observing (10) and (13) provided that the same initial and diagonal conditions, i.e. the initial condition 1 and the diagonal condition in Figure 2, the classical PDE theory promises that (u^1, v^1) and (u^2, v^2) coincide in $(R \cup \Delta_1) \times \mathbb{R}^d$.




Figure 2: Uniqueness of the solution in $\Theta[0, \delta]$

Next, let $(u^1(t, t_0, y), v^1(t, t_0, y)) = (u^2(t, t_0, y), v^2(t, t_0, y)) = (g'(t, y), g'_t(t, y))$. Based on the new initial condition, i.e. the initial condition 2 in Figure 2, we consider the following initial value problem:

$$\begin{cases} u_{s}(t,s,y) = (a(\cdot) + \overline{a}(\cdot)) u_{yy}(t,s,y) + (b(\cdot) + \overline{b}(\cdot)) u_{y}(t,s,y) \\ -\overline{a}(\cdot) \int_{s}^{t} v_{yy}(\theta,s,y) d\theta - \overline{b}(\cdot) \int_{s}^{t} v_{y}(\theta,s,y) d\theta \\ + (c(\cdot) + \overline{c}(\cdot)) u(t,s,y) - \overline{c}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f(t,s,y), \end{cases}$$

$$v_{s}(t,s,y) = a(\cdot)v_{yy}(t,s,y) + b(\cdot)v_{y}(t,s,y) + c(\cdot)v(t,s,y) \\ + (a_{t}(\cdot) + \overline{a}_{t}(\cdot)) u_{yy}(t,s,y) + (b_{t}(\cdot) + \overline{b}_{t}(\cdot)) u_{y}(t,s,y) \\ -\overline{a}_{t}(\cdot) \int_{s}^{t} v_{yy}(\theta,s,y) d\theta - \overline{b}_{t}(\cdot) \int_{s}^{t} v_{y}(\theta,s,y) d\theta \\ + (c_{t}(\cdot) + \overline{c}_{t}(\cdot)) u(t,s,y) - \overline{c}_{t}(\cdot) \int_{s}^{t} v(\theta,s,y) d\theta + f_{t}(t,s,y), \end{cases}$$

$$(25)$$

$$(u,v) (t,t_{0},y) = (g',g'_{t})(t,y), \quad t_{0} \leq s \leq t \leq T, \quad y \in \mathbb{R}^{d}.$$

Our previous proof shows that (25) admits a unique v in the set

$$\mathcal{V}' = \left\{ v \in \Theta_{[t_0, t_0 + \delta']}^{(2+\alpha)} : v(t, t_0, y) = g'_t(t, y), [v - g']_{[t_0, t_0 + \delta']}^{(2+\alpha)} \le R' \right\}$$

provided that R' is large enough and δ' is small enough. Considering R' larger than $[v^1 - g']_{[t_0,t_0+\delta']}^{(2+\alpha)}$ and $[v^2 - g']_{[t_0,t_0+\delta']}^{(2+\alpha)}$, we have $v^1 = v^2$ in $\Delta_2 \times \mathbb{R}^d$. Hence, for any $y \in \mathbb{R}^d$, v^1 equals to v^2 in $\{(t,s): t_0 \le t \le t_0 + \delta', 0 \le s \le t_0]\} \cup \Delta_2$. This contradicts the definition of t_0 . Consequently, $t_0 = \delta$ and $v^1 = v^2$.

Finally, the unique fixed point v determines uniquely a function u(t, s, y) via (18). Moreover, given $v \in \Theta_{[0,\delta]}^{(2+\alpha)}$, it is clear that

$$[u]_{[0,\delta]\times\mathbb{R}^d}^{(2+\alpha)} \le c\left([v]_{[0,\delta]\times\mathbb{R}^d}^{(2+\alpha)} + [f]_{[0,\delta]\times\mathbb{R}^d}^{(\alpha)} + [g]_{[0,\delta]\times\mathbb{R}^d}^{(2+\alpha)}\right) < \infty.$$

Therefore, there exists a unique solution pair $(u, v) \in \Theta_{[0,\delta]}^{(2+\alpha)} \times \Theta_{[0,\delta]}^{(2+\alpha)}$ to (15) in $\Delta[0, \delta] \times \mathbb{R}^d$. \Box

Remark 2.1 (Extension of solutions to a larger time interval). Until now, we have proven the local existence of (u, v) in $R_1 = \{0 \le s \le t \le \delta\}$ in Figure 3. Hence, we can determine the




Figure 3: Extension from $\Delta[0, \delta]$ to a larger time interval

diagonal condition for $s \in [0, \delta]$. Then the nonlocal equations (10) and (13) reduce to classical PDEs with a parameter t. Therefore, we can extend uniquely our solution (u, v) from R_1 to $R_1 \cup R_2 = \{0 \le s \le \delta, s \le t \le T\}$ in Figure 3. Subsequently, we acquire a new initial condition at $s = \delta$ for $t \in [s, T]$. Taking δ as an initial time and $(u(t, \delta, y), v(t, \delta, y))$ as initial datum, one can extend the solution to a larger time intervals $R_1 \cup R_2 \cup R_3$ and then $R_1 \cup R_2 \cup R_3 \cup R_4$ as illustrated in Figure 3. Hence, we can extend uniquely the solution from $\Delta[0, \delta]$ to $\Delta[0, \overline{\delta}]$. The procedure could be repeated indefinitely, up to a *maximally defined solution* $(u, v) : \Delta[0, \sigma] \times \mathbb{R}^d \to \mathbb{R}$, belonging to $\Theta_{[0,\sigma]}^{(2+\alpha)} \times \Theta_{[0,\sigma]}^{(2+\alpha)}$ for any $\sigma < \tau$. The time region $\Delta[0, \tau]$ is maximal in the sense that if $\tau < \infty$, then there does not exist any solution of (15) belonging to $\Theta_{[0,\tau]}^{(2+\alpha)} \times \Theta_{[0,\tau]}^{(2+\alpha)}$. Next, for the nonlocal system (15) of (u, v) and a small enough $\delta \in [0, T]$, we give a Schaudertype estimate. It not only shows the stability of the solutions to (10) with respect to the data (f, g), but also establishes a foundation for the further analysis of nonlocal fully nonlinear equations in the next section.

Corollary 2.1. Suppose that all coefficient functions and f of (10) belong to $\Omega_{[0,T]}^{(\alpha)}$ and assume that $g \in \Omega_{[0,T]}^{(2+\alpha)}$. Then there exist $\delta > 0$ and a unique $u \in \Omega_{[0,\delta]}^{(2+\alpha)}$ satisfying (10) in $\Delta[0,\delta] \times \mathbb{R}^d$. Furthermore, the following holds

$$\sup_{t \in [0,\delta]} \left\{ |u(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} + |u_{t}(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} \right\}$$

$$\leq c \left(\sup_{t \in [0,\delta]} \left\{ |f(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^{d}}^{(\alpha)} + |f_{t}(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^{d}}^{(\alpha)} \right\} + \sup_{t \in [0,\delta]} \left\{ |g(t,\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)} + |g_{t}(t,\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)} \right\} \right)$$
(26)

which implies that

$$\|u\|_{[0,\delta]}^{(2+\alpha)} \le c \left(\|f\|_{[0,\delta]}^{(\alpha)} + \|g\|_{[0,\delta]}^{(2+\alpha)} \right).$$
(27)

Proof. As Lemma 2.1 shows, the first component u of the unique solution (u, v) of (15) solves the nonlocal linear equation (10). By noting of $v = u_t$ in (15), it is clear that $u \in \Omega_{[0,\delta]}^{(2+\alpha)}$ thanks to $(u, u_t) \in \Theta_{[0,\delta]}^{(2+\alpha)} \times \Theta_{[0,\delta]}^{(2+\alpha)}$. Moreover, by (15), we have

$$|u(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} \leq c \left(\delta^{1-\frac{\alpha}{2}} \sup_{t\in[0,\delta]} |u_{t}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} + \sup_{t\in[0,\delta]} |f(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(\alpha)} + \sup_{t\in[0,\delta]} |g(t,\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)} \right)$$

and

$$\begin{aligned} |u_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} &\leq c \left(|u(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} + \delta^{1-\frac{\alpha}{2}} \sup_{t\in[0,\delta]} |u_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} \\ &+ \sup_{t\in[0,\delta]} |f_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} + \sup_{t\in[0,\delta]} |g_t(t,\cdot)|_{\mathbb{R}^d}^{(2+\alpha)} \right). \end{aligned}$$

Consequently, for a small enough δ , it holds

which leads to (26) and (27).

Remark 2.2 (Stability of solutions with respect to data). The inequalities in Corollary 2.1 can be used to show that the map from data (f, g) to solutions of (10) is continuous in the $\|\cdot\|_{[0,\delta]}^{(2+\alpha)}$ topology. Specifically, let u and \hat{u} correspond to (f, g) and (\hat{f}, \hat{g}) satisfying these assumptions of Theorem 2.1, respectively. Then, we have

$$\|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)} \le c \left(\|f - \widehat{f}\|_{[0,\delta]}^{(\alpha)} + \|g - \widehat{g}\|_{[0,\delta]}^{(2+\alpha)} \right).$$

3. Nonlocal Fully Nonlinear Parabolic PDEs

After studying nonlocal linear PDEs (10) and acquiring a prior estimate of solutions (27), we further study nonlocal fully nonlinear PDEs:

$$\begin{cases} u_{s}(t,s,y) = F(t,s,y,u(t,s,y),u_{y}(t,s,y),u_{yy}(t,s,y), \\ u(s,s,y),u_{y}(s,s,y),u_{yy}(s,s,y)), \\ u(t,0,y) = g(t,y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^{d}. \end{cases}$$
(28)

where the mapping F is nonlinear with respect to all its arguments, s and y are temporal and spatial variables, respectively, while t could be considered as an external temporal parameter, and the temporal variables (s, t) are defined in a triangular region $\Delta[0, T]$.

To leverage the results of nonlocal linear PDEs in Section 2, we require certain regularity assumptions on F and g. Generally speaking, we need that the initial condition $g \in \Omega_{[0,T]}^{(2+\alpha)}$ and the nonlinear F maps the functions (about u) from $\Omega_{[0,T]}^{(2+\alpha)}$ to $\Omega_{[0,T]}^{(\alpha)}$. More specifically, we suppose that the real-valued function $\pi \mapsto F(\pi)$ is defined in $\Pi = \Delta[0,T] \times \mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d \times \mathbb{R} \times \mathbb{R}^d \times \mathbb{S}^d$ and satisfies that

(i) (Ellipticity condition) for any $\pi = (t, s, y, u, p, q, l, m, n) \in \Pi$ and $\xi = (\xi_1, \xi_2, \dots, \xi_n) \in \mathbb{R}^n \setminus \{0\}$,

$$\sum_{i,j=1}^{d} \frac{\partial F}{\partial q_{ij}}(t,s,y,u,p,q,l,m,n)\xi_i\xi_j > 0,$$
(29)

$$\sum_{i,j=1}^{d} \left(\frac{\partial F}{\partial q_{ij}} + \frac{\partial F}{\partial n_{ij}} \right) (t, s, y, u, p, q, l, m, n) \xi_i \xi_j > 0;$$
(30)

(ii) (Hölder continuity) for every $\delta \ge 0$ and $\overline{\pi} = (u, p, q, l, m, n)$, there exists a positive constant K > 0 such that

$$\sup_{(t,\overline{\pi})} \left\{ \left| \mathcal{F}(t,\cdot,\cdot,\overline{\pi}) \right|_{[0,\delta] \times \mathbb{R}^d}^{(\alpha)} \right\} = K;$$
(31)

(iii) (Lipschitz continuity) for any $(t_1, s, y, \overline{\pi}_1), (t_2, s, y, \overline{\pi}_2) \in \Pi$, there exists a positive constant L > 0 such that

$$|\mathcal{F}(t_1, s, y, \overline{\pi}_1) - \mathcal{F}(t_2, s, y, \overline{\pi}_2)| \le L \left(|t_1 - t_2| + |\overline{\pi}_1 - \overline{\pi}_2| \right),$$
(32)

where \mathcal{F} represents F itself and some of its first- and second-order derivatives ($F_{\mathcal{X}}$ and $F_{\mathcal{X}\mathcal{Y}}$), which are denoted by " $\sqrt{}$ " in Table 1 and Table 2.

X	t	s	y	u	p	q	l	m	n
$F_{\mathcal{X}}$				\checkmark				\checkmark	

Table 1: First-order derivatives of F required to be Hölder and Lipschitz continuous

$ \begin{array}{c c} \hline F_{\mathcal{X}\mathcal{Y}} & \mathcal{Y} \\ \hline \mathcal{X} & \end{array} $	t	s	y	u	p	q	l	m	n
t									
S									
y									
u				\checkmark	\checkmark	\checkmark		\checkmark	\checkmark
p				\checkmark	\checkmark	\checkmark		\checkmark	\checkmark
q				\checkmark	\checkmark	\checkmark	\checkmark	\checkmark	\checkmark
l				\checkmark	\checkmark	\checkmark			
m					\checkmark				
n				\checkmark	\checkmark	\checkmark			

Table 2: Second-order derivatives of F required to be Hölder and Lipschitz continuous

Before we present our main result, we stress that the standard linearization methods are not applicable for the nonlocal case. In the setting of local parabolic PDEs, [24] introduced a so-called

"quasi-linearization method" and studied local existence for fully nonlinear parabolic problems by transforming fully nonlinear equations into quasi-linear systems. Furthermore, [25, 26] utilized a variant of this method to investigate fully nonlinear PDEs. The linearization method, which we have used in Theorem 3.1 below, is substantially inspired by [27, 23]. Although there are some previous works on how to linearize nonlinear equations, it is still difficult to extend their methods from local to nonlocal cases, mainly due to the limited progress on studying the nonlocal linear PDEs (10).

Next, we prove the well-posedness of nonlocal fully nonlinear PDEs (28).

Theorem 3.1. Suppose that (29)-(32) hold and assume that $g \in \Omega_{[0,T]}^{(2+\alpha)}$. Then, there exist $\delta > 0$ and a unique $u \in \Omega_{[0,\delta]}^{(2+\alpha)}$ satisfying (28) in $\Delta[0,\delta] \times \mathbb{R}^d$.

Proof. We prove the theorem again with fixed-point arguments.

(Existence) The solution of (28) is found via a fixed point of the operator Λ defined in the set

$$\mathcal{U} = \left\{ u \in \Omega_{[0,\delta]}^{(2+\alpha)} : u(t,0,y) = g(t,y), \|u - g\|_{[0,\delta]}^{(2+\alpha)} \le R \right\}$$

and $\Lambda(u) = U$, where U is the solution of

$$\begin{cases} U_{s}(t,s,y) = \mathcal{L}U + F(t,s,y,u(t,s,y),u_{y}(t,s,y),u_{yy}(t,s,y), \\ u(s,s,y),u_{y}(s,s,y),u_{yy}(s,s,y)) - \mathcal{L}u, \\ U(t,0,y) = g(t,y), \quad 0 \le s \le t \le \delta, \quad y \in \mathbb{R}^{d}. \end{cases}$$
(33)

where \mathcal{L} is a linear operator given by

$$\mathcal{L}u = \sum_{i,j=1}^{d} F_{q_{ij}} \frac{\partial^2 u}{\partial y_i \partial y_j} + \sum_{i=1}^{d} F_{p_i} \frac{\partial u}{\partial y_i} + F_u u + \sum_{i,j=1}^{d} F_{n_{ij}} \left(\frac{\partial^2 u}{\partial y_i \partial y_j} \Big|_{t=s} \right) + \sum_{i=1}^{d} F_{m_i} \left(\frac{\partial u}{\partial y_i} \Big|_{t=s} \right) + F_l \left(u \Big|_{t=s} \right)$$
(34)

with the partial derivatives F_u , F_p , F_q , F_l , F_m and F_n of F(t, s, y, u, p, q, l, m, n) evaluated at the point $(t, 0, y, g(t, y), g_y(t, y), g_{yy}(t, y), g(0, y), g_y(0, y), g_{yy}(0, y))$. It is clear that all coefficients of \mathcal{L} belong to $\Omega_{[0,\delta]}^{(\alpha)}$ and $F - \mathcal{L}u$ in (33) also belongs to $\Omega_{[0,\delta]}^{(\alpha)}$. Hence, by the well-posedness result of nonlocal linear PDEs in Section 2.2, the mapping $\Lambda(u)$ from $\Omega_{[0,\delta]}^{(2+\alpha)}$ to $\Omega_{[0,\delta]}^{(2+\alpha)}$ is well-defined.

Generally speaking, we shall show that for every $u, \widehat{u} \in \mathcal{U},$

$$\|\Lambda(u) - \Lambda(\widehat{u})\|_{[0,\delta]}^{(2+\alpha)} \le C(R)\delta^{\frac{\alpha}{2}} \|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$

Let $u,\widehat{u}\in\mathcal{U}.$ Then $w=\Lambda(u)-\Lambda(\widehat{u})$ satisfies

$$\begin{array}{lll} w_{s}(t,s,y) &=& \mathcal{L}w + F\big(t,s,y,u(t,s,y),u_{y}(t,s,y),u_{yy}(t,s,y),\\ && u(s,s,y),u_{y}(s,s,y),u_{yy}(s,s,y)\big) \\ && -F\big(t,s,y,\widehat{u}(t,s,y),\widehat{u}_{y}(t,s,y),\widehat{u}_{yy}(t,s,y),\\ && \widehat{u}(s,s,y),\widehat{u}_{y}(s,s,y),\widehat{u}_{yy}(s,s,y)\big) - \mathcal{L}(u-\widehat{u}),\\ && w(t,0,y) &=& 0, \quad 0 \leq s \leq t \leq \delta, \quad y \in \mathbb{R}^{d}. \end{array}$$

By (26) and (27), there is c > 0, independent of δ , such that

$$\sup_{t\in[0,\delta]} \left\{ |w(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} + |w_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(2+\alpha)} \right\}$$
$$\leq c \sup_{t\in[0,\delta]} \left\{ |\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} + |\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \right\},$$

where

$$\begin{split} \varphi(t,s,y) &= F(t,s,y,u(t,s,y),u_y(t,s,y),u_{yy}(t,s,y), \\ & u(s,s,y),u_y(s,s,y),u_{yy}(s,s,y)) \\ & -F(t,s,y,\widehat{u}(t,s,y),\widehat{u}_y(t,s,y),\widehat{u}_{yy}(t,s,y), \\ & \widehat{u}(s,s,y),\widehat{u}_y(s,s,y),\widehat{u}_{yy}(s,s,y)) - \mathcal{L}(u-\widehat{u}). \end{split}$$

In order to estimate $|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}$ and $|\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}$, it is convenient to write $\varphi(t,s,y)$ as

an integral representation

$$\begin{aligned} \varphi(t,s,y) & (35) \\ = \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y)) \right) \left(u(t,s,y) - \widehat{u}(t,s,y) \right) d\sigma \\ &+ \int_0^1 \sum_{i=1}^d \left(F_{p_i}(t,s,y,\theta_\sigma(t,s,y)) - F_{p_i}(t,0,y,\theta_0(t,y)) \right) \left(u_{y_i}(t,s,y) - \widehat{u}_{y_i}(t,s,y) \right) d\sigma \\ &+ \int_0^1 \sum_{i,j=1}^d \left(F_{q_{ij}}(t,s,y,\theta_\sigma(t,s,y)) - F_{q_{ij}}(t,0,y,\theta_0(t,y)) \right) \left(u_{y_iy_j}(t,s,y) - \widehat{u}_{y_iy_j}(t,s,y) \right) d\sigma \\ &+ \int_0^1 \left(F_l(t,s,y,\theta_\sigma(t,s,y)) - F_l(t,0,y,\theta_0(t,y)) \right) \left(u(s,s,y) - \widehat{u}(s,s,y) \right) d\sigma \\ &+ \int_0^1 \sum_{i=1}^d \left(F_{m_i}(t,s,y,\theta_\sigma(t,s,y)) - F_{m_i}(t,0,y,\theta_0(t,y)) \right) \left(u_{y_iy_j}(s,s,y) - \widehat{u}_{y_iy_j}(s,s,y) \right) d\sigma \\ &+ \int_0^1 \sum_{i,j=1}^d \left(F_{m_i}(t,s,y,\theta_\sigma(t,s,y)) - F_{m_i}(t,0,y,\theta_0(t,y)) \right) \left(u_{y_iy_j}(s,s,y) - \widehat{u}_{y_iy_j}(s,s,y) \right) d\sigma \end{aligned}$$

with $\theta_0(t,y) = \left(g(t,y), g_y(t,y), g_{yy}(t,y), g(0,y), g_y(0,y), g_{yy}(0,y)\right)$ and

$$\theta_{\sigma}(t,s,y) = \sigma \big(u(t,s,y), u_y(t,s,y), u_{yy}(t,s,y), u(s,s,y), u_y(s,s,y), u_{yy}(s,s,y) \big) \\ + (1-\sigma) \big(\widehat{u}(t,s,y), \widehat{u}_y(t,s,y), \widehat{u}_{yy}(t,s,y), \widehat{u}(s,s,y), \widehat{u}_y(s,s,y), \widehat{u}_{yy}(s,s,y) \big).$$

 $\delta \leq T$ and $y \in \mathbb{R}^d$. It is convenient to add and subtract the following to $\varphi(t, s, y) - \varphi(t, s', y)$:

$$\begin{split} &\int_{0}^{1} F_{u}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u(t,s,y) - \hat{u}(t,s,y)) d\sigma \\ &+ \int_{0}^{1} \sum_{i=1}^{d} F_{p}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u_{y_{i}}(t,s,y) - \hat{u}_{y_{i}}(t,s,y)) d\sigma \\ &+ \int_{0}^{1} \sum_{i,j=1}^{d} F_{q_{ij}}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u_{y_{i}y_{j}}(t,s,y) - \hat{u}_{y_{i}y_{j}}(t,s,y)) d\sigma \\ &+ \int_{0}^{1} F_{l}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u(s,s,y) - \hat{u}(s,s,y)) d\sigma \\ &+ \int_{0}^{1} \sum_{i=1}^{d} F_{m_{i}}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s,s,y)) d\sigma \\ &+ \int_{0}^{1} \sum_{i,j=1}^{d} F_{m_{i}}(t,s',y,\theta_{\sigma}(t,s',y)) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \end{split}$$

Let F_{χ} be any first order derivatives of F with respect to (u, p, q, l, m, n). We have

$$\begin{split} |F_{\chi}(t,s,y,\theta_{\sigma}(t,s,y)) - F_{\chi}(t,s',y,\theta_{\sigma}(t,s',y))| &\leq L \left| \theta_{\sigma}(t,s,y) - \theta_{\sigma}(t,s',y) \right| + K(s'-s)^{\frac{\alpha}{2}} \\ &\leq L \left(\sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(t,s,y) - u_{y_{i}y_{j}}(t,s',y) \right| + \sum_{i,j=1}^{d} \left| u_{y_{i}j}(t,s,y) - u_{y_{i}}(t,s',y) \right| \\ &+ \left| u(t,s,y) - u(t,s',y) \right| + \sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(s,s,y) - u_{y_{i}y_{j}}(s',s',y) \right| \\ &+ \sum_{i=1}^{d} \left| u_{y_{i}}(s,s,y) - u_{y_{i}}(s',s',y) \right| + \left| u(s,s,y) - u(s',s',y) \right| \right) \\ &+ L \left(\sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(t,s,y) - \hat{u}_{y_{i}y_{j}}(t,s',y) \right| + \sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s',s',y) \right| \\ &+ \left| \hat{u}(t,s,y) - \hat{u}(t,s',y) \right| + \sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s',s',y) \right| \\ &+ \left| \hat{u}_{i}(t,s,y) - \hat{u}(t,s',y) \right| + \sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s',s',y) \right| \\ &+ \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s',s',y) \right| + \left| \hat{u}(s,s,y) - \hat{u}(s',s',y) \right| \right) + K(s'-s)^{\frac{\alpha}{2}} \\ &\leq L \left(\left| u(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}(s'-s)^{\frac{\alpha}{2}} + \sup_{\overline{s}\in(s,s')} \left| u_{t}(\overline{s},\cdot,\cdot) \right|_{[0,\overline{s}]\times\mathbb{R}^{d}}(s'-s)^{\frac{\alpha}{2}} \\ &+ \sup_{\overline{s}\in(s,s')} \left| \hat{u}_{t}(\overline{s},\cdot,\cdot) \right|_{[0,\overline{s}]\times\mathbb{R}^{d}}(s'-s)^{\frac{\alpha}{2}} + \left| \hat{u}(s',\cdot,\cdot) \right|_{[0,\overline{s}]\times\mathbb{R}^{d}}(s'-s)^{\frac{\alpha}{2}} \\ &\leq \left(K + L \left(\left\| u \right\|_{[0,\delta]}^{(2+\alpha)} + \left\| \hat{u} \right\|_{[0,\delta]}^{(2+\alpha)} \right) \right) (s'-s)^{\frac{\alpha}{2}} \leq C_{1}(R)(s'-s)^{\frac{\alpha}{2}} \end{aligned}$$

and

$$\begin{split} |F_{\chi}(t,s',y,\theta_{\sigma}(t,s',y)) - F_{\chi}(t,0,y,\theta_{0}(t,y))| &\leq L \left|\theta_{\sigma}(t,s',y) - \theta_{0}(t,y)\right| + K(s'-0)^{\frac{\alpha}{2}} \\ &\leq L \left(\sum_{i,j=1}^{d} \left|u_{y_{i}y_{j}}(t,s',y) - g_{y_{i}y_{j}}(t,y)\right| + \sum_{i=1}^{d} \left|u_{y_{i}}(t,s',y) - g_{y_{i}}(t,y)\right| + |u(t,s',y) - g(t,y)| \\ &+ \sum_{i,j=1}^{d} \left|u_{y_{i}y_{j}}(s',s',y) - g_{y_{i}y_{j}}(0,y)\right| + \sum_{i=1}^{d} \left|u_{y_{i}}(s',s',y) - g_{y_{i}}(0,y)\right| + |u(s',s',y) - g(0,y)| \right) \\ &+ L \left(\sum_{i,j=1}^{d} \left|\widehat{u}_{y_{i}y_{j}}(t,s',y) - g_{y_{i}y_{j}}(t,y)\right| + \sum_{i=1}^{d} \left|\widehat{u}_{y_{i}}(t,s',y) - g_{y_{i}}(t,y)\right| + |\widehat{u}(t,s',y) - g(t,y)| \\ &+ \sum_{i,j=1}^{d} \left|\widehat{u}_{y_{i}y_{j}}(s',s',y) - g_{y_{i}y_{j}}(0,y)\right| + \sum_{i=1}^{d} \left|\widehat{u}_{y_{i}}(s',s',y) - g_{y_{i}}(0,y)\right| + |\widehat{u}(s',s',y) - g(0,y)| \right) \\ &+ K(s'-0)^{\frac{\alpha}{2}} \end{split}$$

$$\leq L \left(\left| \left(u - g \right) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} + \sup_{\overline{s} \in (0,s')} \left| g_t(\overline{s}, \cdot) \right|_{\mathbb{R}^d}^{(2+\alpha)} (s'-0)^1 \right. \\ \left. + \left| u(s', \cdot, \cdot) \right|_{[0,s'] \times \mathbb{R}^d}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} + \left| \left(\widehat{u} - g \right) (t, \cdot, \cdot) \right|_{[0,t] \times \mathbb{R}^d}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} \right. \\ \left. + \sup_{\overline{s} \in (0,s')} \left| g_t(\overline{s}, \cdot) \right|_{\mathbb{R}^d}^{(2+\alpha)} (s'-0)^1 + \left| \widehat{u}(s', \cdot, \cdot) \right|_{[0,s'] \times \mathbb{R}^d}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} \right) + K(s'-0)^{\frac{\alpha}{2}} \right. \\ \leq \left(K + L \left(\left\| u - g \right\|_{[0,\delta]}^{(2+\alpha)} + \left\| \widehat{u} - g \right\|_{[0,\delta]}^{(2+\alpha)} + \left\| g \right\|_{[0,\delta]}^{(2+\alpha)} \right) \right) (s'-0)^{\frac{\alpha}{2}} \leq C_2(R) \delta^{\frac{\alpha}{2}}$$

where L > 0 is a constant which could vary from line to line.

Since u and \widehat{u} satisfy the same initial condition g, we have

$$\begin{split} |\varphi(t,s,y) - \varphi(t,s',y)| \\ &\leq C_1(R)(s'-s)^{\frac{\alpha}{2}} \left(\delta^1 \left| (u-\hat{u})_s(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^d}^{(0)} + \delta^{\frac{1+\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_i}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^d}^{(1+\alpha)} \\ &+ \delta^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{y_iy_j}(t,\cdot,\cdot) \right|_{[0,s]\times\mathbb{R}^d}^{(\alpha)} + \delta^1 \left| (u-\hat{u})_s(s,\cdot,\cdot) \right|_{[0,s]\times\mathbb{R}^d}^{(0)} \\ &+ \delta^{\frac{1+\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_i}(s,\cdot,\cdot) \right|_{[0,s]\times\mathbb{R}^d}^{(1+\alpha)} + \delta^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{y_iy_j}(s,\cdot,\cdot) \right|_{[0,s]\times\mathbb{R}^d}^{(\alpha)} \\ &+ \delta^{\frac{1+\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{s(t,\cdot,\cdot)} \right|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} + (s'-s)^{\frac{1+\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_i}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^d}^{(1+\alpha)} \\ &+ (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{t(\bar{s},\cdot,\cdot)} \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^1 \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_i}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{1+\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_i}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(0)} \\ &+ (s'-s)^1 \sum_{i=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_i}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{1+\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_i}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(1+\alpha)} \\ &+ (s'-s)^1 \sum_{i,j=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_iy_j}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_iy_j}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^1 \sum_{i,j=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_iy_j}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{\alpha}{2}} \sum_{i=1}^d \left| (u-\hat{u})_{y_iy_j}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^1 \sum_{i,j=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_iy_j}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{y_iy_j}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_iy_j}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{y_iy_j}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \sup_{\bar{s}\in(s,s')} \left| (u-\hat{u})_{ty_iy_j}(\bar{s},\cdot,\cdot) \right|_{[0,\bar{s}]\times\mathbb{R}^d}^{(0)} + (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \left| (u-\hat{u})_{y_iy_j}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^d}^{(\alpha)} \\ &+ (s'-s)^{\frac{\alpha}{2}} \sum_{i,j=1}^d \sup$$

which also implies that

$$|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{\infty} \le C_3(R)\delta^{\alpha} ||u-\widehat{u}||_{[0,\delta]}^{(2+\alpha)},\tag{37}$$

since $\varphi(t,0,y) \equiv 0$.

To estimate $|\varphi(t,s,y)-\varphi(t,s,y')|$ it is convenient to add and subtract the following to $\varphi(t,s,y)-\varphi(t,s,y)$

$$\begin{split} \varphi(t,s,y') &: \\ & \int_{0}^{1} \left(F_{u}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{u}(t,0,y',\theta_{0}(t,y'))\right) \times (u(t,s,y) - \hat{u}(t,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i=1}^{d} \left(F_{p_{i}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{p_{i}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}}(t,s,y) - \hat{u}_{y_{i}}(t,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{q_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{q_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(t,s,y) - \hat{u}_{y_{i}y_{j}}(t,s,y)) d\sigma \\ & + \int_{0}^{1} \left(F_{i}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{l}(t,0,y',\theta_{0}(t,y'))\right) \times (u(s,s,y) - \hat{u}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i=1}^{d} \left(F_{m_{i}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{m_{i}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{m_{i}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{m_{i}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,0,y',\theta_{0}(t,y'))\right) \times (u_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y)) d\sigma \\ & + \int_{0}^{1} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')\right) - F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')) - F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')\right) + \int_{0}^{d} \sum_{i,j=1}^{d} \left(F_{n_{ij}}(t,s,y',\theta_{\sigma}(t,s,y')\right) + \int_{0}^{d} \sum_{i,j=1}^{d} \sum_{i,j=1}^{d} \sum_{i,j=1}^{d} \sum_{i,j=1}^{d} \sum_{i,j=1}^{d} \sum_{i,j=1}^{d} \sum_{i$$

For any first order derivatives F_{χ} of F with respect to (u, p, q, l, m, n), we have

$$\begin{split} |F_{\chi}(t,s,y,\theta_{\sigma}(t,s,y)) - F_{\chi}(t,s,y',\theta_{\sigma}(t,s,y'))| + |F_{\chi}(t,0,y,\theta_{0}(t,y)) - F_{\chi}(t,0,y',\theta_{0}(t,y'))| \\ &\leq L \left| \theta_{\sigma}(t,s,y) - \theta_{\sigma}(t,s,y') \right| + L \left| \theta_{0}(t,y) - \theta_{0}(t,y') \right| + 2K |y - y'|^{\alpha} \\ &\leq L \left(\sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(t,s,y) - u_{y_{i}y_{j}}(t,s,y') \right| + \sum_{i=1}^{d} \left| u_{y_{i}}(t,s,y) - u_{y_{i}}(t,s,y') \right| + |u(t,s,y) - u(t,s,y')| \\ &+ \sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(s,s,y) - u_{y_{i}y_{j}}(s,s,y') \right| + \sum_{i=1}^{d} \left| u_{y_{i}}(s,s,y) - u_{y_{i}}(s,s,y') \right| + |u(t,s,y) - u(t,s,y')| \\ &+ L \left(\sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(t,s,y) - \hat{u}_{y_{i}y_{j}}(t,s,y') \right| + \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(t,s,y) - \hat{u}_{y_{i}}(s,s,y) - u(s,s,y') \right| \right) \\ &+ L \left(\sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(t,s,y) - \hat{u}_{y_{i}y_{j}}(t,s,y') \right| + \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s,s,y') \right| + |\hat{u}(s,s,y) - \hat{u}(s,s,y')| \\ &+ \sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(s,s,y) - \hat{u}_{y_{i}y_{j}}(s,s,y') \right| + \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - \hat{u}_{y_{i}}(s,s,y') \right| \\ &+ L \left(\sum_{i,j=1}^{d} \left| g_{y_{i}y_{j}}(s,y) - g_{y_{i}y_{j}}(s,y') \right| + \sum_{i=1}^{d} \left| g_{y_{i}}(s,y) - g_{y_{i}}(s,s,y') \right| + |\hat{u}(s,s,y) - \hat{u}(s,s,y')| \right) \\ &+ L \left(\sum_{i,j=1}^{d} \left| g_{y_{i}y_{j}}(0,y) - g_{y_{i}y_{j}}(0,y') \right| + \sum_{i=1}^{d} \left| g_{y_{i}}(0,y) - g_{y_{i}}(0,y') \right| + |g(0,y) - g(0,y')| \right) \\ &+ \sum_{i,j=1}^{d} \left| g_{y_{i}y_{j}}(0,y) - g_{y_{i}y_{j}}(0,y') \right| + \sum_{i=1}^{d} \left| g_{y_{i}}(0,y) - g_{y_{i}}(0,y') \right| + |g(0,y) - g(0,y')| \right) \\ &+ \sum_{i,j=1}^{d} \left| g_{y_{i}y_{j}}(0,y) - g_{y_{i}y_{j}}(0,y') \right| \\ &+ \left| \hat{u}(t,\cdot) \right|_{[0,4] \times \mathbb{R}^{d}} |y - y'|^{\alpha} \\ &+ \left| \hat{u}(t,\cdot) \right|_{[0,4] \times \mathbb{R}^{d}} |y - y'|^{\alpha} + |\hat{u}(s,\cdot,\cdot) \right|_{[0,s] \times \mathbb{R}^{d}} |y - y'|^{\alpha} \\ &+ \left| g(t,\cdot) \right|_{\mathbb{R}^{d}} |y - y'|^{\alpha} + |g(0,\cdot) \right|_{\mathbb{R}^{d}} |y - y'|^{\alpha} \\ &\leq \left(2K + L \left(\left\| u \right\|_{[0,\delta]}^{(2+\alpha)} + \left\| \hat{u} \right\|_{[0,\delta]}^{(2+\alpha)} + \left\| g \right\|_{[0,\delta]}^{(2+\alpha)} \right) \right) |y - y'|^{\alpha} \\ &\leq C_{4}(R) |y - y'|^{\alpha}, \end{aligned}$$

and for every $y \in \mathbb{R}^d$,

$$\begin{split} |F_{\chi}(t,s,y,\theta_{\sigma}(t,s,y)) - F_{\chi}(t,0,y,\theta_{0}(t,y))| \\ &\leq L \left| \theta_{\sigma}(t,s,y) - \theta_{0}(t,y) \right| + K(s-0)^{\frac{\alpha}{2}} \\ &\leq L \left(\sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(t,s,y) - g_{y_{i}y_{j}}(t,y) \right| + \sum_{i=1}^{d} \left| u_{y_{i}}(t,s,y) - g_{y_{i}}(t,y) \right| \\ &+ \left| u(t,s,y) - g(t,y) \right| + \sum_{i,j=1}^{d} \left| u_{y_{i}y_{j}}(s,s,y) - g_{y_{i}y_{j}}(0,y) \right| \\ &+ \sum_{i=1}^{d} \left| u_{y_{i}}(s,s,y) - g_{y_{i}}(0,y) \right| + \left| u(s,s,y) - g(0,y) \right| \right) \\ &+ L \left(\sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(t,s,y) - g_{y_{i}y_{j}}(t,y) \right| + \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - g_{y_{i}}(t,y) \right| \\ &+ \left| \hat{u}(t,s,y) - g(t,y) \right| + \sum_{i,j=1}^{d} \left| \hat{u}_{y_{i}y_{j}}(s,s,y) - g_{y_{i}y_{j}}(0,y) \right| \\ &+ \sum_{i=1}^{d} \left| \hat{u}_{y_{i}}(s,s,y) - g_{y_{i}}(0,y) \right| + \left| \hat{u}(s,s,y) - g(0,y) \right| \right) + K(s-0)^{\frac{\alpha}{2}} \\ &\leq L \left(\left| \left(u - g \right)(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} + \sup_{\overline{s} \in (0,s)} \left| g_{t}(\overline{s},\cdot) \right|_{\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} \\ &+ \sup_{\overline{s} \in (0,s)} \left| g_{t}(\overline{s},\cdot) \right|_{\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} + \sup_{\overline{s} \in (0,s)} \left| g_{t}(\overline{s},\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} \\ &+ \sup_{\overline{s} \in (0,s)} \left| g_{t}(\overline{s},\cdot) \right|_{\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} + \left| (\widehat{u} - g) \left(t,\cdot,\cdot) \right|_{[0,s] \times \mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} \\ &\leq \left(K + L \left(\left\| u - g \right\|_{[0,\overline{s}]}^{(2+\alpha)} + \left\| \widehat{u} - g \right\|_{[0,\overline{s}]}^{(2+\alpha)} + \left\| g \right\|_{[0,\overline{s}]}^{(2+\alpha)} \right) \right) (s-0)^{\frac{\alpha}{2}} \\ &\leq C_{5}(R) \delta^{\frac{\alpha}{2}}. \end{split}$$

Hence, we have

By (36), (37) and (38), for any $t \in [0, \delta]$, we conclude that

$$|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \le C_7(R)\delta^{\frac{\alpha}{2}} \|u-\hat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$
(39)

(Estimates of $|\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}$) Next, we analyze the Hölder continuity of $\varphi_t(t,s,y)$ with respect to s and y in $[0,t]\times\mathbb{R}^d$. By the integral representation of $\varphi(t,s,y)$ in (35), we write $\varphi_t(t,s,y)$ as

$$\varphi_t(t,s,y) = \sum_{\mu} \left(\frac{\partial}{\partial t} I_{\mu}(t,s,y) \right),$$

where "*I*" represents the integrals in (35) and the subscript $\mu \in \{u, p_i, q_{ij}, l, m_i, n_{ij}; i, j = 1, \ldots, d\}$ indicate the variable, of which *F* is differentiated.

Then, we analyze $\frac{\partial}{\partial t}I_u$ in the case that $\mu = u$ as follows:

$$\begin{split} &\frac{\partial}{\partial t} I_u(t,s,y) \\ &= \int_0^1 \frac{\partial \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right)}{\partial t} \times \left(u(t,s,y) - \hat{u}(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \frac{\partial (u(t,s,y) - \hat{u}(t,s,y))}{\partial t} d\sigma \\ &= \int_0^1 \left\{ \left[F_{ut}(t,s,y,\theta_\sigma(t,s,y)) - F_{ut}(t,0,y,\theta_0(t,y))\right] \\ &+ \left[F_{uu}(t,s,y,\theta_\sigma(t,s,y)) \cdot \left(\sigma u_t(t,s,y) + (1 - \sigma)\hat{u}_t(t,s,y)\right) - F_{uu}(t,0,y,\theta_0(t,y)) \cdot g_{t}(t,y)\right] \\ &+ \sum_{i=1}^n \left[F_{up_i}(t,s,y,\theta_\sigma(t,s,y)) \cdot \left(\sigma u_{y_iy_it}(t,s,y) + (1 - \sigma)\hat{u}_{y_iy_jt}(t,s,y)\right) - F_{up_i}(t,0,y,\theta_0(t,y)) \cdot g_{y_it}(t,y)\right] \\ &+ \sum_{i,j=1}^n \left[F_{uq_{ij}}(t,s,y,\theta_\sigma(t,s,y)) \cdot \left(\sigma u_{y_iy_jt}(t,s,y) + (1 - \sigma)\hat{u}_{y_iy_jt}(t,s,y)\right) - F_{up_i}(t,0,y,\theta_0(t,y)) \cdot g_{y_it}(t,y)\right] \\ &- F_{uq_{ij}}(t,0,y,\theta_0(t,y)) \cdot g_{y_iy_jt}(t,y)\right] \right\} \times (u(t,s,y) - \hat{u}(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times (u_t(t,s,y) - \hat{u}_t(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times (u_t(t,s,y) - \hat{u}_t(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times (u_t(t,s,y) - \hat{u}_t(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times (u_t(t,s,y) - \hat{u}_t(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times (u_t(t,s,y) - \hat{u}_t(t,s,y)) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \left(u_t(t,s,y) - \hat{u}_t(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \left(u_t(t,s,y) - \hat{u}_t(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \left(u_t(t,s,y) - \hat{u}_t(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \left(u_t(t,s,y) - \hat{u}_t(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y,\theta_\sigma(t,s,y)) - F_u(t,0,y,\theta_0(t,y))\right) \times \left(u_t(t,s,y) - \hat{u}_t(t,s,y)\right) d\sigma \\ &+ \int_0^1 \left(F_u(t,s,y) + \int_0^1 \int_0^$$

where the index $\nu \in \{uu, up_i, uq_{ij}; i, j = 1, \dots, d\}$ in the summations.

We are to estimate the $\left|\frac{\partial}{\partial t}I_u(t,s,y) - \frac{\partial}{\partial t}I_u(t,s',y)\right|$ in $|\varphi_t(t,s,y) - \varphi_t(t,s',y)|$ for any $0 \leq 0$ $s \leq s' \leq t \leq \delta \leq T$ and $y \in \mathbb{R}^d$. It can be observed that the estimates of J_1 and J_2 are the same as the ones of φ . We focus on those J_{ν} terms below. In order to estimate $|J_{uu}(t,s,y) - J_{uu}(t,s',y)|$, it is convenient to add and subtract the following to $J_{uu}(t, s, y) - J_{uu}(t, s', y)$:

$$\int_0^1 F_{uu}(t,s',y,\theta_\sigma(t,s',y)) \cdot \left(\sigma u_t(t,s',y) + (1-\sigma)\widehat{u}_t(t,s',y)\right) \times (u(t,s,y) - \widehat{u}(t,s,y)) d\sigma.$$

$$37$$

We note that

$$\begin{split} \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma) \widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma u_{t}(t,s',y) + (1-\sigma) \widehat{u}_{t}(t,s',y) \right) \right| \\ &\leq \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma) \widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma) \widehat{u}_{t}(t,s,y) \right) \right| \\ &+ \left| F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma) \widehat{u}_{t}(t,s,y) \right) \right| \\ &\leq \left(K(s'-s)^{\frac{\alpha}{2}} + L \left| \theta_{\sigma}(t,s,y) - \theta_{\sigma}(t,s',y) \right| \right) \cdot \left(\left| u_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} \right| \\ &+ K \left(\left| u_{ts}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{ts}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2)} \right) (s'-s)^{1} \\ &\leq \left[K(s'-s)^{\frac{\alpha}{2}} + L \left(\left| u(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2+\alpha)} (s'-s)^{\frac{\alpha}{2}} + \sup_{\overline{s} \in (s,s')} \left| u_{t}(\overline{s},\cdot,\cdot) \right|_{[0,\overline{s}] \times \mathbb{R}^{d}}^{(2+\alpha)} (s'-s)^{1} \\ &+ \left| u(s',\cdot,\cdot) \right|_{[0,s'] \times \mathbb{R}^{d}}^{(2+\alpha)} (s'-s)^{\frac{\alpha}{2}} + \left| \widehat{u}_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(2)} (s'-s)^{1} \\ &+ \left| \widehat{u}(s',\cdot,\cdot) \right|_{[0,s'] \times \mathbb{R}^{d}}^{(2+\alpha)} (s'-s)^{\frac{\alpha}{2}} \right) \right] \cdot \left(\left| u_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} \right) \\ &+ K \left(\left| u_{ts}(t,\cdot,\cdot) \right|_{[0,s'] \times \mathbb{R}^{d}}^{(0)} (s'-s)^{\frac{\alpha}{2}} \right) \right] \cdot \left(\left| u_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} \right) \\ &+ K \left(\left| u_{ts}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} (s'-s)^{\frac{\alpha}{2}} \right) \right] \cdot \left(\left| u_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{t}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} \right) \\ &+ K \left(\left| u_{ts}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} + \left| \widehat{u}_{ts}(t,\cdot,\cdot) \right|_{[0,t] \times \mathbb{R}^{d}}^{(0)} \right) \left(s'-s \right)^{1} \\ &\leq C_{8}(R)(s'-s)^{\frac{\alpha}{2}}, \end{aligned}$$

and

$$\begin{split} \left| F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma u_{t}(t,s',y) + (1-\sigma) \widehat{u}_{t}(t,s',y) \right) \\ &- F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ \leq \left| F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma u_{t}(t,s',y) + (1-\sigma) \widehat{u}_{t}(t,s',y) \right) \\ &- F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma g_{t}(t,y) + (1-\sigma) g_{t}(t,y) \right) \right) \right| \\ &+ \left| F_{uu}(t,s',y,\theta_{\sigma}(t,s',y)) \cdot \left(\sigma g_{t}(t,y) + (1-\sigma) g_{t}(t,y) \right) - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right) \\ \leq K \left(\left| (u-g)_{ts}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(0)} + \left| (\widehat{u}-g)_{ts}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(0)} \right) (s'-0)^{1} \\ &+ \left| K(s'-0)^{\frac{\alpha}{2}} + L \left(\left| (u-g)(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} + \sup_{\overline{s}\in(0,s')} \left| g_{t}(\overline{s},\cdot) \right|_{\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{1} \\ &+ \left| u(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} + \left| (\widehat{u}-g)(t,\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} \\ &+ \sup_{\overline{s}\in(0,s')} \left| g_{t}(\overline{s},\cdot) \right|_{\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{1} + \left| \widehat{u}(s',\cdot,\cdot) \right|_{[0,s']\times\mathbb{R}^{d}}^{(2+\alpha)} (s'-0)^{\frac{\alpha}{2}} \right) \right] \cdot \left| g_{t}(t,\cdot) \right|_{\mathbb{R}^{d}}^{(0)} \\ \leq C_{9}(R) \delta^{\frac{\alpha}{2}}. \end{split}$$

Hence, we have

$$|J_{uu}(t,s,y) - J_{uu}(t,s',y)| \leq C_8(R)\delta^1(s'-s)^{\frac{\alpha}{2}} \sup_{t\in[0,\delta]} |(u-\hat{u})_s(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(0)} + C_9(R)\delta^{\frac{\alpha}{2}}(s'-s)^1 \sup_{t\in[0,\delta]} |(u-\hat{u})_s(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(0)}$$

$$\leq C_{10}(R)\delta^{\frac{\alpha}{2}}(s'-s)^{\frac{\alpha}{2}} ||u-\hat{u}||_{[0,\delta]}^{(2+\alpha)}$$
(40)

which implies that (since $J_{uu}(t,0,y) \equiv 0$)

$$|J_{uu}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{\infty} \le C_{11}(R)\delta^{\alpha} ||u-\hat{u}||_{[0,\delta]}^{(2+\alpha)}.$$
(41)

Next, to estimate $|J_{uu}(t,s,y) - J_{uu}(t,s,y')|$, it is convenient to add and subtract

$$\int_{0}^{1} \left[F_{uu}(t,s,y',\theta_{\sigma}(t,s,y')) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) - F_{uu}(t,0,y',\theta_{0}(t,y')) \cdot g_{t}(t,y') \right] \times (u_{t}(t,s,y) - \widehat{u}_{t}(t,s,y)) d\sigma$$

Then we note that

$$\begin{aligned} \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \\ &- F_{uu}(t,s,y',\theta_{\sigma}(t,s,y')) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \\ &+ F_{uu}(t,0,y',\theta_{0}(t,y')) \cdot g_{t}(t,y') \right| \\ \leq \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s,y',\theta_{\sigma}(t,s,y')) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \right| \\ &+ \left| F_{uu}(t,0,y',\theta_{0}(t,y')) \cdot g_{t}(t,y') - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ =: N_{1} + N_{2} \end{aligned}$$

For N_1 , we have

$$\begin{aligned} \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s,y,\theta_{\sigma}(t,s,y')) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \right| \\ &\leq \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \right| \\ &+ \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \right| \\ &- F_{uu}(t,s,y',\theta_{\sigma}(t,s,y')) \cdot \left(\sigma u_{t}(t,s,y') + (1-\sigma)\widehat{u}_{t}(t,s,y') \right) \right| \\ &\leq K \left(|u_{t}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(\alpha)} + |\widehat{u}_{t}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(\alpha)} |y-y'|^{\alpha} \\ &+ \left[K |y-y'|^{\alpha} + L \left(|u(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} + |u(s,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} \\ &+ |\widehat{u}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} + |\widehat{u}_{t}(s,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} \\ &+ \left| \widehat{u}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} + |\widehat{u}_{t}(s,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} \right) \right] \\ &\leq C_{v}(R) |u-u'|^{\alpha} \end{aligned}$$

 $\leq C_{12}(R)|y-y'|^{\alpha},$

For N_2 , we have

$$\begin{aligned} \left| F_{uu}(t,0,y',\theta_{0}(t,y')) \cdot g_{t}(t,y') - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ &\leq \left| F_{uu}(t,0,y',\theta_{0}(t,y')) \cdot g_{t}(t,y') - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y') \right| \\ &+ \left| F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y') - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ &\leq \left(K|y-y'|^{\alpha} + |g(t,\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} + |g(0,\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)} |y-y'|^{\alpha} \right) \cdot |g_{t}(t,\cdot)|_{\mathbb{R}^{d}}^{(0)} \\ &+ K |g_{t}(t,\cdot)|_{\mathbb{R}^{d}}^{(\alpha)} |y-y'|^{\alpha} \\ &\leq C_{13}(R)|y-y'|^{\alpha}. \end{aligned}$$

So we have

$$\left| F_{uu}(t, s, y, \theta_{\sigma}(t, s, y)) \cdot \left(\sigma u_{t}(t, s, y) + (1 - \sigma) \widehat{u}_{t}(t, s, y) \right) - F_{uu}(t, 0, y, \theta_{0}(t, y)) \cdot g_{t}(t, y) - F_{uu}(t, s, y', \theta_{\sigma}(t, s, y')) \cdot \left(\sigma u_{t}(t, s, y') + (1 - \sigma) \widehat{u}_{t}(t, s, y') \right) + F_{uu}(t, 0, y', \theta_{0}(t, y')) \cdot g_{t}(t, y') \right|$$

$$\leq N_1 + N_2 \leq C_{14}(R)|y - y'|^{\alpha}$$

Moreover, we need to estimate

$$\begin{split} \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ \leq \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot \left(\sigma u_{t}(t,s,y) + (1-\sigma)\widehat{u}_{t}(t,s,y) \right) \\ &- F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot g_{t}(t,y) + \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot g_{t}(t,y) \right| \\ + \left| F_{uu}(t,s,y,\theta_{\sigma}(t,s,y)) \cdot g_{t}(t,y) - F_{uu}(t,0,y,\theta_{0}(t,y)) \cdot g_{t}(t,y) \right| \\ \leq K \left(\left| (u-g)_{ts}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(0)} + \left| (\widehat{u}-g)_{ts}(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(0)} \right) (s-0)^{1} \\ + \left| K(s-0)^{\frac{\alpha}{2}} + L \left(\left| (u-g)(t,\cdot,\cdot) \right|_{[0,t]\times\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} + \sup_{\overline{s}\in(0,s)} |g_{t}(\overline{s},\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{1} \\ &+ |u(s,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} + |(\widehat{u}-g)(t,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} \\ &+ \sup_{\overline{s}\in(0,s)} |g_{t}(\overline{s},\cdot)|_{\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{1} + |\widehat{u}(s,\cdot,\cdot)|_{[0,s]\times\mathbb{R}^{d}}^{(2+\alpha)}(s-0)^{\frac{\alpha}{2}} \right) \right] \cdot |g_{t}(t,\cdot)|_{\mathbb{R}^{d}}^{(0)} \\ \leq C_{15}(R)\delta^{\frac{\alpha}{2}}. \end{split}$$

Consequently, we have

$$|J_{uu}(t,s,y) - J_{uu}(t,s,y')| \le C_{14}(R)\delta^{1}|y - y'|^{\alpha} \sup_{t \in [0,\delta]} |(u - \widehat{u})_{s}(t,\cdot,\cdot)|^{(0)}_{[0,t] \times \mathbb{R}^{d}} + C_{15}(R)\delta^{\frac{\alpha}{2}} \sup_{t \in [0,\delta]} |(u - \widehat{u})(t,\cdot,\cdot)|^{(\alpha)}_{[0,t] \times \mathbb{R}^{d}} |y - y'|^{\alpha}$$
(42)
$$\le C_{16}(R)\delta^{\frac{\alpha}{2}}|y - y'|^{\alpha}||u - \widehat{u}||^{(2+\alpha)}_{[0,\delta]}.$$

From (40), (41) and (42), for any $t \in [0, \delta]$, we obtain

$$|J_{uu}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \le C_{17}(R)\delta^{\frac{\alpha}{2}} ||u-\widehat{u}||_{[0,\delta]}^{(2+\alpha)}.$$

Similarly, for any $\nu \in \{uu, up_i, uq_{ij}\}$, we also obtain

$$|J_{\nu}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^{d}}^{(\alpha)} \leq C_{18}(R)\delta^{\frac{\alpha}{2}} ||u-\widehat{u}||_{[0,\delta]}^{(2+\alpha)}.$$

Hence, we have

$$\left|\frac{\partial}{\partial t}I_u(t,\cdot,\cdot)\right|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \le C_{19}(R)\delta^{\frac{\alpha}{2}} \|u-\widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$
42

By the same way, we can also estimate $|\frac{\partial}{\partial t}I_{\mu}(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}$ for $\mu \in \{u, p_i, q_{ij}, l, m_i, n_{ij}\}$:

$$\left|\frac{\partial}{\partial t}I_{\mu}(t,\cdot,\cdot)\right|_{[0,t]\times\mathbb{R}^{d}}^{(\alpha)} \leq C_{20}(R)\delta^{\frac{\alpha}{2}} \|u-\widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$

Hence, for any $t \in [0, \delta]$, we obtain

$$|\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \le C_{21}(R)\delta^{\frac{\alpha}{2}} \|u-\widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$

Together with the estimate of $|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}$ in (39), for any $t\in[0,\delta]$, we have

$$|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} + |\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)} \le C(R)\delta^{\frac{\alpha}{2}} \|u-\widehat{u}\|_{[0,\delta]}^{(2+\alpha)},$$

which indicates that

$$\sup_{t\in[0,\delta]}\left\{|\varphi(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}+|\varphi_t(t,\cdot,\cdot)|_{[0,t]\times\mathbb{R}^d}^{(\alpha)}\right\}\leq C(R)\delta^{\frac{\alpha}{2}}\|u-\widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$

Consequently, we derive a contraction mapping

$$\|\Lambda(u) - \Lambda(\widehat{u})\|_{[0,\delta]}^{(2+\alpha)} \le C(R)\delta^{\frac{\alpha}{2}} \|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)}.$$

(A contraction Λ mapping \mathcal{U} into itself.) Now, we need to choose a suitably large R such that Λ maps \mathcal{U} into itself. If δ and R satisfy

$$C(R)\delta^{\frac{\alpha}{2}} \leq \frac{1}{2}$$

then Λ is a $\frac{1}{2}$ -contraction, and for any $u \in \mathcal{U}$, we have

$$\|\Lambda(u) - g\|_{[0,\delta]}^{(2+\alpha)} \le \frac{R}{2} + \|\Lambda(g) - g\|_{[0,\delta]}^{(2+\alpha)}.$$

The function $G=\Lambda(g)-g$ is the solution of

$$\begin{cases} G_s(t,s,y) = \mathcal{L}G + F(t,s,y,g(t,y),g_y(t,y),g_{yy}(t,y),g(s,y),g_y(s,y),g_{yy}(s,y)), \\ G(t,0,y) = 0, \quad 0 \le s \le t \le \delta, \quad y \in \mathbb{R}^n. \end{cases}$$

It is clear that the term $F \in \Omega_{[0,\delta]}^{(\alpha)}$. By (26) and (27), there exists a constant c > 0 independent of δ , such that

$$\|G\|_{[0,\delta]}^{(2+\alpha)} \le c \sup_{t \in [0,\delta]} \left\{ |\psi(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(\alpha)} + |\psi_t(t,\cdot,\cdot)|_{[0,t] \times \mathbb{R}^d}^{(\alpha)} \right\} \doteq C'$$

where $\psi(t, s, y) = F(t, s, y, g(t, y), g_y(t, y), g_{yy}(t, y), g(s, y), g_y(s, y), g_{yy}(s, y)).$

Hence, we have

$$\|\Lambda(u) - g\|_{[0,\delta]}^{(2+\alpha)} \le \frac{R}{2} + C'.$$

Therefore for a suitably large R, Λ is a contraction mapping \mathcal{U} into itself, and it has a unique fixed point u in \mathcal{U} satisfying

$$\begin{array}{lll} u_s(t,s,y) &=& F\big(t,s,y,u(t,s,y),u_y(t,s,y),u_{yy}(t,s,y),\\ && u(s,s,y),u_y(s,s,y),u_{yy}(s,s,y)\big),\\ u(t,0,y) &=& g(t,y), \quad 0 \leq s \leq t \leq \delta, \quad y \in \mathbb{R}^d. \end{array}$$

(Uniqueness) To complete the proof, we ought to show that u is the unique solution of (28) in $\Omega_{[0,\delta]}^{(2+\alpha)}$. It follows with similar arguments as in the proof of Theorem 2.1 for nonlocal linear PDEs, or by studying directly the following problem satisfied by the difference of two solutions u^1 and u^2 :

$$\left\{\begin{array}{rcl} (u^1-u^2)_s\,(t,s,y)&=&F\bigl(t,s,y,u^1(t,s,y),u^1_y(t,s,y),u^1_{yy}(t,s,y),\\&&u^1(s,s,y),u^1_y(s,s,y),u^1_{yy}(s,s,y)\bigr)\\&&-F\bigl(t,s,y,u^2(t,s,y),u^2_y(t,s,y),u^2_{yy}(t,s,y),\\&&u^2(s,s,y),u^2_y(s,s,y),u^2_{yy}(s,s,y)\bigr),\\&&(u^1-u^2)\,(t,0,y)&=&0,\quad 0\leq s\leq t\leq \delta,\quad y\in \mathbb{R}^d.\end{array}\right.$$

By using the mean value theorem, the right hand side of the equation above can be expressed as a linear combination of $u^1 - u^2$ and its first- and second-order partial derivatives with respect to y. In this way, it is transformed into a nonlocal linear equation studied in the previous section. The uniqueness of solutions of nonlocal linear PDEs promises that $u^1 = u^2$ in $\Delta[0, \delta] \times \mathbb{R}^d$.

Remark 3.1 (Maximally defined solutions). We have proven the local well-posedness of (28) in $\Delta[0, \delta] \times \mathbb{R}^d$ and thus the diagonal condition can be determined for $s \in [0, \delta]$. After that, the

nonlocal fully nonlinear equation (28) reduces to a classical local fully nonlinear PDEs parameterized by t. Then we take δ as initial time and $u(t, \delta, y)$ as initial datum, we can extend the solution to a larger time interval up to the maximal interval. It is analogous to the definition of the maximally defined solution of nonlocal linear PDEs in Remark 2.1. It is noteworthy that the problem of existence at large for arbitrary initial data is a difficult task even in the local fully nonlinear case. The difficulty is caused by the fact that a priori estimate in a very high norm $|\cdot|_{[a,b]\times\mathbb{R}^d}^{(2+\alpha)}$ is needed to establish the existence at large. To this end, there should be severe restrictions on the nonlinearities. More details are discussed in [28, 29].

Remark 3.2 (Stability analysis). Suppose u and \hat{u} correspond to (F, g) and (\hat{F}, \hat{g}) , respectively. Then the following estimate holds:

$$\|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)} \le c \left(\|F - \widehat{F}\|_{[0,\delta]}^{(\alpha)} + \|g - \widehat{g}\|_{[0,\delta]}^{(2+\alpha)} \right)$$

where both F and \widehat{F} are evaluated at the point

$$(t, s, y, u(t, s, y), u_y(t, s, y), u_{yy}(t, s, y), u(s, s, y), u_y(s, s, y), s_{yy}(s, s, y)).$$

To see this, note that the difference between u and \hat{u} satisfies

$$\begin{cases} (u-\widehat{u})_{s}(t,s,y) &= \mathcal{L}(u-\widehat{u}) + F(t,s,y,u(t,s,y),u_{y}(t,s,y),u_{yy}(t,s,y), \\ & u(s,s,y),u_{y}(s,s,y),u_{yy}(s,s,y)) \\ & -\widehat{F}(t,s,y,\widehat{u}(t,s,y),\widehat{u}_{y}(t,s,y),\widehat{u}_{yy}(t,s,y), \\ & \widehat{u}(s,s,y),\widehat{u}_{y}(s,s,y),\widehat{u}_{yy}(s,s,y)), \\ & -\mathcal{L}(u-\widehat{u}) \\ (u-\widehat{u})(t,0,y) &= (g-\widehat{g})(t,y), \quad 0 \le s \le t \le \delta, \quad y \in \mathbb{R}^{d}. \end{cases}$$

Similar to the proof of Theorem 3.1, we have

$$\|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)} \le c \left(\|g - \widehat{g}\|_{[0,\delta]}^{(2+\alpha)} + \|\varphi\|_{[0,\delta]}^{(\alpha)} \right) + \frac{1}{2} \|u - \widehat{u}\|_{[0,\delta]}^{(2+\alpha)},$$

where φ represents $F - \hat{F}$ at u.

/

4. Stochastic Representation of Solutions of Nonlocal PDEs

In this section, we illustrate that the nonlocal PDEs (1) is closely connected with the theory of stochastic differential equations (SDEs). Recall that with the classical Feynman–Kac formula, a solution of a (local) linear parabolic PDE can be formulated as an expectation of a terminal stochastic process driven by a forward SDE, while a solution of a (local) semi-linear (resp. quasi-linear) PDE can be represented by adapted solutions of decoupled (resp. coupled) FBSDEs. In other words, there is a PDE-representation for adapted solutions of coupled FBSDEs and conversely, the FBSDEs can also provide a probabilistic interpretation and a numerical scheme for parabolic PDEs.

Parallel to the elegant connection between local parabolic PDEs and FBSDEs, there is also a key connection between nonlocal PDEs and a flow of FBSDEs (or BSVIEs). By exploring this connection, we can construct computational methods to seek for solutions of nonlocal PDEs numerically.

4.1. Existing Feynman–Kac type results

We first briefly review the existing results that connect a flow of FBSDEs with nonlocal PDEs in the literature. Given a k-dimensional Brownian motion $W(\cdot)$ defined in the completed filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, let us consider a coupled FBSDE of the form:

$$X(s) = y + \int_0^s b(\tau, X(\tau), Y(\tau, \tau)) d\tau + \int_0^s \sigma(\tau, X(\tau), Y(\tau, \tau)) dW(\tau),$$
(43)

$$Y(t,s) = g(t,X(T)) + \int_{s}^{T} h(t,\tau,X(\tau),Y(t,\tau),Y(\tau,\tau),Z(t,\tau))d\tau$$
(44)

$$-\int_{s}^{T} Z^{\top}(t,\tau) dW(\tau), \quad 0 \le t \le s \le T, \quad y \in \mathbb{R}^{d},$$

Getting rid of the diagonal term $Y(\tau, \tau)$ in (43) and (44), the FBSDEs above reduce to classical ones parameterized by $t \in [0, T]$, which essentially form a flow of FBSDEs. However, the dependence on t and $Y(\tau, \tau)$ violates the flow property of FBSDEs as well as results in nonlocality of its Feynman–Kac-type representations. Heuristically speaking, a family of $\mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^{k \times 1}$ -valued random fields

$$(X(\cdot), Y(\cdot, \cdot), Z(\cdot, \cdot)) = \left\{ (X(s), Y(t, s), Z(t, s)) : 0 \le t \le s \le T \right\}$$
46

is called an adapted solution of the FBSDE system (43)-(44) if

- 1. $X(\cdot)$ is \mathbb{F} -adapted and continuous;
- 2. for each fixed $t \in [0, T]$, $Y(t, \cdot)$ is \mathbb{F} -adapted and continuous;
- 3. for each fixed $t \in [0, T]$, $Z(t, \cdot)$ is \mathbb{F} -progressively measurable;
- 4. the system holds in the usual Itô's sense for almost every $t \in [0, T]$.

Some well-posedness results and applications of the system have been studied in the literature. Such a class of BSDEs (44), the generator h and the terminal term g of which depend on t and/or diagonal terms $Y(\tau, \tau)$, is also usually called as backward stochastic Volterra integral equation (BSVIE). The concept of BSVIEs was initially introduced in [30] as an extension of classical BSDEs developed in [31, 32]. Then it was developed in several other studies, including [33, 34, 35, 36, 37, 38, 39]. Among them, [34] extends BSVIEs to incorporate with general terminal function motivated by the optimal control of forward-SVIEs (FSVIEs). The regularity of BSVIEs in Hilbert spaces is studied in [35]. A timely survey on BSVIEs is documented in [40]. Recently, [41, 42] develop a theory of path-dependent BSVIEs in a non-Markovian setting. As for applications, BSVIEs generalize the theory of stochastic differential utility to incorporate with state dependence and solve the corresponding TIC stochastic recursive control problems; see [11, 16, 43]. Moreover, [44, 45] study the applications of BSVIE to dynamic risk measures.

Next, we discuss about several special situations of the coupled system (43)-(44). Under some conditions, they will lead to some Feynman–Kac-type results in the existing literature.

 When b, σ and h are all independent of the diagonal term Y(τ, τ), then the system of (43)-(44) is reduced to a family of classical FBSDEs parameterized by t ∈ [0, T], whose solution is linked with that of the following parameterized semilinear PDEs:

$$\begin{cases} u_s(t,s,y) + \frac{1}{2} \text{tr} \left\{ \sigma(s,y) \sigma(s,y)^\top u_{yy}(t,s,y) \right\} + b^\top(s,y) u_y(t,s,y) \\ + h(t,s,y,u(t,s,y), \sigma^\top(s,y) u_y(t,s,y)) = 0, \\ u(t,T,y) = g(t,y), \quad 0 \le t \le s \le T, \quad y \in \mathbb{R}^d. \end{cases}$$

via the relations Y(t, s) = u(t, s, X(s)) and $Z(t, s) = \sigma^{\top}(s, X(s))u_y(t, s, X(s))$. In particular, when the parameter t is taken over a singleton, the result is proved in [46, 47];

2. When b and σ do not depend on $Y(\tau, \tau)$ and h does not include $Y(t, \tau)$, by letting s = t in (44), [20] links the system with a nonlocal PDE:

$$\begin{cases} u_s(t,s,y) + \frac{1}{2} \operatorname{tr} \left\{ \sigma(s,y) \sigma(s,y)^\top u_{yy}(t,s,y) \right\} + b^\top(s,y) u_y(t,s,y) \\ + h(t,s,y,u(s,s,y),\sigma^\top(s,y) u_y(t,s,y)) = 0, \\ u(t,T,y) = g(t,y), \quad 0 \le t \le s \le T, \quad y \in \mathbb{R}^d. \end{cases}$$

via the relations Y(t,t) = u(t,t,X(t)) and $Z(t,s) = \sigma^{\top}(s,X(s))u_y(t,s,X(s));$

3. When b and σ are independent of $Y(\tau, \tau)$, [18] shows that the system corresponds to the following nonlocal PDE:

$$\begin{cases} u_s(t,s,y) + \frac{1}{2} \text{tr} \left\{ \sigma(s,y) \sigma(s,y)^\top u_{yy}(t,s,y) \right\} + b^\top(s,y) u_y(t,s,y) \\ + h(t,s,y,u(t,s,y),u(s,s,y),\sigma^\top(s,y) u_y(t,s,y)) = 0, \\ u(t,T,y) = g(t,y), \quad 0 \le t \le s \le T, \quad y \in \mathbb{R}^d. \end{cases}$$

indicated by Y(t,s) = u(t,s,X(s)) and $Z(t,s) = \sigma^{\top}(s,X(s))u_y(t,s,X(s)).$

4. Inspired by the previous works, [17] concludes that the investigation of a (coupled) flow of forward-backward SDEs (43)-(44) will be reduced to the study of the nonlocal PDE of the following form:

$$\begin{cases} u_s(t,s,y) + \frac{1}{2} \text{tr} \left\{ \sigma \left(s, y, u(s,s,y) \right) \sigma \left(s, y, u(s,s,y) \right)^\top u_{yy}(t,s,y) \right\} \\ + b^\top \left(s, y, u(s,s,y) \right) u_y(t,s,y) \\ + h \left(t, s, y, u(t,s,y), u(s,s,y), \sigma^\top (s, y, u(s,s,y)) u_y(t,s,y) \right) = 0, \\ u(t,T,y) = g(t,y), \qquad 0 \le t \le s \le T, \quad y \in \mathbb{R}^d. \end{cases}$$

One can observe that the examples (1)-(4) above contain neither a nonlinearity of the highestorder term u(t, s, y) nor a dependence of diagonal terms $u_y(s, s, y)$ and $u_{yy}(s, s, y)$. It is clear that our study about nonlocal PDEs in Section 2 and Section 3 provides a general and unified treatment of this class of Feynman–Kac formulas and presents the necessity and significance of FBSDEs in a more general form than (43)-(44).

4.2. Flow of second-order FBSDEs

In fact, there has been some work connecting classical local fully nonlinear PDEs with FBS-DEs. The pioneering work of [48] introduces the concept of second-order BSDEs (2BSDEs) that extends the Feynman–Kac formula to a fully-nonlinear setting. The key ingredient of the method is to introduce a new process that identifies the Hessian matrix of the solution of the corresponding PDE. [49] provides an alternative formulation of 2BSDEs under a non-dominated family of mutually singular probability measures. By parameterizing the 2FBSDEs in [48] or [49] with t and introducing some diagonal terms to them, the parallel concept of 2FBSVIEs or a flow of 2FBSDEs is highly expected to provide a probabilistic interpretation and numerical scheme for solutions of nonlocal fully nonlinear PDEs (1).

To see this, we further impose regularity of the initial condition and thus we can obtain regularity of solutions of (1) as in the following lemma.

Lemma 4.1. For any k = 1, ..., d, suppose that F_{y_k} , $F_{\mathcal{X}}$ in Table 1, and $F_{\mathcal{X}\mathcal{Y}}$ in Table 2 satisfy conditions (29)-(32). If $g \in \Omega_{[0,T]}^{(3+\alpha)}$, then the unique solution $u \in \Omega_{[0,\delta]}^{(2+\alpha)}$ of (28) satisfies that $u_{y_k} \in \Omega_{[0,\delta]}^{(2+\alpha)}$.

Proof. As Theorem 3.1 presents, the nonlocal fully nonlinear equation (28) admits a unique solution u in $\Omega_{[0,\delta]}^{(2+\alpha)}$. Next, for any $k = 1, 2, \dots, d$, we can find u_{y_k} satisfying:

$$\begin{pmatrix}
\left(\frac{\partial u}{\partial y_k}\right)_s(t,s,y) = \sum_{i,j=1}^d F_{q_{ij}}(\cdot) \left(\frac{\partial u}{\partial y_k}\right)_{y_i y_j}(t,s,y) + \sum_{i=1}^d F_{p_i}(\cdot) \left(\frac{\partial u}{\partial y_k}\right)_{y_i}(t,s,y) \\
+ \sum_{i,j=1}^d F_{n_{ij}}(\cdot) \left(\frac{\partial u}{\partial y_k}\right)_{y_i y_j}(s,s,y) + \sum_{i=1}^d F_{m_i}(\cdot) \left(\frac{\partial u}{\partial y_k}\right)_{y_i}(s,s,y) \\
+ F_u(\cdot) \left(\frac{\partial u}{\partial y_k}\right)(t,s,y) + F_l(\cdot) \left(\frac{\partial u}{\partial y_k}\right)(s,s,y) + F_{y_k}(\cdot)$$

$$\left(\frac{\partial u}{\partial y_k}\right)(t,0,y) = g_{y_k}(t,y), \quad 0 \le s \le t \le T, \quad y \in \mathbb{R}^d.$$
(45)

where all F_{y_k} and $F_{\mathcal{X}}(\cdot)$ are evaluated at

$$(t, s, y, u(t, s, y), u_y(t, s, y), u_{yy}(t, s, y), u(s, s, y), u_y(s, s, y), u_{yy}(s, s, y)).$$

Given $u \in \Omega_{[0,\delta]}^{(2+\alpha)}$, all coefficient functions $F_{\mathcal{X}}(\cdot)$ in (45) and the nonhomogeneous term $F_{y_k}(\cdot)$ belong to $\Omega_{[0,\delta]}^{(\alpha)}$. Moreover, since the initial condition $g_{y_k}(t,y) \in \Omega_{[0,\delta]}^{(2+\alpha)}$ by $g \in \Omega_{[0,T]}^{(3+\alpha)}$, applying Corollary 2.1 to (45) completes the proof. Next, we consider a backward nonlocal fully nonlinear PDE of the form

$$\begin{cases} u_{s}(t,s,y) = F(t,s,y,u(t,s,y),u_{y}(t,s,y),u_{yy}(t,s,y), \\ u(s,s,y),u_{y}(s,s,y),u_{yy}(s,s,y)), \\ u(t,T,y) = g(t,y), \quad t_{0} \le t \le s \le T, \quad y \in \mathbb{R}^{d}. \end{cases}$$
(46)

where F has enough regularities of (29)-(32) and t_0 is suitable in the sense that $[t_0, T]$ is a subset of the time interval for the maximally defined solution of (46). Then the following theorem reveals the relationship between nonlocal fully nonlinear PDEs like (46) and flow of 2FBSDEs (48).

Theorem 4.1. For any k = 1, ..., d, suppose that F_{y_k} , $F_{\mathcal{X}}$ in Table 1, and $F_{\mathcal{X}\mathcal{Y}}$ in Table 2 satisfy conditions (29)-(32), $\sigma(s, y) \in C^{1,2}([t_0, T] \times \mathbb{R}^d)$, and $g \in \Omega^{(3+\alpha)}_{[t_0,T]}$. Then, (46) admits a unique solution u(t, s, y) that is first-order continuously differentiable in s and third-order continuously differentiable with respect to y in $\nabla[t_0, T] \times \mathbb{R}^d$. Moreover, let

$$Y(t,s) := u(t,s,X(s)), \qquad Z(t,s) := \left(\sigma^{\top} u_y\right)(t,s,X(s)), \qquad (47)$$
$$\Gamma(t,s) := \left(\sigma^{\top} \left(\sigma^{\top} u_y\right)_y\right)(t,s,X(s)), \qquad A(t,s) := \mathcal{D}\left(\sigma^{\top} u_y\right)(t,s,X(s)),$$

where $(\sigma^{\top}u)(t,s,y) = \sigma^{\top}(s,y)u(t,s,y)$ and the operator \mathcal{D} is defined by

$$\mathcal{D}\varphi = \varphi_s + \frac{1}{2} \sum_{i,j=1}^d \left(\sigma \sigma^{\top}\right)_{ij} \frac{\partial^2 \varphi}{\partial y_i \partial y_j} + \sum_{i=1}^d b_i \frac{\partial \varphi}{\partial y_i}$$

then the family of random fields $(X(\cdot), Y(\cdot, \cdot), Z(\cdot, \cdot), \Gamma(\cdot, \cdot), A(\cdot, \cdot))$ is an adapted solution of the following flow of 2FBSDEs:

$$X(s) = y + \int_{t_0}^{s} b(\tau, X(\tau)) d\tau + \int_{t_0}^{s} \sigma(\tau, X(\tau)) dW(\tau),$$

$$Y(t, s) = g(t, X(T)) - \int_{s}^{T} \mathbb{F}(t, \tau, X(\tau), Y(t, \tau), Y(\tau, \tau), Z(t, \tau), Z(\tau, \tau), \Gamma(t, \tau), \Gamma(\tau, \tau)) d\tau$$

$$- \int_{s}^{T} Z^{\top}(t, \tau) dW(\tau),$$

$$Z(t, s) = Z(t, t_0) + \int_{t_0}^{s} A(t, \tau) d\tau + \int_{t_0}^{s} \Gamma(t, \tau) dW(\tau), \quad t_0 \le t \le s \le T, \quad y \in \mathbb{R}^d$$
(48)

where \mathbb{F} is defined by

$$\mathbb{F}(t,\tau,X(\tau),Y(t,\tau),Y(\tau,\tau),Z(t,\tau),Z(\tau,\tau),\Gamma(t,\tau),\Gamma(\tau,\tau))$$

$$=\overline{F}(t,\tau,X(\tau),u(t,\tau,X(\tau)),u_{y}(t,\tau,X(\tau)),u_{yy}(t,\tau,X(\tau)),$$

$$u(\tau,\tau,X(\tau)),u_{y}(\tau,\tau,X(\tau)),u_{yy}(\tau,\tau,X(\tau))),$$
(49)

with the definition of \overline{F}

$$\overline{F}(t,\tau,y,u(t,\tau,y),u_y(t,\tau,y),u_{yy}(t,\tau,y),u(\tau,\tau,y),u_y(\tau,\tau,y),u_{yy}(\tau,\tau,y))$$

$$=F(t,\tau,y,u(t,\tau,y),u_y(t,\tau,y),u_{yy}(t,\tau,y),u(\tau,\tau,y),u_y(\tau,\tau,y),u_{yy}(\tau,\tau,y))$$

$$+\frac{1}{2}\sum_{i,j=1}^d \left(\sigma\sigma^{\top}\right)_{ij}(\tau,y)\frac{\partial^2 u}{\partial y_i \partial y_j}(t,\tau,y) + \sum_{i=1}^d b_i(\tau,y)\frac{\partial u}{\partial y_i}(t,\tau,y).$$

Proof. First, under the regularity assumptions of F and g, Corollary 2.1 and Lemma 4.1 guarantee that there exists a unique solution u(t, s, y) of (46), which is first-order continuously differentiable in s and third-order continuously differentiable with respect to y. Consequently, the family of random fields (X, Y, Z, Γ, A) defined by (47) is well-defined (adapted).

Next, we show that the random field solves the flow of 2FBSDEs, i.e. (48). For any fixed $(t,s) \in \nabla[t_0,T]$, we apply the Itô's formula to the map $\tau \to u(t,\tau,X(\tau))$ on [s,T]. Then we have

$$\begin{split} du(t,\tau,X(\tau)) \\ = & \Big[u_s(t,\tau,X(\tau)) + \sum_{i=1}^d b_i(\tau,X(\tau)) \frac{\partial u}{\partial y_i}(t,\tau,X(\tau)) \\ & + \frac{1}{2} \sum_{i,j=1}^d \left(\sigma \sigma^\top \right)_{ij}(\tau,X(\tau)) \frac{\partial^2 u}{\partial y_i \partial y_j}(t,\tau,X(\tau)) \Big] d\tau + u_y^\top(t,\tau,X(\tau)) \sigma(\tau,X(\tau)) dW(\tau) \\ = & \Big[F(t,\tau,X(\tau),u(t,\tau,X(\tau)),u_y(t,\tau,X(\tau)),u_y(t,\tau,X(\tau)), \\ & u(\tau,\tau,X(\tau)),u_y(\tau,\tau,X(\tau)),u_y(\tau,\tau,X(\tau)) \\ & + \frac{1}{2} \sum_{i,j=1}^d \left(\sigma \sigma^\top \right)_{ij}(\tau,X(\tau)) \frac{\partial^2 u}{\partial y_i \partial y_j}(t,\tau,X(\tau)) + \sum_{i=1}^d b_i(\tau,X(\tau)) \frac{\partial u}{\partial y_i}(t,\tau,X(\tau)) \Big] d\tau \\ & + u_y^\top(t,\tau,X(\tau)) \sigma(\tau,X(\tau)) dW(\tau) \\ = & \mathbb{E}(t,\tau,X(\tau),Y(t,\tau),Y(\tau,\tau),Z(t,\tau),Z(\tau,\tau),\Gamma(t,\tau),\Gamma(\tau,\tau)) d\tau \\ & + Z^\top(t,\tau) dW(\tau) \end{split}$$

which indicates $dY(t,\tau) = \mathbb{F}ds + Z^{\top}(t,\tau)dW(\tau)$. Similarly, for any fixed $(t,s) \in \nabla[t_0,T]$, by applying the Itô's formula to $\tau \to (\sigma^{\top}u_y)(t,\tau,X(\tau))$ on $[t_0,s]$, we can also verify that $dZ(t,\tau) = A(t,\tau)d\tau + \Gamma(t,\tau)dW(\tau)$. Hence, (47) is an adapted solution of (48).

We make three important observations about the stochastic system (48): (I) When the generator \mathbb{F} is independent of diagonal terms, i.e. $Y(\tau, \tau)$, $Z(\tau, \tau)$, and $\Gamma(\tau, \tau)$, the flow of FBSDEs (48) are reduced to a family of 2FBSDEs parameterized by t, which is exactly the 2FBSDE in [50] and equivalent to the ones in [48] for any fixed t; (II) (48) is more general than the system of (43)-(44) since it allows for a nonlinearity of $(Y(t, \tau), Z(t, \tau), \Gamma(t, \tau))$ by introducing an additional SDE and also contains their diagonal terms $(Y(\tau, \tau), Z(\tau, \tau), \Gamma(\tau, \tau))$ in almost arbitrary way; (III) Theorem 4.1 shows how to solve the flow of 2FBSDE (48) from the perspective of nonlocal PDEs. Inspired by [48] and [49], the opposite implication of solutions (from 2FBSDEs to PDE) is likely valid by establishing the well-posedness of (48) in the theoretical framework of SDEs. However, it is beyond the scope of this paper, while we will prove the existence and uniqueness of (48) in our future publications.

5. Conclusions

This paper studies nonlocal parabolic PDEs originated from time-inconsistent problems in the theory of stochastic differential equations and control problems. Tailor-made norms and Banach spaces are proposed for the study of nonlocality and time-inconsistency. Based on which, the first key step is to establish the well-posedness of nonlocal linear PDEs. Subsequently, a Schauder-type prior estimate is provided to control the behaviour of solutions of nonlocal linear PDEs. The study of the linear case laid a solid foundation for studying nonlocal fully nonlinear PDEs by the linearization method. By constructing a contraction in the space of possible solutions, the Banach's fixed point theorem promises the existence and uniqueness of nonlocal fully nonlinear equations.

Thanks to the close connections among PDE, FBSDE, and stochastic controls, our wellposedness results of nonlocal PDEs contribute significantly to the development of the related fields: (1) From the perspective of TIC stochastic control problems, the main breakthrough is that the nonlocal fully nonlinear PDEs in this paper allow control enter the diffusion of state process, which extends essentially the restricted results in the existing literature. The solvability of nonlocal fully nonlinear PDEs indicates directly the well-posedness of the related HJB-type equations in TIC problems, including equilibrium HJB equations and HJB systems, both of which are used to identify the subgame perfect equilibrium; (2) In terms of the theory of FBSDEs, our study of nonlocal PDEs provides a general and unified treatment for the Feynman–Kac formulas of a flow of 2FBSDEs (48). The advance in nonlocal PDEs of this paper provides a new insight into the study of the flow of FBSDEs or 2FBSDEs from the perspective of PDEs. Besides of pushing the frontiers of PDE, SDE, and stochastic control theory, our work sheds light on some finance and decision-making theories that involve reference points (initial-time dependence), such as prospect and regret theories.

References

- J. Yong, X. Y. Zhou, Stochastic Controls : Hamiltonian Systems and HJB Equations, 1st Edition, Vol. 43 of Stochastic Modelling and Applied Probability, Springer New York, New York, NY, 1999. doi:10.1007/ 978-1-4612-1466-3.
- [2] H. Markowitz, Portfolio selection, The Journal of Finance 7 (1) (1952) 77-91. doi:10.1111/j.
 1540-6261.1952.tb01525.x.
- [3] S. Frederick, G. Loewenstein, T. O'donoghue, Time discounting and time preference: A critical review, Journal of Economic Literature 40 (2) (2002) 351–401. doi:10.1257/jel.40.2.351.
- [4] D. Kahneman, A. Tversky, Prospect theory: An analysis of decision under risk, Econometrica 47 (2) (1979) 263.
 doi:10.2307/1914185.
- [5] R. H. Thaler, Some empirical evidence on dynamic inconsistency, Economics Letters 8 (3) (1981) 201–207.
 doi:10.1016/0165-1765 (81) 90067-7.
- [6] D. Laibson, Golden eggs and hyperbolic discounting, The Quarterly Journal of Economics 112 (2) (1997) 443–478. doi:10.1162/003355397555253.
- [7] R. H. Strotz, Myopia and inconsistency in dynamic utility maximization, The Review of Economic Studies 23 (3) (1955) 165–180. doi:10.2307/2295722.
- [8] R. A. Pollak, Consistent planning, The Review of Economic Studies 35 (2) (1968) 201. doi:10.2307/ 2296548.
- [9] T. Björk, M. Khapko, A. Murgoci, On time-inconsistent stochastic control in continuous time, Finance and Stochastics 21 (2) (2017) 331–360. doi:10.1007/s00780-017-0327-5.

- [10] J. Yong, Time-inconsistent optimal control problems and the equilibrium HJB equation, Mathematical Control & Related Fields 2 (3) (2012) 271-329. doi:10.3934/mcrf.2012.2.271.
- [11] Q. Wei, J. Yong, Z. Yu, Time-inconsistent recursive stochastic optimal control problems, SIAM Journal on Control and Optimization 55 (6) (2017) 4156–4201. doi:10.1137/16m1079415.
- W. Yan, J. Yong, Time-inconsistent optimal control problems and related issues, in: G. Yin, Q. Zhang (Eds.), Modeling, Stochastic Control, Optimization, and Applications, Vol. 164, Springer International Publishing, 2019, pp. 533–569. doi:10.1007/978-3-030-25498-8_22.
- [13] M. Kac, On distributions of certain Wiener functionals, Transactions of the American Mathematical Society 65 (1) (1949) 1–13. doi:10.1090/s0002-9947-1949-0027960-x.
- [14] C. Hernández, D. Possamaï, Me, myself and i: A general theory of non-Markovian time-inconsistent stochastic control for sophisticated agents, arXiv: 2002.12572 (Feb. 2020). arXiv:2002.12572.
- [15] H. Mei, C. Zhu, Closed-loop equilibrium for time-inconsistent McKean–Vlasov controlled problem, SIAM Journal on Control and Optimization 58 (6) (2020) 3842–3867. doi:10.1137/20ml319796.
- [16] H. Wang, J. Yong, Time-inconsistent stochastic optimal control problems and backward stochastic Volterra integral equations, ESAIM: Control, Optimisation and Calculus of Variations 27 (2021) 22. doi:10.1051/ cocv/2021027.
- [17] Y. Hamaguchi, Small-time solvability of a flow of forward-backward stochastic differential equations, Applied Mathematics & Optimization 84 (1) (2021) 567–588.
- [18] H. Wang, Extended backward stochastic Volterra integral equations, quasilinear parabolic equations, and Feynman-Kac formula, Stochastics and Dynamics 21 (01) (2020) 2150004. doi:10.1142/ s0219493721500040.
- [19] J. Ma, P. Protter, J. Yong, Solving forward-backward stochastic differential equations explicitly a four step scheme, Probability Theory and Related Fields 98 (3) (1994) 339–359. doi:10.1007/bf01192258.
- [20] T. Wang, J. Yong, Backward stochastic Volterra integral equations—representation of adapted solutions, Stochastic Processes and their Applications 129 (12) (2019) 4926–4964. doi:10.1016/j.spa.2018. 12.016.
- [21] A. Friedman, Partial Differential Equations of Parabolic Type, 1st Edition, Prentice-Hall, Englewood Cliffs, N.J., 1964.
- [22] O. A. Ladyženskaja, V. A. Solonnikov, N. N. Ural'ceva, Linear and Quasi-linear Equations of Parabolic Type, 1st Edition, American Mathematical Society, 1968.
- [23] A. Lunardi, Analytic Semigroups and Optimal Regularity in Parabolic Problems, 1st Edition, Springer Basel, 1995.
- [24] S. D. Éidel'man, Parabolic Systems, North-Holland and Pub. Co.Wolters-Noordhoff, Amsterdam, Groningen, 1969.

- [25] S. Khudyaev, The first boundary-value problem for non-linear parabolic equations, in: Doklady Akademii Nauk, Vol. 149, Russian Academy of Sciences, 1963, pp. 535–538.
- [26] N. N. Šopolov, The first boundary value problem for nonlinear parabolic equations of arbitrary order, C. R. Acad. Bulgare Sci. 23 (1970) 899–902.
- [27] S. N. Kruzhkov, A. Castro, M. L. Morales, Schauder-type estimates and existence theorems for the solution of basic problems for linear and nonlinear parabolic equations, in: Doklady Akademii Nauk, Vol. 220, Russian Academy of Sciences, 1975, pp. 277–280.
- [28] N. V. Krylov, Nonlinear Elliptic and Parabolic Equations of the Second Order, 1st Edition, Vol. 7 of Mathematics and its Applications, Springer Netherlands, 1987.
- [29] G. M. Lieberman, Second Order Parabolic Differential Equations, 1st Edition, World Scientific, 1996. doi: 10.1142/3302.
- [30] J. Lin, Adapted solution of a backward stochastic nonlinear Volterra integral equation, Stochastic Analysis and Applications 20 (1) (2002) 165–183. doi:10.1081/sap-120002426.
- [31] J.-M. Bismut, Théorie probabiliste du contrôle des diffusions, Memoirs of the American Mathematical Society 4 (167) (1976). doi:10.1090/memo/0167.
- [32] Étienne Pardoux, S. Peng, Adapted solution of a backward stochastic differential equation, Systems & Control Letters 14 (1) (1990) 55–61. doi:10.1016/0167-6911(90)90082-6.
- [33] A. Aman, M. Nzi, Backward stochastic nonlinear Volterra integral equation with local lipschitz drift, Probability and Mathematical Statistics 25 (1) (2005) 105–127.
- [34] J. Yong, Backward stochastic Volterra integral equations and some related problems, Stochastic Processes and their Applications 116 (5) (2006) 779–795. doi:10.1016/j.spa.2006.01.005.
- [35] V. V. Anh, W. Grecksch, J. Yong, Regularity of backward stochastic Volterra integral equations in Hilbert spaces, Stochastic Analysis and Applications 29 (1) (2010) 146–168. doi:10.1080/07362994.2011.532046.
- [36] Y. Ren, On solutions of backward stochastic Volterra integral equations with jumps in Hilbert spaces, Journal of Optimization Theory and Applications 144 (2) (2010) 319–333. doi:10.1007/s10957-009-9596-2.
- [37] J. Djordjević, S. Janković, On a class of backward stochastic Volterra integral equations, Applied Mathematics Letters 26 (12) (2013) 1192–1197. doi:10.1016/j.aml.2013.07.006.
- [38] J. Djordjević, S. Janković, Backward stochastic Volterra integral equations with additive perturbations, Applied Mathematics and Computation 265 (C) (2015) 903–910. doi:10.1016/j.amc.2015.05.077.
- [39] Y. Hu, B. Øksendal, Linear Volterra backward stochastic integral equations, Stochastic Processes and their Applications 129 (2) (2019) 626–633. doi:10.1016/j.spa.2018.03.016.
- [40] J. Yong, Backward stochastic Volterra integral equations a brief survey, Applied Mathematics-A Journal of Chinese Universities 28 (4) (2013) 383–394. doi:10.1007/s11766-013-3189-4.
- [41] L. Overbeck, J. A. L. Röder, Path-dependent backward stochastic Volterra integral equations with jumps, dif-

ferentiability and duality principle, Probability, Uncertainty and Quantitative Risk 3 (1) (Jun. 2018). doi: 10.1186/s41546-018-0030-2.

- [42] H. Wang, J. Yong, J. Zhang, Path dependent Feynman-Kac formula for forward backward stochastic Volterra integral equations, arXiv: 2004.05825 (Apr. 2020). arXiv:2004.05825.
- [43] Y. Hamaguchi, Extended backward stochastic Volterra integral equations and their applications to timeinconsistent stochastic recursive control problems, Mathematical Control & Related Fields 11 (2) (2021) 197– 242. doi:10.3934/mcrf.2020043.
- [44] J. Yong, Continuous-time dynamic risk measures by backward stochastic Volterra integral equations, Applicable Analysis 86 (11) (2007) 1429–1442. doi:10.1080/00036810701697328.
- [45] H. Wang, J. Sun, J. Yong, Recursive utility processes, dynamic risk measures and quadratic backward stochastic Volterra integral equations, Applied Mathematics & Optimization 84 (1) (2021) 145–190. doi:10.1007/ s00245-019-09641-7.
- [46] S. Peng, A nonlinear Feynman–Kac formula and applications, in: Proceedings of Symposium of System Sciences and Control Theory, World Scientific, 1992, pp. 173–184.
- [47] J. Ma, J. Zhang, Representation theorems for backward stochastic differential equations, The Annals of Applied Probability 12 (4) (2002) 1390–1418. doi:10.1214/aoap/1037125868.
- [48] P. Cheridito, H. M. Soner, N. Touzi, N. Victoir, Second-order backward stochastic differential equations and fully nonlinear parabolic PDEs, Communications on Pure and Applied Mathematics 60 (7) (2007) 1081–1110. doi:10.1002/cpa.20168.
- [49] H. M. Soner, N. Touzi, J. Zhang, Wellposedness of second order backward SDEs, Probability Theory and Related Fields 153 (1-2) (2011) 149–190. doi:10.1007/s00440-011-0342-y.
- [50] T. Kong, W. Zhao, T. Zhou, Probabilistic high order numerical schemes for fully nonlinear parabolic PDEs, Communications in Computational Physics 18 (5) (2015) 1482–1503. doi:10.4208/cicp.240515. 280815a.