

Cachazo-Svrček-Witten Rules for Tree-Level Gluonic Amplitudes Revisited

Wen-Jie Zhang¹, Jun-Bao Wu^{1,2*,†}, Chuan-Jie Zhu³

March 26, 2024

¹*Center for Joint Quantum Studies and Department of Physics, School of Science, Tianjin University, 135 Yaguan Road, Tianjin 300350, P. R. China*

²*Peng Huangwu Center for Fundamental Theory, Hefei, Anhui 230026, P. R. China*

³*College of Mathematics and Physics Science, Hunan University of Arts and Science, Changde 415000, P. R. China*

E-mail: zhangwenjie19@tju.edu.cn, junbao.wu@tju.edu.cn, zhucj@huas.edu.cn

Abstract

We provide a new proof of Cachazo-Svrček-Witten rules for tree-level gluonic amplitudes. As a key step, we explicitly show the cancellation of spurious poles originating from the maximally helicity violating vertices in these rules. To achieve this, we introduce specially-defined two-off-shell-line sub-amplitudes and study their residues at spurious poles.

1 Introduction

Great achievement [1, 2, 3] on scattering amplitudes in gauge theories in recent around two decades was largely inspired by the seminal paper of Witten on relationship between amplitudes and twistor string theory [4]. Tree level MHV (maximally helicity violating) amplitudes were obtained there from integration over moduli space of genus-zero degree-one curves¹ in super twistor space $\mathbb{P}^{3|4}$. The tree non-MHV amplitudes are corresponding to genus-zero curves with degree larger than one. More precisely, N^k MHV amplitudes (amplitudes with $k + 2$ negative-helicity gluons) are related to curves

*Corresponding author.

[†]The unusual ordering of authors instead of the standard alphabetical one in hep-th community is for students to get proper recognition of contribution under the current out-dated practice in China.

¹These curves are D -instantons in B model with target space $\mathbb{P}^{3|4}$.

with degree $d = k + 1$. Later it was found that for non-MHV tree amplitudes, one can either integrate over moduli space of connected curve with degree d or over moduli space of d degree-one curves (completely disconnected instantons). The former choice led to compact Roiban-Spradlin-Volovich (RSV) formula [5, 6, 7] for tree amplitudes, while the later led to the Cachazo-Svrček-Witten (CSW) rules for tree amplitudes [8]. The equivalence of these two choices was proved in [9]. Intermediate prescription which is to integrate over moduli space of m ($1 < m < d$) curves with total degree d was also proposed in [9]. A twistor space recursion relation for these intermediate prescription was given in [10].

CSW introduced off-shell continuation of MHV amplitudes to make them MHV vertices. Together with scalar propagators, these MHV vertices build up MHV diagrams. The calculation of tree amplitudes using MHV diagrams is more efficient than using Feynman diagrams. For reviews including CSW rules, see [11, 12, 13]. The analysis of various physical and spurious singularities in [8], combined with the dimension analysis (or on-shell recursion relation) in [14], gave the first proof of the CSW rules. This proof includes three key ingredients. The first is to show that the amplitude from CSW rules, A_{CSW} , has correct collinear and multi-gluon singularities. The second one is to show that all spurious poles related to off-shell continuation are canceled among themselves. This is equivalent to show that A_{CSW} is Lorentz covariant [8]. The last step is the fact that the n -gluon tree amplitude has mass dimension $4 - n$. This step can be replaced by showing that both A_{CSW} and the amplitudes from Feynman rules, $A_{Feynman}$, satisfy the Britto-Cachazo-Feng-Witten (BCFW) recursion relation [14].

The second step of the proof was displayed in detail for tree next-to-MHV (NMHV) amplitudes involving MHV diagrams with a single propagator [8]. This discussion can be apply to the general case but the proof seems to be quite complicated for $N^k MHV$ amplitudes with large k . In fact, NMHV amplitudes are special, not only because each contributing MHV diagram for them always has exactly one propagator, but also because the spurious poles in such amplitudes are always degenerate according to the classification we will give later in this paper. In this paper, we would like to provide another treatment for this step, based on generalization of the treatment in [15, 16] for googly ($\overline{MHV}$) amplitudes. At first sight, the treatment there strongly depends on the special properties of MHV diagrams contribution to googly amplitudes: there is exactly one MHV vertex with four lines and all other vertices are with three lines. As we will show in the main part of this paper, the treatment here can still be non-trivially generalized to the most general cases to show that the spurious poles are canceled among themselves.

Risager provided another proof [19] of CSW rules based on multi-gluon BCFW-like shifts and obtained on-shell recursion relation². Our approach here is complementary to others. We hope that

²Proof of CSW rules for all tree amplitudes in $\mathcal{N} = 4$ super Yang-Mills theory using all-line shift was done in [20].

our proof would lead to new insight to CSW-like rules for amplitudes in various other theories.

In the next section, we will give a brief introduction to the CSW rules. Section 3 is the main part of this paper and provides our proof of the CSW rules. First we introduce specially-defined two-off-shell-line sub-amplitudes and show that their residues at related spurious pole are proportional to certain one-off-shell-line sub-amplitudes. Based on this result, in subsection 3.2, we show that all spurious poles cancel among themselves by dividing them into groups. In subsection 3.3 we finish our proof based on some known facts. The section 4 is devoted to conclusion and some discussions on possible further applications of our approach. Some technical details are putting in several appendices.

2 A brief introduction to CSW rules based on MHV diagrams

Our conventions follow closely the one in [4], especially the signature is chosen to be $(+ - - -)$. By using the crossing symmetry, we make all momenta of external line gluons to be outgoing. For external gluon line with on-shell momentum p_μ , the following decomposition into bispinors will be chosen,

$$p_{\alpha\dot{\alpha}} \equiv p_\mu \sigma_{\alpha\dot{\alpha}}^\mu = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}}, \quad (1)$$

where $\sigma^\mu = (1_{2 \times 2}, \sigma^i)$, with $1_{2 \times 2}$ two-dimensional identity matrix and σ^i being three Pauli matrices. The following Lorentz invariant anti-symmetric inner products

$$\langle \lambda, \mu \rangle = \epsilon_{\alpha\beta} \lambda^\alpha \mu^\beta, \quad (2)$$

$$[\tilde{\lambda}, \tilde{\mu}] = \epsilon_{\dot{\alpha}\dot{\beta}} \tilde{\lambda}^{\dot{\alpha}} \tilde{\mu}^{\dot{\beta}}, \quad (3)$$

are quite useful. We will also denote $\langle \lambda_r, \lambda_s \rangle$ ($[\tilde{\lambda}_r, \tilde{\lambda}_s]$) by $\langle r, s \rangle$ ($[r, s]$). The spinor index α ($\dot{\alpha}$) will be raised (or lowered) by anti-symmetric tensor $\epsilon^{\alpha\beta}$ ($\epsilon^{\dot{\alpha}\dot{\beta}}$) and $\epsilon_{\alpha\beta}$ ($\epsilon_{\dot{\alpha}\dot{\beta}}$),

$$\lambda^\alpha = \epsilon^{\alpha\beta} \lambda_\beta, \quad \lambda_\alpha = \epsilon_{\alpha\beta} \lambda^\beta, \quad (4)$$

with $\epsilon^{\alpha\beta} \epsilon_{\beta\gamma} = \delta_\gamma^\alpha$.³ We omit similar expressions involving $\tilde{\lambda}^{\dot{\alpha}}$ and $\tilde{\lambda}_{\dot{\alpha}}$.

We will always focus on the partial amplitudes $A(1, 2, \dots, n)$ with color factor stripped. These partial amplitudes are defined by the color decomposition of the full amplitudes $\mathcal{M}(1, 2, \dots, n)$ as

$$\mathcal{M}(1, 2, \dots, n) = \sum_{\sigma \in S_n / \mathbf{Z}_n} \text{Tr}(T^{a_{\sigma(1)}} T^{a_{\sigma(2)}} \dots T^{a_{\sigma(n)}}) A(\sigma(1), \sigma(2), \dots, \sigma(n)), \quad (5)$$

Derivation of CSW rules from Lagrangian was studied in [21, 22, 23, 24, 25] and a twistor-action formulation was provided in [26].

³We can explicitly choose $\epsilon_{12} = -\epsilon_{21} = -\epsilon^{12} = \epsilon^{21} = 1$ for $\epsilon_{\alpha\beta}, \epsilon^{\alpha\beta}, \epsilon_{\dot{\alpha}\dot{\beta}}, \epsilon^{\dot{\alpha}\dot{\beta}}$.

where the summation is over the permutation group S_n acting on the n external gluons modulo the cyclic group $\mathbf{Z}_n$ acting as cyclic permutations. Here, without loss of generality, we choose the gauge group to be $U(N)$. T^{a_i} 's are representation matrices (in the fundamental representation) of the generators of the gauge group, with normalization $\text{Tr}(T^a T^b) = \delta^{ab}$. a_i 's are color indices of i th gluon.

From the above color decomposition, we know that we only need to compute the partial amplitude⁴ $A(\sigma(1), \sigma(2), \dots, \sigma(n))$. But this is still a very complicated object since even for small n like $n = 5$, it contains a large number of possible Lorentz invariant combinations of momenta and polarization vectors (see, for example, Fig. 7 of [27]). Since we have introduced bi-spinor for external on-shell momenta $p_i = \lambda_i \tilde{\lambda}_i$, a partial amplitude is specified by λ , $\tilde{\lambda}$ and helicity h of each gluon. We then denote the partial amplitudes simply by $A_n(1^{h_1}, \dots, n^{h_n})$. To achieve this, polarization vectors are constructed as [28, 29]

$$\varepsilon_{\alpha\dot{\alpha}}^{(-)} = \frac{\lambda_{\alpha} \tilde{\mu}_{\dot{\alpha}}}{[\tilde{\lambda}, \tilde{\mu}]}, \quad \varepsilon_{\alpha\dot{\alpha}}^{(+)} = \frac{\mu_{\alpha} \tilde{\lambda}_{\dot{\alpha}}}{\langle \mu, \tilde{\lambda} \rangle}, \quad (6)$$

where μ and $\tilde{\mu}$ are fixed reference spinors. The superscripts of polarization vectors $(+)$, $(-)$ denotes the positive helicity and negative helicity respectively. μ ($\tilde{\mu}$) can be different for different external gluons. Due to gauge invariance, the final results for amplitudes will not depends on the choices of μ 's and $\tilde{\mu}$'s. This spinor helicity trick greatly simplify the computation of amplitudes.

By tree-level MHV amplitudes, we always mean tree amplitudes with exactly two negative-helicity gluons. The results for the n -gluon MHV amplitude is [30, 31]

$$A_n^{\text{MHV}}(1^+, \dots, r^-, \dots, s^-, \dots, n^+) = \frac{\langle r, s \rangle^4}{\prod_{i=1}^n \langle i, i+1 \rangle}. \quad (7)$$

CSW [8] proposed a new way to compute tree level amplitudes in Yang-Mills theory by using tree MHV diagrams where the whole MHV amplitudes, with suitable off-shell continuation, was treated as vertices (called MHV vertices). Since the vertices will be inserted in MHV diagrams, some legs will be internal. The question about how to define λ for such momentum arises. CSW consider the following off-shell continuation for internal gluon with momentum p

$$\lambda_{\alpha} = p_{\alpha\dot{\alpha}} \tilde{\eta}^{\dot{\alpha}}, \quad (8)$$

where $\tilde{\eta}^{\dot{\alpha}}$ is an arbitrary spinor which is fixed for all of the internal lines in all diagrams contributing to a given partial amplitude. For the external gluons, the associated λ is still obtained from the decomposition in (1).

⁴From now on, we will often use ‘amplitudes’ to mean ‘partial amplitudes’.

In order to construct MHV diagram, we connect MHV vertices by propagators. We need to assign a factor $1/p^2$ for each propagator with internal momentum p . It is worth noting that the two ends of each propagator have opposite helicities. The contribution of each MHV diagrams comes from the product of MHV vertices and scalar propagators. Tree level MHV diagrams must be planar ones with the cyclic order of external line particles be consistent with that of external line particles in partial amplitudes to-be-computed. As using Feynman diagrams, the partial amplitudes are obtained from summation over contributions from all tree MHV diagrams with the correct cyclic order of external gluons.

One of the key feature of the CSW rules is that the final result is independent of $\tilde{\eta}$. The key step of our proof here is to show this by generalizing some steps in the computations of googly amplitudes [15, 16]. Our treatment is different from the one in [8].

3 Proof of the CSW rules

3.1 The two-off-shell-line sub-amplitudes and their spurious poles

Spurious poles in MHV diagrams are due to the off-shell continuation of MHV amplitudes. The study of spurious poles in A_{CSW} plays an important role in our proof. These poles appear due to the off-shell continuation in the CSW rules. From this subsection we begin our study of the residues of these poles.

For our convenience we introduce the following notation⁵

$$p_{a,b} \equiv \sum_{i=a}^b p_i, \quad (9)$$

and as reviewed in the previous section,

$$p_{i\alpha\dot{\alpha}} = p_{i\mu} \sigma_{\alpha\dot{\alpha}}^{\mu} = \lambda_{i\alpha} \tilde{\lambda}_{i\dot{\alpha}}, \quad (10)$$

for on-shell external momentum p_i . For generic external momenta p_i 's, $p_{a,b}$ is off-shell for $a \neq b$ and $b \neq a - 2$. For this $p_{a,b}$, CSW [8] introduced a (holomorphic) λ :

$$\lambda_{p_{a,b}} = \sum_{i=a}^b \lambda_i \phi_i, \quad \phi_i \equiv \tilde{\lambda}_{i\dot{\alpha}} \tilde{\eta}^{\dot{\alpha}}, \quad (11)$$

where $\tilde{\eta}$ is a generic auxiliary (anti-holomorphic) spinor. According to the CSW rules [8], this $\lambda_{p_{a,b}}$ will be used in the internal MHV vertex with an internal line of momentum $p_{a,b}$.

⁵When $a > b$, $\sum_{i=a}^b$ always means $\sum_{i=a}^n + \sum_{i=1}^b$.

For generic external momenta p_i 's, we assume that no two $p_{a,b}$'s corresponding λ 's are proportional to each other for generic $\tilde{\eta}$:

$$\lambda_{p_{a,b}} \not\propto \lambda_{p_{c,d}}, \quad (12)$$

except the special case: $p_{c,d} = p_{b+1,a-1} = -p_{a,b}$. Of course, $\lambda_{p_{a,b}}$ is not proportional to any λ_i 's.

Possible spurious poles appear when $\langle \lambda_{p_{n_1,n_2-1}}, \lambda_{p_{n_2,n_3-1}} \rangle$ (with $n_{i+1} - n_i > 1, i = 1, 2$)⁶ or $\langle \lambda_{p_{n_1,n_2-1}}, \lambda_{n_2} \rangle$ vanishes and appears in the denominator of the expression for an MHV vertex.

We refer to the cases with two λ 's from off-shell momenta, like $\langle \lambda_{p_{n_1,n_2-1}}, \lambda_{p_{n_2,n_3-1}} \rangle$ with $n_2 - n_1 > 1, n_3 - n_2 > 1, n_1 - n_3 > 1$ as non-degenerate and all other cases as degenerate.

For definite n_1, n_2 and n_3 , let us define the momenta P_1, P_2, P_3 ,

$$P_1 \equiv p_{n_1,n_2-1}, \quad (13)$$

$$P_2 \equiv p_{n_2,n_3-1}, \quad (14)$$

$$P_3 \equiv p_{n_3,n_1-1}, \quad (15)$$

and the spinors v_1, v_2, v_3 to be

$$v_1 \equiv \lambda_{P_1} = \lambda_{p_{n_1,n_2-1}} = \sum_{i=n_1}^{n_2-1} \lambda_i \phi_i, \quad (16)$$

$$v_2 \equiv \lambda_{P_2} = \lambda_{p_{n_2,n_3-1}} = \sum_{i=n_2}^{n_3-1} \lambda_i \phi_i, \quad (17)$$

$$v_3 \equiv \lambda_{P_3} = \lambda_{p_{n_3,n_1-1}} = \sum_{i=n_3}^{n_1-1} \lambda_i \phi_i. \quad (18)$$

It is easy to see that momentum conservation leads to

$$\sum_{i=1}^3 v_i = 0, \quad (19)$$

which leads to

$$\langle v_1, v_2 \rangle = \langle v_2, v_3 \rangle = \langle v_3, v_1 \rangle. \quad (20)$$

When $\langle v_1, v_2 \rangle = 0$, there exist $\alpha_1, \alpha_2 \in \mathbf{C}$ and spinor v_0 such that

$$v_i = \alpha_i v_0, \quad i = 1, 2. \quad (21)$$

Then $v_3 = -(\alpha_1 + \alpha_2)v_0$. We now define α_3 to be $\alpha_3 \equiv -(\alpha_1 + \alpha_2)$.

For arbitrary function of v_1, v_2 like $F(v_1, v_2) = \frac{f(v_1, v_2)}{\langle v_1, v_2 \rangle}$, we roughly refer $f(v_1, v_2)|_{v_1=\alpha_1 v_0, v_2=\alpha_2 v_0}$ as 'residue' of $F(v_1, v_2)$ at the pole where $\langle v_1, v_2 \rangle = 0$ and simply denote it as $\text{Res}_{\langle v_1, v_2 \rangle=0} F(v_1, v_2)$.




Figure 1: One-off-shell-line sub-amplitude $A^1(P_1^{h_{P_1}}, n_2, \dots, n_1 - 1)$.




Figure 2: Two-off-shell-line sub-amplitude $A^2(P_1^{h_{P_1}}, P_2^{h_{P_2}}, n_3, \dots, n_1 - 1)$.

As in [15], we introduce two kinds of ‘off-shell amplitudes’ constructed by CSW rules. We define $A^1(P_1^{h_{P_1}}, n_2, \dots, n_1 - 1)$ to be the amplitude with one off-shell leg P_1 , see Fig. 1. The all-but-one-negative-helicity one-off-shell-line amplitudes were computed in [15]. The other off-shell amplitude (see Fig. 2), called $A^2(P_1^{h_{P_1}}, P_2^{h_{P_2}}, n_3, \dots, n_1 - 1)$, is defined as the amplitude constructed from all MHV diagrams with two adjacent off-shell legs P_1, P_2 directly attached to the same MHV vertex and the rest by the standard rules (with the external momenta $P_{1,2}$ and $p_{n_3}, \dots, p_{n_1-1}$).

A special note is in order. Because of the constraint that off-shell legs P_1, P_2 must directly attached to the same MHV vertex, $A^2(P_1^{h_{P_1}}, P_2^{h_{P_2}}, n_3, \dots, n_1 - 1)$ is not a sub-amplitude in the usual sense. They can be used in the computation of the amplitude. We will use them in the analysis of the spurious poles of the amplitude. However sometimes we loosely call it a sub-amplitude. Also notice that we will not give explicitly the helicities of the external gluons since these helicities are fixed.

For generic external momenta, The ‘residue’ of $A^2(P_1^{h_{P_1}}, P_2^{h_{P_2}}, n_3, \dots, n_1 - 1)$ at the spurious pole

⁶The label n_i should be understood as modulo n , as in footnote 5.




Figure 3: Diagram decomposition for $A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1)$.

at $\langle v_1, v_2 \rangle = 0$ is the following,

$$\text{Res}_{\langle v_1, v_2 \rangle = 0} A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1) = \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1((P_1 + P_2)^+, n_3, \dots, n_1 - 1), \quad (22)$$

$$\text{Res}_{\langle v_1, v_2 \rangle = 0} A^2(P_1^+, P_2^-, n_3, \dots, n_1 - 1) = \frac{\alpha_2^3}{\alpha_1 \alpha_3^2} A^1((P_1 + P_2)^-, n_3, \dots, n_1 - 1), \quad (23)$$

$$\text{Res}_{\langle v_1, v_2 \rangle = 0} A^2(P_1^-, P_2^+, n_3, \dots, n_1 - 1) = \frac{\alpha_1^3}{\alpha_2 \alpha_3^2} A^1((P_1 + P_2)^-, n_3, \dots, n_1 - 1), \quad (24)$$

$$\text{Res}_{\langle v_1, v_2 \rangle = 0} A^2(P_1^-, P_2^-, n_3, \dots, n_1 - 1) = 0. \quad (25)$$

The proof of eq. (22) is actually not very difficult. To get a feel of what is involved in the proof, we refer the readers to Appendix A for an example, $A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7)$. $A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1)$ is given by a summation over various diagrams. But in any (connected) diagram there is an MHV vertex with two adjacent lines with $\lambda_{P_1} = v_1$ and $\lambda_{P_2} = v_2$. Because we assume the helicities of P_1 and P_2 both to be plus, the dependence on $v_{1,2}$ will only be:

$$\frac{1}{\langle \lambda_L, v_1 \rangle \langle v_1, v_2 \rangle \langle v_2, \lambda_R \rangle}, \quad (26)$$

where λ_L is the spinor of the corresponding momentum to the nearest left of P_1 , and λ_R is the spinor of the corresponding momentum to the nearest right of P_2 , (of the MHV vertex). For $\langle v_1, v_2 \rangle = 0$ this gives a residue:

$$\frac{\alpha_3^2}{\alpha_1 \alpha_2} \frac{1}{\langle \lambda_L, v_3 \rangle \langle v_3, \lambda_R \rangle}. \quad (27)$$

A more complete proof follows.

From the diagram decomposition (see Fig. 3), we have.

$$\begin{aligned}
A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1) &= \sum_{m_3 \geq 2} \sum'_{l_{1(3)}, \dots, l_{m_3(3)}} \sum_{h_{1(3)}, \dots, h_{m_3(3)}} V_{m_3+2}(P_1^+, P_2^+, q_{1(3)}^{h_{q_1(3)}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}) \\
&\times A(n_3, \dots, n_3 + l_{1(3)} - 1, -q_{1(3)}^{-h_{1(3)}}) \frac{1}{q_{1(3)}^2} A(n_3 + l_{1(3)}, \dots, n_3 + l_{1(3)} + l_{2(3)} - 1, -q_{2(3)}^{-h_{2(3)}}) \\
&\times \frac{1}{q_{2(3)}^2} \times \dots \times A(n_1 - l_{m_3(3)}, \dots, n_1 - 1, -q_{m_3(3)}^{-h_{m_3(3)}}) \frac{1}{q_{m_3(3)}^2}, \tag{28}
\end{aligned}$$

where the symbol $\sum'$ in the second summation means sum over $l_{1(3)}, \dots, l_{m_3(3)}$ under the constraints

$$l_{i(3)} > 0, \quad i = 1, \dots, m_3, \quad \sum_{i=1}^{m_3} l_{i(3)} = n_1 - n_3, \tag{29}$$

and

$$q_{i(3)} = p_{n_3 + \sum_{j=1}^{i-1} l_{j(3)}, n_3 + \sum_{j=1}^i l_{j(3)} - 1} \tag{30}$$

Here and below the vertex like V_{m_3+2} is understood to be vanishing if it does not satisfy the MHV conditions.

The involved MHV vertex $V_{m_3+2}(P_1^+, P_2^+, q_{1(3)}^{h_{q_1(3)}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}})$ is,

$$\begin{aligned}
&V_{m_3+2}(P_1^+, P_2^+, q_{1(3)}^{h_{q_1(3)}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}) \\
&= \frac{\langle \tilde{v}_{j(3)}, \tilde{v}_{k(3)} \rangle^4}{\langle v_1, v_2 \rangle \langle v_2, \tilde{v}_{1(3)} \rangle (\prod_{i=1}^{m_3-1} \langle \tilde{v}_{i(3)}, \tilde{v}_{(i+1)(3)} \rangle) \langle \tilde{v}_{m_3(3)}, v_1 \rangle}, \tag{31}
\end{aligned}$$

with $m_3 \geq 2$. Here $\tilde{v}_{i(3)}$ is the spinor corresponding to the momentum $q_{i(3)}$ based on the CSW rules reviewed in the previous section and we have assumed

$$h_{q_{j(3)}} = h_{q_{k(3)}} = -1, \quad h_{q_{i(3)}} = +1, \quad \text{for } 1 \leq i \leq m_3, i \neq j, k. \tag{32}$$

Notice that $\tilde{v}_{1(3)}$ and $\tilde{v}_{m_3(3)}$ in (31) are just λ_L and λ_R in (26). The ‘residue’ of this vertex at the pole $\langle v_1, v_2 \rangle = 0$ is

$$\begin{aligned}
&\text{Res}_{\langle v_1, v_2 \rangle = 0} V_{m_3+2}(P_1^+, P_2^+, q_{1(3)}^{h_{q_1(3)}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}) \\
&= \frac{\alpha_3^2}{\alpha_1 \alpha_2} \frac{\langle \tilde{v}_{j(3)}, \tilde{v}_{k(3)} \rangle^4}{\langle -v_3, \tilde{v}_{1(3)} \rangle (\prod_{i=1}^{m_3-1} \langle \tilde{v}_{i(3)}, \tilde{v}_{(i+1)(3)} \rangle) \langle \tilde{v}_{m_3(3)}, -v_3 \rangle} \\
&= \frac{\alpha_3^2}{\alpha_1 \alpha_2} V_{m_3+1}(-P_3^+, q_{1(3)}^{h_{q_1(3)}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}). \tag{33}
\end{aligned}$$

We see that the ‘residue’ of MHV vertex at $\langle v_1, v_2 \rangle = 0$ is proportional to another MHV vertex with the number of legs reduced by one. This simple fact inherited from the collinear limit of MHV

amplitudes plays an important role in the proof here. So the residue of $A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1)$ at the same pole is

$$\begin{aligned}
\text{Res}_{\langle v_1, v_2 \rangle=0} A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1) &= \frac{\alpha_3^2}{\alpha_1 \alpha_2} \sum_{m_3 \geq 2} \sum'_{l_{1(3)}, \dots, l_{m_3(3)}} \sum_{h_{1(3)}, \dots, h_{m_3(3)}} \\
&\quad V_{m_3+1}(-P_3^+, q_{1(3)}^{h_{q_{1(3)}}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}) A(n_3, \dots, n_3 + l_{1(3)} - 1, -q_{1(3)}^{-h_{1(3)}}) \frac{1}{q_{1(3)}^2} \\
&\quad \times A(n_3 + l_{1(3)}, \dots, n_3 + l_{1(3)} + l_{2(3)} - 1, -q_{2(3)}^{-h_{2(3)}}) \frac{1}{q_{2(3)}^2} \dots \\
&\quad \times A(n_1 - l_{m_3(3)}, \dots, n_1 - 1, -q_{m_3(3)}^{-h_{m_3(3)}}) \frac{1}{q_{m_3(3)}^2}. \tag{34}
\end{aligned}$$

Notice that we have the following result

$$\begin{aligned}
&\sum_{m_3 \geq 2} \sum'_{l_{1(3)}, \dots, l_{m_3(3)}} \sum_{h_{1(3)}, \dots, h_{m_3(3)}} V_{m_3+1}(-P_3^+, q_{1(3)}^{h_{q_{1(3)}}}, \dots, q_{m_3(3)}^{h_{q_{m_3(3)}}}) \\
&\quad \times A(n_3, \dots, n_3 + l_{1(3)} - 1, -q_{1(3)}^{-h_{1(3)}}) \frac{1}{q_{1(3)}^2} \\
&\quad \times A(n_3 + l_{1(3)}, \dots, n_3 + l_{1(3)} + l_{2(3)} - 1, -q_{2(3)}^{-h_{2(3)}}) \\
&\quad \times \frac{1}{q_{2(3)}^2} \times \dots \times A(n_1 - l_{m_3(3)}, \dots, n_1 - 1, -q_{m_3(3)}^{-h_{m_3(3)}}) \frac{1}{q_{m_3(3)}^2} \\
&= A^1(n_3, \dots, n_1 - 1, -P_3^+), \tag{35}
\end{aligned}$$

from the diagram decomposition for amplitudes $A^1(n_3, \dots, n_1 - 1, -p_{n_3, n_1-1}^+)$ which is the same as Fig. 3 but with the two off-shell momentum lines P_1 and P_2 combined into one momentum line $-P_3 = P_1 + P_2$.⁷ This finishes the proof of eq. (22).

The proofs of the next two identities, eq. (23) and eq. (24), are quite similar. The only point to note is that there is an extra factor $\langle v_{1,2}, \lambda_* \rangle^4$ from the MHV amplitude because one of the off-shell lines has negative helicity. (λ_* is the spinor of the other negative helicity line.) This only modifies the pre-factor of the residue. The proof of the last identities, eq. (25), is trivial because there is an extra factor $\langle v_1, v_2 \rangle^4$ in the numerator and there is no pole.

3.2 The independence of the amplitudes on $\tilde{\eta}$

In this section we will show that A_{CSW} is independent of $\tilde{\eta}$. In another word, the obtained amplitude is Lorentz covariant.

⁷ Similar decomposition was used in [16] to prove that the amplitudes from CSW rules satisfying the charge conjugation identity (also known as the color-ordered reversed relation) and the dual Ward identity (also known as the $U(1)$ decoupling relation).




Figure 4: The diagram contributing to R_3 .

It is not hard to see that A_{CSW} is invariant under the transformation $\tilde{\eta}^{\dot{\alpha}} \rightarrow t\tilde{\eta}^{\dot{\alpha}}$. This can be obtained from the facts that the MHV amplitude (as general amplitudes) will scale by a factor $t_i^{-2h_i}$ when we scale λ_i to $t_i\lambda_i$ for fixed i and that the two ends of each propagator in the MHV diagrams have opposite helicities.

Then if we define $\tilde{t} = \tilde{\eta}^1/\tilde{\eta}^2$, the amplitude from CSW rules can be considered as a function of $\lambda_i, \tilde{\lambda}_i$ and $\tilde{t}$. We will show that the amplitude is a holomorphic function of $\tilde{t} \in \mathbf{C}^*(= \mathbf{C} \cup \{\infty\})$ for generic external momenta. To do this, we just need to prove that the residue for arbitrary $\tilde{t}$ vanishes.

Let us consider the non-degenerate spurious poles mentioned in the previous subsection and show that all such poles of $\tilde{t}$ cancel among themselves. The proof of the cancelation of spurious pole for degenerate case is very similar to the non-degenerate case, and we will discuss these cases briefly in Appendix B.

Consider the poles from $\langle v_1, v_2 \rangle = 0$ with $h_{P_1} = h_{P_2} = +1$, we will show these poles will be canceled by poles from $\langle v_2, v_3 \rangle = 0$ with $h_{P_2} = +1, h_{P_3} = -1$, and the poles from $\langle v_3, v_1 \rangle = 0$ with $h_{P_3} = -1, h_{P_1} = +1$.

The pole from $\langle v_1, v_2 \rangle = 0$ (recall that we have assumed $h_{P_1} = h_{P_2} = +1$) is from the following contributions (see Fig. 4),

$$R_3 = A^1(-P_1^-, n_1, \dots, n_2 - 1) \frac{1}{P_1^2} A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^2(P_1^+, P_2^+, n_3, \dots, n_1 - 1). \quad (36)$$

By using eq. (22), the ‘residue’ of these contributions at $\langle v_1, v_2 \rangle = 0$ is

$$\begin{aligned} \tilde{R}_3 &= \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1(-P_1^-, n_1, \dots, n_2 - 1) \frac{1}{P_1^2} A^1(-P_2^-, n_2, \dots, n_3 - 1) \\ &\times \frac{1}{P_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1). \end{aligned} \quad (37)$$

We now consider the pole at $\langle v_2, v_3 \rangle = 0$ ⁸ with $h_{P_2} = +1, h_{P_3} = -1$. Such pole is from

$$R_1 = A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2} A^2(P_2^+, P_3^-, n_1, \dots, n_2 - 1). \quad (38)$$

Using eq. (23), the residue of the above contributions at $\langle v_2, v_3 \rangle (= \langle v_1, v_2 \rangle) = 0$ is

$$\begin{aligned} \tilde{R}_1 &= \frac{\alpha_3^3}{\alpha_1^2 \alpha_2} A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} \\ &\times A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2} A^1(-P_1^-, n_1, \dots, n_2 - 1), \end{aligned} \quad (39)$$

Now we turn to consider the pole at $\langle v_3, v_1 \rangle = 0$ with $h_{P_3} = -1, h_{P_1} = +1$ from

$$R_2 = A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2} A^1(-P_1^-, n_1, \dots, n_2 - 1) \frac{1}{P_1^2} A^2(P_3^-, P_1^+, n_2, \dots, n_3 - 1). \quad (40)$$

From eq. (24), we know that the residue of the above contributions at $\langle v_3, v_1 \rangle (= \langle v_1, v_2 \rangle) = 0$ is

$$\begin{aligned} \tilde{R}_2 &= \frac{\alpha^3}{\alpha_1 \alpha_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2} A^1(-P_1^-, n_1, \dots, n_2 - 1) \frac{1}{P_1^2} \\ &\times A^1(-P_2^-, n_2, \dots, n_3 - 1). \end{aligned} \quad (41)$$

The sum of $\tilde{R}_1, \tilde{R}_2, \tilde{R}_3$ is

$$\begin{aligned} \sum_{i=1}^3 \tilde{R}_i &= \left(\frac{P_1^2}{\alpha_1} + \frac{P_2^2}{\alpha_2} + \frac{P_3^2}{\alpha_3} \right) \frac{\alpha_3^3}{\alpha_1 \alpha_2} A^1(-P_1^-, n_1, \dots, n_2 - 1) \frac{1}{P_1^2} \\ &\times A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2}. \end{aligned} \quad (42)$$

We will prove in Appendix C that

$$\frac{P_1^2}{\alpha_1} + \frac{P_2^2}{\alpha_2} + \frac{P_3^2}{\alpha_3} = 0, \quad (43)$$

and this leads to

$$\sum_{i=1}^3 \tilde{R}_i = 0. \quad (44)$$

Notice that there are other poles from $\langle v_1, v_2 \rangle = 0$, but these other poles can be divided into the following two groups:

- the poles from $\langle v_1, v_2 \rangle = 0$ with $h_{P_1} = +1, h_{P_2} = -1$, poles from $\langle v_2, v_3 \rangle = 0$ with $h_{P_2} = -1, h_{P_3} = +1$, and the poles from $\langle v_3, v_1 \rangle = 0$ with $h_{P_3} = h_{P_1} = +1$;

⁸Recall that $\langle v_1, v_2 \rangle = \langle v_2, v_3 \rangle = \langle v_3, v_1 \rangle$ even away from the poles.

- the poles from $\langle v_1, v_2 \rangle = 0$ with $h_{P_1} = -1, h_{P_2} = +1$, poles from $\langle v_2, v_3 \rangle = 0$ with $h_{P_2} = h_{P_3} = +1$, and the poles from $\langle v_3, v_1 \rangle = 0$ with $h_{P_3} = +1, h_{P_1} = -1$;

Exactly as what we have done above, one can show that the poles in each group cancel each other.

The above classification of non-degenerate poles at $\langle v_1, v_2 \rangle (= \langle v_2, v_3 \rangle = \langle v_3, v_1 \rangle) = 0$ can be summarized into the following table,

poles	$\langle v_1, v_2 \rangle = 0$	$\langle v_2, v_3 \rangle = 0$	$\langle v_3, v_1 \rangle = 0$
	$h_{P_1} = h_{P_2} = +1$	$h_{P_2} = +1, h_{P_3} = -1$	$h_{P_3} = -1, h_{P_1} = +1$
	$h_{P_1} = +1, h_{P_2} = -1$	$h_{P_2} = -1, h_{P_3} = +1$	$h_{P_3} = h_{P_1} = +1$
	$h_{P_1} = -1, h_{P_2} = +1$	$h_{P_2} = h_{P_3} = +1$	$h_{P_3} = +1, h_{P_1} = -1$

The residues from the helicity configuration in each line cancel among themselves as stated above.

3.3 The Completion of the proof

This part of the proof is not new. The ingredients have appeared in [8] and [14]. After proving that the partial amplitudes from the CSW rules, A_{CSW} , is independent of $\tilde{\eta}$, we only need to consider the pole of A_{CSW} as function of $\lambda, \tilde{\lambda}$. In [8], it was shown that the CSW rules correctly reproduce all the collinear singularities and multi-gluon singularities. So $A_{\text{CSW}} - A_{\text{Feynman}}$ can only be a polynomial of $\lambda_i, \tilde{\lambda}_i$ as pointed in [14], where A_{Feynman} is the tree-level amplitudes from the Feynman rules. In [14], as a step to their proof of CSW rules, it was noticed that the mass dimension of tree amplitudes with n gluon is $4 - n$. We will explicitly show that both A_{CSW} and A_{Feynman} has mass dimension $4 - n$ in Appendix D. This leads to $A_{\text{CSW}} = A_{\text{Feynman}}$ when $n > 4$. Also as mentioned in [14], the case with $n = 4$ can be confirmed directly. This completes the proof.

4 Conclusion

In this paper, we provided a new proof of the CSW rules based on MHV diagrams for gluonic tree amplitudes. Our new contribution is an explicit proof of the cancelation of spurious poles originating from the off-shell continuation of MHV amplitudes. This leads to the conclusion that A_{CSW} is Lorentz invariant. This step is based on the study of spurious poles of specially-defined two-off-shell-line sub-amplitude, and we find that the ‘residue’ is proportional to certain one-off-shell-line sub-amplitude.

It is interesting to study the generalization of our approach to other theories or loop level. Our new approach can be applied to amplitudes from CSW rules involving fermions [17] by similarly generalizing the treatment in [18]. It is interesting to study the generalization of our approach to CSW rules for QED [38] and theories involving massive particles like Higgs particle [39, 40, 41, 42, 43]. The story for gravity amplitudes is much more complicated. Risager’s idea [19] was used in [32] to

propose a CSW-like rules for tree-level graviton amplitudes⁹. However it was found that this proposal only works when the number of graviton n is less than 12 [34, 35], see also further studies in [36, 37]. It is interesting to use our approach to show that the spurious poles cancel among themselves when $n < 12$. This cancelation fails when $n \geq 12$. We hope that these studies will provide hints on how to obtain a CSW-like rules which are valid for all cases.

CSW rules were successfully generalized to one-loop level by reproducing some amplitudes in $\mathcal{N} = 4$ SYM theories [44, 45, 46, 47, 48]. It is interesting by using our approach to check whether the integrand of the planar one-loop amplitudes from CSW rules [44, 45, 46] has no spurious poles or these poles only disappear after loop integration.

Acknowledgments

J.-B. W. would like to thank Bin Chen, Jian-Xin Lu, Gang Yang for very helpful discussions. He would also like to thank Peking University since a key idea was obtained when he walked crossing the street between the east gate and School of Physics. C.-J. Zhu would like to thank Bo Feng for discussions. The work of J.-B. W. and W.-J. Z. was supported by the National Natural Science Foundation of China, Grant No. 11975164, 11935009, 12047502, 11947301, and Natural Science Foundation of Tianjin under Grant No. 20JCYBJC00910. The work of C.-J. Zhu was supported in part by fund from Hunan University of Arts and Science. The diagrams were drawn using JavaDraw package [49, 50].

A Cancelation of non-degenerate spurious poles: an example

For the convenience of the readers, we use the googly amplitude $A(1^+, 2^-, 3^+, 4^-, 5^-, 6^-, 7^-)$ to display the procedure of the cancelation of non-degenerate spurious poles. We consider the situation with $n_1 = 1, n_2 = 3, n_3 = 5$.

The diagrams with a pole at $\langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle = 0$ and satisfying $h_{p_{1,2}} = h_{p_{3,4}} = +1$ give contributions proportional to the two-off-line sub-amplitude $A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7)$:

$$R_3 = A^1(1, 2, -p_{1,2}^-) \frac{1}{p_{1,2}^2} A^1(3, 4, -p_{3,4}^-) \frac{1}{p_{3,4}^2} A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7). \quad (45)$$

⁹Earlier attempts can be found in [33, 16].




Figure 5: The diagrammatic decomposition for $A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7)$.

where

$$A^1(1, 2, -p_{1,2}^-) = \frac{\langle \lambda_2, \lambda_{p_{1,2}} \rangle^3}{\langle 1, 2 \rangle \langle \lambda_{p_{1,2}}, \lambda_1 \rangle}, \quad (46)$$

$$A^1(3, 4, -p_{3,4}^-) = \frac{\langle \lambda_4, \lambda_{p_{3,4}} \rangle^3}{\langle 3, 4 \rangle \langle \lambda_{p_{3,4}}, \lambda_3 \rangle}, \quad (47)$$

$$\begin{aligned} A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7) &= \frac{\langle \lambda_5, \lambda_{p_{6,7}} \rangle^3}{\langle \lambda_{p_{6,7}}, \lambda_{p_{1,2}} \rangle \langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle \langle \lambda_{p_{3,4}}, \lambda_5 \rangle} \frac{1}{p_{6,7}^2} A^1(-p_{6,7}^+, p_6^-, p_7^-) \\ &+ \frac{\langle \lambda_{p_{5,6}}, \lambda_7 \rangle^3}{\langle \lambda_7, \lambda_{p_{1,2}} \rangle \langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle \langle \lambda_{p_{3,4}}, \lambda_{p_{5,6}} \rangle} \frac{1}{p_{5,6}^2} A^1(-p_{5,6}^+, p_5^-, p_6^-), \end{aligned} \quad (48)$$

$$A^1(-p_{6,7}^+, 6, 7) = \frac{\langle 6, 7 \rangle^3}{\langle \lambda_7, \lambda_{p_{6,7}} \rangle \langle \lambda_{p_{6,7}}, \lambda_6 \rangle}, \quad (49)$$

$$A^1(-p_{5,6}^+, 5, 6) = \frac{\langle 5, 6 \rangle^3}{\langle \lambda_6, \lambda_{p_{5,6}} \rangle \langle \lambda_{p_{5,6}}, \lambda_5 \rangle}. \quad (50)$$

Here we have used the diagram decomposition for $A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7)$ as shown in Fig. 5.

The ‘residue’ of $A^2(p_{1,2}^+, p_{3,4}^+, 5, 6, 7)$ at $\langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle = 0$ is

$$\begin{aligned} &\frac{\alpha_3^2}{\alpha_1 \alpha_2} \left(\frac{\langle \lambda_5, \lambda_{p_{6,7}} \rangle^3}{\langle \lambda_{p_{6,7}}, \lambda_{-p_{5,7}} \rangle \langle \lambda_{-p_{5,7}}, \lambda_5 \rangle} \frac{1}{p_{6,7}^2} A^1(-p_{6,7}^+, 6, 7) \right. \\ &\quad \left. + \frac{\langle \lambda_{p_{5,6}}, \lambda_7 \rangle^3}{\langle \lambda_7, \lambda_{-p_{5,7}} \rangle \langle \lambda_{-p_{5,7}}, \lambda_{p_{5,6}} \rangle} \frac{1}{p_{5,6}^2} A^1(-p_{5,6}^+, 5, 6) \right) \\ &= \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1(-p_{5,7}^+, 5, 6, 7). \end{aligned} \quad (51)$$

This gives the complete residues from R_3 :

$$\tilde{R}_3 = \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{3,4}^2} A^1(-p_{5,7}^+, 5, 6, 7). \quad (52)$$

The diagram involving a factor $1/\langle \lambda_{p_{3,4}}, \lambda_{p_{5,7}} \rangle$ with $h_{p_{3,4}} = +1, h_{p_{5,7}} = -1$ is shown in Fig. 6. There is only one diagram contributing to two-off-shell-line sub-amplitude $A^2(p_{3,4}^+, p_{5,7}^-, 1, 2)$, as shown in




Figure 6: The diagram gives contribution R_1 which has a pole at $\langle \lambda_{p_{3,4}}, \lambda_{p_{5,7}} \rangle = 0$.




Figure 7: The diagram contributing to the two-off-shell-line sub-amplitude $A^2(p_{3,4}^+, p_{5,7}^-, 1, 2)$.

Fig. 7. This gives a contribution:

$$\begin{aligned}
 R_1 &= A^2(p_{3,4}^+, p_{5,7}^-, 1, 2) \frac{1}{p_{3,4}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{5,7}^2} A^1(-p_{5,7}^+, 5, 6, 7) \\
 &= \frac{\langle \lambda_{p_{5,7}}, \lambda_2 \rangle^4}{\langle 1, 2 \rangle \langle \lambda_2, \lambda_{p_{3,4}} \rangle \langle \lambda_{p_{3,4}}, \lambda_{p_{5,7}} \rangle \langle \lambda_{p_{5,7}}, \lambda_1 \rangle} \frac{1}{p_{3,4}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{5,7}^2} A^1(-p_{5,7}^+, 5, 6, 7). \quad (53)
 \end{aligned}$$

The ‘residue’ of R_2 at $\langle \lambda_{p_{3,4}}, \lambda_{p_{5,7}} \rangle (= \langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle) = 0$ is

$$\begin{aligned}
 \tilde{R}_1 &= \frac{\alpha_3^3}{\alpha_1^2 \alpha_2} \frac{\langle \lambda_{-p_{1,2}}, \lambda_2 \rangle^3}{\langle 1, 2 \rangle \langle \lambda_{-p_{1,2}}, \lambda_1 \rangle} \frac{1}{p_{3,4}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{5,7}^2} A^1(-p_{5,7}^+, 5, 6, 7) \\
 &= \frac{\alpha_3^3}{\alpha_1^2 \alpha_2} A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{3,4}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{5,7}^2} A^1(-p_{5,7}^+, 5, 6, 7). \quad (54)
 \end{aligned}$$

The diagram involving pole at $\langle \lambda_{p_{5,7}}, \lambda_{p_{1,2}} \rangle = 0$ with $h_{p_{5,7}} = -1, h_{p_{1,2}} = +1$ is shown in Fig. 8. The related two-off-shell-line sub-amplitude appearing here is $A^2(p_{5,7}^-, p_{1,2}^+, 3, 4)$, as shown in Fig. 9. This gives a contribution:

$$\begin{aligned}
 R_2 &= A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{5,7}^+, 5, 6, 7) \frac{1}{p_{5,7}^2} A^2(p_{5,7}^-, p_{1,2}^+, 3, 4) \\
 &= A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{5,7}^+, 5, 6, 7) \frac{1}{p_{5,7}^2} \frac{\langle \lambda_4, \lambda_{p_{5,7}} \rangle^3}{\langle \lambda_{p_{5,7}}, \lambda_{p_{1,2}} \rangle \langle \lambda_{p_{1,2}}, \lambda_3 \rangle \langle 3, 4 \rangle} \quad (55)
 \end{aligned}$$




Figure 8: The diagram gives contribution R_2 which has a pole at $\langle \lambda_{p_{5,7}}, \lambda_{p_{1,2}} \rangle = 0$.




Figure 9: The diagram contributing to the two-off-shell-line sub-amplitude $A^2(p_{5,7}^-, p_{1,2}^+, 3, 4)$.

The ‘residue’ of R_2 at $\langle \lambda_{p_{5,7}}, \lambda_{p_{1,2}} \rangle (= \langle \lambda_{p_{1,2}}, \lambda_{p_{3,4}} \rangle) = 0$ is

$$\begin{aligned} \tilde{R}_2 &= \frac{\alpha_3^3}{\alpha_1 \alpha_2^2} A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{5,7}^+, 5, 6, 7) \frac{1}{p_{5,7}^2} \frac{\langle \lambda_4, \lambda_{-p_{3,4}} \rangle^3}{\langle \lambda_{-p_{3,4}}, \lambda_3 \rangle \langle 3, 4 \rangle} \\ &= \frac{\alpha_3^3}{\alpha_1 \alpha_2^2} A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{5,7}^+, 5, 6, 7) \frac{1}{p_{5,7}^2} A^1(-p_{3,4}^-, 3, 4). \end{aligned} \quad (56)$$

The sum of the above three ‘residues’ are

$$\sum_{i=1}^3 \tilde{R}_i = \left(\frac{p_{1,2}^2}{\alpha_1} + \frac{p_{3,4}^2}{\alpha_2} + \frac{p_{5,7}^2}{\alpha_3} \right) \frac{\alpha_3^3}{\alpha_1 \alpha_2} A^1(-p_{1,2}^-, 1, 2) \frac{1}{p_{1,2}^2} A^1(-p_{3,4}^-, 3, 4) \frac{1}{p_{3,4}^2} A^1(-p_{5,7}^+, 5, 6, 7) \frac{1}{p_{5,7}^2}, \quad (57)$$

which vanishes due to eq. (43) proved in Appendix C.

B The degenerate case

In this appendix, we briefly display the cancelation of spurious poles for the degenerate case. We start with the pole at $\langle \lambda_{n_1}, \lambda_{p_{n_1+1}, n_{3-1}} \rangle = 0$ and assume that $h_{n_1} = +1$. (The case with $h_{n_1} = -1$ can be treated similarly for generic external momenta.) This means that $n_2 = n_1 + 1$. Momentum conservation

$$p_{n_1} + P_2 + P_3 = 0, \quad (58)$$

leads to

$$\lambda_{n_1}\phi_{n_1} + \lambda_{P_2} + \lambda_{P_3} = 0. \quad (59)$$

From this we get

$$\langle \lambda_{n_1}, \lambda_{P_2} \rangle = \langle \lambda_{P_3}, \lambda_{n_1} \rangle = \frac{1}{\phi_{n_1}} \langle \lambda_{P_2}, \lambda_{P_3} \rangle. \quad (60)$$

When $\langle \lambda_{n_1}, \lambda_{P_2} \rangle = 0$, we have that there should exist v_0 such that

$$\lambda_{n_1} = \alpha_1 v_0, \lambda_{P_2} = \alpha_2 v_0, \lambda_{P_3} = \alpha_3 v_0, \quad (61)$$

with the following constraint

$$\phi_{n_1}\alpha_1 + \alpha_2 + \alpha_3 = 0, \quad (62)$$

satisfied.

Let us consider the above pole with $h_{P_2} = +1$. As in the non-degenerate case, we get

$$\text{Res}_{\langle \lambda_{n_1}, \lambda_{P_2} \rangle = 0} V_{m_3+2}(n_1, P_2^+, q_{1(3)}^{h_{1(3)}}, \dots, q_{m_3(3)}^{h_{m_3(3)}}) = \frac{\alpha_3^2}{\alpha_1 \alpha_2} V_{m_3+1}(-P_3^+, q_{1(3)}^{h_{1(3)}}, \dots, q_{m_3(3)}^{h_{m_3(3)}}), \quad (63)$$

which gives

$$\text{Res}_{\langle \lambda_{n_1}, \lambda_{P_2} \rangle = 0} A^1(P_2^+, n_3, \dots, n_1 - 1, n_1) = \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1(-P_3^+, n_3, \dots, n_1 - 1). \quad (64)$$

So the ‘residue’ of

$$R_3 = A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^1(P_2^+, n_3, \dots, n_1 - 1, n_1), \quad (65)$$

at the pole $\langle \lambda_{n_1}, \lambda_{P_2} \rangle = 0$ is

$$\tilde{R}_3 = \frac{\alpha_3^2}{\alpha_1 \alpha_2} A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1). \quad (66)$$

Similarly, the ‘residue’ of

$$R_2 = A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2} A^1(P_3^-, n_1, \dots, n_3 - 1), \quad (67)$$

at the pole $\langle \lambda_{P_3}, \lambda_{n_1} \rangle = 0$ is

$$\tilde{R}_2 = \frac{\alpha_3^3}{\alpha_1 \alpha_2^2} A^1(-P_2^-, n_2, \dots, n_3 - 1) A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2}. \quad (68)$$

So we have

$$\begin{aligned} \tilde{R}_2 + \tilde{R}_3 &= \frac{\alpha_3^3}{\alpha_1 \alpha_2} \left(\frac{P_2^2}{\alpha_2} + \frac{P_3^2}{\alpha_3} \right) \\ &\times A^1(-P_2^-, n_2, \dots, n_3 - 1) \frac{1}{P_2^2} A^1(-P_3^+, n_3, \dots, n_1 - 1) \frac{1}{P_3^2}. \end{aligned} \quad (69)$$

From the proof of the identity eq. (43) in Appendix C, we get

$$\begin{aligned} \frac{P_2^2}{\alpha_2} + \frac{P_3^2}{\alpha_3} &= \frac{1}{\Delta} \left(\tilde{\eta}^2 \sum_{1 \leq i \leq n, i \neq n_1} \tilde{\lambda}_{i1} \langle \lambda_i, v_0 \rangle - \tilde{\eta}^1 \sum_{1 \leq i \leq n, i \neq n_1} \tilde{\lambda}_{i2} \langle \lambda_i, v_0 \rangle \right) \\ &= -\frac{1}{\Delta} \left(\tilde{\eta}^2 \tilde{\lambda}_{n_1 1} \langle \lambda_{n_1}, v_0 \rangle - \tilde{\eta}^1 \tilde{\lambda}_{n_1 2} \langle \lambda_{n_1}, v_0 \rangle \right) = 0, \end{aligned} \quad (70)$$

where we have used that λ_{n_1} is proportional to v_0 at the pole. So finally we get $\tilde{R}_2 + \tilde{R}_3 = 0$. Other cases can be treated in the same way.

C The proof of eq. (43)

From [15], we get

$$P_1^2 = p_{n_1, n_2-1}^2 = \sum_{i=n_1}^{n_2-1} \frac{[i, k]}{\phi_i \phi_k} \langle \lambda_{p_i}, \lambda_{p_{n_1, n_2-1}} \rangle, \quad (71)$$

with k can be any integer satisfying $1 \leq k \leq n$.

Recall that

$$\phi_i = \tilde{\lambda}_{i\dot{\alpha}} \tilde{\eta}^{\dot{\alpha}} = \tilde{\lambda}_{i1} \tilde{\eta}^1 + \tilde{\lambda}_{i2} \tilde{\eta}^2, \quad (72)$$

let us further define

$$\psi_i \equiv \tilde{\lambda}_{i1} \tilde{\eta}^2 - \tilde{\lambda}_{i2} \tilde{\eta}^1, \quad 1 \leq i \leq n. \quad (73)$$

From the above two equations, we can express $\tilde{\lambda}_{i\dot{\alpha}}$ in terms of $\phi_i, \psi_i, \tilde{\eta}^{\dot{\alpha}}$ as

$$\tilde{\lambda}_{i1} = \frac{\tilde{\eta}^1 \phi_i + \tilde{\eta}^2 \psi_i}{(\tilde{\eta}^1)^2 + (\tilde{\eta}^2)^2}, \quad (74)$$

$$\tilde{\lambda}_{i2} = \frac{\tilde{\eta}^2 \phi_i - \tilde{\eta}^1 \psi_i}{(\tilde{\eta}^1)^2 + (\tilde{\eta}^2)^2}. \quad (75)$$

Then we get

$$[i, k] = \tilde{\lambda}_{i1} \tilde{\lambda}_{k2} - \tilde{\lambda}_{i2} \tilde{\lambda}_{k1} = \frac{\psi_i \phi_k - \psi_k \phi_i}{(\tilde{\eta}^1)^2 + (\tilde{\eta}^2)^2}, \quad (76)$$

and so

$$\frac{[i, k]}{\phi_i \phi_k} = \frac{1}{(\tilde{\eta}^1)^2 + (\tilde{\eta}^2)^2} \left(\frac{\psi_i}{\phi_i} - \frac{\psi_k}{\phi_k} \right). \quad (77)$$

Now we define

$$\Delta \equiv (\tilde{\eta}^1)^2 + (\tilde{\eta}^2)^2, \quad \varphi_i = \frac{\psi_i}{\phi_i}, \quad 1 \leq i \leq n, \quad (78)$$

we have

$$\frac{[i, k]}{\phi_i \phi_k} = \frac{\varphi_i - \varphi_k}{\Delta}. \quad (79)$$

From this result and (71), we obtain

$$P_1^2 = \frac{1}{\Delta} \sum_{i=n_1}^{n_2-1} (\varphi_i - \varphi_k) \langle \lambda_{p_i}, \lambda_{p_{n_1, n_2-1}} \rangle. \quad (80)$$

Obviously

$$\sum_{i=n_1}^{n_2-1} \varphi_k \langle \lambda_{p_i}, \lambda_{p_{n_1, n_2-1}} \rangle = \varphi_k \langle \sum_{i=n_1}^{n_2-1} \lambda_{p_i}, \lambda_{p_{n_1, n_2-1}} \rangle = 0. \quad (81)$$

So

$$P_1^2 = \frac{1}{\Delta} \sum_{i=n_1}^{n_2-1} \varphi_i \langle \lambda_{p_i}, v_1 \rangle \quad (82)$$

Then

$$\begin{aligned} \frac{P_1^2}{\alpha_1} &= \frac{1}{\Delta} \sum_{i=n_1}^{n_2-1} \varphi_i \langle \lambda_{p_i}, v_0 \rangle \\ &= \frac{1}{\Delta} \sum_{n_1}^{n_2-1} \varphi_i \phi_i \langle \lambda_i, v_0 \rangle \\ &= \frac{1}{\Delta} \sum_{n_1}^{n_2-1} \psi_i \langle \lambda_i, v_0 \rangle, \end{aligned} \quad (83)$$

where we have used $v_1 = \alpha_1 v_0$ and $\varphi_i = \frac{\psi_i}{\phi_i}$.

Similarly, we have

$$\frac{P_2^2}{\alpha_2} = \frac{1}{\Delta} \sum_{i=n_2}^{n_3-1} \psi_i \langle \lambda_i, v_0 \rangle, \quad (84)$$

$$\frac{P_3^2}{\alpha_2} = \frac{1}{\Delta} \sum_{i=n_3}^{n_1-1} \psi_i \langle \lambda_i, v_0 \rangle, \quad (85)$$

From these results, we get

$$\begin{aligned} \frac{P_1^2}{\alpha_1} + \frac{P_2^2}{\alpha_2} + \frac{P_3^2}{\alpha_3} &= \frac{1}{\Delta} \sum_{i=1}^n \psi_i \langle \lambda_i, v_0 \rangle \\ &= \frac{1}{\Delta} \sum_{i=1}^n \left(\tilde{\lambda}_{i1} \tilde{\eta}^2 - \tilde{\lambda}_{i2} \tilde{\eta}^1 \right) \langle \lambda_i, v_0 \rangle \\ &= \frac{1}{\Delta} \left(\tilde{\eta}^2 \sum_{i=1}^n \tilde{\lambda}_{i1} \langle \lambda_i, v_0 \rangle - \tilde{\eta}^1 \sum_{i=1}^n \tilde{\lambda}_{i2} \langle \lambda_i, v_0 \rangle \right) \\ &= 0, \end{aligned} \quad (86)$$

where momentum conservation $\sum_{i=1}^n \lambda_i \tilde{\lambda}_i = 0$ has been used. This finishes the proof.

D The mass dimensions of A_{CSW} and A_{Feynman}

We would like to show explicitly that both A_{CSW} and A_{Feynman} has mass dimension $4 - n$. We first show that the mass dimension of A_{CSW} is $4 - n$. For an n -gluon tree amplitude, assume $n_{\pm}$ being the number of external gluons with helicity ± 1 , n_p being the number of propagators and n_i being the number of MHV vertices with exactly i lines. From [8, 15], we get

$$n_- = \sum_i n_i + 1, \quad (87)$$

$$n_+ = \sum_i n_i(i - 3) + 1. \quad (88)$$

And it is easy to see that

$$n_p = \sum_i n_i - 1 = n_- - 2. \quad (89)$$

Notice that spinors $\lambda, \tilde{\lambda}$ of external gluon have dimension $[M]^{1/2}$. Then the m -gluon MHV amplitudes has mass dimension $4 - m$. Since A_{CSW} is invariant under $\tilde{\eta}^{\dot{\alpha}} \rightarrow t\tilde{\eta}^{\dot{\alpha}}$, we can assign $\tilde{\eta}^{\dot{\alpha}}$ arbitrary dimension. It is convenient to assign its dimension to be $[M]^{-1/2}$, such that λ for internal line has the same dimension as the λ for the external gluon. Then the MHV vertex with i lines has mass dimension $4 - i$, the same as the one of i -gluon MHV amplitudes. Also notice that every propagator has mass dimension -2 . So the mass dimension of the contribution of a MHV diagram to the amplitude is

$$\left(\sum_i (4 - i)n_i \right) - 2n_p \quad (90)$$

which can be shown to be $4 - n_+ - n_- = 4 - n$ by using (87)-(89).

The fact that A_{Feynman} has mass dimension $4 - n$ can be proved similarly. Denote the number of vertices with i lines by $\tilde{n}_i, i = 3, 4$ and the number of propagators by $\tilde{n}_p$. We know that

$$\tilde{n}_p = \tilde{n}_3 + \tilde{n}_4 - 1, \quad (91)$$

$$n = 3\tilde{n}_3 + 4\tilde{n}_4 - 2\tilde{n}_p. \quad (92)$$

From these two equations, we get

$$\tilde{n}_3 = n - 2\tilde{n}_4 - 2, \quad (93)$$

$$\tilde{n}_p = n - \tilde{n}_4 - 3. \quad (94)$$

Then the mass dimension of a Feynman diagram is¹⁰

$$\tilde{n}_3 - 2\tilde{n}_p = 4 - n. \quad (95)$$

¹⁰Notice that the polarization vectors in eq. (6) are dimensionless.

References

- [1] J. M. Henn and J. C. Plefka, *Scattering Amplitudes in Gauge Theories*, Lect. Notes Phys. **883** (2014), pp.1-195, doi:10.1007/978-3-642-54022-6
- [2] H. Elvang and Y. t. Huang, *Scattering Amplitudes in Gauge Theory and Gravity*, Cambridge University Press.
- [3] N. Arkani-Hamed, J. L. Bourjaily, F. Cachazo, A. B. Goncharov, A. Postnikov and J. Trnka, *Grassmannian Geometry of Scattering Amplitudes*, Cambridge University Press doi:10.1017/CBO9781316091548 [arXiv:1212.5605 [hep-th]].
- [4] E. Witten, *Perturbative gauge theory as a string theory in twistor space*, Commun. Math. Phys. **252** (2004), 189-258 doi:10.1007/s00220-004-1187-3 [arXiv:hep-th/0312171 [hep-th]].
- [5] R. Roiban, M. Spradlin and A. Volovich, *A Googly amplitude from the B model in twistor space*, JHEP **04** (2004), 012 doi:10.1088/1126-6708/2004/04/012 [arXiv:hep-th/0402016 [hep-th]].
- [6] R. Roiban and A. Volovich, *All conjugate-maximal-helicity-violating amplitudes from topological open string theory in twistor space*, Phys. Rev. Lett. **93** (2004), 131602 doi:10.1103/PhysRevLett.93.131602 [arXiv:hep-th/0402121 [hep-th]].
- [7] R. Roiban, M. Spradlin and A. Volovich, *On the tree level S matrix of Yang-Mills theory*, Phys. Rev. D **70** (2004), 026009 doi:10.1103/PhysRevD.70.026009 [arXiv:hep-th/0403190 [hep-th]].
- [8] F. Cachazo, P. Svrček and E. Witten, *MHV vertices and tree amplitudes in gauge theory*, JHEP **09** (2004), 006 doi:10.1088/1126-6708/2004/09/006 [arXiv:hep-th/0403047 [hep-th]].
- [9] S. Gukov, L. Motl and A. Neitzke, *Equivalence of twistor prescriptions for superYang-Mills*, Adv. Theor. Math. Phys. **11** (2007) no.2, 199-231 doi:10.4310/ATMP.2007.v11.n2.a1 [arXiv:hep-th/0404085 [hep-th]].
- [10] I. Bena, Z. Bern and D. A. Kosower, *Twistor-space recursive formulation of gauge-theory amplitudes*, Phys. Rev. D **71** (2005), 045008 doi:10.1103/PhysRevD.71.045008 [arXiv:hep-th/0406133 [hep-th]].
- [11] F. Cachazo and P. Svrček, *Lectures on twistor strings and perturbative Yang-Mills theory*, PoS **RTN2005** (2005), 004 doi:10.22323/1.019.0005 [arXiv:hep-th/0504194 [hep-th]].
- [12] A. Brandhuber, B. Spence and G. Travaglini, *Tree-Level Formalism*, J. Phys. A **44** (2011), 454002 doi:10.1088/1751-8113/44/45/454002 [arXiv:1103.3477 [hep-th]].

- [13] B. Feng and M. Luo, *An Introduction to On-shell Recursion Relations*, Front. Phys. (Beijing) **7** (2012), 533-575 doi:10.1007/s11467-012-0270-z [arXiv:1111.5759 [hep-th]].
- [14] R. Britto, F. Cachazo, B. Feng and E. Witten, *Direct proof of tree-level recursion relation in Yang-Mills theory*, Phys. Rev. Lett. **94** (2005), 181602 doi:10.1103/PhysRevLett.94.181602 [arXiv:hep-th/0501052 [hep-th]].
- [15] C. J. Zhu, *The Googly amplitudes in gauge theory*, JHEP **04** (2004), 032 doi:10.1088/1126-6708/2004/04/032 [arXiv:hep-th/0403115 [hep-th]].
- [16] J. B. Wu and C. J. Zhu, *MHV vertices and scattering amplitudes in gauge theory*, JHEP **07** (2004), 032 doi:10.1088/1126-6708/2004/07/032 [arXiv:hep-th/0406085 [hep-th]].
- [17] G. Georgiou and V. V. Khoze, *Tree amplitudes in gauge theory as scalar MHV diagrams*, JHEP **05** (2004), 070 doi:10.1088/1126-6708/2004/05/070 [arXiv:hep-th/0404072 [hep-th]].
- [18] J. B. Wu and C. J. Zhu, *MHV vertices and fermionic scattering amplitudes in gauge theory with quarks and gluinos*, JHEP **09** (2004), 063 doi:10.1088/1126-6708/2004/09/063 [arXiv:hep-th/0406146 [hep-th]].
- [19] K. Risager, *A Direct proof of the CSW rules*, JHEP **12** (2005), 003 doi:10.1088/1126-6708/2005/12/003 [arXiv:hep-th/0508206 [hep-th]].
- [20] H. Elvang, D. Z. Freedman and M. Kiermaier, *Proof of the MHV vertex expansion for all tree amplitudes in $N=4$ SYM theory*, JHEP **06** (2009), 068 doi:10.1088/1126-6708/2009/06/068 [arXiv:0811.3624 [hep-th]].
- [21] A. Gorsky and A. Rosly, *From Yang-Mills Lagrangian to MHV diagrams*, JHEP **01** (2006), 101 doi:10.1088/1126-6708/2006/01/101 [arXiv:hep-th/0510111 [hep-th]].
- [22] P. Mansfield, *The Lagrangian origin of MHV rules*, JHEP **03** (2006), 037 doi:10.1088/1126-6708/2006/03/037 [arXiv:hep-th/0511264 [hep-th]].
- [23] J. H. Eittle and T. R. Morris, *Structure of the MHV-rules Lagrangian*, JHEP **08** (2006), 003 doi:10.1088/1126-6708/2006/08/003 [arXiv:hep-th/0605121 [hep-th]].
- [24] H. Feng and Y. t. Huang, *MHV Lagrangian for $N=4$ super Yang-Mills*, JHEP **04** (2009), 047 doi:10.1088/1126-6708/2009/04/047 [arXiv:hep-th/0611164 [hep-th]].

- [25] J. H. Eftle, C. H. Fu, J. P. Fudger, P. R. W. Mansfield and T. R. Morris, *S-matrix equivalence theorem evasion and dimensional regularisation with the canonical MHV Lagrangian*, JHEP **05** (2007), 011 doi:10.1088/1126-6708/2007/05/011 [arXiv:hep-th/0703286 [hep-th]].
- [26] R. Boels, L. J. Mason and D. Skinner, *From twistor actions to MHV diagrams*, Phys. Lett. B **648** (2007), 90-96 doi:10.1016/j.physletb.2007.02.058 [arXiv:hep-th/0702035 [hep-th]].
- [27] Z. Bern, *String based perturbative methods for gauge theories*, [arXiv:hep-ph/9304249 [hep-ph]].
- [28] Z. Xu, D. H. Zhang and L. Chang, *Helicity Amplitudes for Multiple Bremsstrahlung in Massless Nonabelian Gauge Theories*, Nucl. Phys. B **291** (1987), 392-428 doi:10.1016/0550-3213(87)90479-2
- [29] F. A. Berends and W. Giele, *The Six Gluon Process as an Example of Weyl-Van Der Waerden Spinor Calculus*, Nucl. Phys. B **294** (1987), 700-732 doi:10.1016/0550-3213(87)90604-3
- [30] S. J. Parke and T. R. Taylor, *An Amplitude for n Gluon Scattering*, Phys. Rev. Lett. **56** (1986), 2459 doi:10.1103/PhysRevLett.56.2459
- [31] F. A. Berends and W. T. Giele, *Recursive Calculations for Processes with n Gluons*, Nucl. Phys. B **306** (1988), 759-808 doi:10.1016/0550-3213(88)90442-7
- [32] N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins and K. Risager, *MHV-vertices for gravity amplitudes*, JHEP **01** (2006), 009 doi:10.1088/1126-6708/2006/01/009 [arXiv:hep-th/0509016 [hep-th]].
- [33] S. Giombi, R. Ricci, D. Robles-Llana and D. Trancanelli, *A Note on twistor gravity amplitudes*, JHEP **07** (2004), 059 doi:10.1088/1126-6708/2004/07/059 [arXiv:hep-th/0405086 [hep-th]].
- [34] M. Bianchi, H. Elvang and D. Z. Freedman, *Generating Tree Amplitudes in $N=4$ SYM and $N=8$ SG*, JHEP **09** (2008), 063 doi:10.1088/1126-6708/2008/09/063 [arXiv:0805.0757 [hep-th]].
- [35] P. Benincasa, C. Boucher-Veronneau and F. Cachazo, JHEP **11** (2007), 057 doi:10.1088/1126-6708/2007/11/057 [arXiv:hep-th/0702032 [hep-th]].
- [36] E. Conde and S. Rajabi, *The Twelve-Graviton Next-to-MHV Amplitude from Risager's Construction*, JHEP **09** (2012), 120 doi:10.1007/JHEP09(2012)120 [arXiv:1205.3500 [hep-th]].
- [37] B. Penante, S. Rajabi and G. Sizov, *CSW-like Expansion for Einstein Gravity*, JHEP **05** (2013), 004 doi:10.1007/JHEP05(2013)004 [arXiv:1212.6257 [hep-th]].

- [38] K. J. Ozeren and W. J. Stirling, *MHV techniques for QED processes*, JHEP **11** (2005), 016 doi:10.1088/1126-6708/2005/11/016 [arXiv:hep-th/0509063 [hep-th]].
- [39] L. J. Dixon, E. W. N. Glover and V. V. Khoze, *MHV rules for Higgs plus multi-gluon amplitudes*, JHEP **12** (2004), 015 doi:10.1088/1126-6708/2004/12/015 [arXiv:hep-th/0411092 [hep-th]].
- [40] Z. Bern, D. Forde, D. A. Kosower and P. Mastrolia, *Twistor-inspired construction of electroweak vector boson currents*, Phys. Rev. D **72** (2005), 025006 doi:10.1103/PhysRevD.72.025006 [arXiv:hep-ph/0412167 [hep-ph]].
- [41] S. D. Badger, E. W. N. Glover and V. V. Khoze, *MHV rules for Higgs plus multi-parton amplitudes*, JHEP **03** (2005), 023 doi:10.1088/1126-6708/2005/03/023 [arXiv:hep-th/0412275 [hep-th]].
- [42] S. D. Badger, E. W. N. Glover, V. V. Khoze and P. Svrček, *Recursion relations for gauge theory amplitudes with massive particles*, JHEP **07** (2005), 025 doi:10.1088/1126-6708/2005/07/025 [arXiv:hep-th/0504159 [hep-th]].
- [43] D. Forde and D. A. Kosower, *All-multiplicity amplitudes with massive scalars*, Phys. Rev. D **73** (2006), 065007 doi:10.1103/PhysRevD.73.065007 [arXiv:hep-th/0507292 [hep-th]].
- [44] A. Brandhuber, B. J. Spence and G. Travaglini, *One-loop gauge theory amplitudes in $N=4$ super Yang-Mills from MHV vertices*, Nucl. Phys. B **706** (2005), 150-180 doi:10.1016/j.nuclphysb.2004.11.023 [arXiv:hep-th/0407214 [hep-th]].
- [45] J. Bedford, A. Brandhuber, B. J. Spence and G. Travaglini, *A Twistor approach to one-loop amplitudes in $N=1$ supersymmetric Yang-Mills theory*, Nucl. Phys. B **706** (2005), 100-126 doi:10.1016/j.nuclphysb.2004.11.031 [arXiv:hep-th/0410280 [hep-th]].
- [46] C. Quigley and M. Rozali, *One-loop MHV amplitudes in supersymmetric gauge theories*, JHEP **01** (2005), 053 doi:10.1088/1126-6708/2005/01/053 [arXiv:hep-th/0410278 [hep-th]].
- [47] M. x. Luo and C. k. Wen, *One-loop maximal helicity violating amplitudes in $N=4$ super Yang-Mills theories*, JHEP **11** (2004), 004 doi:10.1088/1126-6708/2004/11/004 [arXiv:hep-th/0410045 [hep-th]].
- [48] M. x. Luo and C. k. Wen, *Systematics of one-loop scattering amplitudes in $N=4$ super Yang-Mills theories*, Phys. Lett. B **609** (2005), 86-94 doi:10.1016/j.physletb.2005.01.041 [arXiv:hep-th/0410118 [hep-th]].

- [49] D. Binosi and L. Theussl, *JaxoDraw: A Graphical user interface for drawing Feynman diagrams*, Comput. Phys. Commun. **161** (2004), 76-86 doi:10.1016/j.cpc.2004.05.001 [arXiv:hep-ph/0309015 [hep-ph]].
- [50] D. Binosi, J. Collins, C. Kaufhold and L. Theussl, *JaxoDraw: A Graphical user interface for drawing Feynman diagrams. Version 2.0 release notes*, Comput. Phys. Commun. **180** (2009), 1709-1715 doi:10.1016/j.cpc.2009.02.020 [arXiv:0811.4113 [hep-ph]].