

THE NEHARI MANIFOLD APPROACH FOR SINGULAR EQUATIONS INVOLVING THE $p(x)$ -LAPLACE OPERATOR

DUŠAN D. REPOVŠ

Faculty of Education and Faculty of Mathematics and Physics, University of Ljubljana & Institute of Mathematics, Physics and Mechanics, 1000 Ljubljana, Slovenia

Email: dusan.repovs@guest.arnes.si

KAMEL SAOUDI

College of Sciences at Dammam, University of Dammam
31441 Dammam, Kingdom of Saudi Arabia

Email: kmsaoudi@iau.edu.sa

Abstract

We study the following singular problem involving the $p(x)$ -Laplace operator $\Delta_{p(x)}u = \operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$, where $p(x)$ is a nonconstant continuous function,

$$(P_\lambda) \begin{cases} -\Delta_{p(x)}u = a(x)|u|^{q(x)-2}u(x) + \frac{\lambda b(x)}{u^{\delta(x)}} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Here, Ω is a bounded domain in $\mathbb{R}^{N \geq 2}$ with C^2 -boundary, λ is a positive parameter, $a(x), b(x) \in C(\overline{\Omega})$ are positive weight functions with compact support in Ω , and $\delta(x), p(x), q(x) \in C(\overline{\Omega})$ satisfy certain hypotheses (A_0) and (A_1) . We apply the Nehari manifold approach and some new techniques to establish the multiplicity of positive solutions for problem (P_λ) .

Keywords: Nehari manifold, generalized Lebesgue-Sobolev space, topological method, singular equation, $p(x)$ -Laplace operator, multiplicity.

1. INTRODUCTION

The aim of this paper is to study the following inhomogeneous equation

$$(P_\lambda) \begin{cases} -\Delta_{p(x)}u = a(x)|u|^{q(x)-2}u(x) + \frac{\lambda b(x)}{u^{\delta(x)}} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Here, operator $\Delta_{p(x)}u := \operatorname{div}(|\nabla u|^{p(x)-2}\nabla u)$ is the $p(x)$ -Laplacian, $p(x)$ is a non-constant continuous function, Ω is a bounded domain in \mathbb{R}^N ($N \geq 2$) with C^2 -boundary, λ is a positive parameter, $a(x), b(x) \in C(\overline{\Omega})$ are positive weight functions with compact support in Ω , and $\delta(x), p(x), q(x) \in C(\overline{\Omega})$ satisfy the following conditions

Math. Subj. Classif. (2010): 35J20, 35J60, 35J70, 47J10.

$$(A_0) \quad 0 < 1 - \delta(x) < p(x) < q(x) < p^*(x);$$

$$(A_1) \quad 0 < 1 - \delta^- \leq 1 - \delta^+ < p^- \leq p^+ < q^- \leq q^+.$$

Here, $p^*(x) := Np(x)/(N - p(x))$, $\delta^+ := \text{ess sup } \delta(x)$, $\delta^- := \text{ess inf } \delta(x)$, and analogous definitions hold for p^- , p^+ , q^- , and q^+ .

Partial differential equations with variable exponents are a very interesting and active topics. The motivation for this type of problems was stimulated by their various applications in physics - for more details see ACERBI-MINGIONE [1], DIENING [4], and in particular the book RĂDULESCU-REPOVŠ [15]) and the references therein.

Before stating our main result, we review the key literature concerning singular partial differential equations with variable exponents. ZHANG [25] proved the existence of solutions for the purely singular problem. Using variational methods, SAOUDI [17] proved the existence for a superlinear singular equation with variable exponent. FAN [6] investigated the multiplicity of solutions using topological methods. In SAOUDI-GHANMI [20] and SAOUDI ET AL. [23] variational methods were used to establish the multiplicity of solutions for a singular problems with Dirichlet and Neumann conditions, respectively (see also SAOUDI-GHANMI [21]).

The case when p is constant in problem (P_λ) , has recieved more attention and has been approached by differents techniques. For a more general presentation we refer to COCLITE-PALMIERI [2], CRANDALL ET AL. [3], GHERGU-RĂDULESCU [9], GIACOMONI-SAOUDI [10], GIACOMONI ET AL. [11], SAOUDI [18], and SAOUDI-KRATOU [22] and the references therein.

Some interesting papers on the applications of the Nehari manifold method in variable exponent problem have recently been published, see e.g. MASHIYEV ET AL. [13], SAIEDINEZHAD-GHAEMI [16], and SAOUDI [19]. In the present paper, we generalize the results of GIACOMONI ET AL. [11] and SAOUDI [18] to the problem with variable exponent, by using topological methods. Here is the main result of this paper.

Theorem 1.1. *Suppose that condition (A_0) and (A_1) are fulfilled. Then there exists $\lambda_0 > 0$ such that for every $\lambda \in (0, \lambda_0)$, problem (P_λ) has at least two positive solutions.*

This paper is organized as follows. In Section 2, we briefly review the properties of generalized Lebesgue-Sobolev spaces. In Section 3, we prove the necessary lemmas. In Section 4, we prove the existence of a minimum for the functional energy E_λ in \mathcal{N}_λ^+ . In Section 5, we prove the existence of a minimum for the functional energy E_λ in \mathcal{N}_λ^- . Finally, In Section 6, we present the proof of our main result.

2. GENERALIZED LEBESGUE-SOBOLEV SPACES

In this section, we recall definitions of functional spaces with variable exponents and properties of the $p(x)$ -Laplacian operator which will be used later (for more on this topics see RĂDULESCU-REPOVŠ [15], and for other additional information see PAPAGEORGIOU ET AL. [14]). Let

$$L^{p(\cdot)}(\Omega) = \left\{ u \in S(\Omega) : \int_{\Omega} |u(x)|^{p(x)} dx < \infty \right\},$$

with the norm

$$|u|_{p(\cdot)} = |u|_{L^{p(\cdot)}(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \left| \frac{u(x)}{\lambda} \right|^{p(x)} dx \leq 1 \right\}.$$

Then $(L^{p(\cdot)}(\Omega), |\cdot|_{p(\cdot)})$ is a reflexive, uniform convex Banach, separable space - for details see FAN-ZHAO [8, Theorems 1.6, 1.10, 1.14] and RĂDULESCU-REPOVŠ [15].

The variable exponent Sobolev space

$$W^{1,p(\cdot)}(\Omega) = \{u \in L^{p(\cdot)}(\Omega) : |\nabla u| \in L^{p(\cdot)}(\Omega)\},$$

can be equipped with the norm

$$\|u\| = |u|_{p(\cdot)} + |\nabla u|_{p(\cdot)}, \quad \text{for all } u \in W^{1,p(\cdot)}(\Omega).$$

Note that $W_0^{1,p(\cdot)}(\Omega)$ is the closure of $C_0^\infty(\Omega)$ in $W^{1,p(\cdot)}(\Omega)$.

We denote by $L^{q(x)}(\Omega)$ the conjugate space of $L^{p(x)}(\Omega)$, where $\frac{1}{q(x)} + \frac{1}{p(x)} = 1$. For $u \in L^{p(x)}(\Omega)$ and $v \in L^{q(x)}(\Omega)$, the Hölder type inequality

$$\left| \int_{\Omega} u(x)v(x)dx \right| \leq \left(\frac{1}{p^-} + \frac{1}{q^-} \right) |u|_{p(x)} |v|_{q(x)}, \quad (2.1)$$

holds. Recall the following result.

Lemma 2.1. *Consider the mapping $\rho_{p(x)} : L^{p(x)}(\Omega) \rightarrow \mathbb{R}$ defined by*

$$\rho_{p(x)}(u) = \int_{\Omega} |u|^{p(x)} dx,$$

where $(u_n), u \in L^{p(x)}(\Omega)$, and $p^+ < \infty$. Then the following relations hold

$$\|u\|_{L^{p(x)}} > 1 \Rightarrow \|u\|_{L^{p(x)}}^{p^-} \leq \rho_{p(x)}(u) \leq \|u\|_{L^{p(x)}}^{p^+}, \quad (2.2)$$

$$\|u\|_{L^{p(x)}} < 1 \Rightarrow \|u\|_{L^{p(x)}}^{p^+} \leq \rho_{p(x)}(u) \leq \|u\|_{L^{p(x)}}^{p^-}, \quad (2.3)$$

$$\|u_n - u\|_{L^{p(x)}} \rightarrow 0 \text{ if and only if } \rho_{p(x)}(u_n - u) \rightarrow 0. \quad (2.4)$$

We state the Sobolev embedding theorem.

Theorem 2.1 (See FAN ET AL. [7] and KOVÁČIK-RÁKOSNIK [12]). *Let $p \in C(\bar{\Omega})$ with $p(x) > 1$ for each $x \in \bar{\Omega}$ where $\Omega \subset \mathbb{R}^N$ is an open bounded domain with Lipschitz boundary and suppose that $p(x) \leq r(x) \leq p^*(x)$ and $r \in C(\bar{\Omega})$, for all $x \in \bar{\Omega}$. Then the embedding $W^{1,p(x)}(\Omega) \hookrightarrow L^{r(x)}(\Omega)$ is continuous. Also, if $r(x) < p^*(x)$ almost everywhere in $\bar{\Omega}$, then this embedding is compact.*

Let $\rho(x, s)$ be a Carathéodory function satisfying the following condition

$$|\rho(x, s)| \leq A \quad \text{for a.e. } x \in \Omega \text{ and all } s \in [-s_0, s_0], \quad (2.5)$$

where $s_0 > 0$ and A is a constant. Recall the following comparison principle.

Lemma 2.2 (ZHANG [24, Lemma 2.3]). *Let $\rho(x, t)$ be a function satisfying (2.5) and increasing in t . Let $u, v \in W^{1,p(\cdot)}(\Omega)$ satisfy*

$$-\Delta_{p(x)} u + \rho(x, u) \leq -\Delta_{p(x)} v + \rho(x, v), \quad \text{for all } x \in \Omega.$$

and assume that $u \leq v$ on $\partial\Omega$. Then $u \leq v$ in Ω .

Next, we recall the following strong maximum principle.

Theorem 2.2 (SAOUDI-GHANMI [20, Theorem 3.2]). *Suppose that for some $0 < \alpha < 1$, $u, v \in C^{1,\alpha}(\overline{\Omega})$ we have $0 \not\leq u$, $0 \not\leq v$, and*

$$-\Delta_{p(x)}u - \frac{\lambda}{u^{\delta(x)}} = h(x) \geq g(x) = -\Delta_{p(x)}v - \frac{\lambda}{v^{\delta(x)}}, \quad (2.6)$$

with $u = v = 0$ on $\partial\Omega$, where $g, h \in L^\infty(\Omega)$ are such that $0 \leq g < h$ pointwise everywhere in Ω . Assume that

$$\frac{\partial u}{\partial \mathbf{n}} > 0 \quad \frac{\partial v}{\partial \mathbf{n}} > 0 \quad \text{on } \partial\Omega, \quad (2.7)$$

where \mathbf{n} is the inward unit normal on $\partial\Omega$. Then the following strong comparison principle holds:

$$u > v \quad \text{in } \Omega, \quad \text{and there is a positive } \epsilon \text{ such that } \frac{\partial(u-v)}{\partial \mathbf{n}} \geq \epsilon \quad \text{on } \partial\Omega. \quad (2.8)$$

We shall now prove the following result.

Theorem 2.3. *Suppose that the domain Ω has the cone property and consider $p \in C(\overline{\Omega})$. Assume that $b \in L^{\alpha(x)}$, $b(x) > 0$ for $x \in \Omega$, $\alpha \in C(\overline{\Omega})$ and $\alpha^- > 1$, $\alpha_0^- \leq \alpha_0(x) \leq \alpha_0^+$ ($\frac{1}{\alpha(x)} + \frac{1}{\alpha_0(x)} = 1$), $\delta \in C(\overline{\Omega})$, and*

$$0 < 1 - \delta(x) < \frac{\alpha(x) - 1}{\alpha(x)} p^*(x), \quad \text{for all } x \in \overline{\Omega}. \quad (2.9)$$

Then the embedding $W^{1,p(x)}(\Omega) \hookrightarrow L_{b(x)}^{1-\delta(x)}(\Omega)$ is compact. Moreover, there is a constant $c_2 > 0$ such that the following inequality holds

$$\int_{\Omega} b(x) |u|^{1-\delta(x)} dx \leq c_2 (||u||^{1-\delta^-} + ||u||^{1-\delta^+}). \quad (2.10)$$

Proof. The proof of the first assertion is adopted from FAN [6]. Let $u \in W^{1,p(x)}(\Omega)$ and let

$$r(x) = \frac{\alpha(x)}{\alpha(x) - 1} (1 - \delta(x)) = \alpha_0(x) (1 - \delta(x)).$$

Hence, (2.9) implies that $r(x) < p^*(x)$. Therefore, using Theorem (2.1), we obtain $W^{1,p(x)}(\Omega) \hookrightarrow L^{r(x)}(\Omega)$. So, for $u \in W^{1,p(x)}(\Omega)$, we get $|u|^{1-\delta(x)} \in L^{\alpha_0(x)}(\Omega)$. By (2.1),

$$\int_{\Omega} b(x) |u|^{1-\delta(x)} dx \leq c_1 |b|_{\alpha(x)} \left| |u|^{1-\delta(x)} \right| < \infty.$$

This means that $W^{1,p(x)}(\Omega) \subset L^{1-\delta(x)}(\Omega)$.

On the other hand, if $u_n \rightharpoonup 0$ weakly in $W^{1,p(x)}(\Omega)$, then we have that $u_n \rightarrow 0$ strongly in $L^{r(x)}(\Omega)$. Therefore,

$$\int_{\Omega} b(x) |u_n|^{1-\delta(x)} dx \leq c_1 |b|_{\alpha(x)} \left| |u_n|^{1-\delta(x)} \right| \rightarrow 0,$$

hence $|u_n|_{1-\delta(x), b(x)} \rightarrow 0$ and we can conclude that

$$W^{1,p(x)}(\Omega) \hookrightarrow L_{b(x)}^{1-\delta(x)}(\Omega).$$

Next, we shall prove inequality (2.10). First, we have from above

$$\int_{\Omega} b(x) |u|^{1-\delta(x)} dx \leq c_1 |b|_{\alpha(x)} \left| |u|^{1-\delta(x)} \right| < \infty.$$

Since $1 - \delta^- \leq 1 - \delta(x) \leq 1 - \delta^+$ and $|u|^{1-\delta(x)} \leq |u|^{1-\delta^-} + |u|^{1-\delta^+}$, we obtain

$$\int_{\Omega} b(x)|u|^{1-\delta(x)} dx \leq \int_{\Omega} b(x)|u|^{1-\delta^-} dx + \int_{\Omega} b(x)|u|^{1-\delta^+} dx.$$

On the other hand, using (2.1), (2.2), (2.3), (2.4), and condition $p(x) < (1 - \delta^-)\alpha_0(x) \leq (1 - \delta^+)\alpha_0(x) < p^*(x)$, we get

$$\int_{\Omega} b(x)|u|^{1-\delta^-} dx \leq c_2 |b|_{\alpha(x)} \left| |u|^{1-\delta(x)} \right|_{\alpha_0(x)} = c_2 |b|_{\alpha(x)} |u|_{(1-\delta^-)\alpha_0(x)}^{1-\delta^-} \leq c_3 \|u\|^{1-\delta^-}. \quad (2.11)$$

In the same way, one gets

$$\int_{\Omega} b(x)|u|^{1-\delta^+} dx \leq c_4 \|u\|^{1-\delta^+}. \quad (2.12)$$

Hence, using (2.11) and (2.12), we have

$$\int_{\Omega} b(x)|u|^{1-\delta(x)} dx \leq c_5 (\|u\|^{1-\delta^-} + \|u\|^{1-\delta^+}).$$

which completes the proof of Theorem 2.3. \square

Theorem 2.4. *Let $p \in C(\overline{\Omega})$, suppose the boundary of domain Ω has the cone property and let $u \in W^{1,p(x)}(\Omega)$. Then there exist nonnegative constants $c_6, c_7, c_8, c_9 > 0$ such that the following inequalities hold*

$$\begin{aligned} \int_{\Omega} a(x)|u|^{q(x)} dx &\leq \begin{cases} c_6 \|u\|^{q^+} & \text{if } \|u\| > 1, \\ c_7 \|u\|^{q^-} & \text{if } \|u\| < 1. \end{cases} \\ \int_{\Omega} b(x)|u|^{1-\delta(x)} dx &\leq \begin{cases} c_8 \|u\|^{1-\delta^-} & \text{if } \|u\| > 1, \\ c_9 \|u\|^{1-\delta^+} & \text{if } \|u\| < 1. \end{cases} \end{aligned}$$

Proof. Theorem 2.4 follows immediately by MASHIYEV ET AL. [13, Theorem 2.3] and Theorem 2.3. \square

3. SOME NECESSARY LEMMAS

Let us define the functional $E_{\lambda} : W_0^{1,p(x)}(\Omega) \rightarrow \mathbb{R}$ by

$$E_{\lambda}(u) \stackrel{\text{def}}{=} \int_{\Omega} \frac{|\nabla u|^{p(x)}}{p(x)} dx - \int_{\Omega} \frac{a(x)|u|^{q(x)}}{q(x)} dx - \lambda \int_{\Omega} \frac{b(x)(u^+)^{1-\delta(x)}}{1-\delta(x)} dx \quad (3.1)$$

Definition 3.1. *We say that $u \in W_0^{1,p(x)}(\Omega)$ is a generalized solution of the equation*

$$-\Delta_{p(x)} u = a(x)|u|^{q(x)-2}u(x) + \frac{\lambda b(x)}{u^{\delta(x)}} \quad (3.2)$$

if for all $\varphi \in C_0^\infty(\Omega)$ and $\text{ess inf}_K u > 0$ for every compact set $K \subset \Omega$,

$$\int_{\Omega} |\nabla u|^{p(x)-2} \nabla u \nabla \varphi dx = \int_{\Omega} a(x)|u|^{q(x)-1} \varphi dx + \lambda \int_{\Omega} b(x) u^{-\delta(x)} \varphi dx \quad (3.3)$$

for all $\varphi \in C_0^\infty(\Omega)$.

Obviously, every weak solution of problem (P_λ) is also a generalized solution of equation (3.2).

In many problems, such as (P_λ) , E_λ is not bounded below on $W_0^{1,p(x)}(\Omega)$, but it is bounded below on the corresponding Nehari manifold which is defined by

$$\mathcal{N}_\lambda := \{u \in W_0^{1,p(x)}(\Omega) \setminus \{0\} : \langle E'_\lambda(u), u \rangle = 0\}.$$

Then $u \in \mathcal{N}_\lambda$ if and only if

$$\int_\Omega \frac{|\nabla u|^{p(x)}}{p(x)} dx - \int_\Omega \frac{a(x)|u|^{q(x)}}{q(x)} dx - \lambda \int_\Omega \frac{b(x)|u|^{1-\delta(x)}}{1-\delta(x)} dx = 0. \quad (3.4)$$

We note that \mathcal{N}_λ contains every solution of problem (P_λ) .

It is well-known that the Nehari manifold is closely related to the behavior of the functions $\Phi_u : [0, \infty) \rightarrow \mathbb{R}$ defined as $\Phi_u(t) = E_\lambda(tu)$. Such maps are called fiber maps and were introduced by DRABEK-POHOZAEV [5]. For $u \in W_0^{1,p(x)}(\Omega) \setminus \{0\}$, we define

$$\begin{aligned} \Phi_u(t) &= \int_\Omega \frac{t^{p(x)}|\nabla u|^{p(x)}}{p(x)} dx - \int_\Omega \frac{a(x)t^{q(x)}}{q(x)}|u|^{q(x)} dx - \lambda \int_\Omega \frac{b(x)t^{1-\delta(x)}|u|^{1-\delta(x)}}{1-\delta(x)} dx, \\ \Phi'_u(t) &= \int_\Omega t^{p(x)-1}|\nabla u|^{p(x)} dx - \int_\Omega a(x)t^{q(x)-1}|u|^{q(x)} dx - \lambda \int_\Omega b(x)t^{-\delta(x)}|u|^{1-\delta(x)} dx, \\ \Phi''_u(t) &= \int_\Omega (p(x)-1)t^{p(x)-2}|\nabla u|^{p(x)} dx - \int_\Omega a(x)(q(x)-1)t^{q(x)-2}|u|^{q(x)} dx \\ &\quad + \lambda \int_\Omega b(x)\delta(x)t^{-\delta(x)-1}|u|^{1-\delta(x)} dx. \end{aligned}$$

It is easy to see that $tu \in \mathcal{N}_\lambda$ if and only if $\Phi'_u(t) = 0$ and in particular, $u \in \mathcal{N}_\lambda$ if and only if $\Phi'_u(1) = 0$. Thus it is natural to split \mathcal{N}_λ into three parts corresponding to local minima, local maxima and points of inflection defined as follows:

$$\begin{aligned} \mathcal{N}_\lambda^+ &:= \{u \in \mathcal{N}_\lambda : \Phi''_u(1) > 0\} = \left\{tu \in W_0^{1,p(x)}(\Omega) \setminus \{0\} : \Phi'_u(t) = 0, \Phi''_u(t) > 0\right\}, \\ \mathcal{N}_\lambda^- &:= \{u \in \mathcal{N}_\lambda : \Phi''_u(1) < 0\} = \left\{tu \in W_0^{1,p(x)}(\Omega) \setminus \{0\} : \Phi'_u(t) = 0, \Phi''_u(t) < 0\right\}, \\ \mathcal{N}_\lambda^0 &:= \{u \in \mathcal{N}_\lambda : \Phi''_u(1) = 0\} = \left\{tu \in W_0^{1,p(x)}(\Omega) \setminus \{0\} : \Phi'_u(t) = 0, \Phi''_u(t) = 0\right\}. \end{aligned}$$

Our first result is the following.

Lemma 3.1. *E_λ is coercive and bounded below on \mathcal{N}_λ .*

Proof. Let $u \in \mathcal{N}_\lambda$ and $\|u\| > 1$. Then, using (2.2)-(2.4) and the embeddings from Theorem 2.1, we estimate $E_\lambda(u)$ as follows:

$$\begin{aligned} E_\lambda(u) &= \int_\Omega \frac{|\nabla u|^{p(x)}}{p(x)} dx - \int_\Omega \frac{a(x)|u|^{q(x)}}{q(x)} dx - \lambda \int_\Omega \frac{b(x)|u|^{1-\delta(x)}}{1-\delta(x)} dx \\ &\geq \left(\frac{1}{p^+} - \frac{1}{q^-}\right) \int_\Omega |\nabla u|^{p(x)} dx - \lambda \left(\frac{1}{1-\delta^+} - \frac{1}{q^-}\right) \int_\Omega b(x)|u|^{1-\delta(x)} dx \\ &\geq \left(\frac{1}{p^+} - \frac{1}{q^-}\right) \|u\|^{p^-} - \lambda c_8 \left(\frac{1}{1-\delta^+} - \frac{1}{q^-}\right) \|u\|^{1-\delta^+}. \end{aligned}$$

Note that since $0 < \delta^+ < 1$ and $1 - \delta^+ < p^-$, it follows that $E_\lambda(u) \rightarrow \infty$ as $\|u\| \rightarrow \infty$. Therefore E_λ is coercive and bounded below. \square

Lemma 3.2. *Let u be a local minimizer for E_λ on subsets \mathcal{N}_λ^+ or \mathcal{N}_λ^- of \mathcal{N}_λ such that $u \notin \mathcal{N}_\lambda^0$. Then u is a critical point of E_λ .*

Proof. Recall u is a local minimizer for E_λ under the constraint

$$I_\lambda(u) := \langle E'_\lambda(u), u \rangle = 0. \quad (3.5)$$

Hence, using the theory of Lagrange multipliers, we obtain the existence of $\mu \in \mathbb{R}$ such that

$$E'_\lambda(u) = \mu I'_\lambda(u).$$

Therefore,

$$\langle E'_\lambda(u), u \rangle = \mu \langle I'_\lambda(u), u \rangle = \mu \Phi''_u(1) = 0.$$

So, $u \notin \mathcal{N}_\lambda^0$, hence $\Phi''_u(1) \neq 0$. Consequently, $\mu = 0$. The proof of Lemma 3.2 is thus complete. \square

Lemma 3.3. *There exists λ_0 such that for every $0 < \lambda < \lambda_0$, we have $\mathcal{N}_\lambda^\pm \neq \emptyset$ and $\mathcal{N}_\lambda^0 = \{0\}$.*

Proof. First, by Lemma 3.2, we deduce that \mathcal{N}_λ^\pm are nonempty for $\lambda \in (0, \lambda_0)$. Now, suppose that there exists $u \in \mathcal{N}_\lambda^0$ such that $\|u\| > 1$. Using the definition of \mathcal{N}_λ^0 , we obtain

$$\int_\Omega |\nabla u|^{p(x)} dx - \int_\Omega a(x)|u|^{q(x)} dx - \lambda \int_\Omega b(x)|u|^{1-\delta(x)} dx = 0.$$

Combining the above equality with (3.5) and Theorem 2.3 in [13], we get

$$\begin{aligned} 0 &= \langle I'_\lambda(u), u \rangle = \int_\Omega p(x)|\nabla u|^{p(x)} dx - \int_\Omega a(x)q(x)|u|^{q(x)} dx - \lambda \int_\Omega b(x)(1-\delta(x))|u|^{1-\delta(x)} dx \\ &\geq p^- \int_\Omega |\nabla u|^{p(x)} dx - q^+ \int_\Omega a(x)|u|^{q(x)} dx - (1-\delta^+) \left(\int_\Omega |\nabla u|^{p(x)} dx - \int_\Omega a(x)|u|^{q(x)} dx \right) \\ &\geq (p^- - (1-\delta^+)) \int_\Omega |\nabla u|^{p(x)} dx + (1-\delta^+ - q^+) \int_\Omega a(x)|u|^{q(x)} dx. \end{aligned}$$

It now follows from Theorem 2.4 that

$$(p^- - (1-\delta^+))\|u\|^{p^-} + c_{10}(1-\delta^+ - q^+)\|u\|^{q^+} \geq 0,$$

hence

$$\|u\| \geq c_{10} \left(\frac{p^- + \delta^+ - 1}{1 - \delta^+ - q^+} \right)^{\frac{1}{q^+ - p^-}}. \quad (3.6)$$

In the same way, since $u \in \mathcal{N}_\lambda$, we obtain

$$\int_\Omega |\nabla u|^{p(x)} dx - \int_\Omega a(x)|u|^{q(x)} dx - \lambda \int_\Omega b(x)|u|^{1-\delta(x)} dx = 0$$

and since $u \in \mathcal{N}_\lambda^0$, we have

$$p^+ \int_\Omega |\nabla u|^{p(x)} dx - q^- \int_\Omega a(x)|u|^{q(x)} dx - \lambda(1-\delta^+) \int_\Omega b(x)|u|^{1-\delta(x)} dx \geq 0.$$

Therefore

$$\begin{aligned} &p^+ \int_\Omega |\nabla u|^{p(x)} dx - q^- \int_\Omega a(x)|u|^{q(x)} dx \\ &- \lambda(1-\delta^+) \left(\int_\Omega |\nabla u|^{p(x)} dx - \lambda \int_\Omega a(x)|u|^{q(x)} dx \right) \geq 0. \end{aligned}$$

$$= (p^+ - q^+) \int_{\Omega} |\nabla u|^{p(x)} dx + \lambda(q^+ + \delta^+ - 1) \int_{\Omega} b(x) |u|^{1-\delta(x)} dx \geq 0.$$

Now since $\|u\| > 1$, by Theorem 2.4, one has

$$(p^+ - q^+) \|u\|^{p^-} + c_{11} \lambda (q^+ + \delta^+ - 1) \|u\|^{1-\delta^+} \geq 0,$$

and therefore

$$\|u\| \leq c_{11} \left(\lambda \frac{q^+ + \delta^+ - 1}{q^+ - p^-} \right)^{\frac{1}{p^- + \delta^+ - 1}}. \quad (3.7)$$

Using (3.6) and (3.7),

$$c_{11} \left(\lambda \frac{q^+ + \delta^+ - 1}{q^+ - p^-} \right)^{\frac{1}{p^- + \delta^+ - 1}} \geq c_{10} \left(\frac{p^- + \delta^+ - 1}{1 - \delta^+ - q^+} \right)^{\frac{1}{q^+ - p^-}}.$$

we get

$$\lambda \geq \frac{c_{10}}{c_{11}} \left(\frac{p^- + \delta^+ - 1}{1 - \delta^+ - q^+} \right)^{\frac{p^- + \delta^+ - 1}{q^+ - p^-}} \left(\frac{q^+ + \delta^+ - 1}{q^+ - p^-} \right).$$

Then, if λ is small enough,

$$\lambda = \frac{c_{10}}{c_{11}} \left(\frac{p^- + \delta^+ - 1}{1 - \delta^+ - q^+} \right)^{\frac{p^- + \delta^+ - 1}{q^+ - p^-}} \left(\frac{q^+ + \delta^+ - 1}{q^+ - p^-} \right),$$

we obtain $\|u\| < 1$, which is impossible. Therefore, $\mathcal{N}_{\lambda}^0 = \{0\}$ for all $\lambda \in (0, \lambda_0)$. Hence, this completes the proof of Lemma 3.3. \square

4. EXISTENCE OF MINIMIZERS ON \mathcal{N}_{λ}^+

In this section, we shall prove the existence of a minimum for the functional energy E_{λ} in \mathcal{N}_{λ}^+ . We shall also prove that this minimizer is a solution to problem (P_{λ}) .

Theorem 4.1. *There exists $u_{\lambda} \in \mathcal{N}_{\lambda}^+$ satisfying*

$$E_{\lambda}(u_{\lambda}) = \inf_{u \in \mathcal{N}_{\lambda}^+} E_{\lambda}(u),$$

for all $\lambda \in (0, \lambda_0)$.

Proof. Suppose that $\lambda \in (0, \lambda_0)$. Now, E_{λ} is bounded below on \mathcal{N}_{λ} and hence also on \mathcal{N}_{λ}^+ . Therefore there exists a sequence $\{u_n\} \subset \mathcal{N}_{\lambda}^+$, satisfying $E_{\lambda}(u_n) \rightarrow \inf_{u \in \mathcal{N}_{\lambda}^+} E_{\lambda}(u)$, as $n \rightarrow \infty$.

Since E_{λ} is coercive, $\{u_n\}$ is bounded in $W_0^{1,p(x)}(\Omega)$. Therefore we can assume that $u_n \rightharpoonup u_0$ weakly in $W_0^{1,p(x)}(\Omega)$ and by the compact embedding, we obtain

$$u_n \rightharpoonup u_0 \text{ in } L_{b(x)}^{1-\delta(x)}(\Omega)$$

and

$$u_n \rightharpoonup u_0 \text{ in } L_{a(x)}^{q(x)}(\Omega).$$

Now, we shall show that $u_n \rightarrow u_0$ strongly in $W_0^{1,p(x)}(\Omega)$. First, we shall prove that

$$\inf_{u \in \mathcal{N}_{\lambda}^+} E_{\lambda}(u) < 0.$$

Let $u_0 \in \mathcal{N}_\lambda^+$. Then $\phi''_{u_0}(1) > 0$ which gives

$$p^+ \int_{\Omega} |\nabla u|^{p(x)} dx - q^- \int_{\Omega} a(x)|u|^{q(x)} dx - \lambda(1 - \delta^+) \int_{\Omega} b(x)|u|^{1-\delta(x)} dx > 0. \quad (4.1)$$

Moreover, by the definition of the functional energy E_λ , we can write

$$E_\lambda(u) \leq \frac{1}{p^-} \int_{\Omega} |\nabla u|^{p(x)} dx - \frac{1}{q^+} \int_{\Omega} a(x)|u|^{q(x)} dx - \frac{\lambda}{1 - \delta^+} \int_{\Omega} b(x)|u|^{1-\delta(x)} dx. \quad (4.2)$$

Now, we multiply (3.5) by $-(1 - \delta^+)$ and get

$$-(1 - \delta^+) \int_{\Omega} |\nabla u|^{p(x)} dx + (1 - \delta^+) \int_{\Omega} a(x)|u|^{q(x)} dx + \lambda(1 - \delta^+) \int_{\Omega} b(x)|u|^{1-\delta(x)} dx = 0.$$

Invoking the above equality and (4.1), one gets

$$\int_{\Omega} a(x)|u|^{q(x)} dx < \frac{p^+ + \delta^+ - 1}{q^- + \delta^+ - 1} \int_{\Omega} |\nabla u|^{p(x)} dx. \quad (4.3)$$

On the other hand, from (3.5) and (4.2), we obtain

$$E_\lambda(u) \leq \left(\frac{1}{p^-} - \frac{1}{1 - \delta^+}\right) \int_{\Omega} |\nabla u|^{p(x)} dx - \left(\frac{1}{q^-} - \frac{1}{1 - \delta^+}\right) \int_{\Omega} a(x)|u|^{q(x)} dx. \quad (4.4)$$

Then, by (4.3) and (4.4), we get

$$E_\lambda(u) < -\frac{(p^- + \delta^+ - 1)(q^+ - p^-)}{p^- q^+ (1 - \delta^+)} \|u\|^{p^-} < 0.$$

Now, let us assume that $u_n \rightharpoonup u_0$ strongly in $W_0^{1,p(x)}(\Omega)$. Then

$$\int_{\Omega} |\nabla u_0|^{p(x)} dx \leq \liminf_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p(x)} dx.$$

Using the compactness of embeddings, we obtain

$$\begin{aligned} \int_{\Omega} a(x)u_0^{q(x)} dx &= \liminf_{n \rightarrow \infty} \int_{\Omega} b(x)u_n^{q(x)} dx, \\ \int_{\Omega} b(x)u_0^{1-\delta(x)} dx &= \liminf_{n \rightarrow \infty} \int_{\Omega} a(x)u_n^{1-\delta(x)} dx. \end{aligned}$$

Now, by (3.5) and Theorem 2.3 in [13], one has

$$E_\lambda(u_n) \geq \left(\frac{1}{p^-} - \frac{1}{q^+}\right) \int_{\Omega} |\nabla u_n|^{p(x)} dx + \lambda \left(\frac{1}{q^+} - \frac{1}{1 - \delta^+}\right) \int_{\Omega} b(x)|u_n|^{1-\delta(x)} dx.$$

Passing to the limit when n goes to ∞ , we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} E_\lambda(u_n) &\geq \left(\frac{1}{p^-} - \frac{1}{q^+}\right) \lim_{n \rightarrow \infty} \int_{\Omega} |\nabla u_n|^{p(x)} dx \\ &+ \lambda \left(\frac{1}{q^+} - \frac{1}{1 - \delta^+}\right) \lim_{n \rightarrow \infty} \int_{\Omega} b(x)|u_n|^{1-\delta(x)} dx. \end{aligned}$$

Hence, using Theorem 2.3 in [13], we get

$$\inf_{u \in \mathcal{N}^+} E_\lambda(u) > \left(\frac{1}{p^-} - \frac{1}{q^+}\right) \|u_0\|^{p^-} + \lambda c_5 \left(\frac{1}{q^+} - \frac{1}{1 - \delta^+}\right) (\|u_0\|^{1-\delta^-} + \|u_0\|^{1-\delta^+}) > 0$$

since $p^- > 1 - \delta^+ \geq 1 - \delta^-$ and $\|u_0\| > 1$, which gives a contradiction. Therefore, $u_n \rightarrow u_0$ strongly in $W_0^{1,p(x)}(\Omega)$ and $E_\lambda(u_0) = \inf_{u \in \mathcal{N}_\lambda^+} E_\lambda(u)$. This completes the proof of Theorem 4.1. \square

5. EXISTENCE OF MINIMIZERS ON \mathcal{N}_λ^-

In this section, we shall prove the existence of a minimum for the functional energy E_λ in \mathcal{N}_λ^- . We shall also prove that this minimizer is a solution to problem (P_λ) .

Theorem 5.1. *There exists $v_\lambda \in \mathcal{N}_\lambda^-$ such that*

$$E_\lambda(v_\lambda) = \inf_{v \in \mathcal{N}_\lambda^-} E_\lambda(v),$$

for all $\lambda \in (0, \lambda_0)$.

Proof. Suppose that $\lambda \in (0, \lambda_0)$. Since E_λ is bounded below on \mathcal{N}_λ hence also on \mathcal{N}_λ^- . Therefore, there exists a sequence $\{v_n\} \subset \mathcal{N}_\lambda^-$, satisfying $E_\lambda(v_n) \rightarrow \inf_{u \in \mathcal{N}_\lambda^-} E_\lambda(u)$, as $n \rightarrow \infty$. Since E_λ is coercive, $\{v_n\}$ is bounded in $W_0^{1,p(x)}(\Omega)$.

Therefore we can assume that $v_n \rightharpoonup v_0$ weakly in $W_0^{1,p(x)}(\Omega)$ and by the compact embedding, we get

$$v_n \rightharpoonup v_0 \text{ in } L_{b(x)}^{1-\delta(x)}(\Omega)$$

and

$$v_n \rightharpoonup v_0 \text{ in } L_{a(x)}^{q(x)}(\Omega).$$

Now, we shall show $v_n \rightarrow v_0$ strongly in $W_0^{1,p(x)}(\Omega)$. First, we shall prove that

$$\inf_{v \in \mathcal{N}_\lambda^-} E_\lambda(v) > 0.$$

Let $v_0 \in \mathcal{N}_\lambda^-$. Then we have from (3.5),

$$\int_\Omega |\nabla u|^{p(x)} dx - \int_\Omega a(x)|u|^{q(x)} dx - \lambda \int_\Omega b(x)|u|^{1-\delta(x)} dx = 0. \quad (5.1)$$

Moreover, by the definition of the functional energy E_λ , we can write

$$E_\lambda(v) \geq \frac{1}{p^-} \int_\Omega |\nabla v|^{p(x)} dx - \frac{1}{q^+} \int_\Omega a(x)|v|^{q(x)} dx - \frac{\lambda}{1-\delta^+} \int_\Omega b(x)|v|^{1-\delta(x)} dx. \quad (5.2)$$

Hence, from (5.1) and (5.2), one has

$$\begin{aligned} E_\lambda(v) &\geq \frac{1}{p^-} \int_\Omega |\nabla v|^{p(x)} dx - \frac{\lambda}{1-\delta^+} \int_\Omega b(x)|v|^{1-\delta(x)} dx \\ &\quad - \frac{1}{q^+} \left(\int_\Omega |\nabla v|^{p(x)} dx - \lambda \int_\Omega b(x)|v|^{1-\delta(x)} dx \right) \\ &\geq \left(\frac{1}{p^-} - \frac{1}{q^+} \right) \int_\Omega |\nabla v|^{p(x)} dx + \left(\frac{1}{q^+} - \frac{1}{1-\delta^+} \right) \int_\Omega b(x)|v|^{1-\delta(x)} dx \\ &\geq \left(\frac{1}{p^-} - \frac{1}{q^+} \right) \|v\|^{p^-} + \lambda c_8 \left(\frac{1}{q^+} - \frac{1}{1-\delta^+} \right) \|v\|^{1-\delta^+} \\ &\geq \left[\left(\frac{1}{p^-} - \frac{1}{q^+} \right) + \lambda c_8 \left(\frac{1}{q^+} - \frac{1}{1-\delta^+} \right) \right] \|v\|^{p^-} \end{aligned}$$

since $p^- > 1 - \delta^+$.

Hence, if we choose

$$\lambda < \frac{(1-\delta^+)(p^- - q^+)}{c_8 p^+(1-\delta^+ - q^+)},$$

we obtain $E_\lambda(v) > 0$. Moreover, since $\mathcal{N}_\lambda^+ \cap \mathcal{N}_\lambda^- = \emptyset$ and $\inf_{v \in \mathcal{N}_\lambda^+} E_\lambda(v) < 0$, we see that $v \in \mathcal{N}_\lambda^-$.

In the same way, if $v_0 \in \mathcal{N}_\lambda^-$, hence there exists t_0 satisfying $t_0 v_0 \in \mathcal{N}_\lambda^-$ and so $E_\lambda(t_0 v_0) \leq E_\lambda(v_0)$. Moreover, since

$$I'_\lambda(v) = \int_\Omega p(x) |\nabla v|^{p(x)} dx - \int_\Omega a(x) q(x) |v|^{q(x)} dx - \lambda \int_\Omega b(x) (1 - \delta(x)) |v|^{1-\delta(x)} dx,$$

we get

$$\begin{aligned} I'_\lambda(t_0 v_0) &= \int_\Omega p(x) |\nabla t_0 v_0|^{p(x)} dx - \int_\Omega a(x) q(x) |t_0 v_0|^{q(x)} dx - \lambda \int_\Omega b(x) (1 - \delta(x)) |t_0 v_0|^{1-\delta(x)} dx \\ &\leq t_0^{p^+} \int_\Omega |\nabla v_0|^{p(x)} dx - t_0^{q^-} \int_\Omega a(x) |v_0|^{q(x)} dx - \lambda (1 - \delta^+) t_0^{1-\delta^+} \int_\Omega b(x) |v_0|^{1-\delta(x)} dx, \end{aligned}$$

since $1 - \delta^+ < p^+ < q^-$. By the conditions on a and b it follows that $I'_\lambda(t_0 v_0) < 0$, so by definition of \mathcal{N}_λ^- $t_0 v_0 \in \mathcal{N}_\lambda^-$.

Now, let us assume that $v_n \rightharpoonup v_0$ strongly in $W_0^{1,p(x)}(\Omega)$. Using the fact that

$$\int_\Omega |\nabla v_0|^{p(x)} dx \leq \liminf_{n \rightarrow \infty} \int_\Omega |\nabla v_n|^{p(x)} dx$$

one gets

$$\begin{aligned} E_\lambda(t v_0) &\leq \int_\Omega \frac{t^{p(x)} |\nabla v_0|^{p(x)}}{p(x)} dx - \int_\Omega \frac{t^{q(x)} |v_0|^{q(x)}}{q(x)} dx - \lambda \int_\Omega \frac{t^{1-\delta(x)} |v_0|^{1-\delta(x)}}{1 - \delta(x)} dx, \\ &\leq \lim_{n \rightarrow \infty} \left[\frac{t^{p^+}}{p^+} \int_\Omega |\nabla v_n|^{p(x)} dx - \frac{t^{q^-}}{q^+} \int_\Omega |v_n|^{q(x)} dx - \lambda \frac{t^{1-\delta^+}}{1 - \delta^+} \int_\Omega |v_n|^{1-\delta(x)} dx \right], \\ &\leq \lim_{n \rightarrow \infty} E_\lambda(t v_n) \leq \lim_{n \rightarrow \infty} E_\lambda(v_n) = \inf_{v \in \mathcal{N}_\lambda^-} E_\lambda(v), \end{aligned}$$

which contradicts with the fact that $t v_0 \in \mathcal{N}_\lambda^-$. Hence, $v_n \rightarrow v_0$ strongly in $W_0^{1,p(x)}(\Omega)$ and $E_\lambda(v_0) = \inf_{v \in \mathcal{N}_\lambda^-} E_\lambda(v)$. This completes the proof of Theorem 5.1. \square

6. PROOF OF THEOREM 1.1

Proof. By Theorem 4.1 and Theorem 5.1, for all $\lambda \in (0, \lambda_0)$, there exist $u_0 \in \mathcal{N}_\lambda^+$, $v_0 \in \mathcal{N}_\lambda^-$ such that

$$E_\lambda(u_0) = \inf_{u \in \mathcal{N}_\lambda^+} E_\lambda(u)$$

and

$$E_\lambda(v_0) = \inf_{v \in \mathcal{N}_\lambda^-} E_\lambda(v).$$

On the other hand, since $E_\lambda(u_0) = E_\lambda(|u_0|)$ and $|u_0| \in \mathcal{N}_\lambda^+$ and in the same way, $E_\lambda(v_0) = E_\lambda(|v_0|)$ and $|v_0| \in \mathcal{N}_\lambda^-$, we suppose that $u_0, v_0 \geq 0$. Using Lemma 3.2, u_0, v_0 are critical points of E_λ on $W_0^{1,p(x)}(\Omega)$ and thus weak solutions of (P_λ) .

Finally, by the Harnack inequality and by ZHANG-LIU [26], we obtain that u_0, v_0 are nonnegative solutions of (P_λ) .

It remains to prove that the solutions we obtained for Theorem 4.1 and Theorem 5.1 are distinct. Indeed, since $\mathcal{N}_\lambda^- \cap \mathcal{N}_\lambda^+ = \emptyset$, it follows that u_0 and v_0 are different. This completes the proof of Theorem 1.1. \square

ACKNOWLEDGEMENTS

The first author was supported by the Slovenian Research Agency program P1-0292 and grants N1-0114 and N1-0083.

REFERENCES

- [1] E. ACERBI AND G. MINGIONE, *Regularity results for a class of functionals with non-standard growth*, Arch. Rational Mech. Anal. **156** (2001) 121–140.
- [2] M.M. COCLITE AND G. PALMIERI, *On a singular nonlinear Dirichlet problem*, Comm. Partial Differential Equations **14** (1989), 1315–1327.
- [3] M.G. CRANDALL, P.H. RABINOWITZ AND L. TARTAR, *On a Dirichlet problem with a singular nonlinearity*, Comm. Partial Differential Equations **2** (1977), 193–222.
- [4] L. DIENING, *Theoretical and numerical results for electrorheological fluids*, Ph.D. Thesis, University of Freiburg, Freiburg, 2002.
- [5] P. DRABEK AND S.I. POHOZAEV, *Positive solutions for the p -Laplacian: application of the fibering method*, Proc. Royal Soc. Edinburgh Sect A, **127** (1997), 703–726.
- [6] X. L. FAN, *Solutions for $p(x)$ -Laplacian Dirichlet problems with singular coefficients*, J. Math. Anal. Appl. **312** (2005), 464–477.
- [7] X.L. FAN, J.S. SHEN AND D. ZHAO, *Sobolev embedding Theorems for spaces $W^{k,p(x)}(\Omega)$* , J. Math. Anal. Appl. **262** (2001), 749–760.
- [8] X.L. FAN AND D. ZHAO, *On spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$* , J. Math. Anal. Appl. **263** (2001), 424–446.
- [9] M. GHERGU AND V. RĂDULESCU, *Singular Elliptic Problems: Bifurcation and Asymptotic Analysis*, Oxford Lecture Series in Mathematics and its Applications **37**. Oxford University Press, Oxford, 2008.
- [10] J. GIACOMONI AND K. SAOUDI, *Multiplicity of positive solutions for a singular and critical problem*, Nonlinear Anal. **71** (2009), no. 9, 4060–4077.
- [11] J. GIACOMONI, I. SCHINDLER AND P. TAKÁČ, *Sobolev versus Hölder local minimizers and global multiplicity for a singular and quasilinear equation*, Annali della Scuola Normale Superiore di Pisa, Classe di scienze Série V **6** (2007), 117–158.
- [12] O. KOVÁČIK AND J. RÁKOSNIK, *On spaces $L^{p(x)}$ and $W^{k,p(x)}$* , Czechoslovak Math. J. **41** (116) (1991), 592–618.
- [13] R.A. MASHIYEV, S. OGRAS, Z. YUCEDAG AND M. AVCI, *The Nehari manifold approach for Dirichlet problem involving the $p(x)$ -Laplacian equation*, J. Korean Math. Soc. **47** (2010), No. 1. pp. 1–16.
- [14] N.S. PAPAGEORGIOU, V.D. RĂDULESCU, D.D. REPOVŠ, *Nonlinear Analysis - Theory and Methods*, Springer Monographs in Mathematics, Springer, Cham, 2019.
- [15] V.D. RĂDULESCU AND D.D. REPOVŠ, *Partial Differential Equations with Variable Exponents: Variational Methods and Qualitative Analysis*, Chapman and Hall /CRC, Taylor & Francis Group, Boca Raton, FL, 2015.
- [16] S. SAEEDINEZHAD AND M.B. GHAEMI, *The fibering map approach to a quasilinear degenerate $p(x)$ -Laplacian equation*, Bull. Iranian Math. Soc., **41** (2015), 1477–1492.
- [17] K. SAOUDI, *Existence and non-existence of solution for a singular nonlinear Dirichlet problem involving the $p(x)$ -Laplace operator*, J. Adv. Math. Stud. **9** (2016), No. 2, 292–303.
- [18] K. SAOUDI, *Existence and non-existence for a singular problem with variables potentials*, Electronic Journal of Differential equations **2017** (2017), No. 291, 1–9.
- [19] SAOUDI, *A singular elliptic system involving the $p(x)$ -Laplacian and generalized Lebesgue–Sobolev spaces*, International Journal of Mathematics **30**(116):1950064 (2019).
- [20] K. SAOUDI AND A. GHANMI, *A multiplicity results for a singular problem involving the fractional p -Laplacian operator*, Complex Var. Elliptic Equ. **61**:9 (2016), 1199–1216.

- [21] K. SAOUDI AND A. GHANMI, *A multiplicity results for a singular equation involving the $p(x)$ -Laplace operator*, Complex Var. Elliptic Equ. **62** (2017) No. 5, 695-725.
- [22] K. SAOUDI AND M. KRATOU, *Existence of multiple solutions for a singular and quasi-linear equation*, Complex Var. Elliptic Equ. **60** (2015), no.7, 893-925.
- [23] K. SAOUDI, M. KRATOU AND S. AL SADHAN, *Multiplicity results for the $p(x)$ -Laplacian equation with singular nonlinearities and nonlinear Neumann boundary condition*, Intern. J. Differential Equ. **2016** (2016), Article ID 3149482, 14 pp.
- [24] Q.H. ZHANG, *A strong maximum principle for differential equations with nonstandard $p(x)$ -growth conditions*, J. Math. Anal. Appl. **312** (2005), No.1, 24-32.
- [25] Q. ZHANG, *Existence and asymptotic behavior of positive solutions to $p(x)$ -Laplacian equations with singular nonlinearities*, J. Inequal. Appl. 2007, Art. ID 19349, 9 pp.
- [26] X. ZHANG AND X. LIU, *The local boundedness and Harnack inequality of $p(x)$ - Laplace equation*, J. Math. Anal. Appl. **332** (2007), 209-218.