

HIGHER TOPOLOGICAL COMPLEXITY OF HYPERBOLIC GROUPS

SAM HUGHES AND KEVIN LI

ABSTRACT. We prove for non-elementary torsion-free hyperbolic groups Γ and all $r \geq 2$ that the higher topological complexity $\mathrm{TC}_r(\Gamma)$ is equal to $r \cdot \mathrm{cd}(\Gamma)$. In particular, hyperbolic groups satisfy the rationality conjecture on the TC -generating function, giving an affirmative answer to a question of Farber and Oprea. More generally, we consider certain toral relatively hyperbolic groups.

1. INTRODUCTION

Let $r \geq 2$ be an integer. The higher (or sequential) topological complexity $\mathrm{TC}_r(X)$ of a path-connected space X was introduced by Rudyak [Rud10], generalising Farber's topological complexity [Far03]. The motivation for these numerical invariants arises from robotics. They provide a measure of complexity for the motion planning problem in the configuration space X with prescribed initial and final states, as well as $r - 2$ consecutive intermediate states. More precisely, consider the path-fibration $p: X^{[0,1]} \rightarrow X^r$ that maps a path $\omega: [0,1] \rightarrow X$ to the tuple $(\omega(0), \omega(\frac{1}{r-1}), \dots, \omega(\frac{r-2}{r-1}), \omega(1))$. Then $\mathrm{TC}_r(X)$ is defined as the minimal integer n for which X^r can be covered by $n + 1$ many open subsets U_0, \dots, U_n such that p admits a local section over each U_i . If no such n exists, one sets $\mathrm{TC}_r(X) := \infty$. Note that $\mathrm{TC}_2(X)$ recovers the usual topological complexity.

Since the higher topological complexities are homotopy invariants, one obtains interesting invariants of groups Γ by setting $\mathrm{TC}_r(\Gamma) := \mathrm{TC}_r(K(\Gamma, 1))$, where $K(\Gamma, 1)$ is an Eilenberg–MacLane space. The topological complexities $\mathrm{TC}_r(\Gamma)$ have been computed for several classes of groups (see e.g. [Far+19; FM20; Dra20] for $r = 2$, [FO19; AGO20; GGY16] for $r \geq 2$, and references therein). In a celebrated result of Dranishnikov [Dra20] (see also [FM20]), the topological complexity $\mathrm{TC}_2(\Gamma)$ of groups with cyclic centralisers, such as hyperbolic groups, was shown to equal $\mathrm{cd}(\Gamma \times \Gamma)$. Here cd denotes the cohomological dimension. We generalise this result to all higher topological complexities $\mathrm{TC}_r(\Gamma)$ for $r \geq 2$, as well as to a larger class of groups containing certain toral relatively hyperbolic groups. Recall that a collection $\{P_i \mid i \in I\}$ of subgroups of Γ is called *malnormal*, if for all $g \in \Gamma$ and $i, j \in I$ we have $gP_i g^{-1} \cap P_j = \{e\}$, unless $i = j$ and $g \in P_i$.

Theorem A. *Let $r \geq 2$ and let Γ be a torsion-free group with $\mathrm{cd}(\Gamma) \geq 2$. Suppose that Γ admits a malnormal collection of abelian subgroups $\{P_i \mid i \in I\}$ satisfying $\mathrm{cd}(P_i^r) < \mathrm{cd}(\Gamma^r)$ such that the centraliser $C_\Gamma(g)$ is cyclic for every $g \in \Gamma$ that is not conjugate into any of the P_i . Then $\mathrm{TC}_r(\Gamma) = \mathrm{cd}(\Gamma^r)$.*

The preceding theorem was obtained for the case $r = 2$ by the second author in [Li21].

For a space X , the TC -generating function $f_X(t)$ is defined as the formal power series

$$f_X(t) := \sum_{r=1}^{\infty} \mathrm{TC}_{r+1}(X) \cdot t^r.$$

The TC -generating function of a group Γ is set to be $f_\Gamma(t) := f_{K(\Gamma, 1)}(t)$. Recall that a group Γ is said to be of *type F* (or *geometrically finite*) if it admits a finite model for $K(\Gamma, 1)$.

Date: 11th October, 2021.

2020 Mathematics Subject Classification. 55M30, 55R35, 20F67.

Following [FO19], we say that a finite CW-complex X (resp. a group Γ of type F) satisfies the *rationality conjecture* if the TC-generating function $f_X(t)$ (resp. $f_\Gamma(t)$) is a rational function of the form $\frac{P(t)}{(1-t)^2}$, where $P(t)$ is an integer polynomial with $P(1) = \text{cat}(X)$ (resp. $P(1) = \text{cd}(\Gamma)$). Here cat denotes the Lusternik–Schnirelmann category. While a counter-example to the rationality conjecture for finite CW-complexes was found in [FKS20], the rationality conjecture for groups of type F remains open. It is known to hold, e.g. for abelian groups of type F , right-angled Artin groups, fundamental groups of closed orientable surfaces, and Higman’s group (see [FO19, Section 8]). Our result extends the class of groups for which the rationality conjecture holds as follows.

Corollary B. *Let Γ be a group as in Theorem A. If Γ is of type F , then*

$$f_\Gamma(t) = \text{cd}(\Gamma) \frac{(2-t)t}{(1-t)^2}.$$

In particular, the rationality conjecture holds for Γ .

As remarked by Farber and Oprea in [FO19, page 159], it is particularly interesting to determine the validity of the rationality conjecture for the class of hyperbolic groups. We answer their question in the affirmative.

Corollary C. *The rationality conjecture holds for torsion-free hyperbolic groups.*

Acknowledgements. Both authors would like to thank the organisers of the workshop “Advances in Homotopy Theory” held in September 2021 at the Southampton Centre for Geometry, Topology, and Applications which inspired this work. The first author was supported by the Engineering and Physical Sciences Research Council grant number 2127970.

2. BACKGROUND

We recall the notion of a classifying space for a family of subgroups (see e.g. [Lüc05]). A *family* \mathcal{F} of subgroups of a group G is a non-empty set of subgroups that is closed under conjugation and finite intersections. The family consisting only of the trivial subgroup is denoted by \mathcal{TR} . The family $\mathcal{F}\langle\mathcal{H}\rangle$ generated by a set of subgroups \mathcal{H} is the smallest family containing \mathcal{H} . For a family \mathcal{F} of subgroups of G and a subgroup H of G , we denote by $\mathcal{F}|_H$ the family $\{L \cap H \mid L \in \mathcal{F}\}$ of subgroups of H . A *classifying space* $E_{\mathcal{F}}G$ for the family \mathcal{F} is a terminal object in the G -homotopy category of G -CW-complexes with isotropy groups in \mathcal{F} . Note that a model for $E_{\mathcal{TR}}G$ is given by EG and in particular, there exists a G -map $EG \rightarrow E_{\mathcal{F}}G$ that is unique up to G -homotopy.

Let $\mathcal{E} \subset \mathcal{F}$ be two families of subgroups of G . We say that G satisfies condition $(M_{\mathcal{E} \subset \mathcal{F}})$ if every element in $\mathcal{F} \setminus \mathcal{E}$ is contained in a unique maximal element $M \in \mathcal{F} \setminus \mathcal{E}$, and that G satisfies condition $(NM)_{\mathcal{E} \subset \mathcal{F}}$ if additionally M equals its normaliser $N_G(M)$. We denote the Weyl group by $W_G(M) := N_G(M)/M$. The following proposition is a special case of a construction due to Lück and Weiermann [LW12, Corollary 2.8] stated in [Li21, Corollary 2.2].

Theorem 2.1 (Lück–Weiermann). *Let G be a group and $\mathcal{E} \subset \mathcal{F}$ be two families of subgroups. Let $\{M_i \mid i \in I\}$ be a complete set of representatives for the conjugacy classes of maximal elements in $\mathcal{F} \setminus \mathcal{E}$.*

- (i) *If $\mathcal{E} = \mathcal{TR}$ and G satisfies condition $(M_{\mathcal{TR} \subset \mathcal{F}})$, then a model for $E_{\mathcal{F}}G$ is given by the following G -pushout*

$$\begin{array}{ccc} \coprod_{i \in I} G \times_{N_G(M_i)} E(N_G(M_i)) & \longrightarrow & EG \\ \downarrow & & \downarrow \\ \coprod_{i \in I} G \times_{N_G(M_i)} E(W_G(M_i)) & \longrightarrow & E_{\mathcal{F}}G; \end{array}$$

- (ii) If G satisfies conditions $(M_{\mathcal{E} \subset \mathcal{F}})$ and $(NM_{\mathcal{E} \subset \mathcal{F}})$, then a model for $E_{\mathcal{F}}G$ is given by the following G -pushout

$$\begin{array}{ccc} \coprod_{i \in I} G \times_{M_i} E_{\mathcal{E}|_{M_i}} M_i & \longrightarrow & E_{\mathcal{E}}G \\ \downarrow & & \downarrow \\ \coprod_{i \in I} G/M_i & \longrightarrow & E_{\mathcal{F}}G. \end{array}$$

The \mathcal{F} -restricted orbit category $\mathcal{O}_{\mathcal{F}}G$ has G -sets G/H with $H \in \mathcal{F}$ as objects and G -maps as morphisms. Let A be an $\mathcal{O}_{\mathcal{F}}G$ -module, that is a contravariant functor from the orbit category $\mathcal{O}_{\mathcal{F}}G$ to the category of modules. The equivariant cellular cohomology $H_G^*(X; A)$ of a G -CW-complex X with isotropy groups in \mathcal{F} is called *Bredon cohomology* [Bre67]. In particular, Bredon cohomology satisfies the Mayer–Vietoris axiom for G -pushouts.

The (higher) topological complexity $\mathrm{TC}_r(\Gamma)$ of a group Γ for $r \geq 2$ can be characterised in terms of classifying spaces for families [FO19, Theorem 3.1], generalising a result of Farber–Grant–Lupton–Oprea [Far+19, Theorem 3.3] for the case $r = 2$. Consider $G = \Gamma^r$ and let \mathcal{D} be the family of subgroups that is generated by the diagonal subgroup $\Delta(\Gamma) \subset \Gamma^r$.

Theorem 2.2 (Farber–Oprea). *Let Γ be a group and $r \geq 2$. Then $\mathrm{TC}_r(\Gamma)$ equals the infimum of integers n for which the canonical Γ^r -map*

$$E(\Gamma^r) \rightarrow E_{\mathcal{D}}(\Gamma^r)$$

is Γ^r -equivariantly homotopic to a Γ^r -map with values in the n -skeleton $E_{\mathcal{D}}(\Gamma^r)^{(n)}$.

As a consequence [FO19, Theorem 5.1], a lower bound for $\mathrm{TC}_r(\Gamma)$ is given by the supremum of integers n for which the canonical Γ^r -map $E(\Gamma^r) \rightarrow E_{\mathcal{D}}(\Gamma^r)$ induces a non-trivial map in Bredon cohomology

$$H_{\Gamma^r}^n(E_{\mathcal{D}}(\Gamma^r); A) \rightarrow H_{\Gamma^r}^n(E(\Gamma^r); A)$$

for some $\mathcal{O}_{\mathcal{D}}(\Gamma^r)$ -module A .

3. PROOFS

Fix an integer $r \geq 2$. Let Γ be a group and $\Delta: \Gamma \rightarrow \Gamma^r$ be the diagonal map. For $\gamma = (\gamma_1, \dots, \gamma_{r-1}) \in \Gamma^{r-1}$ and a subset $S \subset \Gamma$, we define the subgroup $H_{\gamma, S}$ of Γ^r as

$$H_{\gamma, S} := (\gamma_1, \dots, \gamma_{r-1}, e) \cdot \Delta(C_{\Gamma}(S)) \cdot (\gamma_1^{-1}, \dots, \gamma_{r-1}^{-1}, e).$$

Here $C_{\Gamma}(S)$ denotes the centraliser of S in Γ . For $b \in \Gamma$, we write $H_{\gamma, b}$ instead of $H_{\gamma, \{b\}}$. Denote the element $\underline{e} := (e, \dots, e) \in \Gamma^{r-1}$ and note that $H_{\underline{e}, e} = \Delta(\Gamma)$. The elementary proof of the following lemma is omitted.

Lemma 3.1. *Let $\gamma = (\gamma_1, \dots, \gamma_{r-1}), \delta = (\delta_1, \dots, \delta_{r-1}) \in \Gamma^{r-1}$ and $S, T \subset \Gamma$ be subsets. The following hold:*

- (i) *For every $g = (g_1, \dots, g_r) \in \Gamma^r$ we have*

$$g \cdot H_{\gamma, S} \cdot g^{-1} = H_{\gamma', S'},$$

where $\gamma' = (g_1 \gamma_1 g_r^{-1}, \dots, g_{r-1} \gamma_{r-1} g_r^{-1}) \in \Gamma^{r-1}$ and $S' = \{g_r s g_r^{-1} \in \Gamma \mid s \in S\} \subset \Gamma$;

- (ii) *$H_{\gamma, S} \cap H_{\delta, T} = H_{\gamma, S \cup T \cup \{\delta_1^{-1} \gamma_1, \dots, \delta_{r-1}^{-1} \gamma_{r-1}\}}$;*

- (iii) *$N_{\Gamma^r}(H_{\gamma, S}) = \{(\gamma_1 k_1 h \gamma_1^{-1}, \dots, \gamma_{r-1} k_{r-1} h \gamma_{r-1}^{-1}, h) \in \Gamma^r \mid h \in N_{\Gamma}(C_{\Gamma}(S)), k_1, \dots, k_{r-1} \in C_{\Gamma}(C_{\Gamma}(S))\}$.*

Let $\mathcal{F}_1 \subset \mathcal{D}$ be the families of subgroups of Γ^r defined as

$$\begin{aligned}\mathcal{D} &:= \mathcal{F}\langle\{\Delta(\Gamma)\}\rangle; \\ \mathcal{F}_1 &:= \mathcal{F}\langle\{H_{\gamma,b} \mid \gamma \in \Gamma^{r-1}, b \in \Gamma \setminus \{e\}\}\rangle.\end{aligned}$$

Lemma 3.2. *Let Γ be a group as in Theorem A and let $\underline{e} = (e, \dots, e) \in \Gamma^{r-1}$. Then for $n = \text{cd}(\Gamma^r)$ and every $\mathcal{O}_{\mathcal{D}}(\Gamma^r)$ -module A , we have*

$$H_{\Gamma^r}^n(\Gamma^r \times_{H_{\underline{e},e}} E_{\mathcal{F}_1|_{H_{\underline{e},e}}} (H_{\underline{e},e}); A) = 0.$$

Proof. We have that conditions $(M_{\mathcal{TR} \subset \mathcal{F}_1|_{H_{\underline{e},e}}})$ and $(NM_{\mathcal{TR} \subset \mathcal{F}_1|_{H_{\underline{e},e}}})$ hold for the group $H_{\underline{e},e}$. To see this we follow the same argument as in [Li21, Lemma 3.5], using Lemma 3.1 instead of [Li21, Lemma 3.1]. By Theorem 2.1(ii), we obtain an $H_{\underline{e},e}$ -pushout

$$(1) \quad \begin{array}{ccc} \coprod_{H_{\underline{e},b} \in \mathcal{M}} H_{\underline{e},e} \times_{H_{\underline{e},b}} E(H_{\underline{e},b}) & \longrightarrow & E(H_{\underline{e},e}) \\ \downarrow & & \downarrow \\ \coprod_{H_{\underline{e},b} \in \mathcal{M}} H_{\underline{e},e}/H_{\underline{e},b} & \longrightarrow & E_{\mathcal{F}_1|_{H_{\underline{e},e}}} (H_{\underline{e},e}), \end{array}$$

where \mathcal{M} is a complete set of representatives of conjugacy classes of maximal elements in $\mathcal{F}_1|_{H_{\underline{e},e}} \setminus \mathcal{TR}$. Since $\text{cd}(H_{\underline{e},e}) < n$ and $\text{cd}(H_{\underline{e},b}) < n - 1$ for $b \in \Gamma \setminus \{e\}$, the Mayer–Vietoris sequence for $H_{H_{\underline{e},e}}}^*(-; A)$ applied to the pushout (1) yields the lemma. \square

Proof of Theorem A. Throughout the proof let $n = \text{cd}(\Gamma^r)$. We show that the Γ^r -map $E(\Gamma^r) \rightarrow E_{\mathcal{D}}(\Gamma^r)$ induces a surjective map $H_{\Gamma^r}^n(E_{\mathcal{D}}(\Gamma^r); A) \rightarrow H_{\Gamma^r}^n(E(\Gamma^r); A)$ for every $\mathcal{O}_{\mathcal{D}}(\Gamma^r)$ -module A . Then the theorem follows from Theorem 2.2 and the upper bound $\text{TC}_r(\Gamma) \leq n$ (see e.g. [FO19, (4)]).

First, observe that condition $(M_{\mathcal{TR} \subset \mathcal{F}_1})$ holds and that moreover, for $\gamma \in \Gamma^{r-1}$ and $b \in \Gamma \setminus \{e\}$ there is an isomorphism $N_{\Gamma^r}(H_{\gamma,b}) \cong C_{\Gamma}(b)^r$. To see this we follow the same argument as in [Li21, Lemma 3.5], using Lemma 3.1 instead of [Li21, Lemma 3.1]. It follows that for every $\mathcal{O}_{\mathcal{D}}(\Gamma^r)$ -module A , we have

$$H_{\Gamma^r}^n(\Gamma^r \times_{N_{\Gamma^r}(H_{\gamma,b})} E(N_{\Gamma^r}(H_{\gamma,b})); A) = 0.$$

By Theorem 2.1(i), we obtain a Γ^r -pushout

$$(2) \quad \begin{array}{ccc} \coprod_{H_{\gamma,b} \in \mathcal{M}} \Gamma^r \times_{N_{\Gamma^r}(H_{\gamma,b})} E(N_{\Gamma^r}(H_{\gamma,b})) & \longrightarrow & E(\Gamma^r) \\ \downarrow & & \downarrow \\ \coprod_{H_{\gamma,b} \in \mathcal{M}} \Gamma^r \times_{N_{\Gamma^r}(H_{\gamma,b})} E(W_{\Gamma^r}(H_{\gamma,b})) & \longrightarrow & E_{\mathcal{F}_1}(\Gamma^r), \end{array}$$

where \mathcal{M} is a complete set of representatives of conjugacy classes of maximal elements in $\mathcal{F}_1 \setminus \mathcal{TR}$. Applying the Mayer–Vietoris sequence for $H_{\Gamma^r}^*(-; A)$ to the pushout (2) shows that the induced map $H_{\Gamma^r}^n(E_{\mathcal{F}_1}(\Gamma^r); A) \rightarrow H_{\Gamma^r}^n(E(\Gamma^r); A)$ is surjective.

Second, conditions $(M_{\mathcal{F}_1 \subset \mathcal{D}})$ and $(NM_{\mathcal{F}_1 \subset \mathcal{D}})$ hold by the same argument as in [Li21, Lemma 3.2], using Lemma 3.1 instead of [Li21, Lemma 3.1]. By Theorem 2.1(ii), we obtain a Γ^r -pushout

$$(3) \quad \begin{array}{ccc} \Gamma^r \times_{H_{\underline{e},e}} E_{\mathcal{F}_1|_{H_{\underline{e},e}}} (H_{\underline{e},e}) & \longrightarrow & E_{\mathcal{F}_1}(\Gamma^r) \\ \downarrow & & \downarrow \\ \Gamma^r/H_{\underline{e},e} & \longrightarrow & E_{\mathcal{D}}(\Gamma^r). \end{array}$$

Lemma 3.2 and the Mayer–Vietoris sequence for $H_{\Gamma^r}^*(-; A)$ applied to the pushout (3) yield that the map $H_{\Gamma^r}^n(E_{\mathcal{D}}(\Gamma^r); A) \rightarrow H_{\Gamma^r}^n(E_{\mathcal{F}_1}(\Gamma^r); A)$ is surjective.

Together, the map $H_{\Gamma^r}^n(E_{\mathcal{D}}(\Gamma^r); A) \rightarrow H_{\Gamma^r}^n(E(\Gamma^r); A)$ is surjective for every $\mathcal{O}_{\mathcal{D}}(\Gamma^r)$ -module A . This finishes the proof. \square

Proof of Corollary B. For groups Γ of type F , we have $\text{cd}(\Gamma^r) = r \cdot \text{cd}(\Gamma)$ by [Dra19, Corollary 2.5]. The result now follows from Theorem A. \square

Proof of Corollary C. The result follows from Corollary B using the fact that torsion-free hyperbolic groups are of type F (see e.g. [BH99, Corollary 3.26 on page 470]). \square

REFERENCES

- [AGO20] Jorge Aguilar-Guzmán, Jesús González, and John Oprea. *Right-angled Artin groups, polyhedral products and the TC-generating function*. 2020. arXiv: [2011.04742](https://arxiv.org/abs/2011.04742) [math.AT].
- [Bre67] Glen E. Bredon. *Equivariant cohomology theories*. Lecture Notes in Mathematics, No. 34. Springer-Verlag, Berlin-New York, 1967.
- [BH99] Martin R. Bridson and André Haefliger. *Metric spaces of non-positive curvature*. Vol. 319. Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin, 1999.
- [Dra19] Alexander Dranishnikov. “On dimension of product of groups”. In: *Algebra Discrete Math.* 28.2 (2019), pp. 203–212.
- [Dra20] Alexander Dranishnikov. “On topological complexity of hyperbolic groups”. In: *Proc. Amer. Math. Soc.* 148.10 (2020), pp. 4547–4556. DOI: [10.1090/proc/15094](https://doi.org/10.1090/proc/15094).
- [Far03] Michael Farber. “Topological complexity of motion planning”. In: *Discrete Comput. Geom.* 29.2 (2003), pp. 211–221. DOI: [10.1007/s00454-002-0760-9](https://doi.org/10.1007/s00454-002-0760-9).
- [FKS20] Michael Farber, Daisuke Kishimoto, and Donald Stanley. “Generating functions and topological complexity”. In: *Topology Appl.* 278 (2020), pp. 107235, 5. DOI: [10.1016/j.topol.2020.107235](https://doi.org/10.1016/j.topol.2020.107235).
- [FM20] Michael Farber and Stephan Mescher. “On the topological complexity of aspherical spaces”. In: *J. Topol. Anal.* 12.2 (2020), pp. 293–319. DOI: [10.1142/S1793525319500511](https://doi.org/10.1142/S1793525319500511).
- [FO19] Michael Farber and John Oprea. “Higher topological complexity of aspherical spaces”. In: *Topology Appl.* 258 (2019), pp. 142–160. DOI: [10.1016/j.topol.2019.02.055](https://doi.org/10.1016/j.topol.2019.02.055).
- [Far+19] Michael Farber et al. “Bredon cohomology and robot motion planning”. In: *Algebr. Geom. Topol.* 19.4 (2019), pp. 2023–2059. DOI: [10.2140/agt.2019.19.2023](https://doi.org/10.2140/agt.2019.19.2023).
- [GGY16] Jesús González, Bárbara Gutiérrez, and Sergey Yuzvinsky. “Higher topological complexity of subcomplexes of products of spheres and related polyhedral product spaces”. In: *Topol. Methods Nonlinear Anal.* 48.2 (2016), pp. 419–451. DOI: [10.12775/tmna.2016.051](https://doi.org/10.12775/tmna.2016.051).
- [Li21] Kevin Li. “On the topological complexity of toral relatively hyperbolic groups”. To appear in: *Proc. Amer. Math. Soc.* (2021). arXiv: [2102.13511](https://arxiv.org/abs/2102.13511) [math.AT].
- [Lüc05] Wolfgang Lück. “Survey on classifying spaces for families of subgroups”. In: *Infinite groups: geometric, combinatorial and dynamical aspects*. Vol. 248. Progr. Math. Birkhäuser, Basel, 2005, pp. 269–322. DOI: [10.1007/3-7643-7447-0_7](https://doi.org/10.1007/3-7643-7447-0_7).
- [LW12] Wolfgang Lück and Michael Weiermann. “On the classifying space of the family of virtually cyclic subgroups”. In: *Pure Appl. Math. Q.* 8.2 (2012), pp. 497–555. DOI: [10.4310/PAMQ.2012.v8.n2.a6](https://doi.org/10.4310/PAMQ.2012.v8.n2.a6).
- [Rud10] Yuli B. Rudyak. “On higher analogs of topological complexity”. In: *Topology Appl.* 157.5 (2010), pp. 916–920. DOI: [10.1016/j.topol.2009.12.007](https://doi.org/10.1016/j.topol.2009.12.007).

(S. Hughes) MATHEMATICAL INSTITUTE, UNIVERSITY OF OXFORD, OXFORD OX2 6GG, UK

Email address: sam.hughes@math.ox.ac.uk

(K. Li) SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SOUTHAMPTON, SOUTHAMPTON SO17 1BJ, UK

Email address: kevin.li@soton.ac.uk