

Tighter Monogamy Relations in Multi-Qubit Systems

Yudie Gu¹, Yanmin Yang¹*, Jialing Zhang¹, Wei Chen²

¹ Faculty of Science, Kunming University of Science and Technology,
Kunming, 650500, China

² School of Computer Science and Technology, Dongguan University of Technology,
Dongguan, 523808, China

October 25, 2022

Abstract

In this paper, we present some monogamy relations of multiqubit quantum entanglement in terms of the β th power of concurrence, entanglement of formation and convex-roof extended negativity. These monogamy relations are proved to be tighter than the existing ones, together with detailed examples showing the tightness.

Keywords: Monogamy relations; Concurrence; Entanglement of formation; Convex-roof extended negativity

1 Introduction

Quantum entanglement is widely used as a very important resource in quantum information processing [1, 2, 3, 4]. With the emergence of quantum information theory, quantum entanglement plays a very important role in quantum cryptography, quantum teleportation and measurement based quantum computing. An important issue related to the entanglement metric is the limited shareability of the two-part entanglement in a multipartite entangled qubit system, that is, the single duality of entanglement [5]. Monogamy of entanglement (MoE) plays a very important role in many quantum information and communication processing tasks, such as security proof of quantum cryptography schemes and security analysis of quantum key distribution [6, 7].

For a tripartite quantum state ρ_{ABC} , MoE can be described as $E(\rho_{A|BC}) \geq E(\rho_{AB}) + E(\rho_{AC})$, where $\rho_{AB} = \text{tr}_C(\rho_{ABC})$, $\rho_{AC} = \text{tr}_B(\rho_{ABC})$, $E(\rho_{A|BC})$ denotes the entanglement between systems A and BC. A remarkable result was established by Coffman, Kundu and Wootters (CKW) [8] for three qubits, that was the simultaneous squares satisfy monogamy inequality. Then, the so-called CKW inequality was generalized to any N -qubit system [9]. Interestingly, it is further proved that similar inequalities of polyqubit monogamy can be established for negativity and convex-roof extended negativity (CREN) [10, 11, 12], the entanglement of formation (EoF) [13, 14], Rényi- α entanglement [15, 16] and Tsallis- q entanglement [17].

Our paper is organized as follows. In Sec.2, we present and prove two monogamy inequalities for the β th ($\beta \geq 2$) power of concurrence in N -qubit system. In Sec.3, we give a tighter monogamy relation for the β th ($\beta \geq \sqrt{2}$) power of EoF in $2 \otimes 2 \otimes 2^{N-2}$

*e-mail: ym.yang@kust.edu.cn

system. Then we extend the result to N -qubit system. In Sec.4, the monogamy relation for the β th ($\beta \geq 2$) power of CREN in N -qubit system is discussed. In addition, detailed examples are given to illustrate the tightness. In Sec.5, we summarize our results.

2 Tighter Monogamy Relations using Concurrence

Given a bipartite pure state $|\phi\rangle_{AB}$ on Hilbert space $H_A \otimes H_B$, the concurrence is given by [18, 19, 20]

$$C(|\phi\rangle_{AB}) = \sqrt{2(1 - \text{Tr}(\rho_A^2))}, \quad (1)$$

where ρ_A is the reduced density matrix by tracing over the subsystem B , $\rho_A = \text{Tr}_B(|\phi\rangle_{AB}\langle\phi|)$. For a bipartite mixed state ρ_{AB} , the concurrence is defined by the convex-roof,

$$C(\rho_{AB}) = \min_{\{p_i, |\phi_i\rangle\}} \sum_i p_i C(|\phi_i\rangle_{AB}), \quad (2)$$

where the minimum is taken over all possible pure state decompositions of $\rho_{AB} = \sum_i p_i |\phi_i\rangle\langle\phi_i|$, with $\sum_i p_i = 1$ and $p_i \geq 0$.

For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, the concurrence $C(\rho_{A|B_1 \dots B_{N-1}})$ of the state $\rho_{AB_1 \dots B_{N-1}}$ under bipartite partition A and $B_1 \dots B_{N-1}$ satisfies [21]

$$C^\beta(\rho_{A|B_1 \dots B_{N-1}}) \geq C^\beta(\rho_{AB_1}) + C^\beta(\rho_{AB_2}) + \dots + C^\beta(\rho_{AB_{N-1}}), \quad (3)$$

for $\beta \geq 2$. Furthermore, for an N -qubit mixed state, if $C_{AB_i} \geq C_{A|B_{i+1} \dots B_{N-1}}$ for $i = 1, 2, \dots, m$, and $C_{AB_j} \leq C_{A|B_{j+1} \dots B_{N-1}}$ for $j = m+1, \dots, N-2$, a generalized monogamy relation for $\beta \geq 2$ was presented as [22]

$$\begin{aligned} C^\beta(\rho_{A|B_1 \dots B_{N-1}}) &\geq C^\beta(\rho_{AB_1}) + (2^{\frac{\beta}{2}} - 1)C^\beta(\rho_{AB_2}) + \dots + (2^{\frac{\beta}{2}} - 1)^{m-1}C^\beta(\rho_{AB_m}) \\ &\quad + (2^{\frac{\beta}{2}} - 1)^{m+1}[C^\beta(\rho_{AB_{m+1}}) + \dots + C^\beta(\rho_{AB_{N-2}})] + (2^{\frac{\beta}{2}} - 1)^m C^\beta(\rho_{AB_{N-1}}), \end{aligned} \quad (4)$$

where $1 \leq m \leq N-3$, $N \geq 4$.

In the following, we will show that these monogamy relations for concurrence can be further tightened under some conditions. Before that, we first introduce two lemmas as follows.

Lemma 1. For any $x \in [0, 1]$ and $t \geq 1$, we have

$$\begin{aligned} (1+x)^t &\geq 1 + \frac{t}{2}x + \frac{(t-1)^2}{4}x^2 + (2^t - \frac{t}{2} + \frac{(t-1)^2}{4} - 1)x^t - \frac{(t-1)^2}{2}x^{t+1} \\ &\geq 1 + \frac{t}{2}x + (2^t - \frac{t}{2} - 1)x^t \geq 1 + (2^t - 1)x^t. \end{aligned} \quad (5)$$

Proof. Let us consider the function $f(t, x) = \frac{(1+x)^t - 1 - \frac{t}{2}x - \frac{(t-1)^2}{4}x^2 + \frac{(t-1)^2}{2}x^{t+1}}{x^t}$. Then, $\frac{\partial f(t, x)}{\partial x} = \frac{tx^{t-1}[1 + \frac{t-1}{2}x + \frac{(t-1)^2(t-2)}{4t}x^2 + \frac{(t-1)^2}{2t}x^{t+1} - (1+x)^{t-1}]}{x^{2t}}$. Next, we will prove that

$$1 + \frac{t-1}{2}x + \frac{(t-1)^2(t-2)}{4t}x^2 + \frac{(t-1)^2}{2t}x^{t+1} \leq (1+x)^{t-1}, \quad (6)$$

thus $\frac{\partial f(t, x)}{\partial x} \leq 0$, $f(t, x)$ is a decreasing function of x , i.e. $f(t, x) \geq f(t, 1) = 2^t - \frac{t}{2} + \frac{(t-1)^2}{4} - 1$. It follows that $(1+x)^t \geq 1 + \frac{t}{2}x + \frac{(t-1)^2}{4}x^2 + (2^t - \frac{t}{2} + \frac{(t-1)^2}{4} - 1)x^t - \frac{(t-1)^2}{2}x^{t+1}$.

For the case $1 \leq t \leq 2$, it is obvious that $(1+x)^{t-1} \geq 1 + (t-1)x + \frac{(t-1)(t-2)}{2}x^2$. Besides, we have

$$\begin{aligned} \frac{(t-1)(t-2)}{2}x^2 &= \frac{t-1}{4t}2t(t-2)x^2 = \frac{t-1}{4t}[(t-1)(t-2)x^2 + (t^2+t-2)x^2 - 2tx^2] \\ &\geq \frac{t-1}{4t}[(t-1)(t-2)x^2 + (2t-2)x^{t+1} - 2tx]. \end{aligned}$$

Thus, the Eq. (6) is hold.

For the case $t \geq 2$, it is obvious that $(1+x)^{t-1} \geq 1 + (t-1)x + \frac{(t-1)(t-2)}{4}x^2$. Besides, we have

$$\begin{aligned} \frac{(t-1)(t-2)}{4}x^2 &= \frac{t-1}{4t}t(t-2)x^2 = \frac{t-1}{4t}[(t-1)(t-2)x^2 + (2t-2)x^2 - tx^2] \\ &\geq \frac{t-1}{4t}[(t-1)(t-2)x^2 + 2(t-1)x^{t+1} - 2tx]. \end{aligned}$$

Thus, the Eq. (6) is hold.

On the other hand, Since $x^2 - 2x^{t+1} + x^t \geq 0$ and $\frac{(t-1)^2}{4} \geq 0$, for $t \geq 1$ and $x \in [0, 1]$, we can get $(1+x)^t \geq 1 + \frac{t}{2}x + \frac{(t-1)^2}{4}x^2 + (2^t - \frac{t}{2} + \frac{(t-1)^2}{4} - 1)x^t - \frac{(t-1)^2}{2}x^{t+1} \geq 1 + \frac{t}{2}x + (2^t - \frac{t}{2} - 1)x^t \geq 1 + (2^t - 1)x^t$. \square

Lemma 2. For any mixed state ρ_{ABC} in a $2 \otimes 2 \otimes 2^{N-2}$ system, suppose that $C_{AB} \geq C_{AC}$, we have

$$C_{A|BC}^\beta \geq C_{AB}^\beta + hC_{AC}^\beta + \frac{\beta}{4}C_{AC}^2(C_{AB}^{\beta-2} - C_{AC}^{\beta-2}) + \frac{(\beta-2)^2}{16}C_{AC}^4(C_{AB}^{\beta-4} + C_{AC}^{\beta-4} - 2C_{AC}^{\beta-2}C_{AB}^{-2}), \quad (7)$$

for all $\beta \geq 2$, where $h = 2^{\frac{\beta}{2}} - 1$, $C_{A|BC} = C(\rho_{A|BC})$, analogously for C_{AB} and C_{AC} .

Proof. Since $C_{AB} \geq C_{AC}$, we obtain

$$\begin{aligned} C_{A|BC}^\beta &\geq (C_{AB}^2 + C_{AC}^2)^{\frac{\beta}{2}} = C_{AB}^\beta \left(1 + \frac{C_{AC}^2}{C_{AB}^2}\right)^{\frac{\beta}{2}} \\ &\geq C_{AB}^\beta \left[1 + \frac{\beta}{4} \frac{C_{AC}^2}{C_{AB}^2} + \frac{(\beta-2)^2}{16} \frac{C_{AC}^4}{C_{AB}^4} + \left(2^{\frac{\beta}{2}} - \frac{\beta}{4} + \frac{(\beta-2)^2}{16} - 1\right) \frac{C_{AC}^\beta}{C_{AB}^\beta} - \frac{(\beta-2)^2}{8} \frac{C_{AC}^{\beta+2}}{C_{AB}^{\beta+2}}\right] \\ &= C_{AB}^\beta + hC_{AC}^\beta + \frac{\beta}{4}C_{AC}^2(C_{AB}^{\beta-2} - C_{AC}^{\beta-2}) + \frac{(\beta-2)^2}{16}C_{AC}^4(C_{AB}^{\beta-4} + C_{AC}^{\beta-4} - 2C_{AC}^{\beta-2}C_{AB}^{-2}), \end{aligned} \quad (8)$$

where the first inequality is due to the fact that $C_{A|BC}^2 \geq C_{AB}^2 + C_{AC}^2$ for any $2 \otimes 2 \otimes 2^{N-2}$ tripartite state $\rho_{A|BC}$ [9, 23] and the second inequality is due to Lemma 1. \square

Theorem 1. For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, if $C_{AB_i} \geq C_{A|B_{i+1} \dots B_{N-1}}$, for $i = 1, 2, \dots, N-2$, we have

$$C_{A|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^{N-2} h^{i-1} (C_{AB_i}^\beta + P_{AB_i}) + h^{N-2} C_{AB_{N-1}}^\beta, \quad (9)$$

for all $N \geq 3$, $\beta \geq 2$, where $h = 2^{\frac{\beta}{2}} - 1$, $P_{AB_i} = \frac{\beta}{4}C_{A|B_{i+1} \dots B_{N-1}}^2(C_{AB_i}^{\beta-2} - C_{A|B_{i+1} \dots B_{N-1}}^{\beta-2}) + \frac{(\beta-2)^2}{16}C_{A|B_{i+1} \dots B_{N-1}}^4(C_{AB_i}^{\beta-4} + C_{A|B_{i+1} \dots B_{N-1}}^{\beta-4} - 2C_{A|B_{i+1} \dots B_{N-1}}^{\beta-2}C_{AB_i}^{-2})$.

Proof. Due to Eq. (7), we obtain

$$\begin{aligned}
& C_{A|B_1 \dots B_{N-1}}^\beta \\
& \geq C_{AB_1}^\beta + h C_{A|B_2 \dots B_{N-1}}^\beta + \frac{\beta}{4} C_{A|B_2 \dots B_{N-1}}^2 (C_{AB_1}^{\beta-2} - C_{A|B_2 \dots B_{N-1}}^{\beta-2}) \\
& \quad + \frac{(\beta-2)^2}{16} C_{A|B_2 \dots B_{N-1}}^4 (C_{AB_1}^{\beta-4} + C_{A|B_2 \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_2 \dots B_{N-1}}^{\beta-2} C_{AB_1}^{\beta-2}) \\
& \geq C_{AB_1}^\beta + h \left[C_{AB_2}^\beta + h C_{A|B_3 \dots B_{N-1}}^\beta + \frac{\beta}{4} C_{A|B_3 \dots B_{N-1}}^2 (C_{AB_2}^{\beta-2} - C_{A|B_3 \dots B_{N-1}}^{\beta-2}) \right. \\
& \quad \left. + \frac{(\beta-2)^2}{16} C_{A|B_3 \dots B_{N-1}}^4 (C_{AB_2}^{\beta-4} + C_{A|B_3 \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_3 \dots B_{N-1}}^{\beta-2} C_{AB_2}^{\beta-2}) \right] \\
& \quad + \frac{\beta}{4} C_{A|B_2 \dots B_{N-1}}^2 (C_{AB_1}^{\beta-2} - C_{A|B_2 \dots B_{N-1}}^{\beta-2}) \\
& \quad + \frac{(\beta-2)^2}{16} C_{A|B_2 \dots B_{N-1}}^4 (C_{AB_1}^{\beta-4} + C_{A|B_2 \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_2 \dots B_{N-1}}^{\beta-2} C_{AB_1}^{\beta-2}) \\
& \geq \dots \\
& \geq C_{AB_1}^\beta + h C_{AB_2}^\beta + \dots + h^{N-2} C_{AB_{N-1}}^\beta + h^{N-3} \left[\frac{\beta}{4} C_{AB_{N-1}}^2 (C_{AB_{N-2}}^{\beta-2} - C_{AB_{N-1}}^{\beta-2}) \right. \\
& \quad \left. + \frac{(\beta-2)^2}{16} C_{AB_{N-1}}^4 (C_{AB_{N-2}}^{\beta-4} + C_{AB_{N-1}}^{\beta-4} - 2 C_{AB_{N-1}}^{\beta-2} C_{AB_{N-2}}^{\beta-2}) \right] \\
& \quad + \dots + h \left[\frac{\beta}{4} C_{A|B_3 \dots B_{N-1}}^2 (C_{AB_2}^{\beta-2} - C_{A|B_3 \dots B_{N-1}}^{\beta-2}) \right. \\
& \quad \left. + \frac{(\beta-2)^2}{16} C_{A|B_3 \dots B_{N-1}}^4 (C_{AB_2}^{\beta-4} + C_{A|B_3 \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_3 \dots B_{N-1}}^{\beta-2} C_{AB_2}^{\beta-2}) \right] \\
& \quad + \frac{\beta}{4} C_{A|B_2 \dots B_{N-1}}^2 (C_{AB_1}^{\beta-2} - C_{A|B_2 \dots B_{N-1}}^{\beta-2}) \\
& \quad + \frac{(\beta-2)^2}{16} C_{A|B_2 \dots B_{N-1}}^4 (C_{AB_1}^{\beta-4} + C_{A|B_2 \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_2 \dots B_{N-1}}^{\beta-2} C_{AB_1}^{\beta-2}).
\end{aligned} \tag{10}$$

By the denotation of P_{AB_i} , we complete the proof. \square

Theorem 2. For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, if $C_{AB_i} \geq C_{A|B_{i+1} \dots B_{N-1}}$ for $i = 1, 2, \dots, m$, and $C_{AB_j} \leq C_{A|B_{j+1} \dots B_{N-1}}$ for $j = m+1, \dots, N-2$, $\forall 1 \leq m \leq N-3$, we have

$$C_{A|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^m h^{i-1} (C_{AB_i}^\beta + P_{AB_i}) + h^m \sum_{j=m+1}^{N-2} (h C_{AB_j}^\beta + P_{AB_j}^1) + h^m C_{AB_{N-1}}^\beta, \tag{11}$$

for all $N \geq 4$, $\beta \geq 2$, where $h = 2^{\frac{\beta}{2}} - 1$, $P_{AB_i} = \frac{\beta}{4} C_{A|B_{i+1} \dots B_{N-1}}^2 (C_{AB_i}^{\beta-2} - C_{A|B_{i+1} \dots B_{N-1}}^{\beta-2}) + \frac{(\beta-2)^2}{16} C_{A|B_{i+1} \dots B_{N-1}}^4 (C_{AB_i}^{\beta-4} + C_{A|B_{i+1} \dots B_{N-1}}^{\beta-4} - 2 C_{A|B_{i+1} \dots B_{N-1}}^{\beta-2} C_{AB_i}^{\beta-2})$, $P_{AB_j}^1 = \frac{\beta}{4} C_{AB_j}^2 (C_{A|B_{j+1} \dots B_{N-1}}^{\beta-2} - C_{AB_j}^{\beta-2}) + \frac{(\beta-2)^2}{16} C_{AB_j}^4 (C_{A|B_{j+1} \dots B_{N-1}}^{\beta-4} + C_{AB_j}^{\beta-4} - 2 C_{AB_j}^{\beta-2} C_{A|B_{j+1} \dots B_{N-1}}^{\beta-2})$.

Proof. Due to the proof process of Theorem 1, we can get that

$$C_{A|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^m h^{i-1} (C_{AB_i}^\beta + P_{AB_i}) + h^m C_{A|B_{m+1} \dots B_{N-1}}^\beta. \tag{12}$$

In addition, since $C_{AB_j} \leq C_{A|B_{j+1} \dots B_{N-1}}$ for $j = m+1, \dots, N-2$, hence,

$$\begin{aligned}
& C_{A|B_{m+1} \dots B_{N-1}}^\beta \\
& \geq C_{A|B_{m+2} \dots B_{N-1}}^\beta + h C_{AB_{m+1}}^\beta + \frac{\beta}{4} C_{AB_{m+1}}^2 (C_{A|B_{m+2} \dots B_{N-1}}^{\beta-2} - C_{AB_{m+1}}^{\beta-2}) \\
& \quad + \frac{(\beta-2)^2}{16} C_{AB_{m+1}}^4 (C_{A|B_{m+2} \dots B_{N-1}}^{\beta-4} + C_{AB_{m+1}}^{\beta-4} - 2 C_{AB_{m+1}}^{\beta-2} C_{A|B_{m+2} \dots B_{N-1}}^{\beta-2}) \\
& \geq \sum_{j=m+1}^{N-2} (h C_{AB_j}^\beta + P_{AB_j}^1) + C_{AB_{N-1}}^\beta.
\end{aligned} \tag{13}$$

Combing Eq. (12) and Eq. (13), we can get the inequality (11). \square

Example 1. Consider the three-qubit state $|\psi\rangle_{ABC}$ in generalized Schmidt decomposition form [25, 26]

$$|\psi\rangle_{ABC} = \lambda_0 |000\rangle + \lambda_1 e^{i\varphi} |100\rangle + \lambda_2 |101\rangle + \lambda_3 |110\rangle + \lambda_4 |111\rangle, \tag{14}$$




Figure 1: Dash dotted line, $C_{A|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$); solid line, the lower bound of $C_{A|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$) in Eq. (11); dash line, the lower bound of $C_{A|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$) in [24].

where $\lambda_i \geq 0$, $i = 0, 1, 2, 3, 4$, and $\sum_{i=0}^4 \lambda_i^2 = 1$. A direct calculation shows that $C_{A|BC} = 2\lambda_0\sqrt{\lambda_2^2 + \lambda_3^2 + \lambda_4^2}$, $C_{AB} = 2\lambda_0\lambda_2$ and $C_{AC} = 2\lambda_0\lambda_3$.

Set $\lambda_0 = \frac{\sqrt{2}}{3}$, $\lambda_1 = 0$, $\lambda_2 = \frac{\sqrt{5}}{3}$, $\lambda_3 = \frac{\sqrt{2}}{3}$, $\lambda_4 = 0$. We have $C_{A|BC} = \frac{2\sqrt{14}}{9}$, $C_{AB} = \frac{2\sqrt{10}}{9}$ and $C_{AC} = \frac{4}{9}$. Then $C_{A|BC}^\beta = (\frac{2\sqrt{14}}{9})^\beta \geq C_{AB}^\beta + hC_{AC}^\beta + \frac{\beta}{4}C_{AC}^2(C_{AB}^{\beta-2} - C_{AC}^{\beta-2}) + \frac{(\beta-2)^2}{16}C_{AC}^4(C_{AB}^{\beta-4} + C_{AC}^{\beta-4} - 2C_{AC}^{\beta-2}C_{AB}^{\beta-2}) = (\frac{2\sqrt{10}}{9})^\beta + h(\frac{4}{9})^\beta + \frac{\beta}{4}(\frac{4}{9})^2 \left[(\frac{2\sqrt{10}}{9})^{\beta-2} - (\frac{4}{9})^{\beta-2} \right] + \frac{(\beta-2)^2}{16}(\frac{4}{9})^4 \left[(\frac{2\sqrt{10}}{9})^{\beta-4} + (\frac{4}{9})^{\beta-4} - 2(\frac{4}{9})^{\beta-2}(\frac{2\sqrt{10}}{9})^{\beta-2} \right]$. However, the result in [24] is $C_{AB}^\beta + hC_{AC}^\beta + \frac{\beta}{4}C_{AC}^2(C_{AB}^{\beta-2} - C_{AC}^{\beta-2}) = (\frac{2\sqrt{10}}{9})^\beta + h(\frac{4}{9})^\beta + \frac{\beta}{4}(\frac{4}{9})^2 \left[(\frac{2\sqrt{10}}{9})^{\beta-2} - (\frac{4}{9})^{\beta-2} \right]$. We can see that our results are better than the ones in [24] for $\beta \geq 2$, see FIG. 1.

3 Tighter Monogamy Relations using EoF

Let H_A and H_B be two Hilbert spaces with dimension m and n ($m \leq n$). The entanglement of formation (EoF) of a pure state $|\phi\rangle_{AB}$ on Hilbert space $H_A \otimes H_B$, is defined as [27, 28]

$$E(|\phi\rangle_{AB}) = S(\rho_A) = -\text{Tr}(\rho_A \log_2 \rho_A), \quad (15)$$

where $S(\rho) = -\text{Tr}(\rho \log_2 \rho)$ and $\rho_A = \text{Tr}_B(|\phi\rangle_{AB}\langle\phi|)$. For a bipartite mixed state $|\phi\rangle_{AB}$ on Hilbert space $H_A \otimes H_B$, the EoF is given by

$$E(\rho_{AB}) = \inf_{\{p_i, |\phi_i\rangle\}} \sum_i p_i E(|\phi_i\rangle), \quad (16)$$

where the infimum is taken over all possible pure state decompositions of ρ_{AB} .

Let $g(x) = H(\frac{1+\sqrt{1-x}}{2})$ and $H(x) = -x \log_2 x - (1-x) \log_2 (1-x)$, it is obvious that $g(x)$ is a monotonically increasing function for $0 \leq x \leq 1$, and satisfies

$$g^{\sqrt{2}}(x^2 + y^2) \geq g^{\sqrt{2}}(x^2) + g^{\sqrt{2}}(y^2), \quad (17)$$

where $g^{\sqrt{2}}(x^2 + y^2) = [g(x^2 + y^2)]^{\sqrt{2}}$.

From Eqs. (15) and (16), we have $E(|\phi\rangle) = g(\mathcal{C}^2(|\phi\rangle))$ for $2 \otimes d$ ($d \geq 2$) pure state $|\phi\rangle$. And $E(\rho) = g(\mathcal{C}^2(\rho))$ for arbitrary two-qubit mixed state ρ [29].

Wootters [8] shows that the EoF dose not satisfy the monogamy inequality $E_{AB} + E_{AC} \leq E_{A|BC}$. In [30], the authors shows that EoF is a monotonic function satisfying $E^2(C_{A|B_1 B_2 \dots B_{N-1}}^2) \geq E^2 \sum_{i=1}^{N-1} (C_{AB_i}^2)$. For N -qubit systems, one has [21]

$$E_{A|B_1 B_2 \dots B_{N-1}}^\beta \geq E_{AB_1}^\beta + E_{AB_2}^\beta + \dots + E_{AB_{N-1}}^\beta, \quad (18)$$

for $\beta \geq \sqrt{2}$, where $E_{A|B_1 B_2 \dots B_{N-1}}$ is the EoF of ρ under bipartite partition $A|B_1 B_2 \dots B_{N-1}$, E_{AB_i} is the EoF of the mixed state $\rho_{AB_i} = \text{Tr}_{B_1 \dots B_{i-1}, B_{i+1} \dots B_{N-1}}(\rho)$ for $i = 1, 2, \dots, N-1$.

Lemma 3. For any mixed state ρ_{ABC} in a $2 \otimes 2 \otimes 2^{N-2}$ system, $\beta \geq \sqrt{2}$, if $C_{AB} \geq C_{AC}$, then we have

$$E_{A|BC}^\beta \geq E_{AB}^\beta + h E_{AC}^\beta + \frac{t}{2} E_{AC}^{\sqrt{2}} (E_{AB}^{\beta-\sqrt{2}} - E_{AC}^{\beta-\sqrt{2}}) + \frac{(t-1)^2}{4} E_{AC}^{2\sqrt{2}} (E_{AB}^{\beta-2\sqrt{2}} + E_{AC}^{\beta-2\sqrt{2}} - 2 E_{AC}^{\beta-\sqrt{2}} E_{AB}^{\beta-\sqrt{2}}), \quad (19)$$

where $t = \frac{\beta}{\sqrt{2}}$, $h = 2^t - 1$.

Proof. The proof is similar to the proof of Lemma 2. \square

In fact, the result can be generalized to N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$. The following theorem holds for $\rho_{AB_1 \dots B_{N-1}}$.

Theorem 3. For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, if $C_{AB_i} \geq C_{A|B_{i+1} \dots B_{N-1}}$ for $i = 1, 2, \dots, N-2$, we have

$$E_{A|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^{N-2} h^{i-1} (E_{AB_i}^\beta + Q_{AB_i}) + h^{N-2} E_{AB_{N-1}}^\beta, \quad (20)$$

for $\beta \geq \sqrt{2}$, where $h = 2^t - 1$, $t = \frac{\beta}{\sqrt{2}}$, $Q_{AB_i} = \frac{t}{2} (E_{AB_{i+1}}^{\sqrt{2}} + \dots + E_{AB_{N-1}}^{\sqrt{2}}) (E_{AB_i}^{\beta-\sqrt{2}} - E_{A|B_{i+1} \dots B_{N-1}}^{\beta-\sqrt{2}}) + \frac{(t-1)^2}{4} (E_{AB_{i+1}}^{2\sqrt{2}} + \dots + E_{AB_{N-1}}^{2\sqrt{2}}) [E_{AB_i}^{\beta-2\sqrt{2}} + \dots + E_{AB_{N-1}}^{\beta-2\sqrt{2}} - 2 (E_{A|B_{i+1} \dots B_{N-1}}^{\beta-\sqrt{2}}) E_{AB_i}^{\beta-\sqrt{2}}]$.

Proof. Let $\rho = \sum_i p_i |\psi_i\rangle \langle \psi_i| \in H_A \otimes H_{B_1} \otimes \dots \otimes H_{B_{N-1}}$ be the optimal decomposition of $E_{A|B_1 B_2 \dots B_{N-1}}(\rho)$ for the N -qubit mixed state ρ , we have [22]

$$E_{A|B_1 B_2 \dots B_{N-1}} \geq g(C_{A|B_1 B_2 \dots B_{N-1}}^2). \quad (21)$$

In addition, for $\beta \geq \sqrt{2}$, we have

$$\begin{aligned} & g^\beta(x^2 + y^2) \\ &= \left[g^{\sqrt{2}}(x^2 + y^2) \right]^t \geq \left[g^{\sqrt{2}}(x^2) + g^{\sqrt{2}}(y^2) \right]^t \\ &\geq g^\beta(x^2) + (2^t - 1) g^\beta(y^2) + \frac{t}{2} g^{\sqrt{2}}(y^2) \left[g^{\beta-\sqrt{2}}(x^2) - g^{\beta-\sqrt{2}}(y^2) \right] \\ &\quad + \frac{(t-1)^2}{4} g^{2\sqrt{2}}(y^2) \left[g^{\beta-2\sqrt{2}}(x^2) + g^{\beta-2\sqrt{2}}(y^2) - 2 g^{\beta-\sqrt{2}}(y^2) g^{\beta-\sqrt{2}}(x^2) \right], \end{aligned} \quad (22)$$

where the first inequality is due to Eq. (17), and without loss of generality, we can assume $x^2 \geq y^2$, then the second inequality is obtained from the monotonicity of $g(x)$ and Eq. (5).

Thus, combining Eqs. (21) and (22), we obtain

$$\begin{aligned}
E_{A|B_1B_2\cdots B_{N-1}}^\beta &\geq g^\beta(C_{AB_1}^2 + C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \\
&\geq g^\beta(C_{AB_1}^2) + hg^\beta(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \\
&\quad + \frac{t}{2}g^{\sqrt{2}}(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \left[g^{\beta-\sqrt{2}}(C_{AB_1}^2) - g^{\beta-\sqrt{2}}(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \right] \\
&\quad + \frac{(t-1)^2}{4}g^{2\sqrt{2}}(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \left[g^{\beta-2\sqrt{2}}(C_{AB_1}^2) + g^{\beta-2\sqrt{2}}(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \right. \\
&\quad \left. - 2g^{\beta-\sqrt{2}}(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2)g^{-\sqrt{2}}(C_{AB_1}^2) \right] \\
&\geq g^\beta(C_{AB_1}^2) + hg^\beta(C_{AB_2}^2 + \cdots + C_{AB_{N-1}}^2) \\
&\quad + \frac{t}{2} \left[g^{\sqrt{2}}(C_{AB_2}^2) + \cdots + g^{\sqrt{2}}(C_{AB_{N-1}}^2) \right] \cdot \left[g^{\beta-\sqrt{2}}(C_{AB_1}^2) - g^{\beta-\sqrt{2}}(C_{A|B_2\cdots B_{N-1}}^2) \right] \\
&\quad + \frac{(t-1)^2}{4} \left[g^{2\sqrt{2}}(C_{AB_2}^2) + \cdots + g^{2\sqrt{2}}(C_{AB_{N-1}}^2) \right] \cdot \left[g^{\beta-2\sqrt{2}}(C_{AB_1}^2) + \cdots + g^{\beta-2\sqrt{2}}(C_{AB_{N-1}}^2) \right. \\
&\quad \left. - 2g^{\beta-\sqrt{2}}(C_{A|B_2\cdots B_{N-1}}^2)g^{-\sqrt{2}}(C_{AB_1}^2) \right] \\
&\geq g^\beta(C_{AB_1}^2) + hg^\beta(C_{AB_2}^2) + \cdots + h^{N-2}g^\beta(C_{AB_{N-1}}^2) \\
&\quad + h^{N-3} \cdot \frac{t}{2} \cdot g^{\sqrt{2}}(C_{AB_{N-1}}^2) \left[g^{\beta-\sqrt{2}}(C_{AB_{N-2}}^2) - g^{\beta-\sqrt{2}}(C_{AB_{N-1}}^2) \right] + \cdots \\
&\quad + \frac{t}{2} \left[g^{\sqrt{2}}(C_{AB_2}^2) + \cdots + g^{\sqrt{2}}(C_{AB_{N-1}}^2) \right] \cdot \left[g^{\beta-\sqrt{2}}(C_{AB_1}^2) - g^{\beta-\sqrt{2}}(C_{A|B_2\cdots B_{N-1}}^2) \right] \\
&\quad + h^{N-3} \cdot \frac{(t-1)^2}{4} \cdot g^{2\sqrt{2}}(C_{AB_{N-1}}^2) \left[g^{\beta-2\sqrt{2}}(C_{AB_{N-2}}^2) + g^{\beta-2\sqrt{2}}(C_{AB_{N-1}}^2) \right. \\
&\quad \left. - 2g^{\beta-\sqrt{2}}(C_{AB_{N-1}}^2)g^{-\sqrt{2}}(C_{AB_{N-2}}^2) \right] + \cdots \\
&\quad + \frac{(t-1)^2}{4} \left[g^{2\sqrt{2}}(C_{AB_2}^2) + \cdots + g^{2\sqrt{2}}(C_{AB_{N-1}}^2) \right] \cdot \left[g^{\beta-2\sqrt{2}}(C_{AB_1}^2) + \cdots + g^{\beta-2\sqrt{2}}(C_{AB_{N-1}}^2) \right. \\
&\quad \left. - 2g^{\beta-\sqrt{2}}(C_{A|B_2\cdots B_{N-1}}^2)g^{-\sqrt{2}}(C_{AB_1}^2) \right],
\end{aligned} \tag{23}$$

where we have utilized the Eq. (3) and the monotonicity of $g(x)$ to obtain the first inequality, the third and the forth inequalities are due to the Eq. (17) and the monotonicity of the function $g(x)$.

According to Eq. (21) and the fact that $g(C^2(\rho)) = E(\rho)$ for arbitrary two-qubit mixed state ρ , we obtain Eq. (20). \square

Example 2. Let us consider the state in (14) given in Example 1. Set $\lambda_0 = \frac{\sqrt{6}}{3}, \lambda_1 = 0, \lambda_2 = \frac{\sqrt{2}}{3}, \lambda_3 = \frac{1}{3}, \lambda_4 = 0$, we have $E_{A|BC} = 0.91829$, $E_{AB} = 0.68193$, $E_{AC} = 0.40416$. Then $E_{A|BC}^\beta = (0.91829)^\beta \geq E_{AB}^\beta + hE_{AC}^\beta + \frac{\beta}{2\sqrt{2}}E_{AC}^{\sqrt{2}}(E_{AB}^{\beta-\sqrt{2}} - E_{AC}^{\beta-\sqrt{2}}) + \frac{(\beta-\sqrt{2})^2}{8}E_{AC}^{2\sqrt{2}}(E_{AB}^{\beta-2\sqrt{2}} + E_{AC}^{\beta-2\sqrt{2}} - 2E_{AC}^{\beta-\sqrt{2}}E_{AB}^{\beta-\sqrt{2}}) = (0.68193)^\beta + h(0.40416)^\beta + \frac{\beta}{2\sqrt{2}}(0.40416)^{\sqrt{2}} \left[(0.68193)^{\beta-\sqrt{2}} - (0.40416)^{\beta-\sqrt{2}} \right] + \frac{(\beta-\sqrt{2})^2}{8}(0.40416)^{2\sqrt{2}} \left[(0.68193)^{\beta-2\sqrt{2}} + (0.40416)^{\beta-2\sqrt{2}} - 2(0.40416)^{\beta-\sqrt{2}}(0.68193)^{\beta-\sqrt{2}} \right]$. While the result in [24] is $E_{AB}^\beta + hE_{AC}^\beta + \frac{\beta}{2\sqrt{2}}E_{AC}^{\sqrt{2}}(E_{AB}^{\beta-\sqrt{2}} - E_{AC}^{\beta-\sqrt{2}}) = (0.68193)^\beta + h(0.40416)^\beta + \frac{\beta}{2\sqrt{2}}(0.40416)^{\sqrt{2}} \left[(0.68193)^{\beta-\sqrt{2}} - (0.40416)^{\beta-\sqrt{2}} \right]$. We can see that our results are better than the ones in [24], see FIG. 2.

4 Tighter Monogamy Relations using Negativity

The negativity is a well-known quantifier of bipartite entanglement. Given a bipartite state ρ_{AB} in Hilbert space $H_A \otimes H_B$, the negativity is defined as [31]

$$\mathcal{N}(\rho_{AB}) = \frac{\|\rho_{AB}^{T_A}\| - 1}{2}, \tag{24}$$




Figure 2: Dash dotted line , $E_{A|BC}^\beta$ as a function of β ($\sqrt{2} \leq \beta \leq 10$); solid line, the lower bound of $E_{A|BC}^\beta$ as a function of β ($\sqrt{2} \leq \beta \leq 10$) in Eq. (20); dash line, the lower bound of $E_{A|BC}^\beta$ as a function of β ($\sqrt{2} \leq \beta \leq 10$) in [24].

where $\rho_{AB}^{T_A}$ is the partial transposed matrix of ρ_{AB} with respect to the subsystem A and $\|X\|$ denotes the trace norm of X , i.e. $\|X\| = \text{Tr}\sqrt{XX^\dagger}$. For convenience, we use the definition of negativity as $\|\rho_{AB}^{T_A}\| - 1$ [10].

If a bipartite pure state $|\phi\rangle_{AB}$ with the Schmidt decomposition, $|\phi\rangle_{AB} = \sum_i \sqrt{\lambda_i} |ii\rangle$, $\lambda_i \geq 0$, $\sum_i \lambda_i = 1$, then [10]

$$\mathcal{N}(|\phi\rangle_{AB}) = 2 \sum_{i < j} \sqrt{\lambda_i \lambda_j}. \quad (25)$$

From the definition of concurrence (1), we have

$$C(|\phi\rangle_{AB}) = 2 \sqrt{\sum_{i < j} \lambda_i \lambda_j}. \quad (26)$$

As a consequence, for any bipartite pure state $|\phi\rangle_{AB}$ with Schmidt rank 2, one has $\mathcal{N}(|\phi\rangle_{AB}) = C(|\phi\rangle_{AB})$.

For a mixed state ρ_{AB} , the convex-roof extended negativity (CREN) is given by

$$\mathcal{N}_c(\rho_{AB}) = \min_{\{p_i, |\phi_i\rangle\}} \sum_i p_i \mathcal{N}(|\phi_i\rangle), \quad (27)$$

where the minimum is taken over all possible pure state decomposition of ρ_{AB} . CREN gives a perfect discrimination between PPT bound entangled states and separable states in any bipartite quantum system [32]. It follows that for any $2 \otimes d$ ($d \geq 2$) mixed state ρ_{AB} , we have

$$\mathcal{N}_c(\rho_{AB}) = \min_{\{p_i, |\phi_i\rangle\}} \sum_i p_i \mathcal{N}(|\phi_i\rangle) = \min_{\{p_i, |\phi_i\rangle\}} \sum_i p_i C(|\phi_i\rangle) = C(\rho_{AB}). \quad (28)$$

According to the relation between CREN and concurrence, we have the following results for the lower bound of $\mathcal{N}_{cA|B_1 \dots B_{N-1}}^\beta$.




Figure 3: Dash dotted line , $\mathcal{N}_{cA|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$); solid line, the lower bound of $\mathcal{N}_{cA|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$) in Eq. (30); dash line, the lower bound of $\mathcal{N}_{cA|BC}^\beta$ as a function of β ($2 \leq \beta \leq 10$) in [24].

Theorem 4. For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, if $\mathcal{N}_{cAB_i} \geq \mathcal{N}_{cAB_{i+1} \dots B_{N-1}}$ for $i = 1, 2, \dots, m$, and $\mathcal{N}_{cAB_j} \leq \mathcal{N}_{cA|B_{j+1} \dots B_{N-1}}$ for $j = m+1, \dots, N-2$, $\forall 1 \leq m \leq N-3$, then we have

$$\mathcal{N}_{cA|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^m h^{i-1} (\mathcal{N}_{cAB_i}^\beta + R_{AB_i}) + h^m \sum_{j=m+1}^{N-2} (h \mathcal{N}_{cAB_j}^\beta + R_{AB_j}^1) + h^m \mathcal{N}_{cAB_{N-1}}^\beta, \quad (29)$$

for all $N \geq 4$, $\beta \geq 2$, where $h = 2^{\frac{\beta}{2}} - 1$, $R_{AB_i} = \frac{\beta}{4} \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^2 (\mathcal{N}_{cAB_i}^{\beta-2} - \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-2}) + \frac{(\beta-2)^2}{16} \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^4 (\mathcal{N}_{cAB_i}^{\beta-4} + \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-4} - 2 \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-2} \mathcal{N}_{cAB_i}^{\beta-2})$, $R_{AB_j}^1 = \frac{\beta}{4} \mathcal{N}_{cAB_j}^2 (\mathcal{N}_{cA|B_{j+1} \dots B_{N-1}}^{\beta-2} - \mathcal{N}_{cAB_j}^{\beta-2}) + \frac{(\beta-2)^2}{16} \mathcal{N}_{cAB_j}^4 (\mathcal{N}_{cA|B_{j+1} \dots B_{N-1}}^{\beta-4} + \mathcal{N}_{cAB_j}^{\beta-4} - 2 \mathcal{N}_{cAB_j}^{\beta-2} \mathcal{N}_{cA|B_{j+1} \dots B_{N-1}}^{\beta-2})$.

Theorem 5. For any N -qubit mixed state $\rho_{AB_1 \dots B_{N-1}}$, if $\mathcal{N}_{cAB_i} \geq \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}$ for $i = 1, 2, \dots, N-2$, then we can obtain

$$\mathcal{N}_{cA|B_1 \dots B_{N-1}}^\beta \geq \sum_{i=1}^{N-2} h^{i-1} (\mathcal{N}_{cAB_i}^\beta + R_{AB_i}) + h^{N-2} \mathcal{N}_{cAB_{N-1}}^\beta, \quad (30)$$

for all $N \geq 3$, $\beta \geq 2$, where $h = 2^{\frac{\beta}{2}} - 1$, $R_{AB_i} = \frac{\beta}{4} \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^2 (\mathcal{N}_{cAB_i}^{\beta-2} - \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-2}) + \frac{(\beta-2)^2}{16} \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^4 (\mathcal{N}_{cAB_i}^{\beta-4} + \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-4} - 2 \mathcal{N}_{cA|B_{i+1} \dots B_{N-1}}^{\beta-2} \mathcal{N}_{cAB_i}^{\beta-2})$.

Example 3. Let us consider the state in (14) given in Example 1. We have $\mathcal{N}_{cA|BC} = 2\lambda_0 \sqrt{\lambda_2^2 + \lambda_3^2 + \lambda_4^2}$, $\mathcal{N}_{cAB} = 2\lambda_0 \lambda_2$ and $\mathcal{N}_{cAC} = 2\lambda_0 \lambda_3$. Set $\lambda_0 = \frac{\sqrt{2}}{3}$, $\lambda_1 = 0$, $\lambda_2 = \frac{\sqrt{5}}{3}$, $\lambda_3 = \frac{\sqrt{2}}{3}$, $\lambda_4 = 0$. We have $\mathcal{N}_{cA|BC}^\beta \geq \mathcal{N}_{cAB}^\beta + h \mathcal{N}_{cAC}^\beta + \frac{\beta}{4} \mathcal{N}_{cAC}^2 (\mathcal{N}_{cAB}^{\beta-2} - \mathcal{N}_{cAC}^{\beta-2}) + \frac{(\beta-2)^2}{16} \mathcal{N}_{cAC}^4 (\mathcal{N}_{cAB}^{\beta-4} + \mathcal{N}_{cAC}^{\beta-4} - 2 \mathcal{N}_{cAC}^{\beta-2} \mathcal{N}_{cAB}^{\beta-2}) = (\frac{2\sqrt{10}}{9})^\beta + h(\frac{4}{9})^\beta + \frac{\beta}{4}(\frac{4}{9})^2 \left[(\frac{2\sqrt{10}}{9})^{\beta-2} - (\frac{4}{9})^{\beta-2} \right] + \frac{(\beta-2)^2}{16} (\frac{4}{9})^4 \left[(\frac{2\sqrt{10}}{9})^{\beta-4} + (\frac{4}{9})^{\beta-4} - 2(\frac{4}{9})^{\beta-2} (\frac{2\sqrt{10}}{9})^{\beta-2} \right]$. While the result in [24] is $\mathcal{N}_{cAB}^\beta + h \mathcal{N}_{cAC}^\beta + \frac{\beta}{4} \mathcal{N}_{cAC}^2 (\mathcal{N}_{cAB}^{\beta-2} - \mathcal{N}_{cAC}^{\beta-2}) = (\frac{2\sqrt{10}}{9})^\beta + h(\frac{4}{9})^\beta + \frac{\beta}{4}(\frac{4}{9})^2 \left[(\frac{2\sqrt{10}}{9})^{\beta-2} - (\frac{4}{9})^{\beta-2} \right]$. We can see that our result is better than the one in [24] for $\beta \geq 2$, see FIG. 3.

5 Conclusion

Entanglement monogamy relations are fundamental properties of multipartite entangled states. In this paper, we have provided the multipartite entanglement based on the monogamy relations for β th power of concurrence $C_{A|B_1\dots B_{N-1}}^\beta$ ($\beta \geq 2$), entanglement of formation $E_{A|B_1\dots B_{N-1}}^\beta$ ($\beta \geq \sqrt{2}$) and convex-roof extended negativity $\mathcal{N}_{cA|B_1\dots B_{N-1}}^\beta$ ($\beta \geq 2$). Our monogamy relations have larger lower bounds and are tighter than the existing results [24]. These tighter monogamy inequalities can also provide a finer description of the entanglement distribution. In multi-qubit system, our research results provide a rich reference for future research on multi-party quantum entanglement. Our method can also be applied to the study of other properties of monogamy related to quantum correlations.

Acknowledgments

This work is supported by the Yunnan Provincial Research Foundation for Basic Research, China (Grant No. 202001AU070041), the Research Foundation of Education Bureau of Yunnan Province, China (Grant No. 2021J0054), the Basic and Applied Basic Research Funding Program of Guangdong Province (Grant No. 2019A1515111097), the Natural Science Foundation of Kunming University of Science and Technology (Grant No. KKZ3202007036, KKZ3202007049).

Data Availability Statement

All data generated or analysed during this study are included in this submitted article.

References

- [1] M. Jafarpour, F. Kazemi Hasanvand and D. Afshar, Dynamics of entanglement and measurement-induced disturbance for a hybrid qubit-qutrit system interacting with a spin-chain environment: a mean field approach, *Commun. Theor. Phys.* **67** 27 (2017).
- [2] F. G. Deng, B. C. Ren, and X. H. Li, Quantum hyperentanglement and its applications in quantum information processing, *Sci. Bull.* **62** 46 (2017).
- [3] H. L. Huang, A. K. Goswami, W. S. Bao, and P. K. Panigrahi, Demonstration of essentiality of entanglement in a Deutsch-like quantum algorithm, *Sci. China-Phys. Mech. Astron.* **61** 060311 (2018).
- [4] M. Y. Wang, J. Z. Xu, F. L. Yan, and T. Gao, Entanglement concentration for polarization-spatial-time-bin hyperentangled Bell states, *Europhys. Lett.* **123** 60002 (2018).
- [5] B. Terhal, Is entanglement monogamous? *IBM J. Res. Dev.* **48** 71 (2004).

- [6] C. H. Bennett, Quantum cryptography using any two nonorthogonal states, *Phys. Rev. Lett.* **68** 3121 (1992).
- [7] M. Pawłowski, Security proof for cryptographic protocols based only on the monogamy of Bell's inequality violations, *Phys. Rev. A* **82** 032313 (2010).
- [8] V. Coffman, J. Kundu, and W. K. Wootters, Distributed entanglement, *Phys. Rev. A* **61** 052306 (2000).
- [9] T. J. Osborne and F. Verstraete, General monogamy inequality for bipartite qubit entanglement, *Phys. Rev. Lett.* **96** 220503 (2006).
- [10] J. S. Kim, A. Das, and B. C. Sanders, Entanglement monogamy of multipartite higher-dimensional quantum systems using convex-roof extend negativity, *Phys. Rev. A* **79** 012329 (2009).
- [11] Y. Ou, Violation of monogamy inequality for higher-dimensional objects, *Phys. Rev. A* **75** 034305 (2007).
- [12] Y.M. Yany, W. Chen, G. Li, Z.J. Zheng, Generalized monogamy inequalities and upper bounds of negativity for multiqubit systems, *Phys. Rev. A* **97** 012336 (2018).
- [13] P. Rungta, C. M. Caves, Concurrence-based entanglement measures for isotropic states, *Phys. Rev. A* **67** 012307 (2003).
- [14] X. J. Ren, W. Jiang, Entanglement monogamy inequality in a 2-2-4 system, *Phys. Rev. A* **81** 024305 (2010).
- [15] J. S. Kim and B. C. Sanders, Monogamy of multi-qubit entanglement using Rényi entropy, *J. Phys. A: Math. Theor.* **43** 445305 (2010).
- [16] Y. X. Wang, L. Z. Mu, V. Vedral, and H. Fan, Entanglement Rényi α -entropy, *Phys. Rev. A* **93** 022324 (2016).
- [17] Yu. Luo, and Y. M. Li, Hierarchical polygamy inequality for entanglement of Tsallis q -entropy, *Commun. Theor. Phys.* **69** 532 (2018).
- [18] A. Uhlmann, Fidelity and concurrence of conjugated states, *Phys. Rev. A* **62** 032307 (2000).
- [19] S. Alberverio and S. M. Fei, A note on invariants and entanglements, *J. Opt. B: Quantum Semiclass Opt.* **3** 223 (2001).

- [20] P. Rungta, V. Buzek, C. M. Caves, M. Hillery, and G. J. Milburn, Universal state inversion and concurrence in arbitrary dimensions, *Phys. Rev. A* **64** 042315 (2001).
- [21] X. N. Zhu and S. M. Fei, Entanglement monogamy relations of qubit systems, *Phys. Rev. A* **90** 024304 (2014).
- [22] Z. X. Jin, J. Li, T. Li and S. M. Fei, Tighter monogamy relations in multiqubit systems, *Phys. Rev A* **97** 032336 (2018).
- [23] X. J. Ren and W. Jiang, Entanglement monogamy inequality in a $2 \otimes 2 \otimes 4$ system, *Phys. Rev. A* **81** 024305 (2010).
- [24] J. B. Zhang, Z. X. Jin, S. M. Fei, and Z. X. Wang, Enhanced Monogamy Relations in Multiqubit Systems, *INT J. Theor. Phys.* **59** 2020.
- [25] A. Acin, A. Andrianov, L. Costa, E. Jané, J. I. Latorre, and R. Tarrach, Generalized Schmidt decomposition and classification of three-quantum-bit states, *Phys. Rev. Lett.* **85** 1560 (2000).
- [26] X. H. Gao and S. M. Fei, Estimation of concurrence for multipartite mixed states, *Eur. Phys. J. Spec. Topics* **159** 71 (2008).
- [27] C. H. Bennett, H. J. Bernstein, S. Popescu, and B. Schumacher, Concentrating partial entanglement by local operations, *Phys. Rev. A* **53** 2046 (1996).
- [28] C. H. Bennett, D. P. DiVincenzo, J. A. Smolin, and W. K. Wootters, Mixed-state entanglement and quantum error correction, *Phys. Rev. A* **54** 3824 (1996).
- [29] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, *Phys. Rev. Lett.* **80** 2245 (1998).
- [30] Y. K. Bai, N. Zhang, M. Y. Ye, and Z. D. Wang, Exploring multipartite quantum correlations with the square of quantum discord, *Phys. Rev. A* **88** 012123 (2013).
- [31] G. Vidal and R. F. Werner, Computable measure of entanglement, *Phys. Rev. A* **65** 032314 (2002).
- [32] S. Lee, D. P. Chi, S. D. Oh, and J. S. Kim, Convex-roof extended negativity as an entanglement measure for bipartite quantum systems, *Phys. Rev. A* **68** 062304 (2003).