Improving the Accuracy of the Variational Quantum Eigensolver for Molecular Systems by the Explicitly-Correlated Perturbative [2]_{R12} -Correction

Philipp Schleich,^{1,2,3,*} Jakob S. Kottmann,^{1,4,†} and Alán Aspuru-Guzik^{1,4,3,5,‡}

¹Department of Computer Science, University of Toronto, Canada.

²Applied and Computational Mathematics, Department of Mathematics, RWTH Aachen University, Aachen, Germany

³Vector Institute for Artificial Intelligence, Toronto, Canada.

⁴Chemical Physics Theory Group, Department of Chemistry, University of Toronto, Canada.

⁵Canadian Institute for Advanced Research (CIFAR) Lebovic Fellow, Toronto, Canada

(Dated: June 1, 2022)

We provide an integration of the universal, perturbative explicitly correlated $[2]_{R12}$ -correction in the context of the Variational Quantum Eigensolver (VQE). This approach is able to increase the accuracy of the underlying reference method significantly while requiring no additional quantum resources. The proposed approach only requires knowledge of the one- and two-particle reduced density matrices (RDMs) of the reference wavefunction; these can be measured after having reached convergence in the VQE. This computation comes at a cost that scales as the sixth power of the number of electrons. We explore the performance of the VQE+ $[2]_{R12}$ approach using both conventional Gaussian basis sets and our recently proposed directly determined pair-natural orbitals obtained by multiresolution analysis (MRA-PNOs). Both Gaussian orbital and PNOs are investigated as a potential set of complementary basis functions in the computation of $[2]_{R12}$. In particular the combination of MRA-PNOs with $[2]_{R12}$ has turned out to be very promising – persistently throughout our data, this allowed very accurate simulations at a quantum cost of a minimal basis set. Additionally, we found that the deployment of PNOs as complementary basis can greatly reduce the number of complementary basis functions that enter the computation of the correction at a complexity.

I. INTRODUCTION

Quantum computing is an emergent computational paradigm with the potential to disrupt some areas of scientific and financial computing. While recent advances in the development of the hardware bring the time of error-corrected quantum computers closer [1–3], current quantum computers, often called NISQ (noisy intermediate scale quantum computers), still do not offer practically useful computational advantage given their lack of noise resilience, short coherence times and a rather low, yet steadily increasing, available number of qubits [4].

Popular approaches to NISQ computing are constituted by hybrid quantum-classical procedures, which rely on the interplay of a classical and a quantum computer. One important class of such hybrid quantum-classical algorithms is given by variational quantum algorithms (VQAs) that rely on the definition of a parametric cost function, which is evaluated on a quantum computer, and then optimized numerically on a classical computer.[5, 6] In the context of quantum chemistry, particularly ground-state calculations, this cost function is set to be the energy expectation value of a molecular system. To make algorithms more suitable for NISQ hardware, its weaknesses on the hardware end are usually addressed by er-

ror mitigation techniques, more efficient mappings of the

underlying problem to the quantum computer's language or compilation strategies; see Refs. [6, 7] for an extensive overview. As an alternative, or in particular as an addition to these efforts, one can - just as in classical computing - exploit some physical knowledge to make the simulation more efficient. In this context, one aims to reduce quantum resources (in terms of qubits needed and gate count / circuit depth), usually coming at the cost of some additional computational efforts on a classical machine.[8-12] Efforts to do so include a priori and a posteriori computations, which either pre-modify the input for an existing quantum procedure or make use of the output to obtain an external correction, both with the ultimate goal to increase accuracy while maintaining the amount of quantum resources necessary. Conversely, this can be translated into a reduction of quantum resources for a certain level of accuracy.

In this work, we demonstrate a procedure that allows to reduce the number of qubits required for a certain quantum chemical quantum calculation involving electronic systems, that allow for an efficient output of the one- and two-particle reduced density matrices (1- and 2-RDM). This perturbative correction is based on methods usually summarized as "R12", "F12" or explicitly correlated methods, cf. section II A. Recently, *a priori* modifications inspired by Boys' and Handy's transcorrelated Hamiltonian [13, 14] – where a similarity-transformation of *H* renders $H' = e^{-G}He^{G}$ cusp-free given an appropriate *G* – have been adapted for the use in near-term quantum simulations. McArdle et. al. [15] apply Boys' and Handy's original formulation with a non-Hermitian *H'* within the imaginary time evolution

^{*} E-Mail:philipps@cs.toronto.edu

[†] E-mail:jakob.kottmann@utoronto.ca

[‡] E-mail:aspuru@utoronto.ca

ansatz [16], whereas Motta et. al. [10] implement a regularized Hamiltonian straightforwardly in a VQE simulation. The latter makes use of the canonical transcorrelation introduced in Ref. [17], which modifies the Hamiltonian as $H' = e^{A^{\dagger}}He^A$ with some anti-Hermitian A such that H' is still Hermitian; the Baker-Campbell-Hausdorff-expansion of H' is truncated so that only one- and two-body terms are kept.

On the contrary, we opt to use an *a posteriori* correction based on a perturbative *explicitly correlated* method, namely the spin-free variant of $[2]_{R12}$ [18–20]. This approach is to be contrasted with other post-corrections in form of the family of quantum subspace expansion (QSE) techniques [8, 9, 21]; an application of VQE together with the transcorrelated approach, $[2]_{R12}$ or a so called "CABS singles" correction [22] has already been suggested in Ref. [23]. Brief comments on the distinction of our approach with this one will follow in Section II A 2.

Aside of providing a framework for the application of the explicitly correlated $[2]_{R12}$ -method within the variational quantum eigensolver, the goal of this work is to raise additional awareness of explicitly correlated methods within the quantum computing community. This work builds on a recent Master's thesis [24] and contains updated data.

A comment on notation Within this work, we will make use of the Einstein summation convention, where the indices are clarified in Tab. I. Frequently we label orbitals only with their indices $\{p\} \equiv \{\phi_p\}$. For second-quantized creation and annihilation operators as well as matrix elements, we follow the notation of the works by Kutzelnigg and Mukherjee, which is prevalent in the literature of explicitly correlated methods (defined e.g. in Refs. [25, 26]). Excitation operators and matrix elements up to order two can be found in Tab. I as well. Higher-order excitations and matrix elements can be defined in a similar manner but are not needed in this context. If not stated otherwise, we make use of atomic units $\hbar = e = m_e = 1$ except for Ångstrom for the unit of length.

II. BACKGROUND

For the sake of self-containment, we will first give a highlevel description of explicitly correlated techniques and a historical overview before presenting and motivating the $[2]_{R12}$ correction in more detail. After that, we will propose a workflow to embed $[2]_{R12}$ into the hybrid quantum-classical variational quantum eigensolver routine.

A. Explicit Correlation

1. Basics

It is well-known that the wavefunction of an electronic system exhibits a *correlation cusp* at the points of inter-electronic

coalescence [27-30]. A generic numerical representation of such kinks in the functions with smooth functions imposes problems and leads to a rather high number of basis functions to represent the true wavefunction up to a certain accuracy. Yet it is rather obvious that an explicit incorporation of the inter-electronic distance $r_{12} = |r_1 - r_2|$ helps to overcome this slow convergence, as demonstrated for the simple case of the ground-state of Helium by Hylleraas [31], later again proposed [27, 32] and supported by rigorous work [30, 33, 34]. The essential take-away message can be formulated as follows: Upon occurrence of electronelectron correlation, the deployment of explicitly correlated basis function leads to more accurate energies, given that both the partial-wave expansion [30, 33] as well as a naturalorbital expansion [35] decay faster in such a basis (thus fewer basis functions needed for the same level of accuracy). For a comprehensive overview of explicitly correlated methods, we refer to the reviews [36-39].

However, making use of an explicitly correlated basis in the existing methods (such as CI, leading to so called Hylleraas CI [40]) does not allow an efficient implementation due to the appearance of very high-dimensional integrals [36, 41]. One of the two most prominent ways to exploit explicit correlation is given by perturbative descendants of Kutzelniggs R12theory [30, 42–45], where MP2-R12 can be seen as the basis for all upcoming, more sophisticated, perturbation methods such as $[2]_{R12}$ or the approach in Ref. [46]. The other one consists of regularizing the system throughout a similaritytransformed Hamiltonian via Boys and Handy's transcorrelated approach [13, 47], made more applicable throughout approximations such that the effective Hamiltonian is still Hermitian in Refs. [17, 48]. In a recent benchmark [49], both the perturbative $[2]_{R12}$ -correction, that we will explain in detail in section IIA2 and the canonical transcorrelated approach from Ref. [17] performed comparably well. Here, we focus on the universal perturbative $[2]_{R12}$ method and will provide a discussion to distinguish the approach from the transcorrelated Hamiltonian when applied to quantum algorithms.

Initially, the explicitly correlated basis has been built by correlating one-electron basis functions, achieved through multiplication with the inter-electronic distance r_{12} . Motivated by that fact, that a simple factor of r_{12} does not inherit the correct asymptotics, Gaussian correlation factors $\exp\{-r_{12}^2\}$ (*Gaussian geminals* [50, 51]) and Slater-type correlation factors $\exp\{-r_{12}\}$ [42, 52] have been introduced. Within a study that investigated the performance of different correlation factors, it was found that Slater-type factors often have the most favorable behaviour [53]. Hence following the [2]_{R12} -literature, we use correlation factors and its matrix elements given as follows:

$$f_{12} = -\frac{1}{\gamma} e^{-\gamma r_{12}}, \quad r_{\mu\nu}^{\kappa\lambda} = \langle \mu\nu | f_{12} | \kappa\lambda \rangle.$$
(1)

Note that when using Psi4 [54] in our simulations, the correlation factor f_{12} is represented as a linear combination of

Gaussians.

2. $[2]_{R12}$ -Correction

Further on, we provide a brief outline to $[2]_{R12}$ as in Refs. [18–20], recalling that this is a perturbative explicitly correlated method.

Let \mathcal{H} be the Hilbert space of the true wavefunction of a certain electronic system with N_{el} electrons in Born-Oppenheimer approximation, described by its Hamiltonian operator

$$H = h_p^q a_q^p + \frac{1}{2} g_{pq}^{rs} a_{rs}^{pq},$$
(2)

where h_q^p denotes one-electron integrals and g_{rs}^{pq} twoelectron integrals over the Coulomb interaction:

$$g_{rs}^{pq} = \langle rs | (r_{12})^{-1} | pq \rangle.$$
(3)

Further we have a *reference* solution $|\Phi_{ref}\rangle$ using an arbitrary method to find an approximate solution to the ground-state problem for H using a certain basis consisting of a finite number of one-electron orbitals $\{p\}_{p=1}^{N_{OBS}}$ that span $\mathcal{P} \subset \mathcal{H}$. The perturbative correction we use here aims to correct for the incompleteness of this one-electron basis by excitations into the complementary space $Q = \mathcal{H} \setminus \mathcal{P}$. Note that formally, up to this point, the procedure is equivalent to the Quantum Subspace Expansion [8, 9, 21]. Yet here, the basis for the complementary space is enforced to be explicitly dependent on the inter-electronic coordinate and thus is deemed to exhibit faster convergence; it is tailored to account specifically for completeness due to the correlation cusp. Conversely, we expect that for systems where the dominant error originates from insufficient representation of the correlation cusp, the present method needs less basis functions for a qualitative representation of the complementary space than QSE-based procedures with a primitive basis and is thus likely to be more efficient. On the other hand, the application of QSE is not limited to quantum chemistry [21]. It is worth mentioning that all perturbative methods relying on the RDMs require the evaluation of complex equations (e.g. via the generalized Wick's theorem [25]), as well as a set of non-trivial cumulant approximations to avoid measurement of higher-order RDMs [8, 18, 55, 56]. While the QSE depends on the solution of a generalized eigenvalue problem, $[2]_{R12}$ amounts in the sp-ansatz [57], which fixes amplitudes so that they fulfill the cusp condition as shown in Eq. (8), to a set of tensor contractions.

The $[2]_{R12}$ correction then can formally be derived [62] using Löwdin partitioning [18, 63–65]. Then, one can approximate the full complementary space $Q = \text{span}\{\alpha\}$ using a complementary auxiliary basis set (CABS) $\{\alpha'\}_{\alpha'=1}^{N_{\text{CABS}}}$ with finite N_{CABS} – this CABS basis can be an arbitrary basis of functions and does not need to come from a Gaussian basis set,

Table I: Glossary (operator and index conventions, acronyms sorted by subject matter)

Indices $(\{p\} \equiv \{\phi_p\}, p\rangle \equiv \phi_p\rangle)$	Explanation
$\frac{1}{\{p,q,r,s\}}$	Orbital basis set (OBS) for or-
	bital space \mathcal{P} / reference
$\{i, j\}$	Occupied within OBS
$\{a,b\}$	Unoccupied within OBS
$\{\kappa,\lambda\}\rightsquigarrow\{\kappa',\lambda'\}$	Formally complete basis and finite RI basis to approximate full Hilbert space \mathcal{H} with RI space
$\{\alpha, \beta\} \rightsquigarrow \{\alpha', \beta'\}$	Complementary basis and finite complementary auxiliary basis (CABS) to resolve $Q = \mathcal{H} \setminus \mathcal{P}$
Operators and integrals	Following e.g. [26]
$a^p = a_p^{\dagger}, a_p$	Fermionic creation and annihi- lation operators
$a_q^p = a^p a_q, a_{rs}^{pq} = a^p a^q a_s a_r, \dots$	Particle-number conserving ex- citation operators
$X_q^p = \langle q(1) X(1) p(1) \rangle$	One-electron integrals
$X_{rs}^{pq} = \langle r(1)s(2) X(1,2) p(1)q(2) \rangle$	Two-electron integrals
g ^{pq} grs	Coulomb integrals, cf. Eq. (3)
r_{rs}^{pq}	F12 integrals, cf. Eq. (1)
$\gamma_q^p, \gamma_{rs}^{pq}$	One- and two-particle reduced density matrices (1- and 2-RDM), cf. Eq. (9)
f_{λ}^{κ}	Generalized Fock operator, cf. Eq. (5)
Acronyms	Explanation and Reference
NISQ	Noisy intermediate-scale quan- tum (computing) [4, 6]
VQA/VQE	Variational Quantum Algo- rithms/Eigensolver [5, 58]
PNO	Pair-natural orbitals [11, 59]
MRA	Multiresolution analysis [60]
RDM	Reduced density matrix
QSE	Quantum subspace expansion [8, 9, 21]
OBS	Orbital basis set [61]
CBS	Complete basis set [61]
CABS	Complementary auxiliary basis set [61]
	300 [01]

even if the OBS is built from Gaussians. Note that here, this approximation is formally equivalent to the resolution of the identity approximation in the evaluation of molecular integrals of usual R12/F12 methods (usually abbreviated by RI, insertion of $I = |\kappa\rangle \langle \kappa| \approx |\kappa'\rangle \langle \kappa'|$ to avoid higher-order integrals). We construct the RI basis as $\{\kappa'\} = \{p\} \cup \{\alpha'\}$, i.e., as the union of the orbitals $\{p\}$ resulting from the regular orbital basis and $\{\alpha'\}$ coming from the complementary auxiliary basis. The partitioning into a reference $|\Phi_{\text{ref}}\rangle \in \mathcal{P}$ and a perturbation (to account for incompleteness due to Q) then leads to a correction given by the minimization of a

second-order Hylleraas functional

$$J_{H}^{(2)} = \langle \psi^{(1)} | H^{(0)} | \psi^{(1)} \rangle + 2 \langle \psi^{(0)} | H^{(1)} | \psi^{(1)} \rangle, \quad (4)$$

named after Hylleraas' general approach to perturbation theory [31]. The reference Hamiltonian corresponds to the model Hamiltonian $H^{(0)} = H$, and as zeroth-order state, the solution obtained by the reference method $|\psi^{(0)}\rangle = |\Phi_{ref}\rangle$ is picked. Further, the first-order Hamiltonian is chosen to be a normalized, generalized Fock operator

$$H^{(1)} = F = f^{\kappa}_{\lambda} a^{\lambda}_{\kappa} - E_0 = (h^{\kappa}_{\lambda} + \bar{g}^{\lambda r}_{\kappa s} \gamma^s_r) a^{\lambda}_{\kappa} - E_0, \quad (5)$$

while the appearing 1-RDM corresponds to the reference method (not HF as in the standard Fock operator) [26]. The bar over the Coulomb-tensor denotes antisymmetrization. This induces maximal resemblance to the MP2-R12 method [18]. Explicit correlation is brought into play with the first-order perturbation

$$|\psi^{(1)}\rangle = QR |\Phi_{\rm ref}\rangle, \qquad (6)$$

where Q denotes a projector ensuring that contributions of OBS are projected out as well as only semi-internal firstorder excitations are kept (which was found to be important in contrast to Ref. [46]).[18, 19, 36] Further excitations from occupied to CABS orbitals are excluded, being the origin of one-particle incompleteness, which are to be treated by a "CABS Singles" correction as in Ref. [22]. The correlation operator R is defined here as

$$R = \frac{1}{2} d_{pq}^{rs} r_{\kappa\lambda}^{pq} a_{rs}^{\kappa\lambda}$$
(7)

and generates two-particle correlation, performing excitations into the complementary space Q, which is represented by a set of explicitly correlated functions. Using a set of approximations[66], one could then seek to minimize the Hylleraas functional; this would typically result in an illconditioned linear system [36], which usually is avoided by making use of the sp-Ansatz by Ten-No [57]. In this formulation, the amplitudes d_{ps}^{rs} are fixed such that the cusp conditions both for singlet (~s) and triplet (~P) states are fulfilled. Using spin-free orbitals, the sp-ansatz reads

$$d_{pq}^{rs} = \frac{3}{8}\delta_p^r \delta_q^s + \frac{1}{8}\delta_q^r \delta_p^s.$$
 (8)

This way, $[2]_{R12}$ amounts to a set of tensor contractions, whose evaluation scales as $O(N_{el}^6)$ assuming N_{OBS} , $N_{CABS} \in O(N_{el})$ within the framework of the so called approximation C, introduced in Ref. [67] and applied to $[2]_{R12}$ in Ref. [20]. Instead of restating the rather complex equations, we refer to the original works [18–20]. We further point out that a reduction of the cost can be achieved by correlating only a subset of the reference orbitals that may be chosen according to their occupation number [19, 20], as well as by following ideas from Ref. [68] to lower the order of dependency on the RI dimension. Diagonalization of the 1-RDM to obtain these comes at almost no cost, since they need to be available anyway. The framework of approximation C of $[2]_{R12}$ requires evaluation of additional molecular integrals, as well as only the 1- and 2-RDM with respect to the given reference

$$\gamma_q^p = \langle \Phi_{\rm ref} | a_q^p | \Phi_{\rm ref} \rangle, \quad \gamma_{rs}^{pq} = \langle \Phi_{\rm ref} | a_{rs}^{pq} | \Phi_{\rm ref} \rangle. \tag{9}$$

Insofar, this correction can be applied to universal references, as long as the 1- and 2-RDM can be made available. Here, we exclusively use the spin-free formulation, called SF- $[2]_{R12}$. Consequently, excitations, RDMs etc. are to be understood as spin-free obtained by summation over spin – for ease of notation, we treat this implicitly. A more detailed interpretation of $[2]_{R12}$ can be found in Ref. [24].

We note that the application of the correction is not necessarily restricted to ground state calculations but can be applied to excited states in a similar manner.

B. VQE and VQE + [2]_{R12}

In this work, we rely on the Variational Quantum Eigensolver [58, 69] that can be assigned to the class of variational quantum algorithms, which is one of the most promising classes of algorithms in the NISQ era of quantum computing [5, 6, 23, 70]. Here, one defines an objective function in form of the expected value of the Hamiltonian operator with respect to a parametrized quantum state. The quantum state is composed by a sequence of parametrized unitary operations in form of quantum gates (often called ansatz). Minimization of the objective function yields an approximation to the ground-state energy of the system; the accuracy of this approach greatly depends on the capability of the ansatz to express the true ground state [71–77].

This way, we obtain a reference for the perturbative approach in form of $|\Phi_{ref}\rangle = U(\theta^*) |0\rangle$ with

$$\theta^{\star} = \arg\min_{\theta} \left\langle 0 | U^{\dagger}(\theta) H U(\theta) | 0 \right\rangle \tag{10}$$

and $U(\theta)$ a sequence of parametrized quantum gates. Throughout this work, we will restrict ourselves on Unitary Coupled Cluster-type ansätze (UCC) [11, 58, 71, 72, 78, 79]. Given a basis set {*p*}, a generic UCC-type wavefunction can be obtained by

$$|\psi_{\rm UCC}(\theta)\rangle = e^{\hat{T}-\hat{T}^{\dagger}} |\Phi_{\rm HF}\rangle,$$
 (11)

where $\hat{T} = t_a^i a_i^a + \frac{1}{4} t_{ab}^{ij} a_{ij}^{ab}$ is the usual cluster operator. The amplitudes t_a^i, t_{ab}^{ij} serve as variational parameters in this approach. In practice, a wide range of approximations is raised in order to implement or approximate such a wavefunction [72].

In Fig. 1, we summarized the proposed workflow of a VQE+ $[2]_{R12}$ computation. Upon having chosen the molecule of desire along with a choice of orbital basis and

CABS basis, one can build the fermionic Hamiltonian H_{ferm} in the orbital basis as well as molecular integrals needed for the F12-correction (see Ref. [19, 20]) in the RI-basis. Additionally, one needs to provide a F12-exponent γ , cf. Eq. (1). Within this work, we fixed $\gamma = 1.4$, which overall led to good results. When aiming for the best energy possible, one might perform computations with a set of values and then pick the lowest one if the CBS-limit is not available. In general we found that $[2]_{R12}$ is not very likely to yield energies below the CBS-limit despite it being a non-variational method. Next, one needs to map the fermionic Hamiltonian to a set of qubit operators using an operator mapping. Here, we used Jordan-Wigner, but other mappings can be employed in a similar manner; see e.g. Refs. [6, 16, 23]. The qubit Hamiltonian $H_{\rm qc}$ then serves as input for the VQE computation. After VQE has reached convergence, the 1- and 2-RDM need to be measured, which then are processed together with the molecular integrals in the RI basis to a scalar correction in the energy.

To obtain the necessary 1- and 2-RDM, the VOE measurement procedure needs to be adapted to resolving the full reduced density matrices, coming at a cost $O(N_{OBS}^4)$ for the 2-RDM in the worst case (symmetry conditions and spin-free formulation only allow for a reduction by a constant factor). A wide set of measurement reduction techniques has been developed for the evaluation of Hamiltonian expectation values. These dominated by two classes of such techniques: One relies on the identification of sets of commuting operators that can be measured simultaneously, e.g. Refs. [80–82], which allow for an approximately cubic cost. The other one determines basis rotations that result in a formulation with less non-mutually commutative terms, leading to less than quadratic number of terms that need to be measured separately [83, 84]. A comprehensive overview of advances in this area can be found in Ref. [85]. However, as pointed out in Ref. [85], most of these methods are not directly applicable for the measurement of general fermionic k-RDMs. Using a basis of Majorana operators, Ref. [85] shows that smart operator partitioning can yield a reduction of measurement cost up to $\Omega(N_{OBS}^2)$. An advisable strategy for a VQE+ $[2]_{R12}$ routine then would be to implement one of the measurement reduction techniques that reliably can suppress the cost to lower than quadratic order, and make use of the strategy outlined in Ref. [85] for the final RDM-evaluation. An additional set of interesting approaches improving the measurement of RDMs based on the n-representability conditions is given in Ref. [86]. The efficacy of these approaches has been demonstrated in Ref. [87]. Such techniques are in particular important in light of stochastic errors when sampling the RDMs. As it was investigated in Ref. [88] who applied $[2]_{R12}$ in FCIQMC calculations, sampling noise in the RDMs spreads as well to the correction; such behaviour can be expected to appear as well when extracting RDMs from VQE and thus high fidelity in doing so at moderate cost is desirable.

A. Computational Setup

1. Construction of Complementary Basis

Within this work, we follow the CABS+ procedure introduced in Ref. [61] to generate a complementary basis. This means, given an orbital (reference) basis and complementary basis, the complementary space is specified by orthogonalization of the joint RI basis { κ' }, obtained by the union of orbital and complementary basis, and projecting out the orbital space. Formally, one constructs a coefficient matrix $C_{\alpha'\kappa'}$, which maps from the RI to the complementary set, such that $\sum_{\kappa'} C_{\alpha'\kappa'} \langle \kappa' | p \rangle = 0 \forall p$.

In contrast to that, we comment on the possibility of a "quick & dirty"-approach via the choice of an active space. Here, one starts with a given basis set, that is employed in an active space fashion. That is, there is a set of frozen virtual orbitals which do not contribute to electron correlation effects. Consequently, we find ourselves in the same setting as the work by Takeshita et. al. [9] who make use of QSE to account for the "active space incompleteness". However, we do not present results for this situation since we found, in accordance to Valeev's arguments [61], that corrections obtained by CABS+ are more accurate and are not sensitive to the choice of active space. Still this does depend on the specific intention behind the computation – if a valid active space is chosen, this procedure allows in the case of the virtual QSE to retrieve the expressibility of the full orbital basis [89], and a similar behaviour can be expected for $[2]_{R12}$.

Accordingly it is advised to follow the CABS+ procedure, as done in Ref. [10]. But, if one were to perform an active space calculation (and an appropriate set of orbitals is chosen), the application of $[2]_{R12}$ is expected to be beneficial. While we have no numerical evidence to support this, we expect $[2]_{R12}$ to perform better than similar methods that excite into conventional basis functions, such as the subspace expansion-based approach in Ref. [9], whenever electron-electron cusp effects are dominant, re-emphasizing that the F12 method constructs an explicitly correlated basis that is expected to "converge faster" and hence should require fewer basis functions for a given energy correction.

As basis functions, we will mostly resort to the recently introduced MRA-PNOs (globally orthonormalized, directly determined pair natural orbitals obtained by multiresolution analysis) [11, 59]. This enables us to, on one hand, build up on the existing data thereof, and on the other hand, investigate the potential of a combination of PNOs together with a F12method. Even the most simple MRA-PNO generation at this time generates at least $4N_{diag} + 8N_{offdiag}$ orbitals given a number of diagonal and off-diagonal pairs, which means that in most cases there are a few unused PNOs. This raises the question whether PNOs are suited for a deployment as CABS-basis. Building on the theory, it is to be expected that




Figure 1: Workflow of VQE+ $[2]_{R12}$. Modules marked by green colored can be performed on a quantum processing unit or a classical simulator, while orange modules denote classical pre- and post-operations. Used software packages are marked in the above illustration, and reiterated in section III A 3.

• output: $E_0 + \Delta E^{[2]_{R12}}$

TEQUILA + QULACS OF QPU

explicitly correlated methods also deliver faster convergence in terms of natural orbitals [35], which means we would expect that PNOs built from a MP2-R12 surrogate, as already done in Ref. [59], inherit this behaviour. Such behaviour was also found in Refs. [90, 91]. Further, a basis of natural orbitals can be seen as a least-squares approximation [36, 92] and we again expect the PNOs to be somewhat near-optimal despite the pair-approximation. This means that PNOs in general and in particular MP2-F12 PNOs might be interesting candidates. To that end, we provide some data from Ref. [24] and additional discussion in appendix **B**. Based on these results and the discussion, hereinafter, we will pair VQEresults based on MRA-PNOs for the reference with a PNO-CABS (all with plain MP2 for the surrogate computation), given that they have turned out to outperform PNOs obtained by MP2-F12 if obtained in a self-contained procedure and are also far more compact as an RI-approximation than Gaussian basis functions (The reason herefore is that the reference computation when using PNOs generated by R12 performs worse. While this is expected, we found that in our case, this is so significant that the correction does not fully account for that, although the correction is higher; also see appendix **B**). We emphasize here however that our RI by PNOs is not necessarily perfectly accurate but rather more compact and accurate than a GBS of a comparable size. Additionally the point we want to make is that any PNOs generated that are not used in OBS can increase accuracy when used for the complementary basis.

 H_{qc}

Of course any set of basis functions - be it the most prominent choice of orbitals from Gaussian basis sets (GBS), PNOs from GBS or plane-waves – can be used to apply $[2]_{R12}$. As already mentioned, we employ MRA-PNOs here to be able to compare with previous results in Ref. [11]. Further, as specified therein and in Ref. [93], the MRA-PNOs yield surprisingly good results in bond-breaking regimes despite the MP2 dependency. Additionally, in contrast to Gaussian basis sets, they exhibit faster convergence so that even for small basis set sizes, the dominant error can already be expected due to the cusp. In section III A 3 we will point to open-source, freely available software that we employed and wherein we implemented the computation of the correction; both MRA-PNOs and GBS are ready to be employed.

TEOUILA

Choice of Ansatz in VQE-Calculations 2.

In the subsequent results, we use a classical FCI calculation for systems with Gaussian basis sets (GBS); for the PNObasis, we focus on UCC-type ansätze. However, the use of a specialized F12-basis set in the case of GBS is recommended, as this enables the reference computation to focus on the representation of the cusp-free part of the wavefunction and the correction takes care of the cusp. This work can be understood as an extension to Ref. [11] and thus we aim to show the potential of MRA-PNOs together with $[2]_{R12}$. Consequently the results we show mostly deploy MRA-PNOs as a basis but we showcase a few exemplary simulations using FCI/GBS as a verification that general bases can be made use of.

For all MRA-PNO-based computations, we exploit the pairstructure in form of the SPA-ansatz from Ref. [93], which is repeated here in brevity; also see Ref. [72] for a description thereof. Given a set of HF reference orbitals $\{i\}$, for each (ij)-pair we generate orthonormalized PNOs $\tilde{\mathcal{S}}_{ij} = \bigcup_{\tilde{a}_{ij}=1}^{r_{ij}} \{ |\tilde{a}_{ij} \rangle \}$. As an orthonormalization procedure, we use a Cholesky decomposition, which has turned out deliver superior results as opposed to e.g. symmetric orthonormalization because it changes the "most important" PNOs, ordered by occupation number, the least. Then, we restrict to pair-excitations from the references to the associated PNOs $(U_{\tilde{D}})$ and generalized excitations within each PNO $(U_{G\tilde{D}})$. One might also think of single excitations, which we do not consider in this work. We can express the SPA-UpCCGD

wavefunction as

$$|\text{SPA-UpCCGD}\rangle = U_{\text{GD}}U_{\tilde{\text{D}}}U_{\text{HF}}|0\rangle$$
 (12)

where $|\Phi_0\rangle = |\Phi_{\rm HF}\rangle = U_{\rm HF} |0\rangle$ and

$$U_{\tilde{\mathbf{D}}} = \prod_{i=1}^{N_{el/2}} \prod_{a \in \tilde{\mathcal{S}}_{ii}} \exp\left\{\frac{\theta}{2}\tilde{G}_{ii}^{aa}\right\}$$
(13)

$$U_{\rm G\tilde{D}} = \prod_{i=1}^{N_{\rm el/2}} \prod_{a,b\in\tilde{\mathcal{S}}_{ii}} \exp\left\{\frac{\theta}{2}\tilde{G}_{bb}^{aa}\right\}.$$
 (14)

The pair-excitation generator \tilde{G} is defined as

$$\tilde{G}_{ii}^{aa} = a_{i\uparrow i\downarrow}^{a\uparrow a\downarrow} - \text{h.c.}$$
(15)

In what follows, whenever we use PNOs as a basis, we will employ the SPA-UpCCD ansatz and use "SPA" to denote this. Ref. [11] demonstrated that whenever the PNO approximation is a good representation of the molecular system behind, this PNO-restricted parametrization provides potential for a drastic reduction in number of parameters and CNOT-counts as opposed to a UpCCGSD-wavefunction, which itself is compact in comparison to a full UCCSDparametrization [73]. More detailed demonstrations of the performance of SPA can be found in Ref. [93], aiding to classify it in comparison to popular classical methods.

Further we provide results for a *cheap* and *good* set of PNOs obtained by MRA-PNO computations. The cheap set signifies a minimal version of PNOs, with the least rich class of excitations from Ref. [59] ("dipole+") that uses only the spare orbitals from the generation which one does not opt to use for the orbital space. In this sense, this procedure would be equivalent to using an active space of some GBS and exploiting the spare orbitals in a correction scheme. Apart from that, the good set of PNOs allows ten macro-iterations in their generation - convergence in the PNO-generation usually is achieved beforehand when a certain maximum number of orbitals is specified - and uses a richer excitation ansatz ("multipole") with a prescribed goal of the number of PNOs. We set this number accordingly such that $[2]_{R12}$ converges, which for our purposes can be seen as a measure that the identity is resolved well enough. This method then would be somewhat analogous to adding a RI-optimized CABS to some orbital basis set.

3. Software

We have outlined the software packages that have been used in Fig. 1. The leading framework organizing the workflow for the work presented here is TEQUILA [78], while we rely on the methodology in Ref. [79] to obtain analytical gradients at the cost of only two energy expectation values, independent of excitation rank. As a quantum chemistry backend for classical CI calculations and the construction of a CABS made up by Gaussian orbitals, we use Psi4 [54]. To compute MRA-PNOs, we employed the software package MADNESS [60]. For the CABSfunctionality, our implementation uses modified forks of Psi4 and MADNESS [94]. F12-integrals in Eq. (1) from MADNESS are computed directly as Slater-type ($e^{-\gamma r_{12}}$), when obtained from Psi4 via GBS they are approximated by a linear combination of Gaussians.

Further dependencies include OPENFERMION [95] for the handling of fermionic and qubit operations as well as QULACS [96] as quantum computing backend. Further, automatic differentiation is taken care of by JAX [97]. SCIPY [98] is used for the classical optimization procedures (BFGS). In section A, there is a code sample for the computation of the

correction, both in the case of Gaussian orbitals and MRA-PNOs. Further, we made available a tutorial notebook on the matter on the github repository for TEQUILA [78].

B. Atomic systems

We start by looking at the atomic systems Helium ($N_{\rm el} = 2$) and Beryllium ($N_{\rm el} = 4$) and consider a choice of different bases to obtain an intuition as how well MRA-PNOs+[2]_{R12} perform against a naive procedure with a GBS. This should give us an indication regarding possible savings of quantum resources.

Except for the corrected VQE/MRA-energies, Fig. 2 is identical to Fig. 2 in [11]. Compared to the uncorrected energy, $[2]_{R12}$ provides significantly lower energies, which slightly "overshoot" but still within a chemical accuracy of 1.6 mEh, for the simulations with ten and twelve qubits (we denote number of qubits as N_q). This showcases the non-variational nature of the perturbation method. In the case of Beryllium this is a behaviour that carries through the results in particular for $N_{\rm el} > 2$, although in general no assumptions can be made that this needs to be true - the calculations with 10 qubits are equivalently accurate as the one with 24 qubits. Note that as anticipated in Ref. [11], the result for Beryllium "saturates" at a certain level. This way, the combination of MRA-PNOs as a system-adapted basis and the $[2]_{R12}$ -correction enables a significant reduction in the necessary number of qubits for both Helium and Beryllium.

C. Molecules

Again the shown results here widely follow Ref. [11] to demonstrate the potential of a combination of $[2]_{R12}$ with MRA-PNOs.

In the following, we present potential energy surfaces for the dissociation of H_2 , LiH, BH and BeH₂ (symmetric) in Fig. 3. Recall from section III A 2 that we mostly present outcomes using a MRA/PNO-basis with a distinction in *cheap* and *good*




Figure 2: Atomic systems. Comparison of classical FCI with corrected basis-set-free VQE+[2]_{R12} energies using *good* PNOs. We use a SPA-UpCCD ansatz for VQE [93]. Data for classical computations is from [11]. Note that the corrected results of He are in parts lower than FCI and thus show the non-variational nature of the perturbative correction.

PNOs, i.e., using the left-over, unused orbitals vs. creating a CABS on purpose. In addition, Fig. 4 shows potential energy surfaces for H_2 and LiH in Gaussian basis sets with a specific CABS basis. Moving on, we will discuss the results shown in the figures.

 H_2 First, look at the hydrogen molecule with $N_{\rm el} = 2$ in Figs. 3a,3b,3c. For the cheap PNOs, only 5 PNOs are available, which leaves only one orbital for the CABS in the 4-qubit computation. Hence, the 8 qubit computation (with 4 spatial orbitals) is left with only one PNO to represent the complementary space. As expected, the correction is very small in this case, however it still makes sense to apply it given it only improves the result. Given that there is only one electron pair in this case, the PNO generation can be seen as quite efficient, and the correction for the *cheap* set is rather low. On the other hand, the 8 qubit computation using good PNOs almost reaches the accuracy of the 120 qubit FCIcalculation in the regime of small bond distances. We note that for longer distances, the correction tends towards zero because in case of two basically separated hydrogen atoms, there is no more electron-electron correlation present. Further we point out an additional anomaly in Fig. 3a: At a distance of 3 Ångstrom, the 8 qubit result should lie lower (roughly -1 Eh) - this is due to the MP2 surrogate model for the PNO generation, which has not been able to produce orbitals of the correct symmetry in this case (for more details, see Ref. [24]).

LiH The previous findings mostly can be confirmed by looking at LiH in Figs. 3d,3e,3f. Note that since there are two electron pairs now, there would be 8 PNOs left to represent the CABS in the larger OBS for the *cheap* PNOs. However, the associated correction almost vanished since all these orbitals stem from an off-diagonal pair – since the LiH wavefunction very well separates in a product structure of pair one and two, these orbitals barely contribute, also visible by their low occupation number. The *good* PNOs allow for a considerable correction, with the corrected energy of both the 12 and 20 qubit computation at a similar level. Within the dissociation curve for *cheap* PNOs, the energies for both 12 and 20 qubits at -0.6 Å are far too high – this anomaly has the same origin as the 3.0 Å-point for H₂, and can again be resolved by allowing for another macro-iteration in the orbital determination.

In contrast to diatomic hydrogen, the corrections for systems containing atoms with more than two electrons do not vanish for long bond distances because there is still electron-electron correlation embodied in the individual atoms.

Furthermore, we observe that for both sets of PNOs, the lower-qubit computation reaches the same level as the computation with more qubits. This behaviour shows again in the case of BH.

BH Considering Figs. 3g,3h,3i, we additionally observe that while the uncorrected energies with both 12 and 22 qubits are very close (they differ by roughly 0.01 *E h*, the corrected energies are almost equal, and except for around the equilibrium differ only in the order of milli-Hartrees, where the fewer qubits computation actually performs better because of the higher-dimensional CABS. Additionally, the results for *cheap* and *good* PNOs are much closer than before. This rationale behind this is that with $N_{el} = 6$, with three diagonal and three off-diagonal pairs, there are $4N_{diag} + 8N_{offdiag} = 36$ PNOs readily generated within one macro-iteration, and given they do not differ too much in "quality", this already yields a decent RI-representation. For higher bond distance, this almost levels out, while for lower distances with stronger electron-electron correlation, the larger CABS still pays off.

BeH₂ Results for BeH₂ are shown in Figs. $3j_3k_3l_3$; these have been obtained in a frozen-core approximation with four active reference orbitals. We point out that for larger bond distances, the PNO-approximation in the construction of the quantum circuit does not hold anymore as good. This was already noticed in [11] and overcome using an adaptively

enriched parametrization in the style of ADAPT-VQE [75]. Here, we abstain from doing so and investigate the impact of the correction in such a regime. We see that – as expected – the explicitly correlated correction is not able to account for weak properties of the underlying ansatz.

The last set of potential energy surfaces we look at are comprised by the PES for the hydrogen molecule and lithium hydride in Gaussian basis sets in Fig. 4. They show potential energy surfaces for H₂ and LiH in STO-3G and 6-31G with a specific CABS basis generated by CABS+ [61] to demonstrate that Gaussian basis sets with designated CABS can be used in the same manner. Here, we do not observe any anomalies with respect to the results before but see only consistent behaviour, e.g. the vanishing correction for large bond distances in the case of H₂.

D. NPE + MAX

As a final result, we consider the non-parallelity error (NPE) and maximum error (MAX) defined over the computed PES for the cases of H₂, LiH and BH, following [11]. We compare MRA-PNOs with and without corrections with a corrected DZ-basis (with cc-pVDZ-F12-OPTRI as CABS). We do not use a F12-optimized cc-pVDZ-F12 for the orbital basis to keep the number of qubits closer to the MRA-PNO computations. With $\Delta E(R) = |E(R) - E_{ref}^{FCI}(R)|$, these error metrics are defined as

$$\Delta E_{\text{NPE}} = \max_{R \in \text{PES}} \Delta E(R) - \min_{R \in \text{PES}} \Delta E(R)$$
(16)

$$\Delta E_{\text{MAX}} = \max_{R \in \text{PES}} \Delta E(R). \tag{17}$$

Looking at Fig. 5, we can say that in particular the MAX error is consistently and significantly reduced. Since the explicitly correlated correction is not a constant shift but typically higher for lower bond distances and thus higher electronelectron correlation, the NPE error is not necessarily reduced but also does not increase by much in our computations. This holds in particular for the smaller systems because this effect carries more weight here. Occasional increase in NPE can be traced to higher correction for short bond distances, which holds in particular for smaller systems (compare to discussion of H₂ in section III C). This effect is is less severe for larger systems, as witnessed by BH in Fig. 5. For C₂, Ref. [102] observed a reduction in NPE as well.

IV. CONCLUSION AND OUTLOOK

First and foremost, we have proposed a workflow for the combination of the explicitly correlated $[2]_{R12}$ -correction together with VQE, and provided the software infrastructure to carry out such computations. Based on a few small test systems, we were able to show a significant increase in accuracy when using this correction. Although $[2]_{R12}$ is

a non-variational perturbation method, within out experiments, we only noticed few and low overshoots beyond FCI as long as the reference method produced a somewhat sensible approximation. For non-sensical or very bad reference inputs, the behaviour of the correction also seemed unpredictable. In particular in combination with orthogonalized MRA-PNOs [11, 59] employed in the reference method as well as complementary basis shows considerable promise, and allows to perform the quantum routine at a cost of a minimal basis yielding accuracies comparable to large correlation-consistent basis sets currently unfeasible for quantum algorithms. Of course, PNOs generated by a set of Gaussian orbitals can be considered equivalently. To ease the combination of PNOs with a complementary basis composed of Gaussian basis functions, an interface of the MRA part of MADNESS [60] with e.g. MPQC [103] can be envisioned.

Additionally, exploiting the PNO-structure for the parametrized quantum circuit turned out to be a powerful tool to reduce parameters within the classical optimization subroutine of VQE. For larger systems such as BeH₂ however, this parametrization turned out to loose validity – ways to overcome this are proposed in Ref. [11]. Here we used the separable pair ansatz [93] as an affordable way to construct quantum circuits, illustrating also that $[2]_{R12}$ can not overcome fundamental limitations in the ansatz.

We divided our MRA-PNO computations into a set of *cheap* and good PNOs, based on the quality of the generated PNOs. We note that for small systems, the *cheap* set often does not yield a large enough number of extra PNOs to induce a significant impact, larger systems with more than 1-2 pairs quickly generate a quite rich set of PNOs (see correction for BH in Fig. 3i, where the corrections for both sets of PNOs do not differ much). Further, known physical behaviour has to be kept in mind and monitored when growing PNOs, such as the pair structure of LiH (off-diagonal PNOs barely contribute) or critical geometries along the potential energy surface that tend to degeneracies (compare to the cheap sets of H₂ and LiH in Figs. 3a,3d.). Yet in any case, whenever the reference PNOs are sufficiently good, adding remaining left-over orbitals throughout $[2]_{R12}$ in the spirit of Ref. [9] is beneficial. Beyond that, it would be interesting to consider the constructing geminal-spanning orbitals from Ref. [104] instead of MP2-PNOs using MRA.

While our findings imply a reduction in the number of qubits at the same level of accuracy, due to its perturbative nature we cannot give projections regarding circuit depth, number of entangling gates, or other cost measures. Yet in comparison to the transcorrelated approach [10], the Hamiltonians using a perturbation method remain unchanged and do not exhibit an increased number of terms. On the other hand, the perturbation method does not allow for the same level of flexibility as a regularized Hamiltonian. In the end, the choice between these two approaches depends on a case-by-case basis and might even boil down to a matter of taste. It remains to investigate which of the methods yield more accurate results.




Figure 3: Potential energy surfaces with [2]_{R12} -correction and PNOs as CABS for H₂, LiH, BH, BeH₂, following Ref. [11]. SPA ansatz with doubles excitations only (SPA-UpCCD [93]). For validation of the SPA ansatz compared to classical techniques such as HF, CC, we refere to Ref. [93]




Figure 4: Corrected potential energy surfaces using classical FCI and Gaussian basis sets.




Figure 5: Performance of MRA-PNOs with and without $[2]_{R12}$ in comparison to FCI in a DZ-basis. CABS for MRA are generated using the *good* setup (dim(CABS) as in Fig. 3; for the GBS we used cc-pVDZ-F12-OPTRI, amounting to dim(CABS) of 48 for H₂ and 110 for both LiH, BH).

Within a classical benchmark [49], they performed equally well.

Additionally it would be interesting to examine the influence of noise on $[2]_{R12}$, which has not been considered in this work. To this end, we refer to Ref. [89] who applied the virtual quantum subspace expansion on actual quantum hardware. Beyond that, Refs. [88, 89] proposed a technique to account for measurement noise – similar approaches can be thought of in this case.

Finally, one might think of combinations of the quantum subspace expansion together with $[2]_{R12}$, given that the necessary reduced density matrices are already available. However this does turn out to be not trivial because when applying two different perturbative schemes, one must be careful not to account for the same behaviour twice and "overcompensate". One way to approach this might be to follow the idea of a "CABS singles" correction [19] to additionally account for one-electron incompleteness.

V. ACKNOWLEDGMENTS

This work was supported by the U.S. Department of Energy under Award No. DE-SC0019374.

A.A.-G. acknowledges the generous support from Google, Inc. in the form of a Google Focused Award. A.A.-G. also acknowledges support from the Canada Industrial Research Chairs Program and the Canada 150 Research Chairs Program. P.S. acknowledges support by a fellowship within the IFI program of the German Academic Exchange Service (DAAD). Simulations were performed with computing resources granted by RWTH Aachen University under project thes0850. This research was enabled in part by support provided by Compute Canada. Computations were performed on the niagara supercomputer at the SciNet HPC Consortium[105, 106]. SciNet is funded by: the Canada Foundation for Innovation; the Government of Ontario; Ontario Research Fund - Research Excellence; and the University of Toronto. J.S.K thanks Florian Bischoff for introducing him to explicit correlation and the depths of madness back

in the days. We thank the generous support of Anders G. Frøseth.

- [1] F. Arute, K. Arya, R. Babbush, D. Bacon, J. C. Bardin, R. Barends, R. Biswas, S. Boixo, F. G. Brandao, D. A. Buell, B. Burkett, Y. Chen, Z. Chen, B. Chiaro, R. Collins, W. Courtney, A. Dunsworth, E. Farhi, B. Foxen, A. Fowler, C. Gidney, M. Giustina, R. Graff, K. Guerin, S. Habegger, M. P. Harrigan, M. J. Hartmann, A. Ho, M. Hoffmann, T. Huang, T. S. Humble, S. V. Isakov, E. Jeffrey, Z. Jiang, D. Kafri, K. Kechedzhi, J. Kelly, P. V. Klimov, S. Knysh, A. Korotkov, F. Kostritsa, D. Landhuis, M. Lindmark, E. Lucero, D. Lyakh, S. Mandrà, J. R. McClean, M. McEwen, A. Megrant, X. Mi, K. Michielsen, M. Mohseni, J. Mutus, O. Naaman, M. Neeley, C. Neill, M. Y. Niu, E. Ostby, A. Petukhov, J. C. Platt, C. Quintana, E. G. Rieffel, P. Roushan, N. C. Rubin, D. Sank, K. J. Satzinger, V. Smelyanskiy, K. J. Sung, M. D. Trevithick, A. Vainsencher, B. Villalonga, T. White, Z. J. Yao, P. Yeh, A. Zalcman, H. Neven, and J. M. Martinis, Nature 574, 505 (2019).
- [2] G. AI Quantum, Science 369, 1084 (2020), arXiv:2004.04174.
- [3] H. S. Zhong, H. Wang, Y. H. Deng, M. C. Chen, L. C. Peng, Y. H. Luo, J. Qin, D. Wu, X. Ding, Y. Hu, P. Hu, X. Y. Yang, W. J. Zhang, H. Li, Y. Li, X. Jiang, L. Gan, G. Yang, L. You, Z. Wang, L. Li, N. L. Liu, C. Y. Lu, and J. W. Pan, Science **370**, 1460 (2021), arXiv:2012.01625.
- [4] J. Preskill, Quantum 2, 79 (2018), arXiv:1801.00862.
- [5] M. Cerezo, A. Arrasmith, R. Babbush, S. C. Benjamin, S. Endo, K. Fujii, J. R. McClean, K. Mitarai, X. Yuan, L. Cincio, *et al.*, Nature Reviews Physics, 1 (2021).
- [6] K. Bharti, A. Cervera-Lierta, T. H. Kyaw, T. Haug, S. Alperin-Lea, A. Anand, M. Degroote, H. Heimonen, J. S. Kottmann, T. Menke, *et al.*, arXiv preprint arXiv:2101.08448 (2021).
- [7] S. Endo, Z. Cai, S. C. Benjamin, and X. Yuan, Journal of the Physical Society of Japan 90, 032001 (2021).
- [8] J. R. McClean, M. E. Kimchi-Schwartz, J. Carter, and W. A. De Jong, Physical Review A 95, 1 (2017), arXiv:1603.05681.
- [9] T. Takeshita, N. C. Rubin, Z. Jiang, E. Lee, R. Babbush, and J. R. McClean, Physical Review X 10, 11004 (2020), arXiv:1902.10679.
- [10] M. Motta, T. P. Gujarati, J. E. Rice, A. Kumar, C. Masteran, J. A. Latone, E. Lee, E. F. Valeev, and T. Y. Takeshita, Physical Chemistry Chemical Physics 22, 24270 (2020).
- [11] J. S. Kottmann, P. Schleich, T. Tamayo-Mendoza, and A. Aspuru-Guzik, The Journal of Physical Chemistry Letters 12, 663 (2021), arXiv:2008.02819.
- [12] Z.-J. Zhang, T. H. Kyaw, J. Kottmann, M. Degroote, and A. Aspuru-Guzik, Quantum Science and Technology (2021), 10.1088/2058-9565/abdca4.
- [13] S. F. Boys and N. C. Handy, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences 310, 43 (1969).
- [14] S. F. Boys and N. C. Handy, in *Society*, Vol. 310 (1969) pp. 63–78.
- [15] S. McArdle and D. P. Tew, arXiv preprint arXiv:2006.11181 (2020).
- [16] S. McArdle, T. Jones, S. Endo, Y. Li, S. C. Ben-

jamin, and X. Yuan, npj Quantum Information 5 (2019), 10.1038/s41534-019-0187-2, arXiv:1804.03023.

- [17] T. Yanai and T. Shiozaki, Journal of Chemical Physics 136 (2012), 10.1063/1.3688225.
- [18] M. Torheyden and E. F. Valeev, Journal of Chemical Physics 131 (2009), 10.1063/1.3254836.
- [19] L. Kong and E. F. Valeev, Journal of Chemical Physics 135 (2011), 10.1063/1.3664729.
- [20] L. B. Roskop, L. Kong, E. F. Valeev, M. S. Gordon, and T. L. Windus, Journal of Chemical Theory and Computation 10, 90 (2014).
- [21] J. R. McClean, Z. Jiang, N. C. Rubin, R. Babbush, and H. Neven, Nature Communications 11, 1 (2020), arXiv:1903.05786.
- [22] L. Kong and E. F. Valeev, Journal of Chemical Physics 133, 174126 (2010).
- [23] Y. Cao, J. Romero, J. P. Olson, M. Degroote, P. D. Johnson, M. Kieferová, I. D. Kivlichan, T. Menke, B. Peropadre, N. P. Sawaya, S. Sim, L. Veis, and A. Aspuru-Guzik, Chemical Reviews 119, 10856 (2019), arXiv:1812.09976.
- [24] P. Schleich, Regularization of Quantum Chemistry on Quantum Computers by means of Explicit Correlation, Master's thesis, RWTH Aachen University, http://www.mathcces.rwth-aachen.de/_media/ 3teaching/00projects/2020_ma_schleich.pdf (2020).
- [25] W. Kutzelnigg and D. Mukherjee, Journal of Chemical Physics 107, 432 (1997).
- [26] W. Kutzelnigg, in *Explicitly Correlated Wave Functions in Chemistry and Physics*, edited by J. Rychlewski (Springer, Dordrecht, 2003) pp. 3–90.
- [27] T. Kato, Communications on Pure and Applied Mathematics 10, 151 (1957).
- [28] W. Lakin, The Journal of Chemical Physics 43, 2954 (1965).
- [29] R. T. Pack and W. Byers Brown, The Journal of Chemical Physics 45, 625 (1966).
- [30] W. Kutzelnigg and J. D. Morgan, The Journal of Chemical Physics 96, 4484 (1992).
- [31] E. A. Hylleraas, Zeitschrift für Physik 54, 347 (1929).
- [32] J. O. Hirschfelder, The Journal of Chemical Physics **39**, 3145 (1963).
- [33] C. Schwartz, Methods in Computational Physics 2, 241 (1963).
- [34] R. N. Hill, The Journal of Chemical Physics 83, 1173 (1985).
- [35] W. Kutzelnigg, Physical Chemistry Chemical Physics 10, 3460 (2008).
- [36] L. Kong, F. A. Bischoff, and E. F. Valeev, Chemical Reviews 112, 75 (2012).
- [37] W. Klopper, F. R. Manby, S. Ten-No, and E. F. Valeev, International Reviews in Physical Chemistry 25, 427 (2006).
- [38] S. Ten-no and J. Noga, Wiley Interdisciplinary Reviews: Computational Molecular Science 2, 114 (2012).
- [39] A. Grüneis, S. Hirata, Y. ya Ohnishi, and S. Ten-No, "Perspective: Explicitly correlated electronic structure theory for complex systems," (2017).

- [40] J. S. Sims and S. Hagstrom, Phys. Rev. A 4, 908 (1971).
- [41] E. Cancès, M. Defranceschi, W. Kutzelnigg, C. Le Bris, and Y. Maday, Handbook of Numerical Analysis 10, 3 (2003).
- [42] W. Kutzelnigg, Theoretica Chimica Acta **68**, 445 (1985).
- [43] W. Klopper and W. Kutzelnigg, Chemical Physics Letters 134, 17 (1987).
- [44] W. Klopper and W. Kutzelnigg, The Journal of Chemical Physics 94, 2020 (1991).
- [45] W. Kutzelnigg and W. Klopper, The Journal of Chemical Physics 94, 1985 (1991).
- [46] S. Ten-No, Chemical Physics Letters 447, 175 (2007).
- [47] S. F. Boys and N. C. Handy, Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences 309, 209 (1969).
- [48] H. Luo, Journal of Chemical Physics 133 (2010), 10.1063/1.3505037.
- [49] J. A. Kersten, G. H. Booth, and A. Alavi, Journal of Chemical Physics 145, 054117 (2016), arXiv:1605.07065.
- [50] S. F. Boys and M. V. Wilkes, Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 258, 402 (1960).
- [51] K. Singer and M. V. Wilkes, Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences 258, 412 (1960).
- [52] S. Ten-No, Chemical Physics Letters 398, 56 (2004).
- [53] C. M. Johnson, S. Hirata, and S. Ten-no, Chemical Physics Letters 683, 247 (2017).
- [54] D. G. Smith, L. A. Burns, A. C. Simmonett, R. M. Parrish, M. C. Schieber, R. Galvelis, P. Kraus, H. Kruse, R. Di Remigio, A. Alenaizan, A. M. James, S. Lehtola, J. P. Misiewicz, M. Scheurer, R. A. Shaw, J. B. Schriber, Y. Xie, Z. L. Glick, D. A. Sirianni, J. S. O'Brien, J. M. Waldrop, A. Kumar, E. G. Hohenstein, B. P. Pritchard, B. R. Brooks, H. F. Schaefer, A. Y. Sokolov, K. Patkowski, A. E. Deprince, U. Bozkaya, R. A. King, F. A. Evangelista, J. M. Turney, T. D. Crawford, and C. D. Sherrill, Journal of Chemical Physics 152, 184108 (2020).
- [55] W. Kutzelnigg and D. Mukherjee, Journal of Chemical Physics 110, 2800 (1999).
- [56] W. Kutzelnigg, K. R. Shamasundar, and D. Mukherjee, Molecular Physics 108, 433 (2010).
- [57] S. Ten-No, Journal of Chemical Physics 121, 117 (2004).
- [58] A. Peruzzo, J. McClean, P. Shadbolt, M. H. Yung, X. Q. Zhou, P. J. Love, A. Aspuru-Guzik, and J. L. O'Brien, Nature Communications 5 (2014), 10.1038/ncomms5213.
- [59] J. S. Kottmann, F. A. Bischoff, and E. F. Valeev, Journal of Chemical Physics 152, 074105 (2020).
- [60] R. J. Harrison, G. Beylkin, F. A. Bischoff, J. A. Calvin, G. I. Fann, J. Fosso-Tande, D. Galindo, J. R. Hammond, R. Hartman-Baker, J. C. Hill, J. Jia, J. S. Kottmann, M. J. Ou, J. Pei, L. E. Ratcliff, M. G. Reuter, A. C. Richie-Halford, N. A. Romero, H. Sekino, W. A. Shelton, B. E. Sundahl, W. S. Thornton, E. F. Valeev, Á. Vázquez-Mayagoitia, N. Vence, T. Yanai, and Y. Yokoi, in *Madness: A multiresolution, adaptive numerical environment for scientific simulation*, Vol. 38 (Society for Industrial and Applied Mathematics Publications, 2016) pp. S123–S142.
- [61] E. F. Valeev, Chemical Physics Letters 395, 190 (2004).
- [62] This is one way to derive it. One could also blindly start with some perturbation and the second-order Hylleraas functional, interpreting the RI-approximation in the usual sense to break

down higher-order integrals. But since this approach well showcases where the correction for *basis-set incompleteness* comes from, we choose to sketch it here. A more thorough outline can be found in Ref. [24].

- [63] P.O. Löwdin, Journal of Mathematical Physics 3, 969 (1962).
- [64] E. F. Valeev, Physical Chemistry Chemical Physics **10**, 106 (2008).
- [65] M. Torheyden and E. F. Valeev, Physical Chemistry Chemical Physics 10, 3410 (2008).
- [66] Cumulant approximations [55] as well as the standard, extended and generalized Brillouin conditions [45] and a set of so called screening approximations [18, 19, 64].
- [67] S. Kedžuch, M. Milko, and J. Noga, International Journal of Quantum Chemistry 105, 929 (2005).
- [68] F. Pavošević, P. Pinski, C. Riplinger, F. Neese, and E. F. Valeev, The Journal of Chemical Physics 144, 144109 (2016).
- [69] J. R. McClean, J. Romero, R. Babbush, and A. Aspuru-Guzik, New Journal of Physics 18 (2016), 10.1088/1367-2630/18/2/023023, arXiv:1509.04279.
- [70] S. McArdle, S. Endo, A. Aspuru-Guzik, S. C. Benjamin, and X. Yuan, Reviews of Modern Physics **92** (2020), 10.1103/RevModPhys.92.015003, arXiv:1808.10402.
- [71] J. Romero, R. Babbush, J. R. McClean, C. Hempel, P. J. Love, and A. Aspuru-Guzik, Quantum Science and Technology 4 (2019), 10.1088/2058-9565/aad3e4, arXiv:1701.02691.
- [72] A. Anand, P. Schleich, S. Alperin-Lea, P. W. Jensen, S. Sim, M. Díaz-Tinoco, J. S. Kottmann, M. Degroote, A. F. Izmaylov, and A. Aspuru-Guzik, Chemical Society Reviews (2022).
- [73] J. Lee, W. J. Huggins, M. Head-Gordon, and K. B. Whaley, Journal of Chemical Theory and Computation 15, 311 (2019), arXiv:1810.02327.
- [74] H. R. Grimsley, S. E. Economou, E. Barnes, and N. J. Mayhall, Nature Communications **10** (2019), 10.1038/s41467-019-10988-2, arXiv:1812.11173.
- [75] H. R. Grimsley, D. Claudino, S. E. Economou, E. Barnes, and N. J. Mayhall, Journal of Chemical Theory and Computation 16, 1 (2020), arXiv:1910.10329.
- [76] A. F. Izmaylov, M. Díaz-Tinoco, and R. A. Lang, Physical Chemistry Chemical Physics 22, 12980 (2020), arXiv:2003.07351.
- [77] I. G. Ryabinkin, T.-C. Yen, S. N. Genin, and A. F. Izmaylov, Journal of Chemical Theory and Computation 14, 6317 (2018).
- [78] J. Kottmann, S. Alperin-Lea, T. Tamayo-Mendoza, A. Cervera-Lierta, C. Lavigne, T.-C. Yen, V. Verteletskyi, P. Schleich, A. Anand, M. Degroote, and et al., Quantum Science and Technology (2021), 10.1088/2058-9565/abe567.
- [79] J. S. Kottmann, A. Anand, and A. Aspuru-Guzik, Chem. Sci. 12, 3497 (2021).
- [80] A. F. Izmaylov, T.-C. Yen, R. A. Lang, and V. Verteletskyi, Journal of chemical theory and computation 16, 190 (2019).
- [81] V. Verteletskyi, T. C. Yen, and A. F. Izmaylov, Journal of Chemical Physics 152 (2020), 10.1063/1.5141458, arXiv:1907.03358.
- [82] O. Crawford, B. v. Straaten, D. Wang, T. Parks, E. Campbell, and S. Brierley, Quantum 5, 385 (2021).
- [83] W. J. Huggins, J. R. McClean, N. C. Rubin, Z. Jiang, N. Wiebe, K. B. Whaley, and R. Babbush, npj Quantum Information 7, 1 (2021).
- [84] T.-C. Yen and A. F. Izmaylov, arXiv preprint

arXiv:2007.01234 (2020).

- [85] X. Bonet-Monroig, R. Babbush, and T. E. O'Brien, Phys. Rev. X 10, 031064 (2020).
- [86] N. C. Rubin, R. Babbush, and J. McClean, New Journal of Physics 20, 053020 (2018), arXiv:1801.03524.
- [87] J. F. Gonthier, M. D. Radin, C. Buda, E. J. Doskocil, C. M. Abuan, and J. Romero, arXiv preprint arXiv:2012.04001 (2020).
- [88] G. H. Booth, D. Cleland, A. Alavi, and D. P. Tew, The Journal of chemical physics 137, 164112 (2012).
- [89] M. Urbanek, D. Camps, R. Van Beeumen, and W. A. de Jong, Journal of Chemical Theory and Computation 16, 5425 (2020).
- [90] D. P. Tew, B. Helmich, and C. Hättig, The Journal of chemical physics 135, 074107 (2011).
- [91] G. Schmitz, C. Hättig, and D. P. Tew, Physical Chemistry Chemical Physics 16, 22167 (2014).
- [92] P. O. Löwdin, Physical Review 97, 1474 (1955).
- [93] J. S. Kottmann and A. Aspuru-Guzik, Physical Review A 105, 032449 (2022).
- [94] Psi4: Currently, CABS-functionality is not available in the main repository; a hacky but ready-to-use implementation is accessible at https://github.com/ philipp-q/psi4/tree/ri_space. For MADNESS, we use the forked branch https://github.com/kottmanj/ madness/tree/pno_integrals_cabs, which is described in more detail in Ref. [11]. Installation instructions for MADNESS together with TEQUILA can be found at https: //github.com/kottmanj/madness/tree/tequila. Integration into the main repository is planned here.
- [95] J. R. McClean, N. C. Rubin, K. J. Sung, I. D. Kivlichan, X. Bonet-Monroig, Y. Cao, C. Dai, E. S. Fried, C. Gidney, B. Gimby, P. Gokhale, T. Haner, T. Hardikar, V. Havlíček, O. Higgott, C. Huang, J. Izaac, Z. Jiang, X. Liu, S. Mcardle, M. Neeley, T. O'Brien, B. O'Gorman, I. Ozfidan, M. D. Radin, J. Romero, N. P. Sawaya, B. Senjean, K. Setia, S. Sim, D. S. Steiger, M. Steudtner, Q. Sun, W. Sun, D. Wang, F. Zhang, and R. Babbush, Quantum Science and Technology 5 (2020), 10.1088/2058-9565/ab8ebc, arXiv:1710.07629.
- [96] Y. Suzuki, Y. Kawase, Y. Masumura, Y. Hiraga, M. Nakadai, J. Chen, K. M. Nakanishi, K. Mitarai, R. Imai, S. Tamiya, *et al.*, Quantum 5, 559 (2021).
- [97] J. Bradbury, R. Frostig, P. Hawkins, M. J. Johnson, C. Leary, D. Maclaurin, and S. Wanderman-Milne, URL http://github. com/google/jax 4, 16 (2020).
- [98] P. Virtanen, R. Gommers, T. E. Oliphant, M. Haberland, T. Reddy, D. Cournapeau, E. Burovski, P. Peterson, W. Weckesser, J. Bright, S. J. van der Walt, M. Brett, J. Wilson, K. J. Millman, N. Mayorov, A. R. J. Nelson, E. Jones, R. Kern, E. Larson, C. J. Carey, I. Polat, Y. Feng, E. W. Moore, J. VanderPlas, D. Laxalde, J. Perktold, R. Cimrman, I. Henriksen, E. A. Quintero, C. R. Harris, A. M. Archibald, A. H. Ribeiro, F. Pedregosa, P. van Mulbregt, A. Vijaykumar, A. P. Bardelli, A. Rothberg, A. Hilboll, A. Kloeckner, A. Scopatz, A. Lee, A. Rokem, C. N. Woods, C. Fulton, C. Masson, C. Häggström, C. Fitzgerald, D. A. Nicholson, D. R. Hagen, D. V. Pasechnik, E. Olivetti, E. Martin, E. Wieser, F. Silva, F. Lenders, F. Wilhelm, G. Young, G. A. Price, G.-L. Ingold, G. E. Allen, G. R. Lee, H. Audren, I. Probst, J. P. Dietrich, J. Silterra, J. T. Webber, J. Slavič, J. Nothman, J. Buchner, J. Kulick, J. L. Schön-

berger, J. V. de Miranda Cardoso, J. Reimer, J. Harrington, J. L. C. Rodríguez, J. Nunez-Iglesias, J. Kuczynski, K. Tritz, M. Thoma, M. Newville, M. Kümmerer, M. Bolingbroke, M. Tartre, M. Pak, N. J. Smith, N. Nowaczyk, N. Shebanov, O. Pavlyk, P. A. Brodtkorb, P. Lee, R. T. McGibbon, R. Feldbauer, S. Lewis, S. Tygier, S. Sievert, S. Vigna, S. Peterson, S. More, T. Pudlik, T. Oshima, T. J. Pingel, T. P. Robitaille, T. Spura, T. R. Jones, T. Cera, T. Leslie, T. Zito, T. Krauss, U. Upadhyay, Y. O. Halchenko, Y. Vázquez-Baeza, and S. 1.0 Contributors, Nature Methods **17**, 261 (2020).

- [99] J. S. Dehesa, J. C. Angulo, T. Koga, and K. Matsui, Zeitschrift für Physik D Atoms, Molecules and Clusters 25, 9 (1992).
- [100] H. Nakatsuji, Accounts of Chemical Research **45**, 1480 (2012).
- [101] F. A. Bischoff, Journal of Chemical Physics 141, 184105 (2014).
- [102] L. B. Roskop, E. F. Valeev, E. A. Carter, M. S. Gordon, and T. L. Windus, Journal of Chemical Theory and Computation 12, 3176 (2016), pMID: 27281508, https://doi.org/10.1021/acs.jctc.6b00315.
- [103] C. Peng, C. Lewis, X. Wang, M. Clement, F. Pavosevic, J. Zhang, V. Rishi, N. Teke, K. Pierce, J. Calvin, J. Kenny, E. Seidl, C. Janssen, and E. Valeev, "The Massively Parallel Quantum Chemistry Program (MPQC),".
- [104] F. Pavošević, F. Neese, and E. F. Valeev, The Journal of chemical physics 141, 054106 (2014).
- [105] M. Ponce, R. van Zon, S. Northrup, D. Gruner, J. Chen, F. Ertinaz, A. Fedoseev, L. Groer, F. Mao, B. C. Mundim, et al., in Proceedings of the Practice and Experience in Advanced Research Computing on Rise of the Machines (learning) (2019) pp. 1–8.
- [106] C. Loken, D. Gruner, L. Groer, R. Peltier, N. Bunn, M. Craig, T. Henriques, J. Dempsey, C.-H. Yu, J. Chen, et al., in Journal of Physics-Conference Series, Vol. 256 (2010) p. 012026.
- [107] A. J. May, E. Valeev, R. Polly, and F. R. Manby, Physical Chemistry Chemical Physics 7, 2710 (2005).

Appendix A: Code-Sample to calculate [2]_{R12} with TEQUILA

i<mark>mport</mark> tequila as tq

```
def compute_with_mra_pnos(act, ri, geometry):
    madroot = '''link to madness executable''
    madnessinput = {"pnoint": {'cabs_option': '
    # We use "act" orbitals for the reference
    \hookrightarrow and an RI space of "ri" orbitals
    # The n_pno in mol sets the overall number
    \hookrightarrow of PNOs to be generated, while the n_pno
     \rightarrow 
        in the madnessinput sets the size of
    \hookrightarrow the reference space in the CABS+

→ procedure

    mol = tq.Molecule(name='molecule', geometry
    ← =geometry, basis_set='madness', n_pno=ri
    \hookrightarrow
        , active_orbitals=[i for i in range(act)

→ ], executable=executable, **madnessinput

    → )
    H = mol.make_hamiltonian().simplify()
    U = mol.make_upccgsd_ansatz(name='SPA-UCCD'
```

```
→ )
      E = tq.ExpectationValue(H=H, U=U)
      # HF as initial values
      initial_values = {k: 0.0 for k in E.

→ extract_variables() }

      result = tq.minimize(objective=E, method="
14
      energy = result.energy
      angles = result.angles
16
      # Prepare information to build rdms -- this
         can also be handed over explicitly
      rdminfo = {"U": U, "variables": angles}
18
      # Compute f12 correction, use full CABS
19
      dE = mol.perturbative_f12_correction(**
20
      \hookrightarrow rdminfo)
22
      return energy, dE
24
25 # Quick test with GBS and MRA-PNOs
                                         'H 0.0
  res_pno = compute_with_mra_pnos(2, 5,
      \hookrightarrow 0.0 0.0\nH 0.0 0.0 1.0')
27 print(res_pno) # (-1.12866, -0.00638), dim(RI)
      → =6
```

```
1 import tequila as tq
2 import psi4
3
  def compute_with_gbs(obs, cabs, geometry):
4
       # Currently: Need C1 symmetry for [2]R12
       mol = tq.Molecule(geometry=geometry,
6
      ⇔ basis_set = obs, point_group='c1')
# Let's use psi4's "detci" to determine the
       \hookrightarrow RDMs -- generally, only CI-methods work
       → right now
       psi4_method = 'detci'
8
       # Set the CABS-basis set for the correction
0
       cabs_opt= {'cabs_name': cabs}
10
       # Here, let's compute the RDMs explicity (
11
      \hookrightarrow but one can do the same as above and
       \hookrightarrow hand over this information via keyword
      \hookrightarrow arguments)
       mol.compute_rdms(psi4_method=psi4_method)
       energy = mol.logs[psi4_method].variables["
       ↔ CI TOTAL ENERGY"]
       dE = mol.perturbative_f12_correction(rdm1=
14
       → mol.rdm1, rdm2=mol.rdm2, cabs_type='cabs
      \hookrightarrow +', gamma=1.4, cabs_options=cabs_opt)
15
       return energy, dE
16
17
18
19 # Quick test
20 res_gbs = compute_with_gbs('sto-3g', '6-31g', '

→ H 0.0 0.0 0.0\nH 0.0 0.0 1.0')
 print(res_gbs) # (-1.10115, -0.00174), dim(RI)
     → =6
```

Appendix B: PNOs as CABS

The following data comes from [24].

First, we take a look at the "convergence" of the $[2]_{R12}$ - correction with respect to the dimensionality of the RI space

to allow a judgement whether additional F12-regularization of the MRA-PNOs is beneficial, and how MRA-PNOs perform as a CABS in comparison to designated Gaussian basis sets. Note that we do not consider the RI-error here as this is rather a proof-of-principle for the convergence behaviour of MRA-PNOs vs. GBS for CABS than actual production calculations. Results therefor are depicted in Fig. 6.

In the case of the hydrogen molecule, the best energies are achieved by the GBS-CABS, while the mixed-approach does not improve the result further. This means that the GBS-CABS, cc-pVDZ-F12-OPTRI, is likely to be very well suited as RI in this case, which is expected because it is optimized for this use. The regularized equivalent yields a barely improved result. Looking at the PNO-CABS, we see that the unregularized surrogate performs almost as well as the much larger GBS-CABS, while the PNO-CABS grown by MP2-R12 give a worse result despite a correction of almost the same magnitude using even less orbitals as CABS. The reason thereof is that the reference is different. In the case of MP2 as surrogate, both the PNOs for OBS and CABS are unregularized, and accordingly, for the MP2-F12 CABS, we use OBS that have been generated within the same process coming from MP2-F12. We chose this approach because this way, the PNOs are generated within one consistent procedure and then easily combined by CABS+ [61]. Different combinations, mixing MP2 and MP2-F12 for OBS and CABS, can be thought of as well but lead to a less efficient procedure that creates a fair amount of unnecessary orbitals that are to be projected out.

Taking a look at the converse result for lithium hydride, we see that here, the MP2-F12 options do considerably worse than before. The higher magnitude of the correction is again related to the worse reference energy using PNOs built by MP2-F12. Building a higher-quality CABS in this case might account for that but comes at additional cost. Further the choice of GBS-CABS (def2-SVP-RIFIT) leads to generally low-performing GBS-CABS in this case, which is much less extensive than the we chose for the hydrogen molecule. The noticable improvement by the mixed approach in the case of MP2-F12 supports this. Yet still, the same basis set for LiH would amount to 110 basis functions, which is more than three times than for the PNO-CABS in this case.

To formulate an overall recommendation, we come back to the $[2]_{R12}$ -correction, which in the framework of Ref. [20] scales cubically within the framework of approximation C [67] or quadratically within approximation D [68, 107] in the RI-dimensionality. Ref. [68] proposes a method using PNOs to reduce the cost of MP2-F12 to quasi-linear dependence in the RI-basis except for some inexpensive quadratic terms; a similar approach could be employed to reduce this cost for $[2]_{R12}$, too. Given the seemingly almost equivalent performance of unregularized PNOs for OBS and CABS and the fact, that there are significantly less CABS functions in this case, this approach seems most desirable within this context.




Figure 6: Explicitly correlated [2]_{R12} -correction over the dimensionality of RI basis for H₂ and LiH given a fixed OBS of MRA-PNOs, determined according to the *good* procedure. The final point is annotated with the corrected energy. H₂: cc-pVDZ-F12-OPTRI for GBS-CABS and 3 PNOs added to the CABS in the mixed case. LiH: def2-SVP-RIFIT as GBS-CABS, the mixed procedure adds another 6 PNOs to the CABS.