BOUNDARY VALUE PROBLEMS FOR SEMILINEAR SCHRÖDINGER EQUATIONS WITH SINGULAR POTENTIALS AND MEASURE DATA

MOUSOMI BHAKTA, MOSHE MARCUS, AND PHUOC-TAI NGUYEN

ABSTRACT. We study boundary value problems with measure data in smooth bounded domains Ω , for semilinear equations. Specifically we consider problems of the form $-L_V u + f(u) = \tau$ in Ω and $\operatorname{tr}_V u = \nu$ on $\partial\Omega$, where $L_V = \Delta + V$, $f \in C(\mathbb{R})$ is monotone increasing with f(0) = 0 and $\operatorname{tr}_V u$ denotes the measure boundary trace of u associated with L_V . The potential V is typically a Hölder continuous function in Ω that blows up at a set $F \subset \partial\Omega$ as dist $(x, F)^{-2}$. In general the above boundary value problem may not have a solution. We are interested in questions related to the concept of 'reduced measures', introduced in [4] for V = 0. Our results extend results of [4] and [6] and apply to a larger class of nonlinear terms f. In the case of signed measures, some of the present results are new even for V = 0.

Keywords: reduced measures, boundary trace, harmonic measures, Kato's inequality.

MSC numbers: 35J61, 35J75, 35J10

Contents

1. Introduction	2
2. Some results on sub and supersolutions.	7
2.1. Estimates of L_V harmonic functions and L_V potentials.	7
2.2. Remarks on subsolutions and supersolutions	8
3. The reduced measures for couples of positive measures.	9

⁽M. Bhakta) Department of Mathematics, Indian Institute of Science Education and Research, Pune 411008, India

⁽M. Marcus, Corresponding author) DEPARTMENT OF MATHEMATICS, TECHNION, HAIFA 32000, ISRAEL

⁽P. T. Nguyen) DEPARTMENT OF MATHEMATICS AND STATISTICS, MASARYK UNIVERSITY, BRNO, CZECH REPUBLIC

E-mail addresses: mousomi@iiserpune.ac.in, marcusm@math.technion.ac.il, ptnguyen@math.muni.cz.

4. Signed measures	17
Appendix A.	24
References	27

1. INTRODUCTION

Let Ω be a C^2 bounded domain in \mathbb{R}^N , $N \geq 3$, and let

$$L_V := \Delta + V$$

where $V \in C^{\theta}(\Omega)$, for some $\theta \in (0, 1]$, satisfies the following conditions:

(A1)
$$\exists \bar{a} > 0 : |V(x)| \le \bar{a}\delta(x)^{-2} \quad \forall x \in \Omega,$$
$$\delta(x) := \operatorname{dist}(x, \partial\Omega),$$

(A2)
$$\int_{\Omega} |\nabla \phi|^2 \, dx \ge \int_{\Omega} \phi^2 V \, dx \quad \forall \phi \in H^1_0(\Omega).$$

These conditions imply the existence of a (minimal) Green function G_V and of the Martin kernel K_V for the operator $-L_V$. Related to this, the operator has a ground state that we denote by Φ_V . In the present case Φ_V is a positive eigenfunction of $-L_V$ with eigenvalue $\lambda_V > 0$.

The function Φ_V and the Martin kernel K_V are normalized at a reference point $x_0 \in \Omega$:

$$\Phi_V(x_0) = 1, \quad K_V(x_0, y) = 1 \quad \forall y \in \partial \Omega.$$

Notation. Denote

$$\mathbb{K}_{V}[\nu](x) := \int_{\partial\Omega} K_{V}(x, y) d\nu(y) \quad \forall \nu \in \mathfrak{M}(\partial\Omega),$$
$$\mathbb{G}_{V}[\tau](x) := \int_{\Omega} G_{V}(x, y) d\tau(y) \quad \forall \tau \in \mathfrak{M}(\Omega; \Phi_{V}).$$

Here $\mathfrak{M}(\partial\Omega)$ denotes the space of finite Borel measures on $\partial\Omega$ and $\mathfrak{M}(\Omega; \Phi_V)$ denotes the space of real Borel measures τ in Ω such that $\int_{\Omega} \Phi_V d|\tau| < \infty$. As usual $\mathfrak{M}_+(\partial\Omega)$ and $\mathfrak{M}_+(\Omega; \Phi_V)$ denote the positive cones of these spaces.

A function $u \in L^1_{loc}(\Omega)$ is L_V harmonic (resp. subharmonic, superharmonic) in Ω if $-L_V u = (\text{resp.} \leq, \geq) 0$ in Ω in the distribution sense.

By the Martin representation theorem, for every positive L_V harmonic function u in Ω there exists $\nu \in \mathfrak{M}_+(\partial \Omega)$ such that $u = \mathbb{K}_V[\nu]$.

By the Riesz decomposition lemma, a positive L_V superharmonic u can be represented in the form u = p + h where h is the largest

 $\mathbf{2}$

 L_V harmonic function dominated by u and p is an L_V potential, i.e. a positive L_V superharmonic function which does not dominate any positive L_V harmonic.

A function u is an L_V potential if and only if there exists a positive measure $\tau \in \mathfrak{M}(\Omega; \Phi_V)$ such that $u = \mathbb{G}_V[\tau]$. If τ is a positive Radon measure then either $\mathbb{G}_V[\tau]$ is finite everywhere in Ω or $\mathbb{G}_V[\tau] \equiv \infty$. Moreover, $\mathbb{G}_V[\tau] < \infty$ if and only if $\tau \in \mathfrak{M}(\Omega; \Phi_V)$.

For these and other basic potential theory results we refer the reader to [1]. A brief survey can be found in [15].

In this paper we study boundary value problems of the form

(1.1)
$$-L_V u + f(u) = \tau \quad \text{in } \Omega, \quad \text{tr}_V u = \nu \quad \text{on } \partial\Omega,$$

always assuming that

(1.2)
$$f \in C(\mathbb{R}), \quad f \text{ is non-decreasing}, \quad f(0) = 0$$

and

$$\nu \in \mathfrak{M}(\partial \Omega), \quad \tau \in \mathfrak{M}(\Omega; \Phi_V).$$

Finally $\operatorname{tr}_V u$, the L_V boundary trace of u, is defined as follows.

Definition 1.1. A non-negative Borel function u defined in Ω has an L_V boundary trace $\nu \in \mathfrak{M}(\partial \Omega)$ if

(1.3)
$$\lim_{n \to \infty} \int_{\partial D_n} h u \, d\omega_V^{x_0, D_n} = \int_{\partial \Omega} h d\nu \quad \forall h \in C(\bar{\Omega}),$$

for every uniformly Lipschitz exhaustion $\{D_n\}$ of Ω such that $x_0 \in D_n$ for all n. Here x_0 is the reference point previously mentioned and $\omega_V^{x_0,D_n}$ denotes the harmonic measure for L_V in D_n relative to x_0 . The L_V boundary trace of u is denoted by tr_Vu.

A real Borel function u defined in Ω has an L_V boundary trace if

(1.4)
$$\sup_{n} \int_{\partial D_{n}} |u| \, d\omega_{V}^{x_{0}, D_{n}} < +\infty$$

and (1.3) holds.

When V = 0 this definition reduces to the classical definition of measure boundary trace. Recall that

$$d\omega_n^{x_0} = P_{V,n}(x_0, \cdot)dS$$
 on ∂D_n ,

where $P_{V,n}$ is the Poisson kernel of $-L_V$ in D_n .

By [17, Lemma 2.3], if (A1), (A2) hold, the L_V trace has the following properties:

(1.5) (i)
$$\operatorname{tr}_{V}(\mathbb{K}_{V}[\nu]) = \nu \quad \forall \nu \in \mathfrak{M}(\partial \Omega),$$

(1.5)
$$(ii) \quad \operatorname{tr}_{V}(\mathbb{G}_{V}[\tau]) = 0 \quad \forall \tau \in \mathfrak{M}(\Omega; \Phi_{V}).$$

Notation. (i) Let (λ, σ) and (τ, ν) be two couples of measures in $\mathfrak{M}(\Omega; \Phi_V) \times \mathfrak{M}(\partial\Omega)$. Then $(\lambda, \sigma) \prec (\tau, \nu)$ means $\lambda \leq \tau$ and $\sigma \leq \nu$.

(ii) For
$$\beta > 0$$
, denote

(1.6)
$$D_{\beta} := \{ x \in \Omega : \delta(x) > \beta \}, \quad \Omega_{\beta} := \{ x \in \Omega : \delta(x) < \beta \}, \\ \Sigma_{\beta} := \{ x \in \Omega : \delta(x) = \beta \}.$$

Since Ω is a C^2 bounded domain, there exists $\beta_0 > 0$ such that for every $x \in \Omega_{\beta_0}$ there is a unique point $\sigma(x) \in \partial\Omega$ such that $|x - \sigma(x)| = \delta(x)$, and $x \mapsto \delta(x)$ is in $C^2(\Omega_{\beta_0})$ while $x \mapsto \sigma(x)$ is in $C^1(\Omega_{\beta_0})$.

In addition to (A1) and (A2), we assume that the ground state Φ_V satisfies the following condition:

There exist $a_0 \geq 1$ and $\alpha, \alpha^* > 0$ satisfying

$$0 \le \alpha - \alpha^* < \frac{1}{2},$$

such that for every $a > a_0$ and every $x, z \in \Omega_{\beta_0}$ lying on a normal to $\partial \Omega$:

(C1)
$$a\delta(x) \le \delta(z) \Longrightarrow \frac{\Phi_V(x)}{\Phi_V(z)} \le c(a) \frac{\delta(x)^{\alpha^*}}{\delta(z)^{\alpha}}.$$

Conditions (A1), (A2) and (C1) are assumed, without further mention, throughout the paper.

Definition 1.2. Let $(\tau, \nu) \in \mathfrak{M}(\Omega; \Phi_V) \times \mathfrak{M}(\partial \Omega)$ and $u \in L^1_{loc}(\Omega)$.

(i) u is a solution of (1.1) if $f(u) \in L^1(\Omega; \Phi_V)$, the equation holds in the distribution sense and tr_V $u = \nu$.

(ii) u is a subsolution of (1.1) if $f(u) \in L^1(\Omega; \Phi_V)$, $-L_V u + f(u) \leq \tau$ in the distribution sense and $\operatorname{tr}_V u \leq \nu$.

A supersolution is defined in the same way with the inverse inequalities.

With this definition, u is a solution of (1.1) if and only if (see [17, Lemma 3.1])

(1.7)
$$u + \mathbb{G}_V[f(u)] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu] \quad \text{in } \Omega.$$

If (1.1) has a solution we say that (τ, ν) is a good couple.

If $(0, \nu)$ (respectively $(\tau, 0)$) is a good couple we say that ν (respectively τ) is a good measure.

We are interested in questions related to the notion of 'reduced measure' introduced in [4] (for $L_V = \Delta$). In general, problem (1.1) is not solvable for every couple (τ, ν) .

A great deal of research has been devoted to a precise characterization of good measures or good couples in some specific cases. Most of this research dealt with the equation $-\Delta u + f(u) = 0$ in Ω and in particular with the case $f(t) = |t|^p \operatorname{sign} t$, p > 1 (see [9, 12, 13, 24] for 1 and [14, 22, 23] for every <math>p > 1 and the references therein). See also [2] where the problem was treated for a general class of nonlinearities f that satisfy the Keller - Osserman condition.

More recently the characterization of good measures was studied with respect to the equation $-L_V u + f(u) = 0$ in Ω , mainly when Vis the Hardy potential and $f(t) = |t|^p \operatorname{sign} t$ (see, e.g. [7, 10, 18–20]). The question was also studied in the context of fractional Schrödinger equations (see, e.g. [11]).

The idea of 'reduced measure', introduced in [4], is to provide a reduction process that converges to the 'good' part of τ and ν when (1.1) has no solution.¹

In [4], the authors study problem (1.1) for $L_V = \Delta$, mainly in the case where $\nu = 0, \tau \in \mathfrak{M}(\Omega)$, assuming that f satisfies (1.2) and vanishes on $(-\infty, 0]$. A solution is a function $u \in L^1(\Omega)$ such that $f(u) \in L^1(\Omega)$ and u satisfies (1.1) in the weak sense. To determine the reduced measure the authors consider a sequence of problems

(1.8)
$$-\Delta u + f_n(u) = \tau \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega,$$

where $f_n \in C(\mathbb{R})$ is a non-negative, nondecreasing function, $f_n \uparrow f$ and (1.8) has a solution for every $\tau \in \mathfrak{M}(\Omega)$. (For instance, the functions f_n are bounded.) One of the main results states (see [4, Theorem 4.1]):

Let u_n be the unique solution of (1.8). Then the sequence $\{u_n\}$ decreases and $u^* := \lim u_n$ satisfies

$$-\Delta u^* + f(u^*) = \tau^* \quad \text{in } \Omega, \quad u^* = 0 \quad \text{on } \partial\Omega,$$

where τ^* is the largest good measure dominated by τ .

In [6] a similar result is established in the case $(\tau, \nu) \in \mathfrak{M}(\Omega) \times \mathfrak{M}(\partial \Omega)$ (possibly signed measures) assuming as before that f = 0 on $(-\infty, 0]$. It is also shown that (τ, ν) is good if and only if, both τ and ν are good measures.

¹A related notion of 'reduced limit' was studied in [21], [3].

When τ, ν are signed measures and we drop the assumption 'f = 0 on $(-\infty, 0)$ ', the situation is more complex even in the case V = 0.

The present definition of a solution of (1.1) is necessarily different from that of weak solution used in [4], [6]. But when V = 0 these are essentially equivalent.

In the case where τ and ν are positive our results are similar to those quoted above.

Let u_n denote the solution of the problem,

(1.9)
$$-L_V u + f_n(u) = \tau \quad \text{in } \Omega, \quad \text{tr}_V u = \nu \quad \text{on } \partial \Omega$$

where $f_n \in C(\mathbb{R})$ is non-decreasing, bounded and $(f_n)_{\pm} \uparrow f_{\pm}$.

Theorem 1.3. Assume that f satisfies (1.2) and $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial \Omega)$. For u_n as above: $u_n \downarrow u^{\#}$ and

(1.10)
$$-L_V u^{\#} + f(u^{\#}) = \tau^{\#} \quad in \ \Omega, \quad \text{tr}_V u^{\#} = \nu^{\#}$$

where $(0,0) \prec (\tau^{\#}, \nu^{\#}) \prec (\tau, \nu)$. Moreover, $u^{\#}$ is the largest subsolution of (1.1) and $\tau^{\#}$, $\nu^{\#}$ are the largest good measures dominated by τ and ν respectively.

A corresponding result holds for couples of negative measures (see Remark 3.6).

In the case that τ, ν may be signed measures we prove:

Theorem 1.4. Assume that f satisfies (1.2) and $(\tau, \nu) \in \mathfrak{M}(\Omega; \Phi_V) \times \mathfrak{M}(\partial\Omega)$. Let (λ, σ) be a couple of measures such that

(1.11)
$$-(\tau_{-},\nu_{-})\prec(\lambda,\sigma)\prec(\tau_{+},\nu_{+})$$

and let u_n be the solution of (1.9) with (τ, ν) replaced by (λ, σ) .

Every subsequence of $\{u_n\}$ has a limit point with respect to a.e. convergence. If \tilde{u} is such a limit point then

(1.12)
$$-L_V \tilde{u} + f(\tilde{u}) = \lambda \quad in \ \Omega, \quad \operatorname{tr}_V \tilde{u} = \tilde{o}$$

and

(1.13)
$$(-\tau_{-},-\nu_{-})^{\#} \prec (\tilde{\lambda},\tilde{\sigma}) \prec (\tau_{+},\nu_{+})^{\#}.$$

Moreover, every couple (λ, σ) such that

(1.14)
$$(-\tau_{-}, -\nu_{-})^{\#} \prec (\lambda, \sigma) \prec (\tau_{+}, \nu_{+})^{\#}$$

is a good couple.

This naturally leads to the following question: If (λ, σ) is a good couple in the interval (1.11) does it necessarily satisfy (1.14)?

As shown below, if f vanishes in $(-\infty, 0]$, the answer is positive. In the general case this is an open question.

Theorem 1.5. Assume that f satisfies (1.2) and that f(t) = 0 for $t \leq 0$. Let $(\tau, \nu) \in \mathfrak{M}(\Omega; \Phi_V) \times \mathfrak{M}(\partial \Omega)$. Then (τ, ν) is a good couple if and only if

(1.15)
$$(\tau, \nu) \prec (\tau_+^\#, \nu_+^\#)$$

where $\tau_{+}^{\#}$ and $\nu_{+}^{\#}$ are the largest good measures dominated by τ_{+} and ν_{+} respectively.²

Consequently, (τ, ν) is a good couple with respect to (1.1) if and only if τ and ν separately are good measures.

This result extends [6, Theorem 6].

Our main tools include: two-sided estimates of $\mathbb{G}_V[\tau], \tau \in \mathfrak{M}_+(\Omega; \Phi_V)$, and $\mathbb{K}_V[\nu], \nu \in \mathfrak{M}_+(\partial\Omega)$ [16], the inverse maximum principle [8], Kato's inequality and its extension due to [5] and a result of [4] regarding the diffuse part of reduced measures.

Here is the plan of the paper. In Section 2 we recall the estimates of [16] as well as properties of sub and supersolutions established in [17] that are frequently used in the present paper. In Section 3 we study the problem of reduced couple for (1.1) with positive measures. Theorem 1.3 is a consequence of Theorems 3.1 and 3.5. Section 4 is devoted to problem (1.1) with signed measures. Theorem 1.4 is a consequence of Theorem 4.1 and Proposition 4.2. The section is completed by the proof of Theorem 1.5.

2. Some results on sub and supersolutions.

In this section we gather several results from [16] and [17] that are frequently used in the sequel.

2.1. Estimates of L_V harmonic functions and L_V potentials.

The estimates stated below are derived in [16].

Theorem 2.1. ([16, Theorem 3.1]) Assume that (A1), (A2) and (C1) hold. Then for any $\nu \in \mathfrak{M}_+(\partial\Omega)$,

$$\frac{1}{C} \|\nu\|_{\mathfrak{M}(\partial\Omega)} \leq \int_{\Sigma_{\beta}} \frac{\Phi_{V}}{\delta} \mathbb{K}_{V}[\nu] \, dS \leq C \|\nu\|_{\mathfrak{M}(\partial\Omega)} \quad \forall \beta \in (0, \beta_{0}),$$

where the constant C depends on \bar{a}, Ω and the constants in (C1).

²Note that, when f = 0 on $(-\infty, 0]$, the couple $(-\tau_{-}, -\nu_{-})$ is always good.

Next is an estimate of L_V potentials.

Theorem 2.2. ([16, Theorems 3.3 and 3.4])

(i) Assume (A1) and (A2) hold. Then there exists a constant c depending on \bar{a} , Ω such that, for every $\tau \in \mathfrak{M}_+(\Omega; \Phi_V)$,

$$\frac{1}{c} \int_{\Omega} \Phi_V \, d\tau \le \int_{\Omega} \frac{\Phi_V}{\delta} \mathbb{G}_V[\tau] \, dx.$$

(ii) Assume (A1), (A2) and (C1) hold. Then there exists c' > 0depending on \bar{a} , Ω and the constants in (C1) such that for every $\tau \in \mathfrak{M}_+(\Omega; \Phi_V)$,

$$\int_{\Omega} \frac{\Phi_V}{\delta} \mathbb{G}_V[\tau] \, dx \le c' \int_{\Omega} \Phi_V \, d\tau$$

2.2. Remarks on subsolutions and supersolutions. We list some properties of subsolutions and supersolutions from [17].

Lemma 2.3. Let $w \in L^1_{loc}(\Omega)$ be a non-negative L_V subharmonic function. If tr_Vw = 0 then $w \equiv 0$.

This is a consequence of [17, Corollary 2.6].

Lemma 2.4. ([17, Corollary 2.8]) Let $u \in L^1_{loc}(\Omega)$ and suppose that $-L_V u = \tau \in \mathfrak{M}(\Omega; \Phi_V)$. In addition assume that for some smooth exhaustion $\{D_n\}$ of Ω ,

(2.1)
$$\sup_{n} \int_{\partial D_n} P_{V,n}(x_0, y) |u(y)| \, dS(y) < +\infty.$$

Then $\operatorname{tr}_V u =: \nu$ exists and $u = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$ in Ω .

Remark. If $\operatorname{tr}_V u$ exists then, by definition, (2.1) holds.

If $u \ge 0$, condition (2.1) is not needed. In this case the result (stated below) is a consequence of the Riesz decomposition lemma.

Lemma 2.5. ([17, Lemma 2.11]) Let $u \in L^1_{loc}(\Omega)$ be a positive function such that $-L_V u = \tau \in \mathfrak{M}(\Omega; \Phi_V)$. Then u has an L_V boundary trace, say ν , and $u = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$ in Ω .

Notation. A function u is L_V perfect if $u = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$ for some $\tau \in \mathfrak{M}(\Omega; \Phi_V)$ and $\nu \in \mathfrak{M}(\partial \Omega)$.

Lemma 2.6. ([17, Lemma 3.3]) Suppose that u is an L_V perfect function. If $\operatorname{tr}_V u \leq 0$ then $\operatorname{tr}_V u_+ = 0$.

Consider the equation

(2.2)
$$-L_V u + f(u) = \tau \quad \text{in } \Omega$$

and the boundary value problem

(2.3)
$$-L_V u + f(u) = \tau \quad \text{in } \Omega, \quad \text{tr}_V u = \nu_z$$

where f satisfies (1.2), $\tau \in \mathfrak{M}(\Omega; \Phi_V)$ and $\nu \in \mathfrak{M}(\partial \Omega)$.

Definition 2.7. Let $u \in L^1_{loc}(\Omega)$ be a function such that $f(u) \in L^1_{loc}(\Omega)$.

The function u is a subsolution (supersolution) of (2.2) if $-L_V u + f(u) \leq (\geq)\tau$ in Ω in the distribution sense.

The function u is a subsolution (supersolution) of (2.3) if it is a subsolution (supersolution) of (2.2), $f(u) \in L^1(\Omega; \Phi_V)$ and u has an L_V boundary trace such that $\operatorname{tr}_V u \leq \nu$ (tr $_V u \geq \nu$).

Lemma 2.8. ([17, Lemma 3.1]) Assume that $u \in L^1_{loc}(\Omega)$ and $f(u) \in L^1(\Omega; \Phi_V)$.

If u is a subsolution (resp. supersolution) of problem (2.3) then u is L_V perfect. More precisely, there exist measures $\lambda \in \mathfrak{M}_+(\Omega; \Phi_V)$ and $\sigma \in \mathfrak{M}_+(\partial\Omega)$ such that:

(2.4)
$$u = \mathbb{G}_V[\tau - f(u) - \lambda] + \mathbb{K}_V[\nu - \sigma]$$
$$(u = \mathbb{G}_V[\tau - f(u) + \lambda] + \mathbb{K}_V[\nu + \sigma]).$$

Lemma 2.9. ([17, Lemma 3.4]) (i) Let u_1 (resp. u_2) be a supersolution (resp. subsolution) of (2.3). Then $u_2 \leq u_1$.

(ii) Problem (2.3) has at most one solution.

Lemma 2.10. ([17, Corollary 3.7]) Suppose that u_1 , u_2 are respectively a supersolution and a subsolution of (2.2) such that $f(u_i) \in L^1(\Omega; \Phi_V)$. If u_i has an L_V boundary trace, say ν_i , i = 1, 2, and $\nu_2 \leq \nu_1$ then problem (2.3) has a (unique) solution for every measure ν such that $\nu_2 \leq \nu \leq \nu_1$.

3. The reduced measures for couples of positive measures.

Notation. If f and g are two non-negative functions on a set X, we say that f and g are similar if there exists c > 0 such that

$$\frac{1}{c}f(x) \le g(x) \le cf(x) \quad \forall x \in X.$$

This relation is denoted by $f \sim g$.

Theorem 3.1. Let $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial\Omega)$. Let $\{f_n\}$ be a sequence of continuous, bounded, non-decreasing functions on \mathbb{R} such that $f_n(0) = 0$ and $(f_n)_{\pm} \uparrow f_{\pm}$.

Then the boundary value problem

(3.1)
$$-L_V u + f_n(u) = \tau \quad in \ \Omega, \quad \text{tr}_V u = \nu,$$

has a unique solution $u_n = u_n(\tau, \nu)$. The sequence $\{u_n\}$ is a decreasing sequence of positive functions and its limit $u^{\#} = u^{\#}(\tau, \nu)$ has the following properties.

(a) $u^{\#}$ satisfies

(3.2)
$$\|u^{\#}\|_{L^{1}(\Omega;\Phi_{V}/\delta)} + \|f(u^{\#})\|_{L^{1}(\Omega;\Phi_{V})} \le C(\|\nu\|_{\mathfrak{M}(\partial\Omega)} + \|\tau\|_{\mathfrak{M}(\Omega;\Phi_{V})})$$

and

(3.3)
$$u^{\#} + \mathbb{G}_V[f(u^{\#})] \le \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu] =: \tilde{w} \quad in \ \Omega.$$

(b) There exists a non-negative measure $\tau^{\#} \leq \tau$ such that

(3.4)
$$-L_V u^{\#} + f(u^{\#}) = \tau^{\#} \quad in \ \Omega.$$

(c) $u^{\#}$ has L_V boundary trace $0 \le \nu^{\#} \le \nu$. Thus

(3.5)
$$u^{\#} + \mathbb{G}_V[f(u^{\#})] = \mathbb{G}_V[\tau^{\#}] + \mathbb{K}_V[\nu^{\#}] \quad in \ \Omega.$$

(d) $u^{\#}$ is the largest subsolution of problem

(3.6)
$$-L_V u + f(u) = \tau \quad in \ \Omega, \quad \text{tr}_V u = \nu.$$

In particular, if (3.6) has a solution u then $u^{\#} = u$, $\tau^{\#} = \tau$ and $\nu^{\#} = \nu$.

Proof. The function $\tilde{w} := \mathbb{K}_V[\nu] + \mathbb{G}_V[\tau]$ is a supersolution of the equation

$$-L_V u + f_n(u) = \tau \quad \text{in } \Omega$$

and tr_V $\tilde{w} = \nu$. Obviously $v \equiv 0$ is a subsolution, $v \leq \tilde{w}$ and $f_n(\tilde{w}) \in L^1(\Omega; \Phi_V)$.

Therefore, by Lemma 2.10, there exists a unique solution $u_n = u_n(\tau, \nu)$ of the boundary value problem (3.1) satisfying $0 \le u_n \le \tilde{w}$ in Ω . The solution u_n satisfies

(3.7)
$$u_n + \mathbb{G}_V[f_n(u_n)] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu] = \tilde{w} \quad \text{in } \Omega$$

Put $w := u_{n+1} - u_n$. Then w is L_V perfect and tr $_V w = 0$. Consequently, by Lemma 2.6, tr $_V w_+ = 0$. Furthermore

$$-L_V w + f_{n+1}(u_{n+1}) - f_n(u_n) = 0$$
 in Ω .

By Kato's inequality

$$-L_V w_+ + (f_{n+1}(u_{n+1}) - f_n(u_n)) \operatorname{sign}_+ w \le 0 \quad \text{in } \Omega.$$

In the set $\{x \in \Omega : w_+ \ge 0\}$:

$$f_{n+1}(u_{n+1}) - f_n(u_n) \ge f_{n+1}(u_n) - f_n(u_n) \ge 0.$$

Thus w_+ is L_V subharmonic in Ω . As tr_V $w_+ = 0$, Lemma 2.3 yields $w_+ = 0$, i.e., $u_{n+1} \leq u_n$ in Ω .

(a) Let $u^{\#} = \lim u_n$, then $0 \le u^{\#} \le \tilde{w}$ in Ω . By Dini's Lemma, $f_n(u_n) \to f(u^{\#})$ a.e. in Ω . Therefore, by (3.7) and Fatou's lemma, we obtain (3.3).

By Theorems 2.1 and 2.2, we have

$$\int_{\Omega} \frac{\Phi_V}{\delta} \tilde{w} \, dx = \int_{\Omega} \frac{\Phi_V}{\delta} (\mathbb{K}_V[\nu] + \mathbb{G}_V[\tau]) \, dx \sim \|\tau\|_{\mathfrak{M}(\Omega;\Phi_V)} + \|\nu\|_{\mathfrak{M}(\partial\Omega)}$$

and
$$\int \frac{\Phi_V}{\delta} \mathbb{C}_{\mathbb{T}}[f_{-}(u_{-})] \, dx \sim \int_{\Omega} f_{-}(u_{-}) \Phi_{\mathbb{T}} \, dx$$

$$\int_{\Omega} \frac{\Phi_V}{\delta} \mathbb{G}_V[f_n(u_n)] \, dx \sim \int_{\Omega} f_n(u_n) \Phi_V \, dx$$

Therefore, multiplying (3.7) by Φ_V/δ and integrating over Ω , we obtain the following similarity relations

(3.8)
$$\int_{\Omega} u_n \frac{\Phi_V}{\delta} dx + \int_{\Omega} f_n(u_n) \Phi_V dx$$
$$\sim \int_{\Omega} u_n \frac{\Phi_V}{\delta} dx + \int_{\Omega} \frac{\Phi_V}{\delta} \mathbb{G}_V[f_n(u_n)] dx$$
$$= \int_{\Omega} \tilde{w} \frac{\Phi_V}{\delta} dx \sim \|\tau\|_{\mathfrak{M}(\Omega; \Phi_V)} + \|\nu\|_{\mathfrak{M}(\partial\Omega)}$$

Letting $n \to \infty$ and using Fatou's lemma, we obtain (3.2).

(b) Let $\zeta \in C_c^{\infty}(\Omega)$. Multiplying (3.7) by $-L_V \zeta$ and integrating over Ω , we obtain

$$-\int_{\Omega} u_n L_V \zeta \, dx - \int_{\Omega} \mathbb{G}_V[f_n(u_n)] L_V \zeta \, dx = -\int_{\Omega} \mathbb{G}_V[\tau] L_V \zeta \, dx.$$

We used the fact that $\mathbb{K}_V[\nu]$ is L_V harmonic. Further, for every $\tau \in \mathfrak{M}(\Omega; \Phi_V), -L_V \mathbb{G}_V[\tau] = \tau$, i.e.,

$$-\int_{\Omega} \mathbb{G}_V[\tau] L_V \zeta \, dx = \int_{\Omega} \zeta \, d\tau.$$

Hence

$$-\int_{\Omega} u_n L_V \zeta \, dx + \int_{\Omega} f_n(u_n) \zeta \, dx = \int_{\Omega} \zeta \, d\tau.$$

Recall that $\{u_n\}$ converges to $u^{\#}$ and is dominated by u_1 in $L^1(\Omega; \Phi_V/\delta)$ and $\{f_n(u_n)\}$ converges to $f(u^{\#})$ and - by (3.8) - is bounded in $L^1(\Omega; \Phi_V)$. Consequently, using Fatou's lemma,

$$-C\int_{\Omega}\zeta\,dx \le -\int_{\Omega}u^{\#}L_{V}\zeta\,dx + \int_{\Omega}f(u^{\#})\zeta\,dx \le \int_{\Omega}\zeta\,d\tau$$

for every $0 \leq \zeta \in C_c^{\infty}(\Omega)$ where $C \geq 0$ a constant independent of ζ . Thus there exists a bounded measure $\tau^{\#} \leq \tau$ such that (3.4) holds. Now, in Lemma 3.4 below, it is shown that $\tau^{\#} \geq \tau_d$. This is based only on the assumptions of the present theorem, the definition of $u^{\#}$ as the limit of the decreasing sequence $\{u_n\}$ and what is proved above. Therefore, as $\tau \geq 0$, we conclude that $\tau^{\#} \geq 0$. This completes the proof of part (b).

(c) By (3.3), $u^{\#} \leq \tilde{w}$. Since tr_V \tilde{w} exists and $u^{\#} \geq 0$, it follows that

$$\sup_{n} \int_{\partial D_{n}} P_{V,n}(x_{0}, y) | u^{\#}(y) | dS(y) \leq$$
$$\sup_{n} \int_{\partial D_{n}} P_{V,n}(x_{0}, y) \tilde{w}(y) dS(y) < +\infty$$

By (3.4), as $f(u^{\#}) \in L^1(\Omega; \Phi_V)$, the function $v := u^{\#} + \mathbb{G}_V[f(u^{\#})]$ satisfies $-L_V v = \tau$ and $v \ge 0$. Therefore, by Lemma 2.4 and (1.5), $\operatorname{tr}_V v =: \nu^{\#}$ exists, $\operatorname{tr}_V u^{\#} = \operatorname{tr}_V v$ and (3.5) holds. As $0 \le u^{\#} \le \tilde{w}$,

$$0 \le \nu^{\#} \le \operatorname{tr}_V \tilde{w} = \nu.$$

(d) Let w be a positive subsolution of (3.6). Then

$$-L_V w + f_n(w) \le -L_V w + f(w) \le \tau \quad \text{in } \Omega, \quad \text{tr}_V w \le \nu.$$

On the other hand, we have

$$-L_V u_n + f_n(u_n) = \tau$$
 in Ω , $\operatorname{tr}_V u_n = \nu$.

By Lemma 2.9, $w \leq u_n$ and thus $w \leq u^{\#}$. This proves (d).

Obviously, if (3.6) has a solution u then it is the largest subsolution of the problem so that $u^{\#} = u$.

Definition 3.2. A measure $\tau \in \mathfrak{M}_+(\Omega; \Phi_V)$ is a *good measure* with respect to f if there exists a solution u of equation (2.2) such that $f(u) \in L^1(\Omega; \Phi_V)$.

A couple of measures $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial \Omega)$ is a good couple with respect to f if there exists a solution u of problem (3.6).

The couple $(\tau^{\#}, \nu^{\#})$ that satisfies (3.5) is called the *reduced couple* of (τ, ν) .

Remark 3.3. We note that as a consequence of Theorem 3.1 parts (b) and (c):

(i) For every $\nu \in \mathfrak{M}_+(\partial\Omega)$ the reduced couple of $(0,\nu)$ is $(0,\nu^*)$ where ν^* is the largest good measure dominated by ν .

(ii) For every $\tau \in \mathfrak{M}_+(\Omega; \Phi_V)$ the reduced couple of $(\tau, 0)$ is $(\tau^*, 0)$ where τ^* is the largest good measure dominated by τ .

Notation. (a) Let $\tau \in \mathfrak{M}_+(\Omega; \Phi_V)$. Denote by $\tau^{\#}(\nu)$ the measure $\tau^{\#}$ in Theorem 3.1. In particular $\tau^{\#}(0)$ is the reduced measure of problem

$$L_V u + f(u) = \tau$$
 in Ω , $\operatorname{tr}_V u = 0$.

(b) Let λ be a Borel measure in Ω such that $\lambda = \lambda_{+} - \lambda_{-}$ where λ_{\pm} are positive Radon measures. It is well-known (see e.g. [4]) that λ has a unique representation of the form $\lambda = \lambda_{c} + \lambda_{d}$ where λ_{d} vanishes on sets of (Newtonian) capacity zero while λ_{c} is concentrated on a set of zero capacity. We say that λ_{d} is the *diffuse* part of λ while λ_{c} is the *concentrated* part of λ . If $\lambda = \lambda_{d}$ we say that λ is a diffuse measure. If $\lambda = \lambda_{c}$ we say that λ is a concentrated measure.

For the proof of the next theorem we need a version of [4, Lemma 4.1] suitable for the present problem. The proof is essentially the same as in [4], but some slight modifications are needed. For the convenience of the reader we provide the proof below.

Lemma 3.4. Let $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial \Omega)$. Then under the assumptions and with the notation of Theorem 3.1, we have

(3.9)
$$\tau^{\#} \ge \tau_d \quad and \quad (\tau^{\#})_d = \tau_d.$$

Proof. Let u_n be the solution of (3.1). Then $u_n \ge 0$, the sequence $\{u_n\}$ is decreasing and, as in Theorem 3.1, the function $u^{\#} := \lim u_n$ satisfies

(3.10)
$$-L_V u^{\#} + f(u^{\#}) = \tau^{\#} \quad \text{in } \Omega,$$

where $\tau^{\#}$ is a measure such that $\tau^{\#} \leq \tau$.

Denote $T_k(s) := \min(k, s), s \in \mathbb{R}$. By [5] and [4, Corollary 4.9],

$$\Delta T_k(u_n) \le \chi_{[u_n \le k]}(\Delta u_n)_d + ((\Delta u_n)_c)_+,$$

where χ_A denotes the characteristic function of $A \subset \mathbb{R}^N$. Since u_n satisfies (3.1), we obtain

$$(\Delta u_n)_d = -Vu_n + f_n(u_n) - \tau_d, \quad (\Delta u_n)_c = -\tau_c$$

As $\tau \geq 0$, these relations and the previous inequality yield

$$\Delta T_k(u_n) \le \chi_{[u_n \le k]}(-Vu_n + f_n(u_n) - \tau_d) \quad \text{in } \Omega,$$

and

$$-L_V T_k(u_n) \ge -V T_k(u_n) + \chi_{[u_n \le k]} (V u_n - f_n(u_n) + \tau_d)$$

= $-\chi_{[u_n > k]} V k - \chi_{[u_n \le k]} (f_n(u_n) - \tau_d).$

Since $\chi_{[u_n \leq k]} f_n(u_n) \leq f_n(T_k(u_n))$, it follows that (3.11) $-L_V T_k(u_n) + f_n(T_k(u_n)) \geq -\chi_{[u_n > k]} |V| u_n + \chi_{[u_n \leq k]} \tau_d$ in Ω . Let $\zeta \in C_c^{\infty}(\Omega)$ and $\zeta \ge 0$. Since $0 \le u_n \le u_1$, (3.11) yields

$$-\int_{\Omega} T_k(u_n) L_V \zeta \, dx + \int_{\Omega} f_n(T_k(u_n)) \zeta \, dx \ge$$
$$-\int_{\Omega} \chi_{[u_1 > k]} |V| u_1 \zeta \, dx + \int_{\Omega} \chi_{[u_1 \le k]} \zeta \, d\tau_d.$$

By the dominated convergence theorem, letting $n \to \infty$, we obtain

$$-\int_{\Omega} T_k(u^{\#}) L_V \zeta \, dx + \int_{\Omega} f(T_k(u^{\#})) \zeta \, dx \ge$$
$$-\int_{\Omega} \chi_{[u_1 > k]} |V| u_1 \zeta \, dx + \int_{\Omega} \chi_{[u_1 \le k]} \zeta \, d\tau_d.$$

Finally, letting $k \to \infty$, we obtain

$$-\int_{\Omega} u^{\#} L_V \zeta \, dx + \int_{\Omega} f(u^{\#}) \zeta \, dx \ge \int_{\Omega} \zeta \, d\tau_d.$$

In view of (3.10) this implies $\tau^{\#} \geq \tau_d$ and therefore $(\tau^{\#})_d \geq \tau_d$. As $\tau^{\#} \leq \tau$ we obtain $(\tau^{\#})_d = \tau_d$. This proves (3.9).

Theorem 3.5. Under the assumptions and with the notation of Theorem 3.1, the following statements hold.

(i) For every τ , $\nu^{\#} = \nu^*$ with ν^* as in Remark 3.3. Thus $\nu^{\#}$ does not depend on the data τ .

(ii) For every ν , $\tau^{\#} = \tau^*$ with τ^* as in Remark 3.3. Thus $\tau^{\#}$ does not depend on the boundary data ν .

(iii) Let $(0,0) \prec (\lambda,\sigma) \prec (\tau,\nu)$. Then problem

$$-L_V u + f(u) = \lambda \quad in \ \Omega, \quad \mathrm{tr}_V u = \sigma,$$

has a solution if and only if $(\lambda, \sigma) \prec (\tau^*, \nu^*)$.

Proof. (i) Let $u_n(\tau,\nu)$, $u^{\#}(\tau,\nu)$ and $(\tau^{\#},\nu^{\#})$ be as in Theorem 3.1. For simplicity, we write $u^* = u^{\#}(0,\nu)$ and $u^{\#} = u^{\#}(\tau,\nu)$. Since $0 \leq u_n(0,\nu) \leq u_n(\tau,\nu)$, it follows that $0 \leq u^* \leq u^{\#}$ in Ω . Therefore, using Theorem 3.1(c), we obtain

(3.12)
$$\nu^* = \operatorname{tr}_V u^* \le \operatorname{tr}_V u^\# = \nu^\# \le \nu.$$

Consequently u^* is a subsolution of problem

(3.13)
$$-L_V u + f(u) = 0 \text{ in } \Omega, \quad \text{tr}_V u = \nu^{\#}.$$

Obviously $u^{\#}$ is a supersolution of (3.13). Hence, by Lemma 2.10, there exists a unique solution \bar{v} of problem (3.13) and $0 \leq u^* \leq \bar{v} \leq u^{\#}$. By Theorem 3.1, u^* is the largest solution of the equation $-L_V u + f(u) = 0$ in Ω with L_V boundary trace $\leq \nu$. Thus $\bar{v} \leq u^*$ and consequently $\nu^{\#} = \operatorname{tr}_V \bar{v} \leq \operatorname{tr}_V u^* = \nu^*$. This inequality and (3.12) imply $\nu^{\#} = \nu^*$.

(ii) Here we denote by $R(\tau, \nu)$ the reduced couple of (τ, ν) . For given τ we denote by $\tau^{\#}(\nu)$ the first component of $R(\tau, \nu)$.

In view of Theorem 3.1(d),

$$R(\tau, \nu^{\#}) = R(\tau, \nu), \quad \tau^{\#}(\nu) = \tau^{\#}(\nu^{\#}).$$

Hence, as $\nu^{\#} = \nu^*$,

(3.14)
$$R(\tau,\nu) = R(\tau,\nu^*), \quad u^{\#}(\tau,\nu) = u^{\#}(\tau,\nu^*).$$

Therefore, in this part of the proof we may assume that $\nu = \nu^*$. Step I. If $\nu_1, \nu_2 \in \mathfrak{M}_+(\partial \Omega)$ and $\nu_1 \leq \nu_2$ then

(3.15)
$$\tau^{\#}(\nu_1) \ge \tau^{\#}(\nu_2).$$

Proof. The assumption implies that $\nu_1^* \leq \nu_2^*$. Therefore by (3.14), we may assume that ν_1, ν_2 are good measures with respect to (3.6).

Let λ be a measure such that $0 \leq \lambda \leq \tau$ and suppose that there exists a solution u_2 of

$$-Lu + f(u) = \lambda$$
 in Ω , $\operatorname{tr}_V u = \nu_2$.

Then u_2 is a supersolution of

(3.16)
$$-Lu + f(u) = \lambda \quad \text{in } \Omega, \quad \text{tr}_V u = \nu_1.$$

We also know that there exists $\lambda^{\#} \leq \lambda$ such that the following problem has a solution u_1 :

$$-Lu + f(u) = \lambda^{\#} \quad \text{in } \Omega, \quad \text{tr}_V u = \nu_1.$$

The function u_1 is a subsolution of (3.16) and, by Lemma 2.9, $u_1 \leq u_2$. Hence there exists a solution \bar{v} of (3.16). Clearly, \bar{v} is a subsolution of

$$-L_V u + f(u) = \tau \quad \text{in } \Omega, \quad \text{tr}_V u = \nu_1$$

and therefore, by Theorem 3.1 (d), $\lambda \leq \tau^{\#}(\nu_1)$. As $\lambda = \tau^{\#}(\nu_2)$ satisfies the conditions required above, this implies (3.15).

Step II. With ν^* as before we prove that

Let $u_0^{\#}$ be the solution of

$$-L_V u + f(u) = \tau^{\#}(0)$$
 in Ω , tr_V $u = 0$.

Then $u_0^{\#}$ is a subsolution of (3.6). By Theorem 3.1(d) and (3.14),

$$u_0^{\#} \le u^{\#}(\tau, \nu) = u^{\#}(\tau, \nu^*).$$

Let $w := u^{\#}(\tau, \nu^{*}) - u_{0}^{\#}$ so that $-L_{V}w + f(u^{\#}(\tau, \nu^{*})) - f(u_{0}^{\#}) = \tau^{\#}(\nu^{*}) - \tau^{\#}(0)$ in Ω . As $w \ge 0$, by the inverse maximum principle [8],

$$-(\Delta w)_c = (\tau^{\#}(\nu^*) - \tau^{\#}(0))_c \ge 0.$$

Moreover, by Lemma 3.4,

$$\tau^{\#}(\nu^*)_d = \tau^{\#}(0)_d = \tau_d.$$

Hence $\tau^{\#}(\nu^*) \geq \tau^{\#}(0)$. On the other hand, by step I, $\tau^{\#}(\nu^*) \leq \tau^{\#}(0)$. This proves (3.17).

(iii) This is a simple consequence of statements (i) and (ii) and Theorem 3.1(d).

Remark 3.6. Given a real function h on \mathbb{R} , denote by \hat{h} the function given by

$$\hat{h}(t) := -h(-t) \quad \forall t \in \mathbb{R}$$

Let f_n, τ, ν be as in Theorem 3.1. If w_n is the solution of the boundary value problem

$$-L_V w + \hat{f}_n(w) = \tau \quad \text{in } \Omega, \quad \text{tr}_V w = \nu,$$

then $z_n = -w_n$ satisfies

$$-L_V z_n + f_n(z_n) = -\tau$$
 in Ω , $\operatorname{tr}_V z_n = -\nu$.

Since \hat{f}_n has the same properties as f_n , the sequence $\{w_n\}$ has the same properties as the sequence $\{u_n\}$ in Theorem 3.1.

Accordingly, for $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial \Omega)$ the reduced measures for $-\tau$ and $-\nu$ and the corresponding reduced couple are given by

(3.18)
$$(-\tau)^{\#} := -(\tau_{\hat{f}}^{\#}), \quad (-\nu)^{\#} := -(\nu_{\hat{f}}^{\#}), \\ (-\tau, -\nu)^{\#} := -((\tau, \nu)_{\hat{f}}^{\#}) = ((-\tau)^{\#}, (-\nu)^{\#})$$

(The subscript \hat{f} above indicates that the reduced measure or couple is defined relative to this function.)

In this case $(-\tau)^{\#}$ and $(-\nu)^{\#}$ are the smallest good measures dominating $-\tau$ and $-\nu$ respectively. The relation between the solutions corresponding to the reduced couples is given below

$$u^{\#}(-\tau,-\nu) = -u_{\hat{f}}^{\#}(\tau,\nu).$$

Here again the notation $u_{\hat{f}}^{\#}$ indicates that the reduced couple and the corresponding solution is defined relative to \hat{f} .

In view of this remark, the results of Theorems 3.1 and 3.5, with obvious modifications, also apply to couples of negative measures.

4. Signed measures

Theorem 4.1. Let $\tau \in \mathfrak{M}(\Omega; \Phi_V)$ and $\nu \in \mathfrak{M}(\partial\Omega)$. Let $\tau_1, \tau_2 \in \mathfrak{M}_+(\Omega; \Phi_V)$ and $\nu_1, \nu_2 \in \mathfrak{M}_+(\partial\Omega)$ be measures such that

(4.1)
$$-\tau_1 \leq \tau \leq \tau_2, \quad -\nu_1 \leq \nu \leq \nu_2.$$

Let $\{f_n\}$ be a sequence of functions as in Theorem 3.1. Then the boundary value problem (3.1) has a unique solution $u_n = u_n(\tau, \nu)$.

The following statements hold.

(a) The sequence $\{u_n\}$ is bounded in $W^{1,p}_{loc}(\Omega)$ for every $p \in [1, \frac{N}{N-1})$. Consequently every subsequence has a limit point in the sense of convergence in $L^p_{loc}(\Omega)$ and convergence a.e. in Ω .

If $\{u_{n_k}\}$ is a subsequence of $\{u_n\}$ converging to \tilde{u} a.e. then:

(4.2)

$$(i) \quad u_{n_k} \to \tilde{u} \quad in \ L^1(\Omega; \Phi_V/\delta),$$

$$(ii) \quad f_{n_k}(u_{n_k}) \to f(\tilde{u}) \quad a.e. \ in \ \Omega.$$

(b) The following inequality holds

(4.3)
$$\|\tilde{u}\|_{L^1(\Omega;\Phi_V/\delta)} + \|f(\tilde{u})\|_{L^1(\Omega;\Phi_V)} \le C \sum_{i=1,2} (\|\tau_i\|_{\mathfrak{M}(\Omega;\Phi_V)} + \|\nu_i\|_{\mathfrak{M}(\partial\Omega)}).$$

(c) There exist $\tilde{\tau} \in \mathfrak{M}(\Omega; \Phi_V)$ and $\tilde{\nu} \in \mathfrak{M}(\partial \Omega)$ such that

(4.4)
$$(-\tau_1)^{\#} \le \tilde{\tau} \le \tau_2^{\#}, \quad (-\nu_1)^{\#} \le \tilde{\nu} \le \nu_2^{\#}$$

and

$$\tilde{u} + \mathbb{G}_V[f(\tilde{u})] = \mathbb{G}_V[\tilde{\tau}] + \mathbb{K}_V[\tilde{\nu}] \quad in \ \Omega.$$

Thus \tilde{u} is the solution of the boundary value problem

(4.5)
$$-L_V \tilde{u} + f(\tilde{u}) = \tilde{\tau} \quad in \ \Omega, \quad \operatorname{tr}_V \tilde{u} = \tilde{\nu}$$

Proof. Let $v_{2,n}$ denote the solution of the boundary value problem

(4.6)
$$-L_V v + f_n(v) = \tau_2 \quad \text{in } \Omega, \quad \text{tr}_V v = \nu_2,$$

and $v_{1,n}$ denote the solution of the boundary value problem

(4.7)
$$-L_V v + f_n(v) = -\tau_1 \text{ in } \Omega, \quad \text{tr}_V v = -\nu_1.$$

By Theorem 3.1 and Remark 3.6, $v_{2,n}$ and $-v_{1,n}$ are positive and the following inequalities hold

(4.8)
$$\|v_{i,n}\|_{L^{1}(\Omega;\Phi_{V}/\delta)} + \|f_{n}(v_{i,n})\|_{L^{1}(\Omega;\Phi_{V})} \\ \leq C(\|\tau_{i}\|_{\mathfrak{M}(\Omega;\Phi_{V})} + \|\nu_{i}\|_{\mathfrak{M}(\partial\Omega)}), \quad i = 1, 2.$$

Moreover,

$$v_{1,n} \uparrow v_1^{\#}(-\tau, -\nu) =: w_1, \quad v_{2,n} \downarrow v_2^{\#}(\tau, \nu) =: w_2 \quad \text{a.e. in } \Omega,$$

 $f(v_i^{\#}) \in L^1(\Omega; \Phi_V), \quad i = 1, 2,$

and there exist

$$\tau_2^{\#}, (-\tau_1)^{\#} \in \mathfrak{M}(\Omega; \Phi_V), \quad \nu_2^{\#}, (-\nu_1)^{\#} \in \mathfrak{M}(\partial\Omega)$$

such that

(4.9)
$$-L_V w_2 + f(w_2) = \tau_2^{\#} \quad \text{in } \Omega, \qquad \text{tr}_V w_2 = \nu_2^{\#}, \\ -L_V w_1 + f(w_1) = (-\tau_1)^{\#} \quad \text{in } \Omega, \qquad \text{tr}_V w_1 = (-\nu_1)^{\#}.$$

The monotone convergence of the sequences $\{v_{i,n}\}$ and (4.8) imply

(4.10)
$$v_{i,n} \to w_i \text{ in } L^1(\Omega; \Phi_V/\delta)$$

and, by Dini's lemma,

(4.11)
$$f_n(v_{i,n}) \to f(w_i)$$
 a.e. in Ω , $i = 1, 2$.

(a) By (3.1) and (4.6), we have

(4.12)
$$-L_V(u_n - v_{2,n}) + f_n(u_n) - f_n(v_{2,n}) = \tau - \tau_2 \quad \text{in } \Omega$$
$$\operatorname{tr}_V(u_n - v_{2,n}) = \nu - \nu_2.$$

By (4.1), $\tau - \tau_2 \leq 0$ and $\nu - \nu_2 \leq 0$. Therefore, using Kato's inequality, we obtain $-L_V(u_n - v_{2,n})_+ \leq 0$ and, by Lemma 2.6, tr $_V(u_n - v_{2,n})_+ = 0$. By Lemma 2.3, $(u_n - v_{2,n})_+ = 0$, which implies

$$(4.13) u_n \le v_{2,n} \quad \text{in } \Omega$$

Similarly, by (3.1) and (4.7), we obtain

$$(4.14) v_{1,n} \le u_n \quad \text{in } \Omega$$

(Recall that $v_{1,n} \leq 0$.) Thus

(4.15)
$$|u_n| \le \max\{-v_{1,n}, v_{2,n}\}$$
 in Ω

This inequality and the monotonicity of f_n imply that

(4.16)
$$|f_n(u_n)| \le \max\{f_n(-v_{1,n}), f_n(v_{2,n})\}$$
 in Ω

Inequalities (4.15), (4.16) and (4.8) together with the fact that V is locally bounded in Ω imply that $\{|Vu_n + f_n(u_n)|\}$ is bounded in $L^1_{\text{loc}}(D)$. Consequently $\{u_n\}$ is bounded in $W^{1,p}_{\text{loc}}(\Omega)$ for every $p \in [1, \frac{N}{N-1})$. Therefore $\{u_n\}$ is precompact in $L^p_{\text{loc}}(\Omega)$. Hence there exists a subsequence $\{u_{n_k}\}$) that converges to a function \tilde{u} in $L^p_{\text{loc}}(\Omega)$ and a.e. in Ω .

By the generalized dominated convergence theorem, the convergence of $\{u_{n_k}\}$ to \tilde{u} a.e. in Ω , (4.15) and (4.10) imply (4.2) (i).

The assumption $f_n^{\pm} \uparrow f^{\pm}$ and Dini's lemma imply (4.2) (ii).

To simplify the presentation, in the remainder of the proof we assume that $\{u_n\}$ is a sequence converging a.e. to \tilde{u} .

(b) By (4.15), (4.8) and (4.2)(i), we obtain

$$\begin{split} \|\tilde{u}\|_{L^{1}(\Omega;\Phi_{V}/\delta)} &= \lim_{n \to \infty} \|u_{n}\|_{L^{1}(\Omega;\Phi_{V}/\delta)} \\ &\leq C \sum_{i=1,2} (\|\tau_{i}\|_{\mathfrak{M}(\Omega;\Phi_{V})} + \|\nu_{i}\|_{\mathfrak{M}(\partial\Omega)}). \end{split}$$

By Fatou's lemma, (4.16), (4.8) and (4.2) yield

$$\begin{split} \|f(\tilde{u})\|_{L^{1}(\Omega;\Phi_{V})} &\leq \lim_{n \to \infty} \|f_{n}(u_{n})\|_{L^{1}(\Omega;\Phi_{V})} \\ &\leq \limsup_{n \to \infty} \|\max\{f_{n}(-v_{1,n}), f_{n}(v_{2,n})\}\|_{L^{1}(\Omega;\Phi_{V})} \\ &\leq C\sum_{i=1,2} (\|\tau_{i}\|_{\mathfrak{M}(\Omega;\Phi_{V})} + \|\nu_{i}\|_{\mathfrak{M}(\partial\Omega)}). \end{split}$$

This proves (4.3).

(c) Inequalities (4.13) and (4.14) imply $w_1 \leq \tilde{u} \leq w_2$ a.e. in Ω . Let $\zeta \in C_c^{\infty}(\Omega)$ and $\zeta \geq 0$. By (4.12), (4.17) $-\int_{\Omega} (v_{2,n} - u_n) L_V \zeta \, dx + \int_{\Omega} (f_n(v_{2,n}) - f_n(u_n)) \zeta \, dx = \int_{\Omega} \zeta \, d(\tau_2 - \tau).$

By (4.13), (4.11) and Fatou's lemma,

$$\int_{\Omega} (f(w_2) - f(\tilde{u}))\zeta \, dx \le \liminf_{n \to \infty} \int_{\Omega} (f_n(v_{2,n}) - f_n(u_n))\zeta \, dx.$$

Therefore, by (4.17) and (4.10),

$$(4.18) \quad -\int_{\Omega} (w_2 - \tilde{u}) L_V \zeta dx + \int_{\Omega} (f(w_2) - f(\tilde{u})) \zeta dx \le \int_{\Omega} \zeta d(\tau_2 - \tau).$$

By (4.17), we have

$$\left|\int_{\Omega} (v_{2,n} - u_n) L_V \zeta dx\right| \leq \int_{\Omega} (f_n(v_{2,n}) - f_n(u_n)) \zeta dx + \int_{\Omega} \zeta d(\tau_2 - \tau).$$

By (4.8), (4.15) and (4.16),

$$0 \leq \int_{\Omega} (f_n(v_{2,n}) - f_n(u_n))\zeta \, dx + \int_{\Omega} \zeta d(\tau_2 - \tau)$$

$$\leq c \sup(\zeta/\Phi_V) \Big(\sum_{i=1,2} (\|\tau_i\|_{\mathfrak{M}(\Omega;\Phi_V)} + \|\nu_i\|_{\mathfrak{M}(\partial\Omega)} + \|\tau\|_{\mathfrak{M}(\Omega;\Phi_V)} \Big)$$

$$\leq C \sup(\zeta/\Phi_V).$$

Since

$$\lim_{n \to \infty} \int_{\Omega} (v_{2,n} - u_n) L_V \zeta dx = \int_{\Omega} (w_2 - \tilde{u}) L_V \zeta dx,$$

it follows that

$$\left|\int_{\Omega} (w_2 - \tilde{u}) L_V \zeta dx\right| \le C \sup(\zeta/\Phi_V).$$

Hence, by (4.18), there exists a measure $\lambda \leq \tau_2 - \tau$ such that $\lambda \in \mathfrak{M}(\Omega; \Phi_V)$ and

$$-\int_{\Omega} (w_2 - \tilde{u}) L_V \zeta \, dx + \int_{\Omega} (f(w_2) - f(\tilde{u})) \zeta \, dx = \int_{\Omega} \zeta \, d\lambda \quad \forall \zeta \in C_c^{\infty}(\Omega)$$

or equivalently,

(4.19)
$$-L_V(w_2 - \tilde{u}) + f(w_2) - f(\tilde{u}) = \lambda \quad \text{in } \Omega$$

Consequently, by (4.9),

(4.20)
$$-L_V \tilde{u} + f(\tilde{u}) = \tilde{\tau}$$
 in Ω where $\tilde{\tau} := \tau_2^{\#} - \lambda \ge \tau_2^{\#} - \tau_2 + \tau$.

Next, by (3.1) and (4.7), for $\zeta \in C_c^{\infty}(\Omega)$,

$$-\int_{\Omega} (u_n - v_{1,n}) L_V \zeta \, dx + \int_{\Omega} (f_n(u_n) - f_n(v_{1,n})) \zeta \, dx = \int_{\Omega} \zeta \, d(\tau_1 + \tau).$$

By the same argument as above it follows that there exists a measure $\lambda' \leq \tau_1 + \tau$ such that $\lambda' \in \mathfrak{M}(\Omega; \Phi_V)$ and

(4.21)
$$-L_V(\tilde{u} - w_1) + f(\tilde{u}) - f(w_1) = \lambda' \text{ in } \Omega.$$

Consequently, by (4.9),

(4.22)
$$\tilde{\tau} = \lambda' + (-\tau_1)^{\#} \le \tau_1 + \tau + (-\tau_1)^{\#}.$$

Next we show that $\tilde{\tau}$ satisfies (4.4). By Lemma 3.4,

(4.23)
$$(\tau_2^{\#})_d = (\tau_2)_d, \quad ((-\tau_1)^{\#})_d = -(\tau_1)_d.$$

Therefore, by (4.20), $\tilde{\tau}_d \ge \tau_d$ and by (4.22), $\tilde{\tau}_d \le \tau_d$. Thus (4.24) $\tau_d = \tilde{\tau}_d$.

By (4.19) and (4.20),

$$-L_V(w_2 - \tilde{u}) + f(w_2) - f(\tilde{u}) = \tau_2^{\#} - \tilde{\tau}$$
 in Ω .

Since \tilde{u} and w_2 are diffuse and f(0) = 0, it follows that

(4.25)
$$(-\Delta(w_2 - \tilde{u}))_c = (\tau_2^{\#} - \tilde{\tau})_c.$$

As $w_2 - \tilde{u} \ge 0$, by the inverse maximum principle [8], $(-\Delta(w_2 - \tilde{u}))_c \ge 0$. Consequently,

$$(\tau_2^{\#} - \tilde{\tau})_c \ge 0$$

As $\tau \leq \tau_2$, (4.23), (4.24) yield,

$$(\tau_2^{\#})_d = (\tau_2)_d \ge \tau_d = \tilde{\tau}_d.$$

This inequality and (4.25) imply

(4.26)
$$\tilde{\tau} \le \tau_2^{\#}.$$

Similarly by (4.21) and (4.22),

$$L_V(\tilde{u} - w_1) + f(\tilde{u}) - f(w_1) = \tilde{\tau} - (-\tau_1)^{\#}$$
 in Ω

Since $\tilde{u} - w_1 \ge 0$, another application of the inverse maximum principle yields $(-\Delta(\tilde{u} - w_1))_c \ge 0$ and consequently

(4.27)
$$(\tilde{\tau} - (-\tau_1)^{\#})_c \ge 0.$$

As $-\tau_1 \leq \tau$, (4.23), (4.24) imply

$$\tilde{\tau}_d = \tau_d \ge (-\tau_1)_d = ((-\tau_1)^{\#})_d.$$

This and (4.27) yield

$$\tilde{\tau} \ge (-\tau_1)^\#.$$

Finally, this and (4.26) imply (4.4) with respect to $\tilde{\tau}$.

It remains to show that \tilde{u} has an L_V boundary trace and that the second inequality in (4.4) holds.

By (4.19) $-L_V(w_2 - \tilde{u}) = \mu$ where $\mu \in \mathfrak{M}(\Omega; \Phi_V)$. Since $w_2 - \tilde{u} \ge 0$, by Lemma 2.5, $w_2 - \tilde{u}$ has an L_V boundary trace, say σ , and $w_2 - \tilde{u} = \mathbb{G}_V[\mu] + \mathbb{K}_V[\sigma]$. Obviously $\sigma \ge 0$. Therefore

$$\operatorname{tr}_V \tilde{u} = \operatorname{tr}_V w_2 - \sigma \le \operatorname{tr}_V w_2 = \nu_2^{\#}.$$

Similarly, starting with (4.21) we conclude that there exists $\sigma' \in \mathfrak{M}(\partial\Omega)$ such that $\operatorname{tr}_V(\tilde{u} - w_1) = \sigma' \geq 0$. Therefore

$$\operatorname{tr}_V \tilde{u} = \sigma' + \operatorname{tr}_V w_1 \ge \operatorname{tr}_V w_1 = (-\nu_1)^{\#}.$$

This completes the proof.

The theorem is complemented by the following consequence of [17, Corollary 3.7] (see Lemma 2.10).

Proposition 4.2. Let $(\lambda_i, \sigma_i) \in \mathfrak{M}(\Omega; \Phi_V) \times \mathfrak{M}(\partial\Omega)$, i = 1, 2. Suppose that these are good couples with respect to (1.1) and that $(\lambda_1, \sigma_1) \prec (\lambda_2, \sigma_2)$. Then, every couple (λ, σ) such that

(4.28)
$$(\lambda_1, \sigma_1) \prec (\lambda, \sigma) \prec (\lambda_2, \sigma_2)$$

is a good couple.

Proof. Let v_i be the solution corresponding to the couple (λ_i, σ_i) , i = 1, 2 and let (λ, σ) be as in (4.28). Then v_2 is a supersolution and v_1 a subsolution of equation $-L_V u + f(u) = \lambda$ and $\operatorname{tr}_V v_1 \leq \sigma \leq \operatorname{tr}_V v_2$. Therefore the stated result is a consequence of Lemma 2.10).

Remark. As mentioned before, Theorem 4.1 and Proposition 4.2 imply Theorem 1.4. However we emphasize that, in contrast to Theorem 4.1, in Proposition 4.2 (λ_i, σ_i) may be couples of signed measures.

Proposition 4.3. In addition to the assumptions of Theorem 4.1, assume that (τ_2, ν_2) and $(-\tau_1, -\nu_1)$ are good couples.

Suppose that $(-\tau_1, -\nu_1) \prec (\tau, \nu) \prec (\tau_2, \nu_2)$. (By the previous result, (τ, ν) is a good couple.) Let u be the solution of problem (2.3) and let u_n denote the solution of the 'approximating' problem

$$-L_V u + f_n(u) = \tau \quad in \ \Omega, \quad \text{tr}_V u = \nu.$$

If Φ_V satisfies the additional condition

(4.29)
$$\int_{\Sigma_{\beta}} \Phi_V^2 / \delta \ dS \to 0 \quad as \ \beta \to 0$$

then $u_n \to u$, i.e. $\tilde{u} = u$.

Proof. We use the notation in the proof of Theorem 4.1.

Let $v_{1,n}$ and $v_{2,n}$ be the solutions of (4.7) and (4.6). Then

$$(4.30) v_{1,n} \le u_n \le v_{2,n} \quad \text{in } \Omega.$$

The sequences $\{v_{i,n}\}$, i = 1, 2 satisfy (4.10) and (4.11). In addition, by Proposition A.1 (see Appendix),

(4.31)
$$f_n(v_{i,n}) \to f(w_i) \text{ in } L^1(\Omega; \Phi_V).$$

By (4.30) and the monotonicity of f_n ,

$$f_n(v_{1,n}) \le f_n(u_n) \le f_n(v_{2,n}) \quad \text{in } \Omega.$$

Therefore, taking a subsequence for which (4.2) holds, the (generalized) dominated convergence theorem implies $f_n(u_n) \to f(\tilde{u})$ in $L^1(\Omega; \Phi_V)$ as $n \to \infty$. Hence, by Theorem 2.2,

$$\mathbb{G}_{V}[f_{n}(u_{n})] \to \mathbb{G}_{V}[f(\tilde{u})] \quad \text{in} \ L^{1}(\Omega; \Phi_{V}/\delta).$$

By (4.2), $u_{n} \to \tilde{u}$ in $L^{1}(\Omega; \Phi_{V}/\delta)$. As

$$u_n + \mathbb{G}_V[f_n(u_n)] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$$
 in Ω

we conclude that

$$\tilde{u} + \mathbb{G}_V[f(\tilde{u})] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$$
 in Ω .

Thus \tilde{u} is a solution of (2.3). By uniqueness (Lemma 2.9), $\tilde{u} = u$. \Box

Proof of Theorem 1.5. Since f vanishes on $(-\infty, 0]$, if w is a real function on Ω then

(4.32)
$$f(w) = f(w_{+}) + f(-w_{-}) = f(w_{+}).$$

Suppose that (τ, ν) is a good couple, i.e. (3.6) has a solution u such that $f(u) \in L^1(\Omega; \Phi_V)$.

Let u_n and \tilde{u} be as in Theorem 4.1. In view of (4.32), $\{u_n\}$ is decreasing (by Lemma 2.9) and $\tilde{u} = \lim u_n$.

Let w be a subsolution of (3.6) such that $f(w) \in L^1(\Omega; \Phi_V)$. Then

$$-L_V w + f_n(w) \le -L_V w + f(w_+) \le \tau \quad \text{in } \Omega, \quad \text{tr}_V w \le \nu.$$

As u_n satisfies (3.1), Lemma 2.9 implies that $w \leq u_n$. Thus $w \leq \tilde{u}$ and, in particular,

$$(4.33) u \le \tilde{u} \quad \text{in } \Omega.$$

Let $v := \tilde{u} - u$. Then $v \ge 0$, Δv is a measure and therefore, by the inverse maximum principle,

$$(-\Delta v)_c = (\tilde{\tau} - \tau)_c \ge 0.$$

As in the proof of Theorem 4.1 (see (4.24)) $\tau_d = \tilde{\tau}_d$. Therefore

$$\tau \leq \tilde{\tau}.$$

In addition, by (4.33),

$$\nu = \operatorname{tr}_V u \le \operatorname{tr}_V \tilde{u} = \tilde{\nu}.$$

Thus

(4.34)
$$(\tau, \nu) \prec (\tilde{\tau}, \tilde{\nu}).$$

If $(\tau_1, \nu_1) := (\tau_-, \nu_-)$ and $(\tau_2, \nu_2) := (\tau_+, \nu_+)$ then τ, ν satisfy (4.1) and by Theorem 4.1(c),

(4.35)
$$(\tilde{\tau}, \tilde{\nu}) \prec (\tau_+, \nu_+)^{\#}.$$

Hence, by (4.34), we obtain (1.15).

Conversely, assume that (τ, ν) satisfies (1.15). Recall that every couple of negative measures is good relative to f. Therefore, by Proposition 4.2, the relation

$$-(\tau_{-},\nu_{-})\prec(\tau,\nu)\prec(\tau_{+},\nu_{+})^{\#}$$

implies that (τ, ν) is a good couple.

Project GA22-17403S.

The last assertion of the theorem is obvious.

Acknowledgement. The research of M. Bhakta is partially supported by the DST Swarnajaynti fellowship (SB/SJF/2021-22/09). The research of P.-T. Nguyen was supported by Czech Science Foundation,

Conflict of interest. The authors declare that they have no conflicts of interest to this work.

APPENDIX A.

We prove an auxilliary result that is used in the proof of Proposition 4.3.

Proposition A.1. Assume (A1), (A2), (C1) and (4.29) hold. Suppose that $(\tau, \nu) \in \mathfrak{M}_+(\Omega; \Phi_V) \times \mathfrak{M}_+(\partial \Omega)$ is a good couple of measures. Let u be the corresponding solution of problem (3.6) and u_n be the solution of (3.1). Then

$$u_n \to u$$
 in $L^1(\Omega; \Phi_V/\delta)$, $f_n(u_n) \to f(u)$ in $L^1(\Omega; \Phi_V)$.

The proof is based on the following lemma that was established in [20] for a more restricted class of potentials.

Lemma A.2. Assume (A1), (A2), (C1) and (4.29) hold. Let $\tau \in \mathfrak{M}(\Omega; \Phi_V)$ and let λ_V be the eigenvalue of $-L_V$ corresponding to Φ_V . Then

(a.1)
$$\lambda_V \int_{\Omega} \mathbb{G}_V[\tau] \Phi_V \, dx = -\int_{\Omega} \mathbb{G}_V[\tau] L_V \Phi_V \, dx = \int_{\Omega} \Phi_V \, d\tau.$$

Proof. By linearity, we may assume that $\tau \ge 0$. For $\beta > 0$, put

$$I_{\beta}(\tau) := -\lambda_V \int_{D_{\beta}} \mathbb{G}_V[\tau] \Phi_V \, dx + \int_{D_{\beta}} \Phi_V \, d\tau$$

where $D_{\beta} = \{x \in \Omega : \delta(x) > \beta\}$. To prove (a.1) we show that (a.2) $\lim_{\beta \to 0} I_{\beta}(\tau) = 0.$

By Theorem 2.2 (ii),

$$I_0(\tau) \le C \int_{\Omega} \Phi_V d\tau.$$

Given $\epsilon > 0$ we choose $0 < \gamma$ sufficiently small so that, for $\tau_{\gamma} := \tau \mathbf{1}_{\Omega_{\gamma}}$,

(a.3) $I_0(\tau_\gamma) < \epsilon.$

Therefore it is sufficient to prove

$$\lim_{\beta \to 0} I_{\beta}(\tau - \tau_{\gamma}) = 0.$$

Thus it is sufficient to prove (a.2) when τ has compact support in Ω .

Let $\beta_{\tau} = \frac{1}{2} \text{dist}(\text{supp } \tau, \partial \Omega)$ and let $\beta \in (0, \beta_{\tau})$. Applying Green's theorem in D_{β} , we obtain

$$\lambda_V \int_{D_{\beta}} \mathbb{G}_V[\tau] \Phi_V dx = -\int_{D_{\beta}} \mathbb{G}_V[\tau] L_V \Phi_V dx$$
$$= \int_{D_{\beta}} \Phi_V d\tau + \int_{\Sigma_{\beta}} \frac{\partial \mathbb{G}_V[\tau]}{\partial \mathbf{n}} \Phi_V dS(x) - \int_{\Sigma_{\beta}} \frac{\partial \Phi_V}{\partial \mathbf{n}} \mathbb{G}_V[\tau] dS(x).$$

Thus

(a.4)
$$I(\beta) = -\int_{\Sigma_{\beta}} \frac{\partial \mathbb{G}_{V}[\tau]}{\partial \mathbf{n}} \Phi_{V} dS(x) + \int_{\Sigma_{\beta}} \frac{\partial \Phi_{V}}{\partial \mathbf{n}} \mathbb{G}_{V}[\tau] dS(x).$$

Note that

$$G_V(x,y) \sim \Phi_V(x) \quad \forall (x,y) \in \Omega_{\beta_\tau} \times \operatorname{supp} \tau.$$

Therefore

(a.5)
$$\mathbb{G}_V[\tau](x) = \int_{\Omega} G_V(x, y) d\tau(y) \sim \Phi_V(x), \quad \forall x \in \Sigma_{\beta}.$$

By interior elliptic estimates, for every $x \in \Sigma_{\beta}$,

$$\left|\frac{\partial \Phi_V}{\partial \mathbf{n}}(x)\right| \le C \sup_{|\xi-x| < \beta/4} \Phi_V(\xi) \beta^{-1}.$$

Therefore by Harnack's inequality, we deduce

$$\left. \frac{\partial \Phi_V}{\partial \mathbf{n}}(x) \right| \le C \Phi_V(x) \beta^{-1} \quad \forall x \in \Sigma_\beta.$$

Hence, by (a.5) and assumption (4.29),

$$\lim_{\beta \to 0} \int_{\Sigma_{\beta}} \frac{\partial \Phi_{V}}{\partial \mathbf{n}} \mathbb{G}_{V}[\tau] \, dS(x) = 0.$$

In $D_{\beta_{\tau}}$: $G_V[\tau]$ is L_V harmonic and $G_V[\tau] \sim \Phi_V$. Therefore the same argument as above yields,

(a.6)
$$\lim_{\beta \to 0} \int_{\Sigma_{\beta}} \frac{\partial \mathbb{G}_{V}[\tau]}{\partial \mathbf{n}} \Phi_{V} \, dS(x) = 0.$$

Combining (a.4) – (a.6), we obtain (a.2) for measures τ with compact support. In view of previous remarks, this implies (a.2) for any measure $\tau \in \mathfrak{M}(\Omega; \Phi_V)$. This in turn implies (a.1).

Proof of Proposition A.1. By Theorem 3.1(d), $u = u^{\#}$. From the proof of Theorem 3.1, $u_n \geq 0$ satisfies (3.7), $u_n \downarrow u^{\#} = u$ and $f_n(u_n) \to f(u)$ a.e. in Ω . By (3.7), $u_n \leq \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu] \in L^1(\Omega; \Phi_V/\delta)$. Therefore, by the dominated convergence theorem,

$$u_n \to u$$
 in $L^1(\Omega; \Phi_V / \delta)$.

By Lemma A.2 with τ replaced by $f_n(u_n)$ (recall that f_n is a bounded function) we have,

(a.7)
$$\int_{\Omega} f_n(u_n) \Phi_V \, dx = \lambda_V \int_{\Omega} \mathbb{G}_V[f_n(u_n)] \Phi_V \, dx.$$

Since u_n is the solution of (3.1) it satisfies

$$u_n + \mathbb{G}_V[f_n(u_n)] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu].$$

Multiplying this equality by $\lambda_V \Phi_V$ and using (a.7) we obtain

$$\int_{\Omega} f_n(u_n) \Phi_V dx = \lambda_V \int_{\Omega} \mathbb{G}_V[f_n(u_n)] \Phi_V dx$$
$$= -\lambda_V \int_{\Omega} u_n \Phi_V dx + \lambda_V \int_{\Omega} \mathbb{G}_V[\tau] \Phi_V dx + \lambda_V \int_{\Omega} \mathbb{K}_V[\nu] \Phi_V dx.$$

Hence,

(a.8)
$$\lim_{n \to \infty} \int_{\Omega} f_n(u_n) \Phi_V dx = -\lambda_V \int_{\Omega} u \Phi_V dx + \lambda_V \int_{\Omega} \mathbb{G}_V[\tau] \Phi_V dx + \lambda_V \int_{\Omega} \mathbb{K}_V[\nu] \Phi_V dx.$$

Since u is the solution of (3.6), $f(u) \in L^1(\Omega; \Phi_V)$ and $u + \mathbb{G}_V[f(u)] = \mathbb{G}_V[\tau] + \mathbb{K}_V[\nu]$. In addition, by Lemma A.2,

$$\int_{\Omega} f(u) \Phi_V \, dx = \lambda_V \int_{\Omega} \mathbb{G}_V[f(u)] \Phi_V \, dx.$$

Therefore, as before,

(a.9)
$$\int_{\Omega} f(u)\Phi_V dx$$
$$= -\lambda_V \int_{\Omega} u\Phi_V dx + \lambda_V \int_{\Omega} \mathbb{G}_V[\tau]\Phi_V dx + \lambda_V \int_{\Omega} \mathbb{K}_V[\nu]\Phi_V dx.$$

By (a.8) and (a.9), $||f_n(u_n)||_{L^1(\Omega;\Phi_V)} \to ||f(u)||_{L^1(\Omega;\Phi_V)}$. As $f_n(u_n) \to f(u)$ a.e. in Ω , it follows that

$$f_n(u_n) \to f(u)$$
 in $L^1(\Omega; \Phi_V)$.

The proof is complete.

References

- A. Ancona, Théorie du potentiel sur les graphes et les vari 'et 'es, in Springer Lecture Notes in Math., vol.1427, (ed. P.L. Hennequin) (1988) pp.1--112.
- [2] A. Ancona and M. Marcus, Positive solutions of a class of semilinear equations with absorption and Schrödinger equations, J. Math. Pures Appl. (9) 104 (2015), no. 3, 587–618.
- [3] M. Bhakta, M. and M. Marcus, Reduced limit for semilinear boundary value problems with measure data, J. Differential Equations 256 (2014), no. 8, 2691– 2710.
- [4] H. Brezis, M. Marcus and A. C. Ponce, Nonlinear elliptic equations with measures revisited, Mathematical aspects of nonlinear dispersive equations, Ann. of Math. Stud., 163, Princeton Univ. Press, Princeton, NJ, 2007, 55–109.
- [5] H. Brezis and A. C. Ponce, Kato's inequality when Δu is a measure, C. R. Math. Acad. Sci. Paris, Ser. I 338 (2004) 599--604.
- [6] H. Brezis and A. C. Ponce, *Reduced measures on the boundary*, J. Funct. Anal. 229 (2005), no. 1, 95–120.
- [7] H. Chen and L. Véron, Weak solutions of semilinear elliptic equations with Leray-Hardy potentials and measure data, Math. Eng. 1 (2019), no. 3, 391– 418.
- [8] L. Dupaigne and A. C. Ponce, Singularities of positive supersolutions in elliptic PDEs, Selecta Math. (N.S.) 10 (2004), no. 3, 341–358.
- [9] E. B. Dynkin, Superdiffusions and Positive Solutions of Nonlinear Partial Differential Equations, Amer. Math. Soc., Providence, RI, 2004.
- [10] K. T. Gkikas and L. Véron, Boundary singularities of solutions of semilinear elliptic equations with critical Hardy potentials, Nonlinear Anal. 121 (2015), 469–540.
- [11] D. Gómez-Castro and J. L. Vázquez, The fractional Schrödinger equation with singular potential and measure data, Discrete Contin. Dyn. Syst. 39 (2019), no. 12, 7113–7139.
- [12] J. F. Le Gall, The Brownian snake and solutions of $\Delta u = u^2$ in a domain, Probab. Th. Rel. Fields 102, 393-432 (1995).
- [13] J. F. Le Gall, Spatial branching processes, random snakes and partial differential equations, Birkhäuser, Basel/Boston/Berlin, 1999.

- [14] M. Marcus, Complete classification of the positive solutions of $-\Delta u + u^q = 0$, J. d'Anal. Math. **117** (2012), 187-220.
- [15] M. Marcus, Estimates of Green and Martin kernels for Schródinger operators with singular potential in Lipschitz domains, Ann. I. H. Poincar 'e AN 36 (2019) 1183—1200.
- [16] M. Marcus, Estimates of sub and super solutions of Schrödinger equations with very singular potentials, arXiv:1912.01283 (version 2).
- [17] M. Marcus, Boundary value problems with signed measure data for semilinear Schrödinger equations, arXiv:2305.10370.
- [18] M. Marcus and V. Moroz, V, Moderate solutions of semilinear elliptic equations with Hardy potential under minimal restrictions on the potential, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) 18 (2018), no. 1, 39–64.
- [19] M. Marcus and P.-T. Nguyen, Moderate solutions of semilinear elliptic equations with Hardy potential, Ann. Inst. H. Poincaré Anal. Non Linéaire 34 (2017), no. 1, 69–88.
- [20] M. Marcus and P.-T. Nguyen, Schrödinger equations with singular potentials: linear and nonlinear boundary value problems, Math. Ann. 374 (2019), no. 1-2, 361–394.
- [21] M. Marcus and A. C. Ponce, Reduced limits for nonlinear equations with measures, J. Functional An. 258 (2010), 2316-2372
- [22] M. Marcus and L. Véron, The boundary trace of positive solutions of semilinear elliptic equations: the subcritical case, Arch. Rational Mech. Anal. 144 (1998), 201-231.
- [23] M. Marcus and L. Véron, The boundary trace of positive solutions of semilinear elliptic equations: the supercritical case, J. Math. Pures Appl. (9) 77 (1998), 481-524.
- [24] B. Mselati, Classiffication and probabilistic representation of the positive solutions of a semilinear elliptic equation, (English summary) Mem. Amer. Math. Soc. 168, no. 798, (2004) 121 pp.