

The Einstein metrics with smooth scri

J. Tafel

Institute of Theoretical Physics, University of Warsaw,
Pasteura 5, 02-093 Warsaw, Poland, email: tafel@fuw.edu.pl

Abstract

We consider solutions of the Einstein equations with cosmological constant $\Lambda \neq 0$ admitting conformal compactification with smooth scri \mathcal{S}^+ . Metrics are written in the Bondi-Sachs coordinates and expanded into inverse powers of the affine distance r . Unlike in the case $\Lambda = 0$ all free data are located on the scri. There are linear differential constraints on the Bondi mass and angular momentum aspects. All other components of metrics are defined in a recursive way.

1 Introduction

This paper is a continuation of [1] where the case $\Lambda = 0$ is considered. We assume that metric \tilde{g} satisfying vacuum equations with $\Lambda \neq 0$ is conformally equivalent to metric g which is smooth in a neighbourhood of the scri \mathcal{S}^+ . We consider its Bondi-Sachs form with the affine distance r . Because of this choice our approach slightly differs from the standard one using the luminosity distance (see [2, 3] and references). If metric is expanded into inverse powers of r almost all Einstein equations can be solved in an algebraic way with respect to metric coefficients. Only the mass aspect M and the angular momentum aspect L_A undergo some linear differential constraints on \mathcal{S}^+ . All free data are defined on the scri. The constraints on M and L_A reduce to a single Laplace-Beltrami equation in special coordinates. If metric tends asymptotically to the (anti) de Sitter solution this equation implies preservation of the generalized Bondi energy and the oscillatory (for $\Lambda < 0$) or exponential (for $\Lambda > 0$) behaviour of the linear momentum.

2 Recursive solvability of the Einstein equations

Let (M, g) be a partial conformal compactification of spacetime (\tilde{M}, \tilde{g}) with a boundary $\mathcal{S}^+ = R \times \Sigma$ foliated by 2-dimensional spacelike surfaces. In the standard way [1] in this neighbourhood we introduce a system of the Bondi-Sachs coordinates $u, \Omega = 1/r, x^A$ such that

$$g = du(g_{00}du - 2d\Omega + 2g_{0A}dx^A) + g_{AB}dx^A dx^B, \quad (1)$$

where

$$\hat{g}_{0A} = 0 \quad (2)$$

(the hat denotes value on \mathcal{S}^+). We assume that components $g_{\mu\nu}$ admit the Taylor expansion in Ω of any order. The physical metric is given by

$$\tilde{g} = \Omega^{-2}g = du(\tilde{g}_{00}du + 2dr + 2\tilde{g}_{0A}dx^A) + \tilde{g}_{AB}dx^A dx^B, \quad (3)$$

where $\tilde{g}_{00}, \tilde{g}_{AB}$ are of the order r^2 and $\tilde{g}_{0A} = O(r)$.

We impose on (3) the Einstein equations with cosmological constant $\Lambda \neq 0$

$$\tilde{R}_{\mu\nu} + \Lambda \tilde{g}_{\mu\nu} = 0 . \quad (4)$$

In terms of the compactified metric they are

$$R_{\mu\nu} - 2Y_{\mu\nu} - g_{\mu\nu} Y_{\alpha}{}^{\alpha} = 0 , \quad (5)$$

where

$$Y_{\mu\nu} = -\frac{1}{\Omega} \Omega_{|\mu\nu} + \frac{1}{2\Omega^2} (\Omega_{|\alpha} \Omega^{|\alpha} - \frac{\Lambda}{3}) g_{\mu\nu} \quad (6)$$

and $_{|\mu}$ denotes the covariant derivative defined by g .

Assume that $\tilde{R}_{\mu\nu} + \Lambda \tilde{g}_{\mu\nu}$ is finite on \mathcal{S}^+ . Then $Y_{\mu\nu}$ must be also finite. In order to avoid a second order pole at $\Omega = 0$ one has to assume that

$$g_{00} = -\frac{\Lambda}{3} + a\Omega + b\Omega^2 - 2M\Omega^3 + O(\Omega^4) . \quad (7)$$

Thus, metric induced by g on the scri reads

$$\hat{g} = -\frac{\Lambda}{3} du^2 + \hat{g}_{AB} dx^A dx^B \quad (8)$$

and \mathcal{S}^+ is spacelike if $\Lambda > 0$ and timelike if $\Lambda < 0$. For the (A)dS metric there is $\hat{g}_{AB} = -s_{AB}$. Still we are free to change coordinates u, x^A provided that \hat{g} preserves form (8) modulo a conformal factor. It means that we can impose one condition on \hat{g}_{AB} , e. g. we can try to obtain

$$\det \hat{g}_{AB} = f , \quad f_{,0} = 0 \quad (9)$$

with a prescribed function f . For instance, if $\mathcal{S}^+ = R \times S_2$, where S_2 is the 2-dimensional sphere, a natural condition would be

$$\det \hat{g}_{AB} = \det s_{AB} . \quad (10)$$

Equation (10) is a reminiscent of the Bondi luminosity gauge

$$\det g_{AB} = \det s_{AB} . \quad (11)$$

Regularity of Y_{1A} yields

$$g_{0A} = q_A \Omega^2 + 2L_A \Omega^3 + O(\Omega^4) . \quad (12)$$

A lack of singularities in Y_{AB} is equivalent to the relation

$$\frac{\Lambda}{3} n_{AB} = \hat{g}_{AB,0} - a \hat{g}_{AB} \quad (13)$$

between expansion coefficients of g_{AB}

$$g_{AB} = \hat{g}_{AB} + n_{AB} \Omega + p_{AB} \Omega^2 + O(\Omega^3) . \quad (14)$$

All other components of $Y_{\mu\nu}$ are nonsingular at $\Omega = 0$ under conditions (7)-(13).

Unlike in the case $\Lambda = 0$, equation (13) defines n_{AB} in terms of \hat{g}_{AB} and a . This relation can be further simplified by means of a shift of r leading to

$$a = 0 . \quad (15)$$

We will not assume (15) at this stage since another condition

$$n = 0 , \quad (16)$$

where $n = \hat{g}^{AB}n_{AB}$, may be more convenient. Note that (16) is equivalent to

$$a = \frac{1}{2}(\ln \det \hat{g}_{AB})_{,0} \quad (17)$$

thanks to (13). Note that under (9) conditions (15) and (16) are equivalent. We will test usefulness of the above conditions after obtaining field equations.

Folowing [1] we can reduce a number of the Einstein equations (4) by means of the Bianchi identity. To this end it is sufficient to replace $\tilde{R}_{\mu\nu}$ by $\tilde{R}_{\mu\nu} + \Lambda\tilde{g}_{\mu\nu}$ and $\tilde{G}_{\mu\nu}$ by $\tilde{G}_{\mu\nu} - \frac{1}{2}\Lambda\tilde{g}_{\mu\nu}$ in equation (24) in [1]. In this way one obtains a minimal system of equations given by

$$\tilde{R}_{11}^{(k)} = \tilde{R}_{1A}^{(k)} = 0 , \quad (\tilde{R}_{AB} + \Lambda\tilde{g}_{AB})^{(k)} = 0 \quad (18)$$

and

$$(\tilde{R}_{00} + \Lambda\tilde{g}_{00})^{(2)} = (\tilde{R}_{0A} + \Lambda\tilde{g}_{0A})^{(2)} = 0 , \quad (19)$$

where the subscript (k) denotes coefficient of order k in an expansion with respect to Ω .

Equation $\tilde{R}_{11} = 0$ and $\tilde{R}_{1A} = 0$ are similar in character to those for $\Lambda = 0$. They yield

$$\hat{g}^{AB}g_{AB}^{(k+2)} = \langle g_{AB}^{(l)}, l \leq k+1 \rangle , \quad k \geq 0 , \quad (20)$$

$$g_{0A}^{(k+2)} = \langle g_{\mu\nu}^{(l)}, l \leq k+1 \rangle , \quad k \geq 0 , \quad k \neq 1 , \quad (21)$$

where brackets on the r. h. s. denote expressions depending on functions within the bracket. For $k = 0$ equation (21) reads

$$q_A = -\frac{1}{2}n^B{}_{A|B} + \frac{1}{2}n_{,A} . \quad (22)$$

Equation $\tilde{R}_{1A}^{(1)} = 0$ does not define $g_{0A}^{(3)}$. It is equivalent to

$$(p^B{}_A + \frac{1}{4}nn^A{}_B - \frac{1}{2}n^{BC}n_{AC})_{|B} + \frac{1}{8}(n_{BC}n^{BC} - n^2)_{,A} = 0 , \quad (23)$$

where operations on indices and covariant derivatives are defined by \hat{g}_{AB} . For $\Lambda = 0$ we used in [1] the topology of $\mathcal{S}^+ = R \times S_2$ to show that (23) yields

$$p_{AB} = \frac{1}{8}(n_{AB}n^{AB} - n^2)\hat{g}_{AB} + \frac{1}{4}nn_{AB} . \quad (24)$$

For $\Lambda \neq 0$ equation (24) follows algebraically from $(\tilde{R}_{AB} + \Lambda\tilde{g}_{AB})^{(0)} = 0$. The latter equation also implies

$$b = -\frac{1}{2}\hat{R} + \frac{1}{2}n_{,0} + \frac{1}{4}an + \frac{\Lambda}{8}(n_{AB}n^{AB} - \frac{1}{3}n^2) , \quad (25)$$

where \hat{R} is the Ricci scalar of \hat{g}_{AB} . From the point of view of equations (24) and (25) the most convenient gauge condition is (16). In this gauge n_{AB} is proportional to the traceless part of the exterior curvature of the section $u = \text{const}$ of \mathcal{I}^+ and p_{AB} is proportional to metric of this section.

Equation $(\tilde{R}_{AB} + \Lambda \tilde{g}_{AB})^{(k)} = 0$ with $k = 1$ does not contain any new information. For $k \geq 2$ it can be splitted into its trace (with respect to \hat{g}_{AB}) and a traceless part. The trace part is equivalent to the following composition of equations

$$\hat{g}^{AB}(\tilde{R}_{AB} + \Lambda \tilde{g}_{AB})^{(k)} - \frac{\Lambda(k-3)}{3(k+1)}\tilde{R}_{11}^{(k)} = 0, \quad k \geq 2. \quad (26)$$

It allows to obtain $g_{00}^{(k+2)}$ in terms of lower order coefficients

$$g_{00}^{(k+2)} = \langle g_{\mu\nu}^{(l)}, l \leq k+1 \rangle, \quad k \geq 2. \quad (27)$$

The traceless part of $(\tilde{R}_{AB} + \Lambda \tilde{g}_{AB})^{(k)} = 0$ yields the traceless part of $g_{AB}^{(k)}$ (denoted by a ring)

$$\mathring{g}_{AB}^{(k+2)} = \langle g_{\mu\nu}^{(l)}, l \leq k+1 \rangle, \quad k \geq 2. \quad (28)$$

Note that in the case $\Lambda = 0$ instead of (28) one obtains an expression for $\mathring{g}_{AB,0}^{(k+1)}$ in terms of lower order coefficients [1]. Thus, (28) is the second equation, after (13), which makes a qualitative difference between $\Lambda = 0$ and $\Lambda \neq 0$. For $\Lambda \neq 0$ there is no need of initial data for g_{AB} . Moreover, tensor

$$N_{AB} = \mathring{g}_{AB}^{(3)} \quad (29)$$

(instead of n_{AB}) is arbitrary if $\Lambda \neq 0$.

The last equations to consider are (19). They are equivalent to equations

$$(\tilde{R}_{00} + \Lambda \tilde{g}_{00})^{(2)} - \frac{\Lambda}{6}\hat{g}^{AB}(\tilde{R}_{AB} + \Lambda \tilde{g}_{AB})^{(2)} - \frac{\Lambda^2}{18}\tilde{R}_{11}^{(2)} - \frac{\Lambda}{9}a\tilde{R}_{11}^{(1)} = 0, \quad (30)$$

$$(\tilde{R}_{0A} + \Lambda \tilde{g}_{0A})^{(2)} - \frac{\Lambda}{3}\tilde{R}_{1A}^{(2)} + \frac{\Lambda}{6}\tilde{R}_{11,A}^{(1)} = 0 \quad (31)$$

having the following structure

$$M_{,0} + \frac{3}{4}(\ln |\hat{g}|)_{,0}M + \frac{\Lambda}{2}L^A{}_{|A} = \frac{\Lambda^2}{24}n^{AB}N_{AB} + \langle a, \hat{g}_{AB} \rangle, \quad (32)$$

$$L_{A,0} + \frac{1}{2}(\ln |\hat{g}|)_{,0}L_A + \frac{1}{3}M_{,A} = \frac{\Lambda}{6}N_{AB}{}^{|B} + \langle a, \hat{g}_{AB} \rangle. \quad (33)$$

Given a , \hat{g}_{AB} and N_{AB} equations (32) and (33) constitute a conjugate system of linear equations for M and L_A . Condition (9) allows to simplify them to

$$M_{,0} + \frac{\Lambda}{2}L^A{}_{|A} = f, \quad (34)$$

$$L_{A,0} + \frac{1}{3}M_{,A} = h_A, \quad (35)$$

where f and h_A are known functions. Let us introduce function M' such that

$$M = M'_{,0} . \quad (36)$$

Then (35) yields

$$L_A = -\frac{1}{3}M'_{,A} + \int h_A du \quad (37)$$

and (34) becomes

$$-\frac{6}{\Lambda}M'_{,00} + \hat{\Delta}M' = -\frac{6}{\Lambda}f + 3\nabla^A \int h_A du , \quad (38)$$

where $\hat{\Delta}$ is the covariant Laplace operator with respect to \hat{g}_{AB} . Thus, equations (32) and (33) reduce to a hyperbolic (if $\Lambda < 0$) or elliptic ($\Lambda > 0$) equation for M' . Note that the operator acting on M' is not the Laplace-Beltrami operator of the induced metric (8) on \mathcal{I}^+ because of the ‘‘wrong’’ coefficient $\frac{6}{\Lambda}$.

Let us summarize consequences of the Einstein equations expanded into powers of $1/r$.

Proposition 2.1. *The Einstein metric with $\Lambda \neq 0$ and smooth scri \mathcal{I}^+ is given by*

$$\tilde{g} = du(\tilde{g}_{00}du + 2dr + 2\tilde{g}_{0A}dx^A) + \tilde{g}_{AB}dx^A dx^B , \quad (39)$$

$$\tilde{g}_{00} = -\frac{\Lambda}{3}r^2 + ar + b - \frac{2M}{r} + \Sigma_2^\infty g_{00}^{(k+2)} r^{-k} , \quad (40)$$

$$\tilde{g}_{0A} = q_A + \frac{2L_A}{r} + \Sigma_2^\infty g_{0A}^{(k+2)} r^{-k} , \quad (41)$$

$$\tilde{g}_{AB} = r^2\hat{g}_{AB} + rn_{AB} + p_{AB} + \frac{1}{r}(N_{AB} + N\hat{g}_{AB}) + \Sigma_2^\infty g_{AB}^{(k+2)} r^{-k} , \quad (42)$$

where coefficients are defined in the following way:

- u -dependent metric \hat{g}_{AB} and a traceless (with respect to \hat{g}_{AB}) tensor N_{AB} can be arbitrarily chosen up to a gauge condition, e.g. (9).
- Function a is arbitrary but it can be gauged away, e.g. by means of (17).
- Components n_{AB} , p_{AB} , N , q_A and b are defined by a and \hat{g}_{AB} according to (13), (24), (20), (22) and (25).
- Coefficients M and L_A are defined by their initial values at $u = u_0$ via equations (32) and (33). Under condition (9) these equations reduce to equation (38) for M' and relations (36) and (37).
- All other components are defined in a recursive way by equations (20), (21), (27) and (28).

The main difference between present situation and that for $\Lambda = 0$ (see Proposition 2.1 in [1]) is that now all free data are located at \mathcal{I}^+ . It is interesting that for $\Lambda < 0$ there is no need of the Cauchy data (\mathcal{I}^+ is timelike). Evolution of the Cauchy data given on \mathcal{I}^+ for $\Lambda > 0$ was first obtained by Friedrich [4].

3 Solutions with the (A)dS boundary metric

If metric \tilde{g} tends asymptotically to the (anti) de Sitter solution then

$$\hat{g}_{AB} = -s_{AB} . \quad (43)$$

In this case it follows from equations (7)-(13) and (20)-(25) that in the gauge $a = 0$ there is

$$\tilde{g}_{00} = -\frac{\Lambda}{3}r^2 + 1 - \frac{2M}{r} + O\left(\frac{1}{r^2}\right) , \quad (44)$$

$$\tilde{g}_{0A} = \frac{2L_A}{r} + O\left(\frac{1}{r^2}\right) , \quad (45)$$

$$\tilde{g}_{AB} = -r^2 s_{AB} + \frac{1}{r} N_{AB} + O\left(\frac{1}{r^2}\right) . \quad (46)$$

The r. h. s. of equations (34) and (35) is given by

$$h_A = \frac{\Lambda}{6} N_{AB}{}^{|B} , \quad f = 0 . \quad (47)$$

Let us introduce a tensor N'_{AB} such that

$$N_{AB} = N'_{AB,0} , \quad s^{AB} N'_{AB} = 0 . \quad (48)$$

Equation (37) yields

$$L_A = -\frac{1}{3} M'_{,A} + \frac{\Lambda}{6} N'_{AB}{}^{|B} + \tilde{L}_A , \quad \tilde{L}_{A,0} = 0 . \quad (49)$$

Using decomposition

$$\tilde{L}_A = f_{,A} + \eta^C_A h_{,C} \quad (50)$$

we can incorporate f into M' and h into N'_{AB} except the dipole component of h (see Lemma 3.1 in [1]). Thus, without loss of generality we can assume that

$$\tilde{L}_A = \eta^C_A h_{,C} , \quad h = \alpha^k Y_{1k} , \quad (51)$$

where Y_{lk} are spherical harmonics and $\alpha^k = \text{const}$. In spherical coordinates θ, φ related to the axis defined by α^k one obtains expression

$$\tilde{L}_A dx^A = \alpha \sin^2 \theta d\varphi , \quad (52)$$

which is the angular momentum term known from the Kerr solution. In view of (49) and (51) equation (34) transforms into

$$\frac{6}{\Lambda} M'_{,00} + \Delta M' = -\frac{\Lambda}{2} N'^{AB}{}_{|AB} , \quad (53)$$

where Δ is the standard Laplace operator on the sphere.

If we integrate equation (53) over the sphere we obtain the known result

$$\oint_{S_2} M d\sigma = \text{const} . \quad (54)$$

The l. h. s. of (54) is, modulo 4π , rather unique candidate for the total energy in this case. One can interpret (54) as a lack of gravitational radiation. If we integrate (53) with Y_{1k} we get equation

$$\frac{6}{\Lambda}M'_{k,00} - 2M'_k = 0 \quad (55)$$

for dipole moments

$$M'_k = \oint_{S_2} M' Y_{1k} d\sigma . \quad (56)$$

The Bondi linear momentum P_k is proportional to the time derivatives of M'_k , so it satisfies equation

$$P_{k,00} = \frac{\Lambda}{3}P_k . \quad (57)$$

Hence, it either oscilates when u changes (for $\Lambda < 0$) or behaves in an exponential way (for $\Lambda > 0$). In the second case its square must exceed the total energy either for increasing or decreasing u , so a reasonable physical assumption would be $M_k = 0$. Higher moments of M' satisfy nonhomogeneous equations coming from (53). Their solutions are defined in quadratures up to oscillatory or exponential functions depending on sign of Λ .

An alternative approach to equation (53) is to treat it as an equation for tensor N'_{AB} . We can represent this tensor by two scalar functions f and h [1]

$$N'_{AB} = f_{|AB} - \frac{1}{2}\Delta f s_{AB} + \eta^C_{(A} h_{|B)C} . \quad (58)$$

Substituting (58) into (53) yields

$$\frac{6}{\Lambda}M'_{,00} + \Delta M' = -\frac{\Lambda}{4}(\Delta + 2)\Delta f . \quad (59)$$

Function f exists if conditions (54) and (56) are satisfied. Then we can write

$$f = -\frac{4}{\Lambda}\Delta^{-1}(\Delta + 2)^{-1}\left(\frac{6}{\Lambda}M'_{,00} + \Delta M'\right) . \quad (60)$$

Formula (60) can be realised by decomposition of M' into Y_{lk} . In this approach functions M' and h are arbitrary modulo conditions (54) and (56). We summarize above results in the following proposition.

Proposition 3.1. *The Einstein metric which has a smooth scri \mathcal{I}^+ and tends asymptotically to the (A)dS metric is given by (39) and (44)-(46). Components M , N_{AB} and L_A are defined by solution (M', N'_{AB}) of equation (53) via relations (36), (48), (49) and (52). Higher order components are defined in a recursive way by equations (20), (21), (27) and (28). The Bondi energy is constant and the Bondi linear momentum satisfies (57).*

4 Stationary metrics

If metric considered in section 2 admits a timelike Killing vector K then we can transform it to the form (39) with u -independent coefficients but we cannot

assume (2) (see Section 3 in [1]) The Killing vector is given by $K = \partial_0$. Let us assume (14) and expansions

$$g_{00} = \hat{g}_{00} + a\Omega + b\Omega^2 - 2M\Omega^3 + O(\Omega^4) , \quad (61)$$

$$g_{0A} = \hat{g}_{0A} + v_A\Omega + q_A\Omega^2 + 2L_A\Omega^3 + O(\Omega^4) \quad (62)$$

with coefficients depending on coordinates x^A . Tensor (6) is finite on the boundary iff

$$\hat{g}_{00} = -\frac{\Lambda}{3} + \hat{g}_{0A}\hat{g}_0^A , \quad (63)$$

$$v_A = n_A^B \hat{g}_{0B} \quad (64)$$

and

$$\frac{\Lambda}{3}n_{AB} = -2\hat{g}_{0(A|B)} + (n_{CD}\hat{g}_0^C\hat{g}_0^D - a)\hat{g}_{AB} , \quad (65)$$

where indices A, B are raised by means of \hat{g}^{AB} . Since $\hat{g}_{0A}\hat{g}_0^A \leq 0$ and we exclude $\Lambda = 0$ it follows from (63)-(65) and timelike property of $K = \partial_0$ that

$$\Lambda < 0 , \quad |\hat{g}_{0A}\hat{g}_0^A| \leq \frac{|\Lambda|}{3} . \quad (66)$$

Thus, metrics are of the AdS type.

It is convenient to shift coordinate r in order to obtain

$$n_A^A = 0 . \quad (67)$$

Equation (65) is then equivalent to the following relations defining n_{AB} and a

$$\frac{\Lambda}{3}n_{AB} = -2\hat{g}_{0(A|B)} + \hat{g}_0^C{}_{|C}\hat{g}_{AB} , \quad (68)$$

$$a = n_{CD}\hat{g}_0^C\hat{g}_0^D - \hat{g}_0^A{}_{|A} . \quad (69)$$

Thus, again n_{AB} is proportional to the traceless part of the exterior curvature of the section $u = \text{const}$ of \mathcal{S}^+ .

Transformation of coordinates x^A allows us to obtain

$$\hat{g}_{AB} = -\gamma^2 s_{AB} , \quad (70)$$

where γ is a positive function of x^A . Still there is a freedom of supertranslation

$$u \rightarrow u + f(x^A) \quad (71)$$

which can be used to impose a condition on \hat{g}_{0A} , γ or higher order coefficients in metric g . All the Einstein equations except (19) can be solved algebraically as in the general case in Section 2. Equations (19) are now a complicated system of constraints on M , L_A , γ , \hat{g}_{0A} and N_{AB} , which is linear in M , L_A and N_{AB} .

The space of stationary solutions is much bigger than in the case $\Lambda = 0$ [1]. In general, solutions do not tend asymptotically to the AdS metric. It is so if the boundary metric

$$\hat{g} = -\frac{\Lambda}{3}du^2 - \gamma^2 s_{AB}(dx^A + \hat{g}_0^A du)(dx^B + \hat{g}_0^B du) \quad (72)$$

is conformally equivalent to

$$\hat{g}' = -\frac{\Lambda}{3}du'^2 - s_{AB}dx'^A dx'^B . \quad (73)$$

Metric (73) is conformally flat (note that it coincides with the conformal compactification of the 3-dimensional Minkowski metric). Thus, metric (72) must be also conformally flat. Since $K = \partial_u$ is the Killing field of (72) it is a conformal Killing vector of the flat metric. Such vectors compose the 10-dimensional algebra $so(2, 3)$. Given one of them in terms of the Cartesian coordinates of flat space one can find (at least in principle) coordinates u, x^A such that $K = \partial_u$. Writing flat metric in these coordinates shows what metrics (72) are available if the physical metric \tilde{g} is asymptotically AdS (in transformed coordinates).

5 Summary

In this paper we considered the Einstein metrics admitting a conformal compactification with smooth scri. For nonvanishing cosmological constant the scri is either spacelike ($\Lambda > 0$) or timelike ($\Lambda < 0$). Metrics can be transformed to the Bondi-Sachs form with components which can be expanded into powers of a radial distance. The Einstein equations imply that the expansion coefficients can be expressed by coefficients of lower order (see Proposition 2.1). Beside these recursive formulas there are 3 differential conditions constraining the analog of the Bondi mass and angular momentum aspect. For a special foliation of the scri they can be reduced to one second order equation for one function. All free data are located on the scri.

Asymptotic form of metric simplifies considerably if it tends to the (anti) de Sitter solution (see Proposition 3.1). In this case the total energy calculated according to the Bondi prescription is constant and the total momentum oscillates or behaves exponentially.

Finally we considered solutions with the timelike Killing field. In this case, in contrary to $\Lambda = 0$, there is no big difference in solving the Einstein equations with respect to the general case. However, existence of stationary solutions with nonvanishing asymptotic twist (then $\hat{g}_{0A} \neq 0$) can be important for a notion of the total energy in general case with $\Lambda < 0$. We expect that a crucial ingredient in any definition of the energy should be the mass aspect M related to coordinates such that equations (8) and (11) are satisfied. If the timelike Killing field K is present then transformation between coordinates used in (8) and those in (72) must depend on time u . It seems unavoidable that the total energy also depend on time, what seems unacceptable in a stationary spacetime.

Acknowledgments

This work was partially supported by Project OPUS 2017/27/B/ST2/02806 of Polish National Science Centre (NCN).

References

- [1] Tafel J 2021 Stationary vacuum metrics with smooth null scri, arXiv:2105.09372

- [2] Barnich G and Troessaert 2010 Symmetries of asymptotically flat 4 dimensional spacetimes at null infinity revisited *Phys. Rev. Lett.* **105** 111103
- [3] Compère G, Fiorucci A and Ruzziconi R 2019 The Λ -BMS4 group of dS4 and new boundary conditions for AdS4 *Class. Quantum Grav.* **36** 195017
- [4] Friedrich H 1986 On the Existence of n-Geodesically Complete or Future Complete Solutions of Einstein's Field Equations with Smooth Asymptotic Structure *Commun. Math. Phys.* **107** 587-609