

Multiscale pentagon integrals to all orders

Dhimiter D. Canko^{1,2}, Costas G. Papadopoulos¹ and Nikolaos Syrrakos^{1,3,4*}

¹ Institute of Nuclear and Particle Physics, NCSR ‘Demokritos’, Agia Paraskevi, 15310, Greece

² Department of Physics, University of Athens,
Zographou 15784, Greece

³ Physics Division, National Technical University of Athens,
Athens 15780, Greece

⁴ Physik-Department, Technische Universität München,
James-Frank-Str. 1, 85748 Garching, Germany
*syrrakos@inp.demokritos.gr

January 24, 2022



*15th International Symposium on Radiative Corrections:
Applications of Quantum Field Theory to Phenomenology,
FSU, Tallahassee, FL, USA, 17-21 May 2021
doi:10.21468/SciPostPhysProc.?*

Abstract

We present analytical results for one-loop five-point master integrals with up to three off-shell legs. The method of canonical differential equations along with the Simplified Differential Equations approach is employed. All necessary boundary terms are given in closed form, resulting to solutions in terms of Goncharov Polylogarithms of arbitrary weight. Explicit results up to weight six will be presented.

Contents

1	Introduction	2
2	Computational framework	3
3	Results	4
3.1	Integral families	4
3.2	Differential equations	5
3.3	Solutions	6
4	Conclusion	7
	References	7

1 Introduction

Our best understanding of Nature in its most fundamental level is encoded in the Standard Model (SM) of particle physics, written in the mathematical language of Quantum Field Theory. Currently, experimental data coming mostly from the realm of Cosmology, such as the existence of dark matter, impose major challenges on our fundamental theories, since they are not predicted by the SM. Thus, we have concrete experimental signs that the SM does not suffice to explain Nature at its most fundamental level.

On the other hand, the SM reigns supreme when we compare its predictions to experimental data coming from the LHC. Since collider physics remains the best way to explore the validity of the SM predictions against experimental data, a new precision program has been initiated in order to test our current understanding of particle physics with the highest possible precision, both from an experimental and a theoretical point of view [1]. Our hope is that through the careful comparison of highly precise measured cross sections against equally precise theoretical predictions, deviations from the SM will be discovered in collider experiments, leading to constraints on its possible extensions [2].

It is estimated that the Run 3 of the LHC as well as its expected High Luminosity upgrade will require from a theoretical standpoint at least Next-to-Next-to-Leading Order (NNLO) corrections to the QCD dominated processes [2]. The current precision frontier at NNLO lies at $2 \rightarrow 3$ processes. A key factor in the determination of theoretical predictions for these processes is the calculation of the relevant two-loop Feynman integrals (FI). Despite the ever increasing sophistication of computational packages such as FIESTA4 [3] and pySecDec [4] dedicated to the numerical evaluation of FI, analytic results for FI are still important, especially for physical regions of the phase space, where in many cases a direct numerical evaluation is not possible.

Working in the framework of dimensional regularisation in $d = 4 - 2\epsilon$ space-time dimensions, it can be shown that FI satisfy so-called Integration-By-Parts (IBP) identities [5]. These relations allow us to obtain a minimal set of FI known as Master integrals (MI), that need to be computed for a specific scattering process. Regarding MI relevant for $2 \rightarrow 3$ NNLO calculations, all MI involving massless particles are known [6–12] and implemented in the C++ library `pentagon functions` [13], all planar MI for processes involving one off-shell leg have recently been calculated using a numerical [14] and an analytical [15] approach and even more recently there has been important progress for some of the non-planar topologies involving one off-shell leg, known as *hexaboxes* [16, 17].

Despite these very important accomplishments, at some point we will have to consider processes involving more external massive legs and massive propagators. This will require the calculation of very complicated two-loop MI. In order to gauge the level of mathematical complexity that these MI will present, it is instructive to study first their one-loop counterparts. To that end, in this contribution we will present analytical results for one-loop five-point MI with up to three off-shell legs and massless internal lines. All results are given in terms of Goncharov Polylogarithms (GPLs), a class of special functions which is well understood by now [18–21], up to transcendental weight six, although the computational approach that was used allows one to trivially obtain higher weight analytical results.

2 Computational framework

The modern approach for computing MI analytically is through the use of the method of differential equations (DE) [22–26]. After using IBP identities and identifying a basis $\mathbf{G}$ of MI, one differentiates this basis with respect to all kinematic invariants

$$\frac{\partial}{\partial s_{ij}} \mathbf{G} = \mathbf{A}(\{s_{ij}\}, \epsilon) \mathbf{G} \quad (1)$$

In general the matrix $\mathbf{A}$ can be very complicated. The introduction of the canonical DE [27] brought forth a revolution in the computation of MI [28]. This new approach suggests that instead of using basis $\mathbf{G}$, one can use a special basis, known as a pure basis of MI, $\mathbf{g} = \mathbf{T}\mathbf{G}$ for which the DE has the following form, known as *canonical form*,

$$d\mathbf{g} = \epsilon \sum_a d \log(W_a) \tilde{\mathbf{M}}_a \mathbf{g} \quad (2)$$

The functions W_a are known as *letters* of the so-called *alphabet*, which is the set of all W_a for a specific family of MI. When W_a are *rational* functions of the differential variables, (2) is solved by recursively integrating order-by-order in ϵ in terms of GPLs, that can be defined as iterated integrals in the following way

$$\mathcal{G}(a_1, a_2, \dots, a_n; x) = \int_0^x \frac{dt}{t - a_1} \mathcal{G}(a_2, \dots, a_n; t) \quad (3)$$

$$\mathcal{G}(0, \dots, 0; x) = \frac{1}{n!} \log^n(x) \quad (4)$$

For five-point MI, achieving an analytical solution in terms of GPLs beyond weight three is a non-trivial task. This is due to the fact that several of the letters W_a are algebraic functions of the differential variables, thus prohibiting a direct integration in terms of GPLs. More specifically, these algebraic letters consist of square roots arising from leading singularities of massive three-point functions and the Gram determinant of the five-point external momenta.

A variant of the standard DE method, known as Simplified Differential Equations (SDE) approach [29], has been shown to effectively circumvent the problem of algebraic letters in many cases [15, 30, 31], thus allowing fully analytical solutions in terms of GPLs to be achieved. In the SDE approach, we introduce an external parameter x in the external momenta and derive the DE by differentiating with respect to only that parameter, regardless of the number of scales of the scattering process. When this approach is applied on a pure basis $\mathbf{g}$ of MI, a canonical SDE can be derived. In many cases the new letters W'_b are fully rationalised in x , $W'_b = x - l_b$, yielding the form

$$\partial_x \mathbf{g} = \epsilon \sum_b \frac{1}{x - l_b} \mathbf{M}_b \mathbf{g} \quad (5)$$

In what follows we will refer to l_b as letters in the SDE approach. All kinematic dependence is included in these l_b functions, leaving the residue matrices $\mathbf{M}_b$ to consist solely of rational numbers. The form of (5) allows for its solution to be trivially expressed in terms of GPLs, assuming the necessary boundary terms are obtained. To do so, we employ the method of expansion-by-regions [32], with which we compute the $x \rightarrow 0$ limit for each pure basis element.

3 Results

3.1 Integral families




Figure 1: Top-sector MI for the pentagon families considered in this contribution.

The integral families considered in this contribution, whose top-sector diagrams are depicted in figure 1, are defined as follows

$$G_{a_1 a_2 a_3 a_4 a_5} = \int \frac{d^d k_1}{i\pi^{(d/2)}} \frac{e^{\epsilon \gamma_E}}{\mathcal{D}_1^{a_1} \mathcal{D}_2^{a_2} \mathcal{D}_3^{a_3} \mathcal{D}_4^{a_4} \mathcal{D}_5^{a_5}}, \quad d = 4 - 2\epsilon \quad (6)$$

with

$$\begin{aligned} \mathcal{D}_1 &= -(k_1)^2, \quad \mathcal{D}_2 = -(k_1 + q_1)^2, \quad \mathcal{D}_3 = -(k_1 + q_1 + q_2)^2 \\ \mathcal{D}_4 &= -(k_1 + q_1 + q_2 + q_3)^2, \quad \mathcal{D}_5 = -(k_1 + q_1 + q_2 + q_3 + q_4)^2 \end{aligned} \quad (7)$$

The kinematics for the three families are as follows,

- 1-mass: $\sum_{i=1}^5 q_i = 0, q_i^2 = 0, i = 1, 2, 4, 5, q_3^2 = m_3^2$
- 2-mass: $\sum_{i=1}^5 q_i = 0, q_i^2 = 0, i = 1, 2, 4, q_3^2 = m_3^2, q_5^2 = m_5^2$
- 3-mass: $\sum_{i=1}^5 q_i = 0, q_i^2 = 0, i = 2, 4, q_1^2 = \bar{m}_1^2, q_3^2 = m_3^2, q_5^2 = m_5^2$

We introduce the following x -parametrization¹

$$q_1 = xp_1, \quad q_2 = xp_2, \quad q_3 = p_{123} - xp_{12}, \quad q_4 = p_4, \quad q_5 = -p_{1234} \quad (8)$$

The kinematics in this underline momentum parametrization is

- 1-mass: $\sum_{i=1}^5 p_i = 0, p_i^2 = 0, i = 1, 2, 3, 4, 5$
- 2-mass: $\sum_{i=1}^5 p_i = 0, p_i^2 = 0, i = 1, 2, 3, 4, p_5^2 = m_5^2$
- 3-mass: $\sum_{i=1}^5 p_i = 0, p_i^2 = 0, i = 2, 3, 4, p_1^2 = m_1^2, p_5^2 = m_5^2$

¹We use the abbreviations $p_{ij} = p_i + p_j$ and $p_{ijk} = p_i + p_j + p_k$ and similarly for q later.

Introducing (8) results in a mapping between the kinematic invariants in the original momentum parametrization, q_i , and the underline momentum parametrization $\{x, p_i\}$ for each of the three pentagon families.²

$$\begin{aligned}
& \text{1-mass : } s_{12} = S_{12}x^2, s_{23} = S_{23}x - S_{45}x + S_{45}, s_{34} = x(S_{12}(x-1) + S_{34}), s_{45} = S_{45}, \\
& \quad s_{15} = S_{15}x, m_3^2 = (x-1)(S_{12}x - S_{45}) \\
& \text{2-mass : } m_3^2 = (x-1)(S_{12}x - S_{45}), s_{12} = S_{12}x^2, s_{23} = S_{23}x - S_{45}x + S_{45} \\
& \quad s_{34} = m_5^2(-x) + m_5^2 + x(S_{12}(x-1) + S_{34}), s_{45} = S_{45}, s_{15} = m_5^2(-x) + m_5^2 + S_{15}x \\
& \text{3-mass : } s_{12} = S_{12}x^2, s_{23} = x(m_1^2(x-1) + S_{23}) - S_{45}x + S_{45}, \\
& \quad s_{34} = m_5^2(-x) + m_5^2 + x(S_{12}(x-1) + S_{34}), s_{45} = S_{45}, \\
& \quad s_{15} = x(m_1^2(x-1) + S_{15}) + m_5^2(-x) + m_5^2, \bar{m}_1^2 = m_1^2x^2, m_3^2 = (x-1)(S_{12}x - S_{45})
\end{aligned} \tag{9}$$

3.2 Differential equations

Constructing pure bases for the pentagon families under consideration is by now a trivial exercise. Following the ideas of [11, 14], the top sector basis element at the *integrand* level is of the form

$$\epsilon^2 \frac{\mathcal{P}_{11111}}{\sqrt{\Delta_5}} \tilde{G}_{11111} \tag{10}$$

where $\mathcal{P}_{11111}$ is the Baikov polynomial corresponding to the top sector *integral* G_{11111} for each family, $\tilde{G}_{11111}$ is the top sector *integrand* of each family and $\Delta_5 = \det[q_i \cdot q_j]$ is the Gram determinant of the five-point external momenta. The remaining pure basis elements can be constructed through the study of the leading singularities of their corresponding diagrams [22]. Using Azurite [33] and Kira2 [34] we can identify 13, 15 and 18 MI for the 1-mass, 2-mass and 3-mass pentagon families respectively.

When considering five-point scattering with massless propagators, a number of square roots of the kinematic invariants enter the DE of the corresponding pure bases. These square roots originate from leading singularities of massive three-point functions, which are represented by square roots of the Källén function $\lambda(x, y, z) = x^2 - 2xy - 2xz + y^2 - 2yz + z^2$ and from square roots of the Gram determinants of the five-point external momenta. The existence of these square roots makes the task of solving these canonical DE in terms of GPLs quite challenging, and in some cases even impossible [35].

It turns out however, that for the families considered in this contribution, the SDE approach can overcome these difficulties. Introducing the x -parametrization (8) rationalises all square roots in terms of the differential variable x , allowing us to integrate the canonical DE and express the final result in terms of GPLs. The canonical SDE for each of the three pentagon families has the following form,

$$\partial_x \mathbf{g} = \epsilon \left(\sum_{i=1}^{l_{\max}} \frac{\mathbf{M}_i}{x - l_i} \right) \mathbf{g} \tag{11}$$

where $\mathbf{g}$ is the pure basis for each family, $\mathbf{M}_i$ are the residue matrices corresponding to each letter l_i and $l_{\max}$ is the length of the alphabet in x . The length of the alphabet for each of the three families considered in this subsection is $l_{\max}^{1m} = 11$, $l_{\max}^{2m} = 14$, $l_{\max}^{3m} = 19$. For more information on the structure of these alphabets, we refer the interested reader to [30, 31].

²We use the abbreviations $s_{ij} = q_{ij}^2$, $S_{ij} = p_{ij}^2$.

The last ingredient that is missing to solve (11) are the boundary terms. To find them we follow the techniques developed in [15, 36]. We define the *resummation matrix* $\mathbf{R}$ as follows

$$\mathbf{R} = \mathbf{S} e^{\epsilon \mathbf{D} \log(x)} \mathbf{S}^{-1} \quad (12)$$

where $\mathbf{S}$ and $\mathbf{D}$ are matrices coming from the Jordan decomposition of the residue matrix corresponding to $l_1 = 0$, $\mathbf{M}_1 = \mathbf{S} \mathbf{D} \mathbf{S}^{-1}$. Having expressed the pure bases in terms of MI, $\mathbf{g} = \mathbf{T} \mathbf{G}$, we use the expansion-by-regions method implemented in the `asy` code which is shipped along with FIESTA4 [3], to find their asymptotic limit for $x \rightarrow 0$.

$$G_i \underset{x \rightarrow 0}{=} \sum_j x^{b_j + a_j \epsilon} G_i^{(b_j + a_j \epsilon)} \quad (13)$$

where a_j and b_j are integers and G_i are the individual members of the basis $\mathbf{G}$. As explained in [15], we can construct the relation

$$\mathbf{R} \mathbf{b} = \lim_{x \rightarrow 0} \mathbf{T} \mathbf{G} \Big|_{\mathcal{O}(x^{0+a_j \epsilon})} \quad (14)$$

where $\mathbf{b} = \sum_{i=0}^n \epsilon^i \mathbf{b}_0^{(i)}$ are the boundary terms that we need to compute. The right-hand-side of (14) implies that, apart from the terms $x^{a_i \epsilon}$ coming from (13), we expand around $x = 0$, keeping only terms of order x^0 . This procedure allows us to fix all the necessary boundary terms in closed form, thus allowing us to obtain analytical solutions of (11) in terms of GPLs of arbitrary weight.

3.3 Solutions

In this contribution we provide solutions up to weight six for each of the considered pentagon families, which can be written in the following compact form,

$$\begin{aligned} \mathbf{g} = & \epsilon^0 \mathbf{b}_0^{(0)} + \epsilon \left(\sum \mathcal{G}_i \mathbf{M}_i \mathbf{b}_0^{(0)} + \mathbf{b}_0^{(1)} \right) + \epsilon^2 \left(\sum \mathcal{G}_{ij} \mathbf{M}_i \mathbf{M}_j \mathbf{b}_0^{(0)} + \sum \mathcal{G}_i \mathbf{M}_i \mathbf{b}_0^{(1)} + \mathbf{b}_0^{(2)} \right) + \dots \\ & + \epsilon^6 \left(\mathbf{b}_0^{(6)} + \sum \mathcal{G}_{ijklmn} \mathbf{M}_i \mathbf{M}_j \mathbf{M}_k \mathbf{M}_l \mathbf{M}_m \mathbf{M}_n \mathbf{b}_0^{(0)} + \sum \mathcal{G}_{ijklm} \mathbf{M}_i \mathbf{M}_j \mathbf{M}_k \mathbf{M}_l \mathbf{M}_m \mathbf{b}_0^{(1)} \right. \\ & \left. + \sum \mathcal{G}_{ijkl} \mathbf{M}_i \mathbf{M}_j \mathbf{M}_k \mathbf{M}_l \mathbf{b}_0^{(2)} + \sum \mathcal{G}_{ijk} \mathbf{M}_i \mathbf{M}_j \mathbf{M}_k \mathbf{b}_0^{(3)} + \sum \mathcal{G}_{ij} \mathbf{M}_i \mathbf{M}_j \mathbf{b}_0^{(4)} + \sum \mathcal{G}_i \mathbf{M}_i \mathbf{b}_0^{(5)} \right) \end{aligned} \quad (15)$$

where $\mathcal{G}_{ab\dots} := \mathcal{G}(l_a, l_b, \dots; x)$ represent the GPLs. The $\mathbf{b}_0^{(i)}$ terms, with i indicating the corresponding weight, consist of Zeta functions $\zeta(i)$, logarithms and GPLs of weight i which have as arguments rational functions of the underline kinematic variables S_{ij} .

For all pentagon families we have made heavy use of the `Mathematica` package `PolyLogTools` [37] for the manipulation of the resulting GPLs. In Tables 1 and 2 we provide an analysis of our results for each family, regarding the number of GPLs that appear in each transcendental weight, where the weight is counted as the number of l_i indices of $\mathcal{G}(l_a, l_b, \dots; x)$. These numbers are obtained by gathering all GPLs that appear up to order $\mathcal{O}(\epsilon^6)$ in each integral family, and distinguishing them according to their corresponding weight. For comparison, we perform the same task for the top-sector basis element of each family.

We observe a huge increase in the number of GPLs beyond weight 5, in comparison to lower weights. Despite the fact that the top sector basis elements appear to have lower-weight GPLs, we note that each top-sector basis element starts from $\mathcal{O}(\epsilon^3)$. We also provide numerical results and timing obtained using `handyG` [38, 39] for the top-sector basis element of each family for a Euclidean point in table 3.

Family	W=1	W=2	W=3	W=4
P_{1m}	10 (2)	50 (21)	170 (99)	496 (339)
P_{2m}	9 (0)	54 (16)	204 (106)	628 (406)
P_{3m}	13 (0)	87 (24)	349 (172)	1115 (696)

Table 1: Number of GPLs entering in the solution. Top-sector b.e. in parenthesis.

Family	W=5	W=6
P_{1m}	1322 (959)	2983 (1924)
P_{2m}	1728 (1254)	4341 (2656)
P_{3m}	3145 (2228)	7849 (4656)

Table 2: Number of GPLs entering in the solution. Top-sector b.e. in parenthesis.

4 Conclusion

Through the calculation of pentagon families with up to three massive legs and massless propagators, we have demonstrated the ability of the SDE approach in handling multiscale FI and obtaining analytical results in terms of GPLs. We have presented solutions of the canonical DE for each pentagon family up to weight six, but the closed form of the boundary terms allows one to trivially obtain higher-weight solutions. We have also presented numerical results and timings for the evaluation of our solutions in a Euclidean point. Obtaining fast numerical results through analytical expressions in physical points is an open problem, due to the challenging task of analytically continuing the resulting GPLs when algebraic letters are present in the alphabet. In this contribution we have focused on integrals with massless propagators. Recently a step forward has been made in the study of pentagon integral families involving one internal mass using the SDE approach [31].

Acknowledgements

NS would like to thank the organisers of RADCOR 2021 for the opportunity to present recent research results.

Funding information This research is co-financed by Greece and the European Union (European Social Fund- ESF) through the Operational Program Human Resources Development, Education and Lifelong Learning 2014 - 2020 in the context of the project "Higher order corrections in QCD with applications to High Energy experiments at LHC" -MIS 5047812. NS work was supported by the Excellence Cluster ORIGINS funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy - EXC-2094 - 390783311.

References

- [1] G. Heinrich, "Collider Physics at the Precision Frontier," Phys. Rept. **922** (2021), 1-69 doi:10.1016/j.physrep.2021.03.006 [arXiv:2009.00516 [hep-ph]].

Top-Sec	Time (sec)	Result
g_{13}	1.90759	$0.0944261\epsilon^3 + 0.31615\epsilon^4 + 0.666923\epsilon^5 + 1.09948\epsilon^6$
g_{15}	3.75112	$-0.120811\epsilon^3 - 0.314547\epsilon^4 - 0.616424\epsilon^5 - 0.985647\epsilon^6$
g_{18}	9.27125	$-0.0215131\epsilon^3 - 0.0332408\epsilon^4 - 0.0501992\epsilon^5 - 0.057848\epsilon^6$

Table 3: Numerical computation of GPLs.

- [2] S. Amoroso, P. Azzurri, J. Bendavid, E. Bothmann, D. Britzger, H. Brooks, A. Buckley, M. Calvetti, X. Chen and M. Chiesa, *et al.* “Les Houches 2019: Physics at TeV Colliders: Standard Model Working Group Report,” [[arXiv:2003.01700 \[hep-ph\]](#)].
- [3] A. V. Smirnov, “FIESTA4: Optimized Feynman integral calculations with GPU support,” *Comput. Phys. Commun.* **204** (2016), 189-199 doi:10.1016/j.cpc.2016.03.013 [[arXiv:1511.03614 \[hep-ph\]](#)].
- [4] S. Borowka, G. Heinrich, S. Jahn, S. P. Jones, M. Kerner, J. Schlenk and T. Zirke, “pySecDec: a toolbox for the numerical evaluation of multi-scale integrals,” *Comput. Phys. Commun.* **222** (2018), 313-326 doi:10.1016/j.cpc.2017.09.015 [[arXiv:1703.09692 \[hep-ph\]](#)].
- [5] K. G. Chetyrkin and F. V. Tkachov, “Integration by Parts: The Algorithm to Calculate beta Functions in 4 Loops,” *Nucl. Phys. B* **192** (1981), 159-204 doi:10.1016/0550-3213(81)90199-1.
- [6] C. G. Papadopoulos, D. Tommasini and C. Wever, “The Pentabox Master Integrals with the Simplified Differential Equations approach,” *JHEP* **04** (2016), 078 doi:10.1007/JHEP04(2016)078 [[arXiv:1511.09404 \[hep-ph\]](#)].
- [7] T. Gehrmann, J. M. Henn and N. A. Lo Presti, “Analytic form of the two-loop planar five-gluon all-plus-helicity amplitude in QCD,” *Phys. Rev. Lett.* **116** (2016) no.6, 062001 [erratum: *Phys. Rev. Lett.* **116** (2016) no.18, 189903] doi:10.1103/PhysRevLett.116.062001 [[arXiv:1511.05409 \[hep-ph\]](#)].
- [8] D. Chicherin, J. Henn and V. Mitev, “Bootstrapping pentagon functions,” *JHEP* **05** (2018), 164 doi:10.1007/JHEP05(2018)164 [[arXiv:1712.09610 \[hep-th\]](#)].
- [9] T. Gehrmann, J. M. Henn and N. A. Lo Presti, “Pentagon functions for massless planar scattering amplitudes,” *JHEP* **10** (2018), 103 doi:10.1007/JHEP10(2018)103 [[arXiv:1807.09812 \[hep-ph\]](#)].
- [10] D. Chicherin, T. Gehrmann, J. M. Henn, N. A. Lo Presti, V. Mitev and P. Wasser, “Analytic result for the nonplanar hexa-box integrals,” *JHEP* **03** (2019), 042 doi:10.1007/JHEP03(2019)042 [[arXiv:1809.06240 \[hep-ph\]](#)].
- [11] S. Abreu, B. Page and M. Zeng, “Differential equations from unitarity cuts: nonplanar hexa-box integrals,” *JHEP* **01** (2019), 006 doi:10.1007/JHEP01(2019)006 [[arXiv:1807.11522 \[hep-th\]](#)].
- [12] D. Chicherin, T. Gehrmann, J. M. Henn, P. Wasser, Y. Zhang and S. Zoia, “All Master Integrals for Three-Jet Production at Next-to-Next-to-Leading Order,” *Phys. Rev. Lett.* **123** (2019) no.4, 041603 doi:10.1103/PhysRevLett.123.041603 [[arXiv:1812.11160 \[hep-ph\]](#)].

- [13] D. Chicherin and V. Sotnikov, “Pentagon Functions for Scattering of Five Massless Particles,” JHEP **12** (2020), 167 doi:10.1007/JHEP12(2020)167 [[arXiv:2009.07803 \[hep-ph\]](#)].
- [14] S. Abreu, H. Ita, F. Moriello, B. Page, W. Tschernow and M. Zeng, “Two-Loop Integrals for Planar Five-Point One-Mass Processes,” JHEP **11** (2020), 117 doi:10.1007/JHEP11(2020)117 [[arXiv:2005.04195 \[hep-ph\]](#)].
- [15] D. D. Canko, C. G. Papadopoulos and N. Syrrakos, “Analytic representation of all planar two-loop five-point Master Integrals with one off-shell leg,” JHEP **01** (2021), 199 doi:10.1007/JHEP01(2021)199 [[arXiv:2009.13917 \[hep-ph\]](#)].
- [16] C. G. Papadopoulos and C. Wever, “Internal Reduction method for computing Feynman Integrals,” JHEP **02** (2020), 112 doi:10.1007/JHEP02(2020)112 [[arXiv:1910.06275 \[hep-ph\]](#)].
- [17] S. Abreu, H. Ita, B. Page and W. Tschernow, “Two-Loop Hexa-Box Integrals for Non-Planar Five-Point One-Mass Processes,” [[arXiv:2107.14180 \[hep-ph\]](#)].
- [18] A. B. Goncharov, “Multiple polylogarithms, cyclotomy and modular complexes,” Math. Res. Lett. **5** (1998), 497-516 doi:10.4310/MRL.1998.v5.n4.a7 [[arXiv:1105.2076 \[math.AG\]](#)].
- [19] C. Duhr, H. Gangl and J. R. Rhodes, “From polygons and symbols to polylogarithmic functions,” JHEP **10** (2012), 075 doi:10.1007/JHEP10(2012)075 [[arXiv:1110.0458 \[math-ph\]](#)].
- [20] C. Duhr, “Hopf algebras, coproducts and symbols: an application to Higgs boson amplitudes,” JHEP **08** (2012), 043 doi:10.1007/JHEP08(2012)043 [[arXiv:1203.0454 \[hep-ph\]](#)].
- [21] C. Duhr, “Mathematical aspects of scattering amplitudes,” doi:10.1142/9789814678766_0010 [[arXiv:1411.7538 \[hep-ph\]](#)].
- [22] J. M. Henn, “Lectures on differential equations for Feynman integrals,” J. Phys. A **48** (2015), 153001 doi:10.1088/1751-8113/48/15/153001 [[arXiv:1412.2296 \[hep-ph\]](#)].
- [23] A. V. Kotikov, “Differential equations method: New technique for massive Feynman diagrams calculation,” Phys. Lett. B **254** (1991), 158-164.
- [24] A. V. Kotikov, “Differential equations method: The Calculation of vertex type Feynman diagrams,” Phys. Lett. B **259** (1991), 314-322.
- [25] A. V. Kotikov, “Differential equation method: The Calculation of N point Feynman diagrams,” Phys. Lett. B **267** (1991), 123-127 [Erratum: Phys. Lett. B **295** (1992), 409].
- [26] T. Gehrmann and E. Remiddi, “Differential equations for two loop four point functions,” Nucl. Phys. B **580** (2000), 485-518 [[arXiv:hep-ph/9912329 \[hep-ph\]](#)].
- [27] J. M. Henn, “Multiloop integrals in dimensional regularization made simple,” Phys. Rev. Lett. **110** (2013), 251601 doi:10.1103/PhysRevLett.110.251601 [[arXiv:1304.1806 \[hep-th\]](#)].
- [28] A. V. Kotikov, “Differential equations and Feynman integrals,” [[arXiv:2102.07424 \[hep-ph\]](#)].
- [29] C. G. Papadopoulos, “Simplified differential equations approach for Master Integrals,” JHEP **07** (2014), 088 doi:10.1007/JHEP07(2014)088 [[arXiv:1401.6057 \[hep-ph\]](#)].

- [30] N. Syrrakos, “Pentagon integrals to arbitrary order in the dimensional regulator,” JHEP **06** (2021), 037 doi:10.1007/JHEP06(2021)037 [[arXiv:2012.10635 \[hep-ph\]](#)].
- [31] N. Syrrakos, “One-loop Feynman integrals for $2 \rightarrow 3$ scattering involving many scales including internal masses,” [[arXiv:2107.02106 \[hep-ph\]](#)].
- [32] B. Jantzen, A. V. Smirnov and V. A. Smirnov, “Expansion by regions: revealing potential and Glauber regions automatically,” Eur. Phys. J. C **72** (2012), 2139 doi:10.1140/epjc/s10052-012-2139-2 [[arXiv:1206.0546 \[hep-ph\]](#)].
- [33] A. Georgoudis, K. J. Larsen and Y. Zhang, “Azurite: An algebraic geometry based package for finding bases of loop integrals,” Comput. Phys. Commun. **221** (2017), 203-215 doi:10.1016/j.cpc.2017.08.013 [[arXiv:1612.04252 \[hep-th\]](#)].
- [34] J. Klappert, F. Lange, P. Maierhöfer and J. Usovitsch, “Integral reduction with Kira 2.0 and finite field methods,” Comput. Phys. Commun. **266** (2021), 108024 doi:10.1016/j.cpc.2021.108024 [[arXiv:2008.06494 \[hep-ph\]](#)].
- [35] F. Brown and C. Duhr, “A double integral of dlog forms which is not polylogarithmic,” [[arXiv:2006.09413 \[hep-th\]](#)].
- [36] D. D. Canko and N. Syrrakos, “Resummation methods for Master Integrals,” JHEP **02** (2021), 080 doi:10.1007/JHEP02(2021)080 [[arXiv:2010.06947 \[hep-ph\]](#)].
- [37] C. Duhr and F. Dulat, “PolyLogTools — polylogs for the masses,” JHEP **08** (2019), 135 doi:10.1007/JHEP08(2019)135 [[arXiv:1904.07279 \[hep-th\]](#)].
- [38] L. Naterop, A. Signer and Y. Ulrich, “handyG —Rapid numerical evaluation of generalised polylogarithms in Fortran,” Comput. Phys. Commun. **253** (2020), 107165 doi:10.1016/j.cpc.2020.107165 [[arXiv:1909.01656 \[hep-ph\]](#)].
- [39] J. Vollinga and S. Weinzierl, “Numerical evaluation of multiple polylogarithms,” Comput. Phys. Commun. **167** (2005), 177 doi:10.1016/j.cpc.2004.12.009 [[arXiv:hep-ph/0410259 \[hep-ph\]](#)].