

Highlights

Control of diffusion-driven pattern formation behind a wave of competency

Yue Liu, Philip K. Maini, Ruth E. Baker

- The effects of a wave of competency on diffusion-driven pattern formation
- The selection of directionality and alignment of stripe patterns by controlling the propagation speed of the wave of competency

Control of diffusion-driven pattern formation behind a wave of competency

Yue Liu^{a,b}, Philip K. Maini^a, Ruth E. Baker^a

^a*Mathematical Institute, University of Oxford, Oxford, UK*

^b*Corresponding author: yue.liu@maths.ox.ac.uk*

Abstract

In certain biological contexts, such as the plumage patterns of birds and stripes on certain species of fishes, pattern formation takes place behind a so-called “wave of competency”. Currently, the effects of a wave of competency on the patterning outcome is not well-understood. In this study, we use Turing’s diffusion-driven instability model to study pattern formation behind a wave of competency, under a range of wave speeds. Numerical simulations show that in one spatial dimension a slower wave speed drives a sequence of peak splittings in the pattern, whereas a higher wave speed leads to peak insertions. In two spatial dimensions, we observe stripes that are either perpendicular or parallel to the moving boundary under slow or fast wave speeds, respectively. We argue that there is a correspondence between the one- and two-dimensional phenomena, and that pattern formation behind a wave of competency can account for the pattern organization observed in many biological systems.

Keywords: Turing pattern, diffusion-driven instability, pattern formation, wave of competency, partial differential equations

2000 MSC: 92C15, 37N25, 35B30

1. Introduction

The diffusion-driven instability (DDI) mechanism proposed by Turing [1] has been used as a canonical model for pattern formation in mathematical biology. Originally, the model was proposed as a mechanism for cell fate specification, but it has since been adopted to describe a wide range of biological phenomena. In this framework, cell fate is hypothesised to be determined by underlying spatially pre-patterned chemical cues. Turing termed these chemicals morphogens, and proposed that different rates of diffusion of the morphogens in the system destabilize a spatially homogeneous steady state, leading to spatially inhomogeneous chemical profiles (patterns). The basic applications of DDI models are on fixed domains, with kinetic parameters constant throughout the entire domain (see e.g. [2, Ch. 2.1]).

There have been many extensions to the basic DDI model proposed in order to obtain a richer, and more robust range of patterning behaviours, reflecting those observed in biology. One such extension is domain growth. The most basic form of growth, where the underlying tissue grows uniformly, results in dilution effects on the morphogens, which can lead to interesting behaviours not found on a fixed domain, as reported by [3, 4, 5, 6, 7], among others. However, in many cases, tissue growth happens on a much slower timescale compared to pattern formation, therefore the effects of domain growth are ignored as they are not biologically relevant.

In contrast, it has been hypothesized that, in certain cases, a related mechanism called a “wave of competency” may serve as a more biologically relevant way to generate interesting patterning behaviours. In this setting, the overall domain size is fixed, but it is divided into two subdomains by a propagating front representing the so-called wave of competency.

Patterning can take place only in the subdomain that lies behind the propagating wave of competency, and this subdomain expands as the wave advances. In the context of patterning via a DDI, we assume that the model parameters are such that the sub-system behind the wave of competency is capable of displaying a DDI, while that ahead of the wave is not. The goal of this paper is to understand the behaviour of the patterning sub-system under different speeds of propagation of the wave of competency.

One possible biological interpretation for the wave of competency is as follows. The overall fixed domain represents the entirety of the underlying tissue. The subdomain in front of the wave represents immature tissue that cannot support patterning until it matures further, i.e. becomes “competent”; and the subdomain behind the wave represents mature tissue, forming the effective domain of patterning. Experimental support for the existence of a wave of competency comes from the plumage patterning of birds, for example Bailleul et al. [8] argued that early mesodermal development leads to the formation of feather fields (the patterning subdomains), which expand on a similar time scale to that of pattern formation.

In an earlier study, Jung et al. [9] identified possible molecular candidates encoding the wave of competency for feather formation, and Jiang et al. [10] provided further experimental evidence. Mou et al. [11] examined a similar mechanism that restricts feather formation to one side of a front. The study [8] found that while a DDI model on a fixed domain could reproduce the final pattern of follicles in birds, it was not able to reproduce the correct sequence of emergence of the feather follicles, nor the consistent orientation of the stripe-shaped transient structures known as feather tracts, which exist prior to the formation of follicles. Hence, they proposed a model which combined chemotaxis and DDI mechanisms by coupling the dynamics of two morphogens with the evolution of cell density. In their model, the cell population

undergoes logistic growth and diffusion, in addition to responding chemotactically to one of the morphogens. The morphogen concentrations are governed by a DDI model, which is coupled to the cell density model via the reaction terms. If the cell density is assumed to be a constant parameter, then the DDI system possesses a stable spatially uniform steady state, which becomes unstable when the cell density becomes sufficiently high. This means that the region of high cell density forms the patterning subdomain, which is initially small, but expands outwards under the effects of chemotaxis and diffusion.

Here, we seek to better understand this type of patterning scenario by isolating the effect of increase in the size of the patterning subdomain on the dynamics of a model capable of exhibiting a DDI. Therefore, we consider a simple reaction-diffusion system, and impose a wave of competency travelling at a constant speed across the domain. This wave forms a moving boundary which divides the domain into two subdomains. Within each subdomain, the system behaves as a classical reaction-diffusion system, but the dynamics may differ across the subdomains, with a DDI only possible for the subdomain behind the moving boundary.

In the classical fixed domain setting, typical patterns arising from a DDI in two spatial dimensions take the form of spots, stripes, or labyrinths. We will focus on understanding the effect of a wave of competency on stripe or labyrinthine patterns, since stripe patterns can exhibit greater variation through their directionality and alignment than typically observed in spot patterns. Further, In the natural world, some stripe patterns exhibit certain preferential directions, and stripes may or may not be aligned throughout the domain. To illustrate stripe directionality and alignment, we use pigmentation patterns on fishes as an example. As shown in Fig. 1, the stripes on a zebrafish are always aligned head-to-tail, along the antero-posterior axis, whereas some other fish species, such as the banded trevally and certain

A photograph of a yellow-striped surgeonfish (Acanthurus lineatus) swimming in a coral reef. The fish has a bright yellow body with seven distinct black vertical stripes. It is shown in profile, facing right, with its long, thin tail fin visible. The background consists of a textured, brownish-grey coral surface.

A photograph of two large muskellunge fish lying on a grassy surface. The fish in the foreground is positioned horizontally, showing its side profile. It has a long, slender body with a pattern of dark, irregular spots and blotches over a lighter, silvery-grey background. The head is large and pointed, with a prominent snout. The second fish is partially visible behind the first, also lying horizontally. The background consists of green grass and some dry, brownish vegetation.

(*Gnathanodon speciosus*)

types of angelfish, have their stripes aligned vertically instead, along the dorso-ventral axis. However, certain catfish species can develop stripe patterns that are not strictly aligned, but nonetheless display a preference for directionality. It is still an open question as to the precise mechanisms driving the alignment of these patterns. Here, we will use Turing's DDI model to explore this phenomenon.

There have been several other proposed mechanisms for pattern alignment. Shoji et al.

[15] showed that anisotropic diffusion can account for directionality, while Hiscock and Megason [16] showed that a gradient in the reaction parameters, or anisotropic growth, can do the same, and Nakamasu et al. [17] were able to produce aligned stripes with a specific initial condition. Page et al. [18] considered parameters that vary across space in more complicated ways, and showed that the DDI model, is, in this case, able to produce a larger variety of patterns. We will compare the patterns arising from implementations of these different mechanisms with the one we are proposing in the discussion.

Although the main motivation behind the study of DDI models comes from biology, it is very challenging to experiment directly with proposed DDI systems in this context due to the complexity of living tissues. Therefore, chemical systems have been used to experimentally validate DDI models, since their dynamics are better understood, and more amenable to manipulation. In Konow et al. [19], the authors experimentally observed and simulated the CDIMA (chlorine dioxide-iodine-malonic acid) reaction, which is capable of displaying DDI, on a patterning subdomain bounded by a circular boundary. This boundary is then made to expand radially outward, serving as an analogue of a wave of competency. In both the chemical experiment and numerical simulations of the corresponding mathematical model, Konow et al found that the stripes produced behind such a wave of competency tend to be concentric circles aligned parallel to the moving boundary if the expansion speed is fast, and radial stripes aligned perpendicular to the boundary if the expansion speed is slow.

In this paper, we seek to determine whether this effect is specific to the setting in [19], or if it is generalizable to a wider range of scenarios. We will mainly consider Schnackenberg kinetics [20], a simple model that allows insight into the observed dynamics, as well as the CDIMA kinetics, which enables us to compare our results to those of [19].

In Section 2, we describe the models to be used, and provide details of how the wave of competency and the expanding subdomain are implemented. In Section 3, we use a series of numerical simulations to explore the behaviour of the model, and to identify and distinguish the cases where different wave speeds lead to qualitatively distinct patterns. Finally, we summarize our findings in Section 4.

2. Models and methods

The basic two-morphogen DDI model in two spatial dimensions, with an imposed wave of competency, can be written as the following system of partial differential equations (PDEs),

$$\frac{\partial u}{\partial t} = D_u \nabla^2 u + f(u, v, W), \quad (x, y) \in \Omega, t > 0, \quad (1a)$$

$$\frac{\partial v}{\partial t} = D_v \nabla^2 v + g(u, v, W), \quad (x, y) \in \Omega, t > 0, \quad (1b)$$

$$W = W(x, y, t), \quad (1c)$$

$$\frac{\partial u}{\partial n} = \frac{\partial v}{\partial n} = 0, \quad (x, y) \in \partial\Omega,$$

$$u(x, y, 0) = u_0(x, y), \quad v(x, y, 0) = v_0(x, y),$$

where $u(x, y, t)$ and $v(x, y, t)$ are the concentrations of the morphogens U and V, respectively, D_u and D_v are their respective diffusion coefficients, and f and g the reaction functions, which differ for the two models we consider. We impose zero-flux boundary conditions, since in most applications the morphogen cannot cross the boundary of the domain. The model is said to display a DDI (also known as Turing instability) if it possesses a spatially homogeneous steady state that is stable in the absence of diffusion, but can be destabilized in the presence of diffusion by a spatially non-homogeneous perturbation [1]. A necessary condition for DDI is that the diffusion coefficients must be different, and, without loss of generality, we take

$$D_u \ll D_v.$$

We model the wave of competency as a boundary separating two subdomains, $\Omega'(t)$, the patterning subdomain within the larger domain Ω , and $\Omega \setminus \Omega'(t)$. We assume that $W(x, y, t)$ is a piecewise constant function such that $W = 0$ on $\Omega'(t)$, and $W = W_{\max}$ on $\Omega \setminus \Omega'(t)$. The values for W_{\max} and the other model parameters are chosen to ensure that the system is capable of undergoing a DDI in $\Omega'(t)$, but not in $\Omega \setminus \Omega'(t)$. The domain and subdomains are illustrated in Fig. 2.




Figure 2: Illustration of the effective patterning domain, Ω' , which is bounded by the wave of competency at $x = \rho(t)$ that propagates towards the right with constant speed α . (PF - pattern forming)

In [19], Ω' was chosen to be a circular region with the radius increasing linearly with time, at constant speed α . Naturally, we expect the system to be attracted to its homogeneous steady state in $\Omega \setminus \Omega'$ and Turing patterns to form inside Ω' , with the patterns expanding outwards in synchrony with the expansion of the region Ω' . Both chemical experiments and numerical simulations carried out by [19] observed this expected behaviour. The interesting

observation is that the type of pattern that forms depends on the wave speed, α , which determines the rate of expansion of Ω' . For the CDIMA chemical reaction system (see later in this section) on a circular expanding subdomain, the authors mainly observed the formation of concentric ring patterns at higher wave speeds, and stripes perpendicular to the circular front at slower wave speeds, with a mixture of the two behaviours for intermediate values of α . The same behaviours are observed in numerical simulations of a DDI model for this reaction. A reproduction of the numerical simulation results can be found in Fig. A.10.

The impact of the wave speed, α , upon the pattern structure is the motivation of our study. We hypothesize that the relation between the rate of domain expansion and the rate of pattern formation is the key to distinguishing between the different behaviours.

In order to remove any potential boundary curvature effects, in this work we will simulate the DDI model described in Eq. (1) on a rectangular domain. We take

$$\Omega = [-L_x, L_x] \times [-L_y, L_y], \quad \Omega'(t) = [-L_x, \rho(t)] \times [-L_y, L_y], \quad \rho(t) = \min(-L_x + \alpha t, L_x), \quad (2)$$

and recall that

$$W(x, y, t) = \begin{cases} 0 & \text{if } (x, y) \in \Omega'(t), \\ W_{\max} & \text{otherwise.} \end{cases}$$

This choice of domain is illustrated in Fig. 2. Mathematically, this W , together with Eq. (1), forms a non-autonomous reaction-diffusion system on a fixed domain.

In this paper we consider two specific sets of reaction kinetics: those for the Schnackenberg model, and those that describe the CDIMA reaction, in order to explore whether the effect of the wave of competency depends on the form of the reaction kinetics, or the phase of the patterns. For the Schnackenberg model, the profiles for u and v are out-of-phase, whereas for the CDIMA model they are in-phase.

Schnackenberg model. The model we will primarily focus on is the Schnackenberg model, which was proposed by Schnackenberg [20] for a hypothetical reaction system, as a special case of the Gierer–Meinhardt model [21]. It consists of very simple kinetics, and it is one of the most well-studied DDI models [22]. The non-dimensional reaction terms can be written as

$$f(u, v) = a - u + u^2v + W, \quad (3a)$$

$$g(u, v) = b - u^2v. \quad (3b)$$

Using linear stability analysis and simulations on two-dimensional fixed domains as a guide, we take as default parameters:

$$D_u = 1, \quad D_v = 20, \quad a = 0.05, \quad b \in \{1.4, 1.6\}, \quad L = 100, \quad W_{\max} = 1. \quad (4)$$

In this model, u and v represent the concentrations of some abstract morphogens, and we take W to represent an increase in the production rate of u . The system possesses the following unique homogeneous steady state when W is treated as a constant,

$$u_* = a + b + W, \quad v_* = b/(a + b + W)^2, \quad (5)$$

which cannot exhibit a DDI when $W = W_{\max} = 1$, but can undergo DDI when $W = 0$.

We consider two values for b . On a fixed domain without a wave of competency, with all other parameters taking the values in Eq. (4), selecting $b = 1.4$ tends to give rise to patterns that consist of a mixture of spots and stripes on a fixed square two-dimensional domain, while selecting $b = 1.6$ mostly gives rise to patterns that consist of stripes (see Fig. 7, right-most column). The conditions which favour spots over stripes or *vice versa* have been investigated

on a fixed two-dimensional domain in [23, 24]. Here we are interested to see if imposing a wave of competency can change this behaviour.

CDIMA model. We also consider the kinetics derived for a model of the CDIMA (chlorine dioxide–iodine–malonic acid) chemical reaction taken from Konow et al. [19], which is a modified version of the model by Lengyel and Epstein [25]. The non-dimensional form of the model is

$$f(u, v) = a - u - \frac{4uv}{1 + u^2} - W, \quad (6a)$$

$$g(u, v) = \sigma b \left(u - \frac{uv}{1 + u^2} + W \right), \quad (6b)$$

with default parameter values taken from [19]:

$$D_u = 1, \quad D_v = \sigma, \quad a = 12, \quad b = 0.31, \quad d = 1, \quad \sigma = 50, \quad L_x = L_y = 100, \quad W_{\max} = 1.5. \quad (7)$$

Physically, u and v represent the concentrations of two of the chemical species in the CDIMA reaction, and W represents the amount of illumination applied onto the reactor. This model assumes that the reactant U turns into V at a constant rate, and U and V are consumed together in a reaction that is inhibited by an abundance of U. When $W < a/5$ is treated as a constant, this system possesses a unique positive homogeneous steady state at

$$u_* = \frac{a - 5W}{5}, \quad v_* = \frac{(u_* + W)(1 + u_*^2)}{u_*}, \quad (8)$$

which cannot exhibit a DDI when $W = 0$, but can undergo DDI when $W = W_{\max} = 1.5$.

This model is able to capture the patterning behaviours of the CDIMA chemical system, qualitatively matching the orientation and alignment of the stripes [19]. Since the illumina-

tion can be readily controlled, as opposed to the physical size of the reactor, manipulating the illumination provides an easy way of observing DDI behind a wave of competency.

Artificial noise. Throughout this work, we will add noise to the models throughout their simulation, at each time step. The reasons for this are as follows.

Firstly, there is intrinsic noise present in almost all biological systems, therefore adding noise enables us to better reflect reality. Secondly, there have long been criticisms that in many situations, the DDI model is very sensitive to noise and changes in the initial conditions, and so the patterns it produces are unreliable [26]. By adding noise to model simulations, we can ensure that our observations and conclusions are robust to noise, and applicable to biologically-realistic situations.

Finally, it is known that for the type of models we are analysing, the numerical scheme chosen for their simulation can have an impact on both the transient behaviour and the final pattern [4], with lower-order methods prone to produce aliasing artifacts [27]. In fact, as we will see in Section 3.2, the choice of mesh can also have an impact on the qualitative behaviour. However, with the addition of sufficient noise, the effects of numerical noise can be eliminated.

In the next section, we will present and compare results first from simulations without, then with, on-going noise. Any noise we add to the initial condition will be specified in each case. For noisy simulations, we add noise to u at each grid point and each time step, independently and identically distributed as $N(0, \mu)$, with $\mu = 0.01$. We found that this magnitude of noise is sufficiently strong to break symmetries and provide new behaviours compared to the noiseless case, and sufficiently weak so that it does not overwhelm the

system, i.e. solutions known to be stable on a fixed domain persist in the presence of this noise. See [Appendix B](#) for details.

3. Numerical simulations

In this section, we carry out numerical simulations of the models outlined in [Section 2](#). We use a standard finite difference scheme for the spatial discretization, where we put equally spaced mesh points at $x_n = -L_x + n\Delta x$, with $\Delta x = 0.5$, and similarly for the y direction in two dimensions. For the time stepping, we use an implicit-explicit scheme (IMEX), where we implement the diffusion term using the implicit Crank–Nicolson method, and use the explicit Euler method for the nonlinear reaction term, with $\Delta t = 0.01$ (see [\[27\]](#) for a discussion of its properties). We have verified that all described behaviour still occurs for a much finer time-stepping with $\Delta t = 0.00125$ or a finer grid size with $\Delta x = 0.25$, and that the numerical scheme indeed converges as Δt is decreased. Model simulations are implemented in Matlab, with the code provided at github.com/liuyue002/turing_expanding_domain.

In [Section 3.1](#), we begin with simulating the models on a one-dimensional domain to identify and understand the basic behaviours of the model and guide our choice for the range of wave speeds, and then in [Section 3.2](#) we move to two-dimensional domains in order to explore the full range of possibilities for pattern behaviours. Finally, in [Section 3.3](#) we use a very narrow two-dimensional domain to closely examine the structure of the stripes in the patterns observed in the earlier simulations. The dynamics on this narrow two-dimensional domain are effectively one-dimensional, which allows us to explain the differences between the distinct patterns observed under varying speeds of propagation of the wave of competency in two dimensions, by relating them to behaviours in one dimension.

3.1. Simulations in one spatial dimension

Although the focus of this study is on two-dimensional pattern structures, we begin in one dimension, where the pattern and the transient behaviours are simpler to visualize and understand. We simulate the PDE system (1) on the one-dimensional domain $\Omega = [-L, L]$. Analogous to the two-dimensional case, the patterning subdomain is defined as $\Omega'(t) = [-L, \rho(t)]$, where $\rho(t)$ is given in Eq. (2). For comparison, we also run simulations in the classical (fixed domain) set up without an expanding patterning subdomain, so that $\Omega' = \Omega$ and $W \equiv 0$.

For the noiseless case, we initialize the system at its homogeneous steady state $u = u_*$, $v = v_*$ (Eq. (5)), with a perturbation to u at the left-hand boundary, created by setting $u(-L, 0) = 2u_*$. For simulations with an expanding subdomain, we found that for the range of α we tested, this initial condition produces the same behaviour as for the simulations when a noisy perturbation is added to the initial condition across the whole domain Ω . This is because at the beginning of the simulation, most of the domain is not capable of supporting a pattern, so the noise far from the left-hand boundary simply decays away and has no further impact. The simulation is run up to 100 time units after the patterning subdomain, Ω' , has expanded to cover the entire domain, Ω , which we found, by visual inspection, is sufficient time for the pattern to evolve to its steady state in both models.

In one spatial dimension, the only possible patterns in both the Schnackenberg and CDIMA models are multi-spike solutions which consist of a series of N equally-spaced, tall, localized peaks, known as spikes (Fig. 3). The stability properties of multi-spike solutions for the Schnackenberg model have been well-studied with asymptotic analysis [22]. Notably, for a given domain length, there is a range of integers N for which the N -spike solutions are

stable.




Figure 3: A typical multi-spike steady state solution of the Schnackenberg model (Eq. (3)) in one dimension. Notice that u and v are exactly out-of-phase, so we will only consider the profile of u henceforth. This simulation is produced with parameters from Eq. (4) with $b = 1.4$ on a fixed domain (i.e. $\Omega' = \Omega$, $W \equiv 0$) without on-going noise. The development of the system over time leading up to this pattern is illustrated in Fig. 4(d).

This means that the final pattern, regardless of the wave speed, α , will be a multi-spike solution. Indeed, we found that in one dimension, with expanding patterning subdomains, the final patterns for different wave speeds α are all multi-spike patterns, which are qualitatively similar to the pattern shown in Fig. 3, except with possibly different numbers of spikes. Despite the fact that the form of pattern does not depend on α , we will show that the transient behaviour changes as α varies. This transient behaviour will be relevant when we relate the one-dimensional behaviours to two-dimensional behaviours in Section 3.3, and it is also interesting from a biological perspective since the transient distribution of the morphogens may have consequences on developmental processes. We will first present the transient behaviours for the Schnackenberg model, shown in Fig. 4, and then note the differences between these and those exhibited by the CDIMA model.




Figure 4: Kymograph representation of u in the solution of the Schnackenberg model (Eq. (3)) in one spatial dimension, with parameters from Eq. (4) and $b = 1.4$, in the absence of on-going noise. Each yellow horizontal stripe corresponds to a spike, the dark red stripes correspond to the valleys between them, and regions of solid colours represent places where the system is locally spatially homogeneous. The system is initialized at the homogeneous steady state with a fixed perturbation at the left-hand boundary, as detailed in the text. Notice that there is a clear divide between the patterning subdomain $\Omega'(t)$ on the lower right (orange with yellow/red stripes), and the non-patterning part of the domain $\Omega \setminus \Omega'(t)$ on the upper left (solid yellow), with the dark blue dashed line representing the wave of competency at $x = \rho(t)$ separating them. The green dashed line in (A,D) is $x = -L + \nu t$, where $\nu \approx 1.2428$ is the natural speed of pattern propagation, see text for details. In all four cases, the final pattern is a multi-spike pattern similar to that in Fig. 3. The animations corresponding to these simulations are provided at imgur.com/a/PRUZ0kn.

First, let us consider the behaviour of the system without a wave of competency, that is, with $\Omega' = \Omega$ (Fig. 4(D)), to provide a comparison to later simulations. Here the spikes naturally form in sequence, one after another, at a steady rate. We define τ to be the first time at which $|u(L, \tau) - u_*| > 0.5$, as a measure of the amount of time it takes for the pattern to cover the entire domain. We find $\tau = 160.93$, and with a domain length of $2L = 200$, we thus estimate the speed of natural pattern propagation as $\nu = 2L/\tau \approx 1.2428$. This steady speed of spike propagation is visualized by the green dashed line in Fig. 4(D).

Next, we implement a wave of competency at advancing at a constant speed α , located

at $\rho(t) = -L + \alpha t$. We will consider a range of values of α relative to the natural pattern propagation speed, ν . First we consider the case where the wave of competency moves more quickly than the natural speed of propagation, $\alpha > \nu$, as in Fig. 4(A). In this case, the subdomain Ω' expands so rapidly that spike formation, which propagates at speed ν , lags behind the wave of competency, leaving a part of Ω' (which has width $(\alpha - \nu)t$) just behind the moving boundary momentarily devoid of pattern. This region corresponds to the small region of solid orange in the plot that sits in between the green and blue dashed lines. In this case, we see that the moving boundary has no meaningful impact upon the formation of the pattern. Consequently, we observe that the rate and manner of spike formation, represented by the time of emergence and the shape of the stripes in Fig. 4(A), is nearly identical to what we observe in Fig. 4(D), except near the right-hand boundary ($x = 100$). Here we see that when the front reaches the boundary at $t = 100$, the higher level of u in the vanishing region $\Omega \setminus \Omega'$ acts as a perturbation to the homogeneous steady state, triggering the formation of a sequence of spikes from right to left.

Next, we consider a moderately high wave speed, $0 \ll \alpha < \nu$, as in Fig. 4(B). In this case, the pattern propagates at the same speed as the wave speed α , since pattern formation is restricted to behind the moving front. Here new spikes form via insertion between the moving front and the previous leading spike, and remain mostly static after their formation. In contrast, with an even slower wave speed, $\alpha \ll \nu$, as in Fig. 4(C), there is a leading spike that travels at the same speed, α , as the moving front, and new spikes form behind it by splitting off the leading spike. After the split, the leading spike continues to move in tandem with the moving boundary. For an intermediate range of wave speeds $0.2 < \alpha < 0.3$, we observe a mixture of behaviours, where both peak insertion and splitting occurs. We

therefore define $\alpha_c \approx 0.25$ to be a threshold that approximately separates the two cases.

We denote these observed behaviours cases A, B, and C, and summarize them in Table 1, including the fixed domain case (case D) for comparison. In all of these cases, the final pattern is the same, and we observe sequential formation of spikes from left to right, either at speed ν or α , whichever is slower. However, there are important differences in the transient behaviours. In Fig. 4(A), we have a part of the domain that momentarily remains at the homogeneous steady state after it has become capable of supporting a pattern, whereas in Fig. 4(B,C) the pattern always fills the entirety of Ω' .

The results for the CDIMA model are largely similar to the observations described above for the Schnackenberg model. We were able to estimate $\nu \approx 1.60$ for the CDIMA model for the parameter values in Eq. (7), which separates case A from cases B and C in a similar way. One key difference is that in both cases B and C, the CDIMA model exhibits peak splitting behaviour similar to case C of the Schnackenberg model. It does not appear to exhibit peak insertion behaviours for any of the tested value of α within the relevant range. This means that we can only observe one, instead of two, distinct behaviours for the range $0 < \alpha < \nu$, in the CDIMA model.

Cases	Range of α	Behaviour in 1D simulations	Behaviour in noisy 2D simulations
A	$\alpha > \nu$	Peaks propagate at speed ν (noiseless, Fig. 4(a)) Disordered peak formation (noisy, Fig. 5(a))	Labyrinthine patterns (Fig. 7, 8, first column)
B	$0 < \alpha_c \ll \alpha < \nu$	Peak insertion (Fig. 4(b), 5(b))	Stripes parallel to moving front (Fig. 7, 8, second column)
C	$0 < \alpha \ll \alpha_c$	Peak splitting (Fig. 4(c), 5(c))	Stripes transverse to moving front (Fig. 7, 8, fourth column)
D	$\alpha = 0$ (fixed domain)	Peaks propagate at speed ν (noiseless, Fig. 4(d)) Disordered peak formation (noisy, Fig. 5(d))	Labyrinthine patterns (Fig. 7, 8, fifth column)

Table 1: Summary of the four cases identified for the Schnackenberg model in Section 3.1, and their behaviours observed in simulations on one-dimensional domains (without on-going noise (Fig. 4), and with on-going noise (Fig. 5)), and two-dimensional domains, with on-going noise (Fig. 7, 8). Two critical thresholds separate the cases: ν , the natural speed of pattern propagation defined in Section 3.1, and α_c , which was estimated using a parameter sweep.

Next, we consider the impact of on-going noise, as discussed in Section 2. In case A, as well as the fixed domain case D, we would expect this noise to trigger the formation of new spikes in the region of Ω' where the system would remain close to the homogeneous steady state in the absence of noise, disrupting the orderly sequential formation of spikes. On the

other hand, in cases B and C, we do not expect this additional noise to have a major effect. This is because the pattern always fills the entirety of Ω' , so the noise decays in Ω' because the multi-spike solution is stable, and it also decays in $\Omega \setminus \Omega'$ due to the spatially uniform steady state being stable in that subdomain.

The numerical results from our simulations confirm our expectations. In Fig. 5, we observe that, indeed, the sequence of spike formation is disrupted in case A. Once again, the behaviour in this case closely resembles the behaviour on a fixed domain (D), where the pattern fills the patterning subdomain Ω' , which coincides with entire domain Ω in the latter case. In contrast, in cases B and C, the behaviour remains virtually unchanged compared to the noiseless case in Fig. 4.




Figure 5: Solution of the Schnackenberg model, with the same parameters, wave speeds and initial conditions as in Fig. 4, except with on-going noise. The blue dashed lines indicate the boundary of $\Omega'(t)$. Corresponding animations are provided at imgur.com/a/0dy3kPY. Observe that in (A) and (D), the spikes no longer form in an orderly sequence, whereas (B,C) are virtually identical to the corresponding cases in Fig. 4.

The same behaviours are observed for the CDIMA model in the noisy case, except for the difference noted before, where peak splitting is observed instead of peak insertion in case B.

From the numerical simulations in one dimension, we are able to draw the following conclusions for the Schnackenberg model. First, if the wave speed, α , which determines the rate

of expansion of the patterning subdomain, is faster than the natural speed of pattern propagation, ν (case A), then there will be little interaction between the moving front and the pattern forming mechanism, in which case we expect similar behaviour to the case where the whole domain can undergo DDI. However, for $\alpha < \nu$ we can have very different behaviours. A high wave speed, α , leads to the formation of new spikes via the insertion of peaks behind the moving front (case B), whereas for a sufficiently low wave speed, α , the spikes form via the splitting of a leading, moving spike (case C). Moving forward into two dimensions, these observations suggest that in the presence of on-going noise, case A will lead to disorganized patterns, matching the behaviour on a fixed domain, whereas cases B and C may lead to the formation of more organized, but possibly distinct, patterns, following the wave of competency. These predictions apply to the CDIMA model as well, except we might expect fewer differences in behaviour between cases B and C.

3.2. Simulations in two spatial dimensions

With the knowledge gained from the four cases observed in one dimension, we now investigate whether we observe the same distinct behaviours in two dimensions, and if the cases in one and two dimensions have any correspondence between them. We run all simulations long enough for Ω' to expand to cover the entire domain Ω , plus a further 200 time units. By numerical experimentation, we found that this is sufficient time for the system to reach a steady state. For comparison, we also run simulations where we set $\Omega' = \Omega$ (that is, the whole domain is capable of undergoing DDI immediately), in which case we run the simulations until $t = 200$. We present only the final profile for u in the two-dimensional simulations, since the profile for v is either in- or out-of-phase with respect to u , depending on the kinetic

terms, so the alignment and orientation of the stripes will be the same for both morphogens. We will first briefly mention the observations in the noiseless case, then investigate how the results change with the addition of on-going noise.

For both the Schnackenberg and CDIMA models, we use the domain described in Section 2, with the parameters from Eq. (4) and (7), and we initialize the system by perturbing the homogeneous steady state with a random perturbation, which is uniformly distributed in $(-0.3, 0.3)$ and added to u at each mesh point across the entire domain. Without on-going noise, we found that for the range of wave speeds $0.1 \leq \alpha \leq 2$, the system always displays vertical stripes parallel to the moving boundary for both models (Fig. 6). The reason for this result is that since initially Ω' , the subdomain capable of supporting Turing patterns, is very small, the noise in the initial condition decays, similar to what we have found in Section 3.1, leaving the system essentially at its homogeneous steady state in $\Omega \setminus \Omega'$. This, combined with the regular square mesh employed in the numerical simulations, means that there is no symmetry breaking mechanism in the y -direction. Consequently, the patterns we obtain are simply the multi-spike pattern in the x -direction, but constant in the y -direction. This reduces the system back to the one-dimensional case. While we do observe differences in the transient behaviour as α changes, analogous to the observations in Section 3.1, the final pattern is always the same.




Figure 6: The final pattern produced with the Schnackenberg model on a two-dimensional domain with parameters from Eq. (4) and $b = 1.4$, $\alpha = 0.75$, initialized by perturbing the homogeneous steady state at each mesh point, without any on-going noise. The resulting pattern is a series of perfectly vertical stripes. Similar results were obtained for other values of α in the range $[0.1, 2]$, and for the CDIMA model.

We also attempted to recover the results of [19] for the CDIMA model, which uses a circular domain that grows radially. The simulations without on-going noise were only able to produce circular rings parallel to the growing boundary, and not the other patterns observed in [19]. Once again, this is an artifact of the regularity of the mesh used; [19] used a finite element method with an irregular mesh, which naturally acts as a symmetry-breaking mechanism. However, simulating the model with on-going noise, we can reproduce all patterns from [19] (see Appendix A).

For the simulations with on-going noise, we first simulate the system of PDEs (1) with both sets of kinetics, initialized at the homogeneous steady state. We add the same perturbation to the initial condition as before for consistency, that is, uniformly distributed noise in $(-0.3, 0.3)$ to each mesh point. The results are shown in Fig. 7, where the rows correspond

to the different kinetics, and the columns are arranged in the order of decreasing wave speed, α . In the first column, which is the case $\alpha > \nu$, corresponding to case A in Table 1, we can observe that the pattern initially takes the form of vertical stripes, but quickly transitions to become disordered and labyrinthine, matching the behaviour on a fixed domain (case D), as shown in the fifth column. This is reminiscent of the one-dimensional behaviour for the same case in Fig. 5(A), where the sequential formation of spikes is disrupted and the pattern becomes disordered, matching the behaviour in Fig. 5(D).

In the second column of Fig. 7, we have $0 \ll \alpha < \nu$, corresponding to case B in Table 1. With a moderately high α , for both the CDIMA and Schnackenberg models, multiple repeated simulations with different random seeds produced only stripes aligned vertically, parallel to the moving boundary of the patterning subdomain representing the wave of competency. This strongly suggests that vertical stripes are the only pattern possible in this case.

For sufficiently small values of α , as in the fourth column, corresponding to case C in Table 1, the stripes may be aligned horizontally, or at a slanted angle (most clearly visible in the Schnackenberg $b = 1.4$ pattern). The CDIMA model tends to form a few, large regions where the stripes are in alignment, where there are two main regions, one with vertical stripes, and the other horizontal stripes. In comparison, the Schnackenberg model tends to exhibit a greater number of such regions. For both models, the final pattern is highly dependent on the noise. The number and location of these regions containing aligned stripes, as well as the direction of stripes within them, can be completely different when a different random seed is used. The third column appears to exhibit a mixture of its two neighbouring columns. The Schnackenberg model produced some slanted stripes, but not as many as in the fourth column.

We also observe in Fig. 7 that the Schnackenberg model with $b = 1.4$ is much more prone to producing spots interspersed between stripes than the same model with $b = 1.6$, or the CDIMA model. However, with an expanding patterning subdomain, the model tends to produce more aligned stripes compared to a fixed domain.




Figure 7: Examples of the steady state patterns produced by the CDIMA model (6), and the Schnackenberg model (3), with choice of domain as given in Eq. (2), and the wave speed decreasing from left to right. The last column shows the fixed domain case. For the CDIMA model, $\alpha = 2, 0.75, 0.5, 0.2$. For the Schnackenberg model, $\alpha = 2, 0.75, 0.5, 0.1$. Notice that in the first column, corresponding to case A in Table 1, the pattern begins as vertically aligned stripes on the left side, but quickly transitions to a labyrinthine pattern as pattern formation lags behind the wave of competency. The second column corresponds to case B, and the patterns mostly take the form of vertical stripes. The third column, corresponding to an intermediate regime between cases B and C, exhibits a mixture of the behaviours from the second and fourth columns, while the fourth column (corresponding to case C) contains regions where the stripes are slanted or horizontal. Finally, the last column shows the labyrinthine patterns produced for the case where $\Omega' = \Omega$. The animations showing the evolution of these patterns are provided at imgur.com/a/u2b7qaV.

These results suggest that the vertical stripe pattern is the only stable pattern in case B (moderately high α), while in case C (low α), we observe multi-stability where stripes aligned at a variety of angles are possible, including horizontal stripes. This leads us to ask

whether the horizontal stripes are stable at low α , and if so, how high does α need to be for the horizontal stripes to become unstable?

To answer this question, we simulate the same models, but using initial conditions chosen as horizontal stripes in the form of cosines, with the amplitude and period set to the estimated values from the final steady state of the previous simulations. Specifically, we set

$$u(x, y, 0) = m_u + A_u \cos(qy), \quad v(x, y, 0) = m_v + A_v \cos(qy), \quad (9)$$

where m_u, A_u, m_v, A_v , and q are estimated to match the patterns in the second column of Fig. 7 by measuring the average wave length and amplitude of the final patterns. For the CDIMA model, we used

$$m_u = 2.41, \quad A_u = 1.16, \quad m_v = 6.67, \quad A_v = 0.71, \quad q = 0.91.$$

For the Schnackenberg model, we used

$$m_u = 1.45, \quad A_u = 0.98, \quad m_v = 0.65, \quad A_v = -0.20, \quad q = 0.57.$$




Figure 8: Examples of the steady state patterns produced by the CDIMA model (6), and the Schnackenberg model (3), where the wave speed decreases from left to right. The parameter values and wave speeds are the same as in Fig. 7, except the initial conditions are set to horizontal stripes, as described in the text. In the first column, for high $\alpha > \nu$, we observe disordered labyrinthine patterns, as before. In the second column, at moderately high α , the pattern abruptly switches to vertical stripes for both models, once again suggesting that this is the only stable pattern at this wave speed. In the third column, a medium value of α leads to a mixture of behaviours. In the CDIMA model, the stripes curve smoothly to become more vertical, while the Schnackenberg model with $b = 1.4$ forms an abundance of spots, which separates regions of aligned stripes. However, for $b = 1.6$, a similar pattern without spots is produced by the Schnackenberg model. In the fourth column, where α is low, we see that horizontal stripes persist for both models, although in the Schnackenberg model the stripes may bend away at a slanted angle. Finally, in the fifth column, unsurprisingly, we observe that the horizontal stripes persist in the fixed domain case. The corresponding animations can be found at imgur.com/a/SiasbMa.

The results are shown in Fig. 8. For $\alpha > \nu$ (first column, corresponding to case A), we

observe the same disordered labyrinthine patterns as we did in Fig. 7, since the initial stripe pattern in $\Omega \setminus \Omega'$ quickly decays to the homogeneous steady state, essentially recreating the same initial conditions as in the earlier simulations. For $0 \ll \alpha < \nu$ (second column, case B), both models produce vertical stripes, and the transition from horizontal stripes to vertical stripes is abrupt. Some spots appear near the transition, but in both the CDIMA model and the Schnackenberg model with $b = 1.6$, the spots are eventually absorbed into new stripes.

For $\alpha \ll \nu$ (fourth column, case C), the horizontal pattern persists for the CDIMA model and for the Schnackenberg model with $b = 1.4$. For the Schnackenberg model with $b = 1.6$, the stripes bend away in some parts of the domain and align at an angle, but nonetheless horizontal stripes dominate. Whether or not this bending occurs is noise-dependent and differs between realisations. For medium α (third column), in both the CDIMA model and the Schnackenberg model with $b = 1.6$, the stripes curved away smoothly from horizontal to become more vertical. For the Schnackenberg model with $b = 1.4$, the curved stripes are replaced with spots.

For the fixed domain case (fifth column, case D), the on-going noise (recall that the noise has magnitude $\mu = 0.01$) was unable to destabilize the initial stripe pattern. This is the only scenario where we obtain different outcomes in cases A and D. However, for the fixed domain case, the pattern can be destroyed if the noise strength is increased sufficiently, see [Appendix B](#) for details. Note that in both Fig. 7 and Fig. 8, some of the stripes are not perfectly straight, but rather are a little curved. This is a consequence of the noise added at each time step.

These simulations show a clear correspondence with the cases in one dimension. First, for case A ($\alpha < \nu$), we obtain labyrinthine patterns similar to case D. This is analogous to

the results observed in one dimension, where similar transient behaviours were observed in cases A and D. For case B ($0 \ll \alpha < \nu$), our simulations suggest that the only stable pattern is vertical stripes parallel to the moving front, whereas for case C ($\alpha \ll \nu$), patterns that are transverse to the moving front, including horizontal and slanted stripes, appear to be stable.

Moreover, for the Schnackenberg model, we found that the switch between cases B and C happens at approximately the same critical value of α in both one and two dimensions, suggesting that the distinction between peak insertion and peak splitting in one dimension may correspond to the distinction between vertical stripes and transversal stripes in two dimensions. For the CDIMA model, despite our observations that it exhibits the same qualitative behaviour in cases B and C in one dimension, we find it exhibits different outcomes in two dimensions that largely agree with the behaviour of the Schnackenberg model, where the preferred directionality of the stripes changes. While this is a surprising contrast to the one-dimensional behaviours, it is consistent with the results in [19], where distinct patterns analogous to those we have observed at different values of α were also observed.

While the critical threshold between cases A and B appears to be at $\alpha = \nu$, the threshold between cases B and C is much less clear. We will explore how these two cases can be distinguished, at least for the Schnackenberg model, in the next section.

3.3. Single stripes on a narrow two-dimensional domain

The simulations so far on one- and two-dimensional domains have allowed us to identify three distinct patterning regimes under the effect of a wave of competency, denoted by cases A, B, and C. While a promising candidate for the critical value of α separating cases A and B has been found, we still need to delineate cases B and C.

By examining animations of the simulations in two dimensions, we observe that during the formation of stripes transversal to the moving front (including horizontal and slanted stripes), there is usually a spot-like structure trailing right behind the front, with the rest of the stripe extending behind it. Since this behaviour is observed for horizontal stripes and slanted stripes, we will classify both of these orientations together as the same type of pattern henceforth. In contrast, a vertical stripe forms spontaneously when the space between existing stripes and the front becomes sufficiently large. This behaviour is common to both models. It turns out that, for the Schnackenberg model, the key to distinguishing cases B and C is to look closely at the structure of the stripe as it lengthens.

To do so, we modify the subdomain capable of undergoing a DDI to be a narrow rectangle that elongates over time. That is, we now consider

$$\Omega' = [-L_x, \rho(t)] \times [-0.1L_y, 0.1L_y], \quad (10)$$

where $\rho(t)$ is as before. We will focus on the Schnackenberg model with $b = 1.4$, then describe how the behaviour of the CDIMA model differs. The narrow width of the patterning subdomain means that no more than one horizontal stripe can be supported. We also found that the stripe does not break up into a row of spots for either model, and the final pattern is a single horizontal stripe for all α in the range $[0.1, 0.75]$. Typical snapshots of the pattern can be found in Fig. 9. We will focus on the transient behaviour during the process of stripe formation, just like we have done in one dimension.

In this case, on-going noise is not necessary to capture the interesting part of the behaviours. In fact, we obtain the same qualitative results regardless of the presence of the on-going noise. This is not surprising since, just as discussed for the one-dimensional case

(Section 3.1), with $\alpha < \nu$ the pattern always fills Ω' in the noiseless simulations. Without any freedom to extend the pattern in the y -direction, the noise has little effect on the pattern. Therefore, we will only present the results from simulations without noise.

We illustrate the transient behaviour with a kymograph of u along the horizontal cross-section at $y = 0$, along the ridge of the single stripe. The results are presented in Fig. 9. We have also included a snapshot of the stripe during its formation for visualization.




Figure 9: Simulations of the Schnackenberg model (Eq. (3)) with $b = 1.4$ on a narrow patterning subdomain as in Eq. (10), represented with kymographs of u along the cross-section $y = 0$, and a snapshot of the two-dimensional stripe during its formation. The vertical blue line indicates where the snapshot was taken. For $\alpha = 0.75$, which corresponds to case B, we can observe the formation of a series of spots via insertion between the moving boundary and the trailing stripe. They persist for only a short time before being absorbed into the stripe. In contrast, for $\alpha = 0.1$, which corresponds to case C, we observe a moving spot travelling at the same speed as the moving boundary, forming a travelling wave solution along the cross-section $y = 0$. The corresponding animations are provided at imgur.com/a/haAoUZg.

For sufficiently slow wave speeds, $\alpha \ll \alpha_c$, for some critical threshold α_c , there is a single leading spot right next to the moving boundary, as in the earlier simulations in two dimensions, which moves in tandem with the front, with the rest of the stripes trailing behind it, remaining a constant distance away from the leading spot. This can be seen in Fig. 9 (right), where the two white diagonal lines, representing the location of the front and the

leading spot, are parallel to each other. The solution along the horizontal cross-section at the middle of the domain, $y = 0$, is effectively a travelling wave, moving at the same speed, α , as the moving boundary.

In contrast, for faster wave speeds, $\alpha \gg \alpha_c$, the leading spot remains stationary, and the trailing stripe expands by engulfing the spot. New leading spots are inserted periodically between the moving boundary and the previous leading spot, as the space between them becomes sufficiently large. We can see this in Fig. 9 (left), where the locations of all local maxima stay roughly fixed. We performed a parameter sweep, simulating the system for a range of values of α , and found that travelling-wave-like behaviour is observed for $\alpha < 0.31$, while spot-insertion behaviour occurs for $\alpha > 0.45$. For the intermediate values of α lying between these values, we observe a mixture of behaviours. Our analysis allows us to make a crude estimate of the threshold $\alpha_c \approx 0.37$ for the Schnackenberg model in this context. Notice that this value for α_c is higher than what we found with simulations on one-dimensional domains. This difference is likely due to the diffusion in the y -direction having a non-trivial impact on the dynamics at the cross-section. This value of α_c serves as a rough guide to delineate cases B and C for the Schnackenberg model. In a sufficiently wide two-dimensional domain (as in Section 3.2), we expect stripes parallel to the moving front to form if $\alpha \gg \alpha_c$, and stripes transverse to the moving front to arise if $\alpha \ll \alpha_c$. Just as we have seen in Section 3.2, the boundary between the two cases is not clear-cut, as we observe mixed behaviours for α near α_c in the third column.

For the CDIMA model, in the narrow two-dimensional domain described in Eq. (10), we observe the same travelling wave-like behaviour in both cases B and C, which is consistent with the observations made in one dimension.

4. Discussions

The aim of this work was to explore the impact of a wave of competency upon DDI models of pattern formation, taking as exemplar kinetics the Schnackenberg and CDIMA models. Through extensive numerical exploration, we have shown that the speed of the wave of competency can be used to select the resulting pattern. Specifically, the alignment and directionality of striped patterns in two dimensions can be selected to be either parallel or perpendicular to the wave of competency by appropriate choice of the wave speed. Moreover, this phenomenon holds true for both the CDIMA and Schnackenberg kinetics, with minor differences in the model behaviours in one dimension. The agreement between the two models suggests that our results for two-dimensional behaviours may possibly be generalized to a larger class of models capable of exhibiting a DDI, i.e. that the phenomenon does not depend on the specific form of the reaction terms, nor on whether the morphogens are in- or out-of-phase. Our observations are robust to the addition of noise, which gives us confidence that the results on stripe alignment and directionality are robust.

We have observed three main patterning regimes on expanding patterning subdomains with respect to the value of the wave speed α . These cases (A,B,C), along with the behaviour on the fixed domain (case D) for comparison, are summarized in Table 1. We have defined the natural speed of pattern propagation, ν , as the natural speed at which a pattern spreads from a spatially localised perturbation on a fixed domain. Our first main conclusion, which holds for both models, is that in order for stripe patterns in two dimensions to align in the presence of noise, the wave speed must be slower than the speed at which the pattern propagates naturally, that is $\alpha < \nu$, otherwise we obtain labyrinthine patterns that are

essentially the same as those observed on fixed domains.

Our second main conclusion, again holding for both models, is that given $\alpha < \nu$, a higher α leads to stripes that are parallel to the moving front. These stripes arise as a two-dimensional analogue of peak insertion between the moving front and the existing pattern. In contrast, a lower α permits the formation of stripes that are transverse to the moving boundary, either perpendicular or at a slanted angle. Along the ridges of the stripes, the extension of the stripes as the front moves forward can be seen as a two-dimensional analogue of a travelling wave. For the Schnackenberg model, we were able to obtain an estimate for α_c , the threshold for α that separates these two cases, by observing the structure of the stripe that evolves when the region competent to form a pattern is restricted in the y -direction, so that it admits complicated patterns only in the x -direction. However, the separation between the two patterning regimes is not clear-cut, as we observe mixed behaviours for $\alpha \approx \alpha_c$. Moreover, despite its agreement in behaviour with the Schnackenberg model on broad two-dimensional domains, the CDIMA model does not display the peak splitting behaviour observed in the Schnackenberg model in one-dimensional and narrow two-dimensional domains.

In future, it will be interesting to understand why the two models behave differently on one-dimensional domains, but nonetheless exhibit similar behaviours on non-narrow two-dimensional domains. Another research direction, beyond the scope of this study, is to analytically determine the value of the critical threshold α_c . We speculate that this could be done by approximating the dynamics of the two-dimensional system along a stripe as a modified one-dimensional system, and using a wavelet analysis to determine the existence of a travelling wave solution.

We now compare the pattern selection mechanism investigated in this paper with other

mechanisms proposed in earlier works. In our model, we have chosen a wave of competency that advances in the x -direction only. However, the conclusions regarding alignment of the stripes with respect to the wave remain the same if the pattern subdomain expands isotropically (such as an expanding circle, see [Appendix A](#)), meaning that this mechanism does not rely on any anisotropy. In contrast, for the anisotropic diffusion mechanism considered by Shoji et al. [15], where the two morphogens have potentially differing preferred directions of diffusion, the anisotropy is essential for alignment. Hiscock and Megason [16] considered two mechanisms for pattern alignment, which were the existence of spatial gradients in the kinetic parameters, and anisotropies in diffusion or growth. These mechanisms were able to select between disordered labyrinthine patterns and one particular alignment for the stripes, but not between multiple possible directions, as our mechanism is capable of doing. Therefore, the mechanism we have proposed behaves differently compared to the previously proposed mechanisms for controlling the alignment and directionality of stripes.

Further generalizations of this model are possible. In this paper, the wave of competency has a simple shape, and its movement is independent of the morphogen concentrations. In a real biological system, the wave speed and the shape of the wave front may be coupled with the morphogen concentrations. For example, in [28], the patterning domain is interpreted as the interior of a cell, and the morphogens, through interactions with the actin cytoskeleton, deform the cell. It would be interesting to compare behaviours in this kind of dynamic domain under zero-flux boundary conditions, with behaviours in a domain bounded by a moving wave of competency.

In summary, we have demonstrated that modulation of the speed of propagation of a wave of competency can robustly select patterns in DDI models. These results have implications

for patterning in a number of biological systems where a wave of competency has been either identified, or could exist. Our results also pave the way for future studies that aim to control the directionality and alignment of DDI patterns by modulating the wave speed. More complex patterns may be obtained by also varying the shape of the wave front.

Acknowledgement

YL is supported by the Natural Sciences and Engineering Research Council of Canada (NSERC) through the Postgraduate Scholarships – Doctoral program, reference number PGSD3-535584-2019. REM acknowledges the Royal Society for a Wolfson Research Merit Award.

Appendix A. Simulations on two-dimensional circular patterning subdomain

In this appendix we simulate the CDIMA (Eq. (6)) and Schnackenberg models (Eq. (3)) on a circular patterning subdomain that expands radially at constant speeds, that is,

$$\Omega'(t) = \{(x, y) | x^2 + y^2 \leq \rho(t)^2\}, \quad \rho(t) = \min(\alpha t, L_x/2, L_y/2), \quad (\text{A.1})$$

instead of the rectangular patterning subdomain used in the main text. This choice matches the experimental and simulation set up of Konow et al. [19], which enable us to compare our results.

The steady state patterns are shown in Fig. A.10. For simulations without on-going noise, the resulting patterns are concentric rings regardless of the wave speed α . With on-going noise, for the CDIMA model, we observe mostly concentric rings when the wave speed α is sufficiently fast, and mostly stripes transversal to the circular boundary when the wave speed is low. This is consistent with the observations from [19].

The Schnackenberg model with $b = 1.6$ exhibit mostly the same qualitative behaviour as the CDIMA model. An interesting difference is that, with a low wave speed α , despite producing a pattern similar to the CDIMA model at first, the stripes near the circular boundary eventually evolves to rings once the wave of competency stops, while the stripes in the center of the domain remain in a radial direction. With $b = 1.4$, the Schnackenberg model tends to produce more spots. These are consistent with our observations in [Section 3](#) on a rectangular patterning subdomain, where the vertical stripes parallel to the linear wave front plays the same role as the rings in the circular setting.




Figure A.10: The steady state pattern produced with a circular patterning subdomain, as described in Eq. (A.1). The system is initialized by perturbing the homogeneous steady state with noise. In the first column, without on-going noise, the final pattern takes the form of concentric rings regardless of the value of α (the plot shown is produced with $\alpha = 0.75$). In the second and third columns, with the on-going noise we observe different qualitative behaviours for different values of α .

Appendix B. Simulations with higher magnitude of noise

In this appendix we show the effect of higher level of on-going noise on the pattern. We simulate the Schnackenberg model on a fixed domain (that is, $\Omega' = \Omega$), with $b = 1.4$, and other parameter values from Eq. (4). We initialize the system at horizontal stripes as described in Eq. 9.

When the noise is sufficiently strong, the pattern constantly evolves and never settles to a true steady state, so in Fig. B.11 we chose to present snapshots of the system at $t = 350$, which is on the same timescale as the simulations in the main text. Presumably, if the simulations are run for much longer, then even a weak noise will eventually disrupt the pattern.

We observe that with noise strength $\mu = 0.01$, the initial horizontal pattern persists mostly undisturbed, only becoming slightly bent in certain places. With a stronger noise $\mu = 0.03$, the horizontal stripes are still recognizable but significantly disrupted, and with $\mu = 0.05$ the stripes are entirely unrecognizable, and the pattern evolves to a mess of spots and stripes that constantly change and shift as time continue to progress. We chose $\mu = 0.01$ for the simulations in the main text so that the noise does not overwhelm the system.




Figure B.11: Snapshots of the simulations for the Schnackenberg model on a fixed domain, initialized at horizontal stripes, with $b = 1.4$ and varying noise strength μ , taken at $t = 350$. From left to right: $\mu = 0, 0.01, 0.03, 0.05$.

References

- [1] A. M. Turing, The chemical basis of morphogenesis, Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences 237 (1952) 37–72. doi:[10.1098/rstb.1952.0012](https://doi.org/10.1098/rstb.1952.0012).

- [2] J. D. Murray, *Mathematical Biology II: Spatial Models and Biomedical Applications*, volume 3, Springer New York, 2001.
- [3] I. Barrass, E. J. Crampin, P. K. Maini, Mode transitions in a model reaction-diffusion system driven by domain growth and noise, *Bulletin of Mathematical Biology* 68 (2006) 981–995. doi:[10.1007/s11538-006-9106-8](https://doi.org/10.1007/s11538-006-9106-8).
- [4] A. Madzvamuse, P. K. Maini, Velocity-induced numerical solutions of reaction-diffusion systems on continuously growing domains, *Journal of Computational Physics* 225 (2007) 100–119. doi:[10.1016/j.jcp.2006.11.022](https://doi.org/10.1016/j.jcp.2006.11.022).
- [5] E. J. Crampin, E. A. Gaffney, P. K. Maini, Reaction and diffusion on growing domains: scenarios for robust pattern formation, *Bulletin of Mathematical Biology* 61 (1999) 1093–1120. doi:[10.1006/bulm.1999.0131](https://doi.org/10.1006/bulm.1999.0131).
- [6] E. J. Crampin, W. W. Hackborn, P. K. Maini, Pattern formation in reaction-diffusion models with nonuniform domain growth, *Bulletin of Mathematical Biology* 64 (2002) 747–769. doi:[10.1006/bulm.2002.0295](https://doi.org/10.1006/bulm.2002.0295).
- [7] R. A. Van Gorder, V. Klika, A. L. Krause, Turing conditions for pattern forming systems on evolving manifolds, *Journal of Mathematical Biology* 82 (2021) 1–61. doi:[10.1007/s00285-021-01552-y](https://doi.org/10.1007/s00285-021-01552-y).
- [8] R. Bailleul, C. Curantz, C. D.-T. Dinh, M. Hidalgo, J. Touboul, M. Manceau, Symmetry breaking in the embryonic skin triggers directional and sequential plumage patterning, *PLOS Biology* 17 (2019) e3000448. doi:[10.1371/journal.pbio.3000448](https://doi.org/10.1371/journal.pbio.3000448).

- [9] H.-S. Jung, P. H. Francis-West, R. B. Widelitz, T.-X. Jiang, S. Ting-Berreth, C. Tickle, L. Wolpert, C.-M. Chuong, Local inhibitory action of BMPs and their relationships with activators in feather formation: implications for periodic patterning, *Developmental Biology* 196 (1998) 11–23. doi:[10.1006/dbio.1998.8850](https://doi.org/10.1006/dbio.1998.8850).
- [10] T. Jiang, H. Jung, R. Widelitz, C. Chuong, Self-organization of periodic patterns by dissociated feather mesenchymal cells and the regulation of size, number and spacing of primordia, *Development* 126 (1999) 4997–5009. doi:[10.1242/dev.126.22.4997](https://doi.org/10.1242/dev.126.22.4997).
- [11] C. Mou, F. Pitel, D. Gourichon, F. Vignoles, A. Tzika, P. Tato, L. Yu, D. W. Burt, B. Bed’hom, M. Tixier-Boichard, K. J. Painter, D. J. Headon, Cryptic patterning of avian skin confers a developmental facility for loss of neck feathering, *PLOS Biology* 9 (2011) e1001028. doi:[10.1371/journal.pbio.1001028](https://doi.org/10.1371/journal.pbio.1001028).
- [12] Azul, Zebrafish, 2005. URL: <https://commons.wikimedia.org/wiki/File:Zebrafisch.jpg>, image under CC-0 license.
- [13] R. Zerpe, Golden trevally (gnathanodon speciosus), 2019. URL: <https://www.flickr.com/photos/krokodiver/47728328672/>, image under CC-BY license.
- [14] R. A. Barrio, R. E. Baker, B. Vaughan, K. Tribuzy, M. R. de Carvalho, R. Bassanezi, P. K. Maini, Modeling the skin pattern of fishes, *Physical Review E* 79 (2009) 031908. doi:[10.1103/PhysRevE.79.031908](https://doi.org/10.1103/PhysRevE.79.031908).
- [15] H. Shoji, Y. Iwasa, A. Mochizuki, S. Kondo, Directionality of stripes formed by anisotropic reaction-diffusion models, *Journal of Theoretical Biology* 214 (2002) 549–561. doi:[10.1006/jtbi.2001.2480](https://doi.org/10.1006/jtbi.2001.2480).

- [16] T. W. Hiscock, S. G. Megason, Orientation of Turing-like patterns by morphogen gradients and tissue anisotropies, *Cell Systems* 1 (2015) 408–416. doi:[10.1016/j.cels.2015.12.001](https://doi.org/10.1016/j.cels.2015.12.001).
- [17] A. Nakamasu, G. Takahashi, A. Kanbe, S. Kondo, Interactions between zebrafish pigment cells responsible for the generation of Turing patterns, *Proceedings of the National Academy of Sciences* 106 (2009) 8429–8434. doi:[10.1073/pnas.0808622106](https://doi.org/10.1073/pnas.0808622106).
- [18] K. M. Page, P. K. Maini, N. A. M. Monk, Complex pattern formation in reaction-diffusion systems with spatially varying parameters, *Physica D: Nonlinear Phenomena* 202 (2005) 95–115. doi:[10.1016/j.physd.2005.01.022](https://doi.org/10.1016/j.physd.2005.01.022).
- [19] C. Konow, N. H. Somberg, J. Chavez, I. R. Epstein, M. Dolnik, Turing patterns on radially growing domains: Experiments and simulations, *Physical Chemistry Chemical Physics* 21 (2019) 6718–6724. doi:[10.1039/C8CP07797E](https://doi.org/10.1039/C8CP07797E).
- [20] J. Schnakenberg, Simple chemical reaction systems with limit cycle behaviour, *Journal of Theoretical Biology* 81 (1979) 389–400. doi:[10.1016/0022-5193\(79\)90042-0](https://doi.org/10.1016/0022-5193(79)90042-0).
- [21] A. Gierer, H. Meinhardt, A theory of biological pattern formation, *Kybernetik* 12 (1972) 30–39. doi:[10.1007/BF00289234](https://doi.org/10.1007/BF00289234).
- [22] D. Iron, J. Wei, M. Winter, Stability analysis of Turing patterns generated by the Schnakenberg model, *Journal of Mathematical Biology* 49 (2004) 358–390. doi:[10.1007/s00285-003-0258-y](https://doi.org/10.1007/s00285-003-0258-y).
- [23] B. Ermentrout, J. E. Enderby, Stripes or spots? Nonlinear effects in bifurcation of reaction-diffusion equations on the square, *Proceedings of the Royal Society of London*.

- Series A: Mathematical and Physical Sciences 434 (1991) 413–417. doi:[10.1098/rspa.1991.0100](https://doi.org/10.1098/rspa.1991.0100).
- [24] B. N. Nagorcka, J. R. Mooney, From stripes to spots: Prepatterns which can be produced in the skin by a reaction-diffusion system, *Mathematical Medicine and Biology: A Journal of the IMA* 9 (1992) 249–267. doi:[10.1093/imammb/9.4.249](https://doi.org/10.1093/imammb/9.4.249).
- [25] I. Lengyel, I. R. Epstein, Modeling of Turing structures in the Chlorite–Iodide–Malonic Acid–Starch reaction system, *Science* 251 (1991) 650–652. doi:[10.1126/science.251.4994.650](https://doi.org/10.1126/science.251.4994.650).
- [26] J. Bard, I. Lauder, How well does Turing’s theory of morphogenesis work?, *Journal of Theoretical Biology* 45 (1974) 501–531. doi:[10.1016/0022-5193\(74\)90128-3](https://doi.org/10.1016/0022-5193(74)90128-3).
- [27] S. J. Ruuth, Implicit-explicit methods for reaction-diffusion problems in pattern formation, *Journal of Mathematical Biology* 34 (1995) 148–176. doi:[10.1007/BF00178771](https://doi.org/10.1007/BF00178771).
- [28] Y. Liu, E. G. Rens, L. Edelstein-Keshet, Spots, stripes, and spiral waves in models for static and motile cells, *Journal of Mathematical Biology* 82 (2021) 28. doi:[10.1007/s00285-021-01550-0](https://doi.org/10.1007/s00285-021-01550-0).