

Detecting long-range dependence for time-varying linear models

Lujia Bai

Center for Statistical Science,
Department of Industrial Engineering,
Tsinghua University

Weichi Wu

Center for Statistical Science,
Department of Industrial Engineering,
Tsinghua University

Abstract

We consider the problem of testing for long-range dependence in time-varying coefficient regression models, where the covariates and errors are locally stationary, allowing complex temporal dynamics and heteroscedasticity. We develop KPSS, R/S, V/S, and K/S-type statistics based on the nonparametric residuals. Under the null hypothesis, the local alternatives as well as the fixed alternatives, we derive the limiting distributions of the test statistics. As the four types of test statistics could degenerate when the time-varying mean, variance, long-run variance of errors, covariates, and the intercept lie in certain hyperplanes, we show the bootstrap-assisted tests are consistent under both degenerate and non-degenerate scenarios. In particular, in the presence of covariates the exact local asymptotic power of the bootstrap-assisted tests can enjoy the same order as that of the classical KPSS test of long memory for strictly stationary series. The asymptotic theory is built on a new Gaussian approximation technique for locally stationary long-memory processes with short-memory covariates, which is of independent interest. The effectiveness of our tests is demonstrated by extensive simulation studies and real data analysis.

Keywords: Long-range dependence, Locally stationary process, Spurious long memory, Time-varying models

1 Introduction

Consider the time-varying coefficient linear model

$$y_{i,n} = \mathbf{x}_{i,n}^\top \boldsymbol{\beta}(t_i) + e_{i,n}, \quad i = 1, 2, \dots, n, \quad (1.1)$$

where the covariate vector $\mathbf{x}_{i,n} = (1, x_{i,2,n}, \dots, x_{i,p,n})^\top$ is a p -dimensional *short-range dependent* (SRD) locally stationary time series and $y_{i,n}$ is the response variable, $t_i = i/n$. At each time point t_i , we only observe one realization $(\mathbf{x}_{i,n}, y_{i,n})$ and no repeated measurement is available. The time-varying regression coefficient function $\boldsymbol{\beta}(\cdot)$ is a p -dimensional function with each coordinate a smooth function on $[0, 1]$ and the zero mean error process $(e_{i,n})$ is a possibly *long-range dependent* (LRD) or *long-memory* time series. More precisely, we assume $(e_{i,n})$ is a locally stationary $I(d)$ process i.e., for $1 \leq i \leq n$

$$(1 - \mathcal{B})^d e_{i,n} = u_{i,n}, \quad (1.2)$$

where \mathcal{B} is the lag operator, $d \in [0, 1/2)$ is the *long-memory parameter* and $u_{i,n}$ is a SRD or *short-memory* locally stationary process. The strict definitions of locally stationary and long-memory processes are

¹E-mail addresses: blj20@mails.tsinghua.edu.cn(L.Bai), wuweichi@mail.tsinghua.edu.cn(W.Wu)

deferred to [Section 3](#). The error model (1.2) naturally generalizes classical stationary SRD and LRD processes by allowing their generating mechanism to vary with time. Observe that $(e_{i,n})$ will reduce to the SRD process $(u_{i,n})$ if $d = 0$ and will be a LRD process if $d \in (0, 1/2)$. In fact, when $(u_{i,n})$ is stationary, (1.2) allows the classical stationary long-memory processes (e.g. FARIMA-GARCH models), which have found extensive application in hydrology ([Zhang et al. \(2011\)](#), [Koutsoyiannis \(2013\)](#)), economics and finance ([Caporale and Gil-Alana \(2013\)](#), [Caporale et al. \(2016\)](#)) and many other fields since first introduced by [Hurst \(1951\)](#). Moreover, model (1.1) admits heteroscedasticity, i.e., the dependence of $u_{i,n}$ on $(\mathbf{x}_{r,n})_{r=1}^n$, see [Section 3](#) for more details.

The time-varying regression model (1.1) with time series errors has attracted enormous attention, see for instance [Fan and Zhang \(2000\)](#), [Zhou and Wu \(2010\)](#) and [Chen et al. \(2018\)](#) where the errors are assumed to be SRD, and [Kulik and Wichelhaus \(2012\)](#), [Beran et al. \(2013\)](#) and [Ferreira et al. \(2018\)](#) where LRD errors are considered. The aforementioned research reveals that nonparametric estimators of the time-varying coefficient $\beta(\cdot)$ possess distinct properties under the two scenarios, $d = 0$ and $0 < d < 1/2$. When $d = 0$, consider the local linear estimator of the multivariate coefficient function $\beta(\cdot)$ using the kernel function $K(\cdot)$ and the bandwidth b_n , of which the asymptotic behavior rests on the distributions of $(\mathbf{x}_{i,n})$ and $(e_{i,n})$. In particular, the order of the deviation $|\hat{\beta}(\cdot) - \beta(\cdot)|$ is determined by the long-memory parameter d . For $d = 0$ and $t \in (0, 1)$, [Zhou and Wu \(2010\)](#) shows that under mild conditions,

$$\sqrt{nb_n}(\hat{\beta}(t) - \beta(t) - b_n^2\beta''(t)\mu_2/2) \Rightarrow N(0, \phi_0\Sigma(t)), \quad (1.3)$$

where μ_2 and ϕ_0 are constants determined by $K(\cdot)$ and $\Sigma(t)$ is determined by the moments of the process $(\mathbf{x}_{i,n}e_{i,n})_{i=1}^n$. Meanwhile, for $d > 0$ and $p = 1$, Theorem 7.22 in [Beran et al. \(2013\)](#) shows that for stationary $e_{i,n}$ under regularity conditions,

$$(nb_n)^{1/2-d}(\hat{\beta}(t) - \beta(t) - b_n^2\beta''(t)\mu_2/2) \Rightarrow N(0, V(d)), \quad (1.4)$$

where $V(d) = 2c_f\Gamma(1-2d)\sin(\pi d)\int_{-1}^1\int_{-1}^1 K(x)K(y)|x-y|^{2d-1}dxdy$, and c_f is a constant related to the spectral density of errors. Equation (1.4) shows that for $d > 0$ the convergence rate of $\hat{\beta}(t)$ is $(nb_n)^{d-1/2}$, which is much slower than the well-known $(nb_n)^{-1/2}$ convergence rate as given by (1.3) when $d = 0$. Therefore, a crucial problem of the statistical inference of model (1.1) is to test

$$H_0 : d = 0 \quad \text{versus} \quad H_A : 0 < d < 1/2. \quad (1.5)$$

The testing problem (1.5) for (1.1) is closely related to the existing tests of ‘spurious long memory’, which refers to the phenomenon that in the presence of regime changes, level shifts or certain deterministic trends, a short memory process could exhibit many properties of a long-memory process, known as the ‘spurious long-memory’ effects, see for example [Giraitis et al. \(2001\)](#), [Qu \(2011\)](#), and [McCloskey and Perron \(2013\)](#). These findings motivate the tests for distinguishing genuine and spurious long memory. Among others, [Qu \(2011\)](#), [Preuß and Vetter \(2013\)](#) and [Sibbertsen et al. \(2018\)](#) consider testing the null hypothesis of stationary long memory against spurious long memory. Meanwhile, several tests have been introduced to test the null hypothesis of spurious long memory, for which a prevailing approach is to assume a specific and parametric form of non-stationarity, see [Berkes et al. \(2006\)](#), [Harris et al. \(2008\)](#), and [Davis and Yau \(2013\)](#) among others. Recently, there has been growing interest in detecting long memory in the presence of general non-stationarity, see for example [Dette et al. \(2017\)](#) which considered locally stationary moving average formulation. In practice, by testing (1.5) for model (1.1), we are able to identify a new type of ‘spurious long memory’ resulting from the *misspecification in the conditional mean*. See our data analysis in [Section 6.1](#) where we apply our method to the Hong Kong circulatory and respiratory data.

The goal of the present work is to test the *hypothesis* (1.5) under complex and general temporal dynamics, assuming that $(\mathbf{x}_{i,n})$ and $(u_{i,n})$ belong to the flexible class of locally stationary processes generated by smoothly changing underlying mechanisms. Although some related literature has studied hypothesis (1.5) for linear regression models with deterministic covariates, see for example Harris et al. (2008), to the best of the authors' knowledge, this work is the first instance investigating testing (1.5) for (1.1) in the presence of time series covariates. In the literature, KPSS (Kwiatkowski, Phillips, Schmidt, and Shin, see Lee and Schmidt (1996)), R/S (range over standard deviation, see Hurst (1951)), V/S (rescaled variance, see Giraitis et al. (2003)), and K/S (which has a limiting distribution of Kolmogoroff-Smirnoff form, see Lima and Xiao (2004)) tests have been widely used for long memory detection in stationary processes. In this paper, we develop new KPSS, R/S, V/S and K/S-type tests tailored to the non-stationary time series time-varying regression problem (1.1). The limiting distributions of the test statistics under the null hypothesis, the local and fixed alternatives are then derived. Our results differ from their stationary counterparts due to the following reasons. (1) Under the null hypothesis, it is well-known that the KPSS, R/S, V/S and K/S tests are built on the partial sum process whose convergence rate is $n^{-1/2}$. However, the nonparametric estimate $\hat{\beta}(\cdot)$ induces stochastic errors much larger than $n^{-1/2}$ which will lead to different limiting distributions as well as possible degeneracy. (2) Due to the non-stationary errors and covariates, the partial sum processes cannot be approximated by processes with stationary increments, which makes the test statistics non-pivotal. The major contributions of the paper lie in the following three aspects.

Firstly, our methods are applicable to the locally stationary time series regression, which has found considerable attention in various related fields, see for instance Vogt (2012), Hu et al. (2019), Zhou and Wu (2010) and many others. In particular, the flexible locally stationary framework allows the error processes to display conditional and unconditional heteroscedasticity that has been increasingly investigated (see Harris and Kew (2017) and Cavaliere et al. (2020)) in the context of long-memory models. Both the evolving distributional properties of the locally stationary data and the long-memory properties pose long-standing challenges to the inference of time-varying coefficient linear model (1.1) due to the lack of general Gaussian approximation techniques for non-stationary long-memory processes. For stationary long-memory processes, Gaussian approximation has been studied by for example Dehling and Taqqu (1989). Recently, Wu and Zhou (2018b) developed Gaussian approximation schemes for a class of locally stationary long-memory linear processes. However, their results cannot accommodate regression problems with time series covariates which requires the analysis of distributional properties of the partial sum process of $(\mathbf{x}_{i,n}e_{i,n})_{i=1}^n$. In this paper, we address this issue via a further Gaussian approximation theorem, allowing $(\mathbf{x}_{i,n})$ and $(e_{i,n})$ to be non-stationary SRD and LRD processes, respectively, with flexible dependence between them.

Secondly, we develop effective bootstrap approaches which circumvent the difficult estimation of the non-pivotal limiting distributions of the test statistics under time series non-stationarity. In particular, the test statistics could degenerate when the time-varying mean, variance, long-run variance of errors and covariates, and the intercept lie in certain hyperplanes whose geometry cannot be directly identified from the data. Importantly, regardless of the degeneracy of test statistics, our bootstrap procedures are consistent and possess good finite sample properties. Furthermore, we show that the exact local asymptotic power of the four types of bootstrap-assisted tests can reach the order $O(\log^{-1} n)$ in the presence of time series covariates, no matter whether the test statistics degenerate under the null hypothesis. This rate coincides with Shao and Wu (2007) which studies a similar problem of testing the SRD null hypothesis against LRD local alternatives for strictly stationary series without covariates.

The rest of the paper is organized as follows. Section 2 introduces necessary notation. Section 3 formally states the non-stationary LRD model and the related assumptions. Section 4 provides the test statistics, and establishes the asymptotic results via the new Gaussian approximation theory for the product of non-stationary SRD and LRD processes. Section 5 discusses the bootstrap algorithms.

Section 6 reports the simulation results and the analysis of Hong Kong circulatory and respiratory data. Section 7 provides a brief concluding remark. In Appendix A, we provide the proof of the new Gaussian approximation theory. Detailed proofs, the literature review of KPSS and related tests, the implementation details including the selection of tuning parameters, additional simulation results, data analysis results (including COVID-19 data), and additional algorithms are relegated to the supplement.

2 Notation

For a matrix $\mathbf{A} = (a_{ij})_{1 \leq i \leq n, 1 \leq j \leq m} \in \mathbb{R}^{n \times m}$, let $|\mathbf{A}| = (\sum_{j=1}^m \sum_{i=1}^n a_{ij}^2)^{1/2}$ and write $\mathbf{A} \geq 0$ if \mathbf{A} is semi-positive definite. Let $(\mathbf{A})_{(1,1)}$ denote the element of \mathbf{A} in the first column and first row. Notice that when $m = 1$, \mathbf{A} is a vector. For $\mathbf{A} \geq 0$ with eigendecomposition $\mathbf{A} = \mathbf{Q}\mathbf{D}\mathbf{Q}^\top$ with orthonormal matrix \mathbf{Q} and diagonal matrix \mathbf{D} , the root of \mathbf{A} is defined by $\mathbf{A}^{1/2} = \mathbf{Q}\mathbf{D}^{1/2}\mathbf{Q}^\top$, where $\mathbf{D}^{1/2}$ is the elementwise root of \mathbf{D} . Let \mathbf{I}_p denote the p -dimensional identity matrix. For a random matrix \mathbf{A} , for $q \geq 1$, let $\|\mathbf{A}\|_q = (\mathbb{E}|\mathbf{A}|^q)^{1/q}$ denote the \mathcal{L}^q -norm of the random variable $|\mathbf{A}|$ and write $\|\cdot\| = \|\cdot\|_2$ for short. Write $\mathbf{A} \in \mathcal{L}^q$ if $\|\mathbf{A}\|_q < \infty$. For a function $f(\cdot)$, write $f \in C[0, 1]$ if f is continuous over $[0, 1]$, $f \in C^p[0, 1]$ if the p th order derivative of f is continuous over $[0, 1]$. Write $\mathbf{A} \in C[0, 1]$ and $\mathbf{A} \in C^p[0, 1]$ if each element $A_{ij}(\cdot)$ in $\mathbf{A}(\cdot)$ is in $C[0, 1]$ and $C^p[0, 1]$, respectively. For any kernel function $K(\cdot)$, let $K^*(\cdot)$ denote the jackknife equivalent kernel $2\sqrt{2}K(\sqrt{2}x) - K(x)$. Denote by $\lfloor x \rfloor$ the largest integer smaller or equal to x . For any two positive real sequences a_n and b_n , write $a_n \asymp b_n$ if $\exists 0 < c < C < \infty$ such that $c < \liminf_{n \rightarrow \infty} \frac{a_n}{b_n} < \limsup_{n \rightarrow \infty} \frac{a_n}{b_n} < C$. Let $t \wedge s$ denote the smaller value in t and s . Let \Rightarrow denote convergence in distribution. Write λ as Lebesgue measure on $[0, 1]$. Let $:=$ denote ‘defined as’.

3 Model assumptions

We start by introducing the time series time-varying regression model (1.1) in detail. Recall model (1.1) has the following form

$$y_{i,n} = \mathbf{x}_{i,n}^\top \boldsymbol{\beta}(t_i) + e_{i,n}, \quad i = 1, \dots, n, \quad \text{where } (1 - \mathcal{B})^d e_{i,n} = u_{i,n}, \quad d \in [0, 1/2).$$

We assume that the process $(u_{i,n})_{i=-\infty}^n$ and the covariate process $(\mathbf{x}_{i,n})_{i=1}^n$ have the form

$$u_{i,n} = H(t_i, \mathcal{F}_i), \quad \mathbf{x}_{i,n} = \mathbf{W}(t_i, \mathcal{F}_i), \quad (3.1)$$

where $\mathcal{F}_i = (\varepsilon_{-\infty}, \dots, \varepsilon_i)$, $(\varepsilon_i)_{i \in \mathbb{Z}}$ are *i.i.d.* random variables, H and $\mathbf{W} = (W_1, \dots, W_p)^\top$ are measurable functions such that $H : (-\infty, 1] \times \mathbb{R}^{\mathbb{Z}} \rightarrow \mathbb{R}$, $W_s : [0, 1] \times \mathbb{R}^{\mathbb{Z}} \rightarrow \mathbb{R}$, $2 \leq s \leq p$, while W_1 is fixed to be 1 corresponding to the intercept of the regression. Define $(\varepsilon'_i)_{i \in \mathbb{Z}}$ as an *i.i.d.* copy of $(\varepsilon_i)_{i \in \mathbb{Z}}$, and for $j \geq 0$, let $\mathcal{F}_j^* = (\mathcal{F}_{-1}, \varepsilon'_0, \varepsilon_1, \dots, \varepsilon_{j-1}, \varepsilon_j)$. For any (vector) process $\mathbf{L}(t, \mathcal{F}_i)$, it is said to be \mathcal{L}^q stochastic Lipschitz continuous in the interval \mathcal{I} (denoted by $\mathbf{L} \in \text{Lip}_q(\mathcal{I})$) if for $t_1, t_2 \in \mathcal{I}$, there exists a constant $M > 0$ such that

$$\|\mathbf{L}(t_1, \mathcal{F}_0) - \mathbf{L}(t_2, \mathcal{F}_0)\|_q \leq M|t_1 - t_2|.$$

We say the process $\mathbf{L}(t, \mathcal{F}_i)$ is *locally stationary* (LS) on \mathcal{I} if $\mathbf{L}(t, \mathcal{F}_i) \in \text{Lip}_q(\mathcal{I})$ for some $q \geq 2$. Write $\text{Lip}_q = \text{Lip}_q([0, 1])$ for short. The locally stationary process offers a flexible nonparametric device to characterise the complex temporal dynamics of the error and covariate processes in (3.1), which is based on Bernoulli shift processes and leads to a general framework of nonlinear processes, see Wu (2005). Other formulations of locally stationary processes include Dahlhaus (1997) and Nason et al. (2000). See

Dahlhaus et al. (2019) for a comprehensive review. The physical dependence measure of the nonlinear filter $\mathbf{L} \in \mathcal{L}^q$ ($q > 0$) over the interval \mathcal{I} is defined by

$$\delta_q(\mathbf{L}, k, \mathcal{I}) = \sup_{t \in \mathcal{I}} \|\mathbf{L}(t, \mathcal{F}_k) - \mathbf{L}(t, \mathcal{F}_k^*)\|_q.$$

The physical dependence measure $\delta_q(\mathbf{L}, k, \mathcal{I})$ quantifies the influence of the input ε_0 on the output $\mathbf{L}(t, \mathcal{F}_k)$ over the interval \mathcal{I} . Observe that $\delta_q(\mathbf{L}, k, \mathcal{I}) = 0$ if $k < 0$. Write $\delta_q(\mathbf{L}, k) = \delta_q(\mathbf{L}, k, [0, 1])$ for short. We proceed to define SRD and LRD non-stationary processes for non-stationary time series.

Definition 3.1. *The univariate process $(L(t, \mathcal{F}_i))_{i=-\infty}^\infty, t \in \mathcal{I}$ is said to be SRD if*

$$\sum_{j=-\infty}^\infty \sup_{t, s \in \mathcal{I}} |\text{Cov}(L(t, \mathcal{F}_0), L(s, \mathcal{F}_j))| < \infty,$$

and a LRD process otherwise. The p -dimensional vector process $(\mathbf{L}(t, \mathcal{F}_i))_{i=-\infty}^\infty, t \in \mathcal{I}$ is SRD if each of its component is SRD, and is otherwise LRD.

Definition 3.1 distinguishes the SRD and LRD by the uniform summability of covariance, which naturally extends the traditional definition of long memory in second-order stationary univariate processes, see for example Condition III in Chapter 2 of Pipiras and Taqqu (2017). The uniform long-memory definition has been introduced to define LRD and SRD non-stationary time series in Wu and Zhou (2018b). Various definitions of LRD stationary processes could be found in Pipiras and Taqqu (2017) and definitions of LRD locally stationary processes are discussed in Beran (2009), Dette et al. (2017) and Ferreira et al. (2018) among others. In this paper, we posit the following assumptions.

Assumption 3.1. *The zero mean SRD process $(u_{i,n})_{i=-\infty}^n$ satisfies*

(a1)' $H(t, \mathcal{F}_0) \in \text{Lip}_2(-\infty, 1]$, $\mathbb{E}H(t, \mathcal{F}_0) = 0$, and $\sup_{t \in (-\infty, 1]} \|H(t, \mathcal{F}_0)\|_4 < \infty$.

(a2)' $\delta_4(H, k, (-\infty, 1]) = O(\chi^k)$ for some $\chi \in (0, 1)$.

(a3)' Define the long-run variance function as $\sigma_H^2(t) := \sum_{k=-\infty}^\infty \text{Cov}(H(t, \mathcal{F}_0), H(t, \mathcal{F}_k))$, $t \in (-\infty, 1]$, which satisfies that $\inf_{t \in (-\infty, 1]} \sigma_H^2(t) > 0$ and $\sup_{t \in (-\infty, 1]} \sigma_H^2(t) < \infty$.

(a4)' $\sigma_H^2(\cdot)$ is twice continuously differentiable on $[0, 1]$.

Condition (a1)' imposes the assumptions of local stationarity and finite fourth moment on the innovations $(u_{i,n})$. Condition (a1)' will be satisfied if $\sup_{t \in (-\infty, 1]} \|\frac{\partial}{\partial t} H(t, \mathcal{F}_0)\| < \infty$. Condition (a2)' ensures that the innovations $(u_{i,n})$ are SRD and satisfy geometric measure contraction (GMC). Conditions (a3)' and (a4)' guarantee that the innovations $(u_{i,n})$ have a finite, non-degenerate and smooth long-run variance.

Notice that under the null hypothesis $d = 0$, the error process $(e_{i,n})$ reduces to $(u_{i,n})$, which indicates that $(e_{i,n})$ is a SRD process. When $d > 0$, $(e_{i,n})$ is generated by a binomial weighted combination of $u_{i,n} = H(t_i, \mathcal{F}_i)$ starting from the infinite past ($i = -\infty$). To stress that $(e_{i,n})$ is a LRD process under the alternative hypothesis with respect to d , in the remaining of this article we write $e_{i,n}$ as $e_{i,n}^{(d)}$ when $d > 0$, i.e., $e_{i,n}^{(d)} = (1 - B)^{-d} u_{i,n} = \sum_{k=0}^\infty \psi_k(d) u_{i-k}$, $\psi_j(d) = \Gamma(j + d) / [\Gamma(d) \Gamma(j + 1)]$. We further write $e_{i,n}^{(d)} = H^{(d)}(t_i, \mathcal{F}_i)$, where $H^{(d)}(t, \mathcal{F}_l) = \sum_{k=0}^\infty \psi_k(d) H(t - t_k, \mathcal{F}_{l-k})$. The following Proposition 3.1 elaborates that the physical dependence measure of $(e_{i,n}^{(d)})$ relies on d .

Proposition 3.1. *Under Assumption 3.1, we have uniformly for $l \geq 0$, $0 < d < 1/2$,*

$$\delta_p(H^{(d)}, l, (-\infty, 1]) = O\{(1 + l)^{d-1}\}.$$

Our formulation of $(e_{i,n})$ in (1.2) allows for a wide class of non-stationary SRD and LRD processes under H_0 and H_A , respectively, including the following examples.

Example 3.1 (Linear locally stationary process). *Consider the time-varying FARIMA(p, d, q) model ($0 < d < 1/2$) (recall that $u_{j,n} = H(t_j, \mathcal{F}_j)$, $-\infty \leq j \leq n$)*

$$(1 - \mathcal{B})^d e_{i,n} = u_{i,n}, \quad i = 1, 2, \dots, n, \text{ with } \Phi^p(\mathcal{B}, t)H(t, \mathcal{F}_j) = \Theta^q(\mathcal{B}, t)\varepsilon_j,$$

where $t \in (-\infty, 1]$, $j \in \mathbb{Z}$, $\Phi^p(z, t) = 1 + \phi_1(t)z + \dots + \phi_p(t)z^p$ and $\Theta^q(z, t) = 1 + \theta_1(t)z + \dots + \theta_q(t)z^q$ are polynomials with degrees p and q , and the random variables $(\varepsilon_i)_{i \in \mathbb{Z}}$ are i.i.d. with mean 0 and variance 1. Assume that for $t \in (-\infty, 1]$, $\{\phi_i(t), 1 \leq i \leq p\}$ and $\{\theta_j(t), 1 \leq j \leq q\}$ are twice differentiable, $\Phi^p(z, t)$ and $\Theta^q(z, t)$ do not share the same roots, and $\Phi^p(z, t)$ does not have roots in the unit disk $\{|z| \leq 1\}$. Then there exists real-valued differentiable functions $(a_i(t))_{i \geq 0}$ such that $A(z, t) = \Theta^q(z, t)/\Phi^p(z, t) = \sum_{i=0}^{\infty} a_i(t)z^i$, where for $t \in (-\infty, 1]$, $|a_j(t)|$ and $|a'_j(t)|$ are summable. Consequently, we have the MA(∞) representation

$$e_{i,n} = (1 - \mathcal{B})^{-d} u_{i,n} = (1 - \mathcal{B})^{-d} \sum_{j=0}^{\infty} a_j(t_i) \mathcal{B}^j \varepsilon_i := \sum_{j=0}^{\infty} b_{j,i} \varepsilon_{i-j}. \quad (3.2)$$

When $d = 0$, it follows that $b_{j,i} = a_j(t_i)$. Thus, $(e_{i,n})$ is SRD according to Definition 3.1. When $d > 0$, by Lemma 3.2 of [Kokoszka and Taqqu \(1995\)](#), $b_{j,i} = \sum_{l=0}^j \psi_l a_{j-l}(t_{i-l}) = L_i(j)j^{d-1}$, where $L_i(\cdot)$ is a slowly varying function for each i . Suppose for $1 \leq i \leq n$, $|L_i(j)| \leq L(j)$ for some slowly varying function $L(\cdot)$. Noticing that $\text{Cov}(e_{i,n}, e_{i+k,n}) = \sum_{j=0}^{\infty} b_{j,i} b_{j+k,i+k}$. By Proposition 2.2.9 in [Pipiras and Taqqu \(2017\)](#), $\sup_{1 \leq i \leq n} |\text{Cov}(e_{i,n}, e_{i+k,n})|$ is of order k^{2d-1} for all $i \in \mathbb{Z}$. Hence, (3.2) is LRD according to Definition 3.1.

Example 3.2 (Nonlinear locally stationary process). *Consider the time-varying ARFIMA(p, d, q)-GARCH(1, 1) process $(1 - \mathcal{B})^d e_{i,n} = u_{i,n}$, $1 \leq i \leq n$, where $u_{j,n} = H(t_j, \mathcal{F}_j)$ and*

$$\Phi^p(\mathcal{B}, t)H(t, \mathcal{F}_j) = \Theta^q(\mathcal{B}, t)v_j(t), \quad v_j(t) = \varepsilon_j \sigma_j(t), \quad j \in \mathbb{Z}, \quad t \in (-\infty, 1],$$

where $\sigma_j^2(t) = c(t) + \alpha(t)v_{j-1}^2(t) + \beta(t)\sigma_{j-1}^2(t)$, $(\varepsilon_i)_{i \in \mathbb{Z}}$ are i.i.d. random variables with mean 0 and variance 1, $c(t), \alpha(t), \beta(t)$ are smooth non-negative functions, and $\Phi^p(z, t) = 1 + \phi_1(t)z + \dots + \phi_p(t)z^p$ and $\Theta^q(z, t) = 1 + \theta_1(t)z + \dots + \theta_q(t)z^q$ are polynomials with degrees p and q . Assume that for $t \in (-\infty, 1]$, $\{\phi_i(t), 1 \leq i \leq p\}$, $\{\theta_j(t), 1 \leq j \leq q\}$, $c(t)$, $\alpha(t)$ and $\beta(t)$ are twice differentiable, $\Phi^p(z, t)$ and $\Theta^q(z, t)$ do not share the same roots, and $\Phi^p(z, t)$ does not have roots in the unit disk $\{|z| \leq 1\}$, $(\varepsilon_i)_{i \in \mathbb{Z}} \in \mathcal{L}^8$, $\sup_{t \in (-\infty, 1]} c(t) < \infty$ and $\sup_{t \in (-\infty, 1]} \|\alpha(t)\varepsilon_t^2 + \beta(t)\|_4 < 1$. Then, by Example 2 of [Wu and Zhou \(2011\)](#), $(u_{i,n})_{i=-\infty}^n$ satisfy (a2)' of Assumption 3.1. When $d > 0$, Definition 3.1 can be verified similarly as given in Example 3.1, since $(v_i(t))_{i \in \mathbb{Z}}$ are white noises.

4 Main results

Since $(e_{i,n})$ is not observable in (1.1), we propose to test H_0 based on nonparametric residuals. Specifically, we adopt the local linear approach (see for instance [Fan \(1993\)](#) and [Fan and Gijbels \(1996\)](#)) to estimate $\beta(t)$ in (1.1), i.e.,

$$(\hat{\beta}_{b_n}(t), \hat{\beta}'_{b_n}(t)) = \arg \min_{\boldsymbol{\eta}_0, \boldsymbol{\eta}_1 \in \mathbb{R}^p} \sum_{i=1}^n \{y_{i,n} - \mathbf{x}_{i,n}^\top \boldsymbol{\eta}_0 - \mathbf{x}_{i,n}^\top \boldsymbol{\eta}_1(t_i - t)\}^2 K_{b_n}(t_i - t), \quad (4.1)$$

where $K(\cdot)$ is a kernel function with finite support $[-1, 1]$ and b_n is a bandwidth and $K_{b_n}(\cdot) = K(\cdot/b_n)$. To further eliminate the bias term involving $\beta''(\cdot)$, we use the jackknife bias-corrected estimator in [Wu and Zhao \(2007\)](#) :

$$\tilde{\beta}_{b_n}(t) = 2\hat{\beta}_{b_n/\sqrt{2}}(t) - \hat{\beta}_{b_n}(t). \quad (4.2)$$

Then, we obtain the nonparametric residuals $(\tilde{e}_{i,n})$, i.e., $\tilde{e}_{i,n} = y_{i,n} - \mathbf{x}_{i,n}^\top \tilde{\beta}(t_i)$. For simplicity, define $\tilde{S}_{r,n} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_{i,n}$, $r = \lfloor nb_n \rfloor + 1, \dots, n - \lfloor nb_n \rfloor$. We consider four well-known types of partial sum based test statistics, which are KPSS, R/S, V/S and K/S-type tests built on $(\tilde{e}_{i,n})$.

1. KPSS-type statistic

$$T_n = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \sum_{r=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\tilde{S}_{r,n} \right)^2. \quad (4.3)$$

2. R/S-type statistic $Q_n = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{S}_{k,n} - \min_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{S}_{k,n}$.

3. V/S-type statistic $M_n = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \left\{ \sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \tilde{S}_{k,n}^2 - \frac{1}{n - 2\lfloor nb_n \rfloor} \left(\sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \tilde{S}_{k,n} \right)^2 \right\}$.

4. K/S-type statistic $G_n = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \left| \tilde{S}_{k,n} \right|$.

The above four types of tests have been widely applied to the detection of long memory and many other important problems (e.g. unit root testing) for stationary time series. We refer to Section B of the online supplement for the complete literature review and applications. To the best of our knowledge, all the existing work on the KPSS, R/S, V/S, and K/S tests considers the statistics based on the original series $(e_{i,n})$ or parametric residuals (e.g., the residuals obtained by the removal of the sample mean), and is therefore not applicable to the time-varying coefficient model (1.1). Meanwhile, it is well-known that the nonparametric estimators have a slower convergence rate than the corresponding parametric estimators. Therefore, the asymptotic properties of our nonparametric residual-based KPSS and related tests will be very different from their parametric residual-based or original series-based counterparts; in fact we show that the tests can degenerate under certain scenarios (see Theorem 4.1) and thus bootstrap procedures adaptive to the possible degeneracy are proposed for implementation, see Algorithms 5.1, also Algorithms G.1 and G.2 of the online supplement. In this article, we use the term ‘*KPSS and related tests*’ to represent the four types of tests.

4.1 Assumptions

For the sake of brevity, in this section we only discuss the KPSS-type test statistic in detail, and summarize the results of KPSS-related test statistics in Remark 4.4. In order to investigate the asymptotic properties of T_n defined by (4.3) in the presence of time series covariates, we introduce the following assumptions.

Assumption 4.1. *The kernel function $K(\cdot)$ is continuous, symmetric and supported on $[-1, 1]$.*

Assumption 4.2. *Each coordinate of $\beta(\cdot)$, i.e., $\beta_i(\cdot)$ for $i = 1, \dots, p$, lies in $C^3[0, 1]$.*

Assumption 4.3. *Let $\mathbf{U}(t, \mathcal{F}_i) = H(t, \mathcal{F}_i) \mathbf{W}(t, \mathcal{F}_i)$, $i = 1, \dots, n$, s.t.*

(A1) $\mathbf{U}(t, \mathcal{F}_i) \in \text{Lip}_2$, $\sup_{t \in [0, 1]} \|\mathbf{U}(t, \mathcal{F}_i)\|_4 < \infty$.

(A2) *Short-range dependence:* $\delta_4(\mathbf{U}, k) = O(\chi^k)$ for some $\chi \in (0, 1)$.

(A3) *The smallest eigenvalue of $\Sigma(t) := \sum_{j=-\infty}^{\infty} \text{Cov} \{ \mathbf{U}(t, \mathcal{F}_0), \mathbf{U}(t, \mathcal{F}_j) \}$, $t \in [0, 1]$, (i.e., the long-run covariance matrix) is bounded away from 0 on $[0, 1]$.*

Assumption 4.3 is standard for local linear time series regression, see for instance Zhou and Wu (2010). Notice that the first element of $\mathbf{U}(t, \mathcal{F}_i)$ is $H(t, \mathcal{F}_i)$. When $p = 1$ (the time-varying trend model), Assumption 4.3 reduces to (a3)' in Assumption 3.1 with $t \in [0, 1]$.

Define $\mathbf{M}(t) := \mathbb{E}(\mathbf{W}(t, \mathcal{F}_0)\mathbf{W}(t, \mathcal{F}_0)^\top)$, $t \in [0, 1]$. Notice that the first element of \mathbf{W} is 1. Write $\boldsymbol{\mu}_W(\cdot) := (1, \mu_{W,2}(\cdot), \dots, \mu_{W,p}(\cdot))^\top := \mathbb{E}(\mathbf{W}(t, \mathcal{F}_0))$. For $p \geq 2$, Let $x_{i,j,n}$ denote the j th element in $\mathbf{x}_{i,n}$, and $\mathbf{x}_{i,n}^{(-1)} := (x_{i,2,n}, \dots, x_{i,p,n})^\top$. Define $\boldsymbol{\mu}_W^{(-1)}(\cdot) := (\mu_{W,2}(\cdot), \dots, \mu_{W,p}(\cdot))^\top$, $\mathbf{W}^{(-1)}(\cdot, \cdot) := (W_2(\cdot, \cdot), \dots, W_p(\cdot, \cdot))^\top$.

Assumption 4.4. *The following conditions hold for the covariates when $p \geq 2$:*

(B1) *The smallest eigenvalue of $\mathbf{M}(t)$ is bounded away from 0 on $[0, 1]$.*

(B2) *$\mathbf{M}(\cdot) \in C^1[0, 1]$, $\boldsymbol{\mu}_W(\cdot) \in C^1[0, 1]$.*

(B3) *$\mathbf{W}(t, \mathcal{F}_i) \in \text{Lip}_2$, and $\sup_{t \in [0, 1]} \|\mathbf{W}(t, \mathcal{F}_i)\|_8 < \infty$, $i = 1, \dots, n$.*

(B4) *$\delta_8(\mathbf{W}^{(-1)}, k) = O(\chi^k)$ for some $\chi \in (0, 1)$.*

(B5) *$\mathbb{E}(H(t_j, \mathcal{F}_j) | \mathbf{W}(t_j, \mathcal{F}_j)) = 0$ for $j = 1, 2, \dots, n$, a.s..*

Condition (B1) ensures that there is no multicollinearity among the explanatory variables. Assumption (B2) guarantees that the $\mathbf{M}(\cdot)$ and $\boldsymbol{\mu}_W(\cdot)$ have continuous derivatives. Assumption (B3) requires that the covariates are locally stationary. Condition (B4) imposes that each component of $\mathbf{W}^{(-1)}(\cdot, \cdot)$ is SRD. Condition (B5) assumes that $\mathbf{x}_{i,n}$ is a p -dimensional random vector uncorrelated with innovations, which is necessary for model identification. Our assumptions are very mild in the sense that we allow nonlinearity and heteroscedasticity for the covariates and errors, as well as the correlation between $\mathbf{x}_{i,n}$ and $e_{i,n}$. Assumptions 4.3 and 4.4 can be verified using similar arguments in Zhou and Wu (2009). When $p = 1$, we use the convention that $\mathbf{x}_{i,n}^{(-1)} = \emptyset$, $\mathbf{W}^{(-1)}(\cdot, \cdot) = \emptyset$, and $\boldsymbol{\mu}_W^{(-1)}(\cdot) = \emptyset$, $\mathbf{M}(\cdot) = \boldsymbol{\mu}_W(\cdot) = 1$. Therefore Assumption 4.4 always hold in this case.

Assumption 4.5. $\lambda(\boldsymbol{\Sigma}^{1/2}(u)\mathbf{M}^{-1}(u)\boldsymbol{\mu}_W^\top(u) \neq (\sigma_H(u), 0, \dots, 0)^\top) > 0$.

The test statistics will degenerate under the null hypothesis when the parameters of the regression model (1.1) lie in the hyperplane $\{u \in [0, 1] : \boldsymbol{\Sigma}^{1/2}(u)\mathbf{M}^{-1}(u)\boldsymbol{\mu}_W^\top(u) = (\sigma_H(u), 0, \dots, 0)^\top\}$. Assumption 4.5 excludes such situation.

4.2 Asymptotic theory

The following theorem establishes the asymptotic distribution of the KPSS-type statistic T_n (4.3) under the null hypothesis.

Theorem 4.1. *Let Assumptions 4.1, 4.2, 4.3 and 4.4 be satisfied, assuming $nb_n^6 \rightarrow 0$, $nb_n^{7/2}/(\log n)^4 \rightarrow \infty$, we have that under the null hypothesis:*

(i) *If Assumption 4.5 holds,*

$$T_n \Rightarrow \int_0^1 U^2(t)dt, \quad (4.4)$$

where $U(t)$ is a zero mean continuous Gaussian process with covariance function

$$\begin{aligned} \mathbb{E}(U(r)U(s)) &=: \gamma(r, s) = \int_0^{r \wedge s} \sigma_H^2(u)du - 2 \int_0^{r \wedge s} \{\boldsymbol{\mu}_W^\top(u)\mathbf{M}^{-1}(u)\boldsymbol{\Sigma}^{1/2}(u)\}_1 \sigma_H(u)du \\ &\quad + \int_0^{r \wedge s} \boldsymbol{\mu}_W^\top(u)\mathbf{M}^{-1}(u)\boldsymbol{\Sigma}(u)\mathbf{M}^{-1}(u)\boldsymbol{\mu}_W(u)du, \quad r, s \in [0, 1], \end{aligned}$$

where $\{\cdot\}_1$ denotes the first element of a vector.

(ii) If Assumption 4.5 fails, then $s_1^{-1}(T_n/b_n - s_2) \Rightarrow \chi_1^2$, where s_1 and s_2 are constants, i.e., $s_1 = 2\sigma_H^2(0) \int_0^1 \left(\int_{v-1}^1 K^*(t) dt \right)^2 dv$, and $s_2 = 2 \int_0^1 \sigma_H^2(t) dt \int_0^1 \left(\int_v^1 K^*(t) dt \right)^2 dv$.

Theorem 4.1 reveals that for the time-varying coefficient model with time series covariates, the limiting distribution of T_n depends on the time-varying mean and covariance matrix of the covariates $\mathbf{x}_{i,n}$ as well as the long-run covariance matrix of $\mathbf{x}_{i,n}e_{i,n}$. Theorem 4.1 is very general since it posits neither the specific form of heteroscedasticity nor the parametric form of the errors. When Assumption 4.5 is violated, (ii) shows that T_n degenerates with asymptotic variance $2s_1^2b_n^2$. After standardization, it converges to χ_1^2 in distribution. The results of R/S, V/S and K/S follow similarly. An important scenario that T_n degenerates under H_0 (i.e., $d = 0$) is the following time-varying trend model corresponding to $p = 1$, i.e.,

$$y_{i,n} = \beta_1(t_i) + e_{i,n}, \quad i = 1, \dots, n, \text{ with } (1 - \mathcal{B})^d e_{i,n} = u_{i,n}, \quad d \in [0, 1/2). \quad (4.5)$$

Remark 4.1. Theorem 4.1 and other theoretical results in this paper are valid for $p = 1$, with $\beta(t)$ replaced by $\beta_1(t)$, $\Sigma(t)$ replaced by $\sigma_H^2(t)$, $\mu_W(t)$ replaced by 1.

4.2.1 Gaussian approximation

Since the KPSS and related test statistics are constructed based on the partial sum process, to derive the asymptotic properties of the test statistics under the alternative hypothesis, we first study the Gaussian approximation of $\sum_{i=1}^r \mathbf{x}_{i,n}e_{i,n}^{(d)}$, $1 \leq r \leq n$, which is the partial sum of the product of a SRD and a LRD time series under H_A . Though Gaussian approximation theory for stationary processes (see for instance Wu (2007), Dehling and Taqqu (1989), Wu and Shao (2006) and the reference therein) has been successfully established and widely applied to many fields of statistics, there are only a few results of Gaussian approximation for locally stationary processes. Among them, Wu and Zhou (2011) established a flexible Gaussian approximation framework for locally stationary SRD processes, which has served as a fundamental key to the inference of SRD (piecewise) locally stationary processes and functional time series, see for instance Chen and Song (2015) and Wu and Zhou (2018a). Wu and Zhou (2018b) proposed a Gaussian approximation scheme for a class of locally stationary linear LRD processes. However, all the existing Gaussian approximation approaches are not applicable to the partial sum process of the product series $(\mathbf{x}_{i,n}e_{i,n}^{(d)})$, which serves as the crucial ingredient for establishing the limiting distribution of T_n under H_A . To this end, we shall provide a general Gaussian approximation theorem for the product of LRD and SRD processes. In the remaining of this paper, let $d_n = c/\log n$, where c is a positive constant. We substitute d with d_n to differentiate the notation under the fixed alternatives ($d > 0$) and that under the local alternatives ($d = c/\log n$).

Theorem 4.2. Under Assumptions 3.1 and 4.4, on a richer probability space, we have:

(i) There exists $\mathbf{R}_{k,n} = \sum_{j=0}^{\infty} \mu_W(t_k) \psi_j(d) \sigma_H(t_{k-j}) v_{k-j}$, where the random variables $(v_i)_{i \in \mathbb{Z}}$ are i.i.d. $N(0, 1)$, s.t.

$$\max_{[nb_n]+1 \leq r \leq n-[nb_n]} \left| \sum_{i=[nb_n]+1}^r (\mathbf{x}_{i,n}e_{i,n}^{(d)} - \mathbf{R}_{i,n}) \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d + n^{1+\alpha_0(d-1/2)}),$$

where $\alpha_0 \in (1, 4/3)$ and therefore $n^{1+\alpha_0(d-1/2)} = o(n^{d+1/2})$.

(ii) Further under Assumption 4.3, there exists a sequence of Gaussian processes

$$\tilde{\mathbf{R}}_{i,n} = \sum_{j=1}^{\infty} \mu_W(t_i) \psi_j(d_n) \sigma_H(t_{i-j}) V_{i-j,1} + \Sigma^{1/2}(t_i) \mathbf{V}_i,$$

where \mathbf{V}_i , $1 \leq i \leq n$, are i.i.d. $N(\mathbf{0}, \mathbf{I}_p)$, s.t.

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r (\mathbf{x}_{i,n} e_{i,n}^{(d_n)} - \tilde{\mathbf{R}}_{i,n}) \right| = o_{\mathbb{P}}(n^{1/2}).$$

Since $\|\mathbf{R}_{i,n}\| \asymp n^{d+1/2}$ and $\|\tilde{\mathbf{R}}_{i,n}\| \asymp n^{1/2}$, the approximation errors of Theorem 4.2 (i) and (ii) are asymptotically negligible. It can be also verified that the process $(\mathbf{R}_{k,n})_{k=1}^n$ is a locally stationary LRD Gaussian process defined by Definition 3.1.

Remark 4.2. The results of (i) consist of two parts. The rate $\sqrt{n}(\log n)^d$ is due to the approximation of $\max_{1 \leq s \leq n} |\sum_{k=1}^s (\mathbf{x}_{k,n} - \boldsymbol{\mu}_W(t_k)) e_{k,n}^{(d)}|$, see Proposition A.1. The rate $n^{1+\alpha_0(d-1/2)}$ is due to the approximation of $\max_{1 \leq s \leq n} |\sum_{k=1}^s \boldsymbol{\mu}_W(t_k) e_{k,n}^{(d)} - \mathbf{R}_{k,n}|$, which extends Theorem 2 in Wu and Zhou (2018b), see Proposition A.2. Specifically, we allow the driving shocks $(u_{i,n})_{i=-\infty}^n$ to be both dependent and heteroscedastic, while Wu and Zhou (2018b) assumed $(u_{i,n})_{i=-\infty}^n$ to be independent.

It is worth pointing out that (ii) is not an direct consequence of (i). Letting $d = d_n$ in (i), the rate $\sqrt{n}(\log n)^d$ in (i) will eventually lead to a trivial bound, i.e., $O_{\mathbb{P}}(\sqrt{n})$, which is of the same order as the partial sum $\sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_{i,n} e_{i,n}^{(d_n)}$, $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$.

Based on the Gaussian approximation result, we proceed to study the limiting distributions of KPSS and related statistics under the fixed and local alternatives for the time-varying coefficient model (1.1) with time series covariates. For the sake of brevity, we focus on the KPSS-type statistics. The results for R/S, V/S, and K/S-type statistics can be derived similarly and are summarized in Remark 4.4 and Appendix H.2 in the online supplement.

4.2.2 Fixed alternatives

In the following theorem, we establish the asymptotic distribution of the KPSS-type statistic T_n (4.3) under the fixed alternatives.

Theorem 4.3. Under Assumptions 3.1, 4.1, 4.2 and 4.4, assuming $nb_n^4/(\log n)^2 \rightarrow \infty$, $nb_n^6 \rightarrow 0$, and $\lambda(|\boldsymbol{\mu}_W^{(-1)}(\cdot)| \neq 0) > 0$, then we have under H_A with long-memory parameter d ,

$$T_n \Gamma^2(d+1)/n^{2d} \Rightarrow \int_0^1 U_d^2(t) dt,$$

where $U_d(t)$ is a zero mean continuous Gaussian process with covariance function

$$\mathbb{E}(U_d(r)U_d(s)) := \gamma_d(r, s) = \int_{-\infty}^{r \wedge s} \sigma_H^2(v) \lambda_d(r, v) \lambda_d(s, v) dv, \quad r, s \in [0, 1],$$

where for $v \leq u \in [0, 1]$, $\lambda_d(u, v) = d \int_{(-v)_+}^{(u-v)_+} t^{d-1} (\check{M}_W(t+v) - 1) dt$, $\check{M}_W(t) = \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\mu}_W(t)$, $t \in [0, 1]$.

Theorem 4.3 proves that the test statistic diverges to infinity at the rate of n^{2d} . Furthermore, from Theorem 4.3, we observe that the limiting distribution of T_n under the fixed alternatives is independent of $\boldsymbol{\Sigma}(t)$ except $\sigma_H^2(t)$, which is its $(1, 1)$ component, while under the null hypothesis the limiting distribution relies on all the components of $\boldsymbol{\Sigma}(t)$, see Theorem 4.1. This is because when $d > 0$, the stochastic fluctuation of the SRD components $(\mathbf{x}_{i,n})$ is asymptotic negligible compared to that of $(e_{i,n}^{(d)})$.

Straightforward calculation shows that $\check{M}_W(t) \geq 1$. The condition $\lambda(|\boldsymbol{\mu}_W^{(-1)}(\cdot)| \neq 0) > 0$ ensures $\check{M}_W(t) > 1$ in an interval with positive length such that $\lambda_d(u, v) > 0$ and excludes the scenario that all the stochastic covariates are zero mean during the whole period (see [Remark 5.1](#) for detailed discussion of such scenario) as well as the time-varying trend model, i.e., (4.5) with $p = 1$. When $\lambda_d(u, v) = 0$, we can show that $T_n = o_{\mathbb{P}}(n^{2d})$, i.e., the test statistic is degenerate under H_A .

Remark 4.3. *The error model (1.2) is in fact a Type I fractional $I(d)$ process when $d > 0$. If a locally stationary Type II fractional $I(d)$ error process is considered, i.e., $(1 - \mathcal{B})^d e_{i,n} = u_{i,n} \mathbf{1}(i \geq 1)$, the limiting distribution of T_n is almost the same except that the lower bound of the integral in $\gamma_d(r, s)$ is 0 instead of $-\infty$. We refer to [Marinucci and Robinson \(1999\)](#) for the definition of Type I and Type II fractional $I(d)$ processes.*

4.2.3 Local alternatives

The following theorem presents the asymptotic distribution of the KPSS-type statistic T_n (4.3) under the local alternatives.

Theorem 4.4. *Let the conditions of [Theorem 4.3](#) and [Assumption 4.3](#) hold. Then under H_A with $d_n = c/\log n$ for a constant $c > 0$, we have*

$$T_n \Rightarrow \int_0^1 (U^\circ(t))^2 dt,$$

where $U^\circ(t)$ is a zero mean continuous Gaussian process with covariance function

$$\mathbb{E}(U^\circ(r)U^\circ(s)) =: \gamma^\circ(r, s) = \check{\gamma}(r, s) + \gamma(r, s) + 2\check{\gamma}(r, s), \quad r, s \in [0, 1],$$

where $\gamma(r, s)$ is defined in [Theorem 4.1](#), and

$$\begin{aligned} \check{\gamma}(r, s) &= (e^c - 1) \int_0^{r \wedge s} \sigma_H(t) (\{\boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t)\}_1 - \sigma_H(t)) (\check{M}_W(t) - 1) dt, \\ \check{\gamma}(r, s) &= (e^c - 1)^2 \int_0^{r \wedge s} \sigma_H^2(t) (\check{M}_W(t) - 1)^2 dt, \end{aligned}$$

where $\{\cdot\}_1$ and $\check{M}_W(t)$ are as defined in [Theorem 4.1](#) and [Theorem 4.3](#), respectively.

From [Theorem 4.4](#), we shall see that under the local alternatives $d_n = c/\log n$, the KPSS-type statistic converges to a distribution depending on the mean and covariance matrix of $(\mathbf{x}_{i,n})$, the long-run covariance matrix of $(\mathbf{x}_{i,n} e_{i,n})$, as well as the parameter c . Careful examination of the proof of [Theorem 4.4](#) shows that T_n will converge to the limit in [Theorem 4.1](#) if $d_n \log n = o(1)$, indicating that the exact local power of the KPSS-type test is $O(\log^{-1} n)$ for model (1.1). For the time-varying trend model (4.5), it can be shown that the test statistic is degenerate (since $\lambda(|\boldsymbol{\mu}_W^{(-1)}(\cdot)| \neq 0) = 0$) under local alternatives $d_n = c/\log n$, i.e., $T_n = o_{\mathbb{P}}(1)$. The results for R/S, K/S and V/S-type tests follow similarly.

Remark 4.4. *The limiting behavior of R/S, V/S and K/S-type statistics defined in [Section 4](#) under [Assumption 4.5](#) can be derived by [Theorems 4.1](#), [4.3](#) and [4.4](#) as well as an application of continuous mapping theorem. Their limiting distributions are functions of $U(t)$, $U_d(t)$, $U^\circ(t)$ defined therein, see [Appendix H.2](#) in the online supplement for the exact forms.*

5 The bootstrap-assisted procedure

Section 4 shows that under the null hypothesis, the limiting distributions of KPSS and related test statistics are functions of the Gaussian process $U(t)$ which involves parameters $\boldsymbol{\mu}_W(t), \mathbf{M}(t), \boldsymbol{\Sigma}(t)$ (or $\sigma_H^2(t)$ when $p = 1$). Furthermore, the magnitude and specific form of the limiting distributions depend on whether Assumption 4.5 holds, which is usually unknown in practice. Therefore, it's impossible to obtain the critical values by directly simulating the Gaussian process $U(t)$. In this section, we provide a consistent bootstrap approach Algorithm 5.1 which mimics the asymptotic behavior of T_n under the null hypothesis no matter whether Assumption 4.5 is satisfied and yields valid simulated critical values. In addition, the estimation of $\boldsymbol{\mu}_W(\cdot)$ is not required in our proposed bootstrap tests. More precisely, we employ $\mathbf{x}_{i,n}^\top \hat{\mathbf{M}}^{-1}(t)$ instead of $\hat{\boldsymbol{\mu}}_W^\top(t) \hat{\mathbf{M}}^{-1}(t)$, where $\hat{\boldsymbol{\mu}}_W(t)$ stands for any consistent estimator of $\boldsymbol{\mu}_W(t)$, and the validity of the former construction can be easily verified by noting that in (5.1) of Algorithm 5.1, the Gaussian multiplier is independent of $\mathbf{x}_{i,n}$, $\hat{\mathbf{M}}(\cdot)$ and $\hat{\boldsymbol{\Sigma}}(\cdot)$ where the consistent estimators $\hat{\mathbf{M}}(t)$ and $\hat{\boldsymbol{\Sigma}}(t)$ will be discussed later. Thus, the difference between (5.1) and the counterpart with $\boldsymbol{\mu}_W(t)$ can be controlled by the convolution of standard Gaussian multipliers and the partial sum of the zero mean SRD process containing $(\mathbf{x}_{i,n} - \boldsymbol{\mu}_W(t_i))$. We only discuss Algorithm 5.1 for the KPSS-type test when $p \geq 2$ in detail in this section. The other algorithms, including Algorithm G.1 in the online supplement for $p = 1$ case corresponding to the time-varying trend model (4.5), and Algorithm G.2 for R/S, V/S, K/S-type tests are moved to the supplement.

Algorithm 5.1 The KPSS-type test for time-varying coefficient models

1. Select the smoothing parameters m , b_n , and τ_n , according to Appendix C.
2. Calculate $\tilde{e}_{i,n} = y_{i,n} - \mathbf{x}_{i,n}^\top \tilde{\boldsymbol{\beta}}(t_i)$, $i = 1, 2, \dots, n$, using local linear regression (4.1) and jackknife correction (4.2). Compute the KPSS-type statistic T_n in (4.3).
3. Calculate $\hat{\mathbf{M}}(t)$ and $\hat{\boldsymbol{\Sigma}}(t)$, the estimators of $\mathbf{M}(t)$ and $\boldsymbol{\Sigma}(t)$ defined in (5.3) and (5.4), respectively.
4. Generate B (say 2000) *i.i.d.* copies of $N(\mathbf{0}, \mathbf{I}_p)$ vectors $\mathbf{V}_i^{(r)} = (V_{i,1}^{(r)}, \dots, V_{i,p}^{(r)})^\top$, $1 \leq r \leq B$, then calculate (notice that $\hat{\sigma}_H^2(t) = (\hat{\boldsymbol{\Sigma}}(t))_{1,1}$)

$$\tilde{G}_k^{(r)} = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=[nb_n]+1}^k \mathbf{x}_{i,n}^\top \hat{\mathbf{M}}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \hat{\boldsymbol{\Sigma}}^{1/2}(t_j) \mathbf{V}_j^{(r)} + \sum_{i=[nb_n]+1}^k \hat{\sigma}_H(t_i) V_{i,1}^{(r)}, \quad (5.1)$$

and the bootstrap version of the KPSS-type statistic (4.3)

$$\tilde{T}_n^{(r)} = \frac{1}{n(n - 2[nb_n])} \sum_{s=[nb_n]+1}^{n-[nb_n]} \left(\sum_{k=[nb_n]+1}^s \tilde{G}_k^{(r)} \right)^2. \quad (5.2)$$

5. Let $\tilde{T}_{n,(1)} \leq \tilde{T}_{n,(2)} \leq \dots \leq \tilde{T}_{n,(B)}$ be the ordered statistics of $\{\tilde{T}_n^{(r)}\}_{r=1}^B$. Reject H_0 at level α if $T_n > \tilde{T}_{n,([B(1-\alpha)])}$. Let $B^* = \max\{r : \tilde{T}_{n,(r)} \leq T_n\}$. Then the p -value of the KPSS-type test is $1 - B^*/B$.
-

To implement Algorithm 5.1, we need to obtain the estimators $\hat{\mathbf{M}}(t)$ and $\hat{\boldsymbol{\Sigma}}(t)$. For $\hat{\mathbf{M}}(t)$ we propose the following estimator that uses directly observed covariates $\mathbf{x}_{i,n}$:

$$\hat{\mathbf{M}}(t) = \frac{1}{n\eta_n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top K_{\eta_n}(t_i - t^*), \quad (5.3)$$

where $t^* = \max\{\eta_n, \min(t, 1 - \eta_n)\}$ for some bandwidth $\eta_n \rightarrow 0$, $n\eta_n^2 \rightarrow \infty$. Under Assumptions 4.1 and

4.4, after a careful investigation of Lemma 6 of [Zhou and Wu \(2010\)](#), we have $\sup_{t \in [\eta_n, 1 - \eta_n]} |\hat{\mathbf{M}}(t) - \mathbf{M}(t)| = o_{\mathbb{P}}(1)$, i.e. $\hat{\mathbf{M}}(t)$ is uniformly consistent, see Lemma H.1 in the online supplement for details.

Observing that $\Sigma(\cdot)$ depends on the unobserved error process $(e_{i,n})$. Therefore, a common approach to estimate $\Sigma(\cdot)$ is to utilize $\hat{e}'_{i,n}s$ from the local linear fit (4.1), see for instance [Zhou and Wu \(2010\)](#) and [Vogt and Dette \(2015\)](#). Such plug-in estimate will yield test results that are sensitive to the choices of b_n under H_0 , and even more sensitive under H_A . We also find through our extensive numerical studies (which are not reported in this paper due to limited space) that the power of the test using the plug-in estimator of long-run covariance is often unsatisfactory. Therefore, we adopt a difference-based estimator that does not involves $\hat{e}'_{i,n}s$.

For $p = 1$ we recommend the difference statistic proposed in (4.7) of Section 4.2 of [Dette and Wu \(2019\)](#) for $\sigma_H^2(t)$ which is built on the difference of $y_{i,n}$. For $p \geq 2$, it can be shown that a direct extension of the estimator based on the difference of $(\mathbf{x}_{i,n}y_{i,n})$ is asymptotically biased. Therefore, we adopt the following bias-corrected difference-based estimator proposed by [Bai and Wu \(2023\)](#). Let $\mathbf{Q}_{k,m} = \sum_{i=k}^{k+m-1} \mathbf{x}_{i,n}y_{i,n}$, for $t \in [m/n, 1 - m/n]$,

$$\Delta_j = \frac{\mathbf{Q}_{j-m+1,m} - \mathbf{Q}_{j+1,m}}{m}, \quad \dot{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\Delta_j\Delta_j^\top}{2} \omega(t, j),$$

where $\omega(t, i) = K_{\tau_n}(t_i - t) / \sum_{i=1}^n K_{\tau_n}(t_i - t)$ for some bandwidth τ_n and the kernel function $K(\cdot)$ with support $(-1, 1)$. For $t \in [0, m/n]$, set $\dot{\Sigma}(t) = \dot{\Sigma}(m/n)$. For $t \in (1 - m/n, 1]$, set $\dot{\Sigma}(t) = \dot{\Sigma}(1 - m/n)$. The bias-corrected difference-based estimator $\hat{\Sigma}(t)$ for $t \in [0, 1]$ is then defined as:

$$\hat{\Sigma}(t) = \dot{\Sigma}(t) - \check{\Sigma}(t), \quad \text{where} \quad \check{\Sigma}(t) = \sum_{j=m}^{n-m} \frac{m\hat{\mathbf{A}}_j\hat{\mathbf{A}}_j^\top}{2} \omega(t, j), \quad (5.4)$$

where $\hat{\mathbf{A}}_j = \frac{1}{m} \sum_{i=j-m+1}^j (\mathbf{x}_{i,n}\mathbf{x}_{i,n}^\top \check{\beta}(t_i) - \mathbf{x}_{i+m,n}\mathbf{x}_{i+m,n}^\top \check{\beta}(t_{i+m}))$, $\check{\beta}(t) = \Omega^{-1}(t)\varpi(t)$, where $\Omega(t)$, $\varpi(t)$ are the smoothed versions of $\dot{\Delta}_j := \frac{1}{m} \sum_{i=j-m+1}^j \tilde{\mathbf{X}}_{i,m}\tilde{\mathbf{X}}_{i,m}^\top$ and $\check{\Delta}_j := \frac{1}{m} \sum_{i=j-m+1}^j \tilde{\mathbf{X}}_{i,m}^\top \tilde{\mathbf{Y}}_{i,m}$, i.e., $\Omega(t) = \sum_{j=m}^{n-m} \dot{\Delta}_j \omega(t^*, j)/2$, and $\varpi(t) = \sum_{j=m}^{n-m} \check{\Delta}_j \omega(t^*, j)/2$. To make [Algorithm 5.1](#) operational, it's necessary to select smoothing parameters η_n , τ_n and m for $\hat{\mathbf{M}}(t)$ and $\hat{\Sigma}(t)$. The selection of smoothing parameters is postponed to Appendix C of the online supplement.

5.1 The limiting behavior of the bootstrap tests

In this section, we shall show the asymptotic correctness of the bootstrap test [Algorithm 5.1](#). We shall also prove that the power of [Algorithm 5.1](#) against $d > 0$ can be no less than $(n/m)^{2d}$, and that the exact local power of [Algorithm 5.1](#) can achieve the order $O(\log^{-1} n)$.

We start by defining the long-run cross covariance vector between the locally stationary processes $(\mathbf{U}(t, \mathcal{F}_i))$ and $(H(t, \mathcal{F}_j))$.

Definition 5.1. Define the long-run cross covariance vector $\mathbf{s}_{UH}(t) \in \mathbb{R}^p$ by

$$\mathbf{s}_{UH}(t) = \sum_{j=-\infty}^{\infty} \text{Cov}(\mathbf{U}(t, \mathcal{F}_0), H(t, \mathcal{F}_j)), \quad t \in [0, 1].$$

When $p = 1$, $\mathbf{s}_{UH}(t)$ degenerates into $\sigma_H^2(t)$. We assume the following conditions for $\hat{\Sigma}(t)$ used in [Algorithm 5.1](#). Write $\hat{\Sigma}(t)$ defined in (5.4) as $\hat{\Sigma}_d(t)$ under the fixed alternatives and $\hat{\Sigma}_{d_n}(t)$ under the local alternatives d_n . Let $\mathcal{I} = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$, where $\gamma_n = \tau_n + (m + 1)/n$.

Assumption 5.1. *The long-run variance estimator satisfies the following conditions*

(i) *Under the null hypothesis,*

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}(t) - \Sigma(t) \right| = o_{\mathbb{P}}(b_n / \log^2 n).$$

(ii) *Under the fixed alternatives,*

$$\sup_{t \in \mathcal{I}} \left| m^{-2d} \hat{\Sigma}_d(t) - \Sigma_d(t) \right| = o_{\mathbb{P}}(1),$$

where $\Sigma_d(t) = \kappa_2(d) \sigma_H^2(t) \boldsymbol{\mu}_W(t) \boldsymbol{\mu}_W^\top(t)$, $t \in [0, 1]$, and $\kappa_2(d) = \Gamma^{-2}(d+1) \int_0^\infty (t^d - (t-1)_+^d)(2t^d - (t-1)_+^d - (t+1)^d) dt$.

(iii) *Under the local alternatives $d_n = c / \log n$, $m = \lfloor n^{\alpha_1} \rfloor$, $\alpha_1 \in (0, 1)$,*

$$\sup_{t \in \mathcal{I}} \left| \hat{\Sigma}_{d_n}(t) - \check{\Sigma}(t) \right| = o_{\mathbb{P}}(1),$$

where $\check{\Sigma}(t) := \Sigma(t) + (e^{c\alpha_1} - 1)^2 \sigma_H^2(t) \boldsymbol{\mu}_W(t) \boldsymbol{\mu}_W^\top(t) + (e^{c\alpha_1} - 1)(\mathbf{s}_{UH}(t) \boldsymbol{\mu}_W^\top(t) + \boldsymbol{\mu}_W(t) \mathbf{s}_{UH}^\top(t))$.

It can be shown that the both the plug-in estimator of Zhou and Wu (2010) and the bias-corrected estimator (5.4) satisfies this condition under suitable bandwidth conditions following Theorem 3.1 of Wu and Shao (2006) and the chaining argument of Proposition B.1 of Dette et al. (2019).

Let \tilde{T}_n denote the bootstrap statistic (5.2) generated in one iteration. Recall the definitions of $U(t)$, s_1 , s_2 in Theorem 4.1. Theorem 5.1 gives the limiting distributions of bootstrap statistic \tilde{T}_n .

Theorem 5.1 (Bootstrap under null). *Assume the conditions 4.1, 4.2, 4.3, 4.4 and 5.1 hold, $nb_n^{7/2}/(\log n)^4 \rightarrow \infty$, $nb_n^6 \rightarrow 0$, $\eta_n \rightarrow 0$, $n\eta_n^2 \rightarrow \infty$. Then, under the null hypothesis, we have*

(i) *if Assumption 4.5 holds, then $\tilde{T}_n \Rightarrow \int_0^1 U^2(t) dt$.*

(ii) *if Assumption 4.5 doesn't hold, then $s_1^{-1}(\tilde{T}_n/b_n - s_2) \Rightarrow \chi_1^2$.*

Combining with Theorem 4.1, Theorem 5.1 indicates that the bootstrap test Algorithm 5.1 is asymptotically of level α no matter whether Assumption 4.5 is satisfied. We proceed to investigate the behavior of the bootstrap statistic \tilde{T}_n under fixed and local alternatives. Let $\tilde{U}_d(t)$, $\check{U}(t)$ be zero mean continuous Gaussian processes, with the covariance structures defined in the same way as that of $U(t)$ in (4.4), where $\Sigma(t)$ is replaced by $\Sigma_d(t)$ and $\check{\Sigma}(t)$ in Assumption 5.1, respectively. Let $\sigma_{Hd}^2(t) := (\Sigma_d(t))_{(1,1)}$, $\check{\sigma}_H^2(t) := (\check{\Sigma}(t))_{(1,1)}$.

Theorem 5.2 (Bootstrap under alternatives). *Under the conditions of Theorem 5.1,*

(i) *Suppose $\lambda(\Sigma_d^{1/2}(t) \mathbf{M}^{-1}(t) \boldsymbol{\mu}_W(t) \neq (\sigma_{Hd}(t), 0, \dots, 0)) > 0$ under the fixed alternatives $d > 0$. Then, we have*

$$m^{-2d} \tilde{T}_n \Rightarrow \int_0^1 \tilde{U}_d^2(t) dt,$$

where $\tilde{U}_d(t)$ is a zero-mean continuous Gaussian process with covariance function

$$\begin{aligned} \mathbb{E}(\tilde{U}_d(r) \tilde{U}_d(s)) &= \int_0^{r \wedge s} \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \Sigma_d(t) \mathbf{M}^{-1}(t) \boldsymbol{\mu}_W(t) dt \\ &\quad - 2 \int_0^{r \wedge s} \{\boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \Sigma_d^{1/2}(t)\}_1 \sigma_{Hd}(t) dt + \int_0^{r \wedge s} \sigma_{Hd}^2(t) dt, \quad r, s \in [0, 1]. \end{aligned}$$

(ii) Suppose $\lambda(\check{\Sigma}^{1/2}(t)\mathbf{M}^{-1}(t)\boldsymbol{\mu}_W(t) \neq (\check{\sigma}_H(t), 0, \dots, 0)) > 0$. For the local alternatives $d_n = c/\log n$ with some positive constant c , we have

$$\tilde{T}_n \Rightarrow \int_0^1 \check{U}^2(t)dt,$$

where $\check{U}(t)$ is a zero-mean continuous Gaussian process with covariance function

$$\begin{aligned} \mathbb{E}(\check{U}(r)\check{U}(s)) &= \int_0^{r \wedge s} \boldsymbol{\mu}_W^\top(t)\mathbf{M}^{-1}(t)\check{\Sigma}(t)\mathbf{M}^{-1}(t)\boldsymbol{\mu}_W(t)dt \\ &\quad - 2 \int_0^{r \wedge s} \{\boldsymbol{\mu}_W^\top(t)\mathbf{M}^{-1}(t)\check{\Sigma}^{1/2}(t)\}_1 \check{\sigma}_H(t)dt + \int_0^{r \wedge s} \check{\sigma}_H^2(t)dt, \quad r, s \in [0, 1]. \end{aligned}$$

Theorem 5.2 (i) gives the limiting distribution of \tilde{T}_n/m^{2d} , which means the critical values generated by **Algorithm 5.1** for a level α test diverges at the rate of m^{2d} . Thus, the bootstrap-assisted test is consistent since **Theorem 4.3** demonstrates that the KPSS-type test statistic T_n diverges at the rate n^{2d} which is much faster than m^{2d} . Further together with **Theorem 4.1**, **Theorem 5.2** (i) shows that the bootstrap test **Algorithm 5.1** is asymptotically correct. Notice that the condition $\lambda(\Sigma_d(t)\mathbf{M}^{-1}(t)\boldsymbol{\mu}_W(t) \neq (\sigma_{Hd}(t), 0, \dots, 0)) > 0$ prevents the degeneracy of bootstrap statistics. If it is violated, $\tilde{T}_n = o(m^{2d})$ which will yield higher power than when the condition is fulfilled.

On the other hand, **Theorem 5.2** (ii) and **Theorem 4.4** indicate that the bootstrap test **Algorithm 5.1** is able to detect the local alternatives at the rate of $\log^{-1}n$. Observe that under **Assumption 5.1** (iii) as $c \rightarrow 0$, $|\check{\Sigma}(t) - \Sigma(t)| \rightarrow 0$ and the covariance structure of $\check{U}(t)$ will converge to the covariance structure of $U(t)$. Therefore, the bootstrap test **Algorithm 5.1** has no power when $d_n = o(\log^{-1}n)$, indicating that the proposed test has the exact local power of $O(\log^{-1}n)$ under the condition of **Theorem 5.2**. For stationary time series with unknown constant mean, **Shao and Wu (2007)** has also proved that the KPSS test for long memory has the exact local power $O(\log^{-1}n)$.

An important scenario that T_n degenerates under both null and alternatives is the time-varying trend model (4.5). For this case we provide the bootstrap-assisted KPSS test in **Algorithm G.1** of the online supplement, which is the $p = 1$ version of **Algorithm 5.1** with $\hat{\mathbf{M}}(t) = 1$, and $\hat{\Sigma}(t) = \hat{\sigma}_H^2(t)$ using the difference statistic proposed by **Dette and Wu (2019)**. **Theorem 5.1** (ii) ensures that the level α critical value generated by **Algorithm G.1** converges to the α_{th} quantile of the test statistic T_n under the null hypothesis. Under the alternative hypothesis, the following proposition investigates the power of the test implemented via **Algorithm G.1**.

Proposition 5.1. *Let \tilde{T}_n denote the KPSS-type bootstrap statistic generated from **Algorithm G.1**. Under **Assumptions 3.1, 4.1, 4.2, 5.1**, and the bandwidth conditions $nb_n^4/(\log n)^2 \rightarrow \infty, b_n \rightarrow 0, nb_n/m \rightarrow \infty$, we have the following results:*

(i) *Under the fixed alternatives $d > 0$, $\lim_{n \rightarrow \infty} P(T_n > \tilde{T}_n) = 1$.*

(ii) *Suppose $m = \lfloor n^{\alpha_1} \rfloor$, $nb_n = n^\beta$ for some $\alpha_1, \beta \in (0, 1)$. Then under local alternatives with $d = d_n = c/\log n$ for a sufficiently large constant c , $\lim_{n \rightarrow \infty} P(T_n > \tilde{T}_n) = 1$.*

In addition to the KPSS test, **Algorithm G.1** also provides the bootstrap-assisted V/S, R/S and K/S tests for the time-varying trend model (4.5). Similar conclusions as given in **Proposition 5.1** hold for the power of these tests.

Remark 5.1. *For $p \geq 2$, T_n degenerates under the fixed alternatives if and only if all the stochastic covariates are of mean zero (i.e., $\mu_W^{(-1)}(t) = 0$), see **Section 4.2.2**. Under this condition, we can show that T_n/b_n diverges at the rate of $(nb_n)^{2d}$. Combining with **Theorem 5.2** (i), our bootstrap tests **Algorithm 5.1** and **Algorithm G.2** are consistent if $b_n(nb_n/m)^{2d} \rightarrow \infty$. If $\sum_{i \in \mathbb{Z}} \text{Cov}(H(t, \mathcal{F}_0), H(t, \mathcal{F}_i)W^{(-1)}(t, \mathcal{F}_i)) =$*

0, T_n degenerates under H_0 and H_A if and only if $\mu_W^{(-1)}(t) = 0$. In this case, one could implement [Algorithm 5.1](#) and [Algorithm G.2](#) for R/S, V/S, K/S-type tests via modifying the difference-based long-run covariance estimator: set $\hat{\Sigma}(t)_{(1,l)} = 0$ and $\hat{\Sigma}(t)_{(l,1)} = 0$ for $l = 2, \dots, p$. Then, by a further investigation of the proof to [Theorem 5.2](#), \tilde{T}_n/b_n is $O_{\mathbb{P}}(m^{2d})$ and thus the tests are consistent when $m/(nb_n) \rightarrow 0$. In addition, similar to (5.1), we can also show that the local power of the tests with the modified estimator is of order $O(\log^{-1} n)$.

6 Finite sample performance

In the following simulation studies and data analysis, we examine the size and power performance of the bootstrap-assisted KPSS and related tests [Algorithm 5.1](#), [Algorithm G.1](#) and [Algorithm G.2](#) in the online supplement. The number of bootstrap samples is $B = 2000$ and the number of replications is 1000. The parameters $\mathbf{M}(t)$, $\Sigma(t)$, $\sigma_H^2(t)$ are estimated by $\hat{\mathbf{M}}(t)$, $\hat{\Sigma}(t)$, $\hat{\sigma}_H^2(t)$ in [Section 5](#), with all the smoothing parameters selected by the methods advocated in [Appendix C](#) in the online supplement. Let

$$\mathcal{F}_j = (\dots, \zeta_{j-1}, \zeta_j), \quad \mathcal{G}_j = (\dots, \varepsilon_{j-1}, \varepsilon_j), \quad j = -\infty, \dots, n,$$

where $(\varepsilon_l)_{l \in \mathbb{Z}}, (\zeta_l)_{l \in \mathbb{Z}}$ are *i.i.d.* $N(0, 1)$. We consider the following time-varying coefficient model,

$$y_{i,n} = \beta_1(t_i) + \beta_2(t_i)x_{i,n} + e_{i,n}, \quad i = 1, \dots, n,$$

where $\beta_1(t) = 4 \sin(\pi t)$, $\beta_2(t) = 4 \exp\{-2(t - 0.5)^2\}$, $x_{i,n} = W(t_i, \mathcal{F}_i)$, and $u_{j,n} = H(t_j, \mathcal{F}_j, \mathcal{G}_j)$. First, we consider the following independent model:

- (i) Let $W(t, \mathcal{F}_i) = (0.25 + 0.25 \cos(2\pi t))W(t, \mathcal{F}_{i-1}) + 0.25\zeta_i + (t - 0.5)^2$, $H(t, \mathcal{G}_i) = (0.35 - 0.4(t - 0.5)^2)H(t, \mathcal{G}_{i-1}) + 0.8\varepsilon_i$.

Second, we consider the following heteroscedastic model:

$$H(t, \mathcal{F}_i, \mathcal{G}_i) = B(t, \mathcal{G}_i) \sqrt{1 + W^2(t, \mathcal{F}_i)},$$

where $W(t, \mathcal{F}_i) = (0.1 + 0.1 \cos(2\pi t))W(t, \mathcal{F}_{i-1}) + 0.2\zeta_i + 0.7(t - 0.5)^2$, and $B(t, \mathcal{G}_i)$ is as considered in the following linear and nonlinear scenarios.

- (ii.1) Linear errors: $B(t, \mathcal{G}_i) = (0.3 - 0.4(t - 0.5)^2)B(t, \mathcal{G}_{i-1}) + 0.8\varepsilon_i$.
- (ii.2) Nonlinear errors: $B(t, \mathcal{G}_i) = (0.15 - 0.4(t - 0.5)^2)B(t, \mathcal{G}_{i-1}) + 0.8G(t, \mathcal{G}_i)$, $G(t, \mathcal{G}_i) = \varepsilon_i \sigma_i(t)$, where $\sigma_i^2(t) = 0.9 + 0.1 \cos(\pi/3 + 2\pi t) + (0.1 + 0.2t)G^2(t, \mathcal{G}_{i-1}) + (0.1 + 0.2t)\sigma_{i-1}^2(t)$.

Observe that models (ii.1) and (ii.2) are heteroscedastic models with locally stationary AR(1) and locally stationary GARCH(1,1) errors, respectively. [Table 6.1](#) summarizes the performance of our proposed bootstrap-assisted KPSS, R/S, V/S and K/S-type tests for long memory in models (ii.1) and (ii.2) with different b'_n s. We relegate the simulated sizes of model (i) with different b'_n s to [Table D.2](#) of the online supplement. The empirical sizes of all the four tests are close to their nominal levels and are quite stable when b_n changes within a reasonably wide range. Also, [Table D.1](#) in the online supplement reports the simulated Type I error of the proposed tests with respect to increasing sample sizes. As shown in [Table D.1](#), our procedures for smoothing parameter selection including GCV and MV selection as well as the difference-based long-run variance estimator work very well in the sense that the simulated sizes of all four tests are quite close to their nominal levels in different sample sizes.

Recall the long-memory error process of (1.2), which can be written as $e_{i,n}^{(d)} = (1 - \mathcal{B})^{-d}u_{i,n}$ where \mathcal{B} is the lag operator. [Figure 6.1](#) displays the power performance of the KPSS and related tests for model (ii.1) with nominal level 0.1. The left panel reports simulated rejection rates of KPSS, R/S, V/S, and K/S-type tests as the long memory parameter d increases from 0 to 0.5 with sample size 1500. The power of all four KPSS and related tests increases to 1 as d approaches to 1/2, while the V/S-type test remains

(ii.1)									(ii.2)								
	KPSS		R/S		V/S		K/S			KPSS		R/S		V/S		K/S	
b_n	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	
0.15	4.7	8.6	5.5	9.8	6.4	10.0	4.8	10.8	4.1	9.2	4.8	9.4	5.0	9.7	3.9	8.6	
0.175	5.5	10.4	5.5	11.1	4.8	9.6	5.1	9.3	5.1	8.9	5.0	9.3	5.4	10.4	5.3	9.9	
0.2	5.2	9.9	5.9	10.6	6.1	9.3	4.8	8.4	5.4	9.8	4.9	10.9	5.4	10.7	4.6	9.5	
0.225	5.4	10.2	4.8	9.9	4.8	9.5	5.0	8.9	5.0	9.5	5.7	10.3	4.4	8.9	4.7	10.4	

Table 6.1: Simulated sizes (in %) of KPSS, R/S, V/S and K/S-type tests for model (ii.1) and (ii.2) with the sample size 1000, m and τ_n determined by MV selection.




Figure 6.1: Simulated powers of KPSS and related tests with nominal level 0.1. Left: $n = 1500$ and d increases from 0 to 0.5; Right: $d = 0.4$ and n increases from 200 to 2500.

the most powerful among all tests. The right panel depicts the power performance of the four tests as the sample size grows from 200 to 2500 when $d = 0.4$. The figure shows that the power of each test increases to 1 as the sample size grows, and the V/S-type test performs the best among the four tests whenever the sample size is larger than 400. The power performance of models (i) and (ii.2) are shown in Figure D.1 and Figure D.2 of the online supplement where the V/S-type test achieves the best power under most alternatives and sample sizes considered.

6.1 Analysis of Hong Kong hospital data

Hong Kong circulatory and respiratory data contains daily measurements of pollutants and daily hospital admissions in Hong Kong between January 1st, 1994 and December 31st, 1995. The dataset has been studied by Fan and Zhang (2000), Zhou and Wu (2010), and Wu and Zhou (2018a) among others. They investigated the relationship between the levels of pollutants and the total number of hospital admissions of circulation and respiration, see Figure F.1 in Appendix F of the supplement for the observed series. Under the assumption of *i.i.d.* observations, Fan and Zhang (2000) claimed that sulphur dioxide (SO_2) is not significant. Using the non-stationary model, Zhou and Wu (2010) found all three pollutants (SO_2 , nitrogen dioxide (NO_2) and dust) are significant. In both cases, they assumed the observations were SRD. We shall examine this assumption via the KPSS and related tests. Consider the following time-varying

coefficient model,

$$y_{i,n} = \beta_1(t_i) + \sum_{p=2}^4 \beta_p(t_i) x_{i,p,n} + \varepsilon_i, \quad i = 1, \dots, n, \quad (6.1)$$

where $(y_{i,n})$ is the series of daily total number of hospital admissions of circulation and respiration and $(x_{i,p,n})$, $p = 2, 3, 4$, represent the series of daily levels (in micrograms per cubic meter) of SO_2 , NO_2 and dust, respectively. The sample size is $n = 2 \times 365 = 730$. We first investigate whether the covariates $(x_{i,p,n})$, $p = 2, 3, 4$ are SRD series.

We select the smoothing parameters b_n , η_n , m and τ_n through the methods provided in Appendix C in the online supplement and summarize the selected parameters in Table F.1 in Appendix F of the online supplement. We test for long memory in the three pollutants series via KPSS and related tests, presenting the p -values in Table 6.2. For each pollutant series we fail to reject it is SRD at the significance level 0.05.

	KPSS	R/S	V/S	K/S		KPSS	R/S	V/S	K/S
SO_2	0.358	0.076	0.087	0.543	Dust	0.668	0.103	0.124	0.663
NO_2	0.476	0.818	0.232	0.748	(6.1)	0.461	0.633	0.549	0.649

Table 6.2: The p -values for SO_2 , NO_2 , dust and daily total number of hospital admissions modeled by (6.1).

To test for long memory in the daily hospital admissions, we consider two approaches. The first approach is to model the hospital admissions by the time-varying trend model (4.5) and implement the KPSS and related tests, i.e., the Algorithm G.1 in the online supplement. The KPSS test yields p -values 0.004, and other tests yield p -values smaller than 2×10^{-4} . All four tests reject the null hypothesis of short memory at the significance of 0.05. The second approach is to model the hospital admissions via model (6.1) taking into account three pollutants (SO_2 , NO_2 and dust) and apply the KPSS and related tests, i.e., the Algorithm 5.1 and Algorithm G.2 in the online supplement. The large p -values in the right panel of Table 6.2 show that the KPSS and related tests fail to reject that the total number of hospital admissions is SRD at the significance level 0.05. Although both of the two approaches are asymptotically correct, the misspecification of regression models tends to cause ‘spurious long memory’ in finite samples. Therefore, our results conclude that the SRD assumption for (6.1) adopted by Fan and Zhang (2000), Zhou and Wu (2010), Wu and Zhou (2018a) and many others is reasonable.

7 Conclusion and future work

This paper develops bootstrap-assisted KPSS, R/S, K/S, and V/S-type nonparametric tests to detect long memory in time-varying coefficient linear models where the covariates and errors are allowed to be locally stationary and heteroscedastic. Under the null hypothesis, the fixed and local alternatives, we derive the limiting distributions of those test statistics and bootstrap statistics. In particular, we identify the conditions under which the test statistics degenerate. Although such conditions cannot be directly identified from real data in practice, our proposed bootstrap-assisted KPSS and related tests are always asymptotically correct regardless of such conditions. We also establish the theory of Gaussian approximation to the partial sum process of the product of non-stationary SRD and LRD time series, which is of separate interest and useful for a large class of problems in the analysis of (time-varying) linear models with LRD errors and SRD covariates.

A comprehensive Monte Carlo study supports that our proposed KPSS and related tests have good size and power performance in finite samples and are robust to the choices of smoothing parameters. The tests are applied to Hong Kong circulatory and respiratory data, recognizing ‘spurious long memory’ due

to the misspecification in the conditional mean. Therefore, our proposed methods can be employed as regression diagnostics. In Appendix E of the online supplement, we also apply the proposed tests to the COVID-19 data and identify that the time series of log cumulative confirmed cases of Japan and Ireland are LRD, while the time series of log cumulative confirmed deaths of Japan and Ireland are both SRD.

Recent studies have considered LRD models with time-varying long-memory parameter $d(t)$, see for instance [Dette et al. \(2017\)](#) and [Ferreira et al. \(2018\)](#). Extra simulations in Appendix D.4 of the online supplement evidence that our proposed testing procedures are still consistent against the alternatives with time-varying non-negative $d(t)$, i.e., $d(t) > 0$ for t in a sub-interval of $[0, 1]$. The derivation of the theoretical behavior of the test statistics and the bootstrap procedure with time-varying $d(t)$ is left for rewarding future work. The extension of the proposed tests to $d \geq 1/2$ or $d < 0$ for the null hypothesis (see also [Wu and Shao \(2006\)](#), [Duffy and Kaspas \(2021\)](#)) is also challenging and meaningful.

A Proof of Theorem 4.2

This subsection provides the proof of [Theorem 4.2](#). We first present the following propositions which are needed for the proof. The [Proposition A.1](#) approximates the partial sum of the product series by the partial sum of a LRD process weighted by the *expectation* of a SRD process. In [Proposition A.2](#), we establish the Gaussian approximation scheme for the vector partial sum process of $(\boldsymbol{\mu}_W(t_i)e_{i,n}^{(d)})$. [Proposition A.3](#) is a non-trivial extension of [Proposition A.1](#) under the local alternatives. The proofs of [Proposition A.2](#) and [Proposition A.3](#) are postponed to Appendix H.4 of the online supplements.

Proposition A.1. *Under Assumptions 3.1 and 4.4, we have*

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_{i,n} e_{i,n}^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) e_{i,n}^{(d)} \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d).$$

In [Proposition A.1](#), the rate $O_{\mathbb{P}}(\sqrt{n}(\log n)^d)$ is due to the complicated dependence between the covariate process and the driving shocks $(u_{i,n})$. When $(\mathbf{x}_{i,n})$ and $(e_{i,n}^{(d)})$ are independent, the bound can be further sharpened to $O_{\mathbb{P}}(\sqrt{n})$. The discrepancy manifests the subtle effect of heteroscedasticity under the fixed alternatives.

Remark A.1. *The approximation result in [Proposition A.1](#) is frequently considered in the context of regression with LRD errors. With arguments given in the proof of [Proposition A.1](#), we can show that for any deterministic function $v(\cdot) : \mathbb{R}^p \rightarrow \mathbb{R}$, the partial sum process $\sum_{i=\lfloor nb_n \rfloor + 1}^r v(\mathbf{x}_{i,n}) e_{i,n}^{(d)}$ can be approximated by $\sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(v(\mathbf{x}_{i,n})) e_{i,n}^{(d)}$, i.e.,*

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r v(\mathbf{x}_{i,n}) e_{i,n}^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(v(\mathbf{x}_{i,n})) e_{i,n}^{(d)} \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d).$$

Similar to [Proposition A.1](#), if $\mathbf{x}_{i,n}$ is independent of $e_{i,n}$, the approximation error will be reduced to $O_{\mathbb{P}}(\sqrt{n})$. As a consequence, our result is also in line with the result in Section 7.2.3 of [Beran et al. \(2013\)](#) for the quantity $\sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} v(\mathbf{x}_{i,n}) e_{i,n}^{(d)}$. Furthermore, our proof extends that of Proposition 1 and 2 in [Kulik and Wichelhaus \(2012\)](#) in a non-trivial way, since they focus on the partial sum assuming i.i.d. covariates $\{\mathbf{x}_{i,n}\}$ and i.i.d. errors $\{e_{i,n}\}$. [Kulik and Wichelhaus \(2012\)](#) utilized their Propositions 1 and 2 to estimate the conditional variance in the heteroscedastic model (1.1).

Proposition A.2. Under Assumptions 3.1 and 4.4, on a possibly richer probability space, there exists a sequence of i.i.d.. standard normal $\{v_i\}_{i \in \mathbb{Z}}$ and $\mathbf{R}_{k,n} = \sum_{j=0}^{\infty} \boldsymbol{\mu}_W(t_k) \psi_j \sigma_H(t_{k-j}) v_{k-j}$, $1 \leq k \leq n$, such that

$$\max_{1 \leq s \leq n} \left| \sum_{k=1}^s \left(\boldsymbol{\mu}_W(t_k) e_{k,n}^{(d)} - \mathbf{R}_{k,n} \right) \right| = O_p \left(n^{1+\alpha_0(d-1/2)} \right).$$

where $\alpha_0 \in (1, 4/3)$ and therefore $n^{1+\alpha_0(d-1/2)} = o(n^{d+1/2})$.

Proposition A.3. Under Assumptions 3.1 and 4.4, we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_{i,n} e_{i,n}^{(d_n)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \left\{ \mathbf{x}_{i,n} e_{i,n} + \boldsymbol{\mu}_W(t_i) (e_{i,n}^{(d_n)} - e_{i,n}) \right\} \right| = o_{\mathbb{P}}(\sqrt{n}).$$

In the following proofs for the sake of simplicity, we will omit the index n in $e_{i,n}$, $\mathbf{x}_{i,n}$, $y_{i,n}$, $u_{i,n}$ and use ψ_j to represent $\psi_j(d)$ when we discuss the fixed alternatives and to represent $\psi_j(d_n)$ for the theory of the local alternatives.

A.1 Proof of Theorem 4.2

Result (i) follows from Proposition A.2 and Proposition A.1. With regard to (ii), observe that

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \left\{ \mathbf{x}_i e_i + \boldsymbol{\mu}_W(t_i) (e_i^{(d_n)} - e_i) \right\} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i e_i + \sum_{j=1}^{\infty} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \psi_j u_{i-j}.$$

The proof follows from Proposition A.2, Proposition A.3 and (H.16) of the online supplement. \square

A.2 Proof of Proposition A.1

Let $\tilde{m} = M \log n$, $\tilde{e}_{i,\tilde{m}}^{(d)} = \sum_{j=\tilde{m}+1}^{\infty} \psi_j u_{i-j}$, $\tilde{\mathbf{x}}_{i,\tilde{m}} = \mathbb{E}(\mathbf{x}_i | \varepsilon_i, \dots, \varepsilon_{i-\tilde{m}})$. Firstly, we can approximate $\sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i e_i^{(d)}$ by $\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} e_i^{(d)}$ in that by (H.8) of the online supplement,

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \sum_{i=\lfloor nb_n \rfloor + 1}^r (\mathbf{x}_i - \tilde{\mathbf{x}}_{i,\tilde{m}}) e_i^{(d)} \right\| \leq \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \|\mathbf{x}_i - \tilde{\mathbf{x}}_{i,\tilde{m}}\|_4 \|e_i^{(d)}\|_4 = O(n \chi^{\tilde{m}}). \quad (\text{A.1})$$

For every fixed $j = 1, 2, \dots, \tilde{m}$, notice that $(\tilde{\mathbf{x}}_{i,\tilde{m}} u_{i-j})_{i=1}^n$ is a SRD sequence similar to $(\mathbf{x}_i u_i)_{i=1}^n$. Therefore, we have

$$\begin{aligned} \left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} (e_i^{(d)} - \tilde{e}_{i,\tilde{m}}^{(d)}) \right\| &\leq \sum_{j=0}^{\tilde{m}} \psi_j \left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} u_{i-j} \right\| \\ &= O(\sqrt{n} \tilde{m}^d). \end{aligned} \quad (\text{A.2})$$

For $k \in \mathbb{Z}$, we define the projection operator as $\mathcal{P}_k \cdot = \mathbb{E}(\cdot | \mathcal{F}_k) - \mathbb{E}(\cdot | \mathcal{F}_{k-1})$. Then, we have the following decomposition

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} = \sum_{l=0}^{\tilde{m}} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} \right) + \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} | \mathcal{F}_{i-\tilde{m}-1}) := T_1 + T_2. \quad (\text{A.3})$$

We proceed to show that the T_2 is the leading term. Applying Doob's inequality to the martingale $\sum_{i=\lfloor nb_n \rfloor + 1}^r \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} \right)$, we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |T_1| \right\| \leq 2 \sum_{l=0}^{\tilde{m}} \left\| \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} \right) \right\|.$$

Let $\tilde{\mathbf{x}}_{i,\tilde{m}}^{(i-l)}$, $\tilde{e}_{i,\tilde{m}}^{(d),(i-l)}$ denote the random variables replacing ε_{i-l} in $\tilde{\mathbf{x}}_{i,\tilde{m}}$ and $\tilde{e}_{i,\tilde{m}}^{(d)}$ with its *i.i.d.* copy. We have $\tilde{e}_{i,\tilde{m}}^{(d),(i-l)} = \tilde{e}_{i,\tilde{m}}^{(d)}$ for $l \leq \tilde{m}$, following from the definition of $\tilde{e}_{i,\tilde{m}}^{(d)}$. By Jensen's inequality, for $l \leq \tilde{m}$,

$$\begin{aligned} \left\| \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} \right) \right\|^2 &= \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left\| \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d)} \right) \right\|^2 \\ &\leq \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\left\| \tilde{\mathbf{x}}_{i,\tilde{m}} - \tilde{\mathbf{x}}_{i,\tilde{m}}^{(i-l)} \right\|_4 \left\| \tilde{e}_{i,\tilde{m}}^{(d)} \right\|_4 + \left\| \tilde{\mathbf{x}}_{i,\tilde{m}}^{(i-l)} \right\|_4 \left\| \tilde{e}_{i,\tilde{m}}^{(d)} - \tilde{e}_{i,\tilde{m}}^{(d),(i-l)} \right\|_4 \right)^2 \\ &= \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\left\| \tilde{\mathbf{x}}_{i,\tilde{m}} - \tilde{\mathbf{x}}_{i,\tilde{m}}^{(i-l)} \right\|_4 \left\| \tilde{e}_{i,\tilde{m}}^{(d)} \right\|_4 \right)^2 = O(n\chi^{2l}). \end{aligned}$$

Hence, we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |T_1| \right\| = O(\sqrt{n}). \quad (\text{A.4})$$

Since $\tilde{e}_{i,\tilde{m}}^{(d)}$ is $\mathcal{F}_{i-\tilde{m}-1}$ measurable and $\tilde{\mathbf{x}}_{i,\tilde{m}}$ is independent of $\mathcal{F}_{i-\tilde{m}-1}$, we have

$$T_2 = \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(\tilde{\mathbf{x}}_{i,\tilde{m}} | \mathcal{F}_{i-\tilde{m}-1}) \tilde{e}_{i,\tilde{m}}^{(d)} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,\tilde{m}}^{(d)}. \quad (\text{A.5})$$

Therefore, combining the results from (A.1), (A.2), (A.3), (A.4) and (A.5), we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i e_i^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,\tilde{m}}^{(d)} \right| \right\| = O(\sqrt{n} \tilde{m}^d + n\chi^{\tilde{m}} + \sqrt{n}) = O(\sqrt{n} \tilde{m}^d).$$

Finally, since $\mathcal{P}_k \cdot = \mathbb{E}(\cdot | \mathcal{F}_k) - \mathbb{E}(\cdot | \mathcal{F}_{k-1})$ and $e_i^{(d)} - \tilde{e}_{i,\tilde{m}}^{(d)} = \sum_{j=0}^{\tilde{m}} \psi_j u_{i-j}$, it follows from Doob's maximal inequality and Burkholder's inequality that

$$\begin{aligned} &\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) (e_i^{(d)} - \tilde{e}_{i,\tilde{m}}^{(d)}) \right| \right\| \\ &\leq C_1 \sum_{l=0}^{\infty} \sum_{j=0}^{\tilde{m}} \psi_j \left\| \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \boldsymbol{\mu}_W(t_i) \mathcal{P}_{i-l} u_{i-j} \right\| \\ &\leq C_2 \sqrt{n} \left(\sum_{l=0}^{\tilde{m}} \sum_{j=0}^l \psi_j \delta_2(H, l-j, (-\infty, 1]) + \sum_{l=\tilde{m}+1}^{\infty} \sum_{j=0}^{\tilde{m}} \psi_j \delta_2(H, l-j, (-\infty, 1]) \right) \\ &= O(\sqrt{n} \tilde{m}^d), \end{aligned} \quad (\text{A.6})$$

where C_1, C_2 are sufficiently large constants. The last inequality of (A.6) follows in that for $k < 0$, $\delta_2(H, k, (-\infty, 1]) = 0$, and the big O follows from Lemma 3.2 in Kokoszka and Taqqu (1995). \square

Acknowledgement

Weichi Wu is the corresponding author and gratefully acknowledges NSFC Young Program (no. 11901337) of China.

Supplement to “Detecting long-range dependence for time-varying linear models”

We organize the supplementary material as follows: [Appendix B](#) contains a literature review of KPSS and related test. [Appendix C](#) provides the implementation details including the selection procedures for the smoothing parameters m, b_n, τ_n and η_n in the bootstrap tests. [Appendix D](#) reports additional simulations of KPSS and related tests. [Appendix E](#) analyses the COVID-19 dataset. [Appendix F](#) displays some details in analyzing Hong Kong circulatory and respiratory data. [Appendix G](#) provides algorithms of KPSS and related tests under the time-varying trend model and R/S, V/S and K/S-type tests under the time-varying coefficient model. [Appendix H](#) gives the limiting distributions of R/S, V/S and K/S-type statistics under [Assumption 4.5](#) and the proofs of the results in [Sections 3 and 4](#) of the main article. In [Appendix I](#), we justify the proposed bootstrap procedures and offer the proofs of the results in [Section 5](#) of the main article.

Recall filtration $\mathcal{F}_i = (\varepsilon_{-\infty}, \dots, \varepsilon_i)$ for *i.i.d.* random variables $(\varepsilon_i)_{i \in \mathbb{Z}}$ and the projection operator $\mathcal{P}_k \cdot = \mathbb{E}(\cdot | \mathcal{F}_k) - \mathbb{E}(\cdot | \mathcal{F}_{k-1})$ for $k \in \mathbb{Z}$. Recall that $e_{i,n}^{(d)} = \sum_{j=0}^{\infty} \psi_j(d) u_{i-j,n}$, $e_{i,n}^{(d_n)} = \sum_{j=0}^{\infty} \psi_j(d_n) u_{i-j,n}$. For the sake of simplicity, we use ψ_j to represent $\psi_j(d)$ when we discuss the fixed alternatives and $\psi_j(d_n)$ for the theory of the local alternatives. Recall $t_i = i/n$, and that $K^*(x)$ denotes the jackknife equivalent kernel $2\sqrt{2}K(\sqrt{2}x) - K(x)$. Let “ \Rightarrow ” denote weak convergence, and “ \rightsquigarrow ” denote the convergence of a process. Let $0 \times \infty = 0$, $a_n \sim b_n$ denote $\lim_{n \rightarrow \infty} a_n/b_n = 1$ for real sequences a_n and b_n . For a random variable X and a distribution G , $X \sim G$ is denoted by X follows the distribution G . Let $D[0, 1]$ be the space of real functions on $[0, 1]$ that are right-continuous and have left-hand limits (also named càdlàg functions). In the following proofs, we will omit the index n in $e_{i,n}, \mathbf{x}_{i,n}, y_{i,n}, u_{i,n}$ for simplicity.

B KPSS and related tests

The first test statistic is the KPSS-type statistic T_n defined by

$$T_n = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \sum_{r=\lfloor nb_n \rfloor+1}^{n-\lfloor nb_n \rfloor} \left(\tilde{S}_{r,n} \right)^2.$$

The KPSS test was first introduced by [Kwiatkowski et al. \(1992\)](#) to test for the unit root in level and trend stationary series, complementary to the ADF test. Besides the unit root problem, KPSS-type statistics have been widely and successfully applied to many important hypothesis testing problems, including testing for long memory against the null hypothesis of short memory, see for instance [Lee and Schmidt \(1996\)](#). The same statistic was also used for detecting structural changes (see [MacNeill \(1974\)](#) among others) and examining random walk components in functional time series (see [Kokoszka and Young \(2016\)](#)). The exhaustive account of the applications of KPSS-type tests is almost impossible and we have only listed a small fraction here.

The second test statistic Q_n is the R/S-type statistic defined by

$$Q_n = \max_{\lfloor nb_n \rfloor+1 \leq k \leq n-\lfloor nb_n \rfloor} \tilde{S}_{k,n} - \min_{\lfloor nb_n \rfloor+1 \leq k \leq n-\lfloor nb_n \rfloor} \tilde{S}_{k,n}. \quad (\text{B.1})$$

The R/S test was first introduced by [Hurst \(1951\)](#). [Lo \(1989\)](#) proposed a modified R/S test for long memory, which is robust to the short-range dependence of strictly stationary errors under null. Lo’s test [Lo \(1989\)](#) has been widely applied in finance, see [Cheung and Lai \(1993\)](#) and many others.

The third test statistic is the V/S-type statistic M_n defined by

$$M_n = \frac{1}{n(n-2\lfloor nb_n \rfloor)} \left\{ \sum_{k=\lfloor nb_n \rfloor+1}^{n-\lfloor nb_n \rfloor} \tilde{S}_k^2 - \frac{1}{n-2\lfloor nb_n \rfloor} \left(\sum_{k=\lfloor nb_n \rfloor+1}^{n-\lfloor nb_n \rfloor} \tilde{S}_k \right)^2 \right\}. \quad (\text{B.2})$$

The V/S test is proposed by [Giraitis et al. \(2003\)](#), where the authors found that the V/S test achieved better size and power performance than R/S and KPSS tests when applied to certain financial data.

The fourth test statistic is the K/S-type statistic G_n defined by

$$G_n = \max_{\lfloor nb_n \rfloor+1 \leq k \leq n-\lfloor nb_n \rfloor} |\tilde{S}_k|. \quad (\text{B.3})$$

[Lima and Xiao \(2004\)](#) used the K/S statistic to test for long-range dependence and argued that its behavior under the alternative of long memory hypothesis was similar to that of R/S test.

C Implementation details

In this section, we discuss the selection of proper smoothing parameters m, b_n, τ_n and η_n for the implementation of the bootstrap tests [Algorithm 5.1](#), [Algorithm G.1](#) and [Algorithm G.2](#). The criterion for choosing parameters for [Algorithm G.1](#) is to let $p = 1$ in the following schemes.

To select b_n , we adopt the Generalized Cross Validation (GCV) proposed by [Craven and Wahba \(1978\)](#). For the estimation of $\beta(\cdot)$, we can write $\hat{\mathbf{Y}}(b) = \mathbf{Q}(b)\mathbf{Y}$ for some square matrix \mathbf{Q} , where $\mathbf{Y} = (y_{1,n}, \dots, y_{n,n})^\top$, and $\hat{\mathbf{Y}}(b) = (\hat{y}_{1,n}, \dots, \hat{y}_{n,n})^\top$ is the estimated value of \mathbf{Y} via the bandwidth b , i.e., $\hat{y}_{i,n} = \mathbf{x}_{i,n}^\top \hat{\beta}(t_i)$. Then we select \hat{b}_n by

$$\hat{b}_n = \arg \min_{b \in [b_L^*, b_U^*]} \{ \text{GCV}(b) \}, \quad \text{GCV}(b) = \frac{n^{-1} |\mathbf{Y} - \hat{\mathbf{Y}}|^2}{[1 - \text{tr}\{\mathbf{Q}(b)\}/n]^2},$$

where the selection range $[b_L^*, b_U^*]$ are chosen as follows. The theoretical optimum bandwidth b_n for the local linear estimation [\(4.1\)](#), as discussed in [Zhou and Wu \(2010\)](#), is

$$b_n^* = \left[\frac{\phi_0 \int_0^1 \text{tr}\{\boldsymbol{\Sigma}(t)\} dt}{\mu_2^2 \int_0^1 |\boldsymbol{\beta}''(t)|^2 dt} \right]^{1/5} n^{-1/5} := cn^{-1/5},$$

where $\mu_2 = \int_{\mathbb{R}} x^2 K(x) dx$ and $\phi_0 = \int_{\mathbb{R}} K^2(x) dx$. Let $b_n = n^{-1/5}$, then we obtain the pilot estimator $\tilde{\beta}'(t)$ via the local linear estimation [\(4.1\)](#). Next letting $m = \lfloor n^{4/15} \rfloor$, $\tau_n = n^{-5/29}$, we obtain the pilot estimator of $\boldsymbol{\Sigma}(t)$ via the difference-based approach in [Section 5](#). Thus for Epanechnikov kernel, the lower and upper bound of b_n^* is given by

$$b_L^* = \hat{c}n^{-1/4}, \quad b_U^* = \hat{c}n^{-1/6}, \quad \hat{c} = \left[\frac{15 \sum_{i=1}^n \text{tr}\{\hat{\boldsymbol{\Sigma}}(i/n)\}}{\sum_{i=\lfloor nb_n \rfloor+2}^{n-\lfloor nb_n \rfloor} |\tilde{\beta}'(t_i) - \tilde{\beta}'(t_{i-1})|^2} \right]^{1/5}.$$

For the choice of m and τ_n for estimating $\boldsymbol{\Sigma}(t)$ and $\sigma_H(t)$ in [Section 5](#), as a rule of thumb, we can simply choose $m^* = \lfloor n^{4/15} \rfloor, \tau_n^* = n^{-5/29}$ under which [Assumption 5.1](#) holds. We refer to the results of [Bai and Wu \(2023\)](#). For refinement, we recommend the following extended minimum volatility (MV) method as proposed in Chapter 9 of [Politis et al. \(1999\)](#) which works quite well in our empirical studies. The MV

method has the advantage of robustness under complex dependence structures and does not depend on any parametric assumptions of the time series. To be concrete, we first propose a grid of possible block sizes and bandwidths $\{m_1, m_2, \dots, m_{M_1}\}, \{\tau_1, \tau_2, \dots, \tau_{M_2}\}$. Define the sample variance $s_{m_i, \tau_j}^2(t)$ of the bootstrap statistics as

$$s_{m_i, \tau_j}^2(t) = \frac{1}{99} \sum_{i=1}^{100} \left(\tilde{T}_{n, (i)} - \bar{\tilde{T}}_n \right)^2,$$

where $\tilde{T}_{n, (1)}, \dots, \tilde{T}_{n, (100)}$ are the bootstrap statistics calculated from 100 iterations of [Algorithm 5.1](#) with parameters b_n, m_i and τ_j , and $\bar{\tilde{T}}_n = \sum_{i=1}^{100} \tilde{T}_{n, (i)} / 100$. Then calculate

$$MV(i, j) := \max_{1 \leq k \leq n} \text{SE} \left\{ \cup_{r=-1}^1 \{s_{m_i, \tau_{j+r}}^2(t_k)\} \cup \cup_{r=-1}^1 \{s_{m_{i+r}, \tau_j}^2(t_k)\} \right\},$$

where SE stands for the standard deviation, i.e. the maximand is

$$\frac{1}{4} \left\{ \sum_{r=-1, 1} \left(s_{m_i, \tau_{j+r}}^2(t_k) - \bar{s}_{i, j}^2(t_k) \right)^2 + \sum_{r=-1, 1} \left(s_{m_{i+r}, \tau_j}^2(t_k) - \bar{s}_{i, j}^2(t_k) \right)^2 + \left(s_{m_i, \tau_j}^2(t_k) - \bar{s}_{i, j}^2(t_k) \right)^2 \right\}^{1/2},$$

where

$$\bar{s}_{i, j}^2(t) = \frac{1}{5} \left(\sum_{r=-1, 1} s_{m_i, \tau_{j+r}}^2(t) + \sum_{r=-1, 1} s_{m_{i+r}, \tau_j}^2(t) + s_{m_i, \tau_j}^2(t) \right).$$

Then we select the pair (m_{i^*}, τ_{j^*}) where (i^*, j^*) minimizes $MV(i, j)$. Finally, for η_n , as a rule of thumb, we recommend setting $\eta_n = b_n$, which works reasonably well in our Monte Carlo experiments. The choice of η_n can be also refined by MV methods. Specifically, we can first propose a grid of possible bandwidths $\{\eta_1, \dots, \eta_{M_3}\}$. Denoted by $\hat{\mathbf{M}}_{\eta_i}(t)$ the estimated covariance matrix via [\(5.3\)](#) using $\eta_i, i = 1, 2, \dots, M_3$, and select $\eta = \eta_{j^*}$ where j^* is the minimizer of the following criterion $V^\diamond(i)$,

$$V^\diamond(i) = \max_{1 \leq k \leq n} \sum_{r=-2}^2 \left| \hat{\mathbf{M}}_{\eta_{i+r}}(t_k) - \bar{\mathbf{M}}_{\eta_i}(t_k) \right|^2,$$

where $\bar{\mathbf{M}}_{\eta_i}(t_k) = \sum_{r=-2}^2 \hat{\mathbf{M}}_{\eta_{i+r}}(t_k) / 5$.

D Additional Simulations

D.1 Simulation results of the independent model (i)

This subsection contains the simulation results on the sensibility of simulated sizes on sample sizes of model [\(i\)](#), [\(ii.1\)](#) and [\(ii.2\)](#) (see [Table D.1](#)) and those of independent model [\(i\)](#), including simulated sizes (see [Table D.2](#)) and powers (see [Figure D.1](#)) with the selection procedure described in [Appendix C](#) in the main article.

D.2 Simulation results with heavy-tail innovations

This subsection contains the simulation results on the sensibility of simulated sizes on sample sizes of model [\(ii.1\)](#) with heavy-tail *i.i.d.* innovations $\varepsilon_i \sim t_8$.

Model	$n = 1000$								1500							
	KPSS		R/S		V/S		K/S		KPSS		R/S		V/S		K/S	
	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%	5%	10%
(i)	5.8	11.5	6.9	10.5	5.8	10.6	4.2	7.9	6.0	11.3	4.7	9.0	5.8	9.2	4.6	10.0
(ii.1)	4.6	9.3	3.9	7.4	5.1	9.4	4.3	9.2	4.9	10.5	5.4	10.7	4.6	9.6	4.6	9.2
(ii.2)	4.7	10.0	5.6	9.9	5.1	9.8	5.4	9.9	5.0	10.2	4.9	10.0	4.7	9.9	4.2	9.3

Table D.1: Simulated Type I errors (in %) of KPSS, R/S, V/S and K/S-type tests for model (i), (ii.1) and (ii.2). The bandwidths m and τ_n are determined by MV selection. The bandwidth b_n is selected by GCV.

b_n	KPSS		R/S		V/S		K/S	
	5%	10%	5%	10%	5%	10%	5%	10%
0.15	4.6	9.0	5.8	10.5	4.9	9.3	4.2	8.9
0.175	4.4	10.1	5.5	9.6	5.8	11.1	5.5	9.7
0.2	5.7	10.1	4.8	9.6	5.5	11.0	5.8	10.6
0.225	5.2	10.5	5.8	10.9	5.4	8.8	4.5	10.0

Table D.2: Simulated sizes (in %) of KPSS, R/S, V/S and K/S tests for model (i). The bandwidths m and τ_n are determined by MV selection.




Figure D.1: Simulated powers of KPSS and related tests for (i) with nominal level 0.1. Left: $n = 1500$ and d increases from 0 to 0.5; Right: $d = 0.4$ and the sample size n increases from 200 to 2500.

D.3 Power performance of model (ii.2)

Figure D.2 depicts the power performance under data generating model (ii.2). As shown in the left panel, when the long memory parameter increases, the rejection rates of all the four KPSS and related tests grow to 1. The right panel reports the power performance as the sample size increases. It implies that KPSS and related tests have asymptotic power 1.

	KPSS		R/S		V/S		K/S	
b_n	5%	10%	5%	10%	5%	10%	5%	10%
0.15	5.1	10.6	5.0	10.1	5.0	9.0	4.5	9.7
0.175	4.9	10.1	4.4	8.5	5.6	9.9	4.7	9.7
0.2	5.0	9.9	5.0	9.3	6.1	10.4	5.2	9.6
0.225	5.5	10.4	5.2	10.3	6.4	12.2	4.8	10.9

Table D.3: Simulated sizes (in %) of KPSS, R/S, V/S and K/S tests for model (ii.1) with *i.i.d.* t_8 innovations ε_i . The bandwidths m and τ_n are determined by MV selection.




Figure D.2: Simulated powers of KPSS and related tests for (ii.2) with nominal level 0.1. Left: $n = 2500$ and d increases from 0 to 0.5; Right: $d = 0.4$ and the sample size n increases from 200 to 3000.

D.4 Simulation results of time-varying d

Although our theory is established for d as a constant, we examine numerically the power performance of proposed tests as functions of $F = \int_0^1 d(u)du$. In particular, we consider another configuration of d , i.e.

$$d_2(t) = 0.35 + 0.1 \cos(2\pi t) \quad t \in [0, 1].$$

In Figure D.3, KPSS and related tests display good power performance under models (i), (ii.1) and (ii.2), in that the rejection rates of all tests grow to 1 as the sample size increases.

E Analysis of the COVID-19 infection curve

We investigate the time series of the cumulative confirmed cases and deaths of COVID-19 in Japan and Ireland, all in log-scale. For each series, we consider the sub-series from the date when its number first exceeds 500 to 10/06/2021. Our data is obtained from the European Centre for Disease Prevention and Control (ECDC), where the confirmed cases and deaths are updated daily. The cumulative confirmed cases and deaths of COVID-19 has been modeled by a piecewise linear trend model in Jiang et al. (2020). We consider the time-varying trend model (4.5) and apply Algorithm G.1 to testing whether the series of log cumulative confirmed cases and deaths of COVID-19 in Japan and Ireland are LRD. We test for long memory in the two series of each country using the four KPSS and related tests with the critical values generated by 5000 times of bootstrap. For the smoothing parameters, we apply MV criterion to select m




Figure D.3: Simulated rejection rates for model (i), (ii.1) and (ii.2) under the alternative d_2 , nominal level 0.1, starting from $n = 200$.




Figure E.1: Cumulative confirmed cases (left) and deaths (right) in log-scale of COVID-19 in Japan and Ireland

between $\lfloor \frac{6}{7}n^{4/15} \rfloor$ and $\lfloor \frac{12}{7}n^{4/15} \rfloor$ and τ_n between $\lfloor \frac{26}{29}n^{-5/29} \rfloor$ and $\lfloor \frac{34}{29}n^{-5/29} \rfloor$, where n is the length of the time series. For the cumulative confirmed cases of Japan with $n = 577$, m is selected as 4 for KPSS, R/S, V/S and K/S-type tests and τ_n 's are chosen as 0.350, 0.350, 0.300, 0.350, respectively. For the cumulative confirmed cases of Ireland with $n = 567$, m is selected as 4 for KPSS, R/S, V/S and K/S-type tests and τ_n 's are chosen as 0.350, 0.350, 0.300, 0.350, respectively. For the cumulative confirmed deaths of Japan with $n = 524$, m is selected as 4 and τ_n 's are chosen as 0.355 for four KPSS and related tests. For the cumulative confirmed deaths of Ireland with $n = 538$, m is selected as 4 for KPSS, and V/S-type tests, 9 for R/S and K/S -type tests and τ_n 's are chosen as 0.353, 0.303, 0.303, 0.353, respectively. By GCV criterion, we select b_n as 0.096, 0.085 for cumulative confirmed cases and deaths of Japan, respectively, and 0.090, 0.088 for those of Ireland.

Figure E.1 displays the time series of cumulative confirmed cases and deaths of Japan and Ireland in log-scale, respectively. The p -values of KPSS and related tests are shown in Table E.1. For the cumulative confirmed cases series, all four tests reject the null hypothesis at the significance of 0.05, which indicates significant long-range dependence in the time series of log cumulative confirmed cases of COVID-19 in both countries. On the other hand, all the p -values of the four KPSS and related tests for the cumulative confirmed deaths series of COVID-19 exceed 0.05, which fails to reject that the series of log cumulative confirmed deaths of COVID-19 is SRD in either Japan or Ireland.

	KPSS	R/S	V/S	K/S
cases	0.0012	0.0096	2×10^{-4}	0.0022
deaths	0.4332	1	0.9996	0.8244

	KPSS	R/S	V/S	K/S
cases	0.0198	0.0276	0.0042	0.0032
deaths	0.4524	1	0.7986	0.998

Table E.1: p -values of KPSS and related tests for the cumulative confirmed cases and deaths in Japan (left panel) and Ireland (right panel).

F Details in analyzing Hong Kong circulatory and respiratory data

Figure F.1 shows the sample path of the covariates (SO_2 , NO_2 , dust) and the response (the total number of the hospital admissions) in model (6.1). Table F.1 summarizes the smoothing parameters selected in KPSS and related tests when testing for long memory in the series of SO_2 , NO_2 , dust and total number of hospital admissions modeled by (4.5) and by (6.1).




Figure F.1: Sample paths of the time series of SO_2 , NO_2 , dust and the total number of hospital admissions for the Hong Kong circulatory and respiratory data.

G Bootstrap Algorithms

Algorithm G.1 presents the algorithms of KPSS and related tests for the time-varying trend model without time series covariates. Algorithm G.2 presents the algorithms of R/S, V/S and K/S-type tests for the time-varying coefficient model, and Theorem G.1 investigates the limiting distributions of bootstrap statistics in R/S, V/S and K/S-type tests under null, fixed and the local alternatives.

Algorithm G.1 The bootstrap procedure for KPSS and related tests for the time-varying trend model

1. Select the window size m and bandwidth b_n, τ_n , according to the methods in [Appendix C](#).
2. Calculate $\tilde{e}_i = y_i - \tilde{\beta}_1(t_i), i = 1, 2, \dots, n$, where $\tilde{\beta}_1$ is obtained by local linear regression (4.1) with $p = 1$ and jackknife correction (4.2). Then, compute the KPSS-type statistic T_n (4.3), R/S-type statistic Q_n in (B.1), V/S-type statistic M_n in (B.2), K/S-type statistic G_n in (B.3).
3. Calculate $\hat{\sigma}_H^2(t)$ using the estimator in (4.7) of Section 4.2 of [Dette and Wu \(2019\)](#).
4. Generate B (say 2000) *i.i.d.* copies of $N(0, 1)$ variables $V_i^{(r)}$, for $1 \leq r \leq B$, then calculate

$$\tilde{G}_k^{(r)} = \sum_{i=\lfloor nb_n \rfloor + 1}^k \hat{\sigma}_H(t_i) V_i^{(r)} - \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^k \sum_{j=1}^n \hat{\sigma}_H(t_j) V_j^{(r)} K_{b_n}(t_j - t_i).$$

and the bootstrap version of the KPSS-type statistic (4.3),

$$\tilde{T}_n^{(r)} = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \sum_{s=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\sum_{k=\lfloor nb_n \rfloor + 1}^s \tilde{G}_k^{(r)} \right)^2,$$

the bootstrap version of the R/S-type statistic,

$$\widetilde{\text{RS}}_n^{(r)} = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)} - \min_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)},$$

the bootstrap version of the V/S-type statistic,

$$\widetilde{\text{VS}}_n^{(r)} = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \left\{ \sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} (\tilde{G}_k^{(r)})^2 - \frac{1}{n - 2\lfloor nb_n \rfloor} \left(\sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)} \right)^2 \right\},$$

the bootstrap version of the K/S-type statistic,

$$\widetilde{\text{KS}}_n^{(r)} = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} |\tilde{G}_k^{(r)}|.$$

5. Let $\tilde{T}_{n,(1)} \leq \tilde{T}_{n,(2)} \leq \dots \leq \tilde{T}_{n,(B)}$ be the ordered statistics of $\tilde{T}_n^{(r)}, r = 1, 2, \dots, B$. Let $\widetilde{\text{RS}}_{n,(1)} \leq \widetilde{\text{RS}}_{n,(2)} \leq \dots \leq \widetilde{\text{RS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{RS}}_n^{(r)}\}_{r=1}^B$, $\widetilde{\text{VS}}_{n,(1)} \leq \widetilde{\text{VS}}_{n,(2)} \leq \dots \leq \widetilde{\text{VS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{VS}}_n^{(r)}\}_{r=1}^B$, $\widetilde{\text{KS}}_{n,(1)} \leq \widetilde{\text{KS}}_{n,(2)} \leq \dots \leq \widetilde{\text{KS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{KS}}_n^{(r)}\}_{r=1}^B$. Let $B^* = \max\{r : \tilde{T}_{n,(r)} \leq T_n\}$. Let $B_{\text{RS}}^* = \max\{r : \widetilde{\text{RS}}_{n,(r)} \leq Q_n\}$, $B_{\text{VS}}^* = \max\{r : \widetilde{\text{VS}}_{n,(r)} \leq M_n\}$, $B_{\text{KS}}^* = \max\{r : \widetilde{\text{KS}}_{n,(r)} \leq G_n\}$. Then the p -value of KPSS-type test is $1 - B^*/B$, the p -value of the R/S-type test is $1 - B_{\text{RS}}^*/B$, the p -value of the V/S-type test is $1 - B_{\text{VS}}^*/B$, and the p -value of the K/S-type test is $1 - B_{\text{KS}}^*/B$. Reject H_0 at the level of α for each type of test if its p -value is smaller than α .
-

	KPSS	R/S	V/S	K/S
	b_n	(m, τ_n)	(m, τ_n)	(m, τ_n)
SO ₂	0.250	(16, 0.338)	(16, 0.338)	(16, 0.338)
NO ₂	0.149	(11, 0.288)	(11, 0.338)	(11, 0.288)
Dust	0.144	(7, 0.288)	(6, 0.338)	(7, 0.288)
model (4.5)	0.138	(6, 0.338)	(6, 0.338)	(6, 0.338)
model (6.1)	0.181	(6, 0.338)	(11, 0.288)	(9, 0.288)

Table F.1: Selected smoothing parameters of KPSS and related tests for SO₂, NO₂, dust and daily total number of hospital admissions modeled by (4.5) and by (6.1), respectively.

Theorem G.1. *The bootstrap statistics $\widetilde{\text{RS}}_n, \widetilde{\text{VS}}_n, \widetilde{\text{KS}}_n$ are defined in Algorithm G.2. Then, we have the following results*

(i) *Under the conditions of Theorem 5.1(i), we have under H_0*

$$\widetilde{\text{RS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} U(t) - \inf_{0 \leq t \leq 1} U(t), \quad \widetilde{\text{VS}}_n \Rightarrow \int_0^1 U^2(t)dt - \left(\int_0^1 U(t)dt \right)^2, \quad \widetilde{\text{KS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} |U(t)|,$$

where $U(t)$ is as defined in Theorem 4.1.

(ii) *For the fixed alternatives, under the conditions of Theorem 5.2 (i), we have*

$$m^{-d}\widetilde{\text{RS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} \tilde{U}_d(t) - \inf_{0 \leq t \leq 1} \tilde{U}_d(t),$$

$$m^{-2d}\widetilde{\text{VS}}_n \Rightarrow \int_0^1 \tilde{U}_d^2(t)dt - \left(\int_0^1 \tilde{U}_d(t)dt \right)^2, \quad m^{-d}\widetilde{\text{KS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} |\tilde{U}_d(t)|,$$

where $\tilde{U}_d(t)$ is as defined in (i) of Theorem 5.2.

(iii) *For the local alternatives $d_n = c/\log n$ with some constant $c > 0$, under the conditions of Theorem 5.2 (ii), we have*

$$\widetilde{\text{RS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} \check{U}_\alpha(t) - \inf_{0 \leq t \leq 1} \check{U}_\alpha(t), \quad \widetilde{\text{VS}}_n \Rightarrow \int_0^1 \check{U}_\alpha^2(t)dt - \left(\int_0^1 \check{U}_\alpha(t)dt \right)^2, \quad \widetilde{\text{KS}}_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} |\check{U}_\alpha(t)|,$$

where $\check{U}_\alpha(t)$ is as defined in (ii) of Theorem 5.2.

Theorem G.1 follows from the proofs of Theorem 5.1 and Theorem 5.2 and continuous mapping theorem. Therefore, for the sake of brevity, we omit its proof.

H Proofs and related results of Sections 3 and 4

H.1 Proof of Proposition 3.1

By Lemma 3.2 of Kokoszka and Taqqu (1995), under Assumption 3.1, we have

$$\delta_p(H^{(d)}, l, (-\infty, 1]) \leq \sum_{k=0}^l \psi_k(d) \delta_p(H, l-k, (-\infty, 1]) = O\{(1+l)^{d-1}\}.$$

Algorithm G.2 The bootstrap procedure of R/S, V/S, K/S-type tests for time-varying coefficient models

1. Select the window size m and bandwidth b_n, τ_n , according to the methods in [Appendix C](#).
2. Calculate $\tilde{e}_i = y_i - \mathbf{x}_i^\top \tilde{\beta}(t_i), i = 1, 2, \dots, n$, where $\tilde{\beta}$ is obtained using local linear regression (4.1) and jackknife correction (4.2). Then, compute R/S-type statistic Q_n in (B.1), V/S-type statistic M_n in (B.2), K/S-type statistic G_n in (B.3).
3. Calculate $\hat{\mathbf{M}}(t)$ and $\hat{\Sigma}(t)$ defined in (5.3) and (5.4) of [Section 5](#), respectively.
4. Generate B (say 2000) *i.i.d.* copies of $N(\mathbf{0}, \mathbf{I}_p)$ vectors $\mathbf{V}_i^{(r)} = (V_{i,1}^{(r)}, \dots, V_{i,p}^{(r)})^\top$, for $1 \leq r \leq B$, then calculate (notice that $\hat{\sigma}_H^2(t) = (\hat{\Sigma}(t))_{1,1}$)

$$\tilde{G}_k^{(r)} = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^k \mathbf{x}_{i,n}^\top \hat{\mathbf{M}}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \hat{\Sigma}^{1/2}(t_j) \mathbf{V}_j^{(r)} + \sum_{i=\lfloor nb_n \rfloor + 1}^k \hat{\sigma}_H(t_i) V_{i,1}^{(r)},$$

and the bootstrap version of the R/S-type statistic,

$$\widetilde{\text{RS}}_n^{(r)} = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)} - \min_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)},$$

the bootstrap version of the V/S-type statistic,

$$\widetilde{\text{VS}}_n^{(r)} = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \left\{ \sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} (\tilde{G}_k^{(r)})^2 - \frac{1}{n - 2\lfloor nb_n \rfloor} \left(\sum_{k=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \tilde{G}_k^{(r)} \right)^2 \right\},$$

the bootstrap version of the K/S-type statistic,

$$\widetilde{\text{KS}}_n^{(r)} = \max_{\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} |\tilde{G}_k^{(r)}|.$$

5. Let $\widetilde{\text{RS}}_{n,(1)} \leq \widetilde{\text{RS}}_{n,(2)} \leq \dots \leq \widetilde{\text{RS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{RS}}_n^{(r)}\}_{r=1}^B$, $\widetilde{\text{VS}}_{n,(1)} \leq \widetilde{\text{VS}}_{n,(2)} \leq \dots \leq \widetilde{\text{VS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{VS}}_n^{(r)}\}_{r=1}^B$, $\widetilde{\text{KS}}_{n,(1)} \leq \widetilde{\text{KS}}_{n,(2)} \leq \dots \leq \widetilde{\text{KS}}_{n,(B)}$ be the ordered statistics of $\{\widetilde{\text{KS}}_n^{(r)}\}_{r=1}^B$. Let $B_{\text{RS}}^* = \max\{r : \widetilde{\text{RS}}_{n,(r)} \leq Q_n\}$, $B_{\text{VS}}^* = \max\{r : \widetilde{\text{VS}}_{n,(r)} \leq M_n\}$, $B_{\text{KS}}^* = \max\{r : \widetilde{\text{KS}}_{n,(r)} \leq G_n\}$. Then the p -value of the R/S-type test is $1 - B_{\text{RS}}^*/B$, the p -value of the V/S-type test is $1 - B_{\text{VS}}^*/B$, and the p -value of the K/S-type test is $1 - B_{\text{KS}}^*/B$. Reject H_0 at the level of α for each type of test if its p -value is smaller than α .
-

H.2 Limiting distributions of R/S, V/S, and K/S-type statistics

The limiting behavior of R/S, V/S and K/S-type statistics defined in [Section 4](#) under [Assumption 4.5](#) can be derived by [Theorem 4.1](#), [Theorem 4.3](#) and [Theorem 4.4](#) as well as an application of continuous mapping theorem. Recall the definitions of $U(t)$, $U_d(t)$, $U^\circ(t)$ in [Theorem 4.1](#), [Theorem 4.3](#) and [Theorem 4.4](#).

For the R/S-type statistic defined in [\(B.1\)](#), under H_0 , we have $Q_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} U(t) - \inf_{0 \leq t \leq 1} U(t)$. Under fixed alternatives with long-memory parameter d , we have $Q_n \Gamma(d+1)/n^{d+1/2} \Rightarrow \sup_{0 \leq t \leq 1} U_d(t) - \inf_{0 \leq t \leq 1} U_d(t)$, and under local alternatives with $d_n = c/\log n$, $Q_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} U^\circ(t) - \inf_{0 \leq t \leq 1} U^\circ(t)$.

For the V/S-type statistic defined in [\(B.2\)](#), under H_0 , we have $M_n \Rightarrow \int_0^1 U^2(t)dt - \left(\int_0^1 U(t)dt\right)^2$. Under fixed alternatives with long memory parameter d , we have $M_n \Gamma^2(d+1)/n^{2d} \Rightarrow \int_0^1 U_d^2(t)dt - \left(\int_0^1 U_d(t)dt\right)^2$, and under local alternatives with $d_n = c/\log n$, $M_n \Rightarrow \int_0^1 U^{\circ,2}(t)dt - \left(\int_0^1 U^\circ(t)dt\right)^2$.

For the K/S-type statistic defined in [\(B.3\)](#), under H_0 , we have $G_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} |U(t)|$. Under fixed alternatives with long-memory parameter d , we have $G_n \Gamma(d+1)/n^{d+1/2} \Rightarrow \sup_{0 \leq t \leq 1} |U_d(t)|$, and under local alternatives with $d_n = c/\log n$, $G_n/\sqrt{n} \Rightarrow \sup_{0 \leq t \leq 1} |U^\circ(t)|$.

H.3 Proof of [Theorem 4.1](#)

Before proving [Theorem 4.1](#), we study the covariance between \mathbf{x}_i and $\mathbf{x}_j e_j$.

Proposition H.1. *Let $\bar{\mathbf{x}}_i^\top = \mathbf{x}_i^\top - \boldsymbol{\mu}_W(t_i)$ be a p -dimensional vector with j_{th} entry $\bar{x}_{i,j}$. Let $x_{i,l}$ be l_{th} entry of \mathbf{x}_i . Then under [Assumption 4.3](#) and [4.4](#), $1 \leq l, k \leq p$, we have that*

$$\max_{1 \leq i, j \leq n, 1 \leq k, l \leq p} |\mathbb{E}(\bar{x}_{i,l} x_{j,k} e_j)| = O(\chi^{|i-j|}).$$

Proof of [Proposition H.1](#). Under [Assumption 4.4](#), $\bar{x}_{i,k} = \sum_{m \in \mathbb{Z}} \mathcal{P}_m\{\bar{x}_{i,k}\}$, $x_{j,k} e_j = \sum_{m \in \mathbb{Z}} \mathcal{P}_m\{x_{j,k} e_j\}$. Then, with the orthogonality of \mathcal{P}_j , we have

$$|\mathbb{E}(\bar{x}_{i,l} x_{j,k} e_j)| = \left| \mathbb{E} \left[\sum_{m \in \mathbb{Z}} \mathcal{P}_m\{\bar{x}_{i,l}\} \mathcal{P}_m\{x_{j,k} e_j\} \right] \right| \leq \sum_{m \in \mathbb{Z}} \delta_2(\mathbf{W}, i-m) \delta_2(\mathbf{U}, j-m) = O(\chi^{|i-j|}).$$

The last equality follows from the SRD conditions in [Assumptions 4.3](#) and [4.4](#). \square

Lemma H.1. *Under the condition of [Theorem 4.1](#), we have*

$$\sup_{t \in \mathcal{T}} |\mathbf{S}_{n,0}(t) - \mathbf{M}(t)| = O_{\mathbb{P}}(n^{-1/2} b_n^{-3/4} + b_n^2),$$

where $\mathbf{S}_{n,0} = \frac{1}{nb_n} \sum_{i=1}^n \mathbf{x}_i \mathbf{x}_i^\top K_{b_n}(t_i - t)$.

Proof. Similar to the proof in Lemma 6 of [Zhou and Wu \(2010\)](#), under [Assumption \(B4\)](#), we have

$$\|\mathbf{S}_{n,0}(t) - \mathbb{E}\{\mathbf{S}_{n,0}(t)\}\|_4 = O((nb_n)^{-1/2}).$$

By the chaining argument in Proposition B.1 in [Section B.2](#) in [Dette et al. \(2019\)](#), we have

$$\left\| \sup_{t \in \mathcal{T}} |\mathbf{S}_{n,0}(t) - \mathbb{E}\{\mathbf{S}_{n,0}(t)\}| \right\|_4 = O(n^{-1/2} b_n^{-3/4}).$$

Finally, under (B2) and Assumption 4.1, we have uniformly for $t \in \mathcal{T}$, that

$$\begin{aligned} |\mathbb{E}\{\mathbf{S}_{n,0}(t) - \mathbf{M}(t)\}| &= \left| \frac{1}{nb_n} \sum_{i=1}^n \mathbf{M}(t_i) K_{b_n}(t_i - t) - \mathbf{M}(t) \right| \\ &= \left| \frac{1}{nb_n} \sum_{i=1}^n (\mathbf{M}(t) + \mathbf{M}'(t)(t_i - t) + O(b_n^2)) K_{b_n}(t_i - t) - \mathbf{M}(t) \right| = O((nb_n)^{-1} + b_n^2). \end{aligned}$$

□

H.3.1 Proof of Theorem 4.1 (i)

Define $\mathcal{T} = [b_n, 1 - b_n]$. Under Assumptions 4.1, 4.2, 4.3 and 4.4, replacing Lemma 6 of Zhou and Wu (2010) by Lemma H.1 in the proof of Theorem 3 of Zhou and Wu (2010) yields that

$$\sup_{t \in \mathcal{T}} \left| \tilde{\beta}_{b_n}(t) - \beta(t) - \sum_{i=1}^n \frac{\mathbf{M}^{-1}(t)}{nb_n} \mathbf{x}_i e_i K_{b_n}^*(t_i - t) \right| = O_{\mathbb{P}}(\rho'_n \chi'_n),$$

where $\rho'_n = (nb_n)^{-1/2} \log n + b_n^2$, $\chi'_n = n^{-1/2} b_n^{-3/4} + b_n^2$, $K_{b_n}^*(t_i - t) = 2K_{\frac{b_n}{\sqrt{2}}}(t_i - t) - K_{b_n}(t_i - t)$. Then, uniformly for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$, we have

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i^\top \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{x}_j e_j + \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i + O_{\mathbb{P}}(n \rho'_n \chi'_n). \quad (\text{H.1})$$

Define the following function $G^*(r)$, $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$G^*(r) = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{x}_j e_j + \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i. \quad (\text{H.2})$$

Then combining (H.1) and (H.2), we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| G^*(r) - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i \right| \leq \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\tilde{M}_r| + O_{\mathbb{P}}(n \rho'_n \chi'_n), \quad (\text{H.3})$$

where

$$\tilde{M}_r = \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r (\mathbf{x}_i^\top - \boldsymbol{\mu}_W^\top(t_i)) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{x}_j e_j.$$

We shall show (i) the bound for $\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\tilde{M}_r|$, (ii) the asymptotic behavior of the process $G^*(r)$. We break the proof into several steps. Step 1 derives the maximum bound for $|\tilde{M}_r|$. The Gaussian approximation result of $G^*(r)$ is established in Step 2. In Step 3, we obtain the limiting distribution of $G^*(\lfloor nt \rfloor)/\sqrt{n}$ and its convergence with Skorohod topology on $D[0, 1]$.

Step 1: We shall show that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\tilde{M}_r| = O_{\mathbb{P}}(b_n^{-1}). \quad (\text{H.4})$$

Let $\bar{\mathbf{x}}_i^\top = \mathbf{x}_i^\top - \boldsymbol{\mu}_W(t_i)$ be a p -dimensional vector with j_{th} entry $\bar{x}_{i,j}$. Let $x_{i,l}$ be l_{th} entry of \mathbf{x}_i . For the sake of brevity, let $\mathbf{L}_s = \mathbf{x}_s e_s$ and $L_{s,k}$ be the k_{th} element of \mathbf{L}_s , $M_{l,k}^{-1}(t)$ be the element in the l_{th} row and k_{th} column of $\mathbf{M}^{-1}(t)$, where $1 \leq k, l \leq p$. Assumption 4.4 guarantees $\sup_{t \in [0,1], 1 \leq l, k \leq p} |M_{l,k}^{-1}(t)|$ is bounded. Consider the following m -dependent sequences

$$\tilde{L}_{s,k,m} = \mathbb{E}(x_{s,k} e_s | \varepsilon_s, \dots, \varepsilon_{s-m}), \quad \tilde{\tilde{x}}_{s,k,m} = \mathbb{E}(x_{s,k} - \mathbb{E}(x_{s,k}) | \varepsilon_s, \dots, \varepsilon_{s-m}), 1 \leq k \leq p.$$

Further define

$$\tilde{M}_r^{(m)} = \sum_{k=1}^p \sum_{l=1}^p \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \bar{x}_{i,l} M_{l,k}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \tilde{L}_{j,k,m},$$

and

$$\bar{M}_r^{(m)} = \sum_{k=1}^p \sum_{l=1}^p \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\tilde{x}}_{i,l,m} M_{l,k}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \tilde{L}_{j,k,m}.$$

Write $\tilde{\mathcal{F}}_{s,s-j} = (\varepsilon_{s-j}, \dots, \varepsilon_s)$, $\mathcal{F}_{s,s-j} = (\mathcal{F}_{s-j-1}, \varepsilon_{s-j}^*, \dots, \varepsilon_s)$, where $\{\varepsilon_i^*\}_{i \in \mathbb{Z}}$ are the *i.i.d.* copy of $\{\varepsilon_i\}_{i \in \mathbb{Z}}$. Observe that

$$\tilde{L}_{s,k,m} - L_{s,k} = \sum_{j=m}^{\infty} \{ \mathbb{E}[L_{s,k} | \tilde{\mathcal{F}}_{s,s-j}] - \mathbb{E}[L_{s,k} | \tilde{\mathcal{F}}_{s,s-j-1}] \}$$

is the summation of martingale differences. Let $\tilde{L}_{s,k}^{(i-l)}$ denote changing ε_{i-l} with *i.i.d.* copy ε_{i-l}^* in $\tilde{L}_{s,k}$. Under condition (A2), by triangle inequality (the first inequality (H.5)) and Jensen's inequality (the second inequality (H.6)), we have

$$\|\tilde{L}_{s,k,m} - L_{s,k}\|_4 \leq C \sum_{j=m}^{\infty} \left\| \mathbb{E}[L_{s,k} | \tilde{\mathcal{F}}_{s,s-j}] - \mathbb{E}[L_{s,k} | \tilde{\mathcal{F}}_{s,s-j-1}] \right\|_4 \quad (\text{H.5})$$

$$\leq C \sum_{j=m}^{\infty} \left\| L_{s,k}^{(s-j-1)} - L_{s,k} \right\|_4 = O(\chi^m). \quad (\text{H.6})$$

Then, using Jensen's equality, we have

$$\|\mathcal{P}_{s-j}(\tilde{L}_{s,k,m} - L_{s,k})\|_4 \leq 2\|\tilde{L}_{s,k,m} - L_{s,k}\|_4 = O(\chi^m).$$

At the same time,

$$\|\mathcal{P}_{s-j}(\tilde{L}_{s,k,m} - L_{s,k})\|_4 \leq \|\mathcal{P}_{s-j}\tilde{L}_{s,k,m}\|_4 + \|\mathcal{P}_{s-j}L_{s,k}\|_4 \leq 2\delta_4(U, j) = O(\chi^j).$$

Therefore,

$$\|\mathcal{P}_{s-j}(\tilde{L}_{s,k,m} - L_{s,k})\|_4 = O(\chi^{\max(j,m)}). \quad (\text{H.7})$$

Similarly, we have

$$\|\tilde{\tilde{x}}_{i,l,m} - \bar{x}_{i,l}\|_4 = O(\chi^m), \quad \|\mathcal{P}_{s-j}(\tilde{\tilde{x}}_{i,l,m} - \bar{x}_{i,l})\|_4 = O(\chi^{\max(j,m)}). \quad (\text{H.8})$$

As a consequence, by Burkholder's inequality and (H.7), for some large constant M ,

$$\begin{aligned} & \max_{1 \leq i \leq n} \left\| \sum_{j=1}^n K_{b_n}^* (t_i - t_j) (L_{j,k} - \tilde{L}_{j,k,m}) \right\|_4 \\ & \leq M \max_{1 \leq i \leq n} \sum_{l=0}^{\infty} \left\| \sum_{j=1}^n K_{b_n}^* (t_i - t_j) \mathcal{P}_{j-l} (L_{j,k} - \tilde{L}_{j,k,m}) \right\|_4 = O(\sqrt{nb_n} m \chi^m). \end{aligned} \quad (\text{H.9})$$

Then, by Cauchy inequality and (H.9), it follows that

$$\begin{aligned} & \left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\tilde{M}_r - \tilde{M}_r^{(m)}| \right\| \\ & \leq \sum_{k=1}^p \left\{ \max_{1 \leq i \leq n} \left\| \frac{1}{nb_n} \sum_{j=1}^n K_{b_n}^* (t_i - t_j) (L_{j,k} - \tilde{L}_{j,k,m}) \right\|_4 \times \sum_{i=[nb_n]+1}^{n-[nb_n]} \sum_{l=1}^p \left\| \tilde{x}_{i,l} M_{l,k}^{-1}(t_i) \right\|_4 \right\} \\ & = O\left(p^2 \sqrt{n/b_n} m \chi^m\right). \end{aligned}$$

An elementary calculation using Burkholder's inequality shows that

$$\max_{1 \leq i \leq n} \left\| \sum_{j=1}^n K_{b_n}^* (t_i - t_j) \tilde{L}_{j,k,m} \right\|_4 = O\left(\sqrt{nb_n}\right).$$

Along with equation (H.8), it's straightforward to show that

$$\begin{aligned} & \left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\bar{M}_r^{(m)} - \tilde{M}_r^{(m)}| \right\| \\ & \leq \frac{1}{nb_n} \sum_{k=1}^p \left\{ \sum_{i=[nb_n]+1}^{n-[nb_n]} \sum_{l=1}^p \left\| (\bar{x}_{i,l} - \tilde{x}_{i,l,m}) M_{l,k}^{-1}(t_i) \right\|_4 \times \max_{1 \leq i \leq n} \left\| \sum_{j=1}^n K_{b_n}^* (t_i - t_j) \tilde{L}_{j,k,m} \right\|_4 \right\} \\ & = O\left(p^2 \sqrt{n/b_n} \chi^m\right). \end{aligned}$$

Therefore, $\bar{M}_r^{(m)}$ is an appropriate approximation of \tilde{M}_r , in that

$$\left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\bar{M}_r^{(m)} - \tilde{M}_r| \right\| = O\left(p^2 \sqrt{n/b_n} m \chi^m\right). \quad (\text{H.10})$$

Using the argument similar to Proposition H.1, we have that

$$\max_{[nb_n]+1 \leq r \leq n-[nb_n]} \mathbb{E}(\bar{M}_r^{(m)}) = O\left\{p^2 \sum_{i=1}^n \sum_{j=1}^n \chi^{|i-j|} / (nb_n)\right\} = O\left(p^2 b_n^{-1}\right). \quad (\text{H.11})$$

Now, we proceed to compute the order of $\max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\bar{M}_r^{(m)} - \mathbb{E}(\bar{M}_r^{(m)})|$.

Write $\bar{a}_{n,i}^{(m)} = \sum_{k=1}^p \sum_{l=1}^p \left(\frac{1}{nb_n} \tilde{x}_{i,l,m} M_{l,k}^{-1}(t_i) \sum_{j=1}^n K_{b_n}^* (t_i - t_j) \right) \tilde{L}_{j,k,m}$, then $\bar{M}_r^{(m)} = \sum_{i=[nb_n]+1}^r \bar{a}_{n,i}^{(m)}$.

Observe that $\mathcal{P}_{j-s}(\tilde{x}_{j,l,m} \tilde{L}_{i,k,m}) = 0$, for $s > 2m$. Then, we have

$$\left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\bar{M}_r^{(m)} - \mathbb{E}(\bar{M}_r^{(m)})| \right\| \leq \sum_{s=0}^{2m} \left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} \left| \sum_{i=[nb_n]}^r \mathcal{P}_{i-s} \bar{a}_{n,i}^{(m)} \right| \right\|. \quad (\text{H.12})$$

According to Lemma 3 in [Zhou and Wu \(2010\)](#), by triangle inequality, we have

$$\begin{aligned} \|\mathcal{P}_{i-s}\bar{a}_{n,i}^{(m)}\| &\leq \sum_{k=1}^p \sum_{l=1}^p \left\{ \frac{1}{nb_n} \left\| \tilde{x}_{i,l,m} - \tilde{x}_{i,l,m}^{(i-s)} \right\|_4 \left\| M_{l,k}^{-1}(t_i) \right\| \left\| \sum_{j=1}^n K_{b_n}^*(t_i - t_j) \tilde{L}_{j,k,m} \right\|_4 \right. \\ &\quad \left. + \frac{1}{nb_n} \left\| \tilde{x}_{i,l,m}^{(i-s)} \right\|_4 \left\| M_{l,k}^{-1}(t_i) \right\| \left\| \sum_{j=1}^n K_{b_n}^*(t_i - t_j) \left(\tilde{L}_{j,k,m} - \tilde{L}_{j,k,m}^{(i-s)} \right) \right\|_4 \right\} \\ &= O \left\{ p^2 \left(\frac{\chi^s}{\sqrt{nb_n}} + \frac{m}{nb_n} \right) \right\}. \end{aligned}$$

The last inequality follows from the fact that by Jensen's inequality $\|\tilde{x}_{i,l,m} - \tilde{x}_{i,l,m}^{(i-s)}\|_4 \leq \|\bar{x}_{i,l} - \bar{x}_{i,l}^{(i-s)}\|_4 = O(\chi^s)$, Assumption 4.4, and $\tilde{L}_{j,k,m} - \tilde{L}_{j,k,m}^{(i-s)}$ is zero when $j \leq i-s$ and $j \geq i-s+m$. Then, since $\mathcal{P}_{i-s}\bar{a}_{n,i}^{(m)}$ are martingale differences, by Doob's inequality, we obtain

$$\begin{aligned} \left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} \left| \sum_{i=[nb_n]+1}^r \mathcal{P}_{i-s}\bar{a}_{n,i}^{(m)} \right| \right\| &\leq C \left\| \sum_{i=[nb_n]+1}^{n-[nb_n]} \mathcal{P}_{i-s}\bar{a}_{n,i}^{(m)} \right\| \\ &= O \left\{ p^2 \sqrt{n} \left(\frac{\chi^s}{\sqrt{nb_n}} + \frac{m}{nb_n} \right) \right\}, \end{aligned} \quad (\text{H.13})$$

where C is a positive constant. Plugging (H.13) into inequality (H.12) yields

$$\left\| \max_{[nb_n]+1 \leq r \leq n-[nb_n]} \left| \bar{M}_r^{(m)} - \mathbb{E}(\bar{M}_r^{(m)}) \right| \right\| = O \left\{ p^2 \left(\frac{m^2}{n^{1/2}b_n} + b_n^{-1/2} \right) \right\}. \quad (\text{H.14})$$

Finally, from (H.10), (H.11) and (H.14), when the dimension p is fixed, taking $m = \lfloor \log n \rfloor$, we have proved (H.4).

Therefore, by (H.3) and (H.4), we have

$$\max_{[nb_n]+1 \leq r \leq n-[nb_n]} \left| G^*(r) - \sum_{i=[nb_n]+1}^r \tilde{e}_i \right| = O_{\mathbb{P}} \left(b_n^{-5/4} \log n + nb_n^4 + \sqrt{n} b_n^{5/4} \right),$$

which is of smaller order of \sqrt{n} .

Step 2: Recall $\Sigma(t_i)$ is the long-run covariance matrix of the process $(\mathbf{x}_i e_i)$. Since in our regression we let $\mathbf{x}_{i,1} = 1$ for $1 \leq i \leq n$, $(\Sigma(t_i))_{(1,1)} = \sigma_H^2(t_i)$ is the long-run variance of the process (e_i) . We shall show that there exist *i.i.d.* $N(0, I_p)$, $\mathbf{V}_i = (V_{i,1}, \dots, V_{i,p})^\top$, and

$$\tilde{G}^*(r) = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=[nb_n]+1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \Sigma^{1/2}(t_j) \mathbf{V}_j + \sum_{i=[nb_n]+1}^r \sigma_H(t_i) V_{i,1},$$

such that

$$\max_{[nb_n]+1 \leq r \leq n-[nb_n]} |\tilde{G}^*(r) - G^*(r)| = O_{\mathbb{P}} \left(n^{1/4} \log^2 n \right). \quad (\text{H.15})$$

From Corollary 1 in [Wu and Zhou \(2011\)](#), we have

$$\max_{1 \leq i \leq n} \left| \sum_{j=1}^i \mathbf{x}_j e_j - \sum_{j=1}^i \Sigma^{1/2}(t_j) \mathbf{V}_j \right| = o_{\mathbb{P}} \left(n^{1/4} \log^2 n \right), \quad (\text{H.16})$$

and in the first dimension,

$$\max_{1 \leq i \leq n} \left| \sum_{j=1}^i e_j - \sum_{j=1}^i \sigma_H(t_j) V_{j,1} \right| = o_{\mathbb{P}} \left(n^{1/4} \log^2 n \right). \quad (\text{H.17})$$

Write $\mathbf{m}_{r,j}^\top = \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j)$, then

$$\begin{aligned} \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \tilde{G}^*(r) - G^*(r) \right| &\leq \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} - \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i \right| \\ &+ \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{x}_j e_j - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j \right|. \end{aligned} \quad (\text{H.18})$$

Then, (H.15) follows from (H.16), (H.17), and the summation-by-parts formula.

Step 3: Define $\tilde{G}_{n,b_n}(t) = \tilde{G}^*(\lfloor nt \rfloor) / \sqrt{n}$. We shall show that

$$\tilde{G}_{n,b_n}(t) \rightsquigarrow U(t) \quad \text{on } D[0, 1] \text{ with Skorohod topology.}$$

Under the bandwidth condition $nb_n^3 / (\log n)^2 \rightarrow \infty$, $nb_n^6 \rightarrow 0$, we have from Step 1 and Step 2 that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \tilde{G}^*(r) - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i \right| = o_{\mathbb{P}}(\sqrt{n}).$$

Let $\mu_{W,i}(u)$ denote the i th component of $\boldsymbol{\mu}_W(u)$. Let $m_{r,j,k}$ be the k th element in $\mathbf{m}_{r,j}^\top$, and $\{\cdot\}_k$ be the k th element in the vector. Under condition (B1) and (B3), $\boldsymbol{\mu}_W(t)$ and $\mathbf{M}^{-1}(t)$ are Lipschitz continuous. Since $K^*(t)$ can be non-zero only for $t \in [-1, 1]$, elementary calculation shows

$$m_{r,j,k} = \sum_{i=1}^p \mu_{W,i}^\top(t_j) M_{i,k}^{-1}(t_j) \int_{1-\frac{j}{nb_n}}^{\frac{r-j}{nb_n}} K^*(y) dy + O\left(\frac{1}{nb_n} + b_n\right). \quad (\text{H.19})$$

Consider $0 \leq t_1 \leq t_2 \leq 1$. Let $s = \lfloor nt_1 \rfloor$, $r = \lfloor nt_2 \rfloor$. The covariance of $\tilde{G}_{n,b_n}(t)$ is

$$\begin{aligned} \mathbb{E} \left\{ \frac{\tilde{G}^*(r) \tilde{G}^*(s)}{n} \right\} &= \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{r,j}^\top \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j \sum_{j=1}^n \mathbf{m}_{s,j}^\top \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j \right\} / n \\ &- \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{r,j}^\top \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j \sum_{i=\lfloor nb_n \rfloor + 1}^s \sigma_H(t_i) V_{i,1} \right\} / n \\ &- \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{s,j}^\top \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} \right\} / n \\ &+ \mathbb{E} \left\{ \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} \sum_{i=\lfloor nb_n \rfloor + 1}^s \sigma_H(t_i) V_{i,1} \right\} / n \\ &:= I + II + III + IV. \end{aligned}$$

Without loss of generality, suppose $\lfloor nb_n \rfloor + 1 < s \leq r < n - \lfloor nb_n \rfloor$. Let $M_W(t) = \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}(t) \mathbf{M}^{-1}(t) \boldsymbol{\mu}_W(t)$, we have

$$I = \sum_{j=1}^n \mathbf{m}_{r,j}^\top \boldsymbol{\Sigma}(t_j) \mathbf{m}_{s,j} / n = \int_0^1 M_W(t) \int_{1-\frac{t}{b_n}}^{\frac{r-tn}{nb_n}} K^*(y) dy \int_{1-\frac{t}{b_n}}^{\frac{s-tn}{nb_n}} K^*(y) dy dt + O\left(\frac{1}{nb_n} + b_n\right).$$

Let $M_{WK}(r, s, t) = M_W(t) \int_{1-\frac{t}{b_n}}^{\frac{r-tn}{nb_n}} K^*(y) dy \int_{1-\frac{t}{b_n}}^{\frac{s-tn}{nb_n}} K^*(y) dy$. When $s/(nb_n) \rightarrow \infty$,

$$\int_{2b_n}^{\frac{s-nb_n}{n}} M_{WK}(r, s, t) dt = \int_0^{s/n} M_W(t) dt \left\{ \int_{-1}^1 K^*(y) dy \right\}^2 + O(b_n),$$

Since $\int_{1-\frac{t}{b_n}}^{\frac{s-tn}{nb_n}} K^*(y) dy = 0$ for $t > \frac{s+nb_n}{n}$, we have

$$I = \int_0^{s/n} M_W(r, s, t) dt \left\{ \int_{-1}^1 K^*(y) dy \right\}^2 + O\left(\frac{1}{nb_n} + b_n\right).$$

Similar for the case $s = O(nb_n)$, since

$$\int_0^{s/n} M_{WK}(r, s, t) dt = O(b_n), \quad \int_0^{s/n} M_W(t) dt \left\{ \int_{-1}^1 K^*(y) dy \right\}^2 = O(b_n),$$

we have,

$$I = \int_0^{s/n} M_W(t) dt \left\{ \int_{-1}^1 K^*(y) dy \right\}^2 + O\left(\frac{1}{nb_n} + b_n\right). \quad (\text{H.20})$$

Similar and tedious calculation shows

$$II = - \int_0^{s/n} \{ \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t) \}_1 \sigma_H(t) dt \int_{-1}^1 K^*(y) dy + O\left(\frac{1}{nb_n} + b_n\right). \quad (\text{H.21})$$

For *III*, if $s/(nb_n) \rightarrow \infty$,

$$\begin{aligned} III &= -\frac{1}{n} \sum_{j=\lfloor nb_n \rfloor + 1}^r \{ \mathbf{m}_{s,j}^\top \boldsymbol{\Sigma}(t_j) \}_1 \sigma_H(t_j) \\ &= - \int_{2b_n}^{s/n-b_n} \int_{-1}^1 K^*(y) dy \{ \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t) \}_1 \sigma_H(t) dt \\ &\quad - \int_{s/n-b_n}^{s/n+b_n} \int_{-1}^1 K^*(y) dy \{ \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t) \}_1 \sigma_H(t) dt \\ &\quad - \int_{s/n+b_n}^{r/n} \int_{1-\frac{tn}{nb_n}}^{\frac{s-tn}{nb_n}} K^*(y) dy \{ \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t) \}_1 \sigma_H(t) dt + O\left(\frac{1}{nb_n} + b_n\right) \\ &= - \int_0^{s/n} \{ \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\Sigma}^{1/2}(t) \}_1 \sigma_H(t) dt \int_{-1}^1 K^*(y) dy + O\left(\frac{1}{nb_n} + b_n\right). \end{aligned} \quad (\text{H.22})$$

For the third term in (H.22), consider two cases: $r/n > s/n + b_n$ and $s/n \leq r/n \leq s/n + b_n$. If $s/n \leq r/n \leq s/n + b_n$, $(s/n + b_n) - r/n \leq b_n$. Then the third term in (H.22) is $O(b_n)$. If $r/n > s/n + b_n$,

for $s/n + b_n \leq t \leq r/n$, $\frac{s-tn}{nb_n} \leq -1$. Therefore, this term is 0. By careful investigation, the result still holds if $s = O(nb_n)$.

The calculation of IV is rather straightforward,

$$IV = \int_0^{s/n} \sigma_H^2(t) dt + O\left(\frac{1}{nb_n} + b_n\right). \quad (\text{H.23})$$

Then from the approximation of $I - IV$, we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq s \leq r \leq n - \lfloor nb_n \rfloor} \left| \mathbb{E}\{\tilde{G}^*(s)\tilde{G}^*(r)\}/n - \gamma(s/n, r/n) \right| = O\left(b_n + \frac{1}{nb_n}\right).$$

In addition, define $\tilde{G}^*(s) = \tilde{G}^*(\lfloor nb_n \rfloor)$ if $s < \lfloor nb_n \rfloor + 1$ and $\tilde{G}^*(s) = \tilde{G}^*(n - \lfloor nb_n \rfloor)$ if $n - \lfloor nb_n \rfloor < s \leq n$. By the continuity of γ , we have

$$\sup_{0 \leq t_1 \leq t_2 \leq 1} \left| \mathbb{E}\left\{\tilde{G}^*(\lfloor nt_1 \rfloor)\tilde{G}^*(\lfloor nt_2 \rfloor)\right\}/n - \gamma(t_1, t_2) \right| = O\left(b_n + \frac{1}{nb_n}\right). \quad (\text{H.24})$$

The finite dimension convergence of the Gaussian process $\tilde{G}_{n,b_n}(t)$ to $U(t)$ then follows from the Cramer Wold device.

We proceed to show the tightness of $\tilde{G}_{n,b_n}(t)$. For $1 \leq r \leq s \leq n$, since

$$\tilde{G}^*(s) - \tilde{G}^*(r) = - \sum_{j=1}^n \left(\mathbf{m}_{s,j}^\top - \mathbf{m}_{r,j}^\top \right) \Sigma^{1/2}(t_j) \mathbf{V}_j + \sum_{i=r+1}^s \sigma_H(t_i) V_{i,1},$$

it follows from Burholder's inequality that

$$\begin{aligned} \left\| \tilde{G}^*(s) - \tilde{G}^*(r) \right\|_4^2 &\leq K_0 \left(\sum_{j=1}^n \left\| \left(\mathbf{m}_{s,j}^\top - \mathbf{m}_{r,j}^\top \right) \Sigma^{1/2}(t_j) \mathbf{V}_j \right\|_4^2 + \sum_{i=r+1}^s \left\| \sigma_H(t_i) V_{i,1} \right\|_4^2 \right) \\ &\leq K_1 \sum_{j=1}^n \left(\mathbf{m}_{s,j}^\top - \mathbf{m}_{r,j}^\top \right) \Sigma(t_j) (\mathbf{m}_{s,j} - \mathbf{m}_{r,j}) + K_2(s - r) \end{aligned}$$

where K_0 , K_1 and K_2 are sufficiently large constants. For the first term, by the result in (H.19), we have

$$m_{s,j,k} - m_{r,j,k} = \sum_{i=1}^p \mu_{W,i}^\top(t_j) M_{i,k}^{-1}(t_j) \int_{\frac{r-j}{nb_n}}^{\frac{s-j}{nb_n}} K^*(y) dy + O\left(b_n + \frac{1}{nb_n}\right).$$

Observe that $m_{s,j,k} - m_{r,j,k}$ is zero when $j < r - \lfloor nb_n \rfloor + 1$ and $j > s + \lfloor nb_n \rfloor$. When $r - \lfloor nb_n \rfloor + 1 \leq j \leq s + \lfloor nb_n \rfloor$, if $s - r > 2nb_n$, $m_{s,j,k} - m_{r,j,k}$ is $O(1)$ and otherwise $O\left(\frac{s-r}{nb_n}\right)$. Hence, if $s - r > 2nb_n$,

$$\left\| \tilde{G}^*(s) - \tilde{G}^*(r) \right\|_4^2 \leq K_3(s - r + 2\lfloor nb_n \rfloor) + K_2(s - r) = O(|s - r|), \quad (\text{H.25})$$

while for $s - r \leq 2nb_n$,

$$\left\| \tilde{G}^*(s) - \tilde{G}^*(r) \right\|_4^2 \leq K_4 \frac{(s - r)^2 (s - r + 2\lfloor nb_n \rfloor)}{(nb_n)^2} + K_2(s - r) = O(|s - r|), \quad (\text{H.26})$$

where K_3, K_4 are sufficiently large constants. Hence, for $0 \leq t_1 \leq t \leq t_2 \leq 1$, there exists a sufficiently large constant K , s.t.

$$\begin{aligned} & \mathbb{E} \left\{ |\tilde{G}_{n,b_n}(t) - \tilde{G}_{n,b_n}(t_1)|^2 |\tilde{G}_{n,b_n}(t_2) - \tilde{G}_{n,b_n}(t)|^2 \right\} \\ & \leq \left(\|\tilde{G}^*(\lfloor nt \rfloor) - \tilde{G}^*(\lfloor nt_1 \rfloor)\|_4 \|\tilde{G}^*(\lfloor nt_2 \rfloor) - \tilde{G}^*(\lfloor nt \rfloor)\|_4 \right)^2 / n^2 \\ & \leq K(t_2 - t_1)^2. \end{aligned} \quad (\text{H.27})$$

Equation (13.2) of Billingsley (1999) follows from the continuity of $U(t)$. By Theorem 13.5 in Billingsley (1999), the $\alpha = \beta = 1$ case, we have the tightness of $\tilde{G}_{n,b_n}(t)$. The tightness of $\tilde{G}_{n,b_n}(t)$ and the finite dimension convergence lead to the convergence $\tilde{G}_{n,b_n}(t) \rightsquigarrow U(t)$ on $D[0, 1]$ with Skorohod topology. Finally, by the continuous mapping theorem, we have proved the convergence of T_n to $\int_0^1 U^2(t)dt$. \square

H.3.2 Proof of Theorem 4.1 (ii)

Let $\mathbf{z}(u) := (z_1(u), \dots, z_p(u))^\top$ denote $\Sigma^{1/2}(u)\mathbf{M}^{-1}(u)\boldsymbol{\mu}_W(u)$. Then the covariance structure of the limiting distribution in (i) of Theorem 4.1 can be written as

$$\mathbb{E}(U(r)U(s)) = \int_0^{r \wedge s} \left((\sigma_H(u) - z_1(u))^2 + \sum_{j=2}^p z_j^2(u) \right) du.$$

Therefore, the limiting distribution in Theorem 4.1 degenerates if and only if $\lambda(\mathbf{z}(u) \neq (\sigma_H(u), 0, \dots, 0)^\top) > 0$ holds. We shall show that

$$s_1^{-1}(\tilde{T}_n/b_n - s_2) \Rightarrow \chi_1^2, \quad (\text{H.28})$$

where $s_1 = 2\sigma_H^2(0) \int_0^1 \left(\int_{v-1}^1 K^*(t)dt \right)^2 dv$, and $s_2 = 2 \int_0^1 \sigma_H^2(t)dt \int_0^1 \left(\int_v^1 K^*(t)dt \right)^2 dv$. Recall in Step 2 of Theorem 4.1,

$$\tilde{G}^*(r) = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \Sigma^{1/2}(t_j) \mathbf{V}_j + \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1}.$$

By Step 1 and Step 2 in the proof of Theorem 4.1, on a richer probability space we have

$$\begin{aligned} \sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i - \tilde{G}^*(r) \right| &= O_{\mathbb{P}}(b_n^{-5/4} \log n + nb_n^4 + \sqrt{n}b_n^{5/4} + n^{1/4} \log^2 n) \\ &= o_{\mathbb{P}}(\sqrt{nb_n}/\log n). \end{aligned} \quad (\text{H.29})$$

Define

$$\tilde{G}^\circ(r) = \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} - \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r \frac{K_{b_n}^*(t_i - t_j)}{nb_n} \sigma_H(t_j) V_{j,1}.$$

Since $\lambda(\mathbf{z}(u) \neq (\sigma_H(u), 0, \dots, 0)^\top) = 0$, by Taylor series expansion, we have for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$\begin{aligned} \|\tilde{G}^*(r) - \tilde{G}^\circ(r)\|^2 &= \left\| \sum_{j=1}^n \left\{ \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \left(\boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) - \boldsymbol{\mu}_W^\top(t_j) \mathbf{M}^{-1}(t_j) \right) K_{b_n}^*(t_i - t_j) \right\} \Sigma^{1/2}(t_j) \mathbf{V}_j \right\|^2 \\ &= O(nb_n^3). \end{aligned} \quad (\text{H.30})$$

Observe that for $3\lfloor nb_n \rfloor + 1 \leq r \leq n$,

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \frac{K_{b_n}^*(t_i - t_j)}{nb_n} = \begin{cases} \int_{\frac{\lfloor nb_n \rfloor - j}{nb_n}}^1 K^*(t) dt + O((nb_n)^{-1}), & 1 \leq j \leq r - \lfloor nb_n \rfloor, \\ 1 + O((nb_n)^{-1}), & 2\lfloor nb_n \rfloor + 1 \leq j \leq r - \lfloor nb_n \rfloor, \\ \int_{-1}^{\frac{r-j}{nb_n}} K^*(t) dt + O((nb_n)^{-1}), & r - \lfloor nb_n \rfloor + 1 \leq j \leq r + \lfloor nb_n \rfloor, \\ 0, & \text{otherwise.} \end{cases} \quad (\text{H.31})$$

Therefore, we have for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$\|\tilde{G}^\circ(r)\|^2 = O(nb_n).$$

Define

$$\tilde{T}_n^\circ = \frac{1}{n(n - 2\lfloor nb_n \rfloor)} \sum_{r=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} (\tilde{G}^\circ(r))^2.$$

Since $\tilde{G}^*(r) - \tilde{G}^\circ(r)$ and $\tilde{G}^\circ(r)$ are Gaussian processes, we have

$$|\tilde{T}_n^\circ - T_n| = O_{\mathbb{P}}(b_n). \quad (\text{H.32})$$

Write v_j short for $V_{j,1}$. By (H.31) and some tedious calculation, we have

$$\begin{aligned} \tilde{T}_n^\circ / b_n &= \frac{1}{nb_n} \left(\sum_{j=\lfloor nb_n \rfloor + 1}^{2\lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{j - \lfloor nb_n \rfloor - 1}{nb_n}} K^*(t) dt \right) \sigma_H(t_j) v_j - \sum_{j=1}^{\lfloor nb_n \rfloor} \left(\int_{\frac{j - \lfloor nb_n \rfloor - 1}{nb_n}}^1 K^*(t) dt \right) \sigma_H(t_j) v_j \right)^2 \\ &\quad + \frac{1}{n^2 b_n} \sum_{r=3\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\sum_{j=r - \lfloor nb_n \rfloor + 1}^r \left(\int_{\frac{r-j}{nb_n}}^1 K^*(t) dt \right) \sigma_H(t_j) v_j - \sum_{j=r+1}^{r + \lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{r-j}{nb_n}} K^*(t) dt \right) \sigma_H(t_j) v_j \right)^2 \\ &\quad + O_{\mathbb{P}}(b_n^{1/2}) \\ &:= A_n + S_n + O_{\mathbb{P}}(b_n^{1/2}), \end{aligned}$$

where A_n and S_n are defined in the obvious way. Note that A_n and S_n are independent. Since v_i are *i.i.d.* $N(0, 1)$, elementary calculation shows that

$$A_n / \mathbb{E}A_n \sim \chi_1^2, \quad (\text{H.33})$$

and since K^* is symmetric

$$\mathbb{E}A_n = s_1 + O\left(b_n + \frac{1}{nb_n}\right). \quad (\text{H.34})$$

Combining (H.33) and (H.34), by Slutsky's Theorem we have

$$A_n / s_1 \Rightarrow \chi_1^2, \quad (\text{H.35})$$

where s_1 is as defined in (H.28). Observe that

$$\mathbb{E}S_n = s_2 + O\left(b_n + \frac{1}{nb_n}\right), \text{ and } \mathbb{E}S_n^2 = O(b_n) + (\mathbb{E}S_n)^2,$$

where s_2 is as defined in (H.28). Therefore,

$$|S_n - s_2| \leq |S_n - \mathbb{E}S_n| + |\mathbb{E}S_n - s_2| = O_{\mathbb{P}}\left(b_n^{1/2} + \frac{1}{nb_n}\right). \quad (\text{H.36})$$

Then, combining (H.32), (H.35), and (H.36), Equation (H.28) is proved. \square

H.4 Related proofs of Theorem 4.2

This subsection provides the proof of Proposition A.2 and Proposition A.3, which are used in the proof of Theorem 4.2 in Appendix A of the main paper.

H.4.1 Proof of Proposition A.2

Observe that

$$\sum_{k=1}^s \mu_W(k/n) e_k^{(d)} = \sum_{k=1}^s \sum_{j=0}^{\infty} \mu_W(t_k) \psi_j u_{k-j} = \sum_{j=1}^s u_j \sum_{k=j}^s \mu_W(t_k) \psi_{k-j} + \sum_{j=0}^{\infty} u_{-j} \sum_{k=1}^s \mu_W(t_k) \psi_{k+j}$$

Define $Z_j = \sum_{i=0}^j u_{-i}$ with $Z_j = 0$ when $j < 0$ and $S_j = \sum_{i=1}^j u_i$ with $S_j = 0$ when $j \leq 0$. After a careful check of Corollary 2 of Wu and Zhou (2011), there exist independent variables $v_1, v_2, \dots, v_n \sim N(0, 1)$ and independent Gaussian variables $v_i, i \leq 0$, which are independent of $v_j, j > 0$, such that

$$\zeta_n := \max_{1 \leq i \leq n} \left| \sum_{j=1}^i u_j - \sum_{j=1}^i \sigma_H(t_j) v_j \right| = o_{\mathbb{P}} \left(n^{1/4} \log^2 n \right), \quad (\text{H.37})$$

$$\zeta_n^* := \max_{1 \leq i \leq n} \left| \sum_{j=0}^i u_{-j} - \sum_{j=0}^i \sigma_H(t_{-j}) v_{-j} \right| = o_{\mathbb{P}} \left(n^{1/4} \log^2 n \right). \quad (\text{H.38})$$

Define $\mathbf{R}_{k,n} = \sum_{j=0}^{\infty} \mu_W(k/n) \psi_j \sigma(t_{k-j}) v_{k-j}$, $S_j^* = \sum_{i=1}^j \sigma_H(t_i) v_i$, and $Z_j^* = \sum_{i=0}^j \sigma_H(t_{-i}) v_{-i}$. Then, by the summation-by-parts formula, we have for some integer N ,

$$\begin{aligned} \sum_{k=1}^s \left(\mu_W(k/n) e_k^{(d)} - \mathbf{R}_{k,n} \right) &= \sum_{j=1}^{s-1} \left(\sum_{k=j}^s \mu_W(k/n) \psi_{k-j} - \sum_{k=j+1}^s \mu_W(k/n) \psi_{k-j-1} \right) (S_j - S_j^*) \\ &\quad + (S_s - S_s^*) \mu_W(1) \psi_0 \\ &\quad + \sum_{j=0}^{N-1} \sum_{k=1}^s (\mu_W(k/n) \psi_{k+j} - \mu_W(k/n) \psi_{k+j+1}) (Z_j - Z_j^*) \\ &\quad + (Z_N - Z_N^*) \sum_{k=1}^s \psi_{k+N} \mu_W(k/n) \\ &\quad + \sum_{j=N+1}^{\infty} u_{-j} \sum_{k=1}^s \mu_W(k/n) \psi_{k+j} - \sum_{j=N+1}^{\infty} \sigma_H(t_{-j}) v_{-j} \sum_{k=1}^s \mu_W(k/n) \psi_{k+j} \\ &:= A + B + C + D + E + F. \end{aligned}$$

Let $N = \lfloor n^{\alpha_0} \rfloor + 1$, $\alpha_0 > 1$. Condition (B3) indicates that $\mu_W(t)$ is Lipschitz continuous and $\exists C_2 > 0$, $\sup_{t \in [0,1]} |\mu_W(t)| < C_2$. From (H.37), for some positive constant C_1 , and any $0 < q < 1/4$.

$$\max_{1 \leq s \leq n} |A| \leq C_1 \zeta_n \max_{1 \leq s \leq n} \sum_{j=1}^{s-1} \left| \sum_{k=j}^{s-1} \psi_{k-j} (\mu_W(k/n) - \mu_W((k+1)/n)) + \psi_{n-j} \mu_W(1) \right| = O_{\mathbb{P}}(n^{1/4+q+d}).$$

Similar techniques and (H.38) show that,

$$\max_{1 \leq s \leq n} |C| = O_{\mathbb{P}}(n^{\alpha_0/4+q+\alpha_0 d}), \quad \max_{1 \leq s \leq n} |B| \leq |\mu_W(1)| \zeta_n \psi_0 = O_{\mathbb{P}}(n^{1/4+q}), \quad \max_{1 \leq s \leq n} |D| \leq O_{\mathbb{P}}(n^{\alpha_0/4+q+\alpha_0 d}).$$

Uniformly for $1 \leq s \leq n$, $\tilde{\psi}_{j,s} := \sum_{k=1}^s \boldsymbol{\mu}_W(k/n) \psi_{k+j} = O(n|j+1|^{d-1})$, it follows elementary calculation that

$$\begin{aligned} \left\| \max_{1 \leq s \leq n} |E| \right\|^2 &= \mathbb{E} \left(\sum_{j=N+1}^{\infty} |u_{-j}| \max_s |\tilde{\psi}_{j,s}| \right)^2 \\ &= \sum_{i=N+1}^{\infty} \sum_{j=N+1}^{\infty} \max_{1 \leq s \leq n} |\tilde{\psi}_{j,s}| \max_{1 \leq s \leq n} |\tilde{\psi}_{i,s}| \mathbb{E}(|u_{-j}| |u_{-i}|) = O\left(n^{\alpha_0(2d-1)+2}\right). \end{aligned}$$

Therefore, $\max_{1 \leq s \leq n} |E| = O_{\mathbb{P}}(n^{1+\alpha_0(d-1/2)})$, and $\max_{1 \leq s \leq n} |F| = O_{\mathbb{P}}(n^{1+\alpha_0(d-1/2)})$. Finally, it's straightforward to show that $\alpha_0 = 4(1-q)/3$ is the solution of $1 + \alpha_0(d-1/2) = \alpha_0/4 + q + \alpha_0 d$, and hence $\alpha_0 \in (1, 4/3)$. \square

H.4.2 Proof of Proposition A.3

Observe that

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i(e_i^{(d_n)} - e_i) = \sum_{j=1}^L \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i u_{i-j} \psi_j + \sum_{j=L+1}^{\infty} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i u_{i-j} \psi_j := F_1 + F_2,$$

where F_1 and F_2 are defined in the obvious way. We prove the proposition through the following steps:

(i) Show that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| F_1 - \sum_{j=1}^L \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) u_{i-j} \psi_j \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^{-1/2}). \quad (\text{H.39})$$

(ii) The second step is to prove that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| F_2 - \sum_{j=L+1}^{\infty} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) u_{i-j} \psi_j \right| = O_{\mathbb{P}}(\sqrt{n}(\log n)^{-1/2}). \quad (\text{H.40})$$

Step (i) Let $L = \lfloor (\log n)^{1/2} \rfloor$. Then, we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |F_1| \right\| \leq \sum_{j=1}^L \psi_j \left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i u_{i-j} \right| \right\| = O(L d_n \sqrt{n}) = O(\sqrt{n}(\log n)^{-1/2}). \quad (\text{H.41})$$

Similarly, we can show that

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{j=1}^L \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) u_{i-j} \psi_j \right| \right\| = O(L d_n \sqrt{n}) = O(\sqrt{n}(\log n)^{-1/2}). \quad (\text{H.42})$$

From (H.41) and (H.42), we have shown (H.39).

Step (ii) Define $\tilde{e}_{i,L}^{(d_n)} = \sum_{j=L+1}^{\infty} \psi_j u_{i-j}$. We can write

$$F_2 = \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i \tilde{e}_{i,L}^{(d_n)}, \quad \sum_{j=L+1}^{\infty} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) u_{i-j} \psi_j = \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,L}^{(d_n)}.$$

We approximate F_2 following the proof of [Proposition A.1](#). Let $\tilde{m} = \tilde{m} \log n$, then $\tilde{e}_{i,\tilde{m}}^{(d_n)} = \sum_{j=\tilde{m}+1}^{\infty} \psi_j u_{i-j}$. Recall that $\tilde{\mathbf{x}}_{i,\tilde{m}} = \mathbb{E}(\mathbf{x}_i | \varepsilon_i, \dots, \varepsilon_{i-\tilde{m}})$.

Similar to (A.1) and (A.2), we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| F_2 - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)} \right| \right\| = O(\sqrt{n}(\log n)^{-1/2}), \quad (\text{H.43})$$

i.e. we can approximate F_2 by $\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)}$.

Secondly, recall the following decomposition similar to (A.3),

$$\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)} = \sum_{l=0}^{\tilde{m}} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)} \right) + \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)} | \mathcal{F}_{i-\tilde{m}-1}) = T_{1,r} + T_{2,r}. \quad (\text{H.44})$$

We proceed to show that the $T_{1,r}$ is of smaller order of \sqrt{n} , and $T_{2,r}$ approximates $\sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,L}^{(d_n)}$.

(a) Calculation of $T_{1,r}$. Similar to the calculation of (A.4), for $l \leq \tilde{m}$, we have

$$\left\| \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \mathcal{P}_{i-l} \left(\tilde{\mathbf{x}}_{i,\tilde{m}} \tilde{e}_{i,\tilde{m}}^{(d_n)} \right) \right\|^2 \leq \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \left(\|\tilde{\mathbf{x}}_{i,\tilde{m}} - \tilde{\mathbf{x}}_{i,\tilde{m}}^{(i-l)}\|_4 \|\tilde{e}_{i,\tilde{m}}^{(d_n)}\|_4 \right)^2.$$

Notice that uniformly for $\lfloor nb_n \rfloor + 1 \leq i \leq n - \lfloor nb_n \rfloor$, we have

$$\|\tilde{e}_{i,\tilde{m}}^{(d_n)}\|_4^2 = O \left(\sum_{s=\tilde{m}+1}^{\infty} \left\| \mathcal{P}_{i-s} \sum_{j=\tilde{m}+1}^{\infty} \psi_j u_{i-j} \right\|_4^2 \right) = O \left(\sum_{s=\tilde{m}+1}^{\infty} (s+1)^{2d_n-2} \right) = O((\log n)^{-1}),$$

where the second equality is from a careful check of Lemma 3.2 in [Kokoszka and Taqqu \(1995\)](#). Hence, we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |T_{1,r}| \right\| = O(\sqrt{n}(\log n)^{-1/2}). \quad (\text{H.45})$$

(b) Calculation of $T_{2,r}$. Since $\tilde{e}_{i,\tilde{m}}^{(d_n)}$ is $\mathcal{F}_{i-\tilde{m}-1}$ measurable and $\tilde{\mathbf{x}}_{i,\tilde{m}}$ is independent of $\mathcal{F}_{i-\tilde{m}-1}$, we have

$$T_{2,r} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(\tilde{\mathbf{x}}_{i,\tilde{m}} | \mathcal{F}_{i-\tilde{m}-1}) \tilde{e}_{i,\tilde{m}}^{(d_n)} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbb{E}(\tilde{\mathbf{x}}_{i,\tilde{m}}) \tilde{e}_{i,\tilde{m}}^{(d_n)} = \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,\tilde{m}}^{(d_n)}. \quad (\text{H.46})$$

Since $\mathcal{P}_k \cdot = \mathbb{E}(\cdot | \mathcal{F}_k) - \mathbb{E}(\cdot | \mathcal{F}_{k-1})$, $\tilde{e}_{i,L}^{(d_n)} - \tilde{e}_{i,\tilde{m}}^{(d_n)} = \sum_{j=L+1}^{\tilde{m}} \psi_j u_{i-j}$. Similar to (A.6) and by Taylor's expansion, we have

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| T_{2,r} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W(t_i) \tilde{e}_{i,L}^{(d_n)} \right| \right\| = O(\sqrt{n} \tilde{m} d_n / L) = O(\sqrt{n}(\log n)^{-1/2}).$$

Therefore, combining the results in (H.43), (H.44), (H.45) and (H.46), we have proved (H.40). \square

H.5 Proof of [Theorem 4.3](#)

In order to derive [Theorem 4.3](#), we start by investigating some technical lemmas. [Lemma H.2](#) studies the physical dependence of $\mathbf{U}^{(d)}(t, \mathcal{F}_i)$. [Lemma H.3](#) establishes the convergence rate of local linear estimates under the fixed alternative hypothesis. In [Lemma H.4](#), we derive the uniform Gaussian approximation of the partial sum process of nonparametric residuals. [Lemma H.5](#) involves the limiting distribution of a LRD Gaussian process.

H.5.1 Some technical lemmas

Lemma H.2. Assuming $\sup_{t \in (-\infty, 1]} \|H(t, \mathcal{F}_0)\|_{2p} < \infty$, $\sup_{t \in [0, 1]} \|\mathbf{W}(t, \mathcal{F}_0)\|_{2p} < \infty$, $\delta_{2p}(H, k, (-\infty, 1]) = O(\chi^k)$, $\delta_{2p}(\mathbf{W}, k) = O(\chi^k)$, $\chi \in (0, 1)$, we have

$$\delta_p(\mathbf{U}^{(d)}, k) = O(k^{d-1}).$$

Proof. Note that for $j \leq i$,

$$\delta_p(\mathbf{U}^{(d)}, i - j) \leq \|\mathbf{W}(t_i, \mathcal{F}_i)\|_{2p} \delta_{2p}(H^{(d)}, i - j) + \|H^{(d)}(t_i, \mathcal{F}_{i-j}^*)\|_{2p} \delta_{2p}(\mathbf{W}, i - j). \quad (\text{H.47})$$

By Burkholder's inequality and Proposition 3.1, we have

$$\|H^{(d)}(t_i, \mathcal{F}_i)\|_{2p}^2 \leq M \left\| \sum_{j \in \mathbb{Z}} \left(\mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right)^2 \right\|_p \leq M \sum_{j \in \mathbb{Z}} \left\| \mathcal{P}_j H^{(d)}(t_i, \mathcal{F}_i) \right\|_{2p}^2 = O(1),$$

where M is a sufficiently large constant.

Then by Proposition 3.1 and (H.47), we have proved the desired result. \square

Lemma H.3. Under Assumptions 3.1, 4.1, 4.2 and 4.4, $nb_n^2 \rightarrow \infty$ and $b_n \rightarrow 0$, we have

$$\sup_{t \in \mathcal{T}} \left| \tilde{\beta}_{b_n}^{(d)}(t) - \beta(t) - \sum_{i=1}^n \frac{\mathbf{M}^{-1}(t)}{nb_n} \mathbf{x}_i e_i^{(d)} K_{b_n}^*(i/n - t) \right| = O_{\mathbb{P}}(\rho_n^* \chi'_n),$$

where $\mathcal{T} = [b_n, 1 - b_n]$, $\rho_n^* = (nb_n)^{d-1/2} \log n \mathbf{1}(0 \leq d \leq 1/26) + (nb_n)^{d-1/2} b_n^{-1/2} \mathbf{1}(1/26 < d < 1/2) + b_n^2$ and $\chi'_n = n^{-1/2} b_n^{-3/4} + b_n^2$.

Proof. According to Theorem 4.2, when $d \leq 1/26$, take α_0 in Theorem 4.2 to be $31/24 \in (1, 4/3)$, under the bandwidth condition $nb_n^4/(\log n)^2 \rightarrow \infty$, we have $n^{(\alpha_0-1)(\frac{1}{2}-d)} b_n^{d+1/2} \log n \rightarrow \infty$. Therefore, by Theorem 4.2, we obtain

$$\sup_{t \in \mathcal{T}} \left| \frac{1}{nb_n} \sum_{i=1}^n \mathbf{x}_i e_i^{(d)} K_{b_n}(t_i - t) \right| = O_{\mathbb{P}}((nb_n)^{d-1/2} \log n).$$

By Lemma H.2, similar arguments in Remark 4 of Wu (2007) and an application of Proposition B.1 in Dette et al. (2019), we have

$$\sup_{t \in \mathcal{T}} \left| \frac{1}{nb_n} \sum_{i=1}^n \mathbf{x}_i e_i^{(d)} K_{b_n}(t_i - t) \right| = O((nb_n)^{d-1/2} b_n^{-1/2}).$$

The rest of the proof follows from similar procedures in the proof of Theorem 1 in Zhou and Wu (2010). \square

Lemma H.4. Define $G_d^*(r)$ as a counterpart of $G^*(r)$,

$$G_d^*(r) = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{x}_j e_j^{(d)} + \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i^{(d)}.$$

Under the conditions of [Theorem 4.3](#), we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| G_d^*(r) - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} \right| = O_{\mathbb{P}}(\alpha_n)$$

where $\alpha_n = n^d b_n^{-2} \log n + (nb_n)^{d+1/2} \log n + nb_n^3$ when $d \leq 1/26$, $\alpha_n = n^d b_n^{-2} \log n + n^{d+1/2} b_n^d + nb_n^3$, when $1/26 < d < 1/2$. and $\tilde{e}_i^{(d)}$ is the residual under $I(d)$. Under the bandwidth conditions in [Theorem 4.3](#), $\alpha_n = o(n^{d+1/2})$.

Proof. Similar to [\(H.3\)](#), by [Lemma H.3](#), we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| G_d^*(r) - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} \right| \leq \sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \tilde{M}_r^{(d)} \right| + O_{\mathbb{P}}(n \rho_n^* \chi'_n), \quad (\text{H.48})$$

where

$$\tilde{M}_r^{(d)} = \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r (\mathbf{x}_i^\top - \boldsymbol{\mu}_W^\top(t_i)) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{x}_j e_j^{(d)}.$$

Let $\boldsymbol{\xi}_{r,n}(t_j) := \sum_{i=\lfloor nb_n \rfloor + 1}^r (\mathbf{x}_i^\top - \boldsymbol{\mu}_W^\top(t_i)) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j)$ and $\boldsymbol{\xi}_{r,n}(t_0) = 0$, $\sum_{i=1}^0 \mathbf{x}_i e_i^{(d)} = 0$, where $t_j = j/n$. For simplicity, we omit the index of n in $\boldsymbol{\xi}_{r,n}(t_j)$. Using the summation-by-parts formula, it follows that

$$\tilde{M}_r^{(d)} = \frac{1}{nb_n} \boldsymbol{\xi}_r(1) \sum_{j=1}^n \mathbf{x}_j e_j^{(d)} - \frac{1}{nb_n} \sum_{j=1}^n (\boldsymbol{\xi}_r(t_j) - \boldsymbol{\xi}_r(t_{j-1})) \sum_{i=1}^{j-1} \mathbf{x}_i e_i^{(d)} := Z_1 + Z_2, \quad (\text{H.49})$$

where Z_1 and Z_2 are defined in an obvious way. From the proof of Lemma 6 in [Zhou and Wu \(2010\)](#), we have for any $1 \leq j \leq n$,

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\boldsymbol{\xi}_r(t_j)| = O_{\mathbb{P}}(\sqrt{n}). \quad (\text{H.50})$$

From [Proposition A.1](#) and [Proposition A.2](#), we have

$$\max_{1 \leq r \leq n} \left| \sum_{i=1}^r \mathbf{x}_i e_i^{(d)} \right| = O_{\mathbb{P}}(n^{d+1/2} \log n). \quad (\text{H.51})$$

Therefore,

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |Z_1| = \frac{1}{nb_n} \max_{1 \leq r \leq n} |\boldsymbol{\xi}_r(1)| \left| \sum_{j=1}^n \mathbf{x}_j e_j^{(d)} \right| = O_{\mathbb{P}}(b_n^{-1} n^d \log n). \quad (\text{H.52})$$

Under the continuity of $K_{b_n}^*(\cdot)$, by similar arguments of [\(H.50\)](#), we have for any $1 \leq j \leq n$,

$$\left\| \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left\{ \sum_{j=1}^n |\boldsymbol{\xi}_r(t_j) - \boldsymbol{\xi}_r(t_{j-1})| \right\} \right\|_4 = O(n^{1/2} b_n^{-1}). \quad (\text{H.53})$$

Then, by (H.51) and (H.53), we obtain

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |Z_2| = O_{\mathbb{P}}(b_n^{-2} n^d \log n). \quad (\text{H.54})$$

With (H.49), (H.52) and (H.54), it follows that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\tilde{M}_r^{(d)}| = O_{\mathbb{P}}(n^d b_n^{-2} \log n). \quad (\text{H.55})$$

From (H.55) and (H.48), we have shown the desired result. \square

Lemma H.5. *Let v_i be i.i.d. $N(0, 1)$ random variables,*

$$\mathbf{R}_{k,n} = \sum_{j=0}^{\infty} \boldsymbol{\mu}_W(k/n) \psi_j \sigma_H(t_{k-j}) v_{k-j},$$

and $R_{k,n,1}$ is the first element of $\mathbf{R}_{k,n}$. Define

$$\Upsilon_{r,n} = \sum_{i=\lfloor nb_n \rfloor + 1}^r R_{i,n,1} - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \mathbf{R}_{j,n},$$

and $\Upsilon_{r,n} = \Upsilon_{\lfloor nb_n \rfloor, n}$ for $r < \lfloor nb_n \rfloor + 1$, $\Upsilon_{r,n} = \Upsilon_{n - \lfloor nb_n \rfloor, n}$ for $r > n - \lfloor nb_n \rfloor$. Let

$$\Upsilon_n(t) = \Upsilon_{\lfloor nt \rfloor, n} \Gamma(d+1)/n^{d+1/2}.$$

Under the conditions of [Theorem 4.3](#), for $d \in (0, 1/2)$, we have

$$\Upsilon_n(t) \rightsquigarrow U_d(t) \text{ on } D[0, 1] \text{ with Skorohod topology.}$$

where $U_d(t)$ is a continuous Gaussian process in $C[0, 1]$ with covariance function $(0 \leq r \leq s \leq 1)$

$$\gamma_d(r, s) = \int_{-\infty}^{+\infty} \sigma_H^2(v) \lambda_d(r, v) \lambda_d(s, v) dv,$$

where for $v \leq u$, $u \in [0, 1]$,

$$\lambda_d(u, v) = d \int_{(-v)_+}^{(u-v)_+} t^{d-1} (\check{M}_W(t+v) - 1) dt,$$

and $\check{M}_W(t) = \boldsymbol{\mu}_W^\top(t) \mathbf{M}^{-1}(t) \boldsymbol{\mu}_W(t)$, $t \in [0, 1]$.

Proof. The limiting distribution of $\Upsilon_n(t)$ is derived from the follow procedures. Consider $0 \leq t_1 \leq t_2 \leq 1$. Let $r = \lfloor nt_1 \rfloor$, $s = \lfloor nt_2 \rfloor$.

- (i) Calculate the covariance $\mathbb{E} \left\{ \frac{\Upsilon_{r,n} \Upsilon_{s,n}}{n^{2d+1}/\Gamma^2(d+1)} \right\}$ and establish finite dimensional convergence.
- (ii) Prove tightness condition.
- (iii) Show the uniform convergence on $D[0, 1]$.

First, we investigate the terms in $\Upsilon_{r,n}$, when $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$\Upsilon_{r,n} = \sum_{i=\lfloor nb_n \rfloor + 1}^r R_{i,n,1} - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{R}_{j,n}, \quad (\text{H.56})$$

where $\mathbf{m}_{r,j}^\top = \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j)$.

(i) We can write the first term as

$$\sum_{j=\lfloor nb_n \rfloor + 1}^r R_{j,n,1} = \sum_{j=\lfloor nb_n \rfloor + 1}^r \sum_{k=0}^{\infty} \psi_k \sigma_H(t_{j-k}) v_{j-k} = \sum_{l=-\infty}^r \sigma_H(l/n) v_l \sum_{j=(\lfloor nb_n \rfloor + 1 - l)_+}^{r-l} \psi_j.$$

For the second term, let $\check{m}_{r,j} = \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \boldsymbol{\mu}_W(t_j)$, we have

$$\sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{R}_{j,n} = \sum_{j=1}^n \sum_{k=0}^{\infty} \psi_k \check{m}_{r,j} \sigma_H(t_{j-k}) v_{j-k} = \sum_{l=-\infty}^n \sigma_H(l/n) v_l \sum_{j=(1-l)_+ + l}^n \psi_{j-l} \check{m}_{r,j}.$$

Consider $\lfloor nb_n \rfloor + 1 \leq r \leq s \leq n - \lfloor nb_n \rfloor$, the covariance of the Gaussian process is

$$\begin{aligned} \mathbb{E} \left\{ \frac{\Upsilon_{r,n} \Upsilon_{s,n}}{n^{2d+1} / \Gamma^2(d+1)} \right\} &= \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{R}_{j,n} \sum_{j=1}^n \mathbf{m}_{s,j}^\top \mathbf{R}_{j,n} \right\} / (n^{2d+1} / \Gamma^2(d+1)) \\ &- \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{R}_{j,n} \sum_{i=\lfloor nb_n \rfloor + 1}^s R_{i,n,1} \right\} / (n^{2d+1} / \Gamma^2(d+1)) \\ &- \mathbb{E} \left\{ \sum_{j=1}^n \mathbf{m}_{s,j}^\top \mathbf{R}_{j,n} \sum_{i=\lfloor nb_n \rfloor + 1}^r R_{i,n,1} \right\} / (n^{2d+1} / \Gamma^2(d+1)) \\ &+ \mathbb{E} \left\{ \sum_{i=\lfloor nb_n \rfloor + 1}^r R_{i,n,1} \sum_{i=\lfloor nb_n \rfloor + 1}^s R_{i,n,1} \right\} / (n^{2d+1} / \Gamma^2(d+1)) \\ &:= I + II + III + IV. \end{aligned}$$

We first truncate the summands before $l = -N$, $N = \lfloor n^\alpha \rfloor$, $\alpha \geq 1$. By elementary calculation, we have

$$I = \sum_{l=-N}^{r+\lfloor nb_n \rfloor} \sigma_H^2(l/n) \left(\sum_{i=(1-l)_+ + l}^{r+\lfloor nb_n \rfloor} \psi_{i-l} \check{m}_{r,i} \right) \left(\sum_{i=(1-l)_+ + l}^{s+\lfloor nb_n \rfloor} \psi_{i-l} \check{m}_{s,i} \right) / (n^{2d+1} / \Gamma^2(d+1)) + O((N/n)^{2d-1}),$$

where the last equality follows since $\check{m}_{s,j} = 0$ if $i > s + \lfloor nb_n \rfloor$.

Next, define

$$I^* = \frac{d^2}{n} \sum_{l=-N}^r \sigma_H^2(l/n) \int_{(l/n)_+}^{r/n} (t - l/n)^{d-1} \check{M}_W(t) dt \int_{(l/n)_+}^{s/n} (t - l/n)^{d-1} \check{M}_W(t) dt.$$

We approximate $\sum_{i=(1-l)_+ + l}^{r+\lfloor nb_n \rfloor} \psi_{i-l} \check{m}_{r,i}$ by considering different regions of l , namely $|r - l| \leq nb_n \log n$ and $|r - l| > nb_n \log n$, $-l \leq nb_n \log n$, and $-l > nb_n \log n$. Then, we have

$$I = I^* + O(c_n),$$

where $c_n = b_n^{d+1} \log n + N/(n^2 b_n) + N b_n/n + N/n^{d+1} + (N/n)^{2d-1}$. Let

$$I^\diamond = d^2 \int_{-\infty}^{+\infty} \sigma_H^2(v) \int_{(-v)_+}^{(r/n-v)_+} t^{d-1} \check{M}_W(t+v) dt \int_{(-v)_+}^{(s/n-v)_+} t^{d-1} \check{M}_W(t+v) dt dv.$$

Since $\check{M}_W(t)$ is nonnegative and Lipschitz continuous on $[0, 1]$, we have $I = I^\diamond + O(e_n)$, where $e_n = c_n + (N/n^2)^{-\frac{1}{d-2}} + N^{d+1}/n^{d+2}$. The approximation of II-IV follows similarly. Now taking $1 \leq \alpha < \min\{\frac{1}{6}, d, (d+1)^{-1}\} + 1$, elementary calculation shows $e_n = o(1)$. From the continuity of γ_d , we obtain, for $0 \leq t_1, t_2 \leq 1$,

$$\mathbb{E} \left\{ \frac{\Upsilon_{\lfloor nt_1 \rfloor, n} \Upsilon_{\lfloor nt_2 \rfloor, n}}{n^{2d+1}/\Gamma^2(d+1)} \right\} \rightarrow \gamma_d(t_1, t_2), \quad n \rightarrow \infty. \quad (\text{H.57})$$

Similar to the case under the null hypothesis, the finite dimension convergence of the Gaussian process $\Upsilon_n(t)$ to $U_d(t)$ then follows from the Cramer Wold device and equation (H.57).

(ii) To prove the tightness of $\Upsilon_n(t)$, we extend Lemma 2.1 in [Taqqu \(1975\)](#) to the non-stationary case. We verify equation (13.14) and (13.12) in [Billingsley \(1999\)](#). To verify equation (13.14), we need to establish upper bound for

$$J_n(t, t_1, t_2) := \mathbb{E} |\Upsilon_n(t_2) - \Upsilon_n(t)| |\Upsilon_n(t) - \Upsilon_n(t_1)|, \quad 0 \leq t_1 \leq t \leq t_2 \leq 1.$$

By Cauchy-Schwarz inequality, we have

$$J_n(t, t_1, t_2) \leq \frac{\Gamma^2(d+1)}{n^{2d+1}} \|\Upsilon_{\lfloor nt_2 \rfloor, n} - \Upsilon_{\lfloor nt \rfloor, n}\| \|\Upsilon_{\lfloor nt_1 \rfloor, n} - \Upsilon_{\lfloor nt \rfloor, n}\|. \quad (\text{H.58})$$

We proceed to show that, uniformly for $1 \leq r_1 \leq r_2 \leq n$, and $r_1 = \lfloor nt_1 \rfloor$, $r_2 = \lfloor nt_2 \rfloor$, $t_1, t_2 \in [0, 1]$,

$$\|\Upsilon_{r_2, n} - \Upsilon_{r_1, n}\|^2 = O((r_2 - r_1)^{2d+1}). \quad (\text{H.59})$$

According to (H.56), we have

$$\|\Upsilon_{r_2, n} - \Upsilon_{r_1, n}\|^2 \leq 2 \left(\left\| \sum_{i=r_1+1}^{r_2} R_{i, n, 1} \right\|^2 + \left\| \sum_{j=1}^n (\mathbf{m}_{s, j}^\top - \mathbf{m}_{r, j}^\top) \mathbf{R}_{j, n} \right\|^2 \right).$$

Since $\sum_{l=-\infty}^r \left(\sum_{j=r+1-l}^{s-l} \psi_j \right)^2 = O((s-r)^{2d+1})$, (H.59) follows from similar calculation in (H.25) and (H.26).

Then, combining equation (H.58) and (H.59), there exists a sufficiently large positive constant K s.t.

$$J_n(t, t_1, t_2) \leq K^{2d+1} (t_2 - t)^{d+1/2} (t_1 - t)^{d+1/2} \leq (K t_2 - K t_1)^{2d+1}.$$

Hence, equation (13.14) in [Billingsley \(1999\)](#) is satisfied.

We now verify equation (13.12) in Theorem 13.5 in [Billingsley \(1999\)](#). For $d < 1/2$, we have

$$\int_{-\infty}^{\infty} ((t-v)_+^d - (-v)_+^d)^2 dv \leq t^{2d+1} \int_{-\infty}^{\infty} ((1+s)_+^d - (s)_+^d)^2 ds = O(t^{2d+1}).$$

Let $\gamma_d^*(t_1, t_2) = \|U_d(t_2) - U_d(t_1)\|^2$. Note that $\gamma_d^*(t_1, t_2) = \gamma_d(t_1, t_1) + \gamma_d(t_2, t_2) - 2\gamma_d(t_1, t_2)$. Then, since $\check{M}_W(t)$, $\sigma_H^2(t)$ are bounded and continuous on $[0, 1]$ and $(-\infty, 1]$, respectively. For a large constant C , it

follows that

$$\begin{aligned}
\gamma_d^*(t_1, t_2) &= \int_{-\infty}^{\infty} \sigma_H^2(v) (\lambda_d(t_2, v) - \lambda_d(t_1, v))^2 dv \\
&= \int_{-\infty}^{\infty} \sigma_H^2(v) \left\{ (t_2 - v)_+^d - (t_1 - v)_+^d - d \int_{(t_1 - v)_+}^{(t_2 - v)_+} t^{d-1} \check{M}_W(t + v) dt \right\}^2 dv \\
&\leq C \int_{-\infty}^{\infty} ((\tau - v)_+^d - (-v)_+^d)^2 dv = O(\tau^{2d+1}).
\end{aligned} \tag{H.60}$$

For Equation (13.12) in Billingsley (1999), by (H.60), we have

$$\|U_d(t_2) - U_d(t_1)\|^2 = O((t_2 - t_1)^{2d+1}). \tag{H.61}$$

Then, by Chebyshev's inequality, it follows that for any $\epsilon > 0$,

$$\lim_{\delta \rightarrow 0} P[U_d : |U_d(1) - U_d(1 - \delta)| \geq \epsilon] \leq \lim_{\delta \rightarrow 0} \delta^{2d+1} / \epsilon^2 = 0,$$

which satisfies equation (13.12) in Theorem 13.5 in Billingsley (1999).

(iii) From (i), we obtain the finite dimensional convergence. From (ii), we've proved that the $\Upsilon_n(t)$ is tight. Kolmogorov-Chentsov theorem in Karatzas and Shreve (1988) and (H.61) guarantee that the existence of $U_d(t)$ which has a continuous trajectory. Then, $U_d(t) \in C[0, 1] \subset D[0, 1]$. According to Theorem 13.5 in Billingsley (1999), we have

$$\Upsilon_n(t) \rightsquigarrow U_d(t) \text{ on } D[0, 1] \text{ with Skorohod topology.}$$

□

H.5.2 Proof of Theorem 4.3

Recall that $\mathbf{m}_{r,j}^\top = \frac{1}{nb_n} \sum_{i=[nb_n]+1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j)$ as defined in (H.18). Define

$$G_d^*(r) = - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{x}_j e_j^{(d)} + \sum_{i=[nb_n]+1}^r e_i^{(d)}. \tag{H.62}$$

It follows from Lemma H.4 that

$$\max_{[nb_n]+1 \leq r \leq n - [nb_n]} \left| G_d^*(r) - \sum_{i=[nb_n]+1}^r \tilde{e}_i^{(d)} \right| = O_{\mathbb{P}}(\alpha_n),$$

where α_n is of smaller order of $n^{d+1/2}$. According to Theorem 4.2, similar to the proof of (H.18) using the summation-by-parts formula, there exists a series of *i.i.d.* $N(0, 1)$'s $\{v_i\}_{i \in \mathbb{Z}}$ possibly on a richer probability space, such that

$$\max_{[nb_n]+1 \leq r \leq n - [nb_n]} |G_d^*(r) - \Upsilon_{r,n}| = O_{\mathbb{P}}(\sqrt{n}(\log n)^d + n^{1+\alpha_0(d-1/2)}),$$

where

$$\Upsilon_{r,n} = \sum_{i=[nb_n]+1}^r R_{i,n,1} - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{R}_{j,n},$$

and $\mathbf{R}_{k,n} = \sum_{j=0}^{\infty} \boldsymbol{\mu}_W(k/n) \psi_j \sigma(t_{k-j}) v_{k-j}$, $R_{k,n,1}$ is the first element of $\mathbf{R}_{k,n}$. Since by [Lemma H.5](#) $\|\Upsilon_{n,n}\|$ is of order $n^{d+1/2}$, we have

$$\begin{aligned} \left| T_{n,b_n}^{(d)} - \Xi_{n,b_n} \right| &\leq \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \left(\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} \right)^2 - \Upsilon_{r,n}^2 \right| / n \\ &\leq \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} - \Upsilon_{r,n} \right|^2 / n + 2 \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} - \Upsilon_{r,n} \right| |\Upsilon_{r,n}| / n \\ &= O_{\mathbb{P}} \left\{ n^{d-1/2} \left(\alpha_n + \sqrt{n} (\log n)^d + n^{1+\alpha_0(d-1/2)} \right) \right\} = o_{\mathbb{P}}(n^{2d}). \end{aligned}$$

The second part of the proof follows from [Lemma H.5](#) and the continuous mapping theorem. \square

H.6 Proof of [Theorem 4.4](#)

As a counterpart of [\(H.62\)](#), define

$$G_{d_n}^*(r) = - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{x}_j e_j^{(d_n)} + \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i^{(d_n)}.$$

It follows from [Lemma H.4](#) that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| G_{d_n}^*(r) - \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d_n)} \right| = O_{\mathbb{P}}(b_n^{-2} \log n + (nb_n)^{1/2} \log n + nb_n^3),$$

which is of smaller order of $n^{1/2}$ when $nb_n^4/(\log n)^2 \rightarrow \infty$, $nb_n^6 \rightarrow 0$. According to [Theorem 4.2](#), using the summation-by-parts formula, there exists a series of *i.i.d.* Gaussian vectors namely $\{\mathbf{V}_i\}_{i \in \mathbb{Z}}$ possibly on a richer probability space, such that

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |G_{d_n}^*(r) - \Upsilon_{r,n}^\circ| = o_{\mathbb{P}}(\sqrt{n}),$$

where

$$\Upsilon_{r,n}^\circ = \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{R}_{i,n,1} - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \tilde{\mathbf{R}}_{j,n}, \quad \tilde{\mathbf{R}}_{i,n} = \sum_{j=1}^{\infty} \boldsymbol{\mu}_W(t_i) \psi_j \sigma_H(t_{i-j}) V_{i-j,1} + \boldsymbol{\Sigma}^{1/2}(t_i) \mathbf{V}_i := \mathbf{S}_{i,n} + \boldsymbol{\Sigma}^{1/2}(t_i) \mathbf{V}_i.$$

with $\tilde{R}_{i,n,1}$ and $V_{i,1}$ being the first element of $\tilde{\mathbf{R}}_{i,n}$ and \mathbf{V}_i . Extra define $\Upsilon_{r,n}^\circ = \Upsilon_{\lfloor nb_n \rfloor, n}^\circ$ for $r < \lfloor nb_n \rfloor + 1$, $\Upsilon_{r,n}^\circ = \Upsilon_{n - \lfloor nb_n \rfloor, n}^\circ$ for $r > n - \lfloor nb_n \rfloor$. Then, let

$$\Upsilon_n^\circ(t) = n^{-1/2} \Upsilon_{\lfloor nt \rfloor, n}^\circ.$$

Similar to [Lemma H.5](#), the limiting distribution of $\Upsilon_n^\circ(t)$ is derived from the follow procedures. Consider $0 \leq t_1 \leq t_2 \leq 1$. Let $r = \lfloor nt_1 \rfloor$, $s = \lfloor nt_2 \rfloor$.

- (a) Calculate the covariance $n^{-1} \mathbb{E}\{\Upsilon_{r,n}^\circ \Upsilon_{s,n}^\circ\}$ and establish finite dimensional convergence of $\Upsilon_n^\circ(t)$.
- (b) Prove tightness condition of $\Upsilon_n^\circ(t)$.

Then, by (a) and (b), we have

$$\Upsilon_n^\circ(t) \rightsquigarrow U^\circ(t) \text{ on } D[0, 1] \text{ with Skorohod topology.}$$

Step (a). Let $S_{k,n,1}$ be the first element of $\mathbf{S}_{k,n}$. Observe that

$$\Upsilon_{r,n}^\circ = \tilde{G}^*(r) + \check{\Upsilon}_{r,n},$$

where

$$\tilde{G}^*(r) = \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \Sigma^{1/2}(t_j) \mathbf{V}_j, \quad \check{\Upsilon}_{r,n} = \sum_{i=\lfloor nb_n \rfloor + 1}^r S_{i,n,1} - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \mathbf{S}_{j,n}.$$

The covariance of the Gaussian process is

$$n^{-1} \mathbb{E} \{ \Upsilon_{r,n}^\circ \Upsilon_{s,n}^\circ \} = n^{-1} \mathbb{E} \{ (\tilde{G}^*(r) + \check{\Upsilon}_{r,n})(\tilde{G}^*(s) + \check{\Upsilon}_{s,n}) \}. \quad (\text{H.63})$$

Step a.1: Compute $\mathbb{E}\{\tilde{G}^*(s)\tilde{G}^*(r)\}/n$. According to (H.24) of the proof of Theorem 4.1, we have

$$\max_{1 \leq r \leq s \leq n} \left| \mathbb{E}\{\tilde{G}^*(s)\tilde{G}^*(r)\}/n - \gamma(t_1, t_2) \right| = O\left(b_n + \frac{1}{nb_n}\right). \quad (\text{H.64})$$

Step a.2: Compute $\mathbb{E}\{\check{\Upsilon}_{r,n}\check{\Upsilon}_{s,n}\}/n$. Following similar arguments in Lemma H.5 and some tedious calculation, we have

$$\sup_{0 \leq t_1 \leq t_2 \leq 1} \left| n^{-1} \mathbb{E} \{ \check{\Upsilon}_{\lfloor nt_1 \rfloor, n} \check{\Upsilon}_{\lfloor nt_2 \rfloor, n} \} - \tilde{\gamma}(t_1, t_2) \right| = O((\log n)^{-1/2}). \quad (\text{H.65})$$

Step a.3: Compute $\mathbb{E}\{\check{\Upsilon}_{r,n}\tilde{G}^*(s)\}/n$ and $\mathbb{E}\{\check{\Upsilon}_{s,n}\tilde{G}^*(r)\}/n$. Similar to (H.24) of the proof of Theorem 4.1 and some tedious calculation, we have

$$\sup_{0 \leq t_1 \leq t_2 \leq 1} \left| n^{-1} \mathbb{E} \{ \check{\Upsilon}_{\lfloor nt_1 \rfloor, n} \tilde{G}^*(\lfloor nt_2 \rfloor) \} - \tilde{\gamma}(t_1, t_2) \right| = O((\log n)^{-1/2}).$$

Similarly, we have

$$\sup_{0 \leq t_1 \leq t_2 \leq 1} \left| n^{-1} \mathbb{E} \{ \check{\Upsilon}_{\lfloor nt_2 \rfloor, n} \tilde{G}^*(\lfloor nt_1 \rfloor) \} - \tilde{\gamma}(t_1, t_2) \right| = O((\log n)^{-1/2}). \quad (\text{H.66})$$

Combining (H.63), (H.64), (H.65) and (H.66), we have

$$\sup_{0 \leq t_1 \leq t_2 \leq 1} \left| n^{-1} \mathbb{E} \{ \Upsilon_{\lfloor nt_1 \rfloor, n}^\circ \Upsilon_{\lfloor nt_2 \rfloor, n}^\circ \} - \gamma^\circ(t_1, t_2) \right| = O((\log n)^{-1/2}).$$

The finite dimensional convergence of $\Upsilon^\circ(t)$ to $U^\circ(t)$ then follows from Cramer-Wold device.

Step (b). Since $\sum_{l=1}^\infty ((s-r-1+l)^{d_n} - l^{d_n})^2 = o(s-r)$, (13.4) of Theorem 13.5 in Billingsley (1999) with the $\alpha = \beta = 1$ case follows from (H.27), and calculations similar to step (ii) of the proof of Lemma H.5. Equation (13.2) of Billingsley (1999) follows from the continuity of the covariance structure of $U^\circ(t)$. Therefore, by Theorem 13.5 in Billingsley (1999), we have shown the tightness of $\Upsilon^\circ(t)$. \square

I Theoretical properties of bootstrap tests

I.1 Proof of Theorem 5.1

Proof of (i) Under [Assumption 5.1](#) the proof follows from similar but simpler arguments of [Theorem 5.2](#). We omit it for brevity.

Proof of (ii). Given (ii) of [Theorem 4.1](#), it suffices to show that on a possibly richer probability space,

$$|\tilde{T}_n - T_n| = o_{\mathbb{P}}(b_n). \quad (\text{I.1})$$

Let \tilde{G}_r be the bootstrap statistic in one iteration defined in (5.1) in one simulation round. To prove (I.1), it's sufficient to show that on a possibly richer probability space,

$$\sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \left(\sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i - \tilde{G}_r \right) \right| = o_{\mathbb{P}}(\sqrt{nb_n}/\log n). \quad (\text{I.2})$$

and

$$\sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} |\tilde{G}_r| = O_{\mathbb{P}}(\sqrt{nb_n \log n}). \quad (\text{I.3})$$

Then, the theorem follows from continuous mapping theorem. Recall in Step 2 of [Theorem 4.1](#),

$$\tilde{G}^*(r) = - \sum_{j=1}^n \left(\frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j) \right) \boldsymbol{\Sigma}^{1/2}(t_j) \mathbf{V}_j + \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1}.$$

Recall in (H.29), we have

$$\sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i - \tilde{G}^*(r) \right| = o_{\mathbb{P}}(\sqrt{nb_n}/\log n). \quad (\text{I.4})$$

Following the proof of (I.12), we have

$$\begin{aligned} \sup_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| n^{-1/2}(\tilde{G}_r - \tilde{G}^*(r)) \right| &= O_{\mathbb{P}} \left(q_n \left(r_n + g_n^{1/2} \right) + n^{-1/2} b_n^{-3/2} \right) \\ &= o_{\mathbb{P}}((b_n^{1/2}/\log n)), \end{aligned} \quad (\text{I.5})$$

where $r_n = n^{-1/2} b_n^{-3/4} + b_n^2$, q_n is a sequence goes to infinity at an arbitrarily slow rate, g_n is related to the convergence rate of [Assumption 5.1](#) which is $o(b_n/\log^2 n)$. Therefore, [Equation \(I.2\)](#) is proved by (I.4) and (I.5). Recall that in (H.30), we have

$$\|\tilde{G}^*(r) - \tilde{G}^\circ(r)\|^2 = O(nb_n^3),$$

where $\tilde{G}^\circ(r) = \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_H(t_i) V_{i,1} - \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r \frac{K_{b_n}^*(t_i - t_j)}{nb_n} \sigma_H(t_j) V_{j,1}$, and for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$\|\tilde{G}^\circ(r)\|^2 = O(nb_n).$$

Since $\tilde{G}^*(r) - \tilde{G}^\circ(r)$ and $\tilde{G}^\circ(r)$ are Gaussian processes, [Equation \(I.3\)](#) is proved. \square

Remark I.1. The bootstrap consistency of V/S-type test follows similarly. In the following remark, we take K/S-type test as an example, the result of R/S-type test follows similarly. Let

$$Y = (|\tilde{G}^*(3\lfloor nb_n \rfloor + 1), \dots, \tilde{G}^*(n - \lfloor nb_n \rfloor)|)/\sqrt{nb_n},$$

and

$$\widetilde{\text{KS}}_n = \max_{3\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} |\tilde{G}_k|, \quad \text{KS}_n = \max_{3\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} |\tilde{G}_k|, \quad \widetilde{\text{KS}}_n^* = \max_{3\lfloor nb_n \rfloor + 1 \leq k \leq n - \lfloor nb_n \rfloor} |\tilde{G}_k^*|.$$

Since by elementary calculation using (H.30), (H.31) and degeneracy there exist positive constants c_1 and c_2 such that for $3\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$, $c_1 \leq \|\tilde{G}^*(r)\|^2/(nb_n) \leq c_2$, by Lemma C.1 in Dette and Wu (2021), we have

$$\begin{aligned} \sup_{x \in \mathbb{R}} \left| P(|Y|_\infty > x) - P(\text{KS}_n/\sqrt{nb_n} > x) \right| &\leq \sup_{x \in \mathbb{R}} \left| P(|Y|_\infty > x) - P(\widetilde{\text{KS}}_n/\sqrt{nb_n} > x) \right| \\ &\quad + P(|\widetilde{\text{KS}}_n - \text{KS}_n| > \sqrt{nb_n}\delta) + c\delta\sqrt{\max\{1, \log(n/\delta)\}}, \end{aligned}$$

where the right-hand side converges to 0 as $n \rightarrow \infty$, if we let $\delta = \log^{-1} n$, and use (I.2) and (I.5).

I.2 Proof of Theorem 5.2

Recall that $\mathbf{m}_{r,j}^\top = \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \boldsymbol{\mu}_W^\top(t_i) \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j)$. Define $\tilde{T}_{n,m}^* = \sum_{r=\lfloor nb_n \rfloor + 1}^{n-\lfloor nb_n \rfloor} (\tilde{G}_{r,d}^*)^2 / (n(n - 2\lfloor nb_n \rfloor))$, where

$$\tilde{G}_{r,d}^* = - \sum_{j=1}^n \mathbf{m}_{r,j}^\top \boldsymbol{\Sigma}_d^{1/2}(t_j) \mathbf{V}_j + \sum_{i=\lfloor nb_n \rfloor + 1}^r \sigma_{Hd}(t_i) V_{i,1},$$

with $\boldsymbol{\Sigma}_d(t) = \kappa_2(d) \sigma_H^2(t) \boldsymbol{\mu}_W(t) \boldsymbol{\mu}_W^\top(t)$, $\sigma_{Hd}(t) = (\boldsymbol{\Sigma}_d(t))_{(1,1)}$.

The proof consists of three parts.

- (a) Obtain the limiting distribution of $n^{-1/2} \tilde{G}_{r,d}^*$, $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$.
- (b) Show that conditional on data, $m^{-d} n^{-1/2} \tilde{G}_{r,d}$ and $n^{-1/2} \tilde{G}_{r,d}^*$ converge to the same limit, uniformly for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$.
- (c) Derive the limiting distribution of \tilde{T}_n .

Step (a). By similar arguments as (H.20), (H.21), (H.21), and (H.23) in the proof of Theorem 4.1, by Cramer-Wold device, we have the finite dimensional convergence. The tightness follows similarly as in the final part of the proof of Theorem 4.1. Then, we have $n^{-1/2} \tilde{G}_{r,d}^* \rightsquigarrow \tilde{U}_d(t)$ on $D[0, 1]$ with Skorohod topology.

Step of (b). Let $\mathbf{1}$ denote the indicator function. Let C below denote a sufficiently large constant in the following context. We construct sets independent of $\{\mathbf{V}_j\}_{j=1}^n$ as follows. Let q_n be a sequence of real numbers so that $q_n \rightarrow \infty$ arbitrarily slow. Define

$$W_n := \{\sup_{t \in \mathcal{I}} |\hat{\boldsymbol{\Sigma}}_d^{1/2}(t) m^{-d} - \boldsymbol{\Sigma}_d^{1/2}(t)| \leq g_{1,n}^{1/2} q_n\}, \quad H_n := \{\sup_{t \in \mathcal{B}} \rho(\hat{\mathbf{M}}^{-1}(t) - \mathbf{M}^{-1}(t)) \leq r_n q_n\},$$

where $\mathcal{I} = [\gamma_n, 1 - \gamma_n] \subset (0, 1)$, $\gamma_n = \tau_n + (m+1)/n$, $\mathcal{B} = [\eta_n, 1 - \eta_n]$, $g_{1,n} = o(1)$ related to the convergence rate of Assumption 5.1, $r_n = n^{-1/2} \eta_n^{-3/4} + \eta_n^2$.

Under Assumption 5.1, by Gershgorin's circle theorem and Corollary 1 in Yu et al. (2015), we obtain

$$\lim_{n \rightarrow \infty} \mathbb{P}(W_n) = 1, \quad \lim_{n \rightarrow \infty} \mathbb{P}(H_n) = 1. \quad (\text{I.6})$$

Observe that

$$\begin{aligned}
(m^{-d}\tilde{G}_{r,d} - \tilde{G}_{r,d}^*)\mathbf{1}(W_n \cap H_n) &= \sum_{j=1}^n \left\{ \mathbf{m}_{r,j}^\top \Sigma_d^{1/2}(t_j) - \hat{\mathbf{m}}_{r,j}^\top m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \right\} \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \\
&+ \sum_{i=1}^r (\hat{\sigma}_{Hd}(t_j) m^{-d} - \sigma_{Hd}(t_j)) \mathbf{1}(W_n \cap H_n) V_{i,1} \\
&= \sum_{j=1}^n \left\{ (\mathbf{m}_{r,j}^\top - \hat{\mathbf{m}}_{r,j}^\top) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \right\} \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \\
&+ \left\{ \sum_{j=1}^n \mathbf{m}_{r,j}^\top \left(\Sigma_d^{1/2}(t_j) - m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \right) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right. \\
&\left. + \sum_{i=1}^r (\hat{\sigma}_{Hd}(t_j) m^{-d} - \sigma_{Hd}(t_j)) \mathbf{1}(W_n \cap H_n) V_{i,1} \right\} := J_1 + J_2,
\end{aligned}$$

where J_1 and J_2 are defined in the obvious way. Let $\mathbb{T}_n = [\lfloor n\gamma_n \rfloor + 1, n - \lfloor n\gamma_n \rfloor] \cap [\lfloor n\eta_n \rfloor + 1, n - \lfloor n\eta_n \rfloor]$, and consider $r \in \mathbb{T}_n$. First, we show calculate J_1 by the following two steps. Define

$$\tilde{\mathbf{m}}_{r,j}^\top = \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i^\top \mathbf{M}^{-1}(t_i) K_{b_n}^*(t_i - t_j).$$

(1) We shall show that

$$\left\| \max_{r \in \mathbb{T}_n} \sum_{j=1}^n \left(\hat{\mathbf{m}}_{r,j}^\top - \tilde{\mathbf{m}}_{r,j}^\top \right) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right\| = O(n^{1/2} r_n q_n). \quad (\text{I.7})$$

(2) Then, we shall derive

$$\left\| \max_{r \in \mathbb{T}_n} \sum_{j=1}^n \left(\tilde{\mathbf{m}}_{r,j}^\top - \mathbf{m}_{r,j}^\top \right) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right\| = O(b_n^{-3/2}). \quad (\text{I.8})$$

Proof of (I.7). Observe that the left hand side of (I.7) is a martingale w.r.t $\mathcal{G}_r = \{\mathcal{F}_n, \{V_{i,1}\}_{i=1}^r\}$. By Doob's inequality and Burkholder inequality, for a sufficiently large constant C , we have

$$\begin{aligned}
&\left\| \max_{r \in \mathbb{T}_n} \sum_{j=1}^n \left(\hat{\mathbf{m}}_{r,j}^\top - \tilde{\mathbf{m}}_{r,j}^\top \right) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right\| \\
&= \left\| \max_{r \in \mathbb{T}_n} \left| \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^r \mathbf{x}_i^\top (\hat{\mathbf{M}}^{-1}(t_i) - \mathbf{M}^{-1}(t_i)) \sum_{j=1}^n K_{b_n}^*(t_i - t_j) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right| \right\| \\
&\leq C \left\| \frac{1}{nb_n} \sum_{i=\lfloor nb_n \rfloor + 1}^{n - \lfloor nb_n \rfloor} \mathbf{x}_i^\top (\hat{\mathbf{M}}^{-1}(t_i) - \mathbf{M}^{-1}(t_i)) \sum_{j=1}^n K_{b_n}^*(t_i - t_j) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j \right\| \\
&\leq C \left\{ \sum_{j=1}^n \left\| \left(\hat{\mathbf{m}}_{n,j}^\top - \tilde{\mathbf{m}}_{n,j}^\top \right) m^{-d} \hat{\Sigma}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \right\|^2 \right\}^{1/2} \\
&= O(n^{1/2} r_n q_n).
\end{aligned}$$

Proof of (I.8). Let $\boldsymbol{\mu}_{b_n}^\dagger(t) = \frac{1}{nb_n} \sum_{j=1}^n K_{b_n}^*(t - t_j) m^{-d} \hat{\boldsymbol{\Sigma}}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j$. By summation-by-parts formula,

$$\begin{aligned} \sum_{j=1}^n \left(\tilde{\mathbf{m}}_{r,j}^\top - \mathbf{m}_{r,j}^\top \right) m^{-d} \hat{\boldsymbol{\Sigma}}_d^{1/2}(t_j) \mathbf{1}(W_n \cap H_n) \mathbf{V}_j &= \sum_{i=1}^r (\mathbf{x}_i - \boldsymbol{\mu}_i)^\top \mathbf{M}^{-1}(t_i) \boldsymbol{\mu}_{b_n}^\dagger(r/n) \\ &= \sum_{i=1}^r (\mathbf{x}_i - \boldsymbol{\mu}_i)^\top \mathbf{M}^{-1}(t_i) \boldsymbol{\mu}_{b_n}^\dagger(r/n) \\ &\quad - \sum_{i=1}^r (\boldsymbol{\mu}_{b_n}^\dagger(t_i) - \boldsymbol{\mu}_{b_n}^\dagger(t_{i-1})) \sum_{k=1}^{r-1} (\mathbf{x}_i - \boldsymbol{\mu}_i)^\top \mathbf{M}^{-1}(t_i) \\ &:= Z_1 + Z_2 \end{aligned}$$

Under condition (B3) and (B1), similar to proof of (I.7), by Doob's inequality and Cauchy inequality, we have

$$\left\| \max_{r \in \mathbb{T}_n} |Z_1| \right\| \leq \left\| \left\| \sum_{i=1}^{n - \lfloor nb_n \rfloor} (\mathbf{x}_i - \boldsymbol{\mu}_i)^\top \mathbf{M}^{-1}(t_i) \right\|_4 \left\| \boldsymbol{\mu}_{b_n}^\dagger((n - \lfloor nb_n \rfloor)/n) \right\|_4 \right\| = O(b_n^{-1/2}), \quad (\text{I.9})$$

and by Cauchy inequality, we have

$$\left\| \max_{r \in \mathbb{T}_n} |Z_2| \right\| \leq \left\| \max_{r \in \mathbb{T}_n} \left\| \sum_{i=1}^r (\mathbf{x}_i - \boldsymbol{\mu}_i)^\top \mathbf{M}^{-1}(t_i) \right\|_4 \left\| \max_{r \in \mathbb{T}_n} \sum_{i=1}^r \left\| \boldsymbol{\mu}_{b_n}^\dagger(t_i) - \boldsymbol{\mu}_{b_n}^\dagger(t_{i-1}) \right\|_4 \right\| = O(b_n^{-3/2}), \quad (\text{I.10})$$

where the last equality follows from Doob's inequality, triangle and Burkholder inequality. By (I.9) and (I.10), (I.8) is proved. Similar to the calculation of J_1 , we have

$$\left\| \max_{r \in \mathbb{T}_n} |J_2| \right\| = O(n^{1/2} g_{1,n} q_n). \quad (\text{I.11})$$

Therefore, combining (I.7), (I.8), and (I.11), by triangle inequality,

$$\left\| \max_{r \in \mathbb{T}_n} \left| n^{-1/2} (m^{-d} \tilde{G}_{r,d} - \tilde{G}_{r,d}^*) \mathbf{1}(W_n \cap H_n) \right| \right\| = O(g_{1,n}^{1/2} q_n + r_n q_n + n^{-1/2} b_n^{-3/2}).$$

By Proposition A.1 in Wu and Zhou (2018a), since (I.6), we have

$$\max_{r \in \mathbb{T}_n} \left| n^{-1/2} (m^{-d} \tilde{G}_{r,d} - \tilde{G}_{r,d}^*) \right| = O_{\mathbb{P}}(g_{1,n}^{1/2} q_n + r_n q_n + n^{-1/2} b_n^{-3/2}). \quad (\text{I.12})$$

Step (c) Under the bandwidth condition $nb_n^3 \rightarrow \infty$, by Step (a) and (b), $n^{-1/2} m^{-d} \tilde{G}_{r,d} \rightsquigarrow \tilde{U}_d(t)$ on $D[0, 1]$ with Skorohod topology. Therefore, by continuous mapping theorem, we have

$$m^{-2d} \tilde{T}_n \Rightarrow \int_0^1 \tilde{U}_d^2(t) dt.$$

Under Assumption 5.1, following similar arguments in the proof of result (i) of Theorem 5.2, we have result (ii). \square

I.3 Proof of Proposition 5.1

Proof of (i). Recall $\tilde{G}_{r,d}$ is the bootstrap statistic (5.1) in one iteration under the fixed alternative. It's sufficient to show that,

$$\lim_{n \rightarrow \infty} P \left(\left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} \right| > |\tilde{G}_{r,d}| \text{ for } r = \lfloor nb_n \rfloor + 1, \dots, n - \lfloor nb_n \rfloor \right) = 1.$$

Similar to the steps in the proof of (ii) of Theorem 5.1, substituting $\sigma_H(\cdot)$ by $\hat{\sigma}_d(\cdot)$, using law of total expectation, we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \|\tilde{G}_{r,d}\|^2 = O(nb_n m^{2d}). \quad (\text{I.13})$$

Similar to Lemma H.3, for the time varying trend model,

$$\sup_{t \in \mathcal{T}} \left| \tilde{\beta}_1^{(d)}(t) - \beta_1(t) - \frac{1}{nb_n} \sum_{i=1}^n e_i^{(d)} K_{b_n}^*(i/n - t) \right| = O_{\mathbb{P}}(\rho_n^*/(nb_n)),$$

where $\mathcal{T} = [b_n, 1 - b_n]$, $\rho_n^* = (nb_n)^{d-1/2} \log n \mathbf{1}(0 \leq d \leq 1/26) + (nb_n)^{d-1/2} b_n^{-1/2} \mathbf{1}(1/26 < d < 1/2) + b_n^2$. Then, it follows that

$$\begin{aligned} & \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d)} - \sum_{i=\lfloor nb_n \rfloor + 1}^r \left(e_i^{(d)} - \frac{1}{nb_n} \sum_{j=1}^n e_j^{(d)} K_{b_n}^*(i/n - j/n) \right) \right| \\ &= O_{\mathbb{P}}(\rho_n^*/b_n) = o_{\mathbb{P}}((nb_n)^{1/2+d}). \end{aligned} \quad (\text{I.14})$$

A careful investigation of Proposition A.2 yields when $nb_n^3 \rightarrow \infty$,

$$\begin{aligned} & \max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \max_{r - \lfloor nb_n \rfloor \leq s \leq r + \lfloor nb_n \rfloor} \left| \sum_{k=r - \lfloor nb_n \rfloor}^s (e_k^{(d)} - R_{k,n,1}) \right| = O_{\mathbb{P}}(n^{\alpha_0(d-1/2)+1} b_n + n^{\alpha_0/4+d+q} b_n^d) \\ &= o_{\mathbb{P}}((nb_n)^{d+1/2}), \end{aligned} \quad (\text{I.15})$$

where $\alpha_0 \in (1, 4/3)$, $q > 0$ can be arbitrarily small. Observe that by (H.31), uniformly for $3\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$,

$$\begin{aligned} & \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i^{(d)} - \frac{1}{nb_n} \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r e_j^{(d)} K_{b_n}^*(i/n - j/n) \\ &= \sum_{j=\lfloor nb_n \rfloor + 1}^{2\lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{j - \lfloor nb_n \rfloor}{nb_n}} K^*(t) dt \right) e_j^{(d)} + \sum_{j=r - \lfloor nb_n \rfloor + 1}^r \left(\int_{\frac{r-j}{nb_n}}^1 K^*(t) dt \right) e_j^{(d)} \\ &- \sum_{j=1}^{\lfloor nb_n \rfloor} \left(\int_{\frac{j - \lfloor nb_n \rfloor}{nb_n}}^1 K^*(t) dt \right) e_j^{(d)} - \sum_{j=r+1}^{r+\lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{r-j}{nb_n}} K^*(t) dt \right) e_j^{(d)} + O_{\mathbb{P}}(b_n^{-1}), \end{aligned} \quad (\text{I.16})$$

and similarly uniformly for $\lfloor nb_n \rfloor + 1 \leq r \leq 3\lfloor nb_n \rfloor + 1$,

$$\begin{aligned}
& \sum_{i=\lfloor nb_n \rfloor + 1}^r e_i^{(d)} - \frac{1}{nb_n} \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r e_j^{(d)} K_{b_n}^*(i/n - j/n) \\
&= \sum_{j=\lfloor nb_n \rfloor + 1}^{r-\lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{j-\lfloor nb_n \rfloor}{nb_n}} K^*(t) dt \right) e_j^{(d)} + \sum_{j=r-\lfloor nb_n \rfloor + 1}^r \left(\int_{\frac{r-j}{nb_n}}^1 K^*(t) dt \right) e_j^{(d)} \\
&- \sum_{j=1}^{\lfloor nb_n \rfloor} \left(\int_{\frac{j-\lfloor nb_n \rfloor}{nb_n}}^1 K^*(t) dt \right) e_j^{(d)} - \sum_{j=r+1}^{r+\lfloor nb_n \rfloor} \left(\int_{-1}^{\frac{r-j}{nb_n}} K^*(t) dt \right) e_j^{(d)} + O_{\mathbb{P}}(b_n^{-1}). \tag{I.17}
\end{aligned}$$

Combining (I.15), (I.16) and (I.17), by summation-by-parts formula, we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \left| \sum_{i=\lfloor nb_n \rfloor + 1}^r (e_i^{(d)} - R_{i,n,1}) - \frac{1}{nb_n} \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r K_{b_n}^*(i/n - j/n) (e_j^{(d)} - R_{j,n,1}) \right| = o_{\mathbb{P}}((nb_n)^{d+1/2}). \tag{I.18}$$

Using similar technique in (I.16), it follows from tedious calculation that for $r \geq 2\lfloor nb_n \rfloor$, there exists a positive constant c' such that

$$\left\| \sum_{i=\lfloor nb_n \rfloor + 1}^r R_{i,n,1} - \frac{1}{nb_n} \sum_{j=1}^n \sum_{i=\lfloor nb_n \rfloor + 1}^r K_{b_n}^*(i/n - j/n) R_{j,n,1} \right\|^2 \geq c' (nb_n)^{2d+1}. \tag{I.19}$$

Finally, since $(nb_n)/m \rightarrow \infty$, (i) of proposition follows from (I.13), (I.14), (I.18), and (I.19).

Proof of (ii). Similar to the steps in (I.13), we have

$$\max_{\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor} \|\tilde{G}_{r,d}\|^2 = O(nb_n m^{2d_n}),$$

and uniformly for $\lfloor nb_n \rfloor + 1 \leq r \leq n - \lfloor nb_n \rfloor$, there exists a constant c'' ,

$$\left\| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d_n)} \right\| \geq c'' (nb_n)^{d_n+1/2}.$$

Since $m^{2d_n} = e^{2c\alpha_1}$, and $(nb_n)^{d_n} = e^{2c\beta}$, $\beta > \alpha_1$, when c is sufficiently large, we have

$$\lim_{n \rightarrow \infty} P \left(\left| \sum_{i=\lfloor nb_n \rfloor + 1}^r \tilde{e}_i^{(d_n)} \right| > |\tilde{G}_{r,d_n}| \text{ for } r = \lfloor nb_n \rfloor + 1, \dots, n - \lfloor nb_n \rfloor \right) = 1.$$

□

References

- Bai, L. and Wu, W. (2023). Supplement to “difference based variance estimate in time series nonparametric regression with applications to specification tests”. Manuscript.
- Beran, J. (2009). On parameter estimation for locally stationary long-memory processes. *Journal of Statistical Planning and Inference*, 139(3):900 – 915.

- Beran, J., Feng, Y., Ghosh, S., and Kulik, R. (2013). *Long-Memory Processes*. Springer.
- Berkes, I., Horváth, L., Kokoszka, P., and Shao, Q.-M. (2006). On discriminating between long-range dependence and changes in mean. *The Annals of Statistics*, 34(3):1140 – 1165.
- Billingsley, P. (1999). *Convergence of probability measures*. Wiley.
- Caporale, G. M. and Gil-Alana, L. A. (2013). Long memory and fractional integration in high frequency data on the us dollar/british pound spot exchange rate. *International review of financial analysis*, 29:1–9.
- Caporale, G. M., Gil-Alana, L. A., and Lovcha, Y. (2016). Testing unemployment theories: A multivariate long memory approach. *Journal of Applied Economics*, 19(1):95–112.
- Cavaliere, G., Ørregaard Nielsen, M., and Taylor, A. M. R. (2020). Adaptive inference in heteroscedastic fractional time series models. *Journal of Business & Economic Statistics*, 0(0):1–16.
- Chen, M. and Song, Q. (2015). Simultaneous inference of the mean of functional time series. *Electronic Journal of Statistics*, 9(2):1779 – 1798.
- Chen, X. B., Gao, J., Li, D., and Silvapulle, P. (2018). Nonparametric estimation and forecasting for time-varying coefficient realized volatility models. *Journal of Business & Economic Statistics*, 36(1):88–100.
- Cheung, Y.-W. and Lai, K. S. (1993). Do gold market returns have long memory? *Financial Review*, 28(2):181–202.
- Craven, P. and Wahba, G. (1978). Smoothing noisy data with spline functions. *Numerische mathematik*, 31(4):377–403.
- Dahlhaus, R. (1997). Fitting time series models to nonstationary processes. *Ann. Statist.*, 25(1):1–37.
- Dahlhaus, R., Richter, S., and Wu, W. B. (2019). Towards a general theory for nonlinear locally stationary processes. *Bernoulli*, 25(2):1013 – 1044.
- Davis, R. A. and Yau, C. Y. (2013). Consistency of minimum description length model selection for piecewise stationary time series models. *Electronic Journal of Statistics*, 7(none):381 – 411.
- Dehling, H. and Taqqu, M. S. (1989). The Empirical Process of some Long-Range Dependent Sequences with an Application to U -Statistics. *The Annals of Statistics*, 17(4):1767 – 1783.
- Dette, H., Preuss, P., and Sen, K. (2017). Detecting long-range dependence in non-stationary time series. *Electronic Journal of Statistics*, 11(1):1600 – 1659.
- Dette, H. and Wu, W. (2019). Detecting relevant changes in the mean of nonstationary processes—a mass excess approach. *The Annals of Statistics*, 47(6):3578–3608.
- Dette, H. and Wu, W. (2021). Confidence surfaces for the mean of locally stationary functional time series. *arXiv preprint arXiv:2109.03641*.
- Dette, H., Wu, W., and Zhou, Z. (2019). Change point analysis of correlation in non-stationary time series. *Statistica Sinica*, 29(2):611–643.
- Duffy, J. A. and Kasparis, I. (2021). Estimation and inference in the presence of fractional $d = 1/2$ and weakly nonstationary processes. *The Annals of Statistics*, 49(2):1195 – 1217.
- Fan, J. (1993). Local linear regression smoothers and their minimax efficiencies. *The annals of Statistics*, 21(1):196–216.
- Fan, J. and Gijbels, I. (1996). *Local polynomial modelling and its applications*. Number 66 in Monographs on statistics and applied probability series. Chapman & Hall, London [u.a.].

- Fan, J. and Zhang, W. (2000). Simultaneous confidence bands and hypothesis testing in varying-coefficient models. *Scandinavian Journal of Statistics*, 27(4):715–731.
- Ferreira, G., Piña, N., and Porcu, E. (2018). Estimation of slowly time-varying trend function in long memory regression models. *Journal of Statistical Computation and Simulation*, 88(10):1903–1920.
- Giraitis, L., Kokoszka, P., and Leipus, R. (2001). Testing for long memory in the presence of a general trend. *Journal of Applied Probability*, 38(4):1033–1054.
- Giraitis, L., Kokoszka, P., Leipus, R., and Teyssière, G. (2003). Rescaled variance and related tests for long memory in volatility and levels. *Journal of econometrics*, 112(2):265–294.
- Harris, D. and Kew, H. (2017). Adaptive long memory testing under heteroskedasticity. *Econometric Theory*, 33(3):755–778.
- Harris, D., McCabe, B., and Leybourne, S. (2008). Testing for long memory. *Econometric Theory*, 24(1):143–175.
- Hu, L., Huang, T., and You, J. (2019). Estimation and identification of a varying-coefficient additive model for locally stationary processes. *Journal of the American Statistical Association*, 114(527):1191–1204.
- Hurst, H. E. (1951). Long-term storage capacity of reservoirs. *Trans. Amer. Soc. Civil Eng.*, 116:770–799.
- Jiang, F., Zhao, Z., and Shao, X. (2020). Time series analysis of covid-19 infection curve: A change-point perspective. *Journal of econometrics*.
- Karatzas, I. and Shreve, S. E. (1988). Brownian motion. In *Brownian Motion and Stochastic Calculus*, pages 47–127. Springer.
- Kokoszka, P. and Young, G. (2016). Kpss test for functional time series. *Statistics*, 50(5):957–973.
- Kokoszka, P. S. and Taqqu, M. S. (1995). Fractional arima with stable innovations. *Stochastic Processes and their Applications*, 60(1):19 – 47.
- Koutsoyiannis, D. (2013). Hydrology and change. *Hydrological Sciences Journal*, 58(6):1177–1197.
- Kulik, R. and Wichelhaus, C. (2012). Conditional variance estimation in regression models with long memory. *Journal of Time Series Analysis*, 33(3):468–483.
- Kwiatkowski, D., Phillips, P. C., Schmidt, P., and Shin, Y. (1992). Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of econometrics*, 54(1-3):159–178.
- Lee, D. and Schmidt, P. (1996). On the power of the kpss test of stationarity against fractionally-integrated alternatives. *Journal of econometrics*, 73(1):285–302.
- Lima, L. and Xiao, Z. (2004). Robustness of stationary tests under long-memory alternatives. Technical report, EPGE Brazilian School of Economics and Finance-FGV EPGE (Brazil).
- Lo, A. W. (1989). Long-term memory in stock market prices. Technical report, National Bureau of Economic Research.
- MacNeill, I. B. (1974). Tests for change of parameter at unknown times and distributions of some related functionals on brownian motion. *The Annals of Statistics*, 2(5):950–962.
- Marinucci, D. and Robinson, P. M. (1999). Alternative forms of fractional brownian motion. *Journal of statistical planning and inference*, 80(1-2):111–122.
- McCloskey, A. and Perron, P. (2013). Memory parameter estimation in the presence of level shifts and deterministic trends. *Econometric Theory*, 29(6):1196–1237.

- Nason, G. P., Von Sachs, R., and Kroisandt, G. (2000). Wavelet processes and adaptive estimation of the evolutionary wavelet spectrum. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 62(2):271–292.
- Pipiras, V. and Taqqu, M. S. (2017). *Long-range dependence and self-similarity*, volume 45. Cambridge university press.
- Politis, D. N., Romano, J. P., and Wolf, M. (1999). *Subsampling*. Springer Science & Business Media.
- Preuß, P. and Vetter, M. (2013). Discriminating between long-range dependence and non-stationarity. *Electron. J. Statist.*, 7:2241–2297.
- Qu, Z. (2011). A test against spurious long memory. *Journal of Business & Economic Statistics*, 29(3):423–438.
- Shao, X. and Wu, W. B. (2007). Local asymptotic powers of nonparametric and semiparametric tests for fractional integration. *Stochastic processes and their Applications*, 117(2):251–261.
- Sibbertsen, P., Leschinski, C., and Busch, M. (2018). A multivariate test against spurious long memory. *Journal of Econometrics*, 203(1):33–49.
- Taqqu, M. S. (1975). Weak convergence to fractional brownian motion and to the rosenblatt process. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 31(4):287–302.
- Vogt, M. (2012). Nonparametric regression for locally stationary time series. *Ann. Statist.*, 40(5):2601–2633.
- Vogt, M. and Dette, H. (2015). Detecting gradual changes in locally stationary processes. *The Annals of Statistics*, 43(2):713–740.
- Wu, W. and Zhou, Z. (2018a). Gradient-based structural change detection for nonstationary time series M-estimation. *The Annals of Statistics*, 46(3):1197 – 1224.
- Wu, W. and Zhou, Z. (2018b). Simultaneous quantile inference for non-stationary long-memory time series. *Bernoulli*, 24(4A):2991–3012.
- Wu, W. B. (2005). Nonlinear system theory: Another look at dependence. *Proceedings of the National Academy of Sciences*, 102(40):14150–14154.
- Wu, W. B. (2007). Strong invariance principles for dependent random variables. *The Annals of Probability*, 35(6):2294–2320.
- Wu, W. B. and Shao, X. (2006). Invariance principles for fractionally integrated nonlinear processes. In *Recent developments in nonparametric inference and probability*, pages 20–30. Institute of Mathematical Statistics.
- Wu, W. B. and Zhao, Z. (2007). Inference of trends in time series. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(3):391–410.
- Wu, W. B. and Zhou, Z. (2011). Gaussian approximations for non-stationary multiple time series. *Statistica Sinica*, 21(3):1397–1413.
- Yu, Y., Wang, T., and Samworth, R. J. (2015). A useful variant of the davis–kahan theorem for statisticians. *Biometrika*, 102(2):315–323.
- Zhang, Q., Zhou, Y., Singh, V. P., and Chen, Y. D. (2011). Comparison of detrending methods for fluctuation analysis in hydrology. *Journal of Hydrology*, 400(1):121–132.
- Zhou, Z. and Wu, W. B. (2009). Local linear quantile estimation for nonstationary time series. *The Annals of Statistics*, 37(5B):2696 – 2729.
- Zhou, Z. and Wu, W. B. (2010). Simultaneous inference of linear models with time varying coefficients. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72(4):513–531.