

DBI Inflation and Warped Black Holes

Alexander Y. Kamenshchik^{*1}, Alessandro Tronconi^{†2}, and
Giovanni Venturi^{‡2}

¹Dipartimento di Fisica e Astronomia and INFN, Via Irnerio
46,40126 Bologna, Italy,

L.D. Landau Institute for Theoretical Physics of the Russian
Academy of Sciences, 119334 Moscow, Russia

²Dipartimento di Fisica e Astronomia and INFN, Via Irnerio 46,
40126 Bologna, Italy

Abstract

We study a possible amplification mechanism for the curvature perturbations generated during inflation by a Dirac-Born-Infeld (DBI) inflaton in the presence of a sharp feature in the warp factor. The large growth resulting in the scalar power spectrum is a consequence of the decreasing speed of sound. We obtain analytical approximate expressions for the relevant dynamical quantities both in the slow roll phase and during the transient phase which leads to the amplification. We finally compare our approximations to the exact numerical evolution and give, in such a context, the general rules for building a viable inflationary model leading to, during the subsequent radiation dominated era, an abundant primordial black holes formation.

^{*}Alexander.Kamenshchik@bo.infn.it

[†]Alessandro.Tronconi@bo.infn.it

[‡]Giovanni.Venturi@bo.infn.it

1 Introduction

From the very first LIGO detection of a gravitational wave signal in 2015 [1], the question of whether the black holes which produced such a signal could have a primordial origin was raised. The possibility that primordial black holes (PBHs) indeed exist was investigated many years ago in the seminal papers by Zeldovich and Novikov, and by Hawking [2]. At that time cosmology and astrophysics began to deal with the Dark Matter (DM) problem and PBHs soon turned out to be studied as possible DM candidates. Currently DM is definitely an urgent problem needing a solution since evidence for it is given by several independent observations. Moreover it is now clear that the primordial universe underwent a phase of accelerated expansion called inflation [4] generating the seeds of the large scale structures and the anisotropies in the cosmic microwave background (CMB) [5]. In such a context, it is therefore tempting to investigate whether the existence of PBHs offers a further possibility of improving our comprehension of the evolution of the early universe by relating the mechanisms necessary for PBHs formation to current observations. If PBHs were formed from a gravitational collapse of some over-densities during the radiation dominated era [6], and such over-densities were generated during inflation, then the mass distribution of PBHs on their formation can be related to the features of the inflaton potential in an interval which is not probed by CMB [7]. This therefore provides a way of distinguishing among a plethora of inflationary models leading to degenerate predictions concerning the scales probed by CMB.

The quest for a theory of quantum gravity is one of the major challenges of the last 40 years of theoretical physics. Even if quantum gravitational effects are expected to become manifest at energies above the Planck scale, one still expects that inflation is sensible to trans-planckian physics and quantum gravitational effects may then leave footprints on inflationary observables. In the present paper we investigate a possible, string theory inspired, mechanism leading to PBHs formation. String theory is widely believed to provide a fundamental description of Nature and is particularly relevant at the Planck scale, where the Standard Model and General Relativity are expected to break down.

Despite its formulation based on simple principles, string theory predicts an enormous set of possible low energy physical realisations, the string landscape, and we expect to live in one of these realisations where physical laws have the structure we experience. Different inflationary scenarios are con-

sidered as possible in the string theory context as the scalar fields necessary to play the role of the inflaton are ubiquitous and, for example, are associated with the structure of extra-dimensions. In particular, in the Dirac-Born-Infeld (DBI) scenario [8] the scalar field playing the role of the inflaton parametrises the position of a 3+1 D-brane moving along a six-dimensional “throat” with a warped geometry in the direction of motion. The peculiarity of this scenario consists of the fact that it possesses several non-standard features, some of them putting severe constraints on its viability with respect to the CMB observables related to it and others which appear indeed promising if PBHs formation is considered.

In the present article we study a possible mechanism for inflation and PBHs formation in the context of DBI inflation. The paper is organised as follows. In Section II we present the model, the relevant definitions and the equations governing the homogeneous dynamics. In particular, we introduce the slow roll (SR) parameters and illustrate how the dynamics simplifies during SR. We further discuss the possible existence of inflationary solutions which depart from SR and can be used to describe the evolution in the presence of sharp features in the warp factor. Section III concerns inflationary perturbations: we present the relevant equations and we discuss how a sharp feature in the warp factor may produce an amplification of the curvature perturbations. Finally a toy model is discussed in more detail and the numerical results are presented and explained. In Section IV we illustrate our conclusions.

2 Formalism

Let us consider the following lagrangian density for a minimally coupled DBI inflaton

$$\mathcal{L} = \sqrt{-g} \left[\frac{1}{f} \left(1 - \sqrt{1 + 2fX} - fV \right) \right] \equiv \sqrt{-g}P, \quad (1)$$

where $f = f(\phi)$ is related to the warp factor along a six-dimensional “throat” (having dimension $[f] = [V]^{-1}$), $V(\phi)$ is an arbitrary potential, $X = -\frac{1}{2}(\partial\phi)^2$ and P is the pressure of the scalar field fluid. In order to study the existence of inflationary solutions we consider the homogeneous part of the inflaton, i.e. $\phi = \phi(t)$ with $X = -\frac{1}{2}\dot{\phi}^2$ on a flat FLRW background metric, described by the line element

$$ds_4^2 = dt^2 - a(t)^2 d\vec{x} \cdot d\vec{x}. \quad (2)$$

The homogeneous equations of motion are the standard Einstein equations for a perfect fluid and the corresponding continuity equation, where the pressure of the fluid is defined in (1) and its energy density is

$$\rho = 2XP_{,X} - P = \frac{1}{f} \left[-1 + \frac{1}{\sqrt{1+2fX}} + fV \right]. \quad (3)$$

Unlike canonical models of inflation with a scalar field having $P_{\text{can}} = \frac{1}{2}(\partial\phi)^2 - V$, and a speed of sound c_s constant and equal to the speed of light ($c_s = 1$), in this context the speed of sound is variable and given by

$$c_s^2 = \frac{P_{,X}}{P_{,X} + 2XP_{,XX}} = 1 + 2fX = 1 - f\dot{\phi}^2 \quad (4)$$

with $0 < c_s \leq 1$ [9]. Let us note that a speed of sound very close to 1 is a sufficient condition for recovering the canonical behaviour and we observe that when $f\dot{\phi}^2 \ll 1$ such a condition is satisfied.

The resulting Friedmann equation is

$$H^2 = \frac{1}{3M_{\text{P}}^2 f} \left[-1 + \frac{1}{c_s} + fV \right] \quad (5)$$

and the Klein-Gordon (KG) equation for the inflaton is

$$\ddot{\phi} + 3Hc_s^2\dot{\phi} + \left[\frac{1-c_s}{2} \frac{1+2c_s}{1+c_s} \frac{f_{,\phi}}{f} \right] \dot{\phi}^2 + c_s^3 V_{,\phi} = 0. \quad (6)$$

2.1 Inflationary solutions

Inflation occurs if the slow roll parameter $\epsilon_1 \equiv -\dot{H}/H^2$ is smaller than one. From the continuity equation $\dot{\rho} = -3H(\rho + P)$ one easily obtains

$$\epsilon_1 \equiv -\frac{\dot{H}}{H^2} = \frac{3}{2} \left(1 + \frac{P}{\rho} \right) = \frac{3}{2} \frac{1 - c_s^2}{1 + c_s (fV - 1)}, \quad (7)$$

which corresponds to the second Friedmann equation, and the first Friedmann equation (5) can be cast in the form

$$M_{\text{P}}^2 H^2 \left(3 - \frac{2\epsilon_1}{1 + c_s} \right) = V. \quad (8)$$

When $\epsilon_1 \ll 1$ Eq. (8) takes the standard slow roll (SR) form

$$H^2 \simeq \frac{V}{3M_{\text{P}}^2} \quad (9)$$

independently of the value of the speed of sound c_s .

The second Friedmann equation (7) can be rewritten in the more standard form

$$\epsilon_1 = \frac{\phi_{,N}^2}{2M_{\text{P}}^2 c_s}, \quad (10)$$

where $_{,N}$ is the derivative with respect to the logarithm of the scale factor. Eq. (10) can be further derived leading to

$$\frac{\phi_{,NN}}{\phi_{,N}} \equiv \delta_1 + \delta_2 = \frac{\epsilon_c + \epsilon_2}{2} \quad (11)$$

where $\epsilon_c \equiv c_{s,N}/c_s$, $\epsilon_2 \equiv \epsilon_{1,N}/\epsilon_1$ and $\delta_1 \equiv \phi_{,N}/\phi$, $\delta_2 \equiv \delta_{1,N}/\delta_1$ are the SR parameters associated with the variation of the scalar field. Let us note that the condition $\epsilon_2 \ll 1$ is sufficient to guarantee a long inflationary phase and a negative value of ϵ_2 does not necessarily prevent a long inflationary phase. The derivative of the first Friedmann equation (8) leads to

$$\epsilon_1 \frac{3(1+c_s) - 2\epsilon_1 + \epsilon_2 - \frac{c_s}{1+c_s}\epsilon_c}{3(1+c_s) - 2\epsilon_1} = -n_V \frac{\delta_1}{2} \quad (12)$$

where $n_V \equiv \frac{d \ln V}{d \ln \phi}$. To the leading order in the slow roll approximation, with $|\epsilon_i| \ll 1$ and $|\delta_i| \ll 1$, Eq. (12) simplifies to

$$\epsilon_1 \simeq -n_V \frac{\delta_1}{2} \Rightarrow \epsilon_1 \simeq c_s \frac{M_{\text{P}}^2}{2} \left(\frac{V_{,\phi}}{V} \right)^2 \equiv c_s \epsilon_V \quad (13)$$

and $\epsilon_V \equiv (M_{\text{P}} V_{,\phi}/V)^2/2$ is the SR parameter associated with the flatness of the inflaton potential. For consistency the potential must satisfy $n_V = \mathcal{O}(1)$. Let us note that (13) takes the ‘‘canonical’’ form in the $c_s \rightarrow 1$ limit.

Finally, the Klein-Gordon equation can also be written in terms of the SR parameters as

$$\delta_1 \frac{\left[3c_s^2 - \epsilon_1 + \delta_1 + \delta_2 + \frac{1-c_s}{2} \frac{1+2c_s}{1+c_s} \delta_1 n_f \right]}{3 - \frac{2\epsilon_1}{(1+c_s)}} = -\frac{M_{\text{P}}^2}{\phi^2} c_s^3 n_V \quad (14)$$

where $n_f \equiv \frac{d \ln f}{d \ln \phi}$. On now using (11) and the derivative of the definition (4) one finds

$$\epsilon_c = \frac{c_s^2 - 1}{c_s^2 + 1} (n_f \delta_1 + \epsilon_2 - 2\epsilon_1) \quad (15)$$

and we observe that, for consistency, one also needs $n_f = \mathcal{O}(1)$ during SR. Finally, Eq. (14) simplifies to

$$\epsilon_1 \frac{\left[1 + \frac{\epsilon_2 - 2\epsilon_1}{3(1+c_s)} - \frac{c_s \epsilon_c}{3(1+c_s)^2}\right]^2}{\left[1 - \frac{2\epsilon_1}{3(1+c_s)}\right]^2} = c_s \epsilon_V \quad (16)$$

and we observe that (16) and (13) coincide to the leading order in the SR approximation. Let us note that the SR relation (13) results from (6) when $\dot{\phi} \ll 3Hc_s^2\dot{\phi}$ and $\dot{\phi}^2 f_\phi/f \ll 3Hc_s^2\dot{\phi}$.

From the definition of the speed of sound (4) one has

$$0 = c_s^2 + fH^2\phi_N^2 - 1 = c_s^2 + 2\frac{c_s fV}{3 - \frac{2\epsilon_1}{1+c_s}}\epsilon_1 - 1 \quad (17)$$

and during slow roll inflation, with $\epsilon_1 \simeq c_s \epsilon_V \ll 1$, Eq. (17) becomes

$$c_s^2 \simeq \frac{1}{1 + \frac{2fV\epsilon_V}{3}}. \quad (18)$$

Therefore if $c_s \epsilon_V \ll 1$ and $fV\epsilon_V \ll 1$ the speed of sound is close to unity and

$$c_s \simeq 1 - \frac{fV\epsilon_V}{3}. \quad (19)$$

In contrast if $c_s \epsilon_V \ll 1$ and $fV\epsilon_V \gg 1$, the speed of sound is close to zero and

$$c_s \simeq \sqrt{\frac{3}{2fV\epsilon_V}} \quad (20)$$

and we now observe that $fV \gg \epsilon_V^{-1}$ must also hold for consistency. On now squaring each side of Eq. (20) and using (13) one also finds

$$c_s \epsilon_1 \simeq \frac{3}{2fV} \quad (21)$$

and the more stringent condition $fV \gg \epsilon_1^{-1}$ must hold in order for SR to occur when $c_s \ll 1$. Since ϵ_1 is smaller than unity during inflation the product

fV must therefore be very large.

It can be easily verified by comparing (18) with the definition (4) that

$$\dot{\phi}^2 \simeq \frac{2V\epsilon_V}{3 + 2fV\epsilon_V} \quad (22)$$

is a first-order non-linear dynamical equation for ϕ valid in the SR regime which can be solved as usual. When $fV\epsilon_V \gg 1$ then (22) simplifies to $\dot{\phi}^2 \simeq 1/f$.

Let us note that the same, approximate, solution exists in the $c_s \rightarrow 0$ limit and when, in contrast to the SR approximation, $\ddot{\phi} \gg 3Hc_s^2\dot{\phi}$ and $\ddot{\phi} \gg c_s^3V_{,\phi}$. For such a case the KG equation simplifies to

$$\ddot{\phi} + \frac{1}{2} \frac{f_{,\phi}}{f} \dot{\phi}^2 \simeq 0. \quad (23)$$

In more detail, we observe that, even if the homogeneous evolution departs from slow roll, approximate inflationary solutions can be found. In particular, on considering a transient phase with a fast variation of f , and thus $n_f \gg 1$, the leading contributions in the KG equation are

$$\ddot{\phi} + \left[\frac{1 - c_s}{2} \frac{1 + 2c_s}{1 + c_s} \frac{f_{,\phi}}{f} \right] \dot{\phi}^2 = 0 \quad (24)$$

with the corresponding solution

$$c_s = \frac{1}{1 + A_s f(\phi)}, \quad (25)$$

where A_s is an integration constant. If this transient phase follows a slowly rolling phase then, on comparing (25) and (19) one finds

$$A_s \simeq \frac{\epsilon_{V,t} V_t}{3}, \quad (26)$$

where the subscript t indicates that the right hand side must be evaluated at the onset of the transient phase. Correspondingly one may estimate the relevant SR parameters

$$\delta_1 = s_{\dot{\phi}} \sqrt{\frac{3}{fV}} \frac{M_{\text{P}}}{\phi} \sqrt{1 - \frac{1}{(1 + A_s f)^2}} \quad (27)$$

$$\epsilon_1 = \frac{3A_s}{2V} \frac{2 + A_s f}{1 + A_s f}, \quad \epsilon_2 = -\delta_1 \left(n_V + \frac{n_f}{2 + A_s f} \right), \quad \epsilon_c = -\delta_1 n_f, \quad (28)$$

where $s_{\dot{\phi}}$ is the sign of $\dot{\phi}$.

3 Curvature Perturbations

In the above context, the equation for the scalar perturbations can be cast in the standard form [9]

$$v_k'' + \left(c_s^2 k^2 - \frac{z''}{z} \right) v_k = 0, \quad (29)$$

where v_k is the Fourier transformed Mukhanov-Sasaki field, the prime denotes the derivative with respect to conformal time η and $a d\eta = dt$. The time dependent function z is defined as

$$z \equiv \frac{a\sqrt{\rho + P}}{c_s H} = \frac{aM_{\text{P}}\sqrt{2\epsilon_1}}{c_s} \quad (30)$$

and

$$\frac{z''}{z} = a^2 H^2 \left[2 - \epsilon_1 + (3 - \epsilon_1) \left(\frac{\epsilon_2}{2} - \epsilon_c \right) + \left(\frac{\epsilon_2}{2} - \epsilon_c \right)^2 + \frac{d}{dN} \left(\frac{\epsilon_2}{2} - \epsilon_c \right) \right], \quad (31)$$

where the ϵ_i 's are small during SR inflation but may grow large (apart from ϵ_1) when inflation occurs far from SR (for example during some transient phase with n_f large). The corresponding curvature perturbations are described by $\mathcal{R}_k(\eta) = v_k/z$ and satisfy the equation

$$\mathcal{R}_k'' + 2aH \left(1 + \frac{\epsilon_2}{2} - \epsilon_c \right) \mathcal{R}_k' + c_s^2 k^2 \mathcal{R}_k = 0. \quad (32)$$

For modes outside the horizon (roughly for $c_s k/(aH) \ll 1$) the solution of (32) consists of a constant solution plus a varying solution of the form

$$\mathcal{R}_k = A_0 + A_1 \int^\eta \frac{d\eta'}{z^2}, \quad (33)$$

where the second term in Eq. (33) is either decreasing or increasing depending on the sign of z'/z . Indeed, during slow roll inflation the parameters ϵ_2 and ϵ_c in the friction term are much smaller than one and \mathcal{R}_k freezes at the horizon exit. In contrast if $\epsilon_2/2 - \epsilon_c < -1$, the friction term has a negative sign and \mathcal{R}_k may grow after the horizon exit. An inflationary phase with a negative friction term in (32) is often exploited to build models of inflation wherein a certain interval of modes of the curvature perturbations are amplified. The corresponding over-densities, which form when inflation ends, may

then collapse and originate primordial black holes. For example, in canonical models of inflation ($\epsilon_c = 0$) the presence of an inflection point in the potential [7] leads to a phase of ultra slow roll with $\epsilon_2 \simeq -6$ wherein the amplitude of curvature perturbation is amplified. In non-canonical models of inflation a rapid increase of the speed of sound may be responsible for the amplification [10]. Still some care is due because a very fast growth of c_s may delay or even prevent modes from exiting the horizon and therefore stop amplification. In particular one needs a decreasing $\frac{kc_s}{aH}$ ratio, i.e.

$$\left(\frac{kc_s}{aH}\right)^{-1} \frac{d}{dN} \frac{kc_s}{aH} < 0 \Rightarrow \epsilon_c < 1 - \epsilon_1 \text{ and } \epsilon_c > 1 + \frac{\epsilon_2}{2} \quad (34)$$

must be satisfied and this necessarily requires a careful tuning of the inflationary model.

The resulting power spectrum is given by the expression

$$\mathcal{P}_{\mathcal{R}} = \frac{k^3}{2\pi^2} |\mathcal{R}_k|^2 \Big|_{\text{e.o.i.}} \quad (35)$$

where the subscript indicates that it must be evaluated at the end of inflation (e.o.i.). When the SR conditions are satisfied the curvature perturbations freeze after the horizon exit. In such a case their power spectrum can be evaluated at the horizon exit (h.e.) and is given by

$$\mathcal{P}_{\mathcal{R}} = \frac{1}{8\pi^2 M_{\text{P}}^2} \frac{H^2}{c_s \epsilon_1} \Big|_{\text{h.e.}} \simeq \frac{V}{24\pi^2 M_{\text{P}}^4} \frac{1 + \frac{2}{3} fV \epsilon_V}{\epsilon_V} \Big|_{\text{h.e.}} \quad (36)$$

and thus an amplification or suppression of the spectrum can be realised as a consequence of a corresponding suppression or amplification of the product $\epsilon_1 c_s$ or equivalently by the amplification/suppression of the product fV , provided the SR conditions are not violated. If SR conditions are violated but z'/z is positive at horizon crossing one may still evaluate $\mathcal{P}_{\mathcal{R}}$ at horizon exit which now occurs when $c_s k \sim aH|\gamma|$ and $\gamma \neq 1$ is implicitly defined as $z''/z = 2a^2 H^2 \gamma^2$. The resulting expression is

$$\mathcal{P}_{\mathcal{R}} = \frac{1}{8\pi^2 M_{\text{P}}^2} \frac{H^2 |\gamma|^2}{c_s \epsilon_1} \Big|_{\text{h.e.}} = \frac{1}{12\pi^2} \frac{V |\gamma|^2}{\delta_1^2 \phi^2 M_{\text{P}}^2} \Big|_{\text{h.e.}} \quad (37)$$

and coincides with (36) when $\gamma \simeq 1$. If $|\gamma|$ is large and/or $\epsilon_1 c_s$ is smaller compared to its value at CMB scales the resulting amplitude gets amplified.

Let us note that when modes exit the horizon and freeze, they still remains constant if, at some time, z'/z becomes negative because $A_1 \ll 1$ in (33). In contrast, if A_1 is much larger (and this occurs for modes exiting the horizon when $z'/z < 0$) a significant amplification of the curvature \mathcal{R}_k could be produced if the condition $z'/z < 0$ is valid for a certain time interval (or number of e-folds $\Delta N_{\text{amp}} \geq 1$).

A mechanism leading to an abundant amplification of \mathcal{R}_k was already discussed in a previous paper [11] in the context of non-canonical models of inflation. There the evolution of an inflationary model with a modified kinetic term plus a potential was studied. The presence of the potential is necessary in order to drive a SR phase wherein the perturbations we observe in the CMB are generated. In contrast, the modified kinetic term, which becomes leading at some point during inflation and after the SR phase, is responsible for the amplification of the scalar perturbations and the subsequent formation of PBHs.

As we already discussed, for DBI inflation the inflaton lagrangian has a SR solution at $c_s \sim 1$ and at $c_s \sim 0$ and the curvature perturbations can be therefore amplified in the transition from the first phase to the second. In order for the over-densities to collapse and form a sufficient population of PBHs with mass $m_{\text{bh}} \sim 10^{-15} M_\odot$ (thus possibly forming a large fraction of the Dark Matter budget today) the amplification must increase the power spectrum until the critical threshold $\mathcal{P}_{\mathcal{R}} \sim 10^{-2}$ is reached. The modes affected by such an amplification must exit the horizon roughly at

$$\Delta N(m_{\text{bh}}) \sim 26 - \frac{1}{12} \ln \left(\frac{g_{*k}}{g_{*0}} \right) + \ln \frac{c_{s,\text{bh}} H_*}{c_{s,*} H_{\text{bh}}} \quad (38)$$

e-folds after CMB modes, where g_{*k} , g_{*0} indicates the number of relativistic degrees of freedom at the BHs formation and at present respectively, $c_{s,*}$ and H_* are evaluated at horizon exit for CMB modes and $c_{s,\text{bh}}$ and H_{bh} must be evaluated when the spectrum reaches the critical threshold.

3.1 Model building

We now illustrate how a viable DBI inflationary model which includes an amplification phase may be realised. The features of the scalar inflationary perturbations imprinted in the CMB put severe constrains on the inflaton evolution when these perturbations modes exit the horizon, ~ 60 e-folds before inflation ends. CMB observations [12] constrain the spectral index

$n_s - 1 \sim \mathcal{O}(10^{-2})$, the amplitude of the spectrum $\mathcal{P}_{\mathcal{R}} \sim 2 \cdot 10^{-9}$ and the speed of sound (which can be related to the statistical abundance of non-gaussian features observed in the CMB [13])

$$c_s > 3 \cdot 10^{-2}. \quad (39)$$

If we assume that $c_s \sim 1$ at that time the remaining two constraints to be satisfied are

$$\mathcal{P}_{\mathcal{R}} = \frac{1}{24\pi^2 M_{\text{P}}^4} \frac{V_*}{\epsilon_{V,*}} \sim 2 \cdot 10^{-9}, \quad n_s - 1 \sim \mathcal{O}(\epsilon_{V,*}) \sim \mathcal{O}(10^{-2}), \quad (40)$$

where the subscript $*$ indicates that the quantities are evaluated when CMB scales exit the horizon. This fixes the shape of the potential, i.e. its slope and value at the “beginning” of inflation and the magnitude of f_* (as we already discussed $fV\epsilon_V \ll 1$ is needed in order for c_s to be very close to unity).

Hereafter we shall study how the amplification of the spectrum $\mathcal{P}_{\mathcal{R}}$ may occur as a consequence of a sharp feature in the warp factor f . The sharpness is required in order to realise the amplification needed for an efficient PBHs production in the short amount of time of a few e-folds. The perturbations which exit the horizon ~ 30 e-folds after those imprinted in the CMB would collapse in PBHs of particular interest (see (38)) since they may survive evaporation until today and constitute a large part of the Dark Matter budget.

Let us note that the possible physical origins of the presence of a sharp feature in the warp factor have been studied in several articles [14] and we shall not discuss it further. We shall limit ourselves to analysing its consequences as a possible source of amplification of the curvature perturbations. In order to simplify our model we shall study the consequences of a gaussian feature in an interval with a constant warp factor. Nonetheless one may easily generalise our results to more realistic models.

When the scalar field passes through the region where the feature is present the sound speed decreases and the dynamics may enter a non-canonical phase. Thus, as a consequence, the spectrum of the scalar perturbations gets amplified. One must estimate the duration of such a transient phase and tune the parameters of the model to ensure that the inflaton crosses the feature within a few e-folds. Let therefore consider

$$f = F \left[1 + A e^{-\frac{\alpha}{M_{\text{P}}^2} (\phi - \phi_0)^2} \right], \quad (41)$$

with $\epsilon_{V,*} \ll 1$, $AV_0/V_*\epsilon_{V,0} \gg 1$ where the subscript 0 indicates the value at $\phi = \phi_0$ and A and F are constant. When the inflaton evolves non-canonically

then $|\dot{\phi}| \sim f^{-1/2}$ and

$$\Delta N_g = \int_{t_i}^{t_f} H dt = \int_{\phi_i}^{\phi_f} \frac{H}{\dot{\phi}} d\phi \quad (42)$$

is the number of e-folds needed to cross the (gaussian) feature. Between ϕ_i and ϕ_f the gaussian is much larger than 1 and one has

$$\Delta N_g \simeq \int_{\phi_i}^{\phi_f} \sqrt{\frac{VFA}{3M_{\text{P}}^2}} e^{-\frac{\alpha}{2M_{\text{P}}^2}(\phi-\phi_0)^2} d\phi \sim \int_{-\infty}^{+\infty} \sqrt{\frac{V_0FA}{3M_{\text{P}}^2}} e^{-\frac{\alpha}{2M_{\text{P}}^2}(\phi-\phi_0)^2} d\phi, \quad (43)$$

where in the last integral we approximated V by its value at the maximum of the gaussian, ϕ_0 , and we extended the integral over the whole real axis because the contributions to the integral from the tails of the gaussian function are negligible. We then obtain the following estimate

$$\Delta N_g \sim \sqrt{\frac{2\pi V_0FA}{3\alpha}} \quad (44)$$

which can be expressed in terms of the cosmic time as

$$\Delta t_g = \int_{N_i}^{N_f} \frac{dN}{H} \sim \sqrt{\frac{3M_{\text{P}}^2}{V_0}} \Delta N_g. \quad (45)$$

Furthermore, close to the maximum of the gaussian n_f is $\mathcal{O}(1)$ and SR is restored. The amplitude of the spectrum generated around ϕ_0 is given by

$$\mathcal{P}_{\mathcal{R},0} = \frac{V_0}{24\pi^2 M_{\text{P}}^4} \frac{2f_0 V_0}{3} = \mathcal{P}_{\mathcal{R}} \frac{2f_0 V_0^2 \epsilon_{V,*}}{3V_*}, \quad (46)$$

and in order to obtain an amplification of n orders of magnitude one needs

$$\frac{\mathcal{P}_{\mathcal{R},0}}{\mathcal{P}_{\mathcal{R}}} \simeq \frac{2f_0 V_0^2 \epsilon_{V,*}}{3V_*} \sim 10^n. \quad (47)$$

One may also evaluate the spectrum amplification determined by the transient phase, which still occurs for $\phi \sim \phi_0$ if the gaussian is sharply peaked around ϕ_0 . A rough estimate on using Eqs. (27,28) leads to $|\gamma|^2 \sim \mathcal{O}(n_f^2 \delta_1^2)$ and

$$\mathcal{P}_{\mathcal{R},t} \sim \frac{1}{12\pi^2} \frac{V_0 n_f^2}{\phi_0^2 M_{\text{P}}^2} \Big|_{\text{h.e.}} \sim \frac{1}{12\pi^2} \frac{\alpha V_0}{M_{\text{P}}^4} \Big|_{\text{h.e.}} = \frac{2\alpha V_0 \epsilon_{V,*}}{V_*} \mathcal{P}_{\mathcal{R}}, \quad (48)$$

when $n_f \gg 1$ and with $\max |n_f| \sim \mathcal{O}(\phi_0 M_{\text{P}}^{-1} \sqrt{\alpha})$. In order to obtain an amplification of n orders of magnitude during the transient one then needs

$$\frac{\mathcal{P}_{\mathcal{R},t}}{\mathcal{P}_{\mathcal{R}}} \simeq \frac{2\alpha V_0 \epsilon_{V,*}}{V_*} \sim 10^n. \quad (49)$$

3.2 Numerical Example

We finally illustrate a simple applications of our estimates from the previous section and then compare the resulting curvature perturbations spectrum to the numerical (exact) results. For simplicity we consider an exponential potential driving SR inflation

$$V = V_* \exp\left[\frac{\beta(\phi - \phi_*)}{M_{\text{P}}}\right] \quad (50)$$

with $\beta = 3 \cdot 10^{-1}$, a constant $\epsilon_V = 4.5 \cdot 10^{-2}$ and a warp factor given by the expression (41). The amplitude (40) fixes

$$V_* \sim 2 \cdot 10^{-8} M_{\text{P}}^4 \quad (51)$$

and F should be chosen so as to ensure that the sound speed satisfies the experimental constraints. The sound speed must be close to one when CMB scales exit the horizon and therefore we set $F = 10^6 M_{\text{P}}^{-4}$. With such a setup the field slowly rolls down the potential. If we assume that amplification starts $\Delta N_* \sim 30$ e-folds after CMB scales exit the horizon one has

$$\phi_0 \simeq \phi_* - \beta^2 M_{\text{P}} \Delta N_*. \quad (52)$$

One may therefore estimate (44)

$$\Delta N_g \sim \sqrt{\frac{2\pi F V_* A}{3\alpha}} \exp\left[-\frac{\beta^2 \Delta N_*}{2}\right] \quad (53)$$

and the amount of amplification realised during the transient can be related with the parameter α in the gaussian as follows

$$\frac{\mathcal{P}_{\mathcal{R},t}}{\mathcal{P}_{\mathcal{R}}} = 2\alpha e^{-\beta^2 \Delta N_*} \epsilon_{V,*} \sim 10^n \Rightarrow \alpha \sim 10^{n+2}. \quad (54)$$




Figure 1: *In the figure above the evolution of $\log_{10} c_s^2$ is plotted as a function of N . The solid grey line represents the exact numerical solution of the homogeneous equations, the dashed line represents the SR analytical approximation and the dotted line represents the approximation found for the transient phase. The interval of the transient which includes its beginning and the period of maximum variation of c_s is very well described. Moreover the figure shows that the estimate for ΔN_g is well approximated by (53).*

Near the maximum of the warp factor f , SR is restored and the amount of amplification is the following function of the product $F \cdot A$

$$\frac{\mathcal{P}_{\mathcal{R},0}}{\mathcal{P}_{\mathcal{R}}} = \frac{2}{3} F A V_* e^{-2\beta^2 \Delta N_*} \epsilon_{V,*} \sim 10^n \Rightarrow F \cdot A \sim 3 \cdot 10^{n+12}. \quad (55)$$

Let us note that by comparing the ratio of (55) and (54) with (53) one has

$$\frac{\mathcal{P}_{\mathcal{R},0}}{\mathcal{P}_{\mathcal{R},t}} = \frac{F A V_* e^{-\beta^2 \Delta N_*}}{3\alpha} = \frac{\Delta N_g^2}{2\pi} \quad (56)$$

and we therefore observe that for a transient phase lasting ($\sim \Delta N_g/2$) few e-folds the two expressions lead to the same amount of amplification.

3.3 Numerical Results

For more accurate results and in order to test the analytical approximate expressions obtained, a numerical analysis is required.

The plot in the figure (1) shows the homogeneous function $\log_{10} c_s^2$ as a function of N in order to compare the exact (numerical) evolution of the homogeneous system (solid grey line) and the analytical approximations for the SR evolution (dashed line) and the transient (dotted line). The approximations are very close to the exact expression. Let us note that $c_s^2 - 1$ is proportional to f which is very large in the interval when the gaussian feature is present and thus c_s^2 is very sensitive to very small differences between the exact and the approximate evolutions. Very small differences between the exact evolution of $\phi(N)$ and the corresponding analytical approximations are amplified by several order of magnitude when multiplied by f and, as the figure shows, result in just a few order of magnitude deviations as far as the evolution of c_s is concerned.

In the figure (2) we plotted the spectrum $\log_{10} \mathcal{P}_{\mathcal{R}}$ as a function of $\log_{10} k/a_0$ and a_0 is the arbitrary value of the scale factor at the beginning of the numerical analysis. The solid line represents the exact numerical results while the dots corresponds to the analytical estimate (37) evaluated at horizon crossing, where the horizon exit is estimated for each mode by solving exactly the condition $c_s k = aH|\gamma|$ through the (homogeneous) exact numerical solutions. We set

$$\alpha \simeq 2.9 \cdot 10^5, \quad F = 10^6, \quad F A \simeq 3.5 \cdot 10^{15}, \quad (57)$$

which, according to our estimates, gives an amplification phase of about $\Delta N_0 \sim 4$ e-folds (it takes ~ 8 e-folds for the scalar field to get across the gaussian feature) and an amplification factor of about 3 orders of magnitude. Let us note that (37) well approximates the numerical results before and after the transient stage and when the field enters in the region where $|n_f|$ becomes large (c_s decreases) at the onset of the gaussian feature. Close to the maximum of f the homogeneous evolution enters a short SR phase and finally $|n_f|$ becomes large again (c_s increases) as the field crosses the maximum of the gaussian feature and finally relaxes on the SR attractor with $f = \text{const}$. The figure shows that after the first large variation of f the amplitude $\mathcal{P}_{\mathcal{R}}$ reaches a maximum and then begins oscillating with an increasing frequency. This oscillation is the consequence of modes re-entering the horizon due to the increasing speed of sound. At the horizon re-entry v_k evolves as a superposition of ingoing and outgoing waves

$$v_k \sim \frac{1}{\sqrt{2c_s k}} \left[C_k e^{i \int_{\eta_k}^{\eta} kc_s d\eta'} + D_k e^{-i \int_{\eta_k}^{\eta} kc_s d\eta'} \right], \quad (58)$$

where η_k is the conformal time at re-entry, and therefore its modulus oscillates. Let us note that at early times, before exiting the horizon for the first time, the Bunch-Davies initial condition sets $D_k \rightarrow 0$ for each mode and modulus oscillations are absent. At the end of inflation, the amplitude of the modes outside the horizon is frozen and the oscillation remains imprinted in the perturbations.

The amplification realised with the choice of parameters (57) is just an example of how the presence of a sharp feature in the warp factor may affect the scalar power spectrum at the end of inflation. The example shows that the analytical estimates of the previous sections are quite consistent. A suitable choice of α , F and A will certainly lead to an amplification of 6-7 orders of magnitude, eventually resulting in PBHs formation with a mass depending on ΔN (38). We restricted our analysis to the study of a few order of magnitude of amplification because a precise numerical analysis and a much steeper warp factor needed by a much larger amount of amplification would have required too much time and processor power.

Let us finally note that our results can be easily generalised to different potentials and warp factors. In particular, the steepness required to obtain a large amplification in a short amount of time needs a very localised feature and is therefore very mildly sensitive to the global form of f and V .

4 Conclusions

Primordial black holes have attracted much attention in the last few years since they are viable candidates for dark matter and because the dynamical mechanism behind their formation may probe a range of inflationary scales which are distinct compared to those imprinted in the CMB.

In the present article we generalise the conclusions of a preceding paper [11] wherein a mechanism for the amplification of the scalar power spectrum, possibly leading to PBHs formation at the end of inflation, was proposed for a class of non-canonical inflationary models. In such models the inflaton has a non-standard kinetic term and, as a consequence, the speed of sound associated with the curvature perturbations produced during inflation is non-constant and different from the speed of light. In such a context, for a series of simplified models we illustrated that the presence of a decreasing speed of sound may amplify the scalar inflationary perturbations to a critical threshold required in order to lead to an efficient PBHs formation.




Figure 2: In the above figure $\log_{10} \mathcal{P}_{\mathcal{R}}$ is plotted as a function on $\log_{10}(k/a_0)$. The solid line represents the power spectrum calculated with the exact numerical solution of the Mukhanov-Sasaki equation (29) and the points are the corresponding analytical estimate using the Eq. (37) where the horizon exit is calculated numerically for each mode with the exact (homogeneous) solutions. The points represent the modes which essentially exit the horizon once during their evolution (and some limiting cases for modes which re-enter for a very short amount of time). The oscillation are produced by modes which re-enter the horizon and have enough time to evolve inside the horizon and generate the oscillations before exiting the horizon again. For such modes the estimate (37) fails and the corresponding points are not plotted.

In particular a string inspired model, wherein a DBI scalar field with a potential and minimally coupled to gravity plays the role of the inflaton, is considered and the possible amplification of inflationary perturbations due to a decreasing speed of sound is studied.

Let us note that, analogously to what occurs for Ultra Slow Roll inflation and a fast variation of ϵ_1 , also in the presence of a rapidly increasing speed of sound the curvature perturbations may significantly grow after the horizon exit. The corresponding mechanism of amplification has been studied in the literature and applied to DBI inflation [10] but is different from ours which is based on a decreasing speed of sound.

For simplicity our results were applied to an inflaton with an exponential potential and warp factor $f(\phi)$ everywhere constant except for a tiny inter-

val where a steep gaussian feature is present. The presence of such a feature leads to transient phase with a rapidly decreasing speed of sound and a fast amplification of curvature perturbations. We then studied how the homogeneous evolution of the system can be approximated both during SR and in the presence of a rapidly varying warp factor and we found precise analytical expressions for the spectrum of the perturbations generated during these phases. Finally, our approximations were applied to an explicit model where the comparison between the resulting analytical expressions and their exact numerical estimate showed a very good agreement.

We finally concluded that DBI inflation is a viable candidate for inflation and may lead to significant PBHs formation in the presence of some very sharp feature in the warp factor. Different DBI inflationary models can be studied as a source of PBHs by means of the precise analytical expressions we found and compared to observational constraints related to inflation and DM abundance.

5 Acknowledgements

A.K. is partially supported by the Russian Foundation for Basic Research grant No. 20-02-00411.

References

- [1] B. P. Abbott *et al.* [LIGO Scientific and Virgo], Phys. Rev. Lett. **116** (2016) no.6, 061102 doi:10.1103/PhysRevLett.116.061102 [arXiv:1602.03837 [gr-qc]].
- [2] Y. B. Zel'dovich, I. D., Soviet Astron. AJ (Engl. Transl.), **10** (1967), 602; S. Hawking, Mon. Not. Roy. Astron. Soc. **152** (1971), 75
- [3] G. F. Chapline, Nature **253** (1975) no.5489, 251-252 doi:10.1038/253251a0; P. Meszaros, Astron. Astrophys. **38** (1975), 5-13
- [4] A.A. Starobinsky. Springer. in H.J. De Vega and N. Sanchez (eds.) Current trends in field theory quantum gravity and strings, Lecture Notes in Physics 246 Verlag, Heidelberg, 1986), pp. 107-126. A.D. Linde.

Academic. Particle Physics and Inflationary Cosmology (Harwood New York, 1990).

- [5] V.F. Mukhanov, Sov. Phys. JETP **68**, 1297 (1988); J. M. Maldacena, JHEP **0305** (2003) 013; V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rept. **215** (1992) 203. V.F. Mukhanov, Phys. Lett. B **218**, 17 (1989); J. M. Bardeen, Phys. Rev. D **22**, 1882 (1980). doi:10.1103/PhysRevD.22.1882 M. Sasaki, Prog. Theor. Phys. **70** (1983) 394. doi:10.1143/PTP.70.394
- [6] B. J. Carr, Astrophys. J. **201** (1975) 1. doi:10.1086/153853 M. Sasaki, T. Suyama, T. Tanaka and S. Yokoyama, Class. Quant. Grav. **35** (2018) no.6, 063001 doi:10.1088/1361-6382/aaa7b4 [arXiv:1801.05235 [astro-ph.CO]].
- [7] J. Garcia-Bellido and E. Ruiz Morales, Phys. Dark Univ. **18** (2017) 47 doi:10.1016/j.dark.2017.09.007 [arXiv:1702.03901 [astro-ph.CO]]. G. Ballesteros and M. Taoso, Phys. Rev. D **97** (2018) no.2, 023501 doi:10.1103/PhysRevD.97.023501 [arXiv:1709.05565 [hep-ph]]. M. Cicoli, V. A. Diaz and F. G. Pedro, JCAP **1806** (2018) no.06, 034 doi:10.1088/1475-7516/2018/06/034 [arXiv:1803.02837 [hep-th]]. H. Motohashi and W. Hu, Phys. Rev. D **96** (2017) no.6, 063503 doi:10.1103/PhysRevD.96.063503 [arXiv:1706.06784 [astro-ph.CO]]. C. Germani and T. Prokopec, Phys. Dark Univ. **18** (2017) 6 doi:10.1016/j.dark.2017.09.001 [arXiv:1706.04226 [astro-ph.CO]].
- [8] M. Born and L. Infeld, Proc. Roy. Soc. Lond. A **144** (1934) no.852, 425-451 doi:10.1098/rspa.1934.0059 M. Alishahiha, E. Silverstein and D. Tong, Phys. Rev. D **70** (2004), 123505 doi:10.1103/PhysRevD.70.123505 [arXiv:hep-th/0404084 [hep-th]]; D. Bessada, W. H. Kinney and K. Tzirakis, JCAP **09** (2009), 031 doi:10.1088/1475-7516/2009/09/031 [arXiv:0907.1311 [gr-qc]]. W. H. Kinney and K. Tzirakis, Phys. Rev. D **77** (2008), 103517 doi:10.1103/PhysRevD.77.103517 [arXiv:0712.2043 [astro-ph]]. M. Spalinski, Phys. Lett. B **650** (2007), 313-316 doi:10.1016/j.physletb.2007.05.041 [arXiv:hep-th/0703248 [hep-th]].
- [9] C. Armendariz-Picon, T. Damour and V. F. Mukhanov, Phys. Lett. B **458** (1999) 209 doi:10.1016/S0370-2693(99)00603-6 [hep-th/9904075]. J. Garriga and V. F. Mukhanov, Phys. Lett. B **458** (1999)

- 219 doi:10.1016/S0370-2693(99)00602-4 [hep-th/9904176]. R. Gwyn, M. Rummel and A. Westphal, JCAP **1312** (2013) 010 doi:10.1088/1475-7516/2013/12/010 [arXiv:1212.4135 [hep-th]]. P. Franche, R. Gwyn, B. Underwood and A. Wissanji, Phys. Rev. D **81** (2010) 123526 doi:10.1103/PhysRevD.81.123526 [arXiv:0912.1857 [hep-th]].
- [10] O. Özsoy, S. Parameswaran, G. Tasinato and I. Zavala, JCAP **07** (2018), 005 doi:10.1088/1475-7516/2018/07/005 [arXiv:1803.07626 [hep-th]].
- [11] A. Y. Kamenshchik, A. Tronconi, T. Vardanyan and G. Venturi, Phys. Lett. B **791** (2019), 201-205 doi:10.1016/j.physletb.2019.02.036 [arXiv:1812.02547 [gr-qc]].
- [12] P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.02114 [astro-ph.CO]. P. A. R. Ade *et al.* [Planck Collaboration], arXiv:1502.01589 [astro-ph.CO]. Y. Akrami *et al.* [Planck Collaboration], arXiv:1807.06211 [astro-ph.CO]. N. Aghanim *et al.* [Planck Collaboration], arXiv:1807.06209 [astro-ph.CO].
- [13] J. M. Maldacena, JHEP **0305** (2003) 013 doi:10.1088/1126-6708/2003/05/013 [astro-ph/0210603]. N. Bartolo, E. Komatsu, S. Matarrese and A. Riotto, Phys. Rept. **402** (2004) 103 doi:10.1016/j.physrep.2004.08.022 [astro-ph/0406398]. G. Franciolini, A. Kehagias, S. Matarrese and A. Riotto, JCAP **1803** (2018) no.03, 016 doi:10.1088/1475-7516/2018/03/016 [arXiv:1801.09415 [astro-ph.CO]]. P. A. Klimai and E. V. Bugaev, arXiv:1210.3262 [astro-ph.CO]. V. Atal and C. Germani arXiv:1811.07857 [astro-ph.CO] X. Chen, M. x. Huang, S. Kachru and G. Shiu, JCAP **0701** (2007) 002 doi:10.1088/1475-7516/2007/01/002 [hep-th/0605045]. J. Noller and J. Magueijo, Phys. Rev. D **83** (2011) 103511 doi:10.1103/PhysRevD.83.103511 [arXiv:1102.0275 [astro-ph.CO]].
- [14] V. Miranda, W. Hu and P. Adshead, Phys. Rev. D **86** (2012), 063529 doi:10.1103/PhysRevD.86.063529 [arXiv:1207.2186 [astro-ph.CO]]. R. Bean, X. Chen, G. Hailu, S. H. H. Tye and J. Xu, JCAP **03** (2008), 026 doi:10.1088/1475-7516/2008/03/026 [arXiv:0802.0491 [hep-th]]. V. Miranda and W. Hu, Phys. Rev. D **89** (2014) no.8, 083529 doi:10.1103/PhysRevD.89.083529 [arXiv:1312.0946 [astro-ph.CO]]. V. Miranda and W. Hu, Phys. Rev. D **89** (2014)

no.8, 083529 doi:10.1103/PhysRevD.89.083529 [arXiv:1312.0946 [astro-ph.CO]]. T. Kobayashi and J. Yokoyama, JCAP **02** (2013), 005 [erratum: JCAP **09** (2013), E02] doi:10.1088/1475-7516/2013/02/005 [arXiv:1210.4427 [astro-ph.CO]].