KAM THEORY FOR ACTIVE SCALAR EQUATIONS

ZINEB HASSAINIA, TAOUFIK HMIDI, AND NADER MASMOUDI

ABSTRACT. In this paper, we establish the existence of time quasi-periodic solutions to generalized surface quasi-geostrophic equation $(gSQG)_{\alpha}$ in the patch form close to Rankine vortices. We show that invariant tori survive when the order α of the singular operator belongs to a Cantor set contained in $(0, \frac{1}{2})$ with almost full Lebesgue measure. The proof is based on several techniques from KAM theory, pseudo-differential calculus together with Nash-Moser scheme in the spirit of the recent works [4, 9]. One key novelty here is a refined Egorov type theorem established through a new approach based on the kernel dynamics together with some hidden Töpliz structures.

Contents

Pretace	1
1. Introduction	2
1.1. Periodic vortex patches solutions	3
1.2. Quasi-periodic solutions	4
1.3. Main result and ideas of the proof	5
2. Hamiltonian reformulation	16
2.1. Contour dynamics equation	16
2.2. Conservation laws	18
2.3. Hamiltonian structure and reversibility	19
3. Linearization and frequencies structure	22
3.1. Linearized operator	22
3.2. Structure of the linear frequencies	26
3.2.1. Monotonicity and asymptotic behavior	26
3.2.2. Non-degeneracy and transversality	28
3.3. Linear quasi-periodic solutions	34
4. Functional tools	35
4.1. Function spaces	35
4.2. Classical operations	35
4.3. Modified periodic fractional Laplacian	42
5. Hamiltonian toolbox	44
5.1. Action-angle coordinates	44
5.2. Hamiltonian regularity	47
6. Berti-Bolle approach for the approximate inverse	50
6.1. Linear change of variables	51
6.2. Defect of the symplectic structure	55
6.3. Construction of an approximate inverse	57
7. Toroidal pseudo-differential operators	61
7.1. Operators and reversibility	61
7.2. Symbol class topology	62
7.3. Continuity and law products	65
7.4. Commutators	72
8. New approach related to Egorov theorem	76
8.1. Nonlocal pseudo-differential hyperbolic equation	76
8.2. Töplitz matrices and binomial convolution	81
8.3. Refined Egorov's theorem	83

8.3.1. Tame estimates of the iterated commutators	100
8.3.2. Iterated kernel estimates	108
8.3.3. Iterated Töplitz matrix operators	121
9. Reducibility of the linearized operator	126
9.1. Structure on the normal direction	127
9.2. Asymptotic structure	129
9.3. Reduction of the transport part	133
9.3.1. Transport equation with constant coefficients	133
9.3.2. Straightening of the transport equation	134
9.4. First conjugation of the linearized operator	154
9.5. Reduction of the nonlocal part	162
9.6. Complete reduction up to small errors	169
9.6.1. Frequency localization of operators	170
9.6.2. Localization on the normal direction	174
9.6.3. KAM reduction of the remainder term	179
9.6.4. Approximate inverse in the normal direction	200
10. Proof of the main result	203
10.1. Nash-Moser scheme	204
10.2. Final Cantor set estimates	216
Appendix A. Appendix	223
Acknowledgements:	226
References	227

1. INTRODUCTION

In this paper we are concerned with the generalized surface quasi-geostrophic equations $(gSQG)_{\alpha}$ described by

$$\begin{cases} \partial_t \theta + u \cdot \nabla \theta = 0, \quad (t, x) \in [0, T] \times \mathbb{R}^2, \\ u = -\nabla^{\perp} (-\Delta)^{-1 + \frac{\alpha}{2}} \theta, \\ \theta_{|t=0} = \theta_0. \end{cases}$$
(1.1)

Here u stands for the velocity field, $\nabla^{\perp} = (-\partial_2, \partial_1)$ and $\alpha \in (0, 2)$. The fractional Laplacian $(-\Delta)^{-1+\frac{\alpha}{2}}$ is associated to Riesz potential according to the convolution law

$$(-\Delta)^{-1+\frac{\alpha}{2}}\theta(x) = \frac{C_{\alpha}}{2\pi} \int_{\mathbb{R}^2} \frac{\theta(y)}{|x-y|^{\alpha}} dy, \quad \text{with} \quad C_{\alpha} \triangleq \frac{\Gamma(\alpha/2)}{2^{1-\alpha}\Gamma(\frac{2-\alpha}{2})}.$$
 (1.2)

The limiting case $\alpha \to 0$ corresponds to the 2D Euler equation, with θ representing the fluid vorticity, the midpoint case $\alpha = 1$ corresponds to the surface quasi-geostrophic equation (SQG), with θ denoting the temperature in a rapidly rotating stratified fluid with uniform potential vorticity and the case $\alpha = 2$ produces stationary solutions.

While global regularity of smooth solutions to the 2D Euler equations has been known since long ago [63, 83], only local regularity persistence is proved for the entire range $\alpha \in (0, 2)$, we refer for instance to [25, 26, 28]. The question of whether a finite-time singularity may develop from smooth initial datum remains open. However, certain blow-up scenarios have been ruled out in [29, 30] and solutions with exponential growth (not excluding blow up in finite time) are shown to exist in [56]. On the other hand, examples of a non-trivial global in time smooth solutions have been recently constructed in [2, 24, 49] with different approaches related to bifurcation theory/assisted computer proof and variational principle.

In the setting of weak solutions, it is known that for Euler equations Yudovich solutions exist globally in time and they are unique, see [84]. However, for $\alpha > 0$ global existence persists but the uniqueness issue at the energy level remains open, see for instance [71, 75]. Recently, non-unique

weak solutions with negative Sobolev regularity were constructed in [19]. An interesting sub-class which has been explored in various directions during the past few decades is given by the vortex patches, that is, initial data taking the form of the characteristic function $\theta_0(x) = \chi_D$, where D is a bounded smooth domain. In this framework, the contour dynamics equation offers a suitable tool to construct local solutions in the patch form with smooth boundary, see [25, 41, 76]. The uniqueness in this class for $\alpha \in (0, 1)$, was discussed in [66] and the case $\alpha = 1$ was analyzed in [32].

It is worthy to point out that while the boundary's regularity is globally preserved for $\alpha = 0$ as proved in [17, 27], numerical evidence [31] suggests singularity formation in finite time for $\alpha > 0$. In [67] a new scenario with multiple patches with opposite signs colliding in finite time is established for a modified quasi-geostrophic equation in half plane. A finite-time singularity criterion have been recently provided in [42].

1.1. Periodic vortex patches solutions. Due to the complexity of the motion and the deformation process that the vorticity undergoes, the dynamics of the boundary is hard to track and very little is known concerning the nontrivial global vortex patches solutions for the $(gSQG)_{\alpha}$ equations. Therefore, It is of important interest to look for the emergence of ordered structures for this Hamiltonian equation. In our context ordered structures refers to relative equilibria corresponding to steady solutions relative to some translating or rotating frame of reference.

Historically, relative equilibria was initiated long time ago for the two-dimensional Euler equations with the work Kirchhoff [65] who discovered that an elliptical patch rotates uniformly about it center of mass with constant angular velocity. Variational characterization as critical points of some functional energy was formulated by Kelvin. More implicit examples of uniformly rotating vortex patches with higher symmetry, called V-states, were numerically computed by Deem and Zabusky in [33]. Later, an analytical proof was given by Burbea in [20] and based on the conformal mapping parametrization combined with some tools from local bifurcation theory. Burbea's branches of solutions were extended to global ones in [54]. The regularity and the convexity of the V-states have been investigated in [23, 54, 61].

Similar research has been recently carried out for the $(gSQG)_{\alpha}$ equations. The construction of simply connected V-states was established in [52] for $\alpha \in (0, 1)$ using Burbea's approach. In particular, the bifurcation from the unit disc occurs at the angular velocities,

$$\widetilde{\Omega}_{m}^{\alpha} \triangleq \frac{\Gamma(1-\alpha)}{2^{1-\alpha}\Gamma^{2}(1-\frac{\alpha}{2})} \Big(\frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(2-\frac{\alpha}{2})} - \frac{\Gamma(m+\frac{\alpha}{2})}{\Gamma(m+1-\frac{\alpha}{2})} \Big), \quad m \in \mathbb{N} \setminus \{0,1\}.$$
(1.3)

The remaining case $\alpha \in [1, 2)$ was completely solved by Castro, Córdoba and Gómez-Serrano in [22] and the boundary regularity was discussed in [22, 23].

We point out that a rich investigation has been conducted in the past few years around different topological structures for the V-states and for various nonlinear transport equations. For instance, we mention the existence results of rotating multiply-connected patches [34, 36, 47, 59, 62, 74]. It is shown in particular that rotating patches bifurcate from the annulus and two different branches are assigned to the same symmetry. In addition, formation of small loops are discovered when the two branches are close enough. Boundary effects on the emergence of the V-states were analyzed through the disc example in [35], with important numerical experiments putting in evidence the oscillation of the Burbea's curves. Note also that for Euler equations a second bifurcation of countable branches from the ellipses occurs but the shapes have in fact less symmetry and being at most two-folds. The proof of the existence and analyticity of the boundary has been investigated in [22, 58].

Bifurcation of V-states for the quasi-geostrophic shallow-water equation was accomplished in [37] supplemented with numerical experiments describing the imperfect bifurcation with respect to Rossby deformation length. Another delicate subject related to the construction of non homogeneous periodic solutions around specific radial solutions has been developed in [24, 45]. This offers an excellent starting point to explore whether periodic solutions can be captured around generic radial profiles. The rigidity of radial symmetry properties with respect to the angular velocity was explored in a series of works [40, 48, 57]. Another connected subject is the desingularization of the point vortex system to equilibria of vortex patches/smooth profiles type. This was first studied

by Turkington [80] for Euler equation using variational arguments. Following the same approach Turkington's result was extended for the $(gSQG)_{\alpha}$ equations with $\alpha \in (0, 1)$ in [46]. We notice that this approach does not seem to be efficient to describe neither the topological structure of the patches (for example whether they are connected or not) nor the regularity of the boundary. In [60], the second author and Mateu gave a direct proof showing for $\alpha \in [0, 1)$ the existence of co-rotating and counter-rotating pairs of simply connected smooth patches, using a desingularization of the contour dynamics equations and an application of the implicit function theorem. This approach sounds to be flexible and robust and has been adapted recently by different authors to cover various interesting point vortex configurations associated to multiple models. For instance, it was used by [21] to extend the construction for the case $\alpha \in [1, 2)$ and in [53] for the desingularization of the asymmetric pairs. The same technique was used to desingularize a spatial periodic distribution called Karman vortex street [43] and a similar study was performed for the Thomson polygon [44]. Very recently, a generic system of rotating point vortices was analyzed in [55] and a weak condition on the validity of the desingularisation through the contour dynamics has been found.

1.2. Quasi-periodic solutions. It is worthy to point out that $(gSQG)_{\alpha}$ is a reversible Hamiltonian system and as we shall see in Section 2.3 this structure persists at the level of the contour dynamics equation in the setting of the vortex patches. This system depends on one degree of freedom given by the external parameter α related to the order of the nonlocal operator. Then it is legitimate to explore whether quasi-periodic solutions constructed for the linearized operator at the equilibrium state (obtained by linear superposition of its eigenfunctions) could survive under small perturbation for the nonlinear model when α is selected in a suitable Cantor set.

To fix the terminology, a real-valued function $f : \mathbb{R} \to \mathbb{R}$ is called quasi-periodic (shortened in QP) if there exists a multi-variable function $F : \mathbb{T}^d \to \mathbb{R}$, with $d \ge 1$ such that

$$\forall t \in \mathbb{R}, \quad f(t) = F(\omega t)$$

for some frequency vector $\omega \in \mathbb{R}^d$ which is non-resonant in the sense

$$\forall l \in \mathbb{Z}^d \setminus \{0\}, \quad \omega \cdot l \neq 0, \tag{1.4}$$

where we denote by \mathbb{T}^d the flat torus of dimension d. In the case d = 1, we recover the definition of periodic functions with frequency $\omega \in \mathbb{R}^*$. The persistence of invariant torus is a relevant subject for nearly integrable Hamiltonian systems in finite or infinite dimension spaces. In general, the construction stems from KAM theory whose main query is to device a scheme allowing to avoid resonances that may destroy the invariant torus (which may occur even at the linear level). This is an active area which has been improved and enriched through different important studies along the past few decades.

The main task in this paper is to explore the existence of quasi-periodic solutions for (1.1) in the setting of vortex patches. Even though several equations were subject to KAM studies, it seems that the vortex motion which is an old topic in fluid dynamics has escaped to these studies except for the periodic framework where a lot of results have been obtained as we have mentioned before. One of the main advantage in getting QP solutions is the construction of non trivial global in time solutions around the stationary Rankine vortices for the singular model (1.1).

The literature related to KAM theory is very abundant and substantial progress connected to various geometric and analytical aspects has been accomplished during the past decades. Here, we shall only focus on some specific contributions fitting with the main scope of the paper. This theory was initiated by Kolmogorov [68], Arnold [3] and Moser [72] who proved in finite dimensional space the persistence of invariant tori for small perturbation of integrable hamiltonian systems under suitable non degeneracy and smoothness conditions. Later, more development around lower dimensional elliptic/hyperbolic invariant tori for reversible systems. The extension of KAM theory to infinite dimensional Hamiltonian systems with applications to PDE's like the wave, Schrödinger

and Klein Gordon equations was implemented by Kuksin [69], Wayne [82], Bourgain [18], Kuksin-Pöschel [70], Eliasson-Kuksin [38], Grébert-Kappeler [50], Grébert-Paturel [51], Baldi-Berti-Haus-Montalto [4]. More recent studies can be found in [7, 8, 9, 16, 10].

The complexity of the problem depends on the space dimension and on the structure of the equations. For example in the semi-linear case the nonlinearity can be seen as a bounded perturbation of the linear problem and this simplifies a lot the problem of finding a right approximate inverse of the linearized operator around a state close to the equilibrium. However in the quasi-linear case where the nonlinearity is unbounded and has the same order as the linear part the situation turns to be much more tricky. This is the case for instance in the water-waves equations where several results has been obtained in the past few years on the periodic and quasi-periodic settings [1, 4, 13, 14, 16, 64]. For similar discussion we refer to [5] for the 3D incompressible Euler equations with quasi-periodic forcing and to [15] for the vortex patches solutions to the 2D Euler equations.

Next we shall make general comments on the scheme commonly used to construct quasi-periodic solutions for semi-linear or quasi-linear PDE's that was developed in particular in the papers [4, 9, 16]which is robust and flexible and will be adapted in our framework up to some important technical problems that have been settled differently. The first step is to write down the dynamics according to the action-angle coordinates used to describe the tangential part without affecting the normal part which lives in an infinite dimensional space. Notice that the tangential part is composed by a finite number of excited frequencies satisfying non-resonance condition that constitute the linear non-resonant torus. In linearizing around a state near the equilibrium, one finds an operator with variable coefficients that should be inverted approximately up to different types of small errors, provided that the external parameters belong to a suitable Cantor set defined through various Diophantine conditions. To do that it is, first, convenient to look for an approximate inverse using an intermediate isotropic torus built around the initial one. It has the advantage to transform the linearized operator via symplectic change of coordinates into a triangular system up to an error vanishing when testing against an invariant torus. This perturbation can be later incorporated in Nash-Moser scheme. Then the outcome is that the Hamiltonian has a good normal form structure such that one can almost decouple the dynamics in the phase space in tangential and normal modes. On the tangential part the system can be solved in a triangular way provided we can invert the linearized operator on the normal part up to a small coupling error term, and this is more or less finite dimensional KAM theory. Then, the analysis reduces to invert the linearized operator on the normal part, which can be viewed as a small bounded/unbounded perturbation of a diagonal operator. The main ingredient to do that is to conjugate in a suitable way the linearized operator into a diagonal one with constant coefficients that one could invert. This is the major step which turns out to be highly technical due to the resonances which emerge at all the scales and one should use the degree of freedom of the system in order to avoid it during the KAM scheme by making consecutive excisions ending with a Cantor like set. This allows to build an approximate right inverse to the linearized operator with nice tame estimates that we can combine with a frequency cut-off of the Nash-Moser scheme.

1.3. Main result and ideas of the proof. As we shall discuss later in Section 2.1, the contour dynamics equation describing the vortex patch motion can be written in a more tractable way using polar coordinates. This description is meaningful at least for a short time when the initial patch is sufficiently close to the equilibrium state given by Rankine vortex $\mathbf{1}_{\mathbb{D}}$, where \mathbb{D} is the unit disc of the plane. Thus the boundary ∂D_t can be parametrized as follows

$$w(t,\theta) = R(t,\theta)e^{i\theta} \quad \text{with} \quad R(t,\theta) = (1+2r(t,\theta))^{\frac{1}{2}}.$$
(1.5)

As we shall prove in (2.12), the function r satisfies the following nonlinear transport equation

$$\partial_t r + F_\alpha[r] = 0, \tag{1.6}$$

with

$$F_{\alpha}[r](t,\theta) = \frac{C_{\alpha}}{2\pi} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[\left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} \left(1 + 2r(t,\eta)\right)^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta$$
(1.7)

and

$$A_r(\theta,\eta) = \left| R(t,\theta)e^{\mathrm{i}\theta} - R(t,\eta)e^{\mathrm{i}\eta} \right|.$$

Next, we take a parameter $\Omega \neq 0$ and look for the solutions in the form

$$r(t,\theta) = \tilde{r}(t,\theta + \Omega t), \qquad (1.8)$$

then the equation (1.6) is equivalent to

$$\partial_t \tilde{r} + \Omega \partial_\theta \tilde{r} + F_\alpha[\tilde{r}] = 0. \tag{1.9}$$

Notice that the introduction of the parameter Ω sounds at this level artificial but it will be used later to cancel the trivial degeneracy of the first eigenvalue associated with the linearized operator at the equilibrium state. In the quasi-periodic setting, we should find a non-resonant vector frequency $\omega \in \mathbb{R}^d$ such that the equation (1.9) admits a solution in the form $\tilde{r}(t,\theta) = \hat{r}(\omega t,\theta)$ with $\hat{r}: \mathbb{T}^d \times \mathbb{T} \to \mathbb{R}$ being a smooth $(2\pi)^{d+1}$ -periodic function. Then we can easily check that \hat{r} , still denoted in what follows by r, satisfies

$$\omega \cdot \partial_{\varphi} r + \Omega \partial_{\theta} r + F_{\alpha}[r] = 0$$

The computation of the linearized operator of (1.9) at a given state r is described in Proposition 3.1 and one gets

$$\partial_t h(t,\theta) = \partial_\theta \Big[- \big(\Omega + V_{r,\alpha}(t,\theta)\big) h(t,\theta) + \mathbb{K}_{r,\alpha} h(t,\theta) \Big]$$

where $V_{r,\alpha}(t,\theta)$ is the real function

$$V_{r,\alpha}(t,\theta) \triangleq \frac{C(\alpha)}{2\pi} \int_0^{2\pi} \frac{\partial_\eta \left[\left(1 + 2r(t,\eta)\right)^{\frac{1}{2}} \sin(\eta - \theta) \right]}{\left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} A_r^{\alpha/2}(\theta,\eta)} d\eta$$

and the integral operator $\mathbb{K}_{r,\alpha}$ is defined by

$$\mathbb{K}_{r,\alpha}h(t,\theta) \triangleq \frac{C(\alpha)}{2\pi} \int_0^{2\pi} \frac{h(t,\eta)}{A_r^{\alpha/2}(\theta,\eta)} d\eta$$

At the equilibrium state $r \equiv 0$, we infer from Proposition 3.2 that the linearized operator is a Fourier multiplier, since

$$h(\theta) = \sum_{j \in \mathbb{Z} \setminus \{0\}} h_j e^{ij\theta}, \quad \partial_\theta \Big[- \big(\Omega + V_{0,\alpha}\big) + \mathbb{K}_{0,\alpha} \Big] h(\theta) = -\sum_{j \in \mathbb{Z} \setminus \{0\}} \Omega_j(\alpha) h_j e^{ij\theta}$$
(1.10)

where the frequency Ω_j is defined by

$$\Omega_{j}(\alpha) \triangleq j \left(\Omega + \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{2\sqrt{\pi}\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})} \right) \right)$$

$$= j \left(\Omega + \widetilde{\Omega}_{j}^{\alpha} \right) \cdot$$
(1.11)

Here, Γ stands for the usual Gamma function and $\widetilde{\Omega}_{j}^{\alpha}$ was introduced before in (1.3). Consequently, the elements with zero average of the kernel of the linearized operator at the equilibrium state are given by

$$h(t,\theta) = \sum_{j \in \mathbb{Z} \setminus \{0\}} h_j e^{i(j\theta - \Omega_j(\alpha)t)},$$

supplemented with a suitable decay property of the Fourier coefficients to rend the sum meaningful. Since the Hamiltonian system is reversible then we can use this property to filter trivial resonances coming from opposite frequencies by looking only for the real solutions which are invariant by involution, that is, $h(-t, -\theta) = h(t, \theta)$). Therefore we are led to consider only solutions in the form

$$h(t,\theta) = \sum_{j \in \mathbb{N} \setminus \{0\}} h_j \cos\left(j\theta - \Omega_j(\alpha)t\right).$$

Then, by keeping only a finite number of frequencies, this sum gives rise to quasi-periodic solutions with non-resonant frequency provided that α belongs to a suitable set defined with Diophantine condition, for a precise statement see Lemma 3.3. Our main result concerns the persistence of quasi-periodic solutions for the nonlinear model (1.9) when the perturbation is small enough.

Theorem 1.1. Let $\Omega > 0$, $0 < \underline{\alpha} < \overline{\alpha} < \frac{1}{2}$, $\mathbb{S} \subset \mathbb{N} \setminus \{0\}$, with $\#\mathbb{S} = d \ge 1$. There exists $\varepsilon_0 \in (0, 1)$ small enough with the following properties: For every amplitude $\vec{\varepsilon} = (\varepsilon_i)_{i \in \mathbb{S}} \in (\mathbb{R}^*_+)^d$ satisfying

$$\left|\vec{\varepsilon}\right| \leqslant \varepsilon_{0}$$

there exists a Cantor-like set $C_{\infty} \subset (\underline{\alpha}, \overline{\alpha})$ with asymptotically full Lebesgue measure as $\vec{\varepsilon} \to 0$, i.e.,

$$\lim_{\vec{\varepsilon} \to 0} |\mathsf{C}_{\infty}| = \overline{\alpha} - \underline{\alpha}$$

such that for any $\alpha \in C_{\infty}$, the equation (1.9) admits a time quasi-periodic solution with diophantine frequency vector $\omega_{pe}(\alpha, \vec{\epsilon}) \triangleq (\omega_j(\alpha, \vec{\epsilon}))_{j \in \mathbb{S}} \in \mathbb{R}^d$ and taking the form

$$\tilde{r}(t,\theta) = \sum_{j \in \mathbb{S}} \varepsilon_j \cos\left(j\theta + \omega_j(\alpha, \vec{\varepsilon})t\right) + p\left(\omega_{\rm pe}t, \theta\right)$$

with

$$\omega_{\mathrm{pe}}(\alpha, \vec{\varepsilon}) \xrightarrow[\vec{\varepsilon} \to 0]{} (-\Omega_j(\alpha))_{j \in \mathbb{S}}$$

and $\Omega_i(\alpha)$ is defined in (1.11). In addition, the perturbation p satisfies

$$\|\mathbf{p}\|_{H^s_{\mathrm{even}}(\mathbb{T}^{d+1},\mathbb{R})} \stackrel{=}{_{\vec{\varepsilon}\to 0}} o(|\vec{\varepsilon}|)$$

for s large enough, where the Sobolev spaces $H^s_{even}(\mathbb{T}^{d+1},\mathbb{R})$ are defined in (4.2).

Some remarks are in order.

Remarks 1.1. (i) From this theorem, (1.8) and (1.5) one gets that the boundary shape of the quasi-periodic patch can be parametrized in polar coordinates as follows

 $w(t,\theta) = R(t,\theta)e^{i\theta}$ with $R(t,\theta) = (1+2\tilde{r}(t,\theta+\Omega t))^{\frac{1}{2}}$

and \tilde{r} is expanded as in the theorem. The time evolution of the shape is given by small pulsation around the unit disc and the boundary is localized in an annulus around the unit circle.

- (ii) We obtain global existence of non trivial solutions in the patch form for (1.1) where only a few results on global existence are known in the periodic case, see [2, 22, 23, 24, 35, 52] and the references therein.
- (iii) Some technical problems are behind the limitation of $\overline{\alpha} > 0$ and we cannot go to $\underline{\alpha} = 0$. This is connected with the fact that in the fractional Laplacian (1.2) associated to Riesz potential, the constant $C(\alpha)$ blows up when α goes to zero. We know that this operator converges in a weak sense to $(-\Delta)^{-1}$ whose kernel is of logarithmic type and this change of behavior sounds to be the main obstruction in our result. The second limitation concerns the upper bound $\overline{\alpha}$ which should be smaller than $\frac{1}{2}$. This is related first to the reducibility scheme for the fractional Laplacian part. Second, this limitation is needed in the proof of the refined Egorov theorem in Section 8. We believe that the method developed here could handle the remaining case $\overline{\alpha} \in [\frac{1}{2}, 1)$ but the load is very high due to several nonlocal commutators with positive order that should be reduced to constant coefficients one by one. In addition an adaptation of Egorov type theorem is also required

We shall now sketch the main steps of the proof which will be implemented following standard KAM scheme in the spirit of the preceding works [16, 8] but with different substantial variations as we shall discuss below. We basically use techniques from KAM theory, pseudo-differential operators combined with Nash Moser scheme. This will be done in several steps which are detailed below.

• Action-angle reformulation. We first notice that the equation (1.6) enjoys a Hamiltonian structure and from Proposition 2.1 we may write the contour dynamics equation in the form

$$\partial_t r = \partial_\theta \nabla H(r) \tag{1.12}$$

and the Hamiltonian H can expressed in terms of the kinetic energy of the system and the angular momentum. Close to Rankine vortices, we can write the PDE as a Hamiltonian perturbation of an integrable system given by the linear dynamics at the equilibrium state. Indeed, according to the expansion (1.10) the linearized operator at the equilibrium is given by a Fourier multiplier which gives rise to an integrable system and therefore we can write (1.12) in the form

$$\partial_t r = \partial_\theta \mathcal{L}(\alpha)(r) + X_P(r)$$

where $L(\alpha)$ and the perturbed Hamiltonian vector field X_P are defined by

$$\mathcal{L}(\alpha)(r) = -(\Omega + V_{0,\alpha})r + \mathbb{K}_{0,\alpha}r \quad \text{and} \quad X_P(r) = V_{0,\alpha}\partial_\theta r - \partial_\theta \mathbb{K}_{0,\alpha}r - F_\alpha[r].$$

We point out that $V_{0,\alpha}$ is a constant and the operator $\mathbb{K}_{0,\alpha}$ is a Fourier multiplier. Then we find it convenient to rescale the solution size in the following way $r \rightsquigarrow \varepsilon r$, with ε a small parameter. Thus we derive a new equation that appears as a perturbation of the linearized equation at the equilibrium state. Actually, we find that the new equation takes the form

$$\partial_t r = \partial_\theta \mathcal{L}(\alpha)(r) + \varepsilon X_{P_\varepsilon}(r)$$

where $X_{P_{\varepsilon}}$ is the Hamiltonian vector field defined by $X_{P_{\varepsilon}}(r) \triangleq \varepsilon^{-2}X_{P}(\varepsilon r)$. Then, time quasiperiodic solutions oscillating at the frequency $\omega \in \mathbb{R}^{d}$ are simply periodic solutions to the following equation

$$\omega \cdot \partial_{\varphi} r = \partial_{\theta} \mathcal{L}(\alpha)(r) + \varepsilon X_{P_{\varepsilon}}(r).$$

Here we keep the same notation r for the new profile which depends on the variables $(\varphi, \theta) \in \mathbb{T}^{d+1}$. At this stage, we split the phase space into two parts: the tangential part given by the finite dimension space $\mathbb{H}_{\overline{S}}$ and the normal one $\mathbb{H}_{\overline{S}_0}^{\perp}$, defined in (5.8). The dynamics on the tangential space will be parametrized through the action-angle variables (I, ϑ) leading to a new reformulation through the embedded torus. Indeed, by virtue of (5.12) we may decompose r as follows

$$r(\varphi, \theta) = \underbrace{v(\vartheta, I)(\varphi, \theta)}_{\in \mathbb{H}_{\overline{\mathbb{S}}}} + \underbrace{z(\varphi, \theta)}_{\in \mathbb{H}_{\overline{\mathbb{S}}_{0}}},$$

with

$$v(\vartheta, I)(\varphi, \theta) \triangleq \sum_{j \in \overline{\mathbb{S}}} \sqrt{\mathbf{a}_j^2 + \frac{|j|}{2\pi} I_j(\varphi)} \, e^{\mathrm{i}(\vartheta_j(\varphi) + j\theta)}$$

In this way, we view r as an embedded torus

$$i: \mathbb{T}^{d} \to \mathbb{T}^{d} \times \mathbb{R}^{d} \times \mathbb{H}^{\perp}_{\mathbb{S}_{0}}$$

$$\varphi \mapsto (\vartheta(\varphi), I(\varphi), z(\varphi)).$$

$$(1.13)$$

Therefore, in the new coordinates system the problem reduces to finding an invariant torus with non-resonant frequency vector ω such that

$$\omega \cdot \partial_{\varphi} i(\varphi) = X_{H_{\varepsilon}}(i(\varphi)), \qquad (1.14)$$

where the vector field $X_{H_{\varepsilon}}$ is associated to the new Hamiltonian H_{ε} given in (5.18) by

$$H_{\varepsilon} = -\omega_{\mathrm{Eq}}(\alpha) \cdot I + \frac{1}{2} \langle \mathrm{L}(\alpha) z, z \rangle_{L^{2}(\mathbb{T})} + \varepsilon \mathscr{P}_{\varepsilon} \quad \text{and} \quad \omega_{\mathrm{Eq}}(\alpha) = \left(\Omega_{j}(\alpha)\right)_{j \in \mathbb{S}}$$

We remind that the frequencies $\Omega_j(\alpha)$ are defined in (1.11). Instead of solving the equation (1.14) we shall first solve the relaxed problem

$$\omega \cdot \partial_{\varphi} i(\varphi) = X_{H^{\mathsf{c}}_{\varepsilon}}(i(\varphi)),$$

where the vector field $X_{H_{\varepsilon}^{c}}$ is associated to the modified Hamiltonian H_{ε}^{c} given in (5.20) by

$$H_{\varepsilon}^{\mathsf{c}} = \mathsf{c} \cdot I + \frac{1}{2} \langle \mathcal{L}(\alpha) z, z \rangle_{L^{2}(\mathbb{T})} + \varepsilon \mathscr{P}_{\varepsilon}.$$

We emphasize that the advantage of proceeding in this way is to get one degree of freedom with the vector **c** that will be fixed later to ensure some compatibility assumptions in finding an approximate inverse of the linearized operator. At the end of Nash-Moser scheme, we will adjust the frequency ω in such a way that **c** coincides with the equilibrium frequency $\omega_{Eq}(\alpha)$ and thereby we get a solution to the original Hamiltonian equation. To find solutions to the relaxed problem it suffices to construct zeros (i, c) to the nonlinear functional,

$$\mathcal{F}(i, \mathbf{c}, \omega, \alpha, \varepsilon) \triangleq \omega \cdot \partial_{\varphi} i(\varphi) - X_{H_{\varepsilon}^{\mathbf{c}}}(i(\varphi)) \\ = \begin{pmatrix} \omega \cdot \partial_{\varphi} \vartheta(\varphi) - \mathbf{c} - \varepsilon \partial_{I} \mathcal{P}_{\varepsilon}(i(\varphi)) \\ \omega \cdot \partial_{\varphi} I(\varphi) + \varepsilon \partial_{\theta} \mathcal{P}_{\varepsilon}(i(\varphi)) \\ \omega \cdot \partial_{\varphi} z(\varphi) - \partial_{\theta} \left(\mathcal{L}(\alpha) z(\varphi) + \varepsilon \nabla_{z} \mathcal{P}_{\varepsilon}(i(\varphi)) \right) \end{pmatrix}$$
(1.15)

for given $(\omega, \alpha, \varepsilon) \in \mathbb{R}^d \times [\underline{\alpha}, \overline{\alpha}] \times (0, 1)$. We remark that the flat torus $i_{\text{flat}}(\varphi) = (\varphi, 0, 0)$ is a trivial solution in the particular case $\varepsilon = 0$,

$$\mathscr{F}(i_{\text{flat}}, -\omega_{\text{Eq}}(\alpha), -\omega_{\text{Eq}}(\alpha), \alpha, 0) = 0.$$

Then at this point we are tempted to apply the classical Implicit Function Theorem but unfortunately it does not work because the linearized operator at the equilibrium state is not invertible due to the small divisors problem. We can remedy to this defect by imposing suitable non-resonance conditions on the frequency ω and show that this linearized operator admits at least a right inverse but with loss of regularity. Then the main challenge is to extend this property for the linearized operator associated to any arbitrary small state near the flat torus which turns out to be no longer diagonal and admits variable coefficients affecting the main part of its symbol. This is considered as the main step in the implementation of Nash-Moser scheme. Inverting this operator requires careful attention and delicate analysis of the resonances set and one needs to set up some refined tools from toroidal pseudo-differential operators.

2 Approximate right inverse of the linearized operator. The structure of the linearized operator of the functional \mathcal{F} is given by a complicated linear combination of the tangential and normal parts interacting through variables coefficients and it seems to be out of reach to invert it in only one step. Then the approach developed by Berti and Bolle in [9], which is robust and has been performed in several contexts through different papers [4, 7, 16], consists in linearizing first the functional around an isotropic torus sufficiently close to the original one and then proceed with a canonical conjugation through a symplectic change of coordinates leading to a triangular system (decoupling the tangential and the normal parts) up to small decaying errors, essentially of "type Z" or highly decaying in frequency, that can be incorporated in Nash-Moser scheme. As a by-product of this formalism, to invert this latter triangular system it suffices to get an approximate right inverse for the linearized operator in the normal direction, denoted in what follows by $\widehat{\mathscr{L}}_{\omega}$. We will see in Section 6 that we can bypass the use of isotropic torus using exactly the same formalism. Actually, according to Proposition 6.1, we can conjugate the linearized operator with the transformation described by (6.5)and find at the end a triangular system with small errors mainly of "type Z". The computations are done directly in a straightforward way and where only the Hamiltonian structure of the original system sounds to be crucial to get the final triangular structure. Now, let us emphasize that the transformation (6.5) is not symplectic but this does not matter because, first we are not interested in the persistence of the nonlinear Hamiltonian structure but simply concerned with its linear level. Second, this transformation is almost symplectic up to errors of "type Z". The main advantage with this is to require the invertibility only for the linearized operator at the torus itself and not on a closer isotropic one. By this way, we can avoid the accumulation of different extra errors induced by the isotropic torus that one encounters for example in the estimates of the approximate inverse or in the multiple Cantor sets elaborated along the different reduction steps where the coefficients should be computed at the isotropic torus. Therefore the outcome of this first step is to transform the invertibility problem of the full operator to simply invert the partial one resulting from the normal part and taking, according to Proposition 9.1 and (9.3), the form

$$\widehat{\mathscr{L}}_{\omega} = \Pi_{\mathbb{S}_0}^{\perp} \left(\mathscr{L}_{\varepsilon r, \lambda} - \varepsilon \partial_{\theta} \mathscr{R} \right) \Pi_{\mathbb{S}_0}^{\perp} \quad \text{with} \quad \mathscr{L}_{r, \lambda} h = \omega \cdot \partial_{\varphi} h + \partial_{\theta} \left(V_{r, \alpha}(\varphi, \theta) h - \mathbb{K}_{r, \alpha} h \right)$$

where $\Pi_{\mathbb{S}_0}^{\perp}$ is the orthogonal projection on the normal frequency phase space and \mathscr{R} is a smoothing integral operator in the space variable θ with finite rank coming from the interaction between the tangential and the normal parts induced by the transformation (6.5). By virtue of Lemma 9.1, one has the asymptotic structure,

$$\mathscr{L}_{r,\lambda} = \omega \cdot \partial_{\varphi} + \partial_{\theta} \Big[V_{r,\alpha} - \left(\mathscr{W}_{r,\alpha} \left| \mathbf{D} \right|^{\alpha - 1} + \left| \mathbf{D} \right|^{\alpha - 1} \mathscr{W}_{r,\alpha} \right) + \mathscr{R}_{r,\alpha} \Big]$$

with $\mathscr{W}_{r,\alpha}$ being a smooth function taking non vanishing constant value at the equilibrium state $r \equiv 0$ and the remainder $\mathscr{R}_{r,\alpha}$ is a pseudo-differential operator of order -2 in the spatial variable. The operator $|D|^{\alpha-1}$ is a modified fractional Laplacian of order $\alpha-1$ and whose kernel representation is detailed in (4.19). At the equilibrium state, $\widehat{\mathscr{L}}_{\omega}$ coincides with $\mathscr{L}_{0,\lambda}$ which is a Fourier multiplier that can be formally inverted with loss of algebraic regularity by imposing suitable Diophantine conditions that can be guaranteed by the parameters (ω, α) which should be in a suitable massive Cantor set. However when r is taken small, the perturbation is propagated everywhere and affects all the positive order parts of the operator, that is, the transport part in terms of $V_{r,\alpha}$ and the nonlocal part (of order α) through $\mathscr{W}_{r,\alpha}$. This situation is common to different models that we can encounter in the literature such as the water waves [1, 4, 16]. Another observation that we want to stress concerns the order of the fractional Laplacian which depends on the exterior parameter α . This is a new difficulty compared to the equations subject to KAM studies, where the exterior parameter affects only the coefficients of the operator but not its order. This fact brings slightly more technical difficulties related to the functional calculus aspects. Now the question, which is a central key point in KAM theory applied for PDE, is how one could invert the linearized operator with variable coefficients. For this aim, the intuitive idea that one could implement is to diagonalize the operator using different types of transformations in order to get it conjugated to a Fourier multiplier as in the integrable case. Here in our case, we distinguish three different reductions related to the transport part, the fractional part and the remainder. Next, we intend to shed the light on the different techniques used to perform the reduction steps and isolate the main technical difficulties where some of them are solved with a new approach.

(a) Reduction of the transport part. This procedure has been discussed recently throughout several papers and consists in finding a suitable quasi-periodic symplectic change of coordinates allowing to conjugate the transport part into a new one with constant coefficients [14, 39]. The results related to this point can be found in Section 9.3. As we shall see in Proposition 9.3 we may find an invertible transformation

$$\mathscr{B}h = (1 + \partial_{\theta}\beta(\varphi, \theta))h(\varphi, \theta + \beta(\varphi, \theta))$$

such that for an arbitrary $n \in \mathbb{N}$ and if the parameter λ is restricted to the truncated set defined through the first order Melnikov condition (the notation i below stands for the embeddings torus parametrization associated to r according to (1.13)

$$\mathcal{C}_{\infty,n}^{\kappa,\tau_1}(i) = \bigcap_{\substack{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}\\|l|\leqslant N_n}} \left\{ \lambda = (\omega,\alpha) \in \mathcal{O}; \ \left|\omega \cdot l + jc(\lambda,i)\right| > 4\kappa^{\varrho} \frac{\langle j \rangle}{\langle l \rangle^{\tau_1}} \right\}$$

we have

$$\mathscr{L}^{0}_{r,\lambda} \triangleq \mathscr{B}^{-1} \mathscr{L}_{\varepsilon r,\lambda} \mathscr{B} = \omega \cdot \partial_{\varphi} + c(\lambda, i) \partial_{\theta} - \partial_{\theta} \Big(\mu_{r,\lambda} |\mathbf{D}|^{\alpha - 1} + \mathbf{D}|^{\alpha - 1} \mu_{r,\lambda} \Big) + \partial_{\theta} \mathscr{R}_{r,\lambda} + \mathbf{E}^{0}_{n}, \quad (1.16)$$

with $N_n = N_0^{(\frac{3}{2})^n}$, $N_0 \ge 2, \kappa, \varrho \in (0, 1), \tau_1 > d$, $c(\lambda, i)$ a constant and \mathbf{E}_n^0 being a linear operator satisfying in particular the decay estimate

$$\|\mathbf{E}_{n}^{0}h\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{q,s_{0}+2}^{q,\kappa}.$$

The norms $\|\cdot\|_{s_0}^{q,\kappa}$ of weighted Sobolev spaces are given in Definition 4.1. As to the number μ_2 , it is connected with the regularity of the torus associated to r and can be taken arbitrary large allowing the remainder \mathbb{E}_n^0 to get sufficient frequency decay and thereby evacuate it in the small errors during the Nash-Moser scheme implemented at the final stage for the construction of the solutions to the nonlinear problem. On the other hand, the open set \mathcal{O} is without any importance at this level but later and due to several constraints it should be an open set containing the equilibrium frequencies, that is,

$$\mathcal{O} = (\underline{\alpha}, \overline{\alpha}) \times B(0, R),$$

where the open ball B(0, R) with radius R contains the equilibrium frequency vector set $\{\omega_{Eq}(\alpha), \alpha \in \mathcal{L}\}$ $[\underline{\alpha}, \overline{\alpha}]$. The function $\mu_{r,\lambda}$ is not constant but it is close to the constant $\mu_{0,\lambda}$ arising from the equilibrium state. The operator $\partial_{\theta} \mathcal{R}_{\varepsilon r,\lambda}$ is not highly smoothing but of order -1 and with small size ε . The construction of the change of coordinates follows the classical KAM scheme as in [14, 39] and consists in writing successive approximations implemented through solving the *homological equations*. This step requires to impose Melnikov first order non-resonance condition for the transport part. The advantage of this scheme is to replace the first transport operator, after conjugation, with a diagonal one up to a small quadratic error but still of order one. Then iterating this procedure allows to asymptotically get rid of the errors and reduce completely the operator to a diagonal one. The cost of this scheme is directly reflected onto the set of parameters which becomes smaller and smaller due to the excision imposed by Melnikov conditions. We point out that the final Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i)$ is constructed from the ultimate coefficient $c(\lambda, i)$ (obtained as the limit of a suitable sequence) and it is truncated in the time frequency. The advantage of manipulating truncated Cantor sets, which are actually open sets, arises at least at two different levels. First, during the Nash-Moser scheme where we need to construct classical extensions in the whole set of parameters by cut-off functions for the approximations. Second, it is crucial when we plan to estimate the final Cantor set from which nonlinear solutions emerge. There, we need some stability of the different Cantor sets generated throughout Nash-Moser scheme. This is essential to show that the final Cantor set is massive and asymptotically with full Lebesgue measure.

The next step is to reduce the first nonlocal part in (1.16) which is a toroidal pseudo-differential operator of order α .

(b) Reduction of the fractional Laplacian part for $\alpha < \frac{1}{2}$. This step will be explored in Section 9.5 where we reduce the fractional Laplacian part in (1.16) to a Fourier multiplier through the use of infinite dimensional hyperbolic flows. To be more precise, we show in Proposition 9.4 that we can construct a family of invertible operators $\Psi(\lambda)$ such that when $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i)$

$$\Psi^{-1}\mathscr{L}_{\varepsilon r,\lambda}\Psi \triangleq \mathscr{L}^{1}_{r,\lambda} \triangleq \omega \cdot \partial_{\varphi} + c(\lambda, i)\partial_{\theta} - \mathbf{m}(\lambda, i)\partial_{\theta}|\mathbf{D}|^{\alpha-1} + \mathscr{R}^{1}_{r,\lambda} + \mathbf{E}^{1}_{n}$$
(1.17)

with $m(\lambda, i)$ a constant close to $m_{0,\lambda}$ obtained at the equilibrium state and $\mathscr{R}_{r,\lambda}^1$ is a pseudodifferential operator of order $2\alpha - 1$ in the spatial variable. The operator error \mathbf{E}_n^1 can be estimated in a similar way to \mathbf{E}_n^0 and will be included in the small remainders in the Nash-Moser scheme. Let us now outline the main ideas of the proof which will be done in the spirit of [4]. Starting from the first reduction, we want to conjugate the operator $\mathscr{L}_{r,\lambda}^0$ defined in (1.16) to a new one whose positive order part is a Fourier multiplier. For this aim, we use an infinite-dimensional flow Φ satisfying the autonomous pseudo-differential hyperbolic equation

$$\begin{cases} \partial_t \Phi = \partial_\theta (\rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \rho) \Phi(t) \triangleq \mathbb{A} \Phi(t), \\ \Phi(0) = \mathrm{Id}. \end{cases}$$

where the undetermined function ρ will be fixed later according to a transport equation. Then using Taylor expansion at the first order for $\Phi(-1)\mathcal{L}_{r,\lambda}^0\Phi(1)$ we find that

$$\Phi(-1)\mathscr{L}^{0}_{r,\lambda}\Phi(1) = \mathscr{L}^{0}_{r,\lambda} + [\mathscr{L}^{0}_{r,\lambda}, \mathbb{A}_{\rho}] + \text{l.o.t}$$
$$= \omega \cdot \partial_{\varphi} + c(\lambda, i)\partial_{\theta} + \partial_{\theta} \Big(\widehat{\mu}_{r,\lambda} |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1}\widehat{\mu}_{r,\lambda}\Big) + \mathscr{R},$$

with \mathcal{R} a remainder containing various terms that we shall comment later and

$$\widehat{\mu}_{r,\lambda} \triangleq \left(\omega \cdot \partial_{\varphi} + c(\lambda, i) \partial_{\theta} \right) \rho - \mu_{r,\lambda}.$$

Therefore the canonical choice consists in solving the following linear transport equation in ρ

$$\left(\omega \cdot \partial_{\varphi} + c(\lambda, i)\partial_{\theta}\right)\rho = \mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}$$

where $\langle \mu_{r,\lambda} \rangle_{\varphi,\theta}$ stands for the average of $\mu_{r,\lambda}$ in both variables. This equation can be solved in the periodic setting provided that λ belongs to the Cantor set $\mathscr{O}_{\infty,\infty}^{\gamma,\tau_1}(i)$. Since we are interested in working with the truncated Cantor set $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i)$ then instead of solving the above transport equation, we simply solve the modified one

$$\left(\omega \cdot \partial_{\varphi} + c(\lambda, i)\partial_{\theta}\right)\rho = \prod_{N_n} \left(\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}\right),$$

where Π_N is a projector localizing in frequency at the modes range $|l| \leq N$, see Lemma 9.2. Then according to this lemma we may solve this equation with a loss of regularity but uniformly in n. Notice that this frequency cut-off induces small errors that can be balanced into the error operator denoted by E_n^1 which admits a fast decaying rate. Let us now discuss some important technical aspects in estimating the remainder \mathcal{R} . By referring to (9.202) and (9.195), one should deal with some remainders taking the form $\Phi(-t)\mathcal{R}_0\Phi(t)$ with \mathcal{R}_0 is typically described through negative order commutators between pseudo-differential operators. The delicate point is to find a suitable operator topology for which the next reduction on the remainder term, detailed in (C), works and is compatible with the KAM scheme. This step requires an adequate topology satisfying tame estimates in the scales of Sobolev spaces and obviously the bounded operator topology given in (7.1) is not well adapted and seems to be too weak for this purpose. Then one common way is to use a reinforced topology on the symbols class of order zero described through (7.10) with its weighted variant in (7.14). This was for example used in the papers [4, 16]. Now with this topology in mind we get the required functional tame estimates as it is indicated in Lemma 7.4. However, and this is the main technical issue in this part, it is not at all clear whether the new remainder \mathscr{R} still belongs to this topology with tame estimates type. To be more concrete, if we take an operator \mathscr{R}_0 whose symbol satisfies (7.14), is it true that the conjugation by the flow $\Phi(-t)\mathscr{R}_0\Phi(t)$ remains in the same class with suitable tame estimates? We point out that the hyperbolic flow $\Phi(t)$ which acts continuously on the classical Sobolev spaces or their weighted version, see Proposition 8.1, does not in general satisfy the constraint (7.14) which sounds to be too strong for it. At a formal level, one has

$$\Phi(t) \sim Op\left(e^{i\rho(\varphi,\theta)\xi|\xi|^{\alpha-1}}\right)$$

then the differentiation in φ or θ will generate an unbounded operator with a loss of α derivative. Therefore any attempt to use the law products turns to be in vain and one should implement refined tools and this is one of our main technical contribution in this paper that will be investigated along Section 8. Actually, in Theorem 8.1, which can be understood as a refined Egorov type theorem, we shall give a positive answer to our question for a large class of pseudo-differential operators satisfying some weak constraints on the symbol structure, which is large enough to include all the terms of \mathcal{R} . The proof of this theorem is based on a new approach that will be performed through several steps based on the kernel dynamics combined with pseudo-differential calculus and some algebraic structures related to Töplitz matrix operators introduced in Subsection 8.2. In what follows, we intend to give some insights on the proof of this theorem in the particular case of the norm $\||\Phi(-t)\mathcal{R}_0\Phi(t)||_{0,s,0}$, for the definition see (7.9). Let \mathcal{K}_t denote the kernel of the operator $\mathscr{R}_1(t) \triangleq \Phi(-t) \mathscr{R}_0 \Phi(t) - \mathscr{R}_0$, then it satisfies the following transport equation, see (8.32),

$$\mathscr{L}\mathcal{K}_t(\varphi,\theta,\eta) \triangleq \partial_t \mathscr{K}_t(\varphi,\theta,\eta) - \big(\mathbb{A}_\theta + \mathbb{A}_\eta\big) \mathscr{K}_t(\varphi,\theta,\eta) + \big[\partial_\eta, \mathscr{T}_\eta\big] \mathscr{K}_t(\varphi,\theta,\eta) = K_0(\varphi,\theta,\eta),$$

with K_0 the kernel associated to $[\mathbb{A}, \mathcal{R}_0]$ and

$$\mathbb{A}_{\theta} = \partial_{\theta} \left(\rho(\varphi, \theta) | \mathbf{D}_{\theta} |^{\alpha - 1} + | \mathbf{D}_{\theta} |^{\alpha - 1} \rho(\varphi, \cdot) \right) \triangleq \partial_{\theta} \mathcal{T}_{\theta}.$$

According to the norm definition (7.9) and the relationship between the symbol and the kernel one has for any $n \in \mathbb{N}$ that

$$\|\mathscr{R}_{1}(t)\|\|_{0,n,0} \lesssim \sum_{i=0}^{n} \left(\|\partial_{\chi}^{i} \mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} + \|\partial_{\varphi}^{i} \mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \right), \quad \partial_{\chi} \triangleq \partial_{\theta} + \partial_{\eta}$$

Let us explain how to get the estimates of $\|\partial_{\chi}^{i} \mathcal{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})}$ stated in (8.70). We first establish the following vectorial equation:

$$\mathcal{L}_{n}\mathcal{X}_{n}(t) = \mathcal{M}_{n}\mathcal{X}_{n}(t) + \mathcal{Y}_{n} \quad \text{with} \quad \mathcal{L}_{n} \triangleq \mathscr{L}_{I_{n+1}}, \quad \mathcal{X}_{n}(t) = \begin{pmatrix} \mathcal{K}_{t} \\ \partial_{\chi}\mathcal{K}_{t} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n}\mathcal{K}_{t} \end{pmatrix}, \quad \mathcal{Y}_{n} = \begin{pmatrix} K_{0} \\ \partial_{\chi}K_{0} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n}K_{0} \end{pmatrix}$$

with I_{n+1} is the identity matrix and \mathcal{M}_n being a lower Töplitz triangular matrix operator

$$\mathcal{M}_n = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots & 0 \\ m_{1,1} & 0 & \dots & \dots & 0 & 0 \\ m_{2,2} & m_{1,2} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ m_{n,n} & m_{n-1,n} & \dots & \dots & m_{1,n} & 0 \end{pmatrix}$$

Notice that the entries of this matrix are themselves pseudo-differential operators with strictly positive order α and therefore the matrix \mathcal{M}_n is not dissipative and we cannot at this stage implement energy estimates. For the structure of the entries we refer to (8.45). The key point is that the matrix \mathcal{M}_n is nilpotent of order n + 1 and enjoying a rich Töplitz structure. Thus the suitable quantities to estimate are given by the iterated vectors $\{\mathcal{M}^k \mathcal{X}_n, k = 0, ..., n\}$ and one gets the equations

$$\mathscr{L}_{n}\mathscr{M}_{n}^{k}\mathscr{X}_{n}(t) = -\sum_{j=0}^{k-1} {\binom{k}{j}} \operatorname{Ad}_{\mathscr{M}_{n}}^{k-j}(\widehat{\mathscr{L}}_{n}) \mathscr{M}_{n}^{j}\mathscr{X}_{n}(t) + \mathscr{M}_{n}^{k+1}\mathscr{X}_{n}(t) + \mathscr{M}_{n}^{k}\mathscr{Y}_{n}.$$

The estimate of the first term of the right-hand side is detailed in Lemma 8.3 where we use in a crucial way the Töplitz structure not only for the matrix \mathcal{M}_n but also for the iterated matrices commutators $\operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\widehat{\mathcal{Q}}_n)$. Notice that we get in particular that the entries of each matrix commutator are themselves commutators of scalar pseudo-differential operators of order α , and therefore they are of negative order acting continuously in L^2 . The tame estimates are very subtle and require more algebraic structure related to the number of vanishing lower sub-diagonals in the commutators combined with refined estimates on the commutators as in Lemma 7.5. As to the estimate of $\mathcal{M}_n^k \mathcal{Y}_n$, it is stated in Lemma 8.5 and it is very involved due to the fact that \mathcal{Y}_n is not smooth and it is singular at the diagonal line $\{\theta = \eta\}$. Then when the Töplitz matrices \mathcal{M}_n^k act on \mathcal{Y}_n we should check that the singular contributions are cancelled as for $\partial_{\chi}^k K_0$, in this latter case the situation is easy since the operator $\partial_{\chi} = \partial_{\theta} + \partial_{\eta}$ is local and the diagonal singularity belongs to its kernel. The delicate point is that the coefficients of \mathcal{M}_n^k are very complicated and they are non local pseudo-differential operators and we should prove that the singularity is not affected by the iterated operators. The proof is based on suitable recursive estimates governing the iterated kernels, see Lemma 8.4.

We want to precise that similar problem occurs for gravity-capillary water waves as in [16] where the authors proceed in a different way to deal with $\Re_t \triangleq \Phi(-t)\Re_0\Phi(t)$ based on the construction of an approximate solution to the Heisenberg equation satisfied by $\Re_t \triangleq \Phi(-t)\Re_0\Phi(t)$, that is,

$$\partial_t \mathscr{R}_t = [\mathbb{A}, \mathscr{R}_t].$$

The approximation is given in the spirit of the proof of Egorov theorem by expanding the symbol into a finite sum of symbols with decreasing order associated to iterated commutators. Then each element of this sum belongs to the symbol class topology (7.14), however the error term operator is simply

smoothing in the spatial variable and subject to a loss of regularity in the time variable due to the flow. Then to remedy to this lack of information on the error term the authors introduced in [16] new "abstract" topology classes named \mathscr{D}^k -tame and \mathscr{D}^k -modulo-tame operators, much weaker than (7.14). Compared to our case, we are able in this paper to check that the error operator still belongs to the symbol class topology (7.14), which has not only an interest in itself but it brings different technical simplifications related to the use the functional tools during the next steps. The outcome of this long and technical process is to achieve the reduction (1.17), where the new

The outcome of this long and technical process is to achieve the reduction (1.17), where the new remainder $\mathcal{R}_{r,\lambda}^1$ is a pseudo-differential operator of negative order with a tame estimate in the suitable strong topology over the symbol class. For a precise statement we refer to Proposition 9.4-(ii).

(c) KAM reduction of the remainder. As we have seen in (b), we obtain after two reductions a new operator given by (1.17) whose positive part is diagonal and the remainder is of order zero in all the variables and with small size. Then, in order to diagonalize the new operator up to enough decaying small remainder we shall implement a KAM scheme as for instance in [4, 6, 16]. Notice that before proceeding in that way we should first implement the steps (a) and (b) with the linearized operator restricted on the normal direction, given by (9.228). This step is discussed in Section 9.6, and roughly speaking the localization into the normal direction induces new terms that can be described through finite rank operators with small sizes. More precisely, one gets according to Proposition 9.5 a suitable invertible operators Ψ_{\perp} acting on the normal direction such that on the truncated Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i)$ we have

$$\Psi_{\perp}^{-1}\widehat{\mathscr{L}}_{\omega}\Psi_{\perp} = \left(\omega \cdot \partial_{\varphi} + c(\lambda, i)\partial_{\theta} - m(\lambda, i)\partial_{\theta}|\mathbf{D}|^{\alpha-1}\right)\Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{R}_{r,\lambda}^{2} + \mathbf{E}_{n}^{2} \triangleq \mathscr{L}_{0} + \mathbf{E}_{n}^{2},$$

where the orthogonal projector $\Pi_{\mathbb{S}_0}^{\perp}$ and the set \mathbb{S}_0 are defined in (5.9) and (5.6), respectively. The operator \mathbb{E}_n^2 is similar to \mathbb{E}_n^1 seen in **(b)** and describes different small errors coming from the time truncation of the Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i)$. At this stage we get the following structure

$$\mathscr{L}_0 = \omega \cdot \partial_{\varphi} + \mathscr{D}_0 + \mathscr{R}_0$$

where \mathscr{D}_0 is a diagonal operator and \mathscr{R}_0 is a good pseudo-differential operator of zero order and with small size, and satisfying in addition the reversibility structures. Actually, \mathscr{R}_0 is of order $2\alpha - 1$ in the spatial variable θ . The KAM reduction result is stated in Proposition 9.6 and its ultimate target is to eliminate the remainder \mathscr{R}_0 and transform it into a diagonal part by suitable conjugation of \mathscr{L}_0 . This will be developed in a standard way by constructing successive transformations through the KAM reduction allowing to replace at each step the remainder with a smaller new one provided that we make the suitable parameters excision. This scheme works well if we could solve in a reasonable way the associated homological equation which requires to eliminate resonances at each step by imposing the second order Melnikov condition using the external parameters. The final outcome can be summarized as follows: there exists a reversible invertible linear operator Φ_{∞} defined in the full set of parameters such that on the truncated Cantor set

$$\begin{split} \mathcal{O}_{\infty,n}^{\gamma,\tau_1,\tau_2}(i) &\triangleq \left\{ \lambda = (\omega,\alpha) \in \mathcal{O}_{\infty,n}^{\gamma,\tau_1}(i); \,\forall |l| \leqslant N_n, \, j, j_0 \in \mathbb{S}_0^c, \, (l,j) \neq (0,j_0), \\ \left| \omega \cdot l + \mu_j^\infty(\lambda,i) - \mu_{j_0}^\infty(\lambda,i) \right| > \frac{\kappa \langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \right\} \end{split}$$

we have

$$\Phi_{\infty}^{-1} \mathscr{L}_0 \Phi_{\infty} = \omega \cdot \partial_{\varphi} + \mathscr{D}_{\infty} + \mathsf{E}_n^3 \tag{1.18}$$

where $\mathscr{D}_{\infty} = \left(i\mu_{j}^{\infty}(\lambda, i)\right)_{(l,j)\in\mathbb{Z}^{d}\times\mathbb{S}_{0}^{c}}$ is a diagonal reversible operator and the error operator \mathbb{E}_{n}^{3} is similar to \mathbb{E}_{n}^{2} seen before. In addition, the eigenvalues admit the following asymptotic expansion

$$u_j^{\infty}(\lambda, i) = \Omega_j(\alpha) + j r^1(\lambda, i) + j |j|^{\alpha - 1} r^2(\lambda, i) + r_j^{\infty}(\lambda, i)$$

where Ω_j is defined in (1.11), r^1, r^2 and r_j^{∞} satisfies the frequency decay

$$\max_{k=1,2} |r^k(\lambda,i)| \lesssim \varepsilon \kappa^{-1} \quad \text{and} \quad \sup_{\substack{j \in \mathbb{S}_0^c}} |j|^{1-\epsilon-2\overline{\alpha}} |r_j^{\infty}(\lambda,i)| \leqslant C \varepsilon \kappa^{-2}.$$

Actually, we get more precise estimates related to the parameter differentiation of these perturbed coefficients. In fact, one needs to control their derivatives ∂_{λ}^k for any $0 \leq k \leq q_0 + 1$, where q_0 is the index of non-degeneracy of the equilibrium frequencies constructed in Proposition 3.3. More details will be given later on the measure estimates of the final Cantor set.

The final conclusion of this step is the construction of an approximate inverse for the linearized operator restricted to the normal direction as stated in Theorem 9.1. This yields in turn an approximate inverse for the full linearized operator at an arbitrary state close to the equilibrium, according to Proposition 6.1.

③ Nash-Moser scheme. This is the main purpose of Section 10.1 where we construct solutions for the nonlinear function \mathcal{F} , defined in (1.15), and localized around a finite dimensional linear torus with a small amplitude. We basically follow a modified Nash-Moser scheme as in the papers [4, 12, 16] with slight variations. Notice that the Cantor sets used in getting an approximate inverse of the linearized operator, see Theorem 9.1, are truncated in the time frequency and associated to the final states. We emphasize one time more that working with truncated sets allows to generate classical smooth extensions, with a suitable frequency decay, to the whole set of parameters for the approximations defined a priori on the Cantor sets. Dealing with smooth extensions is useful later in measuring the final Cantor set where we shall implement perturbative arguments based on Rüssmann techniques developed in [77] and extended in [8].

The conclusion of this step is summarized in Corollary 10.1 where solutions to the nonlinear functional (1.15) are constructed provided that the (ω, α) belongs to a final Cantor set given by the intersection of all the intermediate Cantor sets needed during the Nash-Moser scheme. The ultimate point to check, and which achieves the proof of the main theorem, is to show that the final Cantor set is massive.

 \bullet Non-degeneracy and Measure estimates. This is the main goal of Section 10.2 where we prove that the final Cantor set $C_{\infty}^{\kappa,\varepsilon}$ described in Corollary 10.1 is asymptotically with full Lebesgue measure. This set is given by

$$\mathsf{C}^{\kappa,\varepsilon}_{\infty} = \bigcap_{m \in \mathbb{N}} \mathsf{C}^{\kappa}_{m} \quad \text{where} \quad \mathsf{C}^{\kappa}_{m} = \Big\{ \alpha \in (\underline{\alpha},\overline{\alpha}); \ \lambda(\alpha,\varepsilon) \in \mathscr{A}^{\kappa}_{m} \Big\},$$

where the intermediate sets \mathscr{A}_m^{κ} are defined in Proposition 10.1 and constructed along Nash-Moser scheme through non resonance conditions imposed at the approximate torus i_m . To measure this set, we mainly use the stability of the Cantor sets combined with the techniques developed in [8] and [77]. One of the crucial key point is the transversality property stated in Lemma 10.3. It will be first established for the linear frequencies in Proposition 3.3, using the analyticity of the eigenvalues and their asymptotic behavior combined with the poles structure of Gamma function. Then the extension of the transversality assumption to the perturbed frequencies is done using perturbative arguments together with the asymptotic description of the approximate eigenvalues detailed in (10.57), (10.59) and (10.58). Notice that the transversality is connected with the nondegeneracy of the eigenvalues in the sense of the Definition 3.2. For instance, we show that the curve $\alpha \in [0,1) \mapsto (\Omega_{j_1}(\alpha), ..., \Omega_{j_d}(\alpha))$ is not contained in any vectorial plane, that is, if there exists a constant vector $c = (c_1, .., c_d)$ such that

$$\forall \alpha \in [0,1), \quad \sum_{j=1}^{d} c_k \Omega_{j_k}(\alpha) = 0$$

then c = 0. We point out that for most of the equations studied before we may check this property by making Taylor expansion at zero leading to an invertible Vandermonde matrix type. However in the current case the structure of the eigenvalues sounds at the first sight more complicated due to the Gamma quotient $\alpha \in \mathbb{C} \mapsto \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})}$ which has the advantage to be a meromorphic function admitting countable sets of separated poles and zeroes altogether located at the real axis. For instance the zeroes are given by

$$\mathscr{Z}_j \triangleq \big\{ \alpha = 2(n+j+1); \ n \in \mathbb{N} \big\}.$$

Then using the different list of zeroes we find successively $c_d = c_{d-1} = \cdots = c_1 = 0$. In Proposition 10.2, we provide the following lower bound of the Lebesgue measure of the $|C_{\infty}^{\kappa,\varepsilon}|$,

$$\left|\mathsf{C}_{\infty}^{\kappa,\varepsilon}\right| \geqslant \overline{\alpha} - \underline{\alpha} - C\varepsilon^{\eta},$$

for a suitable small number $\eta > 0$. We notice that at this level we make the choice $\kappa = \varepsilon^a$ with a being a sufficiently small number.

2. HAMILTONIAN REFORMULATION

This section is devoted to the contour dynamics equation governing the patch motion. We shall in particular reformulate the equations in the polar coordinates whose validity is guaranteed for any time for periodic or quasi-periodic solutions near the unit circle. As we shall see, this can be transformed into a Hamiltonian reformulation which is essential in the approach used for the construction of quasi-periodic solutions.

2.1. Contour dynamics equation. We denote \mathbb{D} the unit disc of \mathbb{R}^2 endowed with its usual Euclidean structure. Here and in the sequel, we identify \mathbb{C} with \mathbb{R}^2 . In particular, the Euclidean structure of \mathbb{R}^2 is seen in the complex sense through the usual inner product defined for all $z_1 = a_1 + ib_1 \in \mathbb{C}$ and $z_2 = a_2 + ib_2 \in \mathbb{C}$ by

$$z_1 \cdot z_2 \triangleq \langle z_1, z_2 \rangle_{\mathbb{R}^2} = \operatorname{Re}\left(z_1 \overline{z_2}\right) = a_1 a_2 + b_1 b_2.$$

$$(2.1)$$

We point out that the Rankine vortex $\mathbf{1}_{\mathbb{D}}$ (and actually any radial function) is a stationary solution to (1.1). Then to find quasi-periodic vortex patches $t \mapsto \mathbf{1}_{D_t}$ around this trivial solution it is convenient to consider the polar parametrization of the boundary given by:

$$w: \mathbb{R}_{+} \times [0, 2\pi] \mapsto \mathbb{C} \qquad \text{with} \qquad w(t, \theta) \triangleq (1 + 2r(t, \theta))^{\frac{1}{2}} e^{\mathbf{i}\theta}.$$
(2.2)
$$(t, \theta) \mapsto w(t, \theta)$$

Here r is the radial deformation of the patch which is small, namely $|r(t,\theta)| \ll 1$. Taking r = 0 gives a parametrization of the unit circle T. To alleviate the notation we use sometimes

$$R(t,\theta) \triangleq (1+2r(t,\theta))^{\frac{1}{2}}.$$
(2.3)

This particular form of R is required later for the Hamiltonian structure. Let us now write down the contour dynamics equation with the polar coordinates. It is known, see for instance [52], that the particles on the boundary move with the flow and remain at the boundary and therefore in the smooth case one has

$$[\partial_t w(t,\theta) - \mathbf{v}(t,w(t,\theta))] \cdot \mathbf{n}(t,w(t,\theta)) = 0$$

where $\mathbf{n}(t, w(t, \theta))$ is the outward normal vector to the boundary ∂D_t of D_t at the point $w(t, \theta)$. Since one has, up to a real constant of renormalization, $\mathbf{n}(t, w(t, \theta)) = -i\partial_{\theta}w(t, \theta)$, then we get the complex formulation of the contour dynamics motion,

$$\operatorname{Im}\left(\left[\partial_t w(t,\theta) - \mathbf{v}(t,w(t,\theta))\right] \overline{\partial_\theta w(t,\theta)}\right) = 0.$$
(2.4)

The velocity field is given by

$$\mathbf{v}(t,w(t,\theta)) = \frac{C_{\alpha}}{2\pi} \int_{\partial D_t} \frac{1}{|w(t,\theta) - \xi|^{\alpha}} d\xi$$
$$= \frac{C(\alpha)}{2\pi} \int_0^{2\pi} \frac{\partial_{\eta} w(t,\eta)}{|w(t,\theta) - w(t,\eta)|^{\alpha}} d\eta$$
(2.5)

We are interested in patches that are perturbation from the unit disc, therefore, we consider the parametrization

$$w(t,\theta) \triangleq \left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} e^{\mathrm{i}\theta},\tag{2.6}$$

Differentiating $z(t, \theta)$ with respect to t we get

$$\partial_t w(t,\theta) = \left(1 + 2r(t,\theta)\right)^{-\frac{1}{2}} \partial_t r(t,\theta) e^{\mathrm{i}\theta}, \qquad (2.7)$$

and, with respect to the θ ,

$$\partial_{\theta}w(t,\theta) = \left(\left(1 + 2r(t,\theta) \right)^{-\frac{1}{2}} \partial_{\theta}r(t,\theta) + i \left(1 + 2r(t,\theta) \right)^{\frac{1}{2}} \right) e^{i\theta}$$
(2.8)

Combining the two last identities gives

$$\operatorname{Im}\left\{\partial_t w(t,\theta)\partial_\theta \overline{w(t,\theta)}\right\} = -\partial_t r(t,\theta).$$
(2.9)

and

$$\int_{0}^{2\pi} \frac{\operatorname{Im}\left\{\partial_{\eta} w(t,\eta)\partial_{\theta} \overline{w(t,\theta)}\right\}}{|w(t,\theta) - w(t,\eta)|^{\alpha}} d\eta = \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[\left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} \left(1 + 2r(t,\eta)\right)^{\frac{1}{2}} \sin(\eta - \theta)\right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta \cdot \quad (2.10)$$

where

$$A_{r}(\theta,\eta) = 2\Big[1 + r(t,\theta) + r(t,\eta) - (1 + 2r(t,\eta))^{\frac{1}{2}} (1 + 2r(t,\theta))^{\frac{1}{2}} \cos(\eta - \theta)\Big].$$
 (2.11)

Inserting the identities (2.9) and (2.10) into equation (2.4) we get

$$\partial_t r(t,\theta) + \underbrace{\frac{C_{\alpha}}{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[\left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} \left(1 + 2r(t,\eta)\right)^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_r^{\alpha/2}(\theta,\eta)}_{\triangleq F_{\alpha}[r](t,\theta)} d\eta}_{\triangleq 0.$$
(2.12)

we get the vortex patch equation in the polar coordinates

$$\partial_t r(t,\theta) + F_\alpha[r](t,\theta) = 0. \tag{2.13}$$

which is a nonlocal transport nonlinear PDE. We look for solutions of the form

$$r(t,\theta) = \tilde{r}(t,\theta + \Omega t) \tag{2.14}$$

for some $\Omega > 0$. Notice that the introduction of the angular velocity Ω is purely technical and needed later to circumvent the degeneracy of the first frequency of the linearized operator at the equilibrium state. From (2.5) one gets

$$F_{\alpha}[\tilde{r}](t,\theta+\Omega t) = F_{\alpha}[r](t,\theta), \qquad (2.15)$$

Thus, equation (2.13) becomes

$$\partial_t \tilde{r}(t,\theta) + \Omega \partial_\theta \tilde{r}(t,\theta) + F_\alpha[\tilde{r}](t,\theta) = 0.$$
(2.16)

Now, a time quasi-periodic solution of (2.16) is nothing but a solution in the form

$$\tilde{r}(t,\theta) = \hat{r}(\omega t,\theta),$$

where $\widehat{r}: (\varphi, \theta) \in \mathbb{T}^{d+1} \mapsto \widehat{r}(\varphi, \theta) \in \mathbb{R}$ and $\omega \in \mathbb{R}^d$ is a non-resonant vector frequency. Hence in this setting, the equation (2.16) becomes

$$\omega \cdot \partial_{\varphi} \widehat{r}(\varphi, \theta) + \Omega \partial_{\theta} \widehat{r}(\varphi, \theta) + F_{\alpha}[\widehat{r}](\varphi, \theta) = 0.$$

In the sequel, we shall alleviate the notation and denote \hat{r} simply by r and the foregoing equation writes

$$\forall (\varphi, \theta) \in \mathbb{T}^{d+1}, \quad \omega \cdot \partial_{\varphi} r(\varphi, \theta) + \Omega \partial_{\theta} r(\varphi, \theta) + F_{\alpha}[r](\varphi, \theta) = 0.$$
(2.17)

2.2. Conservation laws. This section is devoted to some conservation quantities that will be used to recover the Hamiltonian structure of the dynamical system (2.16). In our context, we shall explore the circulation C, the angular momentum J and the energy E and write them in the patch setting using polar coordinates. Let D(t) be a bounded simply connected region with smooth boundary $\partial D(t)$ and $\omega = \chi_{D(t)}$. We define the circulation C, the angular momentum J and the energy E as follows

$$C(t) \triangleq \int_{D(t)} dA(z), \quad J(t) \triangleq -\frac{1}{2} \int_{D(t)} |z|^2 dA(z) \quad \text{and} \quad E(t) \triangleq \frac{1}{2} \int_{D(t)} \psi(t, z) dA(z), \qquad (2.18)$$

where ψ is the stream function defined by

$$\psi(z) \triangleq -\frac{C(\alpha)}{2\pi} \int_{D(t)} \frac{1}{|z-\zeta|^{\alpha}} dA(\zeta)$$
(2.19)

and dA being the planar Lebesgue measure.

Lemma 2.1. The following assertions hold true.

- (i) The quantities C, J and E are conserved in time.
- (ii) In polar coordinates, they can be expressed in the form

$$\begin{split} C(t) &= \pi + \int_{0}^{2\pi} r(t,\theta) d\theta, \\ J(t) &= -\frac{1}{8} \int_{0}^{2\pi} \left(1 + 2r(t,\theta) \right)^{2} d\theta, \\ E(t) &= -\frac{C(\alpha)}{16\pi \left(1 - \frac{\alpha}{2}\right)^{2}} \int_{0}^{2\pi} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[(1 + 2r(t,\theta))^{\frac{1}{2}} (1 + 2r(t,\eta))^{\frac{1}{2}} \cos(\eta - \theta) \right]}{A_{r}^{\frac{\alpha}{2} - 1}(\eta,\theta)} d\eta d\theta, \\ where \ A_{r}(\theta, \eta) \ is \ defined \ in \ (2, 11) \end{split}$$

where $A_r(\theta, \eta)$ is defined in (2.11)

Proof. (i) This result can be checked by using any flow associated to the velocity field generated by the smooth patches $\mathbf{1}_{D_t}$.

(ii) We first recall Green's formula written in the complex form,

$$\forall z \in \mathbb{C}, \quad \int_D \partial_{\overline{\zeta}} f(\zeta, \overline{\zeta}) dA(\zeta) = \frac{1}{2i} \int_{\partial D} f(\zeta, \overline{\zeta}) d\zeta.$$
(2.20)

In view of (2.18) and (2.20) one has

$$C(t) = \frac{1}{2i} \int_{\partial D(t)} \overline{\zeta} d\zeta, \qquad J(t) = -\frac{1}{8i} \int_{\partial D(t)} |\zeta|^2 \overline{\zeta} d\zeta.$$

From (2.6) we get

$$C(t) = \frac{1}{2i} \int_0^{2\pi} (1 + 2r(t,\theta))^{\frac{1}{2}} e^{-i\theta} \partial_\theta \left[(1 + 2r(t,\theta))^{\frac{1}{2}} e^{i\theta} \right] d\theta$$

= $\pi + \int_0^{2\pi} r(t,\theta) d\theta$.

and

$$J(t) = -\frac{1}{8i} \int_0^{2\pi} (1 + 2r(t,\theta))^{\frac{3}{2}} e^{-i\theta} \partial_\theta \left[(1 + 2r(t,\theta))^{\frac{1}{2}} e^{i\theta} \right] d\theta$$

= $-\frac{1}{8} \int_0^{2\pi} (1 + 2r(t,\theta))^2 d\theta$. (2.21)

As to the energy expression, we shall first reformulate the stream function defined in (2.19) in terms of polar coordinates. Applying (2.20) yields

$$\psi(z) = \frac{\mathrm{i}C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2}\right)} \int_{\partial D(t)} \frac{\zeta - \overline{z}}{|\zeta - z|^{\alpha}} d\zeta.$$
18

Let us now move to the energy E. According to (2.18) and the preceding identity we get

$$E(t) = \frac{iC(\alpha)}{8\pi \left(1 - \frac{\alpha}{2}\right)} \int_{D(t)} \int_{\partial D(t)} \frac{\overline{\zeta} - \overline{z}}{|\zeta - z|^{\alpha}} d\zeta dA(z).$$

Using once again Green's formula (2.20), then we deduce that the contour integral

$$E(t) = \frac{C(\alpha)}{16\pi \left(1 - \frac{\alpha}{2}\right) \left(2 - \frac{\alpha}{2}\right)} \int_{\partial D(t)} \int_{\partial D(t)} \frac{(\overline{\zeta} - \overline{z})^2}{|\zeta - z|^{\alpha}} dz d\zeta.$$

Using (2.6) the last expression becomes

$$E(t) = \frac{C(\alpha)}{16\pi \left(1 - \frac{\alpha}{2}\right)\left(2 - \frac{\alpha}{2}\right)} \int_{0}^{2\pi} \int_{0}^{2\pi} \frac{\left(\overline{z(t,\eta)} - \overline{z(t,\theta)}\right)^{2}}{|z(t,\eta) - z(t,\theta)|^{\alpha}} \partial_{\theta} z(t,\theta) \partial_{\eta} z(t,\eta) d\eta d\theta.$$
(2.22)

From the identity

$$\frac{\left(\overline{z(t,\eta)} - \overline{z(t,\theta)}\right)^2}{|z(t,\eta) - z(t,\theta)|^{\alpha}} \partial_{\theta} z(t,\theta) = -\partial_{\theta} \overline{z(t,\theta)} |z(t,\eta) - z(t,\theta)|^{2-\alpha} - \frac{1}{\left(1 - \frac{\alpha}{2}\right)} \left(\overline{z(t,\eta)} - \overline{z(t,\theta)}\right) \partial_{\theta} |z(t,\eta) - z(t,\theta)|^{2-\alpha}.$$

We may write

$$E(t) = -\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)\left(2-\frac{\alpha}{2}\right)} \int_{0}^{2\pi} \int_{0}^{2\pi} \partial_{\theta} \overline{z(t,\theta)} \partial_{\eta} z(t,\eta) |z(t,\eta) - z(t,\theta)|^{2-\alpha} d\eta d\theta$$
$$- \frac{C(\alpha)}{8\pi \left(1-\frac{\alpha}{2}\right)^{2} \left(2-\frac{\alpha}{2}\right)} \int_{0}^{2\pi} \int_{0}^{2\pi} \left(\overline{z(t,\eta)} - \overline{z(t,\theta)}\right) \partial_{\theta} |z(t,\eta) - z(t,\theta)|^{2-\alpha} \partial_{\eta} z(t,\eta) d\eta d\theta.$$

Integrating the last term by parts gives

$$E(t) = -\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)^2} \int_0^{2\pi} \int_0^{2\pi} |z(t,\eta) - z(t,\theta)|^{2-\alpha} \partial_\theta \overline{z(t,\theta)} \partial_\eta z(t,\eta) d\eta d\theta$$

Since E(t) is a real-valued function then

$$E(t) = -\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)^2} \int_0^{2\pi} \int_0^{2\pi} |z(t,\eta) - z(t,\theta)|^{2-\alpha} \operatorname{Re}\left\{\partial_\theta \overline{z(t,\theta)} \partial_\eta z(t,\eta)\right\} d\eta d\theta.$$

Then, substituting (2.6) in the last expression concludes the proof of the lemma.

2.3. Hamiltonian structure and reversibility. In this section we describe the Hamiltonian structure of the contour dynamic equation (2.12). We emphasize that the Hamiltonian formulation is crucial to approximately reduce the linearized operator to a triangular form using the "approximate inverse" approach that will be developed in Section 6.

Proposition 2.1. The following assertions hold true.

(i) The L^2 -gradient of the angular momentum J is given by

$$\nabla J(r) = -\frac{1}{2} - r(\theta)$$

(ii) The L^2 -gradient of the pseudo energy E is given by

$$\nabla E(r) = \frac{C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2}\right)} \int_0^{2\pi} \frac{1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_\eta \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_r^{\alpha/2}(\theta, \eta)} d\eta,$$

where A_r is given by (2.11).

(iii) The equation (2.12) is Hamiltonian and takes the form

 $\partial_t r = \partial_\theta \nabla H(r)$ with $H \triangleq E + \Omega J.$

Moreover, this equation is reversible with respect to the involution

$$(\mathfrak{S}r)(\theta) \triangleq r(-\theta)$$

namely the Hamiltonian vector field $X_H = \partial_\theta \nabla H$ satisfies

 $X_H \circ \mathfrak{S} = -\mathfrak{S} \circ X_H \,.$

The reversibility property may be equivalently read

$$\left(\partial_t r - X_H\right) \circ \mathscr{S} = -\mathscr{S} \circ \left(\partial_t r - X_H\right),$$

with

$$(\mathscr{G}r)(t,\theta) \triangleq r(-t,-\theta)$$

Thus, if $r(t, \theta)$ is a solution of (2.12) then $(\mathcal{G}r)(t, \theta)$ is also a solution.

Proof. For simplicity we shall denote along this proof $r(t, \theta) = r(\theta)$. (i) Differentiating (2.21) with respect to r in the direction

Differentiating (2.21) with respect to
$$r$$
 in the direction h we get

$$dJ(r)[h] = -\frac{1}{2} \int_{\mathbb{T}} h(\theta) \big(1 + 2r(\theta)\big) d\theta.$$

Then (i) is immediate.

(ii) Recall that the expression of the energy in polar coordinates is given by

$$E = -\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)^2} \int_0^{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[(1+2r(\theta))^{\frac{1}{2}} (1+2r(\eta))^{\frac{1}{2}} \cos(\eta-\theta) \right]}{A_r^{\frac{\alpha}{2}-1}(\eta,\theta)} d\eta d\theta.$$
(2.23)

Differentiating the last expression with respect to r in the direction h gives

$$dE(r)[h] = -\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)^2} \int_0^{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[h(\theta)(1+2r(\theta))^{-\frac{1}{2}}(1+2r(\eta))^{\frac{1}{2}}\cos(\eta-\theta)\right]}{A_r^{\frac{\alpha}{2}-1}(\eta,\theta)} d\eta d\theta$$

$$-\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)^2} \int_0^{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[h(\eta)(1+2r(\theta))^{\frac{1}{2}}(1+2r(\eta))^{-\frac{1}{2}}\cos(\eta-\theta)\right]}{A_r^{\frac{\alpha}{2}-1}(\eta,\theta)} d\eta d\theta$$

$$-\frac{C(\alpha)}{16\pi \left(1-\frac{\alpha}{2}\right)} \int_0^{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[(1+2r(\theta))^{\frac{1}{2}}(1+2r(\eta))^{\frac{1}{2}}\cos(\eta-\theta)\right]}{A_r^{\alpha/2}(\theta,\eta)} dA_r[h](\eta,\theta) d\eta d\theta$$

or A is given by (2.11) and

where A_r is given by (2.11) and

$$dA_r[h](\eta,\theta) = 2h(\theta) + 2h(\eta) - 2\left[h(\theta)\frac{(1+2r(\eta))^{\frac{1}{2}}}{(1+2r(\theta))^{\frac{1}{2}}} + h(\eta)\frac{(1+2r(\theta))^{\frac{1}{2}}}{(1+2r(\eta))^{\frac{1}{2}}}\right]\cos(\eta-\theta) + 2h(\eta) - 2\left[h(\theta)\frac{(1+2r(\eta))^{\frac{1}{2}}}{(1+2r(\theta))^{\frac{1}{2}}}\right]\cos(\eta-\theta) + 2h(\eta) - 2\left[h(\theta)\frac{(1+2r(\theta))^{\frac{1}{2}}}{(1+2r(\theta))^{\frac{1}{2}}}\right]\cos(\eta-\theta) + 2h(\eta) - 2\left[h(\theta)\frac{(1+2r(\theta))^{\frac{1}{2}}}{(1+2r(\theta))^{\frac{1}{2}}}\right]\cos(\eta-\theta) + 2h(\eta) - 2h(\eta) + 2h(\eta)$$

1

By symmetry arguments one may write

$$dE(r)[h] = -\frac{C(\alpha)}{8\pi (1-\frac{\alpha}{2})^2} \int_0^{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 [h(\theta)(1+2r(\theta))^{-\frac{1}{2}}(1+2r(\eta))^{\frac{1}{2}}\cos(\eta-\theta)]}{A_r^{\frac{\alpha}{2}-1}(\eta,\theta)} d\eta d\theta - \frac{C(\alpha)}{4\pi (1-\frac{\alpha}{2})} \int_0^{2\pi} \int_0^{2\pi} h(\theta) \frac{\partial_{\theta\eta}^2 [(1+2r(\theta))^{\frac{1}{2}}(1+2r(\eta))^{\frac{1}{2}}\cos(\eta-\theta)]}{A_r^{\alpha/2}(\theta,\eta)} \times \left(1 - (1+2r(\eta))^{\frac{1}{2}}(1+2r(\theta))^{-\frac{1}{2}}\cos(\eta-\theta)\right) d\eta d\theta.$$

Integrating the first term by parts we get

$$dE(r)[h] = \frac{C(\alpha)}{8\pi \left(1 - \frac{\alpha}{2}\right)} \int_{0}^{2\pi} \int_{0}^{2\pi} h(\theta) \frac{\partial_{\eta} \left[(1 + 2r(\eta))^{\frac{1}{2}} \cos(\eta - \theta) \right]}{(1 + 2r(\theta))^{\frac{1}{2}}} \frac{\partial_{\theta} A_r(\eta, \theta)}{A_r^{\alpha/2}(\theta, \eta)} d\eta d\theta$$
(2.24)

$$-\frac{C(\alpha)}{4\pi(1-\frac{\alpha}{2})} \int_{0}^{2\pi} \int_{0}^{2\pi} h(\theta) \frac{\partial_{\theta\eta}^{2} \left[(1+2r(\theta))^{\frac{1}{2}} (1+2r(\eta))^{\frac{1}{2}} \cos(\eta-\theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} \\ \times \left(1 - \left(1+2r(\eta) \right)^{\frac{1}{2}} \left(1+2r(\theta) \right)^{-\frac{1}{2}} \cos(\eta-\theta) \right) d\eta d\theta \cdot$$

In view of (2.11) one has

$$\partial_{\theta} A_{r}(\eta, \theta) = 2\partial_{\theta} r(\theta) \left(1 - \left(1 + 2r(\theta) \right)^{-\frac{1}{2}} \left(1 + 2r(\eta) \right)^{\frac{1}{2}} \cos(\eta - \theta) \right) - 2 \left(1 + 2r(\theta) \right)^{\frac{1}{2}} \left(1 + 2r(\eta) \right)^{\frac{1}{2}} \sin(\eta - \theta).$$

On the other hand, we have the identity

$$\partial_{\theta\eta}^{2} \left[(1+2r(\theta))^{\frac{1}{2}} (1+2r(\eta))^{\frac{1}{2}} \cos(\eta-\theta) \right] = \partial_{\theta} r(\theta) (1+2r(\theta))^{-\frac{1}{2}} \partial_{\eta} \left[(1+2r(\eta))^{\frac{1}{2}} \cos(\eta-\theta) \right] \\ + (1+2r(\theta))^{\frac{1}{2}} \partial_{\eta} \left[(1+2r(\eta))^{\frac{1}{2}} \sin(\eta-\theta) \right].$$

Inserting the two last identities into (2.24) we find

$$dE(r)[h] = \frac{C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2}\right)} \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} h(\theta) \frac{1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_\eta \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_r^{\alpha/2}(\theta, \eta)} d\eta d\theta \cdot d\theta$$

Consequently,

$$\nabla E(r) = \frac{C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2}\right)} \int_0^{2\pi} \frac{1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_\eta \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_r^{\alpha/2}(\theta, \eta)} d\eta.$$
(2.25)

(iii) Since $H = E + \Omega J$. then by (i) and (ii) we have

$$\nabla H(r) = -\frac{\Omega}{2} \left(1 + 2r(\theta) \right) + \frac{C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2} \right)} \int_0^{2\pi} \frac{1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_\eta \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_r^{\alpha/2}(\theta, \eta)} d\eta.$$

Differentiating the last expression with respect to θ gives

$$\partial_{\theta} \nabla H(r) = -\Omega \,\partial_{\theta} r(\theta) - \frac{C(\alpha)}{4\pi \left(1 - \frac{\alpha}{2}\right)} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[(1 + 2r(\theta))^{\frac{1}{2}} (1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}^{\alpha/2}(\theta, \eta)} d\eta \\ - \frac{\alpha C(\alpha)}{8\pi \left(1 - \frac{\alpha}{2}\right)} \int_{0}^{2\pi} \frac{1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_{\eta} \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}(\eta, \theta)^{\frac{\alpha}{2} + 1}} \partial_{\theta} A_{r}(\eta, \theta) d\eta.$$

Straightforward computations allow to get

$$\left(1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_{\eta} \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right] \right) \partial_{\theta} A_r(\eta, \theta)$$

= $-2 \partial_{\theta\eta}^2 \left[(1 + 2r(\theta))^{\frac{1}{2}} (1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right] A_r(\eta, \theta)$
+ $\left(1 + 2r(\theta) + (1 + 2r(\eta))^{\frac{1}{2}} \partial_{\theta} \left[(1 + 2r(\theta))^{\frac{1}{2}} \sin(\eta - \theta) \right] \right) \partial_{\eta} A_r(\eta, \theta).$

This leads to

$$\partial_{\theta} \nabla H(r) = -\Omega \partial_{\theta} r(\theta) + \frac{(\alpha - 1)C(\alpha)}{4\pi (1 - \frac{\alpha}{2})} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[(1 + 2r(\theta))^{\frac{1}{2}} (1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}^{\alpha/2}(\theta, \eta)} d\eta \\ - \frac{\alpha C(\alpha)}{8\pi (1 - \frac{\alpha}{2})} \int_{0}^{2\pi} \frac{1 + 2r(\theta) + (1 + 2r(\eta))^{\frac{1}{2}} \partial_{\theta} \left[(1 + 2r(\theta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}(\eta, \theta)^{\frac{\alpha}{2} + 1}} \partial_{\eta} A_{r}(\eta, \theta) d\eta.$$

Integrating by parts gives

$$\partial_{\theta}\nabla H(r) = -\Omega \partial_{\theta} r(\theta) - \frac{C(\alpha)}{2\pi} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left[(1+2r(\theta))^{\frac{1}{2}} (1+2r(\eta))^{\frac{1}{2}} \sin(\eta-\theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta.$$
(2.26)

Comparing (2.12), (2.26) concludes the proof of (iii).

(iv) In view of (2.26) one has

$$(\mathfrak{S} \circ X_H)(r)(\theta) = -\Omega \,\partial_\theta r(-\theta) - \frac{1}{2\pi} \int_0^{2\pi} \frac{\partial_{\theta\eta}^2 \left[\left(1 + 2r(-\theta)\right)^{\frac{1}{2}} \left(1 + 2r(\eta)\right)^{\frac{1}{2}} \sin(\eta + \theta) \right]}{A_r(-\theta, \eta)^{\alpha/2}} d\eta \cdot$$

Using the change of variable $\eta \mapsto -\eta$ in the last integral we get

$$(\mathfrak{S} \circ X_H)(r)(-\theta) = -\Omega \,\partial_\theta(\mathfrak{S}r)(\theta) + \frac{1}{2\pi} \int_{-2\pi}^0 \frac{\partial_{\theta\eta}^2 \left[\left(1 + 2(\mathfrak{S}r)(\theta) \right)^{\frac{1}{2}} \left(1 + 2(\mathfrak{S}r)(\eta) \right)^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{\mathfrak{S}r}^{\alpha/2}(\theta, \eta)} d\eta \cdot$$

where we have used the fact that $A_r(-\theta, -\eta) = A_{\mathfrak{S}r}(\theta, \eta)$. Consequently

$$\mathfrak{S} \circ X_H(r)(\theta) = -X_H \circ (\mathfrak{S}r)(\theta).$$
(2.27)

This concludes the proof of the lemma.

The symplectic form induced by the hamiltonian equation

$$\partial_t r = \partial_\theta \nabla H(r) \quad \text{with} \quad H \triangleq E + \Omega J.$$
 (2.28)

is given by

$$\mathscr{W}(r,h) \triangleq \int_0^{2\pi} (\partial_\theta^{-1} r) h d\theta, \quad \partial_\theta^{-1} r = \sum_{j \in \mathbb{Z} \setminus \{0\}} \frac{1}{ij} r_j e^{ij\theta}, \quad \forall r, h \in L^2_0(\mathbb{T}).$$
(2.29)

The corresponding Poisson tensor is $\mathcal{J} \triangleq \partial_{\theta}$, and the Poisson bracket

$$\{F,G\} = \mathcal{W}(X_F, X_G) = \int_0^{2\pi} \nabla F(r) \partial_\theta \nabla G(r) \, d\theta,$$

where ∇F , ∇G denote the L^2 -gradients of the functionals $F, G : L_0^2(\mathbb{T}) \to \mathbb{R}$. Note that the Hamiltonian vector field $X_H = \partial_\theta \nabla H$, associated with the Hamiltonian H, is determined by

$$dH(r)[h] = (\nabla H(r), h)_{L^2(\mathbb{T})} = \mathscr{W}(X_H(r), h), \quad \forall r, h \in L^2_0(\mathbb{T}).$$
(2.30)

3. Linearization and frequencies structure

In this section we compute the linear Hamiltonian PDE obtained from linearizing (2.28) at a given small state r close to the equilibrium. At the latter one, we prove that the linearized operator acts as a Fourier multiplier with symbol related to the gamma functions. We shall also explore some important structures on the equilibrium frequencies related to the monotonicity, the asymptotic behavior for large modes and the non-degeneracy/transversality properties. This last point turns out to be crucial to establish the emergence of linear quasi-periodic solutions for a massive set of exponents $\alpha \in (0, 1)$.

3.1. Linearized operator. We intend to discuss the structure of the linearized operator around a general state close to the equilibrium r = 0. As a byproduct we deduce that at the equilibrium state the linear operator acts as a Fourier multiplier with explicit eigenvalues related to Gamma function. The first main result reads as follows.

Proposition 3.1. The linearized Hamiltonian equation (2.28) at a small function $(t, \theta) \mapsto r$ in the direction $(t, \theta) \mapsto h(t, \theta)$ is given by

$$\partial_t h(t,\theta) = \partial_\theta \Big[- \big(\Omega + V_{r,\alpha}(t,\theta)\big) h(t,\theta) + \mathbb{K}_{r,\alpha} h(t,\theta) \Big]$$

where $V_{r,\alpha}(t,\theta)$ is the real function

$$V_{r,\alpha}(t,\theta) \triangleq \frac{C(\alpha)}{2\pi} \left(1 + 2r(t,\theta) \right)^{-\frac{1}{2}} \int_{0}^{2\pi} \frac{\partial_{\eta} \left[\left(1 + 2r(t,\eta) \right)^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta,$$
(3.1)

the integral operator $\mathbb{K}_{r,\alpha}$ is defined by

$$\mathbb{K}_{r,\alpha}h(t,\theta) \triangleq \frac{C(\alpha)}{2\pi} \int_0^{2\pi} \frac{h(t,\eta)}{A_r^{\alpha/2}(\theta,\eta)} d\eta$$
(3.2)

and $A_r(\theta, \eta)$ is given by (2.11).

Proof. Throughout the proof, and for the sake of simple notation, we remove the time dependency from the functions. The computations of the Gâteaux derivative of the vector field ∇E , given by Proposition 2.1-(ii), at the point r in the direction h are straightforward and standard and we shall only sketch the main lines. Notice that the functional is smooth in a suitable functional setting and therefore its Frechet differential can be recovered from its Gâteaux derivative. From direct computations one gets

$$d\nabla E(r)h(\theta) = \frac{C(\alpha)}{4\pi(1-\frac{\alpha}{2})} \left[-\frac{h(\theta)}{(1+2r(t,\theta))^{\frac{1}{2}}} \int_{0}^{2\pi} \frac{\partial_{\eta} \left[(1+2r(\eta))^{\frac{1}{2}} \sin(\eta-\theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta + \int_{0}^{2\pi} \frac{h(\eta) - (1+2r(\theta))^{\frac{1}{2}} \partial_{\eta} \left[h(\eta)(1+2r(\eta))^{-\frac{1}{2}} \sin(\eta-\theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta - \frac{\alpha}{2} \int_{0}^{2\pi} \frac{C_{r}(\eta,\theta)}{A_{r}(\eta,\theta)^{\frac{\alpha}{2}+1}} \left(h(\eta)B_{r}(\eta,\theta) + h(\theta)B_{r}(\theta,\eta) \right) d\eta \right]$$
(3.3)

with

$$B_{r}(\eta,\theta) \triangleq 2 - 2\left(1 + 2r(t,\eta)\right)^{-\frac{1}{2}} \left(1 + 2r(t,\theta)\right)^{\frac{1}{2}} \cos(\eta - \theta)\right), \tag{3.4}$$

$$C_{r}(\eta,\theta) \triangleq 1 + 2r(\eta) - (1 + 2r(\theta))^{\frac{1}{2}} \partial_{\eta} \big[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \big].$$
(3.5)

From (2.11) and (3.4) one may observe that

$$(1+2r(\eta))B_r(\eta,\theta) + (1+2r(\theta))B_r(\theta,\eta) = 2A_r(\theta,\eta).$$
(3.6)

Plugging the last identity into the last term of (3.3) yields

$$T_{3}(\theta) \triangleq \frac{\alpha}{2} \int_{0}^{2\pi} \frac{C_{r}(\eta,\theta)}{A_{r}(\eta,\theta)^{\frac{\alpha}{2}+1}} \left(h(\eta)B_{r}(\eta,\theta) + h(\theta)B_{r}(\theta,\eta)\right)d\eta$$
$$= \frac{\alpha}{2} \int_{0}^{2\pi} \frac{C_{r}(\eta,\theta)B_{r}(\eta,\theta)}{A_{r}(\eta,\theta)^{\frac{\alpha}{2}+1}} \left(h(\eta) - h(\theta)\frac{1+2r(t,\eta)}{1+2r(t,\theta)}\right)d\eta + \frac{\alpha h(\theta)}{1+2r(t,\theta)} \int_{0}^{2\pi} \frac{C_{r}(\eta,\theta)}{A_{r}(\eta,\theta)^{\frac{\alpha}{2}}}d\eta.$$

In view of (2.11), (3.4), (3.5) and (3.6) one may easily check that

$$C_{r}(\eta,\theta)B_{r}(\eta,\theta) = 2A_{r}(\theta,\eta) - (1+2r(\eta))^{-\frac{1}{2}}(1+2r(\theta))^{\frac{1}{2}}\sin(\eta-\theta)\partial_{\eta}A_{r}(\theta,\eta).$$
(3.7)

Then, by (3.5) and (3.7) we get

$$T_{3}(\theta) = \alpha \int_{0}^{2\pi} \frac{h(\eta)}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta - \frac{\alpha h(\theta)}{\left(1 + 2r(t,\theta)\right)^{\frac{1}{2}}} \int_{0}^{2\pi} \frac{\partial_{\eta} \left[(1 + 2r(\eta))^{\frac{1}{2}} \sin(\eta - \theta) \right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta - \frac{\alpha}{2} \int_{0}^{2\pi} \frac{\left(1 + 2r(\eta)\right)^{-\frac{1}{2}} (1 + 2r(\theta))^{\frac{1}{2}} \sin(\eta - \theta) \partial_{\eta} A_{r}(\theta,\eta)}{A_{r}(\eta,\theta)^{\frac{\alpha}{2} + 1}} \left(h(\eta) - h(\theta) \frac{1 + 2r(t,\eta)}{1 + 2r(t,\theta)} \right) d\eta.$$

Integrating the last term by parts implies

$$T_{3}(\theta) = \alpha \int_{0}^{2\pi} \frac{h(\eta)}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta + \frac{(1-\alpha)h(\theta)}{\left(1+2r(t,\theta)\right)^{\frac{1}{2}}} \int_{0}^{2\pi} \frac{\partial_{\eta} \left[(1+2r(\eta))^{\frac{1}{2}}\sin(\eta-\theta)\right]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta$$

$$-\int_{0}^{2\pi} \frac{(1+2r(\theta))^{\frac{1}{2}} \partial_{\eta} [(h(\eta)(1+2r(\eta))^{-\frac{1}{2}} \sin(\eta-\theta)]}{A_{r}^{\alpha/2}(\theta,\eta)} d\eta.$$

Inserting the last formula into (3.3) gives

$$d\nabla E(r)[h](\theta) = \frac{C(\alpha)}{2\pi} \int_0^{2\pi} \frac{h(\eta)}{A_r^{\alpha/2}(\theta,\eta)} d\eta - \frac{C(\alpha)h(\theta)}{2\pi \left(1 + 2r(t,\theta)\right)^{\frac{1}{2}}} \int_0^{2\pi} \frac{\partial_\eta \left[(1 + 2r(\eta))^{\frac{1}{2}}\sin(\eta - \theta)\right]}{A_r^{\alpha/2}(\theta,\eta)} d\eta.$$

Combining this expression with Proposition 2.1 gives the desired result.

Combining this expression with Proposition 2.1 gives the desired result.

Next, we shall move to the special case r = 0 corresponding to the equilibrium state. We shall particularly show that the linear operator acts as a Fourier multiplier allowing to get explicit form for the spectrum. More precisely, we establish the following result.

Proposition 3.2. The linearized equation of (2.12) at the equilibrium state r = 0 writes

$$\partial_t h = \partial_\theta \,\mathcal{L}(\alpha) \,h = \partial_\theta \nabla H_{\mathcal{L}}(h), \tag{3.8}$$

where $H_{\rm L}(h)$ is the quadratic Hamiltonian

$$H_{\rm L}(h) \triangleq \frac{1}{2} \left({\rm L}(\alpha)h, h \right)_{L^2}, \qquad (3.9)$$

the self-adjoint operator $L(\alpha)$ is given by

$$\mathcal{L}(\alpha) \triangleq -(\Omega + V_{0,\alpha}) + \mathbb{K}_{0,\alpha}, \qquad (3.10)$$

the constant $V_{0,\alpha}$ is given by

$$V_{0,\alpha} \triangleq \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{2\sqrt{\pi}\Gamma\left(1 - \frac{\alpha}{2}\right)} \frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma\left(2 - \frac{\alpha}{2}\right)},\tag{3.11}$$

and the integral operator $\mathbb{K}_{0,\alpha}$ is diagonal: for any $h(t,\theta) = \sum_{j\in\mathbb{Z}} h_j(t)e^{ij\theta}$,

$$\mathbb{K}_{0,\alpha} h(t,\theta) = \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{2\sqrt{\pi} \Gamma(1 - \frac{\alpha}{2})} \sum_{j \in \mathbb{Z}} \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})} h_j(t) e^{\mathbf{i}j\theta} \cdot$$
(3.12)

Moreover, the reversible solutions to the linear equation (3.8) are given by

$$h(t,\theta) = \sum_{j \in \mathbb{N}} h_j \cos\left(j\theta - \Omega_j(\alpha)t\right), \quad \text{with} \quad h_j \in \mathbb{R},$$
(3.13)

with

$$\Omega_j(\alpha) \triangleq j\Omega + \frac{j\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{2\sqrt{\pi}\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})}\right).$$
(3.14)

Proof. According to Proposition 3.1, the linearized equation at r = 0 is given by

$$\partial_t h = \partial_\theta \Big[- \big(\Omega + V_{0,\alpha}\big)h + \mathbb{K}_{0,\alpha}(\alpha)h \Big], \qquad (3.15)$$

where

$$V_{0,\alpha} = \frac{C_{\alpha}}{2^{1+\alpha}\pi} \int_{0}^{2\pi} \frac{\cos(\eta-\theta)}{|\sin\left(\frac{\eta-\theta}{2}\right)|^{\alpha}} d\eta \quad \text{and} \quad \mathbb{K}_{0,\alpha}h(t,\theta) = \frac{C_{\alpha}}{2^{1+\alpha}\pi} \int_{0}^{2\pi} \frac{h(t,\eta)}{|\sin\left(\frac{\eta-\theta}{2}\right)|^{\alpha}} d\eta. \tag{3.16}$$

We shall now recall the following identity, see for instance [81, p.449].

$$\int_{0}^{\pi} \sin^{x}(\eta) e^{iy\eta} d\eta = \frac{\pi e^{iy\frac{\pi}{2}} \Gamma(x+1)}{2^{x} \Gamma(1+\frac{x+y}{2}) \Gamma(1+\frac{x-y}{2})} \quad \forall x > -1.$$
(3.17)

Apply this formula with $x = -\alpha$ and y = 2 yields,

$$V_{0,\alpha} = -\frac{C(\alpha)\Gamma(1-\alpha)}{\Gamma\left(2-\frac{\alpha}{2}\right)\Gamma(-\alpha/2)}$$

Recalling $C_{\alpha} = \frac{\Gamma(\alpha/2)}{2^{1-\alpha}\Gamma(\frac{2-\alpha}{2})}$ and using the identity $z\Gamma(z) = \Gamma(z+1)$ we obtain, for any $\alpha \in (0,1)$,

$$V_{0,\alpha} = \frac{\Gamma(\frac{\alpha}{2})\Gamma(1-\alpha)}{2^{1-\alpha}\Gamma(1-\frac{\alpha}{2})\Gamma\left(2-\frac{\alpha}{2}\right)\Gamma(-\frac{\alpha}{2})}$$
$$= \frac{\Gamma(1+\frac{\alpha}{2})\Gamma(1-\alpha)}{2^{1-\alpha}\Gamma^{2}(1-\frac{\alpha}{2})\Gamma\left(2-\frac{\alpha}{2}\right)}.$$

Therefore from Legendre duplication formula (see (A.2)),

$$2^{\alpha}\sqrt{\pi}\,\Gamma(1-\alpha) = \Gamma\left(\frac{1}{2} - \frac{\alpha}{2}\right)\Gamma\left(1 - \frac{\alpha}{2}\right),\tag{3.18}$$

we obtain the expression of $V_{0,\alpha}$ in (3.11). Next, we shall study the action of the integral operator $\mathbb{K}_{0,\alpha}$ on the scalar function h that we expand in Fourier series as

$$h(t,\theta) = \sum_{j \in \mathbb{Z} \setminus \{0\}} h_j(t) \mathbf{e}_j(\theta) \quad \text{with} \quad \mathbf{e}_j(\theta) = e^{\mathbf{i}j\theta}.$$
(3.19)

By linearity, it suffices to evaluate the operator on the exponential basis. From (3.16) and (3.19) we write in view of change of variables

$$\mathbb{K}_{0,\alpha} \mathbf{e}_j(\theta) = \frac{C(\alpha)}{2^{1+\alpha} \pi} \mathbf{e}_j(\theta) \int_0^{2\pi} \frac{e^{ij\eta}}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta$$

Then, applying the formula (3.17) with $x = -\alpha$ and y = 2j we get

$$\mathbb{K}_{0,\alpha} \mathbf{e}_{j}(\theta) = \frac{(-1)^{j} C(\alpha) \Gamma(1-\alpha)}{\Gamma(1+j-\frac{\alpha}{2}) \Gamma(1-j-\frac{\alpha}{2})} \mathbf{e}_{j}(\theta)$$
$$= \frac{\Gamma(\frac{\alpha}{2})}{2^{1-\alpha} \Gamma(1-\frac{\alpha}{2})} \frac{(-1)^{j} \Gamma(1-\alpha)}{\Gamma(1+j-\frac{\alpha}{2}) \Gamma(1-j-\frac{\alpha}{2})} \mathbf{e}_{j}(\theta)$$

Using the identity, see (A.4),

$$\frac{\Gamma(x+j)}{\Gamma(x)} = (-1)^j \frac{\Gamma(1-x)}{\Gamma(1-x-j)},$$
(3.20)

we obtain in view of (3.18)

$$\mathbb{K}_{0,\alpha} \mathbf{e}_j(\theta) = \frac{\Gamma(1-\alpha)}{2^{1-\alpha}\Gamma^2(1-\frac{\alpha}{2})} \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})} \mathbf{e}_j(\theta)$$
$$= \frac{\Gamma(\frac{1}{2}-\frac{\alpha}{2})}{2\sqrt{\pi}\Gamma(1-\frac{\alpha}{2})} \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})} \mathbf{e}_j(\theta).$$

Finally, we obtain

$$\mathbb{K}_{0,\alpha}h(t,\theta) = \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{2\sqrt{\pi}\,\Gamma(1 - \frac{\alpha}{2})} \sum_{j\in\mathbb{Z}} h_j(t) \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})} \mathbf{e}_j(\theta) \cdot \tag{3.21}$$

Inserting (3.11) and (3.21) into (3.15) gives the Fourier expansion of the solutions in the form

$$h(t,\theta) = \sum_{j \in \mathbb{Z}} h_j(0) e^{-i(\Omega_j(\alpha)t - j\theta)}, \ h_j(0) \in \mathbb{C}.$$
(3.22)

where the frequencies $\Omega_i(\alpha)$ are defined in (3.14). In view of (3.20) one has

$$\frac{(-1)^{j}\Gamma(1-\frac{\alpha}{2})}{\Gamma(1-j-\frac{\alpha}{2})} = \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(\frac{\alpha}{2})} \quad \text{and} \quad \frac{(-1)^{j}\Gamma(\frac{\alpha}{2})}{\Gamma(-j+\frac{\alpha}{2})} = \frac{\Gamma(j+1-\frac{\alpha}{2})}{\Gamma(1-\frac{\alpha}{2})}.$$
(3.23)

It follows that

$$\forall j \in \mathbb{Z}, \quad \frac{\Gamma(-j+\frac{\alpha}{2})}{\Gamma(1-j-\frac{\alpha}{2})} = \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})}.$$
(3.24)

Combining the last identity with the expression of $\Omega_j(\alpha)$, given by (3.14), we conclude that

$$\Omega_{-j}(\alpha) = -\Omega_j(\alpha). \tag{3.25}$$

Therefore, every real-valued reversible solution to (3.15) has the form

$$h(t,\theta) = \sum_{j \in \mathbb{N}} h_j \cos \left(\Omega_j(\alpha) t - j\theta \right), \text{ with } h_j \in \mathbb{R}.$$

This concludes the proof of the lemma.

Remark 3.1. In Fourier expansion $h(t,\theta) = \sum_{j \in \mathbb{Z} \setminus \{0\}} h_j(t) e^{in\theta}$, the quadratic Hamiltonian H_L in

(3.9) writes

$$H_{\mathrm{L}}(h) = -\pi \sum_{j \in \mathbb{Z} \setminus \{0\}} \frac{\Omega_j(\alpha)}{j} |h_j|^2 = -2\pi \sum_{j \in \mathbb{N}} \frac{\Omega_j(\alpha)}{j} |h_j|^2, \qquad (3.26)$$

and the self-adjoint operator $L(\alpha)$ reads

$$\mathcal{L}(\alpha)h(t,\theta) = -\sum_{j\in\mathbb{Z}\setminus\{0\}} \frac{\Omega_j(\alpha)}{j} h_j(t) e^{\mathbf{i}j\theta}.$$
(3.27)

The symplectic form writes

$$\mathcal{W} = \frac{1}{2} \sum_{j \in \mathbb{Z} \setminus \{0\}} \frac{1}{\mathrm{i}j} dr_j \wedge dr_{-j} = \sum_{j \ge 1} \frac{1}{\mathrm{i}j} dr_j \wedge dr_{-j}, \qquad (3.28)$$

the Hamiltonian vector field X_H is

$$[X_H(r)]_j = ij (\partial_{r_{-j}} H)(r), \quad \forall j \neq 0$$

and the Poisson bracket,

$$\{F,G\} = \mathcal{W}(X_F,X_G) = -\sum_{j\in\mathbb{Z}\setminus\{0\}} \mathrm{i}j\big(\partial_{r_j}F\big)(r)\big(\partial_{r_j}G\big)(r).$$

3.2. Structure of the linear frequencies. The main task in this section is to investigate some important structures of the equilibrium frequencies. In the first part we shall be concerned with their monotonicity and explore some asymptotic behavior for large modes. However we shall discuss in the second part the non-degeneracy of these frequencies through the so-called Rüssmann conditions. This is crucial in the measure of the final Cantor set giving rise to quasi-periodic solutions for the linear/nonlinear problem. Actually, this set appears as a perturbation of the Cantor set constructed from the equilibrium eigenvalues and therefore perturbative arguments based on their degeneracy are very useful and will be implemented in Section 10.2.

3.2.1. Monotonicity and asymptotic behavior. In what follows, we intend to establish some basic properties related to the monotonicity and the asymptotic behavior for large modes of the spectrum of the linearized operator at the equilibrium state. Notice that their explicit values are detailed in (3.14). Our result reads as follows.

Lemma 3.1. Let $\alpha \in [0,1)$ and $\Omega > 0$. Then the following holds true.

(i) For any
$$j \in \mathbb{Z}$$
, $\Omega_{-j}(\alpha) = -\Omega_j(\alpha)$.

(ii) The sequence $\left(\frac{\Omega_j(\alpha)}{j}\right)_{j\in\mathbb{N}^*}$ is positive and strictly increasing.

(iii) For any $j \in \mathbb{N}^*$, we have the expansion

$$\begin{split} \Omega_j(\alpha) &= \left(\Omega + V_{0,\alpha}\right) j - W_{0,\alpha} \, j \, \mathbb{W}(j,\alpha) \\ &= \left(\Omega + V_{0,\alpha}\right) j - W_{0,\alpha} \, j^\alpha + O\left(j^{\alpha-2}\right) \, . \end{split}$$

where $V_{0,\alpha}$ is given by (3.11) and

$$W_{0,\alpha} \triangleq \frac{1}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})}, \quad \mathsf{W}(j,\alpha) \triangleq \frac{\Gamma(j + \frac{\alpha}{2})}{\Gamma(j + 1 - \frac{\alpha}{2})}.$$

(iv) For all $j \in \mathbb{Z}$, we have

$$|\Omega_j(\alpha)| \ge \Omega |j|.$$

(v) Given $\overline{\alpha} \in (0,1)$, there exists c > 0 such that

$$\forall \alpha \in [0, \overline{\alpha}], \quad \forall j, j' \in \mathbb{Z}, \quad |\Omega_j(\alpha) \pm \Omega_{j'}(\alpha)| \ge c|j \pm j'|.$$

(vi) Given $\overline{\alpha} \in (0,1)$ and $q_0 \in \mathbb{N}$, there exists C > 0 such that

$$\forall j, j' \in \mathbb{Z}, \quad \max_{q \in [0, q_0]} \sup_{\alpha \in [0, \overline{\alpha}]} \left| \partial_{\alpha}^q \left(\Omega_j(\alpha) - \Omega_{j'}(\alpha) \right) \right| \leqslant C |j - j'|.$$

Proof. (i) It is proved in (3.25).

(ii) Using the identity (A.4) together with (3.14) allows to find the alternative formula

$$\Omega_j(\alpha) = j\Omega + jV_{0,\alpha} \left(1 - \frac{\left(1 + \frac{\alpha}{2}\right)_{j-1}}{\left(2 - \frac{\alpha}{2}\right)_{j-1}} \right), \tag{3.29}$$

where $(x)_j$ denotes Pokhhammer's symbol introduced in (A.3). Now, because $x \mapsto (x)_{j-1}$ is strictly increasing in the set \mathbb{R}_+ , we conclude that for all $\alpha \in [0, 1)$ one has

$$0 < \frac{\left(1 + \frac{\alpha}{2}\right)_{j-1}}{\left(2 - \frac{\alpha}{2}\right)_{j-1}} < 1.$$
(3.30)

Since $x \mapsto \Gamma(x)$ is strictly positive, continuous and has a minimum on \mathbb{R}^*_+ then we can easily show, from (3.11), that there exists $c_1 > 0$ such that

$$\inf_{\alpha \in (0,\overline{\alpha})} V_{0,\alpha} \triangleq c_1 > 0. \tag{3.31}$$

Combining (3.29), (3.30) and (3.31) we obtain

$$\forall j \in \mathbb{N}^*, \quad \Omega_j(\alpha) > j\Omega > 0. \tag{3.32}$$

To prove that $j \mapsto \Omega_j(\alpha)/j$ is strictly increasing it suffices, according to (3.29), to check that the sequence $j \mapsto u_j = \frac{\left(1 + \frac{\alpha}{2}\right)_{j-1}}{\left(2 - \frac{\alpha}{2}\right)_{j-1}}$ is strictly decreasing. This follows from the obvious fact that

$$\forall \alpha \in [0,1), \quad \frac{u_{j+1}}{u_j} = \frac{j + \frac{\alpha}{2}}{j+1 - \frac{\alpha}{2}} < 1.$$

This ends the proof of (ii).

(iii) The asymptotic behavior follows from (3.14) and Lemma A.1.

(iv) It is an immediate consequence of (i) and (3.32).

(v) By the oddness of $j \mapsto \Omega_j(\lambda)$ it is enough to establish the estimate for $j, j' \in \mathbb{N}^*$. We shall first focus on the estimate of the difference, that is,

$$\inf_{\alpha \in [0,\overline{\alpha}]} |\Omega_j(\alpha) - \Omega_{j'}(\alpha)| \ge c|j - j'|, \tag{3.33}$$

for some constant c > 0. The result is obvious when j = j', then we shall restrict the discussion to $j \neq j'$. Without loss of generality we can assume that $j > j' \ge 1$. According to (iii) one may writes

$$\Omega_{j}(\alpha) - \Omega_{j'}(\alpha) = \frac{\Omega_{j}(\alpha)}{j}(j-j') - j' \left(\frac{\Omega_{j'}(\alpha)}{j'} - \frac{\Omega_{j}(\alpha)}{j}\right)$$
$$= \frac{\Omega_{j}(\alpha)}{j}(j-j') - j' W_{0,\alpha} \big(\mathbb{W}(j',\alpha) - \mathbb{W}(j,\alpha) \big) \cdot$$
(3.34)

Then, by (iv) we obtain

$$\Omega_{j}(\alpha) - \Omega_{j'}(\alpha) \ge \Omega(j - j') - j' \sup_{\alpha \in [0,\overline{\alpha}]} \left(W_{0,\alpha} | \mathbb{W}(j',\alpha) - \mathbb{W}(j,\alpha) | \right)$$

Using Taylor formula combined with the estimate in Lemma A.1-(i) we get, for any $\alpha \in [0, \overline{\alpha}]$,

$$\mathbb{W}(j',\alpha) - \mathbb{W}(j,\alpha) \Big| \leqslant C \Big| \int_{j'}^{j} \frac{dx}{x^{2-\alpha}} \Big|$$

$$\leq C|j^{\alpha-1} - (j')^{\alpha-1}|$$

$$\leq C\frac{|j-j'|}{j'j^{1-\alpha}}.$$
(3.35)

It follows that

$$\Omega_j(\alpha) - \Omega_{j'}(\alpha) \ge \left(\Omega - \frac{C}{j^{1-\alpha}}\right)(j-j')$$
.

Therefore there exists j_0 independent of $\alpha \in [0, \overline{\alpha}]$ such that if $j > j_0$ then we get (3.33). It remains to verify this property for $j' < j \leq j_0$. Using the one-to-one property of $j \mapsto \Omega_j(\alpha)$ combined with the continuity of $\alpha \in [0, \overline{\alpha}] \mapsto \Omega_j(\alpha) - \Omega_{j'}(\alpha)$ we get for any $j > j' \geq 1$

$$\inf_{\alpha \in [0,\overline{\alpha}]} \left(\Omega_j(\alpha) - \Omega_{j'}(\alpha) \right) \triangleq c_{jj'} > 0.$$

Consequently

$$\inf_{\substack{j\neq j'\in [\![0,j_0]\!]\\\alpha\in [0,\overline{\alpha}]}} |\Omega_j(\alpha) - \Omega_{j'}(\alpha)| = \inf_{j\neq j'\in [\![0,j_0]\!]} c_{jj'} > 0$$

which implies (3.33).

Let us now move to the estimate of $\Omega_j(\alpha) + \Omega_{j_0}(\alpha)$ for $j, j' \in \mathbb{N}^*$. Since both quantities are positive then using the point (iv) yields

$$\forall \lambda \in [0,\overline{\alpha}], \quad |\Omega_j(\alpha) + \Omega_{j_0}(\alpha)| = \Omega_j(\alpha) + \Omega_{j'}(\alpha) \ge \Omega(j+j') \ge c(j+j').$$

This completes the proof of the desired estimate.

 (\mathbf{vi}) From (iii) we may write

$$\Omega_j(\alpha) - \Omega_{j'}(\alpha) = \left(\Omega + V_{0,\alpha}\right)(j-j') - W_{0,\alpha}\left(j\mathsf{W}(j,\alpha) - j'\mathsf{W}(j',\alpha)\right).$$
(3.36)

Let $q_0 \in \mathbb{N}$ and $q \in [0, q_0]$. Differentiating q times the identity (3.34) in α we find

$$\partial_{\alpha}^{q} (\Omega_{j}(\alpha) - \Omega_{j'}(\alpha)) = \partial_{\alpha}^{q} (\Omega + V_{0,\alpha})(j - j') - (\partial_{\alpha}^{q} W_{0,\alpha}) (j \mathbb{W}(j, \alpha) - j' \mathbb{W}(j', \alpha)) - W_{0,\alpha} \partial_{\alpha}^{q} (j \mathbb{W}(j, \alpha) - j' \mathbb{W}(j', \alpha)).$$
(3.37)

By the mean value theorem combined with the estimate in Lemma A.1-(i) we get,

$$\sup_{\alpha \in [0,\overline{\alpha}]} \left| \partial_{\alpha}^{q} \left(j \mathbb{W}(j,\alpha) - j' \mathbb{W}(j',\alpha) \right) \right| \leq C |j-j'|.$$
(3.38)

Combining (3.36) and (3.38) gives the estimate in (vi).

3.2.2. Non-degeneracy and transversality. Next we shall explore the local structure of a given finite set of eigenvalues associated to the equilibrium state. In particular, we intend to establish their non-degeneracy which is an essential property in measuring some suitable connected sets with Diophantine constraints. We first start with giving the following definition.

Definition 3.1. Let $N \in \mathbb{N}^*$. A function $f \triangleq (f_1, \ldots, f_N) : [\alpha_1, \alpha_2] \to \mathbb{R}^N$ is called non-degenerate if, for any vector $c \triangleq (c_1, \ldots, c_N) \in \mathbb{R}^N \setminus \{0\}$, the scalar function $f \cdot c = f_1c_1 + \cdots + f_dc_N$ is not identically zero on the whole interval $[\alpha_1, \alpha_2]$. This means that the curve of f cannot be contained in an hyperplane.

The first main goal is to prove the following result.

Lemma 3.2. Let $\overline{\alpha} \in (0,1)$, $N \in \mathbb{N}^*$ and $1 \leq j_1 < j_2 < \cdots < j_N$ some fixed integers. Then the functions

$$[0,\overline{\alpha}] \to \mathbb{R}^{N},$$

$$\alpha \mapsto \left(\Omega_{j_{1}}(\alpha), \dots, \Omega_{j_{N}}(\alpha)\right),$$

$$[0,\overline{\alpha}] \to \mathbb{R}^{N+1},$$

$$\alpha \mapsto \left(\Omega + V_{0,\alpha}, \Omega_{j_{1}}(\alpha), \dots, \Omega_{j_{N}}(\alpha)\right)$$

are non-degenerate in the sense of the Definition 3.1, where $\Omega_j(\alpha)$ is given by (3.14) and $V_{0,\alpha}$ is defined in (3.11).

Proof. According to Definition 3.1 one has to prove that, for all $c \in \mathbb{R}^N \setminus \{0\}$, the function

$$\alpha \mapsto g_1(\alpha) \triangleq \sum_{k=1}^N c_k j_k \left[\frac{1}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j_k + \frac{\alpha}{2})}{\Gamma(1 + j_k - \frac{\alpha}{2})} \right) + \Omega \right]$$
(3.39)

is not identically zero on the interval $[0, \overline{\alpha}]$. Suppose, by contradiction, that there exists $c \triangleq (c_1, \ldots, c_N) \in \mathbb{R}^N \setminus \{0\}$ such that

$$\sum_{k=1}^{N} c_k j_k \left[\frac{1}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j_k + \frac{\alpha}{2})}{\Gamma(1 + j_k - \frac{\alpha}{2})} \right) + \Omega \right] = 0, \quad \forall \alpha \in [0, \overline{\alpha}].$$
(3.40)

Since the gamma function has no real zeros but simple poles located at $-\mathbb{N}$ then, the function $\alpha \mapsto \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})}$ can be extended to holomorphic function on $\mathbb{C}\backslash \mathfrak{Q}$ with

$$\mathfrak{Q} \triangleq \left\{ \alpha = 2n+1; \ n \in \mathbb{N} \right\}$$

and for all $j \in \mathbb{N}$, the function $\alpha \mapsto \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(1+j-\frac{\alpha}{2})}$ can be extended to holomorphic function on $\mathbb{C} \setminus \mathscr{P}_j$ with

$$\mathcal{P}_j \triangleq \left\{ \alpha = -2(n+j); \ n \in \mathbb{N} \right\}.$$

Moreover,

 $\mathcal{P}_{j+1} \subset \mathcal{P}_j \subset \dots \subset \mathcal{P}_1, \quad \text{and} \quad \mathcal{P}_j \backslash \mathcal{P}_{j+1} = \{-2j\}. \tag{3.41}$

Therefore, the function $g_1(\alpha)$, defined in (3.39), can be extended to holomorphic function on $\mathbb{C}\backslash \mathscr{P}$, where

$$\mathcal{P} \triangleq \mathcal{P}_1 \cup \mathcal{Q} = \left\{ \alpha = -2(n+1); \ n \in \mathbb{N} \right\} \cup \left\{ \alpha = 2n+1; \ n \in \mathbb{N} \right\}.$$
(3.42)

Thus, according to (3.40), there exists $c \triangleq (c_1, \ldots, c_N) \in \mathbb{R}^N \setminus \{0\}$ such that

$$\sum_{k=1}^{N} c_k j_k \left[\frac{1}{2\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j_k + \frac{\alpha}{2})}{\Gamma(1 + j_k - \frac{\alpha}{2})} \right) + \Omega \right] = 0, \quad \forall \alpha \in \mathbb{C} \setminus \mathscr{P}.$$
(3.43)

Since $\Omega > 0$ and the function $\alpha \mapsto 1/\Gamma(1-\frac{\alpha}{2})$ vanishes at 2 then substituting $\alpha = 2$ in (3.43) gives,

$$\sum_{k=1}^{N} c_k j_k = 0$$

It follows that

$$g_1(\alpha) \triangleq \sum_{k=1}^N c_k j_k \left[\left(\frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(2-\frac{\alpha}{2})} - \frac{\Gamma(j_k+\frac{\alpha}{2})}{\Gamma(1+j_k-\frac{\alpha}{2})} \right) \right] = 0, \quad \forall \alpha \in \mathbb{C} \backslash \mathcal{P}.$$
(3.44)

For given $j \in \mathbb{N}$, the function $\alpha \mapsto 1/\Gamma(1+j-\frac{\alpha}{2})$ has zero set

$$\mathscr{Z}_j \triangleq \left\{ \alpha = 2(n+j+1); \ n \in \mathbb{N} \right\}$$

Moreover, one has

$$\mathscr{Z}_j \subset \mathscr{Z}_{j-1}, \text{ and } \mathscr{Z}_{j-1} \setminus \mathscr{Z}_j = \{2j\}.$$
 (3.45)

Thus, substituting $\alpha = 2j_N$ in (3.44) we get

$$g_1(2j_N) = -c_N j_N \Gamma(2j_N) = 0$$

It follows that

$$c_N = 0.$$

Reproducing the same arguments for $c_{N-1}, c_{N-2}, \ldots, c_1$ respectively we get

$$c_{N-1} = c_{N-2} = \dots = c_1 = 0.$$

This gives the contradiction. In order to conclude the proof of the lemma we have to prove that, for any N, for any $1 \leq n_1 < n_2 < \cdots < n_N$ the function $\alpha \in [0, \overline{\alpha}] \mapsto (\Omega + V_{0,\alpha}, \Omega_{j_1}(\alpha), \dots, \Omega_{j_N}(\alpha)) \in \mathbb{R}^{N+1}$ is non-degenerate according to Definition 3.1. Suppose, by contradiction, that there exists $c = (c_0, c_1, \dots, c_N) \in \mathbb{R}^{N+1} \setminus \{0\}$ such that for all $\alpha \in \mathbb{C} \setminus \mathcal{P}$,

$$c_0 \left(\Omega + \frac{2^{-1}}{\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})} \frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} \right) + \sum_{k=1}^N c_k j_k \left[\frac{2^{-1}}{\sqrt{\pi}} \frac{\Gamma(\frac{1}{2} - \frac{\alpha}{2})}{\Gamma(1 - \frac{\alpha}{2})} \left(\frac{\Gamma(1 + \frac{\alpha}{2})}{\Gamma(2 - \frac{\alpha}{2})} - \frac{\Gamma(j_k + \frac{\alpha}{2})}{\Gamma(1 + j_k - \frac{\alpha}{2})} \right) + \Omega \right] = 0.$$

Substituting $\alpha = 2$ in the last equation gives

$$c_0 + \sum_{k=1}^{N} c_k n_k = 0. aga{3.46}$$

Therefore for all $\alpha \in \mathbb{C} \setminus \mathcal{P}$, where \mathcal{P} is given by (3.42), one has

$$g_2(\alpha) \triangleq c_0 \frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(2-\frac{\alpha}{2})} + \sum_{k=1}^N c_k j_k \left(\frac{\Gamma(1+\frac{\alpha}{2})}{\Gamma(2-\frac{\alpha}{2})} - \frac{\Gamma(j_k+\frac{\alpha}{2})}{\Gamma(1+j_k-\frac{\alpha}{2})}\right) = 0.$$
(3.47)

It follows that

$$g_2(2j_N) = -c_N j_N \Gamma(2j_N).$$

Then by (3.44) we conclude that

$$c_N = 0. (3.48)$$

Arguing as above we deduce we get

$$c_{N-1} = c_{N-2} = \dots = c_1 = 0. \tag{3.49}$$

Finally, inserting (3.48) and (3.49) into (3.46) we find

$$c_0 = 0.$$
 (3.50)

Then the vector c is vanishing and this contradicts the assumption.

Proposition 3.3. Let $\overline{\alpha} \in (0,1)$. There exist $q_0 \in \mathbb{N}$, $\rho_0 > 0$ such that, for all $\alpha \in [0,\overline{\alpha}]$, the following assertions hold true.

(i) For any $l \in \mathbb{Z}^d \setminus \{0\}$ we have

$$\max_{k \leq q_0} |\partial_{\alpha}^k \{ \omega_{\mathrm{Eq}}(\alpha) \cdot l \} | \ge \rho_0 \langle l \rangle.$$

(ii) For any $(j,l) \in (\mathbb{N} \times \mathbb{Z}^d) \setminus \{(0,0)\}$, we have

$$\max_{k \leq q_0} |\partial_{\alpha}^k \{ \omega_{\mathrm{Eq}}(\alpha) \cdot l + j V_{0,\alpha} \} | \ge \rho_0 \langle l \rangle.$$

(iii) For any $(j,l) \in (\mathbb{N} \setminus \mathbb{S}_0) \times \mathbb{Z}^d$, we have

$$\max_{k \leq q_0} |\partial_{\alpha}^k \{ \omega_{\mathrm{Eq}}(\alpha) \cdot l + \Omega_j(\alpha) \} | \ge \rho_0 \langle l \rangle.$$

(iv) For any $l \in \mathbb{Z}$, $j, j' \in \mathbb{N} \setminus \mathbb{S}_0$, with $(j, l) \neq (j', 0)$, we have $\max_{k \leq q_0} |\partial_{\alpha}^k \{ \omega_{\mathrm{Eq}}(\alpha) \cdot l + \Omega_j(\alpha) - \Omega_{j'}(\alpha) \} | \geq \rho_0 \langle l \rangle$

where $\omega_{\text{Eq}}(\alpha)$ and $\Omega_n(\alpha)$ are defined in (5.7) and (3.14).

Proof. (i) Suppose, by contradiction, that for all $q_0 \in \mathbb{N}$, $\rho_0 > 0$ there exist $\alpha \in [0, \overline{\alpha}]$ and $l \in \mathbb{Z}^d \setminus \{0\}$ such that

 $\max_{k \leq q_0} |\partial_{\alpha}^k \{ \omega_{\mathrm{Eq}}(\alpha) \cdot l \} | < \rho_0 \langle l \rangle.$

In particular, for all $n \in \mathbb{N}$, if $\rho_0 = \frac{1}{n+1}$ and $q_0 = n$ then there exist $\alpha_n \in [0, \overline{\alpha}]$ and $l_n \in \mathbb{Z}^d \setminus \{0\}$ such that

$$\max_{k \leq n} \left| \partial_{\alpha}^{k} \left\{ \omega_{\mathrm{Eq}}(\alpha_{n}) \cdot \frac{l_{n}}{\langle l_{n} \rangle} \right\} \right| < \frac{1}{n+1}.$$
(3.51)

The sequences $(\alpha_n)_n \subset [0,\overline{\alpha}]$ and $(c_n)_n \triangleq \left(\frac{l_n}{\langle l_n \rangle}\right)_n \subset \mathbb{R}^d \setminus \{0\}$ are bounded. By compactness and up to an extraction we may assume that

$$\lim_{n \to \infty} \frac{l_n}{\langle l_n \rangle} = \widetilde{c} \neq 0 \quad \text{and} \quad \lim_{n \to \infty} \alpha_n = \widetilde{\alpha}.$$

Passing to the limit in (3.51) for $n \to \infty$ we deduce that

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \{ \omega_{\mathrm{Eq}}(\widetilde{\alpha}) \cdot \widetilde{c} \} = 0 \quad \text{with} \quad \widetilde{c} \neq 0.$$

We conclude that the real analytic function $\alpha \to \omega_{\text{Eq}}(\alpha) \cdot \tilde{c}$ is identically zero. This is in contradiction with Lemma 3.2.

(ii) We shall first check the result for the case $l = 0, j \in \mathbb{N}^*$. Obviously one has from (3.31),

$$\inf_{\alpha \in [0,\overline{\alpha}]} \max_{k \leqslant q_0} \left| \partial_{\alpha}^k (j V_{0,\alpha}(\alpha)) \right| \ge \inf_{\alpha \in [0,\overline{\alpha}]} \left| V_{0,\alpha}(\alpha) \right| \ge \rho_0 \langle l \rangle,$$

for some $\rho_0 > 0$. Next, we shall consider the case $j \in \mathbb{N}$, $l \in \mathbb{Z}^d \setminus \{0\}$. By the triangle inequality combined with the boundedness of ω_{Eq} and the bound (3.31) we find

$$|\omega_{\mathrm{Eq}}(\alpha) \cdot l + jV_{0,\alpha}| \ge |j||V_{0,\alpha}| - |\omega_{\mathrm{Eq}}(\alpha) \cdot l| \ge c|j| - C|l| \ge |l|$$

provided that $|j| \ge C_0 |l|$ for some $C_0 > 0$. Thus, we shall restrict the proof to indices j and l with

$$|j| \le C_0 |l|, \quad j \in \mathbb{N}, \quad l \in \mathbb{Z}^d \setminus \{0\}.$$

$$(3.52)$$

Arguing by contradiction as in the previous case, we may assume the existence of sequences $l_n \in \mathbb{Z}^d \setminus \{0\}, j_n \in \mathbb{N}$ satisfying (3.52) and $\alpha_n \in [0, \overline{\alpha}]$ such that

$$\max_{k \leq n} \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha_{n}) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{j_{n} V_{0,\alpha_{n}}}{\langle l_{n} \rangle} \right) \right| < \frac{1}{1+n}$$

and therefore

$$\forall k \in \mathbb{N}, \quad \forall n \ge k, \quad \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha_{n}) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{j_{n} V_{0,\alpha_{n}}}{\langle l_{n} \rangle} \right) \right| < \frac{1}{1+n}.$$
(3.53)

Since sequences $\{\alpha_n\}, \{b_n\} \triangleq \{\frac{j_n}{\langle l_n \rangle}\}$ and $\{c_n\} \triangleq \{\frac{l_n}{\langle l_n \rangle}\}$ are bounded, then up to an extraction we can assume

$$\lim_{n \to \infty} \alpha_n = \widetilde{\alpha}, \quad \lim_{n \to \infty} b_n = \widetilde{b} \neq 0 \quad \text{and} \quad \lim_{n \to \infty} c_n = \widetilde{c} \neq 0.$$

Hence, letting $n \to +\infty$ in (3.53) and using $\alpha \mapsto V_{0,\alpha}$ is smooth we obtain

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \widetilde{c} + \widetilde{b} V_{0,\alpha} \right)_{|\alpha = \widetilde{\alpha}} = 0.$$

Thus, the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{c} + \tilde{b} V_{0,\alpha}$ with $(\tilde{c}, \tilde{b}) \neq (0, 0)$ is identically zero and this contradicts Lemma 3.2.

(iii) Consider $(l, j) \in \mathbb{Z}^d \times (\mathbb{N} \setminus \mathbb{S}_0)$. Then applying the triangle inequality and Lemma 3.1-(iv), we get

$$|\omega_{\mathrm{Eq}}(\alpha) \cdot l + \Omega_j(\alpha)| \ge |\Omega_j(\alpha)| - |\omega_{\mathrm{Eq}}(\alpha) \cdot l| \ge \Omega_j - C|l| \ge \langle l \rangle$$

provided that $|j| \ge C_0 |l|$ for some $C_0 > 0$. So we shall restrict the proof to integers j with

$$0 \leq j < C_0 \langle l \rangle, \quad j \in \mathbb{N} \setminus \mathbb{S}_0 \quad \text{and} \quad l \in \mathbb{Z}^d \setminus \{0\}.$$
 (3.54)

Arguing by contradiction, one can check that for all $n \in \mathbb{N}$, there exists $l_n \in \mathbb{Z}^d \setminus \{0\}, j_n \notin \mathbb{S}_0$ and $\alpha_n \in [0, \overline{\alpha}]$ such that

$$\max_{k \leq n} \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{\Omega_{j_{n}}(\alpha)}{\langle l_{n} \rangle} \right)_{|\alpha = \alpha_{n}} \right| < \frac{1}{1+n}$$

and therefore

$$\forall k \in \mathbb{N}, \quad \forall n \ge k, \quad \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{\Omega_{j_{n}}(\alpha)}{\langle l_{n} \rangle} \right)_{|\alpha = \alpha_{n}} \right| < \frac{1}{1+n}.$$
(3.55)

Since sequences $\{\alpha_n\}$ and $\{c_n\} \triangleq \left\{\frac{l_n}{\langle l_n \rangle}\right\}$ are bounded, then up to an extraction we can assume

$$\lim_{n \to \infty} \alpha_n = \widetilde{\alpha} \quad \text{and} \quad \lim_{n \to \infty} c_n = \widetilde{c} \neq 0.$$

Now we shall distinguish two cases.

• Case 1: $(l_n)_n$ is bounded. Then from (3.54) and up to an extraction the sequences (l_n) and (j_n) are stationary. So we can assume that for any $n \in \mathbb{N}$, we have $l_n = \tilde{l}$ and $j_n = \tilde{j}$, with $\tilde{l} \in \mathbb{Z}^d \setminus \{0\}$ and $\tilde{j} \in \mathbb{N}$. Hence, taking the limit as $n \to +\infty$ in (3.55) yields

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \tilde{l} + \Omega_{\tilde{j}}(\alpha) \right)_{|_{\alpha = \tilde{\alpha}}} = 0.$$

Thus, the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{l} + \Omega_{\tilde{j}}(\alpha)$ with $(\tilde{l}, 1) \neq (0, 0)$ is identically zero which contradicts Lemma 3.2.

• Case 2: (l_n) is unbounded. Up to a subsequence, we can assume that $\lim_{n\to\infty} |l_n| = \infty$ and $\lim_{n\to\infty} \frac{l_n}{\langle l_n \rangle} = \tilde{c} \in \mathbb{R}^d \setminus \{0\}$. We shall distinguish two subclasses depending on whether the sequence $(j_n)_n$ is bounded or not. When it is bounded, then up to an extraction we may assume that this sequence of integers is stationary. Then, taking the limit $n \to +\infty$ in (3.55), we find

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \omega_{\mathrm{Eq}}(\alpha)_{|_{\alpha = \widetilde{\alpha}}} \cdot \widetilde{c} = 0.$$

As before we conclude that the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{c}$, with $\tilde{c} \neq 0$, is identically zero, which is a contradiction with the Lemma 3.2.

It remains to explore the case where $(j_n)_n$ is unbounded. Then up to an extraction we can assume that $\lim_{n\to\infty} j_n = \infty$. According to Lemma 3.1-(iii) we find

$$\frac{\Omega_{j_n}(\alpha)}{\langle l_n \rangle} = \frac{j_n}{\langle l_n \rangle} \left(\Omega + V_{0,\alpha} - W_{0,\alpha} \mathsf{W}(j_n, \alpha) \right).$$
(3.56)

Moreover, by (3.54), the sequence $\left(\frac{j_n}{\langle l_n \rangle}\right)_n$ is bounded, so up to a subsequence, we can assume that it converges to \tilde{d} . Therefore, differentiating then taking the limit in (3.56) we obtain

$$\lim_{n \to +\infty} \frac{\partial_{\alpha}^{q} \Omega_{j_{n}}(\alpha_{n})}{\langle l_{n} \rangle} = \partial_{\alpha}^{q} \left(\widetilde{d} \left(\Omega + V_{0,\alpha} \right) \right)_{|_{\alpha = \widetilde{\alpha}}},$$

where we have used in the last identity the estimate in Lemma A.1-(i). Hence, taking the limit $j \to +\infty$ in (3.55) we get :

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \widetilde{c} - \widetilde{d} (\Omega + V_{0,\alpha}) \right)_{|_{\alpha = \widetilde{\alpha}}} = 0.$$

Thus, the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{c} - \tilde{d}(\Omega + V_{0,\alpha})$ is identically zero. This is a contradiction with the Lemma 3.2 because $\tilde{c} \neq 0$.

(iv) Consider $l \in \mathbb{Z}^d, j, j' \in \mathbb{N} \setminus \mathbb{S}_0$ with $(l, j) \neq (0, j')$. Then applying the triangle inequality combined with Lemma 3.1-(v), we infer that

$$|\omega_{\mathrm{Eq}}(\alpha) \cdot l + \Omega_j(\alpha) \pm \Omega_{j'}(\alpha)| \ge |\Omega_j(\alpha) \pm \Omega_{j'}(\alpha)| - |\omega_{\mathrm{Eq}}(\alpha) \cdot l| \ge c|j \pm j'| - C|l| \ge \langle l \rangle$$

provided $|j - j'| \ge C_0 |l|$ for some $C_0 > 0$. In this case the desired estimate is trivial. So we shall restrict the proof to integers such that

$$|j \pm j'| < c_0 \langle l \rangle, \quad l \in \mathbb{Z}^d \setminus \{0\}, \quad j, j' \in \mathbb{N} \setminus \mathbb{S}_0.$$
 (3.57)

We will argue by contradiction as in the previous cases. We assume that for all $n \in \mathbb{N}$, there exists $(l_n, j_n) \neq (0, j'_n) \in \mathbb{Z}^{d+1}$ satisfying (3.57) and $\alpha_n \in [0, \overline{\alpha}]$ such that

$$\max_{k \leq n} \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{\Omega_{j_{n}}(\alpha) \pm \Omega_{j_{n}'}(\alpha)}{\langle l_{n} \rangle} \right)_{|\alpha = \alpha_{n}} \right| < \frac{1}{1+n}$$

and therefore

$$\forall k \in \mathbb{N}, \quad \forall n \ge k, \quad \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \frac{l_{n}}{\langle l_{n} \rangle} + \frac{\Omega_{j_{n}}(\alpha) \pm \Omega_{j_{n}'}(\alpha)}{\langle l_{n} \rangle} \right)_{|\alpha = \alpha_{n}} \right| < \frac{1}{1+n}$$
(3.58)

Since the sequences $\left\{\frac{l_n}{\langle l_n \rangle}\right\}_n$ and $\{\alpha_n\}_n$ are bounded, then by compactness we can assume that , $\lim_{n \to \infty} \frac{l_n}{\langle l_n \rangle} = \tilde{c} \neq 0$ and $\lim_{n \to \infty} \alpha_n = \tilde{\alpha}$. As before we distinguish two cases : • Case 1: $(l_n)_n$ is bounded. We shall only focus on the most delicate case associated to the difference $\Omega_{j_n} - \Omega_{j'_n}$. Up to an extraction we may assume that this sequence of integers is stationary, that is, $l_n = \tilde{l}$. Now looking at (3.57) we have two sub-cases depending on whether or not the sequences $(j_n)_n$ and $(j'_n)_n$ are bounded.

• Sub-case (1): $(j_n)_n$ and $(j'_n)_n$ are bounded. Up to an extraction we can assume that they are stationary, that is, $j_n = \tilde{j}, j'_n = \tilde{j}'$. In addition from the assumption we have also $(\tilde{l}, \tilde{j}) \neq (0, \tilde{j}')$ and $\tilde{j}, \tilde{j}' \notin \mathbb{S}_0$. Hence taking the limit $n \to +\infty$ in (3.58), we get

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \tilde{l} + \Omega_{\tilde{j}}(\alpha) - \Omega_{\tilde{j}'}(\alpha) \right)_{|_{\alpha = \tilde{\alpha}}} = 0.$$

Thus, the real analytic function $\alpha \mapsto \Omega(\alpha) \cdot \tilde{l} + \Omega_{\tilde{j}}(\alpha) - \Omega_{\tilde{j}'}(\alpha)$ is identically zero. If $\tilde{j} = \tilde{j}'$ then this contradicts Lemma 3.2 since $\tilde{l} \neq 0$. However in the case $\tilde{j} \neq \tilde{j}' \in \mathbb{N} \setminus \mathbb{S}_0$ this still contradicts this lemma applied with the vector frequency $(\omega_{\text{Eq}}, \Omega_{\tilde{j}}, \Omega_{\tilde{j}'})$ instead of ω_{Eq} .

• Sub-case $@: (j_n)_n$ and $(j'_n)_n$ are unbounded. Then up to an extraction we can assume that $\lim_{n\to\infty} j_n = \lim_{m\to\infty} j'_n = \infty$. Without loss of generality, we can assume that for a given n we have $j_n \ge j'_n$. Then according to Lemma 3.1-(iii) we get the splitting

$$\frac{\partial_{\alpha}^{k}\left(\Omega_{j_{n}}(\alpha)-\Omega_{j_{n}'}(\alpha)\right)}{\langle l_{n}\rangle} = \partial_{\alpha}^{k}\left(\Omega+V_{0,\alpha}\right)\frac{j_{n}-j_{n}'}{\langle l_{n}\rangle} - \partial_{\alpha}^{k}\left(W_{0,\alpha}\mathbb{W}(j_{n},\alpha)\right)\frac{j_{n}-j_{n}'}{\langle l_{n}\rangle} - \frac{j_{n}'}{\langle l_{n}\rangle}\partial_{\alpha}^{k}\left[W_{0,\alpha}\left(\mathbb{W}(j_{n}',\alpha)-\mathbb{W}(j_{n},\alpha)\right)\right].$$
(3.59)

Using Taylor formula combined with Lemma A.1-(i), in a similar way to (3.35), give for any $\alpha \in [0, \overline{\alpha}]$,

$$j'_{n} \left| \partial_{\alpha}^{k} \mathbb{W}(j'_{n}, \alpha) - \partial_{\alpha}^{k} \mathbb{W}(j_{n}, \alpha) \right| \leq C \left| j'_{n} \int_{j'_{n}}^{j_{n}} \frac{dx}{x^{2-\alpha-\epsilon}} \right| \leq C |j_{n} - j'_{n}| j_{n}^{\overline{\alpha}+\epsilon-1}.$$

$$(3.60)$$

Using once again (3.57), up to an extraction, we assume that $\lim_{n\to\infty} \frac{j'_n - j_n}{\langle l_n \rangle} = \tilde{d}$. Therefore, combining (3.59), (3.60) and Lemma A.1-(i), we conclude that

$$\lim_{n \to \infty} \partial_{\alpha}^{k} \left(\frac{\Omega_{j_{n}}(\alpha) - \Omega_{j_{n}'}(\alpha)}{\langle l_{n} \rangle} \right)_{|_{\alpha = \alpha_{n}}} = \widetilde{d} \, \partial_{\alpha}^{k} \big(\Omega + V_{0,\alpha} \big)_{|_{\alpha = \widetilde{\alpha}}}.$$

So taking the limit $n \to +\infty$ in (3.58), yields

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \widetilde{c} + \widetilde{d} \left(\Omega + V_{0,\alpha} \right) \right)_{|_{\alpha = \widetilde{\alpha}}} = 0.$$

Thus, the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{c} + \tilde{d}(\Omega + V_{0,\alpha})$ with $(\tilde{c}, \tilde{d}) \neq (0, 0)$ is identically zero which is in a contradiction with Lemma 3.2.

• Case 2: $(l_n)_n$ is unbounded. Up to an extraction we can assume that $\lim_{n \to \infty} |l_n| = \infty$.

We shall distinguish three sub-cases.

• Sub-case ①. The sequences $(j_n)_n$ and $(j'_n)_n$ are bounded. Up to an extraction they will converge and then taking the limit in (3.58) yields,

$$\forall k \in \mathbb{N}, \quad \partial^k_\alpha \omega_{\mathrm{Eq}}(\bar{\alpha}) \cdot \tilde{c} = 0$$

which leads to a contradiction as before.

• Sub-case 2. The sequences $(j_n)_n$ and $(j'_n)_n$ are both unbounded. This is similar to the sub-case 2 of the case 1.

• Sub-case 3. The sequence $(j_n)_n$ is unbounded and $(j'_n)_n$ is bounded (the symmetric case is

similar). Without loss of generality we can assume that $\lim_{n \to \infty} j_n = \infty$ and $j'_n = \tilde{j}$. By (3.57) and up to an extraction one gets $\lim_{n\to\infty} \frac{j_n \pm j'_n}{|l_n|} = \tilde{d}$. Using (3.59) combined with (3.60) and Lemma A.1-(i) in order to get for any $k \in \mathbb{N}$,

$$\lim_{n \to \infty} \langle l_n \rangle^{-1} \partial_{\alpha}^k \left(\Omega_{j_n}(\alpha) \pm \Omega_{j'_n}(\alpha) - (j_n \pm j'_n) \left(\Omega + V_{0,\alpha} \right) \right)_{|\alpha = \alpha_n} = \lim_{n \to \infty} \partial_{\alpha}^k \left(\frac{(j_n \pm j'_n)}{\langle l_n \rangle} W_{0,\alpha} \mathbb{W}(j_n, \alpha) \pm \frac{j'_n}{\langle l_n \rangle} W_{0,\alpha} \left(\mathbb{W}(j_n, \alpha) - \mathbb{W}(j'_n, \alpha) \right)_{|\alpha = \alpha_n} = 0$$

Hence, taking the limit in (3.58) implies

$$\forall k \in \mathbb{N}, \quad \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot \widetilde{c} + \widetilde{d} (\Omega + V_{0,\alpha}) \right)_{\alpha = \widetilde{\alpha}} = 0$$

Thus, the real analytic function $\alpha \mapsto \omega_{\text{Eq}}(\alpha) \cdot \tilde{c} + \tilde{d}(\Omega + V_{0,\alpha})$ is identically zero with $(\tilde{c}, \tilde{d}) \neq 0$ which contradicts Lemma 3.2. This completes the proof of the lemma.

3.3. Linear quasi-periodic solutions. Note that all the solutions (3.13) for (3.8) are periodic, quasi-periodic or almost periodic in time, with linear frequencies of oscillations $\Omega_i(\alpha)$, depending on the irrationality properties of the frequencies $\Omega_i(\alpha)$ and how many normal mode amplitudes $h_i(0)$ are not zero. We shall prove that if $h_i(0) = 0$ for any index j except at a finite set S (tangential sites) then the linear solutions (3.13) are quasi-periodic in time.

Lemma 3.3. Let $\Omega > 0, \overline{\alpha} \in (0,1), \mathbb{S} \subset \mathbb{N} \setminus \{0\}$, with $\#\mathbb{S} = d \ge 1$. There exists a Cantor-like set $G_0 \subset (0,\overline{\alpha})$ with full Lebesgue measure such that for any $\alpha \in G_0$, the linearized vortex patch equation (3.8) admits time quasi-periodic solutions with a non-resonant frequency vector $\omega_{Eq}(\alpha) \triangleq$ $(\Omega_i(\alpha))_{i\in\mathbb{S}}\in\mathbb{R}^d$ satisfying (1.4) and taking the form

$$h(t,\theta) = \sum_{j \in \mathbb{S}} h_j \cos\left(j\theta - \Omega_j(\alpha)t\right)$$
(3.61)

where $\Omega_i(\alpha)$ are the equilibrium frequencies defined in (3.14).

Proof. We consider the sets

$$C_{0} \triangleq \left\{ \alpha \in (0, \overline{\alpha}); \ \omega_{\mathrm{Eq}}(\alpha) \cdot l \neq 0, \ \forall l \in \mathbb{Z}^{d} \setminus \{0\} \right\},$$
(3.62)

$$(0,\overline{\alpha})\backslash \mathsf{C}_0 \subset \bigcup_{l \in \mathbb{Z}^d \setminus \{0\}} \mathsf{G}_l \quad \text{with} \quad \mathsf{G}_l \triangleq \left\{ \alpha \in (0,\overline{\alpha}); \ |\omega_{\mathrm{Eq}}(\alpha) \cdot l| \leqslant \frac{\kappa}{\langle l \rangle^{\tau_1}} \right\}.$$
(3.63)

Applying Lemma A.3 and Proposition 3.3, we find

$$|\mathbf{G}_l| \lesssim \kappa^{\frac{1}{q_0}} \langle l \rangle^{-1 - \frac{\tau_1 + 1}{q_0}}.$$

It follows from (3.63) that

$$\begin{split} \left| (0,\overline{\alpha}) \backslash \mathbf{C}_{0} \right| &\leq \sum_{l \in \mathbb{Z}^{d} \backslash \{0\}} \left| \mathbf{G}_{l} \right| \\ &\lesssim \sum_{l \in \mathbb{Z}^{d}} \kappa^{\frac{1}{q_{0}}} \left\langle l \right\rangle^{-1 - \frac{\tau_{1} + 1}{q_{0}}} \end{split}$$

Then assuming $\tau_1 > (d-1)q - 1$ we deduce that

$$\left| (0,\overline{\alpha}) \backslash \mathbf{C}_0 \right| \lesssim \kappa^{\frac{1}{q_0}} \tag{3.64}$$

and thus

$$\forall \kappa \in (0,1), \quad \overline{\alpha} \ge \left| \mathsf{C}_0 \right| \ge \overline{\alpha} - C\kappa^{\frac{1}{q_0}}. \tag{3.65}$$
be zero we conclude the proof of Lemma 3.3.

Taking the limit when $\overline{\gamma}$ goes to zero we conclude the proof of Lemma 3.3.

4. Functional tools

In this section we set the functional framework, recall basic definitions and gather some technical results that will be used along the paper.

Notations. We denote $\mathbb{N} \triangleq \{0, 1, ...\}$ and $\mathbb{N}^* \triangleq \{1, 2, ...\}$. Along this paper we shall make use of the following parameters

$$\kappa \in (0,1), \quad (q,d) \in (\mathbb{N}^*)^2, \quad S \ge s \ge s_0 > \frac{d+1}{2} + q, \tag{4.1}$$

where S is a fixed large number.

4.1. Function spaces. We shall introduce various function spaces that will be used frequently throughout this paper. The first one is the classical Sobolev space $H^s(\mathbb{T}^d \times \mathbb{T}; \mathbb{C})$ which is the set of all complex periodic functions $h: \mathbb{T}^d \times \mathbb{T} \to \mathbb{C}$ such that

$$h = \sum_{(l,j)\in\mathbb{Z}^{d+1}} h_{l,j} \mathbf{e}_{l,j}, \quad h_{l,j}\in\mathbb{C}, \qquad \|h\|_{H^s}^2 = \sum_{(l,j)\in\mathbb{Z}^{d+1}} \langle l,j\rangle^{2s} |h_{l,j}|^2,$$

where

 $\mathbf{e}_{l,j}(\varphi,\theta) = e^{\mathrm{i}(l\cdot\varphi+j\cdot\theta)} \quad \text{and} \quad \langle l,j\rangle \triangleq \max(1,|l|,|j|).$

For some reasons connected with the reversibility of the system we need to distinguish the following sub-spaces,

$$H^{s}_{\text{even}} = \left\{ h \in H^{s}(\mathbb{T}^{d+1}; \mathbb{R}); \ h(-\varphi, -\theta) = h(\varphi, \theta), \ \forall (\varphi, \theta) \in \mathbb{T}^{d+1} \right\}$$
(4.2)

and

$$H^{s}_{\text{odd}} = \left\{ h \in H^{s}(\mathbb{T}^{d+1}; \mathbb{R}); \ h(-\varphi, -\theta) = -h(\varphi, \theta), \ \forall (\varphi, \theta) \in \mathbb{T}^{d+1} \right\}$$

We denote $H^{\infty} \triangleq \bigcap_{s \in \mathbb{R}} H^s$ and similarly we define the subspaces H^{∞}_{even} and H^{∞}_{even} .

To implement Nash-Moser scheme for the nonlinear equation we need to measure the dependence of the solutions with respect to the exterior parameters $\lambda = (\omega, \alpha) \in \mathbb{R}^{d+1}$ and for this aim we need to define the weighted Sobolev spaces.

Definition 4.1. Let $\kappa \in (0,1)$, $s \in \mathbb{R}$, $q \in \mathbb{N}$, $d \in \mathbb{N}^*$. Let \mathcal{O} be an open bounded subset of \mathbb{R}^{d+1} . We define the following spaces

$$W^{q,\infty}_{\gamma}(\mathcal{O}, H^s(\mathbb{T}^{d+1})) = \left\{ h: \mathcal{O} \to H^s; \, \|h\|^{q,\kappa}_s < +\infty \right\}$$

with

$$\|h\|_{s}^{q,\kappa} \triangleq \sup_{\substack{\lambda \in \mathcal{O}, \\ |\beta| \leqslant q}} \kappa^{|\beta|} \|\partial_{\lambda}^{\beta} h(\lambda, \cdot)\|_{H^{s-|\beta|}(\mathbb{T}^{d+1})}$$
(4.3)

and

$$W^{q,\infty}_{\gamma}(\mathcal{O},\mathbb{C}) = \left\{ h: \mathcal{O} \to \mathbb{C}; \, \|h\|^{q,\kappa} < +\infty \right\}$$

with

$$\|h\|^{q,\kappa} \triangleq \sup_{\substack{\lambda \in \mathcal{O}, \\ |\beta| \leqslant q}} \kappa^{|\beta|} \sup_{\lambda \in \mathcal{O}} |\partial_{\lambda}^{\beta} h(\lambda)|.$$

$$(4.4)$$

4.2. Classical operations. The main aim of this section is to recall some classical results related the law products and composition laws. In what follows \mathcal{O} denotes an open bounded subset of \mathbb{R}^{d+1} . The firs result is standard and whose proof can be found in [16] and the references therein.

Lemma 4.1. Let $\kappa \in (0,1), q \in \mathbb{N}$ and $d \in \mathbb{N}^*$. Then the following assertions hold true. (i) Let $s \ge 0, s_0 > \frac{d+1}{2}$ and $f, g \in H^s(\mathbb{T}^{d+1}) \cap H^{s_0}(\mathbb{T}^{d+1})$, then

(ii) Let
$$s \ge q, s_0 > \frac{d+1}{2} + q$$
 and $f, g \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^{d+1}))$, then $fg \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^{d+1}))$ with $\|fg\|_s^{q,\kappa} \lesssim \|f\|_{s_0}^{q,\kappa} \|g\|_s^{q,\kappa} + \|f\|_s^{\gamma,q} \|g\|_{s_0}^{q,\kappa}.$

(iii) Let $f, g \in W^{q,\infty}_{\kappa}(\mathcal{O}, \mathbb{C})$, then $fg \in W^{q,\infty}_{\kappa}(\mathcal{O}, \mathbb{C})$ with $\|fg\|^{q,\kappa} \lesssim \|f\|^{q,\kappa} \|g\|^{q,\kappa}$. (iv) Let $s \ge q$ and $(f,g) \in W^{q,\infty}_{\kappa}(\mathcal{O}, \mathbb{C}) \times W^{q,\infty}_{\kappa}(\mathcal{O}, H^{s}(\mathbb{T}^{d+1}))$, then $fg \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^{s}(\mathbb{T}^{d+1}))$ with

 $\|fg\|_s^{q,\kappa} \lesssim \|f\|^{q,\kappa} \|g\|_s^{q,\kappa}.$

Next we shall introduce a cut-off frequency operator which has the advantage to smooth out functions. For $N \in \mathbb{N}^*$, we define the orthogonal projections on the space $H^s(\mathbb{T}^{d+1})$ through

$$\Pi_N h = \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \\ \langle l,j \rangle \leqslant N}} h_{l,j} \mathbf{e}_{l,n} \quad \text{and} \quad \Pi_N^\perp = \mathrm{Id} - \Pi_N.$$

The next result is elementary and can be easily checked.

Lemma 4.2. Let $N \in \mathbb{N}^*$, $s \in \mathbb{R}$ and $\mu \in \mathbb{R}_+$. Then

 $\|\Pi_N h\|_{H^{s+\mu}} \leqslant N^{\mu} \|h\|_{H^s}, \quad \|\Pi_N^{\perp} h\|_{H^s} \leqslant N^{-\mu} \|h\|_{H^{s+\mu}}.$

and

$$\|\Pi_N h\|_{s+\mu}^{q,\kappa} \leqslant N^{\mu} \|h\|_s^{q,\kappa}, \quad \|\Pi_N^{\perp} h\|_s^{q,\kappa} \leqslant N^{-\mu} \|h\|_{s+\mu}^{q,\kappa}$$

Now we shall recall the classical interpolation inequalities, see for instance [16].

Lemma 4.3. Let $\kappa \in (0,1), q \in \mathbb{N}, d \in \mathbb{N}^*$ and $s_1 \leq s \leq s_2 \in \mathbb{R}$ such that $s = \theta s_1 + (1-\theta)s_2$ with $\theta \in [0,1]$. Then, for any $h \in W_{\kappa}^{q,\infty}(\mathcal{O}, H^{s_2})$,

$$\|h\|_{s}^{q,\kappa} \lesssim \left(\|h\|_{q,s_{1}}^{q,\kappa}\right)^{\theta} \left(\|h\|_{q,s_{2}}^{q,\kappa}\right)^{1-\theta}$$

The proof of the next result dealing with some composition laws can be found for instance in [16].

Lemma 4.4. Let $s \ge s_0 > \frac{d+1}{2} + q$ and $\kappa \in (0,1)$. Let $f : \mathcal{O} \times \mathbb{R} \to \mathbb{R}$ be \mathscr{C}^{∞} and $h \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^{d+1}))$ such that $\|h\|_{s_0}^{q,\kappa} \leq C_0$ for and arbitrary $C_0 > 0$ and define the point-wise composition

$$\forall (\lambda, x) \in \mathcal{O} \times \mathbb{T}^{d+1}, \quad F(h)(\lambda, x) \triangleq F(\lambda, h(\lambda, x)).$$

Then $F(h) \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^{d+1}))$ with

$$\|F(h) - F(\lambda, 0)\|_s^{q,\kappa} \leqslant C(s, F, C_0) \|h\|_s^{q,\kappa}$$

We shall also make use of the following classical result that can proved directly from Taylor formula combined with the chain rule (Faá di Bruno's formula) and interpolation inequalities in $W^{k,\infty}$.

Lemma 4.5. Let $q \in \mathbb{N}^*$ and $F : \mathbb{R} \to \mathbb{R}$ be a C^{∞} function with bounded derivatives and $h : \mathcal{O} \to \mathbb{R}$ be a function in $W^{q,\infty}(\mathcal{O})$. Then the composition $F(h) \in W^{q,\infty}(\mathcal{O})$ with

$$||F(h) - F(0)||_{W^{q,\infty}} \leq C(q,F) ||u||_{W^{q,\infty}} \left(1 + ||h||_{L^{\infty}}^{q-1}\right).$$

The next result will be useful later in studying the asymptotic structure of the linearized operator.

Lemma 4.6. Let $\epsilon > 0, d \in \mathbb{N}^*$ and $q \in \mathbb{N}$, then the following assertions hold true.

(i) Let $s \ge 0$, there exists C > 0 such that if $f \in H^{s+\frac{5}{2}+\epsilon}(\mathbb{T}^2)$ and satisfying in addition the symmetry property $f(\theta, \eta) = f(\eta, \theta)$, then there exists $g : \mathbb{T}^2 \to \mathbb{R}$ such that

$$f(\theta,\eta) = \frac{1}{2} \left(f(\theta,\theta) + f(\eta,\eta) \right) + \sin^2 \left(\frac{\eta - \theta}{2} \right) g(\theta,\eta)$$

with

$$\sup_{\eta \in \mathbb{T}} \|g(\cdot, \cdot + \eta\|_{H^s(\mathbb{T})} \leqslant C \|f\|_{H^{s+\frac{5}{2}+\epsilon}(\mathbb{T}^2)}$$

More generally, we have the estimate: for any $p_1, p_2 \in \mathbb{N}$

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} g)(\cdot, \cdot + \eta) \|_{H^s(\mathbb{T})} \lesssim \| f \|_{H^{s+\epsilon+\frac{5}{2}+p_1+p_2}(\mathbb{T}^2)}$$

(ii) Let $s \ge q$, there exists C > 0 such that if $f \in W^{q,\infty}_{\kappa}(\mathcal{O}; H^{s+\frac{5}{2}+\epsilon}(\mathbb{T}^d \times \mathbb{T}^2))$ being a symmetric function: $f(\lambda, \varphi, \theta, \eta) = f(\lambda, \varphi, \eta, \theta)$ then there exists g such that

$$f(\lambda,\varphi,\theta,\eta) = \frac{1}{2} \left(f(\lambda,\varphi,\theta,\theta) + f(\lambda,\varphi,\eta,\eta) \right) + \sin^2 \left(\frac{\eta-\theta}{2} \right) g(\lambda,\varphi,\theta,\eta)$$

with

$$\sup_{\eta \in \mathbb{T}} \left\| g\left(*, \cdot, \bullet, \bullet + \eta \right) \right\|_{s}^{q, \kappa} \leq C \|\Delta_{\theta, \eta} f\|_{s + \frac{1}{2} + \epsilon}^{q, \kappa}$$
$$\leq C \|f\|_{s + \frac{5}{2} + \epsilon}^{q, \kappa}$$

and where $\Delta_{\theta,\eta} = \partial_{\theta}^2 + \partial_{\eta}^2$. More generally, we have the estimate: for any $p_1, p_2 \in \mathbb{N}$

$$\sup_{\eta \in \mathbb{T}} \left\| \left(\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} g \right) \left(*, \cdot, \bullet, \bullet + \eta \right) \right\|_{s}^{q, \kappa} \leqslant C \|f\|_{s+\epsilon+\frac{5}{2}+p_1+p_2}^{q, \kappa}$$

Here the symbols $*, \cdot, \bullet$ denote λ, φ, θ , respectively.

Proof. (i) Let us split f through its Fourier expansion,

$$f(\theta, \eta) = \sum_{j, j' \in \mathbb{Z}} a_{j,k} e^{i(j\theta + j'\eta)}$$

Since f is symmetric then one finds from some algebraic manipulations,

$$f(\theta,\eta) = \frac{1}{2} \sum_{j',j\in\mathbb{Z}} a_{j,j'} \left[e^{i(j\theta+j'\eta)} + e^{i(j\eta+j'\theta)} \right]$$
$$= \frac{1}{2} \left(f(\theta,\theta) + f(\eta,\eta) \right) + f_1(\theta,\eta)$$

where

$$f_1(\theta,\eta) = \frac{1}{2} \sum_{j',j \in \mathbb{Z}} a_{j,j'} \mu_{j,j'}(\theta,\eta) e^{i(j\theta+j'\eta)}$$

$$(4.5)$$

and with

$$\mu_{j,j'}(\theta,\eta) = \left(1 - e^{\mathrm{i}j(\eta-\theta)}\right) \left(1 - e^{-\mathrm{i}j'(\eta-\theta)}\right).$$

Using elementary trigonometric identities combined using Chebyshev polynomials of second order U_n that

$$\mu_{j,j'}(\theta,\eta) = -4\sin\left(j'\frac{\theta-\eta}{2}\right)\sin\left(j\frac{\theta-\eta}{2}\right)e^{\frac{i}{2}(j'-j)(\theta-\eta)}$$
$$= -4\sin^2\left(\frac{\theta-\eta}{2}\right)U_{j'-1}\left(\cos\left(\frac{\theta-\eta}{2}\right)\right)U_{j-1}\left(\cos\left(\frac{\theta-\eta}{2}\right)\right)e^{\frac{i}{2}(j'-j)(\theta-\eta)}$$
$$= -4\sin^2\left(\frac{\theta-\eta}{2}\right)\mu_{j,j'}^1(\theta,\eta)$$

with

$$\mu_{j,j'}^1(\theta,\eta) \triangleq U_{j'-1}\left(\cos\left(\frac{\theta-\eta}{2}\right)\right) U_{j-1}\left(\cos\left(\frac{\theta-\eta}{2}\right)\right) e^{\frac{1}{2}i(j'-j)(\theta-\eta)}.$$
(4.6)

Here, we have changed slightly the convention for U_j , it is defined as the polynomial such that

$$\sin(j\theta) = \sin(\theta)U_j(\cos(\theta)), \forall \theta \in \mathbb{R}.$$

With this convention we have $U_0 \equiv 0$ and it is still defined for negative integer, with the identity: $U_{-j} = -U_j$. Plugging this into (4.5) allows to get

$$f_1(\theta, \eta) = \sin^2\left(\frac{\theta-\eta}{2}\right)g(\theta, \eta)$$

with

$$g(\theta,\eta) = -2\sum_{j',j\in\mathbb{Z}} a_{j,j'} \mu_{j,j'}^1(\theta,\eta) e^{\mathrm{i}(j\theta+j'\eta)}.$$
(4.7)

It follows that

$$g(\theta, \theta + \eta) = -2\sum_{j', j \in \mathbb{Z}} a_{j,j'} \mu_{j,j'}^1(\theta, \theta + \eta) e^{ij' \cdot \eta} e^{i(j+j')\theta}$$

One can check easily that

$$\mu_{j,j'}^{1}(\theta,\theta+\eta) = U_{j'-1}\left(\cos\left(\frac{\eta}{2}\right)\right)U_{j-1}\left(\cos\left(\frac{\eta}{2}\right)\right)e^{\frac{1}{2}i(j-j')\eta}$$

$$= \mu_{j,j'}^{1}(0,\eta).$$
(4.8)

Thus we find after a change of indices

$$g(\theta, \theta + \eta) = -2 \sum_{j', j \in \mathbb{Z}^d} a_{j,j'} \,\mu_{j,j'}^1(0,\eta) \,e^{ij'\cdot\eta} e^{i(j+j')\theta}$$
$$= -2 \sum_{j', j \in \mathbb{Z}} a_{j-j',j'} \,\mu_{j-j',j'}^1(0,\eta) \,e^{ij'\eta} e^{ij\theta}.$$

Therefore

$$\|g(\cdot, \cdot + \eta)\|_{H^s}^2 = 4 \sum_{j \in \mathbb{Z}} \langle j \rangle^{2s} \Big| \sum_{j' \in \mathbb{Z}} a_{j-j',j'} \, \mu_{j-j',j'}^1(0,\eta) e^{\mathbf{i}j'\eta} \Big|^2.$$

Using Cauchy-Schwarz inequality we obtain

$$\begin{split} \Big| \sum_{j' \in \mathbb{Z}} a_{j-j',j'} \, \mu_{j-j',j'}^1(0,\eta) e^{\mathbf{i}j' \cdot \eta} \Big|^2 &\leq \left(\sum_{j' \in \mathbb{Z}} \frac{1}{\langle j' - j \rangle^{1+2\epsilon}} \right) \sum_{j' \in \mathbb{Z}} \langle j' - j \rangle^{1+2\epsilon} |a_{j-j',j'}|^2 \, |\mu_{j-j',j'}^1(0,\eta)|^2 \\ &\lesssim \sum_{j' \in \mathbb{Z}} \langle j' - j \rangle^{1+2\epsilon} |a_{j-j',j'}|^2 \, |\mu_{j-j',j'}^1(0,\eta)|^2. \end{split}$$

Now, from (4.8) we get, for any $k \in \mathbb{N}$ there exists C > 0 such that

$$\forall j \in \mathbb{Z}, \quad \sup_{\theta \in [0,2\pi]} \left| \partial_{\theta}^k (U_j(\cos \theta)) \right| \leq C |j|^{k+1}$$

Combining this with (4.6) together with Leibniz formula yield

$$\sup_{\theta,\eta\in\mathbb{T}} \left(|\partial_{\theta}^{k} \mu_{j,j'}^{1}(\theta,\eta)| + |\partial_{\eta}^{k} \mu_{j,j'}^{1}(\theta,\eta)| \right) \lesssim \left(|j| + |j'| \right)^{k+2}.$$

$$\tag{4.9}$$

Applying this for k = 0 implies

$$\sum_{j' \in \mathbb{Z}} \langle j' - j \rangle^{1+2\epsilon} |a_{j-j',j'}|^2 |\mu_{j-j',j'}^1(0,\eta)|^2 \lesssim \sum_{j' \in \mathbb{Z}} \langle j' - j \rangle^{1+2\epsilon} \left(|j - j'| + |j'| \right)^4 |a_{j-j',j'}|^2.$$

It follows that for $s \ge 0$

$$\begin{split} \|g(\cdot,\cdot+\eta)\|_{H^s}^2 &\lesssim \sum_{j,j'\in\mathbb{Z}} \langle j'-j\rangle^{1+2\epsilon} \left(|j-j'|+|j'|\right)^4 \langle j\rangle^{2s} |a_{j-j',j'}|^2 \\ &\lesssim \sum_{j,j'\in\mathbb{Z}} \langle j\rangle^{1+2\epsilon} \left(|j|+|j'|\right)^4 \langle j'+j\rangle^{2s} |a_{j,j'}|^2 \\ &\lesssim \sum_{j,j'\in\mathbb{Z}} \langle j',j\rangle^{5+2\epsilon+2s} |a_{j,j'}|^2. \end{split}$$

Thus we deduce that

$$\sup_{\eta \in \mathbb{T}} \|g(\cdot, \cdot + \eta)\|_{H^s} \lesssim \|f\|_{H^{s+\frac{5}{2}+\epsilon}}.$$

Implementing the same approach as before and using (4.9) we get

$$\sup_{\eta \in \mathbb{T}} \left(\|\partial_{\theta}^k g(\cdot, \cdot + \eta)\|_{H^s} + \|(\partial_{\eta}^k g)(\cdot, \cdot + \eta)\|_{H^s} \right) \lesssim \|f\|_{H^{s+\frac{5}{2}+k+\epsilon}}$$

The proof of the following estimates can be done in a similar way,

$$\sup_{\eta \in \mathbb{T}} \|\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} g(\cdot, \cdot + \eta)\|_{H^s} \lesssim \|f\|_{H^{s+\frac{5}{2}+p_1+p_2+\epsilon}}.$$

(ii) Splitting as before f into its Fourier expansion

$$f(\lambda,\varphi,\theta,\eta) = \sum_{\substack{j,j' \in \mathbb{Z} \\ l \in \mathbb{Z}^d}} a_{j,j',l}(\lambda) e^{i(j\theta+j'\eta+l\cdot\varphi)}.$$

Then proceeding as in the first point (i) we may obtain

$$f(\lambda,\varphi,\theta,\eta) = \frac{1}{2} \left(f(\lambda,\varphi,\theta,\theta) + f(\lambda,\varphi,\eta,\eta) \right) + \sin^2 \left(\frac{\eta-\theta}{2} \right) g(\lambda,\varphi,\theta,\eta)$$

and g takes a similar form to (4.7)

$$g(\lambda,\varphi,\theta,\eta) = -2\sum_{\substack{j',j\in\mathbb{Z}\\l\in\mathbb{Z}^d}} a_{j,j'}(\lambda)\mu^1_{j,j'}(\theta,\eta)e^{\mathrm{i}(j\theta+j'\eta+l\cdot\varphi)}.$$
(4.10)

The coefficients $\mu^1_{j,j'}$ are given by (4.6). Thus we get

$$g(\lambda,\varphi,\theta,\theta+\eta) = -2\sum_{\substack{j',j\in\mathbb{Z}\\l\in\mathbb{Z}^d}} a_{j,j',l}(\lambda)\mu^1_{j,j'}(0,\eta)e^{\mathrm{i}j'\cdot\eta}e^{\mathrm{i}((j+j')\theta+l\cdot\varphi)}.$$
(4.11)

By changing the indices

$$g(\lambda,\varphi,\theta,\theta+\eta) = -2\sum_{\substack{j',j\in\mathbb{Z}\\l\in\mathbb{Z}^d}} a_{j,j',l}(\lambda)\mu_{j,j'}^1(0,\eta)e^{\mathrm{i}j'\cdot\eta}e^{\mathrm{i}((j+j')\theta+l\cdot\varphi)}$$
$$= -2\sum_{\substack{j',j\in\mathbb{Z}\\l\in\mathbb{Z}^d}} a_{j-j',j',l}(\lambda)\mu_{j-j',j'}^1(0,\eta)e^{\mathrm{i}j'\eta}e^{\mathrm{i}(j\theta+l\cdot\varphi)}.$$

Then for $k\in \mathbb{N}^{d+1}, |k|\leqslant q$

$$\left\|\partial_{\lambda}^{k}g\left(\lambda,\cdot,\cdot,\cdot,\cdot+\eta\right)\right\|_{H^{s-|k|}}^{2}=\sum_{j\in\mathbb{Z},l\in\mathbb{Z}^{d}}\langle j,l\rangle^{2(s-|k|)}\left|\sum_{j'\in\mathbb{Z}}\partial_{\lambda}^{\mathbf{j}}a_{j-j',j',l}(\lambda)\mu_{j-j',j'}^{1}(0,\eta)e^{\mathbf{i}j'\eta}\right|^{2}.$$

Using Cauchy-Schwarz inequality we obtain for any $\epsilon>0$

$$\Big|\sum_{j'\in\mathbb{Z}}\partial_{\lambda}^{k}a_{j-j',j',l}(\lambda)\mu_{j-j',j'}^{1}(0,\eta)e^{\mathrm{i}j'\cdot\eta}\Big|^{2} \lesssim \sum_{j'\in\mathbb{Z}}\langle j'-j\rangle^{1+2\epsilon} \Big|\partial_{\lambda}^{k}a_{j-j',j',l}(\lambda)\mu_{j-j',j'}^{1}(0,\eta)\Big|^{2}.$$

Combining this with (4.9) yields

$$\sum_{j'\in\mathbb{Z}}\langle j'-j\rangle^{1+2\epsilon} \left|\partial_{\lambda}^{k}a_{j-j',j',l}(\lambda)\right|^{2}|\mu_{j-j',j'}^{1}(0,\eta)|^{2} \lesssim \sum_{j'\in\mathbb{Z}}\langle j'-j\rangle^{1+2\epsilon} \left(|j-j'|+|j'|\right)^{4} \left|\partial_{\lambda}^{k}a_{j-j',j',l}(\lambda)\right|^{2}.$$

It follows that for $s \ge q \ge |k|$,

$$\begin{split} \left\|\partial_{\lambda}^{k}g\left(\lambda,\cdot,\cdot,\cdot,\cdot,+\eta\right)\right\|_{H^{s-|k|}}^{2} &\lesssim \sum_{\substack{j,j'\in\mathbb{Z}\\l\in\mathbb{Z}^{d}}} \langle j'-j\rangle^{1+2\epsilon} \left(|j-j'|+|j'|\right)^{4} \langle j,l\rangle^{2(s-|k|)} \left|\partial_{\lambda}^{k}a_{j,j',l}(\lambda)\right|^{2} \\ &\lesssim \sum_{\substack{j,j'\in\mathbb{Z}\\l\in\mathbb{Z}^{d}}} \langle j\rangle^{1+2\epsilon} \left(|j|+|j'|\right)^{4} \langle j'+j,l\rangle^{2(s-|k|)} \left|\partial_{\lambda}^{k}a_{j,j',l}(\lambda)\right|^{2} \\ &\lesssim \sum_{\substack{j,j'\in\mathbb{Z}\\l\in\mathbb{Z}^{d}}} \langle j',j,l\rangle^{2(s-|k|+\frac{5}{2}+\epsilon)} \left|\partial_{\lambda}^{k}a_{j,j',l}(\lambda)\right|^{2}. \end{split}$$
(4.12)

It follows that for any $\lambda \in \mathcal{O}, \eta \in \mathbb{T}$ and $k \in \mathbb{N}^{d+1}, |k| \leq q$

$$\left\|\partial_{\lambda}^{k}g\left(\lambda,\cdot,\bullet,\bullet+\eta\right)\right\|_{H^{s-|k|}\left(\mathbb{T}^{d}\times\mathbb{T}\right)} \lesssim \left\|\partial_{\lambda}^{k}f\right\|_{H^{s-|k|+\frac{5}{2}+\epsilon}\left(\mathbb{T}^{d}\times\mathbb{T}^{2}\right)}.$$

Consequently

$$\sup_{\eta \in \mathbb{T}} \left\| g\left(*, \cdot, \bullet, \bullet + \eta\right) \right\|_{s}^{q, \kappa} \leqslant C \|f\|_{s + \frac{5}{2} + \epsilon}^{q, \kappa}.$$

We can also get from (4.12) that

$$\sup_{\eta \in \mathbb{T}} \left\| g\left(*, \cdot, \bullet, \bullet + \eta\right) \right\|_{s}^{q, \kappa} \leqslant C \|\Delta_{\theta, \eta} f\|_{s + \frac{1}{2} + \epsilon}^{q, \kappa}.$$

$$(4.13)$$

Now let us move to the estimate of $(\partial^p_{\theta} g)$. Applying Leibniz formula to (4.10) yields, for $p \in \mathbb{N}$,

$$\partial_{\theta}^{p}g(\lambda,\varphi,\theta,\eta) = -2\sum_{\substack{0\leqslant m\leqslant p\\l,j\in\mathbb{Z},l\in\mathbb{Z}^{d}}} {p \choose m} a_{j,j'}(\lambda)(\mathrm{i}j)^{p-m} \partial_{\theta}^{m} \mu_{j,j'}^{1}(\theta,\eta) e^{\mathrm{i}(j\theta+j'\eta+l\cdot\varphi)}.$$

It follows that

$$\begin{aligned} (\partial_{\theta}^{p}g)(\lambda,\varphi,\theta,\theta+\eta) &= -2\sum_{m=0}^{p}\sum_{j',j\in\mathbb{Z},l\in\mathbb{Z}^{d}} \binom{p}{m} a_{j,j'}(\lambda)(\mathrm{i}j)^{p-m} (\partial_{\theta}^{m}\mu_{j,j'}^{1})(0,\eta) e^{\mathrm{i}j'\cdot\eta} e^{\mathrm{i}((j+j')\theta+l\cdot\varphi)} \\ &= -2\sum_{m=0}^{p}\sum_{j',j\in\mathbb{Z};l\in\mathbb{Z}^{d}} \binom{p}{m} a_{j-j',j'}(\lambda)(\mathrm{i}j)^{p-m} (\partial_{\theta}^{m}\mu_{j-j',j'}^{1})(0,\eta) e^{\mathrm{i}j'\cdot\eta} e^{\mathrm{i}(j\theta+l\cdot\varphi)} \end{aligned}$$

Then from (4.9) and proceeding as before we get

$$\begin{aligned} \left\|\partial_{\lambda}^{k}(\partial_{\theta}^{p}g)\left(\lambda,\cdot,\cdot,\cdot,\cdot+\eta\right)\right\|_{H^{s-|k|}} &\lesssim \sum_{\substack{j,j'\in\mathbb{Z}\\l\in\mathbb{Z}^{d}}} \langle j'-j\rangle^{1+2\epsilon} \left(|j-j'|+|j'|\right)^{4+2p} \langle j,l\rangle^{2(s-|k|)} \left|\partial_{\lambda}^{k}a_{j-j',j',l}(\lambda)\right|^{2} \\ &\lesssim \sum_{\substack{j,j'\in\mathbb{Z}\\l\in\mathbb{Z}^{d}}} \langle j',j,l\rangle^{2(s-|k|+\frac{5}{2}+\epsilon+p)} \left|\partial_{\lambda}^{k}a_{j,j',l}(\lambda)\right|^{2}. \end{aligned}$$

$$(4.14)$$

Hence, for any $\lambda \in \mathcal{O}, \eta \in \mathbb{T}$ and $p \in \mathbb{N}$,

$$\left\|\partial_{\lambda}^{k}(\partial_{\theta}^{p}g)\left(\lambda,\cdot,\cdot,\cdot,\cdot+\eta\right)\right\|_{H^{s-|k|}(\mathbb{T}^{d}\times\mathbb{T})} \lesssim \left\|\partial_{\lambda}^{k}f\right\|_{H^{s_{1}-|k|+\frac{5}{2}+p+\epsilon}(\mathbb{T}^{d}\times\mathbb{T}^{2})}.$$

Consequently

$$\sup_{\eta \in \mathbb{T}} \left\| (\partial_{\theta}^{p} g) \big(*, \cdot, \bullet, \bullet + \eta \big) \right\|_{s}^{q, \kappa} \leqslant C \| f \|_{s + \frac{5}{2} + p + \epsilon}^{q, \kappa}.$$

Following the same lines by applying in particular Leibniz formula to (4.10) yields, for $p_1, p_2 \in \mathbb{N}$,

$$\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} g(\lambda, \varphi, \theta, \eta) = 2 \sum_{\substack{0 \leqslant m_1, m_2 \leqslant p \\ j', j \in \mathbb{Z} \ l \in \mathbb{Z}^d}} {\binom{p_1}{m_1} \binom{p_2}{m_2} a_{j,j'}(\lambda) (\mathrm{i}n)^{p_1 - m_1} (\mathrm{i}j')^{p_2 - m_2} \partial_{\theta}^{m_1} \partial_{\eta}^{m_2} \mu_{j,j'}^1(\theta, \eta) e^{\mathrm{i}(j\theta + j'\eta + l \cdot \varphi)}.$$

This gives the desired estimate.

Next we shall prove the following lemma which turns out to be very useful.

Lemma 4.7. Let $f: \mathcal{O} \times \mathbb{T}^d \times \mathbb{T} \to \mathbb{C}$ be a smooth function and define $g: \mathcal{O} \times \mathbb{T}^d \times \mathbb{T}^2 \to \mathbb{C}$ by

$$g(\lambda,\varphi,\theta,\eta) = \begin{cases} \frac{f(\lambda,\varphi,\eta+\theta) - f(\lambda,\varphi,\theta)}{\tan\left(\frac{\eta}{2}\right)} & \text{if } \eta \in \mathbb{R} \setminus 2\pi\mathbb{Z} \\ 2\partial_{\theta}f(\lambda,\varphi,\theta) & \text{if } \eta \in 2\pi\mathbb{Z}. \end{cases}$$

Then the followings assertions hold true.

(i) For any $s \in \mathbb{R}$, we have

$$\sup_{\eta \in \mathbb{T}} \|g(*,\cdot,\cdot,\eta)\|_s^{\gamma,q} \lesssim \|f\|_{s+1}^{q,\kappa}.$$

(ii) For any $s \ge 0$

$$\begin{aligned} \|g(*,\cdot,\bullet)\|_{s}^{q,\kappa} &\lesssim \|\partial_{\theta}f\|_{s}^{q,\kappa} \\ &\lesssim \|f\|_{s+1}^{q,\kappa}, \end{aligned}$$

where the symbols $*, \cdot, \cdot$ denote λ, φ and (θ, η) .

Proof. (i) We start with expanding f into its Fourier series,

$$f(\varphi,\theta) = \sum_{(l,j)\in\mathbb{Z}^{d+1}} f_{l,j} e^{\mathrm{i}(l\cdot\varphi+j\theta)}.$$
(4.15)

Therefore we get from the Bessel's identity combined with some trigonometric identities

$$\begin{split} \|g(\cdot,\cdot,\eta)\|_{H^s_{\varphi,\theta}}^2 &= \left\|\sum_{(l,j)\in\mathbb{Z}^{d+1}} f_{l,j} \frac{e^{ij\eta}-1}{\tan\left(\frac{\eta}{2}\right)} e^{i(l\cdot\varphi+j\theta)}\right\|_{H^s_{\varphi,\theta}}^2 \\ &\leqslant \left\|\sum_{(l,j)\in\mathbb{Z}^{d+1}} f_{l,j} e^{i\frac{j\eta}{2}} \frac{\sin\left(\frac{j\eta}{2}\right)}{\tan\left(\frac{\eta}{2}\right)} e^{i(l\cdot\varphi+j\theta)}\right\|_{H^s_{\varphi,\theta}}^2 \\ &\lesssim \sum_{(l,j)\in\mathbb{Z}^{d+1}} \langle l,j\rangle^{2s} |f_{l,j}|^2 \frac{\sin^2\left(\frac{j\eta}{2}\right)}{\sin^2\left(\frac{\eta}{2}\right)}. \end{split}$$

Using the bound $\frac{\sin^2(\frac{j\eta}{2})}{\sin^2(\frac{\eta}{2})} \lesssim |j|$ following from basic properties of Féjer kernel, we infer

$$\begin{aligned} \|g(\cdot,\cdot,\eta)\|_{H^s_{\varphi,\theta}}^2 &\lesssim & \sum_{\substack{(l,j)\in\mathbb{Z}^d\times\mathbb{Z}\\ \lesssim & \|\partial_\theta f\|_{H^s}^2.}} \langle l,j\rangle^{2s} |j|^2 |f_{l,j}|^2 \end{aligned}$$

This ensures the desired result for q = 0.

(ii) We first split f into its Fourier expansion,

$$f(\lambda,\varphi,\theta) = \sum_{(l,j)\in\mathbb{Z}^{d+1}} c_{l,j}(\lambda) e^{\mathrm{i}(l\cdot\varphi+j\theta)}.$$

Consequently

$$g(\lambda,\varphi,\theta,\eta) = \sum_{(l,j)\in\mathbb{Z}^{d+1}} \mu_j(\eta) \, c_{j',j}(\lambda) e^{\mathrm{i}(l\cdot\varphi+j\theta)}, \quad \mu_j(\eta) \triangleq \frac{e^{\mathrm{i}j\eta} - 1}{\tan(\frac{\eta}{2})}.$$

The latter function is \mathscr{C}^{∞} and 2π -periodic and therefore its Fourier expansion takes the form

$$\mu_j(\eta) = \sum_{j' \in \mathbb{Z}} A_{j,j'} e^{\mathbf{i}j'\eta}.$$

It follows that, for any $k \in \mathbb{N}^{d+1}$ with $|k| \leq q$,

$$\partial_{\lambda}^{k}g(\lambda,\varphi,\theta,\eta) = \sum_{(l,j,j')\in\mathbb{Z}^{d+2}} (\partial_{\lambda}^{k}c_{l,j}(\lambda))A_{j,j'}e^{\mathrm{i}(l\cdot\varphi+j\theta+j'\eta)}.$$

Thus we find

$$\|\partial_{\lambda}^{k}g(\lambda,\cdot,\cdot)\|_{H^{s-|k|}}^{2} = \sum_{(l,j,j')\in\mathbb{Z}^{d+2}} \langle l,j',j\rangle^{2(s-|k|)} \left|\partial_{\lambda}^{k}c_{l,j}(\lambda)\right|^{2} |A_{j,j'}|^{2}$$

$$\lesssim \sum_{(l,j)\in\mathbb{Z}^{d+1}} \nu_{l,j} \left|\partial_{\lambda}^{k}c_{l,j}(\lambda)\right|^{2}$$

$$(4.16)$$

with

$$\nu_{l,j} \triangleq \sum_{j' \in \mathbb{Z}} \langle l, j', j \rangle^{2(s-|k|)} |A_{j,j'}|^2.$$

Since $s \ge q \ge |k|$, then we can easily check from the inequalities $(|a| + |b|)^s \le |a|^s + |b|^s$ that

$$\nu_{l,j} \lesssim \sum_{j' \in \mathbb{Z}} \langle l, j \rangle^{2(s-|k|)} |A_{j,j'}|^2 + \sum_{j' \in \mathbb{Z}} \langle j' \rangle^{2(s-|k|)} |A_{j,j'}|^2$$

This can be written in the form

$$\nu_{l,j} \lesssim \langle l,j \rangle^{2(s-|k|)} \|\mu_j\|_{L^2}^2 + \|\mu_j\|_{H^{s-|k|}}^2$$

Next we claim that for any $s \in \mathbb{R}_+$

$$\forall j \in \mathbb{Z}, \quad \|\mu_j\|_{H^s} \lesssim |j|^{\frac{1}{2}+s}. \tag{4.17}$$

Let us give the main ideas of the proof. The case s = 0 can be done in the following way

$$\|\mu_{j}\|_{L^{2}}^{2} \lesssim \int_{0}^{\pi} \frac{\sin^{2}(j\eta/2)}{\sin^{2}(\eta/2)} d\eta$$
$$\lesssim \int_{0}^{\frac{\pi}{2}} \frac{\sin^{2}(j\eta)}{\sin^{2}(\eta)} d\eta.$$

Then using the inequality $\forall \eta \in [0, \frac{\pi}{2}]$, $\sin(\eta) \ge \frac{2\eta}{\pi}$, we find through change of variables

$$\|\mu_j\|_{L^2}^2 \lesssim \int_0^{\frac{\pi}{2}} \frac{\sin^2(j\eta)}{\eta^2} d\eta$$
$$\lesssim |j| \int_0^{|j|\frac{\pi}{2}} \frac{\sin^2(\eta)}{\eta^2} d\eta$$
$$\lesssim |j|.$$

The second step is to establish the estimate for $s \in \mathbb{N}^*$ by direct differentiation and using similar arguments as for s = 0. The extension of the estimate to $s \in \mathbb{R}_+$ can be obtained by interpolation inequalities.

Now using (4.17) we obtain the estimate

$$\begin{split} \nu_{l,j} &\lesssim \langle l,j \rangle^{2(s-|k|)} |j| + \langle j \rangle^{2(s-|k|)} |j| \\ &\lesssim \langle l,j \rangle^{2(s-|k|)} |j|. \end{split}$$

Inserting this estimate into (4.16) yields

$$\begin{aligned} \|\partial_{\lambda}^{k}g(\lambda,\cdot,\bullet)\|_{H^{s-|k|}}^{2} \lesssim \sum_{(l,j)\in\mathbb{Z}^{d+1}} \langle l,j\rangle^{2(s-|k|)} |j| |\partial_{\lambda}^{k}c_{l,j}(\lambda)|^{2} \\ \lesssim \|\partial_{\lambda}^{k}\partial_{\theta}f(\lambda,\cdot,\bullet)\|_{H^{s-|k|}}. \end{aligned}$$

$$(4.18)$$

From (4.18) and the Definition 4.1 we infer

$$\begin{aligned} \|g\|_{s}^{q,\kappa} \lesssim &\|\partial_{\theta}f\|_{s}^{q,\kappa} \\ \lesssim &\|f\|_{s+1}^{q,\kappa}. \end{aligned}$$

This achieves the proof.

4.3. Modified periodic fractional Laplacian. The main goal of this section is to explore some basic properties of a modified fractional Laplacian defined as follows. Let $h : \mathbb{T}^{d+1} \to \mathbb{R}$ and consider $\alpha \in (0, 1)$, we define the modified partial fractional Laplacian by

$$|\mathbf{D}|^{-\alpha}h(\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{h(\varphi,\theta-\eta)}{|\sin\left(\frac{\eta}{2}\right)|^{1-\alpha}} d\eta.$$
(4.19)

As we shall see for instance in (4.23), this operator is different from the standard fractional Laplacian denoted by $\Lambda^{-\alpha} = (-\Delta)^{-\frac{\alpha}{2}}$ but asymptotically in Fourier side they have the same behavior.

To state some results connected with the action of the modified fractional Laplacian we need to introduce Sobolev anisotropic spaces. First, we define the anisotropic Fourier multiplier Λ^{s_1,s_2} as follows: For $s_1, s_2 \in \mathbb{R}$

$$\Lambda^{s_1, s_2} \mathbf{e}_{l,j} = \langle l, j \rangle^{s_1} \langle j \rangle^{s_2} \mathbf{e}_{l,j} \quad \text{with} \qquad \langle j \rangle = \max(1, |j|). \tag{4.20}$$

The anisotropic Sobolev space $H^{s_1,s_2}(\mathbb{T}^{d+1};\mathbb{C})$ is the set of functions $h:\mathbb{T}^{d+1}\to\mathbb{C}$ such that

$$h = \sum_{(l,j)\in\mathbb{Z}^{d+1}} f_{l,j} \mathbf{e}_{l,j}, \quad h_{l,j}\in\mathbb{C} \quad \text{and} \quad \|h\|_{H^{s_1,s_2}}^2 = \sum_{(l,j)\in\mathbb{Z}^{d+1}} \langle l,j\rangle^{2s_1} \langle j\rangle^{2s_2} |h_{l,j}|^2 < \infty.$$

The main goal here is to collect some elementary results related to the action of the modified fractional Laplacian to anisotropic spaces.

Lemma 4.8. Let $s_1, s_2 \in \mathbb{R}$ and $\alpha \in (0, 1)$, then the following properties hold true.

- (i) For any p ∈ N, we have ∂^p_θ: H^{s1,s2} → H^{s1,s2-p} is continuous.
 (ii) We have |D|^{-α}: H^{s1,s2} → H^{s1,s2+α} is continuous
- (iii) For any $k \in \mathbb{N}$, then

$$\partial_{\alpha}^{k} |\mathbf{D}|^{-\alpha} : H^{s_{1},s_{2}} \to H^{s_{1},s_{2}+\alpha-\epsilon}$$

is continuous, with

$$\partial_{\alpha}^{k} |\mathbf{D}|^{-\alpha} h(\varphi, \theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\ln^{k} \left[\sin \left(\eta/2 \right) \right]}{|\sin \left(\frac{\eta}{2} \right)|^{1-\alpha}} h(\varphi, \theta - \eta) d\eta.$$

Proof. (i) This can be easily obtained from

$$h = \sum_{(l,j)\in\mathbb{Z}^{d+1}} h_{n,k} \mathbf{e}_{l,j} \Longrightarrow \partial_{\theta}^{k} h = \sum_{(l,j)\in\mathbb{Z}^{d+1}} (\mathrm{i}j)^{k} h_{l,j} \mathbf{e}_{l,j}.$$

Thus

$$\begin{aligned} \|\partial_{\theta}^{p}h\|_{H^{s_{1},s_{2}-p}}^{2} &= \sum_{(l,j)\in\mathbb{Z}^{+1}} \langle l,j\rangle^{2s_{1}} \langle j\rangle^{2(s_{2}-p)} |j|^{2l} |h_{l,j}|^{2} \\ &\leq \|h\|_{H^{s_{1},s_{2}}}^{2}. \end{aligned}$$

(ii) Using the definition and a change of variables we find

$$|\mathbf{D}|^{-\alpha} \mathbf{e}_{l,j}(\varphi, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{e^{-ij\eta}}{|\sin\left(\frac{\eta}{2}\right)|^{1-\alpha}} d\eta \ \mathbf{e}_{l,j}(\varphi, \theta)$$
$$= \frac{1}{\pi} \int_0^{\pi} \frac{e^{2ij\eta}}{|\sin\eta|^{1-\alpha}} d\eta \ \mathbf{e}_{l,j}(\varphi, \theta).$$

Applying (3.17) and using (3.20) give

$$|\mathbf{D}|^{-\alpha} \mathbf{e}_{l,j}(\varphi,\theta) = \frac{2^{1-\alpha} \Gamma(\alpha)}{\Gamma(\frac{1-\alpha}{2}) \Gamma(\frac{1+\alpha}{2})} \frac{\Gamma(|j| + \frac{1-\alpha}{2})}{\Gamma(|j| + \frac{1+\alpha}{2})} \mathbf{e}_{l,j}(\varphi,\theta)$$
(4.21)

$$=\frac{2^{1-\alpha}\Gamma(\alpha)}{\Gamma(\frac{1-\alpha}{2})\Gamma(\frac{1+\alpha}{2})}\mathbb{W}(|j|,1-\alpha)\mathbf{e}_{l,j}(\varphi,\theta)$$
(4.22)

where W was defined in Lemma 3.1-(iii). According to Lemma A.1 the asymptotic of Wallis quotient for large j is given by

$$\mathsf{W}(|j|, 1-\alpha) = \frac{1}{|j|^{\alpha}} + O\bigg(\frac{1}{|j|^{2+\alpha}}\bigg).$$

It follows that

$$|\mathbf{D}|^{-\alpha}\mathbf{e}_{l,j} = \frac{2^{1-\alpha}\Gamma(\alpha)}{\Gamma(\frac{1+\alpha}{2})\Gamma(\frac{1-\alpha}{2})} \left[\frac{1}{|j|^{\alpha}} + O\left(\frac{1}{|j|^{2+\alpha}}\right)\right] \mathbf{e}_{l,j}.$$
(4.23)

In particular, the operator $|D|^{-\alpha}$ acts like the standard fractional Laplacian $(-\Delta)^{-\frac{\alpha}{2}}$ and one gets

$$\begin{aligned} \|\mathbf{D}\|^{-\alpha}h\|_{H^{s_1,s_2+\alpha}}^2 &\leq C \sum_{(l,j)\in\mathbb{Z}^d\times\mathbb{Z}} \langle (l,j)\rangle^{2s_1} \langle j\rangle^{2(s_2+\alpha)} \langle j\rangle^{-2\alpha} |h_{j,k}|^2 \\ &\leq C \|h\|_{H^{s_1,s_2}}^2. \end{aligned}$$

(iii) We can write by definition and change of variables

$$\partial_{\alpha}^{k} |\mathbf{D}|^{-\alpha} \mathbf{e}_{l,j}(\varphi, \theta) = a_{j} \, \mathbf{e}_{l,j}(\varphi, \theta) \quad \text{with} \quad a_{j} = \frac{1}{\pi} \int_{0}^{\pi} \frac{e^{2ij\eta}}{|\sin\eta|^{1-\alpha}} \ln^{k} \big(\sin\eta\big) d\eta.$$

Let $\delta \in (0, \frac{1}{2})$ and write

$$a_j = \frac{1}{2ij\pi} \int_{\sin\eta \ge \delta} \frac{\left(e^{2ij\eta}\right)'}{|\sin\eta|^{1-\alpha}} \ln^k \left(\sin\eta\right) d\eta + \frac{1}{\pi} \int_{0 \le \sin\eta \le \delta} \frac{e^{2ij\eta}}{|\sin\eta|^{1-\alpha}} \ln^k \left(\sin\eta\right) d\eta.$$

Using integration by parts implies

$$|a_j| \lesssim |j|^{-1} \delta^{\alpha - 1} \ln^k(1/\delta) + |j|^{-1} \int_{\sin\eta \ge \delta} \frac{\left|\ln^k\left(\sin\eta\right)\right|}{|\sin\eta|^{2-\alpha}} d\eta + \frac{1}{\pi} \int_{0\leqslant\sin\eta\leqslant\delta} \frac{\left|\ln^k\left(\sin\eta\right)\right|}{|\sin\eta|^{1-\alpha}} d\eta.$$

Therefore using the fact that for any $\epsilon > 0$, $\sup_{x \in (0,1)} |x|^{\epsilon} |\ln x| < \infty$ we get

$$|a_j| \lesssim |j|^{-1} \delta^{\alpha - 1 - \epsilon} + \delta^{\alpha - \epsilon}$$

By taking $\delta = |j|^{-1}$ we deduce that

$$|a_j| \lesssim |j|^{-\alpha + \epsilon}.$$

This gives the desired result, that is,

$$\left\|\partial_{\alpha}^{k}|\mathbf{D}|^{-\alpha}h\right\|_{H^{s_{1},s_{2}+\alpha-\epsilon}}^{2} \leq C\|h\|_{H^{s_{1},s_{2}}}^{2}$$

The proof of the lemma is now achieved.

5. Hamiltonian toolbox

We shall reformulate the equation (2.28) as a perturbation of the Hamiltonian linear equation given by (3.8). More precisely, we write (2.28) in the form

$$\partial_t r = \partial_\theta \,\mathcal{L}(\alpha)(r) + X_P(r) \tag{5.1}$$

where $L(\alpha)$ is defined in (3.10) and X_P is the Hamiltonian vector field generated by the higher order terms $P \triangleq H_{\geq 3}(r)$, namely

$$X_P(r) = (\Omega + V_{0,\alpha}) \,\partial_\theta r - \partial_\theta \,\mathbb{K}_{0,\alpha} r - F_\alpha[r], \tag{5.2}$$

where the non-local contribution $F_{\alpha}[r]$ is introduced in (2.12), the constant $V_{0,\alpha}$ is defined in (3.11) and the operator $\mathbb{K}_{0,\alpha}$ is given by (3.12).

Since we are going to construct small quasi-periodic solutions with small amplitude then it is convenient to rescale the unknown as follows $r \mapsto \varepsilon r$, leading to

$$\mathscr{H}_{\varepsilon}(r) \triangleq \varepsilon^{-2} H(\varepsilon r) = H_{\mathrm{L}}(r) + \varepsilon P_{\varepsilon}(r) \quad \text{where} \quad P_{\varepsilon} \triangleq \varepsilon^{-3} P(\varepsilon r)$$
(5.3)

and $H_{\rm L}(r)$ is the quadratic Hamiltonian in (3.9). Therefore the equation (2.16) writes

$$\partial_t r = \partial_\theta \operatorname{L}(\alpha) r + \varepsilon X_{P_\varepsilon}(r), \quad \text{with} \quad X_{P_\varepsilon}(r) \triangleq \varepsilon^{-2} X_P(\varepsilon r).$$
 (5.4)

5.1. Action-angle coordinates. We consider finitely many "tangential" sites

$$\mathbb{S} \triangleq \{n_1, \dots, n_d\}, \quad 1 \leqslant n_1 < n_2 < \dots < n_d \tag{5.5}$$

and we introduce the set

$$\overline{\mathbb{S}} \triangleq \left\{ \pm n, \, n \in \mathbb{S} \right\} \quad \text{and} \quad \mathbb{S}_0 \triangleq \overline{\mathbb{S}} \cup \{0\}.$$
(5.6)

Denote the unperturbed tangential and normal frequency vectors by

$$\omega_{\mathrm{Eq}} \triangleq (\Omega_j(\alpha))_{j \in \mathbb{S}}, \qquad \Omega(\alpha) \triangleq (\Omega_j(\alpha))_{j \in \mathbb{Z} \setminus \mathbb{S}_0}$$

$$(5.7)$$

where $\Omega_j(\alpha)$ is given by (3.14). We decompose the phase space

$$L_0^2(\mathbb{T}) \triangleq \left\{ r(\theta) = \sum_{j \in \mathbb{Z} \setminus \{0\}} r_j e^{\mathbf{i}j\theta}; \ \overline{r_j} = r_{-j}, \ \|r\|_s^2 = \sum_{j \in \mathbb{Z} \setminus \{0\}} |r_j|^2 < +\infty \right\}$$

as the direct sum

$$L_{0}^{2}(\mathbb{T}) = \mathbb{H}_{\overline{\mathbb{S}}} \oplus \mathbb{H}_{\overline{\mathbb{S}}_{0}}^{\perp} \quad \text{with} \quad \mathbb{H}_{\overline{\mathbb{S}}} \triangleq \left\{ v = \sum_{j \in \overline{\mathbb{S}}} r_{j} e^{\mathbf{i}j\theta}; \ \overline{r_{j}} = r_{-j} \right\},$$

$$\mathbb{H}_{\mathbb{S}_{0}}^{\perp} \triangleq \left\{ z = \sum_{j \in \mathbb{Z} \setminus \mathbb{S}_{0}} z_{j} e^{\mathbf{i}j\theta} \in L_{0}^{2}(\mathbb{T}) \right\}.$$
(5.8)

The orthogonal projectors $\Pi_{\overline{\mathbb{S}}}$ and $\Pi_{\mathbb{S}_0}^{\perp}$ are defined as follows, for $r \in L^2_0(\mathbb{T})$,

$$\Pi_{\overline{\mathbb{S}}}r \triangleq \sum_{j \in \overline{\mathbb{S}}} r_j e^{\mathbf{i}j\theta}, \qquad \Pi_{\mathbb{S}_0}^{\perp}r \triangleq \sum_{j \in \mathbb{Z} \setminus \mathbb{S}_0} r_j e^{\mathbf{i}j\theta}.$$
(5.9)

On the finite dimensional space $\mathbb{H}_{\overline{S}}$ we shall use a local parametrization through the action-angle variables. Fix some positive amplitudes $(\mathbf{a}_j)_{j\in\mathbb{S}}\in(\mathbb{R}^*_+)^d$ such that $\mathbf{a}_{-j}=\mathbf{a}_j$ and define the finite dimensional torus,

$$\mathbf{T}^{d} \triangleq \left\{ \boldsymbol{\theta} \in \mathbb{R} \mapsto \sum_{j \in \overline{\mathbb{S}}} \mathbf{c}_{j} \mathbf{a}_{j} e^{\mathbf{i} j \boldsymbol{\theta}}; \ \mathbf{c}_{j} \in \mathbb{C}, \ |\mathbf{c}_{j}| = 1, \ c_{j} = \overline{c_{j}} \right\}$$

Now we shall introduce action-angle variables $(\vartheta, I) = ((\vartheta_j)_{j \in \overline{\mathbb{S}}}, (I_j)_{j \in \overline{\mathbb{S}}})$ allowing to describe the phase space around the linear torus \mathbf{T}^d , as follows

$$\begin{cases} r_j = \sqrt{\mathbf{a}_j^2 + \frac{|j|}{2\pi} I_j} e^{\mathbf{i}\vartheta_j}, & \text{if} \quad j \in \overline{\mathbb{S}}, \\ r_j = z_j, & \text{if} \quad j \in \mathbb{Z} \setminus \mathbb{S}_0 \end{cases}$$
(5.10)

where

$$I_{-j} = I_j, \quad \vartheta_{-j} = -\vartheta_j, \quad \vartheta_j, I_j \in \mathbb{R}, \quad j \in \overline{\mathbb{S}}.$$

$$(5.11)$$

Therefore we construct a local parameterization of the phase space around the linear torus \mathbf{T}^d as

$$r(\theta) = \mathbf{A}(\vartheta, I, z) \triangleq v(\vartheta, I) + z \quad \text{where}$$

$$v(\vartheta, I) \triangleq \sum_{j \in \overline{\mathbb{S}}} \sqrt{\mathbf{a}_j^2 + \frac{|j|}{2\pi} I_j} e^{\mathbf{i}(\vartheta_j + j\theta)} \quad \text{and} \quad z = \sum_{j \in \mathbb{Z} \setminus \mathbb{S}_0} r_j e^{\mathbf{i}j\theta}.$$
(5.12)

Remark that in the new coordinates system, $(\vartheta, I) = (-\omega_{\rm Eq}t, 0)$ corresponds to the solution to the linear system (3.8) given by (3.61) which belongs to the torus \mathbf{T}^d . The symplectic 2-form in (3.28) becomes

$$\mathcal{W} = \sum_{n \in \mathbb{S}} d\vartheta_n \wedge dI_n + \frac{1}{2i} \sum_{j \in \mathbb{Z} \setminus \mathbb{S}_0} \frac{1}{j} dr_j \wedge dr_{-j} = \left(\sum_{n \in \mathbb{S}} d\vartheta_n \wedge dI_n\right) \oplus \mathcal{W}_{|\mathbb{H}_{\mathbb{S}_0}^{\perp}}$$
(5.13)

where $\mathscr{W}_{|\mathbb{H}_{\mathbb{S}_0}^{\perp}}$ denotes the restriction of \mathscr{W} to $\mathbb{H}_{\mathbb{S}_0}^{\perp}$. We observe that \mathscr{W} is an exact 2-form as

$$\mathcal{W} = d\Lambda$$

where Λ is the Liouville 1-form

$$\Lambda_{(\vartheta,I,z)}[\widehat{\vartheta},\widehat{I},\widehat{z}] \triangleq -\sum_{n \in \mathbb{S}} I_n \widehat{\vartheta}_n + \frac{1}{2} (\partial_{\theta}^{-1} z, \widehat{z})_{L^2(\mathbb{T})} \,.$$
(5.14)

The Poisson bracket are given by

$$\{F,G\} = \mathcal{W}(X_F, X_G) = \langle \nabla F, \mathbf{J} \nabla G \rangle$$
(5.15)

where $\langle \cdot, \cdot \rangle$ is the inner product, defined by

$$\langle (\vartheta_1, I_1, z_1), (\vartheta_2, I_2, z_2) \rangle = \vartheta_1 \cdot \vartheta_2 + I_1 \cdot I_2 + (z_1, z_2)_{L^2(\mathbb{T})}.$$

The Poisson structure **J** corresponding to \mathcal{W} , defined by the identity (5.15), is the unbounded operator

$$\mathbf{J}: (\vartheta, I, z) \mapsto (I, -\vartheta, \partial_{\theta} z).$$

Now we shall study the Hamiltonian system generated by the Hamiltonian $\mathscr{H}_{\varepsilon}$ in (5.3), in the actionangle and normal coordinates $(\vartheta, I, z) \in \mathbb{T}^{\nu} \times \mathbb{R}^{\nu} \times \mathbb{H}_{\mathbb{S}_0}^{\perp}$. We consider the Hamiltonian $H_{\varepsilon}(\vartheta, I, z)$ defined by

$$H_{\varepsilon} = \mathscr{H}_{\varepsilon} \circ \mathbf{A} \tag{5.16}$$

where **A** is the map defined in (5.12). Since $L(\alpha)$ in (3.10) preserves the subspace $\mathbb{H}_{\mathbb{S}_0}^{\perp}$ then the quadratic Hamiltonian H_L in (3.9) (see (3.26)) in the variables (ϑ, I, z) reads, up to a constant,

$$H_N \circ \mathbf{A} = -\sum_{n \in \mathbb{S}} \Omega_n(\alpha) I_n + \frac{1}{2} (\mathcal{L}(\alpha) \, z, z)_{L^2} = -\omega_{\mathrm{Eq}} \cdot I + \frac{1}{2} (\mathcal{L}(\alpha) \, z, z)_{L^2}$$
(5.17)

where $\omega_{\text{Eq}} \in \mathbb{R}^d$ is the unperturbed tangential frequency vector. By (5.3) and (5.17), the Hamiltonian H_{ε} in (5.16) reads

$$H_{\varepsilon} = \mathcal{N} + \varepsilon \mathcal{P}_{\varepsilon} \quad \text{with} \\ \mathcal{N} \triangleq -\omega_{\mathrm{Eq}} \cdot I + \frac{1}{2} (\mathrm{L}(\alpha) \, z, z)_{L^{2}}, \qquad \mathcal{P}_{\varepsilon} \triangleq P_{\varepsilon} \circ \mathbf{A}.$$

$$(5.18)$$

We look for an embedded invariant torus

$$i: \mathbb{T}^d \to \mathbb{R}^d \times \mathbb{R}^d \times \mathbb{H}^{\perp}_{\mathbb{S}_0}, \quad \varphi \mapsto i(\varphi) \triangleq (\vartheta(\varphi), I(\varphi), z(\varphi)), \tag{5.19}$$

of the Hamiltonian vector field

$$X_{H_{\varepsilon}} \triangleq (\partial_I H_{\varepsilon}, -\partial_{\vartheta} H_{\varepsilon}, \Pi_{\mathbb{S}_0}^{\perp} \partial_{\theta} \nabla_z H_{\varepsilon})$$

filled by quasi-periodic solutions with Diophantine frequency vector ω . Note that for the value $\varepsilon = 0$, the Hamiltonian system

$$\omega \cdot \partial_{\varphi} i(\varphi) = X_{H_{\varepsilon}}(i(\varphi))$$

possesses, for any value of the parameter $\alpha \in [0, 1)$, the invariant torus

$$i_{\text{flat}}(\varphi) = (\varphi, 0, 0)$$

provided that $\omega = -\omega_{\rm Eq}$. Now we consider the family of Hamiltonians,

$$H_{\varepsilon}^{\mathsf{c}} \triangleq \mathcal{N}_{\mathsf{c}} + \varepsilon \mathscr{P}_{\varepsilon}, \quad \mathcal{N}_{\mathsf{c}} \triangleq \mathsf{c} \cdot I + \frac{1}{2} (\mathrm{L}(\alpha) \, z, z)_{L^2},$$

$$(5.20)$$

which depend on the constant vector $\mathbf{c} \in \mathbb{R}^d$. For the value $\mathbf{c} = -\omega_{\mathrm{Eq}}$ we have $H_{\varepsilon}^{\mathbf{c}} = H_{\varepsilon}$. The parameter \mathbf{c} is introduced in order to control the average in the *I*-component of the linearized equations. We look for zeros of the nonlinear operator

$$\mathcal{F}(i, \mathbf{c}) \triangleq \mathcal{F}(i, \mathbf{c}, \omega, \alpha, \varepsilon) \triangleq \omega \cdot \partial_{\varphi} i(\varphi) - X_{H_{\varepsilon}^{\mathbf{c}}}(i(\varphi)) = \omega \cdot \partial_{\varphi} i(\varphi) - (X_{\mathcal{N}_{\mathbf{c}}} + \varepsilon X_{\mathcal{P}_{\varepsilon}})(i(\varphi))$$
$$\triangleq \begin{pmatrix} \omega \cdot \partial_{\varphi} \vartheta(\varphi) - \mathbf{c} - \varepsilon \partial_{I} \mathcal{P}_{\varepsilon}(i(\varphi)) \\ \omega \cdot \partial_{\varphi} I(\varphi) + \varepsilon \partial_{\vartheta} \mathcal{P}_{\varepsilon}(i(\varphi)) \\ \omega \cdot \partial_{\varphi} z(\varphi) - \partial_{\theta} \mathcal{L}(\alpha) z(\varphi) - \varepsilon \partial_{\theta} \nabla_{z} \mathcal{P}_{\varepsilon}(i(\varphi)) \end{pmatrix}$$
(5.21)

where $\vartheta(\varphi) - \varphi$ is a $(2\pi)^d$ -periodic function. We denote by

$$\Im(\varphi) \triangleq i(\varphi) - (\varphi, 0, 0) = (\vartheta(\varphi) - \varphi, I(\varphi), z(\varphi))$$
(5.22)

the periodic component of the torus $\varphi \mapsto i(\varphi)$. Note that the involution \mathfrak{S} , described in Proposition 2.1-(iii), becomes

$$\widetilde{\mathfrak{S}}: (\vartheta, I, z) \mapsto (-\vartheta, I, \mathfrak{S}z).$$
(5.23)

Moreover, we can easily check that the Hamiltonian vector field $X_{H_{\varepsilon}^{c}}$ is reversible with respect $\widetilde{\mathfrak{S}}$. Thus, it is natural to look for reversible solutions of $\mathscr{F}(i, \mathbf{c}) = 0$, namely satisfying,

$$\vartheta(-\varphi) = -\vartheta(\varphi), \quad I(-\varphi) = I(\varphi), \quad z(-\varphi) = (\mathfrak{S}z)(\varphi). \tag{5.24}$$

The norm of the periodic component of the embedded torus (5.22) is

$$\|\mathfrak{I}\|_{s}^{q,\overline{\gamma}} \triangleq \|\vartheta - \mathrm{Id}\|_{s}^{q,\overline{\gamma}} + \|I\|_{s}^{q,\overline{\gamma}} + \|z\|_{s}^{q,\overline{\gamma}}.$$
(5.25)

We emphasize that we have used for the tangential variables, depending only on the variable φ , the norm (4.3) associated to functions with two variables (φ, θ) . In this case the norm $H^s(\mathbb{T}^{d+1})$ coincides with the norm $H^s(\mathbb{T}^d)$. Concerning the index regularity q it is an arbitrary integer that will be fixed only at the end of the paper in Section 10.2 devoted to the final Cantor set measure. There we shall make the choice

$$q \triangleq q_0 + 1. \tag{5.26}$$

with q_0 being the non-degeneracy index provided by Proposition 3.3, which only depends on the linear unperturbed frequencies.

5.2. Hamiltonian regularity. We shall analyze in this section some regularity aspects of the Hamiltonian vector field X_P described by (5.2) as well as the rescaled one associated to $\mathscr{P}_{\varepsilon}$ given by (5.18). The arguments are classical and the computations turn out to be long and straightforward. For this reason we will only sketch the proofs only for some essential points. The first main result reads as follows.

Lemma 5.1. Let (κ, q, s_0, s) satisfying (4.1). There exists $\varepsilon_0 \in (0, 1]$ such that if

$$\|r\|_{s_0+2}^{q,\kappa} \leqslant \varepsilon_0, \tag{5.27}$$

then the vector field X_P , given by (5.2), satisfies the following estimates :

- (i) $||X_P(r)||_s^{q,\kappa} \lesssim ||r||_{s+1}^{q,\kappa}$ (ii) $||d_r X_P(r)[h]||_s^{q,\kappa} \lesssim ||h||_{s+1}^{q,\kappa} + ||r||_{s+2}^{q,\kappa} ||h||_{s_0+1}^{q,\kappa}$ (iii) $||d_r^2 X_P(r)[h,h]||_s^{q,\kappa} \lesssim ||h||_{s+2}^{q,\kappa} ||h||_{s_0+1}^{q,\kappa} + ||r||_{s+2}^{q,\kappa} (||h||_{s_0+1}^{q,\kappa})^2$.

Proof. (i) Recall from (5.2) that

$$X_P(r) = (\Omega + V_{0,\alpha}) \,\partial_\theta r - \partial_\theta \,\mathbb{K}_{0,\alpha} r - F_\alpha[r].$$

where we have by (2.12),

$$F_{\alpha}[r](\varphi,\theta) = \frac{C_{\alpha}}{2\pi} \int_{0}^{2\pi} \frac{\partial_{\theta\eta}^{2} \left(R(\varphi,\theta)R(\varphi,\eta)\sin(\eta-\theta)\right)}{A_{r}^{\frac{\alpha}{2}}(\varphi,\theta,\eta)} d\eta.$$
(5.28)

and from (3.16) and (4.19),

$$\mathbb{K}_{0,\alpha} = 2^{-\alpha} C_{\alpha} |\mathbf{D}|^{\alpha - 1}.$$
(5.29)

We shall first estimate the term $\partial_{\theta} \mathbb{K}_{0,\alpha} r$. Using Lemma 4.8-(iii) combined with Leibniz formula we obtain, for all $\epsilon \in (0,1)$ and $\lambda = (\omega, \alpha) \in \mathcal{O}$,

$$\begin{split} \|\partial_{\lambda}^{k}\partial_{\theta}[|\mathbf{D}|^{\alpha-1}r(\lambda,\boldsymbol{\cdot})]\|_{H^{s-|k|}} &\lesssim_{k} \sum_{j\leqslant k} \|\partial_{\lambda}^{j}|\mathbf{D}|^{\alpha-1}\partial_{\lambda}^{k-j}(\partial_{\theta}r)(\lambda,\boldsymbol{\cdot})\|_{H^{s-|k|}} \\ &\lesssim_{k} \sum_{j\leqslant k} \|\partial_{\lambda}^{k-j}r(\lambda,\boldsymbol{\cdot})\|_{H^{s+\alpha+\epsilon-|k|}}. \end{split}$$

Combining the last estimate together with (5.29) and Definition 4.1 we find

$$\|\partial_{\theta} \mathbb{K}_{0,\alpha} r\|_{s}^{q,\kappa} \lesssim \|r\|_{s+\overline{\alpha}+\epsilon}^{q,\kappa}.$$
(5.30)

Next we shall move to the estimate of $F_{\alpha}[r]$. From (2.11) we deduce the identity

$$A_r(\varphi,\theta,\eta) = \sin^2\left(\frac{\eta-\theta}{2}\right) \left[\left(\frac{R(\varphi,\eta)-R(\varphi,\theta)}{\sin\left(\frac{\eta-\theta}{2}\right)}\right)^2 + 4R(\varphi,\eta)R(\varphi,\theta) \right]$$
$$= \sin^2\left(\frac{\eta-\theta}{2}\right) \left[\left(\frac{R(\varphi,\eta)-R(\varphi,\theta)}{\tan\left(\frac{\eta-\theta}{2}\right)}\right)^2 + \left(R(\varphi,\eta)+R(\varphi,\theta)\right)^2 \right].$$

It follows that

$$C_{\alpha}A_{r}^{-\frac{\alpha}{2}}(\varphi,\theta,\eta) = \left|\sin\left(\frac{\eta-\theta}{2}\right)\right|^{-\alpha}\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta)$$
(5.31)

with

$$\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta) = C_{\alpha} \left[\left(\frac{R(\varphi,\eta) - R(\varphi,\theta)}{\tan\left(\frac{\eta-\theta}{2}\right)} \right)^2 + \left(R(\varphi,\eta) + R(\varphi,\theta) \right)^2 \right]^{-\frac{\omega}{2}}.$$
(5.32)

Applying Lemmata 4.4-4.1-4.7 combined with the smallness condition (9.11), we obtain

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{A}_{r,\alpha}(\cdot, \cdot, \eta + \cdot) - 2^{-\alpha} C_{\alpha}\|_{s}^{q,\kappa} \leqslant C \|r\|_{s+1}^{q,\kappa}.$$
(5.33)

According to (5.31) combined with a change of variables in the expression of $F_{\alpha}[r]$, one gets

$$F_{\alpha}[r](\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{B}_{r,\alpha}^{0}(\varphi,\theta,\eta)}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta$$
(5.34)

with

$$\begin{aligned} \mathscr{B}^{0}_{r,\alpha}(\varphi,\theta,\eta) &\triangleq g_{0}(\varphi,\theta,\eta)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta), \\ g_{0}(\varphi,\theta,\eta) &\triangleq \left[\partial_{\theta}R(\varphi,\theta)(\partial_{\eta}R)(\varphi,\eta+\theta) + R(\varphi,\theta)R(\varphi,\eta+\theta)\right]\sin(\eta) \\ &+ \left[\partial_{\theta}R(\varphi,\theta)R(\varphi,\eta+\theta) - (\partial_{\eta}R)(\varphi,\eta+\theta)R(\varphi,\theta)\right]\cos(\eta). \end{aligned}$$

We can easily check by symmetry that

$$F_{\alpha}[0](\varphi,\theta) = \frac{2^{-\alpha}C_{\alpha}}{2\pi} \int_{0}^{2\pi} \frac{\sin\eta}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta$$

= 0. (5.35)

By the law products in Lemma 4.1, the composition law in Lemma 4.4 and (5.27) we infer

$$\sup_{\eta \in \mathbb{T}} \|g_0(\cdot, \cdot, \eta) - \sin(\eta)\|_s^{q,\kappa} \lesssim \|r\|_{s+1}^{q,\kappa}.$$
(5.36)

Hence using once again the law products of Lemma 4.1, (5.27), (5.33) and (5.36) we find

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{B}^{0}_{r,\alpha}(\cdot, \bullet, \eta) - \mathscr{B}^{0}_{0,\alpha}(\cdot, \bullet, \eta)\|_{s}^{q,\kappa} = \sup_{\eta \in \mathbb{T}} \|\mathscr{B}^{0}_{r,\alpha}(\cdot, \bullet, \eta) - 2^{-\alpha}C_{\alpha}\sin(\eta)\|_{s}^{q,\kappa}$$
$$\lesssim \|r\|_{s+1}^{q,\kappa}.$$
(5.37)

Finally, from (5.34), (5.35), (5.37) and straightforward computations lead to

$$\|F_{\alpha}[r]\|_{s}^{q,\kappa} \lesssim \int_{0}^{2\pi} \frac{(1+|\ln(\sin(\eta/2))|^{q})}{|\sin(\frac{\eta}{2})|^{\frac{1}{2}}} \sup_{\eta \in \mathbb{T}} \|\mathscr{B}_{r,\alpha}^{0}(\cdot, \cdot, \eta) - \mathscr{B}_{0,\alpha}^{0}\|_{s}^{q,\kappa} d\eta$$
$$\lesssim \|r\|_{s+1}^{q,\kappa}.$$
(5.38)

Putting together this with (5.30) and (5.2) achieves the proof of the first point.

(ii) From Proposition 3.1 we have

$$d_r F_{\alpha}[r]h = \partial_{\theta} \left(V_{r,\alpha}h - \mathbb{K}_{r,\alpha}h \right)$$

Thus, by (5.2) we get

$$d_r X_P(r)h = -\partial_\theta \left((V_{r,\alpha} - V_{0,\alpha})h \right) + \partial_\theta \left(\mathbb{K}_{r,\alpha}h - \mathbb{K}_{0,\alpha}h \right).$$
(5.39)

Combining (5.31) with (3.1) and using a change of variables we find

$$V_{r,\alpha}(\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathscr{B}_{r,\alpha}^1(\varphi,\theta,\eta)}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta$$
(5.40)

with

$$\mathcal{B}_{r,\alpha}^{1}(\varphi,\theta,\eta) \triangleq g_{1}(\varphi,\theta,\eta)\mathcal{A}_{r,\alpha}(\varphi,\theta,\eta+\theta),$$
$$g_{1}(\varphi,\theta,\eta) \triangleq \frac{(\partial_{\eta}R)(\varphi,\eta+\theta)\sin(\eta) + R(\varphi,\eta+\theta)\cos(\eta)}{R(\varphi,\theta)}.$$

Proceeding in a similar way to (5.36) and (5.37) we obtain

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{B}^{1}_{r,\alpha}(\cdot, \cdot, \eta) - \mathscr{B}^{1}_{0,\alpha}(\cdot, \cdot, \eta)\|_{s}^{q,\kappa} \lesssim \|r\|_{s+1}^{q,\kappa}$$

Then, straightforward computations lead to

$$\|V_{r,\alpha} - V_{0,\alpha}\|_{s}^{q,\kappa} \lesssim \int_{0}^{2\pi} \frac{(1 + |\ln(\sin(\eta/2))|^{q})}{|\sin(\frac{\eta}{2})|^{\frac{1}{2}}} \sup_{\eta \in \mathbb{T}} \|\mathscr{B}^{1}_{r,\alpha}(\cdot, \bullet, \eta) - \mathscr{B}^{1}_{0,\alpha}(\cdot, \bullet, \eta)\|_{s}^{q,\kappa} d\eta$$
$$\lesssim \sup_{\eta \in \mathbb{T}} \|\mathscr{B}^{1}_{r,\alpha}(\cdot, \bullet, \eta) - \mathscr{B}^{1}_{0,\alpha}(\cdot, \bullet, \eta)\|_{s}^{q,\kappa} d\eta$$
$$\lesssim \|r\|_{s+1}^{q,\kappa}.$$
(5.41)

Using the product laws in Lemma 4.1 together with (5.41) and (5.27), we obtain

$$\begin{aligned} \|\partial_{\theta} \left((V_{r,\alpha} - V_{0,\alpha})h \right) \|_{s}^{q,\kappa} &\lesssim \|V_{r,\alpha} - V_{0,\alpha}\|_{s+1}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa} + \|V_{r,\alpha} - V_{0,\alpha}\|_{s_{0}}^{q,\kappa} \|h\|_{s+1}^{q,\kappa} \\ &\lesssim \|r\|_{s+2}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa} + \|h\|_{s+1}^{q,\kappa}. \end{aligned}$$

Coming back to the second term of (5.39). From (5.31) and (3.2) we have the following identity

$$\mathbb{K}_{r,\alpha}h(\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{h(\varphi,\eta+\theta)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta)}{|\sin(\frac{\eta}{2})|^{\alpha}} d\eta.$$
(5.42)

In a similar way to (5.41) we obtain

$$\begin{aligned} \|\partial_{\theta} \left(\mathbb{K}_{r,\alpha} h - \mathbb{K}_{0,\alpha} h \right) \|_{s}^{q,\kappa} &\lesssim \|\mathbb{K}_{r,\alpha} h - \mathbb{K}_{0,\alpha} h \|_{s+1}^{q,\kappa} \\ &\lesssim \|r\|_{s+2}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa} + \|h\|_{s+1}^{q,\kappa} \end{aligned}$$

This concludes the proof of (ii).

(iii) Differentiating in r the identity (5.39) yields,

_

$$d_r^2 X_P(r)[h,h] = \partial_\theta \left(d_r \left(\mathbb{K}_{r,\alpha} h \right)[h] \right) - \partial_\theta \left(\left(d_r V_{r,\alpha}[h] \right) h \right).$$
(5.43)

For the first term of the right-hand side we write, by (5.42) and (5.32),

$$d_r \big(\mathbb{K}_{r,\alpha} h \big) [h](\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{h(\varphi,\eta+\theta) d_r \mathscr{A}_{r,\alpha} [h](\varphi,\theta,\eta+\theta)}{|\sin(\frac{\eta}{2})|^{\alpha}} d\eta$$

with

$$d_{r}\mathscr{A}_{r,\alpha}[h](\varphi,\theta,\eta) = 2C_{\alpha} \left[\frac{\frac{h(\varphi,\eta)}{R(\varphi,\eta)} - \frac{h(\varphi,\theta)}{R(\varphi,\eta)}}{\tan\left(\frac{\eta-\theta}{2}\right)} \frac{R(\varphi,\eta) - R(\varphi,\theta)}{\tan\left(\frac{\eta-\theta}{2}\right)} + \left(\frac{h(\varphi,\eta)}{R(\varphi,\eta)} + \frac{h(\varphi,\theta)}{R(\varphi,\theta)}\right) \left(R(\varphi,\eta) + R(\varphi,\theta)\right) \right] \\ \times \left[\left(\frac{R(\varphi,\eta) - R(\varphi,\theta)}{\tan\left(\frac{\eta-\theta}{2}\right)}\right)^{2} + \left(R(\varphi,\eta) + R(\varphi,\theta)\right)^{2} \right]^{-\frac{\alpha}{2}-1}.$$

Applying Lemmata 4.4-4.1-4.7 combined with the smallness condition (5.27), we obtain

$$\sup_{\eta \in \mathbb{T}} \|d_r \mathscr{A}_{r,\alpha}[h](\cdot, \bullet, \eta + \bullet)\|_s^{q,\kappa} \lesssim \|h\|_{s+1}^{q,\kappa} + \|h\|_{s_0+1}^{q,\kappa} \|r\|_{s+1}^{q,\kappa}.$$

Then, direct computations yield

$$\|d_r \big(\partial_{\theta} \mathbb{K}_{r,\alpha} h\big)[h]\|_s^{q,\kappa} \lesssim \|d_r \big(\mathbb{K}_{r,\alpha} h\big)[h]\|_{s+1}^{q,\kappa} \\ \lesssim \|h\|_{s+2}^{q,\kappa} \|h\|_{s_0+1}^{q,\kappa} + \|r\|_{s+2}^{q,\kappa} \big(\|h\|_{s_0+1}^{q,\kappa}\big)^2.$$
(5.44)

Finally, differentiating (5.40) with respect to r in the direction h gives

$$d_r V_{r,\alpha}[h](\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{d_r g_1[h](\varphi,\theta,\eta)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta)}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta + \frac{1}{2\pi} \int_0^{2\pi} \frac{g_1(\varphi,\theta,\eta)d_r\mathscr{A}_{r,\alpha}[h](\varphi,\theta,\eta+\theta)}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta.$$

Arguing as for (5.38) and (5.44) we get

$$\|\partial_{\theta} \left(d_{r} V_{r,\alpha}[h]h \right)\|_{s}^{q,\kappa} \lesssim \|h\|_{s+2}^{q,\kappa} \|h\|_{s_{0}+1}^{q,\kappa} + \|r\|_{q,s+2}^{q,\kappa} \left(\|h\|_{s_{0}+1}^{\kappa}\right)^{2}.$$

Combining the last estimate with (5.44) and (5.43) we conclude the proof of (iii).

Finally we state tame estimates for the composition operator induced by the Hamiltonian vector field

$$X_{\mathscr{P}_{\varepsilon}} = (\partial_I \mathscr{P}_{\varepsilon}, -\partial_{\vartheta} \mathscr{P}_{\varepsilon}, \Pi^{\perp}_{\mathbb{S}_0} \partial_{\theta} \nabla_z \mathscr{P}_{\varepsilon})$$

where $\mathscr{P}_{\varepsilon}$ is defined in (5.18).

Lemma 5.2. Let (κ, q, s_0, s) satisfying (4.1). There exists $\varepsilon_0 \in (0, 1)$ such that if

$$\|\mathfrak{I}\|_{s_0+2}^{q,\kappa} \leqslant \varepsilon_0,$$

then the perturbed Hamiltonian vector field $X_{\mathcal{P}_{\varepsilon}}$ satisfies the following tame estimates,

- (i) $\|X_{\mathscr{P}_{\varepsilon}}(i)\|_{s}^{q,\kappa} \lesssim 1 + \|\mathfrak{I}\|_{s+2}^{q,\kappa}$. (ii) $\|d_{i}X_{\mathscr{P}_{\varepsilon}}(i)[\hat{i}]\|_{s}^{q,\kappa} \lesssim \|\hat{i}\|_{s+2}^{q,\kappa} + \|\mathfrak{I}\|_{s+2}^{q,\kappa}\|\hat{i}\|_{s_{0}+2}^{q,\kappa}$.
- (iii) $\|d_i^2 X_{\mathscr{P}_{\varepsilon}}(i)[\hat{i},\hat{i}]\|_s^{q,\kappa} \lesssim \|\hat{i}\|_{s+2}^{q,\kappa} \|\hat{i}\|_{s+2}^{q,\kappa} + \|\mathfrak{I}\|_{s+2}^{q,\kappa} \left(\|\hat{i}\|_{s+2}^{q,\kappa}\right)^2.$

Proof. The proof is a straightforward and slightly long. It is based on Taylor expansion in r around the origin of the Hamiltonian vector field X_P detailed in (5.2). Then the estimates for the rescaled vector field in (5.4) follow from Lemma 5.1, the law products and the following fact

$$\forall a, b \in \mathbb{N}^d, \quad |a| + |b| \leqslant 3, \quad \|\partial^a_{\vartheta} \partial^b_I v(\vartheta, I)\|_s^{q,\kappa} \lesssim 1 + \|\mathfrak{I}\|_s^{q,\kappa}, \tag{5.45}$$

for all $\vartheta, I \in H^s(\mathbb{T}^d, \mathbb{R}^d), \|\vartheta\|_{s_0}, \|I\|_{s_0} \leq 1$. For more details we can refer to Lemma 5.1 of [16] where we explore a similar situation.

6. Berti-Bolle Approach for the Approximate inverse

In order to apply a modified Nash-Moser scheme, we need to construct an approximate right inverse of the linearized operator at an arbitrary torus i_0 close to the flat one and an arbitrary vector-valued function $c_0 : \mathcal{O} \to \mathbb{R}^d$,

$$d_{(i,\mathsf{c})}\mathscr{F}(i_0,\mathsf{c}_0)[\widehat{\imath},\widehat{\mathsf{c}}] = \omega \cdot \partial_{\varphi}\widehat{\imath} - d_i X_{H_{\varepsilon}^{\mathsf{c}_0}}(i_0(\varphi))[\widehat{\imath}] - (\widehat{\mathsf{c}},0,0), \tag{6.1}$$

where \mathcal{F} is the nonlinear operator defined in (5.21).

We follow the strategy presented by Berti and Bolle in [9], which reduces the search of an approximate inverse of (6.1) to the task of inverting the operator in the normal direction. Their main idea consists in linearizing the functional \mathcal{F} around an isotropic torus sufficiently close to the torus i_0 and make use of a convenient symplectic transformation in such a way the linearized equations become "approximately" decoupled through a trianglular system in the action-angle components and the normal ones.

The main novelty related to this part is to bypass the use of isotropic torus and work directly with the original one i_0 . Although the transformation G_0 introduced in (6.5) is not symplectic and the nonlinear Hamiltonian structure is no longer preserved, the conjugation of the linearized operator via the linear change of variables $DG_0(G_0^{-1}(i_0))$ leads a triangular system with small errors of size $Z \triangleq \mathscr{F}(i_0, \mathbf{c}_0)$, see Proposition 6.1 and Lemma 6.2. Yet we emphasize that the Hamiltonian structure of the original system is of paramount importance, namely in the construction of the transformation G_0 leading to the final triangular system. Therefore, along this section we shall focus on this linear change of variable. The estimates that we shall perform are very similar to those in [4, 9, 16], with minor differences due to the accumulation of different extra errors induced by the isotropic torus as for example in the various Cantor sets emerging during the reducibility scheme. Thus, we often refer to these papers for some technical points in order to avoid here too much details.

We shall first fix some notations and definitions that will be used in this section. Let H be an Hilbert space equipped with the inner product $\langle \cdot, \cdot \rangle_{\mathbb{H}}$. Given a linear operator $A \in \mathscr{L}(\mathbb{R}^d, \mathbb{H})$ we define the transposed operator $A^{\top} : \mathbb{H} \to \mathbb{R}^d$ by the duality relation

$$\langle A^{\top}u, v \rangle_{\mathbb{R}^d} = \langle u, Av \rangle_{\mathbb{H}}, \qquad \forall u \in \mathbb{H}, v \in \mathbb{R}^d.$$
 (6.2)

In accordance with the notation introduced in (5.22) we denote by

$$i_0(\varphi) \triangleq (\vartheta_0(\varphi), I_0(\varphi), z_0(\varphi)), \qquad \Im_0(\varphi) \triangleq i_0(\varphi) - (\varphi, 0, 0).$$
 (6.3)

We shall first assume the following hypothesis: the map $\lambda \mapsto \mathfrak{I}_0(\varphi; \lambda)$ is q-times differentiable with respect to the parameters $\lambda = (\omega, \alpha) \in \mathcal{O} \subset \mathbb{R}^d \times [\underline{\alpha}, \overline{\alpha}]$ and there exists $\varepsilon_0 \in (0, 1)$ such that if

$$\|\mathfrak{I}_0\|_{s_0+2}^{q,\kappa} + \|\mathbf{c}_0 - \omega\|^{q,\kappa} \leqslant \varepsilon_0 \tag{6.4}$$

where ε_0 is assumed to be small enough. We point out that this assumption is enough to provide tame estimates on the tangential part. Later on, a stronger assumption will be required for the normal contribution, see (9.278).

We introduce the diffeomorphism $G_0: (\phi, y, w) \to (\vartheta, I, z)$ of the phase space $\mathbb{T}^d \times \mathbb{R}^d \times \mathbb{H}_{\mathbb{S}_0}^\perp$ defined by

$$\begin{pmatrix} \vartheta \\ I \\ z \end{pmatrix} \triangleq G_0 \begin{pmatrix} \phi \\ y \\ w \end{pmatrix} \triangleq \begin{pmatrix} \vartheta_0(\phi) \\ I_0(\phi) + L_1(\phi)y + L_2(\phi)w \\ z_0(\phi) + w \end{pmatrix}$$
(6.5)

where

$$L_{1}(\phi) \triangleq [\partial_{\varphi}\vartheta_{0}(\varphi)]^{-\top},$$

$$L_{2}(\phi) \triangleq -[(\partial_{\vartheta}\widetilde{z}_{0})(\vartheta_{0}(\phi))]^{\top}\partial_{\theta}^{-1} \quad \text{with} \quad \widetilde{z}_{0}(\vartheta) \triangleq z_{0}(\vartheta_{0}^{-1}(\vartheta)).$$
(6.6)

In the new coordinates, i_0 becomes the trivial embedded torus $(\phi, y, w) = (\varphi, 0, 0)$, namely

$$G_0(\varphi, 0, 0) = i_0(\varphi)$$

In what follows we shall denote by

$$\mathbf{u}_0(\varphi) \triangleq G_0^{-1}(i_0)(\varphi) = (\varphi, 0, 0)$$

the trivial torus and $\mathbf{u} = (\phi, y, w)$ the coordinates induced by G_0 in (6.5). We shall also denote by

$$\widetilde{G}_0(\mathbf{u},\mathbf{c}) \triangleq (G_0(\mathbf{u}),\mathbf{c})$$

the diffeomorphism with the identity on the *c*-component.

We quantify how an embedded torus $i_0(\mathbb{T})$ is approximately invariant for the Hamiltonian vector field $X_{H_{\varepsilon}^{c}}$ in terms of the "error function"

$$Z(\varphi) \triangleq (Z_1, Z_2, Z_3)(\varphi) \triangleq \mathscr{F}(i_0, \mathsf{c}_0)(\varphi) = \omega \cdot \partial_{\varphi} i_0(\varphi) - X_{H^{\mathsf{c}}_{\varepsilon}}(i_0(\varphi), \mathsf{c}_0).$$
(6.7)

6.1. Linear change of variables. The main task in this subsection is to conjugate the linearized operator $d_{i,c}\mathcal{F}(i_0, \mathbf{c}_0)$ in (6.1), via the linear change of variables

$$DG_0(\mathbf{u}_0(\varphi))\begin{pmatrix}\widehat{\phi}\\\widehat{y}\\\widehat{w}\end{pmatrix} = \begin{pmatrix}\partial_{\varphi}\vartheta_0(\varphi) & 0 & 0\\\partial_{\varphi}I_0(\varphi) & L_1(\varphi) & L_2(\varphi)\\\partial_{\varphi}z_0(\varphi) & 0 & I\end{pmatrix}\begin{pmatrix}\widehat{\phi}\\\widehat{y}\\\widehat{w}\end{pmatrix},$$
(6.8)

to a triangular system up to a remainder of order O(Z). As it was shown in [9] this transformation is symplectic up to errors of type Z. Our main results reads as follows.

Proposition 6.1. Under the linear change of variables $DG_0(u_0)$ the linearized operator $d_{i,c}\mathcal{F}(i_0, c_0)$ is transformed into

$$[DG_0(\mathbf{u}_0)]^{-1}d_{(i,\mathsf{c})}\mathcal{F}(i_0,\mathsf{c}_0)D\widetilde{G}_0(\mathbf{u}_0)[\widehat{\phi},\widehat{y},\widehat{w},\widehat{\mathsf{c}}] = \mathbb{D}[\widehat{\phi},\widehat{y},\widehat{w},\widehat{\mathsf{c}}] + \mathbb{E}[\widehat{\phi},\widehat{y},\widehat{w}]$$
(6.9)

where

(i) the operator \mathbb{D} has the triangular form

$$\mathbb{D}[\widehat{\phi}, \widehat{y}, \widehat{w}, \widehat{\mathbf{c}}] \triangleq \begin{pmatrix} \omega \cdot \partial_{\varphi} \widehat{\phi} - \left[K_{20}(\varphi) \widehat{y} + K_{11}^{\top}(\varphi) \widehat{w} + L_{1}^{\top}(\varphi) \widehat{\mathbf{c}} \right] \\ \omega \cdot \partial_{\varphi} \widehat{y} + \mathcal{B}(\varphi) \widehat{\mathbf{c}} \\ \omega \cdot \partial_{\varphi} \widehat{w} - \partial_{\theta} \left[K_{11}(\varphi) \widehat{y} + K_{02}(\varphi) \widehat{w} - L_{2}^{\top}(\varphi) \widehat{\mathbf{c}} \right] \end{pmatrix}$$

 $\mathscr{B}(\varphi)$ and $K_{20}(\varphi)$ are $d \times d$ real matrices,

$$\mathscr{B}(\varphi) \triangleq [\partial_{\varphi}\vartheta_{0}(\varphi)]^{\top}\partial_{\varphi}I_{0}(\varphi)L_{1}^{\top}(\varphi) + [\partial_{\varphi}z_{0}(\varphi)]^{\top}L_{2}^{\top}(\varphi),$$

$$K_{20}(\varphi) \triangleq \varepsilon L_1^\top(\varphi)(\partial_{II}\mathscr{P}_{\varepsilon})(i_0(\varphi))L_1(\varphi)$$

 $K_{02}(\varphi)$ is a linear self-adjoint operator of $\mathbb{H}_{\mathbb{S}_0}^{\perp}$, given by

$$\begin{split} K_{02}(\varphi) &\triangleq (\partial_z \nabla_z H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi)) + \varepsilon L_2^{\perp}(\varphi)(\partial_{II} \mathscr{P}_{\varepsilon})(i_0(\varphi))L_2(\varphi) \\ &+ \varepsilon L_2^{\top}(\varphi)(\partial_{zI} \mathscr{P}_{\varepsilon})(i_0(\varphi)) + \varepsilon (\partial_I \nabla_z \mathscr{P}_{\varepsilon})(i_0(\varphi))L_2(\varphi), \end{split}$$

and $K_{11}(\varphi) \in \mathscr{L}(\mathbb{R}^d, \mathbb{H}_{\mathbb{S}_0}^{\perp}),$

$$K_{11}(\varphi) \triangleq \varepsilon L_2^{\top}(\varphi)(\partial_{II}\mathscr{P}_{\varepsilon})(i_0(\varphi))L_1(\varphi) + \varepsilon(\partial_I \nabla_z \mathscr{P}_{\varepsilon})(i_0(\varphi))L_1(\varphi),$$

(ii) the remainder \mathbb{E} is given by

$$\mathbb{E}[\widehat{\phi}, \widehat{y}, \widehat{w}] \triangleq [DG_0(\mathbf{u}_0)]^{-1} \partial_{\varphi} Z(\varphi) \widehat{\phi} + \begin{pmatrix} 0 \\ \mathscr{A}(\varphi) [K_{20}(\varphi) \widehat{y} + K_{11}^{\top}(\varphi) \widehat{w}] - R_{10}(\varphi) \widehat{y} - R_{01}(\varphi) \widehat{w} \\ 0 \end{pmatrix}$$

where $\mathcal{A}(\varphi)$ and $R_{10}(\varphi)$ are $d \times d$ real matrices,

$$\begin{aligned} \mathscr{A}(\varphi) &\triangleq [\partial_{\varphi}\vartheta_{0}(\varphi)]^{\top}\partial_{\varphi}I_{0}(\varphi) - [\partial_{\varphi}I_{0}(\varphi)]^{\top}\partial_{\varphi}\vartheta_{0}(\varphi) + [\partial_{\varphi}z_{0}(\varphi)]^{\top}\partial_{\theta}^{-1}\partial_{\varphi}z_{0}(\varphi), \\ R_{10}(\varphi) &\triangleq [\partial_{\varphi}Z_{1}(\varphi)]^{\top}L_{1}(\varphi), \\ and \ R_{01}(\varphi) &\in \mathscr{L}(\mathbb{H}_{\mathbb{S}_{0}}^{\perp}, \mathbb{R}^{d}), \\ R_{01}(\varphi) &\triangleq -[\partial_{\varphi}Z_{1}(\varphi)]^{\top}L_{2}(\varphi) + [\partial_{\varphi}Z_{3}(\varphi)]^{\top}\partial_{\theta}^{-1}. \end{aligned}$$

Remark 6.1. A priori, it is not clear from the expression of \mathcal{A} that the error term \mathbb{E} is of order O(Z). This is the purpose of Lemma 6.3 where we quantify the size of \mathcal{A} in terms of the error function Z defined in (6.7).

Proof. Under the map G_0 , the nonlinear operator \mathcal{F} in (5.21) is transformed into

$$\mathscr{F}(G_0(\mathfrak{u}(\varphi)), \mathfrak{c}) = \omega \cdot \partial_{\varphi} \big(G_0(\mathfrak{u}(\varphi)) \big) - X_{H^c_{\varepsilon}} \big(G_0(\mathfrak{u}(\varphi)) \big).$$
(6.10)

Differentiating (6.10) at (u_0, c_0) in the direction (\hat{u}, \hat{c}) gives

$$d_{(\mathbf{u},\mathbf{c})}(\mathscr{F} \circ G_0)(\mathbf{u}_0,\mathbf{c}_0)[(\widehat{\mathbf{u}},\widehat{\mathbf{c}})](\varphi) = \omega \cdot \partial_{\varphi} \left(DG_0(\mathbf{u}_0)\widehat{\mathbf{u}} \right) - \partial_{\phi} \left[X_{H^c_{\varepsilon}} \left(G_0(\mathbf{u}(\varphi)) \right) \right]_{\mathbf{u}=\mathbf{u}_0} \widehat{\phi}$$

$$- \partial_y \left[X_{H^c_{\varepsilon}} \left(G_0(\mathbf{u}(\varphi)) \right) \right]_{\mathbf{u}=\mathbf{u}_0} \widehat{y} - \partial_w \left[X_{H^c_{\varepsilon}} \left(G_0(\mathbf{u}(\varphi)) \right) \right]_{\mathbf{u}=\mathbf{u}_0} \widehat{w} - \begin{pmatrix} \widehat{\mathbf{c}} \\ 0 \\ 0 \end{pmatrix}.$$

$$(6.11)$$

From the expression of $DG_0(u_0)$ in (6.8), we obtain

$$\omega \cdot \partial_{\varphi} \left(DG_{0}(\mathbf{u}_{0})[\widehat{\mathbf{u}}](\varphi) \right) = DG_{0}(\mathbf{u}_{0}) \,\omega \cdot \partial_{\varphi} \widehat{\mathbf{u}} + \partial_{\varphi} \left(\omega \cdot \partial_{\varphi} i_{0} \right) \widehat{\phi} + \begin{pmatrix} 0 \\ (\omega \cdot \partial_{\varphi} L_{1}(\varphi)) \widehat{y} + (\omega \cdot \partial_{\varphi} L_{2}(\varphi)) \widehat{w} \\ 0 \end{pmatrix}.$$
(6.12)

In view of (6.6) and (6.7) we have

$$\omega \cdot \partial_{\varphi} L_{1}(\varphi) = -[\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} (\omega \cdot \partial_{\varphi} [\partial_{\varphi} \vartheta_{0}(\varphi)]^{\top}) [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} = -[\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} ([\partial_{\varphi} Z_{1}(\varphi)]^{\top} + [\partial_{\varphi} ((\partial_{I} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)))]^{\top}) [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top}.$$

$$(6.13)$$

By (6.6) we can easily check that

$$\partial_{\varphi} z_0(\varphi) = (\partial_{\vartheta} \tilde{z}_0)(\vartheta_0(\varphi)) \partial_{\phi} \vartheta_0(\varphi)$$
(6.14)

and thus, we may write the operator $L_2(\varphi)$ in term of the matrix $L_1(\varphi)$,

$$L_2(\varphi) = -[\partial_{\varphi}\vartheta_0(\varphi)]^{-\top} [\partial_{\varphi}z_0(\varphi)]^{\top} \partial_{\theta}^{-1} = -L_1(\varphi)[\partial_{\varphi}z_0(\varphi)]^{\top} \partial_{\theta}^{-1}.$$
(6.15)

Then, by (6.13), (6.15) we have

$$\omega \cdot \partial_{\varphi} L_{2}(\varphi) = [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} (\omega \cdot \partial_{\varphi} [\partial_{\varphi} \vartheta_{0}(\varphi)]^{\top}) [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} [\partial_{\varphi} z_{0}(\varphi)]^{\top} \partial_{\theta}^{-1} - [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} [\partial_{\varphi} (\omega \cdot \partial_{\varphi} z_{0})(\varphi)]^{\top} \partial_{\theta}^{-1}$$

and from (6.7) we get

$$\omega \cdot \partial_{\varphi} L_{2}(\varphi) = -[\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} \Big(\Big[\partial_{\varphi} Z_{1}(\varphi) \Big]^{\top} + \Big[\partial_{\varphi} \Big((\partial_{I} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) \Big) \Big]^{\top} \Big) L_{2}(\varphi) \\ - [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} \Big(\Big[\partial_{\varphi} Z_{3}(\varphi) \Big]^{\top} + \Big[\partial_{\varphi} \Big((\nabla_{z} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) \Big) \Big]^{\top} \Big) \partial_{\theta}^{-1}.$$
(6.16)

Putting together (6.12), (6.13) and (6.16) we conclude that

$$\omega \cdot \partial_{\varphi} \left(DG_{0}(\mathbf{u}_{0})[\widehat{\mathbf{u}}](\varphi) \right) = DG_{0}(\mathbf{u}_{0}) \,\omega \cdot \partial_{\varphi} \widehat{\mathbf{u}} + \partial_{\varphi} \left(\omega \cdot \partial_{\varphi} i_{0} \right) \widehat{\phi}
- \begin{pmatrix} 0 \\ [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} \left[\mathscr{C}_{I}(\varphi) L_{1}(\varphi) + R_{10}(\varphi) \right] \widehat{y} \\ 0 \end{pmatrix}
- \begin{pmatrix} 0 \\ [\partial_{\varphi} \vartheta_{0}(\varphi)]^{-\top} \left[\mathscr{C}_{I}(\varphi) L_{2}(\varphi) + \mathscr{C}_{z}(\varphi) \partial_{\theta}^{-1} + R_{01}(\varphi) \right] \widehat{w} \end{pmatrix}, \quad (6.17)$$

where $R_{10}(\varphi)$ and $R_{01}(\varphi)$ are given by (ii) and

$$\mathscr{C}_{I}(\varphi) \triangleq \left[\partial_{\varphi} \left((\partial_{I} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) \right) \right]^{\top} \\ = \left[\partial_{\varphi} I_{0}(\varphi)\right]^{\top} \left(\partial_{II} H_{\varepsilon}^{\mathbf{c}}\right)(i_{0}(\varphi)) + \left[\partial_{\varphi} \vartheta_{0}(\varphi)\right]^{\top} \left(\partial_{\vartheta I} H_{\varepsilon}^{\mathbf{c}}\right)(i_{0}(\varphi)) \\ + \left[\partial_{\varphi} z_{0}(\varphi)\right]^{\top} \left(\partial_{I} \nabla_{z} H_{\varepsilon}^{\mathbf{c}}\right)(i_{0}(\varphi)),$$

$$(6.18)$$

$$\mathscr{C}_{z}(\varphi) \triangleq \left[\partial_{\varphi} \left((\nabla_{z} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) \right) \right]^{\top} \\ = \left[\partial_{\varphi} I_{0}(\varphi)\right]^{\top} (\partial_{zI} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) + \left[\partial_{\varphi} \vartheta_{0}(\varphi)\right]^{\top} (\partial_{z\vartheta} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)) \\ + \left[\partial_{\varphi} z_{0}(\varphi)\right]^{\top} (\partial_{z} \nabla_{z} H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi)).$$
(6.19)

On the other hand, in view (5.21) and (6.5) we obtain

$$\partial_{\phi} \left[X_{H_{\varepsilon}^{c}} \big(G_{0}(\mathfrak{u}(\varphi)) \big) \right]_{\mathfrak{u}=\mathfrak{u}_{0}} \widehat{\phi} = \partial_{\varphi} \left[X_{H_{\varepsilon}^{c}}(i_{0}(\varphi)) \right] \widehat{\phi}, \tag{6.20}$$

$$\partial_{y} \left[X_{H_{\varepsilon}^{\mathsf{c}}} \big(G_{0}(\mathfrak{u}(\varphi)) \big) \right]_{\mathfrak{u}=\mathfrak{u}_{0}} \widehat{y} = \begin{pmatrix} (\partial_{II} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{1}(\varphi) \widehat{y} \\ -(\partial_{I\vartheta} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{1}(\varphi) \widehat{y} \\ \partial_{\theta} \left[(\partial_{I} \nabla_{z} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{1}(\varphi) \widehat{y} \right] \end{pmatrix},$$
(6.21)

$$\partial_{w} \left[X_{H_{\varepsilon}^{\mathsf{c}}} \big(G_{0}(\mathfrak{u}(\varphi)) \big) \right]_{\mathfrak{u}=\mathfrak{u}_{0}} \widehat{w} = \begin{pmatrix} (\partial_{II} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{2}(\varphi) \widehat{w} + (\partial_{zI} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) \widehat{w} \\ -(\partial_{I\vartheta} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{2}(\varphi) \widehat{w} - (\partial_{z\vartheta} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) \widehat{w} \\ \partial_{\theta} \left[(\partial_{I} \nabla_{z} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) L_{2}(\varphi) \widehat{w} + (\partial_{z} \nabla_{z} H_{\varepsilon}^{\mathsf{c}}) (i_{0}(\varphi)) \widehat{w} \right] \end{pmatrix}.$$
(6.22)

Plugging (6.17), (6.20), (6.21) and (6.22) into (6.11) we find

$$d_{(\mathbf{u},\mathbf{c})}(\mathscr{F} \circ G_{0})(\mathbf{u}_{0},\mathbf{c}_{0})[(\widehat{\mathbf{u}},\widehat{\mathbf{c}})] = DG_{0}(\mathbf{u}_{0})\omega \cdot \partial_{\varphi}\widehat{\mathbf{u}} + \partial_{\varphi}[\mathscr{F}(i_{0}(\varphi))]\widehat{\phi} + \begin{pmatrix} -(\partial_{II}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))L_{1}(\varphi)\widehat{y} - [\partial_{\varphi}\vartheta_{0}(\varphi)]^{-\top}[\mathscr{C}_{I}(\varphi)L_{1}(\varphi) + R_{10}(\varphi)]\widehat{y} \\ -\partial_{\theta}(\partial_{I}\nabla_{z}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))L_{1}(\varphi)\widehat{y} \end{pmatrix} + \begin{pmatrix} -(\partial_{II}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))L_{2}(\varphi)\widehat{w} - (\partial_{zI}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))\widehat{w} \\ [(\partial_{I}\vartheta H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))L_{2}(\varphi) + (\partial_{z}\vartheta H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))]\widehat{w} \\ -\partial_{\theta}[(\partial_{I}\nabla_{z}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))L_{2}(\varphi)\widehat{w} + \partial_{\theta}(\partial_{z}\nabla_{z}H_{\varepsilon}^{\mathbf{c}})(i_{0}(\varphi))\widehat{w}] \end{pmatrix} - \begin{pmatrix} \widehat{\mathbf{c}} \\ 0 \\ 0 \end{pmatrix}. \qquad (6.23)$$

According to (6.8) and using the identities (6.14) and (6.15), the inverse of the linear operator $DG_0(\mathbf{u}_0)$ is given by

$$[DG_0(\mathbf{u}_0)]^{-1} = \begin{pmatrix} [\partial_{\varphi}\vartheta_0(\varphi)]^{-1} & 0 & 0\\ -\mathscr{B}(\varphi) & [\partial_{\varphi}\vartheta_0(\varphi)]^\top & [\partial_{\varphi}z_0(\varphi)]^\top \partial_{\theta}^{-1}\\ (\partial_{\vartheta}\widetilde{z}_0)(\vartheta_0(\varphi)) & 0 & I \end{pmatrix}$$
(6.24)

where $\mathscr{B}(\varphi)$ is given by (i). Applying $[DG_0(\mathfrak{u}_0)]^{-1}$ to (6.23) and using the identities (6.18), (6.19) and the fact that

$$\mathscr{B}(\varphi) = \mathscr{A}(\varphi)[\partial_{\varphi}\vartheta_{0}(\varphi)]^{-1} + [\partial_{\varphi}I_{0}(\varphi)]^{\top}, \qquad (6.25)$$

where $\mathscr{A}(\varphi)$ is defined in (ii), we obtain

$$\begin{split} [DG_{0}(\mathbf{u}_{0})]^{-1}d_{(\mathbf{u},\mathbf{c})}(\mathscr{F}\circ G_{0})(\mathbf{u}_{0},\mathbf{c}_{0})[\widehat{\mathbf{u}},\widehat{\mathbf{c}}] &= \omega \cdot \partial_{\varphi}\widehat{\mathbf{u}} + [DG_{0}(\mathbf{u}_{0})]^{-1}\partial_{\varphi}\big[\mathscr{F}(i_{0}(\varphi))\big]\widehat{\phi} \\ &+ \left(\begin{array}{c} -K_{20}(\varphi)\widehat{y} \\ \mathscr{A}(\varphi)K_{20}(\varphi)\widehat{y} - R_{10}(\varphi)\widehat{y} \\ -\partial_{\theta}K_{11}(\varphi)\widehat{y} \end{array}\right) + \left(\begin{array}{c} -K_{11}^{\top}(\varphi)\widehat{w} \\ \mathscr{A}(\varphi)K_{11}^{\top}(\varphi)\widehat{w} - R_{01}(\varphi)\widehat{w} \\ -\partial_{\theta}K_{02}(\varphi)\widehat{w} \end{array}\right) + \left(\begin{array}{c} -L_{1}^{\top}(\phi)\widehat{\mathbf{c}} \\ \mathscr{B}(\varphi)\widehat{\mathbf{c}} \\ \partial_{\theta}L_{2}^{\top}(\varphi)\widehat{\mathbf{c}} \end{array}\right), \end{split}$$

with

$$\begin{split} K_{20}(\varphi) &\triangleq L_1^{\top}(\varphi)(\partial_{II}H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi))L_1(\varphi) \,, \\ K_{11}(\varphi) &\triangleq L_2^{\top}(\varphi)(\partial_{II}H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi))L_1(\varphi) + (\partial_{I}\nabla_z H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi))L_1(\varphi) \,, \\ K_{02}(\varphi) &\triangleq (\partial_z \nabla_z H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi)) + L_2^{\top}(\varphi)(\partial_{II}H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi))L_2(\varphi) + L_2^{\top}(\varphi)(\partial_{zI}H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi)) \\ &+ (\partial_I \nabla_z H_{\varepsilon}^{\mathsf{c}})(i_0(\varphi))L_2(\varphi). \end{split}$$

Finally, by (5.20) we conclude the desired identity and this ends the proof of Proposition 6.1.

The next aim is to estimate the induced composition operators $[DG_0(\mathbf{u}_0)]^{\pm 1}$, given by (6.8), (6.24) and the coefficients R_{10} , R_{01} , K_{20} , K_{11} , K_{11}^{\top} and \mathscr{A} defined in Proposition 6.1. More precisely, arguing as in Lemmata 5.6-5.7 in [16], by using law product of Lemma 4.1-(ii) and the smallness condition (6.4), we obtain the following result.

Lemma 6.1. Let (q, d, κ, s, s_0) as in (4.1), then the following assertions hold true.

(i) The operator $DG_0(\mathbf{u}_0)$ and $[DG_0(\mathbf{u}_0)]^{-1}$ satisfy for all $\widehat{\mathbf{u}} = (\widehat{\phi}, \widehat{y}, \widehat{w})$,

$$\|[DG_0(\mathbf{u}_0)]^{\pm 1}[\widehat{\mathbf{u}}]\|_s^{q,\kappa} \lesssim \|\widehat{\mathbf{u}}\|_s^{q,\kappa} + \|\mathfrak{I}_0\|_{s+1}^{q,\kappa} \|\widehat{\mathbf{u}}\|_{s_0}^{q,\kappa}.$$

(ii) The operators R_{10} and R_{01} satisfy the estimates

$$\|R_{10}\widehat{y}\|_{s}^{q,\overline{\gamma}} \lesssim_{s} \|Z\|_{s+1}^{q,\kappa} + \|Z\|_{s_{0}+1}^{q,\kappa} (\|\Im_{0}\|_{s+1}^{q,\kappa}\|\widehat{y}\|_{s_{0}}^{q,\kappa} + \|\widehat{y}\|_{s}^{q,\kappa}), \\ \|R_{01}\widehat{w}\|_{s}^{q,\overline{\gamma}} \lesssim_{s} \|Z\|_{s+1}^{q,\kappa} \|\widehat{w}\|_{s_{0}}^{q,\kappa} + \|Z\|_{s_{0}+1}^{q,\kappa} (\|\Im_{0}\|_{s+1}^{q,\kappa}\|\widehat{w}\|_{s_{0}}^{q,\kappa} + \|\widehat{w}\|_{s}^{q,\kappa}).$$

(iii) The matrix K_{20} and the operators K_{11}, K_{11}^{\top} satisfy the estimates

$$\begin{aligned} \|K_{20}\|_{s}^{q,\overline{\gamma}} &\lesssim_{s} \varepsilon \left(1 + \|\mathfrak{I}_{0}\|_{s+2}^{q,\gamma}\right), \\ \|K_{11}\widehat{y}\|_{s}^{q,\overline{\gamma}} &\lesssim_{s} \varepsilon \left(\|\widehat{y}\|_{s+2}^{q,\overline{\gamma}} + \|\mathfrak{I}_{0}\|_{s+2}^{q,\overline{\gamma}}\|\widehat{y}\|_{s_{0}+2}^{q,\overline{\gamma}}\right), \\ \|K_{11}^{\top}\widehat{w}\|_{s}^{q,\overline{\gamma}} &\lesssim_{s} \varepsilon \left(\|\widehat{w}\|_{s+2}^{q,\overline{\gamma}} + \|\mathfrak{I}_{0}\|_{s+2}^{q,\overline{\gamma}}\|\widehat{w}\|_{s_{0}+2}^{q,\overline{\gamma}}\right). \end{aligned}$$

(iv) The matrices \mathcal{A} and \mathcal{B} satisfy the estimates

$$\|\mathscr{A}\|_{s}^{q,\overline{\gamma}}+\|\mathscr{B}\|_{s}^{q,\overline{\gamma}}\lesssim_{s}\|\mathfrak{I}_{0}\|_{s+1}^{q,\kappa}.$$

6.2. Defect of the symplectic structure. In this subsection we shall prove that the matrix \mathcal{A} , defined in Proposition 6.1-(ii), is zero at an exact solution on some Cantor like set, up to an exponentially small remainder. Recall that the coefficients of the matrix \mathcal{A} are given by

$$\mathscr{A}_{kj}(\varphi) = \partial_{\varphi_k} I_0(\varphi) \cdot \partial_{\varphi_j} \vartheta_0(\varphi) - \partial_{\varphi_k} \vartheta_0(\varphi) \cdot \partial_{\varphi_j} I_0(\varphi) + \left(\partial_{\theta}^{-1} \partial_{\varphi_k} z_0(\varphi), \partial_{\varphi_j} z_0(\varphi)\right)_{L^2}.$$
(6.26)

We point out that those coefficients coincide with the "lack of isotropy coefficients" introduced in Lemma 5 of [9] where the following identitity is proved

$$\omega \cdot \partial_{\varphi} \mathcal{A}_{jk}(\varphi) = \mathcal{W} \left(\partial_{\varphi} Z(\varphi) \underline{e}_{k}, \partial_{\varphi} i_{0}(\varphi) \underline{e}_{j} \right) + \mathcal{W} \left(\partial_{\varphi} i_{0}(\varphi) \underline{e}_{k}, \partial_{\varphi} Z(\varphi) \underline{e}_{j} \right)$$

$$= \partial_{\varphi_{k}} Z_{2}(\varphi) \cdot \partial_{\varphi_{j}} \vartheta_{0}(\varphi) - \partial_{\varphi_{k}} Z_{1}(\varphi) \cdot \partial_{\varphi_{j}} I_{0}(\varphi) + \left(\partial_{\theta}^{-1} \partial_{\varphi_{k}} Z_{3}(\varphi), \partial_{\varphi_{j}} z_{0}(\varphi) \right)_{L^{2}}$$

$$+ \partial_{\varphi_{k}} I_{0}(\varphi) \cdot \partial_{\varphi_{j}} Z_{2}(\varphi) - \partial_{\varphi_{k}} \vartheta_{0}(\varphi) \cdot \partial_{\varphi_{j}} Z_{1}(\varphi) + \left(\partial_{\theta}^{-1} \partial_{\varphi_{k}} z_{0}(\varphi), \partial_{\varphi_{j}} Z_{3}(\varphi) \right)_{L^{2}}.$$

$$(6.27)$$

Here $(\underline{e}_1, \ldots, \underline{e}_d)$ denotes the canonical basis of \mathbb{R}^d . In order to solve (6.27) we need to discuss some elementary results on the invertibility of Fourier multiplier operator in the presence of a small divisor problem. Given $\kappa \in (0, 1]$ and $\tau_1 > 0$, we introduce the Cantor set

$$\mathsf{DC}(\kappa,\tau_1) \triangleq \left\{ \omega \in \mathbb{R}^d : |\omega \cdot l| \ge \frac{\kappa}{\langle l \rangle^{\tau_1}}, \quad \forall \, l \in \mathbb{Z}^d \setminus \{0\} \right\}.$$
(6.28)

If $\lambda \in DC(\kappa, \tau_1)$ then, for any smooth function $h : \mathcal{O} \times \mathbb{T}^d \to \mathbb{R}$ with zero average, the equation $\omega \cdot \partial_{\varphi} u = h$ has a periodic solution $h : \mathbb{T}^d \to \mathbb{R}$ given by

$$u(\lambda,\varphi) = -\mathrm{i} \sum_{l \in \mathbb{Z}^d \setminus \{0\}} \frac{h_l(\lambda)}{\omega \cdot l} \mathbf{e}_l(\varphi) \quad \text{where} \quad h = \sum_{l \in \mathbb{Z}^d \setminus \{0\}} h_l(\lambda) \mathbf{e}_l, \quad \mathbf{e}_l(\varphi) \triangleq e^{\mathrm{i} l \cdot \varphi}.$$

For all $\omega \in \mathcal{O}$ we define the smooth extension of u by

$$(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1}h \triangleq -i \sum_{l \in \mathbb{Z}^d \setminus \{0\}} \frac{\chi((\omega \cdot l)\kappa^{-1} \langle l \rangle^{\tau_1})h_l(\lambda)}{\omega \cdot l} \mathbf{e}_l,$$
(6.29)

where $\chi \in \mathscr{C}^{\infty}(\mathbb{R}, \mathbb{R})$ is an even positive cut-off function such that

$$\chi(\xi) = \begin{cases} 0 & \text{if } |\xi| \leq \frac{1}{3}, \\ 1 & \text{if } |\xi| \geq \frac{1}{2}. \end{cases}$$
(6.30)

Notice that this operator is well-defined in the whole set of parameters \mathcal{O} and coincides with the formal inverse of $(\omega \cdot \partial_{\varphi})^{-1}$ when the frequency ω belongs to $DC(\kappa, \tau_1)$. On the other hand, we note that at each iterative step of the Nash-Moser iteration -and correspondingly for the reduction of the linearized operator in Section 9- we only require that the frequency vector $\omega \in \mathbb{R}^d$ satisfies finitely many non-resonance diophantine conditions. More precisely we assume at the *n*-th step that ω belongs to the truncated Cantor set

$$\mathsf{DC}_{N_n}(\kappa,\tau_1) \triangleq \left\{ \omega \in \mathbb{R}^d; \ |\omega \cdot l| \ge \frac{\kappa}{\langle l \rangle^{\tau_1}}, \ \forall |l| \le N_n \right\}$$
(6.31)

where for any $n \in \mathbb{N} \cup \{-1\}$ the sequence (N_n) is defined by

$$N_{-1} \triangleq 1, \quad \forall n \in \mathbb{N}, \quad N_n \triangleq N_0^{\left(\frac{3}{2}\right)^n} \quad \text{with} \quad N_0 \ge 2.$$
 (6.32)

We shall make use of the following result.

Lemma 6.2. Let $\kappa \in (0, 1], q \in \mathbb{N}$. Then the following holds true.

(i) For any $s \ge q$ we have

$$\left\| (\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} h \right\|_{s}^{q,\kappa} \leqslant C \kappa^{-1} \|h\|_{s+\tau_{1}(q+1)}^{q,\kappa}.$$

(ii) For any $n \in \mathbb{N}$ and any $\omega \in DC_{N_n}(\kappa, \tau_1)$ we have

$$(\omega \cdot \partial_{\varphi})(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} = \Pi_{N_n},$$

where Π_{N_n} is the orthogonal projection defined by

$$\prod_{N_n} \sum_{l \in \mathbb{Z}^d} h_l \mathbf{e}_l = \sum_{\substack{l \in \mathbb{Z}^d \setminus \{0\} \\ |l| \leqslant N_n}} h_l \mathbf{e}_l$$

Proof. (i) The proof of the first point can be done using Faà di Bruno's formula in a similar way to [4, Lemma 2.5].

(ii) By construction, one has for $\omega \in DC_{N_n}(\kappa, \tau_1)$ and $|l| \leq N_n$,

$$\chi((\omega \cdot l)\kappa^{-1}\langle l \rangle^{\tau_1}) = 1,$$

Thus, according to the explicit extension (6.29),

$$(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} h = -\mathbf{i} \sum_{\substack{l \in \mathbb{Z}^d \setminus \{0\} \\ |l| \leqslant N_n}} \frac{\chi((\omega \cdot l)\kappa^{-1} \langle l \rangle^{\tau_1}) h_l(\lambda)}{\omega \cdot l} \mathbf{e}_l$$
$$= -\mathbf{i} \sum_{\substack{l \in \mathbb{Z}^d \setminus \{0\} \\ |l| \leqslant N_n}} \frac{h_l(\lambda)}{\omega \cdot l} \mathbf{e}_l.$$
(6.33)

Therefore, we obtain

$$(\omega \cdot \partial_{\varphi})(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} h = \sum_{\substack{l \in \mathbb{Z}^d \setminus \{0\}\\|l| \leq N_n}} h_l(\lambda) \mathbf{e}_l$$
$$= \Pi_{N_n} h.$$

This concludes the proof of the lemma.

Now, we are in a position to solve (6.27). The following lemma is proved in [16, Lemma 5.3], but for the sake of completeness we will give some key ingredients of the proof.

Lemma 6.3. The coefficients \mathcal{A}_{jk} , defined in (6.26), admit the decomposition

$$\mathscr{A}_{kj} = \mathscr{A}_{kj}^{(n)} + \mathscr{A}_{kj}^{(n),\perp} \quad with \qquad \mathscr{A}_{kj}^{(n)} \triangleq \Pi_{N_n} \mathscr{A}_{kj} \qquad and \qquad \mathscr{A}_{kj}^{(n),\perp} \triangleq \Pi_{N_n}^{\perp} \mathscr{A}_{kj}, \tag{6.34}$$

where

(i) The function $\mathscr{A}_{kj}^{(n),\perp}$ satisfies for any $s \in \mathbb{R}$,

$$\|\mathscr{A}_{kj}^{(n),\perp}\|_{s}^{q,\kappa} \lesssim_{s,b} N_{n}^{-b} \|\mathfrak{I}_{0}\|_{s+1+b}^{q,\kappa}, \quad \forall b \ge 0.$$

(ii) There exist functions $\mathcal{A}_{kj}^{(n),\text{ext}}$ defined for any $\lambda = (\omega, \alpha) \in \mathcal{O}$, q-times differentiable with respect to λ and satisfying, for any $s \ge s_0$, the estimate

$$\|\mathscr{A}_{kj}^{(n),\text{ext}}\|_{s}^{q,\kappa} \lesssim_{s} \kappa^{-1} \left(\|Z\|_{s+\tau_{1}(q+1)+1}^{q,\kappa} + \|Z\|_{s_{0}+1}^{q,\kappa} \|\mathfrak{I}_{0}\|_{s+\tau_{1}(q+1)+1}^{q,\kappa} \right).$$

Moreover,
$$\mathscr{A}_{kj}^{(n),\text{ext}}$$
 coincides with $\mathscr{A}_{kj}^{(n)}$ on the Cantor set $\text{DC}_{N_n}(\kappa, \tau_1)$.

Proof. The point (i) follows immediately from (6.26), (6.34) and Lemma 4.2. (ii) Applying the projector to the identity (6.27) we obtain

$$\omega \cdot \partial_{\varphi} \mathcal{A}_{jk}^{(n)}(\varphi) = \prod_{N_n} \left[\mathcal{W} \left(\partial_{\varphi} Z(\varphi) \underline{e}_k, \partial_{\varphi} i_0(\varphi) \underline{e}_j \right) + \mathcal{W} \left(\partial_{\varphi} i_0(\varphi) \underline{e}_k, \partial_{\varphi} Z(\varphi) \underline{e}_j \right) \right].$$

Then, by (6.4), Lemma 4.2 and Lemma 4.1 we get

$$\left\| \Pi_{N_n} \left[\mathscr{W} \left(\partial_{\varphi} Z(\varphi) \underline{e}_k, \partial_{\varphi} i_0(\varphi) \underline{e}_j \right) + \mathscr{W} \left(\partial_{\varphi} i_0(\varphi) \underline{e}_k, \partial_{\varphi} Z(\varphi) \underline{e}_j \right) \right] \right\|_s^{q,\kappa} \lesssim_s \left(\|Z\|_{s+1}^{q,\kappa} + \|Z\|_{s_0+1}^{q,\kappa} \|\mathfrak{I}_0\|_{s+1}^{q,\kappa} \right).$$
We define the function $\mathscr{A}_{kj}^{(n),\text{ext}}$ as

$$\mathscr{A}_{kj}^{(n),\text{ext}}(\varphi) \triangleq (\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \prod_{N_n} \left[\mathscr{W} \left(\partial_{\varphi} Z(\varphi) \underline{e}_k, \partial_{\varphi} i_0(\varphi) \underline{e}_j \right) + \mathscr{W} \left(\partial_{\varphi} i_0(\varphi) \underline{e}_k, \partial_{\varphi} Z(\varphi) \underline{e}_j \right) \right].$$
Applying Lemma 6.2 concludes the proof of the Lemma.

6.3. Construction of an approximate inverse. Since the error term \mathbb{E} is zero at an exact solution, up to an exponentially small remainder (see Proposition 6.1-(ii) and Lemma 6.3) on the Cantor set $\mathsf{DC}_{N_n}(\kappa, \tau_1)$, then in order to find an approximate inverse of the linear operator in (6.9) (and so of $d_{(i,c)}\mathscr{F}(i_0, c_0)$) it is sufficient to almost invert the operator \mathbb{D} . The linear system $\mathbb{D}[\hat{u}] = (g_1, g_2, g_3)$ may be solved in a triangular way, first inverting the action-component equation, which is decoupled from the other equations,

$$\omega \cdot \partial_{\varphi} \widehat{y} + \mathscr{B}(\varphi) \widehat{\mathsf{c}} = g_2.$$

As we require only the finitely many non-resonance conditions (6.31), for any $\omega \in \mathbb{R}^d$ we also decompose $\omega \cdot \partial_{\varphi}$ as

$$\omega \cdot \partial_{\varphi} = \mathfrak{D}_{(n)} + \mathfrak{D}_{(n)}^{\perp},$$

$$\mathfrak{D}_{(n)} \triangleq \omega \cdot \partial_{\varphi} \Pi_{N_n} + \Pi_{N_n, \mathbf{g}}^{\perp} \qquad \mathfrak{D}_{(n)}^{\perp} \triangleq \omega \cdot \partial_{\varphi} \Pi_{N_n}^{\perp} - \Pi_{N_n, \mathbf{g}}^{\perp}$$
(6.35)

where

$$\Pi_{N_n,\mathbf{g}}^{\perp} \sum_{l \in \mathbb{Z}^d \setminus \{0\}} h_l \mathbf{e}_l \triangleq \sum_{\substack{l \in \mathbb{Z}^d \setminus \{0\}\\|l| > N_n}} g(l) h_l \mathbf{e}_l$$

and the function $\mathbf{g} : \mathbb{Z}^d \setminus \{0\} \to \{-1, 1\}$ is defined, for all $l = (l_1, \dots, l_d) \in \mathbb{Z}^d \setminus \{0\}$, as the sign of the first non-zero component in the vector l. Thus, it satisfies

$$g(-l) = -g(l) \quad \forall l \in \mathbb{Z}^d \setminus \{0\}.$$

Notice that for the high frequency we use the projector $\Pi_{N_n,g}^{\perp}$ instead of $\Pi_{N_n}^{\perp}$ in order to preserve the reversibility. We shall need the following Lemma.

Lemma 6.4. The following holds true.

(i) The operator $\mathfrak{D}_{(n)}^{\perp}$ satisfies

$$\forall b \ge 0, \, s \in \mathbb{R}, \quad \|\mathcal{D}_{(n)}^{\perp}h\|_s^{q,\kappa} \lesssim N_n^{-b} \|h\|_{s+b+1}^{q,\kappa}.$$

(ii) There exists a family of linear operators $([\mathfrak{D}_{(n)}]^{-1}_{ext})_n$ satisfying, for any $g \in H^s_0(\mathbb{T}^d)$,

$$\forall s \ge s_0, \quad \sup_{n \in \mathbb{N}} \| [\mathcal{D}_{(n)}]_{\text{ext}}^{-1} g \|_s^{q,\kappa} \lesssim \kappa^{-1} \| g \|_{s+q(\tau_1+1)}^{q,\kappa}.$$

Moreover, for all $\omega \in DC_{N_n}(\kappa, \tau_1)$ one has the identity

$$\mathcal{D}_{(n)}[\mathcal{D}_{(n)}]_{\text{ext}}^{-1} = \text{Id}.$$

Proof. The point (i) follows immediately from Lemma 4.2. (ii) We define the operator $[\mathcal{D}_{(n)}]_{\text{ext}}^{-1}$ as

$$[\mathcal{D}_{(n)}]_{\text{ext}}^{-1} \triangleq (\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} + \Pi_{N_n, \frac{1}{g}}^{\perp}.$$
 (6.36)

From (6.33), (6.35) and (6.36) we get, for all $\omega \in DC_{N_n}(\kappa, \tau_1)$,

$$\mathcal{D}_{(n)}[\mathcal{D}_{(n)}]_{\text{ext}}^{-1} = \omega \cdot \partial_{\varphi} \Pi_{N_n} \left[(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} + \Pi_{N_n, \frac{1}{g}}^{\perp} \right] + \Pi_{N_n, g}^{\perp} \left[(\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} + \Pi_{N_n, \frac{1}{g}}^{\perp} \right] \\ = \omega \cdot \partial_{\varphi} (\omega \cdot \partial_{\varphi})_{\text{ext}}^{-1} \Pi_{N_n} + \Pi_{N_n}^{\perp}.$$

Applying Lemma 6.2-(ii) we conclude that

$$\mathcal{D}_{(n)}[\mathcal{D}_{(n)}]_{\text{ext}}^{-1} = \Pi_{N_n} + \Pi_{N_n}^{\perp} = \text{Id}$$

The estimate on $[\mathcal{D}_{(n)}]_{\text{ext}}^{-1}$ follows from (6.36), Lemma 4.2 and Lemma 6.2.

The second step in the resolution of the Linear equation $\mathbb{D}[\hat{u}] = g$, where \mathbb{D} is given by Proposition 6.1-(i), is to solve the last normal-component equation

$$\omega \cdot \partial_{\varphi} \widehat{w} - \partial_{\theta} K_{02}(\varphi) \widehat{w} = g_3 + \partial_{\theta} \left[K_{11}(\varphi) \widehat{y} - L_2^{\dagger}(\varphi) \widehat{\mathsf{c}} \right]$$
57

For this aim we need to find an approximate right inverse of the linearized operator in the normal direction

$$\widehat{\mathscr{L}}_{\omega} \triangleq \Pi_{\mathbb{S}_0}^{\perp} \left(\omega \cdot \partial_{\varphi} - \partial_{\theta} K_{02}(\varphi) \right) \Pi_{\mathbb{S}_0}^{\perp}$$
(6.37)

when the set of parameters is restricted to a Cantor like set. Here the projector $\Pi_{\mathbb{S}_0}^{\perp}$ is the one defined in (5.9). For the sake of clarity we shall give a brief statement about the invertibility in the normal direction; for a precise statement with a detailed description of Cantor like sets see Theorem 9.1.

Proposition 6.2. Let $(\kappa, q, d, \tau_1, \tau_2, s_0)$ satisfy (9.2), (μ_2, s_h) satisfy (9.277) and assume the smallness condition (9.278). Then there exist $\sigma_5 = \sigma_5(\tau_1, \tau_2, d, q) > 0$, a family of linear operator $(T_{\omega,n})_n$ satisfying

$$\forall s \in [s_0, S], \quad \sup_{n \in \mathbb{N}} \|\mathsf{T}_{\omega, n}h\|_s^{q, \kappa} \lesssim \kappa^{-1} \left(\|h\|_{s+\sigma_5}^{q, \kappa} + \|\mathfrak{I}_0\|_{s+\sigma_5}^{q, \kappa} \|h\|_{s_0+\sigma_5}^{q, \kappa} \right)$$
(6.38)

and a Cantor set $G_n = G_n(\kappa, \tau_1, \tau_2, i_0) \subset DC_{N_n}(\kappa, \tau_1) \times (\underline{\alpha}, \overline{\alpha})$ in which we have the decomposition

$$\widehat{\mathcal{L}}_{\omega} = \widehat{\mathbf{L}}_{\omega,n} + \widehat{\mathbf{R}}_n,$$

with

$$\widehat{\mathsf{L}}_{\omega,n}\mathsf{T}_{\omega,n} = \mathrm{Id} \tag{6.39}$$

where the operators $\widehat{L}_{\omega,n}$ and \widehat{R}_n are defined in the whole set \mathcal{O} with the estimates

$$\begin{aligned} \forall s \in [s_0, S], \quad \|\widehat{\mathbf{L}}_{\omega,n}h\|_{s_0}^{q,\kappa} \lesssim \|h\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{q,s+\sigma_5}^{q,\kappa} \|h\|_{s_0+1}^{q,\kappa}, \\ \forall b \in [0, S], \quad \|\widehat{\mathbf{R}}_nh\|_{s_0}^{q,\kappa} \lesssim N_n^{-b} \kappa^{-1} \Big(\|h\|_{s_0+b+\sigma_5}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{s_0+b+\sigma_5}^{q,\kappa} \|h\|_{s_0+\sigma_5}^{q,\kappa} \Big) \\ &+ \varepsilon \kappa^{-4} N_0^{\mu_2} N_n^{-\mu_2} \|h\|_{s_0+\sigma_5}^{q,\kappa}. \end{aligned}$$

According to Proposition 6.1, the identities (6.34)-(6.35) and Proposition 6.2 we may decompose the operator $[DG_0(\mathbf{u}_0)]^{-1}d_{(i,\mathbf{c})}\mathcal{F}(i_0,\mathbf{c}_0)D\widetilde{G}_0(\mathbf{u}_0)$ as follows

$$[DG_0(\mathbf{u}_0)]^{-1}d_{(i,\mathbf{c})}\mathcal{F}(i_0,\mathbf{c}_0)D\widetilde{G}_0(\mathbf{u}_0) = \mathbb{D}_n + \mathbb{E}_n + \mathscr{P}_n + \mathscr{Q}_n$$
(6.40)

with

$$\mathbb{D}_{n}[\widehat{\phi},\widehat{y},\widehat{w},\widehat{c}] \triangleq \begin{pmatrix} \mathfrak{D}_{(n)}\widehat{\phi} - K_{20}\widehat{y} - K_{11}^{\top}\widehat{w} - L_{1}^{\top}\widehat{c} \\ \mathfrak{D}_{(n)}\widehat{y} + \mathfrak{B}\widehat{c} \\ \widehat{L}_{\omega,n}\widehat{w} - \partial_{\theta}[K_{11}\widehat{y} - L_{2}^{\top}\widehat{c}] \end{pmatrix},$$
(6.41)

$$\mathbb{E}_{n}[\widehat{\phi},\widehat{y},\widehat{w}] \triangleq [DG_{0}(\mathfrak{u}_{0})]^{-1}[\partial_{\varphi}Z]\widehat{\phi} + \left(\begin{array}{c}0\\\mathscr{A}^{(n)}\left[K_{20}\widehat{y} + K_{11}^{\top}\widehat{w}\right] - R_{10}\widehat{y} - R_{01}\widehat{w}\\0\end{array}\right)$$
(6.42)

and

$$\mathscr{P}_{n}[\widehat{\phi},\widehat{y},\widehat{w}] \triangleq \begin{pmatrix} \mathscr{D}_{(n)}^{\perp}\widehat{\phi} \\ \mathscr{D}_{(n)}^{\perp}\widehat{y} + \mathscr{A}^{(n),\perp}[K_{20}\widehat{y} + K_{11}^{\top}\widehat{w}] \\ 0 \end{pmatrix}, \quad \mathscr{Q}_{n}[\widehat{\phi},\widehat{y},\widehat{w}] \triangleq \begin{pmatrix} 0 \\ 0 \\ \widehat{\mathsf{R}}_{n}[\widehat{w}] \end{pmatrix}, \quad (6.43)$$

where $\mathscr{A}^{(n)}$ and $\mathscr{A}^{(n),\perp}$ are the matrices with coefficients $\mathscr{A}_{kj}^{(n)}$ and $\mathscr{A}_{kj}^{(n),\perp}$ respectively, see (6.34). Because the coefficients $\mathscr{A}_{jk}^{(n)}$ do not vanish at exact solutions on whole set of parameters \mathscr{O} and the $\mathscr{A}_{kj}^{(n),\text{ext}}$ do (see Lemma 6.3), we shall replace the operator \mathbb{E}_n by the extension

$$\mathbb{E}_{n}^{\mathrm{ext}}[\widehat{\phi},\widehat{y},\widehat{w}] \triangleq [DG_{0}(\mathbf{u}_{0})]^{-1}[\partial_{\varphi}Z]\widehat{\phi} + \left(\begin{array}{c}0\\\mathscr{A}^{(n),\mathrm{ext}}\left[K_{20}(\varphi)\widehat{y} + K_{11}^{\top}\widehat{w}\right] - R_{10}\widehat{y} - R_{01}\widehat{w}\\0\end{array}\right), \quad (6.44)$$

with $\mathscr{A}^{(n),\text{ext}}$ is the matrix with coefficients $\mathscr{A}_{kj}^{(n),\text{ext}}$. Thus, we define the linear operator \mathbb{L}_{ext} as

$$\mathbb{L}_{\text{ext}} = \mathbb{D}_n + \mathbb{E}_n^{\text{ext}} + \mathscr{P}_n + \mathscr{Q}_n.$$
(6.45)

The operator \mathbb{L}_{ext} is defined on the whole set \mathcal{O} and, by construction, coincides with the linear operator in (6.40) on the Cantor set G_n ,

$$\forall \lambda \in \mathbf{G}_n, \quad \mathbb{L}_{\text{ext}} = [DG_0(\mathbf{u}_0)]^{-1} d_{(\mathbf{u},\mathbf{c})} \mathscr{F}(i_0,\mathbf{c}_0) D\tilde{G}_0(\mathbf{u}_0). \tag{6.46}$$

Now, we shall estimate the error terms $\mathbb{E}_n^{\text{ext}}$, \mathscr{P}_n , \mathscr{Q}_n and find an exact inverse of the principal term \mathbb{D}_n . This will be the subject of the following proposition.

Proposition 6.3. Let $(\kappa, q, d, \tau_1, s_0)$ satisfy (9.2), (μ_2, s_h) satisfy (9.277) and assume the smallness condition (9.278). Then the following assertions hold true.

(i) The operator $\mathbb{E}_n^{\text{ext}}$ satisfies the estimate

$$\|\mathbb{E}_{n}^{\text{ext}}[\widehat{\mathbf{u}}]\|_{s_{0}}^{q,\kappa} \lesssim_{s} \|Z\|_{s_{0}+1}^{q,\kappa} \|\widehat{\mathbf{u}}\|_{s_{0}+1}^{q,\kappa}.$$

(ii) For all $b \ge 0$ the operator \mathscr{P}_n satisfies the estimate

$$|\mathscr{P}_{n}[\widehat{\mathfrak{u}}]\|_{s_{0}}^{q,\kappa} \lesssim N_{n}^{-b} \left(\|\widehat{\mathfrak{u}}\|_{s_{0}+2+b}^{q,\kappa} + \|\mathfrak{I}_{0}\|_{s_{0}+2+b}^{q,\kappa}\|\widehat{\mathfrak{u}}\|_{s_{0}+2}^{q,\kappa}\right).$$

(iii) For all $b \in [0, S]$, the operator \mathcal{Q}_n satisfies the estimate

$$\|\mathscr{Q}_{n}[\widehat{\mathbf{u}}]\|_{s_{0}}^{q,\kappa} \lesssim N_{n}^{-b}\kappa^{-1}\Big(\|\widehat{\mathbf{u}}\|_{s_{0}+b+\sigma_{5}}^{q,\kappa} + \varepsilon\kappa^{-3}\|\mathfrak{I}_{0}\|_{s_{0}+b+\sigma_{5}}^{q,\kappa}\|\widehat{\mathbf{u}}\|_{s_{0}+\sigma_{5}}^{q,\kappa}\Big) + \varepsilon\kappa^{-4}N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}}\|\widehat{\mathbf{u}}\|_{s_{0}+\sigma_{5}}^{q,\kappa}.$$

(iv) There exist $\sigma_6 = \sigma_6(\tau_1, \tau_2, d, q) > 0$ and a family of operators $([\mathbb{D}_n]_{\text{ext}}^{-1})_n$ such that for all $g \triangleq (g_1, g_2, g_3)$ satisfying the symmetry

$$g_1(\varphi) = g_1(-\varphi), \quad g_2(\varphi) = -g_2(-\varphi), \quad g_3(\varphi) = -(\mathscr{G}g_3)(\varphi),$$

the function $[\mathbb{D}_n]_{\text{ext}}^{-1}g$ satisfies the estimate, for any $s_0 \leqslant s \leqslant S$,

$$[\mathbb{D}_n]_{\text{ext}}^{-1}g\|_s^{q,\kappa} \lesssim_S \kappa^{-1} \left(\|g\|_{s+\sigma_6}^{q,\kappa} + \|\mathfrak{I}_0\|_{s+\sigma_6}^{q,\kappa} \|g\|_{s_0+\sigma_6}^{q,\kappa} \right)$$

and for all $\lambda \in G_n$ one has

$$\mathbb{D}_n[\mathbb{D}_n]_{\text{ext}}^{-1} = \text{Id}.$$

Proof. (i) From (6.44), Lemma 6.1, Lemma 4.1-(ii), Lemma 6.3-(ii), we get the estimate on $\mathbb{P}_n^{\text{ext}}$. (ii) The estimate on \mathscr{P}_n can be easily obtained from (6.43), Lemma 6.4-(i), Lemma 4.1-(ii), Lemma 6.3-(i), Lemma 6.1-(ii).

- (iii) Follows immediately from (6.43) and Theorem 6.2.
- (iv) We shall look for an exact inverse of \mathbb{D}_n by solving the system

$$\mathbb{D}_{n}[\widehat{\phi},\widehat{y},\widehat{w},\widehat{\mathsf{c}}] = \begin{pmatrix} g_{1} \\ g_{2} \\ g_{3} \end{pmatrix}$$
(6.47)

where (g_1, g_2, g_3) satisfy the reversibility property

$$g_1(\varphi) = g_1(-\varphi), \quad g_2(\varphi) = -g_2(-\varphi), \quad g_3(\varphi) = -(\mathscr{S}g_3)(-\varphi).$$
 (6.48)

We first consider the second equation in (6.47) which takes the form, according to (6.41),

$$\mathfrak{D}_{(n)}\,\widehat{y} = g_2 - \mathscr{B}(\varphi)\widehat{\mathsf{c}}.$$

Since g_2 is odd, the φ -average of the right hand side of this equation is zero then by Lemma 6.4-(iv) its solution on the Cantor set DC_{N_n} is

$$\widehat{y} \triangleq [\mathscr{D}_{(n)}]_{\text{ext}}^{-1} \left(g_2 - \mathscr{B}(\varphi) \widehat{\mathsf{c}} \right).$$
(6.49)

Then we consider the third equation $\widehat{L}_{\omega,n}\widehat{w} = g_3 + \partial_\theta K_{11}(\varphi)\widehat{y} + \partial_\theta L_2^{\top}(\varphi)\widehat{c}$, which, by Theorem 6.2, has a solution

$$\widehat{w} \triangleq \mathsf{T}_{\omega,n} \left(g_3 + \partial_\theta K_{11}(\varphi) \widehat{y} + \partial_\theta L_2^\top(\varphi) \widehat{\mathsf{c}} \right).$$
(6.50)

Finally, we solve the first equation in (6.47), which, substituting (6.49), (6.50), becomes

$$\mathcal{D}_{(n)}\widehat{\phi} = g_1 + M_1(\varphi)\widehat{c} + M_2(\varphi)g_2 + M_3(\varphi)g_3, \qquad (6.51)$$

where

$$M_1(\varphi) \triangleq L_1^{\top}(\varphi) - M_2(\varphi) \mathscr{B}(\varphi) + M_3(\varphi) \partial_{\theta} L_2^{\top}(\varphi), \qquad (6.52)$$

$$M_2(\varphi) \triangleq K_{20}(\varphi) [\mathcal{D}_{(n)}]_{\text{ext}}^{-1} + K_{11}^{\top}(\varphi) \mathsf{T}_{\omega,n} \partial_{\theta} K_{11}(\varphi) [\mathcal{D}_{(n)}]_{\text{ext}}^{-1}, \qquad (6.53)$$

$$M_3(\varphi) \triangleq K_{11}^{\top}(\varphi) \mathsf{T}_{\omega,n} \,. \tag{6.54}$$

In order to solve equation (6.51) we have to choose \hat{c} such that the right hand side has zero average. In fact, by Lemma 6.1, (6.4), (6.38), Lemma 6.4-(ii) we get

$$\|M_2(\varphi)g\|_s^{q,\kappa} + \|M_3(\varphi)g\|_s^{q,\kappa} \lesssim \varepsilon \Big(\|g\|_{s+\sigma}^{q,\kappa} + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\|g\|_{s_0+\sigma}^{q,\kappa}\Big).$$

Then from Lemma 6.1, (9.278), the ϕ -averaged matrix is $\langle M_1 \rangle = \mathrm{Id} + O(\varepsilon \kappa^{-1})$. Therefore, for $\varepsilon \kappa^{-1}$ small enough, $\langle M_1 \rangle$ is invertible and $\langle M_1 \rangle^{-1} = \mathrm{Id} + O(\varepsilon \kappa^{-1}) = O(1)$. Thus we define

$$\widehat{\mathbf{c}} \triangleq -\langle M_1 \rangle^{-1} (\langle g_1 \rangle + \langle M_2 g_2 \rangle + \langle M_3 g_3 \rangle) \,. \tag{6.55}$$

Therefore we deduce that

$$\|\widehat{\mathbf{c}}\|^{q,\kappa} \lesssim \|g\|^{q,\kappa}_{s_0+\sigma} \tag{6.56}$$

With this choice of \hat{c} , equation (6.51) has the solution

$$\widehat{\phi} \triangleq [\mathfrak{D}_{(n)}]_{\text{ext}}^{-1} \left(g_1 + M_1(\varphi) [\widehat{\mathbf{c}}] + M_2(\varphi) g_2 + M_3(\varphi) g_3 \right).$$
(6.57)

Coming back to (6.49) and using Lemma 6.1-(iv), Lemma 6.4-(ii), (6.56) we obtain

$$\|\widehat{y}\|_{s}^{q,\kappa} \lesssim \kappa^{-1} \left(\|g\|_{s+\sigma}^{q,\kappa} + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|g\|_{s_{0}+\sigma}^{q,\kappa} \right).$$

$$(6.58)$$

The bound in (iv) for \hat{w} can be easily obtained from (6.50), (6.56), (6.38), (6.58), Lemma 6.1-(iv) and the smallness condition. Consequently, we find that $\hat{\phi}$ satisfies the estimate in (iv) using (6.57), Lemma 6.1, Lemma 6.4-(ii) and (6.38). Finally, the identity

$$\forall \lambda \in \mathbf{G}_n, \quad \mathbb{D}_n[\mathbb{D}_n]_{\mathrm{ext}}^{-1} = \mathrm{Id}$$

follows from Lemma 6.4 and (6.39).

Finally, we shall prove that the operator

$$T_0 \triangleq T_0(i_0) \triangleq D\widetilde{G}_0(u_0) [\mathbb{D}_n]_{ext}^{-1} [DG_0(u_0)]^{-1}$$
 (6.59)

is an approximate right inverse for $d_{i,c}\mathcal{F}(i_0)$.

Theorem 6.1. (Approximate inverse)

Let $(\gamma, q, d, \tau_1, s_0)$ satisfy (9.2), (μ_2, s_h) satisfy (9.277) and assume (9.278). Then there exists $\overline{\sigma} = \overline{\sigma}(\tau_1, \tau_2, d, q) > 0$ such that for smooth $g = (g_1, g_2, g_3)$, satisfying (6.48), the operator T_0 defined in (6.59) satisfies

$$\forall s \in [s_0, S], \quad \|\mathbf{T}_0 g\|_s^{q,\kappa} \lesssim \kappa^{-1} \left(\|g\|_{s+\overline{\sigma}}^{q,\kappa} + \|\mathfrak{I}_0\|_{s+\overline{\sigma}}^{q,\kappa} \|g\|_{s_0+\overline{\sigma}}^{q,\kappa} \right).$$
(6.60)

Moreover T_0 is an almost-approximate right inverse of $d_{i,c}\mathcal{F}(i_0, c_0)$ on the Cantor set G_n . More precisely, for all $\lambda \in G_n$ one has

$$d_{i,c}\mathcal{F}(i_0, \mathbf{c}_0)\mathbf{T}_0 - \mathbf{Id} = \mathcal{E}_1^{(n)} + \mathcal{E}_2^{(n)} + \mathcal{E}_3^{(n)}, \tag{6.61}$$

where the operators $\mathscr{E}_1^{(n)}$, $\mathscr{E}_2^{(n)}$ and $\mathscr{E}_3^{(n)}$ are defined in the whole set \mathscr{O} with the estimates

$$\begin{split} \|\mathscr{E}_{1}^{(n)}h\|_{s_{0}}^{q,\kappa} &\lesssim \overline{\gamma}^{-1} \|\mathscr{F}(i_{0},\mathsf{c}_{0})\|_{s_{0}+\overline{\sigma}}^{q,\kappa} \|h\|_{s_{0}+\overline{\sigma}}^{q,\kappa}, \\ \|\mathscr{E}_{2}^{(n)}h\|_{s_{0}}^{q,\kappa} &\lesssim_{b} \kappa^{-1}N_{n}^{-b} \left(\|h\|_{s_{0}+\overline{\sigma}+b}^{q,\kappa} + \|\Im_{0}\|_{s_{0}+\overline{\sigma}+b}^{q,\kappa} \|h\|_{s_{0}+\overline{\sigma}}^{q,\kappa}\right), \quad \forall b \ge 0, \\ \|\mathscr{E}_{3}^{(n)}h\|_{s_{0}}^{q,\kappa} &\lesssim N_{n}^{-b}\kappa^{-2} \left(\|h\|_{s_{0}+b+\overline{\sigma}}^{q,\kappa} + \varepsilon\kappa^{-3}\|\Im_{0}\|_{s_{0}+b+\overline{\sigma}}^{q,\kappa} \|h\|_{s_{0}+\overline{\sigma}}^{q,\kappa}\right) + \varepsilon\kappa^{-5}N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}}\|h\|_{s_{0}+\overline{\sigma}}^{q,\kappa}, \quad \forall b \in [0,S]. \end{split}$$

Proof. The bound (6.60) follows from (6.59), Proposition 6.3-(iv) and Lemma 6.1-(i). Then, according to (6.45) and (6.46), on the Cantor set G_n we have the decomposition

$$\begin{aligned} d_{i,c}\mathscr{F}(i_0, \mathbf{c}_0) &= DG_0(\mathbf{u}_0) \, \mathbb{L}_{\text{ext}} \, [D\tilde{G}_0(\mathbf{u}_0)]^{-1} \\ &= DG_0(\mathbf{u}_0) \, \mathbb{D}_n \, [D\tilde{G}_0(\mathbf{u}_0)]^{-1} + DG_0(\mathbf{u}_0) \, \mathbb{E}_n^{\text{ext}} \, [D\tilde{G}_0(\mathbf{u}_0)]^{-1} \\ &+ DG_0(\mathbf{u}_0) \, \mathscr{P}_n \, [D\tilde{G}_0(\mathbf{u}_0)]^{-1} + DG_0(\mathbf{u}_0) \, \mathscr{Q}_n \, [D\tilde{G}_0(\mathbf{u}_0)]^{-1}. \end{aligned}$$

Applying T₀, defined in (6.59), to the right of the last identity we get for all $\lambda = (\omega, \alpha) \in G_n$

$$d_{i,c}\mathcal{F}(i_0, \mathsf{c}_0)\mathrm{T}_0 - \mathrm{Id} = \mathcal{E}_1^{(n)} + \mathcal{E}_2^{(n)} + \mathcal{E}_3^{(n)}$$

with

$$\begin{split} & \mathscr{E}_1^{(n)} \triangleq DG_0(\mathsf{u}_0) \, \mathbb{E}_n^{\text{ext}} \, [D\widetilde{G}_0(\mathsf{u}_0)]^{-1} \mathcal{T}_0, \\ & \mathscr{E}_2^{(n)} \triangleq DG_0(\mathsf{u}_0) \, \mathscr{P}_n \, [D\widetilde{G}_0(\mathsf{u}_0)]^{-1} \mathcal{T}_0, \\ & \mathscr{E}_3^{(n)} \triangleq DG_0(\mathsf{u}_0) \, \mathscr{Q}_n \, [D\widetilde{G}_0(\mathsf{u}_0)]^{-1} \mathcal{T}_0. \end{split}$$

The estimates on $\mathscr{E}_1^{(n)}$, $\mathscr{E}_2^{(n)}$ and $\mathscr{E}_3^{(n)}$ follow from (6.60), Proposition 6.3 and Lemma 6.1-(i).

7. TOROIDAL PSEUDO-DIFFERENTIAL OPERATORS

In this section we shall collect some tools from function spaces and toroidal pseudo-differential operators. In particular, we shall introduce suitable functions spaces, discuss some operator topologies related to pseudo-differential operators and establish some useful commutator estimates connected with modified fractional Laplacian.

7.1. Operators and reversibility. This section is devoted to some discussions on parameterdependent operator norms. Consider a smooth family of bounded operators acting on Sobolev spaces $H^{s}(\mathbb{T}^{d+1})$, that is a smooth map

$$\mathscr{A}: \lambda \in \mathcal{O} \to \mathscr{A}(\lambda) \in \mathscr{L}(H^{s}(\mathbb{T}^{d+1}); H^{s}(\mathbb{T}^{d+1}))$$

of linear continuous operators $\mathscr{A}(\lambda)$: $H^{s}(\mathbb{T}^{d+1}) \to H^{s}(\mathbb{T}^{d+1})$. Then $\mathscr{A}(\lambda)$ can be represented through the infinite matrix $\left(\mathscr{A}_{l_0,j_0}^{l,j}(\lambda)\right)_{\substack{(l,l_0)\in(\mathbb{Z}^d)^2\\(j,j_0)\in\mathbb{Z}^2}}$ with

$$\mathscr{A}(\lambda)\mathbf{e}_{l_{0},j_{0}} = \sum_{(l,j)\in\mathbb{Z}^{d}\times\mathbb{Z}}\mathscr{A}_{l_{0},j_{0}}^{l,j}(\lambda)\mathbf{e}_{l,j}$$

and

$$\mathcal{A}_{l_{0},j_{0}}^{l,j}(\lambda) \triangleq \langle \mathcal{A}(\lambda)\mathbf{e}_{l_{0},j_{0}},\mathbf{e}_{l,j}\rangle_{L^{2}(\mathbb{T}^{d+1})} \\ = \frac{1}{(2\pi)^{d+1}} \int_{\mathbb{T}^{d+1}} \left(\mathcal{A}(\lambda)\mathbf{e}_{l_{0},j_{0}} \right) (\varphi,\theta) \,\mathbf{e}_{-l,-j}(\varphi,\theta) d\varphi d\theta.$$

We say that the operator $\mathscr{A}(\lambda)$ is Töplitz in time (actually in the variable φ) if its Fourier coefficients satisfy,

$$\mathscr{A}_{l_0,j_0}^{l,j}(\lambda) = \mathscr{A}_{0,j_0}^{l-l_0,j}(\lambda).$$

Now we say that an element \mathscr{A} belongs to the function space $W^{q,\infty}_{\kappa}(\mathcal{O};\mathcal{L}(H^s;H^s))$ if and only if

$$\|\mathscr{A}\|_{s}^{q,\kappa} = \max_{\substack{\alpha \in \mathbb{N}^{d} \\ |\alpha| \leq q}} \gamma^{|\alpha|} \sup_{\lambda \in \mathcal{O}} \|\partial_{\lambda}^{\alpha} \mathscr{A}(\lambda)\|_{\mathscr{L}(H^{s-|\alpha|})} < \infty.$$

$$(7.1)$$

The next result deals with the algebra structure of the preceding space.

Lemma 7.1. Let $\mathcal{A}_1, \mathcal{A}_2 \in W^{q,\infty}_{\kappa}(\mathcal{O}; \mathcal{L}(H^s; H^s))$, then $\mathcal{A}_1\mathcal{A}_2 \in W^{q,\infty}_{\kappa}(\mathcal{O}; \mathcal{L}(H^s; H^s))$ with $\|\mathscr{A}_1\mathscr{A}_2\|_s^{q,\kappa} \leqslant C \|\mathscr{A}_1\|_s^{q,\kappa} \|\mathscr{A}_2\|_s^{q,\kappa}.$

We remark that this class of operators is not useful in our case because it is too weak to get tame estimates. Therefore it will be replaced by a stronger one that will be detailed in Section 7.2. Notice also that along this paper the operators and the test functions may depend on the same parameter λ and thus the action of a family $\mathscr{A} = (\mathscr{A}(\lambda))$ on an element $h \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^d))$ is by convention defined through

$$(\mathscr{A}h)(\lambda,\varphi,\theta) \triangleq \mathscr{A}(\lambda)h(\lambda,\varphi,\theta),$$

meaning that we observe both objects at the same point λ . This is justified by the fact that with respect to the equations that we consider here, λ is an external parameter in the model.

In what follows we shall collect some definitions and properties about reversible operators. We recall the following definition, see for instance [4, 13, 14, 16]. We recall the involution \mathscr{S} introduced in Proposition 2.1,

$$\forall (\varphi, \theta) \in \mathbb{T}^{d+1}, \quad (\mathcal{G}h)(\varphi, \theta) = h(-\varphi, -\theta).$$
(7.2)

Definition 7.1. Consider a λ -dependent family of operators $\mathcal{A}(\lambda)$. We say that \mathcal{A} is

(i) real if for all $h \in L^2(\mathbb{T}^{d+1}, \mathbb{C})$, we have

$$\overline{h} = h \implies \overline{\mathcal{A}h} = \mathcal{A}h.$$

(ii) reversible if

$$\mathscr{A}(\lambda) \circ \mathscr{S} = -\mathscr{S} \circ \mathscr{A}(\lambda).$$

(iii) reversibility preserving if

$$\mathscr{A}(\lambda)\circ\mathscr{S}=\mathscr{S}\circ\mathscr{A}(\lambda).$$

We now give a useful characterization based on Fourier coefficients and the proof is straightforward.

Proposition 7.1. Let \mathcal{A} be an operator, then we have the following results.

(i) \mathcal{A} is real if and only if

$$\forall l, l_0, j_0, j \in \mathbb{Z}, \quad \mathscr{A}_{-l_0, -j_0}^{-l, -j} = \overline{\mathscr{A}_{l_0, j_0}^{l, j}}.$$

(ii) \mathcal{A} is reversible if and only if

$$\forall l, l_0, j_0, j \in \mathbb{Z}, \quad \mathscr{A}_{-l_0, -j_0}^{-l, -j} = -\mathscr{A}_{l_0, j_0}^{l, j}.$$

(iii) \mathcal{A} is reversibility-preserving if and only if

$$\forall l, l_0, j_0, j \in \mathbb{Z}, \quad \mathscr{A}_{-l_0, -j_0}^{-l, -j} = \mathscr{A}_{l_0, j_0}^{l, j}.$$

7.2. Symbol class topology. In this section we shall collect some classical results on pseudodifferential operators on the periodic setting. Let $\mathscr{A}: \mathscr{C}^{\infty}(\mathbb{T}) \to \mathscr{C}^{\infty}(\mathbb{T})$ be a linear operator, we define its symbol $\sigma_{\mathcal{A}}$ by

$$\forall (\theta, \xi) \in \mathbb{T} \times \mathbb{Z}, \quad \sigma_{\mathscr{A}}(\theta, \xi) \triangleq \mathbf{e}_{-\xi}(\theta)(\mathscr{A}\mathbf{e}_{\xi})(\theta), \quad \mathbf{e}_{\xi}(\theta) = e^{\mathrm{i}\theta\,\xi}.$$

Then for $h(\theta) = \sum_{\xi \in \mathbb{Z}} h_{\xi} e^{i\theta \cdot \xi}$ we get

$$\mathscr{A}h(\theta) = \sum_{\xi \in \mathbb{Z}} \sigma_{\mathscr{A}}(\theta, \xi) h_{\xi} e^{\mathrm{i}\theta\,\xi}.$$

One also gets the kernel representation

$$\mathscr{A}h(\theta) = \int_{\mathbb{T}} K(\theta, \eta) h(\eta) d\eta, \quad \text{with} \quad K(\theta, \eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} \sigma_{\mathscr{A}}(\theta, \xi) e^{i(\theta - \eta)\xi}.$$

By the Fourier inversion formula we may recover the symbol from the kernel through the formula

$$\sigma_{\mathscr{A}}(\theta,\xi) = \int_{\mathbb{T}} K(\theta,\theta+\eta) e^{\mathrm{i}\eta\xi} d\eta$$

Next, we shall introduce the difference operators in the periodic setting. Consider a discrete function $h : \mathbb{Z} \to \mathbb{C}$ and define the forward difference operator Δ_{ξ} by

$$\Delta_{\xi}h(\xi) = h(\xi+1) - h(\xi) \quad \text{and} \quad \Delta_{\xi}^{\ell} = \underbrace{\Delta_{\xi} \circ \ldots \circ \Delta_{\xi}}_{\ell - \text{ times}}.$$
(7.3)

The backward difference $\overline{\Delta}_{\xi}$ is defined by

$$\overline{\Delta}_{\xi}h(\xi) \triangleq h(\xi) - h(\xi - 1), \quad \overline{\Delta}_{\xi}^{\ell} = \underbrace{\overline{\Delta}_{\xi} \circ \dots \circ \overline{\Delta}_{\xi}}_{\ell - \text{ times}}$$

We will recall some useful identities of the difference operator, see for instance [78, Chapter 3]. First, the discrete Leibniz formula reads as follows,

$$\forall \gamma \in \mathbb{N}, \quad \Delta_{\xi}^{\gamma}(fg)(\xi) = \sum_{0 \leq \beta \leq \gamma} {\gamma \choose \beta} \Delta_{\xi}^{\beta} f(\xi) \Delta_{\xi}^{\gamma-\beta} g(\xi+\beta).$$
(7.4)

The next one is the summation by parts

$$\sum_{\xi \in \mathbb{Z}} f(\xi) \Delta_{\xi}^{\gamma}(g)(\xi) = (-1)^{|\gamma|} \sum_{\xi \in \mathbb{Z}} \left(\overline{\Delta}_{\xi}^{\gamma} f(\xi) \right) g(\xi).$$
(7.5)

The next property is easy to check by induction

$$\Delta_{\xi}^{\gamma} e^{i\xi\eta} = \left(e^{i\eta} - 1\right)^{\gamma} e^{i\xi\eta}.$$
(7.6)

Throughout this paper we shall make use of multi-parameter pseudo-differential operators $\varphi \in \mathbb{T}^d \mapsto \mathscr{A}(\varphi)$. This yields to one-parameter symbol $\sigma_{\mathscr{A}}(\varphi)$ defined in the following way: for a smooth periodic function $h(\varphi, \theta) = \sum_{\xi \in \mathbb{Z}} h_{\xi}(\varphi) e^{i\theta \cdot \xi}$

$$\mathscr{A}(\varphi)h(\varphi,\theta) = \sum_{\xi \in \mathbb{Z}} \sigma_{\mathscr{A}}(\varphi,\theta,\xi)h_{\xi}(\varphi)e^{i\theta\,\xi}.$$
(7.7)

The symbol can be recovered from the operator as follows

$$\forall (\theta, \xi) \in \mathbb{T} \times \mathbb{Z}, \quad \sigma_{\mathscr{A}}(\varphi, \theta, \xi) \triangleq \mathbf{e}_{-\xi}(\theta)(\mathscr{A}(\varphi)\mathbf{e}_{\xi})(\theta), \quad \mathbf{e}_{\xi}(\theta) = e^{\mathrm{i}\theta\,\xi}$$

and the kernel representation takes the form

$$\mathscr{A}(\varphi)h(\varphi,\theta) = \int_{\mathbb{T}} K(\varphi,\theta,\eta)h(\varphi,\eta)d\eta, \quad \text{with} \quad K(\varphi,\theta,\eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} \sigma_{\mathscr{A}}(\varphi,\theta,\xi)e^{\mathrm{i}(\theta-\eta)\xi}.$$

By the Fourier inversion formula we get

$$\sigma_{\mathscr{A}}(\varphi,\theta,\xi) = \int_{\mathbb{T}} K(\varphi,\theta,\theta+\eta) e^{i\eta\xi} d\eta.$$
(7.8)

Let $s, m \in \mathbb{R}, \gamma \in \mathbb{N}$ and define the norm over the symbol class

$$\|\mathscr{A}\|\|_{m,s,\gamma} \triangleq \sup_{\substack{\xi \in \mathbb{Z} \\ 0 \leqslant \ell \leqslant \gamma}} \langle \xi \rangle^{-m+\ell} \|\Delta_{\xi}^{\ell} \sigma_{\mathscr{A}}(\cdot, \cdot, \xi)\|_{H^{s}(\mathbb{T}^{d+1})}.$$
(7.9)

The case $m = \gamma = 0$ plays a central role along the paper, needed especially in the remainder KAM reduction, and for the sake of simple notation we simply write

$$\||\mathcal{A}|||_{s} \triangleq |||\mathcal{A}|||_{0,s,0}.$$

We shall give an equivalent form to this norm using Fourier coefficients which turns out to be very useful later at some points. Define

$$\mathscr{A}_{j'}^{j}(l) = \langle \mathscr{A} \mathbf{e}_{0,j'}, \mathbf{e}_{l,j} \rangle_{L^{2}(\mathbb{T}^{d+1})} \quad \text{with} \quad \mathbf{e}_{l,j}(\varphi, \theta) = e^{\mathrm{i}(l \cdot \varphi + j\theta)}.$$

Then it is easy to check that for any $j \in \mathbb{Z}$ (the operator is Töplitz in φ)

$$\sigma_{\mathscr{A}}(\varphi,\theta,j) = \mathbf{e}_{0,-j}(\varphi,\theta) \big(\mathscr{A} \mathbf{e}_{0,j} \big)(\varphi,\theta)$$

$$= \sum_{(l,j') \in \mathbb{Z}^{d+1}} \mathscr{A}_j^{j'}(l) \mathbf{e}_{l,j'-j}(\varphi, \theta).$$

Consequently

$$\|\sigma_{\mathscr{A}}(\cdot, \cdot, j)\|_{H^s}^2 = \sum_{(l,j')\in\mathbb{Z}^{d+1}} \left|\mathscr{A}_j^{j+j'}(l)\right|^2 \langle l, j'\rangle^{2s},$$

which implies that

$$\|\mathscr{A}\|_{s}^{2} = \sup_{j \in \mathbb{Z}} \|\sigma_{\mathscr{A}}(\cdot, \cdot, j)\|_{H^{s}}^{2}$$
$$= \sup_{j \in \mathbb{Z}} \sum_{(l,j') \in \mathbb{Z}^{d+1}} \langle l, j' \rangle^{2s} |\mathscr{A}_{j}^{j+j'}(l)|^{2}.$$
(7.10)

We shall also make use of weighted pseudo-differential operators $(\lambda, \varphi) \in \mathcal{O} \times \mathbb{T}^d \mapsto \mathscr{A}(\lambda, \varphi)$ where \mathcal{O} is a bounded open set of \mathbb{R}^{d+1} . First the multi-parameter symbol of \mathscr{A} is defined as follows,

$$h(\lambda,\varphi,\theta) = \sum_{\xi\in\mathbb{Z}} h_{\xi}(\lambda,\varphi) e^{\mathrm{i}\theta\,\xi} \Longrightarrow \,\mathscr{A}(\lambda,\varphi)h(\lambda,\varphi,\theta) = \sum_{\xi\in\mathbb{Z}} \sigma_{\mathscr{A}}(\lambda,\varphi,\theta,\xi)h_{\xi}(\lambda,\varphi)e^{\mathrm{i}\theta\,\xi}.$$
 (7.11)

In this case we have the kernel representation,

$$\mathscr{A}(\lambda,\varphi)h(\lambda,\varphi,\theta) = \int_{\mathbb{T}} K(\lambda,\varphi,\theta,\eta)h(\lambda,\varphi,\eta)d\eta$$
(7.12)

with

$$K(\lambda,\varphi,\theta,\eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} e^{i(\theta-\eta)\xi} \sigma_{\mathscr{A}}(\lambda,\varphi,\theta,\xi).$$

Take $m, s \in \mathbb{R}, \gamma \in \mathbb{N}$ and $\kappa \in (0, 1)$ and define the weighted norm

$$\||\mathscr{A}\||_{m,s,\gamma}^{q,\kappa} \triangleq \max_{\substack{\beta \in \mathbb{N}^{d+1} \\ |\beta| \leqslant q}} \kappa^{|\beta|} \sup_{\lambda \in \mathcal{O}} \||\partial_{\lambda}^{\beta} \mathscr{A}\||_{m,s-|\beta|,\gamma}.$$
(7.13)

We shall adopt for the particular case $m = \gamma = 0$ the following notation

$$|||\mathscr{A}|||_{s}^{q,\kappa} \triangleq |||\mathscr{A}|||_{0,s,0}^{q,\kappa}$$

Using (7.10) we get the following characterization,

$$(\||\mathscr{A}\||_{s}^{q,\kappa})^{2} = \max_{|\beta| \leq q} \sup_{\lambda \in \mathcal{O}} \sup_{j \in \mathbb{Z}} \sum_{(l,j') \in \mathbb{Z}^{d+1}} \kappa^{2|\beta|} \langle l, j' \rangle^{2(s-|\beta|)} \left| \partial_{\lambda}^{\beta} \mathscr{A}_{j}^{j+j'}(\lambda, l) \right|^{2}.$$
(7.14)

The next lemma is very useful.

Lemma 7.2. Let $q, \gamma \in \mathbb{N}, m, s \in \mathbb{R}$, then the following assertions hold true.

- (i) Let T_M be a multiplication operator by a real-valued function M, that is, T_Mh = Mh. Then
 (a) If M(λ, -φ, -θ) = M(λ, φ, θ), then T_M is real and reversibility preserving Töplitz in time and space operator.
 - (b) If $M(\lambda, -\varphi, -\theta) = -M(\lambda, \varphi, \theta)$, then T_M is real and reversible Töplitz in time and space operator. Moreover,

$$|||T_M|||_{0,s,\gamma}^{q,\kappa} = ||M||_s^{q,\kappa}.$$

- (ii) Let \mathcal{A} be an integral operator with a real-valued kernel K as in (7.12).
 - (a) If $K(\lambda, -\varphi, -\theta, -\eta) = K(\lambda, \varphi, \theta, \eta)$, then \mathcal{A} is real and reversibility preserving Töplitz in time operator.
 - (b) If $K(\lambda, -\varphi, -\theta, -\eta) = -K(\lambda, \varphi, \theta, \eta)$, then \mathcal{A} is real and reversible Töplitz in time operator. In addition,

$$\|\|\mathscr{A}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \sum_{0 \leqslant \ell \leqslant \gamma} \int_{\mathbb{T}} |\sin(\eta/2)|^{\ell} \|\partial_{\eta}^{\ell}(-\Delta_{\eta})^{-\frac{m}{2}} K(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} d\eta.$$

(iii) Let \mathscr{A} as in (7.12) and $m < -\frac{1}{2}$ then

$$\int_{\mathbb{T}} \left(\|K(\cdot, \boldsymbol{\cdot}, \boldsymbol{\cdot} + \eta)\|_{s}^{q,\kappa} \right)^{2} d\eta \lesssim \left(\|\|\mathcal{A}\|\|_{m,s,0}^{q,\kappa} \right)^{2}.$$

Proof. (i)-(ii) The proofs are starightfoward.

(iii) Recall that the link between the kernel and the symbol is given by

$$K(\varphi, \theta, \theta + \eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} \sigma_{\mathscr{A}}(\varphi, \theta, \xi) e^{-i\eta\xi}.$$

Thus from Bessel identity we get for any $(\varphi, \theta) \in \mathbb{T}^{d+1}$

$$\int_{\mathbb{T}} |K(\varphi, \theta, \theta + \eta)|^2 d\eta = \sum_{\xi \in \mathbb{Z}} |\sigma_{\mathscr{A}}(\varphi, \theta, \xi)|^2.$$

Integrating in (φ, θ) we get

$$\int_{\mathbb{T}} \|K(\cdot, \boldsymbol{\cdot}, \boldsymbol{\cdot} + \eta)\|_{L^{2}_{\varphi, \theta}}^{2} d\eta = \sum_{\xi \in \mathbb{Z}} \|\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi)\|_{L^{2}_{\varphi, \theta}}^{2}$$

Then differentiating in φ, θ and using the same argument as before we get for $m < -\frac{1}{2}$,

$$\int_{\mathbb{T}} \|K(\cdot, \cdot, \cdot + \eta)\|^{2}_{H^{s}_{\varphi, \theta}} d\eta \lesssim \sum_{\xi \in \mathbb{Z}} \|\sigma_{\mathscr{A}}(\cdot, \cdot, \xi)\|^{2}_{H^{s}_{\varphi, \theta}}$$
$$\lesssim \|\|\mathscr{A}\|\|^{2}_{m, s, 0} \sum_{\xi \in \mathbb{Z}} \langle \xi \rangle^{2m}$$
$$\lesssim \|\|\mathscr{A}\|\|^{2}_{m, s, 0}.$$

Similarly we obtain for any $q \in \mathbb{N}$,

$$\int_{\mathbb{T}} \left(\|K(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \right)^{2} d\eta \lesssim \sum_{\xi \in \mathbb{Z}} \left(\|\sigma_{\mathscr{A}}(\cdot, \cdot, \xi)\|_{s}^{q,\kappa} \right)^{2} \\ \lesssim \left(\|\|\mathscr{A}\|\|_{m,s,0}^{q,\kappa} \right)^{2}.$$

This ends the proof.

7.3. Continuity and law products. This section is devoted to some classical results. First, we define the frequency cut-off projectors $(P_N)_{N \in \mathbb{N}^*}$ as follows,

$$(P_N \mathscr{A}(\lambda)) \mathbf{e}_{l_0, j_0} = \sum_{\substack{(l,j) \in \mathbb{Z}^d \times \mathbb{Z} \\ |l-l_0|, |j-j_0| \leqslant N}} \mathscr{A}_{l_0, j_0}^{l, j}(\lambda) \mathbf{e}_{l, j} \quad \text{and} \quad P_N^{\perp} \mathscr{A} = \mathscr{A} - P_N \mathscr{A},$$
(7.15)

where \mathscr{A} is a pseudo-differential operator as in Section 7.2. In the next lemma we shall gather classical results some of the proofs can be found for instance in [16, 78].

Lemma 7.3. Let \mathcal{A} be a pseudo-differential operator as in (7.7), then the following assertions hold. (i) Let $s_0 > d + 1$, then

 $\|\mathscr{A}h\|_{L^2(\mathbb{T}^{d+1})} \lesssim \|\mathscr{A}\|\|_{s_0} \|h\|_{L^2(\mathbb{T}^{d+1})}.$

(ii) If $s \ge 0, s_0 > \frac{d+1}{2}$ then $\|\mathscr{A}h\|_{H^s} \lesssim \|\mathscr{A}\|_{s_0} \|h\|_{H^s} + \|\mathscr{A}\|_{s_0} \|h\|_{H^{s_0}}.$ (iii) Let \mathscr{A} as in (7.11) and $s \ge q, s_0 > \frac{d+1}{2} + q$. Then $\|\mathscr{A}h\|_s^{q,\overline{\gamma}} \lesssim \|\mathscr{A}\|\|_{s_0}^{q,\kappa} \|h\|_s^{q,\kappa} + \||\mathscr{A}\|\|_s^{q,\kappa} \|h\|_{s_0}^{q,\kappa}.$ (iv) Let $N \in \mathbb{N}^*, s \in \mathbb{R}, t \ge 0$, then $|||P_N \mathscr{A}|||_{s+t}^{q,\kappa} \leqslant N^t ||| \mathscr{A} |||_s^{q,\kappa} \quad and \quad ||| P_N^{\perp} \mathscr{A} |||_s^{q,\kappa} \leqslant N^{-t} ||| \mathscr{A} |||_{s+t}^{q,\kappa}.$ (v) Interpolation inequality : Let $q < s_1 \leq s_3 \leq s_2, t \in [0,1]$ with $s_3 = ts_1 + (1-t)s_2$. Then

 $\|\|\mathscr{A}\|\|_{s_3}^{q,\kappa} \leqslant C(q,d) \left(\||\mathscr{A}\|\|_{s_1}^{q,\kappa} \right)^t \left(\||\mathscr{A}\|\|_{s_2}^{q,\kappa} \right)^{1-t}.$

(vi) Adjoint estimate: the L^2 adjoint \mathcal{A}^* satisfies

 $\left\|\left\|\mathscr{A}^*\right\|\right\|_{m,s,0}^{q,\kappa} \lesssim \left\|\left|\mathscr{A}\right|\right\|_{m,s+s_0+|m|,0}^{q,\kappa}.$

Now we recall the following result, see for instance [16].

Lemma 7.4. Let \mathcal{A}, \mathcal{B} as in (7.11) and (κ, d, q, s, s_0) as in (4.1) and consider $m_1, m_2 \in \mathbb{R}, \gamma \in \mathbb{N}$. Then we have

 $\left\|\left|\mathscr{AB}\right|\right\|_{m_1+m_2,s,\gamma}^{q,\kappa} \lesssim \left\|\left|\mathscr{A}\right|\right|\right\|_{m_1,s,\gamma}^{q,\kappa} \left\|\left|\mathscr{B}\right|\right|\right\|_{m_2,s_0+\gamma+|m_1|,\gamma}^{q,\kappa} + \left|\left|\left|\mathscr{A}\right|\right|\right|\right\|_{m_1,s_0,\gamma}^{q,\kappa} \left|\left|\mathscr{B}\right|\right|\right\|_{m_2,s+\gamma+|m_1|,\gamma}^{q,\kappa}.$

In particular we get

$$\| \mathscr{A}\mathscr{B} \|_{s}^{q,\kappa} \lesssim \| \mathscr{A} \| \|_{s_{0}}^{q,\kappa} \| \mathscr{B} \| \|_{s}^{q,\kappa} + \mathscr{A} \| \|_{s}^{q,\kappa} \| \mathscr{B} \| \|_{s_{0}}^{q,\kappa}$$

Now we intend to establish refined estimates needed later to derive tame estimates in Section 8.

Lemma 7.5. Let \mathcal{A}, \mathcal{B} as in (7.11) and (κ, d, q, s, s_0) as in (4.1). Then the following assertions hold true.

$$\begin{array}{l} \text{(i) } Let \ m_1, m_2 \in \mathbb{R}, \ then \ for \ any \ \gamma \in \mathbb{N} \ and \ \epsilon > 0 \\ & \parallel \mathscr{AB} \parallel_{m_1+m_2,s,\gamma}^{q,\kappa} \lesssim \\ & \sum_{0 \leqslant \beta \leqslant \gamma} \left(\parallel \mathscr{A} \parallel_{m_1,s,\beta}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s_0+\frac{1}{2}+m_1^++\epsilon,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{A} \parallel_{m_1,s_0,\beta}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s+\frac{1}{2}+m_1^++\epsilon,\gamma-\beta}^{q,\kappa} \right) \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \left(\parallel \mathscr{A} \parallel_{m_1,s,0}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s_0+\frac{1}{2}+\beta-m_1^-,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{A} \parallel_{m_1,s_0,0}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s+\frac{1}{2}+\beta-m_1^-,\gamma-\beta}^{q,\kappa} \right) \right) \\ \text{(ii) } If \ m_1, m_2 \leqslant 1 \ and \ \gamma \in \mathbb{N}, \ then \ for \ any \ \epsilon > 0 \\ & \parallel [\mathscr{A}, \mathscr{B}] \parallel \mathscr{A}_{m_1,s,1+\beta}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{A} \parallel_{m_1,s_0,1+\beta}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{A} \parallel_{m_1,s,0}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{A} \parallel \mathscr{A} \parallel_{m_1,s_0,0}^{q,\kappa} \parallel \mathscr{B} \parallel_{m_2,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{A} \parallel_{m_2,s,1+\beta}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{B} \parallel_{m_2,s_0,1+\beta}^{q,\kappa} \parallel \mathscr{A} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa} + \parallel \mathscr{B} \parallel_{m_2,s_0,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} + \parallel \mathscr{B} \parallel_{m_2,s_0,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} + \parallel \mathscr{B} \parallel_{m_2,s_0,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s_0+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} + \parallel \mathscr{B} \parallel_{m_2,s_0,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} + \parallel \mathscr{B} \parallel_{m_2,s_0,0}^{q,\kappa} \parallel \mathscr{A} \parallel_{m_2,s,0}^{q,\kappa-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa-\beta} \parallel \mathscr{A} \parallel_{m_1,s+\frac{3}{2}+\epsilon,\gamma-\beta}^{q,\kappa-\beta} + \parallel \mathscr{B} \parallel_{m_2,s,0,0}^{q,\kappa-\beta} \parallel \mathscr{A} \parallel_{m_2,s+\frac{3}{2}+\epsilon,\gamma-\beta} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \parallel \mathscr{B} \parallel_{m_2,s,0}^{q,\kappa-\beta} \parallel \mathscr{A} \parallel_{m_2,s+\frac{3}{2}+\epsilon,\gamma-\beta} + \parallel \mathscr{B} \parallel_{m_2,s,0,0}^{q,\kappa-\beta} \parallel \mathscr{A} \parallel_{m_2,s,0,0}^{q,\kappa-\beta} \parallel \mathscr{A}$$

(iii) Let $m_1 + m_2 \leq 1$ and $\gamma \in \mathbb{N}$, then

$$\begin{split} & \|\| [\mathscr{A},\mathscr{B}] \|\|_{0,s,\gamma}^{q,\kappa} \lesssim \\ & \sum_{0 \leqslant \beta \leqslant \gamma} \|\| \mathscr{A} \|\|_{m_{1},s,1+\gamma-\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s_{0}+\mu,\beta}^{q,\kappa} + \|\| \mathscr{A} \|\|_{m_{1},s_{0},1+\gamma-\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s+\mu,\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \|\| \mathscr{A} \|\|_{m_{1},s,0}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s_{0}+\mu_{\beta},\gamma-\beta}^{q,\kappa} + \|\| \mathscr{A} \|\|_{m_{1},s_{0},0}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s+\mu_{\beta},\gamma-\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \|\| \mathscr{A} \|\|_{m_{1},s_{0}+\mu,\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s,1+\gamma-\beta}^{q,\kappa} + \|\| \mathscr{A} \|\|_{m_{1},s+\mu,\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s_{0},1+\gamma-\beta}^{q,\kappa} \\ & + \sum_{0 \leqslant \beta \leqslant \gamma} \|\| \mathscr{A} \|\|_{m_{1},s_{0}+\overline{\mu}_{\beta},\gamma-\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s,0}^{q,\kappa} + \|\| \mathscr{A} \|\|_{m_{1},s+\overline{\mu}_{\beta},\gamma-\beta}^{q,\kappa} \|\| \mathscr{B} \|\|_{m_{2},s_{0},0}^{q,\kappa}, \end{split}$$

provided that μ, μ_{β} and $\overline{\mu}_{\beta}$ satisfy

$$\mu > \frac{1}{2} + \max\left(0, m_1, m_2, m_1 + m_2\right), \quad \mu_\beta \ge \frac{1}{2} + m_1^+ + m_2 + \beta, \quad \overline{\mu}_\beta \ge \frac{1}{2} + m_1 + m_2^+ + \beta,$$
with the notation $m^+ = \max(m, 0)$ and $m^- = \min(m, 0).$

Proof. (i) We shall implement the proof for the case q = 0 and the general case $q \ge 1$ can be done similarly using Leibniz rule. Indeed, one writes

$$\kappa^{|\beta|}\partial_{\lambda}^{\beta}\mathscr{A}\mathscr{B} = \sum_{0\leqslant\beta'\leqslant\beta} \binom{\beta}{\beta'} \kappa^{|\beta'|} (\partial_{\lambda}^{\beta'}\mathscr{A}) \kappa^{|\beta-\beta'|} \partial_{\lambda}^{\beta-\beta'} \mathscr{B}$$

and then we may apply the case q = 0 in order to get the desired result. This remark concerns also the points (ii) and (iii). Now let us prove the estimate for q = 0. First notice that the symbol of the operator \mathscr{AB} is given by

$$\sigma_{\mathscr{AB}}(\varphi,\theta,\xi) = \sum_{\eta \in \mathbb{Z}} e^{\mathrm{i}\eta \cdot \theta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) \widehat{\sigma}_{\mathscr{B}}(\varphi,\eta,\xi)$$

where $\widehat{\sigma}_{\mathscr{B}}(\varphi, \eta, \xi)$ is the Fourier coefficients of the partial periodic function $\theta \in \mathbb{T} \mapsto \sigma_A(\varphi, \theta, \xi)$. Then one gets the decomposition,

$$\sigma_{\mathscr{A}\mathscr{B}}(\varphi,\theta,\xi) = \sigma^{\mathrm{H}}_{\mathscr{A}\mathscr{B}}(\varphi,\theta,\xi) + \sigma^{\mathrm{L}}_{\mathscr{A}\mathscr{B}}(\varphi,\theta,\xi)$$

with

$$\sigma^{\mathrm{H}}_{\mathscr{AB}}(\varphi,\theta,\xi) = \sum_{\substack{|\eta| \leqslant \frac{|\xi|}{2} \\ |\eta| \geqslant 2|\xi|}} \frac{\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta)}{|\eta|^{\mu}} e^{\mathrm{i}\eta \cdot \theta} \widehat{\Lambda^{\mu}_{\theta}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi)$$

and

$$\sigma^{\mathrm{L}}_{\mathscr{A}\mathscr{B}}(\varphi,\theta,\xi) = \sum_{\frac{|\xi|}{2} < |\eta| < 2|\xi|} \frac{\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta)}{|\eta|^{\mu_{\beta}}} e^{\mathrm{i}\eta \cdot \theta} \widehat{\Lambda^{\mu_{\beta}}_{\theta}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi)$$

Here, $\Lambda_{\theta} = \sqrt{-\Delta_{\theta}}$ and μ, μ_{β} are free real numbers. Let us start with studying the high frequency part. Then using Leibniz formula (7.4) yields

$$\Delta_{\xi}^{\gamma}\sigma_{\mathscr{AB}}^{\mathrm{H}}(\varphi,\theta,\xi) = \sum_{\beta \leqslant \gamma} \sum_{\substack{|\eta| \leqslant \frac{|\xi|}{2} \\ |\eta| \ge 2|\xi|}} \binom{\gamma}{\beta} \frac{\Delta_{\xi}^{\beta}\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta)}{|\eta|^{\mu}} e^{\mathrm{i}\eta \cdot \theta} \Delta_{\xi}^{\gamma-\beta} \widehat{\Lambda_{\theta}^{\mu}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi+\beta).$$

Applying Sobolev law products combined with (7.22), Cauchy-Schwarz inequality and the choice

$$\mu = \frac{1}{2} + m_1^+ + \epsilon, \epsilon > 0$$

we get

$$\begin{aligned} \left\| \Delta_{\xi}^{\gamma} \sigma_{\mathscr{AB}}^{\mathrm{H}}(\cdot, \cdot, \xi) \right\|_{H^{s}(\mathbb{T}^{d+1})} &\lesssim C_{0} \sum_{\substack{|\eta| \leq \frac{|\xi|}{2} \\ |\eta| \geq 2|\xi|}} c_{\eta}(\xi) |\eta|^{-\mu} \langle \xi + \eta \rangle^{m_{1}-\beta} \langle \xi \rangle^{m_{2}-\gamma+\beta} \\ &\lesssim C_{0} \langle \xi \rangle^{m_{1}+m_{2}-\gamma}, \end{aligned}$$
(7.16)

with $\sum_{\eta} c_{\eta}^2(\xi) = 1$ and

$$C_0 = \sum_{0 \leqslant \beta \leqslant \gamma} \left\| \left| \mathscr{A} \right| \right\|_{m_1, s, \beta} \left\| \left| \mathscr{B} \right| \right\|_{m_2, s_0 + \frac{1}{2} + m_1^+ + \epsilon, \gamma - \beta} + \left\| \left| \mathscr{A} \right| \right\|_{m_1, s_0, \beta} \left\| \left| \mathscr{B} \right| \right\|_{m_2, s + \frac{1}{2} + m_1^+ + \epsilon, \gamma - \beta}.$$

Now let us move to the low frequency part. First we have

$$\Delta_{\xi}^{\gamma} \sigma_{\mathscr{AB}}^{\mathrm{L}}(\varphi, \theta, \xi) = \sum_{\beta \leqslant \gamma} \sum_{\frac{|\xi|}{2} < |\eta| < 2|\xi|} \binom{\gamma}{\beta} \frac{\Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\varphi, \theta, \xi + \eta)}{|\eta|^{\mu_{\beta}}} e^{\mathrm{i}\eta \cdot \theta} \Delta_{\xi}^{\gamma - \beta} \widehat{\Lambda_{\theta}^{\mu_{\beta}}} \sigma_{\mathscr{B}}(\varphi, \eta, \xi + \beta).$$

Then by definition we get for $\frac{|\xi|}{2} \leqslant |\eta| \leqslant 2|\xi|$

$$\left\|\Delta_{\xi}^{\beta}\left(\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi + \eta)\right)\right\|_{H^{s}} \lesssim \langle \xi \rangle^{m_{1}^{+}} \parallel \mathscr{A} \parallel _{m_{1}, s, 0}.$$

Combined with the inequality (7.22), proved later, and taking

$$\mu_{\beta} = \frac{1}{2} + \beta - m_1^-$$

we find by virtue of Cauchy-Schwarz inequality

$$\Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{L}}(\cdot, \cdot, \xi) \|_{H^{s}} \lesssim \sum_{\substack{|\xi| \\ 2 \\ 0 \leqslant \beta \leqslant \gamma}} c_{\eta}(\xi) |\eta|^{-\mu_{\beta}} \langle \xi \rangle^{m_{1}^{+} + m_{2} - \gamma + \beta} C_{1,\beta}$$

$$\lesssim C_{1} \langle \xi \rangle^{m_{1} + m_{2} - \gamma} \sum_{\substack{|\xi| \\ 2 \\ < |\eta| < 2|\xi|}} c_{\eta}(\xi) |\eta|^{-\frac{1}{2}}$$

$$\lesssim C_{1} \langle \xi \rangle^{m_{1} + m_{2} - \gamma} \qquad (7.17)$$

with

$$C_1 = \sum_{0 \leqslant \beta \leqslant \gamma} C_{1,\beta}$$

This achieves the proof of the desired result.

(ii) We shall start with using the following decomposition

$$\sigma_{\mathscr{A}\mathscr{B}}(\varphi,\theta,\xi) = \sum_{\eta\in\mathbb{Z}} e^{i\eta\,\theta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) \widehat{\sigma}_{\mathscr{B}}(\varphi,\eta,\xi)$$

$$= \sum_{\eta\in\mathbb{Z}} e^{i\eta\,\theta} \Big(\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) - \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \Big) \widehat{\sigma}_{\mathscr{B}}(\varphi,\eta,\xi)$$

$$+ \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \sigma_{\mathscr{B}}(\varphi,\theta,\xi)$$

$$\triangleq \mathscr{R}_{1}(\varphi,\theta,\xi) + \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \sigma_{\mathscr{B}}(\varphi,\theta,\xi).$$
(7.18)

Let μ be an arbitrary parameter then we have the splitting,

$$\mathscr{R}_{1}(\varphi,\theta,\xi) = \mathscr{R}_{1}^{\mathrm{H}}(\varphi,\theta,\xi) + \mathscr{R}_{1}^{\mathrm{L}}(\varphi,\theta,\xi)$$

with

$$\mathscr{R}_{1}^{\mathrm{H}}(\varphi,\theta,\xi) = \sum_{\substack{|\eta| \leq \frac{|\xi|}{2} \\ |\eta| \geq 2|\xi|}} \frac{\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) - \sigma_{\mathscr{A}}(\varphi,\theta,\xi)}{|\eta|^{\mu}} e^{\mathrm{i}\eta \cdot \theta} \widehat{\Lambda_{\theta}^{\mu}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi)$$

and

$$\mathscr{R}_{1}^{\mathrm{L}}(\varphi,\theta,\xi) = \sum_{\frac{|\xi|}{2} < |\eta| < 2|\xi|} \left(\sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) - \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \right) e^{\mathrm{i}\eta\,\theta} \widehat{\sigma}_{\mathscr{B}}(\varphi,\eta,\xi).$$

Let us start with estimating the first term $\mathscr{R}_1^{\mathrm{H}}$. Concerning the region $|\eta| \leq \frac{|\xi|}{2}$ we may use discrete Taylor formulae, see [78, Chapter 3], leading to the following inequality

$$\left\|\Delta_{\xi}^{\beta}\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi+\eta) - \Delta_{\xi}^{\beta}\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi)\right\|_{H^{s}(\mathbb{T}^{d+1})} \lesssim |\eta| \sup_{|\nu| \leqslant |\eta|} \left\|\Delta_{\xi}^{1+\beta}\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi+\nu)\right\|_{H^{s}(\mathbb{T}^{d+1})}.$$
 (7.19)

Therefore, we get in view of (7.9),

$$\begin{aligned} \left\| \Delta_{\xi}^{\beta} \left[\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi + \eta) - \sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi) \right] \right\|_{H^{s}(\mathbb{T}^{d+1})} &\lesssim |\eta| \sup_{|\nu| \leqslant |\eta|} \langle \xi + \nu \rangle^{m_{1}-1-\beta} ||| \mathscr{A} |||_{m_{1}, s, 1+\beta} \\ &\lesssim |\eta| \langle \xi \rangle^{m_{1}-1-\beta} ||| \mathscr{A} |||_{m_{1}, s, 1+\beta} . \end{aligned}$$

$$(7.20)$$

However in the region $|\eta| \ge 2|\xi|$ one deduces from the triangle inequality combined with (7.9)

$$\begin{aligned} \left\| \Delta_{\xi}^{\beta} \left[\sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi + \eta) - \sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi) \right] \right\|_{H^{s}(\mathbb{T}^{d+1})} &\leq \left\| \Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi + \eta) \right\|_{H^{s}(\mathbb{T}^{d+1})} + \left\| \Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\cdot, \boldsymbol{\cdot}, \xi) \right\|_{H^{s}(\mathbb{T}^{d+1})} \\ &\lesssim \left\| |\mathscr{A}| \right\|_{m_{1}, s, \beta} \left(\langle \xi + \eta \rangle^{m_{1} - \beta} + \langle \xi \rangle^{m_{1} - \beta} \right) \\ &\lesssim \left\| |\mathscr{A}| \right\|_{m_{1}, s, 1 + \beta} \left(\langle \eta \rangle^{m_{1} - \beta} + \langle \xi \rangle^{m_{1} - \beta} \right). \end{aligned}$$
(7.21)

Next, we shall use the following identity where we make appeal to integration by parts and change of variables

$$\begin{split} \Delta_{\xi}^{\beta} e^{i\eta \,\theta} \widehat{\Lambda_{\theta}^{\mu_{1}}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi) &= \frac{1}{2\pi} \int_{\mathbb{T}} \left(\Lambda_{\theta'}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}} \right) (\varphi,\theta',\xi) e^{i\eta(\theta-\theta')} d\theta' \\ &= \frac{1}{2\pi} \int_{\mathbb{T}} \left(\Lambda_{\theta}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}} \right) (\varphi,\theta-\theta',\xi) e^{i\eta\theta'} d\theta'. \end{split}$$

Using Bessel identity in the variable θ one deduces

$$\sum_{\eta \in \mathbb{Z}} \left\| \Delta_{\xi}^{\beta} e^{\mathbf{i}\eta \cdot \widehat{\Lambda_{\theta}^{\mu_{1}}} \sigma_{\mathscr{B}}}(\varphi, \eta, \xi) \right\|_{H^{s}_{\theta}}^{2} = 2\pi \left\| \Lambda_{\theta}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}}(\varphi, \centerdot, \xi) \right\|_{H^{s}_{\theta}}^{2}$$

and integrating in φ yields

$$\sum_{\eta \in \mathbb{Z}} \left\| \Delta_{\xi}^{\beta} e^{i\eta} \cdot \widehat{\Lambda_{\theta}^{\mu_{1}}} \sigma_{\mathscr{B}}(\cdot, \eta, \xi) \right\|_{L^{2}_{\varphi}H^{s}_{\theta}}^{2} = 2\pi \left\| \Lambda_{\theta}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}}(\varphi, \cdot, \xi) \right\|_{L^{2}_{\varphi}H^{s}_{\theta}}^{2}.$$

Similarly we get

$$\sum_{\eta \in \mathbb{Z}} \left\| \Delta_{\xi}^{\beta} e^{i\eta} \cdot \widehat{\Lambda_{\theta}^{\mu_{1}}} \sigma_{\mathscr{B}}(\cdot, \eta, \xi) \right\|_{H_{\varphi}^{s} L_{\theta}^{2}}^{2} = 2\pi \left\| \Lambda_{\theta}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}}(\varphi, \cdot, \xi) \right\|_{H_{\varphi}^{s} L_{\theta}^{2}}^{2}.$$

Consequently

$$\begin{split} \sum_{\eta \in \mathbb{Z}} \left\| \Delta_{\xi}^{\beta} e^{\mathrm{i}\eta \cdot \widehat{\Lambda_{\theta}^{\mu_{1}}} \sigma_{\mathscr{B}}}(\cdot, \eta, \xi) \right\|_{H^{s}_{\varphi, \theta}}^{2} \lesssim \left\| \Lambda_{\theta}^{\mu_{1}} \Delta_{\xi}^{\beta} \sigma_{\mathscr{B}}(\varphi, \cdot, \xi) \right\|_{H^{s}_{\varphi, \theta}}^{2} \\ \lesssim \langle \xi \rangle^{2m_{2} - 2\beta} \parallel \mathscr{B} \parallel_{m_{2}, s + \mu_{1}, \beta}^{2} \end{split}$$

Therefore we find

$$\left\|\Delta_{\xi}^{\beta} e^{i\eta} \cdot \widehat{\Lambda_{\theta}^{\mu}} \sigma_{\mathscr{B}}(\cdot,\eta,\xi)\right\|_{H^{s}} \lesssim c_{\eta}(\xi) \langle \xi \rangle^{m_{2}-\beta} \parallel \mathscr{B} \parallel_{m_{2},s+\mu,\beta}$$

with $\sum_{\eta} c_{\eta}^2(\xi) = 1$, for any $\xi \in \mathbb{Z}$. As a consequence,

$$\|\Delta_{\xi}^{\gamma-\beta}e^{\mathbf{i}\eta\cdot\widehat{\Lambda_{\theta}^{\mu}\sigma}}\mathcal{B}(\cdot,\cdot,\xi+\beta)\|_{H^{s}} \lesssim c_{\eta}(\xi)\langle\xi\rangle^{m_{2}-\gamma+\beta} \parallel \mathcal{B} \parallel_{m_{2},s+\mu,\gamma-\beta}.$$
(7.22)

Hence we get in view of Cauchy-Schwarz inequality and by fixing $\mu = \frac{3}{2} + \epsilon, \epsilon > 0$ and for $m_1 \leqslant 1$

$$\begin{split} \left\| \Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{H}}(\cdot, \cdot, \xi) \right\|_{H^{s}} &\lesssim \mathscr{C}_{0} \sum_{|\eta| \leq \frac{|\xi|}{2}} c_{\eta}(\xi) |\eta|^{-\frac{1}{2} - \epsilon} \langle \xi \rangle^{m_{1} + m_{2} - 1 - \gamma} \\ &+ \mathscr{C}_{0} \sum_{\substack{|\eta| \geq 2|\xi| \\ 0 \leq \beta \leq \gamma}} c_{\eta}(\xi) |\eta|^{-\frac{3}{2} - \epsilon} \langle \xi \rangle^{m_{2} - \gamma + \beta} \left(\langle \eta \rangle^{m_{1} - \beta} + \langle \xi \rangle^{m_{1} - \beta} \right) \\ &\lesssim \mathscr{C}_{0} \langle \xi \rangle^{m_{1} + m_{2} - 1 - \gamma}, \end{split}$$
(7.23)

with

$$\mathscr{C}_{0} = \sum_{0 \leqslant \beta \leqslant \gamma} \|\!\|\mathscr{A}\|\!\|_{m_{1},s,1+\beta} \,\|\!|\mathscr{B}\|\!\|_{m_{2},s_{0}+\frac{3}{2}+\epsilon,\gamma-\beta} + \|\!|\!|\mathscr{A}\|\!\|_{m_{1},s_{0},1+\beta} \,\|\!|\mathscr{B}\|\!\|_{m_{2},s+\frac{3}{2}+\epsilon,\gamma-\beta} \,.$$

It remains to investigate the lower interaction term described through

$$\Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{L}}(\varphi,\theta,\xi) = \sum_{\substack{|\xi| < |\eta| < 2|\xi|, \\ 0 \leqslant \beta \leqslant \gamma}} {\binom{\alpha}{\beta}} \left(\Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) - \Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \right) e^{\mathrm{i}\eta \cdot \theta} \Delta_{\xi}^{\gamma-\beta} \widehat{\sigma}_{\mathscr{B}}(\varphi,\eta,\xi+\beta).$$

It is plain that for any family of real numbers $\{\mu_{\beta}, 0 \leq \beta \leq \gamma\}$, we may write

$$\Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{L}}(\varphi,\theta,\xi) = \sum_{\substack{|\underline{\xi}| < |\eta| < 2|\xi|, \\ 0 \leq \beta \leq \gamma}} {\binom{\alpha}{\beta}} \frac{\Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi+\eta) - \Delta_{\xi}^{\beta} \sigma_{\mathscr{A}}(\varphi,\theta,\xi)}{|\eta|^{\mu_{\beta}}} e^{\mathrm{i}\eta \cdot \theta} \Delta_{\xi}^{\gamma-\beta} \widehat{\Lambda_{\theta}^{\mu_{\beta}}} \sigma_{\mathscr{B}}(\varphi,\eta,\xi+\beta).$$

Using the definition (7.9) we infer for $\frac{|\xi|}{2}\leqslant |\eta|\leqslant 2|\xi|$

$$\left\|\Delta_{\xi}^{\beta}\left(\sigma_{\mathscr{A}}(\cdot, \bullet, \xi + \eta) - \sigma_{\mathscr{A}}(\cdot, \bullet, \xi)\right)\right\|_{H^{s}} \lesssim \langle \xi \rangle^{m_{1}^{+}} \parallel \mathscr{A} \parallel_{m_{1}, s, 0}$$

Combining this estimate with (7.22) and fixing the choice

$$\mu_{\beta} = \frac{3}{2} + \beta - m_1^-$$
 with $m_1^- \triangleq \min(m_1, 0),$

we find

$$\begin{split} \|\Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{L}}(\cdot, \cdot, \xi)\|_{H^{s}} &\lesssim \sum_{\substack{|\xi| < |\eta| < 2|\xi| \\ 0 \le \beta \le \gamma}} c_{\eta}(\xi) |\eta|^{-\mu_{\beta}} \langle \xi \rangle^{m_{1}^{+} + m_{2} - \gamma + \beta} \mathscr{C}_{1,\beta} \\ &\lesssim \langle \xi \rangle^{m_{1} + m_{2} - \gamma - 1} \sum_{\substack{|\xi| \\ 0 \le \beta \le \gamma}} c_{\eta}(\xi) |\eta|^{-\frac{1}{2}} \mathscr{C}_{1,\beta} \\ &\lesssim \mathscr{C}_{1} \langle \xi \rangle^{m_{1} + m_{2} - \gamma - 1} \end{split}$$
(7.24)

with

$$\mathscr{C}_{1,\beta} \triangleq \||\mathscr{A}||_{m_1,s,0} |||\mathscr{B}||_{m_2,s_0+\frac{3}{2}+\beta-m_1^-,\gamma-\beta} + |||\mathscr{A}||_{m_1,s_0,0} |||\mathscr{B}||_{m_2,s+\frac{3}{2}+\beta-m_1^-,\gamma-\beta}$$

and

$$\mathscr{C}_1 \triangleq \sum_{0 \leqslant \beta \leqslant \alpha} \mathscr{C}_{1,\beta}$$

Notice that we have used in the last inequality of (7.24) Cauchy-Schwarz inequality as follows

$$\sum_{\substack{|\xi| \\ 2} < |\eta| < 2|\xi|} c_{\eta}(\xi) |\eta|^{-\frac{1}{2}} \leq \left(\sum_{\substack{|\xi| \\ 2} < |\eta| < 2|\xi|} c_{\eta}^{2}(\xi)\right)^{\frac{1}{2}} \left(\sum_{\substack{|\xi| \\ 2} < |\eta| < 2|\xi|} |\eta|^{-1}\right)^{\frac{1}{2}} \leq \left(\sum_{\substack{|\xi| \\ 2} < |\eta| < 2|\xi|} |\eta|^{-1}\right)^{\frac{1}{2}} \leq C$$

with C being independent of η . Combining (7.23) and (7.24) we get for $m_1 \leq 1$ $\|\Delta_{\xi}^{\gamma} \mathcal{R}_1^{\mathrm{L}}(\cdot, \cdot, \xi)\|_{H^s} \lesssim (\mathscr{C}_0 + \mathscr{C}_1) \langle \xi \rangle^{m_1 + m_2 - \gamma - 1}.$

Consequently, we get

$$\||\mathscr{R}_1^{\mathsf{L}}|||_{m_1+m_2-1,s,\gamma} \lesssim \mathscr{C}_0 + \mathscr{C}_1$$

In a similar way to (7.18) one gets, by exchanging \mathcal{A} and \mathcal{B} , the decomposition

$$\sigma_{\mathscr{B}\mathscr{A}}(\varphi,\theta,\xi) \triangleq \mathscr{R}_2(\varphi,\theta,\xi) + \sigma_{\mathscr{A}}(\varphi,\theta,\xi)\sigma_{\mathscr{B}}(\varphi,\theta,\xi)$$

with the estimate, under the assumption $m_2 \leq 1$,

$$\||\mathscr{R}_2|||_{m_1+m_2-1,s,\gamma} \lesssim \mathscr{C}'_0 + \mathscr{C}'_1$$

where

$$\mathscr{C}_{0}^{\prime} \triangleq \sum_{0 \leqslant \beta \leqslant \gamma} \left\| \left| \mathscr{B} \right| \right\|_{m_{2},s,1+\beta} \left\| \left| \mathscr{A} \right| \right\|_{m_{1},s_{0}+\frac{3}{2}+\epsilon,\gamma-\beta} + \left\| \left| \mathscr{B} \right| \right\|_{m_{2},s_{0},1+\beta} \left\| \left| \mathscr{A} \right| \right\|_{m_{1},s+\frac{3}{2}+\epsilon,\gamma-\beta}.$$

and

$$\mathscr{C}_{1}^{\prime} \triangleq \sum_{0 \leqslant \beta \leqslant \gamma} \left(\| \mathscr{B} \| \|_{m_{2},s,0} \| | \mathscr{A} \| \|_{m_{1},s_{0}+\frac{3}{2}+\beta-m_{2}^{-},\gamma-\beta} + \| | \mathscr{B} \| \|_{m_{2},s_{0},0} \| | \mathscr{A} \| \|_{m_{1},s+\frac{3}{2}+\beta-m_{2}^{-},\gamma-\beta} \right).$$

This achieves the desired estimate for the commutator.

(iii) We proceed in a similar way to the preceding point (ii). Putting together (7.22), (7.20) and (7.21) yields (similarly to (7.23)) under the assumptions

$$\mu > \frac{1}{2} + \max\left(0, m_1, m_2, m_1 + m_2\right), \quad m_1 + m_2 \le 1$$
(7.25)

and in view of Cauchy-Schwarz inequality to

$$\begin{split} \left\| \Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{H}}(\cdot, \boldsymbol{\cdot}, \xi) \right\|_{H^{s}} &\lesssim \mathscr{C}_{2} \sum_{|\eta| \leqslant \frac{|\xi|}{2}} c_{\eta}(\xi) |\eta|^{1-\mu} \langle \xi \rangle^{m_{1}+m_{2}-1-\gamma} \\ &+ \mathscr{C}_{2} \sum_{|\eta| \geqslant 2|\xi| \atop 0 \leqslant \beta \leqslant \gamma} c_{\eta}(\xi) |\eta|^{-\mu} \langle \xi \rangle^{m_{2}-\gamma+\beta} \left(\langle \eta \rangle^{m_{1}-\beta} + \langle \xi \rangle^{m_{1}-\beta} \right) \\ &\lesssim \mathscr{C}_{2} \langle \xi \rangle^{-\gamma}, \end{split}$$

with

$$\mathscr{C}_2 = \sum_{0 \leqslant \beta \leqslant \gamma} \|\!|\!\mathcal{A}|\!|\!|_{m_1,s,1+\beta}|\!|\!|\!\mathcal{B}|\!|\!|_{m_2,s_0+\mu,\gamma-\beta} + \|\!|\!|\!\mathcal{A}|\!|\!|_{m_1,s_0,1+\beta}|\!|\!|\!\mathcal{B}|\!|\!|_{m_2,s+\mu,\gamma-\beta}.$$

As to the lower interaction term, we proceed as in (7.24). One gets in view of (7.22), Cauchy-Schwarz inequality and with the choice

$$\mu_{\beta} = m_1^+ + m_2 + \frac{1}{2} + \beta \tag{7.26}$$

that

$$\begin{split} \|\Delta_{\xi}^{\gamma} \mathscr{R}_{1}^{\mathrm{L}}(\cdot, \boldsymbol{\cdot}, \xi)\|_{H^{s}} &\lesssim \sum_{\substack{|\xi| \ 2||| < 2|\xi| \\ 0 \leqslant \beta \leqslant \gamma}} c_{\eta}(\xi) |\eta|^{-\mu_{\beta}} \langle \xi \rangle^{m_{1}^{+} + m_{2} - \gamma + \beta} \mathscr{C}_{3,\beta} \\ &\lesssim \langle \xi \rangle^{-\gamma} \sum_{\substack{|\xi| \ 2|| < 2|\xi| \\ 0 \leqslant \beta \leqslant \gamma}} c_{\eta}(\xi) |\eta|^{-\mu_{\beta} + m_{1} + m_{2} + \beta} \mathscr{C}_{3,\beta} \\ &\lesssim \mathscr{C}_{3} \langle \xi \rangle^{-\gamma} \end{split}$$

with

$$\begin{split} & \mathscr{C}_3 \triangleq \sum_{0 \leqslant \beta \leqslant \gamma} \mathscr{C}_{3,\beta}, \\ & \mathscr{C}_{3,\beta} \triangleq ||| \mathscr{A} |||_{m_1,s,0} ||| \mathscr{B} |||_{m_2,s_0+\mu_\beta,\gamma-\beta} + ||| \mathscr{A} |||_{m_1,s_0,0} ||| \mathscr{B} |||_{m_2,s+\mu_\beta,\gamma-\beta} \,. \end{split}$$

Putting together the preceding estimates we get under the assumptions (7.25) and (7.26)

$$\begin{split} \|\|\mathscr{R}_1\|\|_{0,s,\gamma} \lesssim \mathscr{C}_2 + \mathscr{C}_3 \\ \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathscr{A}\|\|_{m_1,s,1+\gamma-\beta} \|\|\mathscr{B}\|\|_{m_2,s_0+\mu,\beta} + \|\|\mathscr{A}\|\|_{m_1,s_0,1+\gamma-\beta} \||\mathscr{B}\|\|_{m_2,s+\mu,\beta} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \left(\|\|\mathscr{A}\|\|_{m_1,s,0} \|\|\mathscr{B}\|\|_{m_2,s_0+\mu_\beta,\gamma-\beta} + \|\|\mathscr{A}\|\|_{m_1,s_0,0} \||\mathscr{B}\|\|_{m_2,s+\mu_\beta,\gamma-\beta} \right) \end{split}$$

Similarly, we get

$$\sigma_{\mathscr{B}\mathscr{A}}(\varphi,\theta,\xi) = \sum_{\eta\in\mathbb{Z}} e^{i\eta\,\theta} \Big(\sigma_{\mathscr{B}}(\varphi,\theta,\xi+\eta) - \sigma_{\mathscr{B}}(\varphi,\theta,\xi) \Big) \widehat{\sigma}_{\mathscr{A}}(\varphi,\eta,\xi) \\ + \sigma_{\mathscr{B}}(\varphi,\theta,\xi) \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \\ \triangleq \mathscr{R}_{2}(\varphi,\theta,\xi) + \sigma_{\mathscr{A}}(\varphi,\theta,\xi) \sigma_{\mathscr{B}}(\varphi,\theta,\xi)$$

and \mathcal{R}_2 satisfies under the assumptions (7.25) and (7.26) the estimate

$$\begin{split} \||\mathscr{R}_{2}||_{0,s,\gamma} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} |||\mathscr{B}|||_{m_{1},s,1+\gamma-\beta} |||\mathscr{A}|||_{m_{2},s_{0}+\mu,\beta} + |||\mathscr{B}|||_{m_{1},s_{0},1+\alpha-\beta} |||\mathscr{A}|||_{m_{2},s+\mu,\beta} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \left(|||\mathscr{B}|||_{m_{1},s,0} |||\mathscr{A}|||_{m_{2},s_{0}+\overline{\mu}_{\beta},\gamma-\beta} + ||||\mathscr{B}|||_{m_{1},s_{0},0} |||\mathscr{A}|||_{m_{2},s+\overline{\mu}_{\beta},\gamma-\beta} \right) \end{split}$$

with

$$\overline{\mu}_{\beta} = m_1 + m_2^+ + \frac{1}{2} + \beta.$$

Since $\sigma_{[\mathscr{A},\mathscr{B}]} = \sigma_{\mathscr{A}\mathscr{B}} - \sigma_{\mathscr{B}\mathscr{A}} = \mathscr{R}_1 - \mathscr{R}_2$ then

$$\||\boldsymbol{\sigma}_{[\mathscr{A},\mathscr{B}]}||_{0,s,\gamma} \leq |||\mathscr{R}_1|||_{0,s,\gamma} + |||\,\mathscr{R}_2|||_{0,s,\gamma}$$

and the result follows from the preceding estimates.

7.4. Commutators. Our purpose in this section is to establish some commutator estimates between particular pseudo-differential operators of the following type: For a smooth function h we define the operator

$$\mathbb{A}_{\rho} \triangleq \partial_{\theta} \left(\rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \rho \right)$$

where the action of the fractional Laplacian $|D|^{\alpha-1}$ is given by

$$|\mathbf{D}|^{\alpha-1}\rho(\lambda,\varphi,\theta) \triangleq \frac{1}{2\pi} \int_0^{2\pi} \frac{\rho(\lambda,\varphi,\theta-\eta)}{|\sin\left(\frac{\eta}{2}\right)|^{\alpha}} d\eta, \quad \lambda = (\omega,\alpha) \in \mathcal{O}.$$

In this section \mathcal{O} denotes an open set of \mathbb{R}^{d+1} taking the form

$$\mathcal{O} = \mathscr{U} \times (0, \overline{\alpha}) \subset \mathbb{R}^{d+1}, \overline{\alpha} \in (0, \frac{1}{2})$$

with \mathscr{U} is an open bounded set of \mathbb{R}^d . We shall first detail the action of fractional Laplacian $|\mathbf{D}|^{\alpha-1}$ on the spaces $W^{q,\infty}_{\kappa}(\mathcal{O}, H^s(\mathbb{T}^{d+1}))$ introduced in Definition 4.1.

Lemma 7.6. Let $s \in \mathbb{R}, q, \gamma, \ell \in \mathbb{N}, \kappa \in (0, 1)$, then the following assertions hold true.

(i) For any $\epsilon > 0$, there exists C > 0 such that

$$\||\partial_{\theta}^{\ell}|\mathbf{D}|^{\alpha-1}\||_{\overline{\alpha}+\ell+\epsilon-1,s,\gamma}^{q,\kappa} \leqslant C.$$

(ii) For any $\epsilon > 0$, there exists C > 0

$$\||\partial_{\theta}|\mathbf{D}|^{\alpha-1}|\mathbf{D}|^{\alpha-1}\||_{2\overline{\alpha}+\epsilon-1,s,\gamma}^{q,\kappa} \leqslant C.$$

Proof. (i) According to (4.21) and the formula $x\Gamma(x) = \Gamma(1+x)$ one finds

$$|\mathbf{D}|^{\alpha-1}\mathbf{e}_{n}(\theta) = \frac{2^{\alpha}\Gamma(1-\alpha)}{\Gamma(1-\frac{\alpha}{2})\Gamma(\frac{\alpha}{2})} \frac{\Gamma(|n|+\frac{\alpha}{2})}{\Gamma(|n|+1-\frac{\alpha}{2})} \mathbf{e}_{n}(\theta)$$
$$= \frac{2^{\alpha-1}\alpha\Gamma(1-\alpha)}{\Gamma(1-\frac{\alpha}{2})\Gamma(1+\frac{\alpha}{2})} \frac{\Gamma(|n|+\frac{\alpha}{2})}{\Gamma(|n|+1-\frac{\alpha}{2})} \mathbf{e}_{n}(\theta)$$
$$= \mu(\alpha)\mathbb{W}(|n|,\alpha)\mathbf{e}_{n}(\theta),$$
(7.27)

where $\mathbb{W}(n, \alpha)$ was defined in Lemma 3.1-(iii). Then the symbol of $\mathscr{A} \triangleq \partial_{\theta}^{\ell} |\mathbf{D}|^{\alpha-1}$ is given by

$$\sigma_{\mathscr{A}}(\lambda,\varphi,\theta,n) = (\mathrm{i} n)^{\ell} \mu(\alpha) \mathbb{W}(|n|,\alpha)$$

From classical properties of Gamma function we deduce that μ is smooth in α and

$$\sup_{\substack{0 \le k \le j \\ \alpha \in (0,\overline{\alpha})}} |\mu^{(k)}(\alpha)| < \infty.$$

According to Lemma A.1-(ii) one has

$$\forall n, \gamma \in \mathbb{N}, \quad \max_{\substack{k \in [0,q]\\\alpha \in (0,\overline{\alpha})}} |\partial_{\alpha}^{k} \Delta_{n}^{\gamma} \mathbb{W}(n,\alpha)| \leqslant C(\gamma,q,\epsilon) \langle n \rangle^{\overline{\alpha}+\epsilon-1-\gamma}.$$

Therefore, according to Leibniz rule combined with the preceding estimates we obtain

$$\forall n, \gamma \in \mathbb{N}, \qquad \max_{\substack{k \in [0,q]\\\alpha \in (0,\overline{\alpha})}} \langle n \rangle^{1-\ell-\overline{\alpha}-\epsilon+\gamma} \left| \partial_{\alpha}^{k} \Delta_{n}^{\gamma} \sigma_{\mathscr{A}}(\lambda,\varphi,\theta,n) \right| \leqslant C(\gamma,q,\ell,\epsilon).$$
(7.28)

This implies in view of (7.13)

$$\||\partial_{\theta}^{\ell}|\mathbf{D}|^{\alpha-1}\||_{\overline{\alpha}+\ell+\epsilon-1,s,\gamma}^{q,\kappa} \leqslant C(\gamma,q,\ell,\epsilon),$$

which ensures the desired result. We point out that for q = 0 we have no the loss ϵ . More precisely,

$$\forall \alpha \in (0,\overline{\alpha}), \quad |||\partial_{\theta}^{\ell}|\mathbf{D}|^{\alpha-1}|||_{\alpha+\ell-1,s,\gamma} \leqslant C(\gamma,\ell,\epsilon).$$
(7.29)

(ii) Iterating (7.27) we obtain

$$\partial_{\theta} |\mathbf{D}|^{\alpha-1} |\mathbf{D}|^{\alpha-1} \mathbf{e}_n = \mathrm{i} n \mu^2(\alpha) \, \mathbf{W}^2(|n|, \alpha) \, \mathbf{e}_n.$$

Similarly to (7.28) we deduce that for any $\epsilon > 0$,

$$\forall \, n, \gamma \in \mathbb{N}, \qquad \max_{\substack{k \in [0,q] \\ \alpha \in (0,\overline{\alpha})}} \langle n \rangle^{1-2\overline{\alpha}-\epsilon+\gamma} \big| \partial_{\alpha}^k \Delta_n^{\gamma} \big(n \mu^2(\alpha) \, \mathtt{W}^2(|n|,\alpha) \big| \leqslant C(\gamma,q,\epsilon).$$

It follows from (7.13) that

$$\||\partial_{\theta}|\mathbf{D}|^{\alpha-1}|\mathbf{D}|^{\alpha-1}\||_{2\overline{\alpha}+\epsilon-1,s,\gamma}^{q,\kappa} \leqslant C(\gamma,q,\epsilon),$$

which achieves the proof of Lemma 7.6.

As an application, we shall prove the following result.

Lemma 7.7. Let $\epsilon > 0$ be small enough, $\overline{\alpha} \in (0, \frac{1}{2})$ and (κ, d, q, s, s_0) as in (4.1). Then the following assertions hold true.

(i) For any $\alpha \in (0, \overline{\alpha})$, there exists C > 0 such that

$$\|\mathbb{A}_{\rho}\|_{\alpha,s,\gamma} \leqslant C \|\rho\|_{s+\gamma+1}.$$

(ii) There exists C > 0 such that

$$\||\mathbb{A}_{\rho}\||_{\overline{\alpha}+\epsilon,s,\gamma}^{q,\kappa} \leqslant C \|\rho\|_{s+\gamma+1}^{q,\kappa}.$$

(iii) There exists C > 0 such that

$$\|[\mathbb{A}_{\rho_1}, \mathbb{A}_{\rho_2}]\|_{2\overline{\alpha}-1+\epsilon, s, \gamma}^{q,\kappa} \leq C(\|\rho_1\|_{s_0+3}^{q,\kappa} + \|\rho_2\|_{s_0+3}^{q,\kappa})(\|\rho_1\|_{s+\gamma+3}^{q,\kappa} + \|\rho_2\|_{s+\gamma+3}^{q,\kappa}).$$

(iv) Let \mathscr{A} as in (7.11) and $m \in [-1, 0]$. Then for any $s \ge s_0$ there exits C > 0

$$\| \| [\mathscr{A}, \mathbb{A}_{\rho}] \| \|_{m, s, \gamma}^{q, \kappa} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \| \| \mathscr{A} \| \|_{m, s_{0}+2, 1+\beta}^{q, \kappa} \| \rho \|_{s+4+\gamma-\beta}^{q, \kappa} + \| |\mathscr{A} \| \|_{m, s+2, 1+\beta}^{q, \kappa} \rho \|_{s_{0}+4+\gamma-\beta}^{q, \kappa} + \sum_{0 \leqslant \beta \leqslant \gamma} \| \| \mathscr{A} \| \|_{m, s_{0}+2+\beta, \gamma-\beta}^{q, \kappa} \| \rho \|_{s+1}^{q, \kappa} + \| \| \mathscr{A} \| \|_{m, s+2+\beta, \gamma-\beta}^{q, \kappa} \| \rho \|_{s_{0}+1}^{q, \kappa}.$$

(v) Let
$$\overline{s}_0 > \frac{d+5}{2}, s \ge 0, \ \rho \in H^{s+\overline{s}_0}$$
 and $h \in H^s \cap H^{\overline{s}_0}$. Then
 $\|\partial_{\theta} [\rho, \Lambda^s] h\|_{L^2(\mathbb{T}^{d+1})} \lesssim \|h\|_{H^s(\mathbb{T}^{d+1})} \|\rho\|_{H^{\overline{s}_0}(\mathbb{T}^{d+1})} + \|h\|_{H^{\overline{s}_0}(\mathbb{T}^{d+1})} \|\rho\|_{H^{s+\overline{s}_0}(\mathbb{T}^{d+1})}.$
73

Proof. (i) We write

$$\mathbb{A}_{\rho} = \rho \partial_{\theta} |\mathbf{D}|^{\alpha - 1} + (\partial_{\theta} \rho) |\mathbf{D}|^{\alpha - 1} + (\partial_{\theta} |\mathbf{D}|^{\alpha - 1}) \rho$$
$$\triangleq \mathbb{A}_{\rho, 1} + \mathbb{A}_{\rho, 2} + \mathbb{A}_{\rho, 3}.$$

Denote by T_{ρ} the multiplicative operator defined by $T_{\rho}h = \rho h$ then $\sigma_{T_{\rho}} = \rho$ and therefore

$$|||T_{\rho}|||_{0,s,\gamma}^{q,\kappa} \lesssim ||\rho||_{s}^{q,\kappa}$$

Applying Lemma 7.5-(i) combined with (7.29) we infer for any $s \ge s_0$

$$\|\|\mathbb{A}_{\rho,1}\|\|_{\alpha,s,\gamma} \lesssim \|\rho\|_s$$

Similarly we get

$$\begin{aligned} \|\|\mathbb{A}_{\rho,2}\|\|_{\alpha,s,\gamma} &\lesssim \|\|T_{\partial_{\theta}\rho}\|\|_{0,s,\gamma} \\ &\lesssim \|\rho\|_{s+1}. \end{aligned}$$

For the last term we use once again Lemma 7.5-(i) combined with (7.29) and the fact that $\alpha \in (0, \frac{1}{2})$, in order to get for any $s \ge s_0$

$$\begin{split} \|\|\mathbb{A}_{\rho,3}\|\|_{\alpha,s,\gamma} \lesssim \|\rho\|_{s+\frac{1}{2}+\alpha} + \|\rho\|_{s+\frac{1}{2}+\gamma} \\ \lesssim \|\rho\|_{s+\gamma+1}. \end{split}$$

This achieves the proof of the first point.

(ii) With the notations of the first point one has according to Lemma 7.5-(i) combined with Lemma 7.6 we infer for any $s \ge s_0$

$$\forall s \ge s_0, \quad |||\mathbb{A}_{\rho,1}|||^{q,\kappa}_{\overline{\alpha}+\epsilon,s,\gamma} \lesssim ||\rho||^{q,\kappa}_s.$$

Similar arguments yield

$$\|\|\mathbb{A}_{\rho,2}\|\|_{\overline{\alpha}+\epsilon,s,\gamma}^{q,\kappa} \lesssim \|\|T_{\partial_{\theta}\rho}\|\|_{0,s,\gamma}^{q,\kappa} \\ \lesssim \|\rho\|_{s+1}^{q,\kappa}.$$

For the last term we implement once again Lemma 7.5-(i) combined with Lemma 7.6 and the fact that $\overline{\alpha} \in (0, \frac{1}{2})$, in order to get for any $s \ge s_0$

$$\begin{split} \|\|\mathbb{A}_{\rho,3}\|\|_{\overline{\alpha}+\epsilon,s,\gamma}^{q,\kappa} \lesssim \|\rho\|_{s+\frac{1}{2}+\overline{\alpha}+\epsilon}^{q,\kappa} + \|\rho\|_{s+\frac{1}{2}+\gamma}^{q,\kappa} \\ \lesssim \|\rho\|_{s+\gamma+1}^{q,\kappa}, \end{split}$$

where ϵ is taken small enough. Putting together the preceding estimates gives

$$||| \mathbb{A}_{\rho} |||_{\overline{\alpha}+\epsilon,s,\gamma}^{q,\kappa} \leqslant C ||\rho||_{s+1+\gamma}^{q,\kappa}, \tag{7.30}$$

which completes the proof of the desired result.

(iii) Applying Lemma 7.5-(ii) with $m_1 = m_2 = \overline{\alpha} + \frac{\epsilon}{2}$ and using the estimate (7.30) combined with Sobolev embeddings

$$\begin{split} \|\|[\mathbb{A}_{\rho_{1}},\mathbb{A}_{\rho_{2}}]\|\|_{2\overline{\alpha}-1+\epsilon,s,\gamma}^{q,\kappa} &\lesssim \sum_{0\leqslant\beta\leqslant\gamma} \|\rho_{1}\|_{s+2+\beta}^{q,\kappa} \|\rho_{2}\|_{s_{0}+3+\gamma-\beta}^{q,\kappa} + \|\rho_{1}\|_{s_{0}+2+\beta}^{q,\kappa} \|\rho_{2}\|_{s+3+\gamma-\beta}^{q,\kappa} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\rho_{1}\|_{s+1}^{q,\kappa} \|h_{2}\|_{s_{0}+3+\gamma}^{q,\kappa} + \|h_{1}\|_{s_{0}+1}^{q,\kappa} \|\rho_{2}\|_{s+3+\gamma}^{q,\kappa} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\rho_{2}\|_{s+2+\beta}^{q,\kappa} \|\rho_{1}\|_{s_{0}+3+\gamma-\beta}^{q,\kappa} + \|\rho_{2}\|_{s_{0}+2+\beta}^{q,\kappa} \|\rho_{1}\|_{s+3+\gamma-\beta}^{q,\kappa} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\rho_{2}\|_{s+1}^{q,\kappa} \|\rho_{1}\|_{s_{0}+3+\gamma}^{q,\kappa} + \|\rho_{2}\|_{s_{0}+1}^{q,\kappa} \|\rho_{1}\|_{s+3+\gamma}^{q,\kappa}. \end{split}$$

Hence we find from Sobolev embeddings

$$\|\|[\mathbb{A}_{\rho_1}, \mathbb{A}_{\rho_2}]\|\|_{2\overline{\alpha}-1+\epsilon, s, \gamma}^{q, \kappa} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \left(\|\rho_1\|_{s+3+\beta}^{q, \kappa} \|\rho_2\|_{s_0+3+\gamma-\beta}^{q, \kappa} + \|\rho_1\|_{s_0+3+\beta}^{q, \kappa} \|\rho_2\|_{s+3+\gamma-\beta}^{q, \kappa} \right).$$
(7.31)

Applying interpolation inequality we find

$$\|\rho_1\|_{s+3+\beta}^{q,\kappa} \|\rho_2\|_{s_0+3+\gamma-\beta}^{q,\kappa} \lesssim \left(\|\rho_1\|_{s_0+3}^{q,\kappa} + \|\rho_2\|_{s_0+3}^{q,\kappa}\right) \left(\|\rho_1\|_{s+3+\gamma}^{q,\kappa} + \|\rho_2\|_{s+3+\gamma}^{q,\kappa}\right)$$

which implies by virtue of (7.31)

$$\|[\mathbb{A}_{\rho_1}, \mathbb{A}_{\rho_2}]\|_{2\overline{\alpha}-1+\epsilon, s, \gamma}^{q, \kappa} \lesssim \left(\|\rho_1\|_{s_0+3}^{q, \kappa} + \|\rho_2\|_{s_0+3}^{q, \kappa}\right) \left(\|\rho_1\|_{s+3+\gamma}^{q, \kappa} + \|\rho_2\|_{s+3+\gamma}^{q, \kappa}\right).$$

This ends the proof of the third point.

 $({\bf iv})$ First we write according to the definition and since $\overline{\alpha}+\epsilon-1\leqslant 0$

$$\|\|[\mathscr{A},\mathbb{A}_h]\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\|[\mathscr{A},\mathscr{U}_h]\|\|_{m+\overline{\alpha}+\epsilon-1,s,\gamma}^{q,\kappa}$$

Then applying Lemma 7.5-(ii) combined with (7.30) and Sobolev embeddings

$$\begin{split} \|[\mathscr{A}, \mathbb{A}_{\rho}]\|\|_{m+\overline{\alpha}+\epsilon-1, s, \gamma}^{q,\kappa} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathscr{A}\|\|_{m, s, 1+\beta}^{q,\kappa} \|\rho\|_{s_{0}+3+\gamma-\beta}^{q,\kappa} + \||\mathscr{A}\|\|_{m, s_{0}, 1+\beta}^{q,\kappa} \|\rho\|_{s+3+\gamma-\beta}^{q,\kappa} \\ &+ \||\mathscr{A}\|\|_{m, s, 0}^{q,\kappa} \|\rho\|_{s_{0}+4+\gamma}^{q,\kappa} + \||\mathscr{A}\|\|_{m, s_{0}, 0}^{q,\kappa} \|\rho\|_{s+4+\gamma}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \||\mathscr{A}\|\|_{m, s_{0}+2, \beta}^{q,\kappa} \|\rho\|_{s+2+\gamma-\beta}^{q,\kappa} + \||\mathscr{A}\|\|_{m, s+2, \beta}^{q,\kappa} \|\rho\|_{s_{0}+2+\gamma-\beta}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \||\mathscr{A}\|\|_{m, s_{0}+2+\beta, \gamma-\beta}^{q,\kappa} \|\rho\|_{s+1}^{q,\kappa} + \||\mathscr{A}\|\|_{m, s+2+\beta, \gamma-\beta}^{q,\kappa} \|\rho\|_{s_{0}+1}^{q,\kappa}. \end{split}$$

Applying Sobolev embeddings we infer

$$\begin{split} \| [\mathscr{A}, \mathbb{A}_{\rho}] \|_{m,s,\gamma}^{q,\kappa} &\lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \| \|\mathscr{A}\| \|_{m,s_{0}+2,1+\beta}^{q,\kappa} \| \rho \|_{s+4+\gamma-\beta}^{q,\kappa} + \| |\mathscr{A}| \|_{m,s+2,1+\beta}^{q,\kappa} \| \rho \|_{s_{0}+4+\gamma-\beta}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \| \|\mathscr{A}\| \|_{m,s_{0}+2+\beta,\gamma-\beta}^{q,\kappa} \| \rho \|_{s+1}^{q,\kappa} + \| \|\mathscr{A}\| \|_{m,s+2+\beta,\gamma-\beta}^{q,\kappa} \| \rho \|_{s_{0}+1}^{q,\kappa}. \end{split}$$

This proves the desired result.

(v) Denote $f \triangleq \partial_{\theta} [\rho, \Lambda^s] h$, then its Fourier expansion writes

$$f = \sum_{(l,j)\in\mathbb{Z}^{d+1}} c_{l,j}(f)\mathbf{e}_{l,j}, \quad \mathbf{e}_{l,j}(\varphi,\theta) = e^{\mathrm{i}(l\cdot\varphi+j\theta)}$$

with

$$c_{l,j}(f) = ij \sum_{(l',j') \in \mathbb{Z}^{d+1}} \mu_{l',j'}^{l,j} c_{l',j'}(h) c_{l-l',j-j'}(\rho)$$

and

$$\mu_{l',j'}^{l,j} = \langle l', j' \rangle^s - \langle l, j \rangle^s.$$

Using the classical inequality

$$\forall a, b > 0, \quad \left| a^s - b^s \right| \lesssim_s |a - b| \left(a^{s-1} + |a - b|^{s-1} \right)$$
(7.32)

with $a = \langle l', j' \rangle$ and $b = \langle l, j \rangle$ yields from the triangle inequality

$$\begin{split} \left| \langle l', j' \rangle^{s_1} - \langle l, j \rangle^{s_1} \right| &\lesssim \langle l - l', j - j' \rangle \Big(\langle l', j' \rangle^{s-1} + \langle l - l', j - j' \rangle^{s-1} \Big) \\ &\lesssim \langle l - l', j - j' \rangle \langle l', j' \rangle^{s-1} + \langle l - l', j - j' \rangle^s. \end{split}$$

It follows that

$$|\mu_{l',j'}^{l,j}| \lesssim \langle l-l', j-j' \rangle \langle l', j' \rangle^{s-1} + \langle l-l', j-j' \rangle^s.$$

Combining this inequality with $|j| \leq |j - j'| + |j'|$ and $|j'| \leq \langle l', j' \rangle$ allows to get for $s \ge 0$

$$|j||\mu_{l',j'}^{l,j}| \lesssim \langle l-l',j-j'\rangle^2 \langle l',j'\rangle^{s-1} + \langle l-l',j-j'\rangle^{s+1} + \langle l-l',j-j'\rangle \langle l',j'\rangle^s + \langle l-l',j-j'\rangle^s \langle l',j'\rangle.$$

$$(7.33)$$

Using the convolution laws allows to get

$$\begin{split} \|f\|_{L^{2}} \lesssim \|h\|_{H^{s-1}} \sum_{l,j} \langle l,j \rangle^{2} |c_{l,j}(\rho)| + \|h\|_{L^{2}} \sum_{l,j} \langle l,j \rangle^{s+1} |c_{l,j}(\rho)| \\ + \|h\|_{H^{s}} \sum_{l,j} \langle l,j \rangle |c_{l,j}(\rho) + \|h\|_{H^{1}} \sum_{l,j} \langle l,j \rangle^{s} |c_{l,j}(\rho)|. \end{split}$$

Then using Cauchy-Schwarz inequality leads for any $\epsilon > 0$

$$\begin{split} \|f\|_{L^{2}} \lesssim \|h\|_{H^{s-1}} \|\rho\|_{H^{\frac{d+5}{2}+\epsilon}(\mathbb{T}^{d+1})} + \|h\|_{L^{2}} \|\rho\|_{H^{s+\frac{d+3}{2}+\epsilon}} \\ &+ \|h\|_{H^{s}} \|\rho\|_{H^{\frac{d+3}{2}+\epsilon}(\mathbb{T}^{d+1})} + \|h\|_{H^{1}} \|\rho\|_{H^{s+\frac{d+1}{2}+\epsilon}}. \end{split}$$

To get the desired inequality, it is enough to use Sobolev embeddings and this achieves the proof of Lemma 7.7. $\hfill \Box$

8. New Approach related to Egorov Theorem

This section is devoted to the construction of hyperbolic flows in infinite dimensional spaces associated to nonlocal pseudo-differential operators and explore the conjugation of pseudo-differential operators with this flow. More precisely, we are concerned with solving the evolution equation

$$\begin{cases} \partial_t \Phi = \partial_\theta (\rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \rho) \Phi(t) \triangleq \partial_\theta \mathcal{T} \Phi(t) \triangleq \mathbb{A} \Phi(t), \\ \Phi(0) = \mathrm{Id}, \end{cases}$$
(8.1)

where $|D|^{\alpha-1}$ is the modified fractional Laplacian introduced in (4.19) and $\rho : \mathcal{O} \times \mathbb{T}^{d+1} \to \mathbb{R}$ is a smooth function that enjoys the following symmetry,

$$\forall (\lambda, \varphi, \theta) \in \mathcal{O} \times \mathbb{T}^{d+1}, \quad \rho(\lambda, -\varphi, -\theta) = -\rho(\lambda, \varphi, \theta).$$

We shall discuss throughout Section 8.1 some basic properties of the flow such as its continuity over Sobolev spaces or its Lipschitz dependence with respect to ρ . We point out that this flow will be used later in Proposition 9.4 in order to reduce the nonlocal part of the linearized operator to a Fourier multiplier, and this requirement will definitively fix the function ρ . Afterwards, we shall investigate in Section 8.3 the conjugation of pseudo-differential operators by these hyperbolic flows. In particular, if we take an operator \mathscr{A} of order m satisfying (7.13) and we conjugate it with the flow $\Phi(t)$, that is to consider $\mathscr{A}_t \triangleq \Phi^{-1}(t) \mathscr{A} \Phi(t)$, then from Egorov theorem \mathscr{A}_t remains a pseudo-differential operator with order m. The delicate point is to estimate \mathscr{A}_t with respect to the topology given by (7.13) with suitable tame estimates. The main result in this direction will be explained in Theorem 8.1. This theorem, which is interesting in itself, is of great importance and will be used in Section 9.6.3 during the KAM scheme implemented to reduce the remainder to a diagonal part. There, we should check that the new remainders obtained through the different transformations (transport and the nonlocal reductions) give rise to good operators of order zero and enjoying suitable tame estimates in the topology (7.14). Then the required estimates will stem from Theorem 8.1.

8.1. Nonlocal pseudo-differential hyperbolic equation. In this section we shall be concerned with the continuity of the hyperbolic flows generated by the nonlocal pseudo-differential equation (8.1). At this level, we take for granted the existence of such flows and restrict the discussion to the a priori estimates. Notice that the flow construction, which is omitted here, can be done in a classical way by mollifying the equation and getting the suitable a priori estimates combined with compactness arguments, see for instance [16].

Proposition 8.1. Let $d \in \mathbb{N}^*, q \in \mathbb{N}$ and $s_0 > \frac{d+5}{2}$, then the followings assertions hold true.

(i) Assume that $\|\rho\|_{H^{2s_0}} \lesssim 1$, then for any $|t| \leqslant 1$ and $s \ge 0$

 $\|\Phi(t)h\|_{H^s} + \|\Phi^{\star}(t)h\|_{H^s} \lesssim \|h\|_{H^s} + \|\rho\|_{H^{s+s_0}} \|h\|_{H^{s_0}}.$

Moreover the flow map is reversibility preserving in the sense of Definition 7.1.

(ii) If $\|\rho\|_{2s_0+q+1}^{q,\kappa} \lesssim 1$, then for any $|t| \leqslant 1$ and $s \ge q$

$$\|\Phi(t)h\|_{s}^{q,\kappa} + \|\Phi^{\star}(t)h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{q,\kappa}\|h\|_{s_{0}+q}^{q,\kappa}.$$

Moreover, we have

$$\|(\Phi(t) - \mathrm{Id})h\|_{s}^{q,\kappa} + \|(\Phi^{\star}(t) - \mathrm{Id})h\|_{s}^{q,\kappa} \lesssim \|h\|_{s+1}^{q,\kappa} \|\rho\|_{s_{0}}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{q,\kappa} \|h\|_{s_{0}+1+q}^{q,\kappa}.$$

(iii) Assume that $\max_{i=1,2} \|\rho_i\|_{2s_0+1+q}^{q,\kappa} \lesssim 1$, then for any $s \ge \max(s_0,q)$ and $|t| \le 1$

$$\|\Delta_{12}\Phi(t)h\|_{s}^{q,\kappa} + \|\Delta_{12}\Phi^{\star}(t)h\|_{s}^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s+1}^{q,\kappa} \|h\|_{s+1+q}^{q,\kappa} \Big(1 + \|\rho_{1}\|_{s+s_{0}+1}^{q,\kappa} + \|\rho_{2}\|_{s+s_{0}+1}^{q,\kappa}\Big).$$

Proof. (i) As we have mentioned before, we shall skip the details about the existence of the flow map and just focus on the a priori estimates. Let h be a smooth function and denote by $h_t \triangleq \Phi(t)h$ the unique solution of the pseudo-differential equation

$$\begin{cases} \partial_t h_t = \partial_\theta \mathcal{T}(\varphi, \theta) h_t, \\ h(0) = h. \end{cases}$$
(8.2)

where \mathcal{T} is the self-adjoint operator $\mathcal{T} \triangleq \rho |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1}\rho$. We shall first start with the case s = 0. Using integration by parts and the fact that \mathcal{T} is self-adjoint we get

$$\frac{1}{2}\frac{d}{dt}\|h_t\|_{L^2}^2 = \int_{\mathbb{T}^{d+1}} h_t(\varphi,\theta)\partial_\theta \mathcal{T}h_t(\varphi,\theta)d\varphi d\theta$$
$$= -\int_{\mathbb{T}^{d+1}} h_t(\varphi,\theta)\mathcal{T}\partial_\theta h_t(\varphi,\theta)d\varphi d\theta$$
$$\triangleq I_t.$$

Thus by summation,

$$I_{t} = \frac{1}{2} \int_{\mathbb{T}^{d+1}} h_{t} [\partial_{\theta}, \mathcal{T}] h_{t} d\varphi d\theta$$

$$= \frac{1}{2} \int_{\mathbb{T}^{d+1}} h_{t} (\partial_{\theta} \rho) |\mathbf{D}|^{\alpha - 1} h_{t} d\varphi d\theta + \frac{1}{2} \int_{\mathbb{T}^{d+1}} h_{t} |\mathbf{D}|^{\alpha - 1} ((\partial_{\theta} \rho) h_{t}) d\varphi d\theta.$$

Using Lemma 4.8 we get for $\alpha \in [0, 1)$,

$$|||\mathbf{D}|^{\alpha-1}h_t||_{L^2} \lesssim ||h_t||_{L^2}$$
 and $|||\mathbf{D}|^{\alpha-1}(\partial_{\theta}\rho)h_t||_{L^2} \lesssim ||\partial_{\theta}\rho||_{L^{\infty}}||h_t||_{L^2}.$

Consequently

 $|I_t| \lesssim \|\partial_{\theta}\rho\|_{L^{\infty}} \|h_t\|_{L^2}^2.$

Therefore we get after simplification

$$\frac{d}{dt} \|h_t\|_{L^2} \lesssim \|\partial_\theta \rho\|_{L^\infty} \|h_t\|_{L^2}.$$
(8.3)

Hence we infer from Gronwal inequality

$$\sup_{t \in [-1,1]} \|h_t\|_{L^2} \leqslant \|h\|_{L^2} e^{C \|\partial_\theta \rho\|_{L^\infty}}.$$
(8.4)

To get the desired estimate it is enough to make appeal to Sobolev embeddings. Let us now move to the estimate of $||h_t||_{H^s}$ for $s \ge 0$. Applying the Fourier multiplier Λ^s to (8.2) yields

$$\begin{cases} \partial_t \Lambda^s h_t = \partial_\theta \mathcal{T} \Lambda^s h_t + \left\lfloor \Lambda^s, \partial_\theta \mathcal{T} \right\rfloor h_t \\ h(0) = h \in H^s(\mathbb{T}^{d+1}). \end{cases}$$

Proceeding as before for s = 0 we get similarly to (8.3)

$$\frac{d}{dt} \|\Lambda^s h_t\|_{L^2} \lesssim \|\partial_\theta \rho\|_{L^\infty} \|\Lambda^s h_t\|_{L^2} + \|\left[\Lambda^s, \partial_\theta \mathcal{T}\right] h_t\|_{L^2}.$$
(8.5)

It remains to estimate the commutator which can be written in the form

$$\begin{bmatrix} \Lambda^s, \partial_\theta \mathcal{T} \end{bmatrix} = \partial_\theta \begin{bmatrix} \Lambda^s, \rho \end{bmatrix} |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \partial_\theta \begin{bmatrix} \Lambda^s, \rho \end{bmatrix}$$

Combining Lemma 7.7-(v) with Lemma 4.8 we infer for any $s_0 > \frac{d+5}{2}$

$$\begin{aligned} \partial_{\theta} \big[\Lambda^{s}, \rho \big] |\mathbf{D}|^{\alpha - 1} h_{t} \big\|_{L^{2}} &\lesssim \||\mathbf{D}|^{\alpha - 1} h\|_{H^{s}} \|\rho\|_{H^{s_{0}}} + \||\mathbf{D}|^{\alpha - 1} h\|_{H^{s_{0}}} \|\rho\|_{H^{s + s_{0}}} \\ &\lesssim \|h\|_{H^{s}} \|\rho\|_{H^{s_{0}}} + \|h\|_{H^{s_{0}}} \|\rho\|_{H^{s + s_{0}}} \end{aligned}$$

and in a similar way

 $\left\| |\mathbf{D}|^{\alpha-1} \partial_{\theta} [\Lambda^{s}, \rho] h_{t} \right\|_{L^{2}} \lesssim \|h\|_{H^{s}} \|\rho\|_{H^{s_{0}}} + \|h\|_{H^{s_{0}}} \|\rho\|_{H^{s+s_{0}}}.$

Putting together the preceding estimates we obtain

$$\left\| \left[\Lambda^{s}, \partial_{\theta} \mathcal{T} \right] h_{t} \right\|_{L^{2}} \lesssim \|h_{t}\|_{H^{s}} \|\rho\|_{H^{s_{0}}} + \|h_{t}\|_{H^{s_{0}}} \|\rho\|_{H^{s+s_{0}}}.$$
(8.6)

Inserting (8.6) into (8.5) we obtain from Sobolev embeddings

$$\frac{d}{dt} \|h_t\|_{H^s} \lesssim \|\partial_\theta \rho\|_{L^\infty} \|h_t\|_{H^s} + \|h_t\|_{H^s} \|\rho\|_{H^{s_0}} + \|h_t\|_{H^{s_0}} \|\rho\|_{H^{s+s_0}} \\ \lesssim \|h_t\|_{H^s} \|\rho\|_{H^{s_0}} + \|h_t\|_{H^{s_0}} \|\rho\|_{H^{s+s_0}}.$$
(8.7)

By taking $s = s_0$, and applying Gronwall Lemma we find for $t \in \mathbb{R}$

$$\|h_t\|_{H^{s_0}} \leqslant e^{C|t|\|\rho\|_{H^{2s_0}}} \|h\|_{H^{s_0}}.$$
(8.8)

Applying once again Gronwall lemma to (8.7) and using (8.8) yields

$$\begin{aligned} \|h_t\|_{H^s} &\leqslant e^{C|t|\|\rho\|_{H^{s_0}}} \left(\|h\|_{H^s} + \|\rho\|_{H^{s+s_0}} \left| \int_0^t \|h_{t'}\|_{H^{s_0}} dt' \right| \right) \\ &\leqslant e^{C|t|\|\rho\|_{H^{2s_0}}} \left(\|h\|_{H^s} + |t|\|\rho\|_{H^{s+s_0}} \|h\|_{H^{s_0}} \right). \end{aligned}$$

This can be written in the form

$$\|\Phi(t)h\|_{H^{s}} \leqslant e^{C|t|\|\rho\|_{H^{2s_{0}}}} \left(\|h\|_{H^{s}} + |t|\|\rho\|_{H^{s+s_{0}}} \|h\|_{H^{s_{0}}} \right).$$

$$(8.9)$$

This gives the desired statement when $\|\rho\|_{H^{2s_0}} \lesssim 1$ and $|t| \leq 1$. Next we shall estimate the adjoint operator $\Phi^{\star}(t)$. We can check that $\Phi^{\star}(t)$ satisfies the equation

$$\begin{cases} \partial_t \Phi^{\star}(t) = \Phi^{\star}(t) \mathbb{A}^* \\ \Phi(0) = \mathrm{Id.} \end{cases}$$

Since $\Phi(t)$ commutes with its generator A, that is, $\Phi(t)A = A\Phi(t)$ and then taking the adjoint yields $\Phi^{\star}(t)\mathbb{A}^{\star} = \mathbb{A}^{\star}\Phi^{\star}(t)$. Therefore from the identity $\mathbb{A}^{\star} = -\mathcal{T}\partial_{\theta}$ we find

$$\begin{cases} \partial_t \Phi^*(t) = -(\rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1}\rho)\partial_\theta \Phi^*(t), \\ \Phi(0) = \mathrm{Id.} \end{cases}$$
(8.10)

It follows that the equation of $\Phi^{\star}(t)$ is similar to $\Phi(t)$ and its generator enjoys the same properties as the generator \mathbb{A} . Therefore, following the same steps one gets at the end the same estimate (8.9) for the adjoint operator $\Phi^{*}(t)$. The reversibility preserving can be easily checked from the symmetry of ρ and the uniqueness of the Cauchy problem.

(ii) First, observe that the estimate (8.9) gives the desired result with q = 0. Now, we need to establish a similar estimate for $\partial_{\lambda}^{\beta} \Phi(t)h$, with $\lambda = (\omega, \alpha)$ and $|\beta| \leq q$. We point out that the delicate manipulation concerns the estimates when we want to differentiate with respect to α because the order of the generator \mathcal{T} depends on α and not on ω . Therefore, we shall restrict the discussion to this case and estimate $\partial_{\alpha}^{j} \Phi(t)h$ for $1 \leq j \leq q$. To start, we differentiate (8.2) according to Leibniz formula

$$\partial_t \partial^j_\alpha h_t = \partial_\theta \mathcal{T}(\varphi, \theta) \partial^j_\alpha h_t + \sum_{k=0}^{j-1} \binom{j}{k} \partial_\theta (\partial^{j-k}_\alpha \mathcal{T}) \partial^k_\alpha h_t.$$
(8.11)

Then similarly to (8.5) and (8.6) we deduce for $s \ge q$

$$\frac{d}{dt} \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} \lesssim \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} \|\rho\|_{H^{s_{0}}(\mathbb{T}^{d+1})} + \|\partial_{\alpha}^{j} h_{t}\|_{H^{s_{0}}} \|\rho\|_{H^{s-j+s_{0}}}$$

$$+\sum_{k=0}^{j-1} \left\| \partial_{\theta} (\partial_{\alpha}^{j-k} \mathcal{T}) \partial_{\alpha}^{k} h_{t} \right\|_{H^{s-j}}.$$
(8.12)

Using once again Leibniz formula we may write

$$\left(\partial_{\alpha}^{j-k}\mathcal{T}\right) = \sum_{m=0}^{j-k} {\binom{j-k}{m}} \left((\partial_{\alpha}^{m} \rho) (\partial_{\alpha}^{j-k-m} |\mathbf{D}|^{\alpha-1}) + (\partial_{\alpha}^{j-k-m} |\mathbf{D}|^{\alpha-1}) (\partial_{\alpha}^{m} \rho) \right).$$
(8.13)

Using Lemma 4.1 and Lemma 4.8 we get

$$\begin{aligned} \left\| \partial_{\theta} ((\partial_{\alpha}^{m} \rho) (\partial_{\alpha}^{j-k-m} |\mathbf{D}|^{\alpha-1})g) \right\|_{H^{s-j}} &\lesssim \left\| (\partial_{\alpha}^{m} \rho) (\partial_{\alpha}^{j-k-m} |\mathbf{D}|^{\alpha-1})g \right\|_{H^{s-j+1}} \\ &\lesssim \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s_0}} \|g\|_{H^{s-j+1}} + \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s-j+1}} \|g\|_{H^{s_0}}. \end{aligned}$$

$$\tag{8.14}$$

Similarly we obtain

$$\begin{aligned} \left\| \partial_{\theta} (\partial_{\alpha}^{j-k-m} |\mathbf{D}|^{\alpha-1}) (\partial_{\alpha}^{m} \rho) g \right\|_{H^{s-j}} &\lesssim \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s_{0}}} \|g\|_{H^{s-j+1}} \\ &+ \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s-j+1}} \|g\|_{H^{s_{0}}}. \end{aligned}$$

$$\tag{8.15}$$

Therefore putting together (8.13), (8.14) and (8.15) allows to get,

_

$$\begin{aligned} \left\| \partial_{\theta} (\partial_{\alpha}^{j-k} \mathcal{T}) \partial_{\alpha}^{k} h_{t} \right\|_{H^{s-j}} &\lesssim \sum_{m=0}^{j-k} \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s_{0}}} \left\| \partial_{\alpha}^{k} h_{t} \right\|_{H^{s-j+1}} \\ &+ \sum_{m=0}^{j-k} \left\| \partial_{\alpha}^{m} \rho \right\|_{H^{s-j+1}} \left\| \partial_{\alpha}^{k} h_{t} \right\|_{H^{s_{0}}}. \end{aligned}$$

$$(8.16)$$

Plugging (8.16) into (8.12) yields

$$\frac{d}{dt} \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} \lesssim \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} \|\rho\|_{H^{s_{0}}(\mathbb{T}^{d+1})} + \|\partial_{\alpha}^{j} h_{t}\|_{H^{s_{0}}} \|\rho\|_{H^{s-j+s_{0}}} \\
+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \left(\|\partial_{\alpha}^{m} \rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{k} h_{t}\|_{H^{s-j+1}} + \|\partial_{\alpha}^{m} \rho\|_{H^{s-j+1}} \|\partial_{\alpha}^{k} h_{t}\|_{H^{s_{0}}} \right).$$
(8.17)

Using Gronwall inequality we get for $t \geqslant 0$

$$\begin{split} \|\partial_{\alpha}^{j}h_{t}\|_{H^{s-j}} &\lesssim e^{Ct\|\rho\|_{H^{s_{0}}}} \left[\|\partial_{\alpha}^{j}h\|_{H^{s-j}} + \|\rho\|_{H^{s-j+s_{0}}} \int_{0}^{t} e^{-C\tau\|\rho\|_{H^{s_{0}}}} \|\partial_{\alpha}^{j}h_{\tau}\|_{H^{s_{0}}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \|\partial_{\alpha}^{m}\rho\|_{H^{s_{0}}} \int_{0}^{t} e^{-C\tau\|\rho\|_{H^{s_{0}}}} \|\partial_{\alpha}^{k}h_{\tau}\|_{H^{s-j+1}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \|\partial_{\alpha}^{m}\rho\|_{H^{s-j+1}} \int_{0}^{t} e^{-C\tau\|\rho\|_{H^{s_{0}}}} \|\partial_{\alpha}^{k}h_{\tau}\|_{H^{s_{0}}} d\tau \right]. \end{split}$$

Multiplying by κ^j with $\kappa \in (0, 1)$ we infer

$$\begin{split} \kappa^{j} \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} &\lesssim e^{Ct} \|\rho\|_{H^{s_{0}}} \left[\kappa^{j} \|\partial_{\alpha}^{j} h\|_{H^{s-j}} + \|\rho\|_{H^{s-j+s_{0}}} \kappa^{j} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{j} h_{\tau}\|_{H^{s_{0}}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \kappa^{m} \|\partial_{\alpha}^{m} \rho\|_{H^{s_{0}}} \kappa^{k} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{k} h_{\tau}\|_{H^{s-j+1}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \kappa^{m} \|\partial_{\alpha}^{m} \rho\|_{H^{s-j+1}} \kappa^{k} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{k} h_{\tau}\|_{H^{s_{0}}} d\tau \right]. \end{split}$$

Thus we find from Sobolev embeddings

$$\begin{split} \kappa^{j} \|\partial_{\alpha}^{j} h_{t}\|_{H^{s-j}} &\lesssim e^{Ct} \|\rho\|_{H^{s_{0}}} \left[\kappa^{j} \|\partial_{\alpha}^{j} h\|_{H^{s-j}} + \|\rho\|_{H^{s+s_{0}}} \kappa^{j} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{j} h_{\tau}\|_{H^{s_{0}}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \kappa^{m} \|\partial_{\alpha}^{m} \rho\|_{H^{s_{0}+j-m}} \kappa^{k} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{k} h_{\tau}\|_{H^{s-k}} d\tau \\ &+ \sum_{k=0}^{j-1} \sum_{m=0}^{j-k} \kappa^{m} \|\partial_{\alpha}^{m} \rho\|_{H^{s+1-m}} \kappa^{k} \int_{0}^{t} e^{-C\tau} \|\rho\|_{H^{s_{0}}} \|\partial_{\alpha}^{k} h_{\tau}\|_{H^{s_{0}+j-1-k}} d\tau \right]. \end{split}$$

Consequently, according to the Definition 4.1 we get for any $s \ge q$,

$$\|h_{t}\|_{s}^{q,\kappa} \lesssim e^{Ct}\|\rho\|_{s_{0}}^{0,\kappa} \left[\|h\|_{s}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{0,\kappa} \int_{0}^{t} e^{-C\tau}\|\rho\|_{s_{0}}^{0,\kappa}} \|h_{\tau}\|_{s_{0}}^{q,\kappa} d\tau + \|\rho\|_{s_{0}+q}^{q,\kappa} \int_{0}^{t} e^{-C\tau}\|\rho\|_{s_{0}}^{0,\kappa}} \|h_{\tau}\|_{s}^{q-1,\kappa} d\tau + \|\rho\|_{s+1}^{q,\kappa} \int_{0}^{t} e^{-C\tau}\|\rho\|_{s_{0}}^{0,\kappa}} \|h_{\tau}\|_{s_{0}+q-1}^{q-1,\kappa} d\tau \right].$$

$$(8.18)$$

Assume that

$$\|\rho\|_{2s_0+q+1}^{q,\kappa} \lesssim 1, \tag{8.19}$$

then we get from the foregoing estimate combined with Sobolev embeddings

$$\forall |t| \leq 1, \quad \|h_t\|_s^{q,\kappa} \leq \|h\|_s^{q,\kappa} + \|\rho\|_{s+s_0}^{q,\kappa} \int_0^t \|h_\tau\|_{s_0+q}^{q,\kappa} d\tau + \|\rho\|_{s_0+q}^{q,\kappa} \int_0^t \|h_\tau\|_s^{q,\kappa} d\tau.$$

It follows from Gronwall inequality and (8.19) that

$$\forall |t| \lesssim 1, \, \forall s \ge q, \quad \|h_t\|_s^{q,\kappa} \lesssim \|h\|_s^{q,\kappa} + \|\rho\|_{s+s_0}^{q,\kappa} \int_0^t \|h_\tau\|_{s_0+q}^{q,\kappa} d\tau.$$
(8.20)

By taking $s = s_0 + q$ and using once again Gronwall inequality combined with (8.19) we infer

$$\forall |t| \lesssim 1, \quad \|h_t\|_{s_0+q}^{q,\kappa} \lesssim \|h\|_{s_0+q}^{q,\kappa}.$$

Plugging this inequality into (8.20) yields

$$\forall |t| \lesssim 1, \forall s \ge q, \quad \|h_t\|_s^{q,\kappa} \lesssim \|h\|_s^{q,\kappa} + \|\rho\|_{s+s_0}^{q,\kappa} \|h\|_{s_0+q}^{q,\kappa}$$

Using the flow mapping Φ , the preceding estimate can be written in the form

$$\forall |t| \lesssim 1, \, \forall s \ge q, \quad \|\Phi(t)h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{q,\kappa} \|h\|_{s_{0}+q}^{q,\kappa}.$$
(8.21)

The estimate of Φ^* which satisfies the equation (8.10) can be done in a similar way to $\Phi(t)$. The estimates of $g(t) \triangleq (\Phi(t) - \mathrm{Id})h$ can be done using the following steps. First we write the equation

$$\begin{cases} \partial_t g = \partial_\theta \mathcal{T} g + \partial_\theta \mathcal{T} h, \\ g(0) = 0. \end{cases}$$

Then using Duhamel formula we get

ł

$$g(t) = \int_0^t \Phi(t-\tau) \partial_\theta \mathcal{T} h d\tau.$$

Applying Proposition 8.1-(ii) yields under the assumption (8.19) that

$$\begin{aligned} \forall |t| \leq 1, \quad \|g(t)\|_{s}^{q,\kappa} \lesssim \|\partial_{\theta}\mathcal{T}h\|_{s}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{q,\kappa} \|\partial_{\theta}\mathcal{T}h\|_{s_{0}+q}^{q,\kappa} \\ \lesssim \|\mathcal{T}h\|_{s+1}^{q,\kappa} + \|\rho\|_{s+s_{0}+1}^{q,\kappa} \|\mathcal{T}h\|_{s_{0}+q+1}^{q,\kappa} \end{aligned}$$

Then applying the first result in Lemma 4.1-(ii) and Lemma 4.8, one may get under the smallness condition (8.19)

$$\|g(t)\|_{s}^{q,\kappa} \lesssim \|\rho\|_{s_{0}}^{q,\kappa}\|h\|_{s+1}^{q,\kappa} + \|\rho\|_{s+s_{0}}^{q,\kappa}\|h\|_{s_{0}+q+1}^{q,\kappa}.$$

The estimate $(\Phi^{\star}(t) - \mathrm{Id})h$ can be done in a similar way and this ends the proof of the second point. (iii) First notice that the operator $\Phi_{12} \triangleq \Delta_{12} \Phi$ satisfies the equation

$$\begin{cases} \partial_t \Phi_{12} = \mathbb{A}_1 \Phi_{12}(t) + \mathbb{A}_{12} \Phi_2(t), \\ \Phi(0) = 0, \end{cases}$$

$$(8.22)$$

Then using Duhamel formula we get

$$\Phi_{12}(t)h = \int_0^t \Phi_1(t-\tau)\mathbb{A}_{12}\Phi_2(\tau)hd\tau$$

Consequently, Proposition 8.1-(ii) yields under (8.19)

$$\forall |t| \leq 1, \quad \|\Phi_{12}h\|_{s}^{q,\kappa} \lesssim \int_{0}^{t} \|\mathbb{A}_{12}\Phi_{2}(\tau)h\|_{s}^{q,\kappa} d\tau + \|\rho_{1}\|_{s+s_{0}}^{q,\kappa} \int_{0}^{t} \|\mathbb{A}_{12}\Phi_{2}(\tau)h\|_{s_{0}+q}^{q,\kappa} d\tau.$$

On the other hand using the law products of Lemma 4.1 one gets

$$\|\mathbb{A}_{12}\Phi_2(\tau)h\|_s^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s+1}^{q,\kappa} \|\Phi_2(\tau)h\|_{s_0}^{q,\kappa} + \|\Delta_{12}\rho\|_{s_0}^{q,\kappa} \|\Phi_2(\tau)h\|_{s+1}^{q,\kappa}.$$

Using once again Proposition 8.1-(ii) combined with (4.8) we find for any $\tau \in [-1, 1]$

$$\|\mathbb{A}_{12}\Phi_{2}(\tau)h\|_{s}^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s+1}^{q,\kappa}\|h\|_{s_{0}+q}^{q,\kappa} + \|\Delta_{12}\rho\|_{s_{0}}^{q,\kappa} \Big[\|h\|_{s+1}^{q,\kappa} + \|\rho_{2}\|_{s+1+s_{0}}^{q,\kappa}\|h\|_{s_{0}+q}^{q,\kappa}\Big].$$

Combining the preceding few estimates and using (8.19) we find

$$\forall |t| \leq 1, \quad \|\Phi_{12}(t)h\|_{s_0}^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s_0+q+1}^{q,\kappa} \|h\|_{s_0+q+1}^{q,\kappa}.$$

In addition, we get for any $s \ge s_0$

$$\|\Phi_{12}(t)h\|_{s}^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s+1}^{q,\kappa}\|h\|_{s+1+q}^{q,\kappa} \Big(1+\|\rho_{1}\|_{s+s_{0}+1}^{q,\kappa}+\|\rho_{2}\|_{s+s_{0}+1}^{q,\kappa}\Big).$$

The same estimate holds for $\Phi_{12}^*(t)$ by using similar arguments combined with (8.10). The proof of the Proposition 8.1 is now achieved.

8.2. Töplitz matrices and binomial convolution. We shall discuss some basic properties on Töplitz matrix operators that will play a crucial role later when we will investigate the proof of an Egorov type theorem stated in Theorem 8.1.

Given $n \in \mathbb{N}$ and $k \in \{0, 1, ..., n+1\}$, we denote by $\Delta_k(n)$ the set of lower triangluar matrix operators \mathcal{M} such that

$$\mathcal{M} = \begin{pmatrix} m_{0,0} & 0 & \dots & \dots & \dots & 0 \\ m_{1,1} & m_{0,1} & 0 & \dots & 0 & 0 \\ m_{2,2} & m_{1,2} & m_{0,2} & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ m_{n-1,n-1} & m_{n-2,n-1} & \dots & \dots & m_{0,n-1} & 0 \\ m_{n,n} & m_{n-1,n} & \dots & \dots & m_{1,n} & m_{0,n} \end{pmatrix},$$

$$(8.23)$$

supplemented with the following constraints

$$m_{i,j} = \binom{j}{i} \mu_{\mathcal{M}}(i), \forall i \leq j \in \{0, .., n\} \text{ and } m_{i,j} = 0, \forall 0 \leq i \leq k-1,$$

where $\binom{j}{i}$ are the usual binomial coefficients and $\mu_{\mathcal{M}}(i)$ is a scalar operator that depends only on the index i and not on k. The index k gives the number of vanishing sub-diagonals (including the main diagonal) in the square matrix operator. We call $\mu_{\mathcal{M}}$ the law of \mathcal{M} . Typically, the main examples that we shall encounter during the proof of Theorem 8.1 are pseudo-differential operators of type $\mu_{\mathcal{M}}(i) = \rho^{(i)} |\mathbf{D}|^{\alpha}$, where the index *i* measures the spatial regularity of some suitable functions. Notice that by construction one has the inclusions

$$\Delta_0(n) \supset \Delta_1(n) \supset \ldots \supset \Delta_{n+1}(n) = \{0\}.$$

For further purposes, we need to define for $q \in \mathbb{N}$ the class $\Delta_k(n,q)$ of bloc matrix operators taking the form (8.23) and such that each entry $\mu_{\mathcal{M}}(i)$ is a matrix operator in $\Delta_0(q)$. It is obvious from the definitions that $\Delta_k(n,0) = \Delta_k(n)$. We point out that the introduction of such kind of operators through the index q appears natural during the proof of Theorem 8.1 when we will track the regularity estimates with respect to the external parameters. As to the index n it is associated to the time/spatial regularity.

At this stage, we do not care about the topological structure of the operators lying in the class $\Delta_k(n,q)$, we are simply concerned with some of their algebraic structures. One of the main results discussed below shows that commutators between matrix operators in the class $\Delta_k(n,q)$ still in the same class and whose entries are commutators too. This property fails in general even with standard real matrices, unless we impose additional structures as for example with Töplitz matrices. Notice that getting commutators in the entries is crucial to reduce the order of the operators during the proof of Theorem 8.1.

Lemma 8.1. The following assertions hold true.

(i) Let $n, q \in \mathbb{N}$ and $0 \leq k \leq n+1$, then the set $\Delta_k(n,q)$ is stable under multiplication, that is, for any $\mathcal{M}, \mathcal{N} \in \Delta_k(n,q)$ we have $\mathcal{MN} \in \Delta_k(n,q)$ and its law $\mu_{\mathcal{MN}}$ is given by the binomial convolution,

$$\mu_{\mathcal{M}\mathcal{N}}(i) = \sum_{\ell=0}^{i} {i \choose \ell} \, \mu_{\mathcal{M}}(i-\ell) \mu_{\mathcal{N}}(\ell)$$
$$\triangleq (\mu_{\mathcal{M}} \circledast \mu_{\mathcal{N}}) \, (i).$$

In addition, if $\mathcal{M} \in \Delta_k(n,q), \mathcal{N} \in \Delta_m(n,q)$, then $\mathcal{MN} \in \Delta_{\min(k+m,n+1)}(n,q)$. (ii) For any $\mathcal{M}, \mathcal{N} \in \Delta_k(n,q)$ the law of the commutator $[\mathcal{M}, \mathcal{N}]$ is given by

$$\mu_{[\mathcal{M},\mathcal{N}]}(i) = \sum_{\ell=0}^{i} {i \choose \ell} \left[\mu_{\mathcal{M}}(i-\ell), \mu_{\mathcal{N}}(\ell) \right]$$
$$\triangleq (\mu_{\mathcal{M}} \ltimes \mu_{\mathcal{N}})(i).$$

Furthermore, the law of the iterated commutator $\operatorname{Ad}^n_{\mathscr{M}} \mathcal{N}$ is given by

$$\mu_{\mathrm{Ad}_{\mathcal{M}}^{n}\mathcal{N}} = \underbrace{\mu_{\mathcal{M}} \circledast \mu_{\mathcal{M}} \circledast \dots \circledast \mu_{\mathcal{M}} \circledast \mu_{\mathcal{N}}}_{n \text{ times}}.$$

Remark 8.1. As a byproduct of this lemma, if $\mathscr{M}, \mathscr{N} \in \Delta_0(n)$ defined in (8.23) and their entries $m_{i,j}, n_{k,\ell}$ commute together, that is, $[m_{i,j}, n_{k,\ell}] = 0$, then $[\mathscr{M}, \mathscr{N}] = 0$. In addition, the stability point stated in (i) asserts that the product of two elements contains more vanishing sub-diagonals. In particular, if $\mathscr{M}, \mathscr{N} \in \Delta_1(n)$ and $1 \leq j \leq n+1$ then

$$\mathcal{MN} \in \Delta_2(n) \quad and \quad \mathcal{M}^j \in \Delta_j(n).$$

From the last point we infer that the matrix operator \mathcal{M} is nilpotent and $\mathcal{M}^{n+1} = 0$.

Proof. (i) Let $\mathcal{M} = (\overline{m}_{i,j})_{0 \leq i,j \leq n} \in \Delta_k(n,q)$ and $\mathcal{N} = (\overline{n}_{i,j})_{0 \leq i,j \leq n} \in \Delta_k(n,q)$ and set $\mathcal{P} = \mathcal{M}\mathcal{N}$, then with these notations

$$\forall 0 \leq j \leq i, \quad \overline{m}_{i,j} = m_{i-j,i}, \quad \overline{n}_{i,j} = n_{i-j,i}$$

Thus we may check that

$$\mathcal{P} = \begin{pmatrix} p_{0,0} & 0 & \dots & \dots & 0 \\ p_{1,1} & p_{0,1} & 0 & \dots & 0 & 0 \\ p_{2,2} & p_{1,2} & p_{0,2} & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ p_{n-1,n-1} & p_{n-2,n-1} & \dots & \dots & p_{0,n-1} & 0 \\ p_{n,n} & p_{n-1,n} & \dots & \dots & p_{1,n} & p_{0,n} \end{pmatrix} = (\overline{p}_{i,j})_{0 \leq i,j \leq n}$$

with $\overline{p}_{i,j} = 0$ for i < j, and for $0 \leq j \leq i$ we have

$$\overline{p}_{i,j} = \sum_{\ell=j}^{i} \overline{m}_{i,\ell} \overline{n}_{\ell,j}$$
$$= \sum_{\ell=j}^{i} m_{i-\ell,i} n_{\ell-j,\ell}.$$

Consequently we find through elementary combinatorics formula

$$\overline{p}_{i,j} = \sum_{\ell=j}^{i} {i \choose i-\ell} {\ell \choose \ell-j} \mu_{\mathcal{M}}(i-\ell)\mu_{\mathcal{N}}(\ell-j)$$
$$= {i \choose j} (\mu_{\mathcal{M}} \circledast \mu_{\mathcal{N}}) (i-j)$$

and therefore for $0 \leq i \leq j$

$$p_{i,j} = \overline{p}_{j,j-i}$$
$$= {j \choose i} (\mu_{\mathcal{M}} \circledast \mu_{\mathcal{N}}) (i).$$

Thus the law of \mathcal{MN} is given by the binomial convolution. Now since $\mathcal{M}, \mathcal{N} \in \Delta_k(n,q)$ then by construction $\mu_{\mathcal{M}}(i), \mu_{\mathcal{N}}(i) \in \Delta_0(q)$ and

$$\mu_{\mathcal{M}}(i) = \mu_{\mathcal{N}}(i) = 0, \quad \forall \, 0 \leqslant i \leqslant k-1.$$

This implies that $\mu_{\mathcal{M}\mathcal{N}}(i) \in \Delta_0(q)$ with $\mu_{\mathcal{M}\mathcal{N}}(i) = 0$, for any $0 \leq i \leq k-1$. Thus the bloc matrix operator $\mathcal{M}\mathcal{N} \in \Delta_k(n,q)$. Similarly, if $\mathcal{M} \in \Delta_k(n,q)$ and $\mathcal{N} \in \Delta_m(n,q)$ then from straightforward matrix computations we find

$$\mu_{\mathcal{M}\mathcal{N}}(i) = 0, \quad \forall \, 0 \leqslant i \leqslant \min(k+m-1,n).$$

Hence $\mathcal{MN} \in \Delta_{\min(k+m,n+1)}(n,q)$.

(ii) Let $\mathcal{M}, \mathcal{N} \in \Delta_k(n, q)$ then from the preceding point (i) the commutator $[\mathcal{M}, \mathcal{N}]$ remains in $\Delta_k(n, q)$ and its law is described by

$$\mu_{[\mathcal{M},\mathcal{N}]}(i) = \sum_{\ell=0}^{i} {i \choose \ell} \left[\mu_{\mathcal{M}}(i-\ell), \mu_{\mathcal{N}}(\ell) \right]$$

$$\triangleq \mu_{\mathcal{M}} \circledast \mu_{\mathcal{N}}(i)$$

where $[\mu_{\mathcal{M}}(i-\ell), \mu_{\mathcal{N}}(\ell)]$ is the commutator between the matrix operators $\mu_{\mathcal{M}}(i-\ell)$ and $\mu_{\mathcal{N}}(\ell)$ that belong to $\Delta_0(q)$ and thus

$$[\mu_{\mathcal{M}}(i-\ell),\mu_{\mathcal{N}}(\ell)] \in \mathbf{\Delta}_0(q).$$

Notice that the entries of the matrix operator $[\mu_{\mathcal{M}}(i-\ell), \mu_{\mathcal{N}}(\ell)]$ are linear combination of scalar commutators. This ends the proof of the desired result.

8.3. Refined Egorov's theorem. The main concern of this section is to deal with the conjugation of some class of pseudo-differential operators with the hyperbolic flow given in (8.1) and constructed in Proposition 8.1. We shall establish in Theorem 8.1 below a refined version of the classical Egorov theorem in a suitable topology of pseudo-differential operators of order $m \in [-1, 0]$. Notice that the tame estimates, which are crucial in the KAM reduction and along Nash-Moser scheme, will be delicately established. This part is the hardest technical point raised in this paper and will be performed through a new approach based on both kernel and symbol representations of operators combined with Töplitz structure on some matrices involved in the dynamics of the kernel derivatives. Let us start with fixing the problem that we wish to solve and directly connected with KAM reduction of the remainder that will be carefully investigated in Section 9.6.3. Consider a toroidal pseudo-differential operator $\mathscr{R}_0: \mathscr{C}^{\infty}(\mathbb{T}^{d+1}) \mapsto \mathscr{C}(\mathbb{T}^{d+1})$ with the following kernel representation

$$\mathscr{R}_0 h(\varphi, \theta) = \int_{\mathbb{T}} \mathscr{K}_0(\varphi, \theta, \eta) h(\varphi, \eta) d\eta.$$
(8.24)

This structure is very special and more specifically, the variable φ acts as a local parameter in this description. It allows to cover a large class of operators that turns out to be sufficient for the applications in this paper. The symbol a_0 associated to \mathcal{R}_0 is related to the kernel \mathcal{K}_0 according to the formula (7.8)

$$\sigma_{\mathcal{R}_0}(\varphi,\theta,\xi) = \int_{\mathbb{T}} \mathcal{K}_0(\varphi,\theta,\theta+\eta) e^{i\eta\xi} d\eta.$$

Let $\Phi(t)$ be the flow defined as the solution to the equation (8.1) and whose generator is given by the pseudo-differential operator

$$\mathbb{A} = \partial_{\theta} \left(\rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \rho \right)$$

$$\triangleq \partial_{\theta} \mathcal{J}_{\theta}.$$

$$(8.25)$$

The function ρ is smooth enough and may depend on $(\varphi, \theta) \in \mathbb{T}^{d+1}$ or on the variables $(\lambda, \varphi, \theta) \in \mathcal{O} \times \mathbb{T}^{d+1}$. We remind that the modified fractional Laplacian $|\mathbf{D}|^{\alpha-1}$ was introduced in (4.19) and from which we infer

$$\mathbb{A}h(\varphi,\theta) = \partial_{\theta} \int_{\mathbb{T}} K(\varphi,\theta,\eta)h(\varphi,\eta)d\eta, \quad K(\varphi,\theta,\eta) = \frac{1}{2\pi} \frac{\rho(\varphi,\theta) + \rho(\varphi,\eta)}{|\sin(\frac{\theta-\eta}{2})|^{\alpha}} \cdot$$

Let us precise an elementary observation that will be frequently used later. As one can see the kernel K is singular at the diagonal $\theta = \eta$ but the modified one given by $(\varphi, \theta, \eta) \mapsto K(\varphi, \theta, \theta + \eta)$ is smooth on θ and singular only at $\eta = 0$ which is the integration variable. By this way, the singularity in η which is not so violent can be merely treated by integration. Here we shall fix the rectangular open box

$$\mathcal{O} = \mathscr{U} \times (0, \overline{\alpha}) \quad \text{with} \quad \overline{\alpha} \in \left(0, \frac{1}{2}\right)$$

and \mathscr{U} is an open bounded set of \mathbb{R}^d . The main task here is to analyze the conjugation of \mathscr{R}_0 by the flow $\Phi(t)$, that is to consider $\Phi(t)\mathscr{R}_0\Phi(-t)$, and show that this operator remains in the same class of pseudo-differential operators with the suitable tame estimates. We refer to Section 7.2 for the definition of the class of symbols that will used below. Our main statement reads as follows.

Theorem 8.1. Let $m \in [-1,0], q \in \mathbb{N}$ and \mathcal{R}_0 be the operator in (8.24). Then the following assertions hold true.

(i) Let $s \ge 0$, $s_0 > \frac{d+1}{2}$ and assume the existence of an increasing log-convex function F_0 such that

$$\forall s' \ge 0, \forall \gamma \in \mathbb{N} \quad with \quad s' + \gamma \leqslant s_0 + 27 + s \Longrightarrow |||\mathcal{R}_0|||_{m,s',\gamma} \leqslant F_0(s' + \gamma).$$

Then for any $t \in [0, 1]$

$$|||\Phi(t)\mathscr{R}_0\Phi(-t)|||_{m,s,0} \leqslant C e^{C\mu^{s+1}(0)}\mu(s),$$

with

$$\mu(s) = F_0(s + 27 + s_0) + \|\rho\|_{H^{s+27+s_0}}$$

and C being a constant depending only on s. We also get

$$\||\Phi(t)\mathcal{R}_0\Phi(-t)|||_{m,s,0} \le Ce^{C\mu^{s+1}s(0)}F_0(s+27+s_0)(1+\mu(s))$$

(ii) Let $s \ge 0$, $s_0 > \frac{d+1}{2} + q$ and assume the existence of an increasing log-convex function \overline{F}_0 such that

$$\forall s' \ge 0, \forall \gamma \in \mathbb{N} \quad with \quad s' + \gamma \leqslant s_0 + 27 + s \Longrightarrow ||\mathscr{R}_0||_{m,s',\gamma}^{q,\kappa} \leqslant \overline{F}_0(s' + \gamma).$$

Then for any $t \in [0, 1]$

$$\||\Phi(t)\mathcal{R}_0\Phi(-t)|||_{m,s,0}^{q,\kappa} \leqslant Ce^{C\overline{\mu}^{s+1}(0)}\overline{\mu}(s)$$

with

$$\overline{\mu}(s) = \overline{F}_0(s + 27 + s_0) + \|\rho\|_{s+27+s_0}^{q,\kappa}.$$

We also get

$$|||\Phi(t)\mathcal{R}_0\Phi(-t)|||_{m,s,0}^{q,\kappa} \leqslant C e^{C\overline{\mu}^{s+1}(0)}\overline{F}_0(s+27+s_0) \left(1+\overline{\mu}(s)\right).$$

(iii) Denote $\Delta_{12}\Phi(t)\mathcal{R}_0\Phi(-t) \triangleq \Phi_1(t)\mathcal{R}_0\Phi_1(-t) - \Phi_2(t)\mathcal{R}_0\Phi_2(-t)$ where Φ_i is the flow associated to ρ_i . Let $s_0 > \frac{d+1}{2} + q$ and assume the existence of an increasing log-convex function \overline{F}_0 such for such that

$$\forall s' \ge 0, \forall \gamma \in \mathbb{N} \quad with \quad s' + \gamma \leqslant s_0 + 10 \Longrightarrow |||\mathscr{R}_0|||_{m,s',\gamma}^{q,\kappa} \leqslant \overline{F}_0(s' + \gamma).$$

Then for any $t \in [0, 1]$

$$\||\Delta_{12}\Phi(t)\mathcal{R}_0\Phi(-t)|||_{0,0,0}^{q,\kappa} \leqslant Ce^{C\left(\|\rho_2\|_{H^{s_0+10}} + \|\rho_1\|_{H^{s_0+10}} + \overline{F}_0(s_0+10)\right)} \|\rho_1 - \rho_2\|_{s_0+5}^{q,\kappa}.$$

Before moving to the proof, some remarks are in order.

- (i) The statement of the theorem is restricted for $m \in [-1,0]$ which quite suffi-Remark 8.2. cient in the application during the remainder reduction that will be explored in Section 9.6.3. The extension to the values of m in the complement set is likely possible provided we impose the suitable adaptations.
 - (ii) It is worthy to point out that the constraints on the initial symbol covers a range of values on the parameters s, γ but the conclusion deals only with the estimate with the norm $\|\cdot\|_{m,s,0}$ where $\gamma = 0$. This fact is due to a diffusion mechanism in the regularity persistence of the symbol estimates: this stems in part from the commutator estimates detailed in Lemma 7.5 where we win in the order of the operator but we lose in space-phase variables.
 - (iii) In the statement (iii) the estimates on the difference are measured in the topology $\||\cdot\||_{0,0,0}^{q,\kappa}$ and not with the expected one $\||\cdot\||_{m,s,0}^{q,\kappa}$. This is due to delicate technical points that could be solved with the same approach but with more refined analysis. Notice that, later during the application in the study of the spectrum dependence of the linearized operator, this weak estimate will provide us by interpolation arguments with only a Hölder continuity dependence with respect to the torus i instead of Lipschitz dependence. This slight loss of regularity has no important effects in the arguments used to establish the stability of Cantor sets during the last phase in Nash-Moser scheme related the measure of the final Cantor set associated to the emergence of invariant nonlinear torus.

Proof. Let us briefly explore the main ideas of the proof. We shall write down the kernel equation of the the operator $\Re(t) \triangleq \Phi(t) \Re_0 \Phi(-t)$. Then the symbol estimates are transformed into looking for suitable kernel estimates in weighted Sobolev spaces. This will be implemented by successive differentiations of the time evolution of the kernel equation combined with energy estimates. This procedure leads to a collection of coupled transport equations with non dissipative positive order operators acting as source terms that cannot be directly treated by energy estimates. The key observation is to understand the global structure of the full set of equations that can be recast into a vectorial transport equations with an additional positive order matrix operator. The particularity of this matrix is to be nilpotent and of Töpliz structure as in the definition (8.23). Then we perform energy estimates through a new mixed norm and the use of refined commutator estimates leading to several technical difficulties that will be made clear throughout the proof.

(i) It seems to be convenient to start with proving the result for m = 0 and move later to the case $m \in [-1, 0)$ by making the suitable adaptations.

► Case m = 0. Denote by $\Re(t) \triangleq \Phi(t) \Re_0 \Phi(-t)$ then we can check directly from the flow equation (8.1) that this operator satisfies the Heisenberg equation

$$\partial_t \mathcal{R}(t) = [\mathbb{A}, \mathcal{R}(t)].$$

Now, set $\Re_1(t) \triangleq \Re(t) - \Re_0$, then we deduce from the preceding equation that

$$\begin{cases} \partial_t \mathcal{R}_1(t) = [\mathbb{A}, \mathcal{R}_1(t)] + [\mathbb{A}, \mathcal{R}_0] \\ \mathcal{R}_1(0) = 0. \end{cases}$$
(8.26)

To show the desired estimate for $\Re(t)$ it is enough to prove, according to Sobolev embeddings, that the operator $\Re_1(t)$ satisfies for any $t \in [0, 1]$ and $s \in \mathbb{N}$

$$\|\|\mathscr{R}_{1}(t)\|\|_{0,s,0} \lesssim C\left(1 + F_{0}^{2}(s_{0} + 12)\right)e^{C\|\rho\|_{H^{s_{0}+12}}^{s}}\left(F_{0}(s + 12) + \|\rho\|_{H^{s+12}}\right).$$
(8.27)

Since the variable φ acts as a parameter on the generator \mathbb{A} , then it will act in a similar way on the associated flow $\Phi(t)$. This implies that one may view the operator $\mathcal{R}_1(t)$ as a collection of operators $\mathcal{R}_1(t,\varphi)$ parametrized by φ and acting pointwisely on test functions as $\mathcal{R}_1(t,\varphi)h(\varphi,\cdot)$. As a consequence one gets the following kernel representation of $\mathcal{R}_1(t)$,

$$\mathscr{R}_1(t)h(\varphi,\theta) = \int_{\mathbb{T}} \mathscr{K}_t(\varphi,\theta,\eta)h(\varphi,\eta)d\eta$$

and therefore the dynamics is encoded in the new kernel \mathcal{K}_t that we need to estimate carefully. Coming back to the norm definition (7.9), we may write

$$|||\mathscr{R}_{1}(t)|||_{0,s,0} = \sup_{\xi \in \mathbb{Z}} ||\sigma_{t}(\cdot, \cdot, \xi)||_{H^{s}(\mathbb{T}^{d+1})}$$

with σ_t being the symbol of $\Re_1(t)$ that can be recovered from the kernel according to the identity (7.8)

$$\sigma_t(\varphi, \theta, \xi) = e^{-i\xi\theta} \mathscr{R}_1(t) e^{i\xi\theta}$$
$$= \int_{\mathbb{T}} \mathscr{K}_t(\varphi, \theta, \theta + \eta) e^{i\eta\xi} d\eta.$$
(8.28)

Then applying Cauchy-Schwarz inequality and using a change of variables give with $s = n \in \mathbb{N}$

$$\begin{aligned} \|\|\mathscr{R}_{1}(t)\|\|_{0,s,0} &\leq \int_{\mathbb{T}} \|\mathscr{K}_{t}(\cdot, \cdot, \cdot + \eta)\|_{H^{s}_{\varphi,\theta}} d\eta \\ &\leq \int_{\mathbb{T}} \|\mathscr{K}_{t}(\cdot, \cdot, \cdot + \eta)\|_{H^{s}_{\theta}L^{2}_{\varphi}} d\eta + \int_{\mathbb{T}} \|\mathscr{K}_{t}(\cdot, \cdot, \cdot + \eta)\|_{L^{2}_{\theta}H^{s}_{\varphi}} d\eta \\ &\lesssim \sum_{i=0}^{n} \left(\|\partial^{i}_{\chi}\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} + \|\partial^{i}_{\varphi}\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \right) \end{aligned}$$

with

$$\partial_{\chi} = \partial_{\theta} + \partial_{\eta}$$

The estimates of the two terms of the right-hand side are quite similar and we shall start the first one dealing the mixed derivatives.

• Estimate of $\|\partial_{\mathcal{X}}^{i}\mathcal{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})}$. The main goal is to prove the following estimate: $\forall i \in \{0, 1, ..., n\}$,

$$\forall t \in [0,1], \quad \|\partial_{\chi}^{i} \mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant C\left(1 + G^{i+1}(0)\right) e^{C\|\rho\|_{H^{s_{0}+3}}^{n}} G(1+i)$$
(8.29)

with

$$G(i) \triangleq \left(\|\rho\|_{H^{s_0}} + F_0(s_0) \right) \left(\|\rho\|_{H^{s_0+11+i}} + F_0(s_0+11+i) \right) + \|\rho\|_{H^{s_0+7+i}}.$$

Recall that F_0 is a log-convex function that controls some suitable symbol norm as assumed in Theorem 8.1-(i). To proceed, we shall start with writing the evolution equation governing the kernel \mathcal{K}_t . From the integral representation one gets by definition of the commutator

$$[\mathbb{A}, \mathcal{R}_{1}(t)]h(\varphi, \theta) = \int_{\mathbb{T}} \left(\mathbb{A}_{\theta} \mathcal{K}_{t}(\varphi, \theta, \eta) \right) h(\varphi, \eta) d\eta - \int_{\mathbb{T}} \mathcal{K}_{t}(\varphi, \theta, \eta) \left(\mathbb{A}_{\eta} h \right)(\varphi, \eta) d\eta = \int_{\mathbb{T}} \left(\mathbb{A}_{\theta} \mathcal{K}_{t} - \mathbb{A}_{\eta}^{\star} \mathcal{K}_{t} \right)(\varphi, \theta, \eta) h(\varphi, \eta) d\eta.$$

$$(8.30)$$

The notation \mathbb{A}_{θ} stands for the action of the operator \mathbb{A} in the variable θ , that is,

$$\mathbb{A}_{\theta}h(\varphi,\theta) = \partial_{\theta} \big(\rho(\varphi,\theta) |\mathcal{D}_{\theta}|^{\alpha-1}h + |\mathcal{D}_{\theta}|^{\alpha-1}\rho(\varphi,\cdot)h \big)(\varphi,\theta)$$

and similarly we define \mathbb{A}_{η} . We also denote by $\mathbb{A}_{\eta}^{\star}$ the partial adjoint of \mathbb{A}_{η} with respect to Hilbert structure of $L^{2}_{\eta}(\mathbb{T})$. Then combining Heisenberg equation (8.26) with (8.30) we get the evolution nonlocal PDE for the kernel

$$\begin{cases} \partial_t \mathcal{K}_t(\varphi, \theta, \eta) = \mathbb{A}_{\theta} \mathcal{K}_t(\varphi, \theta, \eta) - \mathbb{A}_{\eta}^* \mathcal{K}_t(\varphi, \theta, \eta) + K_0(\varphi, \theta, \eta) \\ \mathcal{K}_0 = 0, \end{cases}$$

where K_0 is the kernel associated to $[\mathbb{A}, \mathcal{R}_0]$ in the sense

$$[\mathbb{A}, \mathcal{R}_0]h(\varphi, \theta) = \int_{\mathbb{T}} K_0(\varphi, \theta, \eta)h(\varphi, \eta)d\eta$$

and one can check the following relation

$$K_0(\varphi,\theta,\eta) = \mathbb{A}_{\theta} \mathscr{K}_0(\varphi,\theta,\eta) - \mathbb{A}_{\eta}^{\star} \mathscr{K}_0(\varphi,\theta,\eta).$$

By virtue of (8.25), it is clear that the operator \mathcal{T}_{η} is self-adjoint and then we can easily check that

$$\mathbb{A}_{\eta}^{\star} = -\mathbb{A}_{\eta} + \left[\partial_{\eta}, \mathcal{T}_{\eta}\right]. \tag{8.31}$$

Consequently, the kernel equation becomes

$$\partial_t \mathcal{K}_t(\varphi, \theta, \eta) - (\mathbb{A}_\theta + \mathbb{A}_\eta) \mathcal{K}_t(\varphi, \theta, \eta) + [\partial_\eta, \mathcal{T}_\eta] \mathcal{K}_t(\varphi, \theta, \eta) = K_0(\varphi, \theta, \eta).$$
(8.32)

Direct computations yield the identity

$$\mathscr{S}_{0} \triangleq \left[\partial_{\eta}, \mathscr{T}_{\eta}\right] = (\partial_{\eta}\rho)|\mathbf{D}_{\eta}|^{\alpha-1} + |\mathbf{D}_{\eta}|^{\alpha-1}(\partial_{\eta}\rho).$$

$$(8.33)$$

Now, we shall prove the estimate (8.29) for i = 0. For this purpose, we take the $L^2(\mathbb{T}^{d+2})$ inner product of the equation (8.32) with \mathcal{K}_t

$$\begin{split} \frac{d}{dt} \int_{\mathbb{T}^{d+2}} |\mathcal{K}_t(\varphi,\theta,\eta)|^2 d\varphi d\theta d\eta &= \int_{\mathbb{T}^{d+2}} \mathcal{K}_t(\varphi,\theta,\eta) \left(\left[\mathcal{T}_\eta, \partial_\eta \right] - \left[\mathcal{T}_\theta, \partial_\theta \right] \right) \mathcal{K}_t(\varphi,\theta,\eta) d\varphi d\theta d\eta \\ &+ 2 \int_{\mathbb{T}^{d+2}} \mathcal{K}_t(\varphi,\theta,\eta) K_0(\varphi,\theta,\eta) d\varphi d\theta d\eta, \end{split}$$

where we have used by virtue of (8.31) the identity

$$\begin{split} \langle \mathbb{A}_{\theta} \mathscr{K}_{t}, \mathscr{K}_{t} \rangle_{L^{2}(\mathbb{T}^{d+2})} &= \frac{1}{2} \langle (\mathbb{A}_{\theta} + \mathbb{A}_{\theta}^{\star}) \mathscr{K}_{t}, \mathscr{K}_{t} \rangle_{L^{2}(\mathbb{T}^{d+2})} \\ &= \frac{1}{2} \langle [\partial_{\theta}, \mathcal{T}_{\theta}] \mathscr{K}_{t}, \mathscr{K}_{t} \rangle_{L^{2}(\mathbb{T}^{d+2})}. \end{split}$$

It follows that

$$\frac{d}{dt}\int_{\mathbb{T}^{d+2}}|\mathscr{K}_t(\varphi,\theta,\eta)|^2d\varphi d\theta d\eta \lesssim \|\partial_\theta\rho\|_{L^{\infty}}\|\mathscr{K}_t\|_{L^2(\mathbb{T}^{d+2})}^2 + \|\mathscr{K}_t\|_{L^2(\mathbb{T}^{d+2})}^2\|K_0\|_{L^2(\mathbb{T}^{d+2})}.$$

Applying Gronwall inequality yields

$$\|\mathscr{K}_t\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|K_0\|_{L^2(\mathbb{T}^{d+2})} t \, e^{Ct \|\rho\|_{W^{1,\infty}}}.$$
(8.34)

Notice that his energy estimate can be extended to cover a more general case with time dependent source terms. Indeed, let f and g satisfy

$$\mathscr{L}(f) \triangleq \partial_t f + \widehat{\mathscr{L}}(f)$$
$$\triangleq \partial_t f - \widehat{\mathbb{A}}f + \mathscr{S}_0 f = g, \quad \widehat{\mathbb{A}} \triangleq \mathbb{A}_\theta + \mathbb{A}_\eta, \tag{8.35}$$

then for any $t \ge 0$

$$\|f(t)\|_{L^{2}(\mathbb{T}^{d+2})} \leq e^{Ct}\|\rho\|_{W^{1,\infty}} \left(\|f(0)\|_{L^{2}(\mathbb{T}^{d+2})} + \int_{0}^{t} e^{-C\tau}\|\rho\|_{W^{1,\infty}} \|g(\tau)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau\right).$$
(8.36)

The next step aims at extending the estimate (8.34) to higher order regularity. First, we remark that the equation (8.32) can be recast in the form

$$\mathscr{L}(\mathscr{K}_t) = K_0. \tag{8.37}$$

Then applying ∂^i_{χ} to (8.37) yields

$$\mathscr{L}\partial^{i}_{\chi}\mathscr{K}_{t} = \left[\partial^{i}_{\chi},\widehat{\mathbb{A}}\right]\mathscr{K}_{t} - \left[\partial^{i}_{\chi},\mathscr{S}_{0}\right]\mathscr{K}_{t} + \partial^{i}_{\chi}K_{0}.$$
(8.38)

By virtue of the abstract Leibniz rule we obtain

$$\left[\partial_{\chi}^{i},\widehat{\mathbb{A}}\right] = \sum_{k=0}^{i-1} {i \choose k} \operatorname{Ad}_{\partial_{\chi}}^{i-k}(\widehat{\mathbb{A}}) \partial_{\chi}^{k}, \qquad (8.39)$$

where the adjoint mapping Ad_A and its iterated ones Ad_A^i are defined by

$$\operatorname{Ad}_A B = [A, B]$$
 and $\operatorname{Ad}_A^i = \underbrace{\operatorname{Ad}_A \circ \dots \circ \operatorname{Ad}_A}_{i \text{ times}}.$

Using the special structure of the operator $\widehat{\mathbb{A}}$ detailed through (8.25) and (8.35) we can check by induction the following identity, using in particular the general relation $\operatorname{Ad}_{A}^{i} = [A, \operatorname{Ad}_{A}^{i-1}],$

$$\mathrm{Ad}_{\partial_{\chi}}^{k}(\widehat{\mathbb{A}}) = \widehat{\mathbb{A}}_{(k)},$$

with

$$\widehat{\mathbb{A}}_{(k)} = \mathbb{A}_{\theta,(k)} + \mathbb{A}_{\eta,(k)}, \quad \mathbb{A}_{\theta,(k)} \triangleq \partial_{\theta} \left((\partial_{\theta}^{k} \rho) |\mathcal{D}_{\theta}|^{\alpha - 1} + |\mathcal{D}_{\theta}|^{\alpha - 1} (\partial_{\theta}^{k} \rho) \right).$$
(8.40)

Notice that we have also used some commutation relations of the type $[\partial_{\theta}, \mathbb{A}_{\eta,(k)}] = [\partial_{\eta}, \mathbb{A}_{\theta,(k)}] = 0.$ Therefore we get by virtue of (8.39)

$$\left[\partial_{\chi}^{i},\widehat{\mathbb{A}}\right] = \sum_{k=0}^{i-1} {i \choose k} \widehat{\mathbb{A}}_{(i-k)} \partial_{\chi}^{k}.$$
(8.41)

Similarly we get from (8.33)

$$\begin{bmatrix} \partial_{\chi}^{i}, \mathscr{S}_{0} \end{bmatrix} = \sum_{k=0}^{i-1} {i \choose k} \operatorname{Ad}_{\partial_{\chi}}^{i-k}(\mathscr{S}_{0}) \partial_{\chi}^{k}$$
$$= \sum_{k=0}^{i-1} {i \choose k} \mathbb{S}_{(i-k)} \partial_{\chi}^{k}$$
(8.42)

with

$$\mathbb{S}_{(k)} \triangleq (\partial_{\eta}^{(k+1)} \rho) |\mathcal{D}_{\eta}|^{\alpha-1} + |\mathcal{D}_{\eta}|^{\alpha-1} (\partial_{\eta}^{(k+1)} \rho).$$

$$(8.43)$$

Plugging the identities (8.41) and (8.42) into (8.38) gives for any $i \in \{0, 1, ..., n\}$

$$\mathscr{L}\partial_{\chi}^{i}\mathscr{K}_{t} = \sum_{k=0}^{i-1} {i \choose i-k} \left(\widehat{\mathbb{A}}_{(i-k)} - \mathbb{S}_{(i-k)}\right) \partial_{\chi}^{k}\mathscr{K}_{t} + \partial_{\chi}^{j}K_{0}$$
$$= \sum_{k=0}^{i-1} m_{i-k,i} \partial_{\chi}^{k}\mathscr{K}_{t} + \partial_{\chi}^{i}K_{0}$$
(8.44)

with

$$m_{k,i} \triangleq \binom{i}{k} \left(\widehat{\mathbb{A}}_{(k)} - \mathbb{S}_{(k)}\right). \tag{8.45}$$

We observe that for i = 0 the first term in the right hand side disappears. Now, we intend to write the system of equations (8.44) in the matrix form. For this goal, let us introduce the lower triangular matrix operator

$$\mathcal{M}_{n} = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots & 0 \\ m_{1,1} & 0 & \dots & \dots & 0 & 0 \\ m_{2,2} & m_{1,2} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & m_{1,n-1} & 0 & 0 \\ m_{n,n} & m_{n-1,n} & \dots & \dots & m_{1,n} & 0 \end{pmatrix}$$
(8.46)

and the vectors

$$\mathfrak{X}_{n}(t) = \begin{pmatrix} \mathcal{K}_{t} \\ \partial_{\chi} \mathcal{K}_{t} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n} \mathcal{K}_{t} \end{pmatrix}, \quad \mathfrak{Y}_{n} = \begin{pmatrix} K_{0} \\ \partial_{\chi} K_{0} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n} K_{0} \end{pmatrix}.$$
(8.47)

Then (8.44) can be recast in the matrix form as follows

$$\mathscr{L}_n \mathscr{X}_n(t) = \mathscr{M}_n \mathscr{X}_n(t) + \mathscr{Y}_n, \quad \mathscr{L}_n \triangleq \mathscr{L} \mathbf{I}_{n+1}.$$
 (8.48)

It is worthy to mention that the time-independent matrix operator \mathcal{M}_n belongs to the class Δ_1 introduced in belongs is (8.23). This algebraic structure will be of important use along the proof. In particular, one deduces that this matrix is nilpotent with

$$\mathcal{M}_n^{n+1} = 0. \tag{8.49}$$

This property is very important because the entries of \mathcal{M}_n are operators with strictly positive orders that cannot be directly treated by energy estimates but the nilpotent structure gives somehow a cancellation and one may expect to close energy estimates by making appeal to a suitable norms combination. In order to take advantage of the nilpotent structure, we shall apply successively the matrix \mathcal{M}_n to the equation (8.48) and write down the suitable evolution equations for the iterated vectors $\mathcal{M}_n^k \mathcal{X}_n(t)$. To perform this, we first apply the abstract Leibniz rule used in (8.39) leading to

$$\begin{bmatrix} \mathcal{M}_n^k, \mathcal{L}_n \end{bmatrix} = \sum_{j=0}^{k-1} {k \choose j} \operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\mathcal{L}_n) \mathcal{M}_n^j$$
$$= \sum_{j=0}^{k-1} {k \choose j} \operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\widehat{\mathcal{L}}_n) \mathcal{M}_n^j, \quad \widehat{\mathcal{L}}_n \triangleq \widehat{\mathscr{L}} \operatorname{I}_{n+1}$$

We have used in the last line the identity $\operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\widehat{\mathcal{L}}_n) = \operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\mathcal{L}_n)$ which is a consequence of the definition (8.35) and the fact that the matrix \mathcal{M}_n is time-independent. Applying \mathcal{M}_n^k to the equation (8.48) and using the preceding identity allow to get for any $k \in \{1, ., n\}$

$$\mathcal{L}_{n}\mathcal{M}_{n}^{k}\mathfrak{X}_{n}(t) = -\left[\mathcal{M}_{n}^{k}, \mathcal{L}_{n}\right]\mathfrak{X}_{n}(t) + \mathcal{M}_{n}^{k+1}\mathfrak{X}_{n}(t) + \mathcal{M}_{n}^{k}\mathfrak{Y}_{n}$$
$$= -\sum_{j=0}^{k-1} {\binom{k}{j}} \operatorname{Ad}_{\mathcal{M}_{n}}^{k-j}(\widehat{\mathcal{L}}_{n}) \mathcal{M}_{n}^{j}\mathfrak{X}_{n}(t) + \mathcal{M}_{n}^{k+1}\mathfrak{X}_{n}(t) + \mathcal{M}_{n}^{k}\mathfrak{Y}_{n}.$$
(8.50)

Define $\mathscr{Z}_k(t) = \mathscr{M}_n^k \mathscr{Z}_n(t)$, and denote by Y(t,i) the i-th component of the vector Y(t). Then applying (8.36) with (8.50) we deduce for any $k \in \{1, ..., n\}$

$$\begin{aligned} \|\mathscr{Z}_{k}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} &\lesssim e^{Ct} \|\rho\|_{W^{1,\infty}} \left(\int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \sum_{j=0}^{k-1} \| \left(\operatorname{Ad}_{\mathscr{M}_{n}}^{k-j}(\widehat{\mathscr{L}}_{n}) \,\mathscr{M}_{n}^{j} \mathscr{X}_{n} \right)(\tau,i) \|_{L^{2}(\mathbb{T}^{d+2})} d\tau \\ &+ \int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \, \|\mathscr{Z}_{k+1}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau + t \| \mathscr{M}_{n}^{k} \mathscr{Y}_{n}(i) \|_{L^{2}(\mathbb{T}^{d+2})} \right) \end{aligned}$$
(8.51)

and for k = 0 we infer

$$\|\mathscr{Z}_{0}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim e^{Ct\|\rho\|_{W^{1,\infty}}} \left(\int_{0}^{t} e^{-C\tau\|\rho\|_{W^{1,\infty}}} \|\mathscr{Z}_{1}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau + t \|\mathscr{Y}_{n}(i)\|_{L^{2}(\mathbb{T}^{d+2})} \right).$$
(8.52)

At this level we shall make appeal to Lemma 8.3 allowing to get for any $i \in \{2, .., n + 1\}$

$$\left\| \left(\operatorname{Ad}_{\mathcal{M}_{n}}^{k-j}(\mathcal{L}_{n}) \,\mathcal{M}_{n}^{j} \mathcal{X}_{n} \right)(\tau, i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{H^{s_{0}+3}(\mathbb{T}^{d+1})}^{k-j} \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_{0}+3+i-\ell}(\mathbb{T}^{d+1})} \|\mathcal{L}_{j}(\tau, \ell)\|_{L^{2}(\mathbb{T}^{d+2})}.$$

Notice that for i = 1 the first component $\left(\operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\mathcal{L}_n)\mathcal{M}_n^j\mathcal{X}_n\right)(\tau, 1)$ is vanishing because

$$\operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\mathcal{L}_n)\,\mathcal{M}_n^j\in\mathfrak{A}_1(n)$$

which follows from Lemma 8.1. Inserting the foregoing estimate into (8.51) yields

$$\begin{aligned} \|\mathscr{Z}_{k}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} &\lesssim e^{Ct} \|\rho\|_{W^{1,\infty}} \left(\int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \|\mathscr{Z}_{k+1}(t,i)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau + t \|\mathscr{M}_{n}^{k}\mathscr{Y}_{n}(i)\|_{L^{2}(\mathbb{T}^{d+2})} \right. \\ &+ \int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \sum_{j=0}^{k-1} \|\rho\|_{H^{s_{0}+3}(\mathbb{T}^{d+1})}^{k-j} \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_{0}+3+i-\ell}(\mathbb{T}^{d+1})} \|\mathscr{Z}_{j}(\tau,\ell)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau \right). \end{aligned}$$
(8.53)

Let us introduce the quantity

$$\zeta(t,i) \triangleq e^{-Ct \|\rho\|_{W^{1,\infty}}} \sum_{k=0}^{n} \|\mathscr{Z}_k(t,i)\|_{L^2}.$$

Then summing up (8.52) and (8.53) over k = 1, ..., n and using the relation $\mathcal{M}_n^{n+1} = 0$, giving $\mathcal{Z}_{n+1}(t) = 0$, we find

$$\begin{split} \zeta(t,i) \lesssim &\int_{0}^{t} e^{-C\tau \|\rho\|_{W^{1,\infty}}} \sum_{k=1}^{n} \sum_{j=0}^{k-1} \|\rho\|_{H^{s_{0}+3}}^{k-j} \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_{0}+3+i-\ell}} \|\mathscr{L}_{j}(\tau,\ell)\|_{L^{2}(\mathbb{T}^{d+2})} d\tau \\ &+ \int_{0}^{t} \zeta(\tau,i)\|_{L^{2}} d\tau + t \sum_{k=0}^{n} \|\mathscr{M}_{n}^{k} \mathscr{Y}_{n}(i)\|_{L^{2}}. \end{split}$$

Consequently, we get from Fubini's principle that for $i \in \{1, .., n + 1\}$

$$\begin{split} \zeta(t,i) \lesssim & \left(\|\rho\|_{H^{s_0+3}} + \|\rho\|_{H^{s_0+3}}^n \right) \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_0+3+i-\ell}} \int_0^t \zeta(\tau,\ell) d\tau \\ & + \int_0^t \zeta(\tau,i) \|_{L^2} d\tau + t \sum_{k=0}^n \|\mathcal{M}_n^k \mathcal{Y}_n(i)\|_{L^2}. \end{split}$$

Notice that for i = 1 the first sum part in the right-hand side disappears. Then Gronwall inequality gives

$$\zeta(t,i) \leqslant C_0 \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_0+3+i-\ell}} \int_0^t e^{C(t-\tau)} \zeta(\tau,\ell) d\tau + C e^{Ct} \sum_{k=0}^n \|\mathcal{M}_n^k \mathcal{Y}_n(i)\|_{L^2},$$
(8.54)

with

$$C_0 \triangleq C(1 + \|\rho\|_{H^{s_0+3}}^n).$$

For i = 1 we get

$$\zeta(t,1) \leqslant C e^{Ct} \sum_{k=0}^{n} \|\mathcal{M}_{n}^{k} \mathcal{Y}_{n}(1)\|_{L^{2}}.$$
(8.55)

From Lemma 8.5 we have

$$\|\mathcal{M}_n^k \mathcal{Y}_n(i)\|_{L^2(\mathbb{T}^{d+2})} \leqslant C \left(1 + G^k(0)\right) G(i), \tag{8.56}$$

with

$$G(i) \triangleq F_1(s_0 + 5 + i) + \|\rho\|_{H^{s_0 + 6 + |m| + i}},$$
(8.57)

where F_1 should be an increasing log – convex function such that

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow |||[\mathbb{A}, \mathcal{R}_0]|||_{m, s, \gamma} \le F_1(s + \gamma), \tag{8.58}$$

for some fixed $m < -\frac{1}{2}$. The next goal is devoted to an explicit construction of F_1 . According to Lemma 7.5-(ii) applied with q = 0 one gets

$$\begin{aligned} \|\|[\mathbb{A}, \mathcal{R}_{0}]\|\|_{\alpha-1, s, \gamma} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathbb{A}\|\|_{\alpha, s, 1+\beta} \|\|\mathcal{R}_{0}\|\|_{0, s_{0}+2, \gamma-\beta} + \|\|\mathbb{A}\|\|_{\alpha, s_{0}, 1+\beta} \|\|\mathcal{R}_{0}\|\|_{0, s+2, \gamma-\beta} \\ + \sum_{0 \le \beta \le \gamma} \|\|\mathbb{A}\|\|_{\alpha, s, 0} \|\|\mathcal{R}_{0}\|\|_{0, s_{0}+2+\beta, \gamma-\beta} + \|\|\mathbb{A}\|\|_{0, s_{0}, 0} \|\|\mathcal{R}_{0}\|\|_{0, s+2+\beta, \gamma-\beta} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathcal{R}_{0}\|\|_{0, s, 1+\beta} \|\|\mathbb{A}\|\|_{\alpha, s_{0}+2, \gamma-\beta} + \|\|\mathcal{R}_{0}\|\|_{0, s_{0}, 1+\beta} \|\|\mathcal{A}\|\|_{\alpha, s+2, \gamma-\beta} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathcal{R}_{0}\|\|_{0, s, 0} \|\|\mathbb{A}\|\|_{\alpha, s_{0}+2+\beta, \gamma-\beta} + \|\|\mathcal{R}_{0}\|\|_{0, s_{0}, 0} \|\|\mathbb{A}\|\|_{\alpha, s+2+\beta, \gamma-\beta} . \end{aligned}$$

$$(8.59)$$

Using (7.30) one gets for $\alpha \in (0, 1)$

$$\|\|\mathbb{A}\|\|_{\alpha,s,\gamma} \lesssim \|\rho\|_{H^{s+\gamma+1}}.$$
(8.60)

Since \mathcal{R}_0 satisfies

 $\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 3 \Longrightarrow |||\mathscr{R}_0|||_{m,s,\gamma} \le F_0(s + \gamma),$

then plugging this into (8.59) allows to get for any $s \ge s_0, \gamma \in \mathbb{N}$ with $s + \gamma \le s_0 + n + 1$

$$\begin{split} \|\|[\mathbb{A},\mathcal{R}_{0}]\|\|_{\alpha-1,s,\gamma} &\lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s+\beta+2}} F_{0}(s_{0}+2+\gamma-\beta) + \|\rho\|_{H^{s_{0}+\beta+2}} F_{0}(s+2+\gamma-\beta) \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s+1}} F_{0}(s_{0}+2+\gamma) + \|\rho\|_{H^{s_{0}+1}} F_{0}(s+2+\gamma) \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s_{0}+3+\gamma-\beta}} F_{0}(s+1+\beta) + \|\rho\|_{H^{s+3+\gamma-\beta}} F_{0}(s_{0}+1+\beta) \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s_{0}+3+\gamma}} F_{0}(s) + \|\rho\|_{H^{s+3+\gamma}} F_{0}(s_{0}). \end{split}$$

Applying Sobolev embeddings and the monotonicity of F_0 we obtain

$$\|[\mathbb{A}, \mathcal{R}_0]\|_{\alpha-1, s, \gamma} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s+\beta+3}} F_0(s_0 + 2 + \gamma - \beta) + \|\rho\|_{H^{s_0+\beta+3}} F_0(s + 2 + \gamma - \beta).$$
(8.61)

By virtue of the log - convexity of Sobolev norms and the function F_0 we infer

$$\begin{aligned} \|\rho\|_{H^{s+\beta+3}}F_{0}(s_{0}+2+\gamma-\beta) &\leq \|\rho\|_{H^{s_{0}}}^{\delta} \|\rho\|_{H^{s+\gamma+5}}^{1-\delta}F_{0}^{1-\delta}(s_{0})F_{0}^{\delta}(s+5+\gamma) \\ &\leq F_{0}(s_{0})\|\rho\|_{H^{s+\gamma+5}} + F_{0}(s+5+\gamma)\|\rho\|_{H^{s_{0}}} \\ &\leq \left(\|\rho\|_{H^{s_{0}}} + F_{0}(s_{0})\right) \left(\|\rho\|_{H^{s+\gamma+5}} + F_{0}(s+5+\gamma)\right), \end{aligned}$$

$$(8.62)$$

with $\delta \in (0, 1)$. Therefore we obtain

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow |||[\mathbb{A}, \mathcal{R}_0]|||_{\alpha - 1, s, \gamma} \leqslant F_1(s + \gamma), \tag{8.63}$$

with

$$F_1(s) \triangleq C\big(\|\rho\|_{H^{s_0}} + F_0(s_0)\big)\big(\|\rho\|_{H^{s+5}} + F_0(s+5)\big).$$
(8.64)

Coming back to (8.58), one may fix from (8.64) the function F_1 with the constraint $m = \alpha - 1 < -\frac{1}{2}$ which is satisfied since $\alpha \in [0, \frac{1}{2})$. Notice that F_1 is log-convex since the sum of log-convex functions is \log -convex too. Inserting (8.64) into (8.57) allows to make the choice

$$G(i) = C(\|\rho\|_{H^{s_0}} + F_0(s_0))(\|\rho\|_{H^{s_0+11+i}} + F_0(s_0+11+i)) + \|\rho\|_{H^{s_0+7+i}}.$$
(8.65)

$$\forall 1 \leq i \leq k, \quad (\mathcal{M}_n^k \mathcal{Y}_n)(i) = 0.$$

Combining this with (8.56) we find

$$\sum_{k=0}^{n} \|\mathcal{M}_{n}^{k} \mathcal{Y}_{n}(i)\|_{L^{2}(\mathbb{T}^{d+2})} = \sum_{k=0}^{i-1} \|\mathcal{M}_{n}^{k} \mathcal{Y}_{n}(i)\|_{L^{2}(\mathbb{T}^{d+2})}$$
$$\leqslant C (1 + G^{i}(0)) G(i).$$
(8.66)

Now plugging (8.66) into (8.55) allows to get

$$\zeta(t,1) \leq Ce^{Ct} (1+G(0))G(1).$$
 (8.67)

We intend to prove by induction that for any $i \in \{1, .., n+1\}$,

$$\zeta(t,i) \leq 2C \left(1 + G^i(0) \right) G(i) e^{\lambda C_0 t} \quad \text{with} \quad C_0 = C \left(1 + \|\rho\|_{H^{s_0+3}}^n \right), \tag{8.68}$$

where $\lambda \ge 1$ to be fixed later. According to (8.67) this is true for i = 1. Now assume that (8.68) is true from 1 up to i-1 and let us check that it remains true at the order i. According to (8.54) and (8.66) we may write

$$\begin{split} \zeta(t,i) &\leqslant C \big(1 + G^i(0) \big) G(i) e^{Ct} \\ &+ 2C (\lambda - 1)^{-1} \, e^{\lambda C_0 t} \sum_{\ell=1}^{i-1} \big(1 + G^\ell(0) \big) \|\rho\|_{H^{s_0 + 3 + i - \ell}} G(\ell). \end{split}$$

Using interpolation inequality, since G defined in (8.65) is log – convex, we get for $\ell \in \{0, ..., i\}$

$$\begin{aligned} \|\rho\|_{H^{s_0+3+i-\ell}}G(\ell) &\leq G(i-\ell)G(\ell) \\ &\leq G(0)G(i). \end{aligned}$$

Therefore we obtain for any $\lambda \ge 1$

$$\zeta(t,i) \leqslant C \left(1 + G^i(0) \right) G(i) e^{\lambda C_0 t} + 2C(\lambda - 1)^{-1} i e^{\lambda C_0 t} \left(1 + G^i(0) \right) G(i).$$

By fixing $\lambda = 1 + 2n$, we find the estimate (8.68). Hence we obtain for any $i \in \{0, 1, ..., n\}$

$$\begin{aligned} \|\partial_{\chi}^{i} \mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} &= \|\mathscr{X}_{0}(t, i+1)\|_{L^{2}} \\ &\leq e^{Ct \|\rho\|_{W^{1,\infty}}} \zeta(t, i+1) \\ &\leq C_{1} (1+G^{i+1}(0)) e^{C_{1}(1+\|\rho\|_{H^{s_{0}+3}}^{n})^{t}} G(1+i). \end{aligned}$$

$$(8.69)$$

for some constant C_1 that depends only on n. This achieves the proof of (8.29) for $t \in [0, 1]$. Notice that making slight modifications of the preceding computations using in particular the second estimate in (8.62) and the second estimate of Lemma 8.5-(ii) yield instead of (8.69)

$$|\partial_{\chi}^{i}\mathcal{K}_{t}||_{L^{2}(\mathbb{T}^{d+2})} \leq C_{1}(1+G^{i}(0))e^{C_{1}(1+\|\rho\|_{H^{s_{0}+3}}^{n})t}G(1+i)F_{0}(s_{0}+11+i).$$
(8.70)

• Estimate of $\|\partial_{\varphi}^{i}\mathcal{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})}$. It can be done in in a similar way to (8.29) and to avoid redundancy we shall merely explain the main steps to estimate $\partial_{\omega}^{i} \mathcal{K}_{t}$. The goal is to check that for any $i \in$ $\{0, 1, ..., n\},\$

$$\forall t \in [0,1], \|\partial_{\varphi}^{i} \mathcal{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \leq C_{1} (1 + G^{i+1}(0)) e^{C_{1} \|\rho\|_{H^{s_{0}+3}}^{n}} G(1+i).$$

$$(8.71)$$

Proceeding as for (8.44) we may write

$$\mathscr{L}\partial_{\varphi}^{i}\mathscr{K}_{t} = \sum_{k=0}^{i-1} \widehat{m}_{i-k,i} \,\partial_{\varphi}^{k}\mathscr{K}_{t} + \partial_{\varphi}^{i}K_{0}$$

with

$$\widehat{m}_{k,i} \triangleq \begin{pmatrix} i \\ k \end{pmatrix} \left(\widehat{\mathbb{B}}_k - \mathbb{T}_k \right)$$

$$\widehat{\mathbb{B}}_k = \mathbb{B}_{\theta,(k)} + \mathbb{B}_{\eta,(k)}, \quad \mathbb{B}_{\theta,(k)} \triangleq \partial_{\theta} \left((\partial_{\varphi}^k \rho) |\mathcal{D}_{\theta}|^{\alpha - 1} + |\mathcal{D}_{\theta}|^{\alpha - 1} (\partial_{\varphi}^k \rho) \right)$$
(8.72)

and

$$\mathbb{T}_{k} \triangleq (\partial_{\eta} \partial_{\varphi}^{k} \rho) |\mathcal{D}_{\eta}|^{\alpha - 1} + |\mathcal{D}_{\eta}|^{\alpha - 1} (\partial_{\eta} \partial_{\varphi}^{k} \rho).$$

Next, we introduce the matrix operator

$$\widehat{\mathcal{M}}_{n} = \begin{pmatrix} 0 & 0 & \dots & \dots & 0 \\ \widehat{m}_{1,1} & 0 & \dots & \dots & 0 & 0 \\ \widehat{m}_{2,2} & \widehat{m}_{1,2} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \widehat{m}_{n,n} & \widehat{m}_{n-1,n} & \dots & \dots & \widehat{m}_{1,n} & 0 \end{pmatrix}$$

and

$$\widehat{\mathscr{X}}_n(t) = \begin{pmatrix} \mathscr{K}_t \\ \partial_{\varphi} \mathscr{K}_t \\ \vdots \\ \vdots \\ \partial_{\varphi}^n \mathscr{K}_t \end{pmatrix}, \quad \widehat{\mathscr{Y}}_n = \begin{pmatrix} K_0 \\ \partial_{\varphi} K_0 \\ \vdots \\ \vdots \\ \partial_{\varphi}^n K_0 \end{pmatrix}.$$

Then (8.44) can be written in the matrix form as follows

$$\mathscr{L}_n\widehat{\mathscr{X}}_n(t) = \widehat{\mathscr{M}}_n\,\widehat{\mathscr{X}}_n(t) + \widehat{\mathscr{Y}}_n, \quad \mathscr{L}_n \triangleq \mathscr{L}\mathrm{I}_{n+1}.$$

Then we get the same structure as for \mathscr{X}_n and therefore we may implement exactly the same approach to deduce the estimate (8.71). This achieves the estimate of the Theorem 8.1-(i) in the case m = 0.

▶ Case $m \in [-1, 0)$. We shall first remove from $\Re(t)$ the first terms given by Taylor formula, that is, to consider the operator

$$\mathscr{R}_{2}(t) \triangleq \mathscr{R}(t) - \sum_{k=0}^{2} \operatorname{Ad}_{\mathbb{A}}^{k} \mathscr{R}_{0} \frac{t^{k}}{k!},$$

then it is straightforward to check

$$\left\{ \begin{array}{l} \partial_t \mathscr{R}_2(t) = [\mathbb{A}, \mathscr{R}_2(t)] + \mathrm{Ad}_{\mathbb{A}}^3 \mathscr{R}_0 \frac{t^3}{3!} \\ \mathscr{R}_1(0) = 0. \end{array} \right.$$

Then to show the desired estimate it is enough to check it for $\mathscr{R}_2(t)$ and $\sum_{k=0}^2 \operatorname{Ad}_{\mathbb{A}}^k \mathscr{R}_0 \frac{t^k}{k!}$.

• Estimate of $\mathscr{R}_2(t)$. Denote by $\mathscr{K}_{1,t}$ the kernel associated to $\mathscr{R}_2(t)$ and $K_1(t)$ the kernel associated to $\mathrm{Ad}^3_{\mathbb{A}}\mathscr{R}_0\frac{t^3}{3!}$. Then using (8.28) combined with integration by parts and fixing $s = n \in \mathbb{N}$

$$\begin{aligned} \|\|\mathscr{R}_{2}(t)\|\|_{-1,s,0} &\leq \int_{\mathbb{T}} \|\partial_{\eta}\mathscr{K}_{1,t}(\cdot, \cdot, \cdot, \cdot + \eta)\|_{H^{s}_{\varphi,\theta}} d\eta \\ &\lesssim \sum_{i=0}^{n} \left(\|\partial_{\chi}^{i}\partial_{\eta}\mathscr{K}_{1,t}\|_{L^{2}(\mathbb{T}^{d+2})} + \|\partial_{\varphi}^{i}\partial_{\eta}\mathscr{K}_{1,t}\|_{L^{2}(\mathbb{T}^{d+2})} \right). \end{aligned}$$

Set

 $\mathcal{K}_{2,t} \triangleq \partial_\eta \mathcal{K}_{1,t}$ and $K_2 \triangleq \partial_\eta K_1(t)$.

Then differentiating (8.32) with respect to η and using the identity

$$\partial_{\eta} \left(-\mathbb{A}_{\eta} + \left[\partial_{\eta}, \mathcal{T}_{\eta} \right] \right) = -\mathbb{A}_{\eta} \partial_{\eta}$$

yield

$$\partial_t \mathscr{K}_{2,t}(\varphi,\theta,\eta) - \mathbb{A}_{\theta} \mathscr{K}_{2,t}(\varphi,\theta,\eta) - \mathbb{A}_{\eta} \mathscr{K}_{2,t}(\varphi,\theta,\eta) = K_2(\varphi,\theta,\eta)$$

which is similar to the equation (8.32). Notice that the operator associated to the kernel K_2 is given by $-\frac{t^3}{3!} \operatorname{Ad}_{\mathbb{A}}^3 \mathscr{R}_0 \partial_{\theta}$. Then from (8.69), (8.57) and (8.58) we deduce that

$$\forall t \in [0,1], \quad \|\partial_{\chi}^{i} \mathscr{K}_{2,t}\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant C_{1} (1 + G_{1}^{i+1}(0)) e^{C_{1} \|\rho\|_{H^{s_{0}+3}}^{n} t} G_{1}(1+i), \tag{8.73}$$

with

$$G_1(i) \triangleq F(s_0 + 5 + i) + \|\rho\|_{H^{s_0 + 6 + |m| + i}}, \tag{8.74}$$

where F should be an increasing log – convex function such that

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow \||\operatorname{Ad}^3_{\mathbb{A}} \mathscr{R}_0 \partial_\theta||_{m, s, \gamma} \le F(s + \gamma), \tag{8.75}$$

for some fixed $m < -\frac{1}{2}$. The next goal is devoted to an explicit construction of F. According to Lemma 7.5-(i) one has for any $s \ge s_0, \gamma \in \mathbb{N}$

$$\|\|\mathrm{Ad}^{3}_{\mathbb{A}}\mathscr{R}_{0}\partial_{\theta}\|\|_{3\alpha-2,s,\gamma} \lesssim \||\mathrm{Ad}^{3}_{\mathbb{A}}\mathscr{R}_{0}|\|_{3\alpha-3,s,\gamma}$$

Using Lemma 7.5-(ii) combined with (8.60), (8.63), (8.64) and $\alpha \in (0, \frac{1}{2})$

$$\begin{split} \|\|\mathrm{Ad}_{\mathbb{A}}^{2}\mathscr{R}_{0}\|\|_{2\alpha-2,s,\gamma} &\lesssim \sum_{0\leqslant\beta\leqslant\gamma} \|\rho\|_{H^{s+\beta+2}} F_{1}(s_{0}+2+\gamma-\beta) + \|\rho\|_{H^{s_{0}+\beta+2}} F_{1}(s+2+\gamma-\beta) \\ &+ \|\rho\|_{H^{s+1}} F_{1}(s_{0}+2+\gamma) + \|\rho\|_{H^{s_{0}+1}} F_{1}(s+2+\gamma) \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\rho\|_{H^{s_{0}+3+\gamma-\beta}} F_{1}(s+\beta+1) + \|\rho\|_{H^{s+3+\gamma-\beta}} F_{1}(s_{0}+\beta+1) \\ &+ \|\rho\|_{H^{s_{0}+4+\gamma}} F_{1}(s) + \|\rho\|_{H^{s+4+\gamma}} F_{1}(s_{0}). \end{split}$$

Thus by Sobolev embeddings we infer

$$\|\|\mathrm{Ad}_{\mathbb{A}}^{2}\mathcal{R}_{0}\|\|_{2\alpha-2,s,\gamma} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho\|_{H^{s+\beta+4}} F_{1}(s_{0}+2+\gamma-\beta) + \|\rho\|_{H^{s_{0}+\beta+4}} F_{1}(s+2+\gamma-\beta).$$

Then using the log – convexity of Sobolev norms and F_1 we find for $s \ge s_0$ and $0 \le \beta \le \gamma$,

$$\begin{aligned} \|\rho\|_{H^{s+\beta+4}}F_0(s_0+2+\gamma-\beta) &\leq \|\rho\|_{H^{s_0}}^{\delta}\|\rho\|_{H^{s+\gamma+6}}^{1-\delta}F_0^{1-\delta}(s_0)F_0^{\delta}(s+6+\gamma) \\ &\leq \Big(\|\rho\|_{H^{s_0}}+F_1(s_0)\Big)\Big(\|\rho\|_{H^{s+\gamma+6}}+F_1(s+6+\gamma)\Big) \end{aligned}$$

with $\delta \in (0, 1)$. By setting

$$F_2(s) \triangleq C\Big(\|\rho\|_{H^{s_0}} + F_1(s_0)\Big)\Big(\|\rho\|_{H^{s+6}} + F_1(s+6)\Big)$$
(8.76)

we get that F_2 is $\log -$ convex and

$$\|\|\operatorname{Ad}^{2}_{\mathbb{A}}\mathscr{R}_{0}\|\|_{2\alpha-2,s,\gamma} \leqslant F_{2}(s+\gamma).$$

$$(8.77)$$

Implementing the same approach we also get

$$\||\mathrm{Ad}^{3}_{\mathbb{A}}\mathscr{R}_{0}|||_{3\alpha-3,s,\gamma} \lesssim F_{3}(s+\gamma)$$
(8.78)

with

$$F_3(s) \triangleq C\Big(\|\rho\|_{H^{s_0}} + F_2(s_0)\Big)\Big(\|\rho\|_{H^{s+7}} + F_2(s+7)\Big).$$
(8.79)

Putting together (8.64), (8.76) and (8.79) implies

$$F_3(s) \leqslant C \Big(1 + \|\rho\|_{H^{s_0+17}}^3 + F_2^3(s_0+17) \Big) \Big(\|\rho\|_{H^{s+17}} + F_0(s+17) \Big).$$

Define

$$F(s) \triangleq C\left(1 + \|\rho\|_{H^{s_0+17}}^3 + F_0^3(s_0 + 17)\right) \left(\|\rho\|_{H^{s+17}} + F_0(s + 17)\right)$$

we get that F is log – convex and for any $\alpha \in (0, \frac{1}{2})$

$$\|\|\operatorname{Ad}^{3}_{\mathbb{A}}\mathcal{R}_{0}\|_{3\alpha-3,s,\gamma} \leqslant F(s+\gamma).$$

This gives (8.75) with $m = 3\alpha - 3$ and the assumption $m < -\frac{1}{2}$ is satisfied when $\alpha < \frac{1}{2}$. From (8.74) we infer that

$$G_{1}(i) \lesssim \|\rho\|_{H^{s_{0}+9+i}} + F(s_{0}+9+i)$$

$$\lesssim \|\rho\|_{H^{s_{0}+9+i}} + \left(1 + \|\rho\|_{H^{s_{0}+17}}^{3} + F_{0}^{3}(s_{0}+17)\right) \left(\|\rho\|_{H^{s_{0}+26+i}} + F_{0}(s_{0}+26+i)\right).$$

Inserting this inequality into (8.73) yields in view of Sobolev embeddings and for $t \in [0, 1]$,

$$\begin{aligned} \|\partial_{\chi}^{i}\mathcal{H}_{2,t}\|_{L^{2}} &\leq C \big(1 + F_{0}^{4(s+1)}(s_{0} + 27)\big) e^{C \|\rho\|_{H^{s_{0}+27}}^{s}} \Big(F_{0}(s + 27 + s_{0}) + \|\rho\|_{H^{s+27+s_{0}}}\Big) \\ &\leq C e^{C \mu^{s}(0)} \,\mu(s), \end{aligned}$$

with

$$\mu(s) \triangleq F_0(s+27+s_0) + \|\rho\|_{H^{s+27+s_0}}.$$

• Estimate of $\sum_{k=0}^{2} \operatorname{Ad}_{\mathbb{A}}^{k} \mathscr{R}_{0} \frac{t^{k}}{k!}$. First for k = 0 we know by assumption that

$$\||\operatorname{Ad}^{0}_{\mathbb{A}}\mathscr{R}_{0}||_{m,s,\gamma} = |||\mathscr{R}_{0}||_{m,s,\gamma} \leqslant F_{0}(s+\gamma).$$

Concerning k = 2 we use (8.77) which gives, since $2\alpha - 2 \leq -1 \leq m$,

$$\begin{aligned} \|\operatorname{Ad}_{\mathbb{A}}^{2} \mathcal{R}_{0}\|_{m,s,\gamma} &\leq \|\operatorname{Ad}_{\mathbb{A}}^{2} \mathcal{R}_{0}\|_{2\alpha-2,s,\gamma} \\ &\leq F_{2}(s+\gamma). \end{aligned}$$

As to case k = 1 we proceed similarly to (8.59). Indeed, according to Lemma 7.5-(ii) applied with q = 0 one gets for $m \in [-1, 0]$

$$\begin{split} \|\|[\mathbb{A},\mathcal{R}_{0}]\|\|_{\alpha+m-1,s,\gamma} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathbb{A}\|\|_{\alpha,s,1+\beta} \|\|\mathcal{R}_{0}\|\|_{m,s_{0}+2,\gamma-\beta} + \|\|\mathbb{A}\|\|_{\alpha,s_{0},1+\beta} \|\|\mathcal{R}_{0}\|\|_{m,s+2,\gamma-\beta} \\ &+ \sum_{0 \le \beta \le \gamma} \|\|\mathbb{A}\|\|_{\alpha,s,0} \|\|\mathcal{R}_{0}\|\|_{m,s_{0}+2+\beta,\gamma-\beta} + \|\|\mathbb{A}\|\|_{0,s_{0},0} \|\|\mathcal{R}_{0}\|\|_{m,s+2+\beta,\gamma-\beta} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathcal{R}_{0}\|\|_{m,s,1+\beta} \|\|\mathbb{A}\|\|_{\alpha,s_{0}+2,\gamma-\beta} + \|\|\mathcal{R}_{0}\|\|_{m,s_{0},1+\beta} \|\|\mathcal{A}\|\|_{\alpha,s+2,\gamma-\beta} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\|\mathcal{R}_{0}\|\|_{0,s,0} \|\|\mathbb{A}\|\|_{\alpha,s_{0}+3+\beta,\gamma-\beta} + \|\|\mathcal{R}_{0}\|\|_{0,s_{0},0} \|\|\mathbb{A}\|\|_{\alpha,s+3+\beta,\gamma-\beta} \,. \end{split}$$

Therefore implementing the same arguments as for getting (8.63) we obtain

$$\|\|[\mathbb{A}, \mathcal{R}_0]\|\|_{m, s, \gamma} \leq \|\|[\mathbb{A}, \mathcal{R}_0]\|\|_{\alpha + m - 1, s, \gamma}$$
$$\leq F_1(s + \gamma + 1)$$

where F_1 is defined in (8.64).

(ii) The case q = 0 has been done in the first point (i). We observe from (8.28) that for any $0 \leq j \leq q$ and for $s = n \in \mathbb{N}$

$$\begin{aligned} \||\partial_{\lambda}^{j}\mathscr{R}_{1}(t)|\|_{0,s,0} &\leq \int_{\mathbb{T}} \|\partial_{\lambda}^{j}\mathscr{K}_{t}(\cdot, \cdot, \cdot + \eta)\|_{H^{s}_{\varphi,\theta}} d\eta \\ &\lesssim \sum_{i=0}^{n} \left(\|\partial_{\lambda}^{j}\partial_{\chi}^{i}\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} + \||\partial_{\lambda}^{j}\partial_{\varphi}^{i}\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \right). \end{aligned}$$

The estimates of $\partial_{\lambda}^{j} \partial_{\chi}^{i} \mathcal{K}_{t}$ and $\partial_{\lambda}^{j} \partial_{\varphi}^{i} \mathcal{K}_{t}$ can be done in a similar way. Therefore we shall restrict the discussion to the estimate of $\partial_{\lambda}^{j} \partial_{\chi}^{i} \mathcal{K}_{t}$.

• Estimate of $\|\partial_{\lambda}^{j}\partial_{\chi}^{i}\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})}, 0 \leq j \leq q, 0 \leq i \leq n$. By applying ∂_{λ}^{j} to the equation (8.37) we get after straightforward algebraic computations

$$\kappa^{j} \mathscr{L} \partial_{\lambda}^{j} \mathscr{K}_{t} = \sum_{k=0}^{j-1} \overline{m}_{j-k,j} \,\kappa^{k} \partial_{\lambda}^{k} \mathscr{K}_{t} + \kappa^{j} \partial_{\lambda}^{j} K_{0}, \qquad (8.80)$$

where

$$\overline{m}_{k,j} \triangleq \begin{pmatrix} j \\ k \end{pmatrix} \left(\overline{\mathbb{B}}_k - \overline{\mathbb{T}}_k \right)$$
(8.81)

with

$$\overline{\mathbb{B}}_k = \overline{\mathbb{B}}_{\theta,(k)} + \overline{\mathbb{B}}_{\eta,(k)}, \quad \overline{\mathbb{B}}_{\theta,(k)} = \kappa^k \big[\partial_{\lambda}^k, \mathbb{A}_{\theta} \big]$$

and

$$\overline{\mathbb{T}}_k \triangleq \kappa^k \big[\partial_\lambda^k, \mathscr{S}_0 \big].$$

Recall that the operator \mathcal{S}_0 is defined in (8.33). Set

$$X_{q} = \begin{pmatrix} \mathcal{K}_{t} \\ \gamma \partial_{\lambda} \mathcal{K}_{t} \\ \vdots \\ \gamma^{q} \partial_{\lambda}^{q} \mathcal{K}_{t} \end{pmatrix}, \quad Y_{q} = \begin{pmatrix} K_{0} \\ \gamma \partial_{\lambda} K_{0} \\ \vdots \\ \gamma^{q} \partial_{\lambda}^{q} K_{0} \end{pmatrix}$$

and
$$\overline{M}_{q} = \begin{pmatrix} 0 & 0 & \vdots & \vdots & \ddots & 0 \\ \overline{m}_{1,1} & 0 & \vdots & \vdots & \ddots & 0 \\ \overline{m}_{2,2} & \overline{m}_{1,2} & 0 & \vdots & \vdots & 0 \\ \vdots & \vdots & \vdots & \vdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \overline{m}_{1,q-1} & 0 & 0 \\ \overline{m}_{q,q} & \overline{m}_{q-1,q} & \vdots & \vdots & \overline{m}_{1,q} & 0 \end{pmatrix}.$$
(8.82)

Then (8.80) can be written in the matrix form

$$L_q X_q(t) = \overline{M}_q X_q + Y_q, \quad L_q \triangleq \mathscr{L} \mathbf{I}_{q+1}.$$
(8.83)

Define

$$\overline{\mathcal{X}}_{n}(t) = \begin{pmatrix} X_{q} \\ \partial_{\chi} X_{q} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n} X_{q} \end{pmatrix} \quad \text{and} \quad \overline{\mathcal{Y}}_{n} = \begin{pmatrix} Y_{q} \\ \partial_{\chi} Y_{q} \\ \vdots \\ \vdots \\ \partial_{\chi}^{n} Y_{q} \end{pmatrix}.$$
(8.84)

Therefore successive differentiation of (8.83) yields to the matrix form

$$\overline{\mathscr{L}}_n \overline{\mathscr{X}}_n(t) = \overline{\mathscr{M}}_n \overline{\mathscr{X}}_n(t) + \overline{\mathscr{Y}}_n, \quad \overline{\mathscr{L}}_n \triangleq L_q \mathbb{I}_{n+1}$$

with the bloc matrix operator

$$\overline{\mathcal{M}}_{n} = \begin{pmatrix} \overline{M}_{q} & 0 & \dots & \dots & \dots & 0 \\ \overline{M}_{1,1} & \overline{M}_{q} & 0 & \dots & \dots & 0 \\ \overline{M}_{2,2} & \overline{M}_{1,2} & \overline{M}_{q} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \overline{M}_{1,n-1} & \overline{M}_{q} & 0 \\ \overline{M}_{n,n} & \overline{M}_{n-1,n} & \dots & \dots & \overline{M}_{1,n} & \overline{M}_{q} \end{pmatrix} \triangleq \overline{M}_{q} \mathbb{I}_{n+1} + N_{n} \in \Delta_{0}(n,q).$$
(8.85)

and

$$\overline{M}_{k,i} \triangleq \begin{pmatrix} i \\ k \end{pmatrix} \left(\operatorname{Ad}_{\partial_{\chi}}^{k} \overline{M}_{q} - \operatorname{Ad}_{\partial_{\chi}}^{k} \overline{L}_{q} \right), \ \overline{L}_{q} = \widehat{\mathscr{L}} \operatorname{I}_{q+1},$$
(8.86)

where $\widehat{\mathscr{L}}$ was defined in (8.35). Since $\overline{M}_q^{q+1} = 0$ and $N_n^{n+1} = 0$, then

$$\overline{\mathcal{M}}_{n}^{q+n+1} = 0. \tag{8.87}$$

Indeed, we observe that $\overline{\mathcal{M}}_n^{q+n+1}$ can be expanded as the sum of elements in the form

$$\prod_{i=1}^{m} \overline{M}_{q}^{\alpha_{i}} \mathbb{I}_{n+1} N_{n}^{\beta_{i}} \quad \text{with} \quad \sum_{\substack{i=1\\96}}^{m} (\alpha_{i} + \beta_{i}) = n + q + 1.$$

According to Lemma 8.1, since $N_n \in \Delta_1(n,q)$ then $N_n^{\beta_i} \in \Delta_{\beta_i}(n,q)$ and $\overline{M}_q^{\alpha_i} \mathbb{I}_{n+1} N_n^{\beta_i} \in \Delta_{\beta_i}(n,q)$. The same lemma gives

$$P_m \triangleq \prod_{i=1}^m \overline{M}_q^{\alpha_i} \mathbb{I}_{n+1} N_n^{\beta_i} \in \mathbb{A}_{\min(\sum_{i=1}^m \beta_i, n+1)}(n, q).$$

Consequently, if $\sum_{i=1}^{m} \beta_i \ge n+1$ then we get $P_m \in \Delta_{n+1}(n,q) = \{0\}$. In the case $\sum_{i=1}^{m} \beta_i \le n$ we have

necessary $\sum_{i=1}^{m} \alpha_i \ge q+1$. Moreover the bloc entries of P_m is the sum of terms in the form

$$\prod_{i=1}^{m} \overline{M}_{q}^{\alpha_{i}} C_{n,i}, \quad \text{with} \quad C_{n,i} \in \Delta_{0}(q).$$

Since $\overline{M}_q \in \Delta_1(q)$ then applying once again Lemma 8.1 we find successively that $\overline{M}_q^{\alpha_i} \in \Delta_{\min(\alpha_i,q+1)}(q)$, $\overline{M}_q^{\alpha_i}C_{n,i} \in \Delta_{\min(\alpha_i,q+1)}(q)$ and

$$\prod_{i=1}^{m} \overline{M}_{q}^{\alpha_{i}} C_{n,i} \in \mathbf{\Delta}_{\min(\sum_{i=1}^{m} \alpha_{i}, q+1)}(q) = \mathbf{\Delta}_{q+1}(q) = \{0\}$$

Thus we get that $P_m = 0$ and this achieves the proof of (8.87). We point out that similar arguments give the following result

$$\forall k \in \{0, ..., n+1\}, \quad \overline{\mathcal{M}}_n^{q+k} \in \Delta_k(n, q).$$

$$(8.88)$$

Now, in a similar way to (8.50) we deduce for any $k \in \{1, .., n+q\}$

$$\overline{\mathscr{L}}_{n}\overline{\mathscr{M}}_{n}^{k}\overline{\mathscr{X}}_{n}(t) = -\left[\overline{\mathscr{M}}_{n}^{k},\overline{\mathscr{L}}_{n}\right]\mathscr{X}_{n}(t) + \overline{\mathscr{M}}_{n}^{k+1}\overline{\mathscr{X}}_{n}(t) + \overline{\mathscr{M}}_{n}^{k}\overline{\mathscr{Y}}_{n} \\
= -\sum_{j=0}^{k-1} \binom{k}{j} \operatorname{Ad}_{\overline{\mathscr{M}}_{n}}^{k-j}(\overline{\mathscr{L}}_{n})\overline{\mathscr{M}}_{n}^{j}\overline{\mathscr{X}}_{n}(t) + \overline{\mathscr{M}}_{n}^{k+1}\overline{\mathscr{X}}_{n}(t) + \overline{\mathscr{M}}_{n}^{k}\overline{\mathscr{Y}}_{n}.$$
(8.89)

Set $\overline{\mathscr{X}}_k(t) = \overline{\mathscr{M}}_n^k \overline{\mathscr{X}}_n(t)$ and denote by $\overline{\mathscr{X}}_k(t,i)$ the i^{th} bloc vector of size q+1. Then applying the estimate (8.36) we get for any $k \in \{1, ..., n+q\}$

$$\begin{aligned} \|\overline{\mathscr{X}}_{k}(t,i)\|_{L^{2}} &\lesssim e^{Ct} \|\rho\|_{W^{1,\infty}} \left(\int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \sum_{j=0}^{k-1} \|\left(\operatorname{Ad}_{\overline{\mathscr{M}}_{n}}^{k-j}(\overline{\mathscr{X}}_{n}) \overline{\mathscr{M}}_{n}^{j} \overline{\mathscr{X}}_{n}\right)(\tau,i)\|_{L^{2}} d\tau \\ &+ \int_{0}^{t} e^{-C\tau} \|\rho\|_{W^{1,\infty}} \|\overline{\mathscr{X}}_{k+1}(t,i)\|_{L^{2}} d\tau + t \|\overline{\mathscr{M}}_{n}^{k} \overline{\mathscr{Y}}_{n}(i)\|_{L^{2}} \right) \end{aligned}$$
(8.90)

i

and for k = 0 it becomes

$$\|\overline{\mathscr{X}}_0(t,i)\|_{L^2} \lesssim e^{Ct}\|\rho\|_{W^{1,\infty}} \left(\int_0^t e^{-C\tau}\|\rho\|_{W^{1,\infty}} \|\overline{\mathscr{X}}_1(t,i)\|_{L^2} d\tau + t \|\overline{\mathscr{Y}}_n(i)\|_{L^2}\right)$$

By applying Lemma 8.3 we deduce for $i \in \{1, .., n+1\}$

$$\left\| \left(\operatorname{Ad}_{\overline{\mathcal{M}}_{n}}^{k-j}(\overline{\mathcal{L}}_{n}) \,\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\tau, i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{r} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \overline{\mathcal{Z}}_{j}(\tau, \ell) \right\|_{L^{2}(\mathbb{T}^{d+2})}.$$

Then setting

$$\overline{\zeta}(t,i) = e^{-Ct\|\rho\|_{W^{1,\infty}}} \sum_{k=0}^{n+q} \|\overline{\mathscr{Z}}_k(t,i)\|_{L^2}$$

and summing up (8.90) over k, using in a crucial way the relation (8.87), which gives $\overline{\mathscr{X}}_{n+q+1}(t) = 0$, we find

$$\begin{split} \overline{\zeta}(t,i) \lesssim &\int_0^t \sum_{k=1}^{n+q} \sum_{j=0}^{k-1} \left(\|\rho\|_{s_0+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^i \|\rho\|_{s_0+3+i-\ell}^{q,\kappa} \quad \left\| \overline{\mathscr{X}}_j(\tau,\ell) \right\|_{L^2(\mathbb{T}^{d+2})} d\tau \\ &+ \int_0^t \overline{\zeta}(\tau,i) \|_{L^2} d\tau + t \sum_{k=0}^{n+q} \|\overline{\mathscr{M}}_n^k \overline{\mathscr{Y}}_n(i)\|_{L^2}. \end{split}$$

From this we deduce that for $i \in \{1, .., n+1\}$

$$\begin{split} \overline{\zeta}(t,i) \lesssim & \left(\|\rho\|_{s_0+3}^{q,\kappa} + \left(\|\rho\|_{s_0+3}^{q,\kappa} \right)^{n+q} \right) \sum_{\ell=1}^{i} \|\rho\|_{s_0+3+i-\ell}^{q,\kappa} \int_{0}^{t} \overline{\zeta}(\tau,\ell) d\tau \\ & + \int_{0}^{t} \overline{\zeta}(\tau,i) \|_{L^2} d\tau + t \sum_{k=0}^{n+q} \|\overline{\mathcal{M}}_{n}^{k} \overline{\mathcal{Y}}_{n}(i)\|_{L^2}. \end{split}$$

Then Gronwall inequality gives

$$\overline{\zeta}(t,i) \leqslant C_0 \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_0+3+i-\ell}} \int_0^t e^{C(t-\tau)} \overline{\zeta}(\tau,\ell) d\tau + Ct e^{Ct} \sum_{k=0}^{n+q} \|\overline{\mathcal{M}}_n^k \overline{\mathcal{Y}}_n(i)\|_{L^2},$$
(8.91)

with

$$C_0 \triangleq C(1 + (\|\rho\|_{s_0+3}^{q,\kappa})^{n+q}).$$

For i = 1 the preceding inequality becomes

$$\zeta(t,1) \leqslant Cte^{C_0 t} \sum_{k=0}^{n+q} \|\overline{\mathcal{M}}_n^k \overline{\mathcal{Y}}_n(1)\|_{L^2}.$$
(8.92)

From Lemma 8.5-(ii) we have

$$\|\overline{\mathcal{M}}_{n}^{k}\overline{\mathcal{Y}}_{n}(i)\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant C\left(1+\overline{G}^{k}(0)\right)\overline{G}(i), \tag{8.93}$$

with

$$\overline{G}(i) \triangleq \overline{F}_1(s_0 + 5 + i) + \|\rho\|_{s_0 + 6 + |m| + i}^{q,\kappa},$$

where \overline{F}_1 should be an increasing log –convex function such that

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow |||[\mathbb{A}, \mathcal{R}_0]|||_{m, s, \gamma}^{q, \kappa} \le \overline{F}_1(s + \gamma),$$

for some fixed $m < -\frac{1}{2}$. The construction of F_1 can be done in a similar way to that of (8.58) and one gets similarly to (8.63) and (8.64)

$$\overline{F}_1(s) = C\left(\|\rho\|_{s_0}^{q,\kappa} + \overline{F}_0(s_0)\right) \left(\|\rho\|_{s+5}^{q,\kappa} + \overline{F}_0(s+5)\right).$$

Then we conclude using an induction principle as for (8.68).

(iii) With the notation $\Re^i(t) \triangleq \Phi_i(t) \Re_0 \Phi_i(-t) - \Re_0$ and $\Re(t) \triangleq \Re^1(t) - \Re^2(t)$ then we write from Heisenberg equations that

$$\begin{cases} \partial_t \mathscr{R}^i(t) = [\mathbb{A}^i, \mathscr{R}^i(t)] + [\mathbb{A}^i, \mathscr{R}_0] \\ \mathscr{R}(0) = 0, \end{cases}$$
(8.94)

where \mathbb{A}^i is the operator (8.25) associated to ρ_i . Therefore the kernel \mathcal{K}_t of $\mathcal{R}(t)$ satisfies in a similar way to (8.35) and (8.37)

$$\partial_t \mathcal{K}_t + \widehat{\mathscr{L}_1}(\mathcal{K}_t) = (\widehat{\mathscr{L}_2} - \widehat{\mathscr{L}_1})\mathcal{K}_t^2 + K_{1,2}, \qquad (8.95)$$

where $\widehat{\mathscr{L}_i} = -\widehat{\mathbb{A}}f + \mathscr{S}_0$ being the operator associated to ρ_i , \mathscr{K}_t^i the kernel associated to $\mathscr{R}^i(t)$ and $K_{1,2}$ the kernel associated to $[\mathbb{A}^2 - \mathbb{A}^1, \mathscr{R}_0]$. Notice that we have used that the kernel \mathscr{K}_t^i satisfies the equation

$$\partial_t \mathscr{K}_t^i + \widehat{\mathscr{L}_i}(\mathscr{K}_t^i) = K_i \tag{8.96}$$

where K_i is the kernel of $[\mathbb{A}^i, \mathcal{R}_0]$. Applying the energy estimate (8.36) with (8.95) allows to get

$$\|\mathscr{K}_{t}\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant e^{Ct\|\rho_{1}\|_{W^{1,\infty}}} \Big(\int_{0}^{t} \left\| (\widehat{\mathscr{L}_{2}} - \widehat{\mathscr{L}_{1}})\mathscr{K}_{\tau}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} d\tau + t \, \|K_{1,2}\|_{L^{2}(\mathbb{T}^{d+2})} \Big).$$
(8.97)

Then from (8.96) we deduce that

$$\left(\partial_t + \widehat{\mathscr{L}}_2\right)(\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1)\mathscr{K}_t^2 = (\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1)K_2 + \left[\widehat{\mathscr{L}}_2, (\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1)\right]\mathscr{K}_t^2.$$
(8.98)

Applying once again the energy estimate (8.36)

$$\begin{aligned} \left\| (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1}) \mathscr{K}_{t}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} &\lesssim t e^{Ct \|\rho_{2}\|_{W^{1,\infty}}} \left\| (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1}) K_{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} \\ &+ e^{Ct \|\rho_{2}\|_{W^{1,\infty}}} \int_{0}^{t} \left\| [\widehat{\mathscr{L}}_{2}, (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1})] \mathscr{K}_{\tau}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} d\tau. \end{aligned}$$

$$(8.99)$$

Applying (7.31) with $q = \gamma = 0$ and $s = s_0$ we find

$$\| [\widehat{\mathscr{L}}_{2}, (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1})] \|_{0, s_{0}, 0} \lesssim \| \rho_{2} \|_{H^{s_{0}+3}} \| \rho_{1} - \rho_{2} \|_{H^{s_{0}+3}}$$

By virtue of Lemma (7.3)-(i) we deduce that

$$\left\| \left[\widehat{\mathscr{L}}_{2}, (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1}) \right] \mathscr{K}_{t}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \|\rho_{2}\|_{H^{s_{0}+3}} \|\rho_{1} - \rho_{2}\|_{H^{s_{0}+3}} \|\mathscr{K}_{t}^{2}\|_{L^{2}(\mathbb{T}^{d+2})}.$$

$$(8.100)$$

Applying (8.36) we find

$$\|\mathscr{K}_{t}^{2}\|_{L^{2}} \lesssim \|K_{2}\|_{L^{2}} t \, e^{Ct \|\rho_{2}\|_{W^{1,\infty}}}.$$
(8.101)

Recall that K_2 is the kernel of the commutator $[\mathbb{A}^2, \mathscr{R}_0]$ where \mathbb{A}^2 is the nonlocal operator constructed in (8.25) from ρ_2 . Therefore we deduce from (8.63) and (8.64)

$$\| [\mathbb{A}, \mathcal{R}_0] \|_{\overline{\alpha} - 1, s_0, 0} \leqslant C \big(\| \rho_2 \|_{H^{s_0 + 5}} + \overline{F}_0(s_0 + 5) \big)^2, \tag{8.102}$$

Since $\overline{\alpha} - 1 < -\frac{1}{2}$, we obtain from Lemma 7.2-(iii) applied with q = 0

$$\int_{\mathbb{T}} \|K_2(\cdot, \cdot, \cdot + \eta)\|_{s_0}^2 d\eta \lesssim \|[\mathbb{A}, \mathcal{R}_0]\|_{\overline{\alpha} - 1, s, 0}^2$$
$$\lesssim \left(\|\rho_2\|_{H^{s_0 + 5}} + \overline{F}_0(s_0 + 5)\right)^4.$$

Plugging this estimate into (8.101) yields

$$\|\mathcal{K}_{t}^{2}\|_{L^{2}} \lesssim t \, e^{Ct \|\rho\|_{W^{1,\infty}}} \left(\|\rho_{2}\|_{H^{s_{0}+5}} + \overline{F}_{0}(s_{0}+5)\right)^{2}. \tag{8.103}$$

Therefore we deduce from (8.100) and Sobolev embeddings that

$$\left\| \left[\widehat{\mathscr{L}}_{2}, (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1}) \right] \mathscr{K}_{t}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim e^{Ct \|\rho_{2}\|_{H^{s_{0}+3}}} \left(\|\rho_{2}\|_{H^{s_{0}+5}} + \overline{F}_{0}(s_{0}+5) \right)^{2} \|\rho_{1} - \rho_{2}\|_{H^{s_{0}+3}}.$$
(8.104)

Let us now move to the estimate of $(\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1)K_2$. Then applying Lemma 8.4-(iii) applied with q = 0 and using (8.63) and (8.64) yield

$$\int_{\mathbb{T}} \left\| \left(\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1\right) K_2(\cdot, \bullet, \bullet + \eta) \right\|_{H^{s_0}}^2 d\eta \lesssim \overline{F}_0 \left(s_0 + 10 \right) \|\rho_1 - \rho_2\|_{H^{s_0+5}}.$$
(8.105)

Putting together (8.105), (8.100) and (8.99), and using Sobolev embeddings give for $t \in [0, 1]$

$$\left\| (\widehat{\mathscr{L}}_{2} - \widehat{\mathscr{L}}_{1}) \mathscr{K}_{t}^{2} \right\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant C e^{C \left(\|\rho_{2}\|_{H^{s_{0}+10}} + \overline{F}_{0}(s_{0}+10) \right)} \|\rho_{1} - \rho_{2}\|_{H^{s_{0}+5}}.$$
(8.106)

It remains to estimate $||K_{1,2}||_{L^2(\mathbb{T}^{d+2})}$. Then using (8.61) combined with Sobolev embeddings

 $\|\|[\mathbb{A}^1 - \mathbb{A}^2, \mathcal{R}_0]\|\|_{\overline{\alpha} - 1, s_0, 0} \lesssim \|\rho_1 - \rho_2\|_{H^{s_0} + 3} \overline{F}_0(s_0 + 3)).$

By virtue of Lemma 7.2-(iii) and Sobolev embeddings we deduce that

$$\|K_{1,2}\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|\rho_1 - \rho_2\|_{H^{s_0+3}} \overline{F}_0(s_0+3)).$$
(8.107)

Inserting (8.107) and (8.106) into (8.97) implies for any $t \in [0, 1]$

$$\|\mathscr{K}_t\|_{L^2(\mathbb{T}^{d+2})} \leqslant C e^{C\left(\|\rho_2\|_{H^{s_0+10}} + \|\rho_1\|_{H^{s_0+10}} + \overline{F}_0(s_0+10)\right)} \|\rho_1 - \rho_2\|_{H^{s_0+5}}.$$
(8.108)

This achieves the proof for q = 0. However for the genral case we may proceed as in (8.80). Since the analysis is quite similar we shall only restrict the discussion to the main lines. First we start from (8.95) and differentiate it successively with respect to λ giving rise to an analogous system to (8.83). Actually, we obtain

$$L_q X_q(t) = \overline{M}_q X_q + Y_q, \quad L_q \triangleq \mathscr{L} I_{q+1}.$$
(8.109)

where

$$X_q = \begin{pmatrix} \mathcal{H}_t \\ \gamma \partial_\lambda \mathcal{H}_t \\ \vdots \\ \gamma^q \partial_\lambda^q \mathcal{H}_t \end{pmatrix}, \quad Y_q = \begin{pmatrix} K_0 \\ \gamma \partial_\lambda K_0 \\ \vdots \\ \gamma^q \partial_\lambda^q K_0 \end{pmatrix} \quad \text{and} \quad \overline{M}_q = \begin{pmatrix} 0 & 0 & \cdots & \cdots & \cdots & 0 \\ \overline{m}_{1,1} & 0 & \cdots & \cdots & \cdots & 0 \\ \overline{m}_{2,2} & \overline{m}_{1,2} & 0 & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \cdots & \cdots & \overline{m}_{1,q-1} & 0 & 0 \\ \overline{m}_{q,q} & \overline{m}_{q-1,q} & \cdots & \cdots & \overline{m}_{1,q} & 0 \end{pmatrix}.$$

with $K_0 = (\widehat{\mathscr{L}}_2 - \widehat{\mathscr{L}}_1)\mathscr{K}_t^2 + K_{1,2}$ and the entries \overline{m}_{ij} are defined in similar way to (8.81). Then we proceed in a straightforward way following the proof of the first point (i). This achieved the proof.

8.3.1. Tame estimates of the iterated commutators. In this section we shall discuss an important result used during the proof of Theorem 8.1 related to some tame estimates governing iterated commutators of pseudo-differential operators of special structure. First we shall recall some operators seen in the preceding sections. The operator $\hat{\mathcal{L}}_n$ is described through (8.48) and (8.35) and the matrix \mathcal{M}_n is defined in (8.46) and takes the form

$$\mathcal{M}_{n} = \begin{pmatrix} 0 & 0 & \dots & \dots & \dots & 0 \\ m_{1,1} & 0 & \dots & \dots & 0 & 0 \\ m_{2,2} & m_{1,2} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & m_{1,n-1} & 0 & 0 \\ m_{n,n} & m_{n-1,n} & \dots & \dots & m_{1,n} & 0 \end{pmatrix} \in \mathbf{\Delta}_{1}(n).$$

Using (8.45), (8.40) and (8.43) one finds that

$$m_{k,i} = \binom{i}{k} \left(\mu(\varphi, \theta, k) + \nu(\varphi, \eta, k) \right)$$
(8.110)

with

$$\begin{aligned} &\mu(\varphi,\theta,k) \triangleq \partial_{\theta} \left((\partial_{\theta}^{k} \rho) |\mathbf{D}_{\theta}|^{\alpha-1} + |\mathbf{D}_{\theta}|^{\alpha-1} (\partial_{\theta}^{k} \rho) \right) \\ &\nu(\varphi,\eta,k) \triangleq (\partial_{\eta}^{k} \rho) \partial_{\eta} |\mathbf{D}_{\eta}|^{\alpha-1} + |\mathbf{D}_{\eta}|^{\alpha-1} (\partial_{\eta}^{k} \rho) \partial_{\eta}. \end{aligned} \tag{8.111}$$

Define

$$\mathscr{C}_k = \operatorname{Ad}^k_{\mathscr{M}_n}(\mathscr{Q}_n). \tag{8.112}$$

Then we have the recursive relation

$$\mathscr{C}_{k+1} = [\mathscr{M}_n, \mathscr{C}_k],$$

with the initial relation

$$\begin{aligned} \mathscr{C}_0 &= \widehat{\mathscr{L}} \mathbf{I}_{n+1} \\ &= - \big(\mu(\varphi, \theta, 0) + \nu(\varphi, \eta, 0) \big) \mathbf{I}_{n+1} \end{aligned}$$

The last identity comes from the definition of $\widehat{\mathscr{L}}$ seen in (8.35). Direct computations show that

According to Lemma 8.1 one has

$$\mathscr{C}_{k} = \begin{pmatrix} 0 & 0 & \dots & \dots & 0 & 0 \\ c_{1,1}^{k} & 0 & \dots & \dots & 0 & 0 \\ c_{2,2}^{k} & c_{1,2}^{k} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & 0 & 0 \\ c_{n,n}^{k} & c_{n-1,n}^{k} & \dots & \dots & c_{1,n}^{k} & 0 \end{pmatrix} \in \mathbf{\Delta}_{k}(n)$$

$$(8.113)$$

with

$$c_{i,j}^{k} = {j \choose i} \left(\mu_{k}(\varphi,\theta,i) + \nu_{k}(\varphi,\eta,i) \right), \ \forall i \leq j \in \{1,..,n\}$$

$$(8.114)$$

and the law $[\mathcal{M}_n, \mathcal{C}_k]$ is given by

$$\mu_{\mathcal{M}_n} * \mu_{\mathscr{C}_k} = \mu * \mu_k + \nu * \nu_k.$$

Thus we find the recursive relations

$$\mu_{k+1} = \mu \boxtimes \mu_k, \quad \mu_0(\varphi, \theta, i) = -\mu(\varphi, \theta, 0)$$
$$\nu_{k+1} = \nu \boxtimes \nu_k, \quad \nu_0(\varphi, \eta, i) = -\nu(\varphi, \eta, 0).$$

The estimates of μ_{k+1} and ν_{k+1} are similar and we shall restrict the discussion only to the first one. By definition of the convolution law one may write

$$\mu_{k+1}(\varphi,\theta,i) = \sum_{\ell=0}^{i} {i \choose \ell} \left[\mu(\varphi,\theta,i-\ell), \mu_k(\varphi,\theta,\ell) \right].$$

Since $\mathcal{M}_n \in \Delta_1(n)$ and $\mathcal{C}_k \in \Delta_k(n)$ then

$$\forall i \in \{k+1, ..., n\}, \quad \mu_{k+1}(\varphi, \theta, i) = \sum_{\ell=k}^{i-1} {i \choose \ell} \left[\mu(\varphi, \theta, i-\ell), \mu_k(\varphi, \theta, \ell) \right], \quad (8.115)$$

$$\forall i \in \{0, ..., k\}, \quad \mu_{k+1}(\varphi, \theta, i) = 0.$$

Our first main result reads as follows.

Lemma 8.2. Let $\gamma \in \mathbb{N}, s \ge s_0 > \frac{d+1}{2}$, then for any $n \in \mathbb{N}^*, k \in \{1, ..., n\}, i \in \{k, ..., n\}$

$$\|\|\mu_k(i)\|\|_{0,s,\gamma} + \|\|\nu_k(i)\|\|_{0,s,\gamma} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+3}}^k.$$

In addition, we have for any $s_0 > d + 1$

$$\|\mu_k(i)h\|_{L^2(\mathbb{T}^{d+2})} + \|\nu_k(i)h\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{H^{s_0+3+i}(\mathbb{T}^{d+1})} \|\rho\|_{H^{s_0+3}(\mathbb{T}^{d+1})}^k \|h\|_{L^2(\mathbb{T}^{d+2})}.$$

Proof. To alleviate the notation we denote $\mu_k(\varphi, \theta, i) = \mu_k(i)$. The goal is to prove the estimate using the induction principle in k. Let us first check the estimate for k = 1. In this case,

$$\mu_1(\varphi,\theta,i) = \sum_{\ell=k}^{i-1} {i \choose \ell} \left[\mu(\varphi,\theta,i-\ell), \mu_0(\varphi,\theta,\ell) \right]$$
$$= -\sum_{\ell=0}^{i-1} {i \choose \ell} \left[\mu(\varphi,\theta,i-\ell), \mu(\varphi,\theta,0) \right].$$

Applying Lemma 7.5-(iii) with $\mathscr{A} = \mu(i-\ell), \mathscr{B} = \mu(0), q = 0$ and

$$m_1 = m_2 = \alpha \in \left(0, \frac{1}{2}\right), \quad \mu = \frac{3}{2}, \mu_\beta = \overline{\mu}_\beta = \frac{3}{2} + \beta$$

yields

$$\begin{split} \|\|[\mu(i-\ell),\mu(0)]\|\|_{0,s,\gamma} &\leq \sum_{0\leqslant\beta\leqslant\gamma} \|\|\mathscr{A}\|\|_{\alpha,s,1+\gamma-\beta} \|\|\mathscr{B}\|\|_{\alpha,s_0+\mu,\beta} + \|\|\mathscr{A}\|\|_{\alpha,s_0,1+\gamma-\beta} \|\|\mathscr{B}\|\|_{\alpha,s+\mu,\beta} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\|\mathscr{A}\|\|_{\alpha,s,0} \|\|\mathscr{B}\|\|_{\alpha,s_0+\mu_{\beta},\gamma-\beta} + \|\|\mathscr{A}\|\|_{\alpha,s_0,0} \|\|\mathscr{B}\|\|_{\alpha,s+\mu_{\beta},\gamma-\beta} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\|\mathscr{A}\|\|_{\alpha,s_0+\mu,\beta} \|\|\mathscr{B}\|\|_{\alpha,s,1+\gamma-\beta} + \|\|\mathscr{A}\|\|_{\alpha,s+\mu,\beta} \|\|\mathscr{B}\|\|_{\alpha,s_0,1+\gamma-\beta} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\|\mathscr{A}\|\|_{\alpha,s_0+\mu_{\beta},\gamma-\beta} \|\|\mathscr{B}\|\|_{\alpha,s,0} + \|\|\mathscr{A}\|\|_{\alpha,s+\mu_{\beta},\gamma-\beta} \||\mathscr{B}\|\|_{\alpha,s_0,0} \,. \end{split}$$

Using (8.111) and (8.60), we obtain

$$\||\mu(\ell)||_{\alpha,s,\beta} \lesssim \|\rho\|_{H^{s+\beta+\ell+1}}.$$
(8.116)

Therefore, combining the preceding two estimates yields

$$\begin{aligned} \|\|[\mu(i-\ell),\mu(0)]\|\|_{0,s,\gamma} &\leq \sum_{0\leqslant\beta\leqslant\gamma} \|\rho\|_{H^{s+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s_0+\frac{5}{2}+\beta}} + \|\rho\|_{H^{s_0+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s+\frac{5}{2}+\beta}} \\ &+ \|\rho\|_{H^{s+1+i-\ell}} \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma}} + \|\rho\|_{H^{s_0+1+i-\ell}} \|\rho\|_{H^{s+\frac{5}{2}+\gamma}} \\ &+ \sum_{0\leqslant\beta\leqslant\gamma} \|\rho\|_{H^{s_0+\frac{5}{2}+\beta+i-\ell}} \|\rho\|_{H^{s+2+\gamma-\beta}} + \|\rho\|_{H^{s+\frac{5}{2}+\beta+i-\ell}} \|\rho\|_{H^{s_0+2+\gamma-\beta}} \\ &+ \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma+i-\ell}} \|\rho\|_{H^{s+1}} + \|\rho\|_{H^{s+\frac{5}{2}+\gamma+i-\ell}} \|\rho\|_{H^{s_0+1}}. \end{aligned}$$
(8.117)

Using Sobolev embeddings and interpolation inequality we get for $s \geqslant s_0$

$$\begin{aligned} \|\rho\|_{H^{s+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s_0+\frac{5}{2}+\beta}} &\lesssim \|\rho\|_{H^{s+2+\gamma-\beta+i}} \|\rho\|_{H^{s_0+\frac{5}{2}+\beta}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{3}{2}}} \end{aligned}$$

and

$$\begin{aligned} \|\rho\|_{H^{s_0+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s+\frac{5}{2}+\beta}} &\lesssim \|\rho\|_{H^{s_0+2+\gamma-\beta+i}} \|\rho\|_{H^{s+\frac{5}{2}+\beta}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{3}{2}}}. \end{aligned}$$

Similarly we get

$$\begin{aligned} \|\rho\|_{H^{s+1+i-\ell}} \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma}} &\lesssim \|\rho\|_{H^{s+1+i}} \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{1}{2}}} \end{aligned}$$

and

$$\begin{aligned} \|\rho\|_{H^{s_0+1+i-\ell}} \|\rho\|_{H^{s+\frac{5}{2}+\gamma}} &\lesssim \|\rho\|_{H^{s_0+1+i}} \|\rho\|_{H^{s+\frac{5}{2}+\gamma}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{1}{2}}} \end{aligned}$$

On the other hand

$$\begin{aligned} \|\rho\|_{H^{s_0+\frac{5}{2}+\beta+i-\ell}} \|\rho\|_{H^{s+2+\gamma-\beta}} &\lesssim \|\rho\|_{H^{s_0+\frac{5}{2}+\beta+i}} \|\rho\|_{H^{s+2+\gamma-\beta}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{3}{2}}} \end{aligned}$$

and

$$\begin{aligned} \|\rho\|_{H^{s+\frac{5}{2}+\beta+i-\ell}} \|\rho\|_{H^{s_0+2+\gamma-\beta}} &\lesssim \|\rho\|_{H^{s+\frac{5}{2}+\beta+i}} \|\rho\|_{H^{s_0+2+\gamma-\beta}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{3}{2}}}. \end{aligned}$$

As to the last terms in (8.117) we implement similar arguments as before yielding

$$\begin{aligned} \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma+i-\ell}} \|\rho\|_{H^{s+1}} &\lesssim \|\rho\|_{H^{s_0+\frac{5}{2}+\gamma+i}} \|\rho\|_{H^{s+1}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{1}{2}}} \end{aligned}$$

and

$$\begin{aligned} \|\rho\|_{H^{s+\frac{5}{2}+\gamma+i-\ell}} \|\rho\|_{H^{s_0+1}} &\lesssim \|\rho\|_{H^{s+\frac{5}{2}+\gamma+i}} \|\rho\|_{H^{s_0+1}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{1}{2}}}. \end{aligned}$$

Putting together the preceding estimates and using Sobolev embeddings yield

$$\begin{aligned} \| [\mu(i-\ell),\mu(0)] \| \|_{0,s,\gamma} &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+\frac{3}{2}}} \\ &\lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+3}}. \end{aligned}$$

This gives the result of Lemma 8.2 for k = 1. Now let us assume that this result is true at the order k and prove it at the order k + 1. According to the relation (8.115) on gets

$$\|\|\mu_{k+1}(i)\|\|_{0,s,\gamma} \lesssim \sum_{\ell=k}^{i-1} \|\|[\mu(i-\ell),\mu_k(\ell)]\|\|_{0,s,\gamma}.$$
(8.118)

Applying Lemma 7.5-(iii) with $m_1 = \alpha \in (0, \frac{1}{2}), m_2 = 0, q = 0$ and fixing the choice

$$\mu = 1, \quad \mu_{\beta} = \overline{\mu}_{\beta} = 1 + \beta$$

we find

$$\| [\mu(i-\ell), \mu_{k}(\ell)] \|_{0,s,\gamma} \lesssim \sum_{0 \le \beta \le \gamma} \| \|\mu(i-\ell)\|_{\alpha,s,1+\gamma-\beta} \| \|\mu_{k}(\ell)\|_{0,s_{0}+1,\beta}$$

$$+ \sum_{0 \le \beta \le \gamma} \| \|\mu(i-\ell)\|_{\alpha,s_{0},1+\gamma-\beta} \| \|\mu_{k}(\ell)\|_{0,s+1,\beta} + \| \|\mu(i-\ell)\|_{\alpha,s,0} \| \|\mu_{k}(\ell)\|_{0,s_{0}+1+\beta,\gamma-\beta}$$

$$+ \sum_{0 \le \beta \le \gamma} \| \|\mu(i-\ell)\|_{\alpha,s_{0},0} \| \|\mu_{k}(\ell)\|_{0,s_{0},1+\gamma-\beta} + \| \|\mu(i-\ell)\|_{\alpha,s_{0}+1,\beta} \| \|\mu_{k}(\ell)\|_{0,s,1+\gamma-\beta}$$

$$+ \sum_{0 \le \beta \le \gamma} \| \|\mu(i-\ell)\|_{\alpha,s+1,\beta} \| \|\mu_{k}(\ell)\|_{0,s_{0},1+\gamma-\beta} + \| \|\mu(i-\ell)\|_{\alpha,s_{0}+1+\beta,\gamma-\beta} \| \|\mu_{k}(\ell)\|_{0,s,0}$$

$$+ \sum_{0 \le \beta \le \gamma} \| \|\mu(i-\ell)\|_{\alpha,s+1+\beta,\gamma-\beta} \| \|\mu_{k}(\ell)\|_{0,s_{0},0} \triangleq \sum_{0 \le \beta \le \gamma} \sum_{j=1}^{8} \mathscr{I}_{j}^{\beta}.$$

$$(8.119)$$

Now let us estimate the first term \mathscr{I}_1^{β} . One has from the induction assumption and (8.116),

$$\begin{aligned} \mathscr{G}_{1}^{\beta} &\lesssim \|\rho\|_{H^{s+2+\gamma-\beta+i-\ell}} \|\|\mu_{k}(\ell)\|_{0,s_{0}+1,\beta} \\ &\lesssim \|\rho\|_{H^{s+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s_{0}+4+\beta+\ell}} \|\rho\|_{H^{s_{0}+3}}^{k} \end{aligned}$$

Using interpolation inequality we find, since $\ell \leq i - 1$, that for any $s \geq s_0$

$$\|\rho\|_{H^{s+2+\gamma-\beta+i-\ell}}\|\rho\|_{H^{s_0+4+\beta+\ell}} \lesssim \|\rho\|_{H^{s_0+3}}\|\rho\|_{H^{s+3+\gamma+i}}.$$

Consequently,

$$\mathcal{I}_{1}^{\beta} \lesssim \|\rho\|_{H^{s+3+\gamma+i}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.120)

Concerning the second term \mathscr{I}_2^β we write

$$\begin{aligned} \mathscr{I}_{2}^{\beta} \lesssim \|\rho\|_{H^{s_{0}+2+\gamma-\beta+i-\ell}} \|\|\mu_{k}(\ell)\|_{0,s+1,\beta} \\ \lesssim \|\rho\|_{H^{s_{0}+2+\gamma-\beta+i-\ell}} \|\rho\|_{H^{s+4+\ell+\beta}} \|\rho\|_{H^{s_{0}+3}}^{k} \end{aligned}$$

Interpolation inequalities allow to get since $\ell \leqslant i-1$

 $\|\rho\|_{s_0+2+\gamma-\beta+i-\ell}\|\rho\|_{H^{s+4+\ell+\beta}} \lesssim \|\rho\|_{s_0+3}\|\rho\|_{H^{s+3+i+\gamma}}$

yields

$$\mathscr{I}_{2}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.121)

For the third term, we write

$$\begin{aligned} \mathscr{I}_{3}^{\beta} &= ||| \ \mu(i-\ell) \ |||_{\alpha,s,0} \ ||| \mu_{k}(\ell) |||_{0,s_{0}+1+\beta,\gamma-\beta} \\ &\lesssim ||\rho||_{H^{s+1+i-\ell}} ||\rho||_{H^{s_{0}+4+\gamma+\ell}} ||\rho||_{H^{s_{0}+3}}. \end{aligned}$$

Using the interpolation inequality, since $\ell \leq i - 1$, we find that for any $s \geq s_0$

$$\|\rho\|_{H^{s+1+i-\ell}} \|\rho\|_{H^{s_0+4+\gamma+\ell}} \lesssim \|\rho\|_{H^{s_0+2}} \|\rho\|_{H^{s+3+\gamma+i}}.$$

Hence, Sobolev embeddings imply

$$\mathscr{I}_{3}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.122)

Let us move the term $\mathscr{I}_4^\beta.$ Then one has

As before, since $\ell \leqslant i-1$ then using the following interpolation inequality

$$\|\rho\|_{H^{s_0+1+i-\ell}}\|\rho\|_{H^{s+4+\ell+\gamma}} \lesssim \|\rho\|_{H^{s_0+2}}\|\rho\|_{H^{s+3+i+\gamma}}.$$

Therefore we obtain

$$\mathcal{I}_{4}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.123)

As to the term $\mathscr{I}_5^\beta,$ one has

$$\begin{aligned} \mathcal{I}_{5}^{\beta} &= \|\| \mu(i-\ell) \|\|_{\alpha,s_{0}+1,\beta} \|\| \mu_{k}(\ell) \|\|_{0,s,1+\gamma-\beta} \\ &\lesssim \|\rho\|_{H^{s_{0}+2+i-\ell+\beta}} \|\rho\|_{H^{s+4+\ell+\gamma-\beta}} \|\rho\|_{H^{s_{0}+3}}^{k}. \end{aligned}$$

Now we use the interpolation inequality, since $\ell\leqslant i-1,$

$$\|\rho\|_{H^{s_0+2+i-\ell+\beta}}\|\rho\|_{H^{s+4+\ell+\gamma-\beta}} \lesssim \|\rho\|_{H^{s_0+3}}\|\rho\|_{H^{s+3+i+\gamma}}.$$

Hence

$$\mathscr{I}_{5}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k}.$$
(8.124)

For the term $\mathscr{I}_{6}^{\beta},$ one has

$$\mathcal{I}_{6}^{\beta} = \||\mu(i-\ell)|||_{\alpha,s+1,\beta} \||\mu_{k}(\ell)|||_{0,s_{0},1+\gamma-\beta}$$
$$\lesssim \|\rho\|_{H^{s+2+i-\ell+\beta}} \|\rho\|_{H^{s_{0}+4+\ell+\gamma-\beta}} \|\rho\|_{H^{s_{0}+3}}^{k}.$$

From interpolation inequality we infer

$$\|\rho\|_{H^{s+2+i-\ell+\beta}}\|\rho\|_{H^{s_0+4+\ell+\gamma-\beta}} \lesssim \|\rho\|_{H^{s+3+i+\gamma}}\|\rho\|_{H^{s_0+3}}$$

and therefore

$$\mathcal{I}_{6}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.125)

Concerning the term $\mathscr{I}_7^\beta,$ one has by interpolation

$$\begin{aligned} \mathscr{G}_{7}^{\beta} &= \||\mu(i-\ell)|||_{\alpha,s_{0}+1+\beta,\gamma-\beta}|||\mu_{k}(\ell)|||_{0,s,0} \\ &\lesssim \|\rho\|_{H^{s_{0}+2+\gamma+i-\ell}} \|\rho\|_{H^{s+3+\ell}} \|\rho\|_{H^{s_{0}+3}}^{k} \\ &\lesssim \|\rho\|_{H^{s_{0}+2}} \|\rho\|_{H^{s+3+\gamma+i}} \|\rho\|_{H^{s_{0}+3}}^{k}. \end{aligned}$$

Thus

$$\mathscr{I}_{7}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.126)

For the last term, we write

$$\mathcal{I}_{8}^{\beta} = |||\mu(i-\ell)|||_{\alpha,s+1+\beta,\gamma-\beta}|||\mu_{k}(\ell)|||_{0,s_{0},0}$$

$$\lesssim ||\rho||_{H^{s+2+\gamma+i-\ell}} ||\rho||_{H^{s_{0}+3+\ell}} ||\rho||_{H^{s_{0}+3}}^{k}.$$

By interpolation we get

$$\|\rho\|_{H^{s+2+\gamma+i-\ell}}\|\rho\|_{H^{s_0+3+\ell}} \lesssim \|\rho\|_{H^{s+3+i+\gamma}}\|\rho\|_{H^{s_0+2}}.$$

It follows that

$$\mathscr{I}_{8}^{\beta} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_{0}+3}}^{k+1}.$$
(8.127)

Inserting the estimate (8.120)-(8.121)-..-(8.127) into (8.119) we deduce for any $0 \leq \ell \leq i-1$

$$\| [\mu(i-\ell), \mu_k(\ell)] \|_{0,s,\gamma} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+3}}^{k+1}.$$

Plugging this into (8.118) gives

$$\|\|\mu_{k+1}(i)\|\|_{0,s,\gamma} \lesssim \|\rho\|_{H^{s+3+i+\gamma}} \|\rho\|_{H^{s_0+3}}^{k+1},$$

which achieves the induction argument. To prove the second part of Lemma 8.2 it suffices to combine the first part with Lemma 7.3-(i).

Now we shall establish the following result based on Lemma 8.2 and used before in the proof of Theorem 8.1. Notice that we shall use below the same notations introduced in (8.46), (8.47), (8.48), (8.84) and (8.85).

Lemma 8.3. Let $0 \leq j < k \leq n$ and $s_0 > \frac{d}{2} + 1$. Then we have the following assertions.

(i) For any $i \in \{2, ..., n+1\}$

$$\left\| \left(\operatorname{Ad}_{\mathcal{M}_n}^{k-j}(\mathcal{L}_n) \,\mathcal{M}_n^j \mathcal{X}_n \right)(i) \right\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{H^{s_0+3}(\mathbb{T}^{d+1})}^{k-j} \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_0+3+i-\ell}(\mathbb{T}^{d+1})} \left\| \left(\mathcal{M}_n^j \mathcal{X}_n \right)(\ell) \right\|_{L^2(\mathbb{T}^{d+2})}.$$

The notation Y(i) means the the *i*-th component of the vector Y. Notice that for i = 1 the first component of the left hand side member is vanishing.

(ii) For any $i \in \{1, ..., n+1\}$

$$\left\| \left(\operatorname{Ad}_{\overline{\mathcal{M}}_{n}}^{k-j}(\overline{\mathcal{Z}}_{n}) \overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \right)^{k-j} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \right\|_{L^{2}(\mathbb{T}^{d+2})} \leq \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \right\|_{L^{2}(\mathbb{T}^{d+2})} \leq \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \right\|_{L^{2}(\mathbb{T}^{d+2})} \leq \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \leq \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s_{0}+3+i-\ell}^{q,\kappa} \left\| \left(\overline{\mathcal{M}}_{n}^{j} \overline{\mathcal{X}}_{n} \right)(\ell) \right\|_{L^{2}(\mathbb{T}^{d+2})} \right)^{k-j}$$

Here, the notation Y(i) means the the *i*-th bloc vector of size q + 1.

Proof. (i) Using (8.112) and (8.113), we can write

$$\mathscr{C}_{k-j} \triangleq \operatorname{Ad}_{\mathscr{M}_n}^{k-j}(\mathscr{L}_n) = (a_{i,\ell}^{k-j})_{1 \leq i,\ell \leq n+1} \in \Delta_{k-j}(n),$$

with

$$\begin{aligned} a_{i,\ell}^{k-j} &= c_{i-\ell,i-1}^{k-j}, \quad \text{if} \quad i \geqslant \ell + k - j \\ a_{i,\ell}^{k-j} &= 0, \quad \text{otherwise.} \end{aligned}$$

Denote $\mathscr{Z}_j = \mathscr{M}_n^j \mathscr{X}_n$, then we obtain for $0 \leq j < k$

$$\left(\mathscr{C}_{k-j} \mathscr{M}_n^j \mathscr{X}_n \right) (i) = \sum_{\ell=1}^{i+j-k} a_{i,\ell}^{k-j} \mathscr{Z}_j(\ell)$$
$$= \sum_{\ell=1}^{i-1} a_{i,\ell}^{k-j} \mathscr{Z}_j(\ell).$$

According to (8.114) and Lemma 8.2 we deduce

$$\|a_{i,\ell}^{k-j}\mathscr{Z}_j(\ell)\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{H^{s_0+3+i-\ell}(\mathbb{T}^{d+1})} \|\rho\|_{H^{s_0+3}(\mathbb{T}^{d+1})}^{k-j} \|\mathscr{Z}_j(\ell)\|_{L^2(\mathbb{T}^{d+2})}.$$

Therefore we get

$$\left\| \left(\mathscr{C}_{k-j} \mathscr{M}_{n}^{j} \mathscr{X}_{n} \right) (i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{H^{s_{0}+3}(\mathbb{T}^{d+1})}^{k-j} \sum_{\ell=1}^{i-1} \|\rho\|_{H^{s_{0}+3+i-\ell}(\mathbb{T}^{d+1})} \|\mathscr{Z}_{j}(\ell)\|_{L^{2}(\mathbb{T}^{d+2})},$$

which is the desired result.

(ii) Using (8.85) and Lemma 8.1 we can write

$$\overline{\mathscr{C}}_{k-j} \triangleq \operatorname{Ad}_{\overline{\mathscr{M}}_n}^{k-j}(\overline{\mathscr{L}}_n) = (\overline{a}_{i,\ell}^{k-j})_{1 \leqslant i, \ell \leqslant n+1} \in \Delta_0(n,q)$$

with $\overline{a}_{i,\ell}^{k-j}$ a Töplitz matrix belonging to $\Delta_0(q)$. Denote $\overline{\mathscr{X}}_j = \overline{\mathscr{M}}_n^j \overline{\mathscr{X}}_n$. Hence we obtain for $0 \leq j < k$

$$\left(\overline{\mathscr{C}}_{k-j}\overline{\mathscr{M}}_{n}^{j}\overline{\mathscr{X}}_{n}\right)(i) = \sum_{\ell=1}^{i} \overline{a}_{i,\ell}^{k-j}\overline{\mathscr{Z}}_{j}(\ell).$$

The main goal is to prove the following estimate,

$$\|\overline{a}_{i,\ell}^{k-j}\overline{\mathscr{X}}_j(\ell)\|_{L^2(\mathbb{T}^{d+2})} \lesssim \|\rho\|_{s_0+3+i-\ell}^{q,\kappa} \left(\|\rho\|_{s_0+3}^{q,\kappa}\right)^{k-j} \|\overline{\mathscr{X}}_j(\ell)\|_{L^2(\mathbb{T}^{d+2})}.$$
(8.128)

Assume for a while this estimate, then

$$\left\| \left(\overline{\mathcal{C}}_{k-j}\overline{\mathcal{M}}_{n}^{j}\overline{\mathcal{X}}_{n}\right)(i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \lesssim \left(\|\rho\|_{s_{0}+3}^{q,\kappa} \right)^{k-j} \sum_{\ell=1}^{i} \|\rho\|_{s+3+i-\ell}^{q,\kappa} \|\overline{\mathcal{Z}}_{j}(\ell)\|_{L^{2}(\mathbb{T}^{d+2})}.$$

Remind that we have the recursive relation

$$\overline{\mathscr{C}}_{k+1} = \left[\overline{\mathscr{M}}_n, \overline{\mathscr{C}}_k\right],$$

with the initial condition

$$\overline{\mathscr{C}}_0 = L_q \mathbb{I}_{n+1} = -\left(\mu(\varphi, \theta, 0) + \nu(\varphi, \eta, 0)\right) \mathbb{I}_{(n+1)(q+1)}.$$

From (8.85) and (8.86) one has that the law of $\overline{\mathcal{M}}_n$ is given by

$$\mu_{\overline{\mathcal{M}}_n}(i) = \begin{cases} \operatorname{Ad}_{\partial_{\chi}}^i \overline{M}_q - \operatorname{Ad}_{\partial_{\chi}}^i \overline{L}_q, & 1 \leq i \leq n \\ \overline{M}_q, & i = 0 \end{cases}$$
(8.129)

and one can check from straightforward computations based on (8.81) and (8.82) that we have the splitting

$$\mu_{\overline{\mathcal{M}}_n}(i) \triangleq \overline{\mu}(i) + \overline{\nu}(i)$$

where for $1 \leq i \leq n$

$$\overline{\mu}(i) = \begin{pmatrix} \overline{\mu}_{0,0}(i) & 0 & 0 & \dots & \dots & 0 \\ \overline{\mu}_{1,1}(i) & \overline{\mu}_{0,1}(i) & 0 & \dots & \dots & 0 \\ \overline{\mu}_{2,2}(i) & \overline{\mu}_{1,2}(i) & \overline{\mu}_{0,2}(i) & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \overline{\mu}_{1,q-1}(i) & \overline{\mu}_{0,q-1}(i) & 0 \\ \overline{\mu}_{q,q}(i) & \overline{\mu}_{q-1,q}(i) & \dots & \dots & \overline{\mu}_{1,q}(i) & \overline{\mu}_{0,q}(i) \end{pmatrix},$$

$$(8.130)$$

such that for $1 \leq k \leq q$

$$\overline{\mu}_{k,j}(i) = {j \choose k} \operatorname{Ad}_{\partial_{\chi}}^{i} \overline{\mathbb{B}}_{\theta,(k)}$$
$$= {j \choose k} \kappa^{k} [\partial_{\lambda}^{k}, \mathbb{A}_{\theta,(i)}]$$

and for k = 0, using (8.35),

$$\overline{\mu}_{0,j}(i) = -\operatorname{Ad}_{\partial_{\chi}}^{i} \mathscr{L}$$
$$= \mathbb{A}_{\theta,(i)}.$$

In addition for i = 0 we have

$$\overline{\mu}(0) = \begin{pmatrix} 0 & 0 & 0 & \dots & \dots & 0 \\ \overline{\mu}_{1,1}(0) & 0 & 0 & \dots & \dots & 0 \\ \overline{\mu}_{2,2}(0) & \overline{\mu}_{1,2}(0) & 0 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \overline{\mu}_{1,q-1}(0) & 0 & 0 \\ \dots & \dots & \dots & \overline{\mu}_{1,q}(0) & 0 \end{pmatrix}, \quad \overline{\mu}_{k,j}(0) = {j \choose k} \kappa^k [\partial_{\lambda}^k, \mathbb{A}_{\theta,(0)}]. \quad (8.131)$$

In a similar way we get

$$\overline{\nu}(i) = \begin{pmatrix} \overline{\nu}_{0,0}(i) & 0 & \dots & \dots & \dots & 0 \\ \overline{\nu}_{1,1}(i) & \overline{\nu}_{0,1}(i) & 0 & \dots & \dots & 0 \\ \overline{\nu}_{2,2}(i) & \overline{\nu}_{1,2} & \overline{\nu}_{1,2}(i) & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ \dots & \dots & \dots & \overline{\nu}_{1,q-1}(i) & \overline{\nu}_{0,q-1}(i) & 0 \\ \overline{\mu}_{q,q}(i) & \overline{\nu}_{q-1,q}(i) & \dots & \dots & \overline{\nu}_{1,q}(i) & \overline{\nu}_{0,q}(i) \end{pmatrix},$$
(8.132)

where for $1 \leq i \leq n$

$$\overline{\nu}_{k,j}(i) = \begin{cases} \binom{j}{k} \operatorname{Ad}_{\partial_{\chi}}^{i} (\overline{\mathbb{B}}_{\eta,(k)} - \overline{\mathbb{T}}_{k}), & 1 \leq k \leq q\\ \overline{\mathbb{B}}_{\eta,(0)} - \overline{\mathbb{T}}_{0}, & k = 0 \end{cases}$$

and for i = 0 we make the adaptation as before. By applying Lemma 8.1 one finds

$$\overline{\mathscr{C}}_{k} = \begin{pmatrix} \overline{C}_{0,0}^{k} & 0 & \dots & \dots & \dots & 0 \\ \overline{C}_{1,1}^{k} & \overline{C}_{0,1}^{k} & 0 & \dots & \dots & 0 \\ \overline{C}_{2,2}^{k} & \overline{C}_{1,2} & \overline{C}_{0,2}^{k} & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & 0 \\ \dots & \dots & \dots & \overline{C}_{1,n-1}^{k} & \overline{C}_{0,n-1}^{k} & 0 \\ \overline{C}_{n,n}^{k} & \overline{C}_{n-1,n}^{k} & \dots & \dots & \overline{C}_{1,n}^{k} & \overline{C}_{0,n}^{k} \end{pmatrix}.$$

$$(8.133)$$

with

$$\overline{C}_{i,j}^{k} = {j \choose i} \left(\overline{\mu}_{k}(\varphi, \theta, i) + \overline{\nu}_{k}(\varphi, \eta, i) \right), \ \forall i \leq j \in \{1, .., n\}$$

$$(8.134)$$

and the law $\left[\overline{\mathcal{M}}_n,\overline{\mathcal{C}}_k\right]$ is given by

$$(\overline{\mu} + \overline{\nu}) \circledast (\overline{\mu}_k + \overline{\nu}_k) = \overline{\mu} \circledast \overline{\mu}_k + \overline{\nu} \circledast \overline{\nu}_k$$
(8.135)

where in the last identity we use the commutation relations

$$(\overline{\mu} \circledast \overline{\mu}_k)(i) = 0 = (\overline{\nu} \circledast \overline{\nu}_k)(i).$$
(8.136)

Indeed, one has

$$(\overline{\mu} \circledast \overline{\mu}_k)(i) = \sum_{\ell=0}^{i} {\binom{i}{\ell}} \left[\overline{\mu}(i-\ell), \overline{\mu}_k(\ell)\right].$$

Since $\overline{\mu}(i-\ell), \overline{\mu}_k(\ell) \in \Delta_0(q)$ and the entries of the matrix operators $\overline{\mu}(i-\ell)$ and $\overline{\mu}_k(\ell)$ commute mutually then from Remark 8.1 we deduce that $[\overline{\mu}(i-\ell), \overline{\nu}_k(\ell)] = 0$. This proves (8.136). Thus we obtain from (8.135) the recursive relations

$$\overline{\mu}_{k+1} = \overline{\mu} \circledast \overline{\mu}_k, \quad \overline{\mu}_0(i) = -\mu(\varphi, \theta, 0) \mathbf{I}_{q+1}$$
$$\overline{\nu}_{k+1} = \overline{\nu} \circledast \overline{\nu}_k, \quad \overline{\nu}_0(i) = -\nu(\varphi, \theta, 0) \mathbf{I}_{q+1}.$$

The estimates of $\overline{\mu}_{k+1}$ and $\overline{\nu}_{k+1}$ are similar and we shall restrict the discussion only to the first one. Hence we get for $i \in \{0, ..., n\}$

$$\overline{\mu}_{k+1}(i) = \left[\overline{\mu}(0), \overline{\mu}_k(i)\right] + \sum_{\substack{\ell=0\\107}}^{i-1} \binom{i}{\ell} \left[\overline{\mu}(i-\ell), \overline{\mu}_k(\ell)\right]$$

$$= \operatorname{Ad}_{\overline{\mu}(0)} \overline{\mu}_{k}(i) + \sum_{\ell=0}^{i-1} {i \choose \ell} \left[\overline{\mu}(i-\ell), \overline{\mu}_{k}(\ell) \right].$$
(8.137)

Iterating this yields

$$\overline{\mu}_{k+q+1}(i) = \operatorname{Ad}_{\overline{\mu}(0)}^{q+1} \overline{\mu}_k(i) + \sum_{\ell=0}^{i-1} {i \choose \ell} \sum_{j=0}^q \operatorname{Ad}_{\overline{\mu}(0)}^{q-j} \left[\overline{\mu}(i-\ell), \overline{\mu}_{k+j}(\ell) \right]$$

On the other hand, from (8.131) one gets $\overline{\mu}(0) \in \Delta_1(q)$. Therefore applying Lemma 8.1 we deduce that

$$\operatorname{Ad}_{\overline{\mu}(0)}^{m} \overline{\mu}_{k}(i) \in \mathfrak{A}_{\min\{m,q+1\}}(q).$$

In particular we find that $\operatorname{Ad}_{\overline{\mu}(0)}^{q+1}\overline{\mu}_k(i) = 0$ and consequently

$$\overline{\mu}_{k+q+1}(i) = \sum_{\ell=0}^{i-1} {i \choose \ell} \sum_{j=0}^{q} \operatorname{Ad}_{\overline{\mu}(0)}^{q-j} \left[\overline{\mu}(i-\ell), \overline{\mu}_{k+j}(\ell) \right].$$
(8.138)

We point out that from (8.137), the matrix operator $\overline{\mu}_k(i)$ is a Topelitz matrix in $\Delta_0(q)$ and therefore the entries of the commutators matrices $[\overline{\mu}(i-\ell), \overline{\mu}_k(\ell)]$ are linear combination of scalar commutators and then we can use the estimates of Lemma 7.5. Thus straightforward computations with slight adaptations of the proof of Lemma 8.2 allow to check by induction, using in particular the identity (8.138), that for any $k \in \mathbb{N}, j \in \{0, 1, ..., q\}, \gamma \in \mathbb{N}, s \ge s_0$

$$\||\overline{\mu}_{k+j}(i)||_{0,s,\gamma} \lesssim \|\rho\|_{s_0+3+i+j+\gamma}^{q,\kappa} (\|\rho\|_{s_0+3}^{q,\kappa})^{k+j},$$

which gives the estimate (8.128) in view of Lemma 7.3-(i).

8.3.2. Iterated kernel estimates. This section is devoted to some estimates on the iterated kernels used before during the proof of Theorem 8.1. Let us first introduce the problem that we want to solve. Given a sequence of smooth functions $\rho_n : \mathcal{O} \times \mathbb{T}^{d+1} \to \mathbb{R}$ and define the pseudo-differential operator of order $\alpha \in (0, \frac{1}{2})$

$$\mathbb{A}_{\theta,n} = \partial_{\theta} \big(\rho_n(\varphi,\theta) |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} \rho_n(\varphi,\theta) \big).$$

Let $(\varphi, \theta, \eta) \mapsto K_0(\varphi, \theta, \eta)$ be a function with some prescribed regularity, essentially smooth in the variable φ but slightly singular in the diagonal $\theta = \eta$, and consider the iterative kernels

$$K_{n+1} = \left(\mathbb{A}_{\theta,n} + \mathbb{A}_{\eta,n} - \mathbb{S}_n\right) K_n, n \in \mathbb{N}$$

with

$$\mathbb{S}_n \triangleq (\partial_\eta \rho_n) |\mathcal{D}_\eta|^{\alpha - 1} + |\mathcal{D}_\eta|^{\alpha - 1} (\partial_\eta \rho_n).$$

We want to find suitable recursive estimates for the sequence $(K_n)_n$ that allow to deduce later tame estimates. At this stage, it is worthy to point out that, because the initial kernel K_0 may be singular at the diagonal $\theta = \eta$, we cannot deal separately with the operators $\mathbb{A}_{\theta,n}$ and $\mathbb{A}_{\eta,n}$ and one should keep together their collective behavior in order to cancel the diagonal singularity. This can be easily understood with the following simple toy model. Take $K_0(\varphi, \theta, \eta) = \Psi(\theta - \eta)$ and Ψ is singular and let $\mathbb{A}_{\theta} = \partial_{\theta}$. Then both new kernels $\mathbb{A}_{\theta}K_0$ and $\mathbb{A}_{\eta}K_0$ are more singular than K_0 and their iterated kernels $\mathbb{A}_{\theta}^n K_0$ and $\mathbb{A}_{\eta}^n K_0$ become more and more singular. However, it is clear that $(\mathbb{A}_{\theta} + \mathbb{A}_{\eta})K_0 = 0$ which means that the singularity is canceled due to the mutual actions of the involved operators. This example is very particular because we are dealing with differential operators which act point-wisely, which is not the case with our nonlocal pseudo-differential operators. Then to capture this cancellation, we find convenient to move to the symbol representation, find suitable recursive estimates and later come back to the kernel. According to the formula (7.8), the symbol associated to K_n and denoted here by a_n can be recovered through the formula

$$a_n(\varphi,\theta,\xi) = \int_{\mathbb{T}} K_n(\varphi,\theta,\theta+\eta) e^{i\eta\xi} d\eta, \qquad (8.139)$$

with the inverse Fourier formula

$$K_n(\varphi,\theta,\eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} a_n(\varphi,\theta,\xi) e^{i(\theta-\eta)\xi}.$$
(8.140)

Now we shall use the following splitting

$$K_{n+1} = (\mathscr{A}_{\theta,n} + \mathscr{A}_{\eta,n}) K_n + \mathscr{R}_n K_n, \quad n \in \mathbb{N}$$
(8.141)

with

$$\mathscr{A}_{\theta,n} = 2\rho_n(\varphi,\theta)\partial_\theta |\mathcal{D}_\theta|^{\alpha-1}$$

and

$$\begin{aligned} \mathscr{R}_{n} &= 2(\partial_{\theta}\rho_{n})|\mathbf{D}_{\theta}|^{\alpha-1} + (\partial_{\eta}\rho_{n})|\mathbf{D}_{\eta}|^{\alpha-1} \\ &+ \partial_{\theta}\left[|\mathbf{D}_{\theta}|^{\alpha-1},\rho_{n}\right] + \partial_{\eta}\left[|\mathbf{D}_{\eta}|^{\alpha-1},\rho_{n}\right] - |\mathbf{D}_{\eta}|^{\alpha-1}(\partial_{\eta}\rho_{n}) \\ &\triangleq \mathscr{R}_{n,1} + \mathscr{R}_{n,2} + \mathscr{R}_{n,3} + \mathscr{R}_{n,4} + \mathscr{R}_{n,5}. \end{aligned}$$
(8.142)

For the sake of simplicity we shall make in the statement below the following convention: if $\sigma_{\mathscr{A}}$ is the symbol of an operator \mathscr{A} as in (7.12) then we denote by $\||\sigma_{\mathscr{A}}|\|_{m,s,\gamma}^{q,\kappa}$ the same norm $\||\mathscr{A}|\|_{m,s,\gamma}^{q,\kappa}$ defined in (7.13). Our first main result reads as follows.

Lemma 8.4. Let $q \in \mathbb{N}$, $m \in \mathbb{R}$, $\gamma \in \mathbb{N}$ and $s \ge s_0 > \frac{d+1}{2} + q$, then the following assertions hold true.

(i) For any $n \in \mathbb{N}$ we have

$$\begin{split} \|\|a_{n+1}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s+1,\gamma}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+1,\gamma}^{q,\kappa} \\ &+ \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s,\gamma+1}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0,\gamma+1}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho_n\|_{s_0+3+|m|+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s,\beta}^{q,\kappa} + \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s+\gamma-\beta,\beta}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho_n\|_{s+3+|m|+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa} \,. \end{split}$$

(ii) Let $m \in \mathbb{R}$, $s \ge s_0, \gamma, n \in \mathbb{N}$ and assume the existence of an increasing log-convex function F such that for

$$\forall s' \ge s_0, \forall \gamma' \in \mathbb{N} \quad with \quad s' + \gamma' \leqslant s + \gamma + n \Longrightarrow |||a_0|||_{m,s',\gamma'}^{q,\kappa} \leqslant F(s' + \gamma'). \tag{8.143}$$

Then,

$$|||a_n|||_{m,s,\gamma}^{q,\kappa} \lesssim F(s_0+3) \sum_{i=0}^{n-1} ||\rho_i||_{s+n+4+|m|+\gamma}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{n-1} ||\rho_j||_{s_0+3+|m|}^{q,\kappa} + F(s+\gamma+n+4) \prod_{j=0}^{n-1} ||\rho_j||_{s_0+3+|m|}^{q,\kappa},$$

with the convention $\sum_{i=0}^{-1} \dots = 0$ and $\prod_{j=0}^{-1} \dots = 1$.

(iii) Let $m < -\frac{1}{2}$, $s \ge s_0$ and $n \in \mathbb{N}$. Assume that (8.143) occurs, then

$$\left(\int_{\mathbb{T}} \left(\|K_{n}(\cdot, ., .+\eta)\|_{s}^{q,\kappa}\right)^{2} d\eta\right)^{\frac{1}{2}} \lesssim F\left(s_{0}+3\right) \sum_{i=0}^{n-1} \|\rho_{i}\|_{s+n+4+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{n-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + F\left(s+n+4\right) \prod_{j=0}^{n-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}.$$

Proof. (i) First, by the integral representation (4.19) one gets by straightforward computations

$$\partial_{\theta} |\mathbf{D}_{\theta}|^{\alpha - 1} h(\varphi, \theta) = \text{p.v.} \int_{\mathbb{T}} \mathscr{K}_{0}(\theta') h(\varphi, \theta - \theta') d\theta', \quad \mathscr{K}_{0}(\theta) = \frac{-\alpha \operatorname{cotan}(\frac{\theta}{2})}{4\pi |\sin(\frac{\theta}{2})|^{\alpha}}.$$

Then using in particular (8.140) we find

$$\mathcal{A}_{\theta,n}K_n(\varphi,\theta,\eta) = 2\rho_n(\varphi,\theta) \int_{\mathbb{T}} \mathscr{K}_0(\theta')K_n(\varphi,\theta-\theta',\eta)d\theta'$$
$$= \frac{1}{\pi}\rho_n(\varphi,\theta) \sum_{\xi'\in\mathbb{Z}} \int_{\mathbb{T}} \mathscr{K}_0(\theta')a_n(\varphi,\theta-\theta',\xi')e^{\mathrm{i}\,(\theta-\theta'-\eta)\xi'}d\theta'.$$

Similarly, we get

$$\mathcal{A}_{\eta,n}K_n(\varphi,\theta,\eta) = \frac{1}{\pi}\rho_n(\varphi,\eta)\sum_{\xi'\in\mathbb{Z}}\int_{\mathbb{T}}\mathscr{K}_0(\theta')a_n(\varphi,\theta,\xi')e^{\mathrm{i}\,(\theta+\theta'-\eta)\xi'}d\theta'$$
$$= -\frac{1}{\pi}\rho_n(\varphi,\eta)\sum_{\xi'\in\mathbb{Z}}\int_{\mathbb{T}}a_n(\varphi,\theta,\xi')\mathscr{K}_0(\theta')e^{\mathrm{i}\,(\theta-\theta'-\eta)\xi'}d\theta', \tag{8.144}$$

where we have used in the last line the fact that \mathscr{K}_0 is odd. Therefore the symbol associated to $\mathscr{A}_{\theta,n}K_n$ and given by (8.139) takes the form

$$\frac{1}{\pi} \sum_{\xi' \in \mathbb{Z}} \rho_n(\varphi, \theta) \iint_{\mathbb{T}^2} \mathscr{K}_0(\theta') a_n(\varphi, \theta - \theta', \xi') e^{\mathrm{i}\,(\xi - \xi')\eta} e^{-\mathrm{i}\,\theta'\xi'} d\theta' d\eta$$

and the one of $\mathcal{A}_{\eta,n}K_n$ is given by

$$-\frac{1}{\pi}\sum_{\xi'\in\mathbb{Z}}a_n(\varphi,\theta,\xi')\iint_{\mathbb{T}^2}\mathscr{K}_0(\theta')\rho_n(\varphi,\theta+\eta)e^{\mathrm{i}\,(\xi-\xi')\eta}e^{-\mathrm{i}\,\theta'\xi'}d\theta'd\eta$$

Hence the symbol associated to $\mathscr{A}_{\theta,n}K_n+\mathscr{A}_{\eta,n}K_n$ takes the form

$$\frac{1}{\pi} \sum_{\xi' \in \mathbb{Z}} \rho_n(\varphi, \theta) \iint_{\mathbb{T}^2} \mathscr{K}_0(\theta') \left[a_n(\varphi, \theta - \theta', \xi') - a_n(\varphi, \theta, \xi') \right] e^{i(\xi - \xi')\eta} e^{-i\theta'\xi'} d\theta' d\eta
- \frac{1}{\pi} \sum_{\xi' \in \mathbb{Z}} \lambda(\xi') a_n(\varphi, \theta, \xi') \int_{\mathbb{T}} \left[\rho_n(\varphi, \theta + \eta) - \rho_n(\varphi, \theta) \right] e^{i(\xi - \xi')\eta} d\eta
\triangleq \sigma_{n,1}(\varphi, \theta, \xi) + \sigma_{n,2}(\varphi, \theta, \xi)$$

with

$$\lambda(\xi) \triangleq \int_{\mathbb{T}} \mathscr{K}_0(\theta') e^{-\mathrm{i}\,\theta'\xi} d\theta'.$$
(8.145)

One may easily write by integration in η

$$\sigma_{n,1}(\varphi,\theta,\xi) = 2\rho_n(\varphi,\theta) \int_{\mathbb{T}} \mathscr{K}_0(\theta') \big[a_n(\varphi,\theta-\theta',\xi) - a_n(\varphi,\theta,\xi) \big] e^{-\mathrm{i}\,\theta'\xi} d\theta'$$

and

$$\sigma_{n,2}(\varphi,\theta,\xi) = \frac{1}{\pi} \int_{\mathbb{T}} \frac{\rho_n(\varphi,\theta+\eta) - \rho_n(\varphi,\theta)}{e^{\mathrm{i}\eta} - 1} \sum_{\xi' \in \mathbb{Z}} \lambda(\xi+\xi') a_n(\varphi,\theta,\xi+\xi') (1-e^{\mathrm{i}\eta}) e^{-\mathrm{i}\xi'\eta} d\eta.$$

Using the identities (7.6) and (7.5)

$$(1 - e^{\mathrm{i}\eta})e^{-\mathrm{i}\xi'\eta} = \overline{\Delta}_{\xi'}e^{-\mathrm{i}\xi'\eta} \quad \text{and} \quad \sum_{\substack{\xi' \in \mathbb{Z} \\ 110}} \overline{\Delta}_{\xi'}f(\xi')g(\xi') = -\sum_{\substack{\xi' \in \mathbb{Z} \\ \xi' \in \mathbb{Z}}} f(\xi')\Delta_{\xi'}g(\xi')$$

we deduce that

$$\sigma_{n,2}(\varphi,\theta,\xi) = \frac{1}{\pi} \int_{\mathbb{T}} \frac{\rho_n(\varphi,\theta+\eta) - \rho_n(\varphi,\theta)}{1 - e^{\mathrm{i}\eta}} \sum_{\xi' \in \mathbb{Z}} e^{\mathrm{i}\,(\xi-\xi')\eta} \Delta_{\xi'} \big[\lambda(\xi')a_n(\varphi,\theta,\xi')\big] d\eta.$$
(8.146)

According to the relation (8.141) we get that the symbol a_n satisfies the recursive relation

$$a_{n+1}(\varphi,\theta,\xi) = \sigma_{n,1}(\varphi,\theta,\xi) + \sigma_{n,2}(\varphi,\theta,\xi) + \mu_n \tag{8.147}$$

where μ_n is the symbol associated to $\mathcal{R}_n K_n$. Let us start with estimating $\sigma_{n,1}$. For $\gamma \in \mathbb{N}$, we use the Leibniz rule (7.4)

$$\Delta_{\xi}^{\gamma}\sigma_{n,1}(\varphi,\theta,\xi) = 2\rho_n(\varphi,\theta) \sum_{0 \leqslant \beta \leqslant \gamma} {\gamma \choose \beta} \int_{\mathbb{T}} \mathscr{K}_0(\theta') \Delta_{\xi}^{\beta} \left[a_n(\varphi,\theta-\theta',\xi) - a_n(\varphi,\theta,\xi) \right] \Delta_{\xi}^{\gamma-\beta} \left[e^{-\mathrm{i}\,\theta'\xi} \right] (\xi+\beta)d\theta'.$$

Applying (7.6) gives for large ξ (such that $\xi + \beta \neq 0$)

$$\Delta_{\xi}^{\gamma-\beta}[e^{-\mathrm{i}\,\theta'\xi}](\xi+\beta) = \left(e^{-\mathrm{i}\,\theta'}-1\right)^{\gamma-\beta}e^{-\mathrm{i}\,\theta'(\xi+\beta)}$$

$$= i^{\gamma-\beta}\left(e^{-\mathrm{i}\,\theta'}-1\right)^{\gamma-\beta}(\xi+\beta)^{\beta-\gamma}\partial_{\theta'}^{\gamma-\beta}e^{-\mathrm{i}\,\theta'(\xi+\beta)}.$$
(8.148)

Hence, integration by parts allows to get for large ξ

$$\begin{split} \langle \xi \rangle^{-m+\gamma} \Delta_{\xi}^{\gamma} \sigma_{n,1}(\varphi,\theta,\xi) = & 2\rho_n(\varphi,\theta) \sum_{\substack{0 \leq \beta \leq \gamma \\ 0 \leq \varrho \leq \gamma - \beta}} (-i)^{\alpha-\beta} \binom{\gamma}{\beta} \binom{\gamma-\beta}{\varrho} \int_{\mathbb{T}} \partial_{\theta'}^{\gamma-\beta-\varrho} \Big[(1-e^{-\mathrm{i}\theta'})^{\gamma-\beta} \mathscr{K}_{0}(\theta') \Big] \\ & \frac{\langle \xi \rangle^{-m+\gamma}}{(\xi+\beta)^{\gamma-\beta}} \partial_{\theta'}^{\varrho} \Big[\Delta_{\xi}^{\beta} a_n(\varphi,\theta-\theta',\xi) - \Delta_{\xi}^{\beta} a_n(\varphi,\theta,\xi) \Big] e^{-\mathrm{i}\theta'(\xi+\beta)} d\theta' \\ & \triangleq 2\rho_n(\varphi,\theta) \sum_{\substack{0 \leq \beta \leq \gamma \\ 0 \leq \varrho \leq \gamma-\beta}} \binom{\gamma}{\beta} \binom{\gamma}{\varrho} \binom{\gamma-\beta}{\varrho} \mathcal{T}_n^{\beta,\varrho}(\varphi,\theta,\xi). \end{split}$$

From straightforward computations we get for any $\epsilon > 0$

$$\left|\partial_{\theta'}^{\gamma-\beta-\varrho}\left[(1-e^{-\mathrm{i}\theta'})^{\gamma-\beta}\partial_{\alpha}^{j}\mathscr{K}_{0}(\theta')\right]\right| \lesssim |\sin(\theta'/2)|^{\varrho-1-\alpha-\epsilon}.$$

Hence we get from Sobolev embeddings that for $\gamma-\beta \geqslant \varrho \geq 1$

$$\sup_{\substack{\alpha \in (0,\overline{\alpha}), \\ 0 \leqslant j \leqslant q}} \kappa^{j} \left\| \partial_{\alpha}^{j} \mathcal{T}_{n}^{\beta,\varrho}(\cdot, \bullet, \xi) \right\|_{H^{s-j}} \lesssim \sup_{\substack{\alpha \in (0,\overline{\alpha}), \\ 0 \leqslant j \leqslant q}} \int_{\mathbb{T}} |\sin(\theta'/2)|^{-\alpha-\epsilon} \langle \xi \rangle^{-m+\beta} \kappa^{j} \left\| \Delta_{\xi}^{\beta} \partial_{\alpha}^{j} a_{n}(\cdot, \bullet, \xi) \right\|_{H^{s+\varrho-j}} d\theta' \\ \lesssim \||a_{n}\||_{m,s+\varrho,\beta}^{q,\kappa} \\ \lesssim \||a_{n}\||_{m,s+\gamma-\beta,\beta}^{q,\kappa} .$$

For $\rho = 0$ we proceed as follows using in particular Lemma 4.7,

$$\begin{split} \sup_{\substack{\alpha \in (0,\overline{\alpha}), \\ 0 \leq j \leq q}} \kappa^{j} \left\| \partial_{\alpha}^{q} \mathcal{T}_{n}^{\beta,0}(\cdot, \bullet, \xi) \right\|_{H^{s-j}} &\lesssim \sup_{\substack{\alpha \in (0,\overline{\alpha}), \\ 0 \leq j \leq q}} \int_{\mathbb{T}} |\sin(\theta'/2)|^{-\alpha - \epsilon} \langle \xi \rangle^{-m + \beta} \\ &\times \kappa^{j} \left\| \frac{\Delta_{\xi}^{\beta} \partial_{\alpha}^{j} a_{n}(\cdot, \bullet - \theta', \xi) - \Delta_{\xi}^{\beta} \partial_{\alpha}^{j} a_{n}(\cdot, \bullet, \xi)}{1 - e^{-\mathrm{i}\theta'}} \right\|_{H^{s-j}} d\theta' \\ &\lesssim \||a_{n}\||_{m,s+1,\beta}^{q,\kappa} \\ &\lesssim \||a_{n}\||_{m,s+1,\gamma}^{q,\kappa} \,. \end{split}$$

~

Notice that the treatment of the quantities related to the differentiation with respect to the parameter ω instead of α can be done in a similar way and we get the same kind of estimates. Therefore, performing the law products in Lemma 4.1 with the preceding estimates yields

$$\| \sigma_{n,1} \|_{m,s,\gamma}^{q,\kappa} \lesssim \| \rho_n \|_{s_0}^{q,\kappa} \| a_n \|_{m,s+1,\gamma}^{q,\kappa} + \| \rho_n \|_s^{q,\kappa} \| a_n \|_{m,s_0+1,\gamma}^{q,\kappa}$$

+
$$\sum_{0 \leqslant \beta \leqslant \gamma} \left(\| \rho_n \|_{s_0}^{q,\kappa} \| a_n \|_{m,s+\gamma-\beta,\beta}^{q,\kappa} + \| \rho_n \|_s^{q,\kappa} \| a_n \|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa} \right).$$
 (8.149)

Next, we shall move to the estimate of the term $\sigma_{n,2}$ described in (8.146). Define

$$\zeta_n(\varphi,\theta,\xi) = \frac{1}{\pi} \int_{\mathbb{T}} \frac{\rho_n(\varphi,\theta+\eta) - \rho_n(\varphi,\theta)}{1 - e^{i\eta}} e^{i\xi\eta} d\eta, \qquad (8.150)$$

then it is clear that

$$\sigma_{n,2}(\varphi,\theta,\xi) = \sum_{\xi' \in \mathbb{Z}} \zeta_n(\varphi,\theta,\xi-\xi') \Delta_{\xi'} \big[\lambda(\xi') a_n(\varphi,\theta,\xi') \big].$$

From the convolution structure, we infer

$$\Delta_{\xi}^{\gamma}\sigma_{n,2}(\varphi,\theta,\xi) = \sum_{\xi'\in\mathbb{Z}}\zeta_n(\varphi,\theta,\xi-\xi')\Delta_{\xi'}^{1+\gamma}\big[\lambda(\xi')a_n(\varphi,\theta,\xi')\big]$$

On the other hand, using (8.145), (4.19) and (4.21), we may write

$$\lambda(\xi) = \left(\partial_{\theta} |\mathbf{D}|^{\alpha - 1} e^{-i\,\theta\xi}\right)_{|\theta = 0}$$
$$= -i \frac{2^{\alpha} \Gamma(1 - \alpha)}{\Gamma(\frac{\alpha}{2}) \Gamma(1 - \frac{\alpha}{2})} \frac{\xi \, \Gamma\left(|\xi| + \frac{\alpha}{2}\right)}{\Gamma\left(|\xi| + 1 - \frac{\alpha}{2}\right)}$$

From the expansion of Gamma function, similarly to (4.23) and based on Lemma A.1-(ii), one finds by straightforward computations that for any $j, \beta \in \mathbb{N}, \epsilon > 0$

$$\sup_{\alpha \in (0,\overline{\alpha})} \left| \Delta_{\xi}^{\beta} \partial_{\alpha}^{j} \lambda(\xi) \right| \lesssim \langle \xi \rangle^{\overline{\alpha} + \epsilon - \beta}.$$
(8.151)

Consequently, applying Leibniz formula (7.4) leads to

$$\left\|\Delta_{\xi'}^{1+\gamma} \left[\lambda(\xi')a_n(\cdot, \cdot, \xi')\right]\right\|_s^{q,\kappa} \lesssim \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma} \parallel a_n \parallel_{m,s,1+\gamma}^{q,\kappa}$$

and

$$\left\|\Delta_{\xi'}^{1+\gamma}\left[\lambda(\xi')a_n(\cdot,\cdot,\xi')\right]\right\|_s^{q,\kappa} \lesssim \langle\xi'\rangle^{m+\overline{\alpha}+\epsilon} \parallel a_n \parallel_{m,s,0}^{q,\kappa}$$

Integration by parts in (8.150), using Lemma 4.7, implies for any $N \in \mathbb{N}$

$$\langle \xi \rangle^N \| \zeta_n(\cdot, \cdot, \xi) \|_s^{q,\kappa} \lesssim \| \rho_n \|_{s+1+N}^{q,\kappa}.$$

Putting together the preceding estimates with the law products of Lemma 4.1 we find for any $N_1, N_2 \in \mathbb{N}$

$$\begin{split} \left\| \Delta_{\xi}^{\gamma} \sigma_{n,2}(\cdot, \cdot, \xi) \right\|_{s}^{q,\kappa} \lesssim \|\rho_{n}\|_{s_{0}+1+N_{1}}^{q,\kappa} \|\| a_{n} \|\|_{m,s,0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{1}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon} \\ + \|\rho_{n}\|_{s_{0}+1+N_{2}}^{q,\kappa} \|\| a_{n} \|\|_{m,s,1+\gamma}^{q,\kappa} \sum_{\substack{\frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{2}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma} \\ + \|\rho_{n}\|_{s+1+N_{1}}^{q,\kappa} \|\| a_{n} \|\|_{m,s_{0},0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{1}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon} \\ + \|\rho_{n}\|_{s+1+N_{2}}^{q,\kappa} \|\| a_{n} \|\|_{m,s_{0},1+\gamma}^{q,\kappa} \sum_{\substack{\frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi| \\ \frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{2}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma}. \tag{8.152}$$

Since $\overline{\alpha} \in [0,1)$ then one can check that with the choice $N_1 = \gamma + 2 + |m|$ and $N_2 = 2$, then

$$\sup_{\xi \in \mathbb{Z}} \langle \xi \rangle^{-m+\gamma} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_1} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon} < \infty$$

and

$$\sup_{\xi \in \mathbb{Z}} \langle \xi \rangle^{-m+\gamma} \sum_{\frac{1}{2} |\xi| \le |\xi'| \le 2|\xi|} \langle \xi - \xi' \rangle^{-N_2} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma} < \infty.$$

Hence

$$\| \sigma_{n,2} \|_{m,s,\gamma}^{q,\kappa} \lesssim \| \rho_n \|_{s_0+3+\gamma+|m|}^{q,\kappa} \| \| a_n \|_{m,s,0}^{q,\kappa} + \| \rho_n \|_{s_0+3}^{q,\kappa} \| \| a_n \|_{m,s,\gamma+1}^{q,\kappa} + \| \rho_n \|_{s+3+\gamma+|m|}^{q,\kappa} \| \| a_n \|_{m,s_0,0} + \| \rho_n \|_{s+3}^{q,\kappa} \| \| a_n \|_{m,s_0,\gamma+1}^{q,\kappa} .$$

$$(8.153)$$

Next, we shall move to the estimate of the symbol associated to $\Re_{n,1}K_n$ introduced in (8.142). This symbol will be denoted by $\mu_{n,1}$ and can be recovered from the formula (8.139). By direct computations based on (4.19) and (8.140) we can check that

$$\mu_{n,1}(\varphi,\theta,\xi) = 2\partial_{\theta}\rho_n(\varphi,\theta) \int_{\mathbb{T}} \frac{a_n(\varphi,\theta-\theta',\xi)}{|\sin(\theta'/2)|^{\alpha}} e^{-\mathrm{i}\,\theta'\xi} d\theta'$$
$$\triangleq 2\partial_{\theta}\rho_n(\varphi,\theta)\mathcal{T}_n(\varphi,\theta,\xi).$$

Then proceeding as for the term $\sigma_{n,1}$ one gets

$$\langle \xi \rangle^{-m+\gamma} \left\| \Delta_{\xi}^{\gamma} \mathcal{T}_{n}(\cdot, \cdot, \xi) \right\|_{s}^{q,\kappa} \lesssim \sum_{0 \leq \beta \leq \gamma} \left\| \left\| a_{n} \right\| \right\|_{m,s+\gamma-\beta,\beta}^{q,\kappa}$$

and by the law products we infer

$$\|\|\mu_{n,1}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \left(\|\rho_n\|_{s_0+1}^{q,\kappa} \|\|a_n\|\|_{m,s+\gamma-\beta,\beta}^{q,\kappa} + \|\rho_n\|_{s+1}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa} \right).$$
(8.154)

The symbol $\mu_{n,2}$ associated to $\Re_{n,2}K_n$ defined in (8.142) can be recovered from a_n in a similar way to (8.144) and one gets

$$\mu_{n,2}(\varphi,\theta,\xi) = \frac{1}{2\pi} \int_{\mathbb{T}} \partial_{\eta} \rho_n(\varphi,\theta+\eta) \sum_{\xi' \in \mathbb{Z}} \lambda_1(\xi') a_n(\varphi,\theta,\xi') e^{i(\xi-\xi')\eta} d\eta$$
$$= \sum_{\xi' \in \mathbb{Z}} \lambda_1(\xi') a_n(\varphi,\theta,\xi') \zeta_{n,1}(\varphi,\theta,\xi-\xi'), \qquad (8.155)$$

with

$$\zeta_{n,1}(\varphi,\theta,\xi) = \frac{1}{2\pi} \int_{\mathbb{T}} \partial_{\eta} \rho_n(\varphi,\theta+\eta) e^{i\,\xi\eta} d\eta$$

and

$$\lambda_1(\xi) = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{e^{\mathrm{i}\theta'\xi}}{|\sin(\theta'/2)|^{\alpha}} d\theta'.$$

By applying (4.19) and (4.21), we find that

$$\lambda_1(\xi) = \left(|\mathbf{D}|^{\alpha - 1} e^{-\mathbf{i}\,\theta\xi} \right)_{|\theta = 0}$$
$$= \frac{2^{\alpha} \Gamma(1 - \alpha)}{\Gamma(\frac{\alpha}{2}) \Gamma(1 - \frac{\alpha}{2})} \frac{\Gamma(|\xi| + \frac{\alpha}{2})}{\Gamma(|\xi| + 1 - \frac{\alpha}{2})}$$

From Lemma A.1 we get that for any $j, \beta \in \mathbb{N}, \epsilon > 0$

$$\sup_{\alpha \in (0,\overline{\alpha})} \left| \Delta_{\xi}^{\beta} \partial_{\alpha}^{j} \lambda_{1}(\xi) \right| \lesssim \langle \xi \rangle^{\overline{\alpha} - 1 + \epsilon - \beta}.$$
(8.156)

Now, according to the convolution structure in the sum (8.155) we infer

$$\Delta_{\xi}^{\gamma}\mu_{n,2}(\varphi,\theta,\xi) = \sum_{\xi'\in\mathbb{Z}}\zeta_{n,1}(\varphi,\theta,\xi-\xi')\,\Delta_{\xi'}^{\gamma}\big[\lambda_1(\xi')a_n(\varphi,\theta,\xi')\big].$$

Using the law products with (8.156) and performing similar arguments as in (8.152) we get for any $N_1, N_2 \in \mathbb{N}$

$$\begin{split} \left\| \Delta_{\xi}^{\gamma} \mu_{n,2}(\cdot, \cdot, \xi) \right\|_{s}^{q,\kappa} \lesssim &\|\rho_{n}\|_{s_{0}+N_{1}+1}^{q,\kappa} \|\|a_{n}\|\|_{m,s,0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{1}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1} \\ &+ \|\rho_{n}\|_{s_{0}+N_{2}+1}^{\kappa} \|\|a_{n}\|\|_{m,s,\gamma}^{q,\kappa} \sum_{\substack{\frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi| \\ |\xi'| \leq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{2}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma} \\ &+ \|\rho_{n}\|_{s+N_{1}+1}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0},0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{1}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1} \\ &+ \|\rho_{n}\|_{s+N_{2}+1}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0},\gamma}^{q,\kappa} \sum_{\substack{\frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi| \\ |\xi'| \leq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{2}} \langle \xi' \rangle^{m+\overline{\alpha}+\epsilon-1-\gamma}. \end{split}$$

Fixing $N_1 = \gamma + 1 + |m|$ and $N_2 = 2$ then

$$\| \mu_{n,2} \|_{m,s,\gamma}^{q,\kappa} \lesssim \| \rho_n \|_{s_0+2+\gamma+|m|}^{q,\kappa} \| \| a_n \|_{m,s,0}^{q\kappa} + \| \rho_n \|_{s_0+3}^{q,\kappa} \| \| a_n \|_{m,s,\gamma}^{q,\kappa} + \| \rho_n \|_{s+2+\gamma+|m|}^{q,\kappa} \| \| a_n \|_{m,s_0,0}^{q,\kappa} + \| \rho_n \|_{s+3}^{q,\kappa} \| \| a_n \|_{m,s_0,\gamma}^{q,\kappa} .$$

$$(8.157)$$

As to the term $\mathcal{R}_{n,3}K_n$ in (8.142) we first write in view of (4.19)

$$\partial_{\theta} \left[|\mathbf{D}_{\theta}|^{\alpha-1}, \rho_n \right] K_n(\varphi, \theta, \eta) = \frac{1}{2\pi} \partial_{\theta} \int_{\mathbb{T}}^{\mathbf{r}} \frac{\rho_n(\varphi, \theta - \theta') - \rho_n(\varphi, \theta)}{|\sin(\theta'/2)|^{\alpha}} K_n(\varphi, \theta - \theta', \eta) d\theta'$$
$$= \int_{\mathbb{T}} \Psi(\varphi, \theta, \theta') K_n(\varphi, \theta - \theta', \eta) d\theta',$$

with

$$\Psi(\varphi,\theta,\theta') = -\frac{\alpha}{4\pi} \frac{\rho_n(\varphi,\theta-\theta') - \rho_n(\varphi,\theta)}{\sin(\theta'/2)} \frac{\cos(\theta'/2)}{|\sin(\theta'/2)|^{\alpha}} - \frac{\partial_\theta \rho_n(\varphi,\theta)}{2\pi |\sin(\theta'/2)|^{\alpha}}.$$
(8.158)

Therefore

$$\partial_{\theta} \left[|\mathcal{D}_{\theta}|^{\alpha-1}, \rho_n \right] K_n(\varphi, \theta, \eta) = \frac{1}{2\pi} \sum_{\xi \in \mathbb{Z}} e^{i(\theta-\eta)\xi} \int_{\mathbb{T}} \Psi(\varphi, \theta, \theta') a_n(\varphi, \theta - \theta', \xi) e^{-i\theta'\xi} d\theta'$$
$$= \sum_{\xi \in \mathbb{Z}} \mu_{n,3}(\varphi, \theta, \xi) e^{i(\theta-\eta)\xi}$$

where the symbol $\mu_{n,3}$ is given by

$$\mu_{n,3}(\varphi,\theta,\xi) = \frac{1}{2\pi} \int_{\mathbb{T}} \Psi(\varphi,\theta,\theta') a_n(\varphi,\theta-\theta',\xi) e^{-\mathrm{i}\theta'\xi} d\theta'$$

We shall proceed as for $\sigma_{n,1}$ and one first writes by (7.4)

$$\Delta_{\xi}^{\gamma}\mu_{n,3}(\varphi,\theta,\xi) = \frac{1}{2\pi} \sum_{0 \le \beta \leqslant \gamma} {\gamma \choose \beta} \int_{\mathbb{T}} \Psi(\varphi,\theta,\theta') \Delta_{\xi}^{\beta} \left[a_n(\varphi,\theta-\theta',\xi) \right] \Delta_{\xi}^{\gamma-\beta} \left[e^{-\mathrm{i}\,\theta'\xi} \right] (\xi+\beta) d\theta'.$$

Using (8.148) then integration by parts implies

$$\langle \xi \rangle^{-m} \xi^{\gamma} \Delta_{\xi}^{\gamma} \mu_{n,3}(\varphi,\theta,\xi) = \sum_{\substack{0 \leqslant \beta \leqslant \gamma \\ 0 \leqslant \eta \leqslant \gamma - \beta}} \binom{\gamma}{\beta} \binom{\gamma-\beta}{\eta} i^{\gamma-\beta} \frac{\xi^{\gamma-\beta}}{(\xi+\beta)^{\gamma-\beta}} \int_{\mathbb{T}} \partial_{\theta'}^{\gamma-\beta-\eta} \Big[(1-e^{-\mathrm{i}\theta'})^{\gamma-\beta} \Psi(\varphi,\theta,\theta') \Big]$$

$$\langle \xi \rangle^{-m} \xi^{\beta} \partial_{\theta'}^{\eta} \Big[\Delta_{\xi}^{\beta} a_n(\varphi, \theta - \theta', \xi) \Big] e^{-\mathrm{i}\,\theta'(\xi+\beta)} d\theta'.$$

From (8.158) we get through straightforward calculus

$$\left\|\partial_{\theta'}^{\gamma-\beta-\eta}\left[(1-e^{-\mathrm{i}\theta'})^{\gamma-\beta}\Psi(\cdot,\bullet,\theta')\right]\right\|_{s}^{q,\kappa} \lesssim \frac{\|\rho_{n}\|_{s+1+\gamma-\beta-\eta}^{q,\kappa}}{|\sin(\theta'/2)|^{\overline{\alpha}+\epsilon}}.$$

Hence we deduce from the law products stated in Lemma 4.1

$$\|\mu_{n,3}\|_{m,s,\gamma}^{q,\kappa} \lesssim \sum_{\substack{0 \le \beta \le \gamma \\ 0 \le \eta \le \gamma - \beta}} \|\rho_n\|_{s_0+1+\gamma-\beta-\eta}^{q,\kappa} \|a_n\|_{m,s+\eta,\beta}^{q,\kappa} + \sum_{\substack{0 \le \beta \le \gamma \\ 0 \le \eta \le \gamma - \beta}} \|\rho_n\|_{s+1+\gamma-\beta-\eta}^{q,\kappa} \|a_n\|_{m,s_0+\eta,\beta}^{q,\kappa}.$$
(8.159)

Concerning the term $\mathcal{R}_{n,4}K_n$ in (8.142) we first write

$$\partial_{\eta} \left[|\mathcal{D}_{\eta}|^{\alpha-1}, \rho_n \right] K_n(\varphi, \theta, \eta) = \frac{1}{2\pi} \partial_{\eta} \int_{\mathbb{T}} \frac{\rho_n(\varphi, \eta - \eta') - \rho_n(\varphi, \eta)}{|\sin(\eta'/2)|^{\alpha}} K_n(\varphi, \theta, \eta - \eta') d\eta'$$
$$= \int_{\mathbb{T}} \Psi(\varphi, \eta, \eta') K_n(\varphi, \theta, \eta - \eta') d\eta'$$
(8.160)

where Ψ is introduced in (8.158). Therefore the associated symbol is given by

$$\mu_{n,4}(\varphi,\theta,\xi) = \frac{1}{2\pi} \sum_{\xi' \in \mathbb{Z}} a_n(\varphi,\theta,\xi') \int_{\mathbb{T}} \left(\int_{\mathbb{T}} \Psi(\varphi,\theta+\eta,\eta') e^{i\eta'\xi'} d\eta' \right) e^{i(\xi-\xi')\eta} d\eta$$

From the summation by parts (7.5) and Leibniz rule (7.4) we infer

$$\begin{split} \Delta_{\xi}^{\gamma} \mu_{n,4}(\varphi,\theta,\xi) &= \frac{1}{2\pi} \sum_{\xi' \in \mathbb{Z}} \int_{\mathbb{T}} \Delta_{\xi'}^{\gamma} \left[a_n(\varphi,\theta,\xi') \left(\int_{\mathbb{T}} \Psi(\varphi,\theta+\eta,\eta') e^{i\eta'\xi'} d\eta' \right) \right] e^{i(\xi-\xi')\eta} d\eta \\ &= \frac{1}{2\pi} \sum_{\substack{\xi' \in \mathbb{Z} \\ 0 \leqslant \beta \leqslant \gamma}} \binom{\gamma}{\beta} \Delta_{\xi'}^{\beta} a_n(\varphi,\theta,\xi') \int_{\mathbb{T}} \left(\Delta_{\xi'}^{\gamma-\beta} \mathscr{P}(\varphi,\theta,\eta,\xi'+\beta) \right) e^{i(\xi-\xi')\eta} d\eta, \end{split}$$

with

$$\mathscr{P}(\varphi,\theta,\eta,\xi) \triangleq \int_{\mathbb{T}} \Psi(\varphi,\theta+\eta,\eta') e^{\mathrm{i}\eta'\xi'} d\eta'.$$

By virtue of (8.148) combined with integration by parts, we find for any $N \in \mathbb{N}$ and $0 \leq \varrho \leq \gamma - \beta$

$$\begin{split} \left\| \int_{\mathbb{T}} \left(\Delta_{\xi'}^{\gamma-\beta} \mathscr{P}(\varphi,\theta,\eta,\xi'+\beta) \right) e^{-\mathrm{i}(\xi-\xi')\eta} d\eta \right\|_{s}^{q,\kappa} \lesssim \int_{\mathbb{T}} \left\| \partial_{\eta'}^{\varrho} \partial_{\eta}^{N} \left((1-e^{-\mathrm{i}\eta'})^{\varrho} \Psi(\cdot, \cdot+\eta,\eta') \right) \right\|_{s}^{q,\kappa} d\eta' d\eta \\ \times \langle \xi - \xi' \rangle^{-N} \langle \xi' \rangle^{-\varrho}. \end{split}$$

Using (8.158) one obtains

$$\int_{\mathbb{T}} \left\| \partial_{\eta'}^{\varrho} \partial_{\eta}^{N} \left((1 - e^{-\mathrm{i}\eta'})^{\gamma - \beta} \Psi(\cdot, \cdot + \eta, \eta') \right) \right\|_{s}^{q,\kappa} d\eta' d\eta \lesssim \|\rho_{n}\|_{s+1+N+\varrho}^{q,\kappa} \int_{\mathbb{T}} \frac{d\eta'}{|\sin(\eta'/2)|^{\overline{\alpha} + \epsilon}} \lesssim \|\rho_{n}\|_{s+1+N+\varrho}^{q,\kappa}$$

Hence using the law products we get for any $N_1, N_2 \in \mathbb{N}$

$$\begin{split} \left\| \Delta_{\xi}^{\gamma} \mu_{n,4}(\cdot, \cdot, \xi) \right\|_{s}^{q,\kappa} \lesssim \|\rho_{n}\|_{s_{0}+1+N_{1}}^{q,\kappa} \|\|a_{n}\|\|_{m,s,0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{1}} \langle \xi' \rangle^{m} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|a_{n}\|\|_{m,s,\beta}^{q,\kappa} \|\rho_{n}\|_{s_{0}+1+N_{2}+\gamma-\beta}^{q,\kappa} \sum_{\substack{\frac{1}{2}|\xi| \leqslant |\xi'| \leqslant 2|\xi|}} \langle \xi - \xi' \rangle^{-N_{2}} \langle \xi' \rangle^{m-\gamma} \end{split}$$

$$+ \|\rho_n\|_{s+1+N_1}^{q,\kappa} \|\|a_n\|\|_{m,s_0,0}^{q,\kappa} \sum_{\substack{|\xi'| \leq \frac{1}{2}|\xi| \\ |\xi'| \geq 2|\xi|}} \langle \xi - \xi' \rangle^{-N_1} \langle \xi' \rangle^m$$

$$+ \sum_{0 \leq \beta \leq \gamma} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} \|\rho_n\|_{s+1+N_2+\gamma-\beta}^{q,\kappa} \sum_{\frac{1}{2}|\xi| \leq |\xi'| \leq 2|\xi|} \langle \xi - \xi' \rangle^{-N_2} \langle \xi' \rangle^{m-\gamma}.$$

Then making the choice $N_1 = \gamma + 2 + |m|$ and $N_2 = 2$ yields

$$\|\|\mu_{n,4}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\rho_n\|_{s_0+3+\gamma+|m|}^{q,\kappa} \|\|a_n\|\|_{m,s,0}^{q,\kappa} + \|\rho_n\|_{s+3+\gamma+|m|}^{q,\kappa} \|\|a_n\|\|_{m,s_0,0}^{q,\kappa} + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} \|\rho_n\|_{s+3+\gamma-\beta}^{q,\kappa} + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} \|\rho_n\|_{s+3+\gamma-\beta}^{q,\kappa}.$$

$$(8.161)$$

It remains to analyze the last term $\mu_{n,5}$ in (8.142). Similarly to (8.160) we get

$$|D_{\eta}|^{\alpha-1}(\partial_{\eta}\rho) K_{n}(\varphi,\theta,\eta) = \frac{1}{2\pi} \int_{\mathbb{T}} \frac{\partial_{\eta}\rho_{n}(\varphi,\eta-\eta')}{|\sin(\eta'/2)|^{\alpha}} K_{n}(\varphi,\theta,\eta-\eta') d\eta'$$
$$\triangleq \int_{\mathbb{T}} \Psi_{1}(\varphi,\eta,\eta') K_{n}(\varphi,\theta,\eta-\eta') d\eta'. \tag{8.162}$$

Therefore the associated symbol takes the form,

$$\mu_{n,5}(\varphi,\theta,\xi) = \frac{1}{2\pi} \sum_{\xi' \in \mathbb{Z}} a_n(\varphi,\theta,\xi') \int_{\mathbb{T}} \left(\int_{\mathbb{T}} \Psi_1(\varphi,\theta+\eta,\eta') e^{\mathrm{i}\eta'\xi'} d\eta' \right) e^{-\mathrm{i}(\xi-\xi')\eta} d\eta.$$

By reproducing exactly the same lines of the proof of (8.161) we find

$$\|\mu_{n,5}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\rho_{n}\|_{s_{0}+3+\gamma+|m|}^{q,\kappa} \|\|a_{n}\|\|_{m,s,0}^{q,\kappa} \|\rho_{n}\|_{s+3+\gamma+|m|}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0},0}^{q,\kappa} + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|a_{n}\|\|_{m,s_{0},\beta}^{q,\kappa} \|\rho_{n}\|_{s_{0}+3+\gamma-\beta}^{q,\kappa} + \sum_{0 \leqslant \beta \leqslant \gamma} \|\|a_{n}\|\|_{m,s_{0},\beta}^{q,\kappa} \|\rho_{n}\|_{s+3+\gamma-\beta}^{q,\kappa}.$$

$$(8.163)$$

Putting together (8.147), (8.149), (8.153), (8.154), (8.157), (8.159), (8.161) and (8.163) leads to $\||a_{n+1}\||_{m=0}^{q,\kappa} \leq \|\rho_n\|_{m=0}^{q,\kappa} < \|\rho_n\|_{m=0}^{q,\kappa}$

$$\|a_{n+1}\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\rho_{n}\|_{s_{0}+3+\gamma+|m|}^{q,\kappa} \|\|a_{n}\|_{m,s,0}^{q,\kappa} \|\rho_{n}\|_{s+3+\gamma+|m|}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0},0}^{q,\kappa} + \|\rho_{n}\|_{s_{0}}^{q,\kappa} \|\|a_{n}\|_{m,s+1,\gamma}^{q,\kappa} + \|\rho_{n}\|_{s}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0}+1,\gamma}^{q,\kappa} + \|\rho_{n}\|_{s_{0}+3}^{q,\kappa} \|\|a_{n}\|\|_{m,s,\gamma+1}^{q,\kappa} + \|\rho_{n}\|_{q,s+3}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0},\gamma+1}^{q,\kappa} + \sum_{\substack{0 \le \beta \le \gamma \\ 0 \le \eta \le \gamma - \beta}} \|\rho_{n}\|_{s_{0}+3+\gamma-\beta-\eta}^{q,\kappa} \|\|a_{n}\|\|_{m,s_{0}+\eta,\beta}^{q,\kappa} .$$

$$(8.164)$$

Now by interpolation we get for any $0\leqslant\eta\leqslant\gamma-\beta$

$$|||a_n|||_{m,s+\eta,\beta}^{q,\kappa} \le \left(|||a_n|||_{m,s,\beta}^{q,\kappa}\right)^{1-\frac{\eta}{\gamma-\beta}} \left(|||a_n|||_{m,s+\gamma-\beta,\beta}^{q,\kappa}\right)^{\frac{\eta}{\gamma-\beta}}.$$

and

$$\|\rho_n\|_{s_0+3+\gamma-\eta-\beta}^{q,\kappa} \le \left(\|\rho_n\|_{s_0+3+\gamma-\beta}^{q,\kappa}\right)^{1-\frac{\eta}{\gamma-\beta}} \left(\|\rho_n\|_{s_0+3}^{q,\kappa}\right)^{\frac{\eta}{\gamma-\beta}}.$$

Consequently

$$\|\|a_n\|\|_{m,s+\eta,\beta}^{q,\kappa} \|\rho_n\|_{s_0+3+\gamma-\eta-\beta}^{q,\kappa} \leq \left(\|\rho_n\|_{s_0+3+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s,\beta}^{q,\kappa}\right)^{1-\frac{\eta}{\gamma-\beta}} \left(\|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s+\gamma-\beta,\beta}^{q,\kappa}\right)^{\frac{\eta}{\gamma-\beta}} \\ \lesssim \|\rho_n\|_{s_0+3+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s,\beta}^{q,\kappa} + \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s+\gamma-\beta,\beta}^{q,\kappa}.$$

Similar arguments yield

 $\|\rho_n\|_{s+3+\gamma-\beta-\eta}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\eta,\beta}^{q,\kappa} \lesssim \|\rho_n\|_{s+3+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa}.$ Therefore we deduce from (8.164) and Sobolev embeddings

$$\begin{aligned} \|\|a_{n+1}\|\|_{m,s,\gamma}^{q,\kappa} \lesssim \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s+1,\gamma}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+1,\gamma}^{q,\kappa} \\ &+ \|\rho_n\|_{s_0+3}^{q,\kappa} \|\|a_n\|\|_{m,s,\gamma+1}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0,\gamma+1}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho_n\|_{s_0+3+|m|+\gamma-\beta}^{q,\kappa} \|\|a_n\|\|_{m,s,\beta}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa} \\ &+ \sum_{0 \leqslant \beta \leqslant \gamma} \|\rho_n\|_{s+3+|m|+\gamma-\beta}^{\kappa} \|\|a_n\|\|_{m,s_0,\beta}^{q,\kappa} + \|\rho_n\|_{s+3}^{q,\kappa} \|\|a_n\|\|_{m,s_0+\gamma-\beta,\beta}^{q,\kappa} \\ &= \sum_{\ell=1}^6 \mathcal{T}_\ell. \end{aligned}$$

$$(8.165)$$

This achieves the proof of the first point.

(ii) Let $s \ge s_0, \gamma, n \in \mathbb{N}$ be fixed numbers and assume (8.143). We will check that for any integer $k \in [0, n]$ we have the following property: $\forall s' \ge s_0, \gamma' \in \mathbb{N}$ with $s' + \gamma' \le s + \gamma + n - k$, we have

$$|||a_k|||_{m,s',\gamma'}^{q,\kappa} \lesssim F(s_0+3) \sum_{i=0}^{k-1} ||\rho_i||_{s'+\gamma'+k+4+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} ||\rho_j||_{s_0+3+|m|}^{q,\kappa} + F(s'+\gamma'+k+4) \prod_{j=0}^{k-1} ||\rho_j||_{s_0+3+|m|}^{q,\kappa},$$

$$(8.166)$$

where we take for k = 0 the convention $\sum_{i=0}^{-1} \dots = 0$ and $\prod_{j=0}^{-1} = 1$ in such a way that the inequality

remains true for k = 0 according to the assumption (8.143). We shall proceed with finite induction principle over k. Assume that the estimate (8.166) is true at the order $k \leq n-1$ and let us show that it remains true at the order k + 1. Set

$$U_k(s,\gamma) \triangleq F(s_0+3) \sum_{i=0}^{k-1} \|\rho_i\|_{s+\gamma+k+4+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa} + F(s+\gamma+k+4) \prod_{j=0}^{k-1} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa}.$$

We intend to prove that for any $s' \ge s_0, \gamma' \in \mathbb{N}$ with $s' + \gamma' \le s + \gamma + n - k - 1$, we have

$$|||a_{k+1}|||_{m,s',\gamma'}^{q,\kappa} \lesssim U_{k+1}(s',\gamma').$$

Let us estimate each term \mathcal{T}_{ℓ} in (8.165) with exchanging n by k and (s, γ) by (s', γ') . For the first one we simply write in view of (8.166), since $s' + 1 + \gamma' \leq s + \gamma + n - k$,

$$\begin{aligned} \mathscr{T}_{1} &= \|\rho_{k}\|_{s_{0}+3}^{q,\kappa} \|\|a_{k}\|\|_{m,s'+1,\gamma'}^{q,\kappa} \leq F(s_{0}+3) \|\rho_{k}\|_{s_{0}+3}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \\ &+ F(s'+\gamma'+k+5) \|\rho_{k}\|_{s_{0}+3}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \\ &\leq F(s_{0}+3) \sum_{i=0}^{k} \|\rho_{i}\|_{s'+(k+1)+4+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \end{aligned}$$

+
$$F(s' + \gamma' + k + 5) \prod_{j=0}^{k} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa}$$
.

Hence, we obtain

$$\mathcal{T}_1 \leqslant U_{k+1}(s',\gamma').$$

For the second term we write according to (8.166), which can be applied since

$$s_0 + 1 + \gamma' \leqslant s' + 1 + \gamma' \leqslant s + \gamma + n - k,$$

combined with Sobolev embeddings

$$\mathcal{T}_{2} = \|\rho_{k}\|_{s'+3}^{q,\kappa'} \|\|a_{k}\|\|_{m,s_{0}+1,\gamma'}^{q,\kappa} \leq F(s_{0}+3)\|\rho_{k}\|_{s'+3+|m|}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s_{0}+k+5+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + F(s_{0}+\gamma'+k+5)\|\rho_{k}\|_{s'+3+|m|}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}.$$

Applying once again Sobolev embeddings and interpolation inequality we find

$$\begin{split} \|\rho_k\|_{s'+3+|m|}^{q,\kappa} \|\rho_i\|_{s_0+k+5+\gamma'+|m|}^{q,\kappa} &\leq \eta \|\rho_k\|_{s_0+3+|m|}^{q,\kappa} \|\rho_i\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} \\ &+ (1-\eta) \|\rho_i\|_{s_0+3+|m|}^{q,\kappa} \|\rho_k\|_{s'+k+5+\gamma'+|m|}^{q,\kappa}, \\ \eta &= \frac{s'-s_0}{s'-s_0+\gamma'+k+2}. \end{split}$$

Consequently, we deduce that

$$F(s_{0}+3) \|\rho_{k}\|_{s'+3+|m|}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s_{0}+k+5+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}$$

$$\leqslant F(s_{0}+3) \sum_{i=0}^{k} \|\rho_{i}\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}$$

$$\leqslant U_{k+1}(s',\gamma').$$
(8.167)

Now from the log-convexity assumption on F we obtain

$$F(s_0 + \gamma' + k + 5) \leqslant F^{\eta}(s_0 + 3)F^{1-\eta}(s' + \gamma' + k + 5).$$

Combined this with the interpolation inequality

$$\|\rho_k\|_{s'+3+|m|}^{q,\kappa} \le \left(\|\rho_k\|_{s_0+3+|m|}^{q,\kappa}\right)^{1-\eta} \left(\|\rho_k\|_{s'+k+5+\gamma'+|m|}^{q,\kappa}\right)^{\eta}$$

we find

$$F(s_0 + \gamma' + k + 5) \|\rho_k\|_{s'+3+|m|}^{q,\kappa} \leq \eta F(s_0 + 3) \|\rho_k\|_{s'+\gamma'+|m|+k+5}^{q,\kappa} + (1 - \eta)F(s' + \gamma' + k + 5) \|\rho_k\|_{s_0+3+|m|}^{q,\kappa}.$$
(8.168)

It follows that

$$F(s_{0} + \gamma' + k + 5) \|\rho_{k}\|_{s'+3+|m|}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \leq \eta F(s_{0} + 3) \|\rho_{k}\|_{s'+\gamma'+|m|+k+5}^{q,\kappa}$$

$$\times \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + (1-\eta)F(s'+\gamma'+k+5) \prod_{j=0}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}$$

$$\leq \eta F(s_{0} + 3) \sum_{i=0}^{k} \|\rho_{i}\|_{s'+\gamma'+k+5+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}$$

+
$$(1 - \eta)F(s' + \gamma' + k + 5) \prod_{j=0}^{k} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa}$$

Thus we find

$$F(s_0 + \gamma' + k + 5) \|\rho_k\|_{s'+3+|m|}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa} \le U_{k+1}(s',\gamma')$$
(8.169)

Therefore we obtain

$$\mathcal{T}_2 \leqslant 2 U_{k+1}(s', \gamma').$$

Let us now move to \mathcal{T}_3 , then the induction assumption (8.166) gives

$$\mathcal{T}_{3} \leqslant F(s_{0}+3) \sum_{i=0}^{k-1} \|\rho_{i}\|_{s'+\gamma'+k+5+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + F(s'+\gamma'+k+5) \prod_{j=0}^{k} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \leqslant U_{k+1}(s',\gamma').$$

As to \mathcal{T}_4 , we may write using the induction assumption (8.166)

$$\begin{aligned} \mathcal{T}_4 \leqslant F(s_0+3) \|\rho_k\|_{s'+3+|m|}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_i\|_{s_0+k+5+\gamma'+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa} \\ &+ F(s_0+\gamma'+k+5) \|\rho_k\|_{s'+3+|m|}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_j\|_{s_0+3+|m|}^{q,\kappa}. \end{aligned}$$

Applying (8.167) and (8.169) we infer

$$\mathcal{T}_4 \leqslant 2U_{k+1}(s',\gamma').$$

Concerning the term \mathcal{T}_5 in (8.165) we first split it as follows

$$\mathcal{T}_5 = \sum_{0 \leqslant \beta \leqslant \gamma'} \mathcal{T}_5^{1,\beta} + \mathcal{T}_5^{2,\beta}$$

with

$$\begin{aligned} \mathcal{T}_5^{1,\beta} &\triangleq \|\rho_k\|_{s_0+3+|m|+\gamma'-\beta}^{q,\kappa} \parallel a_k \parallel_{m,s',\beta}^{q,\kappa} \\ \mathcal{T}_5^{2,\beta} &\triangleq \|\rho_k\|_{s_0+3}^{q,\kappa} \parallel a_k \parallel_{m,s'+\gamma'-\beta,\beta}^{q,\kappa}. \end{aligned}$$

Let us start with estimating $\mathcal{T}_5^{1,\beta}$. Applying Sobolev embeddings, the monotonicity of F and (8.166)

$$\mathcal{T}_{5}^{1,\beta} \leqslant F(s_{0}+3) \|\rho_{k}\|_{s_{0}+3+|m|+\gamma'-\beta}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s'+k+5+|m|+\beta}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + F(s'+\beta+k+5) \|\rho_{k}\|_{s_{0}+3+|m|+\gamma'-\beta}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}.$$

Using the convexity inequality we infer

$$\begin{aligned} \|\rho_k\|_{s_0+3+|m|+\gamma'-\beta}^{q,\kappa} \|\rho_i\|_{s'+k+5+|m|+\beta}^{q,\kappa} &\lesssim \|\rho_k\|_{s_0+3+|m|}^{q,\kappa} \|\rho_i\|_{s'+k+5+|m|+\gamma'}^{q,\kappa} \\ &+ \|\rho_i\|_{s_0+3+|m|}^{q,\kappa} \|\rho_k\|_{s'+k+5+|m|+\gamma'}^{q,\kappa}. \end{aligned}$$

Similar argument using the log-convexity of F yields

$$F(s'+\beta+k+5) \|\rho_k\|_{H^{s_0+3+|m|+\gamma'-\beta}} \lesssim \|\rho_k\|_{s_0+3+|m|}^{q,\kappa} F(s'+\gamma'+k+5)$$
119

+ $\|\rho_k\|_{s'+\gamma'+k+5+|m|}^{q,\kappa} F(s_0+3).$

Consequently we get

$$\mathcal{T}_5^{1,\beta} \lesssim U_{k+1}(s',\gamma')$$

As to the term $\mathcal{T}_5^{2,\beta}$ we may write in view of (8.166)

$$\begin{aligned} \mathcal{T}_{5}^{2,\beta} \leqslant F(s_{0}+3) \|\rho_{k}\|_{s_{0}+3+|m|}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s'+\gamma'+k+5+|m|}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \\ &+ F(s'+\gamma'+k+5) \|\rho_{k}\|_{s_{0}+3+|m|}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} \\ &\leqslant U_{k+1}(s',\gamma'). \end{aligned}$$

Putting together the preceding estimates yields to

$$\mathcal{T}_5 \lesssim U_{k+1}(s',\gamma').$$

It remains to analyze the last term \mathcal{T}_6 in (8.165) which can be decomposed as follows

$$\mathcal{T}_6 = \sum_{0 \leqslant \beta \leqslant \gamma'} \mathcal{T}_6^{1,\beta} + \mathcal{T}_6^{2,\beta}$$

with

$$\mathcal{T}_{6}^{1,\beta} \triangleq \|\rho_{k}\|_{s'+3+|m|+\gamma'-\beta}^{q,\kappa} \|\|a_{k}\|_{m,s_{0},\beta}^{q,\kappa}$$
$$\mathcal{T}_{6}^{2,\beta} \triangleq \|\rho_{k}\|_{s'+3}^{q,\kappa} \|\|a_{k}\|_{m,s_{0}+\gamma'-\beta,\beta}^{q,\kappa}.$$

To estimate the first one we make appeal to (8.166)

$$\mathcal{T}_{6}^{1,\beta} \leqslant F(s_{0}+3) \|\rho_{k}\|_{s'+3+|m|+\gamma'-\beta}^{q,\kappa} \sum_{i=0}^{k-1} \|\rho_{i}\|_{s_{0}+k+5+|m|+\beta}^{q,\kappa} \prod_{\substack{j=0\\j\neq i}}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa} + F(s_{0}+k+5+\beta) \|\rho_{k}\|_{s'+3+|m|+\gamma'-\beta}^{q,\kappa} \prod_{j=0}^{k-1} \|\rho_{j}\|_{s_{0}+3+|m|}^{q,\kappa}.$$

Then using the convex inequalities

$$\begin{aligned} \|\rho_k\|_{s'+3+|m|+\gamma'-\beta}^{q,\kappa} \|\rho_i\|_{s_0+k+5+|m|+\beta}^{q,\kappa} &\lesssim \|\rho_k\|_{s_0+3+|m|}^{q,\kappa} \|\rho_i\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} \\ &+ \|\rho_k\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} \|\rho_i\|_{s_0+3+|m|}^{q,\kappa} \end{aligned}$$

and

$$\begin{aligned} \|\rho_k\|_{s+3+|m|+\gamma'-\beta}^{q,\kappa} F(s_0+k+5+\beta) &\lesssim \|\rho_k\|_{s_0+3+|m|}^{q,\kappa} F(s+k+5+\gamma') \\ &+ \|\rho_k\|_{s'+k+5+\gamma'+|m|}^{q,\kappa} F(s_0+3), \end{aligned}$$

one deduces that

$$\mathcal{T}_6^{1,\beta} \lesssim U_{k+1}(s',\gamma').$$

The estimate of the last $\mathcal{T}_6^{2,\beta}$ can be done in the same spirit. Finally, putting together the preceding estimates we get: $\forall s' \ge s_0, \gamma' \in \mathbb{N}$ with $s' + \gamma' \le s + \gamma + n - k - 1$,

$$|||a_{k+1}|||_{m,s',\gamma'}^{q,\kappa} \lesssim U_{k+1}(s',\gamma')$$

and this achieves the induction in (8.166). (iii) Since $m < -\frac{1}{2}$ then we can apply Lemma 7.2-(iii) allowing to get

$$\int_{\mathbb{T}} \left(\|K_n(\cdot, \cdot, \cdot + \eta)\|_s^{q,\kappa} \right)^2 d\eta \lesssim \left(\|\|a_n\|\|_{m,s,0}^{q,\kappa} \right)^2.$$

Then it suffices to apply Lemma 8.4-(ii) to get the desired result.

8.3.3. Iterated Töplitz matrix operators. In this section we shall be concerned with the establishment of tame estimates for the quantities $\|(\mathcal{M}_n^k \mathcal{Y}_n)(i)\|_{L^2}$ and $\|(\overline{\mathcal{M}}_n^k \overline{\mathcal{Y}}_n)(i)\|_{L^2}$ encountered before during the proof of Theorem 8.1. Before stating ours result let us recall that the Töplitz matrix operator \mathcal{M}_n was defined in (8.46) whose entries are given in (8.45). More precisely,

$$\mathcal{M}_{n} = \begin{pmatrix} 0 & 0 & \dots & \dots & 0 & 0 \\ m_{1,1} & 0 & \dots & \dots & 0 & 0 \\ m_{2,2} & m_{1,2} & 0 & \dots & \dots & 0 \\ \dots & \dots & \dots & \dots & 0 & 0 \\ m_{n,n} & m_{n-1,n} & \dots & \dots & m_{n,n} & 0 \end{pmatrix}, \quad \mathcal{Y}_{n} = \begin{pmatrix} K_{0} \\ \partial_{\chi} K_{0} \\ \dots \\ \dots \\ \partial_{\chi}^{n} K_{0} \\ \partial_{\chi}^{n} K_{0} \end{pmatrix},$$

with

$$m_{l,j} = \binom{n}{k} \left(\widehat{\mathbb{A}}_{(k)} - \mathbb{S}_{(k)}\right)$$
$$\triangleq \binom{n}{k} \mathcal{D}_k,$$

where

$$\widehat{\mathbb{A}}_{(k)} = \mathbb{A}_{\theta,(k)} + \mathbb{A}_{\eta,(k)}, \quad \mathbb{A}_{\theta,(k)} = \partial_{\theta} \left((\partial_{\theta}^{k} \rho) |\mathcal{D}_{\theta}|^{\alpha - 1} + |\mathcal{D}_{\theta}|^{\alpha - 1} (\partial_{\theta}^{k} \rho) \right)$$

and

$$\mathbb{S}_{(k)} = (\partial_{\eta}^{k+1}\rho)|\mathbf{D}_{\eta}|^{\alpha-1} + |\mathbf{D}_{\eta}|^{\alpha-1}(\partial_{\eta}^{k+1}\rho).$$

Denote by a_0 the symbol associated to K_0 defined through the relation,

$$K_0(\varphi,\theta,\eta) = \sum_{\xi \in \mathbb{Z}} a_0(\varphi,\theta,\xi) e^{i(\theta-\eta)\xi}$$

Concerning the matrices $\overline{\mathcal{M}}_n$, they are defined in (8.85) and the vectors $\overline{\mathcal{Y}}_n$ are given by (8.84) and (8.82). In what follows we shall state the main result of this section where the Töplitz structure of the aforementioned matrices plays a crucial role in getting tame estimates. To be more precise, we intend to establish the following lemma.

Lemma 8.5. Let $n \in \mathbb{N}$ and $m < -\frac{1}{2}$, then the following assertions hold true.

(i) Let $s_0 > \frac{d+1}{2}$ and assume the existence of an increasing log-convex function F_0 such that $\forall s \ge s_0, \gamma \in \mathbb{N}$ with $s + \gamma \le s_0 + n + 1 \Longrightarrow ||a_0||_{m,s,\gamma} \le F_0(s + \gamma).$

Then we have for any $k \in \{0, .., n\}$ and $i \in \{1, .., n+1\}$

$$\begin{aligned} \left\| \mathcal{M}_{n}^{k} \mathcal{Y}_{n}(i) \right\|_{L^{2}(\mathbb{T}^{d+2})} &\leq C \Big(F_{0} \big(s_{0} + 5 \big) + \|\rho\|_{H^{s_{0}+6+|m|}} \Big)^{k} \Big(F_{0} \big(s_{0} + 5 + i \big) + \|\rho\|_{H^{s_{0}+6+|m|+i}} \Big). \end{aligned}$$

$$We also get for k \in \{1, ..., n\},$$

$$\left\|\mathscr{M}_{n}^{k}\mathscr{Y}_{n}(i)\right\|_{L^{2}(\mathbb{T}^{d+2})} \leq C \|\rho\|_{H^{s_{0}+6+|m|}}^{k-1} \|\rho\|_{H^{s_{0}+6+|m|+i}} F_{0}(s_{0}+5+i).$$

and for $k = 0, i \in \{1, ..., n + 1\}$

$$\left\| \mathscr{Y}_{n}(i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \leq CF_{0}\left(s_{0}+5+i\right).$$

(ii) Let $q \in \mathbb{N}, s_0 > \frac{d+1}{2} + q$ and assume the existence of an increasing log-convex function F_0 such that

 $\forall s \ge s_0, \gamma \in \mathbb{N} \quad with \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow |||a_0|||_{m,s,\gamma}^{q,\kappa} \le F_0(s + \gamma).$ Then we have for any $k \in \{0, ..., n + q\}, i \in \{1, ..., n + 1\}$

$$\left\| \overline{\mathcal{M}}_{n}^{k} \overline{\mathcal{Y}}_{n}(i) \right\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant C \Big(F_{0} \big(s_{0} + 5 \big) + \|\rho\|_{s_{0}+6+|m|}^{q,\kappa} \Big)^{k} \Big(F_{0} \big(s_{0} + 5 + i \big) + \|\rho\|_{s_{0}+6+|m|+i}^{q,\kappa} \Big).$$

We also get

 $\left\|\overline{\mathcal{M}}_{n}^{k}\overline{\mathcal{Y}}_{n}(i)\right\|_{L^{2}(\mathbb{T}^{d+2})} \leq C\left(\|\rho\|_{s_{0}+6+|m|}^{q,\kappa}\right)^{k-1} \|\rho\|_{s_{0}+6+|m|+i}^{q,\kappa} F_{0}\left(s_{0}+5+i\right)$

and for k = 0,

$$\left\|\overline{\mathscr{Y}}_{n}(i)\right\|_{L^{2}(\mathbb{T}^{d+2})} \leqslant CF_{0}\left(s_{0}+5+i\right).$$

Here the bloc matrix operator $\overline{\mathcal{M}}_n$ was defined in (8.85) and $\overline{\mathcal{Y}}_n$ in (8.84) and (8.82).

Proof. (i) First remark that $\mathcal{M}_n \in \Delta_1$ and according to Lemma 8.1 we have $\mathcal{M}_n^k \in \Delta_k$ for any $k \in \{1, ..., n\}$ and $\mathcal{M}_n^{n+1} = 0$. In particular, we may write $\mathcal{M}_n^k = (a_{i,j}^k)_{1 \leq i,j \leq n+1}$ with

$$a_{i,j}^k = 0$$
, if $i - j < k$ or $i \leq k$.

In addition, according to Lemma 8.1 each coefficient $a_{i,j}^k$ is a linear combination of the iterative operator

$$\prod_{\ell=1}^{k} \mathcal{D}_{\alpha_{\ell}} \quad \text{such that} \quad \sum_{\ell=1}^{k} \alpha_{\ell} = i - j, \quad \alpha_{\ell} \ge 1.$$
(8.170)

Denote $Z = (Z_1, ..., Z_{n+1}) \triangleq \mathscr{M}_n^k \mathscr{Y}_n$, then

$$Z_i = \sum_{j=1}^{i-k} a_{i,j}^k \partial_\chi^j K_0.$$

Consequently Z_i is a linear combination of

$$\prod_{\ell=1}^{k} \mathcal{D}_{\alpha_{\ell}} \partial_{\chi}^{j} K_{0} \quad \text{such that} \quad \sum_{\ell=1}^{k} \alpha_{\ell} = i - j, \quad 1 \leq j \leq i - k, \quad \alpha_{\ell} \geq 1.$$
(8.171)

Using Lemma 8.4-(iii) with q = 0 and $m < -\frac{1}{2}$, we get for using Sobolev embeddings and change of variables

$$\begin{aligned} \left\| \prod_{\ell=1}^{k} \mathscr{D}_{\alpha_{\ell}} \partial_{\chi}^{j} K_{0} \right\|_{L^{2}_{\varphi,\theta,\eta}} &\lesssim \left(\int_{\mathbb{T}} \left\| \left(\prod_{\ell=1}^{k} \mathscr{D}_{\alpha_{\ell}} \partial_{\chi}^{j} K_{0} \right) (\cdot, \cdot, \cdot + \eta) \right\|_{H^{s_{0}}_{\varphi,\theta}}^{2} d\eta \right)^{\frac{1}{2}} \\ &\lesssim F(s_{0}+3) \sum_{\ell=1}^{k} \|\rho\|_{H^{s_{0}+k+4+|m|+\alpha_{\ell}}} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{H^{s_{0}+3+|m|+\alpha_{p}}} \\ &+ F(xs_{0}+k+4) \prod_{\ell=1}^{k} \|\rho\|_{H^{s_{0}+3+|m|+\alpha_{\ell}}}, \end{aligned}$$
(8.172)

where F is defined as an increasing log-convex function such that if $a_{j,0}$ is the symbol associated to the kernel $\partial_{\chi}^{j} K_{0}$ then

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k \Longrightarrow |||a_{j,0}|||_{m,s,\gamma} \leqslant F(s + \gamma).$$

Recall that a_0 is the symbol associated to K_0 and one gets easily the relation

$$\begin{aligned} \partial_{\chi}^{j} K_{0}(\varphi, \theta, \eta) &= \partial_{\chi}^{j} \sum_{\xi \in \mathbb{Z}} a_{0}(\varphi, \theta, \xi) e^{\mathrm{i}(\theta - \eta)} \\ &= \sum_{\xi \in \mathbb{Z}} \partial_{\theta}^{j} a_{0}(\varphi, \theta, \xi) e^{\mathrm{i}(\theta - \eta)}, \end{aligned}$$

which implies that

$$a_{j,0}(\varphi,\theta,\xi) = \partial^{j}_{\theta} a_{0}(\varphi,\theta,\xi).$$

Thus

$$|||a_{j,0}|||_{m,s,\gamma} \lesssim |||a_0|||_{m,s+j,\gamma}$$

Consequently, if F_0 is an increasing log-convex function such that

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k + j \Longrightarrow |||a_0|||_{m,s,\gamma} \le F_0(s + \gamma)$$

then we get

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k \Longrightarrow |||a_{j,0}|||_{m,s,\gamma} \leqslant F_0(s + \gamma + j) \tag{8.173}$$

and the choice $F(x) \triangleq F_0(x+j)$ gives an increasing and log-convex function. Notice that from (8.171) one has $k+j \leq i \leq n+1$ and therefore it suffices to impose to a_0 the assumption

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow |||a_0|||_{m,s,\gamma} \le F_0(s + \gamma)$$

Consequently, we get from the estimate (8.172)

$$\left\| \prod_{\ell=1}^{k} \mathfrak{D}_{\alpha_{\ell}} \partial_{\chi}^{j} K_{0} \right\|_{L^{2}_{\varphi,\theta,\eta}}^{2} \lesssim F_{0} \left(s_{0} + 3 + j \right) \sum_{\ell=1}^{k} \|\rho\|_{H^{s_{0}+k+4+|m|+\alpha_{\ell}}} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{H^{s_{0}+3+|m|+\alpha_{\ell}}} + F_{0} \left(s_{0} + k + 4 + j \right) \prod_{\ell=1}^{k} \|\rho\|_{H^{s_{0}+3+|m|+\alpha_{\ell}}},$$

$$(8.174)$$

Now we shall deal with the first term of the right hand side. Fix $\ell \in \{1, .., k\}$, since $\alpha_p \ge 1$, then interpolation inequality yields

$$\|\rho\|_{H^{s_0+3+|m|+\alpha_p}} \leqslant \|\rho\|_{H^{s_0+4+|m|}}^{1-\mu_p} \|\rho\|_{H^{s_0+6+|m|+i}}^{\mu_p}, \quad \mu_p = \frac{\alpha_p - 1}{i+2}.$$

Then

$$\begin{split} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{H^{s_{0}+3+|m|+\alpha_{p}}} &\leq \|\rho\|_{H^{s_{0}+4+|m|}}^{\Theta_{1,\ell}} \|\rho\|_{H^{s_{0}+6+|m|+i}}^{\Theta_{2,\ell}},\\ \Theta_{1,\ell} &= \sum_{\substack{p=1\\p\neq\ell}}^{k} (1-\mu_{p}), \quad \Theta_{2,\ell} = \sum_{\substack{p=1\\p\neq\ell}}^{k} \mu_{p}. \end{split}$$

Similarly

$$\|\rho\|_{H^{s_0+4+k+|m|+\alpha_\ell}} \leqslant \|\rho\|_{H^{s_0+4+|m|}}^{1-\mu_\ell} \|\rho\|_{H^{s_0+6+|m|+i}}^{\mu_\ell}, \quad \mu_\ell = \frac{\alpha_\ell + k}{i+2}$$

Hence

$$\begin{aligned} \|\rho\|_{H^{s_0+4+k+|m|+\alpha_{\ell}}} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{H^{s_0+3+|m|+\alpha_{p}}} &\leq \|\rho\|_{H^{s_0+4+|m|}}^{\Theta_{1,\ell}+1-\mu_{\ell}} \|\rho\|_{H^{s_0+6+|m|+i}}^{\Theta_{2,\ell}+\mu_{\ell}} \\ &\leq \|\rho\|_{H^{s_0+6+|m|}}^{\Theta_{1,\ell}+1-\mu_{\ell}} \|\rho\|_{H^{s_0+6+|m|+i}}^{\Theta_{2,\ell}+\mu_{\ell}}. \end{aligned}$$

$$(8.175)$$

Now we may check in view of (8.170)

$$\Theta_{2,\ell} + \mu_{\ell} = \frac{1}{i+2} \sum_{\substack{p=1\\p\neq\ell}}^{k} (\alpha_p - 1) + \frac{\alpha_{\ell} + k}{i+2}$$
$$= \frac{1}{i+2} \sum_{p=1}^{k} \alpha_p + \frac{1}{i+2}$$
$$= \frac{i-j+1}{i+2}.$$

Using the convexity inequality and the monotonicity of ${\cal F}_0$

$$F_0(s_0+3+j) \leqslant F_0^{1-\frac{j}{i+2}}(s_0+3)F_0^{\frac{j}{i+2}}(s_0+5+i)$$
$$\leqslant F_0^{1-\frac{j}{i+2}}(s_0+5)F_0^{\frac{j}{i+2}}(s_0+5+i).$$

Therefore we find

$$F_0(s_0+3+j)\sum_{\ell=1}^k \|\rho\|_{H^{s_0+k+4+|m|+\alpha_\ell}} \prod_{\substack{p=1\\p\neq\ell}}^k \|\rho\|_{H^{s_0+3+|m|+\alpha_p}} \lesssim G_0^{k+\frac{1}{i+2}}(0)G_0^{\frac{i+1}{i+2}}(i) \lesssim G_0^k(0)G_0(i),$$

with

$$G_0(i) = F_0(s_0 + 5 + i) + \|\rho\|_{H^{s_0 + 6 + |m| + i}}$$

where we have used in the last inequality the fact $G_0(0) \leq G_0(i)$. The last term in (8.174) can be treated in a similar way as before. Indeed,

$$\prod_{\ell=1}^{k} \|\rho\|_{H^{s_0+3+|m|+\alpha_{\ell}}} \leqslant \|\rho\|_{H^{s_0+4+|m|}}^{k-\frac{i-j-k}{i+2}} \|\rho\|_{H^{s_0+6+|m|+i}}^{\frac{i-j-k}{i+2}} \\ \leqslant \left(G_0(0)\right)^{k-\frac{i-j-k}{i+2}} \left(G_0(i)\right)^{\frac{i-j-k}{i+2}}.$$
(8.176)

and

$$F_0(s_0+4+j+k) \leqslant F_0^{\frac{1+i-k-j}{i+2}}(s_0+3)F_0^{\frac{1+k+j}{i+2}}(s_0+5+i)$$
$$\leqslant (G_0(0))^{\frac{1+i-k-j}{i+2}}(G_0(i))^{\frac{1+k+j}{i+2}}.$$

Consequently we find

$$F_0(s_0 + k + 4 + j) \prod_{\ell=1}^k \|\rho\|_{H^{s_0+3+|m|+\alpha_\ell}} \lesssim G_0^{k+\frac{1}{i+2}}(0) G_0^{\frac{i+1}{i+2}}(i) \\ \lesssim G_0^k(0) G_0(i).$$

This completes the proof of the first point of (i). As to the second estimate we simply use for $k \ge 1$ the inequality (8.175) combined with Sobolev embeddings and the fact $\Theta_{2,\ell} + \mu_{\ell} < 1$ in order to get into

$$\|\rho\|_{H^{s_0+4+k+|m|+\alpha_{\ell}}} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{H^{s_0+3+|m|+\alpha_p}} \leqslant \|\rho\|_{H^{s_0+6+|m|}}^{k-1} \|\rho\|_{H^{s_0+6+|m|+i}}.$$
(8.177)

Concerning the estimate (8.176) it is replaced in a similar way by

1.

$$\prod_{\ell=1}^{\kappa} \|\rho\|_{H^{s_0+3+|m|+\alpha_{\ell}}} \leqslant \|\rho\|_{H^{s_0+4+|m|}}^{k-1} \|\rho\|_{H^{s_0+6+|m|+i}}.$$
(8.178)

Combining both estimates yields to the desired estimate for $k \ge 1$. Concerning the case k = 0, it can derived from (8.173) and Lemma 8.4-(iii).

(ii) Recall from (8.88) that the matrix $\overline{\mathcal{M}}_n^{q+k} \in \Delta_k(n,q)$ and the set $\Delta_k(n,q)$ was defined before in Section 8.2. In particular $\overline{\mathcal{M}}_n^{k+q} = (M_{i,j}^k)_{1 \leq i,j \leq n+1}$ with

$$M_{i,j}^k = 0$$
, if $i - j < k$ or $i \leq k$.

In addition, according to Lemma 8.1 and (8.129) each matrix entry $M_{i,j}^k$ is a linear combination of the iterative matrix operator

$$\prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_n}(\alpha_\ell) \quad \text{such that} \quad \sum_{\ell=1}^{k+q} \alpha_\ell = i-j, \quad \alpha_\ell \ge 0.$$

From (8.129) one has $\mu_{\overline{\mathcal{M}}_n}(0) = \overline{\mathcal{M}}_q \in \Delta_1$ which is nilpotent with $\mu_{\overline{\mathcal{M}}_n}^{1+q} = 0$ and its main diagonal is vanishing. In addition $\mu_{\overline{\mathcal{M}}_n}(\alpha_\ell) \in \Delta_0$ and therefore we deduce from Lemma 8.1-(i)

$$\#\Big\{\ell, \alpha_\ell = 0\Big\} \geqslant q+1 \Longrightarrow \prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_n}(\alpha_\ell) = 0$$

Consequently $M_{i,j}^k$ is a linear combination of the iterative matrix operator

$$\prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_n}(\alpha_\ell) \quad \text{such that} \quad \sum_{\ell=1}^{k+q} \alpha_\ell = i - j \quad \text{and} \quad \#\left\{\ell, \alpha_\ell \ge 1\right\} \ge k \tag{8.179}$$

Set $\overline{Z} = (\overline{Z}_1, ..., \overline{Z}_2) \triangleq \overline{\mathcal{M}}_n^{k+q} \overline{\mathcal{Y}}_n$, then

$$\overline{Z}_i = \sum_{j=1}^{i-k} M_{i,j}^k \partial_{\chi}^j Y_q.$$

Consequently we get in view of (8.179) that \overline{Z}_i is a linear combination of

$$\prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_n}(\alpha_\ell) \partial_{\chi}^j Y_q \quad \text{such that} \quad \sum_{\ell=1}^{k+q} \alpha_\ell = i - j, \quad \# \Big\{ \ell, \alpha_\ell \ge 1 \Big\} \ge k \tag{8.180}$$

Applying Lemma 8.4-(iii) with slight adaptation we get for $m < -\frac{1}{2}$ and using Sobolev embeddings and change of variables

$$\left\| \prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_{n}}(\alpha_{\ell}) \partial_{\chi}^{j} Y_{q} \right\|_{L^{2}_{\varphi,\theta,\eta}}^{2} \lesssim \int_{\mathbb{T}} \left\| \left(\prod_{\ell=1}^{k+q} \mu_{\overline{\mathcal{M}}_{n}}(\alpha_{\ell}) \partial_{\chi}^{j} Y_{q} \right) (\cdot, \cdot, \cdot + \eta) \right\|_{H^{s_{0}}_{\varphi,\theta}}^{2} d\eta$$
$$\lesssim F(s_{0}+3|) \sum_{\ell=1}^{k} \|\rho\|_{q,s_{0}+k+4+|m|+\alpha_{\ell}}^{q,\kappa} \prod_{\substack{p=1\\p\neq\ell}}^{k} \|\rho\|_{q,s_{0}+3+|m|+\alpha_{p}}^{q,\kappa}$$
$$+ F(s_{0}+k+4) \prod_{\ell=1}^{k} \|\rho\|_{q,s_{0}+3+|m|+\alpha_{\ell}}^{q,\kappa}, \tag{8.181}$$

where F is defined as an increasing log-convex function such that if $\overline{a}_{j,0}$ is the matrix symbol of $\partial_{\chi}^{j} Y_{q}$, where Y_{q} is defined in (8.82), then

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k \Longrightarrow |||\overline{a}_{j,0}|||_{m,s,\gamma} \leqslant F(s + \gamma).$$

Let \overline{a}_0 be the symbol associated to the vector Y_q , then we have the relation

$$\overline{a}_{j,0} = \partial^j_\theta \overline{a}_0$$

Thus

$$\begin{aligned} \|\overline{a}_{j,0}\|\|_{m,s,\gamma} &\lesssim \||\overline{a}_0\|\|_{m,s+j,\gamma} \\ &\lesssim \||a_0\|\|_{m,s+j,\gamma}^{q,\kappa} \end{aligned}$$

Consequently, if F_0 is an increasing log-convex function such that

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k + j \Longrightarrow |||a_0|||_{m,s,\gamma}^{q,\kappa} \le F_0(s + \gamma)$$

then we get

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + k \Longrightarrow |||\overline{a}_{j,0}|||_{m,s,\gamma} \leqslant F_0(s + \gamma + j)$$

and the choice $F(x) = F_0(x+j)$ fits with the required assumptions. Remark that one gets in view of (8.179) that $k+j \leq i \leq n+1$ and therefore it suffices to impose to a_0 the assumption

$$\forall s \ge s_0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant s_0 + n + 1 \Longrightarrow ||a_0||_{m,s,\gamma}^{q,\kappa} \le F_0(s + \gamma)$$

At this stage, we proceed exactly as in the preceding point (i) using interpolation inequalities in (8.181) in order to get the desired result. The estimate of the second estimate of the point (ii) can be done in the same spirit as in (i) using in particular analogous estimates to (8.177) and (8.178). The proof is now complete.

9. Reducibility of the linearized operator

In this section we shall investigate the construction of a right approximate inverse for the linearized operator $\hat{\mathcal{L}}_{\omega}$ at a state i_0 close to the flat torus. This is the most delicate part in the construction of quasi-periodic solutions and needed during the implementation of Nash-Moser scheme . This operator was defined in (6.37) and it takes the form

$$\widehat{\mathscr{L}}_{\omega} = \widehat{\mathscr{L}}_{\omega}(i_0) = \Pi_{\mathbb{S}_0}^{\perp} \left(\omega \cdot \partial_{\varphi} - \partial_{\theta} K_{02}(\varphi) \right) \Pi_{\mathbb{S}_0}^{\perp}$$
(9.1)

Notice that we are dealing with an unbounded operator with variable coefficients and when it is evaluated at the flat torus, we simply find a diagonal operator that can be formally inverted provided that we avoid resonances in a suitable way by imposing suitable Diophantine conditions. The intuitive strategy that we shall follow and which is a common fact on the most studies around this topic is to conjugate the linearized operator \mathscr{L}_{ω} into a constant coefficients operator. This will be implemented in three long steps. First, we focus on the transport part and perform a quasi-periodic change of coordinates allowing to get after a suitable conjugation a new transport part with constant coefficients provided that the parameters α, ω are restricted in a Cantor set. The construction of this transformation is based on KAM reducibility procedure in the spirit of the recent works [16, 12]. Then the final outcome of the first step is a new operator whose transport part is diagonal with unbounded nonlocal perturbation of order α supplemented with a remainder of order zero. As to the the second step, it consists in reducing to a constant coefficient the fractional part of order α using a nonlocal hyperbolic flow constructed in Proposition 8.1, and of course without altering the transport part. The final product of the second step is linear operator whose positive part is a Fourier multiplier with a small non-diagonal bounded perturbation. The main concern of the last step is to reduce to a constant operator the remainder by applying the KAM scheme as in the paper [16]. We emphasize that during this step we need to strengthen the topology of linear continuous operators and work with a suitable topology on pseudo-differential operators described in (7.13). One of the delicate point that one should face at the end of step 2 is to check that the new remainder stays in this strong topology. This issue has been solved in Theorem 8.1 following a new approach based on the kernel dynamics and refined structures on Töplitz matrix operators. We stress that along the different reduction processes we need to impose some non-resonance assumptions that will be finally recast in terms of a Cantor like set.

Next, we shall introduce the parameters $\kappa, q, \tau_1, \tau_2$ and ρ connected with the geometric structure of the Cantor sets that will emerge during the reduction process. We shall also fix some regularity levels that will be made clear during the proofs.

$$\kappa \in (0,1), \quad (q,d) \in (\mathbb{N}^*)^2, \quad S \ge s \ge s_0 > \frac{d+1}{2} + q + 3, \quad \tau_2 > \tau_1 > d.$$
(9.2)

where S is a fixed large number and $\sigma = \sigma(d, \tau_1, \tau_2, q)$ is a positive number related to the loss of regularities during the different steps of the reduction. It depends only on the parameters governing the geometric structure of the involved Cantor sets and it is independent of the regularity. Notice that this implicit number may change during the computations from one line to another. Since the exact value of σ does not matter, then we find it convenient to keep this freedom more than making precise computations on the value of σ which is subject to continuous changes from the step 1 to step 3. We observe that sometimes with some statements we can do better and use weaker constraints than those of (9.2). We shall also need during KAM reduction and Nash-Moser scheme to make appeal to cut-offs in frequency introduced in (6.32) Notice that throughout this section, the constants $N_0 \ge 2$ and $\kappa \in (0, 1)$ are independent free parameters, but during the implementation of Nash-Moser scheme that will be explored later in Section 10, they will be suitably fixed with respect to the small parameter ε . Before starting the reduction process we shall discuss in the next two sections some algebraic and analytical properties of the linearized operator.

9.1. Structure on the normal direction. In this section we shall explore some analytical properties of the the operator $\hat{\mathcal{L}}_{\omega}$. We shall in particular give suitable estimates of its coefficients and view this operator as a small perturbation of order zero of the linearized operator computed in Proposition 3.1, which takes the form

$$\mathscr{L}_{r,\lambda}h = \omega \cdot \partial_{\varphi}h + \partial_{\theta} \big(V_{\alpha,r}(\varphi,\theta)h - \mathbb{K}_{r,\alpha}h \big)$$
(9.3)

where $\lambda = (\omega, \alpha)$ and

$$\mathbb{K}_{r,\alpha}h(\varphi,\theta) \triangleq \frac{C_{\alpha}}{2\pi} \int_{0}^{2\pi} \frac{h(\varphi,\eta)}{A_{r}^{\alpha/2}(\varphi,\theta,\eta)} d\eta, \qquad (9.4)$$

$$V_{r,\alpha}(\varphi,\theta) \triangleq \Omega + \frac{C_{\alpha}}{2\pi R(\varphi,\theta)} \int_0^{2\pi} \frac{\partial_{\eta} \left[R(\varphi,\eta) \sin(\eta-\theta) \right]}{A_r(\varphi,\theta,\eta)^{\alpha/2}} d\eta, \tag{9.5}$$

$$A_r(\varphi,\theta,\eta) \triangleq \left(R(\varphi,\eta) - R(\varphi,\theta) \right)^2 + 4R(\varphi,\eta)R(\varphi,\theta)\sin^2\left(\frac{\eta-\theta}{2}\right), \tag{9.6}$$
$$R(\varphi,\theta) \triangleq \left(1 + 2r(\varphi,\theta)\right)^{\frac{1}{2}}.$$

Proposition 9.1. The operator $\widehat{\mathcal{L}}_{\omega}$ defined in (9.1) takes the form

$$\widehat{\mathscr{L}}_{\omega} = \Pi_{\mathbb{S}_{0}}^{\perp} \left(\mathscr{L}_{\varepsilon r_{0},\lambda} - \varepsilon \partial_{\theta} \mathscr{R} \right) \Pi_{\mathbb{S}_{0}}^{\perp}$$

where the operator $\mathscr{L}_{\varepsilon r_0,\lambda}$ is defined in (9.3), with

$$r_{0}(\varphi, \cdot) \triangleq \mathbf{A}(i_{0}(\varphi)) = \mathbf{A}(\vartheta_{0}(\varphi), I_{0}(\varphi), z_{0}(\varphi))$$
$$= v(\vartheta_{0}(\varphi), I_{0}(\varphi)) + z_{0}(\varphi),$$

supplemented with the reversibility assumption

$$r_0(-\varphi,-\theta) = r_0(\varphi,\theta)$$

and \mathcal{R} is an integral operator as in (7.12) with a kernel K_0 satisfying the property

$$K_0(\lambda, -\varphi, -\theta, -\eta) = K_0(\lambda, \varphi, \theta, \eta).$$

Moreover, under the assumption

$$\|\mathfrak{I}_0\|_{q,s_0}^{\gamma,\kappa} \lesssim 1,\tag{9.7}$$

we have for all $s \ge s_0$,

(i) The function r_0 satisfies the estimates,

$$\|r_0\|_s^{q,\kappa} \lesssim 1 + \|\mathfrak{I}_0\|_s^{q,\kappa} \quad and \quad \|\Delta_{12}r_0\|_s^{q,\kappa} \lesssim \|\Delta_{12}i\|_s^{q,\kappa} + \|\Delta_{12}i\|_{s_0}^{q,\kappa} \max_{\ell=1,2} \|\mathfrak{I}_\ell\|_s^{q,\kappa}.$$

(ii) For all $j \in \mathbb{N}$ the kernel K_0 satisfies the estimates,

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{j} K_{0})(*, \cdot, \cdot, \cdot + \eta) \|_{s}^{q, \kappa} \lesssim 1 + \| \mathfrak{I}_{0} \|_{s+3+j}^{q, \kappa}$$

and

$$\sup_{\eta \in \mathbb{T}} \|\Delta_{12}(\partial_{\theta}^{j} K_{0})(*, \cdot, \cdot, \cdot, \cdot + \eta)\|_{s}^{q, \kappa} \lesssim \|\Delta_{12}i\|_{q, s+3+j} + \|\Delta_{12}i\|_{s_{0}+3}^{q, \kappa} \max_{\ell=1, 2} \|\mathfrak{I}_{\ell}\|_{s+3+j}^{q, \kappa},$$

where $*, \cdot, \bullet$, denote successively the variables λ, φ, θ and $\mathfrak{I}_{\ell}(\varphi) = i_{\ell}(\varphi) - (\varphi, 0, 0)$. In addition, for any function $f, \Delta_{12}f := f(i_1) - f(i_2)$ refers for the difference of f taken at two different states i_1 and i_2 satisfying (9.7).

Proof. To alleviate the notation we shall at several stages of the proof remove the dependance of our functions/operators with respect to (ω, α) and keep it when it is relevant. Recall that the operator \mathscr{L}_{ω} is defined in (9.1). Then we need to describe $K_{02}(\varphi)$. According to Proposition 6.1-(i) and the identity (5.20) we may write

$$K_{02}(\varphi) = \mathcal{L}(\lambda)|_{\mathbb{H}_{\mathbb{S}_0}^{\perp}} + \varepsilon \partial_z \nabla_z \mathscr{P}_{\varepsilon}(i_0(\varphi)) + \varepsilon \mathscr{R}(\varphi), \qquad (9.8)$$

where

$$\begin{split} \mathscr{R}(\varphi) &\triangleq \mathscr{R}_{1}(\varphi) + \mathscr{R}_{2}(\varphi) + \mathscr{R}_{3}(\varphi), \quad \text{with} \quad \mathscr{R}_{1}(\varphi) = L_{2}^{\top}(\varphi)\partial_{I}\nabla_{I}\mathscr{P}_{\varepsilon}(i_{0}(\varphi))L_{2}(\varphi), \\ \mathscr{R}_{2}(\varphi) &= L_{2}^{\top}((\varphi)\partial_{z}\nabla_{I}\mathscr{P}_{\varepsilon}(i_{0}(\varphi)), \\ \mathscr{R}_{3}(\varphi) &= \partial_{I}\nabla_{z}\mathscr{P}_{\varepsilon}(i_{0}(\varphi))L_{2}(\varphi). \end{split}$$

As we shall see all the operators $\mathscr{R}_1(\varphi)$, $\mathscr{R}_2(\varphi)$ and $\mathscr{R}_3(\varphi)$ have finite-dimensional rank. This property is obvious for the operator $L_2(\varphi)$ which send $\mathbb{H}_{\mathbb{S}_0}^{\perp}$ to \mathbb{R}^d and therefore for any $\rho \in \mathbb{H}_{\mathbb{S}_0}^{\perp}$

$$L_2(\varphi)[\rho] = \sum_{k=1}^d \left\langle L_2(\varphi)[\rho], \underline{e}_k \right\rangle_{L^2(\mathbb{T})} \underline{e}_k = \sum_{k=1}^d \left\langle \rho, L_2^{\mathsf{T}}(\varphi)[\underline{e}_k] \right\rangle_{L^2(\mathbb{T})} \underline{e}_k,$$

with $(\underline{e}_k)_{k=1}^d$ being the canonical basis of \mathbb{R}^d . Hence

$$\mathcal{R}_{1}(\varphi)[\rho] = \sum_{\substack{k=1 \\ d}}^{d} \left\langle \rho, L_{2}^{\top}(\varphi)[\underline{e}_{k}] \right\rangle_{L^{2}(\mathbb{T})} \mathbf{A}_{1}(\varphi)[\underline{e}_{k}] \quad \text{with} \quad \mathbf{A}_{1}(\varphi) = L_{2}^{\top}(\varphi)\partial_{I}\nabla_{I}\mathcal{P}_{\varepsilon}(i_{0}(\varphi)),$$
$$\mathcal{R}_{3}(\varphi)[\rho] = \sum_{\substack{k=1 \\ k=1}}^{d} \left\langle \rho, L_{2}^{\top}(\varphi)[\underline{e}_{k}] \right\rangle_{L^{2}(\mathbb{T})} \mathbf{A}_{3}(\varphi)[\underline{e}_{k}] \quad \text{with} \quad \mathbf{A}_{3}(\varphi) = \partial_{I}\nabla_{z}\mathcal{P}_{\varepsilon}(i_{0}(\varphi)).$$

In the same spirit, since $\mathbb{A}_2(\varphi) = \partial_z \nabla_I \mathscr{P}_{\varepsilon}(i_0(\varphi)) : \mathbb{H}_{\mathbb{S}_0}^{\perp} \to \mathbb{R}^d$, then we can write

$$\mathcal{R}_{2}(\varphi)[\rho] = \sum_{k=1}^{d} \left\langle \rho, \mathbf{A}_{2}^{\top}(\varphi)[\underline{e}_{k}] \right\rangle_{L^{2}(\mathbb{T})} L_{2}^{\top}(\varphi)[\underline{e}_{k}].$$

By setting

$$g_{k,1}(\varphi,\theta) = g_{k,3}(\varphi,\theta) = \chi_{k,2}(\varphi,\theta) \triangleq L_2^{\top}(\varphi)[\underline{e}_k](\theta), \quad g_{k,2}(\varphi,\theta) \triangleq \mathbf{A}_2^{\top}(\varphi)[\underline{e}_k](\theta)$$

and

$$\chi_{k,1}(\varphi,\theta) \triangleq \mathbf{A}_1(\varphi)[\underline{e}_k](\theta), \quad \chi_{k,3}(\varphi,\theta) \triangleq \mathbf{A}_3(\varphi)[\underline{e}_k](\theta),$$

one can write the operator $\mathcal R$ in the integral form

$$\begin{aligned} \mathscr{R}\rho(\varphi,\theta) &= \sum_{k'=1}^{3} \sum_{k=1}^{d} \langle \rho(\varphi,\cdot), g_{k,k'}(\varphi,\cdot) \rangle_{L^{2}(\mathbb{T})} \chi_{k,k'}(\varphi,\theta) \\ &= \int_{0}^{2\pi} \rho(\varphi,\eta) K_{0}(\varphi,\theta,\eta) d\eta \end{aligned}$$

with

$$K_0(\varphi,\theta,\eta) \triangleq \sum_{k'=1}^3 \sum_{k=1}^d g_{k,k'}(\varphi,\eta)\chi_{k,k'}(\varphi,\theta).$$

Now we remark that $g_{k,k'}, \chi_{k,k'} \in \mathbb{H}_{\mathbb{S}_0}^{\perp}$ and satisfy the estimates

$$\|g_{k,k'}\|_{s}^{q,\kappa} + \|\chi_{k,k'}\|_{s}^{q,\kappa} \lesssim 1 + \|\mathfrak{I}_{0}\|_{s+3}^{q,\kappa}$$
(9.9)

and

$$\|d_{i}g_{k,k'}[\hat{i}]\|_{s}^{q,\kappa} + \|d_{i}\chi_{k,k'}[\hat{i}]\|_{s}^{q,\kappa} \lesssim \|\hat{i}\|_{s+2}^{q,\kappa} + \|\mathfrak{I}_{0}\|_{s+4}^{q,\kappa}\|\hat{i}\|_{s_{0}+2}^{q,\kappa}.$$
(9.10)

By (9.9), combined with the law products of Lemma 4.1 we deduce that

$$\sup_{\eta \in \mathbb{T}} \|(\partial_{\theta}^{j} K_{0})(\lambda, \omega, \varphi, \theta, \eta + \theta)\|_{s}^{q,\kappa} \lesssim 1 + \|\mathfrak{I}_{0}\|_{s+3+j}^{q,\kappa}$$

Applying the mean value theorem with (9.10), yields

$$\sup_{\eta \in \mathbb{T}} \|\Delta_{12}(\partial_{\theta}^{j} K_{0})(\lambda, \omega, \varphi, \theta, \eta + \theta)\|_{s}^{q, \kappa} \lesssim \|\Delta_{12}i\|_{s+3+j}^{q, \kappa} + \|\Delta_{12}i\|_{s_{0}+3}^{q, \kappa} \max_{\ell=1, 2} \|\mathfrak{I}_{\ell}\|_{s+3+j}^{q, \kappa}$$

This concludes the proof of (ii). The symmetry property of K_0 is a consequence of the definition of r and the reversibility condition (5.24) imposed on the torus i_0 . Then, using (9.8), (5.18) and (5.3) we get

$$\begin{split} K_{02}(\varphi) &= \mathcal{L}(\lambda)|_{\mathbb{H}_{\mathbb{S}_{0}}^{\perp}} + \varepsilon \partial_{z} \nabla_{z} \mathscr{P}_{\varepsilon}(i_{0}(\varphi)) + \varepsilon \mathscr{R}(\varphi) \\ &= \mathcal{L}(\lambda)|_{\mathbb{H}_{\mathbb{S}_{0}}^{\perp}} + \varepsilon \Pi_{\mathbb{S}_{0}}^{\perp} \partial_{r} \nabla_{r} P_{\varepsilon}(\mathbf{A}(i_{0}(\varphi))) + \varepsilon \mathscr{R}(\varphi) \\ &= \Pi_{\mathbb{S}_{0}}^{\perp} \partial_{r} \nabla_{r} \mathscr{H}_{\varepsilon}(\mathbf{A}(i_{0}(\varphi))) + \varepsilon \mathscr{R}(\varphi) \\ &= \Pi_{\mathbb{S}_{0}}^{\perp} \partial_{r} \nabla_{r} H(\varepsilon r_{0}(\varphi, \cdot)) + \varepsilon \mathscr{R}(\varphi) \end{split}$$

where we have used the notation $r_0(\varphi, \cdot) \triangleq \mathbf{A}(i_0(\varphi))$. According to the general form of the linearized operator stated in Proposition 3.1 one has

$$-\partial_{\theta}\partial_{r}\nabla_{r}H(\varepsilon r_{0}(\varphi,\cdot))=\partial_{\theta}\left(V_{\varepsilon r_{0},\alpha}\cdot\right)-\partial_{\theta}\mathbb{K}_{\varepsilon r_{0},\alpha}$$

which implies

$$-\partial_{\theta}K_{02}(\varphi) = \Pi_{\mathbb{S}_{0}}^{\perp} \left(\partial_{\theta} \left(V_{\varepsilon r_{0},\alpha} \cdot \right) - \partial_{\theta}\mathbb{K}_{\varepsilon r_{0},\alpha} \right) - \varepsilon \partial_{\theta}\mathscr{R}(\varphi) \right) \Pi_{\mathbb{S}_{0}}^{\perp}.$$

Plugging this identity into (9.1) gives the desired result. Then by (5.12) and (5.45) we get

$$\begin{aligned} \|r_0\|_s^{q,\kappa} &\lesssim \|v(\vartheta, I_0)\|_s^{q,\kappa} + \|z\|_s^{q,\kappa} \\ &\lesssim 1 + \|\mathfrak{I}_0\|_s^{q,\kappa}. \end{aligned}$$

The bound on $\Delta_{12}r_0$ can be done in a similar way using in particular the mean value theorem. Indeed, one may write

$$\|\Delta_{12}r_0\|_s^{q,\kappa} \lesssim \|\Delta_{12}v(\vartheta, I_0)\|_s^{q,\kappa} + \|\Delta_{12}z\|_s^{q,\kappa}.$$

Applying Taylor formula with (5.45) combined with the law products allow to get

$$\begin{aligned} \|\Delta_{12}v(\vartheta, I_0)\|_s^{q,\kappa} &\lesssim \|\Delta_{12}(\vartheta, I_0)\|_s^{q,\kappa} + \|\Delta_{12}(\vartheta, I_0)\|_{s_0}^{q,\kappa} \max_{\ell=1,2} \|\mathfrak{I}_\ell\|_s^{q,\kappa} \\ &\lesssim \|\Delta_{12}i\|_s^{q,\kappa} + \|\Delta_{12}i\|_{s_0}^{q,\kappa} \max_{\ell=1,2} \|\mathfrak{I}_\ell\|_s^{q,\kappa}, \end{aligned}$$

which implies in turn

$$\|\Delta_{12}r_0\|_s^{q,\kappa} \lesssim \|\Delta_{12}i\|_s^{q,\kappa} + \|\Delta_{12}i\|_{s_0}^{q,\kappa} \max_{\ell=1,2} \|\mathfrak{I}_\ell\|_s^{q,\kappa}.$$

This completes the proof of Proposition 9.1.

9.2. Asymptotic structure. We intend here to provide a suitable decomposition for the linearized operator described by (9.3) for a small state r and estimate its coefficients. For the function spaces tools used below we refer to Section 4 and Section 7.2. However for the reversibility concepts we may consult Definition 7.1. Our main result of this section reads as follows.

Lemma 9.1. Let $q, \kappa \in \mathbb{N}, s_0 > \frac{d+1}{2} + q$, there exists $\varepsilon_0 > 0$ such that for $r \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^{\infty}_{\text{even}}(\mathbb{T}^{d+1}))$ satisfying

$$\|r\|_{s_0}^{q,\kappa} \leqslant \varepsilon_0,\tag{9.11}$$

the operator $\mathscr{L}_{r,\lambda}$ in (9.3) is reversible and takes the form

$$\mathscr{L}_{r,\lambda} = \omega \cdot \partial_{\varphi} + \partial_{\theta} \Big[V_{r,\alpha} - \left(\mathscr{W}_{r,\alpha} \left| \mathbf{D} \right|^{\alpha - 1} + \left| \mathbf{D} \right|^{\alpha - 1} \mathscr{W}_{r,\alpha} \right) + \mathscr{R}_{r,\alpha} \Big]$$

with the following properties.

(i) The functions $\mathscr{W}_{r,\alpha}, V_{r,\alpha} \in W^{q,\infty}_{\kappa}(\mathscr{O}, H^{\infty}_{\text{even}}(\mathbb{T}^{d+1}))$ such that for any $s \ge s_0$

$$\|\mathscr{W}_{r,\alpha} - C_{\alpha} 2^{-\alpha}\|_{s+3}^{q,\kappa} + \|\partial_{\theta} \mathscr{W}_{r,\alpha}\|_{s+2}^{q,\kappa} + \|V_{r,\alpha} - V_{0,\alpha}\|_{s}^{q,\kappa} \lesssim \|r\|_{s+4}^{q,\kappa}$$

where $V_{0,\alpha}$ is defined in (3.11).

(ii) The linear operator $\partial_{\theta}\mathscr{R}_{r,\alpha} : W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{even}}(\mathbb{T}^{d+1})) \to W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{odd}}(\mathbb{T}^{d+1}))$ is continuous with

$$\||\partial_{\theta}\mathscr{R}_{r,\alpha}|||_{-1,s,\gamma}^{q,\kappa} \lesssim \|r\|_{s+5+\gamma}^{q,\kappa}.$$

(iii) Let r_1, r_2 be smooth and satisfying (9.11). Denote by $\Delta_{12}V_{r,\alpha} = V_{r_1,\alpha} - V_{r_2,\alpha}$ and $\Delta_{12}r = r_1 - r_2$. Then for any $s \ge s_0$, we have

$$\|\Delta_{12}V_{r,\alpha}\|_{s}^{q,\kappa} \lesssim \|\Delta_{12}r\|_{s+4}^{q,\kappa} + \|\Delta_{12}r\|_{s_{0}+4}^{q,\kappa} \max_{\ell=1,2} \|r_{\ell}\|_{s+4}^{q,\kappa}$$

Proof. (i) We shall first establish the operator decomposition and move later to the estimates of the coefficients. For this aim, we first use (9.6) in order to get

$$A_r(\varphi,\theta,\eta) = \sin^2\left(\frac{\eta-\theta}{2}\right) \left[\left(\frac{R(\varphi,\eta) - R(\varphi,\theta)}{\sin\left(\frac{\eta-\theta}{2}\right)}\right)^2 + 4R(\varphi,\eta)R(\varphi,\theta) \right].$$

Introduce the function

$$f(\varphi,\theta,\eta) = C_{\alpha} \left[\left(\frac{R(\varphi,\eta) - R(\varphi,\theta)}{\sin\left(\frac{\eta-\theta}{2}\right)} \right)^2 + 4R(\varphi,\eta)R(\varphi,\theta) \right]^{-\frac{\alpha}{2}}.$$

Then it is clear that f is symmetric, that is, $f(\varphi, \theta, \eta) = f(\varphi, \eta, \theta)$ and therefore by using Lemma 4.6-(ii) we deduce that

$$f(\varphi,\theta,\eta) = \mathscr{W}_{r,\alpha}(\varphi,\theta) + \mathscr{W}_{r,\alpha}(\varphi,\eta) + \sin^2\left(\frac{\eta-\theta}{2}\right)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta)$$
(9.12)

with

$$\mathscr{W}_{r,\alpha}(\varphi,\theta) = C_{\alpha} 2^{-\alpha-1} \left(R^2(\varphi,\theta) + \left(\partial_{\theta} R\right)^2(\varphi,\theta) \right)^{-\frac{\alpha}{2}},\tag{9.13}$$

where $\mathscr{A}_{r,\alpha}$ is symmetric and satisfies the estimate

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{A}_{r,\alpha}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \leq C \|\Delta_{\theta,\eta} f\|_{s+\frac{1}{2}+\epsilon}^{q,\kappa}$$
$$\leq C \|f - 2^{-\alpha} C_{\alpha}\|_{s+\frac{5}{2}+\epsilon}^{q,\kappa}.$$
(9.14)

Notice that the notation \cdot is for φ and \cdot stands for θ . From the translation invariance of Lebesgue measure we may write

$$\|f - 2^{-\alpha}C_{\alpha}\|_{s+\frac{5}{2}+\epsilon}^{q,\kappa} = \|\tilde{f} - 2^{-\alpha}C_{\alpha}\|_{s+\frac{5}{2}+\epsilon}^{q,\kappa}$$

with

$$\tilde{f}(\varphi,\theta,\eta) = f(\varphi,\theta,\theta+\eta)$$
$$= C_{\alpha} \left[\left(\frac{R(\varphi,\theta+\eta) - R(\varphi,\theta)}{\sin\left(\frac{\eta}{2}\right)} \right)^2 + 4R(\varphi,\theta+\eta)R(\varphi,\theta) \right]^{-\frac{\alpha}{2}}$$

Applying Lemmata 4.4-4.1-4.7 combined with the smallness condition (9.11), we obtain

$$\|\tilde{f} - 2^{-\alpha}C_{\alpha}\|_{s+\frac{5}{2}+\epsilon}^{q,\kappa} \leqslant C \|r\|_{s+\frac{7}{2}+\epsilon}^{q,\kappa}.$$
(9.15)

Notice that in order to apply Lemma 4.7 it is enough to write

$$\left(\frac{R(\varphi,\theta+\eta)-R(\varphi,\theta)}{\sin\left(\frac{\eta}{2}\right)}\right)^2 = \left(\frac{R(\varphi,\theta+\eta)-R(\varphi,\theta)}{\tan\left(\frac{\eta}{2}\right)}\right)^2 + \left(R(\varphi,\theta+\eta)-R(\varphi,\theta)\right)^2.$$

It follows from (9.14) and (9.15) that

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{A}_{r,\alpha}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \leqslant C \|r\|_{s+\frac{7}{2}+\epsilon}^{q,\kappa}.$$
(9.16)

Similarly, using Lemma 4.6-(ii) we get for any $p, m \in \mathbb{N}$

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p} \partial_{\eta}^{m} \mathscr{A}_{r,\alpha})(\cdot, \cdot, \cdot + \eta) \|_{s}^{q,\kappa} \leqslant C \| r \|_{s+p+m+\frac{7}{2}+\epsilon}^{q,\kappa}.$$
(9.17)

Coming back to (9.12), then we have the following decomposition

$$C_{\alpha}A_{r}^{-\frac{\alpha}{2}}(\varphi,\theta,\eta) = \frac{\mathscr{W}_{r,\alpha}(\varphi,\theta) + \mathscr{W}_{r,\alpha}(\varphi,\eta)}{|\sin(\frac{\theta-\eta}{2})|^{\alpha}} + \frac{\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta)}{|\sin(\frac{\eta-\theta}{2})|^{\alpha-2}}.$$
(9.18)

Plugging this identity into (9.5) and using a change of variables we find

$$V_{r,\alpha}(\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathscr{A}_{r,\alpha}^1(\varphi,\theta,\eta)}{|\sin(\frac{\eta}{2})|^{\alpha}} d\eta$$
(9.19)

with

$$\mathscr{A}_{r,\alpha}^{1}(\varphi,\theta,\eta) = \frac{\partial_{\eta} \left[R(\varphi,\eta+\theta)\sin(\eta) \right]}{R(\varphi,\theta)} \left(\mathscr{W}_{r,\alpha}(\varphi,\theta) + \mathscr{W}_{r,\alpha}(\varphi,\eta+\theta) \right) + \frac{\partial_{\eta} \left[R(\varphi,\eta+\theta)\sin(\eta) \right]}{R(\varphi,\theta)} \sin^{2}(\eta/2) \mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta).$$
(9.20)

Similarly inserting (9.18) into (9.4) yields

$$\mathbb{K}_{r,\alpha}h(\varphi,\theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{h(\varphi,\eta) \left(\mathscr{W}_{r,\alpha}(\varphi,\theta) + \mathscr{W}_{r,\alpha}(\varphi,\eta)\right)}{|\sin(\frac{\eta-\theta}{2})|^{\alpha}} d\eta + \frac{1}{2\pi} \int_0^{2\pi} \frac{h(\varphi,\eta)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta)}{|\sin(\frac{\eta-\theta}{2})|^{\alpha-2}} d\eta.$$

Using the definition of the modified fractional Laplacian in (4.19) we infer

$$\mathbb{K}_{r,\alpha}h(\varphi,\theta) = \mathscr{W}_{r,\alpha}(\varphi,\theta)|\mathbf{D}|^{\alpha-1}h(\varphi,\theta) + |\mathbf{D}|^{\alpha-1}(\mathscr{W}_{r,\alpha}h)(\varphi,\theta) + \mathscr{R}_{r,\alpha}h(\varphi,\theta)$$

with

$$\mathscr{R}_{r,\alpha}h(\varphi,\theta) \triangleq \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta)}{|\sin(\frac{\eta-\theta}{2})|^{\alpha-2}} h(\varphi,\eta) d\eta.$$
(9.21)

Direct computations yield

$$\partial_{\theta}\mathscr{R}_{r,\alpha}h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\theta-\eta}{2})}{|\sin(\frac{\eta-\theta}{2})|^{\alpha}} \mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta) h(\varphi,\eta) d\eta$$
(9.22)

with

$$\mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta) = \partial_{\theta}\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta) \sin\left(\frac{\theta-\eta}{2}\right) + 1 - \frac{\alpha}{2}\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta) \cos\left(\frac{\theta-\eta}{2}\right).$$
(9.23)

Inserting this into (9.3) we find

$$\mathscr{L}_{r,\lambda} = \omega \cdot \partial_{\varphi} + \partial_{\theta} \Big[V_{r,\alpha} - \left(\mathscr{W}_{r,\alpha} \left| \mathbf{D} \right|^{\alpha - 1} + \left| \mathbf{D} \right|^{\alpha - 1} \mathscr{W}_{r,\alpha} \right) + \mathscr{R}_{r,\alpha} \Big]$$
(9.24)

where $V_{r,\alpha}$ is given by (9.19) and $\mathscr{R}_{r,\alpha}$ by (9.21). The next goal is to estimate the coefficients $\mathscr{W}_{r,\alpha}$ and $V_{r,\alpha}$. Concerning the first one, defined in (9.13) one has

$$\mathscr{W}_{r,\alpha}(\varphi,\theta) = C_{\alpha} 2^{-\alpha-1} \left(1 + 2r(\varphi,\theta) + \frac{(\partial_{\theta}r)^2(\varphi,\theta)}{1 + 2r(\varphi,\theta)} \right)^{-\frac{\alpha}{2}}$$

Then from Lemma 4.4 combined with the smallness condition (9.11) we obtain

$$\|\mathscr{W}_{r,\alpha} - C_{\alpha} 2^{-\alpha - 1}\|_{s-1}^{q,\kappa} \leqslant C \|r\|_{s}^{q,\kappa}.$$
(9.25)

Similarly, computing $\partial_{\theta} \mathscr{W}_{r,\alpha}$ and using the law products together with Lemma 4.4 we find

$$\|\partial_{\theta} \mathscr{W}_{r,\alpha}\|_{s-2}^{q,\kappa} \leqslant C \|r\|_{s}^{q,\kappa}$$
131

Let us now move to the estimate of $V_{r,\alpha}$ defined by (9.19) then we find from straightforward computations

$$\begin{aligned} \|V_{r,\alpha} - V_{0,\alpha}\|_{s}^{q,\kappa} &\lesssim \int_{0}^{2\pi} \frac{\|\mathscr{A}_{r,\alpha}^{1} - \mathscr{A}_{0,\alpha}^{1}\|_{s}^{q,\kappa}}{|\sin(\frac{\eta}{2})|^{\frac{1}{2}}} \left(1 + |\ln(\sin(\eta/2))|^{q}\right) d\eta \\ &\lesssim \|\mathscr{A}_{r,\alpha}^{1} - \mathscr{A}_{0,\alpha}^{1}\|_{s}^{q,\kappa}. \end{aligned}$$

Notice that $V_{0,\alpha} = V_0(\alpha)$ which is given by (3.11). Applying Lemma 4.1 to (9.20) combined with (9.16), (9.25) and (9.11) yield

$$\|V_{r,\alpha} - V_0(\alpha)\|_s^{q,\kappa} \le C \|r\|_{s+\frac{7}{2}+\epsilon}^{q,\kappa}.$$
(9.26)

To get the suitable estimate it suffices to use Sobolev embeddings. (ii) Coming back to (9.22) and making the change of variables $\eta \rightarrow \eta + \theta$ we find

$$\partial_{\theta}\mathscr{R}_{r,\alpha}h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\eta}{2})}{|\sin(\eta/2)|^{\alpha}} \mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta+\theta) h(\varphi,\eta+\theta) d\eta,$$

with

$$\mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta+\theta) = -\partial_{\theta}\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta)\,\sin\left(\eta/2\right) + \left(1-\frac{\alpha}{2}\right)\mathscr{A}_{r,\alpha}(\varphi,\theta,\eta+\theta)\,\cos\left(\eta/2\right).\tag{9.27}$$

According to (7.8), the symbol of $\partial_{\theta} \mathscr{R}_{r,\alpha}$ can be recovered from the kernel as follows

$$\sigma_{\partial_{\theta}\mathscr{R}_{r,\alpha}}(\varphi,\theta,\xi) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\eta}{2})}{|\sin(\eta/2)|^{\alpha}} \mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta+\theta) e^{i\eta\xi} d\eta$$

Then integration by parts combined with (7.6) yield

$$\xi^{1+\gamma} \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathscr{R}_{r,\alpha}}(\varphi,\theta,\xi) = \frac{\mathrm{i}^{1+\gamma}}{2\pi} \int_{0}^{2\pi} e^{\mathrm{i}\eta\xi} \partial_{\eta}^{1+\gamma} \left[\frac{\sin(\frac{\eta}{2})(e^{\mathrm{i}\eta}-1)^{\gamma}}{|\sin(\eta/2)|^{\alpha}} \mathscr{A}_{r,\alpha}^{2}(\varphi,\theta,\eta+\theta) \right] d\eta.$$

Therefore, Leibniz rule gives

$$\langle \xi \rangle^{1+\gamma} \left\| \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathscr{R}_{r,\alpha}}(\cdot, \cdot, \xi) \right\|_{H^{s}} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma+1} \int_{0}^{2\pi} \frac{\| \partial_{\eta}^{\beta} \mathscr{A}_{r,\alpha}^{2}(\cdot, \cdot, \cdot + \eta) \|_{H^{s}}}{|\sin(\eta/2)|^{\alpha}} \, d\eta.$$

Similarly we get for any $0 \leq |j| \leq q$ and $\alpha \in (0, \overline{\alpha})$ and by Sobolev embeddings

$$\begin{split} \kappa^{|j|} \langle \xi \rangle^{1+\gamma} \left\| \partial_{\lambda}^{j} \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathscr{R}_{r,\alpha}}(\cdot, \bullet, \xi) \right\|_{H^{s-|j|}} &\lesssim \sum_{\substack{0 \leqslant \beta \leqslant \gamma+1 \\ 0 \leqslant j' \leqslant j}} \kappa^{|j'|} \int_{0}^{2\pi} \frac{\| \partial_{\lambda}^{j'} \partial_{\eta}^{\beta} \mathscr{L}_{r,\alpha}^{2}(\cdot, \bullet, \bullet + \eta) \|_{H^{s-|j'|}}}{|\sin(\eta/2)|^{\frac{1}{2}}} \, d\eta \\ &\lesssim \sum_{0 \leqslant \beta \leqslant \gamma+1} \int_{0}^{2\pi} \frac{\| \partial_{\eta}^{\beta} \mathscr{L}_{r,\alpha}^{2}(\cdot, \bullet, \bullet + \eta) \|_{s}^{q,\kappa}}{|\sin(\eta/2)|^{\frac{1}{2}}} \, d\eta. \end{split}$$

By virtue of (9.23) and (9.17) we get for any $0 \leq \beta \leq 1 + \gamma$

$$\begin{aligned} \|\partial_{\eta}^{\beta}\mathscr{A}^{2}_{r,\alpha}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \leqslant C \|r\|_{s+\frac{7}{2}+\beta+\epsilon}^{q,\kappa} \\ \leqslant C \|r\|_{s+5+\gamma}^{q,\kappa}. \end{aligned}$$
(9.28)

Using the definition (7.13) we obtain

$$\|\partial_{\theta}\mathscr{R}_{r,\alpha}\|\|_{-1,s,\gamma}^{q,\kappa} \lesssim \|r\|_{s+5+\gamma}^{q,\kappa}$$

The symmetry of the functions $V_{r,\alpha}$ and $\mathscr{W}_{r,\alpha}$ follows from the expressions (9.13), (9.19). More precisely, one gets that these functions are even, that is, $V_{r,\alpha}(-\varphi, -\theta) = V_{r,\alpha}(\varphi, \theta)$ with the same property for $\mathscr{W}_{r,\alpha}$. The reversibility of the operator $\partial_{\theta}\mathscr{R}_{r,\alpha}$ in the sense of the Definition 7.1 is quite similar and one gets in view the the structures (9.22) and (9.23),

$$\partial_{\theta} \mathscr{R}_{r,\alpha} \circ \mathscr{S} = -\mathscr{S} \circ \partial_{\theta} \mathscr{R}_{r,\alpha}$$

In particular we get for $h \in H^s_{even}(\mathbb{T}^{d+1})$ that $\partial_\theta \mathscr{R}_{r,\alpha} h \in H^s_{odd}(\mathbb{T}^{d+1})$.

(iii) The proof is quite similar to (9.26). We use the identities (9.19) and (9.20) combined with the estimates

$$|\Delta_{12}\mathscr{W}_{r,\alpha}||_{s-1}^{q,\kappa} \lesssim ||\Delta_{12}r||_{s}^{q,\kappa} + ||\Delta_{12}r||_{s_0}^{q,\kappa} \max_{j=1,2} ||r_j||_{s}^{q,\kappa}$$
(9.29)

and

ŝ

$$\sup_{\eta \in \mathbb{T}} \|\Delta_{12}\mathscr{A}_{r,\alpha}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \leqslant C \|\Delta_{12}r\|_{s+\frac{7}{2}+\epsilon}^{q,\kappa} + \|\Delta_{12}r\|_{s_{0}+\frac{7}{2}+\epsilon}^{q,\kappa} \max_{j=1,2} \|r_{j}\|_{s+\frac{7}{2}+\epsilon}^{q,\kappa}.$$
(9.30)

To get the estimates (9.29) and (9.30) we proceed in a similar way to (9.25) and (9.16). This achieves the proof of Lemma 9.1.

9.3. Reduction of the transport part. In this section, we intend to perform the reduction of the transport part of the linearized operator $\mathscr{L}_{\varepsilon r}$ described in Proposition 9.1. This topic is now well-developed in KAM theory and has been implemented in several papers during the last few years, especially in [5, 12]. The formal statement says that up to a conjugation of the operator $\mathscr{L}_{\varepsilon r_{\delta}}$ by a quasi-periodic symplectic change of variables \mathscr{B} described in (9.34) the leading part becomes a transport operator with constant coefficients. In our context, we shall use the same techniques developed in the aforementioned papers in order to get a slightly different version that fits with our Cantor sets chosen to be related to the final states in the KAM reduction. This section is organized as follows. First we discuss elementary results on the invertibility of Fourier multiplier operator in the presence of a small divisor problem. In the second part we shall deal with the straightening of the transport equation when the coefficients are varying slowly around constant coefficients.

9.3.1. Transport equation with constant coefficients. Take two constants $\kappa, \rho \in (0, 1], \tau_1 > 0$ and let $\lambda = (\omega, \alpha) \in \mathcal{O} \mapsto c_{\lambda} \in \mathbb{R}$ be a smooth function. We introduce the Cantor set

$$\mathbf{C}^{\kappa,\tau_1} = \left\{ \lambda = (\omega, \alpha) \in \mathcal{O}; \ \forall (l,j) \in \mathbb{Z}^{d+1} \setminus \{0\}, \ |\omega \cdot l + jc_\lambda| > \frac{\kappa^{\varrho}}{\langle l \rangle^{\tau_1}} \right\}$$

For $N \in \mathbb{N}^{\star}$ we define the truncated Cantor set

$$\mathbf{C}_{N}^{\kappa,\tau_{1}} = \left\{ \lambda \in \mathcal{O} \, ; \ \forall (l,j) \in \mathbb{Z}^{d+1} \backslash \{0\} \text{ with } |l| \leqslant N, \, |\omega \cdot l + jc_{\lambda}| > \frac{\kappa^{\varrho}}{\langle l \rangle^{\tau_{1}}} \right\}.$$

Given $f: \mathcal{O} \times \mathbb{T}^{d+1} \to \mathbb{R}$ a smooth function with zero average, that can be expanded in Fourier series as follows

$$h = \sum_{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}} h_{l,j}(\lambda)\mathbf{e}_{l,j}, \quad \mathbf{e}_{l,j}(\varphi,\theta) = e^{\mathrm{i}(l\cdot\varphi+j\theta)}.$$

We want to solve the transport equation

$$(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})u = h \tag{9.31}$$

in the periodic setting, that is, $u : \mathbb{T}^{d+1} \to \mathbb{R}$. Then when $\lambda \in C^{\kappa,\tau_1}$ we can solve by Fourier series and invert this operator to get

$$u(\lambda,\varphi,\theta) = -\mathrm{i} \sum_{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}} \frac{h_{l,j}(\lambda)}{\omega \cdot l + jc_{\lambda}} \mathbf{e}_{l,j}.$$

We shall give an extension of this formal inverse for the full range of $\lambda \in \mathcal{O}$. To do that, we define the smooth extension of u by

$$(\omega \cdot \partial_{\varphi} + c_{r,\lambda} \partial_{\theta})_{\text{ext}}^{-1} h \triangleq -i \sum_{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\}} \frac{\chi \big(\omega \cdot l + jc_{\lambda}\big) \kappa^{-\varrho} \langle l \rangle^{\tau_1} \big) h_{l,j}(\lambda)}{\omega \cdot l + jc_{\lambda}} \mathbf{e}_{l,j},$$
(9.32)

where the cut-off function $\chi \in \mathscr{C}^{\infty}(\mathbb{R}, \mathbb{R})$ is defined in (6.30). The following result is classical and we can refer to [4], but for the sake of the completeness we will give some key ingredients of the proof.

Lemma 9.2. Let $\kappa \in (0,1], q \in \mathbb{N}$ and $\varrho \in \left(0, \frac{1}{q+1}\right]$. There exists $\varepsilon_0 \in (0,1)$ such that if $\|c_\lambda\|^{q,\kappa} \leq \varepsilon_0$, then for any $s \geq q$ we have

$$\left\| \left(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta} \right)_{\text{ext}}^{-1} h \right\|_{s}^{q,\kappa} \leqslant C \kappa^{-1} (1 + \|c_{\lambda}\|^{q,\kappa}) \|h\|_{s+\tau_{1}(q+1)}^{q,\kappa}.$$

In addition, for any $N\in\mathbb{N}$ and for any $\lambda\in\mathrm{C}_{N}^{\kappa,\tau_{1}}$ we have

$$(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})_{\text{ext}}^{-1} \Pi_{N} = \Pi_{N},$$

where Π_N is the orthogonal projection defined by

$$\Pi_N \sum_{\substack{(l,j)\in\mathbb{Z}^{d+1}\\|l|\leqslant N}} h_{l,j} \mathbf{e}_{l,j} = \sum_{\substack{(l,j)\in\mathbb{Z}^{d+1}\\|l|\leqslant N}} h_{l,j} \mathbf{e}_{l,j}.$$

Proof. The proof of the first point can be done using Faà di Bruno's formula in a similar way to [4, Lemma 2.5]. As to the identity of the second point, it follows easily from the following observation based on the explicit extension (9.32)

$$(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})_{\text{ext}}^{-1} \Pi_N h = -i \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\}\\|l| \leqslant N}} \frac{\chi \left((\omega \cdot l + jc_{\lambda}) \kappa^{-\varrho} \langle l \rangle^{\tau_1} \right) h_{l,j}(\lambda)}{\omega \cdot l + jc_{\lambda}} \mathbf{e}_{l,j}$$

By construction, one deduces for $\lambda \in \mathsf{C}_N^{\kappa,\tau_1}$ and $|l| \leqslant N$,

$$\chi((\omega \cdot l + jc_{\lambda})\kappa^{-\varrho}\langle l\rangle^{\tau_1}) = 1,$$

which implies that

$$(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})_{\text{ext}}^{-1} \Pi_N h = -i \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ |l| \le N}} \frac{h_{l,j}(\lambda)}{\omega \cdot l + jc_{\lambda}} \mathbf{e}_{l,j}.$$

Therefore, we obtain

$$(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})(\omega \cdot \partial_{\varphi} + c_{\lambda} \partial_{\theta})_{\text{ext}}^{-1} \Pi_{N} h = \mathrm{i} \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ |l| \leq N}} h_{l,j}(\lambda) \mathbf{e}_{l,j}$$
$$= \Pi_{N} h.$$

This concludes the proof of the lemma.

9.3.2. Straightening of the transport equation. This section is devoted to the construction of quasiperiodic change of variables needed to conjugate the transport part of the linearized operator to a Fourier multiplier. Before stating our main result we need to introduce some transformations and explore some of their basic properties that can be found for instance in [5, 14, 39]. Let β : $\mathcal{O} \times \mathbb{T}^{d+1} \to \mathbb{R}^d$ be a smooth function such that $\sup_{\lambda \in \mathcal{O}} \|\beta(\lambda, \cdot)\|_{\text{Lip}} < 1$ then the mapping

$$(\varphi,\theta)\in\mathbb{T}^{d+1}\mapsto(\varphi,\theta+\beta(\lambda,\varphi,\theta))\in\mathbb{T}^{d+1}$$

is a diffeomorphism and its inverse takes the same form

$$(\varphi,\theta) \in \mathbb{T}^{d+1} \mapsto (\varphi,\theta + \widehat{\beta}(\lambda,\omega,\varphi,\theta)) \in \mathbb{T}^{d+1}.$$

The relation between β and $\hat{\beta}$ is described through,

$$y = \theta + \beta(\lambda, \varphi, \theta) \Longleftrightarrow \theta = y + \beta(\lambda, \varphi, y).$$
(9.33)

Now we define the operators

$$\mathscr{B} = (1 + \partial_{\theta}\beta)\mathbf{B},\tag{9.34}$$

with

$$\mathbf{B}h(\lambda,\varphi,\theta) = h\big(\lambda,\varphi,\theta + \beta(\lambda,\varphi,\theta)\big)$$
¹³⁴

Direct computations show that the inverse \mathscr{B}^{-1} keeps the same form, that is,

$$\mathscr{B}^{-1}h(\lambda,\varphi,y) = \left(1 + \partial_y\widehat{\beta}(\lambda,\varphi,y)\right)h(\lambda,\varphi,y + \widehat{\beta}(\lambda,\varphi,y))$$
(9.35)

and

$$\mathbf{B}^{-1}h(\lambda,\varphi,y) = h\big(\lambda,\omega,\varphi,y+\widehat{\beta}(\lambda,\varphi,y)\big).$$
(9.36)

The next algebraic properties follow from straightforward computations.

Lemma 9.3. The following assertions holds true

(i) The action of \mathscr{B}^{-1} on the derivative ∂_{θ} is given by

$$\mathscr{B}^{-1}\partial_{\theta} = \partial_{\theta}\mathbf{B}^{-1}$$

(ii) The conjugation of a transport operator by \mathscr{B} keeps the same structure

$$\mathscr{B}^{-1}\big(\omega\cdot\partial_{\varphi}+\partial_{\theta}(V(\varphi,\theta)\cdot)\big)\mathscr{B}=\omega\cdot\partial_{\varphi}+\partial_{y}\big(\mathscr{V}(\varphi,y)\cdot\big)$$

with

$$\mathscr{V}(\varphi, y) = \mathbf{B}^{-1} \Big(\omega \cdot \partial_{\varphi} \beta(\varphi, \theta) + V(\varphi, \theta) \big(1 + \partial_{\theta} \beta(\varphi, \theta) \big) \Big)$$

(iii) Denote by \mathscr{B}^{\star} the $L^{2}_{\theta}(\mathbb{T})$ -adjoint of \mathscr{B} , then

$$\mathscr{B}^{\star} = \mathbf{B}^{-1}$$
 and $\mathbf{B}^{\star} = \mathscr{B}^{-1}$

The next result deals with some analytical properties of the preceding transformations. The proof can be obtained by adapting the proof accomplished in [39].

Lemma 9.4. Let $q \in \mathbb{N}, \gamma \in (0,1), s \ge s_0 > \frac{d+1}{2} + q + 1$. There exists ε_0 small enough such that if $\|\beta\|_{2s_0}^{q,\kappa} \le \varepsilon_0$, then we have the estimates

$$\|\mathbf{B}^{\pm 1}h\|_{s}^{q,\kappa} \leq \|h\|_{s}^{q,\kappa} \left(1 + C\|\beta\|_{s_{0}}^{q,\kappa}\right) + C\|\beta\|_{s}^{q,\kappa}\|h\|_{s_{0}}^{q,\kappa},$$
$$\|\mathscr{B}^{\pm 1}h\|_{s}^{q,\kappa} \leq \|h\|_{s}^{q,\kappa} \left(1 + C\|\beta\|_{s_{0}}^{q,\kappa}\right) + C\|\beta\|_{s+1}^{q,\kappa}\|h\|_{s_{0}}^{q,\kappa}$$

and

$$\|\widehat{\beta}\|_{s}^{q,\kappa} \lesssim \|\beta\|_{s}^{q,\kappa} \quad and \quad \|\mathbf{B}^{\pm 1}h - h\|_{s}^{q,\kappa} \lesssim \|h\|_{s_{0}}^{q,\kappa}\|\beta\|_{s+1}^{q,\kappa} + \|\beta\|_{s_{0}}^{q,\kappa}\|h\|_{s+1}^{q,\kappa}.$$

Furthermore, let $\beta_1, \beta_2 \in W^{q,\infty,\gamma}(\mathcal{O}, H^{\infty}(\mathbb{T}^{d+1}))$ satisfying the foregoing smallness condition and denote

$$\Delta_{12}\beta = \beta_1 - \beta_2$$
 and $\Delta_{12}\beta = \beta_1 - \beta_2$

Then, we have the estimate

$$\|\Delta_{12}\widehat{\beta}\|_{s}^{q,\kappa} \leqslant C\left(\|\Delta_{12}\beta\|_{s}^{q,\kappa} + \|\Delta_{12}\beta\|_{s_{0}}^{q,\kappa} \max_{k \in \{1,2\}} \|\beta_{i}\|_{s+1}^{q,\kappa}\right).$$
(9.37)

In what follows we intend to state the main result of this section concerning the reduction of the transport part.

Proposition 9.2. Let $\varrho \in \left(0, \frac{1}{q+1}\right]$ and assume (9.2). There exists $\varepsilon_0 > 0$ such that with the conditions

$$\mu_{2} \geqslant \overline{\mu}_{2} \triangleq 4(\tau_{1}q + \tau_{1}) + 3, \qquad s_{l} \triangleq s_{0} + \tau_{1}q + \tau_{1} + 2, \\ \overline{s}_{h} \triangleq \frac{3}{2}\overline{\mu}_{2} + s_{l} + 1, \qquad s_{h} \triangleq \frac{3}{2}\mu_{2} + s_{l} + 1, \\ \sigma_{1} \triangleq \tau_{1}q + \tau_{1} + 7$$

$$(9.38)$$

and

$$\|\mathfrak{I}_0\|_{q,s_h+\sigma_1} \leqslant 1 \quad \text{and} \quad N_0^{\mu_2} \varepsilon \kappa^{-2} \leqslant \varepsilon_0,$$

$$(9.39)$$

there exist

$$\lambda \mapsto c(\lambda, i_0) \in W^{q, \infty, \gamma}(\mathcal{O}, \mathbb{R}) \quad and \quad \beta \in \bigcap_{s \in [s_0, S]} W^{q, \infty, \gamma}(\mathcal{O}, H^s_{\text{odd}})$$

such that with \mathscr{B} as in (9.34) one gets the following results.

(i) The function $c(\lambda, i_0)$ satisfies the following estimate,

$$\|c(\lambda, i_0) - V_{0,\alpha}\|^{q,\kappa} \lesssim \varepsilon \tag{9.40}$$

where $V_{0,\alpha}$ is defined in (3.11).

(ii) The transformations $\mathscr{B}^{\pm 1}, \mathbf{B}^{\pm 1}, \beta$ and $\widehat{\beta}$ satisfy the following estimates for all $s \in [s_0, S]$

$$\begin{aligned} \|\mathscr{B}^{\pm 1}h\|_{s}^{q,\kappa} + \|\mathbf{B}^{\pm 1}h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-1} \|\mathfrak{I}_{0}\|_{s+\sigma_{1}}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}, \qquad (9.41) \\ \|(\mathscr{B} - \mathrm{Id})h\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(\|h\|_{s+1}^{q,\kappa} + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}\right) \end{aligned}$$

and

$$\|\widehat{\beta}\|_{s}^{q,\kappa} \lesssim \|\beta\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}}^{q,\kappa}\right).$$

$$(9.42)$$

(iii) On the Cantor set

$$\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0) = \bigcap_{\substack{(l,j)\in\mathbb{Z}^d\times\mathbb{Z}\setminus\{(0,0)\}\\|l|\leqslant N_n}} \left\{\lambda\in\mathcal{O}; \ \left|\omega\cdot l+jc(\lambda,i_0)\right| > 4\kappa^{\varrho}\frac{\langle j\rangle}{\langle l\rangle^{\tau_1}}\right\}$$

we have

$$\mathscr{B}^{-1}\big(\omega\cdot\partial_{\varphi}+\partial_{\theta}\big(V_{\varepsilon r,\alpha}\cdot\big)\big)\mathscr{B}=\omega\cdot\partial_{\varphi}+c(\lambda,i_{0})\partial_{\theta}+\mathsf{E}_{n}^{0}$$

with \mathbf{E}_n^0 a linear operator satisfying

$$\|\mathbf{E}_{n}^{0}h\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon N_{0}^{\mu_{2}}N_{n+1}^{-\mu_{2}}\|h\|_{s_{0}+2}^{q,\kappa}.$$

The function $V_{\varepsilon r,\alpha}$ was defined in Lemma 9.1.

(iv) Given two tori i_1 and i_2 both satisfying (9.39) (replacing \mathfrak{I}_0 by \mathfrak{I}_1 or \mathfrak{I}_2), then

$$\|\Delta_{12}c(\cdot,i)\|^{q,\kappa} \lesssim \varepsilon \|\Delta_{12}i\|^{q,\kappa}_{\overline{s}_h+4}.$$
(9.43)

In addition, assume that $0 \leq a \leq \frac{3}{2}(\mu_2 - \overline{\mu}_2) + 1 - \tau_1$, then

$$\|\Delta_{12}\beta\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} + \|\Delta_{12}\widehat{\beta}\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma_1+\mathbf{a}}^{q,\kappa}.$$
(9.44)

Before giving the proof, some remarks are in order.

- **Remark 9.1.** (i) The final Cantor set $\mathcal{G}_{\infty,n}^{\kappa,\tau_1}(i_0)$ is constructed over the limit coefficient $c(\lambda, i_0)$ but it is still truncated in the time frequency, that is $|l| \leq N_n$, leading to a residual reminder with enough decay through the parameter μ_2 that can be arbitrarily chosen by selecting the regularity index s_h large enough. This induces a suitable stability property which is crucial during the Nash-Moser scheme achieved with the nonlinear functional.
 - (ii) The estimate (9.44) holds if μ_2 satisfies $\mathbf{a} + \tau_1 1 \leq \frac{3}{2}(\mu_2 \overline{\mu}_2)$ for some $\mathbf{a} \geq 0$. This is slightly stronger than the constraint imposed in (9.38)
 - (iii) The constant 4 that appears in the definition of the Cantor set $\mathbb{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ is used to ensure the inclusion of this set in all the Cantor sets built along the KAM procedure.

Proof. The proof will be done in the same spirit of [14, 39] and based on the construction of successive iterations of linear transformations through quasi-periodic symplectic change of coordinates. Notice that at each step of the scheme we should extract from the reminder of size ε its main diagonal part leading to a new reminder with size ε^2 . This can be done through solving the homological equation which requires non-resonances conditions satisfied by excision of the external parameters $\lambda = (\omega, \alpha)$. Iterating this argument allows to get the desired result with a final Cantor set constructed over all the restrictions coming from the different homological equations. To be more precise, we shall describe this procedure in the KAM step and see later how to implement it.

(i)-(ii) ► KAM step. Assume that we have a transport operator taking the form,

$$(\omega \cdot \partial_{\varphi} + \partial_{\theta} (V + f))h = \omega \cdot \partial_{\varphi}h + \partial_{\theta} ((V + f)h)$$

where the parameter λ belongs to a subset $\mathcal{O}_{-}^{\gamma} \subset \mathcal{O}$ and

 $V = V(\lambda)$ and $f = f(\lambda, \varphi, \theta)$

with f being an even function,

$$f(\lambda, -\varphi, -\theta) = f(\lambda, \varphi, \theta).$$
(9.45)

136

Next, we introduce a symplectic quasi-periodic change of coordinates close to the identity in the form

$$\mathscr{B}h(\lambda,\varphi,\theta) \triangleq (1 + \partial_{\theta}g(\lambda,\varphi,\theta))\mathbf{B}h(\lambda,\varphi,\theta)
= (1 + \partial_{\theta}g(\lambda,\varphi,\theta))h(\lambda,\varphi,\theta + g(\lambda,\varphi,\theta))$$
(9.46)

with $g: \mathcal{O} \times \mathbb{T}^{d+1} \to \mathbb{R}$ being a small function to be adjusted later with respect to f. Then according to Lemma 9.3, we may write for any $N \ge 2$

$$\mathscr{B}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V+f\right)\right)\mathscr{B}=\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(\mathbf{B}^{-1}\left(V+\omega\cdot\partial_{\varphi}g+V\partial_{\theta}g+\Pi_{N}f+\Pi_{N}^{\perp}f+f\partial_{\theta}g\right)\cdot\right).$$
(9.47)

The main goal is to obtain after this transformation a new transport operator in the form

$$\mathscr{B}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V+f\right)\right)\mathscr{B}=\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{+}+f_{+}\right)$$
(9.48)

where

$$V_+ = V_+(\lambda)$$
 and $f_+ = f_+(\lambda, \varphi, \theta)$

with f_+ quadratically smaller than f. Coming back to (9.47) and in order to get rid of the *linear* terms in f, we shall impose to the perturbation g the following *homological equation*

$$\omega \cdot \partial_{\varphi}g + V \partial_{\theta}g + \Pi_N f = \langle f \rangle_{\varphi,\theta} \tag{9.49}$$

where the average of f is defined by

$$\langle f \rangle_{\varphi,\theta} = \frac{1}{(2\pi)^{d+1}} \int_{\mathbb{T}^{d+1}} f(\lambda,\varphi,\theta) d\varphi d\theta$$

To solve the homological equation (9.49), we use Fourier decomposition in order to recover g in the form

$$g(\lambda,\varphi,\theta) = \sum_{\substack{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}\\\langle l,j\rangle\leq N}} \frac{\mathrm{i}\,f_{l,j}(\lambda)}{\omega\cdot l + jV(\lambda)} e^{\mathrm{i}(l\cdot\varphi+j\theta)}.$$
(9.50)

Then at this level we should deal with the small divisors problem which is a classical issue in KAM theory. One way to fix it is to avoid resonances through Diophantine conditions type by restricting the exterior parameters to the following set

$$\mathcal{O}_{+}^{\gamma} \triangleq \bigcap_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ \langle l,j \rangle \leq N}} \left\{ \lambda = (\omega, \alpha) \in \mathcal{O}_{-}^{\gamma}; \ \left| \omega \cdot l + jV(\lambda) \right| > \frac{\kappa^{\varrho}\langle j \rangle}{\langle l \rangle^{\tau_{1}}} \right\}.$$
(9.51)

With this choice we can control the size of the denominators in (9.50) and then expect to get suitable estimates for g with some loss of regularity uniformly in N. Before proceeding with this task we need to construct an extension of g to the whole set \mathcal{O} and for the sake of simplicity it will be still denoted by g. This can be done by extending the Fourier coefficients of g in (9.50) using the cut-off function χ defined in (6.30) in the following way

$$g_{l,j}(\lambda) \triangleq i \frac{\chi \left((\omega \cdot l + jV(\lambda)) (\kappa^{\varrho} \langle j \rangle)^{-1} \langle l \rangle^{\tau_1} \right)}{\omega \cdot l + jV(\lambda)} f_{l,j}(\lambda).$$
(9.52)

In what follows, we shall work with this extension which is a solution to (9.49) when the parameters are restricted to the set \mathcal{O}_{+}^{γ} . We then define

$$V_{+} = V + \langle f \rangle_{\varphi,\theta}$$
 and $f_{+} = \mathbf{B}^{-1} (\Pi_{N}^{\perp} f + f \partial_{\theta} g)$

so that in the set \mathscr{O}_{+}^{γ} , the identity (9.48) holds. Notice that V_{+} and f_{+} are well-defined in the whole set of parameters \mathscr{O} and the function g is smooth since it is generated by a finite number of frequencies. From the assumption (9.45) we get that g is odd and therefore

$$g \in \bigcap_{s \ge 0} W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{odd}}).$$
(9.53)

Next, we intend to estimate the Fourier coefficients $g_{l,j}$ defined in (9.52) which can be written in the form

$$g_{l,j}(\lambda) = ia_{l,j} \chi_1(a_{l,j}A_{l,j}(\lambda)) f_{l,j}(\lambda), \quad A_{l,j}(\lambda) = \omega \cdot l + jV(\lambda),$$

$$a_{l,j} = (\kappa^{\varrho} \langle j \rangle)^{-1} \langle l \rangle^{\tau_1}, \quad \chi_1(x) = \frac{\chi(x)}{x}.$$
(9.54)

Since χ_1 is C^{∞} with bounded derivatives and $\chi_1(0) = 0$, then we may apply Lemma 4.5 giving

$$\forall |\gamma| \leq q, \quad \|g_{l,j}\|_{W^{|\gamma|,\infty}(\mathcal{O})} \lesssim a_{l,j}^2 \|A_{l,j}\|_{W^{|\gamma|,\infty}(\mathcal{O})} \Big(1 + a_{l,j}^{|\gamma|-1} \|A_{l,j}\|_{L^{\infty}(\mathcal{O})}^{|\gamma|-1}\Big).$$

It is straightforward that

$$\begin{aligned} \forall (l,j) \in \mathbb{Z}^{d+1}, \, \forall \gamma \in \mathbb{N}^{d+1}, \quad |\gamma| \leq q, \quad \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} A_{l,j}(\lambda) \right| \lesssim \langle l,j \rangle \left(1 + \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} V(\lambda) \right| \right) \\ \lesssim \kappa^{-|\gamma|} \langle l,j \rangle \left(1 + \|V\|^{q,\kappa} \right). \end{aligned}$$

Hence by assuming

$$\|V\|^{q,\kappa} \leqslant C \tag{9.55}$$

we obtain

$$\forall (l,j) \in \mathbb{Z}^{d+1}, \quad \forall \gamma \in \mathbb{N}^{d+1}, \quad |\gamma| \leqslant q \quad \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} A_{l,j}(\lambda) \right| \lesssim \kappa^{-|\gamma|} \langle l,j \rangle.$$

Therefore we find that

$$\forall |\gamma| \leqslant q, \quad \|g_{l,j}\|_{W^{|\gamma|,\infty}(\mathcal{O})} \lesssim a_{l,j}^2 \kappa^{-|\gamma|} \langle l,j \rangle \left(1 + a_{l,j}^{|\gamma|-1} \langle l,j \rangle^{|\gamma|-1}\right)$$

Since $0 \leqslant a_{l,j} \leqslant \kappa^{-\varrho} \langle l \rangle^{\tau_1}$ then the foregoing estimate gives

$$\forall |\gamma| \leq q, \quad |\partial_{\lambda}^{\gamma} g_{l,j}| \lesssim \kappa^{-\varrho(|\gamma|+1)-|\gamma|} \langle l,j \rangle^{\tau_1(1+|\gamma|)+|\gamma|}.$$
(9.56)

By choosing ρ such that

$$\varrho \leqslant \frac{1}{q+1} \tag{9.57}$$

and using Leibniz rule or the law products, we infer from the Definition 4.1

$$\|g\|_{s}^{q,\kappa} \lesssim \kappa^{-1} \|\Pi_{N}f\|_{s+\tau_{1}q+\tau_{1}}^{q,\kappa}.$$
(9.58)

Assume

$$\kappa^{-1} N^{\tau_1 q + \tau_1 + 1} \| f \|_{s_0}^{q,\kappa} \leqslant \varepsilon_0, \tag{9.59}$$

then combined with (9.58) and Lemma 4.2, we get

$$|g||_{s_0+1}^{q,\kappa} \leqslant C\kappa^{-1}N^{\tau_1q+\tau_1+1} ||f||_{s_0}^{q,\kappa} \leqslant C\varepsilon_0.$$

Hence, taking ε_0 small enough we may guarantee the smallness condition in Lemma 9.4 and get that the linear operator \mathscr{B} is invertible. We now set

$$u = \Pi_N^{\perp} f + f \partial_\theta g.$$

By Lemma 4.1 and (9.58), we deduce for all $s \in [s_0, S]$

$$\|u\|_{s}^{q,\kappa} \leq \|\Pi_{N}^{\perp}f\|_{s}^{q,\kappa} + C\Big(\|f\|_{s_{0}}^{q,\kappa}\|\partial_{\theta}g\|_{s}^{q,\kappa} + \|f\|_{s}^{q,\kappa}\|\partial_{\theta}g\|_{s_{0}}^{q,\kappa}\Big) \\ \leq \|\Pi_{N}^{\perp}f\|_{s}^{q,\kappa} + C\kappa^{-1}N^{\tau_{1}q+\tau_{1}+1}\|f\|_{s_{0}}^{q,\kappa}\|f\|_{s}^{q,\kappa}.$$

Applying Lemma 9.4, Lemma 4.1 and (9.59), we obtain

$$\begin{aligned} \|f_{+}\|_{s}^{q,\kappa} &= \|\mathbf{B}^{-1}(u)\|_{s}^{q,\kappa} \\ &\leqslant \|u\|_{s}^{q,\kappa} + C\Big(\|u\|_{s}^{q,\kappa}\|\widehat{g}\|_{s_{0}}^{q,\kappa} + \|\widehat{g}\|_{s}^{q,\kappa}\|u\|_{s_{0}}^{q,\kappa}\Big) \\ &\leqslant \|u\|_{s}^{q,\kappa} + C\Big(\|u\|_{s}^{q,\kappa}\|g\|_{s_{0}}^{q,\kappa} + \|g\|_{s}^{q,\kappa}\|u\|_{s_{0}}^{q,\kappa}\Big) \\ &\leqslant \|\Pi_{N}^{\perp}f\|_{s}^{q,\kappa} + C\kappa^{-1}N^{\tau_{1}q+\tau_{1}+1}\|f\|_{s_{0}}^{q,\kappa}\|f\|_{s}^{q,\kappa}. \end{aligned}$$

Therefore, Lemma 4.2 yields for $s_0 \leq s \leq \overline{s} \leq S$

$$\|f_{+}\|_{s}^{q,\kappa} \leqslant N^{s-\overline{s}} \|f\|_{q,\overline{s}}^{\gamma,0} + C\kappa^{-1} N^{\tau_{1}q+\tau_{1}+1} \|f\|_{s_{0}}^{q,\kappa} \|f\|_{s}^{q,\kappa}.$$
(9.60)

▶ KAM scheme. Assume that we have constructed V_m and f_m for $m \ge 0$, being well-defined in the whole set of parameters \mathcal{O} and verifying the assumptions (9.55) and (9.59), we want to construct V_{m+1} and f_{m+1} still satisfying (9.55) and (9.59). For this purpose, we shall apply the KAM step with (V, f, V_+, f_+, N) replaced by $(V_m, f_m, V_{m+1}, f_{m+1}, N_m)$. To be more precise, we will prove by induction the existence of a sequence $\{V_m, f_m\}_{m\in\mathbb{N}}$ such that

$$\delta_m(s_l) \leq \delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2}, \qquad \delta_m(s_h) \leq \left(2 - \frac{1}{m+1}\right) \delta_0(s_h)$$
(9.61)

and

$$\|V_m\|^{q,\kappa} \leqslant C, \qquad N_m^{\tau_1 q + \tau_1 + 1} \delta_m(s_0) \leqslant \varepsilon_0, \tag{9.62}$$

supplemented with the symmetry condition

$$f_m(\lambda, -\varphi, -\theta) = f_m(\lambda, \varphi, \theta), \qquad (9.63)$$

where

$$\delta_m(s) \triangleq \kappa^{-1} \|f_m\|_s^{q,\kappa}.$$

We remind that the parameters s_l and s_h in (9.61) are defined in (9.38).

♦ Initialization. Let us check the properties (9.61) and (9.62) with m = 0. In this case we start with the transport operator

$$\omega \cdot \partial_{\varphi} + \partial_{\theta} (V_{\varepsilon r,\alpha} \cdot)$$

and use the following decomposition based on Lemma 9.1

$$V_{\varepsilon r,\alpha} = V_{0,\alpha} + (V_{\varepsilon r,\alpha} - V_{0,\alpha})$$

$$\triangleq V_0(\alpha) + f_0. \tag{9.64}$$

Remind that $V_{r,\alpha}$ and $V_{0,\alpha}$ are defined in (9.5) and (3.11). Applying Lemma 9.1-(i) and Proposition 9.1 yield

$$\delta_{0}(s) = \kappa^{-1} \| V_{\varepsilon r} - V_{0} \|_{s+\kappa}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \| r \|_{s+4}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_{0} \|_{s+4}^{q,\kappa} \right).$$

$$(9.65)$$

Then, using the smallness condition (9.39) we obtain

$$N_0^{\mu_2} \delta_0(s_h) \leqslant C \varepsilon_0. \tag{9.66}$$

Moreover, by (9.5) and the symmetry of r we get

$$f_0(\lambda, -\varphi, -\theta) = f_0(\lambda, \varphi, \theta) \tag{9.67}$$

We consider $\mathcal{O}_0^{\gamma} = \mathcal{O}$ and $N_0 \ge 2$ and we shall check that the assumptions (9.55) and (9.59) are satisfied with V_0 and f_0 . First recall that V_0 is defined by (3.11). From the C^{∞} -regularity of Gamma function, we easily obtain

$$\|V_0\|^{q,\kappa} \leqslant C. \tag{9.68}$$

Then the assumption (9.55) is satisfied for $V = V_0$. Now, applying (9.66) and using the assumption on μ_2 we infer

$$\kappa^{-1} N_0^{\tau_1 q + \tau_1 + 1} \| f_0 \|_{s_0}^{q,\kappa} = N_0^{\tau_1 q + \tau_1 + 1} \delta_0(s_0)$$

$$\leqslant N_0^{\tau_1 q + \tau_1 + 1 - \mu_2} N_0^{\mu_2} \delta_0(s_h)$$

$$\leqslant C \varepsilon_0 N_0^{-1}.$$

Taking N_0 large enough such that

$$CN_0^{-1} \leqslant 1 \tag{9.69}$$

we find

$$\kappa^{-1} N_0^{\tau_1 q + \tau_1 + 1} \|f_0\|_{s_0}^{q,\kappa} \leqslant \varepsilon_0$$

Thus, the condition (9.59) is satisfied for $f = f_0$ which achieves the initialization step.

◆ Induction. Assume that we have constructed V_m and f_m with the properties (9.61), (9.62) and (9.63), and let us construct V_{m+1} and f_{m+1} and check the validity of these constraints at this order. Following the KAM step, we may consider a symplectic quasi-periodic change of variables \mathscr{B}_m in the form

$$\mathscr{B}_m h(\lambda, \varphi, \theta) \triangleq \left(1 + \partial_\theta g_m(\lambda, \varphi, \theta) \right) \mathbf{B}_m h(\lambda, \varphi, \theta) \\= \left(1 + \partial_\theta g_m(\lambda, \varphi, \theta) \right) h(\lambda, \varphi, \theta + g_m(\lambda, \varphi, \theta))$$

with

$$g_m(\lambda,\varphi,\theta) \triangleq \sum_{\substack{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}\\\langle l,j\rangle\leqslant N_m}} i\frac{\chi\left((\omega\cdot l+jV_m(\lambda))(\kappa^{\varrho}\langle j\rangle)^{-1}\langle l\rangle^{\tau_1}\right)}{\omega\cdot l+jV_m(\lambda)}(f_m)_{l,j}(\lambda)\,e^{\mathrm{i}(l\cdot\varphi+j\theta)}$$
(9.70)

where χ is the cut-off function defined in (6.30) and N_m is defined in (6.32). As it was explained during the KAM step, g_m is well-defined on the whole set of parameters \mathcal{O} and when it is restricted to the Cantor set

$$\mathcal{O}_{m+1}^{\kappa} \triangleq \bigcap_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ \langle l,j \rangle \in N_m}} \left\{ \lambda \in \mathcal{O}_m^{\kappa}; \ \left| \omega \cdot l + j V_m(\lambda) \right| > \frac{\kappa^{\varrho}\langle j \rangle}{\langle l \rangle^{\tau_1}} \right\}$$
(9.71)

it solves the following homological equation

 $\omega \cdot \partial_{\varphi} g_m + V_m \partial_{\theta} g_m + \Pi_{N_m} f_m = \langle f_m \rangle_{\varphi,\theta}.$

Therefore in the Cantor set $\mathscr{O}_{m+1}^{\gamma}$, we find the following reduction

$$\mathscr{B}_m^{-1} \Big(\omega \cdot \partial_{\varphi} + \partial_{\theta} (V_m + f_m) \Big) \mathscr{B}_m = \omega \cdot \partial_{\varphi} + \partial_{\theta} (V_{m+1} + f_{m+1})$$
(9.72)

where V_{m+1} and f_{m+1} are defined by

$$\begin{cases} V_{m+1} = V_m + \langle f_m \rangle_{\varphi,\theta} \\ f_{m+1} = \mathbf{B}_m^{-1} \Big(\Pi_{N_m}^{\perp} f_m + f_m \partial_{\theta} g_m \Big). \end{cases}$$
(9.73)

According to (9.63), the function g_m is odd and consequently we deduce that f_{m+1} is even which concludes the symmetry persistence during the scheme. On the other hand, one gets similarly to (9.53) that

$$g_m \in \bigcap_{s \ge 0} W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{odd}})$$
(9.74)

We set

 $\overline{\mathscr{B}}_{-1} = \mathrm{Id} \quad \text{and for } \mathbf{m} \in \mathbb{N}, \quad \overline{\mathscr{B}}_m = \mathscr{B}_0 \circ \mathscr{B}_1 \circ \dots \circ \mathscr{B}_m.$ (9.75)

Then we may check that

$$\overline{\mathscr{B}}_m h(\lambda,\varphi,\theta) \triangleq \left(1 + \partial_\theta \beta_m(\lambda,\varphi,\theta)\right) h(\lambda,\varphi,\theta + \beta_m(\lambda,\varphi,\theta))$$

where $(\beta_m)_{m \in \mathbb{N}}$ is defined by $\beta_{-1} = g_{-1} = 0$ and

$$\beta_0 = g_0 \quad \text{and} \quad \beta_m(\lambda, \varphi, \theta) = \beta_{m-1}(\lambda, \varphi, \theta) + g_m(\lambda, \varphi, \theta + \beta_{m-1}(\lambda, \varphi, \theta)).$$
(9.76)

By a trivial induction using (9.74) and Lemma 4.4 we have

$$\beta_m \in \bigcap_{s \ge 0} W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{odd}}).$$
(9.77)

From Sobolev embeddings, (9.73) and the induction hypothesis (9.61), we have

$$\|V_m - V_{m-1}\|^{q,\kappa} = \|\langle f_{m-1} \rangle_{\varphi,\theta}\|^{q,\kappa} \leq \|f_{m-1}\|_{s_0}^{q,\kappa} = \gamma \delta_{m-1}(s_0) \leq \kappa \delta_0(s_h) N_0^{\mu_2} N_{m-1}^{-\mu_2}.$$
(9.78)

This implies in view of the triangle inequality combined with (9.66) and with ε_0 small enough

$$\begin{aligned} \|V_m\|^{q,\kappa} &\leq \|V_{m-1}\|^{q,\kappa} + \kappa \delta_0(s_h) N_0^{\mu_2} N_{m-1}^{-\mu_2} \\ &\leq \|V_0\|^{q,\kappa} + \kappa \delta_0(s_h) N_0^{\mu_2} \sum_{k=0}^{m-1} N_k^{-\mu_2} \\ &\leq \|V_0\|^{q,\kappa} + \sum_{k=0}^{+\infty} N_k^{-\mu_2}. \end{aligned}$$

On the other hand, by (9.61), (9.66) and since $\mu_2 \ge \tau_1 q + \tau_1 + 2$ in view of (9.2), we deduce by the assumption (9.69)

$$\delta_m(s_0) N_m^{\tau_1 q + \tau_1 + 1} \leqslant \delta_0(s_h) N_0^{\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - \mu_2} \leqslant C \varepsilon_0 N_0^{-1} \leqslant \varepsilon_0.$$
(9.79)

Therefore we obtain in view of (9.68) and (9.79)

$$\sup_{m \in \mathbb{N}} \|V_m\|^{q,\kappa} \leqslant C \quad \text{and} \quad \delta_m(s_0) N_m^{\tau_1 q + \tau_1 + 1} \leqslant \varepsilon_0.$$

At this stage, we may apply the KAM step and the estimate (9.60) writes in our case

$$\delta_{m+1}(s) \leqslant N_m^{s-\overline{s}} \delta_m(\overline{s}) + C N_m^{\tau_1 q + \tau_1 + 1} \delta_m(s) \delta_m(s_0).$$
(9.80)

Applying (9.80) with $s = s_l$ and $\overline{s} = s_h$, we get

$$\delta_{m+1}(s_l) \leqslant N_m^{s_l - s_h} \delta_m(s_h) + C N_m^{\tau_1 q + \tau_1 + 1} \delta_m(s_l) \delta_m(s_0).$$

Putting together (9.61) and the fact that $s_l \ge s_0$ combined with Sobolev embeddings yields

$$\delta_{m+1}(s_l) \leq N_m^{s_l - s_h} \delta_m(s_h) + C N_m^{\tau_1 q + \tau_1 + 1} (\delta_m(s_l))^2$$

$$\leq \left(2 - \frac{1}{m+1}\right) N_m^{s_l - s_h} \delta_0(s_h) + C N_0^{2\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - 2\mu_2} (\delta_0(s_h))^2$$

$$\leq 2 N_m^{s_l - s_h} \delta_0(s_h) + C N_0^{2\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - 2\mu_2} (\delta_0(s_h))^2.$$

If we select our parameters s_l , s_h and μ_2 such that

$$N_m^{s_l - s_h} \leqslant \frac{1}{4} N_0^{\mu_2} N_{m+1}^{-\mu_2} \quad \text{and} \quad C N_0^{2\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - 2\mu_2} \delta_0(s_h) \leqslant \frac{1}{2} N_0^{\mu_2} N_{m+1}^{-\mu_2} \tag{9.81}$$

then

$$\delta_{m+1}(s_l) \leqslant \delta_0(s_h) N_0^{\mu_2} N_{m+1}^{-\mu_2}$$

Notice that by our choice of parameters (9.2), the condition (9.81) is fulfilled provided that

$$4N_0^{-\mu_2} \leqslant 1$$
 and $2C\delta_0(s_h) \leqslant N_0^{-\mu_2}$.

The first condition is automatically satisfied by taking N_0 sufficiently large and the second condition holds by (9.66) provided that $\overline{\varepsilon}$ is small enough. This proves the first statement of the induction in (9.61) and we now turn to the proof of the second statement. Making use of (9.80) with $s = \overline{s}$ and using the induction assumption (9.61), we find

$$\delta_{m+1}(s_h) \leq \delta_m(s_h) \left(1 + CN_m^{\tau_1 q + \tau_1 + 1} \delta_m(s_0) \right)$$
$$\leq \left(2 - \frac{1}{m+1} \right) \delta_0(s_h) \left(1 + CN_0^{\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - \mu_2} \delta_0(s_h) \right).$$

Therefore we deduce from (9.38) and (9.39)

$$\left(2 - \frac{1}{m+1}\right) \left(1 + CN_0^{\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - \mu_2} \delta_0(s_h)\right) \leqslant 2 - \frac{1}{m+2},\tag{9.82}$$

then we find

$$\delta_{m+1}(s_h) \leqslant \left(2 - \frac{1}{m+2}\right) \delta_0(s_h).$$

This estimate achieves the induction argument of (9.61). Now, observe that (9.82) is equivalent to

$$\left(2 - \frac{1}{m+1}\right) C N_0^{\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - \mu_2} \delta_0(s_h) \leqslant \frac{1}{(m+1)(m+2)}$$

By virtue of (9.2) we have $\mu_2 \ge \tau_1 q + \tau_1 + 2$ and thus the preceding condition holds true if

$$CN_0^{\mu_2}N_m^{-1}\delta_0(s_h) \leqslant \frac{1}{(m+1)(m+2)}$$
(9.83)

Recall that $N_0 \ge 2$, and then in view of (6.32) we may find a constant $c_0 > 0$ small enough such that

$$\forall m \in \mathbb{N}, \quad c_0 N_m^{-1} \leqslant \frac{1}{(m+1)(m+2)}$$

Therefore, (9.83) is satisfied provided that

$$CN_0^{\mu_2}\delta_0(s_h) \leqslant c_0. \tag{9.84}$$

By taking ε small enough, we can ensure from (9.66) that

$$CN_0^{\mu_2}\delta_0(s_h) \leqslant C\varepsilon_0$$
$$\leqslant c_0.$$

Thus, the condition (9.84) is satisfied. This completes the proof of (9.61).

◆ Regularity persistence. Applying (9.80) with $\overline{s} = s \in [s_0, S]$, (9.61) and (9.2), we obtain

$$\delta_{m+1}(s) \leqslant \delta_m(s) \left(1 + C N_m^{\tau_1 q + \tau_1 + 1} \delta_m(s_0) \right)$$

$$\leq \delta_m(s) \left(1 + C \delta_0(s_h) N_0^{\mu_2} N_m^{\tau_1 q + \tau_1 + 1 - \mu_2} \right)$$

$$\leq \delta_m(s) \left(1 + C N_m^{-1} \right).$$

Combining this estimate with (9.65), we infer by a trivial induction

$$\delta_m(s) \leqslant \delta_0(s) \prod_{k=0}^{+\infty} \left(1 + CN_k^{-1}\right)$$

$$\leqslant C\delta_0(s)$$

$$\leqslant C\varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+4}^{q,\kappa}\right).$$

(9.85)

Putting together (9.58), the interpolation inequality from Lemma 4.3 and (9.61) leads to

$$\begin{aligned} \|g_m\|_s^{q,\kappa} &\leq C\delta_m(s+\tau_1q+\tau_1) \\ &\leq C\left(\delta_m(s_0)\right)^{\overline{\theta}(s)} \left(\delta_m(s+\tau_1q+\tau_1+1)\right)^{1-\overline{\theta}(s)} \\ &\leq C\delta_0^{\overline{\theta}(s)}(s_h) N_0^{\overline{\theta}(s)\mu_2} N_m^{-\overline{\theta}(s)\mu_2} \delta_m^{1-\overline{\theta}(s)}(s+\tau_1q+\tau_1+1) \end{aligned}$$

with $\overline{\theta}(s) \triangleq \frac{1}{s+\tau_1 q+\tau_1+1-s_0}$. Using (9.85), (9.39) and (9.65)

$$\|g_{m}\|_{s}^{q,\kappa} \leq C\varepsilon\kappa^{-1}\left(1+\|\mathfrak{I}_{0}\|_{s_{h}+4}^{q,\kappa}\right)\left(1+\|\mathfrak{I}_{0}\|_{s+\tau_{1}q+\tau_{1}+5}^{q,\kappa}\right)N_{0}^{\theta(s)\mu_{2}}N_{m}^{-\overline{\theta}(s)\mu_{2}}$$
$$\leq C\varepsilon\kappa^{-1}\left(1+\|\mathfrak{I}_{0}\|_{s+\tau_{1}q+\tau_{1}+5}^{q,\kappa}\right)N_{0}^{\overline{\theta}(s)\mu_{2}}N_{m}^{-\overline{\theta}(s)\mu_{2}}.$$
(9.86)

By (9.76) and Lemma 9.4, we deduce for all $s \in [s_0, S]$

$$\|\beta_m\|_s^{q,\kappa} \le \|\beta_{m-1}\|_s^{q,\kappa} \left(1 + C\|g_m\|_{s_0}^{q,\kappa}\right) + C\left(1 + \|\beta_{m-1}\|_{s_0}^{q,\kappa}\right) \|g_m\|_s^{q,\kappa}.$$
(9.87)

Applying this estimate with $s = s_0$ together with Sobolev embeddings,

$$\|\beta_m\|_{s_0}^{q,\kappa} \leq \|\beta_{m-1}\|_{s_0}^{q,\kappa} \left(1 + C\|g_m\|_{s_0}^{q,\kappa}\right) + C\|g_m\|_{s_0}^{q,\kappa}.$$

In order to obtain a good estimate for β_m , we shall use the following result which is quite easy to prove by induction : Given three positive sequences $(a_n)_{n\in\mathbb{N}}$, $(b_n)_{n\in\mathbb{N}}$ and $(c_n)_{n\in\mathbb{N}}$ such that

$$\forall n \in \mathbb{N}, \quad a_{n+1} \leqslant b_n a_n + c_n$$

Then

$$\forall n \ge 2, \quad a_n \leqslant a_0 \prod_{i=0}^{n-1} b_i + \sum_{k=0}^{n-2} c_k \prod_{i=k+1}^{n-1} b_i + c_{n-1} \\ \leqslant \left(a_0 + \sum_{k=0}^{n-1} c_k\right) \prod_{i=0}^{n-1} b_i.$$

$$(9.88)$$

In particular if $\prod_{n=0}^{\infty} b_n$ and $\sum_{n=0}^{\infty} c_n$ converge then

$$\sup_{n \in \mathbb{N}} a_n \leqslant \left(a_0 + \sum_{n=0}^{\infty} c_n\right) \prod_{n=0}^{\infty} b_i.$$
(9.89)

Since (9.38) ensures that $s_h \ge s_0 + \tau_1 q + \tau_1 + 1$ and $\mu_2 \ge \tau_1 q + \tau_1 + 1$, then by (9.86) and (9.39) we get

$$\begin{aligned} \|g_m\|_{s_0}^{q,\kappa} &\leq C\varepsilon\kappa^{-1}N_0^{\mu_2} \left(1 + \|\mathfrak{I}_0\|_{s_0+\tau_1q+\tau_1+5}^{q,\kappa}\right)N_m^{-\overline{\theta}(s_0)\mu_2} \\ &\leq C\varepsilon_0N_m^{-1}. \end{aligned}$$

By (6.32) and $N_0 \ge 2$ we infer $\sum_{m=0}^{\infty} N_m^{-1} < \infty$. Therefore if ε_0 is small enough in such a way $C\varepsilon_0 \le 1$, and applying (9.89) together with the fact that $\beta_0 = g_0$ we find

$$\sup_{m \in \mathbb{N}} \|\beta_m\|_{s_0}^{q,\kappa} \leq \left(\|\beta_0\|_{s_0}^{q,\kappa} + C \sum_{k=0}^{+\infty} \|g_k\|_{s_0}^{q,\kappa} \right) \prod_{k=0}^{\infty} \left(1 + C \|g_k\|_{s_0}^{q,\kappa} \right) \\ \leq \left(1 + C \sum_{k=0}^{\infty} N_k^{-1} \right) \prod_{k=0}^{\infty} \left(1 + N_k^{-1} \right) \leq C.$$
(9.90)

Plugging this estimate into (9.87) gives for all $s \in [s_0, S]$

$$\|\beta_m\|_s^{q,\kappa} \leq \|\beta_{m-1}\|_s^{q,\kappa} \left(1 + C\|g_m\|_{s_0}^{q,\kappa}\right) + C\|g_m\|_s^{q,\kappa}.$$

As above, applying (9.89) and (9.86), we find

$$\sup_{m\in\mathbb{N}} \|\beta_m\|_s^{q,\kappa} \leq \left(\|\beta_0\|_s^{q,\kappa} + C\sum_{k=0}^\infty \|g_k\|_s^{q,\kappa}\right) \prod_{k=0}^\infty \left(1 + C\|g_k\|_{s_0}^{q,\kappa}\right)$$
$$\leq C\varepsilon\kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\tau_1q+\tau_1+5}^{q,\kappa}\right) \left(1 + N_0^{\overline{\theta}(s)\mu_2}\sum_{k=0}^\infty N_k^{-\overline{\theta}(s)\mu_2}\right)$$

Using Lemma A.2 implies the existence of a constant C > 0 such that

$$\forall s \in [s_0, S], \quad N_0^{\overline{\theta}(s)\mu_2} \sum_{k=0}^{+\infty} N_k^{-\overline{\theta}(s)\mu_2} \leqslant C$$

and this allows to get

$$\forall s \in [s_0, S], \quad \sup_{m \in \mathbb{N}} \|\beta_m\|_s^{q,\kappa} \leqslant C \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\tau_1 q+\tau_1+5}^{q,\kappa}\right). \tag{9.91}$$

By Lemma 9.4, (9.90), (9.58), (9.61) and in view of $s_l \ge s_0 + \tau_1 q + \tau_1 + 2$, we obtain,

$$\|\beta_m - \beta_{m-1}\|_{s_0+2}^{q,\kappa} \leqslant C \|g_m\|_{s_0+2}^{q,\kappa} \left(1 + \|\beta_{m-1}\|_{s_0+2}^{q,\kappa}\right)$$

$$\leq C\delta_m(s_l)$$

$$\leq CN_0^{\mu_2}N_m^{-\mu_2}\delta_0(s_h). \tag{9.92}$$

Hence we get the convergence of the series

$$\sum_{m=0}^{\infty} \|\beta_m - \beta_{m-1}\|_{s_0+2}^{q,\kappa} < \infty.$$

Consequently, the sequence $(\beta_m)_{m\in\mathbb{N}}$ converges strongly towards some function $\beta \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^{s_0+2})$. From the uniform boundedness (9.91) we deduce that $\beta \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s)$ and

$$\forall s \in [s_0, S], \quad \|\beta\|_s^{q,\kappa} \leq \liminf_{m \to \infty} \|\beta_m\|_s^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\tau_1 q+\tau_1+5}^{q,\kappa}\right).$$
(9.93)

Define now the quasi-periodic symplectic change of variables \mathscr{B} associated with β by

 ∞

$$\mathscr{B}h(\lambda,\omega,\varphi,\theta) = (1 + \partial_{\theta}\beta(\lambda,\omega,\varphi,\theta)) \mathbf{B}h(\lambda,\omega,\varphi,\theta) = (1 + \partial_{\theta}\beta(\lambda,\omega,\varphi,\theta)) h(\lambda,\omega,\varphi,\theta + \beta(\lambda,\omega,\varphi,\theta)).$$

Notice that by (9.93), hyptothesis (9.39) and (9.38), we infer

$$\|\beta\|_{s_0}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s_0+\tau_1 q+\tau_1+5}^{q,\kappa}\right) \leqslant C\varepsilon_0.$$

$$(9.94)$$

Hence choosing ε_0 small enough, we get by Lemma 9.4 that \mathscr{B} is an invertible operator, and combined with (9.93) we obtain

$$\|\mathscr{B}^{\pm 1}h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-1} \|\mathfrak{I}_{0}\|_{s+\tau_{1}q+\tau_{1}+6}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}.$$
(9.95)

Notice that the symmetry of β follows from the symmetry of the approximation (β_m) described in (9.77). In addition, one easily gets from (9.92)

$$\|\beta - \beta_{m}\|_{s_{0}+2}^{q,\kappa} \leq \sum_{k=m}^{\infty} \|\beta_{k+1} - \beta_{k}\|_{s_{0}+2}^{q,\kappa}$$
$$\lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} \sum_{k=m+1}^{\infty} N_{k}^{-\mu_{2}}.$$
(9.96)

Applying Lemma A.2 yields

$$\sum_{k=m}^{+\infty} N_k^{-\mu_2} \underset{m \to \infty}{=} O\left(N_m^{-\mu_2}\right).$$
(9.97)

Putting together (9.97) and (9.96), we obtain

$$\|\beta - \beta_m\|_{s_0+2}^{q,\kappa} \lesssim \kappa \delta_0(s_h) N_0^{\mu_2} N_{m+1}^{-\mu_2}.$$
(9.98)

• KAM conclusion. According to (9.78) and (9.97) one may write,

. . .

$$\sum_{m=0}^{\infty} \|V_{m+1} - V_m\|^{q,\kappa} \leqslant \kappa \delta_0(s_h) N_0^{\mu_2} \sum_{m=0}^{\infty} N_m^{-\mu_2} \lesssim \kappa \delta_0(s_h).$$

Thus we get that the sequence $(V_m)_{m\in\mathbb{N}}$ converges towards an element $\lambda \mapsto c(\lambda, i_0) \in W^{q,\infty}_{\kappa}(\mathcal{O}, \mathbb{R})$. We denote c_{i_0} its limit and one gets

$$c_{i_0} = V_0 + \sum_{m \in \mathbb{N}} (V_{m+1} - V_m).$$
(9.99)

Therefore by applying (9.65) we find

$$\|c_{i_0} - V_0\|^{q,\kappa} \leq \sum_{\substack{m=0\\144}}^{\infty} \|V_{m+1} - V_m\|^{q,\kappa}$$

$$\lesssim \varepsilon \left(1 + \| \mathfrak{I}_0 \|_{s_h + 4}^{q, \kappa} \right) \lesssim \varepsilon_0$$

Next, we consider the truncated Cantor set

$$\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0) = \bigcap_{\substack{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}\\|l|\leqslant N_n}} \left\{ \lambda = (\omega,\alpha)\in\mathcal{O}\,;\,\,\left|\omega\cdot l + jc(\lambda,i_0)\right| > \frac{4\kappa^\varrho\langle j\rangle}{\langle l\rangle^{\tau_1}} \right\}$$

and we intend to prove that this set satisfies the inclusion

$$\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0) \subset \bigcap_{m=0}^{n+1} \mathscr{O}_m^{\kappa} = \mathscr{O}_{n+1}^{\kappa}$$
(9.100)

where the intermediate Cantor sets are defined in (9.71). For this goal, we proceed by induction in m. First, notice that by construction we have $\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0) \subset \mathscr{O} \triangleq \mathscr{O}_0^{\kappa}$. Second, assume that $\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0) \subset \mathscr{O}_m^{\kappa}$ for some $m \leqslant n$ and let us check that

$$\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0) \subset \mathcal{O}_{m+1}^{\kappa}.\tag{9.101}$$

Remark that by (9.78) and (9.97)

$$|V_m(\lambda) - c(\lambda, i_0)| \leq ||V_m - c(\cdot, i_0)||^{q,\kappa}$$
$$\leq \sum_{l=m}^{\infty} ||V_{l+1} - V_l||^{q,\kappa}$$
$$\lesssim \kappa \delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2}.$$
(9.102)

Let $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ and $(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\}$ with $0 \leq |l| \leq N_m$. Then $|l| \leq N_n$ and by the triangle inequality combined with (9.102), (9.66) and $\varrho, \kappa \in (0, 1)$ we infer

$$\begin{aligned} |\omega \cdot l + jV_m(\lambda)| \geqslant &|\omega \cdot l + jc(\lambda, i_0)| - |j||V_m(\lambda) - c(\lambda, i_0) \\ \geqslant \frac{4\kappa^{\varrho}\langle j\rangle}{\langle l\rangle^{\tau_1}} - C\langle j\rangle\kappa\delta_0(s_h)N_0^{\mu_2}N_m^{-\mu_2} \\ \geqslant \frac{4\kappa^{\varrho}\langle j\rangle}{\langle l\rangle^{\tau_1}} - C\langle j\rangle\kappa^{\varrho}\varepsilon_0\langle l\rangle^{-\mu_2} \end{aligned}$$

Then for $C\varepsilon_0 \leq 1$ and since $\mu_2 \geq \tau_1$ (in view of (9.2)), we get

$$|\omega \cdot l + jV_m(\lambda)| > \frac{\kappa^{\varrho} \langle j \rangle}{\langle l \rangle^{\tau_1}}$$

which shows that $\lambda \in \mathcal{O}_{m+1}^{\gamma}$ and therefore the inclusion (9.101) is satisfied. (iii) We shall start with the following decomposition

$$\mathcal{B}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\right)\mathcal{B}=\mathcal{B}_{n}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\right)\mathcal{B}_{n}$$
$$+\left(\mathcal{B}^{-1}-\mathcal{B}_{n}^{-1}\right)\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\right)\mathcal{B}$$
$$+\mathcal{B}_{n}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\right)\left(\mathcal{B}-\mathcal{B}_{n}\right)$$

By (9.100), (9.72) and (9.75) we have that on the Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\mathscr{B}_{n}^{-1}\left(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\right)\mathscr{B}_{n}=\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{n+1}+f_{n+1}\right)$$

Hence, in the Cantor set $\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$, the following splitting holds

$$\mathscr{B}^{-1}\Big(\omega\cdot\partial_{\varphi}+\partial_{\theta}\left(V_{0}+f_{0}\right)\Big)\mathscr{B}=\omega\cdot\partial_{\varphi}+c(\lambda,i_{0})\partial_{\theta}+\mathsf{E}_{n}^{0}(\lambda,i_{0})$$

with

$$\mathbf{E}_{n}^{0}(\lambda, i_{0}) \triangleq (V_{n+1} - c(\lambda, i_{0})) \partial_{\theta} + \partial_{\theta} (f_{n+1} \cdot) + (\mathscr{B}^{-1} - \mathscr{B}_{n}^{-1}) \left(\omega \cdot \partial_{\varphi} + \partial_{\theta} (V_{0} + f_{0}) \right) \mathscr{B} + \mathscr{B}_{n}^{-1} \left(\omega \cdot \partial_{\varphi} + \partial_{\theta} (V_{0} + f_{0}) \right) (\mathscr{B} - \mathscr{B}_{n})$$

$$(\mathscr{B} - \mathscr{B}_{n})$$

$$(\mathscr{B} - \mathscr{B}_{n})$$

$$\triangleq \mathbf{E}_{n,1}^{0}(\lambda, i_{0}) + \mathbf{E}_{n,2}^{0}(\lambda, i_{0}) + \mathbf{E}_{n,3}^{0}(\lambda, i_{0}) + \mathbf{E}_{n,4}^{0}(\lambda, i_{0}).$$

Applying the law products in Lemma 4.1 together with (9.102), we find

$$\|\mathbf{E}_{n,1}^{0}h\|_{s_{0}}^{q,\kappa} \lesssim \|V_{n+1} - c(\cdot, i_{0})\|_{s_{0}}^{q,\kappa}\|h\|_{s_{0}+1}^{q,\kappa} \lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}}\|h\|_{s_{0}+1}^{q,\kappa}.$$
(9.103)

Concerning the second term we use Lemma 4.1 together with (9.61) and (9.85)

$$\begin{aligned} \|\mathbf{E}_{n,2}^{0}h\|_{s_{0}}^{q,\kappa} &= \|\partial_{\theta}\left(f_{n+1}h\right)\|_{s_{0}}^{q,\kappa} \\ &\lesssim \kappa \delta_{n+1}(s_{l})\|h\|_{s_{0}+1}^{q,\kappa} \\ &\lesssim \kappa \delta_{0}(s_{h})N_{0}^{\mu_{2}}N_{n+1}^{-\mu_{2}}\|h\|_{s_{0}+1}^{q,\kappa}. \end{aligned}$$
(9.104)

We now move to the estimate of $E_{n,3}^0$. Using Lemma 4.1, we deduce that

$$\begin{aligned} \|\omega \cdot \partial_{\varphi} h + \partial_{\theta} \left(V_{\varepsilon r} h \right) \|_{s_{0}}^{q,\kappa} &\leq \|\omega \cdot \partial_{\varphi} h\|_{s}^{q,\kappa} + \|V_{\varepsilon r} h\|_{s_{0}+1}^{q,\kappa} \\ &\lesssim \|h\|_{s_{0}+1}^{q,\kappa} \left(1 + \|V_{\varepsilon r}\|_{s_{0}+1}^{q,\kappa} \right). \end{aligned}$$

Putting together (9.64), (9.68) and (9.65), we find

$$\|V_{\varepsilon r}\|_{s_0+1}^{q,\kappa} \leq \|V_0\|^{q,\kappa} + \|f_0\|_{s_0+1}^{q,\kappa} \\ \leq C$$

Hence by(9.39), we obtain

$$\|\omega \cdot \partial_{\varphi} h + \partial_{\theta} \left(V_{\varepsilon r} h \right) \|_{s_0}^{q,\kappa} \lesssim \|h\|_{s_0+1}^{q,\kappa}.$$

$$(9.105)$$

Then, by virtue of Taylor Formula,

$$(\mathscr{B} - \mathscr{B}_n)h(\theta) = (1 + \partial_{\theta}\beta(\theta)) \left[h(\theta + \beta(\theta)) - h(\theta + \beta_n(\theta))\right] + \partial_{\theta} \left(\beta(\theta) - \beta_n(\theta)\right) h(\theta + \beta_n(\theta))$$
$$\triangleq (1 + \partial_{\theta}\beta(\theta))(\beta(\theta) - \beta_n(\theta))\mathbf{I}_n h(\theta) + \partial_{\theta}(\beta(\theta) - \beta_n(\theta))\mathbf{B}_n h(\theta),$$

with

$$\mathbf{I}_n h(\theta) \triangleq \int_0^1 \left(\partial_\theta h\right) \left(\theta + \beta_n(\theta) + t(\beta(\theta) - \beta_n(\theta))\right) dt.$$

Combining Lemma 9.4 and Lemma 4.1 yields

$$\begin{aligned} \|\partial_{\theta} \left(\beta - \beta_{n}\right) \mathbf{B}_{n} h\|_{s_{0}+1}^{q,\kappa} \lesssim \|\beta - \beta_{n}\|_{s_{0}+2}^{q,\kappa} \|h\|_{s_{0}+1}^{q,\kappa} \left(1 + \|\beta_{n}\|_{s_{0}+1}^{q,\kappa}\right) \\ \lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{s_{0}}^{q,\kappa} \end{aligned}$$

On the other hand, performing similar arguments as before we find

$$\left\| \left(1 + \partial_{\theta} \beta \right) \left(\beta - \beta_{n} \right) \mathbf{I}_{n} h \right\|_{s_{0}+1}^{q,\kappa} \lesssim \left(1 + \|\beta\|_{s_{0}+2}^{q,\kappa} \right) \|\beta - \beta_{n}\|_{s_{0}+1}^{q,\kappa} \|h\|_{s_{0}+2}^{q,\kappa} \left(1 + \|\beta_{n}\|_{s_{0}+1}^{q,\kappa} + \|\beta - \beta_{n}\|_{s_{0}+1}^{q,\kappa} \right) \\ \lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{s_{0}+2}^{q,\kappa}.$$

Putting together the preceding estimates, it follows that

$$\|(\mathscr{B} - \mathscr{B}_n)h\|_{s_0+1}^{q,\kappa} \lesssim \kappa \delta_0(s_h) N_0^{\mu_2} N_{n+1}^{-\mu_2} \|h\|_{s_0+2}^{q,\kappa}.$$

Thus we find by collecting (9.105), (9.95), (9.65) and (9.39)

$$\begin{aligned} |\mathbf{E}_{n,4}^{0}h||_{s_{0}}^{q,\kappa} &\lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} ||h||_{s_{0}+2}^{q,\kappa} \\ &\lesssim \varepsilon N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} ||h||_{s_{0}+2}^{q,\kappa}. \end{aligned}$$
(9.106)

In a similar way, using in particular (9.35) we find

$$\|\mathbf{E}_{n,3}^{0}h\|_{s_{0}}^{q,\kappa} \lesssim \kappa \delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{s_{0}+2}^{q,\kappa} \lesssim \varepsilon N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{s_{0}+2}^{q,\kappa}.$$
(9.107)

Gathering (9.103), (9.107), (9.106), yields

$$\|\mathbf{E}_{n}^{0}h\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon N_{0}^{\mu_{2}}N_{n+1}^{-\mu_{2}}\|h\|_{s_{0}+2}^{q,\kappa}.$$

(iv) We shall start with the estimate of $\Delta_{12}\beta$. For this aim we notice that, since $\beta_{-1} = 0$,

$$\Delta_{12}\beta = \sum_{m=0}^{\infty} \Delta_{12}(\beta_m - \beta_{m-1}).$$
(9.108)

Hence by the triangule inequality we get

$$\|\Delta_{12}\beta\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \leqslant \sum_{m=0}^{\infty} \|\Delta_{12}(\beta_m - \beta_{m-1})\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa}.$$
(9.109)

By Taylor formula and (9.76), we find by removing the dependance with respect to λ and φ

$$\Delta_{12}\beta_m(\theta) = \Delta_{12}\beta_{m-1}(\theta) + \mathbf{B}_{m-1}^{[1]}(\Delta_{12}g_m)(\theta) + \Delta_{12}\beta_{m-1}(\theta) \int_0^1 (\partial_\theta g_m^{[2]}) \big(\theta + \beta_{m-1}^{[2]}(\theta) + t\Delta_{12}\beta_{m-1}(\theta)\big) dt,$$

which implies that

$$\Delta_{12}(\beta_m - \beta_{m-1})(\theta) = \mathbf{B}_{m-1}^{[1]}(\Delta_{12}g_m)(\theta) + \Delta_{12}\beta_{m-1}(\theta) \int_0^1 (\partial_\theta g_m^{[2]}) (\theta + \beta_{m-1}^{[2]}(\theta) + t\Delta_{12}\beta_{m-1}(\theta)) dt.$$

Thus, by law product in Lemma 4.1, Lemma 9.4 and Sobolev embeddings

$$\begin{split} \|\Delta_{12}(\beta_{m}-\beta_{m-1})\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} &\leqslant \|\Delta_{12}g_{m}\|_{\overline{s}_{h}+\mathbf{p}}^{q,\kappa} \left(1+C\|\beta_{m-1}^{[1]}\|_{s_{0}}^{q,\kappa}\right) + \|\Delta_{12}g_{m}\|_{s_{0}}^{q,\kappa}\|\beta_{m-1}^{[1]}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \\ &+ C\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\|g_{m}^{[2]}\|_{\overline{s}_{h}+\mathbf{a}+1}^{q,\kappa} \left(1+\|\beta_{m-1}^{[2]}\|_{s_{0}}^{q,\kappa}+\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\right) \\ &+ C\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\|g_{m}^{[2]}\|_{s_{0}+1}^{q,\kappa} \left(\|\beta_{m-1}^{[2]}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa}+\|\Delta_{12}\beta_{m-1}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa}\right) \\ &+ C\|\Delta_{12}\beta_{m-1}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa}\|g_{m}^{[2]}\|_{s_{0}+1}^{q,\kappa} \left(1+\|\beta_{m-1}^{[2]}\|_{s_{0}}^{q,\kappa}+\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\right) \end{split}$$

and for all $s \in [s_0, \overline{s}_h + \mathbf{a}]$

$$\begin{split} \|\Delta_{12}\beta_{m}\|_{s}^{q,\kappa} &\leqslant \|\Delta_{12}g_{m}\|_{s}^{q,\kappa} \left(1+C\|\beta_{m-1}^{[1]}\|_{s_{0}}^{q,\kappa}\right) + \|\Delta_{12}g_{m}\|_{s_{0}}^{q,\kappa}\|\beta_{m-1}^{[1]}\|_{s}^{q,\kappa} \\ &+ C\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\|g_{m}^{[2]}\|_{s+1}^{q,\kappa} \left(1+\|\beta_{m-1}^{[2]}\|_{s_{0}}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\right) \\ &+ C\|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\|g_{m}^{[2]}\|_{s_{0}+1}^{q,\kappa} \left(\|\beta_{m-1}^{[2]}\|_{s}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{s}^{q,\kappa}\right) \\ &+ \|\Delta_{12}\beta_{m-1}\|_{s}^{q,\kappa} \left(1+C\|g_{m}^{[2]}\|_{s_{0}+1}^{q,\kappa} \left(1+\|\beta_{m-1}^{[2]}\|_{s_{0}}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\right)\right). \end{split}$$

Here we adopt the generic notation $g^{[k]}$ to denote the value of g at r_k (or equivalently to i_k). By (9.86), (9.39) and (9.38), we have since $\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 6 \leq s_h + \sigma_1$

$$\sup_{m\in\mathbb{N}}\max_{k\in\{1,2\}} \|g_m^{[k]}\|_{\overline{s}_h+\mathbf{a}+1}^{q,\kappa} \leqslant C\varepsilon\kappa^{-1} \left(1+\max_{k\in\{1,2\}} \|\mathfrak{I}_0^{[k]}\|_{\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+6}^{q,\kappa}\right) \\ \leqslant C, \tag{9.110}$$

Actually, we get a more precise estimate

$$\max_{k \in \{1,2\}} \|g_m^{[k]}\|_{q,\overline{s}_h + \mathbf{a} + 1} \leqslant C \varepsilon \kappa^{-1} \left(1 + \max_{k \in \{1,2\}} \|\mathfrak{I}_0^{[k]}\|_{\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 6} \right) N_0^{\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2} N_m^{-\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2} \\ \leqslant C \varepsilon \kappa^{-1} N_0^{\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2} N_m^{-\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2}.$$

$$(9.111)$$

The estimate (9.91) and (9.39) allow to get

$$\sup_{m \in \mathbb{N}} \max_{k \in \{1,2\}} \|\beta_m^{[k]}\|_{q,\overline{s}_h + \mathbf{a}} \leqslant C.$$
(9.112)

Putting together (9.110), (9.112) with the previous two estimates imply

$$\|\Delta_{12}(\beta_m - \beta_{m-1})\|_{\overline{s}_h + \mathbf{a}}^{q,\kappa} \leqslant C\left(\|\Delta_{12}g_m\|_{\overline{s}_h + \mathbf{p}}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{\overline{s}_h + \mathbf{p}}^{q,\kappa}\|g_m^{[2]}\|_{\overline{s}_h + \mathbf{a}+1}^{q,\kappa}\right),\tag{9.113}$$

$$\|\Delta_{12}\beta_m\|_{s_0}^{q,\kappa} \leqslant C \|\Delta_{12}g_m\|_{s_0}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{s_0}^{q,\kappa} \left(1 + C \|g_m^{[2]}\|_{s_0+1}^{q,\kappa}\right)$$
(9.114)

and

$$\|\Delta_{12}\beta_{m}\|_{\bar{s}_{h}+\mathbf{a}}^{q,\kappa} \leq C\left(\|\Delta_{12}g_{m}\|_{\bar{s}_{h}+\mathbf{a}}^{q,\kappa} + \|\Delta_{12}\beta_{m-1}\|_{s_{0}}^{q,\kappa}\|g_{m}^{[2]}\|_{q,\underline{s}_{h}+\mathbf{p}+1}\right) + \|\Delta_{12}\beta_{m-1}\|_{\bar{s}_{h}+\mathbf{a}}^{q,\kappa}\left(1+C\|g_{m}^{[2]}\|_{s_{0}+1}^{q,\kappa}\right).$$
(9.115)

Applying (9.89) with (9.114) combined with the fact that $\beta_0 = g_0$, we conclude that

$$\sup_{m \in \mathbb{N}} \|\Delta_{12}\beta_m\|_{s_0}^{q,\kappa} \leqslant \left(\|\Delta_{12}g_0\|_{s_0}^{q,\kappa} + C\sum_{m=0}^{\infty} \|\Delta_{12}g_m\|_{s_0}^{q,\kappa} \right) \prod_{m=0}^{\infty} \left(1 + \|g_m^{[2]}\|_{s_0+1}^{q,\kappa} \right).$$

Therefore we obtain from (9.111),

$$\sup_{m\in\mathbb{N}} \|\Delta_{12}\beta_m\|_{s_0}^{q,\kappa} \leqslant C \sum_{m=0}^{\infty} \|\Delta_{12}g_m\|_{s_0}^{q,\kappa}.$$

In the same way, from (9.115), using (9.89), (9.111) and the previous estimate, we conclude that

$$\sup_{m\in\mathbb{N}} \|\Delta_{12}\beta_m\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \leqslant C \sum_{m=0}^{\infty} \|\Delta_{12}g_m\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa}.$$

Gathering the previous bounds with (9.113) and (9.111), we finally obtain

$$\|\Delta_{12}(\beta_m - \beta_{m-1})\|_{\overline{s}_h + \mathbf{a}}^{q,\kappa} \lesssim \|\Delta_{12}g_m\|_{\overline{s}_h + \mathbf{a}}^{q,\kappa} + \varepsilon \kappa^{-1} N_0^{\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2} N_m^{-\overline{\theta}(\overline{s}_h + \mathbf{a} + 1)\mu_2} \sum_{k=0}^{+\infty} \|\Delta_{12}g_k\|_{\overline{s}_h + \mathbf{a}}^{q,\kappa}.$$
(9.116)

The next goal is to estimate $\Delta_{12}g_m$. First observe that from (9.70) and (9.54) we can write

$$g_m(\lambda,\varphi,\theta) = \mathrm{i} \sum_{\substack{(l,j)\in\mathbb{Z}^{d+1}\setminus\{0\}\\\langle l,j\rangle\leqslant N_m}} a_{l,j}\,\chi_1\big(a_{l,j}A_{l,j}(\lambda)\big)(f_m)_{l,j}(\lambda)\,e^{\mathrm{i}(l\cdot\varphi+j\theta)}.$$

Then we find

$$\begin{split} \Delta_{12}g_{m} = &\mathbf{i} \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ \langle l,j \rangle \leqslant N_{m}}} a_{l,j} \, \chi_{1} \left(a_{l,j} A_{l,j}^{[1]} \right) (\Delta_{12}f_{m})_{l,j} \, \mathbf{e}_{l,j} \\ &+ \mathbf{i} \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ \langle l,j \rangle \leqslant N_{m}}} a_{l,j} \, \Delta_{12}\chi_{1} \left(a_{l,j} A_{l,j}^{[1]} \right) (f_{m}^{[2]})_{l,j} \, \mathbf{e}_{l,j} \\ &\triangleq \mathbf{I}_{1} + \mathbf{I}_{2}, \end{split}$$

where we recall that the notation $A_{l,j}^{[k]}$ stands for the value of $A_{l,j}$ evaluated at r_k , (or the torus i_k) and the same thing applies for $f_m^{[k]}$. The estimate of the first term \mathbf{I}_1 is quite similar to (9.58) and one gets

$$\|\mathbf{I}_1\|_s^{q,\kappa} \lesssim \kappa^{-1} \|\Pi_{N_m} \Delta_{12} f_m\|_{s+\tau_1 q+\tau_1}^{q,\kappa}$$

As to the second term \mathbf{I}_2 we write in view of Taylor formula

$$\mathbf{I}_{2} = \mathbf{i} \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\}\\ \langle l,j \rangle \leqslant N_{m}}} a_{l,j}^{2} (\Delta_{12} A_{l,j}) \int_{0}^{1} \chi_{1}' \left(a_{l,j} \left[(1-\tau) A_{l,j}^{[1]} + \tau A_{l,j}^{[2]} \right] \right) d\tau \left(f_{m}^{[2]} \right)_{l,j} \mathbf{e}_{l,j}.$$
(9.117)

Then by the same analysis developed to get (9.58), using in particular,

$$\|\Delta_{12}A_{l,j}\|^{q,\kappa} \lesssim \langle l,j\rangle \|\Delta_{12}V_m\|^{q,\kappa}$$

and

$$\|a_{l,j}\Delta_{12}A_{l,j}\|^{q,\kappa} \lesssim \kappa^{-\varrho} \langle l,j \rangle^{1+\tau_1} \|\Delta_{12}V_m\|^{q,\kappa}$$

we find

$$\begin{aligned} \|\mathbf{I}_{2}\|_{s}^{q,\kappa} &\lesssim \kappa^{-q(1+\varrho)} \|\Delta_{12}V_{m}\|^{q,\kappa} \|\Pi_{N_{m}}f_{m}^{[2]}\|_{s+\tau_{1}q+2\tau_{1}+1}^{q,\kappa} \\ &\lesssim \kappa^{-1} \|\Delta_{12}V_{m}\|^{q,\kappa} \|\Pi_{N_{m}}f_{m}^{[2]}\|_{s+\tau_{1}q+2\tau_{1}+1}^{q,\kappa}. \end{aligned}$$

It follows that for any $s \ge s_0$

$$\begin{aligned} \|\Delta_{12}g_m\|_s^{q,\kappa} &\lesssim \kappa^{-1} \|\Pi_{N_m} \Delta_{12}f_m\|_{s+\tau_1q+\tau_1}^{q,\kappa} + \kappa^{-1} \|\Delta_{12}V_m\|^{q,\kappa} \|\Pi_{N_m}f_m^{[2]}\|_{s+\tau_1q+2\tau_1+1}^{q,\kappa} \\ &\lesssim \kappa^{-1}N_m^{\tau_1q+\tau_1} \|\Delta_{12}f_m\|_s^{q,\kappa} + \kappa^{-1} \|\Delta_{12}V_m\|^{q,\kappa} \|\Pi_{N_m}f_m^{[2]}\|_{s+\tau_1q+2\tau_1+1}^{q,\kappa}. \end{aligned}$$
(9.118)

Now, we need to estimate $\Delta_{12} f_m$. Then by setting

$$u_m \triangleq \Pi_{N_m}^{\perp} f_m + f_m \partial_\theta g_m, \qquad (9.119)$$

and making appeal to (9.73) we obtain

$$\Delta_{12} f_{m+1} = (\mathbf{B}_m^{-1})^{[1]} \Delta_{12} u_m + \left(\Delta_{12} \mathbf{B}_m^{-1}\right) u_m^{[2]}$$

with

$$\Delta_{12}u_m = \prod_{N_m}^{\perp} \Delta_{12}f_m + \Delta_{12}f_m \partial_\theta g_m^{[1]} + f_m^{[2]} \partial_\theta \Delta_{12}g_m.$$

Thus we get for any $s \ge s_0$

$$\|\Delta_{12}f_{m+1}\|_{s}^{q,\kappa} \leq \|(\mathbf{B}_{m}^{-1})^{[1]}\Delta_{12}u_{m}\|_{s}^{q,\kappa} + \|(\Delta_{12}\mathbf{B}_{m}^{-1})u_{m}^{[2]}\|_{s}^{q,\kappa}.$$
(9.120)

By Lemma 9.4, (9.36), (9.58) and Lemma 4.1, we have for all $s \ge s_0$

$$\begin{aligned} \|(\mathbf{B}_{m}^{-1})^{[1]}\Delta_{12}u_{m}\|_{s}^{q,\kappa} &\leq \|\Delta_{12}u_{m}\|_{s}^{q,\kappa} \left(1+C\|(\widehat{g}_{m}^{[1]}\|_{s_{0}}^{q,\kappa}\right)+C\|(\widehat{g}_{m}^{[1]}\|_{s}^{q,\kappa}\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa} \\ &\leq \|\Delta_{12}u_{m}\|_{s}^{q,\kappa} \left(1+C\|g_{m}^{[1]}\|_{s_{0}}^{q,\kappa}\right)+C\|g_{m}^{[1]}\|_{s}^{q,\kappa}\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa} \\ &\leq \|\Delta_{12}u_{m}\|_{s}^{q,\kappa} \left(1+C\kappa^{-1}\|f_{m}^{[1]}\|_{s_{0}+\tau_{1}q+\tau_{1}}^{q,\kappa}\right)+C\kappa^{-1}\|f_{m}^{[1]}\|_{s+\tau_{1}q+\tau_{1}}^{q,\kappa}\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa}. \end{aligned}$$

From (9.85) and (9.39), one has

$$\kappa^{-1} \sup_{m \in \mathbb{N}} \max_{k \in \{1,2\}} \|f_m^{[k]}\|_{\overline{s}_h + \mathbf{a}}^{q,\kappa} \leqslant C \varepsilon \kappa^{-1} \left(1 + \max_{k \in \{1,2\}} \|(\mathfrak{I}_0^{[k]}\|_{\overline{s}_h + \mathbf{a} + \sigma_1}^q) \right)$$
$$\leqslant C. \tag{9.121}$$

Thus, by (9.61) and (9.121), we obtain for all $s \in [s_0, \overline{s}_h + a]$

$$\|(\mathbf{B}_{m}^{-1})^{[1]}\Delta_{12}u_{m}\|_{s}^{q,\kappa} \leq \|\Delta_{12}u_{m}\|_{s}^{q,\kappa} \left(1 + CN_{0}^{\overline{\mu}_{2}}N_{m}^{-\overline{\mu}_{2}}\right) + CN_{m}^{\tau_{1}q+\tau_{1}}\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa}.$$
(9.122)

To estimate $\Delta_{12}u_m$ we turn to (9.119) and use Lemma 4.1 in order to get for all $s \ge s_0$

$$\begin{aligned} \|\Delta_{12}u_m\|_s^{q,\kappa} &\leqslant \|\Pi_{N_m}^{\perp}\Delta_{12}f_m\|_s^{q,\kappa} + C\|\Delta_{12}f_m\|_s^{q,\kappa}\|g_m^{[1]}\|_{s_0+1}^{q,\kappa} + C\|\Delta_{12}f_m\|_{s_0}^{q,\kappa}\|g_m^{[1]}\|_{s+1}^{q,\kappa} \\ &+ C\|f_m^{[2]}\|_s^{q,\kappa}\|\Delta_{12}g_m\|_{s_0+1}^{q,\kappa} + C\|f_m^{[2]}\|_{s_0}^{q,\kappa}\|\Delta_{12}g_m\|_{s+1}^{q,\kappa}.\end{aligned}$$

Applying (9.58) yields for all $s \ge s_0$

$$\begin{split} \|\Delta_{12}u_m\|_s^{q,\kappa} &\leqslant \|\Pi_{N_m}^{\perp}\Delta_{12}f_m\|_s^{q,\kappa} + C\kappa^{-1} \|\Delta_{12}f_m\|_s^{q,\kappa} \|f_m^{[1]}\|_{s_0+\tau_1q+\tau_1+1}^{q,\kappa} \\ &+ C\kappa^{-1} \|\Delta_{12}f_m\|_{s_0}^{q,\kappa} \|\Pi_{N_m}f_m^{[1]}\|_{s+\tau_1q+\tau_1+1}^{q,\kappa} + C \|f_m^{[2]}\|_s^{q,\kappa} \|\Delta_{12}g_m\|_{s_0+1}^{q,\kappa} \\ &+ C \|f_m^{[2]}\|_{s_0}^{q,\kappa} \|\Delta_{12}g_m\|_{s+1}^{q,\kappa}. \end{split}$$

Combining this estimate with (9.118), we finally obtain for all $s \ge s_0$

$$\begin{split} \|\Delta_{12}u_m\|_s^{q,\kappa} &\leqslant \|\Pi_{N_m}^{\perp}\Delta_{12}f_m\|_s^{q,\kappa} + C\kappa^{-1}N_m^{\tau_1q+\tau_1+1}\|\Delta_{12}f_m\|_s^{q,\kappa} \max_{k\in\{1,2\}} \|f_m^{[k]}\|_{s_0}^{q,\kappa} \\ &+ C\kappa^{-1}N_m^{\tau_1q+\tau_1+1}\|\Delta_{12}f_m\|_{s_0}^{q,\kappa} \max_{k\in\{1,2\}} \|f_m^{[k]}\|_s^{q,\kappa} \end{split}$$

$$+ C\kappa^{-1} N_m^{\tau_1 q + 2\tau_1 + 1} \max_{k \in \{1,2\}} \|f_m^{[k]}\|_s^{q,\kappa} \max_{k \in \{1,2\}} \|f_m^{[k]}\|_{s_0}^{q,\kappa} \|\Delta_{12} V_m\|^{q,\kappa}$$

For $s = s_0$ we can actually get a better estimate,

$$\begin{split} \|\Delta_{12}u_m\|_{s_0}^{q,\kappa} &\leqslant \|\Pi_{N_m}^{\perp}\Delta_{12}f_m\|_{s_0}^{q,\kappa} + C\kappa^{-1}N_m^{\tau_1q+\tau_1+1}\|\Delta_{12}f_m\|_{s_0}^{q,\kappa}\max_{k\in\{1,2\}}\|f_m^{[k]}\|_{s_0}^{q,\kappa} \\ &+ C\kappa^{-1}N_m^{\tau_1q+2\tau_1+1}\max_{k\in\{1,2\}}\|f_m^{[k]}\|_{s_0}^{q,\kappa}\max_{k\in\{1,2\}}\|f_m^{[k]}\|_{s_0}^{q,\kappa}\|\Delta_{12}V_m\|^{q,\kappa}. \end{split}$$

By (9.61), Lemma 4.2 and (9.121), using in particular that $s_0 + \tau_1 q + 2\tau_1 + 1 \leq \overline{s}_h$, we deduce

$$\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa} \leqslant N_{m}^{s_{0}-\overline{s}_{h}-\mathbf{a}}\|\Delta_{12}f_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} + CN_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+\tau_{1}+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} + CN_{m}^{\tau_{1}q+2\tau_{1}+1}N_{0}^{2\overline{\mu}_{2}}N_{m}^{-2\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}V_{m}\|^{q,\kappa}$$
(9.123)

and

$$\begin{split} \|\Delta_{12}u_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} &\leqslant \|\Delta_{12}f_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \left(1 + CN_{0}^{\mu_{2}}N_{m}^{\tau_{1}q+\tau_{1}+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\right) + CN_{m}^{\tau_{1}q+\tau_{1}+1}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} \\ &+ CN_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+2\tau_{1}+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}V_{m}\|^{q,\kappa} \end{split}$$

where

$$\delta_0^{[1,2]}(s) \triangleq \kappa^{-1} \max_{k \in \{1,2\}} \|f_0^{[k]}\|_s^{q,\kappa}.$$

Hence, inserting the estimate (9.123) into (9.122) implies

$$\|(\mathbf{B}_{m}^{-1})^{[1]}\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa} \leq CN_{m}^{\tau_{1}q+\tau_{1}}\|\Delta_{12}u_{m}\|_{s_{0}}^{q,\kappa} \\ \leq N_{m}^{s_{0}-\overline{s}_{h}+\tau_{1}q+\tau_{1}}\|\Delta_{12}f_{m}\|_{\overline{s}_{h}}^{q,\kappa} + CN_{0}^{\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} \\ + CN_{0}^{2\overline{\mu}_{2}}N_{m}^{-2\overline{\mu}_{2}+\tau_{1}q+\tau_{1}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}V_{m}\|^{q,\kappa}.$$
(9.124)

Similarly and after straightforward computations we get

$$\|(\mathbf{B}_{m}^{-1})^{[1]}\Delta_{12}u_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \leqslant \|\Delta_{12}f_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \left(1 + N_{m}^{s_{0}+\tau_{1}q+\tau_{1}-\overline{s}_{h}} + N_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+\tau_{1}+1-\mu_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\right) + C\left(N_{0}^{\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-\overline{\mu}_{2}} + N_{m}^{\tau_{1}q+\tau_{1}+1}\right)\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} + CN_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+\tau_{1}+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h})\|\Delta_{12}V_{m}\|^{q,\kappa}.$$

$$(9.125)$$

Next we intend to estimate $(\Delta_{12}\mathbf{B}_m^{-1})u_m^{[2]}$. For this aim we write according to Taylor formula,

$$(\Delta_{12}\mathbf{B}_m^{-1})u_m^{[2]}(\theta) = \Delta_{12}\widehat{g}_m(\theta) \int_0^1 \left(\partial_\theta u_m^{[2]}\right) \left(\theta + \widehat{g}_m^{[2]}(\theta) + t\Delta_{12}\widehat{g}_m(\theta)\right) dt.$$

Coming back to (9.119) and using Lemma 4.1, (9.58) and (9.61), we have for all $s \ge s_0$

$$\begin{aligned} \|u_{m}^{[2]}\|_{s}^{q,\kappa} \leq \|\Pi_{N_{m}}^{\perp}f_{m}^{[2]}\|_{s}^{q,\kappa} + C\|f_{m}^{[2]}\|_{s}^{q,\kappa}\|\partial_{\theta}g_{m}^{[2]}\|_{s_{0}}^{q,\kappa} + C\|f_{m}^{[2]}\|_{s_{0}}^{q,\kappa}\|\partial_{\theta}g_{m}^{[2]}\|_{s}^{q,\kappa} \\ \leq \|f_{m}^{[2]}\|_{q,s} \left(1 + C\kappa^{-1}\|f_{m}^{[2]}\|_{s_{0}+\tau_{1}q+\tau_{1}+1}^{q,\kappa}\right) \\ \leq C\|f_{m}^{[2]}\|_{s}^{q,\kappa}. \end{aligned}$$

$$(9.126)$$

Applying Lemma 4.1 and Lemma 9.4, we obtain for all $s \geqslant s_0$

$$\begin{aligned} \| (\Delta_{12} \mathbf{B}_m^{-1}) u_m^{[2]} \|_s^{q,\kappa} \leqslant C \| \Delta_{12} \widehat{g}_m \|_s^{q,\kappa} \| u_m^{[2]} \|_{s_0+1}^{q,\kappa} \left(1 + \| \widehat{g}_m^{[2]} \|_{s_0}^{q,\kappa} + \| \Delta_{12} \widehat{g}_m \|_{s_0}^{q,\kappa} \right) \\ &+ C \| \Delta_{12} \widehat{g}_m \|_{s_0}^{q,\kappa} \| u_m^{[2]} \|_{s+1}^{q,\kappa} \left(1 + \| \widehat{g}_m^{[2]} \|_{s_0}^{q,\kappa} + \| \Delta_{12} \widehat{g}_m \|_{s_0}^{q,\kappa} \right) \\ &+ C \| \Delta_{12} \widehat{g}_m \|_{s_0}^{q,\kappa} \| u_m^{[2]} \|_{s_0+1}^{q,\kappa} \left(\| \widehat{g}_m^{[2]} \|_s^{q,\kappa} + \| \Delta_{12} \widehat{g}_m \|_s^{q,\kappa} \right). \end{aligned}$$

From (9.37), (9.111) and Sobolev embeddings, one obtains for all $s \in [s_0, \overline{s}_h + a]$

$$\|\Delta_{12}\widehat{g}_{m}\|_{s}^{q,\kappa} \leq C\left(\|\Delta_{12}g_{m}\|_{s}^{q,\kappa} + \|\Delta_{12}g_{m}\|_{s_{0}}^{q,\kappa}\max_{k\in\{1,2\}}\|g_{m}^{[k]}\|_{s+1}^{q,\kappa}\right)$$
$$\leq C\|\Delta_{12}g_{m}\|_{s}^{q,\kappa}.$$

Combining the last two estimates with (9.110) and Lemma 9.4 we deduce that for all $s \in [s_0, \overline{s}_h + \mathbf{a}]$

$$\|(\Delta_{12}\mathbf{B}_m^{-1})u_m^{[2]}\|_s^{q,\kappa} \leqslant C \|\Delta_{12}g_m\|_s^{q,\kappa} \|u_m^{[2]}\|_{q,s_0+1} + C \|\Delta_{12}g_m\|_{s_0}^{q,\kappa} \|u_m^{[2]}\|_{s+1}^{q,\kappa}.$$

Using (9.118), (9.126), we obtain for all $s \in [s_0, \overline{s}_h + a]$

$$\begin{aligned} \|(\Delta_{12}\mathbf{B}_{m}^{-1})u_{m}^{[2]}\|_{s}^{q,\kappa} \leqslant C\kappa^{-1}N_{m}^{\tau_{1}q+\tau_{1}}\|\Delta_{12}f_{m}\|_{s}^{q,\kappa} \max_{k\in\{1,2\}} \|f_{m}^{[k]}\|_{s_{0}+1}^{q,\kappa} \\ &+ C\kappa^{-1}N_{m}^{\tau_{1}q+\tau_{1}}\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} \max_{k\in\{1,2\}} \|f_{m}^{[k]}\|_{s+1}^{q,\kappa} \\ &+ C\kappa^{-1}N_{m}^{\tau_{1}q+2\tau_{1}}\|\Delta_{12}V_{m}\|_{s}^{q,\kappa} \max_{k\in\{1,2\}} \|f_{m}^{[k]}\|_{s_{0}+1}^{q,\kappa} \max_{k\in\{1,2\}} \|f_{m}^{[k]}\|_{s+1}^{q,\kappa}. \end{aligned}$$

Hence, by (9.61), (9.85) and (9.121), we have

$$\begin{aligned} \| (\Delta_{12} \mathbf{B}_m^{-1}) u_m^{[2]} \|_{s_0}^{q,\kappa} &\leq C N_0^{\overline{\mu}_2} N_m^{\tau_1 q + \tau_1 - \overline{\mu}_2} \delta_0^{[1,2]} (\overline{s}_h + 1) \| \Delta_{12} f_m \|_{s_0}^{q,\kappa} \\ &+ N_0^{2\overline{\mu}_2} N_m^{\tau_1 q + 2\tau_1 - 2\overline{\mu}_2} \delta_0^{[1,2]} (\overline{s}_h + 1) \| \Delta_{12} V_m \|^{q,\kappa} \end{aligned}$$
(9.127)

and

$$\begin{aligned} \| (\Delta_{12} \mathbf{B}_{m}^{-1}) u_{m}^{[2]} \|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} &\leq C N_{0}^{\overline{\mu}_{2}} N_{m}^{\tau_{1}q+\tau_{1}-\overline{\mu}_{2}} \delta_{0}^{[1,2]} (\overline{s}_{h}+1) \| \Delta_{12} f_{m} \|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \\ &+ C N_{m}^{\tau_{1}q+\tau_{1}} \delta_{0}^{[1,2]} (\overline{s}_{h}+\mathbf{a}+1) \| \Delta_{12} f_{m} \|_{s_{0}}^{q,\kappa} \\ &+ N_{0}^{\overline{\mu}_{2}} N_{m}^{\tau_{1}q+2\tau_{1}-\overline{\mu}_{2}} \delta_{0}^{[1,2]} (\overline{s}_{h}+\mathbf{a}+1) \| \Delta_{12} V_{m} \|^{q,\kappa}. \end{aligned}$$
(9.128)

Putting together (9.120), (9.124) and (9.127) and using Sobolev embeddings yield

$$\begin{split} \|\Delta_{12}f_{m+1}\|_{s_{0}}^{q,\kappa} \leqslant N_{m}^{s_{0}-\overline{s}_{h}+\tau_{1}q+\tau_{1}} \|\Delta_{12}f_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \\ &+ CN_{0}^{\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h}+1)\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} \\ &+ CN_{0}^{2\overline{\mu}_{2}}N_{m}^{-2\overline{\mu}_{2}+\tau_{1}q+2\tau_{1}}\delta_{0}^{[1,2]}(\overline{s}_{h}+1)\|\Delta_{12}V_{m}\|^{q,\kappa}. \end{split}$$
(9.129)

Similarly, we obtain according to (9.120), (9.125) and (9.128)

$$\begin{split} \|\Delta_{12}f_{m+1}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \leqslant \|\Delta_{12}f_{m}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \left(1 + N_{m}^{s_{0}+\tau_{1}q+\tau_{1}-\overline{s}_{h}} + N_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+\tau_{1}+1-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h}+1)\right) \\ &+ C\left(N_{0}^{\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-\overline{\mu}_{2}} + N_{m}^{\tau_{1}q+\tau_{1}+1}\right)\delta_{0}^{[1,2]}(\overline{s}_{h}+\mathbf{a}+1)\|\Delta_{12}f_{m}\|_{s_{0}}^{q,\kappa} \quad (9.130) \\ &+ N_{0}^{\overline{\mu}_{2}}N_{m}^{\tau_{1}q+2\tau_{1}-\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h}+\mathbf{a}+1)\|\Delta_{12}V_{m}\|^{q,\kappa}. \end{split}$$

Next, we introduce the quantities

$$\overline{\delta}_m(s) \triangleq \kappa^{-1} \|\Delta_{12} f_m\|_s^{q,\kappa} \text{ and } \varkappa_m \triangleq \kappa^{-1} \|\Delta_{12} V_m\|^{q,\kappa}.$$

According to (9.73) one deduces that

$$\Delta_{12}V_{m+1} = \Delta_{12}V_m + \langle \Delta_{12}f_m \rangle_{\varphi,\theta} \quad \text{and} \quad \Delta_{12}V_0 = 0,$$

which implies in view of Sobolev embeddings

$$\varkappa_m \leqslant C \sum_{k=0}^{m-1} \overline{\delta}_k(s_0). \tag{9.131}$$

The goal now is to prove by induction that for any $0 \leq a \leq s_h - \overline{s}_h + \sigma_1 - 5$

$$\forall k \leqslant m, \quad \overline{\delta}_k(s_0) \leqslant N_0^{\overline{\mu}_2} N_k^{-\overline{\mu}_2} \nu(\overline{s}_h + \mathbf{a}) \quad \text{and} \quad \overline{\delta}_k(\overline{s}_h + \mathbf{a}) \leqslant \left(2 - \frac{1}{k+1}\right) \nu(\overline{s}_h + \mathbf{a}) \tag{9.132}$$

with

$$\nu(s) \triangleq \overline{\delta}_0(s) + \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{s_0+4}.$$

Notice that the property (9.132) is obvious for m = 0 due to Sobolev embeddings. Let us now assume that (9.132) is true at the order m and let us check it at the order m + 1. By hypothesis of induction (9.132) and (9.131), one has

$$\varkappa_m \leqslant C\nu(\overline{s}_h + \mathbf{a}),\tag{9.133}$$

with C independent of m. By (9.129) and hypothesis of induction (9.132), we can write

$$\begin{split} \overline{\delta}_{m+1}(s_0) &\leqslant N_m^{s_0 - \overline{s}_h + \tau_1 q + \tau_1} \overline{\delta}_m(\overline{s}_h + \mathbf{a}) + C N_0^{\overline{\mu}_2} N_m^{2(\tau_1 q + \tau_1) + 1 - \overline{\mu}_2} \delta_0^{[1,2]}(\overline{s}_h + 1) \overline{\delta}_m(s_0) \\ &+ C N_0^{2\overline{\mu}_2} N_m^{-2\overline{\mu}_2 + \tau_1 q + 2\tau_1} \delta_0^{[1,2]}(\overline{s}_h + 1) \varkappa_m \\ &\leqslant \left[2 N_m^{s_0 - \overline{s}_h + \tau_1 q + \tau_1} + C N_0^{2\overline{\mu}_2} N_m^{2(\tau_1 q + \tau_1) + 1 - 2\overline{\mu}_2} \delta_0^{[1,2]}(\overline{s}_h + 1) \right] \nu(\overline{s}_h + \mathbf{a}). \end{split}$$

From the constraints on \overline{s}_h and $\overline{\mu}_2$ fixed by (9.38) and (6.32) we get

$$2N_m^{s_0-\overline{s}_h+\tau_1q+\tau_1} = 2N_m^{-\frac{3}{2}\overline{\mu}_2-2} = 2N_m^{-2}N_{m+1}^{-\overline{\mu}_2}$$
$$\leqslant 2N_0^{-2}N_{m+1}^{-\overline{\mu}_2}$$
$$\leqslant \frac{1}{2}N_0^{\overline{\mu}_2}N_{m+1}^{-\overline{\mu}_2}.$$

The last inequality occurs provided that

$$4 \leqslant N_0^{\overline{\mu}_2 + 2}$$

which is true since $N_0 \ge 2$ and $\overline{\mu}_2 \ge 2$. Similarly, one gets from (9.65) and (9.38)

$$CN_{0}^{2\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-2\overline{\mu}_{2}}\delta_{0}^{[1,2]}(\overline{s}_{h}+1) \leqslant C\varepsilon\kappa^{-1}N_{0}^{\overline{\mu}_{2}}N_{m}^{2(\tau_{1}q+\tau_{1})+1-\frac{1}{2}\overline{\mu}_{2}}N_{0}^{\overline{\mu}_{2}}N_{m+1}^{-\overline{\mu}_{2}}$$
$$\leqslant C\varepsilon\kappa^{-1}N_{0}^{2(\tau_{1}q+\tau_{1})+1+\frac{1}{2}\overline{\mu}_{2}}N_{0}^{\overline{\mu}_{2}}N_{m+1}^{-\overline{\mu}_{2}}$$
$$\leqslant C\varepsilon\kappa^{-1}N_{0}^{\overline{\mu}_{2}}N_{0}^{-\overline{\mu}_{2}}N_{m+1}^{-\overline{\mu}_{2}}.$$

Therefore, from (9.39) and by taking ε_0 small enough we infer

$$CN_0^{2\overline{\mu}_2}N_m^{2(\tau_1q+\tau_1)+1-2\overline{\mu}_2}\delta_0^{[1,2]}(\overline{s}_h+1) \leqslant \frac{1}{2}N_0^{\overline{\mu}_2}N_{m+1}^{-\overline{\mu}_2}$$

Putting together the preceding estimates we find that

$$\overline{\delta}_{m+1}(s_0) \leqslant N_0^{\overline{\mu}_2} N_{m+1}^{-\overline{\mu}_2} \nu(\overline{s}_h + \mathbf{a}).$$

This achieves the induction of the first statement in (9.132). Now, let us move to the second one. Then we can write by virtue of (9.130), the first statement of (9.132), (9.65) and (9.39)

$$\begin{split} \overline{\delta}_{m+1}(\overline{s}_h + \mathbf{a}) \leqslant &\overline{\delta}_m(\overline{s}_h + \mathbf{a}) \left(1 + N_m^{s_0 + \tau_1 q + \tau_1 - \overline{s}_h} + N_0^{\overline{\mu}_2} N_m^{\tau_1 q + \tau_1 + 1 - \overline{\mu}_2} \right) \\ &+ C \left(N_0^{\overline{\mu}_2} N_m^{2(\tau_1 q + \tau_1) + 1 - \overline{\mu}_2} + N_m^{\tau_1 q + \tau_1 + 1} \right) N_0^{\overline{\mu}_2} N_m^{-\overline{\mu}_2} \varepsilon \kappa^{-1} \nu(\overline{s}_h + \mathbf{a}). \end{split}$$

Notice that we have used according to (9.65) and (9.39) that if $\overline{s}_h + a + 5 \leq s_h + \sigma_1$

$$\delta_0^{[1,2]}(\overline{s}_h + \mathbf{a} + 1) \lesssim \varepsilon \kappa^{-1} \left(1 + \max_{k=1,2} \| \mathfrak{I}_k \|_{\overline{s}_h + \mathbf{a} + 5}^{q,\kappa} \right) \\ \lesssim \varepsilon \kappa^{-1}.$$

By hypothesis of induction (9.132), we obtain

$$\overline{\delta}_{m+1}(\overline{s}_h + \mathbf{a}) \leqslant \left(2 - \frac{1}{m+1}\right) \left(1 + N_m^{s_0 + \tau_1 q + \tau_1 - \overline{s}_h} + N_0^{\overline{\mu}_2} N_m^{\tau_1 q + \tau_1 + 1 - \overline{\mu}_2}\right) \nu(\overline{s}_h + \mathbf{a}) \\ + C \left(N_0^{\overline{\mu}_2} N_m^{2(\tau_1 q + \tau_1) + 1 - \overline{\mu}_2} + N_m^{\tau_1 q + \tau_1 + 1}\right) N_0^{\overline{\mu}_2} N_m^{-\overline{\mu}_2} \varepsilon \kappa^{-1} \nu(\overline{s}_h + \mathbf{a}).$$

In the same way to (9.82) we conclude in view of (9.39)

$$\left(2 - \frac{1}{m+1}\right) \left(1 + N_m^{s_0 + \tau_1 q + \tau_1 - \overline{s}_h} + N_0^{\overline{\mu}_2} N_m^{\tau_1 q + \tau_1 + 1 - \overline{\mu}_2}\right) + C \left(N_0^{\overline{\mu}_2} N_m^{2(\tau_1 q + \tau_1) + 1 - \overline{\mu}_2} + N_m^{\tau_1 q + \tau_1 + 1}\right) N_0^{\overline{\mu}_2} N_m^{-\overline{\mu}_2} \varepsilon \kappa^{-1} \leqslant \left(2 - \frac{1}{m+2}\right).$$

Consequently, we find

$$\overline{\delta}_{m+1}(\overline{s}_h + \mathbf{a}) \leqslant \left(2 - \frac{1}{m+2}\right)\nu(\overline{s}_h + \mathbf{a})$$

which achieves the proof of the second statement in (9.132) at the order m + 1.

Estimate on $\Delta_{12}c_{i_0}$. According to (9.99) and since V_0 is independent of r then

$$\Delta_{12}c_{i_0} = \sum_{m=0}^{+\infty} \Delta_{12}(V_{m+1} - V_m)$$

By (9.73), Sobolev embeddings and implementing (9.132) with a = 0

$$\begin{split} \|\Delta_{12}(V_{m+1} - V_m)\|^{q,\kappa} &= \|\langle \Delta_{12} f_m \rangle_{\varphi,\theta} \|^{q,\kappa} \\ &\leq C\kappa \overline{\delta}_m(s_0) \\ &\leq C\kappa N_0^{\overline{\mu}_2} N_m^{-\overline{\mu}_2} \nu(\overline{s}_h) \end{split}$$

It follows by the preceding estimates combined with Lemma A.2 and (9.137)

$$\begin{aligned} \|\Delta_{12}c_{i_0}\|^{q,\kappa} &\leqslant \sum_{m=0}^{\infty} \|\Delta_{12}(V_{m+1} - V_m)\|^{q,\kappa} \\ &\leqslant C\kappa \,\nu(\overline{s}_h) N_0^{\underline{\mu_2}} \sum_{m=0}^{+\infty} N_m^{-\underline{\mu_2}} \\ &\leqslant C\varepsilon \|\Delta_{12}i\|_{\overline{s}_h+4}^{q,\kappa}. \end{aligned}$$

• Estimates of $\Delta_{12}\beta$ and $\Delta_{12}\hat{\beta}$. Now, inserting the estimates of (9.132) into (9.118), taken with $s = s_0$, we obtain

$$\|\Delta_{12}g_m\|_{s_0}^{q,\kappa} \lesssim \overline{\delta}_m(s_0 + \tau_1 q + \tau_1) + \varkappa_m \delta_m(s_0 + \tau_1 q + 2\tau_1 + 1)$$

Applying Sobolev embeddings, Lemma 4.3, (9.61) and (9.132) we get for some $\overline{\vartheta} \in (0, 1)$

$$\overline{\delta}_m(s_0 + \tau_1 q + \tau_1) \leqslant \overline{\delta}_m(s_0 + \tau_1 q + 2\tau_1 + 1)$$
$$\lesssim \overline{\delta}_m(s_0)^{\overline{\vartheta}} \overline{\delta}_m(\overline{s}_h)^{1 - \overline{\vartheta}}$$
$$\lesssim N_0^{\overline{\vartheta}\mu_2} N_m^{-\overline{\vartheta}\mu_2} \nu(\overline{s}_h)$$

and

$$\begin{split} \delta_m(s_0 + \tau_1 q + 2\tau_1 + 1) &\lesssim \delta_m(s_0)^{\overline{\vartheta}} \delta_m(\overline{s}_h)^{1 - \overline{\vartheta}} \\ &\lesssim N_0^{\overline{\vartheta}\overline{\mu}_2} N_m^{-\overline{\vartheta}\overline{\mu}_2} \delta_0(\overline{s}_h) \\ &\lesssim N_0^{\overline{\vartheta}\overline{\mu}_2} N_m^{-\overline{\vartheta}\overline{\mu}_2}. \end{split}$$

Thus we deduce from the preceding estimates and (9.133)

$$\|\Delta_{12}g_m\|_{s_0}^{q,\kappa} \lesssim N_0^{\overline{\vartheta}\overline{\mu}_2} N_m^{-\overline{\vartheta}\overline{\mu}_2} \nu(\overline{s}_h).$$
(9.134)

Now from (9.118) we infer

$$\|\Delta_{12}g_m\|_{\overline{s}_h+\mathsf{a}+1}^{q,\kappa} \lesssim \overline{\delta}_m(\overline{s}_h+\mathsf{a}+\tau_1q+\tau_1+1) + \varkappa_m\delta_m(\overline{s}_h+\mathsf{a}+\tau_1q+2\tau_1+2).$$
(9.135)

Coming back to (9.132) we infer that under the assumption

$$\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 1 \leqslant s_h + \sigma_1 - 5 \tag{9.136}$$

one gets

$$\begin{split} \overline{\delta}_m(\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 1) &\leqslant 2\nu(\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 1) \\ &\leqslant 2\overline{\delta}_0(\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 1) + 2\varepsilon\kappa^{-1} \|\Delta_{12}i\|_{s_0+4}^{q,\kappa}. \end{split}$$

Similarly to (9.65), one gets from Lemma 9.1-(iii) and Proposition 9.1-(i)

$$\forall s \ge s_0, \quad \overline{\delta}_0(s) = \kappa^{-1} \| \Delta_{12} V_{\varepsilon r} \|_s^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-1} \Big(\| \Delta_{12} i \|_{s+4}^{q,\kappa} + \| \Delta_{12} i \|_{s_0+4}^{q,\kappa} \max_{\ell=1,2} \| r_\ell \|_{s+4}^{q,\kappa}. \Big).$$

$$(9.137)$$

Then under the assumption

$$\overline{s}_h + \mathbf{a} + \tau_1 q + 2\tau_1 + 6 \leqslant s_h + \sigma_1 \tag{9.138}$$

combined with (9.39) and (9.137) we find

$$\overline{\delta}_0(\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 1) \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h + \mathbf{a} + \tau_1 q + \tau_1 + 5}^{q,\kappa}$$

From (9.85), (9.39) and (9.138) we obtain the following estimate

$$\delta_m((\overline{s}_h + \mathbf{a} + \tau_1 q + 2\tau_1 + 2) \leqslant C\varepsilon\kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{\overline{s}_h + \mathbf{a} + \tau_1 q + 2\tau_1 + 6}^{q,\kappa}\right)$$
$$\leqslant C\varepsilon\kappa^{-1}.$$

Plugging the preceding estimates into (9.135) and using (9.133) yield

$$\|\Delta_{12}g_m\|_{\overline{s}_h+\mathbf{a}+1}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+5}^{q,\kappa}.$$
(9.139)

provided that (9.138) is satisfied. By virtue of Lemma 4.3, (9.139) and (9.134) we obtain for some $\overline{\theta} \in (0, 1)$

$$\|\Delta_{12}g_m\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{\overline{\theta}\underline{\mu}_2} N_m^{-\overline{\theta}\underline{\mu}_2} \|\Delta_{12}i\|_{\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+5}^{q,\kappa}.$$
(9.140)

Consequently, we deduce from (9.140) and Lemma A.2

$$\sum_{m=0}^{\infty} \|\Delta_{12}g_m\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+5}^{q,\kappa} N_0^{\overline{\theta}\mu_2} \sum_{m=0}^{\infty} N_m^{-\overline{\theta}\mu_2} \\ \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+5}^{q,\kappa}.$$
(9.141)

Finally, putting together (9.109), (9.116), (9.140) and (9.141), we obtain

$$\|\Delta_{12}\beta\|_{\overline{s}_h+\mathbf{a}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{q,\overline{s}_h+\mathbf{a}+\tau_1q+\tau_1+5}$$

Thus, combining this estimate with (9.37), (9.112) and Sobolev embeddings we find

$$\begin{split} \|\Delta_{12}\hat{\beta}\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} &\lesssim \|\Delta_{12}\beta\|_{\overline{s}_{h}+\mathbf{a}}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{q,\overline{s}_{h}+\mathbf{a}+\tau_{1}q+\tau_{1}+5} \end{split}$$

The proof of Proposition 9.2 is now achieved.

9.4. First conjugation of the linearized operator. The main goal of this section is to explore the conjugation of the operator $\mathcal{L}_{r,\lambda}$ introduced in Lemma 9.1 using the symplectic change of coordinates \mathscr{B} constructed in Proposition 9.2. Before stating our main result, we need to prove the following technical lemma.

Lemma 9.5. Let $\hat{\beta}$ the function constructed in Proposition 9.2, then we have the decomposition

$$\frac{\left|\sin\left(\frac{\theta-\eta}{2}\right)\right|^{\alpha}}{\left|\sin\left(\frac{\theta-\eta+\widehat{\beta}(\varphi,\theta)-\widehat{\beta}(\varphi,\eta)}{2}\right)\right|^{\alpha}} = \frac{1}{2}\left(\left(1+\partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{-\alpha}+\left(1+\partial_{\eta}\widehat{\beta}(\varphi,\eta)\right)^{-\alpha}\right)\right)$$
$$+\sin^{2}\left(\frac{\theta-\eta}{2}\right)\mathscr{K}_{r,\lambda}(\varphi,\theta,\eta),$$

with the following estimate

$$\forall s \in [s_0, S], \quad \sup_{\eta \in \mathbb{T}} \|\mathscr{K}_{r,*}(\cdot, \cdot, \cdot + \eta)\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+4}^{q,\kappa}\right),$$

where the notation * stands for λ , \cdot for φ and \cdot denotes θ . The number σ_1 is given in (9.38).

Proof. We shall start with the following expression which follows from standard trigonometric identities

$$f(\varphi, \theta, \eta) \triangleq \frac{\sin\left(\frac{\theta - \eta + \hat{\beta}(\varphi, \theta) - \hat{\beta}(\varphi, \eta)}{2}\right)}{\sin\left(\frac{\theta - \eta}{2}\right)}$$
$$= \cos\left(\frac{\hat{\beta}(\varphi, \theta) - \hat{\beta}(\varphi, \eta)}{2}\right) + \frac{\sin\left(\frac{\hat{\beta}(\varphi, \theta) - \hat{\beta}(\varphi, \eta)}{2}\right)}{\sin\left(\frac{\theta - \eta}{2}\right)}\cos\left(\frac{\theta - \eta}{2}\right). \tag{9.142}$$

Applying the mean value theorem combined with the periodicity of $\widehat{\beta}$ we infer

$$\frac{\sin\left(\frac{\widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)}{2}\right)}{\sin\left(\frac{\theta - \eta}{2}\right)} \leqslant \frac{\left|\widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)\right|}{2\left|\sin\left(\frac{\theta - \eta}{2}\right)\right|} \leqslant \|\widehat{\beta}\|_{\operatorname{Lip}(\mathbb{T}^{d+1})}.$$

Putting together (9.42) and (9.39) combined with Sobolev embeddings yield

$$\begin{aligned} \|\widehat{\beta}\|_{s_0+1}^{q,\kappa} &\leq C \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s_0+1+\sigma_1}^{q,\kappa}\right) \\ &\leq C \varepsilon \kappa^{-1} \leq C \varepsilon_0, \end{aligned} \tag{9.143}$$

which implies in turn from Sobolev embeddings that $\|\widehat{\beta}\|_{\operatorname{Lip}(\mathbb{T}^{d+1})}$ is small enough. Consequently we deduce from (9.142) that f is strictly positive. Next, we shall use the splitting $f = f_1 + f_2 f_3$ with

$$f_1(\varphi,\theta,\eta) = \cos\left(\frac{\widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)}{2}\right), \quad f_2(\varphi,\theta,\eta) = \frac{\sin\left(\frac{\beta(\varphi,\theta) - \beta(\varphi,\eta)}{2}\right)}{\frac{\widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)}{2}},$$
$$f_3(\varphi,\theta,\eta) = \frac{\widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)}{2\tan\left(\frac{\theta-\eta}{2}\right)}.$$

Then using Lemma 4.4 and (9.42) we obtain for any $s \in [s_0, S]$,

$$\sup_{\eta \in \mathbb{T}} \|f_1(\cdot, \cdot, \cdot + \eta) - 1\|_s^{q,\kappa} \lesssim \|\widehat{\beta}\|_s^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1}^{q,\kappa}\right).$$

Since $x \mapsto \frac{\sin x}{x}$ is \mathscr{C}^{∞} then we get in a similar way to f_1

$$\sup_{\eta \in \mathbb{T}} \|f_2(\cdot, \cdot, \cdot + \eta) - 1\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1}^{q,\kappa}\right).$$

As to the term f_3 we simply apply Lemma 4.7-(i) in order to get in view of Proposition 9.2-(i)

$$\sup_{\eta \in \mathbb{T}} \|f_3(\cdot, \cdot, \cdot + \eta)\|_s^{q,\kappa} \lesssim \|\widehat{\beta}\|_{s+1}^{q,\kappa}$$
$$\lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right).$$

Using the law products stated in Lemma 4.1 combined with (9.39) we deduce that

$$\sup_{\eta \in \mathbb{T}} \|f(\cdot, \bullet, \bullet + \eta) - 1\|_{s}^{q,\kappa} \lesssim \sup_{\eta \in \mathbb{T}} \left[\|f_{1}(\cdot, \bullet, \bullet + \eta) - 1\|_{s}^{q,\kappa} + \|f_{3}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} \right] \\ + \sup_{\eta \in \mathbb{T}} \left[\|f_{2}(\cdot, \bullet, \bullet + \eta) - 1\|_{s_{0}}^{q,\kappa} \|f_{3}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} \right] \\ + \sup_{\eta \in \mathbb{T}} \left[\|f_{2}(\cdot, \bullet, \bullet + \eta) - 1\|_{s}^{q,\kappa} \|f_{3}\|_{s_{0}}^{q,\kappa} \right] \\ \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+1}^{q,\kappa} \right).$$
(9.144)

In particular we get in view of the smallness condition (9.39)

$$\sup_{\eta \in \mathbb{T}} \|f(\cdot, \cdot, \cdot + \eta) - 1\|_{s_0}^{q,\kappa} \leq C \varepsilon \kappa^{-1} \leq C \varepsilon_0.$$

This implies that f is close to 1 when ε_0 is small enough and therefore we may apply Lemma 4.4 to deduce that $f^{-\alpha}$ satisfies similar estimates, that is,

$$\sup_{\eta \in \mathbb{T}} \|f^{-\alpha}(\cdot, \cdot, \cdot + \eta) - 1\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right)$$
(9.145)

Arguing similarly we also get $(\varphi, \theta, \eta) \mapsto f^{-\alpha}(\varphi, \theta, \eta) \in W^{q,\infty}(\mathcal{O}; H^s(\mathbb{T}^{d+2}))$ with

$$\|f^{-\alpha} - 1\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right).$$

$$(9.146)$$

According to (9.142) one easily gets that the function f is symmetric, that is, $f(\varphi, \theta, \eta) = f(\varphi, \eta, \theta)$. Then using Lemma 4.6-(ii) combined with (9.146) we find

$$f^{-\alpha}(\varphi,\theta,\eta) = \frac{1}{2} \left(f^{-\alpha}(\varphi,\theta,\theta) + f^{-\alpha}(\varphi,\eta,\eta) \right) + \sin^2 \left(\frac{\theta-\eta}{2} \right) \mathscr{K}_{r,\lambda}(\varphi,\theta,\eta), \tag{9.147}$$

with the estimate

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{K}_{r,\lambda}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \lesssim \|\Delta_{\theta,\eta}(f^{-\alpha})\|_{s+\frac{1}{2}+\epsilon}^{q,\kappa} \lesssim \|f^{-\alpha} - 1\|_{s+\frac{5}{2}+\epsilon}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\Im_{0}\|_{s+\sigma_{1}+4}^{q,\kappa}\right)$$
(9.148)

On the other hand, we infer from (9.142)

$$f(\varphi, \theta, \theta) = 1 + \partial_{\theta} \widehat{\beta}(\varphi, \theta) \text{ and } f(\varphi, \eta, \eta) = 1 + \partial_{\eta} \widehat{\beta}(\varphi, \eta).$$

Consequently, we find

$$f^{-\alpha}(\varphi,\theta,\eta) = \frac{1}{2} \left(\left(1 + \partial_{\theta} \widehat{\beta}(\varphi,\theta) \right)^{-\alpha} + \left(1 + \partial_{\eta} \widehat{\beta}(\varphi,\eta) \right)^{-\alpha} \right) + \sin^2 \left(\frac{\theta - \eta}{2} \right) \mathscr{K}_{r,\lambda}(\varphi,\theta,\eta)$$

and the proof is now achieved.

Next, we plan to describe the action of the transformation \mathscr{B} on the operator $\mathscr{L}_{r,\lambda}$ introduced in Lemma 9.1 and derive some useful analytical estimates.

Proposition 9.3. Under the same assumptions and notations of Proposition 9.2, one gets for any $n \in \mathbb{N}$ and for any $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\mathscr{L}^{0}_{r,\lambda} \triangleq \mathscr{B}^{-1}\mathscr{L}_{\varepsilon r,\lambda}\mathscr{B} = \omega \cdot \partial_{\varphi} + c(\lambda, i_{0})\partial_{\theta} - \partial_{\theta} \Big(\mu_{r,\lambda} |\mathbf{D}|^{\alpha-1} + \mathbf{D}|^{\alpha-1}\mu_{r,\lambda}\Big) + \partial_{\theta}\mathscr{R}_{r,\lambda} + \mathbf{E}^{0}_{n}, \quad (9.149)$$

where $\mathscr{L}_{r,\lambda}$ was defined in (9.24)-(9.13)-(9.5), \mathscr{B} , $c_{r,\lambda}$ and \mathbb{E}_n^0 satisfy the same estimates of Proposition 9.2. In addition, $\mu_r \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\text{even}})$ for any $s \in [s_0, S]$ with the estimate

$$\forall s \in [s_0, S], \quad \|\mu_{r,\lambda} - 2^{-\alpha - 1} C_\alpha\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right)$$

and the remainder $\partial_{\theta} \mathcal{R}_{r,\lambda}$ is a reversible integral operator with,

$$\forall s \ge s_0, \forall \gamma \in \mathbb{N} \quad with \quad s + \gamma \leqslant S, \quad |||\partial_\theta \mathscr{R}_{r,\lambda}|||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + ||\mathfrak{I}_0||_{s+\sigma_1+7+\gamma}^{q,\kappa}\right).$$

Moreover, there exists $\overline{\sigma}_1 = \sigma(\tau_1, d, q) \ge \sigma_1$ such that for any $\gamma \in \mathbb{N}, s \ge 0$ with $s + \gamma \leqslant \overline{s}_h$

$$\|\Delta_{12}\mu_r\|_{\overline{s}_h}^{q,\kappa} + \|\Delta_{12}\partial_\theta \mathcal{R}_r\|_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\overline{\sigma}_1}^{q,\kappa}$$

Proof. Applying Proposition 9.2 and conjugating by \mathscr{B} the operator $\mathscr{L}_{\varepsilon r,\alpha}$ described in (9.24) we deduce that

$$\mathscr{B}^{-1}\mathscr{L}_{\varepsilon r,\lambda}\mathscr{B} = \omega \cdot \partial_{\varphi} + c(\lambda, i_0)\partial_{\theta} - \mathscr{B}^{-1}\partial_{\theta}\mathscr{T}_r\mathscr{B} + \mathscr{B}^{-1}\partial_{\theta}\mathscr{R}_{\varepsilon r,\lambda}\mathscr{B}, \qquad (9.150)$$

where \mathcal{T}_r denotes the self-adjoint operator

$$\mathcal{T}_r = \mathscr{W}_{\varepsilon r,\alpha} |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} \mathscr{W}_{\varepsilon r,\alpha}$$

and $\mathcal{W}_{r,\alpha}$ is defined in (9.13). From the definition of the fractional Laplacian (4.19) we may write

$$\mathcal{T}_r h(\varphi, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{\mathscr{W}_{\varepsilon r, \alpha}(\varphi, \theta) + \mathscr{W}_{\varepsilon r, \alpha}(\varphi, \eta)}{|\sin(\frac{\eta - \theta}{2})|^{\alpha}} h(\varphi, \eta) d\eta.$$
(9.151)

Applying Lemma 9.3-(i) we infer

$$\mathscr{B}^{-1}\partial_{\theta}\mathscr{T}_{r}\mathscr{B} = \partial_{\theta}\mathbf{B}^{-1}\mathscr{T}_{r}\mathscr{B}$$
$$\triangleq \partial_{\theta}\mathscr{T}_{r}^{1}. \tag{9.152}$$

Then using (9.34) and (9.151) together with the change of variable $\eta \sim \eta + \hat{\beta}(\varphi, \eta)$ we get

$$\mathcal{T}_{r}^{1}h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) h(\varphi,\eta)}{\left|\sin(\frac{\theta-\eta+\widehat{\beta}(\varphi,\theta)-\widehat{\beta}(\varphi,\eta)}{2})\right|^{\alpha}} d\eta$$
(9.153)

where

$$\mathscr{W}^{1}_{r,\alpha}(\varphi,\theta,\eta) \triangleq \mathscr{W}^{0}_{r,\alpha}(\varphi,\theta) + \mathscr{W}^{0}_{r,\alpha}(\varphi,\eta)$$

$$\mathscr{W}^{0}_{r,\alpha}(\varphi,\theta) \triangleq \mathscr{W}_{\varepsilon r,\alpha}(\varphi,\theta+\widehat{\beta}(\varphi,\theta))$$
(9.154)

and $\mathscr{W}_{r,\alpha}$ was introduced in (9.13). To estimate $\mathscr{W}_{r,\alpha}^0$ we first note that

$$\mathscr{W}^0_{0,\alpha}(\varphi,\theta) = C_{\alpha} 2^{-\alpha-1} = \mathscr{W}_{0,\alpha}$$

which is independent of φ and θ . Then applying Lemma 9.4 leads

$$\|\mathscr{W}_{r,\alpha}^{0} - \mathscr{W}_{0,\alpha}^{0}\|_{s}^{q,\kappa} \lesssim \|\widehat{\beta}\|_{s_{0}+1}^{q,\kappa} \|\mathscr{W}_{\varepsilon r,\alpha} - \mathscr{W}_{0,\alpha}\|_{s+1}^{q,\kappa} + \|\widehat{\beta}\|_{s+1}^{q,\kappa} \|\mathscr{W}_{\varepsilon r,\alpha} - \mathscr{W}_{0,\alpha}\|_{s_{0}+1}^{q,\kappa}$$

From Lemma 9.1-(i) combined with Proposition 9.2-(ii) we infer

$$\begin{aligned} \|\mathscr{W}_{r,\alpha}^{0} - C_{\alpha} 2^{-\alpha - 1} \|_{s}^{q,\kappa} &\lesssim \varepsilon \, \|\widehat{\beta}\|_{s_{0}+1}^{q,\kappa} \|r\|_{s+2}^{q,\kappa} + \varepsilon \|\widehat{\beta}\|_{s+1}^{q,\kappa} \|r\|_{s_{0}+2}^{q,\kappa} \\ &\lesssim \varepsilon^{2} \, \kappa^{-1} \left(1 + \|\Im_{0}\|_{s_{0}+\sigma_{1}+1}^{q,\kappa}\right) \, \|r\|_{s+2}^{q,\kappa} + \, \varepsilon^{2} \, \kappa^{-1} \left(1 + \|\Im_{0}\|_{s+\sigma_{1}+1}^{q,\kappa}\right) \, \|r\|_{s_{0}+2}^{q,\kappa}. \end{aligned}$$

Therefore, we deduce from Proposition 9.1-(i), Sobolev embeddings and (9.39)

$$\|\mathscr{W}_{r,\alpha}^{0} - C_{\alpha} 2^{-\alpha-1}\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+1}^{q,\kappa}\right).$$

$$(9.155)$$

Putting together (9.155) with (9.154) yields for any $s \in [s_0, S]$,

$$\|\mathscr{W}_{r,\alpha}^{1} - C_{\alpha} 2^{-\alpha}\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+1}^{q,\kappa}\right).$$
(9.156)

Notice that the Sobolev norm H^s in the preceding quantity concerns all the variables φ, θ, η . By inserting the identity of Lemma 9.5 into (9.153) yields

$$\begin{aligned} \mathcal{T}_{r}^{1}h(\varphi,\theta) =& \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{W}_{r,\alpha}^{2}(\varphi,\theta,\eta)}{\left|\sin\left(\frac{\theta-\eta}{2}\right)\right|^{\alpha}} h(\varphi,\eta) d\eta \\ &+ \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) \,\mathscr{K}_{r,\lambda}(\varphi,\theta,\eta)}{\left|\sin\left(\frac{\theta-\eta}{2}\right)\right|^{\alpha-2}} h(\varphi,\eta) d\eta, \end{aligned} \tag{9.157}$$

with

$$\mathscr{W}_{r,\alpha}^{2}(\varphi,\theta,\eta) = \frac{1}{2} \mathscr{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) \left(\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta) \right)^{-\alpha} + \left(1 + \partial_{\eta}\widehat{\beta}(\varphi,\eta) \right)^{-\alpha} \right).$$
(9.158)

Observe that the function $\mathscr{W}_{r,\alpha}^2$ is smooth and symmetric, $\mathscr{W}_{r,\alpha}^2(\varphi,\theta,\eta) = \mathscr{W}_{r,\alpha}^2(\varphi,\eta,\theta)$, then we apply Lemma 4.6 leading to

$$\mathscr{W}_{r,\alpha}^{2}(\varphi,\theta,\eta) = \frac{1}{2} \left(\mathscr{W}_{r,\alpha}^{2}(\varphi,\theta,\theta) + \mathscr{W}_{r,\alpha}^{2}(\varphi,\eta,\eta) \right) + \sin^{2} \left(\frac{\theta-\eta}{2} \right) \mathscr{K}_{r,\lambda}^{1}(\varphi,\theta,\eta),$$
(9.159)

with the estimate

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} \mathscr{K}^1_{r,\lambda})(\cdot, \cdot, \cdot + \eta) \|_s^{q,\kappa} \lesssim \| \nabla_{\theta,\eta} \mathscr{W}^2_{r,\alpha} \|_{s+\frac{3}{2}+p_1+p_2+\epsilon}^{q,\kappa}.$$
(9.160)

By writing (9.158) in the form

$$\mathcal{W}_{r,\alpha}^{2}(\varphi,\theta,\eta) = \frac{1}{2} \mathcal{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) \left(\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{-\alpha} + \left(1 + \partial_{\eta}\widehat{\beta}(\varphi,\eta)\right)^{-\alpha} - 2 \right) \\ + \left(\mathcal{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) - C_{\alpha}2^{-\alpha}\right) + C_{\alpha}2^{-\alpha}$$

and applying Lemmata 4.1 and 4.4 combined with (9.156) we find under the condition (9.39),

$$\begin{split} \|\nabla_{\theta,\eta}\mathscr{W}_{r,\alpha}^{2}\|_{s+p_{1}+p_{2}+\frac{3}{2}+\epsilon}^{q,\kappa} &\lesssim \|\mathscr{W}_{r,\alpha}^{2} - C_{\alpha}2^{-\alpha}\|_{s+\frac{5}{2}+p_{1}+p_{2}+\epsilon}^{q,\kappa} \\ &\lesssim \|\mathscr{W}_{r,\alpha}^{1}\|_{s+p_{1}+p_{2}+\frac{5}{2}+\epsilon}^{q,\kappa}\|\widehat{\beta}\|_{s_{0}+1}^{q,\kappa} + \|\mathscr{W}_{r,\alpha}^{1}\|_{s_{0}}^{q,\kappa}\|\widehat{\beta}\|_{s+p_{1}+p_{2}+\frac{7}{2}+\epsilon}^{q,\kappa} \\ &+ \varepsilon\kappa^{-1}\left(1 + \|\Im_{0}\|_{s+p_{1}+p_{2}+\frac{5}{2}+\epsilon+\sigma_{1}}^{q,\kappa}\right). \end{split}$$

From (9.156), Proposition 9.2-(ii) and (9.39) we infer after straightforward manipulations

$$\|\nabla_{\theta,\eta}\mathscr{W}_{r,\alpha}^{2}\|_{s+p_{1}+p_{2}+\frac{3}{2}+\epsilon}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} (1+\|\mathfrak{I}_{0}\|_{s+p_{1}+p_{2}+\frac{7}{2}+\epsilon+\sigma}^{q,\kappa}).$$
(9.161)

Inserting (9.161) into (9.160) yields in view of Sobolev embeddings

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2}) \mathscr{K}^1_{r,\lambda}(\cdot, \cdot, \cdot + \eta) \|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+p_1+p_2+\frac{7}{2}+\epsilon+\sigma_1}^{q,\kappa} \right) \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+p_1+p_2+4+\sigma_1}^{q,\kappa} \right).$$
(9.162)

On the other hand, using (9.154) and (9.158) we get

$$\mu_{r,\lambda}(\varphi,\theta) \triangleq \frac{1}{2} \mathscr{W}_{r,\alpha}^{2}(\varphi,\theta,\theta)$$
$$= \frac{\mathscr{W}_{\varepsilon r,\alpha}(\varphi,\theta+\widehat{\beta}(\varphi,\theta))}{\left(1+\partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{\alpha}} \cdot$$
(9.163)

Plugging (9.159) and (9.163) into (9.157) implies

$$\begin{aligned} \mathscr{T}_{r}^{1}h(\varphi,\theta) =& \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mu_{r,\lambda}(\varphi,\theta) + \mu_{r,\lambda}(\varphi,\eta)}{\left|\sin(\frac{\theta-\eta}{2})\right|^{\alpha}} h(\varphi,\eta) d\eta \\ &+ \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{K}_{r,\lambda}^{2}(\varphi,\theta,\eta)}{\left|\sin(\frac{\theta-\eta}{2})\right|^{\alpha-2}} h(\varphi,\eta) d\eta, \end{aligned}$$
(9.164)

with

$$\mathscr{K}_{r,\lambda}^{2}(\varphi,\theta,\eta) \triangleq \mathscr{K}_{r,\lambda}^{1}(\varphi,\theta,\eta) + \mathscr{W}_{r,\alpha}^{1}(\varphi,\theta,\eta) \mathscr{K}_{r,\lambda}(\varphi,\theta,\eta)$$
(9.165)

and where $\mathscr{K}_{r,\lambda}$ is defined in (9.147), $\mathscr{K}_{r,\lambda}^1$ in (9.159) and $\mathscr{W}_{r,\alpha}$ in (9.154). To estimate $\mathscr{K}_{r,\lambda}^2$ we use Lemma 4.1

$$\begin{aligned} \|\mathscr{K}_{r,\lambda}^{2}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} &\lesssim \|\mathscr{K}_{r,\lambda}^{1}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} + \|\mathscr{W}_{r,\alpha}^{1}(\cdot, \bullet, \bullet + \eta)\|_{s_{0}}^{q,\kappa} \|\mathscr{K}_{r,\lambda}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} \\ &+ \|\mathscr{W}_{r,\alpha}^{1}(\cdot, \bullet, \bullet + \eta)\|_{s}^{q,\kappa} \|\mathscr{K}_{r,\lambda}(\cdot, \bullet, \bullet + \eta)\|_{s_{0}}^{q,\kappa}. \end{aligned}$$
(9.166)

Then (9.166), (9.162), (9.148), (9.156) combined with (9.39) imply

$$\sup_{\eta \in \mathbb{T}} \|\mathscr{K}_{r,\lambda}^{2}(\cdot, \cdot, \cdot + \eta)\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+4+\sigma_{1}}^{q,\kappa}\right).$$

$$(9.167)$$

Similarly we also get for any $p_1, o_2 \in \mathbb{N}$

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} \mathscr{K}^2_{r,\lambda})(\cdot, \cdot, \cdot + \eta) \|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+p_1+p_2+4+\sigma_1}^{q,\kappa} \right).$$
(9.168)

Coming back to (9.164) and using the definition (4.19) we deduce that

$$\mathcal{T}_{r}^{1}h(\varphi,\theta) = \left(\mu_{r,\lambda}|\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1}\mu_{r,\lambda}\right)h(\varphi,\theta) + \frac{1}{2\pi}\int_{0}^{2\pi}\frac{\mathscr{K}_{r,\lambda}^{2}(\varphi,\theta,\eta)}{\left|\sin(\frac{\theta-\eta}{2})\right|^{\alpha-2}}h(\varphi,\eta)d\eta$$
$$\triangleq \left(\mu_{r,\lambda}|\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1}\mu_{r,\lambda}\right)h(\varphi,\theta) + \mathscr{R}_{r}^{0,1}h(\varphi,\theta). \tag{9.169}$$

Next, we intend to estimate the functions $\mu_{r,\lambda}$ defined in (9.163). Using (9.13) we get

$$\mu_{0,\lambda}(\varphi,\theta) = \frac{\mathscr{W}_{0,\alpha}(\varphi,\theta)}{\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{\alpha}} = C_{\alpha}2^{-\alpha-1}\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{-\alpha}.$$

In addition, we obtain from (9.163) and (9.154)

$$\mu_{r,\lambda}(\varphi,\theta) - \mathscr{W}_{0,\alpha} = \frac{\mathscr{W}_{r,\alpha}^{0}(\varphi,\theta) - \mathscr{W}_{0,\alpha}^{0}}{\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{\alpha}} + \mathscr{W}_{0,\alpha}\left(\left(1 + \partial_{\theta}\widehat{\beta}(\varphi,\theta)\right)^{-\alpha} - 1\right)$$
$$\triangleq \mathscr{G}_{1} + \mathscr{W}_{0,\alpha}\mathscr{G}_{2}.$$

Now, applying Lemmata 4.1 and 4.4 yields under the smallness condition (9.39),

$$\begin{aligned} \|\mathscr{I}_1\|_s^{q,\kappa} &\lesssim \left(1 + \|\partial_\theta \widehat{\beta}\|_{s_0}^{q,\kappa}\right) \left\|\mathscr{W}_{r,\alpha}^0 - \mathscr{W}_{0,\alpha}^0\right\|_s^{q,\kappa} \\ &+ \left(1 + \|\partial_\theta \widehat{\beta}\|_s^{q,\kappa}\right) \left\|\mathscr{W}_{r,\alpha}^0 - \mathscr{W}_{0,\alpha}^0\right\|_{s_0}^{q,\kappa}. \end{aligned}$$

Therefore we deduce from (9.155), (9.39) and Proposition 9.2-(ii)

$$\|\mathscr{I}_1\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right).$$

$$(9.170)$$

As to the term $\mathscr{W}_{0,\alpha}\mathscr{I}_2$ we shall combine Lemmata 4.1, 4.4 with Proposition 9.2-(ii)

$$\begin{aligned} \|\mathscr{W}_{0,\alpha}\mathscr{I}_2\|_s^{q,\kappa} &\lesssim \|\partial_\theta \widehat{\beta}\|_s^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right) \end{aligned}$$

It follows from the preceding estimates that

$$\|\mu_{r,\lambda} - \mathscr{W}_{0,\alpha}\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+1}^{q,\kappa}\right).$$

Putting together (9.150), (9.152) and (9.169) we get

$$\mathscr{B}^{-1}\mathscr{L}_{\varepsilon r,\lambda}\mathscr{B} = \omega \cdot \partial_{\varphi} + c_{r,\lambda}\partial_{\theta} - \partial_{\theta} \left(\mu_{r,\lambda} |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \mu_{r,\lambda} \right) + \partial_{\theta}\mathscr{R}_{r},$$

with

$$\partial_{\theta} \mathcal{R}_{r} \triangleq -\partial_{\theta} \mathcal{R}_{r}^{0,1} + \mathcal{B}^{-1} \partial_{\theta} \mathcal{R}_{\varepsilon r,\lambda} \mathcal{B}$$
$$= \partial_{\theta} (-\mathcal{R}_{r}^{0,1} + \mathbf{B}^{-1} \mathcal{R}_{\varepsilon r,\lambda} \mathcal{B}).$$
(9.171)

We shall start with estimating $\partial_{\theta} \mathscr{R}_{r}^{0,1}$. Coming back to the definition (9.169) and differentiating in the variable θ we get after straightforward computations

$$\partial_{\theta} \mathcal{R}_{r}^{0,1} h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\eta-\theta}{2}) \mathscr{K}_{r,\lambda}^{3}(\varphi,\theta,\eta)}{\left|\sin(\frac{\theta-\eta}{2})\right|^{\alpha}} h(\varphi,\eta) d\eta, \qquad (9.172)$$

with

e.

$$\mathscr{K}_{r,\lambda}^{3}(\varphi,\theta,\eta) = \frac{\alpha-1}{2} \cos\left((\eta-\theta)/2\right) \mathscr{K}_{r,\lambda}^{2}(\varphi,\theta,\eta) + \sin\left((\eta-\theta)/2\right) \partial_{\theta} \mathscr{K}_{r,\lambda}^{2}(\varphi,\theta,\eta).$$
(9.173)

Making the change of variable $\eta \rightsquigarrow \eta + \theta$ allows to get

$$\partial_{\theta} \mathcal{R}_{r}^{0,1} h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\eta}{2}) \mathscr{K}_{r,\lambda}^{3}(\varphi,\theta,\theta+\eta)}{\left|\sin(\frac{\eta}{2})\right|^{\alpha}} h(\varphi,\theta+\eta) d\eta.$$
(9.174)

From Lemma 4.1 combined with (9.167) we infer

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} \mathscr{K}^3_{r,\lambda})(\cdot, \cdot, \cdot + \eta) \|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+5+\sigma_1}^{q,\kappa} \right).$$
(9.175)

Next, we move to the estimate of the symbol associated to $\partial_{\theta} \mathcal{R}_{r}^{0,1}$ whose expression can be derived from (7.8) and takes the form

$$\sigma_{\partial_{\theta}\mathcal{R}_{r}^{0,1}}(\varphi,\theta,\xi) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\sin(\frac{\eta}{2})\mathscr{K}_{r,\lambda}^{3}(\varphi,\theta,\theta+\eta)}{\left|\sin(\frac{\eta}{2})\right|^{\alpha}} e^{i\eta\xi} d\eta$$

Then integration by parts combined with (7.6) yield

$$\xi^{1+\gamma} \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathcal{R}_{r}^{0,1}}(\varphi,\theta,\xi) = \frac{i^{1+\gamma}}{2\pi} \int_{0}^{2\pi} e^{i\eta\xi} \partial_{\eta}^{1+\gamma} \left[\frac{\sin(\frac{\eta}{2})(e^{i\eta}-1)^{\gamma}}{|\sin(\frac{\eta}{2})|^{\alpha}} \mathscr{K}_{r,\lambda}^{3}(\varphi,\theta,\theta+\eta) \right] d\eta$$

Therefore

$$\langle \xi \rangle^{1+\gamma} \left\| \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathcal{R}_{r}^{0,1}}(\cdot, \boldsymbol{\cdot}, \xi) \right\|_{H^{s}} \lesssim \sum_{0 \leqslant \gamma' \leqslant 1+\gamma} \int_{0}^{2\pi} \frac{\|\partial_{\eta}^{\gamma'} \mathscr{K}_{r,\lambda}^{3}(\cdot, \boldsymbol{\cdot}, \boldsymbol{\cdot} + \eta)\|_{H^{s}}}{|\sin(\eta/2)|^{\alpha}} \, d\eta.$$

Similarly we get for any $|\beta| \leq q$ and $\alpha \in (0, \overline{\alpha})$, using Leibniz formula and Sobolev embeddings,

$$\begin{split} \kappa^{|\beta|} \langle \xi \rangle^{1+\gamma} \big\| \partial_{\lambda}^{\beta} \Delta_{\xi}^{\gamma} \sigma_{\partial_{\theta} \mathcal{R}_{r}^{0,1}}(\cdot, \boldsymbol{\cdot}, \xi) \big\|_{H^{s-|\beta|}} &\lesssim \kappa^{|\beta|} \sum_{\substack{0 \leqslant \gamma' \leqslant 1+\gamma \\ 0 \leqslant \beta' \leqslant \beta}} \int_{0}^{2\pi} \frac{\|\partial_{\lambda}^{\beta'} \partial_{\eta}^{\gamma'} \mathscr{K}_{r,\lambda}^{3}(\cdot, \boldsymbol{\cdot}, \boldsymbol{\cdot} + \eta)\|_{H^{s-|\beta'|}}}{|\sin(\eta/2)|^{\frac{1}{2}}} \, d\eta \\ &\lesssim \sum_{0 \leqslant \gamma' \leqslant 1+\gamma} \int_{0}^{2\pi} \frac{\|\partial_{\eta}^{\gamma'} \mathscr{K}_{r,\lambda}^{3}(\cdot, \boldsymbol{\cdot}, \boldsymbol{\cdot} + \eta)\|_{s}^{q,\kappa}}{|\sin(\eta/2)|^{\frac{1}{2}}} \, d\eta. \end{split}$$

Plugging (9.175) into the preceding estimate and using (7.13) allow to get

$$\||\partial_{\theta} \mathcal{R}_{r}^{0,1}\||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\gamma+\sigma_{1}+6}^{q,\kappa}\right).$$

$$(9.176)$$

Let us now move to the estimate of $\mathscr{B}_r^{-1}\partial_\theta \mathscr{R}_{r,\alpha} \mathscr{B}_r$. Then using (9.21) and Lemma 9.3-(i) we find

$$\mathcal{B}^{-1}\partial_{\theta}\mathcal{R}_{\varepsilon r,\lambda}\mathcal{B} = \partial_{\theta}\mathbf{B}^{-1}\mathcal{R}_{\varepsilon r,\lambda}\mathcal{B}$$
$$\triangleq \partial_{\theta}\mathcal{S}_{r}, \qquad (9.177)$$

with

$$\mathscr{S}_{r}h(\varphi,\theta) = \frac{1}{2\pi} \int_{0}^{2\pi} \frac{\mathscr{A}_{r,\alpha}\left(\varphi,\theta + \widehat{\beta}(\varphi,\theta),\eta + \widehat{\beta}(\varphi,\eta)\right)}{\left|\sin(\frac{\theta - \eta + \widehat{\beta}(\varphi,\theta) - \widehat{\beta}(\varphi,\eta)}{2})\right|^{\alpha - 2}} h(\varphi,\eta) d\eta.$$
(9.178)

Applying Lemma 9.5, which remains true if we replace α with $\alpha - 2$, we get

$$\left| \sin\left(\frac{\theta - \eta + \widehat{\beta}(\varphi, \theta) - \widehat{\beta}(\varphi, \eta)}{2} \right) \right|^{2-\alpha} \triangleq \frac{F_1(\varphi, \theta, \eta)}{\left|\sin\left(\frac{\theta - \eta}{2}\right)\right|^{\alpha - 2}}.$$

The estimate of F_1 is similar to (9.156) and one gets

$$\|F_1\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+1}^{q,\kappa}\right).$$

$$(9.179)$$

The H^s norm of F_1 concerns all the variables. Let us introduce

$$F_2(\varphi,\theta,\eta) = F_1(\varphi,\theta,\eta) \mathscr{A}_{r,\alpha} \big(\varphi,\theta + \widehat{\beta}(\varphi,\theta),\eta + \widehat{\beta}(\varphi,\eta)\big).$$

Then similarly to the estimates (9.166) and (9.168) we deduce that

$$\sup_{\eta \in \mathbb{T}} \|F_2(\cdot, \cdot, \cdot + \eta)\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+4}^{q,\kappa}\right)$$
(9.180)

and

$$\sup_{\eta \in \mathbb{T}} \| (\partial_{\theta}^{p_1} \partial_{\eta}^{p_2} F_2)(\cdot, \cdot, \cdot + \eta) \|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma_1+p_1+p_2+4}^{q,\kappa} \right).$$
(9.181)

In addition, (9.178) becomes

$$\mathscr{S}_r h(\varphi, \theta) = \frac{1}{2\pi} \int_0^{2\pi} \frac{F_2(\varphi, \theta + \eta)}{\left|\sin(\frac{\eta}{2})\right|^{\alpha - 2}} h(\varphi, \theta + \eta) d\eta.$$
(9.182)

Now, we proceed as for (9.174) by combining (9.180), (9.181) and (9.182) in order to get

$$\left\|\left|\mathscr{G}_{r}\right|\right\|_{-2,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \left|\left|\Im_{0}\right|\right|_{s+\sigma_{1}+\gamma+6}^{q,\kappa}\right).$$

On the other hand one gets from (7.12) the following identity

$$\sigma_{\partial_{\theta}\mathcal{S}_r} = \partial_{\theta}\sigma_{\mathcal{S}_r} + i\xi\sigma_{\mathcal{S}_r}.$$

Consequently, we find

$$\begin{aligned} \|\partial_{\theta}\mathscr{G}_{r}\|_{-1,s,\gamma}^{q,\kappa} &\lesssim \||\mathscr{G}_{r}\||_{-2,s+1,\gamma}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+\gamma+7}^{q,\kappa}\right), \end{aligned}$$

which gives in view of (9.177)

$$\| \mathscr{B}^{-1} \partial_{\theta} \mathscr{R}_{\varepsilon r, \lambda} \mathscr{B} \|_{-1, s, \gamma}^{q, \kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s + \sigma_1 + \gamma + 7}^{q, \kappa} \right)$$

Next we shall move to the estimate of the differences. Let us start with $\Delta_{12}\mu_r$ where μ_r is defined in (9.163). The computations are slightly long but they are classical and resemble to the preceding estimates. Then based on (9.44), Lemma 9.4 combined with (9.39) we find

$$\|\Delta_{12}\mu_r\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa},$$

for some σ depending only on τ_1 , d and q. Similar arguments combined with the estimates used for getting (9.175) allow to get

$$\forall s \ge 0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant \overline{s}_h, \quad \sup_{\eta \in \mathbb{T}} \|\partial_{\eta}^{\gamma} \Delta_{12} \mathscr{K}^3_{r,\lambda}(\cdot, \cdot, \cdot + \eta)\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12} i\|_{\overline{s}_h + \sigma}^{q,\kappa}.$$

Consequently, we find in view of (9.172) and implementing similar arguments to (9.176)

$$\forall s \ge 0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant \overline{s}_h, \quad |||\Delta_{12} \partial_\theta \mathcal{R}_r^{0,1}|||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} ||\Delta_{12} i||_{\overline{s}_h + \sigma}^{q,\kappa}.$$

Coming back to (9.182) and implementing the same arguments we find

$$\forall s \ge 0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant \overline{s}_h \quad ||| \Delta_{12} \partial_\theta \mathscr{S}_r |||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} ||\Delta_{12}i||_{\overline{s}_h + \sigma}^{q,\kappa}.$$

Finally, we get from (9.171) and the preceding estimates

$$\forall s \ge 0, \gamma \in \mathbb{N} \quad \text{with} \quad s + \gamma \leqslant \overline{s}_h, \quad |||\Delta_{12}\partial_\theta \mathscr{R}_r|||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} ||\Delta_{12}i||_{\overline{s}_h+\sigma}^{q,\kappa} \tag{9.183}$$

This achieves the proof of Proposition 9.3.

9.5. Reduction of the nonlocal part. The main goal of this section is to reduce to constant coefficients the leading term in (9.149) with positive order. Notice that at this level the transport part is with constant coefficient and we need to find a judicious conjugation in order to reduce the nonlocal part to a Fourier multiplier with the same order, of course without altering the transport part. As we shall see below, this will be done using the infinite dimensional hyperbolic flow Φ described by (8.1) through a specific choice of the density ρ . We emphasize that this kind of reduction was used for water waves in [16]. Our main result can be stated as follows.

Proposition 9.4. There exists $\sigma_2 = \sigma(\tau_1, d, q) \ge \overline{\sigma}_1$ such that under (9.2) and (9.38) and by assuming

$$\|\mathfrak{I}_0\|_{q,s_h+\sigma_2} \leqslant 1 \quad \text{and} \quad N_0^{\mu_2} \varepsilon \kappa^{-2} \leqslant \varepsilon_0, \tag{9.184}$$

the following assertions hold true. There exists $\Psi : \mathfrak{O} \to \mathscr{L}(H^s(\mathbb{T}^{d+1}), H^s(\mathbb{T}^{d+1}))$ a family of reversibility preserving invertible linear bounded operators and such that for any $n \in \mathbb{N}$ and for any λ in the set $\mathfrak{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ defined in Proposition 9.2 we have

$$\Psi^{-1}\mathscr{L}_{\varepsilon r,\lambda}\Psi \triangleq \mathscr{L}_{r,\lambda}^{1} \triangleq \omega \cdot \partial_{\varphi} + c(\lambda, i_{0})\partial_{\theta} - \mathrm{m}(\lambda, i_{0})\partial_{\theta}|\mathrm{D}|^{\alpha-1} + \mathscr{R}_{r,\lambda}^{1} + \mathrm{E}_{n}^{1}, \qquad (9.185)$$

with the following properties.

(i) We have
$$\mathbf{m}(\lambda, i_0) = \frac{2}{(2\pi)^{d+1}} \int_{\mathbb{T}^{d+1}} \mu_{r,\lambda}(\varphi, \theta) d\varphi d\theta$$
 and
 $\|\mathbf{m}(\cdot, i_0) - 2^{-\alpha} C_{\alpha}\|^{q,\kappa} + \|c(\cdot, i_0) - V_{0,\alpha}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1}.$

Moreover

$$\|\Delta_{12}\mathbf{m}(\cdot,i)\|^{q,\kappa} + \|\Delta_{12}c(\cdot,i)\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h + \sigma_2}^{q,\kappa}$$

(ii) The operator $\mathscr{R}^1_{r,\lambda}$ is reversible and for any $s \in [s_0, S]$ and for $\epsilon > 0$ small enough

$$\|\|\mathscr{R}^{1}_{r,\lambda}\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa}+\||\mathscr{R}^{1\star}_{r,\lambda}\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa}\lesssim \varepsilon \kappa^{-2}\left(1+\|\mathfrak{I}_{0}\|_{s+\sigma_{2}}^{q,\kappa}\right)$$

In addition,

$$\|\Delta_{12}\mathscr{R}^{1}_{r,\lambda}h\|^{q,\kappa}_{\overline{s}_{h}} + \|\Delta_{12}\mathscr{R}^{1,\star}_{r,\lambda}h\|^{q,\kappa}_{\overline{s}_{h}} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|^{q,\kappa}_{q,\overline{s}_{h}+\sigma_{2}} \|h\|^{q,\kappa}_{\overline{s}_{h}+\sigma_{2}}$$

and

$$\||\Delta_{12}\mathscr{R}^1_{r,\lambda}|||_{0,0,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{q,\overline{s}_h+\sigma_2}^{q,\kappa}.$$

(iii) The operators $\Psi^{\pm 1}$ satisfy for any $s \ge s_0$

$$\left|\Psi^{\pm 1}h\right|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{2}}^{q,\kappa}\right) \|h\|_{s_{0}}^{q,\kappa}$$

and

$$\begin{split} \big\| (\Psi^{\pm 1} - \mathrm{Id}) h \big\|_{s}^{q,\kappa} + \big\| ((\Psi^{\pm 1})^{\star} - \mathrm{Id}) h \big\|_{s}^{q,\kappa} &\lesssim \varepsilon \kappa^{-2} \|h\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma_{2}}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa} \\ &\|\Delta_{12} \Psi^{\pm 1} h\|_{\overline{s}_{h}}^{q,\kappa} + \|\Delta_{12} (\Psi^{\pm 1})^{\star} h\|_{\overline{s}_{h}}^{q,\kappa} &\lesssim \varepsilon \kappa^{-2} \|\Delta_{12} i\|_{\overline{s}_{h}+\sigma_{2}}^{q,\kappa} \|h\|_{\overline{s}_{h}+\sigma_{2}}^{q,\kappa} . \end{split}$$

(iv) The operator \mathbf{E}_n^1 is reversible and satisfies the estimate

$$\forall s \in [s_0, S], \quad \left\| \mathsf{E}_n^1 h \right\|_{s_0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \Big(N_0^{\mu_2} N_{n+1}^{-\mu_2} + N_n^{s_0-s} \big(1 + \|\mathfrak{I}_0\|_{s+\sigma_2}^{q,\kappa} \big) \Big) \|h\|_{s_0+3}^{q,\kappa}.$$

Proof. We have seen in Proposition 9.3 that for any $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\mathcal{L}_{r,\lambda}^{0} = \mathscr{B}^{-1} \mathcal{L}_{\varepsilon r,\lambda} \mathscr{B} = \omega \cdot \partial_{\varphi} + c(\lambda, i_{0}) \partial_{\theta} - \partial_{\theta} \mathscr{A} + \partial_{\theta} \mathscr{R}_{r,\lambda} + \mathsf{E}_{n}^{0},$$

with $\mathscr{A} \triangleq \mu_{r,\lambda} |\mathsf{D}|^{\alpha - 1} + |\mathsf{D}|^{\alpha - 1} \mu_{r,\lambda}.$ (9.186)

Next, we shall conjugate this operator by the flow $\Phi(t)$ described in Proposition 8.1

$$\begin{aligned} \mathcal{L}_{r,\lambda}^{t} &\triangleq \Phi^{-1}(t) \mathcal{L}_{r,\lambda}^{0} \Phi(t), \quad \text{with} \quad \Phi(t) = e^{t\mathbb{A}}, \\ &= \Psi^{-1}(t) \mathcal{L}_{\varepsilon r,\lambda} \Psi(t), \quad \text{with} \quad \Psi(t) = \mathscr{B} \Phi(t), \end{aligned}$$
(9.187)

where the generator A takes the form

$$\mathbb{A} = \partial_{\theta} \mathcal{T} \quad \text{and} \quad \mathcal{T} = \rho |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \rho, \tag{9.188}$$

with ρ being a smooth function to be chosen later. Then Taylor formula implies

$$\mathscr{L}^{1}_{r,\lambda} = \mathscr{L}^{0}_{r,\lambda} + \int_{0}^{1} \Phi^{-1}(t) [\mathscr{L}^{0}_{r,\lambda}, \mathbb{A}] \Phi(t) dt.$$
(9.189)

Let us evaluate the commutator $[\mathscr{L}^{0}_{r,\lambda},\mathbb{A}]$ using the precise structure of \mathbb{A} . By direct computations we may check that

$$\left[\omega \cdot \partial_{\varphi}, \mathbb{A}\right] = \partial_{\theta} \left(\left(\omega \cdot \partial_{\varphi} \rho \right) |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \left(\omega \cdot \partial_{\varphi} \rho \right) \right)$$

and

$$\left[c(\lambda, i_0)\partial_{\theta}, \mathbb{A}\right] = \partial_{\theta} \Big(\big(c(\lambda, i_0)\partial_{\theta}\rho\big) |\mathsf{D}|^{\alpha-1} + |\mathsf{D}|^{\alpha-1} \big(c(\lambda, i_0)\partial_{\theta}\rho\big) \Big).$$

Consequently,

$$\begin{bmatrix} \omega \cdot \partial_{\varphi} + c(\lambda, i_0) \partial_{\theta}, \mathbb{A} \end{bmatrix} = \partial_{\theta} \left(\widehat{\rho} \, |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \widehat{\rho} \right) \quad \text{with} \quad \widehat{\rho} \triangleq \omega \cdot \partial_{\varphi} \rho + c(\lambda, i_0) \partial_{\theta} \rho. \tag{9.190}$$

Define

$$\widehat{\mathbb{A}} = \partial_{\theta} \big(\widehat{\rho} \, |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \widehat{\rho} \big). \tag{9.191}$$

It follows that

$$\left[\mathscr{L}^{0}_{r,\lambda},\mathbb{A}\right] = \widehat{\mathbb{A}} + \left[-\partial_{\theta}\mathscr{A} + \partial_{\theta}\mathscr{R}_{r,\lambda},\mathbb{A}\right] + \left[\mathbb{E}^{0}_{n},\mathbb{A}\right].$$
(9.192)

Inserting (9.192) into (9.189) yields

$$\mathcal{L}_{r,\lambda}^{1} = \omega \cdot \partial_{\varphi} + c(\lambda, i_{0})\partial_{\theta} - \partial_{\theta}\mathcal{A} + \partial_{\theta}\mathcal{R}_{r,\lambda} + \int_{0}^{1} \Phi(-t)\widehat{\mathbb{A}}\Phi(t)dt + \int_{0}^{1} \Phi(-t) \Big[\partial_{\theta}\mathcal{R}_{r,\lambda} - \partial_{\theta}\mathcal{A}, \mathbb{A}\Big]\Phi(t)dt + \mathbf{E}_{n}^{0} + \int_{0}^{1} \Phi(-t) \big[\mathbf{E}_{n}^{0}, \mathbb{A}\big]\Phi(t)dt.$$
(9.193)

Then using once again Taylor formula we find

$$\Phi(-t)\widehat{\mathbb{A}}\Phi(t) = \widehat{\mathbb{A}} + \int_0^t \Phi(-t') [\widehat{\mathbb{A}}, \mathbb{A}] \Phi(t') dt'.$$

Plugging this identity into (9.193) allows to get

$$\mathcal{L}_{r,\lambda}^{1} = \omega \cdot \partial_{\varphi} + c(\lambda, i_{0})\partial_{\theta} + \left(\widehat{\mathbb{A}} - \partial_{\theta}\mathscr{A}\right) + \mathscr{R}_{r,\lambda}^{1}$$

$$+ \mathsf{E}_{n}^{0} + \int_{0}^{1} \Phi(-t) \big[\mathsf{E}_{n}^{0}, \mathbb{A}\big] \Phi(t) dt,$$
(9.194)

with

$$\mathcal{R}_{r,\lambda}^{1} \triangleq \partial_{\theta} \mathcal{R}_{r,\lambda} + \int_{0}^{1} \Phi(-t) \Big[\partial_{\theta} \mathcal{R}_{r,\lambda} - \partial_{\theta} \mathcal{A}, \mathbb{A} \Big] \Phi(t) dt + \int_{0}^{1} \int_{0}^{t} \Phi(-t') \Big[\widehat{\mathbb{A}}, \mathbb{A} \Big] \Phi(t') dt' dt \\ = \partial_{\theta} \mathcal{R}_{r,\lambda} + \int_{0}^{1} \Phi(-t) \Big[\partial_{\theta} \mathcal{R}_{r,\lambda} - \partial_{\theta} \mathcal{A}, \mathbb{A} \Big] \Phi(t) dt + \int_{0}^{1} (1-t) \Phi(-t) \Big[\widehat{\mathbb{A}}, \mathbb{A} \Big] \Phi(t) dt, \quad (9.195)$$

where we used in the last inequality an integration by parts. Combining (9.186) and (9.191) we may write

$$\widehat{\mathbb{A}} - \partial_{\theta} \mathscr{A} = \partial_{\theta} \Big[(\widehat{\rho} - \mu_{r,\lambda}) |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} (\widehat{\rho} - \mu_{r,\lambda}) \Big].$$
(9.196)

We recall that $\mu_{r,\lambda}$ was defined in (9.163) and in what follows $\langle \mu_{r,\lambda} \rangle_{\varphi,\theta}$ stands for its average in both variables. According to Lemma 9.2, the following functions

$$\rho \triangleq (\omega \cdot \partial_{\varphi} + c(\lambda, i_0) \partial_{\theta})_{\text{ext}}^{-1} (\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}) \quad \text{and} \quad \rho_n \triangleq \Pi_{N_n} \rho \tag{9.197}$$

are well defined in the whole set \mathcal{O} and moreover for any $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\left(\omega \cdot \partial_{\varphi} + c(\lambda, i_0)\partial_{\theta}\right)\rho_n = \prod_{N_n} \left(\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}\right).$$
(9.198)

Therefore, we deduce from the decomposition

$$\rho = \rho_n + \Pi_{N_n}^{\perp} \rho \tag{9.199}$$

and (9.198) that for any $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\left(\omega \cdot \partial_{\varphi} + c(\lambda, i_0) \partial_{\theta} \right) \rho = \Pi_{N_n} \left(\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta} \right) + \left(\omega \cdot \partial_{\varphi} + c(\lambda, i_0) \partial_{\theta} \right) \Pi_{N_n}^{\perp} \rho$$

$$\leq \Pi_{N_n} \left(\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta} \right) + \mathbf{R}_n.$$

$$(9.200)$$

Putting together (9.196), (9.190) and (9.200) yields for $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\widehat{\mathbb{A}} - \partial_{\theta} \mathscr{A} = -2 \langle \mu_{r,\lambda} \rangle_{\varphi,\theta} \partial_{\theta} |\mathbf{D}|^{\alpha-1} - \partial_{\theta} \left[(\Pi_{N_n}^{\perp} \mu_{r,\lambda}) |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} (\Pi_{N_n}^{\perp} \mu_{r,\lambda}) \right] + \partial_{\theta} \big(\mathbf{R}_n |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} \mathbf{R}_n \big).$$
(9.201)

Inserting (9.201) into (9.194) allows to get for any $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$

$$\mathcal{L}_{r,\lambda}^{1} = \Psi^{-1} \mathcal{L}_{\varepsilon r,\lambda} \Psi, \quad \text{with} \quad \Psi \triangleq \Psi(1) = \mathscr{B} \Phi(1)$$
$$= \omega \cdot \partial_{\varphi} + c(\lambda, i_{0}) \partial_{\theta} - \mathbf{m}(\lambda, i_{0}) \partial_{\theta} |\mathbf{D}|^{\alpha - 1} + \mathscr{R}_{r,\lambda}^{1} + \mathbf{E}_{n}^{1}, \tag{9.202}$$

with $\mathbf{m}(\lambda, i_0) \triangleq 2 \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}$ and

$$\mathbf{E}_{n}^{1} \triangleq \mathbf{E}_{n}^{0} + \int_{0}^{1} \Phi(-t) \left[\mathbf{E}_{n}^{0}, \mathbb{A} \right] \Phi(t) dt - \partial_{\theta} \left[(\Pi_{N_{n}}^{\perp} \mu_{r,\lambda}) |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} (\Pi_{N_{n}}^{\perp} \mu_{r,\lambda}) \right] \\ + \partial_{\theta} \left(\mathbf{R}_{n} |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} \mathbf{R}_{n} \right).$$
(9.203)

(i) The estimate of $c(\lambda, i_0)$ was done before in Proposition 9.2-(i). However the estimate of $\mu_{r,\lambda}$ is given in Proposition 9.3. Then the estimate of the average $m_{r,\lambda}$ follows easily from Sobolev embeddings and (9.39)

$$\begin{split} \left\| \mathbf{m}(\cdot, i_{0}) - 2^{-\alpha} C_{\alpha} \right\|^{q,\kappa} &= 2 \left\| \left\langle \mu_{r,\lambda} - 2^{-\alpha-1} C_{\alpha} \right\rangle_{\varphi,\theta} \right\|^{q,\kappa} \\ &\lesssim \left\| \mu_{r,\lambda} - 2^{-\alpha-1} C_{\alpha} \right\|_{s_{0}}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_{0} \|_{s_{0}+\sigma_{1}+1}^{q,\kappa} \right) \\ &\lesssim \varepsilon \kappa^{-1}. \end{split}$$

For the estimate of the difference $m_{r,\lambda}$ we write in view of Proposition 9.3

$$\begin{aligned} \left| \Delta_{12} \mathbf{m}(\cdot, i) \right\|^{q, \kappa} &= 2 \left\| \langle \Delta_{12} \mu_{r, \lambda} \rangle_{\varphi, \theta} \right\|^{q, \kappa} \\ &\lesssim \varepsilon \kappa^{-1} \left\| \Delta_{12} i \right\|_{\overline{s}_h + \overline{\sigma}_1}^{q, \kappa}. \end{aligned}$$

(ii) We shall start with estimating ρ given in (9.197). Then applying Lemma 9.2 gives

$$\|\rho\|_{s}^{q,\kappa} \lesssim \kappa^{-1} \left(1 + \|c(\cdot,i_{0})\|^{q,\kappa}\right) \|\mu_{r,\lambda} - \langle\mu_{r,\lambda}\rangle_{\varphi,\theta} \|_{s+\tau_{1}(q+1)}^{q,\kappa}.$$
(9.204)

Using (9.40) combined with (9.184) yield

$$\|c(\cdot, i_0)\|^{q,\kappa} \lesssim 1. \tag{9.205}$$

On the other hand, we infer from Proposition 9.3

$$\left\|\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle_{\varphi,\theta}\right\|_{s+\tau_1(q+1)}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma_1+\tau_1(1+q)+1}^{q,\kappa}\right)$$

Hence, combining this estimate with (9.205) and (9.204) we deduce that

$$|\rho||_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+\tau_{1}(1+q)+1}^{q,\kappa} \right).$$

$$(9.206)$$

Implementing similar estimates used before to get (9.118), by making appeal in particular to the estimates of $\Delta_{12}c_r$ and $\Delta_{12}\mu_r$ detailed in Proposition 9.3, one gets after tedious and long computations

$$\|\Delta_{12}\rho\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}, \tag{9.207}$$

for some $\sigma = \sigma(\tau_1, d, q)$. The next goal is to estimate $\mathscr{R}^1_{r,\lambda}$ whose expression is detailed in (9.195). To do that, we shall first use the splitting

$$\mathscr{R}^{1}_{r,\lambda} = \partial_{\theta} \mathscr{R}_{r,\lambda} + \widehat{\mathscr{R}}_{1} + \widehat{\mathscr{R}}_{2}, \qquad (9.208)$$

with

$$\widehat{\mathscr{R}}_{1} = \int_{0}^{1} \Phi(-t) \Big[\partial_{\theta} \mathscr{R}_{r,\lambda}, \mathbb{A} \Big] \Phi(t) dt, \qquad (9.209)$$

$$\widehat{\mathscr{R}}_{2} = \int_{0}^{1} \Phi(-t) \Big[\partial_{\theta} \mathscr{A} + (t-1) \widehat{\mathbb{A}}, \mathbb{A} \Big] \Phi(t) dt.$$

Recall from Proposition 9.3 that

$$\||\partial_{\theta}\mathscr{R}_{r,\lambda}\||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{1}+7+\gamma}^{q,\kappa}\right).$$

$$(9.210)$$

Applying Lemma 7.7-(iv) and using (9.188)

$$\begin{split} \| [\partial_{\theta} \mathcal{R}_{r,\lambda}, \mathbb{A}] \|_{-1,s,\gamma}^{q,\kappa} \lesssim \sum_{0 \leqslant \beta \leqslant \gamma} \| |\partial_{\theta} \mathcal{R}_{r,\lambda}| \|_{-1,s_{0}+2,1+\beta}^{q,\kappa} \| \rho \|_{s+4+\gamma-\beta}^{q,\kappa} + \| \partial_{\theta} \mathcal{R}_{r,\lambda}| \|_{-1,s+2,1+\beta}^{q,\kappa} \| \rho \|_{s_{0}+4+\gamma-\beta}^{q,\kappa} \\ + \sum_{0 \leqslant \beta \leqslant \gamma} \| |\partial_{\theta} \mathcal{R}_{r,\lambda}| \|_{-1,s_{0}+2+\beta,\gamma-\beta}^{q,\kappa} \| \rho \|_{s+1}^{q,\kappa} + \| \partial_{\theta} \mathcal{R}_{r,\lambda}| \|_{-1,s+2+\beta,\gamma-\beta}^{q,\kappa} \| \rho \|_{s_{0}+1}^{q,\kappa}. \end{split}$$

Then using (9.206) and (9.210) combined with Sobolev embeddings and (9.184), we may find an explicit $\sigma = \sigma(\tau_1, d, q)$ such that

$$\begin{split} \|\|[\partial_{\theta}\mathscr{R}_{r,\lambda},\mathbb{A}\|\|_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\Im_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\right) + \varepsilon \kappa^{-1} \sum_{0 \leqslant \beta \leqslant \gamma} \|\Im_{0}\|_{s_{0}+\sigma+\beta}^{q,\kappa} \|\Im_{0}\|_{s+\sigma+\gamma-\beta}^{q,\kappa} \\ &+ \varepsilon \kappa^{-1} \sum_{0 \leqslant \beta \leqslant \gamma} \|\Im_{0}\|_{s_{0}+\sigma+\gamma}^{q,\kappa} \|\Im_{0}\|_{s+\sigma}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \left(1 + \|\Im_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\right) + \varepsilon \kappa^{-1} \sum_{0 \leqslant \beta \leqslant \gamma} \|\Im_{0}\|_{s_{0}+\sigma+\beta}^{q,\kappa} \|\Im_{0}\|_{s+\sigma+\gamma-\beta}^{q,\kappa}. \end{split}$$

We point out that along the computations below the value of σ may change from line to line but still only depends on τ_1, d and q. Then using interpolation inequalities we find for any $0 \leq \beta \leq \gamma$ and $s \geq s_0$

$$\|\mathfrak{I}_0\|_{s_0+\sigma+\beta}^{q,\kappa}\|\mathfrak{I}_0\|_{s+\sigma+\gamma-\beta}^{q,\kappa} \lesssim \|\mathfrak{I}_0\|_{s_0+\sigma}^{q,\kappa}\|\mathfrak{I}_0\|_{s+\sigma+\gamma}^{q,\kappa}$$

Thus from the smallness condition (9.184) we infer for any $\gamma \in \mathbb{N}, s \ge s_0$

$$\|\|[\partial_{\theta}\mathcal{R}_{r,\lambda},\mathbb{A}]\|\|_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\right).$$

$$(9.211)$$

Applying Theorem 8.1-(ii) we get for any $s \ge s_0$ and for any $t \in [0, 1]$

$$\|\Phi(-t)[\partial_{\theta}\mathcal{R}_{r,\lambda},\mathbb{A}]\Phi(t)\|_{-1,s,0}^{q,\kappa} \leqslant Ce^{C\overline{\mu}^{s+1}(0)}\overline{\mu}(s)$$
(9.212)

with

$$\overline{\mu}(s) \triangleq \overline{F}_0(s+27+s_0) + \|\rho\|_{s+27+s_0}^{q,\kappa}$$

and

$$\overline{F}_0(s) \triangleq C \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma+4}^{q,\kappa} \right).$$

Notice that $s \mapsto \overline{F}_0(s)$ is log-convex since Sobolev norms enjoy this property. Using (9.206) combined with Sobolev embeddings we deduce

$$\overline{\mu}(s) \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \right)$$

and consequently we get from (9.212) and (9.184) that for any $t \in [0, 1]$

$$\||\Phi(-t)[\partial_{\theta}\mathscr{R}_{r,\lambda},\mathbb{A}]\Phi(t)|||_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} (1+||\mathfrak{I}_{0}||_{s+\sigma}^{q,\kappa}).$$

We remind that the value of σ may change from line to line but still depends only on τ_1 , d and q. Combining this estimate with the definition of $\widehat{\mathscr{R}}_1$ in (9.209) we obtain

$$\|\widehat{\mathscr{R}}_1\|_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|r\|_{s+\sigma}^{q,\kappa}.$$

$$(9.213)$$

Let us move to the estimate of the first member of $\widehat{\mathscr{R}}_2$ in (9.209). Then using Lemma 7.7-(iii) combined with (9.186) and (9.188)

$$\| [\partial_{\theta} \mathcal{A}, \mathbb{A}] \|_{2\overline{\alpha}-1+\epsilon, s, \gamma}^{q, \kappa} \leqslant C \big(\| \mu_{r, \lambda} \|_{s_0+3}^{q, \kappa} + \| \rho \|_{s_0+3}^{q, \kappa} \big) \big(\| \mu_{r, \lambda} \|_{s+\gamma+3}^{q, \kappa} + \| \rho \|_{s+\gamma+3}^{q, \kappa} \big).$$

Applying (9.206) and the estimate of $\mu_{r,\lambda}$ stated in Proposition 9.3 together with Sobolev embeddings and (9.184), one gets for any $s \ge s_0, \gamma \in \mathbb{N}$

$$\begin{split} \|\|[\partial_{\theta}\mathscr{A},\mathbb{A}]\|\|_{2\overline{\alpha}-1+\epsilon,s,\gamma}^{q,\kappa} &\lesssim \quad \varepsilon\kappa^{-2} \big(1+\|\mathfrak{I}_{0}\|_{s_{0}+\sigma}^{q,\kappa}\big) \big(1+\|\mathfrak{I}_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\big) \\ &\lesssim \quad \varepsilon\kappa^{-2} \big(1+\|\mathfrak{I}_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\big). \end{split}$$

Then according to Theorem 8.1-(ii), we deduce that for any $s \ge s_0$ and for any $t \in [0, 1]$

$$\||\Phi(-t)[\partial_{\theta}\mathcal{A},\mathbb{A}]\Phi(t)|||_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant Ce^{C\overline{\mu}^{s+1}(0)}\overline{\mu}(s)$$
(9.214)

with

$$\overline{\mu}(s) = \overline{F}_0(s + 27 + s_0) + \|\rho\|_{s+27+s_0}^{q,\kappa}$$

and

$$\overline{F}_0(s) = C\varepsilon\kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \right).$$

Notice that $\overline{F}_0(s)$ is log-convex. Using (9.206) we find

$$\overline{\mu}(s) \lesssim \varepsilon \kappa^{-2} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma}^{q,\kappa} \right).$$

Therefore we get from (9.214) and (9.184) that for any $t \in [0, 1]$ and $s \ge s_0$

$$\||\Phi(-t)[\partial_{\theta}\mathcal{A},\mathbb{A}]\Phi(t)||_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1+||\mathfrak{I}_{0}||_{s+\sigma}^{q,\kappa}\right).$$
(9.215)

For the second member of $\widehat{\mathscr{R}}_2$ in (9.209), we use Lemma 7.7-(iii) combined with (9.191) and (9.188)

$$\| [\widehat{\mathbb{A}}, \mathbb{A}] \|_{2\overline{\alpha}-1+\epsilon, s, \gamma}^{q, \kappa} \leqslant C \big(\|\rho\|_{s_0+3}^{q, \kappa} + \|\widehat{\rho}\|_{s_0+3}^{q, \kappa} \big) \big(\|\rho\|_{s+\gamma+3}^{q, \kappa} + \|\widehat{\rho}\|_{s+\gamma+3}^{q, \kappa} \big).$$

Applying (9.190) combined with the Lemma 4.1, (9.205) and (9.206) we infer

$$\begin{aligned} \|\widehat{\rho}\|_{s}^{q,\kappa} \lesssim \|\rho\|_{s+1}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} (1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}) \end{aligned}$$

It follows from (9.184) that

$$\|\|[\widehat{\mathbb{A}},\mathbb{A}]\|\|_{2\overline{\alpha}-1+\epsilon,s,\gamma}^{q,\kappa} \leqslant \varepsilon \kappa^{-2} \left(1+\|\mathfrak{I}_0\|_{s+\sigma+\gamma}^{q,\kappa}\right).$$

As before we can use Theorem 8.1-(ii) and get for any $s \ge s_0$ and $t \in [0, 1]$

$$\||\Phi(-t)[\widehat{\mathbb{A}},\mathbb{A}]\Phi(t)||_{2\overline{\alpha}-1+\epsilon,s,0} \leq Ce^{C\overline{\mu}^{s+1}(0)}\overline{\mu}(s)$$

with

$$\overline{\mu}(s) = \overline{F}_0(s + 27 + s_0) + \|\rho\|_{s+27+s_0}^{q,\kappa}$$

and

$$\overline{F}_0(s) = C\varepsilon\kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right).$$

Therefore we deduce for any $t \in [0, 1]$ and $s \ge s_0$

$$\||\Phi(-t)[\widehat{\mathbb{A}},\mathbb{A}]\Phi(t)||_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2}(1+||\mathfrak{I}_0||_{s+\sigma}^{q,\kappa}).$$

$$(9.216)$$

Combining (9.209) with (9.215) and (9.216) we find

$$\|\widehat{\mathscr{R}}_2\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2} \left(1+\|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right).$$
(9.217)

Putting together (9.208), (9.210), (9.213) and (9.217) yields

$$\||\mathscr{R}^{1}_{r,\lambda}\||_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2} \left(1+||\mathfrak{I}_{0}||_{s+\sigma}^{q,\kappa}\right).$$

The estimate for the L^2 adjoint $\mathscr{R}^{1\star}_{r\lambda}$ can be done using Lemma 7.3-(vi)

$$\begin{aligned} \|\|\mathscr{R}_{r,\lambda}^{1\star}\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} &\lesssim \||\mathscr{R}_{r,\lambda}^{1}\|\|_{2\overline{\alpha}-1+\epsilon,s+s_{0}+1,0}^{q,\kappa} \\ &\leqslant C\varepsilon\kappa^{-2} \left(1+\|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\right). \end{aligned}$$

Hence we proved the first part of the second point (ii). For the difference operator $\Delta_{12} \mathscr{R}^1_{r,\lambda}$ we write according to (9.208)

$$\Delta_{12}\mathfrak{R}^{1}_{r,\lambda} = \Delta_{12}\partial_{\theta}\mathfrak{R}_{r,\lambda} + \Delta_{12}\widehat{\mathfrak{R}}_{1} + |||\Delta_{12}\widehat{\mathfrak{R}}_{2},$$

The estimate of $\Delta_{12}\partial_{\theta}\mathcal{R}_{r,\lambda}$ is given in Proposition 9.3. Thus it remains to estimate the last two terms. The arguments are similar for both terms and we shall only discuss how to get the suitable estimate for $\Delta_{12}\widehat{\mathscr{R}}_1$. We start with writing

$$\Delta_{12}\widehat{\mathscr{R}}_1 = \int_0^1 \Phi_1(-t) \Big\{ \Delta_{12} \Big[\partial_\theta \mathscr{R}_{r,\lambda}, \mathbb{A} \Big] \Big\} \Phi_1(t) dt + \int_0^1 \Delta_{12} \Big\{ \Phi(-t) \Big[\partial_\theta \mathscr{R}_{r_2,\lambda}, \mathbb{A}_2 \Big] \Phi(t) \Big\} dt,$$

Let us estimate the first term of the right hand side. Then similarly to (9.211), using in particular the estimate (9.183) combined with the law products we obtain

$$\forall s + \gamma \leqslant \overline{s}_h, \quad |||\Delta_{12}[\partial_\theta \mathcal{R}_{r,\lambda}, \mathbb{A}]|||_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} ||\Delta_{12}i||_{\overline{s}_h+\sigma}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-1} ||\Delta_{12}i||_{\overline{s}_h+\sigma+s+\gamma}^{q,\kappa}$$
(9.218)

Therefore applying Theorem 8.1-(ii) and using (9.218) lead in view of (9.184) to

$$\int_{0}^{1} \| \Phi_{1}(-t)\Delta_{12}[\partial_{\theta}\mathcal{R}_{r,\lambda}, \mathbb{A}]\Phi_{1}(t) \|_{-1,\overline{s}_{h}-s_{0}-27,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \| \Delta_{12}i \|_{2\overline{s}_{h}+\sigma}^{q,\kappa}.$$
(9.219)

On the other hand, we apply Theorem 8.1-(iii) combined with (9.211) and (9.207)

$$\int_{0}^{1} \||\Delta_{12}\Phi(-t)[\partial_{\theta}\mathcal{R}_{r_{2},\lambda},\mathbb{A}_{2}]\Phi(t)\||_{0,0,0}^{q,\kappa} \lesssim \|\Delta_{12}\rho\|_{s_{0}+5}^{q,\kappa}$$
$$\lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa}. \tag{9.220}$$

Then combining (9.219) and (9.220) yields

$$\||\Delta_{12}\widehat{\mathscr{R}}_1||_{0,0,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$
(9.221)

Remind once again that the value of σ may change from line to line and it depends only on τ_1, d and q. The estimates of $\Delta_{12} \mathcal{R}_{r,\lambda}^1 h$ and $\Delta_{12} \mathcal{R}_{r,\lambda}^{1,\star} h$ can be implemented in a straightforward way where we use in particular the preceding estimates combined with Lemma 7.3, Proposition 8.1-(iii) and (9.224).

(iii) Recall that $\Psi = \mathscr{B}\Phi(1)$, then using Proposition 9.2-(ii) yields

$$\|\Psi h\|_{s}^{q,\kappa} \lesssim \|\Phi(1)h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-1} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|\Phi(1)h\|_{s_{0}}^{q,\kappa}$$

Applying Proposition 8.1-(ii) combined with (9.206), the smallness condition (9.184) and Sobolev embeddings we obtain

$$\begin{aligned} \|\Phi(1)h\|_{s}^{q,\kappa} &\lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\right) \|h\|_{s_{0}}^{q,\kappa} \\ &\lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}, \end{aligned}$$

which implies

$$\|\Psi h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}.$$
(9.222)

We point out that one would expect in the preceding estimate to get $||h||_{s_0+q}^{q,\kappa}$ instead of $||h||_{s_0}^{q,\kappa}$. However according to Proposition 8.1-(ii), $s_0 > \frac{d+5}{2}$ and then $s_0 + q > \frac{d+5}{2} + q$. Therefore one may $\frac{167}{167}$

replace $s_0 + q$ by s_0 by virtue of the assumption (9.2). Similar arguments give also

$$\|\Psi^{-1}h\|_{s}^{q,\kappa} \lesssim \|h\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}.$$
(9.223)

On the other hand one has

$$\Psi - \mathrm{Id} = \mathscr{B}(\Phi(1) - \mathrm{Id}) + (\mathscr{B} - \mathrm{Id})$$

For the last part we use Proposition 9.2

$$\left\| \left(\mathscr{B} - \mathrm{Id} \right) h \right\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(\|h\|_{s+1}^{q,\kappa} + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa} \right).$$

The same proposition gives

$$\left\|\mathscr{B}(\Phi(1) - \mathrm{Id})h\right\|_{s}^{q,\kappa} \lesssim \left\|(\Phi(1) - \mathrm{Id})h\right\|_{s}^{q,\kappa} + \varepsilon \kappa^{-1} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\|(\Phi(1) - \mathrm{Id})h\|_{s_{0}}^{q,\kappa}.$$

Applying once again Proposition 8.1 combined with (9.206) and (9.184) yields

$$|\mathscr{B}(\Phi(1) - \mathrm{Id})h||_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} ||h||_{s+1}^{q,\kappa} + \varepsilon \kappa^{-2} ||\mathfrak{I}_{0}||_{s+\sigma}^{q,\kappa} ||h||_{s_{0}}^{q,\kappa}.$$

Putting together the preceding estimates gives for any $s \ge s_0$

$$\left\| \left(\Psi - \mathrm{Id} \right) h \right\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|h\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}.$$

In a similar way we obtain

$$\|(\Psi^{-1} - \mathrm{Id})h\|_{s}^{q,\kappa} + \|((\Psi^{\pm 1})^{\star} - \mathrm{Id})h\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|h\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}}^{q,\kappa}.$$

Let us now move to the estimate of the difference $\Delta_{12}\Psi$ and sketch the main arguments. According to Proposition 8.1-(iii), (9.206), (9.207) and (9.184) one gets

$$\|\Delta_{12}\Phi(1)h\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \|h\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$
(9.224)

To estimate $\Delta_{12}\mathscr{B}$ where \mathscr{B} is defined in (9.34) we first write

$$\Delta_{12}\mathscr{B} = (\partial_{\theta} \Delta_{12}\beta)\mathbf{B}_1 + (1 + \partial_{\theta}\beta_2)\Delta_{12}\mathbf{B}_2$$

For the first term of the right hand side we use the law products combined with Lemma 9.4, Proposition 9.2-(iv) and (9.184)

$$\|(\partial_{\theta}\Delta_{12}\beta)\mathbf{B}_{1}h\|_{\overline{s}_{h}}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \|h\|_{\overline{s}_{h}}^{q,\kappa}.$$

Using the fundamental theorem of calculus we infer

$$\Delta_{12}\mathbf{B}h = (\beta_1 - \beta_2) \int_0^1 (\partial_\theta h) \big(\cdot, (1-\tau)\beta_1 + \tau\beta_2\big) d\tau$$

Then the same arguments as for the preceding estimate allow to get

$$\Delta_{12}\mathbf{B}h\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}\|h\|_{\overline{s}_h+1}^{q,\kappa}.$$

Putting together the foregoing estimates imply

$$\|\Delta_{12}\mathscr{B}h\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \|h\|_{\overline{s}_h+1}^{q,\kappa}.$$
(9.225)

Next we use the identity

$$\Delta_{12}\Psi = (\Delta_{12}\mathscr{B})\Phi_1(1) + \mathscr{B}_2\Delta_{12}\Phi(1) \quad \text{with} \quad \Psi = \mathscr{B}\Phi(1)$$

Applying (9.225) with Proposition 8.1-(ii), (9.206) and (9.184) leads to

$$\begin{aligned} \|(\Delta_{12}\mathscr{B})\Phi_1(1)h\|_{\overline{s}_h}^{q,\kappa} &\lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \|\Phi_1(1)h\|_{\overline{s}_h+1}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \|h\|_{\overline{s}_h+1}^{q,\kappa}. \end{aligned}$$

Similarly, (9.224) combined with (9.41) and (9.184) give

$$\begin{aligned} \|\mathscr{B}_{2}\Delta_{12}\Phi(1)h\|_{\overline{s}_{h}}^{q,\kappa} &\lesssim \|\Delta_{12}\Phi(1)h\|_{\overline{s}_{h}}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \|h\|_{\overline{s}_{h}+\sigma}^{q,\kappa}. \end{aligned}$$

Consequently, we obtain

$$\|\Delta_{12}\Psi h\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \|h\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$

The estimates of $\Delta_{12}\Psi^{-1}$, $\Delta_{12}\Psi^{\star}$ and $\Delta_{12}(\Psi^{\star})^{-1}$ are quite similar to $\Delta_{12}\Psi$. We mainly use the same arguments supplemented in particular with Lemma 9.3-(iii) and Proposition 8.1-(ii)-(iii). (iv) Recall from (9.203) that

$$\mathbf{E}_{n}^{1} = \mathbf{E}_{n}^{0} + \int_{0}^{1} \Phi(-t) \left[\mathbf{E}_{n}^{0}, \mathbb{A} \right] \Phi(t) dt - \partial_{\theta} \left[(\Pi_{N_{n}}^{\perp} \mu_{r,\lambda}) |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} (\Pi_{N_{n}}^{\perp} \mu_{r,\lambda}) \right] \\
+ \partial_{\theta} \left(\mathbf{R}_{n} |\mathbf{D}|^{\alpha-1} + |\mathbf{D}|^{\alpha-1} \mathbf{R}_{n} \right) \\
\triangleq \mathbf{E}_{n}^{0} + \mathbf{E}_{n}^{0,1} - \mathbf{E}_{n}^{0,2} + \mathbf{E}_{n}^{0,3}.$$
(9.226)

The estimate of E_n^0 is stated in Proposition 9.2. For the second one we may use the law products combined with (9.206), Proposition 9.2 and (9.184) in order to get

$$\begin{split} \big\| \big[\mathbf{E}_n^0, \mathbb{A} \big] h \big\|_{s_0}^{q,\kappa} &\lesssim \| \mathbf{E}_n^0 \mathbb{A} h \big\|_{s_0}^{q,\kappa} + \varepsilon \kappa^{-2} \| \mathbf{E}_n^0 h \|_{s_0+1}^{q,\kappa} \\ &\lesssim \varepsilon N_0^{\mu_2} N_{n+1}^{-\mu_2} \| h \|_{q,s_0+3}^{q,\kappa}. \end{split}$$

Combining this estimate with Proposition 8.1 and (9.39) implies

$$\left\|\mathbb{E}_{n}^{0,1}h\right\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon N_{0}^{\mu_{2}}N_{n+1}^{-\mu_{2}}\|h\|_{q,s_{0}+3}^{q,\kappa}.$$

For the estimate of $E_n^{0,2}$, we have from the law products, the projector properties and Proposition 9.3

$$\forall s \geq s_0, \quad \left\| \mathbf{E}_n^{0,2} h \right\|_{s_0}^{q,\kappa} \lesssim \left\| \left(\Pi_{N_n}^{\perp} \left(\mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle \right) \right\|_{s_0+1}^{q,\kappa} \| h \|_{s_0+1}^{q,\kappa} \\ \lesssim N_n^{s-s_0} \| \mu_{r,\lambda} - \langle \mu_{r,\lambda} \rangle \|_{s+1}^{q,\kappa} \| h \|_{s_0+1}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-1} N_n^{s-s_0} \left(1 + \| \Im_0 \|_{s+\sigma}^{q,\kappa} \right) \| h \|_{s_0+1}^{q,\kappa}$$

As to the estimate of $E_n^{0,3}$ in (9.226), it is quite similar to $E_n^{0,2}$, and one gets first

$$\left\| \mathbf{E}_{n}^{0,3}h \right\|_{s_{0}}^{q,\kappa} \lesssim \left\| \mathbf{R}_{n} \right\|_{s_{0}+1}^{q,\kappa} \|h\|_{s_{0}+1}^{q,\kappa}.$$
(9.227)

Then, coming back to the definition of \mathbf{R}_n in (9.200), we obtain by virtue of the estimate $c(\lambda, i_0)$ in Proposition 9.2-(i), (9.39) and (9.206)

$$\forall s \geq s_0, \quad \|\mathbf{R}_n\|_{s_0+1}^{q,\kappa} \lesssim \|\Pi_{N_n}^{\perp}\rho\|_{s_0+2}^{q,\kappa} \\ \lesssim N_n^{s_0-s} \|\Pi_{N_n}^{\perp}\rho\|_{s+2}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} N_n^{s-s_0} (1+\|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}).$$

Plugging this estimate into (9.227) implies

$$\forall s \geq s_0, \quad \left\| \mathbf{E}_n^{0,3} h \right\|_{s_0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} N_n^{s-s_0} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma}^{q,\kappa} \right) \|h\|_{s_0+1}^{q,\kappa}.$$

Combining the preceding estimates yields in view of (9.184)

$$\left\| \mathbb{E}_{n}^{1}h \right\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \|h\|_{q,s_{0}+3}^{q,\kappa} + \varepsilon \kappa^{-2} N_{n}^{s-s_{0}} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \right) \|h\|_{s_{0}+1}^{q,\kappa}.$$

This completes the proof of Proposition 9.4.

9.6. Complete reduction up to small errors. The main concern of this section is to reduce to a Fourier multiplier the operator $\widehat{\mathscr{L}}_{\omega}$ described by (9.1) and Proposition 9.1 and which takes the form (To alleviate the notation we replace therein r_0 by r)

$$\widehat{\mathscr{L}}_{\omega} = \Pi_{\mathbb{S}_0}^{\perp} \left(\mathscr{L}_{\varepsilon r, \lambda} - \varepsilon \partial_{\theta} \mathscr{R} \right) | \Pi_{\mathbb{S}_0}^{\perp}.$$
(9.228)

Notice that the set S_0 is defined in (5.6). According to Proposition 9.4, we have seen that when the parameter λ belongs to the Cantor like set $\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ then

$$\Psi^{-1}\mathscr{L}_{\varepsilon r,\lambda}\Psi = \mathscr{L}_{r,\lambda}^{1} = \omega \cdot \partial_{\varphi} + c(\lambda, i_{0})\partial_{\theta} - \mathrm{m}(\lambda, i_{0})\partial_{\theta}|\mathbf{D}|^{\alpha-1} + \mathscr{R}_{r,\lambda}^{1} + \mathbf{E}_{n}^{1}$$
(9.229)

where $\Psi : \mathcal{O} \to \mathscr{L}(H^s(\mathbb{T}^{d+1}), H^s(\mathbb{T}^{d+1}))$ is a family of invertible linear operators preserving the symmetry and satisfying suitable tame estimates.

In (5.6) we introduced the set $\mathbb{S}_0 = \mathbb{S} \cup (-\mathbb{S}) \cup \{0\}$ and the space H^s_{\perp} is described through (5.8). Now, we consider the orthogonal projector $\Pi_{\mathbb{S}_0} : L^2(\mathbb{T}^{d+1}, \mathbb{C}) \to L^2(\mathbb{T}^{d+1}, \mathbb{C})$ given by

$$h(\varphi,\theta) = \sum_{j \in \mathbb{Z}} h_j(\varphi) \mathbf{e}_j(\theta) \Longrightarrow \Pi_{\mathbb{S}_0} h(\varphi,\theta) = \sum_{j \in \mathbb{S}_0} h_j(\varphi) \, \mathbf{e}_j(\theta), \, \mathbf{e}_j(\theta) = e^{\mathrm{i}j\theta} \tag{9.230}$$

and $\Pi_{\mathbb{S}_0}^{\perp} = \mathrm{Id} - \Pi_{\mathbb{S}_0}$. Define

$$\Psi_{\perp} \triangleq \Pi_{\mathbb{S}_0}^{\perp} \Psi \Pi_{\mathbb{S}_0}^{\perp} : \mathcal{O} \longrightarrow \mathscr{L} \big(H^s_{\perp}(\mathbb{T}^{d+1}), H^s_{\perp}(\mathbb{T}^{d+1}) \big).$$

We shall discuss two points. The first one is related to the invertibility of Ψ_{\perp} with suitable tame estimates. We will see that the operators $\Psi_{\perp}^{\pm 1}$ can be viewed as finite rank perturbations of $\Psi^{\pm 1}$. This allows to get a precise description of the truncated operator

$$\Psi_{\perp}^{-1}\widehat{\mathscr{L}}_{\omega}\Psi_{\perp} = \Psi_{\perp}^{-1}\mathscr{L}_{\varepsilon r,\lambda}^{0}\Psi_{\perp} - \varepsilon \Psi_{\perp}^{-1}\partial_{\theta}\mathscr{R}\Psi_{\perp}.$$
(9.231)

Actually, one deduces that $\Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp}$ is a small finite rank perturbation of the operator $\mathscr{L}_{\varepsilon r,\lambda}$. Then the second main goal is to construct an approximate right inverse of this latter operator with tame estimates. For this aim we first use the reductions implemented before throughout Sections 9.3 and 9.5. By this way, the positive order part of the new operator is a Fourier multiplier. Thus, at this stage it remains to reduce the remainder which of order zero and enjoying a good behavior with respect to suitable norms seen in (7.13). The remainder reduction follows standard approach as in [16, 12].

9.6.1. Frequency localization of operators. We shall be concerned with some analytical properties of operators generated by frequency localization of given operators. Before stating the result, we shall first fix the framework. Let $\Psi : \mathcal{O} \to \mathscr{L}(H^s(\mathbb{T}^{d+1}), H^s(\mathbb{T}^{d+1}))$ be an arbitrary family of invertible linear operators (not necessary those of (9.229)) and let $\mathbb{S}_0 \subset \mathbb{Z}$ be a finite set as in (5.6). Consider the orthogonal projectors $\Pi_{\mathbb{S}_0}$ and $\Pi_{\mathbb{S}_0}^{\perp}$ defined by (9.230) and let us introduce the restricted transformation

$$\Psi_{\perp} \triangleq \Pi_{\mathbb{S}_0}^{\perp} \Psi \Pi_{\mathbb{S}_0}^{\perp} : \mathcal{O} \longrightarrow \mathscr{L} \big(H^s_{\perp}(\mathbb{T}^{d+1}), H^s_{\perp}(\mathbb{T}^{d+1}) \big).$$

Consider for $\lambda \in \mathcal{O}$ the finite-dimensional matrix

۱

$$\mathbf{M}(\lambda,\varphi) = \left(\left\langle \mathbf{e}_m, \left(\Psi^{\star}(\lambda) \right)^{-1} \mathbf{e}_k \right\rangle_{L^2_{\theta}(\mathbb{T})} \right)_{m,k\in\mathbb{S}_0}, \quad \mathbf{e}_k(\theta) = e^{\mathbf{i}k\theta}$$
(9.232)

where Ψ^{\star} is the L^2_{θ} -adjoint of Ψ . The scalar product $\langle \cdot \rangle_{L^2_{\theta}(\mathbb{T})}$ is defined by

$$\langle f,g\rangle_{L^2_{\theta}(\mathbb{T})} = \frac{1}{2\pi} \int_{\mathbb{T}} f(\theta) \overline{g(\theta)} d\theta.$$

Remark that the matrix entries depend only on λ and φ but not on θ because we are taking the average over θ . Our main task is to establish the following result.

Lemma 9.6. The following assertions hold true.

(i) Assume that $[\Psi, \rho] = 0$ for any smooth real function $\rho : \varphi \in \mathbb{T}^d \mapsto \rho(\varphi) \in \mathbb{R}$, and

$$\forall (\lambda, \varphi) \in \mathcal{O} \times \mathbb{T}^d, \quad \det \mathbf{M}(\lambda, \varphi) \neq 0.$$

Then, for any $h \in H^s_+$

$$\Psi_{\perp}^{-1}h = \Psi^{-1}h - \sum_{m \in \mathbb{S}_0} \left\langle h, \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \Psi^{-1} \mathbf{e}_m$$

where the functions g_m are defined via

$$\mathbf{M}^{-1}(\lambda,\varphi) = \left(\alpha_{m,k}(\lambda,\varphi)\right)_{(m,k)\in\mathbb{S}_0^2} \quad and \quad g_m(\lambda,\varphi,\theta) \triangleq \sum_{k\in\mathbb{S}_0} \overline{\alpha_{m,k}}(\lambda,\varphi)\mathbf{e}_k(\theta).$$

(ii) Let Ψ be the operator constructed in Proposition 9.4 and assume (9.184), then

$$\forall s \in [s_0, S], \quad \|\Psi_{\perp}^{\pm 1}g\|_s^{q,\kappa} \lesssim \|g\|_s^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|g\|_{s_0}^{q,\kappa}$$

In addition, we have

$$\|g_m\|_s^{q,\kappa} \lesssim \left(1 + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right)$$

and

$$\max_{m \in \mathbb{S}_0} \|\Delta_{12} g_m\|_{\overline{s}_h}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12} i\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$

Proof. (i) Given $g \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\perp})$, we want to solve the equation

$$f \in W^{q,\infty}_{\kappa}(\mathcal{O}, H^s_{\perp}), \quad \Psi_{\perp}f = \Pi^{\perp}_{\mathbb{S}_0}\Psi\Pi^{\perp}_{\mathbb{S}_0}f = g.$$

This is equivalent to

$$\Psi f = g + h$$
, with $\Pi_{\mathbb{S}_0} h = h$ and $\Pi_{\mathbb{S}_0} f = 0$.

Then we get

$$f = \Psi^{-1}(g+h), \text{ with } \Pi_{\mathbb{S}_0}h = h,$$
 (9.233)

provided that $\Pi_{\mathbb{S}_0} f = 0$, that is,

$$\langle \Psi^{-1}(g+h), \mathbf{e}_k \rangle_{L^2_{\theta}(\mathbb{T})} = 0, \quad \forall k \in \mathbb{S}_0.$$

Making appeal to the identity $(\Psi^{\star})^{-1} = (\Psi^{-1})^{\star}$, this equation is equivalent to

$$\langle g+h, \widehat{\mathbf{e}}_k \rangle_{L^2_{\theta}(\mathbb{T})} = 0, \quad \forall k \in \mathbb{S}_0, \quad \text{with} \quad \widehat{\mathbf{e}}_k \triangleq (\Psi^*(\lambda))^{-1} \mathbf{e}_k.$$
 (9.234)

These constraints will uniquely determine h. Indeed, by expanding h in the form

$$h(\varphi, \theta) = \sum_{m \in \mathbb{S}_0} a_m(\lambda, \varphi) \mathbf{e}_m(\theta),$$

we can transform the preceding system into

$$\sum_{m \in \mathbb{S}_0} a_m(\varphi) \big\langle \mathbf{e}_m, \widehat{\mathbf{e}}_k \big\rangle_{L^2_{\theta}(\mathbb{T})} = - \big\langle g, \widehat{\mathbf{e}}_k \big\rangle_{L^2_{\theta}(\mathbb{T})}, \quad \forall k \in \mathbb{S}_0.$$
(9.235)

By assumption, the matrix $\mathbf{M}(\lambda, \varphi)$ defined in (9.232) is invertible. Therefore the system (9.235) is invertible and one gets a unique solution given by

$$a_m(\lambda,\varphi) = -\sum_{k\in\mathbb{S}_0} \alpha_{m,k}(\lambda,\varphi) \langle g, \widehat{\mathbf{e}}_k \rangle_{L^2_{\theta}(\mathbb{T})} \quad \text{with} \quad \mathbf{M}^{-1}(\lambda,\varphi) \triangleq \left(\alpha_{m,k}(\lambda,\varphi)\right)_{(m,k)\in\mathbb{S}_0^2}$$

This implies

$$h(\varphi,\theta) = -\sum_{m \in \mathbb{S}_0} \left\langle g, \widehat{g}_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \mathbf{e}_m(\theta) \quad \text{with} \quad \widehat{g}_m \triangleq \sum_{k \in \mathbb{S}_0} \overline{\alpha_{m,k}} \,\widehat{\mathbf{e}}_k,$$

where we denote by $\overline{\alpha_{m,k}}$ the complex conjugate of $\alpha_{m,k}$. Using the commutation assumption $[\Psi, \rho] = 0$ we also get $[\Psi^{-1}, \rho] = 0$ and therefore

$$\Psi^{-1}h = -\sum_{m \in \mathbb{S}_0} \left\langle g, \widehat{g}_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \Psi^{-1} \mathbf{e}_m.$$
(9.236)

Notice that one also has $[(\Psi^{\star})^{-1}, \rho] = 0$, which implies in view of (9.234)

$$\widehat{g}_m = (\Psi^*(\lambda))^{-1} g_m \quad \text{with} \quad g_m \triangleq \sum_{k \in \mathbb{S}_0} \overline{\alpha_{m,k}} \, \mathbf{e}_k.$$
 (9.237)

Thus from (9.236) and (9.233) we deduce the formula

$$\Psi_{\perp}^{-1}g = \Psi^{-1}g - \sum_{m \in \mathbb{S}_0} \left\langle g, (\Psi^{\star})^{-1}g_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \Psi^{-1}\mathbf{e}_m.$$
(9.238)

Since $\Pi_{\mathbb{S}_0}^{\perp} g = g$ and $\Pi_{\mathbb{S}_0}^{\perp} g_m = 0$ then $\langle g, g_m \rangle_{L^2_{\mathfrak{a}}(\mathbb{T})} = 0$ and therefore

$$\langle g, (\Psi^{\star})^{-1} g_m \rangle_{L^2_{\theta}(\mathbb{T})} = \langle g, ((\Psi^{\star}(\lambda))^{-1} - \mathrm{Id}) g_m \rangle_{L^2_{\theta}(\mathbb{T})}$$

Plugging this identity into (9.238) yields

$$\Psi_{\perp}^{-1}g = \Psi^{-1}g - \sum_{m \in \mathbb{S}_0} \left\langle g, \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \Psi^{-1} \mathbf{e}_m.$$
(9.239)

This ends the proof of the first point.

(ii) The estimate of Ψ_{\perp} follows from the continuity of the orthogonal projection $\Pi_{\mathbb{S}_0}^{\perp}$ on Sobolev spaces combined with Proposition 9.4-(iii). Let us now move to the estimate of Ψ_{\perp}^{-1} which is more delicate. We shall first estimate in H_{φ}^s the term $\mathbf{I}_m \triangleq \langle g, ((\Psi^*)^{-1} - \mathrm{Id})g_m \rangle_{L^2_{\theta}(\mathbb{T})}$ that appears in the identity (9.239). Using the law products in Lemma 4.1 gives

$$\begin{split} \left\| \mathbf{I}_{m} \right\|_{H^{s}_{\varphi}}^{q,\kappa} \lesssim \int_{\mathbb{T}} \left\| g(\cdot,\theta) \right\|_{H^{s}_{\varphi}}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m}(\cdot,\theta) \right\|_{H^{s}_{\varphi}}^{q,\kappa} d\theta \\ + \int_{\mathbb{T}} \left\| g(\cdot,\theta) \right\|_{H^{s}_{\varphi}}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m}(\cdot,\theta) \right\|_{H^{s}_{\varphi}}^{q,\kappa} d\theta. \end{split}$$

According to Cauchy-Schwarz inequality we find

$$\begin{aligned} \left\| \mathbf{I}_{m} \right\|_{H_{\varphi}^{s}}^{q,\kappa} &\lesssim \left\| g \right\|_{L_{\theta}^{2}H_{\varphi}^{s}}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\|_{L_{\theta}^{2}H_{\varphi}^{s_{0}}}^{q,\kappa} + \left\| g \right\|_{L_{\theta}^{2}H_{\varphi}^{s_{0}}}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\|_{L_{\theta}^{2}H_{\varphi}^{s}}^{q,\kappa} \\ &\lesssim \left\| g \right\|_{s}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\|_{s_{0}}^{q,\kappa} + \left\| g \right\|_{s_{0}}^{q,\kappa} \left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\|_{s}^{q,\kappa}. \end{aligned}$$
(9.240)

Thus using Proposition 9.4-(iii) we get under the smallness condition (9.184)

$$\|((\Psi^{\star})^{-1} - \mathrm{Id})g_m\|_s^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|g_m\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|g_m\|_{s_0+q+1}^{q,\kappa}.$$
(9.241)

At this level we need to establish an estimate for $||g_m||_s^{q,\kappa}$. From (9.237) we infer

$$\|g_m\|_s^{q,\kappa} \lesssim \sum_{k \in \mathbb{S}_0} \|\alpha_{m,k}\|_{H^s_{\varphi}}^{q,\kappa}.$$
(9.242)

Recall that the coefficients $\alpha_{m,k}$ are the entries of the matrix \mathbf{M}^{-1} , where $\mathbf{M} = (c_{m,k})_{m,k\in\mathbb{S}_0}$ is defined in (9.232). Then it follows that

$$c_{m,k}(\varphi) - \delta_{km} = \left\langle \mathbf{e}_k, \left((\Psi^\star)^{-1} - \mathrm{Id} \right) \mathbf{e}_m \right\rangle_{L^2_{\theta}(\mathbb{T})}, \tag{9.243}$$

where δ_{km} is the Kronecker symbol. Consequently, we obtain

$$|c_{m,k} - \delta_{km}||_{H^s_{\varphi}}^{q,\kappa} \leq \|(\Psi^{\star})^{-1} - \mathrm{Id}\big)e_m\|_{L^2_{\theta}H^s_{\varphi}}^{q,\kappa}$$
$$\lesssim \sup_{m \in \mathbb{S}_0} \|(\Psi^{\star})^{-1} - \mathrm{Id}\big)e_m\|_s^{q,\kappa}$$

Applying (9.241) by replacing g_m with e_m gives for any $s \in [s_0, S]$,

$$\max_{k,m\in\mathbb{S}_0} \|c_{m,k} - \delta_{km}\|_{H^s_{\varphi}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right).$$
(9.244)

Finally, we deduce that

$$\mathbf{M}(\lambda,\varphi) = \mathrm{Id} - \mathbf{R}(\lambda,\varphi) \quad \text{with} \quad \|\mathbf{R}\|_{H^s_{\varphi}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right). \tag{9.245}$$

Hence under the smallness condition (9.184) combined with the law products in Lemma 4.1 we get

$$\|\mathbf{M}^{-1} - \mathrm{Id}\|_{H^s_{\varphi}}^{q,\kappa} \leq \sum_{n \geq 1} \|\mathbf{R}^n\|_s^{q,\kappa}$$
$$\leq C\varepsilon\kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right)$$

Therefore we deduce that

$$\max_{m,k\in\mathbb{S}_0} \|\alpha_{m,k}\|_{H^s_{\varphi}}^{q,\kappa} \leqslant C\left(1+\varepsilon\kappa^{-2}\|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right).$$
(9.246)

Inserting (9.246) into (9.242) allows to get

$$\|g_m\|_s^{q,\kappa} \leqslant C\left(1 + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right). \tag{9.247}$$

Putting together (9.247), (9.241) and (9.184) we find

$$\|((\Psi^{\star})^{-1} - \mathrm{Id})g_m\|_s^{q,\kappa} \leqslant C\left(1 + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\right).$$
(9.248)

Plugging (9.248) into (9.240) and using (9.184) yield

$$\left\|\mathbf{I}_{m}\right\|_{H^{s}_{\varphi}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|g\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|g\|_{s_{0}}^{q,\kappa}.$$
(9.249)

Inserting this into (9.239) and using the law products in Lemma 4.1 we obtain

$$\|\Psi_{\perp}^{-1}g\|_{s}^{q,\kappa} \lesssim \|\Psi^{-1}g\|_{s}^{q,\kappa} + \sum_{m \in \mathbb{S}_{0}} \left(\|\mathbf{I}_{m}\|_{s}^{q,\kappa}\|\Psi^{-1}\mathbf{e}_{m}\|_{s_{0}}^{q,\kappa} + \|\mathbf{I}_{m}\|_{s_{0}}^{q,\kappa}\|\Psi^{-1}\mathbf{e}_{m}\|_{s}^{q,\kappa}\right).$$
(9.250)

Thus using Proposition 9.4-(iii) and (9.184), we get successively

$$\|\Psi^{\pm 1}g\|_s^{q,\kappa} \lesssim \|g\|_s^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|g\|_{s_0}^{q,\kappa}.$$

and

$$\max_{m \in \mathbb{S}_0} \|\Psi^{-1} \mathbf{e}_m\|_s^{q,\kappa} \lesssim \|\mathbf{e}_m\|_s^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|\mathbf{e}_m\|_{s_0}^{q,\kappa} \lesssim 1 + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}.$$
(9.251)

Combining the preceding two estimates with (9.250), (9.249) and (9.184) we deduce for $s \in [s_0, S]$

$$\|\Psi_{\perp}^{-1}g\|_{s}^{q,\kappa} \lesssim \|g\|_{s}^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|g\|_{s_{0}}^{q,\kappa}.$$

This completes the proof of the desired estimate. It remains to check the estimate of $\Delta_{12}g_m$. For this purpose we come back to (9.243) which gives

$$\Delta_{12}c_{m,k}(\varphi) = \left\langle \mathbf{e}_k, \Delta_{12}(\Psi^{\star})^{-1}\mathbf{e}_m \right\rangle_{L^2_{\theta}(\mathbb{T})}.$$
(9.252)

Thus we get from Proposition (9.4)-(iii)

$$\sup_{m,k\in\mathbb{S}_0} \|\Delta_{12}c_{m,k}\|_{H_{\varphi}^{\overline{s}_h}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$

This implies from (9.245)

$$\|\Delta_{12}\mathbf{R}\|_{H^{\overline{s}_h}_{\varphi}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$

Now we remark that

$$\Delta_{12}\mathbf{M}^{-1} = \sum_{n \ge 1} \Delta_{12}\mathbf{R}^n.$$

Therefore, the law products in Lemma 4.1 combined with (9.245) and (9.184) allow to get

$$\begin{aligned} |\Delta_{12}\mathbf{M}^{-1}||_{H^{\overline{s}_h}_{\varphi}}^{q,\kappa} \lesssim \|\Delta_{12}\mathbf{R}\|_{H^{\overline{s}_h}_{\varphi}}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \end{aligned}$$

From this we deduce that

$$\max_{m,k\in\mathbb{S}_0} \|\Delta_{12}\alpha_{m,k}\|_{H^{\overline{s}_h}_{\varphi}}^{q,\kappa} \leqslant \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}$$

Hence combining this estimate with (9.237) yields

$$\max_{m \in \mathbb{S}_0} \|\Delta_{12} g_m\|_{\overline{s}_h}^{q,\kappa} \leqslant \varepsilon \kappa^{-2} \|\Delta_{12} i\|_{\overline{s}_h+\sigma}^{q,\kappa}.$$

This achieves the proof of Lemma 9.6.

9.6.2. Localization on the normal direction. The main concern of this section is to describe the localization effects on the normal direction of the operator $\hat{\mathcal{L}}_{\omega}$ described in (9.228). Our main result reads as follows.

Proposition 9.5. With the same notations and assumptions of Propositions 9.1 - 9.2 - 9.3 - 9.4, there exists $\sigma_3 = \sigma(\tau_1, d, q) \ge \sigma_2$ such that the following results hold true.

(i) On the Cantor set $\mathcal{G}_{\infty,n}^{\kappa,\tau_1}(i_0)$ introduced in Proposition 9.2, we have

$$\begin{aligned} \mathscr{L}^{2}_{\omega} &\triangleq \Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp} = \left(\omega \cdot \partial_{\varphi} + c_{r,\lambda} \partial_{\theta} - \mathbf{m}_{r,\lambda} \partial_{\theta} |\mathbf{D}|^{\alpha-1} \right) \Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{R}^{2}_{r,\lambda} + \mathbf{E}^{2}_{n} \\ &\triangleq \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{0} \right) \Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{R}^{2}_{r,\lambda} + \mathbf{E}^{2}_{n} \end{aligned}$$

where $\mathscr{R}^2_{r,\lambda} = \prod_{\mathbb{S}_0}^{\perp} \mathscr{R}^2_{r,\lambda} \prod_{\mathbb{S}_0}^{\perp}$ is reversible and \mathscr{D}_0 is reversible Fourier multiplier with

$$\forall (l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c, \quad \mathscr{D}_0 \mathbf{e}_{l,j} = \mathrm{i}\,\mu_j^0(\lambda, i_0)\mathbf{e}_{l,j},$$

with

$$\mu_j^0(\lambda, i_0) \triangleq \Omega_j(\alpha) + jr^1(\lambda, i_0) - \frac{j\Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})}r^2(\lambda, i_0).$$

In addition, for $k \in \{1, 2\}$

$$\|r^k\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_0\|^{q,\kappa}_{\overline{s}_h} \right), \quad \|\Delta_{12}r^k\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \|\Delta_{12}i\|^{q,\kappa}_{\overline{s}_h+\sigma_3}$$

(ii) The operator \mathbf{E}_n^2 is real reversible and satisfies the estimate

$$\left\|\mathbf{E}_{n}^{2}h\right\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}} + N_{n}^{s-s_{0}}\left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma_{3}}^{q,\kappa}\right)\right) \|h\|_{s_{0}+3}^{q,\kappa}$$

(iii) The operator $\mathscr{R}^2_{r,\lambda}$ is a real and reversible Töplitz in time operator satisfying

$$\forall s \in [s_0, S], \quad ||| \mathscr{R}^2_{r,\lambda} |||_{2\overline{\alpha}-1+\epsilon, s, 0} \lesssim \varepsilon \kappa^{-2} \left(1 + ||\mathfrak{I}_0||_{s+\sigma_3}^{q,\kappa}\right)$$

and

$$\||\Delta_{12}\mathscr{R}^2_{r,\lambda}|||^{q,\kappa}_{0,0,0} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|^{q,\kappa}_{\overline{s}_h+\sigma_3}$$

Proof. (i) According to (9.231) and the decomposition $\mathrm{Id} = \Pi_{\mathbb{S}_0} + \Pi_{\mathbb{S}_0}^{\perp}$ we may write

$$\begin{split} \Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp} &= \Psi_{\perp}^{-1} \mathscr{L}_{\varepsilon r, \lambda} \Psi_{\perp} - \varepsilon \Psi_{\perp}^{-1} \partial_{\theta} \mathscr{R} \Psi_{\perp} \\ &= \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \mathscr{L}_{\varepsilon r, \lambda} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} - \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \mathscr{L}_{\varepsilon r, \lambda} \Pi_{\mathbb{S}_{0}} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} - \varepsilon \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \partial_{\theta} \mathscr{R} \Psi_{\perp} \end{split}$$

Recall from Lemma 9.1 that

$$\begin{aligned} \mathscr{L}_{\varepsilon r,\lambda} &= \omega \cdot \partial_{\varphi} + \partial_{\theta} \Big[V_{\varepsilon r,\alpha} - \left(\mathscr{W}_{\varepsilon r,\alpha} \left| \mathbf{D} \right|^{\alpha - 1} + \left| \mathbf{D} \right|^{\alpha - 1} \mathscr{W}_{\varepsilon r,\alpha} \right) + \mathscr{R}_{\varepsilon r,\lambda} \Big] \\ &\triangleq \omega \cdot \partial_{\varphi} + Y_{r,\lambda}. \end{aligned}$$

Applying (9.229) yields

$$\mathscr{L}_{\varepsilon r,\lambda}\Psi = \Psi \mathscr{L}_{r,\lambda}^1$$

with

$$\mathcal{L}_{r,\lambda}^1 = \omega \cdot \partial_{\varphi} + c(\lambda, i_0)\partial_{\theta} - \mathbf{m}(\lambda, i_0)\partial_{\theta} |\mathbf{D}|^{\alpha - 1} + \mathcal{R}_{r,\lambda}^1 + \mathbf{E}_n^1$$

Therefore, we find

$$\Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp} = \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \Psi \mathscr{L}_{r,\lambda}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} - \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \left(Y_{r,\lambda} - Y_{0,\lambda} \right) \Pi_{\mathbb{S}_{0}} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} - \varepsilon \Psi_{\perp}^{-1} \partial_{\theta} \mathscr{R} \Psi_{\perp}$$

where we have used the identities

$$\Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp} = \Psi_{\perp}^{-1} \text{ and } [\Pi_{\mathbb{S}_{0}}^{\perp}, T] = 0 = [\Pi_{\mathbb{S}_{0}}, T],$$

for any Fourier multiplier T. In particular, we have used tha $Y_{0,\lambda}$ is a Fourier multiplier and thus

$$\Pi_{\mathbb{S}_0}^{\perp} Y_{0,\lambda} \Pi_{\mathbb{S}_0} = 0$$

From the structure of $\mathscr{L}^1_{r,\lambda}$ stated before and the preceding commutation relations we infer

$$\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathscr{L}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} = \Pi_{\mathbb{S}_{0}}^{\perp}\Psi\left(\omega\cdot\partial_{\varphi} + c(\lambda,i_{0})\partial_{\theta} - \operatorname{m}(\lambda,i_{0})\partial_{\theta}|\mathsf{D}|^{\alpha-1} + \mathscr{R}_{r,\lambda}^{1}\right)\Pi_{\mathbb{S}_{0}}^{\perp} + \Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathsf{E}_{n}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}$$
¹⁷⁴

$$=\Psi_{\perp}\left(\omega\cdot\partial_{\varphi}+c(\lambda,i_{0})\partial_{\theta}-\mathbf{m}(\lambda,i_{0})\partial_{\theta}|\mathbf{D}|^{\alpha-1}\right)+\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathcal{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}+\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathbf{E}_{n}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}$$

It follows that

$$\begin{split} \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathscr{L}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} &= \omega \cdot \partial_{\varphi} + c(\lambda,i_{0})\partial_{\theta} - \mathbf{m}(\lambda,i_{0})\partial_{\theta}|\mathbf{D}|^{\alpha-1} + \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} + \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathsf{E}_{n}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} \\ &= \omega \cdot \partial_{\varphi} + c(\lambda,i_{0})\partial_{\theta} - \mathbf{m}(\lambda,i_{0})\partial_{\theta}|\mathbf{D}|^{\alpha-1} + \Pi_{\mathbb{S}_{0}}^{\perp}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} + \Psi_{\perp}^{-1}\Psi\Pi_{\mathbb{S}_{0}}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} \\ &+ \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp}\Psi\mathsf{E}_{n}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}. \end{split}$$

Consequently, we find

$$\begin{split} \Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp} = & \omega \cdot \partial_{\varphi} + c(\lambda, i_{0}) \partial_{\theta} - \mathrm{m}(\lambda, i_{0}) \partial_{\theta} |\mathrm{D}|^{\alpha - 1} + \Pi_{\mathbb{S}_{0}}^{\perp} \mathscr{R}_{r,\lambda}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} + \Psi_{\perp}^{-1} \Psi \Pi_{\mathbb{S}_{0}} \mathscr{R}_{r,\lambda}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} \\ & - \Psi_{\perp}^{-1} \Pi_{\mathbb{S}_{0}}^{\perp} \left(Y_{r,\lambda} - Y_{0,\lambda} \right) \Pi_{\mathbb{S}_{0}} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} - \varepsilon \Psi_{\perp}^{-1} \partial_{\theta} \mathscr{R} \Psi_{\perp} + \Psi_{\perp}^{-1} \Psi \mathsf{E}_{n}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} \\ & \triangleq \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{0} \right) \Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{R}_{r,\lambda}^{2} + \mathsf{E}_{n}^{2}, \end{split}$$
(9.253)

with

$$\mathbf{E}_{n}^{2} \triangleq \Psi_{\perp}^{-1} \Psi \mathbf{E}_{n}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} \quad \text{and} \quad \mathscr{D}_{0} \triangleq c(\lambda, i_{0}) \partial_{\theta} - \mathbf{m}(\lambda, i_{0}) \partial_{\theta} |\mathbf{D}|^{\alpha - 1}$$

Notice that

$$\forall (l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c, \quad \mathscr{D}_0 \mathbf{e}_{l,j} = \mathrm{i}\,\mu_j^0(\lambda, i_0)\mathbf{e}_{l,j},$$

with

$$u_j^0(\lambda, i_0) \triangleq \Omega_j(\alpha) + j r^1(\lambda, i_0) - \frac{j \Gamma(j + \frac{\alpha}{2})}{\Gamma(1 + j - \frac{\alpha}{2})} r^2(\lambda, i_0),$$

where $W_{0,\alpha}$ is defined in Lemma 3.1 and

$$r^1(\lambda, i_0) \triangleq c(\lambda, i_0) - V_{0,\alpha}$$
 and $r^2(\lambda, i_0) \triangleq \frac{\mathrm{m}(\lambda, i_0) - 2^{-\alpha}C_{\alpha}}{W_{0,\alpha}}$.

Hence, the estimates of r^1 and r^2 follow from Proposition 9.4-(i).

(ii) To get the suitable estimates for E_n^2 it suffices to combine Lemma 9.6-(ii) with Proposition 9.4-(iii)-(iv) and (9.184). Then one gets first the estimates

$$\left\|\boldsymbol{\Psi}_{\perp}^{-1}\boldsymbol{\Pi}_{\mathbb{S}_{0}}^{\perp}\boldsymbol{\Psi}\mathbf{E}_{n}^{1}\boldsymbol{\Pi}_{\mathbb{S}_{0}}^{\perp}\right\|_{s_{0}}^{q,\kappa}\lesssim\left\|\mathbf{E}_{n}^{1}\boldsymbol{\Pi}_{\mathbb{S}_{0}}^{\perp}\boldsymbol{h}\right\|_{s_{0}}^{q,\kappa}$$

and

$$\left\|\mathbb{E}_{n}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}h\right\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}}+N_{n}^{s-s_{0}}\left(1+\|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\right)\right)\|h\|_{s_{0}+3}^{q,\kappa}.$$

(iii) As to the term $\mathscr{R}^2_{r,\lambda}$ we shall prove the following estimate,

$$\|\|\mathscr{R}^{2}_{r,\lambda}\|^{q,\kappa}_{2\overline{\alpha}-1+\epsilon,s} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|^{q,\kappa}_{s+\sigma}\right).$$

$$(9.254)$$

► Estimate of $\Pi_{\mathbb{S}_0}^{\perp} \mathscr{R}_{r,\lambda}^1 \Pi_{\mathbb{S}_0}^{\perp}$. One gets according to Proposition 9.4-(ii) and the continuity of the projectors

$$\| \Pi_{\mathbb{S}_{0}}^{\perp} \mathscr{R}_{r,\lambda}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \lesssim \| \mathscr{R}_{r,\lambda}^{1} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \left(1 + \| \mathfrak{I}_{0} \|_{s+\sigma}^{q,\kappa} \right).$$

$$(9.255)$$

► Estimate of $\Psi_{\perp}^{-1}\Psi\Pi_{\mathbb{S}_0}\mathscr{R}^1_{\varepsilon r,\lambda}\Pi_{\mathbb{S}_0}^{\perp}$. Using the identity of Lemma 9.6-(i) yields

$$\Psi_{\perp}^{-1}\Psi\Pi_{\mathbb{S}_{0}}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} = \Pi_{\mathbb{S}_{0}}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp} - \mathscr{T}_{0}\mathscr{S}_{1}$$

$$(9.256)$$

where

$$\mathcal{T}_0 h = \sum_{m \in \mathbb{S}_0} \left\langle h, \left((\Psi^\star)^{-1} - \mathrm{Id} \right) g_m \right\rangle_{L^2_{\theta}(\mathbb{T})} \Psi^{-1} e_m \quad \text{and} \quad \mathcal{S}_1 \triangleq \Psi \Pi_{\mathbb{S}_0} \mathcal{R}^1_{r,\lambda} \Pi^{\perp}_{\mathbb{S}_0}.$$

To estimate the first term, we use the same arguments as for (9.255)

$$\begin{split} \| \Pi_{\mathbb{S}_{0}} \mathscr{R}_{r,\lambda}^{1} \Pi_{\mathbb{S}_{0}}^{\perp} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \lesssim \| \mathscr{R}_{r,\lambda}^{1} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \left(1 + \| \Im_{0} \|_{s+\sigma}^{q,\kappa} \right) \end{split}$$

As to the second term, we write

$$\mathcal{T}_{0}\mathcal{S}_{1}h = \sum_{m \in \mathbb{S}_{0}} \left\langle \mathcal{S}_{1}h, \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\rangle_{L^{2}_{\theta}(\mathbb{T})} \Psi^{-1} e_{m}$$
$$= \sum_{m \in \mathbb{S}_{0}} \left\langle h, \mathcal{S}_{1}^{\star} \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_{m} \right\rangle_{L^{2}_{\theta}(\mathbb{T})} \Psi^{-1} e_{m},$$

where \mathscr{G}_1^{\star} is the adjoint of \mathscr{G}_1 . Observe that this operator is an integral operator taking the form

$$\mathcal{T}_0 \mathscr{S}_1 h(\varphi, \theta) = \int_{\mathbb{T}} \mathscr{K}_1(\varphi, \theta, \eta) h(\varphi, \eta) d\eta$$
(9.257)

with

$$\mathcal{K}_{1}(\varphi,\theta,\eta) \triangleq \sum_{m \in \mathbb{S}_{0}} \mathscr{S}_{1}^{\star} \big((\Psi^{\star})^{-1} - \mathrm{Id} \big) g_{m}(\varphi,\eta) \big(\Psi^{-1} e_{m} \big) (\varphi,\theta).$$

Applying Lemma 7.2-(ii)-(b), there $(-\Delta_{\eta})^{\frac{1}{2}}$ can be replaced by ∂_{η} , combined with Lemma 4.1we get for any $s \ge s_0$

$$\|\|\mathcal{T}_{0}\mathcal{S}_{1}\|\|_{-1,s,0}^{q,\kappa} \lesssim \int_{\mathbb{T}} \|\partial_{\eta}\mathcal{K}_{1}(\cdot,\cdot,\cdot+\eta)\|_{s}^{q,\kappa} d\eta \qquad (9.258)$$

$$\lesssim \sum_{m\in\mathbb{S}_{0}} \|\mathcal{S}_{1}^{\star}((\Psi^{\star})^{-1} - \mathrm{Id})g_{m}\|_{s+1}^{q,\kappa} \|\Psi^{-1}e_{m}\|_{s_{0}}^{q,\kappa}$$

$$+ \sum_{m\in\mathbb{S}_{0}} \|\mathcal{S}_{1}^{\star}((\Psi^{\star})^{-1} - \mathrm{Id})g_{m}\|_{s_{0}+1}^{q,\kappa} \|\Psi^{-1}e_{m}\|_{s}^{q,\kappa}.$$

Since

$$\mathscr{S}_1^{\star} = -\Pi_{\mathbb{S}_0}^{\perp}(\mathscr{R}_{r,\lambda}^1)^{\star}\Pi_{\mathbb{S}_0}\Psi^{\star}.$$

This implies

$$\mathscr{S}_{1}^{\star} \big((\Psi^{\star})^{-1} - \mathrm{Id} \big) g_{m} = -\Pi_{\mathbb{S}_{0}}^{\perp} (\mathscr{R}_{r,\lambda}^{1})^{\star} \Pi_{\mathbb{S}_{0}} \big(\mathrm{Id} - \Psi^{\star} \big) g_{m}.$$
(9.259)

Therefore, by applying Proposition 9.4-(ii), one gets

$$\begin{aligned} \| (\mathscr{R}^{1}_{r,\lambda})^{\star} \|_{0,s,0}^{q,\kappa} \lesssim \| (\mathscr{R}^{1}_{r,\lambda})^{\star} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \left(1 + \| \mathfrak{I}_{0} \|_{s+\sigma}^{q,\kappa} \right) \end{aligned}$$

Then combining this estimate with Lemma 7.3-(iii) and using the continuity of the orthogonal projectors yield

$$\|\mathscr{S}_{1}^{\star}((\Psi^{\star})^{-1} - \mathrm{Id})g_{m}\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\right) \|(\mathrm{Id} - \Psi^{\star})g_{m}\|_{s_{0}}^{q,\kappa} + \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s_{0}+\sigma}^{q,\kappa}\right) \|(\mathrm{Id} - \Psi^{\star})g_{m}\|_{s}^{q,\kappa}.$$

$$(9.260)$$

Putting together Proposition 9.4, the smallness condition (9.184) and (9.247) lead to

$$\left\| \left((\Psi^{\star})^{-1} - \mathrm{Id} \right) g_m \right\|_s^{q,\kappa} \lesssim \|g_m\|_s^{q,\kappa} + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|g_m\|_{s_0}^{q,\kappa}$$

$$\lesssim 1 + \varepsilon \kappa^{-2} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}.$$

$$(9.261)$$

Inserting (9.251) and (9.261) into (9.260) and using (9.184) allow to get

$$\|\mathscr{S}_{1}^{\star}((\Psi^{\star})^{-1} - \mathrm{Id})g_{m}\|_{s}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\right).$$

$$(9.262)$$

Plugging this estimate into (9.258) yields for $s \ge s_0$

$$\left\|\left|\mathcal{T}_{0}\mathcal{S}_{1}\right\|\right\|_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \left\|\mathcal{T}_{0}\right\|_{s+\sigma}^{q,\kappa}\right).$$

$$(9.263)$$

 $\textbf{Estimate of } \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp} \left(Y_{r,\lambda} - Y_{0,\lambda}\right)\Pi_{\mathbb{S}_{0}}\Psi\Pi_{\mathbb{S}_{0}}^{\perp}. \text{ We first write,} \\ \Psi_{\perp}^{-1}\Pi_{\mathbb{S}_{0}}^{\perp} \left(Y_{r,\lambda} - Y_{0,\lambda}\right)\Pi_{\mathbb{S}_{0}}\Psi\Pi_{\mathbb{S}_{0}}^{\perp} \triangleq \Psi_{\perp}^{-1}\mathscr{S}_{2}\Psi\Pi_{\mathbb{S}_{0}}^{\perp}$

with

$$\mathscr{S}_2 \triangleq (Y_{r,\lambda} - Y_{0,\lambda}) \Pi_{\mathbb{S}_0}$$

$$\triangleq \partial_{\theta} \Big(\overline{V}_{\varepsilon r, \alpha} - \big(\overline{\mathscr{W}}_{\varepsilon r, \alpha} \, |\mathbf{D}|^{\alpha - 1} + |\mathbf{D}|^{\alpha - 1} \overline{\mathscr{W}}_{\varepsilon r, \alpha} \big) + \mathscr{R}_{\varepsilon r, \lambda} \Big) \Pi_{\mathbb{S}_{0}}$$

and

$$\overline{V}_{\varepsilon r,\alpha} \triangleq V_{\varepsilon r,\alpha} - V_{0,\alpha} \quad \text{and} \quad \overline{\mathscr{W}}_{\varepsilon r,\lambda} \triangleq \mathscr{W}_{\varepsilon r,\lambda} - \mathscr{W}_{0,\alpha}$$

Recall from (9.21) that

$$(Y_{r,\lambda} - Y_{0,\lambda}) h(\varphi, \theta) = \partial_{\theta} \left(\overline{V}_{\varepsilon r,\alpha}(\varphi, \theta) h(\varphi, \theta) \right) - \partial_{\theta} \int_{\mathbb{T}} \mathscr{K}_{2}(\varphi, \theta, \eta) \rho(\varphi, \eta) d\eta,$$

with

$$\mathscr{K}_{2}(\varphi,\theta,\eta) \triangleq \frac{\overline{\mathscr{W}}_{\varepsilon r,\lambda}(\varphi,\theta) + \overline{\mathscr{W}}_{\varepsilon r,\lambda}(\varphi,\eta)}{|\sin(\frac{\theta-\eta}{2})|^{\alpha}} - \frac{\mathscr{A}_{\varepsilon r,\alpha}(\varphi,\theta,\eta)}{|\sin(\frac{\eta-\theta}{2})|^{\alpha-2}}.$$

Then from elementary computations we find

$$\mathscr{S}_2\rho(\varphi,\theta) = \partial_\theta \int_{\mathbb{T}} \mathscr{K}_3(\varphi,\theta,\eta)\rho(\varphi,\eta)d\eta, \qquad (9.264)$$

with

$$\mathcal{K}_{3}(\varphi,\theta,\eta) \triangleq \overline{V}_{\varepsilon r,\alpha}(\varphi,\theta) D_{\mathbb{S}_{0}}(\theta-\eta) - \int_{\mathbb{T}} \mathcal{K}_{2}(\varphi,\theta,\theta+\eta') D_{\mathbb{S}_{0}}(\theta+\eta'-\eta) d\eta'$$
$$D_{\mathbb{S}_{0}}(\theta) = \sum_{n \in \mathbb{S}_{0}} e^{in\theta}.$$

Consequently, we get from (9.34) and (9.35)

$$(\mathscr{B}^{-1}\mathscr{S}_2\mathscr{B})\rho(\varphi,\theta) = \partial_\theta \int_{\mathbb{T}} \mathscr{K}_4(\varphi,\theta,\eta)\rho(\varphi,\eta)d\eta, \qquad (9.265)$$

with

$$\mathscr{K}_4(\varphi,\theta,\eta) \triangleq \mathscr{K}_3(\varphi,\theta+\widehat{\beta}(\varphi,\theta),\eta+\widehat{\beta}(\varphi,\eta)).$$

Straightforward long computations similar to that used to get the estimates of (9.152) and (9.173) yield for any $\gamma \in \mathbb{N}$

$$\sup_{\eta \in \mathbb{T}} \left\| \partial_{\eta}^{\gamma} \left((\partial_{\theta} \mathscr{K}_{4})(\cdot, \cdot, \cdot, \cdot + \eta) \right) \right\|_{s}^{q, \kappa} \lesssim \varepsilon \kappa^{-1} \|r\|_{s+\sigma+\gamma}^{q, \kappa} \\ \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma+\gamma}^{q, \kappa} \right).$$
(9.266)

Applying Lemma 7.2-(ii) combined with (9.266) we infer

$$\|\|\mathscr{B}^{-1}\mathscr{S}_{2}\mathscr{B}\|\|_{-1,s,\gamma}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma+\gamma}^{q,\kappa}\right).$$

$$(9.267)$$

Thus applying Theorem 8.1 combined with (9.267), (9.206) and (9.184) yields

$$\| \Psi^{-1} \mathscr{S}_2 \Psi \|_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-1} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma}^{q,\kappa} \right) + \varepsilon \kappa^{-2} \| r \|_{s+\sigma}^{q,\kappa}$$

$$\lesssim \varepsilon \kappa^{-2} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma}^{q,\kappa} \right).$$

$$(9.268)$$

Now, proceeding as in (9.256) we obtain

$$\Psi_{\perp}^{-1}\mathscr{S}_{2}\Psi\Pi_{\mathbb{S}_{0}}^{\perp} = \Psi^{-1}\mathscr{S}_{2}\Psi\Pi_{\mathbb{S}_{0}}^{\perp} - \mathscr{T}_{0}\mathscr{S}_{2}\Psi\Pi_{\mathbb{S}_{0}}^{\perp}.$$
(9.269)

It follows that

$$\| \Psi_{\perp}^{-1} \mathscr{S}_{2} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} \|_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \| \mathfrak{I}_{0} \|_{s+\sigma}^{q,\kappa} \right) + \| \mathscr{T}_{0} \mathscr{S}_{2} \Psi \Pi_{\mathbb{S}_{0}}^{\perp} \|_{-1,s,0}^{q,\kappa} .$$
(9.270)

The estimate of the second term in (9.270) is similar to (9.263) with slight modifications and one gets

$$\||\mathcal{T}_0 \mathcal{S}_2 \Psi \Pi_{\mathbb{S}_0}^{\perp}||_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + ||\mathcal{I}_0||_{s+\sigma}^{q,\kappa}\right).$$

$$(9.271)$$

Plugging (9.271) into (9.270) we find

 $\|\!|\!| \Psi_{\perp}^{-1} \mathscr{S}_2 \Psi \Pi_{\mathbb{S}_0}^{\perp} \|\!|_{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \right).$

► Estimate of $\varepsilon \Psi_{\perp}^{-1} \partial_{\theta} \Re \Psi_{\perp}$. The estimate of this term is quite similar to the estimate of $\Psi_{\perp}^{-1} \mathscr{S}_2 \Psi$, since $\partial_{\theta} \Re$ admits an inetgral representation as in (9.264). Then using the estimates of the kernel of $\partial_{\theta} \Re$ stated in Proposition 9.1 we obtain

$$\varepsilon \parallel \!\!\mid \Psi_{\perp}^{-1} \partial_{\theta} \mathscr{R} \Psi_{\perp} \parallel \!\!\mid _{-1,s,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(1 + \lVert \mathfrak{I}_{0} \rVert_{s+\sigma}^{q,\kappa} \right).$$

Putting together the preceding estimates gives (9.254).

▶ Estimate of $\Delta_{12} \mathscr{R}^2_{r,\lambda}$. Recall from (9.253) that the operator $\mathscr{R}^2_{r,\lambda}$ takes the form

$$\mathscr{R}^{2}_{r,\lambda} = \Pi^{\perp}_{\mathbb{S}_{0}} \mathscr{R}^{1}_{r,\lambda} \Pi^{\perp}_{\mathbb{S}_{0}} + \Psi^{-1}_{\perp} \Psi \Pi_{\mathbb{S}_{0}} \mathscr{R}^{1}_{r,\lambda} \Pi^{\perp}_{\mathbb{S}_{0}} - \Psi^{-1}_{\perp} \Pi^{\perp}_{\mathbb{S}_{0}} \left(Y_{r,\lambda} - Y_{0,\lambda} \right) \Pi_{\mathbb{S}_{0}} \Psi \Pi^{\perp}_{\mathbb{S}_{0}} - \varepsilon \Psi^{-1}_{\perp} \partial_{\theta} \mathscr{R} \Psi_{\perp}.$$

$$(9.272)$$

The estimates of the different parts are slightly long and tedious sharing lot of similarities with the preceding discussion. For this reason we shall explain how to proceed for the first two terms and the remaining ones can be treated in a similar way. Concerning the first one we simply use

$$\Delta_{12}\Pi^{\perp}_{\mathbb{S}_0}\mathscr{R}^1_{r,\lambda}\Pi^{\perp}_{\mathbb{S}_0} = \Pi^{\perp}_{\mathbb{S}_0}\Delta_{12}\mathscr{R}^1_{r,\lambda}\Pi^{\perp}_{\mathbb{S}_0}.$$

Then by the continuity of the projectors combined with Proposition 9.4-(ii)

$$\||\Delta_{12}\Pi^{\perp}_{\mathbb{S}_0}\mathscr{R}^1_{r,\lambda}\Pi^{\perp}_{\mathbb{S}_0}\||_{0,0,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{q,\overline{s}_h+\sigma}^{q,\kappa}.$$
(9.273)

As to the second term of $\mathscr{R}^2_{r,\lambda}$, we use the following representation detailed in (9.256) and (9.257)

$$\Psi_{\perp}^{-1}\Psi\Pi_{\mathbb{S}_{0}}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}=\Pi_{\mathbb{S}_{0}}\mathscr{R}_{r,\lambda}^{1}\Pi_{\mathbb{S}_{0}}^{\perp}-\mathscr{T}_{0}\mathscr{S}_{1}$$

where

$$\mathcal{T}_0 \mathscr{S}_1 h(\varphi, \theta) = \int_{\mathbb{T}} \mathscr{K}_1(\varphi, \theta, \eta) h(\varphi, \eta) d\eta$$

and

$$\mathscr{K}_{1}(\varphi,\theta,\eta) \triangleq \sum_{m \in \mathbb{S}_{0}} \mathscr{S}_{1}^{\star} \big((\Psi^{\star})^{-1} - \mathrm{Id} \big) g_{m}(\varphi,\eta) \big(\Psi^{-1} e_{m} \big) (\varphi,\theta).$$

The estimate of the first term is similar to (9.273). For the operator $\mathcal{T}_0\mathcal{S}_1$ we use

$$\||\Delta_{12}\mathcal{T}_0\mathcal{S}_1|\|_{0,s_0,0}^{q,\kappa} \lesssim \int_{\mathbb{T}} \|\Delta_{12}\mathcal{K}_1(\cdot, \cdot, \cdot + \eta)\|_{s_0}^{q,\kappa} d\eta \triangleq \mathbf{I}.$$
(9.274)

From the law products we get

$$\mathbf{I} \lesssim \sum_{m \in \mathbb{S}_{0}} \|\Delta_{12} \mathscr{S}_{1}^{\star} ((\Psi^{\star})^{-1} - \mathrm{Id}) g_{m} \|_{s_{0}+1}^{q,\kappa} \|\Psi_{1}^{-1} \mathbf{e}_{m}\|_{s_{0}}^{q,\kappa} + \sum_{m \in \mathbb{S}_{0}} \|\mathscr{S}_{1,2}^{\star} ((\Psi_{2}^{\star})^{-1} - \mathrm{Id}) g_{m} \|_{s_{0}+1}^{q,\kappa} \|\Delta_{12} \Psi^{-1} \mathbf{e}_{m}\|_{s_{0}}^{q,\kappa} \triangleq \mathbf{I}_{1} + \mathbf{I}_{2}$$

where the notation f_j with j = 1, 2 denotes the value of f at the state i_j . We shall start with the estimate of I_2 . For this purpose, we use Proposition 9.4-(iii) combined with Sobolev embeddings

$$\max_{m \in \mathbb{S}_0} \|\Delta_{12} \Psi^{-1} \mathbf{e}_m\|_{s_0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12} i\|_{\overline{s}_h + \sigma}^{q,\kappa}.$$

Thus we obtain from (9.262) and (9.184)

$$\mathbf{I}_2 \lesssim \varepsilon \kappa^{-2} \|\Delta_{12} i\|_{\overline{s}_h + \sigma}^{q,\kappa}. \tag{9.275}$$

In a similar way, coming back to the identity (9.259) and applying Proposition 9.4-(ii)-(iii) combined with (9.251), (9.261) and (9.184)

$$\mathbf{I}_{1} \lesssim \max_{m \in \mathbb{S}_{0}} \left\| \Delta_{12}(\mathscr{R}_{r,\lambda}^{1,\star} \Pi_{\mathbb{S}_{0}} (\mathrm{Id} - \Psi^{\star}) g_{m}) \right\|_{s_{0}+1}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \| \Delta_{12} i \|_{s_{h}+\sigma}^{q,\kappa}.$$

Therefore inserting the preceding estimate and (9.275) into (9.274) gives

$$\||\Delta_{12}\mathcal{T}_0\mathcal{F}_1||_{0,s_0,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} ||\Delta_{12}i||_{\overline{s}_h+\sigma}^{q,\kappa}.$$

The remaining terms of $\Re_{r,\lambda}^2$ in (9.272) can be analyzed following the same arguments. Then the proof of Proposition 9.5 is achieved.

9.6.3. KAM reduction of the remainder term. According to Proposition 9.5, the operator $\mathscr{L}^2_{\omega} = \Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp}$, which is well defined in the whole set \mathscr{O} , decomposes in the Cantor set $\mathscr{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ described in Proposition 9.2, as

$$\mathcal{L}^{2}_{\omega} = \mathcal{L}_{0} + \mathbf{E}^{2}_{n}$$

$$\triangleq \left(\omega \cdot \partial_{\varphi} + \mathcal{D}_{0}\right) \Pi^{\perp}_{\mathbb{S}_{0}} + \mathcal{R}^{2}_{r,\lambda} + \mathbf{E}^{2}_{n}.$$

$$(9.276)$$

This section is devoted to the reduction of the operator \mathscr{L}_0 . It will be implemented in a classical way using the KAM reduction provided that the exterior parameters belong to a suitable Cantor set. Our main result in this section reads as follows.

Proposition 9.6. With the constraints (9.2) and using the same notations of the preceding Propositions 9.1 - 9.2 - 9.3 - 9.4 - 9.5, the following results hold true. Let

$$\mu_2 \ge \overline{\mu}_2 + 2\tau_2(1+q) \quad and \quad s_h = \frac{3}{2}\mu_2 + s_l + \tau_2(1+q) + 1 \triangleq \frac{3}{2}\mu_2 + \overline{s}_l + 1, \tag{9.277}$$

where $\overline{\mu}_2$ and s_l are defined in (9.38). There exists $\varepsilon_0 > 0$ and $\sigma_4 = \sigma(\tau_1, d, q) \ge \sigma_3$ such that if

$$\mathfrak{I}_0 \|_{q,s_h+\sigma_4} \leqslant 1 \quad \text{and} \quad N_0^{\mu_2} \varepsilon \kappa^{-3-q} \leqslant \varepsilon_0,$$

$$(9.278)$$

then the following assertions hold true.

(i) There exists a family of reversibility preserving invertible linear bounded operator $\Phi_{\infty} : \mathcal{O} \to \mathcal{L}(H^{\perp}_{\mathbb{S}_0}, H^{\perp}_{\mathbb{S}_0})$ satisfying the estimates

$$\forall s \in [s_0, S], \quad \|\Phi_{\infty}^{\pm 1}h\|_s^{q,\kappa} \leqslant C \|h\|_s^{q,\kappa} + C\varepsilon\overline{\gamma}^{-3} \|\mathfrak{I}_0\|_{s+\sigma_4}^{q,\kappa} \|h\|_{s_0}^{q,\kappa}.$$
(9.279)

There exists a family of diagonal operators $\mathscr{L}_{\infty}(\lambda, i_0)$ taking the form

$$\mathscr{L}_{\infty}(\lambda, i_0) = \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{\infty}(\lambda, i_0)\right) \Pi_{\mathbb{S}_0}^{\perp}$$

where $\mathscr{D}_{\infty}(\lambda, i_0)$ is a reversible Fourier multiplier operator given by,

$$\mathscr{A}(l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c, \quad \mathscr{D}_{\infty}(\lambda, i_0)\mathbf{e}_{l,j} = \mathrm{i}\,\mu_j^{\infty}(\lambda, i_0)\mathbf{e}_{l,j},$$

with

$$\forall j \in \mathbb{S}_0^c, \quad \mu_j^\infty(\lambda, i_0) = \mu_j^0(\lambda, i_0) + r_j^\infty(\lambda, i_0)$$

and for any $\epsilon > 0$

$$\sup_{j \in \mathbb{S}_0^c} \|r_j^{\infty}\|^{q,\kappa} \leqslant C\varepsilon\kappa^{-2} \qquad and \qquad \sup_{j \in \mathbb{S}_0^c} |j|^{1-\epsilon-2\overline{\alpha}} \|r_j^{\infty}\|^{q,\kappa} \leqslant C\varepsilon\kappa^{-2} \tag{9.280}$$

such that on the Cantor set

$$\mathcal{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \triangleq \left\{ \lambda = (\omega,\alpha) \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0); \; \forall |l| \leq N_n, \, j, j_0 \in \mathbb{S}_0^c, \, (l,j) \neq (0,j_0), \\ \left| \omega \cdot l + \mu_j^\infty(\lambda,i_0) - \mu_{j_0}^\infty(\lambda,i_0) \right| > 2\frac{\kappa\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \right\}$$

we have

$$\Phi_{\infty}^{-1}(\lambda,\omega,i_0)\mathscr{L}_0\Phi_{\infty}(\lambda,)=\mathscr{L}_{\infty}(\lambda,i_0)+\mathsf{E}_n^3(\lambda,i_0),$$

with $\mathbf{E}_n^3(\lambda, i_0)$ a linear operator satisfying

$$\|\mathbf{E}_{n}^{3}(\lambda, i_{0})h\|_{s_{0}}^{q,\kappa} \leqslant C\varepsilon\kappa^{-3}N_{0}^{\mu_{2}}N_{n+1}^{-\mu_{2}}\|h\|_{s_{0}+3}^{q,\kappa}.$$
(9.281)

Notice that the Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0)$ was introduced in Proposition 9.2, the operator \mathscr{L}_0 and the frequencies $\{\mu_j^0(\lambda)\}$ were described in Proposition 9.5 and (9.276).

(ii) Given two tori i_1 and i_2 satisfying (9.278) (replacing \mathfrak{I}_0 by \mathfrak{I}_1 or \mathfrak{I}_2), then if

$$s_h \geqslant 2\overline{s}_h + 8\tau_2(q+1)$$

 $we \ get$

$$\forall j \in \mathbb{S}_0^c, \quad \|\Delta_{12} r_j^\infty\|^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12} i\|_{\overline{s}_h + \sigma_4}^{q,\kappa} \right)^{\frac{1}{2}}$$

and

$$\forall j \in \mathbb{S}_0^c, \quad \|\Delta_{12}\mu_j^\infty\|^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \big(\|\Delta_{12}i\|_{\overline{s}_h+\sigma_4}^{q,\kappa} \big)^{\frac{1}{2}} |j|,$$

where \overline{s}_h is defined in (9.38).

Proof. (i) Set

$$\delta_0(s) \triangleq \kappa^{-1} \parallel \mathscr{R}_0 \parallel \mathscr{R}_0 \parallel s^{q,\kappa},$$

where $\mathscr{R}_0 \triangleq \mathscr{R}_{r,\lambda}^2$ being the remainder introduced in Proposition 9.5 and the norm $||| \cdot |||_s^{q,\kappa}$ is described in (7.14). According to Proposition 9.5-(iii) one gets

$$\delta_0(s) \leqslant C\varepsilon \kappa^{-3} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \right). \tag{9.282}$$

Then with the notation of (9.2), the smallness condition (9.278) implies that

$$\begin{aligned} \mathbf{N}_{0}^{\mu_{2}} \delta_{0}(s_{h}) &\leq C \mathbf{N}_{0}^{\mu_{2}} \varepsilon \kappa^{-3} \\ &\leq C \varepsilon_{0}. \end{aligned} \tag{9.283}$$

• KAM step. In view of (9.276) we write

$$\mathscr{L}_{0} = \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{0}\right) \Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{R}_{0}, \quad \mathscr{R}_{0} \triangleq \mathscr{R}_{r,\lambda}^{2}$$

$$(9.284)$$

and on the Cantor set $\mathscr{O}_{\infty,n}^{\gamma,\tau_1}(i_0)$ one has the structure

$$\mathscr{L}^2_{\omega} = \mathscr{L}_0 + \mathsf{E}_n^2$$

with \mathscr{D}_0 a diagonal Töplitz operator and \mathscr{R}_0 a real and reversible Töplitz in time operator of zero order and satisfies $\Pi_{\mathbb{S}_0}^{\perp} \mathscr{R}_0 \Pi_{\mathbb{S}_0}^{\perp} = \mathscr{R}_0$. We intend to explain the KAM step that will be implemented later during the KAM scheme in order to reduce completely \mathscr{R}_0 into a diagonal operator. The scheme can cover more general linear operators. Actually, assume that we have a linear operator \mathscr{L} such that on some Cantor set \mathscr{O} one has

$$\mathscr{L} = ig(\omega \cdot \partial_arphi + \mathscr{D}ig) \Pi^\perp_{\mathbb{S}_0} + \mathscr{R},$$

where \mathscr{D} is real and reversible diagonal Töplitz in time operator, that is,

$$\mathscr{D}\mathbf{e}_{l,j} = i\mu_j(\lambda)\mathbf{e}_{l,j}$$
 and $\mu_{-j}(\lambda) = -\mu_j(\lambda).$ (9.285)

We shall also assume that the operator \mathscr{R} is real and reversible Töplitz in time operator of zero order and satisfies $\Pi_{\mathbb{S}_0}^{\perp} \mathscr{R} \Pi_{\mathbb{S}_0}^{\perp} = \mathscr{R}$. Take a linear invertible transformation close to the identity

$$\Phi = \Pi_{\mathbb{S}_0}^{\perp} + \Psi : \mathcal{O} \to \mathcal{L}(H_{\mathbb{S}_0}^{\perp})$$

with Ψ being small and will depend on \mathscr{R} . Then

$$\Phi^{-1}\mathscr{L}\Phi = \Phi^{-1}\Big(\Phi\left(\omega\cdot\partial_{\varphi}+\mathscr{D}\right)\Pi^{\perp}_{\mathbb{S}_{0}} + \left[\omega\cdot\partial_{\varphi}\Pi^{\perp}_{\mathbb{S}_{0}}+\mathscr{D},\Psi\right] + \mathscr{R} + \mathscr{R}\Psi\Big)$$

$$= \left(\omega\cdot\partial_{\varphi}+\mathscr{D}\right)\Pi^{\perp}_{\mathbb{S}_{0}} + \Phi^{-1}\Big(\Big[\left(\omega\cdot\partial_{\varphi}+\mathscr{D}\right)\Pi^{\perp}_{\mathbb{S}_{0}},\Psi\Big] + P_{N}\mathscr{R} + P_{N}^{\perp}\mathscr{R} + \mathscr{R}\Psi\Big),$$

where the projector P_N was defined in (7.15). The basic idea consists in replacing the remainder \mathscr{R} with another quadratic one up to a diagonal part and provided that the parameters $\lambda = (\omega, \alpha)$ belongs to a Cantor set connected with non-resonance conditions associated to the *homological equation*. Iterating this scheme will generate new remainders which become smaller and smaller up to new contributions on the diagonal part and with more excision on the parameters. Then by passing to the limit we expect to diagonalize completely the operators provided that the parameters belong to Cantor set limit. Now the first step is to impose the following homological equation,

$$\left[\left(\omega \cdot \partial_{\varphi} + \mathscr{D}\right)\Pi_{\mathbb{S}_{0}}^{\perp}, \Psi\right] + P_{N}\mathscr{R} = \lfloor P_{N}\mathscr{R} \rfloor \tag{9.286}$$

where $\lfloor P_N \mathscr{R} \rfloor$ is the diagonal part of the operator \mathscr{R} . We point out that the notation $\lfloor \mathscr{R} \rfloor$ with a general operator \mathscr{R} is defined as follows, for all $(l_0, j_0) \in \mathbb{Z}^d \times \mathbb{S}_0^c$,

$$\mathscr{R}\mathbf{e}_{l_{0},j_{0}} = \sum_{(l,j)\in\mathbb{Z}^{d}\times\mathbb{S}_{0}^{c}} \mathscr{R}_{l_{0},j_{0}}^{l,j}\mathbf{e}_{l,j} \Longrightarrow \lfloor \mathscr{R} \rfloor \mathbf{e}_{l_{0},j_{0}} = \mathscr{R}_{l_{0},j_{0}}^{l_{0},j_{0}}\mathbf{e}_{l_{0},j_{0}} = \langle \mathscr{R}\mathbf{e}_{\ell_{0},j_{0}}, \mathbf{e}_{l_{0},j_{0}} \rangle_{L^{2}(\mathbb{T}^{d+1})} \mathbf{e}_{l_{0},j_{0}}.$$
(9.287)

Recall that $\mathbf{e}_{l_0,j_0}(\varphi,\theta) = e^{i(l_0\cdot\varphi+j_0\theta)}$. We define the Fourier coefficients expansion of Ψ in a standard way as follows

$$\Psi \mathbf{e}_{l_0,j_0} = \sum_{(l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c} \Psi_{l_0,j_0}^{l,j} \mathbf{e}_{l,j}.$$

After straightforward computations, using Fourier expansion we find

$$\left[\omega \cdot \partial_{\varphi} \Pi_{\mathbb{S}_{0}}^{\perp}, \Psi\right] \mathbf{e}_{l_{0}, j_{0}} = \mathrm{i} \sum_{(l, j) \in \mathbb{Z}^{d} \times \mathbb{S}_{0}^{c}} \Psi_{l_{0}, j_{0}}^{l, j} \ \omega \cdot (l - l_{0}) \ \mathbf{e}_{l, j}$$

and similarly, since \mathscr{D} is diagonal,

$$[\mathscr{D}, \Psi] \mathbf{e}_{l_0, j_0} = \mathbf{i} \sum_{(l, j) \in \mathbb{Z}^d \times \mathbb{S}_0^c} \Psi_{l_0, j_0}^{l, j} \left(\mu_j(\lambda) - \mu_{j_0}(\lambda) \right) \mathbf{e}_{l, j}.$$

By assumption, \mathscr{R} is a real and reversible Töplitz in time operator. Thus its Fourier coefficients satisfy by Proposition 7.1,

$$\mathscr{R}_{l_{0},j_{0}}^{l,j} \triangleq i r_{j_{0}}^{j}(\lambda, l_{0} - l) \in i \mathbb{R} \quad \text{and} \quad \mathscr{R}_{-l_{0},-j_{0}}^{-l,-j} = -\mathscr{R}_{l_{0},j_{0}}^{l,j}.$$
 (9.288)

Hence Ψ is a solution of (9.286) if and only if

$$\Psi \mathbf{e}_{l_0,j_0} = \sum_{\substack{|\ell-\ell_0| \leq N\\|j-j_0| \leq N}} \Psi_{l_0,j_0}^{l,j} \mathbf{e}_{l,j}$$

and

$$\Psi_{l_0,j_0}^{l,j} \left(\omega \cdot (l-l_0) + \mu_j(\lambda) - \mu_{j_0}(\lambda) \right) = \begin{cases} -r_{j_0}^j(\lambda, l_0 - l) & \text{if } (l,j) \neq (l_0,j_0) \\ 0 & \text{if } (l,j) = (l_0,j_0) \end{cases}$$

This implies that Ψ is a Töplitz in time operator with $\Psi_{l_0,j_0}^{l,j} \triangleq \Psi_{j_0}^j(l_0-l)$. In addition for $(l, j, j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2$ with $|l|, |j-j_0| \leq N$, one gets

$$\Psi_{j_0}^j(\lambda, l) = \begin{cases} \frac{-r_{j_0}^j(\lambda, l)}{\omega \cdot l + \mu_j(\lambda) - \mu_{j_0}(\lambda)} & \text{if } (l, j) \neq (0, j_0) \\ 0 & \text{if } (l, j) = (0, j_0), \end{cases}$$
(9.289)

provided that the denominator is not vanishing. Moreover since $\Pi_{\mathbb{S}_0}^{\perp} \mathscr{R} \Pi_{\mathbb{S}_0}^{\perp} = \mathscr{R}$, then one finds that

$$\forall l \in \mathbb{Z}^d, \forall j \text{ or } j_0 \in \mathbb{S}_0, \quad r_{j_0}^j(\lambda, l) = 0.$$

Consequently one may impose

$$\forall l \in \mathbb{Z}^d, \forall j \text{ or } j_0 \in \mathbb{S}_0, \quad \Psi_{j_0}^j(\lambda, l) = 0,$$

which implies in particular that $\Pi_{\mathbb{S}_0}^{\perp} \Psi \Pi_{\mathbb{S}_0}^{\perp} = \Psi$. Next, to justify the expression given by (9.289) we need to avoid resonances and restrict the parameters to the following set

$$\mathscr{O}_{+}^{\gamma} = \bigcap_{\substack{(l,j,j_{0})\in\mathbb{Z}^{d}\times(\mathbb{S}_{0}^{c})^{2}\\|l|\leqslant N\\(l,j)\neq(0,j_{0})}} \left\{\lambda\in\mathscr{O}, |\omega\cdot l+\mu_{j}(\lambda)-\mu_{j_{0}}(\lambda)| > \frac{\kappa\langle j-j_{0}\rangle}{\langle l\rangle^{\tau_{2}}}\right\}.$$

Then with this restriction the identity (9.289) is well defined and to extend Ψ to the whole set of parameters \mathcal{O} we shall use the cut-off function χ of (6.30) by setting

$$\Psi_{j_0}^j(\lambda, l) = \begin{cases} -\varrho_{j_0}^j(\lambda, l) \ r_{j_0}^j(\lambda, l), & \text{if} \quad (l, j) \neq (0, j_0) \\ 0, & \text{if} \quad (l, j) = (0, j_0), \end{cases}$$
(9.290)

with

$$\varrho_{j_0}^j(\lambda,l) \triangleq \frac{\chi\left((\omega \cdot l + \mu_j(\lambda) - \mu_{j_0}(\lambda))(\kappa\langle j - j_0\rangle)^{-1}\langle l\rangle^{\tau_2}\right)}{\omega \cdot l + \mu_j(\lambda) - \mu_{j_0}(\lambda)}.$$
(9.291)

For the sake of simplicity we shall keep the same notation and identify Ψ to its extension. Notice that the extension (9.290) is smooth and coincides with Ψ on the set \mathscr{O}_{+}^{γ} . In particular, we get from (9.288) and (9.290) that $\Psi_{i_0}^j(l) \in \mathbb{R}$. Moreover, combining (9.291) and (9.285) yields

$$\Psi_{-j_0}^{-j}(-l) = \Psi_{j_0}^j(l).$$

This implies by Proposition 7.1 that Ψ is a real and reversibility preserving operator. Let us define,

$$\mathscr{D}_{+} = \mathscr{D} + \lfloor P_{N}\mathscr{R} \rfloor, \qquad \mathscr{R}_{+} = \Phi^{-1} \left(-\Psi \lfloor P_{N}\mathscr{R} \rfloor + P_{N}^{\perp} \mathscr{R} + \mathscr{R} \Psi \right)$$
(9.292)

and

$$\mathscr{L}_{+} \triangleq \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{+} + \mathscr{R}_{+} \right) \Pi_{\mathbb{S}_{0}}^{\perp}$$

Then on the Cantor set \mathscr{O}_{+}^{γ} , we have

$$\mathscr{L}_{+} = \Phi^{-1} \mathscr{L} \Phi.$$

Notice that these operators are defined in the ambient set \mathcal{O} . Next we intend to estimate $\varrho_{j_0}^{\prime}$ introduced in (9.291), which can be written in the form

$$\varrho_{j_0}^j(\lambda,l) = a\chi_1(aA_{j,j_0}^l(\lambda)), \quad A_{j,j_0}^l(\lambda) = \omega \cdot l + \mu_j(\lambda) - \mu_{j_0}(\lambda), \quad a = (\kappa \langle j - j_0 \rangle)^{-1} \langle l \rangle^{\tau_2}, \quad (9.293)$$

where the function $\chi_1(x) \triangleq \frac{\chi(x)}{x}$ is \mathscr{C}^{∞} with bounded derivatives and $\chi_1(0) = 0$. We shall impose the following condition

$$\exists C > 0, \ \forall j, j_0 \in \mathbb{S}_0^c, \quad \max_{0 \le |\gamma| \le q} \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} \left(\mu_j(\lambda) - \mu_{j_0}(\lambda) \right) \right| \le C \left| j - j_0 \right|. \tag{9.294}$$

Hence we obtain

$$\exists C > 0, \ \forall l \in \mathbb{Z}^d, j, j_0 \in \mathbb{S}_0^c, \quad \max_{0 \le |\gamma| \le q} \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} A_{j,j_0}^l(\lambda) \right| \le C \left(|l| + |j - j_0| \right). \tag{9.295}$$

Similarly to (9.56), we may use Lemma 4.5 in order to get

$$\forall |\gamma| \leqslant q, \quad \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} \varrho_{j_0}^{j}(\lambda, l) \right| \leqslant C \kappa^{-(|\gamma|+1)} \langle l, j - j_0 \rangle^{\tau_2(1+|\gamma|)+|\gamma|}. \tag{9.296}$$

Using the characterization (7.14) combined with (9.290), (9.296) and Leibniz formula yield

$$\begin{split} \|\|\Psi\|\|_{s}^{q,\kappa} &= \left(\max_{|\gamma| \leqslant q} \sup_{\lambda \in \mathcal{O}} \sup_{n \in \mathbb{Z}} \sum_{\substack{|l| \leqslant N \\ |m| \leqslant N}} \kappa^{2|\gamma|} \langle l, m \rangle^{2(s-|\gamma|)} \left| \partial_{\lambda}^{\gamma} \Psi_{n}^{n+m}(l,\lambda) \right|^{2} \right)^{\frac{1}{2}} \\ &\leqslant C \kappa^{-1} \|\|P_{N}\mathscr{R}\|\|_{s+\tau_{2}q+\tau_{2}}^{q,\kappa} \\ &\leqslant C \kappa^{-1} \|\|\mathscr{R}\|\|_{s+\tau_{2}q+\tau_{2}}^{q,\kappa}. \end{split}$$
(9.297)

Assume that

$$C\kappa^{-1} \parallel \mathscr{R} \parallel_{s_0+\tau_2 q+\tau_2}^{q,\kappa} \leqslant \varepsilon_0, \qquad (9.298)$$

1

then applying (9.297) yields

$$\begin{aligned} \|\Psi\|\|_{s_0}^{q,\kappa} &\leq C\kappa^{-1} \|\|\mathscr{R}\|\|_{s_0+\tau_2q+\tau_2}^{q,\kappa} \\ &\leq \varepsilon_0. \end{aligned}$$
(9.299)

This implies that Φ^{-1} is invertible for small ε_0 and

$$\Phi^{-1} = \sum_{n=0}^{+\infty} (-1)^n \Psi^n \triangleq \mathrm{Id} + \Sigma$$

1.00

Applying the law products from Lemma 7.4, Lemma 7.3-(iv), (9.297) and (9.299) allow to get

$$\begin{aligned} \|\Sigma\|\|_{s}^{q,\kappa} &\leq \|\Psi\|\|_{s}^{q,\kappa} \left(1 + \sum_{\substack{n=1\\n=1}}^{+\infty} \left(C \|\|\Psi\|\|_{s_{0}}^{q,\kappa}\right)^{n}\right) \\ &\leq C \kappa^{-1} N^{\tau_{2}q+\tau_{2}} \|\|\mathscr{R}\|\|_{s}^{q,\kappa}. \end{aligned}$$

$$(9.300)$$

Consequently, we conclude under the smallness condition (9.298) that Φ^{-1} is invertible with

$$\| \Phi^{-1} - \operatorname{Id} \| \|_{s}^{q,\kappa} \leqslant C \kappa^{-1} N^{\tau_{2}q + \tau_{2}} \| \| \mathscr{R} \| \|_{s}^{q,\kappa} .$$

$$(9.301)$$

Coming back to (9.292), we may write

$$\mathscr{R}_{+} = P_{N}^{\perp} \mathscr{R} + \Phi^{-1} \mathscr{R} \Psi - \Psi \lfloor P_{N} \mathscr{R} \rfloor + \Sigma \left(P_{N}^{\perp} \mathscr{R} - \Psi \lfloor P_{N} \mathscr{R} \rfloor \right).$$

Hence, by virtue of Lemma 7.3-(iv) and (9.301), we deduce that

$$\|\mathscr{R}_{+}\|_{s}^{q,\kappa} \leq \|P_{N}^{\perp}\mathscr{R}\|_{s}^{q,\kappa} + C \|\Sigma\|_{s}^{q,\kappa} \left(\|P_{N}^{\perp}\mathscr{R}\|_{s_{0}}^{q,\kappa} + \|\Psi\|_{s_{0}}^{q,\kappa}\|\mathscr{R}\|_{s_{0}}^{q,\kappa}\right) + C\left(1 + \|\Sigma\|_{s_{0}}^{q,\kappa}\right) \left(\|\Psi\|_{s}^{q,\kappa}\|\mathscr{R}\|_{s_{0}}^{q,\kappa} + \|\Psi\|_{s_{0}}^{q,\kappa}\|\mathscr{R}\|_{s}^{q,\kappa}\right).$$
(9.302)

Combining Lemma 7.3-(iv), (9.297), (9.299) and (9.301), we obtain for all $S \ge \overline{s} \ge s \ge s_0$,

$$\| \mathscr{R}_{+} \|_{s}^{q,\kappa} \leqslant N^{s-\overline{s}} \| \mathscr{R} \|_{\overline{s}}^{q,\kappa} + C\kappa^{-1}N^{\tau_{2}q+\tau_{2}} \| \mathscr{R} \|_{s_{0}}^{q,\kappa} \| \mathscr{R} \|_{s}^{q,\kappa} .$$

$$(9.303)$$

Recall that S is a fixed arbitrary large number.

▶ Initialization We shall check that the assumptions (9.294) and (9.298) required along the KAM step to reach the final form (9.303) are satisfied with the operator $\mathscr{L} = \mathscr{L}_0$ in (9.284). Indeed, for (9.294) we shall use Lemma 3.1-(iv) in order to get

$$\exists C > 0, \ \forall (j, j_0) \in \mathbb{Z}^2, \quad \max_{0 \le |\beta| \le q} \sup_{\alpha \in (0,\overline{\alpha})} \left| \partial_{\alpha}^{\beta} \left(\Omega_j(\alpha) - \Omega_{j_0}(\alpha) \right) \right| \le C |j - j_0|.$$

$$(9.304)$$

Then applying Proposition 9.5-(i) and (9.278) we find

$$\exists C > 0, \ \forall (j, j_0) \in \mathbb{Z}^2, \quad \max_{0 \le |\beta| \le q} \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\beta} \left(\mu_j^0(\lambda) - \mu_{j_0}^0(\lambda) \right) \right| \le C \left| j - j_0 \right| \left(1 + \varepsilon \kappa^{-q-1} \right) \tag{9.305}$$
$$\le C \left| j - j_0 \right|.$$

As to the second assumption (9.298), we apply Proposition 9.5-(iii) with $\mathscr{R}_0 = \mathscr{R}_{r,\lambda}^2$ combined with (9.278) leading to

$$C\kappa^{-1} \parallel \mathscr{R}_{0} \parallel_{s_{0}+\tau_{2}q+\tau_{2}}^{q,\kappa} \leqslant C\kappa^{-1} \parallel \mathscr{R}_{r,\lambda}^{2} \parallel_{2\overline{\alpha}-1+\epsilon,s_{0}+\tau_{2}q+\tau_{2},0}^{q,\kappa} \\ \lesssim \varepsilon\kappa^{-3} \left(1+ \lVert \mathfrak{I}_{0} \rVert_{s_{h}+\sigma}^{q,\kappa}\right) \\ \leqslant C\varepsilon_{0}.$$

▶ KAM iteration. For given $m \in \mathbb{N}$ assume that we have a linear operator

$$\mathscr{L}_m \triangleq \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_m + \mathscr{R}_m \right) \Pi_{\mathbb{S}_0}^{\perp} \tag{9.306}$$

with \mathscr{D}_m a diagonal real reversible Töplitz operator and \mathscr{R}_m is a real and reversible Töplitz in time operator of zero order and satisfies $\Pi_{\mathbb{S}_0}^{\perp} \mathscr{R}_m \Pi_{\mathbb{S}_0}^{\perp} = \mathscr{R}_m$. We assume that both assumptions (9.294) and (9.298) are satisfied for \mathscr{D}_m and \mathscr{R}_m . Notice that for m = 0 we take the operator \mathscr{L}_0 defined in (9.284). Let $\Phi_m = \mathrm{Id} + \Psi_m$ be a linear invertible operator such that

$$\Phi_m^{-1} \mathscr{L}_m \Phi_m \triangleq \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{m+1} + \mathscr{R}_{m+1} \right) \Pi_{\mathbb{S}_0}^{\perp}, \tag{9.307}$$

with Ψ_m satisfying the homological equation

$$\left[\left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{m}\right)\Pi_{\mathbb{S}_{0}}^{\perp}, \Psi_{m}\right] + P_{N_{m}}\mathscr{R}_{m} = \lfloor P_{N_{m}}\mathscr{R}_{m} \rfloor.$$

Remind that N_m was introduced in (6.32). The diagonal parts $(\mathscr{D}_m)_{m\in\mathbb{N}}$ and the remainders $(\mathscr{R}_m)_{m\in\mathbb{N}}$ are defined similarly to (9.292) by the recursive formulas,

$$\mathscr{D}_{m+1} = \mathscr{D}_m + \lfloor P_{N_m} \mathscr{R}_m \rfloor \quad \text{and} \quad \mathscr{R}_{m+1} = \Phi_m^{-1} \left(-\Psi_m \lfloor P_{N_m} \mathscr{R}_m \rfloor + P_{N_m}^{\perp} \mathscr{R}_m + \mathscr{R}_m \Psi_m \right).$$
(9.308)

Notice that \mathscr{D}_m and $\lfloor P_{N_m} \mathscr{R}_m \rfloor$ are Töplitz Fourier multiplier operators that can be identified to their spectra $(i \mu_j^m)_{j \in \mathbb{S}_0^c}$ and $(i r_j^m)_{j \in \mathbb{S}_0^c}$ in the following sense

$$\forall (l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c, \quad \mathscr{D}_m \mathbf{e}_{l,j} = \mathrm{i}\,\mu_j^m \,\mathbf{e}_{l,j} \quad \text{and} \quad \lfloor P_{N_m} \mathscr{R}_m \rfloor \mathbf{e}_{l,j} = \mathrm{i}\,r_j^m \,\mathbf{e}_{l,j}. \tag{9.309}$$

By construction, we deduce that

$$\mu_j^{m+1} = \mu_j^m + r_j^m. \tag{9.310}$$

Similarly to (9.289) we obtain

$$(\Psi_m)_{j_0}^j(\lambda, l) = \begin{cases} \frac{-r_{j_0,m}^j(\lambda, l)}{\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)} & \text{if } (l, j) \neq (0, j_0) \\ 0 & \text{if } (l, j) = (0, j_0), \end{cases}$$
(9.311)

where the sequence $\{r_{j_0,m}^j(\lambda,l)\}$ describes the Fourier coefficients of \mathscr{R}_m , that is,

$$\mathscr{R}_m \mathbf{e}_{l_0,j_0} = \mathrm{i} \sum_{(l,j) \in \mathbb{Z}^{d+1}} r_{j_0,m}^j (\lambda, l_0 - l) \mathbf{e}_{l,j}.$$

We shall introduce the open Cantor set where the preceding formula has a meaning,

$$\mathscr{O}_{m+1}^{\kappa} = \bigcap_{\substack{(l,j,j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2 \\ |l| \leqslant N_m \\ (l,j) \neq (0,j_0)}} \left\{ \lambda \in \mathscr{O}_m^{\kappa}; \ |\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)| > \kappa \frac{\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \right\}.$$
(9.312)

As in (9.290) and (9.291) we may extend (9.311) as follows

$$(\Psi_m)_{j_0}^j(\lambda,\omega,l) = \begin{cases} -\frac{\chi\left((\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda))(\kappa|j - j_0|)^{-1}\langle l\rangle^{\tau_2}\right)r_{j_0,m}^j(\lambda,l)}{\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)} & \text{if } (l,j) \neq (0,j_0) \\ 0 & \text{if } (l,j) = (0,j_0). \end{cases}$$

We emphasize that working with this extension for Ψ_m allows to extend naturally both \mathscr{D}_{m+1} and the remainder \mathscr{R}_{m+1} provided that the operators \mathscr{D}_m and \mathscr{R}_m are defined in the ambient set of parameters \mathscr{O} . Thus the operator defined by the right-hand side in (9.307) can be extended to the whole set of parameters \mathscr{O} and for simplicity we still denote this extension by \mathscr{L}_{m+1} , that is,

$$\left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{m+1} + \mathscr{R}_{m+1}\right) \Pi_{\mathbb{S}_0}^{\perp} \triangleq \mathscr{L}_{m+1}.$$
(9.313)

This allows to construct by induction the sequence of operators (\mathscr{L}_{m+1}) in the full set \mathcal{O} . Similarly the operator $\Phi_m^{-1}\mathscr{L}_m\Phi_m$ admits an extension in \mathcal{O} by simply extending $\Phi_m^{\pm 1}$. Nevertheless, by construction the identity $\mathscr{L}_{m+1} = \Phi_m^{-1}\mathscr{L}_m\Phi_m$ in (9.307) occurs on the Cantor set $\mathscr{O}_{m+1}^{\kappa}$ and may fail outside this set. We set

$$\delta_m(s) \triangleq \kappa^{-1} \parallel \mathscr{R}_m \parallel_s^{q,\kappa} \tag{9.314}$$

and we intend to prove by induction in $m \in \mathbb{N}$ that

$$\forall m \in \mathbb{N}, \forall s \in [s_0, \overline{s}_l], \, \delta_m(s) \leq \delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2} \quad \text{and} \quad \delta_m(s_h) \leq \left(2 - \frac{1}{m+1}\right) \delta_0(s_h). \tag{9.315}$$

In addition, we should check the validity of the assumptions (9.294) and (9.298) for \mathscr{D}_{m+1} and \mathscr{R}_{m+1} . Observe that by Sobolev embeddings, it is sufficient to prove the first inequality with $s = \overline{s}_l$. The property is trivial for m = 0. Now, assume that the property (9.315) is true for $m \in \mathbb{N}$ and let us check it at the order m + 1. We write

$$\Phi_m^{-1} = \operatorname{Id} + \Sigma_m \quad \text{with} \quad \Sigma_m = \sum_{n=1}^{+\infty} (-1)^n \Psi_m^n.$$
(9.316)

Then proceeding as for (9.300), using in particular (9.297) and (9.299) allows to deduce successively

$$\|\Sigma_m\|\|_{s_0}^{q,\kappa} \leqslant \|\|\Psi_m\|\|_{s_0}^{q,\kappa} \left(1 + \sum_{\substack{n=0\\184}}^{+\infty} \left(C \|\|\Psi_m\|\|_{s_0}^{q,\kappa}\right)^n\right)$$

$$\leq \delta_m(s_0 + \tau_2 q + \tau_2) \left(1 + \sum_{n=0}^{+\infty} \left(C \delta_m(s_0 + \tau_2 q + \tau_2) \right)^n \right)$$

and for any $s \in [s_0, S]$,

$$\| \Sigma_{m} \|_{s}^{q,\kappa} \leq \| \Psi_{m} \|_{s}^{q,\kappa} \left(1 + \sum_{n=0}^{+\infty} \left(C \| \Psi_{m} \|_{s_{0}}^{q,\kappa} \right)^{n} \right) \\ \leq N_{m}^{\tau_{2}q+\tau_{2}} \delta_{m}(s) \left(1 + \sum_{n=0}^{+\infty} \left(C \delta_{m}(s_{0}+\tau_{2}q+\tau_{2}) \right)^{n} \right).$$

Consequently, we get by the induction assumption, since $s_0 + \tau_2 q + \tau_2 \leqslant \overline{s}_l$, $N_m \geq N_0$,

$$\begin{aligned} \|\Sigma_m\|\|_{s_0}^{q,\kappa} &\leq CN_0^{\mu_2}N_m^{-\mu_2}\delta_0(s_h) \left(1 + \sum_{\substack{n=0\\ +\infty}}^{+\infty} \left(CN_0^{\mu_2}N_m^{-\mu_2}\delta_0(s_h) \left(1 + \sum_{\substack{n=0\\ +\infty}}^{+\infty} \left(C\delta_0(s_h)\right)^n\right)\right) \end{aligned}$$

and

$$\begin{split} \|\Sigma_m\|\|_{s}^{q,\kappa} &\leq N_m^{\tau_2 q + \tau_2} \delta_m(s) \left(1 + \sum_{n=0}^{+\infty} \left(CN_0^{\mu_2} N_m^{-\mu_2} \delta_0(s_h)\right)^n\right) \\ &\leq N_m^{\tau_2 q + \tau_2} \delta_m(s) \left(1 + \sum_{n=0}^{+\infty} \left(C\delta_0(s_h)\right)^n\right). \end{split}$$

Thus under the smallness condition (9.283) we obtain for any $s \in [s_0, S]$,

$$\|\Sigma_m\|\|_{s_0}^{q,\kappa} \leqslant CN_0^{\mu_2}N_m^{-\mu_2}\delta_0(s_h) \quad \text{and} \quad \|\Sigma_m\|\|_s^{q,\kappa} \leqslant CN_m^{\tau_2q+\tau_2}\delta_m(s).$$
(9.317)

Notice that one also may obtain

$$\|\Sigma_m\|\|_s^{q,\kappa} \leqslant C\delta_m(s+\tau_2(1+q)).$$
(9.318)

Applying KAM step (9.303) and using Sobolev embeddings, we deduce that

$$\delta_{m+1}(\overline{s}_l) \leqslant N_m^{\overline{s}_l - s_h} \delta_m(s_h) + C N_m^{\tau_2 q + \tau_2} \left(\delta_m(\overline{s}_l) \right)^2$$

Thus using the induction assumption (9.315) yields

$$\begin{aligned} \delta_{m+1}(\overline{s}_l) &\leqslant N_m^{\overline{s}_l - s_h} \left(2 - \frac{1}{m+1} \right) \delta_0(s_h) + C N_m^{\tau_2 q + \tau_2} \delta_0^2(s_h) N_0^{2\mu_2} N_m^{-2\mu_2} \\ &\leqslant 2 N_m^{\overline{s}_l - s_h} \delta_0(s_h) + C N_m^{\tau_2 q + \tau_2} \delta_0^2(s_h) N_0^{2\mu_2} N_m^{-2\mu_2}. \end{aligned}$$

If we select the parameters \overline{s}_l, s_h and μ_2 such that

$$N_m^{\overline{s}_l - s_h} \leqslant \frac{1}{4} N_0^{\mu_2} N_{m+1}^{-\mu_2} \quad \text{and} \quad C N_m^{\tau_2 q + \tau_2} \delta_0(s_h) N_0^{2\mu_2} N_m^{-2\mu_2} \leqslant \frac{1}{2} N_0^{\mu_2} N_{m+1}^{-\mu_2} \tag{9.319}$$

then we find

$$\delta_{m+1}(\overline{s}_l) \leqslant \delta_0(s_h) N_0^{\mu_2} N_{m+1}^{-\mu_2}$$

By elementary arguments based on (6.32) and (9.277) we can check that the assumptions of (9.319) hold true provided that

$$4N_0^{-\mu_2} \leqslant 1$$
 and $2CN_0^{\mu_2}\delta_0(s_h) \leqslant 1.$ (9.320)

The second constraint follows from (9.283), however the first one is automatically satisfied due to $N_0 \ge 2$ and $\mu_2 \ge 2$, according to (9.2). This achieves the first statement of the induction in (9.315). Let us now move to the second estimate in (9.315). Applying the KAM step (9.303) combined with the induction assumptions (9.315)

$$\delta_{m+1}(s_h) \leq \delta_m(s_h) + C N_m^{\tau_2 q + \tau_2} \delta_m(s_0) \delta_m(s_h) \leq \left(2 - \frac{1}{m+1}\right) \delta_0(s_h) \left(1 + C N_0^{\mu_2} N_m^{\tau_2 q + \tau_2 - \mu_2} \delta_0(s_h)\right).$$

Therefore if one has

$$\left(2 - \frac{1}{m+1}\right) \left(1 + CN_0^{\mu_2} N_m^{\tau_2 q + \tau_2 - \mu_2} \delta_0(s_h)\right) \leqslant 2 - \frac{1}{m+2}$$
(9.321)

then we find

$$\delta_{m+1}(s_h) \leqslant \left(2 - \frac{1}{m+2}\right) \delta_0(s_h)$$

which achieves the induction argument of (9.315). Observe from (9.277) that $\mu_2 \ge 2\tau_2(1+q)$, then the condition (9.321) holds true if

$$CN_0^{\mu_2} N_m^{-\tau_2 q - \tau_2} \delta_0(s_h) \leqslant \frac{1}{(2m+1)(m+2)}$$
 (9.322)

Then as $N_0 \ge 2$ we may find a constant $c_0 > 0$ small enough such that

$$\forall m \in \mathbb{N}, \quad c_0 N_m^{-1} \leqslant \frac{1}{(2m+1)(m+2)}$$

Therefore (9.322) is satisfied provided that

$$CN_0^{\mu_2} N_m^{-\tau_2 q - \tau_2 + 1} \delta_0(s_h) \leqslant c_0.$$
(9.323)

From the assumption (9.2) we get in particular $\tau_2 q + \tau_2 - 1 \ge 0$. Then (9.323) is satisfied according to (9.283) when ε_0 is small enough. To achieve the induction proof of (9.315) it remains to check that the assumptions (9.294) and (9.298) are satisfied for \mathscr{D}_{m+1} and \mathscr{R}_{m+1} . First, the assumption (9.298) follows from the first inequality of (9.315) applied at the order m + 1 with $s = s_0 + \tau_2(1+q) \leq \overline{s_l}$ supplemented with (9.283). Second, to check the validity of (9.294) for the eigenvalues of \mathscr{D}_{m+1} , we combine (9.309) and (9.310) and (9.287), in order to find

$$\|\mu_j^{m+1} - \mu_j^m\|^{q,\kappa} = \left\| \left\langle P_{N_m} \mathscr{R}_m \mathbf{e}_{l,j}, \mathbf{e}_{l,j} \right\rangle_{L^2(\mathbb{T}^{d+1})} \right\|^{q,\kappa}$$

Since \mathscr{R}_m is Töplitz then

$$\|\mu_j^{m+1} - \mu_j^m\|^{q,\kappa} = \left\| \left\langle P_{N_m} \mathscr{R}_m \mathbf{e}_{0,j}, \mathbf{e}_{0,j} \right\rangle_{L^2(\mathbb{T}^{d+1})} \right\|^{q,\kappa}.$$

A duality argument combined with Lemma 7.3-(iii) and (9.314) yield

$$\mu_{j}^{m+1} - \mu_{j}^{m} \|^{q,\kappa} \lesssim \left\| \mathscr{R}_{m} \mathbf{e}_{0,j} \right\|_{s_{0}}^{q,\kappa} \langle j \rangle^{-s_{0}} \\\lesssim \left\| |\mathscr{R}_{m} \right\|_{s_{0}}^{q,\kappa} \| \mathbf{e}_{0,j} \|_{H^{s_{0}}} \langle j \rangle^{-s_{0}} \\\lesssim \left\| |\mathscr{R}_{m} \right\|_{s_{0}}^{q,\kappa} = \kappa \, \delta_{m}(s_{0}).$$

$$(9.324)$$

Consequently we deduce from (9.315), (9.282) and (9.278)

$$\|\mu_{j}^{m+1} - \mu_{j}^{m}\|^{q,\kappa} \leq C\kappa \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{m}^{-\mu_{2}}$$

$$\leq C\varepsilon \kappa^{-2} \, N_{0}^{\mu_{2}} N_{m}^{-\mu_{2}}.$$
(9.325)

Since the assumption (9.294) holds true with \mathcal{D}_m , that is,

$$\forall j, j_0 \in \mathbb{S}_0^c, \quad \max_{0 \le |\beta| \le q} \sup_{\lambda \in \mathcal{O}} \left| \partial_\lambda^\beta \left(\mu_j^m(\lambda) - \mu_{j_0}^m(\lambda) \right) \right| \le C \left| j - j_0 \right| \tag{9.326}$$

then we get by the triangle inequality, (9.325) and (9.278)

$$\begin{aligned} \forall j, j_0 \in \mathbb{S}_0^c, \quad \max_{0 \le |\beta| \le q} \sup_{\lambda \in \mathcal{O}} \left| \partial_\lambda^\beta \left(\mu_j^{m+1}(\lambda) - \mu_{j_0}^{m+1}(\lambda) \right) \right| &\leq C \left(1 + \varepsilon \kappa^{-2-q} N_0^{\mu_2} N_m^{-\mu_2} \right) |j - j_0| \\ &\leq C \left(1 + \varepsilon^{\frac{1}{3+q}} N_0^{\mu_2} N_m^{-\mu_2} \right) |j - j_0| \end{aligned}$$

Therefore using the convergence of the series $\sum_{m} N_m^{-\mu_2}$ and (9.325) allows to guarantee the required assumption with the same constant C independently of m and this completes the induction principle. Next, we shall provide some estimates for Ψ_m that will be used later to study the strong convergence. Applying (9.297) combined with Lemma 7.3 and $s_0 + \tau_2(1+q) + 1 \leq \overline{s}_l$ yield

$$\|\Psi_m\|\|_{s_0+1}^{q,\kappa} \leqslant C\kappa^{-1} \|\|P_{N_m}\mathscr{R}_m\|\|_{s_0+\tau_2(q+1)+1}^{q,\kappa}$$

$$\leq C \,\delta_m(\overline{s}_l).$$
 (9.327)

Then we infer from (9.315) and (9.283)

$$\begin{aligned} |\Psi_m||_{s_0+1}^{q,\kappa} &\leq C \,\delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2} \\ &\leq C \,\varepsilon \kappa^{-3} N_0^{\mu_2} \, N_m^{-\mu_2}. \end{aligned} \tag{9.328}$$

Let us now discuss the persistence of higher regularity. Take $s \in [s_0, S]$, then from (9.303) and (9.315) we find

$$\delta_{m+1}(s) \leq \delta_m(s) \Big(1 + C N_m^{\tau_2 q + \tau_2} \delta_m(s_0) \Big) \\ \leq \delta_m(s) \Big(1 + C N_0^{\mu_2} N_m^{\tau_2 q + \tau_2 - \mu_2} \delta_0(s_h) \Big).$$

Putting together this estimate with Lemma A.2, (9.282) and (9.283) we infer

$$\forall s \in [s_0, S], \ \forall m \in \mathbb{N}, \quad \delta_m(s) \leqslant \delta_0(s) \prod_{n=0}^{\infty} \left(1 + C N_0^{\mu_2} N_m^{\tau_2 q + \tau_2 - \mu_2} \delta_0(s_h) \right)$$
$$\leqslant C \delta_0(s) e^{\sum_{n \in \mathbb{N}} C N_0^{\mu_2} N_n^{\tau_2 q + \tau_2 - \mu_2} \delta_0(s_h)}$$
$$\leqslant C \delta_0(s) e^{C N_0^{\tau_2 q + \tau_2} \delta_0(s_h)}$$
$$\leqslant C \varepsilon \kappa^{-3} \left(1 + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \right).$$
(9.329)

Thus (9.297) together with Lemma 7.3 and interpolation inequalities lead to

$$\begin{aligned} |\Psi_m||_s^{q,\kappa} &\leq C\kappa^{-1} ||| P_{N_m} \mathscr{R}_m ||_{s+\tau_2 q+\tau_2}^{q,\kappa} \\ &\leq C \,\delta_m(s+\tau_2 q+\tau_2) \\ &\leq C \,\delta_m^{\overline{\theta}}(s_0) \delta_m^{1-\overline{\theta}}(s+\tau_2 q+\tau_2+1). \end{aligned}$$
(9.330)

with $\overline{\theta} = \frac{1}{s - s_0 + \tau_2(1+q)}$. Plugging (9.315) and (9.329) into (9.330) and using (9.283) we obtain

$$\|\Psi_m\|\|_s^{q,\kappa} \leqslant C \,\delta_0^{\overline{\theta}}(s_h) \delta_0^{1-\overline{\theta}}(s+\tau_2 q+\tau_2+1) N_0^{\mu_2 \overline{\theta}} N_m^{-\mu_2 \overline{\theta}} \\ \leqslant C \,\varepsilon_0^{\overline{\theta}} \delta_0^{1-\overline{\theta}}(s+\tau_2 q+\tau_2+1) N_m^{-\mu_2 \overline{\theta}}.$$

$$(9.331)$$

We observe that one also deduces from (9.318), the second inequality of (9.330) and (9.329) that

$$\forall s \in [s_0, S], \quad \sup_{m \in \mathbb{N}} \left(\|\Sigma_m\|\|_s^{q,\kappa} + \|\|\Psi_m\|\|_s^{q,\kappa} \right) \leqslant C \varepsilon \kappa^{-3} \left(1 + \|\mathfrak{I}_0\|\|_{s+\sigma+\tau_2(1+q)}^{q,\kappa} \right).$$
(9.332)

Hence we find

$$\forall s \in [s_0, S], \quad \sup_{m \in \mathbb{N}} \| \Phi_m^{\pm 1} - \operatorname{Id} \|_s^{q,\kappa} \leqslant C \varepsilon \kappa^{-3} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma+\tau_2(1+q)}^{q,\kappa} \right).$$
(9.333)

► KAM conclusion. Consider the sequence of operators $\left(\widehat{\Phi}_{m}\right)_{m \in \mathbb{N}}$,

$$\widehat{\Phi}_0 \triangleq \Phi_0 \quad \text{and} \qquad \forall m \ge 1, \ \widehat{\Phi}_m \triangleq \Phi_0 \circ \Phi_1 \circ \dots \circ \Phi_m.$$
(9.334)

Then it is clear from the identity $\Phi_m = \mathrm{Id} + \Psi_m$ that $\widehat{\Phi}_{m+1} = \widehat{\Phi}_m + \widehat{\Phi}_m \Psi_{m+1}$. Using the law products of Lemma 7.4 we obtain

$$\||\widehat{\Phi}_{m+1}||_{s_0+1}^{q,\kappa} \leqslant \||\widehat{\Phi}_m||_{s_0+1}^{q,\kappa} \left(1+C ||| \Psi_{m+1} ||_{s_0+1}^{q,\kappa}\right).$$

Then iterating this inequality and using (9.283) and (9.328) yield

$$\||\widehat{\Phi}_{m+1}|||_{s_0+1}^{q,\kappa} \leq \||\widehat{\Phi}_0|||_{s_0+1}^{q,\kappa} \prod_{n=1}^{m+1} \left(1 + C ||| \Psi_n |||_{s_0+1}^{q,\kappa}\right)$$
$$\leq \prod_{n=0}^{\infty} \left(1 + C \varepsilon_0 N_n^{-\mu_2}\right).$$

From the first condition of (9.320) and (6.32) and one gets

$$\|\widehat{\Phi}_{m+1}\|_{\text{O-d},q,s_0+1} \leqslant \prod_{n=0}^{\infty} \left(1 + C \varepsilon_0 4^{-(\frac{3}{2})^n}\right).$$

Since the infinite product converges and for ε_0 small enough then we obtain

$$\sup_{m \in \mathbb{N}} \| \widehat{\Phi}_m \|_{s_0+1}^{q,\kappa} \leqslant 2.$$
(9.335)

Next, we intend to estimate the difference $\widehat{\Phi}_{m+1} - \widehat{\Phi}_m$ and for this aim we use the law products of Lemma 7.4 combined with (9.328) and (9.335)

$$\||\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}|||_{s_{0}+1}^{q,\kappa} \leqslant C ||| \widehat{\Phi}_{m} |||_{s_{0}+1}^{q,\kappa} |||\Psi_{m+1}|||_{s_{0}+1}^{q,\kappa} \leqslant C \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} \, N_{m+1}^{-\mu_{2}}.$$
(9.336)

Consequently, Lemma A.2 gives

$$\sum_{m=0}^{\infty} \||\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}|||_{s_0+1}^{q,\kappa} \leqslant C \,\delta_0(s_h).$$
(9.337)

Then by a completeness argument we deduce that the series $\widehat{\Phi}_0 + \sum_{m \in \mathbb{N}} (\widehat{\Phi}_{m+1} - \widehat{\Phi}_m)$ converges to an element Φ_{∞} . Furthermore, we get in view of (9.336)

$$\||\widehat{\Phi}_{m} - \Phi_{\infty}|||_{s_{0}+1}^{q,\kappa} \leq \sum_{j \geq m} \||\widehat{\Phi}_{j+1} - \widehat{\Phi}_{j}|||_{s_{0}+1}^{q,\kappa}$$
$$\leq C \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} \sum_{j \geq m} N_{j+1}^{-\mu_{2}}$$
$$\leq C \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{m+1}^{-\mu_{2}}.$$
(9.338)

Notice that one also deduces from (9.335)

$$\|\| \Phi_{\infty} \|_{s_0+1}^{q,\kappa} \leqslant 2. \tag{9.339}$$

On the other hand, using (9.337) combined with (9.330) for m = 0 and (9.2)

$$\||\Phi_{\infty} - \mathrm{Id}\||_{s_{0}+1}^{q,\kappa} \leq \sum_{m=0}^{\infty} |||\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}||_{s_{0}+1}^{q,\kappa} + |||\Psi_{0}|||_{s_{0}+1}^{q,\kappa}$$

$$\leq C \,\delta_{0}(s_{h}).$$
(9.340)

Let us now analyze the convergence in higher norms. Take $s \in [s_0, S]$, then using the law products of of Lemma 7.4, (9.328), (9.331) and (9.335) we obtain

$$\|\widehat{\Phi}_{m+1}\|\|_{s}^{q,\kappa} \leq \|\widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} \left(1+C \|\|\Psi_{m+1}\|\|_{s_{0}}^{q,\kappa}\right) + C \|\|\widehat{\Phi}_{m}\|\|_{s_{0}}^{q,\kappa} \|\|\Psi_{m+1}\|\|_{s}^{q,\kappa}$$

$$\leq \|\widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} \left(1+C\varepsilon_{0}N_{m+1}^{-\mu_{2}}\right) + C\delta_{0}^{\overline{\theta}}(s_{h})N_{0}^{\mu_{2}\overline{\theta}}\delta_{0}^{1-\overline{\theta}}(s+\tau_{2}q+\tau_{2}+1)N_{m}^{-\mu_{2}\overline{\theta}}.$$
(9.341)

From the first condition of (9.320) and (6.32) and one gets

$$\prod_{n=0}^{\infty} \left(1 + C \varepsilon_0 N_n^{-\mu_2} \right) \leqslant \prod_{n=0}^{\infty} \left(1 + C \varepsilon_0 4^{-\left(\frac{3}{2}\right)^n} \right) \\ \leqslant 2$$

where the last inequality holds if ε_0 is chosen small enough. Similarly we get

$$\sum_{n=0}^{\infty} N_n^{-\mu_2 \overline{\theta}} < \infty.$$

Then applying (9.89) to (9.341) and using (9.297) yields

$$\sup_{m\in\mathbb{N}} \|\widehat{\Phi}_m\|\|_s^{q,\kappa} \leq C\Big(\|\|\Phi_0\|\|_s^{q,\kappa} + \delta_0^{\overline{\theta}}(s_h)N_0^{\mu_2\overline{\theta}}\delta_0^{1-\overline{\theta}}(s+\tau_2q+\tau_2+1)\Big)$$
$$\leq C\Big(1+\delta_0(s+\tau_2q+\tau_2) + \delta_0^{\overline{\theta}}(s_h)N_0^{\mu_2\overline{\theta}}\delta_0^{1-\overline{\theta}}\big(s+\tau_2q+\tau_2+1\big)\Big).$$

Using interpolation inequalities and (9.283) we find

$$\sup_{m\in\mathbb{N}} \||\widehat{\Phi}_m\||_s^{q,\kappa} \leqslant C\Big(1+\delta_0\big(s+\tau_2q+\tau_2+1\big)\Big).$$
(9.342)

The next goal is to estimate the difference $\||\widehat{\Phi}_{m+1} - \widehat{\Phi}_m||_s^{q,\kappa}$. Then by the law products of Lemma 7.4 combined with the first inequality in (9.328), (9.331), (9.335) and (9.342) one gets

$$\begin{split} \|\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} &\leq C \Big(\|\widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} \|\|\Psi_{m+1}\|\|_{s_{0}}^{q,\kappa} + \|\widehat{\Phi}_{m}\|\|_{s_{0}}^{q,\kappa} \|\|\Psi_{m+1}\|\|_{s}^{q,\kappa} \Big) \\ &\leq C \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} \, N_{m+1}^{-\mu_{2}} \Big(1 + \delta_{0} \Big(s + \tau_{2}q + \tau_{2} + 1 \Big) \Big) \\ &+ C \, \delta_{0}^{\overline{\theta}}(s_{h}) N_{0}^{\mu_{2}\overline{\theta}} \, \delta_{0}^{1-\overline{\theta}} \Big(s + \tau_{2}q + \tau_{2} + 1 \Big) N_{m+1}^{-\mu_{2}\overline{\theta}}. \end{split}$$

Consequently, we obtain in view of Lemma A.2

$$\sum_{m=0}^{\infty} \|\|\widehat{\Phi}_{m+1} - \widehat{\Phi}_m\|\|_s^{q,\kappa} \leq C \,\delta_0(s_h) \left(1 + \delta_0 \left(s + \tau_2 q + \tau_2 + 1\right)\right) \\ + C \,\delta_0^{\overline{\theta}}(s_h) \,\delta_0^{1-\overline{\theta}} \left(s + \tau_2 q + \tau_2 + 1\right).$$

Using interpolation inequalities and the second condition in (9.320) we get

$$\sum_{m=0}^{\infty} \| \widehat{\Phi}_{m+1} - \widehat{\Phi}_{m} \| \|_{s}^{q,\kappa} \leqslant C \Big(\delta_{0}(s_{h}) + \delta_{0} \big(s + \tau_{2}q + \tau_{2} + 1 \big) \Big).$$
(9.343)

Therefore we find from this latter inequality combined with (9.283) and (9.342)

$$\|\Phi_{\infty}\|\|_{s}^{q,\kappa} \leq \sum_{m=0}^{\infty} \|\|\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} + \|\|\widehat{\Phi}_{0}\|\|_{s}^{q,\kappa} \leq C\Big(1 + \delta_{0}\big(s + \tau_{2}q + \tau_{2} + 1\big)\Big).$$
(9.344)

On the other hand, one can easily check that, using (9.343) and the second inequality in (9.330) with m = 0,

$$\|\Phi_{\infty} - \operatorname{Id}\|\|_{s}^{q,\kappa} \leq \sum_{m=0}^{\infty} \|\widehat{\Phi}_{m+1} - \widehat{\Phi}_{m}\|\|_{s}^{q,\kappa} + \|\Psi_{0}\|\|_{s}^{q,\kappa}$$
$$\leq C\Big(\delta_{0}(s_{h}) + \delta_{0}\big(s + \tau_{2}q + \tau_{2} + 1\big)\Big).$$
(9.345)

Applying Lemma 7.3-(iii), we obtain by virtue of (9.339), (9.344) and Sobolev embeddings

$$\begin{split} \|\Phi_{\infty}\rho\|_{s}^{q,\kappa} &\lesssim \|\|\Phi_{\infty}\|\|_{s_{0}}^{q,\kappa}\|\rho\|_{s}^{q,\kappa} + \|\Phi_{\infty}\|\|_{s}^{q,\kappa}\|\rho\|_{s_{0}}^{q,\kappa} \\ &\lesssim \|\rho\|_{s}^{q,\kappa} + \left(1 + \delta_{0}\left(s + \tau_{2}q + \tau_{2} + 1\right)\right)\|\rho\|_{s_{0}}^{q,\kappa} \\ &\lesssim \|\rho\|_{s}^{q,\kappa} + \delta_{0}\left(s + \tau_{2}q + \tau_{2} + 1\right)\|\rho\|_{s_{0}}^{q,\kappa}. \end{split}$$
(9.346)

According to (9.282) we infer

$$\delta_0 \left(s + \tau_2 q + \tau_2 + 1 \right) \leqslant C \varepsilon \kappa^{-3} \left(1 + \| \mathfrak{I}_0 \|_{s+\sigma+\tau_2 q+\tau_2+1}^{q,\kappa} \right).$$

$$(9.347)$$

Inserting (9.347) into (9.346) and using (9.278) combined with Sobolev embeddings and (9.2) gives

$$\|\Phi_{\infty}\rho\|_{s}^{q,\kappa} \lesssim \|\rho\|_{s}^{q,\kappa} + \varepsilon \kappa^{-3} \left(1 + \|\mathfrak{I}_{0}\|_{s+\sigma+\tau_{2}q+\tau_{2}+1}^{q,\kappa}\right) \|\rho\|_{s_{0}}^{q,\kappa}$$
189

$$\lesssim \|\rho\|_{s}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|\rho\|_{s_{0}}^{q,\kappa}.$$
(9.348)

Notice that in the last line we have replaced $\sigma + \tau_2 q + \tau_2 + 1$ by σ which can be justified by taking σ sufficient large.

Similarly to (9.346) we get by applying Lemma 7.3-(iii) combined with (9.345) and (9.340)

$$\begin{split} \left\| \left(\Phi_{\infty} - \mathrm{Id} \right) \rho \right\|_{s}^{q,\kappa} &\lesssim \| |\Phi_{\infty} - \mathrm{Id} \| \|_{s_{0}}^{q,\kappa} \| \rho \|_{s}^{q,\kappa} + \| \Phi_{\infty} - \mathrm{Id} \| \|_{s}^{q,\kappa} \| \rho \|_{s_{0}}^{q,\kappa} \\ &\lesssim \delta_{0}(s_{h}) \| \rho \|_{s}^{q,\kappa} + \left(\delta_{0}(s_{h}) + \delta_{0} \left(s + \tau_{2}q + \tau_{2} + 1 \right) \right) \| \rho \|_{s_{0}}^{q,\kappa} \\ &\lesssim \delta_{0}(s_{h}) \| \rho \|_{s}^{q,\kappa} + \delta_{0} \left(s + \tau_{2}q + \tau_{2} + 1 \right) \| \rho \|_{s_{0}}^{q,\kappa}. \end{split}$$

Then we find from (9.347) and (9.283) combined with Sobolev embeddings and (9.282)

$$\left\| \left(\Phi_{\infty} - \mathrm{Id} \right) \rho \right\|_{s}^{q,\kappa} \lesssim \left(\varepsilon \kappa^{-3} + \delta_{0}(s_{h}) \right) \|\rho\|_{s}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_{0}\|_{s+\sigma+\tau_{2}q+\tau_{2}+1}^{q,\kappa} \|\rho\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-3} \|\rho\|_{s}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_{0}\|_{s+\sigma+\tau_{2}q+\tau_{2}+1}^{s,\kappa} \|\rho\|_{s_{0}}^{q,\kappa} .$$

$$(9.349)$$

The estimates of $\Phi_{\infty}^{-1}, \Phi_{\infty}^{-1} - \widehat{\Phi}_{n}^{-1}$ can be checked using similar arguments.

♦ Next we shall study the asymptotic expansion of the eigenvalues. Summing up in m the estimates (9.325) and using Lemma A.2 we find

$$\sum_{m=0}^{\infty} \|\mu_{j}^{m+1} - \mu_{j}^{m}\|^{q,\kappa} \leq C\kappa \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} \sum_{m=0}^{\infty} N_{m}^{-\mu_{2}} \leq C\kappa \delta_{0}(s_{h}).$$
(9.350)

This shows that for each $j \in \mathbb{S}_0^c$ the sequence $(\mu_j^m)_{m \in \mathbb{N}}$ converges in the space $W^{q,\infty,\gamma}(\mathcal{O},\mathbb{C})$ to an element denoted by $\mu_j^{\infty} \in W^{q,\infty,\gamma}(\mathcal{O},\mathbb{C})$. In addition, for any $m \in \mathbb{N}$, we find in view of (9.325)

$$\|\mu_j^{\infty} - \mu_j^m\|^{q,\kappa} \leq \sum_{n=m}^{\infty} \|\mu_j^{n+1} - \mu_j^n\|^{q,\kappa}$$
$$\leq \kappa \,\delta_0(s_h) N_0^{\mu_2} \sum_{n=m}^{\infty} N_n^{-\mu_2}$$

Therefore we deduce according ton Lemma A.2

$$\sup_{j \in \mathbb{S}_0^c} \|\mu_j^{\infty} - \mu_j^m\|^{q,\kappa} \leqslant C\kappa \delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2}.$$
(9.351)

One also gets

$$\mu_{j}^{\infty} = \mu_{j}^{0} + \sum_{m=0}^{\infty} \left(\mu_{j}^{m+1} - \mu_{j}^{m} \right)$$
$$\triangleq \mu_{j}^{0} + r_{j}^{\infty}$$
(9.352)

where (μ_i^0) is described in Proposition 9.5-(i) and given by

$$\mu_j^0(\lambda, i_0) = \Omega_j(\alpha) + jr^1(\lambda, i_0) + j|j|^{\alpha - 1}r^2(\lambda, i_0).$$

Therefore (9.350) and (9.283) yield

$$\|r_j^{\infty}\|^{q,\kappa} \leq C \kappa \,\delta_0(s_h)$$
$$\leq C \,\varepsilon \kappa^{-2}$$

and this gives the first result in (9.280).

Now, let us consider the diagonal operator \mathscr{D}_∞ defined on the normal modes by

$$\forall (l,j) \in \mathbb{Z}^d \times \mathbb{S}_0^c, \quad \mathscr{D}_\infty \mathbf{e}_{l,j} = \mathrm{i}\mu_j^\infty \mathbf{e}_{l,j}.$$
(9.353)

By virtue of (7.14) we obtain

$$|\mathscr{D}_m - \mathscr{D}_\infty||_{s_0}^{q,\kappa} = \sup_{j \in \mathbb{S}_0^c} \|\mu_j^m - \mu_j^\infty\|^{q,\kappa}$$

which gives by virtue of (9.351)

$$\begin{aligned} \|\mathscr{D}_m - \mathscr{D}_\infty\|\|_{s_0}^{q,\kappa} &\leq C \,\kappa \,\delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2} \\ &\leq C \,\varepsilon \kappa^{-2} \,N_0^{\mu_2} N_m^{-\mu_2}. \end{aligned} \tag{9.354}$$

• In what follows we shall prove that the Cantor set $\mathscr{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0)$ defined in Proposition 9.6 satisfies

$$\mathcal{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \subset \bigcap_{m=0}^{n+1} \mathcal{O}_m^{\kappa} = \mathcal{O}_{n+1}^{\kappa}$$

whereas the intermediate Cantor sets are defined in (9.312). For this goal we shall proceed by finite induction on m with n fixed. First, we notice that by construction $\mathscr{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \subset \mathscr{O} \triangleq \mathscr{O}_0^{\kappa}$. Now assume that $\mathscr{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \subset \mathscr{O}_m^{\kappa}$ for all $m \leq n$ and let us check that

$$\mathscr{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \subset \mathscr{O}_{m+1}^{\kappa}. \tag{9.355}$$

Take $\lambda \in \mathcal{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0)$ and give $(l, j, j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2$ such that $0 \leq |l| \leq N_m$ and $(l, j) \neq (0, j_0)$. Then we may write by the triangle inequality, (9.351), (9.2) and (9.283)

$$\begin{split} |\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)| &\ge |\omega \cdot l + \mu_j^\infty(\lambda) - \mu_{j_0}^\infty(\lambda)| - 2\sup_{j \in \mathbb{S}_0^c} \|\mu_j^m - \mu_j^\infty\|^{q,\kappa} \\ &\ge \frac{2\kappa\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} - C\kappa\delta_0(s_h)N_0^{\mu_2}N_m^{-\mu_2} \\ &\ge 2\frac{\kappa\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} - C\kappa\varepsilon_0\langle l \rangle^{-\mu_2}\langle j - j_0 \rangle. \end{split}$$

Then for ε_0 small enough and using that $\mu_2 \ge \mu_c \ge \tau_2$, which follows from (9.277), we get

$$\left|\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)\right| > \frac{\kappa \langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}}.$$

This shows that $\lambda \in \mathcal{O}_{m+1}^{\kappa}$ and therefore the inclusion (9.355) holds.

★ In what follows the *convergence* of the sequence $(\mathscr{L}_m)_{m\in\mathbb{N}}$ introduced in (9.306) towards the diagonal operator $\mathscr{L}_{\infty} \triangleq \omega \cdot \partial_{\varphi} + \mathscr{D}_{\infty}$, where \mathscr{D}_{∞} is given by (9.353). Making use of (9.354) and (9.315) we find

$$\begin{aligned} \|\mathscr{L}_m - \mathscr{L}_\infty\|\|_{s_0}^{q,\kappa} &\leq \||\mathscr{D}_m - \mathscr{D}_\infty|\|_{s_0}^{q,\kappa} + \||\mathscr{R}_m\|\|_{s_0}^{q,\kappa} \\ &\leq C \,\kappa \,\delta_0(s_h) N_0^{\mu_2} N_m^{-\mu_2}, \end{aligned} \tag{9.356}$$

which implies in particular that

$$\lim_{n \to \infty} \left\| \left| \mathscr{L}_m - \mathscr{L}_\infty \right| \right\|_{s_0}^{q,\kappa} = 0.$$
(9.357)

On the other hand, we remark from (9.334) and (9.307) that

$$\forall \lambda \in \mathscr{O}_{n+1}^{\kappa}, \quad \widehat{\Phi}_n^{-1} \mathscr{L}_0 \widehat{\Phi}_n = \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{n+1} + \mathscr{R}_{n+1} \right) \Pi_{\mathbb{S}_0}^{\perp} \\ = \mathscr{L}_{\infty} + \left(\mathscr{D}_{n+1} - \mathscr{D}_{\infty} + \mathscr{R}_{n+1} \right) \Pi_{\mathbb{S}_0}^{\perp}$$

and therefore for any $\lambda \in \mathscr{O}_{n+1}^{\kappa}$ we may write the decomposition

$$\Phi_{\infty}^{-1}\mathscr{L}_{0}\Phi_{\infty} = \mathscr{L}_{\infty} + \left(\mathscr{D}_{n+1} - \mathscr{D}_{\infty} + \mathscr{R}_{n+1}\right)\Pi_{\mathbb{S}_{0}}^{\perp} + \Phi_{\infty}^{-1}\mathscr{L}_{0}\left(\Phi_{\infty} - \widehat{\Phi}_{n}\right) + \left(\Phi_{\infty}^{-1} - \widehat{\Phi}_{n}^{-1}\right)\mathscr{L}_{0}\widehat{\Phi}_{n} \triangleq \mathscr{L}_{\infty} + \mathsf{E}_{n,1}^{2} + \mathsf{E}_{n,2}^{2} + \mathsf{E}_{n,3}^{2} \triangleq \mathscr{L}_{\infty} + \mathsf{E}_{n}^{3}.$$
(9.358)

To estimate $E_{n,1}^2$ we use (9.354) combined with (9.314), (9.315), (9.282) and (9.278)

$$\|\mathbf{E}_{n,1}^{2}\|\|_{s_{0}}^{q,\kappa} \leqslant C \kappa \,\delta_{0}(s_{h}) N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}} \\ \leqslant C \varepsilon \kappa^{-2} N_{0}^{\mu_{2}} N_{n+1}^{-\mu_{2}}.$$
(9.359)

Putting together Lemma 7.3-(iii) with (9.359) yields

$$\|\mathbf{E}_{n,1}^2h\|_{s_0}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2}N_0^{\mu_2}N_{n+1}^{-\mu_2}\|h\|_{s_0}^{q,\kappa}.$$

Next let us prove the estimates of $E_{n,2}^2$ and $E_{n,3}^2$ defined in (9.358). They can be implemented in a similar way and therefore we shall restrict the discussion to the term $E_{n,2}^2$. Using (9.279) yields

$$\|\mathbf{E}_{n,2}^{2}h\|_{s_{0}}^{q,\kappa} \lesssim \|\mathscr{L}_{0}(\Phi_{\infty}-\widehat{\Phi}_{n})h\|_{s_{0}}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_{0}\|_{s_{0}+\sigma}^{q,\kappa} \|\mathscr{L}_{0}(\Phi_{\infty}-\widehat{\Phi}_{n})h\|_{s_{0}}^{q,\kappa}.$$
(9.360)

Then combining this estimate with (9.360) and (9.278) we obtain

$$\begin{aligned} \|\mathbf{E}_{n,2}^{2}h\|_{s_{0}}^{q,\kappa} &\lesssim \left\|\mathscr{L}_{0}\left(\Phi_{\infty}-\widehat{\Phi}_{n}\right)h\right\|_{s_{0}}^{q,\kappa} \\ &\lesssim \left\|\left(\Phi_{\infty}-\widehat{\Phi}_{n}\right)h\right\|_{s_{0}+3}^{q,\kappa}. \end{aligned}$$

Hence (9.338) together with Lemma 7.3, (9.278) and (9.282) allow to find

$$\begin{split} \|\mathbf{E}_{n,2}^{2}h\|_{s_{0}}^{q,\kappa} &\lesssim \|\|\Phi_{\infty} - \widehat{\Phi}_{n}\|\|_{s_{0}+3}^{q,\kappa} \|h\|_{s_{0}+3}^{q,\kappa} \\ &\leqslant C\,\delta_{0}(s_{h})N_{0}^{\mu_{2}}N_{m+1}^{-\mu_{2}}\|h\|_{s_{0}+3}^{q,\kappa} \\ &\leqslant C\,\varepsilon\kappa^{-3}N_{0}^{\mu_{2}}N_{m+1}^{-\mu_{2}}\|h\|_{s_{0}+3}^{q,\kappa}. \end{split}$$

The estimate of $E_{n,3}^2$ can be done in a similar way to $E_{n,1}^2$ and we get the same estimate. Putting together the foregoing estimates yields (9.281).

 \blacklozenge The next task is to prove the second estimate of (9.280). Define

$$\widehat{\delta}_m(s) \triangleq \kappa^{-1} \|\|\mathscr{R}_m\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa},$$

where the corresponding norm is introduced in (7.13). Then we shall prove by induction on $m \in \mathbb{N}$ that

$$\widehat{\delta}_m(s_0) \leqslant C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \quad \text{and} \quad \widehat{\delta}_m(s_h) \leqslant C \varepsilon \kappa^{-3} \left(2 - \frac{1}{m+1}\right). \tag{9.361}$$

Using Proposition 9.5-(iii) combined with (9.278) one gets

$$\widehat{\delta}_{0}(s_{h}) \leq C \varepsilon \kappa^{-3} \left(1 + \| \mathfrak{I}_{0} \|_{s_{h}+\sigma}^{q,\kappa} \right) \\ \leq C \varepsilon \kappa^{-3}.$$
(9.362)

which shows that (9.361) holds true for m = 0. Now assume that (9.361) is satisfied at the order m and let us check it at the order m + 1. Applying Lemma 7.4 to (9.308) and using (9.316) we obtain the expression

$$\| \mathscr{R}_{m+1} \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leq \| \Phi_m^{-1} P_{N_m}^{\perp} \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} + C \| \Phi_m^{-1} \|_{0,s,0}^{q,\kappa} \| \mathscr{U}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} + C \| \Phi_m^{-1} \|_{0,s_0,0}^{q,\kappa} \| \mathscr{U}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa},$$

$$+ C \| \Phi_m^{-1} \|_{0,s_0,0}^{q,\kappa} \| \mathscr{U}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa},$$

$$(9.363)$$

with

$$\mathcal{U}_m \triangleq -\Psi_m \left\lfloor P_{N_m} \mathscr{R}_m \right\rfloor + \mathscr{R}_m \Psi_m$$

Then putting together (9.363), (9.333) and (9.278) gives

$$\|\|\mathscr{R}_{m+1}\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant \|\|\Phi_m^{-1}P_{N_m}^{\perp}\mathscr{R}_m\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} + C\varepsilon\kappa^{-3}\left(1+\|\mathfrak{I}_0\|\|_{s+\sigma}^{q,\kappa}\right) \|\|\mathscr{U}_m\|\|_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} + C \|\|\mathscr{U}_m\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa}.$$

$$(9.364)$$

Using Lemma 7.4 we deduce that

$$\begin{split} \|\Phi_m^{-1} P_{N_m}^{\perp} \mathscr{R}_m\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leqslant \|P_{N_m}^{\perp} \mathscr{R}_m\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} + C \|\sum_m\|_{0,s,0}^{q,\kappa}\|\mathscr{R}_m\|_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} \\ + C \|\sum_m\|_{0,s_0,0}^{q,\kappa}\|\mathscr{R}_m\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \,. \end{split}$$

Consequently we find from Lemma 7.3-(iv), (9.317) and (9.282)

$$\| \Phi_m^{-1} P_{N_m}^{\perp} \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \leq N_m^{s-\overline{s}} \| \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,\overline{s},0}^{q,\kappa} + C N_m^{\tau_2q+\tau_2} \delta_m(s) \| \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} + C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \| \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa},$$

$$(9.365)$$

and for $s = s_0$, we get in view of (9.282) and (9.278)

$$\| \Phi_m^{-1} P_{N_m}^{\perp} \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} \leqslant N_m^{s_0-\overline{s}} \| \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,\overline{s},0}^{q,\kappa} + C\varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \| \mathscr{R}_m \|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} .$$

$$(9.366)$$

Applying once again Lemma 7.4 yields

 $\|\|\mathcal{U}_m\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa} \lesssim \|\|\Psi_m\|\|_{0,s,0}^{q,\kappa}\|\|\mathscr{R}_m\|\|_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} + \|\|\Psi_m\|\|_{0,s_0,0}^{q,\kappa}\|\|\mathscr{R}_m\|\|_{2\overline{\alpha}-1+\epsilon,s,0}^{q,\kappa}.$ Hence we derive from (9.328) and (9.330)

$$\||\mathcal{U}_m||_{2\overline{\alpha}-1+\epsilon,s_h,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-3} \||\mathscr{R}_m||_{2\overline{\alpha}-1+\epsilon,s_0,0}^{q,\kappa} + \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \||\mathscr{R}_m||_{2\overline{\alpha}-1+\epsilon,s_h,0}^{q,\kappa}$$

In addition, for $s = s_0$, we obtain

$$\left\|\left|\mathcal{U}_{m}\right|\right\|_{2\overline{\alpha}-1+\epsilon,s_{0},0}^{q,\kappa} \lesssim \varepsilon \kappa^{-3} N_{0}^{\mu_{2}} N_{m}^{-\mu_{2}} \left\|\left|\left|\mathcal{R}_{m}\right|\right|\right|_{2\overline{\alpha}-1+\epsilon,s_{0},0}^{q,\kappa}.$$

$$(9.367)$$

Then plugging (9.367) and (9.366) into (9.364) and using (9.278) we find

$$\widehat{\delta}_{m+1}(s_0) \leqslant C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \widehat{\delta}_m(s_0) + C N_m^{s_0 - s_h} \widehat{\delta}_m(s_h).$$

In a similar way, we find

$$\widehat{\delta}_{m+1}(s_h) \leqslant \widehat{\delta}_m(s_h) + C\varepsilon\kappa^{-3}N_0^{\mu_2}N_m^{-\mu_2}\widehat{\delta}_m(s_h) + CN_m^{\tau_2q+\tau_2}\delta_m(s_h)\widehat{\delta}_m(s_0) + \varepsilon\kappa^{-3}\widehat{\delta}_m(s_0).$$

Applying (9.315) yields

that

$$\widehat{\delta}_{m+1}(s_h) \leqslant \widehat{\delta}_m(s_h) + C\varepsilon\kappa^{-3}N_0^{\mu_2}N_m^{-\mu_2}\widehat{\delta}_m(s_h) + C\varepsilon\kappa^{-3}N_m^{\tau_2q+\tau_2}\widehat{\delta}_m(s_0) + \varepsilon\kappa^{-3}\widehat{\delta}_m(s_0).$$

Therefore with the constraints on μ_2 and s_h imposed in (9.277) we achieve the validity of the induction (9.361) at the order m + 1 in a similar way to that of (9.315). Next, we will see how to deduce the second estimate of (9.280). Recall from (9.310) and (9.352)

$$r_j^{\infty} = \sum_{j=1}^{+\infty} r_j^m \quad \text{with} \quad r_j^m = \left\langle P_{N_m} \mathscr{R}_m \mathbf{e}_{0,j}, \mathbf{e}_{0,j} \right\rangle_{L^2(\mathbb{T}^{d+1},\mathbb{C})}.$$

Then for any $a \in \mathbb{R}$ and $j \in \mathbb{Z}^*$ it is clear from integration by parts that

m=0

$$\langle P_{N_m} \mathscr{R}_m \mathbf{e}_{0,j}, \mathbf{e}_{0,j} \rangle_{L^2(\mathbb{T}^{d+1},\mathbb{C})} = \frac{1}{|j|^a} \langle P_{N_m} \Lambda^a \mathscr{R}_m \mathbf{e}_{0,j}, \mathbf{e}_{0,j} \rangle_{L^2(\mathbb{T}^{d+1},\mathbb{C})},$$

with $\Lambda = \sqrt{-\Delta}$. Using a duality argument combined with Lemma 7.3-(iii) we obtain

$$\|\langle P_{N_m}\Lambda^a\mathscr{R}_m\mathbf{e}_{0,j},\mathbf{e}_{0,j}\rangle_{L^2(\mathbb{T}^{d+1},\mathbb{C})}\|^{q,\kappa} \leq C \|\|\Lambda^a\mathscr{R}_m\|^{q,\kappa}_{0,s_0-1,0}.$$

Applying Lemma 7.4 with $a = 1 - 2\overline{\alpha} - \epsilon$ combined with (9.361) yields

$$\|\|\Lambda^{a}\mathscr{R}_{m}\|\|_{0,s_{0}-1,0}^{q,\kappa} \leq C \|\|\mathscr{R}_{m}\|\|_{2\overline{\alpha}-1+\epsilon,s_{0},0}^{q,\kappa}$$
$$\leq C\kappa \,\widehat{\delta}_{m}(s_{0})$$
$$\leq C\varepsilon \kappa^{-2} N_{0}^{\mu_{2}} N_{m}^{-\mu_{2}}.$$

It follows that

$$\|r_j^m\|^{q,\kappa} \leqslant C|j|^{2\overline{\alpha}-1+\epsilon}\varepsilon\kappa^{-2}N_0^{\mu_2}N_m^{-\mu_2}.$$

Consequently, we obtain in view of Lemma A.2

$$\begin{aligned} \|r_{j}^{\infty}\|^{q,\kappa} &\lesssim |j|^{2\overline{\alpha}-1+\epsilon} \varepsilon \kappa^{-2} N_{0}^{\mu_{2}} \sum_{m=0}^{+\infty} N_{m}^{-\mu_{2}} \\ &\lesssim |j|^{2\overline{\alpha}-1+\epsilon} \varepsilon \kappa^{-2}. \end{aligned}$$

This completes the proof of (9.280).

(ii) From (9.308) and (9.316) we can write

$$\mathscr{R}_{m+1} = (\mathrm{Id} + \Sigma_m) U_m$$
¹⁹³

with

$$U_m \triangleq P_{N_m}^{\perp} \mathscr{R}_m + \mathscr{R}_m \Psi_m - \Psi_m \lfloor P_{N_m} \mathscr{R}_m \rfloor.$$
(9.368)

Thus we get from straightforward computations

$$\Delta_{12}U_m = P_{N_m}^{\perp} \Delta_{12}\mathscr{R}_m + (\Delta_{12}\mathscr{R}_m)\Psi_m^{[1]} + \mathscr{R}_m^{[2]}(\Delta_{12}\Psi_m) - (\Delta_{12}\Psi_m) \lfloor P_{N_m}\mathscr{R}_m^{[1]} \rfloor - \Psi_m^{[2]} \lfloor P_{N_m} \Delta_{12}\mathscr{R}_m \rfloor$$
(9.369)

and

$$\Delta_{12}\mathscr{R}_{m+1} = \Delta_{12}U_m + (\Delta_{12}\Sigma_m)U_m^{[1]} + \Sigma_m^{[2]}\Delta_{12}U_m.$$
(9.370)

Where we have used the notation $f^{[k]}$ to denote the value of f at the torus i_k for k = 1, 2. Elementary calculations based on (9.316) give

$$\Delta_{12}\Sigma_m = \Delta_{12}\Phi_m^{-1} = -(\Phi_m^{[2]})^{-1}(\Delta_{12}\Psi_m)(\Phi_m^{[1]})^{-1}.$$

Applying Lemma 7.4 and using (9.332) combined with (9.278) we obtain

$$\forall s \in [s_0, s_h], \quad |||\Delta_{12} \Sigma_m |||_s^{q,\kappa} \lesssim |||\Delta_{12} \Psi_m |||_s^{q,\kappa}.$$

$$(9.371)$$

By virtue of Lemma 7.4, (9.371) and (9.370) we find

$$\|\Delta_{12}\mathscr{R}_{m+1}\|_{s_0}^{q,\kappa} \leq \|\Delta_{12}U_m\|_{s_0}^{q,\kappa} + \|\Delta_{12}\Psi_m\|_{s_0}^{q,\kappa}\|U_m^{[1]}\|_{s_0}^{q,\kappa} + \|\Sigma_m^{[2]}\|_{s_0}^{q,\kappa}\|\Delta_{12}U_m\|_{s_0}^{q,\kappa}$$

$$(9.372)$$

and

$$\begin{split} \|\Delta_{12}\mathscr{R}_{m+1}\|_{s_{h}}^{q,\kappa} &\leq \|\Delta_{12}U_{m}\|_{s_{h}}^{q,\kappa} + \|\Delta_{12}\Psi_{m}\|_{s_{0}}^{q,\kappa} \|U_{m}^{[1]}\|_{s_{h}}^{q,\kappa} \\ &+ \|\Delta_{12}\Psi_{m}\|_{s_{h}}^{q,\kappa} \|U_{m}^{[1]}\|_{s_{0}}^{q,\kappa} + \|\Sigma_{m}^{[2]}\|_{s_{0}}^{q,\kappa} \|\Delta_{12}U_{m}\|_{s_{h}}^{q,\kappa} \\ &+ \|\Sigma_{m}^{[2]}\|_{s_{h}}^{q,\kappa} \|\Delta_{12}U_{m}\|_{s_{0}}^{q,\kappa} \,. \end{split}$$
(9.373)

To estimate $U_m^{[1]}$ (to alleviate the notation we remove the subscript [1]) described by (9.368) we use Lemma 7.4 in order to get

$$\|U_m\|\|_{s_0}^{q,\kappa} \le \|\mathscr{R}_m\|\|_{s_0}^{q,\kappa} + \|\mathscr{R}_m\|\|_{s_0}^{q,\kappa}\|\Psi_m\|\|_{s_0}^{q,\kappa}$$
(9.374)

and

$$\| U_m \|_{s_h}^{q,\kappa} \leq \| \mathscr{R}_m \|_{s_h}^{q,\kappa} + \| \mathscr{R}_m \|_{s_0}^{q,\kappa} \| \Psi_m \|_{s_h}^{q,\kappa} + \| \mathscr{R}_m \|_{s_h}^{q,\kappa} \| \Psi_m \|_{s_0}^{q,\kappa} .$$

$$(9.375)$$

Applying (9.315),(9.282) and (9.328) together with (9.374) we obtain

$$|||U_m|||_{s_0}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2}N_0^{\mu_2}N_m^{-\mu_2}.$$
(9.376)

On the other hand, from the first estimate of (9.330), (9.315) and (9.282) we deduce that

$$\max_{k=1,2} \left\| \Psi_m^{[k]} \right\|_{s_h}^{q,\kappa} \lesssim N_m^{\tau_2 q+q} \delta_m(s_h)$$
$$\lesssim N_m^{\tau_2 q+q} \varepsilon \overline{\gamma}^{-3}.$$
(9.377)

Thus we deduce from (9.375), (9.315) and (9.278)

$$\sup_{m \in \mathbb{N}} |||U_m|||_{s_h}^{q,\kappa} \leqslant C \varepsilon \kappa^{-2}.$$
(9.378)

Inserting (9.376) and (9.378) into (9.373) yields

$$\begin{aligned} \|\Delta_{12}\mathscr{R}_{m+1}\|\|_{s_{h}}^{q,\kappa} &\leq \|\Delta_{12}U_{m}\|\|_{s_{h}}^{q,\kappa} + C\varepsilon\kappa^{-2}\|\Delta_{12}\Psi_{m}\|\|_{s_{0}}^{q,\kappa} \\ &+ C\varepsilon\kappa^{-2}N_{0}^{\mu_{2}}N_{m}^{-\mu_{2}}\|\|\Delta_{12}\Psi_{m}\|\|_{s_{h}}^{q,\kappa} + \|\sum_{m}^{[2]}\|\|_{s_{0}}^{q,\kappa}\||\Delta_{12}U_{m}\|\|_{s_{h}}^{q,\kappa} \\ &+ \|\sum_{m}^{[2]}\|\|_{s_{h}}^{q,\kappa}\|\|\Delta_{12}U_{m}\|\|_{s_{0}}^{q,\kappa} . \end{aligned}$$
(9.379)

From (9.317) and (9.282) we get

$$\||\Sigma_{m}^{[2]}|||_{s_{0}}^{q,\kappa} \leqslant C\varepsilon\kappa^{-3}N_{0}^{\mu_{2}}N_{m}^{-\mu_{2}} \quad \text{and} \quad \||\Sigma_{m}^{[2]}|||_{s_{h}}^{q,\kappa} \leqslant C\varepsilon\kappa^{-3}N_{m}^{\tau_{2}q+q}.$$
(9.380)

Plugging (9.380) into (9.379) implies

$$\| \Delta_{12} \mathscr{R}_{m+1} \|_{s_h}^{q,\kappa} \leq \left(1 + C\varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \right) \| \Delta_{12} U_m \|_{s_h}^{q,\kappa} + C\varepsilon \kappa^{-3} N_m^{\tau_2 q+q} \| \Delta_{12} U_m \|_{s_0}^{q,\kappa} + C\varepsilon \kappa^{-2} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \Psi_m \|_{s_h}^{q,\kappa} + C\varepsilon \kappa^{-2} \| \Delta_{12} \Psi_m \|_{s_0}^{q,\kappa} .$$

$$(9.381)$$

Similarly, by putting together (9.376), (9.380) and (9.372) we infer

$$\|\Delta_{12}\mathscr{R}_{m+1}\|\|_{s_0}^{q,\kappa} \leq \left(1 + C\varepsilon\kappa^{-3}N_0^{\mu_2}N_m^{-\mu_2}\right) \|\|\Delta_{12}U_m\|\|_{s_0}^{q,\kappa} + C\varepsilon\kappa^{-2}N_0^{\mu_2}N_m^{-\mu_2} \|\|\Delta_{12}\Psi_m\|\|_{s_0}^{q,\kappa}.$$
(9.382)

Coming back to (9.369) and using Lemma 7.4 together with Lemma 7.3 we get $\forall s \in [s_0, s_h]$,

$$\begin{split} \||\Delta_{12}U_m|||_s^{q,\kappa} &\leq N_m^{s-s_h} \||\Delta_{12}\mathscr{R}_m||_{s_h}^{q,\kappa} + C \||\Delta_{12}\mathscr{R}_m||_s^{q,\kappa} \max_{k=1,2} \||\Psi_m^{[k]}|||_{s_0}^{q,\kappa} \\ &+ C \||\Delta_{12}\mathscr{R}_m||_{s_0}^{q,\kappa} \max_{k=1,2} \||\Psi_m^{[k]}||_s^{q,\kappa} + C \||\Delta_{12}\Psi_m||_s^{q,\kappa} \max_{k=1,2} \|\mathscr{R}_m^{[k]}\||_{s_0}^{q,\kappa} \\ &+ C \|\Delta_{12}\Psi_m||_{s_0}^{q,\kappa} \max_{k=1,2} \|\mathscr{R}_m^{[k]}||_s^{q,\kappa} \,. \end{split}$$

Putting together the preceding estimate with (9.377), (9.315) and (9.328)

$$\| \Delta_{12} U_m \|_{s_0}^{q,\kappa} \leqslant N_m^{s_0 - s_h} \| \Delta_{12} \mathscr{R}_m \|_{s_h}^{q,\kappa} + C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \mathscr{R}_m \|_{s_0}^{q,\kappa} + C \varepsilon \kappa^{-2} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \Psi_m \|_{s_0}^{q,\kappa}$$

$$(9.383)$$

and

$$\begin{split} \|\Delta_{12}U_{m}\|\|_{s_{h}}^{q,\kappa} &\leq \left(1 + C\varepsilon\kappa^{-3}N_{0}^{\mu_{2}}N_{m}^{-\mu_{2}}\right) \|\|\Delta_{12}\mathscr{R}_{m}\|\|_{s_{h}}^{q,\kappa} + CN_{m}^{\tau_{2}q+q}\varepsilon\kappa^{-3} \|\|\Delta_{12}\mathscr{R}_{m}\|\|_{s_{0}}^{q,\kappa} \\ &+ C\varepsilon\kappa^{-2}N_{0}^{\mu_{2}}N_{m}^{-\mu_{2}} \|\|\Delta_{12}\Psi_{m}\|\|_{s_{h}}^{q,\kappa} + C\varepsilon\kappa^{-2} \|\|\Delta_{12}\Psi_{m}\|\|_{s_{0}}^{q,\kappa} \,. \end{split}$$

Combining the foregoing estimate with (9.381), (9.382), (9.2) and (9.278) allows to get

$$\| \Delta_{12} \mathscr{R}_{m+1} \|_{s_h}^{q,\kappa} \leq \left(1 + C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} + C \varepsilon \kappa^{-3} N_m^{s_0 - s_h + \tau_2(1+q)} \right) \| \Delta_{12} \mathscr{R}_m \|_{s_h}^{q,\kappa} + C N_m^{\tau_2 q + q} \varepsilon \kappa^{-3} \| \Delta_{12} \mathscr{R}_m \|_{s_0}^{q,\kappa} + C \varepsilon \kappa^{-2} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \Psi_m \|_{s_h}^{q,\kappa} + C \varepsilon \kappa^{-2} \| \Delta_{12} \Psi_m \|_{s_0}^{q,\kappa} .$$

$$(9.384)$$

Similarly, by putting together (9.382), (9.383) and (9.278) we infer

$$\| \Delta_{12} \mathscr{R}_{m+1} \|_{s_0}^{q,\kappa} \leq N_m^{s_0-s_h} \| \Delta_{12} \mathscr{R}_m \|_{s_h}^{q,\kappa} + C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \mathscr{R}_m \|_{s_0}^{q,\kappa} + C \varepsilon \kappa^{-2} N_0^{\mu_2} N_m^{-\mu_2} \| \Delta_{12} \Psi_m \|_{s_0}^{q,\kappa} .$$

$$(9.385)$$

In what follows we intend to estimate $\Delta_{12}\Psi_m$ which is slightly delicate. According to (9.297), (9.290) and (9.291) we may write

$$\||\Delta_{12}\Psi_m\||_s^{q,\kappa} = \left(\max_{|\gamma| \le q} \sup_{\lambda \in \mathcal{O}} \sup_{n \in \mathbb{Z}} \sum_{\substack{|l| \le N_m \\ |k| \le N_m}} \kappa^{2|\gamma|} \langle l, k \rangle^{2(s-|\gamma|)} \left| \partial_{\lambda}^{\gamma} \Delta_{12} \Psi_n^{n+k}(\lambda, l) \right|^2 \right)^{\frac{1}{2}}$$
(9.386)

with

$$\Psi_{j_0}^j(\lambda, l) = \begin{cases} -\varrho_{j_0}^j(\lambda, l) \ r_{j_0}^j(\lambda, l), & \text{if} \quad (l, j) \neq (0, j_0) \\ 0, & \text{if} \quad (l, j) = (0, j_0), \end{cases}$$
(9.387)

and

$$\varrho_{j_0}^j(\lambda,l) \triangleq \frac{\chi\left((\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda))(\kappa\langle j - j_0\rangle)^{-1}\langle l\rangle^{\tau_2}\right)}{\omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda)}.$$
(9.388)

Remind from (9.288) that $(r_{j_0}^j(\lambda, l))$ are the Fourier coefficients of $P_{N_m}\mathscr{R}_m$, that is,

$$\operatorname{i} r_{j_0}^j(\lambda, l) = \left\langle P_{N_m} \mathscr{R}_m \mathbf{e}_{0, j_0}, \mathbf{e}_{l, j} \right\rangle_{L^2(\mathbb{T}^{d+1})}.$$

From this latter identity we get

$$i \Delta_{12} r_{j_0}^j(\lambda, l) = \left\langle \Delta_{12} P_{N_m} \mathscr{R}_m \mathbf{e}_{0, j_0}, \mathbf{e}_{l, j} \right\rangle_{L^2(\mathbb{T}^{d+1})}.$$

Now we need to estimate $\Delta_{12} \varrho_{j_0}^j(\lambda, l)$. For this aim, we write

$$\varrho_{j_0}^j(\lambda,l) = b_l^j \chi_1(b_l^j B_{l,j_0}^j(\lambda))$$

with

$$B_{l,j_0}^j(\lambda) = \omega \cdot l + \mu_j^m(\lambda) - \mu_{j_0}^m(\lambda), \quad b_l^j = (\kappa \langle j \rangle)^{-1} \langle l \rangle^{\tau_2}, \quad \chi_1(x) = \frac{\chi(x)}{x}.$$
(9.389)

According to (9.296) one gets

$$\forall |\gamma| \leqslant q, \quad \sup_{\lambda \in \mathcal{O}} \left| \partial_{\lambda}^{\gamma} \varrho_{j_0}^{j}(\lambda, l) \right| \leqslant C \kappa^{-(|\gamma|+1)} \langle l, j - j_0 \rangle^{\tau_2(1+|\gamma|)+|\gamma|}. \tag{9.390}$$

Then using Taylor formula in a similar way to (9.117) we find

$$\Delta_{12}\varrho_{j_0}^j(\lambda,l) = (\Delta_{12}B_{l,j_0}^j) (b_l^j)^2 \int_0^1 \chi_1' \left(b_l^j \left[(1-\tau)B_{l,j_0}^{j,[1]} + \tau B_{l,j_0}^{j,[2]} \right] \right) d\tau.$$
(9.391)

By (9.326) combined with Lemma 4.5 we deduce from straightforward computations that

$$(b_l^j)^2 \left\| \chi_1' \left(b_l^j \left[(1-\tau) B_{l,j_0}^{j,[1]} + \tau B_{l,j_0}^{j,[2]} \right] \right) \right\|^{q,\kappa} \lesssim \kappa^{-2} \langle l, j-j_0 \rangle^{q+\tau_2(q+2)}.$$
(9.392)

Next, we shall move to the estimate of $\Delta_{12}B_{l,j_0}^j$. For this purpose we use (9.310) leading to

$$\mu_j^m = \mu_j^0 + \sum_{n=0}^{m-1} \left\langle P_{N_n} \mathscr{R}_n \mathbf{e}_{0,j}, \mathbf{e}_{0,j} \right\rangle_{L^2(\mathbb{T}^{d+1})}.$$
(9.393)

We recall from Proposition 9.5 and the Gamma quotient introduced in Lemma 3.1

$$\mu_j^0(\lambda, i_0) \triangleq \Omega_j(\alpha) + jr^1(\lambda, i_0) - \mathsf{W}(j, \alpha)r^2(\lambda, i_0).$$

Therefore

$$\Delta_{12}\mu_{j}^{m} = \Delta_{12}\mu_{j}^{0} + \sum_{n=0}^{m-1} \left\langle P_{N_{n}}\Delta_{12}\mathscr{R}_{n}\mathbf{e}_{0,j}, \mathbf{e}_{0,j} \right\rangle_{L^{2}(\mathbb{T}^{d+1})}.$$
(9.394)

Combining (9.389) with (9.394) and (9.393) allows to get

$$\begin{split} \Delta_{12}B_{l,j_0}^{j} &= \Delta_{12}(\mu_{j}^{m} - \mu_{j_0}^{m}) \\ &= (j - j_0)\Delta_{12}r^{1} - \left(\mathbb{W}(j,\alpha) - \mathbb{W}(j_0,\alpha)\right)\Delta_{12}r^{2} \\ &+ \sum_{n=0}^{m-1} \left\langle P_{N_n}\Delta_{12}\mathscr{R}_{n}\mathbf{e}_{0,j}, \mathbf{e}_{0,j}\right\rangle_{L^{2}(\mathbb{T}^{d+1})} - \sum_{n=0}^{m-1} \left\langle P_{N_n}\Delta_{12}\mathscr{R}_{n}\mathbf{e}_{0,j_0}, \mathbf{e}_{0,j_0}\right\rangle_{L^{2}(\mathbb{T}^{d+1})}. \end{split}$$

It follows from Proposition 9.5-(i), Lemma 7.3-(i) and Lemma A.1

$$\left\|\Delta_{12}B_{l,j_0}^{j}\right\|^{q,\kappa} \lesssim |j-j_0|\varepsilon\kappa^{-1}\|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} + \sum_{n=0}^{m-1} \|P_{N_n}\Delta_{12}\mathscr{R}_n\|_{s_0}^{q,\kappa}.$$
(9.395)

Combining (9.391), (9.392), (9.395) together with the law products in Lemma 4.1 we find

$$\|\Delta_{12}\varrho_{j_{0}}^{j}(\cdot,l)\|^{q,\kappa} \lesssim \varepsilon \kappa^{-3} \langle l,j-j_{0} \rangle^{q+\tau_{2}(q+2)+1} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa}$$

$$+ \kappa^{-2} \langle l,j-j_{0} \rangle^{q+\tau_{2}(q+2)} \sum_{n=0}^{m-1} \|P_{N_{n}}\Delta_{12}\mathscr{R}_{n}\|\|_{s_{0}}^{q,\kappa}.$$
(9.396)

By the law products (or Leibniz rule) we find from (9.387)

$$\begin{aligned} \|\Delta_{12}\Psi_{n}^{n+k}(\cdot,l)\|_{W^{q,\infty}} &\lesssim \|\Delta_{12}\varrho_{n}^{n+k}(\cdot,l)\|_{W^{q,\infty}} \max_{\ell=1,2} \|(r_{n}^{n+k})^{[\ell]}(\cdot,l)\|_{W^{q,\infty}} \\ &+ \|\Delta_{12}r_{n}^{n+k}(\cdot,l)\|_{W^{q,\infty}} \max_{\ell=1,2} \|(\varrho_{n}^{n+k})^{[\ell]}(\cdot,l)\|_{W^{q,\infty}}. \end{aligned}$$

Putting together (9.396), (9.400) and using (9.278)

$$\begin{split} \|\Delta_{12}\Psi_{n}^{n+k}(\cdot,l)\|_{W^{q,\infty}} &\lesssim \varepsilon \kappa^{-3-q} \langle l,k \rangle^{q+\tau_{2}(q+2)+1} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \max_{\ell=1,2} \|(r_{n}^{n+k})^{[\ell]}(\cdot,l)\|_{W^{q,\infty}} \\ &+ \kappa^{-q-2} \langle l,k \rangle^{q+\tau_{2}(q+2)} \sum_{n=0}^{m-1} \|P_{N_{n}}\Delta_{12}\mathscr{R}_{n}\|_{s_{0}}^{q,\kappa} \max_{\ell=1,2} \|(r_{n}^{n+k})^{[\ell]}(\cdot,l)\|_{W^{q,\infty}} \\ &+ \kappa^{-(q+1)} \langle l,k \rangle^{\tau_{2}(1+q)+q} \|\Delta_{12}r_{n}^{n+k}(\cdot,l)\|_{W^{q,\infty}}. \end{split}$$

Notice that in the preceding estimate we can replace q by any $|\gamma| \leq q$ and get after a slight modification the estimate

$$\begin{split} |\partial_{\lambda}^{\gamma} \Delta_{12} \Psi_{n}^{n+k}(\lambda,l)| &\lesssim \varepsilon \kappa^{-3-|\gamma|} \left\langle l,k \right\rangle^{|\gamma|+\tau_{2}(|\gamma|+2)+1} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \max_{\ell=1,2} \max_{|\beta| \leqslant |\gamma|} |\partial_{\lambda}^{\beta}(r_{n}^{n+k})^{[\ell]}(\lambda,l)| \\ &+ \kappa^{-|\gamma|-2} \left\langle l,k \right\rangle^{|\gamma|+\tau_{2}(|\gamma|+2)} \sum_{n=0}^{m-1} \|P_{N_{n}} \Delta_{12} \mathscr{R}_{n}\|_{s_{0}}^{q,\kappa} \max_{\ell=1,2} \max_{|\beta| \leqslant |\gamma|} |\partial_{\lambda}^{\beta}(r_{n}^{n+k})^{[\ell]}(\lambda,l)| \\ &+ \kappa^{-(|\gamma|+1)} \left\langle l,k \right\rangle^{\tau_{2}(1+|\gamma|)+|\gamma|} \max_{|\beta| \leqslant |\gamma|} |\partial_{\lambda}^{\beta} \Delta_{12} r_{n}^{n+k}(\lambda,l)|. \end{split}$$

Inserting this estimate into (9.386) yields

$$\begin{split} \| \Delta_{12} \Psi_m \|_{s}^{q,\kappa} &\lesssim \varepsilon \kappa^{-3-q} \| \Delta_{12} i \|_{\overline{s}_h + \sigma}^{q,\kappa} \| P_{N_m} \mathscr{R}_m \|_{s + \tau_2(q+2)+1}^{q,\kappa} \\ &+ \kappa^{-q-2} \| P_{N_m} \mathscr{R}_m \|_{s + \tau_2(q+2)}^{q,\kappa} \sum_{n=0}^{m-1} \| P_{N_n} \Delta_{12} \mathscr{R}_n \|_{s_0}^{q,\kappa} \\ &+ \kappa^{-q-1} \| \Delta_{12} P_{N_m} \mathscr{R}_m \|_{s + \tau_2(1+q)}^{q,\kappa} . \end{split}$$

By setting

$$\varkappa_m(s) \triangleq \kappa^{-1} \sum_{n=0}^{m-1} \||\Delta_{12} \mathscr{R}_n\||_s^{q,\kappa}$$
(9.397)

and using (9.314) we get successively

$$\begin{aligned} \|\Delta_{12}\Psi_m\|\|_{s_0}^{q,\kappa} &\lesssim \varepsilon \kappa^{-2-q} N_m^{\tau_2+1} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa} \delta_m(\overline{s}_l) + \kappa^{-q-1} N_m^{\tau_2} \delta_m(\overline{s}_l) \varkappa_m(s_0) \\ &+ \kappa^{-q-1} N_m^{\tau_2(1+q)} \|\|\Delta_{12}\mathscr{R}_m\|\|_{s_0}^{q,\kappa} \end{aligned}$$
(9.398)

and

$$\|\Delta_{12}\Psi_{m}\|_{s_{h}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2-q} N_{m}^{\tau_{2}(q+2)+1} \delta_{m}(s_{h}) \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} + \kappa^{-q-1} N_{m}^{\tau_{2}(q+2)} \delta_{m}(s_{h}) \varkappa_{m}(s_{0})$$

$$+ \kappa^{-q-1} N_{m}^{\tau_{2}(1+q)} \| \Delta_{12}\mathscr{R}_{m} \|_{s_{h}}^{q,\kappa} .$$
 (9.399)

According to (9.315), (9.282) and (9.278) one obtains

$$\delta_m(\overline{s}_l) \leqslant C \varepsilon \kappa^{-3} N_0^{\mu_2} N_m^{-\mu_2} \quad \text{and} \quad \delta_m(s_h) \leqslant C \varepsilon \kappa^{-3} \lesssim 1 \tag{9.400}$$

Putting together (9.398) and (9.400) and using (9.278) yields

$$\|\Delta_{12}\Psi_{m}\|\|_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} N_{m}^{\tau_{2}+1-\mu_{2}} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} + \kappa^{-1} N_{m}^{\tau_{2}-\mu_{2}} \varkappa_{m}(s_{0}) + \kappa^{-q-1} N_{m}^{\tau_{2}(1+q)} \|\|\Delta_{12}\mathscr{R}_{m}\|\|_{s_{0}}^{q,\kappa}.$$
(9.401)

In a similar way, one gets by plugging (9.400) into (9.399)

$$\|\Delta_{12}\Psi_{m}\|\|_{s_{h}}^{q,\kappa} \lesssim \varepsilon \kappa^{-2-q} N_{m}^{\tau_{2}(q+2)+1} \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} + \varepsilon \kappa^{-q-4} N_{m}^{\tau_{2}(q+2)} \varkappa_{m}(s_{0})$$

$$+ \kappa^{-q-1} N_{m}^{\tau_{2}(1+q)} \|\|\Delta_{12}\mathscr{R}_{m}\|\|_{s_{h}}^{q,\kappa}$$
 (9.402)

Inserting (9.401) into (9.385) yields by virtue of (9.315) and (9.278)

$$\| \Delta_{12} \mathscr{R}_{m+1} \|_{s_0}^{q,\kappa} \leq N_m^{s_0-s_h} \| \Delta_{12} \mathscr{R}_m \|_{s_h}^{q,\kappa} + C \varepsilon \kappa^{-1} N_m^{\tau_2(q+1)-2\mu_2} \| \Delta_{12} i \|_{\overline{s}_h+\sigma}^{q,\kappa} + C N_m^{\tau_2(q+1)-\mu_2} \| \Delta_{12} \mathscr{R}_m \|_{s_0}^{q,\kappa} + C N_m^{\tau_2-2\mu_2} \varkappa_m(s_0).$$

$$(9.403)$$

Similarly, by inserting (9.398), (9.399) into (9.384) and using (9.278) we find

$$\| \Delta_{12} \mathscr{R}_{m+1} \|_{s_h}^{q,\kappa} \leq \left(1 + C N_m^{-\mu_2 + \tau_2(q+1)} + C N_m^{s_0 - s_h + \tau_2(q+1)} \right) \| \Delta_{12} \mathscr{R}_m \|_{s_h}^{q,\kappa} + C \varepsilon \kappa^{-3} N_m^{\tau_2(1+q)+q} \| \Delta_{12} \mathscr{R}_m \|_{s_0}^{q,\kappa} + C \varepsilon \kappa^{-3} N_m^{\tau_2(q+2)-\mu_2} \varkappa_m(s_0) . + C \varepsilon \kappa^{-1} \| \Delta_{12} i \|_{\overline{s_h} + \sigma}^{q,\kappa} N_m^{\tau_2(q+2)+1-\mu_2} .$$

$$(9.404)$$

Introduce

$$\overline{\delta}_m(s) \triangleq \kappa^{-1} \parallel \! \mid \Delta_{12} \mathscr{R}_m \parallel \! \mid_s^{q,\kappa}$$

Then (9.403) and (9.404) become

$$\overline{\delta}_{m+1}(s_0) \leqslant N_m^{s_0 - s_h} \overline{\delta}_m(s_h) + C \varepsilon \kappa^{-2} \parallel \Delta_{12} i \parallel_{\overline{s}_h + \sigma}^{q, \kappa} N_m^{\tau_2(q+1) - 2\mu_2} + C N_m^{\tau_2(q+1) - \mu_2} \overline{\delta}_m(s_0) + C N_m^{\tau_2 - 2\mu_2} \varkappa_m(s_0)$$
(9.405)

and

$$\overline{\delta}_{m+1}(s_h) \leqslant \left(1 + CN_m^{-\mu_2 + \tau_2(q+1)} + CN_m^{s_0 - s_h + \tau_2(q+1)}\right) \overline{\delta}_m(s_h) + C\varepsilon\kappa^{-3}N_m^{\tau_2(q+1) + q}\overline{\delta}_m(s_0) + CN_m^{\tau_2(q+2) - \mu_2} \varkappa_m(s_0) + C\varepsilon\kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h + \sigma}^{q,\kappa} N_m^{\tau_2(q+2) + 1 - \mu_2}.$$
(9.406)

It follows from (9.397) that for any $s \ge s_0$

$$\varkappa_m(s) \leqslant \sum_{k=0}^{m-1} \overline{\delta}_k(s). \tag{9.407}$$

We shall prove by induction in $m \in \mathbb{N}$ that

$$\forall k \leqslant m, \quad \overline{\delta}_k(s_0) \leqslant N_0^{\mu_2} N_k^{-\mu_2} \nu(s_h) \quad \text{and} \quad \overline{\delta}_k(s_h) \leqslant \left(2 - \frac{1}{k+1}\right) \nu(s_h), \tag{9.408}$$

with

$$\nu(s) \triangleq \overline{\delta}_0(s) + \varepsilon \kappa^{-2} \parallel \Delta_{12} i \parallel_{\overline{s}_h + \sigma}^{q,\kappa}.$$
(9.409)

The validity of (9.408) for m = 0 is obvious from Sobolev embeddings. Now let us assume that the property (9.408) holds true at the order m and let us check it at the order m + 1. Then from (9.407) and Lemma A.2 we find an absolute constant C > 0 such that

$$\varkappa_m(s_0) \leqslant C\nu(s_h)$$

Combining this estimate with the induction assumption, (9.405) and (9.406) implies

$$\overline{\delta}_{m+1}(s_0) \leqslant 2N_m^{s_0-s_h}\nu(s_h) + CN_m^{\tau_2(q+1)-2\mu_2}\nu(s_h) + CN_0^{\mu_2}N_m^{\tau_2(q+1)-2\mu_2}\nu(s_h)$$
$$\leqslant 2N_m^{s_0-s_h}\nu(s_h) + CN_0^{\mu_2}N_m^{\tau_2(q+1)-2\mu_2}\nu(s_h)$$
(9.410)

and

$$\overline{\delta}_{m+1}(s_h) \leqslant \left(1 + CN_m^{-\mu_2 + \tau_2(q+1)} + CN_m^{s_0 - s_h + \tau_2(q+1)}\right) \left(2 - \frac{1}{m+1}\right) \nu(s_h) + CN_m^{\tau_2(q+1) + q - \mu_2} \nu(s_h) + CN_m^{\tau_2(q+2) - \mu_2} \nu(s_h) + CN_m^{\tau_2(q+2) + 1 - \mu_2} \nu(s_h) \leqslant \left(1 + CN_m^{-\mu_2 + \tau_2(q+1)}\right) \left(2 - \frac{1}{m+1}\right) \nu(s_h) + CN_m^{\tau_2(q+2) + q - \mu_2} \nu(s_h)$$

where in the last inequality we have used that $s_h \ge s_0 + \mu_2$ according to (9.277). Then with the choice of μ_2 made in (9.277) we deduce in a standard way that

$$\overline{\delta}_{m+1}(s_0) \leqslant N_0^{\mu_2} N_{m+1}^{-\mu_2} \nu(s_h) \quad \text{and} \quad \overline{\delta}_{m+1}(s_h) \leqslant \left(2 - \frac{1}{m+2}\right) \nu(s_h)$$

which achieves the induction. Notice that one can fix in (9.408) the free numbers μ_2 and s_h at their lower bounds given in (9.277), that is, in view of the notations (9.38),

$$\mu_2 = \overline{\mu}_2 + 2\tau_2(1+q) \triangleq \mu_c$$

$$s_h = \frac{3}{2}\overline{\mu}_2 + s_l + 1 + 4\tau_2(1+q) = \overline{s}_h + 4\tau_2(1+q) \triangleq s_c$$

Therefore, one finds in particular

$$\overline{\delta}_m(s_0) \leqslant N_0^{\mu_c} N_m^{-\mu_c} \nu(s_c). \tag{9.411}$$

The next task is to estimate $\Delta_{12}r_j^{\infty}$. Then proceeding as for (9.324) we get by using a duality argument, Lemma 7.3, (9.411) and Lemma A.2

$$\begin{split} \|\Delta_{12}r_{j}^{\infty}\|^{q,\kappa} &\leq \sum_{m=0}^{\infty} \|\langle (P_{N_{m}}\Delta_{12}\mathscr{R}_{m})\mathbf{e}_{0,j}, \mathbf{e}_{0,j}\rangle\|^{q,\kappa} \\ &\leq C\sum_{m=0}^{\infty} \|\Delta_{12}\mathscr{R}_{m}\|\|_{s_{0}}^{q,\kappa} \leq C\kappa\nu(s_{c})\sum_{m=0}^{\infty} N_{0}^{\mu_{c}}N_{m}^{-\mu_{c}} \\ &\leq C\kappa\nu(s_{c}). \end{split}$$

Thus we get from (9.409)

$$\|\Delta_{12}r_j^{\infty}\|^{q,\kappa} \lesssim \|\Delta_{12}\mathscr{R}_0\|\|_{s_c}^{q,\kappa} + \varepsilon \kappa^{-1} \|\|\Delta_{12}i\|\|_{\overline{s}_h+\sigma}^{q,\kappa} .$$

$$(9.412)$$

Combining (9.352) with Proposition 9.5-(i) and using Sobolev embeddings we obtain

$$\forall j \in \mathbb{S}_{0}^{c}, \quad \|\Delta_{12}\mu_{j}^{\infty}\|^{q,\kappa} \leqslant \|\Delta_{12}\mu_{j}^{0}\|^{q,\kappa} + \|\Delta_{12}r_{j}^{\infty}\|^{q,\kappa}$$
$$\lesssim \|\Delta_{12}\mathscr{R}_{0}\|_{s_{c}}^{q,\kappa} + \varepsilon \kappa^{-1}|j| \|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa}.$$
(9.413)

Recall that the operator \mathscr{R}_0 coincides with $\mathscr{R}^2_{r,\lambda}$ defined in Proposition 9.5. Then from Proposition 9.5-(iii) and (9.278) we infer

$$\||\Delta_{12}\mathscr{R}_0||_{0,0,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \|\Delta_{12}i\|_{\overline{s}_h+\sigma}^{q,\kappa}$$

and

$$\||\mathscr{R}_0|||_{0,s_h,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2}.$$

From the latter inequality we easily get by applying the triangle inequality, since $s_h \geqslant 2s_c$

$$\||\Delta_{12}\mathscr{R}_0|||_{0,2s_c,0}^{q,\kappa} \lesssim \varepsilon \kappa^{-2}.$$

Applying the interpolation inequality of Lemma 7.3-(v) we get

$$\begin{split} \||\Delta_{12}\mathscr{R}_{0}||_{0,s_{c},0}^{q,\kappa} \lesssim \left(\||\Delta_{12}\mathscr{R}_{0}||_{0,0,0}^{q,\kappa} \right)^{\frac{1}{2}} \left(\||\Delta_{12}\mathscr{R}_{0}||_{0,2s_{c},0}^{q,\kappa} \right)^{\frac{1}{2}} \\ \lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12}i||_{\overline{s}_{h}+\sigma}^{q,\kappa} \right)^{\frac{1}{2}}. \end{split}$$

Plugging this estimate into (9.412) and (9.413) yields successively,

$$\begin{aligned} \forall j \in \mathbb{S}_{0}^{c}, \quad \|\Delta_{12}r_{j}^{\infty}\|^{q,\kappa} &\lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \right)^{\frac{1}{2}} + \varepsilon \kappa^{-1} \|\|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \\ &\lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \right)^{\frac{1}{2}} \end{aligned}$$

and

$$\begin{aligned} \forall j \in \mathbb{S}_{0}^{c}, \quad \|\Delta_{12}\mu_{j}^{\infty}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa}\right)^{\frac{1}{2}} + \varepsilon \kappa^{-1} |j| \parallel \Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa} \\ \lesssim \varepsilon \kappa^{-2} \left(\|\Delta_{12}i\|_{\overline{s}_{h}+\sigma}^{q,\kappa}\right)^{\frac{1}{2}} |j|. \end{aligned}$$

This ends the proof of Proposition 9.6.

9.6.4. Approximate inverse in the normal direction. The main concern of this section is to find an approximate right inverse of the linearized operator $\widehat{\mathscr{L}}_{\omega}$ defined in (9.1) and detailed in Proposition 9.1 when the set of parameters is restricted to a suitable Cantor like set obtained through successive excisions. The main result reads as follows.

Theorem 9.1. Let $(\gamma, q, d, \tau_1, \tau_2, s_0, s_h, S)$ satisfy (9.2), (9.38), (9.277) and assume the smallness condition (9.278). Then the following assertions hold true.

(i) Consider the operator L_∞ defined in Proposition 9.6, then there exists a family of linear operators (T_n)_{n∈ℕ} defined in the whole set 𝔅 with the estimate

$$\forall s \in [s_0, S], \quad \sup_{n \in \mathbb{N}} \|\mathbf{T}_n \rho\|_s^{q, \kappa} \lesssim \kappa^{-1} \|\rho\|_{s+\tau_1 q+\tau_1}^{q, \kappa}$$

and such that for any $n \in \mathbb{N}$ and in the Cantor set

$$\Lambda_{\infty,n}^{\kappa,\tau_1}(i_0) = \bigcap_{(l,j)\in\mathbb{Z}^d\times\mathbb{S}_0^c} \left\{ \lambda \in \mathcal{O} \, ; \, \forall \, |l| \leqslant N_n, \, j \in \mathbb{S}_0^c, \, |\omega \cdot l + \mu_j^\infty(\lambda,i_0)| > \frac{\kappa\langle j \rangle}{\langle l \rangle^{\tau_1}} \right\}$$

we have

$$\mathscr{L}_{\infty} \mathsf{T}_n = \mathrm{Id} + \mathsf{E}_n^4$$

with

$$\forall s \in [s_0, S], \qquad \|\mathbf{E}_n^4 h\|_{s_0}^{q,\kappa} \lesssim N_n^{s_0-s} \, \kappa^{-1} \, \|h\|_{s+\tau_1 q+\tau_1+1}^{q,\kappa}.$$

(ii) There exist
$$\sigma_5 = \sigma(\tau_1, \tau_2, d, q) \ge \sigma_4$$
 and a family of linear operators $(\mathsf{T}_{\omega,n})_n$ satisfying

$$\forall s \in [s_0, S], \quad \sup_{n \in \mathbb{N}} \|\mathsf{T}_{\omega, n}h\|_s^{q, \kappa} \lesssim \kappa^{-1} \left(\|h\|_{s+\sigma_5}^{q, \kappa} + \|\mathfrak{I}_0\|_{s+\sigma_5}^{q, \kappa} \|h\|_{s_0+\sigma_5}^{q, \kappa} \right)$$
(9.414)

and such that in the Cantor set

$$\mathsf{G}_n(\kappa,\tau_1,\tau_2,i_0) \triangleq \mathscr{O}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0) \cap \Lambda_{\infty,n}^{\kappa,\tau_1}(i_0)$$

we have

$$\widehat{\mathscr{L}}_{\omega}\mathsf{T}_{\omega,n} = \mathrm{Id} + \mathsf{E}_n,$$

with the estimates

$$\forall s \in [s_0, S], \quad \|\mathbf{E}_n h\|_{s_0}^{q,\kappa} \lesssim N_n^{s_0 - s} \kappa^{-1} \Big(\|h\|_{s + \sigma_5}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{s + \sigma_5}^{q,\kappa} \|h\|_{s_0 + \sigma_5}^{q,\kappa} \Big) \\ + \varepsilon \kappa^{-4} N_0^{\mu_2} N_n^{-\mu_2} \|h\|_{s_0 + \sigma_5}^{q,\kappa},$$

where $\widehat{\mathcal{L}}_{\omega}, \mathbb{C}_{\infty,n}^{\kappa,\tau_1}(i_0)$ and $\mathbb{C}_{\infty,n}^{\kappa,\tau_1,\tau_2}(i_0)$ are defined in (9.1) and Proposition 9.2 and Proposition 9.6, respectively.

(iii) On the Cantor set $G_n(\kappa, \tau_1, \tau_2, i_0)$, we have the splitting

$$\widehat{\mathscr{L}}_{\omega} = \widehat{\mathsf{L}}_{\omega,n} + \widehat{\mathsf{R}}_n, \quad with \quad \widehat{\mathsf{L}}_{\omega,n}\mathsf{T}_{\omega,n} = \mathrm{Id} \quad and \quad \widehat{\mathsf{R}}_n = \mathsf{E}_n\widehat{\mathsf{L}}_{\omega}$$

where the operators $\widehat{L}_{\omega,n}$ and \widehat{R}_n are defined in the whole set \mathbb{G} with the estimates

$$\forall s \in [s_0, S], \quad \sup_{n \in \mathbb{N}} \|\widehat{\mathbf{L}}_{\omega, n}h\|_s^{q, \kappa} \lesssim \|h\|_{s+1}^{q, \kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{q, s+\sigma_5}^{q, \kappa} \|h\|_{s_0+1}^{q, \kappa}$$

and

$$\|\widehat{\mathbf{R}}_{n}h\|_{s_{0}}^{q,\kappa} \lesssim N_{n}^{s_{0}-s}\kappa^{-1}\Big(\|h\|_{s+\sigma_{5}}^{q,\kappa_{5}} + \varepsilon\kappa^{-3}\|\Im_{0}\|_{s+\sigma_{5}}^{q,\kappa_{5}}\|h\|_{s_{0}+\sigma_{5}}^{q,\kappa}\Big) + \varepsilon\kappa^{-4}N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}}\|h\|_{s_{0}+\sigma_{5}}^{q,\kappa_{5}}.$$

Proof. (i) From Proposition 9.6 we recall that

$$\mathscr{L}_{\infty}(\lambda,\omega,i_0) = \left(\omega \cdot \partial_{\varphi} + \mathscr{D}_{\infty}(\lambda,i_0)\right) \Pi_{\mathbb{S}_0}^{\perp}.$$

Then we may split this operator as follows

$$\mathscr{L}_{\infty}(\lambda,\omega,i_{0}) = \Pi_{N_{n}}\omega \cdot \partial_{\varphi}\Pi_{N_{n}}\Pi_{\mathbb{S}_{0}}^{\perp} + \mathscr{D}_{\infty}(\lambda,i_{0})\Pi_{\mathbb{S}_{0}}^{\perp} - \Pi_{N_{n}}^{\perp}\omega \cdot \partial_{\varphi}\Pi_{N_{n}}^{\perp}\Pi_{\mathbb{S}_{0}}^{\perp} \triangleq \mathbf{L}_{n} - \mathbf{R}_{n},$$
(9.415)

with $\mathbf{R}_n = \prod_{N_n}^{\perp} \omega \cdot \partial_{\varphi} \prod_{N_n}^{\perp} \prod_{\mathbb{S}_0}^{\perp}$ and the projector \prod_{N_n} is defined by

$$\Pi_N \sum_{l,j} h_{l,j} \mathbf{e}_{l,j} = \sum_{\substack{|l| \leq N \\ j \in \mathbb{Z}}} h_{l,j} \mathbf{e}_{l,j}$$

From this definition and the structure of $\mathscr{D}_{\infty}(\lambda, i_0)$ in Proposition 9.6 we deduce that

$$\mathbf{e}_{-l,-j}\mathbf{L}_{n}\mathbf{e}_{l,j} = \begin{cases} \mathbf{i}\left(\omega \cdot l + \mu_{j}^{\infty}\right); & \text{if} \quad |l| \leq N_{n}, \quad j \in \mathbb{S}_{0}^{c} \\ \mathbf{i}\,\mu_{j}^{\infty}; & \text{if} \quad |l| > N_{n}, \quad j \in \mathbb{S}_{0}^{c}. \end{cases}$$

Define the diagonal operator \mathtt{T}_n by

$$\begin{aligned} \mathbf{T}_{n}h(\lambda,\varphi,\theta) &\triangleq -\mathbf{i} \sum_{\substack{|l| \leq N_{n} \\ j \in \mathbb{S}_{0}^{c}}} \frac{\chi\left((\omega \cdot l + \mu_{j}^{\infty}(\lambda))\kappa^{-1}\langle l\rangle^{\tau_{1}}\right)}{\omega \cdot l + \mu_{j}^{\infty}(\lambda)} h_{l,j}(\lambda) \,\mathbf{e}_{l,j}(\varphi,\theta) \\ &-\mathbf{i} \sum_{\substack{|l| > N_{n} \\ j \in \mathbb{S}_{0}^{c}}} \frac{h_{l,j}(\lambda)}{\mu_{j}^{\infty}(\lambda)} \,\mathbf{e}_{l,j}(\varphi,\theta) \end{aligned}$$

where χ is the cut-off function defined in (6.30) and $(h_{l,j}(\lambda))$ are the Fourier coefficients of h. We recall from Proposition 9.6 that

$$\mu_j^{\infty}(\lambda) = \mu_j^0(\lambda) + r_j^{\infty}(\lambda) \quad \text{with} \quad \sup_{j \in \mathbb{S}_0^c} \|r_j^{\infty}\|^{q,\kappa} \leqslant C \varepsilon \kappa^{-2}.$$

Combining this estimate with Proposition 9.5-(i) we find

$$\forall j \in \mathbb{S}_0^c, \, \|\mu_j^\infty\|^{q,\kappa} \leqslant C|j|$$

Similar arguments based on Lemma 3.1-(iv) give under the smallness condition (9.278)

$$|j| \lesssim \|\mu_j^\infty\|^{0,\kappa} \leqslant \|\mu_j^\infty\|^{q,\kappa}$$

Then implementing in part Lemma 9.2 yields

$$\forall s \ge s_0, \quad \|\mathbf{T}_n h\|_s^{q,\kappa} \lesssim \kappa^{-1} \|h\|_{s+\tau_1 q+\tau_1}^{q,\kappa}.$$

$$(9.416)$$

Moreover, by construction we get

$$\forall \lambda \in \Lambda_{\infty,n}^{\kappa,\tau_1}(i_0), \quad \mathbf{L}_n \mathbf{T}_n = \mathrm{Id}, \tag{9.417}$$

since $\chi(\cdot) = 1$ in the set $\Lambda_{\infty,n}^{\kappa,\tau_1}(i_0)$. It follows according to (9.415) that

$$\forall \lambda \in \Lambda_{\infty,n}^{\kappa,\tau_1}(i_0), \quad \mathscr{L}_{\infty} \mathsf{T}_n = \mathrm{Id} - \mathsf{R}_n \mathsf{T}_n \\ \triangleq \mathrm{Id} + \mathsf{E}_n^4.$$
 (9.418)

Notice that one gets from direct computations that

$$\forall s_0 \leqslant s \leqslant \overline{s}, \quad \|\mathbf{R}_n h\|_s^{q,\kappa} \lesssim N_n^{s-\overline{s}} \|h\|_{\overline{s+1}}^{q,\kappa}.$$

Combining this estimate with (9.416) yields

$$\forall s_0 \leqslant s \leqslant \overline{s}, \quad \|\mathbf{E}_n^4 h\|_s^{q,\kappa} \lesssim N_n^{s-\overline{s}} \|\mathbf{T}_n h\|_{\overline{s}+1}^{q,\kappa} \\ \lesssim N_n^{s-\overline{s}} \kappa^{-1} \|h\|_{\overline{s}+\tau_1 q+\tau_1+1}^{q,\kappa}.$$
(9.419)

This achieves the proof of the first point.

(ii) Let us define

$$\mathbf{T}_{\omega,n} \triangleq \Psi_{\perp} \Phi_{\infty} \mathbf{T}_n \Phi_{\infty}^{-1} \Psi_{\perp}^{-1}.$$
(9.420)

where the operators Ψ_{\perp} and Φ_{∞} are defined in Proposition 9.5 and Proposition 9.6, respectively. Notice that $T_{\omega,n}$ is defined in the whole range of parameters \mathcal{O} . Applying Lemma 9.6-(ii) combined with (9.278) yields

$$\forall s \geqslant s_0, \quad \|\mathsf{T}_{\omega,n}h\|_s^{q,\kappa} \lesssim \|\Phi_{\infty}\mathsf{T}_n\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}h\|_s^{q,\kappa} + \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa}\|\Phi_{\infty}\mathsf{T}_n\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}h\|_{s_0}^{q,\kappa}.$$

By using (9.279) and (9.278) one gets

$$\forall s \in [s_0, S], \quad \|\Phi_{\infty} \mathsf{T}_n \Phi_{\infty}^{-1} \Psi_{\perp}^{-1} h\|_s^{q,\kappa} \lesssim \|\mathsf{T}_n \Phi_{\infty}^{-1} \Psi_{\perp}^{-1} h\|_s^{q,\kappa} + \|\mathfrak{I}_0\|_{q,s+\sigma}^{q,\kappa} \|\mathsf{T}_n \Phi_{\infty}^{-1} \Psi_{\perp}^{-1} h\|_{s_0}^{q,\kappa}.$$

From (9.416) we find

$$\forall s \ge s_0, \quad \|\mathbf{T}_n \Phi_\infty^{-1} \Psi_\perp^{-1} h\|_s^{q,\kappa} \lesssim \kappa^{-1} \|\Phi_\infty^{-1} \Psi_\perp^{-1} h\|_{s+\tau_1 q+\tau_1}^{q,\kappa}$$

Applying (9.279) and (9.278) yields

$$\forall s \in [s_0, S], \quad \|\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}h\|_s^{q,\kappa} \lesssim \|\Psi_{\perp}^{-1}h\|_s^{q,\kappa} + \|\mathfrak{I}_0\|_{s+\overline{\sigma}}^{q,\kappa}\|\Psi_{\perp}^{-1}h\|_{s_0}^{q,\kappa}.$$

Putting together the preceding three estimates with Proposition 9.6-(ii) then we get (9.414). Now combining Propositions 9.5-9.6 we find that that in the Cantor set $\mathcal{O}_{\infty,n}^{\kappa,\tau_1}(i_0) \cap \mathcal{O}_{\infty,n}^{\kappa,\tau_2}(i_0)$

$$\begin{split} \Phi_{\infty}^{-1} \Psi_{\perp}^{-1} \widehat{\mathscr{L}}_{\omega} \Psi_{\perp} \Phi_{\infty} &= \Phi_{\infty}^{-1} \mathscr{L}_{0} \Phi_{\infty} + \Phi_{\infty}^{-1} \mathbf{E}_{n}^{2} \Phi_{\infty} \\ &= \mathscr{L}_{\infty} + \mathbf{E}_{n}^{3} + \Phi_{\infty}^{-1} \mathbf{E}_{n}^{2} \Phi_{\infty}. \end{split}$$

It follows that in the Cantor set $G_n(\kappa, \tau_1, \tau_2, i_0) = \mathcal{O}_{\infty,n}^{\kappa, \tau_1, \tau_2}(i_0) \cap \Lambda_{\infty,n}^{\kappa, \tau_1}(i_0)$ one has by virtue of (9.418) and the identity (9.420)

$$\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}\widehat{\mathscr{L}}_{\omega}\Psi_{\perp}\Phi_{\infty}\mathsf{T}_{n} = \mathrm{Id} + \mathsf{E}_{n}^{4} + \mathsf{E}_{n}^{3}\mathsf{T}_{n} + \Phi_{\infty}^{-1}\mathsf{E}_{n}^{2}\Phi_{\infty}\mathsf{T}_{n}$$

which gives in view of (9.420) the identity

$$\widehat{\mathscr{L}}_{\omega}\mathsf{T}_{\omega,n} = \mathrm{Id} + \Psi_{\perp}\Phi_{\infty}(\mathsf{E}_{n}^{4} + \mathsf{E}_{n}^{3}\mathsf{T}_{n} + \Phi_{\infty}^{-1}\mathsf{E}_{n}^{2}\Phi_{\infty}\mathsf{T}_{n})\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}
\triangleq \mathrm{Id} + \Psi_{\perp}\Phi_{\infty}\mathsf{E}_{n}^{5}\Phi_{\infty}^{-1}\Psi_{\perp}^{-1}
\triangleq \mathrm{Id} + \mathsf{E}_{n}$$
(9.421)

provided that $\lambda \in G_n(\kappa, \tau_1, \tau_2, i_0)$. The estimate of the first term of E_n^5 is given in (9.419). For the second term of E_n^5 we use the estimate (9.281) combined with (9.416) leading to

$$|\mathbf{E}_{n}^{3}\mathbf{T}_{n}h||_{s_{0}}^{q,\kappa} \lesssim \varepsilon \kappa^{-3} N_{0}^{\mu_{2}} N_{n}^{-\mu_{2}} \|\mathbf{T}_{n}h||_{s_{0}+1}^{q,\kappa} \lesssim \varepsilon \kappa^{-4} N_{0}^{\mu_{2}} N_{n}^{-\mu_{2}} \|h\|_{s_{0}+1+\tau_{1}(1+q)}^{q,\kappa}.$$

$$(9.422)$$

To estimate $\Phi_{\infty}^{-1} E_n^2 \Phi_{\infty} T_n$ we combine Proposition 9.5-(ii) with (9.279), (9.416) and (9.278)

$$\|\Phi_{\infty}^{-1}\mathsf{E}_{n}^{2}\Phi_{\infty}\mathsf{T}_{n}h\|_{s_{0}}^{q,\kappa} \leqslant C\varepsilon\kappa^{-2} \Big(N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}} + N_{n}^{s-s_{0}}\big(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\big)\Big)\|\mathsf{T}_{n}h\|_{s_{0}+3}^{q,\kappa} \\ \leqslant C\varepsilon\kappa^{-3}\Big(N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}} + N_{n}^{s-s_{0}}\big(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa}\big)\Big)\|h\|_{s_{0}+\tau_{1}(1+q)+3}^{q,\kappa}.$$
(9.423)

Similarly we get from (9.419), (9.422) and (9.423)

$$\begin{aligned} \|\mathbf{E}_{n}^{5}h\|_{s_{0}}^{q,\kappa} &\lesssim \varepsilon \kappa^{-4} \Big(N_{0}^{\mu_{2}} N_{n}^{-\mu_{2}} + N_{n}^{s-s_{0}} \big(1 + \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \big) \Big) \|h\|_{s_{0}+3+\tau_{1}(1+q)}^{q,\kappa} \\ &+ N_{n}^{s_{0}-s} \kappa^{-1} \|h\|_{s+1+\tau_{1}q+\tau_{1}}^{q,\kappa}. \end{aligned}$$

$$(9.424)$$

Set $\overline{\Psi} \triangleq \Psi_{\perp} \Phi_{\infty}$ then from (9.279), Proposition 9.6-(ii) and using (9.278) we deduce that

$$\forall s \in [s_0, S], \quad \|\overline{\Psi}^{\pm 1}h\|_s^{q,\kappa} \lesssim \|h\|_s^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{s+\sigma}^{q,\kappa} \|h\|_{s_0}^{q,\kappa}.$$
(9.425)

In particular, we get

$$\begin{split} \|\overline{\Psi}^{\pm 1}h\|_{s_0}^{q,\kappa} &\lesssim \|h\|_{s_0}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_0\|_{s_0+\sigma}^{q,\kappa} \|h\|_{s_0}^{q,\kappa} \\ &\lesssim \|h\|_{s_0}^{q,\kappa}. \end{split}$$
(9.426)

Thus we deduce from (9.426) combined with (9.424) and (9.278) we deduce

$$\lesssim \varepsilon \kappa^{-4} \Big(N_0^{\mu_2} N_n^{-\mu_2} + N_n^{s_0 - s} \big(1 + \| \mathfrak{I}_0 \|_{s + \sigma}^{q, \kappa} \big) \Big) \| h \|_{s_0 + 3 + \tau_1(1 + q)}^{q, \kappa} \\ + N_n^{s_0 - s} \kappa^{-1} \| h \|_{s + 1 + \tau_1(1 + q)}^{q, \kappa}.$$

Consequently we obtain from (9.278)

$$\sup_{n \in \mathbb{N}} \|\mathbf{E}_{n}h\|_{s_{0}}^{q,\kappa} \lesssim N_{n}^{s_{0}-s}\kappa^{-1} \Big(\|h\|_{s+\sigma}^{q,\kappa} + \varepsilon\kappa^{-3}\|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}+\sigma}^{q,\kappa} \Big) + \varepsilon\kappa^{-4}N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}}\|h\|_{s_{0}+\sigma}^{q,\kappa}.$$
(9.427)

This achieves the proof of the second point.

(iii) This result follows easily from the preceding point (ii). Indeed, according to (9.421) one may write on the Cantor set $G_n(\kappa, \tau_1, \tau_2, i_0)$

$$\widehat{\mathscr{L}}_{\omega} = \mathsf{T}_{\omega,n}^{-1} + \mathsf{E}_n \mathsf{T}_{\omega,n}^{-1} \tag{9.428}$$

In addition, we get from (9.420) and (9.417)

$$\mathbf{T}_{\omega,n}^{-1} = \Psi_{\perp} \Phi_{\infty} \mathbf{L}_n \Phi_{\infty}^{-1} \Psi_{\perp}^{-1} \\ \triangleq \widehat{\mathbf{L}}_{\omega,n}.$$
(9.429)

This provides the splitting

$$\widehat{\mathscr{L}}_{\omega} = \widehat{L}_{\omega,n} + \widehat{R}_n \quad \text{and} \quad \widehat{R}_n \triangleq E_n \widehat{L}_{\omega,n}.$$

From (9.415) and (9.425) combined with Proposition 9.6 we obtain

$$|\widehat{\mathbf{L}}_{\omega,n}h||_{s}^{q,\kappa} \lesssim \|h\|_{s+1}^{q,\kappa} + \varepsilon \kappa^{-3} \|\mathfrak{I}_{0}\|_{s+\sigma}^{q,\kappa} \|h\|_{s_{0}+1}^{q,\kappa}.$$

Inserting this estimate into (9.427) yields

$$\forall s \in [s_0, S], \quad \sup_{n \in \mathbb{N}} \|\widehat{\mathbf{R}}_n h\|_{q,s}^{q,\kappa} \lesssim \kappa^{-1} \|h\|_{s+\sigma+1}^{q,\kappa} + \varepsilon \kappa^{-4} \|\mathfrak{I}_0\|_{q,s+\sigma}^{q,\kappa} \|h\|_{s_0+\sigma+1}^{q,\kappa}.$$

and

$$\sup_{n \in \mathbb{N}} \|\widehat{\mathbf{R}}_{n}h\|_{s_{0}}^{q,\kappa} \lesssim N_{n}^{s_{0}-s}\kappa^{-1} \Big(\|h\|_{s+\sigma+1}^{q,\kappa} + \varepsilon\kappa^{-3} \|\mathfrak{I}_{0}\|_{s+\sigma+1}^{q,\kappa} \|h\|_{s_{0}+\sigma+1}^{q,\kappa} \Big) \\ + \varepsilon\kappa^{-4}N_{0}^{\mu_{2}}N_{n}^{-\mu_{2}} \|h\|_{s_{0}+\sigma+1}^{q,\kappa}.$$

Thus changing $\sigma + 1$ with σ gives the suitable result.

10. Proof of the main result

The main concern of this section is to implement Nash-Moser scheme in order to construct zeros for the nonlinear functional \mathscr{F} defined in (5.21) when $\varepsilon > 0$ and small enough. We shall prove that solutions do exist provided that the parameters $\lambda = (\omega, \alpha)$ belong to a final Cantor set $\mathsf{G}_{\infty}^{\kappa}$. More precisely, we are able to construct smooth functions $\lambda \in \mathscr{O} \mapsto i_{\infty}(\lambda)$, $\mathsf{c}_{\infty}(\lambda)$ in $W^{q,\infty}(\mathscr{O})$ such that

$$\forall \lambda \in \mathsf{G}_{\infty}^{\kappa}, \quad \mathscr{F}(i_{\infty}(\lambda), \mathsf{c}_{\infty}(\lambda), \lambda, \varepsilon) = 0.$$
(10.1)

The next stage is to find solutions to the origininal Hamiltonian equation (5.1). For this aim, we should adjust the parameters in such a way that $c_{\infty}(\omega, \alpha) = -\omega_{Eq}(\alpha)$, where the latter quantity corresponds to the equilibrium frequency defined in (5.7). This equation is invertible in ω close to the equilibrium frequencies and ω becomes an implicit function of α . Consequently we generate solutions when the parameter α belongs to the Cantor set

$$\mathsf{C}_{\infty}^{\kappa,\varepsilon} = \Big\{ \alpha \in (\underline{\alpha},\overline{\alpha}); \ \mathsf{c}_{\infty}(\omega(\alpha),\alpha) = -\omega_{\mathrm{Eq}} \quad \mathrm{and} \quad (\omega,\alpha) \in \mathsf{G}_{\infty}^{\kappa} \Big\}.$$

The ultimate goal is to measure this last Cantor and show that it is asymptotically with full Lebesgue measure when $\varepsilon \to 0$. This important point will be discussed in Section 10.2 and based on the rigidity of the equilibrium frequencies combined with perturbative arguments.

Г		
L		

10.1. Nash-Moser scheme. The construction of the solutions to the nonlinear equation (10.1) stems from the modified Nash-Moser scheme as in [9, 11] and reproduced later in several papers as for instance in [16, 4]. The scheme is based on the construction of successive approximations belonging to a finite-dimensional space given by

$$E_m \triangleq \left\{ \Im; \ \Pi_m \Im = \Im \right\} \quad \text{with} \quad \Im: \varphi \in \mathbb{T}^d \mapsto \Im(\varphi) = \left(\Theta(\varphi), I(\varphi), z(\varphi) \right)$$

where Π_m is the projector defined

$$h(\varphi,\theta) = \sum_{(l,j)\in\mathbb{Z}^{d+1}} h_{l,j} e^{\mathrm{i}(l\cdot\varphi+j\theta)}, \quad \Pi_m h(\varphi,\theta) = \sum_{|l|+|j|\leqslant N_m} h_{l,j} e^{\mathrm{i}(l\cdot\varphi+j\theta)},$$

and when h depends only on φ it is given by

$$h(\varphi) = \sum_{l \in \mathbb{Z}^d} h_l e^{i l \cdot \varphi}, \quad \Pi_m h(\varphi) = \sum_{|l| \leqslant N_m} h_l e^{i l \cdot \varphi},$$

where the sequence of numbers $(N_m)_m$ is defined in (6.32). We point out that in the preceding sections we have used the same notation Π_m to denote the orthogonal projector localizing only on the time frequency set $\{(l, j) \in \mathbb{Z}^{d+1}, |l| \leq N_m\}$. This latter projector will not be used throughout this section and therefore there is no confusion to fear from this notation in this section.

During the Nash-Moser scheme we need a list of significant parameters $\{a_1, a_2, \overline{a}, \mu_1, \mu_2, s_h\}$ that will be fixed with respect to the geometry of Cantor sets through the parameters τ_1, τ_2 and d. Below, we opt for a particular choice, which is not optimal but fulfills all the required constraints in the Nash-Moser scheme,

$$\begin{aligned}
\overline{a} &= \tau_2 + 2 \\
\mu_1 &= 6\overline{\sigma} + 6 + 3q(\tau_2 + 2) \\
a_1 &= 6q(2 + \tau_2) + 12\overline{\sigma} + 15 \\
a_2 &= 3q(2 + \tau_2) + 6\overline{\sigma} + 9 \\
s_h &= s_0 + 9\overline{\sigma} + \tau_2(5q + 2) + 9q + 12 \\
b_1 &= 2s_h - s_0.
\end{aligned}$$
(10.2)

We will also need the parameter μ_2 introduced in Proposition 9.6 and given by

$$\mu_2 = \frac{2}{3} \left(s_h - \left(s_0 + \tau_1(q+1) + \tau_2(q+1) + 3 \right) \right).$$
(10.3)

Let us now give an insight about the roles played by these numbers in the Nash-Moser scheme, see below Proposition 10.1.

• The number $\overline{\sigma} = \sigma(\tau_1, \tau_2, d)$ is arbitrary large and measures the total loss of regularity in the construction of the approximate inverse according to Theorem 6.1.

♦ The list $\{a_1, a_2\}$ is introduced to basically describe the convergence rates at different lower regularity indexes $s_0, s_0 + \overline{\sigma}$.

• The number μ_1 measures the norm inflation at a higher regularity level related to s_h .

 \Rightarrow The number μ_2 is related to the rate of convergence of the errors in the approximate inverse, see Theorem 6.1.

◆ The number \overline{a} is associated to the enlargement of the intermediate Cantor sets by open sets whose thickness is proportional to $N_m^{-\overline{a}}$. This procedure is needed to extend in a classical way the approximate solutions to the whole set of parameters \mathcal{O} .

We shall also impose the following conditions

$$N_0 \triangleq \kappa^{-1}, \quad \kappa \triangleq \varepsilon^a \quad \text{with} \quad 0 < a < \frac{1}{\mu_2 + q + 3}$$
 (10.4)

The main goal is to prove by induction the following result.

Proposition 10.1 (Nash-Moser). Let $(\tau_1, \tau_2, q, d, s_0, s_h)$ satisfy (9.2), (10.2) and assume (10.4). There exist $C_* > 0$ and $\varepsilon_0 > 0$ such that for any $\varepsilon \in [0, \varepsilon_0]$ and under the second condition in (9.278) we get for all $m \in \mathbb{N}$ the following properties, • $(\mathcal{P}1)_m$ There exists a q-times differentiable function

$$\begin{array}{rccc} W_m: & \mathcal{O} & \to & E_{m-1} \times \mathbb{R}^d \\ & \lambda & \mapsto & (\mathfrak{I}_m, \mathsf{c}_m - \omega) \end{array} \end{array}$$

satisfying

$$W_0 = 0$$
 and $||W_m||_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant C_* \varepsilon \kappa^{-1} N_0^{q\overline{a}}$ for $m \ge 1$.

By setting

 $U_0 = ((\varphi, 0, 0), \omega), \quad U_m = U_0 + W_m \quad and \quad H_m = U_m - U_{m-1} \quad for \quad m \ge 1,$ then we have $\forall s \in [s_0, S],$

$$\|H_1\|_s^{q,\kappa} \leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}} \quad and \quad \|H_k\|_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant C_*\varepsilon\kappa^{-1}N_{k-1}^{-a_2} \quad \forall \ 2\leqslant k\leqslant m.$$

• $(\mathscr{P}2)_m$ Define, $i_m = (\varphi, 0, 0) + \mathfrak{I}_m$, $\kappa_m = \kappa(1 + 2^{-m}) \in [\kappa, 2\kappa]$ and

$$\mathscr{A}_0^{\kappa} = \mathscr{O} \quad and \quad \mathscr{A}_{m+1}^{\kappa} = \mathscr{A}_m^{\kappa} \cap \mathsf{G}_m(\kappa_{m+1}, \tau_1, \tau_2, i_m) \quad \forall m \in \mathbb{N},$$

where $G(\kappa_{m+1}, \tau_1, \tau_2, i_m)$ is defined in Theorem 9.1 and DC_{N_0} is described by (6.31). Consider the open sets

$$\forall r > 0, \quad \mathcal{O}_m^r \triangleq \left\{ \lambda \in \mathcal{O} \, ; \, \operatorname{dist} \left(\lambda, \mathscr{A}_m^{\kappa} \right) < r \, N_m^{-\overline{a}} \right\}$$

where $dist(x, A) = \inf_{y \in A} ||x - y||$. Then we have the following estimate

$$\|\mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} \leqslant C_* \varepsilon N_{m-1}^{-a_1}.$$

• $(\mathcal{P}3)_m \|W_m\|_{b_1+\overline{\sigma}}^{q,\kappa,\theta} \leqslant C_* \varepsilon \kappa^{-1} N_{m-1}^{\mu_1}.$

Remark 10.1. For any open set $O \subset \mathcal{O}$, the norm $\|\cdot\|_s^{q,\kappa,O}$ is defined according to the Definition 4.1 by simply changing the set \mathcal{O} with O.

Proof. The proof will be done using an induction principle.

• Initialization. By construction, $U_0 = ((\varphi, 0, 0), \omega)$ and thus using (5.21) we obtain

$$\mathcal{F}(U_0) = \varepsilon \begin{pmatrix} -\partial_I \mathcal{P}_{\varepsilon}((\varphi, 0, 0)) \\ \partial_{\vartheta} \mathcal{P}_{\varepsilon}((\varphi, 0, 0)) \\ -\partial_{\theta} \nabla_z \mathcal{P}_{\varepsilon}((\varphi, 0, 0)). \end{pmatrix}$$

Thus, using Lemma 5.2-(i), we find

$$\forall s \ge 0, \quad \|\mathscr{F}(U_0)\|_s^{q,\kappa,0} \leqslant C_*\varepsilon \tag{10.5}$$

for some positive constant C_* . The properties $(\mathcal{P}1)_0$, $(\mathcal{P}2)_0$ and $(\mathcal{P}3)_0$ then follow immediately.

9 Induction step. Given $m \in \mathbb{N}^*$ and assume that $(\mathcal{P}1)_k$, $(\mathcal{P}2)_k$ and $(\mathcal{P}3)_k$ are true for any $0 \leq k \leq m$. The goal is to check the validity of theses properties at the order m + 1. To do so, we start first with the construction of the approximation U_{m+1} by using a modified Nash-Moser scheme. We shall introduce the linearized operator of \mathcal{F} at the state (i_m, \mathbf{c}_m)

$$egin{aligned} &L_m &\triangleq L_m(\lambda) \ &= d_{i,\mathsf{c}} \mathscr{F}(i_m(\lambda,\omega),\mathsf{c}_m(\lambda)) \end{aligned}$$

As we shall see later, the constriction of the next approximation U_{m+1} requires an approximate right inverse of L_m . This task is discussed in the preceding sections and a precise statement is stated in Theorem 6.1. To apply this result and get some bounds on U_{m+1} we need to establish first some intermediate results connected with the smallness condition and some Cantor set inclusions.

(a) Smallness/boundedness conditions. We remark that with the conditions (10.2) and (10.3) on the parameters s_h and μ_2 the conditions listed in (9.277) are automatically satisfied and therefore

the results of Proposition 9.6 hold true provided that the boundedness/smallness conditions (9.278) are verified. For the smallness condition, it is satisfied provided that

$$N_0^{\mu_2} \varepsilon \kappa^{-3-q} = \varepsilon^{1-a(\mu_2+3+q)} \leq \varepsilon_0,$$

with ε_0 being a small positive number. This holds true for small ε by virtue of (10.4). Concerning the boundedness condition in (9.278), we recall that $\sigma \ge \sigma_4$ and use Lemma 4.3 leading to

$$\|H_m\|_{s_h+\overline{\sigma}}^{q,\kappa} \lesssim \left(\|H_m\|_{s_0+\overline{\sigma}}^{q,\kappa}\right)^{\frac{1}{2}} \left(\|H_m\|_{b_1+\overline{\sigma}}^{q,\kappa}\right)^{\frac{1}{2}}, \quad b_1 = 2s_h - s_0.$$
(10.6)

On the other hand, by using $(\mathcal{P}1)_m$ we find

$$\forall s \in [s_0, S], \|H_1\|_s^{q,\kappa} \leqslant \frac{1}{2} C_* \varepsilon \kappa^{-1} N_0^{q\overline{a}} \quad \text{and} \quad \|H_m\|_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant C_* \varepsilon \kappa^{-1} N_{m-1}^{-a_2}.$$
(10.7)

Now applying $(\mathcal{P}3)_m$ and $(\mathcal{P}3)_{m-1}$ yields

$$\begin{aligned} \|H_m\|_{b_1+\overline{\sigma}}^{q,\kappa} &\leqslant \|W_m\|_{b_1+\overline{\sigma}}^{q,\kappa} + \|W_{m-1}\|_{b_1+\overline{\sigma}}^{q,\kappa} \\ &\leqslant 2C_*\varepsilon\kappa^{-1}N_{m-1}^{\mu}. \end{aligned}$$

Inserting the foregoing estimates into (10.6) allows to get

$$\|H_m\|_{s_h+\overline{\sigma}}^{q,\kappa} \leqslant CC_*\varepsilon\kappa^{-1}N_{m-1}^{\frac{1}{2}\mu-\frac{1}{2}a_2}$$

However for m = 1 one has

$$\|H_1\|_{s_h+\overline{\sigma}}^{q,\kappa} \leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}}$$

From (10.2), one gets

 $\mu - a_2 \leqslant -2. \tag{10.8}$

Thus, using (10.4) and taking ε small enough, we obtain

$$|W_m||_{s_h+\overline{\sigma}}^{q,\kappa} \leqslant ||H_1||_{s_h+\overline{\sigma}}^{q,\kappa} + \sum_{k=2}^m ||H_k||_{s_h+\overline{\sigma}}^{q,\kappa}$$
$$\leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}} + CC_*\varepsilon\kappa^{-1}\sum_{k=0}^\infty N_k^{-1}$$
$$\leqslant C_*\varepsilon^{1-a(1+q\overline{a})}.$$

Hence by virtue of (10.4) and (10.2) we find

$$a \leqslant \frac{1}{2(1+q\overline{a})} \tag{10.9}$$

which implies in turn that

$$\|W_m\|_{s_h+\sigma}^{q,\kappa} \leqslant C\varepsilon^{\frac{1}{2}} \tag{10.10}$$
$$\leqslant 1,$$

provided that ε is small enough. Another important observation is that $\overline{\sigma}$ can be chosen large enough in order to get $s_0 + \overline{\sigma} \ge \overline{s}_h + \sigma_4$ where \overline{s}_h is defined in (9.38). By this way we get from the second estimate of (10.7) and Sobolev embeddings

$$\|H_m\|_{\overline{s}_h + \sigma_4}^{q,\kappa} \leqslant C_* \varepsilon \kappa^{-1} N_{m-1}^{-a_2}.$$
(10.11)

(b) Set inclusions. Notice that by the previous point and the conditions (10.2) and (10.4) we can perform the reducibility of the linearized operator in the normal directions at the step m. In particular, Propositions 9.2 and 9.6 apply. Thus the sets \mathscr{A}_k^{κ} are well-defined for all $k \leq m+1$ and to develop later suitable estimates we need to establish first the following inclusions

$$\mathscr{A}_{m+1}^{\kappa} \subset \mathcal{O}_{m+1}^{2\kappa} \subset \left(\mathscr{A}_{m+1}^{\frac{\kappa}{2}} \cap \mathcal{O}_{m}^{\kappa} \right).$$
(10.12)

We draw the reader's attention not to confuse κ and k in this part. The first left inclusion in (10.12) is obvious by construction of $O_{m+1}^{2\kappa}$ which is an enlargement of $\mathscr{A}_{m+1}^{\kappa}$. As to the second inclusion, we first claim that

$$\forall k \in \llbracket 0, m \rrbracket, \quad \mathcal{O}_{k+1}^{2\kappa} \subset \mathcal{O}_k^{\kappa}. \tag{10.13}$$

Indeed, since by construction $\mathscr{A}_{k+1}^{\kappa} \subset \mathscr{A}_{k}^{\kappa}$ then taking $\lambda \in O_{k+1}^{2\kappa}$ we have the following estimates

$$\begin{split} \mathtt{dist}\big(\lambda,\mathscr{A}_{k}^{\kappa}\big) &\leqslant \mathtt{dist}\big(\lambda,\mathscr{A}_{k+1}^{\kappa}\big) \\ &< 2\kappa N_{k+1}^{-\overline{a}} = 2\kappa N_{k}^{-\overline{a}} N_{0}^{-\frac{1}{2}\overline{a}} \\ &< \kappa N_{k}^{-\overline{a}} \end{split}$$

provided that $2N_0^{-\frac{1}{2}\overline{a}} < 1$, which is satisfied for N_0 large enough, that is in view of (10.4) for ε small enough. The next task is to prove by induction in k that

$$\forall k \in [\![0, m+1]\!], \quad \mathcal{O}_k^{2\kappa} \subset \mathscr{A}_k^{\frac{\kappa}{2}}. \tag{10.14}$$

Remark that for k = 0, one has $O_0^{2\kappa} = \mathcal{O} = \mathcal{A}_0^{\frac{\kappa}{2}}$. Now assume that (10.14) occurs at the order $k \in [0, n]$ and let us check it at the order k+1. Using (10.13) and the induction assumption (10.14), we get

$$O_{k+1}^{2\kappa} \subset O_k^{\kappa} \subset O_k^{2\kappa} \subset \mathscr{A}_k^{\frac{\kappa}{2}}.$$

Hence, it remains to check that

$$\mathcal{O}_{k+1}^{2\kappa} \subset \mathsf{G}_k\left(\frac{\kappa_{k+1}}{2}, \tau_1, \tau_2, i_k\right)$$

Given $\lambda \in O_{k+1}^{2\kappa}$ there exists $\lambda' = (\omega', \alpha') \in \mathscr{A}_{k+1}^{\kappa}$ such that $\operatorname{dist}(\lambda, \lambda') < 2\kappa N_{k+1}^{-\overline{a}}$. Then for all $(l, j) \in \mathbb{Z}^d \times \mathbb{S}_0^c$ with $|l| \leq N_k$, we get using the triangle inequality and Cauchy-Schwarz inequalities

$$\begin{split} \omega \cdot l + \mu_j^{\infty}(\lambda, i_k) \Big| \geqslant \Big| \omega' \cdot l + \mu_j^{\infty}(\lambda', i_k) \Big| - |\omega - \omega'| |l| - \Big| \mu_j^{\infty}(\lambda, i_k) - \mu_j^{\infty}(\lambda', i_k) \Big| \\ > \frac{\kappa_{k+1}\langle j \rangle}{\langle l \rangle^{\tau_1}} - 2\kappa N_{k+1}^{1-\overline{a}} - \Big| \mu_j^{\infty}(\lambda, i_k) - \mu_j^{\infty}(\lambda', i_k) \Big|. \end{split}$$

Using the Mean Value Theorem yields

$$\begin{aligned} \left|\mu_{j}^{\infty}(\lambda, i_{k}) - \mu_{j}^{\infty}(\lambda', i_{k})\right| &\leq |\lambda - \lambda'| \kappa^{-1} \|\mu_{j}^{\infty}(i_{k})\|^{q,\kappa} \\ &\leq 2N_{k+1}^{-\overline{a}} \|\mu_{j}^{\infty}(i_{k})\|^{q,\kappa}. \end{aligned}$$
(10.15)

Therefore combining (10.15) with the assumption $(\mathcal{P}1)_m$ and the eigenvalues estimates given in Propositions 9.6-9.5 we get in view of (9.278)

$$\left|\mu_{j}^{\infty}(\lambda, i_{k}) - \mu_{j}^{\infty}(\lambda', i_{k})\right| \leqslant C\kappa \langle j \rangle N_{k+1}^{-\overline{a}}.$$
(10.16)

Hence, by (9.278) we get since $|l| \leq N_k$ and $\kappa_{k+1} \geq \kappa$,

$$\begin{aligned} \left| \omega \cdot l + \mu_j^{\infty}(\lambda, i_k) \right| &> \frac{\kappa_{k+1}\langle j \rangle}{\langle l \rangle^{\tau_1}} - C\kappa \langle j \rangle N_{k+1}^{1-\overline{a}} \\ &> \frac{\kappa_{k+1}\langle j \rangle}{\langle l \rangle^{\tau_1}} \left(1 - CN_{k+1}^{\tau_1+1-\overline{a}} \right). \end{aligned}$$

From (10.2) and (9.2) we find $\tau_1 + 2 \leq \overline{a}$ and taking N_0 sufficiently large we get

$$CN_{k+1}^{\tau_1+1-\overline{a}} \leqslant CN_0^{-1} \leqslant \frac{1}{2}$$

so that

$$\left|\omega \cdot l + \mu_j^{\infty}(\lambda, i_k)\right| > \frac{\kappa_{k+1}\langle j \rangle}{2\langle l \rangle^{\tau_1}} \cdot$$

As a consequence, we deduce that $\lambda \in \Lambda_{\infty,k}^{\frac{\kappa_{k+1}}{2},\tau_1}(i_k)$. Let us now check that $\lambda \in \mathcal{O}_{\infty,k}^{\frac{\kappa_{k+1}}{2},\tau_1}(i_k)$ defined in Proposition 9.2 (using the new notation $c_i(\lambda) = c(\lambda, i)$),

$$\begin{split} \omega \cdot l + jc_{i_k}(\lambda) \Big| \geqslant \Big| \omega' \cdot l + jc_{i_k}(\lambda') \Big| - |\omega - \omega'||l| - |j| \Big| c_{i_k}(\lambda) - c_{i_k}(\lambda') \Big| \\ > 4 \kappa_{k+1}^{\varrho} \frac{\langle j \rangle}{\langle l \rangle^{\tau_1}} - 2\kappa N_{k+1}^{1-\overline{\alpha}} - \langle j \rangle \Big| c_{i_k}(\lambda) - c_{i_k}(\lambda') \Big|. \end{split}$$

Using the Mean Value Theorem and (9.40) combined with (9.278) and the assumption $(\mathcal{P}1)_m$

$$\begin{aligned} \left|c_{i_{k}}(\lambda) - c_{i_{k}}(\lambda')\right| &\leq |\lambda - \lambda'| \left(C + \kappa^{-1} \|c_{r,\lambda} - V_{0,\alpha}\|^{q,\kappa}\right) + 2\kappa N_{k+1}^{-\overline{a}} \left(C + C\varepsilon\kappa^{-2}\right) \\ &\leq C\kappa N_{k+1}^{-\overline{a}}.\end{aligned}$$

Hence as before we get from the definition of κ_m and $\varrho \in (0, 1)$

$$\begin{split} \omega \cdot l + c_{i_k}(\lambda) \Big| &> 4 \kappa_{k+1}^{\varrho} \frac{\langle j \rangle}{\langle l \rangle^{\tau_1}} - C \kappa \langle j \rangle N_{k+1}^{1-\overline{a}} \\ &\geqslant 4 \frac{\left(\frac{\kappa_{k+1}}{2}\right)^{\varrho} \langle j \rangle}{\langle l \rangle^{\tau_1}} \left(2^{\varrho} - C 2^{\varrho} N_{k+1}^{\tau_1+1-\overline{a}} \right) \end{split}$$

Since by assumption $\overline{a} \ge \tau_1 + 2$ then taking N_0 sufficiently large we get

$$CN_{k+1}^{\tau_1+1-\overline{a}} \leqslant CN_0^{-1} < 1 - 2^{-\varrho}$$

implying that

$$|\omega \cdot l + jc_{i_k}(\lambda)| > 4\left(\frac{\kappa_{k+1}}{2}\right)^{\varrho} \frac{\langle j \rangle}{\langle l \rangle^{\tau_1}}$$

This shows that $\lambda \in \mathcal{O}_{\infty,k}^{\frac{\kappa_{k+1}}{2},\tau_1}(i_k)$. It remains to check that $\lambda \in \mathcal{O}_{\infty,k}^{\frac{\kappa_{k+1}}{2},\tau_1,\tau_2}(i_k)$ defined in Proposition 9.6. We write by the triangle inequality and the condition $\lambda' \in \mathcal{A}_{m+1}^{\kappa}$

$$\begin{split} \left| \omega \cdot l + \mu_j^{\infty}(\lambda, i_k) - \mu_{j_0}^{\infty}(\lambda, i_k) \right| &\ge \left| \omega' \cdot l + \mu_j^{\infty}(\lambda', i_k) - \mu_{j_0}^{\infty}(\lambda', i_k) \right| - \left| \omega - \omega' \right| \left| l \right| \\ &- \left| \mu_j^{\infty}(\lambda, i_k) - \mu_{j_0}^{\infty}(\lambda, i_k) + \mu_{j_0}^{\infty}(\lambda', i_k) - \mu_j^{\infty}(\lambda', i_k) \right| \\ &> \frac{2\kappa_{k+1}\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} - 2\kappa N_{k+1}^{1-\overline{a}} \\ &- \left| \mu_j^{\infty}(\lambda, i_k) - \mu_{j_0}^{\infty}(\lambda, i_k) + \mu_{j_0}^{\infty}(\lambda', i_k) - \mu_j^{\infty}(\lambda', i_k) \right|. \end{split}$$

Now recall that from Proposition 9.6, we can write

$$\mu_j^{\infty}(\lambda, i_k) = \mu_j^0(\lambda, i_k) + r_j^{\infty}(\lambda, i_k)$$

and then

$$\begin{split} \left| \mu_{j}^{\infty}(\lambda, i_{k}) - \mu_{j_{0}}^{\infty}(\lambda, i_{k}) + \mu_{j_{0}}^{\infty}(\lambda', i_{k}) - \mu_{j}^{\infty}(\lambda', i_{k}) \right| \\ & \leq \left| \mu_{j}^{0}(\lambda, i_{k}) - \mu_{j_{0}}^{0}(\lambda, i_{k}) + \mu_{j_{0}}^{0}(\lambda', i_{k}) - \mu_{j}^{0}(\lambda', i_{k}) \right| \\ & + \left| r_{j}^{\infty}(\lambda, i_{k}) - r_{j}^{\infty}(\lambda', i_{k}) \right| + \left| r_{j_{0}}^{\infty}(\lambda, i_{k}) - r_{j_{0}}^{\infty}(\lambda', i_{k}) \right|. \end{split}$$

Using the Mean Value Theorem, combined with Proposition 9.5, Lemma 3.1-(vi) and (9.278) yield

$$\left| \mu_{j}^{0}(\lambda, i_{k}) - \mu_{j_{0}}^{0}(\lambda, i_{k}) + \mu_{j_{0}}^{0}(\lambda', i_{k}) - \mu_{j}^{0}(\lambda', i_{k}) \right| \leq C \kappa N_{k+1}^{-\overline{a}} \langle j - j_{0} \rangle.$$

Similarly we get in view of (9.280)

$$\begin{aligned} \left| r_j^{\infty}(\lambda, i_k) - r_j^{\infty}(\lambda', i_k) \right| &\leq C \kappa N_{k+1}^{-\overline{a}} \varepsilon \kappa^{-3} \\ &\leq C \kappa N_{k+1}^{-\overline{a}} \langle j - j_0 \rangle. \end{aligned}$$

Putting together the foregoing estimates and the facts that $|l| \leq N_k$ and $\kappa_{k+1} \geq \kappa$ yield

$$\left|\omega \cdot l + \mu_j^{\infty}(\lambda, i_k) - \mu_{j_0}^{\infty}(\lambda, i_k)\right| \ge \kappa_{k+1} \frac{\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \left(2 - CN_{k+1}^{\tau_2 + 1 - \overline{a}}\right)$$

Since $\overline{a} = \tau_2 + 2$, see (10.2), then taking N_0 sufficiently large we find

$$CN_{k+1}^{\tau_2+1-\overline{a}} \leqslant CN_0^{-1} < 1$$

we deduce that

$$\left|\omega \cdot l + \mu_j^{\infty}(\lambda, i_k) - \mu_{j_0}^{\infty}(\lambda, i_k)\right| > \kappa_{k+1} \frac{\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \cdot$$

It follows that $\lambda \in \mathcal{O}_{\infty,k}^{\frac{\kappa_{k+1}}{2},\tau_1,\tau_2}(i_k)$. Consequently, we deduce that $\lambda \in \mathsf{G}_k(\frac{1}{2}\kappa_{k+1},\tau_1,\tau_2,i_k)$ and thus we conclude that $\lambda \in \mathscr{A}_{k+1}^{\frac{\kappa}{2}}$. This ends the proof of (10.14) and therefore the proof of (10.12) is now complete.

(c) Construction of the next approximation. Since the assumption (9.278) us satisfied at the order m we can perform the reduction of the linearized operator stated in Theorem 9.1. Then coming back to Theorem 6.1 applied with L_m we find an operator $\mathbf{T}_m \triangleq \mathbf{T}_m(\lambda)$ well-defined in the whole set of parameters $\lambda \in \mathcal{O}$ with the estimates

$$\forall s \in [s_0, S], \quad \|\mathbf{T}_m h\|_s^{q,\kappa} \lesssim \kappa^{-1} \left(\|h\|_{s+\overline{\sigma}}^{q,\kappa} + \|\mathfrak{I}_m\|_{s+\overline{\sigma}}^{q,\kappa} \|h\|_{s_0+\overline{\sigma}}^{q,\kappa} \right)$$
(10.17)

and

$$\|\mathbf{T}_{m}h\|_{s_{0}}^{q,\kappa} \lesssim \kappa^{-1} \|h\|_{s_{0}+\overline{\sigma}}^{q,\kappa}.$$
(10.18)

In addition, when the parameters are restricted in the Cantor set $G_n(\kappa_{m+1}, \tau_1, \tau_2, i_m)$, \mathbf{T}_m is an approximate right inverse of L_m in the sense of (6.61) which will be useful later. Next, we define the function \tilde{U}_{m+1} as follows

 $\widetilde{U}_{m+1} \triangleq U_m + \widetilde{H}_{m+1} \quad \text{with} \quad \widetilde{H}_{m+1} \triangleq (\widehat{\mathfrak{I}}_{m+1}, \widehat{\mathfrak{c}}_{m+1}) \triangleq -\mathbf{\Pi}_m \mathbf{T}_m \mathbf{\Pi}_m \mathscr{F}(U_m) \in E_m \times \mathbb{R}^d$

where Π_m is defined by

$$\mathbf{\Pi}_m(\mathfrak{I},\alpha) = (\Pi_m\mathfrak{I},\alpha) \quad \text{and} \quad \mathbf{\Pi}_m^{\perp}(\mathfrak{I},\alpha) = (\Pi_m^{\perp}\mathfrak{I},0).$$
(10.19)

Observe that U_m is defined in the full set \mathcal{O} which implies that \widetilde{U}_{m+1} is defined in \mathcal{O} too. However we shall not work with this natural extension but we will localize it around the Cantor set $\mathscr{A}_{m+1}^{\kappa}$ in order to get a good decay. Let us introduce the quadratic function

$$Q_m = \mathscr{F}(\widetilde{U}_{m+1}) - \mathscr{F}(U_m) - L_m \widetilde{H}_{m+1}.$$
(10.20)

Then it is easy to check that

$$\mathcal{F}(\tilde{U}_{m+1}) = \mathcal{F}(U_m) - L_m \mathbf{\Pi}_m \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + Q_m$$

$$= \mathcal{F}(U_m) - L_m \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + L_m \mathbf{\Pi}_m^{\perp} \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + Q_m$$

$$= \mathcal{F}(U_m) - \Pi_m L_m \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + (L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m) \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + Q_m$$

$$= \Pi_m^{\perp} \mathcal{F}(U_m) - \Pi_m (L_m \mathbf{T}_m - \mathrm{Id}) \Pi_m \mathcal{F}(U_m) + (L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m) \mathbf{T}_m \Pi_m \mathcal{F}(U_m) + Q_m.$$
(10.21)

► Estimates of $\mathscr{F}(\widetilde{U}_{m+1}), m \ge 1$. We shall estimate $\mathscr{F}(U_{m+1})$ in norm $\|\cdot\|_s^{q,\kappa,O_{m+1}^{2\kappa}}$ through estimating the right-hand side terms in (10.21). We note that the parameters $\lambda = (\omega, \alpha)$ are located in the small open set $O_{m+1}^{2\kappa}$ containing the Cantor set $\mathscr{A}_{m+1}^{\kappa}$ as indicated in (10.12).

• Estimate of $\Pi_m^{\perp} \mathscr{F}(U_m)$. By Taylor formula applied with (5.21) and Lemma 5.2 combined with (10.5) and $(\mathscr{P}1)_m$, we get

$$\forall s \geq s_0, \quad \|\mathscr{F}(U_m)\|_s^{q,\kappa,\mathcal{O}_m^{\kappa}} \leq \|\mathscr{F}(U_0)\|_s^{q,\kappa,\mathcal{O}_0^{\kappa}} + \|\mathscr{F}(U_m) - \mathscr{F}(U_0)\|_s^{q,\kappa,\mathcal{O}_m^{\kappa}} \\ \lesssim \varepsilon + \|W_m\|_{s+\overline{\sigma}}^{q,\kappa,0}.$$
(10.22)

Therefore we deduce from the smallness condition (10.4) and $(\mathcal{P}1)_m$,

$$\kappa^{-1} \| \mathscr{F}(U_m) \|_{s_0}^{q,\kappa,\mathcal{O}_m^\kappa} \leqslant 1.$$

$$(10.23)$$

By Lemma 4.2 combined with (10.22), we infer since $b_1 = 2s_h - s_0$

$$\begin{aligned} \|\Pi_m^{\perp} \mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} &\leq N_m^{s_0-b_1} \|\mathscr{F}(U_m)\|_{b_1}^{q,\kappa,\mathcal{O}_m^{\kappa}} \\ &\lesssim N_m^{s_0-b_1} \left(\varepsilon + \|W_m\|_{b_1+\overline{\sigma}}^{q,\kappa,\mathcal{O}}\right). \end{aligned}$$

Now, by using $(\mathcal{P}3)_m$ combined with (6.32) and (10.4) we obtain

$$\varepsilon + \|W_m\|_{b_1 + \overline{\sigma}}^{q, \kappa, \theta} \leq \varepsilon \left(1 + C_* \kappa^{-1} N_{m-1}^{\mu_1}\right)$$
$$\leq 2C_* \varepsilon N_m^{1 + \frac{2}{3}\mu_1}. \tag{10.24}$$

Inserting this in the previous estimate and making appeal to (10.12) we finally get

$$\|\Pi_m^{\perp} \mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} \leqslant \|\Pi_m^{\perp} \mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}}$$

$$\lesssim \varepsilon N_m^{s_0+1+\frac{2}{3}\mu_1-b_1}.$$
 (10.25)

Notice that one also gets from (10.22) and (10.24)

$$\|\mathscr{F}(U_m)\|_{b_1+\overline{\sigma}}^{q,\kappa,\mathcal{O}_m^{\kappa}} \leqslant C_*\varepsilon N_m^{\overline{\sigma}+\frac{2}{3}\mu_1+1}.$$
(10.26)

• Estimate of $\Pi_m(L_m\mathbf{T}_m - \mathrm{Id})\Pi_m\mathscr{F}(U_m)$. According to (10.12) we may write

$$\mathcal{O}_{m+1}^{2\kappa} \subset \mathscr{A}_{m+1}^{\frac{\kappa}{2}} \subset \mathfrak{G}_m(\frac{1}{2}\kappa_{m+1},\tau_1,\tau_2,i_m) \cap \mathcal{O}_m^{\kappa}.$$

Then we can apply Theorem 6.1

$$\|\Pi_m (L_m \mathbf{T}_m - \mathrm{Id})\Pi_m \mathcal{F}(U_m)\|_s^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} \leq \|\mathcal{E}_1^{(m)}\Pi_m \mathcal{F}(U_m)\|_s^{q,\kappa,\mathcal{O}_m^{\kappa}} + \|\mathcal{E}_2^{(m)}\Pi_m \mathcal{F}(U_m)\|_s^{q,\kappa,\mathcal{O}_m^{\kappa}} + \|\mathcal{E}_3^{(m)}\Pi_m \mathcal{F}(U_m)\|_s^{q,\kappa,\mathcal{O}_m^{\kappa}} \triangleq \mathbb{I}_1^{(m)}(s) + \mathbb{I}_2^{(m)}(s) + \mathbb{I}_3^{(m)}(s).$$
(10.27)

Let us start with the estimate of the three terms on the right hand side for $m \ge 1$. Applying the estimates of Theorem 6.1 and using $(\mathcal{P}2)_m$

$$\mathbf{I}_{1}^{(m)}(s_{0}) \lesssim \kappa^{-1} \|\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \|\Pi_{m}\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \\
\lesssim \kappa^{-1} N_{m}^{\overline{\sigma}} \|\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \|\mathscr{F}(U_{m})\|_{s_{0}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \\
\lesssim \varepsilon \kappa^{-1} N_{m}^{\overline{\sigma}-\frac{2}{3}a_{1}} \|\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}}.$$
(10.28)

On the other hand, one may write in view of Lemma 4.2, $(\mathcal{P}2)_m$, $(\mathcal{P}3)_m$ and (10.26)

$$\begin{aligned} \|\mathscr{F}(U_m)\|_{s_0+\sigma}^{q,\kappa,\mathcal{O}_m^{\kappa}} &\lesssim N_m^{\overline{\sigma}} \|\mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} + \|\Pi_m^{\perp}\mathscr{F}(U_m)\|_{s_0+\overline{\sigma}}^{q,\kappa,\mathcal{O}_m^{\kappa}} \\ &\lesssim N_m^{\overline{\sigma}} \|\mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} + N_m^{s_0-b_1} \|\mathscr{F}(U_m)\|_{b_1+\overline{\sigma}}^{q,\kappa,\mathcal{O}_m^{\kappa}} \\ &\lesssim \varepsilon N_m^{\overline{\sigma}-\frac{2}{3}a_1} + \varepsilon N_m^{s_0+\overline{\sigma}+\frac{2}{3}\mu_1+1-b_1}. \end{aligned}$$

Putting together the preceding two estimates and using $\varepsilon \kappa^{-1} \leq 1$, which follows from (10.4), give

$$I_1^{(m)}(s_0) \lesssim \varepsilon \left(N_m^{2\sigma - \frac{4}{3}a_1} + N_m^{s_0 + 2\overline{\sigma} + \frac{2}{3}\mu_1 + 1 - \frac{2}{3}a_1 - b_1} \right).$$
(10.29)

Coming back once again to Theorem 6.1

$$\mathbf{I}_{2}^{(m)}(s_{0}) \lesssim \kappa^{-1} N_{m}^{s_{0}-b_{1}} \Big(\|\mathscr{F}(U_{m})\|_{b_{1}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \varepsilon N_{m}^{\overline{\sigma}} \|W_{m}\|_{b_{1}+\overline{\sigma}}^{q,\kappa} \big\|\mathscr{F}(U_{m})\|_{s_{0}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \Big).$$

It follows from $(\mathcal{P}2)_m$, $(\mathcal{P}3)_m$ and (10.26)

$$\begin{aligned} \mathbf{I}_{2}^{(m)}(s_{0}) &\lesssim \varepsilon \kappa^{-1} N_{m}^{s_{0}-b_{1}} \left(N_{m}^{\overline{\sigma}+\frac{2}{3}\mu+1} + N_{m}^{\overline{\sigma}} \kappa^{-1} N_{m-1}^{\mu_{1}} \varepsilon N_{m-1}^{-a_{1}} \right) \\ &\lesssim \varepsilon N_{m}^{\overline{\sigma}+\frac{2}{3}\mu_{1}+2+s_{0}-b_{1}}. \end{aligned}$$

For the third term, we shall use once again Theorem 6.1 allowing to get in view of $(\mathcal{P}1)_m$, $(\mathcal{P}2)_m$, $(\mathcal{P}3)_m$, (10.26) and (9.278)

$$\begin{aligned} \mathbf{I}_{3}^{(m)}(s_{0}) &\lesssim N_{m}^{s_{0}-b_{1}}\kappa^{-2} \Big(\|\Pi_{m}\mathscr{F}(U_{m})\|_{b_{1}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \varepsilon\kappa^{-3} \|W_{m}\|_{b_{1}+\overline{\sigma}}^{q,\kappa} \|\Pi_{m}\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \Big) \\ &+ \varepsilon\kappa^{-5}N_{0}^{\mu_{2}}N_{m}^{-\mu_{2}} \|\Pi_{m}\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa} \\ &\lesssim \varepsilon \Big(N_{m}^{s_{0}-b_{1}+\overline{\sigma}+\frac{2}{3}\mu_{1}+3} + N_{m}^{s_{0}-b_{1}-a_{1}+\overline{\sigma}+\frac{2}{3}\mu_{1}} + N_{m}^{\overline{\sigma}+2-\mu_{2}-\frac{2}{3}a_{1}} \Big). \end{aligned}$$

Therefore

$$\mathbf{I}_3^{(m)}(s_0) \lesssim \varepsilon \Big(N_m^{s_0-b_1+\overline{\sigma}+\frac{2}{3}\mu_1+3} + N_m^{\overline{\sigma}+2-\mu_2-\frac{2}{3}a_1} \Big)$$

Plugging the preceding estimates into (10.27) yields for $m \ge 1$

$$\|\Pi_m (L_m \mathbf{T}_m - \mathrm{Id})\Pi_m \mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} \lesssim \varepsilon \Big(N_m^{2\overline{\sigma} - \frac{4}{3}a_1} + N_m^{s_0 + 2\overline{\sigma} + \frac{2}{3}\mu_1 + 1 - b_1} + N_m^{\overline{\sigma} + 2 - \mu_2 - \frac{2}{3}a_1} \Big).$$
(10.30)

For m = 0 we obtain from slight modifications of (10.28) and (10.5)

$$\mathbf{I}_1^{(0)}(s_0) \lesssim \varepsilon^2 \kappa^{-1}$$

Similarly, we get

$$\mathbf{I}_2^{(0)}(s_0) \lesssim \varepsilon \kappa^{-1}$$

and since $\varepsilon \kappa^{-3} \lesssim 1$ in view of (9.278), then

$$\begin{split} \mathbf{I}_{3}^{(0)}(s_{0}) &\lesssim \varepsilon \kappa^{-2} N_{0}^{s_{0}-b_{1}} + \varepsilon^{2} \kappa^{-5} \\ &\lesssim \varepsilon \kappa^{-2}. \end{split}$$

Hence we find

$$\|\Pi_0(L_0\mathbf{T}_0 - \mathrm{Id})\Pi_0\mathcal{F}(U_0)\|_{s_0}^{q,\kappa,\mathcal{O}_1^{2\kappa}} \lesssim \varepsilon \kappa^{-2}.$$
(10.31)

• Estimate of $(L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m) \mathbf{T}_m \Pi_m \mathscr{F}(U_m)$. Using (5.4) one finds for $H = (\widehat{\mathfrak{I}}, \widehat{\mathfrak{c}})$,

$$L_m H = \omega \cdot \partial_{\varphi} \widehat{\mathfrak{I}} - (0, 0, \partial_{\theta} \mathcal{L}(\alpha) \widehat{w}) - \varepsilon d_i X_{\mathscr{P}_{\varepsilon}}(i_n) \widehat{\mathfrak{I}} - (\widehat{\mathfrak{c}}, 0, 0).$$
(10.32)

As $L(\alpha)$ is a Fourier multiplier then $[\Pi_m^{\perp}, L(\alpha)] = 0$. Therefore we infer from (10.19)

$$\left(L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m\right) H = -\varepsilon \left[d_i X_{P_{\varepsilon}}(i_m, \mathbf{c}_m), \Pi_m^{\perp}\right] \widehat{\mathfrak{I}},$$

According to Lemma 5.2-(ii), Lemma 7.3-(iv) and $(\mathcal{P}1)_m$ we obtain

$$\left\| \left(L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m \right) H \right\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{\kappa}} \lesssim \varepsilon N_m^{s_0-b_1} \left(\|\widehat{\mathfrak{I}}\|_{b_1+1}^{q,\kappa,\mathcal{O}_m^{\kappa}} + \|\mathfrak{I}_m\|_{b_1+\overline{\sigma}}^{q,\kappa,\mathcal{O}_m^{\kappa}} \|\widehat{\mathfrak{I}}\|_{s_0+1}^{q,\kappa,\mathcal{O}_m^{\kappa}} \right).$$

It follows that

$$\mathbf{I}_{4}^{(m)} \triangleq \left\| \left(L_m \mathbf{\Pi}_m^{\perp} - \Pi_m^{\perp} L_m \right) \mathbf{T}_m \Pi_m \mathcal{F}(U_m) \right\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} \lesssim \varepsilon N_m^{s_0-b_1} \| \mathbf{T}_m \Pi_m \mathcal{F}(U_m) \|_{b_1+1}^{\gamma,\mathcal{O}_m^{\kappa}} + \varepsilon N_m^{s_0-b_1} \| \mathfrak{I}_m \|_{b_1+\overline{\sigma}}^{q,\kappa,\mathcal{O}} \| \mathbf{T}_m \Pi_m \mathcal{F}(U_m) \|_{s_0+1}^{q,\kappa,\mathcal{O}_m^{\kappa}}.$$

Therefore putting together (10.17) with Lemma 4.2, Sobolev embeddings, (10.4) and $(\mathcal{P}1)_m$ allows to get

$$\begin{split} \mathbf{I}_{4}^{(m)} &\lesssim \varepsilon \kappa^{-1} N_{m}^{s_{0}-b_{1}} \left[\|\Pi_{m} \mathscr{F}(U_{m})\|_{b_{1}+\overline{\sigma}+1}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \|\mathfrak{I}_{m}\|_{b_{1}+\overline{\sigma}+1}^{q,\kappa,\mathcal{O}} \|\Pi_{m} \mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \right] \\ &+ \varepsilon \kappa^{-1} N_{m}^{s_{0}-b_{1}} \|\mathfrak{I}_{m}\|_{b_{1}+\overline{\sigma}} \left(\|\Pi_{m} \mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}+1}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \|\mathfrak{I}_{m}\|_{s_{0}+\overline{\sigma}+1}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \|\Pi_{m} \mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \right) \\ &\lesssim \varepsilon N_{m}^{s_{0}+2-b_{1}} \left(\|\mathscr{F}(U_{m})\|_{b_{1}+\sigma}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \|W_{m}\|_{q,b_{1}+\overline{\sigma}}^{\gamma,\mathcal{O}} \|\Pi_{m} \mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \right). \end{split}$$

Applying Lemma 4.2 and $(\mathcal{P}2)_m$, we find for $m \in \mathbb{N}$

$$\begin{aligned} \|\Pi_m \mathscr{F}(\widetilde{U}_m)\|_{s_0+\sigma}^{q,\kappa,\mathcal{O}_m^{\kappa}} &\lesssim \varepsilon N_m^{\overline{\sigma}} N_{m-1}^{-a_1} \\ &\lesssim \varepsilon N_m^{\overline{\sigma}-\frac{2}{3}a_1}. \end{aligned}$$

Combining this estimate with (10.26) and $(\mathcal{P}3)_m$, we obtain for $m \in \mathbb{N}$

$$\|(L_m\Pi_m^{\perp} - \Pi_m^{\perp}L_m)\mathbf{T}_m\Pi_m\mathcal{F}(U_m)\|_{s_0}^{q,\kappa,\mathbf{O}_{m+1}^{2\kappa}} \leqslant C_*\varepsilon N_m^{s_0+\overline{\sigma}+\frac{2}{3}\mu_1+3-b_1}.$$
(10.33)

• Estimates if Q_m . Coming back to (10.20) and using Taylor Formula at the second order, one finds

$$Q_m = \int_0^1 (1-t) d_{i,\alpha}^2 \mathcal{F}(U_m + t\widetilde{H}_{m+1})[\widetilde{H}_{m+1}, \widetilde{H}_{m+1}] dt$$

Hence (10.32) and Lemma 5.2-(iii) allow to get

$$\|Q_m\|_{s_0}^{q,\kappa,O_{m+1}^{2\kappa}} \lesssim \varepsilon \left(1 + \|W_m\|_{s_0+2}^{q,\kappa,0} + \|\widetilde{H}_{m+1}\|_{s_0+2}^{q,\kappa,O_m^{\kappa}}\right) \left(\|\widetilde{H}_{m+1}\|_{s_0+2}^{q,\kappa,O_m^{\kappa}}\right)^2.$$
(10.34)

Combining (10.17), (10.22) and (10.23), we have for all $s \in [s_0, S]$,

$$\begin{aligned} \|\widetilde{H}_{m+1}\|_{s}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} &= \|\Pi_{m}\mathbf{T}_{m}\Pi_{m}\mathscr{F}(U_{m})\|_{s}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} \\ &\lesssim \kappa^{-1} \left(\|\Pi_{m}\mathscr{F}(U_{m})\|_{s+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + \|\mathfrak{I}_{m}\|_{s+\overline{\sigma}}^{q,\kappa,\mathfrak{O}} \|\Pi_{m}\mathscr{F}(U_{m})\|_{s_{0}+\overline{\sigma}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \right) \\ &\lesssim \kappa^{-1} \left(N_{m}^{\overline{\sigma}}\|\mathscr{F}(U_{m})\|_{s}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} + N_{m}^{2\overline{\sigma}}\|\mathfrak{I}_{m}\|_{s}^{q,\kappa,\mathfrak{O}}\|\mathscr{F}(U_{m})\|_{s_{0}}^{q,\kappa,\mathcal{O}_{m}^{\kappa}} \right) \\ &\lesssim \kappa^{-1}N_{m}^{2\overline{\sigma}} \left(\varepsilon + \|W_{m}\|_{s}^{q,\kappa,\mathfrak{O}} \right). \end{aligned}$$
(10.35)

Similarly, in view of (10.12), (10.18), $(\mathcal{P}1)_m$ and $(\mathcal{P}2)_m$, we get for $m \in \mathbb{N}$

$$\begin{aligned} \|\widetilde{H}_{m+1}\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} &\lesssim \kappa^{-1} N_m^{\overline{\sigma}} \|\mathscr{F}(U_m)\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} \\ &\lesssim \varepsilon \kappa^{-1} N_m^{\overline{\sigma}} N_{m-1}^{-a_1}. \end{aligned}$$
(10.36)

From (10.10), $(\mathcal{P}1)_m$ and (10.36), we find for $m \in \mathbb{N}$

$$\begin{split} \|W_m\|_{s_0+2}^{q,\kappa,0} + \|\widetilde{H}_{m+1}\|_{s_0+2}^{q,\kappa,O_{m+1}^{2\kappa}} &\leqslant 1 + N_m^2 \|H_{m+1}\|_{s_0}^{q,\kappa,O_{m+1}^{2\kappa}} \\ &\leqslant 1 + C\varepsilon\kappa^{-1}N_m^{\overline{\sigma}+2}N_{m-1}^{-a_1} \\ &\leqslant 1 + C\varepsilon N_{m-1}^{3+\frac{3}{2}\overline{\sigma}-a_1}. \end{split}$$

By virtue of (10.2) one has

$$a_1 \geqslant 3 + \frac{3}{2}\sigma \tag{10.37}$$

and using (10.4) we may write for small ε

$$\|W_m\|_{s_0+2}^{q,\kappa,0} + \|\widetilde{H}_{m+1}\|_{s_0+2}^{q,\kappa,O_{m+1}^{2\kappa}} \leqslant 2$$

Thus, by inserting this estimate and (10.36) into (10.34) we deduce that

$$\begin{aligned} \|Q_m\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} &\lesssim \varepsilon \left(\|H_{m+1}\|_{s_0+2}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} \right)^2 \\ &\lesssim \varepsilon N_m^4 \left(\|H_{m+1}\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}} \right)^2 \\ &\lesssim \varepsilon^3 \kappa^{-2} N_m^{2\overline{\sigma}+4} N_{m-1}^{-2a_1}. \end{aligned}$$

Consequently, we find according to (6.32) and $\varepsilon \kappa^{-1} \lesssim 1$ that for $m \ge 1$

$$\begin{aligned} \|Q_m\|_{s_0}^{q,\kappa,\mathcal{O}_m^{\kappa}} &\lesssim \varepsilon^3 \kappa^{-2} N_m^{2\overline{\sigma}+4-\frac{4}{3}a_1} \\ &\lesssim \varepsilon N_m^{2\overline{\sigma}+4-\frac{4}{3}a_1}. \end{aligned}$$
(10.38)

As to the case m = 0, we come back to (10.35) and make suitable adjustments using in particular (10.5)

$$\|\widetilde{H}_1\|_s^{q,\kappa,\mathcal{O}_0^{\kappa}} \lesssim \kappa^{-1} \|\Pi_0 \mathscr{F}(U_0)\|_{s+\overline{\sigma}}^{q,\kappa,\mathcal{O}_0^{\kappa}} \lesssim \varepsilon \kappa^{-1}.$$
(10.39)

Consequently, the inequality (10.38) becomes for m = 0,

$$\|Q_0\|_{s_0}^{q,\kappa,\mathcal{O}_0^\kappa} \lesssim \varepsilon^3 \kappa^{-2}.$$
(10.40)

• Parameters constraints. Plugging (10.25), (10.30), (10.33) and (10.38), into (10.21) yields for $m \ge 1$

$$\|\mathscr{F}(\widetilde{U}_{m+1})\|_{q,s_0}^{\gamma,\mathcal{O}_{m+1}^{2\kappa}} \leqslant CC_* \varepsilon \Big(N_m^{s_0+2\overline{\sigma}+3+\frac{2}{3}\mu_1-b_1} + N_m^{\overline{\sigma}+2-\frac{2}{3}a_1-\mu_2} + N_m^{2\overline{\sigma}+4-\frac{4}{3}a_1} \Big).$$

We want to check that with the assumptions fixed in (10.2) one gets the conditions,

$$\begin{array}{rcl} CN_m^{s_0+2\overline{\sigma}+\frac{2}{3}\mu_1+3-b_1} &\leqslant & \frac{1}{3}N_m^{-a_1} \\ CN_m^{\overline{\sigma}+2-\frac{2}{3}a_1-\mu_2} &\leqslant & \frac{1}{3}N_m^{-a_1} \\ CN_m^{2\overline{\sigma}+4-\frac{4}{3}a_1} &\leqslant & \frac{1}{3}N_m^{-a_1} \end{array}$$

Actually, they are satisfied by taking N_0 large enough, that is ε small enough, provided that

$$\begin{cases} s_0 + 2\overline{\sigma} + \frac{2}{3}\mu_1 + 4 + a_1 \leqslant b_1 \\ \overline{\sigma} + \frac{1}{3}a_1 + 3 \leqslant \mu_2 \\ 2\overline{\sigma} + 5 \leqslant \frac{1}{3}a_1. \end{cases}$$
(10.41)

Hence it is easy to verify that (10.41) follows immediately from (10.2). Thus, we deduce for $m \ge 1$

$$\begin{aligned} \|\mathscr{F}(\widetilde{U}_{m+1})\|_{q,s_0}^{\gamma,O_{m+1}^{2\kappa}} &\leqslant C C_* N_0^{-1} \varepsilon N_m^{-a_1} \\ &\leqslant C_* \varepsilon N_m^{-a_1}. \end{aligned}$$
(10.42)

Concerning the case m = 0, we insert (10.25), (10.31), (10.33) and (10.40) into (10.21)

$$\|\mathscr{F}(\widetilde{U}_1)\|_{q,s_0}^{\kappa,\mathcal{O}_1^{2\kappa}} \leqslant CC_*\varepsilon \Big(N_0^{s_0+\overline{\sigma}+3+\frac{2}{3}\mu_1-b_1}+\varepsilon\kappa^{-2}+\varepsilon^3\kappa^{-2}\Big).$$

Then by virtue of (10.4) and the assumption on b_1 in (10.41) we get for ε small enough

$$C\left(N_0^{s_0+\overline{\sigma}+3+\frac{2}{3}\mu_1-b_1}+\varepsilon\kappa^{-2}+\varepsilon^3\kappa^{-2}\right)\leqslant 1$$

leading to

$$\|\mathscr{F}(U_1)\|_{q,s_0}^{\gamma,\mathcal{O}_1^{2\kappa}} \leqslant C_*\varepsilon.$$

▶ Extension and achievement of $(\mathcal{P}1)_{m+1}$ - $(\mathcal{P}2)_{m+1}$ - $(\mathcal{P}3)_{m+1}$. We shall now extend \widetilde{H}_{m+1} to the whole set of parameters \mathcal{O} using a suitable cut-off function. For this purpose, we consider a C^{∞} cut-off function $\chi_{m+1}: \mathcal{O} \to [0,1]$ defined by

$$\chi_{m+1}(\lambda) = \begin{cases} 1 & \text{in} & \mathcal{O}_{m+1}^{\kappa} \\ 0 & \text{in} & \mathcal{O} \setminus \mathcal{O}_{m+1}^{2\kappa} \end{cases}$$

and satisfying the additional conditions

$$\forall \alpha \in \mathbb{N}^d, \, |\alpha| \leqslant q, \quad \|\partial_\lambda^\alpha \chi_{m+1}\|_{L^\infty(6)} \lesssim \left(\kappa^{-1} N_m^{\overline{a}}\right)^{|\alpha|}. \tag{10.43}$$

Then we define the extension H_{m+1} of H_{m+1} by

$$H_{m+1}(\lambda) = \begin{cases} \chi_{m+1}(\lambda)\widetilde{H}_{m+1}(\lambda) & \text{ in } & \mathcal{O}_{m+1}^{2\kappa} \\ 0 & \text{ in } & \mathcal{O} \setminus \mathcal{O}_{m+1}^{2\kappa} \end{cases}$$

and the extension U_{m+1} is given by

$$U_{m+1} \triangleq U_m + H_{m+1}$$

Notice that

$$H_{m+1} = \widetilde{H}_{m+1}, \quad \mathscr{F}(U_{m+1}) = \mathscr{F}(\widetilde{U}_{m+1}) \quad \text{in} \quad \mathcal{O}_{m+1}^{\kappa}$$

Therefoe we infer from (10.42)

$$\left\|\mathscr{F}(U_{m+1})\right\|_{q,s_0}^{\gamma,\mathcal{O}_{m+1}^{\kappa}} \leqslant C_*\varepsilon N_m^{-a_1},$$

which achieves the induction of $(\mathcal{P}2)_{m+1}$.

Next, using the law product in Lemma 4.1 together with (10.43) we find - 0...

$$\forall s \in [s_0, S], \quad \|H_{m+1}\|_s^{q,\kappa,0} \lesssim N_m^{q\overline{a}} \|\widetilde{H}_{m+1}\|_{q,s}^{\gamma, \mathcal{O}_{m+1}^{2\kappa}}.$$
(10.44)

Combining (10.44) and (10.36) we deduce for $m \ge 1$

$$\|H_{m+1}\|_{s_0+\overline{\sigma}}^{q,\kappa,0} \leqslant N_m^{q\overline{a}+\overline{\sigma}} \|\widetilde{H}_{m+1}\|_{s_0}^{q,\kappa,\mathcal{O}_{m+1}^{2\kappa}}$$

$$\leqslant CC_*\varepsilon\kappa^{-1}N_m^{q\overline{a}+2\overline{\sigma}-\frac{2}{3}a_1}$$

From the choice done in (10.2) one gets

$$a_2 = \frac{2}{3}a_1 - q\overline{a} - 2\overline{\sigma} - 1 \ge 1 \tag{10.45}$$

and by choosing ε small enough we obtain

$$\|H_{m+1}\|_{s_0+\overline{\sigma}}^{q,\kappa,\theta} \leqslant C N_0^{-1} C_* \varepsilon \kappa^{-1} N_m^{-a_2} \leqslant C_* \varepsilon \kappa^{-1} N_m^{-a_2}.$$
(10.46)

For m = 0 we may combine (10.44) and (10.39) in order to get

$$\|H_1\|_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}}.$$
(10.47)

Let us define

$$W_{m+1} \triangleq W_m + H_{m+1},\tag{10.48}$$

then it is obvious that $U_{m+1} = U_0 + W_{m+1}$. In addition, applying $(\mathcal{P}1)_m$ with (10.46) and (10.47) we find in view of Lemma A.2 and for small ε

$$\|W_{m+1}\|_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant \|H_1\|_{s_0+\overline{\sigma}}^{q,\kappa} + \sum_{k=2}^{m+1} \|H_k\|_{s_0+\overline{\sigma}}^{q,\kappa}$$
$$\leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}} + C_*\varepsilon\kappa^{-1}\sum_{k=0}^{\infty} N_k^{-1}$$
$$\leqslant \frac{1}{2}C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}} + CC_*\varepsilon\kappa^{-1}N_0^{-1} \leqslant C_*\varepsilon\kappa^{-1}N_0^{q\overline{a}}$$

Now, applying (10.35), (10.44) and $(\mathcal{P}3)_m$, we deduce that

$$\begin{split} \|W_{m+1}\|_{b_{1}+\overline{\sigma}}^{q,\kappa} &\leqslant \|W_{m}\|_{b_{1}+\overline{\sigma}}^{q,\kappa} + CN_{m}^{q\overline{a}}\|H_{m+1}\|_{b_{1}+\overline{\sigma}}^{q,\kappa} \\ &\leqslant C_{*}\varepsilon\kappa^{-1}N_{m-1}^{\mu_{1}} + \kappa^{-1}N_{m}^{q\overline{a}+2\overline{\sigma}}\left(\varepsilon + \|W_{m}\|_{b_{1}+\overline{\sigma}}^{q,\kappa,0}\right) \\ &\leqslant CC_{*}\varepsilon\kappa^{-1}N_{m}^{q\overline{a}+2\overline{\sigma}+1+\frac{2}{3}\mu_{1}}. \end{split}$$

According to (10.2) one may check that

$$q\overline{a} + 2\overline{\sigma} + 2 = \frac{\mu_1}{3} \tag{10.49}$$

and therefore we deduce

$$\|W_{m+1}\|_{b_1+\overline{\sigma}}^{q,\kappa,\theta} \leqslant C N_0^{-1} C_* \varepsilon \kappa^{-1} N_m^{\mu_1}$$
$$\leqslant C_* \varepsilon \kappa^{-1} N_m^{\mu_1}$$

for ε small enough and (10.6) and this achieves the proof of $(\mathcal{P}3)_{m+1}$. This achieves the proof of $(\mathcal{P}1)_{m+1}$ and therefore the proof of Proposition 10.1 is now achieved.

As a by-product of Proposition 10.1 we shall construct solutions to the nonlinear equation (10.1) provided that the parameters belong to a suitable Cantor set. Later, we shall investigate in Section 10.2 the Lebesgue measure of this set.

Corollary 10.1. There exists $\varepsilon_0 > 0$ such that for any $\varepsilon \in (0, \varepsilon_0)$ the following assertions hold true. Consider the Cantor set (depending implicitly in ε)

$$\mathbf{G}^{\kappa}_{\infty} \triangleq \bigcap_{m \in \mathbb{N}} \mathscr{A}^{\kappa}_{m}$$

There exists a function (depending implicitly on ε)

$$U_{\infty}: \qquad \mathcal{O} \quad \to \quad \left(\mathbb{T}^{d} \times \mathbb{R}^{d} \times H_{\mathbb{S}}^{\perp}\right) \times \mathbb{R}^{d}$$
$$\lambda = (\omega, \alpha) \quad \mapsto \quad (i_{\infty}(\lambda), \mathbf{c}_{\infty}(\lambda))$$
$$214$$

such that

$$\forall \lambda \in \mathbf{G}_{\infty}^{\kappa} \quad \mathscr{F}(U_{\infty}(\lambda), \lambda, \varepsilon) = 0$$

In addition, the function $\lambda \in \mathcal{O} \mapsto c_{\infty}(\lambda, \varepsilon)$ belongs to $W^{q,\infty}(\mathcal{O})$ with

$$\mathbf{c}_{\infty}(\lambda,\varepsilon) = \omega + \mathbf{r}_{\varepsilon}(\lambda) \quad and \quad \|\mathbf{r}_{\varepsilon}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{q\overline{a}}.$$

Moreover, there exists a function $\alpha \in (\underline{\alpha}, \overline{\alpha}) \mapsto \lambda(\alpha, \varepsilon) \triangleq (\omega(\alpha, \varepsilon), \alpha)$ in $W^{q, \infty}$ with

$$\omega(\alpha,\varepsilon) = -\omega_{\rm Eq}(\alpha) + \bar{\mathbf{r}}_{\varepsilon}(\alpha), \qquad \|\bar{\mathbf{r}}_{\varepsilon}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{q\overline{a}}$$
(10.50)

and

$$\forall \alpha \in \mathsf{C}^{\kappa,\varepsilon}_{\infty}, \quad \mathscr{F}\left(U_{\infty}(\lambda(\alpha,\varepsilon),\varepsilon),\lambda(\alpha,\varepsilon),\varepsilon\right) = 0, \quad \mathsf{c}_{\infty}(\lambda(\alpha,\varepsilon),\varepsilon) = -\omega_{\mathrm{Eq}}(\alpha), \varepsilon \in \mathsf{C}^{\kappa,\varepsilon}_{\infty} \text{ is defined by}$$

where the Cantor set
$$C_{\infty}^{\kappa,\epsilon}$$
 is defined by

$$\mathbf{C}_{\infty}^{\kappa,\varepsilon} = \Big\{ \alpha \in (\underline{\alpha}, \overline{\alpha}), \big(\omega(\alpha, \varepsilon), \alpha \big) \in \mathbf{G}_{\infty}^{\kappa} \Big\}.$$

Proof. Combining
$$(10.48)$$
 and (10.46) , we obtain

$$\|W_{m+1} - W_m\|_{s_0}^{q,\kappa} = \|H_{m+1}\|_{s_0}^{q,\kappa} \leqslant \|H_{m+1}\|_{s_0+\overline{\sigma}}^{q,\kappa} \leqslant C_*\varepsilon\kappa^{-1}N_m^{-a_2}$$

Hence, the sequence $(W_m)_{m \in \mathbb{N}}$ is convergent and then we can define

$$W_{\infty} \triangleq \lim_{m \to \infty} W_m$$
$$\triangleq (\mathfrak{I}_{\infty}, \mathbf{c}_{\infty} - \omega)$$

and

$$U_{\infty} = (i_{\infty}, \mathbf{c}_{\infty}) \triangleq U_0 + W_{\infty} = \left((\varphi, 0, 0), \omega\right) + W_{\infty}$$

According to the point $(\mathcal{P}2)_m$ of Proposition 10.1, we have for small ε

$$\forall \lambda \in \mathsf{G}^{\gamma}_{\infty}, \quad \mathscr{F}\big(U_{\infty}(\lambda,\varepsilon),\lambda,\varepsilon\big) = 0.$$

We recall that \mathscr{F} is given in (5.21) and the Cantor set depends also in ε . Now, applying the point $(\mathscr{P}1)_m$ of Proposition 10.1, we obtain,

$$\mathbf{c}_{\infty}(\lambda,\varepsilon) = \omega + \mathbf{r}_{\varepsilon}(\lambda) \quad \text{with} \quad \|\mathbf{r}_{\varepsilon}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{q\overline{a}}.$$
(10.51)

Let us move to the second result and check the existence of solutions to the original Hamiltonian equation. First recall that the open set \mathcal{O} is given by

$$\mathscr{O} = \mathscr{U} \times (\underline{\alpha}, \overline{\alpha}), \quad \text{with} \quad \mathscr{U} \triangleq B(0, R),$$

with R large enough such that the open ball B(0, R) contains the equilibrium frequency vector $\{\omega_{Eq}(\alpha), \alpha \in [\underline{\alpha}, \overline{\alpha}]\}$ Applying (10.51), we find that for any $\alpha \in (\underline{\alpha}, \overline{\alpha})$ the partial mapping $\omega \mapsto c_{\infty}(\omega, \alpha)$ is invertible form \mathcal{U} into its image $c_{\infty}(\mathcal{U}, \alpha)$ and we have similarly to (9.33)

$$\widehat{\omega} = \mathsf{c}_{\infty}(\omega, \alpha) = \omega + \mathrm{r}_{\varepsilon}(\omega, \alpha) \Longleftrightarrow \omega = \mathsf{c}_{\infty}^{-1}(\widehat{\omega}, \alpha) = \widehat{\omega} + \widehat{\mathrm{r}}_{\varepsilon}(\widehat{\omega}, \alpha).$$

Applying Lemma 9.3 we deduce that $\widehat{\mathbf{r}}_{\varepsilon}$ satisfies the estimate

$$\widehat{\mathbf{r}}_{\varepsilon} \|^{q,\kappa} \lesssim \|\mathbf{r}_{\varepsilon}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{q\overline{a}}.$$
(10.52)

We denote

$$\omega(\alpha,\varepsilon) \triangleq \mathsf{c}_{\infty}^{-1} \big(-\omega_{\mathrm{Eq}}(\alpha), \alpha \big) = -\omega_{\mathrm{Eq}}(\alpha) + \overline{\mathsf{r}}_{\varepsilon}(\alpha) \quad \text{with} \quad \overline{\mathsf{r}}_{\varepsilon}(\alpha) \triangleq \widehat{\mathsf{r}}_{\varepsilon} \big(-\omega_{\mathrm{Eq}}(\alpha), \alpha \big)$$

and

$$\lambda(\alpha,\varepsilon) \triangleq (\omega(\alpha,\varepsilon),\alpha). \tag{10.53}$$

It follows that

$$\forall \, \alpha \in \mathsf{C}^{\kappa,\varepsilon}_{\infty}, \quad \mathscr{F}\big(U_{\infty}(\lambda(\alpha,\varepsilon)), \lambda(\alpha,\varepsilon), \varepsilon\big) = 0,$$

where the Cantor set $C_{\infty}^{\kappa,\varepsilon}$ was defined in Corollary 10.1. This gives nontrivial solutions for the original Hamiltonian equation provided that $\alpha \in C_{\infty}^{\kappa,\varepsilon}$. It remains to check the suitable estimates for

the function $\overline{\mathbf{r}}_{\varepsilon}$. According to Lemma 3.1-(vi) we know that all the derivatives $\partial_{\alpha}^{j}\omega_{\mathrm{Eq}}$ are uniformly bounded on the interval $(\underline{\alpha}, \overline{\alpha})$, then by the chain rule, (10.52) and (10.4) we find

$$\|\overline{\mathbf{r}}_{\varepsilon}\|^{q,\kappa} \lesssim \varepsilon \kappa^{-1} N_0^{q\overline{a}} \quad \text{and} \quad \|\omega(\cdot,\varepsilon)\|^{q,\kappa} \lesssim 1 + \varepsilon \kappa^{-1} N_0^{q\overline{a}} \lesssim 1.$$
(10.54)

The proof of Corollary 10.1 is now complete.

10.2. Final Cantor set estimates. The goal of this section is to give a lower bound Lebesgue measure for the Cantor set $C_{\infty}^{\kappa,\varepsilon}$ constructed in Corollary 10.1 and show that when $\varepsilon \to 0$ it will get a full Lebesgue measure. From Corollary 10.1, the Cantor set $C_{\infty}^{\kappa,\varepsilon}$ can be written in the form

$$\mathbf{C}_{\infty}^{\kappa,\varepsilon} = \bigcap_{m \in \mathbb{N}} \mathbf{C}_{m}^{\kappa} \quad \text{where} \quad \mathbf{C}_{m}^{\kappa} = \left\{ \alpha \in (\underline{\alpha}, \overline{\alpha}) \quad \text{s.t.} \quad \lambda(\alpha, \varepsilon) \in \mathscr{A}_{m}^{\kappa} \right\}, \tag{10.55}$$

where the sets \mathscr{A}_m^{κ} are defined in Proposition 10.1. Our result reads as follows

Proposition 10.2. Let q_0 as in Proposition 3.3 and assume (10.2) and (10.4) with $q = q_0 + 1$. Assume in addition that τ_1, τ_2 and ϱ satisfy

$$\begin{cases} \tau_1 > d q_0, \tau_2 > \frac{\tau_1}{1-2\overline{\alpha}} + d q_0, \\ 0 < \varrho < \min\left(\frac{1}{2} - \overline{\alpha}, \frac{1}{q_0+2}, 2\frac{1-a}{a}\right). \end{cases}$$

Then, there exists C > 0 and ε_0 such that for any $\varepsilon \in (0, \varepsilon_0]$

$$\left|\mathsf{C}_{\infty}^{\kappa,\varepsilon}\right| \geqslant \overline{\alpha} - \underline{\alpha} - C\varepsilon^{\frac{a\varrho}{q_{0}}}$$

In particular, $\lim_{\varepsilon \to 0} \left| \mathsf{C}_{\infty}^{\kappa, \varepsilon} \right| = \overline{\alpha} - \underline{\alpha}.$

Proof. By (10.55), we can write

$$|(\underline{\alpha}, \overline{\alpha}) \setminus \mathbf{C}_{\infty}^{\kappa, \varepsilon}| \leq |(\underline{\alpha}, \overline{\alpha}) \setminus \mathbf{C}_{0}^{\kappa}| + \sum_{m=0}^{\infty} |\mathbf{C}_{m}^{\kappa} \setminus \mathbf{C}_{m+1}^{\kappa}| \\ \triangleq \sum_{m=0}^{\infty} \mathscr{S}_{m},$$
(10.56)

where we use the notation |A| for the Lebesgue measure of a given measurable set A and the fact that by by construction $C_0^{\kappa} = (\underline{\alpha}, \overline{\alpha})$. According to the notations introduced in Proposition 9.6 and Proposition 9.5 combined with (10.53) one may write

$$\mu_{j}^{\infty,m}(\alpha,\varepsilon) \triangleq \mu_{j}^{\infty}\left(\lambda(\alpha,\varepsilon), i_{m}\right)$$
$$= \Omega_{j}(\alpha) + j r^{1,m}(\alpha,\varepsilon) - j \mathbb{W}(j,\alpha) r^{2,m}(\alpha,\varepsilon) + r_{j}^{\infty,m}(\alpha,\varepsilon)$$
(10.57)

with $W(j, \alpha)$ is defined in Lemma 3.1,

$$\ell \in \{1, 2\}, \quad r^{\ell, m}(\alpha, \varepsilon) \triangleq r^{\ell} \big(\lambda(\alpha, \varepsilon), i_m \big)$$

and

$$r_j^{\infty,m}(\alpha,\varepsilon) \triangleq r_j^{\infty} \big(\lambda(\alpha,\varepsilon), i_m \big).$$

Applying Proposition 9.5-(i) combined with Proposition $10.1-(\mathcal{P}1)_m$, (10.11), the chain rule and (10.54) yield

$$0 \leq k \leq q, \quad \sup_{m \in \mathbb{N}} \sup_{\alpha \in (\underline{\alpha}, \overline{\alpha})} |\partial_{\alpha}^{k} r^{\ell, m}(\alpha, \varepsilon)| \lesssim \sup_{m \in \mathbb{N}} \varepsilon \kappa^{-1-k} \left(1 + \|\mathfrak{I}_{m}\|_{\overline{s}_{h}}^{q, \kappa} \right) \\ \lesssim \varepsilon \kappa^{-1-k}.$$
(10.58)

In a similar way, using (9.280), (10.54) and Proposition 10.1- $(\mathcal{P}1)_m$, we find

$$0 \leq k \leq q, \quad \sup_{m \in \mathbb{N}} \sup_{j \in \mathbb{S}_{0}^{c}} \sup_{\alpha \in (\underline{\alpha}, \overline{\alpha})} |j|^{1-\epsilon-2\overline{\alpha}} |\partial_{\alpha}^{k} r_{j}^{\infty, m}(\alpha, \varepsilon)| \leq \kappa^{-k} \sup_{m \in \mathbb{N}} \sup_{j \in \mathbb{S}_{0}^{c}} |j|^{1-\epsilon-2\overline{\alpha}} ||r_{j}^{\infty}(i_{m})||^{q, \kappa} \leq \varepsilon \kappa^{-2-k}.$$

$$(10.59)$$

Coming back to (10.55) and using the Cantor sets introduced in Proposition 9.6, Theorem 9.1 and Proposition 9.2 one obtains by construction that for any $m \in \mathbb{N}$

$$\mathbf{C}_{m}^{\kappa} \backslash \mathbf{C}_{m+1}^{\kappa} = \bigcup_{\substack{|l| \leqslant N_{m}, \\ j \in \mathbb{Z} \backslash \{0\}}} \mathscr{R}_{l,j}^{(0)}(i_{m}) \bigcup_{\substack{|l| \leqslant N_{m} \\ j, j_{0} \in \mathbb{S}_{0}^{c}}} \mathscr{R}_{l,j,j_{0}}(i_{m}) \bigcup_{\substack{|l| \leqslant N_{m} \\ j \in \mathbb{S}_{0}^{c}}} \mathscr{R}_{l,j}^{(1)}(i_{m}),$$
(10.60)

where

$$\begin{split} & \mathscr{R}_{l,j}^{(0)}(i_m) \triangleq \left\{ \alpha \in \mathsf{C}_m^{\kappa}; \ |\omega(\alpha,\varepsilon) \cdot l + jc_m(\alpha,\varepsilon)| \leqslant \frac{4\kappa_{m+1}^{\varepsilon}\langle j \rangle}{\langle l \rangle^{\tau_1}} \right\}, \\ & \mathscr{R}_{l,j,j_0}(i_m) \triangleq \left\{ \alpha \in \mathsf{C}_m^{\kappa}; \ |\omega(\alpha,\varepsilon) \cdot l + \mu_j^{\infty,m}(\lambda,\varepsilon) - \mu_{j_0}^{\infty,m}(\lambda,\varepsilon)| \leqslant \frac{2\kappa_{m+1}\langle j - j_0 \rangle}{\langle l \rangle^{\tau_2}} \right\} \\ & \mathscr{R}_{l,j}^{(1)}(i_m) \triangleq \left\{ \alpha \in \mathsf{C}_m^{\kappa}; \ |\omega(\alpha,\varepsilon) \cdot l + \mu_j^{\infty,m}(\alpha,\varepsilon)| \leqslant \frac{\kappa_{m+1}\langle j \rangle}{\langle l \rangle^{\tau_1}} \right\}, \end{split}$$

with the following definition $c_m(\alpha, \varepsilon) \triangleq c(\lambda(\alpha, \varepsilon), i_m(\lambda(\alpha, \varepsilon), \varepsilon))$, where the coefficient $c(\lambda, i_0)$ was introduced in Proposition 9.2. Let us now move to the estimate of \mathscr{S}_m defined in (10.56). Then using (10.60) together with Lemma 10.1-(iv) and Lemma 10.2, we find

$$\begin{aligned} \mathscr{S}_{m} &\leqslant \sum_{\substack{(l,j) \in \mathbb{Z}^{d+1} \setminus \{0\} \\ |j| \leqslant C_{0}\langle l \rangle, |l| > N_{m-1} \ }} \left| \mathscr{R}_{l,j}^{(0)}(i_{m}) \right| + \sum_{\substack{(l,j,j_{0}) \in \mathbb{Z}^{d} \times (\mathbb{S}_{0}^{c})^{2} \\ |j-j_{0}| \leqslant C_{0}\langle l \rangle, |l| > N_{m-1} \\ \min(|j|,|j_{0}|)^{1-2\overline{\alpha}} - \epsilon_{\leqslant c_{2}\kappa} - \varrho_{\langle l \rangle} \tau_{1}} \right. \\ &+ \sum_{\substack{(l,j) \in \mathbb{Z}^{d} \times \mathbb{S}_{0}^{c} \\ |j| \leqslant C_{0}\langle l \rangle, |l| > N_{m-1} \ }} \left| \mathscr{R}_{l,j}^{(1)}(i_{m}) \right|. \end{aligned}$$

By a direct application of Lemma A.3 combined with Lemma 10.3 we obtain for any $m \in \mathbb{N}$

$$\begin{aligned} |\mathscr{R}_{l,j}^{(0)}(i_m)| &\lesssim \kappa^{\frac{\varrho}{q_0}} \langle j \rangle^{\frac{1}{q_0}} \langle l \rangle^{-1 - \frac{\tau_1 + 1}{q_0}} \\ |\mathscr{R}_{l,j}^{(1)}(i_m)| &\lesssim \kappa^{\frac{1}{q_0}} \langle j \rangle^{\frac{1}{q_0}} \langle l \rangle^{-1 - \frac{\tau_1 + 1}{q_0}} \\ |\mathscr{R}_{l,j,j_0}(i_m)| &\lesssim \kappa^{\frac{1}{q_0}} \langle j - j_0 \rangle^{\frac{1}{q_0}} \langle l \rangle^{-1 - \frac{\tau_2 + 1}{q_0}}. \end{aligned}$$
(10.61)

Notice that if $|j - j_0| \leq C_0 \langle l \rangle$ and $\min(|j|, |j_0|)^{1-2\overline{\alpha}-\epsilon} \lesssim \kappa^{-\varrho} \langle l \rangle^{\tau_1}$, then

$$\max(|j|, |j_0|) = \min(|j|, |j_0|) + |j - j_0| \lesssim \left(\kappa^{-\varrho} \langle l \rangle^{\tau_1}\right)^{\frac{1}{1 - 2\overline{\alpha} - \epsilon}} + C_0 \langle l \rangle \lesssim \kappa^{-\frac{\varrho}{1 - 2\overline{\alpha} - \epsilon}} \langle l \rangle^{\frac{\tau_1}{1 - 2\overline{\alpha} - \epsilon}}$$

Hence using (10.61) allows to get

$$\sum_{m\in\mathbb{N}}\mathscr{S}_{m} \lesssim \kappa^{\frac{1}{q_{0}}} \Big(\sum_{\substack{m\in\mathbb{N}\\|l|>N_{m-1}}} \langle l \rangle^{-\frac{\tau_{1}}{q_{0}}} + \kappa^{-\frac{\varrho}{q(1-2\overline{\alpha}-\epsilon)}} \sum_{\substack{m\in\mathbb{N}\\|l|>N_{m-1}}} \langle l \rangle^{\frac{\tau_{1}}{q_{0}(1-2\overline{\alpha}-\epsilon)} - \frac{\tau_{2}}{q_{0}}} \Big) + \kappa^{\frac{\varrho}{q_{0}}} \sum_{\substack{m\in\mathbb{N}\\|l|>N_{m-1}}} \langle l \rangle^{-\frac{\tau_{1}}{q_{0}}} \\ \lesssim \kappa^{\min\left(\frac{\varrho}{q_{0}}, \frac{1}{q_{0}} - \frac{\varrho}{q_{0}(1-2\overline{\alpha}-\epsilon)}\right)}, \tag{10.62}$$

provided that

$$\tau_1 > d q_0 \quad \text{and} \quad \tau_2 > \frac{\tau_1}{1 - 2\overline{\alpha}} + d q_0,$$
 (10.63)

and ϵ is chosen small enough. Thus, by imposing

$$0 < \varrho < \frac{1}{2} - \overline{\alpha} \tag{10.64}$$

we obtain from (10.62)

$$\sum_{m \in \mathbb{N}} \mathscr{S}_m \lesssim \kappa^{\frac{\varrho}{q_0}}.$$
(10.65)

Inserting (10.65) into (10.56) implies

$$\frac{|(\underline{\alpha}, \overline{\alpha}) \setminus \mathbf{C}_{\infty}^{\kappa, \varepsilon}|}{217} \lesssim \kappa^{\frac{\mathcal{U}}{q_0}},$$

provided that the conditions (10.63) and (10.64) are satisfied. By making the choice

$$\varrho < \min\left(\frac{1}{2} - \overline{\alpha}, \frac{1}{q_0 + 2}, 2\frac{1 - a}{a}\right) \tag{10.66}$$

which is compatible with the condition on ρ given by Proposition 9.2 with $q = q_0 + 1$, (10.64) and Lemma 10.1, we find, since $\kappa = \varepsilon^a$ according to (10.4),

$$|(\underline{\alpha}, \overline{\alpha}) \setminus \mathbf{C}_{\infty}^{\kappa, \varepsilon}| \lesssim \varepsilon^{\frac{a\varrho}{q_0}}.$$

This completes the proof of the proposition.

It remains now to prove Lemma 10.1 and Lemma 10.2 below, used in the proof of Proposition 10.2.

Lemma 10.1. Assume (10.2), (10.4) and $\rho \in \left(0, \frac{2(1-a)}{a}\right)$. Let $m \in \mathbb{N}^*$ and $l \in \mathbb{Z}^d$ such that $|l| \leq N_{m-1}$. Then the following assertions hold true.

- (i) For $j \in \mathbb{Z}$ with $(l, j) \neq (0, 0)$, we get $\mathscr{R}_{l, i}^{(0)}(i_m) = \varnothing$.
- (ii) For $(j, j_0) \in (\mathbb{S}_0^c)^2$ with $(l, j) \neq (0, j_0)$, we get $\mathscr{R}_{l, j, j_0}(i_m) = \varnothing$.
- (iii) For $j \in \mathbb{S}_0^c$ with $(l,j) \neq (0,0)$, we get $\mathscr{R}_{l,j}^{(1)}(i_m) = \varnothing$.
- (iv) For any $m \in \mathbb{N}$,

$$\mathbf{C}_{m}^{\kappa} \backslash \mathbf{C}_{m+1}^{\kappa} = \bigcup_{\substack{(l,j) \in \mathbb{Z}^{d} \times \mathbb{Z} \backslash \{(0,0)\} \\ |l| > N_{m-1}}} \mathcal{R}_{l,j}^{(0)}(i_{m}) \cup \bigcup_{\substack{(l,j,j_{0}) \in \mathbb{Z}^{d} \times (\mathbb{S}_{0}^{c})^{2} \\ |l| > N_{m-1}}} \mathcal{R}_{l,j,j_{0}}(i_{m}) \cup \bigcup_{\substack{(l,j) \in \mathbb{Z}^{d} \times \mathbb{S}_{0}^{c} \\ |l| > N_{m-1}}} \mathcal{R}_{l,j}^{(1)}(i_{m}).$$

Proof. \blacktriangleright The key estimate that will be used for all the points comes from (10.11)

$$\|i_{m} - i_{m-1}\|_{\overline{s}_{h} + \sigma_{4}}^{q,\kappa} \leqslant \|U_{m} - U_{m-1}\|_{\overline{s}_{h} + \sigma_{4}}^{q,\kappa} = \|H_{m}\|_{\overline{s}_{h} + \sigma_{4}}^{q,\kappa} \leqslant C_{*}\varepsilon\kappa^{-1}N_{m-1}^{-a_{2}}.$$
(10.67)

(i) We shall first prove that if $|l| \leq N_{m-1}$ and $(l,j) \neq (0,0)$, then $\mathscr{R}_{l,j}^{(0)}(i_m) \subset \mathscr{R}_{l,j}^{(0)}(i_{m-1})$. Assume for a while this inclusion and let us check how this implies that $\mathscr{R}_{l,j}^{(0)}(i_m) = \emptyset$. According to (10.60) one deduces that

$$\mathscr{R}_{l,j}^{(0)}(i_m) \subset \mathscr{R}_{l,j}^{(0)}(i_{m-1}) \subset \mathsf{C}_{m-1}^{\kappa} \backslash \mathsf{C}_m^{\kappa}.$$

Using once again (10.60) gives o $\mathscr{R}_{l,i}^{(0)}(i_m) \subset \mathsf{C}_m^{\kappa} \setminus \mathsf{C}_{m+1}^{\kappa}$, and therefore we get

$$\mathscr{R}_{l,j}^{(0)}(i_m) \subset (\mathsf{C}_m^{\kappa} \backslash \mathsf{C}_{m+1}^{\kappa}) \cap (\mathsf{C}_{m-1}^{\kappa} \backslash \mathsf{C}_m^{\kappa}) = \varnothing.$$

Let us now turn to the proof of the claim $\mathscr{R}_{l,j}^{(0)}(i_m) \subset \mathscr{R}_{l,j}^{(0)}(i_{m-1})$. To do that, consider a point $\alpha \in \mathscr{R}_{l,j}^{(0)}(i_m)$. Then we get first by construction that $\alpha \in C_m^{\kappa} \subset C_{m-1}^{\kappa}$. In addition, by the triangle inequality and Sobolev embeddings we deduce

$$\begin{aligned} \left| \omega(\alpha,\varepsilon) \cdot l + jc_{m-1}(\alpha,\varepsilon) \right| &\leq \left| \omega(\alpha,\varepsilon) \cdot l + jc_m(\alpha,\varepsilon) \right| + |j| \left| c_m(\alpha,\varepsilon) - c_{m-1}(\alpha,\varepsilon) \right| \\ &\leq 4 \frac{\kappa_{m+1}^{\varrho}|j|}{\langle l \rangle^{\tau_1}} + C|j| \|c(\cdot,i_m) - c(\cdot,i_{m-1})\|^{q,\kappa}. \end{aligned}$$

Thus applying (9.43), (10.67) and (10.4) we obtain with the choice $\kappa = \varepsilon^a$

$$\begin{aligned} \left| \omega(\alpha,\varepsilon) \cdot l + jc_{m-1}(\alpha,\varepsilon) \right| &\leq \frac{4\kappa_{m+1}^{\nu}|j|}{\langle l\rangle^{\tau_1}} + C\varepsilon\kappa^{-1}|j| \|i_m - i_{m-1}\|_{\overline{s}_h+4}^{q,\kappa} \\ &\leq \frac{4\kappa_{m+1}^{\rho}|j|}{\langle l\rangle^{\tau_1}} + C\varepsilon^{2(1-a)}|j|N_{m-1}^{-a_2}. \end{aligned}$$

According to the definition of κ_m in Proposition 10.1- $(\mathcal{P}2)_m$ one gets

$$\exists c_0 > 0, \quad \text{s.t} \quad \forall m \in \mathbb{N}, \quad \kappa_{m+1}^{\varrho} - \kappa_m^{\varrho} \leqslant -c_0 \, \kappa^{\varrho} 2^{-m}.$$

Then under the assumption

$$a_2 > \tau_1$$
 and $2(1-a) - a\varrho > 0$ (10.68)

one finds successively that $\sup_{m\in\mathbb{N}} 2^m N_{m-1}^{-a_2+\tau_1} < \infty$ and when ε is small enough and $|l| \leq N_{m-1}$ we deduce that

$$\begin{aligned} \left| \omega(\alpha,\varepsilon) \cdot l + jc_{m-1}(\alpha,\varepsilon) \right| &\leq \frac{4\kappa_m^{\varrho}|j|}{\langle l \rangle^{\tau_1}} + \frac{|j|\kappa^{\varrho}}{2^m \langle l \rangle^{\tau_1}} \Big(-4c_0 + C\varepsilon^{2(1-a)-a\varrho} 2^m N_{m-1}^{-a_2+\tau_1} \Big) \\ &< \frac{4\kappa_m^{\varrho}|j|^{\varrho}}{\langle l \rangle^{\tau_1}}. \end{aligned}$$

Hence we obtain $\alpha \in \mathcal{R}_{l,j}^{(0)}(i_{m-1})$ and this achieves the proof of the first point. Observe that the first assumption in (10.68) is automatically satisfied from (10.2) since $q \ge 1$.

(ii) Let $j, j_0 \in \mathbb{S}_0^c$ and $(l, j) \neq (0, j_0)$. Imitating the proof of the first point (i), then to get the desired result it is enough to check that $\mathcal{R}_{l,j,j_0}(i_m) \subset \mathcal{R}_{l,j,j_0}(i_{m-1})$. Let $\alpha \in \mathcal{R}_{l,j,j_0}(i_m)$ then from the definition of this set introduced in (10.60) we deduce that $\alpha \in \mathbb{C}_m^{\kappa} \subset \mathbb{C}_{m-1}^{\kappa}$ and

$$\begin{aligned} \left| \omega(\alpha,\varepsilon) \cdot l + \mu_{j}^{\infty,m-1}(\alpha,\varepsilon) - \mu_{j_{0}}^{\infty,m-1}(\alpha,\varepsilon) \right| &\leq \frac{2\kappa_{m+1}\langle j-j_{0}\rangle}{\langle l\rangle^{\tau_{2}}} \\ &+ \left| \mu_{j}^{\infty,m}(\alpha,\varepsilon) - \mu_{j_{0}}^{\infty,m}(\alpha,\varepsilon) - \mu_{j}^{\infty,m-1}(\alpha,\varepsilon) + \mu_{j_{0}}^{\infty,m-1}(\alpha,\varepsilon) \right|. \end{aligned}$$
(10.69)

 Set

$$\varrho_{j,j_0}^m(\alpha,\varepsilon) \triangleq \left| \mu_j^{\infty,m}(\alpha,\varepsilon) - \mu_{j_0}^{\infty,m}(\alpha,\varepsilon) - \mu_j^{\infty,m-1}(\alpha,\varepsilon) + \mu_{j_0}^{\infty,m-1}(\alpha,\varepsilon) \right|$$

Then coming back to (10.57) we deduce that

$$\varrho_{j,j_0}^m(\alpha,\varepsilon) \leqslant |j-j_0| |r^{1,m}(\alpha,\varepsilon) - r^{1,m-1}(\alpha,\varepsilon)| + |j \mathbb{W}(j,\alpha) - j_0 \mathbb{W}(j_0,\alpha)| |r^{2,m}(\alpha,\varepsilon) - r^{2,m-1}(\alpha,\varepsilon)| \\
+ |r_j^{\infty,m}(\alpha,\varepsilon) - r_j^{\infty,m-1}(\alpha,\varepsilon)| + |r_{j_0}^{\infty,m}(\alpha,\varepsilon) - r_{j_0}^{\infty,m-1}(\alpha,\varepsilon)|.$$
(10.70)

According to Proposition 9.5-(i), (10.67) and (10.4) one gets for k = 1, 2

$$\begin{aligned} \left| r^{k,m}(\alpha,\varepsilon) - r^{k,m-1}(\alpha,\varepsilon) \right| &\lesssim \varepsilon \kappa^{-1} \| i_m - i_{m-1} \|_{\overline{s}_h + \sigma_3}^{q,\kappa} \\ &\lesssim \varepsilon^2 \kappa^{-2} N_{m-1}^{-a_2} \\ &\lesssim \varepsilon^{2-2a} \langle j - j_0 \rangle N_{m-1}^{-a_2}. \end{aligned}$$

Next, using Proposition 9.6-(ii), (10.67) and (10.4) and using the fact (which follows from the last assumption in (10.2) since $\overline{\sigma}$ can be taken large enough)

$$s_h \geqslant 2\overline{s}_h + 8\tau_2(q+1)$$

yield

$$\begin{aligned} \left| r_{j}^{\infty,m}(\alpha,\varepsilon) - r_{j}^{\infty,m-1}(\alpha,\varepsilon) \right| &\lesssim \varepsilon \kappa^{-2} \left(\| i_{m} - i_{m-1} \|_{\overline{s}_{h}+\sigma_{4}}^{q,\kappa} \right)^{\frac{1}{2}} \\ &\lesssim \varepsilon^{\frac{3}{2}} \kappa^{-\frac{5}{2}} N_{m-1}^{-\frac{1}{2}a_{2}} \\ &\lesssim \varepsilon^{\frac{3-5a}{2}} \langle j - j_{0} \rangle N_{m-1}^{-\frac{1}{2}a_{2}}. \end{aligned}$$

Plugging the two preceding estimates into (10.70) and using the inequality (3.38) we find

$$\varrho_{j,j_0}^m(\alpha,\varepsilon) \lesssim \varepsilon^{\frac{3-5a}{2}} \langle j - j_0 \rangle N_{m-1}^{-\frac{1}{2}a_2}.$$
(10.71)

Combining (10.71) with (10.69) and using $\kappa_{m+1} = \kappa_m - \varepsilon^a 2^{-m-1}$ allow to get

$$\begin{split} \left| \omega(\alpha,\varepsilon) \cdot l + \mu_j^{\infty,m-1}(\alpha,\varepsilon) - \mu_{j_0}^{\infty,m-1}(\alpha,\varepsilon) \right| &\leq \frac{2\kappa_m \langle j-j_0 \rangle}{\langle l \rangle^{\tau_2}} - \varepsilon^a \langle j-j_0 \rangle 2^{-m} \langle l \rangle^{-\tau_2} \\ &+ C \varepsilon^{\frac{3-5a}{2}} \langle j-j_0 \rangle N_{m-1}^{-\frac{1}{2}a_2}. \end{split}$$

Now we write, since $|l| \leq N_{m-1}$

$$-\varepsilon^{a}2^{-m}\langle l\rangle^{-\tau_{2}} + C\varepsilon^{\frac{3-5a}{2}}N_{m-1}^{-\frac{1}{2}a_{2}} \leqslant \varepsilon^{a}2^{-m}\langle l\rangle^{-\tau_{2}}\Big(-1 + C\varepsilon^{\frac{3-7a}{2}}2^{m}N_{m-1}^{\tau_{2}-\frac{1}{2}a_{2}}\Big).$$

From (10.2) and (10.4) it is obvious that

$$2\tau_2 < a_2$$
 and $a < \frac{3}{7}$ (10.72)

implying in turn for ε small enough

$$\forall m \in \mathbb{N}, \quad -1 + C\varepsilon^{\frac{3-7a}{2}} 2^m N_{m-1}^{\tau_2 - \frac{1}{2}a_2} \leqslant 0.$$

Therefore we obtain

$$\omega(\alpha,\varepsilon) \cdot l + \mu_j^{\infty,m-1}(\alpha,\varepsilon) - \mu_{j_0}^{\infty,m-1}(\alpha,\varepsilon) \Big| \leqslant \frac{2\kappa_m \langle j-j_0 \rangle}{\langle l \rangle^{\tau_2}} \cdot$$

This shows that $\alpha \in \mathcal{R}_{l,i,j_0}(i_{m-1})$, which completes the proof of the second point.

(iii) Let $j \in \mathbb{S}_{0}^{c}, (l, j) \neq (0, 0)$, we shall first prove that if $|l| \leq N_{m-1}$ then $\mathscr{R}_{l,j}^{(1)}(i_{m}) \subset \mathscr{R}_{l,j}^{(1)}(i_{m-1})$. As in the point (i) this implies that $\mathscr{R}_{l,j}^{(1)}(i_m) = \varnothing$. Remind that the set $\mathscr{R}_{l,j}^{(1)}(i_m)$ is defined below the form (10.60). Let $\alpha \in \mathcal{R}_{l,j}^{(1)}(i_m)$ then by construction $\alpha \in C_m^{\kappa} \subset C_{m-1}^{\kappa}$.

Now by the triangle inequality we may write in view of Proposition 9.6-(ii), (10.67) and $\kappa = \varepsilon^a$

$$\begin{split} \left| \omega(\alpha,\varepsilon) \cdot l + \mu_{j}^{\infty,m-1}(\alpha,\varepsilon) \right| &\leq \left| \omega(\alpha,\varepsilon) \cdot l + \mu_{j}^{\infty,m}(\alpha,\varepsilon) \right| + \left| \mu_{j}^{\infty,m}(\alpha,\varepsilon) - \mu_{j}^{\infty,m-1}(\alpha,\varepsilon) \right| \\ &\leq \frac{\kappa_{m+1}\langle j \rangle}{\langle l \rangle^{\tau_{1}}} + C\varepsilon\kappa^{-2} |j| \left(\|i_{m} - i_{m-1}\|_{\overline{s}_{h}+\sigma_{4}}^{q,\kappa} \right)^{\frac{1}{2}} \\ &\leq \frac{\kappa_{m+1}\langle j \rangle}{\langle l \rangle^{\tau_{1}}} + C\varepsilon^{\frac{3-5a}{2}} \langle j \rangle N_{m-1}^{-\frac{1}{2}a_{2}}. \end{split}$$

Since $\kappa_{m+1} = \kappa_m - \varepsilon^a 2^{-m-1}$ and $|l| \leq N_{m-1}$ then

$$\left|\omega(\alpha,\varepsilon)\cdot l+\mu_{j}^{\infty,m-1}(\alpha,\varepsilon)\right| \leqslant \frac{\kappa_{m}\langle j\rangle}{\langle l\rangle^{\tau_{1}}} + \frac{\langle j\rangle\varepsilon^{a}}{2^{m+1}\langle l\rangle^{\tau_{1}}} \left(-1+\varepsilon^{\frac{3-7a}{2}}2^{m+1}N_{m-1}^{\tau_{1}-\frac{1}{2}a_{2}}\right).$$

Putting together (9.2) and (10.72) yields

$$a < \frac{3}{7}$$
 and $2\tau_1 < a_2$

and by taking ε small enough we find that

$$\forall m \in \mathbb{N}, \quad -1 + \varepsilon^{2-4a} 2^{m+1} N_{m-1}^{\tau_1 - a_2} \leqslant 0.$$

This implies in turn that

$$\left|\omega(\alpha,\varepsilon)\cdot l+\mu_{j}^{\infty,m-1}(\alpha,\varepsilon)\right|\leqslant \tfrac{\kappa_{m}\langle j\rangle}{\langle l\rangle^{\tau_{1}}}\cdot$$

It follows that $\lambda \in \mathcal{R}_{l,i}^{(1)}(i_{m-1})$, which completes the proof.

(iv) The proof follows easily from (10.60) and the points Lemma 10.1-(i)-(ii)-(iii).

The next result deals with necessary conditions such that the sets in (10.60) are not empty.

Lemma 10.2. Let $a \in (0, \frac{1}{2})$ and $\tau_2 > \tau_1$, there exists ε_0 such that for any $\varepsilon \in [0, \varepsilon_0]$ and $m \in \mathbb{N}$, the following assertions hold true.

- (i) Let $(l,j) \in \mathbb{Z}^d \times \mathbb{Z} \setminus \{(0,0)\}$. If $\mathscr{R}_{l,i}^{(0)}(i_m) \neq \emptyset$, then $|j| \leq C_0 \langle l \rangle$.
- (ii) Let $(l, j, j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2$. If $\mathcal{R}_{l, j, j_0}(i_m) \neq \emptyset$, then $|j j_0| \leq C_0 \langle l \rangle$.
- (iii) Let $(l, j) \in \mathbb{Z}^d \times \mathbb{S}_0^c$. If $\mathscr{R}_{l,j}^{(1)}(i_m) \neq \emptyset$, then $|j| \leq C_0 \langle l \rangle$. (iv) Let $(l, j, j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2$. If $(\min(|j|, |j_0|))^{1-2\overline{\alpha}-\epsilon} \geq c_2 \kappa^{\varrho} \langle l \rangle^{\tau_1}$, then

$$\mathscr{R}_{l,j,j_0}(i_m) \subset \mathscr{R}_{l,j-j_0}^{(0)}(i_m)$$

with c_2 being a fixed constant and $\epsilon > 0$ is an arbitrary small number.

Proof. (i) Assume $\mathscr{R}_{l,j}^{(0)}(i_m) \neq \emptyset$ then by construction we can find $\alpha \in (\underline{\alpha}, \overline{\alpha})$ such that,

$$c_{m}(\alpha,\varepsilon)||j| \leqslant 4\langle j\rangle \kappa_{m+1}^{\varrho} \langle l\rangle^{-\tau_{1}} + |\omega(\alpha,\varepsilon) \cdot l|$$

$$\leqslant 4\langle j\rangle \kappa_{m+1}^{\varrho} + C\langle l\rangle$$

$$\leqslant C\varepsilon^{a\varrho} \langle j\rangle + C\langle l\rangle, \qquad (10.73)$$

where we have used $\kappa = \varepsilon^a$ and the fact that $(\alpha, \varepsilon) \mapsto \omega(\alpha, \varepsilon)$ is bounded. Applying (9.40) combined with Proposition 10.1- $(\mathcal{P}1)_m$ we write (by taking $\overline{\sigma}$ large enough)

$$|c_m(\alpha,\varepsilon) - V_{\alpha,0}| \leq ||c_m - V_{0,\alpha}||^{q,\kappa} \leq \varepsilon \kappa^{-1} \left(1 + ||\mathfrak{I}_m||^{q,\kappa}_{\overline{s}_h + \sigma_1}\right) \leq \varepsilon^{1-a}.$$
(10.74)

On the other hand, one deduces from (3.11) that

$$\inf_{\alpha \in (\underline{\alpha}, \overline{\alpha})} V_0(\alpha) \triangleq c_1 > 0,$$

and then taking ε small enough, we obtain

$$\inf_{m \in \mathbb{N}} \inf_{\alpha \in (\underline{\alpha}, \overline{\alpha})} |c_m(\alpha, \varepsilon)| \ge \frac{c_1}{2} \cdot$$

Thus, coming back to (10.73) and taking ε small enough we deduce that $|j| \leq C_0 \langle l \rangle$ for some constant $C_0 > 0$.

(ii) First, observe that the case $j = j_0$ is trivial and the conclusion is true, then we shall assume that $j \neq j_0$. If $\Re_{l,j,j_0}(i_m) \neq \emptyset$ then there exists $\alpha \in (\underline{\alpha}, \overline{\alpha})$ such that

$$\begin{aligned} |\mu_{j}^{\infty,m}(\alpha,\varepsilon) - \mu_{j_{0}}^{\infty,m}(\alpha,\varepsilon)| &\leq 2\kappa_{m+1}\langle j - j_{0}\rangle\langle l\rangle^{-\tau_{2}} + |\omega(\alpha,\varepsilon) \cdot l| \\ &\leq 2\kappa_{m+1}\langle j - j_{0}\rangle + C\langle l\rangle \\ &\leq 4\varepsilon^{a}\langle j - j_{0}\rangle + C\langle l\rangle. \end{aligned}$$

By virtue of the triangle inequality, Lemma 3.1-(iv), (10.57), (3.38), (10.58) and (10.59) we find

$$\begin{aligned} |\mu_j^{\infty,m}(\alpha,\varepsilon) - \mu_{j_0}^{\infty,m}(\alpha,\varepsilon)| &\ge |\Omega_j(\alpha) - \Omega_{j_0}(\alpha)| - \left(|r^{1,m}(\alpha,\varepsilon)| + |r^{2,m}(\alpha,\varepsilon)|\right)|j - j_0| \\ &- |r_j^{\infty,m}(\alpha,\varepsilon)| - |r_{j_0}^{\infty,m}(\alpha,\varepsilon)| \\ &\ge \left(C_0 - C\varepsilon^{1-2a}\right)|j - j_0| \ge \frac{C_0}{2}|j - j_0| \end{aligned}$$

provided that ε is small enough. Then combining the preceding inequalities implies for ε small enough that $|j - j_0| \leq C_0 \langle l \rangle$, for some $C_0 > 0$.

(iii) Notice that for j = 0 the conclusion is always true. Now assume for $j \neq 0$ that $\mathscr{R}_{l,j}^{(1)}(i_m) \neq \emptyset$ then there exists $\alpha \in (\underline{\alpha}, \overline{\alpha})$ such that

$$\begin{aligned} |\mu_j^{\infty,m}(\alpha,\varepsilon)| &\leq \kappa_{m+1} \langle j \rangle \langle l \rangle^{-\tau_1} + |\omega(\alpha,\varepsilon) \cdot l| \\ &\leq 2\varepsilon^a |j| + C \langle l \rangle. \end{aligned}$$

Using the definition (10.57) combined with the triangle inequality, Lemma 3.1-(iii), (10.58) and (10.59), we obtain

$$|\mu_j^{\infty,m}(\alpha,\varepsilon)| \ge C_0|j| - |j| (|r^{1,m}(\alpha,\varepsilon)| + |r^{2,m}(\alpha,\varepsilon)|) - |r_j^{\infty,m}(\alpha,\varepsilon)| \ge C_0|j| - C\varepsilon^{1-2a}|j|.$$

Putting together these inequalities yields, provided that $a \in (0, \frac{1}{2})$,

$$(C_0 - C\varepsilon^{1-2a} - 2\varepsilon^a)|j| \leq C\langle l \rangle$$

Therefore by taking ε small enough, we find $|j| \leq C_0 \langle l \rangle$, for some $C_0 > 0$. (iv) The case $j = j_0$ is trivial and follows from the definition (10.60) and the fact $\tau_2 > \tau_1$. Consider $j \neq j_0$ and $\alpha \in \mathcal{R}_{l,j,j_0}(i_m)$ then by definition

$$\left|\omega(\alpha,\varepsilon)\cdot l+\mu_{j}^{\infty,m}(\alpha,\varepsilon)-\mu_{j_{0}}^{\infty,m}(\alpha,\varepsilon)\right| \leqslant \frac{2\kappa_{m+1}\langle j-j_{0}\rangle}{\langle l\rangle^{\tau_{2}}}$$

Combining (10.57) with Lemma 3.1-(iii) gives the asymptotic

$$\mu_j^{\infty,m}(\alpha,\varepsilon) = j c_m(\alpha,\varepsilon) - j|j|^{\alpha-1} \left(W_{0,\alpha} + r^{2,m}(\alpha,\varepsilon) \right) + r_j^{\infty,m}(\alpha,\varepsilon) + O\left(\frac{1}{|j|^{1-\alpha}}\right).$$

Therefore using this expansion with the triangle inequality allow to get

$$\begin{split} \left| \omega(\alpha,\varepsilon) \cdot l + (j-j_0) c_m(\alpha,\varepsilon) \right| &\leq \left| \omega(\alpha,\varepsilon) \cdot l + \mu_j^{\infty,m}(\alpha,\varepsilon) - \mu_{j_0}^{\infty,m}(\alpha,\varepsilon) \right| \\ &+ \left(W_{0,\alpha} + |r^{2,m}(\alpha,\varepsilon)| \right) \left| j |j|^{\alpha-1} - j_0 |j_0|^{\alpha-1} \right| \\ &+ \left| r_j^{\infty,m}(\lambda,\varepsilon) | + |r_{j_0}^{\infty,m}(\lambda,\varepsilon) \right| \\ &+ O\left(\frac{1}{|j|^{1-\alpha}} \right) + O\left(\frac{1}{|j_0|^{1-\alpha}} \right). \end{split}$$

From (10.58) and (3.38) we obtain

$$\left(W_{0,\alpha} + |r^{2,m}(\alpha,\varepsilon)|\right) \left|j|j|^{\alpha-1} - j_0|j_0|^{\alpha-1}\right| \leq C|j-j_0|\left(\frac{1}{|j|^{1-\alpha}} + \frac{1}{|j_0|^{1-\alpha}}\right).$$

Thus combining the last two estimates and (10.59) we find for $j \neq j_0 \in \mathbb{S}_0^c$

$$\begin{aligned} \left| \omega(\alpha,\varepsilon) \cdot l + (j-j_0)c_m(\alpha,\varepsilon) \right| &\leq \frac{2\kappa_{m+1}\langle j-j_0\rangle}{\langle l \rangle^{\tau_2}} + C|j-j_0| \left(\frac{1}{|j|^{1-\alpha}} + \frac{1}{|j_0|^{1-\alpha}} \right) \\ &+ C\varepsilon^{1-2a} \left(\frac{1}{|j|^{1-\epsilon-2\overline{\alpha}}} + \frac{1}{|j_0|^{1-\epsilon-2\overline{\alpha}}} \right) \\ &\leq \frac{2\kappa_{m+1}\langle j-j_0\rangle}{\langle l \rangle^{\tau_2}} + C \frac{|j-j_0|}{(\min(|j|,|j_0|))^{1-2\overline{\alpha}-\epsilon}}. \end{aligned}$$
(10.75)

Since $\rho, \kappa_m \in (0, 1)$ then $\kappa_{m+1} \leq \kappa_{m+1}^{\rho}$. Therefore if we assume (we know that $\tau_2 > \tau_1$)

$$(\min(|j|,|j_0|))^{1-2\overline{\alpha}-\epsilon} \ge \frac{1}{2}C\kappa_{m+1}^{-\varrho}\langle l\rangle^{\tau_1},$$

we deduce from (10.75) that

$$\left|\omega(\alpha,\varepsilon)\cdot l+(j-j_0)c_m(\alpha,\varepsilon)\right| \leq \frac{4\kappa_{m+1}^{\varrho}|j-j_0|}{\langle l\rangle^{\tau_1}}.$$

To get the result it suffices to use $\kappa_{m+1} \in [\kappa, 2\kappa]$. This ends the proof of Lemma 10.2.

The last result to prove deals with the uniform transversality during Nash-Moser scheme.

Lemma 10.3. Let $q = q_0$ and ρ_0 as in Proposition 3.3, and consider the function $(\alpha, \varepsilon) \mapsto \omega(\alpha, \varepsilon)$ stated in (10.50). There exist $\varepsilon_0 > 0$ small enough such that for any $\varepsilon \in [0, \varepsilon_0]$ we have the following assertions.

(i) For all $m \in \mathbb{N}$, $(l, j) \in \mathbb{Z}^{d+1} \setminus \{0\}$ such that $|j| \leq C_0 \langle l \rangle$, we have

$$\inf_{\alpha \in (\underline{\alpha}, \overline{\alpha})} \max_{0 \le k \le q} |\partial_{\alpha}^{k} (\omega(\alpha, \varepsilon) \cdot l + jc_{m}(\alpha, \varepsilon))| \ge \frac{\rho_{0} \langle l \rangle}{2}.$$

(ii) For all $m \in \mathbb{N}$, $(l, j) \in \mathbb{Z}^d \times \mathbb{S}_0^c$ such that $|j| \leq C_0 \langle l \rangle$, we have

$$\inf_{\alpha \in (\underline{\alpha}, \overline{\alpha})} \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} \big(\omega(\alpha, \varepsilon) \cdot l + \mu_{j}^{\infty, m}(\alpha, \varepsilon) \big) \right| \geqslant \frac{\rho_{0} \langle l \rangle}{2}$$

(iii) For all $(l, j, j_0) \in \mathbb{Z}^d \times (\mathbb{S}_0^c)^2$ such that $|j - j_0| \leq C_0 \langle l \rangle$, we have

$$\inf_{\alpha \in (\underline{\alpha}, \overline{\alpha})} \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} \left(\omega(\alpha, \varepsilon) \cdot l + \mu_{j}^{\infty, m}(\alpha, \varepsilon) - \mu_{j_{0}}^{\infty, m}(\alpha, \varepsilon) \right) \right| \geqslant \frac{\rho_{0} \langle l \rangle}{2}$$

Proof. (i) Using the triangle inequality combined with (10.50), (10.74), (10.4), Proposition 3.3-(ii) and the fact that $|j| \leq C_0 \langle l \rangle$ we get

$$\begin{aligned} \max_{0 \le k \le q} |\partial_{\alpha}^{k} \left(\omega(\alpha, \varepsilon) \cdot l + jc_{m}(\alpha, \varepsilon) \right)| &\geqslant \max_{0 \le k \le q} |\partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot l + jV_{0}(\alpha) \right)| \\ &- \max_{k \le q} |\partial_{\alpha}^{k} \overline{\mathbf{r}}_{\varepsilon}(\alpha) \cdot l| - C|j|\varepsilon^{1-a(1+q)} \\ &\geqslant \rho_{0} \langle l \rangle - C\varepsilon^{1-a(1+q+q\overline{a})} \langle l \rangle - C \langle l \rangle \varepsilon^{1-a(1+q)}. \end{aligned}$$

It follows that

$$\max_{0 \leq k \leq q} \left| \partial_{\alpha}^{k} \left(\omega(\alpha, \varepsilon) \cdot l + jc_{m}(\alpha, \varepsilon) \right) \right| \geq \left(\rho_{0} - C\varepsilon^{1 - a(1 + q + q\overline{a})} \right) \left\langle l \right\rangle$$
$$\geq \frac{\rho_{0} \left\langle l \right\rangle}{2}$$

provided that

$$a < \frac{1}{1+q(1+\overline{a})}\tag{10.76}$$

and ε is taken small enough. We observe that the condition (10.76) follows from (10.4) and (10.2). (ii) This proof is similar to that of the first point. Indeed, we combine the triangle inequality, (10.50), (10.57), (10.58), (10.59), (10.4) and Proposition 3.3 with $|j| \leq C_0 \langle l \rangle$ in order to get

$$\begin{split} \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} \big(\omega(\alpha, \varepsilon) \cdot l + \mu_{j}^{\infty, m}(\alpha, \varepsilon) \big) \right| &\geqslant \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha) \cdot l + \Omega_{j}(\alpha) \right) \right| \\ &- \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} (\overline{\mathbf{r}}_{\varepsilon}(\alpha) \cdot l + jr^{1, m}(\lambda, \varepsilon) + j|j|^{\alpha - 1} r^{2, m}(\alpha, \varepsilon) \right) \right| \\ &- \max_{0 \leqslant k \leqslant q} \left| \partial_{\alpha}^{k} r_{j}^{\infty, m}(\alpha, \varepsilon) \right) \Big| \\ &\geqslant \rho_{0} \langle l \rangle - C \varepsilon^{1 - a(1 + q + q\overline{a})} \langle l \rangle - C \varepsilon^{1 - a(2 + q)} |j|. \end{split}$$

Therefore

$$\max_{0 \le k \le q} \left| \partial_{\alpha}^{k} \big(\omega(\alpha, \varepsilon) \cdot l + \mu_{j}^{\infty, m}(\alpha, \varepsilon) \big) \right| \ge \frac{\rho_{0} \langle l \rangle}{2}$$

provided that ε is small enough and the condition (10.76) is satisfied. (iii) The case $j = j_0$ is trivial. Arguing as in the preceding point, using (10.57) combined with Proposition 3.3, (10.54), (10.58), (10.59), (10.4) and the fact that $0 < |j - j_0| \leq C_0 \langle l \rangle$ we have

$$\begin{split} \max_{0\leqslant k\leqslant q} \left|\partial_{\alpha}^{k} \left(\omega(\lambda,\varepsilon)\cdot l + \mu_{j}^{\infty,m}(\alpha,\varepsilon) - \mu_{j_{0}}^{\infty,m}(\lambda,\varepsilon)\right)\right| &\geq \max_{0\leqslant k\leqslant q} \left|\partial_{\alpha}^{k} \left(\omega_{\mathrm{Eq}}(\alpha)\cdot l + \Omega_{j}(\alpha) - \Omega_{j_{0}}(\alpha)\right)\right| \\ &- \max_{0\leqslant k\leqslant q} \left|\partial_{\alpha}^{k} \left(\overline{\mathbf{r}}_{\varepsilon}(\alpha)\cdot l + (j-j_{0})r^{1,m}(\alpha,\varepsilon) + r_{j}^{\infty,m}(\alpha,\varepsilon) - r_{j_{0}}^{\infty,m}(\alpha,\varepsilon)\right)\right| \\ &- \max_{0\leqslant k\leqslant q} \left|\partial_{\alpha}^{k} \left[(j\mathbb{W}(j,\alpha) - j_{0}\mathbb{W}(j_{0},\alpha))r^{2,m}(\alpha,\varepsilon)\right]\right| \\ &\geq \rho_{0}\langle l\rangle - C\varepsilon^{1-a(1+q+q\overline{a})}\langle l\rangle - C\varepsilon^{1-a(2+q)}|j-j_{0}| - C\varepsilon^{1-a(1+q+q\overline{a})}|j-j_{0}| \\ &\geq \frac{\rho_{0}\langle l\rangle}{2} \end{split}$$

for ε small enough. Notice that we have used the following inequality (3.38). This ends the proof of Lemma 10.3.

APPENDIX A. APPENDIX

We intend in this section to recall and establish some results related to Gamma function and used before in some proofs. The function $\Gamma: \mathbb{C} \setminus (-\mathbb{N}) \to \mathbb{C}$ refers to the gamma function which is the analytic continuation to the negative half plane of the usual gamma function defined on the positive half-plane $\{\operatorname{Re} z > 0\}$ by the integral representation

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$$

It satisfies some algebraic identities such as the relation

$$\Gamma(z+1) = z \Gamma(z), \quad \forall z \in \mathbb{C} \setminus (-\mathbb{N}).$$
 (A.1)

or the Legendre duplication formula,

$$\Gamma(z)\Gamma(z+\frac{1}{2}) = 2^{1-2z}\sqrt{\pi}\,\Gamma(2z). \tag{A.2}$$

It is known that Γ function does not vanish and admits simple poles at $\{-n, n \in \mathbb{N}\}$ and the reciprocal gamma function $\frac{1}{\Gamma}$ is an entire function. Moreover, the real function $x \in (0, \infty) \mapsto \Gamma(x)$ reaches its absolute minimum at a point $x_0 \in (7/5, 3/2)$, that is,

$$\forall x > 0, \quad \Gamma(x) \ge \Gamma(x_0) > 0.$$

For $x \in \mathbb{R}$, we denote by $(x)_n$ the Pokhhammer's symbol defined by

$$(x)_n = \begin{cases} x(x+1)\dots(x+n-1), & \text{if } n \ge 1, \\ 1, & \text{if } n = 0. \end{cases}$$
(A.3)

From the identity (A.1) we deduce the relations

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)}$$
 and $(x)_n = (-1)^n \frac{\Gamma(1-x)}{\Gamma(1-x-n)}$, (A.4)

provided all the quantities are well-defined. The next result is very important and has been frequently used. It describes some asymptotic behavior of Gamma quotient (also named Wallis quotient) given by,

$$\alpha \in [0,1], j \in [0,\infty), \quad \mathsf{W}(j,\alpha) \triangleq \frac{\Gamma(j+\frac{\alpha}{2})}{\Gamma(j+1-\frac{\alpha}{2})}.$$

Our result reads as follows.

Lemma A.1. For positive large real number j, we have

$$\mathsf{W}(j,\alpha) = \frac{1}{j^{1-\alpha}} + O\bigl(\tfrac{1}{j^{3-\alpha}} \bigr).$$

Moreover, the following estimates hold true.

(i) For any $m, k \in \mathbb{N}$, there exists a constant C(k, m) such that

$$\forall \alpha \in [0,1], \ \forall j \in [1,\infty), \quad |\partial_{\alpha}^k \partial_j^m \mathbb{W}(j,\alpha)| \leqslant C(k,m) j^{\alpha-1-m} \left(1 + \ln^k(j)\right) \\ \leqslant C(\epsilon,k,m) j^{\alpha+\epsilon-1-m}.$$

(ii) For any $\gamma, k \in \mathbb{N}$, there exists a constant C(k, m) such that

$$\forall \alpha \in [0,1], \; \forall \, j \in \mathbb{N}, \quad |\partial_{\alpha}^{k} \Delta_{j}^{\gamma} \mathbb{W}(j,\alpha)| \leqslant C(\gamma,q,\epsilon) \, \langle j \rangle^{\alpha+\epsilon-1-\gamma}.$$

We recall that the difference operator Δ_j was introduced in (7.3).

Proof. Recall Hermite's formula of the Γ function, see formula (1.10) in [73],

$$\forall \operatorname{Re}\{z\} > 0, \quad \forall a \in [0,1], \quad \log \Gamma(z+a) = \left(z+a-\frac{1}{2}\right)\log(z) - z + \frac{1}{2}\log(2\pi) \\ + \int_0^\infty \left(\frac{e^{(1-a)t}}{e^t - 1} + a - \frac{1}{2} - \frac{1}{t}\right)\frac{e^{-zt}}{t}dt.$$
(A.5)

This gives the following expression for Wallis quotient,

$$\forall j \in [1,\infty), \quad \mathsf{W}(j,\alpha) = \frac{1}{j^{1-\alpha}} \exp\left(\mathsf{w}_1(j,\alpha) + \mathsf{w}_2(j,\alpha)\right) \tag{A.6}$$

with

$$\mathbf{w}_{1}(j,\alpha) \triangleq \int_{0}^{1} \mathbf{K}(\alpha,t) \frac{e^{-jt}}{t} dt, \quad \mathbf{w}_{2}(j,\alpha) \triangleq \int_{1}^{\infty} \mathbf{K}(\alpha,t) \frac{e^{-jt}}{t} dt$$

and
$$\mathbf{K}(\alpha,t) \triangleq \frac{e^{(1-\alpha)t} - 1}{e^{t} - 1} e^{\frac{\alpha}{2}t} + \alpha - 1.$$

It is straightforward that, for all $k \ge 0$ there exist C = C(k) such that

$$\forall \alpha \in [0,1], \quad \forall t \in [1,\infty), \quad |\partial_{\alpha}^{k} \mathbf{K}(\alpha,t)| \leqslant C(1+t^{k}).$$

It follows that $w_2(j, \alpha)$ satisfies the estimate

$$\begin{aligned} \forall j \in [1, \infty), \quad \forall \alpha \in [0, 1], \quad |\partial_{\alpha}^{k} \partial_{j}^{m} \mathtt{w}_{2}(j, \alpha)| &\leq \int_{1}^{\infty} |\partial_{\alpha}^{k} \mathtt{K}(\alpha, t)| t^{m-1} e^{-jt} dt \\ &\leq C \int_{1}^{\infty} \left(1 + t^{k+m-1}\right) e^{-jt} dt \\ &\leq C(k, m) e^{-j}. \end{aligned}$$
(A.7)

On the other hand, using Taylor formula we get

$$\forall X > 0, \quad \frac{X^{1-\alpha} - 1}{X - 1} = (1 - \alpha) \int_0^1 (1 - \tau + \tau X)^{-\alpha} d\tau.$$

Thus by the substitution $X = e^t$ we infer

$$\mathbf{K}(\alpha,t) = (1-\alpha) \int_0^1 \left(\mathbf{F}(\alpha,\tau,t) - 1 \right) d\tau \quad \text{with} \quad \mathbf{F}(\alpha,\tau,t) \triangleq \frac{e^{\frac{\alpha}{2}t}}{\left(1 - \tau + \tau e^t\right)^{\alpha}}$$

Applying Taylor expansion of order two in the variable t, combined with

$$\mathbf{F}(\alpha, \tau, 0) = 1$$
 and $\partial_t \mathbf{F}(\alpha, \tau, 0) = \alpha \left(\frac{1}{2} - \tau\right)$

lead to

$$\begin{split} \int_{0}^{1} \Big(\mathbf{F}(\alpha,\tau,t) - 1 \Big) d\tau &= \alpha \, t \int_{0}^{1} (\frac{1}{2} - \tau) d\tau + t^{2} \int_{0}^{1} \int_{0}^{1} (\partial_{t}^{2} \mathbf{F})(\alpha,\tau,ts)(1-s) d\tau ds \\ &= t^{2} \int_{0}^{1} \int_{0}^{1} (\partial_{t}^{2} \mathbf{F})(\alpha,\tau,ts)(1-s) d\tau ds. \end{split}$$

Consequently, we obtain

$$\mathsf{K}(\alpha,t) = t^2 \mathsf{G}(\alpha,t) \quad \text{with} \quad \mathsf{G}(\alpha,t) \triangleq (1-\alpha) \int_0^1 \int_0^1 (\partial_t^2 \mathsf{F})(\alpha,\tau,ts)(1-s)d\tau ds.$$

Since F is \mathscr{C}^{∞} in $[0,1]^3$, then G should be \mathscr{C}^{∞} in $[0,1]^2$. Thus, for any $k \ge 0$ there exists C > 0such that

$$\forall \alpha, t \in [0, 1], \quad |\partial_{\alpha}^{k} \mathbf{G}(\alpha, t)| \leqslant C$$

It follows that the expression of $w_1(j, \alpha)$ becomes

$$\mathbf{w}_1(j,\alpha) = \int_0^1 \mathbf{G}(\alpha,t) t e^{-jt} dt \tag{A.8}$$

and one may easily check that

$$\forall j \in [1, \infty), \quad \forall \alpha \in [0, 1], \quad |\partial_{\alpha}^{k} \partial_{j}^{m} \mathbf{w}_{1}(j, \alpha)| \leq \int_{0}^{1} |\partial_{\alpha}^{k} \mathbf{G}(\alpha, t)| t^{m+1} e^{-jt} dt$$

$$\leq C(k, m) j^{-2-m}.$$
(A.9)

Putting together (A.6), (A.7) and (A.9) and using Leibniz rule give the desired asymptotic of $W(j, \alpha)$ with the suitable estimates as stated in (i). The same arguments used to establish (i) combined with the identity (7.6) allow to get (ii). Therefore, the proof of Lemma A.1 is achieved.

The following elementary result has been used before at several points.

Lemma A.2. Let $N_0 \ge 2$. Consider the sequence $(N_m)_{m \in \mathbb{N}}$ defined by (6.32). Then for all $\alpha > 0$, we have

$$\sum_{k=m}^{\infty} N_k^{-\alpha} \underset{m \to \infty}{\sim} N_m^{-\alpha}.$$
225

Proof. Let us consider the decreasing continuous function

$$t \in \mathbb{R}^*_+ \mapsto N_0^{-\alpha(\frac{3}{2})^t} = e^{-\alpha \ln(N_0)e^{t \ln \frac{3}{2}}}.$$

Then applying the series-integral comparison test and making a change of variables yield,

$$\sum_{k=m+1}^{\infty} N_k^{-\alpha} \leqslant \int_m^{\infty} e^{-\alpha \ln(N_0)e^{t \ln \frac{3}{2}}} dt$$
$$\leqslant \int_0^{\infty} e^{-\alpha \ln(N_0)e^{u \ln \frac{3}{2}}e^{m \ln \frac{3}{2}}} du.$$
(A.10)

Now remark that

$$N_m^{\alpha} e^{-\alpha \ln(N_0)e^{u \ln \frac{3}{2}}e^{m \ln \frac{3}{2}}} = \exp\left(\alpha \ln(N_0)\left(1 - e^{u \ln\left(\frac{3}{2}\right)}\right)e^{m \ln\left(\frac{3}{2}\right)}\right).$$

Then using

$$\forall u \ge 0, \quad 1 - e^{u \ln\left(\frac{3}{2}\right)} \le -u \ln\left(\frac{3}{2}\right)$$

we find

$$\forall u \ge 0, \quad N_m^{\alpha} e^{-\alpha \ln(N_0)e^{u \ln \frac{3}{2}}e^{m \ln \frac{3}{2}}} \le \exp\left(-\alpha \ln(N_0)u \ln\left(\frac{3}{2}\right)\left(\frac{3}{2}\right)^m\right)$$

Inserting this estimate into (A.10) and using $N_0 \ge 2$ we get an absolute constant C > 0 such

$$\forall u \in \mathbb{R}^*_+, \forall m \in \mathbb{N}, \quad N_m^{\alpha} e^{-\alpha \ln(N_0)e^{u \ln \frac{3}{2}}e^{m \ln \frac{3}{2}}} \leqslant e^{-C\alpha \left(\frac{3}{2}\right)^m u} \in L^1(\mathbb{R}_+).$$

It follows that

$$\sum_{k=m+1}^{\infty} N_k^{-\alpha} \underset{m \to \infty}{=} o\left(N_m^{-\alpha}\right)$$

As a consequence, we find

$$\sum_{k=m}^{\infty} N_k^{-\alpha} = N_m^{-\alpha} + \sum_{k=m+1}^{\infty} N_k^{-\alpha} \underset{m \to \infty}{\sim} N_m^{-\alpha},$$

which achieves the proof of Lemma A.2.

We shall end this section with recalling the following lemma due to Rüsseman [77, Theorem 17.1] and used in a crucial way to establish (10.61).

Lemma A.3. Let $q \in \mathbb{N}$ $(\alpha, \beta) \in (\mathbb{R}^*_+)^2$ and $f : [a, b] \to \mathbb{R}$ be a function of class \mathscr{C}^q such that

$$\inf_{x \in [a,b]} \max_{0 \le j \le q} |f^{(j)}(x)| \ge \beta.$$

Then there exists C > 0 depending only on b - a, q and $||g||_{\mathscr{C}^k}$ such that

$$\left|\left\{x \in [a,b]; |f(x)| \leq \alpha\right\}\right| \leq C \frac{\alpha^{\frac{1}{q}}}{\beta^{1+\frac{1}{q}}}.$$

ACKNOWLEDGEMENTS:

The work of Z. Hassainia and N. Masmoudi is supported by Tamkeen under the NYU Abu Dhabi Research Institute grant of the center SITE. The work of N. M is supported by NSF grant DMS-1716466.

Z. Hassainia and N. Masmoudi would like to thank M. Berti for many discussions about KAM techniques.

References

- T. Alazard and P. Baldi. Gravity capillary standing water waves. Arch. Ration. Mech. Anal. 217 (2015), no. 3, 741–830.
- [2] W. Ao, J. Davila, M. del Pino, M. Musso, and J. Wei. Travelling and rotating solutions to the generalized inviscid surface quasi-geostrophic equation. Trans. Amer. Math. Soc. 374 (2021), 6665–6689.
- [3] V. Arnold. Small denominators and problems of stability of motion in classical mechanics and celestial mechanics. Uspekhi Mat. Nauk 18 (1963), 91–192.
- [4] P. Baldi, M. Berti, E. Haus and R. Montalto. Time quasi-periodic gravity water waves in finite depth. Invent. Math., 214 (2018) no.2, 739–911.
- [5] P. Baldi and R. Montalto. Quasi-periodic incompresible Euler fows in 3D. Adv. Math. 384 (2021), Paper No. 107730, 74 pp.
- [6] P. Baldi, M. Berti and R. Montalto. KAM for quasi-linear and fully nonlinear forced perturbations of Airy equation. Math. Ann. 359 (2014), no. 1-2, 471–536.
- [7] P. Baldi, M. Berti and R. Montalto. KAM for autonomous quasi-linear perturbations of KdV. Ann. Inst. H. Poincaré Analyse Non. Lin. 33 (2016), no. 6, 1589–1638.
- [8] D. Bambusi, M. Berti and E. Magistrelli. Degenerate KAM theory for partial differential equations. Journal Diff. Equations, 250 (2011), no. 8, 3379–3397.
- M. Berti and P. Bolle, A Nash-Moser approach to KAM theory. Fields Institute Communications, special volume "Hamiltonian PDEs and Applications", 255–284, 2015.
- [10] M. Berti, P. Bolle. Quasi-periodic Solutions of Nonlinear Wave Equations on the d-Dimensional Torus. European Mathematical Society, ISSN 3037192119, 9783037192115, 2020.
- [11] M. Berti, P. Bolle and M. Procesi. An abstract Nash-Moser theorem with parameters and applications to PDEs. Ann. Inst. H. Poincaré Anal. Non Linéaire 27 (2010), no. 1, 377–399.
- [12] M. Berti, L. Corsi and M. Procesi. An Abstract Nash Moser Theorem and Quasi-Periodic Solutions for NLW and NLS on Compact Lie Groups and Homogeneous Manifolds. Comm. Math. Phys. 334, no. 3, 1413–1454, 2015.
- [13] M. Berti, L. Franzoi and A. Maspero. Traveling quasi-periodic water waves with constant vorticity. Archive for Rational Mechanics and Analysis, 240 (2021), 99–202.
- [14] M. Berti, L. Franzoi and A. Maspero. Pure gravity traveling quasi-periodic water waves with constant vorticity. arXiv:2101.12006.
- [15] M. Berti, Z. Hassainia and N. Masmoudi. Time quasi-periodic vortex patches. Preprint.
- [16] M. Berti and R. Montalto. Quasi-periodic standing wave solutions of gravity-capillary water waves. MEMO, Volume 263, 1273, Memoires AMS, ISSN 0065-9266, 2020.
- [17] A. L. Bertozzi and P. Constantin. Global regularity for vortex patches. Comm. Math. Phys., 152 (1993), no. 1, 9–28.
- [18] J. Bourgain, Construction of quasi-periodic solutions for Hamiltonian perturbations of linear equations and applications to nonlinear PDE. Internat. Math. Res. Notices (1994), no. 11, 475ff., approx. 21 pp
- [19] T. Buckmaster, S. Shkoller and V. Vicol. Nonuniqueness of weak solutions to the SQG equation. Comm. Pure Appl. Math. 72 (2019), no. 9, 1809–1874.
- [20] J. Burbea. Motions of vortex patches. Lett. Math. Phys. 6 (1982), no. 1, 1–16.
- [21] D. Cao, G. Qin, W. Zhan, C. Zou. Existence and regularity of co-rotating and travelling global solutions for the generalized SQG equation. arXiv:2103.03992
- [22] A. Castro, D. Córdoba and J. Gómez-Serrano. Existence and regularity of rotating global solutions for the generalized surface quasi-geostrophic equations. Duke Math. J. 165 (2016), no. 5, 935–984.
- [23] A. Castro, D. Córdoba and J. Gómez-Serrano. Uniformly rotating analytic global patch solutions for active scalars. Ann. PDE 2 (2016) no.1, 1–34.
- [24] A. Castro, D. Córdoba and J. Gómez-Serrano. Uniformly rotating smooth solutions for the incompressible 2D Euler equations. Arch. Ration. Mech. Anal. 231 (2019), no. 2, 719–785.
- [25] D. Chae, P. Constantin, D. Córdoba, F. Gancedo, and J. Wu. Generalized surface quasi-geostrophic equations with singular velocities. Comm. Pure Appl. Math., 65 (2012), no. 8, 1037–1066.
- [26] D. Chae, P. Constantin and J. Wu. Inviscid models generalizing the two-dimensional Euler and the surface quasigeostrophic equations. Arch. Ration. Mech. Anal., 202 (2011), no. 1, 35–62.
- [27] J.Y. Chemin. Fluides parfaits incompressibles. Astérisque 230, Société Mathématique de France (1995).
- [28] P. Constantin, A. J. Majda, and E. Tabak. Formation of strong fronts in the 2-D quasigeostrophic thermal active scalar. Nonlinearity, 7 (1994), no. 6, 1495–1533.
- [29] D. Córdoba. Nonexistence of simple hyperbolic blow-up for the quasi-geostrophic equation. Ann. of Math., 148 (1998), no. 2, 1135–1152.
- [30] D. Córdoba and C. Fefferman. Growth of solutions for QG and 2D Euler equations. J. Amer. Math. Soc., 15 (2002), no. 3, 665–670.
- [31] D. Córdoba, M. A. Fontelos, A. M. Mancho and J. L. Rodrigo. Evidence of singularities for a family of contour dynamics equations. Proc. Natl. Acad. Sci. USA 102 (2005), 5949–5952.

- [32] A. Córdoba, D. Córdoba and F. Gancedo. Uniqueness for SQG patch solutions. Trans. Amer. Math. Soc. Ser. B, 5 (2018), 1–31.
- [33] G.S. Deem and N. J. Zabusky, Vortex waves : Stationary "V-states", Interactions, Recurrence, and Breaking. Phys. Rev. Lett. 40 (1978), no. 13, 859–862.
- [34] F. de la Hoz, Z. Hassainia and T. Hmidi. Doubly connected V-states for the generalized surface quasi-geostrophic equations. Arch. Ration. Mech. Anal., 220 (2016), no. 3, 1209–1281.
- [35] F. de la Hoz, Z. Hassainia, T. Hmidi and J. Mateu. An analytical and numerical study of steady patches in the disc. Anal. PDE, 9 (2016), no. 7, 1609–1670.
- [36] F. de la Hoz, T. Hmidi, J. Mateu and J. Verdera. Doubly connected V-states for the planar Euler equations. SIAM J. Math. Anal. 48 (2016), no. 3, 1892–1928.
- [37] D. G. Dritschel, T. Hmidi, C. Renault. Imperfect bifurcation for the quasi-geostrophic shallow-water equations. Arch. Ration. Mech. Anal. 231 (2019), no. 3, 1853–1915
- [38] L.H. Eliasson and S.B. Kuksin. KAM for the nonlinear Schrödinger equation. Ann. Math 172 (2010), 371–435.
- [39] R. Feola, F. Giuliani, R. Montalto, and M. Procesi. Reducibility of first order linear operators on tori via Moser's theorem. J. Funct. Anal., 276 (2019), no.3, 932–970.
- [40] L. E. Fraenkel. An introduction to maximum principles and symmetry in elliptic problems. Number 128. Cambridge University Press, 2000.
- [41] F. Gancedo. Existence for the α-patch model and the QG sharp front in Sobolev spaces. Adv. Math., 217 (2008), no. 6, 2569–2598.
- [42] F. Gancedo and N. Patel. On the local existence and blow-up for generalized SQG patches. Ann. PDE 7 (2021), no. 1, 63 pp.
- [43] C. García. Kármán vortex street in incompressible fluid models. Nonlinearity, 33(4):1625–1676, 2020.
- [44] C. García. Vortex patches choreography for active scalar equations. J Nonlinear Sci 31, 75 (2021).
- [45] C. García, T. Hmidi and J. Soler, Non uniform rotating vortices and periodic orbits for the two-dimensional Euler Equations. Arch. for Ration. Mech. and Anal. 238, 929–1085 (2020).
- [46] L. Godard-Cadillac, P. Gravejat, and D. Smets. Co-rotating vortices with n fold symmetry for the inviscid surface quasi-geostrophic equation. arXiv preprint arXiv:2010.08194, 2020.
- [47] J. Gómez-Serrano. On the existence of stationary patches. Advances in Mathematics 343 (2019), 110–140.
- [48] J. Gómez-Serrano, J. Park, J. Shi and Y. Yao. Symmetry in stationary and uniformly-rotating solutions of active scalar equations. Duke Math. J. 170 (2021), no. 13, 2957–3038.
- [49] P. Gravejat and D. Smets. Smooth travelling-wave solutions to the inviscid surface quasi-geostrophic equation. Int. Math. Res. Not. IMRN 2019, no. 6, 1744–1757
- [50] B. Grébert and T. Kappeler. The defocusing NLS equation and its normal form. EMS Series of Lectures in Mathematics. European Mathematical Society (EMS), Zürich, 2014. x+166 pp. ISBN: 978-3-03719-131-6.
- [51] B. Grébert and E. Paturel. KAM for the Klein Gordon equation on S^d. Boll. Unione Mat. Ital. 9 (2016), no. 2, 237–288.
- [52] Z. Hassainia and T. Hmidi. On the V-States for the generalized quasi-geostrophic equations. Comm. Math. Phys. 337 (2015), no. 1, 321–377.
- [53] Z. Hassainia and T. Hmidi. Steady asymmetric vortex pairs for Euler equations. Discrete Contin. Dyn. Syst., 41(2021), no. 4, 1939–1969.
- [54] Z. Hassainia, N. Masmoudi and M. H. Wheeler. Global bifurcation of rotating vortex patches. Comm. Pure Appl Math., 73 (2020), no. 9, 1933–1980.
- [55] Z. Hassainia and M. H. Wheeler. Multipole vortex patch equilibria for active scalar equations. arXiv:2103.06839.
- [56] S. He and A. Kiselev. Small scale creation for solutions of the SQG equation. Duke Math. J. 170 (2021), no. 5, 1027–1041.
- [57] T. Hmidi. On the trivial solutions for the rotating patch model. J. Evol. Equ., 15 (2015), no. 4, 801–816.
- [58] T. Hmidi and J. Mateu. Bifurcation of rotating patches from Kirchhoff vortices. Discrete Contin. Dyn. Syst. 36 (2016), no. 10, 5401–5422.
- [59] T. Hmidi and J. Mateu. Degenerate bifurcation of the rotating patches. Adv. Math. 302 (2016), 799–850.
- [60] T. Hmidi and J. Mateu. Existence of corotating and counter-rotating vortex pairs for active scalar equations. Comm. Math. Phys. 350 (2017), no. 2, 699–747.
- [61] T. Hmidi, J. Mateu and J. Verdera. Boundary Regularity of Rotating Vortex Patches. Arch. Ration. Mech. Anal. 209 (2013), no. 1, 171–208.
- [62] T. Hmidi and C. Renault. Existence of small loops in a bifurcation diagram near degenerate eigenvalues. Nonlinearity, 30 (2017), no. 10, 3821–3852.
- [63] E. Hölder. Uber die unbeschränkte Fortsetzbarkeit einer stetigen ebenen Bewegung in einer unbegrenzten inkompressiblen Flüssigkeit. Math. Z. 37 (1933), 727–738
- [64] G. Iooss, P. I. Plotnikov and J. F. Toland. Standing waves on an infinitely deep perfect fluid under gravity. Arch. Ration. Mech. Anal., 177(3):367–478, 2005
- [65] G. Kirchhoff, Vorlesungen uber mathematische Physik. (Leipzig, 1874).
- [66] A. Kiselev, Y. Yao and A. Zlatoš. Local regularity for the modified SQG patch equation. Comm. Pure Appl. Math. 70 (2017), no. 7, 1253–1315.

- [67] A. Kiselev, L. Ryzhik, Y. Yao and A. Zlatoš. Finite time singularity for the modified SQG patch equation. Ann. of Math. (2) 184 (2016), no. 3, 909–948.
- [68] A.N. Kolmogorov. On the persistence of conditionally periodic motions under a small change of the hamiltonian function. Doklady Akad. Nauk SSSR 98 (1954), 527–530.
- [69] S. B. Kuksin. Nearly integrable infinite-dimensional Hamiltonian systems. Lecture Notes in Mathematics, vol. 1556, Springer-Verlag, Berlin, 1993.
- [70] S. Kuksin and J. Pöschel. Invariant cantor manifolds of quasi-periodic oscillations for a nonlinear schrödinger equation. Annals of Math 2 (1996), no. 143, 149–179.
- [71] F. Marchand. Existence and regularity of weak solutions to the quasi-geostrophic equations in the spaces L^p or $\dot{H}^{-1/2}$. Comm. Math. Phys., 277(1), (2008), 45–67.
- [72] Moser. On invariant curves of area-preserving mappings of an annulus. Nachr. Akad. Wiss., Göttingen, Math. Phys. Kl. (1962), 1–20.
- [73] G. Nemes. Generalization of Binet's Gamma function formulas. Integral Transforms Spec. Funct. 24 (8), 597–606.
- [74] C. Renault. Relative equilibria with holes for the surface quasi-geostrophic equations. J. Differential Equations, 263 (2017), no. 1567–614.
- [75] S. G. Resnick. Dynamical problems in non-linear advective partial differential equations. PhD thesis, University of Chicago, Department of Mathematics, 1995.
- [76] J. L. Rodrigo, On the evolution of sharp fronts for the quasi-geostrophic equation. Comm. Pure Appl. Math., 58, no.6 (2005), 821–866.
- [77] H. Rüssmann. Invariant tori in non-degenerate nearly integrable Hamiltonian systems. Regul. Chaotic Dyn. 6 (2001), no. 2, 119–204.
- [78] M. V. Ruzhansky and V. Turunen. Pseudo-Differential Operators and Symmetries. Background analysis and advanced topics. Pseudo-Differential Operators. Theory and Applications 2. Birkhäuser Verlag, Basel, 2010.
- [79] M. B. Sevryuk. Invariant tori in quasi-periodic non-autonomous dynamical systems via Herman's method. Discrete Contin. Dyn. Syst. 18 (2007), no. 2–3, 569–595.
- [80] B. Turkington. Corotating steady vortex flows with N-fold symmetry. Nonlinear Anal., 9 (1985), no. 4, 351–369.
- [81] Watson. A Treatise on the Theory of Bessel Functions. Cambridge University Press, (1944).
- [82] C. E. Wayne. Periodic and quasi-periodic solutions of nonlinear wave equations via KAM theory. Communications in Mathematical Physics 127 (1990), no. 3, 479–528.
- [83] W. Wolibner. Un theorème sur l'existence du mouvement plan d'un fluide parfait, homogène, incompressible, pendant un temps infiniment long (French). Mat. Z., 37 (1933), 698–726.
- [84] Y. Yudovich. Nonstationary flow of an ideal incompressible liquid. Zh. Vych. Mat., 3, (1963), 1032–1066.

NYUAD RESEARCH INSTITUTE, NEW YORK UNIVERSITY ABU DHABI, PO BOX 129188, ABU DHABI, UNITED ARAB EMIRATES.

Email address: zh14@nyu.edu.

NYUAD RESEARCH INSTITUTE, NEW YORK UNIVERSITY ABU DHABI, PO BOX 129188, ABU DHABI, UNITED ARAB EMIRATES. UNIV RENNES, CNRS, IRMAR – UMR 6625, F-35000 RENNES, FRANCE *Email address*: thmidi@univ-rennes1.fr.

NYUAD RESEARCH INSTITUTE, NEW YORK UNIVERSITY ABU DHABI, PO BOX 129188, ABU DHABI, UNITED ARAB EMIRATES. COURANT INSTITUTE OF MATHEMATICAL SCIENCES, NEW YORK UNIVERSITY, 251 MERCER STREET, NEW YORK, NY 10012, USA.

 $Email \ address: \tt masmoudi@cims.nyu.edu.$