

# Spectral analysis and long-time asymptotics of the generalized AKNS equation

Hong-Yi Zhang, Yu-Feng Zhang

*School of Mathematics, China University of Mining and Technology, Xuzhou, Jiangsu, 221116, People's Republic of China.*

**Abstract** In this paper, we obtain the long-time asymptotics of the generalized AKNS equation via Defit-Zhou method (Nonlinear steepest descent method). The Cauchy problem of the generalized AKNS equation is transformed into the corresponding Riemann-Hilbert problem on the basis of the Lax pair and the scattering matrix. After that Riemann-Hilbert problems are converted through a decomposition of the matrix-valued spectral function and factorizations of the jump matrix for Riemann-Hilbert problem. Finally, by solving the last model problem, the long-time asymptotics of the generalized AKNS equation are derived.

**Keywords** Riemann-Hilbert problem; Generalized AKNS equation; Nonlinear steepest descent method; Long-time asymptotics

## 1. Introduction

The research of nonlinear partial differential equations (NLPDEs) has played an important role in the development of science and technology. Nowadays, NLPDEs can be used to explain some complex physical phenomena, including mathematics, plasma physics, aerodynamics, fluid mechanics, atmospheric oceans, etc [1-8]. Hirota bilinear method[9,10], Darboux transformation[11,12], the inverse scattering transformation[13] and so on are effective method to solve NLPDEs. Especially, the inverse scattering transformation is the first method found to be used to obtain the exact solution of the soliton equation, and it is called an effective method for analyzing the initial value of the integrable evolution equation on a straight line[14-16]. In 1993, Defit and Zhou proposed the famous nonlinear steepest descent method to analyze the long-time asymptotic behavior of integrable evolution equations[17]. And Defit and Zhou analyzed the long-time asymptotic behavior of the solution to the initial value problem of the famous mKdV equation and Schrödinger equation[17,18]. Subsequently, the nonlinear steepest descent method was extended to the study of integrable evolution equations. In recent years, Fan and Geng researched some long-time asymptotic behavior of integrable evolution equations based on the method[19-24].

In this paper, we study the equation derived from the Lax pair given by Yishen Li[25]. The Lax pair is

$$\begin{aligned}\psi_x &= -iz\sigma_3\psi + P\psi, \\ \psi_t &= (\vartheta z^3 + \iota z^2 + \kappa z + \varsigma)\sigma_3\psi + Q\psi,\end{aligned}\tag{1.1}$$

---

<sup>†</sup>Corresponding author: Y.F. Zhang. E-mail: zhangyfcumt@163.com

where  $\psi(z; x, t)$  is a  $2 \times 2$  matrix,  $\sigma_3 = \text{diag}(1, -1)$ , and

$$P = \begin{pmatrix} 0 & u \\ v & 0 \end{pmatrix},$$

$$Q = i\vartheta z^2 P - iz \begin{pmatrix} \frac{i\vartheta}{2} uv & -\frac{i\vartheta}{2} u_x - \iota u \\ \frac{i\vartheta}{2} v_x - \iota v & -\frac{i\vartheta}{2} uv \end{pmatrix} - \begin{pmatrix} \frac{i\vartheta}{4} (uv_x - vu_x) - \frac{\iota}{2} uv & -\frac{i\vartheta}{4} (-u_{xx} + 2u^2v) + \frac{\iota}{2} u_x - i\kappa u \\ -\frac{i\vartheta}{4} (-v_{xx} + 2uv^2) - \frac{\iota}{2} v_x - i\kappa v & -\frac{i\vartheta}{4} (uv_x - vu_x) + \frac{\iota}{2} uv \end{pmatrix}. \quad (1.2)$$

The Lax pair (1.1) derive the following equation:

$$\begin{cases} u_t = -\frac{i\vartheta}{4} (u_{xxx} - 6uvu_x) - \frac{\iota}{2} (u_{xx} - 2u^2v) + i\kappa u_x + 2\varsigma u, \\ v_t = -\frac{i\vartheta}{4} (v_{xxx} - 6uvv_x) + \frac{\iota}{2} (v_{xx} - 2v^2u) + i\kappa v_x - 2\varsigma v. \end{cases} \quad (1.3)$$

(I) Take  $\vartheta = -4i$ ,  $\iota = \kappa = \varsigma = 0$ ,  $v = -1$ , Eq.(1.3) reduces to KdV equation

$$u_t + 6uu_x + u_{xxx} = 0. \quad (1.4)$$

(II) Take  $\vartheta = -4i$ ,  $\iota = \kappa = \varsigma = 0$ ,  $v = -u$ , Eq.(1.3) reduces to generalized KdV equation

$$u_t + 6u^2u_x + u_{xxx} = 0. \quad (1.5)$$

(III) Take  $\iota = -2i$ ,  $\vartheta = \kappa = \varsigma = 0$ ,  $v = \mp \bar{u}$ , Eq.(1.3) reduces to nonlinear Schrödinger equation

$$iu_t + u_{xx} \pm 2u^2\bar{u} = 0, \quad (1.6)$$

where superscript bar denotes complex conjugate.

(IV) Take  $\iota = -2$ ,  $\vartheta = \kappa = \varsigma = 0$ ,  $q_x = uv = (\frac{u_x}{u})_x$ , Eq.(1.3) reduces to Burger equation

$$q_t = 2qq_x - q_{xx}. \quad (1.7)$$

Especially, take  $\vartheta = -i\alpha$  ( $\alpha > 0$ ),  $\iota = \kappa = \varsigma = 0$  and  $v = \bar{u}$ , Eq.(1.3) reduces to the generalized AKNS equation

$$u_t = \frac{\alpha}{4} (-u_{xxx} + 2(|u|^2u)_x), \quad (1.8)$$

where  $u(x, t)$  is complex-valued function of variate  $(x, t)$ .

In this paper, we use Defit-Zhou nonlinear steepest descent method to study long-time asymptotics of Eq.(1.8) with Schwartz decaying initial data

$$u(t = 0, x) = u_0(x) \in \mathcal{S}(\mathbb{R}).$$

We know that long-time asymptotics of Eq.(1.8) has not been researched, the main result is presented in the following theorem.

**Theorem 1.1**  $u(x, t)$  be the solution for the Cauchy problem of the generalized AKNS equation (1.8) with  $u_0 \in \mathcal{S}(\mathbb{R})$ , where  $\mathcal{S}(\mathbb{R})$  represents Schwartz space:  $\mathcal{S}(\mathbb{R}) = \{f \in C^\infty(\mathbb{R}), \|f\|_{\alpha, \beta} = \sup |x^\alpha \partial^\beta f(x)| < \infty, \alpha, \beta \in \mathbb{Z}_+\}$ . For  $x < 0$  and  $\frac{x}{t} \leq C$ , the leading asymptotics of  $u(x, t)$  has the form:

$$u(x, t) = \frac{ve^{\frac{\pi v}{2}}}{\sqrt{6\alpha t z_0 \pi}} ((\delta_B^0)^2 e^{-\frac{3\pi i}{4}} \Gamma(iv) \overline{r(z_0)} - (\delta_B^0)^{-2} e^{\frac{3\pi i}{4}} \Gamma(-iv) r(z_0)) + o(c(z_0)t^{-1} \log t) \quad (1.9)$$

where  $C$  is a constant,  $\Gamma$  is the Gamma function,  $c$  is a rapidly decreasing function, and

$$\begin{aligned}\delta_B^0 &= (12\alpha t(z_0)^3)^{-\frac{iv}{2}} e^{2i\alpha t z_0^3} e^{\chi(z_0)}, \\ z_0 &= \sqrt{-\frac{x}{3\alpha t}}, \quad v = -\frac{1}{2\pi} \log(1 - |r(z_0)|^2), \\ \chi(z_0) &= \frac{1}{2\pi i} \int_{-z_0}^{z_0} \log\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z}.\end{aligned}$$

The outline of this paper is as follows. In section 2, we analyze eigenfunction and spectral function of Eq.(1.8) to construct a original Riemann-Hilbert problem. In section 3, by deforming the jump matrix of the original Riemann-Hilbert problem and extending region, the original Riemann-Hilbert problem is transformed into a model Riemann-Hilbert problem. Then the solution of the model Riemann-Hilbert problem can be expressed by the solution of Weber equation. Finally, we obtain the long-time asymptotics of the Cauchy problem for the generalized AKNS equation.

## 2. The Riemann-Hilbert Problems

In this section, we first make use of the Lax pair of the generalized AKNS equation to construct the matrix Jost solutions. Then the Cauchy problem of the generalized AKNS equation turns into Riemann-Hilbert problem.

We make a transformation

$$\mu(z; x, t) = \psi(z; x, t) e^{izx\sigma_3 + i\alpha z^3 t \sigma_3}, \quad (2.1)$$

Eq.(2.1) satisfy the following Lax pair

$$\begin{aligned}\mu_x &= iz[\mu, \sigma_3] + P\mu, \\ \mu_t &= i\alpha z^3[\mu, \sigma_3] + Q\mu,\end{aligned} \quad (2.2)$$

where  $e^{\sigma_3} = \text{diag}(e, e^{-1})$ ,  $[\mu, \sigma_3] = \mu\sigma_3 - \sigma_3\mu$ . Lax pair (2.2) can be written as following total differential form

$$d(e^{(izx+i\alpha z^3 t)\hat{\sigma}_3} \mu) = e^{(izx+i\alpha z^3 t)\hat{\sigma}_3} [(Pdx + Qdt)\mu], \quad (2.3)$$

where  $e^{\hat{\sigma}_3} u = e^{\sigma_3} u e^{-\sigma_3}$ . To analyze the eigenfunction  $\mu(z; x, t)$ , we choose two special integral paths

$$(-\infty, t) \rightarrow (x, t), \quad \text{and} \quad (\infty, t) \rightarrow (x, t), \quad (2.4)$$

and acquire two eigenfunctions of Lax pair (2.2)

$$\begin{aligned}\mu_1(z; x, t) &= I + \int_{-\infty}^x e^{-iz(x-\xi)\hat{\sigma}_3} P(z; \xi, t) \mu_1(z; \xi, t) d\xi, \\ \mu_2(z; x, t) &= I - \int_x^{\infty} e^{-iz(x-\xi)\hat{\sigma}_3} P(z; \xi, t) \mu_2(z; \xi, t) d\xi.\end{aligned} \quad (2.5)$$

$\mu_1(z; x, t)$  and  $\mu_2(z; x, t)$  have the following asymptotics

$$\begin{aligned}\mu_1(z; x, t), \mu_2(z; x, t) &\rightarrow I, \quad x \rightarrow \pm\infty, \\ \mu_1(z; x, t), \mu_2(z; x, t) &\rightarrow I, \quad z \rightarrow \infty.\end{aligned} \quad (2.6)$$

Lax pair (1.1) is first order linear homogeneous equations,  $\psi_1(z; x, t) = \mu_1(z; x, t)e^{-i\theta(z)\hat{\sigma}_3}$  and  $\psi_2(z; x, t) = \mu_2(z; x, t)e^{-i\theta(z)\hat{\sigma}_3}$  are solution of Lax pair (1.1), then  $\psi_1(z; x, t)$  and  $\psi_2(z; x, t)$  are linearly correlated, thus we have

$$\mu_1(z; x, t) = \mu_2(z; x, t)e^{-i\theta(z)\hat{\sigma}_3}S(z), \quad (2.7)$$

where

$$\theta(z) = zx + \alpha z^3 t, \quad S(z) = \begin{pmatrix} s_{11}(z) & s_{12}(z) \\ s_{21}(z) & s_{22}(z) \end{pmatrix},$$

$S(z)$  is irrelevant to  $x, t$  and it is called as the spectral matrix function. Nextly, we study the analyticity of  $\mu_1(z; x, t)$ ,  $\mu_2(z; x, t)$  and  $S(z)$ . For integral equation (2.5), when  $\xi < x$ , we have

$$e^{-iz(x-\xi)\hat{\sigma}_3}P(z; \xi, t) = \begin{pmatrix} 0 & ue^{-2iz(x-\xi)} \\ \bar{u}e^{2iz(x-\xi)} & 0 \end{pmatrix}, \quad (2.8)$$

$$e^{2iz(x-\xi)} = e^{2i(x-\xi)Re z}e^{-2(x-\xi)Im z}, \quad e^{-2iz(x-\xi)} = e^{-2i(x-\xi)Re z}e^{2(x-\xi)Im z}.$$

By direct calculation, first column of  $\mu_1(z; x, t)$  is analytical in the upper half plane  $\mathbb{C}_+$ , second column of  $\mu_1(z; x, t)$  is analytical in the lower half plane  $\mathbb{C}_-$ ,  $\mu_1(z; x, t)$  can be written as

$$\mu_1 = \begin{pmatrix} \mu_1^{(11)} & \mu_1^{(12)} \\ \mu_1^{(21)} & \mu_1^{(22)} \end{pmatrix} = (\mu_1^+, \mu_1^-). \quad (2.9)$$

In the same way, first column of  $\mu_2(z; x, t)$  is analytical in the lower half plane  $\mathbb{C}_-$  and second column of  $\mu_2(z; x, t)$  is analytical in the upper half plane  $\mathbb{C}_+$ ,  $\mu_2(z; x, t)$  can be written as

$$\mu_2 = \begin{pmatrix} \mu_2^{(11)} & \mu_2^{(12)} \\ \mu_2^{(21)} & \mu_2^{(22)} \end{pmatrix} = (\mu_2^-, \mu_2^+). \quad (2.10)$$

In virtue of  $\text{tr}(P - iz\sigma_3) = 0$  and  $\text{tr}(Q - i\alpha z^3\sigma_3) = 0$ , by using Abel formula, we get

$$\begin{aligned} (\det\psi_j)_x &= \text{tr}(P - iz\sigma_3)\det\psi_j = 0, \\ (\det\psi_j)_t &= \text{tr}(Q - i\alpha z^3\sigma_3)\det\psi_j = 0, \end{aligned} \quad (2.11)$$

then through transformation (2.1), we have

$$\det\mu_j = \det\psi_j \det(e^{i\theta(z)\sigma_3}) = \det\psi_j, \quad (2.12)$$

therefore, we obtain

$$(\det\mu_j)_x = (\det\mu_j)_t = 0. \quad (2.13)$$

Which imply that  $\det\mu_j$  is independent of  $x$  and  $t$ , and  $\mu_j \rightarrow I$ ,  $|x| \rightarrow \infty$ , so we have

$$\det\mu_j = \lim_{|x| \rightarrow \infty} \det\mu_j = \det\left(\lim_{|x| \rightarrow \infty} \mu_j\right) = 1, \quad (2.14)$$

taking determinants on both sides of Eq.(2.7), yield

$$\det S(z) = 1. \quad (2.15)$$

From Eq.(2.14), we know that  $\mu_1(z; x, t)$ ,  $\mu_2(z; x, t)$  are invertible matrix. Based on analyticity of the column vector of  $\mu_1(z; x, t)$ ,  $\mu_2(z; x, t)$ , it can be inferred that first row and second row of  $\mu_1^{-1}(z; x, t)$  is analytical in the lower half plane  $\mathbb{C}_-$  and the upper half plane  $\mathbb{C}_+$  respectively; first row and second row of  $\mu_2^{-1}(z; x, t)$  is analytical in the upper half plane  $\mathbb{C}_+$  and the lower half plane  $\mathbb{C}_-$  respectively. That is

$$\mu_1^{-1} = \begin{pmatrix} \mu_1^{(22)} & -\mu_1^{(12)} \\ -\mu_1^{(21)} & \mu_1^{(11)} \end{pmatrix} = \begin{pmatrix} \hat{\mu}_1^- \\ \hat{\mu}_1^+ \end{pmatrix}, \quad (2.16)$$

$$\mu_2^{-1} = \begin{pmatrix} \mu_2^{(22)} & -\mu_2^{(12)} \\ -\mu_2^{(21)} & \mu_2^{(11)} \end{pmatrix} = \begin{pmatrix} \hat{\mu}_2^+ \\ \hat{\mu}_2^- \end{pmatrix}, \quad (2.17)$$

Through Eq.(2.7), Eq.(2.16) and Eq.(2.17), we get

$$e^{-i\theta(z)\hat{\sigma}_3} S(z) = \mu_2^{-1} \mu_1 = \begin{pmatrix} \hat{\mu}_2^+ \\ \hat{\mu}_2^- \end{pmatrix} (\mu_1^+, \mu_1^-) = \begin{pmatrix} \hat{\mu}_2^+ \mu_1^+ & \hat{\mu}_2^+ \mu_1^- \\ \hat{\mu}_2^- \mu_1^+ & \hat{\mu}_2^- \mu_1^- \end{pmatrix}. \quad (2.18)$$

From Eq.(2.18), we know that  $s_{11}(z)$  is analytical in the upper half plane  $\mathbb{C}_+$ ,  $s_{22}(z)$  is analytical in the lower half plane  $\mathbb{C}_-$ ,  $s_{12}(z)$  and  $s_{21}(z)$  are continuous to the real axis, but not analytical in the upper and lower half plane.

**Theorem 2.1** The eigenfunctions  $\mu_1(z; x, t)$ ,  $\mu_2(z; x, t)$  and spectral matrix function  $S(z)$  have the following symmetry property

$$\begin{aligned} \mu_j(z; x, t) &= \sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1, \quad (j = 1, 2) \\ S(z) &= \sigma_1 \overline{S(\bar{z})} \sigma_1, \end{aligned} \quad (2.19)$$

where  $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ .

**Proof.** From Lax pair (2.2), we have

$$\mu_{j,x}(z; x, t) + iz[\sigma_3, \mu_j(z; x, t)] = P\mu_j(z; x, t), \quad (2.20)$$

replace  $z$  with  $\bar{z}$ , then take the conjugate of Eq.(2.20), get

$$\overline{\mu_{j,x}(\bar{z}; x, t) - iz[\sigma_3, \mu_j(\bar{z}; x, t)]} = \overline{P\mu_j(\bar{z}; x, t)}. \quad (2.21)$$

Multiply the left and right sides of Eq.(2.21) by  $\sigma_1$ , notice that

$$\sigma_1 \sigma_3 \sigma_1 = -\sigma_3, \quad \sigma_1 \overline{P} \sigma_1 = P, \quad (2.22)$$

and

$$[\sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1]_x - iz \sigma_1 \sigma_3 \overline{\mu_j(\bar{z}; x, t)} \sigma_1 + iz \sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_3 \sigma_1 = \sigma_1 \overline{P\mu_j(\bar{z}; x, t)} \sigma_1. \quad (2.23)$$

Therefore we obtain

$$[\sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1]_x + iz[\sigma_3, \sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1] = P \sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1. \quad (2.24)$$

By comparing with Eq.(2.2) and Eq.(2.24), we know that  $\mu_j(z; x, t)$  and  $\sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1$  satisfy the same differential equation and have the same asymptotic property:  $\mu_j(z; x, t)$ ,  $\sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1 \rightarrow I$ ,  $x \rightarrow \infty$ . Consequently,

$$\mu_j(z; x, t) = \sigma_1 \overline{\mu_j(\bar{z}; x, t)} \sigma_1, \quad (2.25)$$

that is

$$\begin{pmatrix} \mu_{11}(z) & \mu_{12}(z) \\ \mu_{21}(z) & \mu_{22}(z) \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \overline{\mu_{11}(\bar{z})} & \overline{\mu_{12}(\bar{z})} \\ \overline{\mu_{21}(\bar{z})} & \overline{\mu_{22}(\bar{z})} \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} \overline{\mu_{22}(\bar{z})} & \overline{\mu_{21}(\bar{z})} \\ \overline{\mu_{12}(\bar{z})} & \overline{\mu_{11}(\bar{z})} \end{pmatrix}. \quad (2.26)$$

By comparing the two sides of the above equation, we have  $\mu_{11}(z) = \overline{\mu_{22}(\bar{z})}$ ,  $\mu_{12}(z) = \overline{\mu_{21}(\bar{z})}$ . Nextly, we analyze the symmetry property of  $S(z)$ . By deforming Eq.(2.7), we obtain

$$S(z) = e^{(izx+i\alpha z^3 t)\sigma_3} (\mu_2^{-1}(z; x, t) \mu_1(z; x, t)), \quad (2.27)$$

we already know that  $\mu_j(z; x, t) = \overline{\sigma_1 \mu_j(\bar{z}; x, t) \sigma_1}$ , then

$$\begin{aligned} \sigma_1 \overline{S(\bar{z})} \sigma_1 &= \sigma_1 e^{(-izx-i\alpha z^3 t)\sigma_3} \overline{\mu_2^{-1}(\bar{z}) \mu_1(\bar{z})} \sigma_1 \\ &= \sigma_1 e^{(-izx-i\alpha z^3 t)\sigma_3} \sigma_1 \sigma_1 \overline{\mu_2^{-1}(\bar{z})} \sigma_1 \sigma_1 \overline{\mu_1(\bar{z})} \sigma_1 \sigma_1 e^{(izx+i\alpha z^3 t)\sigma_3} \sigma_1 \\ &= e^{(izx+i\alpha z^3 t)\sigma_3} \mu_2^{-1}(z) \mu_1(z) e^{(-izx-i\alpha z^3 t)\sigma_3} \\ &= S(z). \end{aligned} \quad (2.28)$$

So we get  $\overline{s_{11}(\bar{z})} = s_{22}(z)$  and  $\overline{s_{12}(\bar{z})} = s_{21}(z)$ .

**Theorem 2.2** The eigenfunction  $\mu_1(z; x, t)$ ,  $\mu_2(z; x, t)$  and spectral matrix function  $S(z)$  also have the following symmetry property

$$\begin{aligned} \sigma_3 \mu_j^H(\bar{z}; x, t) \sigma_3 &= \mu_j^{-1}(z; x, t), \\ \sigma_3 S^H(\bar{z}) \sigma_3 &= S^{-1}(z). \end{aligned} \quad (2.29)$$

where superscript H denotes conjugate transpose and  $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ .

**Proof:** By direct calculation

$$\begin{aligned} &\sigma_3 \mu_j^H(\bar{z}; x, t) \sigma_3 \\ &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \mu_{j,11}(\bar{z}, x, t) & \mu_{j,12}(\bar{z}, x, t) \\ \mu_{j,21}(\bar{z}, x, t) & \mu_{j,22}(\bar{z}, x, t) \end{pmatrix}^H \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \overline{\mu_{j,11}(\bar{z}, x, t)} & \overline{\mu_{j,21}(\bar{z}, x, t)} \\ \overline{\mu_{j,12}(\bar{z}, x, t)} & \overline{\mu_{j,22}(\bar{z}, x, t)} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} \mu_{j,22}(z) & \mu_{j,12}(z) \\ \mu_{j,21}(z) & \mu_{j,11}(z) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \\ &= \begin{pmatrix} \mu_{j,22}(z) & -\mu_{j,12}(z) \\ -\mu_{j,21}(z) & \mu_{j,11}(z) \end{pmatrix} \\ &= \mu_j^{-1}(z; x, t). \end{aligned} \quad (2.30)$$

Therefore, we have  $\overline{\mu_{11}(\bar{z})} = \mu_{22}(z)$ ,  $\overline{\mu_{12}(\bar{z})} = \mu_{21}(z)$ . In addition,

$$\begin{aligned}
& \sigma_3 S^H(\bar{z}) \sigma_3 \\
&= \sigma_3 (e^{(i\bar{z}x + i\alpha\bar{z}^3 t)\sigma_3} \mu_2^{-1}(\bar{z}) \mu_1(\bar{z}) e^{(-i\bar{z}x - i\alpha\bar{z}^3 t)\sigma_3})^H \sigma_3 \\
&= \sigma_3 (e^{(-izx - i\alpha z^3 t)\sigma_3} \overline{\mu_2^{-1}(\bar{z})} \overline{\mu_1(\bar{z})} e^{(izx + i\alpha z^3 t)\sigma_3})^T \sigma_3 \\
&= \sigma_3 e^{(izx + i\alpha z^3 t)\sigma_3} \overline{\mu_1(\bar{z})}^T \overline{\mu_2^{-1}(\bar{z})}^T e^{(-izx - i\alpha z^3 t)\sigma_3} \sigma_3 \\
&= \sigma_3 e^{(izx + i\alpha z^3 t)\sigma_3} \sigma_3 \sigma_3 \mu_1^H(\bar{z}) \sigma_3 \sigma_3 (\mu_2^{-1}(\bar{z}))^H \sigma_3 \sigma_3 e^{(-izx - i\alpha z^3 t)\sigma_3} \\
&= e^{(izx + i\alpha z^3 t)\sigma_3} \mu_1^{-1}(z) \mu_2(z) e^{(-izx - i\alpha z^3 t)\sigma_3} \\
&= S^{-1}(z),
\end{aligned} \tag{2.31}$$

So we get  $\overline{s_{11}(\bar{z})} = s_{22}(z)$  and  $\overline{s_{12}(\bar{z})} = s_{21}(z)$ . Otherwise,

$$S(z) = e^{i(zx + \alpha z^3 t)\hat{\sigma}_3} (\mu_2^{-1}(z; x, t) \mu_1(z; x, t)), \tag{2.32}$$

and as  $t = 0$ ,  $x \rightarrow +\infty$ ,  $\mu_2^{-1}(z; x, t) \rightarrow I$ , so we have

$$\begin{aligned}
S(z) &= \lim_{x \rightarrow \infty} e^{izx\hat{\sigma}_3} \mu_1(z; x, 0) \\
&= \lim_{x \rightarrow \infty} e^{izx\hat{\sigma}_3} \left( I + \int_{-\infty}^x e^{-iz(x-\xi)\hat{\sigma}_3} P(z; \xi, 0) \mu_1(z; \xi, 0) d\xi \right) \\
&= I + \int_{-\infty}^{\infty} e^{iz\xi\hat{\sigma}_3} P(z; \xi, 0) \mu_1(z; \xi, 0) d\xi.
\end{aligned} \tag{2.33}$$

By direct calculation, we have

$$\begin{aligned}
S(z) &= \begin{pmatrix} s_{11}(z) & s_{12}(z) \\ s_{21}(z) & s_{22}(z) \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \int_{-\infty}^{+\infty} \begin{pmatrix} e^{iz\xi} & 0 \\ 0 & e^{-iz\xi} \end{pmatrix} \begin{pmatrix} 0 & u \\ \bar{u} & 0 \end{pmatrix} \begin{pmatrix} \mu_{1,11}(z, x, t) & \mu_{1,12}(z, x, t) \\ \mu_{1,21}(z, x, t) & \mu_{1,22}(z, x, t) \end{pmatrix} \begin{pmatrix} e^{-iz\xi} & 0 \\ 0 & e^{iz\xi} \end{pmatrix} d\xi \\
&= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \int_{-\infty}^{+\infty} \begin{pmatrix} u\mu_{1,21} & ue^{2iz\xi}\mu_{1,22} \\ \bar{u}e^{-2iz\xi}\mu_{1,11} & \bar{u}\mu_{1,12} \end{pmatrix} d\xi,
\end{aligned} \tag{2.34}$$

comparing the left and right sides of the above equation, we obtain  $s_{11}(z) = 1 + \int_{-\infty}^{+\infty} u\mu_{1,21}d\xi$ ,  $s_{21}(z) = \int_{-\infty}^{+\infty} \bar{u}e^{-2iz\xi}\mu_{1,11}d\xi$ . Define a piecewise-analytic function  $m(z; x, t)$  as

$$m(z; x, t) = \begin{cases} (\frac{\mu_1^+}{s_{11}}, \mu_2^+), & \text{Im}z > 0, \\ (\mu_2^-, \frac{\mu_1^-}{s_{22}}), & \text{Im}z < 0. \end{cases} \tag{2.35}$$

**Theorem 2.3** The piecewise-analytic function  $m(z; x, t)$  defined by Eq.(2.35) satisfies the Riemann-Hilbert problem

- $m_{\pm}(z; x, t)$  is analytical in  $\mathbb{C}_{\pm}$ ,
  - $m_+(z; x, t) = m_-(z; x, t)v(z; x, t)$ ,
  - $m_{\pm}(z; x, t) \rightarrow I$ , as  $z \rightarrow \infty$ .
- $$\tag{2.36}$$

$m_+(z; x, t)$  and  $m_-(z; x, t)$  denote the limiting values as  $z$  approaches the contour  $\mathbb{R}$  from the left and the right along the contour respectively, the oriented contour is on  $\mathbb{R}$  in Fig. 1, where

$$v(z; x, t) = \begin{pmatrix} 1 - |r(z)|^2 & -e^{-2it\theta(z)}\overline{r(\overline{z})} \\ e^{2it\theta(z)}r(z) & 1 \end{pmatrix}. \quad (2.37)$$

**Proof.** From Eq.(2.7), we have

$$\begin{aligned} [\mu_1^+, \mu_1^-] &= [\mu_2^-, \mu_2^+] \begin{pmatrix} e^{-it\theta(z)} & 0 \\ 0 & e^{it\theta(z)} \end{pmatrix} \begin{pmatrix} s_{11}(z) & s_{12}(z) \\ s_{21}(z) & s_{22}(z) \end{pmatrix} \begin{pmatrix} e^{it\theta(z)} & 0 \\ 0 & e^{-it\theta(z)} \end{pmatrix} \\ &= [s_{11}(z)\mu_2^- + s_{21}e^{2it\theta(z)}\mu_2^+, s_{12}e^{-2it\theta(z)}\mu_2^- + s_{22}(z)\mu_2^+], \end{aligned} \quad (2.38)$$

comparing the left and right sides of Eq.(2.38), we get

$$\begin{aligned} \mu_1^+ &= s_{11}(z)\mu_2^- + s_{21}(z)e^{2it\theta(z)}\mu_2^+, \\ \mu_1^- &= s_{12}(z)e^{-2it\theta(z)}\mu_2^- + s_{22}(z)\mu_2^+, \end{aligned} \quad (2.39)$$

deforming the above equation, we have

$$\begin{aligned} \frac{\mu_1^+}{s_{11}(z)} &= \frac{1}{s_{11}(z)s_{22}(z)}\mu_2^- + \frac{1}{s_{22}(z)}\frac{s_{21}(z)}{s_{11}(z)}e^{2it\theta(z)}\mu_1^-, \\ \mu_2^+ &= \frac{\mu_1^-}{s_{22}(z)} - \frac{s_{12}(z)}{s_{22}(z)}e^{-2it\theta(z)}\mu_2^-. \end{aligned} \quad (2.40)$$

The above system written in matrix form is equivalent to

$$\left[ \frac{\mu_1^+}{s_{11}(z)}, \mu_2^+ \right] = [\mu_2^-, \frac{\mu_1^-}{s_{22}(z)}] e^{-it\theta(z)\hat{\sigma}_3} \begin{pmatrix} \frac{1}{s_{11}(z)s_{22}(z)} & -\frac{s_{12}(z)}{s_{22}(z)} \\ \frac{s_{21}(z)}{s_{11}(z)} & 1 \end{pmatrix}. \quad (2.41)$$

Let  $r(z) = \frac{s_{21}(z)}{s_{11}(z)}$ , when  $z \in \mathbb{R}$  we have

$$\overline{s_{11}(\overline{z})} = s_{22}(z), \overline{s_{12}(\overline{z})} = s_{21}(z), \quad (2.42)$$

$$-\frac{s_{12}(z)}{s_{22}(z)} = -\frac{\overline{s_{21}(\overline{z})}}{s_{11}(\overline{z})} = -\overline{r(\overline{z})}, \quad (2.43)$$

$$\frac{1}{s_{11}(z)s_{22}(z)} = \frac{s_{11}(z)s_{22}(z) - s_{21}(z)s_{12}(z)}{s_{11}(z)s_{22}(z)} = 1 - \frac{s_{21}(z)}{s_{11}(z)}\frac{s_{12}(z)}{s_{22}(z)} = 1 - r(z)\overline{r(\overline{z})}. \quad (2.44)$$

From Eq.(2.15), we know that

$$\det(S(z)) = s_{11}(z)s_{22}(z) - s_{21}(z)s_{12}(z) = 1, \quad (2.45)$$

divide both sides of Eq.(2.45) by  $s_{11}(z)s_{22}(z)$  and use Eq.(2.42), we obtain

$$r(z)\overline{r(\overline{z})} + \frac{1}{s_{11}(z)s_{11}(\overline{z})} = 1, \quad (2.46)$$

when  $z \in R$ ,

$$|r(z)|^2 + \frac{1}{|s_{11}(z)|^2} = 1, \quad (2.47)$$



so  $|r(z)| < 1$ . By using Eq.(2.42)-(2.44), Eq.(2.41) can be turned into

$$\left[\frac{\mu_1^+}{s_{11}(z)}, \mu_2^+\right] = \left[\mu_2^-, \frac{\mu_1^-}{s_{22}(z)}\right] e^{-it\theta(z)\hat{\sigma}_3} \begin{pmatrix} 1 - r(z)\overline{r(\bar{z})} & -\overline{r(\bar{z})} \\ r(z) & 1 \end{pmatrix}. \quad (2.48)$$

The solution of Riemann-Hilbert problem (2.36) exists and is unique, and the function

$$u(x, t) = \lim_{x \rightarrow \infty} (z\mu)_{12}, \quad (2.49)$$

solves the generalized AKNS equation.




Figure 1. The oriented contour  $\Sigma^{(1)}$ .

### 3. Long-Time Asymptotic Behavior

#### 3.1. Factorization of the jump matrix

Notice that there are two oscillatory terms  $e^{it\theta(z)}$  and  $e^{-it\theta(z)}$  in the jump matrix  $v(z; x, t)$ , take

$$\varphi(z) = i\theta(z) = i\left(\frac{x}{t}z + \alpha z^3\right), \quad (3.1)$$

then

$$\varphi'(z) = i\frac{x}{t} + 3i\alpha z^2, \varphi''(z) = 6i\alpha z \neq 0, \quad (3.2)$$

assume that  $\varphi'(z) = i\frac{x}{t} + 3i\alpha z^2 = 0$ , we get two steady state phase point  $\pm z_0 = \pm\sqrt{-\frac{x}{3\alpha t}}$ . Substituting the expression of  $z_0$  into  $\theta(z)$ , we have

$$\theta(z) = z\frac{x}{t} + \alpha z^3 = z(-3\alpha z_0^2) + \alpha z^3 = \alpha(z^3 - 3z_0^2 z), \quad (3.3)$$

deforming Eq.(3.3), we get

$$\theta(z) = \alpha((z + z_0)^3 - 3z_0(z + z_0)^2 + 2z_0^3), \quad (3.4)$$

and

$$\theta(z) = \alpha((z - z_0)^3 + 3z_0(z - z_0)^2 - 2z_0^3). \quad (3.5)$$

The jump matrix  $v(z; x, t)$  has the lower/upper triangular factorization

$$v(z; x, t) = \begin{pmatrix} 1 & -\overline{r(\bar{z})}e^{-2it\theta(z)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(z)e^{2it\theta(z)} & 1 \end{pmatrix}, \quad (z \rightarrow \infty) \quad (3.6)$$

and the upper/diagonal/lower factorization

$$v(z; x, t) = \begin{pmatrix} 1 & 0 \\ e^{2it\theta(z)}\frac{r(z)}{1-r(z)\overline{r(\bar{z})}} & 1 \end{pmatrix} \begin{pmatrix} 1 - r(z)\overline{r(\bar{z})} & 0 \\ 0 & \frac{1}{1-r(z)\overline{r(\bar{z})}} \end{pmatrix} \begin{pmatrix} 1 & -e^{-2it\theta(z)}\frac{\overline{r(\bar{z})}}{1-r(z)\overline{r(\bar{z})}} \\ 0 & 1 \end{pmatrix}, \quad z \in (-z_0, z_0). \quad (3.7)$$

Now we seek a transformation to eliminate the middle diagonal matrix in  $z \in (-z_0, z_0)$  factorization. Make a transformation

$$m^{(1)} = m\delta^{-\sigma_3}, \quad (3.8)$$

then

$$m_+^{(1)} = m_+\delta_+^{-\sigma_3} = m_-v(z; x, t)\delta_+^{-\sigma_3} = m_-\delta_-^{-\sigma_3}\delta_+^{\sigma_3}v(z; x, t)\delta_+^{-\sigma_3} = m_-^{(1)}\delta_-^{\sigma_3}v(z; x, t)\delta_+^{-\sigma_3} = m_-^{(1)}v^{(1)}(z; x, t), \quad (3.9)$$

so we obtain

$$v^{(1)}(z; x, t) = \delta_-^{\sigma_3}v(z; x, t)\delta_+^{-\sigma_3}. \quad (3.10)$$

For  $z \in (-z_0, z_0)$ ,

$$\begin{aligned} & v^{(1)}(z; x, t) \\ &= \delta_-^{\sigma_3} \begin{pmatrix} 1 & 0 \\ e^{2it\theta(z)} \frac{r(z)}{1-r(z)r(\bar{z})} & 1 \end{pmatrix} \begin{pmatrix} 1-r(z)r(\bar{z}) & 0 \\ 0 & \frac{1}{1-r(z)r(\bar{z})} \end{pmatrix} \begin{pmatrix} 1 & -e^{-2it\theta(z)} \frac{r(\bar{z})}{1-r(z)r(\bar{z})} \\ 0 & 1 \end{pmatrix} \delta_+^{-\sigma_3} \\ &= \begin{pmatrix} 1 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \frac{r(z)}{1-r(z)r(\bar{z})} & 1 \end{pmatrix} \begin{pmatrix} (1-r(z)r(\bar{z}))\delta_- \delta_+^{-1} & 0 \\ 0 & (\frac{1}{1-r(z)r(\bar{z})})\delta_-^{-1} \delta_+ \end{pmatrix} \begin{pmatrix} 1 & -\delta_+^2 e^{-2it\theta(z)} \frac{r(\bar{z})}{1-r(z)r(\bar{z})} \\ 0 & 1 \end{pmatrix}. \end{aligned} \quad (3.11)$$

For  $z \rightarrow \infty$ ,

$$\begin{aligned} & v^{(1)}(z; x, t) \\ &= \delta_-^{\sigma_3} \begin{pmatrix} 1 & -\overline{r(\bar{z})}e^{-2it\theta(z)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ r(z)e^{2it\theta(z)} & 1 \end{pmatrix} \delta_+^{-\sigma_3} \\ &= \begin{pmatrix} 1 & -\delta_-^2 \overline{r(\bar{z})}e^{-2it\theta(z)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \delta_-^{-2} r(z)e^{2it\theta(z)} & 1 \end{pmatrix}. \end{aligned} \quad (3.12)$$

Now we introduce a scale Riemann-Hilbert problem

- $\delta(z)$  is analytical in  $\mathbb{C} \setminus \mathbb{R}$ ,
  - $\delta_+(z) = \delta_-(z)(1 - r(z)r(\bar{z}))$ ,  $z \in (-z_0, z_0)$ ,
  - $\delta_+(z) = \delta_-(z)$ ,  $z \rightarrow \infty$ ,
  - $\delta(z) \rightarrow 1$ ,  $z \rightarrow \infty$ .
- (3.13)

Deforming the second condition of the above Riemann-Hilbert problem, we get

$$(\log \delta(z))_+ - (\log \delta(z))_- = f(z) = \begin{cases} \log(1 - r(z)r(\bar{z})), & z \in (-z_0, z_0) \\ 0, & z \rightarrow \infty \end{cases}. \quad (3.14)$$

According to Plemelj formula, Eq.(3.14) has a unique bounded solution

$$\log \delta(z) = \frac{1}{2\pi i} \int_{-\infty}^{+\infty} \frac{f(\xi)}{\xi - z} d\xi = \frac{1}{2\pi i} \int_{-\infty}^{z_0} \frac{\log(1 - r(\xi)r(\bar{\xi}))}{\xi - z} d\xi, \quad (3.15)$$

we calculate the right end of Eq.(3.15), have

$$\begin{aligned}
& \frac{1}{2\pi i} \int_{-z_0}^{z_0} \frac{\ln(1-|r(z)|^2)}{\xi-z} d\xi \\
&= \frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z} + \frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln(1-|r(z_0)|^2) \frac{d\xi}{\xi-z} \\
&= \frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z} + \frac{1}{2\pi i} \ln(1-|r(z_0)|^2) \int_{-z_0}^{z_0} \frac{d\xi}{\xi-z} \\
&= \frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z} + \frac{1}{2\pi i} \ln(1-|r(z_0)|^2) \ln\left(\frac{z-z_0}{z+z_0}\right),
\end{aligned} \tag{3.16}$$

therefore,

$$\delta(z) = \left(\frac{z-z_0}{z+z_0}\right)^{\frac{1}{2\pi i} \ln(1-|r(z_0)|^2)} e^{\frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z}}. \tag{3.17}$$

For convenience, we write  $\delta(z)$  in the following form

$$\delta(z) = \left(\frac{z-z_0}{z+z_0}\right)^{iv} e^{\chi(z)}, \tag{3.18}$$

where  $v = -\frac{1}{2\pi} \ln(1-|r(z_0)|^2)$  and  $\chi(z) = \frac{1}{2\pi i} \int_{-z_0}^{z_0} \ln\left(\frac{1-|r(\xi)|^2}{1-|r(z_0)|^2}\right) \frac{d\xi}{\xi-z}$ .

Then based on uniqueness, we can derive

$$\delta(z) = (\delta^\dagger(\bar{z}))^{-1}. \tag{3.19}$$

Substitute Eq.(3.19) into Eq.(3.13), for  $z \in R$  we obtain

$$|\delta_-(z)|^2 = \begin{cases} \frac{1}{1-|r|^2}, & |z| < z_0, \\ 2, & |z| > z_0, \end{cases} \tag{3.20}$$

and

$$|\delta_+(z)|^2 = \begin{cases} 1-|r|^2, & |z| < z_0, \\ 2, & |z| > z_0. \end{cases} \tag{3.21}$$

Therefore, according to the maximum principle,

$$|\delta(z)| \leq c < \infty, \tag{3.22}$$

for all  $z \in \mathbb{C}$ . Similarly, it has a similar process in  $z \in (-\infty, z_0)$ . On the basis of the above analysis,  $m^{(1)}$  satisfy a new Riemann-Hilbert problem

- $m^{(1)}(z; x, t)$  is analytical in  $\mathbb{C} \setminus \Sigma^{(1)}$ ,
  - $m_+^{(1)}(z; x, t) = m_-^{(1)}(z; x, t) v^{(1)}(z; x, t)$ ,  $z \in \Sigma^{(1)}$ ,
  - $m^{(1)}(z; x, t) \rightarrow I$ , as  $z \rightarrow \infty$ ,
- (3.23)

the jump matrix  $v^{(1)}(z; x, t)$  is

$$v^{(1)}(z; x, t) = \begin{cases} \begin{pmatrix} 1 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \frac{r(z)}{1-r(z)r(\bar{z})} & 1 \end{pmatrix} \begin{pmatrix} 1 & -\delta_+^2 e^{-2it\theta(z)} \frac{\overline{r(\bar{z})}}{1-r(z)r(\bar{z})} \\ 0 & 1 \end{pmatrix}, & z \in (-z_0, z_0), \\ \begin{pmatrix} 1 & -\delta_-^2 \overline{r(\bar{z})} e^{-2it\theta(z)} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \delta_-^{-2} r(z) e^{2it\theta(z)} & 1 \end{pmatrix}, & z \rightarrow \infty, \end{cases} \tag{3.24}$$

and we have

$$u(x, t) = 2i \lim_{z \rightarrow \infty} (zm^{(1)}(z; x, t))_{12}. \quad (3.25)$$

For the sake of convenience, we introduce a function

$$\rho(z) = \begin{cases} \overline{r(\overline{z})}, & z \rightarrow \infty, \\ \frac{-\overline{r(\overline{z})}}{1-r(z)r(\overline{z})}, & z \in (-z_0, z_0), \end{cases} \quad (3.26)$$

then the jump matrix  $v^{(1)}(z; x, t)$  are unified as

$$v^{(1)}(z; x, t) = \begin{pmatrix} 1 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \overline{\rho(\overline{z})} & 1 \end{pmatrix}^{-1} \begin{pmatrix} 1 & \delta_+^2 e^{-2it\theta(z)} \rho(z) \\ 0 & 1 \end{pmatrix}. \quad (3.27)$$

### 3.2. Extend to the augmented contour

Let  $L$  denote the contour


$$\begin{aligned} L &: z = z_0 + z_0 \beta e^{\frac{3\pi i}{4}} : -\infty < \beta \leq \sqrt{2} \\ &\cup z = -z_0 + z_0 \beta e^{\frac{\pi i}{4}} : -\infty < \beta \leq \sqrt{2}, \end{aligned} \quad (3.28)$$

and

$$\begin{aligned} L_\varepsilon &: z = z_0 + z_0 \beta e^{\frac{3\pi i}{4}} : \varepsilon < \beta \leq \sqrt{2} \\ &\cup z = -z_0 + z_0 \beta e^{\frac{\pi i}{4}} : \varepsilon < \beta \leq \sqrt{2}, \end{aligned} \quad (3.29)$$

where  $0 < \varepsilon < \sqrt{2}$ .

Through the analysis of  $Re(i\theta(z))$ , we get the symbol distribution map of  $Re(i\theta(z))$  and as shown in Fig.2.



**Fig.2** Symbol distribution map of  $Re(i\theta(z))$ .

**Theorem 3.1** The spectral function  $\rho(z)$  has a decomposition on the real number axis as follows

$$\rho(z) = R(z) + h_1(z) + h_2(z), \quad (3.30)$$

$R(z)$ ,  $h_1(z)$  and  $h_2(z)$  satisfy

$$|e^{-2it\theta(z)} h_1(z)| \lesssim \frac{1}{|z - i|^2 t^l}, \quad z \in \mathbb{R}, \quad (3.31)$$

$$|e^{-2it\theta(z)}h_2(z)| \lesssim \frac{1}{|z-i|^{2l}}, \quad z \in L, \quad (3.32)$$

$$|e^{-2it\theta(z)}R(z)| \lesssim ce^{-6\alpha\varepsilon^2\tau}, \quad z \in L_\varepsilon, \quad (3.33)$$

where  $\theta(z) = \alpha(z - z_0)^3 + 3\alpha z_0(z - z_0)^2 - 2z_0^3$ ,  $R(z)$  is piecewise rational,  $h_2(z)$  has an analytic continuation to  $L$ ,  $l$  is a arbitrary positive integer.

**Proof.** Firstly, we consider spectral function  $\rho(z)$  with  $z \in (z_0, +\infty)$ . Replace the function  $\rho(z)$  by a rational function with well-controlled errors, expand  $(z - i)^{m+5}\rho(z)$  in a Taylor series around  $z_0$

$$\begin{aligned} & (z - i)^{m+5}\rho(z) \\ &= (z_0 - i)^{m+5}\rho(z_0) + \frac{((z - i)^{m+5}\rho(z))^{(1)}}{1} \Big|_{z=z_0}(z - z_0) + \dots + \frac{((z - i)^{m+5}\rho(z))^{(m)}}{m!} \Big|_{z=z_0}(z - z_0)^m \\ & \quad + \frac{((\xi - i)^{m+5}\rho(\xi))^{(m+1)}}{(m+1)!}(z - z_0)^{m+1} \\ &= \mu_0 + \mu_1(z - z_0) + \dots + \mu_m(z - z_0)^m + \frac{1}{m!} \int_{z_0}^z ((\xi - i)^{m+5}\rho(\xi))^{(m+1)}(z - \xi)^m d\xi. \end{aligned} \quad (3.34)$$

Take

$$R(z) = \frac{\sum_{j=0}^m \mu_j (z - z_0)^j}{(z - i)^{m+5}} \quad (3.35)$$

and

$$h(z) = \rho(z) - R(z), \quad (3.36)$$

then

$$h(z) = \frac{((\xi - i)^{m+5}\rho(\xi))^{(m+1)}}{(z - i)^{m+5}m!}(z - z_0)^{m+1}, \quad (3.37)$$

we have

$$\frac{d^j h(z)}{dz^j} \Big|_{z=z_0} = 0, \quad 0 \leq j \leq m, \quad (3.38)$$

so

$$\frac{d^j \rho(z)}{dz^j} \Big|_{z=z_0} = \frac{d^j R(z)}{dz^j} \Big|_{z=z_0}, \quad 0 \leq j \leq m. \quad (3.39)$$

$\theta(z) = \frac{z}{t}x + \alpha z^3$ ,  $z_0 = \sqrt{\frac{-x}{3\alpha t}}$ , we have  $-x = 3\alpha t z_0^2$ , therefore

$$t\theta(z) = zx + \alpha z^3 t = -x(-z + \frac{\alpha t z^3}{-x}) = |x|(-z + \frac{\alpha t z^3}{3\alpha t z_0^2}) = |x|(\frac{z^3 - 3z_0^2 z}{3z_0^2}) = |x|\tilde{\theta}(z), \quad (3.40)$$

then  $Re(i\theta(z))$  and  $Re(i\tilde{\theta}(z))$  differ by a constant multiplier. Next define

$$\gamma(z) = \frac{(z - z_0)^n}{(z - i)^{n+2}}, \quad (3.41)$$

by Fourier inverse transformation, we have

$$\left(\frac{h}{\gamma}\right)(z) = \int_{-\infty}^{+\infty} e^{is\theta(z)} \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s, \quad (3.42)$$

where

$$\left(\frac{\hat{h}}{\gamma}\right)(s) = \int_{z_0}^{+\infty} e^{-is\theta(z)} \left(\frac{h}{\gamma}\right)(z) \bar{d}\theta(z) \quad (3.43)$$

and

$$\bar{d}s = \frac{ds}{\sqrt{2\pi}}, \quad \bar{d}\theta(z) = \frac{d\theta(z)}{\sqrt{2\pi}}. \quad (3.44)$$

Due to

$$h(z) = \frac{1}{(z-i)^{m+5}m!} \int_{z_0}^z ((\xi-i)^{m+5}\rho(\xi))^{(m+1)} (z-\xi)^m d\xi, \quad (3.45)$$

then

$$\left(\frac{h}{\gamma}\right)(z) = \frac{(z-i)^{n+2}(z-z_0)^{-n}}{(z-i)^{m+5}m!} \int_{z_0}^z ((\xi-i)^{m+5}\rho(\xi))^{(m+1)} (z-\xi)^m d\xi. \quad (3.46)$$

Take  $\xi = z_0 + \eta(z - z_0)$ , we have

$$\begin{aligned} \left(\frac{h}{\gamma}\right)(z) &= \frac{(z-z_0)^{-n}}{(z-i)^{m+3-n}m!} \int_0^1 [(z_0 + \eta(z-z_0) - i)^{m+5}\rho(z_0 + \eta(z-z_0))]^{(m+1)} (z-z_0 - \eta(z-z_0))^m d(z_0 + \eta(z-z_0)) \\ &= \frac{(z-z_0)^{m+1-n}}{(z-i)^{m+3-n}m!} \int_0^1 [(z_0 + \eta(z-z_0) - i)^{m+5}\rho(z_0 + \eta(z-z_0))]^{(m+1)} (1-\eta)^m d\eta \\ &= \frac{(z-z_0)^{m+1-n}}{(z-i)^{m+3-n}m!} g(z, z_0), \end{aligned} \quad (3.47)$$

let  $m = 4n + 1$ , then

$$\left(\frac{h}{\gamma}\right)(z) = \frac{(z-z_0)^{3n+2}}{(z-i)^{3n+4}} g(z, z_0), \quad (3.48)$$

where  $g(z, z_0) = \frac{1}{m!} \int_0^1 [(z_0 + \eta(z-z_0) - i)^{m+5}\rho(z_0 + \eta(z-z_0))]^{(m+1)} (1-\eta)^m d\eta$ , due to  $\rho(z) \in \mathcal{S}(R)$ , therefore

$$\left| \frac{d^j g(z, z_0)}{dz^j} \right| \lesssim c, \quad z \geq z_0, \quad (3.49)$$

and due to

$$\left| \frac{z-z_0}{z+z_0} \right| \leq 1, \quad z \geq z_0, \quad (3.50)$$

we have

$$\begin{aligned} &\int_{z_0}^{\infty} \left| \left(\frac{d}{d\theta}\right)^j \left(\frac{h}{\gamma}\right)(z) \right|^2 \bar{d}\theta(z) \\ &= \int_{z_0}^{\infty} \left| \left(\frac{1}{3\alpha(z^2 - z_0^2)}\right)^j \left(\frac{h}{\gamma}\right)(z) \right|^2 3\alpha(z^2 - z_0^2) \bar{d}z \\ &\leq c \int_{z_0}^{\infty} \left| \frac{(z-z_0)^{3n+2-3j}}{(z-i)^{3n+4}} \right|^2 (z^2 - z_0^2) \bar{d}z \leq c, \quad (0 < z_0 < M, 0 \leq j \leq \frac{3n+2}{3}). \end{aligned} \quad (3.51)$$

By using Plancherel theorem, we have

$$\int_{-\infty}^{+\infty} (1+s^2)^j \left| \frac{\hat{h}}{\gamma}(s) \right|^2 ds \leq c \leq \infty, \quad (0 < z_0 < M, 0 \leq j \leq \frac{3n+2}{3}). \quad (3.52)$$

Next we split  $h(z)$

$$\begin{aligned} h(z) &= \gamma(z) \int_{-\infty}^{+\infty} e^{is\theta(z)} \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s \\ &= \gamma(z) \int_t^{+\infty} e^{is\theta(z)} \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s + \gamma(z) \int_{-\infty}^t e^{is\theta(z)} \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s \\ &= h_1(z) + h_2(z). \end{aligned} \quad (3.53)$$

According to Eq.(3.41), we have

$$|e^{-2it\theta(z)}h_1(z)| = |e^{-2i|x|\tilde{\theta}(z)}h_1(z)|, \quad (3.54)$$

when  $z \in \mathbb{R}$ ,  $z \geq z_0$ , we have  $Re(i\tilde{\theta}(z)) = 0$ , then  $-2i|x|\tilde{\theta}(z)$  is pure imaginary, so

$$\begin{aligned} & |e^{-2i|x|\tilde{\theta}(z)}h_1(z)| \\ & \leq |\gamma(z)| \int_t^{+\infty} |(\frac{\hat{h}}{\gamma})(s)| \bar{d}s \\ & = |\gamma(z)| \int_t^{+\infty} (1+s^2)^{-\frac{p}{2}}(1+s^2)^{\frac{p}{2}} |(\frac{\hat{h}}{\gamma})(s)| \bar{d}s \\ & \leq |\gamma(z)| \left( \int_t^{+\infty} (1+s^2)^{-p} ds \right)^{\frac{1}{2}} \left( \int_t^{+\infty} (1+s^2)^p |(\frac{\hat{h}}{\gamma})(s)|^2 \bar{d}s \right)^{\frac{1}{2}} \\ & \lesssim |\gamma(z)| \left( \int_t^{+\infty} s^{-2p} ds \right)^{\frac{1}{2}} \\ & \lesssim \frac{1}{|z-i|^2} t^l. \end{aligned} \quad (3.55)$$

For  $|e^{-2i|x|\tilde{\theta}(z)}h_2(z)|$ , on ray  $z = z_0 + z_0\eta e^{-\frac{i\pi}{4}}$ , we have

$$\begin{aligned} & |e^{-2i|x|\tilde{\theta}(z)}h_2(z)| \\ & \leq |e^{-2it\theta(z)}| \cdot \left| \frac{(z-z_0)^n}{(z-i)^{n+2}} \right| \cdot \left| \int_{-\infty}^t e^{is\theta(z)} (\frac{\hat{h}}{\gamma})(s) \bar{d}s \right| \\ & \leq c \left| \frac{(z-z_0)^n}{(z-i)^{n+2}} \right| \cdot |e^{-tRe(i\theta(z))}| \cdot \left| \int_{-\infty}^t e^{i(s-t)\theta(z)} (\frac{\hat{h}}{\gamma})(s) \bar{d}s \right| \\ & \leq c \left| \frac{(z-z_0)^n}{(z-i)^{n+2}} \right| \cdot |e^{-tRe(i\theta(z))}| \cdot \left( \int_{-\infty}^t (1+s^2)^{\frac{1}{2}} \right) \cdot \left( \int_{-\infty}^t ((1+s^2)^2 |(\frac{\hat{h}}{\gamma})(s)|^2 ds)^{\frac{1}{2}} \right) \\ & \leq \frac{z_0^n u^n}{|z-i|^{n+2}} \cdot e^{-tRe(i\theta(z))} \\ & \leq \frac{cz_0^n [(tz_0^3)^{\frac{1}{3}}u]^n \cdot e^{-\frac{\sqrt{2}}{2}\alpha z_0^3 u^3 t}}{|z-i|^{n+2} \cdot (tz_0^3)^{\frac{n}{3}}} \\ & \leq \frac{c}{|z-i|^2 t^{\frac{n}{3}}} \leq \frac{c}{|z-i|^2 t^l}, \end{aligned} \quad (3.56)$$

$$|e^{-2it\theta(z)}[\rho(z)]| \leq ce^{-2tRe(i\theta(z))} \leq ce^{-6\alpha t z_0^3 u^2} \leq ce^{-6\alpha\tau}, \quad (3.57)$$

where  $\tau = tz_0^3 = \left(\frac{-x}{3\alpha t^{\frac{1}{3}}}\right)^{\frac{3}{2}}$ .

For  $\overline{\rho(\bar{z})}$ , take Schwartz conjugate to  $\rho(z)$ , we have

$$\overline{\rho(\bar{z})} = \overline{R(\bar{z})} + \overline{h_1(\bar{z})} + \overline{h_2(\bar{z})}, \quad (3.58)$$

There are similar estimates for  $e^{2it\theta(z)}\overline{h_1(\bar{z})}$ ,  $e^{2it\theta(z)}\overline{h_2(\bar{z})}$  and  $e^{2it\theta(z)}\overline{R(\bar{z})}$  on  $\mathbb{R} \cup \bar{L}$ . Based on the result of theorem 3.1 and Beal-Cofman theorem,  $v^{(1)}(z; x, t)$  has the following decomposition

$$v^{(1)}(z; x, t) = (b_-)^{-1}b_+ = (b_-^a)^{-1}(b_-^o)^{-1}b_+^o b_+^a, \quad (3.59)$$

$b_{\pm}$  has decomposition

$$\begin{aligned} b_+ &= b_+^o b_+^a = (I + \omega_+^o)(I + \omega_+^a), \\ b_- &= b_-^o b_-^a = (I - \omega_-^o)(I - \omega_-^a), \end{aligned} \quad (3.60)$$

where

$$\begin{aligned} \omega_+^o &= \begin{pmatrix} 0 & \delta_+^2(z)e^{-2it\theta(z)}h_1(z) \\ 0 & 0 \end{pmatrix}, \quad \omega_+^a = \begin{pmatrix} 0 & \delta_+^2(z)e^{-2it\theta(z)}[\rho(z)]h_2(z) \\ 0 & 0 \end{pmatrix}, \\ \omega_-^o &= \begin{pmatrix} 0 & 0 \\ \delta_-^2(z)e^{2it\theta(z)}\overline{h_1(\bar{z})} & 0 \end{pmatrix}, \quad \omega_-^a = \begin{pmatrix} 0 & 0 \\ \delta_-^2(z)e^{2it\theta(z)}[\overline{\rho(\bar{z})}]\overline{h_2(\bar{z})} & 0 \end{pmatrix}. \end{aligned}$$

Define the oriented contour  $\Sigma^{(2)}$  as  $\Sigma^{(2)} = \mathbb{R} \cup L \cup \bar{L}$  as in Fig. 3, then to extend the jump matrix to the steep descend line  $L$  and  $\bar{L}$ .  $(b_-^a)^{-1}$  can be analytically extend to  $\bar{L}$ ,  $b_+^a$  can be analytically extend to  $L$ ,  $(b_-^o)^{-1}b_+^o$  do not have the property of analytic continuation but decays fast about time on  $\mathbb{R}$ . We make a transformation

$$m^{(2)} = m^{(1)}\phi, \quad (3.61)$$

where

$$\phi = \begin{cases} I, & z \in \Omega_1 \cup \Omega_2, \\ (b_-^a)^{-1}, & z \in \Omega_3 \cup \Omega_4 \cup \Omega_5, \\ (b_+^a)^{-1}, & z \in \Omega_6 \cup \Omega_7 \cup \Omega_8, \end{cases} \quad (3.62)$$

then Riemann-Hilbert problem in  $\Sigma^{(1)}$  turn into a new Riemann-Hilbert problem in  $\Sigma^{(2)}$ , and  $m^{(2)}$  satisfy

- $m^{(2)}(x, t, x)$  is analytic in  $\mathbb{C} \setminus \Sigma^{(2)}$ ,
  - $m_+^{(2)}(x, t, x) = m_-^{(2)}(x, t, x)v^{(2)}(x, t, x)$ ,  $z \in \Sigma^{(2)}$ ,
  - $m_+^{(2)}(x, t, x) \rightarrow I$ , as  $z \rightarrow \infty$ ,
- (3.63)

where the jump matrix is

$$v^{(2)}(x, t, z) = \begin{cases} (b_-^o)^{-1}b_+^o, & z \in \mathbb{R}, \\ b_+^a, & z \in \bar{L}, \\ (b_-^a), & z \in L. \end{cases} \quad (3.64)$$

The solution of the Eq.(1.8) can be presented as


$$u(x, t) = 2i \lim_{z \rightarrow \infty} (zm^{(2)}(z; x, t))_{12}. \quad (3.65)$$

Firstly, we study the case of  $z \in \Omega_6$ , considering the bound of  $\delta(z)$  in Eq.(3.19), the definition of  $h_2(z)$  and  $[\rho(z)]$  in (3.32) and (3.33), have

$$\begin{aligned} |\delta^2(z)e^{-2it\theta(z)}h_2(z)| &\leq c|\gamma(z)| \cdot |e^{-it\text{Re}(i\theta(z))}| \cdot \left| \int_{-\infty}^t e^{i(s-t)\theta(z)} \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s \right| \\ &\leq \frac{|z - z_0|^n}{|z - i|^{n+2}} \cdot \int_{-\infty}^t \left| \left(\frac{\hat{h}}{\gamma}\right)(s) \bar{d}s \right| \leq \frac{c}{|z - i|^2}, \end{aligned} \quad (3.66)$$

$$|\delta^2 e^{-2it\theta(z)}[\rho(z)]| \leq \frac{c|\sum_{j=0}^k \mu_j(z - z_0)^j|}{|z - i|^{k+5}} \leq \frac{c}{|z - i|^5}. \quad (3.67)$$





**Figure 3.** The oriented contour  $\Sigma^{(2)}$ .

Set

$$v^{(2)}(x, t, z) = (b_-^{(2)})^{-1} b_+^{(2)}, \quad (3.68)$$

define

$$\omega_{\pm}^{(2)} = \pm(b_{\pm}^{(2)} - I), \quad \omega^{(2)} = \omega_+^{(2)} + \omega_-^{(2)}, \quad (3.69)$$

and Cauchy operator

$$(C_{\pm}f)(z) = \frac{1}{2\pi i} \int_{\Sigma^{(2)}} \frac{f(\xi)}{\xi - z_{\pm}} d\xi, \quad z \in \Sigma^{(2)}, \quad f \in \ell^2(\Sigma^{(2)}). \quad (3.70)$$

We know that Cauchy operator  $C_{\pm}$  is a bounded operator from  $\ell^2(\Sigma^{(2)})$  to  $\ell^2(\Sigma^{(2)})$  and  $C_+ + C_- = 1$ .

Define

$$C_{\omega^{(2)}} = C_+(f\omega_-^{(2)}) + C_-(f\omega_+^{(2)}), \quad (3.71)$$

let  $\mu^{(2)} = \mu^{(2)}(z; x, t) \in \ell^2(\Sigma^{(2)}) + \ell^{\infty}(\Sigma^{(2)})$  be the solution of the following basic inverse equation

$$\mu^{(2)} = I + C_{\omega^{(2)}}\mu^{(2)}, \quad (3.72)$$

then

$$m^{(2)}(z; x, t) = I + \frac{1}{2\pi i} \int_{\Sigma^{(2)}} \frac{\mu^{(2)}(\xi; x, t)\omega^{(2)}(\xi; x, t)}{\xi - z} d\xi \quad (3.73)$$

is the solution of the Riemann-Hilbert problem (3.63). So the solution of Eq.(1.8) is

$$\begin{aligned} u(x, t) &= 2i \lim_{z \rightarrow \infty} (zm^{(2)}(z; x, t))_{12} \\ &= 2i \lim_{z \rightarrow \infty} \left( z \left( I + \frac{1}{2\pi i} \int_{\Sigma^{(2)}} \frac{\mu^{(2)}(\xi; x, t)\omega^{(2)}(\xi; x, t)}{\xi - z} d\xi \right) \right)_{12} \\ &= \lim_{z \rightarrow \infty} z \left( \int_{\Sigma^{(2)}} \frac{\mu^{(2)}\omega^{(2)} d\xi}{\xi - z} \frac{1}{\pi} \right)_{12} \\ &= - \left( \int_{\Sigma^{(2)}} \frac{\mu^{(2)}\omega^{(2)}}{\pi} d\xi \right)_{12} \\ &= - \left( \int_{\Sigma^{(2)}} ((I - C_{\omega^{(2)}})^{-1})(\xi)\omega^{(2)} \frac{d\xi}{\pi} \right)_{12}. \end{aligned} \quad (3.74)$$

### 3.3. The third transformation.

In this subsection, the Riemann-Hilbert problem on the contour  $\Sigma^{(2)}$  is converted to Riemann-Hilbert problem on the contour  $\Sigma^{(3)} = \Sigma^{(2)} \setminus (\mathbb{R} \cup L_{\varepsilon} \cup \overline{L_{\varepsilon}})$  and we figure out the estimates of the errors between the two Riemann-Hilbert problems. Let  $\omega^e = \omega^a + \omega^b + \omega^c$ ,  $\omega^a = \omega^2|_{\mathbb{R}}$  is supported on  $\mathbb{R}$  and is composed of terms of

type  $h_1(z)$  and  $\overline{h_1(\overline{z})}$ ,  $\omega^b$  is supported on  $L \cup \overline{L}$  and is composed of the contribution to  $\omega^2$  from terms of type  $h_2(z)$  and  $\overline{h_2(\overline{z})}$ ,  $\omega^c$  is supported on  $L_\varepsilon \cup \overline{L_\varepsilon}$  and is composed of the contribution to  $\omega^2$  from terms of type  $R(z)$  and  $\overline{R(\overline{z})}$ . That is

$$\omega^a = \begin{cases} \omega^{(2)}, & z \in \mathbb{R}, \\ 0, & \text{otherwise,} \end{cases}$$

$$\omega^b = \begin{cases} \begin{pmatrix} 0 & \delta_+^2 e^{-2it\theta(z)} h_2(z) \\ 0 & 0 \end{pmatrix}, & z \in L, \\ \begin{pmatrix} 0 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \overline{h_2(\overline{z})} & 0 \end{pmatrix}, & z \in \overline{L}, \\ 0, & \text{otherwise} \end{cases} \quad (3.75)$$

$$\omega^c = \begin{cases} \begin{pmatrix} 0 & \delta_+^2 e^{-2it\theta(z)} R(z) \\ 0 & 0 \end{pmatrix}, & z \in L_\varepsilon, \\ \begin{pmatrix} 0 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \overline{R(\overline{z})} & 0 \end{pmatrix}, & z \in \overline{L_\varepsilon}, \\ 0, & \text{otherwise.} \end{cases}$$

Define  $\omega^{(3)}$  as

$$\omega^{(2)} = \omega^{(3)} + \omega^{(e)}, \quad (3.76)$$

notice that  $\omega^{(3)} = 0$  on  $\Sigma^{(2)} \setminus \Sigma^{(3)}$ , then we get the following estimates.

**Theorem 3.2** For arbitrary positive integer  $l$ , as  $t \rightarrow \infty$ , we have the estimates

$$\|\omega^a\|_{\ell^\infty(\mathbb{R}) \cap \ell^2(\mathbb{R}) \cap \ell^1(\mathbb{R})} \leq ct^{-l}, \quad (3.77)$$

$$\|\omega^b\|_{\ell^\infty(L \cup \overline{L}) \cap \ell^2(L \cup \overline{L}) \cap \ell^1(L \cup \overline{L})} \leq ct^{-l}, \quad (3.78)$$

$$\|\omega^{(3)}\|_{\ell^2(\Sigma^{(2)})} \leq \tau^{-\frac{1}{4}}, \quad \|\omega^{(3)}\|_{\ell^1(\Sigma^{(2)})} \leq \tau^{-\frac{1}{2}}, \quad (3.79)$$

**Proof.** Firstly, we consider the case of  $\omega^a$ . It is necessary to compute  $|\omega^a|$  if we want to give the estimates for  $\|\omega^a\|_{\ell^p(\mathbb{R})}$ . Notice that the bound of  $\delta(z)$  in (3.19), we have

$$\begin{aligned} |\omega^a| &= (|e^{-2it\theta(z)} \delta^2(z) h_1(z)|^2 + |e^{2it\theta(z)} \delta^{-2}(z) \overline{h_1(\overline{z})}|^2)^{\frac{1}{2}} \\ &\leq |e^{-2it\theta(z)} \delta^2(z) h_1(z)| + |e^{2it\theta(z)} \delta^{-2}(z) \overline{h_1(\overline{z})}| \\ &\lesssim |e^{-2it\theta(z)} h_1(z)| + |e^{2it\theta(z)} \overline{h_1(\overline{z})}| \\ &\lesssim \frac{1}{(|z|^2 - 1)t^l}. \end{aligned} \quad (3.80)$$

Similarly, we can prove that (3.77) also holds by a simple calculation. Next, we prove estimate (3.78), note the definition of (3.35), we have


$$R(z) = \frac{|\sum_{j=0}^m \mu_j (z - z_0)^j|}{|(z - i)^{m+5}|} \lesssim \frac{1}{1 + |z|^5}, \quad (3.81)$$

$Re(i\theta(z)) \geq 3\alpha z_0 u^2$  on  $L : z = z_0 + z_0 \beta e^{\frac{3\pi i}{4}} (-\infty < \beta \leq \infty)$ , we find that

$$|e^{-2it\theta(z)} \delta^2(z) R(z)| \lesssim e^{-6\alpha \varepsilon^2 t z_0^3} \frac{1}{1 + |z|^5}. \quad (3.82)$$

From Proposition 2.23 and Corollary 2.25 in [17], we know that operator  $(1 - C_{\omega(3)})^{-1} : \ell^2(\Sigma^{(2)}) \rightarrow \ell^2(\Sigma^{(2)})$  exists and is uniformly bounded and

$$\|(1 - C_{\omega(3)})^{-1}\|_{\ell^2(\Sigma^{(2)})} \lesssim 1, \quad \|(1 - C_{\omega(2)})^{-1}\|_{\ell^2(\Sigma^{(2)})} \lesssim 1, \quad (3.83)$$



**Figure 4.** The oriented contour  $\Sigma^{(3)}$ .

**Theorem 3.3.** As  $t \rightarrow \infty$ ,

$$\int_{\Sigma^{(2)}} ((1 - C_{\omega(2)})^{-1} I)(\xi) \omega^{(2)}(\xi) d\xi = \int_{\Sigma^{(2)}} ((1 - C_{\omega(3)})^{-1} I)(\xi) \omega^{(3)}(\xi) d\xi + o(t^{-l}). \quad (3.84)$$

**Proof.** It is easy to see that

$$(1 - C_{\omega(2)})^{-1} - (1 - C_{\omega(3)})^{-1} = (1 - C_{\omega(3)})^{-1} (C_{\omega(2)} - C_{\omega(3)}) (1 - C_{\omega(2)})^{-1}, \quad (3.85)$$

then

$$\begin{aligned} & (1 - C_{\omega(2)})^{-1} I \omega^{(2)} - (1 - C_{\omega(3)})^{-1} I \omega^{(3)} \\ &= [(1 - C_{\omega(2)})^{-1} - (1 - C_{\omega(3)})^{-1}] I \omega^{(2)} + (1 - C_{\omega(3)})^{-1} I (\omega^{(2)} - \omega^{(3)}) \\ &= (1 - C_{\omega(3)})^{-1} (C_{\omega(2)} - C_{\omega(3)}) (1 - C_{\omega(2)})^{-1} I \omega^{(2)} + (1 - C_{\omega(3)})^{-1} I \omega^e \\ &= (1 - C_{\omega(3)})^{-1} C_{\omega^e} (I + (1 - C_{\omega(3)})^{-1} C_{\omega(2)}) I \omega^{(2)} + (I + (1 - C_{\omega(3)})^{-1} C_{\omega(3)}) I \omega^e \\ &= (1 - C_{\omega(3)})^{-1} C_{\omega^e} \omega + (1 - C_{\omega(3)})^{-1} C_{\omega^e} (1 - C_{\omega(2)})^{-1} C_{\omega(2)} I \omega^{(2)} + \omega^e + (1 - C_{\omega(3)})^{-1} C_{\omega(3)} I \omega^e. \end{aligned} \quad (3.86)$$

From theorem 3.1 and Eq(3.83), we have

$$\begin{aligned} \|\omega^e\|_{\ell^1(\Sigma^2)} &\leq \|\omega^a\|_{\ell^1(R)} + \|\omega^b\|_{\ell^1(L \cup \bar{L})} + \|\omega^c\|_{\ell^1(L_\varepsilon \cup \bar{L}_\varepsilon)} \\ &\leq C t^{-l} + C t^{-l} + C z_0 \tau^{-l} \leq C z_0 \tau^{-l}, \quad (t^l \geq \frac{c \tau^l}{z_0}). \end{aligned} \quad (3.87)$$

By direct calculation, we get

$$\begin{aligned} & \|(1 - C_{\omega(3)})^{-1} C_{\omega^e} I \omega\|_{\ell^1(\Sigma^{(2)})} \\ &\leq \|(1 - C_{\omega(3)})^{-1} C_{\omega^e} I\|_{\ell^2(\Sigma^{(2)})} \|\omega^{(2)}\|_{\ell^2(\Sigma^{(2)})} \\ &\leq \|(1 - C_{\omega(3)})^{-1}\|_{\ell^2(\Sigma^{(2)})} \|C_{\omega^e} I\|_{\ell^2(\Sigma^{(2)})} + (\|\omega^a\|_{\ell^2(R)} + \|\omega^b\|_{\ell^1(L \cup \bar{L})} + \|\omega^{(3)}\|_{\ell^2(\Sigma^{(2)})}) \\ &\lesssim \|\omega^e\|_{\ell^2(\Sigma^{(2)})} (t^{-l} + t^{-l} + t^{-\frac{1}{4}}), \end{aligned} \quad (3.88)$$

and

$$\begin{aligned}
& \|(1 - C_{\omega^{(3)}})^{-1} C_{\omega^{(3)}} I \omega^e\|_{\ell^1(\Sigma^{(2)})} \\
& \leq \|(1 - C_{\omega^{(3)}})^{-1} C_{\omega^{(3)}} I\|_{\ell^2(\Sigma^{(2)})} \|\omega^e\|_{\ell^2(\Sigma^{(2)})} \\
& \leq \|(1 - C_{\omega^{(3)}})^{-1}\|_{\ell^2(\Sigma^{(2)})} \|C_{\omega^{(3)}} I\|_{\ell^2(\Sigma^{(2)})} \|\omega^e\|_{\ell^2(\Sigma^{(2)})} \\
& \lesssim \|\omega^{(3)}\|_{\ell^2(\Sigma^{(2)})} (\|\omega^a\|_{\ell^2(R)} + \|\omega^b\|_{\ell^1(L \cup \bar{L})}) \lesssim t^{-l-\frac{1}{4}},
\end{aligned} \tag{3.89}$$

therefore,

$$\begin{aligned}
& \|(1 - C_{\omega^{(3)}})^{-1} C_{\omega^e} (1 - C_{\omega^{(2)}})^{-1} C_{\omega^{(2)}} I \omega^{(2)}\|_{\ell^1(\Sigma^{(2)})} \\
& \leq \|(1 - C_{\omega^{(3)}})^{-1} C_{\omega^e} (1 - C_{\omega^{(2)}})^{-1} C_{\omega^{(2)}} I\|_{\ell^2(\Sigma^{(2)})} \|\omega^{(2)}\|_{\ell^2(\Sigma^{(2)})} \\
& \leq \|(1 - C_{\omega^{(3)}})^{-1}\|_{\ell^2(\Sigma^{(2)})} \|(1 - C_{\omega^{(2)}})^{-1}\|_{\ell^2(\Sigma^{(2)})} \|C_{\omega^e}\|_{\ell^2(\Sigma^{(2)})} \|C_{\omega^{(2)}} I\|_{\ell^2(\Sigma^{(2)})} \|\omega^{(2)}\|_{\ell^2(\Sigma^{(2)})} \\
& \leq \|\omega^e\|_{\ell^\infty(\Sigma^{(2)})} \|\omega^{(2)}\|_{\ell^2(\Sigma^{(2)})}^2 \lesssim t^{-l-\frac{1}{2}}.
\end{aligned} \tag{3.90}$$

For  $z_0 < M$ , substituting the above estimates into (3.86), we obtain theorem 3.3.

**Notice** As  $z \in \Sigma^{(2)} \setminus \Sigma^{(3)}$ ,  $\omega^{(3)}(z) = 0$ , let  $C_{\omega^{(3)}}|_{\ell^2(\Sigma^{(3)})}$  denote the restriction of  $C_{\omega^{(3)}}$  to  $\ell^2(\Sigma^{(3)})$ . For simplicity, we write  $C_{\omega^{(3)}}|_{\ell^2(\Sigma^{(3)})}$  as  $C_{\omega^{(3)}}$ . Then

$$\int_{\Sigma^{(2)}} ((1 - C_{\omega^{(3)}})^{-1} I)(\xi) \omega^{(3)}(\xi) d\xi = \int_{\Sigma^{(3)}} ((1 - C_{\omega^{(3)}})^{-1} I)(\xi) \omega^{(3)}(\xi) d\xi. \tag{3.91}$$

**Theorem 3.4** As  $t \rightarrow \infty$ , the solution  $u(x, t)$  for the Cauchy problem of Eq.(1.8) has the asymptotic estimate

$$u(x, t) = -\frac{1}{\pi} \left( \int_{\Sigma^{(3)}} ((1 - C_{\omega^{(3)}})^{-1})(\xi) \omega^{(3)}(\xi) d\xi \right)_{12} + o(t^{-l}). \tag{3.92}$$

**Proof.** A direct consequence of (3.74) and (3.84).

Set

$$m^{(3)} = I + \int_{\Sigma^{(3)}} \frac{\mu^{(3)} \omega^{(3)}(\xi, x, t)}{\xi - z} \frac{d\xi}{2\pi i}, \tag{3.93}$$

where  $\mu^{(3)} = (1 - C_{\omega^{(3)}})^{-1} I$ , the solution (3.92) is equivalent to

$$u(x, t) = 2i \lim_{z \rightarrow \infty} (zm^{(3)})_{12} + o(t^{-l}). \tag{3.94}$$

Note that  $m^{(2)}(z, x, t)$  is the solution of the Riemann-Hilbert problem (3.62), we can construct the following Riemann-Hilbert problem

$$\begin{cases} m_+^{(3)}(z, x, t) = m_-^{(3)}(z, x, t) v^{(3)}(z, x, t), & z \in \Sigma^{(3)} \\ m^{(3)}(z, x, t) \rightarrow I, & z \rightarrow \infty, \end{cases} \tag{3.95}$$

where

$$\begin{aligned}
v^{(3)} &= (b_-^{(3)})^{-1} b_+^{(3)} = (I - \omega_-^{(3)})^{-1} (I + \omega_+^{(3)}), \\
\omega^{(3)} &= \omega_+^{(3)} + \omega_-^{(3)}, \\
\omega_+^{(3)} &= \begin{cases} \begin{pmatrix} 0 & \delta_+^2 e^{-2it\theta(z)} \rho(z) \\ 0 & 0 \end{pmatrix}, & z \in L, \\ 0_{2 \times 2}, & z \in \bar{L}, \end{cases}
\end{aligned}$$

$$\omega_-^{(3)} = \begin{cases} 0_{2 \times 2}, & z \in L, \\ \begin{pmatrix} 0 & 0 \\ \delta_-^{-2} e^{2it\theta(z)} \overline{\rho(\bar{z})} & 0 \end{pmatrix}, & z \in \bar{L}. \end{cases}$$

### 3.4. Noninteraction of disconnected contour components.

Let the contour  $\Sigma^{(3)} = \Sigma_A^{(3)} \cup \Sigma_B^{(3)}$  and  $\omega_{\pm}^{(3)} = \omega_{A\pm}^{(3)} + \omega_{B\pm}^{(3)}$ , where  $\omega_{A\pm}^{(3)}(z) = 0$  for  $z \in \Sigma_A^{(3)}$ . The operator  $C_{\omega_A^{(3)}}$  and  $C_{\omega_B^{(3)}}$ :  $\ell^2(\Sigma^{(3)}) + \ell^\infty(\Sigma^{(3)}) \rightarrow \ell^2(\Sigma^{(3)})$  are defined in definition (3.71).

**Lemma 3.5**

$$\begin{aligned} \|C_{\omega_B^{(3)}} C_{\omega_A^{(3)}}\|_{\ell^2(\Sigma^{(3)})} &= \|C_{\omega_A^{(3)}} C_{\omega_B^{(3)}}\|_{\ell^2(\Sigma^{(3)})} \lesssim c(z_0) \tau^{-\frac{1}{2}}, \\ \|C_{\omega_B^{(3)}} C_{\omega_A^{(3)}}\|_{\ell^\infty(\Sigma^{(3)}) \rightarrow \ell^2(\Sigma^{(3)})}, \|C_{\omega_A^{(3)}} C_{\omega_B^{(3)}}\|_{\ell^\infty(\Sigma^{(3)}) \rightarrow \ell^2(\Sigma^{(3)})} &\lesssim c(z_0) \tau^{-\frac{3}{4}}. \end{aligned}$$

**Proof.** See Lemma 3.5 in [17].

**Theorem 3.6**

As  $\tau \rightarrow \infty$ , we have

$$\begin{aligned} &\int_{\Sigma^{(3)}} ((1 - C_{\omega^{(3)}})^{-1} I)(\xi) \omega^{(3)}(\xi) d\xi \\ &= \int_{\Sigma_A^{(3)}} (1 - C_{\omega_A^{(3)}})^{-1} I(\xi) \omega_A^{(3)}(\xi) d\xi + \int_{\Sigma_B^{(3)}} (1 - C_{\omega_B^{(3)}})^{-1} I(\xi) \omega_B^{(3)}(\xi) d\xi + o\left(\frac{c(z_0)}{\tau}\right). \end{aligned} \quad (3.96)$$

**Proof.** From identity

$$\begin{aligned} &(1 - C_{\omega_A^{(3)}} - C_{\omega_B^{(3)}})(1 + C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} + C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1}) \\ &= 1 - C_{\omega_B^{(3)}} C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} - C_{\omega_A^{(3)}} C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1}, \end{aligned} \quad (3.97)$$

we have

$$\begin{aligned} (1 - C_{\omega^{(3)}})^{-1} &= 1 + C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1} + [1 + C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1}] \\ &\quad [1 - C_{\omega_B^{(3)}} C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} - C_{\omega_A^{(3)}} C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1}]^{-1} \\ &\quad [C_{\omega_B^{(3)}} C_{\omega_A^{(3)}}(1 - C_{\omega_A^{(3)}})^{-1} + C_{\omega_A^{(3)}} C_{\omega_B^{(3)}}(1 - C_{\omega_B^{(3)}})^{-1}]. \end{aligned} \quad (3.98)$$

By using theorem (3.2), Eq.(3.84) and Lemma 3.5, we obtain theorem (3.6). For convenience, we write the restriction  $C_{\omega_A^{(3)}}|_{\ell^2(\Sigma_A^{(3)})}$  as  $C_{\omega_A^{(3)}}$ , similar for  $C_{\omega_B^{(3)}}$ .

**Lemma 3.7** As  $\tau \rightarrow \infty$ ,

$$u(x, t) = -\frac{1}{\pi} \left( \int_{\Sigma_A^{(3)}} ((1 - C_{\omega_A^{(3)}})^{-1})(\xi) \omega_A^{(3)}(\xi) d\xi \right)_{12} - \frac{1}{\pi} \left( \int_{\Sigma_B^{(3)}} ((1 - C_{\omega_B^{(3)}})^{-1})(\xi) \omega_B^{(3)}(\xi) d\xi \right)_{12} + o\left(\frac{c(z_0)}{\tau}\right). \quad (3.99)$$

### 3.5. Rescaling and further reduction of the RH problems.


Extend the contours  $\Sigma_A^{(3)}$  and  $\Sigma_B^{(3)}$  to the contour

$$\hat{\Sigma}_A^{(3)} : z = -z_0 + z_0 \eta e^{\pm \frac{\pi i}{4}} (\eta \in \mathbb{R}) \text{ and } \hat{\Sigma}_B^{(3)} : z = z_0 + z_0 \eta e^{\pm \frac{3\pi i}{4}} (\eta \in \mathbb{R}),$$

define  $\hat{\omega}_A^{(3)}$  and  $\hat{\omega}_B^{(3)}$  as


$$\hat{\omega}_{A\pm}^{(3)} = \begin{cases} \omega_{A\pm}^{(3)}(z), & z \in \Sigma_A^{(3)}, \\ 0, & z \in \hat{\Sigma}_A^{(3)} \setminus \Sigma_A^{(3)}, \end{cases} \quad (3.100)$$

$$\hat{\omega}_{B\pm}^{(3)} = \begin{cases} \omega_{B\pm}^{(3)}(z), & z \in \Sigma_B^{(3)}, \\ 0, & z \in \hat{\Sigma}_B^{(3)} \setminus \Sigma_B^{(3)}. \end{cases} \quad (3.101)$$



**Figure 4.** The oriented contour  $\hat{\Sigma}^{(3)}$ .

Let  $\Sigma_A$  and  $\Sigma_B$  denote the contour  $z = z_0 \eta e^{\pm \frac{\pi i}{4}}$  ( $\eta \in \mathbb{R}$ ).



**Figure 5.** The oriented contour  $\Sigma$ .

Define the scaling operator as

$$\begin{aligned} N_A &: \ell^2(\hat{\Sigma}_A^{(3)}) \rightarrow \ell^2(\Sigma_A), \\ f(z) &\rightarrow (N_A f)(z) = f\left(\frac{z}{\sqrt{12\alpha t z_0}} - z_0\right), \end{aligned} \quad (3.102)$$

$$\begin{aligned} N_B &: \ell^2(\hat{\Sigma}_B^{(3)}) \rightarrow \ell^2(\Sigma_B), \\ f(z) &\rightarrow (N_B f)(z) = f\left(\frac{z}{\sqrt{12\alpha t z_0}} + z_0\right). \end{aligned} \quad (3.103)$$

Take  $\omega_A = N_A \hat{\omega}_{A\pm}^{(3)}$  and  $\omega_B = N_B \hat{\omega}_{B\pm}^{(3)}$ , the bounded operator  $C_{\omega_A} : \ell^2(\Sigma_A)(\ell^2(\Sigma_B)) \rightarrow \ell^2(\Sigma_A)(\ell^2(\Sigma_B))$ . For a  $2 \times 2$  matrix-valued function  $f$ , a direct calculation shows

$$N C_{\hat{\omega}_A}^{(3)} f = N(C_+(f \hat{\omega}_{A-}^{(3)}) + C_-(f \hat{\omega}_{A+}^{(3)})) = C_{\hat{\omega}_A}^{(3)} N f, \quad (3.104)$$

which means

$$C_{\hat{\omega}_A}^{(3)} = N_A^{-1} C_{\omega_A} N_A. \quad (3.105)$$

Similarly, we get

$$C_{\hat{\omega}_B}^{(3)} = N_B^{-1} C_{\omega_B} N_B. \quad (3.106)$$

From the definition of  $\omega_A$ , it is easy to see that

$$\omega_A = \omega_{A^+} = \begin{pmatrix} 0 & N_A(\delta_+^2(z)e^{-2it\theta(z)}\rho(z)) \\ 0 & 0 \end{pmatrix} \quad (3.107)$$

on  $L_A$ , and

$$\omega_A = \omega_{A^-} = \begin{pmatrix} 0 & 0 \\ N_A(\delta_-^{-2}(z)e^{2it\theta(z)}\overline{\rho(\bar{z})}) & 0 \end{pmatrix} \quad (3.108)$$

on  $\overline{L_A}$ . By using formulae (3.4) and (3.16), we obtain

$$N_A(\delta_A(z)e^{-it\theta(z)}) = \delta_A^0 \delta_A^1(z), \quad (3.109)$$

where

$$\delta_A^0 = e^{-2i\alpha\tau}(12\alpha\tau)^{\frac{iv}{2}} e^{\chi(-z_0)}$$

and

$$\delta_A^1(z) = z^{-iv} \left( \frac{z_0}{\frac{z}{\sqrt{12\alpha\tau z_0}} - 2z_0} \right) e^{-iv} e^{\frac{iz^2}{4}(1-z(108\alpha\tau)^{-\frac{1}{2}})} e^{\chi(\frac{z}{\sqrt{12\alpha\tau z_0}} - z_0) - \chi(-z_0)}.$$

In addition, through formulae (3.5) and (3.16), we get

$$N_B(\delta_B(z)e^{-it\theta(z)}) = \delta_B^0(z)\delta_B^1(z), \quad (3.110)$$

where

$$\delta_B^0 = e^{2i\alpha\tau}(12\alpha\tau)^{-\frac{iv}{2}} e^{\chi(z_0)}$$

and

$$\delta_B^1(z) = z^{iv} \left( \frac{z_0}{\frac{z}{\sqrt{12\alpha\tau z_0}} + 2z_0} \right) e^{iv} e^{-\frac{iz^2}{4}(1+z(108\alpha\tau)^{-\frac{1}{2}})} e^{\chi(\frac{z}{\sqrt{12\alpha\tau z_0}} + z_0) - \chi(z_0)}.$$

Notice that  $\delta_A^0$  and  $\delta_B^0$  is independent of  $z$ , furthermore

$$|\delta_A^0| = 1, \quad |\delta_B^0| = 1. \quad (3.111)$$

Here we construct  $\omega_0$  on the contour  $\Sigma_A$  and  $\Sigma_B$  respectively. Set  $v^{A0} = (I - \omega_{A^0-})^{-1}(I + \omega_{A^0+})$ , where

$$\omega_{A^0} = \omega_{A^0+} = \begin{cases} \begin{pmatrix} 0 & (\delta_A^0)^2(-z)^{-2iv} e^{\frac{iz^2}{2}\overline{r(-z_0)}} \\ 0 & 0 \end{pmatrix}, & z \in \Sigma_A^4, \\ \begin{pmatrix} 0 & (\delta_A^0)^2(-z)^{-2iv} e^{\frac{iz^2}{2}\frac{-\overline{r(-z_0)}}{1-|r(z_0)|^2}} \\ 0 & 0 \end{pmatrix}, & z \in \Sigma_A^2, \end{cases} \quad (3.112)$$

$$\omega_{A^0} = \omega_{A^0-} = \begin{cases} \begin{pmatrix} 0 & 0 \\ (\delta_A^0)^{-2}(-z)^{2iv} e^{-\frac{iz^2}{2}\overline{r(-z_0)}} & 0 \end{pmatrix}, & z \in \Sigma_A^3, \\ \begin{pmatrix} 0 & 0 \\ (\delta_A^0)^{-2}(-z)^{2iv} e^{-\frac{iz^2}{2}\frac{r(z_0)}{1-|r(z_0)|^2}} & 0 \end{pmatrix}, & z \in \Sigma_A^1, \end{cases} \quad (3.113)$$

$$\delta_A^0 = e^{-2i\alpha\tau} (12\alpha\tau)^{\frac{i\nu}{2}} e^{\chi(-z_0)}. \quad (3.114)$$

Due to Lemma 3.35 in [17], we have

$$\|\omega_A - \omega_A^0\|_{\ell^\infty(\Sigma_A) \cap \ell^2(\Sigma_A) \cap \ell^1(\Sigma_A)} \lesssim t^{-\frac{1}{2}} \log t.$$

As  $t \rightarrow \infty$ , we have

$$\begin{aligned} & \int_{\Sigma_A^{(3)}} ((1 - C_{\omega_A^{(3)}})^{-1} I) \omega_A^{(3)}(\xi) d\xi \\ &= \int_{\hat{\Sigma}_A^{(3)}} ((1 - C_{\hat{\omega}_A^{(3)}})^{-1} I) \hat{\omega}_A^{(3)}(\xi) d\xi \\ &= \int_{\hat{\Sigma}_A^{(3)}} N_A^{-1} ((1 - C_{\omega_A})^{-1} N_A I)(\xi) \hat{\omega}_A^{(3)}(\xi) d\xi \\ &= \int_{\hat{\Sigma}_A^{(3)}} ((1 - C_{\omega_A})^{-1} I) ((\xi + z_0) \sqrt{12\alpha t}) N_A \hat{\omega}_A^{(3)}((\xi + z_0) \sqrt{12\alpha t}) d\xi \\ &= \frac{1}{\sqrt{12\alpha t}} \int_{\Sigma_A} ((1 - C_{\omega_A})^{-1} I)(\xi) \omega_A(\xi) d\xi \\ &= \frac{1}{\sqrt{12\alpha t}} \int_{\Sigma_A} ((1 - C_{\omega_A^0})^{-1} I)(\xi) \omega_A^0(\xi) d\xi + o\left(\frac{\log t}{t}\right). \end{aligned} \quad (3.115)$$

Similarly,

$$\begin{aligned} & \int_{\Sigma_B^{(3)}} ((1 - C_{\omega_B^{(3)}})^{-1} I) \omega_B^{(3)}(\xi) d\xi \\ &= \frac{1}{\sqrt{12\alpha t}} \int_{\Sigma_B} ((1 - C_{\omega_B^0})^{-1} I)(\xi) \omega_B^0(\xi) d\xi + o\left(\frac{\log t}{t}\right). \end{aligned} \quad (3.116)$$

On the basis of above analysis, we have

$$\begin{aligned} u(x, t) &= -\left(\frac{1}{\sqrt{12\alpha t}} \int_{\Sigma_A} ((1 - C_{\omega_A^0})^{-1} I)(\xi) \omega_A^0(\xi) \frac{d\xi}{\pi}\right)_{12} \\ &\quad - \left(\frac{1}{\sqrt{12\alpha t}} \int_{\Sigma_B} ((1 - C_{\omega_B^0})^{-1} I)(\xi) \omega_B^0(\xi) \frac{d\xi}{\pi}\right)_{12} \\ &\quad + o\left(\frac{\log t}{t}\right). \end{aligned} \quad (3.117)$$

For  $z \in \Sigma_A$ , set

$$M^{A0}(z, x, t) = I + \int_{\Sigma_A} \frac{((1 - C_{\omega_A^0})^{-1} I)(\xi) \omega_A^0(\xi) d\xi}{\xi - z} \frac{d\xi}{2\pi i}, \quad (3.118)$$

then  $M^{A0}(z, x, t)$  is the solution of the following Riemann-Hilbert problem

$$\begin{cases} M_+^{A0}(z, x, t) = M_-^{A0}(z, x, t) v^{A0}(z, x, t), & z \in \Sigma_A, \\ M^{A0}(z, x, t) \rightarrow I, & z \rightarrow \infty. \end{cases} \quad (3.119)$$

$M^{A0}(z) = I + \frac{M_1^{A0}}{z} + o(z^{-2})$ ,  $z \rightarrow \infty$ , we have  $M_1^{A0} = -\int_{\Sigma_A} ((1 - C_{\omega_A^0})^{-1} I) \omega_A^0(\xi) \frac{d\xi}{2\pi i}$ .

For  $z \in \Sigma_B$ , set

$$M^{B0}(z, x, t) = I + \int_{\Sigma_B} \frac{((1 - C_{\omega_B^0})^{-1} I)(\xi) \omega_B^0(\xi) d\xi}{\xi - z} \frac{d\xi}{2\pi i}. \quad (3.120)$$



then  $M^{B0}(z, x, t)$  is the solution of the following Riemann-Hilbert problem

$$\begin{cases} M_+^{B0}(z, x, t) = M_-^{B0}(z, x, t)v^{B0}(z, x, t), & z \in \Sigma_B, \\ M^{B0}(z, x, t) \rightarrow I, & z \rightarrow \infty. \end{cases} \quad (3.121)$$

$M^{B0}(z) = I + \frac{M_1^{B0}}{z} + o(z^{-2})$ ,  $z \rightarrow \infty$ , we have  $M_1^{B0} = -\int_{\Sigma_B} ((1 - C_{\omega_B^0})^{-1}I)\omega_B^0(\xi)\frac{d\xi}{2\pi i}$ .  $v^{B0} = (I - \omega_{B^0-})^{-1}(I + \omega_{B^0+})$ , where

$$\omega_B^0 = \omega_{B^0+} = \begin{cases} \begin{pmatrix} 0 & (\delta_B^0)^2 z^{2iv} e^{-\frac{iz^2}{2} \overline{r(z_0)}} \\ 0 & 0 \end{pmatrix}, & z \in \Sigma_B^1, \\ \begin{pmatrix} 0 & (\delta_B^0)^2 z^{2iv} e^{-\frac{iz^2}{2} \frac{-\overline{r(z_0)}}{1-|r(z_0)|^2}} \\ 0 & 0 \end{pmatrix}, & z \in \Sigma_B^3, \end{cases} \quad (3.122)$$

$$\omega_B^0 = \omega_{B^0-} = \begin{cases} \begin{pmatrix} 0 & 0 \\ (\delta_B^0)^{-2} z^{-2iv} e^{\frac{iz^2}{2} r(z_0)} & 0 \end{pmatrix}, & z \in \Sigma_B^2, \\ \begin{pmatrix} 0 & 0 \\ (\delta_B^0)^{-2} z^{-2iv} e^{\frac{iz^2}{2} \frac{r(z_0)}{1-|r(z_0)|^2}} & 0 \end{pmatrix}, & z \in \Sigma_B^4, \end{cases} \quad (3.123)$$

$$\delta_B^0 = e^{2i\alpha\tau} (12\alpha\tau)^{-\frac{iv}{2}} e^{\chi(z_0)}. \quad (3.124)$$

By using (3.112)-(3.114) and (3.122)-(3.124), we have

$$v^{A0}(z) = \overline{(v^{B0})(-\bar{z})}. \quad (3.125)$$

According to the uniqueness of the solution,

$$M^{A0}(z) = \overline{(M^{B0})(-\bar{z})}, \quad (3.126)$$

therefore

$$M_1^{A0} = -\overline{(M_1^{B0})}, \quad (3.127)$$

then we have

$$u(x, t) = \frac{i}{\sqrt{3\alpha t z_0}} (M_1^{B0} - \overline{M_1^{B0}})_{12}. \quad (3.128)$$

### 3.6. Solving the model problem.

In this subsection, our work is devoted to compute  $(M_1^{B0})_{12}$  explicitly. Firstly, we introduce the transformation  $\psi(z) = H(z)z^{iv\sigma_3} e^{-\frac{iz^2}{4}\sigma_3}$ ,  $H(z) = (\delta_B^0)^{-\sigma_3} M_B^0(z)(\delta_B^0)^{\sigma_3}$ , then

$$\begin{aligned} \psi_+(z) &= (\delta_B^0)^{-\sigma_3} M_{B+}^0(z)(\delta_B^0)^{\sigma_3} z^{iv\sigma_3} e^{-\frac{iz^2}{4}\sigma_3} \\ &= (\delta_B^0)^{-\sigma_3} M_{B-}^0(z)v^{B0}(z)(\delta_B^0)^{\sigma_3} z^{iv\sigma_3} e^{-\frac{iz^2}{4}\sigma_3} \\ &= \psi_-(z)v(z_0) \end{aligned} \quad (3.129)$$

where  $v(z_0) = z^{-iv\hat{\sigma}_3} e^{\frac{iz^2}{4}\hat{\sigma}_3} (\delta_B^0)^{-\hat{\sigma}_3} v^{B_0}$ . Notice that the jump matrix is irrelevant to  $z$  on each ray, so

$$\frac{d\psi_+(z)}{dz} = \frac{d\psi_-(z)}{dz} v(z_0) \quad (3.130)$$

and

$$\left(\frac{d\psi_+(z)}{dz}\right)\psi_+^{-1}(z) = \frac{d\psi_-(z)}{dz} v(z_0) v^{-1}(z_0) \psi_-^{-1}(z) = \frac{d\psi_-(z)}{dz} \psi_-^{-1}(z), \quad (3.131)$$

therefore,  $\frac{d\psi}{dz} \psi^{-1}(z)$  has no jump discontinuity along any of the ray. Moreover, from the relation between  $\psi(z)$  and  $H(z)$ , we have

$$\begin{aligned} \frac{d\psi}{dz} \psi^{-1}(z) &= \frac{dH(z)}{dz} H^{-1}(z) - \frac{i}{2} z H(z) \sigma_3 H^{-1}(z) \\ &= -\frac{i}{2} z \sigma_3 - \frac{i}{2} (\delta_B^0)^{-\hat{\sigma}_3} [M_1^{B_0}, \sigma_3]. \end{aligned} \quad (3.132)$$

It follows by the Liouville's theorem that

$$\frac{d\psi(z)}{dz} + \frac{i}{2} z \sigma_3 \psi(z) = \beta \psi(z), \quad (3.133)$$

where  $\beta = -\frac{i}{2} (\delta_B^0)^{-\hat{\sigma}_3} [M_1^{B_0}, \sigma_3] = \begin{pmatrix} 0 & \beta_{12} \\ \beta_{21} & 0 \end{pmatrix}$ . So we have

$$M_1^{B_0}{}_{12} = -i(\delta_B^0)^2 \beta_{12}. \quad (3.134)$$

Set

$$\psi(z) = \begin{pmatrix} \psi_{11}(z) & \psi_{12}(z) \\ \psi_{21}(z) & \psi_{22}(z) \end{pmatrix},$$

from Eq.(3.133), we get

$$\frac{d^2 \psi_{11}(z)}{dz^2} = \left(-\frac{i}{2} - \frac{z^2}{4} + \beta_{12} \beta_{21}\right) \psi_{11}(z), \quad (3.135)$$

$$\beta_{12} \psi_{21}(z) = \frac{d\psi_{11}(z)}{dz} + \frac{i}{2} z \psi_{11}(z), \quad (3.136)$$

$$\frac{d^2 \psi_{22}(z)}{dz^2} = \left(\frac{i}{2} - \frac{z^2}{4} + \beta_{12} \beta_{21}\right) \psi_{22}(z), \quad (3.137)$$

$$\beta_{21} \psi_{12}(z) = \frac{d\psi_{11}(z)}{dz} - \frac{i}{2} z \psi_{11}(z). \quad (3.138)$$

We know that Weber's equation

$$d^2 g(\xi) + \left(a + \frac{1}{2} - \frac{\xi^2}{4}\right) g(\xi) = 0 \quad (3.139)$$

has the solution

$$g(\xi) = c_1 D_a(\xi) + c_2 D_a(-\xi), \quad (3.140)$$

where  $D_a(\cdot)$  denotes the standard parabolic-cylinder function and satisfies

$$\frac{dD_a(\xi)}{d\xi} + \frac{\xi}{2} D_a(\xi) - a D_{a-1}(\xi) = 0, \quad (3.141)$$

$$D_a(\pm\xi) = \frac{\Gamma(a+1)e^{\frac{i\pi a}{2}}}{\sqrt{2\pi}}D_{-a-1}(\pm i\xi) + \frac{\Gamma(a+1)e^{-\frac{i\pi a}{2}}}{\sqrt{2\pi}}D_{-a-1}(\mp i\xi). \quad (3.142)$$

According to [26], as  $\xi \rightarrow \infty$ , we have

$$D_a(\xi) = \begin{cases} \xi^a e^{-\frac{\xi^2}{4}} (1 + o(\xi^{-2})), & |\arg \xi| < \frac{3\pi}{4}, \\ \xi^a e^{-\frac{\xi^2}{4}} (1 + o(\xi^{-2})) - \frac{\sqrt{2\pi}}{\Gamma(-a)} e^{a\pi i + \frac{\xi^2}{4}} \xi^{-a-1} (1 + o(\xi^{-2})), & \frac{\pi}{4} < \arg \xi < \frac{5\pi}{4}, \\ \xi^a e^{-\frac{\xi^2}{4}} (1 + o(\xi^{-2})) - \frac{\sqrt{2\pi}}{\Gamma(-a)} e^{-a\pi i + \frac{\xi^2}{4}} \xi^{-a-1} (1 + o(\xi^{-2})), & -\frac{5\pi}{4} < \arg \xi < -\frac{\pi}{4}, \end{cases} \quad (3.143)$$

where  $\Gamma(\cdot)$  is the Gamma function. Setting  $a = i\beta_{12}\beta_{21}$ , we have

$$\psi_{11}(z) = c_3 D_a(e^{-\frac{3\pi i}{4}} z) + c_4 D_a(e^{\frac{\pi i}{4}} z) \quad (3.144)$$

and

$$\beta_{12}\psi_{22}(z) = c_5 D_{-a}(e^{\frac{3\pi i}{4}} z) + c_6 D_{-a}(e^{-\frac{\pi i}{4}} z). \quad (3.145)$$

When  $\arg(z) \in (-\frac{\pi}{4}, \frac{\pi}{4})$  and  $z \rightarrow \infty$ , we have  $\psi_{11}(z) \rightarrow z^{iv} e^{-\frac{iz^2}{4}}$ ,  $\psi_{22}(z) \rightarrow z^{-iv} e^{\frac{iz^2}{4}}$ . By using Eq.(3.143) and (3.144), we obtain

$$c_3 = 0, \quad c_4 = e^{\frac{\pi v}{4}}, \quad a = iv, \quad \psi_{11}(z) = e^{\frac{\pi v}{4}} D_a(e^{\frac{\pi i}{4}} z). \quad (3.146)$$

Similarly, by using Eq.(3.143) and (3.145), we get

$$c_5 = 0, \quad c_6 = \beta_{12} e^{\frac{\pi v}{4}}, \quad a = iv, \quad \beta_{12}(z)\psi_{22}(z) = \beta_{12}(z) e^{\frac{\pi v}{4}} D_{-a}(e^{-\frac{\pi i}{4}} z). \quad (3.147)$$

Considering Eq.(3.133) and (3.136), we obtain

$$\beta_{21}\psi_{21}(z) = a e^{\frac{\pi(v+i)}{4}} D_{a-1}(e^{\frac{\pi i}{4}} z). \quad (3.148)$$

When  $\arg(z) \in (-\frac{3\pi}{4}, -\frac{\pi}{4})$  and  $z \rightarrow \infty$ , we have  $\psi_{11}(z) \rightarrow z^{iv} e^{-\frac{iz^2}{4}}$ ,  $\psi_{22}(z) \rightarrow z^{-iv} e^{\frac{iz^2}{4}}$ , which implies

$$\psi_{11}(z) = e^{\frac{\pi v}{4}} D_a(e^{\frac{\pi i}{4}} z) \quad (3.149)$$

and

$$\beta_{12}\psi_{22}(z) = \beta_{12} e^{-\frac{3\pi v}{4}} D_{-a}(e^{\frac{3\pi i}{4}} z), \quad (3.150)$$

consequently,

$$\psi_{12}(z) = \beta_{12} e^{\frac{\pi v}{4}} e^{-\frac{3\pi i}{4}} D_{-a-1}(e^{-\frac{\pi i}{4}} z). \quad (3.151)$$

Along the ray  $\arg z = -\frac{\pi}{4}$ ,  $\psi_+(z) = \psi_-(z) \begin{pmatrix} 0 & \overline{r(z_0)} \\ 0 & 0 \end{pmatrix}$  and  $\beta_{12} e^{\frac{\pi(v-3i)}{4}} D_{-a-1}(e^{-\frac{\pi i}{4}} z) = e^{\frac{\pi v}{4}} D_a(e^{\frac{\pi i}{4}} \overline{r(z_0)}) + \beta_{12} e^{\frac{\pi(i-3v)}{4}} D_{-a-1}(e^{\frac{3\pi i}{4}} z)$ . From Eq.(3.142), we have

$$D_a(e^{\frac{\pi i}{4}} z) = \frac{\Gamma(a+1)e^{\frac{i\pi a}{2}}}{\sqrt{2\pi}} D_{-a-1}(e^{\frac{3\pi i}{4}} z) + \frac{\Gamma(a+1)e^{-\frac{i\pi a}{2}}}{\sqrt{2\pi}} D_{-a-1}(e^{-\frac{\pi i}{4}} z). \quad (3.152)$$

Therefore, we obtain

$$\beta_{12} = \frac{e^{\frac{3\pi i}{4} + \frac{\pi v}{2}} \Gamma(1+a)}{\sqrt{2\pi}} \overline{r(z_0)} = \frac{e^{(-\frac{3\pi i}{4} + \frac{\pi v}{2})} v \Gamma(iv)}{\sqrt{2\pi}} \overline{r(z_0)}. \quad (3.153)$$

Finally, theorem (1.1) gives the long-time asymptotics of Eq.(1.8).

## 4. Conclusion

In this paper, we study long-time asymptotics of the generalized AKNS equation by Defit-Zhou steepest descent method, in which the Schwartz initial data  $u_0(x) \in \mathcal{S}(\mathbb{R})$  is inquired. The corresponding result is given in theorem 1.1.

In short order, we plan to consider weaker weighted Sobolev initial data  $u_0(x) \in H^{1,1}(\mathbb{R})$  and  $u_0(x) \in H^{2,2}(\mathbb{R})$  of long-time asymptotics to some NLPDEs.

## Acknowledgements

This work was supported by the National Natural Science Foundation of China (grant No.11971475).

## Data Availability

Data sharing is not applicable to this article as no new data were created or analyzed in this study.

## References

- [1] S. K. El-Labany, W. M. Moslem, E. I. El-Awady et al., Nonlinear dynamics associated with rotating magnetized electron-positron-ion plasmas, *Physics Letters A*, 2010, 375(2): 159-164.
- [2] M. Alam, M. Rahman, R. Islam et al., Application of the new extended (G'/G)-expansion method to find exact solutions for nonlinear partial differential equation, *Computational Methods for Differential Equations*, 2015, 3(1): 59-69.
- [3] Y. Zhang, Q. Liu, Z. Qiao, et al., Fifth-order b-family Novikov (FObFN) model with pseudo-peakons and multi-peakons, *Modern Physics Letters B*, 2019, 33(18).
- [4] A. Chabchoub, N. Hoffmann, M. Onorato, et al., Super rogue waves: observation of a higher-order breather in water waves, *Physical Review X*, 2012, 2(1): 011015.
- [5] A. Yokus, H. M. Baskonus, T. A. Sulaiman et al., Numerical simulation and solutions of the two-component second order KdV evolutionary system, *Numerical Methods for Partial Differential Equations*, 2018, 34(1): 211-227.
- [6] C. M. Khaliq, I. E. Mhlanga, Travelling waves and conservation laws of a (2+1)-dimensional coupling system with Korteweg-de Vries equation, *Applied Mathematics and Nonlinear Sciences*, 2018, 3(1): 241-254.
- [7] Y. Zhang, J. Mei, X. Zhang, et al., Symmetry properties and explicit solutions of some nonlinear differential and fractional equations, *Applied Mathematics and Computation*, 2018: 408-418.
- [8] K. Dysthe, H. E. Krogstad, P. Müller, Oceanic rogue waves, *Annual Review of Fluid Mechanics*, 2008, 40: 287-310.
- [9] J. Chen, Y. Chen, B. Feng, et al., Rational solutions to two- and one-dimensional multicomponent Yajima-Oikawa systems, *Physics Letters A*, 2015, 379(24-25): 1510-1519.
- [10] A. Abdeljabbar, W. Ma, A. Yildirim, Determinant solutions to a (3+1)-dimensional generalized KP equation with variable coefficients, *Chinese Annals of Mathematics, Series B*, 2012, 33(5): 641-650.
- [11] J. Zhang, L. Wang, C. Liu, Superregular breathers, characteristics of nonlinear stage of modulation instability induced by higher-order effects, *Proceedings of the Royal Society A Mathematical Physical and Engineering Sciences*, 2017, 473(2199): 20160681.
- [12] X. Wang, Y. Li, Y. Chen, Generalized Darboux transformation and localized waves in coupled Hirota equations, *Wave Motion*, 2014, 51(7): 1149-1160.

- [13] E. K. Sklyanin, Method of the inverse scattering problem and the nonlinear quantum Schrödinger equation, Soviet Physics Doklady, 1979, 24.
- [14] M. J. Ablowitz, D. J. Kaup, A. C. Newell, and H. Segur, The inverse scattering transform-Fourier analysis for nonlinear problems, Studies in Applied Mathematics, 1974, 53: 2499-315.
- [15] C. S. Gardner, J. M. Greene, M. D. Kruskal, R. M. Miura, Method for solving the Korteweg-de Vries equation, Physical Review Letters, 1967, 19: 1095-1097.
- [16] V. E. Zakharov, A. B. Shabat, A scheme for integrating the nonlinear equations of mathematical physics by the method of the inverse scattering problem, Functional Analysis and Its Applications, 1974, 8(3):226-235.
- [17] P. A. Deift, X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Annals of Mathematics, 1993, 137:295-368.
- [18] P. A. Deift, X. Zhou, Long-time asymptotics for integrable nonlinear wave equation, in "Important developments in soliton theory", Springer Ser, Nonlinear Dynamics, Springer, Berlin, 1993, 181-204.
- [19] J. Xu, E. Fan, Long-time asymptotics for the Fokas-Lenells equation with decaying initial value problem: Without solitons, Journal of Differential Equations, 2015, 259(3):1098-1148.
- [20] J. Xu, E. Fan, et al., Long-time Asymptotic for the Derivative Nonlinear Schrödinger Equation with Step-like Initial Value, Mathematical Physics, Analysis & Geometry, 2013, 16(3):253-288.
- [21] Q. Cheng, E. Fan, Long-time asymptotics for a mixed nonlinear Schrödinger equation with the Schwartz initial data, Journal of Mathematical Analysis and Applications, 2020, 489(2):124188.
- [22] X. Geng, M. Chen, K. Wang, Long-time asymptotics of the coupled modified Korteweg-de Vries equation, Journal of Geometry and Physics, 2019, 142: 151-167.
- [23] X. Geng, H. Liu, The nonlinear steepest descent method to long-time asymptotics of the coupled nonlinear Schrödinger equation, Journal of Nonlinear Science, 2018, 28(2): 739-763.
- [24] X. Geng, K. Wang, M. Chen, Long-time asymptotics for the spin-1 Gross-Pitaevskii equation, Communications in Mathematical Physics, 2021, 382(1): 585-611.
- [25] Y. Li, Solitons and integrable systems, Shanghai Science & Technology Education Press, 1999.
- [26] E.T. Whittaker, G.N. Watson, A Course of Modern Analysis, fourth ed., Cambridge University Press, Cambridge, 1927.