

On vortex stretching for anti-parallel axisymmetric flows

Kyudong Choi* In-Jee Jeong[†]

October 19, 2021

Abstract

We consider axisymmetric incompressible inviscid flows without swirl in \mathbb{R}^3 , under the assumption that the axial vorticity is non-positive in the upper half space and odd in the last coordinate. This flow setup corresponds to the head-on collision of anti-parallel vortex rings. We prove infinite growth of the vorticity impulse on the upper half space. As direct applications, we achieve $\limsup_{t \rightarrow \infty} \|\omega(t, \cdot)\|_{L^\infty} = \infty$ for certain compactly supported data $\omega_0 \in C^{1,\alpha}(\mathbb{R}^3)$ with $0 < \alpha < 3/17$ and $\|\omega(t, \cdot)\|_{L^p} \gtrsim t^{1/15-}$ with any $2 \leq p \leq \infty$ for patch type data with smooth boundary and profile.

Contents

1	Introduction	2
1.1	Main results	2
1.2	Discussion	4
1.3	Outline of the proof	6
1.4	Notation and conventions	8
2	Preliminaries	8
2.1	Elliptic integrals	8
2.2	Stream function and Biot–Savart law	9
2.3	Axisymmetric flows without swirl	10
3	Key Inequalities	12
3.1	Relation between P and vorticity L^p norms	12
3.2	Estimates on kinetic energy and derivative of P	13
3.3	Monotonicity Lemma	16
4	Growth	17
4.1	Preliminary growth	17
4.2	Enhancing the growth rate	19
5	Enstrophy growth for Navier–Stokes solutions	22

*Department of Mathematical Sciences, Ulsan National Institute of Science and Technology, kchoi@unist.ac.kr
[†]Department of Mathematical Sciences and RIM, Seoul National University, injee.j@snu.ac.kr

1 Introduction

1.1 Main results

In this paper, we consider the evolution of three-dimensional incompressible inviscid fluids under the assumption of *axisymmetric without swirl* in \mathbb{R}^3 . The governing system the axisymmetric Euler equations

$$\partial_t \omega + u \cdot \nabla \omega = \frac{u^r}{r} \omega, \quad u = K[\omega] \quad (1.1)$$

where $u \cdot \nabla := u^r(t, r, z) \partial_r + u^z(t, r, z) \partial_z$ and (r, θ, z) is the cylindrical coordinate system in \mathbb{R}^3 . Here, $\omega = \omega^\theta(t, r, z) e_\theta$ denotes the axial vorticity, and the operator $K[\cdot]$ comes from symmetrizing the Biot–Savart kernel $\nabla \times (-\Delta)^{-1}$; see below for its precise form. It is well-known that for initial data $\omega_0 = \omega_0(r, z)$ satisfying

$$\omega_0, \frac{\omega_0}{r} \in L^1 \cap L^\infty(\mathbb{R}^3), \quad (1.2)$$

there is a unique global-in-time solution $\omega(t, \cdot)$ to (1.1) satisfying $\omega, r^{-1} \omega \in L^\infty([0, T]; L^1 \cap L^\infty(\mathbb{R}^3))$ for any $T > 0$ (by [52], also see [44, 11]).

In addition to the above regularity assumption (1.2), we are concerned with vorticities which are odd in z (“anti-parallel”) and non-positive on the upper half-plane; that is,

$$\omega(t, r, z) = -\omega(t, r, -z), \quad \omega(t, r, z) \leq 0 \quad \text{for } z \geq 0. \quad (1.3)$$

Then, it is not difficult to see that the assumptions in (1.3) are satisfied for the unique solution, if they hold for the initial data ω_0 . Furthermore, (1.1) reduces to a system posed on the upper half-space \mathbb{R}_+^3 , and we shall often regard ω as defined on \mathbb{R}_+^3 . Our first main result ensures infinite growth of the vorticity support for *all* compactly support initial data.

Theorem 1.1. *Together with (1.2) and (1.3), assume that the initial vorticity satisfies*

$$0 < \iint_{[0, \infty)^2} -z \omega_0(r, z) \, dr dz < +\infty \quad \text{and} \quad 0 < \iint_{[0, \infty)^2} -r^2 \omega_0(r, z) \, dr dz < +\infty.$$

Then, the unique global solution $\omega(t, \cdot)$ to (1.1) satisfies that

$$\iint_{[0, \infty)^2} -z \omega(t, r, z) \, dr dz \quad \text{is strictly decreasing in time,}$$

$$\iint_{[0, \infty)^2} -r^2 \omega(t, r, z) \, dr dz \quad \text{is strictly increasing in time,}$$

and for any $\varepsilon > 0$,

$$\iint_{[0, \infty)^2} -r^2 \omega(t, r, z) \, dr dz \geq C_\varepsilon (1+t)^{\frac{2}{15}-\varepsilon} \quad \text{for all } t \geq 0,$$

where $C_\varepsilon > 0$ is a constant depending on ε and ω_0 . In particular, from (1.1) we deduce that

$$\sup\{r : (r, z) \in \text{supp } \omega(t, \cdot)\} \geq C_\varepsilon (1+t)^{\frac{1}{15}-\varepsilon} \quad \text{for all } t \geq 0$$

holds whenever ω_0 is compactly supported in r .

In the proof, we shall specify the dependence of C_ε in ω_0 , which is somewhat complicated. As a direct consequence of the above, we are able to deduce infinite growth of the L^p -norms of the vorticity for $p \geq 2$.

Corollary 1.2. *Let $0 \leq \delta < 1/15$. Assume that for some $p \in [2 - \delta, \infty]$, the initial data satisfies*

$$\|r\omega_0^{-1}\mathbf{1}_{\{\omega_0 < 0\}}\|_{L^{\frac{1-\delta}{1-(2-\delta)/p}}(\mathbb{R}_+^3)} < \infty \quad \text{and} \quad \sup\{r : (r, z) \in \text{supp}(\omega_0(\cdot))\} < \infty \quad (1.4)$$

in addition to the hypotheses of Theorem 1.1. Then, for each $\varepsilon > 0$, we have

$$\|\omega(t, \cdot)\|_{L^p(\mathbb{R}^3)} \geq C_\varepsilon(1+t)^{\frac{2}{15}-\frac{2\delta}{2-\delta}-\varepsilon} \quad \text{for all } t \geq 0,$$

where $C_\varepsilon > 0$ is a constant depending on ε and ω_0 .

Remark 1.3. While the assumption (1.4) may look strange, there are plenty of initial data ω_0 satisfying it. For instance, the above result applies to the class of *vortex patches*, namely

$$\omega(t, r, z) = \sum_{i=1}^n -a_i(t, r, z)\mathbf{1}_{\Omega_i(t)}(r, z) \quad \text{on } (r, z) \in [0, \infty)^2. \quad (1.5)$$

Here, Ω_i are bounded axisymmetric open sets which are disjoint with each other and separated from $\{r = 0\}$, and a_i are some bounded non-negative functions.¹ Then, (1.4) is satisfied for all $p \in [2, \infty]$ (with $\delta = 0$) as long as $a_i(t = 0)$ is bounded from below on $\overline{\Omega}_i$ for all $1 \leq i \leq n$, and we obtain $\max_i \|a_i(t, \cdot)\|_{L^p} \gtrsim t^{\frac{1}{15}-}$ and $\max_i \text{diam } \Omega_i(t, \cdot) \gtrsim t^{\frac{1}{15}-}$.

Remark 1.4. To apply the above result to smoother vorticities, we first observe that compactly supported and continuous f can lie on L^q , $q > 0$ (i.e. $\int |f|^q dx < \infty$) when it touches the boundary of its support by rate x^α with

$$\alpha \cdot q < 1.$$

For instance, in the case $p = \infty$, for each $0 < \gamma < 1/14$, the assumption (1.4) holds for some compactly-supported (away from the symmetry axis $\{r = 0\}$) $C^{1,\gamma}$ vorticities ω_0 (with any choice of $\delta \in (\gamma/(1+\gamma), 1/15)$). As a consequence, the solutions from such data $\omega_0 \in C_c^{1,\gamma}(\mathbb{R}^3 \setminus \{r = 0\})$ satisfy, for some $\alpha = \alpha(\gamma) > 0$,

$$\|\omega(t, \cdot)\|_{L^\infty} \geq C(1+t)^\alpha, \quad t \geq 0.$$

To the best of our knowledge, this is the first construction of initial vorticity ω_0 , which is compactly supported and is more regular than C^1 (so that the velocity u is more regular than C^2), with infinite growth of the vorticity maximum as $t \rightarrow \infty$, for the incompressible Euler equations in \mathbb{R}^3 .

Remark 1.5. Similarly, when $p = 2$, for each $0 < \gamma < 1/28$, there are some vorticities $\omega_0 \in C_c^{0,\gamma}(\mathbb{R}^3 \setminus \{r = 0\})$ satisfying (1.4) so that

$$\|\omega(t, \cdot)\|_{L^2} \geq C(1+t)^\alpha, \quad t \geq 0 \quad (1.6)$$

for some $\alpha > 0$ depending on γ . In Section 5, we connect this infinite enstrophy growth for Euler into enstrophy inflation for Navier–Stokes with small viscosity.

¹It is well-known that for axisymmetric initial data of the form (1.5), C^∞ -regularity of a_i and $\partial\Omega_i$ propagates globally in time ([27, 22, 45]). The associated velocity field is C^∞ -smooth in $\overline{\Omega}_i$.

Both the growth rate of $\beta_1(\delta) := \frac{2-2\delta}{2-\delta}$ and the range of $\delta \in [0, 1/15)$ from Corollary 1.2 can be upgraded, if one takes limsup in time. In the case of L^∞ , we have:

Theorem 1.6. *Assuming that (1.4) holds for $p = +\infty$ and for some $\delta \in [0, 3/20)$ in addition to the hypotheses of Theorem 1.1, we have some $\beta_0 = \beta_0(\delta) > \max\{\beta_1(\delta), 0\}$ satisfying*

$$\limsup_{t \rightarrow \infty} \frac{\|\omega(t, \cdot)\|_{L^\infty(\mathbb{R}^3)}}{(1+t)^{\beta_0-\varepsilon}} = +\infty.$$

for any $\varepsilon > 0$.

A version of Theorem 1.6 for *finite* p can also be derived with some different β_0 and δ depending on p .

Remark 1.7. As in Remark 1.4, the above theorem has a direct consequence:

For each $0 < \gamma < 3/17$, there are initial vorticities $\omega_0 \in C_c^{1,\gamma}(\mathbb{R}^3 \setminus \{r = 0\})$ satisfying

$$\limsup_{t \rightarrow \infty} \|\omega(t, \cdot)\|_{L^\infty} = \infty.$$

1.2 Discussion

Let us present some motivation for studying vorticities of the form (1.3) as well as related previous works.

Head-on collision of antiparallel vortex rings. Our primary motivation for studying axisymmetric vorticity satisfying (1.3) comes from the study of head-on collision of vortex rings ([30, 31, 46, 10, 47, 35, 49, 40, 39, 4, 23, 48]). The interested readers may find the flow visualizations in [26] as well as in aforementioned references. Since a single vortex ring with negative axial vorticity travels downward, two vortex rings with opposite sign will “collide” with each other if the ring with negative vorticity is on the above of the other. This flow setup exhibits several fascinating features including strong vortex stretching, bulging instability, rebound effects, breakdown of axisymmetry and vortex reconnection, and has been studied extensively using experimental, theoretical, and computational methods. The authors in [10] demonstrate that the evolution of the vortex rings goes through three stages, before their breakdown: (i) free-traveling, (ii) vortex stretching, and (iii) viscous dissipation. This is intuitively clear: initially, the rings are separated and they travel towards each other as the interaction is weak. As they become sufficiently close, the rings start to stretch out to radial direction due to incompressibility. Numerical simulations and experiments clearly show that as the Reynolds number becomes large, the rings get stretched further away from the symmetry axis before viscous dissipation becomes dominant. In the current work, we employ the inviscid equation and focus on the vortex stretching phenomenon for large times. The basic question we try to answer is: what is the rate of vorticity growth as the time goes to infinity? To answer this question, one can hope to gain some insights from the simpler case of two-dimensional flows.

Vortex dynamics in the two-dimensional case. Under the odd symmetry assumption in z , axisymmetric Euler flows without swirl share structural similarities with two-dimensional Euler flows subject to the odd-odd symmetry: the planar vorticity ω satisfying $\omega(x_1, x_2) = -\omega(-x_1, x_2) = -\omega(x_1, -x_2)$ in \mathbb{R}^2 . Assuming that the vorticity is non-negative on the first quadrant $(\mathbb{R}_+)^2$, it can

be shown that the overall flow is directed southeast on $(\mathbb{R}_+)^2$. If one takes a single point vortex with positive sign on $(\mathbb{R}_+)^2$ and extend it to \mathbb{R}^2 with odd-odd symmetry, then one can directly compute that the vortex on the first quadrant goes out to the x_1 -axis linearly in time while its second component asymptotes to a positive constant ([53]). Even for general non-negative vorticities on $(\mathbb{R}_+)^2$, a similar result is available ([28]): the center of vorticity $\int_{(\mathbb{R}_+)^2} x_1 \omega(t, x) dx$ grows linearly for all positive times. We note that this odd-odd scenario has been used to prove growth of $|\nabla \omega|$ in two-dimensional Euler flows ([12, 13, 33, 55, 32, 29]).

Dyson model for vortex rings. Returning to axisymmetric flows and the problem of finding the rate of vortex stretching, it is natural to first consider the dynamics of circular vortex filaments (thin-cored vortex rings), which are the axisymmetric versions of point vortices in 2D. The well-known difficulty in the filament case is that for a fixed circulation Γ and distance to the axis R , the self-induction velocity diverges as the core radius a goes to zero: the calculation attributed to Kelvin ([51, 24, 34]) shows that for $a \ll R$,

$$u^z = \frac{\Gamma}{4\pi R} \left(\ln \frac{8R}{a} - \frac{1}{4} + O\left(\frac{a}{R}\right) \right).$$

To handle this difficulty, the usual assumption is that the core of each vortex ring remains circular with a *fixed* radius; this is the *localized induction approximation*. Then, proceeding similarly as in the 2D case, one can derive a system of ordinary differential equations for the motion of thin cored vortex rings (see [34, 53] for instance); it is commonly referred to as Dyson's model in the literature ([15]). Applying Dyson's model to the system of two antiparallel vortex rings, one obtains that the dynamics is essentially the same as the point vortex motion under the odd-odd symmetry, which suggests $\|\omega(t, \cdot)\|_{L^\infty} \sim t$. However, as the rings move away from the symmetry axis, vortex stretching together with conservation of total vorticity $\|r^{-1}\omega(t, \cdot)\|_{L^1}$ forces that the core area must vanish to zero: this shows that the localized induction approximation is **self-inconsistent**. This suggests that the asymptotic behavior as $t \rightarrow \infty$ for antiparallel vortex rings could be different from the two-dimensional case and more difficult to understand.

Childress model of vortex growth. In a series of works ([5, 6, 7]), Childress and his collaborators investigated the exact same problem: what is the rate of vortex stretching in axisymmetric flows under the assumption (1.3)? Due to the conservation of $r^{-1}\omega$ along particle trajectories, for compactly supported vorticities, the question is roughly the same with finding the growth rate of the vorticity support in r . The naive a priori estimates give that the rate is bounded by e^{Ct} , with $C > 0$ depending on the initial data. In [5, 6], Childress shows that this upper bound can be improved to t^2 and $t^{4/3}$ by solving certain maximization problems using the constraint of support volume and kinetic energy, respectively. In these works Childress imposes the conditions (1.3) on vorticity, among others.² The $t^{4/3}$ bound can be seen heuristically as follows (see [6, 7]): assuming $\omega(t, r, z) \simeq R(t)\Omega(a(t)^{-1}(r, z) - (R(t), 0))$ with some profile Ω and $R(t) > a(t) > 0$, the kinetic energy conservation dictates the scaling $R^3 a^4 \sim 1$ while the Euler evolution forces $\dot{R} \sim Ra$. While this ansatz clearly contradicts the conservation of circulation, Childress argues that the formation of a long tail accounts for the loss of circulation. Numerical simulations from [7] suggest that growth rate of $t^{4/3}$ can be indeed achieved, with asymptotic profile Ω given by the Sadvskii vortex.

²We note that the upper bound of t^2 can be proved in general using a rather recent estimate of Feng–Sverak [21], see Lemma 2.2 below.

Growth of vorticity for three-dimensional Euler. There has been significant interest in the construction of Euler flows with vorticity maximum growing in time. The 3D Euler equations in vorticity form are given by

$$\partial_t \omega + u \cdot \nabla \omega = \omega \cdot \nabla u, \quad u = \nabla \times (-\Delta)^{-1} \omega$$

and unlike the 2D case, $\|\omega(t, \cdot)\|_{L^\infty}$ could increase when the vorticity vector aligns with an eigenvector of ∇u with a positive eigenvalue. For smooth and decaying solutions to the 3D Euler equations, it is well-known that finite time singularity formation could occur if and only if $\|\omega(t, \cdot)\|_{L^\infty}$ blows up in finite time. A few blow-up results for finite-energy solutions to 3D Euler exist ([18, 17, 16, 50, 3])³ but in these results either the presence of physical boundaries or lack of smoothness⁴ of vorticity plays an important role in the growth. Even when the vorticity does not blow up, it is an interesting question to understand the possible rate of vortex stretching in various situations. If one considers axisymmetric domains with a boundary, then one can obtain growth of the vorticity maximum along the boundary [54]. Even without physical boundaries, rather simple examples of vorticity growth can be obtained using the so-called $2 + \frac{1}{2}$ dimensional flow construction, but this requires the physical domain to be in the form $\Omega \times \mathbb{T}$ where Ω is a two-dimensional domain ([54]). Even in the case of \mathbb{T}^3 , we are not aware of any results giving infinite vorticity growth for smooth vorticity, not relying on the $2 + \frac{1}{2}$ dimensional geometry (however see [42]). In our previous work, we obtained arbitrarily large but finite growth of $\|\omega(t, \cdot)\|_{L^\infty}$ in \mathbb{R}^3 using perturbations of the Hill's vortex [9]. Very recent numerical computations by Hou suggests finite time singularity for axisymmetric Euler with swirl in the interior of the domain [25].

1.3 Outline of the proof

Let us outline the proof of Theorem 1.1, which is inspired by a related work of Iftimie–Sideris–Gamblin [28]. Among others, the authors prove that for compactly supported and non-negative initial vorticity on the positive quadrant $(\mathbb{R}_+)^2$ (extended to \mathbb{R}^2 by odd-odd symmetry as described above), the 2D Euler solution satisfies

$$\iint_{(\mathbb{R}_+)^2} x_1 \omega(t, x) dx \geq ct, \tag{1.7}$$

with $c = c(\omega_0) > 0$. Their proof consists of two main steps: the first is to establish the monotonicity property

$$\frac{d}{dt} \iint_{(\mathbb{R}_+)^2} x_1 \omega(t, x) dx \geq 0, \quad \frac{d}{dt} \iint_{(\mathbb{R}_+)^2} x_2 \omega(t, x) dx \leq 0.$$

Then, the second step is to prove a lower bound on the kinetic energy $E = \iint_{(\mathbb{R}_+)^2} |u|^2 dx$ of the form

$$E \leq C \left(\iint_{(\mathbb{R}_+)^2} x_2 \omega(t, x) dx \right)^p \left(\frac{d}{dt} \iint_{(\mathbb{R}_+)^2} x_1 \omega(t, x) dx \right)^q$$

³Indeed $\|\omega(t, \cdot)\|_{L^\infty}$ blows up in finite time in all of these cases.

⁴For instance, [16, 50] for the case of \mathbb{R}^3 do not meet the initial condition (1.2).

with some $p, q > 0$ and $C > 0$ depending on the initial data. Then, using conservation of E and monotonicity of $\iint_{(\mathbb{R}_+)^2} x_2 \omega(t, x) dx$, we conclude that

$$\left(\iint_{(\mathbb{R}_+)^2} x_2 \omega_0(x) dx \right)^{-p} \frac{E}{C} \leq \left(\frac{d}{dt} \iint_{(\mathbb{R}_+)^2} x_1 \omega(t, x) dx \right)^q,$$

from which (1.7) follows.

While we follow the above steps, there are some difficulties arising in our case due to the form of the axisymmetric Biot–Savart law and the presence of vortex stretching.

- **Monotonicity lemmas:** It turns out that in the axisymmetric case, the monotone quantities are

$$P(t) = \iint_{(\mathbb{R}_+)^2} -r^2 \omega(t, r, z) dr dz, \quad Z(t) = \iint_{(\mathbb{R}_+)^2} -z \omega(t, r, z) dr dz;$$

the former increases monotonically in time while the later decreases. While this tendency of vortex dynamics is clear from numerous simulations and computations, we were not able to find a rigorous proof in the literature. In the meanwhile, this confirms that the “rebound effect” of vortex rings is a purely viscous phenomenon as argued in [10]; the convection term only brings the antiparallel rings towards each other. Unlike the two-dimensional case, the proof of

$$\frac{d}{dt} \iint_{(\mathbb{R}_+)^2} -z \omega(t, r, z) dr dz \leq 0$$

is highly nontrivial and we could establish it only by using the formulation of the Biot–Savart law in terms of certain elliptic integrals (see Lemma 3.3).

- **Kinetic energy bound:** Following the ideas of Iftimie–Sideris–Gamblin, we try to bound E using $\dot{P} = \frac{d}{dt} P$ and other conserved quantities. However, inspecting the expressions for E and \dot{P} given in Lemma 3.2, it seems impossible to obtain a bound of the form $E \lesssim (\dot{P})^a$ for some $a > 0$, unlike the 2D case. Indeed, E grows faster than \dot{P} if one places the initial vorticity further away from the axis, keeping the L^p norms of $r^{-1} \omega_0$ under control. To handle this issue, we use P itself, which controls the mass of vorticity far from the axis: our key estimate is

$$E \lesssim \dot{P}^{1-\frac{1}{q}} P^{\frac{1}{q}},$$

for any $q < \frac{15}{13}$. Integrating this in time gives the bound $P(t) \gtrsim 1 + t^{\frac{2}{15}-}$.

- **Relation between P and vorticity maximum:** Growth of P implies that, in some averaged sense, the center of vorticity in the radial direction increases. Recalling the Cauchy formula for axisymmetric flows without swirl, this implies growth of the vorticity maximum. Unfortunately, to deduce $\|\omega(t, \cdot)\|_{L^\infty} \rightarrow \infty$ from $P(t) \rightarrow \infty$, we need an additional assumption on the distribution function of the initial vorticity, which is expressed in (1.4): heuristically, the measure of the set on which the vorticity takes very small values cannot be too large. Under this assumption (1.4) with $\delta = 0$, which can be easily achieved by patch-type vorticities, we obtain

$$P^{\frac{1}{2}} \lesssim \|\omega\|_{L^\infty}.$$

This implies $\|\omega(t, \cdot)\|_{L^\infty} \gtrsim t^{\frac{1}{15}-}$. We note that the condition (1.4) with $\delta = 0$ (and $p = \infty$) can be satisfied for some C^γ -data ω_0 whenever $\gamma < 1$ while it fails for typical C^1 -data ω_0 . A little room for $\delta > 0$ in the analysis of Lemma 3.1 provides the existence of Lipschitz data ω_0 with infinite growth.

- **Upgrading growth rate by contradiction:** We perform an additional estimate which enhances the growth rate of $\|\omega(t, \cdot)\|_{L^\infty}$. The idea is to *assume* a hypothetical bound of the form $\|\omega(t, \cdot)\|_{L^\infty} \lesssim t^\beta$, which gives $\|\omega(t, \cdot)\|_{L^\infty} \lesssim P(t)^{\frac{15\beta}{2}+}$ using the lower bound $P(t) \gtrsim 1 + t^{\frac{2}{15}-}$. Then, we can use this new estimate to obtain an improved bound on the kinetic energy:

$$E \lesssim \dot{P}^{1-\frac{1}{q}} P^{(\frac{1}{2} + \frac{45}{56}\beta-)\frac{1}{q}},$$

for any $q < \frac{17}{14}$. This time, we obtain

$$\|\omega(t, \cdot)\|_{L^\infty} \gtrsim P^{\frac{1}{2}} \gtrsim 1 + t^{\frac{1}{\frac{5}{3} + \frac{15}{8}\beta}-}.$$

This is a contradiction to $\|\omega(t, \cdot)\|_{L^\infty} \lesssim t^\beta$ when $\beta < \beta_0 = \frac{1}{45}(\sqrt{670} - 20) \simeq 0.13$.

Remark 1.8. We do not believe that the above growth rate is sharp. One possible improvement in the growth rate could come from showing that $Z(t) \rightarrow 0$ as $t \rightarrow \infty$.

1.4 Notation and conventions

We shall write $\Pi = \{(r, z) : r \geq 0, z \in \mathbb{R}\}$ and $\Pi_+ = \{(r, z) : r, z \geq 0\}$. Similarly, $\mathbb{R}_+^3 = \{x \in \mathbb{R}^3 : x_3 \geq 0\}$. As usual, C will denote various positive absolute constants whose value could vary from a line to another. When C depends on a few parameters, we will sometimes express the dependence using subscripts. Unless otherwise specified, the L^p norms are always taken with respect to the Lebesgue measure on \mathbb{R}^3 .

2 Preliminaries

2.1 Elliptic integrals

It will be convenient to introduce some elliptic integral functions in order to simplify the axisymmetric Biot–Savart law. We begin with

$$\begin{aligned} \mathcal{F}(s) &:= \int_0^\pi \frac{\cos(\theta)}{(2(1 - \cos(\theta)) + s)^{\frac{1}{2}}} d\theta \\ &= \frac{1}{(4+s)^{\frac{1}{2}}} \left((2+s)\mathcal{E}_K\left(\frac{4}{4+s}\right) - (4+s)\mathcal{E}_E\left(\frac{4}{4+s}\right) \right). \end{aligned}$$

Then,

$$\begin{aligned} \mathcal{F}'(s) &= - \int_0^\pi \frac{\cos(\theta)}{2(2(1 - \cos(\theta)) + s)^{\frac{3}{2}}} d\theta \\ &= \frac{1}{2s(4+s)^{\frac{1}{2}}} \left(s\mathcal{E}_K\left(\frac{4}{4+s}\right) - (2+s)\mathcal{E}_E\left(\frac{4}{4+s}\right) \right) \end{aligned} \tag{2.1}$$

and

$$\begin{aligned}\mathcal{F}''(s) &= \int_0^\pi \frac{3 \cos(\theta)}{4(2(1 - \cos(\theta)) + s)^{\frac{3}{2}}} d\theta \\ &= \frac{1}{4s^2(4 + s)^{\frac{3}{2}}} \left(-s(2 + s)\mathcal{E}_K\left(\frac{4}{4 + s}\right) + (16 + 4s + s^2)\mathcal{E}_E\left(\frac{4}{4 + s}\right) \right).\end{aligned}\tag{2.2}$$

Here, the elliptic integrals are defined by

$$\mathcal{E}_K(\alpha) = \int_0^1 (1 - t^2)^{-\frac{1}{2}}(1 - \alpha t^2)^{-\frac{1}{2}} dt,$$

$$\mathcal{E}_E(\alpha) = \int_0^1 (1 - t^2)^{-\frac{1}{2}}(1 - \alpha t^2)^{\frac{1}{2}} dt$$

for $0 \leq \alpha < 1$. The integrals \mathcal{E}_K and \mathcal{E}_E are usually referred to as the *complete elliptic integrals* of the first and second kind, respectively. We note the relations

$$\mathcal{E}'_E(\alpha) = \frac{1}{2\alpha} (\mathcal{E}_E(\alpha) - \mathcal{E}_K(\alpha)), \quad \mathcal{E}'_K(\alpha) = \frac{1}{2\alpha(1 - \alpha)} (\mathcal{E}_E(\alpha) - (1 - \alpha)\mathcal{E}_K(\alpha)),$$

which can be used to verify (2.1) and (2.2).

2.2 Stream function and Biot–Savart law

The stream function for axisymmetric no-swirl flows has the form

$$\psi(r, z) = \frac{1}{2\pi} \iint_{\Pi} \left[\int_0^\pi \frac{r\bar{r} \cos(\theta)}{\sqrt{r^2 - 2r\bar{r} \cos(\theta) + \bar{r}^2 + (z - \bar{z})^2}} d\theta \right] \omega(\bar{r}, \bar{z}) d\bar{r}d\bar{z}.\tag{2.3}$$

Following the notation of Feng–Sverak [20], we define

$$\zeta(r, \bar{r}, z, \bar{z}) := \left(\frac{(r - \bar{r})^2 + (z - \bar{z})^2}{r\bar{r}} \right)^{\frac{1}{2}}.$$

Then, one may perform the θ -integration in (2.3) and obtain a simple expression for ψ :

$$\psi(r, z) = \iint_{\Pi} G(r, \bar{r}, z, \bar{z}) \omega(\bar{r}, \bar{z}) d\bar{r}d\bar{z},$$

with

$$G(r, \bar{r}, z, \bar{z}) := \frac{(r\bar{r})^{\frac{1}{2}}}{2\pi} \mathcal{F}(\zeta^2(r, \bar{r}, z, \bar{z})).$$

With this stream function, the velocity is given by

$$u^r(r, z) = -\frac{1}{r} \partial_z \psi(r, z), \quad u^z(r, z) = \frac{1}{r} \partial_r \psi(r, z).\tag{2.4}$$

One may check directly with (2.4) that

$$\partial_r(ru^r) + \partial_z(ru^z) = 0, \quad \omega = -\partial_r u^z + \partial_z u^r.$$

That is, $u = (u^r, u^z)$ indeed defines an incompressible vector field in \mathbb{R}^3 with associated curl ω . Differentiating G , we obtain convenient integral expressions

$$u^r(r, z) = \iint_{\Pi} \frac{\bar{z} - z}{\pi r^{\frac{3}{2}} \bar{r}^{\frac{1}{2}}} \mathcal{F}'(\zeta^2) \omega(\bar{r}, \bar{z}) \, d\bar{r} d\bar{z},$$

$$u^z(r, z) = \iint_{\Pi} \left(\frac{r - \bar{r}}{\pi r^{\frac{3}{2}} \bar{r}^{\frac{1}{2}}} \mathcal{F}'(\zeta^2) + \frac{\bar{r}^{\frac{1}{2}}}{4\pi r^{\frac{3}{2}}} [\mathcal{F}(\zeta^2) - 2\zeta^2 \mathcal{F}'(\zeta^2)] \right) \omega(\bar{r}, \bar{z}) \, d\bar{r} d\bar{z}.$$

We note that the energy of the flow is given simply by

$$E[\omega] = \pi \iint_{\Pi} \psi(r, z) \omega(r, z) \, dr dz = \frac{1}{2} \iiint_{\Pi^2} (r\bar{r})^{\frac{1}{2}} \mathcal{F}(\zeta^2) \omega(r, z) \omega(\bar{r}, \bar{z}) \, dr dz d\bar{r} d\bar{z}.$$

We remark that $0 \leq E < \infty$ for $r^{-1}\omega \in L^1 \cap L^\infty(\mathbb{R}^3)$ with $r\omega \in L^1(\mathbb{R}^3)$. In this case, the quantity $E[\omega]$ represents the kinetic energy of the flow (after integration by parts):

$$E = \frac{1}{2} \iiint_{\mathbb{R}^3} |u|^2 \, dx$$

(*e.g.* see Lemmas 2.3, 2.4 of [8]).

2.3 Axisymmetric flows without swirl

Let us briefly review the well-posedness theory for axisymmetric flows without swirl. A standard reference is [38].

Global existence and uniqueness with $L^1 \cap L^\infty$. As we have mentioned earlier, there is a unique global-in-time solution to (1.1) under the assumption $\omega_0, r^{-1}\omega_0 \in L^1 \cap L^\infty(\mathbb{R}^3)$. This is a natural extension of Yudovich theory for two-dimensional Euler. The analogy is apparent if one introduces the *relative* vorticity function

$$\xi(t, r, z) = r^{-1}\omega(t, r, z) :$$

(1.1) is simply

$$\partial_t \xi + u \cdot \nabla \xi = 0.$$

Since the velocity is incompressible, formally we have $\|\xi(t, \cdot)\|_{L^p} = \|\xi_0\|_{L^p}$ for all $1 \leq p \leq \infty$. Then, using the estimate (*e.g.* see Lemma 2 of [11])

$$\|r^{-1}u^r\|_{L^\infty} \leq C\|\xi\|_{L^1 \cap L^\infty} = C\|\xi_0\|_{L^1 \cap L^\infty},$$

we obtain a priori estimates for the L^p -norms of the vorticity:

$$\frac{d}{dt} \|\omega\|_{L^p} \leq C_p \|\xi_0\|_{L^1 \cap L^\infty} \|\omega\|_{L^p},$$

$$\|\omega(t, \cdot)\|_{L^p} \leq \|\omega_0\|_{L^p} \exp(Ct\|\xi_0\|_{L^1 \cap L^\infty}).$$

In particular, we see that under the assumption $\xi_0 \in L^1 \cap L^\infty$, higher regularity of the vorticity propagates in time, using the Beale–Kato–Majda criterion ([1]). For global existence and uniqueness, the assumption $\omega_0, \xi_0 \in L^1 \cap L^\infty(\mathbb{R}^3)$ can be weakened; see [11] for instance.

Flow map. Given a solution in the class $\omega, \xi \in L_{t,loc}^\infty(L^1 \cap L^\infty)(\mathbb{R}^3)$, we shall denote $\Phi(t, \cdot)$ to be the associated flow map, which is defined by the unique solution to the ODE

$$\frac{d}{dt}\Phi(t, x) = u(t, \Phi(t, x)), \quad \Phi(0, x) = x$$

(for solvability, *e.g.* see Section 2.4 of [9]). Then, along the flow, we have the important Cauchy formula

$$\frac{\omega(t, \Phi(t, x))}{\Phi^r(t, x)} = \frac{\omega_0(x)}{r} \quad (2.5)$$

which follows from the conservation of ξ along Φ . Here, $\Phi = (\Phi^r, \Phi^z)$ in cylindrical coordinates.

A priori estimate. For later use, we recall a simple estimate.

Lemma 2.1. *Define*

$$X(r, \bar{r}, z, \bar{z}, \theta) := r^2 + \bar{r}^2 - 2r\bar{r}\cos(\theta) + (z - \bar{z})^2.$$

For any $q \in [0, 3)$, we have

$$\sup_{(\bar{r}, \bar{z}) \in \Pi} \iint_{\Pi} \int_0^{2\pi} X^{-\frac{q}{2}} d\theta |\omega(r, z)| dr dz \leq C_q \|r^{-1}\omega\|_{L^1(\mathbb{R}^3)}^{1-\frac{q}{3}} \|r^{-1}\omega\|_{L^\infty(\mathbb{R}^3)}^{\frac{q}{3}}$$

with a constant $C_q > 0$ depending only on q .

Proof. We note that for $x = (r, 0, z)$ and $\bar{x} = (\bar{r}, \theta, \bar{z})$, we have

$$\iint_{\Pi} \int_0^{2\pi} X^{-\frac{q}{2}} d\theta |\omega(r, z)| dr dz = \iiint_{\mathbb{R}^3} \frac{1}{|\bar{x} - x|^q} \frac{|\omega(x)|}{r} dx.$$

The statement is clear when $q = 0$. For $0 < q < 3$, we consider the regions $\{|x - \bar{x}| \leq L\}$ and $\{|x - \bar{x}| > L\}$, for some $L > 0$ to be determined. In the latter region, we have that

$$\iiint_{|x - \bar{x}| > L} \frac{1}{|\bar{x} - x|^q} \frac{|\omega(x)|}{r} dx \leq CL^{-q} \|r^{-1}\omega\|_{L^1}$$

since $q > 0$ and in the former, we obtain

$$\iiint_{|x - \bar{x}| \leq L} \frac{1}{|\bar{x} - x|^q} \frac{|\omega(x)|}{r} dx \leq \|r^{-1}\omega\|_{L^\infty} \iiint_{|x| \leq L} |x|^{-q} dx \leq C_q L^{3-q} \|r^{-1}\omega\|_{L^\infty}$$

since $q < 3$. The choice

$$L = \left(\frac{\|r^{-1}\omega\|_{L^1}}{\|r^{-1}\omega\|_{L^\infty}} \right)^{\frac{1}{3}}$$

gives the desired estimate. □

The t^2 bound on the vorticity maximum. We provide a proof that the vorticity maximum for general axisymmetric flows without swirl cannot grow faster than t^2 in time, assuming that the initial vorticity satisfies

$$\sup\{r : (r, z) \in \text{supp}(\omega_0(\cdot))\} < \infty$$

together with (1.2). We need the following lemma from Feng–Sverak [20].

Lemma 2.2 ([20, Propositions 2.11, 2.13]). *The velocity $u = K[\omega]$ satisfies*

$$\|u\|_{L^\infty} \leq C \|r\omega\|_{L^1}^{\frac{1}{4}} \|r^{-1}\omega\|_{L^1}^{\frac{1}{4}} \|r^{-1}\omega\|_{L^\infty}^{\frac{1}{2}}.$$

Recall our convention that the L^p norms are taken always with respect to the Lebesgue measure on \mathbb{R}^3 . Now to prove the claim, we define $R(t) = \sup\{r : (r, z) \in \text{supp}(\omega(t, \cdot))\}$. We may assume that $R(t)$ is increasing in time by redefining it to be $\sup_{t' \leq t} R(t')$. Then

$$\begin{aligned} \frac{d}{dt} R(t) &\leq \|u^r\|_{L^\infty} \leq C \|r\omega\|_{L^1}^{\frac{1}{4}} \|r^{-1}\omega\|_{L^1}^{\frac{1}{4}} \|r^{-1}\omega\|_{L^\infty}^{\frac{1}{2}} \\ &\leq CR(t)^{\frac{1}{2}} \|r^{-1}\omega\|_{L^1}^{\frac{1}{2}} \|r^{-1}\omega\|_{L^\infty}^{\frac{1}{2}} \\ &= CR(t)^{\frac{1}{2}} \|r^{-1}\omega_0\|_{L^1}^{\frac{1}{2}} \|r^{-1}\omega_0\|_{L^\infty}^{\frac{1}{2}}. \end{aligned}$$

Integrating in time gives

$$R(t) \leq C(1+t)^2. \tag{2.6}$$

Recalling the Cauchy formula (2.5), the t^2 -bound on the support can be translated to a bound on $\|\omega(t, \cdot)\|_{L^\infty}$:

$$\|\omega(t, \cdot)\|_{L^\infty} \leq R(t) \|r^{-1}\omega_0\|_{L^\infty} \leq C(1+t)^2,$$

where $C > 0$ depends on the initial data ω_0 .

3 Key Inequalities

From now on, we shall impose the assumption (1.3) on ω , so that the dynamics reduces to $\Pi_+ = \{(r, z) : r, z \geq 0\}$.

3.1 Relation between P and vorticity L^p norms

We define

$$P(t) := \iint_{\Pi_+} -r^2 \omega(t, r, z) \, dr \, dz = \iiint_{\mathbb{R}_+^3} -r^2 \xi(t, \cdot) \, dx \geq 0.$$

Lemma 3.1. *Let $0 \leq \delta < 1$. Assume that $\|\xi_0^{-1} \mathbf{1}_{\{\xi_0 < 0\}}\|_{L^{\frac{1-\delta}{1-((2-\delta)/p)}}(\mathbb{R}_+^3)} < \infty$ for some $p \in [2-\delta, \infty]$ and $R_0 = \sup\{r : (r, z) \in \text{supp}(\omega_0(\cdot))\} < \infty$. Then, we have*

$$P(t) \leq R(t)^\delta \|\omega(t, \cdot)\|_{L^p(\mathbb{R}_+^3)}^{2-\delta} \|\xi_0^{-1} \mathbf{1}_{\{\xi_0 < 0\}}\|_{L^{\frac{1-\delta}{1-((2-\delta)/p)}}(\mathbb{R}_+^3)}^{1-\delta},$$

where $R(t) = \sup\{r : (r, z) \in \text{supp}(\omega(t, \cdot))\}$.

Proof. We estimate using Hölder's inequality

$$\begin{aligned} \iiint_{\mathbb{R}_+^3} -r^2 \xi \, dx &= \iiint_{\mathbb{R}_+^3} -r^\delta \frac{r^{1-\delta}}{\omega^{1-\delta}} \omega^{2-\delta} \, dx \\ &\leq R(t)^\delta \|\omega(t, \cdot)\|_{L^p(\mathbb{R}_+^3)}^{2-\delta} \left(\iiint_{\{\xi(t, \cdot) < 0\}} \xi(t, \cdot)^{-\frac{1-\delta}{1-((2-\delta)/p)}} \, dx \right)^{1-\frac{(2-\delta)}{p}} \\ &= R(t)^\delta \|\omega(t, \cdot)\|_{L^p(\mathbb{R}_+^3)}^{2-\delta} \|\xi_0^{-1} \mathbf{1}_{\{\xi_0 < 0\}}\|_{L^{\frac{1-\delta}{1-((2-\delta)/p)}}(\mathbb{R}_+^3)}^{1-\delta}. \end{aligned}$$

In the last equality, we have used that the distribution function of $\xi(t, \cdot)$ is invariant in time. \square

It is worth considering the special case in which $\xi_0 = -\lambda \mathbf{1}_{\Omega_0}$ on \mathbb{R}_+^3 for some $\lambda > 0$ and a bounded open set $\Omega_0 \subset \Pi_+ \cap \{r \geq \frac{1}{2}\}$. The associated solution takes the form

$$\xi(t, \cdot) = -\lambda \mathbf{1}_{\Omega(t)}.$$

Denoting $R(t) := \sup\{r : \text{there exists } (r, z) \in \Omega(t)\}$, we immediately obtain from $|\Omega(t)| = |\Omega_0|$ ($|\cdot|$ denotes the three-dimensional Lebesgue measure) that

$$P(t) \leq \lambda |\Omega_0| R^2(t),$$

and combining this with

$$\|\omega(t, \cdot)\|_{L^\infty} \geq \frac{1}{2} \lambda R(t),$$

we arrive at the following precise result:

$$\|\omega(t, \cdot)\|_{L^\infty} \geq \frac{1}{2} (|\Omega_0|^{-1} \|\xi_0\|_{L^\infty} P(t))^{\frac{1}{2}}.$$

3.2 Estimates on kinetic energy and derivative of P

In addition to

$$X(r, \bar{r}, z, \bar{z}, \theta) = r^2 + \bar{r}^2 - 2r\bar{r} \cos(\theta) + (z - \bar{z})^2$$

which has been defined in the above, we set

$$\begin{aligned} \bar{X}(r, \bar{r}, z, \bar{z}, \theta) &= r^2 + \bar{r}^2 + 2r\bar{r} \cos(\theta) + (z - \bar{z})^2, \\ Y(r, \bar{r}, z, \bar{z}, \theta) &= r^2 + \bar{r}^2 - 2r\bar{r} \cos(\theta) + (z + \bar{z})^2, \\ \bar{Y}(r, \bar{r}, z, \bar{z}, \theta) &= r^2 + \bar{r}^2 + 2r\bar{r} \cos(\theta) + (z + \bar{z})^2. \end{aligned}$$

For ease of notation, let us write $\omega = \omega(r, z) \, dr \, dz$ and $\bar{\omega} = \omega(\bar{r}, \bar{z}) \, d\bar{r} \, d\bar{z}$.

Lemma 3.2. *The kinetic energy $E \geq 0$ satisfies*

$$\frac{E}{C} \leq \iiint\iiint_{(\Pi_+)^2} \left[\int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 (z\bar{z})(\cos(\theta))^2}{X^{\frac{1}{2}}\bar{X}Y} d\theta \right] \omega\bar{\omega} \leq CE. \quad (3.1)$$

Moreover, $\dot{P} := \frac{d}{dt}P \geq 0$ satisfies

$$\frac{\dot{P}}{C} \leq \iiint\iiint_{(\Pi_+)^2} \left[\int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 (z + \bar{z})(\cos(\theta))^2}{Y^{\frac{3}{2}}\bar{Y}} d\theta \right] \omega\bar{\omega} \leq C\dot{P}. \quad (3.2)$$

Here, $C > 0$ is an absolute constant.

Proof. We begin with the expression

$$\begin{aligned} E &= \frac{1}{2} \iiint\iiint_{\Pi^2} \left[\int_0^{\pi} \frac{r\bar{r} \cos(\theta)}{X^{\frac{1}{2}}} d\theta \right] \omega\bar{\omega} = \frac{1}{2} \iiint\iiint_{\Pi^2} \left[\int_0^{\frac{\pi}{2}} r\bar{r} \cos(\theta) \left(\frac{1}{X^{\frac{1}{2}}} - \frac{1}{\bar{X}^{\frac{1}{2}}} \right) d\theta \right] \omega\bar{\omega} \\ &= 2 \iiint\iiint_{\Pi^2} \left[\int_0^{\frac{\pi}{2}} \frac{(r\bar{r} \cos(\theta))^2}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})} d\theta \right] \omega\bar{\omega}. \end{aligned}$$

Next, using odd symmetry of ω and $\bar{\omega}$ in z ,

$$E = 4 \iiint\iiint_{(\Pi_+)^2} \left[\int_0^{\frac{\pi}{2}} (r\bar{r} \cos(\theta))^2 \left(\frac{1}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})} - \frac{1}{Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} \right) d\theta \right] \omega\bar{\omega}.$$

We compute

$$\begin{aligned} &\frac{1}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})} - \frac{1}{Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} \\ &= \frac{(Y\bar{Y})^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}} - X^{\frac{1}{2}} - \bar{X}^{\frac{1}{2}}) + ((Y\bar{Y})^{\frac{1}{2}} - (X\bar{X})^{\frac{1}{2}})(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} =: I + II. \end{aligned}$$

We consider

$$II = \frac{(Y\bar{Y})^{\frac{1}{2}} - (X\bar{X})^{\frac{1}{2}}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} = \frac{Y\bar{Y} - X\bar{X}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})((Y\bar{Y})^{\frac{1}{2}} + (X\bar{X})^{\frac{1}{2}})}.$$

Noting that

$$Y\bar{Y} - X\bar{X} = 2z\bar{z}(X + \bar{X} + Y + \bar{Y})$$

and

$$X \leq \bar{X}, \quad Y \leq \bar{Y}, \quad X \leq Y, \quad \bar{X} \leq \bar{Y},$$

we observe that

$$Y\bar{Y} - X\bar{X} \sim z\bar{z}\bar{Y}, \quad X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})((Y\bar{Y})^{\frac{1}{2}} + (X\bar{X})^{\frac{1}{2}}) \sim X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y\bar{Y})^{\frac{1}{2}}.$$

Here, we write $A \sim B$ if there is an absolute constant $C > 0$ such that $C^{-1}A \leq B \leq CA$. This gives

$$II \sim \frac{z\bar{z}\bar{Y}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}(Y\bar{Y})^{\frac{1}{2}}} \sim \frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}(Y\bar{Y})^{\frac{1}{2}}}.$$

Next,

$$I = \frac{Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}} - X^{\frac{1}{2}} - \bar{X}^{\frac{1}{2}}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} =: I_1 + I_2$$

with

$$I_1 = \frac{Y^{\frac{1}{2}} - X^{\frac{1}{2}}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})} = \frac{4z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}(X^{\frac{1}{2}} + \bar{X}^{\frac{1}{2}})(Y^{\frac{1}{2}} + \bar{Y}^{\frac{1}{2}})(Y^{\frac{1}{2}} + X^{\frac{1}{2}})} \sim \frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}}$$

Similarly,

$$I_2 \sim \frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}} \lesssim I_1.$$

Since

$$\frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}\bar{Y}^{\frac{1}{2}}} + \frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}^{\frac{1}{2}}Y^{\frac{1}{2}}(Y\bar{Y})^{\frac{1}{2}}} \sim \frac{z\bar{z}}{X^{\frac{1}{2}}\bar{X}Y},$$

we obtain (3.1).

Next, we compute

$$\begin{aligned} \dot{P} &= - \iint_{\Pi_+} 2ru^r(r, z)\omega(r, z) dr dz \\ &= -\frac{1}{\pi} \iint_{\Pi} \iint_{\Pi_+} \left[\int_0^\pi \frac{(z - \bar{z})r\bar{r} \cos(\theta)}{X^{\frac{3}{2}}} d\theta \right] \omega \bar{\omega} \\ &= \frac{1}{\pi} \iiint_{(\Pi_+)^2} \left[\int_0^\pi \frac{(z + \bar{z})r\bar{r} \cos(\theta)}{Y^{\frac{3}{2}}} d\theta \right] \omega \bar{\omega} \\ &= \frac{4}{\pi} \iiint_{(\Pi_+)^2} \left[\int_0^{\frac{\pi}{2}} (z + \bar{z})(r\bar{r})^2 (\cos(\theta))^2 \frac{\bar{Y}^2 + \bar{Y}Y + Y^2}{(Y\bar{Y})^{\frac{3}{2}}(Y^{\frac{3}{2}} + \bar{Y}^{\frac{3}{2}})} d\theta \right] \omega \bar{\omega}. \end{aligned}$$

Then, we simply note that

$$\bar{Y}^2 + \bar{Y}Y + Y^2 \sim \bar{Y}^2, \quad (Y\bar{Y})^{\frac{3}{2}}(Y^{\frac{3}{2}} + \bar{Y}^{\frac{3}{2}}) \sim Y^{\frac{3}{2}}\bar{Y}^3$$

to derive

$$\dot{P} \sim \iiint_{(\Pi_+)^2} \left[\int_0^{\frac{\pi}{2}} \frac{(z + \bar{z})(r\bar{r})^2 (\cos(\theta))^2}{Y^{\frac{3}{2}}\bar{Y}} d\theta \right] \omega \bar{\omega}.$$

This finishes the proof. □

3.3 Monotonicity Lemma

For simplicity, we shall denote

$$Z(t) := \iint_{\Pi_+} -z\omega(t, r, z) \, drdz.$$

We have already seen that

$$P(t) = \iint_{\Pi_+} -r^2\omega(t, r, z) \, drdz$$

is monotone increasing with t . In the meanwhile, we note that $0 \leq Z, P < +\infty$ for $t > 0$ if they are finite at $t = 0$.

Lemma 3.3. *We have*

$$\dot{Z} := \frac{d}{dt} \iint_{\Pi_+} -z\omega(t, r, z) \, drdz < 0. \quad (3.3)$$

Proof of Lemma 3.3. We begin with

$$\frac{d}{dt} \iint_{\Pi_+} z\omega \, drdz = \iint_{\Pi_+} u^z \omega \, drdz.$$

Symmetrizing the kernel of u^z in $(r, z) \leftrightarrow (\bar{r}, \bar{z})$, we obtain with

$$\mathcal{H}[r, \bar{r}, z, \bar{z}] := -\frac{(r - \bar{r})^2}{2\pi r^{\frac{3}{2}} \bar{r}^{\frac{3}{2}}} \mathcal{F}'(\zeta^2) + \frac{\bar{r}^2 + r^2}{8\pi r^{\frac{3}{2}} \bar{r}^{\frac{3}{2}}} [\mathcal{F}(\zeta^2) - 2\zeta^2 \mathcal{F}'(\zeta^2)]$$

that

$$\iint_{\Pi_+} u^z \omega \, drdz = \iint_{\Pi_+} \iint_{\Pi_+} [\mathcal{H}[r, \bar{r}, z, \bar{z}] - \mathcal{H}[r, \bar{r}, z, -\bar{z}]] \omega(r, z) \omega(\bar{r}, \bar{z}) \, drdzd\bar{r}d\bar{z}.$$

We need to show that $\mathcal{H}[r, \bar{r}, z, \bar{z}] - \mathcal{H}[r, \bar{r}, z, -\bar{z}] > 0$, but since the dependence of \mathcal{H} in z, \bar{z} is only through $(z - \bar{z})^2$ in ζ^2 , it suffices to show that $-\mathcal{F}'(s)$ and $\mathcal{F}(s) - 2s\mathcal{F}'(s)$ are strictly *decreasing* functions of s for all $s \in \mathbb{R}_+$.

(i) $\mathcal{F}'(s)$. We need to check $\mathcal{F}'' > 0$. However, this is clear from the integral representation in (2.2) after symmetrizing θ with $\pi - \theta$.

(ii) $\mathcal{F}(s) - 2s\mathcal{F}'(s)$. We need to prove that

$$(\mathcal{F}(s) - 2s\mathcal{F}'(s))' = -\mathcal{F}'(s) - 2s\mathcal{F}''(s) < 0.$$

Using (2.1) and (2.2), we explicitly check that

$$\mathcal{F}'(s) + 2s\mathcal{F}''(s) = \frac{1}{s(4+s)^{\frac{3}{2}}} \left(s\mathcal{E}_K \left(\frac{4}{4+s} \right) + (4-s)\mathcal{E}_E \left(\frac{4}{4+s} \right) \right)$$

but

$$s\mathcal{E}_K \left(\frac{4}{4+s} \right) + (4-s)\mathcal{E}_E \left(\frac{4}{4+s} \right) = 4\mathcal{E}_E \left(\frac{4}{4+s} \right) + s \left(\mathcal{E}_K \left(\frac{4}{4+s} \right) - \mathcal{E}_E \left(\frac{4}{4+s} \right) \right) > 0$$

since $\mathcal{E}_E > 0$ and $\mathcal{E}_K > \mathcal{E}_E$. This finishes the proof of (3.3). \square

4 Growth

4.1 Preliminary growth

We are now in a position to state and prove the first key proposition.

Proposition 4.1. *We have the following lower bound for all $t \geq 0$:*

$$P(t) \geq \left(C_q (E_0 X_0^{-1})^{\frac{q}{q-1}} t + P_0^{\frac{q}{q-1}} \right)^{\frac{q-1}{q}}, \quad 1 < q < \frac{15}{13},$$

where $C_q > 0$ is a constant depending only on q , E_0 is the kinetic energy, and

$$X_0 = \left(\|\xi_0\|_{L^1}^{4-\frac{10}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{4(q-1)} P_0^{1-q} + \|\xi_0\|_{L^1}^{3-\frac{7}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{2(q-1)} \right)^{\frac{1}{q}}$$

is a constant depending only on the initial data.

Note that the statements of Theorem 1.1 follows from the above proposition. Then, Corollary 1.2 follows from Lemma 3.1 thanks to t^2 -bound (2.6) on support.

Proof. To begin with, we recall the bounds (3.1), (3.2) on E and P from Lemma 3.2. Then, we estimate using Hölder's inequality

$$\begin{aligned} E &\lesssim \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 (z\bar{z})(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X} Y} d\theta \omega \bar{\omega} \\ &\lesssim \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 (z+\bar{z})(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X} Y^{\frac{1}{2}}} d\theta \omega \bar{\omega} \\ &\lesssim (\dot{P})^{1-\frac{1}{q}} J^{\frac{1}{q}} \end{aligned} \tag{4.1}$$

for some $1 < q \leq \frac{3}{2}$ to be determined below, where J is given by

$$\begin{aligned} J &= \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 (z+\bar{z}) \bar{Y}^{q-1} (\cos(\theta))^2}{(X \bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega} \\ &= \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{2(r\bar{r})^2 \bar{z} \bar{Y}^{q-1} (\cos(\theta))^2}{(X \bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega}. \end{aligned}$$

We have used the lower bound of P from Lemma 3.2. We now estimate

$$J \leq C(J_1 + J_2) \tag{4.2}$$

using

$$\bar{Y}^{q-1} \leq C(\bar{X}^{q-1} + Y^{q-1})$$

where

$$J_1 = \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 \bar{z} \bar{X}^{q-1} (\cos(\theta))^2}{(X \bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega}, \quad J_2 = \iiint\limits_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 \bar{z} Y^{q-1} (\cos(\theta))^2}{(X \bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega}. \tag{4.3}$$

Estimate of J_1 . We begin with rewriting it as

$$J_1 = 2 \iint_{\Pi_+} r^2 |\omega(r, z)| Q_1(r, z) dr dz,$$

where

$$\begin{aligned} Q_1(r, z) &:= \iint_{\Pi_+} \int_0^{\frac{\pi}{2}} \frac{\bar{r}^2 \bar{z} (\cos(\theta))^2}{X^{\frac{q}{2}} \bar{X} Y^{\frac{3}{2}-q}} d\theta |\bar{\omega}| \\ &\lesssim \iint_{\Pi_+} \int_0^{\frac{\pi}{2}} \frac{\bar{z}^{2(q-1)} (\cos(\theta))^2}{X^{\frac{q}{2}}} d\theta |\bar{\omega}| \\ &\lesssim \left(\iint_{\Pi_+} \int_0^{\frac{\pi}{2}} \frac{1}{X^{\frac{q}{2(3-2q)}}} d\theta |\bar{\omega}| \right)^{3-2q} \left(\iint_{\Pi_+} \bar{z} |\bar{\omega}| \right)^{2(q-1)}. \end{aligned}$$

When $q < \frac{9}{7}$, we obtain that

$$\|Q_1(r, z)\|_{L^\infty(\Pi_+)} \leq C \|\xi_0\|_{L^1}^{3-\frac{7}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{2(q-1)}, \quad J_1 \leq C \|\xi_0\|_{L^1}^{3-\frac{7}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{2(q-1)} P.$$

Estimate of J_2 . We first bound

$$\begin{aligned} J_2 &= \iiint_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 \bar{z} Y^{q-1} (\cos(\theta))^2}{(X\bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega} \\ &\lesssim \iiint_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{r^{4-2q} \bar{z}^{4(q-1)} (\cos(\theta))^2}{X^{\frac{q}{2}}} d\theta \omega \bar{\omega}. \end{aligned}$$

Let us now estimate

$$\begin{aligned} Q_2(r, z) &:= \iint_{\Pi_+} \int_0^{\frac{\pi}{2}} \frac{\bar{z}^{4(q-1)} (\cos(\theta))^2}{X^{\frac{q}{2}}} d\theta |\bar{\omega}| \\ &\lesssim \left(\iint_{\Pi_+} \int_0^{\frac{\pi}{2}} \frac{1}{X^{\frac{q}{2(5-4q)}}} d\theta |\bar{\omega}| \right)^{5-4q} \left(\iint_{\Pi_+} \bar{z} |\bar{\omega}| \right)^{4(q-1)} \\ &\lesssim \|\xi_0\|_{L^1}^{5-\frac{13}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{4(q-1)} \end{aligned}$$

This requires $q < \frac{15}{13}$. Then using Hölder's inequality

$$\begin{aligned} J_2 &\lesssim \iint_{\Pi_+} r^{4-2q} |\omega|^{2-q} Q_2(r, z) |\omega|^{q-1} dr dz \\ &\lesssim \|Q_2\|_{L^\infty} P^{2-q} \|\xi_0\|_{L^1}^{q-1} \\ &\lesssim \|\xi_0\|_{L^1}^{5-\frac{13}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{4(q-1)} \|\xi_0\|_{L^1}^{q-1} P^{2-q} \\ &\lesssim \|\xi_0\|_{L^1}^{4-\frac{10}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{4(q-1)} P^{1-q} P. \end{aligned}$$

Combining the estimates for J_1 and J_2 , we arrive at

$$E \leq C X_0 (\dot{P})^{1-\frac{1}{q}} P^{\frac{1}{q}}, \quad 1 < q < \frac{15}{13}.$$

Here, we recall that

$$X_0 = \left(\|\xi_0\|_{L^1}^{4-\frac{10}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{4(q-1)} P_0^{1-q} + \|\xi_0\|_{L^1}^{3-\frac{7}{3}q} \|\xi_0\|_{L^\infty}^{\frac{q}{3}} Z_0^{2(q-1)} \right)^{\frac{1}{q}}.$$

Therefore,

$$E_0^{\frac{q}{q-1}} \leq C X_0^{\frac{q}{q-1}} \frac{d}{dt} \left(P^{\frac{q}{q-1}} \right),$$

or

$$P(t) \geq \left(C(E_0 X_0^{-1})^{\frac{q}{q-1}} t + P_0^{\frac{q}{q-1}} \right)^{\frac{q-1}{q}}.$$

This finishes the proof. \square

4.2 Enhancing the growth rate

In this section, we complete the proof of Theorem 1.6. We shall need a few additional lemmas.

Lemma 4.2. *For any $0 < \tau \leq \frac{3}{2}$, we have*

$$\int_0^{\frac{\pi}{2}} \frac{(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X}} d\theta \leq C_\tau \frac{(r\bar{r})^{\tau-\frac{3}{2}}}{d^{2\tau}}, \quad d := \sqrt{(r-\bar{r})^2 + (z-\bar{z})^2}. \quad (4.4)$$

Proof. This comes from the following estimate of \mathcal{F} (see [21, Lemma 2.7])

$$\mathcal{F}(s) \lesssim_\tau s^{-\tau}, \quad \tau \in (0, 3/2].$$

We plug in $s = d^2/(r\bar{r})$ to obtain

$$\sqrt{r\bar{r}} \int_0^\pi \frac{\cos(\theta)}{X^{\frac{1}{2}}} d\theta \lesssim_\tau \frac{(r\bar{r})^\tau}{d^{2\tau}}.$$

Symmetrizing the left hand side,

$$\sqrt{r\bar{r}} \int_0^\pi \frac{\cos(\theta)}{X^{\frac{1}{2}}} d\theta = \sqrt{r\bar{r}} \int_0^{\frac{\pi}{2}} \left(\frac{\cos(\theta)}{X^{\frac{1}{2}}} - \frac{\cos(\theta)}{\bar{X}^{\frac{1}{2}}} \right) d\theta = 4(r\bar{r})^{\frac{3}{2}} \int_0^{\frac{\pi}{2}} \frac{(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X}} d\theta.$$

This gives the lemma. \square

Lemma 4.3. *For any $0 \leq \alpha < 2$, we have*

$$\sup_{(\bar{r}, \bar{z})} \iint_{\Pi_+} \frac{|\omega|}{d^\alpha} dr dz \leq C_\alpha \|\omega\|_{L^1(\Pi_+)}^{1-\frac{\alpha}{2}} \|\omega\|_{L^\infty(\Pi_+)}^{\frac{\alpha}{2}}.$$

Proof. The proof is completely parallel to that for Lemma 2.1 in the above. See also Lemma 2.1. of [28]. \square

Proof of Theorem 1.6. Towards a contradiction, we assume that there exists $M > 0$ such that

$$\sup_{t \geq 1} \frac{\|\omega(t, \cdot)\|_{L^\infty}}{t^\beta} \leq M.$$

Based on this assumption, we now obtain *improved estimates* on J_1 and J_2 (in (4.3)). **In this proof, implicit constants will depend on the initial data and M .**

Claim. For each $1 < q < \frac{11}{8}$, if $\varepsilon > 0$ is sufficiently small, then

$$J_1 \lesssim P^{\frac{1}{2} + \varepsilon} (1 + t^{\frac{1}{2}(q-1+2\varepsilon)\beta}).$$

Similarly, we have

$$J_2 \lesssim P^{\frac{3}{2}-q+\varepsilon} (1 + t^{\frac{1}{2}(q-1+2\varepsilon)\beta}), \quad 1 < q < \frac{17}{14}.$$

We first check the estimate for J_1 . This time, we bound using (4.4)

$$\begin{aligned} J_1 &= \iiint_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 \bar{z} \bar{X}^{q-1} (\cos(\theta))^2}{(X \bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega} \\ &\lesssim \iiint_{(\Pi_+)^2} (r\bar{r})^2 \bar{z}^{2(q-1)} \frac{1}{d^{q-1}} \int_0^{\frac{\pi}{2}} \frac{(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X}} d\theta \omega \bar{\omega} \\ &\lesssim \iiint_{(\Pi_+)^2} (r\bar{r})^2 \bar{z}^{2(q-1)} \frac{1}{d^{q-1}} \frac{(r\bar{r})^{\tau-\frac{3}{2}}}{d^{2\tau}} \omega \bar{\omega}. \end{aligned}$$

Next, we use Hölder's inequality to bound

$$\begin{aligned} &\iiint_{(\Pi_+)^2} (r\bar{r})^2 \bar{z}^{2(q-1)} \frac{1}{d^{q-1}} \frac{(r\bar{r})^{\tau-\frac{3}{2}}}{d^{2\tau}} \omega \bar{\omega} \\ &\lesssim \left(\iiint_{(\Pi_+)^2} (r\bar{r})^2 \omega \bar{\omega} \right)^{\frac{1}{4}(1+2\tau)} \left(\iint_{\Pi_+} \bar{z}^{\frac{8(q-1)}{3-2\tau}} \bar{\omega} \iint_{\Pi_+} \frac{\omega}{d^{(q-1+2\tau)\frac{4}{3-2\tau}}} \right)^{\frac{3}{4}-\frac{\tau}{2}} \end{aligned}$$

We now use Lemma 4.3: for $\tau > 0$ small and $q < \frac{11}{8}$,

$$\sup_{(\bar{r}, \bar{z}) \in \Pi_+} \iint_{\Pi_+} \frac{|\omega|}{d^{\frac{4(q-1+2\tau)}{3-2\tau}}} \lesssim \|\omega\|_{L^1(\Pi_+)}^{1-\frac{2(q-1+2\tau)}{(3-2\tau)}} (Mt^\beta)^{\frac{2(q-1+2\tau)}{(3-2\tau)}} \lesssim t^{\frac{2(q-1+2\tau)}{(3-2\tau)}\beta}, \quad t \geq 1.$$

Then, as long as $\frac{8}{3}(q-1) < 1$, we can take $\tau > 0$ sufficiently small and estimate

$$\iint_{\Pi_+} \bar{z}^{\frac{8(q-1)}{3-2\tau}} |\bar{\omega}| \lesssim Z_0^{\frac{8(q-1)}{3-2\tau}} \|\omega\|_{L^1(\Pi_+)}^{1-\frac{8(q-1)}{3-2\tau}} \lesssim 1.$$

This gives for $t \geq 1$

$$J_1 \lesssim P^{\frac{1}{2}(1+2\tau)} t^{\frac{q-1+2\tau}{2}\beta}.$$

We now proceed to the estimate of J_2 : using (4.4) and Hölder's inequality,

$$\begin{aligned}
J_2 &= \iiint_{(\Pi_+)^2} \int_0^{\frac{\pi}{2}} \frac{(r\bar{r})^2 \bar{z} Y^{q-1} (\cos(\theta))^2}{(X\bar{X}^2)^{\frac{q}{2}} Y^{\frac{3}{2}-q}} d\theta \omega \bar{\omega} \\
&\lesssim \iiint_{(\Pi_+)^2} \frac{(r\bar{r})^{3-q} \bar{z}^{4(q-1)}}{d^{q-1}} \int_0^{\frac{\pi}{2}} \frac{(\cos(\theta))^2}{X^{\frac{1}{2}} \bar{X}} d\theta \omega \bar{\omega} \\
&\lesssim \iiint_{(\Pi_+)^2} \frac{(r\bar{r})^{\frac{3}{2}-q+\tau} \bar{z}^{4(q-1)}}{d^{q-1+2\tau}} \omega \bar{\omega} \\
&\lesssim \left(\iiint_{(\Pi_+)^2} (r\bar{r})^2 \omega \bar{\omega} \right)^{\frac{1}{2}(\frac{3}{2}-q+\tau)} \left(\iint_{\Pi_+} \frac{16(q-1)}{\bar{z}^{1+2q-2\tau}} \bar{\omega} \iint_{\Pi_+} \frac{\omega}{d^{\frac{4(q-1+2\tau)}{1+2q-2\tau}}} \right)^{\frac{1}{2}(\frac{1}{2}+q-\tau)}.
\end{aligned}$$

As long as $q < \frac{17}{14}$, we can take $\tau > 0$ sufficiently small that

$$\iint_{\Pi_+} \frac{16(q-1)}{\bar{z}^{1+2q-2\tau}} |\bar{\omega}| \lesssim 1, \quad \sup_{(\bar{r}, \bar{z}) \in \Pi_+} \iint_{\Pi_+} \frac{|\omega|}{d^{\frac{4(q-1+2\tau)}{1+2q-2\tau}}} \lesssim t^{\frac{2(q-1+2\tau)}{1+2q-2\tau} \beta}.$$

This gives

$$J_2 \lesssim P^{\frac{3}{2}-q+\tau} t^{\frac{q-1+2\tau}{2} \beta}.$$

This finishes the proof of **Claim**.

In the following, $\epsilon > 0$ will denote a sufficiently small constant whose value could change from a line to another. Given **Claim**, we now observe, for any $1 < q < 17/14$,

$$J \lesssim J_1 + J_2 \lesssim P^{\frac{1}{2}+\epsilon} (1+t)^{\frac{1}{2}(q-1+2\epsilon)\beta}$$

since P is increasing (recall (4.2) and the above for J). Recall from Proposition 4.1 that

$$P \gtrsim (1+t)^{\frac{2}{15}-\epsilon}.$$

Hence, combining the previous two inequalities, we deduce

$$J \lesssim P^{\frac{1}{2}+\frac{15}{4}(q-1)\beta+\epsilon}.$$

Returning to the energy estimate (4.1):

$$E \lesssim \dot{P}^{1-\frac{1}{q}} J^{\frac{1}{q}}$$

and applying the above upper bound for J , we obtain that

$$\dot{P} P^{\frac{(1/2)}{q-1} + \frac{15}{4}\beta+\epsilon} \gtrsim 1,$$

which gives upon integration after taking q close to $17/14$ that

$$P \gtrsim t^{\frac{1}{\frac{10}{3} + \frac{15}{4}\beta} - \epsilon}. \tag{4.5}$$

On the other hand, from the assumption on the vorticity maximum, the estimate (2.6) with Lemma 3.1 implies that

$$P(t) \lesssim R(t)^\delta \|\omega(t, \cdot)\|_{L^\infty(\mathbb{R}_+^3)}^{2-\delta} \lesssim (1+t)^{2\delta} t^{(2-\delta)\beta} \lesssim (1+t)^{2\delta+(2-\delta)\beta}, \quad (4.6)$$

where $R(t) = \sup\{r : (r, z) \in \text{supp}(\omega(t, \cdot))\}$. Therefore, we derive a contradiction to (4.5) once

$$2\delta + (2-\delta)\beta < \frac{1}{\frac{10}{3} + \frac{15}{4}\beta}. \quad (4.7)$$

Note that, for each $\delta \in [0, 3/20)$, we denote the unique positive root of

$$(2\delta + (2-\delta)\beta) \cdot \left(\frac{10}{3} + \frac{15}{4}\beta\right) = 1$$

by $\beta_0 = \beta_0(\delta)$. Observing that the previous inequality (4.7) holds for any $\beta \in (0, \beta_0)$, we are done. \square

Remark 4.4. The explicit value of $\beta_0(\delta)$ is

$$\beta_0(\delta) = \frac{8+5\delta}{9(\delta-2)} + \frac{\sqrt{536-1148\delta+845\delta^2}}{9(2-\delta)\sqrt{5}},$$

which is strictly positive whenever $0 \leq \delta < 3/20$. In particular, it satisfies

$$\beta_0(\delta) > \beta_1(\delta) \quad \text{when} \quad 0 \leq \delta < 1/15,$$

with $\beta_1 = \frac{\frac{2}{15}-2\delta}{2-\delta}$ from Corollary 1.2.

5 Enstrophy growth for Navier–Stokes solutions

There has been some interest in the question of possible growth of the enstrophy, L^2 -norm of the vorticity, in 3D Navier–Stokes equations as well as related dissipative systems ([19, 36, 37, 41, 43, 14]). This quantity is particularly interesting as it determines regularity and uniqueness of the Navier–Stokes solutions. So far, only **upper bounds** on the enstrophy are available: in Lu–Doering [37], the authors have obtained the bound

$$\frac{d}{dt} \|\omega(t, \cdot)\|_{L^2} \leq \frac{C}{\nu^3} \|\omega(t, \cdot)\|_{L^2}^3.$$

Interestingly, in [37], it is reported that the maximal growth of the enstrophy seems to be achieved by flows in which a pair of vortex rings are colliding with each other, which are exactly the flows considered in the current work.

In relation to this, let us demonstrate that infinite enstrophy growth for Euler can be translated to **enstrophy inflation** for Navier–Stokes with small viscosity. While this is strictly speaking not necessary, we restrict ourselves to the axisymmetric systems without swirl: in this case, the 3D Navier–Stokes equations simplify to

$$\partial_t \left(\frac{\omega}{r}\right) + u \cdot \nabla \left(\frac{\omega}{r}\right) = \nu \left(\Delta + \frac{2}{r}\partial_r\right) \left(\frac{\omega}{r}\right), \quad u = K[\omega]. \quad (5.1)$$

In the statement below, we assume that ω_0, ω_0^ν are axisymmetric axial vorticities.

Proposition 5.1. *Let $\omega_0 \in L^\infty(\mathbb{R}^3)$ be compactly supported with finite kinetic energy. Furthermore, assume that the corresponding Euler solution (1.1) exists globally in time with $u \in L_{t,loc}^\infty W_x^{1,\infty}$ and*

$$\limsup_{t \rightarrow \infty} \|\omega(t, \cdot)\|_{L^2(\mathbb{R}^3)} = +\infty.$$

Then, for any sequence of initial data $\omega_0^\nu \in L^\infty \cap L^1(\mathbb{R}^3)$ with finite kinetic energy which converges to ω_0 strongly in $L^2(\mathbb{R}^3)$ as $\nu \rightarrow 0^+$, we have that

$$\liminf_{\nu \rightarrow 0^+} \sup_{t \in [0, \infty]} \|\omega^\nu(t, \cdot)\|_{L^2(\mathbb{R}^3)} = +\infty,$$

where ω^ν is the solution to (5.1) with initial data ω_0^ν and viscosity ν .

Observe that Corollary 1.2 (e.g. see (1.6) of Remark 1.4) provides a class of Euler initial data for which the hypothesis of the above result is satisfied. Global in time existence and uniqueness of strong solutions with initial data $u_0 \in H^1(\mathbb{R}^3)$ to axi-symmetric Navier–Stokes equations without swirl (5.1) is well-known ([52]).

Proof. Towards a contradiction, assume that there exists a sequence $\nu_n \rightarrow 0^+$ such that

$$\sup_{n \geq 1} \sup_{t \in [0, \infty]} \|\omega^{\nu_n}(t, \cdot)\|_{L^2(\mathbb{R}^3)} \leq M$$

for some $M > 0$. We take some $T > 0$, and since the sequence $\{\omega^{\nu_n}\}_{n \geq 1}$ is precompact in the weak star topology of $L^\infty([0, T]; L^2(\mathbb{R}^3))$, by passing to a subsequence if necessary, we derive

$$\omega^{\nu_n} \rightharpoonup \hat{\omega}$$

weakly in star in $L^\infty([0, T]; L^2(\mathbb{R}^3))$. It is not difficult to see that $\hat{\omega}$ is a weak solution to (1.1). Furthermore, using that $\omega^\nu(t, \cdot)$ is weakly continuous in time with values in $L^2(\mathbb{R}^3)$, it can be checked that $\hat{\omega}(t, \cdot) \rightarrow \omega_0$ as $t \rightarrow 0^+$ weakly in $L^2(\mathbb{R}^3)$. From weak-strong uniqueness of Euler in the class $u \in L_t^\infty(W^{1,\infty} \cap L^2)_x$ (see [11, 2]), we deduce that $\hat{\omega} = \omega$, and this is a contradiction since weak convergence implies

$$\sup_{t \in [0, \infty]} \|\omega(t, \cdot)\|_{L^2} \leq M$$

for all $T > 0$. □

Acknowledgement

KC has been supported by the National Research Foundation of Korea (NRF-2018R1D1A1B07043065) and by the UBSI Research Fund(1.219114.01) of UNIST. IJ has been supported by the New Faculty Startup Fund from Seoul National University and the Samsung Science and Technology Foundation under Project Number SSTF-BA2002-04.

References

- [1] J. T. Beale, T. Kato, and A. Majda. Remarks on the breakdown of smooth solutions for the 3-D Euler equations. Comm. Math. Phys., 94(1):61–66, 1984.
- [2] Yann Brenier, Camillo De Lellis, and László Székelyhidi, Jr. Weak-strong uniqueness for measure-valued solutions. Comm. Math. Phys., 305(2):351–361, 2011.
- [3] Jiajie Chen and Thomas Y. Hou. Finite time blowup of 2D Boussinesq and 3D Euler equations with $C^{1,\alpha}$ velocity and boundary. Comm. Math. Phys., 383(3):1559–1667, 2021.
- [4] M. Cheng, J. Lou, and T. T. Lim. Numerical simulation of head-on collision of two coaxial vortex rings. Fluid Dyn. Res., 50(6):065513, 24, 2018.
- [5] Stephen Childress. Models of vorticity growth in Euler flows I, Axisymmetric flow without swirl and II, Almost 2-D dynamics. AML reports 05-07 and 06-07, Courant institute of mathematical sciences, 2007.
- [6] Stephen Childress. Growth of anti-parallel vorticity in Euler flows. Phys. D, 237(14-17):1921–1925, 2008.
- [7] Stephen Childress and Andrew D. Gilbert. Eroding dipoles and vorticity growth for Euler flows in \mathbb{R}^3 : the hairpin geometry as a model for finite-time blowup. Fluid Dyn. Res., 50(1):011418, 40, 2018.
- [8] Kyudong Choi. Stability of Hill’s spherical vortex. preprint, arXiv:2011.06808.
- [9] Kyudong Choi and In-Jee Jeong. Filamentation near Hill’s vortex. preprint, arXiv:2107.06035.
- [10] C. Chu, C. Wang, C. Chang, R. Chang, and W. Chang. Head-on collision of two coaxial vortex rings: Experiment and computation. Journal of Fluid Mechanics, 296:39–71, 1995.
- [11] R. Danchin. Axisymmetric incompressible flows with bounded vorticity. Uspekhi Mat. Nauk, 62:73–94, 2007.
- [12] Sergey A. Denisov. Double exponential growth of the vorticity gradient for the two-dimensional Euler equation. Proc. Amer. Math. Soc., 143(3):1199–1210, 2015.
- [13] Sergey A. Denisov. The sharp corner formation in 2D Euler dynamics of patches: infinite double exponential rate of merging. Arch. Ration. Mech. Anal., 215(2):675–705, 2015.
- [14] Charles R. Doering and J. D. Gibbon. Applied analysis of the Navier-Stokes equations. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 1995.
- [15] F. W. Dyson. The potential of an anchor ring. Philosophical Transactions of the Royal Society of London. A, 184:43–95, 1893.
- [16] Tarek M. Elgindi. Finite-time singularity formation for $C^{1,\alpha}$ solutions to the incompressible Euler equations on \mathbb{R}^3 . Ann. of Math., to appear, arXiv:1904.04795.

- [17] Tarek M. Elgindi and In-Jee Jeong. The incompressible Euler equations under octahedral symmetry: singularity formation in a fundamental domain. Adv. Math., to appear, [arXiv:2001.07840](https://arxiv.org/abs/2001.07840).
- [18] Tarek M. Elgindi and In-Jee Jeong. Finite-Time Singularity Formation for Strong Solutions to the Axi-symmetric 3D Euler Equations. Ann. PDE, 5(2):Paper No. 16, 2019.
- [19] Aseel Farhat, Michael Jolly, and Evelyn Lunasin. Bounds on energy and enstrophy for the 3D Navier-Stokes- α and Leray- α models. Commun. Pure Appl. Anal., 13(5):2127–2140, 2014.
- [20] H. Feng and V. Šverák. On the Cauchy problem for axi-symmetric vortex rings. Arch. Ration. Mech. Anal., 215:89–123, 2015.
- [21] R. Finn and D. R. Smith. On the stationary solutions of the Navier-Stokes equations in two dimensions. Arch. Rational Mech. Anal., 25:26–39, 1967.
- [22] Pascal Gamblin and Xavier Saint Raymond. On three-dimensional vortex patches. Bull. Soc. Math. France, 123(3):375–424, 1995.
- [23] Hui Guan, Zhi-Jun Wei, Elizabeth Rumenova Rasolkova, and Chui-Jie Wu. Numerical simulations of two coaxial vortex rings head-on collision. Adv. Appl. Math. Mech., 8(4):616–647, 2016.
- [24] H. Helmholtz. Über Integrale der hydrodynamischen Gleichungen, welche den Wirbelbewegungen entsprechen, volume 55, pages 25–55. 1858.
- [25] Thomas Hou. Potential singularity of the 3D Euler equations in the interior domain. [arXiv:2107.05870](https://arxiv.org/abs/2107.05870).
- [26] <https://www.youtube.com/watch?v=Qxr7tsZUy1c>. Head-on vortex ring collision, 2020.
- [27] Chaocheng Huang. Singular integral system approach to regularity of 3D vortex patches. Indiana Univ. Math. J., 50(1):509–552, 2001.
- [28] Dragoş Iftimie, Thomas C. Sideris, and Pascal Gamblin. On the evolution of compactly supported planar vorticity. Comm. Partial Differential Equations, 24:1709–1730, 1999.
- [29] In-Jee Jeong. Loss of Regularity for the 2D Euler Equations. J. Math. Fluid Mech., 23(4):Paper No. 95, 2021.
- [30] T. Kambe and T. Minota. Acoustic wave radiated by head-on collision of two vortex rings. Proc. R. Soc. Lond. A, 386:277–308, 1983.
- [31] T. Kambe and U. Mya Oo. An axisymmetric viscous vortex motion and its acoustic emission. J. Phys. Soc. Japan, 53:2263–2273, 1984.
- [32] Alexander Kiselev and Chao Li. Global regularity and fast small-scale formation for Euler patch equation in a smooth domain. Comm. Partial Differential Equations, 44(4):279–308, 2019.
- [33] Alexander Kiselev and Vladimir Šverák. Small scale creation for solutions of the incompressible two-dimensional Euler equation. Ann. of Math. (2), 180(3):1205–1220, 2014.

- [34] H. Lamb. Hydrodynamics. Cambridge Univ. Press., 3rd ed. edition, 1906.
- [35] T. T. Lim and T. B. Nickels. Instability and reconnection in the head-on collision of two vortex rings. Nature, 357:225–227, 1992.
- [36] Lu Lu. Bounds on the enstrophy growth rate for solutions of the three-dimensional Navier-Stokes equations. ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)—University of Michigan.
- [37] Lu Lu and Charles R. Doering. Limits on enstrophy growth for solutions of the three-dimensional Navier-Stokes equations. Indiana Univ. Math. J., 57(6):2693–2727, 2008.
- [38] Andrew J. Majda and Andrea L. Bertozzi. Vorticity and incompressible flow, volume 27 of Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2002.
- [39] Y. Oshima. Head-on collision of two vortex rings. J. Phys. Soc. Japan, 44:328–331, 1978.
- [40] A. J. Peace and N. Riley. A viscous vortex pair in ground effect. J. Fluid Mech., 129:409–426, 1983.
- [41] Dmitry Pelinovsky. Enstrophy growth in the viscous Burgers equation. Dyn. Partial Differ. Equ., 9(4):305–340, 2012.
- [42] Richard B. Pelz. Symmetry and the hydrodynamic blow-up problem. J. Fluid Mech., 444:299–320, 2001.
- [43] Diogo Poças and Bartosz Protas. Transient growth in stochastic Burgers flows. Discrete Contin. Dyn. Syst. Ser. B, 23(6):2371–2391, 2018.
- [44] X. Saint Raymond. Remarks on axisymmetric solutions of the incompressible Euler system. Comm. Partial Differential Equations, 19(1-2):321–334, 1994.
- [45] Philippe Serfati. Régularité stratifiée et équation d’Euler 3D à temps grand. C. R. Acad. Sci. Paris Sér. I Math., 318(10):925–928, 1994.
- [46] K. Shariff and A. Leonard. Vortex Rings. Ann. Rev. Fluid Mech., 24:235–279, 1992.
- [47] K. Shariff, A. Leonard, N. J. Zabusky, and J. H. Ferziger. Acoustics and dynamics of coaxial interacting vortex rings. In Vortex Motion (ed. H. Hasimoto and T. Kambe), 1988.
- [48] M. J. Shelley, D. I. Meiron, and S. A. Orszag. Dynamical aspects of vortex reconnection of perturbed anti-parallel vortex tubes. J. Fluid Mech., 246:613–652, 1993.
- [49] S. Stanaway, K. Shariff, and F. Hussain. Head-on collision of viscous vortex rings. Proceedings of the Summer Program, Center for Turbulence Research, 1988.
- [50] Nader Masmoudi Tarek M. Elgindi, Tej-Eddine Ghouli. On the stability of self-similar blow-up for $c^{1,\alpha}$ solutions to the incompressible Euler equations on \mathbb{R}^3 . preprint, arXiv:1910.14071.
- [51] W. Thomson. The translatory velocity of a circular vortex ring. Phil. Mag. (Ser. 4), 34:511–512, 1867.

- [52] M. R. Ukhovskii and V. I. Yudovich. Axially symmetric flows of ideal and viscous fluids filling the whole space. J. Appl. Math. Mech., 32:52–61, 1968.
- [53] Cheng Yang. Vortex motion of the Euler and lake equations. J. Nonlinear Sci., 31(3):Paper No. 48, 21, 2021.
- [54] V. I. Yudovich. On the loss of smoothness of the solutions of the Euler equations and the inherent instability of flows of an ideal fluid. Chaos, 10(3):705–719, 2000.
- [55] Andrej Zlatoš. Exponential growth of the vorticity gradient for the Euler equation on the torus. Adv. Math., 268:396–403, 2015.