

AN ACCURATE ANALYTICAL FIT TO THE GRAVITATIONAL-WAVE INSPIRAL DURATION FOR ECCENTRIC BINARIES

ILYA MANDEL^{1,2,3}

¹Monash Centre for Astrophysics, School of Physics and Astronomy, Monash University, Clayton, Victoria 3800, Australia

²The ARC Center of Excellence for Gravitational Wave Discovery – OzGrav and

³Institute of Gravitational Wave Astronomy and School of Physics and Astronomy, University of Birmingham, Birmingham, B15 2TT, United Kingdom

Draft version January 22, 2022

ABSTRACT

I present an analytical fit to the duration of the inspiral of two point masses driven by gravitational-wave emission. The fit is accurate to within 3% over the entire range of initial eccentricities from 0 to 0.99999.

Peters (1964) derived a very widely used post-Newtonian expression for the time T for two point masses M_1 and M_2 to spiral in through gravitational-wave emission from an initial separation a_0 and initial eccentricity e_0 :

$$T = T_c \frac{48}{19} \left(\frac{1 - e_0^2}{e_0^{12/19} \left(1 + \frac{121}{304} e_0^2 \right)^{870/2299}} \right)^4 \times \int_0^{e_0} \frac{e^{29/19} \left(1 + \frac{121}{304} e^2 \right)^{1181/2299}}{(1 - e^2)^{3/2}} de, \quad (1)$$

where

$$T_c = \frac{5c^5 a_0^4}{256G^3 M_1 M_2 (M_1 + M_2)}. \quad (2)$$

Equation (1) is somewhat awkward to compute in practice because it has apparently singular terms at both $e \rightarrow 0$ and $e \rightarrow 1$, although the complete expression is well-behaved in both limits. Consequently it is common to use two approximations derived by Peters (1964):

$$\tau(e_0 \rightarrow 0) \approx T_c \left(\frac{1 - e_0^2}{\left(1 + \frac{121}{304} e_0^2 \right)^{870/2299}} \right)^4 \quad (3)$$

and

$$\tau(e_0 \rightarrow 1) \approx \frac{768}{425} T_c (1 - e_0^2)^{7/2}. \quad (4)$$

However, as shown in Figure 1, neither fit is particularly accurate in the intervening regime. For example, at $e_0 = 0.8$, the estimate from Eq. (3) is too low by a factor of 2.4, while that from Eq. (4) is too large by 75%. Therefore, although sometimes one of the estimates (3) or (4) is used for all eccentricities, this is quite inaccurate. Some codes that need to compute inspiral durations more accurately (e.g., binary population synthesis codes, Team COMPAS: Riley et al. 2021) instead opt to split up the inspiral duration calculation into three pieces: the low-eccentricity regime where fit (3) is applied, the high-eccentricity regime where fit (4) is applied, and the intermediate regime where the integral (1)




FIG. 1.— The ratios of the fits given by Eqs. (3), (4) and (5) to the complete expression of Eq. (1) as a function of initial eccentricity. The fit proposed here (black) is accurate to within 3% over the full range of initial eccentricities up to 0.99999.

is computed numerically. This, however, becomes rather cumbersome.

Pierro & Pinto (1996) showed that Eq. (1) can be solved exactly with an Appell function, a 2-variable hypergeometric series. However, to remain accurate, the series expansion requires a larger number of terms as eccentricity grows, and is not readily available in many software libraries. Meanwhile, interpolation between pre-computed values of the inspiral duration can be an accurate and computationally efficient solution; however, the extra implementation effort makes it somewhat inconvenient.

The following fit to Eq. (1) is purely analytical and easy to compute:

$$T \approx T_c \left(1 + 0.27e_0^{10} + 0.33e_0^{20} + 0.2e_0^{1000} \right) (1 - e_0^2)^{7/2}. \quad (5)$$

It is accurate to $< 3\%$ over the entire range of initial eccentricities between 0 and 0.99999, i.e., $(1 - e_0) \in [10^{-5}, 1]$. The ratio of this fit to the full solution given by Eq. (1) is shown in Figure 1 in black.

The Peters (1964) solution given in Eq. (1) is an approximation valid in the inspiral-dominated post-Newtonian regime, which is generally the situation of

interest for gravitational-wave sources with non-extreme mass ratios that are captured with periapsis separations significantly larger than the innermost stable circular orbit (see Pierro & Pinto 1996 for a detailed discussion of the regime of applicability). A number of works relaxed one or more of these conditions to obtain more complex expressions for the inspiral duration. For example, Gair et al. (2006) computed corrections to the quadrupole flux from conservative relativistic dynamics in order to obtain durations of eccentric extreme-mass-ratio inspirals (EMRIs). Will & Maitra (2017) included the effect of the spin of the supermassive black hole on the duration of an eccentric EMRI. Zwick et al. (2020) considered the impact of post-Newtonian corrections; however, their fit is less accurate at moderate eccentricities and non-monotonic as a function of eccentricity. Tucker & Will (2021) introduced a fit to the inspiral duration including post-Newtonian correc-

tions through the 4.5 order, which is accurate to within 2% for $100 \leq a_0(1 - e_0^2)c^2/G/(M_1 + M_2) \leq 1000$ and $0 \leq e_0 \leq 0.999$.

Nevertheless, the simple fit to the Peters (1964) gravitational-wave driven coalescence time proposed in Eq. (5) may prove useful in astrophysically relevant settings.

I thank Bence Kocsis, Jonathan Gair, Cole Miller, Karthik Rajeev, Frank Timmes, Alexandria Tucker, Clifford Will, Xingjiang Zhu, and Lorenz Zwick for discussions. I acknowledge support from the Australian Research Council Centre of Excellence for Gravitational Wave Discovery (OzGrav), through project number CE17010004. I am a recipient of the Australian Research Council Future Fellowship FT190100574.

Gair, J. R., Kennefick, D. J., & Larson, S. L. 2006, *Astrophysical Journal*, 639, 999, astro-ph/0508275
 Peters, P. C. 1964, *Physical Review*, 136, 1224
 Pierro, V., & Pinto, I. M. 1996, *Nuovo Cimento B Serie*, 111B, 631
 Team COMPAS: Riley, J. et al. 2021, arXiv e-prints,

2109.10352
 Tucker, A., & Will, C. M. 2021, arXiv e-prints, 2108.12210
 Will, C. M., & Maitra, M. 2017, *Phys. Rev. D*, 95, 064003, 1611.06931
 Zwick, L., Capelo, P. R., Bortolas, E., Mayer, L., & Amaro-Seoane, P. 2020, *MNRAS*, 495, 2321, 1911.06024