

BCS-BEC crossover of atomic Fermi superfluid in a spherical bubble trap

Yan He,¹ Hao Guo,² and Chih-Chun Chien³

¹*College of Physics, Sichuan University, Chengdu, Sichuan 610064, China**

²*Department of Physics, Southeast University, Jiulonghu Campus, Nanjing 211189, China[†]*

³*Department of physics, University of California, Merced, CA 95343, USA.[‡]*

We present a theory of a two-component atomic Fermi gas with tunable attractive interactions on a spherical shell going through the BCS - Bose Einstein condensation (BEC) crossover, inspired by the realizations of spherical bubble traps for ultracold atoms in microgravity. The derivation follows the BCS-Leggett theory to obtain the gap and number equations. The BCS-BEC crossover can be induced by tuning the interaction, and the properly normalized gap and chemical potential exhibit universal behavior regardless of the planar or spherical geometry. Nevertheless, the spherical-shell geometry introduces another way of inducing the crossover by the curvature. The curvature-induced BCS-BEC crossover is made possible by fixing the particle number and interaction strength while shrinking the sphere, causing a reduction to the ratio of the pairing and kinetic energies and pushing the system towards the BCS limit. The saturation of the superfluid density further confirms the ground state is a Fermi superfluid.

Ultracold atoms have offer versatile platforms for studying quantum many-body physics with precise controls and broad tunability [1–8]. While the Bose-Einstein condensation (BEC) has been the foundation behind major research of bosonic atoms [9, 10], pairing between fermionic atoms introduces the BCS-BEC crossover that smoothly interpolate the behavior of fermionic and bosonic superfluids [11–15]. On the BCS side of the crossover at zero temperature, the pairing gap is small with respect to the Fermi energy E_F while the chemical potential μ is near E_F . On the BEC side, the gap is comparable to or larger than E_F while μ becomes negative due to the strong binding of fermions. The mean-field BCS-Leggett theory [16] captures the main feature of the ground state in the crossover.

On the other hand, geometry has played an important role in the study of cold atoms. For example, an atomic superfluid in a harmonic trap carries angular momentum by forming vortices [17], but an atomic superfluid in a ring-shape trap carries angular momentum by its circulating persistent current [18]. Another example is the realizations of 2D planar atomic systems, including 2D superfluids [19, 20], 2D BCS-BEC crossover [21, 22], spin-orbit coupled superfluids [23], phase transitions [24, 25], scale invariance [26], along with many theoretical works [27–32]. Recently, spherical bubble traps for cold atoms in microgravity environment, such as the outer space, have been demonstrated [33, 34]. While there have been theoretical studies of bosonic superfluid on a spherical shell [35–41], showing enhanced transition temperature, vortices, multi-component mixtures, etc., less references can be found on fermionic superfluid on a spherical shell.

Here we present an analysis of the BCS-BEC crossover of a Fermi superfluid on a 2D spherical shell at the level of the BCS-Leggett theory. The dispersion of an ideal Fermi gas on a spherical shell already exhibits interesting features [42], including degeneracy within an angular-

momentum level and jumps between adjacent levels. By considering a contact interaction similar to that in nuclear matter [43], we obtain a mean-field Hamiltonian describing pairing of the fermions on a spherical shell. Implementing the Bogoliubov transformation [43], the gap and number equations on a spherical shell are derived. The solution exhibits the signatures of the BCS-BEC crossover as the attractive interaction increases. When the gap and chemical potential are properly normalized, they exhibit universal behavior that depends only on the interaction but not the curvature, as long as the sphere is large so that the scattering remains a local event.

Nevertheless, the curvature will be shown to influence the Fermi superfluid and induce its own BCS-BEC crossover on a sphere. This is because a bound state always exists in the 2D two-body scattering [44, 45], making it very different from the 3D case. The binding energy is determined by the scattering length that quantifies the interaction strength. In experiments, the size of the spherical bubble trap is expected to be tunable with the particle number fixed, so the particle density increases with the curvature. Since the Fermi energy increases with the particle density, the ratio of the pairing energy indicated by the binding energy and the kinetic energy indicated by the Fermi energy decreases as the spherical shell shrinks, thereby pushing the Fermi superfluid towards the BCS limit even when the interaction is fixed. The curvature-induced BCS-BEC crossover is made possible by the 2D nature and the compactness of the spherical bubble trap, and its realization will offer another elegant example of geometric effects on quantum systems.

Fermions on a spherical shell — We consider a two-component atomic Fermi gas confined in a spherical bubble trap. Assuming the shell is thin, the gas thus lives on the surface of a sphere. For a free Fermi gas confined on

a spherical shell, the energy dispersion is given by [42]

$$\epsilon_l = \frac{\hbar^2}{2mR^2}l(l+1), \quad l = 0, 1, \dots \quad (1)$$

Here m is the mass of the atoms and R is the radius of the sphere. In the following, we will set $\hbar = 1$ and $k_B = 1$. l is the quantum number of the orbital angular momentum. For a fixed l , the magnetic quantum number takes the values $m_z = -l, \dots, l$, and $\sigma = \uparrow, \downarrow$ labels the two spin components. Therefore, there are $2(2l+1)$ degenerate states for the level labeled by l .

After including a two-body interaction term, the Hamiltonian in the grand-canonical ensemble is $H = H_K + H_I$, where $H_K = \sum_{l,m,\sigma} (\epsilon_l - \mu) c_{lm\sigma}^\dagger c_{lm\sigma}$ and

$$H_I = \sum_{l_1, m_1, \dots} V_{1234} c_{l_1 m_1 \sigma_1}^\dagger c_{l_2 m_2 \sigma_2}^\dagger c_{l_3 m_3 \sigma_3} c_{l_4 m_4 \sigma_4} \quad (2)$$

Here $V_{1234} = \langle l_1, m_1; l_2, m_2 | V | l_3, m_3; l_4, m_4 \rangle$ and $c_{lm\sigma}^\dagger$ ($c_{lm\sigma}$) is the creation (annihilation) operator. We also assume equal populations of the two component, so $\mu_\sigma = \mu$. Assuming the two-body potential is rotational invariant, then non-vanishing matrix elements only occur if the magnetic quantum numbers satisfy $m_1 + m_2 = m_3 + m_4$, as indicated by the prime above the \sum .

In the conventional BCS theory, one only considers two-body scattering from $|\pm \mathbf{k}\rangle$ to $|\pm \mathbf{k}'\rangle$. Inspired by such a simplification, we also focus on the initial and final states that can be coupled into $|L = 0, M = 0\rangle$ with spin singlet and ignore other scattering processes. The approximation then leads us to the reduced interaction Hamiltonian $H_I = \sum_{l_1, m_1, l_2, m_2} V_{12} c_{l_1 m_1 \uparrow}^\dagger c_{l_1, -m_1 \downarrow}^\dagger c_{l_2 m_2 \uparrow} c_{l_2, -m_2 \downarrow}$. Here $V_{12} = \langle l_1, m_1; l_1, -m_1 | V | l_2, m_2; l_2, -m_2 \rangle$. The coupling among the angular-momentum states gives $|l, m; l, -m\rangle = \sum_{L=0}^{2l} |l, l; L, 0\rangle \langle l, l; L, 0 | l, m; l, -m\rangle$. Here we only keep the $L = 0$ state and use the fact that $\langle l, l; L, 0 | l, m; l, -m\rangle = (-1)^{l-m} / \sqrt{2l+1}$. The interaction Hamiltonian then becomes $H_I = \sum_{l_1, m_1, l_2, m_2} V_{12,0} c_{l_1 m_1 \uparrow}^\dagger c_{l_1, -m_1 \downarrow}^\dagger c_{l_2 m_2 \uparrow} c_{l_2, -m_2 \downarrow}$. Here $V_{12,0} = \langle l_1, l_1; 0, 0 | V | l_2, l_2; 0, 0 \rangle \frac{(-1)^{l_1-m_1} (-1)^{l_2-m_2}}{\sqrt{(2l_1+1)(2l_2+1)}}$. The factor $(-1)^{l_1-m_1} (-1)^{l_2-m_2}$ inside $V_{12,0}$ can be removed by a canonical transformation $c_{lm} \rightarrow c_{lm}$ and $c_{l-m} \rightarrow (-1)^{l-m} c_{l-m}$. The form of H_I is now suitable for a general mean-field approximation similar to the BCS theory.

BCS theory on a spherical shell — Following the BCS approximation, we make the substitutions $c_{lm\uparrow} c_{l,-m\downarrow} \rightarrow \langle c_{lm\uparrow} c_{l,-m\downarrow} \rangle$ and $c_{lm\uparrow}^\dagger c_{l,-m\downarrow}^\dagger \rightarrow \langle c_{lm\uparrow}^\dagger c_{l,-m\downarrow}^\dagger \rangle$ in the interaction Hamiltonian and obtain the BCS Hamiltonian

$$H_{BCS} = H_K + \sum_{l,m} (-\Delta_l c_{lm\uparrow}^\dagger c_{l,-m\downarrow}^\dagger - \Delta_l c_{lm\uparrow} c_{l,-m\downarrow}) \quad (3)$$

Here the gap function is given by

$$\Delta_j = -\frac{1}{(2j+1)^{1/2}} \sum_{l,m} V_{jl} \frac{1}{(2l+1)^{1/2}} \langle c_{lm\uparrow} c_{l,-m\downarrow} \rangle \quad (4)$$




Figure 1. Universal behavior of interaction-induced BCS-BEC crossover: The normalized gap (left) and chemical potential (right) as a function of $-\ln(k_F a)$ on a 2D plane according to Eq. (7) (red circles) and on the shell of a unit sphere according to Eq. (8) (blue squares) at zero temperature.

and $V_{jl} = \langle j, j; 0, 0 | V | l, l; 0, 0 \rangle$. The BCS Hamiltonian can be diagonalized by the Bogoliubov transformation [43] with

$$c_{lm\uparrow} = u_l \alpha_{lm} - v_l \beta_{lm}, \quad c_{l,-m\downarrow}^\dagger = v_l \alpha_{lm} + u_l \beta_{lm}. \quad (5)$$

The coefficients are given by $u_l^2 = \frac{1}{2}(1 + \frac{\xi_l}{E_l})$ and $v_l^2 = \frac{1}{2}(1 - \frac{\xi_l}{E_l})$, where $\xi_l = \epsilon_l - \mu$ and $E_l = \sqrt{\xi_l^2 + \Delta_l^2}$. The diagonalized Hamiltonian has the form

$$H_{BCS} = \sum_{lm} (\xi_l - E_l) + \sum_{lm} E_l (\alpha_{lm}^\dagger \alpha_{lm} + \beta_{lm}^\dagger \beta_{lm}). \quad (6)$$

In terms of the Bogoliubov transformation, the gap function becomes $\Delta_j = \frac{-1}{(2j+1)^{1/2}} \sum_l V_{jl} (2l+1)^{1/2} u_l v_l [1 - 2f(E_l)]$. Here $f(x) = 1/[\exp(x/T) + 1]$ is the Fermi distribution function. We will further approximate the matrix element V_{jl} before solving the gap equation. Meanwhile, the number equation can be derived from $n = \sum_{l,m,\sigma} \langle c_{lm\sigma}^\dagger c_{lm\sigma} \rangle$. Explicitly, $n = \sum_l (2l+1) \left(1 - \frac{\xi_l}{E_l} + 2\frac{\xi_l}{E_l} f(E_l)\right)$. Solving the gap and number equations gives us Δ and μ of the Fermi gas.

BCS theory on 2D spherical shell — We begin with a brief review of the BCS-BEC crossover on a 2D plane, following Refs. [27, 32]. To handle the bound state from the 2D two-body scattering, a regularization introduces a binding energy $\epsilon_b = -\hbar^2/(ma^2)$, where a is the 2D scattering length. Combining the binding energy with the renormalization of the contact interaction, the coupling constant is expressed in terms of the scattering length via $\frac{1}{g} = \int \frac{d^2 k}{(2\pi)^2} \frac{1}{2\epsilon_{\mathbf{k}} + |\epsilon_b|}$. Here $\epsilon_{\mathbf{k}} = \hbar^2 k^2/(2m)$ is the free-fermion dispersion. The gap and number equations can be simplified to

$$\int \frac{d^2 k}{(2\pi)^2} \left[\frac{1 - f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} - \frac{1}{2\epsilon_{\mathbf{k}} + |\epsilon_b|} \right] = 0, \quad (7)$$

$$n = \int \frac{d^2 k}{(2\pi)^2} \left(1 - \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} + 2 \frac{\xi_{\mathbf{k}}}{E_{\mathbf{k}}} f(E_{\mathbf{k}}) \right).$$

Solving the equations gives Δ and μ once the values of a and T are given.

To simplify the BCS theory on a spherical shell, we implement another approximation of the matrix element V_{jl} by considering only a short-range attractive interaction given by $V = -g\delta(1 - \cos\theta_{12})$, where $\cos\theta_{12} = \cos\theta_1 \cos\theta_2 + \sin\theta_1 \sin\theta_2 \cos(\phi_1 - \phi_2)$. After using a generalization of the Wigner-Eckart theorem [43] as explained in the Supplemental Material, the matrix element becomes $V_{jl} = -g\sqrt{(2j+1)(2l+1)}$. The gap equation is then reduced to $\Delta_j = g \sum_l (2l+1) \frac{\Delta_l}{2E_l} [1 - 2f(E_l)]$. Since the right hand side does not depend on j , we conclude that Δ does not depend on j explicitly. Hence, the gap equation becomes $\frac{1}{g} = \sum_l \frac{2l+1}{2E_l} [1 - 2f(E_l)]$. Since $E_l \propto l(l+1)$, the dominant terms in the summation will behave like $\sum_l \frac{2l+1}{2E_l} \sim \sum_l \frac{2l+1}{l(l+1)} \rightarrow \infty$ due to the contact-interaction approximation. A systematic renormalization scheme, similar to the one in flat space, can be applied to render meaningful physical results.

Following the planar case, the regularization on a 2D spherical shell can be modified as $\frac{1}{g} = \int dl \frac{2l+1}{2\epsilon_l + |\epsilon_b|}$. We assume $\epsilon_b = -\hbar^2/(ma^2)$ due to its localized nature. After the regularization, we obtain the gap and number equations as

$$\int dl (2l+1) \left[\frac{1 - 2f(E_l)}{2E_l} - \frac{1}{2\epsilon_l + |\epsilon_b|} \right] = 0, \quad (8)$$

$$n = \frac{1}{4\pi R^2} \int dl (2l+1) \left(1 - \frac{\xi_l}{E_l} + 2 \frac{\xi_l}{E_l} f(E_l) \right).$$

We mention there is another regularization scheme summarized in the Supplemental Material that produces qualitatively the same results. Moreover, we have approximate the summations by integrals, and a comparison in the Supplemental Material shows that there is no observable difference between the results from the summations and the integrals for reasonably large l .

Numerical results of the BCS-BEC crossover on a 2D spherical shell at zero temperature are shown in Fig. 1, along with the results of the 2D planar case. We plot Δ and μ as a function of $-\ln(k_F a)$ for both cases. For the 2D planar case, E_F and k_F are the Fermi energy and Fermi momentum of a noninteracting Fermi gas with the same density. For the spherical-shell case, we take E_F and $k_F = \sqrt{2mE_F}$ from a noninteracting Fermi gas with the same total particle number. Assuming the largest occupied shell has angular momentum L_m for a free Fermi gas, the total particle number is $N = 2L_m(L_m + 1)$, so $E_F = \frac{L_m(L_m+1)}{2mR^2}$ and $n = N/(4\pi R^2)$. As $-\ln(k_F a)$ increases, the gap increases while the chemical potential decreases, showing the signature of the BCS-BEC crossover. The crossover may be identified as the regime where μ changes sign. Since the scattering length reflects the effective interactions between the fermions, Fig. 1 shows the interaction induced BCS-BEC crossover in two different 2D geometries.




Figure 2. Curvature-induced BCS-BEC crossover on a spherical shell at zero temperature: (Top panels) Gap as a function of $1/R^2$ normalized to E_F (left) and Δ_0 (right). (Bottom panels) Chemical potential as a function of $1/R^2$ normalized to E_F (left) and μ_0 (right). Δ_0 and μ_0 on the right panels are the gap and chemical potential of $N = 220$ fermions on the shell of a unit sphere. Here $-\ln(k_F a) = -0.1$ for the unit sphere, and then a is fixed while $1/R^2$ varies.

Importantly, when normalized according to their respective intrinsic quantities like E_F and k_F , the results of the 2D plane are indistinguishable from those of the spherical shell. This is because the pairing from the contact interaction is a local property of the Fermi gas. As a consequence, properly normalized quantities reflect the same local behavior from the mean-field theory and fail to differentiate the global geometry. For the spherical case, taking different values of N , L_m , and R produces the same universal results of Δ/E_F and μ/E_F . The universal behavior can also be confirmed by the resemblance of the equations of state, Eqs. (7) and (8), when written in the normalized quantities (see the Supplemental Material for details).

Curvature induced BEC-BCS crossover — The universal behavior of Fig. 1 may lead to a false impression that the curvature of the sphere, $1/R^2$, does not play a significant role. However, one may envision that the radius of the spherical bubble trap is tunable and consider a different scenario where the particle number, not the local density, is conserved and compare the physical quantities with different curvatures but the same interaction strength. As the curvature increases, the surface area shrinks and the local particle density increases if the total particle number is fixed. Since E_F increases with the density, it is tempting to claim that the gap will increase with the curvature if Δ/E_F is roughly the same. A careful analysis, however, reveals the opposite and establish a BEC-BCS crossover induced by the curvature.

To demonstrate the curvature effects, we plot μ and Δ as functions of the curvature with fixed particle number and scattering length. As the radius of the sphere shrinks from R_0 , one can see that the gap become smaller with respect to the gap at R_0 . The reason is that as $1/R^2$ in-




Figure 3. Gap (left) and superfluid density (right) as functions of T/E_F of a Fermi superfluid on the shell of a unit sphere. Here $-\ln(k_F a) = -1.35$ and $N = 220$. $T^*/E_F \approx 0.42$ is where Δ approaches zero. The vertical dashed line indicates $T_{BKT}/E_F \approx 0.29$, above which n_s is expected to drop to zero.

creases, the Fermi energy becomes larger. Meanwhile, the effective binding energy is fixed by the scattering length. The ratio $|\epsilon_b|/E_F$ thus decreases with the curvature, resulting in a situation where the kinetic energy dominates the pairing energy and thereby driving the system into the BCS limit.

Thus, there are two ways to sweep a Fermi superfluid across the BCS-BEC crossover on a spherical shell, one by tuning the interaction and the other by tuning the geometry. The first one is an analogue of the 2D planar case, where the particle density is fixed and the scattering length is tuned via magnetic or optical means. The second one requires a compact 2D geometry, where the particle number and scattering length are fixed but the ratio between the binding energy and Fermi energy is tuned by the geometry. We remark that the latter is possible in 2D because the two-body binding energy is always present [44, 45], different from the general 3D case where the binding energy is finite only on the BEC side. We also caution that the calculation of the two-body scattering length assumes the system is locally flat, and the assumption breaks down when the curvature is too large, or when $R \sim O(a)$. Moreover, we note that the BCS-Leggett theory does not take into account the induced interaction [1] and Hartree-Fock energy [46]. The former reduces the transition temperature by a factor in the 3D case, and the latter shifts the chemical potential. As a first attempt to develop the BCS-Leggett theory of Fermi superfluids on a spherical shell, we leave those effects for future, more refined studies.

Furthermore, we evaluate the superfluid density given by

$$n_s = n - \frac{1}{4\pi R^2} \int dl 2(2l+1) \frac{l(l+1)}{2mR^2} \left[-\frac{df(E_l)}{dE_l} \right]. \quad (9)$$

A derivation is shown in the Supplemental Material. As $T \rightarrow 0$, $-\frac{df(x)}{dx}$ approaches the delta function. Since E_l is positive, the delta function can never be satisfied. Therefore, $n_s/n = 1$ at $T = 0$ across the whole interaction-induced BCS-BEC crossover, so the ground state is in-

deed a Fermi superfluid. Nevertheless, in the curvature-induced crossover, the density increases with the curvature because the total particle number is fixed, leading to an interesting scenario where n_s increases while Δ decreases with the curvature according to the upper-right panel of Fig. 2. The disparity of the dependence of n_s and Δ on the curvature has its root in that Δ is associated with thermodynamics while n_s is from linear response to perturbations.

Finite-temperature effects — Finally, we investigate the mean-field theory away from zero temperature by solving the gap and number equations at finite temperatures. In Figure 3, we show Δ and n_s as functions of T . The mean-field transition temperature T^* is the point above which Δ vanishes. The 2D Berezinskii-Kosterlitz-Thouless (BKT) transition [47–49] temperature may be estimated by $\frac{k_B T_{BKT}}{\hbar^2 n_s(T_{BKT})/m} = \frac{\pi}{2}$. For the case shown in Figure 3, T_{BKT} is below T^* , so the BKT transition will preempt the mean-field transition and cause n_s to jump to zero. The BCS-Leggett theory only provides a qualitative description of the crossover at finite temperatures. It has been shown [11–15] that the preformed pairs, which are the analogue of thermal bosons in a Bose gas, lead to a substantially lower T_c on the BEC side. There have been studies of 2D planar Fermi superfluids that include pairing fluctuations [22, 31, 32]. A full treatment of the finite-temperature BCS-BEC crossover on a spherical shell will be worth another publication.

Conclusion — We presented a generalization of the BCS-Leggett theory of the BCS-BEC crossover and analyzed the behavior of a Fermi superfluid on a spherical shell, relevant to future experiments using spherical bubble traps in microgravity. Although the highly degenerate levels and jumps between the levels of an ideal Fermi gas on a spherical shell makes it different from that on a 2D plane, the pairing gap and chemical potential of a Fermi superfluid after proper normalization exhibit universal behavior transcending the underlying geometries. Therefore, the conventional interaction-induced BCS-BEC crossover of Fermi superfluid is also present on a spherical shell. Nevertheless, the spherical geometry introduces the curvature-induced BCS-BEC crossover by fixing the interaction strength and particle number while reducing the size of the spherical shell. The latter type of crossover is due to a suppression of the ratio between the pairing and kinetic energies by the curvature. Our work paves the way towards a systematic investigation of Fermi superfluids with compact geometries, exemplified by the spherical bubble traps.

* heyancp@scu.edu.cn

[†] guohao.ph@seu.edu.cn

[‡] cchien5@ucmerced.edu

- [1] C. J. Pethick and H. Smith, *Bose-Einstein condensation in dilute gases* (Cambridge University Press, Cambridge, UK, 2008), 2nd ed.
- [2] M. Ueda, *Fundamentals and New Frontiers of Bose-Einstein Condensation* (World Scientific Publishing, Singapore, 2010).
- [3] K. Levin, A. Fetter, and D. Stamper-Kurn, eds., *Ultracold Bosonic and Fermionic Gases* (Elsevier, Amsterdam, Netherlands, 2012).
- [4] P. Torma and K. Sengstock, eds., *Quantum Gas Experiments: Exploring Many-Body States* (Imperial College Press, London, UK, 2014).
- [5] L. Pitaevskii and S. Stringari, *Bose-Einstein Condensation and Superfluidity* (Oxford University Press, Oxford, UK, 2016).
- [6] H. T. C. Stoof, D. B. M. Dickerscheid, and K. Gubbels, *Ultracold Quantum Fields* (Springer, Berlin, Germany, 2009).
- [7] C. Gardiner and P. Zoller, *The Quantum World of Ultra-Cold Atoms and Light Book III: Ultra-cold Atoms* (World Scientific Publishing, Singapore, 2017).
- [8] H. Zhai, *Ultracold Atomic Physics* (Cambridge University Press, Cambridge, UK, 2021).
- [9] A. Griffin, D. W. Snoke, and S. Stringari, eds., *Bose-Einstein Condensation* (Cambridge University Press, Cambridge, UK, 1995).
- [10] N. P. Proukakis, D. W. Snoke, and P. B. Littlewood, eds., *Universal Themes of Bose-Einstein Condensation* (Cambridge University Press, Cambridge, UK, 2017).
- [11] M. Inguscio, W. Ketterle, and C. Salomon, eds., *Ultracold Fermi Gases* (IOS Press, Amsterdam, Netherlands, 2008).
- [12] K. Levin, Q. Chen, C.-C. Chien, and Y. He, *Ann. Phys.* **325**, 233 (2010).
- [13] W. Zwerger, ed., *The BCS-BEC Crossover and the Unitary Fermi Gas* (Springer-Verlag, Berlin, Germany, 2012).
- [14] M. Randeria and E. Taylor, *Annu. Rev. Condens. Matter Phys.* **5**, 209 (2014).
- [15] Y. Ohashi, H. Tajima, and P. van Wyk, *Prog. Part. Nucl. Phys.* **111**, 103739 (2020).
- [16] A. J. Leggett, in *Modern Trends in the Theory of Condensed Matter* (Springer-Verlag, Berlin, Germany, 1980), pp. 13–27.
- [17] K. W. Madison, F. Chevy, W. Wohlleben, and J. Dalibard, *Phys. Rev. Lett.* **84**, 806 (2000).
- [18] S. Eckel, J. G. Lee, F. Jendrzejewski, N. Murray, C. W. Clark, C. J. Lobb, W. D. Phillips, M. Edwards, and G. K. Campbell, *Nature* **506**, 200 (2014).
- [19] R. Desbuquois, L. Chomaz, T. Yefsah, J. Leonard, J. Beugnon, C. Weitenberg, and J. Dalibard, *Nat. Phys.* **8**, 645 (2012).
- [20] L. Sobirey, N. Luick, M. Bohlen, H. Biss, H. Moritz, and T. Lompe, *Science* **372**, 844 (2021).
- [21] M. G. Ries, A. N. Wenz, G. Zurn, L. Bayha, I. Boettcher, D. Kedar, P. A. Murthy, M. Neidig, T. Lompe, and S. Jochim, *Phys. Rev. Lett.* **114**, 230401 (2015).
- [22] I. Boettcher, L. Bayha, D. Kedar, P. A. Murthy, M. Neidig, M. G. Ries, A. N. Wenz, G. Zurn, S. Jochim, and T. Enss, *Phys. Rev. Lett.* **116**, 045303 (2016).
- [23] Z. Wu, L. Zhang, W. Sun, X.-T. Xu, B.-Z. Wang, S.-C. Ji, Y. Deng, S. Chen, X.-J. Liu, and J.-W. Pan, *Science* **354**, 83 (2016).
- [24] R. J. Fletcher, M. Robert-de-Saint-Vincent, J. Man, N. Navon, R. P. Smith, K. G. H. Viebahn, and Z. Hadzibabic, *Phys. Rev. Lett.* **114**, 255302 (2015).
- [25] P. A. Murthy, I. Boettcher, L. Bayha, M. Holzmann, D. Kedar, M. Neidig, M. G. Ries, A. N. Wenz, G. Zurn, and S. Jochim, *Phys. Rev. Lett.* **115**, 010401 (2015).
- [26] C.-A. Chen and C.-L. Hung, *Phys. Rev. Lett.* **127**, 023604 (2021).
- [27] S. S. Botelho and C. A. R. Sa de Melo, *Phys. Rev. Lett.* **96**, 040404 (2006).
- [28] G. Bertaina and S. Giorgini, *Phys. Rev. Lett.* **106**, 110403 (2011).
- [29] L. He and X.-G. Huang, *Phys. Rev. Lett.* **108**, 145302 (2012).
- [30] C.-C. Chien, J.-H. She, and F. Cooper, *Ann. Phys.* **347**, 192 (2014).
- [31] L. He, H. Lu, G. Cao, H. Hu, , and X.-J. Liu, *Phys. Rev. A* **92**, 023620 (2015).
- [32] C.-T. Wu, B. M. Anderson, R. Boyack, and K. Levin, *Phys. Rev. Lett.* **115**, 240401 (2015).
- [33] N. Lundblad, R. A. Carollo, C. Lannert, M. J. Gold, X. Jiang, D. Paseltiner, N. Sergay, and D. C. Aveline, *npj Microgravity* **5**, 30 (2019).
- [34] R. Carollo, D. Aveline, B. Rhyno, S. Vishveshwara, C. Lannert, J. Murphree, E. Elliott, J. Williams, R. Thompson, and N. Lundblad, *Observation of ultracold atomic bubbles in orbital microgravity* (2021), arXiv: 2108.05880.
- [35] A. Tononi and L. Salasnich, *Phys. Rev. Lett.* **123**, 160403 (2019).
- [36] A. Tononi, F. Cinti, and L. Salasnich, *Phys. Rev. Lett.* **125**, 010402 (2020).
- [37] N. S. M ller, F. E. A. dos Santos, V. S. Bagnato, and A. Pelster, *New J. Phys.* **22**, 063059 (2020).
- [38] S. J. Bereta, M. A. Caracanhas, and A. L. Fetter, *Phys. Rev. A* **103**, 053306 (2021).
- [39] A. Tononi, A. Pelster, and L. Salasnich, *Topological superfluid transition in bubble-trapped condensates* (2021), arXiv:2104.04585.
- [40] B. Rhyno, N. Lundblad, D. C. Aveline, C. Lannert, and S. Vishveshwara, *Thermodynamics in expanding shell-shaped bose-einstein condensates* (2021), arXiv: 2106.00835.
- [41] A. Andriati, L. Brito, L. Tomio, and A. Gammal, *Stability of a bose condensed mixture on a bubble trap* (2021), arXiv: 2107.04130.
- [42] D. Cricchio, E. Fiordilino, and F. Persico, *Phys. Rev. A* **86**, 013201 (2012).
- [43] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw-Hill Book Company, New York, 1971).
- [44] S. K. Adhikari, *Am. J. Phys.* **54**, 362 (1986).
- [45] F. Werner and Y. Castin, *Phys. Rev. A* **86**, 013626 (2012).
- [46] B. Mihaila, J. F. Dawson, F. Cooper, C.-C. Chien, and E. Timmermans, *Phys. Rev. A* **83**, 053637 (2011).
- [47] V. L. Berezinskii, *Sov. Phys. JETP* **32**, 493 (1971).
- [48] V. L. Berezinskii, *Sov. Phys. JETP* **34**, 610 (1972).
- [49] J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **6**, 1181 (1973).

Calculation of V

The interaction $V = -g\delta(1 - \cos\theta_{12})$ can be expanded by the Legendre polynomials and spherical harmonic functions as

$$\begin{aligned} V &= -g \sum_L (2L+1) P_L(\cos\theta_{12}) \\ &= -4\pi g \sum_{LM} (-1)^M Y_{LM}(\theta_1, \phi_1) Y_{L,-M}(\theta_2, \phi_2). \end{aligned} \quad (10)$$

Here we treat Y_{LM} as an irreducible tensor operator, so the summation

$$\sum_{LM} (-1)^M Y_{LM}(\theta_1, \phi_1) Y_{L,-M}(\theta_2, \phi_2) \quad (11)$$

is actually a tensor product of two irreducible tensor operators, which results in a scalar operator.

According to a more general version of the Wigner-Eckart theorem shown in Eq. (B.33) of Ref. [43], we find that

$$\begin{aligned} \langle l_1 l_1 00 | V | l_2 l_2 00 \rangle &= \\ -4\pi g \sum_L (-1)^{l_1+l_2} \left\{ \begin{matrix} 0 & l_1 & l_1 \\ L & l_2 & l_2 \end{matrix} \right\} \langle l_1 || Y_L || l_2 \rangle^2. \end{aligned} \quad (12)$$

Here the 6j symbol is given by

$$\left\{ \begin{matrix} 0 & l_1 & l_1 \\ L & l_2 & l_2 \end{matrix} \right\} = (-1)^{l_1+l_2} \frac{1}{\sqrt{(2l_1+1)(2l_2+1)}} \quad (13)$$

and reduced matrix element is

$$\begin{aligned} \langle l_1 || Y_L || l_2 \rangle &= \\ (-1)^{l_1} \sqrt{\frac{(2l_1+1)(2L+1)(2l_2+1)}{4\pi}} \begin{pmatrix} l_1 & L & l_2 \\ 0 & 0 & 0 \end{pmatrix}. \end{aligned} \quad (14)$$

In the above equation, the large parentheses denote the 3j symbol, not to be confused with the 6j symbol. After collecting all the results, we find that

$$\begin{aligned} \langle l_1 l_1 00 | V | l_2 l_2 00 \rangle &= -g \sum_L \sqrt{(2l_1+1)(2l_2+1)} \\ &\quad \times (2L+1) \begin{pmatrix} l_1 & L & l_2 \\ 0 & 0 & 0 \end{pmatrix}^2. \end{aligned} \quad (15)$$

Moreover, the normalization condition of CG coefficients lead to

$$\sum_L (2L+1) \begin{pmatrix} l_1 & L & l_2 \\ 0 & 0 & 0 \end{pmatrix}^2 = \sum_L \langle l_1 l_2 00 | l_1 l_2 L 0 \rangle^2 = 1 \quad (16)$$

After some algebra, the matrix element takes the form

$$\langle l_1 l_1 00 | V | l_2 l_2 00 \rangle = -g \sqrt{(2l_1+1)(2l_2+1)}. \quad (17)$$




Figure 4. Gap and chemical potential as functions of $-\ln(k_F a)$ from Eq. (8) (red circles) using the integral and from Eq. (22) (blue squares) using the summation. Here $N = 220$ and $k_F R = 10.5$.

Alternative renormalization scheme

There is another way of regularizing the gap equation of a Fermi superfluid on a 2D plane. This has been shown in Eq. (12) of Ref. [45] as follows.

$$-\frac{1}{g} = \lim_{q \rightarrow 0} \left[-\frac{m}{2\pi} \ln\left(\frac{aqe^\gamma}{2}\right) + \int \frac{d^2 k}{(2\pi)^2} \mathcal{P} \frac{1}{2(\epsilon_{\mathbf{q}} - \epsilon_{\mathbf{k}})} \right] \quad (18)$$

Here γ is the Euler constant and \mathcal{P} denotes the Cauchy principle value. Combining with the gap equation of the 2D Fermi superfluid, one finds

$$\begin{aligned} -\frac{m}{2\pi} \ln\left(\frac{aqe^\gamma}{2}\right) &= \int \frac{d^2 k}{(2\pi)^2} \frac{1 - f(E_{\mathbf{k}})}{2E_{\mathbf{k}}} + \\ &\quad \int \frac{d^2 k}{(2\pi)^2} \mathcal{P} \frac{1}{2(\epsilon_{\mathbf{q}} - \epsilon_{\mathbf{k}})}. \end{aligned} \quad (19)$$

The drawback of this method, however, is that one has to assume an infrared (IR) cutoff q . We have verified that this alternative regularization gives qualitatively the same results as those presented in the main text.

Universal behavior

The universal behavior of the gap and chemical potential in the interaction-induced BCS-BEC crossover comes from the equations of state. We let $b = -\ln(k_F a)$, which is equivalent to $a = e^{-b}/k_F$. For the 2D-plane case, $E_F = k_F^2$ when $\hbar = 1$ and $2m = 1$. The particle number per unit area of a noninteracting Fermi gas is given by $N = 2(\pi k_F^2)/(2\pi)^2$, or $n/k_F^2 = 1/(2\pi)$. Therefore, the gap and number equations of Fermi superfluid on a 2D plane can be written as

$$\begin{aligned} \int dy \left[\frac{1 - f(E_k/E_F)}{2E_k/E_F} - \frac{1}{2y^2 + 2e^{2b}} \right] &= 0. \\ \frac{1}{2\pi} &= \int \frac{dy y}{2\pi} \left(1 - \frac{\xi_k/E_F}{E_k/E_F} + 2 \frac{\xi_k/E_F}{E_k/E_F} f(E_k/E_F) \right). \end{aligned} \quad (20)$$

Here $y = k/k_F$ and only Δ/E_F and μ/E_F show up in E_k/E_F and ξ_k/E_F .

Meanwhile, for a noninteracting Fermi gas on a sphere filled up to the angular-momentum state L_m , we have $n = N/(4\pi R^2)$, $N = 2L_m(L_m + 1)$, $E_F = N/(2R^2)$, and $E_F = k_F^2$ with $2m = 1$. Again, let $b = -\ln(k_F a)$. The equations of state of Fermi superfluid on a spherical shell thus becomes

$$\int dz \left(z + \frac{1}{2L_m} \right) \left[\frac{1 - f(E_l/E_F)}{2E_l/E_F} - \frac{1}{2z^2 + 2e^{2b}} \right] = 0, \quad (21)$$

$$\frac{1}{2\pi} = \int \frac{dz}{2\pi} \left(z + \frac{1}{2L_m} \right) \left(1 - \frac{\xi_l/E_F}{E_l/E_F} + 2 \frac{\xi_l/E_F}{E_l/E_F} f(E_l/E_F) \right).$$

Here $z = l/L_m$ and only Δ/E_F and μ/E_F show up in E_l/E_F and ξ_l/E_F . When $L_m \gg 1$, which is usually the case in many-body systems, the two sets of equations of state, Eqs. (20) and (21), are identical and give the universal behavior of the normalized gap and chemical potential.

Integral vs. summation in the equations

Here we compare the results from the gap and number equations using summation over the angular momentum versus the approximation using integration. The equations with explicit summations are

$$\sum_{l=0}^{L_M} (2l+1) \left[\frac{1 - 2f(E_l)}{2E_l} - \frac{1}{2\epsilon_l + |\epsilon_b|} \right] = 0, \quad (22)$$

$$N = \sum_{l=0}^{L_M} (2l+1) \left(1 - \frac{\xi_l}{E_l} + 2 \frac{\xi_l}{E_l} f(E_l) \right).$$

Here L_M is some cutoff level, which is much larger than the highest occupied shell L_m . We present an example with $L_M = 100$ and $l_m = 10$, which is the counterpart of Fig. 1. After solving the gap and μ using summations, we plot the results in Fig. 4 along with the results from the integrals. One can see that they are virtually identical, thereby justifying the approximation of replacing the summation over the angular momentum with integration.

Derivation of superfluid density

The superfluid density on a spherical shell can be deduced from the expression of the 2D planar case. We remark that the superfluid density is derived from linear response theory [43] instead of thermodynamics. Explicitly, n_s can be extracted from the London equation $j_\mu = -\frac{n_s}{m} A_\mu$, where j_μ and A_μ denote the current and four-potential. From linear response theory, the current can be written as

$$j_\mu(\mathbf{k}, \omega) = -K_{\mu\nu}(\mathbf{k}, \omega) A_\nu(\mathbf{k}, \omega), \quad (23)$$

$$K_{\mu\nu}(\mathbf{k}, \omega) = \frac{n}{m} \delta_{\mu\nu} - i \langle [J_\mu(\mathbf{k}, \omega), J_\nu(-\mathbf{k}, -\omega)] \rangle.$$

Here $J_\mu(\mathbf{k}, \omega)$ is the current operator and $\langle \dots \rangle$ denotes the ensemble average. The current-current correlation function can be obtained by analytical continuation from the corresponding Matsubara formula. In the static and uniform limit with $\omega = 0$ and $\mathbf{k} \rightarrow 0$, the result is simplified to

$$\lim_{\mathbf{k} \rightarrow 0} -i \langle [J_\mu(\mathbf{k}, 0), J_\nu(-\mathbf{k}, 0)] \rangle = \frac{1}{m^2} \sum_{\mathbf{k}} k^2 \frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \delta_{\mu\nu} \quad (24)$$

After collecting all the above results, we find the n_s of BCS theory on a 2D plane as

$$n_s = n - \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} \left[-\frac{\partial f(E_{\mathbf{k}})}{\partial E_{\mathbf{k}}} \right]. \quad (25)$$

To generalize the expression to the spherical case, we make the following replacements:

$$\sum_{\mathbf{k}} \rightarrow \frac{1}{4\pi R^2} \int dl \, 2(2l+1), \quad \epsilon_{\mathbf{k}} \rightarrow \frac{l(l+1)}{2mR^2}, \quad (26)$$

Afterwards, we arrive at $n_s = n - \frac{1}{4\pi R^2} \int dl \, 2(2l+1) \frac{l(l+1)}{2mR^2} \left[-\frac{df(E_l)}{dE_l} \right]$ as shown in the main text.