

# Non-local gravity in bouncing cosmology scenarios

D. Jackson,<sup>a</sup> R. Bufalo,<sup>b,1</sup>

<sup>a</sup>Instituto de Física Teórica, IFT-UNESP,

R. Dr. Bento Teobaldo Ferraz, 271, Várzea da Barra Funda, São Paulo - SP, 01140-070, Brazil

<sup>b</sup>Departamento de Física, Universidade Federal de Lavras

Caixa Postal 3037, 37200-900 Lavras, MG, Brazil

E-mail: [dimas.jackson@unesp.br](mailto:dimas.jackson@unesp.br), [rodrigo.bufalo@ufla.br](mailto:rodrigo.bufalo@ufla.br)

**Abstract.** In this work, we analyzed the improved Deser-Woodard non-local gravity over the background of five different bouncing cosmologies, whose premise is to avoid the initial singular state of the universe. We developed the numerical solution for the non-local distortion function, which encompass the modifications to the Einstein-Hilbert action, using the reconstruction procedure and we have found that three classes of bouncing have a viable cosmological solution. Afterwards, we discussed the physical aspects and outcomes of the evolution of the distortion function throughout the bouncing point for three physical classes specifically: the symmetric bounce, the super bounce and the Big Rip universe bounce.

**Keywords:** cosmic singularity, quantum cosmology, modified gravity.

---

<sup>1</sup>Corresponding author.

---

## Contents

|          |                                       |           |
|----------|---------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                   | <b>1</b>  |
| <b>2</b> | <b>Improved Deser-Woodard gravity</b> | <b>2</b>  |
| <b>3</b> | <b>Application to bouncing models</b> | <b>4</b>  |
| 3.1      | Symmetric Bounce                      | 4         |
| 3.2      | Super Bounce                          | 7         |
| 3.3      | Big Rip                               | 9         |
| <b>4</b> | <b>Nonviable bouncing solutions</b>   | <b>10</b> |
| <b>5</b> | <b>Conclusions</b>                    | <b>11</b> |

---

## 1 Introduction

Despite all of the success in comparison of observational data and consistency of the theory in mathematical terms [1], the Einstein’s theory of General Relativity (GR) present some limitations. A well-known example is the equivalence principle, which is not directly applicable to the initial instant of the universe (in a model of the “big bang” type). Furthermore, there is a break in the necessary conditions for its application due to a singularity of space-time, i. e., this initial state has density, pressure and temperatures approximately infinite [2, 3]. In fact, there is no consensus on how to define metric singularities in GR, unlike electromagnetism for example, in which a singularity of the electromagnetic field can be avoided by defining the field in all the space except the singular point. In the GR there is not a background spacetime to make this type of definition, since it is the spacetime itself which becomes singular [4].

In order to address the aforementioned space-time singularity, we would like to bring attention to the proposal of bouncing cosmologies [5–10], where the cosmic contraction reduces the effective radius of the cosmos to a minimum size which then produces an expanding Universe, replacing effectively the big bang singularity by a nonsingular bouncing geometry in which all curvature invariants become finite. Moreover, bouncing cosmologies have also been considered as strong candidates to the standard inflationary paradigm [11, 12].

The current standard cosmological model, namely  $\Lambda$ -Cold Dark Matter ( $\Lambda$ -CDM), is based in the GR. In this model the universe starts in a singular point and after the early epoch, the vacuum energy dominates driving the accelerated expansion. The presence of unknown energy sources in the  $\Lambda$ -CDM model indicates the possibility that General Relativity is not enough to explain the aforementioned phenomena [13]. Thus, several modified gravitational theories were proposed as attempts to generalize Einstein’s gravitational theory. Usually, the most used approach for the description of such phenomena involve addition of new (global and/or local) degrees of freedom, which can be described in terms of new fields or even by enforcing a symmetry principle.

Nowadays, an interesting point of view used to elaborate a modified gravitational theory are the theories with presence of *non-local effects*,<sup>1</sup> which emerged from the works of D.

---

<sup>1</sup>For a detailed account regarding conceptual non-local aspects in field theories, for instance causality, domain of validity, boundary conditions, etc, see ref. [14] and the references therein for a review.

Dalvit [15] and C. Wetterich [16]. Actually, Wetterich proposed adding a term proportional to  $R\Box^{-1}R$ <sup>2</sup> to the Einstein-Hilbert action and analyzed the cosmological implications of a simple model based on this non-local action. He concluded that his model was not compatible with primordial nucleosynthesis, whereas generalizations consistent with experimental observations would be conceivable. Among the many interesting proposals of non-local modifications to the Einstein-Hilbert action [14, 21, 22], we may refer to the S. Deser and R. Woodard non-local model, where they have developed a theory with more general non-local terms of the form  $Rf(\Box^{-1}R)$  [23]. They have found that this model contains the time interval for the transition from radiation dominated universe to the matter dominated one, but several issues were still open. A decisive test for this model was that the screening mechanism (necessary to avoid the presence of non-local effects in the Solar System scale) is not satisfied, violating thus experimental constraints [14]. Afterwards, Deser and Woodard improved their model [24] in order to satisfy the screening mechanism and also to reproduce the late time accelerated expansion without the necessity of a cosmological constant.

Naturally, due to the rich phenomenological scenario presented by the bouncing cosmology, it is very interesting to explore whether modified gravitational models provide viable cosmological solution in this context, see [25] and the references therein for a review. Actually, the first Deser-Woodard non-local cosmological model provided a viable description of a bouncing universe, which can be smoothly connected to the accelerated expansion scenario [26]. Hence, our proposal consists in examining the viability of bouncing cosmologies within the improved non-local Deser-Woodard framework where we consider the five most studied bouncing cosmology scenarios, namely symmetric bounce, superbounce, oscillatory bouncing cosmology, matter bounce, and the singularity cases [25]. The latter encompasses Big Rip singularities, sudden singularities and the Big Freeze universe, all generated by the same scale factor but with different parameters.

In the present work we analyze the description of bouncing universe within the improved non-local Deser-Woodard model. This is achieved by considering a flat Friedmann-Lemaître-Robertson-Walker (FLRW) geometry in which the bouncing frameworks emerge through the choice of particular forms of the scale factor. In Sec. 2 we review the main features of the improved Deser-Woodard model, in special, focusing in obtaining the equations of motion for the non-local distortion function and the auxiliary fields. In Sec. 3 we determine the solutions of three different classes of bouncing: symmetric, super bounce and big rip, and analyze how they are physically viable cosmological scenarios. At last, in Sec. 4 we discuss the issues related with the oscillatory and matter bounces, and also with sudden singularities and big freeze universe within the improved Deser-Woodard model, and establish the main points which render a nonphysical cosmological evolution. We present our final remarks and perspectives in Sec. 5.

## 2 Improved Deser-Woodard gravity

The improved non-local Deser-Woodard model [24] minimally modifies the Einstein-Hilbert action, introducing two non-local scalar fields  $(X, Y)$ , such that

$$\mathcal{L} = \frac{1}{16\pi G} R [1 + f(Y)], \quad (2.1)$$

---

<sup>2</sup>It is worth mentioning that this kind of non-local terms are induced in loop corrections of the quantum effective action [17–20].

with the following definitions

$$Y = \square^{-1} g^{\mu\nu} \partial_\mu X \partial_\nu X, \quad (2.2a)$$

$$X = \square^{-1} R. \quad (2.2b)$$

Here  $\square = g^{\mu\nu} D_\mu D_\nu$  is the covariant d'Alembertian operator and  $R$  is the curvature scalar. Along this paper we will refer to the model described by the Lagrangian density (2.1) as DW II. This model was originally proposed to reproduce the late time accelerated expansion of the universe, without the need of the cosmological constant, only via a dynamical mechanism inspired by quantum effective action corrections [24]. Actually, the term  $g^{\mu\nu} \partial_\mu X \partial_\nu X$  is negative in the Solar System scale and positive in the cosmological scale, which makes viable a screening effect. As a consequence of this mechanism, the non-local corrections appear only on the cosmological background and the GR is recovered on strongly bound gravitational systems. Thus, DW II obey the well established Solar System constraints [27]

To localize the action, one can introduce two auxiliary scalar fields  $U$  and  $V$  as Lagrange multipliers,

$$\mathcal{L} = \frac{1}{16\pi G} [R(1 + f(Y)) + U(R - \square X) + V(g^{\mu\nu} \partial_\mu X \partial_\nu X - \square Y)], \quad (2.3)$$

or, equivalently

$$\mathcal{L} = \frac{1}{16\pi G} [R(1 + f(Y) + U) + g^{\mu\nu} E_{\mu\nu}], \quad (2.4)$$

where we have introduced  $E_{\mu\nu} = \partial_\mu U \partial_\nu X + \partial_\mu V \partial_\nu Y + V \partial_\mu X \partial_\nu X$ , by means of simplicity of notation. Hence, considering the set  $(X, Y, U, V)$  as four independent scalar fields, the action  $S = \int d^4x \sqrt{-g} \mathcal{L}$  is regarded as local. Thus, varying the action in respect to each of these fields, result into the following set of equations

$$R - \square X = 0, \quad (2.5a)$$

$$g^{\mu\nu} \partial_\mu X \partial_\nu X - \square Y = 0, \quad (2.5b)$$

$$2D^\mu (V D_\mu X) + \square U = 0, \quad (2.5c)$$

$$R \frac{\partial f(Y)}{\partial Y} - \square V = 0. \quad (2.5d)$$

The gravitational field equations are obtained by varying the action (2.4) in respect to the metric  $g^{\mu\nu}$ :

$$(G_{\mu\nu} - D_\mu D_\nu + g_{\mu\nu} \square) (1 + U + f(Y)) + E_{(\mu\nu)} - \frac{1}{2} g_{\mu\nu} g^{\rho\sigma} E_{\rho\sigma} = 8\pi G T_{\mu\nu}, \quad (2.6)$$

where the energy momentum tensor  $T_{\mu\nu} = (\rho + p) u_\mu u_\nu + p g_{\mu\nu}$  corresponds to the usual baryonic matter and does not include the dark energy source term <sup>3</sup>.

This model reproduces the current accelerated expansion of the universe without cosmological constant for

$$f(Y) \sim e^{1.1(Y+16.7)}. \quad (2.7)$$

---

<sup>3</sup>The indices in parenthesis denote the symmetric part  $E_{(\mu\nu)} = \frac{1}{2} (E_{\mu\nu} + E_{\nu\mu})$ .

This is an exponential fit to the numerical solution obtained through the reconstruction process [24]. The function  $f(Y)$  is named *non-local distortion function* and the procedure of reconstruction consists in requiring that the Friedmann equations of General Relativity should be satisfied by the DW II.

To perform this procedure, the field equations (2.6) have to be expanded over the Friedmann-Lemaître-Robertson-Walker (FLRW) background:

$$ds^2 = dt^2 - a^2(t)dx_i dx^i. \quad (2.8)$$

The d'Alembertian operator acting on a scalar function  $W(t)$ , which depends only on time, is written as

$$\square W(t) = \partial_t^2 W(t) + 3H\partial_t W(t), \quad (2.9)$$

where  $H = \frac{\dot{a}}{a}$  is the Hubble parameter. In this background, the (00) and (ij) components of the modified field equations (2.6) become

$$(3H^2 + 3H\partial_t)(1 + U + f(Y)) + \frac{1}{2}(\dot{X}\dot{U} + \dot{Y}\dot{V} + V\dot{X}^2) = 8\pi G\rho, \quad (2.10a)$$

$$- [2\dot{H} + 3H^2 + \partial_t^2 + 2H\partial_t](1 + U + f(Y)) + \frac{1}{2}(\dot{X}\dot{U} + \dot{Y}\dot{V} + V\dot{X}^2) = 8\pi Gp. \quad (2.10b)$$

Furthermore, subtracting the equations (2.10a) and (2.10b) we find a differential equation for the function  $F(t) \equiv 1 + U(t) + f[Y(t)]$ :

$$[2\dot{H} + 6H^2 + \partial_t^2 + 5H\partial_t]F(t) = 8\pi G(\rho - p). \quad (2.11)$$

Hence, the non local distortion function can be numerically obtained through the relation  $f = F - U - 1$ . We will perform this analysis in the next sections for some different bouncing universe solutions.

### 3 Application to bouncing models


In this section we will discuss the application of the model DW II to different bouncing models. In special, we shall focus in the numerical analysis of the viability of the cosmological evolution near the bouncing point ( $t = 0$ ) of five different bounce classes: symmetric, oscillatory, super bounce, matter bounce and singular bounce settings [25]. However, only the symmetric, super bounce and big rip are in fact viable cosmological scenarios for DW II, see discussion below. Furthermore, these (physical) bouncing models are illustrated in Figure 1 in terms of their scale factor.

#### 3.1 Symmetric Bounce

The symmetric bounce model was studied in the context of the non-local DW I gravity [26], where it was considered as a viable cosmological scenario. The symmetric bounce is characterized by the following scale factor [25]

$$a(t) = a_m e^{h_1 t^2/2}, \quad (3.1)$$

where  $a_m$  is the minimum value of the scale factor and  $h_1$  is a constant, such that the Hubble parameter is  $H(t) = h_1 t$ . This scale factor is shown in panel (a) of the Figure 1. We search



**Figure 1:** The scale factor  $a(t)$  for three different bouncing models: (a) symmetric bounce, (b) super bounce and (c) big rip bounce, respectively.

for vacuum solutions of the equation (2.11) i.e.  $\rho = p = 0$ , so that

$$\left[2\dot{H} + 6H^2 + \partial_t^2 + 5H\partial_t\right] F(t) = 0. \quad (3.2)$$

In the symmetrical bounce class, the differential equation (3.2) has an exact solution, given by

$$F(t) = A_1 e^{-h_1 t^2} + A_2 H_{-1} \left( \sqrt{\frac{h_1}{2}} t \right), \quad (3.3)$$

where  $H_n$  is the Hermite polynomial (with negative index) and  $A_1, A_2$  are constants of integration. Using the relation

$$H_{-1}(x) = \frac{\sqrt{\pi}}{2} e^{x^2} \operatorname{erfc}(x), \quad (3.4)$$

where  $\operatorname{erfc}(x) \equiv 1 - \operatorname{erf}(x) = 1 - \frac{2}{\sqrt{\pi}} \int_0^x e^{-u^2} du$  is the Gauss's complementary error function, the solution for  $F(t)$  in eq. (3.3) can be written in a suitable form as

$$F(t) = e^{-h_1 t^2} \left( A_1 + A_2 \operatorname{erfc} \left( \sqrt{\frac{h_1}{2}} t \right) \right). \quad (3.5)$$

Finally, in order to obtain the non-local distortion function  $f = 1 - U - F$ , it is necessary to find first the solutions for the fields  $X, Y, U$  and  $V$ . Actually, one can obtain a differential equation for the scalar field  $X$  by applying the background FLRW metric (2.9) to the constraint  $\square X = R$ , eq. (2.5a):

$$(\partial_t^2 + 3H\partial_t) X + 6(\dot{H} + 2H^2) = 0. \quad (3.6)$$

Its exact solution is given in terms of well known special functions:

$$X(t) = B_1 + B_2 \operatorname{erf} \left( \sqrt{\frac{3h_1}{2}} t \right) - h_1 t^2 \left( 2 + {}_2F_2 \left( 1, 1; \frac{3}{2}, 2; -\frac{3}{2} h_1 t^2 \right) \right), \quad (3.7)$$

where  ${}_2F_2 \left( 1, 1; \frac{3}{2}, 2; -\frac{3}{2} h_1 t^2 \right)$  is the generalized hyper-geometric function and  $B_1, B_2$  are constants of integration. Next, we can make use of the eq. (2.5b) to write a differential equation for  $Y$  in terms of  $X$  as

$$(\partial_t^2 + 3H\partial_t) Y - \dot{X}^2 = 0, \quad (3.8)$$

but in this case, by making use of the solution eq. (3.7), we can determine only numerical solutions for  $Y$ . Furthermore, from the definition of  $U$  given in (2.5c), over the background metric FLRW, we have

$$(\partial_t + 3H)\dot{U} = -2(\partial_t + 3H)V\dot{X}, \quad (3.9)$$

which can be equivalently expressed as the following relation

$$-2V = \frac{\dot{U}}{\dot{X}}. \quad (3.10)$$

At last, over the background metric FLRW, the expression for  $V$  in (2.5d) becomes

$$(\partial_t^2 + 3H\partial_t)V = -6\left(\dot{H} + 2H^2\right)\frac{df}{dY}. \quad (3.11)$$

However, since the function  $F = 1 + f + U$  was calculated in (3.5), then, we can eliminate  $f$  by

$$\frac{df}{dY} = \frac{\dot{F} - \dot{U}}{\dot{Y}}. \quad (3.12)$$

Thus, replacing (3.12) back into (3.11), it results in the following

$$(\partial_t^2 + 3H\partial_t)V + 6\left(\dot{H} + 2H^2\right)\left(\frac{\dot{F} - \dot{U}}{\dot{Y}}\right) = 0. \quad (3.13)$$

It is important to emphasize that this step imposes the constraint  $\dot{Y} \neq 0$  for all  $t$  in the solution's interval, otherwise  $V$ ,  $U$  and hence  $f$  will diverge everywhere. Hence, substituting  $\dot{U}$  from equation (3.10), one can finally obtain a differential equation for the auxiliary field  $V$ :

$$(\partial_t^2 + 3H\partial_t)V + 6\left(\dot{H} + 2H^2\right)\left(\frac{\dot{F} + 2V\dot{X}}{\dot{Y}}\right) = 0. \quad (3.14)$$

The numerical solution for the field  $V$  provides the necessary information to perform the numerical calculation of  $U$  through the equation (3.10). Henceforth, the non-local distortion function  $f = F - U - 1$  is constructed from the numerical solution for  $U$  and the exact equation (3.5) for  $F$ . The initial conditions used to perform these calculations were

$$F(0) = 3, \quad \dot{F}(0) = 0, \quad (3.15a)$$


$$X(0) = 0, \quad \dot{X}(0) = 0, \quad (3.15b)$$

$$Y(0) = 0, \quad \dot{Y}(6) = 1, \quad (3.15c)$$

$$V(5) = 1, \quad \dot{V}(5) = 1, \quad (3.15d)$$


$$U(0) = 0. \quad (3.15e)$$

The choice of the condition  $\dot{Y} \approx 0$  near the interval's solution endpoint, was necessary in order to avoid that  $\dot{Y}(t)$  assumes the value zero and resulting in divergences in the functions  $U$  and  $V$ . The numerical solution for the distortion function is presented in the Figure 2. In this plot we can see the evolution of the distortion function throughout the collapse and rebirth of the universe. In the symmetric bounce,  $f(t)$  is approximately constant before and after the bounce, suffering a change of values at  $t = 0$ . Observe that the distortion function is positive in the collapsing universe and becomes negative in the expansion scenario.



**Figure 2:** Numerical solution of the non-local distortion function  $f(t)$  in the symmetrical bounce, in terms of the cosmological time.

Finally, the solutions  $f(t)$  and  $Y(t)$  are used to obtain  $f(Y)$ . The numerical solution for  $f(Y)$  is depicted in the Figure 3 and corresponds to the non-local correction to Einstein-Hilbert action, as introduced by equation (2.1). It is important to note that the axes in panel (a) of Figure 3 are out of scale, so, in spite of  $f(Y)$  have a negative slope, this function varies slowly compared to the values of  $Y$ . This can be better seen in the panel (b) of Figure 3, which is in the logarithmic scale. Consequently, the modification arising from the non-local phenomena to the Einstein-Hilbert action in the symmetric bounce, is roughly an additive (very large) constant.



**Figure 3:** Non-local distortion function in the context of the symmetric bouncing, (a) in linear scale and (b) in logarithmic scale.

### 3.2 Super Bounce

Super bounce cosmologies are models of the universe which collapses and grows again through a Big Bang without a singularity, and also the scale factor, at the bouncing point, is identically zero. This cosmology is characterized by the scale factor [25]

$$a(t) = \left( \frac{t - t_s}{t_0} \right)^{\frac{2}{k^2}}. \quad (3.16)$$

Here  $t_s$  is the bouncing time and  $t_0$  is a characteristic time in which the scale factor reaches the value  $a = 1$ . We present this scale factor near the bouncing point  $t = t_s = 0$  in the panel (b) of Figure 1.

The calculation of the non-local distortion function  $f(Y)$  follows the same development as presented in the previous section; nonetheless, in this class of bounce, the fields  $F$ ,  $X$  and  $Y$  have exact solution as it follows:

$$F(t) = |t|^{1/4}C_1 + C_2\sqrt{|t|}, \quad (3.17a)$$

$$X(t) = 4|t|^{1/4}C_3 + C_4 - 3 \ln |t|, \quad (3.17b)$$

$$Y(t) = C_6 + 4|t|^{1/4}(24C_3 + 2|t|^{1/4}C_3^2 + C_5) - 12(3 + 2|t|^{1/4}C_3) \ln |t|, \quad (3.17c)$$

where  $C_k$ ,  $k = 1, 2, 3, 4, 5, 6$ , are constants of integration. Similarly to what happened in the analysis of the symmetric bounce, the auxiliary fields  $U$  and  $V$  have only numerical solution. The initial conditions used in the numerical calculation were

$$F(\pm 1) = 1, \quad \dot{F}(\pm 1) = 1, \quad (3.18a)$$


$$X(\pm 1) = 0, \quad \dot{X}(\pm 1) = 0, \quad (3.18b)$$

$$Y(\pm 6) = 0, \quad \dot{Y}(\pm 6) = 0, \quad (3.18c)$$

$$V(\pm 5) = 1, \quad \dot{V}(\pm 5) = 1, \quad (3.18d)$$


$$U(\pm 5) = 1. \quad (3.18e)$$

As a result, we found the non-local distortion function presented in the panel (a) of the Figure 4, in terms of the cosmological time. In this case  $f(t)$  grows to  $+\infty$  when  $t \rightarrow 0^+$  and goes to  $-\infty$  at  $t \rightarrow 0^-$ . Hence, the distortion function evolution is different at each phase of the universe and present an infinite like discontinuity at the bouncing point.



**Figure 4:** Numerical solution of the non-local distortion function in the super bounce, (a) in terms of the cosmological time and (b) in terms of the field  $Y$ .

Exploring the one-to-one relation between  $Y$  and  $t$ , follow the numerical solution for  $f(Y)$  depicted in the panel (b) of Figure 4. The orange curve shows the evolution of the non-local effects for the past universe (negative  $t$ ) and the blue curve for the present universe (positive  $t$ ). For  $t < 0$ , we observe that the distortion function starts from  $+\infty$  at the starting point of the interval's solution, stabilizes near the origin and decreases as long as  $t \rightarrow 0^-$  ( $Y$  decreases). After the bouncing point,  $f(Y)$  starts from  $+\infty$ , oscillates and go down to  $-\infty$  at the endpoint  $t = 6$  (as  $Y$  increases). The interval of solution used here,  $t \in [-6, 6]$ , can be extended as much as wished. Therefore, the growth of the non-local distortion function, at the extremes, should be seen as occurring in the late time universe.



**Figure 5:** Numerical solution of the non-local distortion function in the Big Rip bounce, (a) in terms of the cosmological time and (b) in terms of the field  $Y$ .

### 3.3 Big Rip

In this section we will analyze the behavior of the DW II gravity under the singularities scenario, which is generated by the scale factor [25]

$$a(t) = A_0 \exp \left[ f_0 \frac{(t - t_s)^{\alpha+1}}{\alpha + 1} \right], \quad (3.19)$$

where  $A_0 > 0$  is a dimensionless constant that describes the scale factor at the bouncing time and  $f_0 > 0$  is a constant with the dimension of the proper time. The parameter  $\alpha$ , if chosen equal to 1 and  $-1$  correspond to the symmetric bounce and super bounce, respectively. Moreover, the choice  $\alpha = 0$  implies that the scale factor  $a$  corresponds to the de Sitter universe. In order to consider the Big Rip cosmology, characterized by a diverging scale factor at the bouncing point  $t = t_s$ , we make the choice  $\alpha < -1$ . This type of initial singularities can be avoided through mechanisms of hydrodynamics or quantum effects and describes an accelerated expansion at late time [25]. Panel (c) of Figure 1 shows the scale factor (3.19) for the Big Rip scenario. For other values of the parameter  $\alpha$ , which results in sudden singularities and the Big Freeze universe, see the discussion in Section 4.

Once again, the determination of  $f(Y)$  follows the same procedure presented in the previous sections, although, only numerical solutions are available for all the fields in this class of bounce. The initial conditions used in the current analysis are presented below:

$$F(\pm 1) = 3, \quad \dot{F}(\pm 1) = -\sqrt{2/\pi}, \quad (3.20)$$

$$X(\pm 1) = 1, \quad \dot{X}(\pm 1) = 2, \quad (3.21)$$

$$Y(\pm 1) = 1, \quad \dot{Y}(\pm 6) = 0, \quad (3.22)$$

$$V(\pm 5) = 1, \quad \dot{V}(\pm 5) = 1, \quad (3.23)$$

$$U(\pm 5) = 1. \quad (3.24)$$

The resulting non-local distortion function is presented in Figure 5, for the time dependence in panel (a) and for the non-local field  $Y$  in panel (b). The function  $f(t)$  is monotonically increasing for  $t \in (-6, 0)$ , have a jump discontinuity at  $t = 0$  and is not monotonic for  $t \in (0, 6)$ .

Note that  $a(t)$  in (3.19) diverges when  $t \rightarrow 0$ , so  $f(t)$  is not defined at the origin. Moreover, the field  $\dot{Y}$  have almost one zero in the interval of the solution, which is fixed by the initial condition. Our choice of this zero point at the end of the interval is crucial to preserve the fields  $U$  and  $V$  from divergences for all values of  $t$ . Hence, the non-local correction function  $f(Y)$  is defined only in a very narrow interval, in this case it is the values of  $Y$  that appear in panel (b) of Figure 5, i.e.  $Y \in [-32.7, 19.0]$  for  $t < 0$  and  $Y \in [-73.9, 28.5]$  for  $t > 0$ .

#### 4 Nonviable bouncing solutions

In the previous section we have analyzed three classes of bouncing cosmologies and shown that they generate different, but viable cosmological solutions in the case of the improved Deser-Woodard model. However, there are other types of bouncing solutions [25]. For instance, we have the oscillatory bounce, which is generated by the scale factor

$$a(t) = A_0 \sin^2 \frac{kt}{t_0}, \quad (4.1)$$

and produces a complex solution for the function  $F(t)$ :

$$F(t) = C_1 e^{-t\sqrt{-\frac{k^2}{t_0^2}}} \csc^5 \left( \frac{kt}{t_0} \right) + C_2 e^{t\sqrt{-\frac{k^2}{t_0^2}}} \frac{\csc^5 \left( \frac{kt}{t_0} \right)}{2\sqrt{-\frac{k^2}{t_0^2}}}. \quad (4.2)$$

It is interesting to observe that this solution is complex only for real values of the parameters  $k$  and  $t_0$ . But this is a necessary condition to ensure that the scale factor (4.1) is a real valued function. Therefore, due to the presence of the complex function (4.2) in the solution of the distortion function  $f(t)$ , the class of oscillatory bounce can not describe a physical and viable cosmology in the DW II non-local gravity.

Another possible class of bouncing cosmology is the matter bounce. This scenario is characterized by the following scale factor

$$a(t) = A_0 \sqrt[3]{\frac{3\rho_c t^2}{2} + 1}, \quad (4.3)$$

where  $0 < \rho_c \ll 1$  is the critical density of the universe and  $A_0$  is a constant. In this case, the solution for  $F(t)$  is real and given in terms of Hypergeometric functions, but the exact solution of the field  $Y(t)$  is a complex function. Thus, matter bouncing universe renders a physically meaningless solution for the DW II gravity.

In the section 3.3 we have discussed the bouncing scenario generated by the scale factor (3.19) with  $\alpha < -1$ , which produces the Big Rip cosmology. Furthermore, we can also analyze the solutions for the scale factor (3.19) for the ranges of  $0 < \alpha < 1$  (which generates sudden singularities) and for  $-1 < \alpha < 0$  (which results in a Big Freeze universe). In both scenarios we simply can not find initial conditions that provide finite results for the fields  $U$  and  $V$ . Furthermore, even imposing that  $\dot{Y}$  becomes zero at the end of the interval, this field always present another zero points. These other zeros cause divergences in the fields  $U$  and  $V$  for every  $t$ , and consequently the non-local modification in the DW II Lagrangian density grows to infinity.

## 5 Conclusions

We have examined the viability of the improved Deser-Woodard non-local gravity model over the background of five different bouncing cosmologies. Among the bounce scenarios considered in our analysis, three of them have shown a physical and viable solution to the non-local distortion function  $f(Y)$ , namely: the symmetric bounce, the super bounce and the Big Rip bounce. The distortion function  $f$  encodes all the non-local information in the departure of the Einstein-Hilbert action and this kind of modified gravity theory is inspired by quantum phenomena.

Although the improved Deser-Woodard model was initially proposed to reproduce the late time accelerated expansion of the universe (present in the  $\Lambda$ CDM model), we have found that this model can also describe the aforementioned bouncing scenarios. The three solutions of  $f(Y)$  that we have obtained by the reconstruction procedure, are presented in the Figures 3, 4, and 5. The initial conditions used in the calculations are such that the time derivative of the field  $Y$  does not have zeros in the interval of solution, which is a necessary condition in order to avoid that  $f(Y)$  diverges for every value of  $t$ .

In regard of the physically admissible bouncing scenarios:

- In the analysis of the symmetric bounce scenario, we have found that the non-local distortion function is approximately constant throughout the cosmological evolution, from the past universe to the present one;
- Moreover, the super bounce generates one distortion function that diverges on the bouncing point and it is discontinuous, being piecewise-defined;
- Subsequently, the Big Rip bouncing has a solution for  $f(Y)$  as a straight line with positive slope in the past universe, being a different curve for the present universe.

On the other hand, the oscillatory bounce and the matter bounce presented a complex solution for the distortion function, remarkably these scenarios are physically meaningless in the context of DW II gravity. Although our reconstruction analysis is performed in a vacuum spacetime, we could expect that the inclusion of matter, for instance radiation, should not change our main results and conclusions (since the distortion function can be obtained by solving the corresponding differential equations).

We believe that the present analysis and physical results of the bouncing solutions (at the level of background cosmology in the DW II gravity) motivate a deeper look at the early Universe cosmology. To further restrict physically relevant models, we need to study the early time perturbations of these bouncing models and examine their implications, for instance, on the cosmic microwave background. This and other topics are now under development and will be reported elsewhere.

## Acknowledgments

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. R.B. acknowledges partial support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq Projects No. 305427/2019-9 and No. 421886/2018-8) and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG Project No. APQ-01142-17).

## References

- [1] C. M. Will, *The confrontation between general relativity and experiment*, *Living Rev. Relativ.* **vol 17** (2014).
- [2] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time*, **vol 1**, Cambridge University Press (1973).
- [3] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*, Macmillan (1973).
- [4] R. Geroch, *What is a singularity in general relativity?* *Ann. of Phys.*, **vol 17**, 3, Elsevier, (1968), pp. 526–540.
- [5] M. Bojowald, *Absence of singularity in loop quantum cosmology*, *Phys. Rev. Lett.*, **vol 86**, pg. 5227-5230, APS, (2001).
- [6] M. Novello and S. E. Perez Bergliaffa, *Bouncing Cosmologies*, *Phys. Reports*, **vol 463**, pg. 127–213, Elsevier, (2008).
- [7] A. Ashtekar and P. Singh, *Loop Quantum Cosmology: A Status Report*, *Class. Quant. Grav.*, **vol 28**, pg. 213001, IOP, (2011).
- [8] D. Battefeld, and P. Peter, *A critical review of classical bouncing cosmologies*, *Phys. Reports*, **vol 571** pg. 1–66, Elsevier, (2015).
- [9] R. Brandenberger and P. Peter, *Bouncing Cosmologies: Progress and Problems*, *Found. Phys.*, **vol 47**, pg. 797-850, Springer, (2017).
- [10] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, *Modified Gravity Theories on a Nutshell: Inflation, Bounce and Late-time Evolution*, *Phys. Rept.*, **vol 692**, pg. 1-104, Elsevier, (2017).
- [11] V. Mukhanov, *Physical Foundations of Cosmology*, Cambridge University Press, Oxford, 2005.
- [12] K. Bamba and S. D. Odintsov, *Inflationary cosmology in modified gravity theories*, *Symmetry*, **vol 7**, pg. 220-240, MDPI, (2015).
- [13] M. Ishak, *Testing general relativity in cosmology*, *Living Rev. Relativ.*, **vol 22**, 1, Springer, (2019).
- [14] E. Belgacem, Y. Dirian, S. Foffa, and M. Maggiore, *Nonlocal gravity. Conceptual aspects and cosmological predictions*, *J. of Cosmo. and Astr. Phys.*, **vol 03**, IOP Publishing, (2018).
- [15] D. A. R. Dalvit and F. D. Mazzitelli, *Running coupling constants, Newtonian potential, and non-localities in the effective action*, *Phys. Rev. D*, **vol 50**, 2, pg. 1001, APS, (1994).
- [16] C. Wetterich, *Effective non-local Euclidean gravity*, *Gen. Relativ. and Gravit.*, **vol 30**, 1, pg. 159–172, Springer, (1998).
- [17] A. Barvinsky and G. Vilkovisky, *The Generalized Schwinger-DeWitt Technique in Gauge Theories and Quantum Gravity*, *Phys. Rept.*, **vol 119**, pg. 1-74, Elsevier, (1985).
- [18] I. L. Buchbinder, S. D. Odintsov, and I. L. Shapiro, *Effective action in quantum gravity*, Bristol, UK: IOP (1992).
- [19] V. Mukhanov and S. Winitzki, *Introduction to quantum effects in gravity*, University Press, Cambridge, (2007).
- [20] I. L. Shapiro, *Effective Action of Vacuum: Semiclassical Approach*, *Class. Quant. Grav.*, **vol 25**, pg. 103001, IOP, (2008).
- [21] M. Maggiore and M. Mancarella, *Nonlocal gravity and dark energy*, *Phys. Rev. D*, **vol 90**, 2, APS, (2014).
- [22] A. Barvinsky, *Serendipitous discoveries in nonlocal gravity theory*, *Phys. Rev. D*, **vol 85**, 10, APS, (2012).

- [23] S. Deser and R. P. Woodard, *Nonlocal cosmology*, *Phys. Rev. Lett.*, **vol 99**, 11, APS, (2007).
- [24] S. Deser and R. P. Woodard, *Nonlocal cosmology II. Cosmic acceleration without fine tuning or dark energy*, *J. of Cosmo. and Astr. Phys.*, **vol 06**, pg. 34, IOP Publishing, (2019).
- [25] M. Caruana, G. Farrugia, and J. L. Said. *Cosmological bouncing solutions in  $f(T, B)$  gravity* *Eur. Phys. J. C* 80.7 pg. 1-20 Springer, (2020).
- [26] C. Y. Chen, P. Chen and S. Park, *Primordial bouncing cosmology in the Deser-Woodard nonlocal gravity*, *Phys. Lett. B*, **vol 796**, pg. 112–116, Elsevier, (2019).
- [27] J. C. Ding and J. B. Deng, *Structure formation in the new Deser-Woodard nonlocal gravity model*, *J. of Cosmo. and Astr. Phys.*, **vol 12**, pg. 054, IOP Publishing, (2019).