

# Stellar models with like–Tolman IV complexity factor

J. Andrade<sup>1</sup>, E. Contreras<sup>a,1</sup>

<sup>1</sup>Departamento de Física, Colegio de Ciencias e Ingeniería, Universidad San Francisco de Quito, Quito, Ecuador.

Received: date / Accepted: date

**Abstract** In this work, we construct stellar models based on the complexity factor as a supplementary condition which allows to close the system of differential equations arising from the Gravitational Decoupling. The assumed complexity is a generalization of the one obtained from the well known Tolman IV solution. We use Tolman IV, Wyman IIa, Durgapal IV and Heintzmann IIa as seeds solutions. Reported compactness parameters of SMC X-1 and Cen X-3 are used to study the physical acceptability of the models. Some aspects related to the density ratio are also discussed.

## 1 Introduction

For a long time, stellar models were considered to be supported by Pascalian fluids (equal principal stresses); approximation which resulted to be appropriate to describe a variety of circumstances. However, now is well-known that for certain ranges of the density there are some physical phenomena which might take place leading to local anisotropy in the configuration. (see Refs. [1–16], for discussions on this point). Among all these possibilities, we could mention: i) intense magnetic field observed in compact objects such as white dwarfs, neutron stars, or magnetized strange quark stars (see, for example, Refs. [17–25]) and ii) viscosity (see [26–33] and references therein). Besides, it has been recently proven that the presence of dissipation, energy density inhomogeneities and shear yield the isotropic pressure condition becomes unstable [34]. Based on these points, the renewed interest in the study of fluids not satisfying the isotropic condition is clear and justifies our present work on the construction of anisotropic models [35–39].

The strategies to construct anisotropic solutions are many but recently, the well known Gravitational Decoupling (GD) [40] by the Minimal Geometric Deformation approach (MGD) (for implementation in  $3 + 1$  and  $2 + 1$  dimensional spacetimes see [41–79]. For recent developments see [80–87]) has been broadly implemented to extend isotropic solutions to anisotropic domains. In static and spherically symmetric spacetimes, there are only three independent Einstein field equations but five unknown, namely two metric functions, the density energy and the radial and tangential pressures. However, the GD demands the assumption of a seed solution which allows to decrease the number of degrees of freedom and, as a consequence, only one extra condition is required to close the system. In this sense, a key point in the implementation of MGD is to provide such an auxiliary condition which could be the mimic constraint for the pressure and the density, regularity condition of the anisotropy function, barotropic equation of state, among others. In this work we use the recently introduced definition of complexity for self-gravitating fluids [88] and, in particular, we propose a like-Tolman IV complexity factor.

This work is organized as follows. The next section is devoted to reviewing the main aspects of GD by MGD. In section 3 we introduce the concept of complexity and obtain an expression for the complexity factor in GD. Then, in section 4 we calculate and generalize the complexity factor from the Tolman IV solution and implement this result with the aim to construct extension of Tolman IV, Wyman IIa, Durgapal IV and Heintzmann IIa. Section 5 is devoted to interpreting and discussing our results and some comments and final remarks are given in the last section.

<sup>a</sup>e-mail: econtreras@usfq.edu.ec

## 2 Gravitational decoupling

In this section we introduce the GD by MGD (for more details, see [40]). Let us start with the Einstein field equations (EFE)

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 8\pi\tilde{T}_{\mu\nu}, \quad (1)$$

with

$$\tilde{T}_{\mu\nu} = T_{\mu\nu}^{(s)} + \alpha\theta_{\mu\nu}, \quad (2)$$

where  $T_{\mu\nu}^{(s)}$  represents the matter content of a known solution<sup>1</sup>, namely the *seed* sector, and  $\theta_{\mu\nu}$  describes an extra source coupled through the parameter  $\alpha$ . Note that, since the Einstein tensor fulfills the Bianchi identities, the total energy-momentum tensor satisfies

$$\nabla_{\mu}\tilde{T}^{\mu\nu} = 0. \quad (3)$$

It is important to point out that, whenever  $\nabla_{\mu}T^{\mu\nu(s)} = 0$ , the condition

$$\nabla_{\mu}\theta^{\mu\nu} = 0, \quad (4)$$

is automatic and as a consequence, there is no exchange of energy-momentum between the seed solution and the extra source  $\theta^{\mu\nu}$  so that the interaction is entirely gravitational.

In a static and spherically symmetric spacetime sourced by

$$T_{\nu}^{\mu(s)} = \text{diag}(\rho^{(s)}, -p_r^{(s)}, -p_t^{(s)}, -p_t^{(s)}) \quad (5)$$

$$\theta_{\nu}^{\mu} = \text{diag}(\theta_0^0, \theta_1^1, \theta_2^2, \theta_2^2), \quad (6)$$

and a metric given by

$$ds^2 = e^{\nu}dt^2 - e^{\lambda}dr^2 - r^2(\theta^2 + \sin^2\theta d\phi^2), \quad (7)$$

Eqs. (1) and (2) lead to

$$\tilde{\rho} = \frac{1}{8\pi} \left[ \frac{1}{r^2} + e^{-\lambda} \left( \frac{\lambda'}{r} - \frac{1}{r^2} \right) \right], \quad (8)$$

$$\tilde{P}_r = \frac{1}{8\pi} \left[ -\frac{1}{r^2} + e^{-\lambda} \left( \frac{\nu'}{r} + \frac{1}{r^2} \right) \right], \quad (9)$$

$$\tilde{P}_t = \frac{1}{32\pi} e^{-\lambda} \left( 2\nu'' + \nu'^2 - \lambda'\nu' + 2\frac{\nu' - \lambda'}{r} \right), \quad (10)$$

where we have defined<sup>2</sup>

$$\tilde{\rho} = \rho^{(s)} + \alpha\theta_0^0, \quad (11)$$

$$\tilde{P}_r = p_r^{(s)} - \alpha\theta_1^1, \quad (12)$$

$$\tilde{P}_t = p_t^{(s)} - \alpha\theta_2^2. \quad (13)$$

<sup>1</sup>In this work we shall use  $c = G = 1$ .

<sup>2</sup>Note that the matter sector has dimensions of a length squared in the units we are using

It is clear that given the non-linearity of Einstein's equations, the decomposition (2) does not lead to two sets of decoupled equations; one for each source involved. Nevertheless, contrary to the broadly held belief, such a decoupling is possible, to some extent, in the context of MGD as we shall demonstrate in what follows.

Let us introduce a geometric deformation in the metric functions given by

$$\nu \longrightarrow \xi + \alpha g, \quad (14)$$

$$e^{-\lambda} \longrightarrow e^{-\mu} + \alpha f, \quad (15)$$

where  $\{f, g\}$  are the so-called decoupling functions and  $\alpha$  is the same free parameter that ‘‘controls’’ the influence of  $\theta_{\mu\nu}$  on  $T_{\mu\nu}^{(s)}$  in Eq. (2). In this work we shall concentrate in the particular case  $g = 0$  and  $f \neq 0$ . Now, replacing (14) and (15) in the system (8-10), we are able to split the complete set of differential equations into two subsets: one describing a seed sector sourced by the conserved energy-momentum tensor,  $T_{\mu\nu}^{(s)}$

$$\rho^{(s)} = \frac{1}{8\pi} \left[ \frac{1}{r^2} + e^{-\mu} \left( \frac{\mu'}{r} - \frac{1}{r^2} \right) \right], \quad (16)$$

$$p_r^{(s)} = \frac{1}{8\pi} \left[ -\frac{1}{r^2} + e^{-\mu} \left( \frac{\nu'}{r} + \frac{1}{r^2} \right) \right], \quad (17)$$

$$p_t^{(s)} = \frac{1}{32\pi} e^{-\mu} \left( 2\nu'' + \nu'^2 - \mu'\nu' + 2\frac{\nu' - \mu'}{r} \right), \quad (18)$$

and the other set corresponding to quasi-Einstein field equations sourced by  $\theta_{\mu\nu}$

$$\theta_0^0 = \frac{1}{8\pi} \left[ -\frac{f}{r^2} - \frac{f'}{r} \right], \quad (19)$$

$$\theta_1^1 = \frac{1}{8\pi} \left[ -f \left( \frac{\nu'}{r} + \frac{1}{r^2} \right) \right], \quad (20)$$

$$\theta_2^2 = \frac{1}{8\pi} \left[ -\frac{f}{4} \left( 2\nu'' + \nu'^2 + 2\frac{\nu'}{r} \right) - \frac{f'}{4} \left( \nu' + \frac{2}{r} \right) \right]. \quad (21)$$

As we have seen, the components of  $\theta_{\mu\nu}$  satisfy the conservation equation  $\nabla_{\mu}\theta_{\nu}^{\mu} = 0$ , namely

$$\theta_1^1 - \frac{\nu'}{2}(\theta_0^0 - \theta_1^1) - \frac{2}{r}(\theta_2^2 - \theta_1^1) = 0. \quad (22)$$

In this work, we consider that the interior configuration is surrounded by the Schwarzschild vacuum so that, on the boundary surface  $\Sigma$ , we require

$$e^{\nu} \Big|_{\Sigma^-} = \left( 1 - \frac{2M}{r} \right) \Big|_{\Sigma^+}, \quad (23)$$

$$e^{\lambda} \Big|_{\Sigma^-} = \left( 1 - \frac{2M}{r} \right) \Big|_{\Sigma^-}, \quad (24)$$

$$\tilde{P}_r(r) \Big|_{\Sigma^-} = \tilde{P}_r(r) \Big|_{\Sigma^+} = 0, \quad (25)$$

which corresponds to the continuity of the first and second fundamental form across that surface of the star.

To conclude this section, we emphasize the importance of GD as a useful tool to find solutions of EFE. As it is well known, in static and spherically symmetric spacetimes sourced by anisotropic fluids, EFE reduce to three equations given by (8), (9) and (10) and five unknowns, namely  $\{\nu, \lambda, \tilde{\rho}, \tilde{P}_r, \tilde{P}_t\}$ . In this regard, two auxiliary conditions must be specified, namely metric conditions, equations of state, etc. Nevertheless, as in the context of GD a seed solution must be given, the number of degrees of freedom reduces from five to four and, as a consequence, only one extra condition is required. In general, this condition have been implemented in the decoupling sector given by Eqs. (19), (20) and (21) as some equation of state which leads to a differential equation for the decoupling function  $f$ . In this work, we take an alternative route to obtain the decoupling function; namely, the complexity factor that we shall introduce in the next section as a supplementary condition of the total solution.

### 3 Complexity of compact sources

Recently, a new definition of complexity for self-gravitating fluid distributions has been introduced in Ref. [88] which is based on the idea that the least complex gravitational system is the one supported by a homogeneous energy density distribution and isotropic pressure. In this direction, there is a scalar associated with the orthogonal splitting of the Riemann tensor [89] in static and spherically symmetric space-times which encodes the intuitive idea of complexity, namely

$$Y_{TF} = 8\pi\Pi - \frac{4\pi}{r^3} \int_0^r \tilde{r}^3 \rho' d\tilde{r}, \quad (26)$$

with  $\Pi \equiv P_r - P_t$ . Also, it can be demonstrated that in terms of Eq. (26) the Tolman mass reads

$$m_T = (m_T)_\Sigma \left( \frac{r}{r_\Sigma} \right)^3 + r^3 \int_r^{r_\Sigma} \frac{e^{(\nu+\lambda)/2}}{\tilde{r}} Y_{TF} d\tilde{r}, \quad (27)$$

so that  $Y_{TF}$  enclose the modifications on the active gravitational mass produced by the energy density inhomogeneity and the anisotropy of the pressure.

It is worth noticing that the vanishing complexity condition ( $Y_{TF} = 0$ ) can be satisfied not only in the simplest case of isotropic and homogeneous system but in all the cases where

$$\Pi = \frac{1}{2r^3} \int_0^r \tilde{r}^3 \rho' d\tilde{r}, \quad (28)$$

which provides a non-local equation of state that can be used as a complementary condition to close the system of EFE (for a recent implementation, see [60], for example). However, given that this condition could fail in some cases in the construction of specific stellar models, non-vanishing values of  $Y_{TF}$  must be supplied. An example of how this can be achieved can be found in [60].

In this work we shall use the complexity factor as a supplementary condition for the total sector so we replace (14), (15) in (26) and use (8), (9) and (10) to obtain

$$\begin{aligned} \frac{\alpha\xi'}{4} f' + \frac{\alpha}{2} \left( \xi'' - \frac{\xi'}{r} + \frac{\xi'^2}{2} \right) f \\ + \frac{e^{-\mu}}{2} \left( \xi'' - \frac{\xi'}{r} + \frac{\xi'^2}{2} - \frac{\mu'\xi'}{2} \right) + Y_{TF} = 0 \end{aligned} \quad (29)$$

Note that as far as the pair  $\{\xi, \mu\}$  is specified (the seed solution), Eq. (29) becomes a differential equation for the decoupling function  $f$  when a value of  $Y_{TF}$  is specified.

### 4 Stellar models with like Tolman IV complexity

In this work, we construct interior solutions based on Tolman IV, Wyman IIa, Durgapal IV and Heintzmann IIa as isotropic seeds in the framework of GD by using the complexity factor as supplementary condition. At first sight, the vanishing complexity seems to be straightforward but it can be demonstrated that such a condition fails for the seeds under consideration in this work. As an alternative, we generalize the complexity factor of the well-known Tolman IV solution.

As it is well-known, Tolman IV reads [90]

$$e^\nu = B_0^2 \left( 1 + \frac{r^2}{A_0^2} \right) \quad (30)$$

$$e^{-\lambda} = \frac{(C_0^2 - r^2)(A_0^2 + r^2)}{C_0^2(A_0^2 + 2r^2)} \quad (31)$$

$$\rho = \frac{3A_0^4 + A_0^2(3C_0^2 + 7r^2) + 2r^2(C_0^2 + 3r^2)}{8\pi C_0^2(A_0^2 + 2r^2)^2} \quad (32)$$

$$p = \frac{C_0^2 - A_0^2 - 3r^2}{8\pi C_0^2(A_0^2 + 2r^2)}, \quad (33)$$

where  $A_0$  and  $C_0$  are constants with dimension of a length and  $B_0$  is a dimensionless constant. Now, replacing (30), (31), (32) and (33) in (26) we arrive at

$$Y_{TF} = \frac{r^2(A_0^2 + 2C_0^2)}{C_0^2(A_0^2 + 2r^2)^2}, \quad (34)$$

which has dimensions of the inverse of a length squared. Note that Eq. (34) can be easily generalized to

$$Y_{TF} = \frac{a_1 r^2}{(a_2 + a_3 r^2)^2} \quad (35)$$

where  $a_1$  and  $a_3$  are arbitrary dimensionless constants and  $a_2$  must be a constant with dimension of a length squared. It should be emphasized that reason for introducing the set  $\{a_1, a_2, a_3\}$  is nothing but to generalize the complexity factor (34). In what follows we shall consider Eq. (35) as the condition to close the system and generate anisotropic models from isotropic seeds.

#### 4.1 Model 1: like–Tolman IV solution

Replacing (30) and (31) in (29) and using (35) we arrive at

$$f = (A_0^2 + r^2) \left[ c_1 + \frac{1}{\alpha} \left( \frac{a_1}{a_3 \zeta(r)} + \frac{A_0^2 + 2C_0^2}{2C_0^2(A_0^2 + 2r^2)} \right) \right], \quad (36)$$

where  $c_1$  is an integration constant with dimensions of the inverse of a length squared and  $\zeta(r)$  is an auxiliary function with dimensions of a length squared (see Appendix, section 8).

It can be shown that to ensure regularity in the matter sector the constant  $c_1$  must satisfy

$$c_1 = \frac{-2a_1 A_0^2 C_0^2 + a_2 a_3 A_0^2 + 2a_2 a_3 C_0^2}{2\alpha a_2 a_3 A_0^2 C_0^2}. \quad (37)$$

Replacing (36) in (15) and using (37) we find

$$e^{-\lambda} = \frac{(A_0^2 + r^2) (2a_1 C_0^2 + a_3 \zeta(r) (2\alpha c_1 C_0^2 - 1))}{2a_3 C_0^2 \zeta(r)}. \quad (38)$$

Now, from (8), (9), (10) we arrive at

$$\tilde{\rho} = \frac{1}{8\pi a_2 A_0^2 \zeta(r)^2} \left[ a_1 A_0^4 (3a_2 + a_3 r^2) + a_1 A_0^2 r^2 (5a_2 + 3a_3 r^2) - 3a_2 \zeta(r)^2 \right] \quad (39)$$

$$\tilde{P}_r = \frac{3a_2 \zeta(r) - a_1 A_0^2 (A_0^2 + 3r^2)}{8\pi a_2 A_0^2 \zeta(r)} \quad (40)$$

$$\tilde{P}_t = \frac{3a_2 \zeta(r)^2 - a_1 a_2 A_0^4 - a_1 A_0^2 r^2 (5a_2 + 3a_3 r^2)}{8\pi a_2 A_0^2 \zeta(r)^2}, \quad (41)$$

Finally, the continuity of the first and the second fundamental form leads to

$$a_1 = \frac{3a_2 \zeta(R)}{A_0^2 (A_0^2 + 3R^2)}. \quad (42)$$

$$\frac{A_0^2}{R^2} = \frac{R - 3M}{M} \quad (43)$$

$$B_0^2 = 1 - \frac{3M}{R}. \quad (44)$$

Note that the free parameters are  $\{R, M, a_2, a_3\}$  (see Appendix 8 where  $a_3$  appears explicitly). It is worth mentioning that from (43) and (44) is clear that compactness satisfies  $M/R < 1/3$ , which corresponds to a more stringent condition when compared to the Buchdahl's limit ( $M/R < 4/9$ ) for isotropic solutions. More precisely, the solutions allowed with this model should be less compact given that the interval  $1/3 \leq M/R < 4/9$  is forbidden.

As we shall see later, the strategy to explore the feasibility of our solution will be to specify the compactness parameter associated with SMC X-1 and Cen X-3 in order to set suitable values for  $a_2$  and  $a_3$  (see section 5 for details)

#### 4.2 Model 2: like–Wyman IIa solution

In this case we consider the Wyman IIa metric [95, 96] with  $n = 2$  as a seed solution which reads

$$e^{\xi(r)} = (A - Br^2)^2 \quad (45)$$

$$e^{-\mu(r)} = 1 + Cr^2 (A - 3Br^2)^{-2/3}, \quad (46)$$

where  $A$  is a dimensionless constant and  $B$  and  $C$  are constants with dimensions of the inverse of a length squared. It is worth mentioning that all the results below will be written in terms of auxiliary functions  $\{\zeta, \chi, \varrho_1, \mathcal{P}_1, \mathcal{P}_2\}$  which are defined in Appendix, section 8.

Following the same procedure that in the previous section we obtain

$$f = \frac{r^2}{2\alpha a_3} \left[ \frac{a_1 (a_3 A + a_2 B)}{a_2 B \zeta(r)} - \frac{2a_3 C}{(A - 3Br^2)^{2/3}} \right] - \frac{a_1}{2\alpha a_3^2} \chi(r), \quad (47)$$

from where

$$e^{-\lambda} = \frac{1}{2a_3^2} \left[ \frac{a_3 r^2 (a_1 a_3 A + a_2 B (a_1 + 2a_3^2))}{a_2 B \zeta(r)} + \frac{2a_2 a_3^2}{\zeta(r)} - a_1 \chi(r) \right], \quad (48)$$

From 8, 9 and 10 the matter sector reads

$$\tilde{\rho} = \frac{a_1}{16\pi a_3^2} \left[ \frac{\chi(r)}{r^2} - \frac{a_3 \varrho_1(r)}{a_2 B \zeta(r)^2} \right] \quad (49)$$

$$\tilde{P}_r = \frac{a_3 r^2 \mathcal{P}_1(r) - a_1 a_2 B (5Br^2 - A) \zeta(r) \chi(r)}{16\pi a_2 a_3^2 Br^2 (Br^2 - A) \zeta(r)} \quad (50)$$

$$\tilde{P}_t = \frac{a_3 \mathcal{P}_2(r) - 4a_1 a_2 B^2 \zeta(r)^2 \chi(r)}{16\pi a_2 a_3^2 B (Br^2 - A) \zeta(r)^2}. \quad (51)$$

Continuity of the first and the second fundamental form leads to

$$a_1 = \frac{8a_2a_3^2B^2R^2(A - 5BR^2)^{-1}\zeta(R)}{[a_3R^2(a_3A + a_2B) - a_2B\zeta(R)\chi(R)]} \quad (52)$$

$$A^2 = \frac{B^2(5M - 2R)^2R^4}{M^2} \quad (53)$$

$$B^2 = \frac{M^2}{4R^5(R - 2M)}. \quad (54)$$

As in the previous case, the free parameters are  $\{R, M, a_2, a_3\}$ . Note that, as  $\{\zeta, \varrho_1, \mathcal{P}_2\}$  have dimensions of a length squared and  $\{\chi, \mathcal{P}_1\}$  are dimensionless (see Appendix 8), all the expressions above are dimensionally correct. It is worth noticing that from (54),  $R > 2M$  which is in accordance with the restriction that any stable configuration should be greater than its Schwarzschild radius. Furthermore, (53) leads to  $M/R \neq 2/5$  or the metric becomes degenerated, namely  $g_{tt} = 0, \forall r$ .

#### 4.3 Model 3: like–Durgapal IV solution

In this case we consider the Durgapal IV metric [95,97] as a seed solution, which reads

$$e^\xi = A(Cr^2 + 1)^4 \quad (55)$$

$$e^{-\mu} = \frac{7 - 10Cr^2 - C^2r^4}{7(Cr^2 + 1)^2} + \frac{Cr^2}{(Cr^2 + 1)^2(1 + 5Cr^2)^{2/5}} \quad (56)$$

where  $A$  is a dimensionless constants and  $C$  is a constant with dimensions of the inverse of a length squared. In what follows, we shall use the auxiliary functions  $\{\zeta, \eta_1, \beta_1, \beta_4, \beta_5, \beta_8, \beta_{11}, \varrho_2, \mathcal{P}_3, \mathcal{P}_4\}$  defined in the Appendix, section 8.

In this case,

$$f = \frac{r^2}{56\alpha a_3^3 C \beta_1(r)^2} \left[ 7a_1 \left( -a_3 C^3 r^2 + \frac{2(a_2 C - a_3)^3}{a_2 \zeta(r)} + 2C^2(2a_2 C - 3a_3) \right) + 8a_3^3 C^2 \left( Cr^2 + 10 - \frac{7B}{(5Cr^2 + 1)^{2/5}} \right) - \frac{3a_1(a_3 - a_2 C)^2 \chi(r)}{4\alpha a_3^4 \beta_1(r)^2} \right], \quad (57)$$

and the radial metric reads

$$e^{-\lambda} = \frac{a_3 \eta_1(r) - 6a_1 a_2 C (a_3 - a_2 C)^2 \zeta(r) \chi(r)}{8a_2 a_3^4 C \beta_1(r)^2 \zeta(r)}. \quad (58)$$

From (8), (9), (10) we obtain

$$\tilde{\rho} = \frac{1}{64\pi a_2 a_3^4 C r^2 \beta_1(r)^3 \zeta(r)^2} \left[ a_3 r^2 \varrho_2(r) - 6a_1 a_2 C \beta_4(r) (a_3 - a_2 C)^2 \zeta(r)^2 \chi(r) \right] \quad (59)$$

$$\tilde{P}_r = \frac{\mathcal{P}_3(r)}{64\pi a_2 a_3^4 C r^2 \beta_1(r)^3 \zeta(r)} \quad (60)$$

$$\tilde{P}_t = \frac{\mathcal{P}_4(r)}{32\pi a_2 a_3^4 C \beta_1(r)^3 \zeta(r)^2}, \quad (61)$$

and the matching conditions lead to

$$a_1 = \frac{8a_2 a_3^4 C^2 R^2 (6 - CR^2 \beta_5(R)) \zeta(R) \beta_8(R)^{-1}}{[a_3 R^2 \beta_{11}(R) + 6a_2 C (a_3 - a_2 C)^2 \zeta(R) \chi(R)]}, \quad (62)$$

$$A^2 = \frac{1 - \frac{2M}{R}}{\left( \frac{M}{4R - 9M} + 1 \right)^4} \quad (63)$$

$$C^2 = \frac{M}{R^2(4R - 9M)}. \quad (64)$$

Note that, as  $\{\zeta, \eta_1, \varrho_2, \mathcal{P}_3\}$  have dimensions of a length squared and  $\{\beta_1, \beta_4, \beta_5, \beta_8, \beta_{11}\}$  are dimensionless (see Appendix 8), all the expressions above have the correct dimensions. In this case, it is clear that from Eqs. (63) and (64), the solution must satisfy  $M/R < 4/9$ , which corresponds to the Buchdahl's limit for isotropic solutions.

#### 4.4 Model 4: like–Heintzmann IIa

In this section we consider the Heintzmann IIa solution [95,98] which metric components are

$$e^\xi = A^2(1 + Br^2)^3 \quad (65)$$

$$e^{-\mu} = 1 - \frac{3Br^2}{2} \frac{1 + (1 + 4Br^2)^{-1/2}}{1 + Br^2} \quad (66)$$

where  $A$  is a dimensionless constant and  $B$  is a constant with dimensions of the inverse of a length squared. In this section we shall use the list of auxiliary functions  $\{\zeta, \eta_2, \gamma_1, \gamma_3, \gamma_6, \gamma_7, \gamma_9, \varrho_3, \mathcal{P}_5, \mathcal{P}_6\}$  defined in the Appendix, section 8.

In this case we have

$$f = \frac{r^2}{6\alpha B \gamma_1(r)} \left[ \frac{9B^2 C}{\sqrt{4Br^2 + 1}} + 3B^2 - \frac{2a_1(2a_2^2 B^2 + a_2 a_3 B \gamma_2(r) + a_3^2)}{a_2 a_3^2 \zeta(r)} - \frac{2a_1(a_3 - a_2 B) \chi(r)}{3a_3^3 \alpha \gamma_1(r)} \right] \quad (67)$$

from where, the radial metric results

$$e^{-\lambda} = \frac{a_3 \eta_2(r) + 2a_1 a_2 B (a_2 B - a_3) \zeta(r) \chi(r)}{3a_2 a_3^3 B \gamma_1(r) \zeta(r)}. \quad (68)$$

From (8), (9) and (10) we arrive at

$$\tilde{\rho} = \frac{2a_1a_2B\gamma_3(r)(a_2B - a_3)\zeta(r)^2\chi(r) + a_3r^2\varrho_3(r)}{24\pi a_2a_3^3Br^2\gamma_1(r)^2\zeta(r)^2} \quad (69)$$

$$\tilde{P}_r = \frac{\mathcal{P}_5(r)}{24\pi a_2a_3^3Br^2\gamma_1(r)^2\zeta(r)} \quad (70)$$

$$\tilde{P}_t = \frac{\mathcal{P}_6(r)}{24\pi a_2a_3^3B\gamma_1(r)^2\zeta(r)^2} \quad (71)$$

Finally, matching conditions lead to

$$a_1 = -\frac{3a_2a_3^3B^2R^2\gamma_7(R)\zeta(R)\gamma_6(R)^{-1}}{a_3R^2\gamma_9(R) - 2a_2B(a_2B - a_3)\zeta(R)\chi(R)} \quad (72)$$

$$A^2 = -\frac{(7M - 3R)^3}{27R(R - 2M)^2} \quad (73)$$

$$B^2 = \frac{M^2}{R^4(3R - 7M)^2}. \quad (74)$$

Note that, as in all the previous cases the free parameters are  $\{R, M, a_2, a_3\}$ . Note that, as  $\{\zeta, \eta_2, \varrho_3, \mathcal{P}_5, \mathcal{P}_6\}$  have dimensions of a length squared and  $\{\gamma_1, \gamma_3, \gamma_6, \gamma_7, \gamma_9\}$  are dimensionless (see Appendix 8), all the expressions above are dimensionally correct. It should be emphasized that,  $M/R < 3/7 < 4/9$  which corresponds to less compact solutions than the allowed by the Buchdahl's limit.

## 5 Discussion


In this section we analyze the results obtained previously in order to verify the physical acceptability of the models [102]. To this end, we shall use the compactness parameters given in Table 1 and set suitable values of  $a_2$  and  $a_3$  in order to discuss to what extent our solutions are suitable to describe the SMC X-1 and Cen X-3 systems.

### 5.1 Metrics


In figures 1 and 2 we show the metric functions for the compactness parameter indicated in the legend. Note that on one hand  $e^\nu$  is a monotonously increasing function with  $e^{\nu(0)} = \text{constant}$ . On the other hand  $e^{-\lambda}$  is monotonously decreasing with  $e^{-\lambda(0)} = 1$ , as expected.

### 5.2 Matter sector

In figures 3, 4 and 5 we show the profile of  $\tilde{\rho}$ ,  $\tilde{P}_r$  and  $\tilde{P}_t$  as a function of the radial coordinates for the values of the parameters in the legend. Note that all the quantities fulfill the physical requirements for all the parameters involved, namely  $\tilde{\rho}$ ,  $\tilde{P}_r$  and  $\tilde{P}_t$  are finite at



**Fig. 1**  $e^\nu$  for Model 1 (a), Model 2 (b) Model 3 (c) and Model 4 (d)



**Fig. 2**  $e^{-\lambda}$  for Model 1 (a), Model 2 (b) Model 3 (c) and Model 4 (d)

the center and decrease monotonously toward the surface. Besides,  $\tilde{P}_t(0) = \tilde{P}_r(0)$  and  $\tilde{P}_t(r) > \tilde{P}_r(r)$  for all  $r > 0$  as expected (see Fig. 6)

### 5.3 Energy conditions and causality

A suitable stellar model must satisfies the dominant energy condition (DEC) in order to avoid violation of causality. The DEC requires


$$\tilde{\rho} - \tilde{P}_r \geq 0 \quad (75)$$

$$\tilde{\rho} - \tilde{P}_t \geq 0. \quad (76)$$

In Figs. 7 and 8 it can be seen that all the solutions satisfy DEC for all the parameters involved.

Compact start	$M/M_\odot$	$R(\text{km})$	$u = M/R$	$\rho(0)/\rho(R)$	$Z(R)$
SMC X-1 [91]	1.04	8.301	0.19803	1.4659 [93]	0.286776
Cen X-3 [92]	1.49	10.8	0.2035	1.915 [94]	0.298592

**Table 1** Physical parameters for the compact starts SMC X-1 and Cen X-3.




**Fig. 3**  $\rho$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .

In figures 9 and 10 we show that the radial and tangential sound velocities are less than unity, as required (we are assuming  $c = 1$ ).

#### 5.4 Redshift and density ratio


In the previous section we have demonstrated that, based on the compactness parameter of both SMC X-1 and Cen X-3 in Table 1, the four models satisfy the ba-



**Fig. 4**  $\tilde{P}_r(r)$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .

sic physical requirements to be considered as suitable interior configurations. Now, with the aim to explore which model is more adequate to describe the compact objects under consideration, in this section we go a step further and study the redshift  $Z = e^{-\nu/2} - 1$  and the density ratio  $\tilde{\rho}(0)/\tilde{\rho}(R)$  to each model and compare our results with the values in Table 1.

In figure 11 we show the redshift  $Z$  as a function of the radial coordinate. Note that  $Z$  decreases outward and its value at the surface is less than the uni-



**Fig. 5**  $\tilde{P}_t(r)$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .


versal bound for solutions satisfying the DEC, namely  $Z_{bound} = 5.211$ .

The values of the density ratio for SMC X-1 reported in [93] is  $\tilde{\rho}(0)/\tilde{\rho}(R) \approx 1.4659$ . Now, from Table 2, we appreciate that Models 3 and 4 fit accurately to SMC X-1. Similarly,  $\tilde{\rho}(0)/\tilde{\rho}(R) \approx 1.915$  for Cen X-3 as appears in [94] so that Models 2, 3 and 4 are good candidates to describe this compact objects.

In summary, Models 3 and 4 might be considered as suitable solutions describing SMC X-1 while models 2, 3 and 4 are the solutions for Cen X-3.

## 6 Conclusions

In this work, we extended four isotropic models to aniso-



**Fig. 6**  $\tilde{P}_t(r) - \tilde{P}_r(r)$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .

tropic domains by Gravitational Decoupling. As a supplementary condition, we used a complexity factor which corresponds to a generalization of the obtained from the well-known Tolman IV solution. We verify the basic physical acceptability conditions; namely: i) the metric functions are regular at the origin. Furthermore  $g_{tt}(0) = constant$  and  $g^{rr}(0) = 1$ , ii) both the density energy and pressures are regular at the origin and decrease monotonously outward, iii) the solutions satisfy the dominant energy condition. All of these conditions were tested after imposing the compactness parameters of both SMC X-1 and Cen X-3 systems. It is worth mentioning that, although all the solutions are well behaved, only some of them can be considered as suitable models for the compact objects under consideration based on the density ratio. More precisely, Models 3 and 4 are more appropriated to describe SMC X-1




Model	$\rho(0)/\rho(R)$
Model 1 ( $a_2 = 0.53, a_3 = 0.57$ )	2.62443
Model 2 ( $a_2 = 2.7, a_3 = 1.6$ )	1.84825
Model 3 ( $a_2 = 0.8, a_3 = 0.2$ )	1.43124
Model 4 ( $a_2 = 1.1, a_3 = 0.39$ )	1.53133

**Table 2** Estimated values of the density ratio for SMC X-1 ( $u = 0.19803$ ).


Model	$\rho(0)/\rho(R)$
Model 1 ( $a_2 = 0.53, a_3 = 0.57$ )	2.49405
Model 2 ( $a_2 = 0.3, a_3 = 0.2$ )	1.97267
Model 3 ( $a_2 = 0.31, a_3 = 0.2$ )	1.92661
Model 4 ( $a_2 = 0.8, a_3 = 0.5$ )	1.92086

**Table 3** Estimated values of the density ratio for Cen X-3 ( $u = 0.2035$ ).




**Fig. 7**  $\tilde{\rho}(r) - \tilde{P}_t(r)$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .


**Fig. 8**  $\tilde{\rho}(r) - \tilde{P}_t(r)$  as a function of  $r$  for Model 1: (a)  $u = 0.19803$ , (b)  $u = 0.2035$ , Model 2: (c)  $u = 0.19803$ , (d)  $u = 0.2035$ , Model 3: (e)  $u = 0.19803$ , (f)  $u = 0.2035$ , Model 4: (g)  $u = 0.19803$ , (h)  $u = 0.2035$ .



**Fig. 9** Sound velocities as a function of  $r$  for compactness factor  $u = 0.19803$ : (a) Radial velocity  $v_r$  and (b) Tangential velocity  $v_t$ . Models 1, 2, 3 and 4 are identified with blue, black, red and green line respectively.



**Fig. 10** Sound velocities as a function of  $r$  for compactness factor  $u = 0.2035$ : (a) Radial velocity  $v_r$  and (b) Tangential  $v_t$ . Models 1, 2, 3 and 4 are identified with blue, black, red and green line respectively.



**Fig. 11**  $Z$  for Model 1 (a) Model 2 (b) Model 3 (c) and Model 4 (d)

while Models 2, 3 and 4 can be used for Cen X-3.

It should be interesting to explore the response of the system against perturbation. However, this and other points are under active consideration to future works.

## 7 Acknowledgements

E.C acknowledges Decanato de Investigación y Creatividad, USFQ, Ecuador, for continuous support.

## 8 Appendix: Auxiliary functions

$$\zeta = a_2 + a_3 r^2$$

$$\chi = \ln \left( 1 + \frac{a_3}{a_2} r^2 \right)$$

$$\eta_1 = 6a_1 a_2^3 C^3 r^2 + a_2^2 a_3 C (3a_1 C r^2 \beta_2(r) + 8a_3^2) \\ + a_2 a_3^2 C r^2 (8a_3^2 - a_1 (C^2 r^4 + 6\beta_3(r))) \\ - 2a_1 a_3^3 r^2$$

$$\eta_2 = a_2^2 B (3a_3^2 - 2a_1 B r^2) - a_1 a_3^2 r^2 \\ + a_2 a_3 B r^2 (3a_3^2 - a_1 \gamma_2(r)),$$

$$\varrho_1 = a_3 A (3a_2 + a_3 r^2) + a_2 B (a_2 - a_3 r^2)$$

$$\varrho_2 = 8a_2 a_3^3 C^2 (C r^2 \beta_5(r) + 6) \zeta(r)^2 + a_1 \mathcal{S}_1(r)$$

$$\varrho_3 = 3a_2 a_3^2 B^2 (B r^2 + 3) \zeta(r)^2 + a_1 \mathcal{S}_2(r)$$

$$\mathcal{P}_1 = a_2 B^2 (5a_1 r^2 + 8a_3 \zeta(r)) - a_1 A B (a_2 - 5a_3 r^2) \\ - a_1 a_3 A^2$$

$$\mathcal{P}_2 = 4a_1 a_3 A B r^2 (2a_2 + a_3 r^2) \\ + a_2 B^2 \left( 4a_2 r^2 (a_1 + 4a_3^2) + a_3 r^4 (a_1 + 8a_3^2) \right. \\ \left. + 8a_2^2 a_3 \right) - a_1 a_2 a_3 A^2.$$

$$\mathcal{P}_3 = 6a_1 a_2 C \beta_8(r) (a_3 - a_2 C)^2 \zeta(r) \chi(r) \\ + 8a_2 a_3^4 C \beta_1^3 \zeta(r) + a_3 \beta_8(r) \left[ -6a_1 a_2^3 C^3 r^2 \right. \\ \left. + a_2^2 a_3 C (-3a_1 C r^2 \beta_9(r) - 8a_3^2) \right. \\ \left. + a_2 a_3^2 C r^2 (a_1 (C r^2 \beta_{10}(r) - 6) - 8a_3^2) \right. \\ \left. + 2a_1 a_3^3 r^2 \right],$$

$$\mathcal{P}_4 = -18a_1 a_2 C^2 (a_3 - a_2 C)^2 \zeta(r)^2 \chi(r) \\ + 24a_2 a_3^4 C^2 \chi(r)^2 + 6a_2^2 a_3^2 C^2 r^2 (C r^2 \beta_7(r) + 3) \\ + a_1 a_3 \left[ 18a_2^4 C^4 r^2 + 9a_3^2 a_3 C^3 r^2 (3C r^2 - 4) \right. \\ \left. - a_2 a_3^3 (8C^4 r^8 + 34C^3 r^6 + 14C r^2 + 1) \right. \\ \left. - 6a_3^4 C r^4 \right].$$

$$\mathcal{P}_5 = a_3 r^2 \left[ a_1 \gamma_6(r) (2a_2^2 B^2 + a_2 a_3 B \gamma_2(r) + a_3^2) \right. \\ \left. + 3a_2 a_3^2 B^2 \gamma_7(r) \zeta(r) \right] \\ - 2a_1 a_2 B \gamma_6(r) (a_2 B - a_3) \zeta(r) \chi(r).$$

$$\mathcal{P}_6 = 15a_2 a_3^3 B^2 \zeta(r)^2 + 10a_1 a_2 B^2 (a_2 B - a_3) \zeta(r)^2 \chi(r) \\ + a_1 a_3 \left[ -10a_2^3 B^3 r^2 + 5a_2^2 a_3 B^2 r^2 \gamma_8(r) - 5a_3^3 B r^4 \right. \\ \left. + a_2 a_3^2 (B r^2 (-9B^2 r^4 + B r^2 - 11) - 1) \right],$$

$$\mathcal{S}_1 = 6a_2^4 C^3 \beta_4(r) + 3a_2^3 a_3 C^2 (3C r^2 \beta_6(r) + 4) \\ + 2a_2^2 a_3^2 C \beta_4(r) (C r^2 \beta_7(r) + 3) - 2a_3^4 r^2 \beta_4(r) \\ + a_2 a_3^3 (C r^2 (C r^2 (C r^2 \beta_3(r) + 48) + 4) + 6)$$

$$\mathcal{S}_2 = -2a_2^3 B^2 \gamma_3(r) + a_2^2 a_3 B (B r^2 \gamma_4(r) - 2) \\ + a_2 a_3^2 (B r^2 (B r^2 \gamma_5(r) + 3) + 3) \\ - a_3^3 r^2 \gamma_3(r)$$

$$\beta_1 = C r^2 + 1 \quad \beta_2 = C r^2 - 4 \quad \beta_3 = C r^2 - 1 \\ \beta_4 = 3C r^2 - 1 \quad \beta_5 = C r^2 + 3 \quad \beta_6 = 3C r^2 - 5 \\ \beta_7 = C r^2 - 9 \quad \beta_8 = 9C r^2 + 1 \quad \beta_9 = C r^2 - 4 \\ \beta_{10} = C r^2 + 6 \\ \beta_{11} = -6a_2^3 C^3 - 3a_2^2 a_3 C^2 \beta_9 + 2a_3^3 \\ + a_2 a_3^2 C (C r^2 \beta_{10} - 6),$$

$$\gamma_1 = B r^2 + 1 \quad \gamma_2 = B r^2 - 2 \quad \gamma_3 = B r^2 - 1 \\ \gamma_4 = 5 - 3B r^2 \quad \gamma_5 = B r^2 + 9 \quad \gamma_6 = 7B r^2 + 1 \\ \gamma_7 = B r^2 - 5 \quad \gamma_8 = 2 - 3B r^2. \\ \gamma_9 = 2a_2^2 B^2 + a_2 a_3 B \gamma_2 + a_3^2.$$

## References

1. R. Chan, L. Herrera and N. O. Santos, *Mon. Not. R. Astr. Soc.* **265**, 533 (1993).
2. L. Herrera and N. O. Santos, *Phys. Rep.* **286**, 53 (1997).
3. L. Herrera, J. Martin and J. Ospino, *J. Math. Phys.* **43**, 4889 (2002).
4. L. Herrera, A. Di Prisco, J. Martin, J. Ospino, N. O. Santos and O. Troconis, *Phys. Rev. D* **69**, 084026 (2004).
5. L. Herrera, J. Ospino, A. Di Prisco, E. Fuenmayor and O. Troconis, *Phys. Rev. D* **79** 064025 (2009).
6. L. Herrera, J. Ospino and A. Di Prisco, *Phys. Rev. D* **77**, 027502 (2008).
7. L. Herrera, N. O. Santos and A. Wang, *Phys. Rev. D* **78**, 084026 (2008).
8. P. H. Nguyen and J. F. Pedraza, *Phys. Rev. D* **88**, 064020 (2013).
9. P. H. Nguyen and M. Lingam, *Monh. Not. R. Astron. Soc.* **436**, 2014 (2013).
10. J. Krisch and E. N. Glass, *J. Math. Phys.* **54**, 082501 (2013).
11. R. Sharma and B. Ratanpal, *Int. J. Mod. Phys. D* **22**, 1350074 (2013).
12. E. N. Glass, *Gen. Relativ. Gravit.* **45**, 2661 (2013).
13. K. P. Reddy, M. Govender and S. D. Maharaj, *Gen. Relativ. Gravit.* **47**, 35 (2015).

14. H. Hernández, D. Suárez–Urango and L. A. Núñez, *Eur. Phys. J. C* **81**, 241 (2021)
15. D. Suárez–Urango, L. A. Núñez and H. Hernández, arXiv:2102.00496
16. D. Suárez–Urango, J. Ospino, H. Hernández and L. A. Núñez, arXiv:2104.08923.
17. J. C. Kemp, J. B. Swedlund, J. D. Landstreet and J. R. P. Angel, *Astrophys. J.* **161**, L77 (1970).
18. G. D. Schmidt and P. S. Schmidt, *Astrophys. J.* **448**, 305 (1995).
19. A. Putney, *Astrophys. J.* **451**, L67 (1995).
20. D. Reimers, S. Jordan, D. Koester, N. Bade, Th. Kohler and L. Wisotzki, *Astron. Astrophys.* **311**, 572 (1996).
21. A. P. Martínez, R. G. Felipe and D. M. Paret, *Int. J. Mod. Phys. D* **19**, 1511 (2010).
22. M. Chaichian, S. S. Masood, C. Montonen, A. Perez Martínez and H. Perez Rojas, *Phys. Rev. Lett* **84**, 5261 (2000).
23. A. Perez Martínez, H. Perez Rojas and H. J. Mosquera Cuesta, *Eur. Phys. J. C* **29**, 111 (2003).
24. A. Perez Martínez, H. Perez Rojas and H. J. Mosquera Cuesta, *Int. J. Mod. Phys. D* **17**, 2107 (2008).
25. E. J. Ferrer, V. de la Incera, J. P. Keith, I. Portillo and P. L. Springsteen, *Phys. Rev. C* **82**, 065802 (2010).
26. N. Andersson, G. Comer and K. Glampedakis, *Nucl. Phys. A* **763**, 212 (2005).
27. B. Sa'd, I. Shovkovy and D. Rischke, *Phys. Rev. D* **75**, 125004 (2007).
28. M. Alford and A. Schmitt, *AIP Conf. Proc.* **964**, 256 (2007).
29. D. Blaschke and J. Berdermann, arXiv: 0710.5243.
30. A. Drago, A. Lavagno and G. Pagliara, *Phys. Rev. D* **71**, 103004 (2005).
31. P. B. Jones, *Phys. Rev. D* **64**, 084003 (2001).
32. E. N. E. van Dalen and A. E. L. Dieperink, *Phys. Rev. C* **69**, 025802 (2004).
33. H. Dong, N. Su and O. Wang, *J. Phys. G* **34**, S643 (2007).
34. L. Herrera, *Phys. Rev. D* **101**, 104024 (2020).
35. I. Lopes, G. Panotopoulos and Á. Rincón, *Eur. Phys. J. Plus* **134** (2019) no.9, 454 doi:10.1140/epjp/i2019-12842-4 [arXiv:1907.03549 [gr-qc]].
36. G. Panotopoulos, Á. Rincón and I. Lopes, *Eur. Phys. J. C* **80** (2020) no.4, 318 doi:10.1140/epjc/s10052-020-7900-3 [arXiv:2004.02627 [gr-qc]].
37. P. Bhar, F. Tello-Ortiz, Á. Rincón and Y. Gomez-Leyton, *Astrophys. Space Sci.* **365** (2020) no.8, 145 doi:10.1007/s10509-020-03859-6
38. F. Tello-Ortiz, M. Malaver, Á. Rincón and Y. Gomez-Leyton, *Eur. Phys. J. C* **80** (2020) no.5, 371 doi:10.1140/epjc/s10052-020-7956-0 [arXiv:2005.11038 [gr-qc]].
39. F. Tello-Ortiz, Á. Rincón, P. Bhar and Y. Gomez-Leyton, *Chin. Phys. C* **44** (2020), 105102 doi:10.1088/1674-1137/aba5f7 [arXiv:2006.04512 [gr-qc]].
40. J. Ovalle, *Phys. Rev. D* **95**, 104019 (2017).
41. J. Ovalle, L. Á. Gergely and R. Casadio, *Class. Quant. Grav.* **32** (2015), 045015.
42. R. Casadio, J. Ovalle and R. da Rocha, *Class. Quant. Grav.* **32** (2015) no.21, 215020
43. J. Ovalle, R. Casadio and A. Sotomayor, *Adv. High Energy Phys.* **2017** (2017), 9756914 doi:10.1155/2017/9756914 [arXiv:1612.07926 [gr-qc]]. LaTeX (EU)
44. J. Ovalle, R. Casadio, R. da Rocha, A. Sotomayor. *Eur. Phys. J. C* **78**, 122 (2018).
45. J. Ovalle, R. Casadio, R. da Rocha, A. Sotomayor and Z. Stuchlik, *EPL* **124**, 20004 (2018).
46. M. Estrada, F. Tello-Ortiz. *Eur. Phys. J. Plus* **133**, 453 (2018) .
47. J. Ovalle, R. Casadio, R. da Rocha, A. Sotomayor, Z. Stuchlik, *Eur. Phys. J. C* **78**, 960 (2018).
48. C. Las Heras, P. Leon. *Fortschr. Phys.* **66**, 1800036 (2018).
49. G. Panotopoulos, Á. Rincón, *Eur. Phys. J. C* **78**, 851 (2018)
50. J. Ovalle, C. Posada, Z. Stuchlik, *Class. Quant. Grav.* **36**, no. 20, 205010 (2019).
51. J. Ovalle, *Phys. Lett. B* **788**, 213 (2019).
52. M. Estrada, R. Prado, *Eur. Phys. J. Plus* **134**, 168 (2019).
53. S. Maurya, F. Tello, *Eur. Phys. J. C* **79**, 85 (2019).
54. C. Las Heras, P. León, *Eur. Phys. J. C* **79**, no. 12, 990 (2019).
55. M. Estrada, *Eur. Phys. J. C* **79**, no. 11, 918 (2019).
56. L. Gabbanelli, J. Ovalle, A. Sotomayor, Z. Stuchlik, R. Casadio, *Eur. Phys. J. C* **79**, 486 (2019).
57. S. Hensh and Z. Stuchlik, *Eur. Phys. J. C* **79**, no. 10, 834 (2019).
58. F. Linares and E. Contreras, *Phys. Dark Univ.* **28**, 100543 (2020)
59. P. León and A. Sotomayor, *Fortsch. Phys.* **67**, 1900077 (2019).
60. R. Casadio, E. Contreras, J. Ovalle, A. Sotomayor, and Z. Stuchlik, *Eur. Phys. J. C* **79**, 826 (2019).
61. S. Maurya, and F. Tello-Ortiz, *Physics of the Dark Universe*, **27**, 100442 (2020).
62. A. Arias, F. Tello-Ortiz and E. Contreras, *Eur. Phys. J. C* **80**, 463 (2020).
63. G. Abellán, V. Torres-Sánchez, E. Fuenmayor, and E. Contreras, *Eur. Phys. J. C* **80**, 177 (2020).
64. F. Tello-Ortiz, *Eur. Phys. J. C* **80**, 413 (2020).
65. A. Rincón, E. Contreras, F. Tello-Ortiz, P. Bargeño, and G. Abellán, *Eur. Phys. J. C* **80**, 490 (2020)
66. J. Ovalle and R. Casadio, *Beyond Einstein Gravity. The Minimal Geometric Deformation Approach in the Brane-World*, Springer International Publishing (2020). DOI:10.1007/978-3-030-39493-6.
67. G. Abellán, Á. Rincón, E. Fuenmayor and E. Contreras, *Eur. Phys. J. Plus* **135** (2020) no.7, 606 doi:10.1140/epjp/s13360-020-00589-0
68. J. Ovalle, R. Casadio, E. Contreras and A. Sotomayor, *Phys. Dark Univ.* **31** (2021), 100744
69. E. Contreras, J. Ovalle and R. Casadio, *Phys. Rev. D* **103**, 044020 (2021).
70. C. L. Heras and P. Leon, *Eur. Phys. J. Plus* **136**, 828 (2021).
71. J. Ovalle, E. Contreras, Z. Stuchlik, *Physical Review D* **103**, 084016 (2021).
72. F. Tello, S. Maurya and P. Bargeño, *Eur. Phys. J. C* **81**, 426 (2021).
73. P. Meert and R. da Rocha, *Nucl. Phys. B* **967**, 115420 (2021) doi:10.1016/j.nuclphysb.2021.115420 [arXiv:2006.02564 [gr-qc]].
74. R. da Rocha, *Phys. Rev. D* **102**, no.2, 024011 (2020) doi:10.1103/PhysRevD.102.024011 [arXiv:2003.12852 [hep-th]].
75. R. da Rocha, *Symmetry* **12**, no.4, 508 (2020) doi:10.3390/sym12040508 [arXiv:2002.10972 [hep-th]].
76. M. Sharif and Q. Ama-Tul-Mughani, *Annals Phys.* **415**, 168122 (2020) doi:10.1016/j.aop.2020.168122 [arXiv:2004.07925 [gr-qc]].
77. M. Sharif and A. Majid, *Astrophys. Space Sci.* **365**, no.2, 42 (2020) doi:10.1007/s10509-020-03754-0

- 
78. M. Sharif and Q. Ama-Tul-Mughani, *Mod. Phys. Lett. A* **35**, no.12, 2050091 (2020) doi:10.1142/S0217732320500911
  79. M. Estrada, arXiv:2106.02166
  80. M. Zubair, M. Amin and H. Azmat, *Phys. Scripta* **96** (2021) no.12, 125008 doi:10.1088/1402-4896/ac237d
  81. H. Azmat and M. Zubair, *Eur. Phys. J. Plus* **136** (2021) no.1, 112 doi:10.1140/epjp/s13360-021-01081-z [arXiv:2106.08384 [gr-qc]].
  82. Q. Muneer, M. Zubair and M. Rahseed, *Phys. Scripta* **96** (2021) no.12, 125015 doi:10.1088/1402-4896/ac1216
  83. M. Zubair and H. Azmat, *Annals Phys.* **420** (2020), 168248 doi:10.1016/j.aop.2020.168248
  84. S. K. Maurya, A. S. A. Kindi, M. R. A. Hatmi and R. Nag, *Results Phys.* **29** (2021), 104674 doi:10.1016/j.rinp.2021.104674
  85. P. Meert and R. da Rocha, [arXiv:2109.06289 [hep-th]].
  86. J. Sultana, *Symmetry* **13** (2021) no.9, 1598 doi:10.3390/sym13091598
  87. S. K. Maurya, A. Pradhan, F. Tello-Ortiz, A. Banerjee and R. Nag, *Eur. Phys. J. C* **81** (2021) no.9, 848 doi:10.1140/epjc/s10052-021-09628-1
  88. L. Herrera, *Phys. Rev. D* **97**, 044010 (2018).
  89. A. García - Parrado Gomez Lobo arXiv:0707.1475v2.
  90. R. C. Tolman, "Static solutions of Einstein's field equations for spheres of fluid." *Physical Review* 55.4: 364 (1939).
  91. M. L. Rawls, et al. "Refined neutron star mass determinations for six eclipsing x-ray pulsar binaries." *The Astrophysical Journal* 730.1: 25 (2011).
  92. V. A. Torres and E. Contreras, "Anisotropic neutron stars by gravitational decoupling." *The European Physical Journal C* 79.10: 1-8 (2019).
  93. S. K. Maurya, A. Banerjee and Y. K. Gupta. "Exact solution of anisotropic compact stars via mass function." *Astrophysics and Space Science* 363.10: 1-9 (2018).
  94. A. K. Prasad, et al. "Relativistic model for anisotropic compact stars using Karmarkar condition." *Astrophysics and Space Science* 364.4: 1-12 (2019).
  95. M. S. R. Delgaty and K. Lake. "Physical acceptability of isolated, static, spherically symmetric, perfect fluid solutions of Einstein's equations." *Computer Physics Communications* 115.2-3: 395-415 (1998).
  96. M. Wyman, "Radially symmetric distributions of matter." *Physical Review* 75.12: 1930 (1949).
  97. M. C. Durgapal, "A class of new exact solutions in general relativity." *Journal of Physics A: Mathematical and General* 15.8: 2637 (1982).
  98. H. Heintzmann, "New exact static solutions of Einsteins field equations." *Zeitschrift für Physik* 228.4: 489-493 (1969).
  99. M. Spivak, *Calculus 2* Edic. Reverte, 1995.
  100. S. Saitoh, "Division by zero calculus." (2017).
  101. S. M. Carroll, *Spacetime and geometry*. Cambridge University Press, 2019.
  102. B. V. Ivanov, "Analytical study of anisotropic compact star models." *The European Physical Journal C* 77.11: 1-12 (2017).
  103. T. Gangopadhyay, et al. "Strange star equation of state fits the refined mass measurement of 12 pulsars and predicts their radii." *Monthly Notices of the Royal Astronomical Society* 431.4: 3216-3221 (2013).
  104. D. M. Pandya, et al. "Anisotropic compact star model satisfying Karmarkar conditions." *Astrophysics and Space Science* 365.2 (2020): 1-10.
  105. S. Maurya, "Extended gravitational decoupling (GD) solution for charged compact star model." *European Physical Journal C* 80.5 (2020): 1-16.