

The six operations in topology

Marco Volpe*

January 9, 2023

Abstract

In this paper we show that the six functor formalism for sheaves on locally compact Hausdorff topological spaces, as developed for example in Kashiwara and Schapira's book *Sheaves on Manifolds*, can be extended to sheaves with values in any closed symmetric monoidal ∞ -category which is stable and bicomplete. Notice that, since we do not assume that our coefficients are presentable or restrict to hypercomplete sheaves, our arguments are not obvious and are substantially different from the ones explained by Kashiwara and Schapira. Along the way we also study locally contractible geometric morphisms and prove that, if $f : X \rightarrow Y$ is a continuous map which induces a locally contractible geometric morphism, then the exceptional pullback functor $f^!$ preserves colimits and can be related to the pullback f^* . At the end of our paper we also show how one can express Atiyah duality by means of the six functor formalism.

Contents

1	Introduction	2
1.1	Linear overview	5
1.2	Acknowledgements	6
2	Sheaves and tensor products	6
2.1	Tensor product of cocomplete ∞ -categories	7
2.2	Sheaves and cosheaves	12
3	Shape theory and shape submersions	19
3.1	Relative shape	19
3.2	Locally contractible geometric morphisms	23
3.3	Shape submersions	27
4	Localization sequences	30
5	Pullbacks with stable bicomplete coefficients	33
5.1	Recollections on covariant Verdier duality	34
5.2	Dualizability of spectral sheaves	39
5.3	The pullback $f_{\mathbb{C}}^*$	41
6	Six functor formalism	44
6.1	The formulas for $f_!^{\mathbb{C}}$	44
6.2	$f_!^{\mathbb{C}}$ when f is a locally contractible geometric morphism	49

*University of Regensburg, Universitätsstraße 31, Regensburg. email: marco.volpe@ur.de. The author was supported by the SFB 1085 (Higher Invariants) in Regensburg, Germany, funded by the DFG (German Science Foundation).

7	Relative Atiyah Duality	52
7.1	Thom spaces and the J-homomorphism	52
7.2	Relative Atiyah duality	56

1 Introduction

Tell all the truth but tell it slant –
 Success in Circuit lies
 Too bright for our infirm Delight
 The Truth's superb surprise
 As Lightning to the Children eased
 With explanation kind
 The Truth must dazzle gradually
 Or every man be blind –

Emily Dickinson

One of the most complete and general reference dealing with the *six functor formalism* for sheaves on topological spaces is Masaki Kashiwara and Pierre Schapira's seminal book *Sheaves on Manifolds* [KS90]. However, for technical reasons related to the construction of derived functors, the authors there restrict themselves to bounded derived categories of sheaves of R -modules, where R is assumed to have finite global dimension. From a modern perspective, considering ∞ -categorical enhancements of derived categories, this can be regarded as the full subcategory of hypercomplete sheaves with values in $D(R)$ spanned by bounded objects.

For many applications though, one would like to be able to consider *non-hypercomplete* sheaves with values in *unbounded* derived categories of *any* ring. On the other hand, more recent papers such as [RS18], [JT17], [Jin19] and [Jin20], justify the need of even further generalizations to sheaves of modules over *ring spectra*, in order to apply the power of six functors to generalized cohomology theories. The work of Voevodsky on *stable motivic homotopy theory* has provided an analog of such constructions in the world of algebraic geometry (see [CD19] for a textbook source on the subject), whereas topologists have succeeded only partially in this direction by introducing *parametrized spectra* (see [MS06]), which correspond to *locally constant* sheaves of spectra. In this paper we exploit the power of the now established theory of ∞ -categories (as developed for example in [Lur09] or [Cis19]) to extend the six functor formalism on locally compact Hausdorff spaces to a much broader setting.

Let $f : X \rightarrow Y$ be a continuous map between locally compact Hausdorff topological spaces and \mathcal{C} is any stable bicomplete (i.e. complete and cocomplete) ∞ -category. For any such map, we construct adjunctions

$$\begin{array}{ccc}
 \text{Shv}(Y; \mathcal{C}) & \begin{array}{c} \xrightarrow{f_{\mathcal{C}}^*} \\ \perp \\ \xleftarrow{f_{\mathcal{C}}^!} \end{array} & \text{Shv}(X; \mathcal{C})
 \end{array}
 \quad
 \begin{array}{ccc}
 \text{Shv}(X; \mathcal{C}) & \begin{array}{c} \xrightarrow{f_{\mathcal{C}}^!} \\ \perp \\ \xleftarrow{f_{\mathcal{C}}^*} \end{array} & \text{Shv}(Y; \mathcal{C}).
 \end{array}$$

To do this, we make use of Lurie's Verdier duality equivalence ([Lur17, Theorem 5.5.5.1])

$$\mathbb{D}_{\mathcal{C}} : \text{Shv}(X; \mathcal{C}) \xrightarrow{\simeq} \text{CoShv}(X; \mathcal{C})$$

where the target is the ∞ -category of \mathcal{C} -valued *cosheaves* on X , i.e. $\text{Shv}(X; \mathcal{C}^{\text{op}})^{\text{op}}$. The adjunctions above are then defined to satisfy natural equivalences

$$\mathbb{D}_{\mathcal{C}} f_{\mathcal{C}}^{\mathcal{C}} \simeq (f_{\mathcal{C}}^{\text{op}})^{\text{op}} \mathbb{D}_{\mathcal{C}} \quad \mathbb{D}_{\mathcal{C}} f_{\mathcal{C}}^{\mathcal{C}} \simeq (f_{\mathcal{C}}^{\text{op}})^{\text{op}} \mathbb{D}_{\mathcal{C}}.$$

Notice that, since we do not require \mathcal{C} to be presentable, the existence of $f_{\mathcal{C}}^*$ is not at all obvious (see the discussion in Remark 2.34), and so we will have to work a bit harder than one might expect. Nevertheless, even though the presentability assumption will be enough for applications, we would like to point out that our efforts to make the results in this paper as general as possible are not vain, and actually lead to many advantages. First of all, with our definition of $f_{\mathcal{C}}^{\mathcal{C}}$, it is basically immediate to verify that the lower shrieks are functorial with respect to compositions of continuous maps (see Lemma 6.2). Moreover, by working with a class of coefficients closed under the operation of passing to the opposite category, we do not break the symmetry which comes from Verdier duality, meaning that whenever we prove some result involving the functors $f_*^{\mathcal{C}}$ and $f_{\mathcal{C}}^*$ that is true for all \mathcal{C} stable and bicomplete, we immediately obtain a dual theorem involving the functors $f_{\mathcal{C}}^{\mathcal{C}}$ and $f_{\mathcal{C}}^!$, and viceversa. Another way to put it is that our formalism applies with no distinctions to sheaves or cosheaves: this will be used in a follow-up paper (see [Vol22]) in which we will prove a duality theorem for constructible sheaves on a conically smooth stratified space, as we will need to extend some results about constructible sheaves, such as homotopy invariance (see [Hai20]) or the exodromy equivalence (see [PT22]), to constructible cosheaves. It is also worth noticing that, even if one would restrict to presentable coefficients, it would nevertheless be desirable to have formulas such as $f_{\mathcal{C}}^{\text{Sp}} \otimes \mathcal{C} \simeq f_{\mathcal{C}}^{\mathcal{C}}$, and to get this one would still need to verify everything we prove in Section 5.

We try to outline the key ingredients in our paper that allow us to work with non-presentable coefficients. As we explained above, the main difficulty lies in showing the existence of the pullback functor $f_{\mathcal{C}}^*$. The main tool we employ to carry out this purpose is Lurie's tensor product of cocomplete ∞ -categories as defined in [Lur17, 4.8.1]: in particular, a property of this tensor product that we will use over and over is that it preserves adjunctions between cocontinuous functors (see Remark 2.2). We show in Lemma 2.12 that it restricts to a monoidal structure on $\text{Cocont}_{\infty}^{st}$ (i.e the ∞ -category of stable cocomplete ∞ -categories with cocontinuous functors between them) and use it to formulate the following theorem.

Theorem 1.1 (Corollary 5.16). Let \mathcal{C} be a stable bicomplete ∞ -category. Then there is an equivalence

$$\text{Shv}(X; \text{Sp}) \otimes \mathcal{C} \simeq \text{Shv}(X; \mathcal{C})$$

where \otimes on the left-hand side denotes the tensor product of stable cocomplete ∞ -categories.

Theorem 1.1 will play a crucial role in what follows, because it will allow us to reduce a lot of arguments involving cocontinuous functors to the case of sheaves of spectra. To prove Theorem 1.1, we start by observing in Theorem 5.15 that the model of \mathcal{K} -sheaves (see [Lur09, Theorem 7.3.4.9]) implies that $\text{Shv}(X; \text{Sp})$ is a strongly dualizable object in $\text{Cocont}_{\infty}^{st}$, where Sp denotes the ∞ -category of spectra (see also [Lur16, Proposition 21.1.7.1]). Hence, for \mathcal{C} any stable and cocomplete ∞ -category, we get an equivalence

$$\text{CoShv}(X; \text{Sp}) \otimes \mathcal{C} \simeq \text{CoShv}(X; \mathcal{C}).$$

Combined with Verdier duality, this gives Theorem 1.1.

Having Theorem 1.1 at hand, the question of constructing $f_{\mathcal{C}}^*$ can be reduced to the case of sheaves of spectra, where we can directly use the existence of sheafification by the presentability of Sp . More precisely, we first observe that any map f can be factored as the composition of a closed immersion, an open immersion and a proper map (see factorization (5.1)), and then prove the existence of a left adjoint to $f_*^{\mathcal{C}}$ in these separate cases. When f is an open immersion this is done easily in Lemma 5.19, and the only non-trivial part consists in the following theorem.

Theorem 1.2 (Lemma 5.14, Proposition 5.18). Let $f : X \rightarrow Y$ be a proper map between locally compact Hausdorff topological spaces. Then the pushforward

$$f_*^{\mathcal{C}} : \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(Y; \mathcal{C})$$

preserves colimits. Furthermore, there is a commutative square

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathcal{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \mathrm{Shv}(X; \mathcal{C}) \\ \downarrow f_*^{\mathcal{S}\mathrm{p}} \otimes \mathcal{C} & & \downarrow f_*^{\mathcal{C}} \\ \mathrm{Shv}(Y; \mathcal{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \mathrm{Shv}(Y; \mathcal{C}). \end{array}$$

The proof of Theorem 1.2 essentially consists of providing a convenient description of $f_*^{\mathcal{C}}$ through the model of \mathcal{K} -sheaves, which is easily seen to preserve colimits and to be compatible with Verdier duality. We then achieve our final goal observing that, since $f_*^{\mathcal{S}\mathrm{p}} \otimes \mathcal{C}$ admits a left adjoint of the form $f_{\mathcal{S}\mathrm{p}}^* \otimes \mathcal{C}$, the same is true for $f_*^{\mathcal{C}}$. In particular by taking f to be the projection $X \rightarrow *$, we see that the global section functor

$$\mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathcal{C}$$

admits a left adjoint. As a consequence, using the results in [Cis19, 6.7], we show in Theorem 5.21 that the inclusion of $\mathrm{Shv}(X; \mathcal{C})$ in \mathcal{C} -valued presheaves on X admits a left adjoint.

The discussion above involves only the four functors $f_*^{\mathcal{C}}$, $f_{\mathcal{C}}^*$, $f_!^{\mathcal{C}}$ and $f_{\mathcal{C}}^!$, but what about the other two? Our first observation is that, a priori, there is no need to require our category of coefficients to have a monoidal structure to make sense of things like projection formulas or Künneth formulas. To be more precise, one can show that there is a functor

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(Y; \mathcal{D}) & \longrightarrow & \mathrm{Shv}(X \times Y; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{D}) \\ (F, G) & \longmapsto & F \boxtimes G \end{array}$$

which preserves colimits in both variables and induces an equivalence

$$\mathrm{Shv}(X; \mathcal{C}) \otimes \mathrm{Shv}(Y; \mathcal{D}) \simeq \mathrm{Shv}(X \times Y; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{D}).$$

Taking $X = Y$ and composing with $\Delta_{\mathcal{S}\mathrm{p}}^* \otimes (\mathcal{C} \otimes \mathcal{D})$, where $\Delta : X \hookrightarrow X \times X$ is the diagonal embedding, we get a variablewise colimit preserving functor denoted as

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(Y; \mathcal{D}) & \longrightarrow & \mathrm{Shv}(X; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{D}) \\ (F, G) & \longmapsto & F \otimes G \end{array}$$

(see Construction 2.25 and Remark 5.22 for more details). For this kind of tensor product of sheaves, we prove the following formulas.

Theorem 1.3 (Corollary 2.31, Proposition 6.12, Proposition 6.11). Let \mathcal{C} and \mathcal{D} be stable and bicomplete ∞ -categories. Then we have the following functorial identifications

$$\begin{aligned} f_{\mathcal{C} \otimes \mathcal{D}}^*(F \otimes G) &\simeq f_{\mathcal{C}}^* F \otimes f_{\mathcal{D}}^* G \\ f_!^{\mathcal{C} \otimes \mathcal{D}}(F \otimes f_{\mathcal{D}}^* G) &\simeq f_!^{\mathcal{C}} F \otimes G \\ (f \times g)_!^{\mathcal{C} \otimes \mathcal{D}}(F \boxtimes G) &\simeq f_!^{\mathcal{C}} F \boxtimes g_!^{\mathcal{D}} G. \end{aligned}$$

In particular, when \mathcal{C} admits a monoidal structure whose tensor preserves colimits in both variables, one obtains a cocontinuous functor

$$\mathrm{Shv}(X; \mathbb{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

whose composition with (2.27) induces a monoidal structure on $\mathrm{Shv}(X; \mathcal{C})$: this way we can deduce all the analogous formulas in the monoidal setting. If \mathcal{C} is also closed, we deduce their dual versions involving the internal homomorphism functor (see Remark 2.29).

We describe one last advantage of our general rendition of the six functor formalism. In Definition 3.12 we define *locally contractible geometric morphisms* (see also [AC21, Definition 3.2.1]). Later, in Definition 3.21, we specify a vast class of continuous maps between topological spaces called *shape submersions* which induce a locally contractible geometric morphism (see Corollary 3.26). Topological submersions are examples of such morphisms, but our definition is much more general in the sense that it does not force the fibers to be topological manifolds. Another illustrating example to keep in mind is that of the unique map $X \rightarrow *$, when X is any CW-complex. An easy implementation of our machinery generalizes [KS90, Proposition 3.3.2] and [Ver65, Section 5] beyond the case of submersive maps.

Theorem 1.4 (Proposition 6.16). Let $f : X \rightarrow Y$ be a map which induces a locally contractible geometric morphism, and let \mathcal{C} be a stable and bicomplete ∞ -category. Then $f_{\mathcal{C}}^!$ admits a right adjoint and we have a formula

$$f_{\mathcal{C}}^! F \otimes f_{\mathcal{D}}^* G \simeq f_{\mathcal{C} \otimes \mathcal{D}}^! (F \otimes G).$$

Then, inspired by parallel results in motivic homotopy theory, we conclude our paper by formulating and proving a relative version of Atiyah duality.

Theorem 1.5 (Corollary 7.14). Let $f : X \rightarrow Y$ be a proper submersion between smooth manifolds. Denote by $\mathbb{S}_X \in \mathrm{Shv}(X; \mathbb{S}\mathrm{p})$ the constant sheaf at the sphere spectrum, and by $\mathrm{Th}(-Tf)$ the sheaf the *Thom spectrum* of the virtual vector bundle $-Tf$. Then $f_{\sharp}(\mathbb{S}_X)$ is strongly dualizable with dual $\mathrm{Th}(-Tf)$.

1.1 Linear overview

We now give a linear overview of the contents of our paper.

In section 2 we recall the definition of Lurie's tensor product of cocomplete ∞ -categories and prove some of its basic properties. In particular, we will interpret the results in [Cis19, 6.7] in terms of this tensor product in Theorem 2.10, show that it preserves the property of being stable in Lemma 2.12, and show that compactly generated stable ∞ -category is a strongly dualizable object in the symmetric monoidal ∞ -category $\mathrm{Cocont}_{\infty}^{st}$ in Proposition 2.14. Afterwards we recall the definition of sheaves and cosheaves with values in a general ∞ -category and explain how Lurie's tensor product can be used to conveniently describe $\mathrm{Shv}(X; \mathcal{C})$ at least when \mathcal{C} is presentable. Most of the results in this section are not original, but we still felt the necessity to spend some time writing them up to make our discussion as self contained and reader friendly as possible.

In section 3 we define for any geometric morphism $\mathcal{Y} \rightarrow \mathcal{X}$ the *relative shape* $\Pi_{\infty}^{\mathcal{X}}(\mathcal{Y})$ as a pro-object of \mathcal{X} and describe explicitly in Proposition 3.3 how this construction can be enhanced to a functor

$$\Pi_{\infty}^{\mathcal{X}} : \mathcal{T}\mathrm{op}/_{\mathcal{X}} \rightarrow \mathrm{Pro}(\mathcal{X})$$

and, even more, to a lax natural transformation between functors $\mathcal{T}\mathrm{op}^{\mathrm{op}} \rightarrow \mathrm{Cat}_{\infty}$ (see Remark 3.7): these coherent structures with which we equip the shape will be used to prove

easily that the shape is homotopy invariant in Corollary 3.4 and later in Proposition 7.7 to show that the Thom spectrum gives a natural transformation of sheaves of \mathbb{E}_∞ -spaces. Later we define locally contractible geometric morphisms and give a characterization in Proposition 3.11 which mimics the one in [Joh02, C3.3] for the locally connected case. We also show that, when f is a geometric morphism induced by a continuous map of topological spaces, the property of being locally contractible is checked more easily. Then we define shape submersions and prove in Lemma 3.25 a base change formula which will imply that they induce locally contractible geometric morphisms (see Corollary 3.26).

In section 4 we follow the approach of [Kha19] to obtain the localization sequences associated to a decomposition of a topological space into an open subset and its closed complement. Also the results here are not so new but, after section 5, they will imply that there is a recollement of $\mathrm{Shv}(X; \mathcal{C})$ associated to any open-closed decomposition of X whenever \mathcal{C} is stable and bicomplete, while this was previously known only for \mathcal{C} presentable.

Section 5 is devoted to Verdier duality, and how it can be used to show that the pushforward $f_*^{\mathcal{C}}$ admits a left adjoint for any \mathcal{C} stable and bicomplete in the way we have sketched at the beginning of the introduction.

In section 6 we develop the six functor formalism: as usual, we prove base change (Proposition 6.9), projection (Proposition 6.12) and Künneth (Proposition 6.11) formulas for $f_!^{\mathcal{C}}$, and discuss the properties of $f_{\mathcal{C}}^!$ when f is a shape submersion in Proposition 6.16.

At last, in section 7 we show how the six functor formalism can be used to express a relative version of Atiyah duality for any proper submersion between smooth manifolds.

1.2 Acknowledgements

First of all, I would like to express my gratitude to my supervisor Denis-Charles Cisinski, who has encouraged me to work on this subject and has accompanied me during the whole process of writing the paper, providing deep insights and help whenever I felt lost. Secondly, I want to thank Andrea Gagna and Edoardo Lanari for being patient and kind enough to answer all my (sometimes meaningless) questions over the past two years. I would also like to thank Denis Nardin, Peter Haine, George Raptis, Benedikt Preis, Marc Hoyois, Clark Barwick, Guglielmo Nocera, Ivan Di Liberti and Nicola Di Vittorio for showing interest in my work, for offering their support and for pushing me to keep learning new mathematics. This paper will be a part of my PhD thesis.

2 Sheaves and tensor products

The goal of this section will be twofold: first we are going to introduce Lurie's tensor product of cocomplete ∞ -categories as defined in [Lur17], and secondly we will recall the definition of sheaves and cosheaves with values in general ∞ -categories. The reason why we want to spend some time discussing this matter, aside from it being interesting on its own, is that in the following sections this tensor product will prove to be an extremely convenient tool to describe some categories of sheaves and functors between them: through it we will be able to produce a vast class of *essential geometric morphisms*, we will extend easily some results regarding sheaves of spaces to sheaves with values in any presentable ∞ -category, and later prove the existence of a sheafification functor when the ∞ -category of coefficients is stable and bicomplete with no presentability assumption, and construct the full six-functor formalism in this setting. Most of the results in this section are not at all original and can be found for example in [Cis19], in [Lur17], or are already well known.

2.1 Tensor product of cocomplete ∞ -categories

For the whole section we will fix two universes \mathbf{V} , \mathbf{U} such that \mathbf{V} is \mathbf{U} -small, and denote by \mathbf{Cocont}_∞ the ∞ -category of \mathbf{U} -small ∞ -categories admitting \mathbf{V} -small colimits, with \mathbf{V} -cocontinuous functors between them. For short, we will call an object of \mathbf{Cocont}_∞ a cocomplete ∞ -category and, for any two cocomplete ∞ -categories \mathcal{C} and \mathcal{D} we will denote by $\mathbf{Fun}_!(\mathcal{C}, \mathcal{D})$ the ∞ -category of functors preserving \mathbf{V} -small colimits.

Let \mathcal{C} and \mathcal{D} be cocomplete. Recall that, by [Lur09, 5.3.6], there exists a cocomplete ∞ -category, denoted by $\mathcal{C} \otimes \mathcal{D}$, and a functor

$$\boxtimes : \mathcal{C} \times \mathcal{D} \rightarrow \mathcal{C} \otimes \mathcal{D},$$

which preserves colimits in both variables and such that precomposing with \boxtimes gives an equivalence

$$(2.1) \quad \mathbf{Fun}_{! \times !}(\mathcal{C} \times \mathcal{D}, \mathcal{E}) \simeq \mathbf{Fun}_!(\mathcal{C} \otimes \mathcal{D}, \mathcal{E})$$

functorial on \mathcal{C} , \mathcal{D} and \mathcal{E} cocomplete, where $\mathbf{Fun}_{! \times !}$ indicates the ∞ -category of bifunctors preserving \mathbf{V} -small colimits in each variable. More precisely, [Lur17, Corollary 4.8.1.4] shows that this operation provides \mathbf{Cocont}_∞ with the structure of a symmetric monoidal ∞ -category, and the inclusion of \mathbf{Cocont}_∞ in \mathbf{Cat}_∞ is lax monoidal, where the latter is equipped with the cartesian monoidal structure. Since we obviously have a functorial equivalence

$$\mathbf{Fun}_{! \times !}(\mathcal{C} \times \mathcal{D}, \mathcal{E}) \simeq \mathbf{Fun}_!(\mathcal{C}, \mathbf{Fun}_!(\mathcal{D}, \mathcal{E})),$$

this monoidal structure is closed.

Remark 2.2. As usual, one may regard \mathbf{Cocont}_∞ as a $(\infty, 2)$ -category. It follows by (2.1) that, for any cocomplete ∞ -category \mathcal{C} , tensoring with \mathcal{C} actually gives rise to a 2-functor. An important consequence of this observation is that tensoring with \mathcal{C} preserves adjunctions of cocontinuous functors, since any adjunction is characterized by the classical triangular identities (see for example [Cis19, Theorem 6.1.23, (v)]).

We will now present a list of results about the tensor product of cocomplete ∞ -categories that will turn out to be very useful later.

Let A be a small ∞ -category, \mathcal{C} cocomplete. For any two objects $a \in A$ and $c \in \mathcal{C}$, denote by $a \boxtimes c = a_! c$ the left Kan extension of c along a (here we are considering a and c as functors $A^{\text{op}} \leftarrow \Delta^0 \rightarrow \mathcal{C}$). Thus we get a functor

$$\begin{aligned} A \times \mathcal{C} &\xrightarrow{y_{A/\mathcal{C}}} \mathbf{Fun}(A^{\text{op}}, \mathcal{C}) \\ (a, c) &\longmapsto a \boxtimes c \end{aligned}$$

which preserves colimits on the \mathcal{C} variable that we will call the *relative Yoneda embedding*. By definition, we have a functorial equivalence

$$\mathbf{Hom}(a \boxtimes c, F) \simeq \mathbf{Hom}(c, F(a))$$

for any $F \in \mathbf{Fun}(A^{\text{op}}, \mathcal{C})$.

Remark 2.3. Recall that in a closed symmetric monoidal ∞ -category, an object x is *strongly dualizable* if and only if the canonical map

$$(2.4) \quad y \otimes \underline{\mathbf{Hom}}(x, 1) \rightarrow \underline{\mathbf{Hom}}(x, y)$$

obtained as adjoint to

$$y \otimes \underline{\mathbf{Hom}}(x, 1) \otimes x \rightarrow y \otimes 1 \xrightarrow{\simeq} y$$

is an equivalence. In the case of \mathbf{Cocont}_∞ , one sees easily that the map (2.4) can be described as induced by

$$\mathcal{D} \times \mathbf{Fun}_!(\mathcal{C}, \mathcal{S}) \simeq \mathbf{Fun}_!(\mathcal{S}, \mathcal{D}) \times \mathbf{Fun}_!(\mathcal{C}, \mathcal{S}) \rightarrow \mathbf{Fun}_!(\mathcal{C}, \mathcal{D})$$

where the last functor is given by composition.

Remark 2.5. By the naturality on x of the map (2.4), we see that the full subcategory spanned by strongly dualizable objects is closed under retracts.

Remark 2.6. Consider the variablewise cocontinuous functor

$$(2.7) \quad \mathbf{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathcal{C} \rightarrow \mathbf{Fun}(A^{\text{op}}, \mathcal{C})$$

obtained as the extension by colimits of the relative Yoneda embedding, and denote by $F \boxtimes c$ the image of a pair $(F, c) \in \mathbf{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathcal{C}$. This induces a cocontinuous functor

$$(2.8) \quad \mathbf{Fun}(A^{\text{op}}, \mathcal{S}) \otimes \mathcal{C} \rightarrow \mathbf{Fun}(A^{\text{op}}, \mathcal{C}).$$

By definition one has identifications

$$\mathbf{Hom}(F \boxtimes c, G) \simeq \mathbf{Hom}(F, \mathbf{Hom}_e(c, G(-)))$$

functorially on $F \in \mathbf{Fun}(A^{\text{op}}, \mathcal{S})$, $G \in \mathbf{Fun}(A^{\text{op}}, \mathcal{C})$ and $c \in \mathcal{C}$, where the hom-space on the right-hand side is taken on the ∞ -category of presheaves of \mathbf{U} -small spaces.

A very convenient way to model the functor (2.7) is as follows. Let $y : \Delta^0 \hookrightarrow \mathcal{S}$ be the Yoneda embedding. Copmbining the fact that y is fully faithful and [Cis19, Proposition 6.4.12], we have

$$y_!c \circ a_!y \simeq a_!(y_!c \circ y) \simeq a_!c$$

and hence we get a commutative triangle

$$\begin{array}{ccc} \mathbf{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathcal{C} & \xrightarrow{\boxtimes} & \mathbf{Fun}(A^{\text{op}}, \mathcal{C}) \\ \downarrow \text{id} \times y_! & \nearrow \circ & \\ \mathbf{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathbf{Fun}_!(\mathcal{S}, \mathcal{C}) & & \end{array}$$

where the vertical arrow is an equivalence and the diagonal one is given by composition. In particular, one deduces that the functor (2.8) can be seen as an instance of (2.4).

For any cocomplete ∞ -category \mathcal{D} , precomposition with $y_{A/c}$ induces a functor

$$\mathbf{Fun}(\mathbf{Fun}(A^{\text{op}}, \mathcal{C}), \mathcal{D}) \rightarrow \mathbf{Fun}(A \times \mathcal{C}, \mathcal{D}) \simeq \mathbf{Fun}(\mathcal{C}, \mathbf{Fun}(A, \mathcal{D}))$$

and since colimits are computed pointwise in functor categories, it restricts to

$$(2.9) \quad \mathbf{Fun}_!(\mathbf{Fun}(A^{\text{op}}, \mathcal{C}), \mathcal{D}) \rightarrow \mathbf{Fun}_!(\mathcal{C}, \mathbf{Fun}(A, \mathcal{D})).$$

Theorem 2.10. The functor (2.9) is an equivalence. In particular, $\mathbf{Fun}(A^{\text{op}}, \mathcal{S})$ is strongly dualizable in the monoidal ∞ -category \mathbf{Cocont}_∞ with dual $\mathbf{Fun}(A, \mathcal{S})$, and thus the functor (2.8) is an equivalence.

Proof. A complete proof of the first statement can be found in [Cis19, 6.7]. The main ingredient of the proof is that, by [Cis19, Lemma 6.7.7], any $F \in \text{Fun}(A^{\text{op}}, \mathcal{C})$ can be written canonically as

$$F = \varinjlim_{c \rightarrow F(a)} a \boxtimes c$$

where the colimit is indexed by the Grothendieck construction of the functor $(a, c) \mapsto \text{Hom}_{\mathcal{C}}(c, F(a))$. Furthermore, even though this indexing category is not small a priori, [Cis19, Lemma 6.7.5] proves that it is finally small. From this one may deduce easily the theorem, in a similar spirit to how one proves that $\text{Fun}(A^{\text{op}}, \mathcal{S})$ is the free cocompletion under small colimits of A .

To prove the last statement, we just observe that we have canonical equivalences

$$\begin{aligned} \text{Fun}_!(\text{Fun}(A^{\text{op}}, \mathcal{C}), \mathcal{D}) &\simeq \text{Fun}_!(\mathcal{C}, \text{Fun}(A, \mathcal{D})) \\ &\simeq \text{Fun}_!(\mathcal{C}, \text{Fun}_!(\text{Fun}(A^{\text{op}}, \mathcal{S}), \mathcal{D})) \\ &\simeq \text{Fun}_!(\text{Fun}(A^{\text{op}}, \mathcal{S}) \otimes \mathcal{C}, \mathcal{D}) \end{aligned}$$

whose composition is given by precomposing with (2.8), and so we may conclude by Remark 2.6. \square

Corollary 2.11. Let $u : A \rightarrow B$ be a functor between small ∞ -categories, and let \mathcal{C} be any cocomplete ∞ -category. Then we have equivalences $u_! \otimes \mathcal{C} \simeq u_!$ and $u^* \otimes \mathcal{C} \simeq u^*$. Here by an abuse of notation we write $u_!$ (u^*) to indicate both left Kan extension (restriction) along u for functors with values in \mathcal{S} and in \mathcal{C} .

Proof. By Remark 2.2 and Theorem 2.10, we have an adjunction $u_! \otimes \mathcal{C} \dashv u^* \otimes \mathcal{C}$ of cocontinuous functors between \mathcal{C} -valued presheaves. By uniqueness of adjoints, it suffices to show that $u^* \otimes \mathcal{C} \simeq u^*$. But this is clear because by Remark 2.6 we have a commutative square

$$\begin{array}{ccc} \text{Fun}(B^{\text{op}}, \mathcal{S}) \times \mathcal{C} & \xrightarrow{u^* \times \text{id}} & \text{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathcal{C} \\ \downarrow \boxtimes & & \downarrow \boxtimes \\ \text{Fun}(B^{\text{op}}, \mathcal{C}) & \xrightarrow{u^*} & \text{Fun}(A^{\text{op}}, \mathcal{C}). \end{array}$$

\square

Denote by $\text{Cocont}_{\infty}^{\text{pt}}$ ($\text{Cocont}_{\infty}^{\text{st}}$) the full subcategory of Cocont_{∞} spanned by pointed (stable) cocomplete ∞ -categories.

Lemma 2.12. Let \mathcal{C} be any pointed (respectively stable) cocomplete ∞ -category, and let \mathcal{D} be cocomplete. Then $\mathcal{C} \otimes \mathcal{D}$ is pointed (respectively stable). In particular, $\text{Cocont}_{\infty}^{\text{pt}}$ (respectively $\text{Cocont}_{\infty}^{\text{st}}$) inherits an obvious monoidal structure from Cocont_{∞} and the inclusion in Cocont_{∞} admits a left adjoint given by tensoring with \mathcal{S}_* (respectively \mathcal{S}_{pt}).

Proof. First of all, notice that $\Delta^0 \otimes \mathcal{D} \simeq \Delta^0$. Since \mathcal{C} is pointed, the zero object $\Delta^0 \rightarrow \mathcal{C}$ is simultaneously a right and a left adjoint of the unique functor $\mathcal{C} \rightarrow \Delta^0$, and thus one may tensor these two adjunctions with \mathcal{D} and obtain by Corollary 2.11 that $\mathcal{C} \otimes \mathcal{D}$ is pointed.

Assume now that \mathcal{C} is stable. Since $\mathcal{C} \otimes \mathcal{D}$ is pointed, one sees easily that the suspension functor for $\mathcal{C} \otimes \mathcal{D}$ is obtained by applying $- \otimes \mathcal{D}$ to the suspension of \mathcal{C} , and so it is in particular an equivalence.

To prove that last part of the statement, it suffices to show that, for any pointed (stable) and cocomplete ∞ -category \mathcal{C} , the evaluation at S^0 (respectively \mathbb{S}) induces an equivalence $\text{Fun}_!(\mathcal{S}_*, \mathcal{C}) \simeq \mathcal{C}$ (respectively $\text{Fun}_!(\mathbb{S}, \mathcal{C}) \simeq \mathcal{C}$), but this follows easily by noticing that $\mathcal{S}_* \simeq$

$\text{Ind}(\mathcal{S}_*^{fin})$ (respectively $\text{Sp} \simeq \text{Ind}(\text{Sp}^{fin})$) and that evaluation at S^0 (respectively \mathcal{S}) induces an equivalence between finitely cocontinuous functors from \mathcal{S}_* (respectively Sp^{fin}) to \mathcal{C} and \mathcal{C} . \square

Remark 2.13. Let A be any small category and \mathcal{C} any object of $\text{Cocont}_\infty^{st}$. By the previous lemma, the functor (2.7) factors as

$$\begin{array}{ccc} \text{Fun}(A^{\text{op}}, \mathcal{S}) \times \mathcal{C} & \xrightarrow{\boxtimes} & \text{Fun}(A^{\text{op}}, \mathcal{C}) \\ \downarrow \Sigma_+^\infty \times \mathcal{C} & \nearrow \boxtimes^{st} & \\ \text{Fun}(A^{\text{op}}, \text{Sp}) \times \mathcal{C} & & \end{array}$$

inducing an equivalence

$$\text{Fun}(A^{\text{op}}, \mathcal{C}) \simeq \text{Fun}(A^{\text{op}}, \text{Sp}) \otimes \mathcal{C}.$$

Moreover, Remark 2.6 implies that one has identifications

$$\text{Hom}(F \boxtimes^{st} c, G) \simeq \text{Hom}(F, \underline{\text{Hom}}_c(c, G(-)))$$

functorially on $F \in \text{Fun}(A^{\text{op}}, \text{Sp})$, $G \in \text{Fun}(A^{\text{op}}, \mathcal{C})$ and $c \in \mathcal{C}$, where $\underline{\text{Hom}}_c(c, -)$ denotes the canonical enrichment of \mathcal{C} in \mathbf{U} -small spectra, and we have a commutative triangle

$$\begin{array}{ccc} \text{Fun}(A^{\text{op}}, \text{Sp}) \times \mathcal{C} & \xrightarrow{\boxtimes^{st}} & \text{Fun}(A^{\text{op}}, \mathcal{C}) \\ \downarrow \text{id} \times y! & \nearrow \circ & \\ \text{Fun}(A^{\text{op}}, \text{Sp}) \times \text{Fun}_!(\text{Sp}, \mathcal{C}) & & \end{array}$$

Proposition 2.14. Let \mathcal{C} be a compactly generated stable ∞ -category. Then \mathcal{C} is a strongly dualizable object of $\text{Cocont}_\infty^{st}$.

Proof. Since \mathcal{C} is stable and compactly generated, it follows that there exists a small stable ∞ -category A with finite colimits such that $\mathcal{C} \simeq \text{Fun}_{ex}(A^{\text{op}}, \text{Sp})$. Thus, since $\text{Fun}_!(\mathcal{C}, \text{Sp}) \simeq \text{Fun}_{ex}(A, \text{Sp})$, to prove the proposition we have to show that, for any \mathcal{D} stable and cocomplete, the canonical functor

$$\text{Fun}_{ex}(A^{\text{op}}, \text{Sp}) \otimes \mathcal{D} \rightarrow \text{Fun}_{ex}(A^{\text{op}}, \mathcal{D})$$

is an equivalence. We first prove that the inclusion $i : \text{Fun}_{ex}(A^{\text{op}}, \mathcal{D}) \hookrightarrow \text{Fun}(A^{\text{op}}, \mathcal{D})$ admits a left adjoint L .

For any $a \in A$, denote by $y^{st}(a)$ the spectrally enriched representable functor associated to a , obtained as usual through the equivalence

$$(2.15) \quad \text{Fun}_{ex}(A^{\text{op}}, \text{Sp}) \simeq \text{Fun}_{lex}(A^{\text{op}}, \mathcal{S}).$$

We define L as the unique (up to a contractible space of choices) cocontinuous functor extending

$$\begin{aligned} A \times \mathcal{D} &\longrightarrow \text{Fun}_{ex}(A^{\text{op}}, \mathcal{D}) \\ (a, x) &\longmapsto y^{st}(a) \boxtimes^{st} x. \end{aligned}$$

Indeed, $y^{st}(a) \boxtimes^{st} x$ is exact as it can be modelled by the composition of two finite colimit preserving functors. When $\mathcal{D} = \text{Sp}$, by (2.15) and the Yoneda lemma, one sees that L is left adjoint to i .

Let \mathcal{D} be any stable cocomplete ∞ -category. To see that L is the desired left adjoint, we observe that for any $F \in \text{Fun}_{ex}(A^{\text{op}}, \mathcal{D})$, $x \in \mathcal{D}$, we have functorial identifications

$$\begin{aligned} \text{Hom}(y^{st}(a) \boxtimes^{st} x, F) &\simeq \text{Hom}(y^{st}(a), \underline{\text{Hom}}_{\mathcal{D}}(x, F(-))) \\ &\simeq \text{Hom}(y(a), \text{Hom}_{\mathcal{D}}(x, F(-))) \\ &\simeq \text{Hom}(a \boxtimes x, F) \end{aligned}$$

where the hom-space on the right-hand side is taken on the ∞ -category of presheaves of \mathbf{U} -small spectra on A , and the second equivalence follows by the fact that F , and hence $\underline{\text{Hom}}_{\mathcal{D}}(x, F(-))$, is exact.

Now notice that $i : \text{Fun}_{ex}(A^{\text{op}}, \mathcal{S}\mathfrak{p}) \hookrightarrow \text{Fun}(A^{\text{op}}, \mathcal{S}\mathfrak{p})$ preserves colimits, and so, by tensoring with \mathcal{D} , one obtains an adjunction between cocontinuous functors

$$\begin{array}{ccc} & L \otimes \mathcal{D} & \\ \text{Fun}(A^{\text{op}}, \mathcal{D}) & \xrightarrow{\quad} & \text{Fun}_{ex}(A^{\text{op}}, \mathcal{S}\mathfrak{p}) \otimes \mathcal{D} \\ & \xleftarrow{\quad i \otimes \mathcal{D} \quad} & \end{array}$$

where $i \otimes \mathcal{D}$ is fully faithful. Since $\text{Fun}_{ex}(A^{\text{op}}, \mathcal{S}\mathfrak{p}) \otimes \mathcal{D}$ and $\text{Fun}_{ex}(A^{\text{op}}, \mathcal{D})$ can be respectively identified with the essential images of $(Li) \otimes \mathcal{D}$ and Li , to conclude the proof it suffices to show that the two functors are naturally equivalent, but this is true because they coincide on objects of the type $a \boxtimes x$. \square

Recall that an ∞ -category \mathcal{C} is called *\mathbf{V} -presentable* (for short, when there is no possibility of confusion we will only write *presentable*) if there exists a \mathbf{V} -small ∞ -category A such that \mathcal{C} is a left Bousfield localization of $\text{Fun}(A^{\text{op}}, \mathcal{S})$ by a \mathbf{V} -small set of morphism in $\text{Fun}(A^{\text{op}}, \mathcal{S})$. If we furthermore assume that the localization functor $\text{Fun}(A^{\text{op}}, \mathcal{S}) \rightarrow \mathcal{C}$ is left exact, we will say that \mathcal{C} is an *∞ -topos*. It follows easily by this definition that any presentable ∞ -category is complete and cocomplete. Presentable categories are equivalently defined as follows. Recall that, for \mathcal{C} any ∞ -category and S a class of morphisms in \mathcal{C} , we define an object $X \in \mathcal{C}$ to be *S -local* if, for every morphism $f : A \rightarrow B$ in S , the induced morphism

$$\text{Hom}_{\mathcal{C}}(B, X) \rightarrow \text{Hom}_{\mathcal{C}}(A, X)$$

is invertible. Then we say that an ∞ -category \mathcal{C} is *\mathbf{V} -presentable* if there exists a \mathbf{V} -small class S of morphisms in $\text{Fun}(A^{\text{op}}, \mathcal{S})$ such that \mathcal{C} is equivalent to the full subcategory of $\text{Fun}(A^{\text{op}}, \mathcal{S})$ spanned by S -local objects.

We denote by Pr_L the full subcategory of Cocont_{∞} spanned by presentable ∞ -categories and $\text{Pr}_R = \text{Pr}_L^{\text{op}}$. Notice that, by the adjoint functor theorem (see for example [Cis19, Proposition 7.11.8]), the morphisms in Pr_L are functors which admit a right adjoint and, consequently, morphisms in Pr_R are functors which admit a left adjoint. We also denote by Top the non full subcategory of Pr_R whose objects are ∞ -topoi and morphisms are functors which admit a left exact left adjoint (such functors are called *geometric morphisms*).

Proposition 2.16. Let \mathcal{C} and \mathcal{D} be two presentable ∞ -categories. Then $\mathcal{C} \otimes \mathcal{D}$ is presentable and there a canonical equivalence $\mathcal{C} \otimes \mathcal{D} \simeq \text{RFun}(\mathcal{C}^{\text{op}}, \mathcal{D})$. In particular, Pr_L inherits a symmetric monoidal structure.

Proof. Let A and B be two small ∞ -categories, S and S' two small sets of morphisms of $\text{Fun}(A^{\text{op}}, \mathcal{S})$ and $\text{Fun}(B^{\text{op}}, \mathcal{S})$ respectively such that \mathcal{C} and \mathcal{D} are equivalent the full subcategories of S and S' -local objects. By Theorem 2.10 we have

$$\begin{aligned} \text{Fun}(A^{\text{op}}, \mathcal{S}) \otimes \text{Fun}(B^{\text{op}}, \mathcal{S}) &\simeq \text{Fun}(A^{\text{op}}, \text{Fun}(B^{\text{op}}, \mathcal{S})) \\ &\simeq \text{Fun}((A \times B)^{\text{op}}, \mathcal{S}). \end{aligned}$$

It then follows from the proof of [Lur17, Proposition 4.8.1.15] that $\mathcal{C} \otimes \mathcal{D}$ can be identified with the full subcategory of $\text{Fun}((A \times B)^{\text{op}}, \mathcal{S})$ spanned by $S \otimes S'$ -local objects, where $S \otimes S'$ is the image of $S \times S'$ through the canonical functor

$$\text{Fun}(A^{\text{op}}, \mathcal{S}) \times \text{Fun}(B^{\text{op}}, \mathcal{S}) \rightarrow \text{Fun}((A \times B)^{\text{op}}, \mathcal{S}).$$

The proof of the last assertion follows by [Lur17, Lemma 4.8.1.16] and [Lur17, Proposition 4.8.1.17]. \square

Remark 2.17. Let \mathcal{C} be a presentable ∞ -category. One can deduce easily for Proposition 2.16 identifications $\mathcal{C} \otimes \mathcal{S}_* \simeq \mathcal{C}_*$ and $\mathcal{C} \otimes \text{Sp} \simeq \text{Sp}(\mathcal{C})$, where \mathcal{C}_* denotes the ∞ -category of *pointed objects* of \mathcal{C} , and $\text{Sp}(\mathcal{C})$ denotes the ∞ -category of *spectrum objects* of \mathcal{C} , i.e. the limit of the tower

$$\dots \xrightarrow{\Omega} \mathcal{C}_* \xrightarrow{\Omega} \mathcal{C}_*$$

where Ω is the usual loop functor. Both these constructions come with canonical functors $\mathcal{C}_* \rightarrow \mathcal{C}$ and $\Omega^\infty : \text{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$, and since \mathcal{C} is presentable one can show that these admit left adjoints $(-)_+ : \mathcal{C} \rightarrow \mathcal{C}_*$ and $\Sigma_+^\infty : \mathcal{C} \rightarrow \text{Sp}(\mathcal{C})$. By construction, we have a factorization

$$\mathcal{C} \xrightarrow{(-)_+} \mathcal{C}_* \xrightarrow{\Sigma_+^\infty} \text{Sp}(\mathcal{C}).$$

In particular we see that, if \mathcal{C} is presentable and pointed (stable), by tensoring $(-)_+ : \mathcal{S} \rightarrow \mathcal{S}_*$ ($\Sigma_+^\infty : \mathcal{S} \rightarrow \text{Sp}$) with \mathcal{C} we obtain an equivalence $\mathcal{C} \otimes \mathcal{S}_* \simeq \mathcal{C}$ ($\mathcal{C} \otimes \text{Sp} \simeq \text{Sp}(\mathcal{C})$). Thus, if \mathcal{C} , \mathcal{D} and \mathcal{E} are presentable ∞ -categories where \mathcal{D} is pointed and \mathcal{E} is stable, we get functorial identifications

$$\begin{aligned} \text{Fun}_!(\mathcal{C} \otimes \mathcal{S}_*, \mathcal{D}) &\simeq \text{Fun}_!(\mathcal{C}, \text{Fun}_!(\mathcal{S}_*, \mathcal{D})) \simeq \text{Fun}_!(\mathcal{C}, \mathcal{D}) \\ \text{Fun}_!(\mathcal{C} \otimes \text{Sp}, \mathcal{E}) &\simeq \text{Fun}_!(\mathcal{C}, \text{Fun}_!(\text{Sp}, \mathcal{E})) \simeq \text{Fun}_!(\mathcal{C}, \mathcal{E}) \end{aligned}$$

induced respectively by precomposing with $(-)_+$ and Σ_+^∞ . Furthermore, we see that $(-)_+ : \mathcal{S} \rightarrow \mathcal{S}_*$ and $\Sigma_+^\infty : \mathcal{S} \rightarrow \text{Sp}$ make \mathcal{S}_* and Sp into idempotent cocomplete ∞ -categories with respect to Lurie's tensor product: by [Lur17, Proposition 4.8.2.9] this implies that there are canonical variablewise cocontinuous symmetric monoidal structures on \mathcal{S}_* and Sp with unit objects given by $S^0 := (*)_+$ and $\mathbb{S} := \Sigma_+^\infty(*)$, and one can show that these coincide with the usual smash products of pointed spaces and spectra. In particular, we see that the functors

$$\mathcal{S} \xrightarrow{(-)_+} \mathcal{S}_* \xrightarrow{\Sigma_+^\infty} \text{Sp}$$

are all monoidal, where \mathcal{S} is equipped with the cartesian monoidal structure.

2.2 Sheaves and cosheaves

We now pass to recalling the definition of sheaves with values in an ∞ -category. Let X be a small ∞ -category equipped with a Grothendieck topology. Recall that there is a small ∞ -category $\text{Cov}(X)$, as defined in [Lur09, Notation 6.2.2.8], which can be described informally as having for objects pairs (x, R) , where $x \in X$ and $R \hookrightarrow y(x)$ is a sieve covering x , and morphisms between (x, R) and (y, R') are just maps $f : x \rightarrow y$ in X such that the restriction of $y(f)$ to R factors through R' . There is an obvious projection $\rho : \text{Cov}(X) \rightarrow X$ which has a section $s : X \rightarrow \text{Cov}(X)$ defined on objects by sending x to $(x, y(x))$.

Definition 2.18. Let \mathcal{C} be a complete ∞ -category. With the same notations as above, we say that a functor $F \in \text{Fun}(X^{\text{op}}, \mathcal{C})$ is a *sheaf* if the unit morphism

$$\rho^* F \rightarrow s_* s^* \rho^* F \simeq s_* F$$

is an equivalence. Dually, for a cocomplete ∞ -category \mathcal{C} , we say that a functor $F \in \text{Fun}(X, \mathcal{C})$ is a *cosheaf* if the counit morphism

$$s_!F \simeq s_!s^*\rho^*F \rightarrow \rho^*F$$

is an equivalence. We denote by $\text{Shv}(X; \mathcal{C})$ ($\text{CoShv}(X; \mathcal{C})$) the full subcategory of $\text{Fun}(X^{\text{op}}, \mathcal{C})$ ($\text{Fun}(X, \mathcal{C})$) spanned by (co)sheaves. When \mathcal{C} is the ∞ -category of spaces \mathcal{S} , we will simply write $\text{Shv}(X)$.

Remark 2.19. More concretely, one can describe a sheaf as a functor F such that for any covering sieve $R \hookrightarrow y(x)$ the canonical morphism

$$F(x) \rightarrow \varprojlim_{y(x') \rightarrow R} F(x')$$

is an equivalence. Notice also that we clearly have an equivalence $\text{CoShv}(X; \mathcal{C}) \simeq \text{Shv}(X; \mathcal{C}^{\text{op}})^{\text{op}}$.

Remark 2.20. It is well known that, for any ∞ -site X , the category $\text{Shv}(X)$ is an ∞ -topos. Unlike the case of 1-topoi, it's still unclear whether any ∞ -topos is equivalent to $\text{Shv}(X)$ for some ∞ -site X (see [Rez19]).

We now give another description of categories of sheaves and cosheaves.

Lemma 2.21. Let X be an ∞ -site, \mathcal{C} be any cocomplete ∞ -category. Then the restriction along the functor $X \xrightarrow{y} \text{Fun}(X^{\text{op}}, \mathcal{S}) \xrightarrow{L} \text{Shv}(X)$ defines an equivalence

$$\text{CoShv}(X; \mathcal{C}) \simeq \text{Fun}_!(\text{Shv}(X), \mathcal{C}),$$

where y is the Yoneda embedding and L is the sheafification functor. Equivalently, a functor $X \rightarrow \mathcal{C}$ is a cosheaf if and only if its extension by colimits $\text{Fun}(X^{\text{op}}, \mathcal{S}) \rightarrow \mathcal{C}$ factors through L . Dually, for any complete ∞ -category \mathcal{C} , we have an equivalence

$$\text{Shv}(X; \mathcal{C}) \simeq \text{Fun}_*(\text{Shv}(X)^{\text{op}}, \mathcal{C}).$$

Proof. Since L commutes with colimits, by the universal property of localizations composition with L embeds $\text{Fun}_!(\text{Shv}(X), \mathcal{C})$ in $\text{Fun}_!(\text{Fun}(X^{\text{op}}, \mathcal{S}), \mathcal{C})$ as the full subcategory of functors sending covering sieves $R \hookrightarrow y(x)$ to equivalences in \mathcal{C} . On the other hand, a functor $F : X \rightarrow \mathcal{C}$ is a cosheaf precisely if there is an equivalence

$$\varprojlim_{y(x') \rightarrow R} F(x') \simeq F(x)$$

for any sieve R on $x \in X$, thus precisely if its extension by colimits $\text{Fun}(X^{\text{op}}, \mathcal{S}) \rightarrow \mathcal{C}$ lies in $\text{Fun}_!(\text{Shv}(X), \mathcal{C})$. \square

We provide a couple of examples of cosheaves.

Example 2.22. (i) Let $f : X \rightarrow Y$ be a continuous map between topological spaces. Recall that this induces a geometric morphism $f : \text{Shv}(X) \rightarrow \text{Shv}(Y)$, which amounts to an adjunction $f^* \dashv f_*$, where $f_* : \text{Shv}(X) \rightarrow \text{Shv}(Y)$ is defined by $\Gamma(U; f_*F) = \Gamma(f^{-1}(U); F)$ for any $U \subseteq Y$. By Lemma 2.21, one may characterize $f^* : \text{Shv}(Y) \rightarrow \text{Shv}(X)$ as the essentially unique $\text{Shv}(X)$ -valued cosheaf on Y with the property that $f^*(y(U)) = y(f^{-1}(U))$.

(ii) Let Top be the 1-category of topological spaces, and $\text{Kan} \hookrightarrow \text{sSet}$ be the full subcategory of all simplicial sets consisting of Kan complexes. Recall that there is a functor $\text{Top} \rightarrow \text{Kan}$ defined by assigning to each topological space X its *singular complex*, i.e. the simplicial set defined by $n \mapsto \text{Hom}_{\text{Top}}(\Delta^n, X)$, where Δ^n is the standard n -simplex. Recall also that, by [Cis19, Theorem 7.8.9], there is a functor $\text{sSet} \rightarrow \mathcal{S}$ which identifies \mathcal{S} as a localization of sSet at the class of weak homotopy equivalences. We define $\text{Sing} : \text{Top} \rightarrow \mathcal{S}$ as the composition of the two functors defined above. It is proven in [Lur17, A.3] that, for any topological space X , the restriction of Sing to $\mathcal{U}(X)$ is indeed a cosheaf: this may be regarded as a non-truncated version of the classical Seifert-Van Kampen theorem. Furthermore, one can also show that Sing is a *hypercomplete cosheaf* (see [Lur17, Lemma A.3.10]): this means that, as cocontinuous functor $\text{Shv}(X) \rightarrow \mathcal{S}$, Sing factors through the *hypercompletion* of $\text{Shv}(X)$.

Corollary 2.23. Let X be an ∞ -site, \mathcal{C} be any presentable ∞ -category. Then the inclusion $\text{CoShv}(X; \mathcal{C}) \hookrightarrow \text{Fun}(X, \mathcal{C})$ admits a right adjoint.

Proof. By [Lur09, Proposition 5.5.3.8] and the previous lemma, the ∞ -category $\text{CoShv}(X; \mathcal{C})$ is presentable. Thus, since $\text{CoShv}(X; \mathcal{C}) \hookrightarrow \text{Fun}(X, \mathcal{C})$ obviously preserves colimits, we may conclude by the adjoint functor theorem. \square

Corollary 2.24. Let X be an ∞ -site, \mathcal{C} be any presentable ∞ -category. Then we have an equivalence $\text{Shv}(X) \otimes \mathcal{C} \simeq \text{Shv}(X; \mathcal{C})$.

Proof. It follows by the adjoint functor theorem that $\text{Fun}_*(\text{Shv}(X)^{\text{op}}, \mathcal{C}) \simeq \text{RFun}(\text{Shv}(X)^{\text{op}}, \mathcal{C})$. Thus, by the previous lemma and by Proposition 2.16, we get the conclusion. \square

Construction 2.25. Let X and Y be two topological spaces. The functor

$$\begin{aligned} \mathcal{U}(X) \times \mathcal{U}(Y) &\longrightarrow \text{Shv}(X \times Y) \\ (U, V) &\longmapsto y(U \times V) \end{aligned}$$

extends by colimits to a functor

$$\text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{S}) \times \text{Fun}(\mathcal{U}(Y)^{\text{op}}, \mathcal{S}) \rightarrow \text{Shv}(X \times Y).$$

Since it clearly sends covering sieves to equivalences in both variables, we obtain a functor

$$\begin{aligned} \text{Shv}(X) \times \text{Shv}(Y) &\longrightarrow \text{Shv}(X \times Y) \\ (F, G) &\longmapsto F \boxtimes G. \end{aligned}$$

More generally, by Corollary 2.24, tensoring with two presentable ∞ -categories \mathcal{C} and \mathcal{D} gives

$$(2.26) \quad \text{Shv}(X; \mathcal{C}) \times \text{Shv}(Y; \mathcal{D}) \rightarrow \text{Shv}(X \times Y; \mathcal{C} \otimes \mathcal{D})$$

for which the image of a pair (F, G) in the domain will still be denoted as $F \boxtimes G$. Let $\Delta : X \rightarrow X \times X$ be the diagonal. By post composing with Δ^* we get a functor denoted by

$$(2.27) \quad \begin{aligned} \text{Shv}(X; \mathcal{C}) \times \text{Shv}(X; \mathcal{D}) &\longrightarrow \text{Shv}(X; \mathcal{C} \otimes \mathcal{D}) \\ (F, G) &\longmapsto F \otimes G := \Delta^*(F \boxtimes G). \end{aligned}$$

Suppose now that \mathcal{C} is equipped with a monoidal structure $\otimes_{\mathcal{C}}$ such that the functor $\otimes_{\mathcal{C}} : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ preserves colimits in each variable. Then we have a cocontinuous functor

$\mathrm{Shv}(X; \mathcal{C} \otimes \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C})$ and by composing with (2.27) we obtain an induced monoidal structure on $\mathrm{Shv}(X; \mathcal{C})$, which will still be denoted by $\otimes_{\mathcal{C}}$. It is straightforward to check that the functor

$$(2.28) \quad \otimes_{\mathcal{C}} : \mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

can be described as

$$(F, G) \mapsto L^{\mathcal{C}}(U \mapsto \Gamma(U; F) \otimes_{\mathcal{C}} \Gamma(U; G))$$

where $L^{\mathcal{C}}$ is the sheafification for \mathcal{C} -valued presheaves. In particular, we have that any $F \in \mathrm{Shv}(X; \mathcal{C})$ induces a colimit preserving functor

$$- \otimes_{\mathcal{C}} F : \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C}).$$

Since $\mathrm{Shv}(X; \mathcal{C})$ is presentable, this has a right adjoint denoted by

$$\underline{\mathrm{Hom}}_X(F, -) : \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C}).$$

This functor supplies $\mathrm{Shv}(X; \mathcal{C})$ with a self-enrichment and for this reason will be called *internal Hom sheaf* functor.

Remark 2.29. Let X and Y be two topological spaces, \mathcal{C} , \mathcal{D} and \mathcal{E} be presentable ∞ -categories, and let $\alpha : \mathrm{Shv}(X) \rightarrow \mathrm{Shv}(Y)$ and $\Phi : \mathcal{C} \otimes \mathcal{D} \rightarrow \mathcal{E}$ be cocontinuous functors. Notice that the functoriality in each variable of the tensor product of cocomplete ∞ -categories gives a commutative diagram

$$\begin{array}{ccccc} \mathrm{Shv}(X \times X; \mathcal{C} \otimes \mathcal{D}) & \xrightarrow{\Delta^*} & \mathrm{Shv}(X; \mathcal{C} \otimes \mathcal{D}) & \xrightarrow{\alpha \otimes (\mathcal{C} \otimes \mathcal{D})} & \mathrm{Shv}(X; \mathcal{C} \otimes \mathcal{D}) \\ \downarrow \Delta^* & & \downarrow \Delta^* & & \downarrow \mathrm{Shv}(Y) \otimes \Phi \\ \mathrm{Shv}(X; \mathcal{C} \otimes \mathcal{D}) & \xrightarrow{\mathrm{Shv}(X) \otimes \Phi} & \mathrm{Shv}(X; \mathcal{E}) & \xrightarrow{\alpha \otimes \mathcal{E}} & \mathrm{Shv}(Y; \mathcal{E}). \end{array}$$

In particular, the diagram above shows that whenever we prove a formula involving the functor (2.27) and operations on sheaves coming from some continuous map, then we may deduce immediately a corresponding formula for the functor (2.28).

We may now formulate the following proposition, which could be interpreted as a sort of Künneth formula (we will actually see later in Proposition 6.11 how one can deduce the Künneth formula from this).

Proposition 2.30. Let X and Y be topological spaces, \mathcal{C} and \mathcal{D} two presentable ∞ -categories, and assume that one of the two is locally compact. Then the functor (2.26) induces an equivalence

$$\mathrm{Shv}(X; \mathcal{C}) \otimes \mathrm{Shv}(Y; \mathcal{D}) \simeq \mathrm{Shv}(X \times Y; \mathcal{C} \otimes \mathcal{D}).$$

Moreover, let $f : X \rightarrow X'$ and $g : Y \rightarrow Y'$ be two continuous maps and assume that at least one among X' and Y' is locally compact. Then we have an equivalence

$$(f \times g)_{\mathcal{C} \otimes \mathcal{D}}^*(F \boxtimes G) \simeq f_{\mathcal{C}}^* F \boxtimes g_{\mathcal{D}}^* G$$

which is functorial on $F \in \mathrm{Shv}(X'; \mathcal{C})$ and $G \in \mathrm{Shv}(Y'; \mathcal{D})$.

Proof. By [Lur09, Proposition 7.3.3.9], for any topological space X and any ∞ -topos \mathcal{Y} , $\mathrm{Shv}(X; \mathcal{Y})$ is a product of $\mathrm{Shv}(X)$ and \mathcal{Y} in the ∞ -category Top . Thus, by the previous corollary combined with [Lur09, Proposition 7.3.1.11], if Y is a locally compact topological space, we have an equivalence

$$\mathrm{Shv}(X) \otimes \mathrm{Shv}(Y) \simeq \mathrm{Shv}(X \times Y).$$

For the second part of the statement, we first observe that by Corollary 2.24 and Proposition 2.16 it suffices to prove the case when $\mathcal{C} = \mathcal{D} = \mathcal{S}$, which amounts to providing a commutative square

$$\begin{array}{ccc} \mathrm{Shv}(X') \times \mathrm{Shv}(Y') & \longrightarrow & \mathrm{Shv}(X' \times Y') \\ \downarrow f^* \times g^* & & \downarrow (f \times g)^* \\ \mathrm{Shv}(X) \times \mathrm{Shv}(Y) & \longrightarrow & \mathrm{Shv}(X \times Y). \end{array}$$

Since both the top right and the down left composition commute with colimits in both variables, one then gets this by Lemma 2.21, Example 2.22 (i) and by observing that

$$(f \times g)^{-1}(U \times V) = f^{-1}(U) \times g^{-1}(V)$$

for any U and V open subsets of X' and Y' respectively. \square

Corollary 2.31 (Monoidality). Let $f : X \rightarrow Y$ be a morphism of locally compact topological spaces, \mathcal{C} and \mathcal{D} two presentable ∞ -categories. Then we have a canonical identification

$$f^*(F \otimes G) \simeq f^*F \otimes f^*G$$

and in particular, when $\mathcal{C} = \mathcal{D}$ is monoidal, by transposition

$$f_* \underline{\mathrm{Hom}}_X(f^*H, K) \simeq \underline{\mathrm{Hom}}_Y(H, f_*K).$$

Proof. The commutativity of the diagram

$$\begin{array}{ccc} \mathrm{Shv}(Y; \mathcal{C}) \times \mathrm{Shv}(Y; \mathcal{D}) & \xrightarrow{(f^*, f^*)} & \mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(X; \mathcal{D}) \\ \otimes \downarrow & & \downarrow \otimes \\ \mathrm{Shv}(Y; \mathcal{C} \otimes \mathcal{D}) & \xrightarrow{f^*} & \mathrm{Shv}(X; \mathcal{C} \otimes \mathcal{D}) \end{array}$$

follows from the commutativity of

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta \downarrow & & \downarrow \Delta \\ X \times X & \xrightarrow{(f, f)} & Y \times Y \end{array}$$

that is trivially verified. The last part follows directly from the previous lemma. \square

Consider now the ∞ -category $\mathrm{Shv}(X; \mathcal{C})$, where X is any ∞ -site and \mathcal{C} is complete and cocomplete. It is natural to ask oneself whether at this level of generality one is still able to obtain a result like Corollary 2.24, at least when the inclusion $\mathrm{Shv}(X; \mathcal{C}) \hookrightarrow \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$ admits a left adjoint. In the rest of the section we will briefly outline the reason why the answer to this question doesn't seem to be affirmative. We start with a general proposition concerning left Bousfield localizations and categories of local objects.

Proposition 2.32. Let \mathcal{C} be an ∞ -category and S a class of morphism in \mathcal{C} . Denote by \mathcal{C}_S the full subcategory of \mathcal{C} spanned by S -local objects, and assume that the inclusion $i : \mathcal{C}_S \hookrightarrow \mathcal{C}$ admits a left adjoint L . Thus, composition with L gives a fully faithful functor

$$(2.33) \quad \mathrm{LFun}(\mathcal{C}_S, \mathcal{D}) \hookrightarrow \mathrm{LFun}(\mathcal{C}, \mathcal{D})$$

whose essential image is given by left adjoints $\mathcal{C} \rightarrow \mathcal{D}$ sending all morphisms in S to equivalences.

Proof. Let W be the class of morphisms in \mathcal{C} which are sent by L to equivalences and denote by \mathcal{A} and \mathcal{A}' the full subcategories of $\mathrm{LFun}(\mathcal{C}, \mathcal{D})$ spanned respectively by left adjoints $\mathcal{C} \rightarrow \mathcal{D}$ sending all morphisms in W to equivalences and left adjoints $\mathcal{C} \rightarrow \mathcal{D}$ sending all morphisms in S to equivalences. By [Cis19, Proposition 7.1.18], we already know that (2.33) is fully faithful and that its essential image is given by \mathcal{A} . It follows immediately by the definition of a local object that L sends all morphisms in S to equivalences, thus we just need to show that \mathcal{A}' is contained in \mathcal{A} .

Consider a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ in \mathcal{A}' with right adjoint $G : \mathcal{D} \rightarrow \mathcal{C}$. By definition of \mathcal{A}' , we have that for every morphisms in $f \in S$ and every $d \in \mathcal{D}$

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(c, G(d)) & \xrightarrow{f} & \mathrm{Hom}_{\mathcal{C}}(c', G(d)) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathrm{Hom}_{\mathcal{D}}(F(c), d) & \xrightarrow{F(f)} & \mathrm{Hom}_{\mathcal{D}}(F(c'), d). \end{array}$$

Thus $G(d)$ is S -local, and hence there exists a functor $G' : \mathcal{D} \rightarrow \mathcal{C}_S$ such that $G = iG'$. Let now f be a morphism in W . By definition of W we have, functorially on $d \in \mathcal{D}$,

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{D}}(F(c), d) & \xrightarrow{F(f)} & \mathrm{Hom}_{\mathcal{D}}(F(c'), d) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathrm{Hom}_{\mathcal{C}}(c, iG'(d)) & \xrightarrow{f} & \mathrm{Hom}_{\mathcal{C}}(c', iG'(d)) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathrm{Hom}_{\mathcal{C}_S}(L(c), G'(d)) & \xrightarrow[\simeq]{L(f)} & \mathrm{Hom}_{\mathcal{C}_S}(L(c'), G'(d)), \end{array}$$

and hence $F(f)$ is invertible, and so we may conclude. \square

We now claim that there exists a class of morphisms S of $\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$ such that $\mathrm{Shv}(X; \mathcal{C})$ can be identified with the full subcategory of S -local objects of $\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$. We define S as the class of morphisms

$$S = \{R \boxtimes M \rightarrow x \boxtimes M \mid R \hookrightarrow y(x) \text{ is a sieve, } M \in \mathcal{C}\}.$$

For any sieve $R \hookrightarrow y(x)$, $M \in \mathcal{C}$ and $F \in \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$, since $R \simeq \varinjlim_{y(x') \rightarrow R} y(x')$, we have a commutative diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(-, F(x)) & \longrightarrow & \mathrm{Hom}_{\mathcal{C}}(-, \varinjlim_{y(x') \rightarrow R} F(x')) \\ \downarrow \simeq & & \downarrow \simeq \\ \mathrm{Hom}_{\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})}(x \boxtimes -, F) & \xrightarrow{i} & \mathrm{Hom}_{\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})}(R \boxtimes -, F) \end{array}$$

where the upper horizontal arrow is induced by the canonical map $F(x) \rightarrow \varprojlim_{y(x') \rightarrow R} F(x')$.

Thus, we see that the upper horizontal arrow is invertible if and only if the lower horizontal one is, and so F is a sheaf if and only if it is S -local. In particular, by the previous proposition, whenever the inclusion $\mathrm{Shv}(X; \mathcal{C}) \hookrightarrow \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$ admits a left adjoint, the ∞ -category $\mathrm{Shv}(X; \mathcal{C})$ is characterized by the universal property

$$\mathrm{LFun}(\mathrm{Shv}(X; \mathcal{C}), \mathcal{D}) \hookrightarrow \mathrm{LFun}_S(\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C}), \mathcal{D})$$

where the right-hand side denotes the ∞ -category of left adjoint functors sending all morphisms in S to equivalence. On the other hand, tensoring the usual sheafification $\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{S}) \rightarrow \mathrm{Shv}(X)$ with \mathcal{C} gives a colimit preserving functor

$$L' : \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C}) \simeq \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{S}) \otimes \mathcal{C} \rightarrow \mathrm{Shv}(X) \otimes \mathcal{C}.$$

Combining the universal property of the tensor product of cocomplete categories and Theorem 2.10, we see that, for any cocomplete ∞ -category \mathcal{D} , precomposition with L' may be factored as

$$\begin{aligned} \mathrm{Fun}_!(\mathrm{Shv}(X) \otimes \mathcal{C}, \mathcal{D}) &\simeq \mathrm{Fun}_!(\mathcal{C}, \mathrm{Fun}_!(\mathrm{Shv}(X), \mathcal{D})) \\ &\hookrightarrow \mathrm{Fun}_!(\mathcal{C}, \mathrm{Fun}_!(\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{S}), \mathcal{D})) \\ &\simeq \mathrm{Fun}_!(\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C}), \mathcal{D}) \end{aligned}$$

and hence identifies $\mathrm{Fun}_!(\mathrm{Shv}(X) \otimes \mathcal{C}, \mathcal{D})$ with the full subcategory of $\mathrm{Fun}_!(\mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C}), \mathcal{D})$ spanned by those functors sending maps in S to equivalences. Hence we obtain a comparison functor

$$\mathrm{Shv}(X) \otimes \mathcal{C} \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

but unless \mathcal{C} is presentable, there is no evident reason why one should expect this to be an equivalence.

Remark 2.34. A close inspection of the proof of [Lur09, Proposition 6.2.2.7] shows that the usual formula for sheafification provides the desired left adjoint whenever \mathcal{C} is bicomplete and, for every $x \in X$ and every sieve $R \hookrightarrow y(x)$, the functor

$$\begin{array}{ccc} \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C}) & \longrightarrow & \mathcal{C} \\ F & \longmapsto & (s_* F)(x, R) \simeq \varprojlim_{y(x') \rightarrow R} F(x') \end{array}$$

is accessible: this will be true automatically for example when \mathcal{C} is presentable, since any functor between presentable ∞ -categories which is a right adjoint is automatically accessible. A similar observation in the case of sheaves with values in ordinary 1-categories can be found in [KS06, 17.4]. However, if we drop the presentability assumption for \mathcal{C} , it is not clear a priori why the inclusion $\mathrm{Shv}(X; \mathcal{C}) \hookrightarrow \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$ should admit a left adjoint.

Remark 2.35. Suppose that \mathcal{C} is such that $\mathrm{Shv}(X; \mathcal{C}) \hookrightarrow \mathrm{Fun}(X^{\mathrm{op}}, \mathcal{C})$ admits a left adjoint L . Hence, for any $x \in X$ and $M \in \mathcal{C}$, by applying L to $x \boxtimes M$ gives an object denoted by M_x with the property that, for any other sheaf F , we have a functorial identification

$$\mathrm{Hom}(M_x, F) \simeq \mathrm{Hom}(M, F(x)).$$

It follows by [Cis19, Proposition 7.1.18] and Theorem 2.10 that, for any cocomplete ∞ -category \mathcal{D} , we have a fully faithful functor

$$\mathrm{Fun}_!(\mathrm{Shv}(X; \mathcal{C}), \mathcal{D}) \hookrightarrow \mathrm{Fun}_!(\mathcal{C}, \mathrm{Fun}(A, \mathcal{D}))$$

and thus any cocontinuous functor with domain $\mathrm{Shv}(X; \mathcal{C})$ is uniquely determined by its values on objects of the type M_x .

3 Shape theory and shape submersions

In this section we will deal with questions related to shape theory from the perspective of higher topos theory: we recommend [Lur17, Appendix A] and [Hoy18] for some good introductory accounts to this subject. We will start by defining a version of shape which is relative to a geometric morphism, and give a detailed description of its functoriality as well as a proof of its homotopy invariance. After that we will define essential and locally contractible geometric morphisms: the first notion refers to morphisms $f : \mathcal{X} \rightarrow \mathcal{Y}$ whose relative shape is constant (as a pro-object on \mathcal{Y}) locally on \mathcal{X} , while the second to essential geometric morphisms satisfying an additional push-pull formula. After that we will define shape submersions, i.e. continuous maps which are locally given by projections $X \times Y \rightarrow Y$, where X is such that the unique geometric morphism $\mathrm{Shv}(X) \rightarrow \mathcal{S}$ is essential. These are proven to satisfy a base change formula, which will imply that they induce locally contractible geometric morphisms.

3.1 Relative shape

For any ∞ -category \mathcal{C} , denote by $\mathrm{Pro}(\mathcal{C})$ the ∞ -category of pro-objects in \mathcal{C} , i.e. the free completion of \mathcal{C} under cofiltered limits. When \mathcal{C} is accessible and admits finite limits, one shows ([Lur09, Proposition 3.1.6]) that $\mathrm{Pro}(\mathcal{C})$ is in fact equivalent to the full subcategory of $\mathrm{Fun}(\mathcal{C}, \mathcal{S})^{\mathrm{op}}$ spanned by the left exact functors.

Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be an accessible functor between presentable ∞ -categories, and let $G : \mathcal{D} \rightarrow \mathrm{Fun}(\mathcal{C}, \mathcal{S})^{\mathrm{op}}$ be the composition of the Yoneda embedding $\mathcal{D} \rightarrow \mathrm{Fun}(\mathcal{C}, \mathcal{S})^{\mathrm{op}}$ with $F^* : \mathrm{Fun}(\mathcal{D}, \mathcal{S})^{\mathrm{op}} \rightarrow \mathrm{Fun}(\mathcal{C}, \mathcal{S})^{\mathrm{op}}$. By the adjoint functor theorem, G factors through \mathcal{C} if and only if F commutes with limits, and when this condition is verified G is a left adjoint to F . If F is only left exact, by the characterization stated above G factors through $\mathrm{Pro}(\mathcal{C})$: in this case, we say that G is the *left pro-adjoint* of F . Notice that, in this situation, G is a genuine left adjoint of the functor $\mathrm{Pro}(F) : \mathrm{Pro}(\mathcal{C}) \rightarrow \mathrm{Pro}(\mathcal{D})$.

Specializing to the case of a geometric morphism between ∞ -topoi $f : \mathcal{X} \rightarrow \mathcal{Y}$, we see that the pullback f^* admits a pro-left adjoint, that we will denote by $f_{\sharp} : \mathcal{X} \rightarrow \mathrm{Pro}(\mathcal{Y})$. More explicitly, for every object $U \in \mathcal{X}$, $f_{\sharp}(U)$ is the pro-object on \mathcal{Y} defined by the assignment

$$V \mapsto \mathrm{Hom}_{\mathcal{X}}(U, f^*(V)).$$

Definition 3.1. Let \mathcal{X} be an ∞ -topos, $f : \mathcal{Y} \rightarrow \mathcal{X}$ a geometric morphism. We define the *shape of \mathcal{Y} relative to \mathcal{X}* as

$$\Pi_{\infty}^{\mathcal{X}}(\mathcal{Y}) := f_{\sharp}1_{\mathcal{Y}},$$

where $1_{\mathcal{Y}}$ is a terminal object of \mathcal{Y} . We will say that f is *constant shape* if $\Pi_{\infty}^{\mathcal{X}}(\mathcal{Y})$ belongs to \mathcal{Y} . In the case where f is the unique geometric morphism $a : \mathcal{Y} \rightarrow \mathcal{S}$, $a_{\sharp}1_{\mathcal{Y}}$ will be denoted just by $\Pi_{\infty}(\mathcal{Y})$ and will be called the *shape* or *fundamental pro- ∞ -groupoid* of \mathcal{Y} .

Remark 3.2. Notice that, as a left exact functor $\mathcal{X} \rightarrow \mathcal{S}$, $f_{\sharp}1_{\mathcal{Y}}$ can be identified with the functor $a_*f_*f^*$, where $a : \mathcal{X} \rightarrow \mathcal{S}$ is the unique geometric morphism.

Proposition 3.3. There exists a functor

$$\Pi_{\infty}^{\mathcal{X}} : \mathrm{Top}/\mathcal{X} \rightarrow \mathrm{Pro}(\mathcal{X})$$

whose values on objects coincides with the shape relative to \mathcal{X} and whose values on morphisms

$$\begin{array}{ccc} \mathcal{Y} & \xrightarrow{g} & \mathcal{Y}' \\ f \searrow & & \swarrow f' \\ & \mathcal{X} & \end{array}$$

is given by the transformation

$$a_*f'_*f'^* \rightarrow a_*f'_*g_*g^*f'^* \simeq a_*f_*f^*$$

induced by the unit of the adjunction $g^* \dashv g_*$.

Proof. Since we clearly have a functor

$$\mathrm{Fun}^{\mathrm{lex}}(\mathcal{X}, \mathcal{X}) \rightarrow \mathrm{Fun}^{\mathrm{lex}}(\mathcal{X}, \mathcal{S}) \simeq \mathrm{Pro}(\mathcal{X})^{\mathrm{op}}$$

given by post composition with a_* , it suffices to prove that there is a functor

$$T : \Pi_{\infty}^{\mathcal{X}} : (\mathcal{T}\mathrm{op}_{/\mathcal{X}})^{\mathrm{op}} \rightarrow \mathrm{Fun}^{\mathrm{lex}}(\mathcal{X}, \mathcal{X})$$

that assigns f_*f^* to any $f : \mathcal{Y} \rightarrow \mathcal{X}$ and at the level of morphisms $\mathcal{Y} \begin{array}{c} \xrightarrow{g} \mathcal{Y}' \\ f \searrow \quad \swarrow f' \\ \mathcal{X} \end{array}$ is given by

the transformation

$$f'_*f'^* \rightarrow f'_*g_*g^*f'^* \simeq f_*f^*$$

induced by the unit of the adjunction $g^* \dashv g_*$. We will proceed through some reduction steps.

First of all, the Yoneda embedding induces a fully faithful functor

$$\mathrm{Fun}(\mathcal{X}, \mathcal{X}) \hookrightarrow \mathrm{Fun}(\mathcal{X}, \mathrm{Fun}(\mathcal{X}^{\mathrm{op}}, \mathcal{S})) \simeq \mathrm{Fun}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X}, \mathcal{S})$$

and thus it suffices to construct a functor

$$(\mathcal{T}\mathrm{op}_{/\mathcal{X}})^{\mathrm{op}} \rightarrow \mathrm{Fun}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X}, \mathcal{S})$$

whose image lies in $\mathrm{Fun}^{\mathrm{lex}}(\mathcal{X}, \mathcal{X})$. By [Cis19, Remark 6.1.5], standard computations with adjunctions of 1-categories show that we also have, functorially on $F, G \in \mathcal{X}$, a commutative square

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{Y}}(f^*F, f^*G) & \xrightarrow{\simeq} & \mathrm{Hom}_{\mathcal{X}}(F, f_*f^*G) \\ g^* \downarrow & & \downarrow \mathrm{unit} \\ \mathrm{Hom}_{\mathcal{Y}'}(g^*f^*F, g^*f^*G) & \xrightarrow{\simeq} & \mathrm{Hom}_{\mathcal{X}}(F, f_*g_*g^*f^*G), \end{array}$$

so it suffices to show that the transformation on the left hand side can be enhanced to a functor $(\mathcal{T}\mathrm{op}_{/\mathcal{X}})^{\mathrm{op}} \rightarrow \mathrm{Fun}(\mathcal{X} \times \mathcal{X}^{\mathrm{op}}, \mathcal{S})$. Recall also that we have a forgetful functor

$$(\mathcal{T}\mathrm{op}_{/\mathcal{X}})^{\mathrm{op}} \simeq (\mathcal{T}\mathrm{op}^{\mathrm{op}})_{\mathcal{X}/} \longrightarrow \mathrm{Cat}_{\infty\mathcal{X}/}$$

$$(f : \mathcal{Y} \rightarrow \mathcal{X}) \longmapsto (f^* : \mathcal{X} \rightarrow \mathcal{Y}).$$

Hence, we will construct a functor

$$\mathrm{Cat}_{\infty\mathcal{X}/} \rightarrow \mathrm{Fun}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X}, \mathcal{S}).$$

Let Cat_{∞} be the full subcategory of sSet spanned by ∞ -categories, and let $\mathrm{LFib}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X})$ be the full subcategory of $\mathrm{sSet}_{/\mathcal{X}^{\mathrm{op}} \times \mathcal{X}}$ spanned by the left fibrations. Since Cat_{∞} is the category of fibrant objects in the Joyal model structure on sSet , it follows by [Cis19, Theorem 7.5.18], [Cis19, Example 7.10.14] and [Cis19, Theorem 3.9.7] that one may regard Cat_{∞} as a localization of Cat_{∞} by the class W of fully faithful and essentially surjective functors. Thus, by [Cis19, Corollary 7.6.13] and [Cis19, Proposition 7.1.7], we get an equivalence $\mathrm{Cat}_{\infty\mathcal{X}/} \simeq \mathrm{Cat}_{\infty\mathcal{X}/}[W^{-1}]$. On the other hand, $\mathrm{LFib}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X})$ is the category of fibrant objects in the covariant model structure on $\mathrm{sSet}_{/\mathcal{X}^{\mathrm{op}} \times \mathcal{X}}$, and so by [Cis19, Theorem 7.5.18], [Cis19, Theorem 7.8.9] and [Cis19, Theorem 4.4.14] we may regard $\mathrm{Fun}(\mathcal{X}^{\mathrm{op}} \times \mathcal{X}, \mathcal{S})$ as the localization of

$\text{LFib}(\mathcal{X}^{\text{op}} \times \mathcal{X})$ by the class W' of fibrewise equivalences. Thus, to produce T it will suffice to provide a functor $\text{Cat}_{\infty\mathcal{X}/} \rightarrow \text{LFib}(\mathcal{X}^{\text{op}} \times \mathcal{X})$ of 1-categories which maps W into W' .

Recall that, for any ∞ -category \mathcal{C} , we have a left fibration $\mathcal{S}(\mathcal{C}) \xrightarrow{(s,t)} \mathcal{C}^{\text{op}} \times \mathcal{C}$, called the *twisted diagonal*, classifying the hom-bifunctor $\mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{S}$ (as it is defined in [Cis19, 5.6.1]). For any functor $f : \mathcal{X} \rightarrow \mathcal{Y}$, consider the left fibration defined by the pullback

$$\begin{array}{ccc} T(f) & \longrightarrow & \mathcal{S}(\mathcal{Y}) \\ \downarrow & \lrcorner & \downarrow (s,t) \\ \mathcal{X}^{\text{op}} \times \mathcal{X} & \xrightarrow{f^{\text{op}} \times f} & \mathcal{Y}^{\text{op}} \times \mathcal{Y} \end{array}$$

which classifies the functor

$$\text{Hom}_{\mathcal{X}}(f(-), f(-)) : \mathcal{X}^{\text{op}} \times \mathcal{X} \rightarrow \mathcal{S}.$$

The functoriality on \mathcal{Y} of the twisted diagonal and the universal property of pullbacks imply that T defines a functor $\text{Cat}_{\infty\mathcal{X}/} \rightarrow \text{LFib}(\mathcal{X}^{\text{op}} \times \mathcal{X})$, as illustrated by the diagram corresponding

to a morphism $\begin{array}{ccc} & \mathcal{X} & \\ f \swarrow & & \searrow f' \\ \mathcal{Y} & \xrightarrow{g} & \mathcal{Y}' \end{array}$

$$\begin{array}{ccccc} T(f) & \longrightarrow & \mathcal{S}(\mathcal{Y}) & & \\ \downarrow \text{dotted} & & \downarrow \text{dotted} & \searrow S(g) & \\ T(f') & \longrightarrow & T(g) & \longrightarrow & \mathcal{S}(\mathcal{Y}') \\ \downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow (s,t) \\ \mathcal{X}^{\text{op}} \times \mathcal{X} & \xrightarrow{f^{\text{op}} \times f} & \mathcal{Y}^{\text{op}} \times \mathcal{Y} & \xrightarrow{g^{\text{op}} \times g} & (\mathcal{Y}')^{\text{op}} \times \mathcal{Y}' \end{array}$$

Moreover, [Cis19, Corollary 5.6.6] implies that T sends any fully faithful functor to a fibrewise equivalence, and hence we can conclude. \square

Recall that, for two maps $Y \xrightarrow[f_1]{f_0} Y'$ over a topological space X , we say that f_0 is *homotopic to f_1 over X* if there exists a map $h : Y \times I \rightarrow Y'$ over X such that $f_t = hi_t$ $t = 0, 1$, where $i_t : Y \hookrightarrow Y \times I$ is the inclusion corresponding to $t \in I$.

Corollary 3.4 (Homotopy invariance). Let Y, Y' be two topological spaces over X , and let $Y \xrightarrow[f_1]{f_0} Y'$ be two homotopic maps over X . Then the functor T induces an equivalence $T(f_0) \simeq T(f_1)$. In particular, T sends homotopy equivalences over X to invertible morphisms in $\text{Fun}(\mathcal{X}, \mathcal{X})^{\text{op}}$.

Proof. Let $p : Y \times I \rightarrow Y$ be the canonical projection. By [Lur17, Lemma A.2.9], we know that p^* is fully faithful, and hence $T(p)$ is invertible. Since $pi_0 = pi_1 = \text{id}_Y$ and T is functorial, we get an equivalence $T(i_0) \simeq T(i_1)$. Thus, since there exists a homotopy h over X such that $f_t = hi_t$ $t = 0, 1$, the functoriality of T gives the desired $T(f_0) \simeq T(f_1)$. \square

Remark 3.5. Recall that, for any ∞ -topos \mathcal{X} , there is a fully faithful functor

$$(3.6) \quad \begin{array}{ccc} \mathcal{X} & \hookrightarrow & \mathcal{T}\text{op}/\mathcal{X} \\ x & \longmapsto & \mathcal{X}/_x. \end{array}$$

Since $\mathcal{T}\text{op}/_{\mathcal{X}}$ has small cofiltered limits, the latter can be extended to a functor

$$\beta : \text{Pro}(\mathcal{X}) \rightarrow \mathcal{T}\text{op}/_{\mathcal{X}}.$$

It is possible to construct the functor

$$\Pi_{\infty}^{\mathcal{X}} : \mathcal{T}\text{op}/_{\mathcal{X}} \rightarrow \text{Pro}(\mathcal{X})$$

directly by showing that there is an equivalence

$$\text{Hom}_{\text{Pro}(\mathcal{X})}(\Pi_{\infty}^{\mathcal{X}}(\mathcal{Y}), Z) \simeq \text{Hom}_{\mathcal{T}\text{op}/_{\mathcal{X}}}(\mathcal{Y}, \beta(Z))$$

which is functorial on $Z \in \text{Pro}(\mathcal{X})$, as it is done in [Lur16, Proposition E.2.2.1]. However, we preferred to prove directly the functoriality of the relative shape, namely because the approach mentioned above leaves unclear how the functor $\Pi_{\infty}^{\mathcal{X}}$ would behave at the level of morphisms, which is needed to have a proof of homotopy invariance as clean and immediate as the one above.

Remark 3.7. As usual, since $\mathcal{T}\text{op}$ has pullbacks, the slice $\mathcal{T}\text{op}/_{\mathcal{X}}$ can be equipped with a contravariantly functorial structure

$$\mathcal{T}\text{op}^{\text{op}} \rightarrow \text{Cat}_{\infty}$$

that can be described for any geometric morphism $g : \mathcal{X} \rightarrow \mathcal{Y}$ by sending an object $(f : \mathcal{Y}' \rightarrow \mathcal{Y}) \in \mathcal{T}\text{op}/_{\mathcal{Y}}$ to the resulting arrow over \mathcal{X} obtained by performing the pullback of f along g . Thus, since by [Lur09, Remark 6.3.5.8] we have for any $y \in \mathcal{Y}$ a canonical pullback square

$$\begin{array}{ccc} \mathcal{X}/_{f^*y} & \longrightarrow & \mathcal{Y}/_y \\ \downarrow & & \downarrow \\ \mathcal{X} & \xrightarrow{f} & \mathcal{Y} \end{array}$$

in $\mathcal{T}\text{op}$, we see that the functor (3.6) is actually natural in \mathcal{X} , where the left hand side is functorial by the usual forgetful $\mathcal{T}\text{op}^{\text{op}} \rightarrow \text{Cat}_{\infty}$ sending a geometric morphism f to f^* . In particular we obtain that

$$\beta : \text{Pro}(\mathcal{X}) \rightarrow \mathcal{T}\text{op}/_{\mathcal{X}}$$

is actually natural in \mathcal{X} . Notice that, if one regards Cat_{∞} as an $(\infty, 2)$ -category, the universal property of $\text{Pro}(\mathcal{X})$ and the definition of the slice imply that β can be seen as a natural transformation between 2-functors. Hence, by [Hau20, Theorem 3.22], by adjunction we may regard the relative shape as a lax natural transformation, where the 2-cells involved may be described as follows: any geometric morphism induces an adjunction

$$\begin{array}{ccc} & \xrightarrow{g_{\#}} & \\ \text{Pro}(\mathcal{X}) & \perp & \text{Pro}(\mathcal{Y}) \\ & \xleftarrow{g^*} & \end{array}$$

and for any commutative square of topoi

$$\begin{array}{ccc} \mathcal{X}' & \xrightarrow{g'} & \mathcal{Y}' \\ \downarrow f' & & \downarrow f \\ \mathcal{X} & \xrightarrow{g} & \mathcal{Y} \end{array}$$

applying g'^* to the unit of the adjunction $f_{\#} \dashv f^*$ induces a natural transformation

$$g'^* \rightarrow g'^* f^* f_{\#} \simeq f'^* g^* f_{\#}$$

and hence by transposition

$$f'_{\#} g'^* \rightarrow g^* f_{\#}$$

called the *base change transformation*, which when evaluated at $1_{y'}$ gives the desired

$$f'_{\#} g'^* 1_{y'} \simeq f'_{\#} 1_{x'} \rightarrow g^* f_{\#} 1_{y'}.$$

3.2 Locally contractible geometric morphisms

We start by recalling the definition of a locally cartesian closed ∞ -category.

Definition 3.8. An ∞ -category \mathcal{C} is *cartesian closed* if it admits finite products and, for any object $c \in \mathcal{C}$, the functor $- \times c : \mathcal{C} \rightarrow \mathcal{C}$ admits a right adjoint.

An ∞ -category \mathcal{C} is *locally cartesian closed* if it has pullbacks and, for any object $c \in \mathcal{C}$, the slice $\mathcal{C}_{/c}$ is cartesian closed, or equivalently, if for any arrow $f : c \rightarrow d$ in \mathcal{C} , the functor $\mathcal{C}_{/d} \rightarrow \mathcal{C}_{/c}$ given by pulling back along f admits a right adjoint called the *dependent product along f* and denoted by $\prod_f : \mathcal{C}_{/c} \rightarrow \mathcal{C}_{/d}$. A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ between locally cartesian ∞ -categories is locally cartesian closed if F commutes with pullbacks and dependent products, i.e. for any arrow $f : c \rightarrow d$ in \mathcal{C} , we have a commutative square

$$\begin{array}{ccc} \mathcal{C}_{/c} & \xrightarrow{\prod_f} & \mathcal{C}_{/d} \\ F \downarrow & & \downarrow F \\ \mathcal{D}_{/Fc} & \xrightarrow{\prod_{Ff}} & \mathcal{D}_{/Fd}. \end{array}$$

We will denote $\text{Cat}_{\infty}^{lcc}$ the subcategory of Cat_{∞} whose objects are locally cartesian closed ∞ -categories with locally cartesian closed functors between them.

Example 3.9. By universality of colimits and adjoint functor theorem, any ∞ -topos is locally cartesian closed.

Let $F \rightarrow G$ be a morphism in $\text{Pro}(\mathcal{Y})$ and let $H \rightarrow f^*G$ be a morphism in $\text{Pro}(\mathcal{X})$. Then we have a canonical commutative square

$$\begin{array}{ccc} f_{\#}(f^*F \times_{f^*G} H) & \longrightarrow & f_{\#}H \\ \downarrow & & \downarrow \\ f_{\#}f^*F & \longrightarrow & f_{\#}f^*G \\ \downarrow & & \downarrow \\ F & \longrightarrow & G \end{array}$$

which determines a unique morphism

$$(3.10) \quad f_{\#}(f^*F \times_{f^*G} H) \rightarrow F \times_G f_{\#}H.$$

Proposition 3.11. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a geometric morphism between ∞ -topoi. Consider the conditions

- (i) f^* admits a left adjoint and, for every $F \rightarrow G$ in \mathcal{Y} and $H \rightarrow f^*G$ in \mathcal{X} , the associated projection morphism is invertible;
- (ii) f^* is locally cartesian closed;
- (iii) f^* admits a left adjoint and, for every F in \mathcal{Y} and H in \mathcal{X} , the associated projection morphism

$$f_{\sharp}(f^*F \times H) \rightarrow F \times f_{\sharp}H$$

is invertible.

Then (i) and (ii) are equivalent. Moreover, if f is induced by a continuous map between topological spaces, then these are also equivalent to (iii).

Proof. We first show that (i) implies (ii). Since f^* commutes with finite limits, it suffices to show that it commutes with dependent products. But, for any $\alpha : F \rightarrow G$ in \mathcal{Y} , the square

$$\begin{array}{ccc} \mathcal{Y}/_F & \xrightarrow{\Pi_\alpha} & \mathcal{Y}/_G \\ f^* \downarrow & & \downarrow f^* \\ \mathcal{X}/_{f^*F} & \xrightarrow{\Pi_{f^*\alpha}} & \mathcal{X}/_{f^*G} \end{array}$$

commutes if and only if the square given by the corresponding left adjoints commutes. This last assertion is equivalent to requiring the projection morphisms to be invertible, and so we are done.

We now show that (ii) implies (i). By the same argument as above, it suffices to prove that f^* admits a left adjoint. Since f^* is cocontinuous and preserves finite limits, we are only left to prove that f^* commutes with infinite products: we will do this by exhibiting products (more generally, limits indexed by small ∞ -groupoids) in any ∞ -topos as a special case of dependent products, so that the result will follow by assumption (ii). Let \mathcal{X} be an ∞ -topos, $\pi : \mathcal{X} \rightarrow \mathcal{S}$ the unique geometric morphism. First of all, we observe that the cocontinuous functors

$$\begin{array}{ccc} \mathcal{S} & \longrightarrow & \mathcal{C}at_{\infty}^{\text{op}} \\ A & \longmapsto & \text{Fun}(A, \mathcal{X}) \end{array} \qquad \begin{array}{ccc} \mathcal{S} & \longrightarrow & \mathcal{C}at_{\infty}^{\text{op}} \\ A & \longmapsto & \mathcal{X}/_{\pi^*A} \end{array}$$

are naturally equivalent, since have equivalences

$$\text{Fun}(\Delta^0, \mathcal{X}) \simeq \mathcal{X} \simeq \mathcal{X}/_{\pi^*\Delta^0}.$$

In particular, if $\alpha : A \rightarrow \Delta^0$ is the unique map, we have a corresponding commutative square

$$\begin{array}{ccc} \mathcal{X}/_{\pi^*\Delta^0} & \xrightarrow{\pi^*A \times -} & \mathcal{X}/_{\pi^*A} \\ \downarrow \simeq & & \downarrow \simeq \\ \text{Fun}(\Delta^0, \mathcal{X}) & \xrightarrow{\text{const}} & \text{Fun}(A, \mathcal{X}) \end{array}$$

where the lower horizontal arrow assigns to an object F of \mathcal{X} the constant functor at F . Thus we obtain an identification of the respective right adjoints, i.e. a commutative square

$$\begin{array}{ccc} \mathcal{X}/_{\pi^*A} & \xrightarrow{\Pi_{\pi^*\alpha}} & \mathcal{X}/_{\pi^*\Delta^0} \\ \downarrow \simeq & & \downarrow \simeq \\ \text{Fun}(A, \mathcal{X}) & \xrightarrow{\text{lim}} & \text{Fun}(\Delta^0, \mathcal{X}) \end{array}$$

which is what we wanted.

Assume now that we have an essential geometric morphism $f : \mathcal{X} = \mathrm{Shv}(X) \rightarrow \mathcal{Y} = \mathrm{Shv}(Y)$ induced by a continuous map $f : X \rightarrow Y$. Clearly (iii) is a special case of (i). Assume then that f satisfies the hypothesis (iii). Let $\alpha : F \rightarrow G$ be a morphism in \mathcal{Y} and let $H \rightarrow f^*G$ be a morphism in \mathcal{X} . Since the projection morphism is a natural transformation between colimit preserving functors and since for any $H \in \mathcal{X}_{/f^*G}$ we have an equivalence $H \simeq \varinjlim_{y(U) \rightarrow G} (H \times_{f^*G} y(f^{-1}U))$, we may assume that $H \rightarrow f^*G$ factors as $H \rightarrow y(f^{-1}U) \rightarrow f^*G$ for some open $U \in \mathcal{U}(Y)$, and hence by the pasting properties of pullbacks we may also assume that $G = y(U)$. Notice that, for any (-1) -truncated object V in a topos \mathcal{Z} and for any other two objects $A, B \in \mathcal{Z}_{/V}$, we have an identification $A \times_V B \simeq A \times B$: this follows because for any other object C mapping both to A and B , we have that $\mathrm{Hom}_{\mathcal{Z}}(C, V)$ is contractible, and thus $\mathrm{Hom}_{\mathcal{Z}}(C, A \times_V B) \simeq \mathrm{Hom}_{\mathcal{Z}}(C, A \times B)$. Thus, since both $y(U)$ and $y(f^{-1}U)$ are (-1) -truncated, we are only left to prove that

$$f_{\#}(f^*F \times H) \rightarrow F \times f_{\#}H$$

is invertible, which is true by assumption. \square

Definition 3.12. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a geometric morphism of ∞ -topoi. We say that f is *essential* if $f_{\#}$ factors through \mathcal{Y} , or equivalently if f^* admits a left adjoint. Furthermore, we say that an essential geometric morphism is *locally contractible* if it satisfies the equivalent conditions (i) and (ii) in Proposition 3.11. We say that a geometric morphism is of *trivial shape* if f^* is fully faithful, or equivalently if the unit transformation $\mathrm{id}_{\mathcal{Y}} \rightarrow f_*f^*$ is an equivalence. When f is the unique geometric morphism $\mathcal{X} \rightarrow \mathcal{S}$, we will say that \mathcal{X} is locally contractible.

Remark 3.13. For a continuous map $f : X \rightarrow Y$ be a continuous map inducing a, essential geometric morphism, one may interpret the condition of being locally contractible geometric morphism as the requirement of a base change for $f_{\#}$ along open immersions. More precisely, let U be an open subset of Y , and consider the pullback square

$$\begin{array}{ccc} f^{-1}(U) & \xrightarrow{f'} & U \\ \downarrow j' & & \downarrow j \\ X & \xrightarrow{f} & Y. \end{array}$$

For any $F \in \mathrm{Shv}(X)$, we have natural equivalences

$$j_{\#}j^*f_{\#}F \xrightarrow{\simeq} f_{\#}F \times y(U) \xrightarrow{\simeq} f_{\#}(F \times y(f^{-1}(U))) \xrightarrow{\simeq} f_{\#}j'_{\#}(j')^*F \xrightarrow{\simeq} j_{\#}f'_{\#}(j')^*F$$

where the second morphism is (3.10). Therefore, since $j_{\#}$ is fully faithful, we obtain a natural equivalence

$$j^*f_{\#} \simeq f'_{\#}(j')^*,$$

and, by transposition, an equivalence

$$f^*j_* \simeq j'_*(f')^*.$$

Example 3.14. (i) Recall that any object $U \in \mathcal{X}$ of an ∞ -topos determines a geometric morphism $j : \mathcal{X}_{/U} \rightarrow \mathcal{X}$. By [Lur09, Proposition 6.3.5.1] j is locally contractible, and $j_{\#} : \mathcal{X}_{/U} \rightarrow \mathcal{X}$ can be described as the usual forgetful functor.

- (ii) By [Lur17, Proposition A.1.9], an ∞ -topos is locally contractible if and only if the unique geometric morphism $\mathcal{X} \rightarrow \mathcal{S}$ is essential, since in this case the projection morphism is automatically invertible.
- (iii) Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be an essential geometric morphism. For any the ∞ -topos \mathcal{Z} , by Remark 2.2 we obtain a geometric morphism $f \otimes \mathcal{Z} : \mathcal{X} \otimes \mathcal{Z} \rightarrow \mathcal{Y} \otimes \mathcal{Z}$ by applying to f the functor $- \otimes \mathcal{Z}$. Since both f^* and f_{\sharp} commute with colimits, the adjunction $f_{\sharp} \dashv f^*$ is preserved by $- \otimes \mathcal{Z}$, and so $f \otimes \text{id}_{\mathcal{Z}}$ is an essential geometric morphism.

Remark 3.15. Let $f : X \rightarrow Y$ be a continuous map. Notice that, since the functor

$$f^{-1} : \mathcal{U}(Y) \rightarrow \mathcal{U}(X)$$

preserves open coverings, for any complete ∞ -category \mathcal{C} we still have a well defined push-forward $f_* : \text{Shv}(X; \mathcal{C}) \rightarrow \text{Shv}(Y; \mathcal{C})$ given as usual by $\Gamma(U; f_* F) = \Gamma(f^{-1}(U); F)$ for all $U \in \mathcal{U}(Y)$. Although at this level of generality there is no reason to expect f_* to have a left adjoint, if f induces an essential geometric morphism at the level of sheaves of spaces, then it actually does. Indeed, recall that there is an equivalence

$$\text{Shv}(X; \mathcal{C}) \simeq \text{Fun}_*(\text{Shv}(X)^{\text{op}}, \mathcal{C}).$$

Through this equivalence and Example 2.22 (i), we can identify $f_* : \text{Fun}_*(\text{Shv}(X)^{\text{op}}, \mathcal{C}) \rightarrow \text{Fun}_*(\text{Shv}(Y)^{\text{op}}, \mathcal{C})$ with precomposition with the opposite of the pullback $f^* : \text{Shv}(Y) \rightarrow \text{Shv}(X)$. Thus, similarly to Remark 2.2, by applying the 2-functor $\text{Fun}_*((-)^{\text{op}}, \mathcal{C})$ to the adjunction between cocontinuous functors $f_{\sharp} \dashv f^*$, we obtain the desired left adjoint.

It is straightforward to check that the composition of two locally contractible geometric morphism is again locally contractible (see [AC21, Corollary 3.2.5]). We observe that the properties of being essential or locally contractible can be checked locally on the source.

Lemma 3.16. Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a geometric morphism, and let $\mathcal{B} \subseteq \mathcal{X}$ which generates \mathcal{X} under colimits. For any object $U \in \mathcal{B}$, consider the composite geometric morphism

$$(3.17) \quad \mathcal{X}_{/U} \longrightarrow \mathcal{X} \xrightarrow{f} \mathcal{Y}.$$

We have the following

- (i) f is essential if and only if (3.17) is of constant shape for any $U \in \mathcal{B}$;
- (ii) f is locally contractible if and only if (3.17) is locally contractible for any $U \in \mathcal{B}$.

Proof. A proof can be found in [AC21, Proposition 3.1.5] and [AC21, Proposition 3.2.6]. \square

Remark 3.18. The content of part (i) in Lemma 3.16 suggests that a valid alternative way to call a geometric morphism whose pullback has a left adjoint could have been *locally of constant shape*. This is actually the approach taken by Lurie in [Lur17, Appendix A]; however, we have decided to stick with the more concise nomenclature which appears also in [Joh02] and [AC21].

Corollary 3.19. Let X be a locally contractible topological space, $a : X \rightarrow *$ the unique map, and assume that $\mathcal{X} = \text{Shv}(X)$ is hypercomplete. Then $\text{Shv}(X)$ is locally contractible and a_{\sharp} is equivalent to the extension by colimits of the cosheaf Sing . Consequently, for sheaves of spectra, the functor $a_{\sharp} : \text{Shv}(X; \mathbb{S}p) \rightarrow \mathbb{S}p$ obtained by applying $- \otimes \mathbb{S}p$ is uniquely determined by the formula $a_{\sharp}(\mathbb{S}_U) = \Sigma_{\mp}^{\infty} U$ for any $U \in \mathcal{U}(X)$.

Proof. Let B be the poset of all contractible open subsets of X . Notice that, even though B in general is not a sieve, by the hypercompleteness assumption of $\mathrm{Shv}(X)$ we get an equivalence

$$\varinjlim_{U \in B/V} y(U) \simeq y(V)$$

for any $V \in \mathcal{U}(X)$ that can be easily checked on stalks. In particular, we see that the full subcategory $\mathrm{Shv}(B) \subseteq \mathrm{Shv}(X)$ generates $\mathrm{Shv}(X)$ under colimits. Thus by Lemma 3.16 it suffices to check that $\mathrm{Shv}(X)_{/U} \simeq \mathrm{Shv}(U)$ is of constant shape for any $U \in B$, but this is true by homotopy invariance of the shape. The last assertion follows immediately by noticing that through the equivalence $\mathcal{S} \otimes \mathrm{Sp} \simeq \mathrm{Sp}$, an object $A \otimes \mathcal{S}$ corresponds (functorially on A) to $\Sigma_{\dagger}^{\infty} A$. \square

Remark 3.20. Beware that the viceversa of Corollary 3.19 is not true. For this reason, to avoid confusion, from now on we will say that a topological space X is essential if $\mathrm{Shv}(X)$ is (or equivalently, if $\mathrm{Shv}(X)$ is locally contractible by Example 3.14 part (ii)).

3.3 Shape submersions

Definition 3.21. A continuous map $f : X \rightarrow Y$ between topological spaces is a *shape submersion* if for every point $x \in X$ there exist an open neighbourhood U of x and a space X' which is essential, such that $f(U)$ is open in Y , U is homeomorphic to $f(U) \times X'$ and the diagram

$$\begin{array}{ccccc} f(U) \times X' & \xrightarrow{\cong} & U & \hookrightarrow & X \\ & \searrow p & \downarrow & & \downarrow f \\ & & f(U) & \hookrightarrow & Y \end{array}$$

commutes, where p is the obvious projection.

Example 3.22. (i) By Corollary 3.19, if X is locally contractible and hypercomplete, then $X \rightarrow *$ is a shape submersion.

(ii) Any topological submersion of fiber dimension n is a shape submersion.

Remark 3.23. (i) It follows easily from the definition that shape submersions are stable under pullbacks of topological spaces.

(ii) If $f : X \rightarrow Y$ is a shape submersion, then the set of open subsets of X of the type $X' \times V$, where V is open in Y and X' is essential, forms a basis for the topology of X . Although this basis is not closed under finite intersections, the set of representable sheaves corresponding to open subsets homeomorphic to the product of an open in Y and an essential space generates $\mathrm{Shv}(X)$ under colimits. To see this, consider

$$\mathcal{B} = \{U \in \mathcal{U}(X) \mid U \cong W, \text{ with } W \in \mathcal{U}(V \times S) \text{ for some } V \in \mathcal{U}(Y), S \text{ essential}\}.$$

The set \mathcal{B} clearly forms a basis closed under finite intersections, and then we have $\mathrm{Shv}(X) \simeq \mathrm{Shv}(\mathcal{B})$ by [Aok20, Appendix A]. Moreover, since open immersions induce essential geometric morphisms, and since $\mathcal{U}(V) \times \mathcal{U}(S)$ forms a basis of $V \times S$ which is closed under finite intersections, we get our claim.

In the particular case of a topological submersion of fiber dimension n , since \mathbb{R}^n is hypercomplete, we have an equivalence $\mathrm{Shv}(\mathbb{R}^n) \simeq \mathrm{Shv}(W)$ where $W \subseteq \mathcal{U}(\mathbb{R}^n)$ is the poset of open balls inside \mathbb{R}^n , and thus, since \mathbb{R}^n is locally compact, we have

$\mathrm{Shv}(V \times \mathbb{R}^n) \simeq \mathrm{Shv}(V) \otimes \mathrm{Shv}(W)$. In particular, the set of representable sheaves corresponding to open subsets homeomorphic to the product of an open in Y and an open ball in \mathbb{R}^n generates $\mathrm{Shv}(X)$ under colimits.

For technical reasons that will be justified in a moment, from now on whenever we have a shape submersion $f : X \rightarrow Y$ we will assume that either Y is locally compact or all spaces of locally constant shape appearing in the basis of X are locally compact.

Lemma 3.24. Any shape submersion $f : X \rightarrow Y$ induces an essential geometric morphism. Thus, for any presentable ∞ -category \mathcal{C} , we obtain an adjunction $f_{\#} \dashv f^*$ for \mathcal{C} -valued sheaves.

Proof. Since f^* is a left adjoint, it is in particular accessible. Hence, by adjoint functor theorem, it suffices to prove that f^* commutes with limits. For any functor $I \rightarrow \mathrm{Shv}(Y)$ we have a canonical map

$$f^*(\varprojlim_{i \in I} F_i) \rightarrow \varprojlim_{i \in I} f^* F_i$$

and it suffices to check that this is an equivalence after restricting to any open subset in the basis of X associated to f . Hence, since the operation of restricting a sheaf to an open subset commutes with limits, we can assume that f is a projection $f : X \times Y \rightarrow Y$ where either X or Y is locally compact and X is essential. Thus, by point (iii) in Example 3.14 and Proposition 2.30, we get that f is essential. The last assertion follows immediately by Corollary 2.24. \square

Lemma 3.25 (Smooth base change). Let \mathcal{C} be a presentable ∞ -category. For every given pullback square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & X \\ g' \downarrow & \lrcorner & \downarrow g \\ Y' & \xrightarrow{f} & Y \end{array}$$

of topological spaces where g and g' are shape submersions, there is a natural equivalence

$$f^* g_{\#} \simeq g'_{\#} f'^*$$

and, by transposition, also

$$g^* f_{*} \simeq f'_* g'^*$$

for \mathcal{C} -valued sheaves.

Proof. First of all, the base change transformation as defined in Remark 3.7 defines a comparison natural transformation. Since all functors appearing are colimit preserving, it suffices to check that the morphism is an equivalence only on a family of objects generating $\mathrm{Shv}(X;)$ by colimits. Hence, applying Remark 3.23 (ii), we see that we can assume that the pullback square is of the type

$$\begin{array}{ccc} X \times Y' & \xrightarrow{\mathrm{id}_X \times f} & X \times Y \\ g' \downarrow & \lrcorner & \downarrow g \\ Y' & \xrightarrow{f} & Y \end{array}$$

where X is essential, g and g' are the canonical projections. By Example 3.14 point (iii) and Proposition 2.30, we have $(\mathrm{id}_X \times f)^* \simeq (\mathrm{id}_X)^* \otimes f^*$, $g_{\#} \simeq a_{\#} \otimes (\mathrm{id}_Y)_{\#}$ and $g'_{\#} \simeq a_{\#} \otimes (\mathrm{id}_{Y'})_{\#}$,

where $a : X \rightarrow *$, and so, since \otimes is a bifunctor, we have

$$\begin{aligned} f^*g_{\sharp} &\simeq f^*(a_{\sharp} \otimes (\text{id}_Y)_{\sharp}) \\ &\simeq a_{\sharp} \otimes f^* \\ &\simeq (a_{\sharp} \otimes (\text{id}_{Y'})_{\sharp})(\text{id}_X^* \otimes f^*) \\ &\simeq g'_{\sharp}(\text{id}_X \times f)^*. \end{aligned}$$

□

Corollary 3.26 (Smooth projection formula). Let $f : X \rightarrow Y$ be a shape submersion, \mathcal{C} and \mathcal{D} two presentable ∞ -categories. Then for any $F \in \text{Shv}(X; \mathcal{C})$ and $G, H \in \text{Shv}(Y; \mathcal{D})$, we have a canonical equivalence

$$f_{\sharp}(F \otimes f^*G) \simeq f_{\sharp}F \otimes G$$

and hence, by transposition, when $\mathcal{C} = \mathcal{D}$ is monoidal, equivalences

$$f_*\underline{\text{Hom}}_X(F, f^*G) \simeq \underline{\text{Hom}}_Y(f_{\sharp}F, G)$$

and

$$f^*\underline{\text{Hom}}_Y(G, H) \simeq \underline{\text{Hom}}_X(f^*G, f^*H).$$

In particular, any shape submersion induces a locally contractible geometric morphism.

Proof. Let $\Gamma_f : X \rightarrow Y$ be the graph of f . We have

$$\begin{aligned} f_{\sharp}(F \otimes f^*G) &\simeq f_{\sharp}\Gamma_f^*(F \boxtimes G) \\ &\simeq \Delta^*(f \times \text{id}_Y)_{\sharp}(F \boxtimes G) \\ &\simeq f_{\sharp}F \otimes G \end{aligned}$$

where the second equivalence follows by applying Lemma 3.25 to the pullback square

$$\begin{array}{ccc} X & \xrightarrow{\Gamma_f} & X \times Y \\ \downarrow f & & \downarrow f \times \text{id}_Y \\ Y & \xrightarrow{\Delta} & Y \times Y. \end{array}$$

The last assertion follows by specializing (2.28) to the case when \mathcal{C} is \mathcal{S} equipped with the cartesian monoidal structure and by the commutativity of the diagram

$$\begin{array}{ccc} F \times f^*G & \longrightarrow & f^*(f_{\sharp}F \times G) \\ \downarrow \simeq & & \downarrow \simeq \\ \Gamma_f^*(F \boxtimes G) & \longrightarrow & \Gamma_f^*(f \times \text{id}_Y)^*(f \times \text{id}_Y)_{\sharp}(F \boxtimes G) \end{array}$$

where the upper horizontal arrow is the one which transposes to the projection morphism and the lower horizontal one transposes to the smooth base change transformation. □

4 Localization sequences

We will now prove a version of the localization theorem for sheaves of spaces (and of spectra) on a topological space X : this essentially states that, for any closed immersion $i : Z \rightarrow X$ with open complement $j : U \rightarrow X$, the inclusions

$$\mathrm{Shv}(Z) \xleftarrow{i_*} \mathrm{Shv}(X) \xleftarrow{j_*} \mathrm{Shv}(U)$$

form a *recollement* (in the sense of [Lur17, Definition A.8.1]). Achieving this goal in our context will be slightly more complicated than in the case of [KS90, Proposition 2.3.6], namely because we don't want to assume all our spaces to be hypercomplete. We will follow instead the strategy outlined in [Kha19]: the main ingredient will be to show that the pushforward $i_* : \mathrm{Shv}(Z) \rightarrow \mathrm{Shv}(X)$ commutes with *contractible* colimits, i.e. colimits indexed by contractible simplicial sets. From this we will be able to reduce to checking the theorem in the case of representable sheaves, which is almost straightforward.

We start by reporting [Kha19, Definition 3.1.5] and [Kha19, Lemma 3.1.6].

Definition 4.1. Let X and Y be essentially small ∞ -sites, and assume that Y admits an initial object \emptyset_Y . A functor $u : X \rightarrow Y$ is *topologically quasi-cocontinuous* if for every covering sieve $R' \hookrightarrow y(u(x))$ in Y , the sieve $R \hookrightarrow y(x)$, generated by morphisms $x' \rightarrow x$ such that either $u(x')$ is initial or $y(u(x')) \rightarrow y(u(x))$ factors through $R' \hookrightarrow y(u(x))$, is a covering in X .

Lemma 4.2. With notation as in the previous definition, let $u : X \rightarrow Y$ be a topologically quasicontinuous functor. Assume that the initial object \emptyset_Y is strict in the sense that for any object $y \in Y$, any morphism $d \rightarrow \emptyset_Y$ is invertible. Assume also that, for any object $y \in Y$, the sieve $\emptyset_{\mathrm{Fun}(Y, \mathcal{S})} \hookrightarrow y(d)$ is a covering in Y if and only if y is initial (where $\emptyset_{\mathrm{Fun}(Y, \mathcal{S})}$ denotes the initial object of $\mathrm{Fun}(Y, \mathcal{S})$). Then the functor $\mathrm{Shv}(Y) \rightarrow \mathrm{Shv}(X)$, given by the assignment $F \mapsto L_X(u^*(F))$, where $L_X : \mathrm{Fun}(X, \mathcal{S}) \rightarrow \mathrm{Shv}(X)$ denotes the sheafification functor, commutes with contractible colimits.

Lemma 4.3. Let $i : Z \hookrightarrow X$ be a closed immersion. Then i_* commutes with contractible colimits.

Proof. By the lemma above and by unraveling the definition of topologically quasi-cocontinuous functor, this amounts to check that, for any $V \in \mathcal{U}(X)$ and any open covering $\{W_i\}_{i \in I} \subseteq \mathcal{U}(Z)$ of $V \cap Z$, the family

$$T = \{U \subseteq V \mid U \cap Z = \emptyset \text{ or } U \cap Z \subseteq W_i \text{ for some } i \in I\} \subseteq \mathcal{U}(X)$$

covers V . But this is clear, because $V \setminus Z \in T$ and any W_i can be written as $W'_i \cap Z$ for some $W'_i \in \mathcal{U}(V)$. \square

Corollary 4.4. Let \mathcal{C} be any pointed presentable ∞ -category. Then the pushforward $i_*^{\mathcal{C}} : \mathrm{Shv}(Z; \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C})$ commutes with all colimits, and thus admits a right adjoint $i_c^{\mathcal{C}} : \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(Z; \mathcal{C})$.

Proof. It suffices to prove the corollary for $\mathcal{C} = \mathcal{S}_*$. Note that it suffices to check that i_* preserves the initial object and commutes with contractible colimits: any $F : I \rightarrow \mathcal{D}$ from a simplicial set I to an ∞ -category \mathcal{D} with an initial object $\emptyset_{\mathcal{D}}$ may be seen as $I \rightarrow \mathcal{D}_{\emptyset_{\mathcal{D}}/}$ and thus corresponds to a functor $\Delta^0 \star I \rightarrow \mathcal{D}$ with the same colimit as F but indexed by weakly contractible simplicial set. But i_* preserves the initial object because $\mathrm{Shv}(Z; \mathcal{S}_*)$ is pointed, and thus we may conclude by the previous lemma. \square

Let $i : Z \hookrightarrow X$ be a closed immersion with open complement $j : U \hookrightarrow X$. For any $F \in \text{Shv}(X)$, consider the functorial commutative square

$$\begin{array}{ccc} j_{\#}j^*(F) & \longrightarrow & F \\ \downarrow & & \downarrow \\ j_{\#}j^*i_*i^*(F) & \longrightarrow & i_*i^*(F), \end{array}$$

where all the morphisms are given by the obvious units and counits. Notice that for any $G \in \text{Shv}(X)$ and $V \in \mathcal{U}(U)$, we have

$$\Gamma(V; i_*G) = \Gamma(U \cap Z; G) \simeq *,$$

and so we can identify $j^*i_* : \text{Shv}(Z) \rightarrow \text{Shv}(U)$ with a constant functor with value the terminal object $y(U) \in \text{Shv}(U)$. Hence the previous square may be written as

$$(4.5) \quad \begin{array}{ccc} j_{\#}j^*(F) & \longrightarrow & F \\ \downarrow & & \downarrow \\ j_{\#}(y(U)) & \longrightarrow & i_*i^*(F). \end{array}$$

Theorem 4.6. The canonical square (4.5) is a pushout.

Proof. Since all functors appearing in (4.5) commute with contractible colimits and any sheaf on X is canonically written as colimit indexed by the contractible category $\mathcal{U}(X)_{/F} = \text{Shv}(X)_{/F} \times_{\text{Shv}(X)} \mathcal{U}(X)$ (it has an initial object), it suffices to prove the theorem when $F = y(V)$ for some $V \in \mathcal{U}(X)$, and hence we just need to show that $i_*i^*(y(V)) \simeq y(U \cup V)$. For any $W \in \mathcal{U}(X)$, we have

$$\begin{aligned} \Gamma(W; i_*i^*(y(V))) &\simeq \Gamma(W; i_*(y(V \cap Z))) \\ &= \Gamma(W \cap Z; y(V \cap Z)) \\ &= \text{Hom}_{\mathcal{U}(Z)}(W \cap Z, V \cap Z) \\ &= \text{Hom}_{\mathcal{U}(X)}(W, V \cup U) \\ &= \Gamma(W; y(V \cup U)), \end{aligned}$$

where the second to last identification follows by the usual exponential adjunction in the boolean algebra of all subsets of X . \square

Corollary 4.7. Let $i : Z \hookrightarrow X$ be a closed immersion with open complement $j : U \hookrightarrow X$, and let $i_*^{\mathcal{C}}, i_{\mathcal{C}}^*, i_{\mathcal{C}}^!, j_{\#}^{\mathcal{C}}$ and $j_{\mathcal{C}}^*$ be the induced pushforward and pullback functors at the level of \mathcal{C} -valued sheaves, where \mathcal{C} is any pointed presentable ∞ -category. Then we get a canonical cofiber sequence

$$(4.8) \quad j_{\#}^{\mathcal{C}}j_{\mathcal{C}}^*F \rightarrow F \rightarrow i_*^{\mathcal{C}}i_{\mathcal{C}}^*F$$

and dually a fiber sequence

$$(4.9) \quad i_*^{\mathcal{C}}i_{\mathcal{C}}^!F \rightarrow F \rightarrow j_{\mathcal{C}}^*j_{\#}^{\mathcal{C}}F.$$

denote

Proof. It suffices to treat only the case of the sequence (4.8). Furthermore, we only need to prove the case of sheaves of pointed spaces, since all functors appearing are colimit preserving and we have a canonical equivalence $\mathcal{S}h\mathbf{v}(X; \mathcal{C}) \simeq \mathcal{S}h\mathbf{v}(X; \mathcal{S}_*) \otimes \mathcal{C}$, where \mathcal{C} is any presentable pointed ∞ -category and X is any topological space. We define the canonical morphisms in (4.8) through counit and unit of the appropriate adjunctions, and we see immediately that the composition of those two morphisms is null-homotopic because $i_{\mathcal{S}_*}^* j_{\#}^{\mathcal{S}_*}$ is equivalent to a constant functor with value the zero object in $\mathcal{S}h\mathbf{v}(Z; \mathcal{S}_*)$.

Let $\alpha : \mathcal{S}h\mathbf{v}(X; \mathcal{S}_*) \rightarrow \mathcal{S}h\mathbf{v}(X)$ be the forgetful functor. A close inspection of the appropriate universal properties shows that, for any $F \in \mathcal{S}h\mathbf{v}(X; \mathcal{S}_*)$, there is a canonical pushout square

$$\begin{array}{ccc} j_{\#}(y(U)) & \longrightarrow & y(X) \\ \downarrow & \lrcorner & \downarrow \\ j_{\#} j^* \alpha(F) & \longrightarrow & \alpha(j_{\#}^{\mathcal{S}_*} j_{\mathcal{S}_*}^* F), \end{array}$$

where the left vertical map is induced by the point of F . Thus, since α reflects pushouts and we have an equivalence $\alpha i_{\mathcal{S}_*}^{\mathcal{S}_*} i_{\mathcal{S}_*}^* \simeq i_* i^* \alpha$, it suffices to prove that the canonical square

$$\begin{array}{ccc} \alpha(j_{\#}^{\mathcal{S}_*} j_{\mathcal{S}_*}^* F) & \longrightarrow & \alpha(F) \\ \downarrow & & \downarrow \\ y(X) & \longrightarrow & i_* i^* \alpha(F). \end{array}$$

induced by applying α to the sequence (4.8) is a pushout. For this purpose, consider the commutative diagram

$$\begin{array}{ccccc} j_{\#}(y(U)) & \longrightarrow & y(X) & & \\ \downarrow & & \downarrow & & \\ j_{\#} j^* \alpha(F) & \longrightarrow & \alpha(j_{\#}^{\mathcal{S}_*} j_{\mathcal{S}_*}^* F) & \longrightarrow & \alpha(F) \\ \downarrow & & \downarrow & & \downarrow \\ j_{\#} j^*(y(X)) & \longrightarrow & y(X) & \longrightarrow & i_* i^* \alpha(F). \end{array}$$

denote The upper left square and the left vertical rectangle are both pushouts, and so also the lower left square is a pushout. But the lower horizontal rectangle is a pushout, and so we can conclude. \square

Corollary 4.10. Consider a pullback square

$$\begin{array}{ccc} Z' & \xrightarrow{f'} & Z \\ s' \downarrow & \lrcorner & \downarrow s \\ X & \xrightarrow{f} & Y \end{array}$$

where f and f' are shape submersions and s (and consequently s') is a closed immersion. Then, for sheaves with values in a pointed presentable ∞ -category, we have a canonical equivalence

$$s_! f'_{\#} \simeq f_{\#} s'_!$$

or equivalently

$$f'^* s^! \simeq s'^! f^*.$$

Proof. Let j and j' be the complement open immersions associated respectively to s and s' . By the localization sequences and by smooth base change, we have a commutative diagram where all the rows are cofiber sequences and all the vertical arrows are invertible

$$\begin{array}{ccccc}
j_{\#}j^*f_{\#} & \longrightarrow & f_{\#} & \longrightarrow & s_!s^*f_{\#} \\
\downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\
j_{\#}f'_{\#}j'^* & \longrightarrow & f_{\#} & \longrightarrow & s_!f'_{\#}s'^* \\
\downarrow \simeq & & \downarrow \simeq & & \downarrow \simeq \\
f_{\#}j'_{\#}j'^* & \longrightarrow & f_{\#} & \longrightarrow & f_{\#}s'_!s'^*.
\end{array}$$

Hence we may conclude by precomposing the dotted equivalences with $s'_!$, since $s'_!$ is fully faithful. \square

Remark 4.11. It is not hard to see that one may deduce from Theorem 4.6 that i_* is fully faithful (this was already proven in [Lur09, Corollary 7.3.2.10]). From this follows immediately that, in the case of sheaves of pointed spaces or of spectra, one has the identification

$$i^! \simeq \text{fib} \left(i^* \xrightarrow{i^*(\text{unit})} i^*j_*j^* \right).$$

Remark 4.12. From Theorem 4.6 one deduces immediately that, at least when \mathcal{C} is presentable stable, the functors i^* and j^* are *jointly conservative*, i.e. a morphism $\alpha : F \rightarrow G$ in $\text{Shv}(X; \mathcal{C})$ is invertible if and only if both $i^*(\alpha)$ and $j^*(\alpha)$ are invertible. This implies in particular that the fully faithful functors i_* and j_* make $\text{Shv}(X; \mathcal{C})$ a *recollement* of $\text{Shv}(Z; \mathcal{C})$ and $\text{Shv}(U; \mathcal{C})$, in the sense of [Lur17, Definition A.8.1]. However, this is true in a much greater generality: see [Hai21] for a proof in the cases when \mathcal{C} is an ∞ -topos or compactly generated. After Corollary 5.16, for stable coefficients, we will also be able to relax the presentability assumption to the more general requirement for \mathcal{C} to admit both limits and colimits.

5 Pullbacks with stable bicomplete coefficients

From now on, unless otherwise specified, all the topological spaces we will deal with will be assumed to be *locally compact and Hausdorff*. This implies that the following are equivalent

1. $f : X \rightarrow Y$ is proper (i.e. the preimage of any compact subset of Y is compact);
2. f is closed with compact fibers;
3. f is universally closed.

Another important consequence of the previous assumption is that any map $X \xrightarrow{f} Y$ can be factored as a composition of a closed immersion (which is in particular proper by the characterization above), an open immersion and a proper map as follows

$$(5.1) \quad \begin{array}{ccc} X \times Y & \xrightarrow{j \times \text{id}_Y} & \overline{X} \times Y \\ \Gamma_f \uparrow & & \downarrow p \\ X & \xrightarrow{f} & Y \end{array}$$

where Γ_f is the graph of f , $X \xrightarrow{j} \overline{X}$ is the inclusion of X into its one point compactification and p is the projection to the second coordinate: this factorization will be used very often

later. Notice that one may also use the Stone-Cech compactification β to produce a functorial factorization

$$(5.2) \quad \begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow j & \searrow p & \downarrow \\ \overline{X} & \xrightarrow{p} & Y \\ \downarrow & \lrcorner & \downarrow \\ \beta X & \longrightarrow & \beta Y \end{array}$$

where j is an open immersion and p is proper. However, in many cases it will turn out to be more convenient to have the proper map in the factorization to be a product projection.

This section contains the central technical ingredients of our paper. We will explain what are the two fundamental facts that allow us to construct pullbacks with non-presentable coefficients: in brief, these are *covariant Verdier duality* (see [Lur17, Theorem 5.5.5.1]) and the dualizability of $\mathrm{Shv}(X; \mathcal{S}\mathrm{p})$ in $\mathrm{Cocont}_{\infty}^{st}$.

We will start by giving an exposition of covariant Verdier Duality. Following [Lur17, Theorem 5.5.5.1], this is an equivalence between the categories of sheaves and cosheaves on a locally compact Hausdorff space. We will essentially review the proof of [Lur17, Theorem 5.5.5.1], and try to clarify a bit the last step. The reader who is aware of this result may safely skip the first part of this section. Later, we will prove that $\mathrm{Shv}(X; \mathcal{S}\mathrm{p})$ is strongly dualizable in the symmetric monoidal ∞ -category $\mathrm{Cocont}_{\infty}^{st}$, by explicitly exhibiting it as a retract in $\mathrm{Cocont}_{\infty}^{st}$ of a compactly generated ∞ -category. This result is not new, as it could be deduced from [Lur16, Proposition 21.1.7.1] (see also [Hoy]): we would like to thank Peter Haine for pointing this out. We then use the above retraction to provide a convenient description of pushforwards along proper maps, that is compatible with tensor products in $\mathrm{Cocont}_{\infty}^{st}$. We conclude the section by showing that $f_{\mathcal{S}\mathrm{p}}^* \otimes \mathcal{C}$ is a left adjoint of $f_*^{\mathcal{C}}$, which is not at all immediate. It will require a use of the factorizations mentioned in the beginning, and a careful combination of all the previously mentioned results.

5.1 Recollections on covariant Verdier duality

We start by recalling the definition of \mathcal{K} -sheaves.

Definition 5.3. Let $\mathcal{K}(X)$ be the poset of compact subsets of a topological space X . Consider the following conditions:

- (i) $F(\emptyset)$ is a terminal object,
- (ii) For every $K, K' \in \mathcal{K}(X)$ the square

$$\begin{array}{ccc} \Gamma(K \cup K'; F) & \longrightarrow & \Gamma(K; F) \\ \downarrow & \lrcorner & \downarrow \\ \Gamma(K'; F) & \longrightarrow & \Gamma(K \cap K'; F) \end{array}$$

is pullback,

- (iii) For every $K \in \mathcal{K}(X)$, the canonical map

$$\varinjlim_{K \Subset K'} \Gamma(K'; F) \rightarrow \Gamma(K; F)$$

is invertible, where $K \Subset K'$ means that K' contains an open neighbourhood of K .

Notice that (i) and (ii) together are equivalent to the sheaf condition for the Grothendieck topology on $\mathcal{K}(X)$ given by finite covering. Hence, we will denote by $\text{Shv}(\mathcal{K}(X); \mathcal{C})$ the full subcategory spanned by presheaves satisfying (i) and (ii). Moreover, we will say that a functor $F : \mathcal{K}(X) \rightarrow \mathcal{C}$ is a \mathcal{K} -sheaf if it satisfies (i), (ii) and (iii). We will denote by $\text{Shv}_{\mathcal{K}}(X; \mathcal{C}) \subseteq \text{Fun}(\mathcal{K}(X)^{\text{op}}, \mathcal{C})$ the full subcategory spanned by \mathcal{K} -sheaves.

It is possible to relate \mathcal{K} -sheaves with usual sheaves. Let M be the union $\mathcal{U}(X) \cup \mathcal{K}(X)$ considered as a poset contained in the power set of X , and let $i : \mathcal{U}(X) \hookrightarrow M$ and $j : \mathcal{K}(X) \hookrightarrow M$ be the corresponding inclusion. We thus get two adjunctions

$$\begin{array}{ccc} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \xrightleftharpoons[i^*]{i_!} & \text{Fun}(M^{\text{op}}, \mathcal{C}) \\ & \perp & \xrightleftharpoons[j_*]{j^*} \\ \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \perp & \text{Fun}(\mathcal{K}(X)^{\text{op}}, \mathcal{C}). \end{array}$$

More explicitly, at the level of objects the functors are given by the formulas

$$\begin{aligned} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) &\xrightarrow{\theta} \text{Fun}(\mathcal{K}(X)^{\text{op}}, \mathcal{C}) \\ F &\longmapsto (K \mapsto \varinjlim_{K \subseteq U} \Gamma(U; F)) \\ \text{Fun}(\mathcal{K}(X)^{\text{op}}, \mathcal{C}) &\xrightarrow{\psi} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) \\ G &\longmapsto (U \mapsto \varprojlim_{K \subseteq U} \Gamma(K; G)). \end{aligned}$$

These two functors actually restrict to an equivalence, assuming \mathcal{C} has limits and colimits, and filtered colimits are exact.

Theorem 5.4. Let \mathcal{C} be a bicomplete ∞ -category where filtered colimits are exact. Then the functors θ and ψ defined above restrict to an equivalence

$$\text{Shv}(X; \mathcal{C}) \simeq \text{Shv}_{\mathcal{K}}(X; \mathcal{C}).$$

Proof. A proof can be found in [Lur09, Theorem 7.3.4.9]. □

Remark 5.5. Since in any stable ∞ -category \mathcal{C} filtered colimits are exact, and since the opposite of any stable ∞ -category is again stable, by Theorem 5.4 we get equivalences

$$\text{Shv}(X; \mathcal{C}) \simeq \text{Shv}_{\mathcal{K}}(X; \mathcal{C})$$

and

$$\text{CoShv}(X; \mathcal{C}) \simeq \text{Shv}(X; \mathcal{C}^{\text{op}})^{\text{op}} \simeq \text{CoShv}_{\mathcal{K}}(X; \mathcal{C})$$

where we define $\text{CoShv}_{\mathcal{K}}(X; \mathcal{C})$ to be $\text{Shv}_{\mathcal{K}}(X; \mathcal{C}^{\text{op}})^{\text{op}}$.

Definition 5.6. Let $F \in \text{Shv}(X; \mathcal{C})$, $U \in \mathcal{U}(X)$, and K any closed subset of X . We define the *sections of F supported at K* and *compactly supported sections of F over U* respectively as

$$\begin{aligned} \Gamma_K(X; F) &:= \text{fib}(\Gamma(X; F) \rightarrow \Gamma(X \setminus K; F)) \\ \Gamma_c(U; F) &:= \varinjlim_{K \subseteq U} \Gamma_K(X; F), \end{aligned}$$

where the colimit ranges over all compact subsets of U .

Remark 5.7. Notice that, if $K \subseteq U$ for some open U , we get a pullback square

$$\begin{array}{ccc} \Gamma(X; F) & \longrightarrow & \Gamma(X \setminus K; F) \\ \downarrow & \lrcorner & \downarrow \\ \Gamma(U; F) & \longrightarrow & \Gamma(U \setminus K; F) \end{array}$$

and hence the fibers of the two horizontal maps coincide: for this reason, we will often also $\Gamma_K(X; F)$ by $\Gamma_K(U; F)$. Furthermore, if $S \subseteq X$ is locally closed, we define

$$\Gamma_S(X; F) := \Gamma_Z(U; j^* F)$$

where $S = U \cap Z$, with U open, Z closed and $j : U \hookrightarrow X$ the inclusion, but we will also use the notation $\Gamma_S(U; F)$.

Remark 5.8. The definition of the sections of a sheaf F on a compact K is functorial both in F and in K : since we have an obvious functor

$$\begin{array}{ccc} \mathcal{K}(X) & \longrightarrow & \text{Fun}(\Delta^1, \mathcal{U}(X)^{\text{op}}) \\ K & \longmapsto & (X \rightarrow X \setminus K) \end{array}$$

we get

$$\begin{array}{ccc} \mathcal{K}(X) \times \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \longrightarrow & \text{Fun}(\Delta^1, \mathcal{C}) \xrightarrow{\text{fib}} \mathcal{C} \\ \downarrow & \nearrow \circ & \\ \text{Fun}(\Delta^1, \mathcal{U}(X)^{\text{op}}) \times \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & & \end{array}$$

where the diagonal arrow is the composition of functors and the right horizontal arrow is given by taking the fiber of an arrow in \mathcal{C} , and so by adjunction we get the desired

$$\begin{array}{ccc} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \longrightarrow & \text{Fun}(\mathcal{K}(X), \mathcal{C}) \\ F & \longmapsto & (K \mapsto \Gamma_K(X; F)). \end{array}$$

Finally, by further composing with the functor ψ defined in Theorem 5.4, we get

$$(5.9) \quad \begin{array}{ccc} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \xrightarrow{\mathbb{D}_{\mathcal{C}}} & \text{Fun}(\mathcal{U}(X), \mathcal{C}) \\ F & \longmapsto & (U \mapsto \Gamma_c(U; F)). \end{array}$$

Theorem 5.10. The functor (5.9) restricts to an equivalence

$$\mathbb{D}_{\mathcal{C}} : \text{Shv}(X; \mathcal{C}) \xrightarrow{\cong} \text{CoShv}(X; \mathcal{C}).$$

Proof. We first prove that, if F is a sheaf, then $\mathbb{D}_{\mathcal{C}}(F)$ is a cosheaf. By virtue of Theorem 5.4, it suffices to prove that the functor

$$K \mapsto \Gamma_K(X; F)$$

is a \mathcal{K} -cosheaf.

- $\Gamma_{\emptyset}(X; F) \simeq 0$ since $F(X) \rightarrow F(X \setminus \emptyset)$ is an equivalence.

- Let $K, K' \in \mathcal{K}(X)$. The square

$$\begin{array}{ccc} \Gamma_{K \cap K'}(X; F) & \longrightarrow & \Gamma_K(X; F) \\ \downarrow & & \downarrow \\ \Gamma_{K'}(X; F) & \longrightarrow & \Gamma_{K \cup K'}(X; F) \end{array}$$

is the fiber of the obvious map between the pullback squares

$$\begin{array}{ccc} \Gamma(X; F) & \longrightarrow & \Gamma(X; F) \\ \downarrow & & \downarrow \\ \Gamma(X; F) & \longrightarrow & \Gamma(X; F) \end{array} \quad \begin{array}{ccc} \Gamma(X \setminus (K \cap K'); F) & \longrightarrow & \Gamma(X \setminus K; F) \\ \downarrow & & \downarrow \\ \Gamma(X \setminus K'; F) & \longrightarrow & \Gamma(X \setminus (K \cup K'); F), \end{array}$$

and so it is a pullback. Thus, since \mathcal{C} is stable, it's also a pushout.

- For any $K \in \mathcal{K}(X)$, we have a map of fiber sequences

$$\begin{array}{ccc} \Gamma_K(X; F) & \xrightarrow{a} & \varinjlim_{K' \in \mathcal{K}'} \Gamma_{K'}(X; F) \\ \downarrow & & \downarrow \\ \Gamma(X; F) & \xrightarrow{b} & \varinjlim_{K' \in \mathcal{K}'} \Gamma(X; F) \\ \downarrow & & \downarrow \\ \Gamma(X \setminus K; F) & \xrightarrow{c} & \varinjlim_{K' \in \mathcal{K}'} \Gamma(X \setminus K'; F). \end{array}$$

To prove that a is an equivalence, it suffices to prove that b and c are. But b is an equivalence because the poset $\{K' \in \mathcal{K}(X) \mid K \subseteq K'\}$ has a contractible nerve (since it is filtered) and c is an equivalence because $\{U \in \mathcal{U}(X) \mid U = X \setminus K' \text{ for some } K' \in \mathcal{K}'\}$ gives an open covering of $X \setminus K$.

We will now prove that $\mathbb{D}_{\text{cop}}^{\text{op}}$ is an inverse of \mathbb{D}_c . By symmetry, it suffices to show that it is a left inverse. Unraveling the definitions and using the equivalence of Theorem 5.4, this amounts to check that we have a cofiber sequence

$$(5.11) \quad \Gamma_c(X \setminus K; F) \longrightarrow \Gamma_c(X; F) \longrightarrow \Gamma(K; F)$$

natural in K and F .

First of all, we show that it suffices to prove that, for any fixed $K \in \mathcal{K}(X)$, $U \in \mathcal{U}(X)$ containing K and with compact closure, and $K' \in \mathcal{K}(X)$ containing U , the sequence

$$(5.12) \quad \Gamma_{K' \setminus U}(X; F) \longrightarrow \Gamma_{K'}(X; F) \longrightarrow \Gamma(U; F),$$

where the first morphism is given by the functoriality of sections supported on a compact and the second one is given by Remark 5.7, is a cofiber sequence. To see this, we start by noticing that the sequence is natural in K' and F , since both morphisms are canonically induced by the restrictions of F . Thus we can pass to the colimit ranging over all compacts $K' \supseteq U$ and get a fiber sequence

$$\varinjlim_{K' \supseteq U} \Gamma_{K' \setminus U}(X; F) \longrightarrow \Gamma_c(X; F) \longrightarrow \Gamma(U; F),$$

since the poset $\{K' \in \mathcal{K}(X) \mid K' \supseteq U\}$ is filtered (it is non-empty because it contains the closure of U) and the inclusion $\{K' \in \mathcal{K}(X) \mid K' \supseteq U\} \subseteq \mathcal{K}(X)$ is cofinal. Since any $K' \supseteq U$ is contained in $(K' \cup \overline{U}) \setminus U$, we get an equivalence

$$\varinjlim_{K' \supseteq U} \Gamma_{K' \setminus U}(X; F) \simeq \varinjlim_{\{K' \mid K' \cap U = \emptyset\}} \Gamma_{K'}(X; F),$$

and hence, adding everything up, we obtain a fiber sequence

$$\varinjlim_{\{K' \mid K' \cap U = \emptyset\}} \Gamma_{K'}(X; F) \longrightarrow \Gamma_c(X; F) \longrightarrow \Gamma(U; F),$$

which is natural in U , since the morphism $\Gamma_c(X; F) \rightarrow \Gamma(U; F)$ clearly is. Hence we can get the desired sequence (5.11) by passing to the colimit ranging over $P = \{U \in \mathcal{U}(X) \mid \overline{U} \in \mathcal{K}(X) \text{ and } U \supseteq K\}$ because $\Gamma(K; F) = \varinjlim_{U \in P} \Gamma(U; F)$ (since open subsets with compact closure form a basis of X), and because we have equivalences

$$\begin{aligned} \varinjlim_{U \in P} \varinjlim_{\{K' \mid K' \cap U = \emptyset\}} \Gamma_{K'}(X; F) &\simeq \varinjlim_{U \in P} \varinjlim_{\{K' \mid K' \cap U = \emptyset\}} \Gamma_{K'}(X; F) \\ &\simeq \varinjlim_{K' \subseteq X \setminus K} \Gamma_{K'}(X; F) \\ &\simeq \Gamma_c(X \setminus K; F) \end{aligned}$$

where the first one follows by [Lur09, Remark 4.2.3.9] and [Lur09, Corollary 4.2.3.10].

We are now left to show that (5.12) is a cofiber sequence. Consider the commutative diagram

$$\begin{array}{ccccc} \Gamma_{K' \setminus U}(X; F) & \longrightarrow & 0 & & \\ \downarrow & & \downarrow & & \\ \Gamma_{K'}(X; F) & \longrightarrow & Z & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow \\ \Gamma(X; F) & \longrightarrow & \Gamma(X \setminus (K' \setminus U); F) & \longrightarrow & \Gamma(X \setminus K'; F) \\ & & \downarrow & & \downarrow \\ & & \Gamma(U; F) & \longrightarrow & 0 \end{array}$$

where $Z := \text{fib}(\Gamma(X \setminus (K' \setminus U); F) \rightarrow \Gamma(X \setminus K'; F))$. Since the middle big horizontal rectangle is a pullback, it follows that also the left middle square is. But since the left vertical rectangle is pullback, then also the upper left square is. Then it suffices to prove that the composition

$$Z \rightarrow \Gamma(X \setminus (K' \setminus U); F) \rightarrow \Gamma(U; F)$$

is an equivalence, but this is clear since the lower right square is pullback because F is a sheaf. \square

5.2 Dualizability of spectral sheaves

Lemma 5.13. Let \mathcal{C} be a bicomplete ∞ -category where filtered colimits are exact. Then there exists a functor $\varphi : \mathrm{Shv}(\mathcal{K}(X); \mathcal{C}) \rightarrow \mathrm{Shv}_{\mathcal{K}}(X; \mathcal{C})$ satisfying the following properties

1. for any $F \in \mathrm{Shv}(\mathcal{K}(X); \mathcal{C})$ and $K \in \mathcal{K}(X)$, we have

$$\Gamma(K; \varphi F) \simeq \varinjlim_{K \Subset K'} \Gamma(K'; F),$$

2. φ preserves filtered colimits,
3. it is right inverse to the inclusion $\mathrm{Shv}_{\mathcal{K}}(X; \mathcal{C}) \hookrightarrow \mathrm{Shv}(\mathcal{K}(X); \mathcal{C})$.

Proof. Let M be the set whose elements are pairs (K, i) with $K \in \mathcal{K}(X)$ and $i = 0, 1$, where we define $(K, i) \leq (K', j)$ if $K' \subseteq K$ and $i = j$ or $K' \Subset K$ and $i < j$. It is easy to see that \leq actually defines a partial order on M . We have two functors

$$\begin{array}{ccc} \mathcal{K}(X)^{\mathrm{op}} & \xrightarrow{i_0} & M & \quad & \mathcal{K}(X)^{\mathrm{op}} & \xrightarrow{i_1} & M \\ K & \longmapsto & (K, 0) & & K & \longmapsto & (K, 1) \end{array}$$

and consider the cocontinuous functor

$$\varphi := (i_1)^*(i_0)_! : \mathrm{Fun}(\mathcal{K}(X)^{\mathrm{op}}, \mathcal{C}) \rightarrow \mathrm{Fun}(\mathcal{K}(X)^{\mathrm{op}}, \mathcal{C}).$$

Unravelling the definition, we see that

$$\Gamma(K; \varphi F) \simeq \varinjlim_{(K', 0) \rightarrow (K, 1)} \Gamma(K'; F) \simeq \varinjlim_{K \Subset K'} \Gamma(K'; F),$$

and so $\varphi F \simeq F$ whenever F is a \mathcal{K} -sheaf. Hence, to conclude the proof, it suffices to show that the essential image of $\varphi|_{\mathrm{Shv}(\mathcal{K}(X); \mathcal{C})}$ is contained in $\mathrm{Shv}_{\mathcal{K}}(X; \mathcal{C})$.

Indeed, for any $F \in \mathrm{Shv}(\mathcal{K}(X); \mathcal{C})$ we have

$$\begin{aligned} \varinjlim_{K \Subset K''} \Gamma(K''; \varphi F) &\simeq \varinjlim_{K \Subset K''} \varinjlim_{K'' \Subset K'} \Gamma(K'; F) \\ &\simeq \varinjlim_{K \Subset K'} \Gamma(K'; F) \\ &\simeq \Gamma(K; \varphi F), \end{aligned}$$

where the second equivalence holds by [Lur09, Remark 4.2.3.9] and [Lur09, Corollary 4.2.3.10] since for any K compact, the full subposet of $\mathcal{K}(X)^{\mathrm{op}}$ spanned by those K' such that $K \Subset K'$ is filtered, and we have

$$\bigcup_{\{K'' | K \Subset K''\}} \{K' | K'' \Subset K'\} = \{K' | K \Subset K'\}.$$

Moreover, since filtered colimits are exact in \mathcal{C} , φF belongs to $\mathrm{Shv}(\mathcal{K}(X); \mathcal{C})$, and so it is a \mathcal{K} -sheaf. \square

Lemma 5.13 is useful to describe conveniently pushforwards of sheaves along proper maps. Let $f : X \rightarrow Y$ be a proper continuous map between topological spaces, and let again \mathcal{C} be

a bicomplete ∞ -category where filtered colimits are exact. Since for any $K \in \mathcal{K}(Y)$, the preimage $f^{-1}(K)$ is compact, we obtain a functor

$$\begin{aligned} \text{Fun}(\mathcal{K}(X)^{\text{op}}, \mathcal{C}) &\xrightarrow{f_+} \text{Fun}(\mathcal{K}(Y)^{\text{op}}, \mathcal{C}) \\ F &\longmapsto (K \mapsto \Gamma(f^{-1}(K); F)). \end{aligned}$$

Notice that the restriction of f_+ to $\text{Shv}_{\mathcal{K}}(X; \mathcal{C})$ lands in $\text{Shv}(\mathcal{K}(Y); \mathcal{C})$, but a priori not in $\text{Shv}_{\mathcal{K}}(Y; \mathcal{C})$, therefore we define

$$f_*^{\mathcal{K}} : \text{Shv}_{\mathcal{K}}(X; \mathcal{C}) \rightarrow \text{Shv}_{\mathcal{K}}(Y; \mathcal{C})$$

as the composition of f_+ restricted to $\text{Shv}_{\mathcal{K}}(X; \mathcal{C})$ and $\varphi : \text{Shv}(\mathcal{K}(Y); \mathcal{C}) \rightarrow \text{Shv}_{\mathcal{K}}(Y; \mathcal{C})$.

Lemma 5.14. Let $f : X \rightarrow Y$ be a proper continuous map between topological spaces, and let \mathcal{C} be bicomplete ∞ -category where filtered colimits are exact. Then there is a commutative diagram

$$\begin{array}{ccc} \text{Shv}(X; \mathcal{C}) & \xrightarrow{\theta} & \text{Shv}_{\mathcal{K}}(X; \mathcal{C}) \\ \downarrow f_* & & \downarrow f_*^{\mathcal{K}} \\ \text{Shv}(Y; \mathcal{C}) & \xrightarrow{\theta} & \text{Shv}_{\mathcal{K}}(Y; \mathcal{C}), \end{array}$$

where θ is as in Theorem 5.4. In particular, f_* preserves filtered colimits, and when \mathcal{C} is stable it preserves all colimits.

Proof. For any $K \in \mathcal{K}(Y)$, we define

$$T = \{U \in \mathcal{U}(X) \mid \exists K' \ni K \text{ with } f^{-1}(K') \subseteq U\}.$$

Notice that if $V \in \mathcal{U}(Y)$ contains K , then there exists an open neighbourhood W of K with compact closure, and thus, since f is proper, $f^{-1}(V) \in T$. In particular we obtain a functor

$$\alpha : \{V \in \mathcal{U}(Y) \mid K \subseteq V\} \rightarrow T$$

which is obviously final. We have

$$\begin{aligned} \Gamma(K; f_*^{\mathcal{K}} \theta F) &\simeq \varinjlim_{K \in K'} \varinjlim_{f^{-1}(K') \subseteq U} \Gamma(U; F) \\ &\simeq \varinjlim_{U \in T} \Gamma(U; F) \\ &\simeq \varinjlim_{K \subseteq V} \Gamma(f^{-1}(V); F) \\ &\simeq \Gamma(K; \theta f_* F) \end{aligned}$$

where the second equivalence follows by [Lur09, Remark 4.2.3.9] and [Lur09, Corollary 4.2.3.10], and the third one since α is final. \square

Another straightforward application of Lemma 5.13 is the following theorem, that will be of crucial importance for what follows.

Theorem 5.15. The ∞ -category $\text{Shv}(X; \mathcal{S}p)$ is a strongly dualizable object in $\text{Cocont}_{\infty}^{st}$. In particular, for any $\mathcal{C} \in \text{Cocont}_{\infty}^{st}$, the canonical functor

$$\text{CoShv}(X; \mathcal{S}p) \otimes \mathcal{C} \rightarrow \text{CoShv}(X; \mathcal{C})$$

is an equivalence.

Proof. By Proposition 2.14, Theorem 5.4 and Remark 2.5, it suffices to show that $\mathrm{Shv}_{\mathcal{K}}(X; \mathbb{S}\mathrm{p})$ is a retract in $\mathrm{Cocont}_{\infty}^{st}$ of a compactly generated ∞ -category. We see that $\mathrm{Shv}(\mathcal{K}(X); \mathbb{S}\mathrm{p})$ is clearly compactly generated, as it is equivalent to $\mathrm{Fun}_{lex}(\mathcal{K}(X)^{\mathrm{op}}, \mathbb{S}\mathrm{p})$. The proof is then concluded by noticing that the inclusion $\mathrm{Shv}_{\mathcal{K}}(X; \mathbb{S}\mathrm{p}) \subseteq \mathrm{Shv}(\mathcal{K}(X); \mathbb{S}\mathrm{p})$ and $\varphi : \mathrm{Shv}(\mathcal{K}(X); \mathbb{S}\mathrm{p}) \rightarrow \mathrm{Shv}_{\mathcal{K}}(X; \mathbb{S}\mathrm{p})$ are exact and preserve filtered colimits since filtered colimits in $\mathbb{S}\mathrm{p}$ are exact, and thus preserves all colimits since $\mathbb{S}\mathrm{p}$ is stable. One then concludes the proof by identifying $\mathrm{CoShv}(X; \mathbb{S}\mathrm{p})$ with the dual of $\mathrm{Shv}(X; \mathbb{S}\mathrm{p})$, via Lemma 2.21. \square

Corollary 5.16. There is a unique equivalence

$$(5.17) \quad \eta : \mathrm{Shv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} \rightarrow \mathrm{Shv}(X; \mathcal{C}).$$

making the diagram

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\eta} & \mathrm{Shv}(X; \mathcal{C}) \\ \downarrow \mathbb{D}_{\mathbb{S}\mathrm{p}} \otimes \mathcal{C} & & \downarrow \mathbb{D}_{\mathcal{C}} \\ \mathrm{CoShv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \longrightarrow & \mathrm{CoShv}(X; \mathcal{C}) \end{array}$$

Proof. The functor η is obtained by composing

$$\mathrm{Shv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} \xrightarrow{\mathbb{D}_{\mathbb{S}\mathrm{p}} \otimes \mathcal{C}} \mathrm{CoShv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} \xrightarrow{\simeq} \mathrm{CoShv}(X; \mathcal{C}) \xrightarrow{\mathbb{D}_{\mathcal{C}}^{-1}} \mathrm{Shv}(X; \mathcal{C}).$$

where the middle map is the one in Theorem 5.15. More concretely, for any $F \in \mathrm{Shv}(X; \mathbb{S}\mathrm{p})$ and $M \in \mathcal{C}$, we have $\eta(F \boxtimes^{st} M) \simeq \mathbb{D}_{\mathcal{C}}^{-1}(M \circ \mathbb{D}_{\mathbb{S}\mathrm{p}} F)$, where M on the right-hand side denotes the essentially unique colimit preserving functor $\mathbb{S}\mathrm{p} \rightarrow \mathcal{C}$ corresponding to M . \square

5.3 The pullback $f_{\mathcal{C}}^*$

Proposition 5.18. Let $f : X \rightarrow Y$ be a proper map, and denote by $f_{*}^{\mathcal{C}} : \mathrm{Shv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(Y; \mathcal{C})$ the pushforward. Then we have a commutative square

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\eta} & \mathrm{Shv}(X; \mathcal{C}) \\ \downarrow f_{*}^{\mathbb{S}\mathrm{p}} \otimes \mathcal{C} & & \downarrow f_{*}^{\mathcal{C}} \\ \mathrm{Shv}(Y; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\eta} & \mathrm{Shv}(Y; \mathcal{C}). \end{array}$$

In particular, $f_{*}^{\mathcal{C}}$ admits a left adjoint which is identified through η with $f_{\mathbb{S}\mathrm{p}}^* \otimes \mathcal{C}$.

Proof. By a slight abuse of notation, denote as $f_{*}^{\mathcal{C}} : \mathrm{CoShv}(X; \mathcal{C}) \rightarrow \mathrm{CoShv}(Y; \mathcal{C})$ the pushforward for cosheaves (i.e. $(f_{*}^{\mathcal{C}\mathrm{op}})^{\mathrm{op}}$). We have that the square

$$\begin{array}{ccc} \mathrm{CoShv}(X; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \mathrm{CoShv}(X; \mathcal{C}) \\ \downarrow f_{*}^{\mathbb{S}\mathrm{p}} \otimes \mathcal{C} & & \downarrow f_{*}^{\mathcal{C}} \\ \mathrm{CoShv}(Y; \mathbb{S}\mathrm{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \mathrm{CoShv}(Y; \mathcal{C}) \end{array}$$

commutes since the horizontal arrows can be modelled by a composition of functors, and thus we are only left to show that the square

$$\begin{array}{ccc} \mathrm{Shv}(X; \mathcal{C}) & \xrightarrow{\mathbb{D}} & \mathrm{CoShv}(X; \mathcal{C}) \\ \downarrow f_{*} & & \downarrow f_{*} \\ \mathrm{Shv}(Y; \mathcal{C}) & \xrightarrow{\mathbb{D}} & \mathrm{CoShv}(Y; \mathcal{C}) \end{array}$$

commutes.

First of all, we show that there exists a natural transformation $f_*\mathbb{D} \rightarrow \mathbb{D}f_*$ even when f is not proper. Fix $V \in \mathcal{U}(Y)$ and a compact $K \subseteq f^{-1}(V)$, so that $f(K)$ is a compact subset of V . For any $F \in \text{Shv}(X; \mathcal{C})$, the commutative triangle

$$\begin{array}{ccc} \Gamma(X \setminus K; F) & \xrightarrow{\quad} & \Gamma(X \setminus f^{-1}(f(K)); F) \\ & \swarrow \quad \searrow & \\ & \Gamma(X; F) & \end{array}$$

provides a morphism

$$\Gamma_K(X; F) \rightarrow \Gamma_{f(K)}(Y; f_*F) \rightarrow \Gamma_c(V; f_*F).$$

Since all morphisms are induced by the restrictions of F , the resulting map is natural in K and V , and hence gives rise to the desired transformation as K varies. Furthermore, when f is proper, each compact $K \subseteq X$ is contained in the compact $f^{-1}f(K)$, so by cofinality we obtain an equivalence

$$\lim_{K \subseteq f^{-1}(V)} \Gamma_K(X; F) \simeq \lim_{C \subseteq V} \Gamma_c(Y; f_*F)$$

where C varies over the compact subsets of V , and thus we may conclude. \square

Lemma 5.19. Let $j : U \hookrightarrow X$ be an open immersion, and denote by $j_{\mathcal{C}}^* : \text{Shv}(X; \mathcal{C}) \rightarrow \text{Shv}(U; \mathcal{C})$ the restriction. Then we have a commutative square

$$\begin{array}{ccc} \text{Shv}(X; \mathcal{S}\text{p}) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(X; \mathcal{C}) \\ \downarrow j_{\mathcal{S}\text{p}}^* \otimes \mathcal{C} & & \downarrow j_{\mathcal{C}}^* \\ \text{Shv}(U; \mathcal{S}\text{p}) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(U; \mathcal{C}). \end{array}$$

In particular, $j_{\mathcal{C}}^*$ admits a left adjoint which is identified through η with $j_{\#}^{\mathcal{S}\text{p}} \otimes \mathcal{C}$.

Proof. By an abuse of notation, denote by $j_{\mathcal{C}}^* : \text{CoShv}(X; \mathcal{C}) \rightarrow \text{CoShv}(U; \mathcal{C})$ the restriction for cosheaves. Again we see that the square

$$\begin{array}{ccc} \text{CoShv}(X; \mathcal{S}\text{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \text{CoShv}(X; \mathcal{C}) \\ \downarrow j_{\mathcal{S}\text{p}}^* \otimes \mathcal{C} & & \downarrow j_{\mathcal{C}}^* \\ \text{CoShv}(U; \mathcal{S}\text{p}) \otimes \mathcal{C} & \xrightarrow{\simeq} & \text{CoShv}(U; \mathcal{C}) \end{array}$$

is obviously commutative, and thus we only have to show that the square

$$\begin{array}{ccc} \text{Shv}(X; \mathcal{C}) & \xrightarrow{\mathbb{D}} & \text{CoShv}(X; \mathcal{C}) \\ \downarrow j^* & & \downarrow j^* \\ \text{Shv}(U; \mathcal{C}) & \xrightarrow{\mathbb{D}} & \text{CoShv}(U; \mathcal{C}) \end{array}$$

commutes. But this follows immediately because by Remark 5.7, and by observing that $\Gamma(V; j^*F) \simeq \Gamma(V; F)$ for any $V \subseteq U$, we have that

$$\Gamma_c(V; j^*F) \simeq \Gamma_c(V; F)$$

functorially on F and V . \square

Corollary 5.20. Let $f : X \rightarrow Y$ be any continuous map. Then the pushforward $f_*^{\mathcal{C}} : \text{Shv}(X; \mathcal{C}) \rightarrow \text{Shv}(Y; \mathcal{C})$ admits a left adjoint $f_{\mathcal{C}}^*$ such that there exists a commutative square

$$\begin{array}{ccc} \text{Shv}(Y; \mathcal{S}p) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(Y; \mathcal{C}) \\ \downarrow f_{\mathcal{S}p}^* \otimes \mathcal{C} & & \downarrow f_{\mathcal{C}}^* \\ \text{Shv}(X; \mathcal{S}p) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(X; \mathcal{C}). \end{array}$$

Proof. Using the factorization 5.1, this follows immediately by Proposition 5.18 and Lemma 5.19. \square

Theorem 5.21. Let $i_{\mathcal{C}} : \text{Shv}(X; \mathcal{C}) \hookrightarrow \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C})$ be the inclusion functor, and let $L_{\mathcal{S}p} : \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{S}p) \rightarrow \text{Shv}(X; \mathcal{S}p)$ be the left adjoint of $i_{\mathcal{S}p}$. Denote by $L_{\mathcal{C}}$ the composition

$$\begin{array}{ccc} \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{S}p) \otimes \mathcal{C} & \xrightarrow{L_{\mathcal{S}p} \otimes \mathcal{C}} & \text{Shv}(X; \mathcal{S}p) \otimes \mathcal{C} \\ \simeq \uparrow & & \downarrow \eta \\ \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C}) & \xrightarrow{L_{\mathcal{C}}} & \text{Shv}(X; \mathcal{C}). \end{array}$$

Then $L_{\mathcal{C}}$ is left adjoint to $i_{\mathcal{C}}$.

Proof. From the proof of Theorem 2.10 and Remark 2.13, we see that we may write any $F \in \text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C})$ as a colimit

$$F \simeq \varinjlim_{M \rightarrow \Gamma(U; F)} U \boxtimes^{st} M$$

where the indexing category is Grothendieck construction of the functor

$$(U, M) \mapsto \text{Hom}_{\mathcal{C}}(M, \Gamma(U; F)).$$

By definition, the presheaf of spectra $U \boxtimes^{st} M$ represents the functor taking sections at U , and thus we get an equivalence

$$\mathbb{S}_U = L_{\mathcal{S}p}(U \boxtimes \mathbb{S}) \simeq j_{\#}^{\mathcal{S}p} a_{\mathcal{S}p}^* \mathbb{S}$$

which is natural in U , where $j : U \hookrightarrow X$ denotes an open inclusion, $a : U \rightarrow *$ the unique map. We have equivalences

$$\begin{aligned} \text{Hom}_{\text{Shv}(X; \mathcal{C})}(L_{\mathcal{C}}F, G) &\simeq \varinjlim_{M \rightarrow \Gamma(U; F)} \text{Hom}_{\text{Shv}(X; \mathcal{S}p) \otimes \mathcal{C}}(j_{\#}^{\mathcal{S}p} a_{\mathcal{S}p}^* \mathbb{S} \boxtimes^{st} M, \eta^{-1}G) \\ &\simeq \varinjlim_{M \rightarrow \Gamma(U; F)} \text{Hom}_{\text{Shv}(X; \mathcal{C})}(j_{\#}^{\mathcal{C}} a_{\mathcal{C}}^* M, G) \\ &\simeq \varinjlim_{M \rightarrow \Gamma(U; F)} \text{Hom}_{\mathcal{C}}(M, \Gamma(U; G)) \\ &\simeq \varinjlim_{M \rightarrow \Gamma(U; F)} \text{Hom}_{\text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C})}(U \boxtimes M, i^{\mathcal{C}}G) \\ &\simeq \text{Hom}_{\text{Fun}(\mathcal{U}(X)^{\text{op}}, \mathcal{C})}(F, i^{\mathcal{C}}G) \end{aligned}$$

where the first equivalence follows by the observations above, and the second is a consequence of Lemma 5.19 and Corollary 5.20. Since all identifications are functorial on F and G , we obtain the thesis. \square

Remark 5.22. After the results in this section, we are now able to extend everything we have proven so far for sheaves with presentable coefficients to sheaves with values in a stable bicomplete ∞ -category. The only detail we have to handle with more care is the functor (2.27): since the tensor product of two stable bicomplete ∞ -categories is not again complete in general, (2.27) will now take values in $\mathrm{Shv}(X; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{D})$. Nevertheless, when \mathcal{C} has a monoidal structure such that its tensor $\otimes_{\mathcal{C}}$ preserves colimits in both variables, the composition of (2.27) with the obvious functor

$$\mathrm{Shv}(X; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

still gives the usual monoidal structure on $\mathrm{Shv}(X; \mathcal{C})$. As a consequence, we see that the equivalences in Corollary 2.31 and Corollary 3.26 still hold in $\mathrm{Shv}(X; \mathcal{S}\mathrm{p}) \otimes (\mathcal{C} \otimes \mathcal{D})$ and in $\mathrm{Shv}(X; \mathcal{C})$ when (2.27) is exchanged with the tensor product in $\mathrm{Shv}(X; \mathcal{C})$ described above.

6 Six functor formalism

In this section, we will define the operations $f_!$ and $f^!$, and prove all the usual formulas that one expects for these functors. A first attempt towards these results for sheaves of spectra can be found in the paper [BL96], even though it almost totally lacks proofs. A proof of the proper base change theorem with unstable coefficients was provided in [Lur09, Corollary 7.3.1.18], and later extended to spectral coefficients in [Hai21]: our only contribute to this theorem is to explain how to extend it to general stable bicomplete coefficients. A novelty of the approach presented in this section is our expression of the formulas involving tensor products, such as projection or Künneth formula. Here we do not a priori require the coefficients of our sheaves to be equipped with a monoidal structure, but rely instead on the tensor product of stable cocomplete ∞ -categories. The advantage of this perspective is that, using the observations in Remark 5.22, it clarifies how to obtain all these formulas for a general stable bicomplete ∞ -category equipped with a closed symmetric monoidal structure. At the end of the section, building up on the our discussion of Section 3 related to shape theory, we will explain how to prove the formula

$$f^!(\mathbf{1}) \otimes f^* \simeq f^!$$

for any map f which induce locally contractible geometric morphisms.

6.1 The formulas for $f_!^{\mathcal{C}}$

Throughout this section, \mathcal{C} is going to be any stable and bicomplete ∞ -category. For any continuous map $f : X \rightarrow Y$, consider the functor

$$(f_*^{\mathcal{C}^{\mathrm{op}}})^{\mathrm{op}} : \mathrm{Shv}(X; \mathcal{C}^{\mathrm{op}})^{\mathrm{op}} = \mathrm{CoShv}(X; \mathcal{C}) \rightarrow \mathrm{Shv}(Y; \mathcal{C}^{\mathrm{op}})^{\mathrm{op}} = \mathrm{CoShv}(Y; \mathcal{C}).$$

Since taking opposite categories switches left with right adjoints, the functor

$$(f_{\mathrm{op}}^*)^{\mathrm{op}} : \mathrm{Shv}(Y; \mathcal{C}^{\mathrm{op}})^{\mathrm{op}} = \mathrm{CoShv}(Y; \mathcal{C}) \rightarrow \mathrm{Shv}(X; \mathcal{C}^{\mathrm{op}})^{\mathrm{op}} = \mathrm{CoShv}(X; \mathcal{C})$$

is right adjoint to $(f_*^{\mathcal{C}^{\mathrm{op}}})^{\mathrm{op}}$. Hence, by Theorem 5.10, we get a corresponding adjunction at the level of sheaves

$$\begin{array}{ccc} & f_!^{\mathcal{C}} & \\ \curvearrowright & & \curvearrowleft \\ \mathrm{Shv}(X; \mathcal{C}) & \perp & \mathrm{Shv}(Y; \mathcal{C}). \\ \curvearrowleft & & \curvearrowright \\ & f_e^! & \end{array}$$

where $f_!^{\mathcal{C}}$ and $f_!^{\mathcal{C}}$ are defined to be the unique functors fitting in commutative squares

$$\begin{array}{ccc} \mathrm{Shv}(Y; \mathcal{C}) & \xrightarrow{f_!^{\mathcal{C}}} & \mathrm{Shv}(Y; \mathcal{C}) \\ \mathbb{D} \downarrow & & \downarrow \mathbb{D} \\ \mathrm{CoShv}(X; \mathcal{C}) & \xrightarrow{(f_*^{\mathcal{C}op})^{\mathrm{op}}} & \mathrm{CoShv}(Y; \mathcal{C}) \end{array} \qquad \begin{array}{ccc} \mathrm{Shv}(Y; \mathcal{C}) & \xrightarrow{f_!^{\mathcal{C}}} & \mathrm{Shv}(Y; \mathcal{C}) \\ \mathbb{D} \downarrow & & \downarrow \mathbb{D} \\ \mathrm{CoShv}(Y; \mathcal{C}) & \xrightarrow{(f_{\mathcal{C}op}^*)^{\mathrm{op}}} & \mathrm{CoShv}(X; \mathcal{C}). \end{array}$$

Definition 6.1. The functors $f_!^{\mathcal{C}}$ and $f_!^{\mathcal{C}}$ constructed as above are called respectively *push-forward with proper support* and *exceptional pullback*. Unless it is required from the context, we will often omit to include \mathcal{C} in subscripts or superscripts in our notation.

More concretely, $f_!$ is the functor uniquely determined by the formula

$$\Gamma_c(U; f_!F) = \Gamma_c(f^{-1}(U); F)$$

for all $U \in \mathcal{U}(Y)$. In particular, when $a : X \rightarrow *$ is the unique map, we get

$$a_!F \simeq \Gamma_c(X; F).$$

In different geometric contexts, when one deals with six functor formalisms, it is common to define the shriek operations making use of appropriate compactifications of maps analogous to (5.1). This approach, however, makes it a bit tricky to verify that $f_!$ behaves well under compositions. An advantage of our definition of $f_!$ is that its functoriality is more or less immediate, as illustrated by the following lemma.

Lemma 6.2. Let LCH be the category of locally compact Hausdorff spaces. Then, for any \mathcal{C} stable and bicomplete, there is a functor

$$\mathrm{Shv}_!(-, \mathcal{C}) : \mathrm{LCH} \rightarrow \mathrm{Cocont}_{\infty}^{st}$$

whose values on an object X is given by $\mathrm{Shv}(X; \mathcal{C})$, and on a morphism f is given by $f_!^{\mathcal{C}}$.

Proof. We first observe that there is a functor

$$\mathrm{Shv}_*(-, \mathcal{C}) : \mathrm{LCH} \rightarrow \mathrm{Cocont}_{\infty}^{st}$$

whose value on a morphism $f : X \rightarrow Y$ is given by the pushforward $f_*^{\mathcal{C}}$. By definition, $f_*^{\mathcal{C}}$ is given by precomposing with the functor $f^{-1} : \mathcal{U}(Y) \rightarrow \mathcal{U}(X)$. Thus, the functoriality of $\mathrm{Shv}_*(-, \mathcal{C})$ descends from the functoriality of internal-homs in Cat_{∞} , which is straightforward in the model of quasi-categories. By passing to opposite categories, we obtain a similar functor $\mathrm{CoShv}_*(-, \mathcal{C})$.

Let J be the interval object for the Joyal model structure (see [Cis19, Definition 3.3.3]), and consider the monomorphism of simplicial sets $Ob(\mathrm{LCH}) \hookrightarrow \mathrm{LCH}$. By Theorem 5.10, we have a functor

$$Ob(\mathrm{LCH}) \times J \cup \mathrm{LCH} \times \{1\} \xrightarrow{\mathbb{D}_{\mathcal{C} \cup \mathrm{CoShv}_*(-, \mathcal{C})}} \mathrm{Cat}_{\infty}.$$

which admits a lifting

$$\begin{array}{ccc} Ob(\mathrm{LCH}) \times J \cup \mathrm{LCH} \times \{1\} & \longrightarrow & \mathrm{Cat}_{\infty} \\ \downarrow & \searrow \text{dotted} & \\ \mathrm{LCH} \times J & & \end{array}$$

since the vertical arrow is a categorical anodyne extension. Thus we get the desired functor by restricting the dotted arrow to $\mathrm{LCH} \times \{0\}$. \square

Lemma 6.3. There exists a natural transformation $f_! \rightarrow f_*$ which is an equivalence when f is proper.

Proof. By Verdier duality, it suffices to construct a natural transformation between $\mathbb{D}f_! \rightarrow \mathbb{D}f_*$ and show it is an equivalence when f is proper, which is the content of the second part of the proof of Proposition 5.18. \square

Corollary 6.4. Let $i : Z \hookrightarrow X$ be a closed immersion. Then the functor $i_{\text{Sp}}^! : \text{Shv}(X; \text{Sp}) \rightarrow \text{Shv}(Z; \text{Sp})$ coincides with the one defined in Corollary 4.4.

Proof. This follows immediately by the previous lemma, since any closed immersion is proper. \square

Lemma 6.5. Let $j : U \hookrightarrow X$ be an open immersion. Then we have $j_! \dashv j^*$ or equivalently $j^* \simeq j^!$.

Proof. By the proof of Lemma 5.19, we have a natural equivalence $\mathbb{D}j^* \simeq \mathbb{D}j^!$, and thus we may conclude. \square

Remark 6.6. A useful consequence of Lemma 6.2, Lemma 6.5, Lemma 6.3 and (5.1) is that the functor $f_!^{\mathcal{C}}$ is uniquely determined by the fact that it is right adjoint to $f_{\mathcal{C}}^*$ when f is proper and left adjoint to $f_{\mathcal{C}}^*$ when f is an open immersion.

From Remark 6.6 we also see that $f_!$ behaves well with respect to tensor products in $\text{Cocont}_{\infty}^{st}$.

Corollary 6.7. Let $f : X \rightarrow Y$ be any continuous map. Then there is a commutative square

$$\begin{array}{ccc} \text{Shv}(X; \text{Sp}) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(X; \mathcal{C}) \\ \downarrow f_!^{\text{Sp} \otimes \mathcal{C}} & & \downarrow f_!^{\mathcal{C}} \\ \text{Shv}(Y; \text{Sp}) \otimes \mathcal{C} & \xrightarrow{\eta} & \text{Shv}(Y; \mathcal{C}). \end{array}$$

Proof. This follows immediately by Remark 6.6, Proposition 5.18 and Lemma 5.19. \square

Remark 6.8. Let $j : U \hookrightarrow X$ be the inclusion of any open subset with compact closure. Then a simple computation involving Lemma 6.3 and the closure of U shows that, for any sheaf F on X , one has

$$\Gamma(U; F) \simeq \Gamma_c(X; j_* j^* F).$$

Since U has compact closure, any closed subset of U can be written as the intersection of U with some compact subset of X , and thus we obtain

$$\begin{aligned} \Gamma_c(X; j_* j^* F) &\simeq \varinjlim_{K \subseteq X} \Gamma_{U \cap K}(U; F) \\ &\simeq \varinjlim_{\overline{S} \subseteq U} \Gamma_S(U; F), \end{aligned}$$

where the last colimit ranges over all closed subsets of U .

Proposition 6.9 (Base change). For every given pullback square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & X \\ g' \downarrow & \lrcorner & \downarrow g \\ Y' & \xrightarrow{f} & Y \end{array}$$

of topological spaces, there is a natural equivalence

$$f^* g! \simeq g'_! f'^*$$

and, by transposition, also

$$g^! f_* \simeq f'_* g'^!$$

Proof. First of all we observe that, since all functors appearing are colimit preserving, by Corollary 6.7 it suffices to prove the proposition in the case of sheaves of spectra.

By Remark 6.6, it suffices to prove the statement in the separate cases when g is an open immersion or a proper map. The open immersion case follows immediately by Lemma 6.5 and Lemma 3.25, while the proper case follows by [Hai21, Corollary 3.2]. For the reader's convenience, let us briefly summarize the strategy of [Hai21].

By [Lur09, Corollary 7.3.1.18] we know that the statement of the theorem is true for sheaves of spaces. Since $\mathcal{S}p$ is compactly generated, by Proposition 2.16 we

$$\begin{aligned} \mathrm{Shv}(X; \mathcal{S}p) &\simeq \mathrm{Fun}_*(\mathcal{S}p^{\mathrm{op}}, \mathrm{Shv}(X)) \\ &\simeq \mathrm{Fun}_{lex}((\mathcal{S}p^\omega)^{\mathrm{op}}, \mathrm{Shv}(X)) \end{aligned}$$

where $\mathcal{S}p^\omega$ denotes the full subcategory of $\mathcal{S}p$ spanned by all compact objects. One checks easily that, for any continuous map $h : W \rightarrow T$ of topological spaces, there is a commutative square

$$\begin{array}{ccc} \mathrm{Shv}(W; \mathcal{S}p) & \xrightarrow{\simeq} & \mathrm{Fun}_{lex}((\mathcal{S}p^\omega)^{\mathrm{op}}, \mathrm{Shv}(W)) \\ \downarrow h_*^{\mathcal{S}p} & & \downarrow h_*^{\mathcal{S}^\circ} \circ (-) \\ \mathrm{Shv}(T; \mathcal{S}p) & \xrightarrow{\simeq} & \mathrm{Fun}_{lex}((\mathcal{S}p^\omega)^{\mathrm{op}}, \mathrm{Shv}(T)), \end{array}$$

where the right hand vertical square denotes a post-composition with the pushforward $h_*^{\mathcal{S}^\circ}$ of sheaves of spaces. Reasoning analogously to Remark 2.2, we see that the functor

$$\mathrm{Fun}_{lex}((\mathcal{S}p^\omega)^{\mathrm{op}}, -)$$

preserves adjunctions between left exact functors, and so we get a similar commutative square involving the pullbacks $h_{\mathcal{S}p}^*$ and $h_{\mathcal{S}^\circ}^*$. Hence, we obtain base change for spectral sheaves by applying $\mathrm{Fun}_{lex}((\mathcal{S}p^\omega)^{\mathrm{op}}, -)$ to Lurie's nonabelian proper base change in [Lur09, Corollary 7.3.1.18]. \square

Remark 6.10. It follows from Remark 2.17 and the definition of the tensor (2.27) that for any topological space X and any bicomplete stable ∞ -category \mathcal{C} , $\mathrm{Shv}(X; \mathcal{C})$ is tensored over $\mathrm{Shv}(X; \mathcal{S}p)$. When there is no possibility of confusion we will denote by $F \otimes G$ the image through the canonical variablewise colimit preserving functor

$$\mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(X; \mathcal{S}p) \rightarrow \mathrm{Shv}(X; \mathcal{C})$$

of a pair (F, G) , and, when $G \in \mathrm{Shv}(X; \mathcal{S}p)$ by $\underline{\mathrm{Hom}}_X(G, F)$ the image of any $F \in \mathrm{Shv}(X; \mathcal{C})$ through the right adjoint of $- \otimes G$.

Let $f : X \rightarrow X'$ and $g : Y \rightarrow Y'$ be morphisms of topological spaces, and let $f \times g : X \times Y \rightarrow X' \times Y'$ be the induced map on the products. For any two stable bicomplete ∞ -categories \mathcal{C} and \mathcal{D} , the variable-wise colimit preserving functor

$$\mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(Y; \mathcal{D}) \xrightarrow{f_! \times g_!} \mathrm{Shv}(X'; \mathcal{C}) \times \mathrm{Shv}(Y'; \mathcal{D}) \xrightarrow{\boxtimes} \mathrm{Shv}(X' \times Y'; \mathcal{S}p) \otimes (\mathcal{C} \otimes \mathcal{D})$$

induces a functor

$$f_! \boxtimes g_! : \mathrm{Shv}(X \times Y; \mathcal{S}p) \otimes (\mathcal{C} \otimes \mathcal{D}) \rightarrow \mathrm{Shv}(X' \times Y'; \mathcal{S}p) \otimes (\mathcal{C} \otimes \mathcal{D}).$$

Proposition 6.11 (Künneth formula). We have a natural equivalence

$$f_! \boxtimes g_! \simeq (f \times g)_!$$

Notice that, since X and Y are locally compact, $f_! \boxtimes g_!$ is the image of the pair $(f_!, g_!)$ through the bifunctor given by the tensor product of cocomplete ∞ -categories.

Proof. By the factorization (5.1), it will suffice to prove the statement when both f , g and $f \times g$ are either open immersions or proper maps. By uniqueness of adjoints and Remark 2.2, both cases will then follow by Proposition 2.30. More precisely, in the open immersions case we use that $f_! \boxtimes g_!$ is left adjoint to $f^* \boxtimes g^*$, while in the proper case we use that $f_! \boxtimes g_!$ is right adjoint to $f^* \boxtimes g^*$, and by Proposition 2.30 we always have an equivalence $f^* \boxtimes g^* \simeq (f \times g)^*$. \square

Proposition 6.12 (Projection formula). Let $f : X \rightarrow Y$ be a morphism of topological spaces, and let \mathcal{C} and \mathcal{D} be two stable and bicomplete ∞ -categories. Then, for any $F \in \text{Shv}(X; \mathcal{C})$ and $G, H \in \text{Shv}(Y; \mathcal{D})$, we have a canonical equivalence

$$f_! F \otimes G \simeq f_!(F \otimes f^* G)$$

or, when $\mathcal{C} = \mathcal{D}$ has a closed symmetric monoidal structure, by transposition

$$f_* \underline{\text{Hom}}_X(F, f^! G) \simeq \underline{\text{Hom}}_Y(f_! F, G)$$

and

$$f^! \underline{\text{Hom}}_Y(G, H) \simeq \underline{\text{Hom}}_X(f^* G, f^! H).$$

Proof. Exactly as for the Corollary 3.26, one may deduce this result from Proposition 6.9 and Proposition 6.11 applied to $(f \times \text{id}_Y)_!$. \square

Corollary 6.13. Let $k : Z \rightarrow X$ be the inclusion of a locally closed subset of X , $F \in \text{Shv}(X; \mathcal{C})$ with \mathcal{C} stable and bicomplete. Let \mathcal{D} be another stable bicomplete ∞ -category, $M \in \mathcal{D}$ any object. Then we have a canonical equivalence

$$k_! k^*(F \otimes M) \simeq F \otimes M_Z.$$

Moreover, when \mathcal{C} has a closed symmetric monoidal structure, we have

$$k_! k^* F \simeq F \otimes_{\mathcal{C}} \mathbb{1}_Z$$

or equivalently

$$k_* k^! F \simeq \underline{\text{Hom}}_X(\mathbb{1}_Z, F),$$

where $\mathbb{1}$ and $\mathbb{1}_Z$ are the monoidal units of \mathcal{C} .

Proof. This is an immediate consequence of Proposition (6.12) and Corollary (2.31). \square

Corollary 6.14. For every given pullback square

$$\begin{array}{ccc} X' & \xrightarrow{f'} & X \\ g' \downarrow & \lrcorner & \downarrow g \\ Y' & \xrightarrow{f} & Y \end{array}$$

of topological spaces where f and f' are shape submersions, there is a natural equivalence

$$g_! f'_\# \simeq f_\# g'_!$$

or equivalently

$$f'^* g^! \simeq g'^! f^*.$$

Proof. First of all we construct a natural transformation. Applying $g'_!$ to the unit of the adjunction $f'_\# \dashv f'^*$ and using base change we obtain

$$g'_! \rightarrow g'_! f'^* f'_\# \simeq f^* g'_! f'_\#$$

and hence by transposition the desired transformation. By factorization (5.1) and Corollary 4.10, we are only left to prove the case of a pullback square

$$\begin{array}{ccc} X \times Y' & \xrightarrow{\text{id}_X \times f} & X \times Y \\ a \times \text{id}_{Y'} \downarrow & \lrcorner & \downarrow a \times \text{id}_Y \\ Y' & \xrightarrow{f} & Y \end{array}$$

where X is compact and $a : X \rightarrow *$ is the unique map, but this follows by Proposition 6.11 and by the functoriality of the tensor product of cocomplete ∞ -categories as follows

$$\begin{aligned} (a \times \text{id}_Y)_!(\text{id}_X \times f)_\# &\simeq (a_! \otimes (\text{id}_Y)_!)((\text{id}_X)_\# \otimes f_\#) \\ &\simeq a_! f_\# \\ &\simeq ((\text{id}_*)_ \# a_!) \otimes (f_\# (\text{id}_{Y'})_!) \\ &\simeq ((\text{id}_*)_ \# \otimes f_\#)(a_! \otimes (\text{id}_{Y'})_!). \end{aligned}$$

□

Remark 6.15. Using (5.2), Lemma 3.25, Corollary 3.26, Corollary 5.20, Proposition 6.9, Proposition 6.12, Corollary 6.14, and [Man22, Proposition A.5.10], one sees that the functor

$$\begin{array}{ccc} \text{LCH}^{\text{op}} & \longrightarrow & \text{Cat}_\infty \\ X & \longmapsto & \text{Shv}(X; \mathcal{C}) \\ (f : X \rightarrow Y) & \longmapsto & f_{\mathcal{C}}^* \end{array}$$

defines a six functor formalism in the sense of [Man22, Definition A.5.9].

6.2 $f_{\mathcal{C}}^!$ when f is a locally contractible geometric morphism

Let $f : X \rightarrow Y$ be a continuous map inducing an essential geometric morphism (see Definition 3.12). In particular, we have an adjunction $f_\# \dashv f^*$ for \mathcal{C}^{op} -valued sheaves, and thus, after passing to opposite categories and applying Theorem 5.10, we get an adjunction

$$\begin{array}{ccc} & f^! & \\ & \curvearrowright & \\ \text{Shv}(Y; \mathcal{C}) & \perp & \text{Shv}(X; \mathcal{C}) \\ & \curvearrowleft & \\ & f_\circ & \end{array}$$

We now want to show that, in the special case when f induces a locally contractible geometric morphism (see Definition 3.12), one the exceptional pullback coincides with the usual pullback up to a twist. In this way, we will vastly generalize the classical formula relating the $f^!$ and f^* when f is a topological submersion.

Proposition 6.16. Let $f : X \rightarrow Y$ be a continuous map inducing a locally contractible geometric morphism, and let \mathcal{C} and \mathcal{D} be two stable and bicomplete ∞ -categories. Then, for any $F \in \mathrm{Shv}(Y; \mathcal{C})$ and $G \in \mathrm{Shv}(Y; \mathcal{D})$, we have a natural equivalence

$$f_{\mathcal{C}}^! F \otimes f_{\mathcal{D}}^* G \simeq f_{\mathcal{C} \otimes \mathcal{D}}^! (F \otimes G)$$

of functors $\mathrm{Shv}(X; \mathcal{C}) \times \mathrm{Shv}(Y; \mathcal{D}) \rightarrow \mathrm{Shv}(X \times Y; \mathcal{C} \otimes \mathcal{D})$. Equivalently, when $\mathcal{C} = \mathcal{D}$ has a closed symmetric monoidal structure, a canonical equivalence for any $K \in \mathrm{Shv}(X; \mathcal{C})$

$$\underline{\mathrm{Hom}}_Y(F, f_{\circ} K) \rightarrow f_* \underline{\mathrm{Hom}}_X(f^! F, K).$$

Proof. For any $F \in \mathrm{Shv}(Y; \mathcal{C})$, $G \in \mathrm{Shv}(Y; \mathcal{D})$ and $H \in \mathrm{Shv}(Y; \mathcal{C} \otimes \mathcal{D})$, we have the map

$$\mathrm{counit} \otimes G : f_! f^! F \otimes G \rightarrow F \otimes G$$

which by adjunction and Proposition 6.12 gives the desired natural transformation

$$f^!(-) \otimes f^*(-) \rightarrow f^!(- \otimes -).$$

Since all functors appearing are cocontinuous, by Remark 6.10 it suffices to prove the invertibility of the map when $\mathcal{C} = \mathcal{D} = \mathrm{Sp}$, and after evaluation on pairs of the type (F, \mathbb{S}_U) , where $F \in \mathrm{Shv}(Y; \mathrm{Sp})$, \mathbb{S} denotes the sphere spectrum, and $j : U \hookrightarrow Y$ is an open subset. Keeping the same notations as in Remark 3.13, we see that by Theorem 5.10 we have an equivalence

$$f_{\mathcal{C}}^! j_!^{\mathcal{C}} \simeq (j')_!^{\mathcal{C}} (f')_e^!.$$

Thus, we get

$$\begin{aligned} f^!(F) \otimes f^*(\mathbb{S}_U) &\simeq f^!(\mathbb{S}_Y) \otimes \mathbb{S}_{f^{-1}(U)} \\ &\simeq j_!^!(f')^! j^*(F) \\ &\simeq f^! j_! j^*(F) \\ &\simeq f^!(F \otimes \mathbb{S}_U), \end{aligned}$$

where the second equivalence follows by Corollary 6.13 and Lemma 6.5, the third by Remark 3.13 and the fourth again by Corollary 6.13. \square

Remark 6.17. We thank Marc Hoyois for pointing out that a result similar to Proposition 6.16 can be found in [Ver65, Section 5]. By adapting our proof of Corollary 3.26, one can actually deduce the theorem in [Ver65] from Proposition 6.16.

Proposition 6.18. Let $f : X \rightarrow Y$ be a topological submersion of fiber dimension n , \mathcal{C} be any stable bicomplete ∞ -category. Then:

- (i) if f is a trivial submersion, then there is a canonical equivalence

$$f^! \simeq \Sigma^n f^*;$$

- (ii) $f^!$ preserves locally constant sheaves;

- (iii) for all $F \in \mathrm{Shv}(Y; \mathcal{C})$ there is a canonical equivalence

$$f^! \mathbb{S}_Y \otimes f^* F \simeq f^! F$$

or equivalently by adjunction, for every $G \in \mathrm{Shv}(X; \mathcal{C})$

$$f_{\#} G \simeq f_!(G \otimes f^! \mathbb{S}_Y).$$

Proof. Since f^* preserves locally constant sheaves, we see that (ii) follows from (i). Hence, we now prove (i). By assumption, f is a projection $p : X \times \mathbb{R}^n \rightarrow X$. By Proposition 6.16, it suffices to show that $p^! \mathbb{S}_X \simeq \Sigma^n \mathbb{S}_{X \times \mathbb{R}^n}$. Since p induces a locally contractible geometric morphism, we have that $p^!$ preserves colimits. Hence, by Proposition 6.11 and the uniqueness of adjoints, we may assume that p is the unique map $a : \mathbb{R}^n \rightarrow *$.

We first show that $p^! M$ is locally constant for each $M \in \mathbb{S}p$. By a standard argument (see [Hai20, Proposition 3.1] or [Vol22, Proposition 3.14]) it suffices to show that for any $U \subseteq \mathbb{R}^n$ euclidean chart, the restriction

$$\Gamma(\mathbb{R}^n; p^! M) \rightarrow \Gamma(U; p^! M)$$

is an equivalence. By adjunction, we know that for any open V there is an equivalence

$$\Gamma(V; p^! M) \simeq \underline{\mathbf{Hom}}_{\mathbb{S}p}(\Gamma_c(V; \mathbb{S}_{\mathbb{R}^n}), M).$$

Thus, it will suffice to prove that the canonical map

$$\Gamma_c(U; \mathbb{S}_{\mathbb{R}^n}) \rightarrow \Gamma_c(\mathbb{R}^n; \mathbb{S}_{\mathbb{R}^n})$$

is invertible. Since any compact subset of a vector space is contained in some compact closed ball, it suffices to show that, for any K compact closed ball in U , the map

$$\Gamma_K(U; \mathbb{S}_{\mathbb{R}^n}) \rightarrow \Gamma_K(\mathbb{R}^n; \mathbb{S}_{\mathbb{R}^n})$$

is invertible. But this follows by homotopy invariance of the shape, since the inclusion $U \setminus K \hookrightarrow \mathbb{R}^n \setminus K$ is a homotopy equivalence. In particular, we also see that the canonical map

$$\Gamma_{\{0\}}(\mathbb{R}; \mathbb{S}_{\mathbb{R}}) \rightarrow \Gamma_c(\mathbb{R}; \mathbb{S}_{\mathbb{R}})$$

is an equivalence.

To conclude the proof of (ii) it suffices to check that the global sections of $p^! M$ are equivalent to $\Sigma^n M$. We start with the case $n = 1$. Arguing as above, this amounts to proving that we have an equivalence

$$\Gamma_{\{0\}}(\mathbb{R}; \mathbb{S}_{\mathbb{R}}) \simeq \Omega \mathbb{S}.$$

By Corollary 3.4, we may identify $\Gamma_{\{0\}}(\mathbb{R}; \mathbb{S}_{\mathbb{R}})$ with the fiber of the diagonal map

$$\Gamma(\mathbb{R}; \mathbb{S}_{\mathbb{R}}) \rightarrow \Gamma(\mathbb{R}; \mathbb{S}_{\mathbb{R}}) \oplus \Gamma(\mathbb{R}; \mathbb{S}_{\mathbb{R}}).$$

It is then a straightforward exercise in pasting pushouts to verify that, for any object A of a stable ∞ -category, the fiber of the diagonal map is equivalent to ΩA . For $n > 1$, we see that by Proposition 6.11 we have

$$\Gamma_c(\mathbb{R}^n; \mathbb{S}_{\mathbb{R}^n}) \simeq \Gamma_c(\mathbb{R}; \mathbb{S}_{\mathbb{R}}) \otimes \cdots \otimes \Gamma_c(\mathbb{R}; \mathbb{S}_{\mathbb{R}}),$$

and thus what we wanted.

To conclude, we observe that (iii) follows by Proposition 6.16 and the fact that any locally constant sheaf whose stalks are invertible spectra is an invertible object with respect to the smash product of sheaves of spectra. \square

7 Relative Atiyah Duality

Recall that, if X is a compact smooth manifold, then the *Atiyah duality* states that $\Sigma_{\mp}^{\infty} X$ is strongly dual to the *Thom spectrum* associated to the virtual vector bundle given by the inverse of the tangent bundle of X , which is denoted by $\mathrm{Th}(-TX)$. In this section we will revisit Atiyah duality using the six functor formalism, in the spirit of motivic homotopy theory.

By what we have achieved up to now, we can see very easily that whenever $f : X \rightarrow Y$ is a proper map inducing a locally contractible geometric morphism, $f_{\sharp}(\mathbb{S}_X) \in \mathrm{Shv}(Y; \mathbb{S}\mathrm{p})$ is strongly dualizable with dual $f_!(\mathbb{S}_X)$. The question is then about identifying $f_!(\mathbb{S}_X)$ with some sheaf theoretic construction reminiscent of Thom spectra, at least in more geometric situations (e.g. when X and Y are manifolds). We will provide a natural transformation

$$\mathrm{Th} : \mathrm{Vect}(X) \rightarrow \mathrm{Pic}(\mathrm{Shv}(X; \mathbb{S}\mathrm{p}))$$

where the left-hand side denotes the ∞ -groupoid of real vector bundles over X , the right-hand side is the ∞ -groupoid of invertible sheaves of spectra, and X ranges through all paracompact Hausdorff spaces. By observing that the left-hand side is a constant sheaf on the site of all paracompact Hausdorff spaces, the natural transformation above will automatically be induced by a functor

$$\mathrm{Vect}(\ast) \rightarrow \mathrm{Pic}(\mathrm{Shv}(X; \mathbb{S}\mathrm{p}))$$

given by one-point compactification. After that, we will show that for any vector bundle E , $\mathrm{Th}(E)$ can be described through six operations analogously to the definition in motivic homotopy theory. The advantage of our perspective on the definition of $\mathrm{Th}(E)$ is that it makes the verification of all its expected properties, such as compatibility with pulling back vector bundles or short exact sequences, essentially trivial. We will then conclude the section by showing that for a submersion f between smooth manifolds, there is an equivalence

$$f^!(\mathbb{S}_Y) \simeq \mathrm{Th}(-T_f)$$

and hence obtaining a generalization of Atiyah duality.

7.1 Thom spaces and the J-homomorphism

Let Vect be the ∞ -groupoid obtained by taking the coherent nerve of the topological groupoid whose objects are finite dimensional real vector spaces, and morphisms are spaces of linear isomorphisms between. Here we consider $\mathrm{GL}_n(\mathbb{R})$ equipped with the usual topology of a manifold (or equivalently the compact-open topology). As usual, one may equip Vect with a symmetric monoidal structure given by sum, and hence we may regard it as an object of $\mathrm{CMon}(\mathbb{S})$. Notice that we have an homotopy equivalence $\mathrm{Vect}_{\mathbb{R}} \simeq \coprod_{n \in \mathbb{N}} \mathrm{BGL}_n(\mathbb{R})$.

Let PH be the category of paracompact Hausdorff spaces with continuous maps between them. Throughout this section, we will assume all topological spaces appearing to be in PH . Recall also that PH can be equipped with a Grothendieck topology with the usual open coverings. Let \mathbf{U} be a universe such that PH is \mathbf{U} -small, and denote by \mathcal{S}' the ∞ -category of \mathbf{U} -small spaces. Hence, the inclusion of $\mathrm{Shv}(\mathrm{PH}; \mathcal{S}') \hookrightarrow \mathrm{Fun}(\mathrm{PH}^{\mathrm{op}}, \mathcal{S}')$ admits a left adjoint.

Definition 7.1. We denote by

$$\mathrm{Vect}_{\mathrm{PH}} : \mathrm{PH}^{\mathrm{op}} \rightarrow \mathrm{CMon}(\mathcal{S}')$$

the sheafification of the constant presheaf with value Vect .

The next lemma will justify our choice of notation.

Lemma 7.2. There have an equivalence

$$\mathrm{Vect}_{\mathrm{PH}}(X) \simeq \mathrm{Sing}(\mathrm{Map}(X, |\mathrm{Vect}|)),$$

which is natural on $X \in \mathrm{PH}$, and Map denotes the mapping space equipped with the compact-open topology. In particular, $\pi_0(\mathrm{Vect}_{\mathrm{PH}}(X))$ is in bijection with the set of equivalence classes of real vector bundles over X .

Proof. This is a consequence of [Lur09, Theorem 7.1.0.1]. The last part of the statement is standard: see for example [Hat17, Theorem 1.16]. \square

Consider the map

$$\mathrm{GL}_n(\mathbb{R}) \rightarrow \mathrm{Hom}_{\mathfrak{S}^*}(\mathbb{S}^n, \mathbb{S}^n) \simeq \Omega^n \mathbb{S}^n$$

given by the functoriality of the one-point compactification. The restriction of the map above to O_n is what's known as the *J-homomorphism*. We thus obtain a functor

$$\mathrm{Vect} \rightarrow \mathfrak{S}_*^{\simeq}$$

that at the level of objects sends a finite dimensional real vector space V to its one-point compactification \overline{V} . Moreover, this is easily seen to be symmetric monoidal, where the right hand side is equipped with the smash product. By post composing with Σ^∞ , we get

$$\mathrm{Vect} \rightarrow \mathrm{Sp}^{\simeq}$$

and since \overline{V} is homoemorphic to a sphere, this factors as

$$\mathrm{Vect} \rightarrow \mathrm{Pic}(\mathrm{Sp}),$$

where $\mathrm{Pic}(\mathrm{Sp})$ is the *Picard ∞ -groupoid* of Sp , i.e. the full subcategory of Sp^{\simeq} spanned by the objects which are invertible with respect to the smash product of spectra.

Let $f : X \rightarrow Y$ be any map of topological spaces. Since the pullback functor $f^* : \mathrm{Shv}(Y; \mathfrak{Sp}) \rightarrow \mathrm{Shv}(X; \mathfrak{Sp})$ is monoidal, we obtain consequently a functor

$$\begin{aligned} \mathrm{PH}^{\mathrm{op}} &\longrightarrow \mathrm{CMon}(\mathfrak{S}) \\ X &\longmapsto \mathrm{Pic}(\mathrm{Shv}(X; \mathfrak{Sp})) \end{aligned}$$

whose global sections are $\mathrm{Pic}(\mathrm{Shv}(*; \mathfrak{Sp})) \simeq \mathrm{Pic}(\mathrm{Sp})$. Since taking spectrum objects and Picard ∞ -groupoids both commute with limits, such functor is a sheaf. Thus, Lemma 7.2 yields a map

$$(7.3) \quad \mathrm{Th} : \mathrm{Vect}_{\mathrm{PH}} \rightarrow \mathrm{Pic}(\mathrm{Shv}(-; \mathfrak{Sp})).$$

in $\mathrm{Shv}(\mathrm{PH}; \mathrm{CMon}(\mathfrak{S}))$.

Definition 7.4. For each vector bundle $E \rightarrow X$, we define $\mathrm{Th}(E)$ to be the image of E through the morphism (7.3).

Remark 7.5. Denote by $\mathrm{CMon}^{gp}(\mathfrak{S})$ the full subcategory of $\mathrm{CMon}(\mathfrak{S})$ spanned by those commutative monoids M such that $\pi_0(M)$ is a group. The inclusion $\mathrm{CMon}^{gp}(\mathfrak{S}) \hookrightarrow \mathrm{CMon}(\mathfrak{S})$ admits a left adjoint, denoted by $(-)^{gp}$, which is called the *group completion*. Since

$$\pi_0(\mathrm{Pic}(\mathrm{Shv}(X; \mathfrak{Sp})))$$

is a group, for any $X \in \text{PH}$ we obtain a morphism

$$\text{Vect}_{\text{PH}}(X)^{gp} \rightarrow \text{Pic}(\text{Shv}(X; \mathbb{S}p)).$$

In particular we see that, for any vector bundle $E \rightarrow X$ it makes sense to define $\text{Th}(-E)$ as the tensor inverse of $\text{Th}(E)$, where $-E$ denotes the inverse of the class of E in $\pi_0(\text{Vect}_{\text{PH}}(X))^{gp}$. Moreover, for any split exact sequence

$$0 \rightarrow E \rightarrow V \rightarrow E' \rightarrow 0$$

we get an equivalence

$$\text{Th}(V) \simeq \text{Th}(E) \otimes \text{Th}(E').$$

Our next goal is to describe $\text{Th}(E)$ in terms of the six operations. Let $p : E \rightarrow X$ be a real vector bundle over X , and denote by $s : X \hookrightarrow E$ its zero section and by $j : E^\times \hookrightarrow E$ its open complement. Consider the sheaf

$$p_{\#}\text{cofib}(j_{\#}j^*\mathbb{S}_E \rightarrow \mathbb{S}_E) \in \text{Shv}(X; \mathbb{S}p)$$

where the morphism $j_{\#}j^*\mathbb{S}_E \rightarrow \mathbb{S}_E$ is the counit. Notice that $p_{\#}$ exists since any vector bundle is obviously a shape submersion, so indeed the definition above makes sense. Notice also that, using Theorem 4.6, one has $p_{\#}\text{cofib}(j_{\#}j^*\mathbb{S}_E \rightarrow \mathbb{S}_E) \simeq p_{\#}s!\mathbb{S}_X$.

We will need the following lemma.

Lemma 7.6. Let

$$\begin{array}{ccc} X' & \longrightarrow & X \\ \downarrow & & \downarrow f \\ Y' & \longrightarrow & Y \end{array}$$

be any pullback square of locally compact Hausdorff spaces. Then the corresponding diagram

$$\begin{array}{ccc} \text{Shv}(X') & \longrightarrow & \text{Shv}(X) \\ \downarrow & & \downarrow \\ \text{Shv}(Y') & \longrightarrow & \text{Shv}(Y) \end{array}$$

is a pullback of ∞ -topoi.

Proof. Using the factorization (5.1), we just need to prove separately the case of an open immersion and a proper map. The first is treated in [Lur09, Remark 6.3.5.8], while the second follows by the proof of [Lur09, Corollary 7.3.1.18]. \square

Proposition 7.7. The assignment $E \mapsto p_{\#}\text{cofib}(j_{\#}j^*\mathbb{S}_E \rightarrow \mathbb{S}_E) \in \text{Shv}(X; \mathbb{S}p)$ defines a natural transformation

$$\text{Vect}_{\text{PH}} \rightarrow \text{Pic}(\text{Shv}(-; \mathbb{S}p))$$

which is naturally equivalent to (7.3).

Proof. First of all, we show that the Thom spectrum induces a natural transformation

$$(7.8) \quad \text{Vect}_{\text{PH}} \rightarrow \text{Shv}(-; \mathbb{S}p)$$

of presheaves of ∞ -categories. Let $p : E \rightarrow X$ be a vector bundle, $p^\times : E^\times \rightarrow X$ be the induced map on the complement of the zero section. Since one can write $\text{Th}(E)$ as

$\text{cofib}(p_{\sharp}^{\times} \mathbb{S}_{E^{\times}} \rightarrow p_{\sharp} \mathbb{S}_E)$, it will suffice to show that the associations $E \mapsto p_{\sharp} \mathbb{S}_E$ and $E \mapsto p_{\sharp}^{\times} \mathbb{S}_{E^{\times}}$ induce natural transformations. Consider a pullback square in Top

$$\begin{array}{ccc} f^* E & \xrightarrow{f'} & E \\ p' \downarrow & \lrcorner & \downarrow p \\ X' & \xrightarrow{f} & X \end{array}$$

where p is a vector bundle. By Lemma 7.6

$$\begin{array}{ccc} \text{Shv}(f^* E) & \xrightarrow{f'} & \text{Shv}(E) \\ p' \downarrow & & \downarrow p \\ \text{Shv}(X') & \xrightarrow{f} & \text{Shv}(X) \end{array}$$

is a pullback square in $\mathcal{T}\text{op}$. Similarly one has that the square

$$\begin{array}{ccc} \text{Shv}(f^* E^{\times}) & \xrightarrow{f'} & \text{Shv}(E^{\times}) \\ p'^{\times} \downarrow & & \downarrow p^{\times} \\ \text{Shv}(X') & \xrightarrow{f} & \text{Shv}(X) \end{array}$$

is a pullback. Hence we have two natural transformations

$$\text{Vect}_{\text{PH}} \rightarrow \mathcal{T}\text{op}/_{\text{Shv}(-)}$$

given respectively by sending a vector bundle $E \rightarrow X$ to $\text{Shv}(E) \rightarrow \text{Shv}(X)$ and to $\text{Shv}(E^{\times}) \rightarrow \text{Shv}(X)$, and thus, by further composing with the relative shape, by Remark 3.7 we obtain lax natural transformations

$$\text{Vect}_{\text{PH}} \rightarrow \text{Pro}(\text{Shv}(-))$$

which factors as

$$\text{Vect}_{\text{PH}} \rightarrow \text{Shv}(-)$$

since any shape submersion induces a locally contractible geometric morphism by Corollary 3.26. Furthermore, by Lemma 3.25 and [Hau20, Theorem 3.22], we see that these are actually natural transformations, and thus, composing with

$$\text{Shv}(-) \rightarrow \text{Shv}(-; \mathcal{S}\text{p})$$

we get the natural transformation (7.8).

We now prove that (7.8) is symmetric monoidal and that it factors through $\text{Pic}(\text{Shv}(-; \mathcal{S}\text{p}))$. Since Vect_{PH} is the constant sheaf associated to Vect , for any sheaf $F \in \text{Shv}(\text{PH}; \text{CMon}(\mathcal{S}))$, we have a commutative diagram

$$\begin{array}{ccc} \text{Hom}(\text{Vect}_{\text{PH}}, \text{Pic}(F)) & \xrightarrow{\cong} & \text{Hom}_{\text{CMon}}(\text{Vect}, \text{Pic}(F(*))) \\ \downarrow & & \downarrow \\ \text{Hom}(\text{Vect}_{\text{PH}}, F) & \xrightarrow{\cong} & \text{Hom}_{\text{CMon}}(\text{Vect}, F(*)) \\ \downarrow & & \downarrow \\ \text{Hom}(\text{Vect}_{\text{PH}}, F) & \xrightarrow{\cong} & \text{Hom}_{\mathcal{S}}(\text{Vect}, F(*)) \end{array}$$

where the horizontal arrows are induced by taking global sections, the upper vertical arrows by the natural transformation $\text{Pic}(F) \rightarrow F$ and the lower vertical arrows by the forgetful functor $\text{CMon} \rightarrow \mathcal{S}$. By Corollary 7.9 we know that, after taking global sections, (7.8) factors through $\text{Pic}(\mathcal{S}\text{p})$, and thus we may conclude by a simple diagram chase. \square

Corollary 7.9. Assume that X is essential. Then, for any vector bundle E over X , $a_{\#}\mathrm{Th}(E)$ is equivalent to the Thom spectrum of E as classically defined.

Proof. Let $b : E \rightarrow *$ and $c : E^{\times} \rightarrow *$ be the unique maps. By Corollary 3.19, we have equivalences

$$\begin{aligned} a_{\#}\mathrm{Th}(E) &\simeq b_{\#}\mathrm{cofib}(j_{\#}j^*\mathbb{S}_E \rightarrow \mathbb{S}_E) \\ &\simeq \mathrm{cofib}(c_{\#}\mathbb{S}_{E^{\times}} \rightarrow b_{\#}\mathbb{S}_E) \\ &\simeq \mathrm{cofib}(\Sigma_{+}^{\infty}E^{\times} \rightarrow \Sigma_{+}^{\infty}E) \end{aligned}$$

and the spectrum on the last line coincides with the usual Thom spectrum of E . \square

Corollary 7.10. Let $p : E \rightarrow X$ be a real vector bundle over X , and denote $s : X \hookrightarrow E$ its zero section. Then $\mathrm{Th}(E)$ is invertible with inverse given by $s^!\mathbb{S}_E$.

Proof. By definition, we already know that $\mathrm{Th}(E)$ is invertible, and thus to compute its inverse we just need to look at its dual $\underline{\mathrm{Hom}}_X(\mathrm{Th}(E), \mathbb{S}_X)$. Then we may conclude by Corollary 3.26 and Proposition 6.12 since

$$\begin{aligned} \underline{\mathrm{Hom}}_X(p_{\#}s^!\mathbb{S}_X, \mathbb{S}_X) &\simeq p_*\underline{\mathrm{Hom}}_E(s^!\mathbb{S}_X, \mathbb{S}_E) \\ &\simeq p_*s_*s^!\mathbb{S}_E \\ &\simeq s^!\mathbb{S}_E. \end{aligned}$$

\square

7.2 Relative Atiyah duality

Theorem 7.11. Let $f : X \rightarrow Y$ be a submersion between smooth manifolds. Then we have an equivalence $f^!\mathbb{S}_Y \simeq \mathrm{Th}(T_f)$, where T_f is the relative tangent bundle of f , defined by the short exact sequence of vector bundles

$$(7.12) \quad 0 \rightarrow T_f \rightarrow TX \rightarrow f^*TY \rightarrow 0.$$

Proof. First of all we prove the case when f is the unique map $a : X \rightarrow *$ and X is a smooth manifold. Choose a closed embedding $i : X \hookrightarrow \mathbb{R}^n$, and let $a' : \mathbb{R}^n \rightarrow *$ be the unique map. By Proposition 6.18, we have $a'^!\mathbb{S} \simeq \Sigma^n\mathbb{S}_{\mathbb{R}^n}$ and thus, since $T\mathbb{R}^n$ is a trivial vector bundle of fiber dimension n , we have $\mathrm{Th}(T\mathbb{R}^n) \simeq \Sigma^n\mathbb{S}_{\mathbb{R}^n} \simeq a'^!\mathbb{S}$. Let $p : N_i \rightarrow X$ be the conormal bundle of the embedding i , defined by the short exact sequence

$$(7.13) \quad 0 \rightarrow TX \rightarrow i^*T\mathbb{R}^n \rightarrow N_i \rightarrow 0,$$

$s : X \hookrightarrow N_i$ its zero section. Let $k : U \hookrightarrow N_i$ be a tubular neighbourhood of X in \mathbb{R}^n . Thus, we get a commutative triangle

$$\begin{array}{ccc} & X & \\ \tilde{s} \swarrow & & \searrow i \\ U & \xrightarrow{g} & \mathbb{R}^n \end{array}$$

where g is an open immersion and \tilde{s} is a closed immersion. Hence we get an equivalence

$$i^!\mathbb{S}_{\mathbb{R}^n} \simeq \tilde{s}^!\mathbb{S}_U.$$

Then, by Proposition 6.18 and (7.13) we have

$$\begin{aligned}
a^! \mathbb{S} &\simeq a^! i^! \mathbb{S} \\
&\simeq a^! \mathbb{S} \otimes i^* \mathrm{Th}(T\mathbb{R}^n) \\
&\simeq \mathrm{Th}(N_i)^{-1} \otimes \mathrm{Th}(i^* T\mathbb{R}^n) \\
&\simeq \mathrm{Th}(TX).
\end{aligned}$$

Suppose now that $f : X \rightarrow Y$ is any submersion between smooth manifolds, $a : X \rightarrow *$ and $b : Y \rightarrow *$ be the unique maps. Then, by Proposition 6.18 and Remark 7.5, we have

$$\begin{aligned}
\mathrm{Th}(T_f) &\simeq \mathrm{Th}(TX) \otimes \mathrm{Th}(f^* TY)^{-1} \\
&\simeq a^! \mathbb{S} \otimes (f^* b^! \mathbb{S})^{-1} \\
&\simeq a^! \mathbb{S} \otimes (f^! b^! \mathbb{S})^{-1} \otimes f^! \mathbb{S}_Y \\
&\simeq f^! \mathbb{S}_Y
\end{aligned}$$

and thus we can conclude. □

Corollary 7.14 (Relative Atiyah Duality). Let $f : X \rightarrow Y$ be a proper map inducing a locally contractible geometric morphism. Then $f_{\sharp} \mathbb{S}_X \in \mathrm{Shv}(Y; \mathbb{S}p)$ is strongly dualizable with dual $f_! \mathbb{S}_X$. Moreover, if X and Y are smooth manifolds and f is a proper submersion, then $f_{\sharp} \mathbb{S}_X$ is strongly dualizable with dual $f_{\sharp} \mathrm{Th}(-T_f)$.

Proof. Since f is proper, by Corollary 3.26 and Proposition 6.12, we have, functorially on $F \in \mathrm{Shv}(Y; \mathbb{S}p)$

$$\begin{aligned}
\underline{\mathrm{Hom}}_Y(f_{\sharp} \mathbb{S}_X, F) &\simeq f_* \underline{\mathrm{Hom}}_X(\mathbb{S}_X, f^* F) \\
&\simeq f_* f^* F \\
&\simeq f_! f^* F \\
&\simeq f_! \mathbb{S}_X \otimes F.
\end{aligned}$$

In particular, when f is a submersion of smooth manifolds, by Proposition 6.18 and the previous theorem, we have $f_! \mathbb{S}_X \simeq f_{\sharp} \mathrm{Th}(-T_f)$. □

Remark 7.15. Let X be a smooth manifold, $a : X \rightarrow *$ the unique map. By specializing the previous corollary to a and Corollary 7.9, we see that we recover the classical Atiyah duality.

References

- [AC21] Avraham Aizenbud and Shachar Carmeli. Relative de rham theory on nash manifolds. *arXiv preprint arXiv:2105.12417*, 2021.
- [Aok20] Ko Aoki. Tensor triangular geometry of filtered objects and sheaves. *arXiv preprint arXiv:2001.00319*, 2020.
- [BL96] Jonathan Block and Andrey Lazarev. Homotopy theory and generalized duality for spectral sheaves. *International Mathematics Research Notices*, 1996(20):983–996, 1996.
- [CD19] Denis-Charles Cisinski and Frédéric Déglise. *Triangulated categories of mixed motives*, volume 6. Springer, 2019.
- [Cis19] Denis-Charles Cisinski. *Higher categories and homotopical algebra*, volume 180. Cambridge University Press, 2019.
- [Cla11] Dustin Clausen. p-adic j-homomorphisms and a product formula. *arXiv preprint arXiv:1110.5851*, 2011.
- [Hai20] Peter J Haine. The homotopy-invariance of constructible sheaves of spaces. *arXiv preprint arXiv:2010.06473*, 2020.
- [Hai21] Peter J. Haine. From nonabelian basechange to basechange with coefficients. *arXiv preprint arXiv:2108.03545*, 2021.
- [Hat17] Allen Hatcher. Vector bundles and K-theory. *available online at <https://pi.math.cornell.edu/~hatcher/VBKT/VB.pdf>*, 2017.
- [Hau20] Rune Haugseng. A fibrational mate correspondence for ∞ -categories. *arXiv preprint arXiv:2011.08808*, 2020.
- [Hoy] M Hovey. K-theory of dualizable categories (after a. efimov), 2018. *Available from the author’s webpage.*
- [Hoy16] Marc Hovey. Is the ∞ -topos $\mathrm{sh}(x)$ hypercomplete whenever x is a cw complex? MathOverflow answer, 2016. <https://mathoverflow.net/q/247061>.
- [Hoy17] Marc Hovey. The six operations in equivariant motivic homotopy theory. *Advances in Mathematics*, 305:197–279, 2017.
- [Hoy18] Marc Hovey. Higher galois theory. *Journal of Pure and Applied Algebra*, 222(7):1859–1877, 2018.
- [Jin19] Xin Jin. A hamiltonian $\coprod_n \mathrm{BO}(n)$ -action, stratified morse theory and the j-homomorphism. *arXiv preprint arXiv:1902.06708*, 2019.
- [Jin20] Xin Jin. Microlocal sheaf categories and the j-homomorphism. *arXiv preprint arXiv:2004.14270*, 2020.
- [Joh02] Peter T Johnstone. *Sketches of an Elephant: A Topos Theory Compendium: Volume 2*. Oxford University Press, 2002.
- [JT17] Xin Jin and David Treumann. Brane structures in microlocal sheaf theory. *arXiv preprint arXiv:1704.04291*, 2017.

- [Kha19] Adeel Khan. The morel–voevodsky localization theorem in spectral algebraic geometry. *Geometry & Topology*, 23(7):3647–3685, 2019.
- [KS90] Masaki Kashiwara and Pierre Schapira. *Sheaves on Manifolds: With a Short History. Les débuts de la théorie des faisceaux. By Christian Houzel*, volume 292. Springer Science & Business Media, 1990.
- [KS06] Masaki Kashiwara and Pierre Schapira. Categories and sheaves. *Grundlehren der Math. Wiss*, 332, 2006.
- [Lur09] Jacob Lurie. *Higher topos theory*. Princeton University Press, 2009. <https://www.math.ias.edu/~lurie/papers/HTT.pdf>.
- [Lur16] Jacob Lurie. Spectral algebraic geometry. 2016. <https://www.math.ias.edu/~lurie/papers/SAG-rootfile.pdf>.
- [Lur17] Jacob Lurie. *Higher algebra*. 2017. <https://www.math.ias.edu/~lurie/papers/HA.pdf>.
- [Man22] Lucas Mann. A p-Adic 6-Functor Formalism in Rigid-Analytic Geometry *arXiv preprint arXiv:2206.02022*, 2022.
- [MS06] J Peter May and Johann Sigurdsson. *Parametrized homotopy theory*. Number 132. American Mathematical Soc., 2006.
- [Pet14] Dan Petersen. What (if anything) unifies stable homotopy theory and grothendieck’s six functors formalism? MathOverflow question, 2014.
- [PT22] Mauro Porta and Jean-Baptiste Teyssier. Topological exodromy with coefficients. *arXiv preprint arXiv:2211.05004*, 2022.
- [Rez19] Charles Rezk. What’s an example of an ∞ -topos not equivalent to sheaves on a grothendieck site? MathOverflow, 2019.
- [RS18] Marco Robalo and Pierre Schapira. A lemma for microlocal sheaf theory in the ∞ -categorical setting. *Publications of the Research Institute for Mathematical Sciences*, 54(2):379–391, 2018.
- [Ver65] Jean-Louis Verdier. Dualité dans la cohomologie des espaces localement compacts. *Séminaire Bourbaki*, 9:337–349, 1965.
- [Vol22] Marco Volpe. Verdier duality on conically smooth stratified spaces. *arXiv preprint arXiv:2206.02728*, 2022.