

Generalized semiconfined harmonic oscillator model with a position-dependent effective mass

C. Quesne*

*Physique Nucléaire Théorique et Physique Mathématique, Université Libre de Bruxelles,
Campus de la Plaine CP229, Boulevard du Triomphe, B-1050 Brussels, Belgium*

Abstract

By using a point canonical transformation starting from the constant-mass Schrödinger equation for the isotonic potential, it is shown that a semiconfined harmonic oscillator model with a position-dependent mass in the BenDaniel-Duke setting and the same spectrum as the standard harmonic oscillator can be easily constructed and extended to a semiconfined shifted harmonic oscillator, which could result from the presence of a uniform gravitational field. A further generalization is proposed by considering a m -dependent position-dependent mass for $0 < m < 2$ and deriving the associated semiconfined potential. This results in a family of position-dependent mass and potential pairs, to which the original pair belongs as it corresponds to $m = 1$. Finally, the potential that would result from a general von Roos kinetic energy operator is presented and the examples of the Zhu-Kroemer and Mustafa-Mazharimousavi settings are briefly discussed.

Keywords: Schrödinger equation; position-dependent mass; harmonic oscillator; point canonical transformation

PACS Nos.: 03.65.Fd, 03.65.Ge

*e-mail: Christiane.Quesne@ulb.be

1 Introduction

There is much interest in the Schrödinger equation wherein the constant mass is replaced by a position-dependent mass (PDM), because the latter has many applications in problems occurring in several fields of physics [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]. As it has been shown [13] that the PDM Schrödinger equation is equivalent to two other unconventional Schrödinger equations, namely the Schrödinger equation resulting from the use of deformed commutation relations [14, 15, 16], as well as that in curved space [17, 18, 19], this has reinforced the interest in its study.

As a consequence, much attention has been devoted to finding exact solutions of PDM Schrödinger equations because they may provide a conceptual understanding of some physical phenomena, as well as a testing ground for some approximation schemes. The generation of PDM and potential pairs leading to such exact solutions has been achieved by various methods (see, e.g., [20] and references quoted therein). One of the most powerful techniques for such a purpose consists in applying a point canonical transformation (PCT) to an exactly solvable constant-mass Schrödinger equation [21, 22]. Recently, such an approach has proved its efficiency again by providing a straightforward generalization [23] of a harmonic oscillator model wherein both the mass and the angular frequency are dependent on the position [24].

The purpose of the present paper is to re-examine a new model of semiconfined harmonic oscillator with a mass that varies with position, which has the striking property of having the same spectrum as the standard harmonic oscillator model [25]. By using the PCT method, we plan to prove that one can find a family of PDM and semiconfined potential pairs corresponding to such a spectrum and to which the original PDM and semiconfined harmonic oscillator pair belongs.

The paper is organized as follows. In Sec. 2, the model of [25] is reviewed and shown to be derivable by applying the PCT technique to the constant-mass isotonic oscillator model [26, 27]. In Sec. 3, an extension of the model is proposed by starting from a more general PDM and determining the associated semiconfined potential. Finally, Sec. 4 contains some comments.

2 Semiconfined harmonic oscillator model and its derivation by the PCT technique

Jafarov and Van der Jeugt recently determined the exact solution of a PDM semi-confined harmonic oscillator model, characterized by the Schrödinger equation [25]

$$\left(-\frac{d}{dx}\frac{1}{M(x)}\frac{d}{dx}+V_{\text{eff}}(x)\right)\psi_n(x)=E_n\psi_n(x), \quad (1)$$

where the kinetic energy operator has the BenDaniel-Duke form [28] and the potential has the harmonic oscillator form

$$V_{\text{eff}}(x)=\frac{1}{4}M(x)\omega^2x^2, \quad (2)$$

except that the mass

$$M(x)=\begin{cases} \left(1+\frac{x}{a}\right)^{-1} & \text{if } -a < x < +\infty, \\ +\infty & \text{if } x \leq -a, \end{cases} \quad (3)$$

with $a > 0$, depends on the position in such a way that $V_{\text{eff}}(-a) = +\infty$ and $\lim_{x \rightarrow +\infty} V_{\text{eff}}(x) = +\infty$.¹

By directly solving the differential equation (1), they found that the spectrum of this semiconfined model is that of the standard harmonic oscillator,

$$E_n = \omega \left(n + \frac{1}{2}\right), \quad n = 0, 1, 2, \dots, \quad (4)$$

with corresponding wavefunctions

$$\psi_n(x) = C_n \left(1 + \frac{x}{a}\right)^{\frac{1}{2}\omega a^2} e^{-\frac{1}{2}\omega a(x+a)} L_n^{(\omega a^2)}\left(\omega a^2 \left(1 + \frac{x}{a}\right)\right), \quad -a < x < +\infty, \quad (5)$$

expressed in terms of Laguerre polynomials $L_n^{(\alpha)}(z)$ and vanishing at $x = -a$ and $x \rightarrow +\infty$, as it should be. Here C_n is a normalization coefficient given by

$$C_n = (\omega a^2)^{\frac{1}{2}(\omega a^2+1)} \sqrt{\frac{n!}{a\Gamma(\omega a^2 + n + 1)}}. \quad (6)$$

¹Note that we have adopted here units wherein $\hbar = 2m_0$ in the original paper.

These results may be alternatively derived by applying a PCT to the constant-mass Schrödinger equation for the isotonic oscillator [26, 27]

$$\left(-\frac{d^2}{du^2} + U(u)\right) \phi_n(u) = \epsilon_n \phi_n(u), \quad (7)$$

where

$$U(u) = \frac{1}{4}\bar{\omega}^2 u^2 + \frac{g}{u^2}, \quad g > 0, \quad 0 < u < +\infty, \quad (8)$$

$$\epsilon_n = \bar{\omega}(2n + \alpha + 1), \quad \alpha = \frac{1}{2}\sqrt{1 + 4g}, \quad (9)$$

and

$$\phi_n(u) \propto u^{\alpha+\frac{1}{2}} e^{-\frac{1}{4}\bar{\omega}u^2} L_n^{(\alpha)}\left(\frac{1}{2}\bar{\omega}u^2\right). \quad (10)$$

A PCT transforming an equation such as (7) into a PDM equation of type (1) [21, 22] consists in making a change of variable

$$u(x) = \bar{a}v(x) + \bar{b}, \quad v(x) = \int^x \sqrt{M(x')} dx', \quad (11)$$

and a change of function

$$\phi_n(u(x)) \propto [M(x)]^{-1/4} \psi_n(x). \quad (12)$$

The potential $V_{\text{eff}}(x)$ and the energy eigenvalues E_n of the PDM Schrödinger equation are then given in terms of the potential and the energy eigenvalues of the constant-mass one by

$$V_{\text{eff}}(x) = \bar{a}^2 U(u(x)) + \frac{M''}{4M^2} - \frac{7M'^2}{16M^3} + \bar{c}, \quad (13)$$

and

$$E_n = \bar{a}^2 \epsilon_n + \bar{c}, \quad (14)$$

where a prime denotes derivation with respect to x and \bar{a} , \bar{b} , \bar{c} are three real constants.

In the present case, from (3) and (11), we directly obtain

$$v(x) = 2a\sqrt{1 + \frac{x}{a}}. \quad (15)$$

and

$$\frac{M''}{4M^2} - \frac{7M'^2}{16M^3} = \frac{1}{16a^2} \left(1 + \frac{x}{a}\right)^{-1} \quad (16)$$

for $-a < x < +\infty$. With the choice $\bar{a} = \sqrt{\frac{\omega}{2\bar{\omega}}}$, $\bar{b} = 0$, we get for the change of variable (11)

$$u(x) = a\sqrt{\frac{2\omega}{\bar{\omega}}} \sqrt{1 + \frac{x}{a}} \quad (17)$$

and the change of function (12), together with (10), leads to

$$\psi_n(x) = C_n \left(1 + \frac{x}{a}\right)^{\alpha/2} e^{-\frac{1}{2}\omega a(x+a)} L_n^{(\alpha)} \left(\omega a^2 \left(1 + \frac{x}{a}\right)\right), \quad (18)$$

where C_n turns out to be

$$C_n = (\omega a^2)^{\frac{1}{2}(\alpha+1)} \sqrt{\frac{n!}{a\Gamma(\alpha+n+1)}}. \quad (19)$$

Furthermore, on assuming $\bar{c} = -\frac{\omega}{2}\alpha$, the transformed potential (13) becomes

$$V_{\text{eff}}(x) = \frac{a\omega^2}{4(x+a)} \left(x + a - \frac{\alpha}{a\omega}\right)^2, \quad (20)$$

with corresponding eigenvalues E_n given by (4).

If we compare these results with those of [25], we notice that we have obtained the same energy spectrum (4), but with generalized potential and wavefunctions, since the latter depend on an extra parameter α absent in [25]. By taking $\alpha = a^2\omega$, the original results are retrieved, but for other values of α , the potential (20) describes a semiconfined shifted harmonic oscillator. Note that such a potential might be interpreted as a semiconfined harmonic oscillator in a uniform gravitational field as was done for a shifted harmonic oscillator with another type of PDM [29, 30].

3 Family of generalized semiconfined oscillator models

A further generalization of the model of [25] can be obtained by changing the PDM (3) into a PDM depending on some parameter m taking values in the interval $0 < m < 2$,

$$M(x) = \begin{cases} \left(1 + \frac{x}{a}\right)^{-m} & \text{if } -a < x < +\infty, \\ +\infty & \text{if } x \leq -a, \end{cases} \quad (21)$$

and determining the associated potential $V_{\text{eff}}(x)$ with the assumption that the starting constant-mass Schrödinger equation remains as given in (7) and (8). The results of Sec. 2 will then correspond to the $m = 1$ special case.

Equations (15) and (16) are now replaced by

$$v(x) = \frac{2a}{2-m} \left(1 + \frac{x}{a}\right)^{1-\frac{m}{2}} \quad (22)$$

and

$$\frac{M''}{4M^2} - \frac{7M'^2}{16M^3} = -\frac{1}{16a^2} m(3m-4) \left(1 + \frac{x}{a}\right)^{m-2}, \quad (23)$$

respectively. On keeping the same values for \bar{a} , \bar{b} , and \bar{c} as in Sec. 2, we get a new change of variable

$$u(x) = \frac{a}{2-m} \sqrt{\frac{2\omega}{\bar{\omega}}} \left(1 + \frac{x}{a}\right)^{1-\frac{m}{2}}, \quad (24)$$

but the resulting energy eigenvalues remain given by (4). From (13), however, the resulting potential turns out to be m -dependent and given by

$$V_{\text{eff}}(x) = \frac{a^m \omega^2}{4(2-m)^2} (x+a)^{2-m} + \frac{[(m-2)\alpha - (m-1)][(m-2)\alpha + m - 1]}{4a^m (x+a)^{2-m}} - \frac{1}{2} \omega \alpha. \quad (25)$$

This is also the case for the wavefunctions, which become

$$\begin{aligned} \psi_n(x) &= C_n \left(1 + \frac{x}{a}\right)^{-\frac{m}{2}(\alpha+1) + \alpha + \frac{1}{2}} e^{-\frac{1}{2(2-m)^2} \omega a^2 \left(1 + \frac{x}{a}\right)^{2-m}} \\ &\times L_n^{(\alpha)} \left(\frac{\omega a^2}{(2-m)^2} \left(1 + \frac{x}{a}\right)^{2-m} \right), \end{aligned} \quad (26)$$

where

$$C_n = \left(\frac{\omega a^2}{(2-m)^2} \right)^{\frac{1}{2}(\alpha+1)} \sqrt{\frac{(2-m)n!}{a\Gamma(\alpha+n+1)}}. \quad (27)$$

The new m -dependent potential (25) will be a semiconfined potential provided it goes to $+\infty$ for $x \rightarrow +\infty$ and $x \rightarrow -a$. The former condition is automatically satisfied, but the latter imposes that

$$\alpha > \frac{m-1}{2-m}, \quad (28)$$




Figure 1: Plot of the semiconfined potential (25) in terms of x for $m = 1$ (black line), $m = \frac{1}{2}$ (red line), and $m = \frac{3}{2}$ (green line). The parameter values are $\omega = 1$, $a = 2$, and $\alpha = 4$.

which implies a restriction for m values such that $\frac{m-1}{2-m} > \frac{1}{2}$, i.e., for those in the interval $\frac{4}{3} < m < 2$. In such a case, the wavefunctions (26) vanish for $x \rightarrow +\infty$ and $x \rightarrow -a$, as it should be. The minimum of the potential occurs for

$$x_{\min} = -a + \left\{ \frac{(2-m)^2}{a^m \omega} \sqrt{\alpha^2 - \left(\frac{m-1}{2-m} \right)^2} \right\}^{1/(2-m)} \quad (29)$$

and is given by

$$(V_{\text{eff}})_{\min} = \frac{1}{2} \omega \left\{ \sqrt{\alpha^2 - \left(\frac{m-1}{2-m} \right)^2} - \alpha \right\}. \quad (30)$$

It is therefore slightly negative, except for $m = 1$ for which it vanishes.

In Fig. 1, we show the dependence of the semiconfined potential (25) on m . The black line corresponds to the original semiconfined harmonic oscillator (2).

4 Comments

In the present paper, we have first shown that the PCT method applied to the constant-mass Schrödinger equation for the isotonic oscillator allows us to easily

retrieve the results of [25] and to extend them in order to describe a semiconfined shifted harmonic oscillator, which might be interpreted as a semiconfined harmonic oscillator in a uniform gravitational field.

In a second step, we have obtained a further generalization by considering a m -dependent PDM for $0 < m < 2$ and by deriving the corresponding semiconfined potential with the same spectrum as the standard harmonic oscillator. We have therefore constructed a family of PDM and potential pairs, to which the original pair belongs as it corresponds to $m = 1$.

In [25], the BenDaniel-Duke ordering [28] was chosen for the momentum and mass operators. One finds, however, in the literature, several other orderings, which are special cases of the von Roos general two-parameter form of the kinetic energy operator [31], for which the Schrödinger operator writes

$$\left\{ -\frac{1}{2} \left[M(x)^\xi \frac{d}{dx} M(x)^\eta \frac{d}{dx} M(x)^\zeta + M(x)^\zeta \frac{d}{dx} M(x)^\eta \frac{d}{dx} M(x)^\xi \right] + V_{\text{vR}}(x) \right\} \psi_n(x) = E_n \psi_n(x), \quad (31)$$

where ξ, η, ζ are some real parameters restricted by the condition $\xi + \eta + \zeta = -1$. In particular, the BenDaniel-Duke ordering corresponds to $\xi = \zeta = 0, \eta = -1$ and the relation between the potentials in (1) and (31) is given by

$$V_{\text{vR}}(x) = V_{\text{eff}}(x) - \frac{1}{2}(1 + \eta) \frac{M''}{M^2} + [\xi(\xi + \eta + 1) + \eta + 1] \frac{M'^2}{M^3}. \quad (32)$$

For the mass chosen in (21), the latter becomes

$$V_{\text{vR}}(x) = V_{\text{eff}}(x) + \left\{ -\frac{1}{2}(1 + \eta)m(m + 1) + [\xi(\xi + \eta + 1) + \eta + 1]m^2 \right\} \frac{(x + a)^{m-2}}{a^m}. \quad (33)$$

It is worth noting, in particular, the Zhu-Kroemer [32] and Mustafa-Mazharimousavi [33] orderings, which pass the de Souza Dutra and Almeida test [34] as good orderings. The former corresponds to $\xi = \zeta = -\frac{1}{2}, \eta = 0$, and leads to replacing (33) by

$$V_{\text{ZK}}(x) = \frac{a^m \omega^2}{4(2 - m)^2} (x + a)^{2-m} + \frac{[(m - 2)\alpha - 1][(m - 2)\alpha + 1]}{4a^m (x + a)^{2-m}} - \frac{1}{2} \omega \alpha, \quad (34)$$

while the latter is associated with $\xi = \zeta = -\frac{1}{4}$, $\eta = -\frac{1}{2}$, and gives rise to

$$V_{\text{MM}}(x) = \frac{a^m \omega^2}{4(2-m)^2} (x+a)^{2-m} + \frac{(m-2)^2(\alpha^2 - \frac{1}{4})}{4a^m (x+a)^{2-m}} - \frac{1}{2} \omega \alpha. \quad (35)$$

These potentials have a behaviour very similar to that of $V_{\text{eff}}(x)$, since they are semiconfined for α restricted to $\alpha > 1/(2-m)$ or for any value of α ($> 1/2$ by definition (9)), respectively. The place of the minimum and its value are given by (29) and (30) provided $\sqrt{\alpha^2 - [(m-1)/(2-m)]^2}$ is replaced by $\sqrt{\alpha^2 - 1/(2-m)^2}$ or $\sqrt{\alpha^2 - 1/4}$.

Acknowledgements

This work was supported by the Fonds de la Recherche Scientifique-FNRS under Grant No. 4.45.10.08.

References

- [1] G. Bastard, *Wave Mechanics Applied to Semiconductor Heterostructures* (Editions de Physique, Les Ulis, 1988)
- [2] C. Weisbuch, B. Vinter, *Quantum Semiconductor Heterostructures* (Academic, New York, 1997)
- [3] L. Serra, E. Lipparini, Spin response of unpolarized quantum dots. *Europhys. Lett.* **40**, 667 (1997)
- [4] P. Harrison, A. Valavanis, *Quantum Wells, Wires and Dots: Theoretical and Computational Physics of Semiconductor Nanostructures* (Wiley, Chichester, 2016)
- [5] M. Barranco, M. Pi, S.M. Gatica, E.S. Hernández, J. Navarro, Structure and energetics of mixed ^4He - ^3He drops. *Phys. Rev. B* **56**, 8997 (1997)
- [6] M.R. Geller, W. Kohn, Quantum mechanics of electrons in crystals with graded composition. *Phys. Rev. Lett.* **70**, 3103 (1993)
- [7] F. Arias de Saavedra, J. Boronat, A. Polls, A. Fabrocini, Effective mass of one ^4He atom in liquid ^3He . *Phys. Rev. B* **50**, 4248(R) (1994)
- [8] A. Puente, L. Serra, M. Casas, Dipole excitation of Na clusters with a non-local energy density functional. *Z. Phys. D* **31**, 283 (1994)
- [9] P. Ring, P. Schuck, *The Nuclear Many Body Problem* (Springer, New York, 1980)
- [10] D. Bonatsos, P.E. Georgoudis, D. Lenis, N. Minkov, C. Quesne, Bohr Hamiltonian with a deformation-dependent mass term for the Davidson potential. *Phys. Rev. C* **83**, 044321 (2011)
- [11] W. Willatzen, B. Lassen, The BenDaniel-Duke model in general nanowire structures. *J. Phys. Condens. Matter* **19**, 136217 (2007)

- [12] N. Chamel, Effective mass of free neutrons in neutron star crust. Nucl. Phys. A **773**, 263 (2006)
- [13] C. Quesne, V.M. Tkachuk, Deformed algebras, position-dependent effective mass and curved spaces: An exactly solvable Coulomb problem. J. Phys. A: Math. Gen. **37**, 4267 (2004)
- [14] A. Kempf, Uncertainty relation in quantum mechanics with quantum group symmetry. J. Math. Phys. **35**, 4483 (1994)
- [15] H. Hinrichsen, A. Kempf, Maximal localization in the presence of minimal uncertainties in positions and in momenta. J. Math. Phys. **37**, 2121 (1996)
- [16] E. Witten, Reflections on the fate of spacetime. Phys. Today **49**, 24 (1996)
- [17] E. Schrödinger, A method of determining quantum-mechanical eigenvalues and eigenfunctions. Proc. R. Ir. Acad. **A46**, 9 (1940)
- [18] E.G. Kalnins, W. Miller Jr., G.S. Pogosyan, Superintegrability and associated polynomial solutions: Euclidean space and the sphere in two dimensions. J. Math. Phys. **37**, 6439 (1996)
- [19] E.G. Kalnins, W. Miller Jr., G.S. Pogosyan, Superintegrability on the two-dimensional hyperboloid. J. Math. Phys. **38**, 5416 (1997)
- [20] C. Quesne, First-order intertwining operators and position-dependent mass Schrödinger equations in d dimensions. Ann. Phys. (NY) **321**, 1221 (2006)
- [21] B. Bagchi, P. Gorain, C. Quesne, R. Roychoudhury, A general scheme for the effective-mass Schrödinger equation and the generation of the associated potentials. Mod. Phys. Lett. A **19**, 2765 (2004)
- [22] C. Quesne, Point canonical transformation versus deformed shape invariance for position-dependent mass Schrödinger equations. SIGMA **5**, 046 (2009)

- [23] C. Quesne, Comment on 'Exact solution of the position-dependent effective mass and angular frequency Schrödinger equation: harmonic oscillator model with quantized confinement parameter'. J. Phys. A: Math. Theor. **54**, 368001 (2021)
- [24] E.I. Jafarov, S.M. Nagiyev, R. Oste, J. Van der Jeugt, Exact solution of the position-dependent effective mass and angular frequency Schrödinger equation: harmonic oscillator model with quantized confinement parameter. J. Phys. A: Math. Theor. **53**, 485301 (2020)
- [25] E.I. Jafarov, J. Van der Jeugt, Exact solution of the semiconfined harmonic oscillator model with a position-dependent effective mass. Eur. Phys. J. Plus **136**, 758 (2021)
- [26] Y. Weissman, J. Jortner, The isotonic oscillator. Phys. Lett. A **70**, 177 (1979)
- [27] D. Zhu, A new potential with the spectrum of an isotonic oscillator. J. Phys. A: Math. Gen. **20**, 4331 (1987)
- [28] D.J. BenDaniel, C.B. Duke, Space-charge effects on electron tunneling. Phys. Rev. **152**, 683 (1966)
- [29] E.I. Jafarov, S.M. Nagiyev, A.M. Jafarova, Quantum-mechanical explicit solution for the confined harmonic oscillator model with the von Roos kinetic energy operator. Rep. Math. Phys. **86**, 25 (2020)
- [30] E.I. Jafarov, S.M. Nagiyev, Angular part of the Schrödinger equation for the Hautot potential as a harmonic oscillator with a coordinate-dependent mass in a uniform gravitational field. Theor. Math. Phys. **207**, 447 (2021)
- [31] O. von Roos, Position-dependent effective masses in semiconductor theory. Phys. Rev. B **27**, 7547 (1983)

- [32] Q.-G. Zhu, H. Kroemer, Interface connection rules for effective-mass wave functions at an abrupt heterojunction between two different semiconductors. *Phys. Rev. B* **27**, 3519 (1983)
- [33] O. Mustafa, S.H. Mazharimousavi, Ordering ambiguity revisited via position dependent mass pseudo-momentum operators. *Int. J. Theor. Phys.* **46**, 1786 (2007)
- [34] A. de Souza Dutra, C.A.S. Almeida, Exact solvability of potentials with spatially dependent effective masses. *Phys. Lett. A* **275**, 25 (2000)