

Optimizing Strongly Interacting Fermionic Hamiltonians

Matthew B. Hastings*

Ryan O’Donnell†

August 21, 2023

Abstract

The fundamental problem in much of physics and quantum chemistry is to optimize a low-degree polynomial in certain anticommuting variables. Being a quantum mechanical problem, in many cases we do not know an efficient classical witness to the optimum, or even to an approximation of the optimum. One prominent exception is when the optimum is described by a so-called “Gaussian state”, also called a free fermion state. In this work we are interested in the complexity of this optimization problem when no good Gaussian state exists. Our primary testbed is the Sachdev–Ye–Kitaev (SYK) model of random degree- q polynomials, a model of great current interest in condensed matter physics and string theory, and one which has remarkable properties from a computational complexity standpoint. Among other results, we give an efficient classical certification algorithm for upper-bounding the largest eigenvalue in the $q = 4$ SYK model, and an efficient quantum certification algorithm for lower-bounding this largest eigenvalue; both algorithms achieve constant-factor approximations with high probability.

1 Introduction

1.1 Classical optimization

Classical constraint satisfaction problems (CSPs) are very well studied in theoretical computer science, from the perspectives of both worst-case complexity and average-case complexity. We know many ingenious approximation algorithms — often based on semidefinite programming (SDP) [RS09] — for efficiently finding solutions within a constant factor of optimal. At the same time, we also have very precise predictions for the computational intractability of achieving certain approximation factors, subject to conjectures like $P \neq NP$ [Hås01], the Unique Games Conjecture [Rag09], and the optimality of Sum-of-Squares (SOS) hierarchies [KMOW17]. CSPs involve finding an assignment (typically a Boolean one) to a large number n of variables so as to satisfy many given constraints, each a predicate involving at most some q variables. As such, they can often be expressed as finding the maximum value of an n -variate degree- q polynomial over assignments from $\{\pm 1\}^n$; in other words (and in the homogeneous case),

$$\max \left\{ \sum_{\substack{S \subseteq [n] \\ |S|=q}} a_S \prod_{j \in S} \chi_j \quad : \quad \chi_j^2 = 1 \ \forall j \right\}. \quad (1)$$

For example, the $q = 2$ case is essentially the “Max-Cut” problem (with edge-weights), and the famous Goemans–Williamson SDP-rounding algorithm and its successors [GW95, CW04] can efficiently find an $O(\log(1/\epsilon))$ -factor approximation to the optimum in any worst-case instance when

*Station Q. and Microsoft Quantum. mahastin@microsoft.com

†Microsoft Quantum and Carnegie Mellon University Computer Science Department. odonnell@cs.cmu.edu

the optimum is at least $\epsilon|a|_1$. Regarding $q = 2$ in the average case, if the a_S 's are chosen to be independent standard Gaussian random variables, then we have the Sherrington–Kirkpatrick model, and Montanari [Mon21] has given an efficient algorithm for finding a $(1-o(1))$ -factor approximation to the optimum, with high probability.

On the other hand (and oversimplifying the story), CSP optimization tends to be far more intractable once $q \geq 3$. For example, work of Håstad and successors [Hås01, MR10], shows that there are worst-case $q = 3$ instances of (1), with $m = O(n)$ nonzero coefficients $a_S \in \{\pm 1\}$, where the optimum is $(1-\epsilon)|a|_1 = (1-\epsilon)m$, but it is NP-hard to find a solution of value at least $m/f(n)$ for some slowly growing $f(n) \rightarrow \infty$ (and likely this cannot even be done in time $2^{n^{.99}}$). Furthermore, for *random* instances of the above form, even though the optimum is typically $O(\sqrt{n})$ we have significant evidence (in the form of SOS hierarchy lower bounds) that any algorithm for certifying the optimum is at most $.99m$ requires $2^{\Omega(n)}$ time [Gri01].

1.2 Fermionic optimization

There exist analogous optimization problems in the quantum setting. Here we must distinguish two cases. In one case, called a “spin system”, the Hilbert space of the quantum system has a tensor product structure, and the analogue of (1) is called the *q-Local Hamiltonian problem*. In the other case, one considers fermionic degrees of freedom, which obey the so-called canonical anticommutation relations.¹ This second case is the focus of our paper. It is well studied, both from a mathematical physics point of view and for applications in quantum chemistry, condensed matter physics, and materials science.

In addition to being describable in physics terms, a fermionic optimization can be posed rather similarly to (1), but with the following differences: First, the indeterminates χ_j are to be assigned *self-adjoint matrices*, rather than numbers, and it is required that they pairwise *anticommute*. Such matrices are known as Majorana operators. Second, the quantity to be maximized is the largest *eigenvalue* of the resulting matrix. Thus a typical fermionic optimization problem may be stated as follows:

$$\max \left\{ \lambda_{\max}(h), \ h = \sum_{\substack{S=\{j_1, \dots, j_q\} \subseteq [n] \\ j_1 < \dots < j_q}} a_S \chi_{j_1} \cdots \chi_{j_q} \quad : \quad \chi_j^* = \chi_j \ \forall j, \ \chi_j^2 = \mathbb{1} \ \forall j, \ \chi_j \chi_k = -\chi_k \chi_j \ \forall j, k \right\}, \quad (2)$$

where $\mathbb{1}$ denotes the identity operator.²

Although (2) as phrased involves searching over all valid matrix-assignments for the χ_j 's, this is not the standard viewpoint in physics. Rather (as is reviewed in Section 2.4), it is without loss of generality to fix a particular canonical assignment of D -dimensional “gamma matrices” γ_j to the χ_j 's, where $D = 2^{n/2}$. Thus h , termed the “Hamiltonian” in physics, may be thought of as an implicitly-represented matrix of exponentially large dimension, and the only task is to determine its largest eigenvalue. To recall some further physics terminology, the expectation value of the matrix h in some pure quantum state $|\psi\rangle$ — i.e., the value of vector $|\psi\rangle$ under h 's quadratic form — is called its “energy”. Usually in physics one is concerned with the smallest possible energy, termed the ground state energy, but we will follow the computer science convention of seeking the maximum eigenvalue. (Of course, these are equivalent upon replacing h by $-h$.)

The particular case of $q = 4$ corresponds to finding the binding energy of a molecule with 2-body interactions, and is therefore one of the most basic computational tasks in quantum chemistry. Here

¹One can of course consider systems with both fermionic and spin degrees of freedom.

²We remark that this problem is only physically natural if q is even. One should also arrange for h to be self-adjoint and hence have real eigenvalues; this occurs, e.g., if q is a multiple of 4 and each a_S is real.

the binding energy is the difference between the ground state energy for some given molecule and the sum of ground state energies of its constituent atoms. ³

1.3 Computational complexity

The $q = 2$ case of fermionic optimization (2) can be solved by an efficient classical algorithm, involving diagonalizing the Hamiltonian. Moreover, the algorithm can find an implicit representation of the optimal state, which will be a so-called (*fermionic*) *Gaussian state* [TD02, Bra05]. However, for $q > 2$ the problem becomes computationally difficult. To see this, one may observe that the collection of operators $i\chi_{2j-1}\chi_{2j}$ (for $j = 1 \dots n/2$) square to the identity and also *commute*. It follows from this that one can (efficiently) encode any n -variable, degree- q instance of a Boolean CSP (as in (1)) by a $2n$ -variable, degree- $2q$ instance of the fermionic problem (2). (Furthermore, in this encoding, commuting Boolean assignments correspond to fermionic Gaussian states.) This reduction shows that the degree-4 fermionic problem is NP-hard. In fact, one may also encode the q -Local Hamiltonian problem by a degree- $2q$ fermionic problem [LCV07], and it therefore follows from [KKR06] that the degree-4 fermionic optimization problem is even QMA-hard.

That said, there are quite good approximation algorithms for degree-2 Boolean constraint satisfaction (via Goemans–Williamson [GW95] and its extensions), whereas for degree-3 and above there are strong inapproximability results. The latter observation shows that degree-6 (and above) fermionic optimization will be hard to approximate well, but leaves open the possibility that the $q = 4$ case (which is most relevant for quantum chemistry) may be a “sweet spot” in terms of complexity.

In this work we focus on such approximation problems. In the case of quantum optimization problems, though, a fundamentally new difficulty arises. In the classical case of (1), one can naturally certify that the optimum is at least some α by providing a witnessing assignment $\chi \in \{\pm 1\}^n$. By contrast, in the quantum case, there may well be no efficient way to classically represent a witness (such as a state $|\psi\rangle$) to the ground state energy, or even to an energy reasonably close to the optimum. Indeed, assuming $\text{QMA} \neq \text{NP}$, this would be implied by the *quantum PCP conjecture* [AAV13] (at least for local tensor-product Hamiltonians), which posits that determining the ground state energy up to a constant factor is QMA-hard. A noncomputational implication of the quantum PCP conjecture is the *NLTS (no low-energy trivial states) conjecture* [FH14], which posits the nonexistence of a certain class of easily described states with energy close to optimum. It is natural to define analogous conjectures, fermionic quantum PCP (fqPCP) and fermionic NLTS (fNLTS), for sparse instances of fermionic optimization problems, but this seems not to have been studied.

Besides trying to certify that the optimum of (2) is at least some α , one may also try to certify that it is at *most* some β . For this task, it is natural to consider SDP relaxations (as was done in [BH13] for the Local Hamiltonian problem), and in particular SOS hierarchies. Most generally one may use the non-commutative SOS hierarchy, developed in, e.g., [HM04, NPA08, DLTW08, PNA10]; it provides increasingly tight upper bounds $\beta_q \geq \beta_{q+2} \geq \beta_{q+4} \geq \dots$ on (2), with β_k being

³A remark regarding applications in quantum chemistry: in many cases one can define a “particle number operator” $i \sum_{j=1}^{n/2} \chi_{2j-1} \chi_{2j}$ which commutes with the Hamiltonian h . In this case, one may replace the Majorana operators with certain linear combinations called creation and annihilation operators so that every term in the Hamiltonian has an equal number of creation and annihilation operators. For practical purposes, this representation with creation and annihilation operators is convenient, as for such Hamiltonians one immediately knows that the expectation value of an operator that does not commute with the particle number vanishes. However, in this paper we will not focus on this case. Interestingly, though, several of the constraints considered in quantum chemistry can be more readily expressed in terms of the Majorana operators. For example, the so-called “ T_1 constraint” is a constraint on expectation values of degree-4 polynomials that follows from positivity of the “degree-6 pseudoexpectation” value, and (in the Majorana language) from the fact that a degree-6 polynomial in Majorana operators with real coefficients is skew-adjoint.

certifiable in deterministic classical $n^{O(k)}$ time. In the specific context of fermionic optimization, a closely related SDP hierarchy has been studied as a way of characterizing which two-particle reduced density matrices can arise from a many-body system (i.e., which collections of degree-4 expectation values $\text{tr}(\rho \gamma_i \gamma_j \gamma_k \gamma_l)$ can arise from a genuine state ρ). See, e.g., [Col63, Erd78, Per78, ME01, Maz12, Kly06]. This so-called “ N -representability problem” is QMA-hard, and is perhaps the fundamental problem of quantum chemistry [LCV07].

1.4 Strongly interacting Hamiltonians, and SYK model

As mentioned, in the case $q = 2$, the fermionic optimization problem (2) is solvable efficiently, with the optimizer being an efficiently-representable “Gaussian state”. Given this fact, previous research (e.g. [BGKT19]) has studied how close Gaussian states can come to achieving optimality in the $q = 4$ case of (2), and how efficiently one can optimize over Gaussian states in this case. In the present work, our main interest is the complexity of optimizing fermionic Hamiltonians when *no* good Gaussian state exists. A prominent example of this, and our main testbed, is the “SYK model” of random fermionic Hamiltonians.

The *Sachdev–Ye–Kitaev (SYK) model* refers to the case of (2) where the coefficients a_S are chosen to be independent random Gaussians (and where q and n are even).⁴ Contrary to the traditional physics normalization, we will scale by $1/\sqrt{\binom{n}{q}}$ so that $|a|_2^2 = \sum_S a_S^2 = 1$ in expectation. The SYK model has attracted considerable interest in physics for being “maximally chaotic”, and for having a ground state that is, in a sense, as far from being a Gaussian state as is possible for a low-degree Hamiltonian. With study dating back to [FW70, BF71], the canonical references for this model are [SY93, Kit15]. A review from a high energy physics perspective is [Ros19]. The physics papers [GV16, GJV18] are rather accessible for those with a mathematics background, and the works [FTW19, FTW18, FTW20] give some entirely mathematical results.

Considerations from mathematical physics suggest that for h drawn from the degree-4 SYK model, the maximum value in (2) — which we will denote by $\text{Opt}(h)$ — is $\Theta(\sqrt{n})$ ⁵ with high probability. Indeed, as we discuss in Section 3.2, even finer-grained details are predicted, including that the leading constant hidden in the $\Theta(\cdot)$ should be $\frac{1}{2\sqrt{2}}$. On the other hand, the only mathematical proofs we are aware of come from [FTW19], from which it follows that $\omega(1) < \text{Opt}(h) \leq \sqrt{\ln 2} \sqrt{n}$ with high probability.

1.5 Summary of our results

We provide a variety of new results concerning efficient optimization of fermionic Hamiltonians. Primary among these are a rigorous proof that $\text{Opt}(h) = \Theta(\sqrt{n})$ (with high probability) for h drawn from the degree-4 SYK model. Moreover, we prove this via efficient *lower and upper bound certification algorithms*. For the upper bound, we show that degree-6 SOS works, and we prove a tighter bound on the numerical constant for degree-8 SOS:

Theorem 1.1. *The degree-8 SOS relaxation, which is solvable in deterministic $\text{poly}(n)$ time, certifies that $\text{Opt}(h) \leq \sqrt{1 + \sqrt{6}} \sqrt{n}$ with high probability over h drawn from the degree-4 SYK model. (Moreover, even the degree-6 SOS relaxation certifies that $\text{Opt}(h) \leq O(\sqrt{n})$ with high probability.)*

⁴In all cases, a prefactor of $i^{\binom{q}{2}}$ should be included to ensure self-adjointness. We remark that although the case of odd q or n is not natural physically, these cases are still mathematically interesting.

⁵We remark that under the conventional normalization for SYK used in physics, this corresponds to ground state energy $-\Theta(n)$.

Remark 1.2. We also show (Theorem 6.6) that degree-4 SOS *fails* to certify anything better than $\text{Opt}(h) \lesssim n$. On the other hand, our certification procedure can be carried out in a natural “fragment” of degree-6 SOS of size $O(n^4)$. It would be interesting to see if the “fragment” introduced here can have practical applications in, e.g., the simulation of quantum chemistry as in [Maz12].

Our lower bound, perhaps surprisingly, is via an efficient *quantum* certification algorithm that creates a variational state of large expectation value. It is an outstanding open problem as to whether there is an efficient classical certification algorithm (recall that Gaussian states are known to only achieve expectation value $O(1)$).

Theorem 1.3. *There is a $\text{poly}(n)$ -time quantum algorithm that, given any degree-4 Hamiltonian h , returns a quantum state ρ . With high probability over h drawn from the degree-4 SYK model, the output state ρ has expectation value $\Omega(\sqrt{n})$.*

Remark 1.4. As we will discuss, this guarantee *should* be thought of as providing a (quantumly-checkable) “certificate”, since for *every* h and ρ , there is a very simple $\text{poly}(n) \cdot \log(1/\delta)$ -time quantum algorithm that estimates the expectation value of h under ρ to any $\pm 1/\text{poly}(n)$ accuracy with confidence $1 - \delta$.

Together, Theorems 1.1 and 1.3 naturally suggest:

Open problem. *Are there efficient quantum/classical algorithms that certify the value of degree-4 SYK instances up to a $(1 \pm o(1))$ -factor (from above or below or both)?*

Open problem. *Are there sparse instances of (2) with $q = O(1)$ that fulfill the fermionic NLTS conjecture⁶*

We now discuss some additional results we prove. First, we give the below nontrivial upper bound for *worst-case* degree-4 Hamiltonians. This theorem is used in establishing Theorem 1.1.

Theorem 1.5. *Let h be as in (2) with $q = 4$, and assume for normalization that $\sum_S a_S^2 = 1$. Then*

$$\text{Opt}(h) \leq \sqrt{\binom{n/2}{2}} \sim \frac{1}{2\sqrt{2}}n,$$

with equality being attained by a certain Hamiltonian (essentially the square of a quadratic) in which all terms commute. (This may be contrasted with the commutative case of (1), where $\text{Opt}(h) = \sqrt{\binom{n}{4}} \sim \frac{1}{\sqrt{24}}n^2$ is attainable.)

The upper bound in this theorem relies only on the following algebraic relation: two monomials $\chi_{j_1}\chi_{j_2}\chi_{j_3}\chi_{j_4}$ and $\chi_{k_1}\chi_{k_2}\chi_{k_3}\chi_{k_4}$ anticommute if and only if $|\{j_1, \dots, j_4\} \cap \{k_1, \dots, k_4\}|$ is odd. Indeed, we prove Theorem 1.5 by studying the optimization problem (2) in a more general context where one doesn’t assume that χ_j and χ_k anticommute for all $j \neq k$, but rather one assumes — for

⁶The NLTS conjecture posits that there exists a family of quantum Hamiltonians which are “spin systems”, in that the Hilbert space has a local tensor product structure, where all terms in the Hamiltonian h have norm $O(1)$, where each spin participates in $O(1)$ terms, and where each term acts on $O(1)$ spins, so that for some sufficiently constant $c > 0$, there are no quantum states ψ which are “low energy” in that the difference between $\langle \psi | h | \psi \rangle$ and $\text{Opt}(h)$ is $\leq cn$, and which are “trivial” in that ψ is given by acting on some product state by a bounded depth quantum circuit. A fermionic analogue of this conjecture would postulate the same property for a system with fermionic degrees of freedom.

some given graph $\Gamma = ([n], E)$ — that they anticommute if $\{j, k\} \in E$ and they commute if $\{j, k\} \notin E$. In this more general setting, we show that for h of degree 1 with $\sum_j a_j^2 = 1$, it holds that $\alpha(\Gamma) \leq \text{Opt}(h)^2 \leq \vartheta(\Gamma)$, where $\alpha(\Gamma)$ denotes the independence number of Γ , and $\vartheta(\Gamma)$ denotes its *Lovász Theta Function* value. Our [Theorem 1.5](#) is then obtained by determining the Lovász Theta Function of an appropriate Kneser-like graph family on vertex set $\binom{[n]}{4}$. The analogous problem for $\binom{[n]}{q}$, $q > 4$, remains open, as does the question of whether there exists h with $\text{Opt}(h)^2 > \alpha(\Gamma)$ strictly.

From [Theorems 1.1](#) and [1.5](#) we infer that for degree-4 h , the optimum value behaves very differently for typical (SYK) instances, versus worst-cases instances. It would seem that $\text{Opt}(h) \approx n$ is possible only if h is (roughly) a sum of a small number of squared quadratics, in which case Gaussian states achieve a near-optimal expectation value (see [Appendix B](#)). We prove some results along these lines, including the following:

Theorem 1.6. *There is a $\text{poly}(n)$ -time classical algorithm with the following property: Given h as in [\(2\)](#) with $q = 4$, $\sum_S a_S^2 = 1$, and the property that the optimum Gaussian state achieves expectation value at least ϵn , the algorithm finds a Gaussian state achieving expectation value $\Omega(\frac{\epsilon}{\log(1/\epsilon)})n$.*

This result generalizes the earlier work of Bravyi, Gosset, König, and Temme [\[BGKT19\]](#) which achieved an $\Omega(\frac{1}{\log n})$ -approximation factor.

2 Preliminaries

2.1 Notation

Notation 2.1. We use **boldface** to denote random variables. We use $\text{SYK}_q(n)$ to denote the Sachdev-Ye-Kitaev model with a given choice of q, n . We use \sim to indicate that a random variable is drawn from a particular distribution, such as $\mathbf{h} \sim \text{SYK}_4(n)$.

Notation 2.2. For a natural number n we write $[n] = \{1, 2, \dots, n\}$.

Notation 2.3. For a parameter $n \rightarrow \infty$, we sometimes write $\text{poly}(n) = n^{O(1)} = O(n^c)$ for an unspecified constant c .

Notation 2.4. $\mathbb{1}$ denotes the identity operator/matrix (where the dimension can always be inferred from context).

Notation 2.5. For an operator A of dimension D we write $\text{Tr}(A)$ for the trace of A , and $\text{tr}(A) = \frac{1}{D} \text{Tr}(A)$ for the normalized trace.

Notation 2.6. If $G = (V, E)$ is a simple undirected graph, then \overline{G} denotes the edge-complement graph $(V, \binom{V}{2} \setminus E)$. Also, K_n denotes the complete graph on n vertices, and hence \overline{K}_n denotes the empty graph on n vertices.

Notation 2.7. We use $|\cdot|_p$ to denote the ℓ_p norm of a vector; if p is not written then this is the ℓ_2 norm by default.

Notation 2.8. $\|\cdot\|_{\text{op}}$ denotes the operator norm of a matrix. However $\|\cdot\|_p$ denotes the Schatten p -norm of a matrix.

Notation 2.9. We use both ket notation, such as $|\phi\rangle$, and vector notation, such as \vec{a} , to denote vectors. Generally we try to use ket notation for pure quantum states and vector notation for everything else, but we make some exceptions. For example, if \vec{a} is a vector, $|a\rangle\langle a|$ is used as convenient notation for the projector onto \vec{a} .

Notation 2.10. We use $*$ to denote Hermitian conjugation of operators as well as to denote a formal adjoint operator on polynomials.

2.2 Quasi-Clifford algebras

We work in the algebra $\mathbb{C}^*\langle\chi\rangle$ of polynomials with complex coefficients over a finite sequence $\chi = (\chi_1, \dots, \chi_n)$ of noncommuting indeterminates. The empty word is denoted by 1. There is a formal adjoint operator $*$ on polynomials, and we will assume that all indeterminates χ_j are self-adjoint, $\chi_j^* = \chi_j$. This $*$ maps coefficients to their complex conjugates, and reverses words.

We furthermore introduce some relations. First, we assume each χ_j is an *involution*; i.e., $\chi_j^2 = 1$ for all j . Next, given an undirected graph $\Gamma = ([n], E)$ on vertex set $[n]$, we introduce the relation $\chi_j\chi_k = -\chi_k\chi_j$ for all edges $\{j, k\} \in E$, and the relation $\chi_j\chi_k = \chi_k\chi_j$ for all nonedges $\{j, k\} \notin E$. We denote the resulting algebra by $\mathfrak{C}(\Gamma)$; it is (a special case of) a *quasi-Clifford algebra*, as introduced by Gastineau-Hills [GH80, GH82]. We will be particularly interested in the complete graph $\Gamma = K_n$ with n even; then $\mathfrak{C}(K_n)$ is the “usual” Clifford algebra in which every pair of distinct indeterminates anticommutes.

Given our relations, it is easy to see that every monomial (product of indeterminates) is equivalent, up to sign change, to a multilinear monomial in which the indeterminates are ordered lexicographically. Thus we can write a general polynomial $h \in \mathfrak{C}(\Gamma)$ as

$$h = h(\chi) = \sum_{S \subseteq [n]} a_S \chi^S, \quad \text{where } \chi^S := \chi_{k_1}\chi_{k_2} \cdots \chi_{k_q} \text{ if } S = \{k_1, k_2, \dots, k_q\} \text{ with } k_1 < k_2 < \cdots < k_q, \quad (3)$$

with coefficients $a_S \in \mathbb{C}$.⁷ Hence $\mathfrak{C}(\Gamma)$ is always finite-dimensional; i.e., it is a “ \dagger -algebra”, in physics terminology. We will frequently be interested in the case where h is self-adjoint, meaning $h^* = h$. In this case, note that each a_S will either be real (if χ^S is equal to its reverse) or purely imaginary (if χ^S equals the negation of its reverse). Finally, we introduce the following notation (see Remark 2.16 for justification):

Notation 2.11. For $h \in \mathfrak{C}(\Gamma)$ as in Equation (3) we write $\text{tr}(h) := a_\emptyset$.

2.3 The completely anticommuting case

Let us collect here some simple facts concerning the case $\Gamma = K_n$, where the indeterminates pairwise anticommute.

Fact 2.12. In $\mathfrak{C}(K_n)$, monomial $(\chi^S)^* = \sigma \chi^S$, where

$$\sigma = (-1)^{\binom{|S|}{2}} = \begin{cases} +1 & \text{if } |S| = 0, 1 \pmod{4}, \\ -1 & \text{if } |S| = 2, 3 \pmod{4}. \end{cases}$$

⁷Notational notes: When discussing physics applications, we will typically write J_S instead of a_S . Also, χ^\emptyset stands for the empty monomial 1, and for $a \in \mathbb{C}$ we write just a instead of $a \cdot 1$.

Consequently, we can write a generic self-adjoint homogeneous degree- q polynomial as

$$h = i \binom{q}{2} \sum_{\substack{S \subseteq [n] \\ |S|=q}} a_S \chi^S, \quad (4)$$

where each a_S is real.

Fact 2.13. Let χ^S and χ^T be monomials in $\mathfrak{C}(K_n)$. Then

$$\chi^S \chi^T = (-1)^{|S| \cdot |T| - |S \cap T|} \chi^T \chi^S.$$

In particular, if $|S| = |T| = q$, then χ^S and χ^T anticommute iff $q - |S \cap T|$ is odd.

Fact 2.14. Let $\ell_a = a_1 \chi_1 + \cdots + a_n \chi_n$, $\ell_b = b_1 \chi_1 + \cdots + b_n \chi_n \in \mathfrak{C}(K_n)$ for $\vec{a}, \vec{b} \in \mathbb{R}^n$. Then $\ell_a \ell_a = |\vec{a}|^2$, and $\ell_a \ell_b = -\ell_b \ell_a$. In particular, if $O \in \mathbb{R}^{n \times n}$ is an orthogonal matrix and $\ell = O\chi$, then ℓ_1, \dots, ℓ_n are self-adjoint, square to 1, and pairwise anticommute. Indeed, they generate an isomorphic copy of $\mathfrak{C}(K_n)$.

2.4 Representations

By a *representation* of $\mathfrak{C}(\Gamma)$, we will mean a $*$ -homomorphism $\pi : \mathfrak{C}(\Gamma) \rightarrow \mathbb{C}^{D \times D}$ (for some finite D) satisfying $\pi(1) = \mathbb{1}$. Informally, π is an assignment of matrices (of some fixed dimension D) to each χ_j such that the assigned matrices satisfy the algebra relations. The representation theory of quasi-Clifford algebras is completely understood; Gastineau-Hills [GH82] showed that $\mathfrak{C}(\Gamma)$ is always a semisimple algebra and determined its irreducible representations (see also the concrete analysis in [Kho08]). We recap it here.

First suppose $n = 2r$ is even and $\Gamma = ([n], \{\{1, 2\}, \{3, 4\}, \dots, \{n-1, n\}\})$, the matching graph on n vertices. Then we have an (irreducible) 2^r -dimensional representation π that assigns

$$\begin{aligned} \chi_1 &= X \otimes \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} \\ \chi_2 &= Y \otimes \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} \\ \chi_3 &= \mathbb{1} \otimes X \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} \\ \chi_4 &= \mathbb{1} \otimes Y \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} \\ &\dots \\ \chi_{n-1} &= \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes X \\ \chi_n &= \mathbb{1} \otimes \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes Y, \end{aligned}$$

where we use the notation $X = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $Y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$, $Z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ for the Pauli matrices. In fact this π yields an isomorphism between $\mathfrak{C}(\Gamma)$ and $\mathbb{C}^{2^r \times 2^r}$.

As a next example, consider $\mathfrak{C}(\overline{K}_n)$, the algebra in which all indeterminates χ_j pairwise commute. Then for each $x \in \{\pm 1\}^n$ we have a 1-dimensional representation π_x in which χ_j is assigned x_j . The direct sum $\pi = \bigoplus_x \pi_x$ is a (reducible) 2^n -dimensional representation, which can alternately be expressed as the assignment

$$\chi_1 = Z \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1}, \quad \chi_2 = \mathbb{1} \otimes Z \otimes \cdots \otimes \mathbb{1}, \quad \dots, \quad \chi_n = \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes Z, \quad (5)$$

and this π yields a $*$ -homomorphism $\mathfrak{C}(\Gamma) \rightarrow \mathbb{C}^{D \times D}$.

We can also combine the previous two constructions. If Γ consists of r disjoint edges and s isolated vertices (so $n = 2r + s$), then there is a $*$ -homomorphism to $\mathbb{C}^{D \times D}$ for $D = 2^{r+s} = 2^{n-r}$, via the representation π which maps the indeterminates to matrices of the form

$$(\mathbb{1} \otimes \cdots \mathbb{1} \otimes X \otimes \mathbb{1} \otimes \cdots \mathbb{1}) \otimes (\mathbb{1} \otimes \cdots \mathbb{1}) \text{ or } (\mathbb{1} \otimes \cdots \mathbb{1} \otimes Y \otimes \mathbb{1} \otimes \cdots \mathbb{1}) \otimes (\mathbb{1} \otimes \cdots \mathbb{1}) \\ \text{or } (\mathbb{1} \otimes \cdots \mathbb{1}) \otimes (\mathbb{1} \otimes \cdots \mathbb{1} \otimes Z \otimes \mathbb{1} \otimes \cdots \mathbb{1}).$$

As a next observation, suppose Γ has adjacency matrix A , regarded as an element of $\mathbb{F}_2^{n \times n}$. Then if we replace the j th row with the j th plus the k th, and do the same columnwise, the resulting A' is the adjacency matrix of a graph Γ' with $\mathfrak{C}(\Gamma') \cong \mathfrak{C}(\Gamma)$, via the isomorphism that has $\chi'_j = i^{A_{jk}} \chi_j \chi_k$ and $\chi'_{j'} = \chi_{j'}$ for $j' \neq j$. By repeating such operations, any Γ can be transformed to another Γ' consisting of r disjoint edges and s isolated vertices, where $2r$ is the \mathbb{F}_2 -rank of the original A . In this way we can obtain an explicit $*$ -homomorphism to a matrix algebra for any particular $\mathfrak{C}(\Gamma)$.

Let us turn to the case of most interest to us, the usual Clifford algebra $\mathfrak{C}(K_n)$, where we assume for convenience that $n = 2r$ is even. In this case, we obtain the well known 2^r -dimensional representation π of $\mathfrak{C}(K_n)$ given by the Weyl–Brauer γ matrices:

$$\gamma_{2k-1} = \underbrace{Z \otimes \cdots \otimes Z}_{k-1 \text{ times}} \otimes X \otimes \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_{r-k \text{ times}}, \quad \gamma_{2k} = \underbrace{Z \otimes \cdots \otimes Z}_{k-1 \text{ times}} \otimes Y \otimes \underbrace{\mathbb{1} \otimes \cdots \otimes \mathbb{1}}_{r-k \text{ times}}, \quad k = 1 \dots r = n/2. \quad (6)$$

Summarizing these results (and applying standard theory, e.g. [Dav96, Cor. III.1.2]):

Proposition 2.15. ([GH82].) *Given graph Γ on n vertices, let $2r$ be the rank of its adjacency matrix over \mathbb{F}_2 , let $s = n - 2r$, and let $D = 2^{n-r}$. Then we may define a canonical D -dimensional representation $\pi : \mathfrak{C}(\Gamma) \rightarrow \mathbb{C}^{D \times D}$. This π_Γ may be expressed as $\bigoplus_{x \in \{\pm 1\}^s} \pi_{\Gamma,x}$, where each $\pi_{\Gamma,x}$ is irreducible and isomorphic to the Weyl–Brauer representation for $\mathfrak{C}(K_{2r})$ from Equation (6) (and hence is of dimension 2^r). Finally, any representation of $\mathfrak{C}(\Gamma)$ is unitarily equivalent to a direct sum $\bigoplus_{j=1}^m \pi_j$, where each π_j is a copy of one of the $\pi_{\Gamma,x}$'s.*

Remark 2.16. It follows from Proposition 2.15 that for any $h \in \mathfrak{C}(\Gamma)$ and any representation π we have $\text{tr}(\pi(h)) = \text{tr}(h)$ (as in Notation 2.11). In particular, $\text{tr}(gh) = \text{tr}(hg)$ always.

2.5 States, pseudostates, and pseudoexpectations

We recall some standard terminology:

Definition 2.17. We say that $h \in \mathfrak{C}(\Gamma)$ is *positive* if and only if it is a (Hermitian-)square, $h = g^*g$ for some $g \in \mathfrak{C}(\Gamma)$.

Definition 2.18. Let $\varrho : \mathfrak{C}(\Gamma) \rightarrow \mathbb{C}$ be a linear functional (which we take to include the condition $\varrho(h^*) = \varrho(h)^*$, so that $\varrho(h) \in \mathbb{R}$ when h is self-adjoint). We say that ϱ is *positive* if $\varrho(h) \in \mathbb{R}^{\geq 0}$ for all positive h . We furthermore say that ϱ is a *state* if $\varrho(1) = 1$.

Under the canonical π_Γ from Proposition 2.15, we see that any linear functional ϱ is of the form $h \mapsto \text{tr}(\rho \pi_\Gamma(h))$ for some self-adjoint $\rho \in \mathbb{C}^{D \times D}$. From this we infer that ϱ is positive if and only if ρ is positive semidefinite, and furthermore ϱ is a state if and only if $\text{tr}(\rho) = 1$ — i.e., if and only if ρ is a quantum state.⁸ Using π_Γ to pull back to $\mathfrak{C}(\Gamma)$, we are led to an equivalent definition:

⁸In this work we use the nonstandard convention in which quantum states ρ have $\text{tr}(\rho) = 1$; this is as opposed to the conventional normalization requiring $\text{Tr}(\rho) = 1$.

Definition 2.19. Abusing terminology, we also say that $\rho \in \mathfrak{C}(\Gamma)$ is a *state* if it is positive and has $\text{tr}(\rho) = 1$. In this case we use the notation $\mathbf{E}_\rho[h] = \text{tr}(\rho h)$, calling this the *expectation of h under ρ* .

Example 2.20. In the case of $\Gamma = \mathfrak{C}(\overline{K}_n)$, where all indeterminates commute, for each $x \in \{\pm 1\}^n$ the following is a state:

$$\rho_x = (1 + x_1 \chi_1)(1 + x_2 \chi_2) \cdots (1 + x_n \chi_n).$$

It satisfies $\mathbf{E}_{\rho_x}[h] = h(x)$. Under the canonical representation we have that $\pi_\Gamma(\rho_x) \in \mathbb{C}^{2^n \times 2^n}$ has a single nonzero entry in the (x, x) position, equal to 2^n .

For the sake of optimization, we will be interested in relaxing the notion of a linear functional ρ being positive. We begin with the following definition:

Definition 2.21. We say that $h \in \mathfrak{C}(\Gamma)$ is a *degree- $2k$ sum of (Hermitian-)squares* (abbreviated *SOS*) if it is expressible as $g_1^* g_1 + \cdots + g_m^* g_m$ for some polynomials $g_1, \dots, g_m \in \mathfrak{C}(\Gamma)$ of degree at most k .

Remark 2.22. Any positive $h \in \mathfrak{C}(\Gamma)$ is evidently degree- $2k$ SOS for some $k \leq n$ (see Equation (3)). Conversely, any SOS h is positive. This can be shown using the representation π_Γ from Proposition 2.15: if h is SOS then $\pi_\Gamma(h)$ is a sum of (Hermitian-)squares of matrices, hence $\pi_\Gamma(h)$ is PSD and thus of the form $G^* G$, hence $g := \pi_\Gamma^{-1}(G)$ has $g^* g = h$. Note, however, that h being low-degree SOS does not necessarily imply that h is the square of a low-degree polynomial.

The following simple fact shows that being degree- $2k$ SOS is characterized by a semidefinite program of size $n^{O(k)}$ (or $2^{O(n)}$ for $k = \Theta(n)$):

Fact 2.23. Let \vec{X} denote the vector whose entries are the monomials χ^S for all $|S| \leq k$. Then $h \in \mathfrak{C}(\Gamma)$ is degree- $2k$ SOS if and only if there is a complex PSD matrix H such that $\langle X | H | X \rangle = h$ within $\mathfrak{C}(\Gamma)$.

Definition 2.24. We define a *degree- $2k$ pseudoexpectation* to be a linear functional $\tilde{\mathbf{E}}[\cdot] : \mathfrak{C}(\Gamma) \rightarrow \mathbb{C}$ with $\tilde{\mathbf{E}}[1] = 1$ and $\tilde{\mathbf{E}}[h] \geq 0$ for all degree- $2k$ SOS h (or equivalently, $\tilde{\mathbf{E}}[g^* g] \geq 0$ for all g of degree at most k). As before, we may always write $\tilde{\mathbf{E}}[h] = \text{tr}(\rho h)$ for some self-adjoint $\rho \in \mathfrak{C}(\Gamma)$. We therefore call a self-adjoint $\rho \in \mathfrak{C}(\Gamma)$ a *degree- $2k$ pseudostate* if it has $\text{tr}(\rho) = 1$ and $\text{tr}(\rho h) \geq 0$ for all degree- $2k$ SOS h (equivalently, $\text{tr}(g \rho g^*) \geq 0$ for all g of degree at most k). In this case we use the notation $\tilde{\mathbf{E}}_\rho[h] = \text{tr}(\rho h)$.

Remark 2.25. Whether or not a $\tilde{\mathbf{E}}[\cdot]$ is a degree- $2k$ pseudoexpectation is completely determined by the values $\tilde{\mathbf{E}}[\chi^S]$ for $|S| \leq 2k$.

In light of the above, we have the following observation which shows that degree- $2k$ pseudoexpectations are characterized by a semidefinite program of size $n^{O(k)}$ (or $2^{O(n)}$ for $k = \Theta(n)$):

Fact 2.26. Let $\tilde{\mathbf{E}}[\cdot]$ be a linear functional on $\mathfrak{C}(\Gamma)$ with $\tilde{\mathbf{E}}[1] = 1$, and let M be the matrix having rows and columns indexed by the sets with $|S| \leq k$, and having (S, T) -entry equal to $\tilde{\mathbf{E}}[(\chi^S)^* \chi^T]$. Then $\tilde{\mathbf{E}}[\cdot]$ is a degree- $2k$ pseudoexpectation if and only if M is PSD.

2.6 Optimization

Given a self-adjoint polynomial $h \in \mathfrak{C}(\Gamma)$, we consider the following optimization problem:

$$\text{Opt}(h) := \max_{\text{states } \rho \in \mathfrak{C}(\Gamma)} \{\mathbf{E}_\rho[h]\}.$$

Recalling π_Γ from [Proposition 2.15](#) and the discussion in the previous section, we equivalently have

$$\begin{aligned}\text{Opt}(h) &= \max\{\text{tr}(\rho\pi_\Gamma(h)) : \rho \in \mathbb{C}^{D \times D}, \rho \text{ PSD}, \text{tr}(\rho) = 1\} \\ &= \max\{\langle \phi | \pi_\Gamma(h) | \phi \rangle : |\phi\rangle \in \mathbb{C}^D, \langle \phi | \phi \rangle = 1\} \\ &= \lambda_{\max}(\pi_\Gamma(h)).\end{aligned}$$

In physics terminology, if h is a Hamiltonian, then $\text{Opt}(h)$ is the ground state energy of $-h$ (under π_Γ). As a further equivalence, from [Proposition 2.15](#) it follows that

$$\text{Opt}(h) = \max\{\lambda_{\max}(\pi(h)) : \text{representations } \pi \text{ of } \mathfrak{C}(\Gamma)\};$$

in other words, the optimization problem seeks the assignment to the indeterminates χ_j (satisfying the (anti-)commutation relations Γ) that produces a matrix with largest possible eigenvalue. We will also sometimes consider the closely related problem of maximizing the operator norm; i.e., finding $\text{Opt}_\pm(h) = \max\{\text{Opt}(h), -\text{Opt}(-h)\}$. In fact, this makes sense even for non-self-adjoint h , if we define

$$\text{Opt}_\pm(h) := \sqrt{\text{Opt}(h^*h)}.$$

Example 2.27. In the case of most interest to us, $\mathfrak{C}(K_n)$ with n even, the representation π_Γ is the one using γ -matrices from [Equation \(6\)](#). This means that given the (coefficients of the) polynomial $h(\chi)$, the task is to find the largest eigenvalue of the implicitly determined $2^{n/2} \times 2^{n/2}$ matrix $h(\gamma)$.

Example 2.28. In the case of $\mathfrak{C}(\overline{K}_n)$, where all indeterminates commute, the task reduces to optimizing over all the 1-dimensional representations π_x (described prior to [Equation \(5\)](#)). That is, we are given a commutative polynomial $h(\chi)$ with real coefficients, and we are optimizing it over the Boolean hypercube,

$$\text{Opt}(h) = \max_{x \in \{\pm 1\}^n} \{h(x)\}.$$

Remark 2.29. Supposing $h = \sum_{|S|=q} a_S \chi^S \in \mathfrak{C}(\Gamma)$ with \vec{a} real, we may ask for a “naive” upper bound on $\text{Opt}_\pm(h)$. In the fully commuting case, $\Gamma = \overline{K}_n$, a reasonable answer is $|\vec{a}|_1$; this quantity is sometimes achievable (e.g., if each a_S is nonnegative, or if $q = 1$), and in the computer science literature $|\vec{a}|_1 = 1$ is a typical normalization when $q = 2$ (“Max-Cut”).

On the other hand, in the fully anticommuting case $\Gamma = K_n$, it seems that normalizing by $|\vec{a}| \equiv |\vec{a}|_2 = 1$ is most natural. For example, when $q = 1$ we have $h^*h = |\vec{a}|^2$ (see [Fact 2.14](#)) and hence $\text{Opt}_\pm(h) = |\vec{a}|$. In addition, normalization by $|\vec{a}|$ arises naturally in our generic upper bound using the Lovász Theta Function, presented in [Section 4](#) (see [Proposition 4.8](#)).

As we see from [Examples 2.27](#) and [2.28](#), given a degree- q polynomial h by its $n^{O(q)}$ coefficients, the naive algorithms for determining $\text{Opt}(h)$ have exponential complexity. This motivates understanding variants of $\text{Opt}(h)$ that are computable in polynomial time, foremost of which is the following “SOS relaxation”, which [Fact 2.26](#) shows is expressible as an SDP:

Definition 2.30. Given a self-adjoint $h \in \mathfrak{C}(\Gamma)$ of degree q , as well as $k \in \mathbb{N}^+$ with $2k \geq q$, we define

$$\text{SOS}_{2k}(h) = \max\{\tilde{\mathbf{E}}_\rho[h] : \tilde{\mathbf{E}}[\cdot] \text{ is a degree-}2k \text{ pseudoexpectation}\}. \quad (7)$$

This is a nonincreasing function of k , with $\text{SOS}_{2n}(h) = \text{Opt}(h)$.

Alternatively, we may seek to certify $\text{Opt}(h) \leq \beta$ by expressing $\beta - h$ as a degree- $2k$ sum of (Hermitian-)squares. Finding the smallest β for which this is possible is the dual SDP for SOS_{2k} , as is well known. Furthermore, using the fact that we have $\chi_j^2 = 1$ for all j , it is not hard to show that there is no duality gap (see, e.g., [JH16], with the noncommutativity of our setting not affecting the proof). Thus:

Theorem 2.31. *Given a self-adjoint $h \in \mathfrak{C}(\Gamma)$ of degree q , as well as $k \in \mathbb{N}^+$ with $2k \geq q$, we have*

$$\text{SOS}_{2k}(h) = \min\{\beta : \beta - h \text{ is degree-}2k \text{ SOS}\}. \quad (8)$$

Fact 2.23 explains why this characterization is also an SDP.

2.7 Complexity theory

Regarding the optimization problem $\text{Opt}(h)$, there are several computational tasks we will want to consider. But first we must cover some minor technical details.

Complexity-theoretic niceties. The input is always considered to be the coefficients of h , explicitly given as rational (complex) numbers, as well as the graph Γ if necessary (i.e., if it isn't considered fixed to be K_n or \overline{K}_n). Most often we have a fixed constant degree q in mind for h , in which case the number of coefficients of h is bounded by $m = O(n^q)$. For the sake of scaling, we typically prefer to assume the normalization $\sum_S |a_S|^2 = 1$ (where h is written as in [Equation \(3\)](#)). With rational coefficients perhaps only $\sum_S |a_S|^2 \approx 1$ will be possible, but anyway we will never be interested in cases other than $1/\text{poly}(n) \leq \sum_S |a_S|^2 \leq \text{poly}(n)$. In light of this (and the fact that $\|\pi(X^S)\| \leq 1$ for all S), perturbing any coefficient a_S by a sufficiently small $\pm 1/\text{poly}(n)$ will have only negligible effect on $\text{Opt}(h)$, and we may therefore assume that all coefficients a_S are expressed using $O(\log n)$ bits. (In particular, when we consider Gaussian coefficients J_S , we may assume they are rounded to $O(\log n)$ bits of precision.) Thus the total input size L will generally be $O(n^q \log n)$, plus $O(n^2)$ if Γ needs to be specified. Finally, when we speak of “computing $\text{Opt}(h)$ ”, or “solving a semidefinite program”, we mean don't mean literally exactly, but rather up to an additive precision of any arbitrarily small $1/\text{poly}(n)$. Indeed, the SDP for $\text{SOS}_{2k}(h)$ can be computed to additive precision $\exp(-n^k)$ in $n^{O(k)}$ time.

Given an instance h (and graph Γ), there are several possible computational tasks concerning $\text{Opt}(h)$ (or $\text{Opt}_{\pm}(h)$):

Exact computation. Determine $\text{Opt}(h)$. (But this is QMA-hard.)

Upper-bound certification. Output $\beta \in \mathbb{R}$ such that $\text{Opt}(h) \leq \beta$.

Lower-bound certification. Output $\alpha \in \mathbb{R}$ such that $\text{Opt}(h) \geq \alpha$.

Naturally, one wants β as small as possible and α as large as possible. One might also want to upgrade a lower-bound certification algorithm so that it also provides a *witness* to $\text{Opt}(h) \geq \alpha$. We consider two possibilities:

Implicit lower-bound witnessing. Output a classical description⁹ of a state ρ achieving $\mathbf{E}_{\rho}[h] \geq \alpha$.

⁹Here it is only fair to allow for a fixed “encoding scheme” $\text{ENC} : \{\text{bit-strings}\} \rightarrow \mathfrak{C}(\Gamma)$ with the following properties: (i) $\text{ENC}(x)$ is a valid state for any bit-string x ; (ii) there is an algorithm that, given $h \in \mathfrak{C}(\Gamma)$ by its

Quantum lower-bound witnessing. Output an actual quantum state $\pi_\Gamma(\rho) \in \mathbb{C}^{D \times D}$ of $n - r$ qubits (under the canonical representation from [Proposition 2.15](#)) achieving $\mathbf{E}_\rho[h] \geq \alpha$.

The reader should bear in mind that for the implicit lower-bound witnessing problem (with fixed encoding scheme), there may not even *be* a state ρ that simultaneously has a $\text{poly}(n)$ -size description and a particularly large value for $\mathbf{E}_\rho[h]$. Similarly, for the quantum lower-bound witnessing problem, for a given h there may not *be* a $\text{poly}(n)$ -size quantum circuit that outputs a ρ with particularly large $\mathbf{E}_\rho[h]$. This latter possibility is related to the fermionic NLTS Conjecture discussed in [Section 1.3](#).

2.7.1 On randomized algorithms and randomized inputs

In this work we will often be interested in *average-case* complexity, where the input for the task is \mathbf{h} drawn from some distribution \mathcal{H} on instances (typically $\mathcal{H} = \text{SYK}_q$ for some q). In this case we should be careful to define the correctness of algorithms.

Except for one case (to be discussed shortly), each certification/witnessing algorithm presented in this paper is deterministic. Such a deterministic algorithm ALG will have the following properties: On all inputs h , algorithm ALG outputs either a *claim* (such as “ $\text{Opt}(h) \leq \sqrt{n}$ ”) or else outputs “FAIL”. And furthermore:

- (i) For *all* inputs h , if $\text{ALG}(h)$ is a claim, then that claim is correct.
- (ii) $\Pr_{\mathbf{h} \sim \mathcal{H}}[\text{ALG}(\mathbf{h}) = \text{“FAIL”}] < \epsilon$ (for some small ϵ).

We emphasize two things: On one hand, the user of the ALG gets 100% confidence in its claims. On the other hand, for a given outcome $\mathbf{h} = h$, a user *cannot* “boost” the probability of a non-FAIL outcome by repeatedly running the algorithm. (We also remark that for the task of implicit lower-bound witnessing, there is never a concern about confidence; we assume that from the output description of ρ one can efficiently and deterministically compute $\mathbf{E}_\rho[h]$.)

The one exception to this deterministic style of algorithm will be our quantum lower-bound witnessing for $\mathbf{h} \sim \text{SYK}_4(n)$. In [Section 8](#) we first show that

$$\Pr_{\mathbf{h} \sim \text{SYK}_4(n)}[\text{Opt}(\mathbf{h}) \geq c_0 \sqrt{n}] \geq 1 - \exp(-\Omega(n))$$

(for some universal constant $c_0 > 0$). Following that we give an efficient quantum “witnessing” algorithm W that, given any input h , applies a sequence of polynomially-many gates to produce a quantum state ρ . One may think of W *hypothesizing* that $\mathbf{E}_\rho[h] \geq c_0 \sqrt{n}$.

Now on one hand, there is no obvious *deterministic* way to test this hypothesis, even with a quantum computer. On the other hand, there *is* an obvious *randomized* certification algorithm \mathbf{V} that runs efficiently on a quantum computer. Given h , algorithm \mathbf{V} simply repeatedly prepares $W(h) = \rho$ and empirically estimates $\mathbf{E}_\rho[h]$. By running \mathbf{V} for $\text{poly}(n) \log(1/\delta)$ time, a user can obtain confidence $1 - \delta$ that the hypothesis $\text{Opt}(h) \geq c_0 \sqrt{n}$ is true (up to any additive $1/\text{poly}(n)$,

coefficients as well as some x , efficiently evaluates $\mathbf{E}_{\text{ENC}(x)}[h]$. In the fully commuting case of $\mathfrak{C}(\overline{K}_n)$, the mapping $x \mapsto \rho$ from [Example 2.20](#) is natural. In the fully anticommuting case of $\mathfrak{C}(K_n)$, there is an analogous natural scheme for encoding the *Gaussian states*, as described in [Section 7](#). The requirement (ii) is necessary as otherwise one could simply declare that the state is $\exp(\beta p) |\phi\rangle$ for large $\beta > 0$ and for ϕ chosen to have nontrivial projection on the eigenspace of maximal eigenvalue; as $\beta \rightarrow \infty$, this state converges to that eigenspace, but we cannot compute expectation values efficiently. Another interesting state that does not seem to meet our requirements is the coupled cluster variational state of [\[HTS21\]](#).

which we can neglect since we aren't specifying c_0 precisely anyway). That is, for any user-selected δ , we can obtain a $\text{poly}(n) \log(1/\delta)$ -time composite quantum algorithm **ALG** that, on input h , either outputs "FAIL", or else outputs a claim of $\text{Opt}(h) \geq c_0 \sqrt{n}$ together with a witnessing quantum state ρ . It has the following properties:

- (i') For *all* inputs h , if **ALG**(h) outputs a claim, then that claim is correct except with probability at most δ (over the internal randomness of **ALG**).
- (ii') $\Pr_{\mathbf{h} \sim \text{SYK}_4(n)}[\mathbf{ALG}(\mathbf{h}) = \text{"FAIL"}] < \exp(-\Omega(n))$.

In summary, with very high probability over the random *input* \mathbf{h} , the algorithm produces a ρ and the claim that $\mathbf{E}_\rho[\mathbf{h}] \geq \Omega(\sqrt{n})$. Furthermore, for *any* input h resulting in a claim, the user can verify this claim with confidence that can be efficiently tuned arbitrarily close to 1.

We end this section by remarking that for $\mathbf{h} \sim \text{SYK}_4(n)$, we currently do not know any efficient *classical* algorithm for certifying $\text{Opt}(\mathbf{h}) \geq \Omega(\sqrt{n})$ (even without a witness) that has properties (i'), (ii') above. Thus we arguably have a distributional task, with classical input and output, that is efficiently solvable by a quantum computer, but for which we currently do not know an efficient classical algorithm.

3 Upper bounds and heuristics for random polynomials

In this section we recall (and generalize) the known upper bound on $\mathbf{E}[\text{Opt}(\mathbf{h})]$ for $\mathbf{h} \sim \text{SYK}_q(n)$, and also sketch the heuristic from mathematical physics that suggests its asymptotic value.

Before this, we recall that analyzing the expected optimum is sufficient for understanding the high-probability optimum. To be precise, note for $\mathbf{p} \sim \text{SYK}_q(n)$ that $\text{Opt}(\mathbf{p})$ and $\text{Opt}_\pm(\mathbf{p})$ are $1/\sqrt{\binom{n}{q}}$ -Lipschitz functions of the random coefficients. Thus standard concentration results for Gaussian measure (see, e.g., [Led01, Ch. 2]) imply the following:

Proposition 3.1. *For $\mathbf{p} \sim \text{SYK}_q(n)$ it holds that*

$$\Pr[|\text{Opt}(\mathbf{p}) - \mathbf{E}[\text{Opt}(\mathbf{p})]| \geq \epsilon] \leq 2 \exp(-\binom{n}{q} \epsilon^2 / 2)$$

(and the same holds for $\text{Opt}_\pm(\mathbf{p})$). E.g., for any constant $q \geq 2$ there is a constant C such that

$$\Pr[|\text{Opt}(\mathbf{p}) - \mathbf{E}[\text{Opt}(\mathbf{p})]| \geq C/\sqrt{n}] \leq \exp(-n).$$

3.1 Upper bounds for random polynomials

On the subject of upper bounds for $\text{Opt}(\mathbf{p})$ when $\mathbf{p} \sim \text{SYK}_q(n)$, the best provable upper bound that is known is $\mathbf{E}[\text{Opt}_\pm(\mathbf{p})] \leq \sqrt{\ln 2} \sqrt{n}$. One way to prove this¹⁰ is to use the Matrix Chernoff Bound [AW02, Oli10]. Since the γ -matrices from Equation (6) are in dimension $D = 2^{n/2}$, and since we have $\|\gamma_{j_1} \cdots \gamma_{j_q}\|_{\text{op}} \leq 1$ always, the bound [Tro12, Thm. 1.2] gives

$$\Pr_{\mathbf{p} \sim \text{SYK}_q(n)}[\text{Opt}(\mathbf{p}) \geq t] \leq 2^{n/2} \exp(-t^2/2).$$

From this, $\mathbf{E}[\text{Opt}(\mathbf{p})] \leq (\sqrt{\ln 2} + o(1))\sqrt{n}$ follows (and we only need \mathbf{p} to have centered, subgaussian random coefficients, of total variance 1). Indeed, Feng, Tian, and Wei [FTW19] obtain the exact

¹⁰As was pointed out to us by several attendees, including Jerry Li and Cristopher Moore, at the Simons Institute workshop *Rigorous Evidence for Information-Computation Trade-offs*.

bound $\mathbf{E}[\text{Opt}_\pm(\mathbf{p})] \leq \sqrt{\ln 2} \sqrt{n}$ via a simple proof, which is possible because the matrices $\gamma_{j_1} \cdots \gamma_{j_q}$ have uniformly bounded operator norm. In the following [Theorem 3.2](#), we give a generalization of this proof, which we will need (ironically) for our *lower* bound in [Section 8](#).

Theorem 3.2. *Given graph $\Gamma = ([n], E)$, suppose that*

$$\mathbf{h}(\chi) = \sum_S \mathbf{J}_S \chi^S \in \mathfrak{C}(\Gamma)$$

is self-adjoint, where the coefficients $(\mathbf{J}_S)_{S \subseteq [n]}$ are real random variables (not necessarily independent). Then provided conditions (i), (ii) below hold, we have

$$\Pr[|\mathbf{h}(1)| \geq \beta] \leq 1/D \implies \mathbf{E}[\text{Opt}_\pm(\mathbf{h})] \leq \beta,$$

where D is the dimension of the canonical representation π_Γ from [Proposition 2.15](#).

The conditions are:

- (i) $\Pr[|\mathbf{h}(1)| \geq \beta] \leq 1/D$ holds due to the moment bound $\mathbf{E}[\mathbf{h}(1)^k]/\beta^k \leq 1/D$ for some even k ;
- (ii) $\mathbf{E}[\mathbf{J}_{S_1} \mathbf{J}_{S_2} \cdots \mathbf{J}_{S_k}] \geq 0 \ \forall S_1, \dots, S_k \subseteq [n]$.

Proof. This follows from

$$\begin{aligned} \mathbf{E}[\text{Opt}_\pm(\mathbf{h})]^k &\leq \mathbf{E}[\text{Opt}_\pm(\mathbf{h})^k] = \mathbf{E}[\|\pi_\Gamma(\mathbf{h})\|^k] \leq \mathbf{E}[\text{Tr}(\pi_\Gamma(\mathbf{h})^k)] = D \mathbf{E}[\text{tr}(\pi_\Gamma(\mathbf{h}^k))] \\ &= D \sum_{S_1, \dots, S_k \subseteq [n]} \mathbf{E}[\mathbf{J}_{S_1} \cdots \mathbf{J}_{S_k}] \text{tr}(\pi_\Gamma(\chi^{S_1} \cdots \chi^{S_k})) \stackrel{\text{(ii)}}{\leq} D \sum_{S_1, \dots, S_k \subseteq [n]} \mathbf{E}[\mathbf{J}_{S_1} \cdots \mathbf{J}_{S_k}] = D \mathbf{E}[\mathbf{h}(1)^k] \stackrel{\text{(i)}}{\leq} \beta^k, \end{aligned}$$

where the inequality marked “(ii)” also uses that each normalized-trace is at most 1, since $\pi_\Gamma(\chi^{S_1} \cdots \chi^{S_k})$ is a product of matrices $\pi_\Gamma(\chi_j)$ that square to the identity and hence have operator norm at most 1. \square

Remark 3.3. In fact, the proof gives $\mathbf{E}[\text{Opt}_\pm(\mathbf{p})^k] \leq \beta^k$, and hence $\Pr[\text{Opt}_\pm(\mathbf{p}) \geq \beta'] \leq (\beta/\beta')^k$ for any $\beta' \geq \beta$.

Corollary 3.4. *For $\mathbf{h} \sim \text{SYK}_q(n)$ we have $\mathbf{E}[\text{Opt}(\mathbf{h})] \leq (\sqrt{\ln 2} + o(1))\sqrt{n}$. More generally, this holds provided the coefficients $(\mathbf{J}_S : |S| = q)$ of \mathbf{h} are independent, centered, and subgaussian with parameter $1/\binom{n}{q}$ (e.g., they may be independent Rademachers, times $1/\sqrt{\binom{n}{q}}$).*

Proof. In the setting of [Theorem 3.2](#) we have $\Gamma = K_n$ and so $D = 2^{n/2}$. Condition (ii) holds by the independence and centeredness ($\mathbf{E}[\mathbf{J}_S] = 0$) of the coefficients. In the usual Gaussian case, $\mathbf{h}(1)$ is just a standard Gaussian random variable, hence it has $\Pr[|\mathbf{h}(1)| \geq \beta] \leq 2^{-n/2}$ for $\beta \sim \sqrt{\ln 2} \sqrt{n}$. Further, this can be shown by taking $k \sim (\ln 2)n$ to be an even integer and using the formula $\mathbf{E}[\mathbf{h}(1)^k] = (k-1)!! \leq \sqrt{2}(k/e)^{k/2}$. The subgaussian case also follows. \square

Remark 3.5. The bound $O(\sqrt{n})$ in [Corollary 3.4](#) is optimal up to constant factors for $q = 2, 4$ (we prove this for $q = 4$ in [Section 8](#)). It is also expected to be optimal up to constant factors for larger even q . However, it is not expected to be optimal for odd q , where it is expected that $\mathbf{E}[\text{Opt}(\mathbf{h})] \leq O(1)$. See the heuristics discussed in [Section 3.2](#).

Remark 3.6. If the coefficients \mathbf{J}_S are not subgaussian but have reasonably small moments, similar bounds can be obtained.

Remark 3.7. For $\text{SYK}_4(n)$, Feng–Tian–Wei [FTW19] obtained the exact bound $\mathbf{E}[\text{Opt}(\mathbf{h})] \leq \sqrt{\ln 2} \sqrt{n}$ by performing the proof with a convex combination of all integer k .

Remark 3.8. As with upper bounds for random classical CSPs based on Chernoff, this upper bound is an excellent example of the sort that does *not* obviously yield SOS proofs, or indeed any kind of succinct certificate. It shows that before we draw \mathbf{h} , we know it is very likely — say, a 99% chance — that $\text{Opt}_{\pm}(\mathbf{h})$ will be at most 100β . However once we *fix* a randomly drawn outcome $\mathbf{h} = h$, there is no obvious certificate that this *particular* h has $\text{Opt}_{\pm}(h) \leq 100\beta$.

Remark 3.9. In [Theorem 3.2](#) we assumed that the coefficients of \mathbf{h} are indexed by *subsets* $S \subseteq [n]$, with χ^S standing for $\chi_{k_1} \cdots \chi_{k_q}$ when $S = \{k_1, \dots, k_q\}$ with $k_1 < \dots < k_q$. However one can see that the proof goes through equally well if S is allowed to stand for an *ordered sequence* from $[n]$. Allowing this may assist in fulfilling condition (ii).

One scenario where we will be using [Theorem 3.2](#) (together with [Remark 3.9](#)) is the following: \mathbf{h} is a certain random polynomial of low degree, where each coefficient is of the form $c\mathbf{G}_{j_1} \cdots \mathbf{G}_{j_r}$, where $c \geq 0$ and $\mathbf{G}_1, \dots, \mathbf{G}_N$ are independent standard Gaussians. In this case, condition (ii) of [Theorem 3.2](#) holds, and we are reduced to understanding tail/moment bounds for polynomials in Gaussians (of the usual, commutative) kind. For this, we will use Latała’s Theorem on Gaussian chaoses [Lat06] (see also [Leh11]):

Theorem 3.10. ([Lat06].) *Let $A \in \mathbb{R}^{N \times \dots \times N}$ be a q -dimensional array of reals, with $A_{j_1 \dots j_q} = 0$ when j_1, \dots, j_q are not all distinct; and, let*

$$f(x_1, \dots, x_N) = \sum_{j_1, \dots, j_q=1}^N A_{j_1 \dots j_q} \prod_{t=1}^q x_{j_t}. \quad (9)$$

If $\mathbf{G} = (\mathbf{G}_1, \dots, \mathbf{G}_N)$ denotes a sequence of independent standard Gaussians, then for any real $k \geq 2$,

$$\|f(\mathbf{G})\|_k = \mathbf{E}[|f(\mathbf{G})|^k]^{1/k} \leq c_q \cdot \max_{\mathcal{P}} \{k^{|\mathcal{P}|/2} \|A\|_{\mathcal{P}}\},$$

where c_q is a constant depending only on q , the maximum is over all partitions $\mathcal{P} = (P_1, \dots, P_s)$ of $[q]$ into nonempty parts, and

$$\|A\|_{(P_1, \dots, P_s)} := \max \left\{ f(x) : x \in \mathbb{R}^N, \sum_{p \in P_r} x_p^2 \leq 1 \ \forall r \in [s] \right\}.$$

Indeed we have $\|f(\mathbf{G})\|_k \leq c'_q \|\tilde{f}(\mathbf{G}^{(1)}, \dots, \mathbf{G}^{(q)})\| \leq c_q \max_{\mathcal{P}} \{k^{|\mathcal{P}|/2} \|A\|_{\mathcal{P}}\}$, where \tilde{f} denotes the decoupled version of [Equation \(9\)](#), in which x_{j_t} is replaced with the new variable $x_{j_t}^{(t)}$ (and $\mathbf{G}^{(1)}, \dots, \mathbf{G}^{(q)}$ are independent sequences of N independent standard Gaussians).

3.2 Heuristics for the SYK optimum

Here we describe the heuristic developed in the physics community for predicting $\text{Opt}(\mathbf{h})$ when $\mathbf{h} \sim \text{SYK}_q(n)$. We follow the discussion in [GJV18]; see also [FTW19, FTW18, FTW20] for partial formalizations of this heuristic.

Naturally, one can attempt to analyze $\text{Opt}_{\pm}(\mathbf{h})$, and indeed the entire typical spectrum of \mathbf{h} , via the (expected) Trace Method. However, the combinatorics arising when computing $\mathbf{E}[\text{tr}(\mathbf{h}^k)]$ are rather intricate. A heuristic one can use is the following. From [Fact 2.13](#) we see that if \mathbf{S} and \mathbf{T}

are chosen randomly and independently from the $N := \binom{n}{q}$ possible monomials, they anticommute with probability

$$p = p(q, n) = \mathbf{Pr}[q - |\mathbf{S} \cap \mathbf{T}| \text{ is odd}] \approx \begin{cases} \mathbf{Pr}[|\mathbf{S} \cap \mathbf{T}| = 1] \approx q^2/n & \text{for } q = o(\sqrt{n}) \text{ even,} \\ \mathbf{Pr}[|\mathbf{S} \cap \mathbf{T}| = 0] \approx 1 - q^2/n & \text{for } q = o(\sqrt{n}) \text{ odd,} \end{cases}.$$

(Of course, one can easily make this asymptotic approximation more precise if desired.) Given this, the key ansatz is to let $\mathbf{\Gamma} \sim \mathfrak{G}(N, p)$ denote an Erdős–Rényi random graph, and to model the N monomials in $\mathfrak{C}(K_n)$ by the N indeterminates of $\mathfrak{C}(\mathbf{\Gamma})$. That is, for

$$\ell = \frac{1}{\sqrt{N}} \sum_{j=1}^N \mathbf{a}_j \chi_j \in \mathfrak{C}(\mathbf{\Gamma}), \quad \mathbf{a}_j \sim \mathcal{N}(0, 1),$$

the heuristic is

$$\mathbf{E}[\text{tr}(\mathbf{h}^k)] \approx \mathbf{E}_{\mathbf{\Gamma}, \mathbf{a}_i\text{'s}}[\text{tr}(\ell^k)]. \quad (10)$$

Estimating the right side of [Equation \(10\)](#) is more computationally tractable for constant k , and the resulting moment estimates match those of the so-called μ -Gaussian (or μ -semicircular) distribution, with orthogonal polynomials the μ -Hermite polynomials (see, e.g., [\[ISV87\]](#)).¹¹ Here

$$\mu = \mu(q, n) := 1 - 2p = \mathbf{E}_{\mathbf{S}, \mathbf{T} \sim \binom{[n]}{q}}[(-1)^{q - |\mathbf{S} \cap \mathbf{T}|}] \in [-1, +1],$$

and one easily check that for $q = O(\sqrt{n})$, up to lower-order terms we have

$$\mu \approx (-1)^q \exp(-2r), \quad r := q^2/n.$$

Thus the heuristic prediction is that the spectral density of $\mathbf{h} \sim \text{SYK}_q(n)$ will resemble that of the μ -Gaussian density, call it ν_μ . This density ν_μ has an explicit analytic formula (see, e.g., [\[BKS97\]](#)), but rather than reproducing it, we will merely note the following facts:

- For $\mu \rightarrow 1$ (corresponding to $q = o(\sqrt{n})$ even), the density ν_μ tends to the standard Gaussian.
- For $\mu \in (-1, 1)$, the support of ν_μ is the interval $[-\frac{2}{\sqrt{1-\mu}}, +\frac{2}{\sqrt{1-\mu}}]$.
- For $\mu = 0$ (equivalently, $q = \omega(\sqrt{n})$), ν_μ is the semicircle density.
- For $\mu \rightarrow -1$ (corresponding to $q = o(\sqrt{n})$ odd), the density ν_μ tends to Rademacher.

Summarizing these heuristics, the belief is that with high probability,

$$\text{for } \mathbf{h} \sim \text{SYK}_q(n), \quad \text{Opt}_\pm(\mathbf{h}) \sim \frac{2}{\sqrt{1 - (-1)^q \exp(-2q^2/n)}} \sim \begin{cases} \frac{\sqrt{2}}{q} \sqrt{n} & \text{for } q = o(\sqrt{n}) \text{ even,} \\ \sqrt{2} & \text{for } q = o(\sqrt{n}) \text{ odd.} \end{cases} \quad (11)$$

¹¹These are usually called q -Hermite polynomials in the combinatorics literature, according to the usual notion of q -deformity. Unfortunately we have a notational clash with the SYK degree “ q ”. In the physics literature this is resolved by writing “ Q -Hermite”, but we follow the “ μ -Hermite” terminology of Speicher [\[Spe92\]](#) instead.

4 Upper bound certification with the Lovász Theta Function

We first recall the independence number of a graph:

Notation 4.1. If $\Gamma = ([n], E)$ is a graph, $\alpha(\Gamma)$ denotes the cardinality of its largest independent set; i.e., the largest $S \subseteq [n]$ such that no $\{j, k\} \in E$ has both $j, k \in S$.

In the context of $\mathfrak{C}(\Gamma)$, an independent set S is one for which the indeterminates $(\chi_j : j \in S)$ pairwise commute. Lovász [Lov79] introduced the following upper bound on $\alpha(\Gamma)$, which is efficiently computable using semidefinite programming:

Notation 4.2. The *Lovász Theta Function* assigns to each graph $\Gamma = ([n], E)$ the number

$$\vartheta(\Gamma) = \max_{\rho \in \mathbb{R}^{n \times n}} \{ \text{Tr}(\rho \mathbb{J}) : \rho \geq 0, \text{Tr}(\rho) = 1, \rho_{jk} = 0 \ \forall \{j, k\} \in E \}, \quad (12)$$

where \mathbb{J} denotes the all-1's matrix.

Fact 4.3. There is also an equivalent dual SDP definition,

$$\vartheta(G) = \min \{ \kappa : \exists \Sigma \in \mathbb{R}^{n \times n}, \Sigma_{jj} = 1 \ \forall j \in [n], \Sigma_{jk} = 0 \ \forall \{j, k\} \notin E, \kappa \Sigma \geq \mathbb{J} \}. \quad (13)$$

Remark 4.4. The “weak duality” bound (12) \leq (13), which is all we will require, is evident from

$$\text{Tr}(\rho \mathbb{J}) \leq \kappa \text{Tr}(\rho \Sigma) = \kappa \text{Tr}(\rho) = \kappa.$$

Fact 4.5. $\alpha(\Gamma) \leq \vartheta(\Gamma)$ always holds: if S is an independent set, and $|S\rangle \in \{0, 1\}^n$ denotes its indicator, then $\rho = \frac{1}{|S|} |S\rangle \langle S|$ achieves $|S|$ in Equation (12).

Given a graph Γ , we will be interested in the maximum that $\text{Opt}_{\pm}(\ell)$ can possibly be among self-adjoint linear polynomials $\ell \in \mathfrak{C}(\Gamma)$ (subject to a scaling constraint on their coefficients). We therefore introduce the following:

Definition 4.6. For a graph $\Gamma = ([n], E)$, we define

$$\Psi(\Gamma) = \max \left\{ \text{Opt}_{\pm}(\ell)^2 = \text{Opt}(\ell^* \ell) = \text{Opt}(\ell^2) : \ell = \sum_{j=1}^n a_j \chi_j \in \mathfrak{C}(\Gamma), \vec{a} \in \mathbb{R}^n, |\vec{a}|^2 = 1 \right\}.$$

The below two propositions now establish the following sandwiching bound:

$$\alpha(\Gamma) \leq \Psi(\Gamma) \leq \vartheta(\Gamma).$$

Proposition 4.7. For any graph $\Gamma = ([n], E)$ we have $\Psi(\Gamma) \geq \alpha(\Gamma)$.

Proof. If S is a maximum independent set in Γ , then the matrices $(\pi_{\Gamma}(\chi_j) : j \in S)$ pairwise commute and hence can be simultaneously diagonalized. Take any common eigenvector; it will have associated eigenvalue $\lambda_j \in \{\pm 1\}$ for $j \in S$. Now $\ell = \frac{1}{\sqrt{|S|}} \sum_{j \in S} \lambda_j \chi_j$ has an eigenvalue of $\sqrt{|S|} = \sqrt{\alpha(\Gamma)}$ under π_{Γ} . \square

Proposition 4.8. For any graph $\Gamma = ([n], E)$ we have $\Psi(\Gamma) \leq \vartheta(\Gamma)$, and in fact this has a degree-2 SOS proof. That is, if $\ell = \sum_{j=1}^n a_j \chi_j \in \mathfrak{C}(\Gamma)$ with $\vec{a} \in \mathbb{R}^n$ and $|\vec{a}| = 1$, then $\text{SOS}_2(\ell^* \ell) \leq \vartheta(\Gamma)$. Moreover, this holds even without assuming the commutation relations $\chi_j \chi_k = \chi_k \chi_j$ for $\{j, k\} \notin E$; i.e., it only uses the anticommutation relations $\chi_j \chi_k = -\chi_k \chi_j$ for $\{j, k\} \in E$.

Proof. Let $\tilde{\mathbf{E}}$ be any degree-2 pseudoexpectation for $\mathfrak{C}(\Gamma)$, which we may also think of as a PSD matrix in $\mathbb{C}^{n \times n}$. Now given $\ell = \sum_{j=1}^n a_j \chi_j$, with $|\vec{a}| = 1$, define the PSD matrix $\rho = |a\rangle\langle a| \circ \tilde{\mathbf{E}}$, where \circ denotes the Hadamard (entry-wise) product; i.e.,

$$\rho_{jk} = a_j a_k \tilde{\mathbf{E}}[\chi_j \chi_k].$$

Then $\text{Tr}(\rho) = |\vec{a}|^2 = 1$ and $\text{Tr}(\rho \mathbb{I}) = \tilde{\mathbf{E}}[\ell^* \ell]$. Furthermore, observe that for $\{j, k\} \in E$ we have $\tilde{\mathbf{E}}[\chi_j \chi_k]^* = \tilde{\mathbf{E}}[\chi_k^* \chi_j^*] = -\tilde{\mathbf{E}}[\chi_j \chi_k]$; thus $\tilde{\mathbf{E}}[\chi_j \chi_k]$ and ρ_{jk} are purely imaginary. If we now let $\hat{\rho}$ be the real part of ρ , we still have all of the following properties —

$$\hat{\rho} \geq 0, \quad \text{Tr}(\hat{\rho}) = 1, \quad \text{Tr}(\hat{\rho} J) = \tilde{\mathbf{E}}[\ell^* \ell]$$

— and now $\hat{\rho}$ also has $\hat{\rho}_{jk} = 0$ for $\{j, k\} \in E$. This establishes $\tilde{\mathbf{E}}[\ell^* \ell] \leq \vartheta(\Gamma)$, as required. \square

Remark 4.9. Regarding tightness of the bounds $\alpha(\Gamma) \leq \Psi(\Gamma) \leq \vartheta(\Gamma)$, the 5-cycle $\Gamma = C_5$ is a case where $\Psi(\Gamma) < \vartheta(\Gamma)$ strictly. In this case, famously $\vartheta(C_5) = \sqrt{5}$ [Lov79], but one can show $\Psi(C_5) = 2 = \alpha(C_5)$. Briefly, in $\mathfrak{C}(C_5)$ we may write any ℓ under π_Γ as

$$\ell = a_1 \cdot X \otimes \mathbb{1} \otimes \mathbb{1} + a_2 \cdot Y \otimes \mathbb{1} \otimes \mathbb{1} + a_3 \cdot X \otimes X \otimes \mathbb{1} + a_4 \cdot \mathbb{1} \otimes Y \otimes \mathbb{1} + a_5 \cdot Y \otimes Z \otimes Z.$$

Then one may compute that the eigenvalues of ℓ^2 are (4 copies each of) $\sigma \pm 2\sqrt{\tau}$, where $\sigma := \sum_{j=1}^5 a_j^2$ and $\tau := \sum_{\{j,k\} \notin E} a_j^2 a_k^2$, with E is the edge-set of C_5 . Our normalization gives $\sigma = 1$, and subject to this it is elementary to show that $\tau \leq \frac{1}{4}$, and hence $\lambda_{\max}(\ell^2) \leq 1 + 2\sqrt{\frac{1}{4}} = 2$.

On the other hand, we do not know a particular Γ such that $\Psi(\Gamma) > \alpha(\Gamma)$ strictly. Indeed, in [Appendix A](#) we consider the first and second derivatives of $\text{Opt}(\ell)$, showing that the first derivative vanishes and the Hessian is negative semidefinite at the ℓ given in the proof of [Proposition 4.7](#).

4.1 Implications for higher-degree polynomials

The Lovász Theta upper bound from [Proposition 4.8](#) can also be used to bound $\text{Opt}_\pm(h)$ for nonlinear polynomials h , simply by treating each monomial appearing in h as a new indeterminate. Let us work this out in the case of greatest interest to us: homogeneous polynomials over pairwise anticommuting indeterminates, as in [Section 2.3](#). Suppose

$$h = i^{(g)} \sum_{\substack{S \subseteq [n] \\ |S|=q}} a_S \chi^S \in \mathfrak{C}(K_n), \quad \text{with } \vec{a} \in \mathbb{R}^N \text{ satisfying } |\vec{a}| = 1, \text{ where } N := \binom{n}{q}. \quad (14)$$

Let us introduce N new indeterminates $\underline{\chi}^S$, where $\underline{\chi}^S$ stands for $i^{(g)} \chi^S$; these are self-adjoint and square to 1. Recalling from [Fact 2.13](#) how these monomials commute/anticommute leads us to define the following generalized Kneser-type graph (see the subsequent [Definition 4.20](#) for a more general definition):

Definition 4.10. Let $KG_{\text{even}}^{(n,q)}$ denote the graph on vertex set $\binom{n}{q}$ in which $\{S, T\}$ is a *nonedge* if and only if the “distance” $d = q - |S \cap T|$ is even.

We may now regard h from [Equation \(14\)](#) as a self-adjoint, linear homogeneous element of $\mathfrak{C}(KG_{\text{even}}^{(n,q)})$, and any upper bound on $\text{Opt}_\pm(h)$ in this setting is a valid upper bound on $\text{Opt}_\pm(h)$ in the original $\mathfrak{C}(K_n)$ setting. (This is because there are only more relations among the actual monomials χ^S , as compared to the new indeterminates $\underline{\chi}^S$.) We may therefore conclude from [Proposition 4.8](#):

Proposition 4.11. *For h as in Equation (14) we have*

$$\text{Opt}_{\pm}(h)^2 = \text{Opt}(h^*h) \leq \text{SOS}_{2q}(h^*h) \leq \vartheta(KG_{\text{even}}^{(n,q)}).$$

This raises the question of determining $\vartheta(KG_{\text{even}}^{(n,q)})$. As far as we are aware, this problem has not been previously studied. The related quantity $\alpha(KG_{\text{even}}^{(n,q)})$ is well-understood, as a special case of work of Deza, Erdős, and Frankl [DEF78] (discussed further in Section 4.2):

Theorem 4.12. ([DEF78].) *For q even and all n sufficiently large, $\alpha(KG_{\text{even}}^{(n,q)}) \leq \binom{n/2}{q/2}$, and this is tight for even n , as $\{I_1 \cup I_2 \cup \dots \cup I_{q/2} : \text{distinct } I_j \in \{\{1, 2\}, \{3, 4\}, \dots, \{n-1, n\}\}\}$ is an independent set.*

Similarly, for q odd and all n sufficiently large, $\alpha(KG_{\text{even}}^{(n,q)}) \leq \binom{(n-1)/2}{(q-1)/2}$, and this is tight for odd n , as $\{I_1 \cup I_2 \cup \dots \cup I_{(q-1)/2} \cup \{n\} : \text{distinct } I_j \in \{\{1, 2\}, \{3, 4\}, \dots, \{n-2, n-1\}\}\}$ is an independent set.

We have strong evidence for the following conjecture (which implies Theorem 4.12):

Conjecture 4.13. *Theorem 4.12 holds also with $\vartheta(KG_{\text{even}}^{(n,q)})$ in place of $\alpha(KG_{\text{even}}^{(n,q)})$.*

If true, this would imply $\text{Opt}_{\pm}(h) \leq O(n^{\lfloor q/2 \rfloor / 2})$ for all h as in Equation (14). Indeed, using the method of Section 4.2 we can prove Conjecture 4.13 for any particular small q (e.g., any $q \leq 10$), with computer assistance. As $q = 4$ is of most interest to us, we explicitly verify the conjecture in this case in Section 4.3, establishing:

Theorem 4.14. $\vartheta(KG_{\text{even}}^{(n,4)}) \leq \binom{n/2}{2}$ for all $n \geq 12$.

Combining this with Proposition 4.11, we conclude:

Corollary 4.15. *For any $h = \sum_{|S|=4} a_S \chi^S \in \mathfrak{C}(K_n)$ with $a \in \mathbb{R}^n$ satisfying $|\vec{a}| = 1$, we have*

$$\text{Opt}_{\pm}(h) \leq \sqrt{\binom{n/2}{2}} \sim \frac{1}{2\sqrt{2}}n,$$

and indeed $\text{SOS}_8(h^2) \leq \binom{n/2}{2}$, for any $n \geq 12$.

Remark: if there is an SOS proof that $h^2 \leq \theta$ for some scalar θ , there is also an SOS proof of the same degree that $\pm h \leq \sqrt{\theta}$.

Remark 4.16. The bound in Corollary 4.15 is exactly optimal for even n . To see this, consider

$$h_1(\chi) = (\chi_1\chi_2 + \chi_3\chi_4 + \dots + \chi_{n-1}\chi_n)^*(\chi_1\chi_2 + \chi_3\chi_4 + \dots + \chi_{n-1}\chi_n) \quad (15)$$

$$= n/2 - 2 \sum \{\chi^{I \cup J} : I, J \in \{\{1, 2\}, \dots, \{n-1, n\}\}, I \prec J\}. \quad (16)$$

Thus

$$h = \frac{h_1 - n/2}{2\sqrt{\binom{n/2}{2}}}$$

is of the required form; i.e., it is degree-4 homogeneous with squared coefficients summing to 1. It now suffices to show that $\text{Opt}(h_1) = (n/2)^2$, as this implies $\text{Opt}(h) = \sqrt{\binom{n/2}{2}}$. Recalling Example 2.27, it suffices to show that

$$\|M\| = n/2, \quad \text{where } M = -i(\gamma_1\gamma_2 + \gamma_3\gamma_4 + \dots + \gamma_{n-1}\gamma_n)$$

and the γ -matrices are as in Equation (6). But $M = Z \otimes \mathbb{1} \otimes \cdots \otimes \mathbb{1} + \mathbb{1} \otimes Z \otimes \cdots \otimes \mathbb{1} + \cdots + \mathbb{1} \otimes \mathbb{1} \otimes \cdots \otimes Z$, which is the $2^{n/2 \times n/2}$ diagonal matrix whose x th entry, for $x \in \{\pm 1\}^{n/2}$, is $\sum_j x_j$. Hence indeed $\|M\| = n/2$.

Remark 4.17. Equation (15) gives a h_1 which is proportional to a square of a quadratic, and it is optimized by a Gaussian state, as discussed later. In Appendix B we show that any h which is a weighted sum of squares of a small number of quadratics is approximately optimized by Gaussian states.

Remark 4.18. Restricting attention to even n , the lower bound $n \geq 12$ in Corollary 4.15 is best possible, since for $n \in \{8, 10\}$ it even holds that $\alpha(KG_{\text{even}}^{(n,4)}) = 14$. To see that $\alpha(KG_{\text{even}}^{(8,4)}) \geq 14$, consider the 7 lines of the Fano plane, though of as elements of $\binom{[7]}{3}$; then add the element 8 to each line, giving a 7 subsets of $\binom{[8]}{4}$ where any two have intersection size exactly 2; finally, include also the 7 complements (within $[8]$) of these subsets. In physics language, we can understand this example as follows: take terms $\chi_1\chi_2\chi_3\chi_4$ and $\chi_5\chi_6\chi_7\chi_8$ with positive coefficients. Restricting the $+1$ eigenspace of these two terms, the system may be described by 2 qubits, with X, Y, Z operators on the first qubit being bilinears in $\chi_1, \chi_2, \chi_3, \chi_4$ and X, Y, Z operators on the second qubit being bilinears in $\chi_5, \chi_6, \chi_7, \chi_8$. The remaining terms in h then are quartic operators whose sum is proportional to $X_1X_2 + Y_1Y_2 + Z_1Z_2$.

4.2 On bounding $\vartheta(KG_{\text{even}}^{(n,q)})$

Notation 4.19. Recall the *Johnson association scheme* $J(n, q)$ has $\binom{n}{q}$ as its points, with S, T belonging to the d th relation, $d = 0 \dots q$, iff $\text{dist}(S, T) := q - |S \cap T| = d$. We write $A_d^{(n,q)}$ for the adjacency matrix of the d th relation, with the superscript dropped when it is understood from context. Note that $A_0 = \mathbb{1}$ and $\sum_{d=0}^q A_d = \mathbb{J}$. For simplicity we assume $n > 2q$, so that A_1, \dots, A_d represent connected graphs.

Definition 4.20. Given n, q , and a set of *nonedge distances* $\mathcal{D} \subseteq \{1, \dots, q\}$, we write $KG_{\mathcal{D}} = KG_{\mathcal{D}}^{(n,q)}$ for the graph with vertex set $\binom{n}{q}$ and adjacency matrix $\sum_{d \notin \mathcal{D}} A_d$.

Example 4.21. The Erdős–Ko–Rado theorem [EKR61] concerns the case $\mathcal{D} = \{1, 2, \dots, q - k\}$, for $k \in \mathbb{N}^+$; we write the associated graph as KG_{EKR_k} . The theorem says that for sufficiently large n it holds that $\alpha(KG_{\text{EKR}_k}) = \binom{n-k}{q-k}$, with the maximum independent sets being of the form $\{S \subseteq [n] : S \supseteq T\}$ for some $|T| = k$.

Example 4.22. We are particularly interested in the case when $\mathcal{D} = \{2, 4, 6, \dots\}$ is the set of even integers; this yields the case of $KG_{\text{even}} = KG_{\text{even}}^{(n,q)}$ discussed in Theorem 4.12.

Deza, Erdős, and Frankl [DEF78] more generally showed the following:

Theorem 4.23. ([DEF78].) *If the elements of \mathcal{D} are $d_1 > d_2 > \dots > d_r$, then $\alpha(KG_{\mathcal{D}}) \leq \prod_{j=1}^r \frac{n-q+d_j}{d_j}$ for sufficiently large n .*

Indeed, Deza–Erdős–Frankl furthermore showed that unless the divisibility criterion $(d_1 - d_2) \mid (d_2 - d_3) \mid \dots \mid (d_{r-1} - d_r) \mid d_r$ is satisfied, one has the asymptotically stronger bound $\alpha(KG_{\mathcal{D}}) \leq O_q(n^{r-1})$.

Recall that for any graph at all we have $\alpha(G) \leq \vartheta(G)$. In the case of graphs coming from association schemes, the Lovász Theta Function reduces to the “linear programming bound” of Delsarte [Del73]. The next three facts review this bound in the context of the Johnson scheme.

Proposition 4.24. ([Del73, p. 48].) The matrices A_0, \dots, A_q are simultaneously diagonalizable, with eigenvalues given by the dual Hahn polynomials:

$$\text{spec}(A_d) = \{\tilde{H}_d(z) : z = 0 \dots q\}, \quad \tilde{H}_d(z) = \sum_{j=0}^d (-1)^{d-j} \binom{q-j}{d-j} \binom{q-z}{j} \binom{n-q+j-z}{j}.$$

In particular, the all-1's vector is an eigenvector of A_d with eigenvalue $\tilde{H}_d(0) = \binom{n-q}{d} \binom{q}{d}$. This is an eigenvalue of multiplicity 1 for all $d > 0$.

Lemma 4.25. Fix any $KG = KG_{\mathcal{D}}^{(n,q)}$. Then in the characterization of $\vartheta(G)$ given in Equation (13), it is equivalent to minimize only over Σ such that Σ_{ST} depends only on $\text{dist}(S, T)$. That is,

$$\vartheta(KG) = \min\{\kappa : \exists (c_e)_{e \notin \mathcal{D}}, \kappa(\mathbb{1} + \sum_{e \notin \mathcal{D}} c_e A_e) \geq \mathbb{J}\}.$$

Proof. Given any feasible Σ for Equation (13) achieving κ , we may replace it by $\hat{\rho} = \text{avg}_{\pi \in S_n} \{P(\pi)^\top \Sigma P(\pi)\}$, where $P(\pi)$ is the permutation matrix on $\binom{n}{q}$ induced by π . It is easy to see that $\hat{\rho}$ continues to be feasible and achieve κ , while also having $\hat{\rho}_{ST}$ depend only on $\text{dist}(S, T)$. \square

Corollary 4.26. Fix any $KG = KG_{\mathcal{D}}^{(n,q)}$. Then we have the following LP characterization:

$$\vartheta(KG) = \min_{(c_e)_{e \notin \mathcal{D}}} \left\{ \binom{n}{q} \middle/ p(0) : p(1), \dots, p(q) \geq 0 \text{ where } p(z) = \sum_{e \notin \mathcal{D}} c_e \tilde{H}_e(z) \right\}.$$

Proof. This follows directly from Proposition 4.24. \square

In the Erdős–Ko–Rado case, Schrijver [Sch81] was the first to show that in fact $\vartheta(KG_{\text{EKR}_k}) = \alpha(KG_{\text{EKR}_k}) = \binom{n-k}{q-k}$ for all $n \geq n_0(q, k)$; Wilson [Wil84] later gave the optimal $n_0(q, k) = (k+1)(q-k+1)$. Schrijver further asked whether Theorem 4.23 could in general be strengthened by replacing $\alpha(KG_{\mathcal{D}})$ with $\vartheta(KG_{\mathcal{D}})$. Our Conjecture 4.13 is the special case of this question when $\mathcal{D} = \{2, 4, 6, \dots\}$. In the next section, we show how to verify the conjecture for $q = 4$; similar techniques can be used to verify the conjecture for any particular small q .

4.3 Proving Theorem 4.14

We now explicitly prove Conjecture 4.13 in the case of $q = 4$.

Proof of Theorem 4.14. By Corollary 4.26 we have

$$\vartheta(KG_{\text{even}}^{(n,4)}) = \min_{c_1, c_3} \left\{ \binom{n}{4} \middle/ p(0) : p(1), p(2), p(3), p(4) \geq 0 \text{ where } p(z) = c_1 \tilde{H}_1(z) + c_3 \tilde{H}_3(z) \right\}.$$

The minimizer will occur when two of $p(1), p(2), p(3), p(4)$ are 0. We first consider fixing $p(4) = 0$. It is not hard to compute that $p(4) = 1 - 4c_1 - 4c_3$, and hence $p(4) = 0$ provided

$$c_1 = \frac{1}{4} - c_3.$$

Fixing this choice, and also writing $n' = n - 4$, we may compute:

$$\begin{aligned} p(0) &= 1 + 4n'(\frac{1}{4} - c_3) + 4\binom{n'}{3}c_3 \\ p(1) &= 1 + (3n' - 4)(\frac{1}{4} - c_3) + \left(\binom{n'-1}{3} - 3\binom{n'-1}{2}\right)c_3 \\ p(2) &= 1 + (2n' - 6)(\frac{1}{4} - c_3) - \left(2\binom{n'-2}{2} - 2(n' - 2)\right)c_3 \\ p(3) &= 1 + (n' - 6)(\frac{1}{4} - c_3) - (3(n' - 3) - 1)c_3. \end{aligned}$$

We now consider fixing $p(2) = 0$ by choosing

$$c_3 = \frac{1}{2(n' - 4)}.$$

This yields

$$p(0) = \frac{(n-1)(n-3)}{3}, \quad p(1) = \frac{(n-2)(n-4)(n-12)}{12(n-8)}, \quad p(3) = \frac{(n-4)(n-6)}{4(n-8)}.$$

Provided $n \geq 12$, we get $p(1), p(3) \geq 0$ and thereby establish

$$\vartheta(KG_{\text{even}}^{(n,4)}) \leq \binom{n}{4} / \left(\frac{(n-1)(n-3)}{3} \right) = \binom{n/2}{2}. \quad \square$$

5 Efficient upper bound certification for degree-4 SYK in degree-8 Sum-of-Squares

Let us consider the algebra over indeterminates χ_1, \dots, χ_n and τ_1, \dots, τ_n in which we initially assume only the following relations:

$$\chi_j^* = \chi_j, \quad \tau_j^* = \tau_j, \quad \chi_j^2 = 1 \quad \forall j \in [n]. \quad (17)$$

In particular, we do not necessarily assume that $\tau_j \tau_k = -\tau_k \tau_j$ or $\tau_j^2 = 1$. (The anti/commutation of the χ_j 's will be irrelevant.) Consider further the self-adjoint polynomial

$$h = \frac{i}{\sqrt{n}} \sum_{m=1}^n \tau_m \chi_m.$$

By the matrix form of Cauchy–Schwarz we can immediately conclude that

$$\text{Opt}_{\pm}(h) \leq \sqrt{\text{Opt}\left(\sum_{m=1}^n \tau_m^2\right)} \sqrt{\text{Opt}\left(\frac{1}{n} \sum_{m=1}^n \chi_m^2\right)} = \sqrt{\text{Opt}\left(\sum_{m=1}^n \tau_m^2\right)},$$

but let us deduce this slightly differently for the sake of SOS:

Proposition 5.1. *Let $\beta > 0$, and suppose there is a degree- k ($k \geq 2$) SOS proof of $\sum_{m=1}^n \tau_m^2 \leq \beta^2$. Then there is also one of $\pm h \leq \beta$.*

Proof. The inequalities are obtained by choosing $\alpha = \sqrt{\beta/\sqrt{n}}$ in

$$\frac{1}{2}(i\alpha^{-1}\tau_m \pm \alpha\chi_m)^*(i\alpha^{-1}\tau_m \pm \alpha\chi_m) \geq 0 \quad \implies \quad \mp i\tau_m \chi_m \leq \frac{1}{2\alpha^2}\tau_m^2 + \frac{\alpha^2}{2}\chi_m^2,$$

and summing over m . \square

Lemma 5.2. For $\vec{a} \in \mathbb{R}^{\binom{m}{3}}$, let $g = i \sum_{\substack{S \subseteq [m] \\ |S|=3}} a_S \chi^S \in \mathfrak{C}(K_m)$. Then

$$g^2 = |\vec{a}|^2 + \sum_{\substack{U \subseteq [m] \\ |U|=4}} \left(\sum_{\ell \in [m] \setminus U} \sum_{\substack{\{S,T\}: U=S \cup T \\ |S|=|T|=2}} \pm 2a_{S \cup \{\ell\}} a_{T \cup \{\ell\}} \right) \chi^U$$

for certain signs \pm .

Proof. It is clear that g^2 will have terms only of even degree. Further, since it is self-adjoint, these terms can only be of degree-0 or degree 4 (cf. [Fact 2.12](#)). The lemma now follows by explicit computation. \square

We now introduce some matrix and tensor notation. Let $h = \sum_{\substack{S \subseteq [n] \\ |S|=4}} J_S \chi^S$ for some given choice of J_S . Define a 4-index tensor J_{ijkl} with entries $J_{ijkl} = J_{\{i,j,k,l\}}$ for $i < j < k < l$ and for other orderings of i, j, k, l let π be a permutation on 4 elements so that $\pi(i) < \pi(j) < \pi(k) < \pi(l)$ and then $J_{ijkl} = J_{\pi(i), \pi(j), \pi(k), \pi(l)} \text{sign}(\pi)$. Conversely, given an arbitrary tensor J_{ijkl} define $h(J) \equiv \sum_{i,j,k,l \text{ distinct}} J_{ijkl} \chi_i \chi_j \chi_k \chi_l$, where the sum is over *distinct* i, j, k, l . (If J is completely anti-symmetric then there is no need to restrict to distinct i, j, k, l , but for general J this restriction matters.) From this tensor J_{ijkl} define an n^2 -by- n^2 matrix J^{mat} by regarding i, j as row indices and k, l as column indices.

Having introduced this notation we have:

Theorem 5.3. There is a constant $C > 0$ such that for any $h = \sum_{\substack{S \subseteq [n] \\ |S|=4}} J_S \chi^S$ we have $\text{SOS}_8(h) \leq Cn \|J^{\text{mat}}\|_4$, where $\|\dots\|_4$ denotes the Schatten 4-norm.

Proof. Throughout this proof we use C', C'', \dots to denote arbitrary constants. Write $h = \frac{1}{\sqrt{n}} \sum_i \chi_i \tau_i$, where we define $\tau_i = \frac{\sqrt{n}}{4} [h, \chi_i]$, i.e., up to constant factors $\tau_i = \sqrt{n} \sum_{j,k,l} J_{ijkl} \chi_j \chi_k \chi_l$. Remark: the reason for multiplying τ_i by \sqrt{n} and then dividing by \sqrt{n} is that it is a more convenient normalization for random instances.

We will show that there is a degree-8 proof that $\sum_{m=1}^n \tau_m^2 \leq C'n^2 \|(J^{\text{mat}})^2\|_2$. Then by [Proposition 5.1](#), there is a degree-8 proof that $h \leq Cn \|J^{\text{mat}}\|_4$.

By [Lemma 5.2](#), τ_m^2 is a sum of degree-0 and degree-4 terms. In fact, we may express these terms conveniently using matrix notation. We claim that

$$\sum_m \tau_m^2 = \frac{3!}{(4!)^2} n |J|^2 + \frac{3!}{(4!)^2} n h((J^{\text{mat}})^2), \quad (18)$$

where $h((J^{\text{mat}})^2)$ is defined by squaring J^{mat} , regarding it as a matrix, then regarding the resulting matrix as a 4-index tensor as an input to the function $h(\cdot)$.

By, [Corollary 4.15](#), $h((J^{\text{mat}})^2)$ is bounded by $C'n|(J^{\text{mat}})^2|$. Remark: to prove this in some more detail, take $(J^{\text{mat}})^2$, regard it as a 4-index tensor, then project onto the subspace which is fully anti-symmetric of permutations of indices, and then apply [Corollary 4.15](#). However, this projection cannot increase the ℓ_2 norm.

Thus, $\text{SOS}_8(h) \leq \sqrt{C'n^2|(J^{\text{mat}})^2| + C''n|J|^2}$. Since $n|J|^2 \leq n^2|(J^{\text{mat}})^2|$, the theorem follows.

To verify Equation (18), we have

$$\sum_m \tau_m^2 = \frac{n}{(4!)^2} \sum_m \sum_{jkl} \sum_{opq} J_{mjkl} J_{mopq} \chi_j \chi_k \chi_l \chi_o \chi_p \chi_q.$$

We break this sum up into several cases, depending on the number of indices j, k, l which match indices o, p, q , i.e., on $|\{j, k, l\} \cap \{o, p, q\}|$. If 0 or 2 indices match, then the resulting term is degree 6 or 2 and these terms vanish. So, we may consider the case that 1 or 3 indices match. The sum of terms with 3 indices matching is equal to

$$\frac{3!}{(4!)^2} n |J|^2,$$

where to derive we may pick an arbitrary permutation for indices matching (i.e., assume $j = o, k = p, l = q$) and then sum over the $3!$ permutations.

The sum of terms with 1 index matching is equal to

$$\frac{9}{(4!)^2} n \sum_{m,j} \sum_{k,l,p,q \text{ distinct}} J_{mjkl} J_{mj pq} \chi_k \chi_l \chi_p \chi_q,$$

where again we picked an arbitrary choice of indices matching (here picking $j = o$) and summed over the 9 such choices. \square

Theorem 5.3 has the immediate corollary that $\mathbf{E}[\text{SOS}_8(\mathbf{h})] = O(\sqrt{n})$ as one may estimate $\|J^{\text{mat}}\|_4$. However, we can tighten the constants using a slightly different choice of variables τ as in the following theorem:

Theorem 5.4. *Let $\mathbf{h} \sim \text{SYK}_4(n)$, where $n \geq 12$; i.e., for independent standard Gaussians \mathbf{J}_S ,*

$$\mathbf{h} = \frac{1}{\sqrt{\binom{n}{4}}} \sum_{\substack{S \subseteq [n] \\ |S|=4}} \mathbf{J}_S \chi^S.$$

Then $\mathbf{E}[\text{SOS}_8(\mathbf{h})] \leq \sqrt{1 + \sqrt{6}} \sqrt{n}$.

Proof. Write $\mathbf{h} = -\frac{i}{\sqrt{n}} \sum_{m=1}^n \boldsymbol{\tau}_m \chi_m$, where

$$\boldsymbol{\tau}_m = \frac{2i}{\sqrt{\binom{n-1}{3}}} \sum_{\substack{T \subseteq [m-1] \\ |T|=3}} \mathbf{J}_{T \cup \{m\}} \chi^T.$$

We remark that, up to normalization, the $\boldsymbol{\tau}_m$'s are independently distributed as $\text{SYK}_3(m-1)$. Applying Lemma 5.2 to $\boldsymbol{\tau}_m$, we get

$$\sum_{m=1}^n \tau_m^2 = \frac{4}{\binom{n-1}{3}} \sum_{\substack{S \subseteq [n] \\ |S|=4}} \mathbf{J}_S^2 + \sum_{m=1}^n \sum_{\substack{U \subseteq [m-1] \\ |U|=4}} \mathbf{c}_{m,U} \chi^U = \frac{4}{\binom{n-1}{3}} \sum_{\substack{S \subseteq [n] \\ |S|=4}} \mathbf{J}_S^2 + \sum_{\substack{U \subseteq [n] \\ |U|=4}} \sum_{m > \max(U)} \mathbf{c}_{m,U} \chi^U, \quad (19)$$

where

$$\mathbf{c}_{m,U} := \frac{8}{\binom{n-1}{3}} \sum_{\ell \in [m-1] \setminus U} \sum_{\substack{\{S,T\}: U=S \cup T \\ |S|=|T|=2}} \pm \mathbf{J}_{S \cup \{\ell, m\}} \mathbf{J}_{T \cup \{\ell, m\}}. \quad (20)$$

Thus using [Corollary 4.15](#) we get

$$\begin{aligned} \mathbf{E} \left[\text{SOS}_8 \left(\sum_{m=1}^n \tau_m^2 \right) \right] &\leq \frac{4}{\binom{n-1}{3}} \mathbf{E} \sum_{\substack{S \subseteq [n] \\ |S|=4}} \mathbf{J}_S^2 + \mathbf{E} \sqrt{\sum_{\substack{U \subseteq [n] \\ |U|=4}} \left(\sum_{m > \max(U)} \mathbf{c}_{m,U} \right)^2} \sqrt{\binom{n/2}{2}} \\ &\leq n + \sqrt{\mathbf{E} \sum_{\substack{U \subseteq [n] \\ |U|=4}} \left(\sum_{m > \max(U)} \mathbf{c}_{m,U} \right)^2} \sqrt{\binom{n/2}{2}}. \end{aligned} \quad (21)$$

To bound the expectation in [Equation \(21\)](#), note that for a fixed U , the random variables $\mathbf{c}_{m,U}$ have mean zero and are independent across m (since $\mathbf{c}_{m,U}$ depends only on \mathbf{J}_R for $R \ni m$). Thus

$$\mathbf{E} \sum_{\substack{U \subseteq [n] \\ |U|=4}} \left(\sum_{m > \max(U)} \mathbf{c}_{m,U} \right)^2 = \sum_{\substack{U \subseteq [n] \\ |U|=4}} \sum_{m > \max(U)} \mathbf{E}[\mathbf{c}_{m,U}^2] = \sum_{m=1}^n \sum_{\substack{U \subseteq [m-1] \\ |U|=4}} \mathbf{E}[\mathbf{c}_{m,U}^2].$$

Further, for fixed m and U , the summands $\mathbf{J}_{S \cup \{\ell, m\}} \mathbf{J}_{T \cup \{\ell, m\}}$ in $\mathbf{c}_{m,U}$ are products of *distinct* pairs of standard Gaussians, and hence

$$\mathbf{E}[\mathbf{c}_{m,U}^2] = \frac{64}{\binom{n-1}{3}^2} \cdot (m-5) \cdot 3$$

(and we see it did not matter what the \pm signs were). Thus the expectation in [Equation \(21\)](#) is

$$= \sum_{m=1}^n \binom{m-1}{4} \cdot \frac{64}{\binom{n-1}{3}^2} \cdot (m-5) \cdot 3 = \frac{960 \binom{n}{6}}{\binom{n-1}{3}^2},$$

and we conclude that

$$\mathbf{E} \left[\text{SOS}_8 \left(\sum_{m=1}^n \tau_m^2 \right) \right] \leq n + \sqrt{\frac{960 \binom{n}{6}}{\binom{n-1}{3}^2}} \sqrt{\binom{n/2}{2}} = n + \sqrt{6n^2 \cdot \frac{(n-4)(n-5)}{(n-1)(n-3)}} \leq (1 + \sqrt{6})n.$$

The result now follows from [Proposition 5.1](#). □

Remark 5.5. The factor $\sqrt{1 + \sqrt{6}} \approx 1.86$ in this theorem may be compared to the provable factor of $\sqrt{\ln 2} \approx .83$ of [\[FTW19\]](#) ([Remark 3.7](#)) and the conjectural factor of $\frac{1}{2\sqrt{2}} \approx .35$ (see [Equation \(11\)](#)).

6 Lower degree Sum-of-Squares: certification at degree 6 and fooling at degree 4

We have shown that, with high probability, degree-8 sum-of-squares certifies an upper bound to the largest eigenvalue of h for SYK_4 which is within a constant factor of \sqrt{n} . In this section we consider lower degree sum-of-squares.

We show that also a similar statement holds for degree-6 sum-of-squares: with high probability, it also certifies an upper bound to the largest eigenvalue of h for SYK_4 which is within a constant

factor of \sqrt{n} , albeit with a weaker constant. Indeed, a certain “fragment” of degree-6 sum-of-squares is able to prove this, and this “fragment” requires only a constant factor increase in matrix size over degree-4.

Let us explain the reason for the interest in this “fragment”. The interest is *not* to give a faster certifier; indeed, [Theorem 5.3](#) requires only evaluating a certain Schatten 4-norm which can be done faster than one can solve this “fragment”. Rather, the interest is that we hope that this “fragment” will be useful in general, not just for SYK₄. In addition to certifying SYK₄, it can also prove everything that SOS₄ can. In contrast, the Schatten 4-norm bound on its own likely does not give any particularly useful bound for typical problems of interest in quantum chemistry.

On the other hand, we will show that degree-4 sum-of-squares is not able to certify an $O(\sqrt{n})$ upper bound.

6.1 Degree-6 Sum-of-Squares for SYK₄

Let us define a certain “fragment” of SOS₆. We call this a “fragment” because it is defined by a positive semidefinite matrix that includes sum, but not all, of the information in SOS₆.

We call this fragment SOS_{4,2}; the reason for this notation will become clear. Briefly, SOS_{4,2} is defined by considering expectation values of monomials up to degree 4 in variables χ , as well as expectation variables of monomials of degree 1 in χ and degree 1 in τ , and finally expectation values of monomials of degree 2 in τ , subject to appropriate anticommutation constraints. The notation 4,2 in SOS_{4,2} is intended to denote this total degree at most 4 in χ and at most 2 in τ . Note that all expectation values of odd total degree in τ, χ may be assumed to vanish.

We can also formally repeat our previous definitions with slight modification:

Definition 6.1. We say that h is a *degree-4,2 sum of (Hermitian-)squares* (abbreviated *SOS*) if it is expressible as $g_1^*g_1 + \dots + g_m^*g_m$ for some polynomials g_1, \dots, g_m of degree at most 2 in χ and at most 1 in τ .

Definition 6.2. We define a *degree-4,2 pseudoexpectation* to be a linear functional $\tilde{\mathbf{E}}[\cdot] : \mathfrak{C}(\Gamma) \rightarrow \mathbb{C}$ with $\tilde{\mathbf{E}}[1] = 1$ and $\tilde{\mathbf{E}}[h] \geq 0$ for all degree-4,2 SOS h . We therefore call a self-adjoint $\rho \in \mathfrak{C}(\Gamma)$ a *degree-4,2 pseudostate* if has $\text{tr}(\rho) = 1$ and $\text{tr}(\rho h) \geq 0$ for all degree-4,2 SOS h .

We characterize this by the following positive semidefinite matrix:

Fact 6.3. Let $\tilde{\mathbf{E}}[\cdot]$ be a linear functional on $\mathfrak{C}(\Gamma)$ with $\tilde{\mathbf{E}}[1] = 1$. Let M be a matrix having rows and columns each indexed by a pair of sets (S_χ, S_τ) , with $S_\chi, S_\tau \subset [n]$, and with $|S_\chi| \leq 2$ and $|S_\tau| \leq 1$. Let M have $((S_\chi, S_\tau), (T_\chi, T_\tau))$ -entry equal to $\tilde{\mathbf{E}}[(\chi^{S_\chi})^*(\tau^{S_\tau})^*\chi^T\tau^{T_\tau}]$, where τ^{T_τ} is defined in the obvious way as the product of τ_i for $i \in T_\tau$. Then $\tilde{\mathbf{E}}[\cdot]$ is a degree-4,2 pseudoexpectation if and only if M is PSD.

Fact 6.4. The space of M for arbitrary $\tilde{\mathbf{E}}[\cdot]$ is subject to certain linear constraints.

We will write $M_{s,t}$ to denote certain submatrices of total degree at most s in χ and t in τ .

First, the submatrix $M_{4,0}$ indexed by rows and columns where $S_\tau = T_\tau = \emptyset$ is subject to the same linear constraints as for degree-4 pseudoexpectations: antisymmetry under permutation of indices, and constraints imposed by $\chi^2 = 1$.

Second, the entries of $M_{1,1}$ are linearly determined by entries of $M_{4,0}$. Indeed, the pseudoexpectation of $\chi\tau_j$ for any i, j is a linear function of $M_{4,0}$.

Third, the entries of $M_{0,2}$ which are pseudoexpectations of $\tau_i\tau_j$ are subject to a constraint that $\tilde{\mathbf{E}}[\tau_i\tau_j + \tau_j\tau_i]$ is linear determined by entries of $M_{4,0}$. Indeed, $\{\tau_i, \tau_j\}$ is degree at most 4 in χ .

Fact 6.5. Every degree-6 pseudoexpectation is a degree-4, 2 pseudoexpectation. This follows since every degree-4, 2 SOS is a degree-6 SOS since τ has degree 3.

Similarly, given any degree-6 pseudoexpectation, and corresponding PSD matrix of expectation values M , the matrix of expectation values for the corresponding degree-4, 2 pseudoexpectation is obtained by projecting M into an appropriate subspace.

Theorem 6.6. *Let $\mathbf{h} \sim \text{SYK}_4(n)$. Then, $\text{SOS}_6(\mathbf{h}) \leq \text{SOS}_{4,2}(\mathbf{h})$ and with high probability $\text{SOS}_{4,2}(\mathbf{h}) \leq O(\sqrt{n})$.*

Proof. The fact that $\text{SOS}_6(\mathbf{h}) \leq \text{SOS}_{4,2}(\mathbf{h})$ follows from [Fact 6.5](#).

The proof that $\mathbf{E}[\text{SOS}_{4,2}(\mathbf{h})] \leq O(\sqrt{n})$ is similar to that of [Theorem 5.3](#), and we use the same definition for τ_m . The only new thing required is to show that with high probability $\text{SOS}_{4,2}[\sum_{m=1}^n \tau_m^2] = O(n)$.

However, the linear constraints [Fact 6.4](#) show that $\text{SOS}_{4,2}$ still knows [Equation \(18\)](#). What we need to show is that $\text{SOS}_{4,2}$ can bound the right-hand side of this equation.

Indeed, we will show that SOS_4 can bound this. Of course, SOS_4 can bound the degree-0 terms.

Regard the space of N^2 -by- N^2 matrices as a vector space, and label rows of such matrices by a pair $i, j \in [n]$ and label columns by a pair $k, l \in [n]$. Then, define a projector $\pi_{\text{distinct}}(\cdot)$ which projects onto the subspace where i, j, k, l are all distinct. The pseudo-expectation of the sum of degree-4 terms, which is proportional to $nh((J^{\text{mat}})^2)$, equals (in a slight abuse of notation)

$$\text{Tr}(n\pi_{\text{distinct}}((J^{\text{mat}})^2)M_{4,0}).$$

The abuse of notation is that a matrix such as J^{mat} is size n^2 -by- n^2 (though it vanishes when $i = j$ or $k = l$), while $M_{4,0}$ is size $\binom{n}{2}$ -by- $\binom{n}{2}$, so one has essentially twice as many rows and columns as the other. However, we will freely take the trace of one kind of matrix with the other by extending $M_{4,0}$ to a matrix of size n^2 -by- n^2 in the obvious way using anti-symmetry, i.e., if $M_{4,0}$ has some given coefficient $(M_{4,0})_{S,T}$ for $|S| = |T| = 2$, then we extend it to a matrix with coefficients $(M_{4,0})_{i_1, i_2; j_1, j_2}$ which equals $\pm(M_{4,0})_{\{i_1, i_2\}, \{j_1, j_2\}}$ where the sign depends on the permutation needed to make $i_1 < i_2$ and $j_1 < j_2$.

However,

$$\text{Tr}(n\pi_{\text{distinct}}((J^{\text{mat}})^2)M_{4,0}) \leq \text{Tr}(n(J^{\text{mat}})^2M_{4,0}),$$

as the terms where $i, j = k, l$ or $i, j = l, k$ contribute positively.

Further, $M_{4,0}$ is a PSD matrix with trace $O(n^2)$. Hence, this pseudoexpectation is bounded by $O(n^2)\|n(J^{\text{mat}})^2\|_{\text{op}} = O(n^3)\|J^{\text{mat}}\|_{\text{op}}^2$. Standard random matrix theory results show that with high probability $\|J^{\text{mat}}\|_{\text{op}} = O(1/n)$. This completes the proof¹².

Remark: it is interesting to compare this degree-4 bound on z , namely by $O(n^3)\|J^{\text{mat}}\|_{\text{op}}^2$, to the bound we used in the proof of [Theorem 5.3](#), namely bounding z by $O(n^2)\|(J^{\text{mat}})^2\|_2$. The bound used in [Theorem 5.3](#) is tighter, but they have the same scaling for a matrix such as J^{mat} where “most” eigenvalues are of order $O(1/n)$. \square

¹²To give the bound on $\|J^{\text{mat}}\|_{\text{op}}$ in more detail, such a matrix is a matrix with random Gaussian entries, which are independent up to the anti-symmetry. We can draw matrices from that distribution by taking a random Gaussian matrix with *independent* entries (and the same variance up to constant factor) and summing over signed permutations of the indices to project onto the totally antisymmetric subspace. Then, using standard random matrix bounds on the operator norm of the matrix with independent entries and a triangle inequality gives us the desired bound on $\|J^{\text{mat}}\|_{\text{op}}$. We use similar bounds elsewhere and [\[HTS21\]](#) uses the same bound.

6.2 Fooling Degree q

Theorem 6.7. *Let $h = \sum_S$ be an instance of SYK_q for q even. Let $\rho = 1 + Ch$, where C is a scalar. Then with high probability $\text{tr}(\rho h) = C \cdot (1 + o(1))$. Further, if $|C|$ is sufficiently small compared to $n^{q/4}$, then with high probability, ρ is a degree- q pseudostate.*

Consequently, for $\mathbf{h} \sim \text{SYK}_4$, with high probability $\text{SOS}_4(\mathbf{h}) = \Omega(n)$.

Proof. Let \vec{a} be the vector of coefficients of terms in h . Then, $\text{tr}(\rho h) = C|\vec{a}|^2$ and with high probability $|\vec{a}|^2 = 1 + o(1)$.

To verify that it is a degree- q -pseudostate, we must verify that $\text{tr}(g\rho g^*) \geq 0$ for all g of degree at most $q/2$. Note that $\text{tr}(\rho m)$ vanishes for any m which is homogeneous of degree not equal to 0 or q . Hence, it suffices to verify that the inequality holds when g is homogeneous of degree 0 (which is immediate) and when g is homogeneous of degree $q/2$.

In case that g is homogeneous of degree- $q/2$,

$$g = \sum_{\substack{S \subseteq [n] \\ |S|=q/2}} b_S \chi^S,$$

we have

$$\text{tr}(g\rho g^*) = \sum_{\substack{S \subseteq [n] \\ |S|=q/2}} |b_S|^2 + C \text{tr}(ghg^*). \quad (22)$$

To evaluate $\text{tr}(ghg^*)$, we give a representation in terms of matrices (essentially the same representation is used in [HTS21]).

Define a matrix m whose rows and columns are indexed by subsets of $[n]$ with cardinality $q/2$. The coefficient $m_{S,T}$ (where $S, T \subset [n]$ and $|S| = |T| = n/2$) is defined to vanish unless $S \cup T = \emptyset$, in which case $m_{S,T} = \text{tr}(\chi^S h \chi^T) = \pm a_{S \cup T}$. The sign may be worked out using the anticommutation relations, of course. Then, $\text{tr}(ghg^*) = \langle b | m | b \rangle$.

However, standard random matrix theory bounds show that with high probability $\|m\|_{\text{op}} = O(n^{-q/4})$ and so indeed if $|C|$ is sufficiently small compared to $n^{q/4}$, then with high probability, ρ is a degree- q pseudostate. \square

7 Gaussian states

This section will focus exclusively on $\mathfrak{C}(K_n)$, where all indeterminates pairwise anticommute. We will discuss *Gaussian states* [TD02, Bra05], a reasonably broad class of states ρ (including all Slater determinant states) for which: (i) there is an efficient representation of ρ ; (ii) given this representation, and some polynomial $h \in \mathfrak{C}(K_n)$, there is an efficient procedure for computing $\mathbf{E}_\rho[h]$. In particular, these states may be used for classical (implicit) lower-bound witnessing algorithms, as described in Section 2.7.

A succinct characterization of Gaussian states is that they are precisely those states that are optimizers for homogeneous degree-2 Hamiltonians [Bra05]. To define them more explicitly, first suppose $\psi_1, \dots, \psi_m \in \mathfrak{C}(\Gamma)$ are self-adjoint, commuting involutions ($\psi_j^2 = 1$). Then for any $\lambda_1, \dots, \lambda_m \in \{\pm 1\}$, the elements $1 + \lambda_j \psi_j$ are positive and commuting, and hence

$$\rho = \prod_{j=1}^m (1 + \lambda_j \psi_j)$$

is a state (note that the order of the factors is irrelevant). Indeed, this was already pointed out in [Example 2.20](#), just with different letters. Furthermore, the same is true if we generalize by allowing $\lambda_1, \dots, \lambda_m \in [-1, 1]$. Additionally, if $\Gamma = K_n$ with n even, we may take $\psi_1 = i\chi_1\chi_2$, $\psi_2 = i\chi_3\chi_4$, etc. Even more generally, we may augment this construction with the orthogonal transformation described in [Fact 2.14](#). These generalizations lead us to the class of Gaussian states:

Definition 7.1. Consider the setting of $\mathfrak{C}(K_n)$ with n even. Let $\lambda_1, \dots, \lambda_{n/2} \in [-1, 1]$, let $O \in \mathbb{R}^{n \times n}$, and let $\ell = O\chi$. Then any element

$$\rho = \prod_{j=1}^{n/2} (1 + i\lambda_j \ell_{2j-1} \ell_{2j}) \in \mathfrak{C}(K_n) \quad (23)$$

is called a *Gaussian state*.

Example 7.2. For example, if $O = \mathbb{1}$ and $\lambda_j = -1$ for each j , we have the Gaussian state

$$\prod_{j=1}^{n/2} (1 - i\chi_{2j-1}\chi_{2j}).$$

Under the canonical representation using the γ -matrices from [Equation \(6\)](#), this is

$$\prod_{j=1}^{n/2} (1 + \underbrace{Z \otimes \dots \otimes Z}_j \otimes \underbrace{\mathbb{1} \otimes \dots \otimes \mathbb{1}}_{n/2-j}) = 2^{n/2} |00 \dots 0\rangle \langle 00 \dots 0|.$$

We will also be interested in the case when the condition $|\lambda_j| \leq 1$ is not enforced:

Definition 7.3. We call $\rho \in \mathfrak{C}(K_n)$ a *Gaussian pseudostate* if it has the form [Equation \(23\)](#), but with any $\lambda_1, \dots, \lambda_{n/2} \in \mathbb{R}$ allowed. In general we will write $\tilde{\mathbf{E}}_\rho[h] = \text{tr}(\rho h)$, with $\tilde{\mathbf{E}}_\rho[\cdot]$ being at worst a degree-0 pseudoexpectation. (If ρ is known to be a proper state we will switch to the notation $\mathbf{E}_\rho[\cdot]$.)

Definition 7.4. A Gaussian pseudostate as in [Equation \(23\)](#) is characterized by its *covariance matrix* $\Sigma \in \mathbb{R}^{n \times n}$, which is the real antisymmetric matrix defined by $\Sigma = O^\top \Lambda O$, where Λ is block-diagonal with 2×2 blocks of the form $\begin{pmatrix} 0 & \lambda_j \\ -\lambda_j & 0 \end{pmatrix}$; it satisfies $\Sigma_{jk} = \tilde{\mathbf{E}}_\rho[i\chi_j\chi_k]$ for all $1 \leq j, k \leq n$. Conversely, any real antisymmetric Σ defines a Gaussian pseudostate which we will denote by ρ_Σ ; this is a proper Gaussian state provided $\|\Sigma\|_{\text{op}} \leq 1$.

Definition 7.5. We introduce the terminology *Gaussian pseudostate* for the case when $|\lambda_j| \leq 1$ is not necessarily enforced. A Gaussian pseudostate is characterized by its *covariance matrix* $\Sigma \in \mathbb{R}^{n \times n}$, which is the real antisymmetric matrix defined by $\Sigma = O^\top \Lambda O$, where Λ is block-diagonal with 2×2 blocks of the form $\begin{pmatrix} 0 & \lambda_j \\ -\lambda_j & 0 \end{pmatrix}$; it satisfies $\Sigma_{jk} = \tilde{\mathbf{E}}_\rho[iX_jX_k]$ for all $1 \leq j, k \leq n$. Conversely, any real antisymmetric Σ defines a Gaussian pseudostate which we will denote by ρ_Σ ; this is a proper Gaussian state provided $\|\Sigma\|_{\text{op}} \leq 1$.

Fact 7.6. For the Gaussian pseudostate ρ_Σ , and $S \subseteq \binom{[n]}{4}$, we have

$$\tilde{\mathbf{E}}_{\rho_\Sigma}[\chi^S] = -(\Sigma_{S_1S_2}\Sigma_{S_3S_4} - \Sigma_{S_1S_3}\Sigma_{S_2S_4} + \Sigma_{S_1S_4}\Sigma_{S_2S_3}),$$

as is easily verified. More generally, the pseudomoments may be computed by the fermionic Wick formula: for $S \subseteq [n]$ we have $\mathbf{E}_{\rho_\Sigma}[\chi^S] = i^{\binom{|S|}{2}} \text{Pf}(\Sigma_{S,S})$, where Pf denotes the Pfaffian and $\Sigma_{S,S}$ is the submatrix of Σ with rows/columns indexed by S .

Example 7.7. Recalling the degree-4 polynomials h from [Remark 4.16](#), which exactly saturates the bound $\text{Opt}_{\pm}(h) \leq \sqrt{\binom{n/2}{2}}$ from [Corollary 4.15](#), it is easy to see that the optimum is achieved by a Gaussian state, $\rho = (1 + i\chi_1\chi_2)(1 + i\chi_3\chi_4) \cdots (1 + i\chi_{n-1}\chi_n)$.

7.1 Prior work on optimization and Gaussian states

Bravyi, Gosset, König, and Temme [\[BGKT19\]](#) investigated the task of finding the optimal Gaussian state for a given homogeneous degree-4 Hamiltonian with real coefficients.¹³ Let us define:

Definition 7.8. Given self-adjoint $h \in \mathfrak{C}(K_n)$, we define

$$\text{Opt}_{\text{Gauss}}(h) = \max\{\mathbf{E}_{\rho}[h] : \rho \text{ is a Gaussian state}\} = \max_{\substack{\Sigma \in \mathbb{R}^{n \times n} \\ \text{antisymmetric} \\ \|\Sigma\|_{\text{op}} \leq 1}} \{\mathbf{E}_{\rho_{\Sigma}}[h]\}.$$

For h degree-4 homogeneous with real coefficients, the work [\[BGKT19\]](#) introduced a polynomial-size SDP relaxation of $\text{Opt}_{\text{Gauss}}(h)$,

$$\text{SDP}_{4,\text{Gauss}}(h) \geq \text{Opt}_{\text{Gauss}}(h),$$

which we recall in [Definition 7.17](#) (in [Section 7.2](#)). Concerning it, they proved several results, including the following:

Theorem 7.9. ([\[BGKT19, So09\]](#).) Applying the SDP-rounding algorithm of So [\[So09\]](#) gives a randomized polynomial-time rounding algorithm for $\text{SDP}_{4,\text{Gauss}}(h)$ that with high probability finds a Gaussian state achieving at least $\frac{1}{O(\log n)} \text{SDP}_{4,\text{Gauss}}(h)$.

We present and generalize the above result in [Section 7.2](#).

Theorem 7.10. ([\[BGKT19\]](#).) $\text{SDP}_{4,\text{Gauss}}(h) \geq \frac{1}{n} \text{Opt}(h)$ for all degree-4 homogeneous $h \in \mathfrak{C}(K_n)$.

Remark 7.11. Inspecting the proof of [Theorem 7.10](#), one sees it implies the stronger result $\text{SDP}_{4,\text{Gauss}}(h) \geq \frac{1}{n} \text{SOS}_4(h)$.

We also note the following result shown by Haldar, Tavakol, and Scaffidi [\[HTS21\]](#):¹⁴

Theorem 7.12. ([\[HTS21\]](#).) For $\mathbf{h} \sim \text{SYK}_4(n)$, with high probability $\text{Opt}_{\text{Gauss}}(\mathbf{h}) \leq O(1)$.

Remark 7.13. Inspecting the proof of [Theorem 7.12](#), one sees it implies the stronger result $\text{SDP}_{4,\text{Gauss}}(\mathbf{h}) \leq O(1)$, with high probability.

Remark 7.14. One may also prove: for $\mathbf{h} \sim \text{SYK}_4(n)$, with high probability $\text{Opt}_{\text{Gauss}}(\mathbf{h}) = \Omega(1)$. We define two n -by- n matrices. Let g_0 be the block-diagonal matrix with the first $n/4$ blocks being $\begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}$, and the remaining $n/4$ blocks vanishing. Let g_1 be the matrix defined as follows: the k, l entry of j is only nonvanishing if $k > n/2$ and $l > n/2$, in which case the entry is equal to $\sum_{i < j \leq n/2} (g_0)_{i,j} J_{\{i,j,k,l\}}$ for $k < l$ and has the opposite sign for $k > l$. Finally, consider the covariance matrix $g_0 + Cg_1$, where C is a scalar. The entries of the matrix g_1 are independent¹⁵ Gaussian random variables, and the nonvanishing entries have variance $\Theta(1/n^3)$, so standard random matrix theory results show that $\|g_1\|_{\text{op}} = O(1/n)$. Hence we may take $C = \Theta(n)$ and have this covariance matrix define a Gaussian state. One may verify that the expectation of h for this covariance matrix is $C \cdot \Theta(1/n)$, so for $C = \Theta(n)$ this is $\Theta(1)$.

¹³In fact their work also applied to polynomials with a mix of degree-2 and degree-4 terms; but, they showed a reduction to the homogeneous degree-4 case, and we stick to this for simplicity.

¹⁴The eigenvalue bound for random Gaussian matrices that they rely on is standard.

¹⁵Up to the antisymmetry of the matrix, of course.

Remark 7.15. As we show in [Theorem 6.7](#), for $\mathbf{h} \sim \text{SYK}_4(n)$, with high probability $\text{SOS}_4(n) \geq \Omega(n)$. In this case $\text{SDP}_{4,\text{Gauss}}(\mathbf{h}) \geq \Omega(1)$ by [Remark 7.11](#), and so $\text{SDP}_{4,\text{Gauss}}(\mathbf{h}) = \Theta(1)$ by [Remark 7.13](#).

Bravyi–Gosset–König–Temme end their work [\[BGKT19\]](#) by asking whether for all h one might have $\text{SDP}_{4,\text{Gauss}}(h) \geq \frac{1}{C} \text{Opt}(h)$, for some universal constant C . Our [Theorem 7.18](#) below shows that this is true when $\text{Opt}(h)$ is within a constant factor of the upper bound from [Corollary 4.15](#) (namely n times the 2-norm of h ’s coefficients). Unfortunately, the inequality cannot be true in general: any such C must be at least $\Theta(\sqrt{n})$. This follows from [Theorem 7.12](#) together with the result (see our [Theorem 8.1](#)) that $\text{Opt}(\mathbf{h}) \geq \Omega(\sqrt{n})$ with high probability for $\mathbf{h} \sim \text{SYK}_4(n)$. Whether the worst-case factor between $\text{Opt}(h)$ and $\text{Opt}_{\text{Gauss}}(h)$ is more like \sqrt{n} or more like n (or something in between), we leave as an open question.

7.2 Rounding $\text{SDP}_{4,\text{Gauss}}(h)$

Let us now describe the SDP relaxation for degree-4 Hamiltonians introduced in [\[BGKT19\]](#).

Notation 7.16. Given $A \in \mathbb{C}^{m \times m}$, let $\text{Vec}(A) \in \mathbb{C}^m \otimes \mathbb{C}^m$ denote the vector formed by stacking the columns of A , in order. Conversely, let $\text{Mat} : \mathbb{C}^m \otimes \mathbb{C}^m \rightarrow \mathbb{C}^{m \times m}$ denote the inverse operation. We remark that

$$\text{Tr}_1(\text{Vec}(A)\text{Vec}(B)^\top) = A^\top B, \quad \text{Tr}_2(\text{Vec}(A)\text{Vec}(B)^\top) = AB^\top,$$

where Tr_j denotes tracing out the j th tensor component of $\mathbb{C}^m \otimes \mathbb{C}^m$.

Definition 7.17. Let $\Sigma \in \mathbb{R}^{n \times n}$ be antisymmetric with $\|\Sigma\|_{\text{op}} \leq 1$. Define $R = \text{Vec}(\Sigma)\text{Vec}(\Sigma)^\top$. Note that we have the following properties:

1. $R \geq 0$;
2. $\text{Tr}_1(R) \leq \mathbb{1}$ (since $\text{Tr}_1(R) = \Sigma^\top \Sigma$ and $\|\Sigma\|_{\text{op}} \leq 1$);
3. $\text{Tr}_2(R) \leq \mathbb{1}$ (similarly, since $\text{Tr}_2(R) = \Sigma \Sigma^\top$);
4. $\langle j_1 k_1 | R | j_2 k_2 \rangle$ changes sign if j_1, k_1 are interchanged, or if j_2, k_2 are interchanged (since Σ is antisymmetric).

Furthermore, if $h = \sum_{S \in \binom{[n]}{4}} a_S \chi^S \in \mathfrak{C}(K_n)$ (with n even), and we introduce the matrix

$$H = - \sum_{S \in \binom{[n]}{4}} a_S (|S_1 S_2\rangle \langle S_3 S_4| - |S_1 S_3\rangle \langle S_2 S_4| + |S_1 S_4\rangle \langle S_2 S_3|),$$

then $\mathbf{E}_{\rho_\Sigma}[h] = \text{tr}(\rho_\Sigma h) = \text{Tr}(RH)$. We thereby define

$$\text{SDP}_{4,\text{Gauss}}(h) = \max\{\text{Tr}(RH) : R \text{ satisfying properties 1–4 above}\},$$

which is a semidefinite programming relaxation of $\text{Opt}_{\text{Gauss}}(p)$.

Given this SDP, [\[BGKT19\]](#) observe that the efficient randomized rounding algorithm of So [\[So09\]](#) establishes $\text{Opt}_{\text{Gauss}}(p) \geq \frac{1}{O(\log n)} \text{SDP}_{4,\text{Gauss}}(p)$. So’s work is a kind of “noncommutative analogue” of the $\frac{1}{O(\log n)}$ -factor approximation algorithm for quadratic programming due to Nesterov [\[Nes98\]](#), Nemirovsky–Roos–Terlaky [\[NRT99\]](#), and Megretski [\[Meg01\]](#). In the “commutative case”, said

algorithm was generalized by Charikar and Wirth [CW04] to show that there is an $\frac{1}{O(\log(1/\epsilon))}$ -factor approximation for Max-Cut whenever the optimum is within a factor of ϵ of the “naive upper bound” (see Remark 2.29). Here we are able to generalize the Charikar–Wirth result to our noncommutative setting:

Theorem 7.18. *There is an efficient randomized algorithm that, given $h = \sum_{S \in \binom{[n]}{4}} a_S \chi^S$ with real $\vec{a} \in \mathbb{R}^{\binom{[n]}{4}}$ satisfying $|\vec{a}| = 1$ and $\text{SDP}_{4, \text{Gauss}}(p) = \epsilon n$, with high probability outputs the covariance matrix Σ of a Gaussian state satisfying*

$$\mathbf{E}_{\rho_\Sigma}[h] = \text{tr}(\rho_\Sigma h) \geq \frac{\epsilon}{O(\log(1/\epsilon))} n.$$

Remark 7.19. From Corollary 4.15 we know that $\epsilon \leq \frac{1}{2\sqrt{2}}$ always. It’s also trivial to show $\epsilon \geq 1/n^3$, say, meaning the $O(\log(1/\epsilon))$ factor is never worse than $O(\log n)$, recovering Theorem 7.9. In fact, taking care in our proof shows that this approximation factor need not be worse than $(1 + o(1))\sqrt{2 \ln n}$.

Remark 7.20. It is straightforward to observe that our algorithm can be derandomized in $n^{O(\log(1/\epsilon))}$ time, which is polynomial in the case $\epsilon = \Omega(1)$ of most interest. The algorithm employs n^2 Rademacher random variables, and the analysis only needs 4-wise uniformity for Inequality (25) (see e.g. [AH11, Thm. 4.3]), 2-wise uniformity for Inequality (28), and $(p+2)$ -wise uniformity for Inequality (29), with $p = O(\log(1/\epsilon))$. Thus the derandomization follows from [ABI86].

Proof. By standard semidefinite programming methods [GLS81], one can efficiently find a feasible solution R achieving $\text{SDP}_{4, \text{Gauss}}(p) = \epsilon n$.¹⁶ We now describe a randomized rounding algorithm that finds the required Σ with constant probability. We may boost the success probability arbitrarily close to 1 by repeating the algorithm constantly many times and choosing the best result achieved.

Since $R \geq 0$ we may write a Cholesky factorization $R = LL^\top$ which, by property 4 in Definition 7.17 will satisfy

$$\langle k, j | L = -\langle j, k | L \quad \forall j, k \in [n]. \quad (24)$$

Now define

$$\Sigma = \sigma \text{Mat}(L\vec{x}) = \sum_{j,k=1}^n x_{jk} \sigma A^{(jk)}, \quad A^{(jk)} := \text{Mat}(L | jk),$$

where σ is a positive constant to be chosen later, and \vec{x} is a vector of n^2 independent Rademacher random variables. Note that each $A^{(jk)}$ is antisymmetric, by Equation (24); hence Σ is always antisymmetric and thus defines a Gaussian pseudostate. Further,

$$\mathbf{E}[\text{Vec}(\Sigma)\text{Vec}(\Sigma)^\top] = \sigma^2 L \mathbf{E}[|\mathbf{x}\rangle\langle\mathbf{x}|] L^\top = \sigma^2 R,$$

from which we see that

$$\mathbf{E}[\text{tr}(\rho_\Sigma h)] = \sigma^2 \text{Tr}(RH) = \sigma^2 \text{SDP}_{4, \text{Gauss}}(h) = \sigma^2 \cdot \epsilon n.$$

Since $\text{tr}(\rho_\Sigma h)$ is a quadratic polynomial in Rademachers, it follows (e.g. [O’D14, Thm. 9.24]) that

$$\text{tr}(\rho_\Sigma h) \geq \sigma^2 \cdot \epsilon n \text{ with probability at least } e^{-4}/4 > .004. \quad (25)$$

¹⁶See the technicalities discussed in Section 2.7.

(Remark: So [So09] uses a more complicated construction for Σ that ensures $\text{tr}(\rho_\Sigma h) = \sigma^2 \text{SDP}_{4, \text{Gauss}}(h)$ holds with certainty. He then shows $\|\Sigma\|_{\text{op}} \leq 1$ with high probability with the choice $c \sim \frac{1}{\sqrt{2e \ln n}}$. Our methods below would also show this even for $c \sim \frac{1}{\sqrt{2 \ln n}}$.)

We now need to study the eigenvalues of Σ . It will be mildly more convenient to study the eigenvalues of

$$\mathbf{G} := i\Sigma = \sum_{j,k=1}^n \mathbf{x}_{jk} B^{(jk)}, \quad \text{where } B^{(jk)} := \sigma i A^{(jk)},$$

since the matrices $B^{(jk)}$ are self-adjoint. Note that

$$\sum_{j,k=1}^n A^{(jk)\top} A^{(jk)} = \sum_{j,k=1}^n \text{Tr}_1(L |jk\rangle \langle jk| L^\top) = \text{Tr}_1(R) \leq 1$$

(and similarly $\sum_{j,k=1}^n A^{(jk)} A^{(jk)\top} \leq \mathbb{1}$), and this is equivalent to

$$\sum_{j,k=1}^n (B^{(jk)})^2 \leq \sigma^2 \mathbb{1}. \quad (26)$$

At this point one can begin to analyze \mathbf{G} using standard theory for sums of random self-adjoint matrices; e.g., $\mathbf{Pr}[\lambda_{\max}(\mathbf{G}) \geq t] \leq ne^{-t^2/(2\sigma^2)}$ holds [Tro12].

Say \mathbf{G} has (real) eigenvalues $\lambda_1, \dots, \lambda_n$ and associated orthonormal eigenvectors $\vec{\phi}_1, \dots, \vec{\phi}_n$. Our algorithm may form

$$\hat{\mathbf{G}} = \sum \{ \lambda_j |\phi_j\rangle \langle \phi_j| : |\lambda_j| \leq 1 \},$$

which is also self-adjoint, and then $\hat{\Sigma} = i\hat{\mathbf{G}}$. This $\hat{\Sigma}$ is the covariance matrix of a Gaussian state $\rho_{\hat{\Sigma}}$, and by [Fact 7.6](#) we have

$$\mathbf{err} := |\text{tr}(\rho_\Sigma h) - \text{tr}(\rho_{\hat{\Sigma}} h)| = \left| \sum_{S \in \binom{[n]}{4}} a_S (e_1(S) - e_2(S) + e_3(S)) \right|, \quad (27)$$

where

$$e_1(S) = \Sigma_{S_1 S_2} \Sigma_{S_3 S_4} - \hat{\Sigma}_{S_1 S_2} \hat{\Sigma}_{S_3 S_4}$$

and $e_2(S), e_3(S)$ are defined similarly (with subscripts $S_1 S_3, S_2 S_4$ and $S_1 S_4, S_2 S_3$, respectively). By Cauchy–Schwarz,

$$\mathbf{err}^2 \leq \left(\sum_S a_S^2 \right) \left(\sum_S (e_1(S) - e_2(S) + e_3(S))^2 \right) \leq 3 \sum_S e_1(S)^2 + 3 \sum_S e_2(S)^2 + 3 \sum_S e_3(S)^2.$$

We will bound the first sum on the right, with the other two being bounded similarly. Since

$$(rs - \widehat{r}\widehat{s})^2 = (r(s - \widehat{s}) + \widehat{s}(r - \widehat{r}))^2 \leq 2r^2(s - \widehat{s})^2 + 2\widehat{s}^2(r - \widehat{r})^2,$$

we get

$$\sum_S e_1(S)^2 \leq 2 \sum_S \Sigma_{S_1 S_2}^2 (\Sigma_{S_3 S_4} - \hat{\Sigma}_{S_3 S_4})^2 + 2 \sum_S \hat{\Sigma}_{S_3 S_4}^2 (\Sigma_{S_1 S_2} - \hat{\Sigma}_{S_1 S_2})^2.$$

Here, the first expression on the right is at most

$$2 \left(\sum_{1 \leq j < k \leq n} \Sigma_{jk}^2 \right) \left(\sum_{1 \leq j < k \leq n} (\Sigma_{jk} - \widehat{\Sigma}_{jk})^2 \right) = \frac{1}{2} \|\Sigma\|_2^2 \|\Sigma - \widehat{\Sigma}\|_2^2,$$

where we used the antisymmetry of $\Sigma, \widehat{\Sigma}$. We can use similar considerations to bound the second expression, and will also use the evident fact that $\|\widehat{\Sigma}\|_2 \leq \|\Sigma\|_2$. Putting everything together, we conclude

$$\mathbf{err}^2 \leq \frac{9}{2} \|\Sigma\|_2^2 \|\Sigma - \widehat{\Sigma}\|_2^2 = \frac{9}{2} \|\mathbf{G}\|_2^2 \|\mathbf{G} - \widehat{\mathbf{G}}\|_2^2.$$

One easily calculates that $\mathbf{E}[\|\mathbf{G}\|_2^2] = \text{Tr}(\sum_{jk} (B^{(jk)})^2) \leq \sigma^2 n$ (by [Inequality \(26\)](#)), and therefore by Markov we have

$$\mathbf{err}^2 \leq C \sigma^2 n \cdot \|\mathbf{G} - \widehat{\mathbf{G}}\|_2^2 \text{ except with probability at most } .001, \quad (28)$$

where C is a universal constant.

To analyze this final expression we appeal to Khintchine's inequality with operator coefficients (originally due to Lust-Piquard [\[LP86\]](#), improved in [\[LPP91\]](#)), which tells us that for all $p \geq 1$,

$$\mathbf{E}[\|\mathbf{G}\|_p^p] \leq (C' \sqrt{p})^p \left\| \sqrt{\sum_{jk} (B^{(jk)})^2} \right\|_p^p \leq (C' \sqrt{p} \sigma)^p n,$$

where C' is another universal constant, and we used [Inequality \(26\)](#) again. Note that

$$\|\mathbf{G} - \widehat{\mathbf{G}}\|_2^2 = \sum_{j=1}^n \lambda_j^2 \cdot \mathbf{1}[|\lambda_j| \geq 1] \leq \sum_{j=1}^n \lambda_j^2 \cdot |\lambda_j|^p = \|\mathbf{G}\|_{p+2}^{p+2},$$

regardless of p . We choose p so that $\sqrt{p+2} = (eC'\sigma)^{-1}$ (so we need to also choose $\sigma \leq (\sqrt{3}eC')^{-1}$ so that $p \geq 1$). With this choice, we get

$$\mathbf{E}[\|\mathbf{G} - \widehat{\mathbf{G}}\|_2^2] \leq \mathbf{E}[\|\mathbf{G}\|_{p+2}^{p+2}] \leq \exp\left(-\frac{1}{C''\sigma^2}\right) n.$$

Employing Markov again, and combining with [Inequality \(28\)](#), we get

$$\mathbf{err} \leq C_1 \sigma \exp\left(-\frac{1}{C_2 \sigma^2}\right) n \text{ except with probability at most } .002. \quad (29)$$

Finally, from [Inequality \(25\)](#) and [eq. \(27\)](#) we have shown that the algorithm constructs Gaussian state $\rho_{\widehat{\Sigma}}$ satisfying

$$\text{tr}(\rho_{\widehat{\Sigma}} h) \geq \left(\sigma^2 \cdot \epsilon - C_1 \sigma \exp\left(-\frac{1}{C_2 \sigma^2}\right) \right) n$$

with probability at least .002. The proof is completed by taking $\sigma = c/\sqrt{\log(1/\epsilon)}$ for a sufficiently small constant $c > 0$. \square

8 Quantum lower-bound witnessing for degree-4 SYK

The main content of this section is verifying the correct order of magnitude for the largest eigenvalue of the SYK₄ model:

Theorem 8.1. *For $\mathbf{h} \sim \text{SYK}_4(n)$, we have $\text{Opt}(\mathbf{h}) \geq \Omega(\sqrt{n})$ with high probability.*

Remark 8.2. Recall the physics prediction is that $\text{Opt}(\mathbf{h})$ concentrates around $\frac{1}{2\sqrt{2}}\sqrt{n}$. We have not attempted to optimize the implied constant in the $\Omega(\cdot)$ of our theorem.

Having shown this theorem, it will follow straightforwardly from its proof that there is an efficient quantum *witnessing* algorithm that establishes this lower bound.

Actually, it will be more convenient to work with a “2-colored” (or “1-variable decoupled”) version of the SYK model, which we define below; this 2-colored version is a specific case of the models considered in [GR17]. In [Section 8.2](#) we will see an easy way to extend our results for the 2-colored variant to the standard SYK model.

Definition 8.3. We define the 2-colored SYK₄ model as follows. Given even $n_1, n \in \mathbb{N}$, we work in $\mathfrak{C}(K_{n_1+n})$, with the indeterminates named $\varphi_1, \dots, \varphi_{n_1}, \chi_1, \dots, \chi_n$. We write $\mathbf{h} \sim \text{SYK}_4^{2\text{col}}(n_1, n)$ to denote that \mathbf{h} is constructed as

$$\mathbf{h} = \frac{\mathbf{i}}{\sqrt{n}} \sum_{m=1}^n \tau_m \chi_m, \quad (30)$$

where each $\tau_m \sim \text{SYK}_3(\varphi_1, \dots, \varphi_{n_1})$ independently; i.e.,

$$\tau_m = \frac{\mathbf{i}}{\sqrt{\binom{n_1}{3}}} \sum_{\substack{S \subseteq [n_1] \\ |S|=3}} \mathbf{J}_S^{(m)} \varphi^S, \quad (31)$$

where each $\mathbf{J}_S^{(m)}$ is an independent standard Gaussian.

Remark 8.4. One can infer from the proof of [Theorem 8.8](#) that, for n a multiple of 4,

$$\mathbf{E}_{\mathbf{h} \sim \text{SYK}_4(n)} [\text{Opt}(\mathbf{h})] \geq \frac{3\sqrt{3}}{8} \mathbf{E}_{\mathbf{h}_1 \sim \text{SYK}_4^{2\text{col}}(\frac{3}{4}n, \frac{1}{4}n)} [\text{Opt}(\mathbf{h}_1)].$$

Remark 8.5. In the remainder of this section we may sometimes tacitly assume that n or n_1 is a multiple of 2 or 4 or 8. This is purely for convenience, and the reader may easily verify these assumptions are inessential.

8.1 The main variational argument

Theorem 8.6. *For $\mathbf{h} \sim \text{SYK}_4^{2\text{col}}(n_1, n)$ with $n_1 = \Theta(n)$ even, we have $\text{Opt}(\mathbf{h}) \geq \Omega(\sqrt{n})$ except with probability at most $\exp(-\Omega(n))$.*

Proof. Initially we will work in slightly more generality, with a setup similar to that of [Section 5](#). Specifically, we assume a self-adjoint polynomial

$$h = \frac{\mathbf{i}}{\sqrt{n}} \sum_{m=1}^n \tau_m \chi_m$$

where initially we only need to assume that all indeterminates τ_j and χ_k are self-adjoint, and that each τ_j anticommutes with each χ_k . We will be motivated by the idea that the τ_j ’s “almost”

satisfy $\tau_j^2 = 1$ and $\tau_j \tau_k = -\tau_k \tau_j$. To this end¹⁷, we artificially adjoin to our algebra additional self-adjoint indeterminates $\sigma_1, \dots, \sigma_n$ that *do* satisfy $\sigma_j^2 = 1$ and $\sigma_j \sigma_k = -\sigma_k \sigma_j$, and that additionally anticommute with all χ_j 's and τ_j 's. Now if

$$h' := \frac{i}{\sqrt{n}} \sum_{m=1}^n \sigma_m \chi_m,$$

it would be easy to optimize h' ; we could choose the Gaussian state

$$\rho_0 := (1 - i\sigma_1 \chi_1) \cdots (1 - i\sigma_n \chi_n)$$

which achieves $\text{tr}(\rho_0 h') = \sqrt{n}$ (similarly to [Example 7.7](#)). The idea is now to “rotate” the σ_j 's toward the τ_j 's by introducing the skew-adjoint

$$\zeta := \tau_1 \sigma_1 + \cdots + \tau_n \sigma_n$$

and then defining (for $\theta \in \mathbb{R}$)

$$\rho_\theta := \text{Ad}_{e^{-\theta\zeta}}(\rho_0), \quad \text{where } \text{Ad}_U(g) = UgU^{-1}. \quad (32)$$

(Here, to make sense of $e^{-\theta\zeta}$, recall that we will ultimately take the τ_j 's to be polynomials in the additional anticommuting indeterminates $\varphi_1, \dots, \varphi_{n_1}$, n_1 even; then all indeterminates will be residing in the algebra $\mathfrak{C}(K_{n_1+2n})$. This algebra is isomorphic to $\mathbb{C}^{D \times D}$ for $D = n_1/2 + n$, and the matrix exponential is defined therein.) This ρ_θ is always a state, since $i\zeta$ is self-adjoint and hence $e^{-\theta\zeta}$ is unitary. We now have

$$\text{tr}(\rho_\theta h) = \text{tr}(\text{Ad}_{e^{-\theta\zeta}}(\rho_0)h) = \text{tr}(\rho_0 \text{Ad}_{e^{i\theta\zeta}}(h)) = \text{tr}(\rho_0 h_\theta), \quad \text{where } h_\theta := \text{Ad}_{e^{\theta\zeta}}(h). \quad (33)$$

We may now use the following standard Baker–Campbell–Hausdorff-type formula:

Proposition 8.7. *For $[\cdot, \cdot]$ denoting the commutator, we have*

$$h_\theta = h + \theta[\zeta, h] + \theta^2 \int_0^1 (1-s) \text{Ad}_{e^{s\theta\zeta}}([\zeta, [\zeta, h]]) \, ds.$$

Proof. (Sketch.) This formula is Taylor-expanding $A(\theta) = \text{Ad}_{e^{\theta\zeta}}$ around $\theta = 0$, using that $A'(\theta) = aA(\theta)$ for $a(\cdot) = [\zeta, \cdot]$ (see, e.g., [\[Ros02, Ch. 1.2\]](#)), and also using that $[\zeta, A(\theta)(g)] = A(\theta)([\zeta, g])$ since $A(\theta)$ and ζ commute. \square

Putting [Proposition 8.7](#) into [Equation \(33\)](#) and using $\text{tr}(\rho_0 h) = 0$, we infer

$$\begin{aligned} \text{tr}(\rho_\theta h) &= \theta \text{tr}(\rho_0 [\zeta, h]) + \theta^2 \mathbf{E}_{s \sim [0,1]} \left[(1-s) \text{tr}(\rho_{s\theta} [\zeta, [\zeta, h]]) \right] \\ &\geq \theta \text{tr}(\rho_0 [\zeta, h]) - \theta^2 \text{Opt}_\pm([\zeta, [\zeta, h]]), \end{aligned} \quad (34)$$

where we used the triangle inequality.

¹⁷The construction here perhaps be regarded as a quantum version of the classical construction in [\[Has19\]](#): we start with some state (in that paper a random classical state, and here a Gaussian state), and apply some transformation (there a local update, here a unitary) to improve the expectation value of the objective in the state up to linear order in some parameter. In both constructions, the problem is to control higher order corrections in that parameter, though this is done very differently in both cases.

Now is the point at which we will specialize to the 2-colored SYK situation from the theorem statement (and [Definition 8.3](#)), with h replaced by \mathbf{h} as in [Equation \(30\)](#), and with the τ_j 's being drawn from $\text{SYK}_3(n_1)$ as in [Equation \(31\)](#). (We will also write ρ_θ in place of ρ_θ , as it has a dependence on the random τ_j 's.) We have

$$[\zeta, \mathbf{h}] = \frac{i}{\sqrt{n}} \sum_{j,k=1}^n [\tau_j \sigma_j, \tau_k \chi_k],$$

and one can see that only the $j = k$ terms survive when $\text{tr}(\rho_0 \cdot)$ is applied. We deduce

$$\text{tr}(\rho_0 [\zeta, \mathbf{h}]) = \frac{1}{\sqrt{n}} \sum_{k=1}^n \text{tr}(\{\tau_k, \tau_k\}),$$

where $\{\cdot, \cdot\}$ denotes the anticommutator. It is not hard to compute that

$$\text{tr}(\{\tau_k, \tau_k\}) = \frac{2}{\binom{n_1}{3}} \sum_{\substack{S \subseteq [n_1] \\ |S|=3}} (\mathbf{J}^{(k)})^2,$$

and hence

$$\mathbf{E}[\text{tr}(\rho_0 [\zeta, \mathbf{h}])] = 2\sqrt{n}, \quad \mathbf{Pr}[\text{tr}(\rho_0 [\zeta, \mathbf{h}]) < \sqrt{n}] \leq \exp(-\Omega(n_1^3)), \quad (35)$$

the inequality here being a standard tail-bound for chi-squared random variables.

To complete the analysis of [Inequality \(34\)](#), we must now estimate $\text{Opt}_\pm([\zeta, [\zeta, \mathbf{h}]])$. As this involves a lot of bookkeeping, we will stop keeping track of constants and henceforth use a softer approach. Recall that we are assuming $C_1 n \leq n_1 \leq C_2 n_2$ for some constants $C_1, C_2 > 0$. We henceforth use C_3, C_4, \dots to denote either unspecified positive constants, or else quantities like $\binom{n}{3}/n^3$ which — while technically dependent on n — are nevertheless bounded above and below by universal positive constants.

For the sake of symmetry, we now introduce new independent mean-zero Gaussians \mathbf{G}_{rsta} , for $r, s, t \in [n_1]$, $a \in [n]$, where \mathbf{G}_{rsta} has variance 1 if r, s, t are all distinct, and otherwise \mathbf{G}_{rsta} has variance 0. Then (using the fact that the sum and difference of independent Gaussians is Gaussian) it is equivalent in distribution to reexpress each τ_a as

$$\tau_a = \frac{C_3 i}{n^{3/2}} \sum_{r,s,t} \mathbf{G}_{rsta} \varphi_r \varphi_s \varphi_t.$$

Now

$$[\zeta, [\zeta, \mathbf{h}]] = \frac{C_4}{n^5} \sum_{r,s,t,u,v,w,x,y,z,a,b,d} [\mathbf{G}_{rsta} \varphi_r \varphi_s \varphi_t \sigma_a, [\mathbf{G}_{uvwb} \varphi_u \varphi_v \varphi_w \sigma_b, \mathbf{G}_{xyzd} \varphi_x \varphi_y \varphi_z \chi_d]].$$

Considering the inner commutator, we have (using [Fact 2.13](#)) that $\mathbf{G}_{uvwb} \varphi_u \varphi_v \varphi_w \sigma_b$ and $\mathbf{G}_{xyzd} \varphi_x \varphi_y \varphi_z \chi_d$ anticommute iff they share an odd number of indeterminates. In particular this inner commutator is nonzero only if $|\{u, v, w\} \cap \{x, y, z\}| \geq 1$. Furthermore, such a surviving commutator only anticommutes with $\mathbf{G}_{rsta} \varphi_r \varphi_s \varphi_t \sigma_a$ provided there is another indeterminate in common, meaning $|\{r, s, t\} \cap (\{u, v, w\} \triangle \{x, y, z\})| + 1[a = b] \geq 1$. Thus

$$[\zeta, [\zeta, \mathbf{h}]] = \frac{C_5}{n^5} \sum_{r,s,\dots,d} f(r, s, \dots, d) \mathbf{G}_{rsta} \mathbf{G}_{uvwb} \mathbf{G}_{xyzd} \varphi_r \varphi_s \varphi_t \sigma_a \varphi_u \varphi_v \varphi_w \sigma_b \varphi_x \varphi_y \varphi_z \chi_d$$

for some 0-1 indicator f , where f depends only on certain equalities and non-equalities among its arguments and has the property

$$f(r, s, \dots, d) = 0 \text{ unless } |\{u, v, w\} \cap \{x, y, z\}| \geq 1 \text{ and } (|\{r, s, t\} \cap (\{u, v, w\} \triangle \{x, y, z\})| + 1[a = b] \geq 1). \quad (36)$$

We are now in a position to apply [Theorem 3.2](#), together with [Remarks 3.3](#) and [3.9](#); these imply

$$\mathbf{E}[\text{Opt}_{\pm}([\zeta, [\zeta, \mathbf{h}]]) \leq C_0 \sqrt{n}, \quad \mathbf{Pr}[\text{Opt}_{\pm}([\zeta, [\zeta, \mathbf{h}]]) \leq \tfrac{1}{2} C_0 \sqrt{n}] \leq \exp(-\Omega(n)) \quad (37)$$

(for some constant $C_0 < \infty$) provided we can prove via the k th-moment method ($k = \Theta(n)$) that

$$\mathbf{Pr}[|P(\mathbf{G})| \geq C_0 \sqrt{n}] \leq \exp(-C_6 n), \quad P(\mathbf{G}) := \frac{C_5}{n^5} \sum_{r, s, \dots, d} f(r, s, \dots, d) \mathbf{G}_{rsta} \mathbf{G}_{uvwb} \mathbf{G}_{xyzd}. \quad (38)$$

(Here C_0 may depend on C_1, \dots, C_6). Once we have this, we can put [Inequality \(37\)](#) and [Equation \(35\)](#) into [Inequality \(34\)](#), and then

$$\theta = \frac{1}{C_0} \implies \mathbf{E}[\text{tr}(\boldsymbol{\rho}_{\theta} \mathbf{h})] \geq \frac{1}{C_0} \sqrt{n}, \quad \mathbf{Pr}\left[\text{tr}(\boldsymbol{\rho}_{\theta} \mathbf{h}) \leq \frac{1}{2C_0} \sqrt{n}\right] \leq \exp(-\Omega(n)), \quad (39)$$

as needed.

The remainder of the proof is therefore devoted to establishing [Inequality \(38\)](#), by the k th-moment method. (We will in fact choose $k = n$, which we assume is an even integer.) Notice that in the sum defining P , the presence of f forces at least two indices to match; hence this sum effectively has only $O(n^{10})$ terms, not $O(n^{12})$. Thus $\mathbf{E}[P(\mathbf{G})^2] \leq O(1)$, and so $\mathbf{Pr}[|P(\mathbf{G})| \geq C_0 \sqrt{n}] \leq \exp(-\Theta(n))$ is plausible — but only if $P(\mathbf{G})$ has Gaussian-like tails, despite being a degree-3 polynomial in Gaussians.

Let us now make some further simplifications. First, recall that \mathbf{G}_{rsta} is defined to be 0 if r, s, t are not all distinct. We may drop this assumption, and assume that *every* \mathbf{G}_{rsta} is a standard Gaussian, by redefining $f(r, s, \dots, d)$ so that *it* is 0 unless $|\{r, s, t\}| = |\{u, v, w\}| = |\{x, y, z\}| = 3$. Next, this $f(r, s, \dots, d)$ is some Boolean function of all possible Kronecker delta functions δ_{jk} and their negations $(1 - \delta_{jk})$, for $j, k \in \{r, s, \dots, d\}$. By using inclusion-exclusion, we may therefore write

$$P(\mathbf{G}) = \sum_{\ell=1}^{C_7} \pm P_{\ell}(\mathbf{G}), \quad P_{\ell}(\mathbf{G}) := \frac{C_5}{n^5} \sum_{r, s, \dots, d} f_{\ell}(r, s, \dots, d) \mathbf{G}_{rsta} \mathbf{G}_{uvwb} \mathbf{G}_{xyzd},$$

where each f_{ℓ} is a product only of Kronecker deltas δ_{jk} (and *not* their negations). Furthermore, because each f_{ℓ} must enforce [Equation \(36\)](#), it must be *viable*, by which we mean it must include at least $\delta_{jk} \delta_{\ell m}$ for indices $j \in \{r, s, t, a\}$, $k, \ell \in \{u, v, w, b\}$ *distinct*, and $m \in \{x, y, z, d\}$. By a union bound, and adjusting constants, it therefore suffices to achieve

$$\mathbf{Pr}[|P_{\ell}(\mathbf{G})| \geq C_0 \sqrt{n}] \leq \exp(-C_8 n)$$

by the n th-moment method, for all viable P_{ℓ} . Given that we will use the moment method, observe that if the summands in some $P_{\ell}(\mathbf{G})$ are a subset of those in some other $P_{\ell'}(\mathbf{G})$, it suffices for us to handle $P_{\ell'}(\mathbf{G})$; this is because the latter's even moments are only larger (using the fact that Gaussian monomials have nonnegative expectation). Thus it suffices for us to handle the viable $P_{\ell}(\mathbf{G})$'s that include a *minimal* number of Kronecker deltas, namely 2. As we now have symmetry between the indices in our \mathbf{G} 's, it therefore remains to handle one particular minimal case; say, $v = t$ and $x = u$, meaning

$$P_0(\mathbf{G}) := \frac{C_5}{n^5} \sum_{r, s, t, u, w, y, z, a, b, d} \mathbf{G}_{rsta} \mathbf{G}_{utwb} \mathbf{G}_{uyzd}.$$

To obtain the required tail bound of the form $\Pr[|P_0(\mathbf{G})| \geq C_0\sqrt{n}] \leq \exp(-C_8n)$, it remains to establish the following inequality:

$$\|P_0(\mathbf{G})\|_n \leq O(\sqrt{n}). \quad (40)$$

To show this we would like to use the decoupling aspect of [Theorem 3.10](#); however we don't yet have the property “ $A_{j_1 \dots j_q} = 0$ when j_1, \dots, j_q are not all distinct”. Thus we separate out the various equality/nonequality cases for $(r, s, t, a), (u, t, w, b), (u, y, z, d)$, writing

$$P_0(\mathbf{G}) = R(\mathbf{G}) + Q_1(\mathbf{G}) + Q_2(\mathbf{G}) + Q_3(\mathbf{G}) + P'_0(\mathbf{G}), \quad (41)$$

where

$$\begin{aligned} R(\mathbf{G}) &:= \frac{C_5}{n^5} \sum_{r,s,t,a} \mathbf{G}_{rsta}^3, & Q_1(\mathbf{G}) &:= \frac{C_5}{n^5} \sum_{\substack{r,s,t,u,w,a,b \\ (r,s,t,a),(u,t,w,b), \text{ distinct}}} \mathbf{G}_{rsta} \mathbf{G}_{utwb}^2, \\ Q_2, Q_3 &\text{ defined similarly,} & P'_0(\mathbf{G}) &:= \frac{C_5}{n^5} \sum_{\substack{r,s,t,u,w,y,z,a,b,d \\ (r,s,t,a),(u,t,w,b),(u,y,z,d) \text{ distinct}}} \mathbf{G}_{rsta} \mathbf{G}_{utwb} \mathbf{G}_{uyzd}. \end{aligned}$$

Bounding R is very easy; we can use, say, hypercontractivity [[Jan97](#), Thm. 5.10] to deduce

$$\|R(\mathbf{G})\|_n \leq \sqrt{n-1}^3 \|R(\mathbf{G})\|_2 \leq O(n^{-3.5}) \cdot \left\| \sum_{r,s,t,a} \mathbf{G}_{rsta}^3 \right\|_2 = O(n^{-3.5}) \cdot \sqrt{O(n^4)} = O(n^{-1.5}) \ll O(\sqrt{n}).$$

As for Q_1 (and similarly Q_2, Q_3), we can first drop the condition “ $(r, s, t, a), (u, t, w, b)$ distinct” in its definition for simplicity, since (as previously noted) additional terms can only make even moments larger. Then, using the decoupling aspect of [Theorem 3.10](#), it suffices to obtain the bound

$$\left\| \sum_{r,s,t,a,u,w,b} \mathbf{G}'_{rsta} \mathbf{G}_{utwb}^2 \right\|_n \leq O(n^{5.5}), \quad (42)$$

where $(\mathbf{G}')_{rsta}$ are new independent standard Gaussians. This is equivalent to

$$\left\| \sum_{t=1}^n \mathbf{G}''_t \mathbf{q}_t^2 \right\|_n^n = \mathbf{E}_{\mathbf{q}_t} \mathbf{E}_{\mathbf{G}''_t} \left[\left(\sum_{t=1}^n \mathbf{G}''_t \mathbf{q}_t \right)^n \right] \leq O(n^4)^n,$$

where $(\mathbf{G}'')_t$ are independent standard Gaussians, and $(\mathbf{q}_t)_t$ are independent chi-squared random variables with n^3 degrees of freedom. For fixed outcomes $\mathbf{q}_t = \chi_t$, the random variable $\sum_t \mathbf{G}''_t \chi_t$ is a mean-zero Gaussian with variance $\nu^2 := \sum_t \chi_t^2$, and hence its n th moment is bounded by $\nu^n \cdot \sqrt{n}^n$. Thus to verify [Inequality \(42\)](#) it suffices to show

$$\mathbf{E}[\nu^n] \leq O(n^{3.5})^n. \quad (43)$$

Now we use the following well known tail bound for the \mathbf{q}_t 's (see, e.g. [[LM00](#), Lem. 1]):

$$\Pr[\mathbf{q}_t \geq (4c+1)n^3] \leq \exp(-cn^3) \quad \forall c \geq 1.$$

A union bound now easily implies

$$\Pr[\nu \geq 5cn^{3.5}] \leq \exp(-\tfrac{1}{2}cn^3) \quad \forall c \geq 1 \quad \implies \quad \Pr[\nu' \geq c] \leq \exp(-\tfrac{1}{2}n^3)^c, \quad \nu' := \nu/(5n^{3.5}),$$

and from this one easily confirms [Inequality \(43\)](#) and hence [Inequality \(42\)](#):

$$\mathbf{E}[(\nu')^n] \leq 1 + \sum_{j \in \mathbb{N}} 2^{(j+1)n} \cdot \Pr[2^j \leq \nu' \leq 2^{j+1}] \leq 1 + \sum_{j \in \mathbb{N}} 2^{(j+1)n} \cdot \exp(-\tfrac{1}{2}n^3)^{2^j} \leq O(1).$$

Finally, to complete the proof of [Inequality \(40\)](#), it remains to handle P'_0 from [Equation \(41\)](#); i.e., to bound $\|P'_0(\mathbf{G})\|_n \leq O(\sqrt{n})$. Using the decoupling aspect of [Theorem 3.10](#), it in turn suffices to bound the n th norm of

$$\begin{aligned} P_0(\mathbf{G}^{(1)}, \mathbf{G}^{(2)}, \mathbf{G}^{(3)}) &= \frac{C_5}{n^5} \sum_{r,s,t,u,w,y,z,a,b,d} \mathbf{G}_{rsta}^{(1)} \mathbf{G}_{utwb}^{(2)} \mathbf{G}_{uyzd}^{(3)} \\ &= \frac{C_9}{n} \sum_{t,u} \mathbf{H}_t \mathbf{K}_{tu} \mathbf{L}_u, \end{aligned}$$

where $(\mathbf{H}_t)_t, (\mathbf{K}_{tu})_{t,u}, (\mathbf{L}_u)_u$ (for $t, u \in [n]$) are new independent standard Gaussians. [Theorem 3.10](#) now implies

$$\|P'_0(\mathbf{G})\|_n \leq \frac{C_{10}}{n} \left(\sqrt{n} \|A\|_{\{1,2,3\}} + n \|A\|_{\{1,2\},\{3\}} + n \|A\|_{\{1,3\},\{2\}} + n \|A\|_{\{2,3\},\{1\}} + n^{3/2} \|A\|_{\{1\},\{2\},\{3\}} \right),$$

where $A \in \{0,1\}^{n \times n^2 \times n}$ has $A_{h,jk,\ell} = 1$ iff $h = j$ and $k = \ell$. To obtain the desired tail bound of the form $\Pr[|P_0(\mathbf{G})| \geq C_0 \sqrt{n}] \leq \exp(-C_8 n)$, it suffices to show $\|P_0\|_n \leq O(\sqrt{n})$. But this is true, as one can easily compute

$$\begin{aligned} \|A\|_{\{1,2,3\}} &= n, & \|A\|_{\{1,2\},\{3\}} &= \|A\|_{\{2,3\},\{1\}} = \|\mathbb{1}_{n \times n} \otimes \underbrace{(1, \dots, 1)}_n\|_{\text{op}} = \sqrt{n}, \\ \|A\|_{\{1,3\},\{2\}} &= \|\mathbb{1}_{n^2 \times n^2}\|_{\text{op}} = 1, \end{aligned}$$

and

$$\|A\|_{\{1\},\{2\},\{3\}} = \max_{\substack{x,z \in \mathbb{R}^n, y \in \mathbb{R}^{n^2} \\ \|x\|, \|y\|, \|z\| \leq 1}} \sum_{t,u} x_t y_{tu} z_u \leq \max_{x,y,z} \sqrt{\sum_{t,u} x_t^2 z_u^2} \sqrt{\sum_{t,u} y_{tu}^2} = 1,$$

using Cauchy–Schwarz. □

8.2 Efficient quantum lower-bound witnessing

The argument based on variational states used to prove [Theorem 8.6](#) can be readily turned into an efficient quantum lower-bound witnessing algorithm of the type discussed in [Section 2.7.1](#). (In the remainder of this section we will identify elements of $\mathfrak{C}(\Gamma)$, $\Gamma = K_N$, with their representations in $\mathbb{C}^{N/2 \times N/2}$ under π_Γ .)

Theorem 8.8. *For a universal constant $c_0 > 0$, there is a $\text{poly}(n)$ -time, $O(n)$ -space quantum algorithm W that, on input a degree-4 homogeneous $h \in \mathfrak{C}(K_n)$, outputs a quantum state $\rho \in \mathbb{C}^{D \times D}$ (with $D = 2^{n/2}$ as in [Proposition 2.15](#)). Except with probability $\exp(-\Omega(n))$ over the draw of $\mathbf{h} \sim \text{SYK}_4(n)$, it holds for $\rho = W(\mathbf{h})$ that $\text{tr}(\rho \mathbf{h}) \geq c_0 \sqrt{n}$.*

Remark 8.9. As described in [Section 2.7.1](#), there is a $\text{poly}(n) \log(1/\delta)$ -time randomized quantum procedure that, given *any* worst-case h and corresponding $\rho = W(h)$, certifies that $\text{tr}(\rho h) \geq c_0 \sqrt{n}$ (up to any additive $1/\text{poly}(n)$) whenever it holds, except with probability at most δ .

To prove [Theorem 8.8](#), we first prove [Proposition 8.10](#) below, which establishes it for the 2-colored SYK model analyzed in [Theorem 8.6](#). Afterwards, extending to the usual SYK model is not difficult.

Proposition 8.10. *[Theorem 8.8](#) holds with $\mathbf{h} \sim \text{SYK}_4^{2\text{col}}(\frac{3}{4}n, \frac{1}{4}n)$ in place of $\mathbf{h} \sim \text{SYK}_4(n)$.*

Proof. The input h to the algorithm resides in an algebra with $\frac{3}{4}n + \frac{1}{4}n$ anticommuting indeterminates, which we will simply call χ_1, \dots, χ_n (rather than $\varphi_1, \dots, \varphi_{\frac{3}{4}n}, \chi_1, \dots, \chi_{\frac{1}{4}n}$). These are represented by the γ -matrices from [Equation \(6\)](#), over an $n/2$ -qubit Hilbert space.

Now the target variational state ρ_θ defined in [Equation \(32\)](#) (with θ chosen as in [Equation \(39\)](#)) in fact resides in $\mathfrak{C}(\Gamma)$ for $\Gamma = K_{n+\frac{1}{4}n}$, since it involves the additional indeterminates $\sigma_1, \dots, \sigma_{n/4}$. To accommodate this the algorithm will actually work in the enlarged Hilbert space with $n/8$ additional qubits. But since the input h does not depend on the σ_j 's, it's easy to see that if the algorithm manages to construct the quantum state ρ_θ , it may as its final step simply discard (trace out) its last $n/8$ qubits, leaving a quantum state ρ that has $\text{tr}(\rho h) = \text{tr}(\rho_\theta h)$.

Thus we only need to show the algorithm can construct ρ_θ . Indeed, by slightly adjusting the constant c_0 in the statement of [Theorem 8.8](#), it suffices for it to construct ρ_θ up to sufficiently small $1/\text{poly}(n)$ error in trace distance. The first step of the algorithm is to construct the Gaussian state ρ_0 . This may be done very efficiently; this state is maximally mixed on the first $3n/8$ qubits, and on the remaining $n/4$ qubits it is a stabilizer state [[Got97](#)] with a simple description. (Indeed, had the last $n/2$ indeterminates been ordered as $\chi_{3n/4}, \sigma_1, \chi_{3n/4+1}, \sigma_2, \dots$, then ρ_0 would simply be $2^{n/8} |00 \dots 0\rangle \langle 00 \dots 0|$ on these qubits, as in [Example 7.2](#).)

It remains for the algorithm to rotate this state by $\exp(-\theta\zeta)$ (up to polynomially small error). This rotation can be accomplished efficiently: it is the same as Hamiltonian evolution under the Hamiltonian $-i\zeta$ for time θ . This Hamiltonian evolution can be done in polynomial time to any desired inverse polynomial error, for example by Trotter–Suzuki evolution, since the “Hamiltonian” $-i\zeta$ is the sum of polynomially many terms (namely, the monomials in the Majorana operators), and unitary evolution under a given term can be done efficiently as each term is a product of Pauli operators. More efficient algorithms exist which can provide a polynomial speedup in n and exponential speedup in error [[BBN19](#)]. \square

It now remains to move from the 2-colored SYK model to the usual one. For this, let us partition $\binom{[n]}{4} = A_{\text{in}} \sqcup A_{\text{out}}$, where

$$A_{\text{in}} = \{\{j_1, j_2, j_3, j_4\} : j_1 < j_2 < j_3 \leq \frac{3}{4}n < j_4\}, \quad A_{\text{out}} = \binom{[n]}{4} \setminus A_{\text{in}},$$

and let us also write any homogeneous degree-4 polynomial h as $h_{\text{in}} + h_{\text{out}}$, where h_{in} (respectively, h_{out}) contains the monomials of h corresponding to indices $S \in A_{\text{in}}$ (respectively, A_{out}). It is then easy to see that for $\mathbf{h} \sim \text{SYK}_4(n)$ it holds that \mathbf{h}_{in} is distributed as $\text{SYK}_4^{2\text{col}}(\frac{3}{4}n, \frac{1}{4}n)$ up to a scaling factor; specifically, we have

$$c \cdot \mathbf{h}_{\text{in}} \sim \text{SYK}_4^{2\text{col}}(\frac{3}{4}n, \frac{1}{4}n) \quad \text{for } c := \sqrt{\frac{\binom{n}{4}}{\binom{3n/4}{3}(n/4)}} \leq \frac{8}{3\sqrt{3}}. \quad (44)$$

We may now establish [Theorem 8.8](#).

Theorem 8.8. Let W be the quantum algorithm that, on input $h \in \mathfrak{C}(K_n)$ homogeneous of degree 4, applies the algorithm from [Proposition 8.10](#) to $c \cdot h_{\text{in}}$, where c is as in [Equation \(44\)](#). Suppose now

that

$$\mathbf{h} = \frac{1}{\sqrt{\binom{n}{4}}} \sum_{\substack{S \subseteq [n] \\ |S|=4}} \mathbf{J}_S \chi^S \sim \text{SYK}_4(n).$$

Then as discussed, $c \cdot \mathbf{h}_{\text{in}}$ is distributed as $\text{SYK}_4^{2\text{col}}(\frac{3}{4}n, \frac{1}{4}n)$, and hence the output $\boldsymbol{\rho}$ of [Proposition 8.10](#)'s algorithm satisfies

$$\Pr_{\mathbf{J}_S: S \in A_{\text{in}}} [\text{tr}(\boldsymbol{\rho} \mathbf{h}_{\text{in}}) \geq \frac{c'_0}{c} \sqrt{n}] \geq 1 - \exp(-\Omega(n)) \quad (45)$$

(for some constant $c'_0 > 0$). On the other hand, for *any* fixed outcome $\boldsymbol{\rho} = \rho$ we have

$$\mathbf{E}_{\mathbf{J}_S: S \in A_{\text{out}}} [\text{tr}(\rho \mathbf{h}_{\text{out}})] = \mathbf{E} \left[\frac{1}{\sqrt{\binom{n}{4}}} \sum_{S \in A_{\text{out}}} \text{tr}(\rho \chi^S) \mathbf{J}_S \right] = 0.$$

Moreover, since $|\text{tr}(\rho \chi^S)| \leq 1$ for each S , Gaussian concentration implies that

$$\Pr_{\mathbf{J}_S: S \in A_{\text{out}}} [|\text{tr}(\rho \mathbf{h}_{\text{out}})| \geq \frac{c'_0}{2c} \sqrt{n}] \leq \exp(-\Omega(n)). \quad (46)$$

Using

$$\text{tr}(\boldsymbol{\rho} \mathbf{h}) = \text{tr}(\boldsymbol{\rho} \mathbf{h}_{\text{in}}) + \text{tr}(\boldsymbol{\rho} \mathbf{h}_{\text{out}})$$

and taking $c_0 = \frac{c'_0}{2c}$, [Inequalities \(45\)](#) and [\(46\)](#) complete the proof. \square

Open problems

Here we briefly list some open problems not previously mentioned in the paper:

- Can efficient certification algorithms, within a constant factor, be given for the SYK model with $q = 3$ or $q = 6$?
- For SYK of degree 6, can one show that constant-degree SOS is strongly fooled (i.e., degree- k SOS fails to certify an $O(\sqrt{n})$ upper bound, for any constant k)?
- Do our certification algorithms translate to the *sparse* SYK model [\[XSSS20, GJRV21\]](#), in which only $\Theta(n)$ coefficients are chosen to be nonzero?
- What can be said about the closely-related Erdős–Schröder model [\[ES14\]](#) of quantum spin glasses?
- For which deterministic SYK-like models can we prove SOS bounds? For example, the eigenvalue bounds in [\[KMPT18\]](#) can be proven within SOS.
- Can one give evidence that classical algorithms cannot efficiently certify $\text{Opt}(\mathbf{p}) \geq \Omega(\sqrt{n})$ (with high probability) for $\mathbf{p} \sim \text{SYK}_4(n)$?
- What can be said about optimization — even of degree-1 Hamiltonians — within $\mathfrak{C}(\Gamma)$ when $\Gamma \sim G(n, p)$ is an Erdős–Rényi random graph?
- What is the largest possible ratio between $\text{Opt}(h)$ and $\text{Opt}_{\text{Gauss}}(h)$ for $h \in \mathfrak{C}(K_n)$ of degree 4? We know it is at most $O(n)$ and at least $\Omega(\sqrt{n})$ (the latter because of typical $\text{SYK}_4(n)$ instances).

- In the same way that [Theorem 7.18](#) extends [\[CW04\]](#), can we extend [\[GW95\]](#) by showing that in the context of [Theorem 7.18](#), if $\text{Opt}(h) \geq (1 - \delta)\sqrt{\binom{n/2}{2}}$ then an efficient algorithm can deliver a Gaussian state achieving expectation value at least $(1 - \delta')\sqrt{\binom{n/2}{2}}$, for some δ' with $\delta' \rightarrow 0$ as $\delta \rightarrow 0$ (e.g., $\delta' = O(\sqrt{\delta})$)?

Acknowledgments

The second author would like to thank Pravesh Kothari, Sidhanth Mohanty, Tselil Schramm, and William Slofstra for their helpful comments. The authors would also like to thank Maarten Stroeks for pointing out the need to handle R, Q_1, Q_2, Q_3 in [Equation \(41\)](#).

References

- [AAV13] Dorit Aharonov, Itai Arad, and Thomas Vidick. The quantum PCP conjecture. *ACM SIGACT News*, 44(2):47–79, 2013. [1.3](#)
- [ABI86] Noga Alon, László Babai, and Alon Itai. A fast and simple randomized parallel algorithm for the maximal independent set problem. *Journal of Algorithms*, 7(4):567–583, 1986. [7.20](#)
- [AH11] Per Austrin and Johan Håstad. Randomly supported independence and resistance. *SIAM Journal on Computing*, 40(1):1–27, 2011. [7.20](#)
- [AW02] Rudolf Ahlswede and Andreas Winter. Strong converse for identification via quantum channels. *Transactions on Information Theory*, 48(3):569–579, 2002. [3.1](#)
- [BBN19] Ryan Babbush, Dominic Berry, and Hartmut Neven. Quantum simulation of the Sachdev–Ye–Kitaev model by asymmetric qubitization. *Physical Review A*, 99(4):040301, 2019. [8.2](#)
- [BF71] Oriol Bohigas and Jorge Flores. Two-body random Hamiltonian and level density. *Physics Letters B*, 34(4):261–263, 1971. [1.4](#)
- [BGKT19] Sergey Bravyi, David Gosset, Robert König, and Kristan Temme. Approximation algorithms for quantum many-body problems. *Journal of Mathematical Physics*, 60(3):032203, 2019. [1.4](#), [1.5](#), [7.1](#), [7.1](#), [7.9](#), [7.10](#), [7.1](#), [7.2](#), [7.2](#)
- [BH13] Fernando Brandão and Aram Harrow. Product-state approximations to quantum ground states. In *Proceedings of the 45th Annual ACM Symposium on Theory of Computing (STOC)*, pages 871–880, 2013. [1.3](#)
- [BKS97] Marek Bożejko, Burkhard Kümmerer, and Roland Speicher. q -Gaussian processes: non-commutative and classical aspects. *Communications in Mathematical Physics*, 185(1):129–154, 1997. [3.2](#)
- [Bra05] Sergey Bravyi. Lagrangian representation for fermionic linear optics. *Quantum Information & Computation*, 5(3):216–238, 2005. [1.3](#), [7](#)
- [Col63] A. John Coleman. Structure of fermion density matrices. *Reviews of Modern Physics*, 35(3):668–686, 1963. [1.3](#)

- [CW04] Moses Charikar and Anthony Wirth. Maximizing quadratic programs: extending Grothendieck’s inequality. In *Proceedings of the 45th annual Symposium on Foundations of Computer Science (FOCS)*, pages 54–60. IEEE, 2004. 1.1, 7.2, 8.2
- [Dav96] Kenneth Davidson. *C*-algebras by example*. American Mathematical Society, 1996. 2.4
- [DEF78] Michel Deza, Paul Erdős, and Peter Frankl. Intersection properties of systems of finite sets. *Proceedings of the London Mathematical Society. Third Series*, 36(2):369–384, 1978. 4.1, 4.12, 4.2, 4.23
- [Del73] Philippe Delsarte. An algebraic approach to the association schemes of coding theory. *Philips Research Reports. Supplements*, (10):vi+97, 1973. 4.2, 4.24
- [DLTW08] Andrew Doherty, Yeong-Cherng Liang, Ben Toner, and Stephanie Wehner. The quantum moment problem and bounds on entangled multi-prover games. In *Proceedings of the 23rd Annual IEEE Conference on Computational Complexity (CCC)*, pages 199–210, 2008. 1.3
- [EKR61] Paul Erdős, Chao Ko, and Richard Rado. Intersection theorems for systems of finite sets. *Quarterly Journal of Mathematics. Oxford. Second Series*, 12:313–320, 1961. 4.21
- [Erd78] Robert Erdahl. Representability. *International Journal of Quantum Chemistry*, 13(6):697–718, 1978. 1.3
- [ES14] László Erdős and Dominik Schröder. Phase transition in the density of states of quantum spin glasses. *Mathematical Physics, Analysis and Geometry. An International Journal Devoted to the Theory and Applications of Analysis and Geometry to Physics*, 17(3-4):441–464, 2014. 8.2
- [FH14] Michael Freedman and Matthew Hastings. Quantum systems on non- k -hyperfinite complexes: a generalization of classical statistical mechanics on expander graphs. *Quantum Information & Computation*, 14(1-2):144–180, 2014. 1.3
- [FTW18] Renjie Feng, Gang Tian, and Dongyi Wei. Spectrum of SYK model II: Central limit theorem. Technical Report 1806.05714, arXiv, 2018. 1.4, 3.2
- [FTW19] Renjie Feng, Gang Tian, and Dongyi Wei. Spectrum of SYK model. *Peking Mathematical Journal*, 2(1):41–70, 2019. 1.4, 3.1, 3.7, 3.2, 5.5
- [FTW20] Renjie Feng, Gang Tian, and Dongyi Wei. Spectrum of SYK model III: large deviations and concentration of measures. *Random Matrices. Theory and Applications*, 9(2):2050001, 24, 2020. 1.4, 3.2
- [FW70] James French and Samuel Wong. Validity of random matrix theories for many-particle systems. *Physics Letters B*, 33(7):449–452, 1970. 1.4
- [GH80] Humphrey Gastineau-Hills. *Systems of orthogonal designs and quasi Clifford algebras*. PhD thesis, University of Sydney, 1980. 2.2
- [GH82] Humphrey Gastineau-Hills. Quasi Clifford algebras and systems of orthogonal designs. *Journal of the Australian Mathematical Society*, 32(1):1–23, 1982. 2.2, 2.4, 2.15

- [GJRV21] Antonio García-García, Yiyang Jia, Dario Rosa, and Jacobus Verbaarschot. Sparse Sachdev–Ye–Kitaev model, quantum chaos, and gravity duals. *Physical Review D*, 103(10):106002, 2021. [8.2](#)
- [GJV18] Antonio García-García, Yiyang Jia, and Jacobus Verbaarschot. Exact moments of the Sachdev–Ye–Kitaev model up to order $1/N^2$. *Journal of High Energy Physics*, 2018(4):1–43, 2018. [1.4](#), [3.2](#)
- [GLS81] Martin Grötschel, László Lovász, and Alexander Schrijver. The ellipsoid method and its consequences in combinatorial optimization. *Combinatorica*, 1(2):169–197, 1981. [7.2](#)
- [Got97] Daniel Gottesman. *Stabilizer codes and quantum error correction*. California Institute of Technology, 1997. arXiv:quant-ph/9705052. [8.2](#)
- [GR17] David Gross and Vladimir Rosenhaus. A generalization of Sachdev–Ye–Kitaev. *Journal of High Energy Physics*, 2017(2):1–38, 2017. [8](#)
- [Gri01] Dima Grigoriev. Linear lower bound on degrees of Positivstellensatz calculus proofs for the parity. *Theoretical Computer Science*, 259(1-2):613–622, 2001. [1.1](#)
- [GV16] Antonio García-García and Jacobus Verbaarschot. Spectral and thermodynamic properties of the Sachdev–Ye–Kitaev model. *Physical Review D*, 94(12):126010, 2016. [1.4](#)
- [GW95] Michel Goemans and David Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *Journal of the ACM*, 42:1115–1145, 1995. [1.1](#), [1.3](#), [8.2](#)
- [Hås01] Johan Håstad. Some optimal inapproximability results. *Journal of the ACM*, 48(4):798–859, 2001. [1.1](#), [1.1](#)
- [Has09] Matthew Hastings. Making almost commuting matrices commute. *Communications in Mathematical Physics*, 291(2):321–345, 2009. [B](#)
- [Has19] Matthew Hastings. Duality in quantum quenches and classical approximation algorithms: pretty good or very bad. *Quantum*, 3:201, 2019. [17](#)
- [HM04] J. William Helton and Scott McCullough. A Positivstellensatz for non-commutative polynomials. *Transactions of the American Mathematical Society*, 356(9):3721–3737, 2004. [1.3](#)
- [HTS21] Arijit Halder, Omid Tavakol, and Thomas Scaffidi. Variational wave functions for Sachdev–Ye–Kitaev models. *Physical Review Research*, 3(2):023020, 2021. [9](#), [12](#), [6.7](#), [7.1](#), [7.12](#)
- [ISV87] Mourad Ismail, Dennis Stanton, and Gérard Viennot. The combinatorics of q -Hermite polynomials and the Askey-Wilson integral. *European Journal of Combinatorics*, 8(4):379–392, 1987. [3.2](#)
- [Jan97] Svante Janson. *Gaussian Hilbert Spaces*. Cambridge University Press, 1997. [8.1](#)
- [JH16] Cédric Josz and Didier Henrion. Strong duality in Lasserre’s hierarchy for polynomial optimization. *Optimization Letters*, 10(1):3–10, 2016. [2.6](#)

- [Kho08] Tanya Khovanova. Clifford algebras and graphs. Technical Report 0810.3322, arXiv, 2008. 2.4
- [Kit15] Alexei Kitaev. A simple model of quantum holography. In *KITP Strings Seminar and Entanglement*, volume 12, page 26, 2015. <https://online.kitp.ucsb.edu/online/entangled15/kitaev/>. 1.4
- [KKR06] Julia Kempe, Alexei Kitaev, and Oded Regev. The complexity of the local Hamiltonian problem. *SIAM Journal on Computing*, 35(5):1070–1097, 2006. 1.3
- [Kly06] Alexander Klyachko. Quantum marginal problem and N -representability. 36(1):72–86, 2006. 1.3
- [KMOW17] Pravesh Kothari, Ryuhei Mori, Ryan O’Donnell, and David Witmer. Sum of squares lower bounds for refuting any CSP. In *Proceedings of the 49th annual ACM Symposium on Theory of Computing (STOC)*, pages 132–145, 2017. 1.1
- [KMPT18] Igor Klebanov, Alexey Milekhin, Fedor Popov, and Grigory Tarnopolsky. Spectra of eigenstates in fermionic tensor quantum mechanics. *Physical Review D*, 97(10):106023, 2018. 8.2
- [Lat06] Rafał Łatała. Estimates of moments and tails of Gaussian chaoses. *The Annals of Probability*, 34(6):2315–2331, 2006. 3.1, 3.10
- [LCV07] Yi-Kai Liu, Matthias Christandl, and Frank Verstraete. Quantum computational complexity of the N -representability problem: QMA complete. *Physical review letters*, 98(11):110503, 2007. 1.3
- [Led01] Michel Ledoux. *The Concentration of Measure Phenomenon*. Mathematical Surveys and Monographs. American Mathematical Society, 2001. 3
- [Leh11] Joseph Lehec. Moments of the Gaussian chaos. In *Séminaire de Probabilités XLIII*, volume 2006 of *Lecture Notes in Mathematics*, pages 327–340. Springer, Berlin, 2011. 3.1
- [LM00] Béatrice Laurent and Pascal Massart. Adaptive estimation of a quadratic functional by model selection. *The Annals of Statistics*, 28(5):1302–1338, 2000. 8.1
- [Lov79] László Lovász. On the Shannon capacity of a graph. *IEEE Transactions on Information Theory*, 25(1):1–7, 1979. 4, 4.9
- [LP86] Françoise Lust-Piquard. Inégalités de Khintchine dans C_p ($1 < p < \infty$). *Comptes Rendus des Séances de l’Académie des Sciences. Série I. Mathématique*, 303(7):289–292, 1986. 7.2
- [LPP91] Françoise Lust-Piquard and Gilles Pisier. Noncommutative Khintchine and Paley inequalities. *Arkiv för Matematik*, 29(2):241–260, 1991. 7.2
- [Maz12] David Mazziotti. Structure of fermionic density matrices: Complete N -representability conditions. *Physical Review Letters*, 108(26):263002, 2012. 1.3, 1.2
- [ME01] David Mazziotti and Robert Erdahl. Uncertainty relations and reduced density matrices: Mapping many-body quantum mechanics onto four particles. *Physical Review A*, 63(4):042113, 2001. 1.3

- [Meg01] Alexandre Megretski. Relaxations of quadratic programs in operator theory and system analysis. In *Systems, approximation, singular integral operators, and related topics (Bordeaux, 2000)*, volume 129 of *International Workshop on Operator Theory and Applications*, pages 365–392. Birkhäuser, Basel, 2001. 7.2
- [Mon21] Andrea Montanari. Optimization of the sherrington–kirkpatrick hamiltonian. *SIAM Journal on Computing*, (0):FOCS19–1, 2021. 1.1
- [MR10] Dana Moshkovitz and Ran Raz. Two-query PCP with subconstant error. *Journal of the ACM*, 57(5):29, 2010. 1.1
- [Nes98] Yurii Nesterov. Global quadratic optimization via conic relaxation. Technical Report 9860, CORE, Catholic University of Louvain, Discussion Paper, 1998. 7.2
- [NPA08] Miguel Navascués, Stefano Pironio, and Antonio Acín. A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations. *New Journal of Physics*, 10(7):073013, 2008. 1.3
- [NRT99] Arkadi Nemirovski, Cornelis Roos, and Tamás Terlaky. On maximization of quadratic form over intersection of ellipsoids with common center. *Mathematical Programming*, 86(3, Series A):463–473, 1999. 7.2
- [O’D14] Ryan O’Donnell. *Analysis of Boolean functions*. Cambridge University Press, 2014. 7.2
- [Oli10] Roberto Imbuzeiro Oliveira. Sums of random Hermitian matrices and an inequality by Rudelson. *Electronic Communications in Probability*, 15:203–212, 2010. 3.1
- [Per78] Jerome Percus. The role of model systems in the few-body reduction of the N -fermion problem. *International Journal of Quantum Chemistry*, 13(1):89–124, 1978. 1.3
- [PNA10] Stefano Pironio, Miguel Navascués, and Antonio Acín. Convergent relaxations of polynomial optimization problems with noncommuting variables. *SIAM Journal on Optimization*, 20(5):2157–2180, 2010. 1.3
- [Rag09] Prasad Raghavendra. *Approximating NP-hard problems: efficient algorithms and their limits*. PhD thesis, University of Washington, 2009. 1.1
- [Ros02] Wulf Rossmann. *Lie Groups: An introduction through linear groups*. Oxford University Press, 2002. 8.1
- [Ros19] Vladimir Rosenhaus. An introduction to the SYK model. *Journal of Physics A: Mathematical and Theoretical*, 52(32):323001, 2019. 1.4
- [RS09] Prasad Raghavendra and David Steurer. How to round any csp. In *Proceedings of the 50th Annual IEEE Symposium on Foundations of Computer Science (FOCS)*, pages 586–594, 2009. 1.1
- [Sch81] Alexander Schrijver. Association schemes and the Shannon capacity: Eberlein polynomials and the Erdős–Ko–Rado theorem. In *Algebraic methods in graph theory, Vol. I, II (Szeged, 1978)*, volume 25 of *Colloq. Math. Soc. János Bolyai*, pages 671–688. North-Holland, Amsterdam-New York, 1981. 4.2

- [So09] Anthony So. Improved approximation bound for quadratic optimization problems with orthogonality constraints. In *Proceedings of the 20th annual Symposium on Discrete Algorithms (SODA)*, pages 1201–1209. SIAM, Philadelphia, PA, 2009. [7.9](#), [7.2](#), [7.2](#)
- [Spe92] Roland Speicher. A noncommutative central limit theorem. *Mathematische Zeitschrift*, 209(1):55–66, 1992. [11](#)
- [SY93] Subir Sachdev and Jinwu Ye. Gapless spin-fluid ground state in a random quantum Heisenberg magnet. *Physical Review Letters*, 70(21):3339–342, 1993. [1.4](#)
- [TD02] Barbara Terhal and David DiVincenzo. Classical simulation of noninteracting-fermion quantum circuits. *Physical Review A*, 65(3):032325, 2002. [1.3](#), [7](#)
- [Tro12] Joel Tropp. User-friendly tail bounds for sums of random matrices. *Foundations of Computational Mathematics*, 12(4):389–434, 2012. [3.1](#), [7.2](#)
- [Wil84] Richard Wilson. The exact bound in the Erdős-Ko-Rado theorem. *Combinatorica*, 4(2-3):247–257, 1984. [4.2](#)
- [XSSSS20] Shenglong Xu, Leonard Susskind, Yuan Su, and Brian Swingle. A sparse model of quantum holography. Technical Report 2008.02303, arXiv, 2020. [8.2](#)

A Derivatives of $\text{Opt}(\ell)$ for ℓ supported on an independent set

In [Section 4](#), we have shown the bound

$$\alpha(\Gamma) \leq \Psi(\Gamma) \leq \vartheta(\Gamma)$$

for the Ψ function.

Here we further consider the question of whether there is some graph Γ such that $\Psi(\Gamma) > \alpha(\Gamma)$. As partial evidence that these two functions might coincide, we prove the following.

Lemma A.1. *Let S be a maximal independent set in Γ . Regard $\text{Opt}(\sum_j a_j \chi_j)$ as a function of the vector \vec{a} on the sphere $|\vec{a}| = 1$. Let \vec{a}_0 be the vector with $(\vec{a}_0)_j = 0$ for $j \notin S$ and $(\vec{a}_0)_j = \frac{1}{\sqrt{|S|}} \lambda_j$ for $j \in S$, where the signs $\lambda_j \in \{\pm 1\}$ are chosen so that some common eigenvector of the χ_j has associated eigenvalue $\lambda_j \in \{\pm 1\}$. Hence, $\ell_0 = \sum_j (\vec{a}_0)_j \chi_j$ is the ℓ considered in [Proposition 4.7](#).*

Then, the first derivative of this function with respect to \vec{a} vanishes at $\vec{a} = \vec{a}_0$, and the Hessian matrix (i.e., matrix of second derivatives w.r.t. some coordinates on the sphere $|J| = 1$) is negative semidefinite.

Remark: we say that the matrix is negative semidefinite rather than that we are at a local optimum as the matrix may have a zero eigenvalue. Indeed, consider the case of two operators, X, Z which anticommute. Then $\ell = \cos(\theta)X + \sin(\theta)Z$ has largest eigenvalue equal to 1 for all θ .

Proof. A convenient set of coordinates on the sphere in a neighborhood of a given $\vec{a} = \vec{a}_0$ is to consider vectors $d\vec{a}$ normal to \vec{a}_0 and let $\vec{a} = (\vec{a}_0 + d\vec{a})/|\vec{a}_0 + d\vec{a}|$. Let

$$E = \text{Opt}\left(\sum_{\alpha} ((\vec{a}_0)_j + d\vec{a}_j) O_j\right).$$

Note that the argument of the optimum in E is not normalized to have ℓ_2 norm equal to 1. We have

$$\text{Opt}\left(\frac{\vec{a}_0 + d\vec{a}}{|\vec{a}_0 + d\vec{a}|}\right) = \frac{E}{|\vec{a}_0 + d\vec{a}|}.$$

The vanishing of the first derivative is immediate: operators χ_k for $k \notin S$ anticommute with at least one operator χ_j for some $j \in S$ and so the expectation value of χ_k vanishes in any eigenstate of ℓ_0 of maximal eigenvalue.

After some calculus we find to second order, for $d\vec{a}$ normal to \vec{a}_0 , that

$$\frac{E}{|\vec{a}_0 + d\vec{a}|} = \sqrt{|S|} + \sum_{j,k} M_{j,k} d\vec{a}_j d\vec{a}_k - \sum_{j,k} \frac{E}{2} \delta_{j,k} d\vec{a}_j d\vec{a}_k$$

with

$$M_{j,k} = \frac{1}{2} \partial_j \partial_k E,$$

where we write ∂_j for brevity for $\partial_{d\vec{a}_j}$.

So, we must show that the largest eigenvalue of M in the given subspace (i.e., for vectors normal to \vec{a}_0) is bounded by $E/2 = \sqrt{|S|}/2$.

We have, for $j, k \notin S$ that

$$\frac{1}{2} \partial_j \partial_k E = \langle \phi, \chi_j (E - \ell_0)^{-1} \chi_k \phi \rangle, \quad (47)$$

where ϕ is an eigenstate ℓ_0 with maximal eigenvalue. This follows from standard second-order perturbation theory results.

We claim that M is block-diagonal. Each block corresponds to a choice of $T \subset S$ and k is in a given block iff χ_k anticommutes with χ_m for all $m \in T$ and commutes with χ_m for all $m \in S \setminus T$. Indeed, the state $\chi_k \phi$ is an eigenstate of χ_m for each $m \in S$ with eigenvalue ± 1 (the sign is -1 if $m \in T$ and $+1$ otherwise), and $E - \ell_0$ commutes with all such χ_l , so $\partial_j \partial_k E$ vanishes between blocks.

Consider a given block for given T . The operator $E - \ell_0$ equals

$$\frac{2|T|}{\sqrt{|S|}}$$

in that block. Hence, the largest eigenvalue of M in that block is bounded by $\frac{\sqrt{|S|}}{2|T|}$ times the cardinality of that block. Hence it is less than $\sqrt{|S|}/2$ unless the cardinality of that block is larger than $|T|$. However, in that case, the independent set was not maximal: we may remove all elements of T from S and add all elements of the given block to increase the independent set. \square

B Low-rank and Gaussian states

Suppose

$$p = \sum_{\alpha=1}^k \lambda_{\alpha} Q_{\alpha}^2, \quad (48)$$

where α is some index, k is an integer called the *rank*, λ_{α} is a real scalar and Q_{α} is a self-adjoint quadratic operator.

Let $Q_{\alpha} = \sum_{j,k} \chi_j (K_{\alpha})_{jk} \chi_k$, where K_{jk} is a self-adjoint anti-symmetric matrix (hence, K is pure imaginary). Let us fix $|K_{\alpha}|_2 = 1$. Hence, $\|Q_{\alpha}\|_{\text{op}} = O(n^{1/2})$.

Here we consider the low rank case, where $k = O(1)$. Assume also for normalization that $\sum_{\alpha} |\lambda_{\alpha}| = O(1)$.

We have

$$[Q_\alpha, Q_\beta] = 4 \sum_{jk} ([K_\alpha, K_\beta])_{jk} \chi_j \chi_k.$$

The ℓ_2 norm of $[K_\alpha, K_\beta]$ is bounded (by a triangle inequality and Cauchy-Schwarz) by $2|K_\alpha|_2|K_\beta|_2 \leq 2$, though probably tighter bounds are possible. Hence, by results before, $\|[Q_\alpha, Q_\beta]\|_{\text{op}} = O(n^{1/2})$.

Similarly,

$$\begin{aligned} \|[Q_\alpha, p]\|_{\text{op}} &\leq \sum_{\beta=1}^k |\lambda_\beta| \|[Q_\beta^2, Q_\alpha]\|_{\text{op}} \\ &\leq \sum_{\beta=1}^k |\lambda_\beta| \cdot O(n). \end{aligned}$$

Consider the re-scaled operators $n^{-1/2}Q_\alpha$ and $n^{-1}p$. These operators all have operator norm $O(1)$ and the commutator of any two such operators is $O(n^{-1/2})$.

By section VII of [Has09], we can define a POVM (positive operator-valued measure) that approximately measures the K different operators $n^{-1/2}Q_\alpha$ and $n^{-1}p$ on some arbitrary quantum state ρ . In particular (dropping the re-scaling by $n^{-1/2}$ and n^{-1} from here on), the measurement returns scalars q_α such that the average over outcomes of the expectation value $(Q_\alpha - q_\alpha)^2$ in the resulting state is $o(n)$. Further, for any α , the expectation value $\text{tr}(p\rho)$ is within $o(n)$ of the average of $\sum \lambda_\alpha q_\alpha^2$ over measurement outcomes. Let ρ then be the state that maximizes $\text{tr}(p\rho)$. Apply this POVM. The expectation value of p in the resulting state (called ρ_{outcome}), averaged over outcomes, is within $o(n)$ of $\text{tr}(p\rho)$. Hence, the average over measurement outcomes of $\sum_\alpha \lambda_\alpha \text{tr}(\rho_{\text{outcome}} Q_\alpha)^2$ is within $o(n)$ of $\text{tr}(p\rho)$.

So, there is some state σ for which $\sum_\alpha \lambda_\alpha \text{tr}(\sigma Q_\alpha)^2$ is within $o(n)$ of $\text{tr}(p\rho)$. Let us optimize $\sum_\alpha \lambda_\alpha \text{tr}(\sigma Q_\alpha)^2$ over states σ . An argument with Lagrange multipliers shows that the maximum is attained on a Gaussian state. Further, on this Gaussian state, one may verify that $\text{tr}(p\sigma)$ is within $o(n)$ of $\sum_\alpha \lambda_\alpha \text{tr}(\sigma Q_\alpha)^2$ and hence within $o(n)$ of $\text{tr}(p\rho)$.