

Maximum entropy principle approach to a non-isothermal Maxwell-Stefan diffusion model

Benjamin Anwasia^{a,b}, Srboľjub Simić^{c,*}

^a*Universidade do Minho, Centro de Matemática, Campus de Gualtar, 4710-057 Braga, Portugal*

^b*New York University Abu Dhabi, Abu Dhabi, United Arab Emirates*

^c*Department of Mathematics and Informatics, Faculty of Sciences, University of Novi Sad, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia*

Abstract

In this study we apply the maximum entropy principle to derive the properly scaled velocity distribution function of Boltzmann equations for mixtures, which leads to a non-isothermal Maxwell-Stefan diffusion model. We also analyze the entropy balance law and derive the kinetic entropy production from the scaled distribution function.

Keywords: diffusion, Maxwell-Stefan model, maximum entropy principle

2010 MSC: 35Q20, 82C40, 76R50

1. Introduction

Diffusion is a process of motion of one specie relative to another. If the motion occurs relative to a background medium, the process is usually described by the Fick's law or some of its generalizations. When cross-diffusion is more prominent, like in gaseous mixtures, the Maxwell-Stefan diffusion model is more appropriate [1]. It turned out that cross-diffusion models of Maxwell-Stefan kind can be used as a starting point for diverse applications, ranging from engineering [2] to medicine [3], and they also inspired new approaches to diffusion phenomena [4].

Along with applications, mathematical analysis of Maxwell-Stefan equations emerged as equally important but relatively new subject [5, 6]. It covers the aspects of existence and uniqueness of solutions [7, 8], large-time asymptotics [9] and relaxation limits [10]. Our concern, however, will be the formal derivation of Maxwell-Stefan diffusion equations and analysis of their dissipative character.

From the macroscopic point of view, Maxwell-Stefan diffusion model consists of mass balance equations for the constituents, and a kind of momentum balance relations adjoined with a closure relation due to their linear dependence. It can be derived either in the continuum framework [11, 12], or starting from the Boltzmann equations (BEs) for mixtures [13, 14, 15, 16]. In the latter approach, crucial step in derivation of Maxwell-Stefan equations is the asymptotic limit enabled through appropriate scaling of the governing equations.

In this study we shall take the kinetic approach to Maxwell-Stefan equations. It is based upon *formal* diffusive scaling of the BEs, and crucial *assumption* that velocity distribution functions have the following form:

$$f_i^\alpha(t, \mathbf{x}, \mathbf{v}) = \rho_i^\alpha \left(\frac{m_i}{2\pi k_B T} \right)^{3/2} \exp \left(-\frac{m_i |\mathbf{v} - \alpha \mathbf{u}_i^\alpha|^2}{2k_B T} \right), \quad i = 1, \dots, S, \quad (1)$$

* Authors acknowledge financial support (B.A.) from the COST Action, CA18232 MAT-DYN-NET, supported by the "European Cooperation in Science and Technology" through the "Short Term Scientific Mission" at the Department of Mathematics and Informatics, University of Novi Sad, Serbia, and (S.S.) from the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant No. 451-03-9/2021-14/200125).

*Corresponding author

Email addresses: id6226@alunos.uminho.pt (Benjamin Anwasia), ssimic@uns.ac.rs (Srboľjub Simić)

for some small parameter $\alpha \ll 1$, where $\mathbf{v} \in \mathbb{R}^3$ are the particle velocities, $\mathbf{u}_i^\alpha \in \mathbb{R}^3$ are the macroscopic species velocities, m_i are the species atomic masses, k_B is the Boltzmann constant, ρ_i^α are the species mass densities and T is the mixture temperature.

Instead of assuming, our aim is to *derive* the scaled velocity distribution function (1) appropriate for description of diffusion processes. To that end, we shall apply the maximum entropy principle (MEP) in the scaled (dimensionless) form (Theorem 1). Moreover, we shall put our analysis in a broader context of non-isothermal processes. We shall also derive the entropy balance law in the scaled form and compute the entropy production (Theorem 2) for the non-isothermal Maxwell-Stefan model, starting from the scaled velocity distribution function. Computation of the velocity distribution function from MEP will be *exact*. Therefore, we shall derive the moment equations (Maxwell-Stefan model) and the entropy balance law for the mixture of monatomic Maxwellian particles, and in such a way retain the exact character of the results.

2. The Boltzmann equation for inert mixtures

We consider a mixture of monatomic ideal inert gases: a gaseous mixture of non-reactive species consisted of particles endowed with translational degrees of freedom only. For each species \mathcal{A}_i with $i = 1, 2, 3, \dots, S$, let $f_i := f_i(t, \mathbf{x}, \mathbf{v})$ be its distribution function, representing the densities of particles of species \mathcal{A}_i which at time $t \in \mathbb{R}^+$ are located at position $\mathbf{x} \in \mathbb{R}^3$ and have velocity $\mathbf{v} \in \mathbb{R}^3$. In the absence of external forces, the evolution of the non-reactive mixture can be described by the following system of BEs:

$$\begin{aligned} \frac{\partial f_i}{\partial t} + \mathbf{v} \cdot \frac{\partial f_i}{\partial \mathbf{x}} = & \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{S}^2} [f_i(\mathbf{v}') f_i(\mathbf{v}_*) - f_i(\mathbf{v}) f_i(\mathbf{v}_*)] B_i(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} dv_*}_{Q_i^m(f_i, f_i)} \\ & + \sum_{\substack{j=1 \\ j \neq i}}^S \underbrace{\int_{\mathbb{R}^3} \int_{\mathbb{S}^2} [f_i(\mathbf{v}') f_j(\mathbf{v}_*) - f_i(\mathbf{v}) f_j(\mathbf{v}_*)] B_{ij}(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} dv_*}_{Q_{ij}^b(f_i, f_j)}, \end{aligned} \quad (2)$$

for $i = 1, 2, 3, \dots, S$, where the first and second terms on the right-hand side describe collisions between particles of the same species and between particles of different species, respectively. They are called the mono-species and bi-species collision operators with \mathbf{v}' , \mathbf{v}_* defined as follows:

$$\mathbf{v}' = \frac{1}{m_i + m_j} (m_i \mathbf{v} + m_j \mathbf{v}_* + m_j |\mathbf{v} - \mathbf{v}_*| \boldsymbol{\sigma}), \quad \mathbf{v}_* = \frac{1}{m_i + m_j} (m_i \mathbf{v} + m_j \mathbf{v}_* - m_i |\mathbf{v} - \mathbf{v}_*| \boldsymbol{\sigma}), \quad (3)$$

where $\boldsymbol{\sigma} \in \mathbb{S}^2$ is an arbitrary unit vector. The formulas given in equation (3) represent all the possible solutions $(\mathbf{v}', \mathbf{v}_*) \in \mathbb{R}^3 \times \mathbb{R}^3$ of the system of equations composed of the momentum and kinetic energy conservation law, which are respectively defined as follows:

$$m_i \mathbf{v} + m_j \mathbf{v}_* = m_i \mathbf{v}' + m_j \mathbf{v}_*, \quad \frac{1}{2} m_i |\mathbf{v}|^2 + \frac{1}{2} m_j |\mathbf{v}_*|^2 = \frac{1}{2} m_i |\mathbf{v}'|^2 + \frac{1}{2} m_j |\mathbf{v}_*'|^2. \quad (4)$$

2.1. Scaled system of Boltzmann equations

To write the system (2) of BEs in dimensionless form, we first choose macroscopic scales for the independent variables t , \mathbf{x} and \mathbf{v} associated with the distribution function. To this end, let τ , L and T_0 respectively be the macroscopic time, space and reference temperature scales. These macroscopic scales define two macroscopic velocity scales: $u_0 = \frac{L}{\tau}$ and $c_0 = \sqrt{\frac{5k_B T_0}{3m_0}}$, where u_0 is the speed at which some macroscopic portion of the gas is transported over a distance L in time τ , while c_0 is the speed of sound in a monatomic gas with m_0 the average atomic mass of the mixture. Using the macroscopic scales: τ , L and c_0 , one can define the dimensionless time, space, velocity, distribution function and cross-sections:

$$\hat{t} = \frac{t}{\tau}, \quad \hat{\mathbf{x}} = \frac{\mathbf{x}}{L}, \quad \hat{\mathbf{v}} = \frac{\mathbf{v}}{c_0}, \quad \hat{f}_i(\hat{t}, \hat{\mathbf{x}}, \hat{\mathbf{v}}) = \frac{L^3 c_0^3}{N} f_i(t, \mathbf{x}, \mathbf{v}), \quad \hat{B}_i(\hat{\mathbf{v}}, \hat{\mathbf{v}}_*, \hat{\boldsymbol{\sigma}}) = \frac{1}{c_0 4\pi r^2} B_i(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}), \quad (5)$$

respectively, and rewrite the system (2) of BEs in dimensionless form in which only two parameters appear—Mach number Ma which multiplies $\partial \hat{f}_i / \partial \hat{t}$, and a reciprocal of the Knudsen number Kn which multiplies the scaled collision operators:

$$\text{Ma} = \frac{u_0}{c_0}, \quad \text{Kn} = \frac{\text{mean free path}}{\text{macroscopic length scale}} = \frac{L^3}{N \times 4\pi r^2} \times \frac{1}{L}.$$

In the sequel we shall assume that $\text{Ma} = \text{Kn} = \alpha \ll 1$, which amounts to diffusive scaling of the BEs (2). Removing the hats for simplicity and adding superscript α to the distribution function so as to differentiate the dimensionless system of BEs from equation (2), we obtain:

$$\alpha \frac{\partial f_i^\alpha}{\partial t} + \mathbf{v} \cdot \frac{\partial f_i^\alpha}{\partial \mathbf{x}} = \frac{1}{\alpha} Q_i^m(f_i^\alpha, f_i^\alpha) + \frac{1}{\alpha} \sum_{\substack{j=1 \\ j \neq i}}^S Q_{ij}^b(f_i^\alpha, f_j^\alpha). \quad (6)$$

2.2. Moment Equations

Multiplying the scaled system of BEs (6) by an arbitrary function of the velocity variable $\psi_i(\mathbf{v})$, and then integrate the resulting equation over all values of the velocity \mathbf{v} gives the moment equations:

$$\alpha \frac{\partial}{\partial t} \int_{\mathbb{R}^3} \psi_i(\mathbf{v}) f_i^\alpha d\mathbf{v} + \frac{\partial}{\partial \mathbf{x}} \int_{\mathbb{R}^3} \mathbf{v} \psi_i(\mathbf{v}) f_i^\alpha d\mathbf{v} = \frac{1}{\alpha} \int_{\mathbb{R}^3} \psi_i(\mathbf{v}) Q_i^m(f_i^\alpha, f_i^\alpha) d\mathbf{v} + \frac{1}{\alpha} \sum_{\substack{j=1 \\ j \neq i}}^S \int_{\mathbb{R}^3} \psi_i(\mathbf{v}) Q_{ij}^b(f_i^\alpha, f_j^\alpha) d\mathbf{v}, \quad (7)$$

where the integrals on the right-hand side are the weak formulation of the mono-species and bi-species collision operators respectively defined as follows:

$$\begin{aligned} \int_{\mathbb{R}^3} \psi_i(\mathbf{v}) Q_i^m(f_i^\alpha, f_i^\alpha) d\mathbf{v} = & -\frac{1}{4} \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} \left[f_i^\alpha(\mathbf{v}') f_i^\alpha(\mathbf{v}_*) - f_i^\alpha(\mathbf{v}) f_i^\alpha(\mathbf{v}_*) \right] \\ & \times [\psi_i(\mathbf{v}') + \psi_i(\mathbf{v}_*) - \psi_i(\mathbf{v}) - \psi_i(\mathbf{v}_*)] B_i(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} d\mathbf{v}_* d\mathbf{v}, \end{aligned} \quad (8)$$

$$\int_{\mathbb{R}^3} \psi_i(\mathbf{v}) Q_{ij}^b(f_i^\alpha, f_j^\alpha) d\mathbf{v} = \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \int_{\mathbb{S}^2} f_i^\alpha(\mathbf{v}) f_j^\alpha(\mathbf{v}_*) [\psi_i(\mathbf{v}') - \psi_i(\mathbf{v})] B_{ij}(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} d\mathbf{v}_* d\mathbf{v}. \quad (9)$$

3. Maximum entropy principle

The maximum entropy principle (MEP) determines the approximate velocity distribution function, compatible with selected moments, through maximization of the kinetic entropy [17, 18, 19, 20, 21]. Our aim is to apply MEP and *derive* the properly scaled velocity distribution function which leads to a non-isothermal Maxwell-Stefan model. To that end we introduce the following

Definition 1. Kinetic entropy $H(t, \mathbf{x})$ is defined as:

$$H(t, \mathbf{x}) := \sum_{i=1}^S H_i(t, \mathbf{x}), \quad H_i(t, \mathbf{x}) := -k_B \int_{\mathbb{R}^3} f_i \log(b_i f_i) d\mathbf{v}, \quad (10)$$

where b_i is the dimensional constant which makes the argument of the log function dimensionless. Moments of the distribution functions, representing partial mass, momentum and energy densities, are defined as:

$$\begin{aligned} \rho_i(t, \mathbf{x}) &:= \int_{\mathbb{R}^3} m_i f_i d\mathbf{v}, \quad \rho_i(t, \mathbf{x}) \mathbf{u}_i(t, \mathbf{x}) := \int_{\mathbb{R}^3} m_i \mathbf{v} f_i d\mathbf{v}, \\ \rho_i(t, \mathbf{x}) |\mathbf{u}_i(t, \mathbf{x})|^2 + 3\rho_i(t, \mathbf{x}) \frac{k_B}{m_i} T(t, \mathbf{x}) &:= \int_{\mathbb{R}^3} m_i |\mathbf{v}|^2 f_i d\mathbf{v}, \end{aligned} \quad (11)$$

where ρ_i is the mass density and \mathbf{u}_i is the mean velocity of the specie \mathcal{A}_i , while T is the common kinetic temperature.

It will be assumed that all the species have the same kinetic temperature T . This simplifying assumption is appropriate for non-isothermal diffusion problems, see [22].

In the sequel we shall use the dimensionless macroscopic variables:

$$\hat{\rho}_i = \frac{L^3}{m_i N} \rho_i, \quad \hat{\mathbf{u}}_i = \frac{\mathbf{u}_i}{u_0}, \quad \hat{T} = \frac{T}{T_0}, \quad \hat{H}_i = \frac{L^3}{k_B N} H_i, \quad \hat{b}_i = \frac{N}{L^3 c_0^3} b_i. \quad (12)$$

and drop the hat notation for convenience. The variational problem (MEP) may now be formulated in dimensionless form as follows.

Maximum entropy principle. Find the velocity distribution functions $f_i^\alpha(t, \mathbf{x}, \mathbf{v})$ which maximize the kinetic entropy:

$$H^\alpha(t, \mathbf{x}) = - \int_{\mathbb{R}^3} \sum_{i=1}^n f_i^\alpha \log(b_i f_i^\alpha) d\mathbf{v} \rightarrow \max, \quad (13)$$

subject to the following constraints:

$$\begin{aligned} \rho_i^\alpha(t, \mathbf{x}) &= \int_{\mathbb{R}^3} f_i^\alpha d\mathbf{v}, \quad \alpha \rho_i^\alpha(t, \mathbf{x}) \mathbf{u}_i^\alpha(t, \mathbf{x}) = \int_{\mathbb{R}^3} \mathbf{v} f_i^\alpha d\mathbf{v}, \\ \alpha^2 \rho_i^\alpha(t, \mathbf{x}) |\mathbf{u}_i^\alpha(t, \mathbf{x})|^2 + 3 \rho_i^\alpha(t, \mathbf{x}) \frac{3}{5} \frac{m_0}{m_i} T(t, \mathbf{x}) &= \int_{\mathbb{R}^3} |\mathbf{v}|^2 f_i^\alpha d\mathbf{v}. \end{aligned} \quad (14)$$

We would like to note that small parameter α naturally appears in dimensionless constraints (14) due to scaling introduced in previous Section.

Theorem 1. *The velocity distribution functions which solve the variational problem (13)-(14) have the following form:*

$$f_i^\alpha(t, \mathbf{x}, \mathbf{v}) = \left(\frac{5}{3m_0} \right)^{3/2} \rho_i^\alpha(t, \mathbf{x}) \left(\frac{m_i}{2\pi T(t, \mathbf{x})} \right)^{3/2} \exp \left(- \frac{5}{3m_0} \frac{m_i |\mathbf{v} - \alpha \mathbf{u}_i^\alpha(t, \mathbf{x})|^2}{2T(t, \mathbf{x})} \right). \quad (15)$$

PROOF. Proof of the Theorem 1 relies on the method of multipliers. Starting from (13)-(14) we define extended functional:

$$\mathcal{H}^\alpha(t, \mathbf{x}) := \int_{\mathbb{R}^3} \sum_{i=1}^n \left(f_i^\alpha \log(b_i f_i^\alpha) + \lambda_i^{(0)} f_i^\alpha + \boldsymbol{\lambda}_i^{(1)} \cdot \mathbf{v} f_i^\alpha + \lambda_i^{(2)} |\mathbf{v}|^2 f_i^\alpha \right) d\mathbf{v} \equiv \int_{\mathbb{R}^3} \mathcal{L}(\mathbf{v}, f_i^\alpha, \nabla_{\mathbf{v}} f_i^\alpha) d\mathbf{v},$$

where $\lambda_i^{(0)}(t, \mathbf{x}) \in \mathbb{R}$, $\boldsymbol{\lambda}_i^{(1)}(t, \mathbf{x}) \in \mathbb{R}^3$ and $\lambda_i^{(2)}(t, \mathbf{x}) \in \mathbb{R}$ are the unknown multipliers. Since the density function \mathcal{L} does not depend on $\nabla_{\mathbf{v}} f_i^\alpha$, necessary condition for extremum is reduced to $\partial \mathcal{L} / \partial f_i^\alpha = 0$, $i = 1, \dots, S$, which yields S uncoupled equations that can be solved for f_i^α :

$$f_i^\alpha = \frac{1}{b_i} \exp \left[- \left(1 + \lambda_i^{(0)} + \boldsymbol{\lambda}_i^{(1)} \cdot \mathbf{v} + \lambda_i^{(2)} |\mathbf{v}|^2 \right) \right].$$

Plugging the last expression into the constraints (14), after rather straightforward computation the following relations which determine the multipliers are obtained:

$$\lambda_i^{(2)} = \frac{5}{3} \frac{m_i}{m_0} \frac{1}{2T}, \quad \frac{\lambda_i^{(1)}}{2\lambda_i^{(2)}} = -\alpha \mathbf{u}_i^\alpha, \quad \frac{1}{b_i} \exp \left(-1 - \lambda_i^{(0)} \right) \exp \left(\frac{|\boldsymbol{\lambda}_i^{(1)}|^2}{4\lambda_i^{(2)}} \right) \left(\frac{\pi}{\lambda_i^{(2)}} \right)^{3/2} = \rho_i^\alpha.$$

The proof is completed by returning these expressions into the maximizer.

4. Non-isothermal Maxwell-Stefan diffusion equations

The Maxwell-Stefan diffusion equations for the mixture of Maxwell molecules were derived in [13], and in the non-isothermal setting in [22], while in [16] the case of reactive mixture of polyatomic gases was considered. Since our interest is on inert mixtures, we shall give the non-isothermal Maxwell-Stefan diffusion equations here for the completeness. They can be obtained from the moment equations (7) by using the scaled distribution function (15) obtained by MEP. The mass/concentration and momentum balance laws for the species are obtained taking $\psi_i(\mathbf{v}) = 1$ and $\psi_i(\mathbf{v}) = \mathbf{v}$, respectively. The energy conservation law for the mixture is obtained taking $\psi_i(\mathbf{v}) = |\mathbf{v}|^2/2$ and summing up all the energy balance laws for the species. The resulting equations read:

$$\begin{aligned} \frac{\partial \rho_i^\alpha}{\partial t} + \frac{\partial}{\partial \mathbf{x}}(\rho_i^\alpha \mathbf{u}_i^\alpha) &= 0, \\ \alpha^2 \left[\frac{\partial}{\partial t}(\rho_i^\alpha \mathbf{u}_i^\alpha) + \frac{\partial}{\partial \mathbf{x}}(\rho_i^\alpha \mathbf{u}_i^\alpha \otimes \mathbf{u}_i^\alpha) \right] + \frac{3m_0}{5m_i} \frac{\partial}{\partial \mathbf{x}}(\rho_i^\alpha T) &= 2\pi \sum_{\substack{j=1 \\ j \neq i}}^S \|b_{ij}\|_{L^1} \frac{m_j}{m_i + m_j} \rho_i^\alpha \rho_j^\alpha (\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha) \\ \frac{\partial}{\partial t} \sum_{i=1}^S \left(\alpha^2 \frac{1}{2} \rho_i^\alpha |\mathbf{u}_i^\alpha|^2 + \frac{3m_0}{5m_i} \frac{3}{2} \rho_i^\alpha T \right) + \frac{\partial}{\partial \mathbf{x}} \sum_{i=1}^S \left[\left(\alpha^2 \frac{1}{2} \rho_i^\alpha |\mathbf{u}_i^\alpha|^2 + \frac{3m_0}{5m_i} \frac{5}{2} \rho_i^\alpha T \right) \mathbf{u}_i^\alpha \right] &= 0. \end{aligned} \quad (16)$$

Taking the limit as $\alpha \rightarrow 0^+$, one obtains the non-isothermal Maxwell-Stefan diffusion equations:

$$\begin{aligned} \frac{\partial \rho_i}{\partial t} + \frac{\partial}{\partial \mathbf{x}}(\rho_i \mathbf{u}_i) &= 0, \\ \frac{3m_0}{5m_i} \frac{\partial}{\partial \mathbf{x}}(\rho_i T) &= 2\pi \sum_{\substack{j=1 \\ j \neq i}}^S \|b_{ij}\|_{L^1} \frac{m_j}{m_i + m_j} \rho_i \rho_j (\mathbf{u}_j - \mathbf{u}_i) \\ \frac{3}{2} \frac{\partial}{\partial t} \sum_{i=1}^S (\rho_i T) + \frac{5}{2} \frac{\partial}{\partial \mathbf{x}} \sum_{i=1}^S (\rho_i \mathbf{u}_i T) &= 0, \end{aligned} \quad (17)$$

where

$$\rho_i = \lim_{\alpha \rightarrow 0^+} \rho_i^\alpha \quad \text{and} \quad \mathbf{u}_i = \lim_{\alpha \rightarrow 0^+} \mathbf{u}_i^\alpha.$$

Equations (16) and (17) are up to scaling completely equivalent to the equations derived in [13, 22].

5. Entropy balance law

To derive the macroscopic entropy balance law from the BEs (2), we first have to define the entropy flux and the entropy production.

Definition 2. Entropy flux $\Phi(t, \mathbf{x})$ and entropy production $D(t, \mathbf{x})$ are defined as:

$$\begin{aligned} \Phi(t, \mathbf{x}) &:= \sum_{i=1}^S \Phi_i(t, \mathbf{x}), \quad \Phi_i(t, \mathbf{x}) := -k_B \int_{\mathbb{R}^3} \mathbf{v} f_i \log(b_i f_i) d\mathbf{v}, \\ D(t, \mathbf{x}) &:= \sum_{i=1}^S D_i(t, \mathbf{x}), \quad D_i(t, \mathbf{x}) := -k_B \int_{\mathbb{R}^3} Q_i^m(f_i, f_i) \log(b_i f_i) d\mathbf{v} - k_B \sum_{\substack{j=1 \\ j \neq i}}^S \int_{\mathbb{R}^3} Q_{ij}^b(f_i, f_j) \log(b_i f_i) d\mathbf{v}. \end{aligned}$$

By introducing dimensionless quantities:

$$\hat{\Phi}_i = \frac{L^3}{k_B N c_0} \Phi_i, \quad \hat{D}_i = \frac{L^6}{k_B N^2 4\pi r^2 c_0} D_i,$$

and dropping hat notation for convenience, one may determine the dimensionless partial entropy fluxes and partial entropy productions:

$$\begin{aligned}\Phi_i^\alpha(t, \mathbf{x}) &= - \int_{\mathbb{R}^3} \mathbf{v} f_i^\alpha \log(b_i f_i^\alpha) d\mathbf{v}, \\ D_i^\alpha(t, \mathbf{x}) &= - \int_{\mathbb{R}^3} Q_i^m(f_i^\alpha, f_i^\alpha) \log(b_i f_i^\alpha) d\mathbf{v} - \sum_{\substack{j=1 \\ j \neq i}}^S \int_{\mathbb{R}^3} Q_{ij}^b(f_i^\alpha, f_j^\alpha) \log(b_i f_i^\alpha) d\mathbf{v}.\end{aligned}\quad (18)$$

Setting $\psi_i(\mathbf{v}) = -\log(b_i f_i^\alpha)$ in (7), one obtains the following scaled entropy balance law:

$$\alpha \frac{\partial H^\alpha}{\partial t} + \frac{\partial \Phi^\alpha}{\partial \mathbf{x}} = \frac{1}{\alpha} D^\alpha. \quad (19)$$

It is a matter of straightforward computation to show that dimensionless entropy densities and fluxes read:

$$H_i^\alpha(t, \mathbf{x}) = \left\{ \log \left[b_i \left(\frac{5}{3m_0} \right)^{\frac{3}{2}} \rho_i^\alpha \left(\frac{m_i}{2\pi T} \right)^{\frac{3}{2}} \right] - \frac{3}{2} \right\} \rho_i^\alpha, \quad \Phi_i^\alpha(t, \mathbf{x}) = \alpha H_i^\alpha(t, \mathbf{x}) \mathbf{u}_i^\alpha \quad (20)$$

However, computation of the entropy production is summarized in the following Theorem.

Theorem 2. *The dimensionless partial entropy productions read:*

$$D_i^\alpha = \alpha^2 \frac{10\pi}{3Tm_0} \sum_{\substack{j=1 \\ j \neq i}}^S \|b_{ij}\|_{L^1} \frac{m_i m_j \rho_i^\alpha \rho_j^\alpha}{m_i + m_j} \left[\frac{m_i \mathbf{u}_i^\alpha + m_j \mathbf{u}_j^\alpha}{m_i + m_j} \cdot (\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha) - \mathbf{u}_i^\alpha \cdot (\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha) \right], \quad (21)$$

while the dimensionless total entropy production have the form:

$$D^\alpha = \sum_{i=1}^S D_i^\alpha = \alpha^2 \frac{5\pi}{3Tm_0} \sum_{i=1}^S \sum_{\substack{j=1 \\ j \neq i}}^S \left[\|b_{ij}\|_{L^1} \frac{m_i m_j}{m_i + m_j} \rho_i^\alpha \rho_j^\alpha |\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha|^2 \right]. \quad (22)$$

PROOF. To obtain the partial entropy, we need to evaluate the integrals given in equation (18) and we will focus on the case Maxwell molecules. This means that the collision cross-sections depend on the $\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}$ through the deviation angle $\theta \in [0, \pi]$ between $\mathbf{v} - \mathbf{v}_*$ and $\boldsymbol{\sigma}$ only. Specifically,

$$B_{ij}(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) = b_{ij} \left(\frac{\mathbf{v} - \mathbf{v}_*}{|\mathbf{v} - \mathbf{v}_*|} \cdot \boldsymbol{\sigma} \right) = b_{ij}(\cos \theta).$$

Moreover, we assume that b_{ij} is an even function and is such that $b_{ij} \in L^1(-1, 1)$ following Grad's angular cut-off assumption. Therefore,

$$\int_{\mathbb{S}^2} B_{ij}(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} = 2\pi \|b_{ij}\|_{L^1} \quad \text{and} \quad \int_{\mathbb{S}^2} \boldsymbol{\sigma} B_{ij}(\mathbf{v}, \mathbf{v}_*, \boldsymbol{\sigma}) d\boldsymbol{\sigma} = 0, \quad (23)$$

see [13]. Furthermore, using the definition of the species distribution functions (15) in $\log(b_i f_i^\alpha)$, one can rewrite equation (18) as follows:

$$\begin{aligned}D_i^\alpha &= \frac{5m_i}{6Tm_0} k_B \underbrace{\int_{\mathbb{R}^3} |\mathbf{v}|^2 Q_i^m(f_i^\alpha, f_i^\alpha) d\mathbf{v}}_A - \alpha \frac{10m_i k_B}{6Tm_0} \mathbf{u}_i^\alpha \cdot \underbrace{\int_{\mathbb{R}^3} \mathbf{v} Q_i^m(f_i^\alpha, f_i^\alpha) d\mathbf{v}}_B \\ &+ \frac{5m_i}{6Tm_0} k_B \sum_{\substack{j=1 \\ j \neq i}}^S \underbrace{\int_{\mathbb{R}^3} |\mathbf{v}|^2 Q_{ij}^b(f_i^\alpha, f_j^\alpha) d\mathbf{v}}_E - \alpha \frac{10m_i k_B}{6Tm_0} \mathbf{u}_i^\alpha \cdot \sum_{\substack{j=1 \\ j \neq i}}^S \underbrace{\int_{\mathbb{R}^3} \mathbf{v} Q_{ij}^b(f_i^\alpha, f_j^\alpha) d\mathbf{v}}_F\end{aligned}\quad (24)$$

$$+k_B \left\{ \frac{\alpha^2 5 m_i}{6 T m_0} |\mathbf{u}_i^\alpha|^2 - \log \left[b_i \left(\frac{5}{3 m_0} \right)^{\frac{3}{2}} \rho_i^\alpha \left(\frac{m_i}{2 \pi T} \right)^{\frac{3}{2}} \right] \right\} \left(\underbrace{\sum_{j=1, j \neq i}^S \int_{\mathbb{R}^3} Q_{ij}^b(f_i^\alpha, f_j^\alpha) d\mathbf{v}}_{\mathcal{G}} + \underbrace{\int_{\mathbb{R}^3} Q_i^m(f_i^\alpha, f_i^\alpha) d\mathbf{v}}_{\mathcal{H}} \right).$$

The integrals \mathcal{A} , \mathcal{B} , \mathcal{G} and \mathcal{H} all vanish, that is

$$\mathcal{A} = 0 \quad \mathcal{B} = 0, \quad \mathcal{G} = 0 \quad \text{and} \quad \mathcal{H} = 0. \quad (25)$$

To show that the integrals \mathcal{H} and \mathcal{G} both vanish, we use the weak formulation of the mono-species and bi-species collision operators given in equations (8) and (9), respectively, with $\psi(\mathbf{v}) = 1$. To show that the integrals \mathcal{A} and \mathcal{B} both vanish, we use the weak formulation of the mono-species collision operator (8) with $\psi(\mathbf{v}) = |\mathbf{v}|^2$ and $\psi(\mathbf{v}) = \mathbf{v}$, respectively, together with the conservation laws of kinetic energy and linear momentum for mono-species collisions: equations (4)₂ and (4)₁ with m_j replaced by m_i . Furthermore, using the weak form of the bi-species collision operator (9) with $\psi(\mathbf{v}) = |\mathbf{v}|^2$ and $\psi(\mathbf{v}) = \mathbf{v}$, respectively, together with equations (3) and (23), one obtains that the values of the integrals \mathcal{E} and \mathcal{F} are as given below:

$$\mathcal{E} = \alpha^2 4\pi \|b_{ij}\|_{L^1} \frac{m_j \rho_i^\alpha \rho_j^\alpha}{(m_i + m_j)^2} (m_i \mathbf{u}_i^\alpha + m_j \mathbf{u}_j^\alpha) \cdot (\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha), \quad \mathcal{F} = \alpha \frac{m_j}{m_i + m_j} 2\pi \|b_{ij}\|_{L^1} \rho_i^\alpha \rho_j^\alpha (\mathbf{u}_j^\alpha - \mathbf{u}_i^\alpha). \quad (26)$$

Substituting equations (25) and (26) into (24) gives (21). Summing equation (21) over all species gives (22).

References

- [1] R. Krishna, J. Wesselingh, The maxwell-stefan approach to mass transfer, *Chemical Engineering Science* 52 (6) (1997) 861–911.
- [2] R. Krishna, J. M. van Baten, Describing diffusion in fluid mixtures at elevated pressures by combining the maxwell-stefan formulation with an equation of state, *Chemical Engineering Science* 153 (2016) 174–187.
- [3] L. Boudin, D. Götz, B. Grec, Diffusion models of multicomponent mixtures in the lung, in: *ESAIM: Proceedings*, Vol. 30, EDP Sciences, 2010, pp. 90–103.
- [4] M. Hirschler, W. Säckel, U. Niekén, On maxwell-stefan diffusion in smoothed particle hydrodynamics, *International Journal of Heat and Mass Transfer* 103 (2016) 548–554.
- [5] D. Bothe, On the maxwell-stefan approach to multicomponent diffusion, in: *Parabolic problems*, Springer, 2011, pp. 81–93.
- [6] M. Herberg, M. Meyries, J. Prüss, M. Wilke, Reaction–diffusion systems of maxwell-stefan type with reversible mass-action kinetics, *Nonlinear Analysis* 159 (2017) 264–284.
- [7] X. Chen, A. Jüngel, Analysis of an incompressible navier–stokes–maxwell-stefan system, *Communications in Mathematical Physics* 340 (2) (2015) 471–497.
- [8] H. Hutridurga, F. Salvarani, Existence and uniqueness analysis of a non-isothermal cross-diffusion system of maxwell-stefan type, *Applied Mathematics Letters* 75 (2018) 108–113.
- [9] E. S. Daus, A. Jüngel, B. Q. Tang, Exponential time decay of solutions to reaction-cross-diffusion systems of maxwell-stefan type, *Archive for rational mechanics and analysis* 235 (2) (2020) 1059–1104.
- [10] F. Salvarani, A. J. Soares, On the relaxation of the maxwell-stefan system to linear diffusion, *Applied Mathematics Letters* 85 (2018) 15–21.
- [11] V. Giovangigli, *Multicomponent Flow Modeling*, Birkhäuser Boston Inc., Boston, MA, 1999.
- [12] D. Bothe, W. Dreyer, Continuum thermodynamics of chemically reacting fluid mixtures, *Acta Mechanica* 226 (6) (2015) 1757–1805.
- [13] L. Boudin, B. Grec, F. Salvarani, The maxwell-stefan diffusion limit for a kinetic model of mixtures, *Acta Applicandae Mathematicae* 136 (1) (2015) 79–90.
- [14] L. Boudin, B. Grec, V. Pavan, The maxwell-stefan diffusion limit for a kinetic model of mixtures with general cross sections, *Nonlinear Analysis* 159 (2017) 40–61.
- [15] B. Anwasia, P. Gonçalves, A. J. Soares, On the formal derivation of the reactive maxwell-stefan equations from the kinetic theory, *EPL (Europhysics Letters)* 129 (4) (2020) 40005.
- [16] B. Anwasia, M. Bisi, F. Salvarani, A. J. Soares, On the maxwell-stefan diffusion limit for a reactive mixture of polyatomic gases in non-isothermal setting, *Kinetic and Related Models* 13 (1) (2020) 63–95.
- [17] M. Kogan, *Rarefied Gas Dynamics*, Plenum Press, New York, NY, USA, 1969.
- [18] I. Müller, T. Ruggeri, Extended Thermodynamics, Vol. 37 of *Springer Tracts in Natural Philosophy*, Springer-Verlag New York, 1993.
- [19] W. Dreyer, Maximisation of the entropy in non-equilibrium, *Journal of Physics A: Mathematical and General* 20 (18) (1987) 6505.
- [20] C. D. Levermore, Moment closure hierarchies for kinetic theories, *Journal of statistical Physics* 83 (5) (1996) 1021–1065.
- [21] M. Pavić, T. Ruggeri, S. Simić, Maximum entropy principle for rarefied polyatomic gases, *Physica A: Statistical Mechanics and its Applications* 392 (6) (2013) 1302–1317.
- [22] H. Hutridurga, F. Salvarani, Maxwell-stefan diffusion asymptotics for gas mixtures in non-isothermal setting, *Nonlinear Analysis* 159 (2017) 285–297.