

# COMBINATORIAL AND TOPOLOGICAL ASPECTS OF PATH POSETS, AND MULTIPATH COHOMOLOGY

LUIGI CAPUTI, CARLO COLLARI, AND SABINO DI TRANI

**ABSTRACT.** A cohomology theory for digraphs, called multipath cohomology, has been recently introduced by the authors. Its construction uses the so-called path poset, i.e. the poset of disjoint oriented simple paths in a digraph. The aim of this paper is to explore some combinatorial properties of the path posets, in order to provide new insights and computations. In particular, we develop acyclicity criteria for multipath cohomology, and compute it for oriented linear graphs. To conclude, we interpret multipath cohomology as the cohomology of a certain simplicial complex, and we investigate which spaces may or may not arise this way.

## 1. INTRODUCTION

Nowadays, cohomology theories have become extremely important tools in Mathematics. Cohomology theories of directed graphs (digraphs, for short), in particular, are of central interest for the scientific community, and recent years have witnessed a growth in the popularity of this subject – see, for example, [Hap89, Iva94, CYY01, ABM05, TW12, GLMY20, GLS21]. This is mostly due to applications to various research domains (in neuroscience and neurological disorders [LCK<sup>+</sup>11, CPH21], in endoscopy analysis [DEL<sup>+</sup>16], angiography [BMM<sup>+</sup>14], pulmonary diseases [BBD<sup>+</sup>18], finance [Gid17], fingerprint classification [GGM19], image classification [DMV17], to name few), and to the emergence of new tools in Topological Data Analysis employing homological methods, as, for example, persistent homology.

The list of desirable properties a graph cohomology must satisfy depend on the personal perspective; nonetheless, there are some which almost always are required, namely: non-triviality, functoriality, computability, and detection of combinatorial properties. A general framework for constructing cohomology theories of digraphs, including *multipath cohomology*, was introduced in [CCDT21]. In the same paper, it was also shown that multipath cohomology is non-trivial and functorial. The aim of this work is to show that also the last two properties in the list are satisfied. To achieve this result, we analyse topological and combinatorial properties of the path poset, which is of one of the main ingredients in the definition of multipath cohomology. In particular, we study how the combinatorics of the path poset may affect multipath cohomology. As a by-product, we obtain a number of techniques which can be used to explicitly compute  $H_\mu^k(-; \mathbb{K})$ . A first application of such techniques is the full computation of the multipath cohomology for oriented linear graphs – cf. Definition 5.1;

**Theorem 1.1** (Theorem 5.11). *Let  $L$  be an oriented linear graph and  $\mathbb{K}$  be a field. Then, there exist integers  $h, k_1, \dots, k_h$ , which are combinatorially determined only by  $L$ , such that the cohomology of  $L$  is trivial if  $k_i \equiv 0 \pmod{3}$  and  $1 \leq i < h$ , otherwise it decomposes as the tensor product of graded modules, as follows*

$$H_\mu^{*+h-1}(L; \mathbb{K}) = H_\mu^*(A_{3\lfloor k_1/3 \rfloor}; \mathbb{K}) \otimes \cdots \otimes H_\mu^*(A_{3\lfloor k_{h-1}/3 \rfloor}; \mathbb{K}) \otimes H_\mu^*(A_{k_h}; \mathbb{K}),$$

where each factor

$$H_\mu^k(A_n; \mathbb{K}) = \begin{cases} \mathbb{K} & \text{if } n = 3(k-1) + 2 \text{ or } n = 3k, \\ 0 & \text{otherwise} \end{cases}$$

is the multipath cohomology group of an alternating graph  $A_n$  – cf. Figure 1.



FIGURE 1. An alternating graph on  $n$  vertices. The edge between  $v_{n-1}$  and  $v_n$  can be oriented either way depending on the parity of  $n$ .

This result, together with the methods involved in its proof, shows that the multipath cohomology of a digraph captures relevant combinatorial information. Among other implications, our computations reveal the non-triviality of multipath cohomology.

**Theorem 1.2** (Proposition 4.21). *The multipath cohomology  $H_\mu^*(-; \mathbb{K})$  with coefficients in a field  $\mathbb{K}$  can be supported in arbitrarily high degree, and can be of arbitrary high dimension.*

Our interest in the combinatorial properties of the path poset is not limited to the purpose of understanding multipath cohomology groups, but it extends to its connections with the so-called monotone properties. A property of (di)graphs is called *monotone* if it is preserved under deletion of edges – e.g. being acyclic, being a forest *etc.* – and the study of the homotopy type of simplicial complexes associated to monotone properties of (di)graphs is a central topic in topology; see, for instance, the classical papers [Vas93, BW99, Koz99, BBL<sup>+</sup>99], as well as the more recent works [BH17, Mat19, PS18]. The study of a simplicial complex associated to the path poset fits into this framework.

In the last part of the paper, and more precisely in Section 6, we describe the relationship between multipaths and simplicial complexes, and some of the topological properties reflected in multipath cohomology. In fact, we show that for a given digraph  $G$ , there exists a simplicial complex  $X(G)$  whose (reduced) simplicial cohomology is the multipath cohomology of  $G$ .

**Theorem 1.3** (Theorem 6.8). *The multipath cohomology  $H_\mu^n(G; \mathbb{K})$  of  $G$  is isomorphic to the ordinary homology  $\tilde{H}^{n-1}(X(G); \mathbb{K})$ .*

In virtue of this theorem, we reinterpret some of our combinatorial results – for instance, we reprove a Mayer-Vietoris theorem for multipath cohomology. Then, we turn to the question of which simplicial complexes arise as multipath complexes. Among others, we observe that all wedges of spheres of the same dimension can be obtained in this way.

**Proposition 1.4** (Example 6.12 and Proposition 6.15). *Let  $P_n$  be the coherently oriented polygonal graph – cf. Figure 3. Then, the multipath complex  $X(P_n)$  is homeomorphic to  $\mathbb{S}^{n-1}$ . Therefore, we have*

$$H_\mu^i(P_n; \mathbb{K}) \cong \tilde{H}^{i-1}(\mathbb{S}^{n-1}; \mathbb{K}) \cong \begin{cases} \mathbb{K} & \text{if } i = n, \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, for all  $k, n \in \mathbb{N}$ , there exists  $G_{k,n}$  such that  $X(G_{k,n})$  is homotopic to  $\bigvee^k \mathbb{S}^n$ .

It remains open the problem of whether all wedges of spheres, or more complicated spaces, can be realised as  $X(G)$  – cf. Questions 6.16 and 6.18.

**On the combinatorial techniques.** The construction of multipath cohomology with arbitrary coefficients (algebras and bimodules) is quite abstract. In its general definition, one associates to a digraph  $G$  a certain poset, called *path poset*, consisting of all the (possibly disjoint) simple oriented paths in  $G$ , ordered by inclusion; then, using the fact that every poset can be seen as a category, certain abstract categorical constructions provide what we have called the multipath cohomology of  $G$ . In this process, the combinatorial information provided by the graph is somehow obscured. However, when restricting to coefficients in the base ring, instead of a general algebra, computations can be carried over the path poset. It becomes therefore useful to develop combinatorial tools to study path posets. We rely on the gluing construction  $\nabla$  (cf. Definition 3.3) to provide the description of a path poset in terms of simpler path posets.

Given a vertex  $v$  in  $G$ , order the  $k \geq 2$  edges with target  $v$ . Denote by  $G_v^k$  the graph obtained by deleting all such edges, and by  $G_v^{(h)}$  the graph obtained by deleting them all, but the  $h$ -th. A crucial combinatorial result is the following decomposition theorem – cf. Theorem 3.13.

**Theorem 1.5** (Theorem 3.13). *If the vertex  $v$  is the target of  $k \geq 2$  edges, then*

$$P(G) \cong \nabla_{P(G_v^k)} (P(G_v^{(1)}), \dots, P(G_v^{(k)})) .$$

*In other words, the path poset  $P(G)$  is isomorphic to an iterated gluing of the path posets of the subgraphs  $G_v^{(1)}, \dots, G_v^{(k)}$  over the path poset of  $G_v^k$ .*

A similar description applies in the case in which  $v$  is source  $k \geq 2$  edges. The decomposition in Theorem 1.5, together with a Mayer-Vietoris type result for multipath cohomology, provides useful acyclicity criteria; for example, assume that a digraph  $G$  satisfies the following conditions:

- (1) there exists a vertex  $v$  that is the target (or source) of  $k \geq 2$  edges;
- (2) all subgraphs  $G_v^{(1)}, \dots, G_v^{(k)}, G_v^k$  of  $G$  have trivial cohomology.

Then,  $H_\mu^*(G; \mathbb{K}) = 0$ . A different criterion (see Criterion B) uses the notion of  $\nu$ -equivalence, which, roughly speaking, is an embedding of graphs where the valence of vertices is preserved away from a certain set (cf. Definition 4.8).

**Criterion B.** Let  $G$  be a digraph and assume that there exists a  $\nu$ -equivalence  $\phi: H_{n,m} \rightarrow G$  away from  $w_1, \dots, w_n$  and  $x_1, \dots, x_m$ , where  $H_{n,m}$  is the graph illustrated in Figure 9. If  $(\phi(v_0), \phi(v_1))$  is not contained in any coherently oriented cycle<sup>1</sup> of  $G$ , then  $H_\mu^*(G; \mathbb{K}) = 0$ .

The above criteria and techniques can be applied to get explicitly the cohomology of a number of graphs, some of which are listed in Table 1. Further applications of our combinatorial results are the computation of the multipath cohomology for all oriented linear graphs (cf. Theorem 1.1), and the proof that multipath cohomology can have arbitrary dimension and can be supported in arbitrarily high degrees (cf. Theorem 1.2).

Finally, we develop a deletion-contraction type result for multipath cohomology with coefficients in an algebra  $A$ , namely Theorem 5.12. This result allows us to give a recursive formula for the (graded) Euler characteristic of  $A_n$ , which shows how the complexity of multipath cohomology increases in case  $A \neq R$ , and to reprove an analogue one due to Przytycki [Prz10, Lemma 3.3].

**Proposition 1.6** (Corollary 5.14). *Let  $R$  be an integral domain. For each (unital)  $R$ -algebra  $A$ , we have  $H_\mu^*(\mathbb{I}_n; A) = H_\mu^0(\mathbb{I}_n; A)$ , and*

$$\text{rank}_R(H_\mu^*(\mathbb{I}_n; A)) = \text{rank}_R(H_\mu^0(\mathbb{I}_n; A)) = \begin{cases} \text{rank}_R(A)(\text{rank}_R(A) - 1)^n & n \geq 1, \\ \text{rank}_R(A) & n = 0. \end{cases}$$

Proposition 1.6 could be proved, in case  $A$  is commutative, by using the deletion-contraction long exact sequence in chromatic homology [HGR05].

**Organisation of the paper.** In Section 2, we review some basic notions about digraphs and posets, as well as the construction of multipath cohomology. The main combinatorial results are given in Section 3, where we give a description of the path poset associated to a digraph in terms of gluing operations and cones. In particular, we prove our main decomposition result: Theorem 3.13. In Section 4, we proceed with some applications of the developed combinatorial methods to multipath cohomology; for instance, we show that the cone construction between posets is reflected on the level of cochain complexes as a mapping cone, we prove a Mayer-Vietoris type result, and provide acyclicity criteria for multipath cohomology to vanish. Section 5 is entirely devoted to linear graphs; we first give a complete characterisation of the

<sup>1</sup>That is the image of a regular morphism  $P_n \rightarrow G$ , for some  $n$ .









Digraph $G$	$H_\mu^0(G; \mathbb{K})$	$H_\mu^1(G; \mathbb{K})$	$H_\mu^2(G; \mathbb{K})$	$H_\mu^i(G; \mathbb{K}), i > 2$
	$\mathbb{K}$	0	0	0
	0	0	0	0
	0	0	$\mathbb{K}$	0
	0	$\mathbb{K}^{n-1}$	0	0
	0	0	$\mathbb{K}^{(n-1)(m-1)}$	0
	0	0	$\mathbb{K}^2$	0
	0	0	$\mathbb{K}$	$H_\mu^3(G; \mathbb{K}) \cong \mathbb{K}$
	0	0	0	$H_\mu^n(P_n; \mathbb{K}) \cong \mathbb{K}$

TABLE 1. Some digraphs and their respective multipath cohomologies.

cohomology groups of an oriented linear graph and then compute, in some cases, the associated Euler (graded) characteristics. We conclude the paper with Section 6, where we focus on the topological aspects of multipath cohomology. We show that multipath cohomology is simplicial, reprove Mayer-Vietoris in this setting and, finally, provide some further examples.

**Conventions.** Typewriter font, e.g.  $G, H$ , etc., are used to denote *finite* graphs (both directed and unoriented). All base rings are assumed to be unital and commutative, and algebras are assumed to be associative. Unless otherwise stated,  $R$  denotes a principal ideal domain,  $\mathbb{K}$  is a field,  $A$  is a unital  $R$ -algebra, and all tensor products  $\otimes$  are assumed to be over the base ring  $R$ . Given a cochain complex  $C^*$ , we denote by  $C^*[i]$  the shifted complex  $C^{*+i}$ . General references for graph theory, algebra, and algebraic topology are [Wes05], [Lan02], and [Koz08], respectively.

**Acknowledgements.** The authors extend their felicitations to I. C. Luigi Caputi acknowledges support from the École Polytechnique Fédérale de Lausanne via a collaboration agreement with the University of Aberdeen. Sabino Di Trani was partially supported by GNSAGA - INDAM group during the writing of this paper.

## 2. BASIC NOTIONS

In this section, we review some basic notions on (directed) graphs and posets and recall the construction of multipath cohomology, cf. [CCDT21]. In Subsection 2.4, we specialise the general construction to the case of multipath cohomology with coefficients in a field  $\mathbb{K}$  – *en lieu* of a general algebra  $A$ . The advantage of such choice will be apparent in Sections 4 and 5,

where we employ the combinatorial methods developed in Section 3 to compute the multipath cohomology of certain digraphs.

**2.1. Digraphs.** We start with reviewing some basic properties of directed graphs (see, for instance, [Wes05] for a complete introduction about this topic).

**Definition 2.1.** A *directed graph*  $G$ , often shortened to *digraph*, is a pair of finite sets  $(V, E)$ , where  $E \subseteq \{V \times V \setminus \{(v, v) \mid v \in V\}\}$ . The elements of  $V$  are called *vertices*, while the elements of  $E$  are called *edges* of  $G$ .

Unless otherwise stated, we will refer to digraphs, simply, as graphs. When dealing with (un)directed graphs, i.e. graphs for which the edges are not oriented, the adjective “(un)oriented” will be explicitly stated. The forthcoming definitions for digraphs apply verbatim to unoriented graphs by discarding the orientation of the edges – i.e. by replacing ordered pairs of vertices with unordered pairs.

*Remark 2.2.* Two vertices  $v$  and  $w$  in a digraph can share at most two edges:  $(v, w)$  and  $(w, v)$ . We will also use, in Section 6.3, a notion of (directed) graphs with multiple edges (more than two) between two vertices; these graphs are called multigraph or quivers.

By definition, an edge of a digraph is an ordered set of two distinct vertices, say  $e = (v, w)$ . The vertex  $v$  is called the *source* of  $e$ , while the vertex  $w$  is called the *target* of  $e$ . The source and target of an edge  $e$  will be denoted by  $s(e)$  and  $t(e)$ , respectively. If a vertex  $v$  is either a source or a target of an edge  $e$ , we will say that  $e$  is *incident to*  $v$ .

*Notation 2.3.* In the follow-up we may also deal with more than one graph at the time. In such cases, the set of vertices and edges of a digraph  $G$  will be denoted by  $V(G)$  and  $E(G)$ , respectively.

**Definition 2.4.** A *morphism of digraphs* from  $G_1$  to  $G_2$  is a function  $\phi: V(G_1) \rightarrow V(G_2)$  such that:

$$e = (v, w) \in E(G_1) \implies \phi(e) := (\phi(v), \phi(w)) \in E(G_2).$$

A morphism of digraphs is called *regular* if it is injective as a function.

A *sub-graph*  $H$  of a graph  $G$  is a graph such that  $V(H) \subseteq V(G)$  and  $E(H) \subseteq E(G)$ , and in such case we write  $H \leq G$ . If  $H \leq G$  and  $H \neq G$  we say that  $H$  is a *proper sub-graph* of  $G$ , and we write  $H < G$ .

**Definition 2.5.** If  $H \leq G$  and  $V(H) = V(G)$  we say that  $H$  is a *spanning sub-graph* of  $G$ .

Given a spanning proper sub-graph  $H \leq G$ , we can find an edge  $e \in E(G) \setminus E(H)$ . We use the following notation:

*Notation 2.6.* The spanning sub-graph of  $G$  obtained from  $H$  by adding an edge  $e$  is simply denoted by  $H \cup e$ .

**2.2. Posets.** In this subsection we review some basic notions about partially ordered sets.

**Definition 2.7.** A (strict) *partial order* on a set  $S$  is a transitive relation  $\triangleleft$  such that at most one among  $x \triangleleft y$ ,  $y \triangleleft x$ , and  $x = y$ , holds. A *partially ordered set*, or simply *poset*, is a pair  $(S, \triangleleft)$  consisting of a set  $S$  and a partial order  $\triangleleft$  on  $S$ . A morphism of posets  $f: (S, \triangleleft) \rightarrow (S', \triangleleft')$  is a monotonic map of sets; that is, a function  $f: S \rightarrow S'$  such that  $x \triangleleft y$  implies  $f(x) \triangleleft' f(y)$ .

Before moving forward, it is useful to recall two examples of posets.

*Example 2.8.* The *standard Boolean poset*  $\mathbb{B}(n)$  (of size  $2^n$ ) is the poset  $(\wp(\{0, \dots, n-1\}), \subseteq)$ , where  $\wp$  denotes the power set – i.e. the set of all subsets. A poset is called a *Boolean poset*, if it is isomorphic to the standard Boolean poset  $\mathbb{B}(n)$ , for some  $n$ .

*Example 2.9.* Let  $G$  be a digraph with  $n$  edges. The poset  $(SSG(G), <)$  of *spanning subgraphs* of  $G$  is given by all the spanning subgraphs of  $G$  with order relation given by the property of being a subgraph. The associated covering relation  $\prec$  can be described as follows:

$$H \prec H' \iff \exists e \in E(H') \setminus E(H) : H' = H \cup e .$$

Then,  $(SSG(G), <)$  is a Boolean poset isomorphic to  $\mathbb{B}(n)$  – see also [CCDT21, Example 2.14].

Given a partial order  $\triangleleft$  on a set  $S$ , there is an associated *covering relation*  $\tilde{\triangleleft}$ , given by  $x \tilde{\triangleleft} y$  if, and only if,  $x \triangleleft y$  and there is no  $z$  such that  $x \triangleleft z$ ,  $z \triangleleft y$ . In order to visually represent posets associated to digraphs, we use covering relations and the associated Hasse graphs:

**Definition 2.10.** The *Hasse (di-)graph*  $\text{Hasse}(S, \triangleleft)$  of a poset  $(S, \triangleleft)$  is the graph whose vertices are the elements of  $S$  and such that  $(x, y)$  is an edge if, and only if,  $x \tilde{\triangleleft} y$ .

Each morphism of digraphs  $\phi: \text{Hasse}(S, \triangleleft) \rightarrow \text{Hasse}(S', \triangleleft')$  induces a morphism of posets  $\phi: (S, \triangleleft) \rightarrow (S', \triangleleft')$ . We remark that not all morphisms of posets arise this way.

Recall that, given a poset  $(S, \triangleleft)$ , a *sub-poset* is a subset  $S' \subseteq S$  with the order relation  $\triangleleft_{|S' \times S'}$  induced by  $\triangleleft$ .

**Definition 2.11.** A sub-poset  $(S', \triangleleft_{|S' \times S'})$  is called *downward closed* with respect to  $(S, \triangleleft)$ , if whenever  $h \triangleleft h'$  and  $h' \in S$ , then  $h \in S$ .

Essential to the construction of multipath cohomology are the following properties:

**Definition 2.12.** Let  $(S, \triangleleft)$  be a poset and  $(S', \triangleleft_{|S' \times S'})$  be a sub-poset of  $(S, \triangleleft)$ .

- (1) We say that  $(S, \triangleleft)$  is *squared* if for each triple  $x, y, z \in S'$  such that  $z$  covers  $y$  and  $y$  covers  $x$ , there is a unique  $y' \neq y$  such that  $z$  covers  $y'$  and  $y'$  covers  $x$ . Such elements  $x, y, y'$ , and  $z$  will be called a *square* in  $S$ .
- (2) We say that  $(S', \triangleleft_{|S' \times S'})$  is *faithful* if the covering relation in  $S'$  induced by  $\triangleleft_{|S' \times S'}$  is the restriction of the covering relation in  $S$  induced by  $\triangleleft$ ;

Note that Boolean posets are squared and that the property of being squared or faithful is preserved under intersections [CCDT21, Proposition 2.21]. Furthermore, downward closed sub-posets are faithful, and each downward closed sub-poset of a squared poset is also squared.

If  $(S, \triangleleft_S)$  and  $(S', \triangleleft_{S'})$  are posets, then their *product poset* is the set  $S \times S'$  endowed with the relation

$$(1) \quad (x_1, x_2) \triangleleft_{S \times S'} (y_1, y_2) \iff (x_1, x_2) = (y_1, y_2) \text{ or } x_1 \triangleleft_S y_1 \text{ and } x_2 \triangleleft_{S'} y_2.$$

The definition of product poset is essential to introduce the cone of a poset. Let  $P$  be a poset.

**Definition 2.13.** The *cone of  $P$* , denoted by  $\mathfrak{Cone} P$ , is the product poset  $P \times \mathbb{B}(1)$ .

As  $\mathbb{B}(1)$  is (isomorphic to) the poset on the set  $\{0, 1\}$  with the relation  $0 < 1$ , the cone  $\mathfrak{Cone} P$  can also be seen as  $P \times \{0, 1\}$ .

**2.3. Path posets.** We introduce one of the main tools in the definition of multipath cohomology of directed graphs, the *path poset* associated to a directed graph  $G$ .

**Definition 2.14.** A *simple path* in  $G$  is a sequence of edges  $e_1, \dots, e_n$  of  $G$  such that  $s(e_{i+1}) = t(e_i)$  for  $i = 1, \dots, n - 1$ , and no vertex is encountered twice, i.e. if  $s(e_i) = s(e_j)$  or  $t(e_i) = t(e_j)$ , then  $i = j$ , and is not a cycle, i.e.  $s(e_1) \neq t(e_n)$ .

Let  $G$  be a graph and  $|G|$  its geometric realisation as a CW-complex – cf. [Hat00, Appendix A] and Section 6. We say that a graph is *connected*, if  $|G|$  is connected as CW-complex. Likewise, a *connected component* of  $G$  is a sub-graph  $H$  of  $G$  whose realisation  $|H|$  is connected.

*Remark 2.15.* If a connected graph  $G$  admits an ordering of the edges with respect to which it is a simple path, then it is isomorphic to graph  $I_n$ , for some  $n$ . The explicit isomorphism is given by the morphism of digraphs  $\phi : V(G) \rightarrow V(I_n) : s(e_i) \mapsto v_{i-1}, t(e_n) \mapsto v_n$ .

We are interested in taking disjoint sets of simple paths; following [TW12], we call them multipaths:

**Definition 2.16.** A *multipath* of  $G$  is a spanning sub-graph such that each connected component is either a vertex or its edges admit an ordering such that it is a simple path.

The set of multipaths of  $G$  has a natural partially ordered structure:

**Definition 2.17.** The *path poset* of  $G$  is the poset  $(P(G), <)$  associated to  $G$ , that is, the set of multipaths of  $G$  ordered by the relation of “being a sub-graph”.

Observe that the order relation makes sense, because each multipath is a subgraph of  $G$ .

*Remark 2.18.* The path poset  $P(G)$  is a downward closed subposet of a Boolean poset – the Boolean poset of all spanning subgraphs. Therefore, it is faithful and squared.

We use the following notation.

*Notation 2.19.* When the partial order on  $P(G)$  is not specified, we will always implicitly assume it to be the order relation  $<$ . Moreover, with abuse of notation, we will also write  $P(G)$  instead of  $(P(G), <)$ .

We now provide some examples of path posets – cf. [CCDT21, Section 2].

*Example 2.20.* Consider the coherently oriented linear graph  $I_n$  with  $n$  edges, illustrated in Figure 2. In this case, all the spanning sub-graphs are also multipaths, that is  $P(I_n) = SSG(I_n)$ . In particular, it follows that  $(P(I_n), <)$  is isomorphic to the Boolean poset  $\mathbb{B}(n)$ .




FIGURE 2. The coherently oriented linear graph  $I_n$ .

*Example 2.21.* Consider the coherently oriented polygonal graph  $P_n$  with  $n+1$  edges, illustrated in Figure 3. In this case all spanning sub-graphs, but the polygon itself, are multipaths. Equivalently, we have  $(P(P_n) \cup \{P_n\}, <) = (SSG(P_n), <)$ . In particular,  $(P(P_n), <)$ , is isomorphic to the Boolean poset  $\mathbb{B}(n)$  minus its maximum.




FIGURE 3. The coherently oriented polygonal graph  $P_n$ .

**2.4. Multipath cohomology.** Given a special type of poset coherently assigned to each digraph, and a choice of a sign assignment on it (see Definition 2.22), a cohomology theory for directed graphs has been introduced by the authors in [CCDT21]. We review here the general definition first, and then we specialise it to the case of coefficients in a field  $\mathbb{K}$ .

Let  $\mathbb{Z}_2$  be the cyclic group on two elements.

**Definition 2.22.** A sign assignment on a poset  $(S, \triangleleft)$  is an assignment of elements  $\epsilon_{x,y} \in \mathbb{Z}_2$  to each pair of elements  $x, y \in S$  with  $x \triangleleft y$ , such that the equation

$$(2) \quad \epsilon_{x,y} + \epsilon_{y,z} \equiv \epsilon_{x,y'} + \epsilon_{y',z} + 1 \pmod{2}$$

holds for each square  $x \triangleleft y, y' \triangleleft z$ .

*Remark 2.23.* A sign assignment on  $(S, \triangleleft)$  can be seen as a map  $\epsilon: E(\text{Hasse}(S, \triangleleft)) \rightarrow \mathbb{Z}_2$  on the edges  $E(\text{Hasse}(S, \triangleleft))$  of  $\text{Hasse}(S, \triangleleft)$ , such that Equation (2) holds for each square  $x \triangleleft y, y' \triangleleft z$  of  $S$ .

*A priori*, the existence of a sign assignment on a given poset is not clear. A cohomological sufficient condition for a sign assignment on a given poset to exist is exposed in [CCDT21, Section 3.2]. For the spanning sub-graphs poset – or, better, for Boolean posets – and their sub-posets, a sign assignment can be easily described:

*Example 2.24.* Let  $G$  be a graph with a fixed total ordering  $\triangleleft$  on the set of edges  $E(G)$ . Recall that  $H \prec H'$  in  $SSG(G)$  if, and only if,  $H' = H \cup e$ . Then, we can define a sign assignment on the poset  $SSG(G)$  as follows:

$$\epsilon(H, H') := \#\{e' \in E(H) \mid e' \triangleleft e\} \pmod{2},$$

where  $H' = H \cup e$  [Kho00].

More generally, one may ask when two sign assignments on a given poset are isomorphic. We refrain here from giving the definition of isomorphism of sign assignments – cf. [CCDT21, Definition 3.13] – but we point out that in cases of interest to us, all sign assignments are isomorphic – cf. [CCDT21, Theorem 3.16 & Corollary 3.17]:

*Remark 2.25.* Let  $P$  be a downward closed sub-poset of  $SSG(G)$ . Then, any two sign assignments  $\epsilon$  and  $\epsilon'$  on  $P$  are isomorphic. In particular, any two sign assignments on a path poset  $P(G)$  are isomorphic.

Recall that the *length* of a graph  $G$ , denoted by  $\text{lengt}(G)$ , is the number of edges in  $G$ .

**Definition 2.26.** Let  $P$  be a finite poset with a minimum  $m$ . The *level*  $\ell(x)$  of  $x \in P$  is the minimal length among all simple paths joining  $x$  to the minimum in  $\text{Hasse}(P)$ .

If  $P = P(G)$  is the path poset of a graph, then the level and the length coincide. More generally, if  $P$  is a faithful sub-poset of  $SSG(G)$ , the notion of level of an element  $H \in P$  can be extended as follows:

$$\ell(H) = \#E(H) + \#V(H) - \min\{\#E(H') + \#V(H') \mid H' \in P\}.$$

For the rest of the section  $R$  denotes a commutative ring with identity, and  $A$  an associative unital  $R$ -algebra. An *ordered digraph* is a digraph with a fixed well-ordering<sup>2</sup> of the vertices.

Let  $G$  be an ordered graph and let  $v_0 \in V(G)$  be the minimum with respect to the given ordering. Given a multipath  $H < G$ , to each connected component of  $H$  we associate a copy of  $A$ . Then we take the ordered tensor product. More concretely, if  $c_0 < \dots < c_k$  is the set of ordered connected components of  $H$ , we define:

$$(3) \quad \mathcal{F}_A(H) := A_{c_0} \otimes_R \dots \otimes_R A_{c_k},$$

<sup>2</sup>Every non-empty sub-set has a minimal element.



where all the modules are labelled by the respective component.

Assume  $H' = H \cup e$ . We define the source  $s(e, H)$  (resp. target  $t(e, H)$ ) of  $e$  in  $H$  as the index of the connected component of  $H$  containing the source (resp. target) of  $e$ . Denote by  $c_0, \dots, c_k$  the ordered components of  $H$ , denote by  $c'_0, \dots, c'_{k-1}$  the ordered components of  $H'$ , and assume that the addition of  $e$  merges  $c_i$  and  $c_j$ . Then, for each  $h = 0, \dots, k-1$ , there is a natural identification

$$(4) \quad c'_h = \begin{cases} c_h & \text{if } 0 \leq h < i \text{ or } i < h < j; \\ c_i \cup e \cup c_j & \text{if } h = i; \\ c_{h+1} & \text{if } j \leq h < k. \end{cases}$$

for some  $0 \leq i < j \leq k$ . Using this identification, we define  $\mu_{H \prec H'}: \mathcal{F}_A(H) \longrightarrow \mathcal{F}_A(H')$  as

$\mu_{H \prec H'}(a_0 \otimes \dots \otimes a_k) = a_0 \otimes \dots \otimes a_{s(e,H)-1} \otimes a_{s(e,H)} \cdot a_{t(e,H)} \otimes a_{s(e,H)+1} \otimes \dots \otimes \widehat{a_{t(e,H)}} \otimes \dots \otimes a_{k-1} \otimes a_k$  where  $\widehat{a_{t(e,H)}}$  indicates that  $a_{t(e,H)}$  is missing. Let  $\epsilon$  be a sign assignment on  $P(G)$ . We can now define the cochain groups

$$C_\mu^n(G; A) := \bigoplus_{\substack{H \in P(G) \\ \ell(H) = n}} \mathcal{F}_A(H)$$

together with the differential

$$d^n = d_\mu^n := \sum_{\substack{H \in P(G) \\ \ell(H) = n}} \sum_{\substack{H' \in P(G) \\ H \prec H'}} (-1)^{\epsilon(H, H')} \mu_{H \prec H'}.$$

It has been proved that  $(C_\mu^*(G; A), d^*)$  is a cochain complex [CCDT21, Theorem 4.10]. Furthermore, the path poset  $P(G)$  is squared and faithful by Remark 2.18, hence the homology groups of  $(C_\mu^*(G; A), d^*)$  do not depend on the sign assignment  $\epsilon$  used in the definition of  $(C_\mu^*(G; A), d^*)$  and on the choice of the ordering on  $V(G)$  [CCDT21, Corollary 3.18 & Proposition 4.11]. We are ready to give the definition of the multipath cohomology of a directed graph:

**Definition 2.27.** The *multipath cohomology*  $H_\mu^*(G; A)$  of a digraph  $G$  with coefficients in an algebra  $A$  is the homology of the cochain complex  $(C_\mu^*(G; A), d^*)$ .

Observe that, when  $A$  is the ring  $R$ , the tensor products in Equation (3) simply give  $\mathcal{F}_R(H) = R_{c_0} \otimes_R \dots \otimes_R R_{c_k} \cong R$ , for each multipath  $H \prec G$ . The isomorphism between the tensor powers of  $R$  and  $R$  itself is given by multiplication  $\mu$ , therefore the differential can be written as

$$d^n = \sum_{\substack{H \in P(G) \\ \ell(H) = n}} \sum_{\substack{H' \in P(G) \\ H \prec H'}} (-1)^{\epsilon(H, H')} \text{Id}_{H \prec H'}.$$

Identifying  $\mathcal{F}_R(H)$  with a copy of  $R$  gives us a set of linearly independent generators  $\{b_H\}_H$  (as free  $R$ -module) for  $C_\mu^*(G; R)$  indexed by multipaths.

*Remark 2.28.* If  $\phi: G_1 \rightarrow G_2$  is a regular morphism of digraphs, then it induces (functorially) a morphism of posets  $P\phi: P(G_1) \rightarrow P(G_2)$ , as it sends multipaths of  $G_1$  to multipaths of  $G_2$ . We obtain a (contravariant) morphism of cochain complexes

$$\phi^*: C_\mu^*(G_2; A) \rightarrow C_\mu^*(G_1; A)$$

where we fixed a sign assignment on  $P(G_2)$  and we considered the sign on  $P(G_1)$  by restriction.

We conclude the section with the computation of the multipath cohomology of the coherently oriented linear graph;

*Example 2.29.* Consider the coherently oriented linear graph  $I_n$  of length  $n$ , illustrated in Figure 2. Then, its multipath poset  $(P(I_n), <)$  is a Boolean poset, and its multipath cohomology  $H_\mu^*(I_n; \mathbb{K})$  is trivial [CCDT21, Example 4.20].

### 3. A COMBINATORIAL DESCRIPTION OF PATH POSETS

The aim of this section is to give a combinatorial description of the path poset associated to a directed graph  $G$ ; we show that the path poset can be constructed by gluing together simpler path posets associated to (suitable) subgraphs of  $G$ .

We start by recalling two results on the poset structure of the path posets associated to coherently oriented linear graphs  $I_n$  – cf. Figure 2 – and polygons  $P_n$  – cf. Figure 3. Example 2.29 shows that the path poset of the coherently oriented linear graph is a Boolean poset, and Example 2.21 shows that the path poset of the coherently oriented polygonal graph is isomorphic to a Boolean poset with the maximum element removed. These are the only graphs giving such path posets – cf. [CCDT21, Proposition 2.34 & 2.35];

**Proposition 3.1.** *Let  $G$  be a connected graph of length  $n > 2$ .*

- (1) *If  $P(G)$  has a maximum then we have that  $P(G) \cong \mathbb{B}(n)$  and  $G \cong I_n$ . In particular, a connected graph has a Boolean path poset if, and only if, it is isomorphic to  $I_n$ , for some  $n$ .*
- (2) *If  $P(G)$  is isomorphic to  $\mathbb{B}(n)$  minus its maximum, then  $G \cong P_{n-1}$ .*

Sometimes, we might deal with disconnected graphs. A straightforward observation is that the multipath cohomology of disconnected graphs is the tensor product of the multipath cohomologies:

*Remark 3.2.* Let  $G$  be the disjoint union of connected digraphs  $G_1, \dots, G_n$ . Then, the path poset  $P(G)$  is the product  $P(G_1) \times \dots \times P(G_n)$  of the path posets on  $G_1, \dots, G_n$  – cf. Equation (1) – and the multipath cohomology of  $G$  splits as the following graded tensor product:

$$H_\mu^*(G; \mathbb{K}) = H_\mu^*(G_1; \mathbb{K}) \otimes \dots \otimes H_\mu^*(G_n; \mathbb{K}).$$

In particular, if  $H_\mu^*(G_i; \mathbb{K}) = 0$  for some  $i \in \{1, \dots, n\}$ , then  $H_\mu^*(G; \mathbb{K}) = 0$ .

We introduce a gluing operation for directed graphs.

**Definition 3.3** (Gluing). Let  $G, G_1, G_2$  be directed graphs, and  $\iota_1: G \rightarrow G_1$  and  $\iota_2: G \rightarrow G_2$  be regular morphisms of digraphs. The *gluing* of  $G_1$  and  $G_2$  along  $G$  is the directed graph  $\nabla_G(G_1, G_2)$  defined as follows:

- (1)  $V(\nabla_G(G_1, G_2)) := V(G_1) \sqcup V(G_2) / \sim$ , where  $x \sim y$  if, and only if, either  $x = y$  or  $x \in \iota_1(G)$ ,  $y \in \iota_2(G)$ , and  $\iota_1^{-1}(x) = \iota_2^{-1}(y)$ ;
- (2)  $([v], [w]) \in E(\nabla_G(G_1, G_2))$  if, and only if, there exist  $v' \in [v]$ ,  $w' \in [w]$ , and  $i \in \{1, 2\}$  such that  $(v', w') \in E(G_i)$ , where  $[\cdot]$  denotes an equivalence class with respect to  $\sim$ .

Roughly speaking,  $\nabla_G(G_1, G_2)$  is the graph obtained from  $G_1$  and  $G_2$  by identifying the vertices and edges belonging to the image of  $G$ .

When clear from the context and for ease of notation, we denote the edge  $([v], [w])$  in the set  $E(\nabla_G(G_1, G_2))$  as  $(v, w)$ .

*Remark 3.4.* For a given graph  $G$ , the operation  $\nabla_G(-, -)$  is commutative and associative up to isomorphism of digraphs.

Let **Digraph** be the category of digraphs and regular morphisms of digraphs. We can reinterpret the gluing as a categorical pushout:

*Remark 3.5.* The operation  $\nabla_G(-, -)$  represents the categorical push-out – cf. [Mac71, Section III.3] – in the category **Digraph**. Since  $\nabla_G(G_1, G_2)$  is an object of **Digraph** and since the inclusions of  $G_1$  and  $G_2$  in  $\nabla_G(G_1, G_2)$  are regular morphisms of digraphs, we have a commutative

square

$$\begin{array}{ccc} \mathbf{G} & \xrightarrow{\iota_1} & \mathbf{G}_1 \\ \downarrow \iota_2 & & \downarrow j_1 \\ \mathbf{G}_2 & \xrightarrow{j_2} & \nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2) \end{array}$$

in **Digraph**. Note that  $V(\nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2))$  is the push-out of  $V(\mathbf{G}_1)$  and  $V(\mathbf{G}_2)$  along  $V(\mathbf{G})$  in the category **Set** of sets. Now, given another digraph  $\mathbf{G}'$  such that the square

$$\begin{array}{ccc} \mathbf{G} & \xrightarrow{\iota_1} & \mathbf{G}_1 \\ \downarrow \iota_2 & & \downarrow j'_1 \\ \mathbf{G}_2 & \xrightarrow{j'_2} & \mathbf{G}' \end{array}$$

commutes, then we get a function  $V(\nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2)) \rightarrow V(\mathbf{G}')$  since  $V(\nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2))$  is a push-out in **Set**. Such a function extends to a map of digraphs by definition of morphism of digraphs, which is injective as it is composition of injective functions. This is enough to show that  $\nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2)$  is the categorical push-out in **Digraph**.

Before proceeding further, we establish a notation for gluings over more than two graphs.

*Notation 3.6.* If  $\mathbf{G}'$  is a digraph such that there exist two morphisms of digraphs  $\iota: \mathbf{G}' \rightarrow \nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2)$  and  $j: \mathbf{G}' \rightarrow \mathbf{G}_3$  we define:

$$\nabla_{\mathbf{G}', \mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3) := \nabla_{\mathbf{G}'}(\nabla_{\mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2), \mathbf{G}_3) .$$

In general, if  $\mathbf{G}_1, \dots, \mathbf{G}_n$  is a family of digraphs such that for every  $k < n$  there exist a (regular) morphisms of digraphs  $\iota_k: \mathbf{G} \rightarrow \mathbf{G}_k$  we will denote  $\nabla_{\mathbf{G}, \dots, \mathbf{G}}(\mathbf{G}_1, \dots, \mathbf{G}_n)$  by  $\nabla_{\mathbf{G}}(\mathbf{G}_1, \dots, \mathbf{G}_n)$ .

Note that  $\nabla_{\mathbf{G}}(\mathbf{G}_1, \dots, \mathbf{G}_n)$  does not depend, up to isomorphism of digraphs, on the order of the digraphs  $\mathbf{G}_1, \dots, \mathbf{G}_n$ , whereas  $\nabla_{\mathbf{G}', \mathbf{G}}(\mathbf{G}_1, \mathbf{G}_2, \mathbf{G}_3)$  might.

**Definition 3.7.** The gluing of two posets  $P_1$  and  $P_2$  along a common subposet  $P$  is the poset, denoted by  $\nabla_P(P_1, P_2)$ , whose Hasse diagram is the gluing of  $\text{Hasse}(P_1)$  and  $\text{Hasse}(P_2)$  along  $\text{Hasse}(P)$ .

*Remark 3.8.* Observe that the gluing does not commute with the operation of taking path posets.

Let  $\mathbf{G}_1$  be the graph  $v_0 \rightarrow v_1 \rightarrow v_2$  and  $\mathbf{G}_2$  the graph  $v_1 \rightarrow v_2 \rightarrow v_3$  and consider the gluing  $\mathbf{G}$  of  $\mathbf{G}_1$  and  $\mathbf{G}_2$  over  $v_1 \rightarrow v_2$ . Then, the poset  $P(\mathbf{G})$  is isomorphic to the Boolean poset  $\mathbb{B}(3)$ , whose Hasse diagram is (the 1-skeleton of) a 3-dimensional cube. On the other hand, the Hasse diagram of  $\nabla_{P(\mathbf{G})}(P(\mathbf{G}_1), P(\mathbf{G}_2))$  is the gluing of two copies of  $\text{Hasse}(\mathbb{B}(2))$  along a copy of  $\text{Hasse}(\mathbb{B}(1))$  – that is two (empty) squares attached along an edge.

Recall that the cone of a poset  $P$ , denoted by  $\mathfrak{Cone} P$ , is the product poset  $P \times \mathbb{B}(1)$ .

*Remark 3.9.* The covering relations in  $\mathfrak{Cone} P$  can be described in a very explicit way. An element  $(a, i)$  is covered by  $(b, j)$  if and only if  $i = j$  and  $a \triangleleft b$  or  $i < j$  and  $a = b$ .




FIGURE 4. Linear source and sink.

**Definition 3.10.** Let  $G$  be a digraph and let  $v \in V(G)$  be a vertex. We say that  $v$  is a *sink* (resp. a *source*) of  $G$  if,  $e \in E(G)$  and  $e$  incident to  $v$  implies that  $v$  is a target (resp. a source) of  $e$ . A digraph  $G$  with  $n$  edges is a *sink* (resp. a *source*) over  $n$  vertices if it has a sink (resp. source)  $v$ , and every edge in  $G$  is incident to it.

Some straightforward examples of cone constructions follow.

*Example 3.11.* The path poset of a linear sink on three vertices – cf. Figure 4 (b) – is the gluing of two cones over the empty multipath – cf. Figure 5.




FIGURE 5. The path poset of the linear sink on three vertices. Each coloured arrow represents a cone over a point, and these are glued together on a point.

*Example 3.12.* The path poset of the graph  $I_2$  is a cone over the path poset of  $I_1$  – cf. Figure 6.




FIGURE 6. The path poset of  $I_2$ . In orange we denoted the relations coming from the coning process.

Let  $G$  be a digraph. We now relate the path poset  $P(G)$  to the path posets associated to some subgraphs of  $G$ . First, we need to fix some more notation. For a vertex  $v \in V(G)$ , consider the set  $E_v$  of edges  $e_1, \dots, e_n$  in  $G$  incident to  $v$ , ordered so that  $v = t(e_i)$ , for  $i = 1, \dots, k$ , and  $v = s(e_j)$ , for  $j = k + 1, \dots, n$ . Denote by  $G_v^k$  the graph obtained by deleting from  $G$  the edges  $e_1, \dots, e_k$ , and set  $G_v^{(h)} := G_v^k \cup e_h$ .

**Theorem 3.13.** *If the vertex  $v$  is a target for  $k \geq 2$  edges, then*

$$P(G) \cong \nabla_{P(G_v^k)} (P(G_v^{(1)}), \dots, P(G_v^{(k)})) .$$

*In other words, the path poset  $P(G)$  is isomorphic to an iterated gluing of the path posets of the subgraphs  $G_v^{(1)}, \dots, G_v^{(k)}$  over the path poset of  $G_v^k$ .*

*Proof.* We recall that, if  $G'$  is a subgraph of a digraph  $G$ , then every multipath in  $P(G')$  can be seen as a multipath in  $P(G)$ . This means that  $P(G')$  can be seen as an downward closed (and, in particular, faithful) subposet of  $P(G)$ .

To prove the theorem, we want to produce an isomorphism of posets

$$P(G) \cong \nabla_{P(G_v^k)} (P(G_v^{(1)}), \dots, P(G_v^{(k)})) .$$

First, we start by identifying the underlying sets and, then, we proceed with proving that the respective poset structures are isomorphic.

For the rest of the proof, denote by  $T$  the set  $\{e \in E(G) \mid t(e) = v\}$ . Let  $H$  be a multipath in  $P(G)$ . We have two possible cases:

Case 1:  $H$  does not contain any edge in  $T$ . Then, all simple paths in  $H$  are simple paths in  $G_v^k$ , and thus  $H \in P(G_v^k)$ .

Case 2:  $H$  contains an edge  $e_h \in T$ . In this case  $H$  cannot contain any other  $e_j \in T$ . Therefore, we have  $H \in P(G_v^{(h)}) \setminus P(G_v^k)$ .

On the other hand, observe that any multipath  $H \in \nabla_{P(G_v^k)} \left( P(G_v^{(1)}), \dots, P(G_v^{(k)}) \right)$  can be identified with either an element of  $P(G_v^{(j)}) \setminus P(G_v^k)$ , for some  $j \in \{1, \dots, k\}$ , or with an element of the poset  $P(G_v^k)$ . Thus, we have a way to uniquely identify  $H$  with an element of  $P(G)$  and, consequently, the underlying sets of  $P(G)$  and  $\nabla_{P(G_v^k)} \left( P(G_v^{(1)}), \dots, P(G_v^{(k)}) \right)$ . With abuse of notation, we denote this element again by  $H$ .

Now, we want to prove that the general multipath  $H$  covers the same elements both in  $P(G)$  and  $\nabla_{P(G_v^k)} \left( P(G_v^{(1)}), \dots, P(G_v^{(k)}) \right)$ . This is obvious if  $H$  is a multipath of  $P(G_v^k)$ . Assume that  $H$  is in  $P(G) \setminus P(G_v^k)$ . Then, there exists a unique  $e_j \in T$  such that  $e_j \in H$ . A multipath covered by  $H$  is then a multipath in  $P(G_v^{(j)})$  and consequently the same covering relations hold in  $\nabla_{P(G_v^k)} \left( P(G_v^{(1)}), \dots, P(G_v^{(k)}) \right)$ . Finally, if  $H \in \nabla_{P(G_v^k)} \left( P(G_v^{(1)}), \dots, P(G_v^{(k)}) \right)$ , then all the elements covered by  $H$  are contained in  $P(G_v^{(j)})$  and it is possible to conclude the proof because  $P(G_v^{(j)})$  is an downward closed subposet of  $P(G)$ .  $\square$

Note that the decomposition provided in the theorem can not be applied only in the special cases of (disjoint unions of) coherently oriented linear graphs and polygons.

*Remark 3.14.* A decomposition of  $P(G)$  similar to the one provided by Theorem 3.13 can be found using edges with source (instead of target) vertex  $v$ , by an orientation reversing argument. In fact, denoting by  $G^{\text{op}}$  the graph obtained by  $G$  reversing the orientation of its edges, we have that  $P(G) \cong P(G^{\text{op}})$ .

Theorem 3.13 is one of the main tools used in the computations and applications carried on in the next few sections; in fact, it simplifies the poset structure of the path posets, leading to quicker computations for the multipath cohomology groups with coefficients in a field.

#### 4. APPLICATIONS TO MULTIPATH COHOMOLOGY

The combinatorial analysis of path posets investigated in the previous section translates in some remarkable properties of multipath cohomology. In the first part of this section we provide some results concerning the cohomology of cones and gluings of digraphs; in particular, we see the connection with the algebraic cone of cochain complexes. We proceed with proving a Mayer-Vietoris type theorem for multipath cohomology. We apply these techniques to computations of multipath cohomology groups in some examples, together with some aciclicity criteria.

**4.1. The cohomology of the cone construction.** The aim of this subsection is to prove that the gluing and the cone of digraphs have effects on multipath cohomology. We start by proving that a graph  $G'$  whose path poset is isomorphic to a cone  $\mathfrak{Cone} P(G)$ , for some digraph  $G$ , has trivial cohomology.

*Remark 4.1.* A poset  $P = (S, \triangleleft)$  can be seen as a category  $\mathbf{P}$  with set of objects  $S$  and a morphism  $x \rightarrow y$  between  $x$  and  $y$  if, and only if,  $x \triangleleft y$  or  $x = y$ . For a functor  $\mathcal{F}: \mathbf{P} \rightarrow \mathbb{K}\text{-Vect}$  to the category of vector spaces over  $\mathbb{K}$ , with some mild assumptions on  $P$ , there are well defined cohomology groups  $H^*(\mathbf{P}; \mathcal{F})$  of  $\mathbf{P}$  with coefficients in  $\mathcal{F}$  – cf. [CCDT21, Theorem 3.7] – which, when  $P = P(G)$  is the path poset of a digraph  $G$ , gives the multipath cohomology.

We denote by  $(C^*(P; \mathbb{K}), \partial_P)$  the cochain complex  $(C_{\mathcal{F}_{\mathbb{K}}}^*(P), \partial)$  associated to a poset  $P$  (and level  $\ell$ ) and to the functor  $\mathcal{F}_{\mathbb{K}}$  assigning a copy of  $\mathbb{K}$  to each object of  $\mathbf{P}$  and the identity  $\mathbb{K} \rightarrow \mathbb{K}$  to each arrow. Analogously, we denote by  $H^*(P; \mathbb{K})$  the cohomology groups of the cochain complex  $(C^*(P; \mathbb{K}), \partial_P)$ . Recall that for a map  $f: (A, \partial_A) \rightarrow (B, \partial_B)$  of cochain complexes, the mapping cone  $\text{Cone} f$  is the cochain complex defined in degree  $n$  as  $(\text{Cone} f)^n := A^{n+1} \oplus B^n$  and differential

$$\partial := \begin{pmatrix} \partial_A[1] & 0 \\ f[1] & \partial_B \end{pmatrix}$$

where  $\partial_A[1]$  and  $f[1]$  represent the differential  $\partial_A$  and the morphism  $f$  shifted by one – see, for instance, [Gel99, Chapter 4, §2.2], and also [Wei94].

**Theorem 4.2.** *Let  $G$  be a digraph. Then, we get an isomorphism of cochain complexes*

$$C^*(\mathfrak{Cone} P(G); \mathbb{K}) \cong \text{Cone} \left( \text{Id}_{C_{\mu}^*(G; \mathbb{K})} \right) [-1]$$

where  $\text{Cone} \left( \text{Id}_{C_{\mu}^*(G; \mathbb{K})} \right)$  represents the mapping cone of the identity map on the cochain complex  $C_{\mu}^*(G; \mathbb{K})$ . Consequently, we have  $H^*(\mathfrak{Cone} P(G); \mathbb{K}) = 0$ .

To simplify the notation, we drop the reference to  $\mathbb{K}$  in the proof of the theorem.

*Proof.* Recall that the cone of a poset  $P$  is the product poset  $P \times \mathbb{B}(1)$  – cf. Definition 2.13. Consider the partition  $P(G) \times \{0\} \sqcup P(G) \times \{1\} = \mathfrak{Cone} P(G)$ . Furthermore, we also have that  $\ell_{\mathfrak{Cone} P(G)}(H, i) = \ell_{P(G) \times \{i\}}(H, i) + i = \ell_{P(G)}(H) + i$ , for  $i \in \{0, 1\}$ . As a consequence, we have the isomorphism of graded  $\mathbb{K}$ -vector spaces

$$C^*(\mathfrak{Cone} P(G)) \cong C^*(P(G) \times \{0\}) \oplus C^*(P(G) \times \{1\})[-1]$$

where  $C^*(P(G))[-1]$  denotes the complex  $C^*(P(G))$  shifted by one. In turn, we have an isomorphism of posets  $P(G) \cong P(G) \times \{i\}$ , for  $i = 0, 1$ , given by the identification  $H \mapsto (H, i)$ . These identifications induce the isomorphism of graded  $\mathbb{K}$ -vector spaces

$$C^*(P(G) \times \{0\}) \oplus C^{*-1}(P(G) \times \{1\}) \cong C_{\mu}^*(G) \oplus C_{\mu}^{*-1}(G) = \text{Cone} \left( \text{Id}_{C_{\mu}^*(G; \mathbb{K})} \right) [-1].$$

Now, we have to show that the above isomorphism commutes with the differentials. The differential  $\partial$  of  $\text{Cone} \left( \text{Id}_{C_{\mu}^*(G; \mathbb{K})} \right) [-1]$  is defined as

$$\partial := \begin{pmatrix} \partial_{\mu} & 0 \\ \text{Id}_{C_{\mu}^*(G; \mathbb{K})} & \partial_{\mu}[-1] \end{pmatrix}$$

where  $(\partial_{\mu}[-1])^n := (-1)^n \partial_{\mu}^{n-1}$ . The differential of  $C^*(\mathfrak{Cone} P(G))$ , can be explicitly written:

$$\partial_{\mathfrak{Cone} P(G)}(e_{(H, i)}) = (1 - i)e_{(H, 1)} + \sum_{H \prec H'} (-1)^{\epsilon(H, H') + i} e_{(H', i)}$$

where  $\epsilon(H, H')$  is a sign assignment on  $P(G)$  and  $e_{(H, i)}$  is the generator of  $C^*(P(G) \times \{i\})$  associated to the multipath  $H$  in  $G$ . It is easy to check that  $\epsilon'((H, i), (H', j)) := \epsilon(H, H') + i$  is a sign assignment on  $\mathfrak{Cone} P(G)$ .

The isomorphism  $C^*(\mathfrak{Cone} P(G)) \cong \text{Cone} \left( \text{Id}_{C_{\mu}^*(G; \mathbb{K})} \right) [-1]$  described above commutes with these differentials, concluding the proof of the first part of the statement. The second part follows from the classical properties of the mapping cone of chain complexes [Wei94].  $\square$

We observe that the second part of Theorem 4.2 can be alternatively proved using discrete Morse theory – see [Koz08, Chapter 11] for an introduction; if we consider the edges in  $\text{Hasse}(\mathfrak{Cone} P(G))$  with source in  $P(G) \times \{0\}$  and target in  $P(G) \times \{1\}$ , then these form an acyclic matching ([Koz08, Definition 11.1]) whose edges are incident to all vertices of the

graph  $\text{Hasse}(\mathbf{Cone} P(\mathbb{G}))$ . It follows from the definitions and [Koz08, Theorem 11.24] that the homology of  $C^*(\mathbf{Cone} P(\mathbb{G}); \mathbb{K})$  is trivial.

**4.2. A Mayer-Vietoris theorem.** The goal of this subsection is to prove a result which is the analogue, in the framework of multipath cohomology, of the classical Mayer-Vietoris theorem for simplicial homology. In the classical statement, given a decomposition of a topological space as union of two subspaces, there is an induced long exact sequence of (co-)homology groups featuring also their intersections. In the setting of multipath cohomology, the rôle played by unions of topological spaces is given by the gluing of posets. In Subsection 6.3, we will see how this result can also be deduced from the classical Mayer-Vietoris theorem.

Before stating the theorem, we recall a result of functoriality for multipath cohomology. Note that, for  $\phi: \mathbb{G}' \rightarrow \mathbb{G}$ , we have an induced morphism of posets  $P\phi: P(\mathbb{G}') \rightarrow P(\mathbb{G})$ ; more precisely, to a multipath  $H \in P(\mathbb{G}_1)$  we associate the subgraph  $P\phi(H)$  of  $\mathbb{G}_2$  defined by  $V(P\phi(H)) = V(\mathbb{G})$  and  $E(P\phi(H)) = \{\phi(e) \mid e \in E(H)\}$  [CCDT21, Remark 2.33].

**Proposition 4.3** ([CCDT21, Proposition 6.11]). *Let  $\phi: \mathbb{G}' \rightarrow \mathbb{G}$  be regular morphism of digraphs. The inclusion of  $P(\mathbb{G}')$  in  $P(\mathbb{G})$  induces the following short exact sequence of complexes*

$$0 \rightarrow C^*(P_\phi(\mathbb{G}', \mathbb{G}); \mathbb{K}) \left[ - \min_{x \in P_\phi(\mathbb{G}', \mathbb{G})} \{\ell(x)\} \right] \rightarrow C_\mu^*(\mathbb{G}; \mathbb{K}) \rightarrow C_\mu^*(\mathbb{G}'; \mathbb{K}) \rightarrow 0$$

where  $P_\phi(\mathbb{G}', \mathbb{G})$  is the poset  $P(\mathbb{G}) \setminus P(\phi)(P(\mathbb{G}'))$ .

In particular, the above proposition gives us a map between the multipath cochain complex of a graph  $\mathbb{G}$ , and the multipath cochain complex of a sub-graph  $\mathbb{G}_1$ . Such maps play a key role in the following the Mayer-Vietoris theorem for multipath cohomology.

**Theorem 4.4.** *Let  $\mathbb{G}, \mathbb{G}_1, \mathbb{G}_2$  be directed graphs, and  $i_1: \mathbb{G} \rightarrow \mathbb{G}_1$  and  $i_2: \mathbb{G} \rightarrow \mathbb{G}_2$  be regular morphisms of digraphs. Then, we have a short exact sequence of cochain complexes*

$$(5) \quad 0 \rightarrow C^*(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K}) \xrightarrow{I^*} C_\mu^*(\mathbb{G}_1; \mathbb{K}) \oplus C_\mu^*(\mathbb{G}_2; \mathbb{K}) \xrightarrow{J^*} C_\mu^*(\mathbb{G}; \mathbb{K}) \rightarrow 0$$

inducing the long exact sequence

$$\cdots \rightarrow H_\mu^{i-1}(\mathbb{G}; \mathbb{K}) \rightarrow H^i(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K}) \rightarrow H_\mu^i(\mathbb{G}_1; \mathbb{K}) \oplus H_\mu^i(\mathbb{G}_2; \mathbb{K}) \rightarrow H_\mu^i(\mathbb{G}; \mathbb{K}) \rightarrow \cdots$$

of cohomology groups.

*Proof.* We first observe that, as a consequence of the definition of gluing – cf. Definition 3.3 –  $P(\mathbb{G}_1)$  and  $P(\mathbb{G}_2)$  are isomorphic to subposets of  $\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2))$ ; call  $j_1$  and  $j_2$  these isomorphisms. The inclusions of graphs  $i_1: \mathbb{G} \rightarrow \mathbb{G}_1$  and  $i_2: \mathbb{G} \rightarrow \mathbb{G}_2$  induce morphisms of posets  $\iota_1: P(\mathbb{G}) \rightarrow P(\mathbb{G}_1)$  and  $\iota_2: P(\mathbb{G}) \rightarrow P(\mathbb{G}_2)$ . All the resulting morphisms fit into the following commutative square of posets

$$\begin{array}{ccc} P(\mathbb{G}) & \xrightarrow{j_1} & P(\mathbb{G}_1) \\ j_2 \downarrow & & \downarrow \iota_1 \\ P(\mathbb{G}_2) & \xrightarrow{\iota_2} & \nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)) \end{array}$$

and, induce a commutative diagram

$$\begin{array}{ccc} C^*(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K}) & \xrightarrow{i_1^*} & C_\mu^*(\mathbb{G}_1; \mathbb{K}) \\ \downarrow i_2^* & & \downarrow j_1^* \\ C_\mu^*(\mathbb{G}_2; \mathbb{K}) & \xrightarrow{j_2^*} & C_\mu^*(\mathbb{G}; \mathbb{K}) \end{array}$$

of cochain complexes.

Now, for every  $n \in \mathbb{N}$ , consider the maps

$$I^n := i_1^n \oplus i_2^n : C^n(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K}) \rightarrow C_\mu^n(\mathbb{G}_1; \mathbb{K}) \oplus C_\mu^n(\mathbb{G}_2; \mathbb{K}),$$

and

$$J^n := j_1^n - j_2^n : C_\mu^n(\mathbb{G}_1; \mathbb{K}) \oplus C_\mu^n(\mathbb{G}_2; \mathbb{K}) \rightarrow C_\mu^n(\mathbb{G}; \mathbb{K}).$$

We proceed with proving that the sequence of complexes in Equation (5) is exact. The cochain complexes  $C^*(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K})$ ,  $C_\mu^n(\mathbb{G}_1; \mathbb{K})$ , and  $C_\mu^n(\mathbb{G}_2; \mathbb{K})$  have bases indexed by the elements of the corresponding poset (namely,  $\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2))$ ,  $P(\mathbb{G}_1)$ , and  $P(\mathbb{G}_2)$ , respectively). We denote by  $b_H$  the element of each of these bases corresponding the multipath  $H$ .

With this notation, a generic element  $x$  of  $C^n(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K})$  is of the form

$$x = \sum_{H \in P(\mathbb{G})} \alpha_H b_{i_1 \circ j_1(H)} + \sum_{H' \in P(\mathbb{G}_1) \setminus j_1(P(\mathbb{G}))} \beta_{H'} b_{i_1(H')} + \sum_{H'' \in P(\mathbb{G}_2) \setminus j_2(P(\mathbb{G}))} \gamma_{H''} b_{i_2(H'')}$$

with  $\ell(H) = \ell(H') = n$ . Note that  $i_1 \circ j_1 = i_2 \circ j_2$ , thus  $b_{i_1 \circ j_1(H)} = b_{i_2 \circ j_2(H)}$  for each  $H \in P(\mathbb{G})$ .

We are now ready to verify that  $I^n$  is injective. With respect to the basis described above, we can write

$$I^n(x) = \left( \sum_{H \in P(\mathbb{G})} \alpha_H b_{j_1(H)} + \sum_{H' \in P(\mathbb{G}_1) \setminus j_1(P(\mathbb{G}))} \beta_{H'} b_{H'}, \sum_{H \in P(\mathbb{G})} \alpha_H b_{j_2(H)} + \sum_{H'' \in P(\mathbb{G}_2) \setminus j_2(P(\mathbb{G}))} \gamma_{H''} b_{H''} \right).$$

It follows that  $I^n(x) = 0$  if and only if all the coefficients  $\alpha_H$ ,  $\beta_{H'}$  and  $\gamma_{H''}$ , and thus  $x$ , are zero.

It is left to show that  $J^n$  is surjective and that  $\text{Ker}(I^n) = \text{Im}(J^n)$ . First, we write  $J^n$  explicitly, as follows;

$$J^n \left( \sum_{H' \in P(\mathbb{G}_1)} \alpha_{H'} b_{H'}, \sum_{H'' \in P(\mathbb{G}_2)} \beta_{H''} b_{H''} \right) = \sum_{H \in P(\mathbb{G})} (\alpha_{j_1(H)} - \beta_{j_2(H)}) b_H.$$

Now, for  $J^n(x)$  to be zero, we must have that  $\alpha_{j_1(H)} = \beta_{j_2(H)}$ , for all  $H \in P(\mathbb{G})$ , independently on the values of  $\alpha_{H'}$  and  $\beta_{H''}$  for  $H' \notin j_1(P(\mathbb{G}))$ , and  $H'' \notin j_2(P(\mathbb{G}))$ . It follows that the kernel  $J^n$  is precisely the image of  $I^n$ . Finally,  $J^n$  is clearly surjective, concluding the proof.  $\square$

The following observation is straightforward:

*Remark 4.5.* Under the assumptions of Theorem 4.4, if  $\mathbb{G}_1$  and  $\mathbb{G}_2$  have trivial multipath cohomology, then  $H^n(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), P(\mathbb{G}_2)); \mathbb{K})$  is isomorphic to  $H_\mu^{n-1}(\mathbb{G}; \mathbb{K})$  for all  $n$ .

By iterated applications of the Mayer-Vietoris long exact sequence for the multipath cohomology, we obtain the corollary:

**Corollary 4.6.** *Let  $\mathbb{G}_1, \dots, \mathbb{G}_n$  be digraphs and suppose that for all  $j \in \{1, \dots, n\}$  there exists a regular morphism  $i_j : \mathbb{G} \rightarrow \mathbb{G}_j$ . If all the cohomology groups  $H_\mu^*(\mathbb{G}; \mathbb{K})$  and  $H_\mu^*(\mathbb{G}_j; \mathbb{K})$  vanish for all  $j$ , then*

$$H^*(\nabla_{P(\mathbb{G})}(P(\mathbb{G}_1), \dots, P(\mathbb{G}_n)); \mathbb{K}) = 0.$$



In the next subsection, we apply the results shown in this subsection to obtain vanishing criteria for multipath cohomology.

**4.3. Acyclicity Criteria and Remarkable Examples.** The aim of this subsection is to find sufficient conditions on a graph  $G$ , for  $H_\mu^*(G; \mathbb{K})$  to be trivial. That is, we want to establish some acyclicity criteria for multipath cohomology – cf. Criterion A and Criterion B.

A first vanishing criterion is a direct consequence of Corollary 4.6. We first recall some notation from Theorem 3.13. Let  $v \in V(G)$ , and consider the set  $E_v = \{e_1, \dots, e_n\} \subset E(G)$  of edges incident to  $v$ . We order  $E_v$  in such a way that  $v = t(e_i)$ , for  $i = 1, \dots, k$ , and  $v = s(e_j)$ , for  $j = k + 1, \dots, n$ . Denote by  $G_v^k$  the graph obtained by deleting from  $G$  the edges  $e_1, \dots, e_k$ , and set  $G_v^{(h)} := G_v^k \cup e_h$  for  $h \leq k$ .

**Criterion A.** Assume that a digraph  $G$  satisfies the following conditions:

- (1) there exists a vertex  $v$  that is the target (or source) of  $n > 2$  edges;
- (2) all subgraphs  $G_v^{(1)}, \dots, G_v^{(n)}, G_v^k$  of  $G$  have trivial cohomology.

Then,  $H_\mu^*(G; \mathbb{K}) = 0$ .

*Proof.* By Theorem 3.13, we have the isomorphism of posets

$$P(G) \cong \nabla_{P(G_v^k)} (P(G_v^{(1)}), \dots, P(G_v^{(k)})) .$$

Then, the statement follows from Corollary 4.6.  $\square$

At a first glance Criterion A looks quite technical. However, it can easily be applied in practice.

*Example 4.7.* Let  $G$  be the graph shown in Figure 7. Recall from Example 2.29 that a coherent linear graph has trivial cohomology. Then, by applying Criterion A, we get that  $H_\mu^*(G; \mathbb{K}) = 0$ .



FIGURE 7. A graph with  $n$  edges with target  $v$  glued to a linear graph.

Criterion A says that we can infer a vanishing result for multipath cohomology by looking at smaller pieces in the graph. The remaining criteria have a slightly different flavour; they are based on the existence of suitably “embedded” subgraphs. To formalise this statement we need the notion of  $\nu$ -equivalence.

**Definition 4.8.** A morphism of directed graphs  $\phi: G \rightarrow G'$  is a  $\nu$ -equivalence away from a (possibly empty) set of vertices  $V \in V(G)$ , if the valence of  $v \in V(G)$  is the same as the valence of  $\phi(v) \in V(G')$ , for every  $v \in V(G) \setminus V$ .

*Remark 4.9.* A  $\nu$ -equivalence  $\phi: G \rightarrow G'$  away from an empty set of vertices, is just the inclusion of  $G$  as a connected component of  $G'$ .

*Example 4.10.* Let  $G$  be the graph in Figure 8, and denote by  $I_3$  the linear digraph illustrated in Figure 2, with vertices labelled as in the aforementioned figure. Consider the morphisms of digraphs

$$\phi_b, \phi_r: I_3 \rightarrow G,$$

defined as follows:  $\phi_b(v_i) = w_i$ , for each  $i \in \{0, 1, 2, 3\}$ , and

$$\phi_r(v_i) = \begin{cases} w_1 & \text{if } i = 0; \\ z_i & \text{if } i = 1, 2; \\ w_2 & \text{if } i = 3. \end{cases}$$

Then,  $\phi_b$  is a  $\nu$ -equivalence away from  $v_1$  and  $v_2$ , while  $\phi_r$  is a  $\nu$ -equivalence away from  $v_0$  and  $v_3$ .




FIGURE 8. A graph  $G$  and two copies of  $I_3$  inside it. The morphism of digraphs  $\phi_r$ , whose image is the red copy of  $I_3$ , is a  $\nu$ -equivalence away from  $v_0$  and  $v_3$ , while morphism of digraphs  $\phi_b$ , whose image is the blue copy of  $I_3$ , is a  $\nu$ -equivalence away from  $v_1$  and  $v_2$ .

In order to state our next criterion we need a new family of graphs  $H_{n,m}$ , illustrated in Figure 9.




FIGURE 9. The graph  $H_{n,m}$ .

**Criterion B.** Let  $G$  be a digraph and assume that there exists a  $\nu$ -equivalence  $\phi: H_{n,m} \rightarrow G$  away from  $w_1, \dots, w_n$  and  $x_1, \dots, x_m$ . If  $(\phi(v_0), \phi(v_1))$  is not contained in any coherently oriented cycle<sup>3</sup> of  $G$ , then  $H_\mu^*(G; \mathbb{K}) = 0$ .

*Proof.* Denote by  $e$  the edge  $(\phi(v_0), \phi(v_1))$  of  $\phi(H_{n,m})$  and set  $G' := G \setminus \{e\}$ . We want to show that the poset  $P(G)$  is isomorphic to the cone  $\mathfrak{Cone} P(G')$ . Consider the following subset of the path poset:  $P(G)_0 := \{H \in P(G) \mid e \notin H\}$  and  $P(G)_1 := \{H \in P(G) \mid e \in H\}$  endowed with the poset structure induced by  $P(G)$ . We claim that  $P(G)_i \cong P(G') \times \{i\} \subset \mathfrak{Cone} P(G')$ , for  $i = 0, 1$ . The isomorphism  $P(G)_0 \cong P(G')$  is clear. Now, to identify  $P(G)_1$  with  $P(G') \times \{1\}$  we observe that:

- (1) the edge  $e$  is coherently oriented with the other edges in  $\phi(H_{n,m})$ ,
- (2) no multipath contains two edges of the form  $(\phi(x_i), \phi(v_1))$  nor contains two edges of the form  $(\phi(v_1), \phi(w_i))$ ,
- (3) and  $e$  is not contained in a coherently oriented cycle.

It follows that  $H \cup e$  is a multipath for every  $H \in P(G)_0$ , which implies that the map sending  $H \in P(G)_0$  to  $H \cup e \in P(G)_1$  is well-defined. Note that this map is also a bijection, and that it preserves the inclusions, i.e. if  $H \prec H' \in P(G)_0$  then  $H \cup e \prec H' \cup e \in P(G)_1$ . Consequently, we have a sequence of isomorphism of posets  $P(G)_1 \cong P(G)_0 \cong P(G') \times \{0\} \cong P(G') \times \{1\}$ . To complete the proof that  $P(G) \cong \mathfrak{Cone} P(G')$  we have to check that, under the above chain of identifications, a covering relation between two elements  $H \in P(G)_0$  and  $H' \in P(G)_1$  corresponds uniquely to a covering relation between the corresponding elements  $(H, 0) \in P(G') \times \{0\}$  and  $(H' \setminus \{e\}, 1) \in P(G') \times \{1\}$ . This follows directly from Remark 3.9 and, as a consequence, the posets  $P(G)$  and  $\mathfrak{Cone} P(G')$  are isomorphic. From Theorem 4.2, it follows  $H_\mu^*(G; \mathbb{K}) = 0$ .  $\square$

<sup>3</sup>That is the image of a regular morphism  $P_n \rightarrow G$ , for some  $n$ .

*Remark 4.11.* Criterion B also holds if either  $n = 0$  or  $m = 0$ . In these cases, we say that the graph  $G$  has a *coherent tail*. With this terminology, we can restate the special case of Criterion B when either  $n = 0$  or  $m = 0$  as follows; if  $G$  has a coherent tail, then  $H_\mu^*(G; \mathbb{K}) = 0$ .

We apply the vanishing criteria to compute  $H_\mu^*(-; \mathbb{K})$  in some interesting cases.

*Example 4.12.* An *arborescent graph* (or *arborescence*) is a directed graph in which there is a vertex  $r$ , called *root*, and there is exactly one directed path from  $r$  to any other vertex. If an arborescent graph  $T$  has a vertex at distance 2 from the root (i.e. the unique path joining them has length 2), then up to orientation reversing of the edges there is a  $\nu$ -equivalence  $H_{0,m} \rightarrow T$  away from  $v_0$ , for some  $m > 0$ . Applying Criterion B, we get  $H_\mu^*(T; \mathbb{K}) = 0$ .




FIGURE 10. The graph  $O_n$ .

*Example 4.13.* Let  $O_n$  be the graph in Figure 10, and set  $v = v_3$ ,  $e_1 = (v_3, v_2)$ , and  $e_2 = (v_3, v_1)$ . Using the same notation of Criterion A, we have that  $G_v^2$  and  $G_v^{(1)}$  have a coherent tail, while there is a  $\nu$ -equivalence  $I_3(\cong H_{1,1}) \rightarrow G_v^{(2)}$  whose image is the sub-graph with edges  $(v_3, v_1)$ ,  $(v_1, v_0)$ ,  $(v_0, v_2)$ . Thus, by Criterion B,  $G_v^2$ ,  $G_v^{(1)}$ , and  $G_v^{(2)}$  have trivial cohomology. By Criterion A, it follows that  $H_\mu^*(O_n; \mathbb{K}) = 0$ .

Another class of graphs, important to us, is given by the dandelion graphs:

**Definition 4.14.** Let  $D_{n,m}$  the graph on  $(n + m + 1)$  vertices, and  $(m + n)$  edges defined as follows:

- (1)  $V(D_{n,m}) = \{v_0, w_1, \dots, w_n, x_1, \dots, x_m\}$ ;
- (2)  $E(D_{n,m}) = \{(w_i, v_0), (v_0, x_j) \mid i = 1, \dots, n; j = 1, \dots, m\}$ .

The digraph  $D_{n,m}$  is called a *dandelion graph*.




FIGURE 11. The graph  $D_{3,2}$ .

In other words we have a single  $(m + n)$ -valent vertex  $v_0$ , all remaining vertices are univalent, there are  $n$  edges with target  $v_0$ , and there are  $m$  edges with source  $v_0$  – cf. Figure 11.

*Remark 4.15.* If we reverse the orientation of all the edges in  $D_{k,n-k}$  we obtain  $D_{n-k,k}$ . Then, we have an isomorphism  $P(D_{k,n-k}) \cong P(D_{n-k,k})$ ; hence,  $H_\mu^n(D_{k,n-k}; \mathbb{K}) \cong H_\mu^n(D_{n-k,k}; \mathbb{K})$  for all  $n$ .

*Example 4.16.* The dandelion digraph  $D_{n,0}$  is a source with  $n$  edges and the dandelion digraph  $D_{0,n}$  is a sink with  $n$  edges – cf. Definition 3.10. The dandelion digraph  $D_{1,1}$  is the 2-step graph  $I_2$ .

We proceed by proving some interesting properties of the dandelion graphs.

*Remark 4.17.* If  $G$  is a source or a sink with  $n \geq 2$  edges, then  $\dim H_\mu^1(G; \mathbb{K}) = n - 1$  and  $\dim H_\mu^i(G; \mathbb{K}) = 0$  for  $i \neq 1$ ; in fact the path poset of a sink (or a source) with  $n$  edges is given by a single multipath of length 0, and  $n$  multipaths of length 1. It follows that, in the case at hand, the multipath chain complex is very simple:

$$0 \rightarrow C_\mu^0(D_{0,n}; \mathbb{K}) \cong \mathbb{K} \xrightarrow{d^0} C_\mu^1(D_{0,n}; \mathbb{K}) \cong \mathbb{K}^n \rightarrow 0.$$

Furthermore, the map  $d^0$  is injective (since  $d^0$  is the map  $x \mapsto (\pm x, \dots, \pm x)$  for an appropriate choice of signs), giving trivial cohomology in degree 0 and a cohomology group of dimension  $n - 1$  in degree 1.

An immediate consequence of Remark 4.17 is that we can have multipath cohomology groups of arbitrary dimension (as  $\mathbb{K}$ -vector space).

**Proposition 4.18.** *Let  $n \geq 1$  be an integer. Then,  $H_\mu^*(D_{1,n-1}; \mathbb{K}) = 0$ .*

*Proof.* Note that  $D_{1,0} = I_1$ , and that  $D_{1,n-1} = H_{0,n-1}$ . In the former case, the cohomology is trivial by direct computation. In the latter case, the cohomology is trivial by Criterion A, as  $D_{1,n-1}$  has a coherent tail – cf. Remark 4.11.  $\square$

**Proposition 4.19.** *Let  $n > 2$  and  $k > 0$  be two integers such that  $n > k$ . Then, we have*

$$H_\mu^i(D_{k,n-k}; \mathbb{K}) \cong \begin{cases} \mathbb{K}^{(k-1)(n-k-1)} & \text{if } i = 2 \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* We proceed by induction on  $n$ . If  $n = 3$ , Proposition 4.18 and Remark 4.15 imply  $H_\mu^*(D_{2,1}; \mathbb{K}) \cong H_\mu^*(D_{1,2}; \mathbb{K}) = 0$ . Now, set  $G = D_{k,n-k}$  and  $v = v_0$ . By Theorem 3.13, we have

$$P(D_{k,n-k}) \cong \nabla_{P(D_{0,n-k})} \underbrace{(P(D_{1,n-k}), \dots, P(D_{1,n-k}))}_{k \text{ copies}} \stackrel{(*)}{\cong} \nabla_{P(D_{0,n-k})} (P(D_{1,n-k}), P(D_{k-1,n-k}))$$

where the isomorphism marked with  $(*)$  follows from the associativity of the gluing (and from Theorem 3.13). By applying Theorem 4.4 and the inductive hypothesis, it follows that  $H_\mu^i(D_{k,n-k}; \mathbb{K}) = 0$  for  $i \geq 3$ , and that the sequence

$$0 \rightarrow H_\mu^1(D_{0,n-k}; \mathbb{K}) \rightarrow H_\mu^2(D_{k,n-k}; \mathbb{K}) \rightarrow H_\mu^2(D_{k-1,n-k}; \mathbb{K}) \rightarrow 0,$$

is exact. The assertion is now immediate from the fact that the dimension function is additive on short exact sequences.  $\square$

We conclude the section, showing that there exist directed graphs with multipath cohomology of arbitrary rank in arbitrary degree.

**Lemma 4.20.** *Given a digraph  $G'$  with a vertex  $w$  of valence 1, there exists a digraph  $G$  such that  $H_\mu^*(G'; \mathbb{K}) \cong H_\mu^{*-1}(G; \mathbb{K})$ .*

*Proof.* Let  $e \in E(G')$  be the only edge incident to  $w$ . We define  $G$  as follows; if  $s(e) = w$ , glue a linear sink over  $w$  to  $G'$  – cf. Figure 5, otherwise glue a linear source.

In the notations of Figures 12 and 13, by Theorem 3.13 the path poset of  $G$  is the gluing of the path posets  $P(G_w^{(1)})$  and  $P(G_w^{(2)})$  over  $P(G')$ . By Remark 4.11 the subgraphs  $G_w^{(1)}$  and  $G_w^{(2)}$  have trivial cohomology; hence, by Remark 4.5, we have  $H_\mu^*(G; \mathbb{K}) = H_\mu^{*-1}(G'; \mathbb{K})$ .  $\square$

Observe that the digraph  $G$  constructed in Lemma 4.20 has again (at least) one vertex of valence 1 and consequently the construction can be iterated.

**Proposition 4.21.** *For all  $i, n \in \mathbb{N}$ , there exists a digraph  $G$  such that  $\dim_{\mathbb{K}} (H_\mu^i(G; \mathbb{K})) = n$ .*




FIGURE 12. Configurations of a digraph  $G$  with subgraphs  $\nu$  equivalent away from  $v_0$  to  $D_{1,2}$  and  $D_{2,1}$ .




FIGURE 13. The graph  $G'$  as subgraph of  $G$  in the two cases we consider.

*Proof.* The multipath cohomology of a sink graph  $G$  with  $n + 1$  edges is concentrated in degree one, where it is  $H_\mu^1(G; \mathbb{K}) \cong \mathbb{K}^n$ . By applying iteratively Lemma 4.20, we obtain digraphs  $G$  with  $\dim_{\mathbb{K}} (H_\mu^i(G; \mathbb{K})) = n$  for every  $i$ .  $\square$

## 5. ORIENTED LINEAR GRAPHS

This section is devoted to the study of the multipath cohomology of oriented linear graphs. Firstly, we focus on the case of coefficients in a field  $\mathbb{K}$ . In this case, we achieve a complete description of their cohomology groups. Then, we analyse the general case of coefficients in a graded algebra  $A$ , and we prove some recursive formulae for the graded Euler characteristic.

**5.1. Multipath cohomology of linear graphs.** We start with the definition of oriented linear graphs and alternating graphs.

**Definition 5.1.** An *oriented linear graph*  $L$  (on  $n$  vertices) is a directed graph with vertices  $\{v_0, \dots, v_{n-1}\}$ , such that, for all  $i \in \{1, \dots, n - 1\}$ , exactly one among  $(v_i, v_{i-1})$  and  $(v_{i-1}, v_i)$  belongs to  $E(L)$ , and there are no other edges. An oriented linear graph  $L$  is called *alternating* if whenever  $(v_{i-1}, v_i) \in E(L)$  for some  $i < n - 1$ , we have  $(v_{i+1}, v_i) \in E(L)$  and, analogously, if  $(v_i, v_{i-1}) \in E(L)$  then  $(v_i, v_{i+1}) \in E(L)$ . We denote by  $A_n$  an alternating linear graph on  $n + 1$  vertices.

Observe that the alternating graph  $A_n$  is unique up to orientation reversing.

**Definition 5.2.** A vertex of an oriented linear graph  $L$  is called *unstable* if it is both a source and a target, and *stable* otherwise. We denote by  $SV(L)$  the set of stable vertices of  $L$ .

Our aim is to show that the cohomology of an oriented linear graph  $L$  is related to the number of stable vertices in  $L$ , and their relative distance.

**Definition 5.3.** Let  $v_i, v_j$  be vertices of an oriented linear graph  $L$ . The distance  $d(v_i, v_j)$  is the length of the unique simple path, if it exists, between them, and it is set to  $-\infty$  otherwise.

For an oriented linear graph  $L$  the property  $D(k)$  is defined as follows:

$$(6) \quad D(k) : \quad \forall v, w \in SV(L), \quad d(v, w) \leq k .$$




FIGURE 14. The graphs  $G$ ,  $G'$ ,  $G''$ , and the relative orientations of  $e_1$  and  $e_2$ .

A disjoint union of oriented linear graphs satisfies the property  $D(k)$  if each component does. Observe that the set of oriented linear graphs is filtered by the above property; each linear graph satisfies  $D(k)$  for some  $k$ , and if  $L$  satisfies  $D(k)$ , then it also satisfies  $D(k+1)$ . Furthermore, the alternating graphs satisfy the property  $D(1)$ , and they are the only connected graphs satisfying it. The graph  $I_n$  satisfies the property  $D(n)$ .

*Remark 5.4.* If an oriented linear graph  $L$  satisfies the property  $D(n)$  for  $n > 2$ , then  $L$  has trivial cohomology. In fact, if there exists a pair of stable vertices at distance greater than 2, then there exists a  $\nu$ -equivalence  $f: I_3 \rightarrow L$  away from  $v_0, v_3 \in V(I_3)$ ; by Criterion B,  $H_\mu^*(L, \mathbb{K}) = 0$ .

*Remark 5.5.* If  $L$  satisfies the property  $D(n)$  then each of its subgraphs also satisfies  $D(n)$ .

By Remark 5.4, a complete description of the multipath cohomology of oriented linear graphs can be achieved by studying graphs satisfying the property  $D(2)$ . As a first step, we start with oriented linear graphs satisfying  $D(1)$ . Since the cohomology of a disjoint union of linear graphs is the tensor product over its components (cf. Remark 3.2), we restrict to the case of connected ones, i.e. the alternating graphs.

**Theorem 5.6.** *Let  $A_n$  be an alternating graph. Then, we have the following isomorphisms*

$$(7) \quad H_\mu^*(A_n, \mathbb{K}) \cong \begin{cases} H_\mu^*(A_{n-1}, \mathbb{K}) & \text{if } n \equiv 0 \pmod{3}, \\ 0 & \text{if } n \equiv 1 \pmod{3}, \\ H_\mu^{*-1}(A_{n-2}, \mathbb{K}) & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

depending on the congruence class of  $n$  modulo 3.

*Proof.* We use the notation illustrated in Figure 14, with  $G = A_n$ . By Theorem 3.13, the path poset  $P(A_n)$  is isomorphic to  $\nabla_{P(G'')}(P(G \setminus \{e_2\}), P(G'))$ . Observe that  $P(G \setminus \{e_2\})$  is a cone over  $P(G'')$  and that we have isomorphisms of graphs  $G' \cong A_{n-1}$  and  $G'' \cong A_{n-2}$ . Consequently, there is an induced isomorphism

$$P(A_n) = P(G) \cong \nabla_{P(A_{n-2})}(\mathbf{Cone} P(A_{n-2}), P(A_{n-1}))$$

of posets. Since the cohomology groups  $H_\mu^*(\mathbf{Cone} P(A_{n-2}); \mathbb{K})$  are all trivial, from Theorem 4.4 (applied to the gluing of  $\mathbf{Cone} P(A_{n-2})$  and  $P(A_{n-1})$ ) we obtain the following exact sequence:

$$\cdots \rightarrow H_\mu^i(P(A_{n-1}); \mathbb{K}) \rightarrow H_\mu^i(P(A_{n-2}); \mathbb{K}) \rightarrow H_\mu^{i+1}(P(A_n), \mathbb{K}) \rightarrow H_\mu^{i+1}(P(A_{n-1}); \mathbb{K}) \rightarrow \cdots$$

A direct computation shows that the cohomology of the graphs  $A_n$  for  $n < 5$  agrees with the isomorphisms in Equation (7). The assertion now follows by an induction argument.  $\square$

As a consequence, it is possible to obtain a precise description of the ranks of cohomology groups for alternating graphs:

**Corollary 5.7.** *Let  $A_n$  be an alternating graph. Then:*

$$\dim_{\mathbb{K}} H_{\mu}^k(A_n; \mathbb{K}) = \begin{cases} 1 & \text{if } n = 3(k-1) + 2 \text{ or } n = 3k, \\ 0 & \text{otherwise.} \end{cases}$$

Before proceeding with our analysis of the cohomology of oriented linear graphs, we need the following definition.

**Definition 5.8.** Given a oriented linear graph  $L$ , its *reduction*  $\text{Red } L$  is the (possibly disconnected) spanning subgraph of  $L$ , obtained as follows; for each maximal simple path on (the ordered set of) vertices  $\{v_h, \dots, v_{h+m}\}$ , delete all edges, but the one between  $v_{h+m-1}$  and  $v_{h+m}$ .

Note that if a maximal simple path is an edge of  $L$ , then it is still an edge of  $\text{Red } L$ .

*Example 5.9.* The reduction of  $I_n$  is the spanning subgraph of  $I_n$  with only edge  $(v_{n-1}, v_n)$ . The reduction  $\text{Red } L$  is isomorphic to  $L$  if, and only if,  $L$  is alternating.

Observe that, by construction, the digraph  $\text{Red } L$  is the disjoint union of  $h$  connected components, where  $h-1$  is the number of edges deleted during the process of reduction.

We can linearly order the connected components of  $\text{Red } L$  according to their minimal-index vertex. Denote by  $C_i$  the  $i$ -th component with respect to this order. Notice that  $\text{Red } L$  satisfies  $D(1)$ , thence for each  $i \in \{1, \dots, h\}$  there is a  $k_i \geq 0$  such that  $C_i \cong A_{k_i}$ .

**Lemma 5.10.** *Let  $L$  be an oriented linear graph satisfying the property  $D(2)$ , and let  $A_{k_1}, \dots, A_{k_h}$  be the connected components of  $\text{Red } L$ . If  $k_h \equiv 1 \pmod{3}$ , then  $H_{\mu}^*(L; \mathbb{K}) = 0$ .*

*Proof.* Note that if  $L$  satisfies  $D(1)$ , then it is an alternating graph, and the statement follows by Proposition 5.6. Suppose  $L$  satisfies  $D(2)$ , but not  $D(1)$  and denote by  $w_0, \dots, w_{k_h}$  the vertices of  $A_{k_h}$ . First, observe that, by Definition 5.8, if  $k_h = 1$  or if  $|E(L)| = k_h + 1$ , then the linear graph  $L$  has a coherent tail. Thus,  $H_{\mu}^*(L; \mathbb{K}) = 0$  by Remark 4.11.

In all the other cases, up to orientation reversing, the graph  $L$  contains a subgraph as in Figure 15. We can apply Theorem 3.13 choosing the vertex  $x$  illustrated in Figure 15, and



FIGURE 15. The sub-graph  $A_{k_h}$  inside  $L$ . The edge  $(w_{k_h-1}, w_{k_h})$  can be oriented either way depending on the parity of  $k_h$ .

obtain the isomorphism

$$P(L) \cong \nabla_{P(L''')} (P(L'), P(L'')) ,$$

where  $L' = L \setminus \{(w_0, x)\}$ ,  $L'' = L \setminus \{(v, x)\}$  and  $L''' = L \setminus \{(w_0, x), (v, x)\}$ . Now, Remark 4.11 implies  $H_{\mu}^*(L''; \mathbb{K}) = 0$ . Furthermore,  $H_{\mu}^*(L'; \mathbb{K}) = H_{\mu}^*(L'''; \mathbb{K}) = 0$ , since  $L'$  and  $L'''$  have  $A_{k_h}$  as a connected component – cf. Remark 3.2 and Proposition 5.6; in fact,  $w_{k_h}$  is univalent, since  $k_h \neq 0$ . The statement now follows from Theorem 4.4.  $\square$

Denote by  $\lfloor x \rfloor$  the integer part of  $x$ .

**Theorem 5.11.** *Let  $L$  be an oriented linear graph satisfying the property  $D(2)$ , but not  $D(1)$ . Denote by  $A_{k_1}, \dots, A_{k_h}$  the connected components of  $\text{Red } L$ . Then:*

- (1) *if there exists an index  $j \in \{1, \dots, h-1\}$  such that  $k_j \equiv 0 \pmod{3}$ , then  $H_{\mu}^*(L, \mathbb{K}) = 0$ ;*



(2) otherwise, the cohomology groups of  $L$  decompose as

$$H_{\mu}^{*+h-1}(L; \mathbb{K}) = H_{\mu}^*(A_{3\lfloor k_1/3 \rfloor}; \mathbb{K}) \otimes \cdots \otimes H_{\mu}^*(A_{3\lfloor k_{h-1}/3 \rfloor}; \mathbb{K}) \otimes H_{\mu}^*(A_{k_h}; \mathbb{K}),$$

where  $\otimes$  here denotes the graded tensor product over  $\mathbb{K}$ .

*Proof.* We linearly order the edges  $e_1, \dots, e_{h-1}$  in  $E(L) \setminus E(\text{Red } L)$ , i.e. the edges in the complement of the components  $A_{k_1}, \dots, A_{k_h}$ , according to their minimal-index vertex. The addition of  $e_i$  to  $\text{Red } L$  merges the component  $A_{k_i}$  with the component  $A_{k_{i+1}}$ .

(1) To prove the first item, observe that if  $k_j \equiv 0 \pmod{3}$ , then either  $L$  has a coherent tail (when  $k_1 = 0$ ) or, up to orientation reversing, it contains a subgraph as in Figure 16, where the vertices  $w_0, \dots, w_3$  are in  $A_{k_j}$ . Now, if  $v$  is an univalent vertex of the graph  $L$ ,



FIGURE 16. A possible sub-graph of  $L$ .

then  $L$  has a coherent tail of length two and, again, the cohomology groups  $H_{\mu}^*(L, \mathbb{K})$  are trivial. In all remaining cases,  $L$  contains a subgraph as in Figure 17. We are in the




FIGURE 17. Another possible sub-graph of  $L$ .

hypothesis of Theorem 3.13, choosing the vertex  $v$ , to decompose  $P(L)$  as

$$P(L) \cong \nabla_{P(L''')} (P(L'), P(L'')) ,$$

where  $L' = L \setminus \{(v, x)\}$ ,  $L'' = L \setminus \{(v, y)\}$ , and  $L''' = L \setminus \{(v, x), (v, y)\}$ .

The first assertion follows from Criterion A:  $H_{\mu}^*(L''; \mathbb{K}) = 0$  because  $L''$  contains a coherent tail, and the cohomologies  $H_{\mu}^*(L'; \mathbb{K})$  and  $H_{\mu}^*(L'''; \mathbb{K})$  are both trivial in virtue of Lemma 5.10 and Remark 3.2.

(2) The proof of second statement proceeds by induction. We are in the case  $k_j \not\equiv 0 \pmod{3}$  for  $j \in \{1, \dots, h-1\}$ .

The graph obtained from  $L$  by deleting  $e_1$  has two connected components  $L_1 = A_{k_1}$  and  $L_2$ . Observe that  $L_1$  and  $L_2$  are oriented linear graphs satisfying the property  $D(2)$  – cf. Remark 5.5. Since  $k_1 \neq 0$ , we have that  $L_1$  has at least one edge. Furthermore, by definition of the reduction, we can assume that  $e_1$  is contained in a linear subgraph of the form  $\bullet \xrightarrow{e} \bullet \xleftarrow{e_1} \bullet$  or  $\bullet \xleftarrow{e} \bullet \xrightarrow{e_1} \bullet$  with  $e \in E(L_1)$ . By Theorem 3.13, we have the decomposition

$$P(L) \cong \nabla_{P((L_1 \sqcup L_2) \setminus \{e\})} (P(L_1 \sqcup L_2), P(L \setminus \{e\})) .$$

Observe now that  $P(L \setminus \{e\}) \cong \mathbf{Cone} P((L_1 \sqcup L_2) \setminus \{e\})$ , hence

$$P(L) \cong \nabla_{P(A_{k_1-1} \sqcup L_2)} (P(A_{k_1} \sqcup L_2), \mathbf{Cone} P(A_{k_1-1} \sqcup L_2)) .$$

Using Theorem 4.4, we get the following exact sequence

$$(8) \quad \cdots \rightarrow H_{\mu}^i(A_{k_1-1} \sqcup L_2; \mathbb{K}) \rightarrow H_{\mu}^{i+1}(L; \mathbb{K}) \rightarrow H_{\mu}^{i+1}(A_{k_1} \sqcup L_2; \mathbb{K}) \rightarrow H_{\mu}^{i+1}(A_{k_1-1} \sqcup L_2; \mathbb{K}) \rightarrow \cdots$$

In the following, set first  $k_1 = 3j + 1$ . By Corollary 5.7 and Remark 3.2, the cohomology group  $H_{\mu}^i(A_{k_1} \sqcup L_2; \mathbb{K})$  is trivial. Analogously, the group  $H_{\mu}^i(A_{k_1-1} \sqcup L_2; \mathbb{K})$  is isomorphic to the product  $H_{\mu}^j(A_{k_1-1}; \mathbb{K}) \otimes H_{\mu}^{i-j}(L_2; \mathbb{K})$ . In the case  $k_1 = 3j + 2$  instead, the group  $H_{\mu}^i(A_{k_1} \sqcup L_2; \mathbb{K})$  is isomorphic to the product  $H_{\mu}^{j+1}(A_{k_1}; \mathbb{K}) \otimes H_{\mu}^{i-j-1}(L_2; \mathbb{K})$ , and  $H_{\mu}^i(A_{k_1-1} \sqcup L_2; \mathbb{K})$  is trivial.



As a consequence, using the Mayer Vietoris sequence in Equation (8), we have

$$H_\mu^{i+1}(\mathbb{L}; \mathbb{K}) \cong \begin{cases} H_\mu^j(\mathbf{A}_{k_1-1}; \mathbb{K}) \otimes H_\mu^{i-j}(\mathbb{L}_2; \mathbb{K}) & \text{if } k_1 = 3j + 1, \\ H_\mu^{j+1}(\mathbf{A}_{k_1}; \mathbb{K}) \otimes H_\mu^{i-j}(\mathbb{L}_2; \mathbb{K}) & \text{if } k_1 = 3j + 2 \end{cases}$$

If  $k_1 = 3j + 1$ , we can rewrite the first isomorphism, using that the multipath cohomology of  $\mathbf{A}_{k_1-1} = \mathbf{A}_{3j}$  is concentrated in cohomological degree  $j$ , as follows:

$$H_\mu^{i+1}(\mathbb{L}; \mathbb{K}) \cong (H_\mu^j(\mathbf{A}_{k_1-1}, \mathbb{K}) \otimes H_\mu^{i-j}(\mathbb{L}_2, \mathbb{K})) \cong \bigoplus_{r+s=i} (H_\mu^r(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^s(\mathbb{L}_2, \mathbb{K}))$$

On the other hand, if  $k_1 = 3j + 2$ , by Theorem 5.6 and Corollary 5.7, we obtain the isomorphism

$$H_\mu^i(\mathbb{L}; \mathbb{K}) \cong H_\mu^{j+1}(\mathbf{A}_{k_1}, \mathbb{K}) \otimes H_\mu^{i-j-1}(\mathbb{L}_2, \mathbb{K}) \cong H_\mu^j(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^{i-j-1}(\mathbb{L}_2, \mathbb{K}).$$

Finally, we have also that

$$H_\mu^j(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^{i-j-1}(\mathbb{L}_2, \mathbb{K}) \cong \bigoplus_{r+s=i-1} (H_\mu^r(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^s(\mathbb{L}_2, \mathbb{K}))$$

since the multipath cohomology of  $\mathbf{A}_{k_1-2} = \mathbf{A}_{3j}$  is concentrated in degree  $j$ . We have shown that, in either case, we have

$$H_\mu^{*+1}(\mathbb{L}; \mathbb{K}) \cong \bigoplus_{r+s=*} (H_\mu^r(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^s(\mathbb{L}_2, \mathbb{K})) = (H_\mu^*(\mathbf{A}_{3j}, \mathbb{K}) \otimes H_\mu^*(\mathbb{L}_2, \mathbb{K})).$$

The statement now follows by induction. □

**5.2. Graded characteristic of linear graphs.** We now analyse the cohomology of linear graphs from a different perspective; instead of considering multipath cohomology with coefficient in a field, we fix a principal ideal domain  $R$  as a base ring, and take coefficients in a unital  $R$ -algebra  $A$ . We are interested in analysing the (graded) Euler characteristic in the case where  $A$  is graded. Firstly, let us prove a general result.

**Theorem 5.12.** *Let  $G$ ,  $G'$ , and  $G''$  be three digraphs as illustrated in Figure 14, and let  $A$  be a unital  $R$ -algebra. Then, one of the following holds:*

- (1) *if the edges  $e_1$  and  $e_2$  are coherently oriented, as in Figure 14d, then the sequence*

$$0 \rightarrow C_\mu^{*-1}(G'; A) \longrightarrow C_\mu^*(G; A) \longrightarrow C_\mu^*(G'; A) \otimes A \rightarrow 0$$

*is exact;*

- (2) *if the edges  $e_1$  and  $e_2$  are not coherently oriented, see Figure 14e, then the sequence*

$$0 \rightarrow C_\mu^{*-1}(G''; A) \otimes A \longrightarrow C_\mu^*(G; A) \longrightarrow C_\mu^*(G'; A) \otimes A \rightarrow 0$$

*is exact;*

*where all tensor products are over  $R$ .*

*Proof.* Before dwelling into the details of each case, we first discuss the general picture. Denote for simplicity by  $P$ ,  $P'$ , and  $P''$  the path posets of  $G$ ,  $G'$ , and  $G''$ , respectively. We have an embedding of  $G'$  into  $G$  that induces (as a spanning subgraph, cf. Figure 14c) an injective morphism of poset  $\iota: P' \rightarrow P$ . We remark that  $\iota(P')$  is a downward closed faithful subposet of  $P$ , and that the minimal length of an element in  $P \setminus \iota(P')$  is 1. By [CCDT21, Proposition 6.12], we have the following sequence of cochain complexes

$$0 \rightarrow C_{\mathcal{F}_{A,A}}^{*-1}(P \setminus \iota(P')) \longrightarrow C_{\mathcal{F}_{A,A}}^*(P) \longrightarrow C_{\mathcal{F}_{A,A}}^*(P') \otimes A \rightarrow 0$$

which is exact. By definition, we have that  $C_{\mathcal{F}_{A,A}}^*(P) = C_\mu^*(G; A)$  and  $C_{\mathcal{F}_{A,A}}^*(P') = C_\mu^*(G'; A)$ .

To conclude it is enough to identify the complex  $C_{\mathcal{F}_{A,A}}^*(P \setminus \iota(P'))$ . Observe that the elements of  $P \setminus \iota(P')$  are precisely the multipaths in  $G$  containing  $e_1$ . The proof splits now in two cases;

- (1) if  $e_1$  and  $e_2$  are coherently oriented, then  $H \cup \{e_1\} \in P \setminus \iota(P')$  for each  $H \in \iota(P')$ . Thus, we have an order-preserving bijection between  $P(G')$  and  $P \setminus \iota(P')$ , which also preserves the number of connected components. Therefore, we obtain the isomorphism

$$C_{\mathcal{F}_{A,A}}^*(P \setminus \iota(P')) \cong C_{\mathcal{F}_{A,A}}^*(P') = C_{\mu}^*(G'; A)$$

of cochain complexes;

- (2) if  $e_1$  and  $e_2$  are not coherently oriented, then a multipath which contains  $e_1$  cannot contain  $e_2$ . Thus, we have an identification between multipaths in  $P \setminus \iota(P)$  and multipath in  $G'' \cup \{e_1\}$ , which gives the isomorphisms

$$C_{\mathcal{F}_{A,A}}^*(P \setminus \iota(P')) \cong C_{\mathcal{F}_{A,A}}^*(P'') \otimes A = C_{\mu}^*(G''; A) \otimes A.$$

The tensor factor  $\otimes A$  arises from an extra connected component in each element of the poset  $P \setminus \iota(P')$  with respect to the corresponding element in  $P''$  (namely the edge  $e_1$ ).

The statement is now immediate from the above identifications.  $\square$

The short exact sequence in Theorem 5.12 (1) holds also in a slightly different case;

*Remark 5.13.* Assume that the edges  $e_1$ ,  $e_2$ , and  $e_3$  in  $G$  form a path, and that  $e_2$  is not contained in any coherently oriented cycle in  $G$ , then the sequence in Theorem 5.12 (1) holds. In this case, the role of  $G'$  is played by the graph obtained from  $G$  by contracting the “middle edge”  $e_2$ . Then, we have that

$$P(G) \setminus P(G \setminus \{e_2\}) \cong P(G') \quad \text{and} \quad P(G) \setminus P(G \setminus \{e_2\}) \cong P(G \setminus \{e_2\}),$$

where the first identification is given by contracting  $e_2$ , while the second is given by deleting it. Note that, in the former case, the number of connected components of multipaths is preserved, while in the second case a multipath  $H \in P(G) \setminus P(G \setminus \{e_2\})$  is sent to a multipath with one more connected component. At this point, the same reasoning as in the proof of Theorem 5.12 provides the desired exact sequence.

We can use Theorem 5.12 to reprove a result of Przytycki [Prz10] which computes the cohomology of  $I_n$  (cf. [CCDT21, Corollary 5.7]). We observe that Przytycki obtains this result using the chromatic polynomial as intermediate step, while we prove it directly by induction. Moreover, the following corollary, if  $A$  is commutative, can also be proved as an application of the deletion-contraction exact sequence for the chromatic homology [HGR05, Theorem 3.2].

**Corollary 5.14.** *Let  $I_n$  be the coherently oriented linear graph (cf. Figure 2). For each (unital)  $R$ -algebra  $A$ , we have*

$$H_{\mu}^*(I_n; A) = H_{\mu}^0(I_n; A),$$

and

$$\text{rank}_R(H_{\mu}^*(I_n; A)) = \text{rank}_R(H_{\mu}^0(I_n; A)) = \begin{cases} \text{rank}_R(A)(\text{rank}_R(A) - 1)^n & n \geq 1 \\ \text{rank}_R(A) & n = 0 \end{cases}.$$

*Proof.* The statement is true for  $I_0$  and it is easily proved for  $I_1$ ; in fact, we have

$$0 \rightarrow C_{\mu}^0(I_1; A) = A \otimes A \xrightarrow{d^0} C_{\mu}^1(I_1; A) = A \rightarrow 0,$$

where  $d^0$  is the map  $a \otimes b \mapsto ab$ , which is surjective since  $A$  is unital.

We proceed by induction. Assume that the statement is true for  $n = k - 1$ . Then, we can apply Theorem 5.12 to  $G = I_k$ ,  $G' = I_{k-1}$ , and  $e = (v_0, v_1)$  – cf. Figure 2. From the inductive hypothesis, we obtain the exact sequences

$$0 = H_{\mu}^{i-1}(I_{k-1}; A) \rightarrow H_{\mu}^i(I_k; A) \rightarrow H_{\mu}^i(I_{k-1}; A) = 0, \quad \text{for } i > 1,$$

and

$$(9) \quad 0 \rightarrow H_\mu^0(\mathbb{I}_k; A) \rightarrow H_\mu^0(\mathbb{I}_{k-1}; A) \otimes A \xrightarrow{\delta^*} H_\mu^0(\mathbb{I}_{k-1}; A) \rightarrow H_\mu^1(\mathbb{I}_k; A) \rightarrow 0,$$

where  $\delta^*(x \otimes 1) = x$ . As a consequence we have

- (1)  $H_\mu^i(\mathbb{I}_k; A) = 0$ , for all  $i > 1$ ;
- (2) the exact sequence in Equation (9), since  $\delta^*$  is surjective, splits into the exact sequences

$$0 \rightarrow H_\mu^0(\mathbb{I}_k; A) \rightarrow H_\mu^0(\mathbb{I}_{k-1}; A) \otimes A \xrightarrow{\delta^*} H_\mu^0(\mathbb{I}_{k-1}; A) \rightarrow 0$$

and

$$0 \rightarrow H_\mu^1(\mathbb{I}_k; A) \rightarrow 0.$$

Since the rank is additive on short exact sequences<sup>4</sup>, this concludes the proof.  $\square$

Let  $R$  be a PID, and  $A^* = \bigoplus_{i \in \mathbb{Z}} A^i$  be a finitely generated  $\mathbb{Z}$ -graded  $R$ -algebra.

**Definition 5.15.** The *graded dimension* of  $A$  is the Laurent polynomial

$$\text{qdim}(A^*) := \sum_{i \in \mathbb{Z}} \text{rank}_R(A^i) q^i \in \mathbb{Z}[q, q^{-1}],$$

where  $\text{rank}_R(M)$  indicates the maximal number of non-torsion, linearly independent, elements.

A graded algebra has, by definition, an homogeneous multiplication. As a consequence, the multipath cochain complex inherits from  $A^*$  a second  $\mathbb{Z}$ -grading, which is preserved by the differential. This gives the multipath cohomology the structure of bi-graded cohomology theory. Define the *graded Euler characteristic* of a graph  $G$  (with respect to  $A^*$ ) as

$$\chi_{\text{gr}}(G; A^*) = \sum_{i, j \in \mathbb{Z}} (-1)^i \text{rank}_R(H_\mu^{i, j}(G; A^*)) q^j \in \mathbb{Z}[q, q^{-1}].$$

Note that if we evaluate  $\text{qdim}(A^*)$  (resp.  $\chi_{\text{gr}}$ ) in  $q = 1$ , we obtain the rank of  $A^*$  as  $R$ -module (resp. the usual Euler characteristic  $\chi$ ). It is well-known that the Euler characteristic is additive under exact sequences, and the same holds for the graded Euler characteristic<sup>5</sup>. With the above notation in place, the following corollary is an immediate consequence of Theorem 5.12.

**Corollary 5.16.** *Let  $G$ ,  $G'$ , and  $G''$  be three digraphs as illustrated in Figure 14. Let  $A^*$  be a finitely generated free  $\mathbb{Z}$ -graded  $R$ -algebra with graded dimension  $\alpha = \text{qdim}(A^*)$ .*

- (1) *If the edges  $e_1$  and  $e_2$  are coherently oriented (cf. Figure 14d), then*

$$\chi_{\text{gr}}(G; A^*) = (\alpha - 1) \chi_{\text{gr}}(G'; A^*).$$

- (2) *If the edges  $e_1$  and  $e_2$  are not coherently oriented (cf. Figure 14e), then*

$$\chi_{\text{gr}}(G; A^*) = \alpha (\chi_{\text{gr}}(G'; A^*) - \chi_{\text{gr}}(G''; A^*)).$$

Corollary 5.16 allows us to compute the (graded) characteristic of any oriented linear graph, recursively. We provide, as an example of this process, the graded characteristic of some alternating graphs.

<sup>4</sup>To see this one can tensor for the quotient field, or localise – cf. [BH98, Definition 1.4.2 and Proposition 1.4.5].

<sup>5</sup>For each fixed value  $j$  of the second grading we have a short exact sequence of chain complexes. Notice that  $C_\mu^{i, j}(G; A) \neq 0$  for finitely many values of  $i$  and  $j$ . Thence, we can re-arrange the sum and write

$$\chi_{\text{gr}}(G; A^*) = \sum_{j \in \mathbb{Z}} \chi(C_\mu^{*, j}(G; A)) q^j.$$

Now, additivity follows from the additivity of the (usual) Euler characteristic.

*Example 5.17.* Let  $A_n$  be the alternating graph on  $n$  vertices, with  $n \geq 3$ ; by Corollary 5.16 the graded characteristic can be expressed as:

$$\chi_{\text{gr}}(A_n; A^*) = \alpha (\chi_{\text{gr}}(A_{n-1}; A^*) - \chi_{\text{gr}}(A_{n-2}; A^*)).$$

For  $n = 0$  and  $n = 1$ , the graded characteristic of  $A_n$  are  $\text{qdim}(A^*) =: \alpha$  and  $\alpha(\alpha - 1)$ , respectively. Some further examples are listed in Table 2.

$n$	$\chi_{\text{gr}}(A_n; A^*)$
0	$\alpha$
1	$\alpha(\alpha - 1)$
2	$\alpha^2(\alpha - 2)$
3	$\alpha^2(\alpha^2 - 3\alpha + 1)$
4	$\alpha^3(\alpha - 1)(\alpha - 3)$
5	$\alpha^3(\alpha^3 - 5\alpha^2 + 6\alpha - 1)$
6	$\alpha^4(\alpha - 2)(\alpha^2 - 4\alpha + 2)$
7	$\alpha^4(\alpha - 1)(\alpha^3 - 6\alpha^2 + 9\alpha - 1)$
8	$\alpha^5(\alpha^2 - 5\alpha + 5)(\alpha^2 - 3\alpha + 1)$
9	$\alpha^5(\alpha^5 - 9\alpha^4 + 28\alpha^3 - 35\alpha^2 + 15\alpha - 1)$
10	$\alpha^6(\alpha - 1)(\alpha - 2)(\alpha - 3)(\alpha^2 - 4\alpha + 1)$
11	$\alpha^6(\alpha^6 - 11\alpha^5 + 45\alpha^4 - 84\alpha^3 + 70\alpha^2 - 21\alpha + 1)$

TABLE 2. The graded Euler characteristic of some alternating linear graphs.

The usual Euler characteristic over  $R = \mathbb{K}$  can be obtained by evaluating the graded Euler characteristic in  $\alpha = 1$ . Thus, by Corollary 5.7,  $(\alpha - 1)$  divides  $\chi_{\text{gr}}(A_n; A^*)$  if, and only if,  $n \equiv 1$  modulo 3. Observe that our computations are in perfect accordance with this fact.

Despite not having a closed formula for the graded Euler characteristics of alternating graphs, we can compute its associated generating function  $A(t) = \sum \chi_{\text{gr}}(A_n) t^n \in (\mathbb{Z}[\alpha])[[t]]$  (cf. [Wil06]). In fact, if we denote by  $S_h$  the classical shift for power series

$$S_h \left( \sum_{n=0}^{\infty} c_n t^n \right) = \sum_{n=0}^{\infty} c_{n+h} t^n,$$

from Theorem 5.12 one obtains the relation  $S_2(A(t)) = \alpha(S_1(A(t)) - A(t))$ . It is also immediate to see that

$$S_1(A(t)) = \frac{A(t) - \alpha}{t}.$$

Iterating  $S_1$  and using the fact that  $A(1) = A(0)(\alpha - 1)$  one easily obtains that the ordinary generating function of the graded Euler characteristic of alternating graphs is

$$A(t) = \frac{\alpha(1 - t)}{1 - \alpha t(1 - t)}.$$

Furthermore, in some special cases, we can actually obtain a closed formula.

**Corollary 5.18.** *Let  $A^*$  be a free  $\mathbb{Z}$ -graded  $R$ -algebra with graded dimension  $\alpha \in \mathbb{Z}[q, q^{-1}]$ . If  $I_n$  is the coherently oriented linear graph of length  $n$ , then*

$$\chi_{\text{gr}}(I_n; A^*) = \alpha(\alpha - 1)^n, \text{ for } n > 0,$$

and  $\chi_{\text{gr}}(I_0; A^*) = \alpha$ .

From Remark 5.13 and Corollary 5.16 it follows that  $(\alpha - 1)^x$  divides  $\chi_{\text{gr}}(L)$ , where  $x$  is the number of edges incident only to vertices which are either univalent or unstable. This, similar divisibility properties, and the decomposition shown in Theorem 5.6, seem to hint to the fact that  $\chi_{\text{gr}}(L)$  is sensible to “dynamical properties” of the graph.

*Question 5.19.* Does it exist a closed formula for  $\chi_{\text{gr}}(\mathbb{L})$ , with  $\mathbb{L}$  a linear graph, which features only dynamical data (e.g. stable and unstable vertices, change of stability *etc.*) and the polynomial of an alternating  $\chi_{\text{gr}}(\mathbb{A}_n)$  ( $n$  also depending on dynamical data)?

## 6. RELATIONS WITH SIMPLICIAL HOMOLOGY

Let  $\mathbb{G}$  be a digraph and  $P(\mathbb{G})$  be its path poset. In this section we give a topological description of multipath cohomology. More specifically, we see that multipath cohomology  $H_\mu^*(\mathbb{G}; R)$  is the ordinary cohomology of a certain simplicial complex  $X(\mathbb{G})$  associated to  $\mathbb{G}$  – cf. Theorem 6.8. For instance, this approach leads to a reinterpretation of the Mayer-Vietoris exact sequence for multipath cohomology in topological terms. In the last subsection, we discuss which simplicial complexes can be realised as  $X(\mathbb{G})$  for some  $\mathbb{G}$ .

**6.1. Background material.** Recall that an  $n$ -cell is a topological space homeomorphic to an  $n$ -dimensional disc and that a *CW-complex* is a (Hausdorff) topological space  $X$  obtained by inductively gluing  $n$ -cells – cf. [Hat00, Chapter 0]:

- (i) start with a discrete space  $X^0$ , called 0-skeleton, whose points are the 0-cells of  $X$ ;
- (ii) form the  $n$ -skeleton  $X^n$  from  $X^{n-1}$  by gluing  $n$ -dimensional cells  $E_\alpha^n$  via (continuous) characteristic maps  $\phi_\alpha: \partial(E_\alpha^n) \rightarrow X^{n-1}$  from the boundary  $\partial(E_\alpha^n)$  of  $E_\alpha^n$  to the  $(n-1)$ -skeleton. More precisely,  $X^n$  is defined as the quotient space of  $X^{n-1} \sqcup_\alpha E_\alpha^n$  by the identification  $x \sim \phi_\alpha(x)$  for  $x \in \partial(E_\alpha^n)$ .
- (iii)  $X$  is defined as  $X := \cup_n X^n$ , and endowed with the weak topology.

A *regular CW-complex* is a CW-complex for which all the characteristic maps are homeomorphisms, i.e. for every  $n$ , all the  $\phi_\alpha(E_\alpha^n)$  are homeomorphic to closed  $n$ -discs – cf. [Bjo84, Section 3].

Let  $X$  be a CW-complex. The *face poset*  $\mathcal{F}(X)$  of  $X$  is the poset on the set of cells of  $X$ , ordered by containment and augmented with a minimum element  $\hat{0}$  corresponding to the empty cell. We will see that the path poset  $P(\mathbb{G})$  of a graph  $\mathbb{G}$  is the face poset of a certain CW-complex. We first need a definition:

**Definition 6.1** ([Bjo84, Definition 2.1]). A poset  $P$  with at least two elements is said to be a *CW-poset* if it has a minimum  $\hat{0}$ , and, for all  $x \in P \setminus \hat{0}$ , the (geometric realisation<sup>6</sup> of the) interval  $(\hat{0}, x) := \{z \in P \mid \hat{0} < z < x\}$  is homeomorphic to a sphere.

*Example 6.2.* A Boolean poset is a CW-poset. More generally, by [Bjo84, Proposition 2.6 (b)], every downward closed subposet of a Boolean poset is a CW-poset. By Remark 2.18, the path poset  $P(\mathbb{G})$  is a downward closed subposet of a Boolean poset, hence it is a CW-poset.

The following result gives a description of the CW-posets in terms of CW-complexes:

**Lemma 6.3** ([Bjo84, Proposition 3.1]). *A poset  $P$  is a CW-poset if, and only if, it is isomorphic to the face poset of a regular CW-complex.*

As a consequence of Example 6.2 and Lemma 6.3, for  $P(\mathbb{G})$  the path poset of a digraph  $\mathbb{G}$ , there exists a regular CW-complex  $X(\mathbb{G})$  whose face poset is isomorphic to  $P(\mathbb{G})$ . We can actually be more specific. Recall that an (*augmented abstract*) *simplicial complex*  $K$  on a vertex set  $V$  is  $K \subseteq \wp(V)$  such that:

- (i)  $\{v\} \in K$  for all  $v \in V$ ;
- (ii) if  $\sigma \in K$ , then  $\sigma$  is finite;
- (iii) if  $\sigma \in K$  and  $\tau \subseteq \sigma$ , then  $\tau \in K$ .

<sup>6</sup>By geometric realisation of a poset we mean the geometric realisation of its order complex (i.e. the abstract simplicial complex whose faces are the totally ordered sub-posets of our poset), see [Koz08, Chapter 9].

An  $n$ -simplex in  $K$  is  $\sigma \in K$  with  $n + 1$  elements. In particular, the empty set  $\emptyset$  is the unique  $(-1)$ -simplex in  $K$ , while the vertices are all the 0-simplices in  $K$ . In other words, an augmented abstract simplicial complex  $K$  on a vertex set  $V$  is a downward closed sub-poset of  $(\wp(V), \subset)$ , whose elements are finite sets, containing  $\{\{v\}\}_{v \in V}$ . Note that if  $V$  is finite, then condition (ii) is automatically satisfied.

**Definition 6.4.** Given a graph  $G$  its *multipath complex* is the augmented abstract simplicial complex  $X(G)$  on  $E(G)$  whose simplices are given by the set of edges of each multipath.

Since each spanning sub-graph of a multipath is a multipath, it is clear that  $X(G)$  is indeed an augmented abstract simplicial complex. Furthermore, we have the following observation.

*Remark 6.5.* A morphism of digraphs induces a morphism between the corresponding multipath complexes, which sends multipaths of length 1 (edges in the graph) to multipaths of length 1 (edges in the graph). Maps between simplicial complexes preserving the inclusion and sending 0-simplices to 0-simplices are called *simplicial maps*. It follows that for each morphism of digraphs  $\phi: G \rightarrow G'$  there is an associated simplicial map  $X(\phi): X(G) \rightarrow X(G')$ . Clearly, we have that  $X(\text{id}_G) = \text{id}_{X(G)}$ , and that  $X(\phi \circ \psi) = X(\phi) \circ X(\psi)$ . In more abstract terms, the multipath complex defines a functor  $X: \mathbf{Digraph} \rightarrow \mathbf{SimpComp}$  from the category of digraphs, and morphisms of digraphs, to the category of augmented abstract simplicial complexes, and simplicial maps.

We can give a more explicit description of  $X(G)$ .

*Construction of  $X(G)$ .* We construct the CW-complex which has  $X(G)$  as face poset by explicitly describing its cells and their gluing maps. Our construction here follows the proof that simplicial complexes are CW-complexes, and it gives a concrete way to obtain a CW complex from a path poset.

We start by associating to the empty multipath of  $G$ , i.e. the set of vertices of  $G$ , the empty simplex, i.e. the  $(-1)$ -skeleton of  $X(G)$ . We build the complex  $X(G)$  by attaching  $n$ -cells to the discrete set  $X(G)^0 := E(G)$ , i.e. the 0-skeleton of  $X(G)$  is given by the set of edges of  $G$ .

Suppose to have iteratively constructed the  $(n - 1)$ -skeleton  $X(G)^{(n-1)}$  of  $X(G)$ . Each multipath  $H$  of length  $n + 1$  – i.e. with  $n + 1$  edges – is identified by edges  $e_{i_0}, \dots, e_{i_n}$  of  $G$ ; hence, by  $n + 1$  points of  $X(G)^0$ . We associate to  $H$  the (abstract)  $n$ -dimensional simplex  $\Delta_H^n := [e_{i_0}, \dots, e_{i_n}]$ , which is an  $n$ -cell of  $X(G)^n$ . In this way, each multipath  $H$  of length  $n$  gives an  $n$ -cell of  $X(G)$ . Note that the multipath  $H$  can be obtained from exactly  $n + 1$  (sub-)multipaths of length  $n$ , say  $H_0, \dots, H_n$ , by adding an edge; more precisely,  $H_j$  is the multipath of  $G$  with edges  $e_{i_1}, \dots, \widehat{e_{i_j}}, \dots, e_{i_n}$ , where  $\widehat{e_{i_j}}$  indicates that the edge  $e_{i_j}$  is not counted. The multipaths  $H_0, \dots, H_n$  correspond to  $(n - 1)$ -cells of  $X(G)$ , and, moreover, the boundary  $\partial(\Delta_H^n)$  of  $\Delta_H^n$  can be identified with the union of its faces  $\Delta_{H_j}^{n-1}$ . The characteristic map of  $\Delta_H^n$  is then defined by gluing each face  $[e_{i_0}, \dots, \widehat{e_{i_j}}, \dots, e_{i_n}]$  in  $\partial(\Delta_H^n)$  with the corresponding  $(n - 1)$ -cell in  $X(G)^{(n-1)}$ .

More generally, we glue the simplices  $\{\Delta_H^n\}_{\text{length}(H)=n+1}$  to  $X(G)^{(n-1)}$  by identifying the facets of each  $\Delta_H^n$  with the simplices  $\Delta_{H_0}, \dots, \Delta_{H_n}$ , concluding the construction of the  $n$ -skeleton, hence of  $X(G)$ .

Observe that maximal simplices in  $X(G)$  correspond to maximal multipaths in  $P(G)$ . Therefore, to construct the geometric realisation of  $X(G)$  we can proceed by finding the maximal multipaths in  $P(G)$ , look at the intersection of each pair of maximal multipath, then glue the simplices associated to maximal multipaths along the faces determined by their intersections.

*Example 6.6.* In this example we explicitly describe the geometric realisation of the simplicial complex  $X(G)$ , for  $G$  the H-shaped digraph illustrated in Figure 18. Observe that we have five 0-cells  $E_{01}^0, E_{14}^0, E_{21}^0, E_{34}^0$ , and  $E_{54}^0$ , where the cell  $E_{ij}^0$  corresponds to the edge  $(v_i, v_j)$ . Each

1-cell in  $X(\mathbb{G})$  corresponds to a multipath of length two. Thus, there are precisely six 1-cells  $E_{01,14}^1, E_{01,34}^1, E_{01,54}^1, E_{21,14}^1, E_{21,34}^1,$  and  $E_{21,54}^1$ . The 1-cell  $E_{x,y}^1$  bounds the 0-cells  $E_x^0$  and  $E_y^0$ .



FIGURE 18. The  $H$  digraph and its geometric realisation.

Let  $K$  be an augmented abstract simplicial complex. The  $n$ -th reduced simplicial chain group  $\tilde{C}_n(K; R)$  (with coefficients in  $R$ ) is the free  $R$ -module generated by all the  $n$ -simplices in  $K$ .

*Notation 6.7.* Let  $K$  be an augmented abstract simplicial complex. We assume, from now on, a linear ordering  $v_1 < v_2 < \dots < v_k$  of the 0-simplices of  $K$  to be fixed. Given an  $n$ -simplex in  $K$ , say  $\sigma = \{v_{i_0}, \dots, v_{i_n}\}$ , we denote  $\sigma$  as  $[v_{i_0}, \dots, v_{i_n}]$ , where  $i_0 < i_2 < \dots < i_n$ , and define

$$[v_{i_s(0)}, \dots, v_{i_s(n)}] := (-1)^{\text{sgn}(s)} [v_{i_0}, \dots, v_{i_n}],$$

for each  $s$  in the symmetric group over  $\{0, \dots, n\}$ .

The  $n$ -th simplicial differential is defined as

$$\delta_n: \tilde{C}_n(K; R) \longrightarrow \tilde{C}_{n-1}(K; R): [v_{i_0}, \dots, v_{i_n}] \longmapsto \sum_{j=0}^n (-1)^j [v_{i_0}, \dots, \widehat{v_{i_j}}, \dots, v_{i_n}],$$

where the hat  $\widehat{x}$  indicates that  $x$  is missing. Simplicial cohomology is given by applying the Hom functor, that is, the  $n$ -th reduced simplicial co-chain group is

$$\tilde{C}^m(K; R) = \text{Hom}_R(\tilde{C}_m(K; R); R)$$

and, given  $f \in C^{m+1}(K; R)$ , we define the co-boundary map  $\delta$  by setting  $\delta^n(f) = f(\delta_n(\sigma))$ .

**6.2. Multipath cohomology is simplicial.** Let  $X(\mathbb{G})$  be the multipath complex associated to the digraph  $\mathbb{G}$ , and let  $\mathbb{K}$  be a field.

**Theorem 6.8.** *The multipath cohomology  $H_\mu^n(\mathbb{G}; \mathbb{K})$  of  $\mathbb{G}$  is isomorphic to  $\tilde{H}^{n-1}(X(\mathbb{G}); \mathbb{K})$ , that is, the reduced (simplicial) cohomology of  $X(\mathbb{G})$ .*

Note that in the isomorphism between the multipath cohomology of  $\mathbb{G}$  and the simplicial cohomology of  $X(\mathbb{G})$  there is a shift of degree one.

*Proof.* Let  $X$  be a finite simplicial complex. The reduced simplicial cochain complex  $\tilde{C}^m(X; \mathbb{K})$  can be seen as the vector space over  $\mathbb{K}$  generated by the duals of all simplices (including the empty simplex) of  $X$ , with co-boundary map  $\delta$  given by

$$\delta(\sigma^*) = \sum_{\tau} (-1)^{\epsilon'(\tau, \sigma)} \tau^*,$$

where  $\tau$  ranges among all simplices admitting  $\sigma$  as a face, and  $\epsilon'(\tau, \sigma)$  is 0 or 1 depending on whether or not the orientation of  $\sigma$  matches with the orientation induced by  $\tau$  – for a more detailed construction, the reader can consult [Koz08, Section 3.4.3].

By construction, the simplices of  $X(\mathbb{G})$  correspond to multipaths in  $\mathbb{G}$ ; more precisely, the points of  $X(\mathbb{G})^0$  are the edges of  $\mathbb{G}$ , and the multipath  $\mathbb{H}$  identified by the edges  $e_{i_1}, \dots, e_{i_n}$  corresponds to the simplex  $[e_{i_1}, \dots, e_{i_n}]$ . The vector space  $C_\mu^n(\mathbb{G}; \mathbb{K})$  has one generator  $b_{\mathbb{H}}$  for each multipath  $\mathbb{H}$  of length  $n$ , and the differential is given by

$$d(b_{\mathbb{H}}) = \sum_{\mathbb{H}' \supset \mathbb{H}} (-1)^{\epsilon(\mathbb{H}', \mathbb{H})} b_{\mathbb{H}'},$$

for a certain sign assignment  $\epsilon$ . It follows that the correspondence  $b_{\mathbb{H}} \mapsto [e_{i_1}, \dots, e_{i_n}]$ , where  $\mathbb{H}$  is identified by the edges  $e_{i_1}, \dots, e_{i_n}$ , extends to an isomorphism of graded vector spaces which commutes with the differentials up to a sign. Note that a multipath of length  $n$  corresponds to an  $(n - 1)$ -dimensional simplex, which gives the shift in cohomological degree.

We conclude by observing that  $\epsilon'$  is a sign assignment on the face poset of  $X(\mathbb{G})$ , which is isomorphic to  $P(\mathbb{G})$ ; in fact, the statement now follows from the uniqueness (up to isomorphism) of the sign assignment on the path posets [CCDT21, Corollary 3.17].  $\square$

An alternative way to construct a simplicial complex  $X'(\mathbb{G})$ , starting with the path poset  $P(\mathbb{G})$  and with same cohomology groups as  $X(\mathbb{G})$ , is to use the *order complex*  $\Delta(P(\mathbb{G}))$  of  $P(\mathbb{G})$ , see for instance [Wac06]. It is now a standard fact that, for a simplicial complex  $X$ , the realisation of the order complex  $\Delta(\mathcal{F}(X))$  of the face poset of  $X$  is the barycentric subdivision of  $X$  – cf. [Wac06]. Consequently,  $X'(\mathbb{G})$  is the barycentric subdivision of  $X(\mathbb{G})$ . Furthermore, the simplicial (co)homology of the barycentric subdivision of a simplicial complex  $X$  is the simplicial (co)homology of  $X$ . As an immediate consequence we have the following corollary.

**Corollary 6.9.** *The multipath cohomology  $H_\mu^*(\mathbb{G}; \mathbb{K})$  is the reduced (simplicial) cohomology of the order complex  $\Delta(P(\mathbb{G}))$  associated to  $P(\mathbb{G})$ .*

Observe that the isomorphism of Theorem 6.8 is well-behaved in a functorial sense:

*Remark 6.10.* By Remark 6.4, associating the simplicial complex  $X(\mathbb{G})$  to a graph  $\mathbb{G}$  is functorial with respect to morphisms of digraphs. Therefore, since taking cohomology groups is functorial with respect to simplicial maps, the isomorphism in cohomology provided by Theorem 6.8 induces a natural isomorphism of functors  $\eta: H_\mu^*(-; \mathbb{K}) \Rightarrow \tilde{H}^{*-1}(-; \mathbb{K}) \circ X$ . More concretely, given a morphism of digraphs  $\phi: \mathbb{G} \rightarrow \mathbb{G}'$ , we obtain the following square

$$\begin{array}{ccc} H_\mu^*(\mathbb{G}'; \mathbb{K}) & \xrightarrow{\phi^*} & H_\mu^*(\mathbb{G}; \mathbb{K}) \\ \eta_{\mathbb{G}'} \downarrow & & \downarrow \eta_{\mathbb{G}} \\ \tilde{H}^{*-1}(X(\mathbb{G}'); \mathbb{K}) & \xrightarrow{\phi^*} & \tilde{H}^{*-1}(X(\mathbb{G}); \mathbb{K}) \end{array}$$

that is commutative, with vertical arrows which are isomorphisms. This, in particular, extends the functoriality result [CCDT21, Theorem 1.3] also to non-regular morphisms of digraphs.

**6.3. Mayer-Vietoris from the topological viewpoint.** In this subsection we reinterpret the Mayer-Vietoris-type theorem for multipath cohomology, that is Theorem 4.4, using the simplicial description given in Theorem 6.8.

Let  $X$  be a simplicial complex and assume that there are sub-complexes<sup>7</sup>  $Y_1$  and  $Y_2$  such that:

- (1)  $X = Y_1 \cup Y_2$ ;
- (2) their intersection of  $Y_1$  and  $Y_2$  is a sub-complex of  $X$ ,  $Y_1$ , and  $Y_2$ ;

then, we have a long exact sequence in (reduced) simplicial cohomology

$$\dots \rightarrow \tilde{H}^i(X; \mathbb{K}) \rightarrow \tilde{H}^i(Y_1; \mathbb{K}) \oplus \tilde{H}^i(Y_2; \mathbb{K}) \rightarrow \tilde{H}^i(Y_1 \cap Y_2; \mathbb{K}) \rightarrow \tilde{H}^{i+1}(Y_1 \cap Y_2; \mathbb{K}) \rightarrow \dots$$

<sup>7</sup>A *sub-complex*  $Y$  of a simplicial complex  $X$  is a subset of  $X$  which is itself a simplicial complex.



called *Mayer-Vietoris sequence* [Hat00, Chapter 2.2]. Intuitively, this sequence “describes” the cohomology of the space  $X$  in terms of the cohomology of  $Y_1$ ,  $Y_2$ , and  $Y_1 \cap Y_2$ .

In Theorem 4.4, we obtained a similar sequence for path posets associated to regular morphisms of digraphs. We can actually use the correspondence between multipath and simplicial cohomologies to re-prove this result; we can interpret the path poset of a graph  $G$  as the face poset of a simplicial complex  $X(G)$ . Furthermore, a downward closed sub-poset  $S$  of  $P(G)$  corresponds to a (face poset of a) sub-complex of  $X(G)$  – the correspondence being described by taking the simplices of  $X(G)$  given by the multipaths belonging to  $S$ . By Theorem 6.8, since we have  $H_\mu^*(G; \mathbb{K}) \cong \tilde{H}^{*-1}(X(G); \mathbb{K})$ , we can replace the cohomology of spaces with the multipath cohomology of the corresponding graph with the corresponding multipath cohomology, obtaining the sequence in Theorem 4.4. To a more intimate level, and in a more abstract language, this is due to the fact that the gluing construction in Definition 3.3 represents the pushout in **Digraph** – cf. Remark 3.5.

*Remark 6.11.* The Mayer-Vietoris long exact sequence in homology is classically obtained using the long exact sequence of the pair applied to homotopy pushouts [Rot88, Theorem 6.3]. Since the gluing construction in Definition 3.3 gives a pushout diagram of graphs, it is reasonable to think that a Mayer-Vietoris long exact sequence can be obtained also for multipath cohomology. In fact, let  $G, G_1, G_2$  as in Definition 3.3; by Remark 3.5, the square

$$\begin{array}{ccc}
 P(G) & \xrightarrow{i_1} & P(G_1) \\
 \downarrow i_2 & & \downarrow j_1 \\
 P(G_2) & \xrightarrow{j_2} & P(G_1) \nabla_{P(G)} P(G_2)
 \end{array}$$

(♠)

is a pushout square. Observe that also the square

$$\begin{array}{ccc}
 X(G) & \xrightarrow{\iota_1} & X(G_1) \\
 \downarrow \iota_2 & & \downarrow j_1 \\
 X(G_2) & \xrightarrow{j_2} & X(G_1) \coprod_{X(G)} X(G_2)
 \end{array}$$

(◇)

is a pushout square of simplicial complexes, where  $X \coprod_Z Y$  indicates the gluing of  $X$  and  $Y$  along  $Z \hookrightarrow X, Y$ . Furthermore, taking the face poset takes the square (◇) to the pushout square (♠); i.e.  $\mathcal{F}(X(G_1) \coprod_{X(G)} X(G_2)) = P(G_1) \nabla_{P(G)} P(G_2)$ . As inclusions of simplicial complexes are cofibrations, the square (◇) is (equivalent to) a homotopy pushout square. Hence, Diagram (◇) gives rise to the Mayer-Vietoris long exact sequence in reduced cohomology. Naturality of Theorem 6.8, as in Remark 6.10, now gives the Mayer-Vietoris long exact sequence in multipath cohomology. In the light of the above discussion, we can think informally that “inclusions as downward closed sub-posets are cofibrations in the category of posets, and the corresponding nablas are actually homotopy pushouts” – for a more precise treatment of the homotopy theory of posets, see for example [Rap10].

**6.4. Realisability of path posets and examples.** It is interesting to understand which simplicial complexes can be realised as  $X(G)$  for some digraph  $G$ . In this subsection we investigate this problem and some of its consequences. First, we observe that all the spheres can be realised.

*Example 6.12.* Let  $n \geq 1$  be a natural number. The boundary of the  $n$ -dimensional standard simplex  $\Delta^n$  is homeomorphic to the  $(n-1)$ -dimensional sphere  $\mathbb{S}^{n-1}$ ; we argue that  $\partial\Delta^n$  is the simplicial complex  $X(P_n)$ , where  $P_n$  is the coherently oriented polygonal graph on  $n$  vertices as illustrated in Figure 3. By [CCDT21, Section 2.3], the path poset  $P(P_n)$  is isomorphic (as posets) to the Boolean poset  $\mathbb{B}(n+1)$  with its maximum removed. Note that the  $(n+1)$ -Boolean poset  $\mathbb{B}(n+1)$  is the face poset  $\mathcal{F}(\Delta^n)$  of  $\Delta^n$ . Hence, it follows that  $P(P_n)$  is the

face poset of  $\Delta^n$  minus its unique  $(n + 1)$ -dimensional cell, which turn to be precisely the sphere  $\partial\Delta^n \cong \mathbb{S}^{n-1}$ . As a consequence, by Theorem 6.8, we get

$$H_\mu^i(P_n; \mathbb{K}) \cong \tilde{H}^{i-1}(\mathbb{S}^{n-1}; \mathbb{K}) \cong \begin{cases} \mathbb{K} & \text{if } i = n, \\ 0 & \text{otherwise} \end{cases}$$

the computations of the cohomology of  $P_n$ , with coefficients in  $\mathbb{K}$ , for every  $n \in \mathbb{N}$ . Alternatively, this could have been obtained as a consequence of the isomorphism with Hochschild homology of  $\mathbb{K}$  – cf. [CCDT21, Proposition 1.4].

Another example is given by the dandelion graphs; their multipath complexes (or, better, their geometric realisations) are homotopy equivalent to wedges of 1-dimensional spheres.

*Example 6.13.* Recall that the dandelion graph  $D_{n,m}$  is the digraph with a single  $(m + n)$ -valent vertex  $v_0$ , all remaining vertices are univalent, there are  $n$  edges with target  $v_0$ , and there are  $m$  edges with source  $v_0$  – cf. Figure 11. In this example we assume  $nm > 0$ .

The multipath complex of  $D_{n,m}$  is easily described as follows; we have two sets  $e_1, \dots, e_n$  and  $e'_1, \dots, e'_m$  of 0-cells in  $X(D_{n,m})$ , and each  $e_i$  is joined with each  $e'_j$  by a 1-cell, and there are no other 1- or higher cells. Thus, (the realisation of)  $X(D_{n,m})$  is the complete bipartite graph  $K_{n,m}$ . It follows that

$$H_\mu^i(D_{n,m}; \mathbb{K}) \cong \tilde{H}^{i-1}(K_{n,m}; \mathbb{K}) \cong \begin{cases} \mathbb{K}^{(n-1)(m-1)} & i = 2, \\ 0 & \text{otherwise.} \end{cases}$$

A quick sketch on how to get the last isomorphism. Since  $K_{n,m}$  is a connected 1-dimensional simplicial complex, we have that: all the reduced homology groups are trivial except (possibly) in degree 1, and  $\dim(\tilde{H}^{i-1}(K_{n,m}; \mathbb{K})) = \text{rank}(\pi_1(K_{n,m}))$ . Note that, since both homology and  $\pi_1$  are homotopy invariants, we can contract any 1-cells without changing the fundamental group. Contracting one edge in  $K_{n,m}$ , say the 1-cell  $e$  bounding  $e_1$  and  $e'_1$ , we obtain a space  $X$  which is (homotopic to) a wedge of 1-spheres. The 0-cells in  $X$  are: a vertex  $v$ , obtained by contracting  $e$ ,  $e_1, \dots, e_n$ ,  $e'_1, \dots$ , and  $e'_n$ . Now to compute the rank of  $\pi_1(K_{n,m})$  it is sufficient to count the (non-singular simple) loops in  $X$  starting and ending in  $v$ ; these are precisely all triangles with vertices  $v$ ,  $e_i$ , and  $e'_j$ , for all  $i, j > 1$ , which are exactly  $(n - 1)(m - 1)$ .

It is yet not clear whether multipath cohomology can be supported in different degrees. In the next example we show that this is also possible.

*Example 6.14.* Consider the graph  $Q$  to be the coherently oriented polygon  $P_3$  with a diagonal, as illustrated in Figure 19. The path poset associated to  $Q$  has multipaths of length at most 3,




FIGURE 19. The coherently oriented polygon  $P_3$  with a diagonal.

corresponding to having at most 2-simplices in the realisation of  $X(Q)$ . The realisation of the multipath complex associated to  $P_3$  is isomorphic to the 2-sphere  $\mathbb{S}^2$ , and adding the diagonal results in adding two 1-cells to  $\mathbb{S}^2$ . A depiction of the geometric realisation of  $X(Q)$  is given in Figure 20; we have decorated the vertices in the realisation corresponding to the edges  $(v_1, v_2)$  and  $(v_1, v_3)$  of  $Q$ , all the others being interchangeable in the realisation. The simplicial com-




FIGURE 20. The geometric realisation of  $X(G)$ , where  $G$  is the graph in Figure 19. In green is illustrated the boundary of a tetrahedron.

plex  $X(G)$  is homotopically equivalent to the wedge of  $\mathbb{S}^2$  with a copy of  $\mathbb{S}^1$ , showing that the multipath cohomology

$$H_\mu^*(Q; \mathbb{K}) \cong \begin{cases} \mathbb{K} & \text{for } n = 2, \\ \mathbb{K} & \text{for } n = 3, \\ 0 & \text{otherwise.} \end{cases}$$

of  $Q$  is non-zero in degrees 2 and 3 by Theorem 6.8.

Let  $G'$  be a digraph with a univalent vertex  $w$ . In Lemma 4.20, we have proved that there exists a graph  $G$ , obtained by gluing a linear sink or source to  $w$  (cf. Figure 12), such that  $H_\mu^*(G; \mathbb{K}) \cong H_\mu^{*+1}(G'; \mathbb{K})$ . Classically, this is the property of the suspension<sup>8</sup>  $\Sigma X$ . In the proof of the next proposition we will see that these two constructions are related. Let  $\bigvee^k \mathbb{S}^n$  be the wedge of  $k$   $n$ -dimensional spheres.

**Proposition 6.15.** *For all  $k, n \in \mathbb{N}$ , there exists a graph  $G_{k,n}$  such that*

$$\bigvee^k \mathbb{S}^n \simeq |X(G_{k,n})|$$

where  $|X(G_{k,n})|$  is the geometric realisation of the (abstract) simplicial complex  $X(G_{k,n})$ .

*Proof.* Let  $D_{n,m}$  be a dandelion graph, cf. Definition 4.14 and Figure 11. The geometric realisation of  $G_{k,0} := D_{k+1,0}$  consists of  $k+1$  points, which can be seen as a wedge of  $k$  0-spheres. In Example 6.13, we have shown that  $X(D_{k+1,2})$  for  $k > 0$  is homotopy equivalent to a wedge of  $k$  1-spheres. In order to get a wedge  $\bigvee^k \mathbb{S}^n$  of higher dimensional spheres, it is enough to iteratively glue sinks or sources to  $G_{k,1} := D_{k+1,2}$ .

More precisely, we fix  $k$  and proceed by induction on  $n$ . We assume to have already constructed a graph  $G_{k,n}$ , whose associated multipath complex is homotopic to a wedge of  $k$   $n$ -dimensional spheres, and that it has a univalent vertex, say  $v$ . Now, we glue a source or a sink, depending on whether  $v = s(e)$  or  $v = t(e)$ , for some edge  $e \in E(G_{k,n})$  – cf. Lemma 4.20 and Figure 12. Define  $G_{k,n+1}$  to be the graph obtained this way. Note that  $G_{k,n+1}$  has at least two univalent vertices. Now, by Theorem 3.13 applied to  $v$ , the path poset of  $G_{k,n+1}$  is the gluing of two copies of  $\mathfrak{Cone} P(G_{k,n})$  (which are  $P((G_{k,n})_v^{(1)})$  and  $P((G_{k,n})_v^{(2)})$ , in the notation of Theorem 3.13) glued along  $P(G_{k,n})$ . This can be illustrated using the following pushout diagram

$$\begin{array}{ccc} P(G_{k,n}) & \xrightarrow{j_1} & \mathfrak{Cone} P(G_{k,n}) \\ j_2 \downarrow & & \downarrow \iota_1 \\ \mathfrak{Cone} P(G_{k,n}) & \xrightarrow{\iota_2} & G_{k,n+1} \end{array}$$

<sup>8</sup>the topological space  $\Sigma X$  obtained from  $X$  by taking the cylinder  $X \times [0, 1]$ , and collapsing each of the faces  $X \times \{0\}$  and  $X \times \{1\}$  to a point. Alternatively, the suspension can be seen as two copies of the cone  $(X \times [0, 1]) / (x, 1) \sim (y, 1)$  glued along  $X \times \{0\}$ .

By passing to the geometric realisations of multipath complexes (see also Remark 6.11), we obtain the pushout diagram

$$\begin{array}{ccc} |X(\mathbf{G}_{k,n})| \simeq \bigvee^k \mathbb{S}^n & \xrightarrow{\iota_1} & \text{Cone}(\bigvee^k \mathbb{S}^n) \simeq * \\ \iota_2 \downarrow & & \downarrow j_1 \\ * \simeq \text{Cone}(\bigvee^k \mathbb{S}^n) & \xrightarrow{j_2} & \Sigma(\bigvee^k \mathbb{S}^n) \simeq |X(\mathbf{G}_{k,n+1})| \end{array}$$

of topological spaces, where  $*$  denotes the one point space. As the suspension of a wedge on  $k$   $n$ -spheres is homotopy equivalent to a wedge of  $k$   $(n+1)$ -spheres, the statement follows.  $\square$

The proposition says that wedges of spheres of the same dimension can be realised, up to homotopy, as multipath complexes of certain digraphs. It is therefore natural to ask whether wedges of spheres of different dimensions can also be realised as (geometric realisations of) multipath complexes. We have seen, in Example 6.14, that the wedge  $\mathbb{S}^1 \vee \mathbb{S}^2$  can be realised, but it is not clear whether all wedges of spheres can be:

*Question 6.16.* Can we realise, up to homotopy, all wedges of spheres? For example, can we realise a space homotopic to  $\mathbb{S}^1 \vee \dots \vee \mathbb{S}^1 \vee \mathbb{S}^0 \vee \dots \vee \mathbb{S}^0$ ?

We observe here that, if we allow the graphs to be disconnected, then it is possible to obtain also the topological join  $X(\mathbf{G}) * X(\mathbf{G}')$  of  $X(\mathbf{G})$  and  $X(\mathbf{G}')$ ; concretely, this can be realised as  $X(\mathbf{G} \sqcup \mathbf{G}')$  – compare [Koz08, Definition 2.16], and Remark 3.2. Note that the join of simplicial complexes commutes with the geometric realisation [Koz08, Equation (2.9)]. In particular, we can realise the suspension  $\Sigma X(\mathbf{G})$  of  $X(\mathbf{G})$  also as the multipath complex  $X(\mathbf{G} \sqcup \mathbf{A}_2)$  [Koz08, Example 2.32.(1)]. Realising joins allows us to obtain more complicated wedges of spheres giving a partial answer to the question above.

*Example 6.17.* The join commutes, up to homotopy, with the wedge operation, and it is well-known that  $\mathbb{S}^n * \mathbb{S}^m \simeq \mathbb{S}^{n+m+1}$  – cf. [Bel]. An immediate consequence of these facts is that

$$\bigvee_{i=1}^k \mathbb{S}^{n_i} * \bigvee_{j=1}^h \mathbb{S}^{m_j} \simeq \bigvee_{i=1}^k \bigvee_{j=1}^h \mathbb{S}^{n_i+m_j+1}.$$

For instance, we have shown in Example 6.14 that we can realise  $\mathbb{S}^2 \vee \mathbb{S}^1$  as  $|X(\mathbf{Q})|$ , where  $\mathbf{Q}$  is the square  $\mathbf{P}_3$  with a diagonal. Therefore, we have that

$$\mathbb{S}^5 \vee \mathbb{S}^4 \vee \mathbb{S}^4 \vee \mathbb{S}^3 \simeq (\mathbb{S}^2 \vee \mathbb{S}^1) * (\mathbb{S}^2 \vee \mathbb{S}^1) \simeq |X(\mathbf{Q} \sqcup \mathbf{Q})|.$$

Since all spaces we are able to realise are, up to homotopy, wedges of spheres, the following question arises naturally;

*Question 6.18.* Can we realise spaces which are not homotopic to wedges of spheres, e.g. the real projective spaces?

Before further discussing about the realisability of simplicial complexes via path posets, we want to extend the category of digraphs. Recall that a *multigraph*, or *quiver*, is 4-uple  $(V, E, s, t)$  where  $V$  and  $E$  are finite sets, whose elements are called *vertices* and *edges* respectively, while

$$s, t: E \longrightarrow V$$

are the *source* and *target* functions. To avoid self-loops we require  $t(e) \neq s(e)$ , for all  $e \in E$ . Clearly, all digraphs can be seen as multigraphs. The definition of path poset and multipath cohomology extends verbatim to multigraphs. Similarly, we have an isomorphism between constant coefficients multipath cohomology and simplicial homology of  $X(\mathbf{G})$ .




FIGURE 21. A 1-dimensional complex  $X$  whose face poset is not a path poset.

**Proposition 6.19.** *There are simplicial complexes whose face poset cannot be realised as the path poset of any multigraph. In other words, the functor  $X: \mathbf{Digraph} \rightarrow \mathbf{SimpComp}$  (cf. Remark 6.5), or better its extension to multigraphs, is not (essentially<sup>9</sup>) surjective on objects.*

*Proof.* Consider Figure 21, it contains the picture of a geometric realisation of the simplicial complex, say  $X$ . The face poset of  $X$  consists of: the empty simplex  $\emptyset$ , five 0-simplices, and six 1-simplices.

Assume, by the means of contradiction, that there exists a multigraph  $G$  such that  $X = X(G)$ . It follows that  $G$  has five edges, six multipaths of length two, and no multipaths of higher length. Note that we cannot read the number of vertices from the path poset (and thence from  $X(G)$ ). Moreover, adding a vertex with no edges does not change the path poset. Therefore, without loss of generality, we might assume that  $G$  has no connected components which are vertices.

We claim that the edges  $e_1, e_2$ , and  $e_5$ , form a coherently oriented triangle in  $G$ . We start by noticing that the sub-graph  $T$  of  $G$  given by the edges  $e_1, e_2$ , and  $e_5$  is connected. Either all pairs of edges among  $e_1, e_2$ , and  $e_5$  share a vertex, in which case  $T$  is connected, or there are  $e_i$  and  $e_j$ , with  $i \neq j$  and  $i, j \in \{1, 2, 5\}$ , do not share any vertex. In the latter case, both must share at least one vertex with  $e_k$ ,  $k \neq i, j$  and  $k \in \{1, 2, 5\}$ . If not, we would have a multipath of length 3 given by  $e_1, e_2, e_5$ . It follows that  $T$  is connected. The path poset of  $T$  is (isomorphic to) the sub-poset of  $P(G)$  given by multipaths with edges  $e_1, e_2$ , or  $e_5$ . Pasting all the pieces together, we have that  $T$  is a connected graph with 3 edges whose path poset is a Boolean poset minus its maximum; consequently,  $T$  must be a coherently oriented triangle as claimed – cf. Proposition 3.1.

The same reasoning as above proves that the edges  $e_3, e_4$ , and  $e_5$ , also form a coherently oriented triangle in  $G$ . It follows that  $G$  must consist of two coherently oriented triangles glued at least along one edge (namely  $e_5$ ). Now we can enumerate all possible  $G$ 's; these are illustrated, up to orientation reversal and exchanging in the roles of  $e_1$  and  $e_2$ , and of  $e_3$  and  $e_4$ , in Figure 22.




FIGURE 22. Possible configurations for the graph  $G$ .

In case (a) we have two multipaths of length 3, while in case (b) the 0-cells corresponding to  $e_4$  and  $e_3$  must be in the boundary of three 1-cells in  $X(G) = X$ . In either case we get a contradiction, and the statement follows.  $\square$

<sup>9</sup>That is up to (simplicial, in our case) isomorphism.

Note that, while the face poset of the simplicial complex in Figure 21 cannot be realised as a path poset, we can realise the face poset of an homotopy equivalent space as a path poset (e.g.  $P(D_{2,3})$ , cf. Example 6.13).

## REFERENCES

- [ABM05] R. Aharoni, E. Berger, and R. Meshulam. Eigenvalues and homology of flag complexes and vector representations of graphs. *GAFa, Geom. funct. anal.*, 15:555–566, 2005.
- [BBD<sup>+</sup>18] J. Brodzki, F. Belchí, R. Djukanovic, J. Conway, M. Pirashvili, and M. Bennett. Lung topology characteristics in patients with chronic obstructive pulmonary disease. *Scientific Reports*, 8, 03 2018.
- [BBL<sup>+</sup>99] E. Babson, A. Björner, S. Linusson, J. Shareshian, and V. Welker. Complexes of not  $i$ -connected graphs. *Topology*, 38(2):271–299, 1999.
- [Bel] J. Belk. Topological join and wedge sum of spheres. Mathematics Stack Exchange. URL:<https://math.stackexchange.com/q/45036> (version: 2011-06-13).
- [BH98] W. Bruns and J. Herzog. *Cohen-Macaulay Rings*. Cambridge studies in advanced mathematics ; no. 39. Cambridge University Press, Cambridge, 1998.
- [BH17] B. Braun and W. K. Hough. Matching and independence complexes related to small grids. *Electron. J. Comb.*, 24(4):research paper p18, 20, 2017.
- [Bjo84] A. Bjorner. Posets, regular CW complexes and Bruhat order. *European Journal of Combinatorics*, 5(1):7–16, 1984.
- [BMM<sup>+</sup>14] P. Bendich, J. Marron, E. Miller, A. Pieloch, and S. Skwerer. Persistent homology analysis of brain artery trees. *The Annals of Applied Statistics*, 10, 11 2014.
- [BW99] A. Björner and V. Welker. Complexes of directed graphs. *SIAM J. Discrete Math.*, 12:413–424, 10 1999.
- [CCDT21] L. Caputi, C. Collari, and S. Di Trani. Multipath cohomology of directed graphs, 2021. Available on ArXiv:2108.02690.
- [CPH21] L. Caputi, A. Pidnebesna, and J. Hlinka. Promises and pitfalls of topological data analysis for brain connectivity analysis. *NeuroImage*, 238:118245, 2021.
- [CYY01] B. Chen, S.-T. Yau, and Y.-N. Yeh. Graph homotopy and Graham homotopy. *Discrete Mathematics*, 241(1):153–170, 2001. Selected Papers in honor of Helge Tverberg.
- [DEL<sup>+</sup>16] O. Dunaeva, H. Edelsbrunner, A. Lukyanov, M. Machin, D. Malkova, R. Kuvaev, and S. Kashin. The classification of endoscopy images with persistent homology. *Pattern Recognition Letters*, 83:13 – 22, 2016. Geometric, topological and harmonic trends to image processing.
- [DMV17] T. K. Dey, S. Mandal, and W. Varcho. Improved Image Classification using Topological Persistence. In Matthias Hullin, Reinhard Klein, Thomas Schultz, and Angela Yao, editors, *Vision, Modeling and Visualization*. The Eurographics Association, 2017.
- [Gel99] Manin Yu. I. Gelfand, S. I. *Homological algebra*. Springer, Berlin 3B New York, 1999.
- [GGM19] N. Giansiracusa, R. Giansiracusa, and C. Moon. Persistent homology machine learning for fingerprint classification. pages 1219–1226, 12 2019.
- [Gid17] M. Gidea. Topology data analysis of critical transitions in financial networks. *SSRN Electronic Journal*, 01 2017.
- [GLMY20] A. A. Grigor’yan, Yong Lin, Y. V. Muranov, and S.-T. Yau. Path complexes and their homologies. *Journal of Mathematical Sciences*, 248:564–599, 2020.
- [GLS21] D. Govc, R. Levi, and J. P. Smith. Complexes of tournaments, directionality filtrations and persistent homology. *Journal of Applied and Computational Topology*, 5:313–317, 2021.
- [Hap89] D. Happel. Hochschild cohomology of finite-dimensional algebras. *Lecture Notes in Math*, 1404:108–126, 1989.
- [Hat00] A. Hatcher. *Algebraic topology*. Cambridge Univ. Press, Cambridge, 2000.
- [HGR05] L. Helme-Guizon and Y. Rong. A categorification for the chromatic polynomial. *Algebraic & Geometric Topology*, 5(4):1365 – 1388, 2005.
- [Iva94] A. V. Ivashchenko. Contractible transformations do not change the homology groups of graphs. *Discrete Mathematics*, 126(1):159–170, 1994.
- [Kho00] M. Khovanov. A categorification of the Jones polynomial. *Duke Math. J.*, 101(3):359–426, 2000.
- [Koz99] D.N. Kozlov. Complexes of directed trees. *Journal of Combinatorial Theory, Series A*, 88(1):112–122, 1999.
- [Koz08] D. N. Kozlov. *Combinatorial algebraic topology*, volume 21 of *Algorithms and computation in mathematics*. Springer, Berlin
- [Lan02] S. Lang. *Algebra.*, volume 211. New York, NY: Springer, 2002.

- [LCK<sup>+</sup>11] H. Lee, M.K. Chung, H. Kang, B.-N. Kim, and D.S. Lee. Discriminative persistent homology of brain networks. *Proceedings - International Symposium on Biomedical Imaging*, pages 841–844, 03 2011.
- [Mac71] S. MacLane. *Categories for the Working Mathematician*. Springer-Verlag, New York, 1971. Graduate Texts in Mathematics, Vol. 5.
- [Mat19] T. Matsushita. Matching complexes of small grids. *Electron. J. Comb.*, 26(3):research paper p3.1, 8, 2019.
- [Prz10] J. H. Przytycki. When the theories meet: Khovanov homology as Hochschild homology of links. *Quantum Topol.*, 1(2):93–109, 2010.
- [PS18] G. Paolini and M. Salvetti. Weighted sheaves and homology of Artin groups. *Algebraic & Geometric Topology*, 18(7):3943 – 4000, 2018.
- [Rap10] G. Raptis. Homotopy theory of posets. *Homology, Homotopy and Applications*, 12(2):211 – 230, 2010.
- [Rot88] J. Rotman. *An Introduction to Homological Algebra*, volume 119 of *Graduate Texts in Mathematics*. Springer, New York, NY, 1988.
- [TW12] P. Turner and E. Wagner. The homology of digraphs as a generalisation of Hochschild homology. *Journal of Algebra and Its Applications*, 11(02):1250031, 2012.
- [Vas93] V. A. Vassiliev. *Complexes of Connected Graphs*, pages 223–235. Birkhäuser Boston, Boston, MA, 1993.
- [Wac06] M. Wachs. Poset topology: Tools and applications. *Geometric Combinatorics*, 13, 03 2006.
- [Wei94] C. A. Weibel. *An introduction to homological algebra*, volume 38 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1994.
- [Wes05] D. B. West. *Introduction to graph theory*. New Delhi: Prentice-Hall of India, 2005.
- [Wil06] H. S. Wilf. *Generatingfunctionology*. A. K. Peters, Ltd., USA, 2006.

LC: INSTITUTE OF MATHEMATICS, UNIVERSITY OF ABERDEEN, AB24 3UE, UK  
*Email address:* luigi.caputi@abdn.ac.uk

CC: NEW YORK UNIVERSITY ABU DHABI, POBOX 129188, UAE  
*Email address:* carlo.collari@nyu.edu, carlo.collari.math@gmail.com

SDT: DIPARTIMENTO DI MATEMATICA “G. CASTELNUOVO”, SAPIENZA UNIVERSITÀ DI ROMA, IT  
*Email address:* sabino.ditrani@uniroma1.it