p-Laplacian wave equations in non-cylindrical domains *

Lingyang Liu[†] and Hang Gao[‡]

Abstract

This paper is devoted to studying the stability of p-Laplacian wave equations with strong damping in non-cylindrical domains. The method of proof based on some estimates for time-varying coefficients rising from moving boundary and a modified Kormonik inequality. Meanwhile, by selecting appropriate auxiliary functions, finally we obtain the polynomial stability (p > 2) and exponential stability (p = 2) for such systems in some unbounded development domains.

Key Words. p-Laplacian wave equation, stability, strong damping, non-cylindrical domain

1 Introduction and preliminaries

In this paper, we consider the following quasi-linear hyperbolic system with strongly damped terms

$$\begin{cases}
 u_{tt} - \operatorname{div}(|\nabla u|^{p-2}\nabla u) - \Delta u_t = 0 & \text{in } \widehat{Q}, \\
 u = 0 & \text{on } \widehat{\Sigma}, \\
 u(x,0) = u_0(x), \ u_t(x,0) = u_1(x) & \text{in } \Omega_0,
\end{cases}$$
(1.1)

where \widehat{Q} is a non-cylindrical domain in \mathbb{R}^2 , $\widehat{\Sigma}$ is the lateral boundary of \widehat{Q} , $\Omega_t = \widehat{Q} \cap (\mathbb{R} \times \{t\})$, u is the state variable, (u_0, u_1) is any given initial value and $p \geq 2$.

We assume that Q is increasing with time, that is,

$$\Omega_t \subset \Omega_s, \quad t < s$$
 (1.2)

Moreover, we assume the following regular property

if
$$u \in W_0^{1,p}(\Omega)$$
 and $u|_{\Omega-\Omega_t} = 0$, then $u|_{\Omega_t} \in H_0^1(\Omega_t)$. (1.3)

Lemma 1.1 ([2]). Suppose that (1.2) - (1.3) hold. Then given $u_0 \in W_0^{1,p}(\Omega_0)$, $u_1 \in L^2(\Omega_0)$, there exists a function $u: \widehat{Q} \to \mathbb{R}$ such that

$$u \in L^{\infty}(0,T;W_0^{1,p}(\Omega_t)),$$

^{*}This work is supported by the NSF of China under grants 11371084, 11471070 and 11171060.

 $^{^\}dagger$ School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China. E-mail address: liulv938@nenu.edu.cn

[‡]School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China. E-mail address: hanggao2013@126.com

$$u_{t} \in L^{\infty}(0, T; L^{2}(\Omega_{t})) \cap L^{2}(0, T; H_{0}^{1}(\Omega_{t})),$$

$$u(0) = u_{0} \quad and \quad u_{t}(0) = u_{1} \quad a.e. \ in \ \Omega_{0},$$

$$u_{tt} - \operatorname{div}(|\nabla u|^{p-2}\nabla u) - \Delta u_{t} = 0 \quad in \ L^{2}(0, T; W^{-1, p'}(\Omega_{t})).$$

Define the energy of system (1.1) by

$$E(t) = \frac{1}{2}||u_t||_2^2 + \frac{1}{p}||\nabla u||_p^p.$$

Proposition 1.1. E(t) is non-increasing with respect to the time variable.

Proof. It is easy to see that

$$\frac{d}{dt}E(t) = \frac{1}{2}\frac{d}{dt}||u_t||_2^2 + \frac{1}{p}\frac{d}{dt}||\nabla u||_p^p.$$

Since

$$\frac{d}{dt} \int_{\Omega_t} u(x,t) dx = \int_{\Omega_t} \frac{\partial}{\partial t} u(x,t) dx - \int_{\Gamma_t} u(x,t) v_t d\sigma,$$

where $v = (v_x, v_t)^{\top}$ is the external unit normal on $\widehat{\Sigma}$ relative to \widehat{Q} , we have

$$\begin{split} \frac{1}{2} \frac{d}{dt} ||u_t||_2^2 &= \int_{\Omega_t} u_t u_{tt} dx - \frac{1}{2} \int_{\Gamma_t} u_t^2 v_t d\sigma \\ &= \int_{\Omega_t} u_t \left[\operatorname{div} \left(|\nabla u|^{p-2} \nabla u \right) + \Delta u_t \right] dx - \frac{1}{2} \int_{\Gamma_t} u_t^2 v_t d\sigma \\ &= \int_{\Gamma_t} u_t |\nabla u|^{p-2} \nabla u v_x d\sigma - \int_{\Omega_t} |\nabla u|^{p-2} \nabla u \nabla u_t dx \\ &+ \int_{\Gamma_t} u_t \nabla u_t v_x d\sigma - \int_{\Omega_t} |\nabla u_t|^2 - \frac{1}{2} \int_{\Gamma_t} u_t^2 v_t d\sigma, \end{split}$$

and

$$\frac{1}{p}\frac{d}{dt}||\nabla u||_p^p = \int_{\Omega_t} |\nabla u|^{p-2}\nabla u\nabla u_t dx - \frac{1}{p}\int_{\Gamma_t} |\nabla u|^p v_t d\sigma.$$

Hence

$$\frac{d}{dt}E(t) = \int_{\Gamma_t} \left(u_t |\nabla u|^{p-2} \nabla u v_x - \frac{1}{p} |\nabla u|^p v_t - \frac{1}{2} u_t^2 v_t + u_t \nabla u_t v_x \right) d\sigma - \int_{\Omega_t} |\nabla u_t|^2 dx.$$

On the other side, from u = 0 on Σ , one gets $\nabla_{x,t}u = \frac{\partial u}{\partial v}v$, that is, $\nabla u = \frac{\partial u}{\partial v}v_x$, $u_t = \frac{\partial u}{\partial v}v_t$. Besides, noticing that $u_t = 0$ on Σ , we arrive at

$$\frac{d}{dt}E(t) = -\int_{\Omega_t} |\nabla u_t|^2 dx.$$

Lemma 1.2 ([1]). Let $E: \mathbb{R}^+ \to \mathbb{R}^+$ be a non-increasing function and $\phi: \mathbb{R}^+ \to \mathbb{R}^+$ be a strictly increasing C^1 function such that

$$\phi(0) = 0$$
 and $\phi(t) \to +\infty$ as $t \to +\infty$.

If there exist constants $q \ge 0$ and A > 0 such that

$$\int_{S}^{+\infty} E^{q+1}(t)\phi'(t)dt \le \frac{1}{A}E^{q}(0)E(S), \quad 0 \le S < +\infty,$$

then we have

$$E(t) \le E(0) \left(\frac{1+q}{1+qA\phi(t)}\right)^{\frac{1}{q}} \quad \forall t \ge 0, \quad for \ q > 0;$$

and

$$E(t) \le E(0)e^{1-A\phi(t)} \quad \forall t \ge 0, \quad for \ q = 0.$$

Proposition 1.2. For any $t \geq 0$, it holds that

$$||\nabla u(t)||_2^2 \le |\Omega_t|^{1-\frac{2}{p}} ||\nabla u(t)||_p^2$$

$$||u(t)||_2^2 \le |\Omega_t|^{3-\frac{2}{p}} ||\nabla u(t)||_p^2$$

Proof. By Hölder's inequality, one has

$$\int_{\Omega_t} u^2(t)dx = \int_{\Omega_t} \left[\int_0^x u_y(t)dy \right]^2 dx \le |\Omega_t|^2 \int_{\Omega_t} u_x^2(t)dx,$$

that is

$$||u(t)||_2^2 \le |\Omega_t|^2 ||\nabla u(t)||_2^2$$
.

Moreover

$$||\nabla u(t)||_2 = \left[\int_{\Omega_t} |\nabla u(t)|^2 dx \right]^{\frac{1}{2}} \le |\Omega_t|^{\frac{1}{2} - \frac{1}{p}} \left[\int_{\Omega_t} |\nabla u(t)|^p dx \right]^{\frac{1}{p}} = |\Omega_t|^{\frac{1}{2} - \frac{1}{p}} ||\nabla u||_p.$$

Thus

$$||\nabla u(t)||_2^2 \le |\Omega_t|^{1-\frac{2}{p}} ||\nabla u(t)||_p^2$$

$$||u(t)||_2^2 \le |\Omega_t|^{3-\frac{2}{p}}||\nabla u(t)||_p^2.$$

2 Main results and proofs

The main result of this paper is stated as follows.

Theorem 2.1. Let k > 0, $0 < \gamma < 1$ and m > 0. For domain \widehat{Q} with $|\Omega_t| = (1 + kt)^{\frac{1-\gamma}{m}}$, we have (1) when p > 2, and $m \ge \max\left\{2, \frac{(1-\alpha)p}{\alpha p-1}, \frac{(\frac{1}{2}+\alpha)p-1}{(\frac{1}{2}-\alpha)p+1}\right\}$ for $\frac{1}{p} < \alpha < \frac{1}{2} + \frac{1}{p}$, the energy of system (1.1) satisfies

$$E(t) \le \left[\frac{C(1+\beta) \left(E^{\beta}(0) + 1 \right)}{\beta} \right]^{\frac{1}{\beta}} \left[(1+kt)^{\gamma} - 1 \right]^{-\frac{1}{\beta}}, \quad \beta = \frac{p-2}{p};$$

(2) when p = 2, and $m \ge 2$, the energy of system (1.1) satisfies

$$E(t) \le E(0)e^{1-\frac{1}{C}[(1+kt)^{\gamma}-1]}$$
.

Here and henceforth, C is some positive constant.

Proof. Multiply the first equation in (1.1) by $E^{\beta}(t)\phi'(t)u$, and integrating it over $\Omega_t \times (S,T)$, we get

$$0 = \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)u \left(u_{tt} - \operatorname{div}(|\nabla u|^{p-2}\nabla u) - \Delta u_{t}\right) dxdt$$

$$= \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)u u_{tt} dxdt - \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)u \operatorname{div}(|\nabla u|^{p-2}\nabla u) dxdt$$

$$- \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)u \Delta u_{t} dxdt$$

$$\triangleq I_{1} + I_{2} + I_{3}$$

Calculating I_i (i = 1, 2, 3), respectively, we have

$$I_{1} = \int_{\Omega_{T}} E^{\beta}(T)\phi'(T)u(T)u_{t}(T)dx - \int_{\Omega_{S}} E^{\beta}(S)\phi'(S)u(S)u_{t}(S)dx$$
$$-\beta \int_{S}^{T} E^{\beta-1}(t)E'(t) \int_{\Omega_{t}} \phi'(t)uu_{t}dxdt - \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi''(t)uu_{t}dxdt$$
$$-\int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)u_{t}^{2}dxdt,$$

$$I_2 = \int_S^T \int_{\Omega_*} E^{\beta}(t) \phi'(t) |\nabla u|^p dx dt,$$

and

$$I_3 = \int_S^T \int_{\Omega_t} E^{\beta}(t) \phi'(t) \nabla u \nabla u_t dx dt.$$

In view of

$$I_1 + I_2 + I_3 = 0$$
,

one has

$$I_2 = -I_1 - I_3.$$

By the definition of E(t), one gets

$$||\nabla u||_p^p = pE(t) - \frac{p}{2}||u_t||_2^2.$$
(2.1)

Applying (2.1) to I_2 , we obtain

$$\int_{S}^{T} E^{\beta}(t)\phi'(t) \left[pE(t) - \frac{p}{2} \int_{\Omega_{t}} u_{t}^{2} dx \right] dt = -I_{1} - I_{3}.$$

Substituting I_1 and I_3 into the above equality, we get

$$p \int_{S}^{T} E^{\beta+1}(t)\phi'(t)dt$$

$$= \underbrace{\left(\frac{p}{2}+1\right) \int_{S}^{T} E^{\beta}(t)\phi'(t) \int_{\Omega_{t}} u_{t}^{2}dxdt}}_{T_{1}}$$

$$- \int_{\Omega_{T}} E^{\beta}(T)\phi'(T)u(T)u_{t}(T)dx + \int_{\Omega_{S}} E^{\beta}(S)\phi'(S)u(S)u_{t}(S)dx}$$

$$+ \beta \int_{S}^{T} E^{\beta-1}(t)E'(t) \int_{\Omega_{t}} \phi'(t)uu_{t}dxdt + \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi''(t)uu_{t}dxdt$$

$$T_{3}$$

$$- \int_{S}^{T} \int_{\Omega_{t}} E^{\beta}(t)\phi'(t)\nabla u\nabla u_{t}dxdt. \qquad (2.2)$$

We estimate T_i (i = 1, 2, 3, 4, 5), respectively. First, we estimate the value of T_1 . Using Poincaré's inequality, we have

$$T_1 \le \left(\frac{p}{2} + 1\right) \int_S^T E^{\beta}(t) \phi'(t) |\Omega_t|^2 \int_{\Omega_t} |\nabla u_t|^2 dx dt.$$

Recall that ϕ' is a non-negative function in \mathbb{R}^+ . If it satisfies

$$\phi'(t)|\Omega_t|^2 \le C, (2.3)$$

then according to $E'(t) = -\int_{\Omega_t} |\nabla u_t|^2 dx$, we deduce

$$T_1 \le C\left(\frac{p}{2} + 1\right) \int_S^T E^{\beta}(t) \left[-E'(t) \right] dt.$$

Sine E is non-increasing, further

$$T_1 \le C\left(\frac{p}{2} + 1\right) \frac{1}{\beta + 1} E^{\beta + 1}(S).$$
 (2.4)

Secondly, we estimate the value of T_2 . Using Cauchy-Schwarz's inequality and the conclusion in Proposition1.2: $||u(t)||_2^2 \le |\Omega_t|^{3-\frac{2}{p}}||\nabla u(t)||_p^2$, we get

$$T_{2} \leq \int_{\Omega_{T}} \left| E^{\beta}(T)\phi'(T)u(T)u_{t}(T) \right| dx + \int_{\Omega_{S}} \left| E^{\beta}(S)\phi'(S)u(S)u_{t}(S) \right| dx$$

$$\leq E^{\beta}(T) \left(||u_{t}(T)||_{2}^{2} + {\phi'}^{2}(T)||u(T)||_{2}^{2} \right) + E^{\beta}(S) \left(||u_{t}(S)||_{2}^{2} + {\phi'}^{2}(S)||u(S)||_{2}^{2} \right)$$

$$\leq E^{\beta}(T) \left(||u_{t}(T)||_{2}^{2} + {\phi'}^{2}(T)|\Omega_{T}|^{3-\frac{2}{p}} ||\nabla u(T)||_{p}^{2} \right)$$

$$+ E^{\beta}(S) \left(||u_{t}(S)||_{2}^{2} + {\phi'}^{2}(S)|\Omega_{S}|^{3-\frac{2}{p}} ||\nabla u(S)||_{p}^{2} \right).$$

If $\phi'(t)$ satisfies

$$\phi'^{2}(t)|\Omega_{t}|^{3-\frac{2}{p}} \le C,$$
 (2.5)

then by

$$||u_t(t)||_2^2 \le 2E(t)$$
 and $||\nabla u(t)||_p^2 \le p^{\frac{2}{p}} E^{\frac{2}{p}}(t)$,

we obtain

$$T_2 \le C \left[E^{\beta+1}(S) + p^{\frac{2}{p}} E^{\beta+\frac{2}{p}}(S) \right].$$
 (2.6)

Thirdly, we estimate the value of T_3 . By Cauchy-Schwarz's inequality, it follows that

$$T_{3} \leq \beta \int_{S}^{T} E^{\beta-1}(t) \left[-E'(t) \right] \int_{\Omega_{t}} \left| \phi'(t) u u_{t} \right| dx dt$$

$$\leq \beta \int_{S}^{T} E^{\beta-1}(t) \left[-E'(t) \right] \left[\int_{\Omega_{t}} u_{t}^{2} dx + \int_{\Omega_{t}} \phi'^{2}(t) u^{2} dx \right] dt.$$

Under the assumption (2.5), similar to the estimation of T_2 , we derive

$$T_{3} \leq C\beta \left(\int_{S}^{T} E^{\beta}(t) \left[-E'(t) \right] dt - p^{\frac{2}{p}} \int_{S}^{T} E^{\beta + \frac{2}{p} - 1}(t) E'(t) dt \right). \tag{2.7}$$

Next, estimate the value of T_4 . Using Young's inequality with ε , for T_4 , we have the following estimation

$$T_4 \leq \int_S^T E^{\beta}(t) \int_{\Omega_t} |\phi''(t)uu_t| dxdt$$

$$\leq \varepsilon \int_S^T E^{\beta}(t) |\phi''(t)|^{\alpha p} \int_{\Omega_t} |u|^p dxdt + \varepsilon^{-\frac{q}{p}} \int_S^T E^{\beta}(t) |\phi''(t)|^{(1-\alpha)q} \int_{\Omega_t} |u_t|^q dxdt,$$

where $\frac{1}{p} + \frac{1}{q} = 1$, $0 < \alpha < 1$ and $\varepsilon > 0$.

Noticing

$$\int_{\Omega_t} |u|^p dx \le |\Omega_t|^p \int_{\Omega_t} |\nabla u|^p dx,$$

and

$$\int_{\Omega_t} |u_t|^q dx \le |\Omega_t|^q \int_{\Omega_t} |\nabla u_t|^q dx \le |\Omega_t|^{1+\frac{q}{2}} \left(\int_{\Omega_t} |\nabla u_t|^2 dx \right)^{\frac{q}{2}}.$$

we get

$$T_4 \leq \varepsilon p \int_S^T |\phi''(t)|^{\alpha p} |\Omega_t|^p E^{\beta+1}(t) dt + \varepsilon^{-\frac{q}{p}} \int_S^T E^{\beta}(t) |\phi''(t)|^{(1-\alpha)q} |\Omega_t|^{1+\frac{q}{2}} \left[-E'(t) \right]^{\frac{q}{2}} dt.$$

Applying Young's inequality with η to the second term at the right end of the above inequality, we deduce

$$\varepsilon^{-\frac{q}{p}} \int_{S}^{T} E^{\beta}(t) |\phi''(t)|^{(1-\alpha)q} |\Omega_{t}|^{1+\frac{q}{2}} \left[-E'(t) \right]^{\frac{q}{2}} dt
\leq \varepsilon^{-\frac{q}{p}} \eta \int_{S}^{T} \left[E^{\beta}(t) |\phi''(t)|^{(1-\alpha)q} |\Omega_{t}|^{1+\frac{q}{2}} \right]^{\frac{1}{1-\frac{q}{2}}} dt + \varepsilon^{-\frac{q}{p}} \eta^{1-\frac{2}{q}} \int_{S}^{T} \left[-E'(t) \right] dt.$$

If

$$|\phi''(t)|^{\alpha p}|\Omega_t|^p \le C\phi'(t),\tag{2.8}$$

and

$$\left[|\phi''(t)|^{(1-\alpha)q} |\Omega_t|^{1+\frac{q}{2}} \right]^{\frac{1}{1-\frac{q}{2}}} \le C\phi'(t), \tag{2.9}$$

hold, then it follows that

$$T_4 \leq C\varepsilon p \int_S^T E^{\beta+1}(t)\phi'(t)dt + C\varepsilon^{-\frac{q}{p}}\eta \int_S^T E^{\frac{\beta}{1-\frac{q}{2}}}(t)\phi'(t)dt$$
$$+\varepsilon^{-\frac{q}{p}}\eta^{1-\frac{2}{q}} \int_S^T \left[-E'(t)\right]dt. \tag{2.10}$$

Finally, we estimate T_5 . By Young's inequality with δ , one has

$$T_{5} \leq \int_{S}^{T} E^{\beta}(t)\phi'(t) \int_{\Omega_{t}} |\nabla u \nabla u_{t}| \, dxdt$$

$$\leq \int_{S}^{T} E^{\beta}(t)\phi'(t) \left(\delta \int_{\Omega_{t}} |\nabla u|^{p} \, dx + \delta^{-\frac{q}{p}} \int_{\Omega_{t}} |\nabla u_{t}|^{q} \, dx\right) dt$$

$$\leq \delta p \int_{S}^{T} E^{\beta+1}(t)\phi'(t) dt + \delta^{-\frac{q}{p}} \int_{S}^{T} E^{\beta}(t)\phi'(t) \int_{\Omega_{t}} |\nabla u_{t}|^{q} \, dxdt.$$

Using Hölder's inequality, formula $E'(t) = -\int_{\Omega_t} |\nabla u_t|^2 dx$ and Young's inequality with η , from the second term at the right end of the above inequality, one gets

$$\delta^{-\frac{q}{p}} \int_{S}^{T} E^{\beta}(t)\phi'(t) \int_{\Omega_{t}} |\nabla u_{t}|^{q} dx dt$$

$$\leq \delta^{-\frac{q}{p}} \int_{S}^{T} E^{\beta}(t)\phi'(t) |\Omega_{t}|^{1-\frac{q}{2}} \left(\int_{\Omega_{t}} |\nabla u_{t}|^{2} dx \right)^{\frac{q}{2}} dt$$

$$= \delta^{-\frac{q}{p}} \int_{S}^{T} E^{\beta}(t)\phi'(t) |\Omega_{t}|^{1-\frac{q}{2}} \left[-E'(t) \right]^{\frac{q}{2}} dt$$

$$\leq \delta^{-\frac{q}{p}} \eta \int_{S}^{T} \left[E^{\beta}(t) \phi'(t) |\Omega_{t}|^{1-\frac{q}{2}} \right]^{\frac{1}{1-\frac{q}{2}}} dt + \delta^{-\frac{q}{p}} \eta^{1-\frac{2}{q}} \int_{S}^{T} \left[-E'(t) \right] dt.$$

Assume

$$\phi'^{\frac{1}{1-\frac{q}{2}}}(t)|\Omega_t| \le C\phi'(t), \tag{2.11}$$

Then we conclude that

$$T_{5} \leq \delta p \int_{S}^{T} E^{\beta+1}(t)\phi'(t)dt + C\delta^{-\frac{q}{p}}\eta \int_{S}^{T} E^{\frac{\beta}{1-\frac{q}{2}}}(t)\phi'(t)dt + \delta^{-\frac{q}{p}}\eta^{1-\frac{2}{q}} \int_{S}^{T} \left[-E'(t)\right] dt.$$
(2.12)

In order to make (2.3), (2.5), (2.8), (2.9) and (2.11) true, let

$$\phi'(t) \sim |\Omega_t|^{-m}, \quad m \ge 2. \tag{2.13}$$

At this time, it is easy to check that (2.3), (2.5) and (2.11) hold. In the following, for $\phi'(t)$ like (2.13), we verify that (2.8) and (2.9) is also true. Without loss of generality,

let $\phi'(t) = C_1 |\Omega_t|^{-m}$. It is easy to get

$$\phi''(t) = -C_1 m |\Omega_t|^{-m-1} \frac{d}{dt} |\Omega_t| = -C_1^{-\frac{1}{m}} m \phi'^{\frac{m+1}{m}}(t) \frac{d}{dt} |\Omega_t|.$$

Furthermore

$$|\phi''(t)|^{\alpha p}|\Omega_t|^p = C_1^{-\frac{\alpha p}{m}} m^{\alpha p} \phi'^{\left(\frac{m+1}{m}\right)\alpha p}(t) \left| \frac{d}{dt} |\Omega_t| \right|^{\alpha p} |\Omega_t|^p.$$

(2.8) holds, if $\left|\frac{d}{dt}|\Omega_t|\right| \leq C$ and

$$\phi'^{\left[\left(\frac{m+1}{m}\right)\alpha p-1\right]}(t)|\Omega_t|^p \le C.$$

This, by (2.13), is equivalent to

$$|\Omega_t|^{-m\left[\left(\frac{m+1}{m}\right)\alpha p-1\right]+p} \le C.$$

If $|\Omega_t|$ is infinitely increasing, then one has $-m\left[\left(\frac{m+1}{m}\right)\alpha p-1\right]+p\leq 0$. It follows that

$$m \ge \frac{(1-\alpha)p}{\alpha p - 1}, \quad \alpha > \frac{1}{p}.$$

On the other hand,

$$\left[|\phi''(t)|^{(1-\alpha)q}|\Omega_t|^{1+\frac{q}{2}}\right]^{\frac{1}{1-\frac{q}{2}}} = m^{\frac{(1-\alpha)q}{1-\frac{q}{2}}}\phi'^{\left(\frac{m+1}{m}\right)\frac{(1-\alpha)q}{1-\frac{q}{2}}}(t)\left|\frac{d}{dt}|\Omega_t|\right|^{\frac{(1-\alpha)q}{1-\frac{q}{2}}}|\Omega_t|^{\frac{1+\frac{q}{2}}{1-\frac{q}{2}}}.$$

(2.9) holds, if $\left|\frac{d}{dt}|\Omega_t|\right| \leq C$ and

$$\phi'^{\left[\left(\frac{m+1}{m}\right)\frac{(1-\alpha)q}{1-\frac{q}{2}}-1\right]}(t)|\Omega_t|^{\frac{1+\frac{q}{2}}{1-\frac{q}{2}}} \leq C.$$

This, by (2.13), is equivalent to

$$|\Omega_t|^{-m\left[\left(\frac{m+1}{m}\right)\frac{(1-\alpha)q}{1-\frac{q}{2}}-1\right]+\frac{1+\frac{q}{2}}{1-\frac{q}{2}}} \le C.$$

If $|\Omega_t|$ is infinitely increasing, then one has $-m\left[\left(\frac{m+1}{m}\right)\frac{(1-\alpha)q}{1-\frac{q}{2}}-1\right]+\frac{1+\frac{q}{2}}{1-\frac{q}{2}}\leq 0$, $q=\frac{p}{p-1}$. It follows that

$$m \ge \frac{\left(\frac{1}{2} + \alpha\right)p - 1}{\left(\frac{1}{2} - \alpha\right)p + 1}, \quad \alpha < \frac{1}{2} + \frac{1}{p}.$$

To sum up, when

$$m \ge \max\left\{2, \frac{(1-\alpha)p}{\alpha p - 1}, \frac{\left(\frac{1}{2} + \alpha\right)p - 1}{\left(\frac{1}{2} - \alpha\right)p + 1}\right\}, \quad \frac{1}{p} < \alpha < \frac{1}{2} + \frac{1}{p},\tag{2.14}$$

 $\phi'(t)$ like(2.13) satisfies (2.3), (2.5), (2.8), (2.9) and (2.11).

Let $\frac{\beta}{1-\frac{q}{2}} = \beta + 1$ and ε , η , δ be sufficiently small in (2.10) and (2.12), combining (2.2), (2.4), (2.6) and (2.7), we end up with

$$\int_{S}^{T} E^{\beta+1}(t)\phi'(t)dt \le C \left[E^{\beta+1}(S) + E(S) \right] \le C \left[E^{\beta}(0) + 1 \right] E(S).$$

Let $T \to +\infty$, we get

$$\int_{S}^{+\infty} E^{\beta+1}(t)\phi'(t)dt \le C\left[E^{\beta+1}(S) + E(S)\right] \le \frac{C\left[E^{\beta}(0) + 1\right]}{E^{\beta}(0)}E^{\beta}(0)E(S).$$

By Lemma 1.2 and choosing $\phi(t)=(1+kt)^{\gamma}-1,\ k>0,\ 0<\gamma<1,$ one obtains

$$E(t) \le E(0) \left(\frac{1+\beta}{1+\beta A\phi(t)} \right)^{\frac{1}{\beta}} \le E(0) \left(\frac{1+\beta}{\beta A} \right)^{\frac{1}{\beta}} \phi(t)^{-\frac{1}{\beta}},$$

where $A = \frac{E^{\beta}(0)}{C(E^{\beta}(0)+1)}$.

More precisely,

$$E(t) \le \left[\frac{C(1+\beta) \left(E^{\beta}(0) + 1 \right)}{\beta} \right]^{\frac{1}{\beta}} \left[(1+kt)^{\gamma} - 1 \right]^{-\frac{1}{\beta}}, \quad \beta = \frac{p-2}{p}, \quad p > 2.$$

At this time, $\phi'(t) = k\gamma(1+kt)^{\gamma-1}$. Letting $\phi'(t) = k\gamma|\Omega_t|^{-m}$, one has

$$|\Omega_t| = (1 + kt)^{\frac{1 - \gamma}{m}},$$

where m is given in (2.14).

If p=2, then let $\beta=0$. In this case, it is required that $\phi'(t)$ satisfies

$$\phi'(t)|\Omega_t|^2 \le C, (2.15)$$

$$\phi'^2(t)|\Omega_t|^2 \le C, (2.16)$$

$$|\phi''(t)|^{2\alpha}|\Omega_t|^2 \le C\phi'(t),$$
 (2.17)

$$|\phi''(t)|^{2(1-\alpha)}|\Omega_t|^2 \le C, (2.18)$$

where (2.18) is a little different compared to the former (p > 2), but it is more simpler.

Let $\phi'(t) \sim |\Omega_t|^{-m}$, $m \geq 2$. It is easy to see that $\phi'(t)$ satisfies (2.15) and (2.16). If $\left|\frac{d}{dt}|\Omega_t|\right| \leq C$, then $\phi''(t) \sim \phi'^{\frac{m+1}{m}}(t)$.

(2.17) holds, if

$$\phi'(t)^{\left[2\alpha\left(\frac{m+1}{m}\right)-1\right]}|\Omega_t|^2 \le C.$$

Further

$$|\Omega_t|^{-m\left[2\alpha\left(\frac{m+1}{m}\right)-1\right]+2} \le C.$$

If $|\Omega_t|$ is infinitely increasing, then one has $-m\left[2\alpha\left(\frac{m+1}{m}\right)-1\right]+2\leq 0$. It follows that

$$m \ge \frac{2(1-\alpha)}{2\alpha-1}, \quad \alpha > \frac{1}{2}.$$

(2.18) holds, if

$$\phi'(t)^{2(1-\alpha)\left(\frac{m+1}{m}\right)}|\Omega_t|^2 \le C.$$

Similarly, we get

$$m \ge \frac{\alpha}{1-\alpha}$$
.

Let $\alpha = \frac{2}{3}$. We deduce for $m \geq 2$, it holds that

$$\int_{S}^{+\infty} E(t)\phi'(t)dt \le CE(S).$$

Also choosing $\phi(t) = (1 + kt)^{\gamma} - 1$, k > 0, $0 < \gamma < 1$, and by Lemma 1.2, we obtain

$$E(t) \le E(0)e^{1-\frac{1}{C}[(1+kt)^{\gamma}-1]}.$$

At this time, $\phi'(t) = k\gamma(1+kt)^{\gamma-1}$. Letting $\phi'(t) = k\gamma|\Omega_t|^{-m}$, we see that

$$|\Omega_t| = (1+kt)^{\frac{1-\gamma}{m}}, \quad m \ge 2.$$

Remark 2.1. If $|\Omega_t|$ is bounded, then

(1) for p > 2, choosing $\phi(t) = t$, one has

$$E(t) \le E(0) \left(\frac{1+\beta}{1+\beta At}\right)^{\frac{1}{\beta}}, \quad \beta = \frac{p-2}{p};$$

(2) for p = 2, putting $\beta = 0$, and also choosing $\phi(t) = t$, one gets

$$E(t) \le E(0)e^{1-\frac{1}{C}t}.$$

3 Further works

In this article, we get a decay estimate $E(t) \leq E(0)e^{1-\frac{1}{C}[(1+kt)^{\gamma}-1]}$ for the energy of system (1.1) in the case of domains with $|\Omega_t| = (1+kt)^{\frac{1-\gamma}{m}}$, k > 0, $0 < \gamma < 1$, $m \geq 2$. However, the method we adopt here can not be applied to the case of domains with $|\Omega_t| = (1+kt)^{\alpha}$, $\alpha > \frac{1}{2}$. In a forthcoming paper, we are going to study the stability of p-Laplacian wave equations in a more general domain.

References

- [1] Benaissa A, Mokeddem S, Decay estimates for the wave equation of p-Laplacian type with dissipation of m-Laplacian type, Math. Methods Appl. Sci. 30 (2007), no. 2, 237–247.
- [2] Ferreira J, Raposo C A, Santos M L, Global existence for a quasilinear hyperbolic equation in a noncylindrical domain, Int. J. Pure Appl. Math. 29 (2006), no. 4, 457–467.