

Analytic derivation of the non-linear gluon distribution function

G.R.Boroun*

Department of Physics, Razi University, Kermanshah 67149, Iran

(Dated: January 24, 2022)

In the present article, two analytical solutions based on the Laplace transforms method for the linear and non-linear gluon distribution functions have been presented at low values of x . These linear and non-linear methods are presented based on the solutions of the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) evolution equation and the Gribov-Levin-Ryskin Mueller-Qiu (GLR-MQ) equation at the leading-order accuracy in perturbative QCD respectively. The gluon distributions are obtained directly in terms of the parametrization of structure function $F_2(x, Q^2)$ and its derivative and compared with the results from the parametrization models. The n_f changes at the threshold are considered in the numerical results. The effects of the non-linear corrections are visible as Q^2 decreases and vanish as Q^2 increases. The nonlinear corrections tame the behavior of the gluon distribution function at low x and Q^2 in comparison with the parametrization models.

I. Introduction

In recent years, the study and consider of solutions of the Dokshitzer-Gribov- Lipatov-Altarelli-Parisi (DGLAP) evolution equations [1] based on Laplace transforms method have been considered by many authors [2-4]. Firstly authors in Ref.[5] showed that it is possible to solve the leading order (LO) DGLAP evolution equations directly from the parametrization of the proton structure function $F_2(x, Q^2)$ based on Laplace transforms method. These evolution equations in Ref.[5] are a set of integro-differential equations which can be used to evolve the quark and gluon distributions. The parton distribution functions in hadrons play a key role in the Standard Model processes and searches for new physics in future accelerators. The structure function of the proton measured experimentally in deep inelastic scattering processes then traditionally the gluon and quark distribution functions where have been determined simultaneously by fitting experimental data on the proton structure function at small values of the Bjorken variable x . At LO approximation the proton structure function is expressed through the quark density as $F_2(x, Q^2) = \sum_{i=1}^{n_f} e_i^2 x [q(x, Q^2) + \bar{q}(x, Q^2)]$, where n_f is the number of flavors.

Authors in Ref.[5] derived an explicit expression for the gluon distribution function $G(x, Q^2) = xg(x, Q^2)$ in the proton in terms of the proton structure function $F_2(x, Q^2)$ by solving the LO DGLAP equation for the Q^2 evolution of $F_2(x, Q^2)$ analytically. In particular, accurate knowledge of gluon distribution functions at small x and small Q^2 will play a vital role in the electron-proton future colliders. Indeed, the non-linear corrections (NLC) play an important role in the small x and

small Q^2 regions at the Large Hadron electron Collider (LHeC) and Future Circular Collider hadron-electron (FCC-he) [6]. The non-linear corrections of the gluon recombination to the parton distributions have been calculated by Gribov-Levin- Ryskin (GLR) and Mueller-Qiu (MQ) in [7] based on the Abramovsky-Gribov-Kancheli (AGK) cutting rules in the double leading logarithmic approximation (DLLA). It is known that the gluon recombination effects reduce the growth of the gluon distribution function, therefore cannot be negligible at the small x and Q^2 regions. Indeed all possible $g + g \rightarrow g$ ladder recombinations are resummed to leading order of the parameter $\alpha_s \ln(1/x) \ln(Q^2/Q_0^2)$ where leads to saturation of the gluon density at this region.

In the following sections, we present two analytic methods where determine $G(x, Q^2)$ directly from $F_2^p(x, Q^2)$ and its derivative into $\ln Q^2$. The first one is a review for the linear evolution equations based on the Laplace transforms method for the DGLAP evolution equation. The second one is the same method for the GLR-MQ equation which presents the non-linear corrections to the gluon distribution function directly from the proton structure function and its derivative. These methods lead to equivalent results without the intervening differential equation.

The non-linear corrections emerge from the recombination of two gluon ladders where modify the evolution equation of singlet quark distribution by an extra non-linear term. This non-linear term add to the linear DGLAP evolution equation by the following form

$$\frac{\partial F_2(x, Q^2)}{\partial \ln Q^2} = \frac{\partial F_2(x, Q^2)}{\partial \ln Q^2} \Big|_{DGLAP} - \langle e^2 \rangle \frac{27\alpha_s^2(Q^2)}{160R^2 Q^2} [xg(x, Q^2)]^2 + HT, \quad (1)$$

*Electronic address: grboroun@gmail.com; boroun@razi.ac.ir

where

$$\frac{\partial F_2(x, Q^2)}{\partial \ln Q^2} \Big|_{DGLAP} = \frac{\alpha_s(Q^2)}{4\pi} [P_{qq}(x) \otimes F_2(x, Q^2) + \langle e^2 \rangle P_{qg}(x) \otimes G(x, Q^2)] \quad (2)$$

The splitting functions P_{ij} are the Altarelli-Parisi splitting kernels at one loop correction, and $\langle e^2 \rangle$ is the average of the charge e^2 for the active quark flavors, $\langle e^2 \rangle = n_f^{-1} \sum_{i=1}^{n_f} e_i^2$. We take the $n_f = 3$ for $\mu^2 < m_c^2$, $n_f = 4$ for $m_c^2 < \mu^2 < m_b^2$ and $n_f = 5$ for $\mu^2 > m_b^2$ and adjust the QCD parameter Λ at each heavy quark mass threshold. The correlation length \mathcal{R} determine the size of the non-linear term. This value depends on how the gluon ladders are coupled to the nucleon or on how the gluons are distributed within the nucleon. The \mathcal{R} is approximately equal to $\simeq 5 \text{ GeV}^{-1}$ if gluons are populated across the proton and it is equal to $\simeq 2 \text{ GeV}^{-1}$ if gluons have the hotspot like structure. Here the higher dimensional gluon distribution (i.e., higher twist) is assumed to be zero.

Now we review the method of extracting the gluon distribution from the parametrization of the proton structure function and its derivative in the linear and non-linear corrections using the Laplace transforms method in next sections respectively.

II. Linear Formalism

By considering the variable changes $\nu \equiv \ln(1/x)$ and $w \equiv \ln(1/y)$, one can rewrite the DGLAP evolution equation (i.e., Eq.2) in s -space as

$$\frac{\partial f_2(s, Q^2)}{\partial \ln Q^2} = \Phi_f(s, Q^2) f_2(s, Q^2) + \langle e^2 \rangle \Theta_f(s, Q^2) g(s, Q^2), \quad (3)$$

where the Laplace-transform of the distribution functions read

$$\begin{aligned} \mathcal{L}[\widehat{\mathcal{F}}_2(\nu, Q^2); s] &= f_2(s, Q^2), \\ \mathcal{L}[\widehat{G}(\nu, Q^2); s] &= g(s, Q^2) \end{aligned} \quad (4)$$

where

$$\begin{aligned} \widehat{\mathcal{F}}_2(\nu, Q^2) &= F_2(e^{-\nu}, Q^2), \\ \widehat{G}(\nu, Q^2) &= G(e^{-\nu}, Q^2). \end{aligned} \quad (5)$$

and the coefficient functions Φ and Θ in s -space are given by

$$\begin{aligned} \Theta_f(s, Q^2) &= n_f \frac{\alpha_s(Q^2)}{2\pi} \left(\frac{1}{1+s} - \frac{2}{2+s} + \frac{2}{3+s} \right), \\ \Phi_f(s, Q^2) &= \frac{\alpha_s(Q^2)}{4\pi} \left[4 - \frac{8}{3} \left(\frac{1}{1+s} + \frac{1}{2+s} \right) + 2S(s) \right]. \end{aligned} \quad (6)$$

In the above equation the quantity $S(s)$ is related with the Euler $\Psi(s+1)$ function as $S(s) = \Psi(s+1) + \gamma_E$ where $\Psi(s)$ is defined by $\Psi(s) = \frac{d}{ds} \ln \Gamma(s)$. Here $\Psi(x)$ is the digamma function and $\gamma_E = 0.5772156..$ is Euler constant. We introduce the notion of the so-called nested sums [8] throughout the rest of the paper where the function $S(s)$ is defined

$$S(s) = -\ln(2) - \sum_{l=0}^{\infty} \frac{(-1)^{l+1}}{s+l+1}. \quad (7)$$

In Fig.1 we consider the expansion of Eq.(7) by using




FIG. 1: Sensitivity of the function $S(s)$ verses s for different values of l .

different points. We observe that for $l \geq 10$, the results are almost equivalent and have the smooth behavior for $s > 10$. In this article we shall widely use the notation $l = 10$.

The LO solution of the gluon distribution in s -space in Eq.(3) reads

$$g(s, Q^2) = k(s, Q^2) Df_2(s, Q^2) - h(s, Q^2) f_2(s, Q^2) \quad (8)$$

where

$$\begin{aligned} Df_2(s, Q^2) &= \partial f_2(s, Q^2) / \partial \ln Q^2, \\ k(s, Q^2) &= 1 / (\langle e^2 \rangle \Theta_f(s, Q^2)), \\ h(s, Q^2) &= \Phi_f(s, Q^2) / (\langle e^2 \rangle \Theta_f(s, Q^2)). \end{aligned} \quad (9)$$

The inverse Laplace transform of coefficients k and h in Eq.(9) are defined by the following forms

$$\begin{aligned}
k(\nu, Q^2) &\equiv \mathcal{L}^{-1}[k(s, Q^2); \nu] = \frac{\pi}{2 \langle e^2 \rangle \alpha_s} \{ \delta'(\nu) + 3\delta(\nu) - \exp(-\frac{3}{2}\nu) [2 \cos(\frac{1}{2}\sqrt{7}\nu) + \frac{6}{7}\sqrt{7} \sin(\frac{1}{2}\sqrt{7}\nu)] \}, \\
h(\nu, Q^2) &\equiv \mathcal{L}^{-1}[h(s, Q^2); \nu] = \frac{1}{\langle e^2 \rangle} \{ -(\frac{1}{2} + \frac{2}{3} \ln 2) \delta'(\nu) - (\frac{1}{6} + 2 \ln 2) \delta(\nu) + \exp(-\frac{3}{2}\nu) [3.606 \cos(\frac{1}{2}\sqrt{7}\nu) \\
&\quad + 1.371 \sin(\frac{1}{2}\sqrt{7}\nu)] + \frac{1}{2} \exp(-4\nu) - \frac{8}{7} \exp(-5\nu) + \frac{20}{11} \exp(-6\nu) - \frac{5}{2} \exp(-7\nu) + \frac{35}{11} \exp(-8\nu) \\
&\quad - \frac{112}{29} \exp(-9\nu) + \frac{168}{37} \exp(-10\nu) - \frac{120}{23} \exp(-11\nu) \}. \tag{10}
\end{aligned}$$

Finally the gluon distribution function directly is obtained from the parameterization of the structure function

$F_2(x, Q^2)$ and its derivatives by the following form

$$\begin{aligned}
G(x, Q^2) &= \frac{\pi}{2 \langle e^2 \rangle \alpha_s} \{ \frac{\partial DF_2(x, Q^2)}{\partial \ln x} + 3DF_2(x, Q^2) - \int_x^1 \frac{dy}{y} DF_2(y, Q^2) (\frac{x}{y})^{\frac{3}{2}} [2 \cos(\frac{1}{2}\sqrt{7} \ln(\frac{y}{x})) + \frac{6}{7}\sqrt{7} \sin(\frac{1}{2}\sqrt{7} \ln(\frac{y}{x}))] \} \\
&\quad - \frac{1}{\langle e^2 \rangle} \{ (\frac{1}{2} + \frac{2}{3} \ln 2) \frac{\partial F_2(x, Q^2)}{\partial \ln x} + (\frac{1}{6} + 2 \ln 2) F_2(x, Q^2) - \int_x^1 \frac{dy}{y} F_2(y, Q^2) (\frac{x}{y})^{\frac{3}{2}} [3.606 \cos(\frac{1}{2}\sqrt{7} \ln(\frac{y}{x})) \\
&\quad + 1.371 \sin(\frac{1}{2}\sqrt{7} \ln(\frac{y}{x}))] - \int_x^1 \frac{dy}{y} F_2(y, Q^2) (\frac{1}{2}(\frac{x}{y})^4 - \frac{8}{7}(\frac{x}{y})^5 + \dots - \frac{120}{23}(\frac{x}{y})^{11}) \}. \tag{11}
\end{aligned}$$

The parameterization of the structure function $F_2(x, Q^2)$, which describes fairly well the available experimental data [9] on the reduced cross sections in a full accordance with the Froissart predictions in a range of the kinematical variables x and Q^2 , $x \leq 0.1$ and $0.15 \text{ GeV}^2 < Q^2 < 3000 \text{ GeV}^2$, suggested by authors in Ref.[10]. This parametrization and its derivative read

$$F_2(x, Q^2) = D(Q^2)(1-x)^n \sum_{m=0}^2 A_m(Q^2) L^m, \tag{12}$$

and

$$\begin{aligned}
DF_2(x, Q^2) \equiv \frac{\partial F_2(x, Q^2)}{\partial \ln Q^2} &= F_2(x, Q^2) \left[\frac{\partial \ln D(Q^2)}{\partial \ln Q^2} \right. \\
&\quad \left. + \frac{\partial \ln \sum_{m=0}^2 A_m(Q^2) L^m}{\partial \ln Q^2} \right],
\end{aligned}$$

where the effective parameters are defined in Appendix A and Table I. Consequently, one can obtain the gluon distribution into the effective coefficients obtained from a combined fit of the H1 and ZEUS collaborations data.

III. Non-Linear Formalism

Now, the above procedure is used to derive the non-linear corrections to the gluon distribution function directly from the parameterization of the proton structure

function and its derivatives. The resulting modified structure function should now be driven by non-linear DGLAP evolution in s -space in a limited approach. In ν -space, we have defined the Laplace transform of the $\mathcal{L}[\widehat{G}^2(\nu, Q^2); s]$ to be less than $g^2(s, Q^2) \equiv [G(s, Q^2)]^2$. Indeed $\mathcal{L}[\widehat{G}^2(\nu, Q^2); s] < \mathcal{L}[\widehat{G}(\nu, Q^2); s]^2$.¹ Therefore in this limit, we take the Laplace transform of (1), by follow

$$\begin{aligned}
\frac{\partial f_2(s, Q^2)}{\partial \ln Q^2} &\simeq \Phi_f(s) f_2(s, Q^2) + \langle e^2 \rangle \Theta_f(s) g(s, Q^2) \\
&\quad - \langle e^2 \rangle \zeta g^2(s, Q^2), \tag{13}
\end{aligned}$$

¹ The standard parametrization of the gluon distribution function at low x introduced by

$$G(x, Q^2) = f(Q^2)x^{-\delta}$$

where the low x behavior could well be more singular. By considering the variable change $\nu \equiv \ln(1/x)$, one can rewrite the gluon distribution in s -space as

$$\begin{aligned}
\mathcal{L}[\widehat{G}^2(\nu, Q^2); s] &= \frac{f(Q^2)^2}{(s-2\delta)}, \\
\mathcal{L}[\widehat{G}(\nu, Q^2); s]^2 &= \frac{f(Q^2)^2}{(s-\delta)^2}.
\end{aligned}$$

We observe that the function $\mathcal{L}[\widehat{G}^2(\nu, Q^2); s]$ is always lower than $\mathcal{L}[\widehat{G}(\nu, Q^2); s]^2$ for low s values in a wide range of Q^2 values. According to this result, we use from this limited approach for solving the quadratic equation in s -space.

where $\zeta = \frac{27\alpha_s^2(Q^2)}{160\mathcal{R}^2Q^2}$. The non-linear gluon distribution function is defined by a quadratic equation in s -space in the following form

$$g^2(s, Q^2) - k(s, Q^2)g(s, Q^2) + h(s, Q^2) = 0, \quad (14)$$

where

$$\begin{aligned} h(s, Q^2) &= \frac{1}{\langle e^2 \rangle \zeta} [Df_2(s, Q^2) - \Phi_f(s, Q^2)f_2(s, Q^2)], \\ k(s, Q^2) &= \Theta_f(s, Q^2)/\zeta. \end{aligned} \quad (15)$$

One can easily solve this equation and extract the non-linear gluon distribution in s -space as

$$g(s, Q^2) = \frac{1}{2}k(s, Q^2)[1 \pm (1 - \frac{4h(s, Q^2)}{k^2(s, Q^2)})^{1/2}]. \quad (16)$$

The quadratic equation (16) has two roots, which the negative root reads

$$\begin{aligned} g(s, Q^2) &= \frac{h(s, Q^2)}{k(s, Q^2)}[\text{Linear Term}] + \left\{ \frac{h^2(s, Q^2)}{k^3(s, Q^2)} \right. \\ &\quad \left. + 2\frac{h^3(s, Q^2)}{k^5(s, Q^2)} + \dots \right\}[\text{Non - Linear Terms}]. \end{aligned} \quad (17)$$

The non-linear terms in Eq.(17) are coefficients in the form $\sum_{n=1} \zeta^n$, where ζ is around the order $\mathcal{O}(\sim 10^{-3})$ at $Q = 1$ GeV and $\mathcal{R} = 2$ GeV $^{-1}$, so this series is convergent when $n \rightarrow \infty$.

We now re-derive our analytic solution using the inverse Laplace transforms method for $g(s, Q^2)$ (i.e., Eq.(17)). The inverse Laplace transform of terms in Eq.(17) are given by the following form

$$\begin{aligned} G^{\text{NLC}}(x, Q^2) &= \text{Eq.(10)} + \mathcal{L}^{-1}\left[\frac{h^2(s, Q^2)}{k^3(s, Q^2)} \right. \\ &\quad \left. + 2\frac{h^3(s, Q^2)}{k^5(s, Q^2)} + \dots; \nu\right]. \end{aligned} \quad (18)$$

Therefore the non-linear corrections to the gluon distribution function due to the Laplace transforms method is defined directly from the parametrization of the structure function F_2 and its derivative. Then we consider the positive roots for the non-linear corrections to the gluon distribution in Eq.(16). For $Q^2 < 2$ GeV 2 in the range $10^{-5} < x < 10^{-1}$, the positive roots are dominant in Eq.(16). In this domain, the non-linear gluon distribu-




FIG. 2: The shaded area represents the convergence of positive and negative roots in Eq.(16).

tion is defined by

$$\begin{aligned} G^{\text{NLC}}(x, Q^2) &= \mathcal{L}^{-1}\left[k(s, Q^2) - \frac{h(s, Q^2)}{k(s, Q^2)} - \frac{h^2(s, Q^2)}{k^3(s, Q^2)} \right. \\ &\quad \left. - 2\frac{h^3(s, Q^2)}{k^5(s, Q^2)} + \dots; \nu\right] \\ &= \frac{2\alpha_s}{\pi\zeta} \int_x^1 \frac{dy}{y} \left[\left(\frac{x}{y}\right) - 2\left(\frac{x}{y}\right)^2 + 2\left(\frac{x}{y}\right)^3 \right] \\ &\quad - \text{Eq.(10)} - \mathcal{L}^{-1}\left[\frac{h^2(s, Q^2)}{k^3(s, Q^2)} \right. \\ &\quad \left. + 2\frac{h^3(s, Q^2)}{k^5(s, Q^2)} + \dots; \nu\right]. \end{aligned} \quad (19)$$

In next section, we will describe the numerical results for $G^{\text{NLC}}(x, Q^2)$ by using the analytical solutions, (i.e., Eqs.(18) and (19), at low and high Q^2 values.

IV. Results and Discussions

Now we use the LO approximation of $\alpha_s(Q^2)$ which is defined by

$$\alpha_s(Q^2) = \frac{4\pi}{(11 - \frac{2}{3}n_f)\ln(Q^2/\Lambda^2)},$$

with $n_f = 5$ and $\Lambda_5 = 80.80$ MeV for $Q > m_b$, $n_f = 4$ and $\Lambda_4 = 136.80$ MeV for $m_c < Q < m_b$, and $n_f = 3$ and $\Lambda_3 = 136.80$ MeV for $Q < m_c$ where Λ 's have been extracted with $\alpha_s(m_z^2) = 0.1166$ at the Z -boson mass. By using the Eqs.(11), (18) and (19), we can extract numerically the linear and non-linear gluon distribution inside the proton from the parameterization of the proton structure function and its derivatives. Then

compare them with the results in Refs.[11] and [12].

The results of the calculations based on the linear gluon distribution (i.e., Eq.(11)) are shown in Fig.3. In this figure the straight lines represent the solutions resulted from the Laplace transform technique. We take the $n_f = 3$ for $Q < 1.3$ GeV, $n_f = 4$ for $Q < 4.5$ GeV and $n_f = 5$ for $Q > 4.5$ GeV as the gluon distribution function $G(x, Q^2)$ depends on n_f . In Fig.4, the results for the linear gluon distributions have been shown and compared with the parametrization methods in Refs.[11,12] for $Q^2 = 10$ and 100 GeV². For $Q^2 = 10$ GeV² these results compared with $G_{n_f=4}(x, Q^2)$ in Refs.[11] and [12]². For $Q^2 = 100$ GeV² where the active flavor is $n_f = 5$, we compared our results with those obtained in Ref.[11]³. This figure indicate that the obtained results from the present analysis, based on the Laplace transform technique using the number of active flavors, are in good agreements with the ones obtained from the parametrization methods.

In Fig.5 we plot the Q^2 dependence of the non-linear corrections to the gluon distribution for $\mathcal{R} = 2$ GeV⁻¹ at some representative x and check the compatibility of the non-linear results with the linear gluon distributions. In this figure (i.e., Fig.5) we can see that at large scales, the nonlinear corrections (due to the $1/Q^2$ dependence) relax into the linear gluon distribution functions. Also the non-linear corrections play an important role on gluon distribution as x and Q^2 decreases. In Fig.5 we observe that the dot line separates linear and non-linear behaviors of the gluon distributions. It is seen that these distributions change discontinuously at each threshold but remain constant between thresholds. These results are comparable with Eskola-Honkanen-Kolhinen-Qiu-Salgado (EHKQS), which obtained [14-15] the parton distribution functions using the CTEQ6L [16] with respect to the non-linear GLRMQ evolution equations. At low scales $Q^2 < 2$ GeV², where the positive roots are dominate in the non-linear distributions, we obtained the non-linear behavior of the gluon distribution in Fig.6. In this figure (i.e., Fig.6), we show the non-linear

corrections to the gluon distribution determined from Eqs.(18) and (19) as a function of x for two different values of Q , namely $Q = 1.14$ GeV and 1.30 GeV with respect to the Lpalce transforms method. In these calculations, we use the non-linear gluon distribution with $n_f = 3$ for $Q = 1.14$ GeV and compared the obtained results with the parametrization method in Ref.[11] at the same active flavor number. Also the charm threshold is shown for $Q = 1.30$ GeV with $n_f = 3$ and 4 in Fig.6 and compared the obtained results with the parametrization method in Ref.[11] at each active flavor number. A depletion occurs at $x \leq 10^{-4}$ where these results show that the non-linear behavior of the gluon distribution function is tamed with respect to the positive roots. This taming behavior of non-linear gluon distribution function towards low x at low Q^2 values become significant at the hot spot point. Further the computed values of the gluon distribution with non-linear effects play an increasingly important role at low x and low Q^2 values.

In conclusion, we have presented a certain theoretical model to describe the non-linear corrections to the gluon distribution function based on the Laplace transforms method at low values of x and Q^2 in a limited approach. A detailed analysis has been performed to find an analytical solution of the linear and non-linear gluon distribution functions from the proton structure function and its derivative. The effect of non-linear corrections on the behavior of $G(x, Q^2)$ with decreasing Q^2 become significant at the hot spot point. At high Q^2 values the non-linear corrections relax into the linear gluon distribution function. The nonlinear corrections have been tamed the behavior of the gluon distribution function at $Q^2 < 2$ GeV² in comparison with the linear behavior.

² In Ref.[11] the gluon distribution for $n_f = 4$ is just $\frac{3}{5}G_{n_f=3}(x, Q^2)$, where $G_{n_f=3}(x, Q^2)$ is obtained from a fit to ZEUS data [13] into an expression in both $\ln(Q^2)$ and $\ln(1/x)$ to include the effects of heavy-quark masses. In Ref.[12] authors obtained an analytical solution for $G(x, Q^2)$ using a Froissart bounded structure function for $0 < x \lesssim 0.09$. Those obtained a simple quadratic polynomial in $\ln(1/x)$ with quadratic polynomial coefficients in $\ln(Q^2)$.

³ In Ref.[11] authors obtained the gluon distribution $G(x, Q^2)$ for 5 active quarks (for massless u, d, s and massive c, b quarks) into the massless gluon distribution $G_{n_f=3}(x, Q^2)$, as $G_{n_f=5}(x, Q^2) = \frac{6}{11}G_{n_f=3}(x, Q^2)$. Also authors obtained an excellent fit to the gluon distribution for $n_f = 5$ using a quadratic expression in $\ln 1/x$ and a much more complicated power series in $\ln(Q^2)$ for $x \lesssim 0.05$.

ACKNOWLEDGMENTS

I thank the respectable referee for giving the main idea of this work. I sincerely thank the referee for his/her invaluable comments during the review process one of them my papers in EPJC. The feedback from the referee was very important for me.

Appendix A

The explicit expression for the proton structure function suggested in Ref.[10] is defined by the following form

$$F_2^{\gamma p}(x, Q^2) = D(Q^2)(1-x)^n [C(Q^2) + A(Q^2) \ln(\frac{1}{x} \frac{Q^2}{Q^2 + \mu^2}) + B(Q^2) \ln^2(\frac{1}{x} \frac{Q^2}{Q^2 + \mu^2})], \quad (20)$$

where

$$\begin{aligned} A(Q^2) &= a_0 + a_1 \ln(1 + \frac{Q^2}{\mu^2}) + a_2 \ln^2(1 + \frac{Q^2}{\mu^2}), \\ B(Q^2) &= b_0 + b_1 \ln(1 + \frac{Q^2}{\mu^2}) + b_2 \ln^2(1 + \frac{Q^2}{\mu^2}), \\ C(Q^2) &= c_0 + c_1 \ln(1 + \frac{Q^2}{\mu^2}), \\ D(Q^2) &= \frac{Q^2(Q^2 + \lambda M^2)}{(Q^2 + M^2)^2}. \end{aligned} \quad (21)$$

Here M is the effective mass and μ^2 is a scale factor. The additional parameters with their statistical errors are given in Table I.

TABLE I: The effective parameters at low x for $0.15 \text{ GeV}^2 < Q^2 < 3000 \text{ GeV}^2$ provided by the following values. The fixed parameters are defined by the Block-Halzen fit to the real photon-proton cross section as $M^2 = 0.753 \pm 0.068 \text{ GeV}^2$, $\mu^2 = 2.82 \pm 0.290 \text{ GeV}^2$ and $c_0 = 0.255 \pm 0.016$ [10].

parameters	value
a_0	$8.205 \times 10^{-4} \pm 4.62 \times 10^{-4}$
a_1	$-5.148 \times 10^{-2} \pm 8.19 \times 10^{-3}$
a_2	$-4.725 \times 10^{-3} \pm 1.01 \times 10^{-3}$
b_0	$2.217 \times 10^{-3} \pm 1.42 \times 10^{-4}$
b_1	$1.244 \times 10^{-2} \pm 8.56 \times 10^{-4}$
b_2	$5.958 \times 10^{-4} \pm 2.32 \times 10^{-4}$
c_1	$1.475 \times 10^{-1} \pm 3.025 \times 10^{-2}$
n	11.49 ± 0.99
λ	2.430 ± 0.153
$\chi^2(\text{goodness of fit})$	0.95

REFERENCES

1. L.N. Lipatov, Sov. J. Nucl. Phys.**20**, 94 (1975); V.N. Gribov, L.N. Lipatov, Sov. J. Nucl. Phys.**15**, 438 (1972); G. Altarelli, G. Parisi, Nucl. Phys. **B126**, 298 (1977); Yu.L. Dokshitzer, Sov. Phys. JETP **46**, 641




FIG. 3: Our results for the linear gluon distribution, using Eq.(10), as a function of x in a wide range of Q^2 values, $1 \text{ GeV}^2 < Q^2 \leq 1000 \text{ GeV}^2$.




FIG. 4: Comparison of the $G(x, Q^2)$ obtained from Eq.(11), with the parametrization methods in Refs. [11,12] at $Q^2 = 10$ and 100 GeV^2 .

(1977).

2. H.Khanpour, A.Mirjalili and S.Atashbar Tehrani, Phys.Rev.C**95**, 035201 (2017); H.Khanpour, M.Goharipour and V.Guzey, Eur.Phys.J.C**78**, 7(2018); S. Mohammad Moosavi Nejad, H.Khanpour, S.Atashbar Tehrani and M.Mahdavi, Phys.Rev.C**94**, 045201 (2016).
3. G.R.Boroun, S.Zarrin and F.Teimoury, Eur.Phys.J.Plus **130**, 214(2015); F.Teimoury Azad-




FIG. 5: Linear and non-linear behavior of the gluon distributions for several fixed values of x in a wide range of Q^2 shows that the effect of the nonlinear terms vanishes as Q^2 increases.




FIG. 6: Non-linear gluon distribution at low Q values due to the positive and negative roots in Eqs.(18) and (19) compared with the linear gluon distribution functions (dashed lines) obtained in Ref.[11] at each active flavor number.

- bakht and G.R.Boroun, *Int.J.Theor.Phys.***57**, 495 (2018); S.Zarrin and G.R.Boroun, *Nucl.Phys.B***922**, 126(2017); F.Teimoury Azadbakht, G.R.Boroun and B.Rezaei, *Int.J.Mod.Phys.E***27**, 1850071 (2018).
4. M.Mottaghizadeh, F.Taghavi Shahri and P.Eslami, *Phys.Lett.B***773**, 375(2017); M.Mottaghizadeh, P.Eslami and F.Taghavi-Shahri, *Int.J.Mod.Phys.A***32**, 1750065(2017); S.Dadfar and S.Zarrin, *Eur.Phys.J.C***80**, 319(2020); H.Hosseinkhani, M.Modarres and N.Olanj, *Int.J.Mod.Phys.A***32**, 1750121 (2017).
 5. M. M. Block, L. Durand, and D. W. McKay, *Phys. Rev. D* **79**, 014031 (2009).
 6. LHeC Collaboration and FCC-he Study Group, P.Agostini et al., CERN-ACC-Note-2020-0002, arXiv:2007.14491 [hep-ex] (2020).
 7. L.V.Gribov, E.M.Levin and M.G.Ryskin, *Phys.Rept.***100**, 1 (1983); A.H.Mueller and J.w.Qiu, *Nucl.Phys.B* **268**, 427 (1986).
 8. A. V. Kotikov and V. N. Velizhanin, arXiv: 0501274 [hep-ph] (2005).
 9. F. D. Aaron et al. (H1 and ZEUS Collaborations), *JHEP* **1001**, 109 (2010).
 10. M. M. Block, L. Durand and P. Ha, *Phys. Rev. D* **89**, 094027 (2014).
 11. Martin M.Block and L.Durand, arXiv:0902.0372 [hep-ph] 2009.
 12. M. M. Block, L. Durand, and D. W. McKay, *Phys. Rev. D* **77**, 094003 (2008).
 13. J. Breitweg et al. (ZEUS), *Phys. Lett. B* **487**, 53 (2000); S. Chekanov et al. (ZEUS), *Eur. Phys. J. C* **21**, 443 (2001).
 14. K. J. Eskola, H. Honkanen, V. J. Kolhinen, J.-W. Qiu and C. A. Salgado, *Nucl. Phys. B* **660**, 211 (2003).
 15. A. Dainese et al., HERA-LHC Workshop, DESY, (2005).
 16. J. Pumplin et al., *JHEP* **07**, 012 (2002).