

# The $\infty$ -category of stabilized Liouville sectors

Oleg Lazarev, Zachary Sylvan, and Hiro Lee Tanaka

October 31, 2022

## Abstract

We prove the surprising fact that the infinity-category of stabilized Liouville sectors is a localization of an ordinary category of stabilized Liouville sectors and strict sectorial embeddings.

From the perspective of homotopy theory, this result continues a trend of realizing geometrically meaningful mapping spaces through the categorically formal process of localizing. From the symplectic viewpoint, these results allow us to reduce highly non-trivial coherence results to much simpler verifications. For example, we prove that the wrapped Fukaya category is coherently functorial on stabilized Liouville sectors: Not only does a wrapped category receive a coherent action from stabilized automorphism spaces of a Liouville sector, spaces of sectorial embeddings map to spaces of functors between wrapped categories in a way respecting composition actions. As a consequence, we observe that wrapped Floer theory for sectors works in families. As we will explain, our methods immediately establish such coherence results for most known sectorial invariants, including Lagrangian cobordisms.

As another application, we show that this infinity-category admits a symmetric monoidal structure, given by direct product of underlying sectors. The existence of this structure relies on a computation—familiar from the foundations of factorization homology—that localizations detect certain isotopies of smooth manifolds. Moreover, we characterize the symmetric monoidal structure using a universal property, again producing a simple-as-possible criterion for verifying whether invariants are both continuously and multiplicatively coherent in a compatible way.

Much of this paper is devoted to the foundational work of rigorously constructing the infinity-category of stabilized Liouville sectors, where any sector  $M$  is identified with its stabilization  $M \times T^*[0, 1]$ . We begin with a verification that our construction is indeed an infinity-category (which relies on new constructions in Liouville geometry), then show that our infinity-category computes the “correct” mapping spaces (employing new simplicial techniques). A significant ingredient is a demonstration that two notions of sectorial maps give rise to homotopy equivalent spaces of maps.

## Contents

<b>1</b>	<b>Introduction</b>	<b>4</b>
1.1	A coherence result . . . . .	5
1.2	The infinity-categories of Liouville sectors . . . . .	7
1.3	Stabilized Liouville sectors via localization . . . . .	9
1.4	Localization exhibits coherent functoriality . . . . .	12
1.5	Multiplicative coherence . . . . .	13
1.6	Results for Liouville sectors with tangential structures . . . . .	16
1.7	The geometric insights . . . . .	16
1.8	Later works and other future directions . . . . .	17
1.9	This paper is $\infty$ -categorical in most of its tools . . . . .	18
1.10	Acknowledgments . . . . .	18

<b>2</b>	<b>Some Liouville geometry</b>	<b>20</b>
2.1	Basic definitions . . . . .	20
2.2	The movie construction . . . . .	26
2.3	The movie construction for maps . . . . .	29
2.4	Shrinking near the boundary, rounding corners . . . . .	30
2.5	Isotopy extension . . . . .	31
2.6	Spaces of sectorial embeddings . . . . .	34
2.7	Spaces of (bordered) deformation embeddings . . . . .	35
<b>3</b>	<b><math>\infty</math>-categorical tools</b>	<b>38</b>
3.1	Simplicial and semisimplicial sets . . . . .	38
3.2	Subdivisions and $\text{Ex}$ . . . . .	39
3.3	$\text{Ex}_{\simeq}$ of $\infty$ -categories . . . . .	40
3.4	Making a homotopy . . . . .	43
3.5	Understanding subdivisions of subdivisions . . . . .	46
<b>4</b>	<b>Simplices and collarings</b>	<b>48</b>
4.1	An unconventional convention . . . . .	48
4.2	Collaring conventions . . . . .	48
4.3	Linearity, locally . . . . .	51
4.4	From continuous to smooth . . . . .	52
4.5	The homotopy equivalence of different deformation embedding spaces . . . . .	55
<b>5</b>	<b><math>\text{Liou}_{\Delta}</math> and <math>\text{Liou}_{\Delta}^{\text{Def}}</math></b>	<b>58</b>
5.1	Collaring movies of deformation embeddings . . . . .	58
5.2	$\text{Liou}_{\Delta}^{\text{Def}}$ (as a semisimplicial set) . . . . .	59
5.3	A 1-simplex . . . . .	60
5.4	A 2-simplex . . . . .	61
5.5	$\text{Liou}_{\Delta}$ (as a semisimplicial set) . . . . .	64
5.6	Change of bases to max-constant simplices . . . . .	66
<b>6</b>	<b>The weak Kan condition</b>	<b>67</b>
6.1	For $n = 2$ . . . . .	68
6.2	For $n \geq 3$ , filling most of the simplex . . . . .	70
6.3	$n \geq 3$ and $j \neq n - 1$ . . . . .	74
6.4	$n \geq 3$ and $j = n - 1$ . . . . .	75
6.5	Proof of Theorem 6.1 and 6.2 . . . . .	76
6.6	$(\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$ is weak Kan . . . . .	76
<b>7</b>	<b><math>\text{Liou}_{\Delta}</math> as an <math>\infty</math>-category</b>	<b>77</b>
7.1	Idempotent equivalences and constant simplices . . . . .	77
7.2	Proof of Theorem A . . . . .	78
7.3	Some of the geometry of $\text{Liou}_{\Delta}$ . . . . .	79
<b>8</b>	<b>Mapping spaces of <math>\text{Liou}_{\Delta}</math></b>	<b>83</b>
8.1	The right mapping spaces . . . . .	83
8.2	$\text{hom}'$ is equivalent to $\text{hom}^R$ . . . . .	85
8.3	Proof of injection on homotopy groups . . . . .	86
8.4	Proof of surjection on homotopy groups . . . . .	89

8.5	Proof of Theorems 8.1 and B. . . . .	93
8.6	An informal argument . . . . .	93
<b>9</b>	<b><math>\mathrm{Liou}_\Delta</math> and <math>\mathrm{Liou}_\Delta^{\mathrm{Def}}</math> are equivalent</b>	<b>95</b>
<b>10</b>	<b>Localization (the proof of Theorem C)</b>	<b>96</b>
10.1	Defining $\beta$ . . . . .	96
10.2	Defining $\alpha$ . . . . .	97
10.3	$\alpha$ as a left inverse . . . . .	98
10.4	$\alpha$ as a right inverse . . . . .	102
10.5	Proof of Lemma 10.22 . . . . .	104
10.6	Completing the proof of Theorem C . . . . .	109
<b>11</b>	<b>The symmetric monoidal structure</b>	<b>110</b>
11.1	Stabilizing commutes with localization . . . . .	110
11.2	Symmetric monoidal localization . . . . .	111
11.3	Symmetric monoidal structures on stabilizations . . . . .	111
11.4	The symmetric monoidal structure . . . . .	112
11.5	Proof of Theorem D . . . . .	113
11.6	Proof of Lemma 11.7 . . . . .	114
<b>12</b>	<b>Equivalent localizations</b>	<b>121</b>
12.1	The wrapped Fukaya category is unchanged by equivalences . . . . .	125
<b>13</b>	<b>Incorporating tangential structures</b>	<b>127</b>
13.1	Preliminaries: The $\infty$ -categories of (stable) vector bundles . . . . .	127
13.2	Tangential structures respect direct products . . . . .	130
13.3	Stabilized Liouville sectors with tangential structures . . . . .	131
13.4	Localization with tangential structures (Proof of Theorem E) . . . . .	132

# 1 Introduction

The theory of Liouville manifolds, and Liouville sectors more generally, has lacked a convenient framework for proving fundamental categorical questions regarding symplectic invariants. For example, is the wrapped Fukaya category functor sensitive to the topology and continuous actions of Liouville embedding spaces?

Prior to the present work, the “obvious” way to try and answer the above Floer-theoretic question would have been to write down (or prove the existence of) a coherent system of auxiliary Floer data, and prove that the associated moduli spaces of pseudoholomorphic curves compactify in such a way as to output the desired  $A_\infty$ -categorical formulas. Such an approach is, while well-established, time-consuming. Not only that, different questions of coherence raise different flavors of difficulties when constructing relevant moduli spaces, requiring ad hoc techniques in each setting—as an example, we invite the reader to ponder how to prove continuous coherences are compatible with multiplicative/Kunneth-type formulas in wrapped Floer theory. A more fundamental scientific issue is the difficulty for an independent user or referee to verify such methods efficiently. Even if most agree “it can be done,” the classical techniques also result in a troubling number who agree “I don’t want to write or read the details.” The works [26, 27] exhibited one method of proving coherence results while avoiding the gnarliest of these details. See also Remarks 1.12 and 1.26.

As it turns out, and as we show in this work and its follower(s), one can avoid such details and prove certain coherence results with minimal Floer theory—i.e., while constructing only minimally-intricate moduli spaces of holomorphic disks.

More fundamentally, we construct here an  $\infty$ -category  $\mathcal{L}iou^\diamond$  of stabilized Liouville sectors (Notation 1.22). This  $\infty$ -category has excellent formal properties, and the results here open the door for applying powerful  $\infty$ -categorical machinery to Liouville symplectic geometry.

**Remark 1.1.** To our readers unfamiliar with  $\infty$ -categories: One would not lose much intuition by pretending that an  $\infty$ -category is simply a category wherein the collection of morphisms may form a space (and not just a set). For example, the collection of spaces can form an  $\infty$ -category in two distinct ways. First, we may think of the collection of continuous functions as simply a set. Second, one may endow the collection of continuous functions with the compact-open topology and enjoy the fact that composition of continuous functions is continuous.

**Warning 1.2.** Throughout this work, the word “stable” refers to the geometric process of increasing dimension. It does not refer to stability in the sense of higher algebra or pretriangulated categories. For example,  $\mathcal{L}iou^\diamond$  is not a stable  $\infty$ -category in the sense of Lurie [18].

First, some preliminaries. Let  $M$  and  $N$  be Liouville sectors (Definition 2.1).

**Definition 1.3.** A smooth, codimension zero, proper embedding  $f : M \rightarrow N$  is a *sectorial embedding* if  $f^*\lambda^N = \lambda^M + dh$  for some compactly supported smooth function  $h$  on  $M$ . A sectorial embedding is *strict* if  $f^*\lambda^N = \lambda^M$ .<sup>1</sup>

We say (a not necessarily strict)  $f$  is a *sectorial equivalence*, or an equivalence of Liouville sectors, if there exist a (not necessarily strict) sectorial embedding  $g : N \rightarrow M$  and isotopies  $g \circ f \sim \text{id}_N$ ,  $f \circ g \sim \text{id}_M$ , through (not necessarily strict) sectorial embeddings.

Sectorial equivalences are a natural class of maps known to leave wrapped Fukaya categories unchanged (see Remark 2.45 of [16], Lemma 3.33 of [9], and Proposition 12.9 below).

---

<sup>1</sup>This is not the notion of strictness used in [2].

Finally, for the sake of exposition, we present an undefined idea: If  $M$  and  $N$  are Liouville sectors, let

$$\mathrm{Emb}^\diamond(M, N) \approx \{f : M \times T^*[0, 1]^k \rightarrow N \times T^*[0, 1]^{(\dim M + 2k - \dim N)/2}, \quad k \gg 0\} \quad (1.1)$$

denote the space of stable (not necessarily strict) sectorial embeddings.

**Warning 1.4.** This space is not well-defined as written above—hence the approximation symbol—because even if  $f$  is a sectorial embedding, its stabilization  $f \times \mathrm{id}_{T^*[0, 1]}$  is typically not a sectorial embedding, as the compact-support condition on  $h$  is not preserved. At the same time, it is natural to contemplate a “stabilized” mapping space: For Weinstein sectors, the wrapped category is known to be unchanged under stabilization [10]. We give a precise formulation of this mapping space in Definition 1.17.

## 1.1 A coherence result

In the introduction, we will denote by capital letters (A, B, C, et cetera) what we view to be the central results .

To begin, we informally state a result that we make formal in Theorem C. It guarantees that certain invariants known only to be sensitive to the *set* of strict embeddings can in fact be made sensitive to the *space* of stable, not-necessarily-strict embeddings.

**Theorem 1.5.** Let  $\mathbb{W}$  be an invariant of Liouville sectors taking values in an  $\infty$ -category admitting filtered colimits and assume the following:

- (a) ( $\mathbb{W}$  is functorial with respect to strict inclusions.) To any *strict* sectorial embedding  $f : M \rightarrow N$ , one can functorially assign a pushforward map  $f_* : \mathbb{W}(M) \rightarrow \mathbb{W}(N)$ .
- (b) ( $\mathbb{W}$  stabilizes.)  $\mathbb{W}$  is equipped with natural maps  $\mathbb{W}(M) \rightarrow \mathbb{W}(M \times T^*[0, 1])$ . This allows us to define the “stabilized” invariant associated to  $M$ , which we call  $\mathbb{W}^\diamond(M)$ .<sup>2</sup>
- (c) ( $\mathbb{W}$  stably inverts equivalences.) If  $f : M \rightarrow N$  is a strict sectorial embedding that happens to be an equivalence of Liouville sectors, then the induced map  $f_* : \mathbb{W}^\diamond(M) \rightarrow \mathbb{W}^\diamond(N)$  is an equivalence.

Then, for any pair of Liouville sectors  $M$  and  $N$ ,  $\mathbb{W}$  induces a continuous map from the space  $\mathrm{Emb}^\diamond(M, N)$  to the space of morphisms from  $\mathbb{W}^\diamond(M)$  to  $\mathbb{W}^\diamond(N)$  and this map extends the assignments (a). Finally, upon varying  $M$  and  $N$ , these maps coherently respect composition.

**Remark 1.6.** In fact, one may replace (c) with the assumption that  $\mathbb{W}$  stably inverts any of three other natural notions of equivalences—trivial inclusions, bordered deformation equivalences, or movie inclusions—and the conclusion of the theorem still holds (see Section 12). We note that, in practice, some invariants may satisfy a stronger assumption than (c): If  $f$  is a strict equivalence, the map  $f_* : \mathbb{W}(M) \rightarrow \mathbb{W}(N)$  may be an equivalence even before passing to the stabilized invariants. This is the case when  $\mathbb{W}$  is the wrapped Fukaya category.

**Remark 1.7.** In many examples of  $\infty$ -categories, a filtered colimit may be modeled by an increasing union along well-behaved inclusions (of possibly infinitely many objects). So one would not lose much intuition by imagining that  $\mathbb{W}^\diamond(M)$  is a union  $\bigcup_{n \geq 0} \mathbb{W}(M \times T^*[0, 1]^n)$  along the

---

<sup>2</sup>Concretely,  $\mathbb{W}^\diamond(M)$  is the colimit of the diagram  $\mathbb{W}(M) \rightarrow \mathbb{W}(M \times T^*[0, 1]) \rightarrow \mathbb{W}(M \times T^*[0, 1]^2) \rightarrow \dots$ . This filtered colimit exists by assumption; see also Remark 1.7

stabilization maps (b). Finally, most invariants take values in familiar  $\infty$ -categories—of groups, modules, categories, sets, spaces, chain complexes et cetera—where such filtered colimits exist, so the assumption that  $\mathbb{W}$  takes values in a setting with filtered colimits is not onerous.

The wrapped Fukaya category<sup>3</sup>  $\mathcal{W}(M)$  and the Lagrangian cobordism  $\infty$ -category<sup>4</sup>  $\mathbf{Lag}(M)$  are both known to satisfy the hypotheses of Theorem 1.5. In fact, for all Liouville sectors,  $\mathbf{Lag}(M) \simeq \mathbf{Lag}^\diamond(M)$ . So we have the following as an example application:

**Theorem 1.8.** Let  $M$  and  $N$  be Liouville sectors. Then

1. There are continuous maps

$$\mathrm{Emb}^\diamond(M, N) \rightarrow \mathrm{Fun}(\mathbf{Lag}(M), \mathbf{Lag}(N)), \quad \mathrm{Emb}^\diamond(M, N) \rightarrow \mathrm{Fun}(\mathcal{W}^\diamond(M), \mathcal{W}^\diamond(N))$$

to the space of functors.

2. Further, these maps can be made to respect composition operations homotopy-coherently.

We emphasize the uniformity of the result. Before this work, one would have had to use Floer-theoretic techniques to prove the wrapped Fukaya category result and Lagrangian cobordism techniques to prove—you get the idea. At the expense of passing to *stabilized* invariants, our results allow one to cordon off all non-uniform treatments to the verification of the hypotheses in Theorem 1.5. Note also that one could consider a chain-complex valued, or space-valued, invariant of sectors (rather than a categorical invariant); Theorem 1.5 still guarantees continuous maps to the relevant mapping spaces. This uniformity is useful for comparing invariants as well—see Corollary 1.25.

The theorem induces a map of  $A_\infty$ -spaces from  $\mathrm{Emb}^\diamond(M, M)$  to  $\mathrm{Fun}(\mathcal{W}^\diamond(M), \mathcal{W}^\diamond(M))$ . (The “homotopy coherence” of the theorem guarantees that these are indeed maps respecting the  $A_\infty$  structures.) It is then automatic that the higher homotopy groups of the former map to the negative Hochschild cohomology groups of  $\mathcal{W}(M)$ . Indeed, observe the isomorphisms

$$\begin{aligned} \pi_i \mathrm{Fun}(\mathcal{W}(M), \mathcal{W}(M)) &\cong HH^{1-i}(\mathcal{W}(M), \mathcal{W}(M)) & (i > 1), \\ \pi_1 \mathrm{Fun}(\mathcal{W}(M), \mathcal{W}(M)) &\cong (HH^0(\mathcal{W}(M), \mathcal{W}(M)))^\times \end{aligned}$$

where the  $\times$  superscript indicates the units of the ring. Then we have:

**Corollary 1.9.** There exists a natural map

$$\pi_i \mathrm{Emb}^\diamond(M, M) \rightarrow HH^{1-i}(\mathcal{W}(M), \mathcal{W}(M)) \quad (i > 0).$$

And for any other sector  $N$ , the following diagram commutes:

$$\begin{array}{ccc} \pi_i \mathrm{Emb}^\diamond(M, N) \times \pi_i \mathrm{Emb}^\diamond(M, M) & \longrightarrow & \pi_i \mathrm{Emb}^\diamond(M, N) \\ \downarrow & & \downarrow \\ \pi_i \mathrm{Fun}(\mathcal{W}^\diamond(M), \mathcal{W}^\diamond(N)) \times \pi_i \mathrm{Fun}(\mathcal{W}^\diamond(M), \mathcal{W}^\diamond(M)) & \longrightarrow & \pi_i \mathrm{Fun}(\mathcal{W}^\diamond(M), \mathcal{W}^\diamond(N)) \end{array}$$

<sup>3</sup>This functoriality is proven in [9], and other foundational works include [1, 3, 31]. We also point out the font difference between  $\mathbb{W}$  (which refers to a general invariant) and  $\mathcal{W}$  (which refers to the wrapped Fukaya category).

<sup>4</sup>In the notation of [25], we set  $\mathbf{Lag}(M) = \mathbf{Lag}_\Lambda(M)$  where  $\Lambda$  is the skeleton of  $M$ .

**Remark 1.10.** This is a stable analogue of results from [26]—indeed, to our knowledge, *ibid.* was the first work which demonstrated that localization techniques in  $\infty$ -category theory (and not just localization along positive wrappings as in [9]) yield geometrically meaningful results in wrapped Floer theory. One major departure from [26] is the absence of analytical arguments regarding  $J$ -holomorphic disks in the present work – indeed, *ibid.* requires a set-up for Floer theory in families, while here the  $\infty$ -categorical techniques allow us to avoid them completely, and simply piggy-back on the results of [9].

**Example 1.11** (Floer theory works in families). In particular, suppose that  $E \rightarrow X$  is a bundle of Liouville sectors over some base space  $X$  with fiber  $F$  and let  $\text{Aut}(F)$  denote the space of (not necessarily strict) Liouville automorphisms. Then the composition

$$X \rightarrow B \text{Aut}(F) \rightarrow B \text{Emb}^\diamond(F, F) \rightarrow B \text{Aut}(\mathbb{W}(F))$$

where the last map is guaranteed to exist by Theorem 1.5, is an invariant of the bundle. In fact, the last part of the composition gives a way to study the topology of Liouville automorphism spaces. For example, when  $\mathbb{W}$  is the wrapped Fukaya category, and noting the natural map from  $B \text{Aut}(\mathbb{W}(F))$  to the shift of Hochschild cochains of  $\mathbb{W}(F)$ , we obtain a Liouville version of the Seidel homomorphism [29] (for all higher homotopy groups, not just  $\pi_1$ , of  $\text{Aut}(F)$ ). Put another way, wrapped Floer theory behaves well in families. As in Remark 1.10, we obtain these results with no need to address compactness of moduli of disks in families, nor to carefully study degenerations of disks in families.

**Remark 1.12** (Another advantage). Because current definitions of ( $\mathbb{Z}$ -linear) Fukaya categories only depend on counting 0-dimensional components of moduli spaces, as do the definitions of most  $A_\infty$  functors, the combinatorics involved in proving coherence results are largely manageable even if tedious. However, for spectrally enriched Floer invariants, which necessarily extract data from all higher-dimensional moduli spaces, far more delicate book-keeping would be necessary to construct families of functors out of families of embeddings. Theorem 1.5 allows us to deduce the coherent functoriality of such spectral invariants simply by verifying “discrete” or “embedding-by-embedding” (as opposed to families of embeddings) properties, which are far easier to establish.

**Remark 1.13.** Finally, many known methods for constructing actions of *unstabilized* sectorial embedding spaces on the above invariants actually factor through the stabilized embedding spaces anyway.<sup>5</sup> So for invariants satisfying the hypotheses of Theorem 1.5, no information is lost upon passage to stabilization.

## 1.2 The infinity-categories of Liouville sectors

Let us now dip our toes into the details—by highlighting our main constructions, and one fundamental property. These details serve the most important philosophical take-away from our work:  $\infty$ -categorical localizations give rise to meaningful objects in symplectic geometry. We refer the reader to Table 1 for a summary of the various  $\infty$ -categories we construct.

The first construction of our work (Definition 5.12) is an  $\infty$ -category

$$\text{Liou}_\Delta$$

whose objects are Liouville sectors, and whose commutative diagrams consist of certain data organized by the combinatorics of barycentric subdivisions.

---

<sup>5</sup>While this expectation is intuitive, the hard work is in translating known constructions—see [26] for example—to the methods used in this paper.

**Example 1.14.** A morphism from  $M_0$  to  $M_1$  is not the data of a sectorial embedding, but equivalent to the data of two *strict* embeddings

$$M_0 \times T^*[0, \epsilon] \rightarrow M_{01} \times T^*[0, 1] \leftarrow M_1 \times T^*[1 - \epsilon, 1]$$

where the left-pointing arrow is demanded to be an isomorphism to its image, where both embeddings are collared in an appropriate way, and where we specify a Liouville structure on  $M_{01} \times T^*[0, 1]$  which may not be a direct product of the Liouville structures of the two factors. The Liouville structure on  $M_{01} \times T^*[0, 1]$  will be such that it determines a 1-parameter family of Liouville structures on  $M_1$  itself, through deformations by derivatives of compactly supported functions.

Note that the above diagram is in the “shape” of the barycentric subdivision of a 1-simplex. Likewise, a  $k$ -simplex in  $\text{Liou}_\Delta$  will consist of a diagram of sectors in the shape of a barycentric subdivision of a  $k$ -simplex. See (7.3) for the case  $k = 2$ .

**Theorem A** (Theorem 7.8).  $\text{Liou}_\Delta$  is an  $\infty$ -category.

The construction of  $\text{Liou}_\Delta$  is the technical heart of our paper. Indeed, even the verification that  $\text{Liou}_\Delta$  is an  $\infty$ -category is not formal, as one requires results from Liouville geometry (the existence of isotopy extensions and the ability to “squeeze” a sector away from its boundary and corner strata). Moreover, we first define  $\text{Liou}_\Delta$  as a semisimplicial set; the geometry just mentioned allows us to use a result of Steimle [30] to conclude  $\text{Liou}_\Delta$  is an  $\infty$ -category, and a result of the third author [33] to deduce the existence of certain functors out of  $\text{Liou}_\Delta$ .

We then prove:

**Theorem B** (Theorem 8.1). Fix two Liouville sectors  $M$  and  $N$ , and let  $\text{Emb}_{\text{Liou}}(M, N)$  denote the space of sectorial embeddings from  $M$  to  $N$ , topologized to permit control near infinity over compact families. (See Definition 2.57.) Then there exists a homotopy equivalence between the (unstabilized) mapping spaces

$$\text{Emb}_{\text{Liou}}(M, N) \simeq \text{hom}_{\text{Liou}_\Delta}(M, N).$$

The underlying philosophy of the theorem is that families of (not necessarily strict) embeddings may be modeled by certain diagrams in the shapes of barycentric subdivisions (consisting only of strict embeddings). The take-away is that one should think of  $\text{Liou}_\Delta$  as a different  $\infty$ -categorical model of the usual space-enriched category of sectors, just as some chain complexes admit equivalent but differently-behaved models.

**Remark 1.15.** It is possible to construct an equivalence of  $\infty$ -categories between  $\text{Liou}_\Delta$  and the space-enriched category of Liouville sectors, but the combinatorial details are quite involved so we leave this for a separate work. Regardless, we give a “moral” argument in Section 8.6 as to why one should expect such an equivalence.

To obtain a stabilization of  $\text{Liou}_\Delta$ , we construct in Definition 5.5 an auxiliary  $\infty$ -category

$$\text{Liou}_\Delta^{\text{Def}}$$

equivalent to  $\text{Liou}_\Delta$  (Theorem 9.3), but whose diagrams allow for families of Liouville structures that are not necessarily related by compactly supported deformations. The constraints imposed by strictness in the barycentric subdivision diagrams, and the freedoms granted by this larger class of deformations, pay off in the following observation: The stabilization assignment  $M \mapsto T^*[0, 1] \times M$



is a functor from  $\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}}$  to itself.<sup>6</sup> So we can now take the colimit (in our setting, an increasing union)

$$(\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}})^\diamond := \mathrm{colim} \left( \mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}} \xrightarrow{T^*[0,1] \times -} \mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}} \xrightarrow{T^*[0,1] \times -} \dots \right). \quad (1.2)$$

Of course, because  $\mathrm{Liou}_\Delta \simeq \mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}}$ , we may also construct an  $\infty$ -category

$$\mathrm{Liou}_\Delta^\diamond := \mathrm{colim} \left( \mathrm{Liou}_\Delta \xrightarrow{T^*[0,1] \times -} \mathrm{Liou}_\Delta \xrightarrow{T^*[0,1] \times -} \dots \right) \quad (1.3)$$

where we have abused notation to let  $T^*[0,1] \times -$  denote the induced endofunctor of  $\mathrm{Liou}_\Delta$  as well.  $\mathrm{Liou}_\Delta^\diamond$  and  $(\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}})^\diamond$  are equivalent as  $\infty$ -categories; we call them both the  $\infty$ -category of stabilized Liouville sectors. (See also Notation 1.22.)

**Remark 1.16.** Concretely, an object of  $\mathrm{Liou}_\Delta^\diamond$  is a pair  $(M, k)$  where  $M$  is a Liouville sector and  $k$  is a non-negative integer, up to the relation  $(M, k) \sim (T^*[0,1]^N \times M, k + N)$ . In particular, there are some objects such as  $(T^*\mathbb{R}^0, k)$  which one might think of as the  $k$ th “destabilization” of  $T^*\mathbb{R}^0$ .

Now let us discuss the morphisms. If there is a morphism from  $(M, k)$  to  $(M', k')$ , it must be that  $\dim(M) + 2k' = \dim(M') + 2k$ . Thus,  $\mathrm{Liou}_\Delta^\diamond$  is a disjoint union of many subcategories, indexed by the difference between  $2k$  and the underlying dimension of  $M$ . By Theorem B, we see that a morphism from  $M$  to  $N$  is, up to contractible choice, the data of a (not necessarily strict) sectorial embedding  $T^*[0,1]^a \times M \rightarrow T^*[0,1]^b \times N$  for  $\dim M + a = \dim N + b$ .

In light of Theorem 1.8 and our ability to stabilize (1.3), the mapping spaces  $\mathrm{hom}_{\mathrm{Liou}_\Delta^\diamond}(M, N)$  are *the* model for  $\mathcal{E}\mathrm{mb}^\diamond(M, N)$  we use in this work. This makes precise the informal description in (1.1):

**Definition 1.17** ( $\mathcal{E}\mathrm{mb}^\diamond$ ). For two Liouville sectors  $M$  and  $N$ , we define

$$\mathcal{E}\mathrm{mb}^\diamond(M, N) := \mathrm{hom}_{\mathrm{Liou}_\Delta^\diamond}(M, N).$$

**Remark 1.18.** Moreover, for any two choices of even integers  $a$  and  $a'$ , the full subcategory of those  $(M, k)$  for which  $\dim M - 2k = a$  is equivalent to the full subcategory of those  $(M', k')$  for which  $\dim M' - 2k' = a'$ . And, by definition of  $\mathbb{W}^\diamond$  (in Theorem 1.5), we see that  $\mathbb{W}^\diamond(M, k) \simeq \mathbb{W}^\diamond(M, 0)$  regardless of  $k$ . Thus, in studying the invariant  $\mathbb{W}^\diamond$ , one may safely consider only the subcategory of  $\mathrm{Liou}_\Delta^\diamond$  for which  $\dim M = 2k$ .

However, the subcategories with  $\dim M \neq 2k$  will become relevant when we consider symmetric monoidal functors out of  $\mathrm{Liou}_\Delta^\diamond$ . See Remark 1.31

### 1.3 Stabilized Liouville sectors via localization

We now come to the one main property:  $(\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}})^\diamond$  is a localization.

**Remark 1.19.** To situate the reader, recall that a functor  $\mathcal{A} \rightarrow \mathcal{B}$  is called a localization if it exhibits  $\mathcal{B}$  as a universal (in this case, initial)  $\infty$ -category that inverts a particular collection of morphisms in  $\mathcal{A}$ . In particular, if  $F : \mathcal{A} \rightarrow \mathcal{C}$  is a functor sending all morphisms in this collection to equivalences in  $\mathcal{C}$ , then  $F$  canonically factors through  $\mathcal{B}$  up to homotopy. A central tool in

<sup>6</sup> In contrast, this assignment does not know what to do for non-strict diagrams of Liouville sectors for the reasons raised in Warning 1.4. Indeed, much of our simplicial finagling began so we could avoid dealing with this issue. We also note the need to pass to a larger class of deformations (not just compactly supported deformations) for the same reason.

$\mathrm{Liou}_{\mathrm{str}}$	A category of Liouville sectors and only strict embeddings. Notation 1.20.	Sees no homotopical information of embedding spaces, but easy to construct functors out of.
$\mathrm{Liou}_{\mathrm{str}}^{\diamond}$	A category of stabilized Liouville sectors and strict embeddings. (1.4)	Easy to define as a colimit because $\mathrm{Liou}_{\mathrm{str}}$ admits an endofunctor called stabilization.
$\mathrm{Liou}_{\Delta}$	An $\infty$ -category of Liouville sectors. Morphism spaces are equivalent to spaces of sectorial embeddings. Definition 5.12.	Diagrams involve only strict embeddings, but can encode deformations of Liouville structures by compactly supported exact deformations.
$\mathrm{Liou}_{\Delta}^{\mathcal{D}\mathrm{ef}}$	An $\infty$ -category of Liouville sectors. Morphism spaces are equivalent to spaces of sectorial embeddings. Definition 5.5.	Diagrams involve only strict embeddings, but can encode deformations of Liouville structures by arbitrary exact deformations. Morphism spaces do not admit obvious composition law.
$(\mathrm{Liou}_{\Delta}^{\mathcal{D}\mathrm{ef}})^{\diamond}$	An $\infty$ -category of stabilized Liouville sectors. (1.2)	Easy to define as a colimit because $\mathrm{Liou}_{\Delta}^{\mathcal{D}\mathrm{ef}}$ admits an endofunctor called stabilization.
$\mathrm{Liou}_{\Delta}^{\diamond}$	An $\infty$ -category of stabilized Liouville sectors. (1.3)	Defined indirectly, as $\mathrm{Liou}_{\Delta}$ does not admit obvious stabilization endofunctor.
$\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\epsilon q^{\diamond})^{-1}]$	Localization of $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$ along (strict) sectorial equivalences	Useful universal property.
$\mathcal{L}\mathrm{iou}^{\diamond}$	A convenient notation denoting any of the three equivalent $\infty$ -categories of stabilized sectors. Notation 1.22.	Notation is clean.
$\mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^{\diamond}$	A version of $\mathcal{L}\mathrm{iou}^{\diamond}$ where stabilized sectors are equipped with tangential data, and morphisms are maps equipped with compatibilities of tangential data. Construction 13.11.	Necessary to articulate functoriality of various versions of Fukaya and microlocal categories.

Table 1: Summary of the various categories and  $\infty$ -categories of Liouville sectors. The horizontal double-lines delineate equivalences classes of  $\infty$ -categories. For example,  $\mathrm{Liou}_{\Delta}$  and  $\mathrm{Liou}_{\Delta}^{\mathcal{D}\mathrm{ef}}$  are equivalent  $\infty$ -categories.

modern homotopy theory is to begin with  $\mathcal{A}$  a discrete category (i.e., a category in the classical sense [20]), and to localize  $\mathcal{A}$  to an  $\infty$ -category with rich topological information. For example, if one localizes the ordinary category of CW complexes along homotopy equivalences, one recovers the  $\infty$ -category of CW complexes (and in particular, the homotopy type of all mapping *spaces* between CW complexes).<sup>7</sup> In the present work, we recover the stabilized sectorial embedding spaces using this process (Theorem C below).

So, what is  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$  a localization of?

**Notation 1.20.** Let  $\mathrm{Liou}_{\mathrm{str}}$  be the category (in the classical sense) whose objects are Liouville sectors, and whose morphisms are strict sectorial embeddings. We stabilize to define a category (in the classical sense)

$$\mathrm{Liou}_{\mathrm{str}}^{\diamond} := \mathrm{colim} \left( \mathrm{Liou}_{\mathrm{str}} \xrightarrow{T^*[0,1] \times -} \mathrm{Liou}_{\mathrm{str}} \xrightarrow{T^*[0,1] \times -} \dots \right) \quad (1.4)$$

of stabilized Liouville sectors, where morphism sets are sets of strict stable sectorial embeddings. (The  $T^*[0,1] \times -$  functor is well-defined on the nose because  $\mathrm{Liou}_{\mathrm{str}}$  only contains strict sectorial embeddings.)

Based on our models, one can straightforwardly construct a functor (Construction 10.3)

$$\mathrm{Liou}_{\mathrm{str}}^{\diamond} \rightarrow (\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}. \quad (1.5)$$

And though sectorial equivalences are of course not invertible in  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$ , their images become invertible up to homotopy in  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$  (Proposition 7.14).

Let  $\mathfrak{eq}^{\diamond} \subset \mathrm{Liou}_{\mathrm{str}}^{\diamond}$  denote the collection of (strict) sectorial equivalences.

**Theorem C.** The functor (1.5) is a localization of  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$  along (strict) sectorial equivalences. More precisely, the induced map

$$\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}] \rightarrow (\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$$

is an equivalence of  $\infty$ -categories.

**Remark 1.21.** Let us explain where stabilization (the  $\diamond$  superscript) enters the picture. The point is that  $\mathrm{Liou}_{\Delta}$  already incorporates movies (Definition 2.43) in defining its morphisms, and receives a natural map from the localization of  $\mathrm{Liou}_{\mathrm{str}}$  – however, to exhibit an inverse to this map, we resort to a construction (Construction 10.11) where the movies in our morphisms must be interpretable as objects. In particular, we must stabilize our sectors to treat objects and movies on the same footing.

Combining with Theorem B, we see that a category (in the classical sense) localizes to give an  $\infty$ -category with rich geometric information—e.g., the homotopy type of spaces of stable sectorial embeddings.

**Notation 1.22** ( $\mathcal{L}\mathrm{iou}^{\diamond}$ ). The three  $\infty$ -categories  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}]$ ,  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$ , and  $\mathrm{Liou}_{\Delta}^{\diamond}$  are hence all equivalent. Each model has their advantages, but notations are all clunky, so we will use the notation

$$\mathcal{L}\mathrm{iou}^{\diamond}$$

to denote  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}]$  (and hence, up to equivalence,  $\mathrm{Liou}_{\Delta}^{\diamond}$  and  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$ ). In fact, there are three other localizations all equivalent to  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}]$  (Theorem 12.8) and we will notate them all by  $\mathcal{L}\mathrm{iou}^{\diamond}$ .

We call  $\mathcal{L}\mathrm{iou}^{\diamond}$  the  $\infty$ -category of stabilized Liouville sectors.

---

<sup>7</sup>To see this, apply the results of Dwyer-Kan [7] to a convenient model category of spaces, and note that Dwyer-Kan localization models  $\infty$ -categorical localizations—e.g., via Proposition 1.2.1 of [12].

## 1.4 Localization exhibits coherent functoriality

Assuming Theorem B and Theorem C, we may now prove Theorem 1.5:

*Proof of Theorem 1.5.* Let us first make precise the hypotheses of Theorem 1.5. The first hypothesis states that  $\mathbb{W}$  is in fact a functor  $\mathrm{Liou}_{\mathrm{str}} \rightarrow \mathcal{D}$ , where  $\mathcal{D}$  is some  $\infty$ -category admitting filtered colimits. The third assumption is the data of a diagram

$$\begin{array}{ccc} & \mathrm{Liou}_{\mathrm{str}} & \xrightarrow{\mathbb{W}} \mathcal{D} \\ T^*[0,1] \times - \uparrow & & \nearrow \mathbb{W} \\ & \mathrm{Liou}_{\mathrm{str}} & \end{array} \quad (1.6)$$

commuting up to specified natural transformation. By passing to the colimit (1.4), one obtains a functor  $\mathbb{W}^\diamond : \mathrm{Liou}_{\mathrm{str}}^\diamond \rightarrow \mathcal{D}$ . By the second assumption, and by the universal property of localization, Theorem C guarantees a factorization up to homotopy

$$\begin{array}{ccc} \mathrm{Liou}_{\mathrm{str}}^\diamond & \xrightarrow{\mathbb{W}^\diamond} & \mathcal{D} \\ \downarrow & \dashrightarrow \exists & \\ \mathcal{L}\mathrm{iou}^\diamond & & \end{array}$$

Because Theorem B tells us that the mapping spaces of  $\mathcal{L}\mathrm{iou}^\diamond$  are precisely the stabilized sectorial embedding spaces (see also Remark 1.16), this dashed functor produces the desired maps

$$\mathrm{Emb}^\diamond(M, N) \simeq \mathrm{hom}_{\mathrm{Liou}_\Delta^\diamond}(M, N) \rightarrow \mathrm{hom}_{\mathcal{D}}(\mathbb{W}^\diamond(M), \mathbb{W}^\diamond(N)).$$

The “coherence” with respect to compositions is a vague term whose specific meaning is that the dashed arrow is a functor.  $\square$

*Proof of Theorem 1.8.* For **Lag**, let  $\mathcal{D}$  be the  $\infty$ -category of stable  $\infty$ -categories; then the hypotheses of Theorem 1.5 are known. For the wrapped Fukaya category, we note that the work [9] establishes a functor from  $\mathrm{Liou}_{\mathrm{str}}$  to the ordinary category of  $A_\infty$ -categories; this in turn maps to the  $\infty$ -category of  $A_\infty$ -categories  $\mathrm{Cat}_{A_\infty}$ —this is the localization of the ordinary category of  $A_\infty$ -categories along quasi-equivalences. Thus one may take  $\mathcal{D} = \mathrm{Cat}_{A_\infty}$  and apply Theorem 1.5.  $\square$

For the record, and now that we have the notation, let us state the precise formulation of Theorem 1.8:

**Theorem 1.23** (Precise restatement of Theorem 1.8). Let  $\mathrm{Cat}_{A_\infty}$  denote the  $\infty$ -category of  $A_\infty$ -categories. Then the wrapped Fukaya category construction, along with the functoriality established in [9], induces a functor of  $\infty$ -categories

$$\mathcal{L}\mathrm{iou}^\diamond \rightarrow \mathrm{Cat}_{A_\infty}$$

which sends a Weinstein sector  $M$  to  $\mathcal{W}(M)$  and a Liouville sector more generally to  $\mathcal{W}^\diamond(M)$ . Likewise, **Lag** defines a functor from  $\mathcal{L}\mathrm{iou}^\diamond$  to the  $\infty$ -category of stable  $\infty$ -categories.

**Remark 1.24.** Of course, there are natural maps from the unstabilized embedding spaces  $\mathrm{Emb}(M, N)$  to the stabilized embedding spaces  $\mathrm{Emb}^\diamond(M, N)$ . So one would expect Theorem 1.5 to imply that, for invariants satisfying the hypotheses of the theorem, any “naturally” continuous extension of  $\mathbb{W}$  to an  $\infty$ -category of unstabilized Liouville sectors will actually factor through the stabilization,

and in particular only be sensitive to stabilized embedding spaces. We refer to this as an expectation in the present work, rather than a theorem, because we do not extend Theorem B to the level of  $\infty$ -categories. (The theorem only equates mapping spaces, without checking compositional compatibilities.) See also Remark 1.15.

**Corollary 1.25** (of Theorem C). Let  $\mathbb{W}$  and  $\mathbb{V}$  be functors  $\mathbf{Liou}_{\text{str}} \rightarrow \mathcal{D}$  equipped with stabilizing natural transformations  $\mathbb{W}(-) \rightarrow \mathbb{W}(- \times T^*[0, 1])$  and  $\mathbb{V}(-) \rightarrow \mathbb{V}(- \times T^*[0, 1])$ . Further assume that one is supplied with a natural transformation  $\eta : \mathbb{W} \rightarrow \mathbb{V}$  compatible with stabilization. Concretely,  $\eta$  is the data of a map  $\Delta^2 \times \Delta^1 \rightarrow \mathcal{Cat}_{\infty}$  where  $\Delta^2 \times \{0\}$  is the diagram (1.6), and likewise  $\Delta^2 \times \{1\}$  is the corresponding diagram for  $\mathbb{V}$ .

Then there is a natural transformation  $\eta^{\diamond} : \mathbb{W}^{\diamond} \rightarrow \mathbb{V}^{\diamond}$  between functors from  $\mathcal{Liou}^{\diamond}$  to  $\mathcal{D}$ . In particular, the diagram of mapping spaces

$$\begin{array}{ccc} \text{Emb}^{\diamond}(M, N) & \longrightarrow & \text{hom}_{\mathcal{D}}(\mathbb{W}^{\diamond}(M), \mathbb{W}^{\diamond}(N)) \\ \downarrow & & \downarrow (\eta_N)_* \\ \text{hom}_{\mathcal{D}}(\mathbb{V}^{\diamond}(M), \mathbb{V}^{\diamond}(N)) & \xrightarrow{(\eta_M)^*} & \text{hom}_{\mathcal{D}}(\mathbb{W}^{\diamond}(M), \mathbb{V}^{\diamond}(N)) \end{array}$$

commutes up to homotopy. In particular, if  $\mathcal{D}$  is tensored over spaces this implies a compatibility of module actions:

$$\begin{array}{ccc} \text{Emb}^{\diamond}(M, N) \otimes \mathbb{W}^{\diamond}(M) & \longrightarrow & \mathbb{W}^{\diamond}(N) \\ \downarrow \eta_M & & \downarrow \eta_N \\ \text{Emb}^{\diamond}(M, N) \otimes \mathbb{V}^{\diamond}(M) & \longrightarrow & \mathbb{V}^{\diamond}(N) \end{array}$$

If  $\eta : \mathbb{W} \rightarrow \mathbb{V}$  is further a natural equivalence of functors from  $\mathbf{Liou}_{\text{str}}$  to  $\mathcal{D}$ , then  $\eta^{\diamond}$  is a natural equivalence of functors from  $\mathcal{Liou}^{\diamond}$  to  $\mathcal{D}$ .

**Remark 1.26.** Again, the power is in only having to check properties of  $\eta$  that avoid mention of families of embeddings. As example applications: (I) If one has a  $G$ -action on  $M \in \mathcal{Liou}^{\diamond}$  by some topological group  $G$ , one sees that the map  $\mathbb{W}^{\diamond}(M) \rightarrow \mathbb{V}^{\diamond}(M)$  is  $G$ -equivariant (by only checking discrete properties of  $\eta$ ). (II) If one has several models for spectrally enriched wrapped Fukaya categories, one need only check that they are equivalent as functors out of  $\mathbf{Liou}_{\text{str}}$  to conclude one may compute mapping-space information about one model in terms of mapping-space information about the other.

## 1.5 Multiplicative coherence

So far, we have only discussed the “continuous” coherence of various invariants (i.e., whether invariants also map mapping spaces coherently to each other). We now turn our attention to multiplicative coherence (i.e., how invariants behave with respect to direct products of sectors—examples include Kunneth formulas).

Let us first make the simple observation that  $\mathbf{Liou}_{\text{str}}$  is symmetric monoidal under direct product. We lift this structure through the following theorem.

**Theorem D.** The functor  $\mathbf{Liou}_{\text{str}} \rightarrow \mathcal{Liou}^{\diamond}$  induces a symmetric monoidal structure on  $\mathcal{Liou}^{\diamond}$ . This symmetric monoidal structure

1. Acts on objects by direct product of Liouville sectors, and
2. Is universal for symmetric monoidal functors out of  $\mathbf{Liou}_{\text{str}}$  that invert sectorial equivalences, and that send  $T^*[0, 1]$  to a monoidally invertible object.

**Remark 1.27.** Let  $\mathcal{D}$  be a symmetric monoidal  $\infty$ -category with monoidal product  $\otimes_{\mathcal{D}}$ . Recall that an object  $X$  is called *(monoidally) invertible* if there exists an object  $Y$  for which  $X \otimes_{\mathcal{D}} Y$  is equivalent to the monoidal unit  $1_{\mathcal{D}}$ .

As an example, if  $\mathcal{D}$  is the  $\infty$ -category of chain complexes over  $R$  with  $\otimes_{\mathcal{D}}$  (a model for) the derived tensor product, then a chain complex equivalent to  $R[n]—R$  concentrated in a single degree—is an invertible object with inverse  $R[-n]$ . In particular,  $R[n]$  for  $n \neq 0$  is an invertible object that is not equivalent to the monoidal unit  $R$ .

**Remark 1.28.** The proof of Theorem D is not formal. The proof hinges on a non-trivial computation relating the space of smooth self-embeddings of  $[0, 1]^3$  to certain mapping spaces in a localization of the category of compact manifolds with corners. (See Lemma 11.7 and Remark 11.8.) Such results—relating the categorically formal process of localization to meaningful mapping spaces—are familiar from the foundations of factorization homology for manifolds [5, 4]. Indeed, we make use of a proof strategy from Lurie’s work [18] (where factorization homology is referred to as topological chiral homology).

After this computation, however, the proof of Theorem D comes down to the simple geometric fact that the  $(123)$  permutation of the coordinates of  $[0, 1]^3$  is isotopic to the identity morphism.

**Remark 1.29.** The first part of Theorem D would, to put it mildly, be a pain in the backside to prove “by hand.” To see why, the reader may appreciate that the direct product of two sectorial embeddings is not a sectorial embedding (as Liouville-flow-equivariance near infinity is violated upon taking direct product); this issue was raised already in Warning 1.4. The standard techniques to create a symmetric monoidal structure in such a setting would (i) construct collections of coCartesian fibrations whose fibers involve spaces of deformations of  $f_1 \times f_2$  to honest sectorial embeddings, and (ii) lose the attention of many potential readers. We avoid (i) (and we hope (ii) has not happened yet).

In contrast, Theorem C allows us to prove Theorem D without contemplating spaces of ways to deform products of sectorial embeddings.

Finally, it is one thing to produce a symmetric monoidal structure, and it is another to characterize it by a universal property. This is the other appealing feature of Theorem D.

**Example 1.30.** Theorem D immediately implies that the space of stable sectorial self-embeddings of a point—i.e., of  $T^*[0, 1]^\infty$ —is an  $E_\infty$ -algebra. (One could also prove this using a standard Eckmann-Hilton argument, but such an argument would get hairy quickly because one must unpack and utilize a particular model of stabilized mapping spaces.)

We remark in passing that this  $E_\infty$ -space is not group-like, and hence does not define a spectrum without passing to its units. (To see the space is not group-like, one simply embeds an exact Lagrangian copy of  $T_0^*\mathbb{R}^N$  as an interesting object in the Fukaya category of  $T^*[0, 1]^N$ , for example, as a zero object; the endofunctor resulting from the induced sectorial embedding is not an equivalence.) This is in contrast to the classical analogue: The space of smooth self-embeddings of  $[0, 1]^\infty$  is homotopy equivalent to the infinite orthogonal group  $O = O(\infty)$  (Proposition 11.11) which is group-like.

**Remark 1.31.** We gave a description of the objects of  $\mathcal{L}iou^\diamond$  (equivalently, of  $\mathcal{L}iou_\Delta^\diamond$ ) in Remark 1.16. Let us remark why the objects with  $2k \neq \dim M$  become crucial for understanding Theorem D.

First, the symmetric monoidal structure on objects is given by  $(M, k) \otimes (M', k') \simeq (M \otimes M', k + k')$  where  $M \otimes M'$  is the manifold  $M \times M'$  equipped with the product Liouville structure (Definition 2.9).

Note that via the inclusion  $\mathrm{Liou}_{\mathrm{str}} \rightarrow \mathcal{L}\mathrm{iou}^\diamond$ , the sector  $T^*[0, 1]$  is identified with the object  $(T^*[0, 1], 0)$ . This object is not the monoidal unit. (The object  $(T^*[0, 1], 1) \sim (\mathbb{R}^0, 0)$  is the unit.) But it is invertible, with inverse  $(\mathbb{R}^0, 1)$ , as

$$(\mathbb{R}^0, 1) \otimes (T^*[0, 1], 0) \simeq (\mathbb{R}^0 \otimes T^*[0, 1], 1 + 0) \simeq (T^*[0, 1], 1) \simeq (\mathbb{R}^0, 0).$$

One sees that stabilization thus renders the operation  $- \times T^*[0, 1]$  invertible, in a symmetric monoidal fashion.

Here is a corollary of Theorem D. It is a symmetric monoidal version of Theorem 1.5.

**Theorem 1.32.** Let  $\mathcal{D}$  be a symmetric monoidal  $\infty$ -category and let  $\mathbb{W} : \mathrm{Liou}_{\mathrm{str}} \rightarrow \mathcal{D}$  be a symmetric monoidal functor sending  $T^*[0, 1]$  to a monoidally invertible object of  $\mathcal{D}$ , and inverting sectorial equivalences. Then  $\mathbb{W}$  canonically extends to a symmetric monoidal functor  $\mathcal{L}\mathrm{iou}^\diamond \rightarrow \mathcal{D}$ .

**Remark 1.33.** Suppose we have a symmetric monoidal functor  $\mathbb{W}$  from  $\mathrm{Liou}_{\mathrm{str}}$  to some symmetric monoidal  $\infty$ -category  $\mathcal{D}$ , and that  $\mathbb{W}$  sends  $T^*[0, 1]$  to an invertible object, while sending sectorial equivalences to equivalences in  $\mathcal{D}$ . By Theorem 1.32, we are guaranteed a symmetric monoidal extension of  $\mathbb{W}$ , which we will denote by  $\widetilde{\mathbb{W}} : \mathcal{L}\mathrm{iou}^\diamond \rightarrow \mathcal{D}$ . Let us also denote  $\mathbb{W}(T^*[0, 1])$  by  $\mathcal{L}$  (to invoke a line bundle—a prototypical example of an invertible non-unit).

The object  $T^*[0, 1]$  of  $\mathrm{Liou}_{\mathrm{str}}$  is then identified with the object  $(T^*[0, 1], 0)$  of  $\mathcal{L}\mathrm{iou}^\diamond$ , so  $\widetilde{\mathbb{W}}(T^*[0, 1], 0) \simeq \mathcal{L}$ . More generally, we have that  $\widetilde{\mathbb{W}}(T^*[0, 1], k) \simeq \mathcal{L}^{\otimes_{\mathcal{D}} 1-k}$ , where the negative tensor powers make sense because  $\mathcal{L}$  is invertible. Even more generally, for any Liouville sector  $M$ , we have that  $\widetilde{\mathbb{W}}(M, k) \simeq \mathbb{W}(M) \otimes_{\mathcal{D}} \mathcal{L}^{\otimes_{\mathcal{D}} -k}$ . Note that these equivalences are specified by the symmetric monoidal structure of the functor  $\widetilde{\mathbb{W}}$ .

Thus, in contrast with the outcome of Theorem 1.5, the data of  $\widetilde{\mathbb{W}}$  is not formally determined by its restriction to the full subcategory of  $\mathcal{L}\mathrm{iou}^\diamond$  with  $\dim M - 2k = 0$ . (See Remark 1.16.)

**Remark 1.34.** In contrast with Theorem 1.5, there is no need to assume that  $\mathcal{D}$  admits filtered colimits, and there is no assumption on  $\mathbb{W}$  admitting a stabilization natural transformation  $\mathbb{W}(-) \rightarrow \mathbb{W}(- \times T^*[0, 1])$ .

Indeed, there is a symmetry-breaking in the literature when the natural transformation  $\mathbb{W}(M) \rightarrow \mathbb{W}(M \times T^*[0, 1])$ , is constructed; in the wrapped Fukaya category context, one typically chooses a cotangent fiber  $T_p^*[0, 1]$  of  $[0, 1]$  and (on objects) the natural transformation is defined as  $L \mapsto L \times T_p^*[0, 1]$ . Meanwhile, the stabilization of Theorem D does not rely on such symmetry-breaking data; instead, the symmetric monoidal structure, together with a check of a *property* (that  $T^*[0, 1]$  is sent to a monoidally invertible object) guarantees the existence of a symmetric monoidal functor from the stabilized localization.

Finally, we note the utility of Theorem D and versatility of  $\mathcal{L}\mathrm{iou}^\diamond$ . Even when the invariant  $\mathbb{W}(T^*[0, 1])$  is not the monoidal unit (hence when  $\mathbb{W}(M \times T^*[0, 1])$  need not necessarily equal  $\mathbb{W}(M)$ ) one can produce a meaningful multiplicative invariant. A simple example is the functor to chain complexes sending  $M$  to its compactly supported and supported-away-from-the-boundary deRham forms, which sends  $T^*[0, 1]$  to a complex equivalent to  $\mathbb{R}$  concentrated in cohomological degree 2. In particular, while invariants of the type in Theorem 1.5 are insensitive to the underlying dimension of a sector  $M$  in  $\mathrm{Liou}_{\mathrm{str}}$ , invariants arising in Theorem 1.32 may still be sensitive to dimension.

Unlike Theorem 1.8, the verifications that  $\mathrm{Lag}$  and  $\mathcal{W}$  may be promoted to be symmetric monoidal are yet to be written in the literature. For  $\mathrm{Lag}$ , the problem is still wide open, even for the Weinstein case. For  $\mathcal{W}$ , the Kunneth formula is proven for wrapped categories of Weinsteins, while a proof of the full symmetric monoidal structure (even for Weinsteins) is not yet recorded,

though the details are not expected to be onerous. Regardless, the reader may appreciate that lifting the *discrete* functors out of  $\mathrm{Liou}_{\mathrm{str}}$  to be symmetric monoidal is far easier than trying to write, by hand, a compatibility between continuous coherences and multiplicative coherences. Theorem 1.32 eliminates the necessity for writing down such a compatibility.

**Corollary 1.35.** Let  $\mathbb{W} : \mathrm{Liou}_{\mathrm{str}} \rightarrow \mathcal{D}$  be a symmetric monoidal functor satisfying the hypotheses of Theorem 1.32. Then whenever  $M$  is an  $E_n$ -algebra in  $\mathcal{L}\mathrm{iou}^\diamond$ ,  $\mathbb{W}(M)$  is an  $E_n$ -algebra in  $\mathcal{D}$ .

**Remark 1.36.** In particular, if  $\mathbb{W}$  lands in  $\infty$ -categories, any  $E_n$ -algebra  $M$  in stabilized Liouville sectors induces an  $E_{n+1}$ -algebra structure on endomorphisms of the unit of  $\mathbb{W}(M)$ . At present, we know no way of producing algebra objects in stabilized Liouville sectors, but in [16], we show that upon localizing with respect to another natural class of sectorial maps (the *subcritical* maps), one can find a plethora of  $E_\infty$ -algebras, and these conjecturally represent symmetric monoidal localizations of the stable homotopy category.

## 1.6 Results for Liouville sectors with tangential structures

Finally, by now we know that we should equip our Liouville sectors with tangential structures to, for example, construct invariants over different base rings. (Without a trivialization of  $\det(TM)^{\otimes 2}$ , for example,  $\mathcal{W}(M)$  is not  $\mathbb{Z}$ -graded.) We prove in this work that all the results stated so far have analogues in the setting where our sectors are equipped with tangential structures. Let us make this precise.

Fix  $E_\infty$ -spaces  $B'$  and  $B$ , and fix  $E_\infty$  maps  $B' \rightarrow B$  and  $BU \rightarrow B$ . Then for any Liouville sector  $M$ , by virtue of its natural map to  $BU$  classifying the compatible-with-symplectic-form complex tangent bundle, we may ask for a  $B'$ -reduction of its  $B$ -tangential structure.

**Example 1.37.** For example, if  $BU \rightarrow BU(1) = B$  classifies  $\det^2$  and  $B'$  is a point, a  $B'$ -reduction amounts to a trivialization of the stable  $\det^2$ -bundle.

We let  $(\mathrm{Liou}_{\mathrm{str}}^\diamond)_{/(B' \rightarrow B)}$  denote the  $\infty$ -category whose objects are Liouville sectors equipped with  $B'$ -reductions of  $B$ -structures, and where morphisms are pairs  $(f, h)$  where  $f$  is a strict sectorial inclusion, and  $h$  is a homotopy of  $B'$ -reductions from the reduction on the domain to the reduction pulled back along  $f$ . We let  $\mathfrak{eq}_{/B' \rightarrow B} \subset (\mathrm{Liou}_{\mathrm{str}}^\diamond)_{/(B' \rightarrow B)}$  denote those  $(f, h)$  where  $f$  is a sectorial equivalence.

Finally, we let  $\mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^\diamond$  denote the  $\infty$ -category of stabilized Liouville sectors with reductions of  $B$ -structures to  $B'$ .

**Theorem E.** The map from the localization

$$(\mathrm{Liou}_{\mathrm{str}}^\diamond)_{/(B' \rightarrow B)}[\mathfrak{eq}_{/B' \rightarrow B}^{-1}] \rightarrow \mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^\diamond$$

is an equivalence of  $\infty$ -categories.

We have an induced symmetric monoidal structure on  $\mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^\diamond$  characterizable by two different universal properties—one induced from the localization, and the other by virtue of being a pullback of symmetric monoidal  $\infty$ -categories. See Section 13.

## 1.7 The geometric insights

This paper emerged as the authors were trying to create the correct formalism for articulating a *symmetric monoidal* localization process for Liouville sectors. (Indeed, the foundations from



the present work allow for the symmetric monoidal localizations—for categories of sectors, and conjecturally of wrapped Fukaya categories themselves—in [16].)

While we have so far conveyed the  $\infty$ -categorical results, let us now speak to the new geometric insights we incorporate in the present work. The categorical results were not at all obligated to exist based on the geometry known to us when we first began this project.

The first innocent observation is that while stabilization  $M \mapsto M \times T^*[0, 1]$  is easy to define for objects (sectors), it does not preserve the standard class of morphisms (sectorial embeddings). At first it seemed natural to thus create an  $\infty$ -category equipped with data of smooth isotopies through sectorial embeddings, to deform *morphisms* to render them sectorial after stabilizing.

However, also floating in the air was the knowledge that carrying the data of a deformation of *Liouville structures* (on objects) was natural for both Fukaya-categorical and symplectic reasons.

By Moser’s Trick, it seems precisely articulable that carrying auxiliary data of deformations of Liouville structures is somehow equivalent to carrying auxiliary data of deforming the morphisms themselves. We do not explore this equivalence, but suffice it to say that—faced with the fork in the road of deforming morphisms or deforming Liouville structures—we settled on carrying deformations of Liouville structures as the  $\infty$ -categorical data. This is how movies, and barycentric subdivision diagrams, came about. Each object  $M \times T^*\Delta^k$  at a barycenter encodes a  $\Delta^k$ -family of Liouville structures on  $M$ .

Then, to preserve closure under direct products, it was natural to consider only maps that were *strict* sectorial embeddings, then identify a class of deformations closed under products. This class had to be made surprisingly large to accommodate both useful examples and closure under products. (See Definition 2.32—in particular, our deformations of Liouville structures are not through only compactly supported deformations.) The key theorem that justifies this choice of class is that the space of strict-morphisms-equipped-with-this-class-of-deformations (where the deformations need not be compactly supported) is homotopy equivalent to the usual space of all sectorial embeddings (which respect the Liouville form up to  $d$  of a *compactly* supported function). See Theorem 4.18 and the equivalences of (2.7). In other words, up to homotopy equivalence, we can recover the “standard” morphism spaces for Liouville sectors. Carrying around deformations does not change the homotopy type of morphism spaces.

Then, we had to prove some generalizations of standard facts from the differential topology of manifolds with corners: isotopy extension, the ability to shrink a Liouville sector to have image away from its own boundary, and the ability to create a single (higher-dimensional) object for which a single map encodes a family of maps for lower-dimensional objects. We call these objects *movies* (Section 2.2), and they generalize a construction we learned from [8].

## 1.8 Later works and other future directions

We will show in later works that  $\mathcal{L}\text{iou}^\diamond$  has finite colimits.

**Remark 1.38.** Note that  $\text{Liou}_\Delta$  (i.e., before stabilizing) does not admit all colimits, for the same reason that the category of sets with injections does not. For example, if  $X$  and  $Y$  are both non-empty and zero-dimensional sectors, there is no categorical coproduct  $X \coprod Y$ .

As we will show in later works, direct product is compatible with finite colimits in each variable. So  $\mathcal{L}\text{iou}^\diamond$  can be Ind-completed to be a presentably symmetric monoidal  $\infty$ -category. Further, building on results of [10] one can show that the wrapped Fukaya category functor preserves all colimits. By the adjoint functor theorem, the wrapped Fukaya category functor admits a right adjoint. In particular, to any stable  $\infty$ -category, we can canonically associate an Ind-stabilized-Liouville-sector.

Let us note that neither the unit nor the counit of this adjunction is expected to be an equivalence. Regardless, the adjunction shows that for certain Ind-Liouville sectors, the space of morphisms into them may be computed purely algebraically as a space of functors. The ability to even state the ideas in the previous paragraphs precisely is, we note, another outcome of the present work.

Another direction of interest is the articulation of the Weinstein version of the tools we use here. Let  $\text{Wein}^\diamond \subset \mathcal{L}\text{iou}^\diamond$  denote the full subcategory of those (stabilized) sectors admitting Weinstein structures.<sup>8</sup> This is of course a natural  $\infty$ -category to study. However, we do not characterize  $\text{Wein}^\diamond$  as a localization of anything, and in particular, we do not give a proof that invariants defined only for Weinstein sectors (and not for Liouville sectors generally) induce compatible maps from stabilized mapping spaces. A notable invariant which, at present, is only defined for Weinstein sectors is the microlocal category  $\mathfrak{Sh}(M)$  of Nadler-Shende [24].

Regardless, it is straightforward to construct a Weinstein version of  $\text{Liou}_\Delta$ —an  $\infty$ -category whose simplices consist of barycentric subdivision diagrams of *Weinstein* sectors with strict inclusions. In a previous draft of this work, we expressed our belief that morphism spaces arising from  $\text{Wein}_\Delta$  are the only candidate we know for defining a rich “space of Weinstein morphisms.” However, work in progress seems to show that this candidate space is homotopy equivalent to a fully faithful space of Liouville embeddings (i.e., a subspace consisting of certain connected components of the Liouville embedding spaces.) Moreover, we expect the methods of this work carry over to show that  $\text{Wein}_\Delta^\diamond$  is a localization of  $\text{Wein}^\diamond$  along Weinstein equivalences. (A Weinstein equivalence is a sectorial embedding  $f : M \rightarrow N$  admitting an inverse-up-to-Weinstein-isotopy  $g : N \rightarrow M$ . More precisely, the isotopies  $g \circ f \sim \text{id}$  and  $f \circ g \sim \text{id}$  have movies—Example 7.17—that admit Weinstein structures.) Thus, for every pair of objects  $X, Y$ , it seems that the natural map  $\text{hom}_{\text{Wein}_\Delta^\diamond}(X, Y) \hookrightarrow \text{hom}_{\mathcal{L}\text{iou}^\diamond}(X, Y)$  is an inclusion of connected components, and that invariants like those of Nadler-Shende receive actions from Liouville embedding spaces whose elements are Liouville-homotopic to maps respecting Weinstein structures.

It seems to remain an important open problem to construct an  $\infty$ -category with meaningful spaces of “Weinstein morphisms” (of which no homotopically useful definition exists at present). A “correct” definition is delicate, analogous to the way that one must carefully define what we mean by families of handle decompositions (Morse functions) to create invariants sensitive to simple homotopy equivalences and K-theoretic data.

## 1.9 This paper is $\infty$ -categorical in most of its tools

While we hope the results of this work will appeal to both homotopy theorists and symplectic geometers, we remark that we rely heavily on  $\infty$ -categorical machinery.

## 1.10 Acknowledgments

We would like to thank Yasha Eliashberg for very helpful conversations. The first author was supported by the NSF postdoctoral fellowship, Award #1705128. The first and second authors were partially supported by the Simons Foundation through grant #385573, the Simons Collaboration on Homological Mirror Symmetry. The third author was supported by a Texas State University Research Enhancement Program grant, the Texas State University Presidential Seminar and Valero

---

<sup>8</sup>We do not define in this paper any notion of maps between Weinstein sectors that are “compatible” with Weinstein structures; for our purposes, a map of Liouville sectors-that-happen-to-admit-Weinstein-structures is simply a (not necessarily strict) sectorial embedding.

Awards, and by an NSF CAREER Grant under Award Number 2044557; he would also like to thank Rune Haugseng and Jacob Lurie for helpful communications.

## 2 Some Liouville geometry

Here we collect some facts about Liouville geometry and establish notation, conventions, and definitions along the way. The main results original to this work are:

- A movie construction, which provides a way to construct a single (higher-dimensional) Liouville sector encoding a family of Liouville structures on a manifold  $M$  (Proposition 2.42). This is a higher-dimensional, and minor, generalization of a construction of Eliashberg [8].
- A “shrinking isotopy” existence theorem assuring that we may always find a self-equivalence of a Liouville sector whose image is bounded away from the sectorial boundary (Section 2.4).
- An isotopy extension theorem for Liouville sectors (Proposition 2.54), and
- That the space of “embeddings equipped with deformation data rendering the embeddings strict” is homotopy equivalent to the space of sectorial embeddings (Section 4.5).

### 2.1 Basic definitions

#### 2.1.1 Liouville sectors

The following agrees with the notion from [10] of a *straightened* Liouville sector with corners. Because sectors will always be straightened in our work, we remove the adjective.

**Definition 2.1** (Liouville sector). Fix an exact symplectic manifold  $(M, \omega = d\lambda)$  possibly with corners, together with the data, for each  $x \in \partial^i M$  in a codimension  $i$  corner  $i \geq 1$ , of a neighborhood  $\text{Nbhd}(x)$  inside  $M$  and a codimension-preserving symplectic submersion

$$\pi_x: \text{Nbhd}(x) \rightarrow T^*[0, 1]^i. \quad (2.1)$$

We say this collection of data is a *Liouville sector* if it satisfies the following:

- (S1)  $\lambda$  has finite type. This means that  $M$  admits a proper, smooth function  $M \rightarrow \mathbb{R}_{\geq 0}$  which, outside some compact subset of  $M$ , is linear with respect to the Liouville flow of  $M$ .
- (S2) Each  $\pi_x$  is flat, and  $\lambda$  is split with respect to  $\{\pi_x\}_{x \in \partial M}$ . This means: Using the flat splitting of  $\text{Nbhd}(x)$  as a product of a fiber of  $\pi_x$  and the image of  $\pi_x$ ,  $\lambda|_{\text{Nbhd}(x)} = \lambda^{\text{fiber}} + \pi_x^* \mathbf{p} d\mathbf{q}$ . (See Remark 2.2.)
- (S3) If  $y \in \partial^j M \cap \text{Nbhd}(x)$  with  $j \leq i$ , then on the overlap we have  $\pi_y = \pi_{yx} \circ \pi_x$ , where  $\pi_{yx}: T^*[0, 1]^i \rightarrow T^*[0, 1]^j$  is a projection to  $j$  components (not necessarily respecting the order of coordinates).

We will usually denote a Liouville sector by  $(M, \lambda)$  or just  $M$ , leaving the family of projections  $\{\pi_x\}$  implicit.

**Remark 2.2** (The splitting). Let us explain the splitting in (S2). When  $\pi: E \rightarrow B$  is a symplectic submersion—so that  $E$  is symplectic and the symplectic form on  $E$  restricts to a symplectic form on each fiber of  $\pi$ —one has a natural horizontal distribution (given by the symplectic orthogonal to the vertical tangent spaces). We say that the submersion is flat if this distribution is integrable, and in particular, defines a flat connection. We assume this of our projection maps, so  $\pi$  canonically splits  $E$  as a smooth manifold, locally on  $E$ . In particular, note that we may always shrink  $\text{Nbhd}(x)$

so that  $\text{Nbhd}(x)$  itself admits a splitting; indeed, the split expression for  $\lambda$  in (S2) assumes we have chosen such a neighborhood for every  $x$ .

To see a sample consequence, let  $v_\lambda$  denote the Liouville vector field (obtained as the symplectic dual to  $\lambda$ ). Condition (S2) guarantees not only a splitting of the symplectic form, but also a splitting of the Liouville form; as a result,  $v_\lambda$  splits. In particular, because each  $\pi_x$  is codimension-preserving, we observe that the Liouville flow—i.e., the flow of  $v_\lambda$ —is tangent to each boundary and corner stratum of  $M$ . Moreover, because the Liouville form on the  $T^*[0, 1]^i$  factor is  $\mathbf{p}d\mathbf{q}$ , any trajectory of the Liouville flow which begins away from  $\partial M$  must remain away from  $\partial M$ .

A further consequence, by shrinking  $\text{Nbhd}(x)$  so the image of  $\pi_x$  is a box, is that the composite projection  $\text{Nbhd}(x) \xrightarrow{\pi_x} T^*[0, 1]^i \xrightarrow{\mathbf{q}} [0, 1]^i$  induces a collaring of the stratum of  $x$  in  $\text{Nbhd}(x)$  by choosing a fiber of  $\mathbf{q} \circ \pi_x$  and writing  $\text{Nbhd}(x) \cong \text{fiber} \times \text{image}(\mathbf{q} \circ \pi_x)$ . This exhibits a collar of  $\text{Nbhd}(x)$  as a smooth manifold with corners.

**Remark 2.3** (Completeness). Let  $\rho : M \rightarrow \mathbb{R}_{\geq 0}$  be a smooth function as in Condition (S1), and let  $v_\lambda$  denote the Liouville vector field, defined by  $d\lambda(v_\lambda, -) = \lambda(-)$ . The linearity in Condition (S1) means that  $v_\lambda(\rho) = \rho$  (outside some compact subset of  $M$ ). This has the following consequences.

First, all large-enough positive real numbers are regular values of  $\rho$ . Because  $\rho$  is proper and takes non-negative values, we thus see that  $K = \rho^{-1}([0, R])$  for large enough  $R$  is some compact, smooth, codimension-zero submanifold-with-corners of  $M$ . Because the Liouville flow is tangent to the boundary and corner strata of  $M$  (Remark 2.2), every point in the complement of  $K$  “flows to infinity” under the Liouville flow. So,  $M$  is the completion of a compact Liouville domain-with-corners (where the contact boundary may also have corners). As a result, one has a well-defined “contact boundary at infinity” (possibly with corners) which we will denote by

$$\partial_\infty M.$$

Moreover, because  $M$  is the completion of a compact domain under positive Liouville flow, the Liouville flow is complete (in both the positive and negative flow directions).

This has the following implication for our submersions. For any  $x \in \partial^i M$ , there is some point  $x'$  in the same codimension  $i$  stratum as  $x$  whose image under the projection  $\pi_{x'}$  is in the zero section of  $T^*[0, 1]^i$ . For example, one can take  $x'$  to be the  $t \rightarrow \infty$  limit of  $x$  under the negative Liouville flow.

Before we further explore Definition 2.1, we recall a fact from smooth topology. For any compact smooth manifold  $X$  with corners and any element  $x \in \partial^i X$  of the codimension  $i$  locus, consider the stratum  $(\partial^i X)_x$  of  $x$ . (Such a stratum is always an  $i$ -dimensional manifold without boundary or corners, always a connected component of  $\partial^i X$ , and often non-compact.) The fact we wish to recall is that  $(\partial^i X)_x$  admits a compactification

$$\overline{(\partial^i X)_x}$$

to a smooth  $i$ -dimensional manifold possibly with corners, equipped with an immersion to  $X$  extending the embedding of  $(\partial^i X)_x$ .

Despite our overline notation, this compactification is not a closure in  $X$ , and the immersion is often not an embedding. One construction of this compactification is as follows. A point  $\tilde{y}$  in the compactification is an equivalence class of a triplet  $(y, u, \phi)$  where  $y$  is a point in the closure of  $(\partial^i X)_x$  inside  $M$ ,  $u$  is an element of  $(\mathbb{R}_{\geq 0})^{\dim M - i}$ , and  $\phi$  is a germ about  $u$  of a codimension-preserving, smooth embedding of some open subset of  $(\mathbb{R}_{\geq 0})^{\dim M - i}$  into the closure of  $(\partial^i X)_x$  inside  $X$ , satisfying  $\phi(u) = y$ . One declares an equivalence  $(y, u, \phi) \sim (y', u', \phi')$  when  $y = y'$  and there exists a germ of a codimension-preserving diffeomorphism  $j$  for which  $j(u) = u'$ .

**Example 2.4.** Let  $X$  be the manifold with corners obtained by smoothing three of the corners of the square  $[0, 1]^2$ . The codimension 2 stratum is its own compactification; it consists of a point. Meanwhile, the codimension 0 stratum (the interior of  $X$ ) compactifies to  $X$  itself.

The codimension 1 case is of most interest. The lone codimension-one stratum  $\partial^1 X$  is diffeomorphic to an open interval, and its compactification  $\overline{\partial^1 X}$  is diffeomorphic to a closed interval (as opposed to the closure of  $\partial^1 X$  in  $X$ , which is homeomorphic to a circle), immersing to the square with both endpoints mapping to the unsmoothed corner.

**Remark 2.5.** Note that the immersion  $\overline{\partial^1 X} \rightarrow X$  restricts to an embedding on the original  $\partial^1 X$ . We will see identical behavior below when a sector  $\overline{F}$  immerses to  $M$ , and the immersion restricts to an embedding on the interior  $F \subset \overline{F}$ .

**Remark 2.6** (Liouville sectors associated to strata). We will now see that every stratum of codimension  $i$  has the structure of a trivial fiber bundle over  $\mathbb{R}^i$ , with fiber  $F$  naturally identified as the interior of a Liouville sector  $\overline{F}$ . This fiber depends not only on the codimension, but also on the stratum itself (i.e., on the connected component within the codimension  $i$  locus).

Let  $K$  be the compact subset from Remark 2.3, and fix an element  $x \in K \cap \partial^i M \subset \partial^i K$ ; we denote the stratum of  $\partial^i K$  containing  $x$  by  $(\partial^i K)_x$ . By compactness of  $K$ , (S3) guarantees a submersion

$$\pi : \text{Nbhd}((\partial^i K)_x) \rightarrow T^*[0, 1]^i$$

defined on a neighborhood of the whole of  $(\partial^i K)_x$  (and not only locally).

As per the discussion preceding this remark, let us compactify  $(\partial^i K)_x$  to a smooth manifold  $\overline{(\partial^i K)_x}$  of dimension  $\dim M - i$ , possibly with boundary and corners, equipped with a natural immersion to  $M$ .  $\overline{(\partial^i K)_x}$  inherits a natural 1-form by pulling back  $\lambda$ , and is compact because  $K$  is. Again by compactness and (S3), the maps  $\{\pi_y\}_{y \in \text{closure of } (\partial^i K)_x \text{ in } M}$  glue together to give a well-defined submersion

$$\overline{\pi} : \overline{(\partial^i K)_x} \rightarrow T_{\{0\}}^*[0, 1]^i \cong \mathbb{R}^i$$

to the cotangent fiber of the origin. Note that (i) there is no well-defined analogue of  $\overline{\pi}$  on the closure of  $(\partial^i K)_x$  inside  $M$ , precisely because of the ambiguity in permuting the coordinates of  $T^*[0, 1]^i$  in (S3), and (ii) the fibers of  $\overline{\pi}$  are compact manifolds with corners.

By Remark 2.3, the image of  $\overline{\pi}$  contains the origin. Let  $\overline{F}_K$  be the fiber of  $\overline{\pi}$  above the origin, and consider its completion  $\overline{F}$  with respect to the Liouville flow (with respect to the pullback of  $\lambda$ ). Then  $\overline{F}$  admits a natural immersion to  $M$  by extending, via the Liouville flow, the immersion of  $\overline{F}_K$ . We observe  $\overline{F}$  is a Liouville sector, where (S1) is satisfied for example by pulling back a  $\rho$  exhibiting (S1) for  $M$ .

Letting  $F$  denote the interior of  $\overline{F}$ , the immersion of  $\overline{F}$  restricts to an embedding of  $F$  to  $M$  (see Remark 2.5). So we may identify  $F$  with the Liouville completion of the fiber of  $\pi$  above the origin of  $T_{\{0\}}^*[0, 1]^i$ . Next, note that the splitting condition (S2) guarantees that the fibers of  $\pi$  are all isomorphic, hence the fibers of  $\overline{\pi}$  are isomorphic; this shows that each codimension  $i$  stratum  $(\partial^i M)_x$  of  $M$  fibers over  $T_{\{0\}}^*[0, 1]^i \cong \mathbb{R}^i$  with fibers isomorphic to  $F$ , though we caution that the only fiber-inclusion into  $M$  commuting with the Liouville flow is the inclusion of the fiber above the origin.

Finally, we note that by realizing  $(\partial^i M)_x$  as a Liouville completion of  $(\partial^i K)_x$ , and using the compactness of  $K$ , we may arrange for each  $\pi_y, y \in (\partial^i M)_x$  to have image with uniform  $q$  coordinate, giving a collaring of the entire stratum:

$$\text{Nbhd}((\partial^i M)_x) \cong F \times T^*[0, q_0]^i \tag{2.2}$$

where  $F$  is the interior of the Liouville sector associated to the stratum of  $x$  and  $q_0 > 0$  is some positive real number.

**Example 2.7.** Consider the case where  $M$  has boundary but no corners. By Remark 2.6, each boundary component of  $M$  has associated to it a canonical Liouville sector  $F$  (where  $F$  may depend on the boundary component, and  $F = \overline{F}$  because  $M$  has no corners).

Letting  $q_0$  be as in (2.2), choose a diffeomorphism from  $[0, q_0)$  to  $[0, 1)$ . Then every boundary component of  $M$  has a neighborhood admitting an identification—as smooth manifolds with boundary equipped with a 1-form—with  $F \times T^*[0, 1)$ . (Where  $F$  again depends on the boundary component.) As a smooth manifold with boundary, we thus have a natural collaring

$$(F \times T_{\{0\}}^*[0, 1)) \times [0, 1).$$

identifying the boundary component of  $M$  with the smooth manifold  $F \times T_{\{0\}}^*[0, 1) \cong F \times \mathbb{R}$ .

**Definition 2.8** (Sectorial boundary). Let  $M$  be a Liouville sector. Then the *sectorial boundary* of  $M$  is the boundary of  $M$  when considered as a smooth manifold with corners—in other words, the union of all boundary and corner strata of  $M$ .

**Definition 2.9** (Products). Given two Liouville sectors  $(M, \lambda^M), (N, \lambda^N)$ , we define the product sector  $M \otimes N$  to be  $(M \times N, \lambda^M + \lambda^N)$  with the induced projection maps  $\{\pi_{(x,y)} = \pi_x \times \pi_y\}$ .

For any integer  $k \geq 1$ , the sector  $M \otimes T^*D^k$  is called a *stabilization* of the sector  $M$ .

**Remark 2.10.** In this work, we will frequently consider product manifolds whose Liouville forms do not split (see Construction 2.34). We reserve the notation  $M \otimes N$  for the split setting.

**Example 2.11.** Let  $Q$  be a smooth manifold with corners. Then for every  $x$  in a codimension  $i$  stratum,  $\text{Nbhd}(x)$  admits a (codimension-preserving) collaring projection  $\text{Nbhd}(x) \rightarrow [0, 1)^i$ . The space of such projections is contractible, and a choice of such projection for each  $x$  induces projections (2.1) on  $T^*Q$ . With this structure, it is standard to see that  $T^*Q$  is a Liouville sector in the sense of Definition 2.1.

For more examples, we refer the reader to Section 2 of [16], and especially to the construction of sectors from stopped domains.

### 2.1.2 Families of exact deformations of Liouville structures

Though most of the literature contemplates 1-parameter families of Liouville structures, we will contemplate families of Liouville structures parametrized by smooth manifolds  $S$ . In fact, we will typically take  $S$  to be a simplex  $\Delta^k$  (which is a smooth manifold with corners).

**Definition 2.12** (Deformations of Liouville structures). Fix a Liouville sector  $(N, \lambda)$ . Let  $S$  be a smooth manifold, possibly with corners, and fix a smooth function  $h : N \times S \rightarrow \mathbb{R}$ . Then we say that the family of 1-forms

$$\{\lambda_s = \lambda + d_N h_s\}_{s \in S} \tag{2.3}$$

is an  $S$ -parametrized *exact family* of Liouville structures, or an exact *deformation* of Liouville structures, if for every  $s \in S$ ,  $(N, \lambda_s)$  is a Liouville sector, and

- (T) There exists a function  $R : N \times S \rightarrow \mathbb{R}_{\geq 0}$  such that, locally on  $S$ ,  $R$  is proper and one can find a compact subset  $K \subset N$  for which  $R_s$  is  $\lambda_s$ -linear outside  $K$ .

Here, by  $R$  being proper locally on  $S$ , we mean that for every  $s \in S$ , there is a compact neighborhood  $A_s$  of  $s$  so that  $R|_{N \times A_s}$  is proper. The local existence of  $K$  means: Let For every  $s \in S$ , there exists an open  $U \subset S$  with  $s \in U$  and a compact subset  $K_s \subset M$  for which  $\lambda_{s'}(X_{R_{s'}})(x) = R_{s'}(x)$  for all  $x \notin K$  and for all  $s' \in U$ .

Here,  $X_{R_{s'}}$  is the Hamiltonian vector field on  $N$  associated to  $R_{s'}$ .

**Remark 2.13.** For every  $s \in S$ ,  $h_s : N \cong N \times \{s\} \rightarrow \mathbb{R}$  is the restriction of  $h$ .  $d_N h_s = dh_s$  is the de Rham derivative of  $h_s$  with respect to the de Rham operator on  $N$ . In Definition 2.12, the subscript  $N$  is used to emphasize that no derivatives are being taken in the  $S$  directions.

**Remark 2.14.** The reader may ask, “a deformation of *what*?” If one chooses a basepoint  $s_0 \in S$ , then one may consider the  $S$ -parametrized family as deforming  $\lambda_{s_0}$ . In the absence of a natural basepoint, “family” may be a better term than deformation. In most of the examples of this paper,  $S$  will be a simplex with ordered vertices, hence there will be a natural basepoint given by the terminal vertex of the simplex.

**Remark 2.15.** Condition (T) is a “tameness-near-infinity” assumption on the family. Let us motivate it. It turns out there are various topologies one could put on the space of possible deformations. We choose to topologize as follows: One filters the set of all possible deformations by the compact subsets of  $K$ —for any  $K$ , we consider only deformations that are all linear outside  $K$ , meaning roughly that we always have a symplectization coordinate outside of  $K$ . For every  $K$ , we give the  $K$ -supported deformation space a weak  $C^\infty$  topology, then we topologize the space of all deformations by taking the direct limit topology (by filtering by the variable  $K$ ).<sup>9</sup> Unwinding definitions, this means that if we have an  $S$ -parametrized family of deformations, then locally on  $S$ , we can find compact  $K$  (which is not uniformly defined over all  $S$ ) controlling the linearity of the family.

This condition did not appear out of thin air. First, it is natural to ask for a single  $K$  to control behavior-at-infinity when  $S$  is compact. Then, the local existence of  $K$  is an inevitable consequence of sheaffifying this desideratum (and hence when incorporating non-compact  $S$ ).

**Remark 2.16.** In most of our examples,  $S$  will be compact. When  $S$  is compact, we may assume  $R : N \times S \rightarrow \mathbb{R}_{\geq 0}$  is itself proper. We can further choose a single  $K$  so that regardless of  $s$ ,  $R_s$  is  $\lambda_s$ -linear outside the fixed  $K$ . Then, if necessary, we may enlarge  $K$  so that  $K$  contains the union of all the skeleta/cores (i.e., the locus of points that do not escape to infinity) of  $(M, \lambda_s)$ . This enlargement ensures that, outside of this  $K$ , any trajectory of the Liouville flow of  $\lambda_s$  will escape toward infinity (as evidenced by the growth of  $R$ ).

**Example 2.17.** Consider an exotic Weinstein structure on  $\mathbb{R}^{2n}$  that contains a standard Weinstein  $\mathbb{R}^{2n}$  in the interior. One can then isotope the “exotic” regions toward infinity while leaving the standard copy in place. One can arrange for this to result in a smooth family of Liouville structures, defined on  $S = [0, 1]$ , from the exotic structure to the standard structure. However, this family does not satisfy Condition (T). In particular, Condition (T) precludes such ill-behaved “isotopies” between exotic and standard structures. (See also Remark 11.24 of [6].)

**Definition 2.18.**  $S$ -parametrized exact deformations (Definition 2.12) will further be called (in order of increasing restrictiveness):

- *Bordered* if for each  $s$ ,  $\lambda_s$  respects the splitting (S2).
- *Interior* if  $\lambda_s$  is constant (i.e.,  $s$ -independent) near  $\partial N$  and bordered.

---

<sup>9</sup>See also Definition 2.57.



- *Compactly supported* if (i) locally in  $S$ , there exists a compact set  $K \subset N$  for which  $\text{supp}(\lambda_s - \lambda_0) \subset K$ , and (ii) the deformation is interior.

**Remark 2.19.** For a deformation  $\{\lambda_s\}$  to be bordered means exactly that  $\{\lambda_s\}$ , in a neighborhood of  $\partial N$ , is a direct product of deformations—a deformation of Liouville structure of the fiber  $F$ , and a constant (non-)deformation of the structure on  $T^*[0, 1]^i$ —with respect to the splitting in (S2).

**Remark 2.20.** If  $\{\lambda_s\}$  is a compactly supported deformation, the functions  $h_s$  can be chosen to vanish near  $\partial N$  and  $\{\lambda_s\}$  is constant (i.e.,  $s$ -independent) near this boundary.

Put another way, we abusively use compactly supported to mean “interior and compactly supported.”

**Remark 2.21.** Suppose  $S$  is compact and the family  $\{\lambda_s\}_{s \in S}$  is compactly supported. Then the compact subset  $K \subset N$  may be chosen uniformly (i.e., independent of  $s \in S$ ).

**Remark 2.22.** When  $S$  is the smooth  $n$ -simplex equipped with a collaring, we will also define what it means for  $\{\lambda_s\}_{s \in S}$  to be collared (Definition 2.40).

### 2.1.3 Various morphisms

We discuss maps between Liouville sectors. Here is the most basic class:

**Definition 2.23.** Fix two Liouville sectors  $(M, \lambda^M)$  and  $(N, \lambda^N)$ . A *sectorial embedding*, or a map of Liouville sectors, is a function  $f : M \rightarrow N$  satisfying the following:  $f$  is proper,  $f$  is a codimension-zero smooth embedding, and there exists some smooth, compactly supported function  $h : M \rightarrow \mathbb{R}$  for which  $f^* \lambda^N = \lambda^M + dh$ .

If  $f$  is further a diffeomorphism, we call  $f$  an *isomorphism* of Liouville sectors.

**Remark 2.24.** By the assumption that  $f$  is proper and codimension zero, one can equally ask for the existence of a compactly supported  $g$  on  $N$  for which  $f^*(\lambda^N + dg) = \lambda^M$  (rather than asking for the compactly supported function to be defined on  $M$ ).

**Definition 2.25.** We call a sectorial embedding  $f$  *strict* if  $f^* \lambda^N = \lambda^M$ . We further call  $f$  a *strict isomorphism* if  $f$  is a diffeomorphism.

**Remark 2.26.** In [16], what we here call a strict sectorial embedding is called a strict proper inclusion.

**Remark 2.27.** Note we allow the sectorial boundary of  $M$  to intersect the sectorial boundary of  $N$ . (Compare with Convention 3.1 of [9].) If  $M$  is contained in the interior of  $N$ , then the complement  $N \setminus i(M)$  is also a Liouville sector, with a strict sectorial embedding into  $N$ ; if  $N$  is further a Weinstein sector, then so is  $N \setminus i(M)$ .

**Remark 2.28.** If  $i$  and  $i'$  are strict sectorial embeddings, so is the product  $i \times i'$ . In contrast, if  $i$  and  $i'$  are sectorial embeddings, then the product may not be (due to the compact support condition).

**Remark 2.29.** Let  $f : M \rightarrow N$  be a sectorial embedding. Outside a compact region,  $f$  strictly respects  $\lambda$ . Because Liouville sectors may, outside a compact set, be modeled as symplectizations of a contact manifold-with-corners (Remark 2.3), we can contemplate the induced map  $\mathbb{R}_t \times \partial_\infty M \rightarrow \mathbb{R}_t \times \partial_\infty N$ , where both domain and codomain have Liouville form given by  $\lambda \cong e^t \lambda|_{\partial_\infty}$ . A standard computation shows that any such map must be of the form  $(t, x) \mapsto (t + \tau(x), y(x))$  where  $y$  is a contact embedding of  $\partial_\infty M$  into  $\partial_\infty N$  for which the contact form is scaled by  $y^*(\lambda^N) = e^{-\tau(x)} \lambda^M$  on  $\partial_\infty M$ . Note that both  $\tau : \partial_\infty M \rightarrow \mathbb{R}$  and  $y$  are independent of  $t$ .

**Definition 2.30.** Let  $f : M \rightarrow N$  be a sectorial embedding. We call  $f$  an *equivalence*, or *sectorial equivalence*, or an *equivalence of Liouville sectors* if there exists a sectorial embedding  $g : N \rightarrow M$  so that  $g \circ f$  and  $f \circ g$  are smoothly isotopic, through sectorial embeddings, to the identity functions of  $M$  and  $N$ , respectively.

**Example 2.31.** Fix two smooth manifolds  $R$  and  $S$ , possibly with corners. Then  $T^*R$  and  $T^*S$ , equipped with the canonical Liouville form  $pdq$ , are Liouville sectors.

Let  $f : S \rightarrow R$  and  $g : R \rightarrow S$  be smooth maps so that  $g \circ f$  is smoothly isotopic to  $\text{id}_S$  and  $f \circ g$  is smoothly isotopic to  $\text{id}_R$ . (An example: Take  $S = D^n$  and  $R = [0, 1]^n$ . We can choose  $f$  and  $g$  to be smooth embeddings given by a sufficient scaling  $x \mapsto tx$  for  $t$  a positive real number.)

Then  $T^*f$  and  $T^*g$  are both sectorial equivalences.

If  $M$  and  $N$  are Liouville sectors, let  $M \otimes T^*R$  and  $N \otimes T^*S$  denote the direct product of sectors (Definition 2.9). If  $\phi : M \rightarrow N$  is a strict isomorphism of Liouville sectors—meaning  $\phi$  is a diffeomorphism and  $\phi^*\lambda^N = \lambda^M$ —then the map

$$\phi \times T^*f : M \otimes T^*R \rightarrow N \otimes T^*S$$

is an equivalence of Liouville sectors.

The above morphisms are specified completely by a single mapping. We now introduce a class of morphisms specified by a mapping, together with a family of Liouville structures in the codomain.

**Definition 2.32.** Let  $(M, \lambda^M)$  and  $(N, \lambda^N)$  be Liouville sectors. Consider a pair  $(f, \{\lambda_s^N\}_{s \in [-1, 1]})$  where  $f$  is a smooth, proper embedding (with a priori no compatibility with  $\lambda_s^N$ ) and  $\{\lambda_s^N\}$  is a 1-parameter exact family of Liouville structures of  $N$  with  $\lambda_1^N = \lambda^N$ .

We say that  $(f, \{\lambda_s^N\})$  is a *bordered, interior, or compactly supported deformation embedding* if

1.  $\{\lambda_s^N\}$  is a bordered, interior, or compactly supported family (respectively),
2. The family  $\lambda_s^N$  is constant (i.e.,  $s$ -independent) near  $s = -1$  and near  $s = 1$ , and
3.  $f^*\lambda_{-1}^N = \lambda^M$ .

**Remark 2.33.** Note that composition of deformation morphisms is cumbersome to define on the nose, if not impossible. For example, given two deformation embeddings  $f_{01} : M_0 \rightarrow M_1$ ,  $f_{12} : M_1 \rightarrow M_2$ , one would have to make a non-canonical choice of extending  $\{(f_{12})_*\lambda_s^{M_1}\}_s$  and implement a Moore-path-space-like concatenation of  $s$ -intervals to define a composition and define a topologically enriched category of Liouville sectors with morphisms given by deformation morphisms.

This at least partially motivates the reason we use  $\infty$ -categorical models in this work. The higher simplices of an  $\infty$ -category *are* the data of all potentially non-canonical choices—so we need not make any choices—and (by definition of  $\infty$ -category) we need not define any compositions that depend on such choices.

## 2.2 The movie construction

The following generalizes a construction of [8, Section 3.3].

**Construction 2.34** (The movie construction). Let  $(M, \lambda)$  be a Liouville sector and let  $S$  be a smooth manifold, possibly with corners. Fix an  $S$ -parametrized exact family of Liouville structures encoded by a smooth function  $h : M \times S \rightarrow \mathbb{R}$  (Definition 2.12). Then the *movie construction* associated to  $h$  is the pair

$$(M \times T^*S, \lambda + \mathbf{pdq} + dh) \tag{2.4}$$

where  $\mathbf{pdq}$  is the standard Liouville form on  $T^*S$ , and we out of sloth identify  $\mathbf{pdq}$  and  $\lambda$  with their pullbacks to  $M \times T^*S$ .

Observe that the movie construction is an exact symplectic manifold. We will see in Proposition 2.42 that the movie construction is always a Liouville sector provided that the deformation encoded by  $h$  is bordered and collared (Definition 2.37) along the corners of  $S$ .

**Remark 2.35.** Setting  $\lambda_s = \lambda + d_M h_s$  as in (2.3), we observe

$$\lambda + \mathbf{pdq} + dh = \lambda_s + \mathbf{pdq} + d_S h.$$

where the last term denotes we only take the deRham derivative of  $h$  in the  $S$  directions. Both sides of the above equation are written informally; the most rigorous expression for this 1-form is

$$\pi_M^* \lambda + \pi_{T^*S}^* \mathbf{pdq} + \pi_{M \times S}^* dh$$

where  $\pi_M$ ,  $\pi_{T^*S}$ , and  $\pi_{M \times S}$  are the projections from  $M \times T^*S$  to  $M$ ,  $T^*S$ , and  $M \times S$  respectively.

**Remark 2.36.** The presentation of the form as  $\lambda + \mathbf{pdq} + dh$  breaks symmetry by apotheosizing a particular Liouville structure  $\lambda$  on  $M$ . Later, we will often take  $S = \Delta^n$  to be a simplex with ordered vertices, and justify the symmetry-breaking by further demanding that  $h_s$  vanishes (as a function on  $M$ ) in a neighborhood of the terminal vertex of  $\Delta^n$ .

The main result we will use about the movie construction is that the construction itself is a Liouville sector (Proposition 2.42), provided that  $h$  is collared appropriately. We say what we mean by an appropriate collar now.

**Definition 2.37** (Collaring). Let  $S$  be a smooth manifold with corners. A *collaring* of  $S$  is the data, for every codimension  $i \geq 1$  boundary stratum  $\partial^i S \subset S$ , of a smooth, codimension-respecting embedding

$$\phi_i : \partial^i S \times [-1, \epsilon_i]^i \rightarrow S, \quad (x, \vec{u}) \mapsto \phi_i(x, \vec{u})$$

for which  $\phi_i(x, 0) = x$ , and for which there exists a  $\delta = \delta(i, j)$  for which

$$\phi_i(\phi_j(x, \vec{u}'), \vec{u}'') = \phi_j(x, \vec{u})$$

whenever  $\phi_j(x, \vec{u}') \in \partial^j S$  and  $|\vec{u}'|, |\vec{u}''| < 1 + \delta$ . Here,  $\vec{u} \in [-1, \epsilon_j]^j$  is a vector obtained by replacing some of the null coordinates of  $\vec{u}'$  by the coordinates of  $\vec{u}''$ , up to some re-ordering of coordinates.

**Remark 2.38.** In short, a collaring is a trivialization of tubular neighborhoods of each stratum using cubical coordinates. The compatibility condition states that these trivializations respect the passage to higher-codimension strata. We will later choose collarings for simplices  $S = \Delta^k$ , and so that the collarings are compatible with respect to natural simplicial face maps. (See Definition 4.3.)

**Remark 2.39.** It would be most natural to not demand that collarings always extend beyond 0. However, here (and everywhere else in this work) we demand that collaring coordinates extend beyond the interval  $[-1, 0]$  to guarantee a *standard* embedding of  $[-1, 0]$  into these collaring neighborhoods. The reasoning behind this is ultimately a coherence/categorical issue: These collars allow us to make sure that our eventual  $\infty$ -categories  $\text{Liou}_\Delta$  and  $\text{Liou}_\Delta^{\text{Def}}$  are comprised of honestly commuting diagrams of collared objects, rather than of data that allows for shrinking isotopies and reparametrizations.

**Definition 2.40** (Collared family). Fix a collaring on  $S$ . We say a smooth function  $h : M \times S \rightarrow \mathbb{R}$  is *collared* (with respect to the collaring on  $S$ ) if, for every codimension  $i \geq 1$  stratum of  $S$ , there is some neighborhood of  $\partial^i S$  so that  $h$  factors through the projection

$$M \times \text{Nbhd}(\partial^i S) \rightarrow M \times \partial^i S.$$

In other words,  $h$  is collared if it is independent of the “normal directions” to  $\partial^i S$  specified by the collaring  $\{\phi_i\}$ .

An  $S$ -parametrized exact family of Liouville structures encoded by a smooth function  $h : M \times S \rightarrow \mathbb{R}$  (Definition 2.12) will also be called collared if  $h$  is collared.

More generally, any structure on  $S$ —such as a function on  $S$ , a family over  $S$ , et cetera—is called collared if it is constant along the normal directions determined by the collaring.

**Example 2.41.** Let  $S = \Delta^I$  be an  $I$ -simplex with a collaring. If  $h : M \times S \rightarrow \mathbb{R}$  is collared, then  $h$  is independent of the  $\Delta^I$ -coordinate near every vertex of  $\Delta^I$ . More precisely, given a vertex  $v$ , there exists a neighborhood  $\text{Nbhd}(v) \subset \Delta^I$  so that  $h|_{M \times \text{Nbhd}(v)}$  factors as

$$M \times \text{Nbhd}(v) \rightarrow M \xrightarrow{h_v} \mathbb{R}$$

for some smooth function  $h_v$ .

**Proposition 2.42.** Let  $(M, \lambda)$  be a Liouville sector and  $S$  a compact manifold-with-corners equipped with a collaring. Fix a bordered, collared exact deformation encoded by a smooth function  $h : M \times S \rightarrow \mathbb{R}$  (Definitions 2.12 and 2.40). Then the movie construction of  $M$  (Construction 2.34) is a Liouville sector.

**Definition 2.43.** When the deformation is bordered (Definition 2.18) and collared (Definition 2.40), we will call the Liouville sector (2.4) an  $S$ -*movie*, or simply a *movie* when  $S$  is clear or irrelevant. When the deformation is further compactly supported, we will call the movie compactly supported as well.

(We will use this terminology most often when  $S$  is a simplex—i.e., we will often consider  $\Delta^I$ -movies.)

*Proof of Proposition 2.42.* Because  $(M, \lambda)$  is a sector, it comes with the data (2.1) of projections  $\pi_x$  near its corner strata. Further, the collaring of  $S$  (Definition 2.37) induces such projections near the corner strata of  $T^*S$ . The products of these define the projections for  $M \times T^*S$ . We must now ensure that these projections, together with the 1-form (2.4), satisfy the properties (S1), (S2), and (S3) from Definition 2.1. We proceed in reverse order.

(S3) is satisfied, as the projection maps are products of maps satisfying (S3).

(S2) is satisfied because the function  $h$  is collared (so that  $d_S h \equiv 0$  near  $\partial T^*S$ , see Definition 2.40), because  $\mathbf{pdq}$  satisfies (S2), and because  $\{\lambda_s\}$  is a bordered deformation (Definition 2.18).

To verify (S1), it suffices to exhibit a compact subset  $A \subset M \times T^*S$  outside of which every element escapes to infinity. For this, because  $S$  is compact, we may assume there is a single compact  $K \subset M$  outside which every  $\lambda_s$ -trajectory escapes to infinity. ( $K$  is independent of  $s$ . See Remark 2.16.)

We define  $A$  to be the compact set of those  $(x, s, \sigma)$  for which  $x \in K$  and  $|\sigma| \leq |d_S h(x, s)|$ .

If we have  $(x, s, \sigma) \in (M \times T^*S) \setminus A$  for which  $x \notin K$ , then by assumption on  $K$ , we know that the Liouville flow of  $(x, s, \sigma)$  (which is given by the flow of  $\lambda_s$  in the  $M$  component) will force the  $x$  component to flow toward infinity, hence force  $(x, s, \sigma)$  to flow toward infinity. On the other hand,

if  $x \in K$ , we know by definition of  $A$  that  $|\sigma| > |d_S(h(x, s))|$ . Noting that the Liouville vector field on  $M \times T^*S$  is given, in the  $T^*S$  component, in local coordinates as

$$\sum \left( p_i + \frac{\partial h}{\partial q_i}(x, s) \right) \partial_{p_i}$$

the definition of  $A$  shows that the standard Liouville flow given by  $\mathbf{p}\partial_{\mathbf{p}}$  dominates. In particular, we see that the  $\sigma$  coordinate of  $(x, s, \sigma)$  will tend to infinity as we flow  $(x, s, \sigma)$  along the Liouville flow of the movie. This proves (S1).  $\square$

**Remark 2.44.** The movie of any collared, bordered deformation is Liouville. However, unless the deformation takes place through Weinstein deformations, the resulting movie is not Weinstein. Nor, as far as we know, does the movie admit any deformation of Liouville structures to be a Weinstein sector.

### 2.3 The movie construction for maps

We have seen that a deformation of Liouville structures may be encoded into a single sector by the movie construction. Likewise, suppose we now have an  $S$ -parametrized collection  $\{y_s : M \rightarrow N\}_{s \in S}$  of (not necessarily strict) sectorial embeddings. We will see here that the data of  $\{y_s\}$  may be converted into the data of a single map of movies, and in fact, a single *strict* map of movies.

We leave the proof of the following as an exercise for the reader. We remind the reader that  $M \otimes T^*S$  is our notation for a sector with the product Liouville structure, while  $M \times T^*S$  is direct product of manifolds with a Liouville structure that may not necessarily be the product Liouville structure. (See Definition 2.9.)

**Proposition 2.45.** Let  $S$  be a compact manifold-with-corners equipped with a collaring. Let  $M \times T^*S$  and  $N \times T^*S$  be movies (associated to bordered and collared families of Liouville structures  $\{\lambda_s^M\}_{s \in S}$ ,  $\{\lambda_s^N\}_{s \in S}$ ; see Definition 2.43). Assume we are given a *strict* sectorial embedding  $\tilde{y} : M \times T^*S \rightarrow N \times T^*S$  for which there exists a function  $y : M \times S \rightarrow N$ ,  $(x, s) \mapsto y_s(x)$ , fitting into the following commutative diagram:

$$\begin{array}{ccc} M \times T^*S & \xrightarrow{\tilde{y}} & N \times T^*S \\ \downarrow & & \downarrow \\ M \times S & \xrightarrow{(x, s) \mapsto (y_s(x), s)} & N \times S. \end{array} \quad (2.5)$$

Then for each  $s \in S$ ,  $y_s$  is a strict sectorial embedding for the Liouville structures  $\lambda_s^M$  and  $\lambda_s^N$ .

Conversely, suppose we have a smooth, collared, continuous family  $\{y_s : M \rightarrow N\}_{s \in S}$  of not-necessarily-strict sectorial embeddings (where for each  $s \in S$ ,  $y_s$  is a sectorial embedding from  $(M, \lambda_s^M)$  to  $(N, \lambda_s^N)$ ). Then there exists (i) a compactly supported function  $g : N \times S \rightarrow \mathbb{R}$  and (ii) a smooth map  $\tilde{y} : M \times T^*S \rightarrow N \times T^*S$  so that (a)  $\tilde{y}$  makes (2.5) commute, and (b)  $\tilde{y}$  is a *strict* sectorial embedding when the domain is given the movie structure associated to  $\{\lambda_s^M\}$ , and when the codomain is given the movie sectorial structure associated to  $\{\lambda_s^N + d(g_s)\}_{s \in S}$ .

**Definition 2.46.** Fix two movies  $M \times T^*S$  and  $N \times T^*S$ . A map  $\tilde{y} : M \times T^*S \rightarrow N \times T^*S$  is called a *movie of embeddings* if

1.  $\tilde{y}$  is a strict sectorial embedding, and
2. There exists a smooth map  $y : M \times S \rightarrow N$  rendering the diagram 2.5 commutative. (By Proposition 2.45, then each  $y_s$  is necessarily a sectorial embedding.)

We will say that  $\tilde{y}$  is the movie of the family of sectorial embeddings  $\{y_s : M \rightarrow N\}_{s \in S}$ .

**Remark 2.47.** Suppose we are given a movie of embeddings  $\tilde{y} : M \times T^*S \rightarrow N \times T^*S$  where  $S$  is collared. Suppose we are further given that the movies of  $M$  and  $N$  are both movies of compactly supported deformations (Definition 2.43). Then the family  $\{y_s : M \rightarrow N\}$ —guaranteed to exist by Proposition 2.45 and obviously uniquely determined by  $\tilde{y}$ —determines a smooth and continuous map from  $S$  to the space  $\text{Emb}_{\text{Liou}}(M, N)$  of sectorial embeddings from  $M$  to  $N$  (Definitions 2.57 and 2.61). The space  $\text{Emb}_{\text{Liou}}(M, N)$  is defined unambiguously because for any  $s, s' \in S$  the Liouville structures  $\lambda_s^M, \lambda_{s'}^M$  are related by  $d$  of a compactly supported function. (In particular, the identity function is a Liouville isomorphism from  $(M, \lambda_s^M)$  to  $(M, \lambda_{s'}^M)$ .) Likewise for  $N$ .

If further  $\tilde{y}$  is assumed collared, then  $\{y_s\}$  defines a smooth and collared and continuous map from  $S$ .

## 2.4 Shrinking near the boundary, rounding corners

**Definition 2.48.** Let  $M$  be any Liouville sector. A *shrinking isotopy* of  $M$  is a smooth family  $\{\sigma_s : M \rightarrow M\}_{s \in [0,1]}$  of strict sectorial embeddings such that  $\sigma_0 = \text{id}_M$  and  $\sigma_s(\partial M) \cap \partial M = \emptyset$  for  $s > 0$ .

**Lemma 2.49.** Every Liouville sector admits a shrinking isotopy.

*Proof.* Fix a Liouville sector  $M$ . After changing the projections  $\pi_x$  near infinity, we can arrange that  $\pi_x$  is Liouville equivariant up to permutation. In particular, there is some  $\varepsilon > 0$  so that the  $\pi_x$  assemble to a collection of surjections

$$\pi^i : \text{Nbhd}(\partial^i X) \rightarrow T^*[0, \varepsilon]^i / S_i$$

with  $S_i$  the symmetric group acting by permutation. This decomposes  $M$  into pieces indexed by maximal nearby codimension. On each piece, the desired isotopy is induced by applying an embedding  $f_t : [0, \varepsilon] \rightarrow [0, \varepsilon]$  to each component which is identity near  $\varepsilon$  and sends 0 to  $\frac{t}{2\varepsilon}$ .  $\square$

**Corollary 2.50.** Every Liouville sector  $X$  is equivalent to a Liouville sector  $X^{rc}$  with smooth boundary, in the sense that there are maps  $X \rightarrow X^{rc}$  and  $X^{rc} \rightarrow X$  whose compositions are isotopic to identity.

Here, *rc* stands for *rounded corners*.

*Proof.* Take a cutoff function  $\kappa : [0, \varepsilon] \rightarrow [0, 1]$  which vanishes at 0 and equals 1 near  $\varepsilon$ , and has strictly positive derivative on  $\kappa^{-1}[0, 1)$ . Define

$$\kappa^i : [0, \varepsilon]^i \rightarrow [0, 1]$$

to be the product of  $\kappa$  on each component.  $\kappa^i$  is a submersion over  $(0, 1)$ , and we can split it by declaring  $\nabla \kappa^i$  to be horizontal, where  $\nabla$  is the gradient with respect to the standard Euclidean metric. This splitting allows us to define the cotangent map  $T^* \kappa^i$ , which is a symplectic submersion over  $T^*(0, 1)$ .

Since  $\kappa^i$  is permutation-invariant, we can consider the composition

$$\pi^{rc}|_{\text{Nbhd}^\circ(\partial^i X)} = T^* \kappa^i \circ \pi^i,$$

where  $\text{Nbhd}^\circ$  refers to a punctured neighborhood and  $\pi^i$  is as in the proof of Lemma 2.49. The local  $\pi^{rc}$  assemble to a global symplectic submersion  $\text{Nbhd}^\circ(\partial X) \rightarrow T^*(0, 1)$  with split Liouville form.

We can now define the rounded-corner sector

$$X^{rc} := X^\circ \setminus (\pi^{rc})^{-1}T^*(0, \tfrac{1}{2}).$$

□

In fact, we will often be more interested in the projection  $\pi^{rc}$  than the rounded sector  $X^{rc}$ , so let us make the following definition.

**Definition 2.51.** For a Liouville sector  $X$ , a *corner-rounding projection* is a map  $\pi^{rc}: \text{Nbhd}^\circ(\partial X) \rightarrow T^*(0, 1)$  such that

(i) The composition

$$\text{Nbhd}^\circ(\partial X) \xrightarrow{\pi^{rc}} T^*(0, 1) \rightarrow (0, 1)$$

extends to a smooth map

$$\text{Nbhd}(\partial X) \rightarrow [0, 1]$$

sending  $\partial X$  to 0.

(ii)  $\pi^{rc}$  is a symplectic fibration over  $T^*(0, 1)$  which splits the Liouville form as

$$\lambda_X = \lambda_F + pdq.$$

When working in a neighborhood of  $\partial X$ , we will sometimes abuse notation and write  $(\pi^{rc})^{-1}T^*[0, a)$  to mean  $\partial X \cup (\pi^{rc})^{-1}T^*(0, a)$ .

**Corollary 2.52.** Every Liouville sector admits a corner-rounding projection. □

## 2.5 Isotopy extension

Let  $M$  and  $N$  be smooth manifolds, and suppose one is given a smooth isotopy  $\{f_t : M \rightarrow N\}_{t \in [0, 1]}$ . The classical isotopy extension theorem guarantees conditions under which  $f_t$  may be extended to an ambient isotopy  $\{g_t : N \rightarrow N\}$  so that  $g_t \circ f_0 = f_t$ . This is also true in families—i.e., when there is an  $(s, t)$ -dependence for  $s$  a smooth parameter.

Here we prove the Liouville analogue when the  $f_{s,t}$  are (codimension zero) sectorial embeddings.

**Remark 2.53.** Even classically, one faces issues when  $M$  and  $N$  are non-compact, or have boundary. A typical example is to consider  $M = [-1, 1]$  and  $N = [-2, 2]$ , and let  $f_t$  be an isotopy that expands  $M$  to become all of  $N$ —e.g.,  $f_t(x) = x + tx$ . Clearly such an isotopy cannot extend to one of  $N$ . However, after shrinking  $N$  in a  $t$ -dependent manner (and hence changing the original isotopy of  $M$ ), one can indeed find an isotopy extension. This is the reason for the shrinking in Proposition 2.54 below.

**Proposition 2.54** (Isotopy extension for sectorial embeddings). Fix

- a smooth manifold  $S$  (possibly with corners),
- $f : M \rightarrow N$  a sectorial embedding,
- a smooth family of sectorial embeddings  $\{\widetilde{j_{s,0}} : N \rightarrow N\}_{s \in S}$ , and
- a smooth family of sectorial embeddings  $\{j_{s,t} : M \rightarrow N\}_{(s,t) \in S \times [0, 1]}$ , for which  $j_{s,0} = \widetilde{j_{s,0}} \circ f$ .

Then the following are true:

- (i) For any smoothly- $t$ -dependent family of shrinking isotopies  $\text{shrink}_t : N \rightarrow N$  with  $\text{shrink}_0 = \text{id}_N$ , there exists a smooth family of sectorial embeddings

$$\{\widetilde{j_{s,t}} : N \rightarrow N\}_{(s,t) \in S \times [0,1]}$$

so that  $\widetilde{j_{s,t}} \circ f = \text{shrink}_t \circ j_{s,t}$ .

- (ii) Suppose there is an  $(s, t)$ -dependent  $\text{Nbhd}(\partial N)_{s,t}$  such that

$$\forall s, t \quad \text{Nbhd}(\partial N)_{s,t} \cap j_{s,t}(M) = \emptyset. \quad (2.6)$$

Then one can choose  $\text{shrink}_t = \text{id}_N$  for all  $t$ . In particular, there exists a smooth family of sectorial embeddings

$$\{\widetilde{j_{s,t}} : N \rightarrow N\}_{(s,t) \in S \times [0,1]}$$

so that  $\widetilde{j_{s,t}} \circ f = j_{s,t}$ .

In words, assuming (2.6), any isotopy of a Liouville subsector  $M$  extends to an isotopy of the parent sector  $N$ , and smoothly so in  $S$ -families.

*Proof.* Consider the  $(s, t)$ -dependent vector fields  $X_{s,t} := \frac{\partial j_{s,t}}{\partial t}$  defined on the images  $j_{s,t}(M)$ . Because each  $j_{s,t}$  is a sectorial embedding, we know there exist compactly supported smooth functions  $h_{s,t} : M \rightarrow \mathbb{R}$  for which  $j_{s,t}^* \lambda^N = \lambda^M + dh_{s,t}$ . Along the images  $j_{s,t}(M) \subset N$ , the Lie derivative of  $\lambda^N$  along  $X_{s,t}$  is given by

$$\mathcal{L}_{X_{s,t}} \lambda^N = \lim_{r \rightarrow 0} \frac{j_{s,t+r}^* \lambda^N - \lambda^N}{r} = \lim_{r \rightarrow 0} \frac{dh_{s,t+r}}{r} = \frac{\partial(dh_{s,t})}{\partial t} = d\left(\frac{\partial h_{s,t}}{\partial t}\right).$$

While Cartan's magic formula gives

$$\mathcal{L}_{X_{s,t}} \lambda^N = d(\iota_{X_{s,t}}(\lambda^N)) + \iota_{X_{s,t}}(d\lambda^N).$$

Thus,

$$\omega(-, X_{s,t}) = d\lambda^N(-, X_{s,t}) = d\left(\lambda^N(X_{s,t}) - \frac{\partial h_{s,t}}{\partial t}\right)$$

In other words, for every  $(s, t)$ , we see that  $X_{s,t}$  is a Hamiltonian vector field with associated Hamiltonian

$$H_{s,t} = \lambda^N(X_{s,t}) - \frac{\partial h_{s,t}}{\partial t}.$$

We can extend each  $H_{s,t}$  to a smooth function  $\widetilde{H_{s,t}} : N \rightarrow \mathbb{R}$  which satisfies  $\widetilde{H_{s,t}} = d\widetilde{H_{s,t}}(Z^N)$  outside of an  $(s, t)$ -dependent compact subset, and we can choose such extensions smoothly with respect to  $(s, t)$ . (Here,  $Z^N$  is the Liouville vector field on  $N$ ; the equation guarantees that the flow of  $X_{\widetilde{H_{s,t}}}$  generates sectorial embeddings.) We thus obtain Hamiltonian vector fields  $X_{\widetilde{H_{s,t}}}$  on  $N$ . If these vector fields were complete on  $N$ , their  $t$ -dependent flows (pre-composed with  $\widetilde{j_{s,0}}$ ) would extend the isotopies  $j_{s,t}$ , as desired.

If we assume (2.6), we can obviously choose the extensions  $\widetilde{H_{s,t}}$  to be supported away from  $\partial N$ . (Note that the support of  $\widetilde{H_{s,t}}$  does not even need to be uniform in the  $(s, t)$  variables.) Thus the associated Hamiltonian flow is complete, and (ii) is proven.



To prove the more general setting in (i), choose a shrinking isotopy  $\text{shrink}_t : N \rightarrow N$  (Section 2.4). So  $\text{shrink}_0 = \text{id}_N$ , and for every  $t > 0$ ,  $\text{shrink}_t$  is a proper embedding with  $\text{shrink}_t(N) \cap \partial N = \emptyset$ . Let  $X_{s,t}$  be the vector field associated to the flow of  $\text{shrink}_t \circ j_{s,t}$  (rather than  $j_{s,t}$ ); as shown above, this is a Hamiltonian vector field (because  $\text{shrink}_t \circ j_{s,t}$  is a sectorial embedding for every  $t$ ), so let  $G_{s,t}$  be the associated Hamiltonian. Then by design, we can choose an extension  $\widetilde{G}_{s,t}$  whose the  $t$ -dependent Hamiltonian flow is complete and extends  $\text{shrink}_t \circ j_{s,t}$ , as desired.  $\square$

In fact, the exact same proof method yields the following:

**Proposition 2.55.** Fix

- a smooth manifold  $S$  (possibly with corners),
- $f : M \rightarrow \widetilde{M}$  a sectorial embedding,
- a smooth family of sectorial embeddings  $\{\widetilde{j}_{s,0} : \widetilde{M} \rightarrow N\}_{s \in S}$ , and
- a smooth family of sectorial embeddings  $\{j_{s,t} : M \rightarrow N\}_{(s,t) \in S \times [0,1]}$ , for which  $j_{s,0} = \widetilde{j}_{s,0} \circ f$ .

Then the following are true:

- (i) There exists a smoothly- $t$ -dependent family of shrinking isotopies  $\text{shrink}_t : N \rightarrow N$  with  $\text{shrink}_0 = \text{id}_N$ , and a smooth family of sectorial embeddings

$$\{\widetilde{j}_{s,t} : \widetilde{M} \rightarrow N\}_{(s,t) \in S \times [0,1]}$$

so that  $\widetilde{j}_{s,t} \circ f = \text{shrink}_t \circ j_{s,t}$ .

- (ii) Suppose there is an  $(s,t)$ -dependent  $\text{Nbhd}(\partial N)_{s,t}$  such that

$$\forall s, t \quad \text{Nbhd}(\partial N)_{s,t} \cap j_{s,t}(M) = \emptyset.$$

Then one can choose  $\text{shrink}_t = \text{id}_N$  for all  $t$ . In particular, there exists a smooth family of sectorial embeddings

$$\{\widetilde{j}_{s,t} : \widetilde{M} \rightarrow N\}_{(s,t) \in S \times [0,1]}$$

so that  $\widetilde{j}_{s,t} \circ f = j_{s,t}$ .

In words, assuming (2.6), any isotopy of a Liouville subsector  $M$  extends to an isotopy of the parent sector  $\widetilde{M}$ , and smoothly so in  $S$ -families.

**Remark 2.56.** At the heart of isotopy extension is the ability to choose Hamiltonians  $\widetilde{H}_{s,t}$ . This flexibility is quite useful. For example, suppose that the original isotopies are collared in some way in the  $(s,t)$  variable—say, so that the isotopies are  $s$ -independent along some open subset of  $S$ , or  $t$ -independent in some neighborhood of 0 or 1 in  $[0,1]$ . Then we may obviously choose  $\widetilde{H}_{s,t}$  with the same collaring properties, so that the chosen isotopy extensions  $\widetilde{j}_{s,t}$  inherit the collaring properties of  $j_{s,t}$ . (For  $t$ -independence, one would have to modify the  $t$ -dependence of the shrinking isotopies  $\text{shrink}_t$  accordingly; this is straightforward.)

## 2.6 Spaces of sectorial embeddings

**Definition 2.57** (The topology on  $\mathcal{E}mb_{\text{Liou}}(M, N)$ ). Fix two Liouville sectors  $M$  and  $N$ , along with a compact subset  $K \subset M$ . We let

$$\mathcal{E}mb_{\text{Liou}}^K(M, N)$$

denote the set of sectorial embeddings  $f : M \rightarrow N$  such that, writing  $f^*\lambda^N = \lambda^M + dh$ ,  $\text{supp } h \subset K$ . The topology is inherited from the collection of all smooth maps equipped with the weak  $C^\infty$  Whitney topology<sup>10</sup>. We topologize the collection of all sectorial embeddings using a direct limit:

$$\mathcal{E}mb_{\text{Liou}}(M, N) := \bigcup_K \mathcal{E}mb_{\text{Liou}}^K(M, N)$$

where the union runs over all compact subsets  $K \subset M$ .

**Remark 2.58.** This topology has the following consequence. Let  $S$  be a manifold. Then for every continuous map  $j : S \rightarrow \mathcal{E}mb_{\text{Liou}}(M, N)$ , we are guaranteed the following analogue of Property (T): For every  $s_0 \in S$ , there exists a neighborhood  $U \subset S$  containing  $s_0$ , and a compact subset  $K \subset M$ , so that whenever  $s \in U$ , we know that  $j_s^*\lambda^N - \lambda^M$  is  $d$  of some function with support in  $K$ . Note that  $K$  is uniform for all  $s \in U$ .

In particular, when  $S$  is compact, there is a single compact  $K$ —i.e., one may take  $U = S$ .

**Notation 2.59** ( $\text{Emb}_{\text{Liou}}(M, N)$ ). We let

$$\text{Emb}_{\text{Liou}}(M, N) := \text{Sing}(\mathcal{E}mb_{\text{Liou}}(M, N))$$

denote the singular complex. Explicitly,  $\text{Emb}_{\text{Liou}}(M, N)$  is the simplicial set for which a  $k$ -simplex is a continuous map  $\Delta^k \rightarrow \mathcal{E}mb_{\text{Liou}}(M, N)$ .

We emphasize the font difference between  $\mathcal{E}mb$  and  $\text{Emb}$ .

**Remark 2.60.** Writing spaces of maps between non-compact manifolds as a colimit over subspaces of maps behaving well on compact objects is a standard procedure in both smooth geometry and in symplectic geometry. See for example Remark 10.5 of [22].

Let us now say what we mean by a smooth map into a mapping space.

**Definition 2.61** (Smooth map into a mapping space). Let  $S, M, N$  be smooth manifolds (possibly with corners) and let  $C^\infty(M, N)$  be the set of smooth maps from  $M$  to  $N$ . Then a function  $S \rightarrow C^\infty(M, N)$  is set to be *smooth* if the induced function  $S \times M \rightarrow N$  is smooth in the usual sense.

**Remark 2.62.** Definition 2.61 is a natural definition for smoothness, and is employed in many other settings, perhaps most notably in the setting of diffeological spaces.

**Remark 2.63** (Smooth and continuous maps into  $\mathcal{E}mb_{\text{Liou}}(M, N)$ ). So fix a smooth manifold  $S$  (possibly with corners) and a function  $S \rightarrow \mathcal{E}mb_{\text{Liou}}(M, N)$ . Suppose this function is smooth in the sense of Definition 2.61, so that  $S \times M \rightarrow N$  is a smooth map. We note that smoothness does not impose the local tameness-at-infinity of Condition (T), i.e., the continuity with respect to the colimit-topology of Definition 2.57.

<sup>10</sup>We can motivate the use of the weak Whitney topology, as opposed to the strong Whitney topology, by noting that the natural map of diffeomorphism spaces  $\text{Diff}(Q, Q') \rightarrow \text{Diff}(T^*Q, T^*Q')$  is not continuous if the codomain is given the strong topology. See related discussions in [27].

For this reason, we will often talk about *smooth and continuous* maps into  $\text{Emb}_{\text{Liou}}(M, N)$  to consider smooth maps satisfying Condition (T). This is of course dissonant with the usual fact that smoothness implies continuity; perhaps a better terminology would have been *smooth and locally tame near infinity*. But this is more linguistic than mathematical. One could have built in the  $K$ -filtered behavior into our definition of smoothness to avoid having to say “smooth and continuous.”

**Notation 2.64.** Given a function  $j : S \rightarrow \text{Emb}_{\text{Liou}}(M, N)$ , we will denote by  $j_s$  the induced map

$$M \cong M \times \{s\} \hookrightarrow M \times S \rightarrow N.$$

We will also write  $j(s, x)$  to denote  $j(s)(x)$ . So for example,  $j_s(x) = j(s, x)$ .

## 2.7 Spaces of (bordered) deformation embeddings

Here, we will define various spaces of deformation embeddings between two given Liouville sectors. In Section 4.5, we will prove these spaces are all homotopy equivalent. To begin, we need a space of deformation of Liouville structures which is sensitive to Condition (T).

**Definition 2.65.** For  $N$  a Liouville sector and  $K \subset N$  a compact subset, define  $\mathbf{L}_K(N)$  to be the space of pairs  $(\lambda_N + dh, r) \in \Omega^1(N) \times C^\infty(N)$  for which  $(N, \omega = d(\lambda + dh))$  is a Liouville sector with respect to the projections  $\pi_x$  of  $N$ , and such that  $r$  is positive and of linear growth with respect to  $\lambda$  outside  $K$ . (The function  $r$  exhibits that  $\lambda$  is finite type as in (S1).)

The *space of bordered modifications* of  $N$  is

$$\mathbf{L}(N) := \left( \bigcup_K \mathbf{L}_K(N) \right) / \sim,$$

where the equivalence relation forgets the second component  $r$ .

As usual, by a *smooth* map  $S \rightarrow \mathbf{L}(N)$ , we mean a function for which the adjoint is smooth function (so that the  $S$ -parametrized family of  $(\lambda, r)$  depends smoothly on  $S$ ).

We now define two homotopy equivalent (in fact, contractible) spaces of deformations.

**Definition 2.66.** Let  $N$  be a Liouville sector. The *space of bordered Liouville deformations* of  $N$  is the subspace

$$\text{Def}(N) \subset C^\infty(\Delta^1, \mathbf{L}(N))_{\lambda_t}$$

for which  $\lambda_t$  agrees with  $\lambda^N$  in a neighborhood of the terminal vertex of  $\Delta^1$  and is  $t$ -independent in a neighborhood of the initial vertex of  $\Delta^1$ .

The *Moore path space of bordered Liouville deformations* of  $N$  is a subspace

$$\text{Def}_{\text{Moore}}(N) \subset (\mathbb{R}_{\geq 0})_T \times C^\infty((\mathbb{R})_t, \mathbf{L}(N))_{\lambda_t}$$

consisting of the following elements. Given

$$(T, (t \mapsto (\lambda_t, r_t))) \in (\mathbb{R}_{\geq 0})_T \times C^\infty((\mathbb{R})_t, \mathbf{L}(N))_{\lambda_t}$$

we demand that  $\lambda_t$  agrees with  $\lambda^N$  for  $t \leq 0$  and is  $t$ -independent for  $t \geq T$ .

We note that  $\text{Def}(N)$  and  $\text{Def}_{\text{Moore}}(N)$  both have obvious notions of smooth maps from a smooth manifold  $B$ .

**Remark 2.67.** The two spaces in Definition 2.66 weakly homotopy equivalent, and in fact contractible; but the utility of the Moore deformation space is the evident concatenation operation. Likewise, the two spaces in Definition 2.68 below are homotopy equivalent.

**Definition 2.68.** The *space of compactly supported Liouville deformations* of  $N$  is the subspace

$$\mathrm{Def}^{\mathrm{cmpct}}(N) \subset \mathrm{Def}(N)$$

for which  $\lambda_t$  is a compactly supported deformation. We likewise have the Moore path space analogue.

The *Moore path space of compactly supported Liouville deformations* of  $N$  is the subspace

$$\mathrm{Def}_{\mathrm{Moore}}^{\mathrm{cmpct}} \subset \mathrm{Def}_{\mathrm{Moore}}(N).$$

These again have obvious notions of smooth maps from a smooth manifold  $B$ .

This allows us to define models for spaces of deformation embeddings.

**Definition 2.69.** The *space of deformation embeddings* of Liouville sectors from  $M$  to  $N$  is the subspace

$$\mathrm{Emb}_{\mathrm{Liou}}^{\mathrm{Def}}(M, N) \subset \mathrm{Def}(N)_{\lambda_t^N} \times C^\infty(M, N)_f$$

such that, near the initial vertex of  $\Delta^1$ , we have that  $f^*\lambda_t^N = \lambda^M$ . Note that  $C^\infty$  is given the weak Whitney topology.

The *Moore path space of deformation embeddings* of  $N$  is the subspace

$$\mathrm{Emb}_{\mathrm{Liou}, \mathrm{Moore}}^{\mathrm{Def}}(M, N) \subset \mathrm{Def}_{\mathrm{Moore}}(N)_{\lambda_t^N} \times C^\infty(M, N)_f$$

for which  $f^*\lambda_0^N = \lambda^M$ .

Again, both spaces have natural notions of smooth maps to them.

We also have the following versions:

**Definition 2.70.** The *space of compactly supported deformation embeddings* of Liouville sectors from  $M$  to  $N$  is the subspace

$$\mathrm{Emb}_{\mathrm{Liou}}^{\mathrm{Def}, \mathrm{cmpct}}(M, N) \subset \mathrm{Emb}_{\mathrm{Liou}}^{\mathrm{Def}}(M, N)$$

such that  $\lambda_t^N$  is a compactly supported family of deformations. Likewise, the *Moore path space of compactly supported deformation embeddings* of  $N$  is the subspace

$$\mathrm{Emb}_{\mathrm{Liou}, \mathrm{Moore}}^{\mathrm{Def}, \mathrm{cmpct}}(M, N) \subset \mathrm{Emb}_{\mathrm{Liou}, \mathrm{Moore}}^{\mathrm{Def}}(M, N)$$

such that  $\lambda_t^N$  is a compactly supported family of deformations.

Again, both spaces have natural notions of smooth maps to them.

We have bombarded the reader with various notions of morphisms between Liouville sectors. Here are the three most important notions in this work:

1. Sectorial embeddings (Definition 2.23), each of which is the datum of a single map  $f$ .
2. Compactly supported deformation embeddings (Definition 2.32), which are maps  $f$  equipped with a compactly supported (and hence bordered) deformation of Liouville structures in the codomain.

3. (Bordered) deformation embeddings (Definition 2.32), which are maps  $f$  equipped with a bordered deformation of Liouville structures in the codomain.

Having defined the spaces of these morphisms, we may observe maps between them as follows:

$$\begin{array}{ccc}
 \mathrm{Emb}_{\mathrm{Liou}, \mathrm{Moore}}^{\mathcal{D}\mathrm{ef}, \mathrm{cmpct}}(M, N) & \longrightarrow & \mathrm{Emb}_{\mathrm{Liou}, \mathrm{Moore}}^{\mathcal{D}\mathrm{ef}}(M, N) \\
 \uparrow \sim & & \uparrow \sim \\
 \mathrm{Emb}_{\mathrm{Liou}}(M, N) & \xleftarrow{\sim} \mathrm{Emb}_{\mathrm{Liou}}^{\mathcal{D}\mathrm{ef}, \mathrm{cmpct}}(M, N) \longrightarrow & \mathrm{Emb}_{\mathrm{Liou}}^{\mathcal{D}\mathrm{ef}}(M, N)
 \end{array} \tag{2.7}$$

The vertical equivalences are induced by the usual equivalences between path spaces and their Moore path space models. The left-pointing equivalence is induced by noting that the space of compactly supported functions is convex, hence contractible.

We will later prove (Theorem 4.18) that the right-pointing arrows are also weak homotopy equivalences.

### 3 $\infty$ -categorical tools

Here we set up notation and introduce one new result: Proposition 3.19. It allows us to replace an  $\infty$ -category  $\mathcal{C}$  with an equivalent  $\infty$ -category  $\mathrm{Ex}_{\simeq}(\mathcal{C})$ , consisting of certain barycentric subdivision diagrams in  $\mathcal{C}$ . The expert reader will recognize that barycentric subdivision diagrams can be used to model localizations; this philosophy will allow us to construct a model of the localization  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{e}q^{\diamond})^{-1}]$  in Sections 5 and 10.

We expect many readers will not be familiar with some  $\infty$ -categorical arguments, so we include references in the footnotes.

**Notation 3.1** (Simplices). Let  $I$  be a finite, non-empty, linearly ordered set. We let  $\Delta^I$  denote the simplicial set given by the nerve of  $I$ . Concretely, a  $k$ -simplex of  $\Delta^I$  is a length  $k$  ascending chain  $i_0 \leq \dots \leq i_k$  in  $I$ . We call  $\Delta^I$  the standard (combinatorial)  $I$ -simplex.

There is likewise a topological space

$$\Delta^I := \{(x_i)_{i \in I} \in \mathbb{R}_{\geq 0}^I \mid \sum_{i \in I} x_i = 1\}. \quad (3.1)$$

While it is sometimes common to write the space by the notation  $|\Delta^I|$  (to avoid double-booking notation) we will not follow this convention, with the exception of Sections 6 and 8, where we will repeatedly use the topological spaces, not just the combinatorial simplices. When the vertical bars  $|\bullet|$  are missing, it will be clear from context whether we are referring to the combinatorial simplex, or the simplex as a topological space.

Note  $\Delta^I$  is also a smooth manifold with corners in the standard way.

Finally, we will often depart from the convention of (3.1) and allow our simplices to have coordinates as small as -1 (see Section 4.1). This is to more easily define collarings of simplices later, but does not in any meaningful way affect the underlying ideas of our constructions.

#### 3.1 Simplicial and semisimplicial sets

**Notation 3.2** ( $\Delta_{\mathrm{inj}}$ ). Recall that  $\Delta$  is the category whose objects are finite, non-empty, linearly ordered sets  $I$ . A morphism is a weakly order-preserving function.

We let  $\Delta_{\mathrm{inj}}$  denote the subcategory of  $\Delta$  with the same objects, but whose morphisms are injections (equivalently, the morphisms must be strictly order-preserving).

**Definition 3.3.** A *simplicial set* is a functor  $\Delta^{\mathrm{op}} \rightarrow \mathbf{Sets}$  to the category of sets. A map of simplicial sets is a natural transformation. A *semisimplicial set* is a functor  $(\Delta_{\mathrm{inj}})^{\mathrm{op}} \rightarrow \mathbf{Sets}$ . A map of semisimplicial sets is a natural transformation.

Fix a simplicial set  $X$ . Given an injection  $I' \rightarrow I$ , we call the induced map  $X_I \rightarrow X_{I'}$  a face map. If  $I' \rightarrow I$  is a surjection, the induced map is called a degeneracy map. Informally, a semisimplicial set is a simplicial set without the degeneracy maps.

We let  $\Lambda_k^n$  delete the  $k$ th  $n$ -horn—informally, this is a simplicial set obtained by deleting the interior, and the face opposite the  $k$ th vertex, of the  $n$ -simplex. A simplicial set  $X$  is called an  $\infty$ -category if it satisfies the weak Kan condition: For any  $0 < k < n$ , any map  $\Lambda_k^n \rightarrow X$  extends to  $\Delta^n$ . Such an extension is often called a filler.

**Remark 3.4.** The distinction between the notion of simplicial and semisimplicial sets will become prominent especially in Section 7.2. This is because, when the simplicial set is an  $\infty$ -category, degeneracy maps encode identity morphisms and the higher coherences of identity morphisms. The

point of constructing a well-behaved *semisimplicial* set is to construct an “ $\infty$ -category” that admits units, but for which we specify none.

It is often the case that one can easily define face maps, but cannot easily define degeneracy maps. This is common in many geometric settings—e.g., cobordism categories, and in settings where strict collaring conditions must be respected (as in our setting—see Definition 5.12).

We now review formal facts from the theory of  $\infty$ -categories that tell us when semisimplicial sets may be promoted to be simplicial sets, and in fact,  $\infty$ -categories.

**Definition 3.5** ([30], Definition 1.1). Given a semisimplicial set  $X$ , an edge  $e$  is *idempotent* if there is a 2-simplex of  $X$  whose three boundary edges are all given by  $e$ . The edge  $e$  is an equivalence if (a) for any horn  $\Lambda_n^n \rightarrow X$  whose last edge is  $e$ , there is a filler  $\Delta^n \rightarrow X$ , and (b) the same holds for any horn  $\Lambda_0^n \rightarrow X$  whose first edge is  $e$ .

**Theorem 3.6.** (Theorem 1.2 of [30].) Let  $X$  be a semisimplicial set satisfying the semisimplicial weak Kan condition, and for which every object admits an idempotent self-equivalence. Then  $X$  admits a simplicial structure with the same underlying face maps, whose degenerate edges are the idempotent self-equivalences, and for which  $X$  is an  $\infty$ -category.

(Theorem 2.1 of *ibid.*) Moreover, fix a semisimplicial subset  $A \subset X$  with a specified simplicial structure. Then one may instead choose a simplicial structure on  $X$  extending the degeneracy maps of  $A$ .

**Theorem 3.7** ([33]). Let  $X$  and  $Y$  be  $\infty$ -categories, and fix  $f_{\text{semi}} : X \rightarrow Y$  a *semisimplicial* map (meaning  $f$  respects face maps, but may not respect degeneracy maps). If  $f_{\text{semi}}$  sends all degenerate edges of  $X$  to equivalences in  $Y$ , then there exists a *simplicial* set map  $f : X \rightarrow Y$  such that, for any simplex  $e$  of  $X$ ,  $f_{\text{semi}}(e)$  is naturally homotopic to  $f(e)$ .

In some sense, the degeneracies of a simplicial set are the “least homotopical” ingredient in the theory of  $\infty$ -categories, as they demand particular choices and *strict* equations to be satisfied by these choices. The above two theorems allow us to bypass this issue, at the level of both individual  $\infty$ -categories and functors between them.

## 3.2 Subdivisions and Ex

We recall some standard constructions in simplicial homotopy theory [14, 11]

**Notation 3.8.** Let  $I$  be a finite, non-empty, linearly ordered set. By  $\mathcal{P}'(I)$ , we mean the collection of all non-empty subsets of  $I$ . We note that  $\mathcal{P}'(I)$  is itself a (non-linear) poset, ordered by the subset relation.

**Notation 3.9** (sd). Let  $\Delta^I$  be the standard  $I$ -simplex. We let  $\text{sd}(\Delta^I)$  denote the simplicial set given by the nerve of  $\mathcal{P}'(I)$ . We call it the *barycentric subdivision* of  $\Delta^I$ . Concretely, a 0-simplex of  $\text{sd}(\Delta^I)$  is an element of  $\mathcal{P}'(I)$ . A  $k$ -simplex is a chain  $A_0 \subset A_1 \subset \dots \subset A_k$  of elements in  $\mathcal{P}'(I)$ , where  $\subset$  need not be a proper inclusion.

We have defined  $\text{sd}$  on simplices. Following custom, we denote the left Kan extension to all simplicial sets by the same notation. Concretely, given a simplicial set  $\mathcal{C}$ , we have

$$\text{sd}(\mathcal{C}) \cong \text{colim}_{(\Delta^k \rightarrow \mathcal{C}) \in \Delta_{/e}} \text{sd}(\Delta^k).$$

Informally,  $\text{sd}(\mathcal{C})$  is the simplicial set obtained from  $\mathcal{C}$  by replacing every  $k$ -simplex of  $\mathcal{C}$  by  $\text{sd}(\Delta^k)$ .

**Definition 3.10.** We will say that an edge  $(A \subset B)$  of  $\text{sd}(\Delta^I)$  is *max-localizing* if  $\max A = \max B$ .

**Remark 3.11.** Consider the function

$$\max : \mathcal{P}'(I) \rightarrow I, \quad A \mapsto \max A.$$

(Here,  $\max A \in A \subset I$  is the maximal element of  $A$  with respect to the linear order induced from  $I$ .) Then  $\max$  is a map of posets. It has a fully faithful right adjoint

$$I \rightarrow \mathcal{P}'(I), \quad i \mapsto [\min I, i] = \{i' \in I \mid \min I \leq i' \leq i\}. \quad (3.2)$$

Thus,  $\max : \text{sd}(\Delta^I) \rightarrow \Delta^I$  is a localization of  $\infty$ -categories<sup>11</sup>. (We have abused notation and used  $\max$  to also denote the induced map of nerves.)

In particular, for any  $\infty$ -category  $\mathcal{C}$ , the map of functor  $\infty$ -categories

$$\text{Fun}(\Delta^I, \mathcal{C}) \rightarrow \text{Fun}(\text{sd}(\Delta^I), \mathcal{C})$$

is fully faithful, with essential image consisting of those functors that send max-localizing edges to equivalences in  $\mathcal{C}$ .<sup>12</sup>

**Warning 3.12.** The right adjoints (3.2) are not natural in the  $I$  variable.

**Definition 3.13** (Ex). Let  $\mathcal{C}$  be a simplicial set. We define  $\text{Ex}(\mathcal{C})$  to be the simplicial set

$$\text{Ex}(\mathcal{C}) : \Delta^{\text{op}} \rightarrow \mathbf{Sets}, \quad I \mapsto \text{hom}_{\mathbf{sSet}}(\text{sd}(\Delta^I), \mathcal{C})$$

so a  $k$ -simplex of  $\mathcal{C}$  is the data of a simplicial set map from  $\text{sd}(\Delta^k)$  to  $\mathcal{C}$ .<sup>13</sup>

**Remark 3.14.** Given any (not necessarily order-respecting) function  $f : I \rightarrow J$ , we have an induced map of posets  $\mathcal{P}'(I) \rightarrow \mathcal{P}'(J)$  by sending a subset  $A \subset I$  to  $f(A) \subset J$ . Note  $f^* : \text{Ex}(\mathcal{C})_J \rightarrow \text{Ex}(\mathcal{C})_I$  is precomposition with this induced map.

**Notation 3.15.** There is a natural transformation  $\max^* : \text{id}_{\mathbf{sSet}} \rightarrow \text{Ex}$ . Explicitly, for any simplicial set  $\mathcal{C}$ , the map

$$\max^* : \mathcal{C} \rightarrow \text{Ex}(\mathcal{C})$$

is induced by precomposition with  $\max : \mathcal{P}'(I) \rightarrow I, A \mapsto \max A$ .

### 3.3 $\text{Ex}_{\simeq}$ of $\infty$ -categories

**Notation 3.16** ( $\text{Ex}_{\simeq}$ ). Let  $\mathcal{C}$  be an  $\infty$ -category. We let  $\text{Ex}_{\simeq}(\mathcal{C}) \subset \text{Ex}(\mathcal{C})$  denote the simplicial set whose  $I$ -simplices consist only of those maps  $\text{sd}(\Delta^I) \rightarrow \mathcal{C}$  that send max-localizing edges to equivalences.

**Notation 3.17** ( $m$ ). The natural map  $\mathcal{C} \rightarrow \text{Ex}(\mathcal{C})$  from Notation 3.15 factors through  $\text{Ex}_{\simeq}(\mathcal{C})$  because the composition  $\text{sd}(\Delta^I) \xrightarrow{\max} \Delta^I \rightarrow \mathcal{C}$  sends max-localizing edges (Definition 3.10) to (degenerate) equivalences in  $\mathcal{C}$ . We will denote this map of simplicial sets by

$$m : \mathcal{C} \rightarrow \text{Ex}_{\simeq}(\mathcal{C}).$$

We note that the assignment  $\mathcal{C} \mapsto \text{Ex}_{\simeq}(\mathcal{C})$  is a functor (from the category of  $\infty$ -categories to the category of simplicial sets) and  $m$  defines a natural transformation from the identity functor.

In Proposition 3.22, we will prove that the  $\text{Ex}_{\simeq}$  functor may be enriched over simplicial sets; moreover, we will see shortly that  $\text{Ex}_{\simeq}(\mathcal{C})$  is an  $\infty$ -category if  $\mathcal{C}$  is (Proposition 3.19). Thus  $\text{Ex}_{\simeq}$  defines a functor from the  $\infty$ -category of  $\infty$ -categories to itself.

<sup>11</sup>Definition 5.2.7.2 of [17].

<sup>12</sup>Proposition 5.2.7.12 of [17].

<sup>13</sup>The  $\text{Ex}$  stands for “extension” according to Kan’s original work [14].



**Notation 3.18.** For this section only, we will use the notation  $\text{Fun}^*$  from time to time. Concretely,  $\text{Fun}(X, \mathcal{C})$  is the simplicial set of maps from  $X$  to  $\mathcal{C}$ , and the notation  $\text{Fun}^*(X, \mathcal{C})$  emphasizes the simplicial structure:  $\text{Fun}^n(X, \mathcal{C})$  is the set of maps  $X \times \Delta^n \rightarrow \mathcal{C}$ . The reason for the  $*$  symbol is that we will soon have expressions such as  $\text{Fun}^*(\Delta^\bullet, \mathcal{C})$  depending on two indices  $*$  and  $\bullet$ , and we want to distinguish their roles.

**Proposition 3.19.** Let  $\mathcal{C}$  be an  $\infty$ -category.

1.  $\text{Ex}_{\simeq}(\mathcal{C})$  is an  $\infty$ -category.
2. The map (Notation 3.17)  $m : \mathcal{C} \rightarrow \text{Ex}_{\simeq}(\mathcal{C})$  is an equivalence of  $\infty$ -categories.

*Proof.* Consider the simplicial space

$$\text{Fun}^*(\Delta^\bullet, \mathcal{C})^\sim$$

where  $*$  is the “space” index and  $\bullet$  is the “simplicial” index. If  $Y$  is an  $\infty$ -category,  $\text{Fun}^*(X, Y)$  is also an  $\infty$ -category (regardless of  $X$ ).<sup>14</sup> The notation  $\text{Fun}^*(X, Y)^\sim \subset \text{Fun}^*(X, Y)$  indicates the largest  $\infty$ -groupoid inside  $\text{Fun}^*(X, Y)$ . In short,  $\text{Fun}^*(\Delta^I, \mathcal{C})^\sim$  is the space (Kan complex) of functors from  $\Delta^I$  to  $\mathcal{C}$ . Then  $\text{Fun}^*(\Delta^\bullet, \mathcal{C})^\sim$  is a complete Segal space because  $\mathcal{C}$  is an  $\infty$ -category.<sup>15</sup>

Let us now define the simplicial space

$$\text{Fun}_{\simeq}^*(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim.$$

For any  $I$ ,  $\text{Fun}_{\simeq}^*(\text{sd}(\Delta^I), \mathcal{C})^\sim \subset \text{Fun}^*(\text{sd}(\Delta^I), \mathcal{C})^\sim$  is declared to be the full Kan complex spanned by those simplicial set maps  $\text{sd}(\Delta^I) \rightarrow \mathcal{C}$  sending max-localizing edges to equivalences in  $\mathcal{C}$ . (In other words,  $\text{Fun}_{\simeq}^*(\text{sd}(\Delta^I), \mathcal{C})^\sim$  consists exactly of those connected components containing functors that send max-localizing edges to equivalences.)

We now claim  $\text{Fun}_{\simeq}^*(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim$  is a complete Segal space. We will see that  $\text{Fun}_{\simeq}^*(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim$  is Reedy fibrant in Lemma 3.20 below. Further, by Remark 3.11, we know that  $\Delta^I$  is a localization of  $\text{sd}(\Delta^I)$  along the max-localizing edges—so the natural map  $\text{Fun}^*(\Delta^I, \mathcal{C})^\sim \rightarrow \text{Fun}_{\simeq}^*(\text{sd}(\Delta^I), \mathcal{C})^\sim$  is a homotopy equivalence of Kan complexes for every  $I$ . Because  $\text{Fun}^*(\Delta^\bullet, \mathcal{C})^\sim$  satisfies the completeness and Segal conditions, the claim is proven.

We witnessed in the previous paragraph that  $\text{Fun}^*(\Delta^\bullet, \mathcal{C})^\sim \rightarrow \text{Fun}_{\simeq}^*(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim$  is an equivalence of complete Segal spaces. On the other hand, if  $X_{\bullet,*}$  is a complete Segal space, it is known that (i)  $X_{\bullet,0}$  is an  $\infty$ -category<sup>16</sup>, and (ii) any equivalence of complete Segal spaces  $W_{\bullet,*} \rightarrow X_{\bullet,*}$  induces an equivalence of  $\infty$ -categories<sup>17</sup>  $W_{\bullet,0} \rightarrow X_{\bullet,0}$ . The proposition follows by noting the equality of simplicial sets

$$\text{Ex}(\mathcal{C})_\bullet = \text{Fun}_{\simeq}^0(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim.$$

□

We claimed the following while proving Proposition 3.19:

**Lemma 3.20.** If  $\mathcal{C}$  is an  $\infty$ -category, then  $\text{Fun}_{\simeq}^*(\text{sd}(\Delta^\bullet), \mathcal{C})^\sim$  is Reedy fibrant.

The proof of Lemma 3.20 will rely on another lemma:

<sup>14</sup>See Proposition 1.2.7.3 of [17].

<sup>15</sup>See Proposition 4.10 of [13], where  $\Gamma(X)_{m,n}$  stands for our  $\text{Fun}^n(\Delta^m, X)^\sim$ .

<sup>16</sup>Corollary 3.6 of [13].

<sup>17</sup>This follows from the fact that  $X_{\bullet,*} \mapsto X_{\bullet,0}$  is right Quillen. (See [13], where the notation  $i_1^*$  denotes this right adjoint.) Because  $i_1^*$  is right Quillen, it preserves weak equivalences between fibrant objects. On the other hand, complete Segal spaces and  $\infty$ -categories are the fibrant objects of the domain and codomain of  $i_1^*$ , respectively.

**Lemma 3.21.** Let  $X$  and  $Y$  be  $\infty$ -categories and  $f : X \rightarrow Y$  a cofibration (meaning  $f_n : X_n \rightarrow Y_n$  is an injection for all  $n \geq 0$ ). Then for any  $\infty$ -category  $\mathcal{C}$ , the induced map of Kan complexes

$$\mathrm{Fun}(Y, \mathcal{C})^\sim \rightarrow \mathrm{Fun}(X, \mathcal{C})^\sim$$

is a Kan fibration.

*Proof.* This is a special case of Lemma 3.1.3.6 of [17]. In the notation and language of loc. cit. (which we use throughout this proof), one sets  $S = \Delta^0$  and  $Z = \mathcal{C}$ . Note that because  $X$  and  $Y$  are  $\infty$ -categories, any map of simplicial sets sends the naturally marked edges of  $X$  and  $Y$  (i.e., the equivalences in  $X$  and  $Y$ ) to the naturally marked edges (the equivalences) of  $\mathcal{C}$ . Thus  $\mathrm{Map}^b(X, \mathcal{C}) = \mathrm{hom}_{\mathbf{sSet}}(X, \mathcal{C})$ , and  $\mathrm{Map}^\sharp(X, \mathcal{C})$  is the largest Kan complex inside  $\mathrm{hom}_{\mathbf{sSet}}(X, \mathcal{C}) = \mathrm{Fun}(X, \mathcal{C})$  (Remark 3.1.3.1 of ibid.).

(There are other proofs using the fact that the Joyal model structure for  $\infty$ -categories is Cartesian closed – see Proposition 4.10 of [13].)  $\square$

*Proof of Lemma 3.20.* It is straightforward to verify that Reedy fibrancy is equivalent to the following statement: For every  $n \geq 0$ , the map of simplicial sets

$$\mathrm{Fun}_\simeq^*(\mathrm{sd}(\Delta^n), \mathcal{C})^\sim \rightarrow \mathrm{Fun}_\simeq^*(\mathrm{sd}(\partial\Delta^n), \mathcal{C})^\sim \quad (3.3)$$

is a Kan fibration. We prove this now.

Note that the map  $\mathrm{sd}(\partial\Delta^n) \rightarrow \mathrm{sd}(\Delta^n)$  is a monomorphism. We also see  $\mathrm{sd}(\partial\Delta^n)$  is an  $\infty$ -category. (For example, the monomorphism identifies  $\mathrm{sd}(\partial\Delta^n)$  as the full subcategory of  $\mathrm{sd}(\Delta^n)$  obtained by excluding the terminal object  $[n]$ .)

It follows from Lemma 3.21 that the induced map on functor spaces  $\mathrm{Fun}^*(\mathrm{sd}(\Delta^n), \mathcal{C})^\sim \rightarrow \mathrm{Fun}^*(\mathrm{sd}(\partial\Delta^n), \mathcal{C})^\sim$  is a Kan fibration. Thus the map (3.3) is also a Kan fibration (as it is the restriction of a Kan fibration to certain connected components).  $\square$

**Proposition 3.22.** The functor  $\mathrm{Ex}$ , from the category of simplicial sets to itself, admits a natural promotion to a functor of  $\mathbf{sSet}$ -enriched categories.

(Concretely, this means that for any two simplicial sets  $\mathcal{C}, \mathcal{D}$  and a map  $\mathcal{C} \times \Delta^m \rightarrow \mathcal{D}$ , we can create a map of simplicial sets  $\mathrm{Ex}(\mathcal{C}) \times \Delta^m \rightarrow \mathrm{Ex}(\mathcal{D})$ , natural in the three variables  $\mathcal{C}, \mathcal{D}$ , and  $\Delta^m$ .)

Likewise, restricting to the full subcategory of  $\infty$ -categories,  $\mathrm{Ex}_\simeq$  may be promoted to a functor of simplicially enriched categories.

**Remark 3.23.** Informally: Proposition 3.22 tells us that  $\mathrm{Ex}_\simeq$  is a functor that respects natural transformations of functors and higher homotopies of functors.

*Proof.* We have the following sequence of natural (in the  $\Delta^k \in \Delta^{\mathrm{op}}$  variable) maps:

$$\begin{aligned} \mathrm{hom}_{\mathbf{sSet}}(\Delta^k, \mathrm{Ex}(\mathcal{C}) \times \Delta^m) &\cong \mathrm{hom}_{\mathbf{sSet}}(\mathrm{sd}(\Delta^k), \mathcal{C}) \times \mathrm{hom}_{\mathbf{sSet}}(\Delta^k, \Delta^m) \\ &\rightarrow \mathrm{hom}_{\mathbf{sSet}}(\mathrm{sd}(\Delta^k), \mathcal{C}) \times \mathrm{hom}_{\mathbf{sSet}}(\mathrm{sd}(\Delta^k), \Delta^m) \\ &\cong \mathrm{hom}_{\mathbf{sSet}}(\mathrm{sd}(\Delta^k), \mathcal{C} \times \Delta^m) \\ &\cong \mathrm{hom}_{\mathbf{sSet}}(\Delta^k, \mathrm{Ex}(\mathcal{C} \times \Delta^m)) \end{aligned}$$

The one non-bijection is given by pulling back along the map  $\max : \mathrm{sd}(\Delta^k) = N(\mathcal{P}'([k])) \rightarrow N([k]) = \Delta^k$ . (See Remark 3.11.) All bijections are justified by universal properties of products, or by adjunction/definition. By Yoneda's Lemma, we thus have a map of simplicial sets

$$\mathrm{Ex}(\mathcal{C}) \times \Delta^m \rightarrow \mathrm{Ex}(\mathcal{C} \times \Delta^m). \quad (3.4)$$

Because  $\text{Ex}$  is a functor, we have a map (natural in all variables) which takes any map  $H : \mathcal{C} \times \Delta^m \rightarrow \mathcal{D}$  to a map  $\text{Ex}(H) : \text{Ex}(\mathcal{C} \times \Delta^m) \rightarrow \text{Ex}(\mathcal{D})$ . Composing (3.4) with  $\text{Ex}(H)$ , we obtain a map of simplicial sets

$$\text{Ex}(\mathcal{C}) \times \Delta^m \rightarrow \text{Ex}(\mathcal{D})$$

natural in all variables.

The final claim follows by noting that all maps involved preserve simplices satisfying the max-localizing condition.  $\square$

### 3.4 Making a homotopy

Here we explicate the existence of a homotopy we use to prove our localization theorem (Theorem C). Throughout, we will utilize standard notation and results from the theory of marked simplicial sets. We refer the reader to Section 3.1 of [17] for details, where the only relevance for us (in the notation of *ibid.*) is the setting where the base simplicial set  $S = \Delta^0$  is a point.

**Remark 3.24** (Marked simplicial sets model localizations). Let us onramp the unfamiliar reader. When  $\mathcal{D}$  is an  $\infty$ -category, a marked simplicial set  $(\mathcal{D}, \mathcal{E})$  is input data for localizing  $\mathcal{D}$  along a collection of edges  $\mathcal{E}$ . Indeed, in the model category of marked simplicial sets, a fibrant replacement for  $(\mathcal{D}, \mathcal{E})$  produces a localization of  $\mathcal{D}$  along  $\mathcal{E}$ .

This last claim follows as a combination of [17, Proposition 3.1.4.1] which shows that any fibrant replacement is an  $\infty$ -category marked by its equivalences, of [17, Proposition 3.1.3.7] which by model-categorical nonsense guarantees any marked simplicial set  $(\mathcal{D}, \mathcal{E})$  admits a fibrant replacement, and of [17, Proposition 3.1.3.3] which states that any fibrant replacement map is a Cartesian equivalence, co-representing the same spaces of functors as  $(\mathcal{D}, \mathcal{E})$ .

**Notation 3.25** ( $\mathcal{C}^\sharp$  and  $\mathcal{D}^\sharp$ , Definition 3.1.1.8 and Section 3.1 of [17]). Let  $\mathcal{C}$  be an  $\infty$ -category.  $\mathcal{C}^\sharp$  indicates the marked simplicial set  $(\mathcal{C}, \mathcal{E})$  with the marking  $\mathcal{E}$  given by the equivalences in  $\mathcal{C}$ . For any simplicial set  $\mathcal{D}$ , we let  $\mathcal{D}^\sharp$  denote the marked simplicial set where every edge of  $\mathcal{D}$  is marked.

**Notation 3.26** ( $\text{Map}^\sharp$ , see 3.1.3 of [17]). Let  $(X, \mathcal{E})$  be a marked simplicial set and  $Y$  an  $\infty$ -category. We let

$$\text{Map}^\sharp((X, \mathcal{E}), Y^\natural)$$

denote the Kan complex where a  $k$ -simplex is a map of marked simplicial sets  $f : (\Delta^k)^\sharp \times (X, \mathcal{E}) \rightarrow Y^\natural$ . Explicitly,  $f$  is a  $k$ -simplex of  $\text{Map}^\sharp(X, Y)$  if and only if, for every edge  $e$  of  $\Delta^k$  and every marked edge  $e'$  of  $X$ , the image  $f(e, e')$  is an equivalence in  $Y$ .

**Notation 3.27** ( $X_+$ ). Let  $X$  be a simplicial set. We let  $X_+$  be the free simplicial set generated by the semisimplicial set underlying  $X$ .

**Remark 3.28.** Informally,  $X_+$  is obtained by adjoining degeneracies freely to each simplex of  $X$ . It is most conveniently modeled as a Kan extension of  $X$  (restricted to  $\Delta_{\text{inj}}$ ) along the inclusion  $\Delta_{\text{inj}} \hookrightarrow \Delta$ .

In particular, the set of edges

$$(X_+)_1 \cong X_1 \coprod X_0 \tag{3.5}$$

decomposes as a disjoint union of the edges arising in  $X$  (including all the degenerate edges of  $X$ ), and of all the degenerate edges in  $X_+$ , which is naturally identified with a copy of the set of vertices of  $X$ .

**Example 3.29.** Let  $\mathbf{C}$  be a category (in the usual, non- $\infty$ -categorical sense) where only identity morphisms admit left or right inverses. For example,  $\mathbf{C}$  may be a poset.

We let  $\mathbf{C}^{\text{nu}}$  denote the non-unital category obtained by removing all identity morphisms from  $\mathbf{C}$ . Note  $\mathbf{C}^{\text{nu}}$  still has a well-defined composition thanks to the assumption that  $\mathbf{C}$  has no inverses.

One then has a semisimplicial set  $N_{\text{semi}}(\mathbf{C}^{\text{nu}})$  given by the semisimplicial nerve of  $\mathbf{C}^{\text{nu}}$ ; concretely,  $N_{\text{semi}}(\mathbf{C}^{\text{nu}})$  is the functor  $\Delta_{\text{inj}}^{\text{op}} \rightarrow \mathbf{Sets}$  sending  $[k]$  to the set of functors  $[k]^{\text{nu}} \rightarrow \mathbf{C}^{\text{nu}}$ . Equivalently,  $N_{\text{semi}}(\mathbf{C}^{\text{nu}})$  is the semisimplicial subset of  $N(\mathbf{C})$  obtained by removing all degenerate simplices.

Then the natural map

$$(N_{\text{semi}}(\mathbf{C}^{\text{nu}}))_+ \rightarrow N(\mathbf{C})$$

from the free simplicial set on  $N_{\text{semi}}(\mathbf{C}^{\text{nu}})$  to the nerve of  $\mathbf{C}$  is an isomorphism of simplicial sets.

Treating the poset  $[k]$  as a category, we see a subexample: The standard  $k$ -simplex  $\Delta^k \cong N([k])$  is in fact a free simplicial set; it is generated by the semisimplicial set represented by  $[k]$  in  $\Delta_{\text{inj}}$ ; this semisimplicial set is otherwise written as  $N_{\text{semi}}([k]^{\text{nu}})$ .

Likewise,  $\text{sd}(\Delta^k)$  is the nerve of a poset (the poset of non-empty subsets of  $[k]$ ). Hence we see that  $\text{sd}(\Delta^k)$  is also a free simplicial set.

**Notation 3.30.** Let

$$\mathcal{E} = s_0^X(X_0) \coprod X_0 \subset X_1 \coprod X_0 \cong (X_+)_1$$

be the set of edges in  $X_+$  that are identified with the degenerate edges in  $X$ , or are degenerate in  $X_+$ . (In the above equation, the last subset inclusion passes through the identification (3.5).) The pair

$$(X_+, \mathcal{E})$$

is a marked simplicial set.

As suggested by our language of “free” simplicial set, there is a “free-forget” adjunction between the category of simplicial sets and the category of semisimplicial sets. The functor sending a semisimplicial set  $\mathcal{D}$  to the free simplicial set generated by  $\mathcal{D}$  is the left adjoint; accordingly, one has a counit map  $X_+ \rightarrow X$ .

**Notation 3.31.** Let  $X$  be an  $\infty$ -category and let  $X_+ \rightarrow X$  be the counit of the adjunction. We naturally obtain a map of marked simplicial sets

$$\epsilon : (X_+, \mathcal{E}) \rightarrow X^{\natural}.$$

**Remark 3.32.** The main result of [33], which we have cited as Theorem 3.7, implies that—when  $X$  is an  $\infty$ -category— $\epsilon$  is a Cartesian equivalence of marked simplicial sets. That is, the induced map

$$\text{Map}^{\sharp}(X^{\natural}, -) \rightarrow \text{Map}^{\sharp}((X_+, \mathcal{E}), -)$$

is a homotopy equivalence of Kan complexes for any  $\infty$ -category plugged into the target  $-$ . (This informally states that  $X^{\natural}$  and  $(X_+, \mathcal{E})$  corepresent the same functor when tested only against  $\infty$ -categories.) This is of course equivalent to the formal statement that  $X$  is the localization of (an  $\infty$ -category replacement of)  $X_+$  along the edges  $\mathcal{E}$  (see Remark 3.24) but the explicit combinatorics of marked simplicial sets brings added power.

In this work, we will find ourselves in the following formal setting: We will have two simplicial sets  $X$  and  $Y$ , and a map of semisimplicial sets

$$h_{\text{semi}} : X \rightarrow Y.$$

The geometry will allow us to conclude that  $h_{\text{semi}}$  sends degenerate edges of  $X$  to equivalences in  $Y$ . Now, by adjunction,  $h_{\text{semi}}$  induces a map of simplicial sets

$$(h_{\text{semi}})_+ : X_+ \rightarrow Y.$$

Because  $h_{\text{semi}}$  sends degeneracies of  $X$  to equivalences in  $Y$ , we see that  $(h_{\text{semi}})_+$  is a map of marked simplicial sets with domain  $(X_+, \mathcal{E})$ ; by Remark 3.32, we are guaranteed a unique (up to contractible choice) map of simplicial sets

$$h : X \rightarrow Y \tag{3.6}$$

equipped with a homotopy-commuting diagram

$$\begin{array}{ccc} X_+ & \xrightarrow{(h_{\text{semi}})_+} & Y \\ \epsilon \downarrow & \nearrow h & \\ X & & . \end{array} \tag{3.7}$$

**Notation 3.33** ( $H$ ). Concretely, this triangle encodes a map of simplicial sets

$$H : \Delta^1 \times X_+ \rightarrow Y$$

whose restriction to  $\Delta^{\{0\}} \times X_+$  is equal to  $(h_{\text{semi}})_+$ , to  $\Delta^{\{1\}} \times X_+$  is equal to  $h \circ \epsilon$ , and for which every edge of the form  $(e, e')$  is sent to an equivalence in  $Y$  (see Notation 3.26).

**Remark 3.34.** This homotopy  $H$  empowers us to do the following.  $h$  is a priori completely abstract; on the other hand, for every simplex  $f : \Delta^k \rightarrow X$  of  $X$ , consider the natural simplex  $f_+ : \Delta^k \rightarrow X_+$  satisfying  $\epsilon f_+ = f$ . ( $f_+$  is the image of  $f$  under the unit map  $X \rightarrow X_+$  of semisimplicial sets.)  $H$  allows us to homotope  $h(f)$  to  $(h_{\text{semi}})_+(f_+) = h_{\text{semi}}(f)$  in a way respecting faces. In fact, we will see that  $H$  allows us to replace arguments involving  $h$  with those involving the more explicit  $h_{\text{semi}}$ .

Our proof of Theorem C will depend on the following Lemma. To state it, let  $W$ ,  $X$  and  $Y$  be  $\infty$ -categories and fix functors  $i : W \rightarrow Y$  and  $j : W \rightarrow X$ . Fix also a semisimplicial set map  $h_{\text{semi}} : X \rightarrow Y$  sending degenerate edges of  $X$  to equivalences in  $Y$ . (This in particular gives rise to a simplicial set map  $h$  as in (3.6).) The Lemma allows us to construct a homotopy-coherent triangle

$$\begin{array}{ccc} W & \xrightarrow{j} & X \\ & \searrow i & \downarrow h \\ & & Y. \end{array}$$

**Remark 3.35.** Note that the case  $W = X$  and  $j = \text{id}_X$

**Lemma 3.36.** Suppose that for every simplex  $f : \Delta^k \rightarrow W$ , one may exhibit a map

$$G_f : \Delta^1 \times \Delta^k \rightarrow Y$$

in such a way that the collection  $\{G_f\}_{k \geq 0, f \in W_k}$  satisfies the following:

- (i) ( $G_f$  homotopes between the maps we want.) For each  $f$ ,  $G_f|_{\Delta^{\{0\}} \times \Delta^k} = i(f)$  and  $G_f|_{\Delta^{\{1\}} \times \Delta^k} = h_{\text{semi}} \circ j(f)$ . Moreover, every edge of the form  $(e, e')$  with  $e$  an arbitrary edge of  $\Delta^1$ , and  $e'$  degenerate in  $\Delta^k$ , is sent to an equivalence in  $Y$ .

(ii) ( $G_f$  respects face maps.) For each  $f$  and every  $0 \leq i \leq k$ ,  $G_f|_{\Delta^1 \times \partial_i \Delta^k} = G_{\partial_i f}$ .

Then there exists a homotopy between  $i$  and  $h \circ j$ .

*Proof.* Because the  $G_f$  respect faces, the collection  $\{G_f\}$  glues together to define a map of simplicial sets

$$G : \Delta^1 \times W_+ \rightarrow Y.$$

Letting  $\epsilon_W : W_+ \rightarrow W$  denote the counit, we note that  $G$  restricted to  $\Delta^{\{0\}} \times W_+$  agrees with  $i \circ \epsilon_W$ . On the other hand, let  $j_+ : W_+ \rightarrow X_+$  denote the map of simplicial sets induced by applying the  $+$  functor to  $j$ . Then the restriction of  $G$  to  $\Delta^{\{1\}} \times W_+$  agrees with  $(h_{\text{semi}})_+ \circ j_+$  because each  $G_f$  restricts to  $h_{\text{semi}} \circ j(f)$  by hypothesis.

Having spelled out the consequences of Conditions (i) and (ii), the rest of the proof is formal. The definition of  $H$  (Notation 3.33) tells us

$$H|_{\Delta^{\{0\}} \times X_+} \circ j_+ = (h_{\text{semi}})_+ \circ j_+$$

so we may concatenate the two homotopies  $G$  and  $H$  as follows:

$$G \bigcup_{(h_{\text{semi}})_+ \circ j_+} (H \circ (\text{id}_{\Delta^1} \times j_+)) : \Lambda_1^2 \times W_+ \rightarrow Y.$$

Because  $G$  and  $H$  are homotopies (meaning the non-degenerate edges of  $\Lambda_1^2$  are taken to equivalences in  $Y$ ), this defines a map

$$\Lambda_1^2 \rightarrow \text{Map}^\sharp((W_+, \mathcal{E}), Y^\natural)$$

where  $(W_+, \mathcal{E})$  is the marked simplicial set defined in Notation 3.30. Because  $Y$  is an  $\infty$ -category, the codomain  $\text{Map}^\sharp((W_+, \mathcal{E}), Y^\natural)$  is a Kan complex, so we may fill the horn  $\Lambda_1^2$ ; the resulting map

$$\Delta^1 \cong \partial_1 \Delta^2 = \Delta^{\{0,2\}} \subset \Delta^2 \rightarrow \text{Map}^\sharp((W_+, \mathcal{E}), Y^\natural)$$

exhibits a homotopy from  $i \circ \epsilon_W$  to  $h \circ \epsilon \circ j_+ = h \circ j \circ \epsilon_W$ . (This last equality uses that  $j$  is a map of simplicial sets.) By Remark 3.32, we conclude there is a homotopy between  $i$  and  $h \circ j$ , as desired.  $\square$

### 3.5 Understanding subdivisions of subdivisions

In Section 10.5 we will confront the subdivision of a subdivision of a simplex. So we collect a convenient combinatorial description.

**Notation 3.37.** We let

$$\mathcal{P}'_{\text{linear}}(Q)$$

be the subset of the power set of  $Q$  consisting of non-empty subsets of  $Q$  that are linear under the poset relation of  $Q$ . We treat  $\mathcal{P}'_{\text{linear}}(Q)$  as a poset via the inclusion relation of subsets.

We leave the proof of the following to the reader.

**Proposition 3.38.** Let  $Q$  be a poset.

$$\text{sd}(Q) \cong N(\mathcal{P}'_{\text{linear}}(Q)).$$

In particular, we note that  $\text{sd}(Q)$  is (the nerve of) a poset, so we have a recursive description of iterated subdivisions of any poset.

**Example 3.39.** Fix the  $I$ -simplex  $\Delta^I \cong N(I)$ . Then a vertex of  $\mathbf{sd}(\mathbf{sd}(I))$  is the data of a strictly increasing chain

$$\vec{A} := A_0 \subset \dots \subset A_k$$

of non-empty subsets of  $I$  – i.e., an injective map of posets

$$[k] \xrightarrow{\vec{A}} \mathcal{P}'(I).$$

An  $l$ -simplex of  $\mathbf{sd}(\mathbf{sd}(I))$  is the data of a diagram of posets

$$[k_0] \hookrightarrow [k_1] \hookrightarrow \dots \hookrightarrow [k_l] \xrightarrow{\vec{A}} \mathcal{P}'(I).$$

Such a simplex is degenerate precisely when at least one of the inclusions  $[k_i] \hookrightarrow [k_{i+1}]$  is an isomorphism.

## 4 Simplices and collarings

Much of homotopy theory is organized by the combinatorics of simplices and the ability to fill horns. However, we encounter two issues in trying to corral sectorial data into the usual simplicial framework. First, there is no natural sense in which a Liouville structure on  $M \times T^*\Delta^n$  “restricts” to a Liouville structure on  $M \times T^*\partial_i\Delta^n$  without choosing normal coordinates to the face  $\partial_i\Delta^n \subset \Delta^n$ . Second, we would also like to naturally embed stabilizations of  $M$  and of  $M \times T^*\partial_i\Delta^n$  into  $M \times T^*\Delta^n$ . For this, we must choose once and for all a *collaring* on our simplices, which allow us to determine compatible normal coordinates along all faces of all simplices.

But introducing collarings creates a degeneracy wrinkle: While it is common in most situations to observe that collarings can be chosen compatibly with face maps, standard degeneracy maps may not obviously respect collarings. So Theorems 3.6 and 3.7 come to the rescue. Thanks to them, we may use collars to encode only semisimplicial objects, and then a posteriori promote semisimplicial data to simplicial data.

This sort of gymnastics is common when producing  $\infty$ -categories of geometric origins [32].

Here, we make explicit the existence of the necessary collarings. Now, our particular construction of collarings (Proposition 4.8) forces us to declare the following unconventional convention:

### 4.1 An unconventional convention

**Notation 4.1.** In all following sections, for any finite, non-empty, linearly ordered set  $I$ , the (non-standard)  $I$ -simplex is the topological space

$$\Delta^I := \left\{ \left( \sum_{i \in I} x_i = 1 \right) \& (\forall i \in I, x_i \geq -1) \right\} \subset \mathbb{R}^I. \quad (4.1)$$

The usual definition of the standard  $I$ -simplex requires that all coordinate be non-negative. Allowing for negative values allows for us to define collarings in Construction 4.6.

The usual inclusions of coordinate hyperplanes induce inclusions of lower-dimensinal simplices as faces of higher-dimensional simplices, and hence define a semisimplicial collection of smooth manifolds with corners.

**Remark 4.2.** By induction, one may choose homeomorphisms between the non-standard  $k$ -simplices in this work to the standard  $k$ -simplices in such a way that all face maps are respected. Fixing such a choice, one obtains a semisimplicial isomorphism between all natural semisimplicial sets defined in terms of our non-standard simplices, and in terms of the standard simplices. (For example, the singular complex of maps  $\Delta^k \rightarrow \mathcal{E}$  for some topological space  $\mathcal{E}$ .)

In short, the particular non-standard definition of the  $k$ -simplex has no deep philosophical value, and we would strongly encourage the reader to just pretend all simplices are standard anyway. We imagine a collaring convention exists surely for the standard simplices; we simply didn’t construct one.

### 4.2 Collaring conventions

**Definition 4.3.** Consider the data of:

- For every pair of linearly ordered, finite, non-empty sets  $I, J$ , and for every injective, order-preserving map  $a : I \rightarrow J$ , a smooth, open embedding

$$\eta = \eta_a : \Delta^I \times [-1, \epsilon)^{J \setminus a(I)} \rightarrow \Delta^J$$

whose restriction to  $\Delta^I \times \{0\}$  is the standard simplicial inclusion induced by  $a$ .



The data  $\{\eta_a\}_{(I,J,a)}$  (a choice of  $\eta_a$  for every  $a : I \rightarrow J$ ) are called a *collaring convention* if: For every pair of injective, order-preserving maps  $a : I \rightarrow J, b : J \rightarrow K$ , the following diagram commutes:

$$\begin{array}{ccc} \Delta^I \times [-1, \epsilon)^{K \setminus ba(I)} & \xlongequal{\quad} & (\Delta^I \times [-1, \epsilon)^{J \setminus a(I)}) \times [-1, \epsilon)^{K \setminus b(J)} \xrightarrow{\eta_a \times \text{id}} \Delta^J \times [-1, \epsilon)^{K \setminus b(J)} \\ & \searrow \eta_{b \circ a} & \downarrow \eta_b \\ & & \Delta^K \end{array}$$

**Remark 4.4.** The interval  $[-1, \epsilon)$  is chosen to be compatible with our stabilization operation. If our stabilization were defined by  $- \times T^*[a, b]$ , the collaring would have domain  $\Delta^I \times [a, b + \epsilon)^c$ .

Note also that the definition of collaring guarantees that  $\eta$  is a map respecting the natural stratifications.

**Warning 4.5.** The strict commutativity of the diagram in Definition 4.3 is a subtle but important point. Indeed, the most “natural” way to collar  $\Delta^I$  inside  $\Delta^J$  may be by taking join coordinates to realize  $\Delta^I \star \Delta^{J \setminus I} \cong \Delta^J$ . However, join constructions are only associative up to natural isomorphism.

For any finite set  $I$  and any subset  $J$ , we construct inclusions

$$\Phi_{I \subset J} : \Delta^I \times [-1, 0]^{J \setminus I} \hookrightarrow \Delta^J$$

as follows.

**Construction 4.6.** First, note that it is easy to build piecewise smooth collars  $\phi_{I \subset J}$ . Indeed, given  $x \in \Delta^I$ , we can define  $\phi_{I \subset J}(x, t) = \phi_{S_x}(x, t)$ , where  $S_x = \{i \mid x_i > 0\}$  and

$$(\phi_S(x, t))_j = \begin{cases} t_j & j \notin I \\ x_j & j \in I \setminus S \\ b_S(x, t)x_j & j \in S. \end{cases}$$

Here,  $b_S(x, t) = \frac{1 - \sum t_j - \sum_{i \notin S} x_i}{\sum_{i \in S} x_i}$  is the unique scalar for which  $\phi_{S_x}(x, t)$  lands in  $\Delta^J$ . The main task is thus to smooth this map while ensuring that the result is still coherent and is still an embedding.

To that end, suppose that we had a permutation-equivariant partition of unity  $\{g_S\}_{S \subset J}$  on  $\Delta^J$  with the following two properties.

- i If  $g_S(x) > 0$ , then  $x_s > 0$  for all  $s \in S$ .
- ii  $g_S(x)$  depends only on the positive components of  $x$ . In particular, its dependence on the negative components factors through their sum.

Consider a path  $\mathbf{t} : [0, 1] \rightarrow [-1, 0]^{J \setminus I}$ . For each  $\tau \in [0, 1]$  and  $S \subset I$ , we get an embedding

$$\phi_S(\bullet, \mathbf{t}(\tau)) : \Delta^I \cap \text{supp}(g_S) \hookrightarrow \Delta_{\mathbf{t}(\tau)}^I,$$

where  $\Delta_{\mathbf{t}(\tau)}^I = \{x \in \Delta^J \mid x_j = \mathbf{t}_j(\tau)\}$ . We can combine these into a smooth vector field along  $\Delta_{\mathbf{t}(\tau)}^I$  by taking the weighted sum

$$\xi(\tau) = \sum_{S \subset I} g_S \cdot \frac{\partial}{\partial \tau} \phi_S(\bullet, \mathbf{t}(\tau)). \quad (4.2)$$

(Note that a given  $x \in \Delta_{\mathbf{t}(\tau)}^I$  will come from different values of  $\bullet$  for different  $S$ . See Remark 4.7.) Define the collar  $\Phi_{I \subset J}(\bullet, \mathbf{t}(1))$  to be the time 1 flow of this vector field  $\xi$ . Then  $\Phi_{I \subset J}$  is coherent

with respect to face maps by condition (i), and it is independent of the path  $\mathbf{t}$  because both  $g_S$  and  $\phi_S$  factor through  $\mathbf{t} \mapsto \sum \mathbf{t}_j$ .

It thus suffices to construct the partition of unity  $\{g_S\}$ . Let us begin by fixing some notation. Given real numbers  $a < b$ , we'll write  $\kappa_a^b: \mathbb{R} \rightarrow [0, 1]$  for a nondecreasing cutoff function with  $\kappa_a^b(a) = 0$  and  $\kappa_a^b(b) = 1$ . For a subset  $I \subset J$  and a point  $x \in \Delta^J$ , we'll write  $x_I = \sum_{i \in I} x_i$ .

For  $S \subset J$  of size  $n > 0$ , define

$$\kappa_S(x) := \kappa_{1 - \frac{1}{2^n}}^{1 - \frac{1}{2^{n+1}}}(x_S) \prod_{s \in S} \kappa_{\frac{1}{2^{n+1}}}^{\frac{1}{2^n}}(x_s),$$

which is nonzero if and only if  $x_S$  is big and all  $x_s$  are not too small. These functions are positive in the right places and have the correct dependence on the signs of the components of  $x$ , but it is possible for them to overlap and sum to a quantity larger than 1. To repair this, set

$$f_S := \kappa_S \cdot \prod_{\substack{S' \neq S \\ |S'| = n}} (1 - \kappa_{S'}),$$

which satisfies  $\sum_{|S|=n} f_S \leq 1$ . Finally, we can inductively define our partition of unity by  $g_\emptyset = 0$  and

$$g_S = f_S \cdot \left(1 - \sum_{|S'| < n} g_{S'}\right).$$

Let us check the necessary properties. Permutation-equivariance and properties (i) and (ii) are automatic from the construction of the  $\kappa_S$ , so the main task is to show that  $\{g_S\}$  is indeed a partition of unity. To that end, consider a point  $x \in \Delta^J$ , set  $S = S_x$  to be the set of positive components, and set  $n = |S|$ .

If  $\kappa_S(x) = 1$ , then we are done, since any other  $S'$  of size  $n$  must index a negative component, so  $\kappa_{S'}$  must vanish. This implies that  $f_S(x) = 1$ , and so  $\sum_{|S'| \leq n} g_{S'}(x) = 1$ .

If  $\kappa_S(x) < 1$ , then since  $x_S \geq 1 > 1 - \frac{1}{2^n}$  there must be some component  $s_1 \in S$  so that  $x_{s_1} < \frac{1}{2^n}$ . That means  $x_{S \setminus \{s_1\}} > 1 - x_{s_1} > 1 - \frac{1}{2^{n-1}}$ , and we can repeat the argument with  $S \setminus \{s_1\}$  (replacing “negative component” in the previous paragraph with “component smaller than  $\frac{1}{2^{n-1+1}} = \frac{1}{2^n}$ ”). The argument terminates for some nonempty  $S \setminus \{x_1, \dots, x_m\}$ , since otherwise we would have

$$x_S < \frac{1}{2^n} + \frac{1}{2^{n-1}} + \dots + \frac{1}{2^1},$$

contradicting  $x_S \geq 1$ .

**Remark 4.7.** Let us write out the precise (but very cluttered) meaning of (4.2). For  $x \in \Delta_{\mathbf{t}(\tau)}^I$ , we have  $x = \phi_S \left( [\phi_S(\bullet, \mathbf{t}(\tau))]^{-1}(x), \mathbf{t}(\tau) \right)$ . The weighted sum is

$$\sum_{S \subset I} g_S(x, \mathbf{t}(\tau)) \cdot D_2 \phi_S \left( [\phi_S(\bullet, \mathbf{t}(\tau))]^{-1}(x), \mathbf{t}(\tau) \right) \circ \frac{\partial \mathbf{t}}{\partial \tau},$$

where  $D_2$  is the partial derivative with respect to the second component.

Thus, we have shown

**Proposition 4.8.** A collaring convention exists.

**Choice 4.9.** From hereon, we fix a collaring convention.

**Remark 4.10.** A collaring convention renders every simplex  $S = \Delta^k$  a collared manifold-with-corners in the sense of Definition 2.37. The convention further provides the data showing that the collarings are compatible with the usual face maps.

**Remark 4.11.** Thanks to a choice of collaring convention, we may ask geometric structure on  $M \times T^*\Delta^k$  known to *split* with respect to the collaring near the boundary—that is, we may demand that structures be a direct product of a structure on  $M \times T^*\Delta^A$ , and on  $T^*[-1, \epsilon]^i$ . (Here,  $A$  is some subset of  $[k]$ .)

### 4.3 Linearity, locally

At the heart of smooth approximation theorems for  $\infty$ -dimensional smooth spaces is the ability to reduce local problems to linear ones. (See for example [27] and [15].) More precisely, suppose we have a continuous collection in an infinite-dimensional space, and we seek to deform it to a smooth collection. A tried-and-true strategy is to write the collection locally as a continuous collection of functions into a linear object (for examples, as a collection of sections of some vector bundle, e.g., as a collection of real-valued functions) and then use the classical smooth approximation theorem for functions to deform these functions into a smooth collection.

We now prove that  $\text{Emb}_{\text{Liou}}$  is amenable to this strategy.

**Proposition 4.12.** Let  $S$  be a manifold (possibly with corners) and let  $j : S \rightarrow \text{Emb}_{\text{Liou}}(M, N)$  be a continuous and collared map. ( $j$  is not assumed smooth.) Then, for every  $s_0 \in S$ , there exist

- (i) some neighborhood  $U$  with  $s_0 \in U$  and
- (ii) a continuous function  $H : U \times N \rightarrow \mathbb{R}$ , whose restriction to each  $\{s\} \times N \subset U \times N$  is smooth,

so that for all  $s \in U$ ,  $j_s = \text{Flow}_1^{X_{H_s}} \circ j_{s_0}$ . In other words,  $j_s$  is obtained from  $j_{s_0}$  by post-composing with the time-1 flow of some time-independent but  $s$ -dependent family of Hamiltonians.

*Proof.* Fixing  $s_0 \in S$ , we may as well assume that  $M \subset N$  and that  $j_{s_0}$  is the canonical inclusion of the subset. Thus the graph of  $j_{s_0}$  may be thought of as a subset of the diagonal of  $N \times N$ , which in particular has a normal neighborhood identified with an open subset  $V \subset T^*N$ . Under this identification, the image of  $j_{s_0}$  is identified with the graph of the zero 1-form of  $N$  (restricted to  $M$ ).

We now claim that for a small-enough neighborhood  $U$  with  $s_0 \in U \subset S$ , and for all  $s \in U$ , the graph of  $j_s$  is contained in  $V$ . Note that if  $M$  and  $N$  are compact, this is a consequence of the weak  $C^\infty$ -topology on smooth mapping spaces. But both  $M$  and  $N$  are non-compact. So invoke the continuity of  $j$  with respect to the colimit topology of  $\text{Emb}_{\text{Liou}}(M, N)$ ; then the standard yoga of shrinking of  $U$ , choosing a single  $K$  as in Remark 2.29, and using the definition of the weak  $C^\infty$ -topology, we may assume that for all  $s \in U$ , (i) the graph of  $j_s(K)$  is contained in  $V$ , and (ii)  $j_s^* \lambda^N = \lambda^M + dh_s$  where the support of  $h_s$  is inside  $K$ .

It is now a standard fact of symplectic geometry that, interpreting the image of  $j_s(K)$  as the graph of a 1-form, we may conclude that the image of each  $j_s(K)$  is the graph of an *exact* 1-form (Remark 4.13).

These exact 1-forms are  $C^0$ -dependent on  $s \in S$  by hypothesis, so by integrating, we may in fact choose a single continuous function  $g : U \times K \rightarrow \mathbb{R}$  for which the graph of  $d_M(g|_{\{s\} \times K})$  is equal to the graph of  $j_s(K)$  (when both are interpreted as subsets of  $V \subset T^*N$ ).

(The following works on every path-connected component of  $K$ , so we may assume  $K$  is path-connected and fix a base point  $x_0 \in K$ . Given that  $j_s(K) = \text{graph}(\alpha_s)$  and  $\alpha_s$  is exact, choose a smooth function  $g_s$  so that  $dg_s = \alpha_s$  and  $g_s(x_0) = 0$ . This recovers  $g_s(x) = \int_\gamma \alpha_s$  by choosing any smooth  $\gamma$  with endpoints  $x_0$  and  $x$ . Because  $\alpha_s$  depends continuously on  $s$  in the  $C^\infty$ -topology,  $g(s, x) := g_s(x)$  is a continuous function.)

Of course, the graph of an exact 1-form  $\alpha = dg$  is the result of flowing the zero-section for time 1 along the Hamiltonian defined by  $g$ .

For each  $s$ , having fixed the value of  $g_s$  on  $K$ , there is a unique extension of  $g_s$  to  $\tilde{g}_s$  on  $M$  compatible with that Liouville flow of  $N$  outside  $K$  (so that, in particular, the time-1 flow  $j_s$  commutes with the Liouville flow outside of  $K$ ). Then  $H(x, s) = \tilde{g}_s(x)$  is smooth in  $x$  and continuous in  $(s, x)$ ; and we are finished.  $\square$

**Remark 4.13.** Let us explain why an exact symplectic embedding near the identity is the graph (in the cotangent bundle) of an exact 1-form. Endow  $M \times M$  with the 1-form  $\lambda \oplus -\lambda$ . On the other hand, because the diagonal  $\Delta_M \subset M \times M$  is a Lagrangian, we have a Weinstein neighborhood  $T^*\Delta_M \supset U \hookrightarrow M \times M$ , so we have another Liouville structure in a neighborhood of  $\Delta_M$  given by pushing forward  $\mathbf{pdq}|_U$ . Now note that along the diagonal  $\Delta_M$ ,  $(\lambda \oplus -\lambda) - \mathbf{pdq} = 0$ ; in particular, the two Liouville forms differ by an exact form along the diagonal. Because we may choose  $U$  to deformation retract to  $\Delta_M$ , we conclude the difference  $(\lambda \oplus -\lambda) - \mathbf{pdq}$  is exact on all of  $U$ .

Now, let us use that the graph  $\text{Graph}(j)$  of an exact symplectic embedding  $j : M \rightarrow M$  is an exact Lagrangian for the form  $\lambda \oplus -\lambda$ . (Since  $j^*\lambda = \lambda + dg$  for some  $g$ , the restriction of  $\lambda \oplus -\lambda$  to  $\text{Graph}(j)$  is also  $dg$ .) By the previous paragraph, it follows that  $\text{Graph}(j)$  is an exact Lagrangian in  $U \subset T^*\Delta_M$  as well. That  $j$  is near to  $\text{id}_M$  means that  $\text{Graph}(j)$ , treated as a brane in  $T^*\Delta_M$ , is the graph of a 1-form, while the exactness of the brane implies it is the graph of an *exact* 1-form.

#### 4.4 From continuous to smooth

We collect some facts that allow us to pass between *continuous* families of smooth maps and *smooth* families of smooth maps. We will use this trick repeatedly in showing that horns can be filled in  $\text{Liou}_\Delta$  later on (Theorem 6.1)—by first constructing *continuous* data filling a horn (guaranteed by general properties of topological spaces like  $\text{Emb}_{\text{Liou}}(M, N)$ ), then homotoping this data to be smooth (because all data in  $\text{Liou}_\Delta$  are required to be smooth).

**Remark 4.14** (From continuous to continuous and collared). Fix a topological space  $\mathcal{E}$  and a continuous map  $f : \Delta^k \rightarrow \mathcal{E}$ . Recall that  $f$  is called *collared* if the value of  $f$  is independent of the chosen coordinates normal to the faces and corners of  $\Delta^k$  (Definition 2.40). In particular, choose a continuous homotopy from  $\Delta^k$  to itself which is equal to the identity along the boundary strata, but collapses the normal coordinates. (Such a homotopy is necessarily not smooth.) Pre-composing  $f$  by this homotopy, we see that  $f$  is continuously homotopic rel boundary to a collared map.

Of course, one may inductively choose these homotopies for the 1-simplex, then the 2-simplex, and so forth, so that these homotopies respect face maps (i.e., respect restriction to boundary faces).

Now consider the inclusion—of the semisimplicial set of all *collared* continuous maps in a fixed space  $\mathcal{E}$ —into the semisimplicial set of all (not necessarily collared) continuous maps to  $\mathcal{E}$ . By Steimle’s theorem (Theorem 3.6) the domain may be promoted to be a simplicial set, and by Theorem 3.7, this semisimplicial set map may be promoted to a map of simplicial sets up to homotopy. In particular, the previous paragraphs show that the induced map of simplicial sets is a weak homotopy equivalence.

Our goal for Section 4.4 is to show that we can continuously homotope a continuous and collared map into a smooth and collared map (Proposition 4.16). We start with a Lemma, which crucially uses that we can convert small families of sectorial embeddings into linear data (Proposition 4.12).

**Lemma 4.15.** Suppose  $j : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, N)$  is a continuous map that is collared along the boundary strata of  $\Delta^k$ , and is smooth when restricted along any face of  $\Delta^k$ . Then there exists a homotopy of  $j$ , rel boundary, to a smooth and collared and continuous<sup>18</sup> map from  $\Delta^k$  to  $\text{Emb}_{\text{Liou}}(M, N)$ .

*Proof.* First, recall that we may locally model  $\text{Emb}_{\text{Liou}}(M, N)$  as consisting of linear data as in Proposition 4.12 which—using the notation of loc. cit.—we now apply to the case  $S = \Delta^k$ .

Choose a fine enough simplicial subdivision  $\mathbb{D}$  of  $\Delta^k$  so that for every simplex  $\sigma$  in the subdivision, the open star<sup>19</sup> of  $\sigma$  is contained in a neighborhood  $U$  as in Proposition 4.12.

We construct the homotopy inductively, starting with the index  $n = -1$ , and setting  $j^{(-1)} = j$ . Assume  $j^{(n)}$  has been given, with the property that in a neighborhood of any  $n$ -simplex of  $\mathbb{D}$ ,  $j^{(n)}$  is smooth.

For every  $(n + 1)$ -simplex  $\sigma$  in  $\mathbb{D}$  and its open star  $U_\sigma$ , fix some point  $s_0 \in \sigma$  so that we may treat the data of  $j^{(n)}|_{U_\sigma}$  as a function  $H_\sigma : U_\sigma \times N \rightarrow \mathbb{R}$ , so that  $j^{(n)}(s) = \text{Flow}_1^{X_{H_\sigma}} \circ j_{s_0}^{(n)}$  for all  $s \in U_\sigma$ . As in Proposition 4.12,  $H_\sigma$  is continuous, but is smooth upon restriction to  $\{s\} \times N$  for any  $s \in U_\sigma$ . By usual smooth approximation, we may continuously homotope  $H_\sigma$  to another function  $G_\sigma : U_\sigma \times N \rightarrow \mathbb{R}$  so that

- (i)  $G_\sigma$  is *smooth* in a tiny neighborhood  $V_\sigma \subset \subset U_\sigma$  containing  $\sigma$ , meaning the induced  $V_\sigma \times N \rightarrow \mathbb{R}$  is smooth,
- (ii)  $G_\sigma$  is equal to  $H_\sigma$  outside of  $V_\sigma$ , and on neighborhoods of any  $m$ -simplex for  $m < n$ . (More precisely, the homotopy can be chosen to be constant outside of  $V_\sigma$  and on neighborhoods of all  $m$ -simplices.)
- (iii) For every  $s \in U_\sigma$ , the time-1 Hamiltonian flow of  $G_\sigma(s, -) : N \rightarrow \mathbb{R}$  is a sectorial embedding for all  $s \in U_\sigma$ .
- (iv) Let  $K \subset M$  be the compact set outside which  $j^{(n)}|_{U_\sigma}$  is strictly compatible with  $\lambda$ . Then the sectorial embeddings induced by  $G_\sigma(s, -)$  are also strictly compatible with  $\lambda$  outside  $K$ .

We have a collection of functions  $\{G_\sigma\}_{\sigma \in \mathbb{D}_{n+1}}$ , and by flowing by time 1, we in turn obtain a collection of maps

$$\{j_\sigma^{(n+1)} : U_\sigma \rightarrow \text{Emb}_{\text{Liou}}(M, N)\}_{\sigma \in \mathbb{D}_{n+1}}.$$

The fact that the value of  $G_\sigma$  and  $H_\sigma$  agree outside  $V_\sigma$ , and on neighborhoods of all lower-dimensional simplices, means that  $j_\sigma^{(n+1)}$  agrees with  $j^{(n)}$  in these regions; in particular, the collection  $\{j_\sigma^{(n+1)}\}_{\sigma \in \mathbb{D}_{n+1}}$  (and the collection of homotopies from  $j^{(n)}|_{U_\sigma}$  to  $j_\sigma^{(n+1)}|_{U_\sigma}$ ) glues together to form a single map (and a single homotopy from  $j^{(n)}$  to)

$$j^{(n+1)} : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}.$$

Again because  $j^{(n+1)}$  does not alter the value of  $j^{(n)}$  near boundaries of open stars, it in particular does not change the value of  $j^{(n)}$  near the boundary of  $\Delta^k$  itself. This proves the “rel boundary”

<sup>18</sup>For the terminology behind imposing both smoothness and continuity, see Remark 2.63.

<sup>19</sup>Given a triangulation, recall that the open star of a subsimplex  $\sigma$  is defined to be the union of the interior of all the simplices that contain  $\sigma$  as a face (including  $\sigma$  itself).

claim for every homotopy, and in particular, that each  $j^{(n+1)}$  is collared. The continuity of  $j^{(n+1)}$  is a result of Condition iv.

After our final inductive step, we have arrived at a smooth map  $j^{(k)} : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, N)$  satisfying all the claims of the Lemma.  $\square$

**Proposition 4.16** (From continuous and collared to smooth and collared). If  $j : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, N)$  is continuous,  $j$  can be homotoped continuously to a collared and smooth and continuous map.

Moreover, suppose that the restriction of  $j$  to some collection of faces of  $\Delta^k$  is smooth and collared. Then  $j$  may be homotoped *rel this collection of faces* to a collared and smooth and continuous map.

*Proof.* By Remark 4.14 we may suppose that  $j$  is both continuous and collared. By induction on dimension of the faces, we may assume that the restrictions  $j|_{\Delta^A}$  to the relevant faces of  $\Delta^k$  are smooth, collared, and continuous by Lemma 4.15. We are finished.  $\square$

We have the following:

**Proposition 4.17.** Let  $\text{Emb}_{\text{Liou}}(M, N)$  be the semisimplicial set given by (forgetting the degeneracies of)  $\text{Sing}(\text{Emb}_{\text{Liou}}(M, N))$  (Notation 2.59). Let

$$\text{Emb}_{\text{Liou}}^{\text{coll}}(M, N)$$

denote the sub-semisimplicial set consisting only of the collared continuous simplices, and

$$\text{Emb}_{\text{Liou}}^{\text{coll}, C^\infty}(M, N)$$

be the further sub-semisimplicial set consisting of *smooth* and collared and continuous simplices. Then each semisimplicial set may be promoted to be a Kan complex, and further, the inclusions of semisimplicial sets induce homotopy equivalences of Kan complexes.

*Proof.* It is well-known that  $\text{Sing}$  of any topological space is a Kan complex, so it is obvious that  $\text{Emb}_{\text{Liou}}(M, N)$  may be promoted to be a Kan complex.

For  $\text{Emb}_{\text{Liou}}^{\text{coll}}(M, N)$ , we may easily verify the hypotheses of Steimle's theorem (Theorem 3.6). For example, for any 0-simplex of the semisimplicial set, the constant edge at that vertex realizes an idempotent edge. That this edge is a self-equivalence is verified by filling the horn in question by a non-collared simplex, then homotoping the non-collared simplex to a collared simplex *rel the horn* (using similar techniques as Remark 4.14). The end result of this homotopy is the filler we seek. So  $\text{Emb}_{\text{Liou}}^{\text{coll}}(M, N)$  satisfies the hypotheses of Steimle's Theorem.

For  $\text{Emb}_{\text{Liou}}^{\text{coll}, C^\infty}(M, N)$ , the constant edges again are idempotents (filled by constant 2-simplices). To see that these edges admit fillers of the relevant horns, use the previous paragraph to fill the horn with a collared and continuous simplex, then use Proposition 4.16 *rel the horn* to homotope this filler to a collared, smooth, and continuous simplex.

Finally, the inclusions of semisimplicial sets obviously respect the constant edges, so the hypotheses of Theorem 3.7 are satisfied, and we obtain maps of simplicial sets

$$\text{Emb}_{\text{Liou}}^{\text{coll}, C^\infty}(M, N) \rightarrow \text{Emb}_{\text{Liou}}^{\text{coll}}(M, N) \rightarrow \text{Emb}_{\text{Liou}}(M, N). \quad (4.3)$$

These maps are bijections on vertices. On higher simplices, Theorem 3.7 guarantees that these simplicial maps respect the original semisimplicial maps up to homotopy. In particular, by repeated application of Lemma 4.15 and Remark 4.14, we see that these inclusions all induce isomorphisms on all homotopy groups. This completes the proof.  $\square$

## 4.5 The homotopy equivalence of different deformation embedding spaces

Recall that we have defined five different spaces of morphisms between Liouville sectors. They are organized in (2.7). The following proves they are all homotopy equivalent.

**Theorem 4.18.** The right-pointing arrows in (2.7) are weak homotopy equivalences.

We will need the following version of Moser's theorem in proving Lemma 4.20.

**Lemma 4.19.** Let

$$\lambda = (T_b, \lambda_{t,b}): B \rightarrow \text{Def}_{\text{Moore}}(N)$$

be a family of bordered (and not necessarily compactly supported) deformations of  $N$  parametrized by some compact manifold-with-corners  $B$ , and let  $E \subset N$  be a Liouville-invariant neighborhood of  $\partial N$  on which  $\lambda$  is trivial. Then there is a family  $\psi_{t,b}: \mathbb{R} \times B \rightarrow \text{Diff}(N)$  which (i) restricts to  $\text{id}_E$ , (ii) agrees with  $\text{id}_N$  for  $t \leq 0$ , (iii) is  $t$ -independent for  $t \geq T_b$ , and such that (iv)  $\psi_{t,b}^* \lambda^N = \lambda_{t,b} + dh_{t,b}$ , where  $h_{t,b}$  is uniformly compactly supported.

*Proof.* The proof is essentially the same as for the usual Moser theorem for Liouville manifolds [6], so we give only a sketch. The key point is that, by definition of  $\mathbf{L}(N)$ , we can find smoothly varying choices of symplectization coordinates  $r_{t,b}$ , which induce smoothly varying contact structures on  $\partial^\infty N$ . Applying Gray's stability theorem in a  $B$ -family, we may immediately assume that  $\lambda_{t,b} = \lambda^N$  outside a uniform compact set. The remaining isotopy is given by the flow of  $\xi_{t,b}$ , where  $\iota_{\xi_{t,b}} d\lambda_{t,b} = \frac{d}{dt} \lambda_{t,b}$ .  $\square$

**Lemma 4.20.** Let  $B$  be a compact manifold-with-corners, and let  $C \subset B$  be a compact submanifold-with-corners which is a smooth deformation retract of a neighborhood of itself (e.g. a closed submanifold or union of disjoint boundary faces). Then any smooth family

$$\Phi: (B, C) \rightarrow \left( \text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}}(M, N), \text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N) \right)$$

is smoothly homotopic rel  $C$  to a family  $B \rightarrow \text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$ .

*Proof of Lemma 4.20.* After reparametrizing by a partial retraction, we may assume that  $\Phi(b)$  lands in the compactly-supported-deformation-embedding space  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$  on some regular neighborhood  $U$  of  $C$ . We will explain how to homotope  $\Phi$  on the complement  $B^c = B \setminus U$ ; the extension to  $U$  is obtained by interpolating the duration over which we perform each step of the construction (in a way that it only occurs in the region where the previous step was fully performed).

Write  $N_{t,b} = (N, \lambda_{t,b})$  for the family of Liouville sectors and  $\Phi_{\text{str}}$  for the family of underlying strict embeddings  $M \hookrightarrow N_{T_b, b}$ . By post-composing  $\Phi_{\text{str}}$  with a family of shrinking isotopies, we may assume that the image of  $\Phi_{\text{str}}$  avoids  $\partial N$ , and by strictness this means that it avoids a  $\lambda_{T_b, b}$ -Liouville-invariant neighborhood of  $\partial N$  for all  $b \in B^c$ . Since all deformations are bordered and  $B$  is compact, this neighborhood can be made uniform in  $B$ , and in fact it can be taken to be  $(\pi^{rc})^{-1}[0, 1)$  for some corner-rounding projection  $\pi^{rc}$ .

Since most of our remaining work will take place in this neighborhood, we will write  $\pi_{t,b}^{rc}$  for the function  $\pi^{rc}$  thought of as a corner-rounding projection on  $N_{t,b}$ . Write  $F_{t,b}$  for the fiber of  $\pi_{t,b}^{rc}$ . By Lemma 4.19, there is a smooth family of symplectomorphisms  $\psi_{t,b}: F_{t,b} \cong F_{0,b}$  so that  $\psi_{t,b}^* \lambda_{0,b} = \lambda_{t,b} + dh_{t,b}$ , where  $h_{t,b}$  is uniformly compactly supported. Crossing with  $T^*(0, 1)$ , extend  $\psi_{t,b}$  to a symplectomorphism

$$\Psi_{t,b}: (\pi_{t,b}^{rc})^{-1} T^*(0, 1) \rightarrow (\pi_{0,b}^{rc})^{-1} T^*(0, 1).$$

To turn off the deformation of  $\partial N$  while preserving the embedding  $\Phi_{\text{str}}$ , we will need to reparametrize the deformation parameter  $t$  as follows. For  $\tau \in [0, 1]$ , write  $s(t, \tau)$  for a concave function which is constant for  $t \geq \frac{1}{\tau} - 1$  and has  $s(t, \tau) = t$  for  $t \leq \frac{1}{\tau} - 2$ . Then, for any  $\tau$ ,  $N_{s(t, \tau), b}$  is  $t$ -independent for  $t \geq T_b + 1$ , so we begin by moving the deformation time  $T_b$  to  $T_b + 1$ . Next, perform the following  $\tau$ -parametrized modifications simultaneously.

- (i) Replace  $N_{t, b}$  with  $N_{s(t, \tau), b}$  on the collar  $(\pi_{s(t, \tau), b}^{rc})^{-1}T^*[0, \frac{1}{4})$ , so that at the end of the modification  $\partial N_{s(t, 1), b} \equiv \partial N$ .
- (ii) Replace the Liouville form  $\lambda_{s(t, \tau), b}$  on  $(\pi_{s(t, \tau), b}^{rc})^{-1}T^*[\frac{1}{4}, \frac{1}{2}]$  with the movie  $\lambda_{s(t, \tau), b} + d\tilde{h}_{t, \tau, b}$ , where  $\tilde{h}_{t, \tau, b}$  interpolates between 0 near  $q = \frac{1}{4}$  and  $h_{s(t, \tau), b} - (\psi_{t, b}^{-1}\psi_{s(t, \tau), b})^*h_{t, b}$  near  $q = \frac{1}{2}$ .
- (iii) Replace the Liouville form on  $(\pi_{s(t, \tau), b}^{rc})^{-1}T^*[\frac{1}{2}, 1)$  with  $\tilde{\Psi}_{t, \tau, b}^*\lambda_{t, b}$ , where  $\tilde{\Psi}_{t, \tau, b}$  is a diffeomorphism interpolating (by integrating an interpolating vector field) between  $\Psi_{t, b}^{-1}\Psi_{s(t, \tau), b}$  near  $q = \frac{1}{2}$  and  $\text{id}_N$  for  $q \geq \frac{3}{4}$ .

After the above modification, we are in a setting where the family of deformation  $N_{t, b}$  is trivial in the Liouville-invariant neighborhood  $(\pi^{rc})^{-1}T^*[0, \frac{1}{4})$  of  $\partial N$ , meaning that we have retracted our family of bordered deformation morphisms to a family of interior deformation morphisms. The passage from interior to compactly supported is another application of Lemma 4.19.  $\square$

*Proof of Theorem 4.18.* Taking  $B$  and  $C$  to be a cylinder and a sphere, respectively, Lemma 4.20 shows that the smooth homotopy groups of  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$  are isomorphic.

Using methods similar to the proof of Proposition 4.17, we in fact have equivalences among all of the following:

- The Kan complex of continuous simplices in  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$ .
- The Kan complex of collared simplices in  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$ .
- The semisimplicial set (made into a Kan complex by Steimle's theorem) of smooth and collared and continuous simplices in  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}, \text{Moore}}^{\text{Def}, \text{cmpct}}(M, N)$ .

(The first of these shows the weak homotopy equivalence, but we record the other equivalences for later use.)

Now, that two other edges in the square of (2.7) are weak homotopy equivalences implies that the bottom horizontal arrow must also be a weak homotopy equivalence. And again, we have a equivalences of the following Kan complexes:

- The Kan complex of continuous simplices in  $\text{Emb}_{\text{Liou}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}}^{\text{Def}, \text{cmpct}}(M, N)$ .
- The Kan complex of collared simplices in  $\text{Emb}_{\text{Liou}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}}^{\text{Def}, \text{cmpct}}(M, N)$ .
- The semisimplicial set (made into a Kan complex by Steimle's theorem) of smooth and collared and continuous simplices in  $\text{Emb}_{\text{Liou}}^{\text{Def}}(M, N)$  and  $\text{Emb}_{\text{Liou}}^{\text{Def}, \text{cmpct}}(M, N)$ .

$\square$



**Remark 4.21.** One can construct a space of not-necessarily-exact, not-necessarily-bordered Liouville deformations still satisfying Condition (T). With minor additions to the proof of Lemma 4.20, the above arguments show that the resulting spaces of generalized deformation embeddings are homotopy equivalent to the standard ones.

Lemma 4.20 has another significant consequence.

**Definition 4.22.** Fix Liouville sectors  $M$  and  $N$ . We say that a (bordered, but not necessarily compactly supported) deformation embedding  $(f, \{\lambda_t^N\}_{t \in \Delta^1})$  is a *deformation equivalence* if there exists another (bordered, but not necessarily compactly supported) deformation embedding

$$(g, \{\lambda_t^M\}_{t \in \Delta^1})$$

so that

1. The pair  $(gf, \widetilde{g_*\{\lambda_t^N\}} \star \{\lambda_t^M\})$  is homotopic to  $\text{id}_M$  and the constant deformation of  $\lambda^M$  – here, “homotopic” means “through a smooth family of (bordered, but not necessarily compactly supported) deformation embeddings.” In particular,  $gf$  is smoothly isotopic to  $\text{id}_M$ .
2. Likewise, we demand the pair  $(fg, \widetilde{f_*\{\lambda_t^M\}} \star \{\lambda_t^N\})$  is homotopic to  $\text{id}_N$  with the constant  $\lambda^N$  deformation, through a smooth family of (bordered, but not necessarily compactly supported) deformation embeddings.

**Remark 4.23.** Let us explain the notation in Definition 4.22. First,  $\widetilde{g_*\{\lambda_t^N\}_t}$  is an extension of the deformation on the image of  $N$  to a deformation on all of  $M$ .

Next, the notation  $\lambda'_t \star \lambda_t$  denotes a family of Liouville forms obtained through concatenation – first through the family  $\lambda_t$ , then through the family  $\lambda'_t$ . In particular, one may think of  $\widetilde{g_*\{\lambda_t^N\}} \star \{\lambda_t^M\}$  as a family parametrized by the horn  $\Delta^1 \cup \Delta^1 \cong \Lambda_1^2$  comprising two edges in a 2-simplex. Then the homotopy required in each item of Definition 4.22 “fills in” this horn to a 2-simplex  $\Delta^2$ , where the remaining edge in  $\Delta^2$  is occupied by a constant family.

See also Remark 2.33.

Definition 4.22 is the notion of equivalence in any  $\infty$ -category whose morphisms are deformation embeddings (through deformations that are bordered but not necessarily compactly supported). Thus, it seems to be a far weaker relation to be deformation equivalent, than to be sectorially equivalent (Definition 2.30). However:

**Corollary 4.24.** Deformation-equivalent Liouville sectors are sectorially equivalent.

*Proof.* Lemma 4.20 supplies us with morphisms in each direction. The composition of these morphisms are homotopic to identity in  $\text{Emb}_{\text{Liou}}^{\text{Def}}$ , so by Theorem 4.18 they are homotopic to the identity in  $\text{Emb}_{\text{Liou}}^{\text{Def, cmpct}}$ .  $\square$

## 5 $\text{Liou}_\Delta$ and $\text{Liou}_\Delta^{\text{Def}}$

Informally, movies of maps (Definition 2.46) arise when we convert a family of morphisms into the data of a single morphism. Unlike the usual definition of homotopy, which traditionally converts a family of maps  $X \rightarrow Y$  into a map obtained by replacing the domain (with  $X \times [0, 1]$ ), we may replace a family of embeddings with a single strict morphism by thickening both the domain and codomain to remain in the world of codimension zero embeddings.

This allows us to construct semisimplicial sets  $\text{Liou}_\Delta$  and  $\text{Liou}_\Delta^{\text{Def}}$  whose  $n$ -simplices are modeled by a diagram of strict sectorial embeddings in the shape of the barycentric subdivision of the  $n$ -simplex (Definition 5.12). The difference between the two is that  $\text{Liou}_\Delta^{\text{Def}}$  allows for higher simplices to encode all bordered and collared deformations, while  $\text{Liou}_\Delta \subset \text{Liou}_\Delta^{\text{Def}}$  demands that deformations further be compactly supported.

The present section constructs  $\text{Liou}_\Delta$  and  $\text{Liou}_\Delta^{\text{Def}}$  and proves some of their basic properties. We will later see that both may be promoted to be an  $\infty$ -category (7.2).

### 5.1 Collaring movies of deformation embeddings

**Notation 5.1.** For  $A$  a set, we let the  $A$ -fold product  $[0, 1]^A$  denote as usual the collection of functions from  $A$  to  $[0, 1]$ . When  $A$  is a finite set, this collection is naturally a topological space, and in fact, a smooth manifold with corners in the usual way. We will apply this notation when the base is any interval – e.g.,  $[0, \epsilon)^A$ .

We caution that  $\Delta^I$ , for  $I$  a finite, non-empty, linearly ordered set, is the  $I$ -simplex; it is not an  $I$ -fold product. (See Notation 4.1.)

**Remark 5.2.** Fix a  $\Delta^I$ -movie  $M_I \times T^*\Delta^I$  (Definition 2.43). Let  $A \subset I$  and let  $\Delta^A \subset \Delta^I$  be the corresponding face. Because the movie is assumed collared, we may write  $M_I \times T^*\Delta^I$  as  $(M_I \times T^*\Delta^A) \otimes T^*[-1, \epsilon_A]^{I \setminus A}$  in a neighborhood of  $M_I \times T^*\Delta^A$ .

This decomposition uniquely determines a Liouville structure on  $M_I \times T^*\Delta^A$  as well, and it follows that  $M_I \times T^*\Delta^A$  is a  $\Delta^A$ -movie.

In the following definition, all deformation embeddings are assumed to be bordered (Definition 2.32) and collared (Definition 2.40) so that we may safely speak of movies (Definition 2.43).

**Definition 5.3.** Fix an injective, order-preserving map  $a : A \rightarrow I$ , along with a  $\Delta^I$ -movie  $M_I \times T^*\Delta^I$  and a  $\Delta^A$ -movie  $M_A \times T^*\Delta^A$ . A sectorial embedding

$$\phi : (M_A \times T^*\Delta^A) \otimes T^*[-1, \epsilon]^{I \setminus A} \rightarrow M_I \times T^*\Delta^I \quad (5.1)$$

is called a *collaring movie* of deformation embeddings if

1. For some  $\epsilon_A$  with  $0 < \epsilon_A < \epsilon$ , there exists a map  $\tilde{y}$  such that  $\phi$  can be written as a composition

$$(M_A \times T^*\Delta^A) \otimes T^*[-1, \epsilon_A]^{I \setminus A} \xrightarrow{\tilde{y} \otimes \text{id}} (M_I \times T^*\Delta^A) \otimes T^*[-1, \epsilon_A]^{I \setminus A} \xrightarrow{\text{id}_{M_I} \times T^*\eta_a} M_I \times T^*\Delta^I$$

where  $\eta_a$  is the collaring from Choice 4.9 (restricted to  $[-1, \epsilon_A] \subset [-1, \epsilon]$ ), and  $T^*\eta_a$  is the induced smooth map on cotangent bundles. (Note that if  $\tilde{y}$  exists, it is unique.)

2.  $\tilde{y}$  is a movie associated to some family of deformation embeddings

$$\{y^s : M_A \rightarrow M_I\}_{s \in \Delta^A}. \quad (5.2)$$

(See Definition 2.46.)

3. Finally, we demand  $\tilde{y}$  itself is collared. This means that for any subset  $A' \subset A$  (with induced injection  $a' : A' \rightarrow I$ ), there exists some  $\epsilon_{A'}$  with  $0 < \epsilon_{A'} < \epsilon$  and a map

$$\tilde{y}_{a'} : M_A \times T^* \Delta^{A'} \rightarrow M_I \times T^* \Delta^{A'}$$

so that  $\tilde{y}$ —when restricted to a neighborhood of  $\Delta^{A'} \times [-1, \epsilon_{A'})^{A \setminus A'} \subset \Delta^A$ —can be written as a composition

$$\begin{aligned} (M_A \times T^* \Delta^{A'}) \otimes T^*[-1, \epsilon_{A'}]^{I \setminus A'} &\xrightarrow{\tilde{y}_{a'} \otimes \text{id}} (M_I \times T^* \Delta^{A'}) \otimes T^*[-1, \epsilon_{A'}]^{I \setminus A'} \\ &\downarrow \text{id}_{M_I} \times T^* \eta_{a'} \\ &M_I \times T^* \Delta^I. \end{aligned} \quad (5.3)$$

(Equivalently, the family  $\{y^s\}_{s \in \Delta^A}$  is collared near every face  $\Delta^{A'}$  of  $\Delta^A$ .)

**Remark 5.4.** Suppose that we are given a collaring movie of deformation embeddings (5.1), and that we further know that  $M_I \times T^* \Delta^I$  and  $M_A \times T^* \Delta^A$  are both movies of *compactly supported* deformations (Definition 2.43). Then  $\phi$  defines a continuous, smooth, and collared map from  $\Delta^A$  to the space of sectorial embeddings from  $M_A$  to  $M_I$  (Remark 2.47).

## 5.2 $\text{Liou}_\Delta^{\text{Def}}$ (as a semisimplicial set)

**Definition 5.5.** We define a semisimplicial set  $\text{Liou}_\Delta^{\text{Def}}$  as follows. An  $I$ -simplex of  $\text{Liou}_\Delta^{\text{Def}}$  is the data of:

- For every non-empty subset  $A \subset I$ , a  $\Delta^A$ -movie  $M_A \times T^* \Delta^A$ . (Note that  $A$  may equal  $I$ .)
- For every proper subset inclusion  $A' \subset A$  of non-empty subsets of  $I$ , a strict sectorial embedding

$$\phi_{A' \subset A} : M_{A'} \times T^* \Delta^{A'} \otimes T^*[-1, \epsilon_{A, A'}]^{A \setminus A'} \rightarrow M_A \times T^* \Delta^A$$

such that  $\phi_{A' \subset A}$  is a collaring movie of deformation embeddings (Definition 5.3).<sup>20</sup>

satisfying the following conditions.

(LD1) For every string of proper inclusions  $A'' \subset A' \subset A$ , we have

$$\phi_{A'' \subset A} = \phi_{A' \subset A} \circ (\phi_{A'' \subset A'} \times \text{id}_{T^*[-1, \epsilon_{A', A''}]^{A' \setminus A''}})$$

where both sides are defined.

(LD2) If  $\max A = \max A'$ , then the function  $M_{A'} \rightarrow M_A$  is a strict isomorphism of Liouville sectors. (Here, the function is the one determined by the collaring near the vertex  $\Delta^{\{\max A\}} \subset \Delta^{A'} \subset \Delta^A \subset \Delta^I$ . We also note that the Liouville forms on both domain and codomain are those determined by the  $\Delta$ -movie structures near the vertex  $\max A$ .)

**Remark 5.6.** Note that given an  $I$ -simplex of  $\text{Liou}_\Delta^{\text{Def}}$ , for all non-empty subsets  $A, A' \neq I$ , we have that  $\dim M_A = \dim M_{A'}$ . (By definition of dimension of a manifold, the empty manifold is a manifold of all dimensions.)

<sup>20</sup>Note that  $\phi_{A' \subset A}$  thus defines a family of smooth embeddings  $M_{A'} \rightarrow M_A$  parametrized by  $\Delta^{A'}$ . Also note that  $[-1, \epsilon]^{A \setminus A'}$  is a cube of dimension  $\#(A \setminus A')$  (Notation 5.1).

**Remark 5.7.** The collaring on our movies—and the collarings on all the simplices of  $\text{Liou}_\Delta^{\text{Def}}$ —guarantees that our movies are sectors, as we saw in Proposition 2.42. Collaring by cubes of the form  $[-1, \epsilon]^N$  for  $\epsilon > 0$  will serve a further purpose in proving Theorem C; namely, the construction in Remark 10.10 is justified by the fact our collarings take place on neighborhoods of the form  $[-1, \epsilon]^N$  for  $\epsilon > 0$ .

The face maps of  $\text{Liou}_\Delta^{\text{Def}}$  are what one would expect: Given an injection  $\iota : J \rightarrow I$  of linearly ordered sets and an  $I$ -simplex  $(\{M_A \times T^*\Delta^A\}_{\emptyset \neq A \subset I}, \{\phi_{A' \subset A}\})$ , the associated  $J$ -simplex is pulled back along  $\iota$ . More precisely, it is given by:

$$(\{M_{\iota(B)} \times T^*\Delta^B\}_{\emptyset \neq B \subset J}, \{\phi_{\iota(B') \subset \iota(B)}|_{M_{\iota(B)} \times T^*\Delta^B}\}).$$

**Remark 5.8.** Fix an element  $s \in \Delta^{A''}$ . Then the composition requirement (LD1) guarantees that

$$y_{A' \subset A}^s \circ y_{A'' \subset A'}^s = y_{A'' \subset A}^s.$$

Here,  $\{y_{A'' \subset A}^s : M_{A''} \rightarrow M_A\}_{s \in \Delta^{A''}}$  is the family of deformation embeddings determined by the collaring movies  $\phi$ . (See (5.3).) Note that in this family, the Liouville structures on  $M_A$  and  $M_{A''}$  are  $s$ -dependent.

### 5.3 A 1-simplex

Let us explicate the definition of a 1-simplex; see also Figure 5.9. In this section, we will use the symbol  $\epsilon = \epsilon_A$  to mean some positive real number, and not indicate the dependency on the choice of a particular set  $A$ .

Following the notation of Definition 5.5, we take  $I$  to be the poset  $[1] = \{0, 1\}$ . Recall that the standard notation  $\Delta^I$  in this case is often abbreviated to  $\Delta^1$  – that is,  $\Delta^{\{0,1\}}$  is the standard 1-simplex.

Then to give an  $I$ -simplex (i.e., a 1-simplex) is to give the data of

- Three Liouville sectors  $M_{\{0\}}$ ,  $M_{\{1\}}$ , and  $M_{\{0,1\}} \times T^*\Delta^{\{0,1\}}$ .
- A strict sectorial embedding

$$\phi_{\{0\} \subset \{0,1\}} : M_{\{0\}} \otimes T^*[-1, \epsilon]^{\{1\}} \rightarrow M_{\{0,1\}} \times T^*\Delta^{\{0,1\}}$$

and


- A strict sectorial embedding

$$\phi_{\{1\} \subset \{0,1\}} : M_{\{1\}} \otimes T^*[-1, \epsilon]^{\{0\}} \rightarrow M_{\{0,1\}} \times T^*\Delta^{\{0,1\}}.$$

.

These data must satisfy the following condition to be a 1-simplex in  $\text{Liou}_\Delta^{\text{Def}}$ :

- $M_{\{0,1\}} \times T^*\Delta^{\{0,1\}}$  must be a  $\Delta^{\{0,1\}}$ -movie (Definition 2.43). In particular, it must have a Liouville sector structure of the form  $\theta_0 \oplus pdq$  along “ $M_{\{0,1\}}$  times  $T^*$  of a neighborhood of the 0th vertex of  $\Delta^{\{0,1\}}$ ,” and likewise of the form  $\theta_1 \oplus pdq$  in a neighborhood of “ $M_{\{0,1\}}$  times  $T^*$  of a neighborhood of the 1st vertex of  $\Delta^{\{0,1\}}$ .” Note  $\theta_0$  and  $\theta_1$  are (not necessarily equal) Liouville structure on  $M_{\{0,1\}}$ .



**Figure 5.9.** The largest rectangle in this image represents  $M_{\{0,1\}} \times T^*\Delta^1$ , with the cotangent direction omitted. The horizontal direction represents the  $\Delta^1$  direction, and the vertical direction represents the  $M_{\{0,1\}}$  direction. We depict a copy of  $M_{\{0\}} \times [-1, \epsilon]_s$ , in light gray, embedding into  $M_{\{0,1\}} \times \Delta^1$ , where the  $M_{\{0\}}$  component is mapped independently of  $s$ . Likewise we depict a copy of  $M_{\{1\}} \times [-1, \epsilon]_s$  embedded into  $M_{\{0,1\}}$  where the  $M_{\{1\}}$  component is mapped independently of  $s$ ; that  $M_{\{1\}}$  fills the entire vertical component is meant to connote that  $M_{\{1\}}$  maps diffeomorphically to  $M_{\{0,1\}}$ . In contrast, that  $M_{\{0\}}$  occupies only a proper subset of the vertical direction shows that  $y_0$  (5.4) need not be a diffeomorphism.

- By the condition that each  $\phi$  be a collaring movie of deformation embeddings, we conclude that  $\phi_{\{1\} \subset \{0,1\}}$  is simply a direct product of functions

$$\phi_{\{i\} \subset \{0,1\}} = y_i \times T^*\eta \quad (5.4)$$

where  $y_i$  is some strict sectorial embedding from  $M_{\{i\}}$  to  $M_{\{0,1\}}$  (with  $M_{\{0,1\}}$  endowed with the Liouville structure  $\theta_i$ ) and  $\eta = \eta_{\{i\} \subset \{0,1\}}$  is a collaring we have already chosen (Choice 4.9).

- Because  $I = \{0, 1\}$ , there are no proper inclusions  $A'' \subset A' \subset A$ , so condition (LD1) is empty.
- By Condition (LD2), the map  $y_1 : M_{\{1\}} \rightarrow M_{\{0,1\}}$  must be a strict *isomorphism* (that is, a diffeomorphism respecting Liouville structures on the nose). In particular, using  $y_1$ , one may think of the movie  $M_{\{0,1\}} \times T^*\Delta^1$  as a movie of Liouville structures on  $M_{\{1\}}$ .

**Remark 5.10.** Thus, a 1-simplex in  $\text{Liou}_\Delta^{\text{Def}}$  may be informally thought of as the data of two sectors  $M_{\{0\}}$  and  $M_{\{1\}}$ , equipped with (i) a smooth codimension zero embedding  $M_{\{0\}} \hookrightarrow M_{\{1\}}$  and (ii) a (bordered and collared) deformation of the Liouville structure on  $M_{\{1\}}$  which renders the embedding a strict sectorial embedding.

## 5.4 A 2-simplex

Now we illustrate what it means to be a 2-simplex in  $\text{Liou}_\Delta^{\text{Def}}$ . We set  $I = [2] = \{0 < 1 < 2\}$ . As before, we use the symbol  $\epsilon = \epsilon_A$  to mean some positive real number, and not indicate the dependency on the choice of a particular set  $A$ .

The data of a 2-simplex of  $\text{Liou}_\Delta^{\text{Def}}$  involves

- Liouville sectors

$$\begin{aligned}
M_{\{0\}} &\cong M_{\{0\}} \times T^* \Delta^{\{0\}}, & M_{\{1\}} &\cong M_{\{1\}} \times T^* \Delta^{\{1\}}, & M_{\{2\}} &\cong M_{\{2\}} \times T^* \Delta^{\{2\}} \\
M_{\{0,1\}} &\times T^* \Delta^{\{0,1\}}, & M_{\{0,2\}} &\times T^* \Delta^{\{0,2\}}, & M_{\{1,2\}} &\times T^* \Delta^{\{1,2\}}, \\
M_{\{0,1,2\}} &\times T^* \Delta^{\{0,1,2\}}. & & & & 
\end{aligned} \tag{5.5}$$

Each of these are movies; for example,  $M_{\{0,2\}} \times T^* \Delta^{\{0,2\}}$  is a  $\Delta^{\{0,2\}}$ -movie, and in particular induces a Liouville structure on  $M_{\{0,2\}}$  near vertex 0 of  $\Delta^2$ , and a Liouville structure vertex 2 of  $\Delta^2$ . Likewise, the Liouville sector structure on  $M_{\{0,1,2\}} \times T^* \Delta^{\{0,1,2\}}$  is induced from some  $\Delta^{\{0,1,2\}}$ -parameter family of Liouville structures on  $M_{\{0,1,2\}}$ . Using the same reasoning as in Remark 5.10, one may think of  $M_{\{0,1,2\}} \times T^* \Delta^{\{0,1,2\}}$  as encoding a  $\Delta^2$ -family of (bordered and collared) deformations of the Liouville structure on  $M_{\{2\}}$ .

- For every cardinality 2 subset  $\{i, j\}$  of  $[2]$  (where  $i$  may be less than, or greater than,  $j$ ), a  $\Delta^{\{i,j\}}$ -simplex of  $\text{Liou}_\Delta^{\text{Def}}$ . Section 5.3 already described what data this entails – Liouville sectors of the form  $M_{\{i\}}$  and  $M_{\{i,j\}} \times T^* \Delta^{\{i,j\}}$ , along with strict sectorial embeddings

$$M_{\{i\}} \otimes T^*[-1, \epsilon]^{\{j\}} \rightarrow M_{\{i,j\}} \times T^* \Delta^{\{i,j\}} \leftarrow M_{\{j\}} \otimes T^*[-1, \epsilon]^{\{i\}}$$

of the form

$$y_{\{i\} \subset \{i,j\}} \times T^* \eta_{\{i\} \subset \{i,j\}}, \quad y_{\{j\} \subset \{i,j\}} \times T^* \eta_{\{j\} \subset \{i,j\}}$$

again for  $\eta$  given by collaring conventions (Choice 4.9). As an example, the manifold  $M_{\{1\}}$  is thus equipped with two sectorial embeddings

$$y_{\{1\} \subset \{1,2\}} : M_{\{1\}} \rightarrow M_{\{1,2\}} \quad \text{and} \quad y_{\{1\} \subset \{0,1\}} : M_{\{1\}} \rightarrow M_{\{0,1\}}.$$

- Again for every cardinality 2 subset  $\{i, j\}$  of  $[2]$ , a strict sectorial embedding


$$\phi_{\{i,j\} \subset [2]} : M_{\{i,j\}} \times T^* \Delta^{\{i,j\}} \otimes T^*[-1, \epsilon]^{[2] \setminus \{i,j\}} \rightarrow M_{\{0,1,2\}} \times T^* \Delta^{\{0,1,2\}}.$$

Note that the domain is isomorphic to  $M_{\{i,j\}} \times T^* \Delta^1 \otimes T^*[-1, \epsilon]$ . Further, Definition 5.5 demands that  $\phi_{\{i,j\} \subset [2]}$  be a collaring movie of deformation embeddings; in particular,  $\phi_{\{i,j\} \subset [2]}$  encodes a  $\Delta^{\{i,j\}}$ -parametrized family of smooth embeddings

$$\{y^s : M_{\{i,j\}} \rightarrow M_{\{0,1,2\}}\}_{s \in \Delta^{\{i,j\}}}.$$

See Figure 5.4.

These maps must satisfy the composition requirement (LD1) (so that, for example,  $\phi_{\{0\} \subset [2]}$  may be recovered from  $\phi_{\{0\} \subset [1]}$  and  $\phi_{\{1\} \subset [2]}$ ) and the max-localizing requirement (LD2) (so that, for example,  $\phi_{\{0,2\} \subset \{0,1,2\}}$  induces an isomorphism from  $M_{\{0,2\}}$  to  $M_{\{0,1,2\}}$ ).



**Figure 5.11.** Depicted is a drawing of a triangular prism modeling the manifold  $M_{\{0,1,2\}} \times T^*\Delta^2$ . The “base” directions (i.e., the directions of the triangle) indicate the directions of  $\Delta^2$ . The vertical direction depicts the  $M_{\{0,1,2\}}$  direction, and the cotangent directions of  $T^*\Delta^2$  are not indicated. The rectangular prism with the light grey top and bottom represents the image of  $\phi_{\{0\}\subset[2]}$ ; the image is isomorphic to a copy of  $M_{\{0\}} \times T^*(\Delta^1 \times [-1, \epsilon])$ . Likewise, the rectangular prism with the dark grey top and bottom represents the image of  $\phi_{\{1\}\subset[2]}$ . Finally, the curvy grey figure represents the image of  $\phi_{\{0,1\}\subset[2]}$  – i.e., the image of an embedding of  $M_{\{0,1\}} \times T^*(\Delta^1 \times [-1, \epsilon])$  into  $M_{\{0,1,2\}} \times T^*\Delta^2$ . The curviness connotes that  $\phi_{\{0,1\}\subset[2]}$  potentially parametrizes a non-constant  $\Delta^1$ -family of embeddings  $y^2$  of  $M_{\{0,1\}}$  into  $M_{\{0,1,2\}}$ .

## 5.5 $\text{Liou}_\Delta$ (as a semisimplicial set)

**Definition 5.12.** We define  $\text{Liou}_\Delta$  to be the subsemisimplicial set of  $\text{Liou}_\Delta^{\text{Def}}$  consisting only of those movies that are associated to compactly supported deformations.

**Remark 5.13.** The consequence of requiring that all movies in sight be compactly supported is that the maps  $\{\tilde{y}_s\}_{s \in \Delta^A}$  are now families of sectorial embeddings for a *fixed* Liouville structure on  $M$  and  $N$  (because any two of  $\lambda_s, \lambda_{s'}$  are equivalent). See Remark 2.47.

**Remark 5.14.** The simplices of  $\text{Liou}_\Delta$  will thus have concretely interpretable geometric data. As we will see,

- A 1-simplex in  $\text{Liou}_\Delta$  will give rise to the data of a (not necessarily strict) sectorial embedding from  $M_0$  to  $M_1$  (Example 5.18).
- A 2-simplex will give rise to a diagram of sectorial embeddings commuting up to isotopy through sectorial embeddings (Proposition 5.19).

Conversely,

- Any (not necessarily strict) sectorial embedding from  $M$  to  $N$  gives rise to a (non-unique, but determined by contractible data) 1-simplex with initial vertex  $M$  and terminal vertex  $N$  (Construction 7.13).
- Any isotopy through sectorial embeddings  $f, g : M \rightarrow N$  will give rise to a (non-unique, but determined up to contractible data) 2-simplex (Example 7.17).

There is a similar interpretation of higher simplices; this is encapsulated in Theorem 8.1.

We remark that the “converse” constructions are postponed until Section 7.3, as we will not have established that  $\text{Liou}_\Delta$  admits degeneracy maps until then. The delay is worth it: Having degeneracy maps available will allow us to articulate what we mean by homotopies of morphisms in  $\text{Liou}_\Delta$  precisely. Because the converse constructions require choices, we wait for the availability of homotopies to be able to assuage the reader that these choices are not consequential (up to homotopy).

**Remark 5.15.** Fix an  $I$ -simplex of  $\text{Liou}_\Delta$  and fix an element  $a \in I$  along with a subset  $A \subset I$ ,  $a \in A$ . Then the inclusion  $\{a\} \subset A$  gives the data of a map

$$\phi_{\{a\} \subset A} : M_{\{a\}} \otimes T^*[-1, \epsilon]^A \rightarrow M_A \times T^*\Delta^A.$$

By the collaring assumption, this in fact determines a smooth embedding

$$y_{\{a\} \subset A} : M_{\{a\}} \rightarrow M_A \tag{5.6}$$

and this is a strict sectorial embedding for the Liouville structure on  $M_A$  determined near the vertex  $\Delta^{\{a\}} \subset \Delta^A$ .

**Remark 5.16.** When further  $a = \max A$ , Condition (LD2) states that  $y_{\{a\} \subset A}$  is a Liouville isomorphism:

$$y_{\{\max A\} \subset A} : M_{\{\max A\}} \xrightarrow{\cong} M_A \tag{5.7}$$

and this isomorphism is strict if  $M_A$  is endowed with this Liouville structure determined near the vertex  $\Delta^{\{a\}} \subset \Delta^A$ .

For this reason, when we write  $M_A$ , we will often consider the manifold  $M_A$  to be equipped with the Liouville structure determined near the vertex  $\Delta^{\max A}$  of  $\Delta^A$ .



**Remark 5.17.** Given the data of all the families  $\{\tilde{y}_{A' \subset A}^s\}_{s \in \Delta^{A'}}$ , and of Liouville structure on each  $M_A$  for  $A \subset I$ , the data of an  $I$ -simplex of  $\text{Liou}_\Delta$  is determined completely by the additional data of compactly supported smooth functions  $h_A : M_A \times \Delta^A \rightarrow \mathbb{R}$ .

Because the space of  $h_A$  is contractible, one should imagine that the families  $\{\tilde{y}_{A' \subset A}^s\}_{s \in \Delta^{A'}}$  are the homotopically relevant “meat” of the information conveyed by a simplex of  $\text{Liou}_\Delta$ .

**Example 5.18.** Fix an  $I$ -simplex of  $\text{Liou}_\Delta$  with  $|I| \geq 2$ , and fix two elements  $a < b$  in  $I$ . Let us study the associated 1-simplex of  $\text{Liou}_\Delta$ . We have the data of maps

$$M_{\{a\}} \otimes T^*[-1, \epsilon_a] \rightarrow M_{\{a,b\}} \times T^*\Delta^{\{a,b\}} \leftarrow M_{\{b\}} \otimes T^*[-1, \epsilon_b].$$

The main difference with a 1-simplex in  $\text{Liou}_\Delta^{\text{Def}}$  (as studied in Section 5.3) is that the movie  $M_{\{a,b\}} \times T^*\Delta^{\{a,b\}}$  is a movie through *compactly supported* deformations of the Liouville structure.

These maps in turn determine (not necessarily strict) sectorial embeddings

$$y_{\{a\} \subset \{a,b\}} : M_{\{a\}} \rightarrow M_{\{a,b\}}, \quad y_{\{b\} \subset \{a,b\}} : M_{\{b\}} \rightarrow M_{\{a,b\}}$$

as in (5.6). In fact, by endowing  $M_{\{a,b\}}$  with the Liouville structure determined near the vertex  $\Delta^{\{b\}} \subset \Delta^{\{a,b\}}$  (Remark 5.16), the map from  $M_{\{b\}}$  is a strict Liouville isomorphism as in (5.7). By composing with the inverse, we thus obtain a (not necessarily strict) sectorial embedding

$$y_{a < b} = (y_{\{b\} \subset \{a,b\}})^{-1} \circ y_{\{a\} \subset \{a,b\}} : M_{\{a\}} \rightarrow M_{\{b\}}. \quad (5.8)$$

Thus, though  $\text{Liou}_\Delta$  has diagrams built out of strict sectorial embeddings, we see that we can detect non-strict sectorial embeddings as edges in  $\text{Liou}_\Delta$ .

Here is the first hint that  $\text{Liou}_\Delta$  detects the topology of Liouville embedding spaces:

**Proposition 5.19.** Fix a (semisimplicial) 2-simplex  $\Delta^2 \rightarrow \text{Liou}_\Delta$ . Letting 0, 1, 2 denote the vertices of  $\Delta^2$ , consider the (not necessarily strict) sectorial embeddings  $y_{0 < 1}, y_{1 < 2}, y_{0 < 2}$  as defined in (5.8).

Then the composition  $y_{1 < 2} \circ y_{0 < 1}$  is isotopic, through (not necessarily strict) sectorial embeddings, to  $y_{0 < 2}$ .

*Proof.* The 2-simplex gives rise to the following commutative diagram<sup>21</sup> of smooth maps (none of which are codimension zero embeddings):

$$\begin{array}{ccccc}
M_{\{0\}} \times \Delta^{\{0\}} & \xrightarrow{y_{\{0\} \subset \{0,1\}} \times \iota} & M_{\{0,1\}} \times \Delta^{\{0,1\}} & \xleftarrow{y_{\{1\} \subset \{0,1\}} \times \iota} & M_{\{1\}} \times \Delta^{\{1\}} \\
\downarrow y_{\{0\} \subset \{0,2\}} \times \iota & & \downarrow (y_{\{0,1\} \subset \{0,1,2\}}^s, s) & & \downarrow y_{\{1\} \subset \{1,2\}} \times \iota \\
& & M_{\{0,1,2\}} \times \Delta^{\{0,1,2\}} & \xleftarrow{(y_{\{1,2\} \subset \{0,1,2\}}^s, s)} & M_{\{1,2\}} \times \Delta^{\{1,2\}} \\
& \nearrow (y_{\{0,2\} \subset \{0,1,2\}}^s, s) & & & \uparrow y_{\{2\} \subset \{1,2\}} \times \iota \\
M_{\{0,2\}} \times \Delta^{\{0,2\}} & \xleftarrow{y_{\{2\} \subset \{0,2\}} \times \iota} & & & M_{\{2\}} \times \Delta^{\{2\}}
\end{array}$$

where we have used the notation from (5.6) and (5.2), and the  $\iota$  are the obvious inclusions of faces of simplices. The diagonal map takes an element  $(x, s) \in M_{\{0,2\}} \times \Delta^{\{0,2\}}$  and maps it to the element  $(y_{\{0,2\} \subset \{0,1,2\}}^s(x), s) \in M_{\{0,1,2\}} \times \Delta^{\{0,1,2\}}$ .

<sup>21</sup>Though we have drawn the diagram rectilinearly, the reader may reshape the diagram to witness that the diagram has the shape of the barycentric subdivision of the 2-simplex.

Moreover, by the definition of a simplex in  $\text{Liou}_\Delta$ , we know that the  $y^s$  form smooth families of (not necessarily strict) sectorial embeddings. So, for example, the middle vertical map encodes an isotopy

$$\{y_{\{0,1\} \subset \{0,1,2\}}^s : M_{\{0,1\}} \rightarrow M_{\{0,1,2\}}\}_{s \in \Delta^{\{0,1\}}}$$

from the sectorial embedding encoded at the vertex  $s = \Delta^{\{0\}}$  to the sectorial embedding encoded at the vertex  $s = \Delta^{\{1\}}$ . In the equations that follow, we will abuse notation and write  $s = 0, 1, 2$  to mean the vertices  $\Delta^{\{0\}}, \Delta^{\{1\}}, \Delta^{\{2\}}$  of the 2-simplex  $\Delta^{\{0,1,2\}}$ . We also let  $\sim$  mean “is smoothly isotopic to, through (not necessarily strict) sectorial embeddings.” Then we have:

$$\begin{aligned} y_{\{1,2\} \subset \{0,1,2\}}^{s=2} \circ y_{\{2\} \subset \{1,2\}} \circ y_{1 < 2} \circ y_{0 < 1} &\sim y_{\{1,2\} \subset \{0,1,2\}}^{s=1} \circ y_{\{2\} \subset \{1,2\}} \circ y_{1 < 2} \circ y_{0 < 1} \\ &= y_{\{1,2\} \subset \{0,1,2\}}^{s=1} \circ y_{\{1\} \subset \{1,2\}} \circ y_{0 < 1} \\ &= y_{\{0,1\} \subset \{0,1,2\}}^{s=1} \circ y_{\{1\} \subset \{0,1\}} \circ (y_{\{1\} \subset \{0,1\}})^{-1} \circ y_{\{0\} \subset \{0,1\}} \\ &= y_{\{0,1\} \subset \{0,1,2\}}^{s=1} \circ y_{\{0\} \subset \{0,1\}} \\ &\sim y_{\{0,1\} \subset \{0,1,2\}}^{s=0} \circ y_{\{0\} \subset \{0,1\}} \\ &= y_{\{0,2\} \subset \{0,1,2\}}^{s=0} \circ y_{\{0\} \subset \{0,2\}} \\ &\sim y_{\{0,2\} \subset \{0,1,2\}}^{s=2} \circ y_{\{0\} \subset \{0,2\}}. \end{aligned}$$

We are finished by post-composing with the inverse diffeomorphism to

$$y_{\{1,2\} \subset \{0,1,2\}}^{s=2} \circ y_{\{2\} \subset \{1,2\}} = y_{\{0,2\} \subset \{0,1,2\}}^{s=2} \circ y_{\{2\} \subset \{0,2\}}.$$

□

## 5.6 Change of bases to max-constant simplices

**Definition 5.20** (Max-constant simplices). Fix an  $I$ -simplex of  $\text{Liou}_\Delta$ . Let us say that the simplex is *max-constant* (as opposed to just max-localizing) if for every subset  $A \subset I$ ,

1.  $M_A = M_{\{\max A\}}$ , and
2. The sectorial isomorphism  $y_{\{\max A\} \subset A} : M_{\{\max A\}} \rightarrow M_A$ —see (5.7)—is the identity map.

There is a natural way to “change bases” to take the data of an  $I$ -simplex of  $\text{Liou}_\Delta$  and render it a max-constant simplex. Given a chain of subsets  $\emptyset \neq A \subset B \subset I$ , consider the map

$$y_{A \subset B} : \Delta^A \rightarrow \text{Emb}_{\text{Liou}}(M_A, M_B)$$

given by the original  $I$ -simplex. This induces a map

$$y'_{A \subset B} : \Delta^A \rightarrow \text{Emb}_{\text{Liou}}(M_{\max A}, M_{\max B})$$

by conjugating by the isomorphisms  $M_{\max A} \cong M_A$  and  $M_B \cong M_{\max B}$  determined by the original  $I$ -simplex. It is straightforward to verify that the collection of maps  $\{y'_{A \subset B}\}_{\emptyset \neq A \subset B \subset I}$  defines a max-constant  $I$ -simplex.

This change of basis has the advantage that a collection of maps  $M_A \rightarrow M_B$  and  $M_{A'} \rightarrow M_{B'}$  may—if  $\max A = \max A'$  and  $\max B = \max B'$ —be considered as a collection of maps  $M_{\max A} \rightarrow M_{\max B}$  in a single mapping space (i.e., with fixed domain and codomain). We will utilize this when proving that  $\text{Liou}_\Delta$  satisfies the weak Kan condition (see Construction 6.18, for example).

## 6 The weak Kan condition

Our goal is to now prove:

**Theorem 6.1.**  $\text{Liou}_\Delta$  satisfies the (semisimplicial) weak Kan condition.

We will, in parallel, prove also the following:

**Theorem 6.2.**  $\text{Liou}_\Delta^{\text{Def}}$  satisfies the (semisimplicial) weak Kan condition.

**Remark 6.3.** The underlying philosophy of establishing the weak Kan property for both semisimplicial sets is identical, but there are technical differences. For  $\text{Liou}_\Delta$ , the only deformations appearing in movies are compactly supported, so all the technical difficulty is in constructing particular families of sectorial embeddings. (The construction of the compactly supported deformations is simple because the space of compactly supported functions is contractible.) For  $\text{Liou}_\Delta^{\text{Def}}$ , horn-filling further requires us to extend families of (collared, bordered, but otherwise arbitrary) Liouville structures from an image of an isotopy to the rest of the codomain. Here, we make extensive use of shrinking isotopies (Section 2.4). These allow us to rearrange families to only alter the Liouville structures away from the boundary of the image, and hence to extend these families to the entire codomain.

**Notation 6.4** ( $\alpha$  and  $\tilde{\alpha}$ ). So fix a map

$$\alpha : \Lambda_j^n \rightarrow \text{Liou}_\Delta \quad (\text{or } \text{Liou}_\Delta^{\text{Def}})$$

of semisimplicial sets, with  $0 < j < n$ . We must extend  $\alpha$  to a map

$$\tilde{\alpha} : \Delta^n \rightarrow \text{Liou}_\Delta \quad (\text{or } \text{Liou}_\Delta^{\text{Def}})$$

of semisimplicial sets.

**Notation 6.5** ( $\widetilde{\text{Emb}_{\text{Liou}}}$ ). By the compact-support condition on movies, one may interpret the data of an  $I$ -simplex in  $\text{Liou}_\Delta$  as giving rise to maps

$$\Delta^B \rightarrow \text{Emb}_{\text{Liou}}(M_B, M_A)$$

for each  $B \subset A \subset I$  (Remark 5.4). This is a map to the space of all sectorial embeddings. Because the identity function defines a sectorial isomorphism  $(M_A, \lambda_s^A) \rightarrow (M_A, \lambda_{s'}^A)$  for any  $s, s' \in \Delta^B$  (and likewise for  $M_B$ ), this space is independent of  $s$ .

Let us now define a space  $\widetilde{\text{Emb}_{\text{Liou}}}(M_B, M_A)$  as the total space of a fibration

$$\widetilde{\text{Emb}_{\text{Liou}}}(M_B, M_A) \rightarrow \Delta^B \tag{6.1}$$

whose fiber above  $s \in \Delta^B$  is the space  $\text{Emb}_{\text{Liou}}((M_B, \lambda_s^B), (M_A, \lambda_s^A))$ .

An  $I$ -simplex in  $\text{Liou}_\Delta^{\text{Def}}$  defines a section of this fibration.

**Remark 6.6.** Thus, to produce the simplices in  $\text{Liou}_\Delta^{\text{Def}}$ , we will have to produce maps from  $\Delta^B$  to  $\widetilde{\text{Emb}_{\text{Liou}}}$ , while to produce simplices in  $\text{Liou}_\Delta$ , we must also produce such maps, but we may identify the image with  $\text{Emb}_{\text{Liou}}$ .

In our arguments below, to preserve generality, we will thus often speak of maps to  $\widetilde{\text{Emb}_{\text{Liou}}}$ , with the understanding that all such maps are sections of the fibration (6.1).

**Notation 6.7** (The values of  $\alpha$  and  $\tilde{\alpha}$ ). To give  $\tilde{\alpha}$ , we must make explicit  $\tilde{\alpha}(A)$ —that is, we must specify the movies  $M_A \times T^*\Delta^A$ —for every  $A \subset [n]$ . Of course, given  $\alpha$ , this amounts to specifying the movies  $\tilde{\alpha}([n])$  and  $\tilde{\alpha}([n] \setminus \{j\})$ .

Moreover, for every  $B \subset A$ , we must specify collaring movies  $\phi_{B \subset A}$ . By Remark 5.4, the data of such collaring movies is equivalent to the data of a function from  $\Delta^B$  to the space of smooth function from  $M_B$  to  $M_A$  (where each function at  $s \in \Delta^B$  is a sectorial embedding with respect to the Liouville structures on  $M_B$  and  $M_A$  at  $s$ ). We will denote this smooth function by  $\tilde{\alpha}(B \subset A)$ .

Likewise, for every  $B \subset A$  with  $\Delta^A$  a face of the horn  $\Lambda_j^n$ ,  $\alpha$  defines a map

$$\alpha(B \subset A) : \Delta^B \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_B, M_A)(M_B, M_A).$$

## 6.1 For $n = 2$

We begin with the case  $n = 2$  (and hence  $j = 1$ ). Unlike the higher- $n$  case, we here require no isotopy extension arguments. We begin with the  $\text{Liou}_\Delta$  case.

**Construction 6.8** ( $\tilde{\alpha}$  when  $n = 2$ , for  $\text{Liou}_\Delta$ ). The data of  $\alpha : \Lambda_1^2 \rightarrow \text{Liou}_\Delta$  determine smooth, proper codimension zero embeddings

$$M_{\{0\}} \rightarrow M_{\{0,1\}} \xleftarrow{\cong} M_{\{1\}} \rightarrow M_{\{1,2\}} \xleftarrow{\cong} M_{\{2\}} \quad (6.2)$$

Each  $M_{\{i,j\}}$  is equipped with a compactly supported deformation of Liouville structures; when given the Liouville structure near  $\max\{i, j\}$ , the leftward pointing arrows are isomorphisms of Liouville sectors as in (5.7). To fill  $\alpha$ , let us declare

$$M_{\{0,1,2\}} = M_{\{2\}} \quad (6.3)$$

and define the smooth maps

$$y_{\{0,1\} \subset \{0,1,2\}}^s : M_{\{0,1\}} \rightarrow M_{\{0,1,2\}}, \quad s \in \Delta^{\{0,1\}}$$

as the composition

$$M_{\{0,1\}} \xrightarrow{\cong} M_1 \rightarrow M_{\{1,2\}} \xrightarrow{\cong} M_{\{2\}} = M_{\{0,1,2\}}.$$

(The rightward pointing isomorphisms are inverses of the isomorphisms in (6.2). Note also that this is independent of  $s \in \Delta^{\{0,1\}}$ .) We would like to extend this to a strict sectorial embedding

$$M_{\{0,1\}} \times T^*\Delta^{\{0,1\}} \rightarrow M_{\{0,1,2\}} \times T^*\Delta^{\{0,1\}} \quad (6.4)$$

where the domain of (6.4) is given the Liouville structure specified by  $\alpha$  for the  $\Delta$ -movie  $M_{\{1,2\}} \times T^*\Delta^{\{1,2\}}$ , and we must specify the Liouville structure for the codomain. To do this, we note that because the map  $y_{\{0,1\} \subset \{0,1,2\}}^s$  for  $s = \Delta^{\{1\}}$  is a Liouville isomorphism, one can choose a  $\Delta^{\{0,1\}}$ -dependent family of compactly supported functions  $\tilde{h}_s$  on  $M_{\{0,1,2\}}$

$$\tilde{h} : M_{\{0,1,2\}} \times \Delta^{\{0,1\}} \rightarrow \mathbb{R}, \quad (x, s) \mapsto \tilde{h}_s(x) \quad (6.5)$$

so that, endowing  $M_{\{0,1,2\}} = M_{\{2\}}$  with the Liouville structure  $\lambda^{M_{\{2\}}} + d\tilde{h}_s$ —and giving  $M_{\{0,1,2\}} \times T^*\Delta^{\{0,1\}}$  the induced Liouville structure—(6.4) becomes a sectorial embedding.

Likewise, we define an  $s$ -independent family of smooth maps

$$y_{\{1,2\} \subset \{0,1,2\}}^s : M_{\{1,2\}} \rightarrow M_{\{0,1,2\}}, \quad s \in \Delta^{\{1,2\}}$$

by the composition  $M_{\{1,2\}} \xrightarrow{\cong} M_{\{2\}} = M_{\{0,1,2\}}$ . As before, by specifying an appropriate family of compactly supported functions  $\tilde{h}_s : M_{\{2\}} \rightarrow \mathbb{R}$  for  $t \in \Delta^{\{1,2\}}$

$$\tilde{h} : M_{\{0,1,2\}} \times \Delta^{\{1,2\}} \rightarrow \mathbb{R}, \quad (x, s) \mapsto \tilde{h}_s(x) \quad (6.6)$$

we obtain a sectorial embedding

$$M_{\{1,2\}} \times T^* \Delta^{\{1,2\}} \rightarrow M_{\{0,1,2\}} \times T^* \Delta^{\{1,2\}}. \quad (6.7)$$

Now, because all the structures given by  $\alpha$  are collared, one can smoothly extend the function

$$\tilde{h} : M \times (\Delta^{\{0,1\}} \bigcup_{\Delta^{\{1\}}} \Delta^{\{1,2\}}) \rightarrow \mathbb{R}$$

(which is obtained by gluing together (6.5) and (6.6)) to a function

$$\tilde{h} : M \times \Delta^{\{0,1,2\}} \rightarrow \mathbb{R}, \quad (x, s) \mapsto \tilde{h}_s(x), \quad (s \in \Delta^{\{0,1,2\}}) \quad (6.8)$$

for which  $\tilde{h}$  is collared, and for which each  $\tilde{h}_s$  is compactly supported. We then declare the  $\Delta$ -movie

$$M_{\{0,1,2\}} \times T^* \Delta^{\{0,1,2\}} := M_{\{2\}} \times T^* \Delta^{\{0,1,2\}} \quad (6.9)$$

to be given the Liouville structure induced by adding  $d$  of (6.8) to the Liouville structure of  $M_{\{2\}}$ .

(6.9) defines the value of  $\tilde{\alpha}$  on  $[2] = \{0, 1, 2\}$ . It is also clear that the maps (6.4) and (6.7) extend, by collaring, to specify maps  $\phi_{\{1,2\} \subset \{0,1,2\}}$  and  $\phi_{\{0,1\} \subset \{0,1,2\}}$ . It remains to define  $\tilde{\alpha}$  on  $\{0, 2\}$  and the associated  $\phi$ .

- We declare (Notation 6.7)

$$\tilde{\alpha}(\{0, 2\}) = M_{\{0,2\}} \times T^* \Delta^{\{0,2\}} := M_{\{2\}} \times T^* \Delta^{\{0,2\}} \quad (6.10)$$

with the Liouville form inherited from restricting the form on (6.9) along the edge  $\Delta^{\{0,2\}} \subset \Delta^{\{0,1,2\}}$ . (This restriction is well-defined thanks to the collared choice of (6.8).)

- We declare (see Notation 6.7)  $\tilde{\alpha}(\{2\} \subset \{0, 2\})$  to be (extended from) the identity map  $M_{\{2\}} \rightarrow M_{\{2\}}$ . Note that this is a sectorial embedding because by the collaring of  $\tilde{h}$ , the Liouville form on  $M_{\{0,1,2\}} \times T^* \Delta^2$  agrees with that of  $M_{\{2\}}$  near the vertex  $\Delta^{\{2\}}$ .
- We declare  $\tilde{\alpha}(\{0\} \subset \{0, 2\})$  to be (extended from) the composition of smooth maps

$$M_{\{0\}} \rightarrow M_{\{0,1\}} \xrightarrow{\cong} M_{\{1\}} \rightarrow M_{\{1,2\}} \xrightarrow{\cong} M_{\{2\}} = M_{\{0,2\}}.$$

(Every map here was already given by  $\alpha$ .) By design of  $\tilde{h}$  (6.8) and by our definition of the Liouville structure on (6.10), this is a strict sectorial embedding.

We have defined  $\tilde{\alpha}$ . The data satisfy (LD1) and (LD2) by construction, and clearly extend  $\alpha$ .

**Remark 6.9.** The above construction fails when  $j = 0$  and  $n = 2$  (and likewise for  $j = 2$ ). To see this, fix three sectors  $M_0, M_1, M_2$  and strict sectorial embeddings  $y_{0<2} : M_0 \rightarrow M_2, y_{1<2} : M_1 \rightarrow M_2$  for which  $y_{0<2}$  is not isotopic to any map factoring through  $y_{1<2}$ . (This defines a horn by setting, for example,  $M_{\{i,j\}} \times T^* \Delta^{\{i,j\}} := M_{\{\max\{i,j\}\}} \otimes T^* \Delta^{\{i,j\}}$ .)

Proposition 5.19 guarantees that this horn cannot be filled.

**Construction 6.10** ( $\tilde{\alpha}$  when  $n = 2$ , for  $\text{Liou}_\Delta^{\text{Def}}$ ). We follow the notations of (6.2) and (6.3). Then, by pulling back along under the isomorphism  $M_{\{2\}} \xrightarrow{\cong} M_{\{1,2\}}$ , one obtained a collared family of deformations on  $M_{\{0,1,2\}}$  along the edge  $\Delta^{\{1,2\}}$ . We must now extend this collared deformation to all of  $\Delta^{\{0,1,2\}}$ .

We begin by defining it along the edge  $\Delta^{\{0,1\}}$ . Here, we are already given a deformation of  $M_{\{0,1\}} \cong M_{\{1\}}$ , and  $M_{\{1\}}$  is equipped with a strict sectorial embedding into  $M_{\{0,1,2\}}$  (strict with respect to the Liouville structure of  $M_{\{0,1,2\}}$  at  $s = \Delta^{\{1\}}$ ). By applying a shrinking isotopy to  $M_{\{1\}}$  (and hence to its image inside  $M_{\{0,1,2\}}$ ) as necessary, we obtain a family of embeddings of  $M_{\{1\}} \cong M_{\{0,1\}}$  into  $M_{\{0,1,2\}}$ , parametrized by  $s \in \Delta^{\{0,1\}}$ . The deformations of Liouville structures on  $M_{\{0,1\}}$  pushforward, and these pushforwards may be extended to become deformations of Liouville structures on  $M_{\{0,1,2\}}$  again thanks to the shrinking. (See Remark 6.11.)

So we have defined a deformation of Liouville structures for  $M_{\{0,1,2\}}$  parametrized by the horn  $\Delta^{\{0,1\}} \cup \Delta^{\{1,2\}}$ . Of course the space of deformations is a space, so we may continuously extend this to all of  $\Delta^{\{0,1,2\}}$ ; the usual smooth-approximation trick applies as in Section 4.4 (because the space of deformations is locally modeled by the space of (very small) exact 1-forms), and we may thus ensure that we have a smooth, collared, and continuous  $\Delta^{\{0,1,2\}}$ -parametrized deformation of Liouville structures on  $M_{\{0,1,2\}}$ .

Now it is straightforward to define the embeddings  $M_{\{0\}} \rightarrow M_{\{0,1,2\}}$  at  $s = \Delta^{\{0\}}$ , and declaring  $M_{\{0,2\}} = M_{\{0,1,2\}}$  and restricting the family of Liouville structures on  $M_{\{0,1,2\}}$  to one on  $M_{\{0,2\}}$  along the edge  $\Delta^{\{0,2\}}$ .

**Remark 6.11.** Let us elaborate on how to extend families of deformations from a subsector to a large subsector, after a suitable shrinking isotopy of the subsector. We explain in the context of Construction 6.10, but the remarks here apply more widely.

Fix the image  $A$  of  $M_{\{1\}}$  inside  $M_{\{0,1,2\}}$ , induced by the inclusion  $M_{\{1\}} \hookrightarrow M_{\{1,2\}}$  at  $s = \Delta^{\{1\}}$ . We may identify  $A$  with  $M_{\{0,1\}}$  due to the isomorphism  $M_{\{1\}} \cong M_{\{0,1\}}$  at  $s = \Delta^{\{1\}}$ . For every  $s$  in the edge  $\Delta^{\{0,1\}}$ , we have a Liouville structure  $\lambda_s$  on  $M_{\{0,1\}}$ , which is collared by assumption. This induces a new Liouville structure on  $A$ , by choosing a shrinking of  $A$  (Section 2.4) to have image  $A' \subset A$ , and defining a Liouville structure  $\lambda_s^A$  on  $A$  which equals the pushforward of  $\lambda_s$  onto  $A'$ , and parametrizes between  $\lambda_s$  and  $\lambda_{\Delta^{\{1\}}}$  along the region  $A \setminus A'$ . Now, because  $\lambda_s^A$  agrees with  $\lambda^{M_{\{0,1,2\}}}$  in a neighborhood of  $\partial A$ , we may extend  $\lambda_s^A$  to equal  $\lambda_{\{0,1,2\}}^M$  outside of  $A$ .

## 6.2 For $n \geq 3$ , filling most of the simplex

Fix  $\alpha : \Lambda_j^n \rightarrow \text{Liou}_\Delta$  (or to  $\text{Liou}_\Delta^{\text{Def}}$ ) as in Notation 6.4. Roughly speaking, the construction of  $\tilde{\alpha}$  may be divided into two steps: (a) Defining the maps  $\tilde{\alpha}(A \subset [n])$  for those  $A$  with  $\max A = n$ , then (b) Defining the map  $\tilde{\alpha}(A \subset [n])$  for  $A = [n-1]$ . We establish (a) in the present section, using a strategy illustrated in Table 2 for the case of  $n = 3$  and  $j = 1$ . We leave (b) to Sections 6.3 and 6.4.

**Construction 6.12** (Of  $\tilde{\alpha}$ , Part I). We set  $M_{[n]} := M_{\{n\}}$ .

**Remark 6.13.** More generally, we may choose a sectorial diffeomorphism from  $M_{\{n\}}$  to some other Liouville sector  $Y$ , and declare  $M_{[n]}$  to be given by  $Y$ . This freedom will be important for a later inductive step, so that in fact we can choose  $M_{[n]}$  to be a pre-given  $Y$ .

**Choice 6.14** (A family  $\sigma_{t,u}$  of shrinking isotopies). Consider the geometric realization  $|\Lambda_j^n|$  of the  $j$ th  $n$ -horn. We may up to homeomorphism write  $|\Lambda_j^n|$  as a cone/join

$$|\Lambda_j^n| \cong ([0, 1] \times \partial(\partial_j \Delta^n)) / \sim \cong |\Delta^{\{n\}}| \star \partial(\partial_j \Delta^n)$$




Table 2: Partially filling the horn  $\Lambda_1^3$  by accomplishing (a). The top row of images emphasizes the filling of the 3-simplex; the bottom row of images emphasizes the filling of the face opposite the vertex 1, using the decomposition of this face induced by projecting the horn. The subsets  $B''$  of Construction 6.16 are the edges containing the 3rd vertex. The subsets  $B'$  of Construction 6.17 are the faces of  $\Delta^3$  that are not the 1st face, nor the 0th face. Construction 6.18 defines a piecewise smooth map from the 1st face, then smooths it out to give a smooth map from the 1st face. After this, to complete the horn-filling, one need only (b) define data along the face  $\Delta^{\{0,1,2\}}$ —and more generally, on the face opposite the  $n$ th vertex.

from the  $n$ th vertex, to the boundary of the  $j$ th face. Writing the coordinates of  $[0, 1] \times \partial(\partial_j \Delta^n)$  as pairs  $(t, u)$ , we now choose a  $u$ -family of  $t$ -parametrized shrinking isotopies

$$\sigma_{t,u} : [0, 1] \times \partial(\partial_j \Delta^n) \rightarrow \text{Emb}_{\text{Liou}}(M_{[n]}, M_{[n]}) \quad (6.11)$$

satisfying the following:

1.  $\sigma_{t,u}$  is the identity for all small  $t$  (and hence descends to a map from the join),
2.  $\sigma_{t,u}$  is, along every top-dimensional face of  $|\Delta_j^n|$ , collared and smooth.
3. Finally, we put a restriction on the collaring region of  $\sigma_{t,u}$ . We demand that  $\sigma_{t,u}$  is collared (by the identity) whenever<sup>22</sup> the maps  $\tilde{y}^s$  determined by  $\alpha$  are collared near the vertex  $|\Delta^{\{n\}}|$ , but everywhere else,  $\sigma_{t,u} : M_{[n]} \rightarrow M_{[n]}$  has image bounded away from the sectorial boundary of  $M_{[n]}$ .

**Remark 6.15.** We make precise the collaring restriction 3 on  $\sigma_{t,u}$ . By definition,  $\alpha$  determines collared families  $\tilde{y}_{I'' \subset I'}^s$ , so we know that every pair  $I'' \subset I'$ —with  $n \in I'$  and  $I'$  determining some simplex of  $\Lambda_j^n$ —defines some region  $R_{I'' \subset I'} \subset |\Delta^n|$  (i) containing the  $n$ th vertex and (ii) along which  $\tilde{y}^s$  is independent of  $s \in R_{I'' \subset I'}$ . (Each region is open in the face  $\Delta^{I''}$ , but not necessarily in  $\Lambda_j^n$ .) Let  $R \subset |\Delta^n|$  denote the union of all these regions, where the union is indexed by pairs  $I'' \subset I'$ .

We know that the region  $R$  contains a subset  $R_{\min}$ , which is the union, indexed by  $A$  for which  $\Delta^{\{n\}} \subset \Delta^A \subset \Delta_j^n$ , of the regions  $[0, 1]^{I' \setminus A}$  under the image of the collarings  $\eta$ . (These  $A$ -dependent regions are each obtained by setting  $\epsilon_A = 0$  in (5.3).) This  $R_{\min}$  represents the closed

<sup>22</sup>See Remark 6.15.

subset containing the  $n$ th vertex along which everything in sight must be collared. We are then guaranteed that—because each  $\epsilon_A > 0$ — $R$  contains some open neighborhood  $U$  of  $R_{\min}$ . (This  $U$  is open and  $R$ , and may be chosen so that its closure is contained in the interior of  $R$ .)

So here is the precise statement: We demand that the family  $\sigma_{t,u}$  is constant along  $R_{\min}$ , and that for each  $(t,u)$  whose image is not in  $R$ ,  $\sigma_{t,u} : M_{[n]} \rightarrow M_{[n]}$  does not intersect the sectorial boundary of  $M_{[n]}$  (hence, by properness, has image bounded away from the sectorial boundary).

**Construction 6.16** (Of  $\tilde{\alpha}$ , Part II). Let  $B'' \subset [n]$  be a subset of cardinality  $n - 1$  containing  $n$ . We now construct maps

$$\tilde{\alpha}_{B'',B'} : \Delta^{B''} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]}) \quad (6.12)$$

for all  $B'' \subset B' \subset [n]$  with  $B'$  a top-dimensional simplex of  $\Lambda_j^n$ .<sup>23</sup> The goal is to choose (6.12) so that the composite

$$\Delta^{B''} \xrightarrow{\tilde{\alpha}_{B'',B'} \times \alpha(B'' \subset B')} \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]}) \times \widetilde{\text{Emb}}_{\text{Liou}}(M_{B''}, M_{B'}) \xrightarrow{\circ} \widetilde{\text{Emb}}_{\text{Liou}}(M_{B''}, M_{[n]}) \quad (6.13)$$

is independent of the choice of  $B'$ . (Given  $B''$ , there are at most two  $B'$  that fit into a string of proper inclusions  $B'' \subset B' \subset [n]$ .)

For this, consider the diffeomorphism

$$(\tilde{y}_{\{n\} \subset B'}^{\{n\}})^{-1} : M_{B'} \rightarrow M_{\{n\}} = M_{[n]}$$

determined by  $\alpha$ , and the composition

$$\begin{array}{ccc} \Delta^{B''} & \xrightarrow{\sigma_{t,u} \times (\tilde{y}_{\{n\} \subset B'}^{\{n\}})^{-1} \times \alpha(B'' \subset B')} & \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]}) \times \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]}) \times \widetilde{\text{Emb}}_{\text{Liou}}(M_{B''}, M_{B'}) \\ & \searrow & \downarrow \circ \\ & & \widetilde{\text{Emb}}_{\text{Liou}}(M_{B''}, M_{[n]}) \end{array} \quad (6.14)$$

where  $\sigma_{t,u}$  is the family of isotopies from Choice 6.14. We define this composition to be the value of  $\tilde{\alpha}$  on the inclusion  $B'' \subset [n]$ ; that is, this composition defines

$$\tilde{\alpha}(B'' \subset [n]).$$

(To make sense of this in the case of  $\text{Liou}_{\Delta}^{\text{Def}}$ , we must also of course construct deformations of Liouville structures on  $M_{[n]}$  parametrized by  $\Delta^{B''}$ . Once we have defined the maps  $\tilde{\alpha}(B'' \subset [n])$ , it is straightforward to construct such a deformation by extending from the images of  $M_{B''}$ . See Remark 6.11. On the other hand, in the case of  $\text{Liou}_{\Delta}$ , where everything is compactly supported, we have defined a map  $\Delta^{B''} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{B''}, M_{[n]})$  on the nose without any need for choices of deformations of the codomain.)

This is a  $\Delta^{B''}$ -family of (not necessarily strict) sectorial embeddings of  $M_{B''}$  into  $M_{[n]}$ , which near the  $n$ th vertex of  $\Delta^{B''}$  is constant, and is given by

$$(\tilde{y}_{\{n\} \subset B''}^{\{n\}})^{-1} : M_{B''} \rightarrow M_{\{n\}} = M_{[n]}.$$

By Isotopy Extension (Proposition 2.55), there also exists a family of maps  $\tilde{\alpha}_{B'',B'}$  as in (6.12) so that the composition (6.13) equals the map  $\tilde{\alpha}(B'' \subset [n])$  given by the composition in (6.14).

We fix such a choice. Of course, by the usual yoga, we may assume everything in sight is collared and smooth and continuous. To summarize, we have now defined  $\tilde{\alpha}$  on the inclusion  $B'' \subset [n]$  for every  $B'' \subset [n]$  of cardinality  $n - 1$  containing  $n$ ; and also defined maps as in (6.12).

<sup>23</sup>In other words,  $B'$  is a proper subset of  $[n]$  of cardinality  $n$ , and  $B' \cup \{j\} \neq [n]$ .



**Construction 6.17** (Of  $\tilde{\alpha}$ , Part III). Now fix  $B' \subset [n]$  to be a subset of cardinality  $n$  for which  $B' \cup \{j\} \neq [n]$  and  $B' \cup \{n\} \neq [n]$ . By Construction 6.16, we have a map

$$\bigcup_{B'' \subset B'} \tilde{\alpha}_{B'', B'} : \bigcup_{B'' \subset B'} \Delta^{B''} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]})$$

where the index runs over all  $B'' \subset B'$  for which  $B''$  has cardinality  $n-1$  and contains  $n$ . Recognizing the union of  $\Delta^{B''}$  as the horn  $\Lambda_n^{B'}$  obtained by deleting from  $\Delta^{B'}$  the face opposite the vertex  $\Delta^{\{n\}}$ , we thus have a map

$$\Lambda_n^{B'} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]}). \quad (6.15)$$

It is smooth, continuous, and collared one very simplex of  $\Lambda_n^{B'}$ .

Now, the space of Liouville structures on  $M_{[n]}$  is a space, and in particular, horns may be filled. Thus we may extend the  $\Lambda_n^{B'}$ -family of Liouville structures on  $M_{[n]}$  implicit in (6.15) to a simplex  $\Delta^{B'}$  worth of Liouville structures. As usual, we may choose this family to be collared, smooth, and continuous.

By the usual lifting properties of fibrations, we may extend (6.15) to a continuous map

$$\tilde{\alpha}(B' \subset [n]) : \Delta^{B'} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{B'}, M_{[n]}). \quad (6.16)$$

Then, altering this map if necessary, we may assume (6.16) is actually smooth and collared and continuous by Proposition 4.16.

To summarize, we have now defined  $\tilde{\alpha}$  on the inclusions  $B' \subset [n]$  for all facets  $\Delta^{B'} \subset \Delta^n$  for which  $\Delta^{B'} \neq \partial_n \Delta^n$ , and  $\Delta^{B'} \neq \partial_j \Delta^n$ .

**Construction 6.18** (Of  $\tilde{\alpha}$ , Part IV). We have a natural isomorphism of simplicial sets

$$\bigcup_{B'} \partial_n \Delta^{B'} \cong \Lambda_j^{n-1}.$$

The lefthand side's union is indexed over those  $B' \subset [n]$  for which  $B' \neq [n] \setminus \{j\}$ , and for which  $B' \neq [n-1]$ .  $\partial_n \Delta^{B'}$  is the simplicial set given by the face opposite the vertex  $\Delta^{\{n\}} \subset \Delta^{B'}$ . By Section 5.6, and because we know  $\max B' = n$  for all  $B'$ , we may change bases and assume that the maps (6.16) define sections

$$\Delta^{B'} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]}) \quad (6.17)$$

for all choices of  $B'$ . By construction, it follows that these maps agree along intersections of the various  $\Delta^{B'}$ , so they glue to give a single map

$$\bigcup_{B'} \partial_n \Delta^{B'} \cong \Lambda_j^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]}).$$

Again by extending the  $\Lambda_j^{n-1}$ -family of Liouville structures to a  $\Delta^{n-1}$ -family, then using the fact that  $\widetilde{\text{Emb}}_{\text{Liou}}$  is a Kan fibration over the space of Liouville structures, we may fill the horn to yield a map

$$\Delta^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]}) \quad (6.18)$$

where we have used Remark 6.11 again to know what we mean by Liouville structures on  $M_{[n]}$  over  $s \in \Delta^{n-1}$ . We may as usual assume everything in sight is smooth, collared, and continuous.

By gluing (6.18) and (6.17) we obtain a map

$$\left( \bigcup_{B'} \Delta^{B'} \right) \bigcup_{\Lambda_j^{n-1}} \Delta^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]}). \quad (6.19)$$

We now note that, at the level of geometric realizations, there is a piecewise smooth homeomorphism<sup>24</sup>

$$\left( \bigcup_{B'} |\Delta^{B'}| \right) \bigcup_{|\Lambda_j^{n-1}|} |\Delta^{n-1}| \cong |\partial_j \Delta^n|.$$

Thus, the map (6.19) induces a piecewise smooth map from  $|\partial_j \Delta^n|$  to  $\widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]})$ . Importantly, the map (6.19) is collared along all the boundaries and corners; thus, Proposition 4.16 allow us to continuously homotope the piecewise smooth map to an honest smooth map, and collared along the boundary of  $|\partial_j \Delta^n|$ . Thus, we have a map

$$\partial_j \Delta^n \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n]}, M_{[n]})$$

that is smooth and continuous and collared according to our collaring conventions. By undoing our change of basis, we obtain a map

$$\tilde{\alpha}([n] \setminus \{j\} \subset [n]) : \partial_j \Delta^n \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{[n] \setminus \{j\}}, M_{[n]}) \quad (6.20)$$

(End of Construction 6.18.)

To this point, we have constructed  $\tilde{\alpha}(B \subset [n])$  for all facets  $\Delta^B \subset \Delta^n$  that contain the vertex  $\Delta^{\{n\}}$ . It remains to define  $\tilde{\alpha}$  on the face opposite the  $n$ th vertex. Unraveling the definition of a simplex of  $\text{Liou}_{\Delta}^{\text{Def}}$ , and again assuming that both  $\alpha$  and  $\tilde{\alpha}$  as max-constant for simplicity (Section 5.6), this means we must produce a map

$$\tilde{\alpha}([n-1] \subset [n]) : \Delta^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}). \quad (6.21)$$

This will be the goal of Sections 6.3 and 6.4.

### 6.3 $n \geq 3$ and $j \neq n-1$

Assume  $j \neq n-1$ . Again we assume that all simplices of  $\alpha$  are max-constant (Section 5.6). Then  $\alpha$  assigns

1. to the inclusion  $[n-2] \subset [n] \setminus \{n-1\}$  a map

$$\alpha([n-2] \subset [n] \setminus \{n-1\}) : \Delta^{n-2} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-2\}}, M_{\{n\}})$$

(This is the point at which we are using that  $j \neq n-1$ .) By post-composing with (the restriction to  $\Delta^{n-2} \subset \Delta^{n-1}$  of) the map  $\tilde{\alpha}([n] \setminus \{n-1\} \subset [n])$ , we obtain a map

$$\Delta^{n-2} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-2\}}, M_{\{n\}}). \quad (6.22)$$

---

<sup>24</sup>This can be seen, for example, by projecting the horn  $\Lambda_j^n$  to the face  $|\partial_j \Delta^n|$  along a vector normal to  $|\partial_j \Delta^n|$  and passing through the vertex  $|\Delta^{\{j\}}|$ .

2. to the inclusion  $[n-2] \subset [n-1]$  a map

$$\alpha([n-2] \subset [n-1]) : \Delta^{n-2} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-2\}}, M_{\{n-1\}}), \quad (6.23)$$

3. and, for any  $A \subset [n]$  with  $\max A = n-1$ , and for any  $i \neq j$ , to the inclusion  $A \subset [n] \setminus \{i\}$  a map

$$\alpha(A \subset [n] \setminus \{i\}) : \Delta^A \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}).$$

By Construction 6.16, if we post-compose with  $\tilde{\alpha}([n] \setminus \{i\} \subset [n])$ , the induced map from  $\Delta^A$  to  $\widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}})$  is independent of the choice of  $i$ . In particular, by gluing together those  $A$  for which  $\Delta^{A \setminus \{n-1\}} \subset \partial \Delta^{n-2}$ , we obtain a map

$$\partial \Delta^{n-2} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}). \quad (6.24)$$

By construction, we are guaranteed that the composition of (6.23) with (6.24) equals the map (6.22) when restricted to  $\partial \Delta^{n-2}$ .

By isotopy extension, we may thus extend (6.24) to a map

$$\Delta^{n-2} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}). \quad (6.25)$$

Now, by again composing the data of (i)  $\alpha([n-1] \setminus \{i\} \subset [n] \setminus \{i\})$  for  $i \neq j, n-1$ , with (ii)  $\tilde{\alpha}([n] \setminus \{i\} \subset [n])$ , we obtain maps

$$\Delta^{[n-1] \setminus \{i\}} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}), \quad i \neq j, n. \quad (6.26)$$

By construction, the maps (6.25) and (6.26) agree on their overlaps, so we have a map

$$\Lambda_j^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}).$$

Because  $\widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}})$  is a Kan complex, we may fill this horn; and again Proposition 4.16 allows us to deform this filler to be smooth and continuous and collared. The result is a map

$$\tilde{\alpha}([n-1] \subset [n]) : \Delta^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}})$$

which, as we have indicated by the notation, defines  $\tilde{\alpha}([n-1] \subset [n])$ . This finishes our construction of the desired map (6.21) when  $j \neq n-1$ . (By construction, all the composition requirements are met, so  $\tilde{\alpha}$  indeed defines a simplex of  $\text{Liou}_\Delta$ .)

#### 6.4 $n \geq 3$ and $j = n-1$

This is an easier case than the previous one; as it turns out, we have fewer constraints because  $\alpha$  does not make any assignment to the inclusion  $[n-2] \subset [n] \setminus \{n-1\}$ . Indeed, by post-composing the maps

$$\alpha([n-1] \setminus \{i\} \subset [n] \setminus \{i\}) : \Delta^{[n-1] \setminus \{i\}} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}})$$

with (the restrictions of) the maps

$$\tilde{\alpha}([n] \setminus \{i\} \subset [n]) : \partial_i \Delta^n \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n\}}, M_{\{n\}}), \quad i \neq n-1, n$$

from Construction 6.17, we have maps

$$\Delta^{[n-1]\setminus\{i\}} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}), \quad i \neq n-1$$

which agree on the overlaps. In other words, we obtain a single map

$$\Lambda_{n-1}^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}}).$$

We may again use the Kan property of the codomain to extend this to a map from  $\Delta^{n-1}$ , and we may deform this map to be smooth and collared. We declare the resulting map

$$\Delta^{n-1} \rightarrow \widetilde{\text{Emb}}_{\text{Liou}}(M_{\{n-1\}}, M_{\{n\}})$$

to be the desired map (6.21). It is straightforward, while tedious, to check that all composition requirements are met.

## 6.5 Proof of Theorem 6.1 and 6.2

The  $n = 2$  case was accomplished in Section 6.1. For  $n \geq 3$ , unwinding the definition of  $\tilde{\alpha}$ , the horn is filled when we produce maps (6.16), (6.20), and (6.21), satisfying the compatibilities of agreeing along faces and agreeing with  $\alpha$ . This was accomplished by construction.

## 6.6 $(\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$ is weak Kan

Simplices of  $\text{Liou}_{\Delta}^{\text{Def}}$  encode sectorial embeddings  $M_B \times T^*\Delta^B \times T^*[-1, \epsilon) \rightarrow M \times T^*\Delta^A$ . All structures are preserved by taking the direct product with  $T^*[0, 1]$  on the left, so we have a functor

$$T^*[0, 1] \times - : \text{Liou}_{\Delta}^{\text{Def}} \rightarrow \text{Liou}_{\Delta}^{\text{Def}}, \quad M \mapsto T^*[0, 1] \times M.$$

We thus define the semisimplicial set  $(\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$  as an increasing union, as in (1.2).

**Corollary 6.19.** The semisimplicial set  $(\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$  satisfies the (semisimplicial) weak Kan condition.

*Proof.* Any (semisimplicial) map of  $\Lambda_j^n$  into a filtered colimit will factor through a finite stage of the filtration. Thus the corollary reduces to filling horns  $\Lambda_i^n \rightarrow \text{Liou}_{\Delta}^{\text{Def}}$ , which is Theorem 6.2.  $\square$

**Remark 6.20.** Here we see a key difference between  $\text{Liou}_{\Delta}$  and  $\text{Liou}_{\Delta}^{\text{Def}}$ . For the former,  $T^*[0, 1] \times M$  is not a functor, as the compact-support condition for deformations is not preserved under product with (non-compact)  $T^*[0, 1]$ . On the other hand, we saw in Section 5.5 that  $\text{Liou}_{\Delta}$  admits straightforward comparisons to the usual theory of sectorial embeddings and their isotopies; it is more geometrically tractable than  $\text{Liou}_{\Delta}^{\text{Def}}$ . But it does not stabilize easily.

## 7 $\text{Liou}_\Delta$ as an $\infty$ -category

### 7.1 Idempotent equivalences and constant simplices

**Lemma 7.1.** Every object  $M \in \text{Liou}_\Delta, \text{Liou}_\Delta^{\text{Def}}$  admits an idempotent equivalence—i.e., an edge from  $M$  to  $M$  that is an idempotent equivalence (Definition 3.5).

*Proof.* Given an object  $M$ , define a 1-simplex  $e_M$  by declaring

$$M_{\{0,1\}} \times T^* \Delta^1 := M \otimes T^* \Delta^1$$

and set the maps  $y_{\{i\} \subset [1]} := \text{id}_M$ . We claim this is an idempotent self-equivalence. It is obviously idempotent by witnessing the 2-simplex whose  $\Delta$ -movies, for any  $A \subset [2]$ , are given by

$$M_{\{A\}} \times T^* \Delta^A := M \otimes T^* \Delta^A$$

and whose morphisms  $y_{\{A \subset B\}}^s := \text{id}_M$  are  $s$ -independent. This exhibits a 2-simplex with all three faces equaling  $e_M$ , so this shows that  $e_M$  is idempotent.

(a) of Definition 3.5: By assumption, a horn  $\alpha$  gives us identifications  $M_{\{n\}} = M_{\{n-1\}} = M$ . Thus, by symmetry,  $\alpha$  is equivalent to the data of a map  $\Lambda_{n-1}^n \rightarrow \text{Liou}_\Delta$  (or  $\text{Liou}_\Delta^{\text{Def}}$ ). Concretely: Consider the bijection  $\sigma : [n] \rightarrow [n]$  that swaps  $n-1$  and  $n$ , leaving all other elements alone. (This is not order-preserving.) We have an induced map on subsets by  $A \mapsto \sigma(A)$ . Then we can define a map

$$\alpha^\sigma : \Lambda_{n-1}^n \rightarrow \text{Liou}_\Delta, \quad A \mapsto \alpha(\sigma(A)) = M_{\sigma(A)}, \quad \phi_{A \subset B} \mapsto \phi_{\sigma(A) \subset \sigma(B)}.$$

Because the collaring functions are symmetric with respect to permutation actions, these  $\phi$  are all collared. It is clear that  $\alpha^\sigma$  still sends sequences of inclusions to compositions, the assignment satisfies (LD1). And (LD2) is still satisfied because  $M_{\{n\}} = M_{\{n-1\}}$ . Now, because  $\alpha^\sigma$  is an inner horn, it may be filled by a map  $\widetilde{\alpha}^\sigma$  from  $\Delta^n$  (Theorems 6.1 and 6.2). By pre-composing the data of  $\widetilde{\alpha}^\sigma$  with the automorphism of  $T^* \Delta^n$  induced by  $\alpha$ , we obtain a simplex  $\widetilde{\alpha}$  filling  $\alpha$ .

The proof for (b) of Definition 3.5 is similar.  $\square$

**Corollary 7.2.** Every object of the semisimplicial set  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  admits an idempotent equivalence.

*Proof.* Any map of  $\Lambda_0^n$  or  $\Lambda_{n-1}^n$  to  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  factors through some finite stage of the filtered colimit.  $\square$

**Definition 7.3** (Constant simplices). Given any object (i.e., vertex)  $M$  of  $\text{Liou}_\Delta$  (or  $\text{Liou}_\Delta^{\text{Def}}$ ), we define the associated *constant  $I$ -simplex* as follows:

- For every subset  $I' \subset I$ , we declare the  $\Delta^{I'}$ -movie to be  $M \otimes T^* \Delta^{I'}$ .
- We declare, for every inclusion  $I'' \subset I'$ , the collared movie  $\phi_{I'' \subset I'}$  to be the collared movie associated to the constant movie of the identity function. In other words, we set  $\phi_{I'' \subset I'} = \text{id}_M \times T^* \eta_{I'' \subset I'}$ .

**Remark 7.4.** The idempotent edge from Lemma 7.1 is precisely the constant edge associated to the object  $M$ .

**Remark 7.5.** The collection of constant simplices defines a semisimplicial subset of  $\text{Liou}_\Delta$  with the same underlying set of vertices.

Moreover, the collection of constant simplices may obviously be promoted to a simplicial set isomorphic to a disjoint union of  $\Delta^0$ .

## 7.2 Proof of Theorem A

We treat categories as simplicial sets (and semisimplicial sets) by taking nerves (and forgetting degeneracies). Recall the category  $\mathbf{Liou}_{\text{str}}$ .

**Construction 7.6.** We now define a map of semisimplicial sets

$$\mathbf{Liou}_{\text{str}} \rightarrow \mathbf{Liou}_{\Delta}$$

(and hence a map  $\mathbf{Liou}_{\text{str}} \rightarrow \mathbf{Liou}_{\Delta}^{\text{Def}}$  by recalling that  $\mathbf{Liou}_{\Delta}$  is a sub-semisimplicial-set of  $\mathbf{Liou}_{\Delta}^{\text{Def}}$ ). Recall that an  $n$ -simplex of (the nerve of)  $\mathbf{Liou}_{\text{str}}$  is determined by (i) Liouville sectors  $(M_i, \lambda_i)$  for  $i = 0, 1, \dots, n$ , and (ii) for every  $i < j$ , a strict embedding

$$f_{i,j} : M_i \rightarrow M_j \tag{7.1}$$

satisfying  $f_{i,k} = f_{j,k} \circ f_{i,j}$ . Given such data, we define an  $n$ -simplex of  $\mathbf{Liou}_{\Delta}$  as follows.

1. For any subset  $I' \subset [n]$ ,

$$M_{I'} \times T^* \Delta^{I'} := M_{\max I'} \otimes T^* \Delta^{I'}.$$

2. For any  $a : I'' \subset I'$ , we define

$$\phi_{I'' \subset I'} = f_{\max I'', \max I'} \times T^* \eta_a.$$

**Remark 7.7.** This construction is somewhat easier to conceptualize after passing to stabilized versions of  $\mathbf{Liou}_{\text{str}}$  and  $\mathbf{Liou}_{\Delta}$  – see Examples 10.4 and 10.5.

**Theorem 7.8.** The semisimplicial set  $\mathbf{Liou}_{\Delta}$  may be promoted to be a simplicial set satisfying the following properties:

- (a) The face maps are unchanged.
- (b) The map of semisimplicial sets  $\mathbf{Liou}_{\text{str}} \rightarrow \mathbf{Liou}_{\Delta}$  (Construction 7.6) is a map of simplicial sets.
- (c) The degeneracies of constant simplices are constant simplices (Definition 7.3).
- (d)  $\mathbf{Liou}_{\Delta}$  is an  $\infty$ -category.

**Notation 7.9.** From now on, we also let  $\mathbf{Liou}_{\Delta}$  denote the simplicial set guaranteed by Theorem 7.8. This abuse of notation will not cause too much confusion—often, the important structure will be whether a *map* to/from  $\mathbf{Liou}_{\Delta}$  is semisimplicial or simplicial.

**Remark 7.10.** The last item of Theorem 7.8 is Theorem A from the introduction. Theorem 7.8 elaborates on the various properties that the  $\infty$ -category  $\mathbf{Liou}_{\Delta}$  satisfies.

*Proof of Theorem 7.8.* By Theorem 6.1,  $\mathbf{Liou}_{\Delta}$  satisfies the semisimplicial weak Kan condition. By Lemma 7.1, every object admits an idempotent self-equivalence. Thus Theorem 3.6 shows that we can satisfy (b) and (c) by taking  $A$  (see the notation in Theorem 3.6) to be the the image of  $\mathbf{Liou}_{\text{str}}$  (which also contains all constant simplices as in Remark 7.5). We finally note that, indeed, Construction 7.6 is compatible with the idempotent self-equivalences from the proof of Lemma 7.1.  $\square$

**Theorem 7.11.** The semisimplicial set  $\mathbf{Liou}_{\Delta}^{\text{Def}}$  may be promoted to be a simplicial set satisfying the following properties:

- (a) The face maps are unchanged.
- (b) The map of semisimplicial sets  $\mathbf{Liou}_\Delta \rightarrow \mathbf{Liou}_\Delta^{\mathcal{D}\text{ef}}$  is a map of simplicial sets when  $\mathbf{Liou}_\Delta$  is given the simplicial set structure from Theorem 7.8. (In particular, the degeneracies of constant simplices are constant simplices.)
- (c)  $\mathbf{Liou}_\Delta^{\mathcal{D}\text{ef}}$  is an  $\infty$ -category.

*Proof.* Identical to the proof of Theorem 7.8, except one takes  $A$  to be  $\mathbf{Liou}_\Delta$ .  $\square$

### 7.3 Some of the geometry of $\mathbf{Liou}_\Delta$

Concrete geometric constructions in  $\mathbf{Liou}_\Delta$  can be given  $\infty$ -categorical interpretations. We have already seen in Proposition 5.19 that 2-simplices in  $\mathbf{Liou}_\Delta$  encode isotopies (through sectorial embeddings) of sectorial embeddings.

And in Example 5.18, we saw that 1-simplices encode (not necessarily strict) sectorial embeddings. We now record the converse construction.

**Remark 7.12.** We could have introduced the following construction back in Section 5.5, but we have waited until we established the  $\infty$ -categorical structure (namely, degeneracies) of  $\mathbf{Liou}_\Delta$ . The reason is so that we can then state Proposition 7.14, which requires a notion of homotopy to state precisely (and hence, in turn, requires us to know the degenerate edges of  $\mathbf{Liou}_\Delta$ ).

**Construction 7.13.** Let  $f : M_0 \rightarrow M_1$  be a sectorial embedding (Definition 2.23) so that  $f^*\lambda_1 = \lambda_0 + dh$  for some compactly supported  $h$ . Then for every

- smooth, compactly supported function  $\tilde{h} : M_1 \rightarrow \mathbb{R}$  extending  $f_*h$ , and
- smooth, collared, weakly increasing function  $\chi : \Delta^1 \rightarrow \mathbb{R}_{\geq 0}$  interpolating from 1 to 0 (so that  $\chi$  equals 1 at the initial vertex of  $\Delta^1$ , and equals 0 at the terminal vertex),

one obtains an edge of  $\mathbf{Liou}_\Delta$  by declaring

$$M_{01} \times T^*\Delta^1 = (M_1 \times T^*\Delta^1, \lambda_1 \oplus pdq + d(\chi(q)\tilde{h})), \quad (7.2)$$

and

- $(M_0 \times T^*[-1, 0] \rightarrow M_{01} \times T^*\Delta^1) = f \times T^*\eta_{0 \subset 01}$ ,
- $(M_1 \times T^*[-1, 0] \rightarrow M_{01} \times T^*\Delta^1) = \text{id} \times T^*\eta_{1 \subset 01}$ .

**Proposition 7.14.** Fix a map  $f : M_0 \rightarrow M_1$  for which  $f^*\lambda_1 = \lambda_0 + dh$  for some compactly supported function  $h : M_0 \rightarrow \mathbb{R}$ .

- (a) For any two choices of  $\tilde{h}$  and  $\chi$ , Construction 7.13 results in homotopic edges of  $\mathbf{Liou}_\Delta$ . (Informally: Construction 7.13 is independent of the choices of  $\chi, \tilde{h}$  up to homotopy.)
- (b) Further, if  $f$  is smoothly isotopic through sectorial embeddings to  $f'$ , then Construction 7.13 for  $f$  and  $f'$  result in homotopic edges in  $\mathbf{Liou}_\Delta$ . (Informally: Construction 7.13 is compatible with the relation of isotopy.)
- (c) Given another sectorial embedding  $g : M_1 \rightarrow M_2$ , there exists a triangle in  $\mathbf{Liou}_\Delta$  whose 02, 01, and 12 edges are associated to  $g \circ f$ ,  $f$ , and  $g$ , respectively. (Informally: Construction 7.13 respects composition up to homotopy.)

- (d) In particular, if  $f$  is a Liouville equivalence, then the edge associated to  $f$  is an equivalence in  $\text{Liou}_\Delta$ .

*Proof.* (a) It suffices to construct a 2-simplex in  $\text{Liou}_\Delta$  for which the edge from 0 to 1 is a degenerate edge for  $M_0$ , the edge from 1 to 2 is constructed using  $\chi$  and  $h$ , and the edge from 0 to 2 is constructed using some other choices  $\chi'$  and  $h'$ .

Let  $M_{01} \times T^*\Delta^1$  denote the Liouville sector constructed out of  $\chi$  and  $\tilde{h}$  (7.2). We let  $(M_{01} \times T^*\Delta^1)'$  denote the Liouville sector constructed out of  $\chi'$  and  $\tilde{h}'$ . By definition, we have smooth embeddings

$$(M_{01} \times T^*\Delta^1 \times T^*[-1, 0]) \rightarrow M_1 \times T^*\Delta^2 \leftarrow (M_{01} \times T^*\Delta^1 \times T^*[-1, 0])'$$

each given by  $\text{id} \times T^*\eta$  (where the rightward pointing  $\eta$  is induced by the inclusion  $\{1, 2\} \subset \{0, 1, 2\}$  and the leftward pointing  $\eta$  is induced by the inclusion  $\{0, 2\} \subset \{0, 1, 2\}$ ). Because everything in sight is collared, we may extend (the pushforwards of) both  $\chi\tilde{h}$  and  $\chi'\tilde{h}'$  to a smooth, compactly supported function  $\tilde{H} : M_1 \times \Delta^2 \rightarrow \mathbb{R}$ ; moreover, by the smooth contractibility of the space of collared and compactly supported smooth functions, we may further demand that  $\tilde{H}$  is collared by a function  $h_{01} : M_1 \times \Delta^1 \rightarrow \mathbb{R}$  along the edge from 0 to 1 for which  $f^*h_{01} = h$ . By declaring the inclusion  $M_0 \times T^*\Delta^1 \rightarrow M_1 \times T^*\Delta^2$  to be given by  $f \times T^*\eta_{01 \subset 012}$ , the totality of the data in this paragraph exhibits the desired two-simplex.

(b) Let  $f_s$  denote a homotopy between two sectorial embeddings  $f = f_0$  and  $f' = f_1$ . Without loss of generality, we may assume that  $f_s$  is parametrized by  $s \in \Delta^1$ , and that  $f_s$  is collared (and in particular,  $s$ -independent near the endpoints of  $\Delta^1$ ).

We may construct a strict sectorial embedding

$$\tilde{f} : M_0 \otimes T^*\Delta^1 \rightarrow M_1 \times T^*\Delta^1$$

where the Liouville structure on the codomain depends on the family  $f_s$ , and is given by  $\lambda_1 \oplus pdq + H$  for some smooth, compactly supported function  $H : M_1 \times \Delta^1 \rightarrow \mathbb{R}$ . By identifying  $\Delta^1$  with the 2nd face of  $\Delta^2$  (i.e., with the face spanning the 0th and 1st vertices) and again by the collaring condition of everything in sight, we may extend the trio of  $H$ ,  $\chi\tilde{h}$  and  $\chi'\tilde{h}'$  to a single smooth, compactly supported function  $\tilde{H} : M_1 \times \Delta^2 \rightarrow \mathbb{R}$ . (Here,  $\chi\tilde{h}$  is pushed forward along the inclusion of the 12 face, constant along the collaring, and likewise  $\chi'\tilde{h}'$  is pushed forward along the inclusion of the 02 face.) The totality of the data in this paragraph exhibits a 2-simplex collared by  $\text{id}_{M_0}$  (along the 01 edge) and the edges associated to  $f$  and  $f'$  from Construction 7.13; in other words, we have constructed a homotopy from  $f$  to  $f'$ .

(c) We define  $M_{012} = M_2$  as a Liouville sector, and define a smooth embedding

$$M_{01} \times T^*\Delta^1 := M_1 \times T^*\Delta^1 \rightarrow M_2 \times T^*\Delta^2$$

by  $g \times T^*\eta_{01 \subset 012}$ . By our collaring assumptions, the three pushforwards

$$(g \times \eta_{01 \subset 012}) * \chi_{01} \tilde{h}_{01}, \quad (\text{id}_{M_2} \times \eta_{12 \subset 012}) * \chi_{12} \tilde{h}_{12}, \quad (\text{id}_{M_2} \times \eta_{02 \subset 012}) * \chi_{02} \tilde{h}_{02}$$

(which are smooth functions defined on subsets of  $M_2 \times \Delta^2$ ) agree along their overlaps; we can choose a smooth, compactly supported function  $H : M_2 \times \Delta^2 \rightarrow \mathbb{R}$  extending all three. This defines our Liouville sector  $M_{012} \times T^*\Delta^2 = (M_2 \times T^*\Delta^2, \lambda_2 \oplus pdq + dH)$ , and the three face inclusions are defined by

$$g \times T^*\eta_{01 \subset 012} \quad \text{id}_{M_2} \times T^*\eta_{12 \subset 012} \quad \text{id}_{M_2} \times T^*\eta_{02 \subset 012}.$$

(d) This is obvious in light of (b) and (c).  $\square$



**Remark 7.15.** When every sectorial embedding in sight is *strict*, we may choose every extension to be constant in the simplicial directions; by declaring every  $h$  and  $H$  to be zero (and hence removing the dependence on  $\chi$ ) the proof of Proposition 7.14 in fact recovers the functor of  $\infty$ -categories from  $N(\text{Liou}_{\text{str}})$  to  $\text{Liou}_{\Delta}$  from Theorem 7.8.

**Remark 7.16.** Indeed, one can think of an edge in  $\text{Liou}_{\Delta}$  as a choice of embedding  $f : M_0 \rightarrow M_1$ , along with an interpolation of  $\lambda_1$  to an extension of  $f_*\lambda_0$  along compactly supported functions  $h_t : M_1 \rightarrow \mathbb{R}$ . The rough idea is that the forgetful map  $(f, \{h_t\}) \mapsto f$  has contractible fibers (the space of  $h$  is convex) so that the space of edges  $(f, h_t)$  in  $\text{Liou}_{\Delta}$  is the space of (not necessarily strict) Liouville embeddings  $f$ .

**Example 7.17** (Isotopies). The converse to Proposition 5.19 is also true—any isotopy gives rise to a 2-simplex in  $\text{Liou}_{\Delta}$ .

Let us illustrate this in a simple example: Given two Liouville sectors  $M$  and  $N$ , and a smooth isotopy of sectorial embeddings  $\{f_t : M \rightarrow N\}_{t \in [0,1]}$ , we will produce a 2-simplex in  $\text{Liou}_{\Delta}$  that “encodes” this data (in a way depending on some choices).

We choose

$$M_{\{0\}} = M_{\{0,1\}} = M_{\{1\}} = M$$

with the maps  $M_{\{i\}} \otimes T^*[-1, \epsilon] \rightarrow M_{\{0,1\}} \times T^*\Delta^{\{0,1\}}$  given by  $\text{id}_M \times T^*\eta_{\{i\} \subset \{0,1\}}$  for  $i = 0, 1$ . We also choose

$$M_{\{2\}} = M_{\{0,2\}} = M_{\{1,2\}} = M_{\{0,1,2\}} = N$$

choosing all relevant maps among these to be given by  $\text{id}_N \times T^*\eta$ .

Finally, let us convert  $\{f_t\}_{t \in [0,1]}$  to a collared family  $\{\tilde{y}^s : M \rightarrow N\}_{s \in \Delta^{\{0,1\}}}$  by choosing some orientation-respecting  $\phi : \Delta^{\{0,1\}} \rightarrow [0, 1]$  that sends some open neighborhood of the initial vertex  $\Delta^{\{0\}}$  to  $0 \in [0, 1]$ , and some open neighborhood of the vertex  $\Delta^{\{1\}}$  to  $1 \in [0, 1]$ .

Then we are finished by

- declaring the map  $y_{\{0\} \subset \{0,2\}}^0 M = M_{\{0\}} \rightarrow M_{\{0,2\}} = N$  to be the map  $f_0$ ,
- declaring the map  $y_{\{1\} \subset \{1,2\}}^1 : M = M_{\{1\}} \rightarrow M_{\{1,2\}} = N$  to be the map  $f_1$ ,
- declaring the family  $\{y_{\{0,1\} \subset \{0,1,2\}}^s : M_{\{0,1\}} \rightarrow M_{\{0,1,2\}}\}_{s \in \Delta^{\{0,1\}}}$  to be  $y_{\{0,1\} \subset \{0,1,2\}}^s = f_{\phi(s)}$ , and
- Choosing a collared, compactly supported smooth function  $N \times \Delta^2 \rightarrow \mathbb{R}$  so that all embeddings are strict (similar to Construction 7.13).

We may informally draw this 2-simplex as follows:

$$\begin{array}{ccc}
 & M_2 & \\
 \swarrow \sim & & \searrow \sim \\
 M_{02} & \xRightarrow{\sim} & M_{012} \xleftarrow{\sim} M_{12} \\
 \nearrow & & \nwarrow \\
 M_0 & \xrightarrow{\quad} & M_{01} \xleftarrow{\quad} M_1
 \end{array}
 =
 \begin{array}{ccc}
 & N & \\
 \swarrow = & & \searrow = \\
 N & \xRightarrow{=} & N \xleftarrow{=} N \\
 \nearrow f_0 & & \nwarrow f_1 \\
 M & \xrightarrow{=} & M \xleftarrow{=} M
 \end{array}
 \quad (7.3)$$

Above, the double arrows  $\Rightarrow$  indicate the data of a family of morphisms, and arrows labeled by an equality sign indicate the identity morphism, or a constant family of identity morphisms.

**Remark 7.18.** There is likewise a way to construct a 2-simplex out of the data of  $\{f_t\}$  for which the 1st and 2nd vertices are given by  $N$ , while only the 0th vertex is given by  $M$ . (This is in contrast to Example 7.17, whose 2-simplex has the property that the 1st vertex is given by  $M$ .) Such a construction involves extending the isotopy  $f_t$  to an isotopy of  $N$  to itself. This is possible thanks to isotopy extension (Proposition 2.54).

We leave the details to the reader, but point out the following: If  $\mathbf{Liou}_\Delta$  is to be promoted to be an  $\infty$ -category, then there should be no appreciable homotopic difference between the space of 2-simplices where the 0, 1 edge is constant, and the space of 2-simplices whose 1, 2 edge is constant. We see here the importance of isotopy extension in even having a hope for this to be true.

## 8 Mapping spaces of $\text{Liou}_\Delta$

We have already seen that edges in  $\text{Liou}_\Delta$  encode (not necessarily strict) sectorial embeddings, and that triangles in  $\text{Liou}_\Delta$  encode isotopies among (not necessarily strict) sectorial embeddings. See Example 5.18 and Proposition 5.19.

Using higher-dimensional analogues of the construction in Example 7.17, we now prove that morphism spaces of  $\text{Liou}_\Delta$  are homotopy equivalent to the spaces of (not necessarily strict) sectorial embeddings:

**Theorem 8.1.** Fix two Liouville sectors  $M$  and  $N$ . There exists a homotopy equivalence of simplicial sets

$$\text{Emb}_{\text{Liou}}(M, N) \simeq \text{hom}_{\text{Liou}_\Delta}^R(M, N).$$

Here,  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$  is the Kan complex modeling the space of morphisms from  $M$  to  $N$  (Notation 8.2 below), and  $\text{Emb}_{\text{Liou}}(M, N)$  is the singular complex of the space of sectorial embeddings (Notation 2.59).

### 8.1 The right mapping spaces

Now that we have established  $\text{Liou}_\Delta$  as an  $\infty$ -category (Theorem 7.8), we may speak of its morphism spaces.

**Notation 8.2.** We recall the “space<sup>25</sup> of right morphisms” for  $\text{Liou}_\Delta$ ,

$$\text{hom}_{\text{Liou}_\Delta}^R(M, N).$$

$\text{hom}_{\text{Liou}_\Delta}^R(M, N)$  is the simplicial set whose  $k$ -simplices are given by maps  $\Delta^{k+1} \rightarrow \text{Liou}_\Delta$  whose restriction to the vertex  $\Delta^{\{k+1\}}$  is given by  $N$  and whose restriction to  $\Delta^{\{0, \dots, k\}}$  is the degenerate simplex at  $M$ .

**Example 8.3.** Fix a (not necessarily exact) sectorial embedding  $f : M \rightarrow N$ . Then one may construct a 1-simplex in  $\text{Liou}_\Delta$  as in Example 5.18. In particular, these  $f$  are examples of zero-simplices in  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ .

**Example 8.4.** Fix  $f$  as in Example 8.3. Now suppose that we have a  $k - 1$ -simplex  $\gamma$  in  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$  for which the boundary  $\partial\Delta^{k-1}$  is sent to the degenerate simplex at  $f$ . (By definition of simplicial homotopy groups, such a thing defines an element  $[\gamma] \in \pi_{k-1}(\text{hom}_{\text{Liou}_\Delta}^R(M, N), f)$ .) Let us parse this data.

$\gamma$ , by definition, is the data of a  $k$ -simplex in  $\text{Liou}_\Delta$  whose  $k$ th face is the degenerate simplex at the object  $M$ , and whose other faces are the degenerate simplices  $s_0 \circ \dots \circ s_0(f)$  at the morphism  $f$ . Thus,  $\gamma$  is determined by

- A choice of Liouville sector  $Q = \gamma([k])$  (together with Liouville structure on  $Q \times T^*\Delta^k$ ),
- For every  $k \in I \subset [k]$ , choice of map  $\gamma(I \subset [k]) : \Delta^I \rightarrow \text{Emb}_{\text{Liou}}(N, Q)$  (so that the associated movie defines a strict embedding of  $N \times T^*\Delta^I \times [-1, \epsilon]^{[k] \setminus I}$  into  $Q \times T^*\Delta^k$ ), and
- For every  $I \subset [k - 1]$ , a choice of map  $\gamma(I \subset [k]) : \Delta^I \rightarrow \text{Emb}_{\text{Liou}}(M, Q)$  (with the analogous strict embedding requirement as above),

---

<sup>25</sup>See the discussion around Proposition 1.2.2.3 of [17].

satisfying the composition requirements. Because of the composition requirements, the above data may be summarized.  $\gamma$  is determined by  $Q$  (together with Liouville structure on  $Q \times T^*\Delta^k$ ), along with

(i) A map

$$\Phi_N^Q : \Lambda_k^k \rightarrow \text{Emb}_{\text{Liou}}(N, Q),$$

and

(ii) a map

$$\Phi_M^Q : \Delta^{k-1} \rightarrow \text{Emb}_{\text{Liou}}(M, Q)$$

satisfying the equation

$$\Phi_N^Q|_{\partial\Delta^{k-1}} \circ f = \Phi_M^Q|_{\partial\Delta^{k-1}}$$

where  $f$  is treated as a constant map  $\partial\Delta^{k-1} \rightarrow \text{Emb}_{\text{Liou}}(M, N)$ . (Note also that  $\partial\Delta^{k-1}$  is a subsimplicial set of  $\Lambda_k^k$ .)

By definition of  $\text{Liou}_\Delta$ , every  $\Phi$  above, when restricted to any subsimplex, must be smooth and collared and continuous.

Note also that  $N$  is necessarily diffeomorphic to  $Q$  by (LD2); the diffeomorphism is given by evaluating the map  $\Phi_N^Q$  at the maximal vertex of  $\Lambda_k^k$ . In particular, note that  $\Phi_N^Q$  encodes a (contractible) family of embeddings all isotopic to the just-mentioned diffeomorphism.

**Notation 8.5** ( $\text{hom}'_{\text{Liou}_\Delta}(M, N)$ ). Let

$$\text{hom}'_{\text{Liou}_\Delta}(M, N) \subset \text{hom}^R_{\text{Liou}_\Delta}(M, N) \quad (8.1)$$

denote the semisimplicial subset of those  $(k+1)$ -simplices in  $\text{Liou}_\Delta$  satisfying the following:

- If  $I' \subset [k+1]$  does not contain  $k+1$ , then the  $\Delta^{I'}$ -movie is given by  $M \otimes T^*\Delta^{I'}$ .
- If  $I' \subset [k+1]$  does contain  $k+1$ , then the corresponding  $\Delta^{I'}$ -movie has underlying manifold  $N \times T^*\Delta^{I'}$ . (As a sector, this object need not necessarily be a tensor.)
- If  $I'' \subset I'$  and neither contain  $k+1$ , then the associated map  $\phi_{I'' \subset I'}$  is given by  $\text{id}_M \times T^*\eta_{I'' \subset I'}$ .
- Likewise, if  $I'' \subset I'$  and both contain  $k+1$ , then the map  $\phi_{I'' \subset I'}$  is given by  $\text{id}_N \times T^*\eta_{I'' \subset I'}$ .

Note that if  $I''$  does not contain  $k+1$  but  $I'$  does, then  $\phi_{I'' \subset I'}$  encodes some  $\Delta^{I''}$ -family of sectorial embeddings from  $M$  to  $N$ .

**Notation 8.6.** Let  $\widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N)$  denote the semisimplicial set where a  $k$ -simplex is a pair of

1. A collared, smooth map  $f : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, N)$ , and
2. A  $\Delta^{k+1}$ -parametrized family of compactly supported smooth functions  $\{h^s : N \rightarrow \mathbb{R}\}_{s \in \Delta^{k+1}}$ . Notice this determines a  $\Delta^{k+1}$ -movie  $N \times T^*\Delta^{k+1}$  associated to  $\{h^s\}$ . We demand that the pair  $(f, \{h^s\})$  satisfies the following:

- (a) The family  $\{h^s\}$  is collared.
- (b) For every subset  $I \subset [k]$ , there exists a (necessarily unique) strict sectorial embedding

$$M \otimes T^*(\Delta^I \times [-1, 0]^{[k+1] \setminus I}) \rightarrow N \times T^*\Delta^{k+1}$$

exhibiting a movie associated to  $f|_{\Delta^I}$ , and

(c) For every  $s \in \Delta^{k+1}$  near the vertex  $\Delta^{\{k+1\}}$ ,  $h^s \equiv 0$ .

**Remark 8.7.** There is a natural isomorphism of semisimplicial sets between  $\widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N)$  and  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$ . For example, given a pair  $(f, \{h^s\})$ , the restrictions of  $f$  to the faces of  $\Delta^k$  determine maps  $\phi_{I'' \subset I'}$ , while the restrictions of  $\{h^s\}$  to the faces of  $\Delta^{k+1}$  determine the Liouville structures on the various  $\Delta$ -movies.

**Remark 8.8.** We have a forgetful map of semisimplicial sets.

$$\widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N) \rightarrow \text{Emb}_{\text{Liou}}^{\text{coll}}(M, N) \quad (8.2)$$

Using the same methods as Proposition 4.17 (and noting the convexity of the space of compactly supported functions  $h$ ), we see that  $\widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N)$  admits idempotent autoequivalences for all its vertices. By utilizing Theorem 3.6 again, we conclude that (8.2) results in a map of *simplicial* sets, and both the domain and codomain are Kan complexes.

**Proposition 8.9.** The map (8.2) is a homotopy equivalence of simplicial sets.

*Proof.* The map is a Kan fibration because we can relatively fill horns simply by choosing collared extensions of  $\{h^s\}_{s \in \Delta_j^n}$  to  $\{h^s\}_{s \in \Delta^n}$ . The fibers are contractible because of the convexity of the space of compactly supported functions.  $\square$

## 8.2 $\text{hom}'$ is equivalent to $\text{hom}^R$

Combining Remarks 8.7 and 8.8, we conclude:

**Proposition 8.10.**  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$  admits a simplicial structure lifting the face maps inherited from  $\text{hom}^R_{\text{Liou}_\Delta}(M, N)$ .

Now, the inclusion (8.1) is a map of semisimplicial sets, but may not be a map of simplicial sets. This is because the degeneracy maps of  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$  are induced from  $\widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N)$ , while the degeneracy maps of  $\text{hom}^R_{\text{Liou}_\Delta}(M, N)$  are induced from the degeneracy maps of  $\text{Liou}_\Delta$ .

However, it is straightforward to see that, under the semisimplicial set map (8.1), degenerate edges of  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$  are sent to equivalences in  $\text{hom}^R_{\text{Liou}_\Delta}(M, N)$ . Theorem 3.7 guarantees that the semisimplicial set map may then be deformed to a *simplicial* set map

$$\text{hom}'_{\text{Liou}_\Delta}(M, N) \rightarrow \text{hom}^R_{\text{Liou}_\Delta}(M, N) \quad (8.3)$$

with the same assignment on objects as (8.1), and with the property that the images of (8.3) are naturally homotopic to the images of (8.1). (We used this result in Construction 10.11 as well.)

We now prove:

**Lemma 8.11.** The map (8.3) is a homotopy equivalence of simplicial sets.

Because both sides of (8.3) are Kan complexes, it suffices to show that the induced maps on  $\pi_0$ , and on  $\pi_k$  (along all connected components), are bijections.

*Proof of surjection on  $\pi_0$ .* Let us show that the inclusion  $\text{hom}'_{\text{Liou}_\Delta}(M, N) \subset \text{hom}^R_{\text{Liou}_\Delta}(M, N)$  is a surjection on  $\pi_0$ .

For this, fix an arbitrary vertex  $F$  of  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ . By definition of an edge of  $\text{Liou}_\Delta$ ,  $F$  produces the data of a  $\Delta^1$ -movie of the form  $X \times T^*\Delta^1$ , along with two morphisms

$$f : M \rightarrow (X, \lambda^0) \quad \phi : N \rightarrow (X, \lambda^1)$$

where we have emphasized the Liouville form  $\lambda^t$  of  $X_{01}$ , where  $\lambda^t$  is the form on  $X_{01}$  at time  $t \in \Delta^1 \cong [0, 1]$ . By definition of 1-simplex in  $\text{Liou}_\Delta$ , we know that  $\phi$  is an isomorphism of Liouville sectors. (In particular, a diffeomorphism.)

We produce a 2-simplex in  $\text{Liou}_\Delta$  which we illustrate informally using the following diagram:

$$\begin{array}{ccccc} & & N = N_2 & & \\ & \swarrow \phi & \downarrow \text{id} & \searrow \text{id} & \\ & X & \rightleftarrows N & \rightleftarrows N & \\ & \swarrow \phi^{-1} & \downarrow \phi^{-1} & \searrow \text{id} & \\ f \nearrow & & & & \nwarrow \phi^{-1} f \\ M = M_0 & \xrightarrow{\text{id}} & M & \xleftarrow{\text{id}} & M = M_1 \end{array} \quad (8.4)$$

Note that the  $\{0, 1\}$  edge (the bottom edge) is a constant 1-simplex at the object  $M$ . In particular, the 2-simplex indeed depicts an edge in  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ . The  $\{0, 2\}$  edge of the diagram is the given vertex  $F$  of  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ . The  $\{1, 2\}$  edge of the diagram is the vertex in  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$  to which we claim the given vertex is homotopic.

We first describe the underlying smooth maps. All the “double” arrows of the form  $\implies$  would, in general, depict a 1-simplex family of morphisms. Here, they are all constant families given by the morphisms as labeled in the diagram. (So for example, we have a constant 1-simplex family of morphisms from  $X$  to  $N$  given by  $\phi^{-1}$ .)

To finish, we describe the Liouville structures on the  $\Delta$ -movies  $X \times T^*\Delta^{\{0,2\}}$ ,  $N \times T^*\Delta^{\{1,2\}}$ , and  $N \times T^*\Delta^{\{0,1,2\}}$ . The Liouville structure on the first is the one specified by  $F$ . On  $N \times T^*\Delta^{\{1,2\}}$ , we choose a family of Liouville structures on  $N$  which equals the given structure on  $N$  near the vertex  $\Delta^{\{2\}}$ , and equals the structure induced by  $\phi^{-1}$  near the vertex  $\Delta^{\{1\}}$ .

Finally, given the maps above, the Liouville structure on  $N \times T^*\Delta^{\{0,1,2\}}$  is uniquely specified along the boundary of  $\partial\Delta^{\{0,1,2\}}$  by requiring that all morphisms in sight be strict. Now, the space of compactly supported smooth functions on  $N$  is convex (hence contractible) so we may choose any filling of this family of Liouville structures to all of  $\Delta^{\{0,1,2\}}$  to exhibit a  $\Delta^{\{0,1,2\}}$ -movie structure on  $N \times T^*\Delta^{\{0,1,2\}}$ .  $\square$

### 8.3 Proof of injection on homotopy groups

*Proof.* We choose some element  $f \in \text{hom}'(M, N)$  and show that the map of homotopy groups based at  $f$

$$\pi_{k-1}(\text{hom}'_{\text{Liou}_\Delta}(M, N), f) \rightarrow \pi_{k-1}(\text{hom}_{\text{Liou}_\Delta}^R(M, N), f), \quad k \geq 2$$

is an injection. Similar techniques as what follows also demonstrate that the induced map on  $\pi_0$  is an injection.

**Remark 8.12** (Parsing the hypotheses). Fix  $\gamma_0, \gamma_1$  representing elements on  $\pi_{k-1}(\text{hom}'_{\text{Liou}_\Delta}(M, N), f)$  and assume their images are homotopic (rel  $f$ ) in  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ . By definition of simplicial homotopy groups, and by the definition of  $\text{hom}^R$ , we conclude there exists a simplex

$$F : \Delta^{k+1} \rightarrow \text{Liou}_\Delta$$

satisfying the usual boundary conditions. By definition of simplices in  $\text{Liou}_\Delta$ ,  $F$  in particular gives rise to maps

$$\begin{aligned}\Phi_M^N : \partial\Delta^k &\rightarrow \text{Emb}_{\text{Liou}}(M, N) \\ \Phi_N^Q : \Lambda_{k+1}^{k+1} &\rightarrow \text{Emb}_{\text{Liou}}(N, Q) \\ \Phi_M^Q : \Delta^k &\rightarrow \text{Emb}_{\text{Liou}}(M, Q)\end{aligned}\tag{8.5}$$

which are all piecewise smooth and collared, and which satisfy

$$\Phi_N^Q|_{\partial\Delta^k} \circ \Phi_M^N = \Phi_M^Q|_{\partial\Delta^k}.$$

Because  $F$  exhibits a homotopy (rel  $f$ ) from  $\gamma_0$  to  $\gamma_1$ , we observe

$$\Phi_M^N|_{\partial_i\Delta^k} = \begin{cases} \gamma_0 \circ (\partial_0\Delta^k \cong \Delta^{k-1}) & i = 0 \\ \gamma_1 \circ (\partial_1\Delta^k \cong \Delta^{k-1}) & i = 1 \\ \underline{f} & i \neq 0, 1 \end{cases}\tag{8.6}$$

where  $\underline{f}$  is the degenerate  $(k-1)$ -simplex mapping to  $\text{Emb}_{\text{Liou}}(M, N)$  with value  $f$ , and the pre-compositions are by the unique simplicial isomorphisms. Finally,  $F$  also specifies a Liouville isomorphism

$$\phi : N \rightarrow Q \quad \text{at the vertex } \Delta^{\{k+1\}}.\tag{8.7}$$

Now that we have parsed the hypotheses, our task is to construct a smoothly collared map

$$\Gamma : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, N)\tag{8.8}$$

satisfying the boundary conditions (8.6) (with  $\Gamma|_{\partial_i\Delta^k}$  replacing  $\Phi_M^N|_{\partial_i\Delta^k}$ ). We refer the reader to Figure 1 for a visual guide to the construction of  $\Gamma$ , which we now make precise.

Consider the maps

$$\begin{aligned}\underline{\Phi}_M^N : \partial\Delta^k \times \Delta^1 &\rightarrow \partial\Delta^k \xrightarrow{\Phi_M^N} \text{Emb}_{\text{Liou}}(M, N) \\ \underline{\Phi}_N^Q : \partial\Delta^k \times \Delta^1 &\rightarrow \partial\Lambda_{k+1}^{k+1} \xrightarrow{\Phi_N^Q} \text{Emb}_{\text{Liou}}(N, Q)\end{aligned}$$

where the unlabeled maps are the projections to the  $\partial\Delta^k$  factor; realizing the horn as a quotient of a cylinder along  $\partial\Delta^k \times \Delta^{\{1\}}$ ; and to a point, respectively. Composition defines a map

$$\Gamma_a := \phi^{-1} \circ \underline{\Phi}_N^Q \circ \underline{\Phi}_M^N : \partial\Delta^k \times \Delta^1 \rightarrow \text{Emb}_{\text{Liou}}(M, N).$$

Because  $\Phi_N^Q$  is identical to  $\phi$  (8.7) near the vertex  $\Delta^{\{k+1\}}$ ,  $\Gamma_a$  has the property that its restriction to  $\partial\Delta^k \cong \partial\Delta^k \times \Delta^{\{1\}}$  satisfies the boundary conditions (8.6).

We now “cap off”  $\Gamma_a$  along  $\partial\Delta^k \times \Delta^{\{0\}}$ . For this, choose a piecewise smooth homeomorphism  $p : |\Lambda_{k+1}^{k+1}| \xrightarrow{\cong} |\Delta^k|$ , for example by projection along a vector normal to  $|\Delta^k|$  and passing through the vertex  $|\Delta^{\{k+1\}}|$ . We choose a collared smoothing  $(\Phi_N^Q)'$  of the composition  $\Phi_N^Q \circ p^{-1}$ , and define

$$\Gamma_c := \phi^{-1} \circ (\Phi_N^Q)' \circ \phi^{-1} \circ \Phi_M^Q : |\Delta^k| \rightarrow \text{Emb}_{\text{Liou}}(N, Q).$$

To continue the cap-off process, we note that

$$\Gamma_a|_{\partial\Delta^k \times \Delta^{\{0\}}} = \phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \circ \Phi_M^N \quad \text{and} \quad \Gamma_c|_{\partial\Delta^k} = \phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \circ \phi^{-1} \circ \Phi_M^Q|_{\partial\Delta^k}\tag{8.9}$$




Figure 1: An informal drawing, for  $k = 2$ , of the maps  $\Gamma_a, \Gamma_b, \Gamma_c$ , and  $\Gamma$  in proving the injectivity on homotopy groups.



where the latter equality follows from the collaring assumption on  $(\Phi_N^Q)'$ . In Lemma 8.13 below, we show that these two are smoothly homotopic as maps  $|\partial\Delta^k| \rightarrow \text{Emb}_{\text{Liou}}(M, N)$ . Thus we conclude the existence of a smooth map

$$\Gamma_b : |\partial\Delta^k| \times |\Delta^1| \rightarrow \text{Emb}_{\text{Liou}}(M, N)$$

realizing this smooth homotopy. Because everything in sight is assumed collared, we obtain a gluing

$$\Gamma_c \bigcup \Gamma_b \bigcup \Gamma_a : |\Delta^k| \bigcup \left( |\partial\Delta^k| \times |\Delta^1| \right) \bigcup |\partial\Delta^k \times \Delta^1| \rightarrow \text{Emb}_{\text{Liou}}(M, N).$$

Now choose a diffeomorphism

$$|\Delta^k| \xrightarrow{\cong} |\Delta^k| \bigcup \left( |\partial\Delta^k| \times |\Delta^1| \right) \bigcup |\partial\Delta^k \times \Delta^1|$$

preserving the collaring of  $\Gamma_a$  near the boundary of the domain  $|\Delta^k|$ . We define  $\Gamma$  (8.8) to be the composition of this diffeomorphism with  $\Gamma_c \bigcup \Gamma_b \bigcup \Gamma_a$ . We see that  $\Gamma$  satisfies the boundary conditions (8.6) because  $\Gamma_a$  does.

We conclude by noting that, given  $\Gamma$ , one now is free to choose a Liouville structure on  $N \times T^*\Delta^{k+1}$  rendering  $\Gamma$  a family of strict embeddings.  $\square$

**Lemma 8.13.** There exists a smooth homotopy  $\phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \sim \Phi_N^Q|_{\partial\Delta^k} \circ \phi^{-1}$  among maps from  $\partial\Delta^k$  to  $\text{Emb}_{\text{Liou}}(M, N)$ . In particular, there exists a homotopy between the two maps in (8.9).

*Proof.* We have the obvious projection  $\partial\Delta^k \times \Delta^1 \rightarrow \Delta_{k+1}^{k+1}$ , again by collapsing  $\partial\Delta^k \times \Delta^{\{1\}}$ . Composing the projection with  $\Phi_N^Q$  from (8.5), we thus witness a homotopy from  $\Phi_N^Q|_{\partial\Delta^k}$  to  $\phi$  (by abuse of notation, this stands for the constant family  $\partial\Delta^k$ -parametrized family with value  $\phi$ ). We thus have the string of homotopies

$$\phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \sim \phi^{-1}\phi = \text{id}_N = \phi\phi^{-1} \sim \Phi_N^Q|_{\partial\Delta^k} \circ \phi^{-1}. \quad (8.10)$$

We may prove the second claim as follows. The following are all homotopies and equalities as maps from  $\partial\Delta^k$  to  $\text{Emb}_{\text{Liou}}(M, N)$ :

$$\begin{aligned} \Gamma_c|_{\partial\Delta^k} &= \phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \circ \phi^{-1} \circ \Phi_M^Q|_{\partial\Delta^k} \\ &\sim \text{id}_N \circ \phi^{-1} \circ \Phi_M^Q|_{\partial\Delta^k} \\ &= \phi^{-1} \circ \Phi_N^Q|_{\partial\Delta^k} \circ \Phi_M^N|_{\partial\Delta^k} \\ &= \Gamma_a|_{\partial\Delta^k \times \Delta^{\{0\}}} \end{aligned}$$

The lone homotopy is from (8.10), while the non-trivial equality is by the composition assumption of being a simplex in  $\text{Liou}_\Delta$ . (See (LD1).)  $\square$

**Remark 8.14.** Because  $\Phi_N^Q|_{\partial\Delta^k}$  is homotopic, as a family, to  $\phi$ , the above lemma is just a homotopical version of the (obvious) statement that any isomorphism commutes with its inverse.

## 8.4 Proof of surjection on homotopy groups

In the present section, we will use  $|\Delta^k|$  to mean the topological space, and  $\Delta^k$  to mean the simplicial set. (Both are rightly called  $k$ -simplices.)

*Proof.* Let  $\gamma_0$  be a representative of an element of  $\pi_{k-1}(\text{hom}^R(M, N), f)$ . We may assume that  $f$  is in the image of the map  $\text{hom}'_{\text{Liou}_\Delta}(M, N) \rightarrow \text{hom}^R_{\text{Liou}_\Delta}(M, N)$  by the surjectivity on  $\pi_0$ .

We must exhibit an element  $\gamma_1$  in  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$ , together with a simplex exhibiting a homotopy from  $\gamma_0$  to  $\gamma_1$ . Before we do this, let us parse the hypotheses.

**Remark 8.15** (Parsing the data of  $\gamma_0$ ). As usual, we will ignore the data of the Liouville structures on the  $\Delta$ -movies. By definition, the data of  $\gamma_0$  is a  $k$ -simplex in  $\text{Liou}_\Delta$  whose  $k$ th face is the degenerate  $(k-1)$ -simplex associated to  $M$ , and whose other faces are the degenerate simplices  $s_0 \circ \dots \circ s_0(f)$  associated to  $f$ . So, by Example 8.4,  $\gamma_0$  gives rise to the following data:

(i) A Liouville sector  $Q = \gamma_0([k+1])$ .

(ii) A map

$$\Phi_N^Q : \Delta_k^k \rightarrow \text{Emb}_{\text{Liou}}(N, Q) \quad (8.11)$$

which, at the vertex  $\Delta^{\{k\}}$  specifies a diffeomorphism

$$\phi : N \rightarrow Q \quad (8.12)$$

and

(iii) A map

$$\Phi_M^Q : \Delta^{k-1} = \partial_k \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, Q) \quad (8.13)$$

satisfying the condition

$$\Phi_M^Q|_{\partial \Delta^{k-1}} = \Phi_N^Q|_{\partial \Delta^{k-1}} \circ f \quad (8.14)$$

where  $f$  is treated as the constant map  $\partial \Delta^{k-1} \rightarrow \text{Emb}_{\text{Liou}}(M, N)$ .

How do we exhibit  $\gamma_1$ ? By definition,  $\gamma_1$  must be a  $k$ -simplex in  $\text{Liou}_\Delta(M, N)$  satisfying the following boundary conditions:

- (a) The  $k$ th face is the degenerate  $(k-1)$ -simplex generated by the object  $M$ ,
- (b) All other faces are degenerate  $(k-1)$ -simplices  $s_0 \circ \dots \circ s_0(f)$  generated by the morphism  $f$ .
- (c) Moreover, because  $\gamma_1$  is in  $\text{hom}'_{\text{Liou}_\Delta}(M, N)$ , we conclude  $\gamma_1([k]) = N$  and  $\gamma_1(I \subset [k]) \cong \text{id}_N$  for any  $I \subset [k]$  with  $k \in I$ . So the data of  $\gamma_1$  is specified entirely by a map

$$\Psi_M^N : \gamma_1([k-1] \subset [k]) : \Delta^{k-1} \rightarrow \text{Emb}_{\text{Liou}}^{\text{coll}}(M, N)$$

(along with some Liouville structure on  $N \times T^* \Delta^k$ ).

So let us produce  $\Psi_M^N$ . As usual, the extra data of a Liouville structure on  $N \times T^* \Delta^k$  is an afterthought; we ignore it.

**Construction 8.16** ( $\Psi_M^N$ ). Note that we have a map

$$\overline{\Phi_N^Q \circ f} : \Delta^1 \times \partial \Delta^{k-1} \rightarrow \partial \Delta^{k-1} \xrightarrow{\Phi_N^Q|_{\partial \Delta^{k-1}} \circ f} \text{Emb}_{\text{Liou}}(M, Q)$$

where the first arrow is the projection forgetting the  $\Delta^1$  factor. Because of (8.14), and by collaring, we may glue this map with  $\Phi_M^Q$  to obtain a single map

$$\overline{\Phi_N^Q \circ f} \bigcup \Phi_M^Q : \Delta^1 \times \partial \Delta^{k-1} \bigcup_{\partial \Delta^{k-1}} \Delta^{k-1} \rightarrow \text{Emb}_{\text{Liou}}(M, Q).$$

Now choose a diffeomorphism

$$j : |\Delta^{k-1}| \cong |\Delta^1 \times \partial\Delta^{k-1}|$$

respecting the collars. We define

$$\Psi_M^N := \phi^{-1} \circ \left( \overline{\Phi_N^Q \circ f} \bigcup \Phi_M^Q \right) \circ j$$

where  $\phi^{-1}$  is the inverse to (8.12).

**Remark 8.17.** By construction,  $\Psi_N^M$  is equal to  $f$  near the boundary of  $|\Delta^{k-1}|$ . Moreover,  $\Psi_N^M$  satisfies the following:

In some interior region of  $|\Delta^{k-1}|$ ,  $\Psi_N^M$  encodes (up to reparametrization) the family  $\phi^{-1} \circ \Phi_M^Q$ . (8.15)

So now we construct a  $(k+1)$ -simplex  $H$  in  $\text{Liou}_\Delta$  realizing a homotopy from  $\gamma_0$  to  $\gamma_1$  in  $\text{hom}_{\text{Liou}_\Delta}^R(M, N)$ . As it turns out, we will be able to define  $H$  so that  $H([k+1]) = Q$  and so that the embedding

$$H([k+1] \setminus \{0\} \subset [k+1]) : \partial_0 \Delta^{k+1} \rightarrow \text{Emb}_{\text{Liou}}(Q, Q)$$

is the constant map to  $\text{id}_Q$ . Then, unwinding the definitions, we must only specify the following data to specify  $H$ :

- $\Gamma_N^Q : \partial_1 \Delta^{k+1} \rightarrow \text{Emb}_{\text{Liou}}(N, Q)$
- $(\Xi_N^Q)_i : \partial_i \Delta^{k+1} \rightarrow \text{Emb}_{\text{Liou}}(N, Q)$  for  $2 \leq i \leq k$
- $\Gamma_M^Q : \Delta^k = \partial_{k+1} \Delta^{k+1} \rightarrow \text{Emb}_{\text{Liou}}(M, Q)$

(along with a Liouville structure on  $Q \times T^* \Delta^{k+1}$ , which we again ignore because one may be constructed straightforwardly).

**Construction 8.18** ( $\Gamma_N^Q$ ). Because  $\text{Emb}_{\text{Liou}}(N, Q)$  is a Kan complex, we have a horn-filler

$$\begin{array}{ccc} \Lambda_k^k & \xrightarrow{\Phi_N^Q(8.11)} & \text{Emb}_{\text{Liou}}(N, Q) \\ \downarrow & \searrow \widetilde{\Phi_N^Q} & \nearrow \\ \Delta^k & \dashrightarrow & \end{array}$$

Again by Proposition 4.16, we may guarantee that the filler  $\widetilde{\Phi_N^Q}$  is collared. In fact, if we choose this filler using a “project-to- $\partial_k \Delta^k$ -and-interpolate” construction, we may guarantee that the value of  $\widetilde{\Phi_N^Q}$  has the following boundary behavior along  $\partial_k \Delta^k$ :

There is some region in the interior of  $|\partial_k \Delta^k| \cong |\Delta^{k-1}|$  where  $\widetilde{\Phi_N^Q}$  is identical to  $\phi$ . (8.16)

We choose  $\widetilde{\Phi_N^Q}$  so that this region (8.16) contains the region referenced in (8.15). We let  $\Gamma_N^Q$  be the composition

$$\partial_1 \Delta^{k+1} \xrightarrow{\cong} \Delta^k \xrightarrow{\widetilde{\Phi_N^Q}} \text{Emb}_{\text{Liou}}(N, Q)$$

where the first arrow is the unique isomorphism of simplicial sets.

**Construction 8.19**  $((\Xi_N^Q)_i)$ . For every  $2 \leq i \leq k$ , we define the  $(\Xi_N^Q)_i$  to be the composition

$$\partial_i \Delta^{k+1} \cong \Delta^k \xrightarrow{\sigma_0} \Delta^{k-1} \cong \partial_i \Delta^k \xrightarrow{\Phi_N^Q} \text{Emb}_{\text{Liou}}(N, Q)$$

where  $\sigma_0$  is the codegeneracy map, induced by the surjection  $[k] \rightarrow [k-1]$  sending both  $0, 1 \in [k]$  to  $0 \in [k-1]$ .

**Construction 8.20**  $(\Gamma_M^Q)$ . Based on our previous choices, the value of  $\Gamma_M^Q : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, Q)$  is completely constrained on  $\partial \Delta^k$ . For example, on  $\partial_1 \Delta^k$ , the composition requirement for a  $(k+1)$ -simplex in  $\text{Liou}_\Delta$  demands that we have

$$\Gamma_M^Q|_{\partial_1 \Delta^k} = \Gamma_N^Q|_{\partial_1 \Delta^k} \circ \Phi_M^N|_{\partial_1 \Delta^k \cong \Delta^{k-1}}$$

We make an important observation. Because the regions of (8.16) was chosen to contain the region of (8.15), one can choose a diffeomorphism

$$|\partial_1 \Delta^k| \cong |\Delta^{k-1}| \bigcup |\partial \Delta^{k-1} \times \Delta^1| \quad (8.17)$$

so that

$$\Gamma_M^Q|_{\partial_1 \Delta^k} \text{ encodes a homotopy } h \text{ from } \Phi_M^Q|_{\partial \Delta^{k-1}} \text{ to itself.}$$

In particular, by choosing a reverse homotopy  $\bar{h}$ , there is a map

$$H'_1 : \partial \Delta^{k-1} \times \Delta^2 \rightarrow \text{Emb}_{\text{Liou}}(M, Q) \iff \Delta^2 \rightarrow \text{hom}_{\text{sSet}}(\partial \Delta^{k-1}, \text{Emb}_{\text{Liou}}(M, Q))$$

where the 2-simplex has boundary conditions  $h$ ,  $\bar{h}$ , and a constant map along the edges  $\Delta^{\{0,1\}}$ ,  $\Delta^{\{1,2\}}$ , and  $\Delta^{\{0,2\}}$ , respectively. Informally, the two-simplex thus encodes a homotopy from the concatenation  $\bar{h} \# h$  to the constant homotopy at  $\Phi_M^Q|_{\partial \Delta^{k-1}}$ .

Now let  $H'_2$  be the composition

$$H'_2 : \Delta^{k-1} \times \Delta^1 \rightarrow \Delta^{k-1} \xrightarrow{\Phi_M^Q} \text{Emb}_{\text{Liou}}(M, Q)$$

and choose a piecewise smooth, appropriately collared diffeomorphism

$$l : |\Delta^{k-1} \times \Delta^1| \bigcup |\partial \Delta^{k-1} \times \Delta^2| \cong |\Delta^{k-1} \times \Delta^1| \quad (8.18)$$

where the union  $|\Delta^k \times \Delta^{\{0\}}| \bigcup |\partial \Delta^k \times \Delta^{\{0,1\}}|$  of the domain is mapped to the face  $|\Delta^k \times \Delta^{\{0\}}|$  of the codomain. (See Figure 2.) Now we note that the map

$$(H'_2 \bigcup H'_1) \circ l^{-1} : |\Delta^{k-1} \times \Delta^1| \rightarrow \text{Emb}_{\text{Liou}}(M, Q) \quad (8.19)$$

is constant along the  $\Delta^1$ -direction of  $|\partial \Delta^{k-1} \times \Delta^1|$ . Consequently, (8.19) factors through a quotient

$$|\Delta^{k-1} \times \Delta^1| \rightarrow |\Delta^k|$$

(for example, by writing  $|\Delta^k|$  as a cone on  $|\Delta^{k-1}|$ ); we may again apply Proposition 4.16 to smooth and collar the resulting map. We define the resulting factorization to be

$$\Gamma_M^Q : \Delta^k \rightarrow \text{Emb}_{\text{Liou}}(M, Q).$$

We leave to the reader the details of verifying that all compositional requirements are met. □




Figure 2: The piecewise smooth isomorphism (8.18)

## 8.5 Proof of Theorems 8.1 and B.

*Proof.* We have the following homotopy equivalences of Kan complexes:

$$\begin{array}{ccccc} \widetilde{\text{Emb}}_{\text{Liou}}^{\text{coll}}(M, N) & \xrightarrow{\text{Rem 8.7}} & \text{hom}'_{\text{Liou}_\Delta}(M, N) & \xrightarrow{\text{Lem 8.11}} & \text{hom}_{\text{Liou}_\Delta}^R(M, N) \\ \text{Prop 8.9} \downarrow & & & & \\ \text{Emb}_{\text{Liou}}^{\text{coll}}(M, N) & \xrightarrow{\text{Prop 4.17}} & \text{Emb}_{\text{Liou}}(M, N). & & \end{array}$$

□

**Remark 8.21.** We point out one subtlety, which is that the simplicial set structure used in Lemma 8.11 is not necessarily the simplicial set structure used in Proposition 8.9; more specifically, the isomorphism of semisimplicial sets in Remark 8.7 need not respect degeneracy maps.

However, if two Kan complexes have the same underlying semisimplicial sets (i.e., with differing degeneracy maps but with equal face maps), they are automatically homotopy equivalent.<sup>26</sup> So the reader should interpret the arrow labeled by Remark 8.7 not as “the” map constructed in the remark, but as a homotopy equivalence guaranteed by it.

## 8.6 An informal argument

In Theorem 8.1, we proved that  $\text{Liou}_\Delta$  has morphism spaces that are homotopy equivalent to the usual spaces of sectorial embeddings. One can do better. Let us give an informal argument, using the language of constructible sheaves, as to why  $\text{Liou}_\Delta$  is equivalent to the usual  $\infty$ -category of sectors with spaces of sectorial embeddings. (The only difference between this “better” result and Theorem 8.1 is the exhibition of a *functor* resulting in the equivalences of morphism spaces guaranteed in Theorem 8.1.)

Fix an  $I$ -simplex of  $\text{Liou}_\Delta$ . For all  $\emptyset \neq A \subset I$ , If one forgets the sectorial structures on  $M_A \times T^*\Delta^A$  (and only remembers the sectorial structure on  $M_A$ ), one obtains families of sectorial embeddings  $\Delta^A \rightarrow \text{Emb}_{\text{Liou}}(M_A, M_B)$ . We note that the assignment

- To every subsimplex  $\Delta^A$  of  $\Delta^I$ , an object  $M_A$ , and
- To every inclusion  $\Delta^A \subset \Delta^B$ , a family of maps  $\Delta^A \rightarrow \text{Emb}_{\text{Liou}}(M_A, M_B)$

<sup>26</sup>See for example Theorem 1.4 of [30], which proves a more general result—two  $\infty$ -categories with the same underlying semisimplicial sets are equivalent.

determines a constructible sheaf on  $\Delta^I$ , where  $\Delta^I$  is stratified in the usual way.

**Remark 8.22.** The stratification is as follows. Given a point  $x \in \Delta^I \subset \mathbb{R}^I$ ,  $x$  is assigned to the set of those  $i \in I$  for which  $x_i > 0$ . This assignment from  $x$  to subsets of  $I$  defines a stratification on  $\Delta^I$  by endowing the power set of  $I$  the Alexandroff topology.

For more on this perspective on stratified spaces, we refer the reader to [18, Appendix A], [5] and [34].

Moreover, because we demand that our  $I$ -simplices have a max-localizing property, it is formal that this constructible sheaf is constructible when considered as a sheaf on  $\Delta^I$  with the “max stratification” (where now  $\Delta^I$  is stratified by the poset  $I$  itself, and  $x$  is sent to the element  $\max\{i, x_i > 0\}$ ). It is easy to see that the  $\infty$ -category of constructible sheaves on  $\Delta^I$  with the max stratification is equivalent to the  $\infty$ -category of functors from  $I$  to the  $\infty$ -category in which the sheaf is taking values.

Put another way, we see that every  $I$ -simplex of  $\mathrm{Liou}_\Delta$  gives rise to an  $I$ -simplex in the usual  $\infty$ -category of Liouville sectors<sup>27</sup>. Thus, by varying the  $I$ -variable,  $\mathrm{Liou}_\Delta$  admits a map to the  $\infty$ -category of Liouville sectors; this assignment is redundant in that different Liouville structures on  $M_A \times T^*\Delta^A$  (with a fixed structure near  $\max A$ ) may be sent to the same simplices; but the space of these Liouville structures (with fixed structure near  $\max A$ ) is contractible because, by assumption, these deformations are all compactly supported.

Let us make one remark:  $\mathrm{Liou}_\Delta$  does not see all constructible sheaves on the simplex. This is because we demand that along max-localizing inclusions, the morphism  $M_A \rightarrow M_B$  be a *diffeomorphism* along the  $\max A = \max B$  vertex, as opposed to just a sectorial equivalence. However, it is straightforward to verify that, for any constructible sheaf consisting of max-localizing morphisms that are sectorial equivalences (as opposed to sectorial isomorphisms), one can construct a constructible sheaf on a higher-dimensional simplex parametrizing to a simplex whose max-localizing morphisms are all sectorial *isomorphisms* along the relevant vertices.

This is an informal argument. We included our informal argument to give the reader peace of mind that, indeed,  $\mathrm{Liou}_\Delta$  has a moral reason to be equivalent to the  $\infty$ -category of Liouville sectors. The combinatorics to make the argument rigorous will be left for another work.

---

<sup>27</sup>This may be obtained, for example, by topologically enriching the category of Liouville sectors, then taking the homotopy coherent nerve.

## 9 $\text{Liou}_\Delta$ and $\text{Liou}_\Delta^{\mathcal{D}\text{ef}}$ are equivalent

**Notation 9.1** ( $\text{Emb}_{\text{Liou}}^{\mathcal{D}\text{ef}}(M, N)$ ). Fix Liouville sectors  $M$  and  $N$ . We let  $\text{Emb}_{\text{Liou}}^{\mathcal{D}\text{ef}}(M, N)$  denote the singular complex of the space of deformation embeddings  $\mathcal{E}\text{mb}_{\text{Liou}}^{\mathcal{D}\text{ef}}(M, N)$ .

We have the following analogue of Theorem 8.1:

**Theorem 9.2.** There is a homotopy equivalence of simplicial sets

$$\text{Emb}_{\text{Liou}}^{\mathcal{D}\text{ef}}(M, N) \simeq \text{hom}_{\text{Liou}_\Delta^{\mathcal{D}\text{ef}}}(M, N).$$

*Proof.* The proof is nearly identical to that of Theorem 8.1; the same strategy to extending (non-compactly-supported) families of Liouville structure of an embedded image to the rest of the codomain as in Remark 6.11 applies. For the reader's sake, we do not reproduce the arguments.  $\square$

Finally, we have:

**Theorem 9.3.** The inclusion  $\text{Liou}_\Delta \rightarrow \text{Liou}_\Delta^{\mathcal{D}\text{ef}}$  (which is a map of  $\infty$ -categories by Theorem 7.11) is an equivalence of  $\infty$ -categories.

*Proof.* The inclusion is a bijection on objects, so it remains to show that it induces a homotopy equivalence of mapping spaces. For this, fix two sectors  $M$  and  $N$  and note the equivalences

$$\text{hom}_{\text{Liou}_\Delta}^R(M, N) \xrightarrow[\sim]{Th\ 8.1} \text{Emb}_{\text{Liou}}(M, N) \xrightarrow[\sim]{Th\ 4.18} \text{Emb}_{\text{Liou}}^{\mathcal{D}\text{ef}}(M, N) \xrightarrow[\sim]{Th\ 9.2} \text{hom}_{\text{Liou}_\Delta^{\mathcal{D}\text{ef}}}^R(M, N).$$

It is a tedious but straightforward verification that, at the level of  $\pi_0$  and of homotopy groups, the composition of these equivalences equals the map induced by the inclusion  $\text{Liou}_\Delta \rightarrow \text{Liou}_\Delta^{\mathcal{D}\text{ef}}$ .  $\square$

## 10 Localization (the proof of Theorem C)

In this section, we will prove Theorem C—that  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$  is equivalent to the localization  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathbf{eq}^{\diamond})^{-1}]$ .

Recall the category of stabilized sectors  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$  from Notation 1.20. There, in defining  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$ , we stabilized  $\mathrm{Liou}_{\mathrm{str}}$  by the object  $T^*[0, 1]$ . Because  $T^*[0, 1]$  and  $T^*[-1, 0]$  are obviously isomorphic objects (for example, induced by the translation diffeomorphism from  $[0, 1]$  to  $[-1, 0]$ ), we adopt a different model here. (See Remark 10.9 for why prefer  $[-1, 0]$  over  $[0, 1]$ ). Stabilizing by either interval obviously yields equivalent  $\infty$ -categories, so we abuse notation:

**Notation 10.1.** We define

$$\mathrm{Liou}_{\mathrm{str}}^{\diamond} := \mathrm{colim} \left( \mathrm{Liou}_{\mathrm{str}} \xrightarrow{T^*[-1, 0] \times -} \mathrm{Liou}_{\mathrm{str}} \xrightarrow{T^*[-1, 0] \times -} \dots \right) \quad (10.1)$$

and

$$(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond} := \mathrm{colim} \left( \mathrm{Liou}_{\Delta}^{\mathrm{Def}} \xrightarrow{T^*[-1, 0] \times -} \mathrm{Liou}_{\Delta}^{\mathrm{Def}} \xrightarrow{T^*[-1, 0] \times -} \dots \right). \quad (10.2)$$

**Remark 10.2.** The techniques of this section come down to combinatorial arguments enabling Constructions 10.16 and 10.29. Indeed, it should be true that for any  $\infty$ -category  $\mathcal{C}$  with a notion of “simplex object”, and with a class of morphisms  $\mathbf{eq}$  that pick out inclusions of faces of a simplex, the methods here carry over to show that the localization  $\mathcal{C}[\mathbf{eq}^{-1}]$  is equivalent to an  $\infty$ -category  $\mathcal{C}_{\Delta}$  built out of certain barycentric subdivision diagrams in  $\mathcal{C}$ .

### 10.1 Defining $\beta$

**Construction 10.3** ( $j$ ). Recall the map  $\mathrm{Liou}_{\mathrm{str}} \rightarrow \mathrm{Liou}_{\Delta}^{\mathrm{Def}}$  from Construction 7.6; it is in fact a map of  $\infty$ -categories (i.e., simplicial sets) by combining Theorems 7.8 and 7.11. This map is compatible with taking direct product with a fixed sector; in particular, we may pass to the stabilizations to obtain a map of  $\infty$ -categories

$$j : \mathrm{Liou}_{\mathrm{str}}^{\diamond} \rightarrow (\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}.$$

**Example 10.4** ( $j$  on 1-simplices). Fix a 1-simplex  $f_{01} : X_0 \rightarrow X_1$  in  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$ . Then  $j(f_{01})$  is a 1-simplex in  $(\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond}$  whose underlying diagram may be drawn as follows:

$$X_0 \cong X_0 \times T^*[-1, 0] \xrightarrow{f_{01} \times T^*\eta} X_1 \times T^*\Delta^1 \xleftarrow{\mathrm{id}_{X_1} \times T^*\eta} X_1 \times T^*[-1, 0] \cong X_1.$$

To reduce clutter, we will suppress the  $T^*[-1, 0]$  factors of stabilizations, and the  $\mathrm{id}_{X_1}$  factors of functions, to depict the above diagram as

$$X_0 \xrightarrow{f_{01} \times T^*\eta} X_1 \times T^*\Delta^1 \xleftarrow{T^*\eta} X_1.$$

**Example 10.5** ( $j$  on 2-simplices). Fix a 2-simplex in  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$ —this is the data of sectors  $X_0, X_1, X_2$  and morphisms  $f_{01}, f_{12}, f_{02}$  with  $f_{02} = f_{12} \circ f_{01}$  (up to stabilization). Then  $j$  of this 2-simplex may be depicted as the following barycentric subdivision diagram:

$$\begin{array}{ccccc}
 & & X_2 & & \\
 & \swarrow T^*\eta & \downarrow T^*\eta & \searrow T^*\eta & \\
 X_2 \times T^*\Delta^1 & \xrightarrow{T^*\eta} & X_2 \times T^*\Delta^2 & \xleftarrow{T^*\eta} & X_2 \times T^*\Delta^1 \\
 \uparrow f_{02} \times T^*\eta & & \uparrow f_{12} \times T^*\eta & & \uparrow f_{12} \times T^*\eta \\
 X_0 & \xrightarrow{f_{01} \times T^*\eta} & X_1 \times T^*\Delta^1 & \xleftarrow{T^*\eta} & X_1
 \end{array}$$



Note we are suppressing the identifications  $X_0 = X_0 \times T^*[-1, 0] = X_0 \times T^*[-1, 0]^2$ . We are also suppressing subscripts on  $\eta$  – that  $\eta \circ \eta = \eta$  should be interpreted as  $\eta_{J \subset K} \circ \eta_{I \subset J} = \eta_{I \subset K}$ , for example.

**Proposition 10.6.** Let  $\mathfrak{eq} \subset \mathrm{Liou}_{\mathrm{str}}^\diamond$  denote those strict (stabilized) sectorial embeddings which are also equivalences of Liouville sectors (Definition 2.30). Then the functor  $j$  (Construction 10.3) factors, up to homotopy, through the localization  $\mathrm{Liou}_{\mathrm{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]$ .

*Proof.* Any equivalence of Liouville sectors is rendered homotopy-invertible in  $\mathrm{Liou}_\Delta$  (Proposition 7.14), and hence in  $\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}}$  (Theorem 7.11(b)). Further, any morphism which is homotopy-invertible in a finite stage (such as  $\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}}$ ) of an increasing union of  $\infty$ -categories (such as  $(\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}})^\diamond$ ) remains homotopy-invertible in the union. Thus the claim follows by the universal property of localizations.  $\square$

**Notation 10.7**  $(\iota, j, \beta)$ . By Proposition 10.6, letting  $\iota$  denote the localization map, we have a homotopy-commuting diagram of  $\infty$ -categories

$$\begin{array}{ccc} \mathrm{Liou}_{\mathrm{str}}^\diamond & \xrightarrow{j} & (\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}})^\diamond \\ \downarrow \iota & \nearrow \beta & \\ \mathrm{Liou}_{\mathrm{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}] & & \end{array} \quad (10.3)$$

where  $\beta$  is a morphism guaranteed by the universal property of localizations.

Theorem C is the claim that  $\beta$  is an equivalence. To see  $\beta$  is an equivalence, we will exhibit an equivalence  $\alpha$  for which  $\alpha \circ \beta$  is also an equivalence.

## 10.2 Defining $\alpha$

**Remark 10.8.** Let  $\mathrm{sd}(I)$  be the barycentric subdivision of  $I$  (thought of as a simplicial set). The data of an  $I$ -simplex of  $\mathrm{Liou}_\Delta$  determines a functor

$$\mathrm{sd}(I) \rightarrow \mathrm{Liou}_{\mathrm{str}}^\diamond$$

which assigns to a subset  $I' \subset I$  the object

$$M_{I'} \times T^* \Delta^{I'} = T^*[-1, 0]^n \otimes M_{I'} \times T^* \Delta^{I'} \cong M_{I'} \times T^* \Delta^{I'} \otimes T^*[-1, 0]^n$$

of  $\mathrm{Liou}_{\mathrm{str}}^\diamond$ .

Likewise, the data of an  $I$ -simplex of  $\mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}}$  determines a functor  $\mathrm{sd}(I) \rightarrow \mathrm{Liou}_{\mathrm{str}}^\diamond$ .

**Remark 10.9.** Remark 10.8 is the reason that we stabilize by  $T^*[-1, 0]$  instead of  $T^*[0, 1]$ ; our collarings  $\eta$  were by  $[-1, 0]$ . In turn, we chose to collar by  $[-1, 0]$  because of the particulars of Construction 4.6. If the collarings were given by embeddings of  $[a, b]^n$ , then we would stabilize by  $T^*[a, b]$ .

**Remark 10.10.** Remark 10.8 exhibits maps of semisimplicial sets

$$\mathrm{Liou}_\Delta \rightarrow \mathrm{Ex}(\mathrm{Liou}_{\mathrm{str}}^\diamond), \quad \mathrm{Liou}_\Delta^{\mathcal{D}\mathrm{ef}} \rightarrow \mathrm{Ex}(\mathrm{Liou}_{\mathrm{str}}^\diamond).$$

(By abuse of notation, the codomain is the semisimplicial set obtained by forgetting the degeneracy maps of the simplicial set  $\text{Ex}(\text{Liou}_{\text{str}}^\diamond)$ .) In fact, we may identify the stabilization  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  with a semisimplicial subset of  $\text{Ex}(\text{Liou}_{\text{str}}^\diamond)$ , so we obtain a map

$$(\text{Liou}_\Delta^{\text{Def}})^\diamond \rightarrow \text{Ex}(\text{Liou}_{\text{str}}^\diamond) \quad (10.4)$$

of semisimplicial sets. We caution, however, that the image of  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  is not a simplicial subset. This is because the image inside  $\text{Ex}(\text{Liou}_{\text{str}}^\diamond)$  is not closed under degeneracy maps.

**Construction 10.11** ( $\alpha$  and  $\alpha_{\text{semi}}$ ). By post-composing (10.4) with  $\text{Ex}(\iota)$ , we obtain a map of semisimplicial sets  $(\text{Liou}_\Delta^{\text{Def}})^\diamond \rightarrow \text{Ex}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}])$ . Condition (LD2) guarantees that this map factors through  $\text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}])$ —this is because if  $y : M \rightarrow N$  is a strict sectorial isomorphism, then any sectorial embedding of the form  $y \otimes T^* \eta : M \otimes T^*[-1, 0]^k \rightarrow N \otimes T^* \Delta^k$  is an equivalence of Liouville sectors (Example 2.31). And more generally, any map of the form  $\phi_{A' \subset A}$  is also a sectorial equivalence when  $\max A' = \max A$ . See Remark 12.7.

Thus, we have a map of semisimplicial sets

$$\alpha_{\text{semi}} : (\text{Liou}_\Delta^{\text{Def}})^\diamond \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}]). \quad (10.5)$$

On the other hand, the degeneracy maps of the simplicial set  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  are such that the map  $j : \text{Liou}_{\text{str}}^\diamond \rightarrow (\text{Liou}_\Delta^{\text{Def}})^\diamond$  is a simplicial set map (Construction 10.3). This guarantees that the semisimplicial set map (10.5) sends degenerate edges of the domain to equivalences (in fact, degenerate edges) in the codomain. So Theorem 3.7 guarantees the existence of a map of *simplicial* sets

$$\alpha : (\text{Liou}_\Delta^{\text{Def}})^\diamond \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}]).$$

There is moreover a homotopy (unique up to contractible choice) from  $(\alpha_{\text{semi}})_+$  to  $\alpha \circ \epsilon$ , as in (3.7) for  $h = \alpha$ .

**Remark 10.12.** The map  $\alpha_{\text{semi}}$  in (10.5) is explicit. By definition, a  $k$ -simplex in  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$  is a diagram in the shape of a barycentric subdivision of  $k$ -simplex; a  $k$ -simplex in the codomain is also some diagram in the shape of a barycentric subdivision. The construction above is nothing more than a simple check that every such  $k$ -simplex in the domain is an example of a  $k$ -simplex in the codomain.

Moreover, every statement about  $\alpha$  will be deduced by providing explicit constructions involving  $\alpha_{\text{semi}}$ .

### 10.3 $\alpha$ as a left inverse

**Lemma 10.13.** The composition

$$\text{Liou}_{\text{str}}^\diamond \xrightarrow{j} (\text{Liou}_\Delta^{\text{Def}})^\diamond \xrightarrow{\alpha} \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}])$$

is homotopic to the composition

$$\text{Liou}_{\text{str}}^\diamond \xrightarrow{\iota} \text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}] \xrightarrow{m} \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\epsilon q^\diamond)^{-1}]).$$

Assuming Lemma 10.13 for the moment, we have the following consequence:

**Lemma 10.14.**  $\alpha \circ \beta$  is an equivalence of  $\infty$ -categories.

*Proof.* We have a diagram in the  $\infty$ -category of  $\infty$ -categories as follows:

$$\begin{array}{ccc}
\mathrm{Liou}_{\mathrm{str}}^{\diamond} & \xrightarrow{j} & (\mathrm{Liou}_{\Delta}^{\mathrm{Def}})^{\diamond} \\
\downarrow \iota & \nearrow \beta & \downarrow \alpha \\
\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}] & \xrightarrow{\alpha \circ \beta} & \mathrm{Ex}_{\sim}(\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}]).
\end{array}$$

The lower-right triangle commutes on the nose as a diagram of simplicial sets, while the upper-left triangle (Notation 10.7) commutes up to homotopy specified by the universal property of localization.

In what follows, we write  $\sim$  to mean “is homotopic to:”

$$m \circ \iota \sim \alpha \circ j \sim \alpha \circ \beta \circ \iota.$$

The first homotopy is by Lemma 10.13. The next homotopy follows because we know that  $j$  is homotopic to  $\beta \circ \iota$  by definition (Notation 10.7). By the universal property of localizations, we see that  $\alpha \circ \beta$  is homotopic to  $m$ . We conclude by noting that  $m$  is an equivalence (Proposition 3.19).  $\square$

The rest of Section 10.3 is devoted to proving Lemma 10.13. As mentioned in Remark 3.34, the map  $\alpha$  is rather inexplicit, while  $\alpha_{\mathrm{semi}}$  has a known simplex-by-simplex description. So with the goal of utilizing Lemma 3.36, we construct a map

$$G_f : \Delta^1 \times \Delta^k \rightarrow \mathrm{Ex}_{\sim}(\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}])$$

for every  $k$ -simplex  $f : \Delta^k \rightarrow \mathrm{Liou}_{\mathrm{str}}^{\diamond}$ , satisfying the hypotheses of Lemma 3.36.

**Notation 10.15** ( $A$ ). Recall that a top-dimensional simplex of  $\Delta^1 \times \Delta^k$  is uniquely determined by any of the following equivalent data:

1. A length  $k+1$  chain in the product poset  $[k] \times [1]$ . By noting that any such chain must be of the form

$$(0, 0) < (0, 1) < \dots < (0, j) < (1, j) < (1, j+1) < \dots < (1, k) \quad (10.6)$$

we see that the data of this chain is in turn equivalent to:

2. A choice of element  $j \in [k]$ . In turn, a choice of  $j$  determines:
3. A choice of convex subset  $A \subset [k]$  for which  $\min A = 0$ . Here, convexity means if  $p, q \in A$  then  $[p, q] \subset A$ . Any such  $A$  is of the form  $A = \{i \mid 0 \leq i \leq j\}$  for some  $j \in [n]$ .

**Construction 10.16** ( $G_f$ ). Fix a  $k$ -simplex  $f$  in  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}$  – this fixes, up to stabilization by finitely many dimensions, a choice of Liouville sector

$$X_i$$

for each  $i \in [k]$ , and strict sectorial embeddings

$$f_{i,j} : X_i \rightarrow X_j$$

for each  $i < j$ , satisfying  $f_{j,h} f_{i,j} = f_{i,h}$ . (In the case  $i = j$ , we have  $f_{ii} = \mathrm{id}_{X_i}$ .) We fix also a top-dimensional simplex  $\Delta^{k+1} \xrightarrow{A} \Delta^1 \times \Delta^k$ . Note that  $A$  determines the surjection

$$s_j : [k+1] \rightarrow [k]$$

via the composition  $[k+1] \xrightarrow{A} [1] \times [k] \rightarrow [k]$  (see Notation 10.15).

Given the data of  $(A, f)$ , we associate:

- To every non-empty subset  $I \subset [k+1]$ , the (stabilized class of the) Liouville sector

$$X_{\max s_j(I)} \times T^* \Delta^{I/A}$$

where  $I/A$  refers to the linear poset obtained from  $I$  by declaring any two elements in  $I \cap A$  to be equivalent. (There is an induced poset structure on this quotient set by virtue of  $A$  being convex.) The simplex  $\Delta^{I/A}$  is defined for any linear poset  $I/A$  as in (4.1) .

- To every inclusion  $I \subset J \subset [k+1]$  with  $I$  non-empty, the strict sectorial embedding

$$f_{\max s_j(I), \max s_j(J)} \times T^* \eta : X_{\max s_j(I)} \times T^* \Delta^{I/A} \times T^* [-1, 0]^{(J/A) \setminus (I/A)} \rightarrow X_{\max s_j(J)} \times T^* \Delta^{J/A}. \quad (10.7)$$

Here,  $\eta$  is the collaring associated to the inclusion  $I/A \hookrightarrow J/A$  (see Definition 4.3 and Choice 4.9).

We leave it to the reader to verify that for every  $(A, f)$ , the above data define a map

$$G_f|_A : \text{sd}(\Delta^{k+1}) \rightarrow \text{Liou}_{\text{str}}^\diamond$$

and hence a map to  $\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]$ . It is somewhat abusive to write  $G_f|_A$  at this stage as we have not defined  $G_f$  yet; but we justify this notation now: One straightforwardly checks that, by fixing  $f$  and varying  $A$ , the  $G_f|_A$  glue together to define a single map

$$\text{colim}_A(G_f|_A) : \text{colim}_A \Delta^{k+1} \cong \Delta^1 \times \Delta^k \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]).$$

And we define

$$G_f := \text{colim}_A(G_f|_A).$$

**Example 10.17.** In the case  $k = 0$  (so  $f$  is just a choice of object  $X$  in  $\text{Liou}_{\text{str}}^\diamond$ ),  $G_f$  is the map

$$\Delta^1 \times \Delta^0 \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}])$$

encoded by a diagram  $\text{sd}(\Delta^1) \rightarrow \text{Liou}_{\text{str}}^\diamond$  we depict as follows:

$$X \times T^*[-1, 0]\{(0,0)\} \xrightarrow{\text{id}_X \times T^* \eta} X \times T^* \Delta\{(0,0), (1,0)\} \cong X \times T^* \Delta^1 \xleftarrow{\text{id}_X \times T^* \eta} X \times T^*[-1, 0]\{(1,0)\}$$

For sake of space, and because we are in the stabilization  $\text{Liou}_{\text{str}}^\diamond$ , we will simplify the above diagram and not explicitly encode the relevant posets or stabilizing intervals, and also not indicating the identity components of  $\text{id}_X \times T^* \eta$ :

$$X \xrightarrow{T^* \eta} X \times T^* \Delta^1 \xleftarrow{T^* \eta} X$$

**Example 10.18.** In the case  $k = 1$  (so  $f$  is a choice of strict sectorial embedding  $f_{01} : X_0 \rightarrow X_1$  up to stabilization),  $G_f$  is a map

$$\Delta^1 \times \Delta^1 \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}])$$

encoded by a diagram  $\text{sd}(\Delta^2) \cup \text{sd}(\Delta^2) \rightarrow \text{Liou}_{\text{str}}^\diamond$  whose two halves we informally depict as

$$\begin{array}{ccccc} X_0 & \xrightarrow{T^* \eta} & X_0 \times T^* \Delta^1 & \xleftarrow{T^* \eta} & X_0 \\ f_{01} \times T^* \eta \downarrow & & f_{01} \times T^* \eta \downarrow & & \swarrow f_{01} \times T^* \eta \\ X_1 \times T^* \Delta^1 & \xrightarrow{T^* \eta} & X_1 \times T^* \Delta^2 & \xleftarrow{T^* \eta} & X_1 \times T^* \Delta^1 \\ T^* \eta \uparrow & & \nearrow T^* \eta & & \\ X_1 & & & & \end{array}$$

and

$$\begin{array}{ccccc}
& & & & X_0 \\
& & & \swarrow f_{01} \times T^* \eta & \downarrow f_{01} \\
& & X_1 \times T^* \Delta^1 & \xrightarrow{\cong} & X_1 \times T^* \Delta^1 & \xleftarrow{T^* \eta} & X_1 \\
& \nearrow T^* \eta & & \uparrow \cong & & \uparrow \cong & \\
X_1 & \xrightarrow{T^* \eta} & X_1 \times T^* \Delta^1 & \xleftarrow{T^* \eta} & X_1
\end{array}$$

We note that the arrows labeled by  $\cong$  really are isomorphisms in  $\text{Liou}_{\text{str}}^\diamond$ ; this is because in (10.7), it may be that  $I \subsetneq J$  but that  $I/A \cong J/A$ .

*Proof of Lemma 10.13.* We will apply Lemma 3.36. In terms of the notation there, we will set

$$W = \text{Liou}_{\text{str}}^\diamond, \quad X = (\text{Liou}_\Delta^{\text{Def}})^\diamond, \quad Y = \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]), \quad i = m \circ \iota, \quad h_{\text{semi}} = \alpha_{\text{semi}}$$

(and the map  $j$  in the Lemma is the map  $j$  we are considering from Construction 10.3). We must verify that the collection  $\{G_f\}_{f: \Delta^k \rightarrow \text{Liou}_{\text{str}}^\diamond, k \geq 0}$  satisfies the hypotheses of Lemma 3.36. To this end, fix  $f: \Delta^k \rightarrow \text{Liou}_{\text{str}}^\diamond$ .

We first check condition (i) of Lemma 3.36. The  $k$ -simplex  $\Delta^{\{0\}} \times \Delta^k \subset \Delta^1 \times \Delta^k$  is the  $(k+1)$ st face of the simplex determined by  $A = [k]$ . Construction 10.16 assigns the  $k$ -simplex

$$G_f|_{\Delta^{\{0\}} \times \Delta^k} : \Delta^k \cong \Delta^{\{0\}} \times \Delta^k \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]) \quad (10.8)$$

given by a map

$$\text{sd}(\Delta^k) \rightarrow \text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]$$

described as follows:

- To every  $\emptyset \neq I \subset [k]$ , the (stabilized class of the) Liouville sector  $X_{\max I}$ .
- To every  $\emptyset \neq I \subset J \subset [k]$ , the (stabilized class of the) strict sectorial embedding  $f_{\max I, \max J} : X_{\max I} \rightarrow X_{\max J}$ . Note that when  $\max I = \max J$ , this is simply the identity morphism.

Thus the map (10.8) is precisely  $m \circ \iota(f)$ .

Let us next understand the map

$$G_f|_{\Delta^{\{1\}} \times \Delta^k} : \Delta^k \cong \Delta^{\{1\}} \times \Delta^k \rightarrow \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^\diamond[(\mathfrak{eq}^\diamond)^{-1}]) \quad (10.9)$$

(which corresponds to the choice  $A = [0]$ ). For convenience, we will identify any subset  $I \subset \{1, \dots, k+1\}$  with a subset  $I' \subset [k]$  by sending  $i \mapsto i-1$ . Then (10.9) is a diagram in the shape of a barycentric subdivision one may describe as follows:

- To every  $\emptyset \neq I \subset \{1, \dots, k+1\}$ , the (stabilized class of the) Liouville sector  $X_{\max I'} \times \Delta^{I'}$ .
- To every  $\emptyset \neq I \subset J \subset \{1, \dots, k+1\}$ , the (stabilized class of the) strict sectorial embedding

$$f_{\max I', \max J'} \times T^* \eta : X_{\max I'} \times T^* \Delta^{I'} \times T^* [-1, 0]^{J' \setminus I'} \rightarrow X_{\max J'} \times T^* \Delta^{J'}.$$

Thus the map (10.9) is precisely the barycentric subdivision diagram encoded by  $j(f)$  (see Constructions 7.6 and 10.3). So  $G_f$ , restricted to  $\Delta^{\{1\}} \times \Delta^k$ , recovers  $\alpha_{\text{semi}} \circ j(f)$ .

We have finished checking condition (i) of Lemma 3.36.

We now check condition (ii). Fix  $0 \leq i \leq k$ . Note that the top-dimensional simplices of  $\Delta^1 \times \partial_i \Delta^k$  are indexed by the subsets

$$\partial_i A := A \setminus \{i\} = A \cap ([k] \setminus \{i\}), \quad i \neq \max A.$$

Thus, to check the condition  $G_f|_{\Delta^1 \times \partial_i \Delta^k} = G_{d_i f}$ , it suffices to check—for every  $A = \{0, 1, \dots, \max A\} \subset [k]$  with  $\max A \neq i$ —that

$$(\partial_{A,i})^* G_{f,A} = G_{\partial_i f, \partial_i A}. \quad (10.10)$$

Here,  $\partial_{A,i}$  is the map fitting into the following commutative diagram:

$$\begin{array}{ccc} \Delta^{k+1} & \xrightarrow{A} & \Delta^1 \times \Delta^k \\ \uparrow \partial_{A,i} & & \uparrow \\ \Delta^k & \xrightarrow{\partial_i A} & \Delta^1 \times \partial_i \Delta^k \end{array}$$

Parsing the definitions, we see that the lefthand side of (10.10) is a map  $\mathbf{sd}(\Delta^k) \rightarrow \mathbf{Liou}_{\text{str}}^\diamond$  that assigns, to every subset  $\emptyset \neq I \subset [k]$ , the object

$$X_{\max(s_{\max \partial_i A}(I))} \times T^* \Delta^{I/\partial_i A}.$$

The righthand side of (10.10) assigns to each  $I$  the exact same object. A similar check of morphisms associated to  $I \subset J$  shows that the barycentric subdivision diagrams defined by both sides of (10.10) are identical; this finishes the check of condition (ii), so we conclude by applying Lemma 3.36.  $\square$

## 10.4 $\alpha$ as a right inverse

The purpose of this section is to prove the following:

**Lemma 10.19.** The map

$$\text{Ex}_{\simeq}(\beta) \circ \alpha : (\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond \rightarrow \text{Ex}_{\simeq}((\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond)$$

is homotopic to the map  $m : (\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond \rightarrow \text{Ex}_{\simeq}((\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond)$  from Proposition 3.19. In particular,  $\text{Ex}_{\simeq}(\beta) \circ \alpha$  is an equivalence.

The proof requires some set-up.

**Notation 10.20** ( $\eta$ ). For this section only, we will denote the natural transformation  $\text{Ex}_{\simeq} \hookrightarrow \text{Ex}$  by  $\eta$ . (This is a natural transformation between two functors from the category of simplicial sets to itself.)

**Notation 10.21** ( $M$  and  $M_+$ ). Because every simplex in  $(\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond$  encodes a diagram in  $\mathbf{Liou}_{\text{str}}^\diamond$  in the shape of a barycentric subdivision, we easily obtained the map (10.4) of semisimplicial sets. Post-composing with  $\text{Ex}(j)$  (Constructin 10.3), we obtain a map from  $(\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond$  to  $\text{Ex}((\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond)$ . This composition factors through  $\eta$ , and we notate the factoring morphism by  $M$  to obtain a commuting rectangle of semisimplicial sets as follows:

$$\begin{array}{ccc} (\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond & \xrightarrow{M} & \text{Ex}_{\simeq}((\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond) \\ \downarrow (10.4) & & \downarrow \eta \\ \text{Ex}(\mathbf{Liou}_{\text{str}}^\diamond) & \xrightarrow{\text{Ex}(j)} & \text{Ex}((\mathbf{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^\diamond) \end{array}$$

Let us justify the factorization through  $\text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond})$ . By construction, the map (10.4) hits only those simplices in  $\text{Ex}(\text{Liou}_{\text{str}}^{\diamond})$  whose max-localizing edges are sectorial equivalences (Condition (LD2)). And—applying  $j$ —such edges are homotopy-invertible in  $\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}}$  (combine Proposition 7.14(d) with Theorem 9.3).

Finally, while  $M$  and (10.4) do not respect degeneracy maps, by adjunction they induce maps of simplicial sets with domain  $(\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+}$ . Calling these maps  $M_{+}$  and  $(10.4)_{+}$  respectively, we obtain a commutative diagram of simplicial sets:

$$\begin{array}{ccc} (\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+} & \xrightarrow{M_{+}} & \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}) \\ \downarrow (10.4)_{+} & & \downarrow \eta \\ \text{Ex}(\text{Liou}_{\text{str}}^{\diamond}) & \xrightarrow{\text{Ex}(j)} & \text{Ex}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}) \end{array} \quad (10.11)$$

We now have the notation to state the main combinatorial lemma.

**Lemma 10.22.** The composition

$$(\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+} \xrightarrow{\epsilon} (\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond} \xrightarrow{m} \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond})$$

is homotopic to  $M_{+}$ . That is, there exists an edge in

$$\text{Map}^{\sharp}(((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+}, \mathcal{E}), \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}))$$

from  $m \circ \epsilon$  to  $M_{+}$ .

*Proof of Lemma 10.19, assuming Lemma 10.22.* Consider the following diagram of simplicial sets.

$$\begin{array}{ccc} (\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+} & \xrightarrow{M_{+}} & \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}) \\ \downarrow (10.4)_{+} & & \downarrow \eta \\ \text{Ex}(\text{Liou}_{\text{str}}^{\diamond}) & \xrightarrow{\text{Ex}(j)} & \text{Ex}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}) \\ \downarrow \text{Ex}(\iota) & \nearrow \text{Ex}(\beta) & \uparrow \eta \\ \text{Ex}(\text{Liou}_{\text{str}}^{\diamond}[(\mathfrak{e}\mathfrak{q}^{\diamond})^{-1}]) & & \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}) \\ \uparrow \eta & \xrightarrow{\text{Ex}_{\simeq}(\beta)} & \uparrow \\ \text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^{\diamond}[(\mathfrak{e}\mathfrak{q}^{\diamond})^{-1}]) & & \end{array}$$

(A curved arrow labeled  $(\alpha_{\text{semi}})_{+}$  points from  $(\text{Liou}_{\Delta}^{\mathcal{D}\text{ef}})^{\diamond}_{+}$  to  $\text{Ex}_{\simeq}(\text{Liou}_{\text{str}}^{\diamond}[(\mathfrak{e}\mathfrak{q}^{\diamond})^{-1}])$ .)

There are four cells in the diagram, all of which commute on the nose except for the triangle.

- The homotopy-coherent triangle in the middle is obtained by applying  $\text{Ex}$  to the diagram (10.3) – note that Proposition 3.22 shows us that  $\text{Ex}$  transports homotopy-coherent diagrams to homotopy-coherent diagrams. Thus there is a homotopy of maps of simplicial sets

$$\text{Ex}(\beta) \circ \text{Ex}(\iota) \circ (10.4)_{+} \sim \text{Ex}(j) \circ (10.4)_{+}.$$

- On the other hand,  $(\alpha_{\text{semi}})_{+}$  factors  $\text{Ex}(\iota) \circ (10.4)_{+}$  (as indicated by the curved diagram on the left) so the above homotopy restricts to a homotopy of maps of simplicial sets

$$\text{Ex}(\beta) \circ \eta \circ (\alpha_{\text{semi}})_{+} \sim \text{Ex}(j) \circ (10.4)_{+}.$$

- The top rectangle is (10.11).
- The bottom trapezoid is the naturality of  $\eta$  (Notation 10.20).

By using both the rectangle and the trapezoid, the above homotopy becomes

$$\eta \circ \text{Ex}_{\simeq}(\beta) \circ (\alpha_{\text{semi}})_+ \sim \eta \circ M_+. \quad (10.12)$$

Now, the homotopy encoded in the triangle (10.3) is a homotopy of functors between  $\infty$ -categories; in particular, the induced homotopy (10.12) is a map  $(\text{Liou}_{\Delta}^{\text{Def}})_+^{\diamond} \times \Delta^1 \rightarrow \text{Ex}((\text{Liou}_{\Delta}^{\text{Def}})^{\diamond})$  whose image lies entirely in  $\text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\text{Def}})^{\diamond})$ . So (10.12) restricts to a homotopy

$$\text{Ex}_{\simeq}(\beta) \circ (\alpha_{\text{semi}})_+ \sim M_+$$

respecting the marked edges of  $((\text{Liou}_{\Delta}^{\text{Def}})_+^{\diamond}, \mathcal{E})$ . By Lemma 10.22,  $M_+$  is homotopic to  $m \circ \epsilon$ . Combining with the homotopy  $(\alpha_{\text{semi}})_+ \sim \alpha \circ \epsilon$  (Construction 10.11), we conclude

$$\text{Ex}_{\simeq}(\beta) \circ \alpha \circ \epsilon \sim m \circ \epsilon.$$

Because precomposition with  $\epsilon$  induces a homotopy equivalence of mapping spaces (Remark 3.32), we conclude that  $\text{Ex}_{\simeq}(\beta) \circ \alpha$  is homotopic to  $m$ , as desired.  $\square$

## 10.5 Proof of Lemma 10.22

One may establish Lemma 10.22 via the strategy used to prove Lemma 10.13. Because the techniques are similar, we will simply exposit an understanding of the maps  $M$  and  $m$  and outline the construction; what we do not spell out in detail is the construction of the maps  $\mathbb{G}_{f,k,\vec{A}}^{\vee}$  in (10.13).

**Example 10.23** (A description of  $M$ ). Fix an  $l$ -simplex  $f : \Delta^l \rightarrow (\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$ . Such a simplex is equivalent to the data of a diagram in the shape of  $\text{sd}(\Delta^l)$  which assigns

- To any non-empty subset  $A \subset [l]$ , a Liouville sector  $X_A \times T^*\Delta^A$ , and
- To any inclusion  $A \subset B$ , a morphism

$$f_{A \subset B} \otimes T^*\eta_{A \subset B} : X_A \times T^*\Delta^A \otimes T^*[-1, 0]^{B \setminus A} \rightarrow X_B \times T^*\Delta^B.$$

Then, unwinding the definitions,  $M(f) : \Delta^l \rightarrow \text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\text{Def}})^{\diamond})$  encodes a diagram

$$\text{sd}(\text{sd}(\Delta^l)) \rightarrow \text{Liou}_{\text{str}}^{\diamond}$$

which assigns the following. (See Example 3.39 for the notation.)

- To every pair  $(k, A_0 \subset \dots \subset A_k)$  where  $k \geq 0$  and  $A_k \subset [l]$ , the sector

$$(X_{A_k} \times T^*\Delta^{A_k}) \otimes T^*\Delta^k$$

where the sectorial structure in the parentheses is that specified by  $f$ . Indeed, the object above can be identified as what  $j$  assigns to the barycenter of the simplex in  $\text{sd}(\Delta^l)$  determined by the chain  $A_0 \subset \dots \subset A_k$ .



- To every inclusion  $\delta : [k] \hookrightarrow [k']$  exhibiting  $A_0 \subset \dots \subset A_k$  as a subchain of  $A'_0 \subset \dots \subset A'_{k'}$ , the morphism

$$f_{A_k \subset A'_{k'}} \otimes T^* \eta_{A_k \subset A'_{k'}} \otimes T^*(\delta \times \eta_\delta).$$

Concretely, this is the sectorial inclusion

$$X_{A_k} \times T^* \Delta^{A_k} \otimes T^*[-1, 0]^{A'_{k'} \setminus A_k} \otimes T^*(\Delta^k \times [-1, 0]^{[k'] \setminus \delta([k])}) \rightarrow (X_{A'_{k'}} \times T^* \Delta^{A'_{k'}}) \otimes T^* \Delta^{k'}.$$

Note that  $M$  requires no knowledge of the degeneracy maps of  $(\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$ .

We now move onto understanding  $m$ . Let us first set some notation that will help us describe  $m : \mathcal{C} \rightarrow \mathrm{Ex}(\mathcal{C})$  for a general simplicial set  $m$ .

**Notation 10.24** ( $\sigma_{\vec{A}}$  and  $\delta_{\vec{A}}$ ). Fix an  $l$ -simplex  $f : \Delta^l \rightarrow \mathcal{C}$ . The data of  $m(f) : \Delta^l \rightarrow \mathrm{Ex}(\mathcal{C})$  is equivalent to a map

$$\mathrm{sd}(\Delta^l) \rightarrow \mathcal{C}.$$

We will understand this map by understanding what it does on every  $k$ -simplex  $\Delta^k \rightarrow \mathrm{sd}(\Delta^l)$ .

First, note that a  $k$ -simplex in  $\mathrm{sd}(\Delta^l)$  is the data of a chain

$$\vec{A} = A_0 \subset \dots \subset A_k \subset [l].$$

Define the map

$$\max \vec{A} : [k] \rightarrow [l], \quad i \mapsto \max A_i.$$

This is the composition

$$[k] \xrightarrow{\vec{A}} \mathcal{P}'([l]) \xrightarrow{\max} [l].$$

Then  $\max \vec{A}$  factors uniquely as a surjection  $\sigma$  followed by an injection  $\delta$ :

$$[k] \xrightarrow{\sigma_{\vec{A}}} [l'] \cong \mathrm{image}(\max \vec{A}) \xrightarrow{\delta_{\vec{A}}} [l].$$

(The subscript  $\vec{A}$  is to remind us of the dependence on  $\vec{A}$ .) Then the composite

$$\Delta^k \rightarrow \mathrm{sd}(\Delta^l) \rightarrow \mathcal{C}$$

is the simplex

$$(\sigma_{\vec{A}})^*(\delta_{\vec{A}})^* f$$

(where the pullback along  $\sigma$  and  $\delta$  are defined using the simplicial set structure of  $\mathcal{C}$ ).

**Example 10.25** (A description of  $m$ ). Now we explicate  $m : (\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond \rightarrow \mathrm{Ex}_\simeq((\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond)$ . Fix an  $l$ -simplex  $f : \Delta^l \rightarrow (\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$ .

As indicated in Notation 10.24, it is hard to give an explicit definition of  $m(f)$  when  $\max \vec{A}$  is not an injection, for the degeneracy maps of  $(\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$  are, in general, not concretely given (and instead are known to exist by abstract results). We only know the degeneracy maps of a constant barycentric diagram (Definition 7.3)—these degenerate simplices are also constant by Theorem 7.11(b). So let us do the best we can and describe those  $k$ -simplices in  $\mathrm{sd}(\Delta^l)$  defining  $m(f)$  that do not involve the abstractly defined degeneracies:

- We note that  $\delta_{\vec{A}}^* f$  is a constant barycentric diagram when the restriction of  $f : \Delta^l \rightarrow (\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$  to the image of  $\max \vec{A}$  is constant. In this case, the simplex  $(\sigma_{\vec{A}})^*(\delta_{\vec{A}})^* f$  is also constant. In particular, the map  $\Delta^k \rightarrow \mathrm{sd}(\Delta^l) \rightarrow (\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$  is a diagram that is encoded by a single sector  $X$ . The diagram  $\Delta^k \rightarrow (\mathrm{Liou}_\Delta^{\mathrm{Def}})^\diamond$  sends any  $\emptyset \neq A \subset [k]$  to the sector  $X \otimes T^* \Delta^A$  and all  $A \subset B$  to morphisms of the form  $\mathrm{id} \otimes T^* \eta_{A \subset B}$ .

- The other case we can describe is when  $\delta_{\vec{A}}$  is a bijection. This is the case when  $\max \vec{A} : [k] \rightarrow [l]$  is an injection—i.e., when the chain  $A_0 \subset \dots \subset A_k$  satisfies the property that  $\max A_i$  is strictly less than  $\max A_{i+1}$ . In this case, the chain  $\max A_0 < \max A_1 < \dots < \max A_k$  defines a subsimplex of  $\Delta^l$ , and  $(\sigma_{\vec{A}})^*(\delta_{\vec{A}})^*f$  is simply the restriction of  $f$  to this subsimplex.

**Example 10.26** ( $M$  and  $m$  on a 1-simplex). Fix a 1-simplex  $f : \Delta^1 \rightarrow (\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$ . This is the data of a diagram of (stabilized) Liouville sectors as follows:

$$X_0 \xrightarrow{f_{0 \subset 01} \times T^* \eta} X_{01} \times T^* \Delta^1 \xleftarrow{f_{1 \subset 01} \times T^* \eta} X_1.$$

Then  $M(f)$  is a 1-simplex in  $\text{Ex}_{\simeq}((\text{Liou}_{\Delta}^{\text{Def}})^{\diamond})$  that we obtain by applying the map  $j$  (Construction 10.3) to each edge in the above diagram. The result is a diagram of the shape  $\text{sd}(\text{sd}(\Delta^1))$  in  $\text{Liou}_{\text{str}}^{\diamond}$ , which we draw as follows:


$$X_0 \xrightarrow{(f_{0 \subset 01} \times T^* \eta) \otimes T^* \eta} (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 \xleftarrow{\text{id} \otimes T^* \eta} X_{01} \times T^* \Delta^1 \xleftarrow{\text{id} \otimes T^* \eta} (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 \xleftarrow{(f_{1 \subset 01} \times T^* \eta) \otimes T^* \eta} X_1.$$

The dashes are to conform to the notation in Figure 10.28.

On the other hand,  $m(f)$  can be drawn as follows:

$$X_0 \xrightarrow{f_{0 \subset 01} \times T^* \eta} X_{01} \times T^* \Delta^1 \xleftarrow{f_{1 \subset 01} \times T^* \eta} X_1 \xleftarrow{\text{id} \otimes T^* \eta} X_1 \times T^* \Delta^1 \xleftarrow{\text{id} \otimes T^* \eta} X_1.$$

**Example 10.27** ( $M$  and  $m$  on a 2-simplex). Fix a 2-simplex  $f : \Delta^2 \rightarrow (\text{Liou}_{\Delta}^{\text{Def}})^{\diamond}$ . Both  $M(f)$  and  $m(f)$  define a diagram  $\text{sd}(\text{sd}(\Delta^2)) \rightarrow \text{Liou}_{\text{str}}^{\diamond}$ , which we depict in Figure 10.28.



**Figure 10.28.** On top is the diagram  $\text{sd}(\text{sd}(\Delta^2)) \rightarrow \text{Liou}_{\text{str}}^\diamond$  given by  $M$  of a 2-simplex. The dashed arrows are all of the form  $\text{id} \otimes T^*\eta$ . Not indicated are the morphisms to the objects  $X_{012} \times T^*\Delta^2 \otimes T^*\Delta^2$ , to reduce clutter.

On the bottom is the diagram given by  $m$  of a 2-simplex. The dashed arrows are all of the form  $\text{id} \otimes T^*\eta$ . Simply labeled as “ $s$ ” are centers of barycentric subdivision diagrams that are degenerate simplices in  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$ . (The only degenerate simplices that are explicitly indicated are the constant simplices.)

Despite the kinks in the edges, we hope the reader can detect the outline of the  $\text{sd}(\Delta^2)$  of which we are taking the subdivision.

**Construction 10.29** ( $\mathbb{G}_f$ ). We have fixed an  $l$ -simplex  $f : \Delta^l \rightarrow (\text{Liou}_\Delta^{\text{Def}})^\diamond$ . Fix a  $k$ -simplex  $\vec{A} : \Delta^k \rightarrow \text{sd}(\Delta^l)$ . The definition of  $m(f)$  on this  $k$ -simplex involves using degeneracy maps in  $(\text{Liou}_\Delta^{\text{Def}})^\diamond$ . Because these degeneracy maps are maps whose existence was concluded formally, they are rather inexplicit. As a result, we will construct  $\mathbb{G}_f$  inductively, by the cardinality of the fibers of the map

$$\Delta^k \xrightarrow{\vec{A}} \text{sd}(\Delta^l) \xrightarrow{\max} \Delta^l.$$

Fix an  $l$ -simplex  $f : \Delta^l \rightarrow (\text{Liou}_\Delta^{\text{Def}})^\diamond$ . We may understand the simplices

$$M(f), m(f) : \Delta^l \rightarrow \text{Ex}_{\sim}((\text{Liou}_\Delta^{\text{Def}})^\diamond)$$

by their adjoint maps  $\text{sd}(\Delta^l) \rightarrow (\text{Liou}_\Delta^{\text{Def}})^\diamond$ . To understand these maps, let us note

$$\text{sd}(\Delta^l) \cong \bigcup_{k, \vec{A}} \Delta^k.$$

(This is a formula expressing that a  $k$ -simplex in a barycentric subdivision is the same thing as an increasing length- $k$  chain  $\vec{A}$  of subsimplices.) For each length  $k$  chain  $\vec{A} = A_0 \subset \dots \subset A_k$ , suppose one can construct maps

$$\Delta^1 \times \Delta^k \xrightarrow{\mathbb{G}_{f,k,\vec{A}}^\vee} \text{Liou}_\Delta \quad (10.13)$$

that glue together to define a map

$$\mathbb{G}_f^\vee : \Delta^1 \times \text{sd}(\Delta^l) \cong \Delta^1 \times \left( \bigcup_{(k,\vec{A})} \Delta^k \right) \rightarrow \text{Liou}_\Delta. \quad (10.14)$$

As indicated, we let  $\mathbb{G}_f^\vee$  denote the composite map (of the isomorphism above, together with the gluings of  $\mathbb{G}_{f,k,\vec{A}}^\vee$ ). We then have

$$\begin{array}{ccc} \Delta^1 \times \text{sd}(\Delta^l) & \xrightarrow{\mathbb{G}_f^\vee} & \text{Liou}_\Delta \\ \uparrow \text{max}_{\Delta^1} \times \text{id}_{\text{sd}(\Delta^l)} & & \\ \text{sd}(\Delta^1) \times \text{sd}(\Delta^l) & & \\ \uparrow \text{collapse} & & \\ \text{sd}(\Delta^1 \times \Delta^l) & & \end{array} \quad (10.15)$$

By adjunction, we thus obtain a map  $\Delta^l \times \Delta^1 \rightarrow \text{Ex}_{\sim}((\text{Liou}_\Delta^{\text{Def}})^\diamond)$ . A routine check on edges demonstrates that this map factors through  $\text{Ex}_{\sim}((\text{Liou}_\Delta^{\text{Def}})^\diamond)$ , so we obtain the desired map

$$\mathbb{G}_f : \Delta^1 \times \Delta^l \rightarrow \text{Ex}_{\sim}((\text{Liou}_\Delta^{\text{Def}})^\diamond).$$

**Remark 10.30.** Given any two simplicial sets  $X$  and  $Y$ , there is a collapse map given by the dashed arrow below, defined as the composition of the solid arrows.


$$\begin{array}{ccc} \text{sd}(X) \times \text{sd}(Y) & \xleftarrow{\cong} & \left( \text{colim}_{\Delta^{k'} \rightarrow X} \text{sd}(\Delta^{k'}) \right) \times \text{sd}(Y) \xleftarrow{\cong} \text{colim}_{\Delta^{k'} \rightarrow X} \text{sd}(\Delta^{k'}) \times \text{sd}(Y) \xleftarrow{\cong} \text{colim}_{\Delta^{k'} \rightarrow X} \text{sd}(\Delta^{k'}) \times \left( \text{colim}_{\Delta^{k''} \rightarrow Y} \text{sd}(\Delta^{k''}) \right) \xleftarrow{\cong} \text{colim}_{\Delta^{k'} \rightarrow X} \text{colim}_{\Delta^{k''} \rightarrow Y} \text{sd}(\Delta^{k'}) \times \text{sd}(\Delta^{k''}) \\ \uparrow \text{dashed} & & \uparrow \text{colim}_{\Delta^k \rightarrow X \times Y} \text{sd}(\Delta^k) \times \text{sd}(\Delta^k) \\ \text{sd}(X \times Y) & \xleftarrow{\cong} & \text{colim}_{\Delta^k \rightarrow X \times Y} \text{sd}(\Delta^k) \end{array}$$

Every isomorphism in the above diagram is by definition of  $\mathbf{sd}$  of a simplicial set, and by the fact that direct products of simplicial sets commutes with colimits (because, for example, the category of simplicial sets is Cartesian closed).

The two vertical arrows in the right are diagonal maps – the lower-right upward map is induced by the diagonal map of simplicial sets  $W \rightarrow W \times W$  in the case  $W = \mathbf{sd}(\Delta^k)$ , while the upper-right upward map is the inclusion of the diagram of simplices  $\{\Delta^k \rightarrow X \times Y\}$  into the diagram of simplices  $\{\Delta^{k'} \rightarrow X, \Delta^{k''} \rightarrow Y\}$  (by including into the subdiagram where  $k' = k''$ ).

The collapse map in (10.15) is the special case of  $X = \Delta^1, Y = \Delta^l$ .

**Example 10.31.** Let  $l = 1$ . Then the diagram  $\Delta^1 \times \mathbf{sd}(\Delta^l)$  may be drawn as follows:



Fix a simplex  $f : \Delta^l \rightarrow (\mathbf{Liou}_\Delta^{\mathcal{D}\text{ef}})^\diamond$ . The map  $\mathbb{G}_f^\vee : \Delta^1 \times \mathbf{sd}(\Delta^l) \rightarrow (\mathbf{Liou}_\Delta^{\mathcal{D}\text{ef}})^\diamond$  from (10.14) represents a diagram  $\text{sub}(\Delta^1 \times \mathbf{sd}(\Delta^l)) \rightarrow \mathbf{Liou}_{\text{str}}^\diamond$ ; we can draw the parts of this diagram corresponding to the arrows in (10.16) (and not, say, the 2-simplices) as follows:

$$\begin{array}{ccccccc}
 X_0 & \xrightarrow{(f_{0 \subset 01} \times T^* \eta) \otimes T^* \eta} & (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 & \xleftarrow{\text{id} \otimes T^* \eta} & X_{01} \times T^* \Delta^1 & \xrightarrow{\text{id} \otimes T^* \eta} & (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 & \xleftarrow{(f_{1 \subset 01} \times T^* \eta) \otimes T^* \eta} & X_1 \\
 \downarrow & & \swarrow & & \downarrow & & \searrow & & \downarrow \\
 X_0 \otimes T^* \Delta^1 & & (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 & & X_{01} \times T^* \Delta^1 \otimes T^* \Delta^1 & & (X_{01} \times T^* \Delta^1) \otimes T^* \Delta^1 & & X_1 \otimes T^* \Delta^1 \\
 \uparrow & \nearrow & & & \uparrow & & \nwarrow & \uparrow & \\
 X_0 & \xrightarrow{f_{0 \subset 01} \times T^* \eta} & X_{01} \times T^* \Delta^1 & \xleftarrow{f_{1 \subset 01} \times T^* \eta} & X_1 & \xrightarrow{\text{id} \otimes T^* \eta} & X_1 \times T^* \Delta^1 & \xleftarrow{\text{id} \otimes T^* \eta} & X_1
 \end{array}$$

## 10.6 Completing the proof of Theorem C

We have the following corollary of Lemmas 10.14 and 10.19.

**Corollary 10.32.** The maps  $\alpha$  and  $\beta$  are equivalences of  $\infty$ -categories.

*Proof.* It is clear that  $\alpha$  is essentially surjective. Lemma 10.19 shows that  $\alpha$  induces an injection on  $\pi_* \text{hom}$ . Since Lemma 10.14 says  $\alpha \circ \beta$  is an equivalence, we see  $\alpha$  induces a surjection on  $\pi_* \text{hom}$ . Thus  $\alpha$  is an equivalence of  $\infty$ -categories. Since  $\alpha$  and  $\alpha \circ \beta$  are both equivalences, we conclude that  $\beta$  is an equivalence as well.  $\square$

*Proof of Theorem C.*  $\beta$  from Notation 10.7 is the map induced by the universal property of localization. Now invoke Corollary 10.32.  $\square$

## 11 The symmetric monoidal structure

The goal of this section is to construct the symmetric monoidal structure on  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}]$ .

### 11.1 Stabilizing commutes with localization

Let  $\mathcal{C}$  be an arbitrary  $\infty$ -category, and fix a collection of morphisms  $S \subset \mathcal{C}$ . We denote by  $\mathcal{C}[S^{-1}]$  the localization.

**Notation 11.1.** We let  $\diamond : \mathcal{C} \rightarrow \mathcal{C}$  be an endofunctor and define

$$\mathcal{C}^{\diamond} := \mathrm{colim} \left( \mathcal{C} \xrightarrow{\diamond} \mathcal{C} \xrightarrow{\diamond} \dots \right)$$

Assuming that  $\diamond(S) \subset S$ , we define

$$\mathcal{C}[S^{-1}]^{\diamond} := \mathrm{colim} \left( \mathcal{C}[S^{-1}] \xrightarrow{\diamond} \mathcal{C}[S^{-1}] \xrightarrow{\diamond} \dots \right).$$

Then:

**Lemma 11.2.** The natural functor  $\mathcal{C}^{\diamond} \rightarrow \mathcal{C}[S^{-1}]^{\diamond}$  is a localization of  $\mathcal{C}^{\diamond}$ . In fact, letting  $a_j : \mathcal{C} \rightarrow \mathcal{C}^{\diamond}$  denote the natural morphism from the  $j$ th copy of  $\mathcal{C}$  appearing in the colimit, and letting  $S^{\diamond} \subset \mathcal{C}^{\diamond}$  denote the collection of morphisms

$$S^{\diamond} = \bigcup_j a_j(S)$$

the map  $\mathcal{C}^{\diamond} \rightarrow \mathcal{C}[S^{-1}]^{\diamond}$  realizes an equivalence

$$\mathcal{C}^{\diamond}[(S^{\diamond})^{-1}] \xrightarrow{\sim} \mathcal{C}[S^{-1}]^{\diamond}$$

*Proof.* We  $\mathcal{WCat}_{\infty}$  be the  $\infty$ -category of  $\infty$ -categories marked by systems. Concretely, an object is a pair  $(\mathcal{C}, S)$  of an  $\infty$ -category  $\mathcal{C}$  together and a collection of edges  $S$  in  $\mathcal{C}$  containing all equivalences of  $\mathcal{C}$ , and closed under homotopy and composition. (For more details, see Construction 4.1.3.1 of [18].) In what follows, we also let  $\mathcal{C}^{\sim}$  denote the largest  $\infty$ -groupoid contained in  $\mathcal{C}$ —it is the Kan complex generated by all objects of  $\mathcal{C}$  and by all equivalences of  $\mathcal{C}$ . The functor

$$\mathcal{WCat}_{\infty} \rightarrow \mathcal{WCat}_{\infty}, \quad (\mathcal{C}, S) \mapsto (\mathcal{C}[S^{-1}], \mathcal{C}[S^{-1}]^{\sim})$$

—which sends the pair  $(\mathcal{C}, S)$  to the localization  $\mathcal{C}[S^{-1}]$ , marked by all equivalences of the localization—may be modeled as a fibrant replacement functor in the (left proper, combinatorial) model category of marked simplicial sets. The image of this localization functor is equivalent to the  $\infty$ -category of  $\infty$ -categories; the functor is in fact an idempotent left adjoint to the inclusion of the  $\infty$ -category of  $\infty$ -categories. (See Proposition 4.1.3.2 of [18].) So it preserves colimits. We conclude by noting that

$$(\mathcal{C}^{\diamond}, S^{\diamond}) = \mathrm{colim} \left( (\mathcal{C}, S) \xrightarrow{\diamond} (\mathcal{C}, S) \xrightarrow{\diamond} \dots \right)$$

as marked simplicial sets. □

**Corollary 11.3.** We have an equivalence  $\mathrm{Liou}_{\mathrm{str}}^{\diamond}[(\mathfrak{eq}^{\diamond})^{-1}] \xrightarrow{\sim} \mathrm{Liou}_{\mathrm{str}}[\mathfrak{eq}^{-1}]^{\diamond}$ .

## 11.2 Symmetric monoidal localization

Let  $\mathcal{C}^\otimes$  be a symmetric monoidal  $\infty$ -category with underlying  $\infty$ -category  $\mathcal{C}$ . Fix also a collection of morphisms  $S \subset \mathcal{C}$ .

**Definition 11.4.** We say that  $S$  is *compatible* with a symmetric monoidal structure  $\otimes$  if for all objects  $M \in \mathcal{C}$  and all morphisms  $f \in S$ , we have that  $\mathrm{id}_M \otimes f$  and  $f \otimes \mathrm{id}_M$  are both in  $S$ .

**Proposition 11.5.** Suppose  $S$  is compatible with  $\otimes$ . Then there is an induced symmetric monoidal structure on  $\mathcal{C}[S^{-1}]$  so that the functor  $\mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$  may be promoted to a symmetric monoidal functor.

In fact, more is true: for any symmetric monoidal  $\infty$ -category  $\mathcal{D}^\otimes$ , the  $\infty$ -category of symmetric monoidal functors  $\mathrm{Fun}^\otimes(\mathcal{C}[S^{-1}]^\otimes, \mathcal{D}^\otimes)$  is identified with the full subcategory of  $\mathrm{Fun}^\otimes(\mathcal{C}^\otimes, \mathcal{D}^\otimes)$  sending morphisms in  $S$  to equivalences in  $\mathcal{D}$ . This identification is given by composing with the symmetric monoidal functor  $\mathcal{C}^\otimes \rightarrow \mathcal{C}[S^{-1}]^\otimes$

*Proof.* This is a consequence of Proposition 4.1.3.4 of [18].  $\square$

## 11.3 Symmetric monoidal structures on stabilizations

Let  $\mathcal{C}^\otimes$  be a symmetric monoidal  $\infty$ -category with underlying  $\infty$ -category  $\mathcal{C}$ , and an object  $T$ . One may contemplate whether the colimit

$$\mathrm{colim} \left( \mathcal{C} \xrightarrow{T \otimes -} \mathcal{C} \xrightarrow{T \otimes -} \dots \right) \quad (11.1)$$

—which may<sup>28</sup> rightfully be called  $\mathcal{C}[T^{-1}]$ — admits a symmetric monoidal structure for which  $T$  becomes an invertible object. (That is, for which there exists some object  $U$  such that  $T \otimes U$  is equivalent to the tensor unit.)

Inspired by observations of Voevodsky [36, Section 4], and following ideas of Toën-Vezzosi [35], Robalo gives the following criterion.

**Proposition 11.6** (Corollary 4.24 of [28]). Let  $\mathcal{C}^\otimes$  be a symmetric monoidal  $\infty$ -category with underlying  $\infty$ -category  $\mathcal{C}$ . Let  $T \in \mathcal{C}$  and let

$$\tau : T \otimes T \otimes T \rightarrow T \otimes T \otimes T$$

denote the cyclic permutation map (123) determined by the symmetric monoidal structure. If  $\tau$  is homotopic to the identity of  $T \otimes T \otimes T$ , then

1. The colimit (11.1) inherits a symmetric monoidal structure from  $\mathcal{C}$  for which the functor

$$j : \mathcal{C} \rightarrow \mathrm{colim} \left( \mathcal{C} \xrightarrow{T \otimes -} \mathcal{C} \xrightarrow{T \otimes -} \dots \right)$$

can be made symmetric monoidal.

2. Moreover,  $j$  is universal among symmetric monoidal functors that send  $T$  to an invertible object.

---

<sup>28</sup>We will not use the notation  $\mathcal{C}[T^{-1}]$  in the rest of this work, to avoid confusion with notation such as  $\mathfrak{eq}^{-1}$ . The philosophy of our work centers around compositionally inverting morphisms, not multiplicatively inverting objects.

*Proof.* For the reader's convenience, let us recall that any symmetric monoidal  $\infty$ -category is a commutative (that is,  $E_\infty$ ) algebra in the  $\infty$ -category of  $\infty$ -categories (where the monoidal structure is direct product of  $\infty$ -categories). It thus makes sense to speak of modules over any symmetric monoidal  $\infty$ -category. The original reference is Section 4.5 of [18]; we also recommend Section 3 of [28].

We follow the notation of [28]. In Proposition 4.1 of *ibid.*, the universal symmetric monoidal  $\infty$ -category obtained from  $\mathcal{C}$  by inverting  $T$  is constructed—it is denoted by  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes(\mathcal{C}^\otimes)$ . In fact,  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes$  is more generally a left adjoint taking any commutative  $\mathcal{C}^\otimes$ -algebra  $A$  to another  $\mathcal{C}^\otimes$ -algebra  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes(A)$  for which the map  $\mathcal{C} \rightarrow A$  sends  $T$  to an invertible object.

The same construction works for  $\mathcal{C}^\otimes$ -modules: Pulling back along the canonical map  $\mathcal{C}^\otimes \rightarrow \mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes(\mathcal{C}^\otimes)$ , one witnesses a fully faithful embedding from the  $\infty$ -category of  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes(\mathcal{C}^\otimes)$ -modules to that of  $\mathcal{C}^\otimes$ -modules. The left adjoint produces, out of any  $\mathcal{C}^\otimes$ -module  $M$ , the universal  $\mathcal{C}^\otimes$ -module

$$M \rightarrow \mathcal{L}_{(\mathcal{C}^\otimes, T)}(M)$$

rendering  $T \otimes -$  an autoequivalence (Proposition 4.2 of *ibid.*). We note that—by the usual passage between algebras and modules—in the special case that  $M = \mathcal{C}$  (where the module action is the canonical one of  $\mathcal{C}^\otimes$  acting on itself), there is a natural equivalence

$$\mathcal{L}_{(\mathcal{C}^\otimes, T)}^\otimes(\mathcal{C}^\otimes) \simeq \mathcal{L}_{(\mathcal{C}^\otimes, T)}(\mathcal{C}). \quad (11.2)$$

We shall now finally use the assumption on  $\tau$  being homotopic to  $\text{id}_{T^\otimes 3}$ . Given a module  $M$  over  $\mathcal{C}^\otimes$ , the colimit

$$\text{Stab}_T(M) := \text{colim} \left( M \xrightarrow{T \otimes -} M \xrightarrow{T \otimes -} \dots \right)$$

is another  $\mathcal{C}^\otimes$ -module with the property that the endofunctor  $T \otimes -$  is an equivalence (Proposition 4.21 of *ibid.*; this requires  $\tau \sim \text{id}_{T^\otimes 3}$ ). Thus, we obtain a canonical map

$$\mathcal{L}_{(\mathcal{C}^\otimes, T)}(M) \rightarrow \text{Stab}_T(M). \quad (11.3)$$

We claim this map is an equivalence; this is more or less the exact same proof as in Corollary 4.24 of *ibid.* and we reproduce it here for the reader's convenience.

By adjunction, we have a factorization  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}(M) \rightarrow \mathcal{L}_{(\mathcal{C}^\otimes, T)}(\text{Stab}_T M) \rightarrow \text{Stab}_T(M)$ . Because  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}$  is a left adjoint, it commutes with colimits, hence  $\text{Stab}_T \mathcal{L}_{(\mathcal{C}^\otimes, T)} \simeq \mathcal{L}_{(\mathcal{C}^\otimes, T)} \text{Stab}_T$ . This results in a factorization

$$\mathcal{L}_{(\mathcal{C}^\otimes, T)}(M) \rightarrow \text{Stab}_T \mathcal{L}_{(\mathcal{C}^\otimes, T)}(M) \xrightarrow{\sim} \mathcal{L}_{(\mathcal{C}^\otimes, T)}(\text{Stab}_T(M)) \rightarrow \text{Stab}_T(M).$$

The last arrow is an equivalence because  $\text{Stab}_T M$  is already a module for whom  $T \otimes -$  is an equivalence. The first arrow is an equivalence because  $T \otimes -$  is an autoequivalence of  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}(M)$ , meaning the sequence colimit  $\text{Stab}_T$  recovers  $\mathcal{L}_{(\mathcal{C}^\otimes, T)}(M)$  itself.

Combining the equivalences (11.2) and (11.3), and by invoking the known universal property for the lefthand side of (11.2), we are finished.  $\square$

## 11.4 The symmetric monoidal structure

We will prove the following lemma shortly; it is at the interface of geometry and localizations:



**Lemma 11.7.** Consider the object  $T^*[0, 1] \times T^*[0, 1] \times T^*[0, 1]$  in  $\mathbf{Liou}_{\text{str}}$  (and hence in  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$ ). Denote by  $T^*\tau$  the natural automorphism given by the cyclic order 3 permutation (123) of coordinates:

$$\tau : ((q_1, p_1), (q_2, p_2), (q_3, p_3)) \mapsto ((q_3, p_3), (q_1, p_1), (q_2, p_2)).$$

(This map is part of the symmetric monoidal structure in  $\mathbf{Liou}_{\text{str}}$ , hence in  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$ .) Then in  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$ ,  $T^*\tau$  is homotopic to the identity.

**Remark 11.8.** Lemma 11.7 is the only non-formal ingredient of exhibiting a symmetric monoidal structure on  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$ . The main issue is that there is no straightforward geometric description of the morphism spaces in  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$  (in contrast to  $\mathbf{Liou}_{\text{str}}^\diamond[(\mathfrak{e}\mathfrak{q}^\diamond)^{-1}]$ ). All we have is the definition as a localization. Thus, we must somehow argue that the categorically formal process of localization recovers the geometric information of (123) being isotopic to the identity.

## 11.5 Proof of Theorem D

Assume Lemma 11.7 for the moment.

*Proof of Theorem D.* Clearly,  $\mathbf{Liou}_{\text{str}}$  has a symmetric monoidal structure given by direct products—of objects, and of morphisms. (Here we are using that all morphisms in  $\mathbf{Liou}_{\text{str}}$  are *strict* sectorial embeddings.)

By Proposition 11.5, we thus obtain a symmetric monoidal structure on the  $\infty$ -category

$$\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$$

and the symmetric monoidal functor

$$\mathbf{Liou}_{\text{str}} \rightarrow \mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$$

has the universal property that any symmetric monoidal functor  $\mathbf{Liou}_{\text{str}} \rightarrow \mathcal{D}$  sending Liouville equivalences to equivalences in  $\mathcal{D}$  will admit a symmetric monoidal, homotopy-coherent factorization through  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$ .

Assuming Lemma 11.7, we see that  $T = T^*[0, 1]$  satisfies the hypotheses of Proposition 11.6. Invoking Proposition 11.6, we conclude that  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]^\diamond = \text{Stab}_{T^*[0, 1]} \mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$  thus<sup>29</sup> inherits a symmetric monoidal structure, and the symmetric monoidal functor

$$\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}] \rightarrow \mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]^\diamond$$

is universal for symmetric monoidal functors out of  $\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]$  that send  $T^*[0, 1]$  to an invertible object.

Finally, we have an equivalence of  $\infty$ -categories as follows:

$$\mathbf{Liou}_{\text{str}}[\mathfrak{e}\mathfrak{q}^{-1}]^\diamond \xrightarrow[\sim]{\text{Cor 11.3}} \mathbf{Liou}_{\text{str}}^\diamond[(\mathfrak{e}\mathfrak{q}^\diamond)^{-1}]$$

The result follows by endowing  $\mathcal{L}\mathbf{iou}^\diamond \simeq \mathbf{Liou}_{\text{str}}^\diamond[(\mathfrak{e}\mathfrak{q}^\diamond)^{-1}]$  with the induced symmetric monoidal structure.  $\square$

---

<sup>29</sup>Here, we are utilizing notation from Section 11.1 and from the Proof of Proposition 11.6.

## 11.6 Proof of Lemma 11.7

In this proof, we will replace  $T^*[0, 1]$  with  $T^*[-1, 1]$  for convenience. (We would like the origin to be in the interior of the interval.)

Let  $\mathbf{Mfld}$  denote the category (in the classical sense) of smooth manifolds, possibly with corners, and whose morphisms are smooth codimension zero embeddings; it is symmetric monoidal under direct product. We let  $\mathbf{Mfld}^{\text{cmpct}} \subset \mathbf{Mfld}$  denote the full subcategory of *compact* smooth manifolds, possibly with corners. There is a symmetric monoidal functor

$$\mathbf{Mfld}^{\text{cmpct}} \rightarrow \mathbf{Liou}_{\text{str}}, \quad Q \mapsto T^*Q. \quad (11.4)$$

Because this functor is symmetric monoidal, it sends the permutation map  $\tau$  of  $[-1, 1] \times [-1, 1] \times [-1, 1]$  to  $T^*\tau$ .

**Definition 11.9.** Let  $X$  and  $Y$  be objects of  $\mathbf{Mfld}^{\text{cmpct}}$ . We say that a smooth, codimension zero embedding  $\bar{f} : X \rightarrow Y$  is an *isotopy equivalence* if there exist a smooth, codimension zero embedding  $\bar{g} : Y \rightarrow X$  and isotopies  $\bar{f}\bar{g} \sim \text{id}_Y, \bar{g}\bar{f} \sim \text{id}_X$ .

**Example 11.10.** If  $X = Y = [-1, 1]^n$  and if  $\bar{f}$  is orientation-preserving, then  $\bar{f}$  is an isotopy equivalence. This is a result of the following stronger fact:

**Proposition 11.11.** Fix  $n \geq 0$  and a real number  $0 < \epsilon \leq 1$ . The following spaces are all homotopy equivalent:

1. The space of smooth, codimension-zero embeddings  $\bar{f} : [-1, 1]^n \rightarrow [-1, 1]^n$ .
2. The space of those  $\bar{f}$  such that the image of  $\bar{f}$  is contained in  $[-\epsilon, \epsilon]^n$ .
3. The space of smooth, codimension-zero embeddings  $\bar{j} : [-1, 1]^n \rightarrow \mathbb{R}^n$ .
4. The space of those  $\bar{j}$  which send the origin to the origin.
5. The space of those  $\bar{j}$  which are (restrictions of) linear maps.
6. The space of those  $\bar{f} : [-1, 1]^n \rightarrow [-1, 1]^n$  which preserve the origin.

In particular, all these spaces are homotopy equivalent to the orthogonal group  $O(n)$ .

Now fix an orientation on  $[-1, 1]^n$  and  $\mathbb{R}^n$ . If we consider the subspaces of the above consisting only of orientation-preserving maps, they are all homotopy equivalent to  $SO(n)$ .

*Proof.* 1  $\sim$  2: There is a smooth isotopy from  $[-1, 1]^n$  to  $[-\epsilon, \epsilon]^n$ , for example by scaling by  $s$  as we take  $s$  from 1 to  $\epsilon$ .

2  $\sim$  3: Let  $F_\epsilon$  denote the space of all smooth, codimension-zero embeddings  $[-1, 1]^n \rightarrow \mathbb{R}^n$  with image contained in  $[-\epsilon, \epsilon]$ . We have weak homotopy equivalences of simplicial sets as follows:

$$\text{Sing}(F_\epsilon) \xrightarrow{\sim} \text{colim} (\text{Sing} F_\epsilon \hookrightarrow \text{Sing} F_{2\epsilon} \hookrightarrow \dots) \xrightarrow{\cong} \text{Sing}(3).$$

The first arrow is an equivalence because the inclusions  $\text{Sing}(F_{n\epsilon}) \rightarrow \text{Sing}(F_{(n+1)\epsilon})$  are trivial cofibrations of simplicial sets. (The colimit is modeled as a honest “union” of simplicial sets; because all arrows in the colimit are cofibrations, this also models the homotopy colimit.)  $\text{Sing}(3)$  refers to the singular complex of the space 3; so the final arrow is an isomorphism because  $[-1, 1]^n$  is compact.

3  $\sim$  4: Given  $\bar{j}$ , consider the  $s$ -dependent isotopy  $(s, x) \mapsto \bar{j}(x) - s\bar{j}(0)$  for  $x \in [-1, 1]^n$ .

4  $\sim$  5: Consider the  $s$ -dependent isotopy  $(s, x) \mapsto \frac{1}{s} \bar{j}(sx)$ . (This is the usual trick of homotoping a map to its derivative at the origin.)

4  $\sim$  6: Let  $F'_\epsilon$  denote the collection of those  $j : [-1, 1]^n \rightarrow \mathbb{R}^n$  which preserve the origin, and which have image inside the cube  $[-\epsilon, \epsilon]^n$ . Because scaling respects the origin, we have a sequential diagram of trivial cofibrations

$$F'_\epsilon \hookrightarrow F'_{2\epsilon} \hookrightarrow F'_{4\epsilon} \hookrightarrow \dots$$

whose (homotopy) colimit is equivalent to 4. When  $\epsilon = 1$ , we see that the inclusion of 6 as the first term of the sequential diagram exhibits a homotopy equivalence to 4.

5  $\sim O(n)$ : 5 gives us the equivalence to the space  $GL(n)$ , which in turn is homotopy equivalent to  $O(n)$  by Gram-Schmidt.

The last claim of the proposition (regarding orientation-preserving versions) is proven by running through the exact same isotopies.  $\square$

**Remark 11.12.** In fact, the obvious inclusion  $6 \rightarrow 1$  factors the homotopy equivalence from 6 to 4, so we see that this obvious inclusion is a homotopy equivalence.

**Notation 11.13.** We let  $\mathfrak{eq} \subset \mathbf{Mfld}^{\text{cmpct}}$  denote the collection of isotopy equivalences. Given an object  $X$  of  $\mathbf{Mfld}^{\text{cmpct}}$ , we let  $\mathbf{Mfld}_{/X}^{\text{cmpct}}$  denote the slice category, and let

$$\mathfrak{eq}_{/X} \subset \mathbf{Mfld}_{/X}^{\text{cmpct}}$$

be the collection of morphisms obtained by pulling back  $\mathfrak{eq}$  along the forgetful functor  $\mathbf{Mfld}_X^{\text{cmpct}} \rightarrow \mathbf{Mfld}^{\text{cmpct}}$ .

Concretely, an object of  $\mathbf{Mfld}_{/X}^{\text{cmpct}}$  is the data of a smooth, codimension zero embedding  $j : A \rightarrow X$ , and a morphism from  $j$  to  $j'$  is the data of a smooth, codimension zero embedding  $f : A \rightarrow A'$  satisfying  $j'f = j$ . Such a morphism is in  $\mathfrak{eq}_{/X}$  if and only if  $f$  is an isotopy equivalence.

**Remark 11.14.** The map  $\mathbf{Mfld}^{\text{cmpct}} \rightarrow \mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}]$  induces a diagram

$$\begin{array}{ccc} \mathbf{Mfld}_{/X}^{\text{cmpct}} & \longrightarrow & \mathbf{Mfld}_{/X}^{\text{cmpct}}[\mathfrak{eq}^{-1}] \\ \downarrow & & \downarrow \\ \mathbf{Mfld}^{\text{cmpct}} & \longrightarrow & \mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}] \end{array}$$

hence, by the universal property of localizations, we obtain a map

$$\mathbf{Mfld}_{/X}^{\text{cmpct}}[(\mathfrak{eq}_{/X})^{-1}] \rightarrow \mathbf{Mfld}_{/X}^{\text{cmpct}}[\mathfrak{eq}^{-1}]. \quad (11.5)$$

To prove Lemma 11.7, we will prove that  $\tau$  is homotopic to  $\text{id}_{[-1,1]^3}$  in the localization  $\mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}]$ . The strategy to accomplish this follows techniques we learned from Lemma 5.4.5.10 of [18]<sup>30</sup>, with slight modifications stemming from a small complication:  $[-1, 1]^3$  (because of its corners) does not admit a cover by open disks. The reason for considering “ex”-subscripted versions of smooth objects below is to overcome this issue.

Fix a smooth manifold  $M$  with  $\dim M = n$ , and fix a smooth, codimension-zero embedding  $i : M \hookrightarrow \bar{M}$  where  $\bar{M}$  is a smooth *compact* manifold (possibly with corners), and for which  $i(M) = \bar{M} \setminus \partial \bar{M}$ . (Here,  $\partial \bar{M}$  is the locus of all boundary and corner strata of  $\bar{M}$ .) The prototypical

<sup>30</sup>Another approach would be to utilize techniques from Proposition 2.22 of [4].

example for us is  $i : (-1, 1)^N \hookrightarrow [-1, 1]^N$ . (And in fact, all examples for us will be diffeomorphic to this example.)

We also fix an orientation on  $M$ .

Given this data, we define the following:

**Notation 11.15.** We let  $B_M^\delta$  denote the category where an object is the data of a smooth, orientation-preserving, codimension zero embedding  $j : (-1, 1)^n \hookrightarrow M$  with the property that  $j$  extends to a smooth embedding<sup>31</sup>  $\bar{j} : [-1, 1]^n \hookrightarrow \bar{M}$ . A morphism from  $j$  to  $j'$  is the data of a smooth map  $f : (-1, 1)^n \rightarrow (-1, 1)^n$  for which  $j'f = j$ . (Note it is then automatic that  $f$  extends to a smooth embedding  $\bar{f} : [-1, 1]^n \rightarrow [-1, 1]^n$ .)

**Remark 11.16.** There is a functor  $B_M^\delta \rightarrow \mathbf{Mfld}_{/\bar{M}}^{\text{cmpct}}$  by sending  $j : (-1, 1)^n \rightarrow M$  to  $\bar{j}$ . We note that every morphism  $f$  necessarily maps to an isotopy equivalence  $\bar{f}$  (Example 11.10) so we have an induced map of localizations

$$|B_M^\delta| \rightarrow \mathbf{Mfld}_{/\bar{M}}^{\text{cmpct}}[(\mathbf{eq}_{/\bar{M}})^{-1}]. \quad (11.6)$$

Here,  $|B_M^\delta|$  is the Kan complex obtained by localizing  $B_M^\delta$  with respect to every morphism; in other words, it is the  $\infty$ -groupoid completion of the category  $B_M^\delta$ .

**Notation 11.17.** We let  $\text{Disk}_{\text{ex}}(\bar{M})$  denote the poset whose elements are open subsets  $U \subset M$  that are equal to the image of some  $j$  as above. (Equivalently, these are open subsets  $U \subset M$  for which the closures  $\bar{U}$  (taken in  $\bar{M}$ ) are diffeomorphic to the cube  $[-1, 1]^n$ .) The poset relation is the usual inclusion relation of subsets.

**Remark 11.18.** There is a functor

$$p : B_M^\delta \rightarrow \text{Disk}_{\text{ex}}(\bar{M}) \quad (11.7)$$

—called take the image of  $j$ —and this functor is an equivalence of categories. In fact,  $p$  is a coCartesian fibration classifying a functor from  $\text{Disk}_{\text{ex}}(\bar{M})$  to the category of preorders. More precisely, thinking of every preorder as a category, and hence as a simplicial set (by taking the nerve), this coCartesian fibration classifies the map

$$\text{Disk}_{\text{ex}}(\bar{M}) \rightarrow \mathbf{sSet}, \quad U \mapsto p^{-1}(U). \quad (11.8)$$

.

**Notation 11.19** ( $\text{Emb}_{\text{ex}}^{\text{or}}$ ). Given oriented manifolds  $A$  and  $B$ , let  $\text{Emb}^{\text{or}}(A, B)$  be the space of smooth, codimension zero orientation-preserving embeddings, and—given further the data of smooth compactifications  $A \subset \bar{A}, B \subset \bar{B}$  to manifolds with corners—let  $\text{Emb}_{\text{ex}}^{\text{or}}(\bar{A}, \bar{B}) \subset \text{Emb}^{\text{or}}(A, B)$  denote the subspace consisting of those  $j : A \rightarrow B$  that smoothly extend to an embedding  $\bar{A} \rightarrow \bar{B}$ .

**Lemma 11.20.** Fix an embedding  $j : (-1, 1)^n \rightarrow M$  which extends to a smooth embedding  $\bar{j} : [-1, 1]^n \rightarrow \bar{M}$ . Let  $\text{Emb}_{\text{ex}}^{\text{or}, 0}((-1, 1)^n, (-1, 1)^n) \subset \text{Emb}_{\text{ex}}^{\text{or}}((-1, 1)^n, (-1, 1)^n)$  denote the subspace consisting of those  $f$  that respect the origin.

<sup>31</sup>That  $\bar{j}$  is smooth, is an injection, and has derivative everywhere given by a linear isomorphism, is all important in what follows. For example, that  $\bar{j}$  is an injection allows us to identify the closure of the image of  $j$  with the image of  $\bar{j}$ .

The diagram

$$\begin{array}{ccc} \mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or},0}((-1,1)^n, (-1,1)^n) & \xrightarrow{j^0} & \mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, M) \\ \downarrow & & \downarrow \mathrm{ev}_0 \\ * & \xrightarrow{j(0)} & M \end{array} \quad (11.9)$$

is a homotopy pullback square.

*Proof.* This is a version of Remark 5.4.1.11 [18] for manifolds with corners. We provide the arguments for the reader's convenience.

Letting  $C(t) := (-t, t)^n$  with choice of compactification  $[-t, t]^n$ , we have a filtered diagram of homotopy equivalences parametrized by  $t \in (0, 1]$

$$n \mapsto \mathrm{SingEmb}_{\mathrm{ex}}^{\mathrm{or}}(C(1/2^n), M).$$

We let  $\mathrm{Germ}(M)$  denote the colimit simplicial set—informally, a vertex of  $\mathrm{Germ}(M)$  is the data of a point  $x \in M$  and the germ of an embedding of a cube sending the origin of the cube to  $x$ . It is easy to see that  $\mathrm{Germ}(M)$  is a Kan complex. Restriction defines a homotopy equivalence

$$\mathrm{SingEmb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, M) \xrightarrow{\sim} \mathrm{Germ}(M). \quad (11.10)$$

Now choose an element  $j \in \mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, M)$ . The commutative diagram

$$\begin{array}{ccc} \mathrm{Germ}((-1,1)^n) & \longrightarrow & \mathrm{Germ}(M) \\ \downarrow & & \downarrow \\ \mathrm{Sing}((-1,1)^n) & \xrightarrow{j} & \mathrm{Sing}(M) \end{array} \quad (11.11)$$

is obviously a pullback diagram, and the vertical arrows are Kan fibrations. (Here, we note the importance of having taken  $\mathrm{Sing}(M)$  rather than  $\mathrm{Sing}(\overline{M})$ .) Thus (11.11) is a homotopy pullback diagram.

By applying the equivalence (11.10) we see that the diagram (11.9) is equivalent to the diagram (11.11), and we are finished.  $\square$

**Notation 11.21** ( $\mathcal{C}_M$  and  $B_M$ ). We define a topologically enriched category  $\mathcal{C}_M$  with two objects— $(-1,1)^n$  and  $M$ —and morphism spaces as follows:

$$\mathrm{hom}((-1,1)^n, M) := \mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, M), \quad \mathrm{hom}((-1,1)^n, (-1,1)^n) := \mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, (-1,1)^n),$$

and

$$\mathrm{hom}(M, M) = *, \quad \mathrm{hom}(M, (-1,1)^n) = \emptyset.$$

We let  $B\mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}$  denote the full subcategory spanned by  $(-1,1)^n$ . We then define  $B_M$  to be the fiber product  $\infty$ -category

$$B_M := N(B\mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}) \times_{N(\mathcal{C}_M)} N(\mathcal{C}_M)_{/M}.$$

**Remark 11.22.** Informally, this is the  $\infty$ -category whose objects are orientation-preserving, codimension-zero embeddings  $j : (-1,1)^n \rightarrow M$  that smoothly extend to the chosen compactifications. A morphism from  $j$  to  $j'$  is the data of a map  $f : (-1,1)^n \rightarrow (-1,1)^n$  in  $\mathrm{Emb}_{\mathrm{ex}}^{\mathrm{or}}((-1,1)^n, (-1,1)^n)$  together with an isotopy from  $j'f$  to  $j$ —an isotopy through smooth embeddings that extend to the compactifications.

$B_M$  is in fact a Kan complex.

**Proposition 11.23.**  $B_M$  is homotopy equivalent to  $M$ . In particular, if  $M \cong (-1, 1)^n$  with chosen compactification  $\overline{M} \cong [-1, 1]^n$ , then  $B_M$  is weakly contractible.

*Proof.* Our proof of this parallels Remark 5.4.5.2 of [18], with extra caution sprinkled in (because we are dealing with manifolds with corners).

Let  $\mathcal{C}'_M \subset \mathcal{C}_M$  denote the subcategory for which

$$\mathrm{hom}_{\mathcal{C}'_M}((-1, 1)^n, (-1, 1)^n) \subset \mathrm{hom}_{\mathcal{C}_M}((-1, 1)^n, (-1, 1)^n)$$

consists only of those embeddings that respect the origin (and with all other hom spaces unchanged).

First, we see that inclusion  $\mathcal{C}'_M \hookrightarrow \mathcal{C}_M$  is a (Dwyer-Kan) equivalence of Kan-complex-enriched categories. This follows from Remark 11.12.

Thus, defining  $B'_M = N(\mathrm{hom}_{\mathcal{C}'_M}((-1, 1)^n, (-1, 1)^n)) \times_{N(\mathcal{C}'_M)} N(\mathcal{C}_M)_{/M}$ , the induced map  $B'_M \rightarrow B_M$  is an equivalence.

On the other hand, let  $\mathcal{C}''_M$  denote the simplicially enriched category with the same objects as  $\mathcal{C}'_M$ , but where the only endomorphism of  $(-1, 1)^n$  is the identity, and where we declare  $\mathrm{hom}_{\mathcal{C}''_M}((-1, 1)^n, M) = M$  (with other hom spaces unchanged). Then “evaluation at the origin” defines a functor  $\mathcal{C}'_M \rightarrow \mathcal{C}''_M$ . Though this is certainly not an equivalence, we claim that the induced map

$$B'_M \rightarrow B''_M \tag{11.12}$$

is. Here,  $B''_M$  is defined analogously to  $B_M$  and  $B'_M$ :

$$B''_M := \Delta^0 \times_{N(\mathcal{C}''_M)} N(\mathcal{C}''_M)_{/M}.$$

To see (11.12) is an equivalence, choose an object  $j : (-1, 1)^n \rightarrow M$  of  $B'_M$ . This choice induces a diagram

$$\begin{array}{ccccc} \mathrm{hom}_{\mathcal{C}'_M}((-1, 1)^n, (-1, 1)^n) & \xrightarrow{j \circ} & \mathrm{hom}_{\mathcal{C}'_M}((-1, 1)^n, M) & \longrightarrow & B'_M \\ \downarrow & & \downarrow \mathrm{ev}_0 & & \downarrow \\ * = \mathrm{hom}_{\mathcal{C}''_M}((-1, 1)^n, (-1, 1)^n) & \xrightarrow{j(0)} & M = \mathrm{hom}_{\mathcal{C}''_M}((-1, 1)^n, M) & \longrightarrow & B''_M \end{array}$$

where each row is a homotopy fiber sequence. The left square is a homotopy pullback square by Lemma 11.20.

We thus observe a diagram

$$\begin{array}{ccccc}
& \Omega_j B'_M & \xrightarrow{\quad} & * & \\
\swarrow \sim & \downarrow & & \swarrow & \downarrow \\
\Omega_{j(0)} M & \xrightarrow{\quad} & * & \xrightarrow{\quad} & * \\
\downarrow & \downarrow \text{hom}_{\mathcal{C}'_M}((-1,1)^n, (-1,1)^n) & \xrightarrow{j \circ} & \text{hom}_{\mathcal{C}_{M'}}((-1,1)^n, M) & \downarrow \\
& \downarrow & & \downarrow \text{ev}_0 & \downarrow \\
& * & \xrightarrow{\quad} & M & \downarrow \\
& \downarrow & & \downarrow & \downarrow \\
& * & \xrightarrow{\quad} & B'_M & \downarrow \\
& \downarrow & & \downarrow & \downarrow \\
& * & \xrightarrow{\quad} & B''_M & \downarrow
\end{array}$$

where the topmost face is a homotopy pullback square; in particular, the map  $\Omega_j B'_M \rightarrow \Omega_{j(0)} M$  is a homotopy equivalence (as indicated).

So it suffices to show that the map  $B'_M \rightarrow M$  induces a bijection on connected components. This is obvious, and our proof is finished.  $\square$

**Notation 11.24** ( $q$  and  $\tilde{B}$ ). Note that if  $U$  is any object of  $\text{Disk}_{\text{ex}}(\overline{M})$ , we by definition have a smooth manifold with corners  $\overline{U}$  defined by taking the closure of  $U$ . This results in a functor

$$\text{Disk}_{\text{ex}}(\overline{M}) \rightarrow \mathbf{sSets}, \quad U \mapsto B_U. \quad (11.13)$$

We let  $q : \tilde{B} \rightarrow \text{Disk}_{\text{ex}}(\overline{M})$  denote the coCartesian fibration classifying this functor. (The map  $q$  can be concretely modeled applying the relative nerve construction of [17, Section 3.2.5] to the functor (11.13).) So for example, a vertex of  $B_U$  is a pair  $(U, j : (-1, 1)^n \rightarrow U)$  where  $j$  extends to a smooth embedding  $[-1, 1]^n \rightarrow \overline{U}$ .

Now let  $p$  be the map from (11.7). The inclusion  $p^{-1}(U) \rightarrow B_U$  (from the strictly commuting diagrams to the isotopy-commuting diagrams) is natural in  $U$ , so defines a map of coCartesian fibrations:

$$\begin{array}{ccc}
B_M^\delta & \xrightarrow{\quad} & \tilde{B} \\
& \searrow p & \swarrow q \\
& \text{Disk}_{\text{ex}}(\overline{M}) &
\end{array}$$

**Lemma 11.25.**  $p$  and  $q$  are weak homotopy equivalences of simplicial sets. In particular, the map  $B_M^\delta \rightarrow \tilde{B}$  is a weak homotopy equivalence of simplicial sets.

*Proof.* That  $p$  is a weak equivalence is obvious because  $p$  is an equivalence of categories.  $q$  is a coCartesian fibration, so to show it is a weak equivalence, it suffices to show that  $q$  has contractible fibers. This follows from Proposition 11.23.  $\square$

**Notation 11.26** ( $\psi'$ ). There is also the natural functor

$$\psi' : \tilde{B} \rightarrow B_M \quad (11.14)$$

obtained by post-composing any  $j$  with the inclusion  $U \subset M$ .

**Lemma 11.27.**  $\psi'$  is a weak homotopy equivalence.

*Proof.* One can copy-paste the (second half of) the proof of Lemma 5.4.5.10 in [18]. The only important note is that we have deliberately taken our  $U$  to be open so that the collection of  $U$  forms an open cover of  $M$  (which is a smooth manifold with no boundary or corners).  $\square$

*Proof of Lemma 11.7.* By Lemmas 11.25 and 11.27 we obtain a composition of weak homotopy equivalences  $B_M^\delta \rightarrow \tilde{B} \rightarrow B_M$ , and in particular, a homotopy equivalence of Kan complexes

$$|B_M^\delta| \rightarrow B_M.$$

Choosing an inverse to this homotopy equivalence, consider the composition

$$B_M \xrightarrow{\sim} |B_M^\delta| \xrightarrow{(11.6)} (\mathbf{Mfld}_{/\overline{M}}^{\text{cmpct}})[(\mathfrak{eq}_{/\overline{M}})^{-1}] \xrightarrow{(11.5)} \mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}]_{/\overline{M}}.$$

When  $M = [-1, 1]^3$ , in  $B_M$  there is an edge given by an isotopy-commuting diagram

$$\begin{array}{ccc} [-1, 1]^3 & \xrightarrow{\text{id}} & [-1, 1]^3 \\ & \searrow \tau & \swarrow \text{id} \\ & [-1, 1]^3 & \end{array}$$

because  $\tau$  is orientation-preserving. Its image in the slice  $\mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}]_{/[-1, 1]^3}$  exhibits a homotopy in  $\mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}]$  from  $\tau$  to  $\text{id}_{[-1, 1]^3}$ .

Thus, the natural functor of  $\infty$ -categories  $\mathbf{Mfld}^{\text{cmpct}}[\mathfrak{eq}^{-1}] \rightarrow \mathbf{Liou}_{\text{str}}[\mathfrak{eq}^{-1}]$ —induced by (11.4)—shows that  $T^*\tau$  is homotopic to  $\text{id}_{T^*[-1, 1]^3}$  in  $\mathbf{Liou}_{\text{str}}[\mathfrak{eq}^{-1}]$ .  $\square$



## 12 Equivalent localizations

There are four natural candidates for morphisms to invert (i.e., to declare to be equivalences) between Liouville sectors:

1. (Bordered) deformation equivalences (Definition 4.22),
2. Sectorial equivalences (Definition 2.30),
3. Trivial inclusions (Definition 12.1 below),
4. Movie inclusions (Definition 12.2 below).

Here we prove Theorem 12.8, which shows that the localizations of  $\text{Liou}_{\text{str}}$  with respect to the first two are equivalent, and that localization with respect to all four are equivalent upon stabilization.

**Definition 12.1** (Trivial inclusions, following Section 2.4 of [9]). A *trivial inclusion* is a sectorial embedding  $f : M \rightarrow N$  for which there exists:

1. A smooth family of 1-forms  $\{\lambda_t\}_{t \in [0,1]}$  on  $M$  rendering each  $(M, \lambda_t)$  a Liouville sector, and
2. A smooth isotopy  $\{f_t : M \rightarrow N\}_{t \in [0,1]}$

such that

- (i) For all  $t$ ,  $f_t : (M, \lambda_t) \rightarrow N$  is a sectorial embedding,
- (ii)  $f_0 = f$  and  $\lambda_0 = \lambda_M$ , and
- (iii)  $f_1(M) = N$ . (In light of the other conditions, this condition is equivalent to demanding that  $f_1$  is an isomorphism of Liouville sectors between  $(M, \lambda_1)$  and  $N$ .)

We emphasize here that the  $\lambda_t$  for differing  $t$  may not be related to each other by compactly supported deformations. In particular, the  $f_t$  do not need to form an isotopy through sectorial embeddings for a fixed Liouville structure on  $M$ .

**Definition 12.2** (Movie inclusions). Let  $S = [0, 1]$ , fix some  $1 > \epsilon > 0$  and declare  $S_0 = [0, \epsilon]$  or  $S_0 = [1 - \epsilon, 1]$ . Finally, fix a collared, bordered,  $S$ -parametrized exact family of Liouville structures on  $M$ , and let  $M \times T^*S$  be the associated movie (Construction 2.34) and assume that the  $S$ -parametrized exact family is constant in  $S_0 \subset S$ . There is then a unique Liouville structure on  $M$  for which the inclusion  $M \otimes T^*S_0 \hookrightarrow M \times T^*S$  is a strict sectorial embedding.

For any choice of  $\epsilon$ , either choice of  $S_0$ , and any choice of exact family as above, we call the strict sectorial embedding  $M \otimes T^*S_0 \hookrightarrow M \times T^*S$  a *movie inclusion of intervals*.

**Remark 12.3.** A prototypical trivial cofibration of topological spaces is an inclusion  $X \times \{0\} \hookrightarrow X \times [0, 1]$ . We view movie inclusions as the sectorial analogue of such “endpoint inclusions.”

**Example 12.4.** Fix  $0 < a_i < b_i < 1$  for  $i = 1, \dots, n$ . By allowing ourselves to vary  $M$ , the codimension zero inclusion

$$T^*([a_1, b_1] \times \dots \times [a_n, b_n]) \hookrightarrow T^*[0, 1]^n$$

may be written as a composition of movie inclusions and Liouville isomorphisms. For example, when  $n = 1$ , the inclusion  $[a, b] \hookrightarrow [0, 1]$  may be factored as such a composition by doing the following. Choose

- Any diffeomorphism  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  taking  $[a, b]$  to the interval  $[a, 1]$  (and fixing  $a$ ).
- Any diffeomorphism  $\psi : [\phi^{-1}(0), b] \rightarrow [0, b]$  which restricts to the identity along  $[a, b]$ .

Consider the diagram

$$\begin{array}{ccccccc}
 \mathbb{R} & \xrightarrow[\cong]{\phi} & \mathbb{R} & \xlongequal{\quad} & \mathbb{R} & \xrightarrow[\cong]{\phi^{-1}} & \mathbb{R} \\
 \uparrow \subset & & \uparrow \subset & & \uparrow \subset & & \uparrow \subset \\
 [a, b] & \xrightarrow[\cong]{} & [a, 1] & \xrightarrow{\iota} & [0, 1] & \xrightarrow[\cong]{} & [\phi^{-1}(0), b] & \xrightarrow[\cong]{\psi} & [0, b] & \xrightarrow{\iota} & [0, 1]
 \end{array}$$

where each  $\iota$  is a standard inclusion. Then the bottom horizontal arrows, upon passage to cotangent bundles, exhibit the standard inclusion  $T^*[a, b] \hookrightarrow T^*[0, 1]$  as a composition of movie inclusions and Liouville isomorphisms.

For  $n = 2$ , note that  $T^*$  of the inclusion  $[a_1, 1] \times [a_2, 1] \hookrightarrow [0, 1]^2$  can be written as a composition of two movie inclusions, where we change which of the two components of the square are treated as the  $M$  component (in the notation of Definition 12.2). This is what we mean by “varying  $M$ .”

Now suppose  $j : [0, 1]^n \hookrightarrow [0, 1]^n$  is any smooth, codimension zero embedding (that need not respect the stratification of the cubes). Then we may find a commutative diagram of embeddings as follows:

$$\begin{array}{ccccc}
 [0, 1]^n & \xrightarrow{j} & [0, 1]^n & \xrightarrow{h} & [0, 1]^n \\
 h' \downarrow & & j' \nearrow & \cong & \\
 [0, 1]^n & & & & 
 \end{array}$$

Here, the arrows labeled  $h$  and  $h'$  are compositions of the form  $[0, 1]^n \cong [\epsilon, 1 - \epsilon]^n \hookrightarrow [0, 1]^n$  – i.e., diffeomorphisms followed by a standard inclusion – so that the images of  $h$  and  $h'$  are each bounded away from the boundary of  $[0, 1]^n$ . Then there exists a *diffeomorphism*  $j' : [0, 1]^n \rightarrow [0, 1]^n$  extending the top horizontal composition in the diagram. That  $hj = j'h'$  shows that, upon inverting  $T^*h$  and  $T^*h'$  (in particular, upon inverting movie inclusions) the arbitrary embedding  $j$  may be obtained up to homotopy as a composition of  $T^*$  of diffeomorphisms and movie inclusions.

In particular, inverting movie inclusions renders  $T^*$  of isotopy equivalences of cubes into homotopy-invertible morphisms. (This will also be a consequence of Theorem 12.8 below.)

**Proposition 12.5.** Let  $\mathcal{C}$  be an  $\infty$ -category and let  $S \subset T$  be two collections of morphisms in  $\mathcal{C}$ . Assume every morphism of  $T$  becomes invertible in the localization  $\mathcal{C}[S^{-1}]$  (more precisely, that the image of  $T$  under the map  $\mathcal{C} \rightarrow \mathcal{C}[S^{-1}]$  is contained in the collection of equivalences of  $\mathcal{C}[S^{-1}]$ ). Then the natural map

$$\mathcal{C}[S^{-1}] \rightarrow \mathcal{C}[T^{-1}]$$

is an equivalence of  $\infty$ -categories.

*Proof.* We first set some notation: Let  $i_S$  and  $i_T$  denote the localization maps from  $\mathcal{C}$  to  $\mathcal{C}[S^{-1}]$  and  $\mathcal{C}[T^{-1}]$ , respectively. The natural map  $h : \mathcal{C}[S^{-1}] \rightarrow \mathcal{C}[T^{-1}]$ , by the universal property of localizations, induces a homotopy  $i_T \sim h \circ i_S$ .

By hypothesis, the universal property of localizations guarantees

1. a map  $j : \mathcal{C}[T^{-1}] \rightarrow \mathcal{C}[S^{-1}]$ ,
2. a homotopy  $j \circ i_T \sim i_S$ , and hence (by pulling back along  $i_S$ )

3. a homotopy  $j \circ h \sim \text{id}_{\mathcal{C}[S^{-1}]}$ .

Similar reasoning shows that  $h \circ j \sim \text{id}_{\mathcal{C}[T^{-1}]}$ .  $\square$

**Proposition 12.6.** Let  $f : X \rightarrow Y$  be a (not necessarily strict) sectorial embedding.

- (a)  $f$  is a sectorial equivalence if and only if  $f$  lifts to (i.e., may be equipped with data realizing  $f$  as being) a bordered deformation equivalence.
- (b) If  $f$  is a trivial inclusion, then  $f$  is a bordered deformation equivalence (hence, by (a), a sectorial equivalence).
- (c) If  $f$  is a movie inclusion, then  $f$  is a bordered deformation equivalence (hence, by (a), a sectorial equivalence).

We note that the proof of this proposition will not rely on any of the results of Section 10—this is logically necessary, as Section 10 relies on Remark 12.7 below, which in turn relies on the proof methods of the present proposition.

*Proof.* (a): It is obvious that any sectorial equivalence is a bordered deformation equivalence (by choosing, for example, compactly supported deformations). So now suppose  $f : M \rightarrow N$  is a bordered deformation equivalence. This means the image of  $f$  under the composition  $\text{Liou}_{\text{str}} \rightarrow \text{Liou}_{\Delta} \rightarrow \text{Liou}_{\Delta}^{\text{Def}}$  is an equivalence in  $\text{Liou}_{\Delta}^{\text{Def}}$ . Moreover, the inclusion  $\text{Liou}_{\Delta} \rightarrow \text{Liou}_{\Delta}^{\text{Def}}$  is an equivalence of  $\infty$ -categories by Theorem 9.3, so the existence of inverse data exhibiting  $f$  as an equivalence in  $\text{Liou}_{\Delta}^{\text{Def}}$  implies the existence of inverse data exhibiting  $f$  as an equivalence in  $\text{Liou}_{\Delta}$ ; but by applying Proposition 5.19, the 2-simplices exhibiting  $f$  as an equivalence in  $\text{Liou}_{\Delta}$  supply an inverse map  $g : N \rightarrow M$  and isotopies of sectorial embeddings demonstrating that  $f$  is a sectorial equivalence.

(b): Given the data exhibiting  $f : M \rightarrow N$  as a trivial inclusion (Definition 12.1), one may construct data exhibiting  $f$  as a bordered deformation equivalence. In the notation of Definition 4.22, one may for example choose  $g = f_1^{-1}$ , and bordered deformations extending  $(f_t)_*(\lambda_t^M)$ .

(c): We use the notation from Definition 12.2. Let  $\lambda_0$  denote the Liouville structure on  $M$  associated to  $S_0$ , and choose an isotopy  $\{j_t : S_0 \rightarrow [0, 1]\}_{0 \leq t \leq 1}$  where  $j_0$  is the inclusion and  $j_1$  is a diffeomorphism. From hereon, by the sector  $M$ , we mean the sector  $(M, \lambda_0)$ . Then  $j_t$  induces an isotopy, through exact sectorial embeddings, from  $\text{id}_M \times T^*j_0$  to  $\text{id}_M \times T^*j_1$ . There is an obvious inverse (up to isotopy) from  $M \otimes T^*[0, 1]$  to  $M \otimes T^*S_0$ , so  $\text{id}_M \times T^*j_0$  is a bordered deformation equivalence from  $M \otimes T^*S_0$  to  $M \otimes T^*[0, 1]$ . (In fact, the deformations of Liouville structures may be chosen to be constant.)

Now note that, by definition of movie inclusion, there is a bordered deformation  $\{\lambda_s^{M \times T^*[0, 1]}\}_{s \in [0, 1]}$  of Liouville forms from  $\lambda_0 + \mathbf{pdq}$  to the movie Liouville form (2.4) on  $M \times T^*[0, 1]$ . Thus, the “identity map”  $M \otimes T^*[0, 1] \rightarrow M \times T^*[0, 1]$  (from the stabilization of  $M$  to the movie of  $M$ ), coupled with  $\{\lambda_s^{M \times T^*[0, 1]}\}_{s \in [0, 1]}$ , is a non-strict bordered deformation equivalence. The inverse is given by the “identity map” along with the reverse deformation  $\{\lambda_{1-s}^{M \times T^*[0, 1]}\}_{s \in [0, 1]}$ , where obviously the  $s$ -concatenation of the reverse and forward deformations may be deformed to the constant  $s$ -parametrized deformation.

Now we see that the strict map  $\text{id}_M \times T^*j_0$  from  $M \otimes T^*S_0$  to the movie  $M \times T^*[0, 1]$  is a bordered deformation equivalence—one simply composes and concatenates the inverse data from the previous two paragraphs to exhibit the bordered deformation inverse to  $\text{id}_M \times T^*j_0$ .  $\square$

**Remark 12.7.** When  $\max A' = \max A$ , we claim that the map  $\phi_{A' \subset A}$  is an equivalence of Liouville sectors. (This was used crucially in Construction 10.11.) To see why this claim is true, note we have a commutative diagram

$$\begin{array}{ccc}
& & M_{\max A'} \otimes T^* \Delta^{A'} \otimes T^*[-1, 0]^{A \setminus A'} \\
& \nearrow \sim & \downarrow \sim \\
M_{\max A'} \otimes T^*[-1, 0]^A & \longrightarrow & M_{A'} \times T^* \Delta^{A'} \otimes T^*[-1, 0]^{A \setminus A'} \\
\downarrow \cong & & \downarrow \phi_{A' \subset A} \\
M_{\max A} \otimes T^*[-1, 0]^A & \longrightarrow & M_A \times T^* \Delta^A \\
& \searrow \sim & \uparrow \sim \\
& & M_A \otimes T^* \Delta^A.
\end{array}$$

The claim follows if we can prove that both (unlabeled) horizontal arrows are sectorial equivalences. This follows from more or less the same method as in Proposition 12.6 (c). Namely, the inclusions of  $[-1, 0]^A$  into  $\Delta^{A'} \times [-1, 0]^{A \setminus A'}$  and  $\Delta^A$  are both isotopy equivalences of smooth manifolds with corners. (Warning: this does not mean the inclusions are isotopic to diffeomorphisms; see Definition 11.9.) Thus, the horizontal arrows are sectorial equivalences if the codomains are given the constant movie structure. One can then apply a bordered deformation from the constant movie Liouville structure to the movie structures of the original codomains of the horizontal arrows. As in the proof of Proposition 12.6 (c), this exhibits the horizontal arrows as bordered deformation equivalences, and hence sectorial equivalences by Proposition 12.6 (a).

**Theorem 12.8.** In  $\text{Liou}_{\text{str}}$ , let  $\mathbf{eq}_{\text{triv}}$  be the collection of strict trivial inclusions,  $\mathbf{eq}$  the collection of strict sectorial equivalences,  $\mathbf{eq}_{\text{movie}}$  the collection of (strict) movie inclusions, and  $\mathbf{eq}_{\mathcal{D}\text{ef}}$  the collection of strict sectorial embeddings that may be promoted to bordered deformation equivalences. Then the natural maps

- (a)  $\text{Liou}_{\text{str}}[\mathbf{eq}^{-1}] \rightarrow \text{Liou}_{\text{str}}[(\mathbf{eq}_{\mathcal{D}\text{ef}})^{-1}]$ ,
- (b)  $\text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}^{\diamond})^{-1}] \rightarrow \text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\mathcal{D}\text{ef}}^{\diamond})^{-1}]$ ,
- (c)  $\text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\text{triv}}^{\diamond})^{-1}] \rightarrow \text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\mathcal{D}\text{ef}}^{\diamond})^{-1}]$ , and
- (d)  $\text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\text{movie}}^{\diamond})^{-1}] \rightarrow \text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\mathcal{D}\text{ef}}^{\diamond})^{-1}]$

are equivalences of  $\infty$ -categories. (Note that the final three equivalences are stabilized, while the first is not. Also note that the above natural maps are guaranteed by Proposition 12.6.)

In particular, whether one localizes  $\text{Liou}_{\text{str}}^{\diamond}$  by trivial inclusions, by sectorial equivalences, by bordered deformation equivalences, or by movie inclusions, one obtains an  $\infty$ -category equivalent to  $\mathcal{L}\text{iou}^{\diamond}$ .

*Proof.* (a) This follows from Proposition 12.6 (a) and Proposition 12.5.

(b) This follows from the above by applying Lemma 11.2. (See also Corollary 11.3.)

The rest of this proof is devoted to establishing (c) and (d).

By Proposition 12.5 and the equivalence (b), we are reduced to showing that stabilized sectorial equivalences become homotopy invertible in  $\mathcal{D}$ , where  $\mathcal{D}$  is either of  $\text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\text{triv}}^{\diamond})^{-1}]$  or  $\text{Liou}_{\text{str}}^{\diamond}[(\mathbf{eq}_{\text{movie}}^{\diamond})^{-1}]$ .

So fix a strict sectorial equivalence  $f : M \rightarrow N$ . By definition, there is a (not necessarily strict) sectorial embedding  $g : N \rightarrow M$  and isotopies  $gf \sim \text{id}_M, fg \sim \text{id}_N$  through (not necessarily strict) sectorial embeddings. Endow  $N$  with a form  $\lambda_0^N$  for which  $g$  is a strict map to  $M$ . Then, the isotopy  $fg \sim \text{id}_N$  and Proposition 2.45 guarantee a strict map from  $(N, \lambda_0^N) \otimes T^*[0, 1]$  to a movie  $N \times T^*[0, 1]$  realizing this isotopy, fitting into the following commutative diagram in  $\text{Liou}_{\text{str}}$ :

$$\begin{array}{ccccc}
(N, \lambda_0^N) \otimes T^*[0, \epsilon] & & \xrightarrow{\text{id}_N \times T^* \iota_0} & & \\
\text{id}_N \times T^* \iota_0 \downarrow \simeq & & \simeq & & \\
(N, \lambda_0^N) \otimes T^*[0, 1] & \xrightarrow{\text{Prop 2.45}} & & N \times T^*[0, 1] & \\
\text{id}_N \times T^* \iota_1 \uparrow \simeq & & & \simeq \uparrow \text{id}_N \times T^* \iota_1 & \\
(N, \lambda_0^N) \otimes T^*[1 - \epsilon, 1] & \xrightarrow{g \times \text{id}} M \otimes T^*[1 - \epsilon, 1] & \xrightarrow{f} & N \otimes T^*[1 - \epsilon, 1] &
\end{array}$$

Here,  $\iota_0 : [0, \epsilon] \rightarrow [0, 1]$  and  $\iota_1 : [1 - \epsilon, 1] \rightarrow [0, 1]$  are the standard inclusions. Note also we are identifying  $f$  with its stabilization. The arrows labeled by  $\simeq$  are simultaneously movie inclusions and trivial inclusions, so are homotopy invertible in  $\mathcal{D}$ . It follows that in  $\mathcal{D}$ ,  $f$  admits a right homotopy inverse.

Likewise, one has a commutative diagram as follows in  $\text{Liou}_{\text{str}}$ :

$$\begin{array}{ccccc}
M \otimes T^*[0, \epsilon] & & \xrightarrow{\text{id}_M \times T^* \iota_0} & & \\
\text{id}_M \times T^* \iota_0 \downarrow \simeq & & \simeq & & \\
M \otimes T^*[0, 1] & \xrightarrow{\text{Prop 2.45}} & & M \times T^*[0, 1] & \\
\text{id}_M \times T^* \iota_1 \uparrow \simeq & & & \simeq \uparrow \text{id}_M \times T^* \iota_1 & \\
M \otimes T^*[1 - \epsilon, 1] & \xrightarrow{f} N \otimes T^*[1 - \epsilon, 1] & \xrightarrow{g \times \text{id}} & (M, \lambda_1^M) \otimes T^*[1 - \epsilon, 1] &
\end{array}$$

Here, we have chosen a form  $\lambda_1^M$  to render  $g$  strict, along with a movie structure on  $M \times T^*[0, 1]$  to render all maps strict and to invoke Proposition 2.45. The arrows labeled by  $\simeq$  again become equivalences in  $\mathcal{D}$ , and it follows that  $f$  admits a left homotopy inverse.

$f$  is thus an equivalence in  $\mathcal{D}$ . □

## 12.1 The wrapped Fukaya category is unchanged by equivalences

We have the following consequence of Theorem 12.8:

**Proposition 12.9.** Fix Liouville sectors  $M$  and  $N$ .

- (a) Let  $f : M \rightarrow N$  be a sectorial equivalence. Then the induced map on the stabilized wrapped Fukaya categories  $f_* : \mathcal{W}^\diamond(M) \rightarrow \mathcal{W}^\diamond(N)$  is an equivalence of  $A_\infty$ -categories.
- (b) Let  $f : M \rightarrow N$  be a bordered deformation equivalence. Then the induced map on the stabilized wrapped Fukaya categories  $f_* : \mathcal{W}^\diamond(M) \rightarrow \mathcal{W}^\diamond(N)$  is an equivalence of  $A_\infty$ -categories.
- (c) Now suppose  $M$  and  $N$  are Weinstein sectors, and that  $f : M \rightarrow N$  is a bordered deformation equivalence. Then the induced map on wrapped Fukaya categories is an equivalence.

*Proof.* The work [9] exhibited the wrapped Fukaya category functor

$$\text{Liou}_{\text{str}} \rightarrow A_\infty \text{Cat} \tag{12.1}$$

and showed that trivial inclusions induce equivalences of  $A_\infty$ -categories. Thus, letting  $\text{Cat}_{A_\infty}$  be the localization of  $A_\infty\text{Cat}$  along quasi-equivalences, one has an induced functor  $\mathcal{W}$  as below:

$$\text{Liou}_{\text{str}} \rightarrow \text{Liou}_{\text{str}}[\mathbf{eq}_{\text{triv}}^{-1}] \xrightarrow{\mathcal{W}} \text{Cat}_{A_\infty} .$$

There are natural stabilization transformations  $\mathcal{W}(M) \rightarrow \mathcal{W}(M \times T^*[0, 1])$ , so we have an induced functor

$$\text{Liou}_{\text{str}}^\diamond \rightarrow \text{Cat}_{A_\infty}, \quad M \mapsto \mathcal{W}^\diamond(M) = \text{hocolim}_k \mathcal{W}(M \times T^*[0, 1]^k)$$

factoring through  $\text{Liou}_{\text{str}}^\diamond[(\mathbf{eq}_{\text{triv}}^\diamond)^{-1}] \simeq \text{Liou}_{\text{str}}[\mathbf{eq}_{\text{triv}}^{-1}]^\diamond$ . Thus (a) and (b) follow from Theorem 12.8 (b) and (c).

(c) This follows from (b) by noting that the natural maps  $\mathcal{W}(M) \rightarrow \mathcal{W}(M \times T^*[0, 1])$  are equivalences for Weinstein sectors, and hence so are the maps  $\mathcal{W}(M) \rightarrow \mathcal{W}^\diamond(M)$ .  $\square$

**Remark 12.10.** Note that the proof of Proposition 12.9 required no mention of holomorphic disks – indeed, the only requisite arguments concerning holomorphic disks are fully contained in exhibiting the functor (12.1). The universal property of localizations, together with the geometry of *sectors* (without reference to branes or holomorphic disks) established in Theorem 12.8, allowed us to conclude the proposition. While it is not too onerous to prove Proposition 12.9 by directly using Floer-theoretic methods, any such proof would first pass through the geometry used in proving Theorem 12.8, then require additional arguments about branes and holomorphic disks (by riffing on arguments already used in [9]’s Lemma 3.33, for example).

## 13 Incorporating tangential structures

To define a  $\mathbb{Z}$ -graded (as opposed to two-periodic) wrapped Fukaya category, one must trivialize a natural map from a Liouville sector to  $BU(1)$ . One must also trivialize certain tangentially classified obstructions to render the coefficient ring  $\mathbb{Z}$ . Likewise, the construction of Nadler-Shende's microlocal category requires a trivialization of the stable normal Gauss map. So it is natural and necessary to consider not only Liouville sectors, but Liouville sectors equipped with particular tangential structures (such as trivializations).

Here we give a description of the  $\infty$ -categories of Liouville sectors equipped with tangential structures, paying particular attention to the symmetric monoidal structures we can endow upon them. We omit some details that rely on classical differential-topological constructions (such as equipping a bundle with a locally finite collection of local trivializations, and observing that the space of such choices is contractible upon allowing for common refinements).

**Notation 13.1.** Throughout this section,  $*$  refers to the one-point space.

### 13.1 Preliminaries: The $\infty$ -categories of (stable) vector bundles

First, suppose that  $\mathcal{C}$  is a symmetric monoidal category, and fix an object  $B \in \mathcal{C}$ . It is not the case that the slice category  $\mathcal{C}_{/B}$  (an object of which is the data of an object  $X$  equipped with a map  $X \rightarrow B$ ) has an induced symmetric monoidal structure.

**Recollection 13.2.** However,  $\mathcal{C}_{/B}$  can be rendered symmetric monoidal each time  $B$  is given the structure of a commutative algebra object. Informally, the monoidal product  $(X \rightarrow B) \otimes (Y \rightarrow B)$  is encoded by the composition  $X \otimes Y \rightarrow B \otimes B \rightarrow B$ . This holds true when  $\mathcal{C}$  is a symmetric monoidal  $\infty$ -category and  $B$  is given an  $E_\infty$ -algebra structure. (See Notation 2.2.2.3 and Theorem 2.2.2.4 of [18]; in the notation of loc. cit., we are concerned with the case  $S = N(\text{Fin}_*)$ .)

**Example 13.3.** We let  $\text{Top}$  denote the  $\infty$ -category of topological spaces. We let  $BO$  denote the usual classifying space of stable vector bundles, which we model concretely as the colimit of infinite Grassmannians:

$$BO := \text{colim} (BO(1) \rightarrow BO(2) \rightarrow \dots), \quad BO(n) = \{n\text{-planes in } \mathbb{R}^\infty\}.$$

We let  $\text{Top}_{/BO}$  denote the slice  $\infty$ -category. Concretely, an object of  $\text{Top}_{/BO}$  is the data of a topological space  $X$  equipped with a map  $f : X \rightarrow BO$ . A 1-simplex in  $\text{Top}_{/BO}$  is the data of a homotopy-commuting diagram

$$\begin{array}{ccc} X & \xrightarrow{g} & X' \\ & \searrow f & \swarrow f' \\ & BO & \end{array}$$

together with the data of the homotopy  $f' \circ g \sim f$ . Endowing  $BO$  with the standard infinite-loop-space structure (which models the direct sum of stable vector bundles), one obtains a symmetric monoidal structure on  $\text{Top}_{/BO}$ .

**Remark 13.4.** Suppose one has a model for the infinite-loop space structure on a space  $B$  using the language of operads à la May and Boardman-Vogt. For example, one might use the linear isometries operad to model the infinite-loop space structure on  $BO$  or  $BU$ . (See for example Chapter I of [21].) Then one can endow the usual topologically enriched category  $\text{Top}_{/B}$  with the structure

of an  $E_\infty$ -algebra in topologically enriched categories. Taking the homotopy coherent nerve<sup>32</sup>, one obtains a symmetric monoidal  $\infty$ -category  $N(Top/B)^\otimes$ —concretely, the  $E_\infty$ -algebra structure, via the  $\infty$ -categorical Grothendieck construction (otherwise known as the unstraightening), models an  $\infty$ -category equipped with a coCartesian fibration over the (nerve of the multicategory of the particular model of the)  $E_\infty$ -operad, which in turn is fibered over  $N(\mathcal{F}in_*)$ . One can then exhibit an equivalence of symmetric monoidal  $\infty$ -categories

$$N(Top/B)^\otimes \simeq \mathcal{J}op_B^\otimes$$

where the righthand side is the symmetric monoidal  $\infty$ -category from Recollection 13.2.

We only outline the proof: Both sides are most naturally modeled as having a coCartesian fibration not to  $N(\mathcal{F}in_*)$ , but to the  $\infty$ -operad modeling (for example) the linear isometries operad. By definition of the nerve construction, one observes that every simplex of  $N(Top/B)^\otimes$  defines a simplex of  $\mathcal{J}op_B^\otimes$  by the characterization of simplices of  $\mathcal{J}op_B^\otimes$  given in Notation 2.2.2.3 of [18]. This map of simplicial sets is obviously essentially surjective and respects coCartesian edges; it remains to show that the map is an equivalence of  $\infty$ -categories, which follows from a straightforward computation of mapping spaces. (To compute mapping spaces on the lefthand side, one may invoke that  $Top/B$  arises as the nerve of a simplicial model category, then use the usual formula for mapping spaces in a multicategory associated to a symmetric monoidal category. The righthand side is computed using the usual fibration sequence for computing morphism spaces in slice categories<sup>33</sup>, then the formula for computing mapping spaces in an  $\infty$ -operad<sup>34</sup>.)

Let  $\mathcal{V}ecBun_{\mathbb{R}}$  denote the topologically enriched category whose objects are vector bundles  $(E \rightarrow X)$ , and whose morphisms are pairs  $(f : X \rightarrow X', \tilde{f} : E \rightarrow E')$  where  $f$  is continuous and  $\tilde{f}$  is a bundle map (meaning it induces isomorphisms along fibers) over  $f$ . As an example, the object  $(\mathbb{R}^n \rightarrow *)$  given by the trivial rank  $n$  bundle over a point has endomorphism space homeomorphic to the Lie group  $GL(n)$ . For simplicity, we will assume that every  $X$  is (homotopy equivalent to) a CW complex—this will later allow us to use paracompactness arguments.

We endow  $\mathcal{V}ecBun_{\mathbb{R}}$  with the symmetric monoidal structure induced by Whitney sum and direct product of spaces; so the product of  $(E \rightarrow X)$  and  $(E' \rightarrow X')$  is given by the bundle  $p_X^*E \oplus p_{X'}^*E'$  over  $X \times X'$ . We let

$$(\mathcal{V}ecBun_{\mathbb{R}})^\oplus$$

denote the symmetric monoidal  $\infty$ -category obtained by taking the homotopy coherent nerve. (We note the change in font from  $\mathcal{V}ecBun$  to  $\mathcal{V}ecBun$ .) There is a stabilizing endofunctor

$$\diamond : \mathcal{V}ecBun_{\mathbb{R}} \rightarrow \mathcal{V}ecBun_{\mathbb{R}}, \quad (E \rightarrow X) \mapsto ((E \oplus \underline{\mathbb{R}}) \rightarrow X)$$

which is modeled by taking a product with the object  $(\mathbb{R} \rightarrow *)$ . (Here,  $\underline{\mathbb{R}}$  denotes the trivial vector bundle.) Obviously, the cyclic permutation action on  $(\mathbb{R}^{\oplus 3} \rightarrow *)$ —which may be identified with an orientation-preserving element of  $O(3)$ —is homotopic to the identity. Thus the colimit of the diagram of  $\infty$ -categories

$$\mathcal{V}ecBun_{\mathbb{R}} \xrightarrow{\diamond} \mathcal{V}ecBun_{\mathbb{R}} \xrightarrow{\diamond} \dots$$

inherits a symmetric monoidal structure by Proposition 11.6.

<sup>32</sup>By a version of Proposition 2.1.1.27 of [17], any symmetric monoidal, topologically enriched category yields a symmetric monoidal  $\infty$ -category by taking the homotopy coherent nerve.

<sup>33</sup>For example, Proposition 5.5.5.12 of [17].

<sup>34</sup>For example, (2) of Definition 2.1.1.10 of [18].



**Notation 13.5.** We let

$$(\mathrm{VecBun}_{\mathbb{R}}^{\diamond})^{\oplus}$$

denote the resulting symmetric monoidal  $\infty$ -category. When considering complex vector bundles, we likewise let

$$(\mathrm{VecBun}_{\mathbb{C}}^{\diamond})^{\oplus}$$

denote the symmetric monoidal  $\infty$ -category of CW complexes equipped with stable complex vector bundles.

**Remark 13.6.** The functor of taking a product with  $(\mathbb{R} \rightarrow *)$  is equivalent to taking a product with  $(\mathbb{R} \rightarrow T^*[0, 1])$  because  $T^*[0, 1] \rightarrow *$  is an equivalence in the  $\infty$ -category  $\mathcal{T}\mathrm{op}$ . Thus,  $\mathrm{VecBun}^{\diamond}$  admits a model by stabilizing with respect to direct product with  $(\mathbb{R} \rightarrow T^*[0, 1])$ . This model is more on-the-nose compatible with the stabilization of  $\mathrm{Liou}_{\mathrm{str}}$ .

**Remark 13.7.** Concretely, an object of  $\mathrm{VecBun}_{\mathbb{R}}^{\diamond}$  is the data of a pair  $((E \rightarrow X), k)$ —where  $(E \rightarrow X)$  is a real vector bundle over a CW base, and  $k \geq 0$  is some integer—up to an equivalence relation. The relation identifies  $((E \rightarrow X), k)$  with the object  $((E \oplus \underline{\mathbb{R}} \rightarrow X), k + 1)$ , where  $\underline{\mathbb{R}}$  is the trivial rank one real vector bundle over  $X$ .

In particular, there are objects  $((\mathbb{R} \rightarrow *), k)$  which one may think of as a stable rank one vector bundle destabilized  $k$  times. Each of these has a monoidal inverse given by  $((\mathbb{R}^{\oplus k-1} \rightarrow *), 0)$ .

**Lemma 13.8.** There exists an equivalence of symmetric monoidal  $\infty$ -categories

$$(\mathrm{VecBun}_{\mathbb{R}}^{\diamond})^{\oplus} \simeq \mathcal{T}\mathrm{op}_{/BO}^{\otimes}, \quad (\mathrm{VecBun}_{\mathbb{C}}^{\diamond})^{\oplus} \simeq \mathcal{T}\mathrm{op}_{/BU}^{\otimes}.$$

*Sketch of proof.* There are several proof methods. Because this is a standard result, we sketch one proof method and omit the details. First, we recall that all  $(E \rightarrow X) \in \mathrm{VecBun}_{\mathbb{R}}$  are assumed to have a CW base, and that every CW complex is paracompact.

$\mathrm{VecBun}_{\mathbb{R}}^{\diamond}$  can be written as the base of a right fibration

$$p : \widetilde{\mathrm{VecBun}_{\mathbb{R}}^{\diamond}} \rightarrow \mathrm{VecBun}_{\mathbb{R}}^{\diamond}.$$

The vertices of  $\widetilde{\mathrm{VecBun}_{\mathbb{R}}^{\diamond}}$  consist of vector bundles  $(E \rightarrow X)$  equipped with data of local trivializations subordinate to a locally finite cover<sup>35</sup>, along with partitions of unity sufficient to define a map  $j : E \rightarrow \mathbb{R}^{\infty}$  for which  $j|_{E_x}$  is a linear injection for all  $x \in X$ . Higher simplices are encoded by common refinements of such covering data, along with homotopies of trivializations and of the partitions of unity. (In particular, the ability to construct edges using refinements of covers is one utility of the model of  $\infty$ -categories; it would be more difficult to construct a topologically enriched category incorporating such refinements.)

The fibers of  $p$  are contractible, so  $p$  is an equivalence. Modeling  $BO$  as a colimit of Grassmannians of planes in  $\mathbb{R}^{\infty}$ , we obtain a functor from  $\widetilde{\mathrm{VecBun}_{\mathbb{R}}^{\diamond}}$  to  $\mathcal{T}\mathrm{op}_{/BO}^{\otimes}$ ; on an object, we assign the map taking  $x \in X$  to  $j(E_x)$ . (The basic idea of this construction is age-old; see for example Lemma 5.3 of [23].) The details omitted are the constructions of  $\widetilde{\mathrm{VecBun}_{\mathbb{R}}^{\diamond}}$ , the symmetric monoidal  $\infty$ -category structure on  $\mathrm{VecBun}_{\mathbb{R}}^{\diamond}$  and the symmetric monoidal enrichment of the map to  $\mathcal{T}\mathrm{op}_{/BO}$ .  $\square$

---

<sup>35</sup>Here we are using paracompactness.

## 13.2 Tangential structures respect direct products

**Lemma 13.9.** There exists a symmetric monoidal functor

$$(\mathbf{Liou}_{\text{str}})^{\otimes} \rightarrow (\mathbf{VecBun}_{\mathbb{C}})^{\oplus}$$

which sends a Liouville sector  $X$  to  $(TX \rightarrow X)$  with some choice of almost-complex structure compatible with  $\omega_X$ .

*Proof.* There is a symmetric monoidal functor from  $\mathbf{Liou}_{\text{str}}$  to  $\mathbf{VecBun}_{\mathbb{R}}$  sending any Liouville sector to its tangent bundle. Now note the following two functors to the  $\infty$ -category of spaces:

$$(\mathbf{Liou}_{\text{str}})^{\text{op}} \rightarrow \mathcal{T}\text{op}, \quad X \mapsto \{\omega_X\text{-compatible almost-complex structures } J\}$$

and

$$\mathbf{VecBun}_{\mathbb{R}}^{\text{op}} \rightarrow \mathcal{T}\text{op}, \quad (E \rightarrow X) \mapsto \{\text{endomorphisms } J \text{ of } E \text{ that square to } -1.\}$$

The first of these functors sends every  $X$  to a contractible space, as is well-known; hence it admits a symmetric monoidal promotion. The second is a lax symmetric monoidal functor. The following diagram commutes up to natural transformation:

$$\begin{array}{ccc} (\mathbf{Liou}_{\text{str}})^{\text{op}} & \xrightarrow{X \mapsto TX} & \mathbf{VecBun}_{\mathbb{R}}^{\text{op}} \\ & \searrow X \mapsto \{J\} & \swarrow (E \rightarrow X) \mapsto \{J\} \\ & \mathcal{T}\text{op} & \end{array}$$

The natural transformation is the inclusion of  $\omega$ -compatible  $J$  into the space of all  $J$ , and can obviously be upgraded to a lax monoidal natural transformation. Because the above functors are (lax) symmetric monoidal, the fibrations classifying these functors are  $\infty$ -operads, and the lax symmetric monoidal natural transformation results in a diagram of  $\infty$ -operads:

$$\begin{array}{ccccc} \widetilde{(\mathbf{Liou}_{\text{str}})^{\otimes}} & \longrightarrow & \widetilde{\mathbf{VecBun}_{\mathbb{R}}^{\oplus}} & \xrightarrow{\sim} & (\mathbf{VecBun}_{\mathbb{C}})^{\oplus} \\ \downarrow \sim & & \downarrow & & \\ (\mathbf{Liou}_{\text{str}})^{\otimes} & \longrightarrow & \mathbf{VecBun}_{\mathbb{R}}^{\oplus} & & \end{array}$$

References for the  $\infty$ -operadic claims include Example 2.4.2.4, Proposition 2.4.2.5, and Remark 2.4.2.7 (to deal with the contravariance of the functors we classify) of [18].

We note that, as  $\infty$ -categories, the total space  $\widetilde{\mathbf{VecBun}_{\mathbb{R}}^{\oplus}}$  of the classifying fibration is equivalent to  $\mathbf{VecBun}_{\mathbb{C}}$ . The equivalence extends (as indicated by the rightmost horizontal arrow in the diagram) to an equivalence of symmetric monoidal  $\infty$ -categories. We also note that the left downward arrow is an equivalence of  $\infty$ -categories because the fibration has contractible fibers. It is then straightforward to see that all the unlabeled arrows in the diagram are maps of symmetric monoidal  $\infty$ -categories (not just maps of  $\infty$ -operads).

The desired symmetric monoidal functor is the composition of an inverse to the left vertical arrow with the top horizontal arrows.  $\square$

**Lemma 13.10.** There exists a symmetric monoidal functor

$$(\mathbf{Liou}_{\text{str}})^{\otimes} \rightarrow \mathcal{T}\text{op}_{/BU}^{\otimes} \tag{13.1}$$

which sends a Liouville sector  $X$  to some map  $(X \rightarrow BU)$  classifying the stable tangent bundle of  $X$  (equipped with a compatible almost-complex structure).

*Proof.* Combine Lemma 13.8 with Lemma 13.9, composing with the natural functor  $(\mathrm{VecBun}_{\mathbb{C}})^{\oplus} \rightarrow (\mathrm{VecBun}_{\mathbb{C}}^{\diamond})^{\oplus}$  along the way.  $\square$

### 13.3 Stabilized Liouville sectors with tangential structures

The functor (13.1) sends sectorial equivalences to homotopy equivalences, so one obtains a symmetric monoidal map from the localization (Proposition 11.5):

$$(\mathrm{Liou}_{\mathrm{str}}[\mathfrak{eq}^{-1}])^{\otimes} \rightarrow (\mathcal{T}\mathrm{op}_{/BU})^{\otimes}.$$

Because the object  $T^*[0, 1]$  is sent to an invertible object in the target<sup>36</sup>, we obtain a symmetric monoidal functor

$$(\mathcal{L}\mathrm{iou}^{\diamond})^{\otimes} := (\mathrm{Liou}_{\mathrm{str}}[\mathfrak{eq}^{-1}]^{\diamond})^{\otimes} \rightarrow \mathcal{T}\mathrm{op}_{/BU}^{\otimes}.$$

(This follows from Theorem D, or from Proposition 11.6. See also Notation 1.22.)

At this moment, we may contemplate tangential structures as follows:

**Construction 13.11**  $(\mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^{\diamond})$ . Fix  $E_{\infty}$ -spaces  $B$  and  $B'$ , along with  $E_{\infty}$  maps  $BU \rightarrow B \leftarrow B'$ .

We define the  $\infty$ -category of stable Liouville sectors equipped with a reduction of induced  $\Omega B$ -structures to  $\Omega B'$ -structures as the following pullback  $\infty$ -category:

$$\begin{array}{ccc} \mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^{\diamond} & \xrightarrow{\quad\quad\quad} & \mathcal{T}\mathrm{op}_{/(B' \rightarrow B)} \\ \downarrow & & \downarrow \\ \mathcal{L}\mathrm{iou}^{\diamond} & \xrightarrow{\quad\quad\quad} \mathcal{T}\mathrm{op}_{/BU} \xrightarrow{\quad\quad\quad} & \mathcal{T}\mathrm{op}_{/B}. \end{array}$$

(The bottom horizontal arrow is depicted as a composition for explicitness.) Here,  $\mathcal{T}\mathrm{op}_{/(B' \rightarrow B)}$  is the  $\infty$ -category whose objects are topological spaces  $X$  equipped with (i) Maps to  $B'$  and  $B$ , and (ii) a homotopy between  $X \rightarrow B' \rightarrow B$  and  $X \rightarrow B$ .

By remembering the  $E_{\infty}$ -structure of the maps  $BU \rightarrow B$  and  $B' \rightarrow B$ , one may promote this pullback diagram to be a pullback of symmetric monoidal  $\infty$ -categories, thereby endowing  $\mathcal{L}\mathrm{iou}_{/(B' \rightarrow B)}^{\diamond}$  with a symmetric monoidal structure. (See Recollection 13.2.)

**Example 13.12** (Reductions to  $\Omega B'$ ). Let  $B = BU$ . Fix an  $E_{\infty}$  map  $B' \rightarrow BU$  from some  $E_{\infty}$  space  $B'$ . Then one can contemplate stabilized Liouville sectors with a reduction of the stable complex structure of its tangent bundle to some  $\Omega B'$ -structure. The associated  $\infty$ -category of such sectors is the pullback of  $\infty$ -categories

$$\begin{array}{ccc} \mathcal{L}\mathrm{iou}_{/(B' \rightarrow BU)}^{\diamond} & \xrightarrow{\quad\quad\quad} & \mathcal{T}\mathrm{op}_{/(B' \rightarrow BU)} \\ \downarrow & & \downarrow \\ \mathcal{L}\mathrm{iou}^{\diamond} & \xrightarrow{\quad\quad\quad} & \mathcal{T}\mathrm{op}_{/BU}. \end{array}$$

Informally,  $(\mathcal{L}\mathrm{iou}^{\diamond})_{/(B' \rightarrow BU)}$  is the  $\infty$ -category of stabilized Liouville sectors equipped with a reduction of the stable complex structure bundle's structure group to  $\Omega B'$ . Taking  $B' = *$  to be the trivial/initial  $E_{\infty}$ -space, this is the data of a stable trivialization of the complex tangent bundle. One could also choose  $B' = BG$  for some infinite-loop space  $G$  (e.g.,  $GL(\infty)$ ) in which case objects of  $\mathcal{L}\mathrm{iou}_{/(B' \rightarrow BU)}^{\diamond}$  are stabilized Liouville sectors with a reduction of the stable complex structure of its tangent bundle to a  $G$ -structure.

<sup>36</sup>In fact, the image of  $T^*[0, 1]$  under this functor is (equivalent to) the unit—it is a/the trivial stable vector bundle on  $* \simeq T^*[0, 1]$ .

**Example 13.13** (Trivializations of induced structures). This time, fix an  $E_\infty$  map  $BU \rightarrow B$  and let  $B' = *$ . One may then contemplate the following pullback diagram, where the bottom horizontal arrow is depicted as a composition for explicitness:

$$\begin{array}{ccc} \mathcal{L}\mathrm{iou}^\diamond_{/(* \rightarrow B)} & \xrightarrow{\quad} & \mathcal{T}\mathrm{op}_{/(* \rightarrow B)} \\ \downarrow & & \downarrow \\ \mathcal{L}\mathrm{iou}^\diamond & \xrightarrow{\quad} \mathcal{T}\mathrm{op}_{/BU} \longrightarrow \mathcal{T}\mathrm{op}_{/B} & \end{array}$$

An object of  $(\mathcal{L}\mathrm{iou}^\diamond)_{* \rightarrow B}$  is a stable Liouville sector  $X$  equipped with a null-homotopy of the map  $X \rightarrow B$ . For example, when  $BU \rightarrow BU/BO \simeq B(U/O) =: B$  is chosen to be the map classifying the stable Lagrangian Grassmanian bundle, such a null-homotopy exhibits a stable trivialization of this bundle.

**Example 13.14.** There is another example relevant in the conjectural construction of (wrapped) Floer theory with various coefficient ring spectra.

For example, choose a commutative coefficient ring spectrum  $R$ . Then  $B\mathrm{Pic}(R)$ —the classifying space of rank-one  $R$ -gerbes—inherits an action of  $B\mathrm{Pic}(\mathbb{S})$  by virtue of  $\mathbb{S}$  being the initial ring. One can take

- $BU \rightarrow B = B^2\mathrm{Pic}(\mathbb{S})$  to be a two-fold delooping of the complex  $J$  homomorphism, and
- $B' = B\mathrm{Pic}(R)/B\mathrm{Pic}(\mathbb{S}) \rightarrow B^2\mathrm{Pic}(\mathbb{S})$  to be the map induced by the  $B\mathrm{Pic}(\mathbb{S})$ -action on  $B\mathrm{Pic}(R)$ .

See, for example, Section 1.3 of [19].

**Remark 13.15.** For any choice of  $E_\infty$  maps  $BU \rightarrow B \leftarrow B'$ , the map  $\mathcal{L}\mathrm{iou}^\diamond_{/(B' \rightarrow B)} \rightarrow \mathcal{L}\mathrm{iou}^\diamond$  is a right fibration. To put this remark in context, recall that a right fibration is a map that classifies contravariant functor from the base to the  $\infty$ -category of spaces—see [17, Section 2.1]. In our case, the contravariant functor may be informally describe as sending an object  $X \in \mathcal{L}\mathrm{iou}^\diamond$  to the space of ways to reduce the stable  $\Omega B$  bundle to have structure group  $\Omega B'$ .

### 13.4 Localization with tangential structures (Proof of Theorem E)

There is also the obvious strict analogue of  $\mathcal{L}\mathrm{iou}^\diamond_{/(B' \rightarrow B)}$ :

**Construction 13.16** ( $(\mathrm{Liou}^\diamond_{\mathrm{str}})_{/(B' \rightarrow B)}$ ). Fix  $E_\infty$ -spaces  $B$  and  $B'$ , along with  $E_\infty$  maps  $BU \rightarrow B \leftarrow B'$ . We define the following pullback  $\infty$ -category:

$$\begin{array}{ccc} (\mathrm{Liou}^\diamond_{\mathrm{str}})_{/(B' \rightarrow B)} & \xrightarrow{\quad} & \mathcal{T}\mathrm{op}_{/(B' \rightarrow B)} \\ \downarrow & & \downarrow \\ \mathrm{Liou}^\diamond_{\mathrm{str}} & \xrightarrow{\quad} \mathcal{T}\mathrm{op}_{/BU} \longrightarrow \mathcal{T}\mathrm{op}_{/B} & \end{array}$$

(The bottom horizontal arrow is depicted as a composition for explicitness.)

Note that  $(\mathrm{Liou}^\diamond_{\mathrm{str}})_{/(B' \rightarrow B)}$  is typically not a category (in the usual sense), but is an  $\infty$ -category. Informally, this  $\infty$ -category only detects (stabilized) strict sectorial embeddings and no topology of the space of sectorial embeddings, but detects the homotopy type of lifts of  $B$ -structures to  $B'$ .

**Remark 13.17** (Symmetric monoidal structure). As in Construction 13.11, one may promote this pullback diagram to be a pullback of symmetric monoidal  $\infty$ -categories, endowing  $(\mathbf{Liou}_{\text{str}}^\diamond)_{/(B' \rightarrow B)}$  with a symmetric monoidal structure.

By the universal property of pullbacks, the functor  $\mathbf{Liou}_{\text{str}}^\diamond \rightarrow \mathcal{L}\mathbf{iou}^\diamond$  (which factors the map to  $\mathcal{T}\mathbf{op}_{/BU}$ ) induces a functor

$$(\mathbf{Liou}_{\text{str}}^\diamond)_{/(B' \rightarrow B)} \rightarrow \mathcal{L}\mathbf{iou}_{/(B' \rightarrow B)}^\diamond. \quad (13.2)$$

**Theorem 13.18** (Theorem E). The map (13.2) is a localization. In fact, letting  $\mathfrak{eq}_{/B' \rightarrow B} \subset (\mathbf{Liou}_{\text{str}}^\diamond)_{/(B' \rightarrow B)}$  be the pullback of  $\mathfrak{eq} \subset \mathbf{Liou}_{\text{str}}^\diamond$ , the induced map

$$(\mathbf{Liou}_{\text{str}}^\diamond)_{/(B' \rightarrow B)}[\mathfrak{eq}_{/B' \rightarrow B}^{-1}] \rightarrow \mathcal{L}\mathbf{iou}_{/(B' \rightarrow B)}^\diamond$$

is an equivalence of  $\infty$ -categories.

*Proof.* This follows more generally from the following fact: If  $\mathcal{C} \rightarrow \mathcal{D}$  is a localization, and  $\mathcal{D}' \rightarrow \mathcal{D}$  is a right fibration, then the pullback induces a localization  $\mathcal{C}' \rightarrow \mathcal{D}'$ .

Here is a proof. The formation of colimits of  $\infty$ -categories commutes with pulling back along right fibrations. So the question immediately reduces to the case where  $\mathcal{C}$  consists of a single morphism and  $\mathcal{D}$  is obtained by inverting it; this can be modeled by the projection  $\Delta^1 \rightarrow \Delta^0$ . In this case the right fibration  $\mathcal{D}' \rightarrow \mathcal{D}$  is constant, so the result is trivial.  $\square$

## References

- [1] Mohammed Abouzaid. On the wrapped Fukaya category and based loops. *J. Symplectic Geom.*, 10(1):27–79, 2012.
- [2] Mohammed Abouzaid and Paul Seidel. Altering symplectic manifolds by homologous recombination. arXiv:1007.3281, 2010.
- [3] Mohammed Abouzaid and Paul Seidel. An open string analogue of Viterbo functoriality. *Geom. Topol.*, 14(2):627–718, 2010.
- [4] David Ayala, John Francis, and Hiro Lee Tanaka. Factorization homology of stratified spaces. *Sel. Math., New Ser.*, 23(1):293–362, 2017.
- [5] David Ayala, John Francis, and Hiro Lee Tanaka. Local structures on stratified spaces. *Adv. Math.*, 307:903–1028, 2017.
- [6] Kai Cieliebak and Yakov Eliashberg. *From Stein to Weinstein and back – Symplectic geometry of affine complex manifolds* –, volume 59 of *AMS Colloquium Publications*. American Mathematical Society, Providence, RI, 2012.
- [7] W. G. Dwyer and D. M. Kan. Function complexes in homotopical algebra. *Topology*, 19(4):427–440, 1980.
- [8] Yakov Eliashberg. Weinstein manifolds revisited. In *Modern geometry. A celebration of the work of Simon Donaldson*, pages 59–82. Providence, RI: American Mathematical Society (AMS), 2018.
- [9] Sheel Gantra, John Pardon, and Vivek Shende. Covariantly functorial wrapped Floer theory on Liouville sectors. arXiv:1706.03152, 2017.
- [10] Sheel Gantra, John Pardon, and Vivek Shende. Sectorial descent for wrapped Fukaya categories. arXiv:1809.03427v2, 2018.
- [11] Paul G. Goerss and John F. Jardine. *Simplicial homotopy theory*. Modern Birkhäuser Classics. Birkhäuser Verlag, Basel, 2009. Reprint of the 1999 edition [MR1711612].
- [12] Vladimir Hinich. Dwyer-Kan localization revisited. *Homology Homotopy Appl.*, 18(1):27–48, 2016.
- [13] André Joyal and Myles Tierney. Quasi-categories vs Segal spaces. In *Categories in algebra, geometry and mathematical physics*, volume 431 of *Contemp. Math.*, pages 277–326. Amer. Math. Soc., Providence, RI, 2007.
- [14] Daniel M. Kan. On c. s. s. complexes. *Am. J. Math.*, 79:449–476, 1957.
- [15] Hiroshi Kihara. Smooth Homotopy of Infinite-Dimensional  $C^\infty$ -Manifolds. Available at <https://arxiv.org/abs/2002.03618>, 2020.
- [16] Oleg Lazarev, Zachary Sylvan, and Hiro Lee Tanaka. Localization and flexibilization in symplectic geometry. arxiv:2109.06069, 2021.
- [17] Jacob Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.

- [18] Jacob Lurie. Higher algebra. Available at <http://www.math.harvard.edu/~lurie/papers/higheralgebra.pdf>, 2012.
- [19] Jacob Lurie. Rotation Invariance in Algebraic K-Theory. Available at <https://www.math.ias.edu/~lurie/papers/Waldhaus.pdf>, 2015.
- [20] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [21] J. Peter May.  *$E_\infty$  ring spaces and  $E_\infty$  ring spectra. With contributions by Frank Quinn, Nigel Ray, and Jorgen Tornehave*, volume 577. Springer, Cham, 1977.
- [22] Dusa McDuff and Dietmar Salamon. *Introduction to symplectic topology*. Oxford: Clarendon Press, 1995.
- [23] John W. Milnor and James D. Stasheff. *Characteristic classes*, volume 76. Princeton University Press, Princeton, NJ, 1974.
- [24] David Nadler and Vivek Shende. Sheaf quantization in Weinstein symplectic manifolds. arXiv:2007.10154, 2021.
- [25] David Nadler and Hiro Lee Tanaka. A stable  $\infty$ -category of Lagrangian cobordisms. arXiv:1109.4835, 2011.
- [26] Yong-Geun Oh and Hiro Lee Tanaka. Continuous and coherent actions on wrapped Fukaya categories. arXiv:1911.00349, 2019.
- [27] Yong-Geun Oh and Hiro Lee Tanaka. Smooth approximation for classifying spaces of diffeomorphism groups. arxiv:2003.06033, 2020.
- [28] Marco Robalo. Noncommutative Motives I: A Universal Characterization of the Motivic Stable Homotopy Theory of Schemes, 2013.
- [29] P. Seidel.  $\pi_1$  of symplectic automorphism groups and invertibles in quantum homology rings. *Geom. Funct. Anal.*, 7(6):1046–1095, 1997.
- [30] Wolfgang Steimle. Degeneracies in quasi-categories. *J. Homotopy Relat. Struct.*, 13(4):703–714, 2018.
- [31] Zachary Sylvan. On partially wrapped Fukaya categories. *J. Topol.*, 12(2):372–441, 2019.
- [32] Hiro Lee Tanaka. The Fukaya category pairs with Lagrangian cobordisms. arxiv:1607.04976, 2016.
- [33] Hiro Lee Tanaka. Functors (between  $\infty$ -categories) that aren’t strictly unital. *J. Homotopy Relat. Struct.*, 13(2):273–286, 2018.
- [34] Hiro Lee Tanaka. Cyclic structures and broken cycles. arXiv:1907.03301, 2019.
- [35] Bertrand Toën and Gabriele Vezzosi. *Homotopical algebraic geometry. II: Geometric stacks and applications*, volume 902. Providence, RI: American Mathematical Society (AMS), 2008.
- [36] Vladimir Voevodsky.  $A^1$ -homotopy theory. *Doc. Math.*, Extra Vol.:579–604, 1998.