

Double Proton Transfer in Hydrated Formic Acid Dimer: Interplay of Spatial Symmetry and Solvent-Generated Force on Reactivity

Kai Töpfer, Silvan Käser, and Markus Meuwly*

*Department of Chemistry, University of Basel, Klingelbergstrasse 80, CH-4056 Basel,
Switzerland.*

E-mail: m.meuwly@unibas.ch

Abstract

The double proton transfer (DPT) reaction in hydrated formic acid dimer (FAD) is investigated at molecular-level detail. For this, a global and reactive machine learned (ML) potential energy surface (PES) is developed to run extensive (more than 100 ns) mixed ML/MM molecular dynamics (MD) simulations in explicit molecular mechanics (MM) solvent at MP2-quality for the solute. Simulations with fixed - as in a conventional empirical force field - and conformationally fluctuating - as available from the ML-based PES - charge models for FAD shows significant impact on the competition between DPT and dissociation of FAD into two formic acid monomers. With increasing temperature the barrier height for DPT in solution changes by about 10 % (~ 1 kcal/mol) between 300 K and 600 K. The rate for DPT is largest, $\sim 1 \text{ ns}^{-1}$, at 350 K and decreases for higher temperatures due to destabilisation and increased probability for dissociation of FAD. The water solvent is found to promote the first proton transfer by exerting a favourable solvent-induced Coulomb force along the O-H \cdots O hydrogen bond whereas the second proton transfer is significantly controlled by the O-O separation and other conformational degrees of freedom. Double proton transfer in hydrated FAD is found to involve a subtle interplay and balance between structural and electrostatic factors.

Introduction

Understanding the molecular details of chemical reactions in solution is one of the challenges in physical and general chemistry at large.¹⁻¹⁰ These processes are at the center of interest of organic, inorganic and biological chemistry as the environment in which they take place can determine reaction products, their rates and propensities. However, the direct and atomistically resolved experimental characterization by which the solvent influences reactions is challenging because simultaneous control over temporal (i.e. “rates”) and spatial (“collocation” and local solvation) aspects of the process along the progression coordinate is rarely possible. On the other hand, computer simulations have matured to a degree that

allows the study of realistic processes at a sufficiently high level of theory including the environment.^{11,12}

More specifically, polar solvents such as water, can stabilize the transitions state (TS) of the reactant.^{4,13,14} For the Claisen rearrangement the origin of the catalytic effect of the water solvent is the differentially stronger hydrogen bonding to the TS structure (allyl phenyl ether) compared with that to the reactant structure because the TS is more polar which lowers the reaction barrier.^{15,16} Further stabilization of the TS and lowering of the reaction barrier by the water solvent can be affected by solvent polarizability and dipole-dipole interaction between solute in its TS structure and the solvent.¹⁶ The explicit description of solvent effects by full *ab initio* MD simulations for reactive systems in solution is computationally very demanding and can be typically only followed on the picosecond time scale.^{17,18} To avoid this, the (reactive) solute can be treated quantum mechanically (QM - ranging from tight-binding to post-Hartree-Fock methods) whereas the remaining degrees of freedom are described by an empirical force field.¹⁹⁻²¹ Alternatively, reactive force fields have been developed which allow to follow chemical reactions in solution on extended time scales.^{1,22,23} More recently, another possibility has emerged by developing a machine learned (ML) representation for the solute that can be used together with an implicit^{24,25} or explicit description of the environment at an empirical level. The present work develops such a scheme and applies it to formic acid dimer in solution.

Formic acid dimer (FAD) has been subject of several computational²⁶⁻⁴¹ and experimental studies.^{36,42-48} In the gas phase, formic acid exists as hydrogen bonded dimers⁴⁹ making it a prototype for complexes with hydrogen bonds such as enzymes or DNA base pairs.⁵⁰ Such conformations allow double proton transfer (DPT) to occur which leads to broadening of the line shape for the O-H frequency in IR-spectra.³⁶ The dissociation energy of FAD into two formic acid monomers (FAMs) has been determined from spectroscopy and statistical thermodynamics to be $D_0 = -59.5(5)$ kJ/mol (~ -14.22 kcal/mol)⁴⁷ which compares with

-14.23 ± 0.08 kcal/mol from recent computations.⁴¹

Another quantity that has been accurately measured experimentally is the tunneling splitting for which a recent value of 331.2 MHz is available from microwave spectroscopy.⁵¹ From this, a barrier for DPT of 2559 cm^{-1} (30.6 kJ/mol or 7.3 kcal/mol) from analysis of a 3D model was determined.⁵¹ The tunneling splitting from the microwave data compares with values of 0.011 cm^{-1} (340.8 MHz)⁵² and 0.016 cm^{-1} (473.7 MHz)^{53,54} from infrared spectroscopy. Computationally, a high-quality (CCSD(T)-F12a/haTZ), full-dimensional PES for FAD has been determined and represented as permutationally invariant polynomials which features a DPT barrier of 2853 cm^{-1} (8.16 kcal/mol).³⁵ The computed splitting on this PES from using a ring-polymer instanton approach was 0.014 cm^{-1} (420 MHz).³⁷

Given its thorough characterization in the gas phase, FAD is also an interesting system to study reactions in solution. In particular the balance between DPT and the dissociation of FAD into two monomers provides an attractive aspect of the system to be explored also as a proxy for DNA base-pair dimers.^{55,56} The homologous series of carboxylic acids (formic, acetic, propionic, butyric) in water have also served as model systems for studying the balance and role of hydrogen bonding, hydrophobicity and entropy changes which is of particular relevance in the context of protein folding and stability.⁵⁷

For FAD in pure water detailed experimental results are scarce. Early work reported that dimerization is most relevant for the low-concentration range for all carboxylic acids investigated, including formic acid.⁵⁸ This was subsequently challenged and measurements in 3 M NaCl by potentiometric titrations were interpreted as leading to singly H-bonded, extended formic acid oligomers.⁵⁹ The hydration of formic acid in pure water was also investigated with neutron diffraction experiments for higher formic acid concentrations.⁶⁰ At all conditions studied, a pronounced peak for the $\text{O} \cdots \text{O}$ separation with a maximum at 2.7 \AA was reported

which, together with a peak for the $\text{OH}\cdots\text{O}$ angle at 180° , supports formation of singly H-bonded or cyclic FAD. It is of interest to note that the extended structures proposed by Scheraga are incompatible with the neutron scattering results.⁶⁰ Surface scattering experiments found that FAD colliding with the liquid water surface almost completely inserts in its cyclic form into bulk water even at low collision energies.⁶¹ This indicates that H-bond breaking in FAD is not essential for uptake. Whether or not and for how long FAD remains in its cyclic form inside water was, however, not determined in these studies.⁶¹ Accompanying *ab initio* MD simulations reported that the cyclic dimer has fewer FAD-water hydrogen bonds (i.e. is “more hydrophobic”) than the branched, singly H-bonded FAD isomer.⁶² Finally, a study considering clathrate formation involving formic acid and water interpreted the measured infrared spectra as partially originating from water-mediated cyclic formic acid dimers in which the two formic acid monomers are H-bonded by a bridging water molecule.⁶³ An independent MD simulation for FAD in water reported⁵⁷ a negligible stabilization free energy for cyclic FAD of 0.6 kcal/mol which is surprising given the substantial gas phase stability of FAD of -16.8 kcal/mol.^{33,41} Very recent Raman spectroscopy measurements reported formation of both, singly and cyclic H-bonded FADs.⁶⁴ Taken together, the available studies suggest that FAD on and in pure water can exist as an equilibrium between cyclic and singly H-bonded dimers. Therefore, the present work will focus on the dynamics of cyclic FAD in pure water.

The aim of the present work is to characterize DPT dynamics in FAD in aqueous solution. First, the methods and the ML/MM embedding are described. Next, the quality of the PES is reported and the structural dynamics, free energies, and rates together with conformational coordinates relevant for DPT of FAD in solution are discussed. This is followed by an analysis of the solvent distributions for reactant and transition state structures, and the effect of the solvent-generated electrical field. Finally, the results are discussed in a broader context.

Methods

This section describes the generation of the potential energy surface for the simulation of FAD in water and the molecular dynamics (MD) simulations performed within the Atomic Simulation Environment (ASE).⁶⁵ In the following the potential energy of FAD is represented as a Neural Network (NN),⁶⁶ referred to as “ML” or “PhysNet” whereas solvent water is described with the TIP3P model (MM).⁶⁷ First, the reference data generation, the training of the NN-based PES and the embedding of the ML (FAD) part in the MM environment (water) is described. This is followed by a description of the MD simulations and their analysis.

Potential Energy Surface

All electronic structure calculations were carried out at the MP2/aug-cc-pVTZ level of theory using the MOLPRO software package.⁶⁸ The reference data included structures, energies, forces, and dipole moments for FAM (5000 structures), FAD (47069), and a number (6000) of related fragments (“amons”)⁶⁹ including H₂, CO, H₂O, CH₄, CH₂O and CH₃OH. Structures were generated from MD simulations, carried out at the PM7 level of theory using ASE. The “amons” were only used for training the NN and help in generalizing the ML model. Additional FAD structures were generated from simulations in solution (akin to adaptive sampling) by using PhysNet trained only on gas-phase structures.^{41,70} The energies of the FAD structures cover a range up to ~ 100 kcal/mol above the global minimum compared with the TS for DPT which is ~ 8 kcal/mol above the global minimum. For the final training the 58069 reference structures were split into training (49300), validation (5800) and testing (2969) sets. The parameters of PhysNet were fitted to reproduce *ab initio* energies, forces and dipole moments of the training set.⁶⁶ Including FAD conformations exhibiting two, one or no H-bond in the training set has been shown essential to ensure the correct propensity towards the global minimum conformation of FAD with two formed H-bonds.

PhysNet computes the total potential energy E for given coordinates \vec{x} and nuclear charges \mathbf{Z} of N atoms from atomic energies E_i and pairwise electrostatic and dispersion interactions according to

$$E = \sum_{i=1}^N E_i + k_e \sum_{i=1}^N \sum_{j>i}^N \frac{q_i q_j}{r_{ij}} + E_{D3}. \quad (1)$$

The prediction of the atomic energies E_i and charges q_i is based on feature vectors that encode the local chemical environment of each atom $i \in N$ to all atoms j within a cutoff radius that was $r_{\text{cut}} = 10 \text{ \AA}$ in the present work.⁷¹ For charge conservation, the partial charges q_i are scaled to the correct total charge of the system and used for the calculation of the electrostatic interaction energy where k_e is the Coulomb constant and r_{ij} is the distance between atoms i and j . For small distances the electrostatic energy is damped to avoid instabilities due to the singularity at $r_{ij} = 0$.⁶⁶ The initial parameters for the dispersion correction E_{D3} are the standard values recommended for the revPBE level of theory.⁷² Since the atomic features, electrostatic and dispersion interaction depend only on pairwise distances and are combined by summation, the PES is invariant to translation, rotation and permutation of equivalent atoms. The forces \vec{F} with respect to the atomic coordinates \vec{x} are computed by reverse mode automatic differentiation provided by Tensorflow.^{73,74} For further details, the reader is referred to Ref. 66.

For the water solvent the TIP3P model⁶⁷ as implemented in the GPAW program package⁷⁵ is used with a cutoff range of 10 \AA due to its improved performance compared to that of ASE. The interaction between formic acid and the water solvent includes van-der-Waals and electrostatic terms which were treated with a cutoff at 14 \AA or half the simulation box edge length at most. The van-der-Waals interactions are calculated using the Lennard-Jones-Potential and the parameters from CGenFF⁷⁶ for formic acid.

The electrostatic interactions between the TIP3P atomic charges of water and the NN-predicted atomic charges of FAD are computed within PhysNet. This allows exploitation of

reverse mode automatic differentiation to calculate the necessary derivatives of the fluctuating atomic charges from the atomic coordinates and ensures energy conservation. For the van-der-Waals and electrostatic interactions between the “high level” treatment (here PhysNet) and MM atoms a switch function $s(r)$

$$s(r) = \begin{cases} 1 & \text{for } r \leq r_1 \\ 1 - 6y^5 + 15y^4 - 10y^3 & \text{for } r_1 < r < r_2 \\ 0 & \text{for } r \geq r_2 \end{cases} \quad (2)$$

$$y = \frac{r - r_1}{r_2 - r_1} \quad (3)$$

was used to set the interactions zero in the range from $r_1 = 13 \text{ \AA}$ to $r_2 = 14 \text{ \AA}$, where r is the separation between ML and MM atoms, respectively. Additionally, the electrostatic Coulomb potential V_C is shifted to zero at the cutoff distance of $r_{\text{cut}} = 14 \text{ \AA}$ following the shifted force method to prevent the inconsistent gradient at the cutoff point.⁷⁷ While the van-der-Waals interaction between both species become negligible at that cutoff distances, this is not the case for the Coulomb forces between two charges q_i and q_j of atoms i from the ML part and j from the MM part separated by r_{ij} .

$$V_C = \sum_i^{N_{\text{ML}}} \sum_j^{N_{\text{MM}}} V_{C,ij}^{\text{shifted}}(r_{ij}) \quad (4)$$

$$V_{C,ij}^{\text{shifted}}(r_{ij}) = s(r_{ij}) \cdot \left(V_{C,ij}(r_{ij}) - V_{C,ij}(r_{\text{cut}}) - \frac{\partial V_{C,ij}}{\partial r} \Big|_{r_{\text{cut}}} \cdot (r_{ij} - r_{\text{cut}}) \right) \quad (5)$$

$$V_{C,ij}(r_{ij}) = \frac{1}{4\pi\epsilon} \frac{q_i \cdot q_j}{r_{ij}} \quad (6)$$

An ASE calculator class for the PhysNet model was written as interface to the ASE program package. Further extensions are implemented to enable constant atomic charges in formic acid only for the calculation of the electrostatic potential between the ML and MM part

of the simulation setup. In addition, custom energy functions were added to the potential energy to perform biased simulations.⁷⁸

Molecular Dynamics Simulations

All MD simulations were performed using Python packages of a modified version of ASE (3.20.1).⁶⁵ Simulations are initialized with FAD in its minimum conformation in a cubic simulation box with edge length 28.036 Å containing 729 water molecules which corresponds to a density of 0.997 g/cm³. The density is not adjusted for simulations at different temperatures. All simulations were run with periodic boundary conditions. To maintain the solute near the center of the simulation box a center-of-mass harmonic constraint with a force constant of 0.23 kcal/mol/Å² was applied on formic acid molecules when moving further than 10 Å from the center of the simulation box. RATTLE-type holonomic constraints⁷⁹ as implemented in the GPAW program package⁸⁰ were applied to the oxygen-hydrogen and hydrogen-hydrogen distances of the water molecules. The simulations are carried out in the *NVT* ensemble using the Langevin propagator with a time step of $\Delta t = 0.2$ fs and a friction coefficient of 10 ps⁻¹.

First, the structure of the system was minimized to release strain. A heating run is performed for 10 ps to reach the respective target temperatures of 300 K, 350 K, 400 K, 450 K, 500 K and 600 K. As no rigorous implementation of a barostat is available in ASE, simulations at constant volume but increased temperature were carried out to qualitatively assess the influence of increased pressure on the DPT dynamics. The system equilibrates within 5 ps during the heating runs. Equilibration is followed by production simulations of 100 ps each, accumulating to a total of 16 ns simulation time at each temperature. Positions and momenta are stored (1) every 10 fs for the complete trajectory and (2) every 1 fs in a time window of $\Delta t = [-20, +50]$ fs around every attempt of a reaction registered during the simulation (see Analysis).

The implementation is validated by examining energy conservation for a 100 ps *NVE* MD simulation at 600 K. The total energies at each time step are within a standard deviation of $6.1 \cdot 10^{-3}$ kcal/mol from the average total energy with a maximum deviation of $\pm 25.2 \cdot 10^{-3}$ kcal/mol and no drift is found. This confirms that the forces are correctly implemented.

To determine the impact of the fluctuating atomic charges available from PhysNet on the dynamics and energetics, MD simulations were also performed by fixing the atomic charges to those of the TS structure of FAD to retain the symmetry of the system before and after PT. These fixed atomic charges were used for the solute-solvent interaction only. For the internal interaction of the ML part (FAD) fluctuating charges were retained in order to correctly represent the PES for the isolated solute. Finally, as FAD was found to be largely hydrophobic from earlier simulations^{57,62} and to probe the effect of the electrostatic interactions between solute and solvent on DPT, simulations were also carried out with zeroed charges on all atoms of the solute.

Analysis

Proton transfer (PT) is a transient process. To characterize a PT event it is often useful to use a geometric criterion.⁸¹ Here, the criterion was the distance between the hydrogen and the oxygen atoms O_A and O_B of the respective H-bond. A PT in one H-bond was identified by a sign change of the progression (or reaction) coordinate $\xi = d(H - O_A) - d(H - O_B)$. Further, the oxygen atom bonded to the hydrogen atom is the donor oxygen (O_{don}) and the second oxygen atom in the H-bond is the acceptor oxygen (O_{acc}), see Figure 1. Successful and attempted DPT always involve two consecutive PTs, see Figure S1. For “successful” DPT the two PTs include a “first” and a “second” hydrogen atom H_A and H_B in both H-bonds,

respectively, whereas for an “attempted” DPT both PTs involve only hydrogen atom H_A . DPT is considered to be successful when a sign change in both progression coordinates ξ_1 and ξ_2 occurs. Attempted - as opposed to successful - DPT events are those for which only one forth and back PT occurs along the first H-bond without PT along the second H-bond.

In addition to unbiased simulations, biased simulations were carried out at different temperatures for FAD in the gas phase and in solution. For this, an artificial harmonic potential $V_{\text{bias}}(\xi_i) = \frac{1}{2} \sum_{i=1}^2 (k_{\text{bias}} \cdot (\xi_i - \xi_{i,e})^2)$ involving both progression coordinates ξ_1 and ξ_2 was added to the total potential energy. For FAD in solution the force constant was $k_{\text{bias}} = 16.1$ kcal/mol/Å² and $5 \cdot 10^6$ snapshots were recorded from 1 ns trajectories at each temperature. Both constraining harmonic potentials are centered at the TS coordinate, $\xi_{1,\text{TS}} = \xi_{2,\text{TS}} = 0$, and the magnitude of the force constant was chosen to lift the potential of the equilibrium conformation ($\xi_{\text{eq}} = \pm 0.66$ Å) of FAD roughly to the level of the TS conformation. Additional tests were run with lower values of k_{bias} . The resulting potentials of mean force (PMFs) were found to be insensitive of this choice, see Figure S2.

Results

Quality of the Potential Energy Surface

Figure 1 shows the quality of the ML PES by comparing reference MP2 energies with those predicted from the trained PhysNet model for formic acid dimer for 2969 test structures. The energies of the smaller fragments (“amons”) are outside the energy range of the graph but are included in the root mean squared error (RMSE) of 0.336 kcal/mol and the Pearson coefficient is $R^2 = 1 - 2 \cdot 10^{-6}$. The energy deviation between the MP2 energy and the PhysNet model prediction is smaller than $3 \cdot 10^{-3}$ kcal/mol for the minimum conformation of FAD and the TS for DPT.




Figure 1: Correlation between the MP2/aug-cc-pVTZ reference and the NN-predicted energies from PhysNet for the test set. The 2969 randomly chosen structures which were not used during training contain samples of FAM, FAD, and their “amons”. The RMSE between reference and NN-predicted energies is 0.336 kcal/mol with a Pearson coefficient of $R^2 = 1 - 2 \cdot 10^{-6}$. The red cross indicates the energy for the TS. A sketch of FAD with atom labels is shown in the upper left corner. The vector pointing from O_{don} towards O_{acc} , \vec{R}_{OO} , is also reported.

The MP2 method is a good compromise between speed and accuracy. For example, the barrier for DPT of 6.7 kcal/mol compares with a value of 7.2 kcal/mol from a morphed MMPT-MP2 force field that is consistent with infrared spectroscopy³⁶ and a barrier height of 7.3 kcal/mol from analysis of microwave spectroscopy data.⁵¹ Calculations at the CCSD(T) level of theory report somewhat higher barriers of 7.9 kcal/mol and 8.2 kcal/mol, depending on the basis set and additional F12 corrections used.^{35,36} Furthermore, the dissociation of the dimer into two

monomers is of interest. In the gas phase the PhysNet model yields $D_e = -16.8$ kcal/mol which is also the value from the reference MP2/aug-cc-pVTZ calculations. *Ab initio* calculations at the higher CCSD(T)/aug-cc-pVTZ level of theory find $D_e = -16.8$ kcal/mol^{33,41} and -16.0 kcal/mol at the basis set limit.³³

Next, the energetics of increasingly hydrated FAD is determined from the present ML/MM energy function, from the CGenFF⁷⁶ parametrization of FAD together with the TIP3P water model, and from electronic structure calculations. For this, 50 snapshots from a 2 ns ML/MD simulation with FAD in its dimeric structure were extracted. The 15 water molecules closest to the center of mass of FAD were retained for each snapshot. Electronic structure calculations for this analysis were carried out at the B3LYP+D3/aug-cc-pVDZ level of theory⁸²⁻⁸⁴ because MP2/aug-cc-pVTZ calculations for FAD surrounded by up to 15 water molecules are computationally too demanding. The total interaction energies for FAD-(H₂O)_{*n*} (*n* = 1, ..., 15) complexes were determined. Water molecules were retained in increasing order of their distance from FAD. The results are reported in Figures S3 and S4 and the analysis shows that the interaction energies with 15 water molecules from the MM/ML energy function differ on average by less than 1 kcal/mol from the reference B3LYP+D3 calculations whereas CGenFF underestimates the reference results by about 4 kcal/mol.

The 50 snapshots were also analyzed from retaining the 4, 8, and 12 closest water molecules and correlating the MM/ML and CGenFF interaction energies with those from the reference B3LYP+D3 calculations (Figures S5 to S7). Mean absolute errors between MM/ML and DFT reference calculations are 1.13, 2.06 and 2.85 kcal/mol compared with 2.42, 3.36 and 3.62 kcal/mol for CGenFF. It is also worth to point out that CGenFF finds a dimer stabilization energy of only 9 kcal/mol in the gas phase compared with high-level electronic structure calculations that yield -16.8 kcal/mol^{33,41} and -17.4 kcal/mol at the B3LYP+D3/aug-cc-pVDZ level which is -0.6 kcal/mol lower than for the MP2 and CCSD(T) levels of theory with

the aug-cc-pVTZ basis set. Overall, the comparison with the B3LYP+D3 results validates the quality of the ML/MM energy function whereas the CGenFF parametrization is found to considerably underestimate the stability of FAD.

Structural Dynamics in Solution

Figure 2 reports the propensity of FAD to exist as a doubly (blue circle) or singly (violet square) H-bonded dimer or as two separate monomers (red cross) in water. This is consistent with what has been inferred from experiments.^{60,61} For the equilibrium conformation the $O_{\text{don}}-O_{\text{acc}}$ distance is $\sim 2.66 \text{ \AA}$. An H-bond in FAD is considered “broken” if the $O_{\text{don}}-O_{\text{acc}}$ distance exceeds 4 \AA . MD simulations with fluctuating atomic charges on FAD show (Figure 2A) that the propensity for a doubly H-bonded FAD decreases from 97.0 % at 300 K to 0.4 % at 600 K. The probability for the singly H-bonded dimer increases from 0.8 % at 300 K to a maximum of 4.9 % at 400 K and decreases again to 0.7 % at 600 K. At the temperature with the highest DPT rate (350 K) both H-bonds are formed for 82.4 % of the propagation time.

In conventional force fields the partial charges are fixed⁷⁶ and do not change with conformation. PhysNet provides geometry-dependent charges, see Figure S8, which fluctuate by $\pm 0.05 e$ to $\pm 0.1 e$ around their mean. The average charges for the C, H_C and one of the oxygen atoms from PhysNet are similar to those in CGenFF whereas for the H_O and the second oxygen they differ by about $0.1 e$. It is also found that the charge distributions from PhysNet differ for cyclic and branched FAD. Given these differences it is of interest to determine the differences between the simulations analyzed with conformationally fluctuating charges and those with fixed partial charges on the FAD. The charges assigned to the atoms are those of the TS structure to ensure a symmetric atomic charge distribution before and after successful DPT. The distribution of formed H-bonds for FAD in water with fluctuating and fixed charges in Figure 2 shows that the propensity for H-bond formation significantly depends on fixing




Figure 2: DPT rates and distribution of the number of formed H-bonds in the simulation of FAD in water at different temperatures using fluctuating atomic charges (panel A) and fixed atomic charges (panel B). The green numbers atop the bars show the respective rate and absolute number of observed DPT events. The markers show the propensity of FAD to have two formed H-bonds (blue circle), to be partially dissociated by one (purple square) or completely separated into two formic acid monomers (red cross). The lines are to guide the eye.

the charges. With the fixed charge model the probability for the doubly H-bonded dimer to exist is larger at all temperatures from 300 K (98.7%) to 600 K (3.3%) compared with the fluctuating charge model (97.0% and 0.4% at 300 K and 600 K, respectively). In addition, the probability for singly H-bonded FAD and two separate monomers at 600 K differs with the fixed charge model with 6.4% and 90.3%, respectively, from the fluctuating charge model with 0.7% and 98.8%. Thus, a fixed charge model with charges from the TS structure predicts a higher propensity for FAD to exist in solution compared to the fluctuating charge model.

The higher propensity for dimer dissociation in solution using fluctuating rather than fixed charges correlates with the differences in the magnitudes of the atomic charges between the two charge models. From the equilibrium conformation of FAD towards separated FAMs the polarization along the atoms in the H-bond increases. The fluctuating charges along

the $\text{O}_{\text{don}}\text{-H}\cdots\text{O}_{\text{acc}}$ bond change from $[-0.39, +0.32, -0.43] e$ in the equilibrium dimer conformation to $[-0.42, +0.36, -0.45] e$ for separated monomers, respectively, which compares with $[-0.40, +0.30, -0.40] e$ in the fixed charge models with charges taken from the TS structure. Comparable amplitudes for charge transfer ($\delta q = 0.025 e$) have been found for the $\text{HO}_{\text{don}}\text{-H}\cdots\text{O}_{\text{acc}}\text{H}_2$ hydrogen bond in bulk water when the energy function was fit to match experimental frequencies and intensities of the infrared and Raman bands.⁸⁵ The larger absolute values of the atomic charges in the fluctuating charge model (differences of up to 20 % and 13 % for the hydrogen and oxygen atoms of the H-bonds relative to the charges in the TS structure, respectively) lead to stronger electrostatic interactions with the surrounding solvent. This also impacts on the conformational sampling and the reaction barrier height.

To evaluate the effect of the water solvent on dissociation, NVT simulations for FAD in the gas phase were performed using the same setup as for the simulations in solution. The presence of both H-bonds is a prerequisite for DPT to occur as the top of the barrier becomes inaccessible if even one H-bond is broken. Thus, dissociation of the FAD competes with DPT and therefore governs the DPT rate. In the gas phase it is found that - similar to the situation in solution - the propensity towards broken H-bonds increases for higher temperatures. At 300 K both H-bonds are formed for 99.4 % of the propagation time. This fraction decreases at higher temperatures: 27.3 % and 2.1 % at 500 K and 600 K, respectively.

Free Energies of Activation and Rates for DPT

Biased simulations were performed to investigate the free energy barrier for DPT. Figure 3 shows the 1-dimensional cuts along the constrained progression coordinate ξ . The 2-dimensional free energy surfaces $G(\xi_1, \xi_2)$ are given in Figure S9. The energies around the equilibrium $\xi_{\text{eq}} = \pm 0.66 \text{ \AA}$ are set to zero. Biased simulations are carried out for fluctuating and fixed charges. Between 300 K and 600 K the free energy barrier increases from

7.45 kcal/mol to 8.13 kcal/mol for simulations using the fluctuating charge model. With fixed charges the barriers change from 7.53 kcal/mol to 8.39 kcal/mol. Thus, as a function of temperature, the barrier increases by 0.68 kcal/mol with fluctuating charges, compared with 0.85 kcal/mol with fixed charges. In other words, the *change* of the barrier height as a function of temperature differs by 25 % from fluctuating to fixed charges on solution.

Simulations were also carried out in the *NVE* ensemble for FAD in the gas phase and yield a free energy barrier for DPT of 8.25 kcal/mol at a temperature of 310 K which was determined from the kinetic energy of all atoms in this biased simulation. Earlier classical *ab initio* metadynamics simulations with the same reaction coordinates as those used in the present work at the BLYP level with a plane wave basis for FAD in the gas phase reported a free energy barrier height of 4.7 kcal/mol at 300 K compared with the potential barrier height of 5.0 kcal/mol.³² This differs qualitatively from the present results for which the free energy barrier in the gas phase increases from the potential barrier of 6.7 kcal/mol.

To quantify the electrostatic contribution of the solute/solvent interactions to the free energy barrier, biased simulations of FAD in solution were performed with the atomic charges of FAD set to zero. At 300 K the free energy of activation is 7.46 kcal/mol and increases by +0.87 kcal/mol to 8.33 kcal/mol at 600 K. This compares with 7.45 and 7.53 kcal/mol with fluctuating and fixed charges at 300 K, respectively, which increase to 8.13 and 8.39 kcal/mol at 600 K. At 300 K the contribution of fluctuating electrostatics is thus -0.01 kcal/mol as opposed to $+0.07$ kcal/mol if the charges are fixed and a reduction by -0.20 kcal/mol and an increase of $+0.06$ kcal/mol at 600 K for fluctuating and fixed charges, respectively. This finding suggests that FAD behaves largely as a hydrophobic solute for which direct electrostatic interactions with the solvent are less relevant.

The T -dependence to the free energy ($-T\Delta S$) suggests that the change ΔS between the re-




Figure 3: Free energy profile for DPT in FAD with fluctuating (panel A) and fixed charges (panel B) for different temperatures in solution. The free energy curve follows the minimum energy path of the 2-dimensional free energy surface of antiparallel reactive coordinates $\xi_1 = -\xi_2$. The average temperature of FAD in solution in the NVT simulations are given in the legend and the standard deviation ranges from ± 6 K at 300 K to ± 13 K at 600 K. The PMF is symmetrized around $\xi_1 = \xi_2 = 0$ to increase data sampling as the artifacts at large absolute values of ξ result from insufficient sampling especially for biased simulations at low temperatures.

actant and the TS state is negative for all charge models used but the magnitude of ΔS differs. This difference is accommodated in the solvent ordering around the solute. Linear interpolation of the free energy barriers at the different temperatures yields an activation enthalpy and activation entropy of $[\Delta U, \Delta S] = [6.87 \text{ kcal/mol}, -2.1 \text{ cal/mol/K}]$ for fluctuating charges, $[6.62 \text{ kcal/mol}, -2.9 \text{ cal/mol/K}]$ for fixed charges, and $[6.70 \text{ kcal/mol}, -2.7 \text{ cal/mol/K}]$ for zero charges.

Because the constraints in the biased simulations favour FAD over two separated monomers (and the dimer has been found to dissociate at higher temperatures for free dynamics) it is mandatory to also run and analyze unbiased simulations that allow DPT to occur in solution. For this, an aggregate of 16 ns simulations was run and analyzed at each temperature. As the barrier for DPT is relatively high, DPT is a rare event. DPT rates in unbiased simulations are highest at 350 K for both fluctuating and fixed charge models, see Figures 2A and B. With the fluctuating charge model (Figure 2A) the rate is 0.9 ns^{-1} at 350 K and decreases to 0.4 ns^{-1} at 500 K, while the dimer is only present for 82.4 % and 6.4 % of the propagation time, respectively. Hence, the true rates are ~ 6.5 times higher at 500 K compared with those at 350 K. Within transition state theory energy barriers of $\sim 8 \text{ kcal/mol}$ from the biased simulations correspond to rates of $\sim 0.1 \text{ ns}^{-1}$ which is consistent with results from unbiased simulations shown in Figure 2.

With the fixed charge model a similar increase of the DPT rates is observed. At 300 K, no DPT was observed during 8 ns of propagation time whereas for $T = 350 \text{ K}$ and 500 K they are 1.0 ns^{-1} and 0.5 ns^{-1} , respectively. As the simulation time with fixed charges is 8 ns - shorter than simulations with the fluctuating charge model - the rates for DPT are based on fewer events with larger uncertainties which are not quantified in this work.

Conformational Coordinates Involved in DPT

Conformationally relevant degrees of freedom to consider are the $\text{O}_{\text{don}}\text{-O}_{\text{acc}}$ and the CO bond distances. The involvement of the $\text{O}_{\text{don}}\text{-O}_{\text{acc}}$ separation becomes apparent from noting that it contracts from $d_{\text{eq}}(\text{OO}) = 2.66 \text{ \AA}$ in the global minimum to $d_{\text{TS}}(\text{OO}) = 2.41 \text{ \AA}$ in the TS. In other words, compressing the O-O separation towards the TS structure facilitates (D)PT. The CO bond in the COH moiety has bond order 1 ($\text{BO} = 1$) whereas the second CO bond

is formally a double bond ($\text{BO} = 2$). This bonding pattern reverses after DPT and for the TS structure all bond orders are 1.5.

The first PT leads to a short-lived, metastable ion-pair with one protonated formic acid monomer and the corresponding anion, see Figure S1. Subsequently, the transferred hydrogen atom returns either to where it came from (attempted DPT) within an average delay time of 4.8 ± 2.9 fs at 350 K or the hydrogen atom along the second H-bond transfers (successful DPT) with an average delay time of 5.9 ± 2.6 fs. At 450 K the analysis of unbiased simulations find delay times of 5.1 ± 3.9 fs and 8.2 ± 7.1 fs for attempted and successful DPT, respectively. This is consistent with results from earlier work that reported an average delay time of ~ 8 fs for successful DPT.³⁰

Figures 4A and D compare the instantaneous $\text{O}_{\text{don}}\text{--O}_{\text{acc}}$ separations at the time of the second PT. The probability distribution functions $P(d_{\text{OO,A}}, d_{\text{OO,B}})$ for the O–O separations in unbiased simulations at 350 K and 400 K peak at (2.75 ± 0.18) Å and (2.78 ± 0.22) Å, respectively. These distances shorten significantly to (2.40 ± 0.05) Å and (2.42 ± 0.06) Å for successful DPT; see Figure 4. For successful DPT both O–O separations are between 2.4 Å and 2.6 Å whereas for attempted DPT the first O–O separation ranges from 2.3 Å to 2.5 Å but the second oxygen donor-acceptor distance covers a range between 2.4 and 2.9 Å. This suggests that a more symmetrical geometry at the time of the first PT increases the probability for successful DPT during the second step, i.e. if the distances between *both* oxygen atoms in the H-bonds are simultaneously contracted.

Next, the CO bond lengths are analysed. Figure 4B, C, E, and F shows the distribution of the CO_{acc} double and CO_{don} single bond distances from all MD simulations at 350 K and 400 K. The maximum peak positions are clearly distinguishable and demonstrate the single and double bond character of the CO bonds with $\text{BO} = 1$ and $\text{BO} = 2$, respectively. In




Figure 4: Bond distance distributions in FAD from simulations at 350 K (A-C) and 400 K (D-F). The blue colormap (panels A and D) shows the probability distribution of the $\text{O}_{\text{don}}\text{-O}_{\text{acc}}$ distances $P(d_{\text{OO,A}}, d_{\text{OO,B}})$ in both H-bonds. Oxygen-oxygen separations d_{OO} for successful (green diamonds) and attempted DPT events (red circles) at the time of the second PT are shown for both hydrogen bonds. The blue and red distributions (panels B, C, E, and F) show the distribution $P(d_{\text{CO,acc}})$ and $P(d_{\text{CO,don}})$ for the complete simulations. Distributions $P(d_{\text{CO,A}})$ (blue cross) and $P(d_{\text{CO,B}})$ (red plus) at the time of the second PT are reported for successful (B, E) or attempted DPT (C, F). For successful DPT they peak at the same CO separation (symmetric) whereas for attempted DPT they overlap less.

addition, the histograms for the CO distances at the second PT of a successful (panels B and E) and attempted (panels C and F) DPTs are reported. For successful DPT the distributions overlap for both temperatures and $\text{BO} = 1.5$ whereas for attempted DPT they are clearly non-overlapping and closer to $\text{BO} = 1$ and $\text{BO} = 2$, respectively.

The correlation between narrowly overlapping distance distributions of the CO single and double bond and the higher probability for successful DPT can be explained by the respective




Figure 5: Scan of the potential curve along the reactive coordinate ξ_1 of the hydrogen atom in the H-bond of FAD in the TS conformation. The three cases are for different elongations of the CO bond involving the first H-bond. CO distances are for the (A) TS conformation $d_{\text{TS}}(\text{CO})$ (solid line, blue), (B) average CO single bond separation $d_1(\text{CO})$ (dotted line, orange) and (C) average CO double bond length $d_2(\text{CO})$ (dashed line, green). The minimum of the potential for the TS conformation is shifted to 0 for reference.

OH bond potential in the protonated formic acid monomer. Figure 5 shows the PhysNet potential along the progression coordinate ξ_1 of a hydrogen atom in the TS of FAD. With all other bond lengths frozen in the TS conformation, a scan along ξ_1 is performed for three different distances of the CO bond involved in the first H-bond as in A) the TS conformation, B) a CO single bond and C) CO double bond (see illustration in Figure 5). The remaining CO bonds are kept frozen at the TS bond length of 1.26 Å. For $d_{\text{TS}}(\text{CO}) = 1.26$ Å (TS conformation with BO = 1.5) the potential minimum along the reactive coordinate is at $\xi_1 = 0$ Å, which changes if the CO bond contracts (B) or extends (C). The OH-stretch

potential is energetically more attractive in the covalent bonding range if the CO bond has $BO = 1$ rather than $BO = 2$. Vice versa, the OH-stretch potential becomes repulsive at shorter CO distances as in CO double bonds. Thus, hydrogen abstraction from an oxygen donor atom involved in a CO double bond is energetically more favourable compared to that involved in a CO single bond.

Solvent Distribution

MD simulations also provide information about the solvent distribution along the reaction path. Figure 6 reports the radial and distribution of the water oxygen atoms in the vicinity of the hydrogen atoms in the H-bonds of FAD from simulations at 300 K and 400 K. The solvent distributions are obtained either from unbiased MD simulations with FAD primarily in its reactant state and from biased simulations with FAD constrained close to its TS conformation by applying harmonic constraints along ξ_1 and ξ_2 with a force constant of $k = 115 \text{ kcal/mol/\AA}^2$. Analysis of the solvent distribution in the TS compared with the reactant structure is of interest to assess the amount of solvent reorganization that is required to reach the TS.

The solvent radial distribution functions around the hydrogen atoms of the H-bonds in Figures 6A and B exhibit a weak double peak between 4 and 6 Å. The maxima indicate the first solvation shell around FAD and the double peak structure is more pronounced for simulations with FAD in its TS structure. In the biased simulations FAD is more rigid which leads to a more structured solvent distribution. The radial distribution also shows a broad maximum around 8 Å that can be associated with a second solvation shell. The solvent radial distribution from simulation at 400 K (orange line in Figure 6A) features an additional peak around 2 Å that represents H-bond formation between branched, i.e. non-cyclic FAD, separated formic acid monomers and water molecules. Such H-bonds become available in




Figure 6: The radial distribution function $g(d)$ for the water-oxygen atoms around hydrogen atoms H_A and H_B of FAD in the reactant (A) and TS (B) structure from the from ML/MM (blue line) and CGenFF (dashed green line) simulations at 300 K. Panel A also shows $g(d)$ from one entire ML/MM simulation at 400 K (orange line) including formic acid dominantly as cyclic dimer and separated monomers (see Figure 2A). Panels C and D report the solvent distribution functions $P(x, y)$ around FAD from ML/MM MD simulations in its reactant (C) and TS (D) structures, respectively, and panel E reports the difference $\Delta P(x, y) = P_D - P_C$ between the two.

simulation at temperatures higher than 300 K as the probability for almost exclusively cyclic FAD decreases significantly (see Figure 2A).

For a more comprehensive characterization of the changes in the hydration of FAD between product/reactant geometries and the TS structure, 2-dimensional solvent distributions were determined. Figures 6C and D show the actual distributions projected onto the xy -plane which is the average plane containing the solute. Figure 6E reports the solvent density difference around FAD in its reactant and TS conformation, i.e. the difference density from panels C and D in Figure 6. In the 2d-histograms the first and second solvation shells are clearly visible as was also found from the radial distribution functions (Figure 6A and B). There is also a dent for the first solvation shell at $(-4, 0)$ Å and $(4, 0)$ Å along the horizontal axis joining the hydrogen atoms along the H-bonds of FAD.

The solvent distribution difference in Figure 6E indicates a shift of the first solvation shell closer to the side of the CH group of FAD for the biased simulations due to the short monomer-monomer distance in the TS structure compared with the reactant state. Hence, the solvent probability in this region is higher for FAD in its TS than in its reactant state. This is consistent with the shorter monomer-monomer distance of FAD in the TS conformation. The monomer-monomer contraction in the TS also flattens the dent towards the hydrogen atoms in the H-bond as shown by the blue area that indicates a lower solvent distribution density around FAD in the TS than the reactant state conformation. Finally, the solvent distribution difference exhibits the symmetry of FAD which indicates the convergence of the simulations.

Influence of the Solvent-Generated Electrical Field

Finally, DPT events are analyzed from the perspective of the alignment of the H-transfer motif relative to the solvent-generated electric field. Although cyclic FAD was found to be rather hydrophobic, the long ranged nature of the solvent-generated electric field can still significantly impact H-transfer along the progression coordinate for DPT. The effect of external fields on chemical reactions has been demonstrated from both, experiments⁸⁶ and computations.⁸⁷

In the following the electric field at the position of the transferring hydrogen atoms along the hydrogen bond was analyzed. For this, the force on the transferring hydrogen atom at the time of PT was determined as the linear interpolation of the forces for the 1 fs-frame before and after PT. The “time of PT” was determined by the sign change of the linearly interpolated reactive coordinate of the respective H-bond along which PT occurs. In Figures 7 and S10 the first PT - to which hydrogen atom H_A is attributed - is associated with the horizontal x -axis. In a successful DPT, the second PT is performed by hydrogen atom H_B (vertical y -axis) whereas in an attempted DPT the second PT is also performed by H_A

(back-transfer).




Figure 7: Correlation between the magnitude and direction of the electrostatic component of the solvent-generated Coulomb force \vec{F}_C^{MM} at the position of the transferring hydrogen atom during the first PT that leads to successful (green) or attempted (red) DPT; for definitions see Analysis. Results from unbiased (A and B) and biased simulations (C and D) are shown separately. The panels show the results from simulation at (A and C) 350 K and (B and D) 400 K. The forces on the hydrogen atom H_A that has completed the first PT are given along the horizontal axis. On the vertical axis the forces on the hydrogen atom H_B are shown for either supporting or inhibiting the second PT (for successful DPT or attempted DPT, respectively). By definition, the force vector always points towards O_{acc} in the H-bond. Effective electrostatic force on the hydrogen atom for a PT towards O_{acc} in the respective H-bond is “supportive”, and “inhibiting” otherwise. For the same analysis involving the second PT, see Figure S10.

Figures 7 and S10 report the projection of the solvent-generated (MM) Coulomb (C) force $F_{C,A}^{\text{MM}}$ along the $O_{\text{don}}-H_A \cdots O_{\text{acc}}$ and $O_{\text{don}}-H_B \cdots O_{\text{acc}}$ hydrogen bonds for successful and attempted DPTs during the first and second PT, respectively. As per definition, a positive force always supports PT of a hydrogen atom whereas a negative force has an inhibitory effect. The analysis was carried out for the second PT independent of whether transfer of H_B did (successful DPT) or did not (attempted DPT with H_A transferring back) occur.

In the unbiased simulations, the first PT is typically accompanied by a supportive Coulomb force. This effect is more pronounced at higher temperatures ($T = 400$ K, see Figure 7B) than at lower ones (see Figure 7A). Biased simulations increase the probability for DPT and provide improved statistics which also allows to fit 2-dimensional Gauss distributions to the data (Figures 7C and D). These results demonstrate that for the first PT the field along the H_A -bond is typically supportive. Interestingly, for an attempted DPT the field along the first H_A -bond at the time of PT is supportive but $F_{C,B}^{MM}$ along the second H_B -bond is more inhibiting. This applies to both, unbiased and biased simulations, see red distributions in Figure 7. Regarding Figure 7 the gravity center of the average solvent-generated Coulomb force is $[\bar{F}_{C,A}^{MM}, \bar{F}_{C,B}^{MM}] = [0.24, -0.23]$ and $[0.86, -1.50]$ kcal/mol/Å for successful and attempted DPT at 350 K, respectively, and shifts to $[0.51, -0.56]$ and $[-0.93, -1.65]$ kcal/mol/Å at 400 K. In biased simulations the gravity centers are at $[0.45, -0.43]$ and $[0.56, -0.78]$ kcal/mol/Å, $[0.56, -0.55]$ and $[-0.51, -0.81]$ kcal/mol/Å accordingly. Hence, for successful DPT the opposing electrostatic field along the second H-bond is considerably smaller in magnitude than for attempted DPT in all cases.

For the second PT the solvent-generated Coulomb force along both H-bonds is in general inhibiting (red distributions in Figure S10). In unbiased simulations at 350 K the center of gravity of the average solvent-generated Coulomb force is at $[\bar{F}_{C,A}^{MM}, \bar{F}_{C,B}^{MM}] = [-0.21, -0.22]$ and $[-0.84, -1.52]$ kcal/mol/Å for successful and attempted DPT, respectively. This shifts to $[-0.52, -0.58]$ and $[-0.92, -1.64]$ kcal/mol/Å at 400 K. In biased simulations the gravity centers are at $[-0.45, -0.42]$ and $[-0.56, -0.78]$ kcal/mol/Å at 350 K and $[-0.56, -0.54]$ and $[-0.51, -0.81]$ kcal/mol/Å at 400 K. Except for the sign change in the projection of the Coulomb force on the hydrogen atoms of the H_A -bond the center of gravity does not change significantly between the first and second PT in successful and attempted DPT.

Comparison of the solvent-generated Coulomb force for successful and attempted DPT further

establishes that the field of the water molecules at the position of the transferring hydrogen atom does not change appreciably within ~ 14 fs, which is the maximum time between first and second PT observed in the simulations at 350 K. This is expected as the rotational reorientation time of water in solution is on the order of several picoseconds.⁸⁸ The time evolution of the Coulomb force along the H-bonds is shown in Figures S11 and S12 for an example of a successful and attempted DPT, respectively.

In summary, successful DPT for FAD in water is often accompanied by a mildly supporting solvent-generated Coulomb force along the first PT in the H_A -bond whereas for the second PT in the H_B -bond the force is inhibiting but smaller in magnitude than for the back transfer along the first H_A -bond. As the inhibiting force along the H_B -bond increases, back-transfer along the first H_A -bond becomes favoured and results in attempted DPT with FAD returning to its initial reactant conformation, see Figure S1. The findings complete the picture that the probability for successful DPT primarily depends on the conformation of FAD itself at the first PT event. At conformation of high symmetry and low reaction barriers for PT events in both H-bonds, the solvent-generated Coulomb field impacts the probability of performing one PT in both H-bonds each (successful DPT) or a forth and back PT along one H-bond (attempted DPT).

Discussion and Conclusions

In this work double proton transfer in cyclic FAD in solution was characterized from extensive ML/MM MD simulations (total of ~ 100 ns for all temperatures and all charge models considered) using an energy function at MP2/aug-cc-pVTZ level quality for the solute akin to a QM/MM treatment at this level of quantum chemical theory. To put the present simulations in context it is worthwhile to note that 100 ps of a ML/MM MD simulation

(with ML trained at the MP2/aug-cc-pVTZ level) on 1 CPU takes ~ 5 days for the $5 \cdot 10^5$ energy and force evaluations. At the MP2/aug-cc-pVTZ level of theory the $5 \cdot 10^5$ energy evaluations alone, i.e. without calculating forces, on 1 CPU would take at least 4200 days. Clearly, QM/MM simulation at such a level of theory are not feasible at present and in the foreseeable future. It is further worth to mention that the PhysNet representation captures bond polarization through geometry-dependent, fluctuating charges. On the other hand, intramolecular polarization of FAD by the water solvent is not included but is expected to be small due to the hydrophobicity of cyclic FAD.

Spontaneous DPT is observed on the nanosecond time scale at 300 K to 350 K with a maximum rate of 0.9 ns^{-1} at 350 K. The total ML/MM energy function used was validated vis-a-vis B3LYP+D3 calculations and agrees to within $\sim 1 \text{ kcal/mol}$ for hydrated FAD with up to 15 surrounding water molecules. One particularly interesting aspect of the dynamics of FAD in solution is the fact that dissociation into a branched, singly H-bonded structure or into two separate formic acid monomers competes with DPT. The present results (see Figure 2) suggest that on the $\sim 14 \text{ ns}$ time scale the probability to form branched, singly H-bonded structures is considerably smaller than for cyclic FAD to exist (which decreases as T increases) or that of forming two separated monomers (which increases as T increases). Simulations at higher temperature allow to qualitatively assess conditions at higher pressure which has been used to induce proton transfer in other systems.^{89,90}

Results from previous simulation studies are somewhat ambiguous. Recent *ab initio* MD simulations using the BLYP functional including dispersion correction point towards pronounced hydrophobicity of cyclic FAD⁶² as was also found from earlier simulations using a fully empirical force field.⁵⁷ Both are consistent with the present work. On the other hand, simulations with this non-reactive, fully empirical force field report facile transitions from FAD to branched structures.⁵⁷ Such branched structures were also implied from experiments

which were, however, carried out in 3 molal NaCl.⁵⁹ Analysis of the T -dependent free energies for DPT yields a change in solvation entropy of $\Delta S = -2.1$ cal/mol/K. It is of interest to juxtapose this value with earlier work that rather focused on dissociation of FAD into two separated monomers for which $\Delta S_D = -3$ cal/mol/K was reported also using the TIP3P water model.⁵⁷ However, it is quite likely that the two entropy values which differ by about 30 % are not directly comparable as dimerization is driven by the different arrangements of solvent molecules around FAD and the FAMs whereas for DPT the reorganization is less pronounced; see Figure 6E.

As the calculated pK_D underestimates the experimentally measured one by 1 to 2 units (0.25 vs. 1.2 to 2.1),⁵⁷ it is expected that the energy function used in these earlier free energy simulations for cyclic FAD⁵⁷ underestimate the enthalpic contribution by about one order of magnitude. The reported stabilization from umbrella sampling simulations using the CHARMM22 force field was $\Delta G \sim -0.5$ kcal/mol. Hence, to be consistent with the experimentally determined pK_D , the enthalpic contribution to the stabilization of FAD in solution should rather be -5 kcal/mol or larger. Adding this to the stabilization energy for FAD in the gas phase using the CGenFF energy function (-9 kcal/mol) yields an estimated stabilization of -14 kcal/mol or larger which is more consistent with results from recent high-level electronic structure calculations which find a gas-phase stability for cyclic FAD of -16 kcal/mol.^{33,41} In other words, the unreactive, empirical CGenFF energy function considerably underestimates the stabilization of cyclic FAD in the gas phase and in solution which is also confirmed by the results in Figures S3 to S7 and leads to ready dissociation of FAD into two FAMs from such simulations.⁵⁷

Successful DPT is found to chiefly depend on the conformation of the cyclic FAD. In particular, increased symmetry with simultaneously contracted O-O distances in both H-bonds and CO bonds with $BO \approx 1.5$ lowers the reaction barrier for PT along both H-bonds. Successful DPT

requires the solvent induced electrostatic force to be only mildly inhibiting $\sim -0.5 \text{ kcal/mol/\AA}$ along the second H-bond. If the inhibiting force originating from the solvent is too strong along the second H-bond, attempted DPT dominates. For DPT, the MD simulations show an average delay between two PT events of ~ 5 and ~ 6 fs for attempted and successful DPT at 350 K, respectively, and the solvent-generated Coulomb field assists the first PT. A time difference of ~ 6 fs between the successive PT events for successful DPT compares with a time scale of ~ 155 fs for the $\text{O}_{\text{don}}-\text{O}_{\text{acc}}$ vibration. This suggests that DPT in hydrated FAD is essentially concerted and not stepwise; both PTs occur within one $\text{O}_{\text{don}}-\text{O}_{\text{acc}}$ vibration period. For a stepwise DPT the intermediate state (both protons on one FAM) would need to stabilize, i.e. have a lifetime, which is, however, not what is found here.

From a chemical perspective, one point of particular note is the successful and correct description of the change between bond orders 1 and 2 for the CO bond in PhysNet. Conventional empirical force fields encode the bond order in the equilibrium separation of the bonded term which does not allow easily for changes in the bond order depending on changes in the chemical environment. Although there are examples for capturing such effects⁹¹ making provisions for it in the context of empirical force fields is cumbersome. As Figures 4B, C, E, and F demonstrate, a NN-trained energy function based on reference electronic structure calculations successfully captures such chemical effects.

In summary, a ML/MM MD scheme was implemented and applied to DPT for FAD in solution. The results from ~ 100 ns of ML/MM MD simulations show that DPT and dissociation into two FAMs compete depending on temperature. FAD is predominantly hydrophobic which agrees with earlier findings, and the rate for DPT of $\sim 0.1 \text{ ns}^{-1}$ is consistent between biased and unbiased simulations. Extending the present work to other reaction types and protein-ligand binding will provide deeper chemical understanding and improved models suitable for statistically significant sampling to give molecular level insight into processes

in the condensed phase. Contrary to straight AIMD simulations at the correlated level, an ML/MM MD ansatz is feasible even for nanosecond time scale simulations.

Supporting Information

The supporting information reports additional graphics S1-S14.

Data Availability Statement

The data needed for the PhysNet representation of the MP2 PES is available at <https://github.com/MMunibas/fad.git>.

Acknowledgments

This work was supported by the Swiss National Science Foundation grants 200021-117810, 200020-188724, NCCR MUST, and the University of Basel which is gratefully acknowledged. This project received funding from the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie grant agreement No 801459 - FP-RESOMUS.

References

- (1) Warshel, A.; Weiss, R. M. An Empirical Valence Bond Approach for Comparing Reactions in Solutions and in Enzymes. *J. Am. Chem. Soc.* **1980**, *102*, 6218–6226.
- (2) Åqvist, J.; Warshel, A. Simulation of Enzyme Reactions Using Valence Bond Force Fields and Other Hybrid Quantum/classical Approaches. *Chem. Rev.* **1993**, *93*, 2523–2544.
- (3) Gajewski, J.; Brichford, N. Secondary Deuterium Kinetic Isotope Effects in the Aqueous

- Claisen Rearrangement: Evidence against an Ionic Transition State. *J. Am. Chem. Soc.* **1994**, *117*, 3165–3166.
- (4) Brickel, S.; Meuwly, M. Molecular Determinants for Rate Acceleration in the Claisen Rearrangement Reaction. *J. Phys. Chem. B* **2019**, *123*, 448–456.
- (5) Meuwly, M. Machine Learning for Chemical Reactions. *Chem. Rev.* **2021**, *121*, 10218–10239.
- (6) Kamerlin, S. C.; Haranczyk, M.; Warshel, A. Are Mixed Explicit/implicit Solvation Models Reliable for Studying Phosphate Hydrolysis? A Comparative Study of Continuum, Explicit and Mixed Solvation Models. *ChemPhysChem* **2009**, *10*, 1125–1134.
- (7) Plech, A.; Wulff, M.; Bratos, S.; Mirloup, F.; Vuilleumier, R.; Schotte, F.; Anfinrud, P. A. Visualizing Chemical Reactions in Solution by Picosecond X-ray Diffraction. *Phys. Rev. Lett.* **2004**, *92*, 125505.
- (8) Hu, H.; Yang, W. Free Energies of Chemical Reactions in Solution and in Enzymes with Ab Initio Quantum Mechanics/molecular Mechanics Methods. *Ann. Rev. Phys. Chem.* **2008**, *59*, 573–601.
- (9) Kim, K. H.; Kim, J. G.; Nozawa, S.; Sato, T.; Oang, K. Y.; Kim, T.; Ki, H.; Jo, J.; Park, S.; Song, C. et al. Direct Observation of Bond Formation in Solution with Femtosecond X-ray Scattering. *Nature* **2015**, *518*, 385–389.
- (10) Meuwly, M. Reactive Molecular Dynamics: From Small Molecules to Proteins. *WIREs Comput. Mol. Sci.* **2019**, *9*, e1386.
- (11) Nerenberg, P. S.; Head-Gordon, T. New Developments in Force Fields for Biomolecular Simulations. *Curr. Opin. Struct. Biol.* **2018**, *49*, 129–138.
- (12) Koner, D.; Salehi, S. M.; Mondal, P.; Meuwly, M. Non-conventional Force Fields for

- Applications in Spectroscopy and Chemical Reaction Dynamics. *J. Chem. Phys.* **2020**, *153*, 010901.
- (13) Guest, J. M.; Craw, J. S.; Vincent, M. A.; Hillier, I. H. The Effect of Water on the Claisen Rearrangement of Allyl Vinyl Ether: Theoretical Methods Including Explicit Solvent and Electron Correlation. *J. Chem. Soc., Perkin Trans. 2* **1997**, 71.
- (14) Jung, Y.; Marcus, R. A. On the Theory of Organic Catalysis “on Water”. *J. Am. Chem. Soc.* **2007**, *129*, 5492–5502, PMID: 17388592.
- (15) White, W. N.; Wolfarth, E. F. The O-claisen Rearrangement. Viii. Solvent Effects. *J. Org. Chem.* **1970**, *35*, 2196.
- (16) Acevedo, O.; Armacost, K. Claisen Rearrangements: Insight into Solvent Effects and “on Water” Reactivity from QM/MM Simulations. *J. Am. Chem. Soc.* **2010**, *132*, 1966.
- (17) van Keulen, S. C.; Solano, A.; Rothlisberger, U. How Rhodopsin Tunes the Equilibrium between Protonated and Deprotonated Forms of the Retinal Chromophore. *J. Chem. Theo. Comp.* **2017**, *13*, 4524–4534, PMID: 28731695.
- (18) El Hage, K.; Brickel, S.; Hermelin, S.; Gaulier, G.; Schmidt, C.; Bonacina, L.; van Keulen, S. C.; Bhattacharyya, S.; Chergui, M.; Hamm, P. et al. Implications of Short Time Scale Dynamics on Long Time Processes. *Struct. Dyn.* **2017**, *4*, 061507.
- (19) Mulholland, A. J.; Lyne, P. D.; Karplus, M. Ab Initio QM/MM Study of the Citrate Synthase Mechanism. A Low-barrier Hydrogen Bond Is Not Involved. *J. Am. Chem. Soc.* **2000**, *122*, 534–535.
- (20) Senn, H. M.; Thiel, W. QM/MM Methods for Biomolecular Systems. *Angew. Chem. Intern. Ed.* **2009**, *48*, 1198–1229.
- (21) Groenhof, G. In *Biomolecular Simulations: Methods and Protocols*; Monticelli, L., Salonen, E., Eds.; Humana Press: Totowa, NJ, 2013; pp 43–66.

- (22) van Duin, A. C. T.; Dasgupta, S.; Lorant, F.; Goddard III, W. A. . ReaxFF: A Reactive Force Field for Hydrocarbons. *J. Phys. Chem. A* **2001**, *105*, 9396–9409.
- (23) Nagy, T.; Yosa Reyes, J.; Meuwly, M. Multisurface Adiabatic Reactive Molecular Dynamics. *J. Chem. Theo. Comp.* **2014**, *10*, 1366–1375.
- (24) Ang, S. J.; Wang, W.; Schwalbe-Koda, D.; Axelrod, S.; Gómez-Bombarelli, R. Active Learning Accelerates Ab Initio Molecular Dynamics on Reactive Energy Surfaces. *Chem* **2021**, *7*, 738–751.
- (25) Bösel, L.; Thürlmann, M.; Riniker, S. Machine Learning in QM/MM Molecular Dynamics Simulations of Condensed-phase Systems. *J. Chem. Theo. Comp.* **2021**, *17*, 2641–2658.
- (26) Pan, Y.; McAllister, M. A. Characterization of Low-barrier Hydrogen Bonds. 1. Microsolvation Effects. An Ab Initio and DFT Investigation. *J. Am. Chem. Soc.* **1997**, *119*, 7561–7566.
- (27) Lim, J.-H.; Lee, E. K.; Kim, Y. Theoretical Study for Solvent Effect on the Potential Energy Surface for the Double Proton Transfer in Formic Acid Dimer and Formamidinium Dimer. *J. Phys. Chem. A* **1997**, *101*, 2233–2239.
- (28) Miura, S.; Tuckerman, M. E.; Klein, M. L. An Ab Initio Path Integral Molecular Dynamics Study of Double Proton Transfer in the Formic Acid Dimer. *J. Chem. Phys.* **1998**, *109*, 5290–5299.
- (29) Kohanoff, J.; Koval, S.; Estrin, D. A.; Laria, D.; Abashkin, Y. Concertedness and Solvent Effects in Multiple Proton Transfer Reactions: The Formic Acid Dimer in Solution. *J. Chem. Phys.* **2000**, *112*, 9498–9508.
- (30) Ushiyama, H.; Takatsuka, K. Successive Mechanism of Double-proton Transfer in Formic Acid Dimer: A Classical Study. *J. Chem. Phys.* **2001**, *115*, 5903–5912.

- (31) Kalescky, R.; Kraka, E.; Cremer, D. Local Vibrational Modes of the Formic Acid Dimer—the Strength of the Double Hydrogen Bond. *Mol. Phys.* **2013**, *111*, 1497–1510.
- (32) Ivanov, S. D.; Grant, I. M.; Marx, D. Quantum Free Energy Landscapes from Ab Initio Path Integral Metadynamics: Double Proton Transfer in the Formic Acid Dimer Is Concerted but Not Correlated. *J. Chem. Phys.* **2015**, *143*, 124304.
- (33) Miliordos, E.; Xantheas, S. S. On the Validity of the Basis Set Superposition Error and Complete Basis Set Limit Extrapolations for the Binding Energy of the Formic Acid Dimer. *J. Chem. Phys.* **2015**, *142*, 094311.
- (34) Tew, D. P.; Mizukami, W. Ab Initio Vibrational Spectroscopy of Cis-and Trans-formic Acid from a Global Potential Energy Surface. *J. Phys. Chem. A* **2016**, *120*, 9815–9828.
- (35) Qu, C.; Bowman, J. M. An Ab Initio Potential Energy Surface for the Formic Acid Dimer: Zero-point Energy, Selected Anharmonic Fundamental Energies, and Ground-State Tunneling Splitting Calculated in Relaxed 1–4-mode Subspaces. *Phys. Chem. Chem. Phys.* **2016**, *18*, 24835–24840.
- (36) Mackeprang, K.; Xu, Z.-H.; Maroun, Z.; Meuwly, M.; Kjaergaard, H. G. Spectroscopy and Dynamics of Double Proton Transfer in Formic Acid Dimer. *Phys. Chem. Chem. Phys.* **2016**, *18*, 24654–24662.
- (37) Richardson, J. O. Full-and Reduced-dimensionality Instanton Calculations of the Tunneling Splitting in the Formic Acid Dimer. *Phys. Chem. Chem. Phys.* **2017**, *19*, 966–970.
- (38) Qu, C.; Bowman, J. M. High-dimensional Fitting of Sparse Datasets of CCSD(T) Electronic Energies and MP2 Dipole Moments, Illustrated for the Formic Acid Dimer and Its Complex IR Spectrum. *J. Chem. Phys.* **2018**, *148*, 241713.
- (39) Qu, C.; Bowman, J. M. Quantum and Classical Ir Spectra of (HCOOH)₂, (DCOOH)₂

- and (DCOOD)₂ Using Ab Initio Potential Energy and Dipole Moment Surfaces. *Faraday Discuss.* **2018**, *212*, 33–49.
- (40) Qu, C.; Bowman, J. M. Ir Spectra of (HCOOH)₂ and (DCOOH)₂: Experiment, VSCF/VCI, and Ab Initio Molecular Dynamics Calculations Using Full-dimensional Potential and Dipole Moment Surfaces. *J. Phys. Chem. Lett.* **2018**, *9*, 2604–2610.
- (41) Käser, S.; Meuwly, M. Transfer Learned Potential Energy Surfaces: Accurate Anharmonic Vibrational Dynamics and Dissociation Energies for the Formic Acid Monomer and Dimer. *Phys. Chem. Chem. Phys.* **2022**,
- (42) Ito, F.; Nakanaga, T. A Jet-cooled Infrared Spectrum of the Formic Acid Dimer by Cavity Ring-down Spectroscopy. *Chem. Phys. Lett.* **2000**, *318*, 571–577.
- (43) Freytes, M.; Hurtmans, D.; Kass, S.; Liévin, J.; Vander Auwera, J.; Campargue, A.; Herman, M. Overtone Spectroscopy of Formic Acid. *Chem. Phys.* **2002**, *283*, 47–61.
- (44) Georges, R.; Freytes, M.; Hurtmans, D.; Kleiner, I.; Vander Auwera, J.; Herman, M. Jet-cooled and Room Temperature FTIR Spectra of the Dimer of Formic Acid in the Gas Phase. *Chem. Phys.* **2004**, *305*, 187–196.
- (45) Zielke, P.; Suhm, M. Raman Jet Spectroscopy of Formic Acid Dimers: Low Frequency Vibrational Dynamics and Beyond. *Phys. Chem. Chem. Phys.* **2007**, *9*, 4528–4534.
- (46) Xue, Z.; Suhm, M. A. Probing the Stiffness of the Simplest Double Hydrogen Bond: The Symmetric Hydrogen Bond Modes of Jet-cooled Formic Acid Dimer. *J. Chem. Phys.* **2009**, *131*, 054301.
- (47) Kollipost, F.; Larsen, R. W.; Domanskaya, A.; Nörenberg, M.; Suhm, M. Communication: The Highest Frequency Hydrogen Bond Vibration and an Experimental Value for the Dissociation Energy of Formic Acid Dimer. *J. Chem. Phys.* **2012**, *136*, 151101.

- (48) Nejad, A.; Suhm, M. A. Concerted Pair Motion Due to Double Hydrogen Bonding: The Formic Acid Dimer Case. *J. Ind. Inst. Sci.* **2020**, *100*, 5–19.
- (49) Reutemann, W.; Kieczka, H. *Ullmann's Encyclopedia of Industrial Chemistry*; American Cancer Society, 2011.
- (50) Balabin, R. M. Polar (acyclic) Isomer of Formic Acid Dimer: Gas-phase Raman Spectroscopy Study and Thermodynamic Parameters. *J. Phys. Chem. A* **2009**, *113*, 4910–4918.
- (51) Li, W.; Evangelisti, L.; Gou, Q.; Caminati, W.; Meyer, R. The Barrier to Proton Transfer in the Dimer of Formic Acid: A Pure Rotational Study. *Angew. Chem. Intern. Ed.* **2019**, *58*, 859–865.
- (52) Zhang, Y.; Li, W.; Luo, W.; Zhu, Y.; Duan, C. High Resolution Jet-cooled Infrared Absorption Spectra of (HCOOH)₂, (HCOOD)₂, and HCOOH-HCOOD Complexes in 7.2 μm Region. *J. Chem. Phys.* **2017**, *146*, 244306.
- (53) Ortlieb, M.; Havenith, M. Proton Transfer in (HCOOH)₂: An IR High-resolution Spectroscopic Study of the Antisymmetric C–O Stretch. *J. Phys. Chem. A* **2007**, *111*, 7355–7363.
- (54) Goroya, K. G.; Zhu, Y.; Sun, P.; Duan, C. High Resolution Jet-cooled Infrared Absorption Spectra of the Formic Acid Dimer: A Reinvestigation of the C–O Stretch Region. *J. Chem. Phys.* **2014**, *140*, 164311.
- (55) Zoete, V.; Meuwly, M. Double Proton Transfer in the Isolated and DNA-embedded Guanine-cytosine Base Pair. *J. Chem. Phys.* **2004**, *121*, 4377–4388.
- (56) Arabi, A. A.; Matta, C. F. Effects of Intense Electric Fields on the Double Proton Transfer in the Watson–crick Guanine–Cytosine Base Pair. *J. Phys. Chem. B* **2018**, *122*, 8631–8641.

- (57) Chen, J.; Brooks, C. L.; Scheraga, H. A. Revisiting the Carboxylic Acid Dimers in Aqueous Solution: Interplay of Hydrogen Bonding, Hydrophobic Interactions, and Entropy. *J. Phys. Chem. B* **2008**, *112*, 242–249.
- (58) Katchalsky, A.; Eisenberg, H.; Lifson, S. Hydrogen Bonding and Ionization of Carboxylic Acids in Aqueous Solutions. *J. Am. Chem. Soc.* **1951**, *73*, 5889–5890.
- (59) Schrier, E. E.; Pottle, M.; Scheraga, H. A. The Influence of Hydrogen and Hydrophobic Bonds on the Stability of the Carboxylic Acid Dimers in Aqueous Solution. *J. Am. Chem. Soc.* **1964**, *86*, 3444–3449.
- (60) Soffientini, S.; Bernasconi, L.; Imberti, S. The Hydration of Formic Acid and Acetic Acid. *J. Mol. Liq.* **2015**, *205*, 85–92.
- (61) Sobyra, T. B.; Melvin, M. P.; Nathanson, G. M. Liquid Microjet Measurements of the Entry of Organic Acids and Bases into Salty Water. *J. Phys. Chem. C* **2017**, *121*, 20911–20924.
- (62) Hänninen, V.; Murdachaew, G.; Nathanson, G. M.; Gerber, R. B.; Halonen, L. Ab Initio Molecular Dynamics Studies of Formic Acid Dimer Colliding with Liquid Water. *Phys. Chem. Chem. Phys.* **2018**, *20*, 23717–23725.
- (63) Tarakanova, E.; Voloshenko, G.; Kislina, I.; Mayorov, V.; Yukhnevich, G.; Lyashchenko, A. Composition and Structure of Hydrates Formed in Aqueous Solutions of Formic Acid. *J. Struct. Chem.* **2019**, *60*, 255–267.
- (64) Dou, Z.; Wang, L.; Hu, J.; Fang, W.; Sun, C.; Men, Z. Hydrogen Bonding Effect on Raman Modes of Formic Acid-water Binary Solutions. *J. Mol. Liq.* **2020**, *313*, 113595.
- (65) Larsen, A. H.; Mortensen, J. J.; Blomqvist, J.; Castelli, I. E.; Christensen, R.; Dulak, M.; Friis, J.; Groves, M. N.; Hammer, B.; Hargus, C. et al. The Atomic Simulation Environ-

- ment – a Python Library for Working with Atoms. *J. Phys. Condens. Matter* **2017**, *29*, 273002.
- (66) Unke, O. T.; Meuwly, M. Physnet: A Neural Network for Predicting Energies, Forces, Dipole Moments, and Partial Charges. *J. Chem. Theo. Comp.* **2019**, *15*, 3678–3693.
- (67) Jorgensen, W. L.; Chandrasekhar, J.; Madura, J. D.; Impey, R. W.; Klein, M. L. Comparison of Simple Potential Functions for Simulating Liquid Water. *J. Chem. Phys.* **1983**, *79*, 926–935.
- (68) Werner, H.-J.; Knowles, P. J.; Manby, F. R.; Black, J. A.; Doll, K.; Heßelmann, A.; Kats, D.; Köhn, A.; Korona, T.; Kreplin, D. A. et al. The Molpro Quantum Chemistry Package. *J. Chem. Phys.* **2020**, *152*, 144107.
- (69) Huang, B.; von Lilienfeld, O. A. Quantum Machine Learning Using Atom-in-molecule-based Fragments Selected on the Fly. *Nat. Chem.* **2020**, *12*, 945–951.
- (70) Behler, J. Constructing High-Dimensional Neural Network Potentials: A Tutorial Review. *Int. J. Quantum. Chem.* **2015**, *115*, 1032–1050.
- (71) Unke, O. T.; Meuwly, M. A Reactive, Scalable, and Transferable Model for Molecular Energies from a Neural Network Approach Based on Local Information. *J. Chem. Phys.* **2018**, *148*, 241708.
- (72) Grimme, S.; Ehrlich, S.; Goerigk, L. Effect of the Damping Function in Dispersion Corrected Density Functional Theory. *J. Comput. Phys.* **2011**, *32*, 1456–1465.
- (73) Abadi, M.; Agarwal, A.; Barham, P.; Brevdo, E.; Chen, Z.; Citro, C.; Corrado, G. S.; Davis, A.; Dean, J.; Devin, M. et al. TensorFlow: Large-scale Machine Learning on Heterogeneous Systems. 2015; <http://tensorflow.org/>, Software available from tensorflow.org.

- (74) Baydin, A. G.; Pearlmutter, B. A.; Radul, A. A.; Siskind, J. M. Automatic Differentiation in Machine Learning: A Survey. *J. Mach. Learn. Res.* **2018**, *18*, 1–43.
- (75) Mortensen, J. J.; Hansen, L. B.; Jacobsen, K. W. Real-space Grid Implementation of the Projector Augmented Wave Method. *Phys. Rev. B* **2005**, *71*, 035109.
- (76) Vanommeslaeghe, K.; Hatcher, E.; Acharya, C.; Kundu, S.; Zhong, S.; Shim, J.; Darian, E.; Guvench, O.; Lopes, P.; Vorobyov, I. et al. CHARMM General Force Field: A Force Field for Drug-like Molecules Compatible with the CHARMM All-Atom Additive Biological Force Fields. *J. Comput. Phys.* **2010**, *31*, 671–690.
- (77) Spohr, E. Effect of Electrostatic Boundary Conditions and System Size on the Interfacial Properties of Water and Aqueous Solutions. *J. Chem. Phys.* **1997**, *107*, 6342–6348.
- (78) Torrie, G. M.; Valleau, J. P. Nonphysical Sampling Distributions in Monte Carlo Free-energy Estimation: Umbrella Sampling. *J. Chem. Phys.* **1977**, *23*, 187–199.
- (79) Andersen, H. C. Rattle: A “Velocity” Version of the Shake Algorithm for Molecular Dynamics Calculations. *J. Comput. Phys.* **1983**, *52*, 24–34.
- (80) Enkovaara, J.; Rostgaard, C.; Mortensen, J. J.; Chen, J.; Dulak, M.; Ferrighi, L.; Gavnholt, J.; Glinsvad, C.; Haikola, V.; Hansen, H. A. et al. Electronic Structure Calculations with GPAW: A Real-space Implementation of the Projector Augmented-wave Method. *J. Phys.: Condens. Matter* **2010**, *22*, 253202.
- (81) Meuwly, M.; Karplus, M. Simulation of Proton Transfer along Ammonia Wires: An “Ab Initio” and Semiempirical Density Functional Comparison of Potentials and Classical Molecular Dynamics. *J. Chem. Phys.* **2002**, *116*, 2572–2585.
- (82) Becke, A. D. Density-functional Thermochemistry. Iii. The Role of Exact Exchange. *J. Chem. Phys.* **1993**, *98*, 5648–5652.

- (83) Lee, C.; Yang, W.; Parr, R. G. Development of the Colle-Salvetti Correlation-energy Formula into a Functional of the Electron Density. *Phys. Rev. B* **1988**, *37*, 785–789.
- (84) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A Consistent and Accurate Ab Initio Parametrization of Density Functional Dispersion Correction (DFT-D) for the 94 Elements H-Pu. *J. Chem. Phys.* **2010**, *132*, 154104.
- (85) Sidler, D.; Meuwly, M.; Hamm, P. An Efficient Water Force Field Calibrated against Intermolecular THz and Raman Spectra. *J. Chem. Phys.* **2018**, *148*, 244504.
- (86) Huang, X.; Tang, C.; Li, J.; Chen, L.-C.; Zheng, J.; Zhang, P.; Le, J.; Li, R.; Li, X.; Liu, J. et al. Electric Field-induced Selective Catalysis of Single-molecule Reaction. *Science advances* **2019**, *5*, eaaw3072.
- (87) Shaik, S.; De Visser, S. P.; Kumar, D. External Electric Field Will Control the Selectivity of Enzymatic-like Bond Activations. *J. Am. Chem. Soc.* **2004**, *126*, 11746–11749.
- (88) Ropp, J.; Lawrence, C.; Farrar, T.; Skinner, J. Rotational Motion in Liquid Water Is Anisotropic: A Nuclear Magnetic Resonance and Molecular Dynamics Simulation Study. *J. Am. Chem. Soc.* **2001**, *123*, 8047–8052.
- (89) Martins, D. M. S.; Middlemiss, D. S.; Pulham, C. R.; Wilson, C. C.; Weller, M. T.; Henry, P. F.; Shankland, N.; Shankland, K.; Marshall, W. G.; Ibberson, R. M. et al. Temperature- and Pressure-induced Proton Transfer in the 1:1 Adduct Formed between Squaric Acid and 4,4'-bipyridine. *J. Am. Chem. Soc.* **2009**, *131*, 3884–3893.
- (90) Ma, Z.; Li, J.; Sun, C.; Zhou, M. High Pressure Spectroscopic Investigation on Proton Transfer in Squaric Acid and 4,4'-bipyridine Co-crystal. *Sci. Rep.* **2017**, *7*, 4677.
- (91) Xu, Z.-H.; Meuwly, M. Vibrational Spectroscopy and Proton Transfer Dynamics in Protonated Oxalate. *J. Phys. Chem. A* **2017**, *121*, 5389–5398.

Supporting Information: Double Proton Transfer in Hydrated Formic Acid Dimer: Interplay of Spatial Symmetry and Solvent-Generated Force on Reactivity

Kai Töpfer, Silvan Käser and Markus Meuwly*

*Department of Chemistry, University of Basel, Klingelbergstrasse 80 , CH-4056 Basel,
Switzerland.*

E-mail: m.meuwly@unibas.ch




Figure S1: Schematic reaction profile of DPT in FAD. The intermediate ion-pair configuration after the first PT in the H_A -bond is shown square brackets. The second PT occurs either in the H_B -bond resulting in a successful DPT or in the H_A -bond again labelled as attempted DPT. Note the inversion of the covalent O-H bonds (blue line) in a successful DPT where the attempted DPT leads to the original configuration.




Figure S2: 1-dimensional free energy profiles from the 2-dimensional free energy surfaces at different force constants for the artificial harmonic potential in biased simulation of FAD in solution to constraint FAD around the TS conformation.




Figure S3: Average interaction energy and fluctuation around it between FAD and increasing number of closest water molecules N_{water} from 50 randomly chosen snapshots from a 2 ns simulation at 300 K. The standard deviation is given by the dashed error bars and the light grey lines shows the hydration energy of the chosen snapshots. The interaction energies are determined by (left panel) DFT (B3LYP+D3/aug-cc-pVTZ, BSSE corrected), (center panel) the current ML/MM model and (right panel) with the CHARMM program package with the CGenFF force field.




Figure S4: Average interaction energy and fluctuation around it between FAD and increasing number of closest water molecules N_{water} in 50 randomly chosen snapshots from 300 K simulations. The standard deviation is given by the dashed error bars which are slightly shifted for better visualization. The interaction energies are determined by DFT (B3LYP+D3/aug-cc-pVTZ, BSSE corrected), the current ML/MM model and with the CHARMM program package with the CGenFF force field.




Figure S5: Correlation between DFT and model potential energies for FAD surrounded by the 4 closest water molecules (relative to the FAD center of mass) for 50 randomly chosen snapshots from 300 K simulations. The DFT energy of the energetically lowest structure is shifted to zero and the model energies are shifted to minimize the mean absolute error (MAE) with the DFT energies. The linear regression for ML/MM vs. DFT and CGenFF vs. DFT are the dashed blue and red lines, respectively. The slope of the linear fit is given in the legend. The solid black line is the ideal 1:1 correlation and the MAE reported is with respect to the black solid line.




Figure S6: Correlation between DFT and model potential energies for FAD surrounded by the 8 closest water molecules (relative to the FAD center of mass) for 50 randomly chosen snapshots from 300 K simulations. The DFT energy of the energetically lowest structure is shifted to zero and the model energies are shifted to minimize the mean absolute error (MAE) with the DFT energies. The linear regression for ML/MM vs. DFT and CGenFF vs. DFT are the dashed blue and red lines, respectively. The slope of the linear fit is given in the legend. The solid black line is the ideal 1:1 correlation and the MAE reported is with respect to the black solid line.




Figure S7: Correlation between DFT and model potential energies for FAD surrounded by the 12 closest water molecules (relative to the FAD center of mass) for 50 randomly chosen snapshots from 300 K simulations. The DFT energy of the energetically lowest structure is shifted to zero and the model energies are shifted to minimize the mean absolute error (MAE) with the DFT energies. The linear regression for ML/MM vs. DFT and CGenFF vs. DFT are the dashed blue and red lines, respectively. The slope of the linear fit is given in the legend. The solid black line is the ideal 1:1 correlation and the MAE reported is with respect to the black solid line.




Figure S8: Predicted atomic charge probability distributions $P(q)$ from the trained PhysNet model for FAD. The distributions are shown for the carbon atoms C, carbon-bonded hydrogen atoms H_C, oxygen atoms O and oxygen-bonded hydrogen atoms H_O. Blue: $P(q)$ for 10000 conformations of cyclic-FAD from gas phase MD simulations; orange: $P(q)$ for 10000 conformations of non-cyclic FAD. The vertical black dashed lines mark the atomic charges for the corresponding atom types from the CGenFF force field.¹




Figure S9: 2-dimensional free energy surfaces $G(\xi_1, \xi_2)$ for DPT in FAD in solution with fluctuating charges for different temperatures.




Figure S10: Correlation between the magnitude and direction of the electrostatic component of the solvent force F_C^{MM} at the position of the transferring hydrogen atom during the second PT for successful (green) and attempted (red) DPT. For definitions see Analysis section. Results from unbiased (A and B) and US simulations (C and D) are shown separately from simulations at (A and C) 350 K and (B and D) 400 K. The forces on the hydrogen atom H_A involved in the first PT are given on the horizontal axis. In the vertical axis the forces on the hydrogen atom H_B are shown performing either the second PT or no PT for successful or attempted DPT, respectively. By definition, the force vector always points towards O_{acc} in the H-bond after the first PT. An electrostatic force on the hydrogen atom towards O_{acc} in the respective H-bond is labelled as “supportive”, and “inhibiting” otherwise.




Figure S11: Time sequence covering 1 ps of reporting the solvent-generated Coulomb force F_C^{MM} on the hydrogen atoms in both H-bonds before and after the first (top panel) and second PT (bottom panel) for a successful DPT at 350 K. The black solid and dashed lines follow the position of the O and H atoms in the respective H-bond. The two center panels provide the F_C^{MM} sequence in both H-bonds across a 70 fs window around the DPT. The red colormap denotes a Coulomb force on the respective H atom parallel and the blue colormap antiparallel along the H-bond vector defined between both oxygen atoms. The horizontal solid and dashed lines follow the position of the O and H atoms in the respective H-bond. The vertical dashed line at 0 fs is the time point of the first PT in the H_A -bond and dotted line of the second PT in the H_B -bond.




Figure S12: Time sequence covering 1 ps of reporting the solvent-generated Coulomb force F_{C}^{MM} on the hydrogen atoms in both H-bonds before and after the first (top panel) and second PT (bottom panel) for an attempted DPT event at 350 K. The black solid and dashed lines follow the position of the O and H atoms in the respective H-bond. The two center panels provide the F_{C}^{MM} sequence in both H-bonds across a 70 fs window around the DPT. The red colormap denotes a Coulomb force on the respective H atom parallel and the blue colormap antiparallel along the H-bond vector defined between both oxygen atoms. The horizontal passing solid and dashed line shows the position of the O and H atoms in the respective H-bond. The vertical dashed line at 0 fs is the time point of the first PT in the H_{A} -bond and dotted line of the second PT in the H_{B} -bond.




Figure S13: Correlation between difference in O–O separation $\Delta d_{\text{OO},\text{AB}} = d_{\text{OO},\text{A}} - d_{\text{OO},\text{B}}$ and difference in solvent-generated Coulomb force $\Delta F_{\text{C},\text{AB}}^{\text{MM}} = F_{\text{C},\text{A}}^{\text{MM}} - F_{\text{C},\text{B}}^{\text{MM}}$ of both H-bonds for the first PT at 350 K and 400 K of unbiased simulations, respectively. The average force difference for attempted DPT is indicated as the dashed red line. Manifestly, the average $\Delta d_{\text{OO},\text{AB}} < 0$ for attempted DPT whereas $\Delta d_{\text{OO},\text{AB}} \sim 0$, i.e. the structure is symmetric, for successful DPT.




Figure S14: Correlation between difference in O–O separation $\Delta d_{\text{OO},\text{AB}} = d_{\text{OO},\text{A}} - d_{\text{OO},\text{B}}$ and difference in solvent-generated Coulomb force $\Delta F_{\text{C},\text{AB}}^{\text{MM}} = F_{\text{C},\text{A}}^{\text{MM}} - F_{\text{C},\text{B}}^{\text{MM}}$ of both H-bonds for the second PT at 350 K and 400 K of unbiased simulations, respectively. The average force difference for attempted DPT is indicated as the dashed red line. Manifestly, the average $\Delta d_{\text{OO},\text{AB}} < 0$ for attempted DPT whereas $\Delta d_{\text{OO},\text{AB}} \sim 0$, i.e. the structure is symmetric, for successful DPT. Note the slight displacements in position and forces compared with Figure S13 as the first and second PT are separated by a few femtoseconds.

References

- (1) Vanommeslaeghe, K.; Hatcher, E.; Acharya, C.; Kundu, S.; Zhong, S.; Shim, J.; Darian, E.; Guvench, O.; Lopes, P.; Vorobyov, I. et al. CHARMM General Force Field: A Force Field for Drug-like Molecules Compatible with the CHARMM All-Atom Additive Biological Force Fields. *J. Comput. Phys.* **2010**, *31*, 671–690.