

# Connes spectral distance and nonlocality of generalized noncommutative phase spaces

Bing-Sheng Lin<sup>1,2,†</sup>, Tai-Hua Heng<sup>3</sup>

<sup>1</sup>School of Mathematics, South China University of Technology,  
Guangzhou 510641, China

<sup>2</sup>Laboratory of Quantum Science and Engineering,  
South China University of Technology, Guangzhou 510641, China

<sup>†</sup>Email: sclbs@scut.edu.cn

<sup>3</sup>School of Physics and Material Science, Anhui University, Hefei 230601, China

October 25, 2021

## Abstract

We study the Connes spectral distance of quantum states and analyse the nonlocality of the  $4D$  generalized noncommutative phase space. By virtue of the Hilbert-Schmidt operatorial formulation, we obtain the Dirac operator and construct a spectral triple corresponding to the noncommutative phase space. Based on the ball condition, we obtain some constraint relations about the optimal elements, and then calculate the Connes spectral distance between two Fock states. Due to the noncommutativity, the spectral distances between Fock states in generalized noncommutative phase space are shorter than those in normal phase space. This shortening of distances implies some kind of nonlocality caused by the noncommutativity. We also find that these spectral distances in  $4D$  generalized noncommutative phase space are additive and satisfy the normal Pythagoras theorem. When the noncommutative parameters equal zero, the results return to those in normal quantum phase space.

## 1 Introduction

The ideas of noncommutative spacetime started in 1947 [1]. In the 1980's, Connes formulated the mathematically rigorous framework of noncommutative geometry [2]. In the past decades, there has been much interest in the study of physics in noncommutative spaces [3–11]. A noncommutative spacetime also appeared in string theory, namely in the quantization of open string [3]. The noncommutativity of spacetime also plays an important role in quantum gravity [12, 13]. The concept of noncommutative spacetime is also applied in condensed matter physics, such as the integer quantum Hall effect [14]. Usually, only spatial noncommutativity is considered. But many researchers have also studied models in which a noncommutative geometry is defined on the whole phase space [15–24]. Noncommutativity between momenta can be naturally considered as a consequence

of noncommutativity between positions, as momenta are defined to be the partial derivatives of the action with respect to the position coordinates.

Because of the noncommutativity, there are no traditional point and distance in a noncommutative space. One can study states in noncommutative spaces. In a noncommutative space, a pure state is the analog of a traditional point in a normal commutative space. One can calculate some kinds of distance measures between the states, such as the Connes spectral distance [25]. The Connes spectral distances in some kinds of noncommutative spaces have already been studied in the literatures [26–37]. For example, Cagnache *et. al.* have studied the Connes spectral distance in the Moyal plane [27]. They explicitly computed Connes spectral distance between the pure states which corresponding to eigenfunctions of the quantum harmonic oscillators. Scholtz and his collaborators have developed the Hilbert-Schmidt operatorial formulation [32–34], they studied the Connes spectral distances of harmonic oscillator states and also coherent states in Moyal plane and fuzzy space. Barrett *et. al.* also used Monte Carlo simulation to study spectral distances in the fuzzy spaces [36].

In the present work, we study the Connes spectral distance of the quantum states in  $4D$  generalized noncommutative phase space (NCPS). We find that the Connes spectral distances of the quantum states in noncommutative phase space are shorter than those in normal phase space. This means that the noncommutativity of the generalized noncommutative phase space can lead to some kind of nonlocality. This nonlocality can equivalently make the distances shorter.

This paper is organized as follows. In Sec. 2, we construct a spectral triple corresponding to the  $4D$  generalized noncommutative phase space. Using the Hilbert-Schmidt operatorial formulation, we construct a boson Fock space and a quantum Hilbert space, and obtain the Dirac operator. In Sec. 3, we review the definition of Connes spectral distance. Based on the ball condition, we derive some constraint relations about the optimal elements. The Connes spectral distances between Fock states in generalized noncommutative phase space are calculated in Sec. 4. Some discussions and conclusions are given in Sec. 5. Some details of calculations of the Dirac operator and the Connes spectral distances are given in Appendixes.

## 2 Spectral triple of generalized noncommutative phase space

Now let us consider the following  $4D$  generalized noncommutative phase space [38], the position operators  $\hat{X}_i$  and the momentum operators  $\hat{P}_i$  satisfy the following commutation relations,

$$[\hat{X}_1, \hat{P}_1] = [\hat{X}_2, \hat{P}_2] = i\hbar, \quad [\hat{X}_1, \hat{X}_2] = i\mu, \quad [\hat{P}_1, \hat{P}_2] = i\nu, \quad (1)$$

and others vanish. Here  $\mu, \nu$  are some real parameters. In the present work, we only consider the case  $\mu, \nu > 0$ . For simplicity, we can use the following transformations [39],

$$\tilde{X}_i = \sqrt[4]{\frac{\nu}{\mu}} \hat{X}_i, \quad \tilde{P}_i = \sqrt[4]{\frac{\mu}{\nu}} \hat{P}_i, \quad (2)$$

and we have

$$[\tilde{X}_1, \tilde{P}_1] = [\tilde{X}_2, \tilde{P}_2] = i\hbar, \quad [\tilde{X}_1, \tilde{X}_2] = i\theta, \quad [\tilde{P}_1, \tilde{P}_2] = i\theta, \quad (3)$$

and others vanish. Here  $\theta = \sqrt{\mu\nu}$ . Usually we also assume  $\theta \ll \hbar$ .

In a normal  $4D$  quantum phase space, the coordinate operators  $\hat{x}_i, \hat{p}_i$  satisfy the following commutation relations,

$$[\hat{x}_i, \hat{p}_j] = i\delta_{ij}\hbar, \quad i, j = 1, 2, \quad (4)$$

and others vanish. Similarly, we can denote

$$\tilde{x}_i = \sqrt[4]{\frac{\nu}{\mu}} \hat{x}_i, \quad \tilde{p}_i = \sqrt[4]{\frac{\mu}{\nu}} \hat{p}_i, \quad (5)$$

and there is the similar commutation relation  $[\tilde{x}_i, \tilde{p}_j] = i\delta_{ij}\hbar$ . Define the creation and annihilation operators

$$\hat{a}_i = \frac{1}{\sqrt{2\hbar}}(\tilde{x}_i + i\tilde{p}_i), \quad \hat{a}_i^\dagger = \frac{1}{\sqrt{2\hbar}}(\tilde{x}_i - i\tilde{p}_i), \quad (6)$$

these operators satisfy the commutation relations  $[\hat{a}_i, \hat{a}_j^\dagger] = \delta_{ij}$ , and  $[\hat{a}_i, \hat{a}_j] = [\hat{a}_i^\dagger, \hat{a}_j^\dagger] = 0$ . One can define the Fock states as follows,

$$|n\rangle_i = \frac{1}{\sqrt{n!}}(\hat{a}_i^\dagger)^n|0\rangle, \quad n = 0, 1, 2, \dots, \quad (7)$$

where  $|0\rangle$  is the vacuum state, and  $\hat{a}_i|0\rangle = 0$ . These Fock states  $|n\rangle_i$  satisfy the relations

$$\hat{a}_i|n\rangle_i = \sqrt{n}|n-1\rangle_i, \quad \hat{a}_i^\dagger|n\rangle_i = \sqrt{n+1}|n+1\rangle_i. \quad (8)$$

We also have the resolution of the identity,

$$\sum_{n=0}^{\infty} |n\rangle_i \langle n| = \mathbb{I}_i. \quad (9)$$

Furthermore, one can define the boson Fock space as follows,

$$\mathcal{F} = \text{span} \{ |m, n\rangle := |m\rangle_1 \otimes |n\rangle_2, \quad m, n = 0, 1, 2, \dots \}. \quad (10)$$

There are the relations,

$$\begin{aligned} \sum_{n=0}^{\infty} |m, n\rangle \langle m, n| &= |m\rangle_1 \langle m| \otimes \mathbb{I}_2, & \sum_{m=0}^{\infty} |m, n\rangle \langle m, n| &= \mathbb{I}_1 \otimes |n\rangle_2 \langle n|, \\ \sum_{m,n=0}^{\infty} |m, n\rangle \langle m, n| &= \mathbb{I}_1 \otimes \mathbb{I}_2. \end{aligned} \quad (11)$$

By virtue of the Hilbert-Schmidt operatorial formulation [40], one can construct the quantum Hilbert space as follows,

$$\mathcal{Q} = \text{span} \{ |m_1, m_2\rangle \langle n_1, n_2| \}. \quad (12)$$

The elements  $\psi$  of the quantum Hilbert space  $\mathcal{Q}$  are denoted by  $|\psi\rangle$ , and the inner product is defined as

$$(\phi|\psi) := \text{tr}_{\mathcal{F}}(\phi^\dagger\psi) = \sum_{m,n=0}^{\infty} \langle m, n | \phi^\dagger\psi | m, n \rangle, \quad (13)$$

where  $\text{tr}_{\mathcal{F}}(\cdot)$  denotes the trace over  $\mathcal{F}$ .

In general, a noncommutative space corresponds to a spectral triple  $(\mathcal{A}, \mathcal{H}, \mathcal{D})$  [2], where  $\mathcal{A}$  is an involutive algebra acting on a Hilbert space  $\mathcal{H}$ , and  $\mathcal{D}$  is the Dirac operator on  $\mathcal{H}$ . One can construct a spectral triple  $(\mathcal{A}, \mathcal{H}, \mathcal{D})$  corresponding to the generalized noncommutative phase space (3) as follows,

$$\mathcal{A} = \mathcal{Q}, \quad \mathcal{H} = \mathcal{F} \otimes \mathbb{C}^4, \quad (14)$$

and an element  $e \in \mathcal{A}$  acts on  $\Psi = (|\psi_1\rangle, |\psi_2\rangle, |\psi_3\rangle, |\psi_4\rangle)' \in \mathcal{H}$  through the diagonal representation  $\pi$  as

$$\pi(e)\Psi = \pi(e) \begin{pmatrix} |\psi_1\rangle \\ |\psi_2\rangle \\ |\psi_3\rangle \\ |\psi_4\rangle \end{pmatrix} = \begin{pmatrix} e & 0 & 0 & 0 \\ 0 & e & 0 & 0 \\ 0 & 0 & e & 0 \\ 0 & 0 & 0 & e \end{pmatrix} \begin{pmatrix} |\psi_1\rangle \\ |\psi_2\rangle \\ |\psi_3\rangle \\ |\psi_4\rangle \end{pmatrix} = \begin{pmatrix} e|\psi_1\rangle \\ e|\psi_2\rangle \\ e|\psi_3\rangle \\ e|\psi_4\rangle \end{pmatrix}. \quad (15)$$

The Dirac operator for the  $4D$  generalized noncommutative phase space (3) can be defined as (see Appendix A for more details)

$$\mathcal{D} = \beta \begin{pmatrix} 0 & 0 & -\hat{A}_2^\dagger & -\hat{A}_1^\dagger \\ 0 & 0 & \hat{A}_1 & -\hat{A}_2 \\ -\hat{A}_2 & \hat{A}_1^\dagger & 0 & 0 \\ -\hat{A}_1 & -\hat{A}_2^\dagger & 0 & 0 \end{pmatrix}, \quad (16)$$

where

$$\beta = \frac{\sqrt{\hbar + \sqrt{\hbar^2 - \theta^2}}}{\sqrt{\hbar^2 - \theta^2}}, \quad \hat{A}_i = \hat{a}_i - it\varepsilon_{ij}\hat{a}_j, \quad (17)$$

and

$$t = \frac{\theta}{\hbar + \sqrt{\hbar^2 - \theta^2}}, \quad \varepsilon = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (18)$$

Obviously, there is  $0 < t \ll 1$ .

When the noncommutative parameters  $\mu, \nu \rightarrow 0$ , there is  $\theta \rightarrow 0$ , and  $t \rightarrow 0$ . The operators  $\hat{A}_i$  will return to the operators  $\hat{a}_i$ , and the Dirac operator  $\mathcal{D}$  (16) will return to the Dirac operator of normal  $4D$  quantum phase space.

### 3 Connes spectral distance and optimal elements

Let us consider the case where the quantum states  $\omega$  are normal and bounded, so they are representable by density matrices  $\rho$  [32]. The action of the state  $\omega$  on an element  $e \in \mathcal{A}$  can be written as

$$\omega(e) = \text{tr}_{\mathcal{F}}(\rho e). \quad (19)$$

Suppose the quantum states  $\omega$  and  $\omega'$  correspond to the density matrices  $\rho$  and  $\rho'$ , respectively. The Connes spectral distance between two states  $\omega$  and  $\omega'$  is [25]

$$d(\omega, \omega') \equiv \sup_{e \in B} |\omega(e) - \omega'(e)| = \sup_{e \in B} |\text{tr}_{\mathcal{F}}(\rho e) - \text{tr}_{\mathcal{F}}(\rho' e)|, \quad (20)$$

where

$$B = \left\{ e \in \mathcal{A} : \|\mathcal{D}, \pi(e)\|_{op} \leq 1 \right\}, \quad (21)$$

and the operator norm is defined as

$$\|a\|_{op} \equiv \sup_{\psi \in \mathcal{H}} \frac{\|a\psi\|}{\|\psi\|}, \quad \|a\|^2 \equiv (a, a) = \text{tr}_{\mathcal{F}}(a^\dagger a). \quad (22)$$

The inequality in (21) is the so-called ball condition. Using the Dirac operator  $\mathcal{D}$  (16), the commutator  $[\mathcal{D}, \pi(e)]$  for a Hermitian element  $e \in \mathcal{A}$  is

$$[\mathcal{D}, \pi(e)] = \beta \begin{pmatrix} 0 & D_1 \\ -D_1^\dagger & 0 \end{pmatrix}, \quad (23)$$

where

$$D_1 = \begin{pmatrix} [\hat{A}_2, e]^\dagger & [\hat{A}_1, e]^\dagger \\ [\hat{A}_1, e] & -[\hat{A}_2, e] \end{pmatrix}. \quad (24)$$

After some straightforward calculations, one can obtain

$$\|[\mathcal{D}, \pi(e)]^\dagger [\mathcal{D}, \pi(e)]\|_{op} = \beta^2 \max \left\{ \|D_1^\dagger D_1\|_{op}, \|D_1 D_1^\dagger\|_{op} \right\}. \quad (25)$$

From the above expression (24), we have

$$D_1^\dagger D_1 = \begin{pmatrix} [\hat{A}_2, e][\hat{A}_2, e]^\dagger + [\hat{A}_1, e]^\dagger [\hat{A}_1, e] & [\hat{A}_2, e][\hat{A}_1, e]^\dagger - [\hat{A}_1, e]^\dagger [\hat{A}_2, e] \\ [\hat{A}_1, e][\hat{A}_2, e]^\dagger - [\hat{A}_2, e]^\dagger [\hat{A}_1, e] & [\hat{A}_1, e][\hat{A}_1, e]^\dagger + [\hat{A}_2, e]^\dagger [\hat{A}_2, e] \end{pmatrix}. \quad (26)$$

So there is <sup>1</sup>

$$\|D_1^\dagger D_1\|_{op} \geq \sup_{\phi \in \mathcal{F}, \langle \phi | \phi \rangle = 1} \langle \phi | [\hat{A}_1, e]^\dagger [\hat{A}_1, e] + [\hat{A}_2, e][\hat{A}_2, e]^\dagger | \phi \rangle. \quad (28)$$

It is known that for any bounded operator  $a$ , there is

$$\|a\|_{op}^2 = \|a^\dagger\|_{op}^2 = \|a^\dagger a\|_{op}. \quad (29)$$

Using the ball condition in (21), there is the following inequality,

$$1 \geq \|[\mathcal{D}, \pi(e)]\|_{op}^2 = \|[\mathcal{D}, \pi(e)]^\dagger [\mathcal{D}, \pi(e)]\|_{op} = \beta^2 \|D_1^\dagger D_1\|_{op}. \quad (30)$$

So from (28) and (30), for any Hermitian element  $e \in B$ , one can obtain

$$\sup_{\phi \in \mathcal{F}, \langle \phi | \phi \rangle = 1} \langle \phi | [\hat{A}_1, e]^\dagger [\hat{A}_1, e] + [\hat{A}_2, e][\hat{A}_2, e]^\dagger | \phi \rangle \leq \frac{1}{\beta^2}. \quad (31)$$

---

<sup>1</sup>For any operator  $a$  with the matrix elements  $a_{ij}$  in some orthonormal bases, there is the following Bessel's inequality [34],

$$\|a\|_{op}^2 \geq \sum_i |a_{ij}|^2 \geq |a_{ij}|^2. \quad (27)$$

Similarly, we also have the following inequalities,

$$\begin{aligned}
\sup_{\phi \in \mathcal{F}, \langle \phi | \phi \rangle = 1} \langle \phi | [\hat{A}_1, e] [\hat{A}_1, e]^\dagger + [\hat{A}_2, e]^\dagger [\hat{A}_2, e] | \phi \rangle &\leq \frac{1}{\beta^2}, \\
\sup_{\phi \in \mathcal{F}, \langle \phi | \phi \rangle = 1} \langle \phi | [\hat{A}_1, e]^\dagger [\hat{A}_1, e] + [\hat{A}_2, e]^\dagger [\hat{A}_2, e] | \phi \rangle &\leq \frac{1}{\beta^2}, \\
\sup_{\phi \in \mathcal{F}, \langle \phi | \phi \rangle = 1} \langle \phi | [\hat{A}_1, e] [\hat{A}_1, e]^\dagger + [\hat{A}_2, e] [\hat{A}_2, e]^\dagger | \phi \rangle &\leq \frac{1}{\beta^2}.
\end{aligned} \tag{32}$$

Now consider an element  $e \in B$  which can be expressed as

$$e = \sum_{i,j,k,l=0}^{\infty} C_{k,l}^{i,j} |i, j\rangle \langle k, l|. \tag{33}$$

Since Hermitian elements can give the supremum in the Connes spectral distance function [41], we only need to consider the element  $e$  being Hermitian. So there are  $C_{k,l}^{i,j} = (C_{i,j}^{k,l})^*$ , and  $C_{i,j}^{i,j}$  are real numbers. Using the relations (8), after some straightforward calculations, we have

$$\begin{aligned}
[\hat{A}_1, e] &= [\hat{a}_1 - it\hat{a}_2, e] \\
&= \sum_{i,j,k,l=0}^{\infty} \left( \left( C_{k,l}^{i+1,j} \sqrt{i+1} - C_{k-1,l}^{i,j} \sqrt{k} \right) \right. \\
&\quad \left. - it \left( C_{k,l}^{i,j+1} \sqrt{j+1} - C_{k,l-1}^{i,j} \sqrt{l} \right) \right) |i, j\rangle \langle k, l|,
\end{aligned} \tag{34}$$

and then

$$\begin{aligned}
[\hat{A}_1, e] [\hat{A}_1, e]^\dagger &= \sum_{i,j=0}^{\infty} \left( \sum_{k,l=0}^{\infty} \left| \left( C_{k,l}^{i+1,j} \sqrt{i+1} - C_{k-1,l}^{i,j} \sqrt{k} \right) \right. \right. \\
&\quad \left. \left. - it \left( C_{k,l}^{i,j+1} \sqrt{j+1} - C_{k,l-1}^{i,j} \sqrt{l} \right) \right|^2 \right) |i, j\rangle \langle i, j| + n.d..
\end{aligned} \tag{35}$$

Here “*n.d.*” denotes the sum of terms  $|k, l\rangle \langle i, j|$  with  $k \neq i$  and/or  $l \neq j$ , and these terms will not affect the calculations in the following content. So we can just simply ignore these terms.

Similar to Refs. [27, 34], one can just consider the optimal elements being diagonal elements. So we can set  $C_{k,l}^{i,j} = 0$  if  $k \neq i$  and/or  $l \neq j$ , and denote  $c_{i,j} \equiv C_{i,j}^{i,j}$ . For convenience, we also denote

$$E_{i,j} \equiv c_{i,j} - c_{i-1,j}, \quad F_{i,j} \equiv c_{i,j} - c_{i,j-1}, \tag{36}$$

and

$$G_{i,j} \equiv iE_{i,j}^2 = i(c_{i,j} - c_{i-1,j})^2, \quad H_{i,j} \equiv jF_{i,j}^2 = j(c_{i,j} - c_{i,j-1})^2. \tag{37}$$

So the above equation (35) can be rewritten as

$$\begin{aligned}
[\hat{A}_1, e][\hat{A}_1, e]^\dagger &= \sum_{i,j=0}^{\infty} ((i+1)(c_{i+1,j} - c_{i,j})^2 \\
&\quad + t^2(j+1)(c_{i,j+1} - c_{i,j})^2) |i, j\rangle \langle i, j| + n.d. \\
&= \sum_{i,j=0}^{\infty} ((i+1)E_{i+1,j}^2 + t^2(j+1)F_{i,j+1}^2) |i, j\rangle \langle i, j| + n.d. \\
&= \sum_{i,j=0}^{\infty} (G_{i+1,j} + t^2 H_{i,j+1}) |i, j\rangle \langle i, j| + n.d. .
\end{aligned} \tag{38}$$

Similarly, there is

$$[\hat{A}_1, e]^\dagger [\hat{A}_1, e] = \sum_{i,j=0}^{\infty} (G_{i,j} + t^2 H_{i,j}) |i, j\rangle \langle i, j| + n.d. . \tag{39}$$

For the operator  $\hat{A}_2$ , we also have

$$\begin{aligned}
[\hat{A}_2, e] &= \sum_{i,j,k,l=0}^{\infty} \left( (C_{k,l}^{i,j+1} \sqrt{j+1} - C_{k,l-1}^{i,j} \sqrt{l}) \right. \\
&\quad \left. + it (C_{k,l}^{i+1,j} \sqrt{i+1} - C_{k-1,l}^{i,j} \sqrt{k}) \right) |i, j\rangle \langle k, l|,
\end{aligned} \tag{40}$$

and

$$\begin{aligned}
[\hat{A}_2, e][\hat{A}_2, e]^\dagger &= \sum_{i,j=0}^{\infty} (H_{i,j+1} + t^2 G_{i+1,j}) |i, j\rangle \langle i, j| + n.d. , \\
[\hat{A}_2, e]^\dagger [\hat{A}_2, e] &= \sum_{i,j=0}^{\infty} (H_{i,j} + t^2 G_{i,j}) |i, j\rangle \langle i, j| + n.d. .
\end{aligned} \tag{41}$$

From the inequalities (31) and (32), using the results (38), (39) and (41), one can obtain the following relations,

$$\begin{aligned}
(G_{i+1,j} + H_{i,j}) + t^2(G_{i,j} + H_{i,j+1}) &\leq \frac{1}{\beta^2}, \quad (1+t^2)(G_{i+1,j} + H_{i,j+1}) \leq \frac{1}{\beta^2}, \\
(G_{i,j} + H_{i,j+1}) + t^2(G_{i+1,j} + H_{i,j}) &\leq \frac{1}{\beta^2}, \quad (1+t^2)(G_{i,j} + H_{i,j}) \leq \frac{1}{\beta^2}.
\end{aligned} \tag{42}$$

So in the calculation of Connes spectral distance (20), in general, one only need to consider the so-called optimal elements,

$$e_o = \sum_{i,j=0}^{\infty} c_{i,j} |i, j\rangle \langle i, j|, \tag{43}$$

where the coefficients  $c_{i,j}$  satisfy the above constraint relations (42). The Connes spectral distance between the quantum states  $\omega$  and  $\omega'$  is obtained by

$$d(\omega, \omega') = |\text{tr}_{\mathcal{F}}(\rho e_o) - \text{tr}_{\mathcal{F}}(\rho' e_o)|. \tag{44}$$

## 4 Connes spectral distance between Fock states in NCPS

Using the spectral triple  $(\mathcal{A}, \mathcal{H}, \mathcal{D})$  constructed above, one can calculate the Connes spectral distances between Fock states  $|n_1, n_2\rangle$ . First, let us consider the adjacent Fock states  $|m+1, n\rangle$  and  $|m, n\rangle$ , and the corresponding density matrices are  $\rho_{m+1,n} = |m+1, n\rangle\langle m+1, n|$  and  $\rho_{m,n} = |m, n\rangle\langle m, n|$ , respectively. The spectral distance is

$$\begin{aligned} d(\omega_{m+1,n}, \omega_{m,n}) &= \sup_{e \in B} |\text{tr}_{\mathcal{F}}(\rho_{m+1,n}e) - \text{tr}_{\mathcal{F}}(\rho_{m,n}e)| \\ &= \sup_{e \in B} |\langle m+1, n|e|m+1, n\rangle - \langle m, n|e|m, n\rangle| \\ &= \sup_{e \in B} |c_{m+1,n} - c_{m,n}| = \sup_{e \in B} |E_{m+1,n}|. \end{aligned} \quad (45)$$

From (42), we have

$$\begin{aligned} (1+t^2)(G_{m+1,n} + H_{m,n+1}) &\leq \frac{1}{\beta^2}, \quad (1+t^2)(G_{m+1,n} + H_{m+1,n}) \leq \frac{1}{\beta^2}, \\ (G_{m+1,n} + H_{m+1,n+1}) + t^2(G_{m+2,n} + H_{m+1,n}) &\leq \frac{1}{\beta^2}, \\ (G_{m+1,n} + H_{m,n}) + t^2(G_{m,n} + H_{m,n+1}) &\leq \frac{1}{\beta^2}. \end{aligned} \quad (46)$$

So in order to attain the supremum of  $|c_{m+1,n} - c_{m,n}| = |E_{m+1,n}|$ , there should be

$$H_{m,n+1} = H_{m+1,n} = 0, \quad (47)$$

and

$$(1+t^2)G_{m+1,n} = (1+t^2)(m+1)E_{m+1,n}^2 = \frac{1}{\beta^2}, \quad (48)$$

or

$$|c_{m+1,n} - c_{m,n}| = |E_{m+1,n}| = \frac{1}{\beta\sqrt{(1+t^2)(m+1)}}. \quad (49)$$

We can also set

$$H_{m,n} = H_{m+1,n+1} = 0, \quad (50)$$

so there is

$$F_{m,n+1} = F_{m+1,n} = F_{m,n} = F_{m+1,n+1} = 0. \quad (51)$$

Without loss of generality, we can assume  $E_{m+1,n} \geq 0$ . There is

$$\begin{aligned} E_{m+1,n} &= F_{m+1,n+1} + E_{m+1,n} = c_{m+1,n+1} - c_{m,n} \\ &= E_{m+1,n+1} + F_{m,n+1} = E_{m+1,n+1}. \end{aligned} \quad (52)$$

Similarly, one can obtain

$$E_{m+1,i} = \frac{1}{\beta\sqrt{(1+t^2)(m+1)}}, \quad F_{m,i+1} = F_{m+1,i+1} = 0, \quad (53)$$

where  $i = 0, 1, 2, \dots$



For example, one can choose

$$c_{m,i} = 0, \quad c_{m+1,i} = \frac{1}{\beta\sqrt{(1+t^2)(m+1)}}, \quad (54)$$

and the optimal element is

$$\begin{aligned} e_o &= \sum_{i=0}^{\infty} c_{m+1,i} |m+1, i\rangle \langle m+1, i| \\ &= \frac{1}{\beta\sqrt{(1+t^2)(m+1)}} \sum_{i=0}^{\infty} |m+1, i\rangle \langle m+1, i| \\ &= \frac{1}{\beta\sqrt{(1+t^2)(m+1)}} |m+1\rangle_1 \langle m+1| \otimes \mathbb{I}_2. \end{aligned} \quad (55)$$

So the Connes distance between the states  $|m+1, n\rangle$  and  $|m, n\rangle$  is

$$\begin{aligned} d(\omega_{m+1,n}, \omega_{m,n}) &= |\text{tr}_{\mathcal{F}}(\rho_{m+1,n} e_o) - \text{tr}_{\mathcal{F}}(\rho_{m,n} e_o)| \\ &= |\langle m+1, n | e_o | m+1, n \rangle - \langle m, n | e_o | m, n \rangle| \\ &= \frac{1}{\beta\sqrt{(1+t^2)(m+1)}} \\ &= \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}} \frac{1}{\sqrt{m+1}}. \end{aligned} \quad (56)$$

Similarly, the Connes spectral distance between Fock states  $|m, n+1\rangle$  and  $|m, n\rangle$  is

$$d(\omega_{m,n+1}, \omega_{m,n}) = \frac{1}{\beta\sqrt{(1+t^2)(n+1)}} = \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}} \frac{1}{\sqrt{n+1}}. \quad (57)$$

Next, let us consider the Fock states  $|m, n\rangle$  and  $|m+k, n\rangle$ ,  $k > 1$ , and the corresponding Connes spectral distance is

$$\begin{aligned} d(\omega_{m+k,n}, \omega_{m,n}) &= \sup_{e \in B} |\text{tr}_{\mathcal{F}}(\rho_{m+k,n} e) - \text{tr}_{\mathcal{F}}(\rho_{m,n} e)| = \sup_{e \in B} |c_{m+k,n} - c_{m,n}| \\ &= \sup_{e \in B} |c_{m+k,n} - c_{m+k-1,n} + c_{m+k-1,n} - c_{m+k-2,n} \\ &\quad + \dots + c_{m+1,n} - c_{m,n}| \\ &= \sup_{e \in B} |E_{m+k,n} + E_{m+k-1,n} + \dots + E_{m+1,n}|. \end{aligned} \quad (58)$$

From (42), we have

$$\begin{aligned} (1+t^2)(G_{m+2,n} + H_{m+1,n+1}) &\leq \frac{1}{\beta^2}, \quad (1+t^2)(G_{m+1,n} + H_{m+1,n}) \leq \frac{1}{\beta^2}, \\ (G_{m+2,n} + H_{m+1,n}) + t^2(G_{m+1,n} + H_{m+1,n+1}) &\leq \frac{1}{\beta^2}, \\ (G_{m+1,n} + H_{m+1,n+1}) + t^2(G_{m+2,n} + H_{m+1,n}) &\leq \frac{1}{\beta^2}. \end{aligned} \quad (59)$$

So in order to attain the supremum of  $|c_{m+k,n} - c_{m,n}| = |E_{m+k,n} + E_{m+k-1,n} + \dots + E_{m+1,n}|$ , one must let  $|c_{m+2,n} - c_{m,n}| = |E_{m+2,n} + E_{m+1,n}|$  as large as possible. There should be  $H_{m+1,n} = H_{m+1,n+1} = 0$ , and

$$\begin{aligned} G_{m+2,n} + t^2 G_{m+1,n} &= \frac{1}{\beta^2}, & G_{m+1,n} + t^2 G_{m+2,n} &= \frac{1}{\beta^2}, \\ (1+t^2)G_{m+2,n} &= \frac{1}{\beta^2}, & (1+t^2)G_{m+1,n} &= \frac{1}{\beta^2}. \end{aligned} \quad (60)$$

Therefore we have

$$G_{m+1,n} = G_{m+2,n} = \frac{1}{\beta^2(1+t^2)}. \quad (61)$$

Similarly, in order to attain the supremum of  $|c_{m+k,n} - c_{m,n}|$ , there must be

$$H_{m+i,n} = H_{m+i,n+1} = 0, \quad G_{m+i,n} = \frac{1}{\beta^2(1+t^2)}, \quad i = 1, 2, \dots, k, \quad (62)$$

so

$$c_{m+i,n} - c_{m+i-1,n} = E_{m+i,n} = \frac{1}{\beta\sqrt{1+t^2}} \frac{1}{\sqrt{m+i}}. \quad (63)$$

For example, one can choose

$$c_{m+p,i} = \frac{1}{\beta\sqrt{1+t^2}} \sum_{j=1}^{m+p} \frac{1}{\sqrt{j}} = \zeta_{0;m+p} \frac{1}{\beta\sqrt{1+t^2}}, \quad (64)$$

where  $i = 0, 1, 2, \dots, p = 0, 1, \dots, k$ , and the function  $\zeta_{p;q}$  is

$$\zeta_{p;q} \equiv \sum_{i=p+1}^q \frac{1}{\sqrt{i}} = \zeta\left(\frac{1}{2}, p+1\right) - \zeta\left(\frac{1}{2}, q+1\right), \quad (65)$$

where  $\zeta(s, q)$  is the Hurwitz zeta function

$$\zeta(s, q) \equiv \sum_{i=0}^{\infty} \frac{1}{(q+i)^s}. \quad (66)$$

Obviously, there is  $\zeta_{i;j} = -\zeta_{j;i}$ , and therefore  $\zeta_{i;i} = 0$ . We also have  $\zeta_{i;j} - \zeta_{i;k} = \zeta_{k;j}$ . It is easy to verify that these coefficients  $c_{i,j}$  (64) satisfy the relations (62).

Suppose  $e_o$  to be the optimal element with the coefficients  $c_{i,j}$  (64) which can attain the supremum in (58). There is

$$\begin{aligned} e_o &= \sum_{p=0}^k \sum_{i=0}^{\infty} c_{m+p,i} |m+p, i\rangle \langle m+p, i| \\ &= \frac{1}{\beta\sqrt{1+t^2}} \sum_{p=0}^k \zeta_{0;m+p} |m+p\rangle \langle m+p| \otimes \mathbb{I}_2, \end{aligned} \quad (67)$$

and the Connes spectral distance between the states  $|m+k, n\rangle$  and  $|m, n\rangle$  is

$$\begin{aligned} d(\omega_{m+k,n}, \omega_{m,n}) &= |\text{tr}_{\mathcal{F}}(\rho_{m+k,n} e_o) - \text{tr}_{\mathcal{F}}(\rho_{m,n} e_o)| \\ &= |\langle m+k, n | e_o | m+k, n \rangle - \langle m, n | e_o | m, n \rangle| \\ &= \frac{1}{\beta\sqrt{1+t^2}} \sum_{i=1}^k \frac{1}{\sqrt{m+i}} = \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}} \sum_{i=1}^k \frac{1}{\sqrt{m+i}} \\ &= \zeta_{m;m+k} \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}}. \end{aligned} \quad (68)$$

Since these distances do not depend on  $n$ , for any integers  $n \neq n'$ , we always have  $d(\omega_{m+k,n}, \omega_{m,n}) = d(\omega_{m+k,n'}, \omega_{m,n'})$ . Obviously, there is  $d(\omega_{m+k,n}, \omega_{m,n}) = d(\omega_{m,n}, \omega_{m+k,n})$ .

It is easy to see that, for  $m, n, k, l \geq 0$ , these distances satisfy

$$d(\omega_{m+k+l,n}, \omega_{m,n}) = d(\omega_{m+k,n}, \omega_{m,n}) + d(\omega_{m+k+l,n}, \omega_{m+k,n}). \quad (69)$$

This means that these distances are additive.

Similarly, the Connes spectral distance between Fock states  $|m, n+k\rangle$  and  $|m, n\rangle$  is

$$d(\omega_{m,n+k}, \omega_{m,n}) = \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}} \sum_{i=1}^k \frac{1}{\sqrt{n+i}} = \zeta_{n;n+k} \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}}. \quad (70)$$

We also have  $d(\omega_{m,n+k}, \omega_{m,n}) = d(\omega_{m',n+k}, \omega_{m',n})$  for  $m \neq m'$ .

Compare with the result in Refs. [27, 32, 34], one can find that the Connes spectral distances between Fock states in generalized noncommutative phase space have one more factor  $\sqrt{1 - \theta^2/\hbar^2} = \sqrt{1 - \mu\nu/\hbar^2}$  than those in normal quantum phase space. So in general, the Connes spectral distances in generalized noncommutative phase space are shorter than those in normal quantum phase space. This is intuitive and reasonable, because the noncommutativity of the space will lead to some kind of nonlocality. The partial nonlocality can equivalently make the distances shorter.

Obviously, when  $\theta \rightarrow \hbar$ , these Connes spectral distances will become zero. This will cause some singularities.

Furthermore, using the above method, one can find that for any integers  $m, n, k, l \geq 0$ , the Connes spectral distance between the Fock states  $|m, n\rangle$  and  $|k, l\rangle$  is (see Appendix B for more details)

$$\begin{aligned} d(\omega_{k,l}, \omega_{m,n}) &= \frac{1}{\beta\sqrt{1+t^2}} \sqrt{\zeta_{m;k}^2 + \zeta_{n;l}^2} \\ &= \frac{\sqrt{\hbar^2 - \theta^2}}{\sqrt{2\hbar}} \sqrt{\zeta_{m;k}^2 + \zeta_{n;l}^2}. \end{aligned} \quad (71)$$

It is easy to verify that, these spectral distances between Fock states in NCPS also satisfy the Pythagoras theorem,

$$d(\omega_{m+k,n+l}, \omega_{m,n}) = \sqrt{d(\omega_{m+k,n}, \omega_{m,n})^2 + d(\omega_{m,n+l}, \omega_{m,n})^2}. \quad (72)$$

This also coincides with the result in Ref. [30].

## 5 Discussions and conclusions

In this paper, we study the Connes spectral distance of Fock states in  $4D$  generalized noncommutative phase space. By virtue of the Hilbert-Schmidt operatorial formulation, we obtain the Dirac operator and a spectral triple corresponding to the generalized noncommutative phase space. Based on the ball condition, we obtain some constraint relations about the optimal elements. Using these constraint

relations, we construct the corresponding optimal elements and derive the Connes spectral distance between two Fock states.

We find that the spectral distances in noncommutative phase space are shorter than those in normal phase space. This is intuitive and reasonable, because the noncommutativity of the space will lead to some kind of nonlocality. This nonlocality can make the distances shorter equivalently. We also find that these spectral distances in  $4D$  generalized noncommutative phase space are additive and satisfy the normal Pythagoras theorem. When the noncommutative parameter  $\theta = \sqrt{\mu\nu}$  equals zero, the results return to those in normal quantum phase space. But if  $\theta \rightarrow \hbar$ , these Connes spectral distances will become zero. This will cause some singularities. Usually, we assume  $\theta \ll \hbar$ .

Here we only study the Connes spectral distances between Fock states. Obviously, these distances depend on the specific quantum states being considered. One can also study the spectral distances between other kinds of pure states and mixed states. In a noncommutative space, a pure state is the analog of a traditional point in a normal commutative space, and the spectral distance between pure states corresponds to the geodesic distance between points. Different kinds of quantum states can form different kinds of abstract spaces, and these spaces have different mathematical structures. So studies of the Connes spectral distances between quantum states can help us to study the mathematical structures of the spaces and also physical properties of the quantum systems.

Our results are significant for the study of the Connes spectral distances of physical systems in noncommutative spaces. Our methods can be used to study other physical systems in other kinds of noncommutative spaces. Using our method, one can also study the Connes spectral distance between quantum states in higher-dimensional noncommutative spaces. But usually the calculations and results will be much more complicated.

## Acknowledgements

This work is partly supported by Key Research and Development Project of Guangdong Province (Grant No. 2020B0303300001), the National Natural Science Foundation of China (Grant No. 11911530750), the Guangdong Basic and Applied Basic Research Foundation (Grant No. 2019A1515011703), the Fundamental Research Funds for the Central Universities (Grant No. 2019MS109) and the Natural Science Foundation of Anhui Province (Grant No. 1908085MA16).

## A Dirac operator for $4D$ generalized noncommutative phase space

Similar to Ref. [34], in order to construct the Dirac operator for the  $4D$  generalized noncommutative phase space (3), one can consider the following extended noncommutative phase space in which the coordinate operators  $\tilde{X}_i$ ,  $\tilde{Y}_i$  and  $\tilde{P}_i$ ,  $\tilde{Q}_i$  satisfy the following commutation relations,

$$\begin{aligned} [\tilde{X}_i, \tilde{P}_j] &= [\tilde{Y}_i, \tilde{Q}_j] = i\delta_{ij}\hbar, & [\tilde{X}_i, \tilde{Y}_j] &= i\delta_{ij}\lambda, & [\tilde{Q}_i, \tilde{P}_j] &= i\delta_{ij}\lambda, \\ [\tilde{X}_1, \tilde{X}_2] &= [\tilde{Y}_1, \tilde{Y}_2] = i\theta, & [\tilde{P}_1, \tilde{P}_2] &= [\tilde{Q}_1, \tilde{Q}_2] = i\theta, \end{aligned} \quad (73)$$

where  $\lambda$  is some parameter.

It is easy to verify that, the following unitary representation acting on the quantum Hilbert space  $\mathcal{Q}$  satisfies the above noncommutative relations (73),

$$\begin{aligned}\tilde{X}_i|\phi\rangle &= \alpha|(\tilde{x}_i - t\varepsilon_{ij}\tilde{p}_j)\phi\rangle, & \tilde{P}_i|\phi\rangle &= \alpha|(\tilde{p}_i + t\varepsilon_{ij}\tilde{x}_j)\phi\rangle, \\ \tilde{Y}_i|\phi\rangle &= \frac{\alpha}{\sqrt{\hbar^2 - \theta^2}} \left[ \lambda|(\tilde{p}_i - t\varepsilon_{ij}\tilde{x}_j)\phi\rangle - \sqrt{\hbar^2 + \lambda^2 - \theta^2}|\phi(\tilde{x}_i + t\varepsilon_{ij}\tilde{p}_j)\rangle \right], \\ \tilde{Q}_i|\phi\rangle &= \frac{\alpha}{\sqrt{\hbar^2 - \theta^2}} \left[ \lambda|(\tilde{x}_i + t\varepsilon_{ij}\tilde{p}_j)\phi\rangle + \sqrt{\hbar^2 + \lambda^2 - \theta^2}|\phi(\tilde{p}_i - t\varepsilon_{ij}\tilde{x}_j)\rangle \right],\end{aligned}\quad (74)$$

where

$$\alpha = \frac{\sqrt{\hbar + \sqrt{\hbar^2 - \theta^2}}}{\sqrt{2\hbar}}. \quad (75)$$

The Dirac operator  $\mathcal{D}$  for the generalized noncommutative phase space (3) can be written as [5],

$$\mathcal{D} = \frac{1}{\lambda}\gamma^1\tilde{Y}_1 - \frac{1}{\lambda}\gamma^2\tilde{Q}_1 + \frac{1}{\lambda}\gamma^3\tilde{Y}_2 - \frac{1}{\lambda}\gamma^4\tilde{Q}_2, \quad (76)$$

where  $\gamma^k$ 's are the Euclidean Dirac matrices satisfying  $\gamma^k\gamma^l + \gamma^l\gamma^k = 2\delta_{kl}\mathbb{I}_4$ , for example,

$$\begin{aligned}\gamma^1 &= \begin{pmatrix} 0 & 0 & 0 & i \\ 0 & 0 & i & 0 \\ 0 & -i & 0 & 0 \\ -i & 0 & 0 & 0 \end{pmatrix}, & \gamma^2 &= \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}, \\ \gamma^3 &= \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & -i \\ -i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \end{pmatrix}, & \gamma^4 &= \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix}.\end{aligned}\quad (77)$$

So the Dirac operator (76) can be expressed as,

$$\mathcal{D} = \frac{1}{\lambda} \begin{pmatrix} 0 & 0 & i\tilde{Y}_2 - \tilde{Q}_2 & i\tilde{Y}_1 - \tilde{Q}_1 \\ 0 & 0 & i\tilde{Y}_1 + \tilde{Q}_1 & -i\tilde{Y}_2 - \tilde{Q}_2 \\ -i\tilde{Y}_2 - \tilde{Q}_2 & -i\tilde{Y}_1 + \tilde{Q}_1 & 0 & 0 \\ -i\tilde{Y}_1 - \tilde{Q}_1 & i\tilde{Y}_2 - \tilde{Q}_2 & 0 & 0 \end{pmatrix}. \quad (78)$$

After some straightforward calculations, one can obtain the commutator  $[\mathcal{D}, \pi(e)]$  acting on an element  $\Phi \in \mathcal{Q} \otimes \mathbb{C}^4$  as

$$\begin{aligned}[\mathcal{D}, \pi(e)]\Phi &= [\mathcal{D}, \pi(e)] \begin{pmatrix} |\phi_1\rangle \\ |\phi_2\rangle \\ |\phi_3\rangle \\ |\phi_4\rangle \end{pmatrix} \\ &= \beta \begin{pmatrix} 0 & 0 & [-\hat{a}_2^\dagger + it\hat{a}_1^\dagger, e] & [-\hat{a}_1^\dagger - it\hat{a}_2^\dagger, e] \\ 0 & 0 & [\hat{a}_1 - it\hat{a}_2, e] & [-\hat{a}_2 - it\hat{a}_1, e] \\ [-\hat{a}_2 - it\hat{a}_1, e] & [\hat{a}_1^\dagger + it\hat{a}_2^\dagger, e] & 0 & 0 \\ [-\hat{a}_1 + it\hat{a}_2, e] & [-\hat{a}_2^\dagger + it\hat{a}_1^\dagger, e] & 0 & 0 \end{pmatrix} \begin{pmatrix} |\phi_1\rangle \\ |\phi_2\rangle \\ |\phi_3\rangle \\ |\phi_4\rangle \end{pmatrix} \\ &= \beta \begin{pmatrix} 0 & 0 & [-\hat{A}_2^\dagger, e] & [-\hat{A}_1^\dagger, e] \\ 0 & 0 & [\hat{A}_1, e] & [-\hat{A}_2, e] \\ [-\hat{A}_2, e] & [\hat{A}_1^\dagger, e] & 0 & 0 \\ [-\hat{A}_1, e] & [-\hat{A}_2^\dagger, e] & 0 & 0 \end{pmatrix} \begin{pmatrix} |\phi_1\rangle \\ |\phi_2\rangle \\ |\phi_3\rangle \\ |\phi_4\rangle \end{pmatrix}.\end{aligned}\quad (79)$$

So regarding  $\Phi$  as a test function, one can identify the Dirac operator  $\mathcal{D}$  as

$$\mathcal{D} = \beta \begin{pmatrix} 0 & 0 & -\hat{A}_2^\dagger & -\hat{A}_1^\dagger \\ 0 & 0 & \hat{A}_1 & -\hat{A}_2 \\ -\hat{A}_2 & \hat{A}_1^\dagger & 0 & 0 \\ -\hat{A}_1 & -\hat{A}_2^\dagger & 0 & 0 \end{pmatrix}. \quad (80)$$

## B Calculations of Connes distance between Fock states $|m, n\rangle$ and $|m+k, n+l\rangle$

First, let us consider the distance between the states  $|m+1, n+1\rangle$  and  $|m, n\rangle$ ,

$$\begin{aligned} d(\omega_{m+1, n+1}, \omega_{m, n}) &= \sup_{e \in B} |\text{tr}_{\mathcal{F}}(\rho_{m+1, n+1} e) - \text{tr}_{\mathcal{F}}(\rho_{m, n} e)| \\ &= \sup_{e \in B} |c_{m+1, n+1} - c_{m, n}| = \sup_{e \in B} |E_{m+1, n+1} + F_{m, n+1}|. \end{aligned} \quad (81)$$

From (42), there is

$$\begin{aligned} (1+t^2)(G_{m+1, n} + H_{m, n+1}) &\leq \frac{1}{\beta^2}, & (1+t^2)(G_{m+1, n+1} + H_{m+1, n+1}) &\leq \frac{1}{\beta^2}, \\ G_{m+1, n+1} + H_{m, n+1} &\leq \frac{1}{\beta^2}, & G_{m+1, n} + H_{m+1, n+1} &\leq \frac{1}{\beta^2}. \end{aligned} \quad (82)$$

In order to attain the supremum of  $|E_{m+1, n+1} + F_{m, n+1}|$ , one must let  $G_{m+1, n+1} + H_{m, n+1}$  as large as possible. But if

$$G_{m+1, n+1} + H_{m, n+1} > \frac{1}{(1+t^2)\beta^2}, \quad (83)$$

then compare with (82), there must be

$$G_{m+1, n} < G_{m+1, n+1}, \quad H_{m+1, n+1} < H_{m, n+1}. \quad (84)$$

Without loss of generality, we can assume that  $E_{m+1, n+1}, F_{m, n+1} \geq 0$ . Therefore there are

$$E_{m+1, n} < E_{m+1, n+1}, \quad F_{m+1, n+1} < F_{m, n+1}. \quad (85)$$

But

$$E_{m+1, n} + F_{m+1, n+1} = c_{m+1, n+1} - c_{m, n} = E_{m+1, n+1} + F_{m, n+1}, \quad (86)$$

this is in contradiction with the relations (85). This means that the inequality (83) is not true.

So there is

$$G_{m+1, n+1} + H_{m, n+1} \leq \frac{1}{(1+t^2)\beta^2}, \quad (87)$$

In order to attain the supremum of  $|E_{m+1, n+1} + F_{m, n+1}|$ , there must be

$$G_{m+1, n+1} + H_{m, n+1} = \frac{1}{(1+t^2)\beta^2}, \quad (88)$$

and

$$G_{m+1, n} = G_{m+1, n+1}, \quad H_{m+1, n+1} = H_{m, n+1}, \quad (89)$$

or

$$E_{m+1,n} = E_{m+1,n+1}, \quad F_{m+1,n+1} = F_{m,n+1}. \quad (90)$$

So the Connes spectral distance between the states  $|m+1, n+1\rangle$  and  $|m, n\rangle$  is

$$\begin{aligned} d(\omega_{m+1,n+1}, \omega_{m,n}) &= \sup_{e \in B} |E_{m+1,n+1} + F_{m,n+1}| \\ &= \sup_{e \in B} \left| \frac{1}{\sqrt{m+1}} E_{m+1,n+1} \sqrt{m+1} + \frac{1}{\sqrt{n+1}} F_{m,n+1} \sqrt{n+1} \right| \\ &\leq \sup_{e \in B} \sqrt{\frac{1}{m+1} + \frac{1}{n+1}} \sqrt{(m+1)E_{m+1,n+1}^2 + (n+1)F_{m,n+1}^2} \\ &= \sup_{e \in B} \sqrt{\frac{1}{m+1} + \frac{1}{n+1}} \sqrt{G_{m+1,n+1} + H_{m,n+1}} \\ &= \sqrt{\frac{1}{m+1} + \frac{1}{n+1}} \sqrt{\frac{1}{(1+t^2)\beta^2}}. \end{aligned} \quad (91)$$

In the above inequality, we have used the Cauchy-Schwartz inequality,

$$(x_1 y_1 + x_2 y_2)^2 \leq (x_1^2 + x_2^2)(y_1^2 + y_2^2), \quad (92)$$

where the equality holds if  $x_1 y_2 = x_2 y_1$ .

So one can set

$$(m+1)E_{m+1,n+1} = (n+1)F_{m,n+1}, \quad (93)$$

and there should be

$$d(\omega_{m+1,n+1}, \omega_{m,n}) \leq \frac{1}{\beta\sqrt{1+t^2}} \sqrt{\frac{1}{m+1} + \frac{1}{n+1}}. \quad (94)$$

Furthermore, let us consider the Connes spectral distance between two Fock states  $|m, n\rangle$  and  $|m+k, n+l\rangle$ . For simplicity, we assume  $k, l > 1$ . The corresponding Connes spectral distance is

$$\begin{aligned} d(\omega_{m+k,n+l}, \omega_{m,n}) &= \sup_{e \in B} |\text{tr}_{\mathcal{F}}(\rho_{m+k,n+l} e) - \text{tr}_{\mathcal{F}}(\rho_{m,n} e)| \\ &= \sup_{e \in B} |c_{m+k,n+l} - c_{m,n}| = \sup_{e \in B} |c_{m+k,n+l} - c_{m+k,n} + c_{m+k,n} - c_{m,n}| \\ &= \sup_{e \in B} \left| \sum_{i=1}^l (c_{m+k,n+i} - c_{m+k,n+i-1}) + \sum_{j=1}^k (c_{m+j,n} - c_{m+j-1,n}) \right| \\ &= \sup_{e \in B} \left| \sum_{i=1}^l F_{m+k,n+i} + \sum_{j=1}^k E_{m+j,n} \right|. \end{aligned} \quad (95)$$

From (42), in order to attain the supremum of  $|c_{m+k,n+l} - c_{m,n}|$ , there must be

$$\begin{aligned} (1+t^2)(G_{m+2,n+1} + H_{m+1,n+2}) &= \frac{1}{\beta^2}, \quad (1+t^2)(G_{m+1,n+1} + H_{m+1,n+1}) = \frac{1}{\beta^2}, \\ (G_{m+2,n+1} + H_{m+1,n+1}) + t^2(G_{m+1,n+1} + H_{m+1,n+2}) &= \frac{1}{\beta^2}, \\ (G_{m+1,n+1} + H_{m+1,n+2}) + t^2(G_{m+2,n+1} + H_{m+1,n+1}) &= \frac{1}{\beta^2}. \end{aligned} \quad (96)$$

So there is

$$G_{m+2,n+1} = G_{m+1,n+1}, \quad H_{m+1,n+1} = H_{m+1,n+2}. \quad (97)$$

Similarly, one can obtain

$$\begin{aligned} G_{m+i,n+j} &= G_{m+1,n}, & 1 \leq i \leq k, 0 \leq j \leq l, \\ H_{m+i,n+j} &= H_{m,n+1}, & 0 \leq i \leq k, 1 \leq j \leq l, \end{aligned} \quad (98)$$

or

$$E_{m+i,n+j} = \sqrt{\frac{m+1}{m+i}} E_{m+1,n}, \quad F_{m+i,n+j} = \sqrt{\frac{n+1}{n+j}} F_{m,n+1}. \quad (99)$$

We also have

$$(1+t^2)(G_{m+1,n} + H_{m,n+1}) = \frac{1}{\beta^2}, \quad (100)$$

or

$$(m+1)E_{m+1,n}^2 + (n+1)F_{m,n+1}^2 = \frac{1}{(1+t^2)\beta^2}. \quad (101)$$

Therefore the Connes spectral distance between the Fock states  $|m, n\rangle$  and  $|m+k, n+l\rangle$  is

$$\begin{aligned} d(\omega_{m+k,n+l}, \omega_{m,n}) &= \sup_{e \in B} \left| \sum_{i=1}^k E_{m+i,n} + \sum_{j=1}^l F_{m,n+j} \right| \\ &= \sup_{e \in B} \left| \sqrt{m+1} E_{m+1,n} \sum_{i=1}^k \frac{1}{\sqrt{m+i}} + \sqrt{n+1} F_{m,n+1} \sum_{j=1}^l \frac{1}{\sqrt{n+j}} \right| \\ &= \sup_{e \in B} \left| \sqrt{m+1} E_{m+1,n} \zeta_{m;m+k} + \sqrt{n+1} F_{m,n+1} \zeta_{n;n+l} \right| \\ &\leq \sup_{e \in B} \sqrt{(m+1)E_{m+1,n}^2 + (n+1)F_{m,n+1}^2} \sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2} \\ &= \frac{1}{\beta \sqrt{1+t^2}} \sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}. \end{aligned} \quad (102)$$

Here we have used the Cauchy-Schwartz inequality (92) in the above inequality, and the equality holds if

$$\sqrt{m+1} E_{m+1,n} \zeta_{n;n+l} = \sqrt{n+1} F_{m,n+1} \zeta_{m;m+k}. \quad (103)$$

From (101) and (103), one can derive

$$E_{m+1,n} = \frac{1}{\beta \sqrt{1+t^2}} \frac{1}{\sqrt{m+1}} \frac{\zeta_{m;m+k}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}} = c_0 \frac{1}{\sqrt{m+1}}, \quad (104)$$

$$F_{m,n+1} = \frac{1}{\beta \sqrt{1+t^2}} \frac{1}{\sqrt{n+1}} \frac{\zeta_{n;n+l}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}} = d_0 \frac{1}{\sqrt{n+1}}, \quad (105)$$

where

$$c_0 = \frac{1}{\beta \sqrt{1+t^2}} \frac{\zeta_{m;m+k}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}}, \quad d_0 = \frac{1}{\beta \sqrt{1+t^2}} \frac{\zeta_{n;n+l}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}}. \quad (106)$$



So in order to attain the supremum of  $|c_{m+k,n+l} - c_{m,n}|$ , for example, one can choose

$$\begin{aligned} c_{p,q} &= c_0 \sum_{i=1}^p \frac{1}{\sqrt{i}} + d_0 \sum_{j=1}^q \frac{1}{\sqrt{j}} = c_0 \zeta_{0;p} + d_0 \zeta_{0;q} \\ &= \frac{1}{\beta \sqrt{1+t^2}} \frac{\zeta_{0;p} \zeta_{m;m+k} + \zeta_{0;q} \zeta_{n;n+l}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}}, \end{aligned} \quad (107)$$

where  $p = m, m+1, \dots, m+k$  and  $q = n, n+1, \dots, n+l$ . Obviously, there is

$$E_{i+1,j} = c_{i+1,j} - c_{i,j} = \frac{1}{\sqrt{i+1}} c_0, \quad F_{i,j+1} = c_{i,j+1} - c_{i,j} = \frac{1}{\sqrt{j+1}} d_0. \quad (108)$$

The corresponding optimal element  $e_o$  is

$$\begin{aligned} e_o &= \sum_{p=m}^{m+k} \sum_{q=n}^{n+l} c_{p,q} |p, q\rangle \langle p, q| \\ &= \frac{1}{\beta \sqrt{1+t^2}} \sum_{p=m}^{m+k} \sum_{q=n}^{n+l} \frac{\zeta_{0;p} \zeta_{m;m+k} + \zeta_{0;q} \zeta_{n;n+l}}{\sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}} |p, q\rangle \langle p, q|, \end{aligned} \quad (109)$$

and the spectral distance between the Fock states  $|m+k, n+l\rangle$  and  $|m, n\rangle$  is

$$\begin{aligned} d(\omega_{m+k,n+l}, \omega_{m,n}) &= |\text{tr}_{\mathcal{F}}(\rho_{m+k,n+l} e_o) - \text{tr}_{\mathcal{F}}(\rho_{m,n} e_o)| \\ &= |\langle m+k, n+l | e_o | m+k, n+l \rangle - \langle m, n | e_o | m, n \rangle| \\ &= c_0 \zeta_{m;m+k} + d_0 \zeta_{n;n+l} \\ &= \frac{1}{\beta \sqrt{1+t^2}} \sqrt{\zeta_{m;m+k}^2 + \zeta_{n;n+l}^2}. \end{aligned} \quad (110)$$

It is easy to verify that, these results (109) and (110) are also true for  $k \leq 0$  and/or  $l \leq 0$ .

## References

- [1] H. S. Snyder, “Quantized space-time.” *Phys. Rev.* **71** 38 (1947).
- [2] A. Connes, *Noncommutative geometry* (Academic Press, New York, 1994).
- [3] N. Seiberg, E. Witten, “String theory and noncommutative geometry.” *J. High Energy Phys.* **09** 032 (1999).
- [4] M. R. Douglas, N. A. Nekrasov, “Noncommutative field theory.” *Rev. Mod. Phys.* **73** 977 (2001).
- [5] V. Gayral, J. M. Gracia-Bondía, B. Iochum, T. Schücker, J. C. Várilly, “Moyal planes are spectral triple.” *Commun. Math. Phys.* **246** 569–623 (2004).

- [6] M. Chaichian, P. Prešnajder, A. Tureanu, “New concept of relativistic invariance in noncommutative space-time: Twisted Poincaré symmetry and its implications.” *Phys. Rev. Lett.* **94** 151602 (2005).
- [7] M. M. Ettefaghi, M. Haghighat, “Massive neutrino in noncommutative space-time.” *Phys. Rev. D* **77** 056009 (2008).
- [8] B. S. Lin, T. H. Heng, W. Chen, “Quantum field theory with a minimal length induced from noncommutative space.” *Commun. Theor. Phys.* **61** 605 (2014).
- [9] X. Calmet, C. Fritz, “Inflation on a non-commutative space-time.” *Phys. Lett. B* **747** 406 (2015).
- [10] J. Couch, S. Eccles, W. Fischler, M.-L. Xiao, “Holographic complexity and noncommutative gauge theory.” *J. High Energy Phys.* **03** 108 (2018).
- [11] A. Muhuri, D. Sinha, S. Ghosh, “Entanglement induced by noncommutativity: anisotropic harmonic oscillator in noncommutative space.” *Eur. Phys. J. Plus* **136** 35 (2021).
- [12] S. Doplicher, K. Fredenhagen, J. E. Roberts, “Spacetime quantization induced by classical gravity.” *Phys. Lett. B* **331** 39 (1994).
- [13] B. M. Zupnik, “Reality in noncommutative gravity.” *Class. Quantum Grav.* **24** 15 (2007).
- [14] A. P. Polychronakos, “Quantum Hall states on the cylinder as unitary matrix Chern-Simons theory.” *J. High Energy Phys.* **06** 070 (2001).
- [15] C. Duval, P. A. Horváthy, “The exotic Galilei group and the ‘Peierls substitution’.” *Phys. Lett. B* **479** 284 (2000).
- [16] V. P. Nair, A. P. Polychronakos, “Quantum mechanics on the noncommutative plane and sphere.” *Phys. Lett. B* **505** 267 (2001).
- [17] R. Banerjee, “A novel approach to noncommutativity in planar quantum mechanics.” *Mod. Phys. Lett. A* **17** 631 (2002).
- [18] J. Z. Zhang, “Fractional angular momentum in non-commutative spaces.” *Phys. Lett. B* **584** 204 (2004).
- [19] K. Li, J. H. Wang, C. Y. Chen, “Representation of noncommutative phase space.” *Mod. Phys. Lett. A* **20** 2165 (2005).
- [20] C. Bastos, O. Bertolami, N. C. Dias, J. N. Prata, “Phase-space noncommutative quantum cosmology.” *Phys. Rev. D* **78** 023516 (2008).
- [21] P. A. Horváthy, L. Martina, P. C. Stichel, “Exotic Galilean Symmetry and Non-Commutative Mechanics.” *SIGMA* **6** 060 (2010).
- [22] S. C. Jing, B. S. Lin, “A new kind of representations on noncommutative phase space.” *Phys. Lett. A* **372** 7109 (2008).

- [23] B. S. Lin, T. H. Heng, “Energy spectra of the harmonic oscillator in a generalized noncommutative phase space of arbitrary dimension.” *Chin. Phys. Lett.* **28** 070303 (2011).
- [24] Kh. P. Gnatenko, O. V. Shyiko, “Effect of noncommutativity on the spectrum of free particle and harmonic oscillator in rotationally invariant non-commutative phase space.” *Mod. Phys. Lett. A* **33** 1850091 (2018).
- [25] A. Connes, “Compact metric spaces, Fredholm modules and hyperfiniteness.” *Ergodic Theory Dynam. Systems* **9** 207-220 (1989).
- [26] G. Bimonte, F. Lizzi, G. Sparano, “Distances on a lattice from non-commutative geometry.” *Phys. Lett. B* **341** 139-146 (1994).
- [27] E. Cagnache, F. D’Andrea, P. Martinetti, J.-C. Wallet, “The spectral distance on the Moyal plane.” *J. Geom. Phys.* **61** 1881-1897 (2011).
- [28] P. Martinetti, L. Tomassini, “Noncommutative geometry of the Moyal plane: translation isometries, Connes’ distance on coherent states, Pythagoras equality.” *Commun. Math. Phys.* **323** 107–141 (2013).
- [29] F. D’Andrea, F. Lizzi, J.C. Várilly, “Metric Properties of the Fuzzy Sphere.” *Lett. Math. Phys.* **103** 183–205 (2013).
- [30] F. D’Andrea, P. Martinetti, “On Pythagoras theorem for products of spectral triples.” *Lett. Math. Phys.* **103** 469–492 (2013).
- [31] N. Franco, J.-C. Wallet, “Metrics and causality on Moyal planes.” *Contemporary Mathematics* **676** 147-173 (2016).
- [32] F. G. Scholtz, B. Chakraborty, “Spectral triplets, statistical mechanics and emergent geometry in non-commutative quantum mechanics.” *J. Phys. A: Math. Theor.* **46** 085204 (2013).
- [33] Y. Chaoba Devi, S. Prajapat, A. K. Mukhopadhyay, B. Chakraborty, F. G. Scholtz, “Connes distance function on fuzzy sphere and the connection between geometry and statistics.” *J. Math. Phys.* **56** 041707 (2015).
- [34] Y. Chaoba Devi, K. Kumar, B. Chakraborty, F. G. Scholtz, “Revisiting Connes’ finite spectral distance on noncommutative spaces: Moyal plane and fuzzy sphere.” *Int. J. Geo. Methods Mod. Phys.* **15** 1850204 (2018).
- [35] K. Kumar, B. Chakraborty, “Spectral distances on the doubled Moyal plane using Dirac eigenspinors.” *Phys. Rev. D* **97** 086019 (2018).
- [36] J. W Barrett, P. Druce, L. Glaser, “Spectral estimators for finite non-commutative geometries.” *J. Phys. A: Math. Theor.* **52** 275203 (2019).
- [37] A. Chakraborty, B. Chakraborty, “Spectral distance on Lorentzian Moyal plane.” *Int. J. Geo. Methods Mod. Phys.* **17** 2050089 (2020).
- [38] B. S. Lin, J. Xu, T. H. Heng, “Induced entanglement entropy of harmonic oscillators in non-commutative phase space.” *Mod. Phys. Lett. A* **34** 1950269 (2019).

- [39] B. S. Lin, S. C. Jing, “Deformed squeezed states in noncommutative phase space.” *Phys. Lett. A* **372** 4880 (2008).
- [40] F. G. Scholtz, L. Gouba, A. Hafver, C. M. Rohwer, “Formulation, interpretation and application of non-commutative quantum mechanics.” *J. Phys. A: Math. Theor.* **42** 175303 (2009).
- [41] B. Iochum, T. Krajewski, P. Martinetti, “Distances in finite spaces from noncommutative geometry.” *J. Geom. Phys.* **37** 100–125 (2001).