

On matrix Painlevé-4 equations. Part 2: Isomonodromic Lax pairs

I.A. Bobrova*, V.V. Sokolov^{†‡}

13 August 2021

Abstract

For all non-equivalent matrix systems of Painlevé-4 type found by authors in [4], isomonodromic Lax pairs are presented. Limiting transitions from these systems to matrix Painlevé-2 equations are found.

Keywords: Matrix Painlevé equations, isomonodromic Lax pairs

1 Introduction

It is known that the scalar system

$$\begin{cases} u' &= -u^2 + 2uv - 2zu + c_1, \\ v' &= -v^2 + 2uv + 2zv + c_2, \end{cases} \quad (1)$$

is equivalent to the Painlevé-4 equation

$$y'' = \frac{y'^2}{2y} + \frac{3}{2}y^3 + 4zy^2 + 2(z^2 - \gamma)y + \frac{\delta}{y}, \quad (2)$$

where $y(z) = u(z)$, $\gamma = 1 + \frac{1}{2}c_1 - c_2$, $\delta = -\frac{1}{2}c_1^2$. The prime $'$ means the derivative with respect to the variable $z \in \mathbb{C}$.

In the paper [4], matrix generalizations of system (1) of the form

$$\begin{cases} u' &= -u^2 + 2uv + \alpha(uv - vu) - 2zu + b_1u + ub_2 + b_3v + vb_4 + b_5 \\ v' &= -v^2 + 2vu + \beta(vu - uv) + 2zv + c_1v + vc_2 + c_3u + uc_4 + c_5, \end{cases} \quad (3)$$

were investigated. In system (3) the coefficients α , β are scalar and others are constant $n \times n$ -matrices. Here and below, we consider matrices over the field \mathbb{C} . The authors found all non-equivalent systems of the form (3) that satisfy the matrix Painlevé–Kovalevskaya test [2]. They can be written as follows:

$$\begin{cases} u' &= -u^2 + uv + vu - 2zu + h_1u + \gamma_1 \mathbb{I}, \\ v' &= -v^2 + vu + uv + 2zv - vh_1 + \gamma_2 \mathbb{I}, \end{cases} \quad \mathbb{P}_4^0$$

*National Research University “Higher School of Economics”, Moscow, Russian Federation.

[†]L.D. Landau Institute for Theoretical Physics, Chernogolovka, Russian Federation.

[‡]Federal University of ABC, Santo André, Sao Paulo, Brazil. E-mail: vsokolov@landau.ac.ru

$$\begin{cases} u' &= -u^2 + 2uv - 2zu + h_2, \\ v' &= -v^2 + 2uv + 2zv + h_2 + \gamma_1 \mathbb{I}, \end{cases} \quad \mathbf{P}_4^1$$

$$\begin{cases} u' &= -u^2 + 2uv - 2zu + h_2, \\ v' &= -v^2 + 3uv - vu + 2zv + h_1u + 2h_2 + \gamma_1 \mathbb{I}. \end{cases} \quad \mathbf{P}_4^2$$

Here $\gamma_1, \gamma_2 \in \mathbb{C}$, h_i are arbitrary matrices and two constant matrices h_1, h_2 in \mathbf{P}_4^2 are connected by the relation $[h_2, h_1] = -2h_1$.

In the present paper we construct for each of systems $\mathbf{P}_4^0 - \mathbf{P}_4^2$ an isomonodromic Lax representation of the form

$$A_z = B_\zeta + [B, A], \quad (4)$$

where ζ is the spectral parameter, $A(z, \zeta)$, $B(z, \zeta)$ are some 2×2 -matrices with entries being polynomials in u, v, z . We assume that the ζ -dependence of A and B is given by

$$A = A_1\zeta + A_0 + A_{-1}\zeta^{-1}, \quad B = B_1\zeta + B_0, \quad (5)$$

where A_1 and B_1 are constant diagonal matrices.

Remark 1. *It is clear that in the scalar case a pair of the form (5) admits the shift*

$$A \mapsto A + \kappa_1\zeta + \kappa_2 + \kappa_3\zeta^{-1} + \kappa_4z, \quad B \mapsto B + \kappa_4\zeta + \kappa_5f(u, v, z) + \kappa_6, \quad (6)$$

where κ_i are arbitrary constant scalar matrices, $f(u, v, z)$ is an arbitrary function. In addition, the scaling $\zeta \mapsto \text{const } \zeta$ and a conjugation of A and B by a constant diagonal matrix are allowed.

Section 3 is devoted to limiting transitions from the matrix Painlevé-4 systems $\mathbf{P}_4^0 - \mathbf{P}_4^2$ to the matrix Painlevé-2 equations $\mathbf{P}_2^0 - \mathbf{P}_2^2$ found in [1]. Extending these transitions to the Lax pairs from Section 2, we obtain Lax pairs for the Painlevé-2 equations.

2 Isomonodromic Lax pairs for systems $\mathbf{P}_4^0 - \mathbf{P}_4^2$

The \mathbf{P}_4 equation (2) admits several equivalent isomonodromic representations (4) with different dependence on the spectral parameter ζ (see for example [16]).

An isomonodromic pair of the form (5) for the scalar equation (2) was found in [14]. More precisely, the matrices

$$B(z, \zeta) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \zeta + \begin{pmatrix} 0 & w \\ 2(u - \theta_1)w^{-1} & 0 \end{pmatrix},$$

$$A(z, \zeta) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \zeta + \begin{pmatrix} z & w \\ 2(u - \theta_1)w^{-1} & -z \end{pmatrix} + \begin{pmatrix} -u + \theta_0 & -\frac{1}{2}vw \\ 2u(u - 2\theta_0)v^{-1}w^{-1} & z - \theta_0 \end{pmatrix} \zeta^{-1}$$

define the representation (5) for the system

$$\begin{cases} u' &= -2u^2v^{-1} - uv + 4\theta_0uv^{-1} + \theta_1v, \\ v' &= -4u + v^2 + 2zv + 4\theta_0, \\ w' &= -w(v + 2z), \end{cases}$$

which is equivalent to the P_4 equation (2) for $y(z) = v(z)$. The key point of our approach is the observation that the transformation

$$u \mapsto 2uv, \quad z \mapsto -z, \quad B \mapsto g B^T g^{-1} + g' g^{-1}, \quad A \mapsto -g A^T g^{-1},$$

where $g = \text{diag}\left(w^{-\frac{1}{2}}, w^{\frac{1}{2}}\right)$, brings this Lax pair to a polynomial pair for system (1) with

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \zeta + \begin{pmatrix} \frac{1}{2}v - z & uv + c_1 \\ 1 & -\frac{1}{2}v + z \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \zeta + \begin{pmatrix} z & -uv - c_1 \\ -1 & -z \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}c_2 & -u^2v - c_2u \\ v & -uv - \frac{1}{2}c_2 \end{pmatrix} \zeta^{-1}. \end{aligned} \quad (7)$$

As a result of the shift $z \mapsto z + \tau$ we obtain a pair for the system

$$\begin{cases} u' &= -u^2 + 2uv - 2zu + 2\tau u + c_1, \\ v' &= -v^2 + 2uv + 2zv - 2\tau v + c_2. \end{cases} \quad (8)$$

with matrices A and B given by

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \zeta + \begin{pmatrix} r_0v + r_1u - z + \tau + \kappa_6 & uv + c_1 \\ 1 & (r_0 - 1)v + r_1u + z - \tau + \kappa_6 \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \zeta + \begin{pmatrix} z - \tau + \kappa_2 & -uv - c_1 \\ -1 & -z + \tau + \kappa_2 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}c_2 + \kappa_3 & -u^2v - c_2u \\ v & -uv - \frac{1}{2}c_2 + \kappa_3 \end{pmatrix} \zeta^{-1}. \end{aligned} \quad (9)$$

The parameters $r_0, r_1, \kappa_2, \kappa_3, \kappa_6$ were implemented by a shift described in Remark 1.

We use a procedure of non-abelinization of the latter pair to the matrix case. Namely, we form an ansatz for the matrix pair replacing parameters and variables u and v by non-commutative ones such that in the case of 1×1 -matrices this ansatz coincides with (9). In particular, we replace the commutative product uv by $r uv + (1 - r) vu$ with unknown number r and so on.

Note that when the parameters in (9) become matrices, no shifts like (6) can remove them.

2.1 Case P_4^0

In the case of 1×1 -matrices the P_4^0 system has the form (8), where $\tau = \frac{1}{2}h_1$, $c_1 = \gamma_1$, $c_2 = \gamma_2$. Thus, the non-abelian ansatz becomes

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} r_0v + r_1u - z\mathbb{I} + \frac{1}{2}h_1 + \kappa_6 & r_2uv + (1 - r_2)vu + \gamma_1\mathbb{I} \\ \mathbb{I} & (r_0 - 1)v + r_1u + z\mathbb{I} - \frac{1}{2}h_1 + \kappa_6 \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z\mathbb{I} - \frac{1}{2}h_1 + \kappa_2 & -r_3uv - (1 - r_3)vu - \gamma_1\mathbb{I} \\ -\mathbb{I} & -z\mathbb{I} + \frac{1}{2}h_1 + \kappa_2 \end{pmatrix} \\ &\quad + \frac{1}{2} \begin{pmatrix} r_4uv + (1 - r_4)vu + \frac{1}{2}\gamma_2\mathbb{I} & -r_6u^2v - r_7uvu - (1 - r_6 - r_7)vu^2 - \gamma_2u \\ v & -r_5uv - (1 - r_5)vu - \frac{1}{2}\gamma_2\mathbb{I} \end{pmatrix} \zeta^{-1}. \end{aligned}$$

Substituting it in (5), we obtain two possible Lax pairs given by

$$\begin{aligned} \kappa_2 = \frac{h_1}{2}, \quad \kappa_6 = \frac{h_1}{2}, \quad r_0 = r_2 = r_3 = r_4 = r_7 = 1, \quad r_1 = r_5 = r_6 = 0, \\ \text{or } \kappa_2 = -\frac{h_1}{2}, \quad \kappa_6 = \frac{h_1}{2}, \quad r_4 = r_7 = 1, \quad r_0 = r_1 = r_2 = r_3 = r_5 = r_6 = 0. \end{aligned}$$

In the first case the matrices B and A read

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} v - z\mathbb{I} + h_1 & uv + \gamma_1\mathbb{I} \\ & \mathbb{I} & & z\mathbb{I} \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z\mathbb{I} & -uv - \gamma_1\mathbb{I} \\ -\mathbb{I} & -z\mathbb{I} + h_1 \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}\gamma_2\mathbb{I} & -uvu - \gamma_2u \\ v & -vu - \frac{1}{2}\gamma_2\mathbb{I} \end{pmatrix} \zeta^{-1}. \end{aligned} \quad (10)$$

To get the second pair, one can apply the following symmetry of the \mathbf{P}_4^0 system:

$$u \mapsto e^{-zh_1} u^T e^{zh_1}, \quad v \mapsto e^{-zh_1} v^T e^{zh_1}$$

to (10) and then conjugate the resulting Lax pair by the matrix $s = \begin{pmatrix} 0 & -\mathbb{I} \\ \mathbb{I} & 0 \end{pmatrix}$ to bring it to the form (9).

2.2 Case \mathbf{P}_4^1

In the case $\alpha = 0, \beta = -2$ we have $\tau = 0, c_1 = h_2, c_2 = h_2 + \gamma_1$ and the ansatz is given by

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} r_0v + r_1u - z\mathbb{I} & r_2uv + (1 - r_2)vu + h_2 \\ & \mathbb{I} & & (r_0 - 1)v + r_1u + z\mathbb{I} \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z\mathbb{I} & -r_3uv - (1 - r_3)vu - h_2 \\ -\mathbb{I} & -z\mathbb{I} \end{pmatrix} \\ &\quad + \frac{1}{2} \begin{pmatrix} r_4uv + (1 - r_4)vu & -r_6u^2v - r_7uvu - (1 - r_6 - r_7)vu^2 \\ +\frac{1}{2}h_2 + \frac{1}{2}\gamma_1\mathbb{I} + \kappa_3 & -r_8h_2u - (1 - r_8)uh_2 - \gamma_1u \\ v & -r_5uv - (1 - r_5)vu - \frac{1}{2}h_2 - \frac{1}{2}\gamma_1\mathbb{I} + \kappa_3 \end{pmatrix} \zeta^{-1}. \end{aligned}$$

From (5) it follows that

$$\begin{aligned} \kappa_3 = \frac{h_2}{2}, \quad r_1 = r_2 = r_3 = r_4 = r_7 = r_8 = 1, \quad r_0 = r_5 = r_6 = 0, \\ \text{or } \kappa_3 = -\frac{h_2}{2}, \quad r_2 = r_3 = r_4 = r_5 = r_6 = 1, \quad r_0 = r_1 = r_7 = r_8 = 0. \end{aligned}$$

In the second case the matrices B and A have the form

$$\begin{aligned} B(z, \zeta) &= \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} -z\mathbb{I} & uv + h_2 \\ & \mathbb{I} & & -v + z\mathbb{I} \end{pmatrix}, \\ A(z, \zeta) &= \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z\mathbb{I} & -uv - h_2 \\ -\mathbb{I} & -z\mathbb{I} \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}\gamma_1\mathbb{I} & -u^2v - u(h_2 + \gamma_1\mathbb{I}) \\ v & -uv - h_2 - \frac{1}{2}\gamma_1\mathbb{I} \end{pmatrix} \zeta^{-1}. \end{aligned} \quad (11)$$

The first pair should be equivalent to this, but we have not yet been able to find an appropriate gauge transformation.

2.3 Case \mathbf{P}_4^2

The case $\alpha = 0$, $\beta = -3$ is different from the previous two since for 1×1 -matrices the commutation relation $[h_2, h_1] = -2h_1$ implies $h_1 = 0$. If $h_1 = 0$ then in the scalar equation (8) we have $\tau = 0$, $c_1 = h_2$, $c_2 = 2h_2 + \gamma_1$. Substituting the ansatz

$$B(z, \zeta) = \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} r_0 v + r_1 u - z \mathbb{I} & r_2 uv + (1 - r_2)vu + h_2 \\ \mathbb{I} & (r_0 - 1)v + r_1 u + z \mathbb{I} \end{pmatrix},$$

$$A(z, \zeta) = \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z \mathbb{I} & -r_3 uv - (1 - r_3)vu - h_2 \\ -\mathbb{I} & -z \mathbb{I} \end{pmatrix}$$

$$+ \frac{1}{2} \begin{pmatrix} r_4 uv + (1 - r_4)vu & -r_6 u^2 v - r_7 uvu - (1 - r_6 - r_7)vu^2 \\ +h_2 + \frac{1}{2}\gamma_1 \mathbb{I} + \kappa_3 & -r_8 h_2 u - (2 - r_8)uh_2 - \gamma_1 u \\ v & -r_5 uv - (1 - r_5)vu - h_2 - \frac{1}{2}\gamma_1 \mathbb{I} + \kappa_3 \end{pmatrix} \zeta^{-1}$$

into (5), we obtain

$$\kappa_3 = 0, \quad r_1 = r_2 = r_3 = r_4 = r_5 = r_6 = r_8 = 1, \quad r_0 = r_7 = 0,$$

and therefore

$$B(z, \zeta) = \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} u - z \mathbb{I} & uv + h_2 \\ \mathbb{I} & u - v + z \mathbb{I} \end{pmatrix}, \quad (12)$$

$$A(z, \zeta) = \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z \mathbb{I} & -uv - h_2 \\ -\mathbb{I} & -z \mathbb{I} \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + h_2 + \frac{1}{2}\gamma_1 \mathbb{I} & -u^2 v - h_2 u - uh_2 - \gamma_1 u \\ v & -uv - h_2 - \frac{1}{2}\gamma_1 \mathbb{I} \end{pmatrix} \zeta^{-1}.$$

To implement h_1 into these matrices, we use the method of undetermined coefficients, combining it with some considerations of homogeneity. It is clear that the right hand sides of systems $\mathbf{P}_4^0 - \mathbf{P}_4^2$ are homogeneous polynomials with weights

$$w(u) = w(v) = w(z) = w(h_1) = 1, \quad w(h_2) = w(\gamma_1) = 2.$$

In the Lax pairs found above we have

$$w(A_j\{1, 1\}) = w(A_j\{2, 2\}) = w(B_j\{1, 1\}) = w(B_j\{2, 2\}) = 1 - j,$$

$$w(A_j\{1, 2\}) = w(B_j\{1, 2\}) = 2 - j, \quad w(A_j\{2, 1\}) = w(B_j\{2, 1\}) = -j,$$

where $j = 0, -1$. Suppose that this is true also for the Lax pair in the case $h_1 \neq 0$. All monomials of weight 1, 2 and 3 that contain h_1 are the following:

$$\begin{array}{ll} 1 : & h_1; \\ 2 : & h_1^2, \\ & zh_1, uh_1, h_1 u, vh_1, h_1 v; \\ 3 : & h_1^3, \\ & zh_1^2, uh_1^2, h_1 uh_1, h_1^2 u, vh_1^2, h_1 vh_1, h_1^2 v, \\ & z^2 h_1, h_1 u^2, uh_1 u, u^2 h_1, h_1 v^2, vh_1 v, v^2 h_1, \\ & \gamma_1 h_1, h_1 h_2, h_2 h_1, uvh_1, vuh_1, h_1 uv, h_1 vu, uh_1 v, vh_1 u, \\ & zh_1 u, zuh_1, zh_1 v, zvh_1. \end{array}$$

Adding to the matrices (12) arbitrary linear combinations of these monomials of proper weights, we constitute a new ansatz with 57 unknown coefficients. Substituting it to (4) and solving the corresponding large but simple algebraic system, we obtain the Lax pair for P_4^2 with

$$\begin{aligned}
B(z, \zeta) &= \begin{pmatrix} \mathbb{I} & 0 \\ 0 & -\mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} u - z\mathbb{I} & uv + \frac{1}{2}h_1u + h_2 \\ \mathbb{I} & u - v + z\mathbb{I} \end{pmatrix}, \\
A(z, \zeta) &= \begin{pmatrix} -\mathbb{I} & 0 \\ 0 & \mathbb{I} \end{pmatrix} \zeta + \begin{pmatrix} z\mathbb{I} & -uv - \frac{1}{2}h_1u - h_2 \\ -\mathbb{I} & -z\mathbb{I} \end{pmatrix} \\
&\quad + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}h_1u + h_2 + \frac{1}{2}\gamma_1\mathbb{I} & -u^2v - \frac{1}{2}uh_1u - \frac{1}{2}h_1uv + zh_1u \\ & -h_2u - uh_2 - \gamma_1u - \frac{1}{2}h_1h_2 \\ & v & -uv - \frac{1}{2}h_1u - h_2 - \frac{1}{2}\gamma_1\mathbb{I} \end{pmatrix} \zeta^{-1}.
\end{aligned} \tag{13}$$

Additional monomials with h_1 are marked in blue.

3 Limiting transitions in Lax pairs

In the scalar case, the scheme of degenerations of the Painlevé equations [8] and their isomonodromic Lax pairs is well-known. In particular, there exists a limiting transition from the equation P_4 to the equation P_2 . We construct similar transitions from the systems $P_4^0 - P_4^2$ and their Lax pairs found in Section 2 to the matrix P_2 equations

$$\begin{aligned}
y'' &= 2y^3 + xy + by + yb + \alpha\mathbb{I}, & \alpha \in \mathbb{C}, \quad b \in \text{Mat}_n(\mathbb{C}), & P_2^0 \\
y'' &= [y, y'] + 2y^3 + xy + a, & a \in \text{Mat}_n(\mathbb{C}), & P_2^1 \\
y'' &= 2[y, y'] + 2y^3 + xy + by + yb + a, & a, b \in \text{Mat}_n(\mathbb{C}), \quad [b, a] = 2b, & P_2^2
\end{aligned}$$

discovered in the paper [1]. As a result, we obtain also Lax pairs for P_2 equations equivalent to those presented in [1].

In the scalar case, one can apply to the system (1) the following transformation:

$$\begin{aligned}
z &= \frac{1}{4}\varepsilon^{-3} - \varepsilon x, & u(z) &= -\frac{1}{4}\varepsilon^{-3} - \varepsilon^{-1}f(x), & v(z) &= -2\varepsilon g(x), & (14) \\
c_1 &= -\frac{1}{16}\varepsilon^{-6}, & c_2 &= 2\theta, & \theta &\in \mathbb{C},
\end{aligned}$$

to bring it to the form

$$\begin{cases} f' &= 2\varepsilon^2(2fg - xf) - f^2 + g - \frac{1}{2}x, \\ g' &= 2\varepsilon^2(-g^2 + xg) + 2fg + \theta. \end{cases}$$

Passing to the limit $\varepsilon \rightarrow 0$, we obtain the system

$$\begin{cases} f' &= -f^2 + g - \frac{1}{2}x, \\ g' &= 2fg + \theta, \end{cases} \tag{15}$$

which is equivalent to the Painlevé-2 equation

$$y'' = 2y^3 + xy + \left(\theta - \frac{1}{2}\right)$$

for $y(x) = f(x)$.

This degeneration procedure can be extended to the Lax pair (7). Conjugating (7) by the matrix $\tilde{g} = \begin{pmatrix} 1 & -u \\ 0 & 1 \end{pmatrix}$, we get

$$B(z, \zeta) = \begin{pmatrix} 1 & 2u \\ 0 & -1 \end{pmatrix} \zeta + \begin{pmatrix} -u + \frac{1}{2}v - z & 0 \\ 1 & u - \frac{1}{2}v + z \end{pmatrix},$$

$$A(z, \zeta) = \begin{pmatrix} -1 & -2u \\ 0 & 1 \end{pmatrix} \zeta + \begin{pmatrix} u + z & -uv + u^2 + 2zu - c_1 \\ -1 & -u - z \end{pmatrix} + \frac{1}{2} \begin{pmatrix} \frac{1}{2}c_2 & 0 \\ v & -\frac{1}{2}c_2 \end{pmatrix} \zeta^{-1}.$$

Let us make the transformation (14), supplemented by

$$\zeta = 2\varepsilon\lambda, \quad (16)$$

in the relation (4). This leads to the ε -dependent pair

$$B(x, \lambda) = \begin{pmatrix} -2\varepsilon^2 & \varepsilon^{-1} + 4\varepsilon f \\ 0 & 2\varepsilon^2 \end{pmatrix} \lambda + \begin{pmatrix} -f + \varepsilon^2 g - \varepsilon^2 x & 0 \\ -\varepsilon & f - \varepsilon^2 g + \varepsilon^2 x \end{pmatrix},$$

$$A(x, \lambda) = \begin{pmatrix} -4\varepsilon^2 & 2(\varepsilon^{-1} + 4\varepsilon f) \\ 0 & 4\varepsilon^2 \end{pmatrix} \lambda$$

$$+ \begin{pmatrix} -2f - 2\varepsilon^2 x & -4\varepsilon f(g - x) + \varepsilon^{-1}(2f^2 - g + x) \\ -2\varepsilon & 2f + 2\varepsilon^2 x \end{pmatrix} + \begin{pmatrix} \frac{1}{2}\theta & 0 \\ -\varepsilon g & -\frac{1}{2}\theta \end{pmatrix} \lambda^{-1},$$

in which, after conjugation by the matrix $g = \text{diag}(1, \varepsilon^{-1})$, one can pass to the limit $\varepsilon \rightarrow 0$ to obtain the pair

$$B(x, \lambda) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -f & 0 \\ -1 & f \end{pmatrix}, \quad (17)$$

$$A(x, \lambda) = \begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -2f & 2f^2 - g + x \\ -2 & 2f \end{pmatrix} + \begin{pmatrix} \frac{1}{2}\theta & 0 \\ -g & -\frac{1}{2}\theta \end{pmatrix} \lambda^{-1}$$

for the P_2 system (15). Here we replace z by x and ζ by λ in the formula (4). The pair (17) is just the Harnad–Tracy–Widom pair [11] up to a scaling.

We are going to describe similar relations between matrix systems $P_4^0 - P_4^2$ and P_2 equations $P_2^0 - P_2^2$. All these equations have the form

$$y'' = \kappa[y, y'] + 2y^3 + xy + b_1 y + y b_2 + a, \quad \kappa \in \mathbb{C}, \quad a, b_1, b_2 \in \text{Mat}_n(\mathbb{C}).$$

Such a matrix equation can be rewritten as the system of two equations

$$\begin{cases} f' &= -f^2 + g - \frac{1}{2}x\mathbb{I} - c_1, \\ g' &= 2gf + \beta[g, f] + c_2 f + f c_3 + c_4, \end{cases} \quad \beta \in \mathbb{C}, \quad c_i \in \text{Mat}_n(\mathbb{C}). \quad (18)$$

Here $f(x) = y(x)$, $\kappa = -1 - \beta$, $b_1 = c_2 + (2 + \beta)c_1$, $b_2 = c_3 - \beta c_1$, and $a = c_4 - \frac{1}{2}\mathbb{I}$.

3.1 Case $\alpha = \beta = -1$

Let us apply transformations (14), (16) taking together with

$$h_1 = 4\varepsilon b, \quad \gamma_1 = -\frac{1}{16}\varepsilon^{-6}, \quad \gamma_2 = 2\theta, \quad b \in \text{Mat}_n(\mathbb{C}), \quad \theta \in \mathbb{C}$$

to the pair (10). Then, after the passage to the limit $\varepsilon \rightarrow 0$, we obtain matrices

$$\begin{aligned} B(x, \lambda) &= \begin{pmatrix} 0 & \mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -f & 0 \\ -\mathbb{I} & f \end{pmatrix}, \\ A(x, \lambda) &= \begin{pmatrix} 0 & 2\mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -2f & 2f^2 - g + x\mathbb{I} + 2b \\ -2\mathbb{I} & 2f \end{pmatrix} + \begin{pmatrix} \frac{1}{2}\theta\mathbb{I} & 0 \\ -g & -\frac{1}{2}\theta\mathbb{I} \end{pmatrix} \lambda^{-1}, \end{aligned} \tag{19}$$

which define the Lax pair for the P_2^0 system

$$\begin{cases} f' &= -f^2 + g - \frac{1}{2}x\mathbb{I} - b, \\ g' &= fg + gf + \theta\mathbb{I} \end{cases} \tag{20}$$

of the form (18).

3.2 Case $\alpha = 0, \beta = -2$

As in the previous case, for the limiting transition in the pair (11), we supplement the change of variables (14), (16) with the following transformations:

$$h_2 = 2a - \frac{1}{16}\varepsilon^{-6}\mathbb{I}, \quad \gamma_1 = \frac{1}{16}\varepsilon^{-6}, \quad a \in \text{Mat}_n(\mathbb{C}),$$

and perform the shift of the form (6) with $\kappa_3 = \frac{1}{64}\varepsilon^{-6}\mathbb{I}$. Then in the limit $\varepsilon \rightarrow 0$ we get the pair

$$\begin{aligned} B(x, \lambda) &= \begin{pmatrix} 0 & \mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -f & 0 \\ -\mathbb{I} & f \end{pmatrix}, \\ A(x, \lambda) &= \begin{pmatrix} 0 & 2\mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -2f & 2f^2 - g + x\mathbb{I} \\ -2\mathbb{I} & 2f \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ -g & -fg + gf - a \end{pmatrix} \lambda^{-1}, \end{aligned}$$

for the P_2^1 system

$$\begin{cases} f' &= -f^2 + g - \frac{1}{2}x\mathbb{I}, \\ g' &= 2fg + a \end{cases}$$

of the form (18). This system is equivalent to the P_2^1 equation for $y(x) = f(x)$.

Remark 2. *The system is also equivalent to the equation*

$$w'' = \frac{1}{2}(w' - a)w^{-1}(w' + a) + 2w^2 - xw,$$

for the variable $w(x) = g(x)$. In the scalar case it coincides with the P_{34} equation

$$w'' = \frac{1}{2}((w')^2 - a^2)w^{-1} + 2w^2 - xw, \quad a \in \mathbb{C}.$$

3.3 Case $\alpha = 0, \beta = -3$

Consider the transformations (14), (16) padded with formulas

$$h_2 = a - \frac{1}{8}\varepsilon^{-2}b - \frac{1}{2}\gamma_1\mathbb{I}, \quad h_1 = -\frac{8}{3}\varepsilon b, \quad \gamma_1 = \frac{1}{8}\varepsilon^{-6}, \quad \kappa_6 = \frac{1}{4}\varepsilon^{-2}\mathbb{I}, \quad a, b \in \text{Mat}_n(\mathbb{C}),$$

and with the shift $g \mapsto g + \frac{2}{3}b$. Then the pair (13) in the limit $\varepsilon \rightarrow 0$ takes the form

$$\begin{aligned} B(x, \lambda) &= \begin{pmatrix} 0 & \mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} 0 & -\frac{1}{3}b \\ -\mathbb{I} & 2f \end{pmatrix}, \\ A(x, \lambda) &= \begin{pmatrix} 0 & 2\mathbb{I} \\ 0 & 0 \end{pmatrix} \lambda + \begin{pmatrix} -2f & 2f^2 - g + x\mathbb{I} + \frac{2}{3}b \\ -2\mathbb{I} & 2f \end{pmatrix} \\ &\quad + \begin{pmatrix} \frac{2}{3}bf + \frac{1}{2}a & \frac{1}{3}bg - \frac{2}{3}bf^2 - \frac{1}{3}xb - \frac{4}{9}b^2 \\ -g + \frac{2}{3}b & -fg + gf - \frac{4}{3}bf + \frac{2}{3}fb - \frac{1}{2}a \end{pmatrix} \lambda^{-1}. \end{aligned}$$

The relation (4) is equivalent to the system

$$\begin{cases} f' &= -f^2 + g - \frac{1}{2}x\mathbb{I} - b, \\ g' &= 3fg - gf - 2[f, b] + a, \end{cases} \quad [b, a] = 2b,$$

of the form (18) corresponding to the \mathbf{P}_2^2 equation for $y(x) = f(x)$.

Thus, we found the Lax pairs for matrix Painlevé equations $\mathbf{P}_2^0 - \mathbf{P}_2^2$. Our pairs are gauge equivalent to those presented in [1]. In particular, the Irfan pair [13, 1]

$$B(x, \mu) = \begin{pmatrix} \mu\mathbb{I} & -f \\ -f & -\mu\mathbb{I} \end{pmatrix},$$

$$A(x, \mu) = -4B(x, \mu)\mu + \begin{pmatrix} 2f^2 + x\mathbb{I} + 2b & -2f^2 + 2g - x\mathbb{I} - 2b \\ 2f^2 - 2g + x\mathbb{I} + 2b & -2f^2 - x\mathbb{I} - 2b \end{pmatrix} + \left(\theta - \frac{1}{2}\right) \begin{pmatrix} 0 & \mathbb{I} \\ \mathbb{I} & 0 \end{pmatrix} \mu^{-1}$$

for system \mathbf{P}_2^0 (20) can be reduced to (19) by conjugation with the matrix $g = \begin{pmatrix} \mu^{-\frac{1}{2}}\mathbb{I} & -\mu^{\frac{1}{2}}\mathbb{I} \\ \mu^{-\frac{1}{2}}\mathbb{I} & \mu^{\frac{1}{2}}\mathbb{I} \end{pmatrix}$ (the so-called Fabri transform, see [15]), where $\lambda = -\mu^2$.

Using the Fabri transformations it is not difficult to convert our pairs to pairs of the Flashka-Newell type [7]. However, they look more cumbersome.

4 Conclusion

4.1 \mathbf{P}_4 -equation with non-commutative independent variable

The systems $\mathbf{P}_4^0 - \mathbf{P}_4^2$ as well as their Lax pairs found in Section 2 are polynomial and therefore make sense if we replace the matrix variables with elements of an associative unital algebra \mathcal{A} over \mathbb{C} . In the case of systems $\mathbf{P}_4^0, \mathbf{P}_4^1$ we can take for this algebra the trivial central extension of the free associative algebra with generators u, v , and h_i by the element z . For system \mathbf{P}_4^2 the algebra is generated by z, u, v, h_1, h_2 and by the relation $[h_2, h_1] = -2h_1$.

In this approach a system (3) is replaced by the corresponding derivation D of \mathcal{A} , which acts on the generators as

$$D(b_i) = D(c_i) = 0, \quad D(u) = -u^2 + 2uv + \alpha(uv - vu) - 2zu + b_1u + ub_2 + b_3v + vb_4 + b_5,$$

$$D(v) = -v^2 + 2vu + \beta(vu - uv) + 2zv + c_1v + vc_2 + c_3u + uc_4 + c_5, \quad D(z) = 1.$$

Formula (4) means that $D(A) = B_\zeta + [B, A]$. It would be interesting to find additional commutator relations that can be imposed on the generators of \mathcal{A} in the case of systems $P_4^0 - P_4^2$ [18, 17, 12].

For systems of Painlevé type with non-commutative independent variable (cf. [19]) the algebra \mathcal{A} is the free associative algebra with generators \bar{z} , u , and v . Since in such models we have $D(\bar{z}) = 1$, in the matrix representation the non-commutative variable \bar{z} may be replaced by $z\mathbb{I} + b$, where z is the commutative independent variable and b is an arbitrary constant matrix. Therefore, any system with the non-commutative \bar{z} that has a Lax representation (4) generates the corresponding system with commutative z . The converse is not always true. However, if in the pair (10) we add the term $(-\zeta - z)\mathbb{I}$ to B and $(\zeta - z)\mathbb{I}$ to A by a shift (6), the result can be written as

$$B(\bar{z}, \zeta) = \begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix} \zeta + \begin{pmatrix} v - 2\bar{z} & uv + \gamma_1 \\ 1 & 0 \end{pmatrix},$$

$$A(\bar{z}, \zeta) = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix} \zeta + \begin{pmatrix} 0 & -uv - \gamma_1 \\ -1 & -2\bar{z} \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}\gamma_2 & -uvu - \gamma_2u \\ v & -vu - \frac{1}{2}\gamma_2 \end{pmatrix} \zeta^{-1},$$

where $\bar{z} = z\mathbb{I} - \frac{1}{2}h_1$. This is the Lax pair for the P_4^0 system

$$\begin{cases} u' &= -u^2 + uv + vu - 2\bar{z}u + \gamma_1, \\ v' &= -v^2 + vu + uv + 2v\bar{z} + \gamma_2, \end{cases} \quad \gamma_1, \gamma_2 \in \mathbb{C},$$

with non-commutative independent variable \bar{z} .

For slightly more general system [4]

$$\begin{cases} u' &= -u^2 + uv + vu + (k-2)\bar{z}u - k u\bar{z} + \gamma_1, \\ v' &= -v^2 + vu + uv + k\bar{z}v - (k-2)v\bar{z} + \gamma_2, \end{cases}$$

where $k \in \mathbb{C}$, the Lax pair has the form

$$B(\bar{z}, \zeta) = \begin{pmatrix} 0 & 0 \\ 0 & -2 \end{pmatrix} \zeta + \begin{pmatrix} v + (k-2)\bar{z} & uv + \gamma_1 \\ 1 & k\bar{z} \end{pmatrix},$$

$$A(\bar{z}, \zeta) = \begin{pmatrix} 0 & 0 \\ 0 & 2 \end{pmatrix} \zeta + \begin{pmatrix} 0 & -uv - \gamma_1 \\ -1 & -2\bar{z} \end{pmatrix} + \frac{1}{2} \begin{pmatrix} uv + \frac{1}{2}\gamma_2 & -uvu - \gamma_2u \\ v & -vu - \frac{1}{2}\gamma_2 \end{pmatrix} \zeta^{-1}.$$

4.2 Outlook

Possibly, our approach can be applied to the classification of matrix polynomial systems of P_6 -type. It is natural to expect that the limiting transitions (see Section 3) relate them to other matrix Painlevé type systems. Besides, one can try to generalize recent results related to matrix P_1 and P_2 equations [9, 10, 12, 6, 5]. In these papers hierarchies of higher matrix Painlevé equations, non-commutative monodromy surfaces (for a matrix P_2 -equation there exist five non-commutative monodromy surfaces [3]), auto-Bäcklund transformations, and non-commutative orthogonal polynomials are investigated.

Acknowledgements

The authors are grateful to V. Adler, V. Poberezhny and V. Roubtsov for useful discussions. The research of the second author was carried out under the State Assignment 0029-2021-0004 (Quantum field theory) of the Ministry of Science and Higher Education of the Russian Federation. The first author was partially supported by the International Laboratory of Cluster Geometry NRU HSE, RF Government grant № 075-15-2021-608.

References

- [1] V. E. Adler and V. V. Sokolov. On matrix Painlevé II equations. *Theoret. and Math. Phys.*, 207(2):188–201, 2021. [arXiv:2012.05639](#).
- [2] S. P. Balandin and V. V. Sokolov. On the Painlevé test for non-Abelian equations. *Physics letters A*, 246(3-4):267–272, 1998.
- [3] M. Bertola, M. Cafasso, and V. Rubtsov. Noncommutative Painlevé equations and systems of Calogero type. *Communications in Mathematical Physics*, 363(2):503–530, 2018. [arXiv:1710.00736](#).
- [4] I. A. Bobrova and V. V. Sokolov. On matrix Painlevé-4 equations. Part 1: Painlevé-Kovalevskaya test. *arXiv preprint arXiv:2107.11680*, 2021.
- [5] A. Branquinho, A. Moreno, and M. Mañas. Matrix biorthogonal polynomials: Eigenvalue problems and non-Abelian discrete Painlevé equations: a Riemann–Hilbert problem perspective. *Journal of Mathematical Analysis and Applications*, 494(2):124605, 2021.
- [6] M. Cafasso and D. Manuel. Non-commutative Painlevé equations and Hermite-type matrix orthogonal polynomials. *Communications in Mathematical Physics*, 326(2):559–583, 2014.
- [7] H. Flaschka and A. C. Newell. Monodromy- and spectrum-preserving deformations I. *Communications in Mathematical Physics*, 76(1):65–116, 1980.
- [8] B. Gambier. Sur les équations différentielles du second ordre et du premier degré dont l’intégrale générale est à points critiques fixes. *Acta Mathematica*, 33(1):1–55, 1910.
- [9] P. R. Gordoa, A. Pickering, and Z. N. Zhu. Bäcklund transformations for a matrix second Painlevé equation. *Physics Letters A*, 374(34):3422–3424, 2010.
- [10] P. R. Gordoa, A. Pickering, and Z. N. Zhu. On matrix Painlevé hierarchies. *Journal of Differential Equations*, 261(2):1128–1175, 2016.
- [11] J. Harnad, C. A. Tracy, and H. Widom. Hamiltonian structure of equations appearing in random matrices. In *Low-Dimensional Topology and Quantum Field Theory*, pages 231–245. Springer, 1993. [arXiv:9301051](#).
- [12] K. Inamasu and H. Kimura. Matrix hypergeometric functions, semi-classical orthogonal polynomials and quantum Painlevé equations. *Integral Transforms and Special Functions*, 32(5-8):528–544, 2021.
- [13] M. Irfan. Lax pair representation and Darboux transformation of noncommutative Painlevé’s second equation. *Journal of Geometry and Physics*, 62(7):1575–1582, 2012. [arXiv:1201.0900](#).

- [14] M. Jimbo and T. Miwa. Monodromy preserving deformation of linear ordinary differential equations with rational coefficients. II. *Physica D: Nonlinear Phenomena*, 2(3):407–448, 1981.
- [15] N. Joshi, A. V. Kitaev, and P. A. Treharne. On the linearization of the first and second Painlevé equations. *Journal of Physics A: Mathematical and Theoretical*, 42(5):055208, 2009. [arXiv:0806.0271v1](#).
- [16] N. Joshi, A.V. Kitaev, and P.A. Treharne. On the linearization of the Painlevé III–VI equations and reductions of the three-wave resonant system. *Journal of Mathematical Physics*, 48(10):103512, 2007. [arXiv:0706.1750v3](#).
- [17] A. V. Mikhailov. Quantisation ideals of nonabelian integrable systems. *Russian Mathematical Surveys*, 75(5):978–980, 2020. [arXiv:2009.01838](#).
- [18] H. Nagoya, B. Grammaticos, A. Ramani, et al. Quantum Painlevé equations: from Continuous to discrete. *SIGMA. Symmetry, Integrability and Geometry: Methods and Applications*, 4:051, 2008.
- [19] V. S. Retakh and V. V. Rubtsov. Noncommutative Toda Chains, Hankel Quasideterminants and Painlevé II Equation. *Journal of Physics. A, Mathematical and Theoretical*, 43(50):505204, 2010. [arXiv:1007.4168](#).