

Homotopy momentum sections on multisymplectic manifolds

Yuji Hirota^{a *} and Noriaki Ikeda^{b †}

^a*School of Veterinary Medicine, Azabu University,
Sagamihara, Kanagawa 252-5201, Japan*

^b*Department of Mathematical Sciences, Ritsumeikan University
Kusatsu, Shiga 525-8577, Japan*

September 29, 2022

Abstract

We introduce a notion of a homotopy momentum section on a Lie algebroid over a pre-multisymplectic manifold. A homotopy momentum section is a generalization of the momentum map with a Lie group action and the momentum section on a pre-symplectic manifold, and is also a generalization of the homotopy momentum map on a multisymplectic manifold. We show that a gauged nonlinear sigma model with Wess-Zumino term with Lie algebroid gauging has the homotopy momentum section structure.

*E-mail: hirotaATazabu-u.ac.jp

†E-mail: nikedatase.ritsumeikan.ac.jp

Contents

1	Introduction	2
2	Preliminary: Lie algebroid	4
2.1	Lie algebroid	4
2.2	Lie algebroid differential and connection	6
3	Homotopy momentum section	9
3.1	Examples	11
4	Weak homotopy momentum section and equivariant E-differential form	14
5	Relation to homotopy momentum map and weak homotopy momentum map	16
6	Gauged nonlinear sigma model with Wess-Zumino term	19
7	Conclusion and discussion	24
A	Geometry of Lie algebroid	25
B	Supergeometry of Lie algebroid	27

1 Introduction

The moment(um) map is a fundamental object in symplectic geometry inspired by conserved quantities and symmetries in the analytical mechanics. In many cases, for a Lie group action on a symplectic manifold, there exists a corresponding function called a Hamiltonian function. Such a space is called a Hamiltonian G -space and the geometric structure is described by the momentum map theory.

In this paper, a generalization of a momentum map theory on a multisymplectic manifold is discussed. There are many versions of a momentum map theory on a multisymplectic manifold, such as a momentum map differential form [11, 20], a multimomentum map [34, 35],

a homotopy momentum map [‡] [18], a weak (homotopy) momentum map [21, 22], etc. Some recent developments about moment maps on multisymplectic manifolds are, for instance, [40, 41].

Recently, Blohmann and Weinstein [5] have proposed a generalization of a momentum map and a Hamiltonian G -space with a Lie algebra (Lie group) action to a Lie algebroid, inspired by the analysis of the Hamiltonian formalism of general relativity [6]. A Lie algebroid [33] is a widely analyzed generalization of a Lie algebra. A Lie algebroid is an infinitesimal object of a Lie groupoid analogous to the fact that a Lie algebra is an infinitesimal object of a Lie group. The generalization of a momentum map is called a momentum section and the corresponding space is called a Hamiltonian Lie algebroid. One of authors gave new examples in the constrained Hamiltonian mechanics and sigma models, and generalized a momentum section theory [27]. Moreover a momentum section has been generalized to Courant algebroid actions [28].

In this paper, we propose a generalization of momentum sections to a pre-multisymplectic manifold and analyze its properties. We call it a homotopy momentum section since it is also a generalization of a homotopy momentum map. A Lie algebroid called a homotopy Hamiltonian Lie algebroid is introduced as a generalization of a Hamiltonian G -space. Moreover a Lie algebroid generalization of a weak momentum map is defined.

We introduce a homotopy momentum section and weak homotopy momentum section in Sections 3 and 4. In Section 5, relations of a homotopy momentum section on an action Lie algebroid with a homotopy momentum map and a weak homotopy momentum map are discussed. If $n = 1$, i.e., a base manifold is a symplectic manifold and a Lie algebroid is the action Lie algebroid, a homotopy momentum section is a momentum map, thus is also equivalent to a homotopy momentum map. If $n \geq 2$, on a multisymplectic manifold, a homotopy momentum section is not necessarily a homotopy momentum map. However, we prove that both are equivalent if a natural condition is satisfied on the action Lie algebroid. It is emphasized that the condition, *equivariance*, of (weak) homotopy momentum sections are important for comparisons. We also discuss the relation between a weak homotopy momentum section and a weak homotopy momentum map on an action Lie algebroid.

In Section 6, we show that a nonlinear sigma model with WZ term in n dimensions

[‡]It is called a *homotopy* moment map in the paper [18].

has a homotopy momentum section structure if we consider gauging with the Lie algebroid gauge symmetry. The nonlinear sigma model is an important physical model which has many physical applications [44]. One of the applications is the T-duality of string theory [9, 10]. If we add the WZ term, the target space has a pre-multisymplectic structure additional to a Riemannian structure. Gauging of the nonlinear sigma model with a Lie algebroid was analyzed in [7, 8, 12, 13, 14, 15, 27, 45]. If we take the Hull-Spence ansatz for gauging of the nonlinear sigma model, and suppose that the gauge symmetry is a Lie algebroid, the gauge invariant condition of the gauged action functional requires existence of a homotopy momentum section on the target pre-multisymplectic manifold.

This paper is organized as follows. In Section 2, a Lie algebroid and related notion are prepared and notation is fixed. In Section 3, a homotopy momentum section is defined and some examples are discussed. In Section 4, a weak homotopy momentum section and equivariance are introduced. In Section 5, relations between a (weak) homotopy momentum section, and a (weak) homotopy momentum map are discussed. In Section 6, the gauged nonlinear sigma model with WZ term is considered and a homotopy momentum section structure in this model is showed. Section 7 is devoted to discussion and outlook. In the Appendix, some formulas are summarized.

2 Preliminary: Lie algebroid

In this section, we explain the background geometry of this paper. Formulas of a Lie algebroid, differentials and connections are discussed.

2.1 Lie algebroid

Definition 2.1 Let E be a vector bundle over a smooth manifold M . A Lie algebroid $(E, \rho, [-, -])$ is a vector bundle E with a bundle map $\rho : E \rightarrow TM$ called the anchor map, and a Lie bracket $[-, -] : \Gamma(E) \times \Gamma(E) \rightarrow \Gamma(E)$ satisfying the Leibniz rule,

$$[e_1, fe_2] = f[e_1, e_2] + \rho(e_1)f \cdot e_2, \quad (1)$$

where $e_i \in \Gamma(E)$ and $f \in C^\infty(M)$.

A Lie algebroid is a generalization of a Lie algebra and the space of vector fields on a smooth manifold.

Example 2.1 (Lie algebra) Let a manifold M be one point $M = \{pt\}$. Then a Lie algebroid is a Lie algebra \mathfrak{g} .

Example 2.2 (Vector fields) If a vector bundle E is a tangent bundle TM and $\rho = \text{id}$, then a bracket $[-, -]$ is a normal Lie bracket on the space of vector fields $\mathfrak{X}(M)$ and $(TM, \text{id}, [-, -])$ is a Lie algebroid. It is called a tangent Lie algebroid.

Example 2.3 (Action Lie algebroid) Suppose a smooth action of a Lie group G to a smooth manifold M , $M \times G \rightarrow M$. The differential of the map induces an infinitesimal action of the Lie algebra \mathfrak{g} of G on a manifold M . Since \mathfrak{g} acts as a differential operator on M , the differential of the map determines a bundle map $\rho : M \times \mathfrak{g} \rightarrow TM$. Consistency of a Lie bracket requires that ρ is a Lie algebra morphism such that

$$[\rho(e_1), \rho(e_2)] = \rho([e_1, e_2]), \quad (2)$$

and we obtain a Lie algebroid $(E = M \times \mathfrak{g}, \rho, [-, -])$. Here the bracket in left hand side of (2) is a Lie bracket of vector fields. This Lie algebroid is called an *action Lie algebroid*.

Example 2.4 (Poisson structure) An important nontrivial Lie algebroid is a Lie algebroid induced from a Poisson structure.

A bivector field $\pi \in \Gamma(\wedge^2 TM)$ is called a Poisson structure if $[\pi, \pi]_S = 0$, where $[-, -]_S$ is a Schouten bracket on the space of multivector fields, $\Gamma(\wedge^\bullet TM)$. For a Poisson bivector field π , a bundle map is defined as $\pi^\sharp : T^*M \rightarrow TM$ by $\langle \pi^\sharp(\alpha), \beta \rangle = \pi(\alpha, \beta)$ for all $\beta \in \Omega^1(M)$.

Let (M, π) be a Poisson manifold. Then, a Lie algebroid structure is induced on T^*M . $\pi^\sharp : T^*M \rightarrow TM$ is the anchor map, and a Lie bracket on $\Omega^1(M)$ is defined by the so called Koszul bracket,

$$[\alpha, \beta]_\pi = \mathcal{L}_{\pi^\sharp(\alpha)}\beta - \mathcal{L}_{\pi^\sharp(\beta)}\alpha - d(\pi(\alpha, \beta)), \quad (3)$$

where $\alpha, \beta \in \Omega^1(M)$. $(T^*M, -\pi^\sharp, [-, -]_\pi)$ is a Lie algebroid.

Example 2.5 (Twisted Poisson structure) If a bivector field $\pi \in \Gamma(\wedge^2 TM)$ and $H \in \Omega^3(M)$ satisfy

$$\frac{1}{2}[\pi, \pi]_S = \langle \otimes^3 \pi, H \rangle, \quad (4)$$

$$dH = 0, \quad (5)$$

(M, π, H) is called a twisted Poisson manifold [31, 39, 42, 30]. Here $\langle \otimes^3 \pi, H \rangle$ is defined by $\langle \otimes^3 \pi, H \rangle(\alpha_1, \alpha_2, \alpha_3) := H(\pi^\sharp(\alpha_1), \pi^\sharp(\alpha_2), \pi^\sharp(\alpha_3))$ for $\alpha_i \in \Omega^1(M)$.

If we take the same bundle map, $\pi^\sharp : T^*M \rightarrow TM$ as in Example 2.4, and a Lie bracket on $\Omega^1(M)$,

$$[\alpha, \beta]_{\pi, H} = \mathcal{L}_{\pi^\sharp(\alpha)}\beta - \mathcal{L}_{\pi^\sharp(\beta)}\alpha - d(\pi(\alpha, \beta)) + \iota_{\pi^\sharp(\alpha)}\iota_{\pi^\sharp(\beta)}H, \quad (6)$$

for $\alpha, \beta \in \Omega^1(M)$. Then, $(T^*M, -\pi^\sharp, [-, -]_{\pi, H})$ is a Lie algebroid.

One can refer to many other examples, for instance, in [33].

2.2 Lie algebroid differential and connection

Consider the spaces of exterior products of sections of E^* called the space of E -differential forms, $\Gamma(\wedge^\bullet E^*)$. A *Lie algebroid differential* ${}^E d : \Gamma(\wedge^m E^*) \rightarrow \Gamma(\wedge^{m+1} E^*)$ such that $({}^E d)^2 = 0$ is defined as follows.

Definition 2.2 A Lie algebroid differential ${}^E d : \Gamma(\wedge^m E^*) \rightarrow \Gamma(\wedge^{m+1} E^*)$ is defined by

$$\begin{aligned} {}^E d\alpha(e_1, \dots, e_{m+1}) &= \sum_{i=1}^{m+1} (-1)^{i-1} \rho(e_i) \alpha(e_1, \dots, \check{e}_i, \dots, e_{m+1}) \\ &+ \sum_{1 \leq i < j \leq m+1} (-1)^{i+j} \alpha([e_i, e_j], e_1, \dots, \check{e}_i, \dots, \check{e}_j, \dots, e_{m+1}), \end{aligned} \quad (7)$$

where $\alpha \in \Gamma(\wedge^m E^*)$ and $e_i \in \Gamma(E)$.[§]

We introduce an ordinary connection on the vector bundle E , $\nabla : \Gamma(E) \rightarrow \Gamma(T^*M \otimes E)$, which is a \mathbb{R} -linear map satisfying the Leibniz rule,

$$\nabla(fe) = f\nabla e + (df) \otimes e, \quad (8)$$

[§]In Equation (7), indices i, j are not indices of local coordinates on M but counting of elements of $\Gamma(E)$.

for $e \in \Gamma(E)$ and $f \in C^\infty(M)$. A dual connection on E^* is defined by the equation,

$$d\langle \mu, e \rangle = \langle \nabla \mu, e \rangle + \langle \mu, \nabla e \rangle, \quad (9)$$

for all sections $\mu \in \Gamma(E^*)$ and $e \in \Gamma(E)$.

The connection and the dual connection are extended to the *exterior covariant derivative* on the space of differential forms taking a value in $E^{\otimes m} \otimes E^{*\otimes n}$, $\Omega^l(M, E^{\otimes m} \otimes E^{*\otimes n}) = \Gamma(M, \wedge^l T^*M \otimes E^{\otimes m} \otimes E^{*\otimes n})$. The exterior covariant derivative is denoted by the same notation ∇ . It increases the degree of the order of $\wedge^l T^*M$ by one. i.e., $\nabla : \Omega^l(M, E^{\otimes m} \otimes E^{*\otimes n}) \rightarrow \Omega^{l+1}(M, E^{\otimes m} \otimes E^{*\otimes n})$.

On a Lie algebroid, another derivation is defined.

Definition 2.3 An *E-connection* on a vector bundle E' with respect to the Lie algebroid E is a \mathbb{R} -linear map ${}^E\nabla : \Gamma(E') \rightarrow \Gamma(E^* \otimes E')$ satisfying

$${}^E\nabla_e(fe') = f{}^E\nabla_e e' + (\rho(e)f)e', \quad (10)$$

for $e \in \Gamma(E)$, $e' \in \Gamma(E')$ and $f \in C^\infty(M)$.

The ordinary connection is regarded as an E -connection for $E = TM$, $\nabla = {}^{TM}\nabla$.

The *standard E-connection* on E , ${}^E\nabla : \Gamma(E) \rightarrow \Gamma(E^* \otimes E)$ is defined by

$${}^E\nabla_e e' := \nabla_{\rho(e)} e', \quad (11)$$

for $e, e' \in \Gamma(E)$. It is also denoted by $\rho^\nabla(e)e' = {}^E\nabla_e e'$ and called the covariantized anchor map. For the tangent bundle $E' = TM$, there is the following canonical E -connection on TM . If a normal connection ∇ on E as a vector bundle is given, a (canonical) E -connection on a tangent bundle, ${}^E\nabla : \Gamma(TM) \rightarrow \Gamma(E^* \otimes TM)$ is defined by

$${}^E\nabla_e v := \mathcal{L}_{\rho(e)} v + \rho(\nabla_v e) = [\rho(e), v] + \rho(\nabla_v e), \quad (12)$$

where $e \in \Gamma(E)$ and $v \in \mathfrak{X}(M)$. It is also called the opposite connection. For a 1-form $\alpha \in \Omega^1(M)$, the E -connection is given by

$${}^E\nabla_e \alpha := \mathcal{L}_{\rho(e)} \alpha + \langle \rho(\nabla e), \alpha \rangle. \quad (13)$$

The E -connection on a k -form $\alpha \in \Omega^k(M) = \Gamma(\wedge^k T^*M)$, ${}^E\nabla : \Gamma(\wedge^k T^*M) \rightarrow \Gamma(E^* \otimes \wedge^k T^*M)$ is given by applying the Leibniz rule recursively,

$${}^E\nabla_e(\beta\gamma) = ({}^E\nabla_e\beta)\gamma - \beta({}^E\nabla_e\gamma), \quad (14)$$

for $\beta \in \Omega^1(M)$ and $\gamma \in \Omega^{k-1}(M)$. Throughout our paper, the canonical E -connection is used as the E -connection unless it is written.

Given an E -connection ${}^E\nabla$, the Lie algebroid differential ${}^E d$ can be generalized to the operation on the space $\Gamma(\wedge^k E' \otimes \wedge^m E^*)$, ${}^E d^\nabla : \Gamma(\wedge^k E' \otimes \wedge^m E^*) \rightarrow \Gamma(\wedge^k E' \otimes \wedge^{m+1} E^*)$. For instance, see [1]. We set $E' = T^*M$ in the general formula since only this case appears in our paper. The concrete equation is as follows.

Definition 2.4 For $\Omega^k(M, \wedge^m E^*) = \Gamma(\wedge^k T^*M \otimes \wedge^m E^*)$, the E -exterior covariant derivative ${}^E d^\nabla : \Omega^k(M, \wedge^m E^*) \rightarrow \Omega^k(M, \wedge^{m+1} E^*)$ is defined by

$$\begin{aligned} {}^E d^\nabla \alpha(e_1, \dots, e_{m+1}) &:= \sum_{i=1}^{m+1} (-1)^{i-1} {}^E\nabla_{e_i}(\alpha(e_1, \dots, \check{e}_i, \dots, e_{m+1})) \\ &\quad + \sum_{1 \leq i < j \leq m+1} (-1)^{i+j} \alpha([e_i, e_j], e_1, \dots, \check{e}_i, \dots, \check{e}_j, \dots, e_{m+1}), \end{aligned} \quad (15)$$

for $\alpha \in \Omega^k(M, \wedge^m E^*)$ and $e_i \in \Gamma(E)$.

Note that the E -exterior covariant derivative increases the order of $\wedge^m E^*$.

In Section 6, we use the following E -covariant derivative,

$${}^E\nabla_e \alpha(e_1, \dots, e_m) = (\mathcal{L}_{\rho(e)}^\nabla \alpha)(e_1, \dots, e_m), \quad (16)$$

where

$$(\mathcal{L}_{\rho(e)}^\nabla \alpha) = (\iota_{\rho(e)} \nabla + \nabla \iota_{\rho(e)}) \alpha. \quad (17)$$

is the covariantized Lie derivative. Comparing to Eq. (7), \mathcal{L}^∇ is also regarded as an extension of the anchor map ρ . We also use the following notation for the covariantized exterior anchor map as $\rho^\nabla : \Gamma(E) \times \Omega^k(M, \wedge^m E^*) \rightarrow \Omega^k(M, \wedge^m E^*)$, $\rho^\nabla : (e, \alpha) \mapsto \rho^\nabla(e)\alpha$,

$$(\rho^\nabla(e)\alpha)(e_1, \dots, e_m) = (\mathcal{L}_{\rho(e)}^\nabla \alpha)(e_1, \dots, e_m), \quad (18)$$

for $e \in \Gamma(E)$ and $\alpha \in \Omega^k(M, \wedge^m E^*)$. Note that $\rho^\nabla(e)f = \rho(e)f$ for $f \in C^\infty(M)$. Moreover, it is useful to notice the manifestly covariant expression of the Lie bracket on $\Gamma(E)$,

$$[e_1, e_2] = -T(e_1, e_2) + \rho^\nabla(e_1)e_2 - \rho^\nabla(e_2)e_1. \quad (19)$$

where $T \in \Gamma(E \oplus \wedge^2 E^*)$ is the E -torsion defined by

$$T(e_1, e_2) := {}^E\nabla_{e_1}e_2 - {}^E\nabla_{e_2}e_1 - [e_1, e_2]. \quad (20)$$

Eq. (18) and the Lie bracket $[-, -]$ satisfy the following covariantized identities of the Lie algebroid,

$$[\rho^\nabla(e_1), \rho^\nabla(e_2)] = \rho^\nabla([e_1, e_2]), \quad (21)$$

$$[e_1, fe_2] = f[e_1, e_2] + \rho^\nabla(e_1)f \cdot e_2. \quad (22)$$

Eq. (22) is easily obtained from $\rho^\nabla(e_1)f = \rho(e_1)f$ and (21) is proved from (22) and the Jacobi identity of the Lie bracket.

A boundary operator on the exterior algebra of the vector bundle E , $\Gamma(\wedge^\bullet E)$, is introduced. Let $e_1 \wedge \dots \wedge e_m \in \Gamma(\wedge^m E)$ be a decomposable element of $\Gamma(\wedge^m E)$ with $e_i \in \Gamma(E)$. Then, a Lie algebroid homology operator is given by

$${}^E\partial_m(e_1 \wedge \dots \wedge e_m) = \sum_{1 \leq i < j \leq m} (-1)^{i+j} [e_i, e_j] \wedge e_1 \wedge \dots \wedge \check{e}_i \wedge \dots \wedge \check{e}_j \wedge \dots \wedge e_m, \quad (23)$$

for $m > 1$. Here $\Gamma(\wedge^0 E) \simeq C^\infty(M)$. The corresponding homology operator on $\Gamma(\wedge^\bullet E)$ is defined by ${}^E\partial = \sum_{k=1}^{\text{rk}E} \partial_k$. The Lie algebroid homology operator ${}^E\partial$ satisfies $({}^E\partial)^2 = 0$.

One can refer to [1, 17, 19] about the general theory of connections on a Lie algebroid. Local coordinate expressions of formulas in a Lie algebroid are summarized in the Appendix.

3 Homotopy momentum section

In this section, we introduce a homotopy momentum section on a pre-multisymplectic manifold.

Definition 3.1 A pre- n -plectic manifold is a pair (M, ω) , where M is a smooth manifold and ω is a closed $(n+1)$ -form on M .

A pre- n -plectic manifold is also called a pre-multisymplectic manifold for $n \geq 2$. If ω is nondegenerate, i.e., if $\iota_v \omega = 0$ for a vector field $v \in \mathfrak{X}(M)$ implies $v = 0$, a pre- n -plectic manifold is called an n -plectic manifold. A pre-1-plectic manifold is a pre-symplectic manifold and a 1-plectic manifold is nothing but a symplectic manifold.

We introduce a series of k -forms taking values in $\wedge^{n-k} E^*$, $\mu_k \in \Omega^k(M, \wedge^{n-k} E^*)$, where $k = 0, \dots, n-1$. Recall two derivations introduced in Section 2.2. The exterior covariant derivative ∇ and the E -exterior covariant derivative ${}^E d^\nabla$ act on the space $\Omega^k(M, \wedge^{n-k} E^*)$.

Definition 3.2 A formal sum $\mu = \sum_{k=0}^{n-1} \mu_k$ is called a *homotopy momentum section* if μ satisfies

$$(\nabla + {}^E d^\nabla)\mu = -\sum_{k=0}^n \iota_\rho^{n+1-k} \omega. \quad (24)$$

Here, $\iota_\rho^k \omega \in \Omega^{n-k}(M, \wedge^k E^*)$ is given by

$$\begin{aligned} \iota_\rho^k \omega(v_{k+1}, \dots, v_{n+1})(e_1, \dots, e_k) &= \iota_{\rho(e_1)} \dots \iota_{\rho(e_k)} \omega(v_{k+1}, \dots, v_{n+1}) \\ &:= \omega(\rho(e_k), \dots, \rho(e_1), v_{k+1}, \dots, v_{n+1}), \end{aligned} \quad (25)$$

for $e_1, \dots, e_k \in \Gamma(E)$ and $v_{k+1}, \dots, v_{n+1} \in \mathfrak{X}(M)$.

Expanding the equation by form degree of both sides, Equation (24) is the following $n+1$ equations,

$$\nabla \mu_{n-1} = -\iota_\rho^1 \omega, \quad (26)$$

$$\nabla \mu_{n-2} + {}^E d^\nabla \mu_{n-1} = -\iota_\rho^2 \omega, \quad (27)$$

\vdots

$$\nabla \mu_{k-1} + {}^E d^\nabla \mu_k = -\iota_\rho^{n+1-k} \omega, \quad (28)$$

\vdots

$${}^E d^\nabla \mu_0 = {}^E d \mu_0 = -\iota_\rho^{n+1} \omega. \quad (29)$$

In general, neither the square of the operator $\nabla + {}^E d^\nabla$, ∇ , nor ${}^E d^\nabla$ is zero; i.e., they are not coboundary operators. In this paper, we do not impose the condition such that $\nabla^2 \mu = 0$ for the definition of a homotopy momentum section, since these conditions are not needed for existence of a homotopy momentum section μ .

A homotopy Hamiltonian Lie algebroid is defined as follows.

Definition 3.3 A Lie algebroid E with a connection ∇ and a homotopy momentum section $\mu \in \bigoplus_{k=0}^{n-1} \Omega^k(M, \wedge^{n-k} E^*)$ is called *homotopy Hamiltonian* if Equation (24) and $\nabla \iota_\rho \omega := \nabla \iota_\rho^1 \omega = 0$ are satisfied.

Remark 3.1 If the connection ∇ is flat, i.e., $\nabla^2 = 0$, the E -covariant derivative satisfies $({}^E d^\nabla)^2 = 0$, and $[\nabla, {}^E d^\nabla] = 0$, $(\Omega^\bullet(M, \wedge^\bullet E^*), \nabla, {}^E d^\nabla)$ gives a double complex. However in general, both ∇ and ${}^E d^\nabla$ are not necessarily differentials.

3.1 Examples

In this section, some examples are listed. Many geometric structures are regarded as an example of the homotopy momentum section.

Example 3.1 (Momentum map) A homotopy momentum section is a generalization of the momentum map on a symplectic manifold with a Lie group action.

Let (M, ω) be a symplectic manifold with an action of a Lie group G on M . ω is a symplectic form, i.e., a nondegenerate closed 2-form. As explained in Example 2.3, the Lie group action induces an action Lie algebroid structure on a trivial bundle $E = M \times \mathfrak{g}$ with a Lie algebra \mathfrak{g} of G . The anchor map is induced from the Lie algebra action $\mathfrak{g} \curvearrowright TM$. Since $E = M \times \mathfrak{g}$ is the trivial bundle, we can take the trivial connection $\nabla = d$ with the de Rham differential d . For $\alpha \in \Gamma(\wedge^m E^*) = C^\infty(M, \wedge^m \mathfrak{g}^*)$ and $e_i \in \mathfrak{g}$, the E -exterior covariant derivative becomes

$${}^E d^\nabla \alpha(e_1, \dots, e_{m+1}) = \text{ad}_\rho^* \alpha(e_1, \dots, e_{m+1}) + d_{CE} \alpha(e_1, \dots, e_{m+1}), \quad (30)$$

where ad^* is the action on α induced from the Lie algebra action on M ,

$$\begin{aligned} \text{ad}_\rho^* \alpha(e_1, \dots, e_{m+1}) &:= \sum_{i=1}^{m+1} (-1)^{i-1} \text{ad}_{e_i}^* \alpha(e_1, \dots, \check{e}_i, \dots, e_{m+1}) \\ &= \sum_{i=1}^{m+1} (-1)^{i-1} \rho(e_i) \alpha(e_1, \dots, \check{e}_i, \dots, e_{m+1}), \end{aligned} \quad (31)$$

for $e_i \in \mathfrak{g}$, and d_{CE} is the Chevalley-Eilenberg differential,

$$d_{CE} \alpha(e_1, \dots, e_{m+1}) = \sum_{1 \leq i < j \leq m+1} (-1)^{i+j} \alpha([e_i, e_j], e_1, \dots, \check{e}_i, \dots, \check{e}_j, \dots, e_{m+1}). \quad (32)$$

satisfying $(d_{CE})^2 = 0$.

Take $n = 1$ in the definition of the homotopy momentum section. A 1-plectic form ω is nothing but a symplectic form, and μ has only one component $\mu = \mu_0 \in C^\infty(M, \mathfrak{g}^*)$ which is a function taking a value in \mathfrak{g}^* . Then, Equation (24) reduces to two equations,

$$d\mu_0 = -\iota_\rho\omega, \quad (33)$$

$${}^E d\mu_0 = -\iota_\rho^2\omega. \quad (34)$$

Substituting $e_1, e_2 \in \mathfrak{g}$ in Equation (34), the left hand side is

$${}^E d\mu_0(e_1, e_2) = \rho(e_1)\langle\mu_0, e_2\rangle - \rho(e_2)\langle\mu_0, e_1\rangle - \langle\mu_0, [e_1, e_2]\rangle. \quad (35)$$

Using Equation (33) and $d\mu_0(e) = -\iota_\rho\omega(e)$, we obtain $\iota_{\rho(e_2)}d\mu_0(e_1) = -\iota_{\rho(e_2)}\omega(\rho(e_1))$. Using this equation, the second term of the right hand side in Eq. (35) becomes

$$\rho(e_2)\langle\mu_0, e_1\rangle = \iota_{\rho(e_2)}d\mu_0(e_1) = -\iota_{\rho(e_2)}\omega(\rho(e_1)) = (\iota_\rho^2\omega)(e_1, e_2). \quad (36)$$

Therefore substituting (35) and (36) to (34), we obtain the equation, $\langle\mu_0, [e_1, e_2]\rangle = \rho(e_1)\langle\mu_0, e_2\rangle$. i.e.,

$$\mu_0([e_1, e_2]) = \text{ad}_{e_1}^*\mu_0(e_2). \quad (37)$$

for $e_1, e_2 \in \mathfrak{g}$. Therefore, Equation (24) is equivalent to Equations (33) and (37). Equation (37) means that the Lie algebra action is equivariant on μ_0 . Two equations shows that μ_0 is a momentum map of the Lie group action on M .

Equation (37) can also be written as

$$d_{CE}\mu_0(e_1, e_2) = -(\iota_\rho^2\omega)(e_1, e_2), \quad (38)$$

in view of Equation (33).

Example 3.2 (Momentum map on multisymplectic manifold) Let $n \geq 2$ and (M, ω) be an n -plectic manifold with an n -plectic form ω . Suppose an action of a Lie group G on M like the previous example 3.1. Similar to Example 2.3, the Lie group action induces an action Lie algebroid on a trivial bundle $E = M \times \mathfrak{g}$, where \mathfrak{g} is a Lie algebra of G .

If we set $\mu_k = 0$ for $k = 0, \dots, n-2$, μ has only one component, an $n-1$ form $\mu = \mu_{n-1} \in \Omega^{n-1}(M, \mathfrak{g}^*)$. Since $\nabla = d$ again, Equation (24) reduces to two equations,

$$d\mu_{n-1} = -\iota_\rho\omega. \quad (39)$$

$${}^E d\mu_{n-1} = -\iota_\rho^2\omega. \quad (40)$$

Similar to Example (3.1), Equations (39) and (40) give the equation,

$$\mu_{n-1}([e_1, e_2]) = \text{ad}_{e_1}^* \mu_{n-1}(e_2). \quad (41)$$

for $e_1, e_2 \in \mathfrak{g}$. Equations (39) and (40) are equivalent to Equations (39) and (41). Two equations means that μ_{n-1} is a momentum map of the Lie group action on an n -plectic manifold M [20].

Example 3.3 (Momentum section) We take $n = 1$ and a general Lie algebroid E . Since $n = 1$, the base manifold M is a pre-symplectic manifold. Similar to the momentum map, a homotopy momentum section μ is only one component $\mu = \mu_0 \in \Gamma(E^*)$. Equation (24) reduces to two equations,

$$\nabla\mu_0 = -\iota_\rho\omega, \quad (42)$$

$${}^E d\mu_0 = -\iota_\rho^2\omega. \quad (43)$$

Equations (42) and (43) are equivalent to the definition of a bracket-compatible momentum section on a Lie algebroid over a pre-symplectic manifold introduced by Blohmann and Weinstein [5]. See also [32]. Many examples of momentum sections which are not momentum maps have been discussed in [5]. One can refer to more examples of momentum sections appearing in physical theories in [27].

In some cases, only parts of Equations (24) in the definition of homotopy momentum sections appear. Especially the 0-form part of Equation (29) is important itself.

Example 3.4 (Twisted Poisson structure) We consider a twisted Poisson manifold M in Example 2.5. Then, as explained in Example 2.5, the twisted Poisson structure gives a Lie algebroid structure on T^*M . Using the Lie algebroid differential ${}^E d$ given by this Lie algebroid, Equation (4) of the twisted Poisson structure is rewritten as

$${}^E d\pi = -\iota_\pi^3 H. \quad (44)$$

It means that $\mu_0 = \pi$ satisfies Equation (29) with $H = \omega$ for $n = 2$. Here $\mu_1 = 0$.

Example 3.5 (twisted R -Poisson structure) A twisted R -Poisson manifold (M, π, J, H) [16] is regarded as a higher dimensional generalization of the twisted Poisson structure. Let $\pi \in \Gamma(\wedge^2 TM)$ be a Poisson bivector field on M . i.e., π is a bivector field satisfying $[\pi, \pi]_S = 0$. H is a closed $(n + 1)$ -form, and $J \in \Gamma(\wedge^n TM)$ is an n -multivector field. [¶] The R -Poisson structure (π, J, H) is defined by the triple satisfying the equation

$$[\pi, J]_S = \langle \otimes^{n+1} \pi, H \rangle. \quad (45)$$

The Poisson bivector field π induces a Lie algebroid structure on T^*M as written in Example 2.4. Equation (45) is equivalent to the equation,

$${}^E dJ = -\iota_\pi^{n+1} H, \quad (46)$$

where the Lie algebroid differential ${}^E d$ is one with respect to the Lie algebroid T^*M induced from the Poisson structure. The condition (46) is nothing but Equation (29) with $\omega = H$, $\mu_0 = J$ and $\mu_k = 0$ for $k = 1, \dots, n - 1$. The twisted R -Poisson structure is also regarded as a special case of the homotopy momentum section. Refer to [29] for a generalization of the twisted R -Poisson structure to a general Lie algebroid and a corresponding topological sigma model.

4 Weak homotopy momentum section and equivariant E -differential form

In this section, we analyze the Lie algebroid over an n -plectic manifold in details for $n \geq 2$, and consider a generalization of the weak homotopy momentum map [22]. Moreover we introduce a condition, *equivariant*, which is nontrivial only for $n \geq 2$.

A pairing of $\Gamma(\wedge^\bullet E)$ and $\Gamma(\wedge^\bullet E^*)$, $\langle -, - \rangle : \Gamma(\wedge^\bullet E) \times \Gamma(\wedge^\bullet E^*) \rightarrow \mathbb{R}$ is induced from the natural pairing of E and E^* . Let u_i and u^i be basis on E and E^* . Then, for an E -differential form $\alpha = \frac{1}{m!} \alpha_{a_1 \dots a_m} u^{a_1} \wedge \dots \wedge u^{a_m} \in \Gamma(\wedge^m E^*)$ and a decomposable element of

[¶]An n -multivector field is denoted by J in this paper though it is denoted by R in [16].

$\Gamma(\wedge^\bullet E)$, $e_1 \wedge \dots \wedge e_m = \frac{1}{m!} e_1^{a_1} \dots e_m^{a_m} u_{a_1} \wedge \dots \wedge u_{a_m}$ for $e_i^{a_i} \in \mathbb{R}$, the pairing is given by

$$\alpha(e_1, \dots, e_m) = \langle \alpha, e_1, \dots, e_m \rangle = \frac{1}{m!} \alpha_{a_1 \dots a_m} e_1^{a_1} \dots e_m^{a_m}. \quad (47)$$

An m -th *Lie kernel* $\mathcal{P}_m \subset \Gamma(\wedge^m E)$ for a Lie algebroid is a subset of $\Gamma(\wedge^m E)$ defined by the kernel of ${}^E\partial_m$,

$$\mathcal{P}_m = \{w \in \Gamma(\wedge^m E) | w \in \text{Ker}({}^E\partial_m)\}, \quad (48)$$

where ${}^E\partial_m$ is the Lie algebroid homology operator defined in Equation (23). A (total) Lie kernel is defined by $\mathcal{P} = \bigoplus_{k=1}^{\text{rank} E} \mathcal{P}_k$.

Let $\mu = \sum_{k=0}^{n-1} \mu_k$, where μ_k be a map $\mu_k : \mathcal{P}_{n-k} \rightarrow \Omega^k(M)$. An element $\mu \in \Omega^k(M, \wedge^{n-k} E^*)$ is also regarded as the map $\mu_k : \mathcal{P}_{n-k} \rightarrow \Omega^k(M)$. We define a generalization of a weak homotopy momentum map.

Definition 4.1 Let E be a Lie algebroid over a pre- n -plectic manifold M . $\mu : \mathcal{P} \rightarrow \sum_{k=0}^{n-1} \Omega^k(M)$ is called a *weak homotopy momentum section* if μ satisfies

$$\nabla \mu = - \sum_{k=1}^{n-1} \iota_\rho^{n+1-k} \omega. \quad (49)$$

Next, we introduce the notion of equivariance for sections of $\wedge^m E^*$ as follows:

Definition 4.2 $\alpha \in \Omega^l(M, \wedge^m E^*)$ is called *equivariant* with respect to a Lie algebroid $(E, \rho, [-, -])$ if it satisfies

$${}^E\nabla_e(\alpha(e_1, \dots, e_m)) = \sum_{i=1}^m (-1)^{i-1} \alpha([e, e_i], e_1, \dots, \check{e}_i, \dots, e_m), \quad (50)$$

for $e, e_i \in \Gamma(E)$ ^{||}.

In the action Lie algebroid, the connection is trivial $\nabla = d$. Thus Equation (50) reduces to

$$\rho(e)\alpha(e_1, \dots, e_m) = \sum_{i=1}^m (-1)^{i-1} \alpha([e, e_i], e_1, \dots, \check{e}_i, \dots, e_m). \quad (51)$$

For a momentum map μ , Equation (51) follows from Equation (37) in Example 3.1.

^{||}For $m = 0$, the righthand side of Eq. (50) is understood as zero.

Proposition 4.3 *A momentum map μ on a symplectic manifold M with a Lie group action is equivariant in the sense of Definition 4.2.*

Let M be a general n -plectic manifold and E be a Lie algebroid. Then if α is equivariant, applying Eq. (50), the E -exterior covariant derivative (7) is simplified to

$${}^E d^\nabla \alpha(e_1, \dots, e_{m+1}) = - \sum_{1 \leq i < j \leq m+1} (-1)^{i+j} \alpha([e_i, e_j], e_1, \dots, \check{e}_i, \dots, \check{e}_j, \dots, e_{m+1}). \quad (52)$$

Thus, the following formula holds.

Lemma 4.4 *Let $\alpha \in \Gamma(\wedge^m E^*)$ be equivariant. Then, for a decomposable element of a Lie kernel, $e_1 \wedge \dots \wedge e_{m+1} \in \mathcal{P}_{m+1}$,*

$${}^E d^\nabla \alpha(e_1, \dots, e_{m+1}) = 0. \quad (53)$$

In fact, the left hand side of (53) is Equation (52), and it is zero since $e_1 \wedge \dots \wedge e_m$ is ∂ -closed since it is a decomposable element of the Lie kernel. The lemma holds since general elements of \mathcal{P}_{m+1} are linear combinations of decomposable elements.

Compare to the definition of the homotopy momentum section (24) and the weak homotopy momentum section (49). A homotopy momentum map and a weak homotopy momentum map have been compared in [36]. If $\mu \in \bigoplus_{k=0}^{n-1} \Omega^k(M, \wedge^{n-k} E^*)$ satisfies Equation (24) and is equivariant, μ satisfies Equation (49) from Lemma 4.4. i.e.,

Proposition 4.5 *Let a homotopy momentum section μ be equivariant, i.e., satisfies Eq. (50). Then, μ is a weak homotopy momentum section.*

5 Relation to homotopy momentum map and weak homotopy momentum map

In this section, we discuss the relation between a homotopy momentum section in this paper and a homotopy momentum map in [18]. The correspondence of weak versions is also discussed.

Let (M, ω) be an n -plectic manifold. Suppose an action of a Lie group G on M . The action of G induces the corresponding infinitesimal Lie algebra action on M as vector fields,

$\rho : \mathfrak{g} \rightarrow TM$. We consider two differentials on the space, $\Omega^k(M, \wedge^{n-k} \mathfrak{g}^*) = \Omega^k(M) \otimes \wedge^{n-k} \mathfrak{g}^*$, $d := d \otimes 1$ and $d_{CE} := 1 \otimes d_{CE}$ where d is the de Rham differential on $\Omega^k(M)$ and d_{CE} is the Chevalley-Eilenberg differential on $\wedge^{n-k} \mathfrak{g}^*$. Introduce a k -form taking a value in $\wedge^{n-k} \mathfrak{g}^*$, $\hat{\mu}_k \in \Omega^k(M, \wedge^{n-k} \mathfrak{g}^*)$, where $k = 0, \dots, n-1$, and their formal sum, $\hat{\mu} = \sum_{k=0}^{n-1} \hat{\mu}_k$.

Definition 5.1 A homotopy momentum map $\hat{\mu}$ is defined by **

$$(d + d_{CE})\hat{\mu} = \sum_{k=0}^n (-1)^{n-k+1} \iota_\rho^{n+1-k} \omega. \quad (54)$$

Expanded by the form degree, Equation (54) becomes the following n equations,

$$d\hat{\mu}_{n-1} = -\iota_\rho^1 \omega, \quad (55)$$

$$d\hat{\mu}_{n-2} + d_{CE}\hat{\mu}_{n-1} = \iota_\rho^2 \omega, \quad (56)$$

\vdots

$$d\hat{\mu}_{k-1} + d_{CE}\hat{\mu}_k = (-1)^{n-k+1} \iota_\rho^{n+1-k} \omega, \quad (57)$$

\vdots

$$d_{CE}\hat{\mu}_0 = (-1)^{n+1} \iota_\rho^{n+1} \omega. \quad (58)$$

For $n = 1$, a homotopy momentum map is equivalent to a momentum map. In fact, in this case, ω is a symplectic form and $\hat{\mu} = \hat{\mu}_0 \in C^\infty(M, \mathfrak{g}^*)$ has only one element. Equation (54) reduces to two equations,

$$d\hat{\mu}_0 = -\iota_\rho \omega, \quad (59)$$

$$d_{CE}\hat{\mu}_0 = \iota_\rho^2 \omega. \quad (60)$$

Let $X_e = \rho(e)$ be a vector field on M given by an element e of the Lie algebra \mathfrak{g} . Since $d_{CE}\hat{\mu}_0(e_1, e_2) = -\hat{\mu}_0([e_1, e_2])$ and $\iota_\rho^2 \omega(e_1, e_2) = -\omega(\rho(e_1), \rho(e_2)) = -\omega(X_1, X_2)$, Equation (60) is equivalent to

$$\hat{\mu}_0([e_1, e_2]) = \omega(X_1, X_2), \quad (61)$$

which means that $\hat{\mu}_0$ is an (anti)-homomorphism from \mathfrak{g} to TM . Equations (59) and (61) show that $\hat{\mu}_0$ is a momentum map on the symplectic manifold M . For $n = 1$, both a homotopy

**Sign factors of the equation are different from [18]. Sign factors are arranged to original ones by proper redefinitions.

momentum map and a homotopy momentum section in Example 3.1 (a momentum section) give a momentum map on a symplectic manifold. Therefore, the result is summarized as follows.

Proposition 5.2 *For $n = 1$, M is a symplectic manifold. Then, a homotopy momentum section on an action Lie algebroid $M \times \mathfrak{g}$ is equivalent to a homotopy momentum map and a momentum map on M with the Lie algebra action \mathfrak{g} .*

We consider the $n \geq 2$ case. As we see in Example 3.1, the Lie group action induces an action Lie algebroid structure on the trivial bundle $E = M \times \mathfrak{g}$. Then, the E -exterior covariant derivative becomes ${}^E d^\nabla = \text{ad}_\rho^* + d_{CE}$ from Equation (30).

Suppose that the homotopy momentum section μ on the action Lie algebroid is equivariant. Then, from Equation (52), we obtain ${}^E d^\nabla = -d_{CE}$. Thus, Equation (24) of the homotopy momentum section reduces to

$$(d - d_{CE})\mu = - \sum_{k=0}^n l_\rho^{n+1-k} \omega. \quad (62)$$

If we take $\hat{\mu}_k = (-1)^{n-k+1} \mu_k$, Equation (62) coincides with Equation (55)–(58). Therefore we obtain the following relation between a homotopy momentum section and a homotopy moment map for a Lie algebra action.

Theorem 5.3 *Let $n \geq 2$ and a Lie algebroid be the action Lie algebroid $E = M \times \mathfrak{g}$ with a Lie algebra \mathfrak{g} . Then, If the homotopy momentum section μ for the action Lie algebroid E is equivariant, it is a homotopy momentum map for \mathfrak{g} .*

Next, we discuss relations between a weak homotopy momentum section and a weak homotopy momentum map. Let $\wedge^m \mathfrak{g}$ be an exterior algebra of the Lie algebra \mathfrak{g} . A Lie algebra homology operator $\partial_m : \Gamma(\wedge^m \mathfrak{g}) \rightarrow \Gamma(\wedge^{m-1} \mathfrak{g})$ is defined by

$$\partial_m(e_1 \wedge \dots \wedge e_m) = \sum_{1 \leq i < j \leq m} (-1)^{i+j} ([e_i, e_j] \wedge e_1 \wedge \dots \wedge \check{e}_i \wedge \dots \wedge \check{e}_j \wedge \dots \wedge e_m). \quad (63)$$

for a decomposable element $e_1 \wedge \dots \wedge e_m \in \wedge^m \mathfrak{g}$ with $e_i \in \mathfrak{g}$. The Lie algebra homology operator on $\wedge^\bullet \mathfrak{g}$ is defined by $\partial = \sum_m \partial_m$.

An m -th Lie kernel $\mathcal{P}_m \subset \Gamma(\wedge^m \mathfrak{g})$ for a Lie algebra \mathfrak{g} is defined by the kernel of a Lie algebra homology operator ∂_m ,

$$\mathcal{P}_m = \{w \in \Gamma(\wedge^m \mathfrak{g}) | w \in \text{Ker}(\partial_m)\}, \quad (64)$$

and a (total) Lie kernel is defined by $\mathcal{P} = \bigoplus_{m=1}^{\dim \mathcal{P}} \mathcal{P}_m$.

Consider a map $\widehat{\mu}_k : \mathcal{P} \rightarrow \Omega^k(M)$, where $k = 0, \dots, n-1$. The map $\widehat{\mu}_k : \mathcal{P} \rightarrow \Omega^k(M)$ is regarded as $\widehat{\mu}_k \in \Omega^k(M, \wedge^{n-k} \mathcal{P}^*)$. For the formal sum of $\widehat{\mu}_k$, $\widehat{\mu} = \sum_{k=0}^{n-1} \widehat{\mu}_k$, a weak homotopy momentum map $\widehat{\mu}$ is defined by [21]

$$d\widehat{\mu} = \sum_{k=0}^n (-1)^{n-k+1} \iota_{\rho}^{n+1-k} \omega, \quad (65)$$

A homotopy momentum map $\widehat{\mu}$ is always a weak homotopy momentum map since $d_{CE}\widehat{\mu} = 0$ if the homotopy momentum map $\widehat{\mu}$ is restricted to \mathcal{P} .

We consider a weak homotopy momentum section in the definition 4.1 on the action Lie algebroid. A homotopy momentum section μ is regarded as a map from the Lie kernel \mathcal{P} to $\Omega^\bullet(M)$. A connection is $\nabla = d$ on the action Lie algebroid. Then, the Lie algebroid homology operator ${}^E\partial_m$ in Equation (23) with $\nabla = d$ is the same as the Lie algebra homology operator ∂_m (63). Thus Equation (49) and (65) is equivalent if we set $\widehat{\mu}_k = (-1)^{n-k+1} \mu_k$. Therefore, we obtain

Theorem 5.4 *Let $n \geq 2$ and a Lie algebroid be the action Lie algebroid. Then, if μ is a weak homotopy momentum section, it is a weak homotopy momentum map.*

6 Gauged nonlinear sigma model with Wess-Zumino term

In this section, we show that a homotopy momentum section appears in the Lie algebroid gauging of an n -dimensional nonlinear sigma model with Wess-Zumino (WZ) term. A nonlinear sigma model is a mechanics on the mapping space from an $(n+1)$ -dimensional smooth manifold Ξ with n -dimensional boundary $\Sigma = \partial\Xi$ to a d -dimensional manifold M . Suppose that a target space M is a d -dimensional pre- n -plectic Riemannian manifold with a metric g . Moreover suppose a Lie algebroid E over M . The claim is that if gauge invariance of the action functional is required, we need an additional condition, existence of homotopy momentum sections on E .

Suppose a metric g , and a closed $(n+1)$ -form H , which is a pre- n -plectic form. Next we introduce an $(n+1)$ -dimensional worldvolume Ξ with boundary $\Sigma = \partial\Xi$, and a metric γ on

Σ . Let $X : \Xi \rightarrow M$ be a map from Ξ to M . The action functional of the n dimensional nonlinear sigma model with a Wess-Zumino term is as follows,

$$\begin{aligned}
S &= \int_{\Sigma} \frac{1}{2} X^* g(dX, *dX) + \int_{\Xi} X^* H \\
&= \int_{\Sigma} \frac{1}{2} g_{ij}(X) dX^i \wedge *dX^j + \int_{\Xi} \frac{1}{(n+1)!} H_{i_1 \dots i_{n+1}}(X) dX^{i_1} \wedge \dots \wedge dX^{i_{n+1}} \\
&= \int_{\Sigma} d^n \sigma \frac{1}{2} g_{ij}(X) \sqrt{\gamma} \gamma^{\mu\nu} \partial_{\mu} X^i \partial_{\nu} X^j \\
&\quad + \int_{\Xi} \frac{1}{(n+1)!} d^{n+1} \sigma H_{i_1 \dots i_{n+1}}(X) \epsilon^{M_1 \dots M_{n+1}} \partial_{M_1} X^{i_1} \wedge \dots \wedge \partial_{M_{n+1}} X^{i_{n+1}}, \quad (66)
\end{aligned}$$

where i, j are indices for local coordinates on M , μ, ν, \dots are indices for local coordinates on Σ , M, N are indices for local coordinates on Ξ . σ^M is the local coordinate on Ξ . $X^*g = g(X)$ is a pullback of the metric g and $X^*H = H(X) = \frac{1}{(n+1)!} H_{i_1 \dots i_{n+1}}(X) dX^{i_1} \wedge \dots \wedge dX^{i_{n+1}}$ in the second term is a pullback of the pre- n -plectic form H on M . $*$ is the Hodge star with respect to the metric γ on Σ . $n = 2$ case is most important since it is the string sigma model with NS-flux 3-form H . Mainly we take expressions with local coordinate indices of the target space M like the second expression in Equation (66).

If M has an isometry with respect to a Lie group G action, its infinitesimal Lie algebra \mathfrak{g} acts on M as a set of vector fields given by a homomorphism $\rho : M \times \mathfrak{g} \rightarrow TM$. Then, the action of the Lie algebra on X is given by ρ as

$$\delta X^i = \rho_a^i(X) \epsilon^a, \quad (67)$$

where $\rho := \rho_a^i(x) e^a \partial_i$ for the basis e^a of \mathfrak{g}^* . ϵ^a is a gauge parameter, which is a function on Ξ taking a value in \mathfrak{g} . Since ρ must be a Lie algebra homomorphism, it satisfies Equation (2) in the action Lie algebroid.

Gauge invariance of the action functional S under the transformation (67) imposes the following conditions of H and g ,

$$\mathcal{L}_{\rho(\epsilon)} g = 0, \quad (68)$$

$$\mathcal{L}_{\rho(\epsilon)} H = d\beta(\epsilon), \quad (69)$$

where β is an arbitrary n -form taking a value in E^* .

Next, we consider *gauging* of the n -dimensional sigma model (66) by introducing a connection D on Σ and its connection 1-form A . Gauging is a deformation of the theory by

introducing connections (gauge fields) on the worldvolume. The trivial bundle $E = M \times \mathfrak{g}$ is replaced to a general vector bundle E with the fiber \mathfrak{g} . A Lie algebra structure on \mathfrak{g} , which gives an action Lie algebroid on $E = M \times \mathfrak{g}$, is replaced to a Lie algebroid structure on E .

Introduce a 1-form $A \in \Omega^1(\Sigma, X^*E)$ on Σ taking a value in the pullback of E . Moreover introduce a connection ∇ on E and its connection 1-form $\omega_{ai}^b(x)dx^i \otimes e^a \otimes e_b$. Take local coordinate expressions of the anchor map and the Lie bracket as

$$\rho(e_a) := \rho_a^i(x)\partial_i \quad (70)$$

$$[e_a, e_b] := C_{ab}^c(x)e_c. \quad (71)$$

We consider the following target space covariant gauge transformations, ^{††}

$$\delta X^i = \rho_a^i(X)\epsilon^a, \quad (72)$$

$$\delta A^a = d\epsilon^a + C_{bc}^a(X)A^b\epsilon^c + \omega_{bi}^a(X)\epsilon^b DX^i. \quad (73)$$

Here $DX^i = dX^i - \rho_a^i(X)A^a$ is the covariant derivative of X^i on Σ . In fact, Equations (72) and (73) are covariant under diffeomorphisms and coordinate transformations on the target bundle E [38].

In order to make the action functional to be invariant under gauge transformations (72) and (73), we deform the action functional S (66) adding some terms based on so called following Hull-Spence type ansatz [24, 25],

$$S_{LA} = S_g + S_H + S_\mu, \quad (74)$$

where

$$S_g = \int_\Sigma \frac{1}{2} g_{ij} DX^i \wedge *DX^j \quad (75)$$

$$S_H = \int_\Sigma \frac{1}{(n+1)!} H_{i_1 \dots i_{n+1}}(X) dX^{i_1} \wedge \dots \wedge dX^{i_{n+1}}, \quad (76)$$

$$S_\mu = \int_\Sigma \sum_{k=0}^{n-1} \frac{1}{k!(n-k)!} \tilde{\mu}_{i_1 \dots i_k a_{k+1} \dots a_n}^{(k)}(X) dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n}, \quad (77)$$

where $\tilde{\mu}^{(k)}$ is the pullback of a k -form on M taking a value in $\wedge^{n-k} E^*$, $\tilde{\mu}^{(k)} \in X^* \Omega^k(M, \wedge^{n-k} E^*)$. In our case, the gauge structure is not a Lie algebra but a Lie algebroid. Requiring (74) is

^{††}We can consider more general gauge transformations, for instance, by adding the term $\phi_{bi}^a(X)\epsilon^b *DX^i$ to the right hand side of Equation (73) [13, 14, 15]. We set this term zero in this paper.

invariant under gauge transformations (72) and (73), we obtain geometric conditions for a metric g , H and $\tilde{\mu}^{(k)}$. ^{‡‡}

The condition of the metric g is independent of conditions for H and $\tilde{\mu}^{(k)}$ since $\delta S_g = 0$ is required. We obtain the following condition for g ,

$${}^E\nabla g = 0, \quad (78)$$

where ${}^E\nabla$ is the E -connection induced on $TM \otimes TM$. We can check that if the connection is trivial, it Equation (78) reduces to Equation (68).

Next we fix conditions of H and $\tilde{\mu}^{(k)}$ to make the action functional (74) gauge invariant. This condition is $\delta(S_H + S_\mu) = 0$. The gauge transformation of S_μ is

$$\begin{aligned} \delta S_\mu &= \int_\Sigma \sum_{k=0}^n \frac{1}{k!(n-k)!} \epsilon^b \rho_b^j \partial_j \tilde{\mu}_{i_1 \dots i_k a_{k+1} \dots a_n}^{(k)}(X) dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \epsilon^b \partial_{i_1} \rho_b^j \tilde{\mu}_{j i_2 \dots i_k a_{k+1} \dots a_n}^{(k)}(X) dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=1}^n \frac{1}{(k-1)!(n-k)!} \rho_b^j \tilde{\mu}_{j i_2 \dots i_k a_{k+1} \dots a_n}^{(k)}(X) d\epsilon^b \wedge dX^{i_2} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=0}^{n-1} (-1)^k \frac{1}{k!(n-k-1)!} \tilde{\mu}_{i_1 \dots i_k b a_{k+2} \dots a_n}^{(k)}(X) d\epsilon^b \wedge dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+2}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=0}^{n-1} \frac{1}{k!(n-k-1)!} \tilde{\mu}_{i_1 \dots i_k c a_{k+2} \dots a_n}^{(k)} C_{a_{k+1} b}^c \epsilon^b dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=0}^{n-1} \frac{1}{k!(n-k-1)!} \tilde{\mu}_{i_1 \dots i_k c a_{k+2} \dots a_n}^{(k)} \omega_{b i_{k+1}}^c \epsilon^b dX^{i_1} \wedge \dots \wedge dX^{i_{k+1}} \wedge A^{a_{k+2}} \wedge \dots \wedge A^{a_n} \\ &+ \sum_{k=0}^{n-1} \frac{1}{k!(n-k-1)!} (-1) \tilde{\mu}_{i_1 \dots i_k c a_{k+2} \dots a_n}^{(k)} \omega_{b j}^c \rho_{a_{k+1}}^j \epsilon^b dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n}. \end{aligned} \quad (79)$$

Since $\delta H = \mathcal{L}_{\rho(\epsilon)} H$ from the gauge transformation of X , (72), the gauge transformation of S_H is

$$\begin{aligned} \delta S_H &= \int_\Xi \mathcal{L}_{\rho(\epsilon)} H = \int_\Xi d\iota_{\rho(\epsilon)} H = \int_\Sigma \iota_{\rho(\epsilon)} H \\ &= \frac{1}{n!} \int_\Sigma \epsilon^a \rho_a^j H_{j i_2 \dots i_{n+1}}(X) dX^{i_2} \wedge \dots \wedge dX^{i_{n+1}}, \end{aligned} \quad (80)$$

^{‡‡}For simplicity of notation, we use the same notation for geometric quantities on M and E and their pullbacks by the map X .

where $dH = 0$ and $\partial\Xi = \Sigma$ are used. Since the sum of coefficient functions of Equations (79) and (80) must vanish, we obtain the following conditions for H and $\tilde{\mu}^{(k)}$. From terms including $d\epsilon^b$, we obtain the algebraic condition for $\tilde{\mu}^{(k-1)}$ and $\tilde{\mu}^{(k)}$ for $k = 1, \dots, n-1$,

$$\tilde{\mu}^{(k-1)}(e_k, \dots, e_n) = (-1)^k \iota_{\rho(e_k)} \tilde{\mu}^{(k)}(e_{k+1}, \dots, e_n) + \text{Cycl}(e_k, \dots, e_n). \quad (81)$$

Comparing coefficients of terms, $\epsilon^b dX^{i_1} \wedge \dots \wedge dX^{i_k} \wedge A^{a_{k+1}} \wedge \dots \wedge A^{a_n}$, we obtain differential equations for $\tilde{\mu}^{(k)}$ and H ,

$$\iota_{\rho(e)} H + \mathcal{L}_{\rho(e)} \tilde{\mu}^{(n)} + \langle \omega(e), \tilde{\mu}^{(n-1)} \rangle = 0, \quad (k = n) \quad (82)$$

$$\begin{aligned} & \sum_{l=k+1}^n (-1)^{l-k-1} \mathcal{L}_{\rho(e_l)} \tilde{\mu}^{(k)}(e_{k+1}, \dots, \check{e}_l, \dots, e_n) \\ & - \sum_{k+1 \leq l < m \leq n} (-1)^{l+m-1} \tilde{\mu}^{(k)}([e_l, e_m], e_{k+1}, \dots, \check{e}_l, \dots, \check{e}_m, \dots, e_n) \\ & + \sum_{l=k}^n (-1)^{l-k} \langle \omega(e_l), \tilde{\mu}^{(k-1)} \rangle(e_k, \dots, \check{e}_l, \dots, e_n) \\ & - \sum_{l \neq m} (-1)^{l+m-1} \langle \iota_{\rho(e_m)} \omega(e_l), \tilde{\mu}^{(k)} \rangle(e_{k+1}, \dots, \check{e}_l, \dots, \check{e}_m, \dots, e_n) = 0, \quad (k = 1, \dots, n-1) \end{aligned} \quad (83)$$

$${}^E d\tilde{\mu}^{(0)}(e_1, \dots, e_n) - \sum_{l=1}^n (-1)^{l-1} \langle \iota_{\rho(e_l)} \omega, \tilde{\mu}^{(0)} \rangle(e_1, \dots, \check{e}_l, \dots, e_n) = 0, \quad (k = 0). \quad (84)$$

First we absorb the n -form $\tilde{\mu}^{(n)}$ to H by defining $\tilde{H} = H + d\tilde{\mu}^{(n)}$ in Equation (82), and next, using Equation (81) step by step, Equations (82)–(84) are rearranged to

$$\nabla \tilde{\mu}^{(n-1)} = -(-1)^{n-1} \iota_{\rho} \tilde{H}, \quad (85)$$

$$\nabla \tilde{\mu}^{(n-2)} + (-1)^{n-1} {}^E d^\nabla \tilde{\mu}^{(n-1)} = -(-1)^{(n-2)(n-1)} \iota_{\rho}^2 \tilde{H}, \quad (86)$$

\vdots

$$\nabla \tilde{\mu}^{(k-1)} + (-1)^k {}^E d^\nabla \tilde{\mu}^{(k)} = -(-1)^{(k-1)\dots(n-1)} \iota_{\rho}^{n+1-k} \tilde{H}, \quad (87)$$

\vdots

$${}^E d\tilde{\mu}^{(0)} = -\iota_{\rho}^{n+1} \tilde{H}. \quad (88)$$

Here the E -exterior covariant derivative ${}^E d^\nabla$ in Equations (85)–(88) is given by Eq. (16). If we define

$$\mu_k := (-1)^{(k+1)\dots(n-1)} \tilde{\mu}^{(k)}, \quad (89)$$

and

$$\omega(\rho(e_1), \dots, \rho(e_k), v_{k+1}, \dots, v_{n+1}) := (-1)^{n-k} \tilde{H}(\rho(e_k), \dots, \rho(e_1), v_{k+1}, \dots, v_{n+1}), \quad (90)$$

Equations (85)–(88) become Equations (26)–(29) in the definition of the homotopy momentum section. Here $e_1, \dots, e_k \in \Gamma(E)$ and $v_{k+1}, \dots, v_{n+1} \in \mathfrak{X}(M)$. Equation (81) is

$$\mu_{k-1} = \iota_\rho \mu_k, \quad (91)$$

for $k = 1, \dots, n - 1$. We summarize our result.

Theorem 6.1 *We consider the n -dimensional nonlinear sigma model with $(n+1)$ -dimensional WZ term. If we consider gauging by a Lie algebroid, the theory has a homotopy Hamiltonian Lie algebroid structure.*

i.e. Suppose the gauge action functional S_{LA} is Equation (74), the action functional S_{LA} is gauge invariant if and only if $\mu_k := (-1)^{(k+1)\dots(n-1)} \tilde{\mu}^{(k)}$ are homotopy momentum sections with Equation (91).

7 Conclusion and discussion

We have defined a homotopy momentum section on a Lie algebroid over a pre-multisymplectic manifold. It is a simultaneous generalization of a momentum map on a symplectic manifold, a homotopy momentum map on a pre-multisymplectic manifold, and a momentum section on a symplectic manifold.

We analyzed relations between (weak) homotopy momentum sections and (weak) homotopy momentum maps on an action Lie algebroid. For it, the equivariant condition has been introduced. If μ is equivariant, a homotopy momentum section is a homotopy momentum map. On the other hand, a weak homotopy momentum section is always a weak homotopy momentum map without the equivariance condition.

As an important application, we showed that the n -dimensional gauged nonlinear sigma model with WZ term with Lie algebroid gauging has a homotopy momentum section and a homotopy Hamiltonian Lie algebroid structure.

One of important applications of the momentum map theory is the equivalent cohomology and the symplectic reduction [37]. These important applications are left for future works.

There is a discussion about reductions on a multisymplectic manifold in the Lie group setting [2].

The connection of a homotopy momentum section with the gauged nonlinear sigma model will provide possibility to analyze sigma models, field theories and their quantizations using the homotopy momentum section theory. Physical applications such as string theory can be discussed in this context.

Acknowledgments

The authors would like to thank Aliaksandr Hancharuk, Sylvain Lavau, Thomas Strobl, Alan Weinstein and Marco Zambon for useful discussion and comments. This work was supported by the research promotion program for acquiring grants in-aid for scientific research(KAKENHI) in Ritsumeikan university and JSPS KAKENHI Grant Number 22K03323.

A Geometry of Lie algebroid

We summarize notation, formulas and their local coordinate expressions of geometry of a Lie algebroid.

Let $(E, \rho, [-, -])$ be a Lie algebroid over a smooth manifold M . x^i is a local coordinate on M , $e_a \in \Gamma(E)$ is a basis of sections of E and $e^a \in \Gamma(E^*)$ is a dual basis of sections of E^* . i, j , etc. are indices on M and a, b , etc. are indices on the fiber of E . Local coordinate expressions of the anchor map and the Lie bracket are $\rho(e_a)f = \rho_a^i(x)\partial_i f$ and $[e_a, e_b] = C_{ab}^c(x)e_c$, where $f \in C^\infty(M)$ and $\partial_i = \frac{\partial}{\partial x^i}$. Then, identities of ρ and C induced from the Lie algebroid condition are

$$\rho_a^j \partial_j \rho_b^i - \rho_b^j \partial_j \rho_a^i = C_{ab}^c \rho_c^i, \quad (92)$$

$$C_{ad}^e C_{bc}^d + \rho_a^i \partial_i C_{bc}^e + \text{Cycl}(abc) = 0. \quad (93)$$

Let ∇ be an ordinary connection on the vector bundle E . The canonical E -connection ${}^E\nabla : \Gamma(TM) \rightarrow \Gamma(TM \otimes E^*)$ on the space of vector fields $\mathfrak{X}(M) = \Gamma(TM)$ is defined by

$${}^E\nabla_e v := \mathcal{L}_{\rho(e)}v + \rho(\nabla_v e) = [\rho(e), v] + \rho(\nabla_v e), \quad (94)$$

where $e \in \Gamma(E)$ and $v \in \Gamma(TM)$. The standard E -connection on E is

$${}^E\nabla_e \alpha := \nabla_{\rho(e)} \alpha. \quad (95)$$

for $\alpha \in \Gamma(\wedge^m E^*)$. The dual E -connection ${}^E\nabla$ on the space of differential forms $\Omega^1(M)$ is obtained from the equation,

$${}^E d\langle v, \alpha \rangle = \langle {}^E\nabla_e v, \alpha \rangle + \langle v, {}^E\nabla \alpha \rangle, \quad (96)$$

for a vector field v and a 1-form α . For a 1-form α , it is given by

$${}^E\nabla_e \alpha := \mathcal{L}_{\rho(e)} \alpha + \langle \rho(\nabla e), \alpha \rangle. \quad (97)$$

Let $\omega = \omega_{ai}^b dx^i \otimes e^a \otimes e_b$ be a connection 1-form. For the basis vectors, the covariant derivatives are $\nabla e_a = -\omega_{ai}^b dx^i \otimes e_b$ and $\nabla e^a = \omega_{bi}^a dx^i \otimes e^b$. Local coordinate expressions of covariant derivatives and the standard E -covariant derivative on TM and E are

$$\nabla_i \alpha^a = \partial_i \alpha^a + \omega_{bi}^a \alpha^b, \quad (98)$$

$$\nabla_i \beta_a = \partial_i \beta_a - \omega_{ai}^b \beta_b, \quad (99)$$

$${}^E\nabla_a v^i = \rho_a^j \partial_j v^i - \partial_j \rho_a^i v^j + \rho_b^i \omega_{aj}^b v^j, \quad (100)$$

$${}^E\nabla_a \alpha_i = \rho_a^j \partial_j \alpha_i + \partial_i \rho_a^j \alpha_j - \rho_b^j \omega_{ai}^b \alpha_j. \quad (101)$$

The covariant derivative of an l -form taking a value in $E^{\otimes m} \otimes E^{*\otimes n}$, $\alpha \in \Omega^l(M, E^{\otimes m} \otimes E^{*\otimes n})$ is given by

$$\nabla_j \alpha_{k_1 \dots k_l b_1 \dots b_n}^{a_1 \dots a_m} = \partial_j \alpha_{k_1 \dots k_l b_1 \dots b_n}^{a_1 \dots a_m} + \sum_{i=1}^m \omega_{cj}^{a_i} \alpha_{k_1 \dots k_l b_1 \dots b_n}^{a_1 \dots a_{i-1} c a_{i+1} \dots a_m} - \sum_{i=1}^n \omega_{b_i j}^c \alpha_{k_1 \dots k_l b_1 \dots b_{i-1} c b_{i+1} \dots b_n}^{a_1 \dots a_m}. \quad (102)$$

An E -torsion, a curvature, an E -curvature and a basic curvature, $T \in \Gamma(E \oplus \wedge^2 E^*)$, $R \in \Omega^2(M, E \oplus E^*)$, ${}^E R \in \Gamma(\wedge^2 E^* \oplus E \oplus E^*)$, and $S \in \Omega^1(M, \wedge^2 E^* \oplus E)$, are defined by

$$R(s, s') := [\nabla_s, \nabla_{s'}] - \nabla_{[s, s']}, \quad (103)$$

$$T(s, s') := {}^E\nabla_s s' - {}^E\nabla_{s'} s - [s, s'], \quad (104)$$

$${}^E R(s, s') := [{}^E\nabla_s, {}^E\nabla_{s'}] - {}^E\nabla_{[s, s']}, \quad (105)$$

$$\begin{aligned} S(s, s') &:= \mathcal{L}_s(\nabla s') - \mathcal{L}_{s'}(\nabla s) - \nabla_{\rho(\nabla s)} s' + \nabla_{\rho(\nabla s')} s \\ &\quad - \nabla[s, s'] = (\nabla T + 2\text{Alt } \iota_\rho R)(s, s'), \end{aligned} \quad (106)$$

for $s, s' \in \Gamma(E)$.

B Supergeometry of Lie algebroid

Lie algebroids are described by means of \mathbb{Z} -graded geometry [43]. It is the so called Q -manifold description. A graded manifold $E[1]$ for a vector bundle E are shifted vector bundle spanned by local coordinates x^i , ($i = 1, \dots, \dim M$) on the base manifold M of degree zero, and local coordinates q^a , ($a = 1, \dots, \text{rank } E$) on the fiber of degree one, respectively. Degree one coordinate q^a has the property, $q^a q^b = -q^b q^a$. E -differential forms which are sections of $\wedge^\bullet E^*$ are identified functions on the graded manifold $E[1]$, i.e., $C^\infty(E[1]) \simeq \Gamma(\wedge^\bullet E^*)$, where the degree one odd coordinate q^a is identified to the basis e^a of sections of E^* . The differential operator of degree -1 , $\frac{\partial}{\partial q^a}$, is the derivation satisfying $\frac{\partial}{\partial q^a} q^b = \delta_a^b$, which is a linear operator on $C^\infty(E[1])$ satisfying the Leibniz rule.

There exists a degree plus one (odd) vector field Q on $E[1]$:

$$Q = \rho_a^i(x) q^a \frac{\partial}{\partial x^i} - \frac{1}{2} C_{ab}^c(x) q^a q^b \frac{\partial}{\partial q^c}. \quad (107)$$

where $\rho_a^i(x)$ and $C_{ab}^c(x)$ are local functions of x . Then, the odd vector field Q satisfies

$$Q^2 = 0, \quad (108)$$

if and only if ρ, C are the anchor map and the structure function of a Lie algebroid on E . Identifying $C^\infty(E[1]) \simeq \Gamma(\wedge^\bullet E^*)$, Q is the Lie algebroid differential ${}^E d$.

The precise correspondence of Q with ${}^E d$ is as follows. For e^a , the basis of E^* , the map $j : \Gamma(\wedge^\bullet E^*) \rightarrow C^\infty(E[1])$ is given by the map of basis, $j : (x^i, e^a) \mapsto (x^i, q^a)$. The differential ${}^E d$ on $\Gamma(\wedge^\bullet E^*)$ is defined by the pullback using the map j , ${}^E d = j^* Q$.

Note that Q is itself covariant even when no connection is introduced. In fact, introducing a connection we can rewrite Q to the manifestly covariant form as

$$\begin{aligned} Q &= \rho_a^i(x) q^a \left(\frac{\partial}{\partial x^i} - \omega_{bi}^c q^b \frac{\partial}{\partial q^c} \right) + \frac{1}{2} T_{bc}^a(x) q^b q^c \frac{\partial}{\partial q^a} \\ &= \rho_a^i(x) q^a \nabla_i + \frac{1}{2} T_{bc}^a(x) q^b q^c \frac{\partial}{\partial q^a}. \end{aligned} \quad (109)$$

Let $\alpha = \frac{1}{m!} \alpha_{a_1 \dots a_m}(x) e^{a_1} \wedge \dots \wedge e^{a_m} \in \Gamma(\wedge^m E^*)$ be an E -differential form. Then, the supergeometry description is $\underline{\alpha} := j_* \alpha = \frac{1}{m!} \alpha_{a_1 \dots a_m}(x) q^{a_1} \dots q^{a_m}$. The Lie algebroid differential ${}^E d \alpha$ can be calculated by ${}^E d \alpha = j^* Q \underline{\alpha}$.

Next, we consider the Lie algebroid homology operator ${}^E \partial$. Let $u = u^a(x) e_a \in \Gamma(E)$ be a section of E , where e_a is a basis of the fiber. Then, the corresponding graded vector field \underline{u} is

defined by $\underline{u} = u^a(x)\partial/\partial q^a$. Analogous to the Cartan calculus, the following *derived bracket* $[[\underline{u}, Q], \underline{v}] \underline{\alpha}$ for $u, v \in \Gamma(E)$ gives the ordinary Lie bracket taking the pullback to the nonsuper vector bundle E ,

$$\iota_{[u,v]}\alpha = j^*([[u, Q], v] \underline{\alpha}). \quad (110)$$

Therefore, in the super formulation, the Lie algebroid homology operator is calculated as

$${}^E\partial_m(e_1 \wedge \dots \wedge e_m) = j^* \sum_{1 \leq i < j \leq m} (-1)^{i+j} [[\underline{e}_i, Q], \underline{e}_j] \underline{e}_1 \dots \check{\underline{e}}_i \dots \check{\underline{e}}_j \dots \underline{e}_m, \quad (111)$$

where $\underline{e}_1 \dots \check{\underline{e}}_i \dots \check{\underline{e}}_j \dots \underline{e}_m$ in the right hand side is super products of \underline{e}_i . For $f \in C^\infty(M)$, the formula of the Lie algebroid homology operator is

$${}^E\partial_m(fe_1 \wedge \dots \wedge e_m) = f^E\partial_m(e_1 \wedge \dots \wedge e_m) + \sum_{i=1}^m (-1)^{i-1} (\rho(e_i)f)(e_1 \wedge \dots \wedge \check{e}_i \wedge \dots \wedge e_m). \quad (112)$$

References

- [1] C. A. Abad, M. Crainic, Representations up to homotopy of Lie algebroids, J. Reine Angew. Math. 663 (2012), 91-126, arXiv:0901.0319 [math.DG]
- [2] C. Blacker, Reduction of multisymplectic manifolds, Lett. Math. Phys. 111 (2021) 30, arXiv:2002.10062 [math.SG]
- [3] A. D. Blaom. Geometric structures as deformed infinitesimal symmetries, Trans. Amer. Math. Soc. **358** (2006) 3651. arXiv:math/0404313 [math.DG]
- [4] C. Blohmann, The homotopy momentum map of general relativity, arXiv:2103.07670 [math.SG]
- [5] C. Blohmann, and A. Weinstein, Hamiltonian Lie algebroids, [arXiv:1811.11109 [math.SG]].
- [6] C. Blohmann, M. C. B. Fernandes and A. Weinstein, Groupoid symmetry and constraints in general relativity, Commun. Contemp. Math. **15** (2013) 1250061 [arXiv:1003.2857 [math.DG]].

- [7] P. Bouwknegt, M. Bugden, C. Klimcik and K. Wright, Hidden isometry of “T-duality without isometry”, *JHEP* **1708** (2017) 116 [arXiv:1705.09254 [hep-th]].
- [8] M. Bugden, A Tour of T-duality: Geometric and Topological Aspects of T-dualities, [arXiv:1904.03583 [hep-th]].
- [9] T. H. Buscher, A symmetry of the string background field equations, *Phys. Lett. B* **194** (1987) 59.
- [10] T. H. Buscher, Path integral derivation of quantum duality in nonlinear sigma models, *Phys. Lett. B* **201** (1988) 466.
- [11] J. F. Carinena, M. Crampin and L. A. Ibort, On the multisymplectic formalism for first order field theories, *Differ. Geom. Appl.* **1** (1992) 345.
- [12] A. Chatzistavrakidis, A. Deser and L. Jonke, T-duality without isometry via extended gauge symmetries of 2D sigma models, *JHEP* **1601** (2016) 154 [arXiv:1509.01829 [hep-th]].
- [13] A. Chatzistavrakidis, A. Deser, L. Jonke and T. Strobl, Strings in singular spacetimes and their universal gauge theory, *Annales Henri Poincare* **18** (2017) no.8, 2641 doi:10.1007/s00023-017-0580-3 [arXiv:1608.03250 [math-ph]].
- [14] A. Chatzistavrakidis, A. Deser, L. Jonke and T. Strobl, Beyond the standard gauging: gauge symmetries of Dirac sigma models, *JHEP* **1608** (2016) 172 [arXiv:1607.00342 [hep-th]].
- [15] A. Chatzistavrakidis, A. Deser, L. Jonke and T. Strobl, Gauging as constraining: the universal generalised geometry action in two dimensions, *PoS CORFU* **2016** (2017) 087 [arXiv:1705.05007 [hep-th]].
- [16] A. Chatzistavrakidis, Topological field theories induced by twisted R-Poisson structure in any dimension, [arXiv:2106.01067 [hep-th]].
- [17] M. Crainic, R. L. Fernandes, Lectures on integrability of Lie brackets. *Geometry and Topology Monographs* 17 (2011) 1-107, *Geom. Topol. Publ.*, Coventry

- [18] M. Callies, Y. Fregier, C. L. Rogers and M. Zambon, Homotopy moment maps, *Adv. Math.* **303** (2016) 954 [arXiv:1304.2051 [math.DG]].
- [19] J. P. Dufour and N. T. Zung, *Poisson Structures and their normal forms*, Birkhauser, Progress in Mathematics, 242 (2005).
- [20] M. J. Gotay, J. Isenberg, J. E. Marsden, and R. Montgomery, Momentum maps and classical relativistic fields. Part 1: Covariant Field Theory, physics/9801019 [math-ph].
- [21] J. Herman, Noether’s theorem in multisymplectic geometry, *Differential geometry and its applications* 56 (2018) 260-294, arXiv:1705.05818 [math.SG]
- [22] J. Herman, Weak moment maps in multisymplectic geometry, arXiv:1807.01641 [math.SG].
- [23] P. J. Higgins, and K. Mackenzie, Algebraic constructions in the category of Lie algebroids, *J. Algebra* **129** (1990) 194-230.
- [24] C. M. Hull and B. J. Spence, The gauged nonlinear σ model with Wess-Zumino term, *Phys. Lett. B* **232** (1989), 204-210
- [25] C. M. Hull and B. J. Spence, The geometry of the gauged sigma model with Wess-Zumino term, *Nucl. Phys. B* **353** (1991) 379.
- [26] N. Ikeda, Lectures on AKSZ Sigma Models for Physicists, *Noncommutative Geometry and Physics 4, Workshop on Strings, Membranes and Topological Field Theory: 79-169*, 2017, World scientific, Singapore. [arXiv:1204.3714 [hep-th]].
- [27] N. Ikeda, Momentum sections in Hamiltonian mechanics and sigma models, *SIGMA* **15** (2019), 076 [arXiv:1905.02434 [math-ph]].
- [28] N. Ikeda, Momentum section on Courant algebroid and constrained Hamiltonian mechanics, *J. Geom. Phys.* **170** (2021), 104350 [arXiv:2104.12091 [math-ph]].
- [29] N. Ikeda, Higher dimensional Lie algebroid sigma model with WZ term, *Universe* 2021, 7(10), 391, Special Issue “Dualities and Geometry”, [arXiv:2109.02858 [hep-th]].

- [30] N. Ikeda and T. Strobl, BV and BFV for the H-twisted Poisson sigma model, *Annales Henri Poincaré* **22** (2021) no.4, 1267-1316 [arXiv:1912.13511 [hep-th]].
- [31] C. Klimcik and T. Strobl, WZW - Poisson manifolds, *J. Geom. Phys.* **43** (2002), 341-344 [arXiv:math/0104189 [math.SG]].
- [32] A. Kotov and T. Strobl, Lie algebroids, gauge theories, and compatible geometrical structures, *Rev. Math. Phys.* **31** (2018) no.04, 1950015 [arXiv:1603.04490 [math.DG]].
- [33] K. Mackenzie, Lie groupoids and Lie algebroids in differential geometry, LMS Lecture Note Series **124**, Cambridge U. Press, 1987.
- [34] T. B. Madsen and A. Swann, Multi-moment maps, *Adv. Math.* **229** (2012) 2287 [arXiv:1012.2048 [math.DG]].
- [35] T. B. Madsen and A. Swann, Closed forms and multi-moment maps, *Geom. Dedicata* **165** (2013) no.1, 25 [arXiv:1110.6541 [math.DG]].
- [36] L. Mammadova, L. Ryvkin, On the extension problem for weak moment maps, *Homology, Homotopy and Applications* **24** (2022) 143-159,
- [37] J. Marsden, A. Weinstein, Reduction of symplectic manifolds with symmetry, *Reports on Math. Phys.*, 5 (1974), 121-130.
- [38] C. Mayer and T. Strobl, Lie algebroid Yang Mills with matter fields, *J. Geom. Phys.* **59** (2009), 1613-1623 [arXiv:0908.3161 [hep-th]].
- [39] J. S. Park, Topological open p-branes, [arXiv:hep-th/0012141 [hep-th]].
- [40] L. Ryvkin and T. Wurzbacher, Existence and unicity of co-moments in multisymplectic geometry, *Differ. Geom. Appl.* **41** (2015), 1-11
- [41] L. Ryvkin, T. Wurzbacher and M. Zambon, Conserved quantities on multisymplectic manifolds, *Journal Of The Australian Mathematical Society* 108 (2020) 120 - 144
- [42] P. Ševera and A. Weinstein, Poisson geometry with a 3 form background, *Prog. Theor. Phys. Suppl.* **144** (2001), 145-154 [arXiv:math/0107133 [math.SG]].

- [43] A. Vaintrob, Lie algebroids and homological vector fields, *Uspekhi Mat. Nauk*, **52/2 314** (1997) 161.
- [44] J. Wess and B. Zumino, Consequences of anomalous Ward identities, *Phys. Lett. B* **37** (1971), 95-97 doi:10.1016/0370-2693(71)90582-X
- [45] K. Wright, Lie algebroid gauging of non-linear sigma models, *J. Geom. Phys.* **146** (2019) 103490 [arXiv:1905.00659 [math.DG]].