

NON-DEGENERATE KILLING HORIZONS IN ANALYTIC VACUUM SPACETIMES

KLAUS KRÖNCKE AND OLIVER L. PETERSEN

ABSTRACT. We prove a geometric characterization of all possible 4-dimensional real analytic vacuum spacetimes near non-degenerate Killing horizons. It is known that any such horizon admits a canonically induced real analytic Riemannian metric with a Killing vector field of constant length. In this paper we prove the converse statement: Every real analytic Riemannian 3-manifold, which admits a Killing vector field of constant length, gives rise to a unique (up to isometry) real analytic vacuum spacetime near a non-degenerate Killing horizon. Our result applies in particular to bifurcate horizons (e.g. black hole event horizons) and non-degenerate compact Cauchy horizons in real analytic vacuum spacetimes, since classical results of Hawking and Moncrief-Isenberg (and recent generalizations thereof) show that such horizons are non-degenerate Killing horizons.

CONTENTS

1. Introduction	1
1.1. Killing horizon initial data	3
1.2. Main results	4
2. Applications	7
2.1. Bifurcate horizons (e.g. black hole event horizons)	7
2.2. Compact Cauchy horizons	9
3. Asymptotic expansion	12
3.1. Geometric gauge: the null time function	13
3.2. The statement	14
3.3. The first derivative	15
3.4. The higher derivatives	18
4. Geometric uniqueness	29
4.1. Construction of the candidate isometry	29
4.2. Proof of geometric uniqueness	30
5. Existence	31
5.1. Moncrief's existence theorem	31
5.2. Proof of existence	32
References	34

1. INTRODUCTION

A Killing horizon is a lightlike hypersurface in a spacetime with a Killing vector field, such that the Killing vector field is lightlike at the horizon. In this paper, we are interested in the following question:

2020 Mathematics Subject Classification. Primary 53C50; Secondary 35L80; 83C05.

Key words and phrases. Non-degenerate Killing horizon, bifurcate horizon, compact Cauchy horizon, black hole uniqueness.

What geometry could 4-dimensional vacuum spacetimes have in a neighborhood of a non-degenerate Killing horizon?

We answer this question completely, assuming the spacetime metric and the horizon are real analytic, by introducing a new characteristic Cauchy problem for Einstein's vacuum equation on spacetimes with non-degenerate Killing horizons, prescribing data on the horizon. Our main result is (see Theorem 1.7 and Theorem 1.8 below):

The real analytic 4-dimensional vacuum spacetimes with a non-degenerate Killing horizon are near the horizon in one-to-one-correspondence (up to isometry) with the real analytic Riemannian 3-manifolds admitting a Killing vector field of constant length.

The Riemannian metric and its Killing vector field is induced by the geometry of the vacuum spacetime at the horizon and by the spacetime Killing vector field, in a way we describe in the following section. We thus obtain a complete geometric characterization of all possible neighborhoods of non-degenerate Killing horizons in 4-dimensional real analytic vacuum spacetimes.

The simplest examples of real analytic Riemannian 3-manifolds with a Killing vector field of constant length is the product manifold

$$\mathbb{R} \times K,$$

where K is *any* real analytic surface. If $K = S^2$, our result produces the Schwarzschild spacetime with a certain mass, see Example 2.2. Given two such surfaces K_1 and K_2 , our result implies that the resulting vacuum spacetimes are isometric near the horizon if and only if K_1 and K_2 are isometric.

The Riemannian manifold producing the Kerr spacetime is also

$$\mathbb{R} \times S^2,$$

but the Riemannian metric is *not* the product metric. The Killing vector field is parallel to the \mathbb{R} -component, but the orthogonal complement is a *non-integrable* distribution, see Example 2.3. This is in fact very similar to the Taub-NUT spacetime, where the Riemannian manifold is a 3-dimensional Berger sphere, with the Killing vector field generating the S^1 -action, see Example 2.11.

The main novelty in this paper is the identification of the correct *geometric* initial data for the characteristic Cauchy problem for vacuum spacetimes with non-degenerate Killing horizons. The main technical challenge, which takes up most of this paper, is to show that the full asymptotic expansion of the spacetime metric is given by our initial data in a geometric way. We have to work in a gauge allowing to show that if the initial data is isometric, then the resulting asymptotic expansions of the spacetime at the horizon are isometric (in an appropriate sense). Real analyticity then proves that the candidate isometry actually is an isometry.

The gauge we use was introduced by the second author in [Pet21b] and is called the *null time function*. Recall that a natural foliation of a neighborhood near a non-characteristic hypersurface is obtained by considering geodesics normal to the hypersurface. This is not possible for characteristic/lightlike hypersurfaces, since the normal vector field is tangent to the horizon. However, as observed in [Pet21b], there is in fact a canonically given transversal vector field along non-degenerate Killing horizons (or more generally to horizons with constant non-zero surface gravity). The null time function is constructed by considering geodesics emanating from the horizon in this unique transversal direction. This way we may easily identify the foliations for isometric spacetimes and use this observation to construct the candidate isometry when proving the geometric uniqueness.

Classical work by Moncrief [Mon82, Mon84] shows how to produce real analytic vacuum spacetimes from data on the horizon. Moncrief prescribes a function ϕ ,

a one-form $\mathring{\beta}$ and a lightlike metric \mathring{g} at the horizon, which are all assumed to be independent of one coordinate (which will be the Killing direction). It does in general not seem to be possible to determine from his initial data if two such spacetimes will be isometric near the horizon or not. Indeed, we show that one can in fact always put $\mathring{\phi} = 0$, since any other choice just reproduces isometric spacetimes, see Remark 5.3. We use Moncrief's theorem in order to construct local solutions on coordinate patches and then use our geometric uniqueness theorem to glue these solutions together to a solution globally along the horizon.

Our characterization is reminiscent of the local well-posedness of the Cauchy problem in General Relativity [YFB52]:

For every Riemannian manifold equipped with a symmetric 2-tensor satisfying the relativistic constraint equations, there is a unique (up to isometry) local globally hyperbolic vacuum spacetime such that the first and second fundamental form on a Cauchy hypersurface is the given data.

The main difference is that we only construct (analytic) vacuum spacetimes with a non-degenerate Killing horizon. The point is, however, that it is much easier to find Riemannian manifolds with a Killing vector field of constant length, than to solve the relativistic constraint equations.

Let us finally discuss how our results relate to the black hole uniqueness conjecture, saying that the subextremal Kerr spacetime is the only 4-dimensional smooth stationary asymptotically flat non-degenerate vacuum black hole spacetime. Under the assumption of real analyticity, Hawking has proven the conjecture, see [Haw72, HE73]. In the smooth setting, not assuming analyticity, much less is known about stationary vacuum black holes in general, other than that they are non-degenerate Killing horizons. This is a celebrated theorem by Alexakis-Ionescu-Klainerman [AIK10]. We may thus apply our main tool in this paper, Theorem 3.9, which *does not* assume real analyticity and conclude that one in fact can compute the full asymptotic expansion of the metric at the horizon, even without assuming axisymmetry of the black hole. Our results therefore give new information about the geometry of potential counterexamples to the black hole uniqueness conjecture, which conceivably could be an ingredient in ruling them out.

1.1. Killing horizon initial data. Though our main results are proven assuming real analyticity, important intermediate results hold in the smooth category. We therefore introduce all objects assuming only smoothness. Let (M, g) be a smooth spacetime, i.e. a time-oriented connected Lorentzian manifold of dimension $n+1 \geq 2$, with a smooth lightlike hypersurface $\mathcal{H} \subset M$.

Definition 1.1. A smooth Killing vector field W on M , such that

$$W|_{\mathcal{H}}$$

is lightlike and tangent to \mathcal{H} , is called a *horizon Killing vector field* on M . A smooth lightlike hypersurface \mathcal{H} , with a horizon Killing vector field W , is called a *Killing horizon* in M .

Assuming the curvature condition

$$\text{Ric}(W|_{\mathcal{H}}, X) = 0, \tag{1}$$

for all $X \in T\mathcal{H}$, the existence of a horizon Killing vector field implies that the surface gravity is constant, i.e. that there is a constant $\kappa \in \mathbb{R}$, called *surface gravity*, such that

$$\nabla_W W|_{\mathcal{H}} = \kappa W|_{\mathcal{H}}. \tag{2}$$

For a proof of this statement, we refer to [PV21, Rem. 1.9, Lem. B.1].

Definition 1.2. Assume that (1) is satisfied. We say that \mathcal{H} is a *non-degenerate Killing horizon*, with respect to W , if $\kappa \neq 0$.

By replacing W with $-W$ if necessary, we may always assume that $\kappa > 0$:

Assumption 1.3. We assume throughout this paper that $\mathcal{H} \subset M$ is a smooth non-degenerate Killing horizon with $\kappa > 0$.

Note that

$$g(\nabla_X W, Y)|_{\mathcal{H}} = \frac{1}{2} \mathcal{L}_W g(X, Y)|_{\mathcal{H}} - \frac{1}{2} g(W, [X, Y])|_{\mathcal{H}} = 0$$

for all $X, Y \in T\mathcal{H}$, which implies that $\nabla_X W|_{\mathcal{H}}$ is normal to the lightlike hypersurface \mathcal{H} and must hence be a multiple of the lightlike direction $W|_{\mathcal{H}}$ at each point. In other words, there is a unique smooth one-form ω on \mathcal{H} , such that

$$\nabla_X W|_{\mathcal{H}} = \omega(X)W|_{\mathcal{H}}. \quad (3)$$

It immediately follows that

$$\mathcal{L}_{W|_{\mathcal{H}}} \omega = 0. \quad (4)$$

Define the $(0, 2)$ -tensor σ at any $p \in \mathcal{H}$ as

$$\sigma(X, Y) := g(X, Y) + \omega(X)\omega(Y), \quad (5)$$

for all $X, Y \in T_p \mathcal{H}$. Since

$$\omega(W|_{\mathcal{H}}) = \kappa > 0$$

by assumption, we note that σ is positive definite, i.e. σ is a *Riemannian metric* on \mathcal{H} . Let us also introduce the notation

$$V := W|_{\mathcal{H}}. \quad (6)$$

Definition 1.4. We call (\mathcal{H}, σ, V) as defined in (5) and (6) the *induced initial data*.

We have thus shown that each vacuum spacetime with a smooth non-degenerate Killing horizon gives an induced initial data set (\mathcal{H}, σ, V) in a geometric way. Note the following two important properties of σ and V :

- Firstly, we have

$$\mathcal{L}_V \sigma = \mathcal{L}_W g|_{\mathcal{H}} + \mathcal{L}_V \omega \otimes \omega + \omega \otimes \mathcal{L}_V \omega = 0,$$

i.e. V is a Killing vector field with respect to σ .

- Secondly, note that

$$\sigma(V, V) = \kappa^2$$

is constant, i.e. the length of V with respect to σ is constant.

1.2. Main results. We have already discussed *induced initial data* on any non-degenerate Killing horizon in Definition 1.4. Conversely, the following turns out to be the correct notion of *initial data*:

Definition 1.5. Let (\mathcal{H}, σ) be a Riemannian manifold, equipped with a Killing vector field V of constant length, i.e.

$$\mathcal{L}_V \sigma = 0$$

and

$$\sigma(V, V)$$

is constant. We say that (\mathcal{H}, σ, V) is an *initial data set for the characteristic Cauchy problem for vacuum spacetimes with a non-degenerate Killing horizons* or simply an *initial data set* for short.

Note that all constructions above make sense when replacing smooth with real analytic.

Remark 1.6. If a nowhere vanishing vector field V on a manifold \mathcal{H} is a Killing vector field with respect to a Riemannian metric $\hat{\sigma}$, then (\mathcal{H}, σ, V) is an initial data set, with

$$\sigma := \frac{\hat{\sigma}}{\hat{\sigma}(V, V)}.$$

In other words, it suffices to find Riemannian manifolds with nowhere vanishing Killing vector fields in order to solve our constraint equation.

1.2.1. *Existence.* Our first main result says that any 3-dimensional real analytic initial data gives rise to a 4-dimensional real analytic vacuum spacetime with a non-degenerate Killing horizon:

Theorem 1.7. *Let (\mathcal{H}, σ, V) be real analytic initial data set, in the sense of Definition 1.5, where \mathcal{H} is 3-dimensional. Then there is a real analytic 4-dimensional spacetime (M, g) , a real analytic Killing vector field W on M and a real analytic non-degenerate Killing horizon*

$$\iota : \mathcal{H} \hookrightarrow M,$$

with respect to W , such that

- (\mathcal{H}, σ, V) is the induced initial data as in Definition 1.4,
- $d\iota(V) = W|_{\iota(\mathcal{H})}$.

1.2.2. *Geometric uniqueness.* Our second main result says that two real analytic vacuum spacetimes with non-degenerate Killing horizons are isometric, with an isometry respecting the horizon Killing vector fields, precisely when there is a real analytic isometry of the initial data, respecting the Riemannian Killing vector fields:

Theorem 1.8. *Let (M, g) and (\hat{M}, \hat{g}) be real analytic vacuum spacetimes. Assume that*

$$\iota : \mathcal{H} \hookrightarrow M,$$

$$\hat{\iota} : \hat{\mathcal{H}} \hookrightarrow \hat{M}$$

are real analytic non-degenerate Killing horizons with horizon Killing vector fields W and \hat{W} . Let (\mathcal{H}, σ, V) and $(\hat{\mathcal{H}}, \hat{\sigma}, \hat{V})$ be the induced initial data, in the sense of Definition 1.5. If there is a real analytic isometry

$$\varphi : (\mathcal{H}, \sigma) \rightarrow (\hat{\mathcal{H}}, \hat{\sigma}),$$

with real analytic inverse, such that

$$d\varphi(V) = \hat{V},$$

then there are open neighborhoods $\iota(\mathcal{H}) \subset \mathcal{U} \subset M$ and $\hat{\iota}(\hat{\mathcal{H}}) \subset \hat{\mathcal{U}} \subset \hat{M}$ and a real analytic isometry

$$\Phi : (\mathcal{U}, g|_{\mathcal{U}}) \rightarrow (\hat{\mathcal{U}}, g|_{\hat{\mathcal{U}}})$$

with

$$d\Phi(W) = \hat{W},$$

such that

$$\Phi \circ \iota = \hat{\iota} \circ \varphi.$$

Remark 1.9. Theorem 1.8 requires no restriction on the spacetime dimension.

1.2.3. *Motivating the initial data.* Let us give a brief motivation why the initial data (σ, V) are correct for this characteristic Cauchy problem, by showing how the spacetime metric at the horizon and the one-form ω are constructed from (σ, V) . Given an initial data set (\mathcal{H}, σ, V) , note that the quadratic form

$$g(X, Y) := \sigma(X, Y) - \frac{\sigma(X, V)\sigma(Y, V)}{\sigma(V, V)}$$

for any $X, Y \in T_p\mathcal{H}$ is a *lightlike* metric at \mathcal{H} , i.e. $g(X, X) \geq 0$ and

$$g(X, X) = 0 \Leftrightarrow X = V.$$

Moreover, defining

$$\omega(X) := \frac{\sigma(X, V)}{\sqrt{\sigma(V, V)}},$$

we have

$$\sigma(X, Y) = g(X, Y) + \omega(X)\omega(Y).$$

The quadratic form g will be the induced lightlike metric at the horizon in the resulting spacetime, and ω will be the one-form defined by (3).

1.2.4. *Relation to earlier results.* This paper was greatly inspired by remarkable ideas of Moncrief in [Mon82, Mon84]. Moncrief's result [Mon82, Thm. 2] corresponds to our existence theorem Theorem 1.7 when \mathcal{H} is the three-torus T^3 with V is one of the standard coordinate vector fields with closed integral curves. Similarly, his result [Mon84, Thm. 2] corresponds to our existence theorem Theorem 1.7 when \mathcal{H} is the three-sphere S^3 and V is the generator of a fiber in the Hopf fibration

$$S^3 \rightarrow S^2.$$

As already explained, Moncrief does not prove any geometric uniqueness statement. In fact, as can be seen in the proof of Lemma 5.2 below, Moncrief's initial data sometimes give isometric solutions. Heuristically, Moncrief specifies 6 functions on the horizon, but the correct space of initial data turns out to be parametrized by only 5 functions, see Remark 5.3. More precisely, we may set his initial datum $\check{\phi}$ to zero. All other choices of $\check{\phi}$ lead to spacetimes which are isometric to a spacetime obtained with $\check{\phi} = 0$.

It is a classical topic in general relativity to show that horizons in vacuum spacetimes are Killing horizons, under weak assumptions. This line of argument was initiated by Hawking as the main novelty in his black hole uniqueness theorem [Haw72, HE73]. It is now known that any bifurcate horizon in a real analytic vacuum spacetime of any dimension is necessarily a non-degenerate Killing horizon, see the work of Chruściel-Costa [CC08], Moncrief-Isenberg [MI08] and Hollands-Ishibashi-Wald [HIW07]. In particular, no assumption on the geometry away from the event horizon is needed. As a parallel to this, Moncrief and Isenberg have shown that any real analytic non-degenerate (non-ergodic) compact Cauchy horizon is a Killing horizon [MI83, MI20]. The analyticity assumptions in these theorems has been dropped for stationary black holes by Alexakis-Ionescu-Klainerman in [AIK10, IK13] and for compact Cauchy horizons by the second author in [Pet21a], assuming constant surface gravity, based on [Pet21b, PR18]. We refer to Section 2 for more details on this.

1.2.5. *Structure of the paper.* We begin by presenting our main applications to bifurcate horizons and compact Cauchy horizons in vacuum spacetimes, see Section 2. We also compute the initial data producing the subextremal Kerr spacetime and compute the spacetimes produced by the simplest compact Riemannian manifolds admitting a Killing vector field of constant length.

Section 3 is the technical core of the paper, where we show how to iteratively compute the full asymptotic expansion of the spacetime metric at a non-degenerate Killing horizon in a smooth vacuum spacetime. Indeed, we do not need real analyticity at this step and we only assume smoothness for potential future applications.

In Section 4 we apply the result on the asymptotic expansion to prove geometric uniqueness. The main point here is of course the analyticity of the spacetime metric.

The paper is concluded with Section 5, where we show that the formal solution given by the asymptotic expansion actually converges if the initial data is real analytic. As one might expect, this is not easy to read off directly from the asymptotic expansion. Combining Moncrief's methods in [Mon82, Mon84], in which he solves a Cauchy-Kowalevski problem for PDE with singular coefficients, with our geometric uniqueness Theorem 1.8, proves our existence theorem, Theorem 1.7. We obtain a real analytic solution, which, as we a posteriori conclude, coincides with the asymptotic expansion at the horizon.

Acknowledgements. We would like to thank Vincent Moncrief for very valuable conversations. The first author is grateful for the support of the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the priority programme 2026 *Geometry at Infinity*. The second author is grateful for the support of the DFG – GZ: LI 3638/1-1, AOBJ: 660735 and the Stanford Mathematics Research Center.

2. APPLICATIONS

There are two important classes of results showing that certain horizons in vacuum spacetimes are non-degenerate Killing horizons:

- The case when \mathcal{H} is a bifurcate horizon. This is the union of two intersecting null hypersurfaces. An important example is the union of a past and future event horizon in, say, the subextremal Kerr spacetime.
- The case when \mathcal{H} is compact. This is precisely when \mathcal{H} is a compact non-degenerate Cauchy horizon in a vacuum spacetime (c.f. [Pet21a, Cor. 2.13]). Important examples include the Misner spacetime and the Taub-NUT spacetime.

The main point is: In both these situations, the horizon Killing vector field has been shown to exist under weak assumptions. We recall here these classical rigidity results and combine them with our main results, Theorem 1.7 and Theorem 1.8.

2.1. Bifurcate horizons (e.g. black hole event horizons). The following theorem is known as *Hawking's local rigidity*, see [Haw72, HE73]. We give a general formulation, which is due to Alexakis-Ionescu-Klainerman, see [AIK10, Thm. 1.1]. The latter only assume smoothness of the horizon, as opposed to real analyticity, but we do not need it here. (The analyticity of the Killing vector field in an analytic vacuum spacetime is standard, by elliptic theory.)

Theorem 2.1 (Hawking, Alexakis-Ionescu-Klainerman). *Given a local, regular, bifurcate, non-expanding horizon in a 4-dimensional real analytic vacuum spacetime, there is a Killing vector field in an open neighborhood of the bifurcation surface, which is lightlike at the horizon and tangent to the horizon. In other words, the horizon is a non-degenerate Killing horizon.*

For the precise definition of the objects involved, we refer to [AIK10, Thm. 1.1]. The non-degeneracy conclusion is not explicitly mentioned in [AIK10, Thm. 1.1], but is explained in e.g. [IK09, p. 38]. See also [IK13] for the generalization to higher dimensions (not requiring analyticity). A bifurcate horizon is the union of two smooth (or real analytic) horizons (like the past and future event horizons). Our

main results, Theorem 1.7 and Theorem 1.8, applies to either of the real analytic parts of a bifurcate horizon.

The most important example of spacetimes with bifurcate horizons are the stationary black holes, in particular the family of subextremal Kerr spacetimes. The subextremal Kerr spacetimes satisfy all our assumptions and we would like to compute the induced initial data

$$(\mathcal{H}, \sigma, V),$$

where \mathcal{H} is the future or past event horizon or (half of) the Cauchy horizon. We start with the simpler special case of non-rotating black holes, the Schwarzschild black hole:

Example 2.2 (The Schwarzschild black hole). Given a mass $m \in (0, \infty)$, the Schwarzschild black hole spacetime is given in Eddington-Finkelstein coordinates by

$$M := (0, \infty)_r \times \mathbb{R}_v \times S^2,$$

equipped with the metric

$$g := 2drdv + \left(\frac{2m}{r} - 1 \right) dv^2 + r^2 g_{S^2}.$$

More precisely, (M, g) is the union of the domain of outer communication ($r > 2m$), the event horizon ($r = 2m$) and the interior of the black hole ($r < 2m$). The event horizon

$$\mathcal{H} := \{2m\} \times \mathbb{R} \times S^2$$

is indeed a real analytic lightlike hypersurface with a horizon Killing vector field

$$W := \partial_v.$$

One easily computes that

$$\omega = \frac{1}{4m} dv|_{\mathcal{H}}.$$

Hence the surface gravity is

$$\kappa = \omega(\partial_v|_{\mathcal{H}}) = \frac{1}{4m}.$$

This verifies that \mathcal{H} is a *non-degenerate* Killing horizon for any $m \in (0, \infty)$. The induced initial data (in the sense of Definition 1.4) on $\mathcal{H} \cong \mathbb{R}_v \times S^2$ is therefore

$$\begin{aligned} \sigma &= \frac{1}{16m^2} dv^2 + 4m^2 g_{S^2}, \\ V &= \partial_v|_{\mathcal{H}} \end{aligned}$$

We are now ready to compute the initial data of the significantly more complicated subextremal Kerr spacetime:

Example 2.3 (The subextremal Kerr black hole). Given a mass $m \in (0, \infty)$ and an angular momentum $a \in \mathbb{R}$, such that $0 < |a| < m$, the subextremal Kerr spacetime is given in Eddington-Finkelstein coordinates by

$$M := (0, \infty)_r \times \mathbb{R}_v \times S_{\phi, \theta}^2,$$

equipped with the metric

$$\begin{aligned} g &= 2(dv - a \sin^2(\theta) d\phi)(dr - a \sin^2(\theta) d\phi) \\ &\quad + \left(\frac{2mr}{r^2 + a^2 \cos^2(\theta)} - 1 \right) (dv - a \sin^2(\theta) d\phi)^2 \\ &\quad + (r^2 + a^2 \cos^2(\theta))(d\theta^2 + \sin^2(\theta) d\phi^2). \end{aligned}$$

Setting $a = 0$, we recover the Schwarzschild metric in the previous example. Let

$$r_{\pm} := m \pm \sqrt{m^2 - a^2}.$$

Since $|a| < m$, note that $r_+ \neq r_-$. Now, (M, g) is the union of the domain of outer communication ($r > r_+$), the future outer event horizon ($r = r_+$) and the interior of the black hole, consisting of (half) the inner future event horizon/future Cauchy horizon ($r = r_-$) and the region beyond ($r < r_-$). Both horizons

$$\mathcal{H}^{\pm} := \{r_{\pm}\} \times \mathbb{R} \times S^2$$

are real analytic Killing horizons, but with *different* horizon Killing vector fields

$$W_{\pm} := \partial_v + \frac{a}{r_{\pm}^2 + a^2} \partial_{\phi},$$

i.e. W_{\pm} is lightlike at \mathcal{H}^{\pm} . The surface gravity at either horizon is given by

$$\kappa_{\pm} = \frac{1}{2} \left(\frac{1}{m} - \frac{1}{r_{\pm}} \right),$$

which is non-zero, since $|a| < m$. The induced lightlike metric is

$$g_{\pm} = \frac{\sin^2(\theta)}{r_{\pm}^2 + a^2 \cos^2(\theta)} ((r_{\pm}^2 + a^2) d\phi - a dv)^2 + (r_{\pm}^2 + a^2 \cos^2(\theta)) d\theta^2,$$

and we have

$$\begin{aligned} \omega_{\pm} = \kappa_{\pm} & \frac{r_{\pm}^2 + a^2}{r_{\pm}^2 + a^2 \cos^2(\theta)} (dv - a \sin^2(\theta) d\phi) \\ & - \frac{r_{\pm} a \sin^2(\theta)}{(r_{\pm}^2 + a^2 \cos^2(\theta))^2} ((r_{\pm}^2 + a^2) d\phi - a dv) \\ & - \frac{1}{2} \frac{a^2 \sin(2\theta)}{r_{\pm}^2 + a^2 \cos^2(\theta)} d\theta. \end{aligned}$$

The induced initial data (in the sense of Definition 1.4) on the horizons

$$\mathcal{H}^{\pm} \cong \mathbb{R}_v \times S_{\phi, \theta}^2$$

are

$$\begin{aligned} \sigma_{\pm} &= g_{\pm} + \omega_{\pm} \otimes \omega_{\pm}, \\ V_{\pm} &= \partial_v|_{\mathcal{H}^{\pm}} + \frac{a}{r_{\pm}^2 + a^2} \partial_{\phi}|_{\mathcal{H}^{\pm}}, \end{aligned}$$

where g_{\pm} and ω_{\pm} are given by the expressions above.

2.2. Compact Cauchy horizons. Whereas black hole spacetimes are often assumed to be asymptotically flat, and supposed to model isolated systems, compact Cauchy horizons are naturally associated with *cosmological* spacetimes, where the Cauchy hypersurfaces are compact.

Let us recall the definition of a Cauchy horizon and the properties *compact* Cauchy horizons are known to have. Let M be a real analytic spacetime of dimension $n + 1 \geq 2$ and let $\Sigma \subset M$ be a closed acausal C^0 -hypersurface of M . It is proven in [O'N83, Prop. 14.53] that the domain of dependence

$$D(\Sigma) := \{p \in M \mid \text{any inextendible causal curve through } p \text{ intersects } \Sigma\} \subseteq M$$

is a globally hyperbolic open submanifold of M and that the boundary

$$\partial D(\Sigma) = \mathcal{H}_- \cup \mathcal{H}_+$$

has two connected components \mathcal{H}_- and \mathcal{H}_+ (one past and one future component) which are achronal lightlike C^0 -hypersurfaces in M .

Definition 2.4. We call $\mathcal{H}_{-/ +}$ the past/future Cauchy horizon with respect to Σ .

Assumption 2.5. We assume in this section that \mathcal{H} is a future or past Cauchy horizon, which is *compact* (without boundary).

The following recent theorem was proven independently by Larsson in [Lar15, Cor. 1.43] and Minguzzi in [Min15, Thm. 18]:

Theorem 2.6 (Larsson, Minguzzi). *If M is a real analytic spacetime satisfying the null energy condition (for example vacuum), then \mathcal{H} is a real analytic lightlike submanifold of M .*

In 1983, Moncrief and Isenberg conjectured that any compact Cauchy horizon in a smooth vacuum spacetime is in fact a Killing horizon [MI83]. Under the assumption of real analyticity, Moncrief-Isenberg has done remarkable work on their conjecture throughout the last decades in [IM85, IM92, MI83, MI08], which can by now be summarized in the following theorem:

Theorem 2.7 (Moncrief-Isenberg). *Let \mathcal{H} be a non-degenerate compact Cauchy horizon in a 4-dimensional real analytic vacuum spacetime. Then \mathcal{H} is a non-degenerate real analytic Killing horizon in the sense of Definition 1.1.*

Non-degeneracy here means that at least one generator (lightlike geodesics on the horizon) is incomplete. Recent work by Reiris-Bustamente [RB21] (see also the work by Gurriaran-Minguzzi GM2021) show that compact non-degenerate Cauchy horizons in vacuum spacetimes have constant non-zero surface gravity, even without assuming real analyticity. Strictly speaking, Moncrief and Isenberg did not treat the “ergodic case”, but this was dealt with by the second author of this paper in [Pet21a, Thm. 1.4]. In [Pet21a] the smooth setting is considered, but Killing vector fields in real analytic spacetimes are automatically real analytic by elliptic regularity theory.

Combining the above results with the main results of this paper, we arrive at the following:

Theorem 2.8. *Non-degenerate compact Cauchy horizons in 4-dimensional real analytic vacuum spacetimes M , are in one-to-one correspondence (as in Theorem 1.7 and Theorem 1.8) with real analytic closed Riemannian 3-manifolds (\mathcal{H}, σ) , equipped with a Killing vector field V of constant length.*

Proof. After applying Theorem 1.7 and Theorem 1.8, what remains to check is that the horizon indeed is a Cauchy horizon, but this is proven in [Pet21a, Cor. 2.13]. \square

There are of course many examples of real analytic compact Riemannian manifolds with Killing vector fields of constant length. We refer to the recent work of Bustamente-Reiris [BR21] for a detailed analysis of this. Some particularly simple ones are

$$T^3, \quad S^1 \times S^2, \quad S^3,$$

with the standard metrics and standard nowhere vanishing Killing vector fields. In all these examples, the corresponding vacuum spacetime and Killing horizons are explicitly known. We begin by presenting the case T^3 :

Example 2.9 (The Misner spacetime). The spacetime

$$M := \mathbb{R}_t \times S_x^1 \times T^2,$$

equipped with the real analytic metric

$$g := 2dt dx + \alpha t dx^2 + g_{T^2},$$

for any $\alpha \in \mathbb{R} \setminus \{0\}$, where g_{T^2} is the standard metric on the 2-dimensional torus, is a vacuum spacetime. Choosing

$$\alpha := -2,$$

there is a future compact Cauchy horizon (if ∂_t is future oriented)

$$\mathcal{H} := \{0\} \times S^1 \times T^2$$

and Killing vector field

$$W := \partial_x.$$

In this case

$$\omega = dx|_{\mathcal{H}},$$

and we get the induced initial data on $\mathcal{H} \cong T^3$:

$$\begin{aligned} \sigma &= dx^2|_{\mathcal{H}} + g_{T^2} = g_{T^3}, \\ V &= \partial_x, \end{aligned}$$

which are the standard metric and a standard Killing vector field on the torus.

Let us continue with $S^1 \times S^2$:

Example 2.10 (The quotient Schwarzschild spacetime). For the case of $S^1 \times S^2$, the idea is to construct a locally isometric (discrete) quotient of the Schwarzschild spacetime. Recall from Example 2.2, that

$$(0, \infty)_r \times \mathbb{R}_v \times S^2, \quad g = 2drdv + \left(\frac{2m}{r} - 1\right) dv^2 + r^2 g_{S^2},$$

is the Schwarzschild spacetime with mass $m \in (0, \infty)$. We first make the coordinate change

$$w := \frac{v}{2}$$

to get

$$(0, \infty)_r \times \mathbb{R}_w \times S^2, \quad g = 4drdw + 4\left(\frac{2m}{r} - 1\right) dw^2 + r^2 g_{S^2},$$

Since ∂_w is a Killing vector field, the metric is well-defined on the quotient spacetime

$$M = (0, \infty)_r \times S_w^1 \times S^2.$$

The event horizon at $r = 2m$ now becomes a compact future Cauchy horizon (if ∂_r is future oriented) given by

$$\mathcal{H} := \{r = 2m\} \times S_w^1 \times S^2.$$

Choosing the horizon Killing vector field

$$W := \partial_w,$$

the surface gravity is

$$\kappa = \frac{1}{2m}$$

and

$$\omega = \frac{1}{2m} dw|_{\mathcal{H}}.$$

The induced lightlike metric on the horizon is

$$4m^2 g_{S^2}.$$

Choosing $m = \frac{1}{2}$, we thus have the following initial data on $\mathcal{H} \cong S^1 \times S^2$:

$$\begin{aligned} \sigma &= dv^2 + g_{S^2}, \\ V &= \partial_w|_{\mathcal{H}}, \end{aligned}$$

which is the standard metric and standard (nowhere vanishing) Killing vector field.

We finish the section by discussing S^3 :

Example 2.11 (The Taub-NUT spacetime). Consider the spacetime

$$\mathbb{R} \times S^3$$

equipped with the metric

$$g := 4l dt \sigma_1 + 4l^2 U(t) \alpha_1^2 + (t^2 + l^2)(\alpha_2^2 + \alpha_3^2).$$

Here, $m \in \mathbb{R}$, $l \neq 0$ and

$$U(t) := \frac{(t_+ - t)(t - t_-)}{t^2 + l^2},$$

with $t_{\pm} := m \pm \sqrt{m^2 + l^2}$, and α_i , for $i = 1, 2, 3$, are the standard left-invariant 1-forms on S^3 . There are a future and a past Cauchy horizon, given by

$$\mathcal{H}_{\pm} := \{t = t_{\pm}\} \times S^3.$$

We choose the horizon Killing vector field to be

$$W := E_1,$$

where E_i , for $i = 1, 2, 3$, are the vector fields dual to α_i , $i = 1, 2, 3$. The surface gravity at the horizon \mathcal{H}_{\pm} is

$$\kappa_{\pm} = \pm \frac{2l\sqrt{m^2 + l^2}}{t_{\pm}^2 + l^2}.$$

We get

$$\omega_{\pm} = \kappa_{\pm} \alpha_1$$

and the induced initial data on $\mathcal{H}_{\pm} \cong S^3$ is

$$\begin{aligned} \sigma &= \frac{4l^2(m^2 + l^2)}{(t_{\pm}^2 + l^2)} \alpha_1^2 + (t_{\pm}^2 + l^2)(\alpha_2^2 + \alpha_3^2), \\ V &= E_1, \end{aligned}$$

which is a Berger metric on S^3 . The induced initial data is the unit sphere metric if $m = 0$ and $l = \pm \frac{1}{\sqrt{2}}$.

3. ASYMPTOTIC EXPANSION

The key in our arguments is to show that the full asymptotic expansion of the spacetime metric g at the horizon is *given uniquely* by the initial data, assuming the Ricci curvature vanishes to infinite order at the horizon:

Assumption 3.1. Assume that M is a smooth spacetime of dimension $n + 1 \geq 2$, with a smooth Killing horizon $\mathcal{H} \subset M$, with horizon Killing vector field W , such that

$$\nabla^k \text{Ric}|_{\mathcal{H}} = 0 \tag{7}$$

for all $k \in \mathbb{N}_0$.

We assume throughout this section that Assumption 3.1 is satisfied.

Remark 3.2. Note that we do not assume the metric to be real analytic in this section, we only assume it be smooth. Moreover, the results hold in arbitrary spacetime dimension.

3.1. Geometric gauge: the null time function. Let (\mathcal{H}, σ, V) denote the induced initial data (in the sense of Definition 1.4). Let

$$V^\perp \subset T\mathcal{H}$$

denote the smooth vector bundle of tangent vectors which are orthogonal to V , with respect to σ . For notational convenience, we do in this section not write out the embedding ι and simply write

$$\mathcal{H} \subset M, \quad V = W|_{\mathcal{H}},$$

etc.

Definition 3.3. The unique lightlike smooth vector field L along \mathcal{H} which satisfies

$$g(L, V) = 1, \quad g(L, X) = 0,$$

for all $X \in V^\perp$ is called the *canonical transversal vector field* of \mathcal{H} .

Note that indeed L is not tangent to \mathcal{H} , since

$$0 \neq g(L, V) = g(L, W|_{\mathcal{H}}).$$

We think of L as a natural replacement for the unit normal along a spacelike or timelike hypersurface.

Proposition 3.4. *There is a unique nowhere vanishing lightlike smooth (real analytic, if the metric is real analytic) vector field ∂_t on an open neighborhood $\mathcal{U} \supset \mathcal{H}$ satisfying*

- $\nabla_{\partial_t} \partial_t = 0$,
- $g(\partial_t|_{\mathcal{H}}, V) = 1$,
- $g(\partial_t|_{\mathcal{H}}, X) = 0$ for all $X \in V^\perp$,
- all integral curves of ∂_t intersect \mathcal{H} precisely once.

Moreover, it follows that

$$[W, \partial_t] = 0.$$

Proof. We construct ∂_t as the unique solution to

$$\begin{aligned} \nabla_{\partial_t} \partial_t &= 0, \\ \partial_t|_{\mathcal{H}} &= L, \end{aligned}$$

on a neighborhood $\mathcal{U} \supset \mathcal{H}$. It follows that

$$\partial_t g(\partial_t, \partial_t) = 0,$$

which together with

$$g(\partial_t, \partial_t)|_{\mathcal{H}} = g(L, L)|_{\mathcal{H}} = 0$$

implies that ∂_t is lightlike. It remains to check that $[W, \partial_t] = 0$. For this, we first show that $[W, \partial_t]|_{\mathcal{H}} = 0$. Note first that

$$\begin{aligned} g([W, \partial_t]|_{\mathcal{H}}, L) &= g(\nabla_W \partial_t|_{\mathcal{H}}, L) - g(\nabla_{\partial_t} W|_{\mathcal{H}}, L) \\ &= \frac{1}{2} W g(L, L) - \frac{1}{2} \mathcal{L}_W g(L, L) \\ &= 0. \end{aligned}$$

Note now that $g(\partial_t|_{\mathcal{H}}, X) = g(L, X) = \frac{\omega(X)}{\kappa}$, for all $X \in T\mathcal{H}$. Using (4), we compute

$$\begin{aligned} g([W, \partial_t]|_{\mathcal{H}}, X) &= W g(\partial_t, X)|_{\mathcal{H}} - \mathcal{L}_W g(\partial_t, X)|_{\mathcal{H}} - g(\partial_t, [W, X])|_{\mathcal{H}} \\ &= W g(\partial_t, X)|_{\mathcal{H}} - g(\partial_t, [W, X])|_{\mathcal{H}} \\ &= \frac{W \omega(X)}{\kappa} - \frac{\omega([W, X])}{\kappa} \end{aligned}$$

$$\begin{aligned}
&= \frac{\mathcal{L}_W \omega(X)}{\kappa} \\
&= 0.
\end{aligned}$$

It follows that $[W, \partial_t]|_{\mathcal{H}} = 0$. Now, since $\mathcal{L}_W g = 0$, we note that

$$0 = \mathcal{L}_W (\nabla_{\partial_t} \partial_t) = \nabla_{[W, \partial_t]} \partial_t + \nabla_{\partial_t} [W, \partial_t],$$

which is a linear first order ODE. It thus follows that $[W, \partial_t] = 0$ as claimed. Shrinking \mathcal{U} if necessary, we can ensure that each integral curve of ∂_t intersects \mathcal{H} precisely once. \square

Proposition 3.5 (The null time function). *Let Assumption 3.1 be satisfied. There is a unique smooth (real analytic, if the metric is real analytic) function*

$$t : \mathcal{U} \rightarrow \mathbb{R},$$

such that

$$\begin{aligned}
dt(\partial_t) &= 1, \\
t^{-1}(0) &= \mathcal{H} \cap \mathcal{U}.
\end{aligned}$$

In particular, $dt \neq 0$ everywhere on \mathcal{U} .

Proof. Construct the function t as the eigentime of the integral curves of ∂_t , starting at \mathcal{H} . Then t is smooth and satisfies the assertion. \square

Definition 3.6. We call $t : \mathcal{U} \rightarrow \mathbb{R}$ the *null time function*.

3.2. The statement. Since the null time function t is constructed geometrically, it is very natural to study the asymptotic expansion of the metric g in terms of t , i.e. we formally write

$$g \sim \sum_{m=0}^{\infty} \mathcal{L}_t^m g|_{\mathcal{H}} \frac{t^m}{m!},$$

and iteratively compute $\mathcal{L}_t^m g|_{\mathcal{H}} := \underbrace{\mathcal{L}_{\partial_t} \dots \mathcal{L}_{\partial_t}}_{m \text{ times}} g|_{\mathcal{H}}$ in terms of the initial data (σ, V) .

Remark 3.7. Proposition 3.4 implies that

$$\begin{aligned}
g(\partial_t, V)|_{\mathcal{H}} &= 1, \\
g(X, Y)|_{\mathcal{H}} &= \sigma(X, Y), \\
g(\partial_t, \partial_t)|_{\mathcal{H}} &= g(\partial_t, X)|_{\mathcal{H}} = g(V, V)|_{\mathcal{H}} = g(X, V)|_{\mathcal{H}} = 0,
\end{aligned}$$

for all $X, Y \in V^{\perp}$. Consequently, since $\partial_t|_{\mathcal{H}}$ is determined by the initial data (σ, V) , we conclude that $g|_{\mathcal{H}}$ is determined by (σ, V) .

We note that many components of $\mathcal{L}_t^m g$ automatically vanish:

Lemma 3.8. *By construction of ∂_t , we have*

$$\mathcal{L}_t^m g(\partial_t, \cdot) = 0,$$

for all $m \in \mathbb{N}$.

Proof. For $m = 1$, we note that for any X with $[\partial_t, X] = 0$, we have

$$\mathcal{L}_t g(\partial_t, X) = \partial_t g(\partial_t, X) = g(\nabla_{\partial_t} \partial_t, X) + g(\partial_t, \nabla_{\partial_t} X) = -\frac{1}{2} X g(\partial_t, \partial_t) = 0.$$

The general statement then follows by induction, by noting that

$$\mathcal{L}_t^m g(\partial_t, X) = \partial_t \mathcal{L}_t^{m-1} g(\partial_t, X). \quad \square$$

For the remaining components of $\mathcal{L}_t^m g|_{\mathcal{H}}$, we will prove the following theorem:

Theorem 3.9. *Let Assumption 3.1 be satisfied. Then there are unique (non-linear) differential operators Q_m on \mathcal{H} for $m \in \mathbb{N}$, such that*

$$\mathcal{L}_t^m g(X, Y)|_{\mathcal{H}} = Q_m(\sigma, V)(X, Y),$$

for all $m \in \mathbb{N}$ and all $X, Y \in T\mathcal{H}$. Moreover, we have

$$Q_m(\phi^* \sigma, \phi^* V) = \phi^* Q_m(\sigma, V), \quad (8)$$

for all diffeomorphisms $\phi : \mathcal{H} \rightarrow \mathcal{H}$ and the $Q_m(\sigma, V)$'s are real analytic if σ and V are real analytic.

The condition (8) is very natural, it says that the differential operators Q_m are *diffeomorphism invariant*. Theorem 3.9 would obviously not be true without assuming (7), i.e. that the Ricci curvature vanishes to infinite order. The rest of this section will be devoted to prove Theorem 3.9, i.e. to study the remaining components of $\mathcal{L}_t^m g|_{\mathcal{H}}$.

3.3. The first derivative. We start by computing $Q_1(\sigma, V)$ in Theorem 3.9. It turns out that its components are given by simple explicit formulas. Let ∇^σ denote the Levi-Civita connection with respect to σ and let R^σ and Ric^σ denote the curvature tensor and the Ricci curvature of σ , respectively. We will also use the notation

$$\nabla_t := \nabla_{\partial_t}, \quad \mathcal{L}_t := \mathcal{L}_{\partial_t}.$$

Proposition 3.10. *In terms of the null time function t , we have*

$$\begin{aligned} \mathcal{L}_t g(V, V)|_{\mathcal{H}} &= -2\kappa, \\ \mathcal{L}_t g(V, X)|_{\mathcal{H}} &= 0, \\ \mathcal{L}_t g(X, Y)|_{\mathcal{H}} &= \frac{1}{\kappa} \left(\text{Ric}^\sigma(X, Y) + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_Y^\sigma V) \right) \end{aligned}$$

for all $X, Y \in V^\perp$, where κ is the surface gravity with respect to W , defined in (2). Moreover, we have

$$\nabla_t W|_{\mathcal{H}} = -\kappa \partial_t|_{\mathcal{H}}. \quad (9)$$

Remark 3.11. We conclude that

$$\begin{aligned} Q_1(\sigma, V)(V, V) &= -2\kappa, \\ Q_1(\sigma, V)(V, X) &= 0, \\ Q_1(\sigma, V)(X, Y) &= \frac{1}{\kappa} \left(\text{Ric}^\sigma(X, Y) + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_Y^\sigma V) \right), \end{aligned}$$

which is a diffeomorphism invariant differential operator in (σ, V) . Since we use the convention that $\kappa > 0$, equation (5) implies that

$$\kappa = \omega(V) = \sqrt{\sigma(V, V)}.$$

This proves the assertion in Theorem 3.9 for $m = 1$.

In order to prove Proposition 3.10, we begin with the following lemma:

Lemma 3.12. *We have*

$$\sum_{i,j=2}^n g^{ij} R(X, e_i, e_j, Y)|_{\mathcal{H}} = \text{Ric}^\sigma(X, Y) + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_Y^\sigma V),$$

for any $X, Y \in V^\perp$, where $\{e_2, \dots, e_n\}$ is a basis for $V^\perp \subset T\mathcal{H}$.

Proof. We need to compare ∇ and ∇^σ . Since

$$\sigma(X, Y) = g(X, Y)|_{\mathcal{H}},$$

for all $X \in V^\perp$ and $Y \in T\mathcal{H}$, the Koszul formula implies for all $X, Y, Z \in V^\perp$ that

$$\begin{aligned} 2\sigma(\nabla_X^\sigma Y, Z) &= X\sigma(Y, Z) + Y\sigma(X, Z) - Z\sigma(X, Y) \\ &\quad + \sigma([X, Y], Z) - \sigma([X, Z], Y) - \sigma([Y, Z], X) \\ &= 2g(\nabla_X Y, Z)|_{\mathcal{H}}. \end{aligned}$$

This implies for all $X, Y \in V^\perp$ that

$$\begin{aligned} \nabla_X^\sigma Y &= \sum_{i,j=2}^n \sigma(\nabla_X^\sigma Y, e_i) \sigma^{ij} e_j + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma Y, V) V \\ &= \sum_{i,j=2}^n g(\nabla_X Y, e_i) g^{ij} e_j|_{\mathcal{H}} + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma Y, V) V \\ &= \nabla_X Y|_{\mathcal{H}} - g(\nabla_X Y, \partial_t)|_{\mathcal{H}} V + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma Y, V) V, \end{aligned}$$

where we have used that $g(\nabla_X Y, V)|_{\mathcal{H}} = 0$. The Koszul formula further implies for $X, Y \in V^\perp$ that

$$\begin{aligned} 2\sigma(\nabla_V^\sigma X, Y) &= V\sigma(X, Y) + X\sigma(V, Y) - Y\sigma(V, X) \\ &\quad + \sigma([V, X], Y) - \sigma([V, Y], X) - \sigma([X, Y], V) \\ &= 2g(\nabla_V X, Y)|_{\mathcal{H}} - \sigma([X, Y], V). \end{aligned}$$

We may now compute for all $X, Y \in V^\perp$ that

$$\begin{aligned} R(X, Y, Y, X)|_{\mathcal{H}} &= g(\nabla_X \nabla_Y Y - \nabla_Y \nabla_X Y - \nabla_{[X, Y]} Y, X)|_{\mathcal{H}} \\ &= g(\nabla_X (\nabla_Y^\sigma Y + g(\nabla_Y Y, \partial_t) V - \frac{1}{\kappa^2} \sigma(\nabla_Y^\sigma Y, V) V), X)|_{\mathcal{H}} \\ &\quad - g(\nabla_Y (\nabla_X^\sigma Y + g(\nabla_X Y, \partial_t) V - \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma Y, V) V), X)|_{\mathcal{H}} \\ &\quad - \sum_{i,j=2}^n \sigma([X, Y], e_i) \sigma^{ij} g(\nabla_{e_j} Y, X)|_{\mathcal{H}} - \frac{1}{\kappa^2} \sigma([X, Y], V) g(\nabla_V Y, X)|_{\mathcal{H}} \\ &= g(\nabla_X \nabla_Y^\sigma Y, X)|_{\mathcal{H}} - g(\nabla_Y \nabla_X^\sigma Y, X)|_{\mathcal{H}} \\ &\quad - \sum_{i,j=2}^n \sigma([X, Y], e_i) \sigma^{ij} g(\nabla_{e_j}^\sigma Y, X)|_{\mathcal{H}} \\ &\quad - \frac{1}{\kappa^2} \sigma([X, Y], V) g(\nabla_V Y, X)|_{\mathcal{H}} \\ &= \sigma(\nabla_X^\sigma \nabla_Y^\sigma Y, X) - \sigma(\nabla_Y^\sigma \nabla_X^\sigma Y, X) - \sigma(\nabla_{[X, Y]}^\sigma Y, X) \\ &\quad + \frac{1}{\kappa^2} \sigma([X, Y], V) \sigma(\nabla_V^\sigma Y, X) - \frac{1}{\kappa^2} \sigma([X, Y], V) g(\nabla_V Y, X)|_{\mathcal{H}} \\ &= R^\sigma(X, Y, Y, X) + \frac{1}{2\kappa^2} \sigma([X, Y], V)^2 \\ &= R^\sigma(X, Y, Y, X) + \frac{1}{2\kappa^2} (\sigma(\nabla_Y^\sigma V, X) - \sigma(\nabla_X^\sigma V, Y))^2 \\ &= R^\sigma(X, Y, Y, X) + \frac{2}{\kappa^2} \sigma(\nabla_X^\sigma V, Y)^2, \end{aligned}$$

where we have used that $\mathcal{L}_V \sigma = 0$. It follows that

$$\begin{aligned} \sum_{i,j=2}^n g^{ij} R(X, e_i, e_j, X)|_{\mathcal{H}} &= \sum_{i,j=2}^n \sigma^{ij} R^\sigma(X, e_i, e_j, X) + \frac{2}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_X^\sigma V) \\ &= \text{Ric}^\sigma(X, X) - \frac{1}{\kappa^2} R^\sigma(X, V, V, X) + \frac{2}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_X^\sigma V). \end{aligned}$$

In order to compute $R^\sigma(X, V, V, X)$, we first note that for all $X \in T\mathcal{H}$, we have

$$\begin{aligned} \sigma(\nabla_V^\sigma V, X) &= \mathcal{L}_V \sigma(V, X) - \sigma(\nabla_X^\sigma V, V) \\ &= -\frac{1}{2} X \sigma(V, V) \\ &= 0, \end{aligned}$$

from which we conclude that

$$\nabla_V^\sigma V = 0.$$

We may thus compute for all $X \in V^\perp$ that

$$\begin{aligned} R^\sigma(X, V, V, X) &= \sigma(\nabla_X^\sigma \nabla_V^\sigma V - \nabla_V^\sigma \nabla_X^\sigma V - \nabla_{[X,V]}^\sigma V, X) \\ &= -V \sigma(\nabla_X^\sigma V, X) + \sigma(\nabla_X^\sigma V, \nabla_V^\sigma X) \\ &\quad - \mathcal{L}_V \sigma([X, V], X) + \sigma(\nabla_X^\sigma V, [X, V]) \\ &= -\frac{1}{2} V \mathcal{L}_V \sigma(X, X) + \sigma(\nabla_X^\sigma V, \nabla_X^\sigma V) \\ &= \sigma(\nabla_X^\sigma V, \nabla_X^\sigma V), \end{aligned}$$

which completes the proof. \square

Proof of Proposition 3.10. Using that $g(\partial_t, W)|_{\mathcal{H}} = 1$ and $\nabla_W W|_{\mathcal{H}} = \kappa W|_{\mathcal{H}}$, we get

$$\mathcal{L}_t g(W, W)|_{\mathcal{H}} = 2g(\nabla_W \partial_t, W)|_{\mathcal{H}} = -2g(\partial_t, \nabla_W W)|_{\mathcal{H}} = -2g(\partial_t, \kappa W)|_{\mathcal{H}} = -2\kappa.$$

Let X be a smooth vector field, such that $X|_{\mathcal{H}} \in C^\infty(V^\perp)$ and $[X, \partial_t] = 0$. We compute

$$\begin{aligned} \mathcal{L}_t g(W, X)|_{\mathcal{H}} &= g(\nabla_W \partial_t, X)|_{\mathcal{H}} + g(W, \nabla_X \partial_t)|_{\mathcal{H}} \\ &= -g(\partial_t, \nabla_W X)|_{\mathcal{H}} - g(\nabla_X W, \partial_t)|_{\mathcal{H}} \\ &= -g([W, X], \partial_t)|_{\mathcal{H}} - 2g(\partial_t, \nabla_X W)|_{\mathcal{H}} \\ &= -\omega([W, X]|_{\mathcal{H}})g(V, \partial_t)|_{\mathcal{H}} - 2\omega(X|_{\mathcal{H}})g(\partial_t, W)|_{\mathcal{H}} \\ &= -\omega([W, X]|_{\mathcal{H}}) \\ &= -W|_{\mathcal{H}} \omega(X) + \mathcal{L}_{W|_{\mathcal{H}}} \omega(X|_{\mathcal{H}}) \\ &= 0, \end{aligned}$$

where we in the last step used (4). Let us now prove (9), using Lemma 3.8. For any $X \in C^\infty(V^\perp)$, we compute

$$\begin{aligned} 0 &= \mathcal{L}_W g(X, \partial_t)|_{\mathcal{H}} = g(\nabla_X W, \partial_t)|_{\mathcal{H}} + g(X, \nabla_t W)|_{\mathcal{H}} = g(X, \nabla_t W)|_{\mathcal{H}}, \\ 0 &= \mathcal{L}_W g(\partial_t, \partial_t)|_{\mathcal{H}} = 2g(\nabla_t W, \partial_t)|_{\mathcal{H}}, \\ 0 &= \mathcal{L}_W g(W, \partial_t)|_{\mathcal{H}} = g(\nabla_W W, \partial_t)|_{\mathcal{H}} + g(W, \nabla_t W)|_{\mathcal{H}} = \kappa + g(W, \nabla_t W)|_{\mathcal{H}}. \end{aligned}$$

By Remark 3.7, we conclude that $\nabla_t W = -\kappa \partial_t$. We use this in the final computation of $\mathcal{L}_W g(X, Y)|_{\mathcal{H}}$ for $X, Y \in C^\infty(V^\perp)$. Since $[W, \partial_t] = 0$, we note that

$$\mathcal{L}_W(\mathcal{L}_t g) = \mathcal{L}_t(\mathcal{L}_W g) + \mathcal{L}_{[W, \partial_t]} g = 0.$$

We compute

$$0 = \mathcal{L}_W(\mathcal{L}_t g)(X, Y)$$

$$\begin{aligned}
&= W(\mathcal{L}_t g(X, Y)) - \mathcal{L}_t g([W, X], Y) - \mathcal{L}_t g(X, [W, Y]) \\
&= W(g(\nabla_X \partial_t, Y) + g(X, \nabla_Y \partial_t)) - g(\nabla_{[W, X]} \partial_t, Y) \\
&\quad - g([W, X], \nabla_Y \partial_t) - g(\nabla_{[W, Y]} \partial_t, X) - g([W, Y], \nabla_X \partial_t) \\
&= g(\nabla_W \nabla_X \partial_t - \nabla_{[W, X]} \partial_t, Y) + g(X, \nabla_W \nabla_Y \partial_t - \nabla_{[W, Y]} \partial_t) \\
&\quad + g(\nabla_X \partial_t, \nabla_W Y) + g(\nabla_W X, \nabla_Y \partial_t) \\
&\quad - g([W, X], \nabla_Y \partial_t) - g([W, Y], \nabla_X \partial_t) \\
&= R(W, X, \partial_t, Y) + g(\nabla_X \nabla_W \partial_t, Y) + R(W, Y, \partial_t, X) + g(\nabla_Y \nabla_W \partial_t, X) \\
&\quad + g(\nabla_X \partial_t, \nabla_Y W) + g(\nabla_Y \partial_t, \nabla_X W).
\end{aligned}$$

Evaluating this at \mathcal{H} , with $X, Y \in C^\infty(V^\perp)$, using (9), we get

$$\begin{aligned}
0 &= R(W, X, \partial_t, Y)|_{\mathcal{H}} - \kappa g(\nabla_X \partial_t, Y)|_{\mathcal{H}} \\
&\quad + R(W, Y, \partial_t, X)|_{\mathcal{H}} - \kappa g(\nabla_Y \partial_t, X)|_{\mathcal{H}} \\
&= R(W, X, \partial_t, Y)|_{\mathcal{H}} + R(W, Y, \partial_t, X)|_{\mathcal{H}} - \kappa \mathcal{L}_t g(X, Y)|_{\mathcal{H}} \\
&= -\text{Ric}(X, Y)|_{\mathcal{H}} - \sum_{i,j=2}^n g^{ij} R(e_i, X, e_j, Y)|_{\mathcal{H}} - \kappa \mathcal{L}_t g(X, Y)|_{\mathcal{H}},
\end{aligned}$$

where e_2, \dots, e_n is a basis for V^\perp . The proof is now completed by applying Lemma 3.12 and recalling that $\text{Ric}|_{\mathcal{H}} = 0$. \square

The following corollary will be useful when computing the higher derivatives:

Corollary 3.13. *For all $X, Y \in C^\infty(V^\perp)$, we have*

$$g(\nabla_X \partial_t, Y)|_{\mathcal{H}} = \frac{1}{2\kappa} \left(\text{Ric}^\sigma(X, Y) + \frac{1}{\kappa^2} \sigma(\nabla_X^\sigma V, \nabla_Y^\sigma V) + d\omega(X, Y) \right).$$

Proof. We compute

$$\begin{aligned}
d\omega(X, Y) &= \partial_X \omega(Y) - \partial_Y \omega(X) - \omega([X, Y]) \\
&= -\omega([X, Y]) \\
&= -g([X, Y], \partial_t)|_{\mathcal{H}} \omega(V) \\
&= -\kappa g([X, Y], \partial_t)|_{\mathcal{H}}
\end{aligned}$$

Combining this with

$$\begin{aligned}
g(\nabla_X \partial_t, Y)|_{\mathcal{H}} &= \frac{1}{2} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} + \frac{1}{2} (g(\nabla_X \partial_t, Y) - g(\nabla_Y \partial_t, X))|_{\mathcal{H}} \\
&= \frac{1}{2} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} - \frac{1}{2} g(\partial_t, [X, Y])
\end{aligned}$$

and Proposition 3.10 yields the desired result. \square

3.4. The higher derivatives. The proof of Theorem 3.9 is an iterative construction of the Q_m 's. The proof is constructive, meaning that it in principle is possible to compute the Q_m 's explicitly, just like we did for Q_1 in the previous subsection. Remark 3.11 implies that there is a unique Q_1 such that

$$\mathcal{L}_t g(X, Y)|_{\mathcal{H}} = Q_1(\sigma, V)(X, Y)$$

for all $X, Y \in T\mathcal{H}$. Let us therefore make the following induction assumption:

Induction Assumption 3.14. Fix an $m \in \mathbb{N}$. We assume that there are unique (non-linear) diffeomorphism invariant differential operators Q_1, \dots, Q_m on \mathcal{H} such that

$$\mathcal{L}_t^k g(X, Y) = Q_k(\sigma, V)(X, Y)$$

for all $X, Y \in T\mathcal{H}$ and all $k = 1, \dots, m$.

Remark 3.15. Indeed, by Remark 3.11 we have proven that Induction Assumption 3.14 is satisfied for

$$m = 1.$$

Given Induction Assumption 3.14, the goal of this section is to show the existence of a unique Q_{m+1} such that

$$\mathcal{L}_t^{m+1}g(X, Y)|_{\mathcal{H}} = Q_{m+1}(\sigma, V)(X, Y)$$

for all $X, Y \in T\mathcal{H}$, which by induction would prove Theorem 3.9. Recall that we already know that

$$\mathcal{L}_t^m g(\partial_t, \cdot) = 0,$$

for all $m \in \mathbb{N}$.

3.4.1. *Notation and first identities.* It will be convenient to use the notation

$$\Theta(X) = \nabla_X \partial_t, \quad A(X, Y) := g(\Theta(X), Y).$$

We also define the square of a $(0, 2)$ -tensor T as

$$T^2(X, Y) := \sum_{\alpha, \beta=0}^n g^{\alpha\beta} T(X, e_\alpha) T(e_\beta, Y).$$

Lemma 3.16. *We have the identities*

$$\begin{aligned} \nabla_t A(X, Y) &= -A^2(X, Y) + R(\partial_t, X, \partial_t, Y), \\ \nabla_t \mathcal{L}_t g(X, Y) &= -A^2(X, Y) - A^2(Y, X) + 2R(\partial_t, X, \partial_t, Y), \\ \nabla_t A(X, Y) &= \frac{1}{2} \nabla_t \mathcal{L}_t g(X, Y) - \frac{1}{2} (A^2(X, Y) - A^2(Y, X)). \end{aligned}$$

Proof. Since $\nabla_t \partial_t = 0$, we have

$$\begin{aligned} \nabla_t A(X, Y) &= g(\nabla_{\partial_t, X}^2 \partial_t, Y) \\ &= g(\nabla_{X, \partial_t}^2 \partial_t, Y) + R(\partial_t, X, \partial_t, Y) \\ &= -g(\nabla_{\nabla_X \partial_t} \partial_t, Y) + R(\partial_t, X, \partial_t, Y) \\ &= -g(\Theta(\Theta(X)), Y) + R(\partial_t, X, \partial_t, Y) \\ &= -A^2(X, Y) + R(\partial_t, X, \partial_t, Y). \end{aligned}$$

The second and the third identities follow from the identity

$$\mathcal{L}_t g(X, Y) = A(X, Y) + A(Y, X).$$

□

3.4.2. *The curvature components.* The strategy in the proof of Theorem 3.9 is to show that $\mathcal{L}_t^{m+1}g|_{\mathcal{H}}$ is given uniquely by lower orders $\mathcal{L}_t^k g|_{\mathcal{H}}$ and for $k = 0, \dots, m$, using that derivatives of the Ricci curvature vanish at the horizon. The following lemma will be crucial for that purpose:

Lemma 3.17. *Let $X, Y, Z, W \in C^\infty(T\mathcal{H})$ and fix a $k \in \mathbb{N}$. Then there is a unique way to express*

- (a) $\nabla_t^k A|_{\mathcal{H}}$, for all $j = 0, \dots, k-1$,
- (b) $(\nabla_t^j \mathcal{L}_t g - \mathcal{L}_t^{j+1} g)|_{\mathcal{H}}$, for all $j = 0, \dots, k$,
- (c) $\nabla_t^j R(\partial_t, X, \partial_t, Y)|_{\mathcal{H}}$, for all $j = 0, \dots, k-2$,
- (d) $\nabla_t^j R(X, Y, \partial_t, Z)|_{\mathcal{H}}$, for all $j = 0, \dots, k-1$,
- (e) $\nabla_t^j R(X, Y, Z, W)|_{\mathcal{H}}$, for all $j = 0, \dots, k$,

in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^k g|_{\mathcal{H}}. \tag{10}$$

In the proof of this lemma and in further computations, we will use the following schematic notation:

Notation 3.18. Given two tensors A and B , the notation

$$A * B$$

denotes a tensor, which is given by linear combinations of contractions with respect to the metric g (not derivatives of g). In particular, we may write

$$\nabla^k(A * B) = \sum_{i+j=k} \nabla^i A * \nabla^j B.$$

The proof of Lemma 3.17 will use the following simple observation:

Lemma 3.19. *Let S be a smooth tensor field on M . For any $k \in \mathbb{N}$, the commutator*

$$[\nabla_{t,\dots,t}^k, \nabla]S|_{\mathcal{H}}$$

is determined uniquely in terms of $g|_{\mathcal{H}}$ and

$$\nabla_t^j R|_{\mathcal{H}}, \quad \nabla^j A|_{\mathcal{H}},$$

for $j = 0, \dots, k-1$ and

$$\nabla^j S|_{\mathcal{H}}$$

for $j = 0, \dots, k$.

Proof of Lemma 3.19. Note that for any tensor field S , we have

$$\begin{aligned} [\nabla_t, \nabla]S(X) &= \nabla_{t,X}^2 S - \nabla_X(\nabla_t S) \\ &= R(\partial_t, X)S - \nabla_{\Theta(X)} S. \end{aligned}$$

Using this and $\nabla_t \partial_t = 0$, we may schematically compute the higher commutators as

$$\begin{aligned} [\nabla_{t,\dots,t}^k, \nabla]S &= [(\nabla_t)^k, \nabla]S \\ &= \sum_{i+j=k-1} (\nabla_t)^i [\nabla_t, \nabla](\nabla_t)^j S \\ &= \sum_{i+j=k-1} (\nabla_t)^i R(\partial_t, \cdot) \nabla_{t,\dots,t}^j S - (\nabla_t)^i \nabla_{\Theta(\cdot)} \nabla_{t,\dots,t}^j S \\ &= \sum_{i+j=k-1} \sum_{a+b=i} (\nabla^a R) * (\nabla^{b+j} S) - (\nabla^a \Theta) * (\nabla^{b+j+1} S). \end{aligned}$$

This completes the proof, since $\nabla^k A = g(\nabla^k \Theta(\cdot), \cdot)$. \square

Proof of Lemma 3.17. We start by proving the statement for $k = 1$. By Corollary 3.13, we know that

$$A|_{\mathcal{H}},$$

and hence $\Theta|_{\mathcal{H}}$, are given in terms of $g|_{\mathcal{H}}$ and $\mathcal{L}_t g|_{\mathcal{H}}$, which proves claim (a) for $k = 1$. Claim (b) then follows for $k = 1$ immediately by noting that

$$\nabla_t S = \mathcal{L}_t S + A * S, \tag{11}$$

for any covariant tensor S , from which we get

$$\nabla_t \mathcal{L}_t g|_{\mathcal{H}} = \mathcal{L}_t^2 g|_{\mathcal{H}} + A * \mathcal{L}_t g|_{\mathcal{H}}.$$

Further, we compute

$$\begin{aligned} R(X, Y, \partial_t, Z)|_{\mathcal{H}} &= Xg(\nabla_Y \partial_t, Z)|_{\mathcal{H}} - g(\nabla_Y \partial_t, \nabla_X Z)|_{\mathcal{H}} - Yg(\nabla_X \partial_t, Z)|_{\mathcal{H}} \\ &\quad + g(\nabla_X \partial_t, \nabla_Y Z)|_{\mathcal{H}} - g(\nabla_{[X,Y]} \partial_t, Z)|_{\mathcal{H}} \\ &= \nabla_X A(Y, Z)|_{\mathcal{H}} - \nabla_Y A(X, Z)|_{\mathcal{H}}, \end{aligned}$$

for all $X, Y, Z \in T\mathcal{H}$. Clearly, ∇ can be expressed in terms of $g|_{\mathcal{H}}$ and $\mathcal{L}_t g|_{\mathcal{H}}$ and since X, Y are tangent to \mathcal{H} , we conclude claim (d) for $k = 1$. The curvature component $R(X, Y, Z, W)|_{\mathcal{H}}$ can locally be given in terms of Christoffel symbols and derivatives thereof tangent to \mathcal{H} , which are all given by $g|_{\mathcal{H}}$ and $\mathcal{L}_t g|_{\mathcal{H}}$, proving claim (e) for $j = 0$.

In order to complete the case $k = 1$, we still need to prove (e) for $j = 1$. Using the second Bianchi identity, we compute the derivatives of the third curvature component:

$$\begin{aligned} \nabla_t^j R(X, Y, Z, W)|_{\mathcal{H}} &= \nabla_t^{j-1}(\nabla R)(\partial_t, X, Y, Z, W)|_{\mathcal{H}} \\ &= -\nabla_t^{j-1}(\nabla R)(X, Y, \partial_t, Z, W)|_{\mathcal{H}} - \nabla_t^{j-1}(\nabla R)(Y, \partial_t, X, Z, W)|_{\mathcal{H}} \\ &= -[\nabla_t^{j-1}, \nabla]R(X, Y, \partial_t, Z, W)|_{\mathcal{H}} - \nabla_X \nabla_t^{j-1}R(Y, \partial_t, Z, W)|_{\mathcal{H}} \\ &\quad - [\nabla_t^{j-1}, \nabla]R(Y, \partial_t, X, Z, W)|_{\mathcal{H}} - \nabla_Y \nabla_t^{j-1}R(\partial_t, X, Z, W)|_{\mathcal{H}}, \end{aligned} \quad (12)$$

for $X, Y, Z, W \in T\mathcal{H}$. Using this with $j = 1$ and that we have proven claim (d) for $k = 1$, we conclude claim (e) for $j = 1$. We have thus proven Lemma 3.17 with $k = 1$, which is initial step in our induction on k .

Assume therefore that the assertion is proven for an $k \in \mathbb{N}$, we want to then prove it for $k + 1$. Lemma 3.16 implies that

$$\begin{aligned} \nabla_t^k A|_{\mathcal{H}} &= \frac{1}{2} \nabla_t^k \mathcal{L}_t g|_{\mathcal{H}} + \sum_{a+b=k-1} \nabla_t^a A * \nabla_t^b A|_{\mathcal{H}} \\ &= \frac{1}{2} (\nabla_t^k \mathcal{L}_t g - \mathcal{L}_t^{k+1} g)|_{\mathcal{H}} + \mathcal{L}_t^{k+1} g|_{\mathcal{H}} + \sum_{a+b=k-1} \nabla_t^a A * \nabla_t^b A|_{\mathcal{H}}, \end{aligned}$$

which by the induction assumption is given by

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^{k+1} g|_{\mathcal{H}}.$$

proving claim (a) for $k + 1$. By applying (11), we note that

$$(\nabla_t^{k+1} \mathcal{L}_t g - \mathcal{L}_t^{k+2} g)|_{\mathcal{H}} = \sum_{a+b=k} \nabla_t^a A * \nabla_t^b \mathcal{L}_t g|_{\mathcal{H}}.$$

By claim (a) for $k + 1$, this proves claim (b) for $k + 1$. By Lemma 3.16, we deduce that

$$\nabla_t^{k-1} R(\partial_t, X, \partial_t, Y)|_{\mathcal{H}} = \nabla_t^k A|_{\mathcal{H}} + \sum_{j=0}^{k-1} \nabla_t^j A * \nabla_t^{k-1-j} A(X, Y)|_{\mathcal{H}},$$

proving claim (c) for $k + 1$. To prove claim (d) for $k + 1$, we note that the second Bianchi identity implies similar to (12) that

$$\begin{aligned} \nabla_t^k R(X, Y, \partial_t, Z)|_{\mathcal{H}} &= -[\nabla_t^{k-1}, \nabla]R(X, Y, \partial_t, \partial_t, Z)|_{\mathcal{H}} - \nabla_X \nabla_t^{k-1} R(Y, \partial_t, \partial_t, Z)|_{\mathcal{H}} \\ &\quad - [\nabla_t^{k-1}, \nabla]R(Y, \partial_t, X, \partial_t, Z)|_{\mathcal{H}} - \nabla_Y \nabla_t^{k-1} R(\partial_t, X, \partial_t, Z)|_{\mathcal{H}}, \end{aligned}$$

for any $X, Y, Z \in T\mathcal{H}$. By Lemma 3.19 and claim (c) for $k + 1$, the right hand side is expressed in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^{k+1} g|_{\mathcal{H}},$$

proving claim (d) for $k + 1$. Finally, in order to prove claim (e), we consider equation (12) with $j = k + 1$ and get

$$\nabla_t^{k+1} R(X, Y, Z, W)|_{\mathcal{H}}$$

$$\begin{aligned}
&= -[\nabla_t^k, \nabla]R(X, Y, \partial_t, Z, W)|_{\mathcal{H}} - \nabla_X \nabla_t^k R(Y, \partial_t, Z, W)|_{\mathcal{H}} \\
&\quad - [\nabla_t^k, \nabla]R(Y, \partial_t, X, Z, W)|_{\mathcal{H}} - \nabla_Y \nabla_t^k R(\partial_t, X, Z, W)|_{\mathcal{H}}.
\end{aligned}$$

Again, by Lemma 3.19 and claim (d) for $k+1$, the right hand side is expressed in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^{k+1} g|_{\mathcal{H}},$$

proving claim (e) for $k+1$.

We have thus proven the assertion in Lemma 3.17 for $k+1$, which completes the induction argument. \square

3.4.3. Expressions for the V -components. With the preparations in the previous subsection, we are now able to study the expression

$$\mathcal{L}_t^{m+1} g(\cdot, V)|_{\mathcal{H}}.$$

Proposition 3.20. *Let Assumption 3.1 and Induction Assumption 3.14 be satisfied for an $m \in \mathbb{N}$. Then, for any $X \in C^\infty(T\mathcal{H})$, the expression*

$$\mathcal{L}_t^{m+1} g(X, V)|_{\mathcal{H}}$$

is given uniquely in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^m g|_{\mathcal{H}}. \quad (13)$$

Proof. By Lemma 3.16, we have

$$\nabla_t^m \mathcal{L}_t g(X, V)|_{\mathcal{H}} = \sum_{i+j=m-1} \nabla_t^i A * \nabla_t^j A(X, V)|_{\mathcal{H}} + 2\nabla_t^{m-1} R(\partial_t, X, \partial_t, V)|_{\mathcal{H}}$$

for all $X \in T\mathcal{H}$. The first term on the right hand side is dealt with in Lemma 3.17. The second term is computed as follows:

$$\begin{aligned}
0 &= \nabla_t^{m-1} \text{Ric}(X, \partial_t)|_{\mathcal{H}} \\
&= \nabla_t^{m-1} \text{tr}_g(R(\cdot, X, \partial_t, \cdot))|_{\mathcal{H}} \\
&= \text{tr}_g((\nabla_t^{m-1} R)(\cdot, X, \partial_t, \cdot))|_{\mathcal{H}} \\
&= \nabla_t^{m-1} R(\partial_t, X, \partial_t, V)|_{\mathcal{H}} + \sum_{i,j=2}^n g^{ij} \nabla_t^{m-1} R(e_i, X, \partial_t, e_j)|_{\mathcal{H}}.
\end{aligned}$$

By Lemma 3.17, the sum on the right hand side is given by (13). The proof is completed by applying Lemma 3.17, claim (b). \square

3.4.4. Computation of the commutator. In this section, we will compute the following commutator, which is essential for the remaining part of the proof of Theorem 3.9:

$$[\nabla_t^{m-1}, \square] \mathcal{L}_t g.$$

Proposition 3.21. *Let Assumption 3.1 and Induction Assumption 3.14 be satisfied for an $m \in \mathbb{N}$. The expression*

$$[\nabla_t^{m-1}, \square] \mathcal{L}_t g|_{\mathcal{H}} + 2(m-1)\kappa \mathcal{L}_t^{m+1} g|_{\mathcal{H}}$$

is given uniquely in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^m g|_{\mathcal{H}}. \quad (14)$$

We begin with the following lemma:

Lemma 3.22. *For any tensor field S , we schematically have*

$$\begin{aligned}
[\nabla_t, \nabla]S(X) &= R(\partial_t, X)S - \nabla_{\Theta(X)}S, \\
[\nabla_t, \nabla^2]S &= \nabla R * S + \Theta * R * S + R * \nabla S + \nabla \Theta * \nabla S + \Theta * \nabla^2 S, \\
[\nabla_t, \square]S &= g(\mathcal{L}_t g, \nabla^2)S + \nabla \text{Ric}(\partial_t, \cdot) * S + R * \nabla S + \nabla \Theta * \nabla S.
\end{aligned}$$

Here, we have used the notation

$$g(T, \nabla^2)S := \sum_{\alpha, \beta, \gamma, \delta=0}^n g^{\alpha\beta} g^{\gamma\delta} T(e_\alpha, e_\beta) \nabla_{e_\beta, e_\delta}^2 S.$$

Proof. Note first that

$$\begin{aligned} [\nabla_t, \nabla]S(X) &= \nabla_{t,X}^2 S - \nabla_X(\nabla_t S) \\ &= R(\partial_t, X)S - \nabla_{\Theta(X)} S, \end{aligned}$$

which proves the first formula. For the second formula, write

$$([\nabla_t, \nabla^2]S)(X, Y) = ([\nabla_t, \nabla]\nabla S)(X, Y) + (\nabla[\nabla_t, \nabla])S(X, Y).$$

Inserting the above yields

$$\begin{aligned} [\nabla_t, \nabla]\nabla S(X, Y) &= R(\partial_t, X)\nabla_Y S - \nabla_{R(\partial_t, X)Y} S - \nabla_{\Theta(X), Y}^2 S, \\ \nabla[\nabla_t, \nabla]S(X, Y) &= (\nabla_X R)(\partial_t, Y)S + R(\Theta(X), Y)S + R(\partial_t, Y)\nabla_X S \\ &\quad - \nabla_{X, \Theta(Y)}^2 S - \nabla_{(\nabla_X \Theta)(Y)} S, \end{aligned}$$

proving in particular the second formula. We may now contract over X and Y to get

$$\begin{aligned} [\nabla_t, \square]S &= [\nabla_t, -\text{tr}_g(\nabla^2)]S \\ &= -\text{tr}_g([\nabla_t, \nabla^2])S \\ &= \text{tr}_g((\nabla \cdot R)(\partial_t, \cdot))S + \text{tr}_g\left(\nabla_{\cdot, \Theta(\cdot)}^2 + \nabla_{\Theta(\cdot), \cdot}^2\right)S \\ &\quad + \Theta * R * S + R * \nabla S + \nabla \Theta * \nabla S. \end{aligned}$$

We therefore consider the endomorphism

$$\text{tr}_g((\nabla \cdot R)(\partial_t, \cdot))$$

by using the second Bianchi identity s to write

$$\begin{aligned} g(\text{tr}_g((\nabla \cdot R)(\partial_t, \cdot))X, Y) &= \sum_{\alpha, \beta=0}^n g(\nabla_{e_\alpha} R(\partial_t, e_\beta)X, Y) \\ &= \sum_{\alpha, \beta=0}^n \nabla_{e_\alpha} R(X, Y, \partial_t, e_\beta) \\ &= - \sum_{\alpha, \beta=0}^n \nabla_X R(Y, e_\alpha, \partial_t, e_\beta) - \sum_{\alpha, \beta=0}^n \nabla_Y R(e_\alpha, X, \partial_t, e_\beta) \\ &= \nabla_X \text{Ric}(\partial_t, Y) - \nabla_Y \text{Ric}(\partial_t, X). \end{aligned}$$

Finally, we use the formula

$$\mathcal{L}_t g(X, Y) = g(\Theta(X), Y) + g(X, \Theta(Y))$$

to see that

$$\text{tr}_g\left(\nabla_{\cdot, \Theta(\cdot)}^2 + \nabla_{\Theta(\cdot), \cdot}^2\right)S = g(\mathcal{L}_t g, \nabla^2)S.$$

This completes the proof. \square

Proof of Proposition 3.21. We first note that

$$[\nabla_t^{m-1}, \square]\mathcal{L}_t g|_{\mathcal{H}} = \sum_{i+j=m-2} \nabla_t^i [\nabla_t, \square] \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}}.$$

By applying Lemma 3.17, Lemma 3.22 and Assumption 3.1, we note that the only term in this sum which is not determined by (14) is

$$\sum_{i+j=m-2} \nabla_t^i g(\mathcal{L}_t g, \nabla^2) \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}} = \sum_{i+j=m-2} \sum_{a+b=i} g(\nabla_t^a \mathcal{L}_t g, \nabla_t^b \nabla^2) \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}}.$$

The only term in this sum which is not determined by (14) is

$$\begin{aligned} \sum_{i+j=m-2} g(\mathcal{L}_t g, \nabla_t^i \nabla^2) \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}} &= \sum_{i+j=m-2} g(\mathcal{L}_t g, [\nabla_t^i, \nabla^2]) \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad + \sum_{i+j=m-2} g(\mathcal{L}_t g, \nabla^2) \nabla_t^i \nabla_t^j \mathcal{L}_t g|_{\mathcal{H}} \\ &= \sum_{i+j=m-2} \sum_{a+l=i-1} g(\mathcal{L}_t g, \nabla_t^a [\nabla_t, \nabla^2]) \nabla_t^{l+j} \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad + (m-1) g(\mathcal{L}_t g, \nabla^2) \nabla_t^{m-2} \mathcal{L}_t g|_{\mathcal{H}}. \end{aligned}$$

Again, Lemma 3.22 implies that the only term which is not determined by (14) is

$$(m-1) g(\mathcal{L}_t g, \nabla^2) \nabla_t^{m-2} \mathcal{L}_t g|_{\mathcal{H}} = (m-1) g^{\alpha\gamma} g^{\beta\delta} \mathcal{L}_t g_{\alpha\beta} \nabla_{e_\delta, e_\gamma}^2 \nabla_t^{m-2} \mathcal{L}_t g|_{\mathcal{H}}.$$

By Proposition 3.10 and Lemma 3.22, the only term in this expression which is not given by (14), is the term

$$-2(m-1) \kappa \nabla_t^m \mathcal{L}_t g|_{\mathcal{H}}.$$

Applying Lemma 3.17, claim (b), completes the proof. \square

3.4.5. Expressions for the V^\perp -components. We now turn to the computation of the final components

$$\mathcal{L}_t^{m+1} g(X, Y)|_{\mathcal{H}}, \quad (15)$$

where $X, Y \in V^\perp$. This will use the linearization of the Ricci curvature $\mathcal{L}_t \text{Ric}|_{\mathcal{H}}$, found for example in [Bes08, Thm. 1.174]:

$$2\mathcal{L}_t \text{Ric} = \square_L \mathcal{L}_t g + \mathcal{L}_{\text{div}(\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)}^\# g, \quad (16)$$

where \square_L is the d'Alembert-Lichnerowicz operator, defined as

$$\square_L h := \square h - 2\mathring{R}h,$$

where

$$\mathring{R}h(X, Y) := \text{tr}_g(h(\mathring{R}(\cdot, X)Y, \cdot)).$$

Proposition 3.23. *Let Assumption 3.1 and Induction Assumption 3.14 be satisfied for an $m \in \mathbb{N}$. For $X, Y \in C^\infty(V^\perp)$, the expression*

$$\mathcal{L}_t^{m+1} g(X, Y)|_{\mathcal{H}}$$

is given uniquely in terms of

$$g|_{\mathcal{H}}, \dots, \mathcal{L}_t^m g|_{\mathcal{H}} \quad \text{and} \quad \mathcal{L}_t^{m+1} g(\cdot, V)|_{\mathcal{H}}. \quad (17)$$

The idea in the proof is to differentiate (16):

$$\begin{aligned} 2\nabla_t^{m-1} \mathcal{L}_t \text{Ric}|_{\mathcal{H}} &= \nabla_t^{m-1} \left(\square \mathcal{L}_t g - 2\mathring{R} \mathcal{L}_t g + \mathcal{L}_{\text{div}((\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)}^\# g) \right)|_{\mathcal{H}} \\ &= [\nabla_t^{m-1}, \square] \mathcal{L}_t g|_{\mathcal{H}} + \square \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} - 2\nabla_t^{m-1} \mathring{R} \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad + \nabla_t^{m-1} \mathcal{L}_{\text{div}((\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)}^\# g)|_{\mathcal{H}}. \end{aligned} \quad (18)$$

By Proposition 3.21, we know that

$$[\nabla_t^{m-1}, \square] \mathcal{L}_t g|_{\mathcal{H}} + 2(m-1) \kappa \mathcal{L}_t^{m+1} g|_{\mathcal{H}}$$

is given by (17). The remaining three terms in (18) are treated separately in Lemma 3.24, Lemma 3.25 and Lemma 3.26 below.

Lemma 3.24. *For $X, Y \in C^\infty(V^\perp)$, the expression*

$$\square \nabla_t^{m-1} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} + 2\kappa \mathcal{L}_t^{m+1} g(X, Y)|_{\mathcal{H}},$$

is uniquely determined by (17).

Proof. We compute that

$$\begin{aligned} \square \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} &= - \sum_{i,j=2}^n g^{\alpha\beta} \nabla_{e_\alpha, e_\beta}^2 \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} \\ &= - \nabla_{W,t}^2 \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} - \nabla_{t,W}^2 \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad - \sum_{i,j=2}^n g^{ij} \nabla_{e_i, e_j}^2 \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} \\ &= -2 \nabla_W \nabla_t^m \mathcal{L}_t g|_{\mathcal{H}} + 2 \nabla_{\nabla_W \partial_t} \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad - R(\partial_t, W) \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} - \sum_{i,j=2}^n g^{ij} \nabla_{e_i} \nabla_{e_j} \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} \\ &\quad + \sum_{i,j=2}^n g^{ij} \nabla_{\nabla_{e_i} e_j} \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}}. \end{aligned}$$

By Lemma 3.17 (b) and since $\nabla_{e_i} e_j|_{\mathcal{H}} \in T\mathcal{H}$, the fourth and fifth term are uniquely given by (17). We compute the second term using

$$\nabla_W \partial_t|_{\mathcal{H}} = -\kappa \partial_t|_{\mathcal{H}},$$

by Proposition 3.10, and get

$$2 \nabla_{\nabla_W \partial_t} \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}} = -2\kappa \nabla_t^m \mathcal{L}_t g|_{\mathcal{H}}.$$

Applying Lemma 3.17 (b), we conclude that the second term uniquely given by (17). The first term is computed using that W is a Killing vector field with $[\partial_t, W] = 0$. For $X, Y \in V^\perp$, we have

$$\begin{aligned} -2 \nabla_W \nabla_t^{m+1} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} &= -2 \mathcal{L}_W \nabla_t^{m+1} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} + 2 \nabla_t^{m+1} \mathcal{L}_t g(\nabla_X W, Y)|_{\mathcal{H}} \\ &\quad + 2 \nabla_t^{m+1} \mathcal{L}_t g(X, \nabla_Y W)|_{\mathcal{H}} \\ &= -2 \nabla_t^{m+1} \mathcal{L}_t \mathcal{L}_W g(X, Y)|_{\mathcal{H}} \\ &= 0. \end{aligned}$$

We finally consider the last term. Note first that for $m > 1$, Lemma 3.17 implies that $R|_{\mathcal{H}}$ is determined by (17). By Lemma 3.17 (b), we conclude that in this case, the last term is determined by (17). In case $m = 1$, Proposition 3.10 implies that for any $X, Y \in V^\perp$, we have

$$\begin{aligned} -R(\partial_t, W) \mathcal{L}_t g(X, Y)|_{\mathcal{H}} &= \mathcal{L}_t g(R(\partial_t, W)X, Y)|_{\mathcal{H}} + \mathcal{L}_t g(X, R(\partial_t, W)Y)|_{\mathcal{H}} \\ &= \sum_{i,j=2}^n R(\partial_t, W, X, e_j) g^{ij} \mathcal{L}_t g(e_j, Y)|_{\mathcal{H}} \\ &\quad + \sum_{i,j=2}^n R(\partial_t, W, X, e_i) g^{ij} \mathcal{L}_t g(X, e_j)|_{\mathcal{H}}. \end{aligned}$$

These curvature components are determined (17), by Lemma 3.17 (d). Taken together, this completes the proof. \square

Lemma 3.25. *For all $X, Y \in V^\perp$, the expression*

$$\nabla_t^{m-1}(\mathring{\mathcal{R}}\mathcal{L}_t g)(X, Y)|_{\mathcal{H}} - \kappa \mathcal{L}_t^{m+1} g(X, Y)|_{\mathcal{H}}$$

is given uniquely by (17).

Proof. We compute

$$\begin{aligned} & \nabla_t^{m-1} \mathring{\mathcal{R}}\mathcal{L}_t g(X, Y)|_{\mathcal{H}} \\ &= \sum_{\alpha, \beta, \gamma, \delta=0}^n \sum_{k=0}^{m-1} \binom{m-1}{k} g^{\alpha\beta} g^{\gamma\delta} \nabla_t^k \mathcal{R}(e_\alpha, X, Y, e_\gamma) \nabla_t^{m-1-k} \mathcal{L}_t g(e_\beta, e_\delta)|_{\mathcal{H}} \\ &= \nabla_t^{m-1} \mathcal{R}(\partial_t, X, Y, \partial_t) \mathcal{L}_t g(V, V)|_{\mathcal{H}} \\ &\quad + \sum_{i,j=2}^n \sum_{a,b=2}^n g^{ij} g^{ab} \nabla_t^{m-1} \mathcal{R}(e_i, X, Y, e_a) \mathcal{L}_t g(e_j, e_b)|_{\mathcal{H}} \\ &\quad + \sum_{k=0}^{m-2} \nabla_t^k \mathcal{R} * \nabla_t^{m-1-k} \mathcal{L}_t g(X, Y)|_{\mathcal{H}}. \end{aligned}$$

By Lemma 3.17, the second and the third terms are given by (17). The first term is computed using that $\mathcal{L}_t g(V, V)|_{\mathcal{H}} = -2\kappa$, by Proposition 3.10, and Lemma 3.16:

$$\begin{aligned} \nabla_t^{m-1} \mathcal{R}(\partial_t, X, Y, \partial_t) \mathcal{L}_t g(V, V)|_{\mathcal{H}} &= \kappa \nabla_t^m \mathcal{L}_t g(X, Y)|_{\mathcal{H}} \\ &\quad + \sum_{k=0}^{m-1} \nabla_t^k A * \nabla_t^{m-1-k} A(X, Y)|_{\mathcal{H}}. \end{aligned}$$

By Lemma 3.17 (a), the sum is given uniquely by (17). The proof is hence finished by applying Lemma 3.17 (b). \square

Lemma 3.26. *For any $X, Y \in T\mathcal{H}$, we have*

$$\nabla_t^{m-1} \mathcal{L}_{\text{div}(\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)}^\# g(X, Y)|_{\mathcal{H}}$$

is given uniquely by (17).

Proof. We have

$$\begin{aligned} & \nabla_t^{m-1} \mathcal{L}_{\text{div}(\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)}^\# g(X, Y)|_{\mathcal{H}} \\ &= g(\nabla_{t, \dots, t, X}^m \text{div}(\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)^\#, Y)|_{\mathcal{H}} \\ &\quad + g(X, \nabla_{t, \dots, t, Y}^m \text{div}(\mathcal{L}_t g - \frac{1}{2} \text{tr}_g(\mathcal{L}_t g)g)^\#)|_{\mathcal{H}} \\ &= \sum_{\alpha, \beta=0}^n g^{\alpha\beta} (\nabla_{t, \dots, t, X, e_\alpha}^{m+1} (\mathcal{L}_t g)(e_\beta, Y)|_{\mathcal{H}} + \sum_{\alpha, \beta=0}^n g^{\alpha\beta} \nabla_{t, \dots, t, Y, e_\alpha}^{m+1} (\mathcal{L}_t g)(e_\beta, X)|_{\mathcal{H}} \\ &\quad - \sum_{\alpha, \beta=0}^n g^{\alpha\beta} \nabla_{t, \dots, t, X, Y}^{m+1} (\mathcal{L}_t g)(e_\alpha, e_\beta)|_{\mathcal{H}} \\ &= \nabla_{t, \dots, t, X, t}^{m+1} (\mathcal{L}_t g)(V, Y)|_{\mathcal{H}} + \nabla_{t, \dots, t, Y, t}^{m+1} (\mathcal{L}_t g)(V, X)|_{\mathcal{H}} \\ &\quad + \nabla_{t, \dots, t, X, V}^{m+1} (\mathcal{L}_t g)(\partial_t, Y)|_{\mathcal{H}} + \nabla_{t, \dots, t, Y, V}^{m+1} (\mathcal{L}_t g)(\partial_t, X)|_{\mathcal{H}} \\ &\quad + \sum_{i,j=2}^n g^{ij} \nabla_{t, \dots, t, X, e_i}^{m+1} (\mathcal{L}_t g)(e_j, Y)|_{\mathcal{H}} + \sum_{i,j=2}^n g^{ij} \nabla_{t, \dots, t, Y, e_i}^{m+1} (\mathcal{L}_t g)(e_j, X)|_{\mathcal{H}} \\ &\quad - \sum_{\alpha, \beta=0}^n g^{\alpha\beta} \nabla_{t, \dots, t, X, Y}^{m+1} (\mathcal{L}_t g)(e_\alpha, e_\beta)|_{\mathcal{H}}. \end{aligned} \tag{19}$$

We begin by noting that

$$\nabla_{t,\dots,t,X,t}^{m+1}(\mathcal{L}_t g)(V, Y)|_{\mathcal{H}} = [\nabla_t^{m-1}, \nabla] \nabla_t \mathcal{L}_t g(X, V, Y)|_{\mathcal{H}} + \nabla_X \nabla_t^m \mathcal{L}_t g(V, Y)|_{\mathcal{H}}. \quad (20)$$

Since $X \in T\mathcal{H}$, the second term in (20) is given by (17). We compute the first term in (20) as follows:

$$\begin{aligned} [\nabla_t^{m-1}, \nabla] \nabla_t \mathcal{L}_t g(X, V, Y)|_{\mathcal{H}} &= \sum_{i+j=m-2} \nabla_t^i [\nabla_t, \nabla] \nabla_t^{j+1} \mathcal{L}_t g(X, V, Y)|_{\mathcal{H}} \\ &= \sum_{i+j=m-2} \nabla_t^i (R(\partial_t, \cdot) - \nabla_{\Theta(X)}) \nabla_t^{j+1} \mathcal{L}_t g(X, V, Y)|_{\mathcal{H}} \\ &= \sum_{i+j=m-2} \nabla_t^i R * \nabla_t^{j+1} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} \\ &\quad + \sum_{i+j=m-2} \sum_{a=1}^i \nabla_t^a \Theta * \nabla_t^{i-a} \nabla \nabla_t^{j+1} \mathcal{L}_t g(X, Y)|_{\mathcal{H}} \\ &\quad - \sum_{i+j=m-2} \nabla_{\underbrace{t, \dots, t}_{i \text{ times}}, \underbrace{t, \Theta(X), t, \dots, t}_{j+1 \text{ times}}}^m \mathcal{L}_t g(V, Y)|_{\mathcal{H}}. \end{aligned}$$

Applying Lemma 3.17, note that only the last term is not obviously given by (17), we compute it separately:

$$\begin{aligned} &\sum_{i+j=m-2} \nabla_{\underbrace{t, \dots, t}_{i \text{ times}}, \underbrace{t, \Theta(X), t, \dots, t}_{j+1 \text{ times}}}^m \mathcal{L}_t g(V, Y)|_{\mathcal{H}} \\ &= \sum_{i+j=m-2} [\nabla_t^i, \nabla] \nabla_t^{j+1} \mathcal{L}_t g(\Theta(X), V, Y)|_{\mathcal{H}} + (m-1) \nabla_{\Theta(X), t, \dots, t}^m \mathcal{L}_t g(V, Y)|_{\mathcal{H}}. \end{aligned}$$

Arguing as above for $[\nabla_t^i, \nabla]$, we see that the sum is given by (17). Since $X \in C^\infty(V^\perp)$, we know that $\Theta(X) \in C^\infty(T\mathcal{H})$, which implies that the second term in the above expression is given by (17). Thus we have proven that the first and second terms in (19) are given by (17).

We turn to the remaining terms in (19). Note first that

$$\begin{aligned} \nabla_{t,\dots,t,\cdot,\cdot}^{m+1} \mathcal{L}_t g &= [\nabla_t^{m-1}, \nabla^2] \mathcal{L}_t g + \nabla^2 \nabla_t^{m-1} \mathcal{L}_t g \\ &= [\nabla_t^{m-1}, \nabla] \nabla \mathcal{L}_t g + \nabla [\nabla_t^{m-1}, \nabla] \mathcal{L}_t g + \nabla^2 \nabla_t^{m-1} \mathcal{L}_t g \\ &= [[\nabla_t^{m-1}, \nabla], \nabla] \mathcal{L}_t g + 2 \nabla [\nabla_t^{m-1}, \nabla] \mathcal{L}_t g + \nabla^2 \nabla_t^{m-1} \mathcal{L}_t g. \end{aligned}$$

Analogous to above, one checks that $[\nabla_t^{m-1}, \nabla] \mathcal{L}_t g$ and $\nabla_t^{m-1} \mathcal{L}_t g$ are given by (17). Hence for $X, Y \in C^\infty(T\mathcal{H})$, we have $\nabla_X Y \in T\mathcal{H}$ and we conclude that

$$2 \nabla_X [\nabla_t^{m-1}, \nabla] \mathcal{L}_t g(Y)|_{\mathcal{H}} + \nabla_{X,Y}^2 \nabla_t^{m-1} \mathcal{L}_t g|_{\mathcal{H}}$$

is given by (17). We also note that

$$[[\nabla_t^{m-1}, \nabla], \nabla] \mathcal{L}_t g = \sum_{i+j=m-2} [\nabla_t^i R * \nabla_t^j, \nabla] \mathcal{L}_t g + \sum_{i+j=m-2} [\nabla_t^i \Theta * \nabla_{t,\dots,t,\cdot,\cdot}^{j+1}, \nabla] \mathcal{L}_t g.$$

Applying Lemma 3.17 gives, by the same arguments as above, that also this is given by (17). We conclude that for any $X, Y \in C^\infty(T\mathcal{H})$

$$\nabla_{t,\dots,t,X,Y} \mathcal{L}_t g|_{\mathcal{H}}$$

is given by (17). This completes the proof. \square

We may now prove Proposition 3.23:

Proof of Proposition 3.23. Combining equation (18) with Proposition 3.21, Lemma 3.24, Lemma 3.25 and Lemma 3.26, we conclude that for all $X, Y \in V^\perp$

$$2\nabla_t^{m-1}\mathcal{L}_t\text{Ric}(X, Y)|_{\mathcal{H}} + 2(m+1)\kappa\mathcal{L}_t^{m+1}g(X, Y)|_{\mathcal{H}}$$

is given uniquely by (17). Since by Assumption 3.1, we have

$$2\nabla_t^{m-1}\mathcal{L}_t\text{Ric}(X, Y)|_{\mathcal{H}} = 0$$

and $\kappa \neq 0$, we conclude that

$$\mathcal{L}_t^{m+1}g(X, Y)|_{\mathcal{H}}$$

is given uniquely by (17). This completes the proof. \square

We may finally conclude the proof of Theorem 3.9:

Proof of Theorem 3.9. As we pointed out in Remark 3.15, Proposition 3.10 implies that the assertion in Induction Assumption 3.14 for

$$m = 1.$$

We thus assume that Induction Assumption 3.14 is true for a fixed

$$m \in \mathbb{N}$$

and aim to prove the assertion in Induction Assumption 3.14 for

$$m + 1.$$

By Proposition 3.20, we know that there is a unique way to express the components

$$\mathcal{L}_t^{m+1}g(X, V)|_{\mathcal{H}}$$

in terms of $g|_{\mathcal{H}}, \dots, \mathcal{L}_t^m g|_{\mathcal{H}}$. Since by Induction Assumption 3.14, we know that

$$\mathcal{L}_t^k g(X, Y)|_{\mathcal{H}} = Q_k(\sigma, V)(X, Y)$$

for all $k = 0, \dots, m$ and $X, Y \in T\mathcal{H}$, and since Lemma 3.8 implies that

$$\mathcal{L}_t^k g(\partial_t, \cdot)|_{\mathcal{H}} = 0$$

for all $k \in \mathbb{N}$, we conclude that there is a unique $Q_{1,m+1}(\sigma, V)$ such that

$$\mathcal{L}_t^{m+1}g(V, X)|_{\mathcal{H}} = Q_{1,m+1}(\sigma, V)(V, X)$$

for all $X \in T\mathcal{H}$. Similarly, by Proposition 3.23, we conclude that there is a $Q_{2,m+1}(\sigma, V)$ such that

$$\mathcal{L}_t^{m+1}g(X, Y)|_{\mathcal{H}} = Q_{2,m+1}(\sigma, V)(X, Y)$$

for all $X, Y \in V^\perp$. To sum up, we have shown that there is a unique $Q_{m+1}(\sigma, V)$ such that

$$\mathcal{L}_t^{m+1}g(X, Y)|_{\mathcal{H}} = Q_{m+1}(\sigma, V)(X, Y)$$

for all $X, Y \in T\mathcal{H}$. Finally, note that Q_{m+1} constructed this way is diffeomorphism invariant. This completes the induction argument and finishes the proof. \square

4. GEOMETRIC UNIQUENESS

The goal of this section is to prove Theorem 1.8. In this section, objects and maps will be *real analytic*, as opposed to only smooth as in the previous section. We are given two real analytic vacuum solutions

$$\iota : \mathcal{H} \rightarrow (M, g), \quad \hat{\iota} : \hat{\mathcal{H}} \rightarrow (\hat{M}, \hat{g}),$$

where the induced real analytic initial data (\mathcal{H}, σ, V) and $(\hat{\mathcal{H}}, \hat{\sigma}, \hat{V})$ are related by the isometry

$$\varphi : (\mathcal{H}, \sigma) \rightarrow (\hat{\mathcal{H}}, \hat{\sigma})$$

such that

$$d\varphi(V) = \hat{V}.$$

We want to construct open neighborhoods $\mathcal{H} \subset \mathcal{U} \subset M$ and $\hat{\mathcal{H}} \subset \hat{\mathcal{U}} \subset \hat{M}$ and an isometry

$$\Phi : (\mathcal{U}, g|_{\mathcal{U}}) \rightarrow (\hat{\mathcal{U}}, \hat{g}|_{\hat{\mathcal{U}}})$$

such that

$$\Phi|_{\iota(\mathcal{H})} = \hat{\iota} \circ \varphi \circ \iota^{-1} : \iota(\mathcal{H}) \rightarrow \hat{\iota}(\hat{\mathcal{H}}).$$

The question is how Φ should be constructed away from $\iota(\mathcal{H})$? If $\iota(\mathcal{H}) \subset M$ was a spacelike or timelike hypersurface, a natural way to construct Φ near $\iota(\mathcal{H})$ would be to map the unit speed geodesics normal to $\iota(\mathcal{H})$ to the unit speed geodesics normal to $\hat{\iota}(\hat{\mathcal{H}})$. Since $\iota(\mathcal{H}) \subset M$ is a *lightlike* hypersurface, the normal vector field is *tangent* to $\iota(\mathcal{H})$, so that method does not work. However, recall from Proposition 3.4 that there is a *geometrically canonical transversal* vector field $\partial_t|_{\iota(\mathcal{H})}$ along $\iota(\mathcal{H})$ and $\partial_{\hat{t}}|_{\hat{\iota}(\hat{\mathcal{H}})}$ along $\hat{\iota}(\hat{\mathcal{H}})$, which plays an analogous role as a unit normal does to a time-like or space-like hypersurface. This idea together with the iterative determination of the asymptotic expansion, Theorem 3.9, are the keys in our proof of Theorem 1.8.

4.1. Construction of the candidate isometry. We let ∂_t and $\partial_{\hat{t}}$ denote the real analytic vector fields given by Proposition 3.4, defined in open neighbourhoods

$$\mathcal{H} \subset \mathcal{U} \subset M, \quad \hat{\mathcal{H}} \subset \hat{\mathcal{U}} \subset \hat{M},$$

respectively.

Proposition 4.1 (The candidate isometry). *Assume the same as in Theorem 1.8. Shrinking \mathcal{U} and $\hat{\mathcal{U}}$ if necessary, there is a unique real analytic diffeomorphism (with real analytic inverse)*

$$\Phi : \mathcal{U} \rightarrow \hat{\mathcal{U}}$$

such that

$$\Phi|_{\iota(\mathcal{H})} = \hat{\iota} \circ \varphi \circ \iota^{-1}, \tag{21}$$

$$d\Phi(\partial_t) = \partial_{\hat{t}}. \tag{22}$$

Proof. Let $q \in \mathcal{U}$ be given. By construction, there is a unique integral curve of ∂_t , starting at some point $p \in \iota(\mathcal{H})$ and reaching q after eigentime $t(q)$. Let $\hat{p} := \varphi(p) \in \hat{\iota}(\hat{\mathcal{H}})$ and consider the integral curve of $\partial_{\hat{t}}$ in $\hat{\mathcal{U}}$. Let $\hat{q} \in \hat{\mathcal{U}}$ denote point which the integral curve of $\partial_{\hat{t}}$ reaches after time $t(q)$, i.e. the unique point on the integral curve starting in \hat{p} with

$$\hat{t}(\hat{q}) = t(q).$$

In order for this to be defined, we shrink \mathcal{U} if necessary, i.e. assume that p is sufficiently near $\iota(\mathcal{H})$. For each $q \in \mathcal{U}$, set

$$\Phi(q) := \hat{q}.$$

It follows that Φ is real analytic and we may shrink $\hat{\mathcal{U}}$ to make sure that Φ is bijective. The inverse is real analytic, since it is constructed analogously. \square

Remark 4.2. Note that if we had already proven the assertion in Theorem 1.8, then Φ would necessarily be as in Proposition 4.1. Indeed, condition (21) is required in Theorem 1.8 and any isometry has to map the geometrically defined vector field $\partial_t|_{\iota(\mathcal{H})}$ along $\iota(\mathcal{H})$ to $\partial_{\hat{t}}|_{\hat{\iota}(\hat{\mathcal{H}})}$ along $\hat{\iota}(\hat{\mathcal{H}})$. Recalling that ∂_t and $\partial_{\hat{t}}$ are *geodesic* vector fields, Proposition 4.1 is just saying that the geodesics emanating from $\partial_t|_{\iota(\mathcal{H})}$ have to be mapped to the geodesics emanating from $\partial_{\hat{t}}|_{\hat{\iota}(\hat{\mathcal{H}})}$, which is automatic if Φ is an isometry.

4.2. Proof of geometric uniqueness. We may finally prove Theorem 1.8, which says that Φ is an isometry after shrinking \mathcal{U} and $\hat{\mathcal{U}}$ if necessary.

Proof of Theorem 1.8. We claim that

$$\mathcal{L}_t^m (\Phi^* \hat{g} - g)|_{\iota(\mathcal{H})} = 0 \quad (23)$$

for all $m \in \mathbb{N}_0$. By real analyticity, this will suffice to prove the assertion. We begin with $m = 0$. Note that

$$\begin{aligned} d\Phi(\partial_t) &= \partial_{\hat{t}}, \\ d\Phi(d\iota(V)) &= d\hat{\iota} \circ d\varphi(V) \\ &= d\hat{\iota}(\hat{V}) \end{aligned}$$

and

$$d\Phi|_{d\iota(V^\perp)} = d\hat{\iota} \circ d\varphi \circ (d\iota)^{-1} : d\iota(V^\perp) \rightarrow d\hat{\iota}(\hat{V}^\perp) \quad (24)$$

is an isomorphism. What remains is therefore to check that (24) is an isometry. Given $X, Y \in V^\perp$, we use the assumption $\varphi^* \hat{\sigma} = \sigma$ to compute

$$\begin{aligned} \Phi^* \hat{g}(d\iota(X), d\iota(Y)) &= \hat{g}(d\Phi \circ d\iota(X), d\Phi \circ d\iota(Y)) \\ &= \hat{g}(d\hat{\iota} \circ d\varphi(X), d\hat{\iota} \circ d\varphi(Y)) \\ &= \hat{\sigma}(d\varphi(X), d\varphi(Y)) \\ &= \sigma(X, Y) \\ &= g(d\iota(X), d\iota(Y)). \end{aligned}$$

This completes the proof of (23) when $m = 0$. Let us now turn to the case $m \geq 1$. Proposition 4.1 implies that

$$\mathcal{L}_t \Phi^* = \Phi^* \mathcal{L}_{d\Phi(\partial_t)} = \Phi^* \mathcal{L}_{\partial_{\hat{t}}} = \Phi^* \mathcal{L}_{\hat{t}}$$

and hence

$$\mathcal{L}_t^m \Phi^* = \Phi^* \mathcal{L}_{\hat{t}}^m$$

for all $m \in \mathbb{N}_0$. The claim (23) can thus be rewritten as

$$\Phi^* \mathcal{L}_{\hat{t}}^m \hat{g}|_{\iota(\mathcal{H})} = \mathcal{L}_t^m g|_{\iota(\mathcal{H})}. \quad (25)$$

We first prove that

$$\Phi^* \mathcal{L}_{\hat{t}}^m \hat{g}(\partial_t, \cdot)|_{\iota(\mathcal{H})} = \mathcal{L}_t^m g(\partial_t, \cdot)|_{\iota(\mathcal{H})}. \quad (26)$$

The right hand side of this is vanishing by Lemma 3.8. The left hand side is computed using Proposition 4.1 as

$$\begin{aligned} \Phi^* \mathcal{L}_{\hat{t}}^m \hat{g}(\partial_t, \cdot)|_{\iota(\mathcal{H})} &= \mathcal{L}_{\hat{t}}^m \hat{g}(d\Phi(\partial_t), d\Phi(\cdot))|_{\Phi(\iota(\mathcal{H}))} \\ &= \mathcal{L}_{\hat{t}}^m \hat{g}(\partial_{\hat{t}}, d\Phi(\cdot))|_{\hat{\iota}(\hat{\mathcal{H}})}, \end{aligned}$$

which is vanishing as well by Lemma 3.8, proving (26). We now prove that

$$\iota^* \Phi^* \mathcal{L}_{\hat{t}}^m \hat{g} = \iota^* \mathcal{L}_t^m g, \quad (27)$$

which together with (26) would prove (25) and thus prove (23). By Proposition 4.1 and Theorem 3.9, we have

$$\begin{aligned}
\iota^* \Phi^* \mathcal{L}_t^m \hat{g} &= (\Phi \circ \iota)^* \mathcal{L}_t^m \hat{g} \\
&= (\hat{\iota} \circ \varphi)^* \mathcal{L}_t^m \hat{g} \\
&= \varphi^* \hat{\iota}^* \mathcal{L}_t^m \hat{g} \\
&= \varphi^* Q_m(\hat{\sigma}, \hat{V}) \\
&= Q_m(\varphi^* \hat{\sigma}, \varphi^* \hat{V}) \\
&= Q_m(\sigma, V) \\
&= \iota^* \mathcal{L}_t^m g,
\end{aligned}$$

which completes the proof of (27). This completes the proof of (23). The assertion in Theorem 1.8 now follows by real analyticity. \square

5. EXISTENCE

The goal of this section is to prove Theorem 1.7. Moncrief proved a local version of this in [Mon82], which we recall here in the form we need it. We combine this with our geometric uniqueness result, Theorem 1.8, in order to allow for arbitrary topology and conclude Theorem 1.7.

5.1. Moncrief's existence theorem. We use a slightly different notation than in [Mon82] to better match the notation of the present paper:

Theorem 5.1 (A version of Moncrief's existence theorem). *Let $\mathcal{V} \subset \mathbb{R}^2$ be an open region with coordinates x_2, x_3 . Given a real analytic function $\check{\phi}$, a real analytic one-form $\check{\beta}$ and a real analytic Riemannian metric \check{h} on \mathcal{V} , there is a unique real analytic vacuum metric of the form*

$$g := e^{-2\phi} \left(-\frac{N^2 d\tau^2}{4\tau} + h_{ab} dx^a dx^b \right) + \tau e^{2\phi} \left(\frac{1}{2\tau} d\tau + \frac{1}{2} dx^1 + \beta_a dx^a \right)^2, \quad (28)$$

on

$$M := (-\epsilon, \epsilon)_\tau \times S_{x_1}^1 \times \mathcal{V}_{x_2, x_3},$$

where repeated Latin indices are summed over 2, 3 and where

$$N := \frac{e^{2\check{\phi}}}{\sqrt{\det(\check{h}_{ij})}} \sqrt{\det(h_{ij})}$$

and where ϕ, β and h are a real analytic function, one-form and Riemannian metric, respectively, such that

$$\begin{aligned}
\partial_{x_1} \phi &= 0, & \mathcal{L}_{\partial_{x_1}} \beta &= 0, & \mathcal{L}_{\partial_{x_1}} h &= 0, \\
\phi|_{\tau=0, x_1=c} &= \check{\phi}, & \beta|_{\tau=0, x_1=c} &= \check{\beta}, & h|_{\tau=0, x_1=c} &= \check{h},
\end{aligned}$$

for any $c \in S^1$. In particular, the spacetime

$$(M, g)$$

is a real analytic vacuum spacetime with a non-degenerate Killing horizon

$$\{\tau = 0\} \times S_{x_1}^1 \times \mathcal{V}_{x_2, x_3}.$$

Proof. In case we replace \mathcal{V} with T^2 , this is precisely the statement in Theorem [Mon82, Thm. 2]. However, Moncrief proves the statement of Theorem 5.1 as an intermediate step, see [Mon82, p. 90]. \square

5.2. Proof of existence. We use Theorem 5.1 to prove a local version of our Theorem 1.7:

Lemma 5.2. *Let (\mathcal{H}, σ, V) be real analytic initial data, in the sense of Definition 1.5, with a 3-dimensional \mathcal{H} . For every $p \in \mathcal{H}$, there is an open subset*

$$p \in \mathcal{U} \subseteq \mathcal{H}$$

and a real analytic spacetime (M, g) , a real analytic Killing vector field W on M and a real analytic non-degenerate Killing horizon

$$\iota : \mathcal{U} \hookrightarrow M,$$

with respect to W , such that

- $(\mathcal{U}, \sigma|_{\mathcal{U}}, V|_{\mathcal{U}})$ is the induced initial data as in Definition 1.4,
- $d\iota(V) = W|_{\iota(\mathcal{U})}$.

Proof. Let \mathcal{U} be a coordinate neighborhood around p , such that

$$\partial_{x_1} = -\frac{V}{2\sqrt{\sigma(V, V)}}.$$

We get

$$\sigma|_{\mathcal{U}} = \sigma_{ij} dx^i dx^j,$$

where σ_{ij} are independent of x_1 . We may thus extend σ to a metric on

$$S^1_{x_1} \times \mathcal{V}_{x_2, x_3},$$

by the same formula, since σ_{ij} are independent of x_1 . We get an isometric embedding

$$\mathcal{U} \hookrightarrow S^1_{x_1} \times \mathcal{V}_{x_2, x_3},$$

which respects the Killing vector field. Note that it suffices to prove the assertion for the latter, so we can without loss of generality assume that

$$\mathcal{U} = S^1_{x_1} \times \mathcal{V}_{x_2, x_3}.$$

Recall that we want

$$\sigma|_{\mathcal{U}} = \iota^* g|_{\mathcal{U}} + \omega^2|_{\mathcal{U}}.$$

Recall from Section 1.2.3 that σ and V determine $\iota^* g$ and ω uniquely at every point on the horizon. We therefore need to compute what $\iota^* g$ and ω^2 are in Moncrief's expression (28). Setting $\tau = 0$ and disregarding the components containing $d\tau$ in g , we get

$$\iota^* g = e^{-2\tilde{\phi}} \tilde{h}_{ab} dx^a dx^b. \quad (29)$$

In order to compute ω , we recall the defining equation

$$\begin{aligned} \omega(X)V &= \nabla_X W|_{\mathcal{H}} \\ &= -\frac{1}{2\sqrt{\sigma(V, V)}} \nabla_{\sum_{i=1}^3 X^i \partial_{x_i}} \partial_{x_1}|_{\tau=0} \\ &= \sum_{i=1}^3 X^i \Gamma_{i1}^1 V. \end{aligned}$$

We compute

$$\begin{aligned} \Gamma_{i1}^1|_{\tau=0} &= \frac{1}{2} \sum_{\alpha=0}^3 g^{1\alpha} (\partial_i g_{\alpha 1} + \partial_1 g_{i\alpha} - \partial_\alpha g_{i1})|_{\tau=0} \\ &= \frac{1}{2} g^{10} (\partial_i g_{01} - \partial_0 g_{i1})|_{\tau=0}, \end{aligned}$$

where $x_0 := \tau$. Using that

$$g_{10}|_{\tau=0} = \frac{e^{2\dot{\phi}}}{4}, \quad g_{1i}|_{\tau=0} = 0,$$

for $i = 1, 2, 3$, we note that

$$1 = \sum_{\alpha=0}^3 g_{1\alpha} g^{\alpha 1}|_{\tau=0} = \frac{e^{2\dot{\phi}}}{4} g^{01}|_{\tau=0},$$

from which it follows that

$$g^{01}|_{\tau=0} = 4e^{-2\dot{\phi}}, \quad \partial_i g_{01}|_{\tau=0} = \frac{e^{2\dot{\phi}}}{2} \partial_i \dot{\phi}.$$

It remains to compute, for $a = 2, 3$,

$$\begin{aligned} -\partial_0 g_{11}|_{t=0} &= -\partial_\tau \left(\frac{\tau e^{2\dot{\phi}}}{4} \right) |_{\tau=0} = -\frac{e^{2\dot{\phi}}}{4} \\ -\partial_0 g_{a1}|_{t=0} &= -\partial_\tau \left(\frac{\tau \beta_a}{2} \right) |_{\tau=0} = -\frac{\dot{\beta}_a}{2}. \end{aligned}$$

Putting this together, we conclude that

$$\omega = -\frac{1}{2} dx^1 + (\partial_2 \dot{\phi} - \dot{\beta}_2 e^{-2\dot{\phi}}) dx^2 + (\partial_3 \dot{\phi} - \dot{\beta}_3 e^{-2\dot{\phi}}) dx^3. \quad (30)$$

Recall that $\iota^* g$ and ω are given and we want to construct solutions to (29) and (30). Note, however, that this system is underdetermined. Indeed, if we choose $\dot{\phi} = 0$, there is a unique way of choosing \dot{h}_{ab} and $\dot{\beta}_a$ satisfying (29) and (30). By Theorem 5.1, we conclude that there is a spacetime of the form (28), such that

$$\sigma|_{\mathcal{U}} = \iota^* g|_{\mathcal{U}} + \omega^2|_{\mathcal{H}}.$$

We finally note that

$$\kappa = \omega(V) = 2\sqrt{\sigma(V, V)} \omega(\partial_{x_1}) = \sqrt{\sigma(V, V)}.$$

This completes the proof. \square

Remark 5.3. We proved in Theorem 1.8 that the initial data (\mathcal{H}, σ, V) characterizes the vacuum spacetime uniquely near the horizon. As can be seen in the proof of Lemma 5.2, there are many choices of initial data in Theorem 5.1 that lead to isometric (i.e. the same) vacuum spacetime, i.e. we can without loss of generality set $\dot{\phi} = 0$.

We may now prove Theorem 1.7 by patching together Moncrief's local existence result in a cover of the horizon of open subsets, using our Theorem 1.8:

Proof of Theorem 1.7. We are given initial data

$$(\mathcal{H}, \sigma, V),$$

and we would like to define the real analytic spacetime metric g on the manifold

$$\mathbb{R} \times \mathcal{H},$$

in an open neighborhood of $\{0\} \times \mathcal{H}$. Let $p \in \mathcal{H}$. By Lemma 5.2, there is an open neighborhood $\mathcal{V}_p \subset \mathcal{H}$ of p and a spacetime M_p such that

$$\iota_p : \mathcal{V}_p \rightarrow M_p$$

is a non-degenerate Killing horizon with induced initial data (as in Definition 1.4)

$$(\mathcal{V}_p, \sigma|_{\mathcal{V}_p}, V|_{\mathcal{V}_p}).$$

By shrinking \mathcal{V}_p and M_p if necessary, Proposition 3.4 and Proposition 3.5 provide a diffeomorphism

$$M_p \cong (-\epsilon_p, \epsilon_p) \times \mathcal{V}_p$$

for a small $\epsilon_p > 0$, such that ∂_t is tangent to the curves $(-\epsilon, \epsilon) \times \{x\}$, with $x \in \mathcal{V}_p$. We can now use this to *formally* construct the vacuum metric g in an open neighborhood of

$$\{0\} \times \mathcal{H} \subset \mathbb{R} \times \mathcal{H}.$$

It remains to check that this is well-defined, i.e. if $q \in \mathcal{V}_p$ with $q \neq p$, that the metrics on

$$(-\epsilon_p, \epsilon_p) \times \mathcal{V}_p \cap (-\epsilon_q, \epsilon_q) \times \mathcal{V}_q$$

coincide. We know that the initial data coincide on \mathcal{V}_p and \mathcal{V}_q . It thus follows by Theorem 1.8 that the resulting spacetimes are isometric in a neighbourhood of any point on the horizon. This isometry is explicitly given by the null time function, which means that the isometry is the identity map

$$\text{id} : (-\epsilon_p, \epsilon_p) \times \mathcal{V}_p \cap (-\epsilon_q, \epsilon_q) \times \mathcal{V}_q \rightarrow (-\epsilon_p, \epsilon_p) \times \mathcal{V}_p \cap (-\epsilon_q, \epsilon_q) \times \mathcal{V}_q.$$

In other words, the metrics coincide and g is well-defined. This concludes the proof. \square

REFERENCES

- [AIK10] S. Alexakis, A. D. Ionescu, and S. Klainerman, *Hawking's local rigidity theorem without analyticity*, Geom. Funct. Anal. **20** (2010), no. 4, 845–869.
- [Bes08] A. L. Besse, *Einstein manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition.
- [BR21] I. Bustamante and M. Reiris, *A classification theorem for compact Cauchy horizons in vacuum spacetimes*, Gen. Relativity Gravitation **53** (2021), no. 3, Paper No. 36, 10.
- [CC08] P. T. Chruściel and J. L. Costa, *On uniqueness of stationary vacuum black holes*, Astérisque **321** (2008), 195–265.
- [YFB52] Y. Fourès-Bruhat, *Théorème d'existence pour certains systèmes d'équations aux dérivées partielles non linéaires.*, Acta Math. **88** (1952), 141–225.
- [GM21] S. Gurriaran and E. Minguzzi, *Surface gravity of compact non-degenerate horizons under the dominant energy condition*, Preprint: arXiv:2108.04056 (2021).
- [Haw72] S. W. Hawking, *Black holes in general relativity*, Comm. Math. Phys. **25** (1972), 152–166.
- [HE73] S. W. Hawking and G. F. R. Ellis, *The large scale structure of space-time*, Cambridge University Press, London-New York, 1973. Cambridge Monographs on Mathematical Physics, No. 1.
- [HIW07] S. Hollands, A. Ishibashi, and R. M. Wald, *A higher dimensional stationary rotating black hole must be axisymmetric*, Comm. Math. Phys. **271** (2007), no. 3, 699–722.
- [IK09] A. D. Ionescu and S. Klainerman, *On the uniqueness of smooth, stationary black holes in vacuum*, Invent. Math. **175** (2009), no. 1, 35–102.
- [IK13] ———, *On the local extension of Killing vector-fields in Ricci flat manifolds*, J. Amer. Math. Soc. **26** (2013), no. 2, 563–593.
- [IM85] J. Isenberg and V. Moncrief, *Symmetries of cosmological Cauchy horizons with exceptional orbits*, J. Math. Phys. **26** (1985), no. 5, 1024–1027.
- [IM92] ———, *On spacetimes containing Killing vector fields with nonclosed orbits*, Classical Quantum Gravity **9** (1992), no. 7, 1683–1691.
- [Lar15] E. Larsson, *Smoothness of compact horizons*, Ann. Henri Poincaré **16** (2015), no. 9, 2163–2214.
- [Min15] E. Minguzzi, *Area theorem and smoothness of compact Cauchy horizons*, Comm. Math. Phys. **339** (2015), no. 1, 57–98.
- [Mon82] V. Moncrief, *Neighborhoods of Cauchy horizons in cosmological spacetimes with one Killing field*, Ann. Physics **141** (1982), no. 1, 83–103.
- [Mon84] ———, *The space of (generalized) Taub-NUT spacetimes*, J. Geom. Phys. **1** (1984), no. 1, 107–130.
- [MI83] V. Moncrief and J. Isenberg, *Symmetries of cosmological Cauchy horizons*, Comm. Math. Phys. **89** (1983), no. 3, 387–413.

- [MI08] ———, *Symmetries of higher dimensional black holes*, Classical Quantum Gravity **25** (2008), no. 19, 195015, 37.
- [MI20] ———, *Symmetries of cosmological Cauchy horizons with non-closed orbits*, Comm. Math. Phys. **374** (2020), no. 1, 145–186.
- [O’N83] B. O’Neill, *Semi-Riemannian geometry*, Pure and Applied Mathematics, vol. 103, Academic Press, San Diego, 1983. With applications to relativity.
- [Pet21a] O. L. Petersen, *Extension of Killing vector fields beyond compact Cauchy horizons*, Adv. Math. **391** (2021), Paper No. 107953.
- [Pet21b] ———, *Wave equations with initial data on compact Cauchy horizons*, Anal. PDE, to appear (2021).
- [PR18] O. L. Petersen and I. Rácz, *Symmetries of vacuum spacetimes with a compact Cauchy horizon of constant non-zero surface gravity*, Preprint: arXiv: 1809.02580 (2018).
- [PV21] O. L. Petersen and A. Vasy, *Analyticity of quasinormal modes in the Kerr and Kerr-de Sitter spacetimes*, Preprint: arXiv:2104.04500 (2021).
- [RB21] M. Reiris and I. Bustamente, *On the existence of Killing fields in smooth spacetimes with a compact Cauchy horizon*, Classical Quantum Gravity **38** (2021), no. 7.

UNIVERSITY OF HAMBURG, DEPARTMENT OF MATHEMATICS, BUNDESSTRASSE 55, 20146 HAMBURG, GERMANY

Email address: klaus.kroencke@uni-hamburg.de

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, CA 94305-2125, USA

Email address: oliverlp@stanford.edu