# How to Find New Characteristic-Dependent Linear Rank Inequalities using Secret Sharing

Victor Peña-Macias\* Departamento de Matemáticas Universidad Nacional de Colombia Bogotá, Colombia

#### Abstract

Determining information ratios of access structures is an important problem in secret sharing. Information inequalities and linear rank inequalities play an important role for proving bounds. Characteristic-dependent linear rank inequalities are rank inequalities which are true over vector spaces with specific field characteristic. In this paper, using ideas of secret sharing, we show a theorem that produces characteristic-dependent linear rank inequalities. These inequalities can be used for getting lower bounds on information ratios in linear secret sharing.

#### Mathematics Subject Classification: 68P30

**Keywords:** Secret sharing, cryptography, access structure, linear rank inequalities, matroids.

## **1** Introduction

Secret sharing is a cryptographic protocol that consists of distributing a *secret* in several messages or *shares* within a group of participants, in such a way that if a group of participants with access to the secret shares their messages, they can discover the secret; but if a group of participants that does not have access to the secret shares their messages, they cannot get any information about the secret [2, 7, 10, 14]. A specific protocol with this property is called the *secret sharing scheme*, and the collection of participants with access to the secret is called the *access structure*. The efficiency of a scheme is measured by *information ratios* which relates the size of the secret and the size of the shares. In secret sharing, it is important to build efficient secret sharing schemes on an access structure. Therefore, determining the best information ratio, known as the *optimal information ratio*, is an important task.

<sup>\*</sup>E-mail: vbpenam@unal.edu.co

A *linear rank inequality* is a linear inequality that is always satisfied by ranks (dimensions) of subspaces of a vector space over any field. *Information inequalities* are a sub-class of linear rank inequalities [15]. A *characteristic-dependent linear rank inequality* is like a linear rank inequality but this is always satisfied by vector spaces over fields of certain characteristic and does not in general hold over other characteristics [3, 11, 12]. Information inequalities have been useful to estimate lower bounds on the optimal information ratio of access structures, and linear rank inequalities have been useful to estimate lower bounds on the optimal information ratio of access structures in *linear secret sharing schemes*, i.e. when secret sharing schemes have a linear structure [4]. To the best of our knowledge, characteristic-dependent linear rank inequalities have not been used for determining bounds in linear secret sharing schemes in specific finite fields, but due to the nature of distinguishing finite fields according to their characteristics, these inequalities can be useful. One area where these inequalities have been useful for determing bounds is in *newtork coding* [1, 3, 5, 11, 13].

**Contributions.** In [7], Jafari and S. Khazaei developed a technique for proving lower bounds on access structures in linear secret sharing schemes on finite fields with a specific characteristic. They present their technique using access structures or matroid ports associated with the Fano and non-Fano matroids. We note that this technique can be improved in order for producing characteristicdependent linear rank inequalities that also imply lower bounds on information ratios in linear secret sharing. For any binary matrix whose determinant is greater than 1, we studied some matroid ports associated with the representable matroid of this matrix. Since the matrix defines different matroids according to the field where it is defined, we get different access structures; we establish the properties that depend on the characteristic of the finite field associated with the matrix. These properties serve as a guide to define conditions and inequalities that must satisfy vector spaces of a specific field characteristic. Then, using the vector deletion technique of Blasiak et al. in [1], which was improved in [11, 12], we get a theorem that produces characteristic-dependent linear rank inequalities. We emphasize that this theorem produces a pair of inequalities as long as there is a binary matrix whose determinant is greater than 1. We also show a class of matrices that satisfy this property and produce  $2\lfloor \frac{n-1}{2} \rfloor - 4$  inequalities for each  $n \ge 7$  and we compute some lower bounds of optimal information ratios associated with matroid ports of these matrices over specific fields.

**Organization of the work.** In section 2 and 3, we study concepts of information theory and secret sharing. In section 4, we show our method for producing characteristic-dependent linear rank inequalities; this method is summarized with a theorem. In section 5, some inequalities are produced and we estimate lower bounds on information ratios of some access structures in linear secret sharing.

#### **2** Information Theory

Let  $A_1, \ldots, A_n$  be vector subspaces of a finite dimensional vector space V over a finite field  $\mathbb{F}$ . Let  $\sum A_i$  be the span of  $A_i, i \in I$ . We are interested in tuples of random variables associated to these

vector spaces, for details about the usual construction of these variables, see [10, 12, 15]; we refer as *linear random variables*. There is a correspondence between entropy of linear random variables and dimension of vector spaces [15, Theorem 2]. So, we identify the entropy of linear random variables with the dimension of the associated spaces, i.e.

$$\mathrm{H}\left(A_{i}: i \in I\right) := \dim\left(\sum A_{i}\right).$$

The *mutual information* of  $A_1$  and  $A_2$  is given by

$$I(A_1; A_2) := \dim (A_1 \cap A_2).$$

The codimension of  $A_1$  in V is given by  $\operatorname{codim}_V(A_1) = \dim(V) - \dim(A_1)$ , we have

$$\mathrm{H}\left(A_{1} \mid A_{2}\right) := \operatorname{codim}_{A_{1}}\left(A_{1} \cap A_{2}\right).$$

The conditional mutual information is expressed as

$$I(A_1; A_2 | A_3) := \dim (A_1 + A_3) - \operatorname{codim}_{A_1} (A_1 \cap A_2 \cap A_3).$$

The following definition is given to fix ideas about inequalities and vector spaces.

**Definition 1.** Let P be a proper subset of prime numbers and  $I_1, \ldots, I_k \subseteq [n]$ . Let  $\alpha_i \in \mathbb{R}$ , for  $1 \le i \le k$ . Consider a linear inequality of the form  $\sum \alpha_i \operatorname{H} (A_j : j \in I_i) \ge 0$ . The inequality is called a *characteristic-dependent linear rank inequality* if it holds for all vector spaces  $A_1, \ldots, A_n$  over a finite field whose characteristic is in P, and does not in general hold over other characteristics. Besides, the inequality is called a *linear rank inequality* if it holds for all vector spaces.

The following linear rank inequality is called Ingleton's inequality [6]. For any  $A_1$ ,  $A_2$ ,  $A_3$ ,  $A_4$  subspaces of a finite dimensional vector space,

$$I(A_1; A_2) \le I(A_1; A_2 | A_3) + I(A_1; A_2 | A_4) + I(A_3; A_4).$$

The following inequality [3] is a characteristic-dependent linear rank inequalities over fields with characteristic other than 2:

$$2 \operatorname{H} (A_{1}) + \operatorname{H} (A_{2}) + 2 \operatorname{H} (A_{3}) \leq \operatorname{H} (B_{1}) + \operatorname{H} (B_{2}) + \operatorname{H} (B_{3}) + \operatorname{H} (C)$$
  
+2 \overline{H} (A\_{1} | B\_{1}, C) + \overline{H} (A\_{2} | B\_{2}, C) + 2 \operatorname{H} (A\_{3} | A\_{1}, B\_{2})  
+3 \overline{H} (B\_{2} | B\_{1}, B\_{3}) + 3 \operatorname{H} (C | A\_{3}, B\_{3}) + 5 \operatorname{H} (B\_{3} | A\_{1}, A\_{2}) + 5 \operatorname{H} (B\_{1} | A\_{2}, A\_{3})  
+5 (\overline{H} (A\_{1}) + \overline{H} (A\_{2}) + \operatorname{H} (A\_{3}) - \operatorname{H} (A\_{1}, A\_{2}, A\_{3})).

Let  $V = A_1 \oplus \cdots \oplus A_n$  and take a vector subspace C of V such that

$$A_1 + \dots + A_{i-1} + C + A_{i+1} + \dots + A_n$$

is a direct sum for each *i*. We say that  $(A_1, \ldots, A_n, C)$  is a tuple of complementary vector spaces. Every vector of V has a unique representation as a sum of elements of  $A_1, \ldots, A_n$ . Therefore,  $\pi_I$  denotes the *I*-projection function  $V \twoheadrightarrow \bigoplus_{i \in I} A_i$  given by

$$x = \sum x_i \mapsto \sum_{i \in I} x_i.$$

**Proposition 2.** For any tuple  $(A_1, \ldots, A_n, C)$  of complementary vector spaces, we have

$$\mathrm{H}\left(\pi_{I}\left(C\right)\right) = \mathrm{H}\left(C\right) \leq \mathrm{H}\left(A_{i}\right),$$

for all i and  $\emptyset \neq I \subseteq [n]$ .

*Proof.* See proposition 6 in [12].

**Lemma 3.** Let  $A_1, \ldots, A_n$  and C be vector subspaces of a finite dimensional vector space V, such that

- $C \leq \sum A_i$ .
- $C \cap \sum_{i \neq k} A_i = O$  for all k.

Then, there exist subspaces  $\bar{A}_i \leq A_i$ , for i = 1, ..., n, such that

- (i)  $\bar{A}_k \cap \sum_{i \neq k} A_i = O$  for all k.
- (ii)  $(\bar{A}_1, \ldots, \bar{A}_n, C)$  is a tuple of complementary vector spaces.
- (iii)  $H(C) = H(\overline{A}_i)$  for all *i*.

*Proof.* In case C = O, we take  $\bar{A}_i = O$  for all *i*. Otherwise, we assume  $A_1, \ldots, A_n, C \neq O$ . Let  $(e_i)$  be a basis of C, we remark that each  $e_i$  can be written as  $\sum_j e_i^j$  with  $e_i^j \in A_i$  by hypothesis. Define  $\bar{A}_i = \langle e_i^j : j \rangle$ . For proving (i), we take

$$x = \sum_{i} \alpha_i e_k^i \in \bar{A}_k \cap \sum_{i \neq k} A_i.$$

We define  $\sum_{i} \alpha_i e_i \in C$  for getting

$$\sum_{i} \alpha_i e_i = \sum_{i} \alpha_i \sum_{j} e_i^j$$

$$= \sum_{i} \alpha_{i} e_{k}^{i} + \sum_{j} \sum_{i \neq k} \alpha_{i} e_{i}^{j}$$
$$= x + \sum_{j} \sum_{i \neq k} \alpha_{i} e_{i}^{j}.$$

Thus,  $\sum_{i} \alpha_{i} e_{i} \in C \cap \sum_{i \neq k} A_{i}$  which implies that  $\sum_{i} \alpha_{i} e_{i} = O$  by hypothesis. Since  $(e_{i})$  is a basis,  $\alpha_{i} = 0$ , i.e. x = O. Hence, (i) is true. In particular, this implies that  $\overline{A}_{k} \cap \sum_{i \neq k} \overline{A}_{i} = O$ , and by definition  $C \leq \sum_{i} \overline{A}_{i}$ . It follows that (ii) is true. We also note  $\overline{A}_{i}$  is generated by at most H(C)-vectors; therefore, by Proposition 2 and (ii), we have (iii) is true.  $\Box$ 

**Lemma 4.** For any vector subspaces  $A_1, \ldots, A_n$  of a finite dimensional vector space V, we have

$$\sum_{i} \operatorname{H}(A_{i}) - \operatorname{I}(A_{1}; \cdots; A_{n}) \leq \sum_{1 < i} \operatorname{H}(A_{1}, A_{i})$$

*Proof.* The proof is by induction. The case n = 2 gives a straightforward information identity. We suppose the case n - 1 holds, and we show the case n is true,

$$\begin{split} \sum_{i} \operatorname{H} \left( A_{i} \right) - \operatorname{I} \left( A_{1}; \cdots; A_{n} \right) &= \operatorname{H} \left( A_{n} \right) + \operatorname{H} \left( A_{1} \cap \cdots \cap A_{n-1} \right) \\ - \operatorname{I} \left( A_{1} \cap \cdots \cap A_{n-1}; A_{n} \right) + \sum_{i \leq n-1} \operatorname{H} \left( A_{i} \right) - \operatorname{I} \left( A_{1}; \cdots; A_{n-1} \right) \\ &\leq \operatorname{H} \left( A_{1} \cap \cdots \cap A_{n-1}, A_{n} \right) + \sum_{1 < i \leq n-1} \operatorname{H} \left( A_{1}, A_{i} \right) \text{ [from cases } n = 2 \text{ and } n - 1 \text{]} \\ &\leq \operatorname{H} \left( A_{1}, A_{n} \right) + \sum_{1 < i \leq n-1} \operatorname{H} \left( A_{1}, A_{i} \right) \\ &= \sum_{1 < i} \operatorname{H} \left( A_{1}, A_{i} \right). \end{split}$$

## **3** Secret Sharing

Secret Sharing is an important component in many kinds of cryptographic protocols [2, 4, 10]. In a *secret sharing scheme*, a *secret value* is distributed into *shares* among a set of *participants* in such a way that only the *qualified sets* of participants can recover the secret value.

**Definition 5.** An access structure, denoted by  $\Gamma$  on a set of participants P, is a monotone increasing family of subsets of P. Consider a special participant  $p \notin P$ , called dealer. A secret sharing scheme on P with access structure  $\Gamma$  is a tuple of random variables  $\Sigma := (S_x)_{x \in Q}$ , where  $Q = P \cup p$ , such that the following properties are satisfied:

(i)  $H(S_p) > 0.$ 

- (ii) If  $A \in \Gamma$ , then  $H(S_p \mid S_A) = 0$ .
- (iii) If  $A \notin \Gamma$ , then I  $(S_p; S_A) = 0$ .

The random variable  $S_p$  is the *secret value*, and the *shares* received by the participants are given by the random variables  $S_x$ ,  $x \in P$ . A set of participants A is said to be *qualified* if  $A \in \Gamma$ ; and it is said to be *non-qualified* if  $A \notin \Gamma$ . A *minimal qualified set* is a qualified set such that any proper subset is non-qualified. It is clear that an access structure is determined by the family min  $\mathcal{F}$  of its minimal qualified sets.

**Definition 6.** The *information ratio*  $\sigma(\Sigma)$  of the secret sharing scheme  $\Sigma$  is given by

$$\sigma\left(\Sigma\right) = \max_{x \in P} \frac{\mathrm{H}\left(S_{x}\right)}{\mathrm{H}\left(S_{p}\right)}.$$

The optimal information ratio  $\sigma(\Gamma)$  of an access structure  $\Gamma$  is the infimum of the information ratios of all secret sharing schemes for  $\Gamma$ . The optimal information ratio using only tuples of linear random variables is denoted by  $\lambda(\Gamma)$ . When we want to specify the characteristic of the field  $\mathbb{F}$  or some characteristic field condition, the optimal information ratio is denoted by  $\lambda_{char}(\mathbb{F})(\Gamma)$ .

We study the following classes of linear programming problems which are useful for calculating bounds on optimal information ratios [4].

**Problem 7.** For any access structure  $\Gamma$  on a set P with leader  $p \notin P$ , the optimal solution  $\kappa(\Gamma)$  of the linear programming problem is to calculate min (v) such that

- (i)  $v \ge f(x)$  for each  $x \in P$ .
- (ii)  $f(X \cup p) = f(X)$  for each  $X \subseteq P$  with  $X \in \Gamma$ .
- (iii)  $f(X \cup p) = f(X) + 1$  for each  $X \subseteq P$  with  $X \notin \Gamma$ .
- (iv) Information inequalities.

Given a secret sharing scheme  $\Sigma = (S_x)_{x \in Q}$ , with access structure  $\Gamma$ , we consider the mapping given by  $h(X) := H(S_X)$ , for every  $X \subseteq Q$ . We define  $f = \frac{1}{h(p)}h$ . The function f satisfies the conditions of problem 8. Therefore, f is a feasible solution and we have

$$\kappa\left(\Gamma\right) \leq \sigma\left(\Gamma\right).$$

When we add linear rank inequalities in (iv), we have a linear programming problem whose optimal solution, denoted by  $\kappa^*(\Gamma)$  holds

$$\kappa^{*}\left(\Gamma\right) \leq \lambda\left(\Gamma
ight)$$
 .

When we add characteristic-dependent linear rank inequalities the optimal solution is denoted by  $\kappa^*_{char}(\mathbb{F})$  ( $\Gamma$ ), we obtain

$$\kappa_{\operatorname{char}(\mathbb{F})}^{*}(\Gamma) \leq \lambda_{\operatorname{char}(\mathbb{F})}(\Gamma)$$




Figure 1: A matrix over GF(p) and Fano matroid.

**Definition 8.** A secret sharing scheme  $\Sigma = (S_x)_{x \in Q}$  is said to be *ideal* if its information ratio is equal to 1. An access structure that admits an ideal secret sharing scheme is called *ideal access structure*.

Matroids are related to secret sharing, for concepts associated, see [9].

**Definition 9.** Given a matroid  $\mathcal{M} = (Q, r)$ , where Q is the ground set and r is the rank function. The *port* of the matroid  $\mathcal{M}$  at  $p \in Q$  is the access structure on P = Q - p whose qualified sets are the sets  $X \subseteq P$  satisfying  $r(X \cup p) = r(X)$ .

The following result connects ideal secret sharing and matroids.

**Theorem 10.** Let  $\Sigma = (S_x)_{x \in Q}$  be an ideal secret sharing scheme on P with access structure  $\Gamma$ . Then, the mapping given by  $f(X) = \frac{H(S_X)}{H(S_p)}$  for each  $X \subseteq Q$  is the rank function of a matroid  $\mathcal{M}$  with ground set Q. Moreover,  $\Gamma$  is the port of the matroid  $\mathcal{M}$  at p.

As a consequence, every ideal access structure is a matroid port. A known result about  $\kappa$  and matroid ports is as follows.

**Theorem 11.** Let  $\Gamma$  be an access structure. Then,  $\Gamma$  is a matroid port if and only if  $\kappa(\Gamma) = 1$ . Moreover,  $\kappa(\Gamma) \geq \frac{3}{2}$  if  $\Gamma$  is not a matroid port. **Example 12.** The port of the Fano matroid at *c*, according figure 1, is given by the minimum qualified sets

$$\min \mathcal{F} := \{a_1b_1, a_2b_2, a_3b_3, a_1a_2a_3, a_1b_2b_3, b_1a_2b_3, b_1b_2a_3\}$$

The columns of a matrix of representation of Fano matroid define an ideal linear secret sharing scheme over fields whose characteristic is two, and therefore the ports of the Fano matroid are ideal. We have:

•  $\sigma(\mathcal{F}) = \lambda(\mathcal{F}) = \lambda_{\operatorname{char}(\mathbb{F})=2}(\mathcal{F}) = 1,$ 

• 
$$\kappa(\Gamma) = \kappa^*(\Gamma) = \kappa^*_{\operatorname{char}(\mathbb{F})=2}(\Gamma) = 1.$$

It is more hard for showing [7, 16]:

• 
$$\lambda_{\operatorname{char}(\mathbb{F})\neq 2}(\mathcal{F}) = \kappa^*_{\operatorname{char}(\mathbb{F})\neq 2}(\Gamma) = \frac{4}{3}$$

## **4 Producing inequalities**

Consider any  $n \times n$  binary matrix  $B = (b_{ii})$ , we write the *i*-th column as  $e_{S_i}$  where

$$S_i = \{j : b_{ji} = 1\}$$

We then define the sets:

$$\mathcal{B}' := \{ e_{S_i} : 1 < |S_i| < n \} ,$$
$$\mathcal{B}'' := \{ e_{S_i} : |S_i| = 1 \} ,$$
$$\mathcal{B}''' := \{ e_{S_i} : |S_i| = n \} .$$

In the following, we suppose that  $|\det(B)| = t > 1$ , for some  $t \in \mathbb{N}$  and  $\mathcal{B}'''$  is empty.

These matrices can be used to define matroid ports which are ideal over some fields; they are representable matroids.

We consider  $n + |\mathcal{B}'|$  participants labeled as follows

$$P := \{a_{e_i} : i \in [n]\} \cup \{b_{e_{S_j}} : e_{S_j} \in \mathcal{B}'\}.$$

We remark that n participants are labeled using the canonical basis  $(e_i)$  and  $|\mathcal{B}'|$  participants are labeled using the columns of B in  $\mathcal{B}'$ .

Now, consider any access structure on  $n + |\mathcal{B}'|$  participants such that:

• The following set is a subclass of the collection of minimal qualified sets,

$$\left\{ (a_{e_i})_{i \notin S_j} b_{e_{S_j}} : e_{S_j} \in \mathcal{B}' \right\} \cup \left\{ a_{e_1} \cdots a_{e_n} \right\}.$$

• The following set is a subclass of the class of non-qualified sets,

$$\left\{ (a_{e_i})_{i \in S_j} b_{e_{S_j}} : e_{S_j} \in \mathcal{B}' \right\}.$$

Let  $P_B$  be the subset of participants labeled by the columns of B. We produce two different classes of access structures according to add  $P_B$  to the subclass of minimal qualified sets or to the subclass of non-qualified sets. There are several access structures with these properties, in the next section we show an example.

The classes of access structures defined above can be used as a guide for determining properties or conditions that must be satisfied by associated vector spaces in order to derive some inequalities. Each vector space can be thought of as follows:

- $A_{e_i}$  is associated to  $a_{e_i}$ .
- $B_{e_i}$  is associated to  $b_{e_i}$ .
- C is associated to the dealer  $c \notin P$ .

**Proposition 13.** Let  $A_{e_i}$ , for  $i \in [n]$ ,  $B_{e_{S_j}}$ , for  $e_{S_j} \in \mathcal{B}'$ , and C be vector subspaces of a vector space V such that

- 
$$C \leq A_{[e_n]} \cap \left(\sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}}\right)$$
 for each  $j$ .

-  $C \cap A_{[e_n]-e_i} = O$  for each i.

- 
$$C \cap \left(\sum_{i \notin S_j, i \neq k} A_{e_i} + B_{e_{S_j}}\right) = O$$
 for each  $e_{S_j} \in \mathcal{B}$  and  $k \notin S_j$ .

Then, we have vector subspaces  $\bar{A}_{e_i} \leq A_{e_i}$ ,  $i \in [n]$ ;  $A_{e_i}^{S_j} \leq A_{e_i}$  and  $\hat{B}_{e_{S_j}} \leq B_{e_{S_j}}$  for each  $e_{S_j} \in \mathcal{B}'$  and  $i \notin S_j$  such that

- $(\bar{A}_{e_1}, \cdots, \bar{A}_{e_n}, C)$  is a tuple of complementary vector spaces.
- $(A_{e_i}^{S_j}, \hat{B}_{e_{S_j}}, C : i \notin S_j)$ , for each  $e_{S_j} \in \mathcal{B}'$ , is a tuple of complementary vector spaces.
- The dimension of any of these subspaces is H(C).
- These subspaces are unique except isomorphisms.

This proposition is a consequence of Lemma 3 and expresses the fact that in the access structure the participants  $a_{e_1} \cdots a_{e_n}$ ,  $(a_{e_i})_{i \notin S_i} b_{e_{S_i}}$ , for all  $e_{S_j} \in \mathcal{B}'$ , are minimal qualified sets.

The following linear mapping is well-defined under hypothesis of previous proposition:

$$\varphi_B : C \to \bigoplus_i \frac{\bar{A}_{e_i}}{(\bigcap \mathcal{A}_{e_i}) \cap \bar{A}_{e_i}}, \\ c \longmapsto \varphi_B (c) := \sum_i [a_i]_{(\bigcap \mathcal{A}_{e_i}) \cap \bar{A}_{e_i}},$$

where  $c = \sum_{i} a_i$  with  $a_i \in \overline{A}_{e_i}$  and  $\mathcal{A}_{e_i} := \left\{ A_{e_i}^{S_j} : i \notin S_j \text{ for some } j \right\}$ ; we take  $\mathcal{A}_{e_i}$  as  $\{O\}$ , in case that  $i \in S_j$  for all j. We remark that there is a correspondence between  $\mathcal{A}_{e_i}$  and the subset of columns of B given by  $\mathcal{B}_{e_i} := \{e_{S_j} : i \notin S_j\}$ ; we take  $\mathcal{B}_{e_i}$  as  $\{O\}$  in case that  $i \in S_j$  for all j.

**Lemma 14.** For any vector subspaces  $A_{e_1}, \ldots, A_{e_n}$ ,  $B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|\mathcal{B}'|}}}}$  and C of a finite dimensional vector space V such that

(i) 
$$C \leq A_{[e_n]} \cap \left(\sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}}\right)$$
 for each  $e_{S_j} \in \mathcal{B}'$ .

(ii)  $C \cap A_{[e_n]-e_i} = O$  for each *i*.

(iii) 
$$C \cap \left(\sum_{i \notin S_j, i \neq k} A_{e_i} + B_{e_{S_j}}\right) = O$$
 for each  $e_{S_j} \in \mathcal{B}$  and  $k \notin S_j$ .

Then,

$$\left[1 + \sum_{i} |\mathcal{B}_{e_{i}}|\right] \operatorname{H}(C) \leq \sum_{i} |\mathcal{B}_{e_{i}}| \operatorname{H}(A_{e_{i}}) + \operatorname{H}(\ker(\varphi_{B})).$$

*Proof.* From mapping  $\varphi_B$ , we can derive the inequality

$$\mathrm{H}\left(\frac{C}{\mathrm{ker}\left(\varphi_{B}\right)}\right) \leq \sum_{i} \mathrm{H}\left(\frac{\bar{A}_{e_{i}}}{\left(\bigcap \mathcal{A}_{e_{i}}\right) \cap \bar{A}_{e_{i}}}\right).$$

So

$$\mathrm{H}(C) - \mathrm{H}(\mathrm{ker}(\varphi_B)) \leq \sum_{i} \left[ \mathrm{H}(\bar{A}_{e_i}) - \mathrm{I}(\bar{A}_{e_i}; \bigcap \mathcal{A}_{e_i}) \right]$$

Then

$$\begin{split} \mathrm{H}\left(C\right) - \mathrm{H}\left(\ker\varphi_{B}\right) + \sum_{i} \sum_{e_{S_{j}}\in\mathcal{B}_{e_{i}}} \mathrm{H}\left(A_{e_{i}}^{S_{j}}\right) &\leq \sum_{i} \left[\mathrm{H}\left(\bar{A}_{e_{i}}\right) + \sum_{e_{S_{j}}\in\mathcal{B}_{e_{i}}} \mathrm{H}\left(A_{e_{i}}^{S_{j}}\right) - \mathrm{I}\left(\bar{A}_{e_{i}};\bigcap\mathcal{A}_{e_{i}}\right)\right], \\ &\leq \sum_{i} \sum_{e_{S_{j}}\in\mathcal{B}_{e_{i}}} \mathrm{H}\left(\bar{A}_{e_{i}},A_{e_{i}}^{S_{j}}\right), \text{ [from Lemma 4]}. \end{split}$$

Since  $\operatorname{H}\left(A_{e_{i}}^{S_{j}}\right) = \operatorname{H}(C), \sum_{e_{S_{j}}\in\mathcal{B}_{e_{i}}}1 = |\mathcal{B}_{e_{i}}| \text{ and } \bar{A}_{e_{i}}, A_{e_{i}}^{S_{j}} \leq A_{e_{i}}, \text{ we get}$  $\operatorname{H}(C) - \operatorname{H}\left(\ker\left(\varphi_{B}\right)\right) + \sum_{i}|\mathcal{B}_{e_{i}}|\operatorname{H}(C) \leq \sum_{i}|\mathcal{B}_{e_{i}}|\operatorname{H}(A_{e_{i}}),$ 

which implies the desired inequality.

**Lemma 15.** For any vector subspaces  $A_{e_1}, \ldots, A_{e_n}$ ,  $B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|\mathcal{B}'|}}}}$  and C of a finite dimensional vector space V over a finite field  $\mathbb{F}$  whose characteristic does not divide t, such that

(i) 
$$C \leq A_{[e_n]} \cap \left(\sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}}\right)$$
, for each  $e_{S_j} \in \mathcal{B}'$ .

(ii) 
$$C \cap A_{[e_n]-e_i} = C \cap \left(\sum_{i \in S_j} A_{e_i} + B_{e_{S_j}}\right) = C \cap \left(\sum_{i \notin S_j, i \neq k} A_{e_i} + B_{e_{S_j}}\right) = O$$
, for all  $e_{S_j} \in \mathcal{B}'$  and  $k \notin S_j$ .

(iii)  $C \cap \mathcal{B} = O$ , where  $\mathcal{B}$  is the sum of all vector subspaces indexed by the columns of B. Then

$$\ker\left(\varphi_B\right)=O.$$

*Proof.* We take  $c = \sum_{i} a_i \in C$  such that  $\varphi_B(c) = O$ , where  $a_i \in \overline{A}_i$ . We have to show c = O. By definition of  $\varphi_B$ ,

$$a_i \in \bar{A}_{e_i} \cap \left(\bigcap_{e_{S_j} \in \mathcal{B}_{e_i}} A_{e_i}^{S_j}\right)$$

Hence,  $a_i \in A_{e_i}^{S_j}$  for all  $e_{S_j} \in \mathcal{B}_{e_i}$ , and therefore

$$\sum_{i \notin S_j} a_i \in \sum_{i \notin S_j} A_{e_i}^{S_j}.$$

From (i) in Lemma 3, there exists  $b_j \in \hat{B}_{e_{S_j}}$  for each  $e_{S_j} \in \mathcal{B}'$  such that  $\sum_{i \notin S_j} a_i + b_j \in C$ . Hence,

$$\sum_{i \in S_j} a_i - b_j = \sum_i a_i - \left(\sum_{i \notin S_j} a_i + b_j\right) \in C \cap \left(\sum_{i \in S_j} A_{e_i} + B_{e_{S_j}}\right).$$

From (ii), this implies

$$\sum_{i \in S_j} a_i = b_j \text{ for all } e_{S_j} \in \mathcal{B}'.$$

These equalities define the following linear system of equations

$$B^{T}\begin{pmatrix}a_{1}\\\vdots\\a_{n}\end{pmatrix} = \begin{pmatrix}b_{1}\\\vdots\\b_{|\mathcal{B}'|}\\b^{1}\\\vdots\\b^{|\mathcal{B}''|}\end{pmatrix},$$
(1)

where the vectors  $b^1, \ldots, b^{|\mathcal{B}''|}$  are omitted when  $\mathcal{B}''$  is empty; in other case,  $b^i := a_i$ , for  $e_i \in \mathcal{B}''$ . Since char (F) does not divide  $t = |\det(B)|$ , the matrix  $B^T$  is non-singular. Therefore, each  $a_i$  can be written as a linear combination of  $b_1, \ldots, b_{|\mathcal{B}'|}, b^1, \ldots, b^{|\mathcal{B}''|}$ , which implies that  $c \in \mathcal{B}$ . From (iii), we get c = O.

**Corollary 16.** For any vector subspaces  $A_{e_1}, \ldots, A_{e_n}, B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|\mathcal{B}'|}}}}$  and C of a finite dimensional vector space V over a finite field  $\mathbb{F}$ , such that

(i) 
$$C \leq A_{[e_n]} \cap \left(\sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}}\right)$$
, for all  $e_{S_j} \in \mathcal{B}'$ .  
(ii)  $C \cap A_{[e_n]-e_i} = C \cap \left(\sum_{i \in S_j} A_{e_i} + B_{e_{S_j}}\right) = C \cap \left(\sum_{i \notin S_j, i \neq k} A_{e_i} + B_{e_{S_j}}\right) = O$ , for all  $e_{S_j} \in \mathcal{B}'$  and  $k \notin S_j$ .

Then, the mapping

$$\phi_B^k : \ker (\varphi_B) \to B_{e_{S_k}}$$
$$c \longmapsto \phi_B^k (c) := \sum_{i \in S_k} a_i = b_k$$

is an one-to-one well-defined linear function for each  $e_{S_k} \in \mathcal{B}'$ . Also, if the k-th column of B is a linear combination of the columns of the submatrix of B denoted by  $B_X$ ,  $k \notin X$ . Then,

$$\phi_B^k\left(\ker\left(\varphi_B\right)\right) \subseteq \sum_{e_i \in B_X \cap \mathcal{B}''} A_{e_i} + \sum_{e_{S_i} \in B_X \cap \mathcal{B}'} B_{e_{S_i}}.$$

*Proof.* We can follow line-by-line the proof of the previous lemma to obtain that there exists a unique  $b_k \in \left(\sum_{i \in S_k} \bar{A}_i\right) \cap \hat{B}_{e_{S_k}} \subseteq B_{e_{S_k}}$ . So  $\phi_B^k$  is well-defined. Since the written of each  $c \in \ker(\varphi_B)$  is unique,  $\phi_B^k$  is also an one-to-one linear mapping. Also, if the k-th column of B is a linear combination of the columns of  $B_X$ , from equation (1), we have that  $b_k$  is a linear combination of  $(b_i)_{i \in B_X} \cup (b^i)_{i \in B_X}$ . Therefore,  $b_k \in \sum_{e_i \in B_X \cap \mathcal{B}'} A_{e_i} + \sum_{e_{S_i} \in B_X \cap \mathcal{B}'} B_{e_{S_i}}$ .

We finally show a theorem that can produce characteristic-dependent linear rank inequalities as long as there are suitable binary matrices.

**Theorem 17.** For a  $n \times n$  binary matrix B such that  $\mathcal{B}''' = \emptyset$  and  $|\det(B)| = t \in \mathbb{N}$ , t > 1. Let  $A_{e_1}, \ldots, A_{e_n}, B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|\mathcal{B}'|}}}}$  and C be vector subspaces of a finite dimensional vector space V over  $\mathbb{F}$ . We have

- The following inequality is a characteristic-dependent linear rank inequality over fields whose characteristic does not divide t,

$$\begin{split} \mathrm{H}\left(C\right) &\leq \frac{1}{1 + \sum_{i} |\mathcal{B}_{e_{i}}|} \sum_{i} |\mathcal{B}_{e_{i}}| \,\mathrm{H}\left(A_{e_{i}}\right) + \mathrm{H}\left(C \mid A_{[e_{n}]}\right) + \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}} : e_{i} \in \mathcal{B}'', e_{S_{j}} \in \mathcal{B}'\right) \\ &+ \sum_{i} \mathrm{I}\left(C; A_{[e_{n}] - e_{i}}\right) + \sum_{e_{S_{h}} \in \mathcal{B}', i \notin S_{h}} \mathrm{I}\left(C; A_{e_{j}}, B_{e_{S_{h}}} : j \notin S_{h}, j \neq i\right) \\ &+ \sum_{e_{S_{j}} \in \mathcal{B}'} \left[\mathrm{H}\left(C \mid A_{e_{i}}, B_{e_{S_{j}}}, i \notin S_{j}\right) + \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}} : i \in S_{j}\right)\right]. \end{split}$$

- Fixed  $k \in [n]$  such that  $e_{S_k} \in \mathcal{B}'$ . The following inequality is a characteristic-dependent linear rank inequality over fields whose characteristic divides t,

$$\begin{split} \mathrm{H}\left(C\right) &\leq \frac{1}{2 + \sum_{i} |\mathcal{B}_{e_{i}}|} \left[ \sum_{i} |\mathcal{B}_{e_{i}}| \,\mathrm{H}\left(A_{e_{i}}\right) + \mathrm{H}\left(B_{e_{S_{k}}}\right) \right] + \mathrm{H}\left(C \mid A_{e_{i}}, B_{e_{S_{j}}} : e_{i} \in \mathcal{B}'', e_{S_{j}} \in \mathcal{B}'\right) \\ &+ \mathrm{H}\left(C \mid A_{[e_{n}]}\right) + \sum_{e_{S_{i}} \in \mathcal{B}'} \mathrm{H}\left(C \mid A_{e_{j}}, B_{e_{S_{i}}} : j \notin S_{i}\right) \\ &+ \sum_{i} \mathrm{I}\left(C; A_{[e_{n}] - e_{i}}\right) + \sum_{e_{S_{h}} \in \mathcal{B}', i \notin S_{h}} \mathrm{I}\left(C; A_{e_{j}}, B_{e_{S_{h}}} : j \notin S_{h}, j \neq i\right) \\ &+ \sum_{e_{S_{j}} \in \mathcal{B}'} \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}} : i \in S_{j}\right) + \sum_{i} \mathrm{I}\left(C; A_{e_{j}}, B_{e_{S_{i}}} : e_{S_{j}} \in \mathcal{B}', e_{j} \in \mathcal{B}'', j \neq i\right). \end{split}$$

The inequalities do not in general hold over fields whose characteristic is different to the mentioned. Counter examples would be in  $V = \operatorname{GF}(p)^n$ , take the vector spaces  $A_{e_i} = \langle e_i \rangle$ ,  $e_i \in [e_n]$ ,  $B_{e_{S_j}} = \langle e_{S_j} \rangle$ ,  $e_{S_j} \in \mathcal{B}'$ , and  $C = \langle \sum e_i \rangle$  Then, when p divides t, the first inequality does not hold; and when p does not divide t, the second inequality does not hold. *Proof.* To prove the first inequality: let  $\mathbb{F}$  be a finite field whose characteristic does not divide t. Let

$$C^{\langle 0 \rangle} := C \cap A_{[e_n]} \cap \left[ \bigcap_{e_{S_j} \in \mathcal{B}'} \left( \sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}} \right) \right].$$

We have

$$\mathrm{H}\left(C \mid C^{\langle 0 \rangle}\right) \leq \mathrm{H}\left(C \mid A_{[e_{n}]}\right) + \sum_{e_{S_{j}} \in \mathcal{B}'} \mathrm{H}\left(C \mid A_{e_{i}}, B_{e_{S_{j}}} : i \notin S_{j}\right).$$

Recursively, for  $i \in [n]$ , denote by  $C^{\langle i \rangle}$ , a subspace of  $C^{\langle i-1 \rangle}$  which is a complementary space to  $\sum_{j \neq i} A_{e_j}$  in

$$C^{\langle i-1\rangle} + \sum_{j \neq i} A_{e_j}.$$

We have

$$\mathrm{H}\left(C^{\langle i-1\rangle} \mid C^{\langle i\rangle}\right) \leq \mathrm{I}\left(C; A_{e_j} : j \neq i\right).$$

Let  $C_{e_{S_{j_1}}}^{[0]} := C^{\langle n \rangle}$  and recursively, for each  $i \notin S_{j_1}$ , we denote by  $C_{e_{S_{j_1}}}^{[i]}$ , a subspace of  $C_{e_{S_{j_1}}}^{[i-1]}$  which is a complementary space to  $\sum_{j \notin S_{j_1}, j \neq i} A_{e_j} + B_{e_{S_{j_1}}}$  in

$$C_{e_{S_{j_1}}}^{[i-1]} + \sum_{j \notin S_{j_1}, j \neq i} A_{e_j} + B_{e_{S_{j_1}}}.$$

We have

$$\mathbf{H}\left(C_{e_{S_{j_{1}}}}^{[i-1]} \mid C_{e_{S_{j_{1}}}}^{[i]}\right) \leq \mathbf{I}\left(C; A_{e_{j}}, B_{e_{S_{j_{1}}}} : j \notin S_{j_{1}}, j \neq i\right).$$

In a similar way, we define  $C_{e_{S_{j_2}}}^{[0]} = C_{e_{S_{j_1}}}^{[n]}, \ldots, C_{e_{S_{j_{|\mathcal{B}'}|}}}^{[0]} = C_{e_{S_{j_{|\mathcal{B}'}|-1}}}^{[n]}$  until to find a subspace  $C^{(0)} := C_{e_{S_{j_{|\mathcal{B}'}|}}}^{[n]}$  that holds

$$\operatorname{H}\left(C^{\langle n \rangle} \mid C^{(0)}\right) \leq \sum_{e_{S_{h}} \in \mathcal{B}', i \notin S_{h}.} \operatorname{I}\left(C; A_{e_{j}}, B_{e_{S_{h}}}: j \notin S_{h}, j \neq i\right).$$

Recursively, for i, with  $e_{S_i} \in \mathcal{B}'$ , we denote by  $C^{(i)}$ , a subspace of  $C^{(i-1)}$  which is a complementary space to  $\sum_{j \in S_i} A_{e_j} + B_{e_{S_i}}$  in

$$C^{(i-1)} + \left(\sum_{j \in S_i} A_{e_j} + B_{e_{S_i}}\right).$$

We also have

$$H(C^{(i-1)} | C^{(i)}) \le I(C^{(i-1)}; A_{e_j}, B_{e_{S_i}} : j \in S_i)$$

Define by  $\hat{C},$  a subspace of  $C^{(|\mathcal{B}'|)}$  which is a complementary space to

$$\mathcal{B} = \left(\sum_{e_{S_i} \in \mathcal{B}'} B_{e_{S_i}}\right) + \left(\sum_{e_i \in \mathcal{B}''} A_{e_i}\right)$$

in  $C^{(|\mathcal{B}'|)} + \mathcal{B}$ . We have

$$\operatorname{H}\left(C^{(|\mathcal{B}'|)} \mid \hat{C}\right) \leq \operatorname{I}\left(C; A_{e_i}, B_{e_{S_j}} : e_i \in \mathcal{B}'', e_{S_j} \in \mathcal{B}'\right).$$

Hence,

$$\begin{split} \operatorname{H}\left(C \mid \hat{C}\right) &= \operatorname{H}\left(C \mid C^{\langle 0 \rangle}\right) + \operatorname{H}\left(C^{\langle 0 \rangle} \mid C^{\langle n \rangle}\right) + \operatorname{H}\left(C^{\langle n \rangle} \mid C^{\langle 0 \rangle}\right) \\ &+ \operatorname{H}\left(C^{\langle 0 \rangle} \mid C^{(|\mathcal{B}'|)}\right) + \operatorname{H}\left(C^{\langle |\mathcal{B}'|\rangle} \mid \hat{C}\right) \\ &\leq \operatorname{H}\left(C \mid A_{[e_n]}\right) + \sum_{e_{S_j} \in \mathcal{B}'} \operatorname{H}\left(C \mid A_{e_i}, B_{e_{S_j}} : i \notin S_j\right) \\ &+ \sum_{i} \operatorname{I}\left(C; A_{[e_n] - e_i}\right) + \sum_{e_{S_h} \in \mathcal{B}', i \notin S_h} \operatorname{I}\left(C; A_{e_j}, B_{e_{S_h}} : j \notin S_h, j \neq i\right) \\ &+ \sum_{e_{S_j} \in \mathcal{B}'} \operatorname{I}\left(C; A_{e_i}, B_{e_{S_j}} : i \in S_j\right) + \operatorname{I}\left(C; A_{e_i}, B_{e_{S_j}} : e_i \in \mathcal{B}'', e_{S_j} \in \mathcal{B}'\right). \end{split}$$

Since  $A_{e_1}, \ldots, A_{e_n}, B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|B'|}}}}$  and  $\hat{C}$  satisfy hypothesis in Lemma 15, we have ker  $(\varphi_B) = O$ . Therefore, as these spaces also satisfy hypothesis in Lemma 14, it follows

$$\left[1 + \sum_{i} |\mathcal{B}_{e_{i}}|\right] \operatorname{H}\left(\hat{C}\right) \leq \sum_{i} |\mathcal{B}_{e_{i}}| \operatorname{H}\left(A_{e_{i}}\right).$$

Using the last two inequalities, we can obtain the described inequality:

$$\begin{split} \mathrm{H}\left(C\right) &-\mathrm{H}\left(C\mid A_{[e_{n}]}\right) - \sum_{e_{S_{j}}\in\mathcal{B}'}\mathrm{H}\left(C\mid A_{e_{i}}, B_{e_{S_{j}}}: i\notin S_{j}\right) \\ &-\sum_{i\in[n]}\mathrm{I}\left(C; A_{[e_{n}]-e_{i}}\right) - \sum_{e_{S_{h}}\in\mathcal{B}', i\notin S_{h}.}\mathrm{I}\left(C; A_{e_{j}}, B_{e_{S_{h}}}: j\notin S_{h}, j\neq i\right) \\ &-\sum_{e_{S_{j}}\in\mathcal{B}'}\mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}}: i\in S_{j}\right) - \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}}: e_{i}\in\mathcal{B}'', e_{S_{j}}\in\mathcal{B}'\right) \leq \mathrm{H}\left(\hat{C}\right) \\ &\leq \frac{1}{1+\sum_{i}|\mathcal{B}_{e_{i}}|}\sum_{i}|\mathcal{B}_{e_{i}}|\,\mathrm{H}\left(A_{e_{i}}\right). \end{split}$$

To prove the second inequality, let  $k \in [n]$  such that  $e_{S_k} \in \mathcal{B}'$  and let  $\mathbb{F}$  be a finite field whose characteristic divides t. Let

$$C^{[0]} := C \cap \mathcal{B} \cap A_{[e_n]} \cap \left[ \bigcap_{e_{S_j} \in \mathcal{B}'} \left( \sum_{i \notin S_j} A_{e_i} + B_{e_{S_j}} \right) \right].$$

We apply to  $C^{[0]}$  the same argument applied to space  $C^{\langle 0 \rangle}$  in the proof of the previous inequality, we therefore obtain a subspace  $C^{\{0\}} := C^{(|\mathcal{B}'|)}$ . Recursively, for  $i \in [n]$ , we denote by  $C^{\{i\}}$ , a subspace of  $C^{\{i-1\}}$  which is a complementary space to

$$\left(\sum_{e_{S_j}\in\mathcal{B}', j\neq i} B_{e_{S_j}}\right) + \left(\sum_{e_j\in\mathcal{B}'', j\neq i} A_{e_j}\right)$$

in

$$C^{\{i-1\}} + \left(\sum_{e_{S_j} \in \mathcal{B}', j \neq i} B_{e_{S_j}}\right) + \left(\sum_{e_j \in \mathcal{B}'', j \neq i} A_{e_j}\right);$$

we have

$$\operatorname{H}\left(C^{\{i-1\}} \mid C^{\{i\}}\right) \leq \operatorname{I}\left(C; A_{e_j}, B_{e_{S_i}} : e_{S_j} \in \mathcal{B}', e_j \in \mathcal{B}'', j \neq i\right)$$

We define  $\tilde{C} := C^{\{n\}}$  and the following inequality is true

$$H\left(C \mid \tilde{C}\right) = H\left(C \mid C^{\{0\}}\right) + H\left(C^{\{0\}} \mid \tilde{C}\right)$$

$$\leq H\left(C \mid A_{e_i}, B_{e_{S_j}} : e_i \in \mathcal{B}'', e_{S_j} \in \mathcal{B}'\right) + H\left(C \mid A_{[e_n]}\right) + \sum_{e_{S_j} \in \mathcal{B}'} H\left(C \mid A_{e_i}, B_{e_{S_j}} : i \notin S_j\right)$$

$$+ \sum_i I\left(C; A_{[e_n]-e_i}\right) + \sum_{e_{S_h} \in \mathcal{B}', i \notin S_h.} I\left(C; A_{e_j}, B_{e_{S_h}} : j \notin S_h, j \neq i\right)$$

$$+ \sum_{e_{S_j} \in \mathcal{B}'} I\left(C; A_{e_i}, B_{e_{S_j}} : i \in S_j\right) + \sum_i I\left(C; A_{e_j}, B_{e_{S_i}} : e_{S_j} \in \mathcal{B}', e_j \in \mathcal{B}'', j \neq i\right).$$

$$(2)$$

We remark that vector subspaces  $A_{e_1}, \ldots, A_{e_n}, B_{e_{S_{j_1}}}, \ldots, B_{e_{S_{j_{|B'|}}}}$  and  $\tilde{C}$  satisfy hypothesis in Lemma 14. Thus,

$$\left[1 + \sum_{i} |\mathcal{B}_{e_{i}}|\right] \operatorname{H}\left(\tilde{C}\right) \leq \sum_{i} |\mathcal{B}_{e_{i}}| \operatorname{H}\left(A_{e_{i}}\right) + \operatorname{H}\left(\ker\left(\varphi_{B}\right)\right).$$
(3)

As B is singular over fields whose characteristic divides t, without loss generality, we suppose that there exists a submatrix  $B_X$  such that  $e_{S_k}$  is a linear combination of the columns of  $B_X$ . So, from Corollary 16,

$$H\left(\ker\left(\varphi_{B}\right)\right) \leq I\left(B_{e_{S_{k}}}; A_{e_{i}}, B_{e_{S_{j}}}: e_{i} \in B_{X} \cap \mathcal{B}'', e_{S_{j}} \in B_{X} \cap \mathcal{B}'\right).$$

$$\tag{4}$$

We note that  $\tilde{C} \leq \mathcal{B}$  and  $\tilde{C} \cap \mathcal{B}_Y = O$ , for all  $B_Y$ , where  $\mathcal{B}_Y$  is the sum of all vector subspaces indexed by the columns of a proper submatrix  $B_Y$  of B. From Lemma 3, taking  $C := \tilde{C}$ ,  $A_i := A_{e_i}$ ,  $A_j := B_{e_{S_j}}$  according to  $e_i \in \mathcal{B}''$  or  $e_{S_j} \in \mathcal{B}'$ , the inequality

$$\mathrm{H}\left(\tilde{C}\right) + \mathrm{I}\left(B_{e_{S_{k}}}; A_{e_{i}}, B_{e_{S_{j}}}: e_{i} \in \mathcal{B}'', e_{S_{j}} \in \mathcal{B}', S_{j} \neq S_{k}\right) \leq \mathrm{H}\left(B_{e_{S_{k}}}\right)$$

$$[\text{we note that } \bar{B}_{e_{S_{k}}} \cap \mathcal{B}_{B-k} = O \text{ and } \mathrm{H}\left(\tilde{C}\right) = \mathrm{H}\left(\bar{B}_{e_{S_{k}}}\right)]$$

which implies

$$\operatorname{H}\left(\tilde{C}\right) + \operatorname{I}\left(B_{e_{S_{k}}}; A_{e_{i}}, B_{e_{S_{i}}}: e_{i} \in B_{X} \cap \mathcal{B}'', e_{S_{i}} \in B_{X} \cap \mathcal{B}'\right) \leq \operatorname{H}\left(B_{e_{S_{k}}}\right).$$

Using inequality (4), we have

$$\operatorname{H}\left(\tilde{C}\right) + \operatorname{H}\left(\ker\left(\varphi_{B}\right)\right) \leq \operatorname{H}\left(B_{e_{S_{k}}}\right)$$

Therefore, from inequality (3),

$$\left[2 + \sum_{i} |\mathcal{B}_{e_{i}}|\right] \operatorname{H}\left(\tilde{C}\right) \leq \sum_{i} |\mathcal{B}_{e_{i}}| \operatorname{H}\left(A_{e_{i}}\right) + \operatorname{H}\left(B_{e_{S_{k}}}\right).$$

From this and inequality (2), we obtain the desired inequality:

$$\begin{split} \mathrm{H}\left(C\right) - \mathrm{H}\left(C \mid A_{[e_{n}]}\right) &- \sum_{e_{S_{j}} \in \mathcal{B}'} \mathrm{H}\left(C \mid A_{e_{i}}, B_{e_{S_{j}}} : i \notin S_{j}\right) - \mathrm{H}\left(C \mid A_{e_{i}}, B_{e_{S_{j}}} : e_{i} \in \mathcal{B}'', e_{S_{j}} \in \mathcal{B}'\right) \\ &- \sum_{i} \mathrm{I}\left(C; A_{[e_{n}]-e_{i}}\right) - \sum_{e_{S_{h}} \in \mathcal{B}', i \notin S_{h}} \mathrm{I}\left(C; A_{e_{j}}, B_{e_{S_{h}}} : j \notin S_{h}, j \neq i\right) \\ &- \sum_{e_{S_{j}} \in \mathcal{B}'} \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}} : i \in S_{j}\right) - \sum_{i} \mathrm{I}\left(C; A_{e_{i}}, B_{e_{S_{j}}} : e_{S_{j}} \in \mathcal{B}', e_{i} \in \mathcal{B}'', j \neq i\right) \\ &\leq \mathrm{H}\left(\tilde{C}\right) \leq \frac{1}{2 + \sum_{i} |\mathcal{B}_{e_{i}}|} \left(\sum_{i} |\mathcal{B}_{e_{i}}| \operatorname{H}\left(A_{e_{i}}\right) + \mathrm{H}\left(B_{e_{S_{k}}}\right)\right). \end{split}$$

$B_1 \cdots B_{t+1} A_{t+2} \cdots A_k$	M(n,t)
$(0 \cdots 1 0 \cdots$	0
1 : 1 0 :	$0 \qquad \qquad a_1 \ \cdots \ a_{t+1} \ b_1 \ \cdots \ b_{t+1} \ c$
: : : : :	$0 \qquad \qquad \begin{pmatrix} 1 & \cdots & 0 & 0 & \cdots & 1 & 1 \end{pmatrix}$
$1  \vdots  1  0  \vdots$	$0 \qquad \qquad 0  \vdots  \vdots  1  1$
$1  \vdots  0  0  \vdots$	
1 : 1 1 :	$0 \qquad 0  \vdots  0  1  \vdots  1  1$
$1  \vdots  \vdots  0  \vdots$	$0 \qquad \qquad$
1 : 1 : :	0
$\begin{pmatrix} 1 & \cdots & 1 & 0 & \cdots \end{pmatrix}$	1 /

Figure 2: A family of matrices  $B_{M(n,t)}^t$  whose determinant is  $\pm t$  and a family of representable matroids.

#### 5 **Examples and applications**

We now produce some characteristic-dependent linear rank inequalities using a convenient class of matrices. Let  $n \ge 7$  and t integer such that  $2 \le t \le \lfloor \frac{n-1}{2} \rfloor - 1$  and M(n,t) = n - t - 2. In Theorem 17, we take square matrices  $B_{M(n,t)}^t$  as described in figure 2 on the left side with column vectors of the form  $A_i := A_{e_i} = e_i$ ,  $B_i := B_{e_{[M(n,t)]-i}} = c - e_i$  and  $c = \sum_{i \in [M(n,t)]} e_i$ . We have  $\left| \det \left( B_{M(n,t)}^t \right) \right| = t$ ,  $\left| \mathcal{B}_{B_{M(n,t)}^t}^t \right| = t + 1$ ,  $\left| \mathcal{B}_{B_{M(n,t)}^t}^{"} \right| = M(n,t) - t - 1$ ,  $\left| \mathcal{B}_{e_i} \right| = 1$  for  $i \in [t+1]$ and  $\left| \mathcal{B}_{e_i} \right| = 0$  for  $i \in [t+2]M(n,t)$ . Let  $A_i = A_{e_i} = A_{e_i} = P_{e_i}$  and C he matrices

and  $|\mathcal{B}_{e_i}| = 0$  for  $i \in [t+2, M(n,t)]$ . Let  $A_1, A_2, \ldots, A_{M(n,t)}, B_1, B_2, \ldots, B_{t+1}$  and C be vector subspaces of a finite dimensional vector space V over a finite field  $\mathbb{F}$ . We have

- A characteristic-dependent linear rank inequality over fields whose characteristic does not divide t,

$$H(C) \leq \frac{1}{t+2} \sum_{i \in [t+1]} H(A_i) + I(C; B_{[t+1]}, A_{[t+2,M(n,t)]}) + H(C \mid A_{[M(n,t)]})$$
  
+ 
$$\sum_{i \in [M(n,t)]} I(C; A_{[M(n,t)]-i}) + \sum_{i \in [t+1]} [I(C; B_i) + H(C \mid A_i, B_i) + I(C; A_{[M(n,t)]-i}, B_i)].$$

- A characteristic-dependent linear rank inequality over fields whose characteristic divides t,

$$H(C) \leq \frac{1}{t+3} \left[ \sum_{i \in [t+1]} H(A_i) + H(B_1) \right] + H(C \mid B_{[t+1]}, A_{[t+2,M(n,t)]}) + H(C \mid A_{[M(n,t)]})$$

$$+ \sum_{i \in [M(n,t)]} I(C; A_{[M(n,t)]-i}) + \sum_{i \in [t+1]} \left[ I(C; B_i) + H(C \mid A_i, B_i) + I(C; A_{[M(n,t)]-i}, B_i) \right]$$

$$+\sum_{i\in[t+1]} \mathrm{I}\left(C; B_{[t+1]-i}, A_{[t+2,M(n,t)]}\right) + \sum_{i\in[t+2,M(n,t)]} \mathrm{I}\left(C; B_{[t+1]}, A_{[t+2,M(n,t)]-i}\right).$$

*Remark* 18. We produce a class of  $\lfloor \frac{n-1}{2} \rfloor - 2$  inequalities that are true over finite sets of primes and another class of  $\lfloor \frac{n-1}{2} \rfloor - 2$  inequalities that are true over co-finite sets of primes.

Let  $t \in \mathbb{N}$ , t > 1 and let  $\mathbb{F}$  be a finite field. We use the port at c of the representable matroid obtained from the matrix in Figure 2 on the right side; the set of participants is  $P = \{a_1, \ldots, a_{t+1}, b_1, \ldots, b_{t+1}\}$ , with dealer p = c.

When char ( $\mathbb{F}$ ) divides *t*, the following set is a subclass of the minimal qualified set:

$$\{a_1b_1,\ldots,a_{t+1}b_{t+1},a_1\cdots a_{t+1},a_1b_2\cdots b_{t+1},b_1a_2b_3\cdots b_{t+1},\ldots,b_1\cdots b_ta_{t+1}\},\$$

and the following set is a subclass of the non-qualified set:

$$\{b_1a_2\cdots a_{t+1}, a_1b_2a_3\cdots a_{t+1}, \dots, a_1\cdots a_tb_{t+1}\} \cup \{b_1\cdots b_{t+1}\}$$

When char ( $\mathbb{F}$ ) does not divide t, the following set is a subclass of the minimal qualified set:

$$\{a_1b_1, \dots, a_{t+1}b_{t+1}, a_1 \cdots a_{t+1}, a_1b_2 \cdots b_{t+1}, b_1a_2b_3 \cdots b_{t+1}, \dots, b_1 \cdots b_ta_{t+1}\} \cup \{b_1 \cdots b_{t+1}\}$$

and the following set is a subclass of the non-qualified set:

$$\{b_1a_2\cdots a_{t+1}, a_1b_2a_3\cdots a_{t+1}, \ldots, a_1\cdots a_tb_{t+1}\}.$$

So, we have defined two types of access structures using matroid ports. Let  $\mathcal{F}_t$  be an access structure of the first type and let  $\mathcal{N}_t$  be an access structure of the second type. We remark that  $\mathcal{F}_2$  is a port of Fano matroid and  $\mathcal{N}_2$  is a port of non-Fano matroid. Some characteristic-dependent linear rank inequalities are used for getting lower bounds on the linear information ratio over specific fields of these access structures. Taking n = 2t + 3 and M(n, t) = t + 1 in previous inequalities, we obtain two classes of constraints. The first must be satisfied by linear secret sharing schemes over fields whose characteristic does not divide t and the second must be satisfied by linear secret sharing schemes over fields whose characteristic divides t. We have the following proposition.

**Proposition 19.** Let  $t \in \mathbb{N}$ , t > 1 and let  $\mathbb{F}$  be a finite field. For any  $\mathcal{F}_t$  and  $\mathcal{N}_t$ , we have:

- $\sigma(\mathcal{F}_t) = \lambda_{\operatorname{char}(\mathbb{F})|t}(\mathcal{F}_t) = \kappa(\mathcal{F}_t) = \kappa^*_{\operatorname{char}(\mathbb{F})|t}(\mathcal{F}_t) = 1.$
- $\sigma\left(\mathcal{N}_{t}\right) = \lambda_{\operatorname{char}(\mathbb{F})\nmid t}\left(\mathcal{N}_{t}\right) = \kappa\left(\mathcal{N}_{t}\right) = \kappa_{\operatorname{char}(\mathbb{F})\nmid t}^{*}\left(\mathcal{N}_{t}\right) = 1.$
- $\lambda_{\operatorname{char}(\mathbb{F})|t}(\mathcal{F}_t) \ge \kappa^*_{\operatorname{char}(\mathbb{F})|t}(\mathcal{F}_t) \ge \frac{t+2}{t+1}$ .
- $\lambda_{\operatorname{char}(\mathbb{F})|t}(\mathcal{N}_t) \ge \kappa^*_{\operatorname{char}(\mathbb{F})|t}(\mathcal{N}_t) \ge \frac{t+3}{t+2}.$

*Proof.* It is clear that these access structures are ideal over fields where the associated matroid are representable. So, we have the optimal information ratios are equal to 1 over these fields. It remains for proving the last two items. Taking  $A_i = a_i$ ,  $B_i = b_i$  and C = c in the linear programming problem 7 with the constraints valid over fields whose characteristic does not divide t, The access structure  $\mathcal{F}_t$  holds that  $f(a_i) \leq v$ ,  $f(a_i) \leq v$ ,  $f(\emptyset) = 0$ , f(c) = 1,  $f(c \mid a_{[t+1]}) = f(c \mid a_i, b_i) = f(c; a_{[M(n,t)]-i}, b_i) = f(c; a_{[t+1]-i}) = f(c \mid a_i, b_i) = f(c; a_{[t+1]-i}, b_i) = 0$ . Thus, using the constraint obtained from the characteristic-dependent linear rank inequality n = 2t + 3 and M(n,t) = t + 1, we get

$$1 = f(c) \le \frac{1}{t+2} \sum_{i \in [t+1]} f(a_i) \le \frac{t+1}{t+2} v.$$

Therefore,  $\kappa^*_{\operatorname{char}(\mathbb{F})\nmid t}(\mathcal{F}_t) \geq v \geq \frac{t+2}{t+1}$ . In a similar way, we show the other inequalities.

## Acknowledgments

The author thanks the support provided by COLCIENCIAS in Conv. 727 and Carles Padró for the doctoral stay in Barcelona.

#### References

- A. Blasiak, R. Kleinberg and E. Lubetzky. Lexicographic Products and the Power of non-Linear Network Coding. IEEE Symposium on Foundations of Computer Science, pp. 609-618, 2011.
- [2] E. F. Brickell and D. M. Davenport, On the Classification of Ideal Secret Sharing, J. of Cryptology, 4:123-134, 1991.
- [3] R. Dougherty, C. Freiling and K. Zeger. Achievable Rate Regions for Network Coding. IEEE Transactions on Information Theory, 61(5): 2488–2509, 2015. Also in ArXiv 1311.4601, 2013.
- [4] O. Farràs, T. Kaced, S. Martín and C. Padró, Improving the Linear Programming Technique in the Search for Lower Bounds in Secret Sharing, 2018.
- [5] E. F. Freiling. Characteristic Dependent Linear Rank Inequalities and Applications to Network Coding. Dissertation for the Doctoral Degree. University of California, San Diego, 2014.
- [6] A. W. Ingleton. Representation of Matroids. Combinatorial Mathematics and its Applications, pp. 149-167, Oxford, 1969.
- [7] A. Jafari and S. Khazaei, On Abelian Secret Sharing: duality and separation, 2019.

- [8] S. Martín, C. Padró and A. Yang, Secret Sharing, Rank Inequalities, and Information Inequalities, 2015.
- [9] J. G. Oxley. Matroid Theory. Oxford University Press, New York, 1992.
- [10] C. Padró. Lecture Notes in Secret Sharing. Cryptology ePrint Archive: Report 674, 2012.
- [11] V. Peña-Macias and H. Sarria, Characteristic-Dependent Linear Rank Inequalities via Complementary Vector Spaces, Journal of Information & Optimization Sciences, DOI:10.1080/02522667.2019.1668157, 2020. 42(2): 345-369, 2021.
- [12] V. Peña-Macias and H. Sarria, Characteristic-Dependent Linear Rank Inequalities in 21 variables, Revista Academia Colombiana de Ciencias Exactas, Físicas y Naturales, 43(169): 765-770, 2019.
- [13] V. Peña-Macias and H. Sarria, Linear Programming Problems in Network Coding and Closure Operators via Partitions, Revista Selecciones Matemáticas, Universidad Nacional de trujillo, 6(2): 269-274, 2019.
- [14] P. D. Seymour, On Secret-Sharing Matroids, Journal of Combinatorial Theory, Series B, 56: 69-73, 1992.
- [15] A. Shen, D. Hammer, A. E. Romashchenko and N.K. Vereshchagin. Inequalities for Shannon Entropy and Kolmogorov Complexity. Journal of Computer and Systems Sciences, 60: 442-464, 2000.
- [16] D. R. Stinson, Decomposition Constructions for Secret-Sharing Schemes. IEEE Transactions on Information Theory, 40(1): 118-125, 1994.