# Ramsey-type results for path covers and path partitions

Shuya Chiba<sup>1</sup>\*

Michitaka Furuya<sup>2†</sup>

<sup>1</sup>Applied Mathematics, Faculty of Advanced Science and Technology, Kumamoto University,

> 2-39-1 Kurokami, Kumamoto 860-8555, Japan <sup>2</sup>College of Liberal Arts and Sciences,

> > Kitasato University,

1-15-1 Kitasato, Minami-ku, Sagamihara, Kanagawa 252-0373, Japan

#### Abstract

A family  $\mathcal{P}$  of subgraphs of G is called a *path cover* (resp. a *path partition*) of G if  $\bigcup_{P \in \mathcal{P}} V(P) = V(G)$  (resp.  $\dot{\bigcup}_{P \in \mathcal{P}} V(P) = V(G)$ ) and every element of  $\mathcal{P}$  is a path. The minimum cardinality of a path cover (resp. a path partition) of G is denoted by pc(G) (resp. pp(G)). In this paper, we characterize the forbidden subgraph conditions assuring us that pc(G) (or pp(G)) is bounded by a constant. Our main results introduce a new Ramsey-type problem.

*Key words and phrases.* path cover number, path partition number, forbidden subgraph, Ramsey number

AMS 2020 Mathematics Subject Classification. 05C38, 05C55.

### 1 Introduction

All graphs considered in this paper are finite, simple, and undirected. For terms and symbols not defined in this paper, we refer the reader to [2].

Let G be a graph. Let V(G) and E(G) denote the vertex set and the edge set of G, respectively. For a vertex  $x \in V(G)$ , let  $N_G(x)$  denote the neighborhood of x in G; thus  $N_G(x) = \{y \in V(G) : xy \in E(G)\}$ . For a subset X of V(G), let  $N_G(X) = (\bigcup_{x \in X} N_G(x)) \setminus X$ , and let G[X] (resp. G - X) denote the subgraph of G induced by X (resp.  $V(G) \setminus X$ ). Let  $\alpha(G)$  denote the independence number of

<sup>\*</sup>e-mail:schiba@kumamoto-u.ac.jp

<sup>&</sup>lt;sup>†</sup>e-mail:michitaka.furuya@gmail.com

G, i.e., the maximum cardinality of an independent set of G. Let  $K_n$ ,  $P_n$  and  $K_{1,n}$  denote the *complete graph* of order n, the *path* of order n and the *star* of order n+1, respectively. For two positive integers  $n_1$  and  $n_2$ , the *Ramsey number*  $R(n_1, n_2)$  is the minimum positive integer R such that any graph of order at least R contains a clique of cardinality  $n_1$  or an independent set of cardinality  $n_2$ .

For two graphs G and H, G is said to be H-free if G contains no induced copy of H. For a family  $\mathcal{H}$  of graphs, a graph G is said to be  $\mathcal{H}$ -free if G is H-free for every  $H \in \mathcal{H}$ . In this context, the members of  $\mathcal{H}$  are called forbidden subgraphs. For two families  $\mathcal{H}_1$  and  $\mathcal{H}_2$  of graphs, we write  $\mathcal{H}_1 \leq \mathcal{H}_2$  if for every  $H_2 \in \mathcal{H}_2$ , there exists  $H_1 \in \mathcal{H}_1$  such that  $H_1$  is an induced subgraph of  $H_2$ . The relation " $\leq$ " between two families of forbidden subgraphs was introduced in [7]. Note that if  $\mathcal{H}_1 \leq \mathcal{H}_2$ , then every  $\mathcal{H}_1$ -free graph is also  $\mathcal{H}_2$ -free.

Let  $\mathcal{A}$  be a family of graphs. A family  $\mathcal{P}$  of subgraphs of G is called an  $\mathcal{A}$ -cover of G if  $\bigcup_{P \in \mathcal{P}} V(P) = V(G)$  and each element of  $\mathcal{P}$  is isomorphic to a graph belonging to  $\mathcal{A}$ . Note that some elements of an  $\mathcal{A}$ -cover of G might have common vertices. An  $\mathcal{A}$ -cover  $\mathcal{P}$  of G is called an  $\mathcal{A}$ -partition of G if the elements of  $\mathcal{P}$  are pairwise vertexdisjoint. A  $\{P_i : i \geq 1\}$ -cover (resp. a  $\{P_i : i \geq 1\}$ -partition) of G is called a path cover (resp. a path partition) of G. Since  $\{G[\{x\}] : x \in V(G)\}$  is a path partition of G (and so a path cover of G), the minimum cardinality of a path cover (or a path partition) of any graph is well-defined. The value min $\{|\mathcal{P}| : \mathcal{P}$  is a path cover of G} (resp. min $\{|\mathcal{P}| : \mathcal{P}$  is a path partition of G}), denoted by pc(G) (resp. pp(G)), is called the path cover number (resp. the path partition number) of G. It is trivial that  $pc(G) \leq pp(G)$ . Since a graph G has a Hamiltonian path if and only if pp(G) = 1, the decision problem for the path partition number is a natural generalization of the Hamiltonian path problem. In fact, it has been widely studied in, for example, [13–17]. Throughout this paper, we implicitly use the following fact.

**Fact 1.1** Let G be a graph, and let  $\{X_1, X_2, \ldots, X_m\}$  be a partition of V(G). Then  $pc(G) \leq \sum_{1 \leq i \leq m} pc(G[X_i])$  and  $pp(G) \leq \sum_{1 \leq i \leq m} pp(G[X_i])$ .

In this paper, we focus on the following conditions concerning a family  $\mathcal H$  of forbidden subgraphs:

- (A1) There exists a constant  $c_1 = c_1(\mathcal{H})$  such that  $pc(G) \leq c_1$  for every connected  $\mathcal{H}$ -free graph G.
- (A2) There exists a constant  $c_2 = c_2(\mathcal{H})$  such that  $pp(G) \leq c_2$  for every connected  $\mathcal{H}$ -free graph G.

Our main aim is to characterize the finite families  $\mathcal{H}$  of connected graphs satisfying (A1) or (A2).




Figure 1: Graphs  $K_m^*$ ,  $F_{m,n}^{(1)}$ ,  $F_{m,n}^{(2)}$ ,  $F_{m,n}^{(3)}$  and  $F_{m,n}^{(4)}$ 

Let m and n be two positive integers. We define five graphs which will be used as forbidden subgraphs in our main result (see Figure 1).

- Let  $K_m^*$  denote the graph with  $V(K_m^*) = \{x_i, y_i : 1 \le i \le m\}$  and  $E(K_m^*) = \{x_i x_j : 1 \le i < j \le m\} \cup \{x_i y_i : 1 \le i \le m\}.$
- Let  $A := \{x_1, x_2\} \cup \{y_i : 1 \le i \le m\} \cup \{z_i : 1 \le i \le n\}$ . We define four graphs as follows:
  - Let  $F_{m,n}^{(1)}$  denote the graph on A such that  $E(F_{m,n}^{(1)}) = \{x_1x_2, x_1y_1, x_1z_1\} \cup \{y_iy_{i+1}: 1 \le i \le m-1\} \cup \{z_iz_{i+1}: 1 \le i \le n-1\}.$
  - Let  $F_{m,n}^{(2)}$  is the graph obtained from  $F_{m,n}^{(1)}$  by adding the edge  $y_1 z_1$ .
  - Let  $F_{m,n}^{(3)}$  denote the graph on A such that  $E(F_{m,n}^{(3)}) = \{x_1y_1, x_1z_1, x_2y_1, x_2z_1\} \cup \{y_iy_{i+1} : 1 \le i \le m-1\} \cup \{z_iz_{i+1} : 1 \le i \le n-1\}.$
  - Let  $F_{m,n}^{(4)}$  is the graph obtained from  $F_{m,n}^{(3)}$  by adding the edge  $y_1 z_1$ .

Our first main result is the following, which is proved in Section 2.

**Theorem 1.2** Let  $\mathcal{H}$  be a finite family of connected graphs. Then the following hold:

- (i) The family  $\mathfrak{H}$  satisfies (A1) if and only if  $\mathfrak{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$  for an integer  $n \geq 2$ .
- (ii) The family  $\mathfrak{H}$  satisfies (A2) if and only if  $\mathfrak{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}, F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$  for an integer  $n \geq 2$ .

Our motivation derives from two different lines of research. The first one is forbidden subgraph conditions for the existence of a Hamiltonian path. Now we focus on the condition that

every connected  $\mathcal{H}$ -free graph (of sufficiently large order) has a Hamiltonian path (1)

for a family  $\mathcal{H}$  of connected graphs. Duffus et al. [3] proved  $\mathcal{H} = \{K_{1,3}, K_3^*\}$  satisfies (1), and Faudree and Gould [5] showed that if a family  $\mathcal{H}$  satisfying (1) consists of two connected graphs, then  $\mathcal{H} \leq \{K_{1,3}, K_3^*\}$ . Thereafter a series by Gould and Harris [10–12] characterized the families  $\mathcal{H}$  of connected graphs with  $|\mathcal{H}| = 3$  satisfying (1). Since a graph has a Hamiltonian path if and only if its path cover number (or its path partition number) is exactly one, it is natural to study the forbidden subgraph conditions assuring us that the path cover/partition number is bounded by a constant as a next step. Our main result gives a complete solution for the problem in a sense.

Our second motivation is an analysis of gap between minimum  $\mathcal{A}$ -covers and minimum  $\mathcal{A}$ -partitions. A path cover/partition, which are main topic in this paper, is just one of examples of  $\mathcal{A}$ -cover/partition problems, and there also exist many other cover/partition problems. One of representative other examples is the case where  $\mathcal{A}$  is the family of all stars, where we regard  $K_1$  as one of stars. If we define the star cover number and the star partition number in the same way as pc(G)and pp(G), we can easily verify that the values are always equivalent. (Indeed, the star cover number also equals to the *domination number*, which is one of classical invariants in graph theory. The forbidden subgraph conditions assuring us that the domination number is bounded by a constant were characterized in [8].) On the other hand, as it is evident from Theorem 1.2, there is a gap between the path cover number and the path partition number. By Theorem 1.2, we discover that  $F_{n,n}^{(3)}$  and  $F_{n,n}^{(4)}$  play an important role for essential structures giving such a gap.

We also obtain an analogy of Theorem 1.2 considering a cycle cover/partition problem. A  $\{K_1, K_2, C_i : i \geq 3\}$ -cover (resp. a  $\{K_1, K_2, C_i : i \geq 3\}$ -partition) of G is called a *cycle cover* (resp. a *cycle partition*) of G. The value min $\{|\mathcal{P}| :$  $\mathcal{P}$  is a cycle cover of  $G\}$  (resp. min $\{|\mathcal{P}| : \mathcal{P} \text{ is a cycle partition of } G\}$ ), denoted by cc(G) (resp. cp(G)), is called the *cycle cover number* (resp. the *cycle partition number*) of G. Since trees (or graphs having a vertex of degree one) has no  $\{C_i : i \geq 3\}$ -cover, one sometimes focuses on cycle covers/partitions of general graphs instead of  $\{C_i : i \geq 3\}$ -covers/partitions (see, for example, [4,6]). In Section 3, as the second result, we characterize the families  $\mathcal{H}$  of forbidden subgraphs satisfying one of the following:

- (A'1) There exists a constant  $c_1 = c_1(\mathcal{H})$  such that  $cc(G) \leq c_1$  for every connected  $\mathcal{H}$ -free graph G.
- (A'2) There exists a constant  $c_2 = c_2(\mathcal{H})$  such that  $cp(G) \le c_2$  for every connected  $\mathcal{H}$ -free graph G.

**Theorem 1.3** Let  $\mathcal{H}$  be a family of connected graphs. Then the following are equivalent.

- (i) The family  $\mathcal{H}$  satisfies (A'1).
- (ii) The family  $\mathcal{H}$  satisfies (A'2).
- (iii) For an integer  $n \ge 2$ ,  $\mathcal{H} \le \{K_{1,n}, K_n^*, P_n\}$ .

We conclude this section by defining a new Ramsey-type concept concerning the path cover/partition number. Let  $\mathcal{H}$  be a family of graphs. The path cover Ramsey number  $R^{\mathrm{pc}}(\mathcal{H})$  (resp. the path partition Ramsey number  $R^{\mathrm{pp}}(\mathcal{H})$ ) is the minimum positive integer R such that any connected graph G with  $\mathrm{pc}(G) \geq R$  (resp.  $\mathrm{pp}(G) \geq R$ ) contains an induced copy of an element of  $\mathcal{H}$ , where  $R^{\mathrm{pc}}(\mathcal{H}) = \infty$  (resp.  $R^{\mathrm{pp}}(\mathcal{H}) = \infty$ ) if such an integer does not exist. Then it follows from Theorem 1.2 that the following hold:

- (P1) For a finite family  $\mathcal{H}$  of connected graphs,  $R^{\mathrm{pc}}(\mathcal{H})$  is a finite number if and only if  $\mathcal{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$  for an integer  $n \geq 2$ .
- (P2) For a finite family  $\mathcal{H}$  of connected graphs,  $R^{\text{pp}}(\mathcal{H})$  is a finite number if and only if  $\mathcal{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}, F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$  for an integer  $n \geq 2$ .

Note that  $R^{\rm pc}(\mathcal{H}) = 2$  if and only if  $R^{\rm pp}(\mathcal{H}) = 2$ . As we mentioned above, it is known that  $R^{\rm pc}(\{K_{1,3}, K_3^*\}) = 2$  and the study of triples  $\{H_1, H_2, H_3\}$  of connected graphs with  $R^{\rm pc}(\{H_1, H_2, H_3\}) = 2$  is completed. Since the  $K_{1,3}$ -freeness tends to give an important structure to many Hamiltonian properties, one might be interested in a relationship between such new Ramsey-type values and  $K_{1,3}$ -freeness. Here we focus on the values  $R^{\rm pc}(\mathcal{H})$  and  $R^{\rm pp}(\mathcal{H})$  for the case where  $\mathcal{H}$  contains  $K_{1,3}$ . Note that for positive integers m and n with  $m+n \geq 3$ , all of  $F_{m,n}^{(1)}, F_{m,n}^{(3)}$  and  $F_{m,n}^{(4)}$  contain  $K_{1,3}$  as an induced copy. Thus if  $K_{1,3} \in \mathcal{H}$ , then

$$R^{\mathrm{pc}}(\mathcal{H}) = R^{\mathrm{pc}}(\mathcal{H} \setminus \{F_{m,n}^{(1)}, F_{m,n}^{(3)}, F_{m,n}^{(4)} : m \ge 1, \ n \ge 1, \ m + n \ge 3\})$$

and

$$R^{\rm pp}(\mathcal{H}) = R^{\rm pp}(\mathcal{H} \setminus \{F_{m,n}^{(1)}, F_{m,n}^{(3)}, F_{m,n}^{(4)} : m \ge 1, \ n \ge 1, \ m+n \ge 3\}).$$

Considering (P1) and (P2), we leave the following open problem which will be a next interesting target on this concept for readers.

**Problem 1** For positive integers p, q and r with  $p \ge 3$  and  $q+r \ge 4$  and for a family  $\mathcal{H}$  of graphs with  $\mathcal{H} \le \{K_{1,3}, K_p^*, F_{q,r}^{(2)}\}$ , determine the value  $R^{\mathrm{pc}}(\mathcal{H})$  and  $R^{\mathrm{pp}}(\mathcal{H})$ .

# 2 Proof of Theorem 1.2

### 2.1 The "if" parts of Theorem 1.2

In this subsection, we prove the following theorem, which implies that the "if" parts of Theorem 1.2 hold.

**Theorem 2.1** Let  $n \ge 2$  be an integer. Then the following hold:

- (i) There exists a constant  $c_1 = c_1(n)$  depending on n only such that  $pc(G) \le c_1$ for every connected  $\{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$ -free graph G.
- (ii) There exists a constant  $c_2 = c_2(n)$  depending on n only such that  $pp(G) \le c_2$ for every connected  $\{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}, F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$ -free graph G.

The following lemma is well-known (or it is also obtained from a result on digraph by Gallai and Milgram [9]). So many readers can skip are advised to skip the proof.

**Lemma 2.2** For a graph G,  $pc(G) \le pp(G) \le \alpha(G)$ .

*Proof.* Since a path partition of G is also a path cover of G, we have  $pc(G) \le pp(G)$ .

Let  $\mathcal{P}$  be a path partition of G with  $|\mathcal{P}| = pp(G)$ , and write  $\mathcal{P} = \{Q_i : 1 \leq i \leq pp(G)\}$ . For each i with  $1 \leq i \leq pp(G)$ , let  $x_i$  be an endvertex of  $Q_i$ . If  $x_i x_j \in E(G)$  for some  $1 \leq i < j \leq pp(G)$ , then the graph Q obtained from  $Q_i$  and  $Q_j$  by joining the edge  $x_i x_j$  is a path, and hence  $\mathcal{P}' = (\mathcal{P} \setminus \{Q_i, Q_j\}) \cup \{Q\}$  is a path partition of G with  $|\mathcal{P}'| = pp(G) - 1$ , which contradicts the definition of the path partition number. Thus  $\{x_i : 1 \leq i \leq pp(G)\}$  is an independent set of G, and hence  $pp(G) \leq \alpha(G)$ .  $\Box$ 

**Lemma 2.3** Let  $n \ge 2$  and  $\alpha \ge 1$  be integers. Let G be a  $\{K_{1,n}, K_n^*\}$ -free graph, and let X be a subset of V(G) with  $\alpha(G[X]) \le \alpha$ . Then  $\alpha(G[N_G(X)]) \le (n - 1)R(n, \alpha + 1) - 1$ .

Proof. By way of contradiction, we suppose that there exists a subset Y of  $N_G(X)$ such that Y is an independent set of G and  $|Y| = (n-1)R(n, \alpha + 1)$ . Take a subset  $X_0$  of X with  $Y \subseteq N_G(X_0)$  so that  $|X_0|$  is as small as possible. If  $|X_0| \leq R(n, \alpha + 1) - 1$ , then  $\frac{|Y|}{|X_0|} \geq \frac{(n-1)R(n,\alpha+1)}{R(n,\alpha+1)-1} > n-1$ , and hence there exists a vertex  $x_0 \in X_0$  with  $|N_G(x_0) \cap Y| \geq n$ , which contradicts the  $K_{1,n}$ -freeness of G. Thus  $|X_0| \geq R(n, \alpha + 1)$ . Since  $\alpha(G[X_0]) \leq \alpha(G[X]) \leq \alpha$ , this implies that there exists




Figure 2: Path P and sets  $X_i$  and Y

a subset  $X_1$  of  $X_0$  such that  $X_1$  is a clique of G and  $|X_1| = n$ . By the minimality of  $X_0$ ,  $(N_G(x) \cap Y) \setminus N_G(X_0 \setminus \{x\}) \neq \emptyset$  for every  $x \in X_0$ . For each  $x \in X_0$ , let  $y_x \in (N_G(x) \cap Y) \setminus N_G(X_0 \setminus \{x\})$ . Then  $X_1 \cup \{y_x : x \in X_1\}$  induces a copy of  $K_n^*$  in G, which contradicts the  $K_n^*$ -freeness of G.  $\Box$ 

In the remainder of this subsection, we fix an integer  $n \ge 2$  and a connected  $\{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$ -free graph G. Set  $n_0 = \max\{\lceil \frac{n^2 - n - 2}{2} \rceil, n\}$ . Take a longest induced path P of G, and write  $P = u_1 u_2 \cdots u_m$ . Let  $X_0 = \{u_i : 1 \le i \le n_0 \text{ or } m - n_0 + 1 \le i \le m\}$  and  $Y = N_G(V(P) \setminus X_0) \setminus (X_0 \cup N_G(X_0))$ . Note that if  $|V(P)| \le 2n_0$ , then  $X_0 = V(P)$  and  $Y = \emptyset$ . We further remark that  $N_G(y) \cap V(P) \subseteq \{u_i : n_0 + 1 \le i \le m - n_0\}$  for every  $y \in Y$  (and in the remainder of this subsection, we frequently use the fact without mentioning). For each i with  $n_0 + 1 \le i \le m - n_0$ , let  $Y_i = \{y \in Y : \min\{j : n_0 + 1 \le j \le m - n_0, yu_j \in E(G)\} = i\}$ . Now we recursively define the sets  $X_i$   $(i \ge 1)$  as follows: Let  $X_1 = N_G(X_0) \setminus V(P)$ , and for i with  $i \ge 2$ , let  $X_i = N_G(X_{i-1}) \setminus (V(P) \cup Y \cup (\bigcup_{1 \le j \le i-1} X_j))$  (see Figure 2). Then  $X_1 \cap Y = \emptyset$  and  $X_1 \cup Y = N_G(V(P))$ .

#### **Lemma 2.4** We have $X_{2n_0} = \emptyset$ .

Proof. Suppose that  $X_{2n_0} \neq \emptyset$ . Let  $x_{2n_0} \in X_{2n_0}$ . Then we can recursively take a vertex  $x_{2n_0-i} \in N_G(x_{2n_0-i+1}) \cap X_{2n_0-i}$  for i with  $1 \leq i \leq 2n_0$ . Note that  $x_0 = u_k$  for some k with  $1 \leq k \leq n_0$  or  $m - n_0 + 1 \leq k \leq m$ . By symmetry, we may assume that  $1 \leq k \leq n_0$ . Under this condition, we choose k so that k is as large as possible. Since  $x_0x_1 \cdots x_{2n_0}$  is an induced path of G having  $2n_0 + 1$  vertices, it follows from the maximality of P that  $|V(P)| \geq 2n_0 + 1$ . In particular,  $V(P) \setminus X_0 \neq \emptyset$ .




Figure 3: An example for s = 5

If  $N_G(x_1) \cap (V(P) \setminus X_0) = \emptyset$ , then  $x_{2n_0} x_{2n_0-1} \cdots x_1 u_k u_{k+1} \cdots u_{m-n_0}$  is an induced path of G having  $2n_0 + m - n_0 - k + 1 \ (\geq m + 1)$  vertices, which contradicts the maximality of P. Thus  $N_G(x_1) \cap (V(P) \setminus X_0) \neq \emptyset$ .

Now we consider an operation recursively defining integers  $j_1, j_2, \ldots$  with  $1 \leq j_p \leq m$   $(p \geq 1)$  and  $j_1 < j_2 < \cdots$  as follows (see Figure 3): Let  $j_1 = \min\{j : 1 \leq j \leq m, x_1u_j \in E(G)\}$ . For  $p \geq 2$ , we assume that the integer  $j_{p-1}$  has defined. If  $\{j : j_{p-1} + 2 \leq j \leq m, x_1u_j \in E(G)\} \neq \emptyset$ , we let  $j_p = \min\{j : j_{p-1} + 2 \leq j \leq m, x_1u_j \in E(G)\}$ ; otherwise, we finish the operation. Let  $S = \{u_{j_p} : p \geq 1\}$ , and set s = |S|. Let  $j^* = \max\{j : 1 \leq j \leq m, x_1u_j \in E(G)\}$ . Note that  $j^* \in \{j_s, j_s + 1\}$ . Since  $j_p \geq j_{p-1}+2$ , S is an independent set of G. Since G is  $K_{1,n}$ -free and  $\{x_1, x_2\} \cup S$  induces a copy of  $K_{1,s+1}$  in G, we have  $s + 1 \leq n - 1$ .

For the moment, suppose that s = 1. Since  $N_G(x_1) \cap \{u_j : 1 \le j \le n_0\} \ne \emptyset$  and  $N_G(x_1) \cap (V(P) \setminus X_0) \ne \emptyset$ , this forces  $N_G(x_1) = \{u_{n_0}, u_{n_0+1}\}$ . Then

$$\{x_1, x_2, u_{n_0}, u_{n_0-1}, \dots, u_{n_0-n+1}, u_{n_0+1}, u_{n_0+2}, \dots, u_{n_0+n}\}$$

induces a copy of  $F_{n,n}^{(2)}$  in G, which is a contradiction. Thus  $s \ge 2$ .

Let  $Q_1 = u_1 u_2 \cdots u_{j_1}$  and  $Q_{s+1} = u_{j^*} u_{j^*+1} \cdots u_m$  be subpaths of P. For p with  $2 \leq p \leq s$ , let  $Q_p = u_{j_{p-1}+2} u_{j_{p-1}+3} \cdots u_{j_p}$  be a subpath of P. Then  $V(P) \setminus (\bigcup_{1 \leq p \leq s+1} V(Q_p)) = \{u_{j_p+1} : 1 \leq p \leq s-1\}$ , and hence

$$2n_0 + 1 \le |V(P)|$$

$$= \left| V(P) \setminus \left( \bigcup_{1 \le p \le s+1} V(Q_p) \right) \right| + \left| \bigcup_{1 \le p \le s+1} V(Q_p) \right|$$

$$\le (s-1) + \sum_{1 \le p \le s+1} |V(Q_p)|$$

$$= \sum_{1 \le p \le s+1} (|V(Q_p)| + 1) - 2.$$

This implies that  $\sum_{1 \le p \le s+1} (|V(Q_p)|+1) \ge 2n_0+3 \ge 2\lceil \frac{n^2-n-2}{2}\rceil+3 \ge n^2-n+1$ . If  $|V(Q_p)| \le n-1$  for all p with  $1 \le p \le s+1$ , then  $n^2-n+1 \le \sum_{1 \le p \le s+1} (|V(Q_p)|+1) \le (s+1)n \le (n-1)n$ , which is a contradiction. Thus  $|V(Q_q)| \ge n$  for some q with  $1 \le q \le s+1$ .

Note that  $|N_G(x_1) \cap V(Q_q)| = 1$ . Write  $N_G(x_1) \cap V(Q_q) = \{u_j\}$ . If  $q \neq s+1$ , then  $j \in \{j_p : 1 \leq p \leq s\}$ ; otherwise,  $j = j^* (\in \{j_s, j_{s+1}\})$ . Since  $u_j$  is an endvertex of  $Q_q$ , there exists a subpath Q of  $Q_q$  such that  $u_j$  is an endvertex of Q and |V(Q)| = n. Since  $|S| \geq 2$ , we can take a vertex  $v \in S$  as follows: If  $q \neq s+1$ , let  $v \in S \setminus \{u_j\}$ ; otherwise (i.e.,  $j = j^*$ ), let  $v = u_{j_1}$ . Then by the definition of  $Q_p$   $(1 \leq p \leq s+1)$ ,  $N_G(v) \cap V(Q_q) = \emptyset$ . Since  $2n_0 \geq 2n \geq n+1$ , the vertices  $x_i$  with  $2 \leq i \leq n+1$  have been defined, and hence this implies that  $\{x_1, v, x_2, x_3, \dots, x_{n+1}\} \cup V(Q)$  induces a copy of  $F_{n,n}^{(1)}$  in G, which is a contradiction.  $\Box$ 

**Lemma 2.5** Let *i* be an integer with  $n_0 + 1 \le i \le m - n_0$ , and let  $y \in Y_i$ . Then the following hold:

- (i) If  $yu_{i+1} \notin E(G)$ , then  $N_G(y) \cap V(P) = \{u_i, u_{i+2}\}$ .
- (ii) We have  $Y_{m-n_0} = \emptyset$ .
- (iii) If G is  $F_{n,n}^{(3)}$ -free, then  $yu_{i+1} \in E(G)$ .

Proof.

(i) Suppose that  $yu_{i+1} \notin E(G)$  and  $N_G(y) \cap V(P) \neq \{u_i, u_{i+2}\}$ . Let  $k = \max\{j : n_0 + 1 \le j \le m - n_0, yu_j \in E(G)\}$ . If k = i (i.e.,  $N_G(y) \cap V(P) = \{u_i\}$ ), then

$$\{u_i, y, u_{i-1}, u_{i-2}, \dots, u_{i-n}, u_{i+1}, u_{i+2}, \dots, u_{i+n}\}$$

induces a copy of  $F_{n,n}^{(1)}$  in G, which is a contradiction. Since  $yu_{i+1} \notin E(G)$  and  $N_G(y) \cap V(P) \neq \{u_i, u_{i+2}\}$ , this forces  $k \geq i+3$ . Then

$$\{u_i, u_{i+1}, u_{i-1}, u_{i-2}, \dots, u_{i-n}, y, u_k, u_{k+1}, \dots, u_{k+n-2}\}$$

induces a copy of  $F_{n,n}^{(1)}$  in G, which is a contradiction.

- (ii) By (i), if there exists a vertex  $y \in Y_{m-n_0}$ , then it follows that  $yu_{m-n_0+1} \in E(G)$ or  $N_G(y) \cap V(P) = \{u_{m-n_0}, u_{m-n_0+2}\}$ , and in particular,  $N_G(y) \cap X_0 \neq \emptyset$ , which contradicts the definition of Y. Thus we have  $Y_{m-n_0} = \emptyset$ .
- (iii) Suppose that G is  $F_{n,n}^{(3)}$ -free and  $yu_{i+1} \notin E(G)$ . Then it follows from (i) that  $N_G(y) \cap V(P) = \{u_i, u_{i+2}\}$ , and hence

$$\{y, u_{i+1}, u_i, u_{i-1}, \dots, u_{i-n+1}, u_{i+2}, u_{i+3}, \dots, u_{i+n+1}\}$$

induces a copy of  $F_{n,n}^{(3)}$  in G, which is a contradiction.  $\Box$ 

**Lemma 2.6** We have  $V(G) = V(P) \cup N_G(V(P)) \cup (\bigcup_{2 \le i \le 2n_0 - 1} X_i).$ 

Proof. Suppose that  $V(G) \neq V(P) \cup N_G(V(P)) \cup (\bigcup_{2 \leq i \leq 2n_0-1} X_i)$ . Since G is connected, there exists a vertex  $z \in V(G) \setminus (V(P) \cup N_G(V(P)) \cup (\bigcup_{2 \leq i \leq 2n_0-1} X_i))$  adjacent to a vertex  $y \in V(P) \cup N_G(V(P)) \cup (\bigcup_{2 \leq i \leq 2n_0-1} X_i)$  in G. By Lemma 2.4 and the definition of  $X_i$  and Y, this implies that  $y \in Y$ . Let i be the integer such that  $y \in Y_i$ . Then by Lemma 2.5(ii),  $n_0 + 1 \leq i \leq m - n_0 - 1$ . Let  $k = \max\{j : n_0 + 1 \leq j \leq m - n_0, yu_j \in E(G)\}$ . By Lemma 2.5(ii),  $k \geq i + 1$ . If k = i + 1, then

 $\{y, z, u_i, u_{i-1}, \dots, u_{i-n+1}, u_{i+1}, u_{i+2}, \dots, u_{i+n}\}$ 

induces a copy of  $F_{n,n}^{(2)}$  in G; if  $k \ge i+2$ , then

$$\{y, z, u_i, u_{i-1}, \dots, u_{i-n+1}, u_k, u_{k+1}, \dots, u_{k+n-1}\}$$

induces a copy of  $F_{n,n}^{(1)}$  in G. In either case, we obtain a contradiction.  $\Box$ 

Now we recursively define the values  $\alpha_i$   $(i \ge 0)$  as follows: Let  $\alpha_0 = 2\lceil \frac{n_0}{2} \rceil$ , and for *i* with  $i \ge 1$ , let  $\alpha_i = (n-1)R(n, \alpha_{i-1} + 1) - 1$ .

**Lemma 2.7** For an integer *i* with  $i \ge 0$ ,  $\alpha(G[X_i]) \le \alpha_i$ .

Proof. We proceed by induction on *i*. If  $|V(P)| \leq 2n_0$ , then  $G[X_0]$  equals to *P*, and hence  $\alpha(G[X_0]) = \alpha(P) = \lceil \frac{|V(P)|}{2} \rceil \leq n_0 \leq \alpha_0$ ; if  $|V(P)| \geq 2n_0 + 1$ , then  $G[X_0]$ consists of two components each of which is a path of order  $n_0$ , and hence  $\alpha(G[X_0]) = 2\lceil \frac{n_0}{2} \rceil = \alpha_0$ . In either case, we have  $\alpha(G[X_0]) \leq \alpha_0$ . Thus we may assume that  $i \geq 1$ , and suppose that  $\alpha(G[X_{i-1}]) \leq \alpha_{i-1}$ . Since  $X_i \subseteq N_G(X_{i-1})$ , it follows from Lemma 2.3 that  $\alpha(G[X_i]) \leq \alpha(G[N_G(X_{i-1})]) \leq (n-1)R(n, \alpha_{i-1}+1) - 1 = \alpha_i$ , as desired.  $\Box$ 

Note that the value  $\sum_{1 \le i \le 2n_0 - 1} \alpha_i$  is a constant depending on *n* only. Thus, considering Lemmas 2.2, 2.6 and 2.7, it suffices to show that

- $pc(G[V(P) \cup Y])$  is bounded by a constant depending on n only, and
- if G is  $\{F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$ -free, then  $pp(G[V(P) \cup Y])$  is bounded by a constant depending on n only.

Hence the following lemma completes the proof of Theorem 2.1.

**Lemma 2.8** (i) We have  $pc(G[V(P) \cup Y]) \le max\{3n - 6, 1\}$ .

(ii) If G is  $\{F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$ -free, then there exists a Hamiltonian path of  $G[V(P) \cup Y]$ , i.e.,  $pp(G[V(P) \cup Y]) = 1$ . *Proof.* If  $Y = \emptyset$ , then P is a Hamiltonian path of  $G[V(P) \cup Y]$ , and hence  $pc(G[V(P) \cup Y]) = pp(G[V(P) \cup Y]) = 1$ . Thus we may assume that  $Y \neq \emptyset$ . By Lemma 2.5(ii),  $Y_{m-n_0} = \emptyset$ .

We first prove (i). Fix an integer i with  $n_0+1 \leq i \leq m-n_0-1$ . Let  $Y_{i,1} = \{y \in Y_i : yu_{i+1} \in E(G)\}$  and  $Y_{i,2} = Y_i \setminus Y_{i,1}$ . Then by Lemma 2.5(i),  $N_G(y) \cap V(P) = \{u_i, u_{i+2}\}$  for all  $y \in Y_{i,2}$ . Let  $j \in \{1,2\}$ . If there exists an independent set  $U \subseteq Y_{i,j}$  of G with |U| = n - 1, then  $\{u_{i-1}, u_i\} \cup U$  induces a copy of  $K_{1,n}$  in G, which is a contradiction. Thus  $\alpha(G[Y_{i,j}]) \leq n - 2$ . Since  $Y \neq \emptyset$ , i.e.,  $Y_{p,q} \neq \emptyset$  for some p and q with  $n_0 + 1 \leq p \leq m - n_0 - 1$  and  $q \in \{1,2\}$ , this implies that  $n \geq 3$ . By Lemma 2.2, there exists a path partition  $\mathcal{P}_{i,j} = \{Q_{i,j}^{(1)}, Q_{i,j}^{(2)}, \ldots, Q_{i,j}^{(s_{i,j})}\}$  of  $G[Y_{i,j}]$  with  $s_{i,j} \leq n - 2$ , where  $\mathcal{P}_{i,j} = \emptyset$  and  $s_{i,j} = 0$  if  $Y_{i,j} = \emptyset$ . For an integer t with  $1 \leq t \leq n - 2$ , if  $t \leq s_{i,j}$ , let  $R_{i,j}^{(t)}$  be the path  $u_i vQ_{i,j}^{(t)}wu_{i+j}$ , where  $\{v, w\}$  is the set of endvertices of  $Q_{i,j}^{(t)}$ ; otherwise, let  $R_{i,j}^{(t)}$  be the path between  $u_i$  and  $u_{i+j}$  on P (i.e.,  $R_{i,1}^{(t)} = u_i u_{i+1}$  and  $R_{i,2}^{(t)} = u_i u_{i+1} u_{i+2}$ ). We define the value  $\xi_2$  (resp.  $\xi_3$ ) with  $\xi_2 = m - n_0$  or  $\xi_2 = m - n_0 - 1$  (resp.  $\xi_3 = m - n_0 - 1$  or  $\xi_3 = m - n_0$ ) according as m is odd or even. Let

$$R_{1}^{(t)} = u_{1}u_{2}\cdots u_{n_{0}+1}R_{n_{0}+1,1}^{(t)}u_{n_{0}+2}R_{n_{0}+2,1}^{(t)}u_{n_{0}+3}\cdots u_{m-n_{0}-1}R_{m-n_{0}-1,1}^{(t)}u_{m-n_{0}}u_{m-n_{0}+1}\cdots u_{m},$$

$$R_{2}^{(t)} = u_{1}u_{2}\cdots u_{n_{0}+1}R_{n_{0}+1,2}^{(t)}u_{n_{0}+3}R_{n_{0}+3,2}^{(t)}u_{n_{0}+5}\cdots u_{\xi_{2}-2}R_{\xi_{2}-2,2}^{(t)}u_{\xi_{2}}u_{\xi_{2}+1}\cdots u_{m},$$
 and
$$R_{3}^{(t)} = u_{1}u_{2}\cdots u_{n_{0}+2}R_{n_{0}+2,2}^{(t)}u_{n_{0}+4}R_{n_{0}+4,2}^{(t)}u_{n_{0}+6}\cdots u_{\xi_{3}-2}R_{\xi_{3}-2,2}^{(t)}u_{\xi_{3}}u_{\xi_{3}+1}\cdots u_{m}.$$

Then we easily verify that  $\{R_a^{(t)} : a \in \{1, 2, 3\}, 1 \le t \le n - 2\}$  is a path cover of  $G[V(P) \cup Y]$  having cardinality at most 3(n-2), which proves (i).

Next we prove (ii). Suppose that G is  $\{F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$ -free. We start with the following claim.

Claim 2.1 For an integer i with  $n_0 + 1 \le i \le m - n_0 - 1$ ,  $\{u_i, u_{i+1}\} \cup Y_i$  is a clique of G.

Proof. Suppose that there exist two vertices  $y, y' \in \{u_i, u_{i+1}\} \cup Y_i$  with  $yy' \notin E(G)$ . By the definition of  $Y_i$  and Lemma 2.5(iii), every vertex in  $Y_i$  is adjacent to both  $u_i$ and  $u_{i+1}$  in G. Thus  $y, y' \in Y_i$ . Recall that  $N_G(Y) \cap V(P) \subseteq \{u_j : n_0 + 1 \leq j \leq m - n_0\}$ . Let  $k = \max\{j : n_0 + 1 \leq j \leq m - n_0, N_G(u_j) \cap \{y, y'\} \neq \emptyset\}$ . We may assume that  $yu_k \in E(G)$ . Note that  $k \geq i + 1$ . If k = i + 1, then

$$\{y, y', u_i, u_{i-1}, \dots, u_{i-n+1}, u_{i+1}, u_{i+2}, \dots, u_{i+n}\}$$

induces a copy of  $F_{n,n}^{(4)}$  in G, which is a contradiction. Thus  $k \ge i+2$ . If  $y'u_k \in E(G)$ , then

$$\{y, y', u_i, u_{i-1}, \dots, u_{i-n+1}, u_k, u_{k+1}, \dots, u_{k+n-1}\}$$

induces a copy of  $F_{n,n}^{(3)}$  in G; if  $y'u_k \notin E(G)$ , then

$$\{u_i, y', u_{i-1}, u_{i-2}, \dots, u_{i-n}, y, u_k, u_{k+1}, \dots, u_{k+n-2}\}$$

induces a copy of  $F_{n,n}^{(1)}$  in G. In either case, we obtain a contradiction.

For an integer i with  $n_0 + 1 \le i \le m - n_0 - 1$ , it follows from Claim 2.1 that there exists a Hamiltonian path  $R_i$  of  $G[\{u_i, u_{i+1}\} \cup Y_i]$  with the endvertices  $u_i$  and  $u_{i+1}$ . Then

 $u_{1}u_{2}\cdots u_{n_{0}+1}R_{n_{0}+1}u_{n_{0}+2}R_{n_{0}+2}u_{n_{0}+3}\cdots u_{m-n_{0}-1}R_{m-n_{0}-1}u_{m-n_{0}}u_{m-n_{0}+1}\cdots u_{m}$ is a Hamiltonian path of  $G[V(P)\cup(\bigcup_{n_{0}+1\leq i\leq m-n_{0}-1}Y_{i})]$  (=  $G[V(P)\cup Y]$ ), as desired.  $\Box$ 

### 2.2 The "only if" parts of Theorem 1.2

Let  $s \ge 2$  and  $t \ge 3$  be integers, and let  $Q_i = u_i^{(1)} u_i^{(2)} \cdots u_i^{(t)}$   $(1 \le i \le s)$  be s pairwise vertex-disjoint paths. We define four graphs.

- Let  $H_{s,t}^{(1)}$  be the graph obtained from the union of the paths  $Q_1, \ldots, Q_s$  by adding 2(s-1) vertices  $v_i, w_i$   $(1 \le i \le s-1)$  and 3(s-1) edges  $v_i w_i, v_i u_i^{(t)}, v_i u_{i+1}^{(1)}$   $(1 \le i \le s-1)$ .
- Let  $H_{s,t}^{(2)}$  be the graph obtained from  $H_{s,t}^{(1)}$  by adding s-1 edges  $u_i^{(t)}u_{i+1}^{(1)}$   $(1 \le i \le s-1)$ .
- Let  $H_{s,t}^{(3)}$  be the graph obtained from the union of the paths  $Q_1, \ldots, Q_s$  by adding 2(s-1) vertices  $v_i, w_i \ (1 \le i \le s-1)$  and 4(s-1) edges  $v_i u_i^{(t)}, v_i u_{i+1}^{(1)}, w_i u_{i+1}^{(t)}, w_i u_{i+1}^{(1)} \ (1 \le i \le s-1)$ .
- Let  $H_{s,t}^{(4)}$  be the graph obtained from  $H_{s,t}^{(3)}$  by adding s-1 edges  $u_i^{(t)}u_{i+1}^{(1)}$   $(1 \le i \le s-1)$ .

**Lemma 2.9** We have  $pc(H_{s,t}^{(1)}) = pc(H_{s,t}^{(2)}) = \lceil \frac{s+1}{2} \rceil$ .

*Proof.* Note that  $u_1^{(1)}, u_s^{(t)}, w_i$   $(1 \le i \le s-1)$  have degree one in  $H_{s,t}^{(2)}$ . Since a path contains at most two vertices of degree at most one,  $pc(G) \ge \lceil \frac{l}{2} \rceil$  for every graph G where l is the number of the vertices of G having degree one. In particular, we have

$$\operatorname{pc}(H_{s,t}^{(2)}) \ge \left\lceil \frac{s+1}{2} \right\rceil.$$
(2)

If s is odd, let

$$\mathcal{P} = \left\{ H_{s,t}^{(1)} - \{ w_j : 1 \le j \le s - 1 \}, \ w_{2i-1}v_{2i-1}u_{2i}^{(1)}Q_{2i}u_{2i}^{(t)}v_{2i}w_{2i} : 1 \le i \le \frac{s-1}{2} \right\};$$

if s is even, let

$$\mathcal{P} = \left\{ H_{s,t}^{(1)} - \{ w_j : 1 \le j \le s - 1 \}, \ H_{s,t}^{(1)}[\{w_{s-1}\}], \ w_{2i-1}v_{2i-1}u_{2i}^{(1)}Q_{2i}u_{2i}^{(t)}v_{2i}w_{2i} : 1 \le i \le \frac{s-2}{2} \right\}.$$

Then we verify that  $\mathcal{P}$  is a path cover of  $H_{s,t}^{(1)}$  with  $|\mathcal{P}| = \lceil \frac{s+1}{2} \rceil$ . Furthermore, since  $H_{s,t}^{(1)}$  is a spanning subgraph of  $H_{s,t}^{(2)}$ , a path cover of  $H_{s,t}^{(1)}$  is also a path cover of  $H_{s,t}^{(2)}$ , and hence  $\operatorname{pc}(H_{s,t}^{(2)}) \leq \operatorname{pc}(H_{s,t}^{(1)}) \leq \lceil \frac{s+1}{2} \rceil$ . This together with (2) leads to the desired conclusion.  $\Box$ 

**Lemma 2.10** We have  $pp(H_{s,t}^{(3)}) = pp(H_{s,t}^{(4)}) = s$ .

*Proof.* We first prove that

$$pp(H_{s,t}^{(4)}) \ge s. \tag{3}$$

Let  $\mathcal{P}$  be a path partition of  $H_{s,t}^{(4)}$ . It suffices to show that  $|\mathcal{P}| \geq s$ . For each i with  $1 \leq i \leq s$ , let  $R_i$  be the unique element of  $\mathcal{P}$  containing  $u_i^{(2)}$ . We remark that  $R_i$  might equal to  $R_j$  for some  $1 \leq i < j \leq s$ . Let  $I = \{i : 1 \leq i \leq s-1, R_i = R_{i+1}\}$ , and write  $I = \{i_1, i_2, \ldots, i_h\}$  with  $i_1 < i_2 < \ldots < i_h$  where h = 0 if  $I = \emptyset$ . For integers i and i' with  $1 \leq i < i' \leq s$ , any paths of  $H_{s,t}^{(4)}$  joining  $u_i^{(2)}$  and  $u_{i'}^{(2)}$  contain every vertex in  $\{u_j^{(2)} : i < j < i'\}$ . This implies that if  $R_i = R_{i'}$  with  $1 \leq i < i' \leq s$ , then i' - i + 1 paths  $R_j$   $(i \leq j \leq i')$  are equal. In particular, we have  $|\{R_i : 1 \leq i \leq s\}| = s - h$ .

Fix an integer l with  $1 \leq l \leq h$ . Then for every path R of  $H_{s,t}^{(4)}$  joining  $u_{i_l}^{(2)}$  and  $u_{i_l+1}^{(2)}$ , we easily verify that

- $\{u_{i_l}^{(t)}, u_{i_l+1}^{(1)}\} \subseteq V(R)$ , and
- $v_{i_l} \notin V(R)$  or  $w_{i_l} \notin V(R)$ .

Since  $v_{i_l}w_{i_l} \notin E(H_{s,t}^{(4)})$ , this implies that there exists an element  $R'_{i_l}$  of  $\mathcal{P}$  such that either  $V(R'_{i_l}) = \{v_{i_l}\}$  or  $V(R'_{i_l}) = \{w_{i_l}\}$ . Therefore

$$\begin{aligned} |\mathcal{P}| &\geq |\{R_i : 1 \leq i \leq s\} \cup \{R'_{i_j} : 1 \leq j \leq h\}| \\ &= |\{R_i : 1 \leq i \leq s\}| + |\{R'_{i_j} : 1 \leq j \leq h\}| \\ &= (s-h) + h \\ &= s, \end{aligned}$$

which proves (3).

Since

$$\mathcal{P}' = \{H_{s,t}^{(3)} - \{w_j : 1 \le j \le s - 1\}, \ H_{s,t}^{(3)}[\{w_i\}] : 1 \le i \le s - 1\}$$

is a path partition of  $H_{s,t}^{(3)}$  with  $|\mathcal{P}'| = s$ . Furthermore, since  $H_{s,t}^{(3)}$  is a spanning subgraph of  $H_{s,t}^{(4)}$ , a path partition of  $H_{s,t}^{(3)}$  is also a path partition of  $H_{s,t}^{(4)}$ , and hence  $pp(H_{s,t}^{(4)}) \leq pp(H_{s,t}^{(3)}) \leq s$ . This together with (3) leads to the desired conclusion.

Now we prove the following proposition, which gives the "only if" parts of Theorem 1.2.

**Proposition 2.11** Let  $\mathcal{H}$  be a finite family of connected graphs.

- (i) If  $\mathcal{H}$  satisfies (A1), then  $\mathcal{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$  for an integer  $n \geq 2$ .
- (ii) If  $\mathcal{H}$  satisfies (A2), then  $\mathcal{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}, F_{n,n}^{(3)}, F_{n,n}^{(4)}\}$  for an integer  $n \geq 2$ .

*Proof.* Since  $\mathcal{H}$  is a finite family, the value  $p = \max\{|V(H)| : H \in \mathcal{H}\}$  is well-defined. If  $p \leq 2$ , then the desired conclusions trivially hold. Thus we may assume that  $p \geq 3$ .

We first suppose that  $\mathcal{H}$  satisfies (A1), and show that (i) holds. There exists a constant  $c_1 = c_1(\mathcal{H})$  such that  $pc(G) \leq c_1$  for every connected  $\mathcal{H}$ -free graph G. Since  $pc(K_{1,2c_1+1}) = c_1 + 1$  and  $pc(K_{2c_1+1}^*) = c_1 + 1$ , neither  $K_{1,2c_1+1}$  nor  $K_{2c_1+1}^*$  is  $\mathcal{H}$ -free. This implies that

$$\mathcal{H} \le \{K_{1,2c_1+1}, K_{2c_1+1}^*\}.$$
(4)

For each  $i \in \{1, 2\}$ , it follows from Lemma 2.9 that  $pc(H_{2c_1,p}^{(i)}) = \lceil \frac{2c_1+1}{2} \rceil = c_1+1$ , and hence  $H_{2c_1,p}^{(i)}$  is not  $\mathcal{H}$ -free, i.e.,  $H_{2c_1,p}^{(i)}$  contains an induced subgraph  $A_i$  isomorphic to an element of  $\mathcal{H}$ . Since  $|V(A_i)| \leq p$ , we have

- $|\{j: 1 \le j \le 2c_1, V(A_i) \cap V(Q_j) \ne \emptyset\}| \le 2$ , and
- $|\{j: 1 \le j \le 2c_1 1, V(A_i) \cap \{v_j, w_j\} \ne \emptyset\}| \le 1.$

This implies that  $A_i$  is an induced copy of  $F_{p,p}^{(i)}$ , and hence

$$\mathcal{H} \le \{F_{p,p}^{(1)}, F_{p,p}^{(2)}\}.$$
(5)

Let  $n = \max\{2c_1 + 1, p\}$ . Then by (4) and (5),  $\mathcal{H} \leq \{K_{1,n}, K_n^*, F_{n,n}^{(1)}, F_{n,n}^{(2)}\}$ , which proves (i).

Next we suppose that  $\mathcal{H}$  satisfies (A2), and show that (ii) holds. There exists a constant  $c_2 = c_2(\mathcal{H})$  such that  $pp(G) \leq c_2$  for every connected  $\mathcal{H}$ -free graph G. Since  $pp(G) \geq pc(G)$  for all graphs G,  $\mathcal{H}$  also satisfies (A1). Hence by (i), there exists an integer  $m \geq 2$  such that

$$\mathcal{H} \le \{K_{1,m}, K_m^*, F_{m,m}^{(1)}, F_{m,m}^{(2)}\}.$$
(6)

For each  $i \in \{3, 4\}$ , it follows from Lemma 2.10 that  $pp(H_{c_2+1,p}^{(i)}) = c_2 + 1$ , and hence  $H_{c_2+1,p}^{(i)}$  is not  $\mathcal{H}$ -free, i.e.,  $H_{c_2+1,p}^{(i)}$  contains an induced subgraph  $B_i$  isomorphic to an element of  $\mathcal{H}$ . Since  $|V(B_i)| \leq p$ , we have

- $|\{j: 1 \le j \le c_2 + 1, V(B_i) \cap V(Q_j) \ne \emptyset\}| \le 2$ , and
- $|\{j: 1 \le j \le c_2, V(B_i) \cap \{v_j, w_j\} \ne \emptyset\}| \le 1.$

This implies that  $B_i$  is an induced copy of  $F_{p,p}^{(i)}$ , and hence

$$\mathcal{H} \le \{F_{p,p}^{(3)}, F_{p,p}^{(4)}\}.$$
(7)

Let  $n' = \max\{m, p\}$ . Then by (6) and (7),  $\mathcal{H} \leq \{K_{1,n'}, K_{n'}^*, F_{n',n'}^{(1)}, F_{n',n'}^{(2)}, F_{n',n'}^{(3)}, F_{n',n'}^{(4)}\}$ , which proves (ii).  $\Box$ 

## 3 Proof of Theorem 1.3

In this section, we prove Theorem 1.3. We start with the following lemma, which is an analogy of Lemma 2.2.

**Lemma 3.1** For a graph G,  $cp(G) \le R(\alpha(G) + 1, \alpha(G) + 1) - 1$ .

Proof. Let  $\mathcal{P}$  be a cycle partition of G with  $|\mathcal{P}| = cp(G)$ , and write  $\mathcal{P} = \{Q_i : 1 \leq i \leq cp(G)\}$ . By way of contradiction, suppose that  $cp(G) \geq R(\alpha(G) + 1, \alpha(G) + 1)$ . For each integer i with  $1 \leq i \leq cp(G)$ , we define vertices  $x_i$  and  $y_i$  of  $Q_i$  as follows: If either  $Q_i \simeq K_2$  or  $Q_i$  is a cycle, let  $x_i$  and  $y_i$  be vertices of  $Q_i$  with  $x_iy_i \in E(Q_i)$ ; if  $Q_i \simeq K_1$ , let  $x_i = y_i = u$  where u is the unique vertex of  $Q_i$ . For integers i and j with  $1 \leq i < j \leq cp(G)$ , if  $\{x_ix_j, y_iy_j\} \subseteq E(G)$ , then we easily verify that there exists a spanning subgraph Q of  $G[V(Q_i) \cup V(Q_j)]$  such that either  $Q \simeq K_2$  or Q is a cycle, and hence  $\mathcal{P}' = (\mathcal{P} \setminus \{Q_i, Q_j\}) \cup \{Q\}$  is a cycle partition of G with  $|\mathcal{P}'| = cp(G) - 1$ , which contradicts the definition of the cycle partition number. Thus if  $x_ix_j \in E(G)$ , then  $y_iy_j \notin E(G)$ .

Let K be the complete graph on  $\{1, 2, ..., cp(G)\}$ , and color all edges of K by red or blue as follows: For integers i and j with  $1 \le i < j \le cp(G)$ , if  $x_i x_j \notin E(G)$ , we color the edge ij of K by red; if  $x_i x_j \in E(G)$  and  $y_i y_j \notin E(G)$ , we color the edge ij of K by blue. Since  $|V(K)| = cp(G) \ge R(\alpha(G) + 1, \alpha(G) + 1)$ , there exists a monochromatic clique I of K with  $|I| = \alpha(G) + 1$ . If I is a red clique of K, then  $\{x_i : i \in I\}$  is an independent set of G; if I is a blue clique of K, then  $\{y_i : i \in I\}$  is an independent set of G. In either case, we obtain a contradiction.  $\Box$ 

The following lemma was implicitly proved in [1]. (To keep the paper selfcontained, we give its proof.) **Lemma 3.2 (Choi et al. [1])** Let  $n \ge 2$  be an integer. There exists a constant c = c(n) depending on n only such that  $\alpha(G) \le c$  for every connected  $\{K_{1,n}, K_n^*, P_n\}$ -free graph G.

*Proof.* Let x be a vertex of G, and for an integer i with  $i \ge 0$ , let  $X_i$  be the set of vertices y of G such that the distance between x and y in G is exactly i. Note that  $X_0 = \{x\}$  and  $X_1 = N_G(x)$ . Since G is  $P_n$ -free,  $X_i = \emptyset$  for all  $i \ge n - 1$ . Since G is connected, this implies that

$$V(G) = \bigcup_{0 \le i \le n-2} X_i.$$
(8)

We recursively define the values  $\alpha_i$   $(i \ge 0)$  as follows: Let  $\alpha_0 = 1$ , and for *i* with  $i \ge 1$ , let  $\alpha_i = (n-1)R(n, \alpha_{i-1} + 1) - 1$ .

We prove that

$$\alpha(G[X_i]) \le \alpha_i \text{ for an integer } i \text{ with } 0 \le i \le n-2.$$
(9)

We proceed by induction on *i*. Since  $\alpha(G[X_0]) = 1 = \alpha_0$ , we may assume that  $i \ge 1$ and  $\alpha(G[X_{i-1}]) \le \alpha_{i-1}$ . Since  $X_i \subseteq N_G(X_{i-1})$ , it follows from Lemma 2.3 that  $\alpha(G[X_i]) \le \alpha(G[N_G(X_{i-1})]) \le (n-1)R(n, \alpha_{i-1}+1) - 1 = \alpha_i$ , as desired.

By (8) and (9), we have  $\alpha(G) \leq \sum_{0 \leq i \leq n-2} \alpha(G[X_i]) \leq \sum_{0 \leq i \leq n-2} \alpha_i$ . Since the value  $\sum_{0 \leq i \leq n-2} \alpha_i$  is a constant depending on n only, we obtain the desired conclusion.  $\Box$ 

Proof of Theorem 1.3. By the definition of cycle cover and cycle partition, "(ii)  $\implies$  (i)" clearly holds.

We show that "(iii)  $\implies$  (ii)" holds. Let  $n \ge 2$  be an integer, and let c = c(n) be the constant as in Lemma 3.2. It suffices to show that there exists a constant  $c_1 = c_1(n)$  depending on n only such that  $cp(G) \le c_1$  for every connected  $\{K_{1,n}, K_n^*, P_n\}$ free graph G. By the definition of c(n), we have  $\alpha(G) \le c$ . This together with Lemma 3.1 leads to  $cp(G) \le R(\alpha(G) + 1, \alpha(G) + 1) - 1 \le R(c + 1, c + 1) - 1$ . Since R(c+1, c+1) - 1 is a constant depending on n only, we obtain the desired conclusion.

Finally, we show that "(i)  $\implies$  (iii)" holds, which completes the proof of Theorem 1.3. Suppose that a family  $\mathcal{H}$  of connected graphs satisfies (A'1). Then there exists a constant  $c_1 = c_1(\mathcal{H})$  such that  $cc(G) \leq c_1$  for every connected  $\mathcal{H}$ free graph G. Since  $cc(K_{1,c_1+1}) = c_1 + 1$ ,  $cc(K_{c_1+1}^*) = c_1 + 1$  and  $cc(P_{2c_1+1}) =$  $\lceil \frac{2c_1+1}{2} \rceil = c_1 + 1$ , none of  $K_{1,c_1+1}, K_{c_1+1}^*$  and  $P_{2c_1+1}$  is  $\mathcal{H}$ -free. This implies that  $\mathcal{H} \leq \{K_{1,c_1+1}, K_{c_1+1}^*, P_{2c_1+1}\}$ , and hence  $\mathcal{H} \leq \{K_{1,2c_1+1}, K_{2c_1+1}^*, P_{2c_1+1}\}$ , which leads (iii).  $\Box$ 

### Acknowledgment

This work was partially supported by JSPS KAKENHI Grant number JP20K03720 (to S.C) and JSPS KAKENHI Grant number JP18K13449 (to M.F).

## References

- [1] I. Choi, M. Furuya, R. Kim and B. Park, A Ramsey-type theorem for the matching number regarding connected graphs, Discrete Math. **343** (2020), 111648.
- [2] R. Diestel, "Graph Theory" (5th edition), Graduate Texts in Mathematics 173, Springer, Berlin (2017).
- [3] D. Duffus, R.J. Gould and M.S. Jacobson, Forbidden subgraphs and the hamiltonian theme, The Theory and Applications of Graphs, pp. 297–316, Wiley, New York (1981).
- [4] H. Enomoto and H. Li, Partition of a graph into cycles and degenerated cycles, Discrete Math. 276 (2004), 177–181.
- [5] R.J. Faudree and R.J. Gould, Characterizing forbidden pairs for Hamiltonian properties, Discrete Math. 173 (1997), 45–60.
- [6] S. Fujita, Degree conditions for the partition of a graph into cycles, edges and isolated vertices, Discrete Math. 309 (2009), 3534–3540.
- S. Fujita, K. Kawarabayashi, C. L. Lucchesi, K. Ota, M. Plummer and A. Saito, A pair of forbidden subgraphs and perfect matchings, J. Combin. Theory Ser. B 96 (2006), 315–324.
- [8] M. Furuya, Forbidden subgraphs for constant domination number, Discrete Math. Theor. Comput. Sci. 20(1) (2018), Paper No. 19.
- [9] T. Gallai and A.N. Milgram, Verallgemeinerung eines graphentheoretischen Satzes von Redei, Acta Sci. Math. 21 (1960), 181–186.
- [10] R.J. Gould and J.M. Harris, Forbidden triples of subgraphs and traceability, Congr. Numer. 108 (1995), 183–192.
- [11] R.J. Gould and J.M. Harris, Traceability in graphs with forbidden triples of subgraphs, Discrete Math. 189 (1998), 123–132.
- [12] R.J. Gould and J.M. Harris, Forbidden triples and traceability: a characterization, Discrete Math. 203 (1999), 101–120.

- [13] J. Han, On vertex-disjoint paths in regular graphs, Electron. J. Combin. 25 (2018), Paper No. 2.12.
- [14] S. Ishizuka, Closure, path-factors and path coverings in claw-free graphs, Ars Combin. 50 (1998), 115–128.
- [15] C. Magnant and D.M. Martin, A note on the path cover number of regular graphs, Australas. J. Combin. 43 (2009), 211–217.
- [16] B. Reed, Paths, stars and the number three, Combin. Probab. Comput. 5 (1996), 277–295.
- [17] G. Yu, Covering 2-connected 3-regular graphs with disjoint paths, J. Graph Theory 88 (2018), 385–401.