

A new group of doubly charmed molecule with T -doublet charmed meson pair

Fu-Lai Wang^{1,2,*}, Rui Chen^{3,†} and Xiang Liu^{1,2,4,‡§}

¹*School of Physical Science and Technology, Lanzhou University, Lanzhou 730000, China*

²*Research Center for Hadron and CSR Physics, Lanzhou University and Institute of Modern Physics of CAS, Lanzhou 730000, China*

³*Key Laboratory of Low-Dimensional Quantum Structures and Quantum Control of Ministry of Education, Department of Physics and Synergetic Innovation Center for Quantum Effectss and Applications, Hunan Normal University, Changsha 410081, China*

⁴*Lanzhou Center for Theoretical Physics, Key Laboratory of Theoretical Physics of Gansu Province, and Frontiers Science Center for Rare Isotopes, Lanzhou University, Lanzhou 730000, China*

Inspired by the observation of the doubly charmed tetraquark state T_{cc}^+ at the LHCb Collaboration, we systematically study the S -wave interactions between a pair of charmed mesons (D_1 , D_2^*) in the T -doublet, where we adopt the one-boson-exchange model and consider both the S - D wave mixing and coupled channel effects. Our numerical results suggest that the prime doubly charmed molecular tetraquark candidates are the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$, the S -wave $D_1 D_2^*$ states with $I(J^P) = 0(1^+, 2^+, 3^+)$, and the S -wave $D_2^* D_2^*$ states with $I(J^P) = 0(1^+, 3^+)$. The S -wave $D_1 D_1$ state with $I(J^P) = 1(2^+)$, the S -wave $D_1 D_2^*$ state with $I(J^P) = 1(3^+)$, and the S -wave $D_2^* D_2^*$ state with $I(J^P) = 1(4^+)$ may be the possible isovector doubly charmed molecular tetraquark candidates. We expect the experiments like the LHCb or Belle II collaborations to search for these predicted doubly charmed molecular tetraquarks near the $D_1 D_1$, $D_1 D_2^*$, and $D_2^* D_2^*$ thresholds in the future.

I. INTRODUCTION

Very recent, the LHCb Collaboration reported the first observation of the doubly charmed tetraquark state T_{cc}^+ in the $D^0 D^0 \pi^+$ invariant mass spectrum [1]. Obviously, it is beyond the conventional hadron since the T_{cc}^+ has minimal quark component $cc\bar{u}\bar{d}$. In Ref. [2], the pole parameters $\hat{s} = m_{\text{pole}} - i\Gamma_{\text{pole}}/2$ for the T_{cc}^+ state related to the $D^0 D^{*+}$ mass threshold are

$$\begin{aligned}\delta m_{\text{pole}} &= -360 \pm 40_{-0}^{+4} \text{ keV}/c^2, \\ \Gamma_{\text{pole}} &= 48 \pm 2_{-14}^{+0} \text{ keV}.\end{aligned}$$

Additionally, its spin-parity quantum number is $J^P = 1^+$. After the observation of the T_{cc}^+ state, a series of theoretical studies tried to understand its inner structures and properties within the doubly charmed molecular tetraquark [3–20] and compact doubly charmed tetraquark pictures [21–24]. In fact, before the above LHCb experiment, several theoretical groups predicted the existence of the DD^* doubly charmed molecular tetraquark state [25–36] and compact doubly charmed tetraquark states [37–65], and suggested experimentalists to carry out the search for them.

The importance of the observed T_{cc} is equal to that of the $X(3872)$. In 2003, the $X(3872)$ discovered by the Belle Collaboration [66] inspired extensive discussion of assigning it as the $D\bar{D}^*$ charmoniumlike molecular state with $I(J^{PC}) = 0(1^{++})$ [67–73]. Obviously, it is a start point of exploring hidden-charm molecular tetraquark state since more hidden-charm molecular tetraquark explanations to the charmoniumlike $X/Y/Z$ states have been proposed [74–80], which also

stimulated the study of the hidden-charm molecular pentaquarks [81–86] and the observation of the P_c states [87, 88] (see Table I for a concise review).

TABLE I: A summary of experimental observations and the corresponding hidden-charm and doubly charmed molecular tetraquark explanations.

Hidden-charm molecular tetraquark systems	
$H\bar{H}$ [67–73, 89–102]	$X(3872)$ [66], $Z_c(3900)$ [103], $X(3915)$ [104], $Z(3930)$ [105], $Y(3940)$ [106], $Y(4008)$ [107], $Z_c(4020)$ [108], $Z_c(4025)$ [109], $Z^+(4051)$ [110]
$H\bar{T}$ [99, 111–116]	$Z^+(4248)$ [110], $Y(4260)$ [117], $Y(4360)$ [118], $Y(4390)$ [119], $Z^+(4430)$ [120]
$H\bar{S}$ [121–123]	$Z^+(4248)$ [110], $Z^+(4430)$ [120]
$T\bar{T}$ [99, 124]	
$S\bar{S}$ [125]	
Doubly charmed molecular tetraquark systems	
HH [3–18, 25–36]	T_{cc}^+ [1]
HT [9, 126]	

As we can see, there are several common features between the $X(3872)$ and T_{cc} states. Their masses are very close to the threshold of the DD^* pair, and the long-range interaction for the DD^* state with $I(J^{PC}) = 0(1^{++})$ is exactly same as the DD^* state with $I(J^P) = 0(1^+)$ [3]. Inspired by the $X(3872)$, a zoo of the hidden-charm molecular tetraquarks has been formed [89]. Along this line, the T_{cc}^+ observation must initiate a parallel zoo of the doubly charmed molecular tetraquarks. Searching for the doubly charmed molecular tetraquarks not only deepens our understanding to the interactions between charmed meson pair, but also may provide some hints to shed light on the interactions between charmed meson and anti-charmed meson.

*Corresponding author

*Electronic address: wangfl2016@lzu.edu.cn

†Electronic address: chenrui@hunnu.edu.cn

§Electronic address: xiangliu@lzu.edu.cn

In the heavy quark spin symmetry [127], the charmed mesons with the same light quark spin-parity quantum number j_l^P can be categorized into a doublet, such as $H[j_l^P = 1/2^-] = (D, D^*)$, $S[j_l^P = 1/2^+] = (D_0^*, D_1')$, and $T[j_l^P = 3/2^+] = (D_1, D_2^*)$. Here, we need to mention that the D_1 and D_2^* states are abbreviation for the $D_1(2420)$ and $D_2^*(2460)$ states in the Particle Data Group (PDG), respectively. In Refs. [25–36], the authors predicted the possible doubly charmed molecular tetraquark candidates composed of a pair of charmed mesons in the H -doublet. Recently, we performed a systematic study of the HT -type meson-meson interactions, and predicted a series of possible HT -type doubly charmed molecular tetraquark candidates [126].

The newly reported doubly charmed tetraquark state T_{cc} makes the study of the interactions between two charmed mesons an intriguing and important research issue [1, 2], we expect that there exists a zoo of the doubly charmed molecular tetraquarks in the hadron spectroscopy. In fact, the present research status of the doubly charmed molecular tetraquark states is similar to that of the hidden-charm molecular tetraquark states around 2003 [75], the theorists should pay more attention to making the reliable prediction of various types of the doubly charmed molecular tetraquark states and give more abundant suggestions to searching for the doubly charmed molecular tetraquarks accessible at the forthcoming experiment, by which we hope that the crucial information can encourage experimental colleagues to focus on the doubly charmed molecular tetraquark states. For continuing the exploration of doubly charmed molecular tetraquark family, in this work, we focus on the S -wave interactions between a pair of charmed mesons in the T -doublet by adopting the one-boson-exchange (OBE) model. In realistic calculation, both the S - D wave mixing effect and the coupled channel effect are considered. With these obtained OBE effective potentials, we can further solve the coupled channel Schrödinger equation to search for the bound state solutions. By this way, we study how charmed and anti-charmed mesons in the T doublet form the TT -type doubly charmed molecular tetraquark states. Finally, several possible TT -type doubly charmed molecular tetraquark candidates are predicted, which is the important step to construct the family of the doubly charmed molecular tetraquarks, where the provided crucial information is helpful to the experimental search for the possible doubly charmed molecular tetraquark candidates in the future experiment.

This paper is organized as follows. After the introduction in Sec. I, we illustrate the detailed deduction of the S -wave interactions between a pair of charmed mesons in the T -doublet by adopting the OBE model in Sec. II. With this preparation, we present the corresponding bound state properties of the S -wave $D_1 D_1$, $D_1 D_2^*$, and $D_2^* D_2^*$ systems, and further conclude whether there exist possible TT -type doubly charmed molecular tetraquark candidates in Sec. III. This work ends with the summary in Sec. IV.

II. INTERACTIONS OF THE S -WAVE TT SYSTEMS

In this section, we present the detailed derivation of the OBE effective potentials for these discussed S -wave TT -type doubly charmed tetraquark systems, and we include the contributions from the light scalar meson (σ), pseudoscalar meson (π/η), and vector meson (ρ/ω) exchange processes. In this work, we treat the D_1 and D_2^* states as the P -wave charmed mesons to study the doubly charmed molecular tetraquark candidates [128]. Here, we need to point out that the D_1 and D_2^* states as the hadronic molecular states were also studied in the past decades [80, 129–132].

A. The flavor and spin-orbital wave functions

Before deducing the effective potentials, we first construct the flavor and spin-orbital wave functions for these discussed TT -type doubly charmed tetraquark systems. The isospin quantum numbers for the TT -type doubly charmed tetraquark systems can be either 0 or 1, and we collect the relevant flavor wave functions $|I, I_3\rangle$ for the TT systems in Table II [4, 32, 35, 126].

TABLE II: Flavor wave functions $|I, I_3\rangle$ for the TT systems. Here, I and I_3 stand for their isospin and the third component of these discussed TT systems, respectively. For the S -wave $D_1 D_2^*$ system, the factor $(-1)^{J_{D_1 D_2^*}-3}$ comes from the exchange of two charmed mesons [112, 116, 124, 126, 133].

$ I, I_3\rangle$	$D_1 D_1$ system	$D_2^* D_2^*$ system
$ 1, 1\rangle$	$ D_1^+ D_1^+\rangle$	$ D_2^{*+} D_2^{*+}\rangle$
$ 1, 0\rangle$	$\frac{1}{\sqrt{2}}(D_1^0 D_1^+\rangle + D_1^+ D_1^0\rangle)$	$\frac{1}{\sqrt{2}}(D_2^{*0} D_2^{*+}\rangle + D_2^{*+} D_2^{*0}\rangle)$
$ 1, -1\rangle$	$ D_1^0 D_1^0\rangle$	$ D_2^{*0} D_2^{*0}\rangle$
$ 0, 0\rangle$	$\frac{1}{\sqrt{2}}(D_1^0 D_1^+\rangle - D_1^+ D_1^0\rangle)$	$\frac{1}{\sqrt{2}}(D_2^{*0} D_2^{*+}\rangle - D_2^{*+} D_2^{*0}\rangle)$
$ I, I_3\rangle$	$D_1 D_2^*$ system	
$ 1, 1\rangle$	$\frac{1}{\sqrt{2}}(D_1^+ D_2^{*+}\rangle + (-1)^{J_{D_1 D_2^*}-3} D_2^{*+} D_1^+\rangle)$	
$ 1, 0\rangle$	$\frac{1}{2}[(D_1^0 D_2^{*+}\rangle + D_1^+ D_2^{*0}\rangle) + (-1)^{J_{D_1 D_2^*}-3} (D_2^{*+} D_1^0\rangle + D_2^{*0} D_1^+\rangle)]$	
$ 1, -1\rangle$	$\frac{1}{\sqrt{2}}(D_1^0 D_2^{*0}\rangle + (-1)^{J_{D_1 D_2^*}-3} D_2^{*0} D_1^0\rangle)$	
$ 0, 0\rangle$	$\frac{1}{2}[(D_1^0 D_2^{*+}\rangle - D_1^+ D_2^{*0}\rangle) - (-1)^{J_{D_1 D_2^*}-3} (D_2^{*+} D_1^0\rangle - D_2^{*0} D_1^+\rangle)]$	

These allowed quantum numbers for the S -wave $D_1 D_1$ and $D_2^* D_2^*$ states include

$$D_1 D_1[I(J^P)] : 1(0^+), 0(1^+), 1(2^+), \\ D_2^* D_2^*[I(J^P)] : 1(0^+), 0(1^+), 1(2^+), 0(3^+), 1(4^+).$$

The spin-orbital wave functions $|^{2S+1}L_J\rangle$ for the $D_1 D_1$, $D_1 D_2^*$,

and $D_2^* D_2^*$ systems can be expressed as

$$\begin{aligned} D_1 D_1 : |^{2S+1} L_J\rangle &= \sum_{m,m',m_S,m_L} C_{1m,1m'}^{S,m_S} C_{S m_S, L m_L}^{J,M} \epsilon_m^\mu \epsilon_{m'}^\nu |Y_{L,m_L}\rangle, \\ D_1 D_2^* : |^{2S+1} L_J\rangle &= \sum_{m,m',m_S,m_L} C_{1m,2m'}^{S,m_S} C_{S m_S, L m_L}^{J,M} \epsilon_m^\mu \epsilon_{m'}^{\lambda\mu\nu} |Y_{L,m_L}\rangle, \\ D_2^* D_2^* : |^{2S+1} L_J\rangle &= \sum_{m,m',m_S,m_L} C_{2m,2m'}^{S,m_S} C_{S m_S, L m_L}^{J,M} \zeta_m^{\mu\nu} \zeta_{m'}^{\lambda\rho} |Y_{L,m_L}\rangle, \end{aligned}$$

respectively. Here, $C_{ab,cd}^{e,f}$ is the Clebsch-Gordan coefficient, and $|Y_{L,m_L}\rangle$ stands for the spherical harmonics function. ϵ_m^μ ($m = 0, \pm 1$) and $\zeta_m^{\mu\nu}$ ($m = 0, \pm 1, \pm 2$) denote the polarization vector and tensor for the axial-vector charmed meson D_1 and the tensor charmed meson D_2^* , respectively. In the static limit, their explicit expressions can be written as [134]

$$\begin{aligned} \epsilon_0^\mu &= (0, 0, 0, -1), \quad \epsilon_\pm^\mu = \frac{1}{\sqrt{2}} (0, \pm 1, i, 0), \\ \zeta_m^{\mu\nu} &= \sum_{m_1, m_2} C_{1m_1, 1m_2}^{2,m} \epsilon_{m_1}^\mu \epsilon_{m_2}^\nu. \end{aligned}$$

B. The effective Lagrangians

In the following, we adopt the effective Lagrangian approach to deduce the OBE effective potentials for the $D_1 D_1$, $D_1 D_2^*$, and $D_2^* D_2^*$ systems. Based on the heavy quark symmetry, the chiral symmetry, and the hidden gauge symmetry [135–139], the compact form of the relevant effective Lagrangians can be constructed as [140]

$$\mathcal{L}_{TT\sigma} = g_\sigma'' \langle T_a^{(Q)\mu} \sigma \bar{T}_{a\mu}^{(Q)} \rangle, \quad (2.1)$$

$$\mathcal{L}_{TT\mathbb{P}} = ik \langle T_b^{(Q)\mu} \mathcal{A}_{ba} \gamma_5 \bar{T}_{a\mu}^{(Q)} \rangle, \quad (2.2)$$

$$\begin{aligned} \mathcal{L}_{TT\mathbb{V}} &= i\beta'' \langle T_b^{(Q)\lambda} v^\mu (\mathcal{V}_\mu - \rho_\mu)_{ba} \bar{T}_{a\lambda}^{(Q)} \rangle \\ &\quad + i\lambda'' \langle T_b^{(Q)\lambda} \sigma^{\mu\nu} F_{\mu\nu}(\rho)_{ba} \bar{T}_{a\lambda}^{(Q)} \rangle, \end{aligned} \quad (2.3)$$

where the four velocity is $v = (1, 0)$ in the static approximation. Surperfield $T_a^{(Q)\mu}$ can be expressed as a linear combination of the axial-vector heavy flavor meson $P_1^{(Q)}$ with $I(J^P) = 1/2(1^+)$ and the tensor heavy flavor meson $P_2^{(Q)}$ with $I(J^P) = 1/2(2^+)$, which reads as [140]

$$T_a^{(Q)\mu} = \frac{1 + \not{v}}{2} \left[P_{2a}^{*(Q)\mu\nu} \gamma_\nu - \sqrt{\frac{3}{2}} P_{1a\nu}^{(Q)} \gamma_5 \left(g^{\mu\nu} - \frac{\gamma^\nu (\gamma^\mu - v^\mu)}{3} \right) \right],$$

where $P_1^{(c)} = (D_1^0, D_1^+)$, $P_2^{*(c)} = (D_2^{*0}, D_2^{*+})$, and the normalization relations for these discussed charmed mesons are $\langle 0 | D_1^+ | c \bar{q}(1^+) \rangle = \epsilon^\mu \sqrt{m_{D_1}}$ and $\langle 0 | D_2^{*+} | c \bar{q}(2^+) \rangle = \xi^{\mu\nu} \sqrt{m_{D_2^*}}$. Its conjugate field $\bar{T}_a^{(Q)\mu}$ is written as $\bar{T}_a^{(Q)\mu} = \gamma^0 T_a^{(Q)\mu\dagger} \gamma^0$.

In Eqs. (2.1)–(2.3), we define the axial current \mathcal{A}_μ , the vector current \mathcal{V}_μ , the vector meson field ρ_μ , and the vector meson field strength tensor $F_{\mu\nu}(\rho)$, i.e.,

$$\begin{aligned} \mathcal{A}_\mu &= \frac{1}{2} (\xi^\dagger \partial_\mu \xi - \xi \partial_\mu \xi^\dagger), \quad \mathcal{V}_\mu = \frac{1}{2} (\xi^\dagger \partial_\mu \xi + \xi \partial_\mu \xi^\dagger), \\ \rho_\mu &= \frac{ig_V}{\sqrt{2}} \mathbb{V}_\mu, \quad F_{\mu\nu}(\rho) = \partial_\mu \rho_\nu - \partial_\nu \rho_\mu + [\rho_\mu, \rho_\nu], \end{aligned}$$

with $\xi = \exp(i\mathbb{P}/f_\pi)$. Here, \mathbb{P} and \mathbb{V}_μ stand for the pseudoscalar meson and vector meson matrices, respectively, which have the forms of

$$\begin{aligned} \mathbb{P} &= \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\sqrt{\frac{2}{3}}\eta \end{pmatrix}, \\ \mathbb{V}_\mu &= \begin{pmatrix} \frac{\rho^0}{\sqrt{2}} + \frac{\omega}{\sqrt{2}} & \rho^+ & K^{*+} \\ \rho^- & -\frac{\rho^0}{\sqrt{2}} + \frac{\omega}{\sqrt{2}} & K^{*0} \\ K^{*-} & \bar{K}^{*0} & \phi \end{pmatrix}_\mu, \end{aligned}$$

respectively. Once expanding the effective Lagrangians in Eqs. (2.1)–(2.3), we can further obtain the concrete effective Lagrangians between the charmed mesons in the T -doublet and the light mesons, which can be explicitly expressed as

$$\mathcal{L}_{D_1 D_1 \sigma} = -2g_\sigma'' D_{1a\mu} D_{1a}^{\mu\dagger} \sigma, \quad (2.4)$$

$$\mathcal{L}_{D_1 D_1 \mathbb{P}} = -\frac{5ik}{3f_\pi} \epsilon^{\mu\nu\rho\tau} v_\tau D_{1b\nu} D_{1a\mu}^\dagger \partial_\rho \mathbb{P}_{ba}, \quad (2.5)$$

$$\begin{aligned} \mathcal{L}_{D_1 D_1 \mathbb{V}} &= -\sqrt{2}\beta'' g_V (v \cdot \mathbb{V}_{ba}) D_{1b\mu} D_{1a}^{\mu\dagger} \\ &\quad + \frac{5\sqrt{2}i\lambda'' g_V}{3} (D_{1b}^\mu D_{1a}^{\nu\dagger} - D_{1b}^\nu D_{1a}^{\mu\dagger}) \partial_\mu \mathbb{V}_{bav}, \end{aligned} \quad (2.6)$$

$$\mathcal{L}_{D_2^* D_2^* \sigma} = 2g_\sigma'' D_{2a}^{*\mu\nu} D_{2a\mu\nu}^{\dagger} \sigma, \quad (2.7)$$

$$\mathcal{L}_{D_2^* D_2^* \mathbb{P}} = \frac{2ik}{f_\pi} \epsilon^{\mu\nu\rho\tau} v_\nu D_{2b\alpha\tau}^* D_{2a\rho}^{*\dagger} \partial_\mu \mathbb{P}_{ba}, \quad (2.8)$$

$$\begin{aligned} \mathcal{L}_{D_2^* D_2^* \mathbb{V}} &= \sqrt{2}\beta'' g_V (v \cdot \mathbb{V}_{ba}) D_{2b}^{*\lambda\nu} D_{2a\lambda\nu}^{\dagger} + 2\sqrt{2}i\lambda'' g_V \\ &\quad \times (D_{2b}^{*\lambda\nu} D_{2a\lambda}^{\mu\dagger} - D_{2b\lambda}^{*\mu} D_{2a}^{\lambda\nu\dagger}) \partial_\mu \mathbb{V}_{bav}, \end{aligned} \quad (2.9)$$

$$\mathcal{L}_{D_1 D_2^* \sigma} = \sqrt{\frac{2}{3}} ig_\sigma'' \epsilon^{\mu\nu\rho\tau} v_\rho (D_{2a\mu\tau}^* D_{1a\nu}^\dagger - D_{1a\nu} D_{2a\mu\tau}^{\dagger}) \sigma, \quad (2.10)$$

$$\mathcal{L}_{D_1 D_2^* \mathbb{P}} = -\sqrt{\frac{2}{3}} \frac{k}{f_\pi} (D_{2b}^{*\mu\lambda} D_{1a\mu}^\dagger + D_{1b\mu} D_{2a}^{*\mu\lambda\dagger}) \partial_\lambda \mathbb{P}_{ba}, \quad (2.11)$$

$$\begin{aligned} \mathcal{L}_{D_1 D_2^* \mathbb{V}} &= \frac{i\beta'' g_V}{\sqrt{3}} \epsilon^{\lambda\alpha\rho\tau} v_\rho (v \cdot \mathbb{V}_{ba}) (D_{2b\lambda\tau}^* D_{1a\alpha}^\dagger - D_{1b\alpha} D_{2a\lambda\tau}^{\dagger}) \\ &\quad + \frac{2\lambda'' g_V}{\sqrt{3}} [3\epsilon^{\mu\lambda\nu\tau} v_\lambda (D_{2b\alpha\tau}^* D_{1a}^{\alpha\dagger} + D_{1b}^\alpha D_{2a\alpha\tau}^{\dagger}) \partial_\mu \mathbb{V}_{bav} \\ &\quad + 2\epsilon^{\lambda\alpha\rho\nu} v_\rho (D_{2b\lambda}^{*\mu} D_{1a\alpha}^\dagger + D_{1b\alpha} D_{2a\lambda}^{\mu\dagger}) \\ &\quad \times (\partial_\mu \mathbb{V}_{bav} - \partial_\nu \mathbb{V}_{ba\mu})]. \end{aligned} \quad (2.12)$$

In this work, we estimate all the coupling constants in the quark model (see Refs. [141–144] for more details). The values for the involved coupling constants are $g_\sigma'' = 0.76$, $k = 0.59$, $f_\pi = 132$ MeV, $\beta'' = 0.90$, $\lambda'' = 0.56$ GeV $^{-1}$, and $g_V = 5.83$ [141]. The masses of these involved hadrons are $m_\sigma = 600.00$ MeV, $m_\pi = 137.27$ MeV, $m_\eta = 547.86$ MeV, $m_\rho = 775.26$ MeV, $m_\omega = 782.66$ MeV, $m_{D_1} = 2422.00$ MeV, and $m_{D_2^*} = 2463.05$ MeV [145]. As the important input parameters within the OBE model, the information about the coupling constants is crucial when studying the existence pos-

sibility of the hadronic molecular states. However, the coupling constants with the charmed meson in T -doublet cannot be fixed based on the experimental data at present, and we estimate these coupling constants by the quark model in this work [142], which is often adopted to determine the coupling constants. Because of the uncertainties of the coupling constants, we should be cautious when studying the existence possibility of the hadronic molecular states. Here, we need to indicate our results of the existence possibility of the hadronic molecular states at the qualitative level, but the uncertainties of the coupling constants do not significantly affect our qualitative conclusions.

C. The effective potentials

Based on the above preparation, we can further deduce the effective potentials to judge whether these discussed doubly-charmed molecular tetraquark states exist or not. As indicated in Ref. [141], there exists the standard strategy to judge the possibility of the existence of the investigated hadronic molecular state within the OBE model, which is given in Fig. 1.




FIG. 1: (color online) The standard strategy to judge the possibility of the existence of the investigated hadronic molecular state within the OBE model.

For extracting the effective potential in the coordinate space, which is shown in the upper part of the Fig. 1, we will give a brief introduction in the following. First, we can write down the scattering amplitude $\mathcal{M}^{h_1 h_2 \rightarrow h_3 h_4}(q)$ of the scattering process $h_1 h_2 \rightarrow h_3 h_4$ by exchanging the light mesons under the effective Lagrangian approach. The relevant Feynman diagram is given in Fig. 2, and the scattering amplitude $\mathcal{M}^{h_1 h_2 \rightarrow h_3 h_4}(q)$ can be written as

$$i\mathcal{M}^{h_1 h_2 \rightarrow h_3 h_4}(q) = \sum_{m=\sigma, \pi, \eta} i\Gamma_{(\mu)}^{h_1 h_3 m} P_m^{(\mu\nu)} i\Gamma_{(\nu)}^{h_2 h_4 m}, \quad (2.13)$$

where the interaction vertices $\Gamma_{(\mu)}^{h_1 h_3 m}$ and $\Gamma_{(\nu)}^{h_2 h_4 m}$ can be extracted from the former effective Lagrangians.

And then, we adopt the Breit approximation [146, 147] and the nonrelativistic normalizations to obtain the relations between the effective potential in the momentum space $\mathcal{V}_E^{h_1 h_2 \rightarrow h_3 h_4}(q)$ and the corresponding scattering amplitude




FIG. 2: The relevant Feynman diagram for the scattering process $h_1 h_2 \rightarrow h_3 h_4$.

$\mathcal{M}^{h_1 h_2 \rightarrow h_3 h_4}(q)$, i.e.,

$$\mathcal{V}_E^{h_1 h_2 \rightarrow h_3 h_4}(q) = -\frac{\mathcal{M}^{h_1 h_2 \rightarrow h_3 h_4}(q)}{\sqrt{\prod_i 2m_i} \sqrt{\prod_f 2m_f}}, \quad (2.14)$$

where $m_{i(f)}$ are the masses of the initial (final) states. Finally, we can perform the Fourier transformation to deduce the effective potential in the coordinate space $\mathcal{V}_E^{h_1 h_2 \rightarrow h_3 h_4}(r)$, i.e.,

$$\mathcal{V}_E^{h_1 h_2 \rightarrow h_3 h_4}(r) = \int \frac{d^3 q}{(2\pi)^3} e^{i\mathbf{q} \cdot \mathbf{r}} \mathcal{V}_E^{h_1 h_2 \rightarrow h_3 h_4}(q) \mathcal{F}^2(q^2, m_E^2). \quad (2.15)$$

In order to compensate the off-shell effect of the exchanged particles, we introduce the form factor $\mathcal{F}(q^2, m_E^2)$ at every interaction vertex, $\mathcal{F}(q^2, m_E^2) = (\Lambda^2 - m_E^2)/(\Lambda^2 - q^2)$. Here, Λ , q , and m_E are the cutoff parameter, the four-momentum, and the mass of the exchanged light mesons, respectively. According to the experience of studying the deuteron, a reasonable cutoff value is taken around 1.00 GeV [28, 148], in this cutoff region, we can reproduce the masses of the three P_c states [81–86] and the T_{cc} state [3–9, 24, 32, 33, 35, 36] in the hadronic molecular picture.

After getting the effective potentials in the coordinate space for the discussed systems by the above three typical steps, we can search for the bound state solutions by numerically solving the coupled channel Schrödinger equation,

$$-\frac{1}{2\mu} \left(\nabla^2 - \frac{\ell(\ell+1)}{r^2} \right) \psi(r) + V(r)\psi(r) = E\psi(r), \quad (2.16)$$

where $\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r}$, and $\mu = \frac{m_1 m_2}{m_1 + m_2}$ is the reduced mass for the discussed systems. In our calculation, the obtained bound state solutions mainly include the binding energy E and the radial wave function $\psi(r)$, with which we can further calculate the root-mean-square radius r_{RMS} and the probability of the individual channel P_i .

The total OBE effective potentials for all the discussed TT -type doubly charmed tetraquarks can be expressed as

$$\mathcal{V}_1 = -g_{\sigma}''^2 O_1 Y_{\sigma} - \frac{25k^2}{108f_{\pi}^2} (O_2 \mathcal{Z}_r + O_3 \mathcal{T}_r) \left(\frac{GY_{\pi}}{2} + \frac{Y_{\eta}}{6} \right)$$

$$\begin{aligned}
& + \frac{1}{2} \beta'^2 g_V^2 O_1 \left(\frac{\mathcal{G}Y_\rho}{2} + \frac{Y_\omega}{2} \right) \\
& - \frac{25}{54} \lambda'^2 g_V^2 (2O_2 \mathcal{Z}_r - O_3 \mathcal{T}_r) \left(\frac{\mathcal{G}Y_\rho}{2} + \frac{Y_\omega}{2} \right) \quad (2.17)
\end{aligned}$$

for the $D_1 D_1 \rightarrow D_1 D_1$ process,

$$\begin{aligned}
\mathcal{V}_2 = & -g_\sigma'^2 \frac{O_4 + O'_4}{2} Y_\sigma \\
& - \frac{5k^2}{18f_\pi^2} \left(\frac{O_5 + O'_5}{2} \mathcal{Z}_r + \frac{O_6 + O'_6}{2} \mathcal{T}_r \right) \left(\frac{\mathcal{G}Y_\pi}{2} + \frac{Y_\eta}{6} \right) \\
& + \frac{1}{2} \beta'^2 g_V^2 \frac{O_4 + O'_4}{2} \left(\frac{\mathcal{G}}{2} Y_\rho + \frac{1}{2} Y_\omega \right) \\
& - \frac{2}{3} \lambda'^2 g_V^2 \left(2 \frac{O_5 + O'_5}{2} \mathcal{Z}_r - \frac{O_6 + O'_6}{2} \mathcal{T}_r \right) \left(\frac{\mathcal{G}Y_\rho}{2} + \frac{Y_\omega}{2} \right) \quad (2.18)
\end{aligned}$$

for the $D_1 D_2^* \rightarrow D_1 D_2^*$ process,

$$\begin{aligned}
\mathcal{V}_3 = & \frac{k^2}{18f_\pi^2} \left(\frac{O_7 + O'_7}{2} \mathcal{Z}_r + \frac{O_8 + O'_8}{2} \mathcal{T}_r \right) \left(\frac{\mathcal{G}Y_{\pi 0}}{2} + \frac{Y_{\eta 0}}{6} \right) \\
& + \frac{\lambda'^2 g_V^2}{9} \left(2 \frac{O_7 + O'_7}{2} \mathcal{Z}_r - \frac{O_8 + O'_8}{2} \mathcal{T}_r \right) \left(\frac{\mathcal{G}Y_{\rho 0}}{2} + \frac{Y_{\omega 0}}{2} \right) \quad (2.19)
\end{aligned}$$

for the $D_1 D_2^* \rightarrow D_2^* D_1$ process, and

$$\begin{aligned}
\mathcal{V}_4 = & -g_\sigma'^2 O_9 Y_\sigma - \frac{k^2}{3f_\pi^2} (O_{10} \mathcal{Z}_r + O_{11} \mathcal{T}_r) \left(\frac{\mathcal{G}Y_\pi}{2} + \frac{Y_\eta}{6} \right) \\
& + \frac{1}{2} \beta'^2 g_V^2 O_9 \left(\frac{\mathcal{G}Y_\rho}{2} + \frac{Y_\omega}{2} \right) \\
& - \frac{2}{3} \lambda'^2 g_V^2 (2O_{10} \mathcal{Z}_r - O_{11} \mathcal{T}_r) \left(\frac{\mathcal{G}Y_\rho}{2} + \frac{Y_\omega}{2} \right) \quad (2.20)
\end{aligned}$$

for the $D_2^* D_2^* \rightarrow D_2^* D_2^*$ process. Here, the constant \mathcal{G} is taken as -3 for the isoscalar system and 1 for the isovector system, respectively. For the convenience, we define the following expressions

$$\begin{aligned}
\mathcal{Z}_r &= \frac{1}{r^2} \frac{\partial}{\partial r} r^2 \frac{\partial}{\partial r}, \quad \mathcal{T}_r = r \frac{\partial}{\partial r} \frac{1}{r} \frac{\partial}{\partial r}, \\
Y_i &= \frac{e^{-m_i r} - e^{-\Lambda_i r}}{4\pi r} - \frac{\Lambda_i^2 - m_i^2}{8\pi \Lambda_i} e^{-\Lambda_i r}. \quad (2.21)
\end{aligned}$$

Here, $m_0 = \sqrt{m^2 - q_0^2}$ and $\Lambda_0 = \sqrt{\Lambda^2 - q_0^2}$ with $q_0 = m_{D_2^*} - m_{D_1}$.

In the above OBE effective potentials, we define several relevant operators $O_k^{(\prime)}$, which include

$$\begin{aligned}
O_1 &= (\epsilon_3^\dagger \cdot \epsilon_1) (\epsilon_4^\dagger \cdot \epsilon_2), \quad O_2 = (\epsilon_3^\dagger \times \epsilon_1) \cdot (\epsilon_4^\dagger \times \epsilon_2), \\
O_3 &= S(\epsilon_3^\dagger \times \epsilon_1, \epsilon_4^\dagger \times \epsilon_2, \hat{r}), \\
O_4 &= \mathcal{A}(\epsilon_3^\dagger \cdot \epsilon_1) (\epsilon_{4a}^\dagger \cdot \epsilon_{2c}) (\epsilon_{4b}^\dagger \cdot \epsilon_{2d}), \\
O'_4 &= \mathcal{A}(\epsilon_4^\dagger \cdot \epsilon_2) (\epsilon_{3a}^\dagger \cdot \epsilon_{1c}) (\epsilon_{3b}^\dagger \cdot \epsilon_{1d}),
\end{aligned}$$

$$\begin{aligned}
O_5 &= \mathcal{A}(\epsilon_{4a}^\dagger \cdot \epsilon_{2c}) [(\epsilon_3^\dagger \times \epsilon_1) \cdot (\epsilon_{4b}^\dagger \times \epsilon_{2d})], \\
O'_5 &= \mathcal{A}(\epsilon_{3a}^\dagger \cdot \epsilon_{1c}) [(\epsilon_4^\dagger \times \epsilon_2) \cdot (\epsilon_{3b}^\dagger \times \epsilon_{1d})], \\
O_6 &= \mathcal{A}(\epsilon_{4a}^\dagger \cdot \epsilon_{2c}) S(\epsilon_3^\dagger \times \epsilon_1, \epsilon_{4b}^\dagger \times \epsilon_{2d}, \hat{r}), \\
O'_6 &= \mathcal{A}(\epsilon_{3a}^\dagger \cdot \epsilon_{1c}) S(\epsilon_4^\dagger \times \epsilon_2, \epsilon_{3b}^\dagger \times \epsilon_{1d}, \hat{r}), \\
O_7 &= \mathcal{A}(\epsilon_{3a}^\dagger \cdot \epsilon_1) (\epsilon_4^\dagger \cdot \epsilon_{2c}) (\epsilon_{3b}^\dagger \cdot \epsilon_{2d}), \\
O'_7 &= \mathcal{A}(\epsilon_{4a}^\dagger \cdot \epsilon_2) (\epsilon_3^\dagger \cdot \epsilon_{1c}) (\epsilon_{4b}^\dagger \cdot \epsilon_{1d}), \\
O_8 &= \mathcal{A}(\epsilon_{3a}^\dagger \cdot \epsilon_1) (\epsilon_4^\dagger \cdot \epsilon_{2c}) S(\epsilon_{3b}^\dagger, \epsilon_{2d}, \hat{r}), \\
O'_8 &= \mathcal{A}(\epsilon_{4a}^\dagger \cdot \epsilon_2) (\epsilon_3^\dagger \cdot \epsilon_{1c}) S(\epsilon_{4b}^\dagger, \epsilon_{1d}, \hat{r}), \\
O_9 &= \mathcal{B}(\epsilon_{3a}^\dagger \cdot \epsilon_{1c}) (\epsilon_{3b}^\dagger \cdot \epsilon_{1d}) (\epsilon_{4e}^\dagger \cdot \epsilon_{2g}) (\epsilon_{4f}^\dagger \cdot \epsilon_{2h}), \\
O_{10} &= \mathcal{B}(\epsilon_{3a}^\dagger \cdot \epsilon_{1c}) (\epsilon_{4e}^\dagger \cdot \epsilon_{2g}) [(\epsilon_{3b}^\dagger \times \epsilon_{1d}) \cdot (\epsilon_{4f}^\dagger \times \epsilon_{2h})], \\
O_{11} &= \mathcal{B}(\epsilon_{4e}^\dagger \cdot \epsilon_{1c}) (\epsilon_{4e}^\dagger \cdot \epsilon_{2g}) S(\epsilon_{3b}^\dagger \times \epsilon_{1d}, \epsilon_{4f}^\dagger \times \epsilon_{2h}, \hat{r}) \quad (2.22)
\end{aligned}$$

with $S(\mathbf{x}, \mathbf{y}, \hat{r}) = 3(\hat{r} \cdot \mathbf{x})(\hat{r} \cdot \mathbf{y}) - \mathbf{x} \cdot \mathbf{y}$, $\mathcal{A} = \sum_{a,b}^{c,d} C_{1a,1b}^{2,a+b} C_{1c,1d}^{2,c+d}$, and $\mathcal{B} = \sum_{a,b,c,d}^{e,f,g,h} C_{1a,1b}^{2,a+b} C_{1c,1d}^{2,c+d} C_{1e,1f}^{2,e+f} C_{1g,1h}^{2,g+h}$. The operator matrix elements $\langle f | O_k^{(\prime)} | i \rangle$ are collected in Appendix A.

III. NUMERICAL RESULTS AND DISCUSSIONS

In this section, we solve the coupled channel Schrödinger equations to search for the bound state solutions for the S -wave $D_1 D_1$, $D_1 D_2^*$, and $D_2^* D_2^*$ systems. Our numerical calculations are presented in three cases. First, we perform a single channel analysis with the OBE effective potentials. After that, we further introduce the S - D wave mixing effect and the coupled channel effect, and give the numerical analysis again. With these steps, we can reveal the roles of the S - D wave mixing effect and the coupled channel effect in the formation of the TT -type doubly charmed molecular tetraquark states.

When we consider the S - D wave mixing effect and the coupled channel effect, the relevant spin-orbit wave functions $|^{2S+1}L_J\rangle$ for the $D_1 D_1$, $D_1 D_2^*$, and $D_2^* D_2^*$ systems are

$$\begin{aligned}
(I, J^P) &= (1, 0^+) : D_1 D_1 |^1\mathbb{S}_0\rangle / |^5\mathbb{D}_0\rangle, \quad D_2^* D_2^* |^1\mathbb{S}_0\rangle / |^5\mathbb{D}_0\rangle; \\
(I, J^P) &= (0, 1^+) : D_1 D_1 |^3\mathbb{S}_1\rangle / |^3\mathbb{D}_1\rangle, \quad D_1 D_2^* |^3\mathbb{S}_1\rangle / |^{3,5,7}\mathbb{D}_1\rangle, \\
&\quad D_2^* D_2^* |^3\mathbb{S}_1\rangle / |^{3,7}\mathbb{D}_1\rangle; \\
(I, J^P) &= (1, 1^+) : D_1 D_2^* |^3\mathbb{S}_1\rangle / |^{3,5,7}\mathbb{D}_1\rangle; \\
(I, J^P) &= (0, 2^+) : D_1 D_2^* |^5\mathbb{S}_2\rangle / |^{3,5,7}\mathbb{D}_2\rangle; \\
(I, J^P) &= (1, 2^+) : D_1 D_1 |^5\mathbb{S}_2\rangle / |^{1,5}\mathbb{D}_2\rangle, \quad D_1 D_2^* |^5\mathbb{S}_2\rangle / |^{3,5,7}\mathbb{D}_2\rangle, \\
&\quad D_2^* D_2^* |^5\mathbb{S}_2\rangle / |^{1,5,9}\mathbb{D}_2\rangle; \\
(I, J^P) &= (0, 3^+) : D_1 D_2^* |^7\mathbb{S}_3\rangle / |^{3,5,7}\mathbb{D}_3\rangle, \quad D_2^* D_2^* |^7\mathbb{S}_3\rangle / |^{3,7}\mathbb{D}_3\rangle; \\
(I, J^P) &= (1, 3^+) : D_1 D_2^* |^7\mathbb{S}_3\rangle / |^{3,5,7}\mathbb{D}_3\rangle; \\
(I, J^P) &= (1, 4^+) : D_2^* D_2^* |^9\mathbb{S}_4\rangle / |^{5,9}\mathbb{D}_4\rangle.
\end{aligned}$$

Within the OBE model, the bound state properties are depend on the cutoff values, but the cutoff parameter cannot

be determined exactly without relevant experiment data. According to the experience of studying the deuteron, we believe the cutoff parameter around 1.0 GeV is the reasonable input [28, 148]. Thus, we attempt to find the loosely bound state solutions of the S -wave TT systems by varying the cutoff parameter in the range around 1.0 GeV in this work. In addition, when judging whether the bound state is an ideal hadronic molecular candidate, we expect that the typical root mean square radius should be larger than the size of all the included component hadrons, and the reasonable binding energy is around several MeV to several tens MeV only, which is due to the hadronic molecular state is a loosely bound state [75]. Therefore, we consider the root mean square radius is greater than 0.7 fm and the binding energy is less than 20.0 MeV when presenting the loosely bound state solutions of the S -wave TT systems in the present work. In short, a loosely bound state with the cutoff parameter closed to 1.0 GeV can be suggested as a good hadronic molecular candidate in realistic calculation.

A. The S -wave $D_1 D_1$ system

When we take the cutoff value Λ in the range of $0.8 < \Lambda < 2.5$ GeV, we cannot find the bound state solutions for the S -wave $D_1 D_1$ state with $I(J^P) = 1(0^+)$, even if we include the S - D wave mixing effect and the coupled channel effect.

In Fig. 3, we present the cutoff parameter Λ dependence of the binding energy E , the root-mean-square (RMS) radius r_{RMS} , and the probabilities for different components P_i for the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$ after considering the single channel, the S - D wave mixing effect, and the coupled channel effect. As we can see, for the single $|^3S_1\rangle$ channel analysis, the binding energy appears when taking the cutoff value $\Lambda \sim 1.23$ GeV. With increasing the cutoff value, its binding becomes deeper. When the cutoff value Λ is taken as 1.44 GeV, the binding energy increases to be 18.00 MeV, and the RMS radius decreases to be 0.80 fm. Thus, we suggest that the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$ can be a good doubly charmed molecular tetraquark candidate when we only consider the contribution of the $|^3S_1\rangle$ channel. After considering the S - D wave mixing effect, the bound state solutions are very similar to those in the single channel analysis, but the cutoff value Λ becomes smaller if obtaining the same binding energy. Thus, the S - D wave mixing effect is helpful to form the $D_1 D_1$ bound state with $I(J^P) = 0(1^+)$. Here, we find that the probability for the D -wave component for this bound state is very tiny, which is less than 1%. When we perform the coupled channel analysis from the $D_1 D_1|^3S_1\rangle$, $D_1 D_2|^3S_1\rangle$, and $D_2^* D_2^*|^3S_1\rangle$ interactions, we can obtain the loosely bound state solutions with the cutoff value Λ around 1.12 GeV, which is a little smaller than those in the single channel and the S - D wave mixing analysis. Obviously, the coupled channel effect can help to generate this bound state. When the binding energy is less than a dozen of MeV, the dominant channel is the $D_1 D_1|^3S_1\rangle$ component, followed by the $D_1 D_2|^3S_1\rangle$ and $D_2^* D_2^*|^3S_1\rangle$ channels. As the increasing of the cutoff value, the $D_1 D_2|^3S_1\rangle$ and $D_2^* D_2^*|^3S_1\rangle$ channels be-

come much more important. To summary, the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$ can be a good doubly charmed molecular tetraquark candidate, and the S - D wave mixing effect and the coupled channel effect play a minor role in generating the S -wave $D_1 D_1$ bound state with $I(J^P) = 0(1^+)$.




FIG. 3: (color online) The cutoff parameter Λ dependence of the bound state solutions for the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$ when performing (a) the single channel analysis, (b) the S - D wave mixing analysis, and (c) the coupled channel analysis.

For the S -wave $D_1 D_1$ state with $I(J^P) = 1(2^+)$, there does not exist the bound state solutions with the cutoff parameter around 0.80 to 2.50 GeV after performing both of the single channel analysis and the S - D wave mixing analysis. However, when we take into account the coupled channel effect by including the $D_1 D_1|^5S_2\rangle$, $D_1 D_2^*|^5S_2\rangle$, and $D_2^* D_2^*|^5S_2\rangle$ channels, we can find weakly binding with the cutoff value larger than 2.30 GeV, which is much larger than the cutoff value in the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$. Thus, the S -wave $D_1 D_1$ state with $I(J^P) = 1(2^+)$ may be the possible doubly charmed molecular tetraquark candidate. As shown in Fig. 4, we present the bound state solutions for the S -wave $D_1 D_1$ state with $I(J^P) = 1(2^+)$ after considering the coupled channel effects, where the $D_1 D_1|^5S_2\rangle$ is the dominant channel with the probability over 80%.

Because the hadrons are not point-like particles, we usually introduce the form factor at each interaction vertexes to describe the off-shell effect of the exchanged light mesons and reflect the inner structure effect of the discussed hadrons. As a general rule, the monopole form factor $\mathcal{F}(q^2, m_E^2) = (\Lambda^2 - m_E^2)/(\Lambda^2 - q^2)$ is often adopted to discuss the hadron-hadron interactions within the OBE model, which may reduce the strength of the vector meson exchange interactions com-




FIG. 4: The cutoff parameter Λ dependence of the bound state solutions for the S -wave D_1D_1 state with $I(J^P) = 1(2^+)$ by performing the coupled channel analysis.

pared with the π exchange interaction. However, the contribution of the vector meson exchange interactions may not be suppressed by the factor $(\Lambda^2 - m_E^2)$ [149]. In order to further test our conclusions, we use other kind of form factor $\mathcal{F}(q^2) = \Lambda^2/(\Lambda^2 - q^2)$ to discuss the bound state properties [150], which is not suppress the contribution of the vector meson exchange interactions. Here, we would like to mention that the cutoff values around 1.0 GeV in the form factor $\mathcal{F}(q^2, m_E^2) = (\Lambda^2 - m_E^2)/(\Lambda^2 - q^2)$ and 0.5 GeV in the form factor $\mathcal{F}(q^2) = \Lambda^2/(\Lambda^2 - q^2)$ are the reasonable input parameters to study the hadronic molecular candidates [150]. In Table III, we give the bound state properties for the single S -wave D_1D_1 states with the form factors $\mathcal{F}(q^2, m_E^2) = (\Lambda^2 - m_E^2)/(\Lambda^2 - q^2)$ and $\mathcal{F}(q^2) = \Lambda^2/(\Lambda^2 - q^2)$. For the S -wave D_1D_1 state with $I(J^P) = 0(1^+)$, we can obtain the bound state solution by setting the cutoff values around 1.24 GeV in the form factor $\mathcal{F}(q^2, m_E^2)$ and 0.63 GeV in the form factor $\mathcal{F}(q^2)$, which reflects that the S -wave D_1D_1 state with $I(J^P) = 0(1^+)$ can be viewed as a good doubly charmed molecular tetraquark candidate when using the two types of form factors. For the S -wave D_1D_1 states with $I(J^P) = 1(0^+, 2^+)$, we cannot find the bound state solutions with the two types of form factors until we increase the cutoff value to be around 2.5 GeV, which indicates that the S -wave D_1D_1 states with $I(J^P) = 1(0^+, 2^+)$ as the hadronic molecular candidates are no priority if we use the two types of form factors. Based on the analysis mentioned above, it is clear that the qualitative conclusion of the existence priority of the S -wave D_1D_1 hadronic molecular states does not change when increasing the contribution of the vector meson exchange interactions.

In the light pseudoscalar meson matrix \mathbb{P} , there exists the $\eta - \eta'$ mixing. In the following, we test the effect of the $\eta - \eta'$ mixing in forming the bound states. If we consider the $\eta - \eta'$ mixing, the light pseudoscalar meson matrix \mathbb{P} in the effective Lagrangians is defined as

$$\mathbb{P} = \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \alpha\eta + \beta\eta' & \pi^- \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \alpha\eta + \beta\eta' \end{pmatrix}, \quad (3.1)$$

where the parameters α and β can be related to the mixing angle θ by

$$\alpha = \frac{\cos\theta - \sqrt{2}\sin\theta}{\sqrt{6}}, \quad \beta = \frac{\sin\theta + \sqrt{2}\cos\theta}{\sqrt{6}}. \quad (3.2)$$

TABLE III: Bound state properties for the single S -wave D_1D_1 states with different form factors $\mathcal{F}(q^2, m_E^2) = (\Lambda^2 - m_E^2)/(\Lambda^2 - q^2)$ and $\mathcal{F}(q^2) = \Lambda^2/(\Lambda^2 - q^2)$. Here, “ \times ” indicates no bound state solutions until we increase the cutoff parameter to be around 2.5 GeV.

Form factors	$\mathcal{F}(q^2, m_E^2)$			$\mathcal{F}(q^2)$		
(I, J^P)	$\Lambda(\text{GeV})$	$E(\text{MeV})$	$r_{\text{RMS}}(\text{fm})$	$\Lambda(\text{GeV})$	$E(\text{MeV})$	$r_{\text{RMS}}(\text{fm})$
$(1, 0^+)$	\times	\times	\times	\times	\times	\times
$(0, 1^+)$	1.24	-0.42	4.18	0.63	-0.35	4.60
	1.35	-6.80	1.27	0.82	-5.62	1.49
	1.45	-19.50	0.81	1.00	-19.10	0.88
$(1, 2^+)$	\times	\times	\times	\times	\times	\times

Here, we use the mixing angle $\theta = -19.1^\circ$ [151]. In Table IV, we present the numerical results both without and with considering the $\eta - \eta'$ mixing for the single S -wave D_1D_1 state with $I(J^P) = 0(1^+)$. As presented in Table IV, if we take the same cutoff value without and with considering the $\eta - \eta'$ mixing, we notice that the binding energies increase less than 4.0 MeV for the single S -wave D_1D_1 state with $I(J^P) = 0(1^+)$. Thus, the $\eta - \eta'$ mixing does not obviously affect the bound state properties, which is due to the contribution from the $\eta(\eta')$ exchange interaction is small compared with the π exchange interaction [152].

TABLE IV: Bound state properties for the single S -wave D_1D_1 state with $I(J^P) = 0(1^+)$ without and with considering the $\eta - \eta'$ mixing.

Cases	Without the $\eta - \eta'$ mixing		With the $\eta - \eta'$ mixing	
$\Lambda(\text{GeV})$	$E(\text{MeV})$	$r_{\text{RMS}}(\text{fm})$	$E(\text{MeV})$	$r_{\text{RMS}}(\text{fm})$
1.24	-0.42	4.18	-0.16	5.27
1.35	-6.80	1.27	-5.21	1.42
1.45	-19.50	0.81	-15.88	0.88

B. The S -wave $D_1D_2^*$ system

In Table V, we present the obtained bound state properties for the S -wave $D_1D_2^*$ system by performing the single channel, S - D wave mixing, and coupled channel analysis. For simplicity, we collect typical loosely bound state solutions for three groups, i.e.,

- For the $D_1D_2^*$ system with $I(J^P) = 0(1^+)$, the binding energy appears at the cutoff parameter around 1.04 GeV, when we take into account the single $|\bar{3}\mathbb{S}_1\rangle$ channel. When further adding the contribution from the D -wave channels, it is easy to form the $D_1D_2^*$ bound state with $I(J^P) = 0(1^+)$. The probability of the $|\bar{3}\mathbb{S}_1\rangle$ channel is over 99%, where the S - D wave mixing effect can be ignored in forming the S -wave $D_1D_2^*$ bound state

TABLE V: Bound state properties for the S -wave $D_1 D_2^*$ system. The cutoff Λ , the binding energy E , and the root-mean-square radius r_{RMS} are in units of GeV, MeV, and fm, respectively. Here, we label the major probability for the corresponding channels in a bold manner.

Single channel analysis				
$I(J^P)$	Λ	E	r_{RMS}	
$0(1^+)$	1.04	-0.29	4.59	
	1.12	-6.01	1.33	
	1.20	-19.53	0.81	
$0(2^+)$	1.45	-0.42	4.20	
	1.62	-6.91	1.27	
	1.78	-20.03	0.81	
$1(3^+)$	2.26	-0.50	3.42	
	2.28	-4.92	1.08	
	2.30	-11.50	0.71	
S - D wave mixing analysis				
$I(J^P)$	Λ	E	r_{RMS}	$P(^3S_1/^3D_1/^5D_1/^7D_1)$
$0(1^+)$	1.04	-0.42	4.14	99.89 /0.01/ $o(0)$ /0.10
	1.12	-6.37	1.30	99.83 /0.02/ $o(0)$ /0.15
	1.20	-20.05	0.81	99.85 /0.02/ $o(0)$ /0.13
$I(J^P)$	Λ	E	r_{RMS}	$P(^5S_2/^3D_2/^5D_2/^7D_2)$
$0(2^+)$	1.42	-0.26	4.83	99.82 / $o(0)$ /0.18/ $o(0)$
	1.59	-6.10	1.35	99.62 / $o(0)$ /0.38/ $o(0)$
	1.76	-19.47	0.82	99.60 / $o(0)$ /0.40/ $o(0)$
$I(J^P)$	Λ	E	r_{RMS}	$P(^7S_3/^3D_3/^5D_3/^7D_3)$
$0(3^+)$	2.47	-0.26	5.01	98.81 /0.05/ $o(0)$ /1.14
	2.49	-0.30	4.84	98.75 /0.05/ $o(0)$ /1.20
	2.50	-0.33	4.75	98.72 /0.05/ $o(0)$ /1.23
	2.26	-0.60	3.17	99.96 /0.01/ $o(0)$ /0.03
$1(3^+)$	2.28	-5.12	1.06	99.97 /0.01/ $o(0)$ /0.02
	2.30	-11.75	0.70	99.98 /0.01/ $o(0)$ /0.01
Coupled channel analysis				
$I(J^P)$	Λ	E	r_{RMS}	$P(D_1 D_2^*/D_2^* D_2^*)$
$0(1^+)$	1.02	-0.40	4.18	99.00 /1.00
	1.09	-6.67	1.24	93.87 /6.13
	1.15	-19.07	0.79	86.86 /13.14
	1.75	-0.22	4.98	93.96 /6.14
$0(3^+)$	1.85	-5.20	1.25	66.77 /33.23
	1.95	-16.30	0.73	48.78/51.22

with $I(J^P) = 0(1^+)$. By performing the coupled channel analysis with the $D_1 D_2^*|^3S_1\rangle$ and $D_2^* D_2^*|^3S_1\rangle$ channels, we can obtain the loosely bound state solutions with the cutoff value $\Lambda = 1.02$ GeV, and the probability for the $D_1 D_2^*|^3S_1\rangle$ channel is over 85%.

- For the $D_1 D_2^*$ system with $I(J^P) = 0(2^+)$, the bound state solutions appear at the cutoff parameter larger than 1.45 GeV for the single channel case. In the S - D wave mixing case, the corresponding cutoff parameter becomes smaller if getting the same binding energy, where the D -wave contribution is not obvious.
- For the $D_1 D_2^*$ system with $I(J^P) = 0(3^+)$, we can find a shallow binding energy with the cutoff value Λ around 2.50 GeV after considering the S - D wave mixing effect. However, taking this cutoff range, there is no bound state solutions for the single channel case. If we further consider the coupled channels like $D_1 D_2^*|^7S_3\rangle$ and $D_2^* D_2^*|^7S_3\rangle$, the bound state solutions appear with the cutoff parameter around 1.75 GeV, and the $D_1 D_2^*$ channel is the dominant component when the binding energy is less than 12.00 MeV. Thus, the coupled channel effect plays an important role for forming the S -wave $D_1 D_2^*$ bound state with $I(J^P) = 0(3^+)$.
- In the cutoff range $0.8 < \Lambda < 2.5$ GeV, we cannot find the bound state solutions for the $D_1 D_2^*$ systems with $I(J^P) = 1(1^+, 2^+)$ even if we introduce the S - D wave mixing effect and the coupled channel effect.
- For the $D_1 D_2^*$ system with $I(J^P) = 1(3^+)$, we obtain the loosely bound state solutions with the binding energy around several to several tens MeV, where the RMS radius is around several fm in the cutoff value larger than 2.20 GeV. Here, the bound state solutions are almost same for the single channel case and the S - D wave mixing case.

To summarize, we can predict several possible doubly charmed molecular tetraquark candidates, such as the S -wave $D_1 D_2^*$ states with $I(J^P) = 0(1^+)$, $0(2^+)$, and $0(3^+)$. The $D_1 D_2^*$ state with $I(J^P) = 1(3^+)$ cannot be excluded as the possible doubly charmed molecular tetraquark candidate. In addition, we find the coupled channel effect plays an essential role in the formation of the $D_1 D_2^*$ bound state with $I(J^P) = 0(3^+)$.

C. The S -wave $D_2^* D_2^*$ system

In Fig. 5, we present the relevant bound state properties for the S -wave $D_2^* D_2^*$ system by performing the single channel and S - D wave mixing analysis.

For the S -wave $D_2^* D_2^*$ state with $I(J^P) = 0(1^+)$, there exists the bound state solutions with the small binding energy and the suitable RMS radius at the cutoff parameter around 1.04 GeV for the single channel analysis. Compared to the results for the single channel case, the bound state properties in the S - D wave mixing analysis change slightly.




FIG. 5: The cutoff parameter Λ dependence of the bound state solutions for the S -wave $D_2^*D_2^*$ system by performing the single channel and S - D wave mixing analysis.

For the S -wave $D_2^*D_2^*$ state with $I(J^P) = 0(3^+)$, the OBE effective interactions are a little weaker than those in the S -wave $D_2^*D_2^*$ state with $I(J^P) = 0(1^+)$. When the cutoff value is taken to be larger than 1.87 GeV, we can obtain the bound state solutions for the single channel analysis. After considering the mixing effect of the $|\mathbb{S}_3\rangle$, $|\mathbb{D}_3\rangle$, and $|\mathbb{D}_3\rangle$ channels, there exists the loosely bound state solutions at the cutoff value $\Lambda > 1.64$ GeV, and the probabilities for the D -wave components are less than 2%.

For the S -wave $D_2^*D_2^*$ states with $I(J^P) = 1(0^+, 2^+)$, their OBE effective potentials are weak attractive or repulsive by varying the cutoff parameter in the range of 0.80 to 2.50 GeV, these interactions are not strong enough to form the bound states. For the $D_2^*D_2^*$ state with $I(J^P) = 1(4^+)$, we can obtain

the loosely bound state solutions with the cutoff value around 2.25 GeV or even larger for both the single channel case and the S - D wave mixing effect case, and the S - D wave mixing effect plays a rather minor role to form the S -wave $D_2^*D_2^*$ bound state with $I(J^P) = 1(4^+)$. Here, we need to specify that the cutoff parameter is a little larger than the reasonable value around 1.00 GeV [28, 148], and we conclude that the $D_2^*D_2^*$ state with $I(J^P) = 1(4^+)$ may be viewed as the possible doubly charmed molecular tetraquark candidate.

Based on the above numerical results, the $D_2^*D_2^*$ states with $I(J^P) = 0(1^+)$ and $0(3^+)$ can be good doubly charmed molecular tetraquark candidates, and the $D_2^*D_2^*$ state with $I(J^P) = 1(4^+)$ as the possible doubly charmed molecular tetraquark candidate can be also acceptable.

In the following, we discuss how the binding energies of the S -wave D_1D_1 , $D_1D_2^*$, and $D_2^*D_2^*$ states can be affected by the S -wave DD , DD^* , and D^*D^* states. For example, when we restudy the bound state properties for the S -wave D_1D_1 state with $I(J^P) = 0(1^+)$, we consider the S -wave DD^* , D^*D^* , and D_1D_1 scattering processes. As shown in the Table VI, we present the binding energies E for the single S -wave D_1D_1 state with $I(J^P) = 0(1^+)$ in the second column, and the mass gaps $\Delta M = M_{\text{resonance}} - M_{D_1D_1}$ between the mass of the obtained resonance and the D_1D_1 threshold after considering the S -wave $DD^*/D^*D^*/D_1D_1$ scattering processes in the third column. From the Table VI, we can find the masses for the S -wave D_1D_1 bound state with $I(J^P) = 0(1^+)$ are almost the same with the masses for the obtained resonance when we take the same cutoff values. It indicates that the lower DD^* and D^*D^* states barely affect the binding energies for the S -wave D_1D_1 state with $I(J^P) = 0(1^+)$, which is due to the mass threshold of the D_1D_1 channel is far from that of the $D^{(*)}D^{(*)}$ channels around 1.0 GeV.

TABLE VI: Bound state properties for the single S -wave D_1D_1 state with $I(J^P) = 0(1^+)$ without and with considering the scattering states.

$\Lambda(\text{GeV})$	$E(\text{MeV})$	$M_{\text{resonance}} - M_{D_1D_1}(\text{MeV})$
1.26	-1.01	-1.08
1.35	-6.80	-6.90
1.42	-14.96	-15.10

IV. SUMMARY

In the past decades, the study of the exotic hadronic state has become an influential and attractive research field for the hadron physics. Benefited from the accumulation of experimental data with the high precision, the LHCb Collaboration reported a new structure T_{cc}^+ existing in the $D^0D^0\pi^+$ invariant mass spectrum [1], where the T_{cc}^+ state can be regarded as the DD^* doubly charmed molecular state with $J^P = 1^+$ [3–20]. This new observation provides us a good opportunity to construct the family of the doubly charmed molecular tetraquarks.

In this work, we systematically study the S -wave interactions between a pair of charmed mesons in the T -doublet, where we can predict possible doubly charmed molecular tetraquarks. For the interactions between charmed meson pair in the T -doublet, we consider both the long-range contribution from the pseudoscalar meson exchange and the short-range and medium-range contributions from the vector and scalar meson exchanges. In the realistic calculation, we explore the roles of the single channel, the S - D wave mixing effect, and the coupled channel effect to form the possible doubly charmed molecular tetraquark states, simultaneously.

Our numerical results show that the S -wave $D_1 D_1$ state with $I(J^P) = 0(1^+)$, the S -wave $D_1 D_2^*$ states with $I(J^P) = 0(1^+, 2^+, 3^+)$, and the S -wave $D_2^* D_2^*$ states with $I(J^P) = 0(1^+, 3^+)$ can be recommended as the prime doubly charmed molecular tetraquark candidates, which are consistent with the theoretical predictions in Ref. [9]. From the numerical results, we also can find that the binding properties of these most promising doubly charmed molecular tetraquark candidates are not significantly dependent on the cutoff values. Meanwhile, the S -wave $D_1 D_1$ state with $I(J^P) = 1(2^+)$, the S -wave $D_1 D_2^*$ state with $I(J^P) = 1(3^+)$, and the S -wave $D_2^* D_2^*$ state with $I(J^P) = 1(4^+)$ may be the secondary doubly charmed molecular tetraquark candidates. Their allowed decay modes include (i) two charmed mesons, (ii) a doubly charmed baryon plus a light anti-baryon, (iii) two charmed mesons plus one light meson, and (iv) two charmed mesons plus one photon.

Experimental search for these predicted doubly charmed molecular tetraquark candidates is an interesting and important research topic. With the accumulation of the Run II and Run III data [153], the LHCb Collaboration has the potential to hunt for these predicted doubly charmed molecular tetraquarks.

ACKNOWLEDGMENTS

F. L. Wang would like to thank J. Z. Wang for very helpful discussions. This work is supported by the China National Funds for Distinguished Young Scientists under Grant No. 11825503, National Key Research and Development Program of China under Contract No. 2020YFA0406400, the 111 Project under Grant No. B20063, and the National Natural Science Foundation of China under Grant No. 12047501.

Appendix A: The operator matrix elements $\langle f | \mathcal{O}_k^{(\prime)} | i \rangle_{[J]}$

For the operators $\mathcal{O}_k^{(\prime)}$ ($k = 1, \dots, 11$), they should be sandwiched by the relevant spin-orbit wave functions $|^{2S+1}L_J\rangle$. In Table VII, we present the operator matrix elements $\langle f | \mathcal{O}_k^{(\prime)} | i \rangle_{[J]}$ [124].

-
- [1] R. Aaij *et al.* (LHCb Collaboration), Observation of an exotic narrow doubly charmed tetraquark, [arXiv:2109.01038](#).
 - [2] R. Aaij *et al.* (LHCb Collaboration), Study of the doubly charmed tetraquark T_{cc}^+ , [arXiv:2109.01056](#).
 - [3] N. Li, Z. F. Sun, X. Liu and S. L. Zhu, Perfect DD^* molecular prediction matching the T_{cc} observation at LHCb, *Chin. Phys. Lett.* **38**, 092001 (2021).
 - [4] R. Chen, Q. Huang, X. Liu and S. L. Zhu, Another doubly charmed molecular resonance $T_{cc}^{*+}(3876)$, [arXiv:2108.01911](#).
 - [5] H. Ren, F. Wu and R. Zhu, Hadronic molecule interpretation of T_{cc}^+ and its beauty-partners, [arXiv:2109.02531](#).
 - [6] Q. Xin and Z. G. Wang, Analysis of the axialvector doubly charmed tetraquark molecular states with the QCD sum rules, [arXiv:2108.12597](#).
 - [7] X. Chen, Doubly heavy tetraquark states $cc\bar{u}\bar{d}$ and $bb\bar{u}\bar{d}$, [arXiv:2109.02828](#).
 - [8] M. Albaladejo, T_{cc}^+ coupled channel analysis and predictions, [arXiv:2110.02944](#).
 - [9] X. K. Dong, F. K. Guo and B. S. Zou, A survey of heavy-heavy hadronic molecules, [arXiv:2108.02673](#).
 - [10] V. Baru, X. K. Dong, M. L. Du, A. Filin, F. K. Guo, C. Hanhart, A. Nefediev, J. Nieves and Q. Wang, Effective range expansion for narrow near-threshold resonances, [arXiv:2110.07484](#).
 - [11] M. L. Du, V. Baru, X. K. Dong, A. Filin, F. K. Guo, C. Hanhart, A. Nefediev, J. Nieves and Q. Wang, Coupled-channel approach to T_{cc}^+ including three-body effects, [arXiv:2110.13765](#).
 - [12] Y. Kamiya, T. Hyodo and A. Ohnishi, Femtoscopic study on DD^* and $D\bar{D}^*$ interactions for T_{cc} and $X(3872)$, [arXiv:2203.13814](#).
 - [13] M. Padmanath and S. Prelovsek, Evidence for a doubly charm tetraquark pole in DD^* scattering on the lattice, [arXiv:2202.10110](#).
 - [14] S. S. Agaev, K. Azizi and H. Sundu, Hadronic molecule model for the doubly charmed state T_{cc}^+ , [arXiv:2201.02788](#).
 - [15] H. W. Ke, X. H. Liu and X. Q. Li, Possible molecular states of $D^{(*)}D^{(*)}$ and $B^{(*)}B^{(*)}$ within the Bethe-Salpeter framework, *Eur. Phys. J. C* **82** (2022) no.2, 144.
 - [16] M. J. Zhao, Z. Y. Wang, C. Wang and X. H. Guo, Investigation of the possible $D\bar{D}^*/B\bar{B}^*$ and $DD^*/\bar{B}\bar{B}^*$ molecule states, [arXiv:2112.12633](#).
 - [17] C. Deng and S. L. Zhu, T_{cc}^+ and its partners, *Phys. Rev. D* **105** (2022) no.5, 054015.
 - [18] N. Santowsky and C. S. Fischer, Four-quark states with charm quarks in a two-body Bethe-Salpeter approach, [arXiv:2111.15310](#).
 - [19] L. R. Dai, R. Molina and E. Oset, Prediction of new T_{cc} states of D^*D^* and $D_s^*D^*$ molecular nature, *Phys. Rev. D* **105** (2022) no.1, 016029.
 - [20] A. Feijoo, W. H. Liang and E. Oset, $D^0 D^0 \pi^+$ mass distribution in the production of the T_{cc} exotic state, *Phys. Rev. D* **104** (2021) no.11, 114015.
 - [21] T. Guo, J. Li, J. Zhao and L. He, Mass spectra of doubly heavy tetraquarks in an improved chromomagnetic interaction model, *Phys. Rev. D* **105** (2022) no.1, 014021.
 - [22] X. Z. Weng, W. Z. Deng and S. L. Zhu, Doubly heavy tetraquarks in an extended chromomagnetic model, *Chin. Phys. C* **46** (2022) no.1, 013102.
 - [23] Y. Kim, M. Oka and K. Suzuki, Doubly heavy tetraquarks in a

TABLE VII: The operator matrix elements $\langle f|\mathcal{O}_k^{(i)}|i\rangle_{[J]}$ for the operators $\mathcal{O}_k^{(i)}$ ($k = 1, \dots, 11$) in the effective potentials.

$\langle f \mathcal{O}_k^{(i)} i\rangle$	$\langle f \mathcal{O}_k^{(i)} i\rangle_{[J]}$		
$\langle D_1 D_1 \mathcal{O}_1 D_1 D_1 \rangle$	Diag(1, 1) _[0]	Diag(1, 1) _[1]	Diag(1, 1, 1) _[2]
$\langle D_1 D_1 \mathcal{O}_2 D_1 D_1 \rangle$	Diag(2, -1) _[0]	Diag(1, 1) _[1]	Diag(-1, 2, -1) _[2]
$\langle D_1 D_1 \mathcal{O}_3 D_1 D_1 \rangle$	$\begin{pmatrix} 0 & \sqrt{2} \\ \sqrt{2} & 2 \end{pmatrix}_{[0]}$	$\begin{pmatrix} 0 & -\sqrt{2} \\ -\sqrt{2} & 1 \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & \sqrt{\frac{2}{5}} & -\sqrt{\frac{14}{5}} \\ \sqrt{\frac{2}{5}} & 0 & -\frac{2}{\sqrt{7}} \\ -\sqrt{\frac{14}{5}} & -\frac{2}{\sqrt{7}} & -\frac{3}{7} \end{pmatrix}_{[2]}$
$\langle D_1 D_2^* \mathcal{O}_4 D_1 D_2^* \rangle$	Diag(1, 1, 1, 1) _[1]	Diag(1, 1, 1, 1) _[2]	Diag(1, 1, 1, 1) _[3]
$\langle D_2^* D_1 \mathcal{O}_4' D_2^* D_1 \rangle$	Diag(1, 1, 1, 1) _[1]	Diag(1, 1, 1, 1) _[2]	Diag(1, 1, 1, 1) _[3]
$\langle D_1 D_2^* \mathcal{O}_5 D_1 D_2^* \rangle$	Diag($\frac{3}{2}, \frac{3}{2}, \frac{1}{2}, -1$) _[1]	Diag($\frac{1}{2}, \frac{3}{2}, \frac{1}{2}, -1$) _[2]	Diag(-1, $\frac{3}{2}, \frac{1}{2}, -1$) _[3]
$\langle D_2^* D_1 \mathcal{O}_5' D_2^* D_1 \rangle$	Diag($\frac{3}{2}, \frac{3}{2}, \frac{1}{2}, -1$) _[1]	Diag($\frac{1}{2}, \frac{3}{2}, \frac{1}{2}, -1$) _[2]	Diag(-1, $\frac{3}{2}, \frac{1}{2}, -1$) _[3]
$\langle D_1 D_2^* \mathcal{O}_6 D_1 D_2^* \rangle$	$\begin{pmatrix} 0 & \frac{3}{5\sqrt{2}} & \sqrt{\frac{6}{5}} & \frac{\sqrt{\frac{21}{5}}}{5} \\ \frac{3}{5\sqrt{2}} & -\frac{3}{10} & \sqrt{\frac{3}{5}} & -\frac{\sqrt{\frac{3}{7}}}{5} \\ \sqrt{\frac{6}{5}} & \sqrt{\frac{3}{5}} & \frac{1}{2} & \frac{2}{\sqrt{35}} \\ \frac{\sqrt{\frac{21}{5}}}{5} & -\frac{\sqrt{\frac{3}{7}}}{5} & \frac{2}{\sqrt{35}} & \frac{48}{35} \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & -\frac{3\sqrt{2}}{5} & -\sqrt{\frac{7}{10}} & \frac{\sqrt{7}}{5} \\ -\frac{3\sqrt{2}}{5} & \frac{3}{10} & \frac{3}{\sqrt{35}} & -\frac{3\sqrt{\frac{2}{7}}}{5} \\ -\sqrt{\frac{7}{10}} & \frac{3}{\sqrt{35}} & -\frac{3}{14} & \frac{4\sqrt{\frac{2}{35}}}{7} \\ \frac{\sqrt{7}}{5} & -\frac{3\sqrt{\frac{2}{7}}}{5} & \frac{4\sqrt{\frac{2}{35}}}{7} & \frac{12}{35} \end{pmatrix}_{[2]}$	$\begin{pmatrix} 0 & \frac{3}{5\sqrt{2}} & -\frac{1}{\sqrt{5}} & -\frac{4\sqrt{3}}{5} \\ \frac{3}{5\sqrt{2}} & -\frac{3}{35} & -\frac{6\sqrt{\frac{2}{5}}}{7} & -\frac{6\sqrt{6}}{35} \\ -\frac{1}{\sqrt{5}} & -\frac{6\sqrt{\frac{2}{5}}}{7} & -\frac{4}{7} & \frac{\sqrt{\frac{3}{5}}}{7} \\ -\frac{4\sqrt{3}}{5} & -\frac{6\sqrt{6}}{35} & \frac{\sqrt{\frac{3}{5}}}{7} & -\frac{22}{35} \end{pmatrix}_{[3]}$
$\langle D_2^* D_1 \mathcal{O}_6' D_2^* D_1 \rangle$	$\begin{pmatrix} 0 & \frac{3}{5\sqrt{2}} & -\sqrt{\frac{6}{5}} & \frac{\sqrt{\frac{21}{5}}}{5} \\ \frac{3}{5\sqrt{2}} & -\frac{3}{10} & -\sqrt{\frac{3}{5}} & -\frac{\sqrt{\frac{3}{7}}}{5} \\ -\sqrt{\frac{6}{5}} & -\sqrt{\frac{3}{5}} & \frac{1}{2} & -\frac{2}{\sqrt{35}} \\ \frac{\sqrt{\frac{21}{5}}}{5} & -\frac{\sqrt{\frac{3}{7}}}{5} & -\frac{2}{\sqrt{35}} & \frac{48}{35} \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & \frac{3\sqrt{2}}{5} & -\sqrt{\frac{7}{10}} & -\frac{\sqrt{7}}{5} \\ \frac{3\sqrt{2}}{5} & \frac{3}{10} & -\frac{3}{\sqrt{35}} & -\frac{3\sqrt{\frac{2}{7}}}{5} \\ -\sqrt{\frac{7}{10}} & -\frac{3}{\sqrt{35}} & -\frac{3}{14} & -\frac{4\sqrt{\frac{2}{35}}}{7} \\ -\frac{\sqrt{7}}{5} & -\frac{3\sqrt{\frac{2}{7}}}{5} & -\frac{4\sqrt{\frac{2}{35}}}{7} & \frac{12}{35} \end{pmatrix}_{[2]}$	$\begin{pmatrix} 0 & \frac{3}{5\sqrt{2}} & \frac{1}{\sqrt{5}} & -\frac{4\sqrt{3}}{5} \\ \frac{3}{5\sqrt{2}} & -\frac{3}{35} & \frac{6\sqrt{\frac{2}{5}}}{7} & -\frac{6\sqrt{6}}{35} \\ \frac{1}{\sqrt{5}} & \frac{6\sqrt{\frac{2}{5}}}{7} & -\frac{4}{7} & -\frac{\sqrt{\frac{3}{5}}}{7} \\ -\frac{4\sqrt{3}}{5} & -\frac{6\sqrt{6}}{35} & -\frac{\sqrt{\frac{3}{5}}}{7} & -\frac{22}{35} \end{pmatrix}_{[3]}$
$\langle D_2^* D_1 \mathcal{O}_7 D_1 D_2^* \rangle$	Diag($\frac{1}{6}, \frac{1}{6}, \frac{1}{2}, 1$) _[1]	Diag($\frac{1}{2}, \frac{1}{6}, \frac{1}{2}, 1$) _[2]	Diag(1, $\frac{1}{6}, \frac{1}{2}, 1$) _[3]
$\langle D_1 D_2^* \mathcal{O}_7' D_2^* D_1 \rangle$	Diag($\frac{1}{6}, \frac{1}{6}, \frac{1}{2}, 1$) _[1]	Diag($\frac{1}{2}, \frac{1}{6}, \frac{1}{2}, 1$) _[2]	Diag(1, $\frac{1}{6}, \frac{1}{2}, 1$) _[3]
$\langle D_2^* D_1 \mathcal{O}_8 D_1 D_2^* \rangle$	$\begin{pmatrix} 0 & -\frac{23}{15\sqrt{2}} & -2\sqrt{\frac{2}{15}} & -\frac{\sqrt{\frac{7}{6}}}{5} \\ -\frac{23}{15\sqrt{2}} & \frac{23}{30} & -\frac{2}{\sqrt{15}} & \frac{1}{5\sqrt{21}} \\ 2\sqrt{\frac{2}{15}} & \frac{2}{\sqrt{15}} & \frac{1}{2} & -\frac{2}{\sqrt{35}} \\ -\frac{\sqrt{\frac{7}{6}}}{5} & \frac{1}{5\sqrt{21}} & \frac{2}{\sqrt{35}} & \frac{24}{35} \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & -\frac{2\sqrt{2}}{5} & -\sqrt{\frac{7}{10}} & -\frac{\sqrt{7}}{5} \\ \frac{2\sqrt{2}}{5} & -\frac{23}{30} & -\frac{2}{\sqrt{35}} & \frac{\sqrt{\frac{7}{6}}}{5} \\ -\sqrt{\frac{7}{10}} & \frac{2}{\sqrt{35}} & -\frac{3}{14} & -\frac{4\sqrt{\frac{2}{35}}}{7} \\ \frac{\sqrt{7}}{5} & \frac{\sqrt{\frac{7}{6}}}{5} & \frac{4\sqrt{\frac{2}{35}}}{7} & \frac{6}{35} \end{pmatrix}_{[2]}$	$\begin{pmatrix} 0 & -\frac{1}{5\sqrt{2}} & -\frac{1}{\sqrt{5}} & -\frac{2\sqrt{3}}{5} \\ -\frac{1}{5\sqrt{2}} & \frac{23}{105} & \frac{4\sqrt{\frac{2}{5}}}{7} & \frac{2\sqrt{6}}{35} \\ \frac{1}{\sqrt{5}} & -\frac{4\sqrt{\frac{2}{5}}}{7} & -\frac{4}{7} & -\frac{\sqrt{\frac{3}{5}}}{7} \\ -\frac{2\sqrt{3}}{5} & \frac{2\sqrt{6}}{35} & \frac{\sqrt{\frac{3}{5}}}{7} & -\frac{11}{35} \end{pmatrix}_{[3]}$
$\langle D_1 D_2^* \mathcal{O}_8' D_2^* D_1 \rangle$	$\begin{pmatrix} 0 & -\frac{23}{15\sqrt{2}} & 2\sqrt{\frac{2}{15}} & -\frac{\sqrt{\frac{7}{6}}}{5} \\ -\frac{23}{15\sqrt{2}} & \frac{23}{30} & \frac{2}{\sqrt{15}} & \frac{1}{5\sqrt{21}} \\ -2\sqrt{\frac{2}{15}} & -\frac{2}{\sqrt{15}} & \frac{1}{2} & \frac{2}{\sqrt{35}} \\ -\frac{\sqrt{\frac{7}{6}}}{5} & \frac{1}{5\sqrt{21}} & -\frac{2}{\sqrt{35}} & \frac{24}{35} \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & \frac{2\sqrt{2}}{5} & -\sqrt{\frac{7}{10}} & \frac{\sqrt{7}}{5} \\ -\frac{2\sqrt{2}}{5} & -\frac{23}{30} & \frac{2}{\sqrt{35}} & \frac{\sqrt{\frac{7}{6}}}{5} \\ -\sqrt{\frac{7}{10}} & -\frac{2}{\sqrt{35}} & -\frac{3}{14} & \frac{4\sqrt{\frac{2}{35}}}{7} \\ -\frac{\sqrt{7}}{5} & \frac{\sqrt{\frac{7}{6}}}{5} & -\frac{4\sqrt{\frac{2}{35}}}{7} & \frac{6}{35} \end{pmatrix}_{[2]}$	$\begin{pmatrix} 0 & -\frac{1}{5\sqrt{2}} & \frac{1}{\sqrt{5}} & -\frac{2\sqrt{3}}{5} \\ -\frac{1}{5\sqrt{2}} & \frac{23}{105} & -\frac{4\sqrt{\frac{2}{5}}}{7} & \frac{2\sqrt{6}}{35} \\ -\frac{1}{\sqrt{5}} & \frac{4\sqrt{\frac{2}{5}}}{7} & -\frac{4}{7} & \frac{\sqrt{\frac{3}{5}}}{7} \\ -\frac{2\sqrt{3}}{5} & \frac{2\sqrt{6}}{35} & -\frac{\sqrt{\frac{3}{5}}}{7} & -\frac{11}{35} \end{pmatrix}_{[3]}$
$\langle D_2^* D_2^* \mathcal{O}_9 D_2^* D_2^* \rangle$	Diag(1, 1) _[0]	Diag(1, 1, 1) _[1]	Diag(1, 1, 1, 1) _[2]
$\langle D_2^* D_2^* \mathcal{O}_{10} D_2^* D_2^* \rangle$	Diag(1, 1, 1) _[3]	Diag(1, 1, 1) _[4]	
$\langle D_2^* D_2^* \mathcal{O}_{10}' D_2^* D_2^* \rangle$	Diag($\frac{3}{2}, \frac{3}{4}$) _[0]	Diag($\frac{5}{4}, \frac{5}{4}, 0$) _[1]	Diag($\frac{3}{4}, \frac{3}{4}, \frac{3}{4}, -1$) _[2]
$\langle D_2^* D_2^* \mathcal{O}_{11} D_2^* D_2^* \rangle$	Diag($0, \frac{5}{4}, 0$) _[3]	Diag(-1, $\frac{3}{4}, -1$) _[4]	
$\langle D_2^* D_2^* \mathcal{O}_{11}' D_2^* D_2^* \rangle$	$\begin{pmatrix} 0 & \frac{3\sqrt{\frac{7}{10}}}{2} \\ \frac{3\sqrt{\frac{7}{10}}}{2} & \frac{15}{14} \end{pmatrix}_{[0]}$	$\begin{pmatrix} 0 & -\frac{13}{10\sqrt{2}} & \frac{3\sqrt{7}}{10} \\ -\frac{13}{10\sqrt{2}} & \frac{13}{20} & -\frac{3}{5\sqrt{14}} \\ \frac{3\sqrt{7}}{10} & -\frac{3}{5\sqrt{14}} & \frac{36}{35} \end{pmatrix}_{[1]}$	$\begin{pmatrix} 0 & \frac{3\sqrt{\frac{7}{2}}}{10} & -\frac{3\sqrt{\frac{5}{14}}}{2} & \frac{9}{5\sqrt{14}} \\ \frac{3\sqrt{\frac{7}{2}}}{10} & 0 & -\frac{3}{2\sqrt{5}} & 0 \\ -\frac{3\sqrt{\frac{5}{14}}}{2} & -\frac{3}{2\sqrt{5}} & -\frac{45}{196} & -\frac{18}{49\sqrt{5}} \\ \frac{9}{5\sqrt{14}} & 0 & -\frac{18}{49\sqrt{5}} & 0 \end{pmatrix}_{[2]}$
	$\begin{pmatrix} 0 & \frac{3\sqrt{3}}{10} & -\frac{3\sqrt{3}}{5} \\ \frac{3\sqrt{3}}{10} & \frac{13}{70} & -\frac{18}{35} \\ -\frac{3\sqrt{3}}{5} & -\frac{18}{35} & -\frac{33}{70} \end{pmatrix}_{[3]}$	$\begin{pmatrix} 0 & \frac{3}{\sqrt{70}} & -\sqrt{\frac{11}{7}} \\ \frac{3}{\sqrt{70}} & \frac{15}{49} & -\frac{3\sqrt{\frac{35}{2}}}{49} \\ -\sqrt{\frac{11}{7}} & -\frac{3\sqrt{\frac{35}{2}}}{49} & \frac{65}{98} \end{pmatrix}_{[4]}$	

- chiral-diquark picture, [arXiv:2202.06520](#).
- [24] S. S. Agaev, K. Azizi and H. Sundu, Newly observed exotic doubly charmed meson T_{cc}^+ , *Nucl. Phys. B* **975** (2022), 115650.
 - [25] R. Molina, T. Branz and E. Oset, A new interpretation for the $D_s^*(2573)$ and the prediction of novel exotic charmed mesons, *Phys. Rev. D* **82** (2010), 014010.
 - [26] A. V. Manohar and M. B. Wise, Exotic $QQ\bar{q}\bar{q}$ states in QCD, *Nucl. Phys. B* **399** (1993), 17-33.
 - [27] T. E. O. Ericson and G. Karl, Strength of pion exchange in hadronic molecules, *Phys. Lett. B* **309** (1993), 426-430.
 - [28] N. A. Tornqvist, From the deuteron to deusons, an analysis of deuteron-like meson-meson bound states, *Z. Phys. C* **61** (1994), 525-537.
 - [29] D. Janc and M. Rosina, The $T_{cc} = DD^*$ molecular state, *Few Body Syst.* **35**, 175-196 (2004).
 - [30] G. J. Ding, J. F. Liu and M. L. Yan, Dynamics of Hadronic Molecule in One-Boson Exchange Approach and Possible Heavy Flavor Molecules, *Phys. Rev. D* **79** (2009), 054005.
 - [31] S. Ohkoda, Y. Yamaguchi, S. Yasui, K. Sudoh and A. Hosaka, Exotic mesons with double charm and bottom flavor, *Phys. Rev. D* **86** (2012), 034019.
 - [32] N. Li, Z. F. Sun, X. Liu and S. L. Zhu, Coupled-channel analysis of The possible $D^{(*)}D^{(*)}$, $\bar{B}^{(*)}\bar{B}^{(*)}$ and $D^{(*)}\bar{B}^{(*)}$ molecular states, *Phys. Rev. D* **88**, 114008 (2013).
 - [33] H. Xu, B. Wang, Z. W. Liu and X. Liu, DD^* potentials in chiral perturbation theory and possible molecular states, *Phys. Rev. D* **99**, 014027 (2019).
 - [34] L. Tang, B. D. Wan, K. Maltman and C. F. Qiao, Doubly Heavy Tetraquarks in QCD Sum Rules, *Phys. Rev. D* **101** (2020) no.9, 094032.
 - [35] M. Z. Liu, T. W. Wu, M. Pavon Valderrama, J. J. Xie and L. S. Geng, Heavy-quark spin and flavor symmetry partners of the $X(3872)$ revisited: What can we learn from the one-boson exchange model?, *Phys. Rev. D* **99**, 094018 (2019).
 - [36] Z. M. Ding, H. Y. Jiang and J. He, Molecular states from $D^{(*)}\bar{D}^{(*)}/B^{(*)}\bar{B}^{(*)}$ and $D^{(*)}D^{(*)}/\bar{B}^{(*)}\bar{B}^{(*)}$ interactions, *Eur. Phys. J. C* **80**, 1179 (2020).
 - [37] L. Heller and J. A. Tjon, On the Existence of Stable Dimesons, *Phys. Rev. D* **35**, 969 (1987).
 - [38] J. Carlson, L. Heller and J. A. Tjon, Stability of Dimesons, *Phys. Rev. D* **37**, 744 (1988).
 - [39] B. Silvestre-Brac and C. Semay, Systematics of $L = 0$ $q^2\bar{q}^2$ systems, *Z. Phys. C* **57**, 273-282 (1993).
 - [40] C. Semay and B. Silvestre-Brac, Diquonia and potential models, *Z. Phys. C* **61**, 271-275 (1994).
 - [41] M. A. Moinester, How to search for doubly charmed baryons and tetraquarks, *Z. Phys. A* **355**, 349-362 (1996).
 - [42] S. Pepin, F. Stancu, M. Genovese and J. M. Richard, Tetraquarks with color blind forces in chiral quark models, *Phys. Lett. B* **393**, 119-123 (1997).
 - [43] B. A. Gelman and S. Nussinov, Does a narrow tetraquark $cc\bar{u}\bar{d}$ state exist?, *Phys. Lett. B* **551**, 296-304 (2003).
 - [44] J. Vijande, F. Fernandez, A. Valcarce and B. Silvestre-Brac, Tetraquarks in a chiral constituent quark model, *Eur. Phys. J. A* **19**, 383 (2004).
 - [45] F. S. Navarra, M. Nielsen and S. H. Lee, QCD sum rules study of $QQ\bar{u}\bar{d}$ mesons, *Phys. Lett. B* **649**, 166-172 (2007).
 - [46] J. Vijande, E. Weissman, A. Valcarce and N. Barnea, Are there compact heavy four-quark bound states?, *Phys. Rev. D* **76**, 094027 (2007).
 - [47] D. Ebert, R. N. Faustov, V. O. Galkin and W. Lucha, Masses of tetraquarks with two heavy quarks in the relativistic quark model, *Phys. Rev. D* **76**, 114015 (2007).
 - [48] S. H. Lee and S. Yasui, Stable multi-quark states with heavy quarks in a diquark model, *Eur. Phys. J. C* **64**, 283-295 (2009).
 - [49] Y. Yang, C. Deng, J. Ping and T. Goldman, S-wave $QQ\bar{q}\bar{q}$ state in the constituent quark model, *Phys. Rev. D* **80**, 114023 (2009).
 - [50] S. Q. Luo, K. Chen, X. Liu, Y. R. Liu and S. L. Zhu, Exotic tetraquark states with the $qq\bar{Q}\bar{Q}$ configuration, *Eur. Phys. J. C* **77**, no.10, 709 (2017).
 - [51] M. Karliner and J. L. Rosner, Discovery of doubly-charmed Ξ_{cc} baryon implies a stable $(bb\bar{u}\bar{d})$ tetraquark, *Phys. Rev. Lett.* **119**, no.20, 202001 (2017).
 - [52] E. J. Eichten and C. Quigg, Heavy-quark symmetry implies stable heavy tetraquark mesons $Q_i Q_j \bar{q}_k \bar{q}_l$, *Phys. Rev. Lett.* **119**, no.20, 202002 (2017).
 - [53] Z. G. Wang, Analysis of the axialvector doubly heavy tetraquark states with QCD sum rules, *Acta Phys. Polon. B* **49**, 1781 (2018).
 - [54] W. Park, S. Noh and S. H. Lee, Masses of the doubly heavy tetraquarks in a constituent quark model, *Nucl. Phys. A* **983**, 1-19 (2019).
 - [55] P. Junnarkar, N. Mathur and M. Padmanath, Study of doubly heavy tetraquarks in Lattice QCD, *Phys. Rev. D* **99**, no.3, 034507 (2019).
 - [56] C. Deng, H. Chen and J. Ping, Systematical investigation on the stability of doubly heavy tetraquark states, *Eur. Phys. J. A* **56**, no.1, 9 (2020).
 - [57] L. Maiani, A. D. Polosa and V. Riquer, Hydrogen bond of QCD in doubly heavy baryons and tetraquarks, *Phys. Rev. D* **100**, no.7, 074002 (2019).
 - [58] G. Yang, J. Ping and J. Segovia, Doubly-heavy tetraquarks, *Phys. Rev. D* **101**, no.1, 014001 (2020).
 - [59] Y. Tan, W. Lu and J. Ping, Systematics of $QQ\bar{q}\bar{q}$ in a chiral constituent quark model, *Eur. Phys. J. Plus* **135**, no.9, 716 (2020).
 - [60] Q. F. Lü, D. Y. Chen and Y. B. Dong, Masses of doubly heavy tetraquarks $T_{QQ'}$ in a relativized quark model, *Phys. Rev. D* **102**, no.3, 034012 (2020).
 - [61] E. Braaten, L. P. He and A. Mohapatra, Masses of doubly heavy tetraquarks with error bars, *Phys. Rev. D* **103**, no.1, 016001 (2021).
 - [62] D. Gao, D. Jia, Y. J. Sun, Z. Zhang, W. N. Liu and Q. Mei, Masses of doubly heavy tetraquark states with isospin $= \frac{1}{2}$ and 1 and spin-parity 1^{++} , [arXiv:2007.15213](#).
 - [63] J. B. Cheng, S. Y. Li, Y. R. Liu, Z. G. Si and T. Yao, Double-heavy tetraquark states with heavy diquark-antiquark symmetry, *Chin. Phys. C* **45**, no.4, 043102 (2021).
 - [64] S. Noh, W. Park and S. H. Lee, The Doubly-heavy Tetraquarks ($qq'\bar{Q}\bar{Q}'$) in a Constituent Quark Model with a Complete Set of Harmonic Oscillator Bases, *Phys. Rev. D* **103**, 114009 (2021).
 - [65] R. N. Faustov, V. O. Galkin and E. M. Savchenko, Heavy tetraquarks in the relativistic quark model, *Universe* **7**, no.4, 94 (2021).
 - [66] S. K. Choi *et al.* [Belle Collaboration], Observation of a Narrow Charmonium-Like State in Exclusive $B^\pm \rightarrow K^\pm \pi^+ \pi^- J/\psi$ Decays, *Phys. Rev. Lett.* **91**, 262001 (2003).
 - [67] C. Y. Wong, Molecular states of heavy quark mesons, *Phys. Rev. C* **69**, 055202 (2004).
 - [68] E. S. Swanson, Short range structure in the $X(3872)$, *Phys. Lett. B* **588**, 189 (2004).
 - [69] M. Suzuki, The $X(3872)$ boson: Molecule or charmonium, *Phys. Rev. D* **72**, 114013 (2005).
 - [70] Y. R. Liu, X. Liu, W. Z. Deng, and S. L. Zhu, Is $X(3872)$ really

- a molecular state?, *Eur. Phys. J. C* **56**, 63 (2008).
- [71] C. E. Thomas and F. E. Close, Is $X(3872)$ a molecule?, *Phys. Rev. D* **78**, 034007 (2008).
- [72] X. Liu, Z. G. Luo, Y. R. Liu, and S. L. Zhu, $X(3872)$ and other possible heavy molecular states, *Eur. Phys. J. C* **61**, 411 (2009).
- [73] I. W. Lee, A. Faessler, T. Gutsche, and V. E. Lyubovitskij, $X(3872)$ as a molecular DD^* state in a potential model, *Phys. Rev. D* **80**, 094005 (2009).
- [74] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C. P. Shen, C. E. Thomas, A. Vairo and C. Z. Yuan, The XYZ states: Experimental and theoretical status and perspectives, *Phys. Rep.* **873**, 1 (2020).
- [75] H. X. Chen, W. Chen, X. Liu, and S. L. Zhu, The hidden-charm pentaquark and tetraquark states, *Phys. Rep.* **639**, 1 (2016).
- [76] X. Liu, An overview of XYZ new particles, *Chin. Sci. Bull.* **59**, 3815 (2014).
- [77] A. Hosaka, T. Iijima, K. Miyabayashi, Y. Sakai, and S. Yasui, Exotic hadrons with heavy flavors: X , Y , Z , and related states, *Prog. Theor. Exp. Phys.* **2016**, 062C01 (2016).
- [78] Y. R. Liu, H. X. Chen, W. Chen, X. Liu, and S. L. Zhu, Pentaquark and tetraquark states, *Prog. Part. Nucl. Phys.* **107**, 237 (2019).
- [79] S. L. Olsen, T. Skwarnicki, and D. Zieminska, Nonstandard heavy mesons and baryons: Experimental evidence, *Rev. Mod. Phys.* **90**, 015003 (2018).
- [80] F. K. Guo, C. Hanhart, U. G. Meißner, Q. Wang, Q. Zhao, and B. S. Zou, Hadronic molecules, *Rev. Mod. Phys.* **90**, 015004 (2018).
- [81] J. J. Wu, R. Molina, E. Oset and B. S. Zou, Prediction of narrow N^* and Λ^* resonances with hidden charm above 4 GeV, *Phys. Rev. Lett.* **105**, 232001 (2010).
- [82] M. Karliner and J. L. Rosner, New Exotic Meson and Baryon Resonances from Doubly-Heavy Hadronic Molecules, *Phys. Rev. Lett.* **115**, 122001 (2015).
- [83] W. L. Wang, F. Huang, Z. Y. Zhang and B. S. Zou, $\Sigma_c \bar{D}$ and $\Lambda_c \bar{D}$ states in a chiral quark model, *Phys. Rev. C* **84**, 015203 (2011).
- [84] Z. C. Yang, Z. F. Sun, J. He, X. Liu and S. L. Zhu, The possible hidden-charm molecular baryons composed of anti-charmed meson and charmed baryon, *Chin. Phys. C* **36**, 6 (2012).
- [85] J. J. Wu, T. S. H. Lee and B. S. Zou, Nucleon Resonances with Hidden Charm in Coupled-Channel Models, *Phys. Rev. C* **85**, 044002 (2012).
- [86] R. Chen, X. Liu, X. Q. Li and S. L. Zhu, Identifying exotic hidden-charm pentaquarks, *Phys. Rev. Lett.* **115**, 132002 (2015).
- [87] R. Aaij *et al.* [LHCb Collaboration], Observation of J/ψ Resonances Consistent with Pentaquark States in $\Lambda_b^0 \rightarrow J/\psi K^- p$ Decays, *Phys. Rev. Lett.* **115**, 072001 (2015).
- [88] R. Aaij *et al.* [LHCb Collaboration], Observation of a narrow pentaquark state, $P_c(4312)^+$, and of two-peak structure of the $P_c(4450)^+$, *Phys. Rev. Lett.* **122**, no. 22, 222001 (2019).
- [89] F. L. Wang, X. D. Yang, R. Chen and X. Liu, Correlation of the hidden-charm molecular tetraquarks and the charmonium-like structures existing in the $B \rightarrow XYZ + K$, *arXiv:2103.04698*.
- [90] X. Liu and S. L. Zhu, $Y(4143)$ is probably a molecular partner of $Y(3930)$, *Phys. Rev. D* **80**, 017502 (2009).
- [91] L. Zhao, L. Ma and S. L. Zhu, The recoil correction and spin-orbit force for the possible $B^* \bar{B}^*$ and $D^* \bar{D}^*$ states, *Nucl. Phys. A* **942**, 18 (2015).
- [92] Y. R. Liu and Z. Y. Zhang, The Bound state problem of S-wave heavy quark meson-antimeson systems, *Phys. Rev. C* **80**, 015208 (2009).
- [93] Y. Liu and I. Zahed, Heavy Exotic Molecules with Charm and Bottom, *Phys. Lett. B* **762**, 362 (2016).
- [94] L. R. Dai, G. Y. Wang, X. Chen, E. Wang, E. Oset and D. M. Li, The $B^+ \rightarrow J/\psi \omega K^+$ reaction and $D^* \bar{D}^*$ molecular states, *Eur. Phys. J. A* **55**, 36 (2019).
- [95] Z. F. Sun, J. He, X. Liu, Z. G. Luo and S. L. Zhu, $Z_b(10610)^\pm$ and $Z_b(10650)^\pm$ as the $B^* \bar{B}$ and $B^* \bar{B}^*$ molecular states, *Phys. Rev. D* **84**, 054002 (2011).
- [96] M. Z. Liu, D. J. Jia and D. Y. Chen, Possible hadronic molecular states composed of S-wave heavy-light mesons, *Chin. Phys. C* **41**, 053105 (2017).
- [97] Y. C. Yang, Z. Y. Tan, J. Ping and H. S. Zong, Possible $D^{(*)} \bar{D}^{(*)}$ and $B^{(*)} \bar{B}^{(*)}$ molecular states in the extended constituent quark models, *Eur. Phys. J. C* **77**, 575 (2017).
- [98] Y. J. Zhang, H. C. Chiang, P. N. Shen and B. S. Zou, Possible S-wave bound-states of two pseudoscalar mesons, *Phys. Rev. D* **74**, 014013 (2006).
- [99] X. K. Dong, F. K. Guo and B. S. Zou, A survey of heavy-antiheavy hadronic molecules, *Progr. Phys.* **41**, 65 (2021).
- [100] Z. F. Sun, Z. G. Luo, J. He, X. Liu, and S. L. Zhu, A note on the $B^* \bar{B}$, $B^* \bar{B}^*$, $D^* \bar{D}$, $D^* \bar{D}^*$ molecular states, *Chin. Phys. C* **36**, 194 (2012).
- [101] E. Wang, H. S. Li, W. H. Liang and E. Oset, Analysis of the $\gamma\gamma \rightarrow D\bar{D}$ reaction and the $D\bar{D}$ bound state, *Phys. Rev. D* **103** (2021) no.5, 054008.
- [102] L. Dai, G. Toledo and E. Oset, Searching for a $D\bar{D}$ bound state with the $\psi(3770) \rightarrow \gamma D^0 \bar{D}^0$ decay, *Eur. Phys. J. C* **80** (2020) no.6, 510.
- [103] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmonium-like Structure in $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ at $\sqrt{s} = 4.26$ GeV, *Phys. Rev. Lett.* **110**, 252001 (2013).
- [104] B. Aubert *et al.* (BaBar Collaboration), Observation of $Y(3940) \rightarrow J/\psi \omega$ in $B \rightarrow J/\psi \omega K$ at BABAR, *Phys. Rev. Lett.* **101**, 082001 (2008).
- [105] S. Uehara *et al.* (Belle Collaboration), Observation of a χ'_{c2} candidate in $\gamma\gamma \rightarrow D\bar{D}$ production at BELLE, *Phys. Rev. Lett.* **96**, 082003 (2006).
- [106] K. Abe *et al.* (Belle Collaboration), Observation of a near-threshold $\omega J/\psi$ mass enhancement in exclusive $B \rightarrow K \omega J/\psi$ decays, *Phys. Rev. Lett.* **94**, 182002 (2005).
- [107] C. Z. Yuan *et al.* (Belle Collaboration), Measurement of $e^+e^- \rightarrow \pi^+\pi^- J/\psi$ cross-section via initial state radiation at Belle, *Phys. Rev. Lett.* **99**, 182004 (2007).
- [108] M. Ablikim *et al.* (BESIII Collaboration), Observation of a Charged Charmoniumlike Structure $Z_c(4020)$ and Search for the $Z_c(3900)$ in $e^+e^- \rightarrow \pi^+\pi^- h_c$, *Phys. Rev. Lett.* **111**, 242001 (2013).
- [109] M. Ablikim *et al.* (BESIII Collaboration), Observation of a charged charmoniumlike structure in $e^+e^- \rightarrow (D^* \bar{D}^*)^\pm \pi^\mp$ at $\sqrt{s} = 4.26$ GeV, *Phys. Rev. Lett.* **112**, 132001 (2014).
- [110] R. Mizuk *et al.* (Belle Collaboration), Observation of two resonance-like structures in the $\pi^+ \chi_{c1}$ mass distribution in exclusive $\bar{B}^0 \rightarrow K^- \pi^+ \chi_{c1}$ decays, *Phys. Rev. D* **78**, 072004 (2008).
- [111] F. Close and C. Downum, On the possibility of Deeply Bound Hadronic Molecules from single Pion Exchange, *Phys. Rev. Lett.* **102**, 242003 (2009).
- [112] M. T. Li, W. L. Wang, Y. B. Dong and Z. Y. Zhang, A Study of One S- and One P-Wave Heavy Meson Interaction in a Chiral Quark Model, *Commun. Theor. Phys.* **63**, 63 (2015).
- [113] M. T. Li, W. L. Wang, Y. B. Dong and Z. Y. Zhang, A Study of P-wave Heavy Meson Interactions in A Chiral Quark Model, *arXiv:1303.4140*.

- [114] F. Close, C. Downum and C. E. Thomas, Novel Charmonium and Bottomonium Spectroscopies due to Deeply Bound Hadronic Molecules from Single Pion Exchange, *Phys. Rev. D* **81**, 074033 (2010).
- [115] J. He and D. Y. Chen, Interpretation of $Y(4390)$ as an isoscalar partner of $Z(4430)$ from $D^*(2010)\bar{D}_1(2420)$ interaction, *Eur. Phys. J. C* **77**, 398 (2017).
- [116] W. Zhu, Y. R. Liu and T. Yao, Is $J^{PC} = 3^{-+}$ molecule possible?, *Chin. Phys. C* **39**, 023101 (2015).
- [117] B. Aubert *et al.* (BaBar Collaboration), Observation of a Broad Structure in the $\pi^+\pi^-J/\psi$ Mass Spectrum Around 4.26-GeV/ c^2 , *Phys. Rev. Lett.* **95**, 142001 (2005).
- [118] B. Aubert *et al.* (BaBar Collaboration), Evidence of a broad structure at an invariant mass of 4.32-GeV/ c^2 in the reaction $e^+e^- \rightarrow \pi^+\pi^-\psi_{2S}$ measured at BaBar, *Phys. Rev. Lett.* **98**, 212001 (2007).
- [119] M. Ablikim *et al.* (BESIII Collaboration), Evidence of Two Resonant Structures in $e^+e^- \rightarrow \pi^+\pi^-h_c$, *Phys. Rev. Lett.* **118**, 092002 (2017).
- [120] S. K. Choi *et al.* (Belle Collaboration), Observation of a Resonance-Like Structure in the $\pi^+\psi'$ Mass Distribution in Exclusive $B \rightarrow K\pi^+\psi'$ Decays, *Phys. Rev. Lett.* **100**, 142001 (2008).
- [121] L. L. Shen, X. L. Chen, Z. G. Luo, P. Z. Huang, S. L. Zhu, P. F. Yu, and X. Liu, The molecular systems composed of the charmed mesons in the $H\bar{S} + h.c.$ doublet, *Eur. Phys. J. C* **70**, 183 (2010).
- [122] X. Liu, Y. R. Liu, W. Z. Deng, and S. L. Zhu, Is $Z^+(4430)$ a loosely bound molecular state?, *Phys. Rev. D* **77**, 034003 (2008).
- [123] X. Liu, Y. R. Liu, W. Z. Deng, and S. L. Zhu, $Z^+(4430)$ as a $D_1^*D^*(D_1D^*)$ molecular state, *Phys. Rev. D* **77**, 094015 (2008).
- [124] R. Chen, X. Liu, Y. R. Liu, and S. L. Zhu, Predictions of the hidden-charm molecular states with four-quark component, *Eur. Phys. J. C* **76**, 319 (2016).
- [125] B. Hu, X. L. Chen, Z. G. Luo, P. Z. Huang, S. L. Zhu, P. F. Yu, and X. Liu, Possible heavy molecular states composed of a pair of excited charm-strange mesons, *Chin. Phys. C* **35**, 113 (2011).
- [126] F. L. Wang and X. Liu, Investigating new type of doubly charmed molecular tetraquarks composed of the charmed meson in H -doublet and the charmed meson in T -doublet, [arXiv:2108.09925](https://arxiv.org/abs/2108.09925).
- [127] M. B. Wise, Chiral perturbation theory for hadrons containing a heavy quark, *Phys. Rev. D* **45**, R2188 (1992).
- [128] H. X. Chen, W. Chen, X. Liu, Y. R. Liu and S. L. Zhu, A review of the open charm and open bottom systems, *Rept. Prog. Phys.* **80** (2017) no.7, 076201.
- [129] E. E. Kolomeitsev and M. F. M. Lutz, On Heavy light meson resonances and chiral symmetry, *Phys. Lett. B* **582** (2004), 39-48.
- [130] F. K. Guo, P. N. Shen and H. C. Chiang, Dynamically generated 1^+ heavy mesons, *Phys. Lett. B* **647** (2007), 133-139.
- [131] D. Gamermann and E. Oset, Axial resonances in the open and hidden charm sectors, *Eur. Phys. J. A* **33** (2007), 119-131.
- [132] R. Molina, H. Nagahiro, A. Hosaka and E. Oset, Scalar, axial-vector and tensor resonances from the ρD^* , ωD^* interaction in the hidden gauge formalism, *Phys. Rev. D* **80** (2009), 014025.
- [133] Y. R. Liu, Heavy quark spin selection rules in meson-antimeson states, *Phys. Rev. D* **88**, 074008 (2013).
- [134] H. Y. Cheng and K. C. Yang, Charmless hadronic B decays into a tensor meson, *Phys. Rev. D* **83**, 034001 (2011).
- [135] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Light vector resonances in the effective chiral Lagrangian for heavy mesons, *Phys. Lett. B* **292**, 371 (1992).
- [136] R. Casalbuoni, A. Deandrea, N. Di Bartolomeo, R. Gatto, F. Feruglio, and G. Nardulli, Phenomenology of heavy meson chiral Lagrangians, *Phys. Rep.* **281**, 145 (1997).
- [137] T. M. Yan, H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin, and H. L. Yu, Heavy quark symmetry and chiral dynamics, *Phys. Rev. D* **46**, 1148 (1992); [*Phys. Rev. D* **55**, 5851E (1997)].
- [138] M. Bando, T. Kugo, and K. Yamawaki, Nonlinear realization and hidden local symmetries, *Phys. Rep.* **164**, 217 (1988).
- [139] M. Harada and K. Yamawaki, Hidden local symmetry at loop: A new perspective of composite gauge boson and chiral phase transition, *Phys. Rep.* **381**, 1 (2003).
- [140] G. J. Ding, Are $Y(4260)$ and $Z_2^+(4250)$ D_1D or D_0D^* hadronic molecules? *Phys. Rev. D* **79**, 014001 (2009).
- [141] F. L. Wang, R. Chen, Z. W. Liu, and X. Liu, Probing new types of P_c states inspired by the interaction between S -wave charmed baryon and anti-charmed meson in a \bar{T} doublet, *Phys. Rev. C* **101**, 025201 (2020).
- [142] D. O. Riska and G. E. Brown, Nucleon resonance transition couplings to vector mesons, *Nucl. Phys. A* **679**, 577 (2001).
- [143] F. L. Wang, R. Chen, Z. W. Liu, and X. Liu, Possible triple-charm molecular pentaquarks from $\Xi_{cc}D_1/\Xi_{cc}D_2^*$ interactions, *Phys. Rev. D* **99**, 054021 (2019).
- [144] Y. R. Liu and M. Oka, $\Lambda_c N$ bound states revisited, *Phys. Rev. D* **85**, 014015 (2012).
- [145] P. A. Zyla *et al.* [Particle Data Group], Review of Particle Physics, *PTEP* **2020**, 083C01 (2020).
- [146] G. Breit, The effects of retardation on the interaction of two electrons, *Phys. Rev.* **34**, 553 (1929).
- [147] G. Breit, The fine structure of HE as a test of the spin interactions of two electrons, *Phys. Rev.* **36**, 383 (1930).
- [148] N. A. Tornqvist, On deusons or deuteron-like meson-meson bound states, *Nuovo Cimento Soc. Ital. Fis.* **107A**, 2471 (1994).
- [149] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, Chiral Lagrangians for Massive Spin 1 Fields, *Phys. Lett. B* **223** (1989), 425-432.
- [150] R. Chen, A. Hosaka and X. Liu, Heavy molecules and one- σ/ω -exchange model, *Phys. Rev. D* **96** (2017) no.11, 116012.
- [151] D. Y. Chen, X. Liu and T. Matsuki, η transitions between charmonia with meson loop contributions, *Phys. Rev. D* **87** (2013) no.5, 054006.
- [152] R. Chen, J. He and X. Liu, Possible strange hidden-charm pentaquarks from $\Sigma_c^{(*)}\bar{D}_s^*$ and $\Xi_c^{(*)}\bar{D}^*$ interactions, *Chin. Phys. C* **41** (2017) no.10, 103105.
- [153] R. Aaij *et al.* (LHCb Collaboration), Physics case for an LHCb Upgrade II-Opportunities in flavour physics, and beyond, in the HL-LHC era, [arXiv:1808.08865](https://arxiv.org/abs/1808.08865).