

Towards NNLO corrections to muon-electron scattering

Syed Mehedi Hasan^{1*}

1 INFN, Sezione di Pavia, Via A. Bassi 6, 27100 Pavia, Italy
 * syedmehe@pv.infn.it

November 2, 2021



*15th International Symposium on Radiative Corrections:
 Applications of Quantum Field Theory to Phenomenology,
 FSU, Tallahassee, FL, USA, 17-21 May 2021
 doi:10.21468/SciPostPhysProc.?*

Abstract

The recently proposed MUonE experiment at CERN aims at providing a novel determination of the leading order hadronic contribution to the muon anomalous magnetic moment through the study of elastic muon-electron scattering at relatively small momentum transfer. The anticipated accuracy of the order of 10 ppm demands for high-precision predictions, including all the relevant radiative corrections. To aid the effort, the theoretical formulation for the fixed order NNLO QED corrections are described and the virtual NNLO leptonic corrections are calculated with complete mass effects.¹

Contents

1	Introduction	2
2	NNLO photonic corrections to muon-electron scattering	3
3	Vertex and box-like irreducible double virtual NNLO leptonic corrections to muon-electron scattering	4
4	Conclusion	6
References		8

¹* speaker

1 Introduction

The anomalous magnetic moment of the muon, $a_\mu = (g - 2)_\mu / 2$, is a fundamental quantity in particle physics. This quantity has been recently measured by the Fermilab Muon $g - 2$ Experiment (E989) with relative precision of 0.46 ppm [1] for positive muons, which is an improvement with respect to the 0.54 ppm precision obtained by the E821 experiment at the Brookhaven National Laboratory [2].

The muon anomaly is due to loop corrections coming from the QED, weak and strong sector of the Standard Model (SM) [3, 4]. There is a long-standing puzzle that the discrepancy between the measured value and theoretical value of $g - 2$ currently exceeds 3σ level. Moreover it grows up to 4.2σ when combined with previous measurements using both μ^+ and μ^- beams [1]. The present status of the SM theoretical prediction has been recently reviewed in ref. [5].

The E34 experiment, which is under development at J-PARC [6, 7] and future runs of the E989 experiment at Fermilab [8, 9], are expected to bring the relative precision below the level of 0.2 ppm. Naturally a major effort from the theory side is called for to reduce the uncertainty in the SM prediction, most of which comes from the non-perturbative strong interaction effects as given by the leading order hadronic correction, a_μ^{HLO} , and the hadronic light-by-light contribution. There is at present general consensus that the latter has a small impact on the theoretical uncertainty on a_μ [5].

The traditional computation of a_μ^{HLO} is performed via dispersion integral of the hadron production cross section in electron-positron annihilation at low energies [5, 10–12]. An alternative evaluation of the a_μ^{HLO} can be carried out using Lattice QCD calculation [13–26]. Recently the calculation carried out by the BMW collaboration shows a more precise determination of a_μ^{HLO} [27] with a central value larger than the dispersive approach and in closer agreement with BNL and FNAL [1, 2] experiment outcome. To resolve the tension, alternative approaches for the evaluation of a_μ^{HLO} are more than welcome.

A novel approach to derive a_μ^{HLO} from a measurement of the effective electromagnetic coupling constant in the space-like region via scattering data, has been proposed in ref. [28]. The elastic scattering of high-energy muons on atomic electrons has been identified as an ideal process [29] and as a result a new experiment, MUonE, has been proposed at CERN [30–33]. This new determination of a_μ^{HLO} can be competitive with the dispersive approach, if the uncertainty in the measurement of the μe differential cross section is of the order of 10 ppm. This challenge of MUonE requires very high-precision predictions for μe scattering, which includes all the relevant radiative corrections, implemented in fully-fledged Monte Carlo (MC) tools. The necessary perturbative accuracy is at next-to-next-to leading order in QED (NNLO QED), combined with the leading logarithmic contributions due to multiple photon radiation. A comprehensive review of the recent developments along this line, as of the beginning of 2020, can be found in ref. [34]. The complete calculation at NLO accuracy and the evaluation of its phenomenological impact on MUonE observables can be found in ref. [35]. At NNLO accuracy there has been significant progress recently [36–46].²

In the following we summarize our effort along NNLO photonic corrections and NNLO leptonic corrections to muon-electron scattering, which are the subject matter of ref. [40] and ref. [46] respectively.

²It is worth reminding the studies of refs. [47, 48] which investigate the robustness of the MUonE measurements to possible New Physics effects

2 NNLO photonic corrections to muon-electron scattering

We define NNLO photonic corrections by removing the contribution due to the fermionic loop (for example, in Figure 1, diagram 1d is removed), which forms a gauge invariant subset. Further justification is shown in favour of a stand-alone analysis for this type of contribution in ref. [40]. All the calculations are exact except the double-box like diagrams, where YFS inspired approximation [49] is used to capture the correct IR structure. In other words, the approximation misses only the non-IR remnant of the two-loop diagrams where at least two photons connect the electron and muon lines.




Figure 1: Virtual QED corrections to the electron line in $\mu e \rightarrow \mu e$ scattering. One-loop correction (diagram 1a); sample topologies for the two-loop corrections (diagrams 1b-1d). The blob in diagram 1d denotes an electron loop insertion. On-shell scheme counterterms are understood.




Figure 2: Sample diagrams for the one-loop QED corrections to single photon emission (diagrams 2a-2b); sample diagrams for the double bremsstrahlung process (diagrams 2c-2d).

We studied the numerical impact of QED NNLO photonic corrections to observables assuming the following event selections (Setup 2 and Setup 4 of Ref. [35]):

1. $\theta_e, \theta_\mu < 100$ mrad and $E_e > 1$ GeV (i.e. $t_{ee} \lesssim -1.02 \cdot 10^{-3}$ GeV 2). The angular cuts model the typical acceptance conditions of the experiment and the electron energy threshold is imposed to guarantee the presence of two charged tracks in the detector (Setup 1);

2. the same criteria as above, with the additional acoplanarity cut $|\pi - |\phi_e - \phi_\mu|| \leq 3.5$ mrad. This event selection is considered in order to mimic an experimental cut which allows to stay close to the elasticity curve given by the tree-level relation between the electron and muon scattering angles (Setup 2).

The input parameters for the simulations are set to

$$\alpha = 1/137.03599907430637 \quad m_e = 0.510998928 \text{ MeV} \quad m_\mu = 105.6583715 \text{ MeV}$$

and the considered differential observables are the following ones:

$$\frac{d\sigma}{dt_{ee}}, \quad \frac{d\sigma}{dt_{\mu\mu}}, \quad \frac{d\sigma}{d\theta_e}, \quad \frac{d\sigma}{d\theta_\mu}, \quad \frac{d\sigma}{dE_e}, \quad \frac{d\sigma}{dE_\mu}, \quad (1)$$

where $t_{ee} = (p_2 - p_4)^2$, $t_{\mu\mu} = (p_1 - p_3)^2$, (θ_e, E_e) and (θ_μ, E_μ) are the scattering angle and the energy, in the laboratory frame, of the outgoing electron and muon, respectively.

At this point it is useful to define two quantities as follows:

$$\Delta_{\text{NNLO}}^i = 100 \times \frac{d\sigma_{\text{NNLO}}^i - d\sigma_{\text{NLO}}^i}{d\sigma_{\text{LO}}}, \quad (2)$$

where $i = e, \mu, f$ implies electron-line only, muon-line only and the full set of NNLO corrections. Similarly we define:

$$\Delta_{\text{NLO}}^i = 100 \times \frac{d\sigma_{\text{NLO}}^i - d\sigma_{\text{LO}}}{d\sigma_{\text{LO}}}. \quad (3)$$

Fig. 3 shows relative contribution, differential in the squared momentum transfer measured along the electron line, of the exact NNLO corrections from one leptonic line. Whereas Fig. 4 shows the relative contribution of the full(approximate) NNLO corrections. Similar analysis are performed with respect to the other listed observables [40].

3 Vertex and box-like irreducible double virtual NNLO leptonic corrections to muon-electron scattering

The calculation of the complete fixed-order NNLO QED corrections which include at least one leptonic pair, with all the relevant virtual and real contributions, summing over all contributing lepton flavours, is carried out in ref. [46]. We name this class of contribution as the NNLO leptonic corrections. Here we only report the vertex and box-like irreducible double virtual case, represented by Fig. 5 and 6 respectively.

We use dispersion relation (DR) technique to calculate the above mentioned contributions. An amplitude involving a photon in a loop with a vacuum polarisation (VP) insertion, due to lepton ℓ , can be extracted by replacing the photon propagator [50–55] as follows

$$\frac{-ig_{\mu\nu}}{q^2 + i\epsilon} \rightarrow \frac{-ig_{\mu\delta}}{q^2 + i\epsilon} i(q^2 g^{\delta\lambda} - q^\delta q^\lambda) \Pi_\ell(q^2) \frac{-ig_{\lambda\nu}}{q^2 + i\epsilon}, \quad (4)$$

where the renormalised $\Pi(q^2)$ is obtained from its imaginary part using the subtracted DR

$$\Pi_\ell(q^2) = -\frac{q^2}{\pi} \int_{4m_\ell^2}^{\infty} \frac{dz}{z} \frac{\text{Im } \Pi_\ell(z)}{q^2 - z + i\epsilon} \quad (5)$$




Figure 3: Red histograms: NNLO corrections, according to Eq. (2), along the electron line, for the $\mu^\pm e^- \rightarrow \mu^\pm e^-$ process, as a function of the squared momentum transfer measured along the electron line, t_{ee} ; blue histogram: $10 \times$ NNLO corrections along the muon line. Left panel: differential distribution in the presence of acceptance cuts only (Setup 1); right panel: the same as in the left panel with the additional acoplanarity cut of Setup 2. The insets report the NLO corrections, according to Eq. (3); the blue line represents directly Eq. (3) without any multiplicative factor.

where

$$\text{Im}\Pi_\ell(z) = -\frac{\alpha}{3}R_\ell(z), \quad (6)$$

$$R_\ell(z) = \left(1 + \frac{4m_\ell^2}{2z}\right)\sqrt{1 - \frac{4m_\ell^2}{z}}, \quad (7)$$

m_ℓ representing the mass of the lepton blob. The $q^\delta q^\lambda$ term in eq. (4) does not contribute because of gauge invariance. Hence the two-loop calculation can be performed starting from the one-loop amplitude with the replacement

$$\frac{-ig_{\mu\nu}}{q^2 + i\epsilon} \rightarrow -ig_{\mu\nu}\left(\frac{\alpha}{3\pi}\right)\int_{4m_\ell^2}^{\infty} \frac{dz}{z} \frac{1}{q^2 - z + i\epsilon} \left(1 + \frac{4m_\ell^2}{2z}\right) \sqrt{1 - \frac{4m_\ell^2}{z}}. \quad (8)$$

Furthermore, we have analytically evaluated the vertex form factors (Fig. 5) by evaluating master integrals (MI) and found agreement with the DR method.

Seven master integrals, depicted in figure 7, are evaluated. We used differential equation methods [56–58] and chose a suitable basis such that the dimensional regularisation parameter




Figure 4: Approximation (based on YFS) to the full NNLO corrections as a function of the squared momentum transfer measured along the electron line, t_{ee} . Green histogram refers to the process $\mu^- e^- \rightarrow \mu^- e^-$ while the violet histogram refers to $\mu^+ e^- \rightarrow \mu^+ e^-$. Left panel: differential distribution in the presence of acceptance cuts only (Setup 1); right panel: the same as in the left panel with the additional acoplanarity cut of Setup 2. The insets report the NLO corrections, according to Eq. (3).

ϵ factorises from the kinematics. These integrals were then encoded in a *dlog*-form, also called canonical form [59], which we obtained using the Magnus algorithm [60, 61]. Finally the master integrals are written in terms of the generalised polylogarithms [62–64]. The explicit results are available in the ancillary files of the arXiv version of ref. [46].

4 Conclusion

We have analyzed the NNLO photonic corrections to $\mu^\pm e^- \rightarrow \mu^\pm e^-$ scattering. The phenomenological results are obtained with a related MC code, named MESMER. The motivation of the study is due to the recent proposal of measuring the effective electromagnetic coupling constant in the space-like region by using this particular process (MUonE experiment). From the measurement of the hadronic contribution to the running of α_{QED} , the leading hadronic contribution to the muon anomaly can be derived as an alternative approach to the standard time-like evaluation of a_μ^{HLO} .

The NNLO leptonic virtual contribution have been calculated with dispersion relation techniques including all finite mass effects. For the class of vertex corrections, the numerical results have been cross-checked with exact analytical result, valid for arbitrary internal and external masses, finding excellent agreement.




Figure 5: NNLO irreducible vertex diagrams. Vertex correction along the electron line (left) and along the muon line (right).




Figure 6: NNLO Box diagrams. Direct diagrams (a)-(b) and crossed diagrams (c)-(d).




Figure 7: The master integral topologies $\mathcal{T}_{1\dots 7}$ for fermionic loop induced NNLO vertex correction. The solid single line represents a propagator with mass m_1 and the double line represents a propagator with mass m_2 , while the dashed line represents a massless propagator with momentum squared equal to t . The dots represent additional powers of the propagator.

Acknowledgements

We are grateful to the MUonE colleagues for many useful discussions. SMH acknowledges INFN Sezione di Pavia for providing the necessary funding and hospitality to carry out the research.

References

- [1] B. Abi *et al.*, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, Phys. Rev. Lett. **126**(14), 141801 (2021), doi:[10.1103/PhysRevLett.126.141801](https://doi.org/10.1103/PhysRevLett.126.141801), 2104.03281.
- [2] G. W. Bennett *et al.*, *Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL*, Phys. Rev. **D73**, 072003 (2006), doi:[10.1103/PhysRevD.73.072003](https://doi.org/10.1103/PhysRevD.73.072003), hep-ex/0602035.
- [3] F. Jegerlehner and A. Nyffeler, *The Muon g-2*, Phys. Rept. **477**, 1 (2009), doi:[10.1016/j.physrep.2009.04.003](https://doi.org/10.1016/j.physrep.2009.04.003), 0902.3360.
- [4] F. Jegerlehner, *The Anomalous Magnetic Moment of the Muon*, Springer Tracts Mod. Phys. **274**, pp. 1 (2017), doi:[10.1007/978-3-319-63577-4](https://doi.org/10.1007/978-3-319-63577-4).
- [5] T. Aoyama *et al.*, *The anomalous magnetic moment of the muon in the Standard Model*, Phys. Rept. **887**, 1 (2020), doi:[10.1016/j.physrep.2020.07.006](https://doi.org/10.1016/j.physrep.2020.07.006), 2006.04822.
- [6] H. Iinuma, *New approach to the muon g-2 and EDM experiment at J-PARC*, J. Phys. Conf. Ser. **295**, 012032 (2011), doi:[10.1088/1742-6596/295/1/012032](https://doi.org/10.1088/1742-6596/295/1/012032).
- [7] T. Mibe, *Measurement of muon g-2 and EDM with an ultra-cold muon beam at J-PARC*, Nucl. Phys. Proc. Suppl. **218**, 242 (2011), doi:[10.1016/j.nuclphysbps.2011.06.039](https://doi.org/10.1016/j.nuclphysbps.2011.06.039).
- [8] G. Venanzoni, *The New Muon g-2 experiment at Fermilab*, Nucl. Part. Phys. Proc. **273-275**, 584 (2016), doi:[10.1016/j.nuclphysbps.2015.09.087](https://doi.org/10.1016/j.nuclphysbps.2015.09.087), 1411.2555.
- [9] J. Grange *et al.*, *Muon (g-2) Technical Design Report* (2015), [1501.06858](https://arxiv.org/abs/1501.06858).
- [10] F. Jegerlehner, *The Muon g-2 in Progress*, Acta Phys. Polon. **B49**, 1157 (2018), doi:[10.5506/APhysPolB.49.1157](https://doi.org/10.5506/APhysPolB.49.1157), 1804.07409.
- [11] A. Keshavarzi, D. Nomura and T. Teubner, *The muon $g - 2$ and $\alpha(M_Z^2)$: a new data-based analysis*, Phys. Rev. **D97**(11), 114025 (2018), doi:[10.1103/PhysRevD.97.114025](https://doi.org/10.1103/PhysRevD.97.114025), 1802.02995.
- [12] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, *A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to $\alpha(m_Z^2)$* , Eur. Phys. J. C **80**(3), 241 (2020), doi:[10.1140/epjc/s10052-020-7792-2](https://doi.org/10.1140/epjc/s10052-020-7792-2), [Erratum: Eur.Phys.J.C 80, 410 (2020)], [1908.00921](https://arxiv.org/abs/1908.00921).
- [13] M. Della Morte, A. Francis, V. Gülpers, G. Herdoiza, G. von Hippel, H. Horch, B. Jäger, H. B. Meyer, A. Nyffeler and H. Wittig, *The hadronic vacuum polarization contribution to the muon $g-2$ from lattice QCD*, JHEP **10**, 020 (2017), doi:[10.1007/JHEP10\(2017\)020](https://doi.org/10.1007/JHEP10(2017)020), [1705.01775](https://arxiv.org/abs/1705.01775).

- [14] B. Chakraborty *et al.*, *Strong-Isospin-Breaking Correction to the Muon Anomalous Magnetic Moment from Lattice QCD at the Physical Point*, Phys. Rev. Lett. **120**(15), 152001 (2018), doi:[10.1103/PhysRevLett.120.152001](https://doi.org/10.1103/PhysRevLett.120.152001), [1710.11212](https://arxiv.org/abs/1710.11212).
- [15] S. Borsanyi *et al.*, *Hadronic vacuum polarization contribution to the anomalous magnetic moments of leptons from first principles*, Phys. Rev. Lett. **121**(2), 022002 (2018), doi:[10.1103/PhysRevLett.121.022002](https://doi.org/10.1103/PhysRevLett.121.022002), [1711.04980](https://arxiv.org/abs/1711.04980).
- [16] H. B. Meyer and H. Wittig, *Lattice QCD and the anomalous magnetic moment of the muon*, Prog. Part. Nucl. Phys. **104**, 46 (2019), doi:[10.1016/j.ppnp.2018.09.001](https://doi.org/10.1016/j.ppnp.2018.09.001), [1807.09370](https://arxiv.org/abs/1807.09370).
- [17] T. Blum, P. A. Boyle, V. Gülpers, T. Izubuchi, L. Jin, C. Jung, A. Jüttner, C. Lehner, A. Portelli and J. T. Tsang, *Calculation of the hadronic vacuum polarization contribution to the muon anomalous magnetic moment*, Phys. Rev. Lett. **121**(2), 022003 (2018), doi:[10.1103/PhysRevLett.121.022003](https://doi.org/10.1103/PhysRevLett.121.022003), [1801.07224](https://arxiv.org/abs/1801.07224).
- [18] D. Giusti, F. Sanfilippo and S. Simula, *Light-quark contribution to the leading hadronic vacuum polarization term of the muon $g - 2$ from twisted-mass fermions*, Phys. Rev. D **98**(11), 114504 (2018), doi:[10.1103/PhysRevD.98.114504](https://doi.org/10.1103/PhysRevD.98.114504), [1808.00887](https://arxiv.org/abs/1808.00887).
- [19] D. Giusti, V. Lubicz, G. Martinelli, F. Sanfilippo and S. Simula, *Electromagnetic and strong isospin-breaking corrections to the muon $g - 2$ from Lattice QCD+QED*, Phys. Rev. D **99**(11), 114502 (2019), doi:[10.1103/PhysRevD.99.114502](https://doi.org/10.1103/PhysRevD.99.114502), [1901.10462](https://arxiv.org/abs/1901.10462).
- [20] D. Giusti and S. Simula, *Lepton anomalous magnetic moments in Lattice QCD+QED*, PoS **LATTICE2019**, 104 (2019), doi:[10.22323/1.363.0104](https://doi.org/10.22323/1.363.0104), [1910.03874](https://arxiv.org/abs/1910.03874).
- [21] E. Shintani and Y. Kuramashi, *Hadronic vacuum polarization contribution to the muon $g - 2$ with 2+1 flavor lattice QCD on a larger than $(10 \text{ fm})^4$ lattice at the physical point*, Phys. Rev. D **100**(3), 034517 (2019), doi:[10.1103/PhysRevD.100.034517](https://doi.org/10.1103/PhysRevD.100.034517), [1902.00885](https://arxiv.org/abs/1902.00885).
- [22] C. T. H. Davies *et al.*, *Hadronic-vacuum-polarization contribution to the muon's anomalous magnetic moment from four-flavor lattice QCD*, Phys. Rev. D **101**(3), 034512 (2020), doi:[10.1103/PhysRevD.101.034512](https://doi.org/10.1103/PhysRevD.101.034512), [1902.04223](https://arxiv.org/abs/1902.04223).
- [23] A. Gérardin, M. Cè, G. von Hippel, B. Hörrz, H. B. Meyer, D. Mohler, K. Ott nad, J. Wilhelm and H. Wittig, *The leading hadronic contribution to $(g - 2)_\mu$ from lattice QCD with $N_f = 2 + 1$ flavours of $O(a)$ improved Wilson quarks*, Phys. Rev. D **100**(1), 014510 (2019), doi:[10.1103/PhysRevD.100.014510](https://doi.org/10.1103/PhysRevD.100.014510), [1904.03120](https://arxiv.org/abs/1904.03120).
- [24] C. Aubin, T. Blum, C. Tu, M. Golterman, C. Jung and S. Peris, *Light quark vacuum polarization at the physical point and contribution to the muon $g - 2$* , Phys. Rev. D **101**(1), 014503 (2020), doi:[10.1103/PhysRevD.101.014503](https://doi.org/10.1103/PhysRevD.101.014503), [1905.09307](https://arxiv.org/abs/1905.09307).
- [25] D. Giusti and S. Simula, *Ratios of the hadronic contributions to the lepton $g - 2$ from Lattice QCD+QED simulations*, Phys. Rev. D **102**(5), 054503 (2020), doi:[10.1103/PhysRevD.102.054503](https://doi.org/10.1103/PhysRevD.102.054503), [2003.12086](https://arxiv.org/abs/2003.12086).
- [26] C. Lehner and A. S. Meyer, *Consistency of hadronic vacuum polarization between lattice QCD and the R-ratio*, Phys. Rev. D **101**, 074515 (2020), doi:[10.1103/PhysRevD.101.074515](https://doi.org/10.1103/PhysRevD.101.074515), [2003.04177](https://arxiv.org/abs/2003.04177).

- [27] S. Borsanyi *et al.*, *Leading hadronic contribution to the muon magnetic moment from lattice QCD*, Nature **593**(7857), 51 (2021), doi:[10.1038/s41586-021-03418-1](https://doi.org/10.1038/s41586-021-03418-1), [2002.12347](https://arxiv.org/abs/2002.12347).
- [28] C. M. Carloni Calame, M. Passera, L. Trentadue and G. Venanzoni, *A new approach to evaluate the leading hadronic corrections to the muon g-2*, Phys. Lett. **B746**, 325 (2015), doi:[10.1016/j.physletb.2015.05.020](https://doi.org/10.1016/j.physletb.2015.05.020), [1504.02228](https://arxiv.org/abs/1504.02228).
- [29] G. Abbiendi *et al.*, *Measuring the leading hadronic contribution to the muon g-2 via μe scattering*, Eur. Phys. J. **C77**(3), 139 (2017), doi:[10.1140/epjc/s10052-017-4633-z](https://doi.org/10.1140/epjc/s10052-017-4633-z), [1609.08987](https://arxiv.org/abs/1609.08987).
- [30] G. Abbiendi *et al.*, *Letter of Intent: the MUonE project*, Tech. Rep. CERN-SPSC-2019-026, SPSC-I-252, CERN, Geneva (2019).
- [31] G. Abbiendi *et al.*, *Results on multiple Coulomb scattering from 12 and 20 GeV electrons on carbon targets*, JINST **15**(01), 01 (2020), doi:[10.1088/1748-0221/15/01/P01017](https://doi.org/10.1088/1748-0221/15/01/P01017), [1905.11677](https://arxiv.org/abs/1905.11677).
- [32] G. Ballerini *et al.*, *A feasibility test run for the MUonE project*, Nucl. Instrum. Meth. A **936**, 636 (2019), doi:[10.1016/j.nima.2018.10.148](https://doi.org/10.1016/j.nima.2018.10.148).
- [33] G. Abbiendi *et al.*, *A study of muon-electron elastic scattering in a test beam*, JINST **16**(06), P06005 (2021), doi:[10.1088/1748-0221/16/06/P06005](https://doi.org/10.1088/1748-0221/16/06/P06005), [2102.11111](https://arxiv.org/abs/2102.11111).
- [34] P. Banerjee *et al.*, *Theory for muon-electron scattering @ 10 ppm: A report of the MUonE theory initiative*, Eur. Phys. J. C **80**(6), 591 (2020), doi:[10.1140/epjc/s10052-020-8138-9](https://doi.org/10.1140/epjc/s10052-020-8138-9), [2004.13663](https://arxiv.org/abs/2004.13663).
- [35] M. Alacevich, C. M. Carloni Calame, M. Chiesa, G. Montagna, O. Nicrosini and F. Piccinini, *Muon-electron scattering at NLO*, JHEP **02**, 155 (2019), doi:[10.1007/JHEP02\(2019\)155](https://doi.org/10.1007/JHEP02(2019)155), [1811.06743](https://arxiv.org/abs/1811.06743).
- [36] P. Mastrolia, M. Passera, A. Primo and U. Schubert, *Master integrals for the NNLO virtual corrections to μe scattering in QED: the planar graphs*, JHEP **11**, 198 (2017), doi:[10.1007/JHEP11\(2017\)198](https://doi.org/10.1007/JHEP11(2017)198), [1709.07435](https://arxiv.org/abs/1709.07435).
- [37] S. Di Vita, S. Laporta, P. Mastrolia, A. Primo and U. Schubert, *Master integrals for the NNLO virtual corrections to μe scattering in QED: the non-planar graphs*, JHEP **09**, 016 (2018), doi:[10.1007/JHEP09\(2018\)016](https://doi.org/10.1007/JHEP09(2018)016), [1806.08241](https://arxiv.org/abs/1806.08241).
- [38] T. Engel, C. Gnendiger, A. Signer and Y. Ulrich, *Small-mass effects in heavy-to-light form factors*, JHEP **02**, 118 (2019), doi:[10.1007/JHEP02\(2019\)118](https://doi.org/10.1007/JHEP02(2019)118), [1811.06461](https://arxiv.org/abs/1811.06461).
- [39] T. Engel, A. Signer and Y. Ulrich, *A subtraction scheme for massive QED*, JHEP **01**, 085 (2020), doi:[10.1007/JHEP01\(2020\)085](https://doi.org/10.1007/JHEP01(2020)085), [1909.10244](https://arxiv.org/abs/1909.10244).
- [40] C. M. Carloni Calame, M. Chiesa, S. M. Hasan, G. Montagna, O. Nicrosini and F. Piccinini, *Towards muon-electron scattering at NNLO*, JHEP **11**, 028 (2020), doi:[10.1007/JHEP11\(2020\)028](https://doi.org/10.1007/JHEP11(2020)028), [2007.01586](https://arxiv.org/abs/2007.01586).
- [41] P. Banerjee, T. Engel, A. Signer and Y. Ulrich, *QED at NNLO with McMule*, SciPost Phys. **9**, 027 (2020), doi:[10.21468/SciPostPhys.9.2.027](https://doi.org/10.21468/SciPostPhys.9.2.027), [2007.01654](https://arxiv.org/abs/2007.01654).

- [42] R. Bonciani *et al.*, *The two-loop four-fermion scattering amplitude in QED* (2021), [2106.13179](#).
- [43] M. Fael, *Hadronic corrections to μ -e scattering at NNLO with space-like data*, JHEP **02**, 027 (2019), doi:[10.1007/JHEP02\(2019\)027](https://doi.org/10.1007/JHEP02(2019)027), [1808.08233](#).
- [44] M. Fael and M. Passera, *Muon-Electron Scattering at Next-To-Next-To-Leading Order: The Hadronic Corrections*, Phys. Rev. Lett. **122**(19), 192001 (2019), doi:[10.1103/PhysRevLett.122.192001](https://doi.org/10.1103/PhysRevLett.122.192001), [1901.03106](#).
- [45] M. Heller, *Planar two-loop integrals for μ e scattering in QED with finite lepton masses* (2021), [2105.08046](#).
- [46] E. Budassi, C. M. C. Calame, M. Chiesa, C. L. Del Pio, S. M. Hasan, G. Montagna, O. Nicrosini and F. Piccinini, *NNLO virtual and real leptonic corrections to muon-electron scattering* (2021), [2109.14606](#).
- [47] P. S. B. Dev, W. Rodejohann, X.-J. Xu and Y. Zhang, *MUonE sensitivity to new physics explanations of the muon anomalous magnetic moment*, JHEP **05**, 053 (2020), doi:[10.1007/JHEP05\(2020\)053](https://doi.org/10.1007/JHEP05(2020)053), [2002.04822](#).
- [48] A. Masiero, P. Paradisi and M. Passera, *New physics at the MUonE experiment at CERN*, Phys. Rev. D **102**(7), 075013 (2020), doi:[10.1103/PhysRevD.102.075013](https://doi.org/10.1103/PhysRevD.102.075013), [2002.05418](#).
- [49] D. R. Yennie, S. C. Frautschi and H. Suura, *The infrared divergence phenomena and high-energy processes*, Annals Phys. **13**, 379 (1961), doi:[10.1016/0003-4916\(61\)90151-8](https://doi.org/10.1016/0003-4916(61)90151-8).
- [50] A. H. Hoang, J. H. Kuhn and T. Teubner, *Radiation of light fermions in heavy fermion production*, Nucl. Phys. B **452**, 173 (1995), doi:[10.1016/0550-3213\(95\)00308-F](https://doi.org/10.1016/0550-3213(95)00308-F), [hep-ph/9505262](#).
- [51] S. Actis, M. Czakon, J. Gluza and T. Riemann, *Virtual hadronic and leptonic contributions to Bhabha scattering*, Phys. Rev. Lett. **100**, 131602 (2008), doi:[10.1103/PhysRevLett.100.131602](https://doi.org/10.1103/PhysRevLett.100.131602), [0711.3847](#).
- [52] S. Actis, J. Gluza and T. Riemann, *Virtual Hadronic Corrections to Massive Bhabha Scattering*, Nucl. Phys. B Proc. Suppl. **183**, 174 (2008), doi:[10.1016/j.nuclphysbps.2008.09.100](https://doi.org/10.1016/j.nuclphysbps.2008.09.100), [0807.0174](#).
- [53] J. H. Kuhn and S. Uccirati, *Two-loop QED hadronic corrections to Bhabha scattering*, Nucl. Phys. B **806**, 300 (2009), doi:[10.1016/j.nuclphysb.2008.08.002](https://doi.org/10.1016/j.nuclphysb.2008.08.002), [0807.1284](#).
- [54] R. Barbieri, J. A. Mignaco and E. Remiddi, *Electron form-factors up to fourth order. 1.*, Nuovo Cim. A **11**, 824 (1972), doi:[10.1007/BF02728545](https://doi.org/10.1007/BF02728545).
- [55] R. Barbieri, J. A. Mignaco and E. Remiddi, *Electron form factors up to fourth order. 2.*, Nuovo Cim. A **11**, 865 (1972), doi:[10.1007/BF02728546](https://doi.org/10.1007/BF02728546).
- [56] A. V. Kotikov, *Differential equations method: New technique for massive Feynman diagrams calculation*, Phys. Lett. B **254**, 158 (1991), doi:[10.1016/0370-2693\(91\)90413-K](https://doi.org/10.1016/0370-2693(91)90413-K).
- [57] E. Remiddi, *Differential equations for Feynman graph amplitudes*, Nuovo Cim. A **110**, 1435 (1997), [hep-th/9711188](#).

- [58] T. Gehrmann and E. Remiddi, *Differential equations for two loop four point functions*, Nucl. Phys. B **580**, 485 (2000), doi:[10.1016/S0550-3213\(00\)00223-6](https://doi.org/10.1016/S0550-3213(00)00223-6), hep-ph/[9912329](https://arxiv.org/abs/hep-ph/9912329).
- [59] J. M. Henn, *Multiloop integrals in dimensional regularization made simple*, Phys. Rev. Lett. **110**, 251601 (2013), doi:[10.1103/PhysRevLett.110.251601](https://doi.org/10.1103/PhysRevLett.110.251601), [1304.1806](https://arxiv.org/abs/1304.1806).
- [60] S. Di Vita, P. Mastrolia, U. Schubert and V. Yundin, *Three-loop master integrals for ladder-box diagrams with one massive leg*, JHEP **09**, 148 (2014), doi:[10.1007/JHEP09\(2014\)148](https://doi.org/10.1007/JHEP09(2014)148), [1408.3107](https://arxiv.org/abs/1408.3107).
- [61] M. Argeri, S. Di Vita, P. Mastrolia, E. Mirabella, J. Schlenk, U. Schubert and L. Tancredi, *Magnus and Dyson Series for Master Integrals*, JHEP **03**, 082 (2014), doi:[10.1007/JHEP03\(2014\)082](https://doi.org/10.1007/JHEP03(2014)082), [1401.2979](https://arxiv.org/abs/1401.2979).
- [62] E. Remiddi and J. A. M. Vermaasen, *Harmonic polylogarithms*, Int. J. Mod. Phys. A **15**, 725 (2000), doi:[10.1142/S0217751X00000367](https://doi.org/10.1142/S0217751X00000367), hep-ph/[9905237](https://arxiv.org/abs/hep-ph/9905237).
- [63] T. Gehrmann and E. Remiddi, *Numerical evaluation of two-dimensional harmonic polylogarithms*, Comput. Phys. Commun. **144**, 200 (2002), doi:[10.1016/S0010-4655\(02\)00139-X](https://doi.org/10.1016/S0010-4655(02)00139-X), hep-ph/[0111255](https://arxiv.org/abs/hep-ph/0111255).
- [64] J. Vollinga and S. Weinzierl, *Numerical evaluation of multiple polylogarithms*, Comput. Phys. Commun. **167**, 177 (2005), doi:[10.1016/j.cpc.2004.12.009](https://doi.org/10.1016/j.cpc.2004.12.009), hep-ph/[0410259](https://arxiv.org/abs/hep-ph/0410259).