Experimental investigation of ground-state properties of ⁷H with transfer reactions

M. Caamaño^{a,*}, T. Roger^b, A. M. Moro^c, G. F. Grinyer^{b,1}, J. Pancin^b, S. Bagchi^{d,2}, S. Sambi^e, J. Gibelin^f,
B. Fernández–Domínguez^a, N. Itagaki^g, J. Benlliure^a, D. Cortina–Gil^a, F. Farget^{b,3}, B. Jacquot^b,
D. Pérez–Loureiro^{b,4}, B. Pietras^a, R. Raabe^e, D. Ramos^{a,5}, C. Rodríguez Tajes^b, H. Savajols^b, M. Vandebrouck^{h,6}

^aIGFAE – Universidade de Santiago de Compostela, E-15706 Santiago de Compostela, Spain

^bGANIL, CEA/DSM–CNRS/IN2P3, BP 55027, F–14076 Caen Cedex 5, France

^eInstituut voor Kernen Stralingsfysica, KU Leuven, B–3001 Leuven, Belgium

^fLPC Caen, Université de Caen Basse–Normandie–ENSICAEN–CNRS/IN2P3, F–14050 Caen Cedex, France

⁸Yukawa Institute for Theoretical Physics, Kyoto University, Kitashirakawa Oiwake–Cho, Kyoto 606–8502, Japan

^hIPN Orsay, Université Paris Sud, IN2P3 – CNRS, F–91406 Orsay Cedex, France

Abstract

The properties of nuclei with extreme neutron–to–proton ratios, far from those naturally occurring on Earth, are key to understand nuclear forces and how nucleons hold together to form nuclei. ⁷H, with six neutrons and a single proton, is the nuclear system with the most unbalanced neutron–to–proton ratio known so far. However, its sheer existence and properties are still a challenge for experimental efforts and theoretical models. Here we report experimental evidences on the formation of ⁷H as a resonance, detected with independent observables, and the first measurement of the structure of its ground state. The resonance is found at ~0.7 MeV above the ³H+4n mass, with a narrow width of ~0.2 MeV and a $1/2^+$ spin and parity. These data are consistent with a ⁷H as a ³H core surrounded by an extended four-neutron halo, with a unique four-neutron decay and a relatively long half-life thanks to neutron pairing; a prime example of new phenomena occurring in what would be the most pure-neutron nuclear matter we can access in the laboratory.

Keywords: 7H, Hydrogen resonance, Active target

1. Introduction

Experimentally, the use of nuclear reactions with unstable nuclei permits to explore regions of the nuclear chart far from stability and beyond the limits of particle binding by systematically adding or removing nucleons [1]. This is particularly feasible for light nuclei, where the binding limits and large neutron-to-proton ratios can be reached by adding few nucleons to stable isotopes. In hydrogen, experiments have already obtained evidences of ^{4,5,6,7}H isotopes, a chain of four resonances beyond the neutron dripline [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27]. These systems are found to decay by neutron emission and thus believed to be built on a core of ³H surrounded by neutrons confined by a centrifugal barrier and grouped in pairs. The systematic study of such a chain helps to understand the evolution of nuclear phenomena, as the effect of neutron pairing and the properties of dilute nuclear matter [28], away from stability and into the nuclear continuum. At the end of this chain we find the super-heavy ⁷H isotope, the nucleus

^cUniversidad de Sevilla, E–41080 Sevilla, Spain

^dKVI–CART, University of Groningen, NL–9747 AA, Groningen, The Netherlands

^{*}Corresponding author

Email addresses: manuel.fresco@usc.es (M. Caamaño), roger@ganil.fr (T. Roger)

¹Present address: Department of Physics, University of Regina, Regina, SK S4S 0A2, Canada

²Present address: Indian Institute of Technology (Indian School of Mines), Dhanbad, Jharkhand - 826004, India

³Present address: LPC Caen, Université de Caen Basse-Normandie-ENSICAEN-CNRS/IN2P3, F-14050 Caen

Cedex, France

⁴Present address: Canadian Nuclear Laboratories, Canada

⁵Present address: GANIL, CEA/DSM–CNRS/IN2P3, BP 55027, F–14076 Caen Cedex 5, France

⁶Present address: Irfu, CEA, Université Paris–Saclay, 91191 Gif–sur–Yvette, France




Figure 1: Schematic drawing of the detection set-up. A typical protontransfer reaction producing 7 H with a 19 F nucleus is also shown.

with the most unbalanced neutron-to-proton ratio of the nuclear chart. ⁷H is expected to display unusual characteristics and open questions. Neutron pairing may render ⁷H the least unstable of the chain despite being the most neutron-rich, forcing it to decay directly to ³H by emitting four neutrons simultaneously [24, 29]. However, other theoretical models predict the resonance to be \sim 3 MeV above the ³H+4n mass [30, 31], and thus favouring a sequential neutron emission over a unique simultaneous decay. Concerning its structure, a recent model based on Antisymmetrised Molecular Dynamics (AMD) describes the four outer neutrons of ⁷H grouped in pairs, and acting as two bosons held together by their interaction with the ³H core in a di-neutron condensate [31]; an exclusive feature of 7 H among the known hydrogen resonances since a di-neutron condensate needs at least two neutron pairs, hence four valence neutrons. The behaviour of these neutrons can also help to understand the influence and strength of T=3/2 threebody neutron forces, which are key ingredients of the Equation of State that rules the properties and structure of compact stars and supernovae [32, 33]. The AMD model also predicts that the most probable configuration keeps the di-neutron pairs in a region where the nuclear density of the system drops to 0.02 fm⁻³, almost 10 times lower than the saturation density of nuclear matter. These conditions of density and extreme neutron-to-proton ratio are unique in a nuclear system that can be studied in the laboratory but similar to those expected in a neutron-star crust [34, 35].

From the experimental point of view, the access to these nuclear systems is challenging: their production probabilities are very small; their short lifetimes,




Figure 2: Kinematics of target-like recoil. The figure shows the range of target-like products as a function of their recoil angle for events measured in coincidence with a single ³H. The colour lines show reference kinematics of one-proton transfer with ¹⁹F (red) and ¹²C (green). Q_F and Q_C correspond to the Q-values of proton-transfer channels forming a ³H+4n system with ¹⁹F and ¹²C targets, respectively. The white dashed lines enclose the data displayed in Fig. 3(a).

of the order of 10^{-22} s, prevent direct measurements; and their multi-particle decays complicate the identification of the resonances. This is particularly true for ⁷H, with few experiments with very low statistics reporting evidences on its formation. Meanwhile, its main characteristics, even its existence, are yet to be precisely determined. The first evidence was found in p(⁸He,⁷H)pp knock-out reactions [24]: a sharp increase in the energy distribution of the beam-like product just above the ³H+4n mass was interpreted as hint of the formation of ⁷H. Later, other indications were reported with ²H(⁸He,⁷H)³He reactions: A candidate resonance was found between 1 and 3 MeV above the ³H+4n mass [36], while a recent campaign suggests a ⁷H ground state somewhere around 2 MeV and a width below 300 keV and, interestingly, possible excited states at ~6 MeV [26, 27]. Other experiments using the same reaction channel show less conclusive evidences [37, 38]. A setup similar to the one presented here was used to identify a resonance peak with a width of ~0.1 MeV, placed at ~0.6 MeV above the ${}^{3}H+4n$ mass [25] in ¹²C(⁸He,⁷H)¹³N reactions. As to its structure, there are no data on the ⁷H spin and parity.

2. Experimental set-up

In order to contribute to these sparse results, we have explored binary, one-proton transfer reactions between a ⁸He beam and carbon and fluorine targets to investigate the formation and properties of ⁷H. In 2011, a ⁸He beam with 10^4 particles per second was produced and accelerated to 15.4A MeV in the SPIRAL facilities at GANIL (France) before being directed to an experimental set-up based on the MAYA activetarget [39] and the CATS beam-tracking detectors [40]. The MAYA active-target detector works essentially as a Time, Charge-Projection Chamber where the filling gas also plays the role of reaction target. In this case, we use 176 mbar of a 90%–10% molar mix of helium and CF₄, with equivalent thickness of 4.2·10¹⁹ atoms/cm² of ¹⁹F and 1.1·10¹⁹ atoms/cm² of ¹²C. This mixture allowed enough electron multiplication while maintaining a small stopping power of the reaction products. Figure 1 shows a schematic drawing of the experimental setup with a typical ¹⁹F(⁸He,⁷H)²⁰Ne transfer event.

In a proton-transfer event producing ⁷H, the trajectory of the ⁸He projectile is measured by the CATS beam-tracking detectors before entering MAYA. Once inside, the ⁸He projectile interacts with either a ¹²C or ¹⁹F nucleus and transfers a proton, yielding a ¹³N or ²⁰Ne target-like recoil and a ⁷H resonance that, less than 10^{-20} s after the reaction, decays into a ³H nucleus and four neutrons. The trajectory of the ¹³N or ²⁰Ne targetlike is imaged in the segmented pad plane of MAYA, where the angle and range are measured with typical uncertainties of 1.2° and 16 mm, respectively. The ³H scattered at forward angles is identified in a dE-E telescope composed of a first layer of 20 5×5-cm², 75- μ m thick silicon detectors and a second layer of 80 2.5×2.5 cm², 1-cm thick CsI crystal detectors. In this setup, neutrons are not detected. Unreacted beam projectiles are collected before the dE-E telescope in a 2×2 cm² aluminium beam-stopper. The experimental acceptance and uncertainties were verified with the elastic measurement, while the target thickness and beam current were confirmed with the elastic cross-section.

3. Data Analysis

3.1. Identification of ^{7}H resonance production

The selection of ⁷H candidate events is done with the simultaneous measurement of the reaction charged products. A measured proton-transfer reaction with ¹²C or ¹⁹F consists of a single ³H detected in the dE–E telescope in coincidence with a single trajectory from the target-recoil product, projected on the pad plane. This selection rejects reactions with the helium atoms in the gas, where none of the products ionises enough to induce an image on the pad plane.

Fusion and high-excitation breakup channels with more than two ionising products and/or more than two




Figure 3: Experimental evidence of the ⁷H formation. (a) The MM distribution of the data selected between the white dashed lines in Fig. 2 (black dots) is shown in one-proton transfer reactions with ¹⁹F (red lower axis) or ¹²C (green top axis), and after subtraction of background from incomplete events (see text). (b) Measured range distribution of target–like products with a recoil angle between 45° and 54° in the laboratory frame and after subtraction of background from incomplete events. In both panels, the production of ⁷H (solid lines) and non-resonant backgrounds (N-R, dashed lines) are shown from reactions with ¹⁹F and ¹²C targets (red and green lines, respectively).

products on the dE–E telescope are also rejected. However, some breakup events may mimic one-proton transfer if only a ³H is detected in the dE–E telescope and a track is recorded in MAYA while the rest of the products are missed. The distribution of these incomplete events is obtained from those detected with a ³H in coincidence with any other product in the dE–E telescope and a track inside MAYA. The resulting distribution is subtracted from the set of ⁷H candidate events. This contamination of breakup channels amounts to less than 10% of single ³H and track events, and it follows a smooth behaviour without peaks or recognisable features. Concerning random coincidences between events, they are below 0.1%.

One-proton transfer reactions correlate the angle and range of the target-like recoils following kinematic lines that depend on the mass of the nuclei involved, including ⁷H. Figure 2 shows this correlation for the measured data, along kinematic lines corresponding to two reference values for the ⁷H mass. The accumulation of events around the kinematic lines of Q_F and Q_C in Fig. 2 is consistent with the production of ⁷H in both channels. A second hint is the abrupt decrease of events we can see around 80 mm along the Q_F line. This is due to the angular acceptance in the detection of the scattered ³H and it would only affect binary reactions, reinforcing the identification of the accumulation of data around the line as transfer reactions producing ⁷H.

Since 7 H decays immediately after being formed, its characteristics can be indirectly observed with the missing-mass (MM) method, in which the mass of the undetected participant (7 H, in this case) is deduced from the kinematics of the remaining participants in the reaction. Figure 3(a) shows the MM spectrum of the beam-like product from one-proton transfer reactions with respect to the mass of the ³H+4n subsystem for data within the largest c.m. angular region of the ¹⁹F(⁸He,³H+4n)²⁰Ne channel that is not cut by acceptance, shown with white dashed lines in Fig. 2. Our experimental setup does not perform element separation of the target-like recoils, thus the lower and top horizontal axis of Fig. 3(a) show the MM as calculated for ¹⁹F and ¹²C targets, respectively. The events accumulated around the Q_F and Q_C kinematic lines shown in Fig. 2 would appear in Fig. 3(a) as two peaks: one around zero in the MM spectra of ¹⁹F targets and another around zero for ¹²C targets.

The first peak lies in a clean kinematical region: only low-lying states of ⁷H or the lower tail of a 3-body nonresonant (N-R) continuum (which would not be a peak) can populate it. In addition, the angle and energy of the ³H detected in coincidence were checked to be within the limits expected from a possible ⁷H decay. Other channels, such as partially reconstructed breakup channels, are subtracted from our selections, as discussed previously. This peak is a safe candidate for the resonant formation of ⁷H with ¹⁹F targets. The peak that would correspond to the formation of ⁷H with ¹²C targets is less obvious: it occupies a region also populated by N-R background from both targets and other multiparticle transfer channels. A fit to possible contributions other than ⁷H and its N-R continuum sets an upper limit of 0.2 mb/sr to their production.

3.2. Resonance mass and half-life

The values of the resonance mass and width are obtained by fitting a simulation of the main experimental observables to the collected data. The simulation is folded with the experimental resolutions and measurement conditions, and includes the production of ⁷H with both ¹⁹F and ¹²C targets, and also N-R events. Besides these channels, other multi-particle transfer reactions were also considered. The measured distributions, and in particular the observed widths, are dominated by the experimental uncertainty, which can reach 10 MeV, depending on the kinematical region, and determines the final uncertainty on the measurement of the resonance parameters.

In order to describe the peak associated with ⁷H, we use a Breit–Wigner probability distribution following the prescription of [25]. The mass and width of the resonance, as well as the scaling factor, were treated as free parameters and extracted from a log-likelihood minimisation between the simulation and the measured range

distribution in the angular region between 45° and 54° , shown in Fig. 3(b). This region is mainly populated by the production of ⁷H and N-R components, with a very small contribution from the ¹²C(⁸He,⁷H)¹³N channel, making it especially well-suited for a clean fit.

The results of the fit describe a low-lying, narrow resonance state with a mass of $0.73^{+0.58}_{-0.47}$ MeV above the ³H+4n mass and a width of $0.18^{+0.47}_{-0.16}$ MeV. The mass value reinforces the notion of ⁷H as the least unstable of the known hydrogen resonances, less than 1 MeV close to being bound, despite being the most neutronrich. The reinforced stability brought by neutron pairing also gives a narrow width to ⁷H, which translates into a half-life of $5 \cdot 10^{-21}$ s, an order of magnitude longer than the other hydrogen resonances.

Compared to previous experiments, our measured mass and width are in good agreement with the $0.57^{+0.42}_{-0.21}$ and $0.09^{+0.94}_{-0.06}$ MeV reported in [25]. The value estimated in [26], 1.8±0.5, is larger but not too dissimilar from our result while ref. [27] reports the largest value at 2.2 ± 0.5 MeV. It is worthy to note that in each of these references, not more than 10 ⁷H events were collected, less than one order of magnitude lower compared to the present work. Evidences of excited states of ⁷H were reported around 6 MeV in refs. [26, 27]. In the case of the ¹⁹F channel, these states may be hidden in the tail of the peak associated with the ⁷H ground state, although a fit to a tentative state around 6 MeV gives no statistically significant population above 0.1 mb/sr. From the theory side, calculations tend to heavier ⁷H masses. In particular, models based on a di-neutron condensate from AMD [29] and on hyperspherical functions methods [30] describe a resonance with a mass of ~3 MeV [31]. Concerning the resonance width, ref. [29] finds a strong correlation with its mass and predicts values below 1 keV for masses around 1 MeV, two orders of magnitude below our measured width.

3.3. Angular distribution and structure

The capabilities of MAYA to measure low energy products and the relatively high intensity of the ⁸He beam have allowed us to collect more than 200 events assigned to ⁷H formation, significantly more than previous experiments. The average production cross-section with ¹⁹F is 2.7 ± 0.5 mb/sr between 4° and 18° in the centre of mass reference frame (c.m.), whereas ¹²C yields $1.2^{+0.5}_{-0.6}$ mb/sr between 6° and 27°. Besides the corresponding statistical uncertainty, systematic uncertainties from the number of incoming projectiles and target thickness are around 0.7%, while the uncertainty on the position and width of the resonance contribute




Figure 4: (a) Measured differential cross-section of the ${}^{19}F+{}^{8}He$ proton-transfer (black dots) compared with DWBA calculation of a proton transfer to a $1/2^+$ state in ${}^{7}H$ and a ${}^{20}Ne$ in its 0^+ ground state (red line) and other spin combinations (yellow, green, and blue lines). (b) Measured differential cross-section of the ${}^{12}C+{}^{8}He$ proton-transfer channel (black dots) compared with a scaled DWBA calculation performed with a $1/2^+$ ${}^{7}H$ and the $1/2^-$ ground state of ${}^{13}N$ (green line). In both panels, vertical error bars include statistical uncertainty and the effect of the uncertainty in the parameters of the resonance.

with ~10% of the cross-section value. The ¹²C channel was also measured in [25], reporting $0.04^{+0.06}_{-0.03}$ mb/sr between 10° and 48°. When evaluated in the same angular region, our measurement averages to $0.4^{+0.2}_{-0.3}$ mb/sr, a larger value although still compatible within 1.1 of standard deviation.

Previous experiments mostly used ²H(⁸He,⁷H)³He reactions, obtaining results that vary with the beam energy and angular coverage. At 15.3 AMeV, values below 0.1 mb/sr were measured in a wide 0° -50° region in c.m. [36]. Around 25 AMeV, ref. [37] estimates a cross-section below 0.02 mb/sr in 9°-21° in c.m., while ref. [26] reports 0.025 mb/sr between 17° and 27°, and in ref. [27] even seems to increase beyond 0.04 mb/sr for angles below 10° in c.m. When increasing the beam energy to 42 AMeV, ref. [38] finds a similar 0.03 mb/sr between 6° and 14° in c.m. In addition to these transfer measurements, the pioneering work of ref. [24] calculates the production of ⁷H around 0.01 mb/sr·MeV with proton knock-out reactions. While a precise comparison is difficult due to the different reaction mechanisms and angular coverage, and their low statistics, the complete list of results seems to suggest a dependence of the cross-section with the target size. An explanation for this behaviour is beyond the scope of this work. However, these results, together with the proposition in ref. [27] of an extreme peripheral character of the ⁷H ground-state population, suggest that a deeper understanding of the reaction mechanism leading to the formation of ⁷H may be needed to reproduce the ensemble of experimental data.

The improved statistics have also permitted to measure the c.m. angular distribution of the ⁷H production with both targets. Figure 4(a) shows that the angular distribution of the ¹⁹F(⁸He,⁷H)²⁰Ne channel follows a clear oscillating pattern, with distinct minima. This behaviour is a strong indication of the formation of two well-defined systems in the output channel: ²⁰Ne and the ⁷H resonance; it offers a further, independent evidence of its production. The measurement of the angular distribution of ¹²C(⁸He,⁷H)¹³N, shown in Fig. 4(b), suffered from large statistical and systematic uncertainties due to the contribution of competing channels, as discussed previously. The relatively featureless behaviour and its uncertainties do not allow a clear assignment of spin and parity but only a rough assessment of the mean differential cross-section.

The angular distribution is compared in Fig. 4 with different DWBA calculations of the cross-section made with the FRESCO [41] code. In these calculations, we have used shell-model spectroscopic factors for the $\langle {}^{20}\text{Ne}|{}^{19}\text{F}\rangle$ overlaps, using the WBT effective interaction by Warburton and Brown [42]. The ⁷H nuclear density was obtained from AMD calculations assuming a di-neutron condensate structure around a ³H core [31]. The resulting cross-sections were folded with the experimental uncertainties and further scaled to match the experimental data. The scaling factor was found to vary between 4.5 ± 2.8 and 12.7 ± 6.1 , depending on the prescription for the nuclear density of ⁸He. We explore the possibility of populating the 0^+ ground state or the 2^+ first excited state of ²⁰Ne, and for ⁷H, we consider either a $1/2^+$ state with a ³H core in its ground state and four outer neutrons, or a $3/2^{-}$ state with an excited ³H core. The relative amplitude of the oscillations and the positions of the minima are best reproduced with a proton transfer to the 0^+ ground state of 20 Ne and a $1/2^+$ ⁷H resonance. The agreement between the calculations and the data suggests that the di-neutron pairs keep a separation similar to the one predicted by the AMD calculations. Concerning channel mixing, a tentative fit to evaluate a possible mixture of transfer to 0^+ and 2^+ states in ²⁰Ne gives a probability of less than 10% towards the excited 2^+ state. The preference for the transfer to the 0⁺ state is surprisingly different from DWBA calculations, which predict a more probable transfer to the 2⁺ state in ²⁰Ne. The data suggest that either this is not the case or that the resonance has a lower probability of being fully formed in the output channel with the 2⁺ state in ²⁰Ne. As suggested in ref. [27], the very radially extended and, thus, very "fragile" nature of the ⁷H ground state may play a role to understand this observation and the set of cross-section measurements.

4. Summary

In conclusion, we have measured evidences of the formation of the ⁷H resonance with a larger statistical significance than previous attempts via two different reaction channels and performed the characterisation of its ground state with two independent observables: the resonance MM distribution and the angular distribution. From these observables, we have obtained a new determination of the mass and width of ⁷H, and, for the first time, an assignment of spin and parity to its ground state. Together, these results depict the super-heavy ⁷H nucleus as an extended pure-neutron shell around a ³H core in a $1/2^+$ ground state. However, the same neutron pairing that allows this large neutron configuration also renders the ⁷H nucleus a long-lived and almost-bound resonance, despite being the system with what would be the largest neutron-to-proton ratio in the nuclear chart known today.

Acknowledgments

The authors thank Navin Alahari for the useful discussions and the careful reading of the manuscript, and Miguel Marqués for his help in the data analysis. The authors are deeply thankful to the technical staff at GANIL for their support and help. This work has been supported by the European Community FP7-Capacities -Integrated Infrastructure Initiative- contract ENSAR n° 262010, and by the Spanish Ministerio de Economía y Competitividad under contracts FPA2009-14604-C02-01 and FPA2012-39404-C02-01. M.C. acknowledges the support by the Spanish Ministerio de Economía y Competitividad through the Programmes "Ramón y Cajal" with the grant number RYC-2012-11585 and "Juan de la Cierva" with the grant number JCI2009-05477. A.M.M. is partially supported by the project Ref. P20_01247, funded by the Consejería de Economía, Conocimiento, Empresas y Universidad, Junta de Andalucía (Spain) and by "ERDF A way of making Europe."

Competing interests

The authors declare no competing interests.

References

 P. G. Hansen, Beyond the neutron drip line, Nature 328 (1987) 476.

- [2] R. Cohen, et al., Two-body breakups following π⁻ absorption in lithium: evidence for the production of H⁴, Phys. Lett. 14 (1964) 242.
- [3] R. C. Minehart, et al., Pion capture in ⁶Li and ⁷Li, the formation of ⁴H, a search for ⁵h, Phys. Rev. 177 (1969) 1455.
- [4] T. C. Meyer, A study of particle unstable ⁴H, Nucl. Phys. A 324 (1979) 335.
- [5] U. Sennhauser, et al., Spectroscopy of single and correlated charged particles emitted following bound pion absorption in ⁶Li and ⁷li, Nucl. Phys. A 386 (1982) 429.
- [6] R. Franke, et al., Search for highly excited states in light nuclei with three-body reactions, Nucl. Phys. A 433 (1985) 351.
- [7] D. Milijanić, et al., ⁴H and (n, α x) reactions on ⁶li and ⁷li, Phys. Rev. C 33 (1986) 2204.
- [8] A. V. Belozyorov, et al., Search for ⁴H, ⁵H and ⁶H nuclei in the ¹¹B-induced reaction on ⁹Be, Nucl. Phys. A 460 (1986) 352.
- [9] S. Blagus, et al., ⁴H nucleus and the ²H(t,tp)n reaction, Phys. Rev. C 44 (1991) 325.
- [10] S. I. Sidorchuk, et al., Resonance states of hydrogen nuclei ⁴H and ⁵H obtained in transfer reactions with exotic beams, Nucl. Phys. A 719 (2003) 229c.
- [11] M. Meister, et al., Searching for the ⁵H resonance in the t + n + n system, Nucl. Phys. A 723 (2003) 13.
- [12] S. I. Sidorchuk, et al., Experimental study of ⁴H in the reactions ²H(t,p) and ³H(t,d), Phys. Lett. B 594 (2004) 54.
- [13] Y. B. Gurov, et al., Spectroscopy of superheavy hydrogen isotopes ⁴H and ⁵H, Eur. Phys. J. A 24 (2005) 231.
- [14] T. A. Tombrello, Phase-shift analysis of t(n, n)t, Phys. Rev. 143 (1965) 772.
- [15] P. G. Young, et al., Search for the ground state of ⁵H by means of the ³H(t,p) reaction, Phys. Rev. 173 (1968) 949.
- [16] A. A. Korsheninnikov, et al., Superheavy hydrogen ⁵H, Phys. Rev. Lett. 87 (2001) 092501.
- [17] M. Meister, et al., The t+n+n system and ${}^{5}H$, Phys. Rev. Lett. 91 (2003) 162504.
- [18] M. S. Golovkov, et al., Evidences for resonance states in ⁵H, Phys. Lett. B 566 (2003) 70.
- [19] M. S. Golovkov, et al., Observation of excited states in ⁵H, Phys. Rev. Lett. 93 (2004) 262501.
- [20] S. V. Stepantsov, et al., ⁵H and ⁵He nuclear systems studied by means of the ⁶He+²H reaction, Nucl. Phys. A 738 (2004) 436.
- [21] M. S. Golovkov, et al., Correlation studies of the ⁵H spectrum, Phys. Rev. C 72 (2005) 064612.
- [22] G. M. Ter-Akopian, et al., New insights into the resonance states of ⁵H and ⁵He, Eur. Phys. J. A 25 (2005) 315.
- [23] A. H. Wuosmaa, et al., Ground-state properties of ⁵H from the ⁶He(d,³He)⁵H reaction, Phys. Rev. C 95 (2017) 014310.
- [24] A. A. Korsheninnikov, et al., Experimental evidence for the existence of ⁷H and for a specific structure of 8He, Phys. Rev. Lett. 90 (2003) 082501.
- [25] M. Caamaño, et al., Resonance state in ⁷H, Phys. Rev. Lett. 99 (2007) 062502.
- [26] A. A. Bezbakh, et al., Evidence for the first excited state of ⁷H, Phys. Rev. Lett. 124 (2020) 022502.
- [27] I. A. Muzalevskii, et al., Resonant states in ⁷H: Experimental studies of the ²H(⁸He, ³He) reaction, Phys. Rev. C 103 (2021) 044313.
- [28] K. Hagino, H. Sagawa, J. Carbonell, P. Schuck, Coexistence of bcs- and bec-like pair structures in halo nuclei, Phys. Rev. Lett. 99 (2007) 022506.
- [29] M. S. Golovkov, et al., Estimates of the ⁷H width and lower decay energy limit, Phys. Lett. B 588 (2004) 163.
- [30] N. K. Timofeyuk, Hyperspherical harmonics with orthogonal symmetry in the shell model approach, and its application to light nuclei, Phys. Rev. C 69 (2004) 034336.

- [31] S. Aoyama, N. Itagaki, Di-neutron correlations in ⁷H, Phys. Rev. C 80 (2009) 021304(R).
- [32] A. Akmal, et al., Equation of state of nucleon matter and neutron star structure, Phys. Rev. C 58 (1998) 1804.
- [33] M. Oertel, et al., Equations of state for supernovae and compact stars, Rev. Mod. Phys. 89 (2017) 015007.
- [34] N. Chamel, P. Haensel, Physics of neutron star crusts, Living Rev. Relativity 11 (2008) 10.
- [35] F. Gulminelli, A. R. Raduta, Unified treatment of subsaturation stellar matter at zero and finite temperature, Phys. Rev. C 92 (2015) 055803.
- [36] S. Fortier, et al., Search for resonances in ⁴n, ⁷H and ⁹He via transfer reactions, in: International Symposium on Exotic Nuclei, Vol. 912 of AIP Conference Proceedings, 2007, p. 3.
- [37] G. M. Ter-Akopian, et al., Neutron excess nuclei of hydrogen and helium at acculinna, Eur. Phys. J. Special Topics 150 (2007) 61.
- [38] E. Y. Nikolskii, et al., Search for ⁷H in ²H+⁸He collisions, Phys. Rev. C 81 (2010) 064606.
- [39] C. E. Demonchy, et al., Maya: An active-target detector for binary reactions with exotic beams, Nucl. Instr. and Meth. A 583 (2007) 341.
- [40] S. Ottini-Hustache, et al., Cats, a low pressure multiwire proportionnal chamber for secondary beam tracking at ganil, Nucl. Instr. and Meth. A 431 (1999) 475.
- [41] I. J. Thompson, Coupled reaction channels calculations in nuclear physics, Comput. Phys. Rep. 7 (1988) 167.
- [42] E. K. Warburton, B. A. Brown, Effective interactions for the 0p1s0d nuclear shell-madel space, Phys. Rev. C 46 (1992) 923.