

TOPICS ON FERMI VARIETIES OF DISCRETE PERIODIC SCHRÖDINGER OPERATORS

WENCAI LIU

ABSTRACT. This is a survey of recent progress on the irreducibility of Fermi varieties, rigidity results and embedded eigenvalue problems of discrete periodic Schrödinger operators.

1. INTRODUCTION

The geometry of Fermi varieties plays critical roles in studying many topics of periodic Schrödinger operators. In this article, we will first discuss the irreducibility of the Fermi variety as an analytic set and then study rigidity results (inverse problems) and embedded eigenvalue problems of discrete periodic Schrödinger operators. We refer readers to an earlier survey article [32] for other topics, techniques, and results of both continuous and discrete periodic Schrödinger operators.

1.1. Fermi, Bloch and Floquet varieties. Given $q_i \in \mathbb{Z}_+$, $i = 1, 2, \dots, d$, let $\Gamma = q_1\mathbb{Z} \oplus q_2\mathbb{Z} \oplus \dots \oplus q_d\mathbb{Z}$. We say that a function $V : \mathbb{Z}^d \rightarrow \mathbb{C}$ is Γ -periodic (or just periodic) if for any $\gamma \in \Gamma$ and $n \in \mathbb{Z}^d$, $V(n + \gamma) = V(n)$.

For $n = (n_1, n_2, \dots, n_d) \in \mathbb{Z}^d$, denote by $\|n\|_1 = \sum_{i=1}^d |n_i|$. Let Δ be the discrete Laplacian on lattices \mathbb{Z}^d , namely

$$(\Delta u)(n) = \sum_{n' \in \mathbb{Z}^d, \|n' - n\|_1 = 1} u(n').$$

Consider the discrete Schrödinger operator on $\ell^2(\mathbb{Z}^d)$

$$H_0 = -\Delta + V. \tag{1}$$

In this article, we always assume that q_i , $i = 1, 2, \dots, d$, are relatively prime, V is periodic and H_0 is the discrete periodic Schrödinger operator given by (1).

Date: February 18, 2022.

2010 Mathematics Subject Classification. 39A12 (primary); 14H10, 35J10, 47A75 (secondary).

Key words and phrases. Fermi variety, Bloch variety, analytic variety, algebraic variety, irreducibility, periodic Schrödinger operator, embedded eigenvalue, unique continuation, rigidity theorem, isospectrality, Floquet isospectrality, Fermi isospectrality.

Let $\{\mathbf{e}_j\}$, $j = 1, 2, \dots, d$, be the standard basis in \mathbb{Z}^d :

$$\mathbf{e}_1 = (1, 0, \dots, 0), \mathbf{e}_2 = (0, 1, 0, \dots, 0), \dots, \mathbf{e}_d = (0, 0, \dots, 0, 1).$$

Definition 1. The Bloch variety $B(V)$ of $-\Delta + V$ consists of all pairs $(k, \lambda) \in \mathbb{C}^{d+1}$ for which there exists a non-zero solution of the equation

$$(-\Delta u)(n) + V(n)u(n) = \lambda u(n), n \in \mathbb{Z}^d, \quad (2)$$

satisfying the so called Floquet-Bloch boundary condition

$$u(n + q_j \mathbf{e}_j) = e^{2\pi i k_j} u(n), j = 1, 2, \dots, d, \text{ and } n \in \mathbb{Z}^d, \quad (3)$$

where $k = (k_1, k_2, \dots, k_d) \in \mathbb{C}^d$.

Definition 2. Given $\lambda \in \mathbb{C}$, the Fermi surface (variety) $F_\lambda(V)$ is defined as the level set of the Bloch variety:

$$F_\lambda(V) = \{k : (k, \lambda) \in B(V)\}.$$

Definition 3. Let $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ and $z = (z_1, z_2, \dots, z_d)$. The Floquet variety is defined as

$$\mathcal{F}_\lambda(V) = \{z \in (\mathbb{C}^*)^d : z_j = e^{2\pi i k_j}, j = 1, 2, \dots, d, k \in F_\lambda(V)\}. \quad (4)$$

Clearly,

$$F_\lambda(V) = \{k \in \mathbb{C}^d : (e^{2\pi i k_1}, e^{2\pi i k_2}, \dots, e^{2\pi i k_d}) \in \mathcal{F}_\lambda(V)\}.$$

Namely, $z \in (\mathbb{C}^*)^d \in \mathcal{F}_\lambda(V)$ if and only if the equation

$$(-\Delta u)(n) + V(n)u(n) = \lambda u(n), n \in \mathbb{Z}^d \quad (5)$$

with the boundary condition

$$u(n + q_j \mathbf{e}_j) = z_j u(n), j = 1, 2, \dots, d, \text{ and } n \in \mathbb{Z}^d \quad (6)$$

has a non-zero solution.

1.2. Algebricity of the Floquet variety. Introduce a fundamental domain W for Γ :

$$W = \{n = (n_1, n_2, \dots, n_d) \in \mathbb{Z}^d : 0 \leq n_j \leq q_j - 1, j = 1, 2, \dots, d\}.$$

By writing out $-\Delta + V$ as acting on the $Q = q_1 q_2 \dots q_d$ dimensional space $\{u(n), n \in W\}$, the eigen-equation (5) and (6) ((2) and (3)) translates into the eigenvalue problem for a $Q \times Q$ matrix $\mathcal{D}_V(z)$ ($D_V(k)$). Let $\mathcal{P}_V(z, \lambda)$ ($P_V(k, \lambda)$) be the determinant of $\mathcal{D}_V(z) - \lambda I$ ($D_V(k) - \lambda I$).

It is easy to see that

$$F_\lambda(V) = \{k \in \mathbb{C}^d : P_V(k, \lambda) = 0\}, \quad (7)$$

and

$$\mathcal{F}_\lambda(V) = \{z \in (\mathbb{C}^*)^d : \mathcal{P}_V(z, \lambda) = 0\}. \quad (8)$$

Example: $d = 1$

$$D_V(k) = \begin{pmatrix} V(1) & -1 & 0 & \cdots & 0 & e^{-2\pi i k} \\ -1 & V(2) & -1 & \cdots & 0 & 0 \\ 0 & -1 & V(3) & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & V(q_1 - 1) & -1 \\ e^{2\pi i k} & 0 & 0 & \cdots & -1 & V(q_1) \end{pmatrix}.$$

It is not difficult to see that $\mathcal{P}_V(z, \lambda)$ is a polynomial in variables λ and $z_1, z_1^{-1}, z_2, z_2^{-1}, \dots, z_d, z_d^{-1}$. See Lemma 1.1 for a more precise description. In other words, $\mathcal{P}_V(z, \lambda)$ is a Laurent polynomial of λ and z_1, z_2, \dots, z_d . Therefore, the Floquet variety $\mathcal{F}_\lambda(V)$ is essentially an algebraic set.

1.3. Discrete Floquet transform. Define the discrete Fourier transform $\hat{V}(l)$, $l \in W$ by

$$\hat{V}(l) = \frac{1}{Q} \sum_{n \in W} V(n) \exp \left\{ -2\pi i \left(\sum_{j=1}^d \frac{l_j n_j}{q_j} \right) \right\}.$$

For convenience, we extend $\hat{V}(l)$ to a Γ -periodic function on \mathbb{Z}^d , namely, for any $l \equiv m \pmod{\Gamma}$,

$$\hat{V}(l) = \hat{V}(m).$$

Let

$$\tilde{\mathcal{D}}_V(z) = \tilde{\mathcal{D}}_V(z_1, z_2, \dots, z_d) = \mathcal{D}_V(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}), \quad (9)$$

and

$$\tilde{\mathcal{P}}_V(z, \lambda) = \det(\tilde{\mathcal{D}}_V(z, \lambda) - \lambda I) = \mathcal{P}_V(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}, \lambda). \quad (10)$$

Let

$$\rho_{n_j}^j = e^{2\pi i \frac{n_j}{q_j}},$$

where $0 \leq n_j \leq q_j - 1$, $j = 1, 2, \dots, d$.

By the standard discrete Floquet transform (e.g., [32, 33, 40]), one has

Lemma 1.1. *Let $n = (n_1, n_2, \dots, n_d) \in W$ and $n' = (n'_1, n'_2, \dots, n'_d) \in W$. Then $\tilde{\mathcal{D}}_V(z)$ is unitarily equivalent to $A + B_V$, where A is a diagonal matrix with entries*

$$A(n; n') = - \left(\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) \right) \delta_{n, n'} \quad (11)$$

and

$$B_V(n; n') = \hat{V}(n_1 - n'_1, n_2 - n'_2, \dots, n_d - n'_d). \quad (12)$$

In particular,

$$\tilde{\mathcal{P}}_V(z, \lambda) = \det(A + B_V - \lambda I).$$

From Lemma 1.1, we have that $\mathcal{P}_V(z, \lambda)$ is a polynomial in variables λ and $z_1, z_1^{-1}, \dots, z_d, z_d^{-1}$, with highest degree terms (up to a \pm sign),

$$\lambda^Q, z_1^{\frac{Q}{q_1}}, z_1^{-\frac{Q}{q_1}}, z_2^{\frac{Q}{q_2}}, z_2^{-\frac{Q}{q_2}}, \dots, z_d^{\frac{Q}{q_d}}, z_d^{-\frac{Q}{q_d}}.$$

1.4. Spectral bands. When V is a real periodic function, H_0 is a self-adjoint operator and $D_V(k)$ is a Hermitian matrix for any $k \in \mathbb{R}^d$. For each $k \in \mathbb{R}^d$, $D_V(k)$ has Q eigenvalues and order them in non-decreasing order

$$\lambda_V^1(k) \leq \lambda_V^2(k) \leq \dots \leq \lambda_V^Q(k). \quad (13)$$

We call $\lambda_V^m(k)$ the m -th (spectral) band function, $m = 1, 2, \dots, Q$. Let

$$[a_m, b_m] = [\min_{k \in \mathbb{R}^d} \lambda_V^m(k), \max_{k \in \mathbb{R}^d} \lambda_V^m(k)], m = 1, 2, \dots, Q.$$

Clearly, a_m and b_m , $m = 1, 2, \dots, Q$, depend on the potential V . We drop the dependence for simplicity. $[a_m, b_m]$, $m = 1, 2, \dots, Q$, are referred to as spectral bands. By the standard Floquet theory [33], we have

$$\sigma_{ac}(H_0) = \sigma(H_0) = \cup_{m=1}^Q [a_m, b_m], \sigma_{pp}(H_0) = \sigma_{sc}(H_0) = \emptyset. \quad (14)$$

For the zero potential, one has that

$$\sigma_{ac}(-\Delta) = \sigma(-\Delta) = \cup_{m=1}^Q [a_m, b_m] = [-2d, 2d], \sigma_{pp}(-\Delta) = \sigma_{sc}(-\Delta) = \emptyset. \quad (15)$$

2. IRREDUCIBILITY OF FERMI AND BLOCH VARIETIES

2.1. Irreducibility of Fermi and Bloch varieties. In this section, we will discuss the irreducibility of both Fermi and Bloch varieties as analytic sets.

Definition 4. A subset $\Omega \subset \mathbb{C}^k$ is called an analytic set if for any $x \in \Omega$, there is a neighborhood $U \subset \mathbb{C}^k$ of x , and analytic functions f_1, f_2, \dots, f_p in U such that

$$\Omega \cap U = \{y \in U : f_1(y) = 0, f_2(y) = 0, \dots, f_p(y) = 0\}.$$

Definition 5. An analytic set Ω is said to be irreducible if it can not be represented as the union of two non-empty proper analytic subsets.

From (7), one can see that the Fermi/Bloch variety is a principal analytic set, namely, it is determined by a single analytic function. Since the Floquet-Bloch boundary condition is unchanged under the shift: $k \rightarrow k + \mathbb{Z}^d$, it is convenient to study Fermi and Bloch varieties modulo periodicity. It is widely believed that the Bloch/Fermi variety (modulo periodicity) is always irreducible for periodic Schrödinger operators, which has been formulated as conjectures:

Conjecture 1. [33, Conjecture 5.17] The Bloch variety $B(V)$ is irreducible (modulo periodicity).

Conjecture 2. [33, Conjecture 5.35] [35, Conjecture 12] Let $d \geq 2$. Then $F_\lambda(V)/\mathbb{Z}^d$ is irreducible, possibly except for finitely many $\lambda \in \mathbb{C}$.

We remark that in Conjecture 1, the irreducibility of Bloch variety modulo periodicity means for any two irreducible components Ω_1 and Ω_2 of $B(V)$, there exists $k \in \mathbb{Z}^d$ such that $\Omega_1 = (k, 0) + \Omega_2$. In Conjecture 2, for fixed λ , $F_\lambda(V)/\mathbb{Z}^d$ is irreducible means for any two irreducible components Ω_1 and Ω_2 of $F_\lambda(V)$, there exists $k \in \mathbb{Z}^d$ such that $\Omega_1 = k + \Omega_2$.

Conjectures 1 and 2 have been mentioned in many articles [2–4, 15, 31, 36]. See Conjecture 13 in [35] for a “generic” version of Conjecture 2.

The irreducibility of Fermi and Bloch varieties was well understood for $d = 2, 3$ about 30 years ago. When $d = 2$, Bättig [1] proved that the Bloch variety $B(V)$ is irreducible (modulo periodicity). In [15], Gieseke, Knörrer and Trubowitz proved that $F_\lambda(V)/\mathbb{Z}^2$ is irreducible except for finitely many values of λ . When $d = 3$, Bättig [3] proved that the Fermi variety $F_\lambda(V)/\mathbb{Z}^3$ is irreducible for every $\lambda \in \mathbb{C}$. We refer readers to [6, 13, 39] and references therein for more recent related results.

In [1–4, 15], proofs heavily depend on the construction of toroidal and directional compactifications of Fermi and Bloch varieties.

Recently, we introduced a new approach to study the Fermi variety and in particular proved both conjectures [40]. For any $d \geq 3$, we proved that the Fermi variety at every level is irreducible (modulo periodicity). For $d = 2$, we proved that the Fermi variety at every level except for the average of the potential is irreducible (modulo periodicity). We also proved that for any $d \geq 2$ the Bloch variety is irreducible (modulo periodicity).

Our approach focuses on the study of the Laurent polynomial $\mathcal{P}(z, \lambda)$. This approach turns out to be quite robust and has already been used to study the irreducibility of Bloch varieties of discrete periodic operators on general lattices [12].

Before stating the main results in [40], some preparations are necessary.

A Laurent polynomial of a single term is called monomial, i.e., $C z_1^{a_1} z_2^{a_2} \cdots z_k^{a_k}$, where $a_j \in \mathbb{Z}$, $j = 1, 2, \dots, k$, and C is a non-zero constant.

Definition 6. We say that a Laurent polynomial $h(z_1, z_2, \dots, z_k)$ is irreducible if it can not be factorized non-trivially, that is, there are no non-monomial Laurent polynomials $f(z_1, z_2, \dots, z_k)$ and $g(z_1, z_2, \dots, z_k)$ such that $h = fg$.

Remark 1. When h is a polynomial, the definition of irreducibility in Def.6 is slightly different (allowing a difference of monomial) from the traditional definition (a polynomial is called irreducible if it can not be factorized into two non-constant polynomials). For example, the polynomial $z^2 + z$ is irreducible according to Def.6.

Denote by $[V]$ the average of V :

$$[V] = \frac{1}{Q} \sum_{n \in W} V(n).$$

Now we are ready to state the main results in [40].

Theorem 2.1. [40] *Let $d \geq 3$. Then for any $\lambda \in \mathbb{C}$, the Laurent polynomial $\mathcal{P}(z, \lambda)$ (as a function of z) is irreducible.*

Theorem 2.2. [40] *Let $d = 2$. Then the Laurent polynomial $\mathcal{P}_V(z, \lambda)$ (as a function of z) is irreducible for any $\lambda \in \mathbb{C}$ except for $\lambda = [V]$. Moreover, if $\mathcal{P}(z, [V])$ is reducible, $\mathcal{P}(z, [V])$ has exactly two non-trivial irreducible factors (count multiplicity).*

Theorem 2.3. [40] *Let $d \geq 2$. Then the Laurent polynomial $\mathcal{P}(z, \lambda)$ (as a function of both z and λ) is irreducible. In particular, the Bloch variety $B(V)$ is irreducible (modulo periodicity).*

Remark 2. (1) In [40], we actually proved that when $\mathcal{P}_V(z, [V])$ is reducible, $\mathcal{P}_V(z, [V]) = \mathcal{P}_0(z, 0) = ((-1)^{q_2} z_1^{q_2} + (-1)^{q_2} z_1^{-q_2} + (-1)^{q_1} z_2^{q_1} + (-1)^{q_1} z_2^{-q_1})$, where 0 is the zero potential.
 (2) Theorem 2.3 immediately follows from Theorems 2.1 and 2.2 and some simple facts of the Laurent polynomial $\mathcal{P}(z, \lambda)$.

Corollary 2.4. [40] *For any $d \geq 3$, the Fermi variety $F_\lambda(V)/\mathbb{Z}^d$ is irreducible for any $\lambda \in \mathbb{C}$. For $d = 2$, the Fermi variety $F_\lambda(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$ except for $\lambda = [V]$. Moreover, if $F_{[V]}(V)/\mathbb{Z}^2$ is reducible, it has exactly two irreducible components.*

Corollary 2.5. [40] *For any $d \geq 3$, the Floquet variety $\mathcal{F}_\lambda(V)$ is irreducible for any $\lambda \in \mathbb{C}$. For $d = 2$, the Floquet variety $\mathcal{F}_\lambda(V)$ is irreducible for any $\lambda \in \mathbb{C}$ except for $\lambda = [V]$. Moreover, if $\mathcal{F}_{[V]}(V)$ is reducible, it has exactly two irreducible components.*

2.2. Proof of Theorems 2.1 and 2.2 with the zero potential. In order to better illustrate our ideas in [40], we first prove Theorems 2.1 and 2.2 with the zero potential. By Lemma 1.1, for the zero potential,

$$\tilde{\mathcal{P}}_0(z, \lambda) = \mathcal{P}_0(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}, \lambda) = (-1)^Q \prod_{n \in W} \left(\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda \right). \quad (16)$$

Denote by μ_{q_j} the multiplicative group of q_j roots of unity, $j = 1, 2, \dots, d$. Let $\mu = \mu_{q_1} \times \mu_{q_2} \times \dots \times \mu_{q_d}$.

For any $\rho = (\rho^1, \rho^2, \dots, \rho^d) \in \mu$, we define a natural action on \mathbb{C}^d

$$\rho \cdot (z_1, z_2, \dots, z_d) = (\rho^1 z_1, \rho^2 z_2, \dots, \rho^d z_d).$$

Proof of Theorem 2.1 with the zero potential. Assume for some $\lambda \in \mathbb{C}$, $\mathcal{P}_0(z, \lambda) = f(z)g(z)$ and both f and g are non-monomial Laurent polynomials. Let

$$\tilde{f}(z) = f(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}), \tilde{g}(z) = g(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}). \quad (17)$$

By (16) and (17), one has that

$$\tilde{f}(z)\tilde{g}(z) = (-1)^Q \prod_{n \in W} \left(\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda \right). \quad (18)$$

When $d \geq 3$, for any $n \in W$ and fixed $\lambda \in \mathbb{C}$, $\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda$ is irreducible as a function of z . By the assumption that q_1, q_2, \dots, q_d are relatively prime, we have that for any $n \in W, n' \in W$ with $n \neq n'$, $\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda$ is not multiple of $\sum_{j=1}^d \left(\rho_{n'_j}^j z_j + \frac{1}{\rho_{n'_j}^j z_j} \right) + \lambda$. Since both $\tilde{f}(z)$ and $\tilde{g}(z)$ are unchanged under the action μ , we have that if $\tilde{f}(z)$ (or $\tilde{g}(z)$) has one factor $\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda$, then $\tilde{f}(z)$ (or $\tilde{g}(z)$) will have a factor

$$\prod_{n \in W} \left(\sum_{j=1}^d \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda \right).$$

This contradicts (18). \square

Proof of Theorem 2.2 with the zero potential. When $\lambda \neq 0$, it is easy to see that for any $n \in W$, $\sum_{j=1}^2 \left(\rho_{n_j}^j z_j + \frac{1}{\rho_{n_j}^j z_j} \right) + \lambda$ is irreducible as a function of z . By the similar proof of Theorem 2.1 with the zero potential, one has that $\mathcal{P}_0(z, \lambda)$ is irreducible. It suffices to prove the case that $\lambda = 0$. In this case, by (16),

$$\begin{aligned} \tilde{\mathcal{P}}_0(z, 0) &= (-1)^Q \prod_{n \in W} \left(\rho_{n_1}^1 z_1 + \rho_{n_2}^2 z_2 + \frac{1}{\rho_{n_1}^1 z_1} + \frac{1}{\rho_{n_2}^2 z_2} \right) \\ &= (-1)^Q \left(\prod_{n \in W} (\rho_{n_1}^1 z_1 + \rho_{n_2}^2 z_2) \right) \left(\prod_{n \in W} \left(1 + \frac{1}{\rho_{n_1}^1 \rho_{n_2}^2 z_1 z_2} \right) \right). \end{aligned} \quad (19)$$

Since both $\prod_{n \in W} (\rho_{n_1}^1 z_1 + \rho_{n_2}^2 z_2)$ and $\prod_{n \in W} \left(1 + \frac{1}{\rho_{n_1}^1 \rho_{n_2}^2 z_1 z_2} \right)$ are unchanged under μ , one has that there exist a polynomial $f(z_1, z_2)$ and a Laurent polynomial $g(z_1, z_2)$ such that

$$f(z_1^{q_1}, z_2^{q_2}) = \prod_{n \in W} (\rho_{n_1}^1 z_1 + \rho_{n_2}^2 z_2)$$

and

$$g(z_1^{q_1}, z_2^{q_2}) = \prod_{n \in W} \left(1 + \frac{1}{\rho_{n_1}^1 \rho_{n_2}^2 z_1 z_2} \right).$$

By the similar proof of Theorem 2.1, both f and g are irreducible. By (19), one has that $\mathcal{P}_0(z, 0) = (-1)^Q f(z)g(z)$. This completes the proof. \square

2.3. Ideas of the proof of Theorems 2.1 and 2.2. In this subsection, we review the ideas of proof of Theorems 2.1 and 2.2 in [40]. Let $A_1 = (0, 0, \dots, 0)$ (namely $z_1 = z_2 = \dots = z_d = 0$) and $A_2 = (0, 0, \dots, 0, \infty)$ (namely $z_1 = z_2 = \dots = z_{d-1} = 0, z_d^{-1} = 0$).

Assume that $\mathcal{P}(z, \lambda) = \prod_{j=1}^m f_j(z)$, where f_j is a (non-monomial) irreducible Laurent polynomial. Let

$$\tilde{f}_j(z) = f_j(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}), j = 1, 2, \dots, m \quad (20)$$

and hence

$$\tilde{\mathcal{P}}(z, \lambda) = \prod_{j=1}^m \tilde{f}_j(z). \quad (21)$$

Step 1.

Let

$$Z_{f_j} = \{z \in (\mathbb{C}^*)^d : f_j(z) = 0\}.$$

Let $z_1 = z_0^2, z_2 = z_3 = \dots = z_{d-1} = z_0$ and $z_0 \rightarrow 0$. Solving the equation $\tilde{\mathcal{P}}(z, \lambda) = 0$ and by (11), we have that either $z_d \rightarrow 0$ or $z_d^{-1} \rightarrow 0$. This implies that for any $j = 1, 2, \dots, m$, the closure (the closure is taken in $(\mathbb{C} \cup \{\infty\})^d$) of Z_{f_j} contains either A_1 or A_2 .

Step 2.

We are going to define “asymptotics” of $\mathcal{P}(z, \lambda) / \tilde{\mathcal{P}}(z, \lambda)$ at $A_1 = (0, 0, \dots, 0)$ and $A_2 = (0, \dots, 0, \infty)$. Let $\tilde{h}_1(z)$ be the lowest (homogeneous) degree component of $\tilde{\mathcal{P}}(z, \lambda)$, more precisely,

$$\tilde{h}_1(z) = (-1)^Q \prod_{n \in W} \left(\sum_{j=1}^d \frac{1}{\rho_{n_j}^j z_j} \right). \quad (22)$$

Let $\tilde{h}_2(z)$ be the lowest (homogeneous) degree component of $\tilde{\mathcal{P}}(z, \lambda)$ with respect to variables $z_1, z_2, \dots, z_{d-1}, z_d^{-1}$, more precisely,

$$\tilde{h}_2(z) = (-1)^Q \prod_{n \in W} \left(\rho_{n_d}^d z_d + \sum_{j=1}^{d-1} \frac{1}{\rho_{n_j}^j z_j} \right). \quad (23)$$

Since both $\tilde{h}_1(z)$ and $\tilde{h}_2(z)$ are unchanged under the action μ , there exist Laurent polynomials $h_1(z)$ and $h_2(z)$ such that

$$\tilde{h}_j(z) = h_j(z_1^{q_1}, z_2^{q_2}, \dots, z_d^{q_d}), j = 1, 2.$$

Like previous arguments in Section 2.2, it is not difficult to see that both $h_1(z)$ and $h_2(z)$ are irreducible.

Step 3.

Steps 1 and 2 allow us to conclude that $\mathcal{P}(z, \lambda)$ has at most two non-trivial irreducible factors. Assume indeed, $\mathcal{P}(z, \lambda)$ has two non-trivial irreducible factors, saying $f_1(z)$ and $f_2(z)$. Without loss of generality, assume the closure of Z_{f_j} contains A_j , $j = 1, 2$. From Step 2, one can see that $\tilde{f}_j(z)$ has “asymptotics” $\tilde{h}_j(z)$, $j = 1, 2$, namely up to monomials,

$$\tilde{f}_1(z) = \tilde{h}_1(z) + \text{higher degree terms of } z_1, z_2, \dots, z_{d-1}, z_d \quad (24)$$

and

$$\tilde{f}_2(z) = \tilde{h}_2(z) + \text{higher degree terms of } z_1, z_2, \dots, z_{d-1}, z_d^{-1}. \quad (25)$$

Finally, degree arguments enable us to show that two irreducible factors can only happen when $d = 2$ and both higher degree terms in (24) and (25) vanish, namely, up to monomials,

$$\tilde{f}_j(z) = \tilde{h}_j(z), j = 1, 2. \quad (26)$$

From (26), it is not difficult to obtain that $d = 2$ and $\lambda = [V]$.

We remark that the higher degree terms in (24) and (25) are with respect to different variables ($z_1, z_2, \dots, z_{d-1}, z_d$ or $z_1, z_2, \dots, z_{d-1}, z_d^{-1}$) and they may differ by monomials. Those obstacles make the degree arguments quite challenging.

Define the polynomial $\mathcal{P}_1(z, \lambda) = (-1)^Q z_1^{\frac{Q}{q_1}} z_2^{\frac{Q}{q_2}} \dots z_d^{\frac{Q}{q_d}} \mathcal{P}(z, \lambda)$. During the proof in [40], we need to constantly work between the polynomial $\mathcal{P}_1(z, \lambda)$ and the Laurent polynomial $\mathcal{P}(z, \lambda)$.

2.4. Open problems. Let $d = 2$. Assume that the Fermi variety $F_\lambda(V)$ is reducible at the average energy level $\lambda = [V]$. Does it imply the constancy of the potential V ?

It is not difficult to see that there exist non-constant complex valued functions V such that the Fermi variety is reducible at the energy level $[V]$. See [45] for example. However, for real-valued potentials, people believe the constant potential is the only case that the Fermi variety $F_\lambda(V)$ is reducible at some energy level, which has been formulated as a Conjecture by Gieseke, Knörrer and Trubowitz in the early 1990s [15].

Conjecture 3. [15, p.43] *Assume that V is a non-constant real periodic potential on \mathbb{Z}^2 . Then the Fermi variety $F_\lambda(V)/\mathbb{Z}^2$ is irreducible for any $\lambda \in \mathbb{C}$.*

In a recent note, we proved Conjecture 3 for separable potentials [45] (see next section for the definition of separable functions).

Let Δ_c be the continuous Laplacian on $L^2(\mathbb{R}^d)$:

$$(\Delta_c u)(x) = \sum_{j=1}^d \frac{\partial^2 u}{\partial^2 x_j}, x = (x_1, x_2, \dots, x_d) \in \mathbb{R}^d.$$

Let $V_c : \mathbb{T}^d = \mathbb{R}^d / \mathbb{Z}^d \rightarrow \mathbb{C}^d$ be a periodic function. Like discrete periodic Schrödinger operators, we can define the Fermi and Bloch varieties of continuous periodic Schrödinger operators $-\Delta_c + V_c$. Both Fermi and Bloch varieties are (principal) analytic sets [33]. Conjectures 1 and 2 are formulated for both continuous and discrete Schrödinger operators in [33, 35]. For continuous periodic Schrödinger operators, Knörrer and Trubowitz proved that the Bloch variety is irreducible (modulo periodicity) when $d = 2$ [31].

Motivated by Theorems 2.1 and 2.2 and Conjecture 3 for discrete periodic Schrödinger operators, we conjecture that

Conjecture 4. *Let $d \geq 3$. Then the Fermi variety of $-\Delta_c + V_c$ is irreducible (modulo periodicity) for any $\lambda \in \mathbb{C}$.*

Conjecture 5. *Let $d = 2$. Then the Fermi variety of $-\Delta_c + V_c$ is irreducible (modulo periodicity) for any $\lambda \in \mathbb{C} \setminus \{[V_c]\}$, where $[V_c] = \int_{\mathbb{T}^2} V_c(x) dx$.*

Conjecture 6. *Assume that V_c is a non-constant real periodic potential on \mathbb{R}^2 . Then the Fermi variety of $-\Delta_c + V_c$ is irreducible (modulo periodicity) for any $\lambda \in \mathbb{C}$.*

3. FERMISPECTRALITY

3.1. Isospectrality. In this section, we discuss the inverse problem of (2) and (3) with real periodic potentials. Two Γ -periodic potentials V and Y are called Floquet isospectral if

$$\sigma(D_V(k)) = \sigma(D_Y(k)), \text{ for any } k \in \mathbb{R}^d. \quad (27)$$

We say a function V on \mathbb{Z}^d is (d_1, d_2, \dots, d_r) separable (or simply separable), where $\sum_{j=1}^r d_j = d$, if there exist functions V_j on \mathbb{Z}^{d_j} , $j = 1, 2, \dots, r$, such that for any $(n_1, n_2, \dots, n_d) \in \mathbb{Z}^d$,

$$\begin{aligned} V(n_1, n_2, \dots, n_d) = & V_1(n_1, \dots, n_{d_1}) + V_2(n_{d_1+1}, n_{d_1+2}, \dots, n_{d_1+d_2}) \\ & + \dots + V_r(n_{d_1+d_2+\dots+d_{r-1}+1}, \dots, n_{d_1+d_2+\dots+d_r}). \end{aligned} \quad (28)$$

We say $V : \mathbb{Z}^d \rightarrow \mathbb{R}$ is completely separable if V is $(1, 1, \dots, 1)$ separable. When there is no ambiguity, we write (28) as $V = \bigoplus_{j=1}^r V_j$.

Our interest is first motivated by several questions asked by Eskin, Ralston and Trubowitz [9, 10], and Gordon and Kappeler [16] :

Q'1. If Y and V are Floquet isospectral, and Y is (completely) separable, is V (completely) separable?

Q'2. Assume that both $Y = \bigoplus_{j=1}^d Y_j$ and $V = \bigoplus_{j=1}^d V_j$ are completely separable. If V and Y are Floquet isospectral, are the one-dimensional potentials V_j and Y_j Floquet isospectral (up to possible translations)?

While Q'1 and Q'2 were formulated in the continuous case [9, 10, 16, 33], the same questions apply to the discrete case. For both continuous and discrete periodic Schrödinger operators, Q'1 and Q'2 have been partially answered by Eskin-Ralston-Trubowitz [9, 10], Gordon-Kappeler [16] and Kappeler [28]. We refer readers to [8–10, 16–18, 26–28, 33, 50, 59] and references therein for precise descriptions of those results and other related developments.

By (13), one has that

$$P_V(k) = \prod_{m=1}^Q (\lambda_V^m(k) - \lambda). \quad (29)$$

Floquet isospectrality of V and Y says that for any $k \in \mathbb{R}^d$, eigenvalues of eigen-equations (2) and (3) with potentials V and Y are the same. This implies that Floquet isospectrality of V and Y is equivalent to $F_\lambda(V) = F_\lambda(Y)$ for any $\lambda \in \mathbb{C}$ ($P_V(k) = P_Y(k)$ for any k ; Bloch varieties of V and Y are the same).

In [4], Bättig, Knörrer and Trubowitz studied several rigidity problems based on the Fermi variety when $d = 3$. In particular, they proved that for continuous periodic Schrödinger operators, if at some energy level, Fermi varieties of a potential V and the zero potential are the same, then V is zero. This motivated us to study the isospectrality based on Fermi varieties in [44], where *Fermi isospectrality* was first introduced.

Definition 7. [44] *Let V and Y be two Γ -periodic functions. We say V and Y are Fermi isospectral if $F_{\lambda_0}(V) = F_{\lambda_0}(Y)$ for some $\lambda_0 \in \mathbb{C}$.*

Motivated by Q'1 and Q'2, and the work of Bättig, Knörrer and Trubowitz [4], we asked two questions in [44]:

Q'3. If Y and V are *Fermi isospectral*, and Y is (completely) separable, is V (completely) separable?

Q'4. Assume that both $Y = \bigoplus_{j=1}^r Y_j$ and $V = \bigoplus_{j=1}^r V_j$ are separable. If V and Y are *Fermi isospectral*, are the lower dimensional potentials V_j and Y_j Floquet isospectral?

In the same paper [44], we established several rigidity theorems of Fermi isospectrality for discrete periodic Schrödinger operators. In particular, we answered Q'3 and Q'4 affirmatively for any dimension $d \geq 3$, and thus answered Q'1 and Q'2 as well.

3.2. Fermi isospectrality.

Theorem 3.1. [44] *Let $d \geq 3$. Assume that V and Y are Fermi isospectral, and Y is (d_1, d_2, \dots, d_r) separable, then V is (d_1, d_2, \dots, d_r) separable.*

As an immediate corollary, we have

Corollary 3.2. [44] *Let $d \geq 3$. Assume that V and Y are Fermi isospectral, and Y is completely separable, then V is completely separable.*

Theorem 3.3. [44] *Let $d = d_1 + d_2$ with $d_1 \geq 2$ and $d_2 \geq 1$. Assume that both V and Y are (d_1, d_2) separable, namely, there exist V_1, Y_1 on \mathbb{Z}^{d_1} and V_2, Y_2 on \mathbb{Z}^{d_2} such that $V = V_1 \oplus V_2$ and $Y = Y_1 \oplus Y_2$. Assume that Y and V are Fermi isospectral. Then, up to a constant, V_2 and Y_2 are Floquet isospectral.*

As a corollary of Theorems 3.1 and 3.3, we have

Corollary 3.4. [44] *Let $d \geq 3$. Assume that V and Y are Fermi isospectral, and $Y = \bigoplus_{j=1}^d Y_j$ is completely separable. Then $V = \bigoplus_{j=1}^d V_j$ is completely separable. Moreover, up to a constant Y_j and V_j , $j = 1, 2, \dots, d$, are Floquet isospectral.*

Theorem 3.5. [44] *Let $d \geq 3$. Assume that V and the zero potential are Fermi isospectral. Then V is zero.*

Remark 3. *For discrete periodic Schrödinger operators, Floquet isospectrality implies Fermi isospectrality, so all results in this section hold if we replace the assumption “Fermi isospectrality of Y and V ” with “Floquet isospectrality of Y and V ”.*

Let us talk about ideas of the proof in [44]. By the irreducibility results (Theorems 2.1 and 2.2), we have that if V and Y are Fermi isospectral, then $\mathcal{P}_V(z, \lambda) = \mathcal{P}_Y(z, \lambda)$ for some $\lambda \in \mathbb{C}$. Starting here, the proof in [44] focuses on the study of the Laurent polynomial $\mathcal{P}_V(z, \lambda)$ and $\mathcal{P}_Y(z, \lambda)$. We first prove

Theorem 3.6. [44] *Assume that V and Y are Fermi isospectral. Then*

$$[V] = [Y] \quad (30)$$

and for all possible $z \in \mathbb{C}^d$,

$$\sum_{n \in W, n' \in W} \frac{|\hat{V}(n - n')|^2}{(\sum_{j=1}^d \rho_{n_j}^j z_j)(\sum_{j=1}^d \rho_{n'_j}^j z_j)} \equiv \sum_{n \in W, n' \in W} \frac{|\hat{Y}(n - n')|^2}{(\sum_{j=1}^d \rho_{n_j}^j z_j)(\sum_{j=1}^d \rho_{n'_j}^j z_j)}. \quad (31)$$

The following proof depends on the study of linearly dependence/independence among hyperplanes $\sum_{j=1}^d \rho_{n_j}^j z_j = 0$, $n \in W$ and characterization of separable functions in the Fourier space. See [44] for details.

3.3. Open problems. Note that in Theorems 3.1, 3.3 and 3.5, we assume $d \geq 3$. It is natural to ask the following questions:

Q1. Let $d = 2$. If Y and V are Fermi isospectral, and Y is separable, is V separable?

- Q2. Let $d = 2$. Assume that both V and Y are separable, namely, there exist V_1, Y_1 on \mathbb{Z} and V_2, Y_2 on \mathbb{Z} such that $V = V_1 \oplus V_2$ and $Y = Y_1 \oplus Y_2$. Assume that Y and V are Fermi isospectral. Is it true that up to a constant, V_1 (V_2) and Y_1 (Y_2) are Floquet isospectral?
- Q3. Let $d = 2$. Assume that V and the zero potential are Fermi isospectral. Is it true that V is zero?

By Remark 2, one can see that the positive answer to Q3 implies Conjecture 3. From [45], we have that the positive answer to Q1 implies that to Q3.

4. EMBEDDED EIGENVALUES

4.1. Absence of embedded eigenvalues of perturbed periodic Schrödinger operators. Let us consider a perturbed periodic Schrödinger operators:

$$H = H_0 + V = -\Delta + V + v,$$

where V is periodic and v is a decaying function.

Recall that (see (14)) the spectrum of $H_0 = -\Delta + V$ has the band structure and there are no eigenvalues. We are interested in characterizing the perturbation v such that $-\Delta + V + v$ has no eigenvalues embedded into the spectral bands, which is referred to as the embedded eigenvalue problem.

For $d = 1$, the existence/absence of embedded eigenvalues has been understood very well [30, 38, 43, 46, 48, 49, 53, 55]. Problems of the existence of embedded eigenvalues in dimension $d \geq 2$ are a lot more complicated. The techniques of the generalized Prüfer transformation and oscillatory integrals developed for $d = 1$ are not available.

In [35], Kuchment and Vainberg introduced a new approach to study the embedded eigenvalue problem for perturbed periodic operators. It employs the analytic structure of the Fermi variety, unique continuation results, and techniques of several complex variables theory.

Condition 1: Given $\lambda \in \bigcup(a_m, b_m)$, we say that λ satisfies Condition 1 if any irreducible component of the Fermi variety at energy level λ contains an open analytic hypersurface of dimension $d - 1$ in \mathbb{R}^d .

Remark 4. For λ in the interior of a spectral band, the irreducibility of the Fermi variety at energy level λ implies Condition 1 for the same λ [33, 35, 36].

Theorem 4.1. [35] Let $d = 2, 3$, $H_0 = -\Delta_c + V_c$ and $H = -\Delta_c + V_c + v_c$. Assume that there exist constants $C > 0$ and $\gamma > 4/3$ such that

$$|v_c(x)| \leq Ce^{-|x|^\gamma}, x \in \mathbb{R}^d.$$

Assume the Condition 1 for some $\lambda \in \bigcup(a_m, b_m)$, where $\bigcup(a_m, b_m)$ is the spectral band of $-\Delta_c + V_c$. Then this λ can not be an eigenvalue of $H = -\Delta_c + V_c + v_c$.

The restriction on $d = 2, 3$ and the critical exponent $4/3$ arise from a quantitative unique continuation result. Suppose u is a solution of

$$-\Delta_c u + \tilde{V}u = 0 \text{ in } \mathbb{R}^d,$$

where $|\tilde{V}| \leq C$, $|u| \leq C$ and $u(0) = 1$. From the unique continuation principle, u cannot vanish identically on any open set. The quantitative result states [5]

$$\inf_{|x_0|=R} \sup_{|x-x_0|\leq 1} |u(x)| \geq e^{-CR^{4/3} \log R}. \quad (32)$$

A similar version of (32) was established in [51] (also see Remark 2.6 in [14]), namely, there is no non-trivial solution of $(-\Delta_c + \tilde{V})u = 0$ such that

$$|u(x)| \leq e^{-C|x|^{4/3}} \text{ for any } C > 0. \quad (33)$$

For complex-valued potentials \tilde{V} , the critical exponent $4/3$ in (32) and (33) is optimal by an example of Meshkov [51]. The Landis' conjecture states that the critical exponent is 1 for real potentials. Recently, the Landis' conjecture was proved when $d = 2$ by Logunov-Malinnikova-Nadirashvili-Nazarov [47]. Following the proof of Theorem 4.1 by Kuchment-Vainberg and the result of Logunov-Malinnikova-Nadirashvili-Nazarov [47], one could prove the following result.

Theorem 4.2. *Let $d = 2$. Assume that v_c is real and there exist constants $C > 0$ and $\gamma > 1$ such that*

$$|v_c(x)| \leq Ce^{-|x|^\gamma}, x \in \mathbb{R}^2.$$

Assume the Condition 1 for some $\lambda \in \bigcup(a_m, b_m)$, where $\bigcup(a_m, b_m)$ is the spectral band of $-\Delta_c + V_c$. Then this λ can not be an eigenvalue of $H = -\Delta_c + V_c + v_c$.

The unique continuation principle for discrete Laplacians is well known not to hold (see e.g., [21, 37]). In [40], we observed that a weak unique continuation result is sufficient for Kuchment-Vainberg's arguments in [35]. Moreover, the critical component can be improved from " $4/3$ " to " 1 ". Therefore, we established the discrete version of Theorem 4.1 for any dimension in [40].

Theorem 4.3. [40] *Assume V is a real-valued periodic function. Let $d \geq 2$. Assume that there exist constants $C > 0$ and $\gamma > 1$ such that the complex-valued function $v : \mathbb{Z}^d \rightarrow \mathbb{C}$ satisfies*

$$|v(n)| \leq Ce^{-|n|^\gamma}, n \in \mathbb{Z}^d. \quad (34)$$

Assume the Condition 1 for some $\lambda \in \bigcup_{m=1}^Q(a_m, b_m)$. Then this λ can not be an eigenvalue of $H = -\Delta + V + v$.

The irreducibility result (established in Theorems 2.1 and 2.2, and Remark 2) and Remark 4 allow us to prove that

Lemma 4.4. [40] *Assume V is a real-valued periodic function. Then the Condition 1 holds for every $\lambda \in \bigcup_{m=1}^Q(a_m, b_m)$.*

By Theorem 4.3 and Lemma 4.4, we have that

Theorem 4.5. [40] *Assume V is a real-valued periodic function. If there exist constants $C > 0$ and $\gamma > 1$ such that the complex-valued function $v : \mathbb{Z}^d \rightarrow \mathbb{C}$ satisfies*

$$|v(n)| \leq Ce^{-|n|^\gamma}, \quad (35)$$

then $H = -\Delta + V + v$ does not have any embedded eigenvalues, i.e., for any $\lambda \in \bigcup_{m=1}^Q(a_m, b_m)$, λ is not an eigenvalue of H .

Assume that V is zero. We can think that V is a periodic function on \mathbb{Z}^d with any q_1, q_2, \dots, q_d . Recall (15).

Lemma 4.6. [19, Lemmas 1.2 and 1.3] *Let $d \geq 2$. Then*

- *for any $\lambda \in (-2d, 2d) \setminus \{0\}$, $\lambda \in (a_m, b_m)$ for some $1 \leq m \leq Q$,*
- *if at least one of q_j 's is odd, then $0 \in (a_m, b_m)$ for some $1 \leq m \leq Q$.*

For $d = 2$, Lemma 4.6 was also proved in [7].

Theorem 4.5 and Lemma 4.6 imply

Corollary 4.7. [40] *Assume that there exist some $C > 0$ and $\gamma > 1$ such that*

$$|v(n)| \leq Ce^{-|n|^\gamma}. \quad (36)$$

Then $\sigma_p(-\Delta + v) \cap (-2d, 2d) = \emptyset$.

Remark 5. (1) *When $d = 1$, stronger results (replace (35) and (36) with $v(n) = \frac{o(1)}{1+|n|}$ as $n \rightarrow \infty$) in Theorem 4.5 and Corollary in 4.7 have been proved [43, 46, 55].*

(2) *Under a stronger assumption that v has compact support, Isozaki and Morioka proved that $\sigma_p(-\Delta + v) \cap (-2d, 2d) = \emptyset$ [20].*

Remark 6. (1) *For periodic (quantum) graph operators, the reducibility of Fermi varieties may occur. (quantum) graph operators may lack unique continuation results. There are examples showing that embedded eigenvalues (even with compactly supported eigenfunctions) happen under compactly supported perturbations [13, 34, 36, 37, 56, 57].*

(2) *For the continuous fourth order differential operator D^4 on $L^2(\mathbb{R})$, under a compactly supported perturbation v_c , $D^4 + v_c$ can have embedded eigenvalues [54].*

(3) *For the continuous fourth order differential operator on periodic hexagonal lattices, the reducibility of Fermi varieties occurs [11].*

4.2. Open problems. We list a few questions here.

Q4. Let v_c be a real function on \mathbb{R}^d . Assume that the Landis' conjecture holds and the Condition 1 holds for some $\lambda \in \bigcup(a_m, b_m)$. Assume that there exist constants $C > 0$ and $\gamma > 1$ such that

$$|v_c(x)| \leq Ce^{-|x|^\gamma}.$$

Then following the argument of Kuchment-Vainberg [35], λ can not be an eigenvalue of $H = -\Delta_c + V_c + v_c$. We ask whether or not we could find an alternative proof which does not depend on Landis' conjecture or Condition 1.

- Q5. For continuous Schrödinger operators, when the periodic function $V_c = 0$, Kato [29] proved that $-\Delta_c + v_c$ does not have embedded eigenvalues if $\frac{o(1)}{1+|x|}$ as $x \rightarrow \infty$ (the actual result is quantitative). Motivated by results of Kato and one dimensional case (see (1) in Remark 5), we ask is it true that $-\Delta + V + v$ ($-\Delta_c + V_c + v_c$) does not have any embedded eigenvalues if $|v(n)| \leq \frac{C}{|n|^K}$, $n \in \mathbb{Z}^d$ ($|v_c(x)| \leq \frac{C}{|x|^K}$, $x \in \mathbb{R}^d$) for some $K, C > 0$? We should mention that for the discrete case, this is open even for $V = 0$.
- Q6. Construct properly decaying functions v (v_c) such that $-\Delta + V + v$ ($-\Delta_c + V_c + v_c$) has embedded eigenvalues. Here “properly decaying” means exponentially decaying, algebraic decaying or something similar. When $d = 1$, there are many examples of perturbed periodic Schrödinger operators with (dense)embedded eigenvalues [22–25, 41–43, 46, 52, 58]. The construction is quite challenging when $d \geq 2$ and the underlying periodic functions (V or V_c) are non-zero. For the continuous case, when $V_c = 0$, it reduces to an one dimensional problem by choosing radial functions $v_c(x) = v_c(r)$, where $r = \sqrt{x_1^2 + x_2^2 + \cdots + x_d^2}$. For the discrete case, when $V = 0$, it can be done by taking separable functions $v(n) = v_1(n_1) + v_2(n_2) + \cdots + v_d(n_d)$.

ACKNOWLEDGMENTS

W. Liu was supported by NSF DMS-2000345 and DMS-2052572.

REFERENCES

- [1] D. Bättig. *A toroidal compactification of the two dimensional Bloch-manifold*. PhD thesis, ETH Zurich, 1988.
- [2] D. Bättig. A directional compactification of the complex Fermi surface and isospectrality. In *Séminaire sur les Équations aux Dérivées Partielles, 1989–1990*, pages Exp. No. IV, 11. École Polytech., Palaiseau, 1990.
- [3] D. Bättig. A toroidal compactification of the Fermi surface for the discrete Schrödinger operator. *Comment. Math. Helv.*, 67(1):1–16, 1992.
- [4] D. Bättig, H. Knörrer, and E. Trubowitz. A directional compactification of the complex Fermi surface. *Compositio Math.*, 79(2):205–229, 1991.
- [5] J. Bourgain and C. E. Kenig. On localization in the continuous Anderson-Bernoulli model in higher dimension. *Invent. Math.*, 161(2):389–426, 2005.
- [6] N. Do, P. Kuchment, and F. Sottile. Generic properties of dispersion relations for discrete periodic operators. *J. Math. Phys.*, 61(10):103502, 19, 2020.

- [7] M. Embree and J. Fillman. Spectra of discrete two-dimensional periodic Schrödinger operators with small potentials. *J. Spectr. Theory*, 9(3):1063–1087, 2019.
- [8] G. Eskin. Inverse spectral problem for the Schrödinger equation with periodic vector potential. *Comm. Math. Phys.*, 125(2):263–300, 1989.
- [9] G. Eskin, J. Ralston, and E. Trubowitz. On isospectral periodic potentials in \mathbf{R}^n . *Comm. Pure Appl. Math.*, 37(5):647–676, 1984.
- [10] G. Eskin, J. Ralston, and E. Trubowitz. On isospectral periodic potentials in \mathbf{R}^n . II. *Comm. Pure Appl. Math.*, 37(6):715–753, 1984.
- [11] M. Ettehad and B. Hatinoglu. On the Spectra of Periodic Elastic Beam Lattices: Single-Layer Graph. *arXiv preprint arXiv:2110.05466*, 2021.
- [12] J. Fillman, W. Liu, and R. Matos. Irreducibility of the Bloch variety for finite-range Schrödinger operators. *arXiv preprint arXiv:2107.06447*, 2021.
- [13] L. Fisher, W. Li, and S. P. Shipman. Reducible Fermi surface for multi-layer quantum graphs including stacked graphene. *Comm. Math. Phys.*, 385(3):1499–1534, 2021.
- [14] R. Froese, I. Herbst, M. Hoffmann-Ostenhof, and T. Hoffmann-Ostenhof. L^2 -lower bounds to solutions of one-body Schrödinger equations. *Proc. Roy. Soc. Edinburgh Sect. A*, 95(1-2):25–38, 1983.
- [15] D. Gieseke, H. Knörrer, and E. Trubowitz. *The geometry of algebraic Fermi curves*, volume 14 of *Perspectives in Mathematics*. Academic Press, Inc., Boston, MA, 1993.
- [16] C. S. Gordon and T. Kappeler. On isospectral potentials on tori. *Duke Math. J.*, 63(1):217–233, 1991.
- [17] C. S. Gordon and T. Kappeler. On isospectral potentials on flat tori. II. *Comm. Partial Differential Equations*, 20(3-4):709–728, 1995.
- [18] V. Guillemin. Inverse spectral results on two-dimensional tori. *J. Amer. Math. Soc.*, 3(2):375–387, 1990.
- [19] R. Han and S. Jitomirskaya. Discrete Bethe-Sommerfeld conjecture. *Comm. Math. Phys.*, 361(1):205–216, 2018.
- [20] H. Isozaki and H. Morioka. A Rellich type theorem for discrete Schrödinger operators. *Inverse Probl. Imaging*, 8(2):475–489, 2014.
- [21] S. Jitomirskaya. Ergodic Schrödinger operators (on one foot). In *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon's 60th birthday*, volume 76 of *Proc. Sympos. Pure Math.*, pages 613–647. Amer. Math. Soc., Providence, RI, 2007.
- [22] S. Jitomirskaya and W. Liu. Noncompact complete Riemannian manifolds with dense eigenvalues embedded in the essential spectrum of the Laplacian. *Geom. Funct. Anal.*, 29(1):238–257, 2019.
- [23] E. Judge, S. Naboko, and I. Wood. Eigenvalues for perturbed periodic Jacobi matrices by the Wigner–von Neumann approach. *Integral Equations Operator Theory*, 85(3):427–450, 2016.

- [24] E. Judge, S. Naboko, and I. Wood. Embedded eigenvalues for perturbed periodic Jacobi operators using a geometric approach. *J. Difference Equ. Appl.*, 24(8):1247–1272, 2018.
- [25] E. Judge, S. Naboko, and I. Wood. Spectral results for perturbed periodic Jacobi matrices using the discrete Levinson technique. *Studia Math.*, 242(2):179–215, 2018.
- [26] T. Kappeler. On isospectral periodic potentials on a discrete lattice. I. *Duke Math. J.*, 57(1):135–150, 1988.
- [27] T. Kappeler. On isospectral potentials on a discrete lattice. II. *Adv. in Appl. Math.*, 9(4):428–438, 1988.
- [28] T. Kappeler. Isospectral potentials on a discrete lattice. III. *Trans. Amer. Math. Soc.*, 314(2):815–824, 1989.
- [29] T. Kato. Growth properties of solutions of the reduced wave equation with a variable coefficient. *Comm. Pure Appl. Math.*, 12:403–425, 1959.
- [30] A. Kiselev, C. Remling, and B. Simon. Effective perturbation methods for one-dimensional Schrödinger operators. *J. Differential Equations*, 151(2):290–312, 1999.
- [31] H. Knörrer and E. Trubowitz. A directional compactification of the complex Bloch variety. *Comment. Math. Helv.*, 65(1):114–149, 1990.
- [32] H. Krueger. Periodic and limit-periodic discrete Schrödinger operators. *arXiv preprint arXiv:1108.1584*, 2011.
- [33] P. Kuchment. An overview of periodic elliptic operators. *Bull. Amer. Math. Soc. (N.S.)*, 53(3):343–414, 2016.
- [34] P. Kuchment and O. Post. On the spectra of carbon nano-structures. *Comm. Math. Phys.*, 275(3):805–826, 2007.
- [35] P. Kuchment and B. Vainberg. On absence of embedded eigenvalues for Schrödinger operators with perturbed periodic potentials. *Comm. Partial Differential Equations*, 25(9-10):1809–1826, 2000.
- [36] P. Kuchment and B. Vainberg. On the structure of eigenfunctions corresponding to embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 268(3):673–686, 2006.
- [37] P. A. Kuchment. On the Floquet theory of periodic difference equations. In *Geometrical and algebraical aspects in several complex variables (Cetraro, 1989)*, volume 8 of *Sem. Conf.*, pages 201–209. EditEl, Rende, 1991.
- [38] P. Kurasov and S. Naboko. Wigner-von Neumann perturbations of a periodic potential: spectral singularities in bands. *Math. Proc. Cambridge Philos. Soc.*, 142(1):161–183, 2007.
- [39] W. Li and S. P. Shipman. Irreducibility of the Fermi surface for planar periodic graph operators. *Lett. Math. Phys.*, 110(9):2543–2572, 2020.
- [40] W. Liu. Irreducibility of the Fermi variety for discrete periodic Schrödinger operators and embedded eigenvalues. *Geom. Funct. Anal.* to appear.

- [41] W. Liu. Criteria for eigenvalues embedded into the absolutely continuous spectrum of perturbed Stark type operators. *J. Funct. Anal.*, 276(9):2936–2967, 2019.
- [42] W. Liu. Sharp bounds for finitely many embedded eigenvalues of perturbed Stark type operators. *Math. Nachr.*, 293(9):1776–1790, 2020.
- [43] W. Liu. Criteria for Embedded Eigenvalues for Discrete Schrödinger Operators. *Int. Math. Res. Not. IMRN*, (20):15803–15832, 2021.
- [44] W. Liu. Fermi isospectrality for discrete periodic Schrödinger operators. *arXiv preprint arXiv:2106.03726*, 2021.
- [45] W. Liu. Fermi isospectrality of discrete periodic Schrödinger operators with separable potentials on \mathbb{Z}^2 . Preprint. 2021.
- [46] W. Liu and D. C. Ong. Sharp spectral transition for eigenvalues embedded into the spectral bands of perturbed periodic operators. *J. Anal. Math.*, 141(2):625–661, 2020.
- [47] A. Logunov, E. Malinnikova, N. Nadirashvili, and F. Nazarov. The landis conjecture on exponential decay. *arXiv preprint arXiv:2007.07034*, 2020.
- [48] M. Lukic. Schrödinger operators with slowly decaying Wigner-von Neumann type potentials. *J. Spectr. Theory*, 3(2):147–169, 2013.
- [49] M. Lukic and D. C. Ong. Wigner-von Neumann type perturbations of periodic Schrödinger operators. *Trans. Amer. Math. Soc.*, 367(1):707–724, 2015.
- [50] H. P. McKean and E. Trubowitz. Hill’s operator and hyperelliptic function theory in the presence of infinitely many branch points. *Comm. Pure Appl. Math.*, 29(2):143–226, 1976.
- [51] V. Z. Meshkov. On the possible rate of decrease at infinity of the solutions of second-order partial differential equations. *Mat. Sb.*, 182(3):364–383, 1991.
- [52] S. Naboko. Dense point spectra of Schrödinger and dirac operators. *Theoretical and Mathematical Physics*, 68(1):646–653, 1986.
- [53] S. Naboko and S. Simonov. Zeroes of the spectral density of the periodic Schrödinger operator with Wigner-von Neumann potential. *Math. Proc. Cambridge Philos. Soc.*, 153(1):33–58, 2012.
- [54] V. G. Papanicolaou. Examples of fourth-order scattering-type operators with embedded eigenvalues in their continuous spectra. *arXiv preprint arXiv:2110.00739*, 2021.
- [55] F. S. Roze-Beketov. A finiteness test for the number of discrete levels which can be introduced into the gaps of the continuous spectrum by perturbations of a periodic potential. *Dokl. Akad. Nauk SSSR*, 156:515–518, 1964.
- [56] S. P. Shipman. Eigenfunctions of unbounded support for embedded eigenvalues of locally perturbed periodic graph operators. *Comm. Math. Phys.*, 332(2):605–626, 2014.
- [57] S. P. Shipman. Reducible Fermi surfaces for non-symmetric bilayer quantum-graph operators. *J. Spectr. Theory*, 10(1):33–72, 2020.

- [58] B. Simon. Some Schrödinger operators with dense point spectrum. *Proc. Amer. Math. Soc.*, 125(1):203–208, 1997.
- [59] A. Waters. Isospectral periodic torii in dimension 2. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 32(6):1173–1188, 2015.

(Wencai Liu) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

Email address: liuwencai1226@gmail.com; wencail@tamu.edu