

On the M2–Brane Index on Noncommutative Crepant Resolutions

Michele Cirafici

*Dipartimento di Matematica e Geoscienze, Università di Trieste,
Via A. Valerio 12/1, I-34127 Trieste, Italy,
Institute for Geometry and Physics (IGAP), via Beirut 2/1, 34151, Trieste, Italy
INFN, Sezione di Trieste, Trieste, Italy
Email: michelecirafici@gmail.com*

Abstract

On certain M-theory backgrounds which are a circle fibration over a smooth Calabi-Yau the quantum theory of M2 branes can be studied in terms of the K-theoretic Donaldson-Thomas theory on the threefold. We extend this relation to noncommutative crepant resolutions. In this case the threefold develops a singularity and classical smooth geometry is replaced by the algebra of paths of a certain quiver. K-theoretic quantities on the quiver representation moduli space can be computed via toric localization and result in certain rational functions of the toric parameters. We discuss in particular the case of the conifold and certain orbifold singularities.

September 3, 2022

Contents

1	Introduction	1
2	The M2-brane index for local threefolds	2
2.1	BPS states in type IIA	3
2.2	The M2-brane index at large radius	4
3	Noncommutative algebraic geometry	7
3.1	Quivers and noncommutative crepant resolutions	8
3.2	Noncommutative Donaldson-Thomas theory	9
4	The noncommutative conifold	11
4.1	Geometry and BPS states	11
4.2	Toric action and fixed points	12
4.3	Localization	14
4.4	Some explicit fixed points	15
4.5	A conjecture for the full partition function	18
4.6	Refined and unrefined limits	19
5	Orbifold Singularities	21
5.1	BPS states and McKay quivers	21
5.2	Localization on quiver moduli spaces	22
5.3	The case of A-fibered singularities	24
5.4	The case $\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2$	29
5.5	The $\mathbb{C}^3/\mathbb{Z}_3$ orbifold	32
6	Discussion	33
6.1	Towards M2 branes on NCCRs	33
6.2	Some open problems	35

1 Introduction

One of the most interesting aspects of string/M-theory is that it is capable of describing quantum degrees of freedom in spacetimes where the manifold structure breaks down. For example in the moduli spaces of vacua of Calabi-Yau compactifications one can generically find points where the manifold develops a singularity. The local dynamics near such a point is often captured by a non-compact threefold. For example the quintic contains a conifold point, where a rigid rational curve shrinks to zero size. The local dynamics can be studied using strings on the resolved conifold, by sending to zero the Kähler volume of the curve.

The situation is less clear in M-theory where we lack a complete description of the fundamental degrees of freedom. In [28, 30] it was proposed that certain aspects of the theory, namely the supersymmetric index of the membrane, can be computed explicitly using K-theoretic Donaldson-Thomas theory on a smooth Calabi-Yau. Computing an index is equivalent to enumerating supersymmetry protected states with certain quantum numbers and therefore provides a window on the fundamental degrees of freedom and their supersymmetric excitations.

In this note we study K-theoretic Donaldson-Thomas theory defined on noncommutative crepant resolutions of Calabi-Yau singularities as a model for the M2-brane index in M-theory. Recent progress in K-theoretic countings of BPS states have been made in the context of smooth geometries and fourfolds [2, 3, 12, 13, 16, 19, 25, 29, 31, 34] as well as defects in quantum field theory [5].

In many cases when a toric threefold develops a singularity the local geometry is replaced by algebraic quantities. This is the case for the conifold or for resolved orbifolds, when the orbifold group is subgroup of $SU(3)$. In this examples there exists a noncommutative algebra A which contains the local singularity as its center. The algebra itself plays the role of a resolution of the singularity and it is called a noncommutative crepant resolution [36]. D-branes can be consistently studied in such backgrounds. Indeed the derived category of modules over A is derived equivalent to the derived category of coherent sheaves over the smooth threefold. In other words such backgrounds are required for the consistency of the compactification.

The algebra A can be seen as the Jacobian path algebra of a quiver with relations. The low energy dynamics of BPS states can be described in terms of a certain quantum mechanical model associated with this quiver. When one restricts attention to BPS states formed by a gas of D2/D0 branes bounded to a single D6 brane the computation can be carried on exactly. The presence of the D6 brane modifies the quiver via a framing node. The toric symmetry of the original threefold descend to the framed quantum mechanics model and its partition function, the index of D-brane states, can be computed directly via toric localization in many cases [35, 7].

For toric threefolds the enumeration of BPS bound states in the type II string at large radius and on noncommutative crepant resolutions are related by wall crossing [18, 27]. Consistency requires that a similar relation should hold in K-theoretic enumerative problems. The large radius index for membranes can be computed within the formalism of [30]. In this proposal the enumeration of supersymmetric membrane states is again given in terms of a melting crystal, where each crystal configuration is weighted by a certain function. The result depends explicitly on the toric parameters of the large radius threefold.

We expect that a similar computation should exist on M-theory backgrounds where the toric threefold develops a singularity and the geometric setup is replaced by the algebra A . In this note we develop a formalism to carry out these computations and conjecture explicit closed forms for the membrane partition function, as plethystic exponentials of rational functions. We postpone the analysis of the wall crossing behaviour of the membrane index to a follow up paper [11].

This note is organized as follows. In Section 2 we review BPS states on local threefolds and the M2-brane index. Section 3 contains some background material on noncommutative crepant resolutions. In Section 4 and 5 we compute the M2-brane index for the noncommutative conifold and orbifold singularities respectively. In Section 6 we conclude with a discussion and some open problems.

To make the paper easier to read the computations are detailed explicitly in the supporting MATHEMATICA files.

2 The M2-brane index for local threefolds

In this section we briefly review BPS states on local threefolds and the M2-brane index construction of [30]. BPS bound states for toric Calabi-Yau are captured by the localization of a certain gauge

theory living on the D6 brane worldvolume [17, 33]. In particular for bound states with a single D6 branes, this is Donaldson-Thomas theory of ideal sheaves. The proposal of [30] is that the K-theoretic version of this story captures the membrane contribution to the index of M-theory.

2.1 BPS states in type IIA

A concrete way to set up BPS counting problems on toric threefolds is to consider the type IIA string on a compact Calabi-Yau and then take a local limit. After the first step the resulting low energy theory is $\mathcal{N} = 2$ supergravity and its BPS states can be engineered by wrapping D-branes on holomorphic cycles of the Calabi-Yau geometry. The theory has a moduli space of vacua which parametrizes complexified Kähler deformations of the Calabi-Yau manifold. Which BPS states are physically realized as stable states depends on the Kähler moduli. The Kähler moduli space has an intricate chamber structure. As the Kähler moduli are varied one will generically cross walls of marginal stability and the physical BPS spectrum will change according to the wall-crossing formula for generalized Donaldson-Thomas invariants [20].

Ordinary Donaldson-Thomas theory concerns the large radius chamber, where supersymmetric BPS states are bound states of D6-D4-D2-D0 branes labelled by a charge vector $\gamma \in \Gamma = H^{\text{even}}(X, \mathbb{Z})$ associated via Poincaré duality to holomorphic cycles in $H_{\text{even}}(X, \mathbb{Z})$. Choosing an electro-magnetic splitting one can write

$$\gamma = p^0 + P^A D_A + Q_A \tilde{D}^A + q_0 dV \quad (2.1)$$

where $\{D_A\}$ is a basis of $H^2(X; \mathbb{Z})$, $\{\tilde{D}^A\}$ the dual basis of $H^4(X; \mathbb{Z})$ and dV the volume element. The lattice of charges is endowed with a symplectic product

$$\langle \gamma, \gamma' \rangle_{\Gamma} = \int_X \gamma \wedge (-1)^{\text{deg}/2} \gamma' . \quad (2.2)$$

In the large radius limit one can neglect worldsheet instanton corrections and the holomorphic central charge is given by

$$Z_X(\gamma; t) = - \int_X \gamma \wedge e^{-t} \quad (2.3)$$

where $t = B + iJ$ is the complexified Kähler modulus consisting of the background supergravity two-form B -field and the Kähler $(1, 1)$ -form J of X . Physically the theory can be understood from the point of view of the D6-branes as a $U(n)$ supersymmetric Yang-Mills theory which localizes onto solutions of the Donaldson-Uhlenbeck-Yau equations.

At this point of the moduli space the theory is well defined in full generality for an arbitrary charge vector and a compact threefold [15]. For toric threefolds and a particular choice of the charge vector, a single D6 brane and no D4 branes, the gauge theory becomes abelian and the enumerative problem of Donaldson-Thomas invariants is equivalent to the enumerative problem of Gromov-Witten invariants, and therefore to the topological string [22]. As the moduli are varied we will cross walls of marginal stability and the enumerative problem will change accordingly.

To properly define non-compact D6 branes one needs to take the limit from a compact Calabi-Yau; then this construction results in a certain extension of the local Kähler moduli space [18]. By moving through a path in this space of parameters we will generically cross many walls of marginal stability

producing new BPS state counting problems. In this paper we will only focus on one such chamber, the noncommutative crepant resolution chamber. A more detailed study of wall-crossing properties will appear elsewhere [11].

2.2 The M2-brane index at large radius

The above discussion can be lifted to M-theory by considering a background \tilde{Z} which is a S^1 bundle over a Calabi-Yau 5-fold Z . To begin with, one can pick Z to be $X \times \mathbb{C}^2$, with X a smooth toric Calabi-Yau threefold. Then such an M-theory background can be explicitly constructed as the total space of this fibration and it includes a factor with a Taub-NUT metric, whose $U(1)$ isometry is identified with the M-theory circle S^1 . The S^1 fibration induces a \mathbb{C}_q^\times action which acts with weights q and q^{-1} on the \mathbb{C}^2 factor and leaves $X \subset Z$ fixed.

The fundamental insight of [30] is that the M2 brane index for M-theory on \tilde{Z} can be recast in terms of the K-theoretic Donaldson-Thomas theory of Z . This can be understood as a generalization of the ordinary moduli problem of BPS states counting. Roughly speaking this is usually formulated in terms of a moduli space M obtained as the locus $s^{-1}(0) \subset \tilde{M}$ where s is a section of \mathcal{E} , the obstruction bundle. The structure sheaf \mathcal{O}_M is the 0-th cohomology of the Koszul complex

$$0 \rightarrow \bigwedge^{rk} \mathcal{E}^\vee \rightarrow \cdots \rightarrow \bigwedge^2 \mathcal{E}^\vee \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O}_{\tilde{M}} \rightarrow 0. \quad (2.4)$$

For the purpose of virtual countings it is better to introduce the virtual sheaf $\mathcal{O}_M^{\text{vir}}$ as [1]

$$\mathcal{O}_M^{\text{vir}} = [\bigwedge^{rk} \mathcal{E}^\vee \rightarrow \cdots \rightarrow \bigwedge^2 \mathcal{E}^\vee \rightarrow \mathcal{E}^\vee \rightarrow \mathcal{O}_{\tilde{M}}]. \quad (2.5)$$

However in physics we typically have a complex constructed out of a Dirac operator $\not{D} = \bar{\partial} + \bar{\partial}^*$ which acts on the spin bundles

$$\mathcal{S}_\pm = \mathcal{K}_M^{1/2} \otimes \bigotimes_{n \text{ even/odd}} \Omega_M^{0,n} \quad (2.6)$$

where $\mathcal{K}_M^{1/2}$ is the square root of the canonical bundle. In this setup the relevant object is the symmetrized structure sheaf $\hat{\mathcal{O}}_M^{\text{vir}} = (-1)^n \mathcal{K}_{\text{vir}}^{1/2} \otimes \mathcal{O}_M^{\text{vir}}$ where now $\mathcal{K}_{\text{vir}} = \kappa^{\dim \tilde{M}/2} \otimes \mathcal{K}_{\tilde{M}}|_M$ is the determinant of the virtual tangent bundle $T_M^{\text{vir}} = T_{\tilde{M}} - \kappa \otimes T_M^*|_M$. In more general situations this sheaf can be generalized and the relevant object to consider is the tensor product of $\hat{\mathcal{O}}^{\text{vir}}$ with a tautological bundle. See [32] for a recent review.

To be concrete we will now specialize to the case \mathbb{C}^3 . In this case the relevant moduli space is the Hilbert scheme of points $\text{Hilb}(\mathbb{C}^3, n)$. We can see $\text{Hilb}(\mathbb{C}^3, n)$ as the locus in $\text{Hilb}(\text{free}_3, n)$ (the space of $n \times n$ matrices $B_1, B_2, B_3 \in \text{End}(\mathbb{C}^3)$ together with a cyclic vector $v \in \mathbb{C}^3$ and modulo the action of $GL(n)$) cut out by the critical locus $s = \partial W = 0$, with

$$W = \text{tr}(B_1 B_2 B_3 - B_1 B_3 B_2) \quad (2.7)$$

The toric action on Z lifts to $\text{Hilb}(\mathbb{C}^3, n)$ by rescaling: the torus $\mathbb{T} = (t_1, t_2, t_3)$ acts as $B_i \rightarrow B_i t_i$

for $i = 1, 2, 3$. The K-theoretic Donaldson-Thomas partition function is then defined as

$$Z_{\mathbb{C}^3}(q, \{t_i\}) := \chi \left(\mathbb{M}, \widehat{\mathcal{O}}^{\text{vir}} \right) = \sum_n (-q)^n \chi \left(\text{Hilb}(\mathbb{C}^3, n), \widehat{\mathcal{O}}^{\text{vir}} \right) \quad (2.8)$$

where $\mathbb{M} = \bigsqcup_n \text{Hilb}(\mathbb{C}^3, n)$. The counting parameter q can be introduced by a minor modification of the symmetrized virtual structure sheaf.

This partition function can be computed directly via virtual localization as follows. The information on deformations and obstructions of the moduli space is encoded in the virtual tangent space

$$T_\pi^{\text{vir}} = \sum a_i - \sum b_i = \sum w_i - \sum \kappa/w_i \quad (2.9)$$

where a_i and b_i are generic placeholders for the toric weights of the deformations and the obstructions. In the case at hand these have the form w_i and κ/w_i , with w_i monomials in the toric weights. In localized K-theory we can write

$$\mathcal{O}_{\text{Hilb}(\mathbb{C}^3, n)}^{\text{vir}} = \sum_{|\pi|=n} \mathcal{O}_{I_\pi} \prod \frac{1 - \frac{w}{\kappa}}{1 - w^{-1}} \quad (2.10)$$

where fixed points in $\text{Hilb}(\mathbb{C}^3, n)$ are labelled by monomial ideals I_π associated with three dimensional partitions π with a fixed number of boxes $|\pi| = n$.

The symmetrized virtual structure sheaf is now $\widehat{\mathcal{O}}^{\text{vir}} = \mathcal{O}^{\text{vir}} \otimes (\mathcal{K}^{\text{vir}})^{1/2}$ where

$$\mathcal{K}^{\text{vir}} = \frac{\det \text{Obs}}{\det \text{Def}} = \frac{\det(T^* \widetilde{M} \otimes \kappa)}{\det T \widetilde{M}}. \quad (2.11)$$

In the case at hand for $\text{Hilb}(\mathbb{C}^3, n)$

$$T_{\widetilde{M}} = (\mathbb{C}^3 - 1) \otimes \overline{V} \otimes V + V \quad (2.12)$$

and therefore

$$T_\pi^{\text{vir}} = T_{\widetilde{M}} - \kappa \otimes T_{\widetilde{M}}^* = V - t_1 t_2 t_3 \overline{V} - V \otimes \overline{V} (1 - t_1)(1 - t_2)(1 - t_3) \quad (2.13)$$

where $\kappa = t_1 t_2 t_3$ and

$$V = \sum_{(i,j,k) \in \pi} t_1^{-i} t_2^{-j} t_3^{-k} \quad (2.14)$$

is the character of the subscheme \mathcal{O}_Z associated with the \mathbb{T} -fixed ideal sheaf \mathcal{J}_π .

Here the square root can be computed as the top exterior power of the dual of the virtual tangent bundle and has the structure of ratio of determinants of obstructions and deformations

$$\mathcal{K}_{\text{vir}}^{1/2} = \det^{-1/2} T_\pi^{\text{vir}} = \frac{\det^{1/2} \text{Obs}}{\det^{1/2} \text{Def}} = \frac{\prod_i \sqrt{\frac{\kappa}{w_i}}}{\prod_i \sqrt{w_i}} \quad (2.15)$$

The net effect of the twist by $\mathcal{K}^{1/2}$ is to replace the contribution of each fixed point by

$$\frac{1 - w/\kappa}{1 - w^{-1}} \rightarrow \frac{(k/w)^{1/2} - (w/k)^{1/2}}{w^{1/2} - w^{-1/2}} = \widehat{\mathbf{a}}(w - \kappa/w) \quad (2.16)$$

We have defined the Okounkov function

$$\widehat{\mathbf{a}}(w) = \frac{1}{w^{1/2} - w^{-1/2}} \quad (2.17)$$

which enjoys the properties $\widehat{\mathbf{a}}(w_1 + w_2) = \widehat{\mathbf{a}}(w_1)\widehat{\mathbf{a}}(w_2)$ and in particular $\widehat{\mathbf{a}}(2w) = \widehat{\mathbf{a}}(w)^2$.

At a \mathbb{T} -fixed point the virtual tangent space decomposition can be written schematically as in (2.9); then by using the Okounkov function we can write

$$\widehat{\mathbf{a}}(T_\pi^{\text{vir}}) = \prod_w \frac{(\kappa/w)^{1/2} - (w/\kappa)^{1/2}}{w^{1/2} - w^{-1/2}} \quad (2.18)$$

so that

$$\widehat{\mathcal{O}}_{\text{Hilb}(\mathbb{C}^3, n)}^{\text{vir}} = \sum_{|\pi|=n} (-1)^n \mathcal{O}_{I_\pi} \widehat{\mathbf{a}}(T_\pi^{\text{vir}}) \quad (2.19)$$

in localized K-theory. Following [30] the full result can be written in closed form

$$Z_{\mathbb{C}^3}(q; \{t_i\}) = \sum_n (-q)^n \chi(\text{Hilb}(\mathbb{C}^3, n), \widehat{\mathcal{O}}^{\text{vir}}) = \mathbf{S}^\bullet \frac{\prod_{i < j \leq 3} (t_i t_j)^{1/2} - (t_i t_j)^{-1/2}}{\prod_i^5 (t_i^{1/2} - t_i^{-1/2})} \quad (2.20)$$

as the generating function of the degree 0 K-theoretic Donaldson-Thomas invariants of \mathbb{C}^3 . In the right hand side the toric weights t_4 and t_5 act on the \mathbb{C}^2 side of the fivefold $Z = \mathbb{C}^5$. They are related to the left hand side parameters by $t_4 = q/\sqrt{\kappa}$ and $t_5 = 1/(q\sqrt{\kappa})$, with $\kappa = t_1 t_2 t_3$.

The symmetrization operation \mathbf{S}^\bullet , or plethystic exponential, is defined as follows. We introduce the Nekrasov function

$$\mathcal{F}_{\mathbb{C}^3}[t_1, t_2, t_3, t_4, t_5] := \frac{\prod_{i < j \leq 3} (t_i t_j)^{1/2} - (t_i t_j)^{-1/2}}{\prod_i^5 (t_i^{1/2} - t_i^{-1/2})} \quad (2.21)$$

and then set

$$\mathbf{S}^\bullet \mathcal{F}_{\mathbb{C}^3}[t_1, t_2, t_3, t_4, t_5] := \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} \mathcal{F}_{\mathbb{C}^3}[t_1^n, t_2^n, t_3^n, t_4^n, t_5^n] \right) \quad (2.22)$$

The full computation gives (schematically)

$$Z_{\mathbb{C}^3}(q; \{t_i\}) = \sum (-q)^{|\pi|} \prod_i \frac{\sqrt{\frac{\kappa}{w_i}} - \sqrt{\frac{w_i}{\kappa}}}{\sqrt{w_i} - \frac{1}{\sqrt{w_i}}} \quad (2.23)$$

where now the w_i are the arguments of $\widehat{\mathbf{a}}$ above and again we set $t_4 = q/\sqrt{\kappa}$ and $t_5 = 1/(q\sqrt{\kappa})$.

A generating function for refined Donaldson-Thomas invariants can be obtained from the full K-theoretic result [30]. To this end one picks a subtorus of \mathbb{T} defined by sending the toric weights t_1, t_2, t_3 to infinity or to zero in such a way that their product κ is kept constant. The result will in

principle depend on the choice of such a torus.

Before explaining this let us quickly review the various refinements of the MacMahon function. One can actually define a one parameter family of refined MacMahon functions

$$M_\delta(q_1, q_2) = \prod_{i,j}^{\infty} \left(1 - q_1^{i-1/2+\delta/2} q_2^{j-1/2-\delta/2}\right)^{-1}. \quad (2.24)$$

For example, $\delta = -1$ is the choice of [30] while the choice of $\delta = 0$ is the one made in [14]. One can introduce new variables $q_1 = q\sqrt{\kappa}$ and $q_2 = q/\sqrt{\kappa}$ and write

$$M_\delta(q, \sqrt{\kappa}) = \prod_{i,j}^{\infty} \left(1 - q^{i+j-1} \sqrt{\kappa}^{j-i-\delta}\right)^{-1}. \quad (2.25)$$

A concrete way to obtain the refined limit is for example to choose a subtorus

$$t_1 \longrightarrow x^{N-1} \quad t_3 \longrightarrow x \quad t_2 \longrightarrow 1/x^N \quad (2.26)$$

with $N \in \mathbb{Z}$ very large, and then send $x \longrightarrow 0$. With this choice we get

$$Z_{\mathbb{C}^3}^{ref}(q; \kappa) = 1 - q\sqrt{\kappa} + q^2(1 + 2\kappa) - q^3 \frac{1 + 2\kappa + 3\kappa^2}{\sqrt{\kappa}} + \dots \quad (2.27)$$

which agrees with the first terms of the refined MacMahon function

$$M(q\sqrt{\kappa}, q/\sqrt{\kappa})_{\delta=1} = M(q/\sqrt{\kappa}, q\sqrt{\kappa})_{\delta=-1} \quad (2.28)$$

Finally the cohomological limit is obtained by sending $\kappa \longrightarrow 1$.

The \mathbb{C}^3 partition function can be in principle generalized to any toric manifold, constructed by gluing together copies of \mathbb{C}^3 . To each copy one can associate a vertex with boundary conditions given by the asymptotics at infinity of a plane partition. The full vertex is not known in closed form except in particular cases [19]. The partition function can however be obtained order by order.

In this perspective the results of [30] correspond to the lift to M-theory of the BPS state counting problem formulated in the large radius chamber. In the rest of the article we will extend this construction to a different chamber.

3 Noncommutative algebraic geometry

In this Section we quickly review some basic elements of noncommutative algebraic geometry in the context of local Calabi-Yau threefolds. Certain toric singularities admit crepant resolutions which can be understood as moduli spaces of representations of a noncommutative algebra, the Jacobi algebra of a quiver with superpotential [36]. Such an algebra is an example of a noncommutative crepant resolution. We will briefly explain the relation between such resolutions and quivers and then discuss their role in defining noncommutative Donaldson-Thomas theory. The reader is referred to the reviews [9, 4] for a more in depth discussion.

3.1 Quivers and noncommutative crepant resolutions

Several aspects of BPS invariants on local threefolds can be understood algebraically via quiver representation theory [35, 7, 8, 6]. A quiver is a finite directed graph identified by the two finite sets \mathbf{Q}_0 and \mathbf{Q}_1 , the nodes and the arrows, and by the two maps $s, t : \mathbf{Q}_1 \rightarrow \mathbf{Q}_0$ that to an arrow $a \in \mathbf{Q}_1$ associate its starting vertex $s(a) \in \mathbf{Q}_0$ and its terminal vertex $t(a) \in \mathbf{Q}_0$.

The algebra of paths $\mathbb{C}\mathbf{Q}$ is the unital associative \mathbb{C} -algebra spanned by all paths in the quiver with product given by concatenation of paths if possible and zero otherwise. The Jacobian algebra $\mathbf{A} = \mathbb{C}\mathbf{Q}/\langle \mathbf{R} \rangle$ is the algebra of paths modulo the ideal generated by a set of relations \mathbf{R} , intended as \mathbb{C} -linear combinations of paths in $\mathbb{C}\mathbf{Q}$. We will be interested in the case where the relations \mathbf{R} are derived from a superpotential. A superpotential is a function $W : \mathbf{Q}_1 \rightarrow \mathbb{C}\mathbf{Q}$, given by a sum of cyclic monomials, and the ideal of relation is given by $\mathbf{R} = \langle \partial_a W \mid a \in \mathbf{Q}_1 \rangle$.

Representation of a quiver are defined by assigning a complex vector space V_i to each node $i \in \mathbf{Q}_0$ and a morphism $B_a \in \text{Hom}(V_{s(a)}, V_{t(a)})$ to each arrow $a \in \mathbf{Q}_1$. The dimension vector of the representation has components $d_i = \dim V_i$. If the quiver has a superpotential we require such morphisms to be compatible with the relations $\partial W = 0$. The category of quiver representations $\text{rep}(\mathbf{Q}, W)$ is equivalent to the category $\mathbf{A} - \text{mod}$ of left \mathbf{A} -modules. The representation space is defined by fixing a dimension vector $d \in \mathbb{N}^{\mathbf{Q}_0}$ as

$$\text{Rep}_d(\mathbf{Q}) = \bigoplus_{a: i \rightarrow j} \text{Hom}_{\mathbb{C}}(V_i, V_j) \quad (3.1)$$

and similarly $\text{Rep}_d(\mathbf{Q}, W)$ denotes the subscheme obtained by imposing the relations $\partial W = 0$. In most physical applications the objects of interest are isomorphism class of representations; that is the orbits with respect to the action of the group $G_d = \prod_{i \in \mathbf{Q}_0} GL(V_i, \mathbb{C})$. In this case we are let to the quotient stack $\mathbf{M}_d(\mathbf{Q}) = \text{Rep}_d(\mathbf{Q}, W)/G_d$.

In many physical situations one is interested in framed quivers and their representations. A framed quiver can be obtained for example by adding an extra vertex $\{\bullet\}$ together with an additional arrow a_\bullet such that $s(a_\bullet) = \bullet$ and $t(a_\bullet) = i_0$, where $i_0 \in \mathbf{Q}_0$ is a reference node of \mathbf{Q} . The new quiver $\widehat{\mathbf{Q}}$ has $\widehat{\mathbf{Q}}_0 = \mathbf{Q}_0 \cup \{\bullet\}$ and $\widehat{\mathbf{Q}}_1 = \mathbf{Q}_1 \cup \{\bullet \xrightarrow{a_\bullet} i_0\}$. Similarly one can define the Jacobian algebra of a framed quiver, framed representations and their moduli spaces.

Simple modules D_v are associated to each vertex $v \in \mathbf{Q}_0$ and have $V_v = \mathbb{C}$ and $V_w = 0$ for $w \neq v$. Physically such modules describe fractional branes. Let e_v be the trivial path at v of length zero. Then $P_v := e_v \mathbf{A}$ is the subspace of the path algebra which is generated by all paths that begin at vertex v . These are projective objects in the category $\text{rep}(\mathbf{Q}, W)$. Then a projective resolution of the simple module D_v is given by

$$\dots \longrightarrow \bigoplus_{w \in \mathbf{Q}_0} P_w^{\oplus d_{w,v}^k} \longrightarrow \dots \longrightarrow \bigoplus_{w \in \mathbf{Q}_0} P_w^{\oplus d_{w,v}^1} \longrightarrow P_v \longrightarrow D_v \longrightarrow 0 \quad (3.2)$$

where

$$d_{w,v}^p = \dim \text{Ext}_{\mathbf{A}}^p(D_v, D_w) . \quad (3.3)$$

In particular $d_{w,v}^0 = \delta_{w,v}$ since D_v are simple objects; $d_{w,v}^1$ gives the number of arrows in \mathbf{Q}_1 from

vertex w to vertex v ; $d_{w,v}^2$ gives the number of relations expressed as sum of paths from vertex w to vertex v ; $d_{w,v}^3$ is the number of relations among the relations, and so on.

The path algebra of a quiver can be thought of as a noncommutative analog of a polynomial algebra, while quiver representations are a noncommutative analog of coherent sheaves. In the case of local threefolds this can be made more precise. The relevant object which describes BPS states on the threefold is $D^b(\text{coh}(X))$, the homotopy category of complexes of sheaves where we invert all quasi-isomorphisms. Complexes of sheaves represent branes wrapping cycles in the Calabi-Yau geometry, and different complexes which give rise to the same physical state have to be identified.

If X arise as a crepant resolution of a singularity, then it also admits a noncommutative crepant resolution [36]. Crepant resolutions and noncommutative crepant resolutions are derived equivalent. The noncommutative resolution is a certain algebra A for which $D^b(\text{coh}(X)) = D^b(A - \text{mod})$. The equivalence is induced by the functor $\text{Hom}^\bullet(\mathcal{F}, -)$ where \mathcal{F} is a *tilting sheaf* \mathcal{F} (a generator with $\text{Hom}^\bullet(\mathcal{F}, \mathcal{F}[i]) = 0$ for $i > 0$).

For local Calabi-Yau threefolds X which are crepant resolutions of a singularity, the algebra A is the Jacobian algebra of a quiver. This is precisely the quiver which captures the low energy description of supersymmetric bound states of D-branes probing the singularity. In particular its center $Z(A)$ is the coordinate ring of the singularity and its moduli spaces of representations are resolutions of the singularity. Furthermore A is a Calabi-Yau 3-algebra.

In this paper we will limit ourselves to the simplest cases which arise as singular limits of toric Calabi-Yau. In this case the following construction holds: consider a singular threefold and let X be a (canonical) crepant resolution. Assume we can find a tilting bundle $\mathcal{T} \in \text{coh}(X)$, that is a vector bundle such that $\text{Ext}^i(\mathcal{T}, \mathcal{T}) = 0$ for $i > 0$ and which generates $D(\text{coh}(X))$. Then $\Lambda = \text{End}_X(\mathcal{T})$ is a noncommutative crepant resolution.

3.2 Noncommutative Donaldson-Thomas theory

On a noncommutative crepant resolution we can define BPS states. The low energy effective theory of a D2–D0 system is described by a supersymmetric quiver quantum mechanics with superpotential. Geometrically the quiver is derived through the endomorphisms of a tilting object. To incorporate configurations with non-vanishing D6-brane charge consider framed quivers. In practice we add an extra vertex $\{\bullet\}$ and an additional arrow a_\bullet from the new vertex to a reference vertex v_0 of \mathbb{Q} .

We denote the new quiver by $\hat{\mathbb{Q}}$ and its vertex and arrow sets by

$$\hat{\mathbb{Q}}_0 = \mathbb{Q}_0 \cup \{\bullet\} \quad \text{and} \quad \hat{\mathbb{Q}}_1 = \mathbb{Q}_1 \cup \{\bullet \xrightarrow{a_\bullet} v_0\}. \quad (3.4)$$

The representation theory of quivers as summarized above can be easily extended to framed quivers. More generically one can introduce a framing node with many arrows going out and incoming from the unframed quiver. If such arrows form new loops one can introduce a framed superpotentials \hat{W} . In this note we will not consider this case (for us it will be $W = \hat{W}$), but it will be relevant to the study of wall-crossing. We will also only consider framed representations where the framing node is one dimensional.

This new quiver has its own path algebra \hat{A} . Representations are now constructed by specifying an n_v -dimensional vector space V_v on each node $v \in \mathbb{Q}_0$, while the framing node \bullet always carries

a one-dimensional vector space \mathbb{C} . The moduli space $\mathcal{M}_{\mathbf{n}}(\hat{\mathbf{Q}}, v_0)$ of stable representations with dimension vector $d = (d_v)_{v \in \mathbf{Q}_0}$ is compact and well behaved [26] and enjoys a symmetric perfect obstruction theory [35, 26] which follows from the general results of [1].

One can define the noncommutative Donaldson-Thomas invariant as the weighted Euler characteristic of this moduli space as

$$\mathrm{DT}_{\mathbf{n}, v_0}(\mathbf{A}) := \chi(\mathcal{M}_{\mathbf{n}}(\hat{\mathbf{Q}}, v_0), \nu_{\mathbf{A}}) , \quad (3.5)$$

which physically represents the index of BPS bound states constituted by a single D6 brane and a gas of D2-D0 states.

Physically D6-D2-D0 BPS states are identified stable representations of $\hat{\mathbf{Q}}$. One interprets the simple representations as elementary BPS constituents which interact via the superpotential \hat{W} . When a representation of $(\hat{\mathbf{Q}}, W)$ with dimensions $d_i = \dim V_i$ exists and is stable, this is interpreted physically as the existence of a bound state of charge $\gamma = \sum_i d_i \gamma_i$. The stability condition is determined by a central charge function, a linear map $Z : K(\mathrm{rep}(\hat{\mathbf{Q}}, W)) \rightarrow \mathbb{C}$. We say that state of charge γ_r described by a representation $R \in \mathrm{rep}(\hat{\mathbf{Q}}, W)$ is stable if for any proper sub-representation $S \in \mathrm{rep}(\hat{\mathbf{Q}}, W)$ associated to a state of charge γ_s , we have that $\arg Z_{\gamma_s}(u) < \arg Z_{\gamma_r}(u)$.

For these invariants we define a cohomological partition function associated with the algebra \mathbf{A}

$$Z_{\mathbf{A}}^{\mathrm{coho}}(\mathbf{p}, v_0) = \sum_{n_v \in \mathbb{Z}} \mathrm{DT}_{\mathbf{n}, v_0}(\mathbf{A}) \prod_{v \in \mathbf{Q}_0} p_v^{n_v} . \quad (3.6)$$

where we have introduced a $|\mathbf{Q}_0|$ dimensional vector \mathbf{p} to keep track of the nodes upon which the representations are based. This enumerative problem can be studied via a combinatorial algorithm obtained by applying the virtual localization formula on the representation moduli space, with respect to a certain natural toric action. The fixed points of this action are ideals in the path algebra \mathbf{A} , which admit a graphical representation in terms of certain combinatorial arrangements known as *pyramid partitions*.

It follows from a result by Behrend-Fantechi [1] that for a symmetric perfect obstruction theory of the kind constructed in [35, 26] the contribution of an isolated fixed point to the Euler characteristic (3.5) is just a sign determined by the parity of the dimension of the tangent space at the fixed point.

Therefore if all fixed points π are isolated we obtain

$$\mathrm{DT}_{\mathbf{n}, v_0}(\mathbf{A}) = \sum_{\pi \in \mathcal{M}_{\mathbf{n}}(\hat{\mathbf{Q}}, v_0)^{\mathbb{T}}} (-1)^{\dim T_{\pi} \mathcal{M}_{\mathbf{n}}(\hat{\mathbf{Q}}, v_0)} . \quad (3.7)$$

While this formula has a concrete combinatorial description, at first sight its physical interpretation is less clear because one loses track of the relation between the D-brane charges defined at large radius and the combinatorial objects. The physical states at the singularity have charges which are non-trivial functions of the D0 and D2 brane charges at large radius. The relevant dynamical variables at the singularities should be interpreted as fractional branes, stuck at the singularity. This change of variables is known in some models [7].

4 The noncommutative conifold

In this Section we will study the K-theoretic DT theory for the conifold quiver. We will briefly review the relevant quiver and its representation theory and then compute the M2-brane index by a direct localization computation. We then provide a conjecture for the full partition function and discuss some of its limits.

4.1 Geometry and BPS states

We will now consider the conifold geometry. At large radius it is given by the total space of the bundle $X = \mathcal{O}(-1) \oplus \mathcal{O}(-1) \rightarrow \mathbb{P}^1$. In this phase the topological string captures the BPS invariants corresponding to bound states of a single D6 brane with a gas of D0/D2 branes, with a strong B-field [17]. In this limit the problem of computing BPS invariants correspond to the study of Donaldson-Thomas invariants associated with the Hilbert scheme of points and curves $\text{Hilb}(X; n, \beta)$, with n the number of D0 branes and β the class of the curve wrapped by the D2 branes.

We will be interested in the singular limit where the size of the \mathbb{P}^1 is sent to zero. In the noncommutative phase the conifold geometry is replaced by the representation theory of the Klebanov-Witten quiver

$$\circ \begin{array}{c} \xrightarrow{a_1, a_2} \\ \xleftarrow{b_1, b_2} \end{array} \bullet \quad (4.1)$$

with superpotential $W = a_1 b_1 a_2 b_2 - a_1 b_2 a_2 b_1$. We will denote by $\mathbb{C}\mathcal{Q}$ the algebra of paths, defined as the algebra generated by the arrow set, with product given by the concatenation of arrows whenever possible. The Jacobian path algebra is $\mathbf{A} = \mathbb{C}\mathcal{Q}/\langle \partial W \rangle$. The center of the path algebra $Z(\mathbf{A})$ is spanned by the variables

$$\begin{aligned} x_1 &= a_1 b_1 + b_1 a_1 \\ x_2 &= a_2 b_2 + b_2 a_2 \\ x_3 &= a_1 b_2 + b_2 a_1 \\ x_4 &= a_2 b_1 + b_1 a_2 \end{aligned} \quad (4.2)$$

so that $\text{Spec } R = \text{Spec } \mathbb{C}[x_1, x_2, x_3, x_4]/(x_1 x_2 - x_3 x_4)$ is the ring of functions on the conifold singularity. Therefore the path algebra of the Klebanov-Witten quiver sees the conifold singularity as its center. The path algebra itself is a noncommutative crepant resolution. The category of left \mathbf{A} modules $\mathbf{A} - \text{mod}$ is equivalent to the category of quiver representations $\text{rep}(\mathcal{Q}, W)$. Large radius and noncommutative phases are related by $D^b(\mathbf{A} - \text{mod}) \simeq D^b(\text{coh } X)$.

We would like to understand BPS states in the noncommutative phase. The study of BPS states for the conifold quiver was done in [35]. To model D-brane bound states one adds a framing node to the quiver, corresponding to a single D6 brane

$$\begin{array}{c} \star \\ \searrow \\ \circ \begin{array}{c} \xrightarrow{a_1, a_2} \\ \xleftarrow{b_1, b_2} \end{array} \bullet \end{array} \quad (4.3)$$

BPS invariants associated with bound states of D-branes now correspond to noncommutative Donaldson-Thomas invariants associated with this quiver [35]. We want to generalize this computation to the K-theory setting.

4.2 Toric action and fixed points

Consider the representation space

$$\text{Rep}(\mathbf{Q}, v_0) = \bigoplus_{(v \rightarrow w) \in \mathbf{Q}_1} \text{Hom}_{\mathbb{C}}(V_v, V_w) \oplus \text{Hom}(V_0, \mathbb{C}) \quad (4.4)$$

The last factor is associated to the framing and determines a vector $v_0 \in V_0$. A suitable stability condition which selects the noncommutative crepant resolution chamber is given by only considering *cyclic modules*, that is modules which are generated by v_0 [35, 9]. Now we consider the open subvariety $\text{Rep}^c(\mathbf{Q}, v_0)$ defined by the condition that the module is cyclic. Similarly we can consider the representation space $\text{Rep}(\mathbf{Q}, v_0; \mathbf{W})$ defined by imposing the F-term relations on $\text{Rep}(\mathbf{Q}, v_0)$, and the corresponding subvariety of cyclic modules $\text{Rep}^c(\mathbf{Q}, v_0; \mathbf{W})$. Finally the relevant moduli space is

$$\mathcal{M}_{d_\circ, d_\bullet} = \left[\text{Rep}^c(\mathbf{Q}, v_0; \mathbf{W}) \Big/ GL(d_\circ, \mathbb{C}) \times GL(d_\bullet, \mathbb{C}) \right]. \quad (4.5)$$

We will denote a (representative of an isomorphism class of a) cyclic module as (v_\circ, M) .

This moduli space carries a natural toric action which can be used to classify fixed points. The classification is carried out in Proposition 2.5.1 of [35]; since the proof is constructive, we summarize it here in some detail. The only difference is that we do not impose that the determinant of the toric action is trivial. We have the following tori

- \mathbb{T}_F . This is the natural torus which rescales each arrow by a phase, which we will denote by $\{t_{a_1}, t_{a_2}, t_{b_1}, t_{b_2}\}$. Note that this torus also preserves the F-term relations and therefore acts naturally on the moduli space.
- \mathbb{T}_G . This is the ‘‘gauge’’ torus, induced by the $GL(d_\circ, \mathbb{C}) \times GL(d_\bullet, \mathbb{C})$ action which changes the bases of V_\circ and V_\bullet . This group contains a $(\mathbb{C}^*)^2$ factor, whose elements $(\mu_\circ, \mu_\bullet) \in (\mathbb{C}^*)^2$ act on representation maps by conjugation, say $\mu_\circ X_a \mu_\bullet^{-1}$ or $\mu_\bullet X_b \mu_\circ^{-1}$. However the induced gauge torus is just $\mathbb{T}_G = \mathbb{C}^* = (\mathbb{C}^*)^2 / \mathbb{C}^*$, since diagonal elements act trivially. We take this torus acting by weights $(-1, -1, 1, 1)$ (since it must act on the b ’s in the opposite way as on the a ’s). This torus however does not influence the fixed point classification.
- $\mathbb{T}_{F, \mathbf{W}}$. This is the sub-torus of \mathbb{T}_F with the weight condition $t_{a_1} t_{a_2} t_{b_1} t_{b_2} = \kappa$. In [35] the condition is $\kappa = 1$
- Finally the torus $\mathbb{T}_{\mathbf{W}} = \mathbb{T}_{F, \mathbf{W}} / \mathbb{T}_G$.

Now we proceed to study the fixed points of $\mathcal{M}_{d_\circ, d_\bullet}$. The result of [35] is that such fixed points correspond to ideals in \mathbf{A} . To begin with, a cyclic module (v_\circ, M) is canonically associated to an ideal (its annihilator) I , as $M = \mathbf{A}/I$. In particular since v_\circ is based at the vertex \circ , this ideal has the form $I = I_\circ \oplus P_\bullet$, where $P_\bullet = \mathbf{A} e_\bullet$ are the projectives based on the vertex \bullet .

Note that $\mathbb{T}_W = \mathbb{T}_{F,W}/\mathbb{T}_G$ -fixed points of $\mathcal{M}_{d_\circ, d_\bullet}$ correspond to $\mathbb{T}_{F,W}$ -fixed ideals I in A . The reason is that the generators of I_0 are sum of path algebra monomials with the same starting point and endpoint. Therefore on each generator the gauge torus acts by multiplication by a constant. Since I_0 is an ideal, it does not change under this operation. In other words the gauge torus \mathbb{T}_G acts trivially on I and only the action of the covering torus $\mathbb{T}_{F,W}$ matters. Finally we have to look for $\mathbb{T}_{F,W}$ -fixed ideals I , that is ideals generated by eigenvectors of $\mathbb{T}_{F,W}$.

It is shown in [35] that such ideals are in one to one correspondence with pyramid partitions. Pyramid partitions are defined as follows. First we define a pyramid arrangement, which is a three dimensional infinite pyramid; the tip of this pyramid is shown in Figure 1. The first layer (starting from the top of the pyramid) of this configuration corresponds to the vector v_\circ , the second layer corresponds to the two ways we can reach the node \bullet starting from \circ , that is a_1v_\circ and a_2v_\circ , and so on. The shape of the pyramid partition is determined by the superpotential relations $\partial W = 0$.




Figure 1: Pyramid arrangement. We show the tip of an infinite pyramid; white and black stones are associated with the two nodes of the KW quiver. One moves from one stone to another via the arrows of the quiver; the toric weight of the arrows is shown explicitly.

Given a pyramid arrangement, one defines a pyramid partition π by the following condition: π is a configuration of stones such that for each stone in π , the ones immediately above it (of a different color) are in π as well. Note that this is basically the same definition of a Young diagram, which guarantees that the complement of a pyramid partition is an ideal. Examples of pyramid partitions are shown in Figure 2.




Figure 2: Conifold fixed points. Each fixed point corresponds to a pyramid partition, a configuration of stones which can be consistently removed from the pyramid arrangement.

Now we argue that pyramid partitions corresponds to fixed points.

First consider a module M . By definition it is spanned by vectors of the form

$$\{v_\circ, a_1 v_\circ, a_2 v_\circ, b_1 a_1 v_\circ, b_2 a_1 v_\circ, \dots\}. \quad (4.6)$$

Note that by cyclicity, if a term like $b_1 a_1 v_\circ$ is present, then also $a_1 v_\circ$ and v_\circ must be present. This is just the condition that M defines an ideal I via $M = A/I$. From this set of vectors we remove those which are linearly dependent when imposing the F-term relations $\partial W = 0$. Then to each of the remaining vectors, which now form a basis, we associate a stone in a pyramid partition. The partition has the correct form because of the equations of motion.

Conversely, consider a partition π . To each stone we assign a basis vector $v_{i,a}$, with $a = \circ, \bullet$. Then we collect all the $v_{i,a}$ with the same color for a and use them as a spanning set to construct a vector space V_a , whose dimension is the number of stones with that color. The arrows of the quiver induce maps between these vectors, which obey the relations $\partial W = 0$ by construction. Finally the corresponding module is $M = \mathbb{C} \oplus V_\circ \oplus V_\bullet$, where the first factor corresponds to the framing node.

This complete the classification of fixed points in terms of pyramid partitions [35]. Note that to compute the K-theoretic partition function it is not enough to enumerate the combinatorial configurations, since we need explicitly the isotypical decomposition of the vector spaces V_\circ and V_\bullet in toric weight spaces. This information can be obtained, for example, from the spanning set (4.6). This is why we have summarized the discussion of [35] in such detail.

4.3 Localization

To each fixed point π we associate the decomposition in weight spaces

$$V_\circ = \sum_{(i,j,k,l) \in \pi_\circ} t_{a_1}^{-i} t_{a_2}^{-j} t_{b_1}^{-k} t_{b_2}^{-l} \quad (4.7)$$

$$V_\bullet = \sum_{(i,j,k,l) \in \pi_\bullet} t_{a_1}^{-i} t_{a_2}^{-j} t_{b_1}^{-k} t_{b_2}^{-l} \quad (4.8)$$

where the two sets (π_\circ, π_\bullet) run over stones of the respective color. The weights are those of the generators of the cyclic module, as explained above.

To use the localization formula we need the deformation/obstruction theory around each fixed point. This is given by the virtual tangent space. The virtual tangent space can be constructed in terms of the tangent bundle. The tangent bundle has the form “arrows” - “gauge transformations” :

$$T = (t_{a_1} + t_{a_2}) \otimes \bar{V}_\circ \otimes V_\bullet + (t_{b_1} + t_{b_2}) \otimes \bar{V}_\bullet \otimes V_\circ + V_\circ - \bar{V}_\circ \otimes V_\circ - \bar{V}_\bullet \otimes V_\bullet \quad (4.9)$$

where the “arrows” carry their toric weight, and we do not consider gauge transformations on the

framing node. Then we have

$$\begin{aligned}
T^{vir} &= T - \kappa \otimes T^* & (4.10) \\
&= (t_{a_1} + t_{a_2}) \otimes \bar{V}_\circ \otimes V_\bullet + (t_{b_1} + t_{b_2}) \otimes \bar{V}_\bullet \otimes V_\circ + V_\circ - \bar{V}_\circ \otimes V_\circ - \bar{V}_\bullet \otimes V_\bullet \\
&\quad - t_{a_1} t_{a_2} t_{b_1} t_{b_2} \left((t_{a_1}^{-1} + t_{a_2}^{-1}) \otimes V_\circ \otimes \bar{V}_\bullet + (t_{b_1}^{-1} + t_{b_2}^{-1}) \otimes V_\bullet \otimes \bar{V}_\circ + V_\circ - \bar{V}_\circ \otimes V_\circ - \bar{V}_\bullet \otimes V_\bullet \right)
\end{aligned}$$

with $\kappa = t_{a_1} t_{a_2} t_{b_1} t_{b_2}$.

We define our partition function as

$$Z_{NCC}(q_0, q_i, \{t_i\}) = \chi(\mathcal{M}, \hat{\mathcal{O}}^{vir}) = \sum_{(d_\circ, d_\bullet)} q_0^{d_\circ} q_1^{d_\bullet} \chi \left(\mathcal{M}_{d_\circ, d_\bullet}, (\text{sgn}) \mathcal{O}_{\mathcal{M}_{d_\circ, d_\bullet}}^{vir} \otimes \mathcal{K}_{vir}^{1/2} \right) \quad (4.11)$$

Here (sgn) refers to the fact that in defining the prefactor of the symmetrized virtual structure sheaf we have the freedom of choosing a sign. The possibilities we will use in the following are $\text{sgn} = 1, (-1)^{d_\circ}, (-1)^{d_\bullet}$; but this is just a convention. For the moment we will drop this sign and reinstate it only at the end. We have defined $\mathcal{K}_{vir}^{1/2} = \det^{-1/2} T^{vir}$ and the full moduli space is given by the disjoint union $\mathcal{M} = \cup_{d_\circ, d_\bullet} \mathcal{M}_{d_\circ, d_\bullet}$.

After expanding and cancelling terms, the virtual tangent space assumes schematically the form $T^{vir} = \text{Def} - \text{Obs} = \sum_i w_i - \sum_i \kappa/w_i$, where w_i denotes a generic toric weight. Then the localized virtual structure sheaf has the form

$$\begin{aligned}
\mathcal{O}_{\mathcal{M}_{d_\circ, d_\bullet}}^{vir} \otimes \mathcal{K}_{vir}^{1/2} &= \sum_{|\pi_\circ|=d_\circ, |\pi_\bullet|=d_\bullet} \hat{a}(T_{\pi_\circ, \pi_\bullet}^{vir}) \mathcal{O}_{I_{\pi_\circ, \pi_\bullet}} \\
&= \sum_{|\pi_\circ|=d_\circ, |\pi_\bullet|=d_\bullet} \prod_w \frac{\left(\frac{\kappa}{w}\right)^{1/2} - \left(\frac{w}{\kappa}\right)^{1/2}}{(w)^{1/2} - (w)^{-1/2}} \mathcal{O}_{I_{\pi_\circ, \pi_\bullet}} \quad (4.12)
\end{aligned}$$

in K-theory. The sum runs over all the fixed points with d_\circ white stones and d_\bullet black stones. The role of the twist by $\mathcal{K}_{vir}^{1/2}$ is to transform the usual K-theory weights into $\hat{a}(T^{vir})$.

4.4 Some explicit fixed points

Here we collect some explicit results of the localization computation. Higher order computations become quickly rather technical. The interested reader can find some of them in the supporting MATHEMATICA files.

1. Fixed point with $d = (d_\circ, d_\bullet) = (1, 0)$. This is the first fixed point in Figure 2. It has $V_\circ = 1$ and $V_\bullet = \emptyset$. Then $T^{vir} = 0$, which means that the deformations exactly cancel the obstructions. Since the contribution of a fixed point is the ration of obstructions and deformations, we simply have $\hat{a}(T^{vir}) = 1$.
2. Fixed points with $d = (1, 1)$. There are two fixed points, corresponding to the representations with $V_\circ = 1, V_\bullet = 1/t_{a_1}$ and $V_\circ = 1, V_\bullet = 1/t_{a_2}$, as \mathbb{T} -modules. The corresponding pyramid partitions are shown in Figure 2. In the first case

$$T_1^{vir} = \frac{t_{a_2}}{t_{a_1}} + t_{a_1} t_{b_1} - t_{a_2} t_{b_1} + t_{a_1} t_{b_2} - t_{a_2} t_{b_2} - t_{a_1}^2 t_{b_1} t_{b_2} \quad (4.13)$$

which contributes

$$\hat{a}(T_1^{vir}) = \frac{\left(\frac{\kappa}{t_{a_2}/t_{a_1}}\right)^{1/2} - \left(\frac{t_{a_2}/t_{a_1}}{\kappa}\right)^{1/2}}{(t_{a_2}/t_{a_1})^{1/2} - (t_{a_2}/t_{a_1})^{-1/2}} \frac{\left(\frac{\kappa}{t_{a_1}t_{b_1}}\right)^{1/2} - \left(\frac{t_{a_1}t_{b_1}}{\kappa}\right)^{1/2}}{(t_{a_1}t_{b_1})^{1/2} - (t_{a_1}t_{b_1})^{-1/2}} \frac{\left(\frac{\kappa}{t_{a_1}t_{b_2}}\right)^{1/2} - \left(\frac{t_{a_1}t_{b_2}}{\kappa}\right)^{1/2}}{(t_{a_1}t_{b_2})^{1/2} - (t_{a_1}t_{b_2})^{-1/2}}. \quad (4.14)$$

Similarly for the second fixed point

$$T_2^{vir} = \frac{t_{a_1}}{t_{a_2}} - t_{a_1}t_{b_1} + t_{a_2}t_{b_1} - t_{a_1}t_{b_2} + t_{a_2}t_{b_2} - t_{a_2}^2t_{b_1}t_{b_2} \quad (4.15)$$

we find

$$\hat{a}(T_2^{vir}) = \frac{\left(\frac{\kappa}{t_{a_1}/t_{a_2}}\right)^{1/2} - \left(\frac{t_{a_1}/t_{a_2}}{\kappa}\right)^{1/2}}{(t_{a_1}/t_{a_2})^{1/2} - (t_{a_1}/t_{a_2})^{-1/2}} \frac{\left(\frac{\kappa}{t_{a_2}t_{b_1}}\right)^{1/2} - \left(\frac{t_{a_2}t_{b_1}}{\kappa}\right)^{1/2}}{(t_{a_2}t_{b_1})^{1/2} - (t_{a_2}t_{b_1})^{-1/2}} \frac{\left(\frac{\kappa}{t_{a_2}t_{b_2}}\right)^{1/2} - \left(\frac{t_{a_2}t_{b_2}}{\kappa}\right)^{1/2}}{(t_{a_2}t_{b_2})^{1/2} - (t_{a_2}t_{b_2})^{-1/2}}. \quad (4.16)$$

Summing the two contributions gives

$$\left(-\frac{(\kappa t_{a_1} - t_{a_2})(t_{a_1}t_{b_1} - \kappa)(t_{a_1}t_{b_2} - \kappa)}{\kappa^{3/2}(t_{a_1} - t_{a_2})(t_{a_1}t_{b_1} - 1)(t_{a_1}t_{b_2} - 1)} - \frac{(t_{a_1} - \kappa t_{a_2})(t_{a_2}t_{b_1} - \kappa)(t_{a_2}t_{b_2} - \kappa)}{\kappa^{3/2}(t_{a_1} - t_{a_2})(t_{a_2}t_{b_1} - 1)(t_{a_2}t_{b_2} - 1)} \right) \quad (4.17)$$

3. Fixed point with $d = (1, 2)$. There is only one, shown in Figure 2, with $V_\circ = 1$ and $V_\bullet = 1/t_{a_1} + 1/t_{a_2}$. We see that $T^{vir} = 0$, so its contribution is just $\hat{a}(T^{vir}) = 1$
4. Fixed points with $d = (2, 1)$. There are four fixed point, which we write as tuples $\{V_\circ, V_\bullet\}$

$$\left\{ \frac{1}{t_{a_1}t_{b_1}} + 1, \frac{1}{t_{a_1}} \right\} \quad \left\{ \frac{1}{t_{a_1}t_{b_2}} + 1, \frac{1}{t_{a_1}} \right\} \quad \left\{ \frac{1}{t_{a_2}t_{b_1}} + 1, \frac{1}{t_{a_2}} \right\} \quad \left\{ \frac{1}{t_{a_2}t_{b_2}} + 1, \frac{1}{t_{a_2}} \right\} \quad (4.18)$$

The corresponding virtual tangent spaces are

$$T_1^{vir} = \frac{t_{a_2}}{t_{a_1}} + t_{a_1}t_{b_1} - t_{a_1}t_{a_2}t_{b_1}^2 - t_{a_2}t_{b_2} + \frac{t_{b_2}}{t_{b_1}} - t_{a_1}^2t_{b_1}t_{b_2}, \quad (4.19)$$

$$T_2^{vir} = \frac{t_{a_2}}{t_{a_1}} - t_{a_2}t_{b_1} + \frac{t_{b_1}}{t_{b_2}} + t_{a_1}t_{b_2} - t_{a_1}^2t_{b_1}t_{b_2} - t_{a_1}t_{a_2}t_{b_2}^2, \quad (4.20)$$

$$T_3^{vir} = \frac{t_{a_1}}{t_{a_2}} + t_{a_2}t_{b_1} - t_{a_1}t_{a_2}t_{b_1}^2 - t_{a_1}t_{b_2} + \frac{t_{b_2}}{t_{b_1}} - t_{a_2}^2t_{b_1}t_{b_2}, \quad (4.21)$$

$$T_4^{vir} = \frac{t_{a_1}}{t_{a_2}} - t_{a_1}t_{b_1} + \frac{t_{b_1}}{t_{b_2}} + t_{a_2}t_{b_2} - t_{a_2}^2t_{b_1}t_{b_2} - t_{a_1}t_{a_2}t_{b_2}^2. \quad (4.22)$$

The full contribution of the four fixed points is

$$\begin{aligned} & -\frac{(\kappa t_{a_1} - t_{a_2})(-\kappa + t_{a_1}t_{b_1})(\kappa t_{b_1} - t_{b_2})}{\kappa^{3/2}(t_{a_1} - t_{a_2})(-1 + t_{a_1}t_{b_1})(t_{b_1} - t_{b_2})} - \frac{(t_{a_1} - \kappa t_{a_2})(-\kappa + t_{a_2}t_{b_1})(\kappa t_{b_1} - t_{b_2})}{\kappa^{3/2}(t_{a_1} - t_{a_2})(-1 + t_{a_2}t_{b_1})(t_{b_1} - t_{b_2})} \\ & -\frac{(\kappa t_{a_1} - t_{a_2})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_1}t_{b_2})}{\kappa^{3/2}(t_{a_1} - t_{a_2})(-t_{b_1} + t_{b_2})(-1 + t_{a_1}t_{b_2})} - \frac{(t_{a_1} - \kappa t_{a_2})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_2}t_{b_2})}{\kappa^{3/2}(t_{a_1} - t_{a_2})(-t_{b_1} + t_{b_2})(-1 + t_{a_2}t_{b_2})}. \end{aligned} \quad (4.23)$$

5. Fixed points with $d = (2, 2)$. There are eight fixed points (one of which is shown in Figure 2), we list them in the form $\{V_\circ, V_\bullet\}$

$$\begin{aligned}
& \left\{ \frac{1}{t_{a_1} t_{b_1}} + 1, \frac{1}{t_{a_1}^2 t_{b_1}} + \frac{1}{t_{a_1}} \right\} \left\{ \frac{1}{t_{a_1} t_{b_2}} + 1, \frac{1}{t_{a_1}^2 t_{b_2}} + \frac{1}{t_{a_1}} \right\} \left\{ \frac{1}{t_{a_2} t_{b_1}} + 1, \frac{1}{t_{a_2}^2 t_{b_1}} + \frac{1}{t_{a_2}} \right\} \\
& \left\{ \frac{1}{t_{a_2} t_{b_2}} + 1, \frac{1}{t_{a_2}^2 t_{b_2}} + \frac{1}{t_{a_2}} \right\} \left\{ \frac{1}{t_{a_1} t_{b_1}} + 1, \frac{1}{t_{a_1}} + \frac{1}{t_{a_2}} \right\} \left\{ \frac{1}{t_{a_1} t_{b_2}} + 1, \frac{1}{t_{a_1}} + \frac{1}{t_{a_2}} \right\} \\
& \left\{ \frac{1}{t_{a_2} t_{b_1}} + 1, \frac{1}{t_{a_1}} + \frac{1}{t_{a_2}} \right\} \left\{ \frac{1}{t_{a_2} t_{b_2}} + 1, \frac{1}{t_{a_1}} + \frac{1}{t_{a_2}} \right\}
\end{aligned} \tag{4.24}$$

The overall contribution is

$$\begin{aligned}
& \frac{(\kappa t_{a_1} - t_{a_2})(-\kappa + t_{a_1} t_{b_1})(-t_{a_2} + \kappa t_{a_1}^2 t_{b_1})(-\kappa + t_{a_1}^2 t_{b_1}^2)(\kappa t_{b_1} - t_{b_2})(-\kappa + t_{a_1} t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_1} t_{b_1})^2(1 + t_{a_1} t_{b_1})(-t_{a_2} + t_{a_1}^2 t_{b_1})(t_{b_1} - t_{b_2})(-1 + t_{a_1} t_{b_2})} \\
& + \frac{(t_{a_1} - \kappa t_{a_2})(-\kappa + t_{a_1} t_{b_1})(-\kappa + t_{a_2} t_{b_1})(\kappa t_{a_1} - t_{a_2}^2 t_{b_1})(\kappa t_{b_1} - t_{b_2})(-\kappa t_{a_2} t_{b_1} + t_{a_1} t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_1} t_{b_1})(-1 + t_{a_2} t_{b_1})(t_{a_1} - t_{a_2}^2 t_{b_1})(t_{b_1} - t_{b_2})(-t_{a_2} t_{b_1} + t_{a_1} t_{b_2})} \\
& + \frac{(\kappa t_{a_1} - t_{a_2})(-\kappa + t_{a_1} t_{b_1})(-\kappa t_{a_2} + t_{a_1}^2 t_{b_1})(-\kappa + t_{a_2} t_{b_1})(\kappa t_{b_1} - t_{b_2})(\kappa t_{a_1} t_{b_1} - t_{a_2} t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_1} t_{b_1})(-t_{a_2} + t_{a_1}^2 t_{b_1})(-1 + t_{a_2} t_{b_1})(t_{b_1} - t_{b_2})(t_{a_1} t_{b_1} - t_{a_2} t_{b_2})} \\
& + \frac{(t_{a_1} - \kappa t_{a_2})(-\kappa + t_{a_2} t_{b_1})(t_{a_1} - \kappa t_{a_2}^2 t_{b_1})(-\kappa + t_{a_2}^2 t_{b_1}^2)(\kappa t_{b_1} - t_{b_2})(-\kappa + t_{a_2} t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_2} t_{b_1})^2(1 + t_{a_2} t_{b_1})(t_{a_1} - t_{a_2}^2 t_{b_1})(t_{b_1} - t_{b_2})(-1 + t_{a_2} t_{b_2})} \\
& + \frac{(\kappa t_{a_1} - t_{a_2})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_1} t_{b_2})(-t_{a_2} t_{b_1} + \kappa t_{a_1} t_{b_2})(-\kappa t_{a_2} + t_{a_1}^2 t_{b_2})(-\kappa + t_{a_2} t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-t_{b_1} + t_{b_2})(-1 + t_{a_1} t_{b_2})(-t_{a_2} t_{b_1} + t_{a_1} t_{b_2})(-t_{a_2} + t_{a_1}^2 t_{b_2})(-1 + t_{a_2} t_{b_2})} \\
& + \frac{(t_{a_1} - \kappa t_{a_2})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_1} t_{b_2})(-\kappa + t_{a_2} t_{b_2})(t_{a_1} t_{b_1} - \kappa t_{a_2} t_{b_2})(\kappa t_{a_1} - t_{a_2}^2 t_{b_2})}{\kappa^3(t_{a_1} - t_{a_2})(-t_{b_1} + t_{b_2})(-1 + t_{a_1} t_{b_2})(t_{a_1} t_{b_1} - t_{a_2} t_{b_2})(-1 + t_{a_2} t_{b_2})(t_{a_1} - t_{a_2}^2 t_{b_2})} \\
& - \frac{(\kappa t_{a_1} - t_{a_2})(-\kappa + t_{a_1} t_{b_1})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_1} t_{b_2})(-t_{a_2} + \kappa t_{a_1}^2 t_{b_2})(-\kappa + t_{a_1}^2 t_{b_2}^2)}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_1} t_{b_1})(t_{b_1} - t_{b_2})(-1 + t_{a_1} t_{b_2})^2(1 + t_{a_1} t_{b_2})(-t_{a_2} + t_{a_1}^2 t_{b_2})} \\
& - \frac{(t_{a_1} - \kappa t_{a_2})(-\kappa + t_{a_2} t_{b_1})(-t_{b_1} + \kappa t_{b_2})(-\kappa + t_{a_2} t_{b_2})(t_{a_1} - \kappa t_{a_2}^2 t_{b_2})(-\kappa + t_{a_2}^2 t_{b_2}^2)}{\kappa^3(t_{a_1} - t_{a_2})(-1 + t_{a_2} t_{b_1})(t_{b_1} - t_{b_2})(-1 + t_{a_2} t_{b_2})^2(1 + t_{a_2} t_{b_2})(t_{a_1} - t_{a_2}^2 t_{b_2})}
\end{aligned} \tag{4.25}$$

6. Fixed points with $d = (3, 1)$. They are

$$\begin{aligned}
V_\circ &= 1 + 1/(t_{a_1} t_{b_1}) + 1/(t_{a_1} t_{b_2}) & V_\bullet &= 1/t_{a_1} \\
V_\circ &= 1 + 1/(t_{a_2} t_{b_1}) + 1/(t_{a_2} t_{b_2}) & V_\bullet &= 1/t_{a_2}
\end{aligned} \tag{4.26}$$

7. Fixed points with $d = (2, 3)$

$$\begin{aligned}
V_\circ &= 1 + 1/(t_{a_1} t_{b_2}) & V_\bullet &= 1/t_{a_1} + 1/(t_{a_1}^2 t_{b_2}) + 1/t_{a_2} \\
V_\circ &= 1 + 1/(t_{a_1} t_{b_1}) & V_\bullet &= 1/t_{a_1} + 1/(t_{a_1}^2 t_{b_1}) + 1/t_{a_2} \\
V_\circ &= 1 + 1/(t_{a_2} t_{b_2}) & V_\bullet &= 1/t_{a_2} + 1/(t_{a_2}^2 t_{b_2}) + 1/t_{a_1} \\
V_\circ &= 1 + 1/(t_{a_2} t_{b_1}) & V_\bullet &= 1/t_{a_2} + 1/(t_{a_2}^2 t_{b_1}) + 1/t_{a_1}
\end{aligned} \tag{4.27}$$

and so on. The rest of the fixed points up to $(d_\circ, d_\bullet) = (4, 4)$ can be found in the supporting MATHEMATICA file.

4.5 A conjecture for the full partition function

We can use the partial knowledge given by the few fixed points listed above to propose a conjecture for the full K-theoretic partition function. As explained above we will keep open the freedom to chose the sign of the counting variables. We define the following function associated to the noncommutative conifold

$$\begin{aligned} \mathcal{F}_{NCC}[t_{a_1}, t_{a_2}, t_{b_1}, t_{b_2}, t_4, t_5, q_1] &= \frac{\frac{1}{q_1} + q_1}{\left(-\frac{1}{\sqrt{t_4}} + \sqrt{t_4}\right) \left(-\frac{1}{\sqrt{t_5}} + \sqrt{t_5}\right)} \\ &+ \frac{\left(\sqrt{\frac{\kappa t_{a_1}}{t_{a_2}}} - \sqrt{\frac{t_{a_2}}{\kappa t_{a_1}}}\right) \left(\sqrt{\frac{\kappa}{t_{a_1} t_{b_1}}} - \sqrt{\frac{t_{a_1} t_{b_1}}{\kappa}}\right) \left(\sqrt{\frac{\kappa}{t_{a_1} t_{b_2}}} - \sqrt{\frac{t_{a_1} t_{b_2}}{\kappa}}\right)}{\left(-\frac{1}{\sqrt{\frac{t_{a_2}}{t_{a_1}}}} + \sqrt{\frac{t_{a_2}}{t_{a_1}}}\right) \left(-\frac{1}{\sqrt{t_{a_1} t_{b_1}}} + \sqrt{t_{a_1} t_{b_1}}\right) \left(-\frac{1}{\sqrt{t_{a_1} t_{b_2}}} + \sqrt{t_{a_1} t_{b_2}}\right)} \frac{1}{\left(-\frac{1}{\sqrt{t_4}} + \sqrt{t_4}\right) \left(-\frac{1}{\sqrt{t_5}} + \sqrt{t_5}\right)} \\ &+ \frac{\left(-\sqrt{\frac{t_{a_1}}{\kappa t_{a_2}}} + \sqrt{\frac{\kappa t_{a_2}}{t_{a_1}}}\right) \left(\sqrt{\frac{\kappa}{t_{a_2} t_{b_1}}} - \sqrt{\frac{t_{a_2} t_{b_1}}{\kappa}}\right) \left(\sqrt{\frac{\kappa}{t_{a_2} t_{b_2}}} - \sqrt{\frac{t_{a_2} t_{b_2}}{\kappa}}\right)}{\left(-\frac{1}{\sqrt{\frac{t_{a_1}}{t_{a_2}}}} + \sqrt{\frac{t_{a_1}}{t_{a_2}}}\right) \left(-\frac{1}{\sqrt{t_{a_2} t_{b_1}}} + \sqrt{t_{a_2} t_{b_1}}\right) \left(-\frac{1}{\sqrt{t_{a_2} t_{b_2}}} + \sqrt{t_{a_2} t_{b_2}}\right)} \frac{1}{\left(-\frac{1}{\sqrt{t_4}} + \sqrt{t_4}\right) \left(-\frac{1}{\sqrt{t_5}} + \sqrt{t_5}\right)} \end{aligned} \quad (4.28)$$

In this formula it is understood that $\kappa = t_{a_1} t_{a_2} t_{b_1} t_{b_2}$. The variable q_1 plays the role of a Kähler modulus. The factor $q_1 + \frac{1}{q_1}$ captures the ‘‘Kähler dependent’’ part of the partition function (meaning that this is the analog of the Kähler modulus in the noncommutative phase).

The remaining terms are precisely copies of the \mathbb{C}^3 partition function $\mathcal{F}_{\mathbb{C}^3}[t_1, t_2, t_3, t_4, t_5]$, with the substitutions

$$\begin{aligned} t_1 &\longrightarrow t_{a_1} t_{b_1} & t_2 &\longrightarrow t_{a_1} t_{b_2} & t_3 &\longrightarrow \frac{t_{a_2}}{t_{a_1}}, \\ t_1 &\longrightarrow t_{a_2} t_{b_1} & t_2 &\longrightarrow t_{a_2} t_{b_2} & t_3 &\longrightarrow \frac{t_{a_1}}{t_{a_2}}. \end{aligned} \quad (4.29)$$

These two copies differ by $t_{a_1} \leftrightarrow t_{a_2}$.

The conjecture is inspired by the cohomological answer [35]

$$Z_{NCC}^{coho}(q, z) = M(-q)^2 \prod_{k \geq 1} \left(1 - (-q)^k z\right)^k \left(1 - (-q)^k z^{-1}\right)^k \quad (4.30)$$

with $q = q_0 q_1$ and $z = q_1$. The ‘‘Kähler’’ dependent part has the $q_1 + 1/q_1$ structure. The remaining part is two copies of the MacMahon function.

We define the full partition function by symmetrization

$$\begin{aligned} Z_{NCC}(t_{a_1}, t_{a_2}, t_{b_1}, t_{b_2}, t_4, t_5, q_1) &= \mathbf{S}^\bullet \mathcal{F}_{NCC}[t_{a_1}, t_{a_2}, t_{b_1}, t_{b_2}, t_4, t_5, q_1] \\ &= \exp \sum_{r \geq 1} \frac{1}{r} \mathcal{F}_{NCC}[t_{a_1}^r, t_{a_2}^r, t_{b_1}^r, t_{b_2}^r, t_4^r, t_5^r, q_1^r] \end{aligned} \quad (4.31)$$

Using MATHEMATICA we find the following relations

$$\begin{aligned} Z_{NCC}(\{t_i\}, q_0 q_1 \kappa^{-1/2}, (q_0 q_1)^{-1} \kappa^{-1/2}, q_1) &= \sum_{(d_\circ, d_\bullet)} (-q_0)^{d_\circ} q_1^{d_\bullet} \chi(\mathcal{M}_{d_\circ, d_\bullet}, \mathcal{O}_{\mathcal{M}_{d_\circ, d_\bullet}}^{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2}) \\ Z_{NCC}(\{t_i\}, q_0 q_1 \kappa^{-1/2}, (q_0 q_1)^{-1} \kappa^{-1/2}, -q_1) &= \sum_{(d_\circ, d_\bullet)} q_0^{d_\circ} (-q_1)^{d_\bullet} \chi(\mathcal{M}_{d_\circ, d_\bullet}, \mathcal{O}_{\mathcal{M}_{d_\circ, d_\bullet}}^{\text{vir}} \otimes \mathcal{K}_{\text{vir}}^{1/2}) \end{aligned} \quad (4.32)$$

where the different signs in the right hand side correspond to the different signs in the prefactor of the virtual structure sheaf. The substitution $t_4 = q_0 q_1 \kappa^{-1/2}$ and $t_5 = (q_0 q_1)^{-1} \kappa^{-1/2}$ is the same as in \mathbb{C}^3 with counting parameter $q = q_0 q_1$. The compact form of (4.28) allows us easily to switch between sign conventions.

4.6 Refined and unrefined limits

To take the refined limit we scale the toric weights to zero or infinity, in such a way that the product κ remains constant [30]. When taking this scaling limit each fixed point scales as

$$\hat{\mathbf{a}}\left(\sum_i w_i - \kappa \sum_i w_i^{-1}\right) \longrightarrow (-\kappa^{1/2})^{\text{Index}} \quad (4.33)$$

where

$$\text{Index} = \#\{i | w_i \longrightarrow 0\} - \#\{i | w_i \longrightarrow \infty\} \quad (4.34)$$

is the number of weights which go to zero minus the number of weights which go to infinity. In our case we take the torus where t_{a_1} , t_{a_2} and t_{b_1} go to zero, the first very fast and the latter two very slowly, while t_{b_2} goes to infinity fast, such that the product of the toric weights is a constant¹.

We compute the limit of the fixed points one by one using MATHEMATICA. Before stating the result we need to introduce a bit of notation, to take into account the multiple refinement of the MacMahon function, as well as different signs in the conventions. Introduce [23]

$$G_{re}[a_0, a_1] = \prod_{j=0}^{a_0-1} \left(1 - \kappa^{-\frac{a_0}{2} + \frac{1}{2} + j} (-y_0)^{a_0} y_1^{a_1}\right) \quad (4.35)$$

$$G_{im}[a_0, a_1, \epsilon, \delta] = \prod_{j=0}^{a_0-1} \left(1 - \kappa^{-\frac{a_0}{2} + \epsilon + j} (-y_0)^{a_0} y_1^{a_1}\right)^{-1} \left(1 - \kappa^{-\frac{a_0}{2} + \delta + j} (-y_0)^{a_0} y_1^{a_1}\right)^{-1} \quad (4.36)$$

¹To be more concrete, we take the subtorus $\{t_{a_1} \rightarrow x^{1000}, t_{a_2} \rightarrow x^{10}, t_{b_1} \rightarrow x^{10}, t_{b_2} \rightarrow 1/x^{1020}\}$ and then send $x \rightarrow 0$. The specific form of the result will depend in general by the chosen subtorus

and consider

$$G(\epsilon, \delta) = \prod_{i \geq 1} G_{re}[i, i-1] G_{re}[i-1, i] G_{im}[i, i, \epsilon, \delta] \quad (4.37)$$

Using this notation our result is $G(\epsilon = 0, \delta = 1)$ which explicitly reads

$$\begin{aligned} Z_{NCC}^{ref}(q_0, q_1; \kappa) &= G(\epsilon = 0, \delta = 1) = \\ &1 + q_0 + \left(-\frac{1}{\sqrt{k}} - \sqrt{k}\right) q_0 q_1 + \left(-\frac{2}{\sqrt{k}} - 2\sqrt{k}\right) q_0^2 q_1 + \left(-\frac{1}{\sqrt{k}} - \sqrt{k}\right) q_0^3 q_1 + q_0 q_1^2 \\ &+ \left(4 + \frac{2}{k} + 2k\right) q_0^2 q_1^2 + \left(6 + \frac{4}{k} + 4k\right) q_0^3 q_1^2 + \left(4 + \frac{2}{k} + 2k\right) q_0^4 q_1^2 + \left(-\frac{2}{\sqrt{k}} - 2\sqrt{k}\right) q_0^2 q_1^3 \\ &+ \left(-\frac{3}{k^{3/2}} - \frac{9}{\sqrt{k}} - 9\sqrt{k} - 3k^{3/2}\right) q_0^3 q_1^3 + \left(-\frac{7}{k^{3/2}} - \frac{15}{\sqrt{k}} - 15\sqrt{k} - 7k^{3/2}\right) q_0^4 q_1^3 \\ &+ \left(6 + \frac{4}{k} + 4k\right) q_0^3 q_1^4 + \left(26 + \frac{5}{k^2} + \frac{18}{k} + 18k + 5k^2\right) q_0^4 q_1^4 + \dots \end{aligned} \quad (4.38)$$

Note that this series is explicitly invariant for $k \leftrightarrow 1/k$.

This should be compared with other results in the literature. For example the refined noncommutative conifold partition function of [14] can be written as $G(1/2, 1/2)$ in this notation, while the motivic partition function computed in [23] is $G(1, 2)$. Interestingly our partition function agrees with a theorem of [23] according to which the MacMahon factor in front is not a square of the same refinement; and also agrees with the result physical expectation that such partition functions should have coefficients which are linear combinations of $SU(N)$ characters. Note however that the differences between the partition functions can all be traced to the ambiguity of defining the MacMahon, arising from the non-compactness of the moduli space, see [25] for a recent discussion. Furthermore our result depends explicitly on the torus chosen to take the refined limit. Different tori will give a priori different results. We won't pursue the issue here further, and hope to return to a more systematic study of the refined limits in the future.

Finally we can rewrite the refined partition function in compact form introducing

$$F_{NCC}^r[q_0, q_1, k] = -\frac{q_0 \left(-\sqrt{k} (q_1^2 + 1) + kq_1 + q_1\right)}{\left(\sqrt{k} - q_0 q_1\right) \left(\sqrt{k} q_0 q_1 - 1\right)} \quad (4.39)$$

So that $\mathbf{S}^\bullet F_{NCC}^r[-q_0, q_1, k]$ gives our refined result.

In the unrefined limit, the noncommutative partition function has the following relation with the large radius partition functions [35]

$$Z_{NCC}^{coho,red}(q, z) = Z_{resolved}^{coho,red}(q, z) Z_{resolved}^{coho,red}\left(q, \frac{1}{z}\right) \quad (4.40)$$

where the reduced partition functions are obtained by removing the MacMahon factors. From this we are lead to conjecture an analog relation between the reduced K-theoretic partition functions. In

particular this produces a prediction for the large radius conifold partition function, as

$$Z'_{DT}(t_4, t_5, q_1) = S^\bullet \frac{q_1}{\left(-\frac{1}{\sqrt{t_4}} + \sqrt{t_4}\right) \left(-\frac{1}{\sqrt{t_5}} + \sqrt{t_5}\right)} \quad (4.41)$$

which agrees with the result of [19], upon the usual change of variables $t_4 = q/\sqrt{\kappa}$ and $t_5 = 1/(q\sqrt{\kappa})$. A more systematic analysis of the relation between the noncommutative chamber and the other chambers will appear elsewhere [11].

5 Orbifold Singularities

Now we will move to orbifold singularities in toric Calabi-Yau threefolds and set up a formalism to compute the K-theoretic Donaldson-Thomas invariants. The counting of BPS states for orbifold singularities is obtained in terms of quivers associated to the singularity via the three dimensional McKay correspondence. From this quiver we set up the localization computation as in the previous sections. For singularities of the form \mathbb{C}^3/Γ with $\Gamma \subset \mathrm{SL}(3, \mathbb{C})$, the formalism reduces to an appropriate Γ projection on the \mathbb{C}^3 results. The results can be interpreted as membrane contributions to the M-theory index at the singularity. We will obtain closed forms of the partition function for certain Γ . The results in this section are based on and generalized those of [7, 8]

5.1 BPS states and McKay quivers

In this Section we will focus on BPS countings associated with quivers which arise from orbifold singularities of the form \mathbb{C}^3/Γ where Γ is a finite subgroup of $\mathrm{SL}(3, \mathbb{C})$. In this case the relevant quiver is the McKay quiver \mathbf{Q}_Γ which is constructed out of the representation theory data of Γ as follows. The set of vertices of \mathbf{Q}_Γ is in one to one correspondence with the irreducible one-dimensional representations ρ_a of Γ . Such representations form a group which we will denote by $\hat{\Gamma}$. The arrow structure of \mathbf{Q}_Γ is dictated by the tensor product decomposition

$$\bigwedge^i Q \otimes \rho_a = \bigoplus_{b \in \hat{\Gamma}} a_{ba}^{(i)} \rho_b, \quad \text{with } a_{ba}^{(i)} = \dim_{\mathbb{C}} \mathrm{Hom}_\Gamma(\rho_b, \bigwedge^i Q \otimes \rho_a), \quad (5.1)$$

where $a_{ab}^{(1)}$ is the number of arrows going from node ρ_b to node ρ_a . Here $Q = \rho_{a_1} \oplus \rho_{a_2} \oplus \rho_{a_3}$ denotes the fundamental three-dimensional representation of Γ . This is defined by the action of Γ on \mathbb{C}^3 . The condition that $\Gamma \subset \mathrm{SL}(3, \mathbb{C})$ implies that $a_1 + a_2 + a_3 = 0$. There are $a_{ab}^{(1)}$ arrows going from node ρ_b to node ρ_a . This construction is the higher dimensional generalization of the ordinary McKay quiver for ALE singularities.

Consider now a representation of the McKay quiver. From the individual vector spaces V_a assigned to the vertices labelled ρ_a we form $V = \bigoplus_{a \in \hat{\Gamma}} V_a \otimes \rho_a^\vee$ (here ρ^\vee is the conjugate representation). The linear maps between vector spaces are determined by $B \in \mathrm{Hom}_\Gamma(V, Q \otimes V)$. In other words the representations of the McKay quiver have a natural Γ -module structure. If we introduce the notation $B_\alpha^{(a)} \in \mathrm{Hom}_{\mathbb{C}}(V_a, V_{a+a_\alpha})$, the relations between the quiver maps can be derived from the superpotential

$$\mathcal{W}_\Gamma = \sum_{\alpha \in \hat{\Gamma}} B_1^{(a+a_2+a_3)} \left(B_2^{(a+a_3)} B_3^{(a)} - B_3^{(a+a_2)} B_2^{(a)} \right). \quad (5.2)$$

The Jacobian algebra $A_\Gamma = \mathbb{C}Q_\Gamma / \langle \partial W_\Gamma \rangle$ is a noncommutative crepant resolution of the singularity \mathbb{C}/Γ .

As showed in [7], from the point of view of the six dimensional topological theory living on the D6 brane the dimensions of the vector spaces $\dim V_a = k_a$ which enter in the isotypical decomposition $V = \bigoplus_{a \in \hat{\Gamma}} V_a \otimes \rho_a^\vee$, correspond to instanton configurations which transform in the irreducible representation ρ_a . Similarly in the framing node decomposition $W = \bigoplus_{a \in \hat{\Gamma}} W_a \otimes \rho_a^\vee$ the vector spaces label boundary conditions at infinity, that is flat gauge fields transforming in a certain representation ρ , which decomposes into irreducible representations ρ_a . In this note we will only consider trivial representations at infinity: the framing node is connected to the quiver only to the vertex associated with the trivial representation.

The formation of D0/D2 bound states can be seen from the point of view of the quiver quantum mechanics which arises from the quantization of the collective coordinates around each instanton configuration; or more precisely from a direct parametrization of the moduli space of ideal sheaves [7]. This quiver quantum mechanics localizes onto fixed points of its BRST operator. This operator can be twisted by the natural toric action on \mathbb{C}^3 , since the \mathbb{T}^3 action and the Γ action commute. Fixed points are again classified by plane partitions π , except that now they carry a Γ action. The reason for this is that the superpotential W_Γ can be seen as the same as the superpotential W where one takes into account also the Γ -action on the representation spaces $\text{Hom}_{\mathbb{C}}(V, V)$. As a consequence the classification of the fixed points in terms of 3d partitions remains the same, except that now each box also carries a Γ -weight, corresponding to the (inverse of the) weight of the monomial in $\mathbb{C}[z_1, z_2, z_3]$ represented by that box.

5.2 Localization on quiver moduli spaces

The relevant quiver representation moduli space is the quotient stack

$$\mathcal{M}_\Gamma(\mathbf{v}, \mathbf{w}) = [\text{Rep}(\hat{Q}_\Gamma; W_\Gamma) / \prod_{r \in \hat{\Gamma}} GL(k_r, \mathbb{C})], \quad (5.3)$$

where $\text{Rep}(\hat{Q}_\Gamma; W)$ is the subset of

$$\text{Hom}_\Gamma(V, Q \otimes V) \oplus \text{Hom}_\Gamma(V, \bigwedge^3 Q \otimes V) \oplus \text{Hom}_\Gamma(W, V) \quad (5.4)$$

cut out by the F-term equations derived from the superpotential W_Γ . Here $\mathbf{v} = \{v_a\}_{a \in \hat{\Gamma}}$ where $v_a = \dim_{\mathbb{C}} V_a$, and similarly for \mathbf{w} . Since we are considering only trivial boundary conditions we can assume $W = \mathbb{C}$. Therefore in the following we will take $w_0 = 1$ as the only non trivial component of \mathbf{w} . This correspond to framing the McKay quiver Q_Γ by adding a single vertex connected to the quiver by a single arrow, which ends on the node associated with the trivial representation. In the gauge theory analogy this corresponds to fixing boundary condition at infinity [7]. Boundary conditions require the gauge connection to be flat and therefore to be labelled by an irreducible representation of Γ . In this case we can choose the trivial representation. Picking a different representation does not change substantially the instanton counting. We will denote $\mathcal{M}_\Gamma(\mathbf{v}, (1, 0, \dots, 0))$ by $\mathcal{M}_\Gamma(\mathbf{v})$.

In instanton counting problems one can construct a local model for the moduli space by studying

a deformation complex, which parametrizes linearized field configurations near solutions of the equations of motion, up to gauge invariance. In the case of orbifold singularities it is given by [7]

$$\begin{array}{ccc} \text{Hom}_\Gamma(V_\pi, V_\pi) & \longrightarrow & \begin{array}{c} \text{Hom}_\Gamma(V_\pi, V_\pi \otimes Q) \\ \oplus \\ \text{Hom}_\Gamma(\mathbb{C}, V_\pi) \\ \oplus \\ \text{Hom}_\Gamma(V_\pi, V_\pi \otimes \wedge^3 Q) \end{array} & \longrightarrow & \begin{array}{c} \text{Hom}_\Gamma(V_\pi, V_\pi \otimes \wedge^2 Q) \\ \oplus \\ \text{Hom}_\Gamma(V_\pi, \mathbb{C} \otimes \wedge^3 Q) \end{array} \end{array} \quad (5.5)$$

If we forget about the Γ action, the above complex becomes precisely the relevant complex for \mathbb{C}^3 , whose space of configurations is given by $\text{Hilb}(\mathbb{C}^3, n)$.

From the deformation complex we can read directly the virtual tangent space

$$(T_\Gamma^{\text{vir}})_\pi = (V_\pi - \overline{V}_\pi t_1 t_2 t_3 + (1 - t_1)(1 - t_2)(1 - t_3) \overline{V}_\pi \otimes V_\pi)^\Gamma. \quad (5.6)$$

Fixed points of the toric action are still classified by 3d partitions, since the orbifold group Γ is a subgroup of the torus \mathbb{T}^3 . The vector space V can be decomposed at a fixed point π as

$$V_\pi = \bigoplus_{a \in \widehat{\Gamma}} (P_{l,a} \otimes \rho_a^\vee) \quad (5.7)$$

The modules $P_{l,a}$ are associated to the decomposition of $\sum_{(n_1, n_2, n_3) \in \pi_l} t_1^{-n_1+1} t_2^{-n_2+1} t_3^{-n_3+1}$ as a Γ -module, while the ρ_a^\vee factors keep track of the representation. The vector spaces V have a decomposition as a sum of monomials in the toric variables, and each monomial has a definite transformation under the orbifold action. Each 3d partition carries an action of the group Γ induced by the fundamental representation $Q = \rho_{a_1} \oplus \rho_{a_2} \oplus \rho_{a_3}$. In plain words each box of the partition is associated to a character of Γ depending on the transformation of the monomial in the toric variables associated with that box under the Γ action.

We define the K-theoretic BPS partition function as

$$Z_\Gamma(\{p_i\}, \{t_i\}) = \chi(\mathcal{M}_\Gamma, \widehat{\mathcal{O}}_\Gamma^{\text{vir}}) \quad (5.8)$$

where additional counting variables $\{p_i\}$ can be introduced to keep track of the dimensions of the representations by a simple modification of the virtual structure sheaf. The moduli space decomposes as a disjoint union

$$\mathcal{M}_\Gamma = \bigsqcup \mathcal{M}_\Gamma(\mathbf{v}). \quad (5.9)$$

The modified virtual structure sheaf is

$$\widehat{\mathcal{O}}_\Gamma^{\text{vir}} = \text{prefactor } \mathcal{O}_\Gamma^{\text{vir}} \otimes \mathcal{K}_{\text{vir}, \Gamma}^{1/2} \quad (5.10)$$

where the prefactor contains counting parameters to label each component of the moduli space, and is a vector of the form $\prod_{i \in \mathbb{Q}_0} p_i^{v_i}$.

As before, all the information about the modified virtual structure sheaf $\widehat{\mathcal{O}}_\Gamma^{\text{vir}}$ is contained in T_Γ^{vir} .

The virtual structure sheaf parametrizes obstructions while $\mathcal{K}_{\text{vir},\Gamma}^{1/2} = \det^{-1/2} T_{\Gamma}^{\text{vir}}$.

Explicitly we can write

$$(T_{\pi}^{\text{vir}})^{\Gamma} = (T_{\pi} \widetilde{M})^{\Gamma} - (\kappa \otimes T_{\pi}^* \widetilde{M})^{\Gamma} = \sum_i a_i - \sum_i b_i = \sum_i w_i^{\Gamma} - \sum_i \frac{\kappa}{w_i^{\Gamma}} \quad (5.11)$$

where the weights w_i^{Γ} are in general monomials in the toric weights which are *invariant* under the Γ -action. Note that since Γ has trivial determinant, the combination $\kappa = t_1 t_2 t_3$ is invariant.

At a fixed point π the virtual structure sheaf can be written in localized K-theory as

$$(\mathcal{O}_{\pi}^{\text{vir}})^{\Gamma} = \mathcal{O}_{\pi} \prod_i \frac{1 - b_i^{-1}}{1 - a_i^{-1}}. \quad (5.12)$$

Similarly

$$\mathcal{K}_{\text{vir},\Gamma} = \frac{\det_{\Gamma}(T^* \widetilde{M} \otimes \kappa)}{\det_{\Gamma} T \widetilde{M}} \quad (5.13)$$

Finally, putting everything together (in localized K-theory)

$$(\widehat{\mathcal{O}}_{\pi}^{\text{vir}})^{\Gamma} = \mathcal{O}_{\pi} \prod_i \frac{1 - b_i^{-1}}{1 - a_i^{-1}} \otimes (K^{\text{vir}})^{1/2} = \mathcal{O}_{\pi} \prod_i \frac{(\kappa/w_i^{\Gamma})^{1/2} - (\kappa/w_i^{\Gamma})^{-1/2}}{(w_i^{\Gamma})^{1/2} - (w_i^{\Gamma})^{-1/2}} \quad (5.14)$$

where again we stress that the weights are Γ -invariant since $\kappa = t_1 t_2 t_3$ is invariant due to the Calabi-Yau condition.

The refined limit is taken precisely as in the affine space case, by sending the toric weights t_i to zero or infinity while keeping their product κ constant. Sending $\kappa \rightarrow 1$ reproduces the cohomological limit [7]

$$Z_{\Gamma}^{\text{cho}}(\{p_i\}) = \sum_{\pi} \prod_{i \in \mathbf{Q}} p_i^{v_i} \int_{[\mathcal{M}_{\Gamma}(\mathbf{v})]^{\text{vir}}} 1 \quad (5.15)$$

5.3 The case of A -fibered singularities

We begin by studying a series of examples of orbifold singularities which are fibrations of ADE singularities over the complex plane. Such singularities have semi-small crepant resolutions which are fibrations of ALE spaces over the complex plane. We will however work in the noncommutative crepant resolution chamber, using quiver representation data to compute BPS partition functions. In this example we will consider the simplest case where $X = \mathbb{C}^2/\mathbb{Z}_p \times \mathbb{C}$ and the singularity is resolved as a noncommutative crepant resolution. In this example we will also denote by Γ the orbifold group acting on the \mathbb{C}^2 factor, hoping that this will not cause confusion.

Geometry and representation theory We will consider orbifolds of the form $X = \mathbb{C}^2/\Gamma \times \mathbb{C}$ where $g \in \Gamma = \mathbb{Z}_n$ acts on $\mathbb{C}[z_1, z_2, z_3]$ as

$$g \cdot (z_1, z_2, z_3) = (\zeta z_1, \zeta^{-1} z_2, z_3) \quad (5.16)$$


where ζ is a n -th root of unity. At large radius such singularities have semi-small crepant resolutions given by trivial fibration of a resolved A_{n-1} singularity over \mathbb{C} . The resolved geometry is toric and the exceptional locus consists of -2 curves which intersect transversally. The intersection matrix of the exceptional curves is the negative of the Cartan matrix of A_{n-1} .

At short distances smooth geometry is replaced by representation theory data. In our case this is given by the McKay quiver associated to the singularity. The nodes of the quiver are in one to one correspondence with the irreducible representations ρ_α of $\Gamma \subset \text{SU}(2) \subset \text{SU}(3)$. The arrows are given by the multiplicities of the decomposition of $Q \otimes \rho_r$ into irreducible representations, where Q denotes the fundamental representation.

For example if we take $\Gamma = \mathbb{Z}_3$, the defining representation is $Q = \rho_1 \oplus \rho_2 \oplus \rho_0$. From (5.1) we now see

$$(a_{rs}^{(1)}) = (a_{rs}^{(2)}) = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}. \quad (5.17)$$

The quiver Q constructed with these data is



The quiver \hat{Q} which is relevant to the problem of computing degeneracies of BPS bound states is obtained by framing Q . The framing node represents a single D6 brane and is connected to Q by a single arrow to the node labelled by v_0 . From a gauge theory perspective the choice of this node corresponds to the choice of a superselection sector where the gauge connection at infinity is flat and in the trivial representation of Γ .

We will first go through a few examples and then make a general conjecture for the partition function for arbitrary $\Gamma = \mathbb{Z}_n$.

BPS partition function for $\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}$ We begin with a simple example. In this case the Γ -action on $\mathbb{C}[z_1, z_2, z_3]$ is

$$g \cdot (z_1, z_2, z_3) = (\zeta z_1, \zeta z_2, z_3) \quad (5.19)$$

with $\zeta^2 = 1$.

At a fixed point the representation spaces decompose

$$V = V_0 + V_1 = \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{(i-j=0 \pmod{2})} + \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{(i-j=1 \pmod{2})} \quad (5.20)$$

we set $|\pi_i| = \dim V_r$ for $r = 0, 1$. These measure the number of boxes in the 3d partition which transform in the conjugacy class labelled by r . In terms of this decomposition the virtual tangent

space has the form

$$(T_\pi^{\text{vir}})^\Gamma = V_0 - \bar{V}_0 t_1 t_2 t_3 - (V_0 \otimes \bar{V}_0 + V_1 \otimes \bar{V}_1)(1 + t_1 t_2)(1 - t_3) - (V_0 \otimes \bar{V}_1 + V_1 \otimes \bar{V}_0)(-t_1 - t_2)(1 - t_3). \quad (5.21)$$

We can now compute the partition function

$$Z_{\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}}(p_1, p_2, \{t_i\}) = \sum_n \sum_{|\pi|=n} p_0^{|\pi_0|} p_1^{|\pi_1|} \hat{a}((T_\pi^{\text{vir}})^\Gamma) \quad (5.22)$$

where the second sum is over all three dimensional partitions with fixed number of boxes. We can see directly the first few terms

$$\begin{aligned} Z_{\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}}(p_1, p_2, \{t_i\}) &= 1 + \frac{p_0(\kappa - t_3)}{\sqrt{\kappa}(-1 + t_3)} + \frac{p_0 p_1(-\kappa + t_1^2)(\kappa t_1 - t_2)(\kappa - t_3)}{\kappa^{3/2}(-1 + t_1^2)(t_1 - t_2)(-1 + t_3)} \\ &+ \frac{p_0 p_1(t_1 - \kappa t_2)(-\kappa + t_2^2)(\kappa - t_3)}{\kappa^{3/2}(t_1 - t_2)(-1 + t_2^2)(-1 + t_3)} + \frac{p_0^2(-\kappa + t_3)(-\kappa + t_3^2)}{\kappa(-1 + t_3)^2(1 + t_3)} + \dots \end{aligned} \quad (5.23)$$

We conjecture that this partition function has a closed form presentation as the symmetrized form of a certain orbifold generalization of Nekrasov's function. Define the function

$$\begin{aligned} \mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5; p_1) &= \left(p_1 + \frac{1}{p_1}\right) \frac{(\kappa/t_3)^{1/2} - (t_3/\kappa)^{1/2}}{\prod_3^5 (t_i^{1/2} - t_i^{-1/2})} \\ &+ \mathcal{F}_{\mathbb{C}^3}(t_1/t_2, t_2^2, t_3, t_4, t_5) + \mathcal{F}_{\mathbb{C}^3}(t_1^2, t_2/t_1, t_3, t_4, t_5) \end{aligned} \quad (5.24)$$

Then we claim that

$$Z_{\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}}(q, p_1, \{t_i\}) = \mathcal{S}^\bullet \mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_2 \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5; p_1) \quad (5.25)$$

with the identifications $\kappa = t_1 t_2 t_3$, $t_4 = q/\sqrt{\kappa}$, $t_5 = 1/(q\sqrt{\kappa})$ and finally $q = -p_0 p_1$. This claim is checked order by order in the supporting MATHEMATICA file.

BPS partition function for $\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}$ Now we consider the orbifold $\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}$ with action

$$g \cdot (z_1, z_2, z_3) = (\zeta z_1, \zeta^{-1} z_2, z_3) \quad (5.26)$$

with $\zeta^3 = 1$. We decompose the representation space as $V = V_0 + V_1 + V_2$ with

$$\begin{aligned} V_0 &= \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{i-j=0 \pmod 3} \\ V_1 &= \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{i-j=1 \pmod 3} \\ V_2 &= \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{i-j=2 \pmod 3} \end{aligned} \quad (5.27)$$

In particular we set $|\pi_r| = \dim_{\mathbb{C}} V_r$, effectively obtained by setting the toric weights to 1.

Then the virtual tangent space decomposes as

$$\begin{aligned}
(T_\pi^{\text{vir}})^{\mathbb{Z}_3} &= V_0 - \bar{V}_0 t_1 t_2 t_3 - (V_0 \bar{V}_0 + V_1 \bar{V}_1 + V_2 \bar{V}_2)(1 + t_1 t_2)(1 - t_3) \\
&\quad - (V_0 \bar{V}_1 + V_1 \bar{V}_2 + V_2 \bar{V}_0)(-t_2)(1 - t_3) \\
&\quad - (V_0 \bar{V}_2 + V_1 \bar{V}_0 + V_2 \bar{V}_1)(-t_1)(1 - t_3).
\end{aligned} \tag{5.28}$$

We can now compute the partition function

$$\mathbb{Z}_{\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}}(\{p_i\}, \{t_i\}) = \sum_n \sum_{|\pi|=n} p_0^{|\pi_0|} p_1^{|\pi_1|} p_2^{|\pi_2|} \hat{\mathbf{a}} \left((T_\pi^{\text{vir}})^{\mathbb{Z}_3} \right). \tag{5.29}$$

We can see directly the first few terms

$$\begin{aligned}
\mathbb{Z}_{\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}}(\{p_i\}, \{t_i\}) &= 1 + \frac{p_0(\kappa - t_3)}{\sqrt{\kappa}(-1 + t_3)} + \frac{p_0 p_1(\kappa - t_3)}{\sqrt{\kappa}(-1 + t_3)} + \frac{p_0 p_2(\kappa - t_3)}{\sqrt{\kappa}(-1 + t_3)} \\
&\quad + \frac{p_0 p_1 p_2 (-\kappa + t_1^3) (\kappa t_1^2 - t_2) (\kappa - t_3)}{\kappa^{3/2} (-1 + t_1^3) (t_1^2 - t_2) (-1 + t_3)} + \frac{p_0 p_1 p_2 (t_1^2 - \kappa t_2) (\kappa t_1 - t_2^2) (\kappa - t_3)}{\kappa^{3/2} (t_1^2 - t_2) (t_1 - t_2^2) (-1 + t_3)} \\
&\quad + \frac{p_0 p_1 p_2 (t_1 - \kappa t_2^2) (-\kappa + t_2^3) (\kappa - t_3)}{\kappa^{3/2} (t_1 - t_2^2) (-1 + t_2^3) (-1 + t_3)} + \frac{p_0^2 p_1 (-\kappa + t_3)^2}{\kappa(-1 + t_3)^2} + \frac{p_0^2 p_2 (-\kappa + t_3)^2}{\kappa(-1 + t_3)^2} \\
&\quad + \frac{p_0^2 (-\kappa + t_3) (-\kappa + t_3^2)}{\kappa(-1 + t_3)^2 (1 + t_3)} - \frac{p_0^3 (-\kappa + t_3) (-\kappa + t_3^2) (-\kappa + t_3^3)}{\kappa^{3/2} (-1 + t_3)^3 (1 + t_3) (1 + t_3 + t_3^2)} + \dots
\end{aligned} \tag{5.30}$$

Now we want to recast the partition function as a symmetrized form of an orbifold function. We define the orbifold function as

$$\begin{aligned}
\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5; p_1, p_2) &= \left(p_1 + \frac{1}{p_1} + p_2 + \frac{1}{p_2} + p_1 p_2 + \frac{1}{p_1 p_2} \right) \frac{(\kappa/t_3)^{1/2} - (t_3/\kappa)^{1/2}}{\prod_3^5 (t_i^{1/2} - t_i^{-1/2})} \\
&\quad + \mathcal{F}_{\mathbb{C}^3}(t_1^3, t_2/t_1^2, t_3, t_4, t_5) + \mathcal{F}_{\mathbb{C}^3}(t_1/t_2^2, t_2^3, t_3, t_4, t_5) + \mathcal{F}_{\mathbb{C}^3}(t_1^2/t_2, t_2^2/t_1, t_3, t_4, t_5)
\end{aligned} \tag{5.31}$$

with the usual identifications $\kappa = t_1 t_2 t_3$, $t_4 = q/\sqrt{\kappa}$, $t_5 = 1/(q\sqrt{\kappa})$ and finally $q = -p_0 p_1 p_2$. Then we can check order by order that

$$\mathbb{Z}_{\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}}(\{p_i\}, \{t_i\}) = \mathcal{S}^\bullet \mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_3 \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5; p_1, p_2) \tag{5.32}$$

Note that

$$\begin{aligned}
&\frac{1}{3} \sum_{r=0}^2 \frac{1}{(1 - w^r t_1)(1 - w^{2r} t_2)(1 - t_3)} \\
&= \frac{1}{(1 - t_1^3)(1 - \frac{t_2}{t_1^2})(1 - t_3)} + \frac{1}{(1 - \frac{t_1}{t_2^2})(1 - t_2^3)(1 - t_3)} + \frac{1}{(1 - \frac{t_1^2}{t_2})(1 - \frac{t_2^2}{t_1})(1 - t_3)}
\end{aligned} \tag{5.33}$$

is the generating function of \mathbb{Z}_3 invariant monomials in the variables t_i . The left hand side gives invariants by construction since it averages the generating function of symmetric monomials in the

toric variables, over all the images of the orbifold action. The right hand side follows from a direct computation. This is the origin of the three copies of the \mathbb{C}^3 generating function above.

General formula These set of results lead us to conjecture a general formula for arbitrary $\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}$. Consider first the decomposition

$$\frac{1}{n} \sum_{r=0}^{n-1} \frac{1}{(1-w^r t_1)(1-w^{2r} t_2)(1-t_3)} = \frac{1}{1-t_3} \sum_{k=0}^{n-1} \frac{1}{\left(1-\frac{t_1^{n-k}}{t_2^k}\right)\left(1-\frac{t_2^{k+1}}{t_1^{n-1+k}}\right)} \quad (5.34)$$

where $w = \exp \frac{2\pi i}{n}$ is an n^{th} root of unity. We define the function

$$\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5) = \sum_{k=0}^{n-1} \mathcal{F}_{\mathbb{C}^3} \left(\frac{t_1^{n-k}}{t_2^k}, \frac{t_2^{k+1}}{t_1^{n-1+k}}, t_3, t_4, t_5 \right) \quad (5.35)$$

which generates Γ -invariant monomials in the toric parameters.

Introduce now the variables $p_{[r,s]} = p_r p_{r+1} \cdots p_s$. We define the reduced function

$$\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}^{(r)}(\{p_i\}, t_1, t_2, t_3, t_4, t_5) = \sum_{0 < r \leq s < n} \left(p_{[r,s]} + \frac{1}{p_{[r,s]}} \right) \frac{(\kappa/t_3)^{1/2} - (t_3/\kappa)^{1/2}}{\prod_{i=3}^5 (t_i^{1/2} - t_i^{-1/2})} \quad (5.36)$$

We claim that

$$Z_{\mathbb{C}^2/\mathbb{Z}_k \times \mathbb{C}}(\{p_i\}, \{t_i\}) = \mathcal{S}^\bullet \left(\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5) + \mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}^{(r)}(\{p_i\}, t_1, t_2, t_3, t_4, t_5) \right) \quad (5.37)$$

with the identifications $\kappa = t_1 t_2 t_3$, $t_4 = q/\sqrt{\kappa}$, $t_5 = 1/(q\sqrt{\kappa})$, $q = -p_0 p_1 \cdots p_{n-1}$.

We have checked this formula order by order for $\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}$ with $n = 2, 3, 4, 5, 6$.

Refined limit Now we take a refined limit by sending $t_3 \rightarrow 0$, $t_1 \rightarrow 0$ in such a way that t_3 approaches zero much faster and $t_2 \rightarrow \infty$ in such a way that the product $\kappa = t_1 t_2 t_3$ remains constant.

First recall that

$$\frac{\left(\frac{\kappa}{w}\right)^{1/2} - \left(\frac{w}{\kappa}\right)^{1/2}}{(w)^{1/2} - (w)^{-1/2}} \rightarrow \begin{cases} -\kappa^{-1/2} & \text{if } w \rightarrow \infty \\ -\kappa^{1/2} & \text{if } w \rightarrow 0 \end{cases} \quad (5.38)$$

Next we have the limits

$$\frac{t_1^{n-k}}{t_2^k} \rightarrow 0, \quad \frac{t_2^{k+1}}{t_1^{n-1+k}} \rightarrow \infty, \quad (5.39)$$

since $0 \leq k \leq n-1$. Therefore the contributions of each ratio to the refined limit exactly compensate each other for each k and the leading term is always determined by the behaviour of t_3 . Therefore

each term in $\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}$ contributes equally. If we set $t_4 = q/\sqrt{\kappa}$, $t_5 = 1/(q\sqrt{\kappa})$, then

$$\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}(t_1, t_2, t_3, t_4, t_5) \longrightarrow n(-\kappa^{1/2}) \frac{1}{(1 - q/\sqrt{\kappa})(1 - 1/(q\sqrt{\kappa}))} = n\kappa^{1/2} \frac{q}{(1 - q\sqrt{\kappa})(1 - q/\sqrt{\kappa})} \quad (5.40)$$

In the function $\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}^{(r)}$ the only term relevant to the refined limit is the t_3 dependent ratio. Overall this function becomes

$$\mathcal{F}_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}^{(r)}(\{p_i\}, t_1, t_2, t_3, t_4, t_5) \longrightarrow \sum_{0 < r \leq s < n} \left(p_{[r,s]} + \frac{1}{p_{[r,s]}} \right) (\kappa^{1/2}) \frac{q}{(1 - q\sqrt{\kappa})(1 - q/\sqrt{\kappa})} \quad (5.41)$$

By writing all the pieces together we find the full refined limit

$$Z_{\mathbb{C}^2/\mathbb{Z}_n \times \mathbb{C}}^{ref}(\{p_i\}; \kappa) = \mathbf{S}^\bullet \left(\left(n + \sum_{0 < r \leq s < n} \left(p_{[r,s]} + \frac{1}{p_{[r,s]}} \right) \right) \kappa^{1/2} \frac{q}{(1 - q\sqrt{\kappa})(1 - q/\sqrt{\kappa})} \right) \quad (5.42)$$

where again $q = -p_0 p_1 \cdots p_{n-1}$.

We can now write this generating function in product formula. First note that

$$\begin{aligned} \mathbf{S}^\bullet x \frac{\kappa^{1/2} q}{(1 - q\sqrt{\kappa})(1 - q/\sqrt{\kappa})} &= \exp \left[- \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \log \left(1 - x q^{i+j-1} \kappa^{1/2(j-i+1)} \right) \right] \\ &= \prod_{i,j=1}^{\infty} \left(1 - x q^{i+j-1} \sqrt{\kappa}^{j-i+1} \right)^{-1} =: M(x, q, \sqrt{\kappa}) \end{aligned} \quad (5.43)$$

which is just a generalization of the MacMahon function. Using repeatedly this identity we find

$$\begin{aligned} Z_{\mathbb{C}^2/\mathbb{Z}_k \times \mathbb{C}}^{ref}(\{p_i\}; \kappa) &= M \left(1, (-p_0 p_1 \cdots p_{n-1}), \sqrt{\kappa} \right)^n \times \\ &\times \prod_{0 < r \leq s < n} M(p_{[r,s]}, (-p_0 p_1 \cdots p_{n-1}), \sqrt{\kappa}) M\left(\frac{1}{p_{[r,s]}}, (-p_0 p_1 \cdots p_{n-1}), \sqrt{\kappa}\right) \end{aligned} \quad (5.44)$$

As in the conifold case, the results disagree with the literature [24] in the choice of refinement of the “degree zero” part, the factor $M \left(1, (-p_0 p_1 \cdots p_{n-1}), \sqrt{\kappa} \right)^n$. However as already mentioned in [24] and discussed above, this has to be expected and is due to the non-compactness of the relevant moduli spaces. Most likely there exist a choice of the refined limit which reproduces the results of [24], but we haven’t managed to find it so far.

5.4 The case $\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2$

Geometry and representation theory In this case the orbifold acts as


$$\begin{aligned} g_1(z_1, z_2, z_3) &= (-z_1, -z_2, z_3) \\ g_2(z_1, z_2, z_3) &= (-z_1, z_2, -z_3) \\ g_3(z_1, z_2, z_3) &= (z_1, -z_2, -z_3) \end{aligned} \quad (5.45)$$

and g_0 the identity. The crepant resolution given by $\text{Hilb}_{\mathbb{Z}_2 \times \mathbb{Z}_2}(\mathbb{C}^3, n)$ has three compact curves and no compact divisors.

The orbifold group $\mathbb{Z}_2 \times \mathbb{Z}_2$ acts on \mathbb{C}^3 and its non trivial weights are $r_1 = (1, 1, 0)$, $r_2 = (1, 0, 1)$ and $r_3 = r_1 + r_2$. There are four corresponding irreducible representations ρ_r and the tensor product decomposition $\rho_r \otimes Q$ with the defining representation gives

$$(a_{rs}^{(1)}) = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 \\ 1 & 1 & 1 & 0 \end{pmatrix}. \quad (5.46)$$

The resulting quiver Q is



The framed quiver \hat{Q} has an extra node connected by a single arrow to the node v_0 .

BPS partition function The representation space decomposes as

$$V_\pi = V_{00} + V_{01} + V_{10} + V_{00} \quad (5.48)$$

with

$$V_{r_1, r_2} = \sum_{(i, j, k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{i+k=r_1 \bmod 2} \delta_{j+k=r_2 \bmod 2} \quad (5.49)$$

As a consequence the virtual tangent space is

$$\begin{aligned} (T_\pi^{\text{vir}})^{\mathbb{Z}_2 \times \mathbb{Z}_2} = & V_{0,0} - V_{0,0}(t_1 t_2 t_3) - \left[(V_{0,0} \bar{V}_{0,0} + V_{1,0} \bar{V}_{1,0} + V_{0,1} \bar{V}_{0,1} + V_{1,1} \bar{V}_{1,1})(1 - t_1 t_2 t_3) \right. \\ & - (V_{1,0} \bar{V}_{0,0} + V_{0,0} \bar{V}_{1,0} + V_{0,1} \bar{V}_{1,1} + V_{1,1} \bar{V}_{0,1})(t_1 - t_2 t_3) \\ & - (V_{0,1} \bar{V}_{0,0} + V_{0,0} \bar{V}_{0,1} + V_{1,0} \bar{V}_{1,1} + V_{1,1} \bar{V}_{1,0})(t_2 - t_1 t_3) \\ & \left. - (V_{0,0} \bar{V}_{1,1} + V_{1,1} \bar{V}_{0,0} + V_{1,0} \bar{V}_{0,1} + V_{0,1} \bar{V}_{1,0})(t_3 - t_1 t_2) \right]. \end{aligned} \quad (5.50)$$

From the virtual tangent space and the classification of fixed points we can compute the partition function, defined as

$$\mathcal{Z}_{\mathbb{C}^3 / \mathbb{Z}_2 \times \mathbb{Z}_2}(\{p_i\}, \{t_i\}) = \sum_n \sum_{|\pi|=n} p_0^{|\pi_{00}|} p_1^{|\pi_{10}|} p_2^{|\pi_{01}|} p_3^{|\pi_{11}|} \hat{a} \left((T_\pi^{\text{vir}})^{\mathbb{Z}_2 \times \mathbb{Z}_2} \right). \quad (5.51)$$

Explicitly the first few terms read

$$\begin{aligned} Z_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}(\{p_i\}, \{t_i\}) &= 1 + p_0 + p_0 p_1 p_2 + p_0 p_1 p_3 + p_0 p_2 p_3 + \frac{p_0 p_1 (\kappa - t_1^2)}{\sqrt{\kappa} (-1 + t_1^2)} + \frac{p_0^2 p_1 (\kappa - t_1^2)}{\sqrt{k} (-1 + t_1^2)} \\ &+ \frac{p_0 p_2 (k - t_2^2)}{\sqrt{\kappa} (-1 + t_2^2)} + \frac{p_0^2 p_2 (\kappa - t_2^2)}{\sqrt{k} (-1 + t_2^2)} + \frac{p_0 p_3 (\kappa - t_3^2)}{\sqrt{\kappa} (-1 + t_3^2)} + \frac{p_0^2 p_3 (\kappa - t_3^2)}{\sqrt{\kappa} (-1 + t_3^2)} + \dots \end{aligned} \quad (5.52)$$

Symmetrized form Now we write down the orbifold function, whose symmetrization reproduces the partition function. Consider first the function

$$\begin{aligned} \mathcal{F}_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}(t_1, t_2, t_3, t_4, t_5) &= \mathcal{F}_{\mathbb{C}^3}\left(t_1^2, t_2^2, \frac{t_3}{t_1 t_2}, t_4, t_5\right) + \mathcal{F}_{\mathbb{C}^3}\left(t_1^2, \frac{t_2}{t_1 t_3}, t_3^2, t_4, t_5\right) \\ &+ \mathcal{F}_{\mathbb{C}^3}\left(\frac{t_1}{t_2 t_3}, t_2^2, t_3^2, t_4, t_5\right) + \mathcal{F}_{\mathbb{C}^3}\left(\frac{t_2 t_3}{t_1}, \frac{t_1 t_3}{t_2}, \frac{t_1 t_2}{t_3}, t_4, t_5\right) \end{aligned} \quad (5.53)$$

Such a function is related to the decomposition

$$\begin{aligned} \frac{1}{4} \sum_{i=0}^3 \frac{1}{(1 - g_i t_1)(1 - g_i t_2)(1 - g_i t_3)} &= \frac{1}{(1 - t_1^2)(1 - t_2^2)\left(1 - \frac{t_3}{t_1 t_2}\right)} + \frac{1}{(1 - t_1^2)(1 - t_3^2)\left(1 - \frac{t_2}{t_1 t_3}\right)} \\ &+ \frac{1}{(1 - t_2^2)(1 - t_3^2)\left(1 - \frac{t_1}{t_2 t_3}\right)} + \frac{1}{\left(1 - \frac{t_1 t_2}{t_3}\right)\left(1 - \frac{t_1 t_3}{t_2}\right)\left(1 - \frac{t_2 t_3}{t_1}\right)} \end{aligned} \quad (5.54)$$

which generated invariant monomials. Secondly define

$$\begin{aligned} \mathcal{F}_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}^{(r)}(\{p_i\}, t_1, t_2, t_3, t_4, t_5) &= \frac{1}{\left(\sqrt{t_4} - \frac{1}{\sqrt{t_4}}\right)\left(\sqrt{t_5} - \frac{1}{\sqrt{t_5}}\right)} \\ &\times \left[\frac{\left(\frac{\sqrt{k}}{t_3} - \frac{t_3}{\sqrt{k}}\right)\left(p_1 p_2 + \frac{1}{p_1 p_2}\right)}{t_3 - \frac{1}{t_3}} + \frac{\left(\frac{\sqrt{k}}{t_2} - \frac{t_2}{\sqrt{k}}\right)\left(p_1 p_3 + \frac{1}{p_1 p_3}\right)}{t_2 - \frac{1}{t_2}} + \frac{\left(\frac{\sqrt{k}}{t_1} - \frac{t_1}{\sqrt{k}}\right)\left(p_2 p_3 + \frac{1}{p_2 p_3}\right)}{t_1 - \frac{1}{t_1}} \right. \\ &\left. + p_1 p_2 p_3 + \frac{1}{p_1 p_2 p_3} + p_1 + \frac{1}{p_1} + p_2 + \frac{1}{p_2} + p_3 + \frac{1}{p_3} \right]. \end{aligned} \quad (5.55)$$

We claim that

$$Z_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}(\{p_i\}, \{t_i\}) = \mathbf{S}^\bullet \left(\mathcal{F}_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}(t_1, t_2, t_3, t_4, t_5) + \mathcal{F}_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}^{(r)}(\{p_i\}, t_1, t_2, t_3, t_4, t_5) \right) \quad (5.56)$$

with the usual identifications $\kappa = t_1 t_2 t_3$, $t_4 = q/\sqrt{\kappa}$, $t_5 = 1/(q\sqrt{\kappa})$, $q = -p_0 p_1 p_2 p_3$.

Refined limit By scaling away the toric weights with the same procedure of A-fibered singularities we find

$$Z_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}^{ref}(\{p_i\}, \kappa) = \mathbf{S}^\bullet \left[\frac{q \kappa^{1/2}}{(1 - \frac{q}{\kappa^{1/2}})(1 - \kappa^{1/2} q)} \left(3 + \frac{1}{\kappa} + \left(p_1 p_2 + \frac{1}{p_1 p_2} \right) + \left(p_2 p_3 + \frac{1}{p_2 p_3} \right) \right) \right. \\ \left. + \frac{1}{\kappa} \left(p_1 p_3 + \frac{1}{p_1 p_3} \right) - \frac{1}{\kappa^{1/2}} \left(p_1 + \frac{1}{p_1} \right) - \frac{1}{\kappa^{1/2}} \left(p_2 + \frac{1}{p_2} \right) - \frac{1}{\kappa^{1/2}} \left(p_3 + \frac{1}{p_3} \right) \right] \quad (5.57)$$

Proceeding as before, we can also write down a product form for the partition function

$$Z_{\mathbb{C}^3/\mathbb{Z}_2 \times \mathbb{Z}_2}^{ref}(\{p_i\}, \kappa) = M(1, q, \sqrt{\kappa})^3 M(1/\sqrt{\kappa}, q, \sqrt{\kappa}) M(p_1 p_2, q, \sqrt{\kappa}) M\left(\frac{1}{p_1 p_2}, q, \sqrt{\kappa}\right) \\ M(p_2 p_3, q, \sqrt{\kappa}) M\left(\frac{1}{p_2 p_3}, q, \sqrt{\kappa}\right) M\left(\frac{1}{\kappa} p_1 p_3, q, \sqrt{\kappa}\right) M\left(\frac{1}{\kappa} \frac{1}{p_1 p_3}, q, \sqrt{\kappa}\right) \\ M\left(\frac{1}{\kappa} p_1, q, \sqrt{\kappa}\right)^{-1} M\left(\frac{1}{\kappa} p_2, q, \sqrt{\kappa}\right)^{-1} M\left(\frac{1}{\kappa} p_3, q, \sqrt{\kappa}\right)^{-1} \\ M\left(\frac{1}{\kappa} \frac{1}{p_1}, q, \sqrt{\kappa}\right)^{-1} M\left(\frac{1}{\kappa} \frac{1}{p_2}, q, \sqrt{\kappa}\right)^{-1} M\left(\frac{1}{\kappa} \frac{1}{p_3}, q, \sqrt{\kappa}\right)^{-1}. \quad (5.58)$$

5.5 The $\mathbb{C}^3/\mathbb{Z}_3$ orbifold

Consider now the example the orbifold $\mathbb{C}^3/\mathbb{Z}_3$. In this case no closed formula is known even in the case of the (unrefined) topological string, and therefore we do not expect one in the K-theory setup.

Geometry and representation theory We let the generator g of \mathbb{Z}_3 act on \mathbb{C}^3 as $g(z_1, z_2, z_3) = (\zeta z_1, \zeta z_2, \zeta z_3)$ with $\zeta^3 = 1$. The resulting orbifold $\mathbb{C}^3/\mathbb{Z}_3$ has a geometric crepant resolution given by the total space of $\mathcal{O}_{\mathbb{P}^2}(-3) \rightarrow \mathbb{P}^2$. This geometry contains a \mathbb{P}^2 as a compact divisor, and three rational curves.

From the representation theory data one can construct the quiver

$$\begin{array}{ccc} & v_0 \bullet & \\ & \swarrow \quad \searrow & \\ v_1 \bullet & & \bullet v_2 \\ & \longleftarrow \quad \longrightarrow & \end{array} \quad (5.59)$$

Indeed in this case $Q = \rho_1 \oplus \rho_1 \oplus \rho_1$, from which we see from (5.1) that

$$a_{ab}^{(1)} = \begin{pmatrix} 0 & 0 & 3 \\ 3 & 0 & 0 \\ 0 & 3 & 0 \end{pmatrix} \quad \text{and} \quad a_{ab}^{(2)} = \begin{pmatrix} 0 & 3 & 0 \\ 0 & 0 & 3 \\ 3 & 0 & 0 \end{pmatrix} \quad (5.60)$$

The framing consists in an extra node connected with a single arrow to v_0 .

BPS partition function The representation space has the decomposition $V = V_0 + V_1 + V_2$, where

$$V_r = \sum_{(i,j,k) \in \pi} t_1^{i-1} t_2^{j-1} t_3^{k-1} \delta_{i+j+k=r \bmod 3} \quad (5.61)$$

The virtual tangent space is given by

$$\begin{aligned} (T_\pi^{\text{vir}})^\Gamma = & V_0 - \bar{V}_0(t_1 t_2 t_3) - \left[(V_0 \bar{V}_0 + V_1 \bar{V}_1 + V_2 \bar{V}_2)(1 - t_1 t_2 t_3) \right. \\ & + (V_0 \bar{V}_2 + V_1 \bar{V}_0 + V_2 \bar{V}_1)(-t_1 - t_2 - t_3) \\ & \left. + (V_0 \bar{V}_1 + V_1 \bar{V}_2 + V_2 \bar{V}_0)(t_1 t_2 + t_1 t_3 + t_2 t_3) \right]. \end{aligned} \quad (5.62)$$

The partition function is now

$$Z_{\mathbb{C}^3/\mathbb{Z}_3}(\{p_i\}, \{t_i\}) = \sum_n \sum_{|\pi|=n} p_0^{|\pi_0|} p_1^{|\pi_1|} p_2^{|\pi_2|} \hat{a}((T_\pi^{\text{vir}})^\Gamma). \quad (5.63)$$

Explicitly the few first terms are

$$\begin{aligned} Z_{\mathbb{C}^3/\mathbb{Z}_3}(\{p_i\}, \{t_i\}) = & 1 + p_0 + \frac{p_0 p_1 (\kappa t_1 - t_2)(\kappa t_1 - t_3)}{\kappa(t_1 - t_2)(t_1 - t_3)} - \frac{p_0 p_1 p_2 (-\kappa + t_1^3)(\kappa t_1 - t_2)(\kappa t_1 - t_3)}{\kappa^{3/2}(-1 + t_1^3)(t_1 - t_2)(t_1 - t_3)} \\ & + \frac{p_0 p_1 (t_1 - \kappa t_2)(\kappa t_2 - t_3)}{\kappa(t_1 - t_2)(t_2 - t_3)} - \frac{p_0 p_1 p_2 (t_1 - \kappa t_2)(-\kappa + t_2^3)(\kappa t_2 - t_3)}{\kappa^{3/2}(t_1 - t_2)(-1 + t_2^3)(t_2 - t_3)} \\ & + \frac{p_0 p_1^2 (\kappa t_1 - t_3)(-\kappa t_2 + t_3)}{\kappa(t_1 - t_3)(-t_2 + t_3)} + \frac{p_0 p_1^2 (t_1 - \kappa t_2)(t_1 - \kappa t_3)}{\kappa(t_1 - t_2)(t_1 - t_3)} + \frac{p_0 p_1^2 (\kappa t_1 - t_2)(t_2 - \kappa t_3)}{\kappa(t_1 - t_2)(t_2 - t_3)} \\ & + \frac{p_0 p_1 (t_1 - \kappa t_3)(-t_2 + \kappa t_3)}{\kappa(t_1 - t_3)(-t_2 + t_3)} - \frac{p_0 p_1 p_2 (t_1 - \kappa t_3)(-t_2 + \kappa t_3)(-\kappa + t_3^3)}{\kappa^{3/2}(t_1 - t_3)(-t_2 + t_3)(-1 + t_3^3)} + \dots \end{aligned} \quad (5.64)$$

6 Discussion

We conclude with some comments and some open problems.

6.1 Towards M2 branes on NCCRs

The results of the previous sections should admit an interpretation as the theory of M2 branes propagating on noncommutative crepant resolutions. Indeed the main goal of the M2-brane index computations is to provide some mathematical understanding of certain degrees of freedom of M-theory. While it is not yet clear how precisely to attain this goal, we can still try to make some progress in understanding the noncommutative theory by transferring large radius results to the singular non-geometric limit. The following discussion will be necessarily somewhat vague.

Roughly speaking the proposal made in [30] for the membrane moduli space consists of the moduli space of maps $f : C \rightarrow Z$ from 1-dimensional Cohen-Macaulay schemes into the smooth Calabi-Yau fivefold, with certain stability conditions that ensure that the Euler characteristic $\chi(\mathcal{O}_C)$ is bounded both from above and from below. More precisely the moduli space $\text{M2}(Z)$ parametrizes such maps together with a certain subsheaf N of the normal sheaf N_f . Such moduli space is endowed with a

symmetrized virtual structure sheaf \mathcal{O}_{M2} . Conjecturally virtual counts of over this moduli space should reproduce the K-theoretical Donaldson-Thomas theory of Z .

Our results should point the way to define the analog object in the noncommutative regime $\text{NCM2}(Z)$, which describe membranes degrees of freedom when Z develops a singularity. This means that it should be possible to interpret our results as a theory of membranes in a noncommutative geometry, possibly obtained as a phase of a membrane worldvolume theory. One could expect that such a theory is related to maps from a curve C into the noncommutative algebra \mathbf{A} which resolves the singularity, where the membrane is wrapping the three manifold $C \times S^1$. The situation is somewhat analog to what happens in the study of three dimensional gauge theories, as discussed for example in [32]. If one considers a supersymmetric gauge theory on $C \times S^1$ then under certain assumptions the Higgs branch has the structure of a Nakajima variety. At low energies the theory reduces to the K-theory of quasimaps $f : C \dashrightarrow \text{Nak}$, defined as section of certain bundles \mathcal{V}, \mathcal{W} on C which obey certain moment map relations $\mu = 0$. Here $\mathcal{V}_i = f^*V_i$ where V_i are the vector spaces which enter in the definition of the Nakajima quiver variety. In our case the relevant target space is not a Nakajima variety but we can still consider the moduli space of representations of the relevant quiver. Indeed this plays the same role in six-dimensional instanton counting that Nakajima varieties play in four dimensional instanton counting [9].

Roughly speaking one would study collections of vector bundles \mathcal{V}_i on C of fixed rank, together with a section $f \in H^0(C, \mathcal{M}^\Gamma)$ which obeys $\mu = 0$, with

$$\mathcal{M}^\Gamma = \text{Hom}_\Gamma(\mathcal{V}, \mathcal{Q} \otimes \mathcal{V}) \oplus \text{Hom}_\Gamma(\mathcal{V}, \wedge^3 \mathcal{Q} \otimes \mathcal{V}) \oplus \text{Hom}_\Gamma(\mathcal{W}, \mathcal{V}) \quad (6.1)$$

where the product of toric weights is interpreted as \hbar . The moment map $\mu = 0$ is the one obtained by the F-term equations derived from the superpotential [7].

The associated enumerative problem appears to be interesting and somewhat natural. However there are two serious drawbacks: to begin with it is not clear at all if it is mathematically well posed; secondly the physical motivations are based on an analogy, but this is rather thin and there is no reason that the worldvolume theory of the membrane will have a phase where the moduli space of vacua is described by quasimaps.

Another option would be to transfer directly the construction of [30] from large radius to the singular phase. In the case of orbifolds we can use the McKay correspondence, or more generically the fact that the derived category $\text{D}^b(\text{coh}X)$ of the smooth threefold is derived equivalent to the derived category of modules over the path algebra of the quiver $\text{D}^b(\mathbf{A} - \text{mod})$. This equivalence is given by a tilting sheaf and under favourable conditions it descends at the level of K-theory. For example for the resolved conifold Y the tilting generator is $T = \mathcal{O}_Y \oplus \pi^* \mathcal{O}_{P^1}(1)$. Then $\mathbb{R}\text{Hom}(T, -) : \text{D}^b(\text{coh}(Y)) \rightarrow \text{D}^b(\mathbf{A} - \text{mod})$ induces the derived equivalence, and $\mathbf{A} = \text{End}_Y(T)$. The tilting generator has the form $T = P_\circ \oplus P_\bullet$ where $P_v = \mathbf{A} e_i$ are the projective modules and the e_i are the idempotents.

The derived equivalence descends at the level of K-theory to the isomorphism $[\text{RF}(-)] : \text{K}(\text{coh}Y) \rightarrow \text{K}(\mathbf{A} - \text{mod})$ (since this discussion is rather qualitative we ignore when opposite categories appear), given by $[\text{RF}(-)] = \sum_i (-1)^i \text{Ext}_Y^i(T, -)$. Quasi-isomorphic complexes map to the same K-theory class. In the large radius limit we are looking at maps f from schemes C to Z . We can replace the

curve with the sheaf $f_*\mathcal{O}_C$, and then via the derived equivalence map the two sheaves $f_*\mathcal{O}_C$ and \mathcal{N} into complexes of \mathbf{A} -modules, with K-theory classes $[\mathrm{RF}(\mathcal{O}_C)]$ and $[\mathrm{RF}(\mathcal{N})]$.

In the case of orbifolds the above procedure is essentially a generalization of the McKay correspondence, reviewed in [9] in a similar context. At the level of K-theory this maps complex of vector bundles which are exact outside the exceptional locus to Γ -equivariant sheaves supported at the origin of \mathbb{C}^3 . Therefore this discussion predicts that the moduli space $\mathrm{NCM2}$ of M2 branes on the noncommutative geometry can be understood as the Γ -fixed locus $\mathrm{M2}_\Gamma(\mathbb{C}^5)$ of the flat space moduli space $\mathrm{M2}(\mathbb{C}^5)$. Note that while the discussion above is rather vague, this proposal is concrete. In particular it should be possible to analyse this moduli space concretely to check the above quasi-map proposal.

We hope to return to this problem in the near future

6.2 Some open problems

We summarize a few more concrete open problems that are currently under investigation

- Perhaps the most interesting problem is a systematic study of the wall-crossing behaviour of the M2 index. In this paper we have computed the index in the noncommutative crepant resolution chamber. We believe our techniques can be extended to other chambers, especially those which arise from this chamber via quiver mutations. Work is in progress in this direction and we hope to report about it soon [11].
- Ordinary Donaldson-Thomas theory associated to quivers can also be rephrased in terms of a dimer model, a certain statistical mechanical model associated with the quiver [33]. It should be possible to generalize this model to the case where the weight of a dimer configuration is given by the Okounkov function of the configuration, computed at a fixed point in the associated quiver moduli space. It would be interesting to understand if such a model has hidden algebraic structures, perhaps related to those uncovered in [21].
- Donaldson-Thomas theory, from a gauge theory point of view, can be generalized to include defects [10]. Roughly speaking the idea is to specify singularities for the gauge connection over a divisor. Such singularities can be classified in term of parabolic sheaves and the Donaldson-Thomas problem is reduced to counting ideal sheaves with a prescribed parabolic structure. Also this problem can be rephrased in terms of representations of a certain quiver, and our techniques certainly apply to that case.
- In case of orbifolds the higher rank Donaldson-Thomas invariants admit a generalization as Donaldson-Thomas invariants of type \mathbf{w} , where \mathbf{w} is the vector which specifies the rank of the D6 brane theory as well as the boundary conditions at infinity [7]. It would be interesting to study the K-theoretic generalization to this setting. In particular it is natural to propose that the same vector \mathbf{w} specifies boundary conditions at infinity for M2 branes.

Acknowledgements

I am grateful to Vasily Pestun for many discussions and collaboration in the early stages of this work. I am a member of INDAM-GNFM and I am supported by INFN via the Iniziativa Specifica GAST. The author also acknowledges the support of IHÉS during a visit. The research of M.C. on this

project has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (QUASIFT grant agreement 677368).

References

- [1] K. Behrend and B. Fantechi, “Symmetric obstruction theories and Hilbert schemes of points on threefolds,” *Algebra Number Theory* **2** (2008) 313-345 [arXiv:math.AG/0512556].
- [2] F. Benini, G. Bonelli, M. Poggi and A. Tanzini, “Elliptic non-Abelian Donaldson-Thomas invariants of \mathbb{C}^3 ,” *JHEP* **07** (2019), 068 [arXiv:1807.08482 [hep-th]].
- [3] G. Bonelli, N. Fasola, A. Tanzini and Y. Zenkevich, “ADHM in 8d, coloured solid partitions and Donaldson-Thomas invariants on orbifolds,” [arXiv:2011.02366 [hep-th]].
- [4] M. Cirafici, “On Framed Quivers, BPS Invariants and Defects,” *Confluentes Mathematici* **9** (2017) 2, 71-99 [arXiv:1801.03778 [hep-th]].
- [5] M. Cirafici, “Quantum Line Defects and Refined BPS Spectra,” *Lett. Math. Phys.* **110** (2019) no.3, 501-531 [arXiv:1902.08586 [hep-th]].
- [6] M. Cirafici, A. Sinkovics and R. J. Szabo, “Cohomological gauge theory, quiver matrix models and Donaldson-Thomas theory,” *Nucl. Phys. B* **809** (2009) 452 [arXiv:0803.4188 [hep-th]].
- [7] M. Cirafici, A. Sinkovics and R. J. Szabo, “Instantons, Quivers and Noncommutative Donaldson-Thomas Theory,” *Nucl. Phys. B* **853** (2011) 508 [arXiv:1012.2725 [hep-th]].
- [8] M. Cirafici, A. Sinkovics and R. J. Szabo, “Instanton counting and wall-crossing for orbifold quivers,” *Annales Henri Poincare* **14** (2013) 1001 [arXiv:1108.3922 [hep-th]].
- [9] M. Cirafici and R. J. Szabo, “Curve counting, instantons and McKay correspondences,” *J. Geom. Phys.* **72** (2013), 54-109 [arXiv:1209.1486 [hep-th]].
- [10] M. Cirafici, “Defects in cohomological gauge theory and Donaldson-Thomas invariants,” *Adv. Theor. Math. Phys.* **20** (2016), 945-1006 [arXiv:1302.7297 [hep-th]].
- [11] M. Cirafici. “Wall-crossing and membranes” *to appear*.
- [12] M. Del Zotto, N. Nekrasov, N. Piazzalunga and M. Zabzine, “Playing with the index of M-theory,” [arXiv:2103.10271 [hep-th]].
- [13] P. Descombes, “Motivic DT invariants from localization,” [arXiv:2106.02518 [math.AG]].
- [14] T. Dimofte and S. Gukov, “Refined, Motivic, and Quantum,” *Lett. Math. Phys.* **91** (2010) 1 [arXiv:0904.1420 [hep-th]].
- [15] S. K. Donaldson and R. P. Thomas, “Gauge theory in higher dimensions”, in *The Geometric Universe; Science, Geometry, And The Work Of Roger Penrose*, Oxford University Press, 1998
- [16] N. Fasola, S. Monavari and A. T. Ricolfi, “Higher rank K-theoretic Donaldson-Thomas theory of points,” *Forum Math. Sigma* **9** (2021), e15 [arXiv:2003.13565 [math.AG]].
- [17] A. Iqbal, N. Nekrasov, A. Okounkov and C. Vafa, *JHEP* **04** (2008), 011 [arXiv:hep-th/0312022 [hep-th]].
- [18] D. L. Jafferis and G. W. Moore, “Wall crossing in local Calabi Yau manifolds,” [arXiv:0810.4909 [hep-th]].

- [19] Y. Kononov, A. Okounkov and A. Osinenko, “The 2-Leg Vertex in K-theoretic DT Theory,” *Commun. Math. Phys.* **382** (2021) no.3, 1579-1599 [arXiv:1905.01523 [math-ph]].
- [20] M. Kontsevich and Y. Soibelman, “Stability structures, motivic Donaldson-Thomas invariants and cluster transformations,” arXiv:0811.2435 [math.AG].
- [21] W. Li and M. Yamazaki, “Quiver Yangian from Crystal Melting,” *JHEP* **11** (2020), 035 [arXiv:2003.08909 [hep-th]].
- [22] D. Maulik, N. Nekrasov, A. Okounkov and R. Pandharipande, *Compos. Math.* **142** (2006) no.05, 1263-1285 doi:10.1112/S0010437X06002302 [arXiv:math/0312059 [math.AG]].
- [23] A. Morrison, S. Mozgovoy, K. Nagao and B. Szendroi, “Motivic Donaldson-Thomas invariants of the conifold and the refined topological vertex,” arXiv:1107.5017 [math.AG].
- [24] A. Morrison and K. Nagao “Motivic Donaldson-Thomas invariants of toric small crepant resolutions,” *Algebra Number Theory* **9** (2015) 767-813 arXiv:1110.5976 [math.AG]
- [25] S. Mozgovoy and B. Pioline, “Attractor invariants, brane tilings and crystals,” [arXiv:2012.14358 [hep-th]].
- [26] S. Mozgovoy and M. Reineke, “On the noncommutative Donaldson-Thomas invariants arising from brane tilings,” *Adv. Math.* **223** (2010) 1521 [arXiv:0809.0117 [math.AG]].
- [27] K. Nagao and H. Nakajima, “Counting invariant of perverse coherent sheaves and its wall-crossing,” [arXiv:0809.2992 [math.AG]].
- [28] N. Nekrasov, “A la recherche de la M-theorie perdue Z theory: Chasing M / f theory,” [arXiv:hep-th/0412021 [hep-th]].
- [29] N. Nekrasov, “Magnificent four,” *Adv. Theor. Math. Phys.* **24** (2020) no.5, 1171-1202 [arXiv:1712.08128 [hep-th]].
- [30] N. Nekrasov and A. Okounkov, “Membranes and Sheaves,” arXiv:1404.2323 [math.AG].
- [31] N. Nekrasov and N. Piazzalunga, “Magnificent Four with Colors,” *Commun. Math. Phys.* **372** (2019) no.2, 573-597 [arXiv:1808.05206 [hep-th]].
- [32] A. Okounkov, “Lectures on K-theoretic computations in enumerative geometry,” arXiv:1512.07363 [math.AG].
- [33] H. Ooguri and M. Yamazaki, “Crystal Melting and Toric Calabi-Yau Manifolds,” *Commun. Math. Phys.* **292** (2009), 179-199 [arXiv:0811.2801 [hep-th]].
- [34] E. Pomoni, W. Yan and X. Zhang, [arXiv:2106.11611 [hep-th]].
- [35] B. Szendroi, “Non-commutative Donaldson-Thomas theory and the conifold,” *Geom. Topol.* **12** (2008) 1171 [arXiv:0705.3419 [math.AG]].
- [36] M. Van den Bergh “Non-commutative crepant resolutions”, *The legacy of Niels Henrik Abel*, 749-770, Springer, Berlin, 2004.