

OPTIMAL WEAK ESTIMATES FOR RIESZ POTENTIALS

LIANG HUANG AND HANLI TANG

ABSTRACT. In this note we prove a sharp reverse weak estimate for Riesz potentials

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \gamma_s v_n^{\frac{n-s}{n}} \|f\|_{L^1} \text{ for } 0 < f \in L^1(\mathbb{R}^n),$$

where $\gamma_s = 2^{-s} \pi^{-\frac{n}{2}} \frac{\Gamma(\frac{n-s}{2})}{\Gamma(\frac{n}{2})}$. We also consider the behavior of the best constant $\mathcal{C}_{n,s}$ of weak type estimate for Riesz potentials, and we prove $\mathcal{C}_{n,s} = O(\frac{\gamma_s}{s})$ as $s \rightarrow 0$.

1. INTRODUCTION

The Riesz potentials (fractional integral operators) I_s , which play an important part in Analysis, are defined by

$$I_s(f)(x) = \gamma_s \int_{\mathbb{R}^n} \frac{f(x-y)}{|y|^{n-s}} dy,$$

where $0 < s < n$ and $\gamma_s = 2^{-s} \pi^{-\frac{n}{2}} \frac{\Gamma(\frac{n-s}{2})}{\Gamma(\frac{n}{2})}$. Such operators were first systematically investigated by M. Riesz [11]. The (L^p, L^q) -boundedness of Riesz potentials were proved by G. Hardy and J. Littlewood [6] when $n = 1$ and by S. Sobolev [12] when $n > 1$. The $(L^1, L^{\frac{n}{n-s},\infty})$ -boundedness were obtained by A. Zygmund [16]. More precisely, they established the following theorem.

Theorem A. *Let $0 < s < n$ and let p, q satisfy $1 \leq p < q < \infty$ and $\frac{1}{p} - \frac{1}{q} = \frac{s}{n}$, then when $p > 1$,*

$$\|I_s(f)\|_{L^q(\mathbb{R}^n)} \leq C(n, p, s) \|f\|_{L^p(\mathbb{R}^n)}.$$

And when $p = 1$,

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} = \sup_{\lambda > 0} \lambda |\{x \in \mathbb{R}^n : |I_s f| > \lambda\}|^{\frac{n-s}{n}} \leq C(n, s) \|f\|_{L^1(\mathbb{R}^n)}.$$

The best constant in the (L^p, L^q) inequality when $p = \frac{2n}{n+s}$, $q = \frac{2n}{n-s}$ was precisely calculated by E. Lieb [8] (see also [4]), and E. Lieb and M. Loss also offered an upper bound of $C(n, p, s)$ (see chapter 4 in [9]).

Although the best constant of (L^p, L^q) estimate for Riesz potentials has been studied for decades, to the best of the authors' knowledge there is no result about the best constant of

2020 *Mathematics Subject Classification.* 42B20.

Key words and phrases. Riesz potentials, sharp constant, optimal estimate .

The first author is supported by National Key Research and Development Program of China (Grant No. 2020YFA0712900) and National Natural Science Foundation of China (Grant No. 11701032)

Corresponding author: Hanli Tang at hltang@bnu.edu.cn.

$(L^1, L^{\frac{n}{n-s}, \infty})$ estimate for Riesz potentials. In this paper, we will provide some estimates for the best constant of the weak type inequality.

In [14](see multilinear case in [15]), the second author setted up the following limiting weak-type behavior for Riesz potentials,

$$\lim_{\lambda \rightarrow 0} \lambda |\{x \in \mathbb{R}^n : |I_s f| > \lambda\}|^{\frac{n-s}{n}} = \gamma_s v_n^{\frac{n-s}{n}} \|f\|_{L^1(\mathbb{R}^n)} \text{ for } 0 < f \in L^1(\mathbb{R}^n),$$

which implies a reverse weak estimate

$$\|I_s(f)\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)} \geq \gamma_s v_n^{\frac{n-s}{n}} \|f\|_{L^1(\mathbb{R}^n)} \text{ for } 0 < f \in L^1(\mathbb{R}^n), \quad (1.1)$$

where v_n is the volume of the unit ball in \mathbb{R}^n . So a natural question that arises here is whether the constant $\gamma_s v_n^{\frac{n-s}{n}}$ is sharp? In the paper, we will give an affirmative answer.

Let $\mathcal{C}_{n,s}$ be the best constant such that the $(L^1, L^{\frac{n}{n-s}, \infty})$ estimate holds for Riesz potentials, i.e.

$$\mathcal{C}_{n,s} = \sup_{f \in L^1(\mathbb{R}^n)} \frac{\|I_s(f)\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)}}{\|f\|_{L^1(\mathbb{R}^n)}}.$$

Then from (1.1), one can directly obtain a lower bound for $\mathcal{C}_{n,s}$,

$$\mathcal{C}_{n,s} \geq \gamma_s v_n^{\frac{n-s}{n}}.$$

Our another goal in this paper is to provide upper and lower bounds of $\mathcal{C}_{n,s}$ to study the behavior of $\mathcal{C}_{n,s}$ as $s \rightarrow 0$. Our approach depends on the weak $L^{\frac{n}{n-s}}$ norm $\|\cdot\|_{L^{\frac{n}{n-s}, \infty}}$ which is defined by

$$\|\cdot\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)} = \sup_{0 < |E| < \infty} |E|^{-\frac{1}{r} + \frac{n-s}{n}} \left(\int_E |f|^r dx \right)^{\frac{1}{r}}, \quad 0 < r < \frac{n}{n-s}.$$

The norm $\|\cdot\|_{L^{\frac{n}{n-s}, \infty}}$ is equivalent to $\|\cdot\|_{L^{\frac{n}{n-s}, \infty}}$. In fact there holds (see Exercise 1.1.12 in [5])

$$\|f\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)} \leq \|\cdot\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)} \leq \left(\frac{n}{n-r(n-s)} \right)^{\frac{1}{r}} \|f\|_{L^{\frac{n}{n-s}, \infty}(\mathbb{R}^n)}. \quad (1.2)$$

Closely related to the Riesz potentials is the centered fractional maximal function, which is defined by

$$M_s f(x) = \sup_{r>0} \frac{1}{|B(x, r)|^{1-\frac{s}{n}}} \int_{B(x, r)} |f(y)| dy, \quad 0 < s < n.$$

M_s satisfy the same (L^p, L^q) and $(L^1, L^{\frac{n}{n-s}, \infty})$ inequality as I_s does, see [1] and [10]. For any positive function f it is easy to see $M_s(f) \leq 1/\gamma(s) v_n^{\frac{s-n}{n}} I_s(f)$. Although the reverse inequality dose not hold in general, B.Muckenhoupt and R.Wheeden [10] proved the two quantities are comparable in L^p norm ($1 < p < \infty$) when f is nonnegative.

Now let us state our main results. First of all we consider the weak estimate of $I_s(f)$ and $M_s(f)$ under the norm $\|\cdot\|_{L^{\frac{n}{n-s}, \infty}}$. Surprisingly identities for the weak type estimate of Riesz potentials and fractional maximal function can be established, which implies the two quantities are comparable in $L^{\frac{n}{n-s}, \infty}$ (quasi)norm when $f \in L^1(\mathbb{R}^n)$ is nonnegative.

Theorem 1.1. *Let $0 < s < n$ and $f \in L^1(\mathbb{R}^n)$. When $1 \leq r < \frac{n}{n-s}$,*

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} \leq \gamma_s v_n^{\frac{n-s}{n}} \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1(\mathbb{R}^n)},$$

and

$$\|M_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} = \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1(\mathbb{R}^n)}.$$

Moreover if $0 < f \in L^1(\mathbb{R}^n)$, then

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} = \gamma_s v_n^{\frac{n-s}{n}} \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1(\mathbb{R}^n)}.$$

Remark 1. *In fact, from the proof one can obtain the reverse weak estimate holds when $0 < r < \frac{n}{n-s}$. More precisely when $0 < r < \frac{n}{n-s}$,*

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} \geq \gamma_s v_n^{\frac{n-s}{n}} \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1(\mathbb{R}^n)}, \text{ if } 0 < f \in L^1(\mathbb{R}^n),$$

and

$$\|M_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} \geq \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1(\mathbb{R}^n)}, \text{ if } f \in L^1(\mathbb{R}^n).$$

Then we prove the following sharp reverse weak estimates for Riesz potentials.

Theorem 1.2. *Let $0 < f \in L^1(\mathbb{R}^n)$, then*

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}(\mathbb{R}^n)} \geq \gamma_s v_n^{\frac{n-s}{n}} \|f\|_{L^1(\mathbb{R}^n)}.$$

And the equality holds when $f = (\frac{a}{b+|x-x_0|^2})^{\frac{n+s}{2}}$, where $a, b > 0$ and $x_0 \in \mathbb{R}^n$.

As a corollary of Theorem 1.1 and Theorem 1.2, we can obtain the following sharp reverse inequality.

Corollary 1.1. *Let $f \in L^1(\mathbb{R}^n)$, then*

$$\|M_s f\|_{L^{\frac{n}{n-s},\infty}} \geq \|f\|_{L^1}.$$

And the equality holds when $f(x) = h(|x - x_0|)$ where h is a radial decreasing function.

At last we offer an upper and a lower bound for $\mathcal{C}_{n,s}$, which implies the behavior of the best constant $\mathcal{C}_{n,s}$ for small s is optimal, i.e. $\mathcal{C}_{n,s} = O(\frac{2s}{s}) = O(1)$ as $s \rightarrow 0$.

Theorem 1.3. *When $n > 2$ and $0 < s < \frac{n-2}{4}$,*

$$\gamma_s v_n^{\frac{n-s}{n}} \frac{n-2-4s}{2s(n-2-s)} \leq \mathcal{C}_{n,s} \leq \gamma_s v_n^{\frac{n-s}{n}} \frac{n}{s}.$$

Remark 2. *Besides using the rearrangement inequality to obtain an upper bound $\gamma_s v_n^{\frac{n-s}{n}} \frac{n}{s}$, we can take the heat-diffusion semi-group as a tool (see the Appendix), which was used by E. Stein and J. Strömberg in [13] to study the $(L^1, L^{1,\infty})$ bound for centered maximal function, to obtain another upper bound which is equal to $O(\gamma_s v_n^{\frac{n-s}{n}} \frac{n}{s}) = O(1)$ as $(s, n) \rightarrow (0, \infty)$.*

2. THE IDENTITY FOR $I_s(f)$ AND $M_s(f)$ IN $\|\cdot\|_{L^{\frac{n}{n-s},\infty}}$

In this section, we will prove Theorem 1.1. Without loss of generality let us assume $\|f\|_{L^1(\mathbb{R}^n)} = 1$. Since $I_s(f) \leq I_s(|f|)$ and $r \geq 1$, using Minkowski inequality one have for any measurable set E with $|E| < \infty$,

$$|E|^{-\frac{1}{r} + \frac{n-s}{n}} \left[\int_E |I_s f(x)|^r dx \right]^{\frac{1}{r}} \leq \gamma_s |E|^{-\frac{1}{r} + \frac{n-s}{n}} \int_{\mathbb{R}^n} \left[\int_E \frac{dx}{|x-y|^{(n-s)r}} \right]^{\frac{1}{r}} |f(y)| dy. \quad (2.1)$$

Then by Hardy Littlewood rearrangement inequality, there holds

$$\int_E \frac{dx}{|x-y|^{(n-s)r}} \leq \int_{E^*} \frac{dx}{|x|^{(n-s)r}} = v_n^{\frac{n-s}{n}} \frac{n}{n-(n-s)r} |E|^{1-\frac{n-s}{n}r}, \quad (2.2)$$

where E^* is the symmetric rearrangement of the set E , i.e. E^* is an open ball centered at the origin whose volume is $|E|$. Therefore by (2.1) and (2.2) one can obtain

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} \leq \gamma_s v_n^{\frac{n-s}{n}} \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1}.$$

Next, let us prove when $0 \leq f \in L^1(\mathbb{R}^n)$ and $0 < r < \frac{n}{n-s}$,

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \gamma_s v_n^{\frac{n-s}{n}} \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \|f\|_{L^1}. \quad (2.3)$$

For any $\epsilon > 0$, choose R large enough such that $\int_{B_R(0)} f(y) dy = 1 - \epsilon$. Let $E = B_{lR}(0)$. Since

$$\int_{B_R(0)} \frac{f(y)}{|x-y|^{n-s}} dy \geq \int_{B_R(0)} \frac{f(y)}{(|x|+R)^{n-s}} dy = (1-\epsilon)(|x|+R)^{s-n},$$

then

$$\begin{aligned} \|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} &\geq \gamma_s |E|^{-\frac{1}{r} + \frac{n-s}{n}} \left[\int_E \left(\int_{B_R(0)} \frac{f(y)}{|x-y|^{n-s}} dy \right)^r dx \right]^{\frac{1}{r}} \\ &\geq \gamma_s |E|^{-\frac{1}{r} + \frac{n-s}{n}} (1-\epsilon) \left[\int_E \frac{dx}{(|x|+R)^{(n-s)r}} \right]^{\frac{1}{r}} \\ &= \gamma_s v_n^{\frac{n-s}{n}} n^{\frac{1}{r}} (1-\epsilon) l^{-\frac{n}{r} + n-s} \left[\int_0^l \frac{t^{n-1}}{(t+1)^{(n-s)r}} dt \right]^{\frac{1}{r}}. \end{aligned}$$

By the fact that this inequality holds for any $l > 0$, then letting $l \rightarrow \infty$, one obtain

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \gamma_s v_n^{\frac{n-s}{n}} (1-\epsilon) \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}},$$

which implies (2.3). And we finish the proof of the identity for Riesz potential.

For fractional maximum function M_s , since

$$\begin{aligned} M_s(f)(x) &\geq \frac{1}{v_n^{\frac{n-s}{n}}(|x| + R)^{n-s}} \int_{|y-x| \leq R+|x|} |f(y)| dy \\ &\geq \frac{1}{v_n^{\frac{n-s}{n}}(|x| + R)^{n-s}} \int_{|y| \leq R} |f(y)| dy = \frac{1-\epsilon}{v_n^{\frac{n-s}{n}}(|x| + R)^{n-s}}, \end{aligned}$$

then one can use the same method to get

$$\|M_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}} \text{ when } 0 < r < \frac{n}{n-s}.$$

On the other hand,

$$\|M_s(f)\|_{L^{\frac{n}{n-s},\infty}} \leq \|1/\gamma(s)v_n^{\frac{s-n}{n}} I_s(|f|)\|_{L^{\frac{n}{n-s},\infty}} = \left(\frac{n}{n-(n-s)r} \right)^{\frac{1}{r}}.$$

Thus one can obtain the desired identity for M_s .

3. THE SHARP REVERSE WEAK ESTIMATE FOR I_s AND M_s

In this section, first we prove the sharp reverse weak estimate for Riesz potentials I_s . By (1.2) and Theorem 1.1, there holds

$$\|I_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \gamma_s v_n^{\frac{n-s}{n}} \|f\|_{L^1}, \quad 0 < f \in L^1(\mathbb{R}^n).$$

Next, we will prove that the equality can be attained by the function $g(x) = (\frac{a}{b+|x-x_0|^2})^{\frac{n+s}{2}}$, where $a, b > 0$ and $x_0 \in \mathbb{R}^n$. Since the translation and dilation of g do not change the ratio $\|I_s(g)\|_{L^{\frac{n}{n-s},\infty}}/\|g\|_{L^1}$, we only need to consider $g(x) = (\frac{2}{1+|x|^2})^{\frac{n+s}{2}}$. In our calculus we will use the stereographic projection, so we will introduce some notations about the stereographic projection here.

The inverse stereographic projection $\mathcal{S} : \mathbb{R}^n \rightarrow \mathbb{S}^n \setminus \{S\}$, where $S = -e_{n+1}$ denotes the southpole, is given by

$$(\mathcal{S}(x))_i = \frac{2x_i}{1+|x|^2}, \quad i = 1, \dots, n, \quad (\mathcal{S}(x))_{n+1} = \frac{1-|x|^2}{1+|x|^2}.$$

Correspondingly, the stereographic projection is given by $\mathcal{S}^{-1} : \mathbb{S}^n \setminus \{S\} \rightarrow \mathbb{R}^n$,

$$(\mathcal{S}^{-1}(\xi))_i = \frac{\xi_i}{1+\xi_{n+1}}, \quad i = 1, \dots, n.$$

And the Jacobian of the (inverse) stereographic projection are

$$\mathcal{J}_{\mathcal{S}}(x) = \left(\frac{2}{1+|x|^2} \right)^n \text{ and } \mathcal{J}_{\mathcal{S}^{-1}}(\xi) = (1+\xi_{n+1})^{-n}.$$

By changing of variables,

$$\begin{aligned}
\|g\|_{L^1} &= \int_{\mathbb{R}^n} \left(\frac{2}{1+|x|^2} \right)^{\frac{n+s}{2}} dx = \int_{\mathbb{S}^n} \left(\frac{2}{1+|\mathcal{S}^{-1}(\xi)|^2} \right)^{\frac{s-n}{2}} d\xi \\
&= \int_{\mathbb{S}^n} (1+\xi_{n+1})^{\frac{s-n}{2}} d\xi = |\mathbb{S}^{n-1}| \int_{-1}^1 (1+t)^{\frac{s-2}{2}} (1-t)^{\frac{n-2}{2}} dt \\
&= \pi^{n/2} 2^{\frac{s+n}{2}} \frac{\Gamma(s/2)}{\Gamma(\frac{s+n}{2})}.
\end{aligned} \tag{3.1}$$

Denote

$$c_{n,s} = \pi^{n/2} 2^{\frac{s+n}{2}} \frac{\Gamma(s/2)}{\Gamma(\frac{s+n}{2})}.$$

Since

$$|\mathcal{S}^{-1}(\xi) - \mathcal{S}^{-1}(\eta)|^2 = \mathcal{J}_{\mathcal{S}^{-1}}(\xi)^{\frac{1}{n}} |\xi - \eta|^2 \mathcal{J}_{\mathcal{S}^{-1}}(\eta)^{\frac{1}{n}}, \text{ for any } \xi, \eta \in \mathbb{S}^n,$$

and

$$\int_{\mathbb{S}^n} \frac{d\eta}{|\xi - \eta|^{n-s}} = \frac{2^s \pi^{n/2} \Gamma(s/2)}{\Gamma(\frac{n+s}{2})} = \frac{c_{n,s}}{2^{\frac{n-s}{2}}} \text{ for any } \eta \in \mathbb{S}^n \text{ (see D.4 in [5])},$$

one can obtain

$$\begin{aligned}
I_s(g)(x) &= \gamma(s) \int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-s}} \left(\frac{2}{1+|y|^2} \right)^{\frac{n+s}{2}} dy \\
&= \gamma(s) \int_{\mathbb{S}^n} \frac{1}{|\mathcal{S}^{-1}(\xi) - \mathcal{S}^{-1}(\eta)|^{n-s}} \left(\frac{2}{1+|\mathcal{S}^{-1}(\eta)|^2} \right)^{\frac{s-n}{2}} d\eta \\
&= \gamma(s) \int_{\mathbb{S}^n} \frac{1}{|\xi - \eta|^{n-s} |J_{\mathcal{S}^{-1}}(\xi)|^{\frac{n-s}{2n}} |J_{\mathcal{S}^{-1}}(\eta)|^{\frac{n-s}{2n}}} \left(\frac{2}{1+|\mathcal{S}^{-1}(\eta)|^2} \right)^{\frac{s-n}{2}} d\eta \\
&= \gamma(s) \int_{\mathbb{S}^n} \frac{d\eta}{|\xi - \eta|^{n-s}} (1+\xi_{n+1})^{\frac{n-s}{2}} = \gamma(s) \frac{c_{n,s}}{(1+|x|^2)^{\frac{n-s}{2}}}.
\end{aligned}$$

Thus for any $\lambda > 0$,

$$|\{I_s(g) > \lambda\}| = v_n \left(\left(\frac{\gamma(s)c_{n,s}}{\lambda} \right)^{\frac{2}{n-s}} - 1 \right)^{\frac{n}{2}}. \tag{3.2}$$

Therefore combining (3.1) and (3.2) one have

$$\frac{\|I_s(g)\|_{L^{\frac{n}{n-s},\infty}}}{\|g\|_{L^1}} = v_n^{\frac{n-s}{n}} \sup_{\lambda>0} \left(\gamma(s)^{\frac{2}{n-s}} - \left(\frac{\lambda}{c_{n,s}} \right)^{\frac{2}{n-s}} \right)^{\frac{n-s}{2}} = \gamma(s) v_n^{\frac{n-s}{n}}.$$

Next let us prove the sharp reverse weak estimate for M_s . By the identity in Theorem 1.1 for M_s and (1.2) one can find for any $f \in L^1$,

$$\|M_s(f)\|_{L^{\frac{n}{n-s},\infty}} \geq \|f\|_{L^1}. \tag{3.3}$$

On the other hand, since $M_s(f) \leq 1/\gamma(s) v_n^{\frac{s-n}{n}} I_s(f)$ and we already proved that the function $g = \left(\frac{a}{b+|x-x_0|^2} \right)^{\frac{n+s}{2}}$ satisfies $\|I_s(g)\|_{L^{\frac{n}{n-s},\infty}} = \gamma(s) v_n^{\frac{n-s}{n}} \|g\|_{L^1}$, then by (3.3) the following equality holds

$$\|M_s(g)\|_{L^{\frac{n}{n-s},\infty}} = \|g\|_{L^1}. \tag{3.4}$$

In fact, one can prove (3.4) holds for any L^1 function $f(x) = h(|x - x_0|)$, where h is a radial decreasing function, by using an approach from [2]. First assume $\|f\|_{L^1} = 1$. Let δ_{x_0} denote the Dirac delta mass placed at the x_0 . It is easy to check that

$$M(\delta_{x_0})(x) = \frac{1}{|B(x, |x|)|},$$

where M is the centered Hardy-Littlewood maximum function. Hence, for every $\lambda > 0$, there holds

$$\lambda |\{x : M(\delta_{x_0})(x) > \lambda\}|^{\frac{n}{n-s}} = 1.$$

Since h is a radial decreasing function with $\|h\|_{L^1} = 1$, then by the Lemma 2.1 in [2], one have

$$M(f)(x) \leq M(\delta_{x_0})(x) \text{ for every } x \in \mathbb{R}^n.$$

Then for any $r > 0$ and $x \in \mathbb{R}^n$,

$$\frac{1}{|B(x, r)|^{1-\frac{s}{n}}} \int_{B(x, r)} f(y) dy \leq \left(\frac{1}{|B(x, r)|} \int_{B(x, r)} f(y) dy \|f\|_{L^1}^{\frac{s}{n-s}} \right)^{\frac{n-s}{n}} \leq (M(\delta_{x_0})(x))^{\frac{n-s}{n}},$$

which implies that

$$\|M_s f\|_{L^{\frac{n}{n-s}, \infty}} \leq 1 = \|f\|_{L^1}. \quad (3.5)$$

Combining this inequality with (3.3), one can obtain the desired result for M_s .

What is noteworthy at the end of the section is that this result is also true for centered Hardy-Littlewood maximal function. That is because using the same method one can prove (3.5) when $s = 0$, i.e. (3.5) is true for centered Hardy-Littlewood maximal function. On the other hand, using the limiting weak type behavior for maximum function in [7], (3.3) is also true for centered Hardy-Littlewood maximal function.

4. THE UPPER AND LOWER BOUND OF $\mathcal{C}_{n,s}$

In this section, we will provide an upper and a lower bound for $\mathcal{C}_{n,s}$. Using Theorem 1.1 and (1.2), we can get an upper bound

$$\mathcal{C}_{n,s} \leq \gamma_s \frac{n}{s} v_n^{\frac{n-s}{n}}.$$

To obtain the lower bound, we will use the following formula (see section 5.10 in [9]). Let $0 < \alpha < n$, $0 < s < n$ and $\alpha + s < n$, then

$$\int_{\mathbb{R}^n} \frac{1}{|x-y|^{n-s}} \frac{1}{|y|^{n-\alpha}} dy = C_{n,\alpha,s} \frac{1}{|x|^{n-s-\alpha}} \quad (4.1)$$

with

$$C_{n,\alpha,s} = \pi^{\frac{n}{2}} \frac{\Gamma(\frac{s}{2}) \Gamma(\frac{\alpha}{2}) \Gamma(\frac{n-s-\alpha}{2})}{\Gamma(\frac{n-s}{2}) \Gamma(\frac{n-\alpha}{2}) \Gamma(\frac{s+\alpha}{2})}.$$

Now assume $n - 2 > 4s$. Choose $f(y) = \frac{1}{|y|^{n-2}} \chi_{(|y| \leq 1)}$ and let us prove

$$\|I_s f\|_{L^{\frac{n}{n-s}, \infty}} \geq \gamma_s \frac{v_n^{\frac{n-s}{n}}}{s} \frac{n-2-4s}{2(n-2-s)} \|f\|_{L^1}.$$

Since $|x| \leq \frac{1}{2}$, $|y| > 1$ implies $|y - x| \geq \frac{|y|}{2}$, using (4.1) with $\alpha = 2$ one have

$$\begin{aligned}
\frac{1}{\gamma_s} I_s(f)(x) &= \int_{\mathbb{R}^n} \frac{1}{|x - y|^{n-s}} f(y) dy = \int_{|y| \leq 1} \frac{1}{|x - y|^{n-s}} \frac{1}{|y|^{n-2}} dy \\
&= \int_{\mathbb{R}^n} \frac{1}{|x - y|^{n-s}} \frac{1}{|y|^{n-2}} dy - \int_{|y| > 1} \frac{1}{|x - y|^{n-s}} \frac{1}{|y|^{n-2}} dy \\
&\geq \int_{\mathbb{R}^n} \frac{1}{|x - y|^{n-s}} \frac{1}{|y|^{n-2}} dy - \int_{|y| > 1} \frac{2^{n-s}}{|y|^{2n-2-s}} dy \\
&= \frac{c}{|x|^{n-s-2}} - d,
\end{aligned} \tag{4.2}$$

where

$$c = \frac{4\pi^{n/2}}{(n-s-2)\Gamma(n/2-1)s} \text{ and } d = \frac{2^{n-s+1}\pi^{n/2}}{(n-s-2)\Gamma(n/2)}.$$

Choose $\lambda_0 = \gamma_s(2^{n-s-2}c - d)$, since $\frac{c}{d} = \frac{n-2}{s} \frac{1}{2^{n-s}} > \frac{1}{2^{n-s-2}}$, then λ_0 is positive. Thus by (4.2), there holds

$$|\{I_s f > \lambda_0\}| \geq |\{|x| \leq 1/2, \frac{c}{|x|^{n-s-2}} - d > \frac{\lambda_0}{\gamma_s}\}| = v_n \left(\frac{1}{2}\right)^n. \tag{4.3}$$

Using the fact $\|f\|_{L^1(\mathbb{R}^n)} = \frac{\omega_{n-1}}{2}$ and (4.3) one can obtain

$$\frac{\|I_\alpha f\|_{\frac{n}{n-s}, \infty}}{\|f\|_{L^1}} \geq \frac{\lambda_0 |\{I_\alpha f > \lambda_0\}|^{\frac{n-s}{n}}}{\|f\|_{L^1}^{\frac{n-s}{n}}} = \lambda_0 v_n^{\frac{n-s}{n}} \frac{\Gamma(n/2)}{2^{n-s}\pi^{n/2}} = \gamma_s \frac{v_n^{\frac{n-s}{n}}}{s} \frac{n-2-4s}{2(n-2-s)}.$$

So we complete the proof of Theorem 1.3 .

APPENDIX

In this Appendix, we give an alternative approach to prove the $(L^1, L^{\frac{n}{n-s}, \infty})$ estimate for Riesz potentials, and at the same time this approach also provide an upper bound for $\mathcal{C}_{n,s}$, which have the same behavior with $\gamma_s v_n^{(n-s)/n} n/s$ as $(s, n) \rightarrow (0, \infty)$. First, we state a lemma (see section 3 in [13], also see the Hopf abstract maximal ergodic theorem in [3]) about the weak estimate of the average of the heat-diffusion semi-group $T^t(f) = P_t * f$, where

$$P_t = (4\pi t)^{-\frac{n}{2}} e^{-\frac{|x|^2}{4t}}.$$

Lemma 4.1. *For any $f \in L^1(\mathbb{R}^n)$, there holds*

$$|\{x \in \mathbb{R}^n : \sup_{s>0} \frac{1}{s} \int_0^s P_t f(x) dt > \lambda\}| \leq \frac{1}{\lambda} \|f\|_{L^1(\mathbb{R}^n)}, \lambda > 0.$$

Now let prove the $(L^1, L^{\frac{n}{n-s}, \infty})$ estimate for Riesz potentials $I_s(f)$, which also can be presented by the following formula related to $T^t(f)$,

$$I_s(f)(x) = \frac{1}{\Gamma(s/2)} \int_0^\infty t^{\frac{s}{2}-1} P_t * f(x) dt.$$

We divide the integral into two parts

$$\int_0^\infty t^{\frac{s}{2}-1} P_t * f(x) dt = J_1(f)(x) + J_2(f)(x),$$

where

$$J_1(f)(x) = \int_0^R t^{\frac{s}{2}-1} P_t * f(x) dt,$$

$$J_2(f)(x) = \int_R^\infty t^{\frac{s}{2}-1} P_t * f(x) dt,$$

for some R to be determined later.

Denote $\mathcal{M}^0 f(x) = \sup_{r>0} \frac{1}{r} \int_0^r P_t * f(x) dt$, then we have

$$\begin{aligned} J_1(f)(x) &= \sum_{i=1}^\infty \int_{2^{-i}R}^{2^{-i+1}R} t^{\frac{s}{2}-1} P_t * f(x) dt \\ &\leq \sum_{i=1}^\infty \int_{2^{-i}R}^{2^{-i+1}R} (2^{-i}R)^{\frac{s}{2}-1} P_t * f(x) dt \\ &\leq \sum_{i=1}^\infty (2^{-i}R)^{\frac{s}{2}-1} 2^{-i+1}R \left(\frac{1}{2^{-i+1}R} \int_0^{2^{-i+1}R} P_t * f(x) dt \right) \\ &\leq 2R^{\frac{s}{2}} \frac{2^{-\frac{s}{2}}}{1 - 2^{-\frac{s}{2}}} \mathcal{M}^0 f(x). \end{aligned} \tag{4.4}$$

On the other hand, by direct computation, we obtain that

$$\begin{aligned} J_2(f)(x) &\leq \int_R^\infty t^{\frac{s}{2}-1} \|P_t\|_{L^\infty} \|f\|_{L^1} dt \\ &\leq \frac{2}{n-s} (4\pi)^{-\frac{n}{2}} R^{\frac{s}{2}-\frac{n}{2}} \|f\|_{L^1}. \end{aligned} \tag{4.5}$$

Combining (4.4) and (4.5), we obtain that

$$I_s(f)(x) \leq \frac{1}{\Gamma(s/2)} \left(2R^{\frac{s}{2}} \frac{2^{-\frac{s}{2}}}{1 - 2^{-\frac{s}{2}}} \mathcal{M}^0 f(x) + \frac{2}{n-s} (4\pi)^{-\frac{n}{2}} R^{\frac{s}{2}-\frac{n}{2}} \|f\|_{L^1} \right) \tag{4.6}$$

for all $R > 0$. The choice of

$$R = \left(\frac{(4\pi)^{-\frac{n}{2}} \|f\|_{L^1}}{\frac{s}{2^{\frac{s}{2}-1}-1} \mathcal{M}^0 f(x)} \right)^{\frac{2}{n}}$$

minimizes the right side of the expression in (4.6). Thus

$$I_s(f)(x) \leq \tau_s(\mathcal{M}^0 f(x))^{\frac{n-s}{n}} \|f\|_{L^1}^{\frac{s}{n}}, \tag{4.7}$$

where

$$\tau_s = 2(4\pi)^{-\frac{s}{2}} (2^{\frac{s}{2}} - 1)^{\frac{s-n}{n}} \frac{n}{n-s} \left(\frac{1}{s}\right)^{\frac{s}{n}} \frac{1}{\Gamma(s/2)}.$$

Now using lemma 4.1 one can see that

$$\begin{aligned}
\lambda |\{I_s f > \lambda\}|^{\frac{n-s}{n}} &\leq \lambda |\{\tau_s(\mathcal{M}^0 f(x))^{\frac{n-s}{n}} \|f\|_{L^1}^{\frac{s}{n}} > \lambda\}|^{\frac{n-s}{n}} \\
&\leq \lambda \left[\left(\frac{\tau_s \|f\|_{L^1}^{\frac{s}{n}}}{\lambda} \right)^{\frac{n}{n-s}} \|f\|_{L^1}^{\frac{n-s}{n}} \right]^{\frac{n-s}{n}} \\
&\leq \tau_s \|f\|_{L^1}.
\end{aligned}$$

Notice that

$$2^{\frac{s}{2}} - 1 > \frac{\ln 2}{2} s \text{ for } s > 0,$$

thus,

$$\tau_s \leq \frac{2}{\ln 2} \left(\frac{1}{4\pi} \right)^{-\frac{s}{2}} \frac{1}{\Gamma(\frac{s}{2} + 1)} \frac{n}{n-s}.$$

So by this approach, one can obtain that $\mathcal{C}_{n,s} \leq \frac{2}{\ln 2} \left(\frac{1}{4\pi} \right)^{-\frac{s}{2}} \frac{1}{\Gamma(\frac{s}{2} + 1)} \frac{n}{n-s}$ and it is easy to check that when $(s, n) \rightarrow (0, \infty)$,

$$\frac{2}{\ln 2} \left(\frac{1}{4\pi} \right)^{-\frac{s}{2}} \frac{1}{\Gamma(\frac{s}{2} + 1)} \frac{n}{n-s} = O(\gamma_s v_n^{\frac{n-s}{n}} \frac{n}{s}).$$

REFERENCES

- [1] R. Adams, *A note on Riesz potentials*. Duke Math. J. **42** (1975), 765–778.
- [2] J. M. Aldaz and J. Pérez Lázaro, *The best constant for the centered maximal operator on radial decreasing functions*. Math. Inequal. Appl. **14** (2011), no. 1, 173–179.
- [3] N. Dunford and J. Schwartz, *Linear Operators. Part I. General Theory*. With the Assistance of W.G. Bade and R.G. Bartle, Reprint of the 1958 original Wiley Classics Library, a Wiley-Interscience Publication, John Wiley & Sons, Inc., New York,
- [4] R. Frank and E. Lieb, *A new, rearrangement-free proof of the Sharp Hardy-Littlewood-Sobolev Inequality*. Spectral Theory, Function Spaces and Inequalities. Oper. Theory: Adv. Appl. **219** (2012), 55–67.
- [5] L. Grafakos, *Classical Fourier Analysis*. Third edition. Graduate Texts in Mathematics, 250. Springer, New York, 2014.
- [6] G. Hardy and J. Littlewood, *Some properties of fractional integral I*. Math. Zeit. (1927), 565–606.
- [7] P. Janakiraman, *Limiting weak-type behavior for singular integral and maximal operators*. Trans. Amer. Math. Soc. **358** (2006), no. 5, 1937–1952.
- [8] E. Lieb, *Sharp constants in the Hardy-Littlewood-Sobolev and related inequalities*. Ann. of Math. (2) **118** (1983), no. 2, 349–374.
- [9] E. Lieb and M. Loss, *Analysis*. Second edition. Graduate Studies in Mathematics, 14. American Mathematical Society, Providence, RI, 2001.
- [10] B. Muckenhoupt and R. Wheeden, *Weighted norm inequalities for fractional integrals*. Trans. Amer. Math. Soc. **192** (1974), 261–274.
- [11] M. Riesz, *L'integrale de Riemann-Liouville et le probleme de Cauchy*. Acta Math. **81** (1949), 1–222.
- [12] S. Sobolev, *On a theorem in functional analysis* (in Russian). Mat. Sob. **46** (1938), 471–497.
- [13] E. Stein and J. Stromberg, *Bagavior of maximal functions in \mathbb{R}^n for largr n* . Ark. Mat. **21** (1983), 259–269.
- [14] H. Tang, *Limiting weak-type behavior for fractional integral operators*(Chinese). Beijing Shifan Daxue Xuebao **52** (2016), no. 1, 5–7.
- [15] H. Tang and G.Wang *Limiting weak type behavior for multilinear fractional integrals*. Nonlinear Anal. **197** (2020), 111858, 13 pp.

- [16] A. Zygmund, *On a theorem of Marcinkiewicz concerning interpolation of operators*. Jour. de Math. Pures et Appliquées **35** (1956), 223–248.

SCHOOL OF SCIENCE, XI'AN UNIVERSITY OF POSTS AND TELECOMMUNICATIONS, XI'AN, 710121, CHINA, E-MAIL: HUANGLIANG10@163.COM

LABORATORY OF MATHEMATICS AND COMPLEX SYSTEMS (MINISTRY OF EDUCATION), SCHOOL OF MATHEMATICAL SCIENCES, BEIJING NORMAL UNIVERSITY, BEIJING, 100875, CHINA, E-MAIL: HLTANG@BNU.EDU.CN