

From quartic anharmonic oscillator to double well potential

Alexander V. Turbiner* and J.C. del Valle†

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,

Apartado Postal 70-543, 04510 México, D.F., Mexico

(ΩDated: January 3, 2022)

Abstract

It is already known that the quantum quartic single-well anharmonic oscillator $V_{ao}(x) = x^2 + g^2 x^4$ and double-well anharmonic oscillator $V_{dw}(x) = x^2(1 - gx)^2$ are essentially one-parametric, their eigenstates depend on a combination $(g^2 \hbar)$. Hence, these problems are reduced to study the potentials $V_{ao} = u^2 + u^4$ and $V_{dw} = u^2(1 - u)^2$, respectively. It is shown that by taking uniformly-accurate approximation for anharmonic oscillator eigenfunction $\Psi_{ao}(u)$, obtained recently, see JPA 54 (2021) 295204 [1] and Arxiv 2102.04623 [2], and then forming the function $\Psi_{dw}(u) = \Psi_{ao}(u) \pm \Psi_{ao}(u - 1)$ allows to get the highly accurate approximation for both the eigenfunctions of the double-well potential and its eigenvalues.

*Electronic address: turbiner@nucleares.unam.mx

†Electronic address: delvalle@correo.nucleares.unam.mx

I. INTRODUCTION

For the one-dimensional quantum quartic single-well anharmonic oscillator $V_{ao}(x) = x^2 + g^2 x^4$ and double-well anharmonic oscillator with potential $V_{dw}(x) = x^2(1 - gx)^2$ the (trans)series in the coupling constant g (which is the Perturbation Theory in powers of g (the Taylor expansion) in the former case of $V_{ao}(x)$ supplemented by exponentially-small terms in g in the latter case of $V_{dw}(x)$) and the semiclassical expansion in $\hbar$ (the Taylor expansion for $V_{ao}(x)$ supplemented by the exponentially small terms in $\hbar$ for $V_{dw}(x)$) for energies *coincide* [3]. This property plays crucially important role in our consideration.

Both the quartic anharmonic oscillator

$$V = x^2 + g^2 x^4, \quad (1)$$

with a single harmonic well at $x = 0$ and the double-well potential

$$V = x^2(1 - gx)^2, \quad (2)$$

with two symmetric harmonic wells at $x = 0$ and $x = 1/g$, respectively, are particular cases of the quartic polynomial potential

$$V = x^2 + agx^3 + g^2 x^4, \quad (3)$$

where g is the coupling constant and a is a parameter. Interestingly, the potential (3) is symmetric for three particular values of the parameter a : $a = 0$ and $a = \pm 2$. All three potentials (1), (2), (3) belong to the family of potentials of the form

$$V = \frac{1}{g^2} \tilde{V}(gx),$$

for which there exists a remarkable property: the Schrödinger equation becomes one-parametric, both the Planck constant $\hbar$ and the coupling constant g appear in the combination $(\hbar g^2)$, see [2]. It can be immediately seen if instead of the coordinate x the so-called classical coordinate $u = (gx)$ is introduced. This property implies that the action S in the path integral formalism becomes g -independent and the factor $\frac{1}{\hbar}$ in the exponent becomes $\frac{1}{\hbar g^2}$ [4]. Formally, the potentials (1)-(2), which enter to the action, appear at $g = 1$, hence, in the form

$$V = u^2 + u^4, \quad (4)$$

$$V = u^2(1-u)^2, \quad (5)$$

respectively. Both potentials are symmetric with respect to $u = 0$ and $u = 1/2$, respectively.

Namely, this form of the potentials will be used in this short Note. This Note is the extended version of a part of presentation in AAMP-18 given by the first author [5].

II. SINGLE-WELL POTENTIAL

In [1] for the potential (4) matching the small distances $u \rightarrow 0$ expansion and the large distances $u \rightarrow \infty$ expansion (in the form of semiclassical expansion) for the phase ϕ in the representation

$$\Psi = P(u) e^{-\phi(u)},$$

of the wave function, where P is a polynomial, it was constructed the following function for the $(2n + p)$ -excited state with quantum numbers (n, p) , $n = 0, 1, 2, \dots$, $p = 0, 1$:

$$\begin{aligned} \Psi_{(approximation)}^{(n,p)} = & \\ & \frac{u^p P_{n,p}(u^2)}{(B^2 + u^2)^{\frac{1}{4}} (B + \sqrt{B^2 + u^2})^{2n+p+\frac{1}{2}}} \\ & \exp \left(- \frac{A + (B^2 + 3)u^2/6 + u^4/3}{\sqrt{B^2 + u^2}} + \frac{A}{B} \right), \end{aligned} \quad (6)$$

where $P_{n,p}$ is some polynomial of degree n in u^2 with positive roots. Here $A = A_{n,p}$, $B = B_{n,p}$ are two parameters of interpolation. These parameters $(-A)$, B are slow-growing with quantum number n at fixed p taking, in particular, the values

$$A_{0,0} = -0.6244, B_{0,0} = 2.3667, \quad (7)$$

$$A_{0,1} = -1.9289, B_{0,1} = 2.5598, \quad (8)$$

for the ground state and the first excited state, respectively. This remarkably simple function (6), see Fig.1 (top), provides 10-11 exact figures in energies for the first 100 eigenstates. Furthermore, the function (6) deviates uniformly for $u \in (-\infty, +\infty)$ from the exact function in $\sim 10^{-6}$.




FIG. 1: Two lowest, normalized to one eigenfunctions of positive/negative parity: for single-well potential (4), see (6) (top) and for double-well potential (5), see (9)(bottom). Potentials shown by black lines.

III. DOUBLE-WELL POTENTIAL: WAVEFUNCTIONS

Following the prescription, usually assigned in folklore to E.M. Lifschitz - one of the authors of the famous Course on Theoretical Physics by L.D. Landau and E.M. Lifschitz - when a wavefunction for single well potential with minimum at $u = 0$ is known, $\Psi(u)$, the wavefunction for double well potential with minima at $u = 0, 1$ can be written as $\Psi(u) \pm \Psi(u - 1)$. This prescription was already checked successfully for the double-well potential (2) in [6] for somehow simplified version of (6), based on matching the small distances $u \rightarrow 0$ expansion and the large distances $u \rightarrow \infty$ expansion for the phase ϕ but ignoring subtleties emerging in semiclassical expansion. Taking the wavefunction (6) one can construct

$$\Psi_{(approximation)}^{(n,p)} = \frac{P_{n,p}(\tilde{u}^2)}{(B^2 + \tilde{u}^2)^{\frac{1}{4}} (\alpha B + \sqrt{B^2 + \tilde{u}^2})^{2n+\frac{1}{2}}} \exp\left(-\frac{A + (B^2 + 3)\tilde{u}^2/6 + \tilde{u}^4/3}{\sqrt{B^2 + \tilde{u}^2}} + \frac{A}{B}\right) D^{(p)}, \quad (9)$$

where $p = 0, 1$ and

$$\begin{aligned} D^{(0)} &= \cosh \left(\frac{a_0 \tilde{u} + b_0 \tilde{u}^3}{\sqrt{B^2 + \tilde{u}^2}} \right), \\ D^{(1)} &= \sinh \left(\frac{a_1 \tilde{u} + b_1 \tilde{u}^3}{\sqrt{B^2 + \tilde{u}^2}} \right). \end{aligned}$$

Here

$$\tilde{u} = u - \frac{1}{2}, \quad (10)$$

$\alpha = 1$ and $A, B, a_{0,1}, b_{0,1}$ are variational parameters. If $\alpha = 0$ as well as $b_{0,1} = 0$ the function (9) is reduced to ones which were explored in [6], see Eqs.(10)-(11). The polynomial $P_{n,p}$ is found unambiguously after imposing the orthogonality conditions of $\Psi_{(approximation)}^{(n,p)}$ to $\Psi_{(approximation)}^{(k,p)}$ at $k = 0, 1, 2, \dots, (n-1)$, here it is assumed that the polynomials $P_{k,p}$ at $k = 0, 1, 2, \dots, (n-1)$ are found beforehand.

IV. DOUBLE-WELL POTENTIAL: RESULTS

In this section we present concrete results for energies of the ground state $(0, 0)$ and of the first excited state $(0, 1)$ obtained with the function (9) at $p = 0, 1$, respectively. The results are compared with the Lagrange-Mesh Method (LMM) [7].

A. Ground State (0,0)

The ground state energy for (5) obtained variationally using the function (9) at $p = 0$ and compared with LMM results [7], where all printed digits (in the second line) are correct,

$$\begin{aligned} E_{var}^{(0,0)} &= 0.932\,517\,518\,401, \\ E_{mesh}^{(0,0)} &= 0.932\,517\,518\,372. \end{aligned}$$

Note that ten decimal digits in $E_{var}^{(0,0)}$ coincide with ones in $E_{mesh}^{(0,0)}$ (after rounding). Variational parameters in (9) take values,

$$\begin{aligned} A &= 2.3237, \\ B &= 3.2734, \\ a_0 &= 2.3839, \\ b_0 &= 0.0605, \end{aligned}$$

cf.(7). Note that b_0 takes a very small value.

B. First Excited State (0,1)

The first excited state energy for (5) obtained variationally using the function (9) at $p = 1$ and compared with LMM results [7], where all printed digits (in the second line) are correct,

$$\begin{aligned} E_{var}^{(0,1)} &= 3.396\,279\,329\,936\,, \\ E_{mesh}^{(0,1)} &= 3.396\,279\,329\,887\,.\end{aligned}$$

Note that ten decimal digits in $E_{var}^{(0,1)}$ coincide with ones in $E_{mesh}^{(0,1)}$ (after rounding). Variational parameters in (9) take values,

$$\begin{aligned} A &= -2.2957\,, \\ B &= 3.6991\,, \\ a_1 &= 4.7096\,, \\ b_1 &= 0.0590\,,\end{aligned}$$

cf.(8). Note that b_1 takes a very small value similar to b_0 .

V. CONCLUSIONS

It is presented the approximate expression (9) for the eigenfunctions in the double-well potential (5). In Non-Linearization procedure [8] it can be calculated the first correction (the first order deviation) to the function (9). It can be shown that for any $u \in (-\infty, +\infty)$ the functions (9) deviate uniformly from the exact eigenfunctions, beyond the sixth significant figure similarly to the function (6) for the single-well case. It increases the accuracy of the simplified function, proposed in [5] with $\alpha = 0$ and $b_{0,1} = 0$, in the domain under the barrier $u \in (0.25, 0.75)$ from 4 to 6 significant figures leaving the accuracy outside of this domain practically unchanged.

Acknowledgments

This work is partially supported by CONACyT grant A1-S-17364 and DGAPA grant IN113819 (Mexico). AVT thanks the PASPA-UNAM program for support during his sab-

batical leave.

-
- [1] A.V. Turbiner, J.C. del Valle,
Anharmonic oscillator: a solution,
J Phys. A **54** (2021) 295204
DOI: 10.1088/1751-8121/ac0733 ;
Talk presented by AVT at CRM, Montreal, Canada (February 23, 2021)
- [2] A.V. Turbiner and E. Shuryak,
On connection between perturbation theory and semiclassical expansion in quantum mechanics,
Arxiv: 2102.04623 (February(version-1) - August(version-2), 2021)
- [3] E. Shuryak and A.V. Turbiner,
Trans-series for the ground state density and Generalized Bloch equation,
Phys Rev D **98** (2018) 105007 (10pp) doi: 10.1103/PhysRevD.98.105007
- [4] M.A. Escobar-Ruiz, E. Shuryak and A.V. Turbiner,
Phys. Rev. D **93** (2016) 105039
doi: 10.1103/PhysRevD.93.105039
- [5] A.V. Turbiner, J.C. del Valle,
Anharmonic oscillator: almost analytic solution,
Talk presented by AVT at AAMP-18 (Sept.1-3), Prague, Czech Republic (September 1, 2021)
- [6] A.V. Turbiner,
Double well potential: perturbation theory, tunneling, WKB (beyond instantons),
Int.Journ.Mod.Phys. A **25**, 647-658 (2010)
DOI: 10.1142/S0217751X10048937
- [7] A. V. Turbiner, J.C. del Valle,
Comment on: Uncommonly accurate energies for the general quartic oscillator, Int. J. Quantum Chem., e26554 (2020), by P. Okun and K. Burke,
Int Journal of Quantum Chemistry **122** (2021) qua.26766 (4pp) DOI: 10.1002/qua.26766
- [8] A.V. Turbiner,
Soviet Phys. - Usp. Fiz. Nauk. **144**, 35-78 (1984),
Sov. Phys. Uspekhi **27**, 668-694 (1984) (English Translation)