

# The set of stable indices of 0-1 matrices with a given order

Zhibing Chen, Zejun Huang\*

College of Mathematics and Statistics, Shenzhen University, Shenzhen 518060, China

## Abstract

The stable index of a 0-1 matrix  $A$  is defined to be the smallest integer  $k$  such that  $A^{k+1}$  is not a 0-1 matrix if such an integer exists; otherwise the stable index of  $A$  is defined to be infinity. We characterize the set of stable indices of 0-1 matrices with a given order.

**Keywords:** 0-1 matrix, digraph, stable index, walk

**Mathematics Subject Classification:** 05C50, 05C20, 15A99

## 1 Introduction and Main Results

Combinatorial problems on the power of nonnegative matrices is an interesting topic in matrix theory. One of the classical problems in this topic is the characterization of the exponents of primitive matrices with a given order. Let  $A$  be a nonnegative matrix. Denote by  $\rho(A)$  the spectral radius of  $A$ . By the famous Perron-Frobenius theorem we know that  $\rho(A)$  is an eigenvalue of  $A$ . If  $A$  has no other eigenvalue of modulus  $\rho(A)$ , then  $A$  is called *primitive*. Frobenius proved that a nonnegative square matrix is primitive if and only if there is a positive integer  $k$  such that  $A^k$  is positive entrywise. The *exponent* of a primitive matrix is the smallest positive integer  $k$  such that  $A^k$  is positive. A natural interesting problem is which numbers are the exponents of nonnegative matrices with a given order, which has been solved by Wielandt [6], Dulmage and Mendelsohn [3], Lewin and Vitek [4], Shao [5] and Zhang [7]. It turns out that the set of exponents of primitive matrices of order  $n$  is not a consecutive set in the interval  $[1, (n-1)^2 + 1]$ , which is quite surprising.

Analogous to the exponent of primitive matrices, Chen, Huang and Yan [2] introduce the concept of *stable index* for 0-1 matrices as follows.

**Definition.** *The stable index of a 0-1 matrix  $A$ , denoted by  $\theta(D)$ , is defined to be the smallest integer  $k$  such that  $A^{k+1}$  is not a 0-1 matrix if such an integer exists; otherwise the stable index of  $A$  is defined to be infinity.*

---

\*Corresponding author ( Email: mathzejun@gmail.com)

The stable index of 0-1 matrices is closely related to directed walks of digraphs. Digraphs in this paper allow loops but do not allow multiple arcs. We follow the terminology on digraphs in [1]. Directed paths, directed cycles and directed walks will be abbreviated as paths, cycles and walks, respectively. A path (walk) with initial vertex  $u$  and terminal vertex  $v$  is called a  $uv$ -path ( $uv$ -walk). A walk (path, cycle) of length  $k$  is called a  $k$ -walk ( $k$ -path,  $k$ -cycle). The number of vertices in a digraph is called its *order*.

Denote by  $M_n\{0, 1\}$  the set of 0-1 matrices of order  $n$ . Given a matrix  $A = (a_{ij}) \in M_n\{0, 1\}$ , we define its digraph as  $D(A) = (\mathcal{V}, \mathcal{A})$  with the vertex set  $\mathcal{V} = \{1, 2, \dots, n\}$  and the arc set  $\mathcal{A} = \{(i, j) : a_{ij} = 1, 1 \leq i, j \leq n\}$ . Conversely, given a digraph  $D = (\mathcal{V}, \mathcal{A})$  with a vertex set  $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$  and an arc set  $\mathcal{A}$ , its *adjacency matrix* is defined as  $A_D = (a_{ij})_{n \times n}$ , where

$$a_{ij} = \begin{cases} 1, & \text{if } (v_i, v_j) \in \mathcal{A}; \\ 0, & \text{otherwise.} \end{cases}$$

Denote by  $A(i, j)$  the  $(i, j)$ -entry of  $A$ . Given  $A \in M_n\{0, 1\}$ ,  $A^k(i, j) = t$  if and only if  $D(A)$  has exactly  $t$  distinct  $ij$ -walks of length  $k$ .

Using adjacency matrices, we have an equivalent definition for the stable index of digraphs as follows.

**Definition.** *The stable index of a digraph  $D$ , denoted by  $\theta(D)$ , is defined to be the smallest integer  $k$  such that  $D$  contains two distinct  $(k+1)$ -walks with the same initial vertex and terminal vertex if such an integer exists; otherwise the stable index of  $D$  is defined to be infinity.*

Given a digraph  $D$ , by the above definitions we have  $\theta(D) = \theta(A_D)$ . Conversely, given a square 0-1 matrix  $A$ , we have  $\theta(A) = \theta(D(A))$ .

Given a positive integer  $n$ , we denote by  $s(n)$  the maximum finite stable index of 0-1 matrices with order  $n$ . The precise value of  $s(n)$  was determined in [2]. For  $n \leq 6$ , we have  $s(2) = 1, s(3) = 3, s(4) = 4, s(5) = 6, s(6) = 7$ ; see [2]. For  $n \geq 7$  we have

**Theorem 1.** [2] *Let  $n \geq 7$  be an integer. Then*

$$s(n) = \begin{cases} \frac{n^2-1}{4}, & \text{if } n \text{ is odd,} \\ \frac{n^2-4}{4}, & \text{if } n \equiv 0 \pmod{4}, \\ \frac{n^2-16}{4}, & \text{if } n \equiv 2 \pmod{4}. \end{cases} \quad (1)$$

Similar with the exponent of primitive matrices, we are interested in the following natural

problem.

**Problem 2.** *Given a positive integer  $n$ , which numbers can be the stable indices of 0-1 matrices (digraphs) of order  $n$ ?*

We solve this problem in this paper. Denote by  $\mathbb{Z}^+$  the set of positive integers and denote by  $\text{LCM}(p, q)$  the least common multiple of two integers  $p$  and  $q$ . For  $k \in \mathbb{Z}^+$ , we use  $[k]$  to represent the set  $\{1, 2, \dots, k\}$ . Let  $\Theta(n)$  be the set of stable indices of 0-1 matrices with order  $n$ . Then  $\Theta(n)$  is also the set of indices of digraphs with  $n$  vertices. Our main result states as follows.

**Theorem 3.** *Let  $n \geq 7$  be an integer. Then*

$$\Theta(n) = [s(n-1) + 1] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\} \cup \{\infty\}.$$


For the sake of convenience, we prove Theorem 3 for digraphs.

## 2 Proof of Theorem 3

Two digraphs  $D_1 = (\mathcal{V}_1, \mathcal{A}_1)$  and  $D_2 = (\mathcal{V}_2, \mathcal{A}_2)$  are *isomorphic*, written  $D_1 \cong D_2$ , if there is a bijection  $\sigma : \mathcal{V}_1 \rightarrow \mathcal{V}_2$  such that  $(u, v) \in \mathcal{A}_1$  if and only if  $(\sigma(u), \sigma(v)) \in \mathcal{A}_2$ . It is clear that two digraphs are isomorphic if and only if their adjacency matrices are permutation similar. We say a digraph  $D$  contains a *copy* of  $H$  if  $D$  has a subgraph isomorphic to  $H$ .

In a digraph  $D$ , if there is a walk from  $u$  to  $v$  for all  $u, v \in \mathcal{V}(D)$ , then  $D$  is said to be *strongly connected*. A digraph is strongly connected if and only if its adjacency matrix is irreducible.

Denote by  $\vec{C}_k$  a  $k$ -cycle. Given an integer  $k \geq 2$  and two disjoint cycles  $\vec{C}_p$  and  $\vec{C}_q$ , let  $\vec{g}(p, k, q)$  be the digraph obtained by adding a  $(k-1)$ -path from a vertex of  $\vec{C}_p$  to a vertex of  $\vec{C}_q$ , which has the following diagram. When  $k = 2$ , we abbreviate  $\vec{g}(p, k, q)$  as  $\vec{g}(p, q)$ .



It is clear that

$$\theta(\vec{g}(p, k, q)) = \text{LCM}(p, q) + k - 2.$$

Denote by  $C_p$  the adjacency matrix of the  $p$ -cycle  $v_1 \cdots v_p v_1$ , which is

$$\begin{bmatrix} & 1 & & \\ & & \ddots & \\ & & & 1 \\ 1 & & & \end{bmatrix}.$$

We need the following lemmas.

**Lemma 4.** [2] *Let  $D$  be a strongly connected digraph of order  $n$ . If  $D$  is a cycle, then  $\theta(D) = \infty$ ; otherwise, we have  $\theta(D) \leq n$ .*

**Lemma 5.** *Let  $A = \begin{pmatrix} C_p & X \\ 0 & C_q \end{pmatrix} \in M_{p+q}\{0,1\}$ , where  $X$  has exactly  $k$  nonzero entries. Then*

$$\theta(A) \leq \lfloor \frac{pq}{k} \rfloor.$$

*Proof.* Notice that

$$A^m = \begin{pmatrix} C_p^m & \sum_{k=0}^{m-1} C_p^k X C_q^{m-1-k} \\ 0 & C_q^m \end{pmatrix} \equiv \begin{pmatrix} C_p^m & B \\ 0 & C_q^m \end{pmatrix}$$

for all positive integer  $m$ . Since  $C_p^k X C_q^{m-1-k}$  only changes the positions of  $X$ 's entries, each  $C_p^k X C_q^{m-1-k}$  has exactly  $k$  entries equal to 1. So the summation of all entries in  $B$  is  $km$ . If  $m > \lfloor pq/k \rfloor$ , then  $km > pq$ , which implies that  $B$  is not a 0-1 matrix. Therefore,  $\theta(A) \leq \lfloor pq/k \rfloor$ .  $\square$

**Corollary 6.** *Let  $p, q, n$  be positive integers such that  $p + q = n$ . Suppose  $D$  is a digraph of order  $n$  containing a copy of  $\vec{g}(p, q)$ . If  $\theta(D) > \max\{n, \lfloor pq/2 \rfloor\}$ , then  $D \cong \vec{g}(p, q)$ .*

*Proof.* Since  $D$  contains a copy of  $\vec{g}(p, q)$ , we may assume

$$A_D = \begin{pmatrix} C_p + A_{11} & A_{12} \\ A_{21} & C_q + A_{22} \end{pmatrix},$$

where  $A_{12} \neq 0$ .

If  $A_{21} \neq 0$ , then  $A_D$  is irreducible. Applying Lemma 4 we have  $\theta(D) \leq n$ , a contradiction. Therefore, we have  $A_{21} = 0$ .

If  $A_{11} \neq 0$ , then since  $C_p + A_{11}$  is irreducible, we have  $\theta(D) \leq \theta(C_p + A_{11}) \leq p$ , a contradiction. Hence,  $A_{11} = 0$ . Similarly, we have  $A_{22} = 0$ .

Now applying Lemma 5, we can deduce that  $A_{12}$  has exactly one nonzero entry. Therefore,  $D \cong \vec{\mathcal{J}}(p, q)$ .  $\square$

**Lemma 7.** *Suppose  $p, q > 0$  are relatively prime numbers. Let  $kq \equiv x_k \pmod{p}$  for  $1 \leq k \leq p-1$ . Then*

$$\{x_1, x_2, \dots, x_{p-1}\} = \{1, 2, \dots, p-1\}.$$

*Proof.* It suffices to prove  $x_u \neq x_v$  for all  $u > v$ . Otherwise, suppose there exist  $u, v \in \{1, 2, \dots, p-1\}$  such that  $u > v$  and  $x_u = x_v$ . Then  $x_u - x_v = (u-v)q$  is divided by  $p$ . Since  $(p, q) = 1$ , we have  $p \mid u-v$ , which contradicts the fact  $1 \leq u-v < p-1$ . This completes the proof.  $\square$

**Corollary 8.** *Suppose  $p, q$  are relatively prime numbers such that  $p > q > 1$ . Then*

$$\min\{u : \text{there exists a nonnegative numbers } v \text{ such that } p-q = uq - vp\} = p-1.$$

*Proof.* Notice that  $(p-1)q \equiv p-q \pmod{p}$ . Applying Lemma 7 we have the conclusion.  $\square$

For the sake of simplicity, we denote by

$$[a, b] = \{x \in \mathbb{Z}^+ : a \leq x \leq b\}.$$

Now we are ready to present the proof of our main result.

*Proof of Theorem 3.* We first show

$$\Theta(n) \subseteq [s(n-1) + 1] \cup \{\text{LCM}(p, q) : p+q = n, p, q \in \mathbb{Z}^+\} \cup \{\infty\}. \quad (2)$$

Suppose  $D$  has finite stable index  $r$ . Then  $D$  has two distinct  $(r+1)$ -walks  $\vec{w}_1$  and  $\vec{w}_2$  from  $x$  to  $y$  for some vertices  $x, y$ . Denote by  $\vec{w}_1 \cup \vec{w}_2$  the union of  $\vec{w}_1$  and  $\vec{w}_2$ , i.e., the subgraph of  $D$  with vertex set  $\mathcal{V}(\vec{w}_1) \cup \mathcal{V}(\vec{w}_2)$  and arc set  $\mathcal{A}(\vec{w}_1) \cup \mathcal{A}(\vec{w}_2)$ .

If the number of cycles in  $\vec{w}_1 \cup \vec{w}_2$  is less than 2, then by the proof of Theorem 1 in [2] we have  $\theta(D) < n$ . If  $\vec{w}_1 \cup \vec{w}_2$  contain two cycles whose vertex sets have nonempty intersection, then the subgraph  $D_1$  induced by these two cycles is strongly connected. Applying Lemma

4, we have  $\theta(D) \leq \theta(D_1) \leq n$ . Now suppose  $\vec{w}_1 \cup \vec{w}_2$  contains two disjoint cycles. We distinguish three cases.

*Case 1.*  $\vec{w}_1 \cup \vec{w}_2$  contains a copy of  $\vec{g}(p, q)$  with  $p + q = n$ . Then applying Corollary 6, we have either  $\theta(D) \leq \max\{n, \lfloor pq/2 \rfloor\}$  or  $D \cong \vec{g}(p, q)$ , while the later case leads to  $\theta(D) = \text{LCM}(p, q)$ . Thus

$$\theta(D) \in [\max\{n, \lfloor n/2 \rfloor \lceil n/2 \rceil / 2\}] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\}.$$

Since  $\max\{n, \lfloor n/2 \rfloor \lceil n/2 \rceil / 2\} \leq s(n-1) + 1$ , we have

$$\theta(D) \in [s(n-1) + 1] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\}.$$

*Case 2.*  $\vec{w}_1 \cup \vec{w}_2$  contains no copy of  $\vec{g}(p, q)$  with  $p + q = n$  and it contains a copy of  $\vec{g}(p, k, q)$  for some positive integers  $p, q, k$  with  $p + q < n$ . Since  $\theta(\vec{g}(p, k, q)) = \text{LCM}(p, q) + k - 2$ , we have  $\theta(D) \leq \text{LCM}(p, q) + k - 2$ . Let

$$\lambda_1(n) = \max\{\theta(\vec{g}(p, k, q)) : p + q + k - 2 \leq n, p + q \leq n - 1\}.$$

Note that

$$s(n) = \max\{\text{LCM}(p, q) : p + q = n\} \quad \text{for } n \geq 7.$$

If  $n = 7$ , since  $s(6) = 7$ , we have

$$\lambda_1(7) = \theta(\vec{g}(2, 4, 3)) = 8 = s(6) + 1.$$

For  $n \geq 8$ , since  $s(n)$  is strictly increasing for  $n \in \mathbb{Z}^+$ , we have

$$\theta(\vec{g}(p, k, q)) = \text{LCM}(p, q) + k - 2 \leq s(p + q) + n - p - q \leq s(n - 1) + 1.$$


Hence,

$$\lambda_1(n) = s(n - 1) + 1 \quad \text{for } n \geq 7$$

and

$$\theta(D) \leq s(n - 1) + 1.$$

*Case 3.*  $\vec{w}_1 \cup \vec{w}_2$  does not contain any copy of  $\vec{g}(p, k, q)$ . Then similarly as in the proof of Theorem 1 in [2],  $\vec{w}_1 \cup \vec{w}_2$  has the following diagram.



Suppose the  $xy$ -paths contained in  $\vec{w}_1$  and  $\vec{w}_2$  have lengths  $l$  and  $t$ , respectively. Then we have

$$\begin{aligned}\theta(D) &= \min\{l + up : l + up = t + vq \text{ for some nonnegative integers } u, v\} - 1 \\ &\leq \min\{l + (q - 1)p, t + (p - 1)q\} - 1.\end{aligned}\quad (3)$$

Next we show

$$\theta(D) \leq s(n - 1) + 1.$$

If  $p = q$ , then we have  $\theta(D) \leq \max\{l, t\} < s(n - 1) + 1$ . Next we assume  $p > q$ . Since  $l \leq n - 1 - q$ ,  $t \leq n - 1 - p$  and  $p + q \leq n - 2$ , by (3) we have

$$\theta(D) \leq t + (p - 1)q - 1 \leq n - 2 - p + (p - 1)q = n - 3 + (p - 1)(q - 1) \equiv h(n, p, q).$$

If  $n$  is odd, then  $\max\{h(n, p, q) : p + q \leq n - 2, p > q\}$  is attained at  $(p, q) = ((n - 1)/2, (n - 3)/2)$ . Hence, by (1) we have

$$\theta(D) \leq h(n, p, q) \leq n - 3 + (n - 3)(n - 5)/4 = (n^2 - 4n + 3)/4 \leq s(n - 1) + 1.$$

If  $n$  is even, then  $\max\{h(n, p, q) : p + q \leq n - 2, p > q\}$  is attained at  $(p, q) = (n/2, n/2 - 2)$ . Hence, by (1) we have

$$\theta(D) \leq h(n, p, q) \leq n - 3 + (n/2 - 1)(n/2 - 3) = (n^2 - 4n)/4 \leq s(n - 1) + 1.$$


Combining all the above cases we have (2). Now we prove

$$[s(n - 1) + 1] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\} \cup \{\infty\} \subseteq \Theta(n). \quad (4)$$

Notice that  $\theta(\vec{g}(p, q)) = \text{LCM}(p, q)$ ,  $\theta(\vec{C}_n) = \infty$  and  $\theta(\vec{K}_n) = 1$ , where  $\vec{K}_n$  is the complete digraph of order  $n$ . It suffices to verify

$$[2, s(n - 1) + 1] \subseteq \Theta(n). \quad (5)$$

Let  $G(p, q)$  be the set of graphs with the following diagrams, where  $\vec{C}_p$  is a  $p$ -cycle  $u_1u_2 \cdots u_pu_1$ ,  $\vec{C}_q$  is a  $q$ -cycle  $v_1v_2 \cdots v_qv_1$ , the arrows from  $x$  to  $\vec{C}_p$  and  $\vec{C}_q$  represents two paths  $xx_1 \cdots x_ru_1$  and  $xy_1 \cdots y_sv_1$ , the arrow from  $\vec{C}_p$  to  $\vec{C}_q$  represents an arc  $u_iv_j$  with  $1 \leq i \leq p$ ,  $1 \leq j \leq q$ .



Let  $G(p, q, l, t)$  be the set of graphs in  $G(p, q)$  such that the length of the path  $xx_1 \cdots x_r u_1 \cdots u_i v_j$  is  $l$  and the length of the path  $xy_1 \cdots y_s v_1 \cdots v_j$  is  $t$ . Then

$$\theta(D) = \min\{l + ap : l + ap = t + bq \text{ for some nonnegative integers } a, b\} - 1$$

for all  $D \in G(p, q, l, t)$ . We denote by  $\theta(G(p, q, l, t))$  the stable index of digraphs in  $G(p, q, l, t)$ .

If  $q = p - 1$  and  $l - t = d < p$ , applying Lemma 7,

$$\begin{aligned} & \min\{l + ap : l + ap = t + bq \text{ for some nonnegative integers } a, b\} \\ &= \min\{l + ap : l - t = bq - ap \text{ for some nonnegative integers } a, b\} \end{aligned}$$

is attained at  $(a, b) = (p - d - 1, p - d)$ . Therefore,

$$\theta(G(p, p - 1, l, t)) = t - 1 + (p - 1)(p - d).$$

Denote by  $f(p, q, d) = \{\theta(G(p, q, l, t)) : l - t = d\}$ . If  $q = p - 1 \geq 1$  and  $d = 1$ , then we have  $1 \leq t \leq \lceil n/2 \rceil - 1$  in  $G(p, q, l, t)$ . Hence,

$$\begin{aligned} f(p, p - 1, 1) &= \{w + (p - 1)^2 : 0 \leq w \leq \lceil n/2 \rceil - 2\} \\ &= [(p - 1)^2, (p - 1)^2 + \lceil n/2 \rceil - 2]. \end{aligned}$$

If  $q = p - 1 \geq 2$  and  $d = 2$ , then we have  $1 \leq t \leq \lfloor n/2 \rfloor - 1$  in  $G(p, q, l, t)$ . Hence,

$$\begin{aligned} f(p, p - 1, 2) &= \{w + (p - 1)(p - 2) : 0 \leq w \leq \lfloor n/2 \rfloor - 2\} \\ &= [(p - 1)(p - 2), (p - 1)(p - 2) + \lfloor n/2 \rfloor - 2]. \end{aligned}$$

Notice that

$$\begin{aligned} (p - 1)^2 &\leq (p - 1)(p - 2) + \lfloor n/2 \rfloor - 1 \quad \text{for } 2 \leq p \leq \lfloor n/2 \rfloor, \\ (p - 1)(p - 2) &\leq (p - 2)^2 + \lceil n/2 \rceil - 1 \quad \text{for } 2 \leq p \leq \lceil n/2 \rceil. \end{aligned}$$



We have

$$\bigcup_{p=3}^{\lfloor n/2 \rfloor} \{f(p, p-1, 1) \cup f(p, p-1, 2)\} = [2, (\lfloor n/2 \rfloor - 1)^2 + \lceil n/2 \rceil - 2] \equiv T(n),$$

which implies  $T(n) \subseteq \Theta(n)$ .

Next we distinguish three cases.

*Case 1.*  $n$  is even, say,  $n = 2m$ . Then

$$(\lfloor n/2 \rfloor - 1)^2 + \lceil n/2 \rceil - 2 = m^2 - m - 1$$

and

$$s(n-1) + 1 = (n^2 - 2n + 4)/4 = m^2 - m + 1.$$

Note that

$$\theta(\vec{g}(m, m-1)) = m^2 - m \quad \text{and} \quad \theta(\vec{g}(m, 3, m-1)) = m^2 - m + 1.$$

We have (5).

*Case 2.*  $n \equiv 1 \pmod{4}$ , say,  $n = 4k + 1$ . Suppose  $m = 2k$ . Then

$$(\lfloor n/2 \rfloor - 1)^2 + \lceil n/2 \rceil - 2 = m^2 - m$$

and

$$s(n-1) + 1 = (n^2 - 2n + 1)/4 = m^2.$$

Since  $m+1$  and  $m-1$  are relatively prime, applying Corollary 8, we have

$$f(m+1, m-1, 2) = \{w + m(m-1) : 0 \leq w \leq m-2\} = [m^2 - m, m^2 - 2].$$

Moreover, since  $m$  is even, we have

$$\theta(\vec{g}(m+1, m-1)) = m^2 - 1 \quad \text{and} \quad \theta(\vec{g}(m+1, 3, m-1)) = m^2 = s(n-1) + 1.$$

Therefore, we have (5).

*Case 3.*  $n \equiv 3 \pmod{4}$ , say,  $n = 4k + 3$ . Suppose  $m = 2k + 1$ . Then

$$(\lfloor n/2 \rfloor - 1)^2 + \lceil n/2 \rceil - 2 = m^2 - m$$

and

$$s(n-1) + 1 = (n^2 - 2n - 11)/4 = m^2 - 3.$$

Since  $m + 2$  and  $m - 2$  are relatively prime, applying Corollary 8 we have

$$f(m + 2, m - 2, 4) = \{w + (m + 1)(m - 2) : 0 \leq w \leq m - 3\} = [m^2 - m - 2, m^2 - 5].$$

On the other hand,

$$\theta(\vec{g}(m + 2, m - 2)) = m^2 - 4 \quad \text{and} \quad \theta(\vec{g}(m + 2, 3, m - 2)) = m^2 - 3 = s(n - 1) + 1.$$

Therefore, we get (5).

Combining (2) and (4), we obtain

$$\Theta(n) = [s(n - 1) + 1] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\} \cup \{\infty\}.$$

This completes the proof. □

*Remark.* Denote by  $\vec{L}_n$  the union of the  $n$ -cycle  $v_1v_2 \cdots v_nv_1$  and the arc  $v_1v_n$ . Then we have

$$\theta(\vec{L}_n) = n \quad \text{for} \quad n \geq 3.$$

Note that

$$\theta(\vec{K}_n) = 1, \theta(\vec{g}(1, 2)) = 2, \theta(\vec{g}(2, 3)) = 6, \theta(\vec{g}(2, 3, 3)) = 7.$$

We have

$$\Theta(n) = [s(n)] \cup \{\infty\} \quad \text{for} \quad 1 \leq n \leq 6.$$

When  $n \geq 7$ , by direct computation we see that  $\Theta(10) = [s(10)] \cup \{\infty\}$  and

$$[s(n - 1) + 1] \cup \{\text{LCM}(p, q) : p + q = n, p, q \in \mathbb{Z}^+\} \neq [s(n)] \quad \text{for} \quad n \neq 10.$$

So  $\Theta(n)$  has gaps in the set  $[s(n)]$  for  $n \in \mathbb{Z}^+ - [6] \cup \{10\}$ .

## Acknowledgement

This work was supported by the National Natural Science Foundation of China (No. 12171323) and the Science and Technology Foundation of Shenzhen City (No. JCYJ20190808174211224, No. JCYJ20210324095813036). The authors are grateful to Professor Xingzhi Zhan for valuable suggestions.

## References

- [1] J.A. Bondy and U.S.R. Murty, Graph Theory, GTM 244, Springer, 2008.
- [2] Z. Chen, Z. Huang, J. Yan, The stable index of 0-1 matrices, Linear Algebra Appl. 600 (2020) 148-160.
- [3] A.L. Dulmage, N.S. Mendelsohn, Gaps in the exponent set of primitive matrices, Illinois J. Math. 8 (1964) 642-656.
- [4] M. Lewin, Y. Vitek, A system of gaps in the exponent set of primitive matrices, Illinois J. Math. 25 (1981) 87-98.
- [5] J.-Y. Shao, On a conjecture about the exponent set of primitive matrices, Linear Algebra Appl. 65 (1985) 91-123.
- [6] H. Wielandt, Unzerlegbare nicht negative matrizen, Math. Z. 52 (1950) 642-648.
- [7] K.M. Zhang, On Lewin and Vitek's conjecture about the exponent set of primitive matrices, Linear Algebra Appl. 96 (1987) 101-108.