

IRREGULAR TRIADS IN 3-UNIFORM HYPERGRAPHS

C. TERRY AND J. WOLF

ABSTRACT. Over the past several years, numerous authors have explored model theoretically motivated combinatorial conditions that ensure that a graph has an efficient regular decomposition in the sense of Szemerédi. In this paper we set out a research program that explores a corresponding set of questions for 3-uniform hypergraphs, a setting in which useful notions of regularity are significantly more intricate.

The main results in this paper concern certain combinatorial properties which arose as natural higher-order generalizations of the order property in parallel work of the authors in the arithmetic setting. Interpreted in the context of 3-uniform hypergraphs, these are tightly connected to the nature of irregular triads. Specifically, we show that a hereditary 3-graph property admits regular decompositions with so-called “linear error” if and only if it does not have the *functional order property* (FOP_2).

Along the way, we show that a hereditary 3-graph property is homogeneous (i.e. all regular triads have density near 0 or near 1) if and only if it has bounded VC_2 -dimension, complementing a recent (non-quantitative) result of Chernikov and Towsner.

We also address several questions arising from prior work on tame regularity in hypergraphs. In particular, we characterize the hereditary properties of 3-uniform hypergraphs admitting the type of regular partitions appearing in work of Fox et al. as those that have finite weak VC -dimension. This is again analogous to a recent non-quantitative result of Chernikov and Towsner.

CONTENTS

1. Introduction	2
Acknowledgements	7
2. Formal statements of problems and results	7
2.1. Quasirandomness and regularity in 3-uniform hypergraphs	8
2.2. Statements of problems	11
2.3. Statements of results	13
2.4. Outline of paper	24
3. Preliminaries	25
3.1. Notions of quasirandomness for graphs and hypergraphs	27
3.2. Lemmas for slicing, intersecting, and combining decompositions	32
3.3. Ramsey facts	36
4. VC_2 -dimension and $\text{disc}_{2,3}$ -homogeneity	37
5. Weak NIP, weak stability, and vdisc_3 -homogeneity	40
5.1. The \sim -classes of weakly stable and weakly NIP properties	41
5.2. Weak VC -dimension and vdisc_3 -homogeneity	42

5.3. Weak stability and non-zero error	47
5.4. A generalized strong stable regularity lemma	50
5.5. Weakly stable properties admit binary error	58
6. The functional order property and linear $\text{disc}_{2,3}$ -error	64
6.1. Encodings and recharacterizations of FOP_2	65
6.2. Removal in almost stable bipartite graphs	70
6.3. A property admits linear $\text{disc}_{2,3}$ -error if and only if it is NFOP_2	74
7. Binary $\text{disc}_{2,3}$ -error	89
7.1. Admitting binary $\text{disc}_{2,3}$ -error is distinct from admitting zero vdisc_3 -error	89
7.2. Proof of Proposition 2.57	92
7.3. Special 3-graphs and the proof of Theorems 2.53 and 2.56	93
7.4. Proof of Theorems 2.53 and 2.56	105
Appendix A. Facts about \sim -classes	126
Appendix B. Lemmas about combining and refining decompositions	130
Appendix C. Ramsey facts	142
Appendix D. General properties of NFOP_2 -formulas	144
References	149

1. INTRODUCTION

Szemerédi's regularity lemma [54] is a theorem with a wide range of applications in graph theory. Roughly speaking, it says that any large graph can be partitioned into a small number of pieces, so that in between most pairs of pieces the graph behaves like a random graph. More precisely, Szemerédi's regularity lemma states that for all $\epsilon > 0$ there is some $M = M(\epsilon)$ so that for every sufficiently large graph $G = (V, E)$, there is some $m \leq M$ and an equipartition of $V = V_1 \cup \dots \cup V_m$ such that all but at most ϵm^2 pairs (V_i, V_j) are ϵ -regular (see Definition 3.2). In order to utilize such a regular decomposition, one needs a so-called *counting lemma*, which says that the number of triangles appearing in a triple of ϵ -regular pairs is about what one would expect in a 3-partite random graph of the same density. An important early application of the counting lemma was the so-called *triangle removal lemma* [20], which says that a graph with a small number of triangles can be made triangle free by removing a small number of edges. We refer the reader to the excellent surveys [10, 32] for more details.

Despite its myriad applications, the regularity lemma is limited by the potential existence of irregular pairs, as well as the enormous size of the bound $M(\epsilon)$, which appears as a tower-type function in Szemerédi's proof. Investigation into these limitations showed them to be necessary. Indeed, the folklore example of the half-graph showed the necessity of the irregular pairs, while a construction of Gowers [25] proved the necessity of tower-type bounds.

On the other hand, under certain assumptions, strengthened versions of the regularity lemma can be proved. For instance, several authors [3, 34] showed that graphs of bounded VC-dimension allow for regular decompositions with very good bounds. These results yield

a structural dichotomy which is best stated in terms of hereditary graph properties (see Theorem 1.1 below). We require a few definitions to state this result.

A *hereditary graph property* is a class \mathcal{H} of finite graphs which is closed under isomorphism and induced subgraphs. The VC-dimension of a graph $G = (V, E)$ is the largest k for which there exist $a_1, \dots, a_k \in V$, and $b_S \in V$ for each $S \subseteq [k]$, such that $a_i b_S \in E$ if and only if $i \in S$. We then say that a hereditary graph property \mathcal{H} has *VC-dimension at most k* if every $G \in \mathcal{H}$ has $VC(G) \leq k$, and \mathcal{H} has *bounded VC-dimension* (or is NIP) if it has VC-dimension at most k for some k . Otherwise, we say \mathcal{H} has *unbounded VC-dimension* (or IP). Theorem 1.1 sums up the work of Alon, Fischer and Newman [3] (part (1)) and Gowers [25] (part (2)), where the bounds quoted are improvements due to Fox et al. [21] (part (1)) and Moshkovitz and Shapira [37] (part (2)).

Theorem 1.1. *Suppose \mathcal{H} is a hereditary graph property. Then one of the following holds.*

- (1) \mathcal{H} has bounded VC-dimension. In this case, if $k = \max\{VC(G) : G \in \mathcal{H}\}$, then for all $\epsilon > 0$, any $G \in \mathcal{H}$ has a regular partition with $O(\epsilon^{-k})$ parts, and where moreover, all regular pairs have density in $[0, \epsilon) \cup (1 - \epsilon, 1]$.
- (2) \mathcal{H} has unbounded VC-dimension. In this case, there exists $\epsilon > 0$ and arbitrarily large $G \in \mathcal{H}$, such that any ϵ -regular partition of G has at least $Tw(\epsilon^{-1/6})$ parts.

A similar dichotomy exists with regards to irregular pairs. A graph $G = (V, E)$ is said to have the *k -order property* if there exist $a_1, \dots, a_k \in V$ and $b_1, \dots, b_k \in V$ so that $a_i b_j \in E$ if and only if $i \leq j$. If G does not have the k -order property, then we say G is *k -stable*. A hereditary graph property \mathcal{H} is called *k -stable* if every $G \in \mathcal{H}$ is k -stable, and \mathcal{H} is *stable* if it is k -stable for some k . Otherwise, \mathcal{H} is called *unstable*.

While it was known since the 1990s [32] that a sufficiently large instance of the order property forces the existence of irregular pairs (part (2) below), it was not until 2011 when Malliaris and Shelah [36] showed that the converse (part (1) below) also holds.

Theorem 1.2. *Suppose \mathcal{H} is a hereditary graph property. Then one of the following holds.*

- (1) \mathcal{H} is stable. In this case, the following holds, with k minimal such that \mathcal{H} is k -stable. For all $\epsilon > 0$, every sufficiently large $G \in \mathcal{H}$ has an equipartition into $O(\epsilon^{-2^{2^k}})$ parts, such that each pair is ϵ -regular with density in $[0, \epsilon) \cup (1 - \epsilon, 1]$.
- (2) \mathcal{H} is unstable. In this case, for all $\epsilon > 0$ and M there are arbitrarily large $G \in \mathcal{H}$ such that every partition of G into at most M parts contains $\Omega(M)$ irregular pairs.

The notions of VC-dimension and stability are both central to model theory, where proving structural results under these and related tameness assumptions is a major theme. We can think of Theorems 1.1 and 1.2 as finitary versions of such results. In particular, any hereditary graph property \mathcal{H} is the class of finite models of a universal theory, $T_{\mathcal{H}}$, in the language of graphs. Therefore, Theorems 1.1 and 1.2 give structural dichotomies for such universal theories, based on whether they are of bounded VC-dimension or, respectively, stable.

Following the work of Malliaris and Shelah, several results appeared that illustrate how certain model theoretic notions yield different kinds of “tame” regularity lemmas [1, 13–15].

During this time, the field also gained an understanding of theorems in the same spirit proved earlier and/or by researchers outside of model theory (e.g. [3, 4, 9, 22, 34]). As a result, tame regularity lemmas are now well known as points of connection between model theory and combinatorics. Indeed, some of this work demonstrates explicit connections between upgraded regularity lemmas and model theoretic tools, such as compact domination, indiscernible sequences, and definability of types (see [36, 44, 45]). This demonstrates that in the setting of graphs, looking for dichotomies in the behavior of hereditary properties with respect to regular partitions leads directly to notions of primary importance in model theory.

Analogues of Szemerédi’s regularity lemma for hypergraphs were considered desirable for many years, with the earliest examples appearing in the late 1980s [31] and early 1990s [18]. This early version of hypergraph regularity involved partitioning the vertex set of a hypergraph so that most triples of vertex parts satisfy a certain quasirandomness condition with respect to the hyperedges. This type of regularity is now often referred to as *weak hypergraph regularity*.

In the early 2000s, more sophisticated notions of hypergraph regularity were conceived, which went hand in hand with powerful counting lemmas and increasingly general removal lemmas [24, 26, 27, 40, 41, 49, 50]. For a detailed history, we refer the reader to [39]. As in the case of graphs, these counting lemmas have found robust applications, including the first fully finitary proof of the multidimensional Szemerédi’s theorem [27] and other, more classical applications in extremal combinatorics [8]. We will refer to this type of regularity as *strong hypergraph regularity*. In the setting of a 3-uniform hypergraph $H = (V, E)$, a strong regular decomposition consists of a partition $\mathcal{P}_{\text{vert}} = \{V_1, \dots, V_t\}$ of the vertex set V , along with, for each $ij \in \binom{[t]}{2}$, a partition of the pairs $K_2[V_i, V_j] = \bigcup_{\alpha \leq \ell} P_{ij}^\alpha$, so that certain quasirandomness properties hold. The essential “unit” in such a regular decomposition is a so-called *triad*, which is a 3-partite graph of the form $(V_i \cup V_j \cup V_k, P_{ij}^\alpha \cup P_{ik}^\beta \cup P_{jk}^\gamma)$. The strong regularity lemma then roughly says that every large 3-uniform hypergraph has an equitable decomposition so that almost all triples of vertices are in a regular triad (for details, see Section 3.1).

Several results mentioned above [1, 14, 22] give improved *weak* regular partitions for hypergraphs under various tameness assumptions. The only previous treatment of strong hypergraph regularity and model theoretic tameness occurs in work of Chernikov and Towsner [15], where higher arity analogues of Theorem 1.1 are proved in the context of strong regularity, using infinitary techniques. The results in [15] are in terms of a higher arity analogue of VC-dimension called VC_k -dimension, first defined by Shelah [52, 53], and subsequently developed in [12], with applications to algebra appearing in [11, 29], and to combinatorics in [58].

Despite the extensive work on model theoretic regularity lemmas of the last ten years, the following basic problem remained open.

Problem 1.3. *Prove an analogue of Theorem 1.2 in the realm of strong regularity for hypergraphs.*

This is an interesting problem from a model theoretic perspective, as it is asking one to develop a higher arity analogue of stability, a problem which has been discussed in the model theory community for several years (see for example [53, 55]).

Question 1.4. *What higher arity notion generalizes stability, in the same way that VC_k -dimension generalizes VC -dimension?*

A convincing answer to Question 1.4 has also remained open. Given the important applications of strong hypergraph regularity in combinatorics, as well as its connections to arithmetic structure [27], one might reasonably expect an answer to Problem 1.3 to lead to the “right” answer to Question 1.4, as well as important new tools and ideas in model theory.

The focus of this paper is Problem 1.3 and Question 1.4 in the setting of 3-uniform hypergraphs, where our contributions are two-fold. First, we prove the first analogue of Theorem 1.2 in the realm of strong hypergraph regularity in terms of a new higher arity generalization of stability (see Theorem 1.7 below). Our work gives substantial evidence that this particular analogue of Theorem 1.2 is especially robust, and the generalization of stability we define is the “right” answer to Question 1.4.

Our second contribution is more complicated. We show that there are several distinct ways to formulate generalizations of Theorem 1.2 in the setting of strong regularity for 3-uniform hypergraphs, suggesting that Problem 1.3 does not have a single answer. We explore these distinct generalizations, related notions of higher arity stability, and specific examples. The sum of these results is the first significant contribution to understanding the full landscape of Problem 1.3. Our work teases apart several potential generalizations of stability via their connections to Problem 1.3, and the examples we give rule out certain natural ideas for how to generalize stability. Indeed, working out these subtle distinctions and uncovering the relevant examples were significant hurdles to correctly addressing Question 1.4, and we view this as a major contribution of this paper.

We will now give a more detailed overview of our results (for full details, see Section 2.2). We begin by taking a slightly different perspective on Theorem 1.2. A naive attempt to generalize Theorem 1.2 to the setting of 3-uniform hypergraphs would be to insist that there are *no* irregular triads. However, we will show that doing so reduces to a question about *weak* regularity, rather than strong regularity (see Theorem 2.46). Therefore, we start by giving a more flexible formulation of that Theorem 1.2, which will illuminate the proper generalization to higher arities. In particular, the following is an exercise which gives an equivalent formulation of part (1) of Theorem 1.2.

Fact 1.5. *The following are equivalent for a hereditary graph property \mathcal{H} .*

- (i) *For all $\epsilon > 0$, there is M so that for all $G = (V, E) \in \mathcal{H}$ with $|V| \geq M$, there is some $t \leq M$ and an equipartition $V = V_0 \cup V_1 \cup \dots \cup V_t$ such that $|V_0| \leq \epsilon|V|$ and for all $1 \leq i \neq j \leq t$, (V_i, V_j) is ϵ -regular.*
- (ii) *For all $\epsilon > 0$, there is M so that for all $G = (V, E) \in \mathcal{H}$ with $|V| \geq M$, there is some $t \leq M$ and an equipartition $V = V_1 \cup \dots \cup V_t$ such that for all $ij \in \binom{[t]}{2}$, (V_i, V_j) is ϵ -regular.*

That (ii) implies (i) is trivial. For the reverse direction, one chooses the parameters carefully in (i), then evenly distributes the error set V_0 among the other parts in the partition to obtain a partition as in (ii). Fact 1.5 shows that having no irregular pairs in a regular partition of a graph can be reinterpreted as saying that there is a single, small set in the vertex partition which is used in every irregular pair. Thinking of sets of vertices as “lower order” objects in a graph, this is informally saying one can use a lower order set to control the irregular pairs. It turns out that this simple reinterpretation is much more conducive to higher arity generalization. We thus reinterpret Problem 1.3 as follows.

Problem 1.6. *Characterize hereditary properties of 3-graphs which have regular partitions where the irregular triads are constrained by “lower order” sets.*

Problem 1.3 is made complicated by the fact that there are several distinct ways to constrain irregular triads using lower order sets. We define three natural ways of constraining the error triads in a regular decomposition, which we will refer to as having zero, binary, and linear $\text{disc}_{2,3}$ -error. We give rough definitions of these here and refer the reader to Definition 2.11 for details. Given a 3-uniform hypergraph $H = (V, E)$ and a decomposition \mathcal{P} with $\mathcal{P}_{\text{vert}} = \{V_1, \dots, V_t\}$, we say that

- (1) \mathcal{P} has *zero* $\text{disc}_{2,3}$ -error if there are no irregular triads in \mathcal{P} with respect to H ;
- (2) \mathcal{P} has *binary* $\text{disc}_{2,3}$ -error if there is a small set $\Sigma \subseteq \binom{[t]}{2}$ such that every irregular triad from \mathcal{P} is contained in some $V_i V_j V_k$ with one of ij , jk , or ik in Σ ; and
- (3) \mathcal{P} has *linear* $\text{disc}_{2,3}$ -error if there is a small set $\Sigma \subseteq \binom{[t]}{3}$ such that every irregular triad from \mathcal{P} is contained in some $V_i V_j V_k$ with $ijk \in \Sigma$.

We then say a hereditary 3-graph property \mathcal{H} admits zero/binary/linear $\text{disc}_{2,3}$ -error if all the hypergraphs in \mathcal{H} have regular decompositions with zero/binary/linear $\text{disc}_{2,3}$ -error, respectively.

Our main theorem provides a complete characterization of the properties which admit linear error in terms of a new generalization of the order property, which we call the *functional order property* (FOP_2 , see Definition 2.48). We state this result below and refer the reader to Section 2.2 for the precise definitions.

Theorem 1.7. *Suppose \mathcal{H} is a hereditary property of 3-uniform hypergraphs. Then the following are equivalent.*

- (1) *There is some k so that no $H \in \mathcal{H}$ has the k -functional order property.*
- (2) *For all $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$, there are T, L, N such that the following hold. For all $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there are $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and \mathcal{P} a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and which has linear $\text{disc}_{2,3}$ -error with respect to H .*

This is the first analogue of Theorem 1.2 to be proved in the setting of strong hypergraph regularity. Our companion results in the arithmetic setting [60] show a correspondence between forbidding the functional order property in groups, and quadratic arithmetic regularity lemmas where the “error part” is supported on a small number of *linear* atoms. These results show that this particular generalization of Theorem 1.2 has strong connections to

algebraic structure. Our proof techniques in this paper will also show that the relationship of the functional order property to stability is analogous to that between VC_2 -dimension and VC-dimension (see Section 6.1).

Our other main results focus on zero and binary $\text{disc}_{2,3}$ -error, and their relationships to various examples and possible generalizations of stability. We define the notion of *weak stability* (Definition 2.40) and show that a hereditary property which is close to being weakly stable admits zero error (Proposition 2.43). We conjecture that the converse holds as well. We also show that the notion of zero error reduces to a problem about weak regularity (Theorem 2.46), and cannot be characterized by stability, as defined in [1, 14]. We show that being close to weakly stable implies that a property admits binary error, but that the converse to this statement is false.

We give two distinct examples of 3-uniform hypergraphs which do not admit regular decompositions with binary error. The first arises from a combinatorial definition called the *hyperplane order property* (HOP_2), and the other from an example of Green and Sanders [28] in elementary abelian p -groups. Based on these examples, we prove a general sufficient condition for when a property cannot admit binary error (Section 7).

The hyperplane order property has a strong geometric appeal as a potential generalization of the order property. The existence of the second example in elementary abelian p -groups, however, shows that HOP_2 is unlikely to be a fundamental definition generalizing the order property. We consider this observation crucial for the future investigation of this sphere of problems.

Our results imply that the notions of zero, binary, and linear error are pairwise distinct. Further, we show that binary error cannot be characterized by stability or weak stability, and cross-cuts the theories of bounded VC-dimension.

The main results of this paper also require several auxiliary results, some of which are of independent interest. These include finitary versions of the results of Chernikov and Towsner [15] for 3-uniform hypergraphs (see Theorems 2.23 and 2.32), as well as a strong form of the stable graph regularity lemma (see Theorem 5.7).

Acknowledgements. The first author would like to thank M. Malliaris for conversations about stable graphs which informed several proofs in this paper. The first author would also like to thank G. Conant for many helpful discussions around the paper's main results. The first author is partially supported by NSF grant DMS-2115518. The authors' collaboration has been supported by several travel grants over the years, including by the London Mathematical Society, the Simons Foundation, and the Association for Women in Mathematics.

2. FORMAL STATEMENTS OF PROBLEMS AND RESULTS

We state our results in full detail in this section (see Section 2.2). This requires the definitions of Section 3.1. The reader familiar with these definitions may jump ahead to Section 2.2.

In the following subsections, we will frequently include translations to model theoretic language to aid the model theoretic reader. Any such undefined terminology is not necessary for the rest of the paper. To facilitate these explanations, we will use the fact that every hereditary 3-graph property \mathcal{H} is the class of finite models of a universal theory $T_{\mathcal{H}}$ in the language $\mathcal{L} = \{R(x, y, z)\}$. Further, we remind the model theoretic reader that given a 3-graph $H = (V, E)$, an induced sub-3-graph of H is simply a substructure of H in the language \mathcal{L} .

Throughout, we refer to 3-uniform hypergraphs as simply *3-graphs*. A *hereditary 3-graph property* is a class of finite 3-graphs closed under induced sub-3-graphs and isomorphism.

2.1. Quasirandomness and regularity in 3-uniform hypergraphs. In this subsection we introduce two notions of quasirandomness for 3-graphs. We largely follow the notation of [24, 39], with some modifications. We begin by recalling the definition of quasirandomness for graphs. For sets V_1, V_2 , we let $K_2[V_1, V_2] = \{xy : x \in V_1, y \in V_2, x \neq y\}$.

Definition 2.1 (disc_2). Suppose $\epsilon > 0$ and $G = (U \cup V, E)$ is a bipartite graph with $|E| = d|U||V|$. We say G has $\text{disc}_2(\epsilon)$ if for all $U' \subseteq U$ and $V' \subseteq V$,

$$||E \cap K_2[U', V']| - d|U'||V'|| \leq \epsilon|U||V|.$$

We say that G has $\text{disc}_2(\epsilon, d')$ if it has $\text{disc}_2(\epsilon)$ and $||E| - d'|U||V|| \leq \epsilon|U||V|$ (here “disc” is short for discrepancy).

The notion of disc_2 -quasirandomness is essentially equivalent to regularity and several other notions of quasirandomness in graphs. We will present some of these in more detail in Section 3.1. Turning to 3-graphs, the most naive generalization of disc_2 yields Definition 2.2 below. Given sets V_1, V_2, V_3 , we let

$$K_3[V_1, V_2, V_3] = \{uvw : u \in V_1, v \in V_2, w \in V_3, u \neq v, u \neq w, v \neq w\}.$$

Definition 2.2 (vdisc_3). Suppose $\epsilon > 0$ and $H = (U \cup V \cup W, E)$ is a 3-partite, 3-graph with $|E| = d|U||V||W|$. We say H has $\text{vdisc}_3(\epsilon)$ if for all $U' \subseteq U$, $V' \subseteq V$, and $W' \subseteq W$,

$$||E \cap K_3[U', V', W']| - d|U'||V'||W'|| \leq \epsilon|U||V||W|.$$

The “v” in vdisc is meant to emphasize that this definition considers quasirandomness with respect to vertex subsets only. This notion of quasirandomness was introduced in the early 1990s [17] as a generalization of quasirandomness for graphs. It has a corresponding regularity lemma, first proved by Chung [17] and stated as Theorem 2.4 below.

Definition 2.3 (vdisc_3 -regular triples and partitions). Suppose $H = (V, E)$ is a 3-graph and $\mathcal{P} = \{V_1, \dots, V_t\}$ is a partition of V .

- (1) Given $V_i V_j V_k \in \binom{\mathcal{P}}{3}$, we say $V_i V_j V_k$ has $\text{vdisc}_3(\epsilon)$ with respect to H if the 3-partite 3-graph $(V_i \cup V_j \cup V_k, E \cap K_3[V_i, V_j, V_k])$ has $\text{vdisc}_3(\epsilon)$.
- (2) We say \mathcal{P} is a $\text{vdisc}_3(\epsilon)$ -regular partition for H if
 - (a) \mathcal{P} is an equipartition, and
 - (b) for all but at most $\epsilon \binom{t}{3}$ many $ijk \in \binom{[t]}{3}$, $V_i V_j V_k$ has $\text{vdisc}_3(\epsilon)$ with respect to H .

Theorem 2.4 (vdisc₃-regularity lemma for 3-graphs). *For all integers $t_0 \geq 1$ and $\epsilon > 0$ there exists $T = T(t_0, \epsilon)$ and $N = N(t_0, \epsilon)$ such that the following holds. If $H = (V, E)$ is a 3-uniform hypergraph with $|V| \geq N$, there is some $t_0 \leq t \leq T$ and a vdisc₃(ϵ)-regular partition for H with t parts.*

It was well known that vdisc₃-quasirandomness does not allow for a general counting lemma. It was shown later [30] that it does allow for counting copies of *linear* hypergraphs (see also [42] for an application of this counting lemma). A general counting lemma was the motivation for a more sophisticated notion of quasirandomness [24, 26], which considers not only sets of vertices but also sets of pairs of vertices. Given a graph $G = (V, E)$, we let $K_3^{(2)}(G)$ denote the set of triples from V forming a triangle in G . In other words,

$$K_3^{(2)}(G) = \left\{ xyz \in \binom{V}{3} : xy, yz, xz \in E \right\}.$$

Given a 3-graph $H = (V, R)$ on the same vertex set, we say that G *underlies* H if $R \subseteq K_3^{(2)}(G)$, i.e. if all the edges of H sit atop a triangle from G . A subgraph $G' \subseteq G$ is a graph (V', E') where $V' \subseteq V$ and $E' \subseteq E$. Our next definition describes when a 3-graph is quasirandom relative to an underlying quasirandom graph.

Definition 2.5 (disc_{2,3}). Let $\epsilon_1, \epsilon_2 > 0$, and suppose $H = (V, E)$ is a 3-graph. Assume $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph underlying H , and suppose d_3 is defined by $|E| = d_3 |K_3^{(2)}(G)|$. We say that (H, G) has disc_{2,3}(ϵ_1, ϵ_2) if there is $d_2 \in (0, 1)$ such that for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has disc₂(ϵ_2, d_2), and for every subgraph $G' \subseteq G$,

$$||E \cap K_3^{(2)}(G')| - d_3 |K_3^{(2)}(G')|| \leq \epsilon_1 d_2^3 |V_1| |V_2| |V_3|.$$

Our notational convention is to use single subscripts of 2 or 3 for notions of strictly binary or strictly ternary quasirandomness (e.g. disc₂ and vdisc₃), and the subscript 2,3 as above to emphasize that disc_{2,3} quasirandomness has both a binary and ternary quasirandomness requirement. Definition 2.5 has several equivalent formulations, and a more complete account of these will appear in Section 3.1. We now state the definition which will serve as a higher order analogue of an equitable partition of a vertex set. In contrast to the partition appearing in Theorem 2.4, Definition 2.6 will include a partition of the vertex set as well as a partition of the set of pairs of vertices.

Definition 2.6 ($((t, \ell, \epsilon_1, \epsilon_2)$ -decomposition). Let V be a vertex set, $t, \ell \in \mathbb{N}^{>0}$, and $\epsilon_1, \epsilon_2 > 0$. A $((t, \ell, \epsilon_1, \epsilon_2)$ -decomposition \mathcal{P} for V consists of a partition $V = V_1 \cup \dots \cup V_t$ and for each $1 \leq i \neq j \leq t$, a collection $P_{ij}^1, \dots, P_{ij}^\ell$ of disjoint subsets of $\binom{V}{2}$ such that the following hold.

- (1) $|V_1| \leq \dots \leq |V_t| \leq |V_1| + 1$,
- (2) For each $1 \leq i \neq j \leq t$, $K_2[V_i, V_j] = \bigcup_{\alpha \leq \ell} P_{ij}^\alpha$,
- (3) For all but $\epsilon_1 |V|^3$ triples $xyz \in \binom{V}{3}$, the following holds. There is some $ijk \in \binom{[t]}{3}$ and $1 \leq \alpha, \beta, \gamma \leq \ell$ such that $xy \in P_{ij}^\alpha$, $xz \in P_{ik}^\beta$, $yz \in P_{jk}^\gamma$, and such that each of $P_{ij}^\alpha, P_{ik}^\beta, P_{jk}^\gamma$ have disc₂($\epsilon_2, 1/\ell$).

In the notation of Definition 2.6, a *triad* of \mathcal{P} is a 3-partite graph of the form

$$G_{ijk}^{\alpha,\beta,\gamma} := (V_i \cup V_j \cup V_k, P_{ij}^\alpha \cup P_{ik}^\beta \cup P_{jk}^\gamma)$$

for some $ijk \in \binom{[t]}{3}$ and $\alpha, \beta, \gamma \leq \ell$.

To ease notation, we define $\text{Triads}(\mathcal{P}) := \{G_{ijk}^{\alpha,\beta,\gamma} : ijk \in \binom{[t]}{3}, \alpha, \beta, \gamma \in [\ell]\}$. When $H = (V, E)$ is a 3-graph, then for each triad $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$, we write

$$H_{ijk}^{\alpha,\beta,\gamma} := (V_i \cup V_j \cup V_k, E \cap K_3^{(2)}(G_{ijk}^{\alpha,\beta,\gamma})).$$

Note that by definition, $H_{ijk}^{\alpha,\beta,\gamma}$ is a 3-partite 3-graph underlied by $G_{ijk}^{\alpha,\beta,\gamma}$. We now define regular triads and decompositions, in analogy with Definition 2.6.

Definition 2.7 ($\text{disc}_{2,3}$ -regular triads and decompositions). Suppose $H = (V, E)$ is a 3-graph and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition for V .

- (1) We say $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H if $(H_{ijk}^{\alpha,\beta,\gamma}, G_{ijk}^{\alpha,\beta,\gamma})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$.
- (2) We say that \mathcal{P} is $(\epsilon_1, \epsilon_2(\ell))$ -regular for H if for all but $\epsilon_1 n^3$ many triples $xyz \in \binom{V}{3}$, there is some $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$, so that $xyz \in K_3^{(2)}(G_{ijk}^{\alpha,\beta,\gamma})$ and which has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H .

Definition 2.7, along with its equivalent formulations (see Section 3.1), are now considered the standard notions of regular triads and decompositions for 3-graphs (see [26, 27, 39, 49]). We now state the regularity lemma for $\text{disc}_{2,3}$ -quasirandomness, which is taken from [24].

Theorem 2.8 ($\text{disc}_{2,3}$ -regularity lemma for 3-graphs). *For all $\epsilon_1 > 0$, every function $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and every $t_0 \in \mathbb{N}$, there exist positive integers T_0 and n_0 such that for any 3-graph $H = (V, E)$ on $n \geq n_0$ vertices, there exists a $(\epsilon_1, \epsilon_2(\ell))$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} for H with $t_0 \leq t \leq T_0$ and $\ell \leq T_0$.*

We end this section by setting up some terminology around irregular triads in the sense of Definition 2.6.

Definition 2.9 (Irregular triads). Suppose $H = (V, E)$ is a 3-uniform hypergraph and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition for V .

- (1) We say $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$ is $\text{disc}_{2,3}$ -regular with respect to (H, \mathcal{P}) if it has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H . Otherwise, we say $G_{ijk}^{\alpha,\beta,\gamma}$ is $\text{disc}_{2,3}$ -irregular.
- (2) We say $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$ is disc_2 -regular with respect to (H, \mathcal{P}) if each of $P_{ij}^\alpha, P_{ik}^\beta, P_{jk}^\gamma$ have $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Otherwise, we say $G_{ijk}^{\alpha,\beta,\gamma}$ is disc_2 -irregular.
- (3) We say $G_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$ is disc_3 -irregular with respect to (H, \mathcal{P}) if it is disc_2 -regular but not $\text{disc}_{2,3}$ -regular with respect to (H, \mathcal{P}) .

Note that in the notation of Definition 2.9, $G_{ijk}^{\alpha,\beta,\gamma}$ is $\text{disc}_{2,3}$ -irregular with respect to (H, \mathcal{P}) if and only if it is disc_2 -irregular with respect to (H, \mathcal{P}) or disc_3 -irregular with respect to (H, \mathcal{P}) .

There are several other notions of quasirandomness for hypergraphs which we do not consider (see [33, 61]). Our main motivation in this paper is to consider $\text{disc}_{2,3}$ -regularity, due to the important applications of its corresponding counting lemma. As our results will show, one of our main problems related to $\text{disc}_{2,3}$ (namely that centering around “zero $\text{disc}_{2,3}$ -error”) will in fact collapse to a problem about vdisc_3 (see Theorem 2.46). For this reason, we will also be considering vdisc_3 -regularity.

2.2. Statements of problems. In this subsection we state formally the problems of interest in this paper. We begin with two basic definitions in the case of vertex-regularity.

Definition 2.10 (zero and binary vdisc_3 -error). Let \mathcal{H} be a hereditary 3-graph property.

- (1) We say \mathcal{H} *admits zero vdisc_3 -error*, if for all $\epsilon > 0$ and $t_0 \geq 1$, there is $T = T(\epsilon, t_0)$ and $N = N(\epsilon, t_0)$ such that for all $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there is an equipartition \mathcal{P} of V with some $t_0 \leq t \leq T$ many parts, such that every triple $XYZ \in \binom{\mathcal{P}}{3}$ has $\text{vdisc}_3(\epsilon)$ with respect to H .
- (2) We say \mathcal{H} *admits binary vdisc_3 -error*, if for all $\epsilon > 0$ and $t_0 \geq 1$, there is $T = T(\epsilon, t_0)$ and $N = N(\epsilon, t_0)$ such that for all $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there is an equipartition \mathcal{P} of V with some $t_0 \leq t \leq T$ many parts, along with a set $\Sigma \in \binom{\mathcal{P}}{2}$ of size at most ϵt^2 , such that every triple $XYZ \in \binom{\mathcal{P}}{3}$ with $XY, YZ, XZ \notin \Sigma$ has $\text{vdisc}_3(\epsilon)$ with respect to H .

When \mathcal{H} fails (1) (respectively, (2)), we say it *requires non-zero (respectively, non-binary) vdisc_3 -error*. Clearly, if \mathcal{H} satisfies (1), it also satisfies (2). We now define similar notions for $\text{disc}_{2,3}$ -regularity.

Definition 2.11 (zero, binary, and linear $\text{disc}_{2,3}$ -error). Suppose \mathcal{H} is a hereditary 3-graph property.

- (1) We say \mathcal{H} *admits zero $\text{disc}_{2,3}$ -error* if for all $t_0, \ell_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are N, T, L such that for every $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there exists $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V , such that every triad in $\text{Triads}(\mathcal{P})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H .
- (2) We say \mathcal{H} *admits binary $\text{disc}_{2,3}$ -error* if for all $t_0, \ell_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are N, T, L such that for every $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there exists $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V such that the following holds. There is a set $\Gamma \subseteq \binom{[t]}{2}$ with $|\Gamma| \leq \epsilon_1 t^2$, such that for every $G_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ with $ij, ik, jk \notin \Gamma$, $G_{ijk}^{\alpha, \beta, \gamma}$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H .
- (3) We say \mathcal{H} *admits linear¹ $\text{disc}_{2,3}$ -error* if for all $t_0, \ell_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are N, T, L such that for every $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there exists $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V such that the following holds. There is a set $\Sigma \subseteq \binom{[t]}{3}$ with $|\Sigma| \leq \epsilon_1 t^3$, such that for every $G_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ with $ijk \notin \Sigma$, $G_{ijk}^{\alpha, \beta, \gamma}$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H .

¹The use of the term “linear” stems from the authors’ work in the arithmetic context, where this type of error set arises from the purely linear part of a quadratic factor, see [60, Corollary 4.30].

If \mathcal{H} fails (1) (respectively (2), (3)), we say \mathcal{H} *requires non-zero (respectively non-binary, non-linear) $\text{disc}_{2,3}$ -error*. By definition, if \mathcal{H} admits zero $\text{disc}_{2,3}$ -error, it also admits binary $\text{disc}_{2,3}$ -error and linear $\text{disc}_{2,3}$ -error. It is also not difficult to see that if \mathcal{H} admits binary $\text{disc}_{2,3}$ -error then it admits linear $\text{disc}_{2,3}$ -error (one just takes Σ to be the set of all triples with at least one pair in Γ). Thus (1) \Rightarrow (2) \Rightarrow (3). The reader may be wondering why the possibility of a linear error appears in Definition 2.11 but not in Definition 2.10. This is because Theorem 2.4 implies that *every* hereditary 3-graph property admits linear vdisc_3 -error.

One can restate parts (1) and (2) of Definition 2.11 in the rough form “sufficiently large elements in \mathcal{H} have regular decompositions with no irregular triads of a certain type.” It is clear that (1) is already in this form. Indeed, informally speaking, (1) says that sufficiently large elements in \mathcal{H} have $\text{disc}_{2,3}$ -regular decompositions with *no* $\text{disc}_{2,3}$ -irregular triads. It turns out that (2) is equivalent to saying that all sufficiently large elements in \mathcal{H} have $\text{disc}_{2,3}$ -regular decompositions with no disc_3 -irregular triads (see Definition 2.9). More specifically, we will prove the following in Section 7.

Theorem 2.12. *\mathcal{H} admits binary $\text{disc}_{2,3}$ -error if and only if for all $t_0, \ell_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are n, T, L such that for every $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there are $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$ and an $(\epsilon_1, \epsilon_2(\ell))$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} for H , such that no triad of \mathcal{P} is disc_3 -irregular with respect to H .*

On the other hand, we do not know of a way to restate (3) using a statement of the form “there are no irregular triads of a certain type.” It is for this reason that we transition away from thinking about “no irregular triads”, and instead aim to *control* irregular triads using lower order sets.

Finally, the reader may be wondering why none of these definitions address disc_2 -irregular triads. It turns out that these are simply not very interesting, since any $\text{disc}_{2,3}$ -regular partition \mathcal{P} can be modified so that it has no disc_2 -irregular triads. This is implicit in the work of Frankl and Rödl [24], but we will also include a proof for completeness. Specifically, we prove the following in Section 3.2.

Proposition 2.13. *For every hereditary 3-graph property \mathcal{H} the following holds. For all $\ell_0, t_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are n_0, T_0, L_0 such that for all $n \geq n_0$, every $G \in \mathcal{H}$ with at least n_0 vertices has an $(\epsilon_1, \epsilon_2(\ell))$ -regular $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition for some $t_0 \leq t \leq T_0$, and $\ell_0 \leq \ell \leq L_0$, such that every triad of \mathcal{P} is disc_2 -regular.*

Thus we are left with three potentially interesting notions with regards to $\text{disc}_{2,3}$ -irregular triads: zero, binary, and linear $\text{disc}_{2,3}$ -error. In summary then, the following are the problems concerning irregular triples/triads in 3-graphs which we will consider in this paper.

Problem 2.14. *Suppose \mathcal{H} is a hereditary 3-graph property.*

- (1) *Characterize when \mathcal{H} admits zero vdisc_3 -error.*
- (2) *Characterize when \mathcal{H} admits binary vdisc_3 -error.*
- (3) *Characterize when \mathcal{H} admits zero $\text{disc}_{2,3}$ -error.*

- (4) Characterize when \mathcal{H} admits binary $\text{disc}_{2,3}$ -error.
- (5) Characterize when \mathcal{H} admits linear $\text{disc}_{2,3}$ -error.

It will turn out that in order to address these problems, we must also consider the following general question, which is also motivated by Theorem 1.1: which hereditary 3-graph properties have regular decompositions where all regular triples/triads have density near 0 or 1? Following [22], we refer to such triads/triples as *homogeneous* (see Definition 2.16). What this question means specifically depends on the notion of regularity one considers. For vdisc_3 -regularity, we consider triples of sets (V_i, V_j, V_k) , while for $\text{vdisc}_{2,3}$ -regularity, we consider triads $(V_i \cup V_j \cup V_k, P_{ij}^\alpha \cup P_{jk}^\beta \cup P_{ik}^\gamma)$.

Definition 2.15. Suppose $H = (V, E)$ is a 3-graph with $|V| = n$ and $\mu > 0$.

- (1) Suppose \mathcal{P} is a partition of V . We say \mathcal{P} is μ -homogeneous with respect to H , if at least $(1 - \mu)\binom{n}{3}$ triples $xyz \in \binom{V}{3}$ satisfy the following. There is some $V_i V_j V_k \in \binom{\mathcal{P}}{3}$ such that $xyz \in K_3[V_i, V_j, V_k]$ and

$$|E \cap K_3[V_i, V_j, V_k]| / |K_3[V_i, V_j, V_k]| \in [0, \mu] \cup (1 - \mu, 1].$$

- (2) Suppose $t, \ell \geq 1$ and \mathcal{P} is a (t, ℓ) -decomposition of V . We say that \mathcal{P} is μ -homogeneous with respect to H if at least $(1 - \mu)\binom{n}{3}$ triples $xyz \in \binom{V}{3}$ satisfy the following. There is some triad $G_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ such that $xyz \in K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})$ and

$$|E \cap K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})| / |K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})| \in [0, \mu] \cup (1 - \mu, 1].$$

Definition 2.16 (vdisc_3 - and $\text{disc}_{2,3}$ -homogeneity). Suppose \mathcal{H} is a hereditary 3-graph property.

- (1) \mathcal{H} is vdisc_3 -homogeneous, if for all $\epsilon > 0$ and t_0 , there are $T = T(\epsilon, t_0)$ and $N = N(\epsilon, t_0)$ such that for all $H \in \mathcal{H}$ on at least N vertices, there is a partition \mathcal{P} of V which is $\text{vdisc}_3(\epsilon)$ -regular and ϵ -homogeneous with respect to H .
- (2) Call \mathcal{H} $\text{disc}_{2,3}$ -homogeneous, if for all $t_0, \ell_0 \geq 1$, $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are T, L, N such that for $H \in \mathcal{H}$ on at least N vertices, there are $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and ϵ_1 -homogeneous with respect to H .

We can now state concrete versions of our homogeneity problems.

Problem 2.17. Suppose \mathcal{H} is a hereditary 3-graph property.

- (1) Characterize when \mathcal{H} is vdisc_3 -homogeneous.
- (2) Characterize when \mathcal{H} is $\text{disc}_{2,3}$ -homogeneous.

2.3. Statements of results. In this section we state the main results of the paper. We will give full answers to Problem 2.17 and Problem 2.14 (5), and partial results for Problem 2.14 (1)-(4).

To state some of our results, we require a few more definitions. First, we can recharacterize stability and VC-dimension in terms of the bipartite counterpart of a graph. Given a graph $G = (V, E)$, let $\text{Bip}(G)$ denote the bipartite graph $(U \cup W, E')$, where $U = \{u_v :$

$v \in V\}$ and $W = \{w_v : v \in V\}$ are disjoint copies of V , and $E' = \{u_v w_{v'} : vv' \in E\}$. Note that the VC-dimension of a graph G is then the largest k such that $\text{Bip}(G)$ contains an induced copy of the k th power set graph,

$$U(k) = (\{a_i : i \in [k]\} \cup \{b_S : S \subseteq [k]\}, \{a_i b_S : i \in S\}).$$

Similarly, a graph G has the k -order property if and only if $\text{Bip}(G)$ contains an induced copy of the *half-graph of height k* ,

$$H(k) = (\{a_i : i \in [k]\} \cup \{b_j : j \in [k]\}, \{a_i b_j : i \leq j\}).$$

We will also use a natural equivalence relation on hereditary properties. Given two 3-graphs $H = (V, E)$ and $H' = (V, E')$ on the same vertex set, we say that H and H' are δ -close if $|E \Delta E'| \leq \delta |V|^3$.

Definition 2.18 (Closeness of hereditary properties). Suppose \mathcal{H} and \mathcal{H}' are hereditary 3-graph properties. We say \mathcal{H} is close to \mathcal{H}' if for all $\delta > 0$, there is $N = N(\delta)$ such that if $n \geq N$, then every $G \in \mathcal{H}$ on at least n vertices is δ -close to an element of \mathcal{H}' with the same vertex set.

We say \mathcal{H} and \mathcal{H}' are close, denoted $\mathcal{H} \sim \mathcal{H}'$, if \mathcal{H} is close to \mathcal{H}' , and \mathcal{H}' is close to \mathcal{H} .

This notion appears in several other places in the literature. For example it is referred to as indistinguishability in [2]. Clearly \sim is an equivalence relation. It turns out that the behavior of a hereditary property with respect to Problems 2.17 and 2.14 depends only on its \sim -equivalence class.

Proposition 2.19. *Suppose \mathcal{H} and \mathcal{H}' are close. Then the following are equivalent.*

- (1) \mathcal{H} is vdisc_3 -homogeneous (is $\text{disc}_{2,3}$ -homogeneous, admits zero/binary vdisc_3 -error, admits zero/binary/linear $\text{disc}_{2,3}$ -error).
- (2) \mathcal{H}' is vdisc_3 -homogeneous (is $\text{disc}_{2,3}$ -homogeneous, admits zero/binary vdisc_3 -error, admits zero/binary/linear $\text{disc}_{2,3}$ -error).

The proof of Proposition 2.19 is straightforward from standard arguments combined with the induced removal lemma of Rödl and Schacht [46], and appears in Appendix A.

To answer Problem 2.17 (2), we characterize $\text{disc}_{2,3}$ -homogeneity in terms of VC_2 -dimension, which we define below. Given a 3-graph $H = (V, E)$, let $\text{Trip}(H)$ denote the 3-partite 3-graph $(X \cup Y \cup Z, E')$, where $X = \{x_v : v \in V\}$, $Y = \{y_v : v \in V\}$, and $Z = \{z_v : v \in V\}$ are disjoint copies of V , and $E' = \{x_v y_{v'} w_{v''} : vv'v'' \in E\}$. For a hereditary 3-graph property \mathcal{H} , we define $\text{Trip}(\mathcal{H}) = \{\text{Trip}(H) : H \in \mathcal{H}\}$. It is straightforward to check that $\text{Trip}(\mathcal{H})$ is also a hereditary 3-graph property.

Definition 2.20 ($V(k)$). Given $k \geq 1$, define $V(k)$ to be the 3-partite 3-graph

$$V(k) = (\{b_1, \dots, b_k\} \cup \{c_1, \dots, c_k\} \cup \{a_S : S \subseteq [k]^2\}, \{a_S b_v c_w : (v, w) \in S\}).$$

In other words, $V(k)$ encodes the power set of the Cartesian product $[k] \times [k]$.

Definition 2.21 (VC_2 -dimension of a 3-graph). For a 3-graph H , the VC_2 -dimension of H , denoted by $\text{VC}_2(H)$, is the largest k such that $\text{Trip}(H)$ contains an induced copy of $V(k)$.

In other words, given a 3-graph $H = (V, E)$, $\text{VC}_2(H)$ is the largest k for which there exists $b_1, \dots, b_k, c_1, \dots, c_k \in V$, and $a_S \in V$ for each $S \subseteq [k]^2$, so that $a_S b_i c_j \in E$ if and only if $(i, j) \in S$.

Given a hereditary 3-graph property \mathcal{H} , we say it has *bounded VC_2 -dimension*² (is NIP_2) if there is some $k \geq 1$ such that $\text{VC}_2(H) \leq k$ for all $H \in \mathcal{H}$. In this case, we define the *VC_2 -dimension of \mathcal{H}* , $\text{VC}_2(\mathcal{H})$, to be $\max\{\text{VC}_2(H) : H \in \mathcal{H}\}$. Otherwise, we say that \mathcal{H} has *unbounded VC_2 -dimension* (has IP_2), and write $\text{VC}_2(\mathcal{H}) = \infty$. This notion has several equivalent formulations, for example the following (see [12]).

Fact 2.22. *\mathcal{H} has unbounded VC_2 -dimension if and only if $\text{Trip}(\mathcal{H})$ contains every finite 3-partite 3-graph.*

This is analogous to the graph setting, where a hereditary graph property \mathcal{H} has unbounded VC-dimension if and only if $\text{Bip}(\mathcal{H})$ contains every finite bipartite graph (this is folklore, and an easy exercise). In the case of graphs, with a bit more work on top of Theorem 1.1, one can show that a hereditary graph property has bounded VC-dimension if and only if it has regular decompositions where all regular pairs are homogeneous. We prove the analogous result for hereditary 3-graph properties and VC_2 -dimension. In particular, we first show the following result for 3-graphs of uniformly bounded VC_2 -dimension (see Section 5).

Theorem 2.23. *For all $k \geq 1$, $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $\ell_0, t_0 \geq 1$, there are $N, T, L \geq 1$ such that the following hold. Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$ and $\text{VC}_2(H) < k$. Then there exist $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and ϵ_1 -homogeneous with respect to H .*

A closely related result was recently proved by Chernikov and Towsner [15]. Specifically, they showed that if an r -uniform hypergraph equipped with a Keisler measure has bounded VC_{r-1} -dimension, then it has a decomposition where all “regular triads” have density near 0 or 1 with respect to said measure [15]. They employ infinitary techniques, and their results are non-quantitative. Our proof of Theorem 2.23, on the other hand, is based on Theorem 2.8 and thus yields the same bounds as can be achieved there (see [38]). Subsequent to the results of Section 5, the first author [58] produced a different proof of Theorem 2.23, in which the bound L in can be taken to be of the form $\epsilon_1^{-O_k(1)}$. See [57] for details.

An immediate corollary of Theorem 2.23 is that any NIP_2 hereditary 3-graph property is $\text{disc}_{2,3}$ -homogeneous. We will show that this is a characterization.

Theorem 2.24. *Suppose \mathcal{H} is a hereditary 3-graph property. Then \mathcal{H} is $\text{disc}_{2,3}$ -homogeneous if and only if \mathcal{H} has bounded VC_2 -dimension.*

Combining [58] with the lower bound construction of Moshkovitz and Shapira [38], we see that bounded VC_2 -dimension is also characterized in terms of the bounds in the regularity lemma.

²For the model theorist, \mathcal{H} has bounded VC_2 -dimension if and only if the edge relation is an NIP_2 -formula in every model of $T_{\mathcal{H}}$.

We now turn to vdisc_3 -homogeneity and Problem 2.17 (1). We begin by stating prior results. Recall that the VC-dimension of a graph G is the largest k such that $U(k)$ appears as an induced subgraph of the bipartite graph $\text{Bip}(G)$. A simple way to extend this notion to the setting of 3-graphs is via the VC-dimension of an auxiliary bipartite graph associated to a 3-graph.

Definition 2.25 (Graph associated to a 3-graph). Suppose $H = (V, E)$ is a 3-graph. The graph associated to H is $\text{Graph}(H) := (V \cup \binom{V}{2}, R)$, where $R = \{ae : \{a\} \cup e \in E\}$.

The VC-dimension of H , $\text{VC}(H)$, is then defined as the VC-dimension of $\text{Graph}(H)$. We say that a hereditary 3-graph property has *unbounded VC-dimension* (or IP) if for all k , there is $H \in \mathcal{H}$ with $\text{VC}(H) \geq k$. Otherwise we say \mathcal{H} has *bounded VC-dimension*³ (or NIP), and let $\text{VC}(\mathcal{H}) = \max\{\text{VC}(H) : H \in \mathcal{H}\}$. In [22], Fox et al. showed that 3-graph properties of bounded VC-dimension are vdisc_3 -homogeneous. In fact, they showed something much stronger, namely that any k -graph with VC-dimension at most d has an equitable ϵ -homogeneous partition with $O_{d,k}(\epsilon^{-d})$ many parts (see [22] for details). Similar results with a weaker bound were also obtained by Chernikov and Starchenko in [14]. Thus, having bounded VC-dimension is sufficient for vdisc_3 -homogeneity. However, it turns out that it is not necessary.

Proposition 2.26. *There is a hereditary 3-graph property \mathcal{H} which is vdisc_3 -homogeneous but which has unbounded VC-dimension.*

The example used to prove Proposition 2.35 is extremely simple. Concretely, we define the infinite 3-graph

$$\mathcal{W}_1 = (\{b_i, c_i : i \in \mathbb{N}\} \cup \{a_S : S \in \mathcal{P}(\mathbb{N})\}, \{a_S b_j c_k : j = k \in S\}),$$

and then let $\mathcal{H}_{\mathcal{W}_1}$ contain all finite 3-graphs isomorphic to a finite induced sub-3-graph of \mathcal{W}_1 . The key feature of this example is that, while the edge relation has unbounded VC-dimension in \mathcal{W}_1 , for any finite induced sub-3-graph H of \mathcal{W}_1 , the copies of $U(k)$ in $\text{Graph}(H)$ are extremely sparse. This contrasts with the following example of a 3-graph, in which copies of $U(k)$ occur more densely.

Definition 2.27 ($\overline{U}(k)$). Define $\overline{U}(k)$ to be the 3-partite 3-graph

$$\overline{U}(k) = (\{a_1, \dots, a_k\} \cup \{b_S : S \subseteq [k]\} \cup \{c_1, \dots, c_k\}, \{a_i b_S c_j : j \in S\}).$$

One can think of $\overline{U}(k)$ as arising from adjoining k new vertices to $U(k)$ to create a 3-graph. We will show that vdisc_3 -homogeneity is characterized by whether or not there is a bound on the size of k for which $\overline{U}(k) \in \text{Trip}(\mathcal{H})$.

Theorem 2.28. *\mathcal{H} is vdisc_3 -homogeneous if and only if there is $k \geq 1$ such that $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$.*

³For the model theorist, \mathcal{H} has bounded VC-dimension if and only if the edge relation is an NIP formula for every partition of the variables, in every model of $T_{\mathcal{H}}$.

Note that if a hereditary 3-graph property \mathcal{H} has VC-dimension less than k , then $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$. However, the converse of this is false, as witnessed by $\mathcal{H}_{\mathcal{W}_1}$.

There are several ways a ternary hypergraph $H = (V, E)$ can have large VC-dimension, based on how elements of $V \cup \binom{V}{2}$ are chosen to build a copy of $U(k)$ in $\text{Graph}(H)$. We now single out a particular way that will be important for stating our results on vdisc_3 -homogeneity.

Definition 2.29 ($U^*(k)$). Given $k \geq 1$, define $U^*(k)$ to be the 3-partite 3-graph

$$U^*(k) = (\{a\} \cup \{b_S : S \subseteq [k]\} \cup \{c_1, \dots, c_k\}, \{abc_j : j \in S\}).$$

Note that $U^*(k)$ arises by adding a single new vertex to $U(k)$ to make a 3-graph. It is easy to see that if $U^*(k) \in \text{Trip}(\mathcal{H})$, then \mathcal{H} has VC-dimension at least k . On the other hand, it is not hard to show that even though the example $\mathcal{H}_{\mathcal{W}_1}$ above has unbounded VC-dimension, $U^*(k) \notin \text{Trip}(\mathcal{H}_{\mathcal{W}_1})$ for any $k > 1$. Thus, not every 3-graph H with large VC-dimension contains a copy of $U^*(k)$ in $\text{Trip}(H)$. For this reason we refer to the analogue of VC-dimension defined below as *weak*.

Definition 2.30 (Weak VC-dimension). Suppose H is a finite 3-graph. The *weak VC-dimension of H* is $\text{WVC}(H) := \max\{k : U^*(k) \text{ is an induced sub-3-graph of } \text{Trip}(H)\}$.

In other words, the weak VC-dimension of $H = (V, E)$ is the largest k for which there exist $a, c_1, \dots, c_k \in V$, and $b_S \in V$ for each $S \subseteq [k]$, so that $ac_i b_S \in E$ if and only if $i \in S$. This is also the largest k such that for some $a \in V$, the graph $(V, N(a))$ has VC-dimension k , where $N(a) = \{bc \in \binom{V}{2} : abc \in E\}$.

We say that a hereditary 3-graph property \mathcal{H} has *bounded WVC-dimension*⁴ (or is WNIP) if there is some k so that $\text{WVC}(H) \leq k$ for all $H \in \mathcal{H}$. In this case, we define the *WVC-dimension of \mathcal{H}* to be $\text{WVC}(\mathcal{H}) = \max\{\text{WVC}(H) : H \in \mathcal{H}\}$. Otherwise, we say \mathcal{H} has *unbounded weak VC-dimension* (or has WIP), and write $\text{WVC}(\mathcal{H}) = \infty$.

It turns out that a hereditary 3-graph property is close to a WNIP property if and only if $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$ for some $k \geq 1$.

Theorem 2.31. *Suppose \mathcal{H} is a hereditary 3-graph property. Then \mathcal{H} is close to some \mathcal{H}' with $\text{WVC}(\mathcal{H}') < \infty$ if and only if there is k such that $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$.*

This gives us another way to identify the \sim -classes which are vdisc_3 -homogeneous.

Theorem 2.32. *The following are equivalent.*

- (1) \mathcal{H} is vdisc_3 -homogeneous.
- (2) $\text{Trip}(\mathcal{H})$ omits $\overline{U}(k)$ for some k .
- (3) There is a property \mathcal{H}' with $\text{WVC}(\mathcal{H}') < \infty$ such that $\mathcal{H} \sim \mathcal{H}'$.

An interesting open question is to determine the optimal bounds corresponding to regularity lemmas for vdisc_3 -homogeneous properties. Specifically, Fox et al. [22] obtained

⁴For the model theorist, \mathcal{H} has bounded WVC-dimension if and only if for every $M \models T_{\mathcal{H}}$, and every $a \in M$, $R(a, x, y)$ is an NIP formula in \mathcal{M} .

polynomial bounds under the assumption of bounded VC-dimension. Can one still obtain such bounds if one only assumes bounded WVC-dimension?

In [15], Chernikov and Towsner produced results similar to Theorem 2.32 via a different proof which uses infinitary techniques, as well as related results for k -uniform hypergraphs satisfying various generalizations of the notion of having bounded weak VC-dimension.

We now turn to Problem 2.14, which asks about irregular triples/triads. We begin with Problem 2.14 (1) concerning zero vdisc_3 -error. Previous work has shown that this problem is related to stability in 3-graphs, as defined below.

Definition 2.33 (Stability of a 3-graph property). Given a 3-graph $H = (V, E)$, we say H is k -stable if $\text{Graph}(H)$ is k -stable.

We say a hereditary 3-graph property \mathcal{H} is *stable*⁵ if there is k such that every $H \in \mathcal{H}$ is k -stable.

It was shown in [1] that k -stable 3-graphs have vdisc_3 -regular decompositions with no irregular triples. Similar results were also obtained in [14] without equitability conditions on the partition. The results in [1] directly imply the following.

Theorem 2.34 (Stable 3-graphs admit zero vdisc_3 -error). *Suppose \mathcal{H} is a hereditary 3-graph property. If \mathcal{H} is stable, then \mathcal{H} admits zero vdisc_3 -error.*

However, it turns out that stability is too strong to give a characterization, as we shall show by way of the following proposition.

Proposition 2.35. *There is a hereditary 3-graph property which admits zero vdisc_3 -error but which is not stable.*

The example used to prove Proposition 2.35 is similar to $\mathcal{H}_{\mathcal{W}_1}$. Concretely, we define the infinite 3-graph

$$\mathcal{W}_2 = (\{a_i, b_i, c_i : i \in \mathbb{N}\}, \{a_i b_j c_k : i \leq j \text{ and } j = k\}),$$

and then let $\mathcal{H}_{\mathcal{W}_2}$ be the class of all finite 3-graphs isomorphic to an induced sub-3-graph of \mathcal{W}_2 . The important quality of this example is that while the edge relation is unstable in \mathcal{W}_2 , given any finite induced sub-3-graph H of \mathcal{W}_2 , the copies of $H(k)$ in $\text{Graph}(H)$ are extremely sparse. The following is a simple example of a 3-graph where half-graphs occur more densely.

Definition 2.36 ($\overline{H}(k)$). Given $k \geq 1$, define $\overline{H}(k)$ to be the 3-graph

$$\overline{H}(k) = (\{a_i, b_i, c_i : i \in [k]\}, \{a_i b_j c_k : j \leq k\}).$$

One can think of $\overline{H}(k)$ as constructed by adjoining k new vertices to $H(k)$ to obtain a 3-graph. We prove that the presence of these 3-graphs is sufficient to imply that a property requires non-zero vdisc_3 -error.

Theorem 2.37. *If $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for every k , then \mathcal{H} requires non-zero vdisc_3 -error.*

⁵For the model theorist, \mathcal{H} is stable if and only if the edge relation is stable, for every bipartition of the variables, in every model of $T_{\mathcal{H}}$.

We conjecture that the converse is true as well.

Conjecture 2.38. *If there is some $k \geq 1$ such that $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$, then \mathcal{H} admits zero vdisc_3 -error.*

We will also give an equivalent formulation of the conjecture by characterizing the \sim -classes $[\mathcal{H}]$ with the property that $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$ for some k . This characterization centers on the 3-graph obtained by adjoining a single vertex to $H(k)$ to obtain a 3-graph.

Definition 2.39 ($H^*(k)$). Given $k \geq 1$, define $H^*(k)$ to be the 3-graph

$$H^*(k) = (\{a\} \cup \{b_i, c_i : i \in [k]\}, \{ab_i c_j : i \leq j\}).$$

Note that if $H^*(k)$ appears as an induced sub-3-graph of G , then G is not k -stable. However, not every G with the k -order property has an induced copy of $H^*(k)$ in $\text{Trip}(G)$ (the 3-graph used to prove Proposition 2.35 is such an example). For this reason, we will name the following property *weak stability*.

Definition 2.40 (Weak stability). A 3-graph $H = (V, E)$ is *weakly k -stable* if and only if $H^*(k)$ is not an induced sub-3-graph of $\text{Trip}(H)$.

A hereditary 3-graph property \mathcal{H} is *weakly stable*⁶ if there is some k such that every $H \in \mathcal{H}$ is weakly k -stable.

In other words, a 3-graph $H = (V, E)$ is weakly k -stable if for all $a \in V$, the graph $H_a := (V, N(a))$ is k -stable, where $N(a) = \{bc : abc \in E\}$. It turns out that weak stability characterizes the \sim -class $[\mathcal{H}]$ such that $\text{Trip}(\mathcal{H})$ omits $\overline{H}(k)$ for some k .

Theorem 2.41. *Suppose \mathcal{H} is a hereditary 3-graph property. The following are equivalent.*

- (1) *There is some $k \geq 1$ such that $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$.*
- (2) *There is some weakly stable hereditary 3-graph property \mathcal{H}' such that $\mathcal{H} \sim \mathcal{H}'$.*

Thus the following (a priori weaker) conjecture is in fact equivalent to Conjecture 2.38.

Conjecture 2.42. *If \mathcal{H} is weakly stable, then \mathcal{H} admits zero vdisc_3 -error.*

We will prove something weaker than Conjecture 2.38 in Section 5.5, namely that a property which is close to a weakly stable property admits binary vdisc_3 -error.

Proposition 2.43. *Suppose \mathcal{H} is close to some weakly stable \mathcal{H}' . Then \mathcal{H} admits binary vdisc_3 -error.*

One tool we will use in the proof of Proposition 2.43 is the following set of conditions that are equivalent to having zero and binary vdisc_3 -error, respectively.

Proposition 2.44. *Suppose \mathcal{H} is a hereditary 3-graph property.*

- (1) *\mathcal{H} admits zero vdisc_3 -error if and only if \mathcal{H} is close to a WNIP property and admits zero $\text{disc}_{2,3}$ -error.*

⁶For the model theorist, \mathcal{H} is weakly stable if and only if for all $\mathcal{M} \models T_{\mathcal{H}}$ and $a \in M$, $R(a, x, y)$ is a stable formula in \mathcal{M} .

- (2) \mathcal{H} admits binary vdisc_3 -error if and only if \mathcal{H} is close to a WNIP property and admits binary $\text{disc}_{2,3}$ -error.

The proof of Proposition 2.44 uses Theorem 2.32 alongside some standard techniques from extremal combinatorics. To prove Proposition 2.43, we use Proposition 2.44, and ideas from the proof of the stable graph regularity lemma of [36], as well as the strong graph regularity lemma (see e.g. [2, 35, 48]). We believe this argument provides useful techniques for showing that a given hereditary 3-graph property admits binary error.

Note that the combination of Proposition 2.44 and Theorem 2.32 immediately implies that any property admitting zero or binary vdisc_3 -error is also vdisc_3 -homogeneous.

Corollary 2.45. *If \mathcal{H} admits zero or binary vdisc_3 -error, then \mathcal{H} is vdisc_3 -homogeneous.*

Proposition 2.44 will also play a role in showing that Problem 2.14 (1) and (2) are in fact the same question. In particular, a property admits zero $\text{disc}_{2,3}$ -error if and only if it admits zero vdisc_3 -error.

Theorem 2.46. *\mathcal{H} admits zero $\text{disc}_{2,3}$ -error if and only if \mathcal{H} admits zero vdisc_3 -error.*

In addition to Proposition 2.44, the proof of Theorem 2.46 will use Theorems 2.32 and 2.37.

We now turn to part (4) of Problem 2.14, which asks for a characterization of the hereditary 3-graph properties admitting linear $\text{disc}_{2,3}$ -error. The motivation for this question, as well as the definition of linear error, stems from [60], where such decompositions occur naturally in the ternary sum graphs associated to certain quadratically structured subsets of \mathbb{F}_p^n . As in that paper, we show that such decompositions are related to the following combinatorial notion.

Definition 2.47 ($F(\ell)$). Given $\ell \geq 1$, let $F(\ell)$ be the 3-graph

$$F(\ell) = (\{a_i^f, b_i, c_i : i \in [\ell], f : [\ell]^2 \rightarrow [\ell]\}, \{a_i^f b_j c_k : k \leq f(i, j)\}).$$

Definition 2.48 (Ternary functional order property (FOP_2)). Given $\ell \geq 1$ and a 3-graph H , we say H has ℓ - FOP_2 if there is an induced copy of $F(\ell)$ in $\text{Trip}(H)$.

In other words, $H = (V, E)$ has ℓ - FOP_2 if and only if there exist $b_1, \dots, b_\ell, c_1, \dots, c_\ell \in V$ and for every function $f : [\ell]^2 \rightarrow [\ell]$, vertices a_1^f, \dots, a_ℓ^f , such that $a_i^f b_j c_k \in E$ if and only if $k \leq f(i, j)$. We then say that a hereditary 3-graph property \mathcal{H} has FOP_2 if for all $\ell \geq 1$, there is $H \in \mathcal{H}$ with ℓ - FOP_2 . Otherwise, we say \mathcal{H} is NFOP_2 .

It is not difficult to see (see [60]) that if \mathcal{H} has FOP_2 then it has unbounded VC-dimension. It is further clear from Fact 2.22 that if \mathcal{H} has unbounded VC_2 -dimension, then it has FOP_2 . We will show this implication is strict in Appendix D. For further properties of general NFOP_2 formulas, including a proof that NFOP_2 is closed under finite boolean combinations, we refer the reader to Appendix D.

In [60] we show that given a set $A \subseteq \mathbb{F}_p^n$, if the 3-graph $\Gamma_A = (\mathbb{F}_p^n, \{xyz : x + y + z \in A\})$ has no ℓ - FOP_2 , then there is a decomposition of \mathbb{F}_p^n defined by a high rank quadratic factor, so that on almost all parts in the decomposition, the set A has density near 0 or

1. Moreover, the parts of this decomposition on which the density of A is bounded away from 0 and 1 are contained in a small number of linearly defined pieces [60, Theorem 1.33], hence the term “linear error”. This structure theorem can be translated directly into the statement that there exists a regular decomposition for Γ_A with linear error, in the sense of Definition 2.11 (see [60, Corollary 4.30]). Here, we prove an analogous result for 3-graphs, namely that if H is a large 3-graph with no ℓ -FOP₂, then it has a $\text{disc}_{2,3}$ -regular decomposition with linear error.

Theorem 2.49. *For all $k \geq 1$, $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$, there are T, L, N such that the following hold. For all $H = (V, E)$ with $|V| \geq N$ such that H has no k -FOP₂, there are $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$, and \mathcal{P} a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and which has linear $\text{disc}_{2,3}$ -error with respect to H .*

Theorem 2.49 implies, by definition, that any NFOP₂ hereditary 3-graph property admits linear $\text{disc}_{2,3}$ -error. We will show that NFOP₂ hereditary 3-graph properties are in fact characterized by the admission of linear $\text{disc}_{2,3}$ -error.

Theorem 2.50. *Suppose \mathcal{H} is a hereditary 3-graph property. Then \mathcal{H} is NFOP₂ if and only if \mathcal{H} admits linear $\text{disc}_{2,3}$ -error.*

We finally arrive at parts (2) and (3) of Problem 2.14, which ask about 3-graph properties admitting binary vdisc_3 - and binary $\text{disc}_{2,3}$ -error, respectively. For this problem, we will give two examples of properties which require binary $\text{disc}_{2,3}$ -error and binary vdisc_3 -error. The first example is the following.

Definition 2.51 (HP(k)). Given $k \geq 1$, define HP(k) to be the 3-partite 3-graph

$$\text{HP}(k) = (\{a_i, b_i, c_i : i \in [k]\}, \{a_u b_v c_w : u + v + w \geq k + 2\}).$$

The 3-graph HP(k) is a natural higher-order analogue of the half graph. Indeed, observe that $H(k)$ is isomorphic to the graph with vertex set $\{a_i, b_i : i \in [k]\}$ and edge set $\{a_u b_v : u + v \geq k + 1\}$. Based on this analogy, it is natural to make the following definition.

Definition 2.52 (Hyperplane order property (HOP₂)). We say a 3-graph H has the k -hyperplane order property of dimension 3 (k -HOP₂) if HP(k) is an induced sub-3-graph of $\text{Trip}(H)$.

A hereditary 3-graph property \mathcal{H} has k -HOP₂ if there is $H \in \mathcal{H}$ which has k -HOP₂. We say that \mathcal{H} has HOP₂ if it has k -HOP₂ for all $k \geq 1$. Otherwise, we say \mathcal{H} has HOP_2 .

Besides the obvious geometric reasons for viewing HOP₂ as a ternary analogue of a half-graph, there are algebraic reasons why Definition 2.51 might appear to be the canonical ternary analogue of the order property. We refer the reader to [60] for an extensive discussion of this. In the hypergraph setting, one compelling piece of evidence is the fact that if a hereditary 3-graph property has HOP₂, then it requires non-binary $\text{disc}_{2,3}$ -error.

Theorem 2.53. *Suppose \mathcal{H} is a hereditary 3-graph property with HOP₂. Then \mathcal{H} requires non-binary $\text{disc}_{2,3}$ -error.*

However, the authors' work on analogous questions in the arithmetic setting [60] identified a second, distinct example whose presence requires non-binary $\text{disc}_{2,3}$ -error. The example is defined as a disjoint union of cosets of a nested chain of subgroups in an elementary abelian p -group.

Definition 2.54 (Green-Sanders examples). Suppose $p \geq 3$. Given $n \geq 1$, define

$$A(p, n) := \{\bar{x} \in \mathbb{F}_p^n : \text{there is } 1 \leq i \leq n \text{ such that } x_i = 1 \text{ and for all } 1 \leq j < i, x_j = 0\}.$$

The *Green-Sanders p -example of dimension n* is the pair $(\mathbb{F}_p^n, A(p, n))$.

In the case when $p = 3$, Green and Sanders [28] showed that every subspace of \mathbb{F}_3^n has a non-trivial Fourier coefficient with respect to $A(3, n)$, and in [60], the authors showed that for any prime $p \geq 3$, the sets $A(p, n) \subseteq \mathbb{F}_p^n$ cannot be written as a union of quadratic atoms with “no error” (see [60]). We refer the reader to the forthcoming note [59] for further examples with similar properties. The sets $A(p, n)$ of Definition 2.54 naturally give rise to 3-partite 3-graphs as follows.

Definition 2.55 ($\text{GS}_p(n)$). We define $\text{GS}_p(n)$ to be the 3-graph

$$\text{GS}_p(n) = (\{a_g, b_g, c_g : g \in \mathbb{F}_p^n\}, \{a_g b_{g'} c_{g''} : g + g' + g'' \in A(p, n)\}).$$

We prove that these examples also require non-binary $\text{disc}_{2,3}$ -error.

Theorem 2.56. *If a hereditary 3-graph property \mathcal{H} has $\text{GS}_p(n) \subseteq \text{Trip}(\mathcal{H})$ for every n , then \mathcal{H} requires non-binary $\text{disc}_{2,3}$ -error.*

The authors show in [60] that $\text{GS}_3(n)$ does not have 4-HOP₂, and thus, NHOP₂ cannot characterize the properties which admit binary $\text{disc}_{2,3}$ -error. Instead, it seems that Theorems 2.53 and 2.56 are distinct instantiations of another, more general combinatorial definition.

Let \mathcal{H}_{HP} be the hereditary 3-graph obtained by closing $\bigcup_{k \geq 1} \text{HP}(k)$ under isomorphism and induced sub-3-graphs, and let $\mathcal{H}_{\text{GS}_p}$ be the hereditary 3-graph property obtained by closing $\bigcup_{n \geq 1} \text{GS}_p(n)$ under isomorphism and induced sub-3-graphs. Theorems 2.53 and 2.56 tell us that if \mathcal{H}_{HP} or $\mathcal{H}_{\text{GS}_p}$ are contained in $\text{Trip}(\mathcal{H})$, then \mathcal{H} requires binary $\text{disc}_{2,3}$ -error. Interestingly, both these examples have bounded (in fact, very small) VC-dimension.

Proposition 2.57. *\mathcal{H}_{HP} has VC-dimension 1 and for all $p \geq 3$, $\mathcal{H}_{\text{GS}_p}$ has VC-dimension 3.*

We will give proofs of these bounds in Section 7. Proposition 2.57 also plays a role in our proofs of Theorems 2.53 and 2.56. In particular, since properties with finite VC-dimension also have finite WVC-dimension, both $\mathcal{H}_{\text{GS}_p}$ and \mathcal{H}_{HP} are WNIP. Thus by Proposition 2.44, in order to show that they require non-binary $\text{disc}_{2,3}$ -error, it suffices to show that both require non-binary vdisc_3 -error. We will do this using a general sufficient condition satisfied by both examples (see Section 7.3).

Proposition 2.57 shows that a hereditary 3-graph property can have bounded VC-dimension while also requiring non-binary $\text{disc}_{2,3}$ -error. On the other hand, we will show

that there are properties admitting binary $\text{disc}_{2,3}$ -error which not only have unbounded VC-dimension, but which contain $\overline{U}(k)$ for all k .

Proposition 2.58. *There is a hereditary 3-graph property \mathcal{H} which admits binary $\text{disc}_{2,3}$ -error, and which contains $\overline{U}(k)$ for all k .*

Thus, whether or not a property admits binary $\text{disc}_{2,3}$ -error has no general relationship to whether or not it has bounded VC-dimension. This might seem surprising at first, but it reflects the fact that that hypergraph regularity is really about ternary structure, while VC-dimension is closely related to binary structure. Proposition 2.58 also shows that a property can admit binary $\text{disc}_{2,3}$ -error while not admitting binary vdisc_3 -error. Indeed, by Theorem 2.32, if a property contains $\overline{U}(k)$ for all k , it is not vdisc_3 -homogeneous, and therefore cannot admit binary vdisc_3 -error. Combining this with Proposition 2.44, we see that admitting binary vdisc_3 -error is strictly stronger than admitting binary $\text{disc}_{2,3}$ -error.

In contrast to this, there is a clear implication between stability and binary $\text{disc}_{2,3}$ -error. In particular, if \mathcal{H} is stable, then by Theorem 2.34, \mathcal{H} admits zero vdisc_3 -error, so by Proposition 2.44, it admits zero $\text{disc}_{2,3}$ -error. If Conjecture 2.42 were true, then a similar argument would show that any weakly stable \mathcal{H} admits zero $\text{disc}_{2,3}$ -error, and hence binary $\text{disc}_{2,3}$ -error. We show directly in this paper (Proposition 2.43) that any weakly stable property admits binary $\text{disc}_{2,3}$ -error (in fact, binary vdisc_3 -error).

Finally, our results imply that admitting any good control over the $\text{disc}_{2,3}$ -irregular triads implies $\text{disc}_{2,3}$ -homogeneity (see Section 7).

Corollary 2.59. *If \mathcal{H} admits zero, binary, or linear $\text{disc}_{2,3}$ -error, then \mathcal{H} is $\text{disc}_{2,3}$ -homogeneous.*

We end the introduction with a discussion of some open problems. As mentioned above, we conjecture that \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ are two manifestations of an underlying combinatorial definition, which will characterize the properties admitting binary $\text{disc}_{2,3}$ -error.

Conjecture 2.60. *There exists a definition XOP_2 (for “Unknown Order Property”) which is defined by an infinite scheme of existential sentences in $\mathcal{L} = \{R(x, y, z)\}$, such that a hereditary 3-graph property admits binary $\text{disc}_{2,3}$ -error if and only if it does not satisfy XOP_2 .*

We also conjecture a connection between this property and quadratic structure in the arithmetic setting. For a discussion of this, we direct the reader to [60].

In addition to the interesting problem above for 3-graphs, several open questions arise from this work with regard to k -graphs for $k > 3$. For example, the results of Fox et al. [22] give strong regularity lemmas for k -graphs with bounded VC-dimension. Our work suggests that an analogue of weak VC-dimension should characterize hereditary k -graph properties with such partitions. Similarly, while Theorem 2.34 was proven in the setting of k -graphs for $k \geq 2$, it remains open to give a characterization of which hereditary k -graph properties admit regular partitions as described there. It is fairly clear that bounded VC_k -dimension (see [12] for the definition) will correspond to the appropriate higher order analogue of $\text{disc}_{2,3}$ -homogeneity. Indeed, all the same tools used in Section 4 have their analogues for

k -graphs. One direction of this has been proved, non-quantitatively, by Chernikov and Towsner [15].

It is also straightforward to define a higher-dimensional analogue of the ternary functional order property.

Definition 2.61 (FOP_k). Given a $(k+1)$ -graph $H = (V, E)$, we say H has ℓ - FOP_k if there are $\{x_j^u : j \in [\ell], u \in [k]\} \subseteq V$ and for every function $f : [\ell]^k \rightarrow [\ell]$, there are $z_1^f, \dots, z_\ell^f \in V$ such that $z_j^f x_{i_1}^1 \dots x_{i_k}^k \in E$ if and only if $j < f(i_1, \dots, i_k)$.

We say a hereditary $(k+1)$ -graph property \mathcal{H} has FOP_k if for all $\ell \geq 1$, there is $H \in \mathcal{H}$ with ℓ - FOP_k . Otherwise, we say \mathcal{H} is NFOP_k .

We conjecture that NFOP_k should have a characterization analogous to Theorem 2.50. There are also natural k -dimensional generalizations of the $\text{HP}(n)$ and $\text{GS}_p(n)$.

Definition 2.62. Given $k, n \geq 1$, define

$$\text{HP}_k(n) = (\{a_i^j : i \in [n], j \in [k]\}, \{a_{i_1}^1 \dots a_{i_k}^k : i_1 < i_2 < \dots < i_k\}).$$

Let $\mathcal{H}_{\text{HP}_k}$ be the closure of $\bigcup_{n \geq 1} \text{HP}_k(n)$ under isomorphism and induced sub- k -graph.

Definition 2.63 ($\text{GS}_p^{(k)}(n)$). Suppose $\ell, k \geq 1$ and $p \geq 3$ is a prime. Define

$$\text{GS}_p^{(k)}(n) = (\{a_g^i : i \in [k], g \in \mathbb{F}_p^n\}, \{a_{g_1}^1 \dots a_{g_k}^k : g_1 + \dots + g_k \in A(p, n)\}).$$

Let $\mathcal{H}_{\text{GS}_p^{(k)}}$ be the closure of $\bigcup_{n \geq 1} \text{GS}_p^{(k)}(n)$ under isomorphism and induced sub- k -graph.

We conjecture that both $\mathcal{H}_{\text{HP}_k}$ and $\mathcal{H}_{\text{GS}_p^{(k)}}$ fail to admit the appropriate generalization of binary $\text{disc}_{2,3}$ -error. Clearly this also leads to a higher order version of Conjecture 2.60.

2.4. Outline of paper. We now give a brief outline of what we do in each section. In Section 3, we set up notation, and give further background on regularity lemmas for 3-graphs, counting lemmas, and other tools we use throughout the paper. In Section 4 we prove Theorem 2.24, the equivalence between $\text{disc}_{2,3}$ -homogeneity and finite VC_2 -dimension. In Section 5, we prove our results about vdisc_3 -homogeneity, weak VC -dimension, and weak stability, including Theorems 2.28, 2.32, 2.31, 2.37, 2.41 and Propositions 2.26, 2.35, 2.43, 2.44, and Corollary 2.45. Section 5 also contains strong and generalized versions of the regularity lemma for stable graphs (see Theorems 5.7 and 5.15). In Section 6 we prove our results on linear error and FOP_2 , including Theorem 2.50. In this section there are also several general results about FOP_2 -formulas and recharacterizations of both FOP_2 and IP_2 hereditary 3-graph properties. Section 6 also contains a strong removal lemma for stable graphs, with an accompanying structure theorem for graphs which contain few half-graphs (see Theorem 6.13 and Lemma 6.14). In 7, we prove our main results regarding examples of properties requiring binary error, including Theorems 2.53 and 2.56. It is in this section that we prove Theorem 2.12, which recharacterizes binary $\text{disc}_{2,3}$ -error in terms of no disc_3 -irregular triads.

3. PRELIMINARIES

This section contains the necessary background on quasirandomness and regularity. We begin with some notation.

Given a positive integer n , let $[n] := \{1, \dots, n\}$. Given a set X and $k \geq 1$, define

$$\binom{X}{k} := \{Y \subseteq X : |Y| = k\}.$$

For sets X, Y, Z , we set

$$\begin{aligned} K_2[X, Y] &= \{\{x, y\} : x \in X, y \in Y, x \neq y\} \text{ and} \\ K_3[X, Y, Z] &= \{\{x, y, z\} : x \in X, y \in Y, z \in Z, x \neq y, y \neq z, x \neq z\}. \end{aligned}$$

When $x, y \in X$ are distinct, we will write xy to denote the set $\{x, y\}$. Similarly, if $x, y, z \in X$ are pairwise distinct, we write xyz to denote the set $\{x, y, z\}$.

A k -uniform hypergraph is a pair (V, E) where V is a set of vertices and $E \subseteq \binom{V}{k}$ is a set of edges. When $k = 2$, a k -uniform hypergraph is a *graph*. To ease notation, throughout the paper, we will call a k -uniform hypergraph a k -graph. Given a k -graph G , $V(G)$ denotes the vertex set of G , $E(G)$ denotes its edge set, and we set $v(G) := |V(G)|$ and $e(G) := |E(G)|$. We say a k -graph $G' = (V', E')$ is a *sub- k -graph of G* if $V' \subseteq V$ and $E' \subseteq E$. In this case we write $G' \subseteq G$. We say G' is an *induced sub- k -graph of G* if $V' \subseteq V$ and $E' = E \cap \binom{V'}{k}$. We will occasionally call a sub- k -graph a *non-induced sub- k -graph* to emphasize what is meant. When $V'' \subseteq V$, we define $G[V''] = (V'', E \cap \binom{V''}{k})$. For a k -graph G'' , an *induced copy of G'' in G* is an induced sub- k -graph of G isomorphic to G'' . A *non-induced copy of G'' in G* is a non-induced sub- k -graph of G isomorphic to G'' .

Suppose $G = (V, E)$ is a k -graph. If $E = \binom{V}{k}$, then $G = (V, E)$ is called *complete*. For any $2 \leq k \leq r$, we say $G = (V, E)$ is *r -partite* if there is a partition $V = V_1 \cup \dots \cup V_r$ such that for all $e \in E$, and $i \in [r]$, $|e \cap V_i| \leq 1$. In this case we write $G = (V_1 \cup \dots \cup V_r, E)$ to denote that G is r -partite with partition $V = V_1 \cup \dots \cup V_r$. When $r = 2$, we say that G is *bipartite*.

Suppose now that $G = (V, E)$ is a graph. For $v \in V$, the *neighborhood of v* is

$$N_G(v) = \{u \in V : uv \in E\}.$$

Given sets $U, W \subseteq V$, set $E_G[U, W] := E \cap K_2[U, W]$, and $e_G(U, W) := |E_G[U, W]|$. The *density* between U and W is the number $d_G(U, W) \in [0, 1]$ such that

$$e_G(U, W) = d_G(U, W)|U||W|.$$

When G is clear from context, we will occasionally drop the subscripts above. We define $G[U, W]$ to be the graph $(U \cup W, E_G[U, W])$. Similarly, if $U, W, Z \subseteq V$,

$$G[U, W, Z] := (U \cup W \cup Z, E_G[U, W] \cup E_G[U, Z] \cup E_G[W, Z]).$$

Suppose that $H = (V, E)$ is a 3-graph. For sets $X, Y, Z \subseteq V$, we define

$$E_H[X, Y, Z] = \{xyz \in E : x \in X, y \in Y, z \in Z\}.$$

Given $v \in V$, $N_H(v) = \{bc \in \binom{V}{2} : abc \in E\}$ is the *neighborhood* of v , and given $uv \in \binom{V}{2}$, $N_G(uv) = \{b \in V : uvb \in E\}$ is the *neighborhood* of uv .

Let $G = (V, E)$ be a graph. For $p \geq 2$, define

$$K_p^{(2)}(G) = \left\{ e \in \binom{V}{p} : e' \in E \text{ for all } e' \in \binom{e}{2} \right\}.$$

In other words, $K_p^{(2)}(G)$ is the collection of complete p -element subgraphs of G . If $H = (V, E')$ is a 3-graph on the same vertex set as G , then we say G *underlies* H if $E' \subseteq K_3^{(2)}(G)$. In other words, G underlies H if every edge of H sits atop a triangle of G . Similarly, we let

$$K_p^{(3)}(H) = \left\{ e \in \binom{V}{p} : e' \in E \text{ for all } e' \in \binom{e}{3} \right\},$$

i.e. $K_p^{(3)}(H)$ is the collection of complete p -element sub-3-graphs of H .

If \mathcal{H} is a class of finite k -graphs, we say \mathcal{H} is *closed under induced sub- k -graphs* if for all $G = (V, E) \in \mathcal{H}$ and $V' \subseteq V$, $G[V'] := (V', E \cap \binom{V'}{k}) \in \mathcal{H}$.

Definition 3.1. Given $k \geq 2$, a *hereditary k -graph property* is a class of finite k -graphs which is closed under isomorphism and induced sub- k -graphs.

When H, H' are k -graphs, we say that H *omits* H' or is *H' -free* if no induced sub- k -graph of H is isomorphic to H' . Given a class of k -graphs \mathcal{F} , let $\text{Forb}(\mathcal{F})$ denote the class of finite k -graphs which omit every element of \mathcal{F} . Every class of the form $\text{Forb}(\mathcal{F})$ is a hereditary k -graph property. Conversely, it is well known that for every hereditary k -graph property \mathcal{H} , there is a class $\mathcal{F}_{\mathcal{H}}$ of finite k -graphs such that $\mathcal{H} = \text{Forb}(\mathcal{F}_{\mathcal{H}})$.

Recall that given a graph $G = (V, E)$, let $\text{Bip}(G)$ denote the bipartite graph $(U \cup W, E')$, where $U = \{u_v : v \in V\}$, $W = \{w_v : v \in V\}$, and $E' = \{u_v w_{v'} : vv' \in E\}$. Similarly, for a 3-graph $H = (V, E)$, let $\text{Trip}(H)$ denote the 3-partite 3-graph $(X \cup Y \cup Z, E')$, where $X = \{x_v : v \in V\}$, $Y = \{y_v : v \in V\}$, $Z = \{z_v : v \in V\}$, and $E' = \{x_v y_{v'} z_{v''} : vv'v'' \in E\}$.

Given a hereditary graph property \mathcal{H} , let $\text{Bip}(\mathcal{H}) = \{\text{Bip}(G) : G \in \mathcal{H}\}$, and given a hereditary 3-graph property \mathcal{H}' , we let $\text{Trip}(\mathcal{H}') = \{\text{Trip}(H) : H \in \mathcal{H}'\}$. Given $n \geq 1$, and a hereditary k -graph property, \mathcal{H} , set $\mathcal{H}_n = \{G \in \mathcal{H} : V(G) = [n]\}$.

For a k -graph H , $\text{age}(H)$ denotes the class of finite k -graphs isomorphic to an induced sub- k -graph of H . Note that $\text{age}(H)$ is always a hereditary k -graph property.

Given a vertex set V , and $t, \ell \geq 1$, a (t, ℓ) -decomposition \mathcal{P} of V consists of a partition $V = V_1 \cup \dots \cup V_t$ and for each $ij \in \binom{[t]}{2}$, a partition $K_2[V_i, V_j] = \bigcup_{\alpha \in [\ell]} P_{ij}^{\alpha}$. Observe that Definition 2.6 is a special kind of (t, ℓ) -decomposition. Given such a \mathcal{P} , we let

$$\mathcal{P}_{\text{vert}} = \{V_1, \dots, V_t\} \text{ and } \mathcal{P}_{\text{edge}} = \left\{ P_{ij}^{\alpha} : ij \in \binom{[t]}{2}, \alpha \in [\ell] \right\}.$$

We will write $G_{ijk}^{\alpha, \beta, \gamma}$ to denote the 3-partite graph

$$G_{ijk}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_k, P_{ij}^{\alpha} \cup P_{ik}^{\beta} \cup P_{jk}^{\gamma}),$$

called a *triad* of \mathcal{P} . When V is the vertex set of a 3-graph $H = (V, E)$, and no confusion might otherwise arise, we will write $H_{ijk}^{\alpha, \beta, \gamma}$ for the 3-partite 3-graph

$$H_{ijk}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_k, E \cap K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})).$$

We say that a triad $G_{ijk}^{\alpha, \beta, \gamma}$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2)$ with respect to H if $(H_{ijk}^{\alpha, \beta, \gamma}, G_{ijk}^{\alpha, \beta, \gamma})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2)$, and we say that $G_{ijk}^{\alpha, \beta, \gamma}$ is ϵ -homogeneous with respect to H if

$$\frac{|E \cap K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})|}{|K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})|} \in [0, \epsilon] \cup (1 - \epsilon, 1].$$

Similarly, given a partition $\mathcal{Q} = \{V_1, \dots, V_t\}$ of V , we say that a triple $V_i V_j V_k$ satisfies $\text{vdisc}_3(\epsilon)$ with respect to H if $H[V_i, V_j, V_k]$ satisfies $\text{vdisc}_3(\epsilon)$, and we say that $V_i V_j V_k$ is ϵ -homogeneous with respect to H if

$$\frac{|E \cap K_3[V_i, V_j, V_k]|}{|K_3[V_i, V_j, V_k]|} \in [0, \epsilon] \cup (1 - \epsilon, 1].$$

Given parameters a and b , we will write $a \ll b$ to mean that a sufficiently small compared to b . Similarly, we write $b \gg a$ to mean b is sufficiently large compared to a . Given $\epsilon > 0$ and $d, r \in \mathbb{R}$, we will write $d = r \pm \epsilon$ to mean $d \in (r - \epsilon, r + \epsilon)$.

3.1. Notions of quasirandomness for graphs and hypergraphs. In this section we give a more complete account of quasirandomness for graphs and 3-graphs. We begin with quasirandomness for graphs, of which there are several equivalent formulations. One of them, disc_2 , was already defined in the introduction, and we now define three more which turn out to be equivalent. Recall that C_4 is the cycle on 4-vertices.

Definition 3.2. Let $\epsilon, d_2 > 0$, and let $G = (V \cup U, E)$ be a bipartite graph with $|E| = d|U||V|$.

- (1) We say that G has $\text{cycle}_2(\epsilon, d_2)$ if $d \in (d_2 - \epsilon, d_2 + \epsilon)$ and G contains at most $(d_2^4 + \epsilon)|U|^2|V|^2$ many non-induced copies of C_4 .
- (2) We say that G has $\text{dev}_2(\epsilon, d_2)$ if $d \in (d_2 - \epsilon, d_2 + \epsilon)$ and the following holds, where $g(u, v) = 1 - d_2$ if $uv \in E$ and $g(u, v) = -d_2$ if $uv \notin E$.

$$\sum_{u_0, u_1 \in U} \sum_{v_0, v_1 \in V} \prod_{i \in \{0, 1\}} \prod_{j \in \{0, 1\}} g(u_i, v_j) \leq \epsilon |U|^2 |V|^2.$$

- (3) We say that G is ϵ -regular if for all $U' \subseteq U$ with $|U'| \geq \epsilon|U|$ and $W' \subseteq W$ with $|W'| \geq \epsilon|W|$,

$$|e_G(U', W') - d|U'||W'|| \leq \epsilon|U||W|.$$

In the notation above, we say simply that G has $\text{disc}_2(\epsilon)$ (respectively $\text{cycle}_2(\epsilon)$, $\text{dev}_2(\epsilon)$) if it has $\text{disc}_2(\epsilon, d_2)$ (respectively $\text{cycle}_2(\epsilon, d_2)$, $\text{dev}_2(\epsilon, d_2)$) for some d_2 .

Given a graph $G = (V, E)$ and pair of subsets $U, V \subseteq G$, we say (U, V) is an ϵ -regular pair (respectively that the pair has $\text{disc}_2(\epsilon)$, $\text{cycle}_2(\epsilon)$, $\text{dev}_2(\epsilon)$) if $G[U, V]$ is ϵ -regular (respectively has $\text{disc}_2(\epsilon)$, $\text{cycle}_2(\epsilon)$, $\text{dev}_2(\epsilon)$). In the late 1980s, Chung and Graham [16] showed that these notions are equivalent in the sense summarized in Proposition 3.3.

Proposition 3.3. *For all $\epsilon, d_2 > 0$, there exists a δ_2 and n_0 such that if $G = (U \cup V, E)$ is a bipartite graph with $|U| = |V| = n \geq n_0$ then the following hold.*

- (1) *If G has $\text{disc}_2(\delta_2, d_2)$, then G has $\text{cycle}_2(\epsilon, d_2)$.*
- (2) *If G has $\text{cycle}_2(\delta_2, d_2)$, then G has $\text{dev}_2(\epsilon, d_2)$.*
- (3) *If G has $\text{dev}_2(\delta, d_2)$, then G is ϵ -regular.*
- (4) *If G is δ_2 -regular with density $d_2 \pm \epsilon$, then G has $\text{cycle}_2(\epsilon, d_2)$.*

These notions of quasirandomness capture the behavior of a random graph. For example, for every $\eta > 0$, a sufficiently large random bipartite graph will have $\text{disc}_2(\eta, 1/2)$ with high probability. This can be used to prove the following fact.

Fact 3.4. *For all $\eta > 0$, there is n_0 such that for all $n \geq n_0$, almost every bipartite graph $G = (U \cup W, E)$ with $|U| = |W| = n$ has $\text{disc}_2(\eta, 1/2)$.*

Quasirandomness is useful because it allows us to count copies of small fixed subgraphs. For example, quasirandom graphs have about the same number of triangles as one would expect in a random graph, see for instance [47, Theorem 18].

Proposition 3.5 (Counting Lemma). *For every $t \geq 2$ and $\gamma > 0$ there exists $\epsilon > 0$ such that the following holds. Let $G = (V_1 \cup \dots \cup V_t, E)$ be a t -partite graph such that for each $1 \leq i \neq j \leq t$, $G[V_i, V_j]$ has $\text{disc}_2(\epsilon, d_{ij})$. Then*

$$\left| |K_t^{(2)}(G)| - \prod_{1 \leq i < j \leq t} d_{ij} \prod_{i=1}^t |V_i| \right| \leq \gamma \prod_{i=1}^t |V_i|.$$

It will be convenient to have a slightly simpler version of this proposition packaged in the following way.

Corollary 3.6. *For all $t \geq 2$ and $\epsilon, r > 0$, there is $\mu = \mu(k, \epsilon, r) > 0$ such that the following hold. Suppose $G = (U, E)$ is a graph and $U = U_1 \cup \dots \cup U_t$ is a partition so that for each $1 \leq i < j \leq t$, $G[U_i, U_j]$ has $\text{disc}_2(\mu, r)$.*

Then the number of $\bar{u} \in \prod_{i \in [t]} U_i$ with $u_i u_j \in E$ for each $ij \in \binom{[t]}{2}$ is $(1 \pm \epsilon) r^{\binom{k}{2}} \prod_{i \in [t]} |U_i|$.

It is an exercise to deduce Corollary 3.6 from Proposition 3.5 (see Appendix B). One important application of counting lemmas are so-called removal lemmas, which say that if a graph G has very few copies of a small graph H , then G can be made H -free by changing a small number of edges. We will use an induced version of this result, due to Alon, Fischer, Krivelevich, and Szegedy [2]. Recall that two graphs G and G' on the same vertex set are δ -close if $|E(G) \Delta E(G')| \leq \delta |V(G)|^2$.

Theorem 3.7 (Induced graph removal). *For any graph H and $\epsilon > 0$, there are N and δ such that if G is a graph on $n \geq N$ vertices, containing at most $\delta n^{v(H)}$ copies of H , then G is ϵ -close to some G' on the same vertex set, which contains no induced copies of H .*

We now turn to notions of quasirandomness for 3-graphs. One notion of 3-graph quasirandomness already defined in the introduction is vdisc_3 , and this can be seen as a natural generalization of disc_2 . As was shown by Kohayakawa, Rödl, and Skokan, this notion of

quasirandomness does not correspond to a general counting lemma, but rather to one which counts copies of *linear hypergraphs* [30].

It is also natural, and for certain applications necessary, to consider notions which take into account the quasirandomness of both the pairs and triples in a 3-graph. The next definition contains three such notions, which are equivalent to one another (but not to vdisc_3 , see [39]).

Definition 3.8. Let $\epsilon > 0$, $n \geq 1$, and suppose $H = (V, E)$ is a 3-uniform hypergraph. Assume $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph underlying H , and assume d_3 is defined by $|E(H)| = d_3 |K_3^{(2)}(G)|$.

- (1) We say that (H, G) has $\text{disc}_{2,3}(\epsilon)$ if for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has $\text{disc}_2(\epsilon)$, and for every subgraph $G' \subseteq G$,

$$||E(H) \cap K_3^{(2)}(G')| - d_3 |K_3^{(2)}(G')|| \leq \epsilon |V_1| |V_2| |V_3|.$$

- (2) We say that (H, G) has $\text{dev}_{2,3}(\epsilon)$ if for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has $\text{disc}_2(\epsilon)$, and

$$\sum_{u_0, u_1} \sum_{v_0, v_1} \sum_{w_0, w_1} \prod_{\epsilon \in \{0,1\}^3} f_{(H,G)}(u_{\epsilon_1}, v_{\epsilon_2}, w_{\epsilon_3}) \leq \epsilon |V_1|^2 |V_2|^2 |V_3|^2,$$

where $f_{(H,G)}(u, v, w) = 1_{K_3^{(2)}(G)}(u, v, w)(1_{E(H)}(u, v, w) - d_3)$.

- (3) We say that (H, G) has $\text{oct}_{2,3}(\epsilon)$ if for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has $\text{oct}_2(\epsilon)$, and H contains at most $(d_3^8 d_2^{12} + \epsilon) |V_1|^2 |V_2|^2 |V_3|^2$ copies of $K_{2,2,2}^{(3)}$.

Like quasirandomness in graphs, these notions reflect the behavior of random 3-graphs. For example, for all $\eta > 0$, a sufficiently large 3-partite 3-graph has $\text{disc}_3(\eta)$ with high probability. This can be used to prove the following well known fact which we will use later on in the paper.

Fact 3.9. For all $\eta > 0$, there is n_0 such that for all $n \geq n_0$ there exists a 3-partite 3-graph $H_n = (U \cup W \cup Z, E)$ such that $(H_n, K_3[U, W, Z])$ has $\text{disc}_3(\eta)$ and density $1/2 \pm \eta$.

As mentioned in the introduction, to obtain a general counting lemma, one requires a more subtle definition, which maintains control over the relative quasirandomness of the edge decomposition.

Definition 3.10. Let $\epsilon_1, \epsilon_2 > 0$, and suppose $H = (V, E)$ is a 3-uniform hypergraph. Assume $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph underlying H , and suppose d_3 is defined by $|E(H)| = d_3 |K_3^{(2)}(G)|$.

- (1) We say that (H, G) has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2)$ if there is $d_2 \in (0, 1]$ such that for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has $\text{disc}_2(\epsilon_2, d_2)$, and for every subgraph $G' \subseteq G$,

$$||E(H) \cap K_3^{(2)}(G')| - d_3 |K_3^{(2)}(G')|| \leq \epsilon_1 d_2^3 |V_1| |V_2| |V_3|.$$

- (2) We say that (H, G) has $\text{dev}_{2,3}(\epsilon_1, \epsilon_2)$ if there is $d_2 \in (0, 1]$ such that for each $1 \leq i < j \leq 3$, the graph $G[V_i, V_j]$ has $\text{disc}_2(\epsilon_2, d_2)$, and

$$\sum_{u_0, u_1} \sum_{v_0, v_1} \sum_{w_0, w_1} \prod_{\epsilon \in \{0,1\}^3} f_{(H,G)}(u_{\epsilon_1}, v_{\epsilon_2}, w_{\epsilon_3}) \leq \epsilon_1 d_2^{12} |V_1|^2 |V_2|^2 |V_3|^2.$$

where $f_{(H,G)}(u, v, w) = 1_{K_3^{(2)}(G)}(u, v, w)(1_{E(H)}(u, v, w) - d_3)$.

- (3) We say that (H, G) has $\text{oct}_{2,3}(\epsilon_1, \epsilon_2)$ if there is $d_2 \in (0, 1]$ such that for each $1 \leq i \neq j \leq 3$, the graph $G[V_i, V_j]$ has $\text{cycle}_2(\epsilon_2, d_2)$, and the number of copies of $K_{2,2,2}^{(3)}$ in H is at most $d_3^8 d_2^{12} |V_1|^2 |V_2|^2 |V_3|^2 + \epsilon_1 d_2^{12} |V_1|^2 |V_2|^2 |V_3|^2$.

In applications, ϵ_2 will usually be much smaller than ϵ_1 . The history of these notions and their corresponding regularity and counting lemmas is explained in full in [39], for example. Concerning the results that we shall use, a regularity and counting lemma for $\text{dev}_{2,3}$ and $\text{oct}_{2,3}$ was first proved by Gowers in [26], while Rödl and Frankl proved a regularity lemma for $\text{disc}_{2,3}$ in addition to a counting lemma for a stronger version of $\text{disc}_{2,3}$ in [24]. However, it has been shown that the notions in Definition 2.5 and Definition 3.10 are all equivalent in the following sense.

Proposition 3.11. *For all $d_3, \epsilon_3 > 0$ there exists $\delta_3 > 0$ such that for all $d_2, \epsilon_2 > 0$ there exists $\delta_2 > 0$ and $n_0 \geq 1$ such that the following hold. Suppose $H = (V, E)$ is a 3-graph. Assume $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph underlying H with $|V_1| = |V_2| = |V_3| = n$. Suppose the density of each $G[V_i, V_j]$ is in $(d_2 - \delta_2, d_2 + \delta_2)$ and $e(H) = d_3 |K_3^{(2)}(G)|$.*

- (1) (Nagle et al. [39]) *If (H, G) has $\text{disc}_{2,3}(\delta_3, \delta_2)$ then it also has $\text{oct}_{2,3}(\epsilon_3, \epsilon_2)$.*
- (2) (Dementieva et al. [19]) *If (H, G) has $\text{oct}_{2,3}(\delta_3, \delta_2)$ then it also has $\text{disc}_{2,3}(\epsilon_3, \epsilon_2)$.*
- (3) (Nagle et al. [39]) *If (H, G) has $\text{oct}_{2,3}(\delta_3, \delta_2)$ then it also has $\text{dev}_{2,3}(\epsilon_3, \epsilon_2)$.*
- (4) (Gowers [26]) *If (H, G) has $\text{dev}_{2,3}(\delta_3, \delta_2)$ then it also has $\text{oct}_{2,3}(\epsilon_3, \epsilon_2)$.*

Proposition 3.11 shows that Definition 2.11 could be equivalently formulated with any of these three notions of quasirandomness.

In order to apply a hypergraph regularity lemma, one needs a corresponding counting lemma. Theorem 3.12 was first proven by Gowers in [26] for $\text{dev}_{2,3}$, but we state it in terms of $\text{disc}_{2,3}$ (via Proposition 3.11).

Theorem 3.12 (Counting lemma for regular 3-graphs). *For all $t \in \mathbb{N}$ and $\xi, d_3 > 0$, there exists $\delta_3 > 0$ such that for every $d_2 > 0$, there exists $\delta_2 > 0$ and $n_0 \in \mathbb{N}$ such that the following holds.*

Let $G = (V, E)$ be a t -partite graph with vertex partition $V = V_1 \cup \dots \cup V_t$, where $|V_1| = \dots = |V_t| = n \geq n_0$, and let $H = (V, R)$ be a 3-uniform hypergraph on vertex set V such that $R \subseteq K_3^{(2)}(G)$. Suppose that for each $ij \in \binom{[t]}{2}$, $G[V_i, V_j]$ has $\text{disc}_2(\delta_2, d_2)$. For each $ijk \in \binom{[t]}{3}$, let $G^{ijk} = G[V_i, V_j, V_k]$, $H^{ijk} = (V_i \cup V_j \cup V_k, K_3^{(2)}(G^{ijk}) \cap R)$, and let d_{ijk} be such that $e(H^{ijk}) = d_{ijk} |K_3^{(2)}(G^{ijk})|$. Suppose that for each $ijk \in \binom{[t]}{3}$, $d_{ijk} \geq d_3$, and

(H^{ijk}, G^{ijk}) satisfies $\text{disc}_3(\delta_3, \delta_2)$. Then

$$\left| |K_t^{(3)}(H)| - d_2^{\binom{t}{2}} dn^t \right| \leq \xi d_2^{\binom{t}{2}} dn^t$$

where $d = \prod_{ijk \in \binom{[t]}{3}} d_{ijk}$.

An induced analogue of the above was proven by Rödl and Schacht in [46].

Theorem 3.13 (Induced Counting Lemma). *For all $t \in \mathbb{N}$ and $\xi, d_3 > 0$, there exists $\delta_3 > 0$ such that for every $d_2 > 0$, there exists $\delta_2 > 0$ and $n_0 \in \mathbb{N}$ such that the following holds.*

Let $F = ([t], R_F)$ be a t -partite, 3-graph. Let $G = (V, E)$ be a t -partite graph with vertex partition $V = V_1 \cup \dots \cup V_t$, where $|V_1| = \dots = |V_t| = n \geq n_0$, and let $H = (V, R)$ be a 3-uniform hypergraph on vertex set V such that $R \subseteq K_3^{(2)}(G)$. Suppose that for each $ij \in \binom{[t]}{2}$, $G[V_i, V_j]$ has $\text{disc}_2(\delta_2, d_2)$. For each $ijk \in \binom{[t]}{3}$, let $G^{ijk} = G[V_i, V_j, V_k]$, $H^{ijk} = (V_i \cup V_j \cup V_k, K_3^{(2)}(G^{ijk}) \cap R)$, and let d_{ijk} be such that $e(H^{ijk}) = d_{ijk} |K_3^{(2)}(G^{ijk})|$.

Suppose that for each $ijk \in R_F$, $d_{ijk} \geq d_3$ and for each $ijk \in \binom{[t]}{3} \setminus R_F$, $d_{ijk} \leq 1 - d_3$. Suppose further that (H^{ijk}, G^{ijk}) satisfies $\text{disc}_{2,3}(\delta_3, \delta_2)$ for each $ijk \in \binom{[t]}{3}$. Let m_F be the number of tuples $(v_1, \dots, v_t) \in \prod_{i=1}^t V_i$ such that $v_i v_j v_k \in R$ if and only if $ijk \in R_F$. Then

$$\left| m_F - d_2^{\binom{t}{2}} dn^t \right| \leq \xi d_2^{\binom{t}{2}} dn^t$$

where $d = \prod_{ijk \in \binom{[t]}{3}} d_{ijk}$.

As an application of Theorem 3.13, Rödl and Schacht [46] proved the following induced removal lemma.

Theorem 3.14 (Induced removal for 3-graphs). *For all $\eta > 0$ and finite 3-graphs G , there are $c > 0$, $C > 0$, and n_0 such that the following holds. Suppose H is a 3-graph on $n \geq n_0$ vertices, and H contains at most $cn^{v(G)}$ -many induced copies of G . Then H is η -close to some H' containing no induced copies of G .*

In fact, Rödl and Schacht's results hold for k -graphs for any $k \geq 3$, but we will only use the case $k = 3$. We will also use the following, even stronger result for hereditary properties, also from [46].

Theorem 3.15 (Induced removal for hereditary properties). *Suppose \mathcal{H} is a hereditary 3-graph property. Then for all $\eta > 0$ there are $c > 0$, $C > 0$, and n_0 such that the following holds. Suppose H is a 3-graph on $n \geq n_0$ vertices, and for every $G \in \mathcal{F}_{\mathcal{H}}$ on at most C vertices, H contains at most $cn^{v(G)}$ copies of G . Then H is η -close to some $H' \in \mathcal{H}_n$.*

This theorem can be used to give another characterization of \sim -classes.

Definition 3.16. Given a hereditary 3-graph property \mathcal{H} , let $\Delta(\mathcal{H})$ denote all finite 3-graphs G such that for all $\epsilon > 0$, there is $N = N(G, \mathcal{H}, \epsilon)$ so that if $n \geq N$ and $H \in \mathcal{H}_n$, then H contains at most $\epsilon n^{v(G)}$ -many induced sub-3-graphs isomorphic to G .

In other words, $\Delta(\mathcal{H})$ is the set of finite 3-graphs G so that large elements of \mathcal{H} cannot contain too many copies of G . Theorem 3.15 implies that this definition characterizes \sim -classes.

Proposition 3.17. $\mathcal{H} \sim \mathcal{H}'$ if and only if $\Delta(\mathcal{H}) = \Delta(\mathcal{H}')$.

For a proof, see [46].

3.2. Lemmas for slicing, intersecting, and combining decompositions. This subsection contains several facts which will be useful for building regular decompositions. While most of these are fairly standard, we include several proofs in Appendix B for the sake of completeness. First, Lemma 3.18 tells us that large sub-pairs in quasirandom graphs are still quasirandom.

Proposition 3.18 (Sub-pairs lemma). *Suppose $G = (A \cup B, E)$ is a bipartite graph and $|E| = d|A||B|$. Suppose $A' \subseteq A$ and $B' \subseteq B$ satisfy $|A'| \geq \gamma|A|$ and $|B'| \geq \gamma|B|$ for some $\gamma \geq \epsilon$, and G satisfies $\text{disc}_2(\epsilon, d)$. Then $G' := (A' \cup B', G[A', B'])$ satisfies $\text{disc}_2(\epsilon', d')$ where $\epsilon' = 2\gamma^{-2}\epsilon$ and $d' \in (d - \gamma^{-2}\epsilon, d + \gamma^{-2}\epsilon)$.*

We refer the reader to [32] for a proof. Lemma 3.19 is a similar result for triples, and says that large sub-triples of $\text{disc}_{2,3}$ -regular triads are still somewhat regular.

Lemma 3.19 (Sub-triples lemma). *For all $k \geq 1$ and $\delta_3 > 0$, there exists $\delta'_3 > 0$ such that for all $d_2, d_3 \in (0, 1]$ and $0 < \delta_2 \leq d_2^3/8$, there are $\delta'_2 > 0$ and $m_0 \geq 1$ so that the following holds. Suppose $V = V_1 \cup V_2 \cup V_3$ where $|V_1|, |V_2|, |V_3| \geq m_0$, and $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph such that for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta'_2, d_2)$. Assume $H = (V, R)$ is a 3-graph, and (H', G) has $\text{disc}_{2,3}(\delta'_2, \delta'_3)$ with density d_3 , where $H' = (V, R \cap K_3^{(2)}(G))$.*

Suppose that for each $i \in [3]$, $V'_i \subseteq V_i$ satisfies $|V'_i| \geq |V_i|/k$. Then (H'', G') has $\text{disc}_{2,3}(\delta_3, \delta_2)$ with density $d'_3 = d_3 \pm \delta_3$, where $G' = G[V'_1 \cup V'_2 \cup V'_3]$, and $H'' = H'[V'_1 \cup V'_2 \cup V'_3]$.

The proof consists of standard arguments and appears in Appendix B. The next couple of propositions tell us that homogeneity (i.e. density near zero or one) implies quasirandomness, both in the $\text{disc}_{2,3}$ and vdisc_3 -sense, thereby generalizing the following easy fact about graphs.

Fact 3.20. *Suppose $H = (U \cup V, R)$ is a bipartite graph. For all $\epsilon > 0$, if $|R \cap K_2[U, V]| \leq \epsilon|K_2[U, V]|$. Then H satisfies $\text{disc}_2(\epsilon)$.*

Proposition 3.21. *If $H = (V_1 \cup V_2 \cup V_3, R)$ is a 3-partite 3-graph and $|R \cap K_3[V_1, V_2, V_3]| \leq \epsilon|V_1||V_2||V_3|$, then H has $\text{vdisc}_3(\epsilon)$.*

Proof. Suppose for each $i \in [3]$, $V'_i \subseteq V_i$. Let d be such that $|R \cap K_3[V_1, V_2, V_3]| = d|K_3[V_1, V_2, V_3]|$. Since $d \leq \epsilon$,

$$|R \cap K_3[V'_1, V'_2, V'_3]| \leq \epsilon|V_1||V_2||V_3| \leq \epsilon|V_1||V_2||V_3| + d|V'_1||V'_2||V'_3|.$$

Clearly $|R \cap K_3[V'_1, V'_2, V'_3]| \geq 0 > d|V'_1||V'_2||V'_3| - \epsilon|V_1||V_2||V_3|$. Thus

$$||R \cap K_3[V'_1, V'_2, V'_3]| - d|V'_1||V'_2||V'_3|| \leq \epsilon|V_1||V_2||V_3|,$$

as desired. \square

Proposition 3.22. *For all $0 < \epsilon < 1/2$ and $d_2 > 0$, there is $\delta_2 > 0$ such that for all $0 < \delta \leq \delta_2$, there is N such that the following holds. Suppose $H = (V_1 \cup V_2 \cup V_3, R)$ is a 3-partite 3-graph on $n \geq N$ vertices, and for each $i, j \in [3]$, $||V_i| - |V_j|| \leq \delta|V_i|$. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph, where for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta, d_2)$, and assume*

$$|R \cap K_3^{(2)}(G)| \leq \epsilon |K_3^{(2)}(G)|.$$

Then (H', G) has $\text{disc}_{2,3}(\delta, \sqrt{\epsilon})$ where $H' = (V_1 \cup V_2 \cup V_3, R \cap K_3^{(2)}(G))$.

Proof. Suppose $0 < \epsilon < 1/2$ and $0 < d_2$. Let δ_1 and N be from Corollary 3.6 for $\gamma = 1/32$, and set $\delta = \min\{\epsilon^2/32, \delta_2/32\}$. Suppose $0 < \delta \leq \delta_2$, $H = (V_1 \cup V_2 \cup V_3, R)$ is a 3-partite 3-graph on $n \geq N$ vertices such that for each $i, j \in [3]$, $||V_i| - |V_j|| \leq \delta|V_i|$. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph, where for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta, d_2)$, and assume

$$|R \cap K_3^{(2)}(G)| \leq \epsilon |K_3^{(2)}(G)|.$$

By Corollary 3.6, $|K_3^{(2)}(G)| = d_2^3(1 \pm \gamma)|V_1||V_2||V_3|$. Suppose $G' \subseteq G$ is a subgraph, and let d be such that $|R \cap K_3^{(2)}(G)| = d|K_3^{(2)}(G)|$. Then

$$|R \cap K_3^{(2)}(G')| \leq \epsilon |K_3^{(2)}(G)| \leq \epsilon d_2^3(1 + \gamma)^3|V_1||V_2||V_3| \leq \sqrt{\epsilon} d_2^3|V_1||V_2||V_3|,$$

where the last inequality is by our choices of γ and since $\epsilon < 1/2$. Using that $\epsilon |K_3^{(2)}(G)| \leq \sqrt{\epsilon} d_2^3|V_1||V_2||V_3|$, $d \leq \epsilon$, and $K_3^{(2)}(G') \subseteq K_2^{(2)}(G)$, we have that

$$|R \cap K_3^{(2)}(G')| \geq 0 \geq d|K_3^{(2)}(G')| - \epsilon |K_3^{(2)}(G)| \geq d|K_3^{(2)}(G')| - \sqrt{\epsilon} d_2^3|V_1||V_2||V_3|.$$

Thus $||R \cap K_3^{(2)}(G')| - d|K_3^{(2)}(G')|| \leq \sqrt{\epsilon} d_2^3|V_1||V_2||V_3|$. This concludes the proof that (H, G) has $\text{disc}_{2,3}(\sqrt{\epsilon}, \delta)$. \square

Our next set of lemmas deals with combining and partitioning quasirandom graphs. First, we observe that adding or subtracting quasirandom graphs results in graphs which are still somewhat quasirandom.

Fact 3.23. *Suppose E_1 and E_2 are disjoint subsets of $K_2[U, V]$.*

- (1) *If $(U \cup V, E_1)$ has $\text{disc}_2(\epsilon_1, d_1)$, $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon_2, d_2)$, then $(U \cup V, E_1 \cup E_2)$ has $\text{disc}_2(\epsilon_1 + \epsilon_2, d_2 + d_1)$.*
- (2) *If $(U \cup V, E_1 \cup E_2)$ has $\text{disc}_2(\epsilon, d)$ and $(U \cup V, E_1)$ has $\text{disc}_2(\epsilon_1, d_1)$, then $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon + \epsilon_1, d - d_1)$.*

For a proof, see Appendix B. Our next lemma is due to Frankl and Rödl [24, Lemma 3.8], and allows us to partition quasirandom graphs into several equally sized, quasirandom subgraphs.

Lemma 3.24. *For all $\epsilon > 0$, $\rho \geq 2\epsilon$, $0 < p < \rho/2$, and $\delta > 0$, there is $m_0 = m_0(\epsilon, \rho, \delta)$ such that the following holds. Suppose $|U| = |V| = m \geq m_0$, and $G = (U \cup V, E)$ is a bipartite graph satisfying $\text{disc}_2(\epsilon)$ with density ρ . Then if $\ell = \lceil 1/p \rceil$, and $\epsilon \geq 10(1/\ell m)^{1/5}$, then there is a partition $E = E_0 \cup E_1 \cup \dots \cup E_\ell$, such that*

- (i) For each $1 \leq i \leq \ell$, $(U \cup V, E_i)$ has $\text{disc}_2(\epsilon)$ with density $\rho p(1 \pm \delta)$, and
- (ii) $|E_0| \leq \rho p(1 + \delta)m^2$.

Further, if $1/p \in \mathbb{Z}$, then $E_0 = \emptyset$.

Lemma 3.24 will come in handy frequently as we build decompositions. An example of such an application is Lemma 3.25 below, which shows that given a decomposition \mathcal{P} of a vertex set V , we can change a small number of elements from $\mathcal{P}_{\text{edge}}$ to obtain a decomposition \mathcal{P}' of V which has no disc_2 -irregular elements in $\mathcal{P}'_{\text{edge}}$.

Lemma 3.25. *For all $\epsilon_1 > 0$ and non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are $\epsilon'_1 > 0$ and $\epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ such that for all $\ell, t \geq 1$, there is N such that the following holds. Suppose V is a set of size $n \geq N$, and \mathcal{P}' is an $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V . Then there is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V such that the following hold.*

- (1) $\mathcal{P}_{\text{vert}} = \mathcal{P}'_{\text{vert}}$,
- (2) Every $P \in \mathcal{P}_{\text{edge}}$ has $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$,
- (3) If $P' \in \mathcal{P}'_{\text{edge}}$ satisfied $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, then $P' \in \mathcal{P}_{\text{edge}}$.

Proof. Fix $\epsilon_1 > 0$ and a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Set $\epsilon'_1 = \epsilon_1^2/4$, and for all $x \in \mathbb{N}$, set $\epsilon'_2(x) = \epsilon_2(x^2)/x^3$. Given $\ell, t \geq 1$, let $N = \max\{tm_0(s\epsilon'_2(\ell), 1 - s/\ell, s\epsilon'_2(\ell)) : 1 \leq s \leq (1 - \sqrt{\epsilon'_1})\ell\}$, where $m_0(\cdot, \cdot, \cdot)$ is from Lemma 3.24.

Suppose V is a set of size $n \geq N$, and \mathcal{P}' is a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V , consisting of $\{V_i : i \in [t]\}$ and $\{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Set $m = n/t$, and define $\Delta = \{P \in \text{Triads}(\mathcal{P}') : P \text{ has } \text{disc}_2(\epsilon'_2(\ell), 1/\ell)\}$. Then let

$$\Omega = \{xyz \in \binom{V}{3} : \text{for some } P \in \text{Triads}(\mathcal{P}') \setminus \Delta, P \cap \{xy, yz, xz\} \neq \emptyset\}.$$

In other words, Ω is the set of triples who intersect a disc_2 -irregular element from $\mathcal{P}'_{\text{edge}}$. Then define

$$\Sigma = \{ij \in \binom{[t]}{2} : |K_2[V_i, V_j] \setminus (\bigcup \Delta)| \geq \sqrt{\epsilon'_1} n^2/t^2\}.$$

Note that if $ij \in \Sigma$, then for all $k \in [t] \setminus \{i, j\}$, $K_3[V_i, V_j, V_k]$ contains at least $\sqrt{\epsilon'_1} n^3/t^3$ many triples from Ω , and thus, at least $(t-2)\sqrt{\epsilon'_1} n^3/t^3$ triples in Ω intersect $K_2[V_i, V_j]$. Since by assumption, $|\Omega| \leq \epsilon'_1 n^3$, this implies $|\Sigma| \leq 3\sqrt{\epsilon'_1} t^2$. We now define a new partition of $K_2[V_i, V_j]$, for each $ij \in \binom{[t]}{2}$. We do this in cases, depending on whether $ij \in \Sigma$ or $ij \notin \Sigma$.

Suppose first $ij \in \Sigma$. In this case, apply Lemma 3.24 to $K_2[V_i, V_j] = E_1$ with $\epsilon = \delta = \epsilon'_2(\ell)$ and $p = 1/\ell$ to obtain a partition $K_2[V_i, V_j] = Q_{ij}^1 \cup \dots \cup Q_{ij}^\ell$ such that for each $1 \leq \alpha \leq \ell$, Q_{ij}^α has $\text{disc}_2(2\epsilon'_2(\ell), 1/\ell)$. By definition of $\epsilon'_2(\ell)$, this shows each such Q_{ij}^α has $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$.

Suppose now $ij \notin \Sigma$. Let $P_{ij}^{\alpha_1}, \dots, P_{ij}^{\alpha_s}$ enumerate the elements of Δ_{ij} and set $E_0 = K_2(V_i, V_j) \setminus \bigcup_{u=1}^s P_{ij}^{\alpha_u}$. If $s = \ell$, then let $Q_{ij}^\alpha = P_{ij}^\alpha$ for each $\alpha \leq \ell$. Assume on the other hand that $s < \ell$. Then since $ij \notin \Sigma$, $|E_0| \leq \sqrt{\epsilon'_1} m^2$, so we must have that $s \geq (1 - \sqrt{\epsilon'_1})\ell$. By Fact 3.23, $E_1 \cup \dots \cup E_s$ has $\text{disc}_2(s\epsilon'_2(\ell), s/\ell)$, so its complement in $K_2[V_i, V_j]$, E_0 , has

$\text{disc}_2(s\epsilon'_2(\ell), 1-s/\ell)$. Setting $\rho_0 = |E_0|/m^2$, this implies $\rho_0 \geq \frac{1}{\ell}(1-\epsilon'_2(\ell))$, and consequently, by definition of $\epsilon'_2(\ell)$, $\rho_0 \geq 2s\epsilon'_2(\ell)$. Let $u = 2\ell(\ell-s)$ and $p = 1/u$. Then

$$p = \frac{1}{2\ell(\ell-s)} < \frac{1}{2(\ell-s)} < \frac{\ell-s}{2\ell} \leq \rho_0/2,$$

where the second inequality is because $s \geq (1-\sqrt{\epsilon'_1})\ell$. Consequently, we may apply Lemma 3.24 to E_0 with this u and p , and $\epsilon = \delta = s\epsilon'_2(\ell)$ to obtain a partition $E_0 = E'_1 \cup \dots \cup E'_u$ where for each $i \in [u]$, E'_i has of density $\frac{1}{u}(1 \pm s\epsilon'_2(\ell))$ and satisfies $\text{disc}_2(s\epsilon'_2(\ell))$. Let $I_1 \cup \dots \cup I_{\ell-s}$ be an equipartition of $[u]$ and for each $i \in [\ell-s]$, set $E''_i = \bigcup_{j \in I_i} E'_j$. By Fact 3.23, each E''_i has $\text{disc}_2(2\ell s\epsilon'_2(\ell), \frac{2(\ell-s)}{u}(1 \pm 2\ell s\epsilon'_2(\ell)))$. By our choice of $\epsilon'_2(\ell)$ and the definition of u , this means each E''_i has $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Now define \mathcal{P} by setting $\mathcal{P}_{\text{vert}} = \{V_1, \dots, V_t\}$ and $\mathcal{P}_{\text{edge}} = \{P_{ij}^{\alpha_1}, \dots, P_{ij}^{\alpha_s}, E''_1, \dots, E''_{\ell-s}\}$. Clearly each element of $\mathcal{P}_{\text{edge}}$ satisfies $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Moreover, by construction, if $P_{ij}^\alpha \in \mathcal{P}'_{\text{edge}}$ had $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$ in \mathcal{P}' , then $P_{ij}^\alpha \in \mathcal{P}_{\text{edge}}$, as desired. \square

As a corollary, we now prove Proposition 2.13, which tells us that every sufficiently large 3-graph has regular decompositions with no disc_2 -irregular triads.

PROOF OF PROPOSITION 2.13: Fix a hereditary 3-graph property \mathcal{H} . Fix $\epsilon_1 > 0$ and a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Let $\epsilon'_1 > 0$ and $\epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ be as in Lemma 3.25 for ϵ_1 and ϵ_2 . Given $t_0, \ell_0 \geq 1$, choose n_0, L_0, T_0 as in Theorem 2.8 for $\epsilon'_1, \epsilon'_2, t_0, \ell_0$, and let n_1 be as in Lemma 3.25 for $\epsilon_1, \epsilon_2, T_0, L_0$. Set $N = \{n_0, n_1\}$.

Suppose $H \in \mathcal{H}$ is a 3-graph on $n \geq N$ vertices. By Theorem 2.8, there are $t_0 \leq t \leq T_0$, $\ell_0 \leq \ell \leq L_0$, and a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition \mathcal{P}' of $V(H)$ which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular with respect to H . By Lemma 3.25, there is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V which has no disc_2 -irregular triads, and so that the elements from $\mathcal{P}'_{\text{edge}}$ which had $\text{disc}_2(\epsilon'_1, \epsilon'_2(\ell))$ in \mathcal{P}' remain unchanged in \mathcal{P} . Consequently, every triad of \mathcal{P}' which had $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$ with respect to (H, \mathcal{P}') is unchanged in \mathcal{P} . Since $\epsilon'_1 \leq \epsilon_1$ and $\epsilon'_2(\ell) \leq \epsilon_2(\ell)$, this implies that \mathcal{P} is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . \square

Our next series of lemmas will be useful for situations where we intersect two different decompositions of a vertex set V . First, we make a definition that describes when one decomposition approximately refines another.

Definition 3.26. Suppose V is a set and \mathcal{P}, \mathcal{Q} are decompositions of V . Then \mathcal{P} is an (ϵ_1, ϵ_2) -approximate refinement of \mathcal{Q} if there exists $\Sigma \subseteq \binom{\mathcal{P}_{\text{vert}}}{2}$ such that the following hold.

- (1) For all $V_i \in \mathcal{P}_{\text{vert}}$ there is $W_j \in \mathcal{Q}_{\text{vert}}$ such that $|V_i \setminus W_j| < \epsilon_1 |V_i|$,
- (2) For all $P \in \mathcal{P}_{\text{edge}}$ with $V_i V_j \notin \Sigma$, there $Q \in \mathcal{Q}_{\text{edge}}$ with $|P \setminus Q| \leq \epsilon_1 |P|$, and such that $P \setminus Q$ has $\text{disc}_2(\epsilon_2)$.
- (3) $|\Sigma| \leq \epsilon_1 \binom{\mathcal{P}_{\text{vert}}}{2}$.

Given two decompositions \mathcal{P} and \mathcal{Q} of the same vertex set V , we can find an approximate common refinement of both which is a $(t', \ell', \epsilon'_1, \epsilon'_2(\ell'))$ -decomposition of V .

Lemma 3.27. For all $\epsilon_1 > 0$, non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are functions $f, g : \mathbb{N}^4 \rightarrow (0, 1]$ such that for all $\ell, t, s, r \geq 1$, there is $N \geq 1$ such that the following holds.

Suppose V is a set of size $n \geq N$, \mathcal{P} is an (ℓ, t) -decomposition of V , and \mathcal{Q} is an (r, s) -decomposition of V . Then there are $\ell' \leq g(r, s, \ell, t)$ and $t' \leq f(r, s, \ell, t)$, and a $(t', \ell', \epsilon_1, \epsilon_2(\ell'))$ -decomposition \mathcal{R} of V such that every $R_{ij}^\alpha \in \mathcal{R}_{\text{edge}}$ has $\text{disc}_2(\epsilon_2(\ell'), 1/\ell')$, and so that \mathcal{R} is an $(\epsilon_1, \epsilon_2(\ell'))$ -approximate refinement of $\mathcal{P} \wedge \mathcal{Q}$.

The proof of Lemma 3.27 appears in Appendix B. Finally, it will be useful to know that approximate refinements of homogeneous decompositions are still fairly homogeneous.

Lemma 3.28. *For all $\epsilon_1 > 0$ and non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are $\epsilon'_1 > 0$, $\epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ and such that for all $\ell, t, \ell', t' \geq 1$, there is N such that the following holds. Suppose $H = (V, E)$ is a 3-graph on $n \geq N$ vertices, and \mathcal{Q} is a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular and ϵ'_1 -homogeneous with respect to H .*

Suppose \mathcal{P} is a $(t', \ell', \epsilon'_1, \epsilon'_2(\ell'))$ -decomposition of V which is a $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement of \mathcal{Q} , witnessed by the set $\Sigma \subseteq \binom{[t']}{2}$. Then \mathcal{P} is a $(t', \ell', \epsilon_1, \epsilon_2(\ell'))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell'))$ -regular and ϵ_1 -homogeneous with respect to H .

For a proof, see Appendix B.

3.3. Ramsey facts. It will be convenient at times to know some simple Ramsey facts about hereditary properties and the combinatorial definitions from the introduction. For example, if G is a 3-graph and $\overline{H}(k) \cong \text{Trip}(G)$, then by definition there are $a_1, \dots, a_k \in V(G)$ and $b_1, \dots, b_k, d \in V(G)$ and so that $a_i b_j d \in E$ if and only if $i \leq j$. However, it is possible that the sets $\{a_1, \dots, a_k\}$ and $\{b_1, \dots, b_k\}$ are not disjoint, so we do not necessarily know the size of the set $\{a_1, \dots, a_k, b_1, \dots, b_k\}$. This can be inconvenient at times, and is the motivation for the following definition.

Definition 3.29. Suppose $H = (U \cup V \cup W, E)$ is a 3-partite 3-graph. A *clean copy* of H is a 3-graph $G = (V, E)$ such that $\text{Trip}(G) \cong H$ and $|V(G)| = |U| + |V| + |W|$.

Using standard Ramsey arguments, we will show that we can usually assume that there are clean copies of the configurations we are interested in. For example, the following is a stronger version of Fact 2.22, the universality property corresponding to IP_2 .

Fact 3.30. *If \mathcal{H} has IP_2 then \mathcal{H} contains a clean copy of every 3-partite 3-graph.*

For a proof, see Appendix C. Similarly, we will use the following facts.

Lemma 3.31. *Suppose \mathcal{H} is a hereditary 3-graph property.*

- (1) *Suppose $H^*(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Then \mathcal{H} contains a clean copy of $H^*(k)$ for all $k \geq 1$.*
- (2) *If \mathcal{H} has FOP_2 , then \mathcal{H} contains a clean copy of $F(k)$ for all $k \geq 1$.*
- (3) *Suppose $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Then \mathcal{H} contains a clean copy of $\overline{H}(k)$ for all $k \geq 1$.*
- (4) *Suppose $U^*(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Then \mathcal{H} contains a clean copy of $U^*(k)$ for all $k \geq 1$.*

For proofs of these facts, see Appendix C. Before being able to state the condition we will use for $\overline{U}(k)$, we need to make one definition.

Definition 3.32. Suppose $n \geq 1$ and $G = (U \cup W, E)$ is a bipartite graph. Set

$$n \otimes G := (\{U \cup W \cup \{c_1, \dots, c_n\}, \{c_i uw : uw \in E\}\}.$$

In other words, $n \otimes G$ is the 3-graph obtained by adjoining n new vertices to the graph G . For example, note that $\overline{H}(k) = k \otimes H(k)$ and $\overline{U}(k) = k \otimes U(k)$. We will show that arbitrarily large $\overline{U}(k)$ come with a universality property for 3-graphs of the form $n \otimes G$.

Lemma 3.33. *Suppose \mathcal{H} is a hereditary 3-graph property and $\overline{U}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Then for every bipartite graph G and $n \geq 1$, \mathcal{H} contains a clean copy of $n \otimes G$.*

Again, we refer the reader to Appendix C for a proof.

4. VC₂-DIMENSION AND disc_{2,3}-HOMOGENEITY

In this section we show that a hereditary 3-graph property is vdisc₃-homogeneous if and only if it has finite VC₂-dimension.

The first step is to show that a sufficiently disc_{2,3}-regular triad in a large 3-graph of bounded VC₂-dimension must have density near 0 or 1 (see Proposition 4.2). The crucial ingredient for this is the following counting lemma for induced 3-partite 3-graphs. It follows easily from Theorem 3.13. For completeness, we include a proof in Appendix B.

Theorem 4.1 (Counting lemma for induced 3-partite 3-graphs). *For all ξ , $d_3 > 0$ and $t \geq 1$, there exists $\delta_3 > 0$ such that for every $d_2 > 0$, there exists $\delta_2 > 0$ and $N \geq 1$ such that the following holds.*

Suppose $t_1, t_2, t_3 \geq 1$, $t_1 + t_2 + t_3 = t$, and $F = (U \cup W \cup Z, R_F)$ is a 3-partite 3-graph with $U = \{u_1, \dots, u_{t_1}\}$, $W = \{w_1, \dots, w_{t_2}\}$, and $Z = \{z_1, \dots, z_{t_3}\}$. Suppose V is a vertex set of size $n \geq N$, $H = (V, R)$ is a 3-graph, and $G = (V, E)$ is a t -partite graph, with vertex partition

$$V = A_1 \cup \dots \cup A_{t_1} \cup B_1 \cup \dots \cup B_{t_2} \cup C_1 \cup \dots \cup C_{t_3},$$

where each part has size $\frac{n}{t}(1 \pm \delta_2)$. For each $(i, j, s) \in [t_1] \times [t_2] \times [t_3]$, set $G^{ijs} = G[A_i, B_j, C_s]$ and $H^{ijs} = (A_i \cup B_j \cup C_s, K_3^{(2)}(G^{ijs}) \cap R)$, and let d_{ijs} be such that $e(H^{ijs}) = d_{ijs} |K_3^{(2)}(G^{ijs})|$. Assume that for each $(i, j, s) \in [t_1] \times [t_2] \times [t_3]$, each of $G[A_i, B_j]$, $G[A_i, C_s]$, and $G[B_j, C_s]$ have disc₂(δ_2, d_2), (H^{ijs}, G^{ijs}) satisfies disc_{2,3}(δ_3, δ_2), and

$$\begin{cases} d_{ijs} \geq d_3 & \text{if } u_i v_j z_s \in R_F, \text{ and} \\ d_{ijs} \leq (1 - d_3) & \text{if } u_i v_j z_s \notin R_F. \end{cases}$$

Then the number of tuples $\bar{a}\bar{b}\bar{c} \in \prod_{i=1}^{t_1} A_i \times \prod_{i=1}^{t_2} B_i \times \prod_{i=1}^{t_3} C_i$ such that $a_i b_j c_s \in E(H)$ if and only if $u_i w_j z_s \in R_F$ is

$$d_2^{(t)} d(1 \pm \xi) \prod_{i=1}^{t_1} |A_i| \prod_{i=1}^{t_2} |B_i| \prod_{i=1}^{t_3} |C_i|,$$

where $d = (\prod_{u_i v_j w_s \in R_F} d_{ijs})(\prod_{u_i v_j w_s \notin R_F} (1 - d_{ijs}))$.

We now prove that $\text{disc}_{2,3}$ -regular triads in 3-graphs of bounded VC_2 -dimension have density near 0 or near 1. We do this by showing that a large $\text{disc}_{2,3}$ -regular triad with intermediate density can be broken up into sub-triads, all of which are regular and of intermediate density (this uses Lemma 3.19). We then apply Theorem 4.1 to the resulting configuration to find a copy of $V(k)$, and thus a contradiction.

Proposition 4.2 ($\text{VC}_2 < \infty$ implies regular triples have trivial density). *For all $k \geq 1$ and $\epsilon \in (0, 1/2)$, there exists a $\delta_3 > 0$ such that for every $d_2 > 0$ there is $\delta_2 > 0$ and M_0 such that the following holds. Suppose $V = V_1 \sqcup V_2 \sqcup V_3$, where V is a set of size $n \geq M_0$, and $|V_i| = \frac{n}{3}(1 \pm \delta_2)$ for each $i \in [3]$. Suppose $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph, and $H = (V, R)$ is a 3-graph with $\text{VC}_2(H) < k$. Assume that for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta_2, d_2)$. Let $H' = (V, R \cap K_3^{(2)}(G))$, and assume (H', G) has $\text{disc}_{2,3}(\delta_2, \delta_3)$. Then*

$$\frac{|R \cap K_3^{(2)}(G)|}{|K_3^{(2)}(G)|} \in [0, \epsilon] \cup (1 - \epsilon, 1].$$

Proof. Fix $k \geq 1$ and $\epsilon \in (0, 1/2)$. First, apply Theorem 4.1 to obtain δ_3''' for $t = 2k + 2^k$, $\xi = 1/2$, and $d_3 = \epsilon/2$. Fix any $d_2 > 0$. Apply Theorem 4.1 to obtain δ_2'' and n_0 for $t = 2^{k^2} + k^2$, ξ , d_3 , δ_3''' , and d_2 . Let $\delta_3'' = \min\{\delta_3''', \epsilon/4\}$ and $\delta_2'' = \min\{\delta_2''', d_2^3/8\}$, and apply Lemma 3.19 to $d_2, d_3, \delta_3'', \delta_2''$ to obtain δ_2', δ_3' and m_0 . Finally, set $\delta_2 = \min\{\delta_2'/(3k2^{1+k^2}), \delta_2''/(3k2^{1+k^2})\}$, $\delta_3 = \delta_3'$, and choose $M_0 \geq \delta_2^{-1} \max\{m_0, n_0\}2^k$ sufficiently large so that $d_2^{(t)}(\epsilon/2)^{(t)}(M_0)^t/2 > 0$.

Now suppose $V = V_1 \cup V_2 \cup V_3$ where $|V| = n \geq M_0$, and for each $i \in [3]$, $|V_i| = \frac{n}{3}(1 \pm \delta_2)$. Assume $G = (V_1 \cup V_2 \cup V_3, E)$ a 3-partite graph, and $H = (V, R)$ is a 3-graph with $\text{VC}_2(H) < k$. Set $H' = (V, R \cap K_3^{(2)}(G))$. Assume that for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta_2, d_2)$, and assume (H', G) has $\text{disc}_{2,3}(\delta_2, \delta_3)$. Suppose towards a contradiction that

$$d := \frac{|R \cap K_3^{(2)}(G)|}{|K_3^{(2)}(G)|} \in [\epsilon, 1 - \epsilon].$$

For each $i \in \{1, 2, 3\}$, let $V_{i,1}, \dots, V_{i,2^k}$ be an equipartition of V_i . Note each $V_{i,j}$ has size $\frac{|V_i|}{2^k} \pm 1 = \frac{n}{3(2^k)}(1 \pm \delta_2 2^{k+1}) = \frac{n}{3(2^k)}(1 \pm \delta_2') \geq \max\{m_0, n_0\}$.

For each $(u, v, w) \in [k] \times [k] \times [2^k]$, set $G_{uvw} = G[V_{1,u}, V_{2,v}, V_{3,w}]$ and

$$H_{uvw} = (V_{1,u} \cup V_{2,v} \cup V_{3,w}, R \cap K_3^{(2)}(G_{uvw})).$$

By Lemma 3.18, each of $G[V_{1,u}, V_{2,v}]$, $G[V_{1,u}, V_{3,w}]$, and $G[V_{2,v}, V_{3,w}]$ has $\text{disc}_2(\delta_2'', d_2)$. By Lemma 3.19, (H_{uvw}, G_{uvw}) has $\text{disc}_3(\delta_2'', \delta_3'')$ with density $d_{uvw} \in [\epsilon - \delta_3'', 1 - \epsilon + \delta_3'']$. Since $\delta_3'' \leq \epsilon/4$, this implies $d_{uvw} \in (3\epsilon/4, 1 - 3\epsilon/4)$.

Since $\delta_2'' \leq \delta_2'''$ and $\delta_3'' \leq \delta_3'''$, Theorem 4.1 implies that there are at least $d_2^{(t)}(\epsilon/2)^{(t)}m^t/2$ many tuples $\bar{v}\bar{w}\bar{z} \in (\prod_{j=1}^{2^k} V_{1,j}) \times (\prod_{j=1}^k V_{2,j}) \times (\prod_{j=1}^k V_{3,j})$ such that $v_i w_j z_s \in R$ if and only if $v_i u_j w_s \in E(V(k))$. By choice of m , $d_2^{(t)}(\epsilon/2)^{(t)}m^t/2 > 0$, so $\text{Trip}(H)$ contains an induced copy of $V(k)$, contradicting that $\text{VC}_2(H) < k$. \square

We are now in a position to prove Theorem 2.23, which shows that 3-graphs with bounded VC_2 -dimension have $\text{disc}_{2,3}$ -homogeneous decompositions.

PROOF OF THEOREM 2.23: Fix $k \geq 1$. Fix $\ell_0, t_0 \geq 1$, $\epsilon_1 > 0$ and a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Apply Proposition 4.2 to $\epsilon = \epsilon_1$ to obtain $\delta_3 > 0$. For each $\ell \in \mathbb{N}$, let $\delta_2(\ell)$ and $M_0(\ell)$ be obtained by applying from Proposition 4.2 to ϵ , δ_3 , and $d_2 = 1/\ell$. Let $\epsilon_2''(x)$ be as in Corollary 3.6 for $t = 3$, ϵ_1^2 and density $1/x$. Let $\delta_2'(x)$ be from Lemma 3.18 for $\epsilon_2''(x)$ and with density $1/x$. Then define $\epsilon_2' : \mathbb{N} \rightarrow (0, 1]$ by setting $\epsilon_2'(\ell) = \min\{\epsilon_2(\ell), \epsilon_2''(\ell), \delta_2(\ell), \delta_2'(\ell)\}$ for each $\ell \in \mathbb{N}$, and let $\epsilon_1' = \min\{\epsilon_1, \delta_3\}$. Choose $T = T(\epsilon_1', \epsilon_2', t_0)$ and $N = N(\epsilon_1', \epsilon_2', t_0)$ from Theorem 2.8, and define $m_0 = \max\{M_0(T), N\}$.

Suppose $H = (V, E) \in \mathcal{H}$ and $|V| \geq m_0$. By Theorem 2.8, there are $t_0 \leq t \leq T$ and $\ell_0 \leq \ell \leq L$ and a $(t, \ell, \epsilon_1', \epsilon_2'(\ell))$ -decomposition, \mathcal{P} , of V which is $(\epsilon_1', \epsilon_2'(\ell))$ -regular with respect to H . We show every $\text{disc}_{2,3}$ -regular triad of \mathcal{P} is ϵ_1 -homogeneous with respect to H . Suppose $G_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$, $H_{ijs}^{\alpha, \beta, \tau} := (V_i \cup V_j \cup V_s, E \cap K_3^{(2)}(G_{ijk}^{\alpha, \beta, \tau}))$, and assume $(H_{ijk}^{\alpha, \beta, \tau}, G_{ijk}^{\alpha, \beta, \tau})$ has $\text{disc}_{2,3}(\delta_3, \delta_2(\ell))$. By Proposition 4.2, the density of edges in this triad is in $[0, \epsilon_1] \cup (1 - \epsilon_1, 1]$. Thus \mathcal{P} is ϵ_1 -homogeneous. Since $\epsilon_1' < \epsilon_1$ and $\epsilon_2'(\ell) < \epsilon_2(\ell)$, \mathcal{P} is also $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . \square

Equipped with Theorem 2.23, we now prove Theorem 2.24, which says that a hereditary 3-graph property is $\text{disc}_{2,3}$ -homogeneous if and only if it has bounded VC_2 -dimension.

PROOF OF THEOREM 2.24: Suppose first that $\text{VC}_2(\mathcal{H}) < \infty$, say $\text{VC}_2(\mathcal{H}) < k$. Fix $\ell_0, t_0 \geq 1$, $\epsilon_1 > 0$ and a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. By Theorem 2.23, there are T, L, N so that for all $H = (V, E) \in \mathcal{H}$ with $|V| \geq m_0$, there are $t_0 \leq t \leq T$ and $\ell_0 \leq \ell \leq L$ and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition, \mathcal{P} , of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and ϵ_1 -homogeneous with respect to H . This shows \mathcal{H} is $\text{disc}_{2,3}$ -homogeneous.

For the converse, suppose $\text{VC}_2(\mathcal{H}) = \infty$. Fix $\ell_0, t_0 \geq 1$ and $\epsilon_1 \in (0, 8^{-4})$. Define $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ as follows. Given $\ell \geq 1$, let $\gamma_\ell = \epsilon_1^3/4\ell^3$. Apply Corollary 3.6 to γ_ℓ to obtain $\epsilon_\ell > 0$. Let $\epsilon_2(\ell)$ be as in Lemma 3.18 for $d = 1/\ell$ and $\delta = \epsilon_\ell \epsilon_1^3/\ell^3$.

Fix any $T \geq t_0$ and $L \geq \ell_0$. Set $\epsilon = \epsilon_2(L)/T^3 L^3$, let n_0 be as in Fact 3.9 for ϵ . We show that for all $n \geq n_0$ there is $H = (V, E) \in \mathcal{H}$ with $|V| \geq n_0$ such that for every $\ell_0 \leq \ell \leq L$ and $t_0 \leq t \leq T$, and every $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V , if \mathcal{P} is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H , then at least $\frac{1}{9}(1 - \epsilon_1^{1/5})\binom{n}{3}$ triples $xyz \in \binom{V}{3}$ are in a triad of \mathcal{P} with density in $(1/8, 7/8)$ (so in particular \mathcal{P} cannot be ϵ_1 -homogeneous with respect to H).

Fix $n \geq n_0$. Let $H_n = (U \cup W \cup Z, E)$ be as in Fact 3.9 for ϵ , where $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_n\}$, and $Z = \{z_1, \dots, z_n\}$. By assumption, $(H_n, K_3[U, W, Z])$ has $\text{disc}_3(\epsilon)$ and density $1/2 \pm \epsilon$. By Fact 3.30, there is a clean copy of H_n in \mathcal{H} . In other words, there is $H = (A \sqcup B \sqcup C, R) \in \mathcal{H}$ where $A = \{a_1, \dots, a_n\}$, $B = \{b_1, \dots, b_n\}$, and $C = \{c_1, \dots, c_n\}$ such that $a_i b_j c_k \in R$ if and only if $u_i w_j z_k \in E$. Let $H' = (A \cup B \cup C, R \setminus (\binom{A}{2} \cup \binom{B}{2} \cup \binom{C}{2}))$, and note H' satisfies $\text{disc}_3(\epsilon)$.

Now suppose $t_0 \leq t \leq T$, $\ell_0 \leq \ell \leq L$ and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition for $V = A \cup B \cup C$. Let $V = V_1 \cup \dots \cup V_t$ be the corresponding vertex partition, and for each $1 \leq i \leq t$, set $A_i = A \cap V_i$, $B_i = B \cap V_i$ and $C_i = C \cap V_i$. Let Σ be the set of $i_1 i_2 i_3 \in \binom{[t]}{3}$ such that for some $\sigma \in S_3$, $\min\{|A_{i_{\sigma(1)}}|, |B_{i_{\sigma(2)}}|, |C_{i_{\sigma(3)}}|\} \geq \sqrt{\epsilon_1} n/t$.

Suppose $i_1 i_2 i_3 \in \Sigma$, and $\alpha, \beta, \gamma \leq \ell$ are such that (H', G') has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ where $G' = G_{i_1 i_2 i_3}^{\alpha, \beta, \gamma}$ and $H' = H_{i_1 i_2 i_3}^{\alpha, \beta, \gamma}$. Let $\sigma \in S_3$ be such that $\min\{|A_{i_{\sigma(1)}}|, |B_{i_{\sigma(2)}}|, |C_{i_{\sigma(3)}}|\} \geq \sqrt{\epsilon_1} n/t$. Set $G'' = G'[A_{i_{\sigma(1)}} \cup B_{i_{\sigma(2)}} \cup C_{i_{\sigma(3)}}]$. By Lemma 3.18, Corollary 3.6, and because each of $A_{i_{\sigma(1)}}$, $B_{i_{\sigma(2)}}$, $C_{i_{\sigma(3)}}$ are large,

$$(1) \quad |K_3^{(2)}(G'')| = \frac{1}{\ell}(1 \pm \epsilon_1^2)|A_{i_{\sigma(1)}}||B_{i_{\sigma(2)}}||C_{i_{\sigma(3)}}|.$$

Since H' has $\text{disc}_3(\epsilon)$,

$$(2) \quad ||R \cap K_3(G'')| - \frac{1}{2}|K_3(G'')|| \leq \epsilon n^3.$$

Since (H', G') has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$, if d is such that $|R \cap K_3^{(2)}(G')| = d|K_3^{(2)}(G')|$, then

$$||R \cap K_3^{(2)}(G'')| - d|K_3^{(2)}(G'')|| \leq \frac{\epsilon_1}{\ell^3}(n/t)^3.$$

By the triangle inequality, (1), and the lower bounds on the sizes of $A_{i_{\sigma(1)}}$, $B_{i_{\sigma(2)}}$, and $C_{i_{\sigma(3)}}$, we can conclude

$$\begin{aligned} |d|K_3^{(2)}(G'')| - \frac{1}{2}|K_3^{(2)}(G'')|| &\leq \frac{\epsilon_1}{\ell^3}(n/t)^3 + \epsilon n^3 \leq 2\epsilon_1^{1/4}|A_{i_{\sigma(1)}}||B_{i_{\sigma(2)}}||C_{i_{\sigma(3)}}|/\ell^3 \\ &\leq 3\epsilon_1^{1/4}|K_3^{(2)}(G'')|. \end{aligned}$$

Thus, $|d - \frac{1}{2}| \leq 3\epsilon_1^{1/4}$, so $d \in (1/8, 7/8)$ since $\epsilon_1 < 8^{-4}$.

Call a set A_i, B_i, C_i *trivial* if it has size at most $\epsilon_1^{1/4} n/t$. For each $i \in [t]$, at most two of A_i, B_i, C_i can be trivial. Consequently, the number of vertices in a trivial set is at most $3\epsilon_1^{1/4} n$. Let Γ be the set of $xyz \in K_3[A, B, C]$ such that none of x, y or z are in a trivial set. Then $|\Gamma| \geq n^3 - 3\epsilon_1^{1/4} n^3$. Since \mathcal{P} is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H , if Γ' is the set of $xyz \in \binom{V}{3}$ in a $\text{disc}_{2,3}$ -regular triad of \mathcal{P} , then $|\Gamma'| \geq \binom{3n}{3} - \epsilon_1(3n)^3 \geq 9n^3(1 - \epsilon_1)$. Therefore $|\Gamma| + |\Gamma'| > \binom{3n}{3} + n^3(1 - 3\epsilon_1^{1/4} - 9\epsilon_1)$, so

$$|\Gamma \cap \Gamma'| \geq n^3(1 - 3\epsilon_1^{1/4} - 9\epsilon_1) \geq n^3(1 - 4\epsilon_1^{1/4}) \geq \binom{|V|}{3} \frac{1}{9}(1 - 4\epsilon_1^{1/4}).$$

Since every $xyz \in \Gamma \cap \Gamma'$ is in a $\text{disc}_{2,3}$ -regular triad $G_{ijk}^{\alpha, \beta, \tau}$ with density in $(1/8, 7/8)$, this shows \mathcal{P} is not ϵ_1 -homogeneous, and thus \mathcal{H} is not $\text{disc}_{2,3}$ -homogeneous. \square

5. WEAK NIP, WEAK STABILITY, AND vdisc_3 -HOMOGENEITY

This section contains the results on weakly stable and WNIP properties. We will first show in Section 5.1 that the \sim -classes of weakly stable (respectively WNIP) properties are exactly those \mathcal{H} for which $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$ for some k (respectively $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$ for some k). In Section 5.2, we prove Theorem 2.32, characterizing the vdisc_3 -homogeneous properties as those close to a WNIP property. In Section 5.3 we show that a property admits zero $\text{disc}_{2,3}$ -error if and only if it admits zero vdisc_3 -error. In the same section we show that any property containing $\overline{H}(k)$ for all $k \geq 1$ requires non-zero $\text{disc}_{2,3}$ -error. In Section 5.4, we prove a strong version of the stable regularity lemma. This will serve as

a warm-up for the proof of Proposition 2.43, and will also play a key role in the proof of Theorem 2.49. Finally, in Section 5.5, we will prove Proposition 2.43, which says that any property which is close to a weakly stable property admits binary $\text{disc}_{2,3}$ -error.

5.1. The \sim -classes of weakly stable and weakly NIP properties. In this subsection, we prove Theorem 2.41, which says that a 3-graph property \mathcal{H} is close to a weakly stable property if and only if for some $k \geq 1$, $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$. The proof of Theorem 2.31 (the analogous result for WNIP) is extremely similar, and is thus relegated to Appendix A.

Given a 3-partite 3-graph F on n vertices, let $\mathcal{B}(F)$ be the set of 3-graphs G with $\text{Trip}(G) \cong F$ and $V(G) = [n]$. In other words, $\mathcal{B}(F)$ is the set of clean copies of F with vertex set $[n]$. Note that $\mathcal{B}(F)$ is finite.

Proof of Theorem 2.41. Suppose $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all k . Fix \mathcal{H}' a weakly stable hereditary 3-graph property. We show that \mathcal{H} is far from \mathcal{H}' .

By assumption, there is $k \geq 1$ such that $H^*(k) \notin \text{Trip}(\mathcal{H}')$. Set $\epsilon = (2k)^{-2k-1}/|\mathcal{B}(H^*(k))|$, and consider any $n \gg \epsilon^{-1}k$. By Lemma 3.31, there is $G \in \mathcal{H}$ a clean copy of $H^*(k)$. In other words, $V(G) = U \sqcup V \sqcup W$, for some $U = \{u_1, \dots, u_n\}$, $V = \{v_1, \dots, v_n\}$, $W = \{w_1, \dots, w_n\}$, and $u_i v_j w_m \in E(G)$ if and only if $j \leq m$. For each $1 \leq i \leq k$, let $V_i = \{v_{ni/k}, \dots, v_{n(i+1)/k}\}$ and $W_i = \{w_{ni/k}, \dots, w_{n(i+1)/k}\}$. Let $m = |U \cup V \cup W|$, and note $m \geq n$. We construct a large set S of induced subgraphs of G as follows.

- Choose $u \in U$. There are n choices.
- For each $i \in [k]$, choose $v'_i \in V_i \setminus \{u\}$. There are at least $(\frac{n}{k} - 1)^k \geq (n/2k)^k$ choices.
- For each $1 \leq i \leq k$, choose $w_i \in W_i \setminus \{u, v'_1, \dots, v'_k\}$. There are at least $(\frac{n}{k} - k - 1)^k \geq (\frac{n}{2k})^k$ ways to do this.
- Put $G[\{u\} \cup \{v'_i : i \in [k]\} \cup \{w_i : i \in [k]\}]$ in S .

Clearly every element of S is isomorphic to an element of $\mathcal{B}(H^*(k))$ and

$$|S| \geq n(n/2k)^{2k} > \epsilon |\mathcal{B}(H^*(k))| n^{2k+1},$$

where the inequality is by definition of ϵ and because n is large. By the pigeonhole principle, there is some $G_n \in \mathcal{B}(H^*(k))$, such that S contains at least ϵn^{2k+1} elements isomorphic to G_n . Consequently, there is some $G' \in \mathcal{B}(H^*(k))$ so that $\mathcal{B}(H^*(k)) = G_n$ for arbitrarily large n . By definition, this implies $G' \in \Delta(\mathcal{H})$. Since $H^*(k) \cong \text{Trip}(G')$ and $H^*(k) \notin \text{Trip}(\mathcal{H}')$, $G' \notin \Delta(\mathcal{H}')$ (in fact $G' \notin \mathcal{H}$). By Proposition 3.17, $\mathcal{H} \approx \mathcal{H}'$.

Suppose on the other hand that $\mathcal{H} \approx \mathcal{H}'$ for every weakly stable \mathcal{H}' . We will use the fact that, by Lemma 3.31, a property \mathcal{H}' is weakly stable if and only if there exists $k \geq 1$ so that \mathcal{H}' contains on clean copy of $H^*(k)$.

Fix $k \geq 1$. By assumption, \mathcal{H} is not close to any weakly stable property, so there is some $\delta > 0$ such that for arbitrarily large n , there is $G(n) \in \mathcal{H}_n$ and $\Gamma(n) \in \mathcal{B}(H^*(k))$, such that $G(n)$ is not $\delta/|\mathcal{B}(H^*(k))|$ -close to $\Gamma(n)$ -free. For each $\Gamma \in \mathcal{B}(H^*(k))$, let $c_k(\Gamma), N_k(\Gamma)$ be as in Theorem 3.14 for Γ and $\delta/|\mathcal{B}(H^*(k))|$, then set $c_k = \min\{c_k(\Gamma) : \Gamma \in \mathcal{B}(H^*(k))\}$ and $N_k = \max\{N_k(\Gamma) : \Gamma \in \mathcal{B}(H^*(k))\}$. Choose any $n \gg k, N_k$. By Theorem 3.14, $G(n)$ contains at least $c_k n^{2k+1}$ induced copies of some $\Gamma(n)$. Say $V(\Gamma(n)) = \{a, b_1, \dots, b_k, c_1, \dots, c_k\}$ and $ab_i c_j \in E(\Gamma(n))$ if and only if $i \leq j$. Let S be the set of induced sub-3-graphs

isomorphic to $\Gamma(n)$ in $G(n)$. For all $H \in S$, let $a^H, b_1^H, \dots, b_k^H, c_1^H, \dots, c_k^H \in V(H)$ be such that $a^H b_i^H c_j^H \in E(G(n))$ if and only if $i \leq j$. By assumption $|S| \geq c_k n^{2k+1}$. Consequently, there is $\bar{e}\bar{f} \in [n]^{2k}$ and $X \subseteq S$ with $|X| \geq \epsilon n$, such that for all $H \in X$, $(b_1^H, \dots, b_k^H, c_1^H, \dots, c_k^H) = \bar{e}\bar{f}$. Let d_1, \dots, d_k be any k distinct elements in X' . Then $d_u e_i f_j \in E$ if and only if $i \leq j$, and consequently, $\text{Trip}(G)$ contains an induced copy of $\overline{H}(k)$, so $\overline{H}(k) \in \text{Trip}(\mathcal{H})$. \square

5.2. Weak VC-dimension and vdisc_3 -homogeneity. In this section we prove Theorem 2.32, characterizing the properties which are vdisc_3 -homogeneous. We begin by showing that if H is a large 3-graph such that $\text{Trip}(H)$ omits $\overline{U}(k)$ for some $k \geq 1$, then there are restrictions on the behavior of $\text{disc}_{2,3}$ -regular triads in any decomposition of H .

Proposition 5.1. *For all $k \geq 1$ and $\epsilon \in (0, 1/2)$ there are $\epsilon_1 > 0$ and a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ such that for all $T, L \geq 1$, there is M such that the following holds.*

Suppose $H = (V, E)$ is a 3-graph on at least M vertices such that $\overline{U}(k)$ is not an induced sub-3-graph of $\text{Trip}(H)$. Suppose $t \leq T$, $\ell \leq L$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . Suppose $ijs \in \binom{[t]}{3}$ is such that at least $(1 - \sqrt{\epsilon_1})|V_i||V_j||V_s|$ many $xyz \in K_3[V_i, V_j, V_s]$ are in a $\text{disc}_{2,3}$ -regular triad of \mathcal{P} . Then $d_{ijs} \in [0, \epsilon) \cup (1 - \epsilon, 1]$ where d_{ijs} is such that $|E \cap K_3[V_i, V_j, V_s]| = d_{ijs}|V_i||V_j||V_s|$.

Proof. Fix $k \geq 1$ and $\epsilon \in (0, 1/2)$. Choose $\delta_3^{4,2}$ by applying Proposition 4.2 to k and ϵ . Then for each $\ell \geq 1$, choose $\delta_2^{4,2}(\ell)$, and M_ℓ by applying Proposition 4.2 to $k, \epsilon, \delta_3^{4,2}$, and $d_2 = 1/\ell$. For each ℓ , let ϵ_ℓ be obtained by applying Corollary 3.6 to ϵ^2/ℓ^3 and density $1/\ell$. Apply Theorem 4.1 to $t = 2k + 2^k$, $\xi = 1/2$, and $d_3 = 1/2$ to obtain $\delta_3^{4,1}$. For each $\ell \geq 1$, let $\delta_2^{4,1}(\ell)$ and m_ℓ be obtained by applying Theorem 4.1 to $t = 2k + 2^k$, $\xi = 1/2$, $d_3 = 1/2$, and $d_2 = 1/\ell$. Now let $\delta_3^{3,19}$ be obtained by applying Lemma 3.19 to $t = 2k + 2^k$, then for each $\ell \geq 1$, let $\delta_2^{3,19}(\ell)$ be obtained by applying Lemma 3.19 to $t = 2k + 2^k$, $\delta_2 = \delta_2'(\ell)$, and $\delta_3 = \delta_3^{4,1}$. Now set $\epsilon_1 = \min\{\delta_3^{3,19}, \delta_3^{4,2}, 3^{-16}, \epsilon^{-10}/10\}$, $\epsilon_2(\ell) = \min\{\delta_2^{3,19}(\ell), \delta_2^{4,2}(\ell), \epsilon_\ell\}$, and let $M = TM_T m_T$.

Suppose H is a 3-graph on at least M vertices such that $\overline{U}(k)$ is not an induced sub-3-graph of $\text{Trip}(H)$. By Fact 2.22, $\text{VC}_2(H) \leq k$. Suppose $t \leq T$, $\ell \leq L$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . For each triad $G_{ijs}^{\alpha, \beta, \tau}$ of \mathcal{P} , let $d_{ijs}^{\alpha, \beta, \tau}$ be such that $|E \cap K_3^{(2)}(G_{ijs}^{\alpha, \beta, \tau})| = d_{ijs}^{\alpha, \beta, \tau} |K_3^{(2)}(G_{ijs}^{\alpha, \beta, \tau})|$. For each $ijs \in \binom{[t]}{3}$, let I_{ijs} be the set of $(\alpha, \beta, \tau) \in [\ell]^3$ such that $(H_{ijk}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \tau})$ satisfies $\text{disc}_{2,3}(\epsilon_2(\ell), \epsilon_1)$. Let

$$\Sigma = \left\{ ijs \in \binom{[t]}{3} : \left| \bigcup_{(\alpha, \beta, \gamma) \in I_{ijs}} K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma}) \right| \geq (1 - \sqrt{\epsilon_1}) |K_3^{(2)}(G_{ijs})| \right\}.$$

Note that by our choice of parameters and by Proposition 4.2, for every $(\alpha, \beta, \gamma) \in I_{ijs}$, there is $\delta_{ijs}^{\alpha, \beta, \tau} \in \{0, 1\}$ such that $|\delta_{ijs}^{\alpha, \beta, \tau} - d_{ijs}^{\alpha, \beta, \tau}| \leq \epsilon$.

Claim 5.2. *For all $ijs \in \Sigma$, if $\tau \neq \tau'$ and $(\alpha, \beta, \tau), (\alpha, \beta, \tau') \in I_{ijs}$, then $\delta_{ijs}^{\alpha, \beta, \tau} = \delta_{ijs}^{\alpha, \beta, \tau'}$.*

PROOF OF CLAIM 5.2: Suppose towards a contradiction there are $(\alpha, \beta, \tau), (\alpha, \beta, \tau') \in I_{ijs}$ with $\tau \neq \tau'$ such that $\delta_{ijs}^{\alpha, \beta, \tau} = 0$ while $\delta_{ijs}^{\alpha, \beta, \tau'} = 1$. Let $m = n/t2^k$, and choose equipartitions $V_i = V_{i,1} \cup \dots \cup V_{i,2^k}$, $V_j = V_{j,1} \cup \dots \cup V_{j,2^k}$, and $V_s = V_{s,1} \cup \dots \cup V_{s,2^k}$. Let $\{S_1, \dots, S_{2^k}\}$ enumerate the elements of $\mathcal{P}([k])$. We define a $(2k + 2^k)$ -partite graph G with vertex set $V' = \bigcup_{u=1}^{2^k} V_{i,u} \cup \bigcup_{u=1}^k V_{j,u} \cup \bigcup_{u=1}^k V_{s,u}$ as follows. For each $1 \leq a, b \leq k$, and $1 \leq u \leq 2^k$, set $G[V_{i,u}, V_{j,a}] = P_{ij}^\alpha$, $G[V_{i,u}, V_{s,b}] = P_{is}^\beta$, and

$$G[V_{j,a}, V_{s,b}] = \begin{cases} P_{js}^{\tau'} & \text{if } a \in S_u \\ P_{js}^\tau & \text{if } a \notin S_u. \end{cases}$$

Now define H' to be the 3-partite 3-graph with vertex set V' and whose edge set is defined as follows.

$$E(H) := \bigcup_{1 \leq a, b \leq k, 1 \leq u \leq 2^k} E \cap K_3^{(2)}(G[V_{i,u}, V_{j,a}, V_{s,b}]).$$

By Lemma 3.19, and our choice of parameters, we have that for all $1 \leq a, b \leq k$ and $1 \leq u \leq 2^k$, $(H', G[V_{i,u}, V_{j,a}, V_{s,b}])$ has $\text{disc}_3(\delta_2^{4,1}(\ell), \delta_3^{4,1})$ and by assumption, the density of edges in H' on $K_3(G[V_{i,u}, V_{j,a}, V_{s,b}])$ is at least ϵ if $a \in S_u$ and at most $1 - \epsilon$ if $a \notin S_u$. But now Theorem 4.1 implies $\text{Trip}(H)$ contains an induced copy of $\overline{U}(k)$, a contradiction. \square

Note that the same argument shows that if $ijs \in \Sigma$ and $(\alpha, \beta', \tau), (\alpha, \beta', \tau) \in I_{ijs}$ for some $\beta \neq \beta'$ (respectively $(\alpha', \beta, \tau), (\alpha', \beta, \tau) \in I_{ijs}$ for some $\alpha \neq \alpha'$), then $\delta_{ijs}^{\alpha, \beta, \tau} = \delta_{ijs}^{\alpha, \beta', \tau}$ (respectively $\delta_{ijs}^{\alpha, \beta, \tau} = \delta_{ijs}^{\alpha', \beta, \tau}$).

Now fix $ijs \in \Sigma$. Let J_{ijs} be the set of $\alpha \in [\ell]$ such that there are at least $(1 - \epsilon_1^{1/4})\ell^2$ many pairs $(\beta, \tau) \in [\ell]^2$, with $(\alpha, \beta, \tau) \in I_{ijs}$. By an easy counting argument, $|J_{ijs}| \geq (1 - \epsilon_1^{1/4})\ell$. Given $\alpha \in J_{ijs}$, let J_{ijs}^α be the set of $\beta \in [\ell]$ such that there are at least $(1 - \epsilon_1^{1/16})\ell$ many $\tau \in [\ell]$ such that $(\alpha, \beta, \tau) \in I_{ijs}$. By an easy counting, $|J_{ijs}^\alpha| \geq (1 - \epsilon_1^{1/16})\ell$. Given $\beta \in J_{ijs}^\alpha$, let $J_{ijs}^{\alpha, \beta}$ be the set of $\tau \in [\ell]$ such that $(\alpha, \beta, \tau) \in I_{ijs}$. By definition, $|J_{ijs}^{\alpha, \beta}| \geq (1 - \epsilon_1^{1/16})\ell$.

We show that there is a $\delta_{ijs} \in \{0, 1\}$ so that for all (α, β, γ) with $\alpha \in J_{ijs}$, $\beta \in J_{ijs}^\alpha$, and $\gamma \in J_{ijs}^{\alpha, \beta}$, $\delta_{ijs}^{\alpha, \beta, \gamma} = \delta_{ijs}$. Suppose $\alpha \in J_{ijs}$. By Claim 5.2, for all $\beta \in J_{ijs}^\alpha$, there is $\delta_{ijs}^{\alpha, \beta} \in \{0, 1\}$ such that for all $\tau \in J_{ijs}^{\alpha, \beta}$, $\delta_{ijs}^{\alpha, \beta, \tau} = \delta_{ijs}^{\alpha, \beta}$. Suppose $\beta \neq \beta' \in J_{ijs}^\alpha$. By definition of J_{ijs}^α , $|J_{ijs}^{\alpha, \beta} \cap J_{ijs}^{\alpha, \beta'}| \geq (1 - 2\epsilon_1^{1/16})\ell$. Thus there is some $\tau \in J_{ijs}^{\alpha, \beta} \cap J_{ijs}^{\alpha, \beta'}$. Claim 5.2 applied to (α, β, τ) and (α, β', τ) implies that $\delta_{ijs}^{\alpha, \beta} = \delta_{ijs}^{\alpha, \beta'}$. Thus there is δ_{ijs}^α , such that for all $\beta \in J_{ijs}^\alpha$, $\delta_{ijs}^{\alpha, \beta} = \delta_{ijs}^\alpha$. Now suppose $\alpha \neq \alpha' \in J_{ijs}$. By definition of J_{ijs} , $|J_{ijs}^\alpha \cap J_{ijs}^{\alpha'}| \geq (1 - 2\epsilon_1^{1/4})\ell$. Thus there is some $\beta \in J_{ijs}^\alpha \cap J_{ijs}^{\alpha'}$. By definition, $|J_{ijs}^{\alpha, \beta} \cap J_{ijs}^{\alpha', \beta}| \geq (1 - 2\epsilon_1^{1/16})\ell$, so there is some $\tau \in J_{ijs}^{\alpha, \beta} \cap J_{ijs}^{\alpha', \beta}$. Now Claim 5.2 applied to (α, β, τ) and (α', β, τ) implies that $\delta_{ijs}^{\alpha, \beta} = \delta_{ijs}^{\alpha', \beta}$, and thus $\delta_{ijs}^\alpha = \delta_{ijs}^{\alpha'}$. We have shown there is δ_{ijs} such that for all $\alpha \in J_{ijs}$, $\delta_{ijs}^\alpha = \delta_{ijs}$. In other words, for all $\alpha \in J_{ijs}$, $\beta \in J_{ijs}^\alpha$, and $\gamma \in J_{ijs}^{\alpha, \beta}$, $\delta_{ijs}^{\alpha, \beta, \gamma} = \delta_{ijs}$.

We now show that the triple (V_i, V_j, V_s) has density near δ_{ijs} in H . For simplicity, let us assume $\delta_{ijs} = 1$ (the case $\delta_{ijs} = 0$ is similar). First, given $\alpha \in J_{ijs}$, the following holds,

where $G_{ijs}^\alpha = (V_i \cup V_j \cup V_s, P_{is}^\alpha \cup K_2[V_i, V_j] \cup K_2[V_j, V_s])$.

$$\begin{aligned} |E \cap K_3^{(2)}(G_{ijs}^\alpha)| &\geq \sum_{\beta \in J_{ijs}^\alpha} \sum_{\tau \in J_{ijs}^{\alpha, \beta}} (1 - \epsilon) |K_3^{(2)}(G_{ijs}^{\alpha, \beta, \tau})| \\ &\geq (1 - \epsilon^{1/16})(1 - \epsilon^{1/4}) \ell^2 \frac{1}{\ell^3} (1 - \ell^2 \epsilon^2 / \ell^2) |V_i| |V_j| |V_s| \\ &\geq (1 - \epsilon^{1/16})(1 - \epsilon^{1/4})(1 - \epsilon^2) \frac{1}{\ell} |V_i| |V_j| |V_s|, \end{aligned}$$

where the second inequality uses Corollary 3.6 and the size bounds on $J_{ijs}^\alpha, J_{ijs}^{\alpha, \beta}$. Thus

$$\begin{aligned} |E \cap K_3[V_i, V_j, V_s]| &\geq |J_{ijs}| (1 - \epsilon^{1/16})(1 - \epsilon^{1/4})(1 - \epsilon^2) \frac{1}{\ell} |V_i| |V_j| |V_s| \\ &\geq (1 - \epsilon^{1/16})(1 - \epsilon^{1/4})^2 (1 - \epsilon^2) |V_i| |V_j| |V_s| \\ &\geq (1 - \epsilon) |V_i| |V_j| |V_s|, \end{aligned}$$

where the last inequality is by choice of ϵ . This shows that for every $ijs \in \Sigma$, the triple (V_i, V_j, V_s) is ϵ -homogeneous with respect to H , as desired. \square

We are now in a position to prove Theorem 2.28 which states that a hereditary 3-graph property \mathcal{H} is vdisc_3 -homogeneous if and only if there is $k \geq 1$ such that $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$.

Proof of Theorem 2.28. Suppose first that there is $k \geq 1$ such that $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$. Fix $0 < \epsilon < 1/100$ and $t_0 \geq 1$. Let δ'_3 and δ''_3 be obtained by applying Proposition 4.2 and Proposition 5.1 respectively to ϵ and k , and set $\delta_3 = \min\{\delta'_3, \delta''_3\}$. Given ℓ , let $\delta'_2(\ell)$, m'_ℓ be obtained by applying Proposition 4.2 to ϵ , k , δ'_3 and $d_2 = 1/\ell$. Given ℓ , let $\delta''_2(\ell)$ and m''_ℓ be obtained by applying Proposition 5.1 to ϵ , k , and δ''_3 with $d_2 = 1/\ell$.

Set $\delta_2(\ell) = \min\{\delta'_2(\ell), \delta''_2(\ell)\}$, and $\epsilon_1 = \min\{\delta_3, \epsilon^2\}$, and define $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ so that $\epsilon_2(\ell) = \min\{\delta_2(\ell), \epsilon/(3\ell)^3\}$. Let $T = T(\epsilon_1, \epsilon_2, t_0, 1)$, $L = L(\epsilon_1, \epsilon_2, t_0, 1)$ and $N_1 = N_1(\epsilon_1, \epsilon_2, t_0, 1)$ be as in Theorem 2.8. Let N_2 be sufficiently large so that $\binom{N_2}{3} \geq N_2^3/10$. Now let $M = \max\{N_1, N_2, m'_T T, m''_T T\}$.

Now suppose $H = (V, E) \in \mathcal{H}$ with $|V| = n \geq M$. By Theorem 2.8, there is $t_0 \leq t \leq T$ and $1 \leq \ell \leq L$ and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . Let Γ be the set of $xyz \in \binom{V}{3}$, such that xyz is not in a $\text{disc}_{2,3}$ -regular triad of \mathcal{P} . Let Σ be the set of $ijs \in \binom{[t]}{3}$ such that $|\Gamma \cap K_3[V_i, V_j, V_s]| \geq \sqrt{\epsilon_1} |V_i| |V_j| |V_s|$. Clearly $|\Sigma| \leq \sqrt{\epsilon_1} t^3$ because \mathcal{P} is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H . By Proposition 5.1, for all $ijs \in \binom{[t]}{3} \setminus \Sigma$, the triple (V_i, V_j, V_k) is ϵ -homogeneous with respect to H . Since $\sqrt{\epsilon_1} \leq \epsilon$, this shows that $\mathcal{P}_{\text{vert}}$ is an ϵ -homogeneous partition with respect to \mathcal{H} . We have now shown that \mathcal{H} is vdisc_3 -homogeneous.

For the converse, suppose $\overline{U}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Fix $\epsilon < 1/8$ and $t_0 \geq 1$, and set $M_0 = \max\{\epsilon^{-1}, t_0\}$. By Fact 3.4, for all $T \geq t_0$ there is n_0 such that for all $n \geq n_0$, there is a bipartite graph $G = (U \cup V, E)$ with $|U| = |V| = n$, and which satisfies $\text{disc}_2(\epsilon^2/T^2, 1/2)$.

Suppose $T \geq t_0$ and $n \geq n_0 M_0$. Fix a bipartite graph $G = (U \cup V, E)$ with $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_n\}$ which satisfies $\text{disc}_2(\epsilon^2/T, 1/2)$. By Lemma 3.33, \mathcal{H}

contains a clean copy of $n \otimes G$. Thus there exists some $H = (V, R) \in \mathcal{H}$, and sets $B = \{b_1, \dots, b_n\}$, $A = \{a_1, \dots, a_n\}$, $C = \{c_1, \dots, c_n\}$ so that $V = A \sqcup B \sqcup C$, and such that $a_s b_i c_j \in R$ if and only if $u_i w_j \in E$. We show there is no equipartition of V into at least t_0 and at most T parts which is ϵ -homogeneous with respect to H .

Suppose towards a contradiction there are $t_0 \leq t \leq T$, an equipartition $\{V_1, \dots, V_t\}$ of V , and $\Sigma \subseteq \binom{[t]}{3}$, such that $|\Sigma| \leq \epsilon t^3$ and such that for all $ijk \in \binom{[t]}{3} \setminus \Sigma$, $|R \cap K_3[V_i, V_j, V_k]|/|K_3[V_i, V_j, V_k]| \in [0, \epsilon) \cup (1 - \epsilon, 1]$. Given $ijs \in \binom{[t]}{3}$, set

$$d_{ijs} = \frac{|E \cap K_3(V_i, V_j, V_s)|}{|V_i||V_j||V_s|}.$$

For each $i \in [t]$, let $A_i = A \cap V_i$, $B_i = B \cap V_i$, and $C_i = C \cap V_i$. Let Σ' be the set of $i_1 i_2 i_3 \in \binom{[t]}{3}$ such that for some $\sigma \in S_3$, $\min\{|A_{i_{\sigma(1)}}|, |B_{i_{\sigma(2)}}|, |C_{i_{\sigma(3)}}|\} \geq \epsilon^{1/4} n/t$.

We show that for all $i_1 i_2 i_3 \in \Sigma'$, $d_{i_1 i_2 i_3} \in (\epsilon, 1 - \epsilon)$. Fix $i_1 i_2 i_3 \in \Sigma'$ and let $\sigma \in S_3$ be such that $\min\{|A_{i_{\sigma(1)}}|, |B_{i_{\sigma(2)}}|, |C_{i_{\sigma(3)}}|\} \geq \epsilon^{1/4} n/t$. Let $U' = \{u_j : b_j \in B_{i_{\sigma(2)}}\}$ and $W' = \{w_j : c_j \in C_{i_{\sigma(3)}}\}$. Then since G has $\text{disc}_2(\epsilon^2/T^2, 1/2)$, the following holds.

$$(3) \quad ||E \cap K_2(U', V')| - |U'||V'|/2| \leq \frac{\epsilon^2}{T^2} n^2.$$

Note that because $i_{\sigma(2)} \neq i_{\sigma(3)}$, $B_{i_{\sigma(2)}}$, $C_{i_{\sigma(3)}}$, and $A_{i_{\sigma(1)}}$ are disjoint, thus

$$\begin{aligned} |R \cap K_3[A_{i_{\sigma(1)}}, B_{i_{\sigma(2)}}, C_{i_{\sigma(3)}}]| &= |A_{i_{\sigma(1)}}| |E \cap K_2[U', V']| \text{ and} \\ |K_3[A_{i_{\sigma(1)}}, B_{i_{\sigma(2)}}, C_{i_{\sigma(3)}}] \setminus R| &= |A_{i_{\sigma(1)}}| |K_2[U', V'] \setminus E| \end{aligned}$$

Combining this with (3), we have the following.

$$\frac{|A_{i_{\sigma(1)}}|}{(n/t)^3} \left(\frac{|U'||V'|}{2} - \frac{\epsilon^2 n^2}{T^2} \right) \leq d_{i_1 i_2 i_3} \leq \frac{|A_{i_{\sigma(1)}}|}{(n/t)^3} \left(\frac{|U'||V'|}{2} + \frac{\epsilon^2 n^2}{T^2} \right).$$

By assumption, $\min\{|A_{i_{\sigma(1)}}|, |U'|, |W'|\} \geq \epsilon^{1/4} n/t$ and $n/t \geq \max\{|A_{i_{\sigma(1)}}|, |U'|, |W'|\}$. Consequently,

$$\epsilon^{3/4}/2 - \epsilon^{5/4} t^2/T^2 \leq d_{ijs} \leq \frac{1}{2} + \frac{\epsilon^2 t^2}{T^2}.$$

Since $t \leq T$ and $\epsilon < 1/8$, this implies $d_{ijs} \in (\epsilon, 1 - \epsilon)$.

We have now shown that for all $ijs \in \Sigma'$, $d_{ijs} \in (\epsilon, 1 - \epsilon)$, while by assumption, for all $ijs \in \Sigma$, $d_{ijs} \in [0, \epsilon) \cup (1 - \epsilon, 1]$. Consequently, $\Sigma' \subseteq \Sigma$, and therefore, $|\Sigma'| \leq \epsilon t^3$.

Given $i \in [t]$, we say A_i , B_i , or C_i is *trivial* if it has size at most $\epsilon^{1/4} |V_i|$. Note that for each $i \in [t]$, at most two of A_i , B_i , or C_i can be trivial. Consequently, if V_0 is the union of all trivial A_i, B_i, C_i , then $|V_0| \leq 2\epsilon^{1/4} |V|$. Note that if $ijs \in \binom{[t]}{3} \setminus \Sigma'$, then $K_3[V_i, V_j, V_s] \subseteq K_3[V_0, V, V]$, and consequently $|\bigcup_{ijs \in \binom{[t]}{3} \setminus \Sigma'} K_3[V_i, V_j, V_s]| \leq 2\sqrt{\epsilon} |V|^3$. But

now we have that

$$\begin{aligned} \binom{|V|}{3} &\leq \left| \bigcup_{ijs \in \binom{[t]}{3} \setminus \Sigma'} K_3[V_i, V_j, V_s] \right| + \left| \bigcup_{ijs \in \Sigma'} K_3[V_i, V_j, V_s] \right| \leq 2\sqrt{\epsilon}|V|^3 + \epsilon t^3(|V|/t)^3 \\ &\leq 3\sqrt{\epsilon}|V|^3. \end{aligned}$$

Clearly this is a contradiction, our assumptions on ϵ and $|V|$ imply $\binom{|V|}{3} > 3\sqrt{\epsilon}|V|^3$. This finishes our proof that there is no equipartition of V into $t_0 \leq t \leq T$ parts which is ϵ -homogeneous with respect to H . This shows that \mathcal{H} is not vdisc_3 -homogeneous. \square

Theorem 2.32 follows immediately from Theorem 2.28 and Theorem 2.31. We end this subsection by proving Proposition 2.44, which gives an equivalent characterization of properties admitting zero/binary vdisc_3 -error. For this, it will be useful to have a uniform way of talking about decompositions with zero and binary vdisc_3 -error.

Definition 5.3. Given $G = (V, E)$ a 3-graph, an ϵ -homogeneous partition of G with zero (respectively, binary) error is an equipartition $V = V_1 \cup \dots \cup V_t$ along with a set $\Sigma \subseteq \binom{[t]}{2}$ such that the following hold.

- (1) $|\Sigma| = 0$ (respectively, $|\Sigma| \leq \epsilon t^2$).
- (2) For all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Sigma$, $\frac{|E \cap K_3[V_i, V_j, V_k]|}{|K_3[V_i, V_j, V_k]|} \in [0, \epsilon) \cup (1 - \epsilon, 1]$.

Observe that a hereditary 3-graph property \mathcal{H} is vdisc_3 -homogeneous with zero (respectively, binary) error if and only if for all $\epsilon > 0$ and $t_0 \geq 1$ there are $T, N \geq 1$ such that for every $G = (V, E) \in \mathcal{H}$ with $|V| \geq N$, there is an equipartition of V into at most T parts which is a ϵ -homogeneous partition of G with zero (respectively, binary error).

We now prove Proposition 2.44, which shows that a hereditary 3-graph property \mathcal{H} admits zero (respectively binary) vdisc_3 -error if and only if \mathcal{H} is close to a WNIP property and admits zero (respectively binary) $\text{disc}_{2,3}$ -error.

PROOF OF PROPOSITION 2.44: Suppose first that \mathcal{H} is a hereditary 3-graph property which is WNIP and admits zero (respectively, binary) $\text{disc}_{2,3}$ -error. Say $\text{WVC}(\mathcal{H}) < k$. By Theorem 2.32, \mathcal{H} is vdisc_3 -homogeneous. Fix $\epsilon > 0$ and $t_0 \geq 1$. Without loss of generality, assume $\epsilon < 1/2$. Choose $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ as in Proposition 5.1 for k and $\epsilon/2$. Let M_1, T, L be as in the definition of \mathcal{H} is admitting zero (respectively binary) $\text{disc}_{2,3}$ -error for $\epsilon_1, \epsilon_2, t_0$ and $\ell_0 = 1$. Let $M_2 = M_2(k, \epsilon, T)$ in Proposition 5.1.

Now suppose $G = (V, E) \in \mathcal{H}$ satisfies $|V| \geq \max\{M_1, M_2\}$. By assumption, there are $t_0 \leq t \leq T$, $1 \leq \ell \leq L$, and \mathcal{P} a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to H , and $\Sigma \subseteq \binom{[t]}{2}$ of size 0 (respectively, of size at most $\epsilon_1 t^2$), such that for every $P_{ijs}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ which is not $\text{disc}_{2,3}$ -regular with respect to H , one of $ij, is, jk \in \Sigma$. By Proposition 5.1, for every $ijs \in \binom{[t]}{3}$ with $ij, js, is \notin \Sigma$, $|E \cap K_3[V_i, V_j, V_s]|/|K_3[V_i, V_j, V_s]| \in [0, \epsilon) \cup (1 - \epsilon, 1]$. Thus $V_1 \cup \dots \cup V_t$ is an ϵ -homogeneous partition of G with zero (respectively, binary) error. This shows that if \mathcal{H} is WNIP and admits zero (respectively, binary) $\text{disc}_{2,3}$ -error, then \mathcal{H} admits zero (respectively, binary)

vdisc₃-error. It follows immediately from Proposition 2.19 that the same holds for \mathcal{H}' close to a WNIP property.

Suppose conversely that \mathcal{H} admits zero (respectively, binary) vdisc₃-error. By Theorem 2.32, \mathcal{H} is close to a WNIP property. Fix $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$. Given $x \in \mathbb{N}$, let $\delta_2(x)$ be the δ_2 from Proposition 3.22 for $\epsilon_1/2\ell_0$ and $d_2 = 1/x$. Let ϵ_2'' be as in Corollary 3.6 for density $1/\ell_0$ and $\min\{\epsilon_2(\ell_0)/\ell_0^3, \delta_2(\ell_0)\}$. Let $\epsilon_2' = \min\{\epsilon_2'', \epsilon_2(\ell_0)/\ell_0, \delta_2(\ell_0)\}$. Let T and N_1 be as in the definition of \mathcal{H} admitting zero (respectively, binary) vdisc₃-error for $\epsilon = \epsilon_1^2/2\ell_0^3$ and t_0 . Let N_2 be the $m(\epsilon_2', \epsilon_2', \epsilon_2', \ell_0)$ from Lemma 3.24, and set $N = \max\{N_1, N_2T\}$.

Suppose $H = (V, E) \in \mathcal{H}$ and $|V| \geq N$. By assumption, there exists $t_0 \leq t \leq T$ and $\mathcal{P} = \{V_1, \dots, V_t\}$ an $(\epsilon_1^2/2\ell_0^3)$ -homogeneous partition for H with zero (respectively, binary) error. For each $ij \in \binom{[t]}{2}$, by Lemma 3.24, there exists a partition $K_2[V_i, V_j] = \bigcup_{\alpha \leq \ell_0} P_{ij}^\alpha$ such that each P_{ij}^α has $\text{disc}_2(\epsilon_2'(\ell_0), 1/\ell_0)$. Suppose $ijk \in \binom{[t]}{3}$ has the property that none of ij, ik, jk are Σ . Then by assumption, there is $u \in \{0, 1\}$ such that $|E^u \cap K_3[V_i, V_j, V_k]| \geq (1 - \epsilon_1^2/2\ell_0^3)|V_i||V_j||V_k|$. By Corollary 3.6, for any $\alpha, \beta, \gamma \leq \ell_0$,

$$|K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})| = \frac{1}{\ell_0^3}(1 \pm \epsilon_2(\ell_0)/\ell_0^3)|V_i||V_j||V_k|,$$

so

$$|E^u \cap K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})| \geq |K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})| - (\epsilon_1^2/2\ell_0^3)|V_i||V_j||V_k| \geq (1 - \epsilon_1^2)|K_3^{(2)}(G_{ijk}^{\alpha, \beta, \gamma})|.$$

By Proposition 3.22, $(H_{ijk}^{\alpha, \beta, \gamma}, G_{ijk}^{\alpha, \beta, \gamma})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell_0))$, as desired. This shows that if \mathcal{H} admits zero (respectively, binary) vdisc₃-error, then \mathcal{H} is close to a WNIP property and admits zero (respectively, binary) vdisc_{2,3}-error. \square

5.3. Weak stability and non-zero error. In this subsection, we prove results concerning weak stability and the notions of zero $\text{disc}_{2,3}$ -error and zero vdisc₃-error. We begin by showing that a property may be unstable but still admit zero $\text{disc}_{2,3}$ -error.

Example 5.4. Let H be the 3-uniform hypergraph with vertex set $\{a_i, b_i, c_i : i \in \mathbb{N}\}$ and edge set $\{a_i b_j c_k : j \leq k\}$. Let $\mathcal{H} = \text{age}(H)$. Then $R(x; y, z)$ is unstable in \mathcal{H} , but \mathcal{H} admits zero $\text{disc}_{2,3}$ -error.

Proof. Fix $\ell_0, t_0 \geq 1$, $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Let $\epsilon_2' : \mathbb{N} \rightarrow (0, 1]$ be such that for all $\ell \geq 1$, $\epsilon_2'(\ell) \leq \min\{\epsilon_2(\ell), \epsilon_1^4/\ell^4, \epsilon_\ell\}$, where ϵ_ℓ comes from applying Corollary 3.6 to $\gamma = \epsilon_1^4/10\ell^3$ and $t = 3$. Let N_1, T, L be as in Proposition 2.13 for $\epsilon_1^4, \epsilon_2'$ and t_0, ℓ_0 . Let N_2 be sufficiently large so that $N_2^2 < \epsilon_1^4 N_2^3 / 2T^3 L^3$. Let $N = \max\{N_1, N_2\}$.

Suppose $H = (V, E) \in \mathcal{H}$, where $|V| = n \geq N$. Then by definition of \mathcal{H} , we can write $V(H) = V_A \cup V_B \cup V_C$ where $V_A = \{a_{i_1}, \dots, a_{i_s}\}$, $V_B = \{b_{j_1}, \dots, b_{j_t}\}$ and $V_C = \{c_{r_1}, \dots, c_{r_m}\}$ for some $s + t + m = n$, and $E(H) = \{(a_{i_u}, b_{j_v}, c_{r_k}) : i_u \leq j_v = r_k\}$. Note $|E(H)| \leq s \max\{t, m\} \leq n^2/4$, where the second inequality is by the AM-GM inequality. Since $n \geq N_1$, there exists $t_0 \leq t \leq T$ and $\ell_0 \leq \ell \leq L$, and a $(t, \ell, \epsilon_1^4, \epsilon_2'(\ell))$ -decomposition \mathcal{P} of V which is $(\epsilon_1^4, \epsilon_2'(\ell))$ -regular with respect to H , and which has no disc_2 -irregular

triads. Fix a triad $G_{ijk}^{\alpha,\beta,\gamma}$ of \mathcal{P} . By assumption, each of $P_{ij}^\alpha, P_{ik}^\beta, P_{jk}^\gamma$ has $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$. Combining Corollary 3.6 with the bound on $E(H)$ implies the following.

$$\frac{|E \cap K_3^{(2)}(G_{ijk}^{\alpha,\beta,\gamma})|}{|K_3^{(2)}(G_{ijk}^{\alpha,\beta,\gamma})|} \leq \frac{n^2}{4|K_3^{(2)}(G_{ijk}^{\alpha,\beta,\gamma})|} \leq \frac{\epsilon_1^4 |V_i| |V_j| |V_k|}{T^3 L^3} \leq \epsilon_1^2,$$

where the second inequality is by our choice of N and Corollary 3.6. By Proposition 3.22, $(H_{ijk}^{\alpha,\beta,\gamma}, G_{ijk}^{\alpha,\beta,\gamma})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$. This shows \mathcal{H} admits zero $\text{disc}_{2,3}$ -error. \square

Our next goal is to show that if a hereditary 3-graph property \mathcal{H} has $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all k , then \mathcal{H} requires non-zero $\text{disc}_{2,3}$ -error. The main reason for this is the following theorem, which shows that any equipartition of the vertex set of $\overline{H}(k)$ has a triple with intermediate density.

We would like to acknowledge here a (no longer available) blog post written by Aleksandar Makelov, which presented a proof that $H(k)$ requires irregular pairs in the graph regularity lemma. A proof of this fact does not appear in the published literature (to the authors' knowledge), and we thus found this blog post a very useful starting point as we developed the following proof, as well the other proofs in this paper showing that certain examples require certain kinds of irregular triples/triads.

Theorem 5.5. *For all $0 < \epsilon_1 \leq 2^{-18}$, and all $T, N \geq 1$, there is $n \geq N$ such that for every $t \leq T$ and equipartition $\mathcal{P} = \{V_1, \dots, V_t\}$ of $V(\overline{H}(n))$, the following holds. There is a triple $ijk \in \binom{[t]}{3}$, and sets $A_i \subseteq V_i$, $B_{j1}, B_{j0} \subseteq V_j$, and $C_{k1}, C_{k0} \subseteq V_k$, each of size at least $\epsilon_1^{1/9}(3n/t)$, such that $K_3[A_i, B_{j1}, C_{k1}] \subseteq E$ and $K_3[A_i, B_{j0}, C_{k0}] \cap E = \emptyset$.*

Proof. Fix $0 < \epsilon_1 < 2^{-18}$ and $T, N \geq 1$. Let $n_0 = \epsilon_1^{-100} T^2$ and assume $n \gg n_0$ and $t \leq T$. Suppose $\mathcal{P} = \{V_1, \dots, V_t\}$ is an equipartition of $V(\overline{H}(n))$. Say $V = V_1 \cup \dots \cup V_t$ is the partition from \mathcal{P} . Fix a labeling $V(\overline{H}(k)) = A \cup B \cup C$ where $A = \{a_i : i \in [n]\}$, $B = \{b_i : i \in [n]\}$, $C = \{c_i : i \in [n]\}$ are such that $b_i c_j \in E(\overline{H}(k))$ if and only if $i \leq j$. For each $1 \leq i \leq t$, define $A_i = A \cap V_i$, $B_i = B \cap V_i$, and $C_i = C \cap V_i$, and set

$$\mathcal{A} = \{i \in [t] : |A_i| \geq \epsilon_1^{1/9} n/t\}, \quad \mathcal{B} = \{i \in [t] : |B_i| \geq \epsilon_1^{1/9} n/t\}, \quad \text{and} \quad \mathcal{C} = \{i \in [t] : |C_i| \geq \epsilon_1^{1/9} n/t\}.$$

Set $A' = \bigcup_{i \in \mathcal{A}} A_i$, $B' = \bigcup_{i \in \mathcal{B}} B_i$, and $C' = \bigcup_{i \in \mathcal{C}} C_i$. Observe that

$$|A \setminus A'| \leq \sum_{i \notin \mathcal{A}} \epsilon_1^{1/9} n/t \leq \epsilon_1^{1/9} n.$$

Similarly, $|B \setminus B'| \leq \epsilon_1^{1/9} n$ and $|C \setminus C'| \leq \epsilon_1^{1/9} n$. Given $i \in [t]$, and $\mu > 0$, let

$$B_i(\mu) = \{j \in [n] : b_j \in B_i \text{ and } \min\{|\{b_1, \dots, b_j\} \cap B_i|, |\{b_{j+1}, \dots, b_n\} \cap B_i|\} \geq \mu |B_i|\}, \text{ and}$$

$$C_i(\mu) = \{j \in [n] : c_j \in C_i \text{ and } \min\{|\{c_1, \dots, c_j\} \cap C_i|, |\{c_{j+1}, \dots, c_n\} \cap C_i|\} \geq \mu |C_i|\},$$

Observe that for any $\mu > 0$ and $i \in [t]$, set

$$(B_i \setminus B_i(\mu)) \subseteq (B_i \cap \{b_1, \dots, b_{\min(B_i(\mu))-1}\}) \cup (B_i \cap \{b_{\max(B_i(\mu))+1}, \dots, b_n\}),$$

and consequently, $|B_i(\mu)| \geq (1 - 2\mu)|B_i|$. A similar argument shows $|C_i(\mu)| \geq (1 - 2\mu)|C_i|$.

Suppose first there exist $i \neq j$ such that $i \in \mathcal{B}$, $j \in \mathcal{C}$, and $B_i(\epsilon_1^{1/9}) \cap C_j(\epsilon_1^{1/8}) \neq \emptyset$. Say $w \in B_i(\epsilon_1^{1/9}) \cap C_j(\epsilon_1^{1/9})$. Set $C_{j0} = C_j \cap \{c_{w+1}, \dots, c_n\}$, $C_{j1} = C_j \cap \{c_1, \dots, c_w\}$, $B_{i0} = B_i \cap \{b_{w+1}, \dots, b_n\}$, and $B_{i1} = B_i \cap \{b_1, \dots, b_w\}$. Fix any $k \in \mathcal{A} \cap ([t] \setminus \{i, j\})$. Then $K_3[A_k, C_{j1}, B_{i1}] \subseteq E$ while $K_3[A_k, C_{j0}, B_{i0}] \cap E = \emptyset$, so we are done.

This leaves us with the case where, for all $i \in \mathcal{B}$ and $j \in \mathcal{C}$, if i and j are distinct, then $B_i(\epsilon_1^{1/9}) \cap C_j(\epsilon_1^{1/9}) = \emptyset$. In this case we have the following.

$$\begin{aligned}
n &\geq \left| \bigcup_{i \in \mathcal{B}} B_i(\epsilon_1^{1/9}) \cup \bigcup_{j \in \mathcal{C}} C_j(\epsilon_1^{1/9}) \right| \geq \sum_{i \in \mathcal{B}} |B_i(\epsilon_1^{1/9})| + \sum_{j \in \mathcal{C}} |C_j(\epsilon_1^{1/9})| - \left| \sum_{s \in \mathcal{B} \cap \mathcal{C}} B_s(\epsilon_1^{1/9}) \cap C_s(\epsilon_1^{1/9}) \right| \\
&\geq (1 - 2\epsilon_1^{1/9}) \left(\sum_{i \in \mathcal{B}} |B_i| + \sum_{j \in \mathcal{C}} |C_j| \right) - \left| \sum_{s \in \mathcal{B} \cap \mathcal{C}} B_s(\epsilon_1^{1/9}) \cap C_s(\epsilon_1^{1/9}) \right| \\
&= (1 - 2\epsilon_1^{1/9}) (|B'| + |C'|) - \left| \sum_{s \in \mathcal{B} \cap \mathcal{C}} B_s(\epsilon_1^{1/9}) \cap C_s(\epsilon_1^{1/9}) \right| \\
&\geq (1 - 2\epsilon_1^{1/9}) 2n(1 - \epsilon_1^{1/9}) - \left| \sum_{s \in \mathcal{B} \cap \mathcal{C}} B_s(\epsilon_1^{1/9}) \cap C_s(\epsilon_1^{1/9}) \right|.
\end{aligned}$$

This implies $\left| \sum_{s \in \mathcal{B} \cap \mathcal{C}} B_s(\epsilon_1^{1/9}) \cap C_s(\epsilon_1^{1/9}) \right| \geq (1 - 2\epsilon_1^{1/9}) 2n(1 - \epsilon_1^{1/9}) - n \geq n(1 - \epsilon_1^{1/18})$. Since n is sufficiently large, this implies there exist $i \neq j$, with both in $i, j \in \mathcal{B} \cap \mathcal{C}$, and such that $|B_i(\epsilon_1^{1/9}) \cap C_i(\epsilon_1^{1/9})| \neq \emptyset$ and $|B_j(\epsilon_1^{1/9}) \cap C_j(\epsilon_1^{1/9})| \neq \emptyset$.

Fix any $u \in B_i(\epsilon_1^{1/9}) \cap C_i(\epsilon_1^{1/9})$ and $v \in B_j(\epsilon_1^{1/9}) \cap C_j(\epsilon_1^{1/9})$. Since $i \neq j$, we know $u \neq v$. Without loss of generality, let us assume $u < v$ (the other case is similar). Define $B_{i1} = B_i \cap \{b_1, \dots, b_u\}$, $C_{i1} = C_i \cap \{c_1, \dots, c_u\}$, $B_{j0} = B_j \cap \{b_{v+1}, \dots, b_n\}$, and $C_{j0} = C_j \cap \{c_{v+1}, \dots, c_n\}$. Choose any $k \in \mathcal{A} \setminus \{i, j\}$. Note that $K_3[A_k, B_{i1}, C_{j1}] \subseteq E$, while $K_3[A_k, B_{i0}, C_{j0}] \cap E = \emptyset$, so we are done. \square

As a corollary, we show that if $\text{Trip}(\mathcal{H})$ contains arbitrarily large $\overline{H}(k)$, then \mathcal{H} requires non-zero $\text{disc}_{2,3}$ -error.

Corollary 5.6. *Suppose \mathcal{H} is a hereditary property of 3-graphs such that $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. Then \mathcal{H} requires non-zero $\text{disc}_{2,3}$ -error.*

Proof. Suppose towards a contradiction there is some \mathcal{H} with $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$, such that \mathcal{H} admits zero $\text{disc}_{2,3}$ -error. Let $\epsilon = 2^{-28}$ (anything small enough for Theorem 5.5 will do). Let ϵ_1 be as in Proposition 4.2 for k and ϵ^2 . Given $\ell \geq 1$, define $\epsilon'_2(\ell)$ as in Corollary 3.6 for $t = 3$, density $1/\ell$ and ϵ^2 . Then define $\epsilon''_2(\ell)$ be as in Lemma 3.18 for density $1/\ell$, $\gamma = \epsilon^{1/9}$, and let $\epsilon'_1, \epsilon'''_2(\ell)$ be as in Lemma 3.19 for $\epsilon_1, \gamma = \epsilon^{1/9}, 3$, density $1/\ell, \epsilon'_2(\ell)$. Set $\epsilon_2(\ell) = \min\{\epsilon'_2(\ell), \epsilon''_2(\ell), \epsilon'''_2(\ell)\}$.

By assumption there are $T, L, N \geq 1$ such that if $G \in \mathcal{H}$ has at least N vertices, then there exists $1 \leq \ell \leq L, 1 \leq t \leq T$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of $V(G)$ which is $(\epsilon_1, \epsilon_2(\ell))$ -regular with respect to G , with zero $\text{disc}_{2,3}$ -error.

Fix $n \geq N$. By Lemma 3.31, there is $H = (V, E) \in \mathcal{H}$ with $|V| = 3n$ and such that $\text{Trip}(G) \cong \overline{H}(n)$. This means $V = A \sqcup B \sqcup C$ where $A = \{a_i : i \in [n]\}$, $B = \{b_i : i \in [n]\}$ and $C = \{c_i : i \in [n]\}$, and $a_i b_j c_k \in E$ if and only if $i \leq j$.

By assumption, there are $1 \leq \ell \leq L$, $1 \leq t \leq T$, and a $(t, \ell, \epsilon'_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V which is $(\epsilon'_1, \epsilon_2(\ell))$ -regular with respect to H with zero $\text{disc}_{2,3}$ -error. By Theorem 5.5, there are $ijk \in \binom{[k]}{3}$ and sets $A_i \subseteq V_i \cap A$, $B_{j1}, B_{j0} \subseteq V_j \cap B$, and $C_{k1}, C_{k0} \subseteq V_k \cap C$, each of size at least $\epsilon^{1/9}(n/t)$, such that $K_3[A_i, B_{j1}, C_{k1}] \subseteq E$ and $K_3[A_i, B_{j0}, C_{k0}] \cap E = \emptyset$. Let $A' = A_i$, $B' = B_{j1} \cup B_{j0}$, $C' = C_{k1} \cup C_{k0}$, and $V' = A' \cup B' \cup C'$. Then define $H'' = (V', E \cap K_3[A', B', C'])$. By Proposition 3.19, for any $1 \leq \alpha, \beta, \gamma \leq \ell$, $(H'', G_{ijk}^{\alpha, \beta, \gamma}[V'])$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon'_2(\ell))$. But we observe that for any $1 \leq \alpha, \beta, \gamma \leq \ell$,

$$\begin{aligned} & \min\{|E \cap K_3(G_{ijk}^{\alpha, \beta, \gamma}[V'])|, |K_3(G_{ijk}^{\alpha, \beta, \gamma}[V']) \setminus E|\} \\ & \geq \min\{|E \cap K_3(G_{ijk}^{\alpha, \beta, \gamma}) \cap K_2[A_i \cup B_{j0} \cup C_{k0}]|, |(K_3(G_{ijk}^{\alpha, \beta, \gamma}) \cap K_3[A_i \cup B_{j1} \cup C_{k1}]) \setminus E|\}, \\ & = \min\{|K_3(G_{ijk}^{\alpha, \beta, \gamma}) \cap K_2[A_i \cup B_{j0} \cup C_{k0}]|, |K_3(G_{ijk}^{\alpha, \beta, \gamma}) \cap K_3[A_i \cup B_{j1} \cup C_{k1}]|\}. \end{aligned}$$

By Lemma 3.18, for each $u \in \{0, 1\}$, each of $P_{ij}^\alpha[A_i, B_{ju}]$, $P_{ik}^\beta[A_i, C_{ku}]$ and $P_{jk}^\gamma[B_{ju}, C_{ku}]$ have $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$. Consequently, by Corollary 3.6, for each $u \in \{0, 1\}$,

$$|K_3(G_{ijk}^{\alpha, \beta, \gamma}) \cap K_2[A_i \cup B_{ju} \cup C_{ku}]| = \frac{1}{\ell^3}(1 \pm \epsilon^2)|A_i||B_{ju}||C_{ku}| \geq \frac{1}{\ell^3}(1 \pm \epsilon^2)\epsilon^{1/3}|V_i||V_j||V_k|.$$

Combining this with the above, we have that

$$\begin{aligned} \min\{|E \cap K_3(G_{ijk}^{\alpha, \beta, \gamma}[V'])|, |K_3(G_{ijk}^{\alpha, \beta, \gamma}[V']) \setminus E|\} & \geq \epsilon^{1/3} \frac{1}{\ell^3}(1 \pm \epsilon^2)\epsilon^{1/3}|A'||B'||C'| \\ & \geq \epsilon^{2/2} |K_2^{(2)}(G_{ijk}^{\alpha, \beta, \gamma}[V'])|, \end{aligned}$$

where the last inequality is by Corollary 3.6. This shows that $(H'', G_{ijk}^{\alpha, \beta, \gamma}[V'])$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$, and has density in $(\epsilon^2, 1 - \epsilon^2)$. By our choice of ϵ_1 and ϵ_2 , this implies H'' has VC_2 -dimension at least k . By definition H'' is an induced sub-3-graph of $\overline{H}(n)$, but it is not difficult to show that $\text{VC}_2(\overline{H}(n)) < 3$. Thus we have arrived at a contradiction. \square

We can now give a very short proof of Theorem 2.46, which shows that a hereditary 3-graph property admits zero $\text{disc}_{2,3}$ -error if and only if it admits zero vdisc_3 -error.

Proof of Theorem 2.46. Suppose first that \mathcal{H} admits zero $\text{disc}_{2,3}$ -error. By Corollary 5.6, there is some k such that $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$. Clearly this implies $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$, so by Theorem 2.31, \mathcal{H} is close to a WNIP property and vdisc_3 -homogeneous. By Proposition 2.44, \mathcal{H} admits zero vdisc_3 -error.

Suppose on the other hand that \mathcal{H} admits zero vdisc_3 -error. By Theorem 2.32, \mathcal{H} admits zero $\text{disc}_{2,3}$ -error. \square

Combining Theorem 2.46 with Corollary 5.6 yields Theorem 2.37, and combining Theorem 2.46 with Example 5.4 yields Proposition 2.35.

5.4. A generalized strong stable regularity lemma. The last goal of this section is to prove Proposition 2.43, which shows that a weakly stable hereditary 3-graph property admits binary vdisc_3 -error and binary $\text{disc}_{2,3}$ -error. The proof will be partially based on the proof of the following strong regularity for stable graphs.

Theorem 5.7. *For all $k \geq 1$, $\epsilon > 0$ and $f : \mathbb{N} \rightarrow (0, 1]$ non-increasing, there is $M = M(\epsilon, f, d)$ such that the following holds.*

Suppose $G = (V, E)$ is a k -stable graph. Then there is $m \leq M$ and a partition $W = W_0 \cup \dots \cup W_m$ such that $|W_0| \leq \epsilon|W|$, and such that for each $1 \leq i, j \leq m$, (W_i, W_j) satisfies $\text{disc}_2(f(m))$ and $|W_i| = |W_j|$.

Note that the degree of quasirandomness in the pairs avoiding W_0 is allowed to depend on the number of parts in the partition. Theorem 5.7 is similar to the general strong regularity lemma (see [2, 5, 35, 48]), however the conclusions here are strictly stronger than can be obtained in general. Indeed Theorem 5.7 implies the existence of a regular partition with no irregular pairs, so $H(n)$ could not have such a partition.

We will prove Theorem 5.7 in this subsection. This serves as a warm-up for the proof of Proposition 2.43 in the next subsection, which is very similar. This will also lead naturally to the proof of a result needed in Section 6, namely Theorem 5.15.

The proof of Theorem 5.7 uses ideas from [36], where the regularity lemma for stable graphs was first proved. We begin with some terminology from that paper. Given a graph $G = (V, E)$, a subset $X \subseteq V$, and a vertex $v \in V$, we say that X is ϵ -good with respect to v if there is some $u = u(v) \in \{0, 1\}$ such that

$$\left| \frac{|N(v) \cap X|}{|X|} - u \right| \leq \epsilon.$$

The set X is called ϵ -good if it is ϵ -good with respect to every $v \in V$. In other words, X is ϵ -good if every vertex is connected to all but at most $\epsilon|X|$ many elements of X , or is non-connected to all but at most $\epsilon|X|$ many elements of X . In [36], Malliaris and Shelah showed that in a stable graph, any large set of vertices can be partitioned into ϵ -good subsets, all of which have about the same size.

Theorem 5.8 (Theorem 5.18 in [36]). *For all $k \geq 1$ and $\epsilon > 0$, there exists $K = K(\epsilon, k)$ and $N = N(k, \epsilon)$ such that the following holds. Suppose $G = (V, E)$ is a k -stable graph and $W \subseteq V$ has size at least N . Then there is an equipartition of W into at most K subsets, all of which are ϵ -good.*

Our strategy will be to prove a strong version of Theorem 5.8, and then combine it with the following symmetry lemma to obtain a strong regularity lemma.

Lemma 5.9 (Symmetry Lemma). *For all $0 < \epsilon < 1/4$ there is n such that the following holds. Suppose $G = (U \cup W, E)$ is a bipartite graph, $|U|, |W| \geq n$, and $U' \subseteq U$, $W' \subseteq W$ satisfy $|U'| \geq (1 - \epsilon)|U|$ and $|W'| \geq (1 - \epsilon)|W|$. Suppose that for all $u \in U'$,*

$$\max\{|N(u) \cap W|, |\neg N(u) \cap W|\} \geq (1 - \epsilon)|W|,$$

and for all $w \in W'$,

$$\max\{|N(w) \cap U|, |\neg N(w) \cap U|\} \geq (1 - \epsilon)|U|.$$

Then $|E|/|U||W| \in [0, 2\epsilon^{1/2}) \cup (1 - 2\epsilon^{1/2}, 1]$.

Proof. For each $\alpha \in \{0, 1\}$, define

$$\begin{aligned} U_\alpha &= \{u \in U : |N^\alpha(u) \cap W| \geq (1 - \mu)|W|\} \text{ and} \\ W_\alpha &= \{w \in W : |N^\alpha(w) \cap U| \geq (1 - \mu)|U|\}. \end{aligned}$$

It suffices to show there is $\alpha \in \{0, 1\}$ such that $|U_\alpha| \geq (1 - \sqrt{\epsilon})|U|$, since then

$$|E^\alpha| \geq (1 - \epsilon)|W||U_\alpha| \geq (1 - \epsilon)(1 - \sqrt{\epsilon})|W||U| \geq (1 - 2\sqrt{\epsilon})|U||W|.$$

Suppose towards a contradiction that $\max\{|U_0|, |U_1|\} < (1 - \sqrt{\epsilon})|U|$. Note that for each $\alpha \in \{0, 1\}$, $|E^\alpha \cap K_2[U_\alpha, W]| \geq (1 - \epsilon)|W||U_\alpha|$, so by standard arguments, there is $W'_\alpha \subseteq W$ of size at least $(1 - \sqrt{\epsilon})|W|$ such that for all $w \in W'_\alpha$, $|N^\alpha(w) \cap U_\alpha| \geq (1 - \sqrt{\epsilon})|U_\alpha|$. Then $W'' = W_0 \cap W_1$ has size at least $(1 - 2\sqrt{\epsilon})|W|$, and for all $w \in W''$,

$$\begin{aligned} \min\{|N(w) \cap U|, |\neg N(w) \cap U|\} &\geq \min\{|N(w) \cap U_1|, |\neg N(w) \cap U_0|\} \\ &\geq \min\{(1 - \sqrt{\epsilon})|U_0|, (1 - \sqrt{\epsilon})|U_1|\} \\ &> \epsilon|U|. \end{aligned}$$

This implies that $W'' \cap W' = \emptyset$. However, this is not possible since $|W'| \geq (1 - \epsilon)|W|$, $|W''| \geq (1 - 2\sqrt{\epsilon})|W|$, and $\epsilon < 1/4$. \square

The phenomenon described in Lemma 5.9 shows up in many places, for example in model theory (see [43, Lemma 2.8]), and in graph theory (see [7, Lemma 27]). An immediate corollary of Lemma 5.9 is that between any two good sets in a graph, the density of edges is always close to 0 or close to 1.

Corollary 5.10 (A pair of good sets is homogeneous). *For all $0 < \epsilon < 1/4$ there is $n \geq 1$ such that the following holds. Suppose $G = (V, E)$ is a graph and $U, W \subseteq V$ each have size at least n . If U and W are both ϵ -good, then $|E \cap K_2[U, W]|/|K_2[U, W]| \in [0, 2\epsilon^{1/2}) \cup (1 - 2\epsilon^{1/2}, 1]$.*

One can now quickly sketch a proof of a stable regularity lemma as follows. Suppose $G = (V, E)$ is a large k -stable graph. By Theorem 5.8, there is an equipartition \mathcal{P} of V into ϵ -good sets. By Corollary 5.10, every pair of parts from \mathcal{P} has density in $[0, 2\epsilon^{1/2}) \cup (1 - 2\epsilon^{1/2}, 1]$. By Fact 3.20, every pair of parts from \mathcal{P} satisfies $\text{disc}_2(2\epsilon^{1/2})$.

We will use a similar strategy to prove Theorem 5.7. The first step is to prove a strong version of Theorem 5.8. We will work with an equivalent characterization of stability based on the Shelah 2-rank (see [51]). To state this equivalence we need some more notation.

Given an integer $d \geq 1$, let $2^d = \{0, 1\}^d$. In other words, 2^d consists of sequences of 0s and 1s of length d . By convention, we let $2^0 = \{<>\}$, where $<>$ denotes the so-called empty string. Then for $d \geq 1$, we set $2^{<d} := \bigcup_{i=0}^{d-1} 2^i$. Given $0 \leq i \leq j$, $\sigma \in 2^i$ and $\tau \in 2^j$, we write $\sigma \triangleleft \tau$ to denote that σ is a proper initial segment of τ , and $\sigma \trianglelefteq$ to denote that either $\sigma \triangleleft \tau$ or $\sigma = \tau$. By convention $<> \trianglelefteq \sigma$, for any $\sigma \in 2^d$ and $d \geq 0$. We think of $2^{<d}$ as a binary branching tree with root $<>$. Given $\sigma \in 2^i$ and $\sigma' \in 2^j$, $\sigma \wedge \sigma'$ denotes the element of 2^{i+j} obtained by adjoining σ' to the end of σ (e.g. $(01) \wedge (001) = (01001)$).

Definition 5.11. Suppose $G = (V, E)$ is a graph. Given $d \geq 1$, a d -tree in G is a tuple $(a_\sigma)_{\sigma \in 2^{<d}}(b_\rho)_{\rho \in 2^d}$ with the property that for all $\sigma \triangleleft \eta$, $b_\sigma a_\eta \in E$ if and only if $\sigma \wedge 1 \trianglelefteq \eta$.

The *rank* of G is

$$\text{rk}(G) := \max \left(\{0\} \cup \{d \geq 1 : \text{there is a } d\text{-tree in } G\} \right).$$

Given a d -tree as in Definition 5.11, we refer to the elements $\{a_\eta : \eta \in 2^d\}$ as the *leaves* and the elements $\{b_\sigma : \sigma \in 2^{<d}\}$ as the *nodes*. There is a fundamental relationship between tree-rank and the order property.

Theorem 5.12 (Shelah [51]).

- (1) For all $k \geq 1$ there is a $d = d(k) \geq 1$ such that the following holds. Suppose $G = (V, E)$ is a graph with rank d , witnessed by $(a_\eta)_{\eta \in 2^d}(b_\sigma)_{\sigma \in 2^{<d}}$. Then there are $a_1, \dots, a_k \subseteq \{a_\eta : \eta \in 2^d\}$ and $b_1, \dots, b_k \in \{b_\sigma : \sigma \in 2^{<d}\}$ such that $a_i b_j \in E$ if and only if $i \leq j$.
- (2) For all $d \geq 1$ there is a $k = k(d) \geq 1$ such that the following holds. Suppose $G = (V, E)$ is a graph with the k -order property, witnessed by $(a_i)_{i \in [k]}(b_i)_{i \in [k]}$. Then there are $\{a_\eta : \eta \in 2^d\} \subseteq \{a_i : i \in [k]\}$ and $\{b_\sigma : \sigma \in 2^{<d}\} \subseteq \{b_i : i \in [k]\}$ such that for all $\sigma \triangleleft \eta$, $a_\eta b_\sigma \in E$ if and only if $\sigma \wedge 1 \trianglelefteq \eta$.

Theorem 5.12 tells us that a graph with sufficiently large rank will have the k -order property, and on the other hand, a graph with a sufficiently long order property will have rank at least d . We now define a notion of rank for subsets of vertices.

Definition 5.13. Suppose $G = (V, E)$. The *rank* of a set $A \subseteq V$ is

$$\text{rk}(A) := \max(\{0\} \cup \{d \geq 1 : \text{there exists a } d\text{-tree in } G \text{ with leaves in } A\}).$$

Note that the rank of A is 0 if there do *not* exist $a_0, a_1 \in A$ and $b_{<} \in V$ such that $b_{<} a_1 \in E$ and $b_{<} a_0 \notin E$. Thus, $\text{rk}(A) = 0$ if and only if A is 0-good in G .

Observe that if A has rank $d > 0$, then for any $b \in V$, one of $N(b) \cap A$ or $(\neg N(b)) \cap A$ must have rank less than d . This observation will be crucial in our proofs. We will also use the easy fact that if $A' \subseteq A$, then $\text{rk}(A') \leq \text{rk}(A)$.

In our first lemma, we show that in any bipartite graph $G = (U \cup W, E)$ with tree rank d , we can partition W into sets of rank less than d , and sets which are somewhat good.

Lemma 5.14. Suppose $d \geq 1$, $\epsilon > 0$ and $f : \mathbb{N} \rightarrow (0, 1]$ is a non-increasing function satisfying $\lim_{n \rightarrow \infty} f(n) = 0$.

Suppose $G = (U \cup W, E)$ is a bipartite graph and $\text{rk}(W) \leq d$. Then there is an integer $t \leq |W|$, a sequence (s_1, \dots, s_t) , and a partition $W = \bigcup_{i \in [t]} \bigcup_{j \in [s_i]} W_{i,j}$ such that the following hold, where for each $1 \leq i \leq t$, $W_i := \bigcup_{j \in [s_i]} W_{i,j}$.

- (1) For each $1 \leq i \leq t$ and $1 \leq j \leq s_i$, $\text{rk}(W_{i,j}) < d$, and

$$|W_{i,j}| = f(i) |W \setminus (\bigcup_{1 \leq i' < i} W_{i'} \cup \bigcup_{j'=1}^{j-1} W_{i,j'})|.$$

- (2) For each $1 \leq i \leq t$, $W \setminus (\bigcup_{i'=1}^i W_{i'})$ is $f(i)$ -good.

Proof. Suppose $G = (U \cup W, E)$ is a bipartite graph in which $\text{rk}(W) \leq d$. We define an integer t , and sequences (s_0, \dots, s_t) , Z_0, \dots, Z_t , and W_1, \dots, W_t as follows.

Step 0: Set $s_0 = 0$, $Z_0 = W$ and $W_0 = \emptyset$.

Step α : Suppose $\alpha > 0$ and we have chosen $Z_0, \dots, Z_{\alpha-1}, W_0, \dots, W_{\alpha-1}$, and for all $0 \leq \alpha' < \alpha$, $s_{\alpha'}$ and $W_{\alpha',1}, \dots, W_{\alpha',s_{\alpha'}}$ so that the following hold.

- For all $0 \leq \alpha' < \alpha$, $W_{\alpha'} = \bigcup_{u=1}^{s_{\alpha'}} W_{\alpha',u}$, and $Z_{\alpha'} = W \setminus (\bigcup_{\beta < \alpha'} W_{\beta})$,
- For each $0 \leq \alpha' < \alpha$ and $1 \leq u < s_{\alpha'}$, $|W_{\alpha',u}| = f(\alpha')|W \setminus Z_{\alpha'-1} \cup (\bigcup_{v=1}^{u-1} W_{\alpha',v})|$,
- For each $0 < \alpha' < \alpha$ and $Z_{\alpha'}$ if $f(\alpha')$ -good.

Substep $(\alpha, 0)$: Let $Z_{\alpha,0} = Z_{\alpha-1} = W \setminus (\bigcup_{\beta < \alpha} W_{\beta})$ and $W_{\alpha,0} = \emptyset$. If $Z_{\alpha,0} = \emptyset$, or $Z_{\alpha,0}$ is 0-good, set $t = \alpha - 1$ and end the construction.

Substep $(\alpha, v+1)$: Suppose $v \geq 0$, and we have defined $W_{\alpha,0}, \dots, W_{\alpha,v}$, $Z_{\alpha,0}, \dots, Z_{\alpha,v}$ so that for each $0 \leq v' \leq v$, $Z_{\alpha,v'} = Z_{\alpha-1} \setminus (\bigcup_{v'=0}^{v'} W_{\alpha,v'})$, and for each $1 \leq v' \leq v$, $\text{rk}(W_{\alpha,v'}) < d$, and $|W_{\alpha,v'}| = f(\alpha)|Z_{\alpha,v'-1}|$. Define

$$U_{\alpha}^{v+1} = \{u \in U : \min\{|N(u) \cap Z_{\alpha,v}|, |\neg N(u) \cap Z_{\alpha,v}^{\alpha}| \} \geq f(\alpha)|Z_{\alpha,v}|\}.$$

If $U_{\alpha}^{v+1} = \emptyset$, set $s_{\alpha} = v$, $W_{\alpha} = \bigcup_{1 \leq v' \leq v} W_{\alpha,v'}$, $Z_{\alpha} = Z_{\alpha,v}$, and go to step $\alpha + 1$.

If $U_{\alpha}^{v+1} \neq \emptyset$, choose any $u \in U_{\alpha}^{v+1}$, and let $X \in \{|N(u) \cap Z_{\alpha,v}^{\alpha}|, |\neg N(u) \cap Z_{\alpha,v}^{\alpha}|\}$ be a set rank less than d (such an X exists because $\text{rk}(W) \leq d$). Define $W_{\alpha,v+1}$ to be any subset of X of size $f(\alpha)|Z_{\alpha,v}^{\alpha}|$. Since $\text{rk}(X) < d$ and $W_{\alpha,v+1} \subseteq X$, we also have $\text{rk}(W_{\alpha,v+1}) < d$. Set $Z_{\alpha,v+1} = Z_{\alpha,v} \setminus W_{\alpha,v+1}$ and go to step $(\alpha, v+2)$.

Since W is finite and $\lim_{n \rightarrow \infty} f(n) = 0$, there will be some $\alpha = t$ where this ends. \square

We now prove a much stronger version of Lemma 5.14. This is the main technical lemma behind Theorem 5.7, and plays a crucial role in the proof of Theorem 2.49.

Theorem 5.15. *Suppose $d \geq 0$. For all $\epsilon > 0$ and non-increasing functions $f : \mathbb{N} \rightarrow (0, 1]$ with $\lim_{n \rightarrow \infty} f(n) = 0$, there is $M \geq 1$ such that the following hold.*

Suppose $m \geq 1$, $G = (U \cup W, E)$ is a bipartite graph, and $W_1 \cup \dots \cup W_m$ is a partition of W so that $\text{rk}(W_i) \leq d$ for all $i \in [m]$.

Then there are $m' \leq M$, and a set $\Omega \subseteq [m]$, such that $|\bigcup_{u \in \Omega} W_u| \leq \epsilon|W|$, and for each $u \in [m] \setminus \Omega$, there are $s_u \leq m'$ and a partition

$$W_u = W_{u,0} \cup W_{u,1} \cup \dots \cup W_{u,s_u},$$

such that $|W_{u,0}| \leq \epsilon|W_u|$, and such that $W_{u,i}$ is $f(m')$ -good for each $1 \leq i \leq s_u$.

Before we give the proof, we point out some important features of this theorem. First, observe that the bound M does not depend on m , the number of parts in the partition we start with. Further, a graph need not be stable to satisfy the hypotheses of Theorem 5.15. Indeed, when m is large, G could have large rank even though each W_i has small rank. We will use Theorem 5.15 in such a scenario in Section 6.

PROOF OF THEOREM 5.15: To ease notation, let I be the set of non-increasing functions $f : \mathbb{N} \rightarrow (0, 1]$ satisfying $\lim_{n \rightarrow \infty} f(n) = 0$. We proceed by induction on $d \geq 0$.

Case $d = 0$: Suppose $d = 0$. Given $\epsilon > 0$, and $f \in I$, set $M = 1$. Suppose $G = (U \cup W, E)$ is a bipartite graph and assume W_1, \dots, W_m partition W such that $\text{rk}(W_i) \leq 0$ for each $i \in [m]$. This means each W_i is 0-good in G . Setting $m' = 1$ and $W'_{i,1} = W_i$ for each $i \in [m]$, we are done.

Case $d > 0$: Suppose now that $d > 0$ and assume by induction we have defined $M(d', f, \epsilon)$ for all $0 \leq d' < d$, $\epsilon > 0$, and $f \in I$.

Fix $\epsilon > 0$ and $f \in I$. Without loss of generality, assume $\epsilon < 1/4$. We inductively define a new function $f' : \mathbb{N} \rightarrow (0, 1]$, and a sequence $(M_i, g_i)_{i \in \mathbb{N}}$, where each $M_i \in \mathbb{N}$, and each $g_i \in I$.

Step 1: Define g_1 by setting $g_1(x) = \epsilon^4 f(2\epsilon^{-4}x)$. Then let $M_1 = M(d-1, g_1, \epsilon^8)$ and $f'(1) = g_1(M_1)/2$.

Step $i+1$: Suppose $i \geq 1$, and assume that for all $1 \leq j \leq i$, we have defined g_j , M_j , and $f'(j)$. Set $P_i = \sum_{j=1}^i \epsilon^{-2} f'(j)^{-1}$ and define g_{i+1} by setting $g_{i+1}(x) = \epsilon^4 f(\epsilon^{-4}(x \cdot (P_i + 1)))$. Then set $M_{i+1} = M(d-1, g_{i+1}, \epsilon^8)$ and $f'(i+1) = f'(i)(i+2)^{-1} g_{i+1}(M_{i+1})$.

This finishes the definition of $(M_i, g_i)_{i \in \mathbb{N}}$ and of f' . It is straightforward to check that by construction, and because $f \in I$, we have that $f' \in I$, and $g_i \in I$ for all $i \geq 1$. Let $S_1 = \lceil \epsilon^{-2} \rceil$ and choose $M(d, \epsilon, f) \gg M(d-1, \epsilon^8, f) \epsilon^{-8} \prod_{i=1}^{S_1} M_i P_i$.

Suppose $G = (U \cup W, E)$ is a bipartite graph and $W = W_1 \cup \dots \cup W_m$ is a partition so that for each $i \in [m]$, $\text{rk}(W_i) \leq d$. Let ℓ be the number of W_i with $\text{rk}(W_i) = d$. If $\ell = 0$, then we are done by the inductive hypothesis.

Thus we may assume $\ell > 0$. By relabeling if necessary, we may assume W_1, \dots, W_ℓ have rank d and $W_{\ell+1}, \dots, W_m$ have rank less than d .

For each $1 \leq u \leq \ell$, apply Lemma 5.14 with parameters d , ϵ^2 , and f' to the set W_u to obtain an integer $t(u) \leq |W_u|$, a sequence $(s_i(u))_{i \in [t(u)]}$, and a partition $W_u = \bigcup_{i \in [t(u)]} \bigcup_{j \in [s_i(u)]} V_{i,j}(u)$ such that the following hold, where for each $1 \leq i \leq t(u)$,

$$V_i(u) := \bigcup_{j \in [s_i(u)]} V_{i,j}(u) \text{ and } Z_{i,j}(u) = W_u \setminus \left(\bigcup_{i'=1}^{i-1} V_{i'}(u) \cup \bigcup_{j'=1}^{j-1} V_{i,j'}(u) \right).$$

- (i) For each $i \in [t(u)]$ and $j \in [s_i(u)]$, $|V_{i,j}(u)| = f'(i)|Z_{i,j}(u)|$ and $\text{rk}(V_{i,j}(u)) < d$,
- (ii) For each $i \in [t(u)]$, $W_u \setminus (\bigcup_{i'=1}^i V_{i'}(u))$ is $f'(i)$ -good.

Let $t = \max\{t(u) : 1 \leq u \leq \ell\}$. Given $u \in [\ell]$, if $t(u) < i \leq t$, set $V_i(u) = \emptyset$. Then for each $1 \leq i \leq t$, define $Y_i = \bigcup_{u=1}^\ell V_i(u)$, and let i_1 be minimal such that $|Y_{i_1}| \leq \epsilon^8 |W|$. Clearly such an i_1 exists and is at most ϵ^{-8} . Then define

$$\Omega_0 = \{u \in [\ell] : |V_{i_1}(u)| > \epsilon^4 |W_u|\}.$$

Note by definition, $\epsilon^4 |\bigcup_{u \in \Omega_0} W_u| < |Y_{i_1}| \leq \epsilon^8 |W|$, so $|\bigcup_{u \in \Omega_0} W_u| \leq \epsilon^4 |W|$.

Now fix $u \in [\ell] \setminus \Omega_0$. Set $J(u) = \{j \in [i_1 - 1] : |V_j(u)| \leq \epsilon^4 f'(i_1 - 1) |W_u|\}$. Observe that by definition,

$$|\bigcup_{j \in J(u)} V_j(u)| \leq |J(u)| \epsilon^4 f'(i_1 - 1) |W_u| \leq (i_1 - 1) \epsilon^4 f'(i_1 - 1) |W_u| < \epsilon^4 |W_u|,$$

where the last inequality is because, by definition, $f'(i_1) < (i_1)^{-1}$. Set $I(u) = [i_1 - 1] \setminus J(u)$, and for each $i \in I(u)$, define

$$T_i(u) = \{j \in [s_i(u)] : |V_{i,j}(u)| \geq \epsilon^2 f'(i) |V_i(u)|\},$$

We then let $t_i(u) = |T_i(u)|$, and observe that $t_i(u)$ is at most $\epsilon^{-2} f'(i)^{-1}$. Now define

$$V_{i,0}(u) = \bigcup_{j \in [s_i(u)] \setminus T_i(u)} V_{i,j}(u).$$

We claim $|V_{i,0}(u)| \leq 2\epsilon^2 |V_i(u)|$. If $V_{i,0}(u) = \emptyset$, this is obvious. If $V_{i,0}(u) \neq \emptyset$, consider $j_0 = \min([s_i(u)] \setminus T_i(u))$. By (i), and since $j_0 \notin [s_i(u)] \setminus T_i(u)$,

$$|V_{i,j_0}(u)| = f'(i) |Z_{i,j_0}(u)| \leq \epsilon^2 f'(i) |V_i(u)|.$$

Consequently $|Z_{i,j_0}(u)| \leq \epsilon^2 |V_i(u)|$. By construction, and minimality of j_0 , we have $V_{i,0}(u) \subseteq V_{i,j_0}(u) \cup Z_{i,j_0}(u)$. This shows $|V_{i,0}(u)| \leq 2\epsilon^2 |V_i(u)|$. We now set

$$V_0(u) = V_{i_1}(u) \cup \left(\bigcup_{i \in J(u)} V_j(u) \right) \cup \bigcup_{i \in I(u)} V_{i,0}(u).$$

By above,

$$|V_0(u)| \leq \epsilon^4 |W_u| + \epsilon^4 |W_u| + \sum_{i \in I(u)} |V_{i,0}(u)| \leq 2\epsilon^4 |W_u| + \sum_{i \in I(u)} 2\epsilon^2 |V_i(u)| \leq 4\epsilon^2 |W_u|.$$

Recall that by (ii), $Z_{i_1}(u) = W_u \setminus (\bigcup_{v=1}^{i_1} V_i(u))$ is $f'(i_1)$ -good. If $|Z_{i_1}(u)| < \epsilon^2 |W_u|$, set $V'_0 = V_0 \cup Z_{i_1}$ and $V_{s+1}(u) = \emptyset$. If $|Z_{i_1}(u)| \geq \epsilon^2 |W_u|$, set $V'_0 = V_0$ and $V_{s+1}(u) = Z_{i_1}$. In either case, $|V'_0| \leq 5\epsilon^2 |W_u|$, and we now have a partition

$$W_u = V'_0(u) \cup V_{s+1}(u) \cup \bigcup_{i \in I_i(u)} \bigcup_{j \in T_i(u)} V_{i,j}(u).$$

Set $S(u) := \sum_{i \in I(u)} t_i(u)$. Since each $t_i(u) \leq (\epsilon^2 f'(i))^{-1}$, we have

$$S(u) \leq \sum_{i=1}^{i_1-1} \epsilon^{-2} f'(i)^{-1} = P_{i_1-1}.$$

Let $R_1(u), \dots, R_{S(u)}(u)$ be an enumeration of $\{V_{i,j}(u) : i \in I(u), j \in T_i(u)\}$.

Now consider the partition $\mathcal{P} = \{R_i(u) : u \in [\ell] \setminus \Omega_0, i \in [S(u)]\} \cup \{W_{\ell+1}, \dots, W_m\}$ of the set $W' := \bigcup_{X \in \mathcal{P}} X$. Note that each element in \mathcal{P} has rank at most $d-1$ in $G' = G[U \cup W']$. By the induction hypothesis, applied to $G' = G[U \cup W']$, there are $\Omega_1 \subseteq \mathcal{P}$ and $m' \leq M_{i_1} = M(t-1, g_{i_1}, \epsilon^8)$ such that the following hold.

- $|\bigcup_{X \in \Omega_1} X| \leq \epsilon^8 |W'|$,
- For each $\ell+1 \leq u \leq m$ with $W_u \notin \Omega_1$, there is an integer $c_u \leq m'$, and a partition

$$W_u = W_{u,0} \cup W_{u,1} \cup \dots \cup W_{u,c_u},$$

so that $|W_{u,0}| \leq \epsilon^8 |W_u|$ and such that for each $v \in [c_u]$, $W_{u,v}$ is $g_{i_1}(m')$ -good in G' ,

- For each $u \in [\ell] \setminus \Omega_0$, and each $i \in [S(u)]$ with $R_i(u) \notin \Omega_1$, there is an integer $b_i(u) \leq m'$ and a partition

$$R_i(u) = R_{i,0}(u) \cup R_{i,1}(u) \cup \dots \cup R_{i,b_i(u)}(u),$$

so that $|R_{i,0}(u)| \leq \epsilon^8 |R_i(u)|$ and such that for each $v \in [b_i(u)]$, $R_{i,v}(u)$ is $g_{i_1}(m')$ -good in G' .

Note that since G is bipartite, and $W' \subseteq W$, any $g_{i_1}(m')$ -good subset of W' in G' is also $g_{i_1}(m')$ -good in G . Let $\Omega_2 = \{u \in [\ell] : |W_u \cap (\bigcup_{X \in \Omega_1} X)| \geq \epsilon^4 |W_u|\}$. Observe that

$$\epsilon^4 \left| \sum_{u \in \Omega_2} W_u \right| \leq \left| \bigcup_{X \in \Omega_1} X \right| \leq \epsilon^8 |W'|,$$

and consequently, $\left| \sum_{u \in \Omega_2} W_u \right| \leq \epsilon^4 |W'| \leq \epsilon^4 |W|$. Now define

$$\Omega = \{u \in [\ell] : u \in \Omega_0 \cup \Omega_2\} \cup \{u \in [\ell + 1, m] : W_u \in \Omega_1\}.$$

Observe that $\left| \bigcup_{X \in \Omega} X \right| \leq \epsilon^4 |W| + \epsilon^8 |W'| < \epsilon |W|$. For each $u \in [\ell] \setminus \Omega$, define

$$W_{u,0} = V'_0(u) \cup \bigcup_{R_{i,v}(u) \in \Omega_1} R_{i,v}(u).$$

Further, if $V_{s+1}(u) = \emptyset$, define $c_u := |\{R_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \setminus \Omega_1|$, and let $W_{u,1}, \dots, W_{u,c_u}$ be an enumeration of $\{R_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \setminus \Omega_1$. On the other hand, if $V_{s+1}(u) \neq \emptyset$, define

$$c_u := 1 + |\{R_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \setminus \Omega_1|,$$

and let $W_{u,1}, \dots, W_{u,c_u}$ enumerate $\{V_{s+1}(u)\} \cup (\{R_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \setminus \Omega_1)$. Since $u \notin \Omega_2$,

$$|W_{u,0}| \leq |V'_0| + \left| \bigcup_{R_{i,v}(u) \in \Omega_1} R_{i,v}(u) \right| \leq 5\epsilon^2 |W_u| + \epsilon^4 |W_u| \leq 6\epsilon^2 |W_u|.$$

Further, by construction, for all $u \in [\ell] \setminus \Omega$, and all $1 \leq v \leq c_u$, $W_{u,v}$ is $\max\{f'(i_1), g_{i_1}(m')\}$ -good, and

$$c_u \leq S(u)m' + 1 \leq P_{i_1-1}m' + 1 \leq m'(P_{i_1-1} + 1) \leq M_{i_1}(P_{i_1-1} + 1).$$

We have now constructed, for all $u \in [m] \setminus \Omega$, a partition

$$W_u = W_{u,0} \cup W_{u,1} \cup \dots \cup W_{u,c_u},$$

where $|W_{u,0}| \leq 6\epsilon^2 |W_u| < \epsilon |W_u|$, and where for all $1 \leq v \leq c_u$, $W_{u,v}$ is μ^* -good, where $\mu^* := \max\{g_{i_1}(m'), f'(i_1)\}$. Further, for each $u \in [m] \setminus \Omega$, $c_u \leq M_{i_1}(P_{i_1-1} + 1) \leq M$. Since by construction, $\mu^* \leq f(M_{i_1}(P_{i_1-1} + 1))$, this finishes the proof. \square

We can now give an easy proof of Theorem 5.7.

PROOF OF THEOREM 5.7: Fix $k \geq 1$, $\epsilon > 0$ and $f : \mathbb{N} \rightarrow (0, 1]$ non-increasing with $\lim_{n \rightarrow \infty} f(n) = 0$. Let $d = d(k)$ be as in part (2) of Theorem 5.12, and define a function $\psi : \mathbb{N} \rightarrow (0, 1]$ by setting $\psi(x) = \epsilon^4 f(\epsilon^{-4}x)^2 / 6x$. Let $M = M(d, \psi, \epsilon^2)$ be as in Theorem 5.15.

Suppose $G = (V, E)$ is a k -stable graph on at least N vertices. Consider $\text{Bip}(G) = (U \cup W, E')$. Note $\text{rk}(W) \leq d$ in $\text{Bip}(G)$, by Theorem 5.12. By Theorem 5.15, there some $m \leq M$ and a partition $W = W_0 \cup \dots \cup W_m$ such that $|W_0| \leq \epsilon^2|W|$ and such that for each $1 \leq i \leq m$, W_i is $\psi(m)$ -good in $\text{Bip}(G)$. By definition of $\text{Bip}(G)$, this yields a partition $V = V_0 \cup \dots \cup V_m$ so that $|V_0| \leq \epsilon^2|V|$ and for each $i \in [m]$, V_i is $\psi(m)$ -good in G . Let $\Omega = \{i \in [m] : |V_i| < \epsilon^2|V|/m\}$, and let $V'_0 = V_0 \cup \bigcup_{i \in \Omega} V_i$. Note

$$|V'_0| \leq \epsilon^2|V| + |\Omega|\epsilon^2|V|/m \leq 2\epsilon^2|V|.$$

Set $K = \lfloor \epsilon^4|V|/m \rfloor$. For each $i \in [m] \setminus \Omega$, choose a partition $V_i = V_{i,0} \cup V_{i,1} \cup \dots \cup V_{i,r_i}$ so that for each $1 \leq v \leq r_i$, $|V_v| = K$ and $|V_{i,0}| < K$. Now let $R_0 = V'_0 \cup \bigcup_{i \in [m] \setminus \Omega} V_{i,0}$, and let $R_1, \dots, R_{m'}$ be an enumeration of $\{V_{i,j} : i \in [m] \setminus \Omega, j \in [r_i]\}$. Then $m' \leq \epsilon^{-4}m$, and

$$|R_0| \leq 2\epsilon^2|V| + mK \leq \epsilon^2|V| + \epsilon^4|V| < \epsilon|V|.$$

For each $1 \leq i \leq m'$, R_i is contained in some $V_{u,v}$ which is $\psi(m)$ -good, so for all $a \in V$, there is some $\delta = \delta(a)$ so that $|N^\delta(a) \cap V_{u,v}| \leq \psi(m)|V_{u,v}|$. Then

$$|N^\delta(a) \cap R_i| \leq \psi(m)|V_{u,v}| = \frac{\psi(m)|V_{u,v}|}{K}|R_i| \leq \epsilon^{-4}m\psi(m)|R_i| < f(m')^2|R_i|/2,$$

where the second inequality is by definition of K , and since $|V_{u,v}| \leq |V|$, and the last inequality is by definition of ψ and because $m' \leq \epsilon^{-4}m$. Thus for each $1 \leq i \leq m'$, R_i is $f(m')^2/2$ -good.

By Corollary 5.10, for each $1 \leq i, j \leq m'$, the pair (R_i, R_j) has density in $[0, f(m')] \cup (1 - f(m'), 1]$. By Proposition 3.20, (R_i, R_j) satisfies $\text{disc}_2(f(m'))$. \square

5.5. Weakly stable properties admit binary error. In this subsection we prove Proposition 2.43, which shows that a weakly stable hereditary 3-graph property admits binary vdisc_3 -error and $\text{disc}_{2,3}$ -error.

Recall that if $H = (V, E)$ is a 3-graph and $a \in V$, then H_a is the graph $(V, N(a))$, where $N(a) = \{bc \in \binom{V}{2} : abc \in E\}$. Recall further that H is weakly k -stable if and only if H_a is k -stable for all $a \in V$. Given a weakly stable $H = (V, E)$, our strategy will be to find strong decompositions of H_a for all $a \in V$, yielding a partition of $V \times V$ with special properties. We will then apply the multicolored regularity lemma to the partition of $V \times V$.

The main technical lemma is Lemma 5.16 below. Its proof is closely based on the proof of Theorem 5.15 of the prior subsection. We will use the following notation in the proof. Given a set V , $a \in V$, and $\alpha \subseteq V \times V$, we will let $\alpha(a) := \{b \in V : (a, b) \in \alpha\}$. Given a 3-graph $H = (V, E)$, $a \in V$, and $C, B \subseteq V$, we let $H_a^{\text{bip}}[C, B]$ be the graph with vertex set $\{u_c : c \in C\} \cup B$ and edge set $\{u_c b : cb \in E(H_a)\}$. Note that if $B' \subseteq B$, and B' is ϵ -good in $H_a^{\text{bip}}[V, B]$, then B' is also ϵ -good in H_a . Further, the rank of B' in $H_a^{\text{bip}}[V, B]$ is the same as the rank of B' in H_a .

Lemma 5.16. *Suppose $d \geq 0$. For all $\epsilon > 0$, and non-increasing functions $f : \mathbb{N} \rightarrow [0, 1]$ with $\lim_{n \rightarrow \infty} f(n) = 0$, there is $M \geq 1$ such that the following hold. Suppose $H = (V, E)$ is a 3-graph, $\alpha \subseteq V \times V$, and $\alpha = \alpha_1 \cup \dots \cup \alpha_m$ is a partition such that for all $a \in V$, and all $i \in [m]$, $\text{rk}(\alpha_i(a)) \leq d$ in H_a .*

Then there exists $1 \leq s \leq M$ and $\Omega \subseteq [m]$ so that $|\bigcup_{u \in \Omega} \alpha_u| \leq \epsilon|V|^2$, and such that for each $u \in [m] \setminus \Omega$, there is $s_u \leq s$, and a partition $\alpha_u = \alpha_{u,0} \cup \alpha_{u,1} \cup \dots \cup \alpha_{u,s_u}$ with $|\alpha_{u,0}| \leq \epsilon|\alpha_u|$, and so that for all $a \in V$ and $1 \leq i \leq s_u$, $\alpha_{u,i}(a)$ is $f(s)$ -good in H_a .

Proof. We proceed by induction on $d \geq 0$. To ease notation, let I be the set of non-increasing functions $f : \mathbb{N} \rightarrow [0, 1]$ with $\lim_{n \rightarrow \infty} f(n) = 0$.

Case $d = 0$: Suppose $d = 0$. Given $\epsilon > 0$ and $f \in I$, set $M = 1$. Suppose $H = (V, E)$ is a 3-graph, $\alpha \subseteq V \times V$, and $\alpha = \alpha_1 \cup \dots \cup \alpha_m$ is a partition such that for all $a \in V$ and $i \in [m]$, $\text{rk}(\alpha_i(a)) \leq 0$ in H_a . This means that for all $a \in V$, each of $\alpha_1(a), \dots, \alpha_m(a)$ are 0-good in H_a . Setting $s = 1$ and $\alpha_{i,1} = \alpha_i$ for each $1 \leq i \leq m$, we are done.

Case $d > 0$: Suppose now that $d > 0$ and assume by induction we have defined $M(d - 1, f, \epsilon)$ for all $\epsilon > 0$ and $f \in I$.

Fix $\epsilon > 0$ and $f \in I$. We inductively define a new function $f' \in I$, and a sequence $(M_i, g_i)_{i \in \mathbb{N}}$, where each $M_i \in \mathbb{N}$ and each $g_i \in I$.

Step 1: Define g_1 by setting $g_1(x) = \epsilon^8 f(2x + 1)$. Choose M_1 by applying the inductive hypothesis to $(d - 1, g_1, \epsilon^9)$, and set $f'(1) = g(M_1)/2$.

Step $i + 1$: Suppose we have defined g_j , M_j , and $f'(j)$ for each $1 \leq j \leq i$. Let $P_i = \sum_{j=1}^i (\epsilon f'(j))^{-1}$, and define g_{i+1} by setting $g_{i+1} = g((P_i + 1) \cdot x)$. Choose M_{i+1} by applying the inductive hypothesis to $(d - 1, g_{i+1}, \epsilon^8)$, and set $f'(i + 1) = g(M_{i+1})(i + 2)^{-1} f'(i)$.

This finishes the definition of f' and our sequence $(M_i, g_i)_{i \in \mathbb{N}}$. Set

$$M = \max\{M_i : i \leq 2\epsilon^{-1}\}.$$

Now suppose $H = (V, E)$ is a 3-graph, $\alpha \subseteq V \times V$, and $\alpha = \alpha_1 \cup \dots \cup \alpha_m$ is a partition such that for all $a \in V$ and $i \in [m]$, $\text{rk}(\alpha_i(a)) \leq d$ in H_a . If $|\alpha| \leq \epsilon|V|^2$, then we are done by setting $s = 1$ and $\Omega = [m]$. So we may assume $|\alpha| > \epsilon|V|^2$.

For each $1 \leq \ell \leq m$, let

$$V_\ell = \{a \in V : \ell\text{-many elements in } \{\alpha_1(a), \dots, \alpha_m(a)\} \text{ have rank } d \text{ in } H_a\}.$$

For each $u \in [m]$, let

$$V_u^* = \{a \in V : \text{rk}(\alpha_u(a)) = d\} \text{ and } \alpha_u^* = \alpha_u \cap (V_u^* \times V).$$

For each $u \in [m]$ and $a \in V_u^*$, Apply Lemma 5.14 with parameters d , ϵ^8 , and f' to the set $\alpha_u(a)$ in $H_a[V, \alpha_u(a)]$ to obtain some $t^a(u) \leq |\alpha_u(a)|$, a sequence $(s_i^a(u))_{i \in [t^a(u)]}$, and a partition $\alpha_u(a) = \bigcup_{i \in [t^a(u)]} \bigcup_{j \in [s_i^a(u)]} V_{i,j}^a(u)$ such that the following hold, where for each $1 \leq i \leq t^a(u)$, $V_i^a(u) := \bigcup_{j \in [s_i^a(u)]} V_{i,j}^a(u)$ and $Z_{i,j}^a(u) = \alpha_u(a) \setminus (\bigcup_{i'=1}^{i-1} V_{i'}^a(u) \cup \bigcup_{j'=1}^{j-1} V_{i,j'}^a(u))$.

- (i) For each $i \in [t^a(u)]$ and $j \in [s_i^a(u)]$, $|V_{i,j}^a(u)| = f'(i)|Z_{i,j}^a(u)|$, and $\text{rk}(V_{i,j}^a(u)) < d$ in H_a ,
- (ii) For each $i \in [t^a(u)]$, $\alpha_u(a) \setminus (\bigcup_{i'=1}^i V_{i'}^a(u))$ is $f'(i)$ -good in H_a .

For each $u \in [m]$, let $t(u) = \max\{t^a(u) : a \in V_u^*\}$. Then for each $i \leq t(u)$, let

$$s_i(u) := \max\{s_i^a(u) : a \in V_u^* \text{ and } i \leq t^a(u)\}.$$

For each $a \in V_u^*$, if $t^a(u) < i \leq t(u)$, set $V_i^a(u) = \emptyset$, and if $i \leq t^a(u)$ but $j > s_i^a(u)$, set $V_{i,j}^a(u) = \emptyset$. For each $i \in [t(u)]$ and $j \in [s_i(u)]$, we now define

$$\beta_{i,j}(u) := \{(a, b) \in V^2 : a \in V_u^* \text{ and } b \in V_{i,j}^a(u)\}.$$

For each $i \in [t(u)]$, set $\beta_i(u) = \bigcup_{j \in [s_i(u)]} \beta_{i,j}(u)$. Finally, we set $t = \max\{t(u) : u \in [m]\}$, and given $u \in [m]$, if $t(u) < i \leq t$, define $\beta_i(u) = \emptyset$.

We can now define, for each $i \in [t]$, $\beta_i = \bigcup_{u \in [m]} \beta_i(u)$. Let i_1 be minimal such that $|\beta_{i_1}| \leq \epsilon^8 |V|^2$. Clearly such an i_1 exists and is at most ϵ^{-8} . We now set

$$\Omega_0 = \{u \in [m] : |\beta_{i_1}(u)| \geq \epsilon^4 |\alpha_u|\}.$$

Note that by definition, $\epsilon^4 |\bigcup_{u \in \Omega_0} \alpha_u| \leq |\beta_{i_1}| \leq \epsilon^8 |V|^2$, so $|\bigcup_{u \in \Omega_0} \alpha_u| \leq \epsilon^4 |V|^2$. For each $u \in [m] \setminus \Omega_0$, set

$$J(u) = \{j \in [i_1 - 1] : |\beta_j(u)| \leq \epsilon^4 f'(i_1 - 1) |\alpha_u|\}.$$

By definition, $|\bigcup_{j \in J(u)} \beta_j(u)| \leq |J(u)| \epsilon^4 f'(i_1 - 1) |\alpha_u| \leq (i_1 - 1) \epsilon^4 f'(i_1 - 1) |\alpha_u| < \epsilon^4 |\alpha_u|$, where the last inequality is because, by construction, $f'(i_1 - 1) < (i_1)^{-1}$. We then set $I(u) = [i_1 - 1] \setminus J(u)$, and for each $i \in I(u)$, let

$$T_i(u) = \{j \in [s_i(u)] : |\beta_{i,j}(u)| \geq \epsilon^2 f'(i) |\beta_i(u)|\}.$$

Setting $t_i(u) = |T_i(u)|$, we have that $t_i(u)$ is at most $\epsilon^{-2} f'(i)^{-1}$. Now define

$$\beta_{i,0}(u) = \bigcup_{j \in [s_i(u)] \setminus T_i(u)} \beta_{i,j}(u).$$

We claim $|\beta_{i,0}(u)| \leq \epsilon^2 |\beta_i(u)|$. If $[s_i(u)] \setminus T_i(u) = \emptyset$, then this is obvious. Otherwise, consider $j_0 = \min([s_i(u)] \setminus T_i(u))$. Then for all $a \in V$, either $\beta_{j_0,0}(u)(a) = \emptyset$, or

$$|\beta_{i,j_0}(u)(a)| = |V_{i,j_0}^a(u)| = f'(i) |Z_{i,j_0}^a(u)|.$$

Therefore if $\zeta_{i,j_0}(u) := \{(a, b) : b \in Z_{i,j_0}^a(u)\}$, then $|\beta_{i,j_0}(u)| = f'(i) |\zeta_{i,j_0}(u)| \leq \epsilon^2 f'(i) |\beta_i(u)|$, where the last inequality is since $j_0 \notin T_i(u)$. Consequently, $|\zeta_{i,j_0}(u)| \leq \epsilon^2 |\beta_i(u)|$. By construction and the minimality of j_0 , for all $a \in V$ and $j \in [s_i(u)] \setminus T_i(u)$,

$$V_{i,j}^a(u) \subseteq V_{i,j_0}^a(u) \cup Z_{i,j_0}^a(u).$$

This implies $\beta_{i,0}(u) \subseteq \beta_{i,j_0}(u) \cup \zeta_{i,j_0}$, and thus $|\beta_{i,0}(u)| \leq 2\epsilon^2 |\beta_i(u)|$. We now define $\beta_0(u) = \beta_{i_1}(u) \cup (\bigcup_{i \in J(u)} \beta_j(u)) \cup \bigcup_{i \in I(u)} \beta_{i,0}$. Then by above,

$$|\beta_0(u)| \leq \epsilon^4 |\alpha_u| + \epsilon^4 |\alpha_u| + \sum_{i \in I(u)} |V_{i,0}| \leq 2\epsilon^4 |\alpha_u| + \sum_{i \in I(u)} 2\epsilon^2 |\beta_i(u)| \leq 3\epsilon^2 |\alpha_u|.$$

Recall that by (ii), $\zeta_{i_1}(u) := \alpha_u \setminus (\bigcup_{v=1}^{i_1} \beta_v(u))$ has the property that for all $a \in V$, $\zeta_{i_1}(u)(a)$ is $f'(i_1)$ -good in H_a . If $|\zeta_{i_1}(u)| < \epsilon^2 |\alpha_u|$, set $\beta'_0(u) = \beta_0(u) \cup \zeta_{i_1}(u)$ and $\beta_{s+1}(u) = \emptyset$. If $|\zeta_{i_1}(u)| \geq \epsilon^2 |\alpha_u|$, set $\beta'_0 = \beta_0$ and $\beta_{s+1}(u) = \zeta_{i_1}(u)$. In either case, $|\beta'_0(u)| \leq 5\epsilon^2 |\alpha_u|$, and we now have a partition

$$\alpha_u^* = \beta'_0(u) \cup \beta_{s+1}(u) \cup \bigcup_{i \in I(u)} \bigcup_{j \in T_i(u)} \beta_{i,j}(u).$$

Set $S(u) := \sum_{i \in I(u)} t_i(u)$. Since each $t_i(u) \leq (\epsilon^2 f'(i))^{-1}$, $S(u) \leq \sum_{i=1}^{i_1-1} \epsilon^{-2} f'(i)^{-1} = P_{i_1-1}$. Let $\gamma_1(u), \dots, \gamma_{S(u)}(u)$ be an enumeration of $\{\beta_{i,j}(u) : i \in I(u), j \in T_i(u)\}$.

For each $u \in [m]$, define

$$\alpha'_u = \alpha_u \cap ((V \setminus V_u^*) \times V),$$

and set $\alpha' = \bigcup_{u \in [m]} \bigcup_{i \in [S(u)]} \gamma_i(u)$. Then we have the following partition of α' .

$$\mathcal{P} = \{\gamma_i(u) : u \in [m], i \in [S(u)]\} \cup \{\alpha'_1, \dots, \alpha'_m\}$$

By construction, for every $a \in V$ and $\gamma \in \mathcal{P}$, $\gamma(a)$ has rank at most $d-1$ in H_a . By the induction hypothesis, there are $\Omega_1 \subseteq \mathcal{P}$ and $m' \leq M_{i_1} = M(d-1, g_{i_1}, \epsilon^8)$ such that the following hold.

- $|\bigcup_{\gamma \in \Omega_1} \gamma| \leq \epsilon^8 |V|^2$,
- For each $u \in [m]$ with $\alpha'_u \notin \Omega_1$, there is an integer $c_u \leq m'$, and a partition

$$\alpha'_u = \alpha'_{u,0} \cup \alpha'_{u,1} \cup \dots \cup \alpha'_{u,c_u},$$

so that $|\alpha'_{u,0}| \leq \epsilon^8 |\alpha'_u|$ and for each $v \in [c_u]$, and $a \in V$, $\alpha'_{u,v}(a)$ is $g_{i_1}(m')$ -good in H_a ,

- For each $u \in [m]$ and $i \in [S(u)]$ with $\gamma_i(u) \notin \Omega_1$, there is an integer $b_i(u) \leq m'$ and a partition

$$\gamma_i(u) = \gamma_{i,0}(u) \cup \gamma_{i,1}(u) \cup \dots \cup \gamma_{i,b_i(u)}(u),$$

so that $|\gamma_{i,0}(u)| \leq \epsilon^8 |\gamma_i(u)|$, and for each $v \in [b_i(u)]$ and $a \in V$, $\gamma_{i,v}(u)(a)$ is $g_{i_1}(m')$ -good in H_a .

Let $\Omega_2 = \{u \in [m] : |\alpha_u \cap (\bigcup_{\gamma \in \Omega_1} \gamma)| \geq \epsilon^4 |\alpha_u|\}$. Observe that

$$\epsilon^4 \sum_{u \in \Omega_2} |\alpha_u| \leq |\bigcup_{\gamma \in \Omega_1} \gamma| \leq \epsilon^8 |V|^2.$$

Consequently, $|\sum_{u \in \Omega_2} \alpha_u| \leq \epsilon^4 |V|^2$. For each $u \in [m] \setminus \Omega_2$, define

$$\Omega_1(u) = (\{\gamma_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \cup \{\alpha'_{u,j} : j \in [c_u]\}) \cap \Omega_1 \text{ and}$$

$$\mathcal{P}(u) = (\{\gamma_{i,v}(u) : i \in [S(u)], v \in [b_i(u)]\} \cup \{\alpha'_{u,j} : j \in [c_u]\}) \setminus \Omega_1.$$

Set $\tau_{u,0} = \beta'_0(u) \cup \bigcup_{\gamma \in \Omega_1(u)} \gamma$. If $\beta_{s+1}(u) = \emptyset$, set $d_u := |\mathcal{P}(u)|$, and let $\tau_{u,1}, \dots, \tau_{u,d_u}$ be an enumeration of $\mathcal{P}(u)$. If $\beta_{s+1}(u) \neq \emptyset$, set $d_u := 1 + |\mathcal{P}(u)|$, and let $\tau_{u,1}, \dots, \tau_{u,d_u}$ enumerate $\{V_{s+1}(u)\} \cup \mathcal{P}(u)$. Since $u \notin \Omega_2$,

$$|\tau_{u,0}| \leq 5\epsilon^2 |\alpha_u| + \epsilon^4 |\alpha_u| \leq 6\epsilon^2 |\alpha_u|,$$

and for all $1 \leq v \leq d_u$, and $a \in V$, $\tau_{u,v}(a)$ is $\max\{f'(i_1), g_{i_1}(m')\}$ -good in H_a . Note

$$d_u \leq S(u)m' + 1 \leq P_{i_1-1}m' + 1 \leq m'(P_{i_1-1} + 1) \leq M_{i_1}(P_{i_1-1} + 1).$$

We have now constructed, for all $u \in [m] \setminus \Omega_2$, a partition

$$\alpha_u = \tau_{u,0} \cup \tau_{u,1} \cup \dots \cup \tau_{u,d_u},$$

where $|\tau_{u,0}| \leq 6\epsilon^2 |\alpha_u| < \epsilon |\alpha_u|$, and for all $1 \leq v \leq d_u$, $\tau_{u,v}$ is μ^* -good, where $\mu^* := \max\{g_{i_1}(m'), f'(i_1)\}$. Since by above, $d_u \leq M_{i_1}(P_{i_1-1} + 1) \leq M$, and by construction, $\mu^* \leq f(M_{i_1}(P_{i_1-1} + 1))$, this finishes the proof. \square

We now use Lemma 5.16 to prove that any weakly k -stable property admits binary vdisc_3 -error.

Corollary 5.17. *If \mathcal{H} is weakly k -stable then it admits binary vdisc_3 -error.*

Proof. Fix $k \geq 1$, $\epsilon > 0$, and $t_0 \geq 1$. Let $d = d(k)$ as in part (1) of Theorem 5.12. Choose $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ as in Proposition 5.1 for ϵ and k . Choose $\epsilon'_1 \ll \epsilon'_1 \ll \epsilon, \epsilon_1, k^{-1}$, and define $\epsilon'_2, \epsilon''_2 : \mathbb{N} \rightarrow (0, 1]$ so that for all x , $\epsilon''_2(x) \ll \epsilon'_2(x) \ll \epsilon_2(x)$. Then let f and g be as in Lemma 3.27 for ϵ'_1, ϵ'_2 . Now define $\psi : \mathbb{N} \rightarrow (0, 1]$ so that $\psi(x) \ll \epsilon''_1 f(t_0 x^{2^{x+1}}, 2^{s+1}, t_0 x^{2^{x+1}}, 2^{x+1})^{-1} g(t_0 x^{2^{x+1}}, 2^{s+1}, t_0 x^{2^{x+1}}, 2^{x+1})^{-1} 2^{(x+1)^2} x^2$, for all x .

Apply Lemma 5.16 to the function ψ , the rank bound d , and ϵ'_1 to obtain M . Then choose $T \gg (\epsilon'_1)^{-1} t_0 \epsilon^{-1} M$ and $N \gg T$.

Suppose $H = ([n], E)$ is a weakly k -stable 3-graph with $n \geq N$. Apply Lemma 5.16 to $\alpha = V \times V$ to find $s \leq M$ and a partition $V \times V = \alpha_0 \cup \dots \cup \alpha_s$ so that $|\alpha_0| \leq \epsilon''_1 n^2$ and such that for all $1 \leq i \leq s$ and $a \in V$, $\alpha_i(a)$ is $\psi(s)$ -good in H_a . Observe that by Lemma 5.9, we know that for all $a \in V$ and $1 \leq i, j \leq s$, there is some $\delta = \delta(i, j, a)$ so that

$$(4) \quad |E_a^\delta \cap K_2[\alpha_i(a), \alpha_j(a)]| \geq (1 - 2\sqrt{\psi(s)}) |K_2[\alpha_i(a), \alpha_j(a)]|.$$

Let $\mathcal{P} = \{V_1, \dots, V_{t_1}\}$ be a minimal partition of V so that if a, a' are in the same part of \mathcal{P} , then $\delta(i, j, a) = \delta(i, j, a')$ for all $1 \leq i, j \leq s$. Note $t_1 \leq 2^{s^2}$.

We now adjust \mathcal{P} to obtain an equipartition. Set $t = (t_0 t_1 \epsilon'_1)^{-1}$, and $K = \lfloor n/t \rfloor$. For each $1 \leq i \leq t_1$, choose a partition $V_i = V_{i,0} \cup V_{i,1} \cup \dots \cup V_{i,r_i}$ so that $|V_{i,0}| < K$ and for each $1 \leq j \leq r_i$, $|V_{i,j}| = K$. Define $V_0 = \bigcup_{i=1}^{t_1} V_{i,0}$ and $V' = V \setminus V_0$. By construction, we have

$$|V_0| \leq \frac{t_1}{t} n \leq \epsilon'_1 n.$$

Choose an equipartition $V_0 = \bigcup_{i=1}^{t_1} \bigcup_{j=1}^{r_i} V_{i,j}^0$, and for each $1 \leq i \leq t_1$ and $1 \leq j \leq r_i$, set $V'_{i,j} = V_{i,j} \cup V_{0,i}$. Clearly $|V'_{i,j} \setminus V_{i,j}| \leq \epsilon''_1 |V'_{i,j}|$. We now define $\mathcal{Q}_{\text{vert}} = \{V'_{i,j} : i \in [t_1], j \in [r_i]\}$ and $t_2 := |\mathcal{Q}_{\text{vert}}|$. Note $t_2 \leq t$. Our next goal is to define a partition of $K_2[X, Y]$ for all $XY \in \binom{\mathcal{Q}_{\text{vert}}}{2}$.

Given $XY \in \binom{\mathcal{Q}_{\text{vert}}}{2}$, we know that XY has the form $V'_{i,j} V'_{i',j'}$ for some $i, i' \in [t_1]$, $j \in [r_i]$, and $j' \in [r_{i'}]$. If $1 \leq i < i' \leq t_1$, then for each $1 \leq j \leq r_i$, $1 \leq j' \leq r_{i'}$, and $0 \leq u \leq s$, define

$$F_u[V'_{i,j}, V'_{i',j'}] = \{vv' \in K_2[V'_{i,j}, V'_{i',j'}] : v' \in \alpha_u(v)\}.$$

Similarly, if $1 \leq i = i' \leq t_1$, then for each $1 \leq j < j' \leq r_i$ and $0 \leq u \leq s$, define

$$F_u[V'_{i,j}, V'_{i,j'}] = \{vv' \in K_2[V'_{i,j}, V'_{i,j'}] : v' \in \alpha_u(v)\}.$$

Now set $\mathcal{Q}_{\text{edge}} = \{F_u[X, Y] : XY \in \binom{\mathcal{Q}_{\text{vert}}}{2}, 0 \leq u \leq s\}$. Then $\mathcal{Q} := (\mathcal{Q}_{\text{vert}}, \mathcal{Q}_{\text{edge}})$ is a $(t_2, s+1)$ -decomposition of V . Given $X, Y, Z \in \binom{\mathcal{Q}_{\text{vert}}}{2}$ and $0 \leq u, v, w \leq s$, let

$$\mathcal{Q}_{XYZ}^{u,v,w} = (X \cup Y \cup Z, F_u[X, Y] \cup F_v[X, Z] \cup F_w[Y, Z]).$$

We claim that for all $XYZ \in \binom{\mathcal{Q}_{vert}}{3}$ and $1 \leq u, v, w \leq s$, there is a $\delta(X, Y, Z, u, v, w) \in \{0, 1\}$ so that

$$|E \cap K_3^{(2)}(Q_{XYZ}^{u,v,w})| \geq (1 - \psi(s)^{1/4})|K_3^{(2)}(Q_{XYZ}^{u,v,w})|.$$

Fix $XYZ \in \binom{\mathcal{Q}_{vert}}{3}$. By construction, there $1 \leq i_1, i_2, i_3 \leq t_2$, $j_1 \in [r_{i_1}]$, $j_2 \in [r_{i_2}]$, and $j_3 \in [r_{i_3}]$ such that $X = V'_{i_1, j_1}$, $Y = V'_{i_2, j_2}$, and $Z = V'_{i_3, j_3}$.

Suppose first $i_1 < i_2, i_3$. Then for all $xy \in F_u[X, Y]_u$, $y \in \alpha_u(x)$ and for all $xz \in F_w[X, Z]$, $z \in \alpha_w(x)$. By (4), there is $\delta = \delta(u, w, x) \in \{0, 1\}$ so that

$$|N_H^\delta(x) \cap K_2[\alpha_u(x), \alpha_w(x)]| \geq (1 - 2\sqrt{\psi(s)})|\alpha_u(x)||\alpha_w(x)|.$$

Let $G' = (X \cup Y \cup Z, \{xyz \in K_3[X, Y, Z] : xy \in F_u[X, Y], xz \in F_w[X, Z]\})$. Then $|E^\delta \cap K_3^{(2)}(Q_{XYZ}^{u,v,w})|$ is at least the following.

$$\begin{aligned} |K_3^{(2)}(Q_{XYZ}^{u,v,w})| - \sum_{x \in X} 2\sqrt{\psi(s)}|\alpha_u(x)||\alpha_w(x)| &\geq |K_3^{(2)}(Q_{XYZ}^{u,v,w})|(1 - 2\sqrt{\psi(s)}t^2) \\ &\geq |K_3^{(2)}(Q_{XYZ}^{u,v,w})|(1 - (\psi(s))^{1/4}), \end{aligned}$$

where the first inequality is since $|\alpha_u(x)||\alpha_w(x)| \leq t^2|Y||Z|$, and the second inequality is by the definition of t and our choice of ψ .

A symmetric argument works if $i_2 < i_1, i_3$ or $i_3 < i_1, i_2$. Suppose now $i_1 = i_2 < i_3$. Since X and Y are distinct, we may assume after relabeling that $j_1 < j_2$. Then we have that for all $xy \in F_u[X, Y]$, $y \in \alpha_u(x)$ and for all $xz \in F_w[X, Z]$, $z \in \alpha_w(x)$. We now finish this case as above, with $\delta = \delta(u, w, x)$. A symmetric argument works if $i_1 = i_3 < i_2$ or $i_2 = i_3 < i_1$. So we are now left with the case where $i_1 = i_2 = i_3$. Since X, Y, Z are pairwise distinct, we must have the j_1, j_2, j_3 are pairwise distinct. By relabeling if necessary, we may assume $j_1 < j_2, j_3$. Then we know that for all $xy \in F_u[X, Y]_u$, $y \in \alpha_u(x)$ and for all $xz \in F_w[X, Z]$, $z \in \alpha_w(x)$. We can thus finish this case as above with $\delta = \delta(u, w, x)$.

We now apply Lemma 3.27 to \mathcal{Q} and \mathcal{P} , where \mathcal{P} is the trivial decomposition of V consisting of $\mathcal{P}_{vert} = \{V\}$ and $\mathcal{P}_{edge} = \{K_2[V, V]\}$. Then we obtain $\ell' \leq f(1, 1, t_2, s + 1)$ and $t' \leq g(1, 1, t_2, s + 1)$, and an $(t', \ell', \epsilon_1', \epsilon_2'(\ell'))$ -decomposition \mathcal{R} of V in which every element of \mathcal{R}_{edge} satisfies $\text{disc}_2(\epsilon_2'(\ell'))$ and which is an $(\epsilon_1', \epsilon_2'(\ell'))$ -approximate refinement of both \mathcal{Q} and \mathcal{P} . Let Σ_1 be as in the definition of \mathcal{R} being an $(\epsilon_1', \epsilon_2'(\ell'))$ -approximate refinement of \mathcal{Q} . Then for all $XY \in \binom{\mathcal{R}_{vert}}{2} \setminus \Sigma_1$, there is $\phi(R_{XY}^\alpha) \in \mathcal{Q}_{edge}$ be so that $|R_{XY}^\alpha \setminus \phi(R_{XY}^\alpha)| \leq \epsilon_1'|R_{XY}^\alpha|$, and so that $R_{XY}^\alpha \setminus \phi(R_{XY}^\alpha)$ has $\text{disc}_2(\epsilon_2'(\ell'))$. Now define

$$\Sigma_2 = \{XY \in \binom{\mathcal{R}_{vert}}{2} : F_0[X, Y] \geq \sqrt{\epsilon_1''}|X||Y|\}.$$

Since $|\alpha_0| \leq \epsilon_1'n^2$ and \mathcal{R}_{vert} is an equipartition of V , we know that $|\Sigma_2| \leq \sqrt{\epsilon_1''}(t')^2$. We now set $\Sigma = \Sigma_1 \cup \Sigma_2$. Note

$$|\Sigma| \leq \epsilon_1'(t')^2 + \sqrt{\epsilon_1''}(t')^2 \leq \epsilon_1 t^2.$$

Now suppose we have $XYZ \in \mathcal{R}_{vert}$ satisfies $XY, YZ, XZ \notin \Sigma$. We claim that XYZ is ϵ'_1 -homogeneous with respect to H . Define

$$\Omega(XY) = \{\alpha \in [\ell'] : \phi(R_{XY}^\alpha) \subseteq F_u[X', Y'] \text{ for some } u \neq 0 \text{ and } X'Y' \in \binom{\mathcal{Q}_{vert}}{2}\}.$$

Since $XY \notin \Sigma_2$, we can deduce that $|\Omega(XY)| \geq (1 - (\epsilon'_1)'^{1/4})\ell'$. We can then define $\Omega(XZ), \Omega(YZ)$ the same way and deduce that $|\Omega(YZ)|, |\Omega(XZ)| \geq (1 - (\epsilon'_1)'^{1/4})\ell'$. Now fix some $\alpha \in \Omega(XY), \beta \in \Omega(XZ), \gamma \in \Omega(YZ)$. Let $X', Y', Z' \in \mathcal{Q}_{vert}$ and $1 \leq u, v, w \leq s$ be such that $\phi(R_{XY}^{\alpha, \beta, \gamma}) = Q_{X'Y'Z'}^{u, v, w}$. By above, there is some $\delta = \delta(X', Y', Z', u, v, w)$ so that

$$|E^\delta \cap K_3^{(2)}(Q_{X'Y'Z'}^{u, v, w})| \geq (1 - (\psi(s))^{1/4})|K_3^{(2)}(Q_{X'Y'Z'}^{u, v, w})|.$$

Let $A = R_{XY}^\alpha \setminus \phi(R_{XY}^\alpha)$, $B = R_{XZ}^\beta \setminus \phi(R_{XZ}^\beta)$ and $C = R_{YZ}^\gamma \setminus \phi(R_{YZ}^\gamma)$. Then each of A, B, C satisfy $\text{disc}_2(\epsilon'_2(\ell'))$ and

$$|A| \leq (\epsilon'_1)^{1/4}|R_{XY}^\alpha|, |B| \leq (\epsilon'_1)^{1/4}|R_{XZ}^\beta|, \text{ and } |C| \leq (\epsilon'_1)^{1/4}|R_{YZ}^\gamma|.$$

Let $G_A = (X \cup Y \cup Z, A \cup R_{YZ}^\gamma \cup R_{XZ}^\gamma)$, $G_B = (X \cup Y \cup Z, R_{XY}^\alpha \cup B \cup R_{XZ}^\gamma)$, and $G_C = (X \cup Y \cup Z, R_{XY}^\alpha \cup R_{YZ}^\gamma \cup C)$. Then $|E^\delta \cap K_3^{(2)}(R_{XYZ}^{\alpha, \beta, \gamma})|$ is at least the following.

$$\begin{aligned} & |K_3^{(2)}(R_{XYZ}^{\alpha, \beta, \gamma}) \setminus K_3^{(2)}(Q_{X'Y'Z'}^{u, v, w})| - (\psi(s))^{1/4}|K_3^{(2)}(Q_{X'Y'Z'}^{u, v, w})| \\ &= |K_3^{(2)}(R_{XYZ}^{\alpha, \beta, \gamma})| - |K_3^{(2)}(G_A)| - |K_3^{(2)}(G_B)| - |K_3^{(2)}(G_C)| - (\psi(s))^{1/4}|K_3^{(2)}(Q_{X'Y'Z'}^{u, v, w})| \\ &\geq |K_3^{(2)}(R_{XYZ}^{\alpha, \beta, \gamma})|(1 - 3(\epsilon'_1)^{1/4} - \frac{(t')^2}{t^2}\psi(s)^{1/4}) \\ &\geq |K_3^{(2)}(R_{XYZ}^{\alpha, \beta, \gamma})|(1 - (\epsilon'_1)^{1/8}), \end{aligned}$$

where the second inequality uses Corollary 3.6, and the last uses the definition of ψ . Thus, we have that for every $\alpha \in \Omega(XY), \beta \in \Omega(XZ), \gamma \in \Omega(YZ)$, $R_{XYZ}^{\alpha, \beta, \gamma}$ is ϵ'_1 -homogeneous with respect to H . Combining this with Proposition 5.1, the size bounds on $\Omega(XY), \Omega(YZ), \Omega(XZ)$ and since $\epsilon'_1 \ll \epsilon_1$, we find that XYZ is in fact ϵ_1 -homogeneous with respect to H . Consequently, XYZ satisfies $\text{vdisc}_3(\epsilon_1)$ with respect to H by Proposition 3.21. Since $|\Sigma| \leq \epsilon_1 t^2$, this shows \mathcal{P} is $\text{vdisc}_3(\epsilon_1)$ -homogeneous with respect to H with binary error, as desired. \square

Proposition 2.43 follows immediately from this and Proposition 2.19.

6. THE FUNCTIONAL ORDER PROPERTY AND LINEAR $\text{disc}_{2,3}$ -ERROR

In this section we prove Theorem 2.50, which says that a hereditary 3-graph property admits linear $\text{disc}_{2,3}$ -error if and only if it is NFOP_2 . We will begin in Section 6.1 by giving equivalent conditions for when a hereditary 3-graph property has FOP_2 or IP_2 . This equivalent formulation will use certain auxiliary edge-colored graphs associated to regular decompositions of 3-graphs. In particular, we will show a correspondence between when a hereditary 3-graph property has IP_2 or FOP_2 with the existence of IP or OP in these auxiliary edge colored graphs, respectively. In Section 6.2, we prove Theorem 6.13, which is a strong removal lemma for graphs containing few d -trees. In Section 6.3, we then prove

our main result of the section, which says that a hereditary 3-graph property admits linear $\text{disc}_{2,3}$ -error if and only if it is NFOP_2 . This proof will leverage the recharacterization from Section 6.1, Theorem 5.15 of the previous section, and the removal lemma, Theorem 6.13.

6.1. Encodings and recharacterizations of FOP_2 . In this subsection we will give additional characterizations of when a hereditary property is NFOP_2 . These recharacterizations will be used later in our proof of Theorem 2.50. Along the way, we will include some related results which demonstrate that a strong analogy FOP_2 plays an analogous role compared to stability, as VC_2 -dimension plays compared to VC-dimension.

The main recharacterization will be presented in terms of auxiliary graphs defined from regular triads in decompositions of 3-graphs.

Definition 6.1. Suppose $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow \mathbb{N}$, $\ell, t \geq 1$, V is a set, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition for V consisting of $\{V_i : i \in [t]\}$ and $\{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Define

$$\begin{aligned} \mathcal{P}_{\text{cnr}} &= \{P_{ij}^\alpha P_{ik}^\beta : ijk \in \binom{[t]}{3}, \alpha, \beta \leq \ell, \text{ and } P_{ij}^\alpha, P_{ij}^\beta \text{ satisfy } \text{disc}_2(\epsilon_2(\ell), 1/\ell)\} \text{ and} \\ \mathcal{P}_{\text{edge}}^* &= \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell, \text{ and } P_{ij}^\alpha \text{ satisfies } \text{disc}_2(\epsilon_2(\ell), 1/\ell)\}. \end{aligned}$$

In the above, cnr stands for “corner.” The motivation for this definition is that given an element $P_{ij}^\alpha \in \mathcal{P}_{\text{edge}}$ and an element $P_{uv}^\beta P_{uw}^\gamma \in \mathcal{P}_{\text{cnr}}$, if $\{v, w\} = \{i, j\}$, the pair $(P_{ij}^\alpha, P_{uv}^\beta P_{uw}^\gamma)$ corresponds to a triad from \mathcal{P} , in which case there is a natural way to define a reduced edge, or a reduced non-edge, between P_{ij}^α and $P_{uv}^\beta P_{uw}^\gamma$. We will now define a bipartite edge colored graph associated to a regular decomposition of a 3-graph, where a bipartite edge-colored graph is a tuple of the form $(A \cup B, E_1, \dots, E_t)$ such that $K_2[A, B] = E_1 \cup \dots \cup E_t$.

Definition 6.2. Suppose $H = (V, E)$ is a 3-graph, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition for V . Given $G_{ijs}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$, let $H_{ijs}^{\alpha, \beta, \gamma} = H|G_{ijs}^{\alpha, \beta, \gamma}$ and $d_{ijs}^{\alpha, \beta, \gamma} = |E \cap K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})| / |K_3^{(2)}(G_{ijk}^{\alpha\beta\gamma})|$. Then define

$$\begin{aligned} \mathbf{E}_1 &= \{P_{ij}^\alpha (P_{jk}^\beta P_{ik}^\gamma) \in K_2[\mathcal{P}_{\text{edge}}^*, \mathcal{P}_{\text{cnr}}] : (H_{ijs}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \gamma}) \text{ has } \text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell)) \text{ and } d_{ijs}^{\alpha, \beta, \gamma} \geq 1/2\}, \\ \mathbf{E}_0 &= \{P_{ij}^\alpha (P_{jk}^\beta P_{ik}^\gamma) \in K_2[\mathcal{P}_{\text{edge}}^*, \mathcal{P}_{\text{cnr}}] : (H_{ijs}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \gamma}) \text{ has } \text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell)) \text{ and } d_{ijs}^{\alpha, \beta, \gamma} < 1/2\}, \text{ and} \\ \mathbf{E}_2 &= K_2[\mathcal{P}_{\text{edge}}^*, \mathcal{P}_{\text{cnr}}] \setminus (\mathbf{E}_1 \cup \mathbf{E}_0), \end{aligned}$$

Note that $\mathbf{E}_0, \mathbf{E}_1, \mathbf{E}_2$ each depend on \mathcal{P} and H , but we suppress these in the notation as they should be clear from context. Note Definition 6.2 associates to a decomposition \mathcal{P} of a 3-graph H , an auxiliary edge-colored bipartite graph, $(\mathcal{P}_{\text{edge}}^* \cup \mathcal{P}_{\text{cnr}}, \mathbf{E}_1, \mathbf{E}_0, \mathbf{E}_2)$. When the decomposition \mathcal{P} is $\text{disc}_{2,3}$ -homogeneous, elements in \mathbf{E}_1 will correspond to triads of density near 1, and elements in \mathbf{E}_0 will correspond to triads of density near 0. In that case, Definition 6.2 becomes an especially useful reflection of the structure in the original 3-graph. This will be one of the main ideas behind our proofs in Section 6.3. We will also consider special copies of certain bipartite graphs made using \mathbf{E}_1 and \mathbf{E}_0 .

Definition 6.3. Suppose $R = (A \cup B, E_R)$ is a bipartite graph, $H = (V, E)$ is a 3-graph, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2)$ -decomposition of V . An (A, B) -encoding of R in (H, \mathcal{P}) consists of

a pair of functions (g, f) where $g : A \rightarrow \mathcal{P}_{cnr}$ and $f : B \rightarrow \mathcal{P}_{edge}^*$ are such that the following hold for some $j_0 k_0 \in \binom{[t]}{2}$.

- (1) $\text{Im}(f) \subseteq \{P_{j_0 k_0}^\alpha : \alpha \in [\ell]\}$, and $\text{Im}(g) \subseteq \{P_{ij_0}^\beta P_{ik_0}^\gamma : i \in [t], \beta, \gamma \in [\ell]\}$, and
- (2) For all $a \in A$ and $b \in B$, if $ab \in E_R$, then $g(a)f(b) \in \mathbf{E}_1$, and if $ab \notin E_R$, then $g(a)f(b) \in \mathbf{E}_0$.

We say $(V_{j_0}, V_{k_0}, \bigcup_{i=u}^k V_{i_u})$ is the *partition corresponding to* (g, f) .

Note that there is an inherent asymmetry in Definition 6.3, since A is mapped into \mathcal{P}_{cnr} while B is mapped into \mathcal{P}_{edge}^* . We will be interested in the cases where $R = H(k)$ and $R = U(k)$, so it will be convenient to set notation for the labeling of their parts. Given $k \geq 1$, we set $U(k) = (A_k \cup C_{\mathcal{P}([k])}, E)$, where $A_k = \{a_i : i \in [k]\}$, $C_{\mathcal{P}([k])} = \{c_S : S \subseteq [k]\}$, and $E = \{a_i c_S : i \in S\}$. Similarly, we set $H(k) = (A_k \cup B_k, E)$ where $A_k = \{a_i : i \in [k]\}$, $B_k = \{b_i : i \in [k]\}$, and $E = \{a_i b_j : i \leq j\}$. To ease notation, an *encoding of* $H(k)$ will always mean an (A_k, B_k) -encoding of $H(k)$, and an *encoding of* $U(k)$ will always mean an $(A_k, C_{\mathcal{P}([k])})$ -encoding of $U(k)$.

We now make a definition for when a hereditary 3-graph property has many encodings of a given bipartite graph R .

Definition 6.4. Suppose $R = (A \cup B, E)$ is a bipartite graph and \mathcal{H} is a hereditary 3-graph property. We say that R is (A, B) -encoded in \mathcal{H} if for all $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are T, L such that for all N , the following holds.

There exist a 3-graph $H = (V, E)$ with $|V| \geq n$, some $t \leq T, \ell \leq L$, and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V , such that there exists an (A, B) -encoding of R in (H, \mathcal{P}) .

We will simply say that $H(k)$ is *encoded in* \mathcal{H} if it is (A_k, B_k) -encoded in \mathcal{H} , and $U(k)$ is *encoded in* \mathcal{H} if it is $(A_k, C_{\mathcal{P}([k])})$ -encoded in \mathcal{H} . We now draw a connection between encodings of $H(k)$ and $U(k)$, and k -FOP₂ and k -IP₂, respectively. Analogues of these results also appear in the algebraic setting in [60].

Theorem 6.5. Suppose \mathcal{H} is a hereditary 3-graph property.

- (1) If $H(k)$ is encoded in \mathcal{H} , then \mathcal{H} has k -FOP₂.
- (2) If $U(k)$ is encoded in \mathcal{H} , then \mathcal{H} has k -IP₂.

The proof of Theorem 6.5 will rely on two facts, Propositions 6.6 and 6.7.

Proposition 6.6. For all $k \geq 1$, there exist $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ such that for all $t, \ell \geq 1$, there is N such that the following hold. Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . If there exists an encoding of $U(k)$ in (H, \mathcal{P}) , then H has k -IP₂.

Proof. Fix $k \geq 1$. Choose δ_3 as in Theorem 4.1 for $m = 2k + 2^{k^2}$, $\xi = 1/2$, and $d_3 = 1/2$. Set $\epsilon_1 = \delta_3^2$, and let $\epsilon_2(x)$ be the function sending x to $\delta_2(x^{-1})$ from Theorem 4.1, applied with δ_3, m, ξ, d_3 . Given $t, \ell \geq 1$, and choose $N \gg t, \ell, \epsilon_1^{-1}, \epsilon_2(\ell)^{-1}$.

Suppose $H = (V, E)$ with $|V| = n \geq N$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . Say $\mathcal{P}_{vert} = \{V_i : i \in [t]\}$ and $\mathcal{P}_{edge} = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Assume there exists an

encoding of $U(k)$ in (H, \mathcal{P}) . Then there are $1 \leq j_0 \neq k_0 \leq t$, some $\{\alpha_S \in [\ell] : S \subseteq [k]\}$, $\{i_u \in [t] : u \in [k]\}$, and $\{(\beta_u, \gamma_u) \in [\ell]^2 : u \in [k]\}$ such that for each $u \in [k]$ and $S \subseteq [k]$, each of $P_{i_u j_0}^{\gamma_u}, P_{i_u k_0}^{\beta_u}, P_{j_0 k_0}^{\alpha_S}$ satisfy $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$ and if $u \in S$ then $(H_{i_u j_0 k_0}^{\alpha_S, \beta_u, \gamma_u}, G_{i_u j_0 k_0}^{\alpha_S, \beta_u, \gamma_u})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ and density at least $1/2$ and if $u \in S$, and density less than $1/2$ if $u \notin S$.

Given $X \subseteq [k]^2$ and $i \in [k]$, let $N_X(i) = \{j \in [k] : (i, j) \in X\}$. For each $u \in [k]$, set $A_u = V_{i_u}$ and $B_u = V_{j_0}$. Then for each $X \subseteq [k]$, let $C_X = V_{k_0}$. Given $X \subseteq [k]^2$ and $u \in [k]$, define $P_{C_X B_u} = P_{j_0 k_0}^{\alpha_{N_X(u)}}$ and $P_{C_X A_u} = P_{i_u k_0}^{\beta_u}$. For $(u, v) \in [k]^2$, define $P_{A_v B_u} = P_{i_u j_0}^{\gamma_u}$.

Given $(u, v) \in [k]^2$ and $X \subseteq [k]^2$, set $G_{uvX} := (A_v \cup B_u \cup C_X, P_{A_v C_X} \cup P_{B_u C_X} \cup P_{A_v B_u})$ (in other words, $G_{uvX} = G_{i_v j_0 k_0}^{\alpha_{N_X(u)}, \beta_v, \gamma_v}$). Then define $H_{uvX} = (A_v \cup B_u \cup C_X, E \cap K_3^{(2)}(G_{vuX}))$. By assumption, (H_{uvX}, G_{uvX}) has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ and density at least $1/2$ if $v \in N_X(u)$ (i.e. if $(u, v) \in X$) and less than $1/2$ if $v \notin N_X(u)$ (i.e. $(u, v) \notin X$). By Theorem 4.1, there are at least $(1/\ell)^{\binom{m}{2}} (1/2)^{\binom{m}{3}} (n/t)^m > 0$ many tuples $(a_u)_{u \in [k]} (b_v)_{v \in [k]} (c_X)_{X \subseteq [k]^2}$ such that $a_u b_v c_X \in E$ if and only if $(u, v) \in X$. This finishes the proof. \square

Note that in Proposition 6.6, we did not just show that H has $k\text{-IP}_2$, but rather the stronger statement that $\text{Trip}(H)$ contains many copies of $V(k)$. We now prove the analogous result for FOP_2 , where $U(k)$ is replaced by $H(k)$ and IP_2 is replaced by FOP_2 . We will give a more detailed conclusion than we did in Proposition 6.6, as we will need these details later on in the section.

Proposition 6.7. *For all $k \geq 1$, there exist $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ such that for all $t, \ell \geq 1$, there is N such that the following hold. Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . Suppose (g, f) is an encoding of $H(k)$ in (H, \mathcal{P}) , with corresponding partition $(V_{j_0}, V_{k_0}, \bigcup_{i=w}^k V_{i_w})$. Then the following hold.*

- (1) *There exist $(c_w)_{w \in [k]} \in \prod_{w=1}^k V_{i_w}$, $\{a_u^f : u \in [k]\} \subseteq V_{j_0}$, and $\{b_v : v \in [k]\} \subseteq V_{k_0}$, such that $a_u^f b_v c_w \in E$ if and only if $w \leq f(u, v)$.*
- (2) *There exist $(c_w)_{w \in [k]} \in \prod_{w=1}^k V_{i_w}$, $\{a_u^f : u \in [k]\} \subseteq V_{k_0}$, and $\{b_v : v \in [k]\} \subseteq V_{j_0}$ such that $a_u^f b_v c_w \in E$ if and only if $w \leq f(u, v)$.*

Proof. Fix $k \geq 1$. Choose δ_3 as in Theorem 4.1 for $m = 2k + k^{k^2}$, $\xi = 1/2$, and $d_3 = 1/2$. Set $\epsilon_1 = \delta_3^2$. Let $s(x)$ be the function sending x to $\delta_2(x^{-1})$ from Theorem 4.1 applied with δ_3, m, ξ, d_3 . For all x , define $\epsilon_2(x) \ll s(x)$. Given t, ℓ , choose $N \gg t, \ell, \epsilon_1^{-1}, \epsilon_2(\ell)^{-1}$.

Suppose $H = (V, E)$ with $|V| = n \geq N$, and \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . Say \mathcal{P} consists of $\mathcal{P}_{\text{vert}} = \{V_i : i \in [t]\}$ and $\mathcal{P}_{\text{edge}} = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Assume there exists an encoding of $H(k)$ in (H, \mathcal{P}) . Then there are $1 \leq j_0 \neq k_0 \leq t$, some $\{(\alpha_u, \beta_u, \gamma_u) \in [\ell]^3 : u \leq k\}$, and $i_1, \dots, i_k \leq t$ such that for each $u \in [k]$, each of $P_{i_u j_0}^{\alpha_u}, P_{i_u k_0}^{\beta_u}, P_{j_0 k_0}^{\gamma_u}$ satisfy $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$ and if $u \leq v$ then $(H_{j_0 k_0 i_u}^{\gamma_v, \alpha_u, \beta_u}, G_{j_0 k_0 i_u}^{\gamma_v, \alpha_u, \beta_u})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ and density at least $1/2$ and if $u > v$ then $(H_{j_0 k_0 i_u}^{\gamma_v, \alpha_u, \beta_u}, G_{j_0 k_0 i_u}^{\gamma_v, \alpha_u, \beta_u})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ and density less than $1/2$.

Let I be the set of functions $f : [k]^2 \rightarrow [k]$. For each $u \in [k]$, set $C_u = V_{i_u}$, $B_u = V_{k_0}$, and for each $f \in I$, set $A_u^f = V_{j_0}$. Then for each $u, v, w \in [k]$ and $f \in I$, define $Q_{A_u^f C_w} = P_{j_0 i_w}^{\alpha_w}$,

$Q_{B_v C_w} = P_{k_0, i_w}^{\beta_w}$ and $Q_{A_u^f B_v} = P_{j_0 k_0}^{\gamma f(u, v)}$. For each $u, v, w \in [k]$ and $f \in I$, let

$$\begin{aligned} G_{uvw}^f &= (A_u^f \cup B_v \cup C_w, Q_{B_v C_w} \cup Q_{A_u^f C_w} \cup Q_{A_u^f B_v}) \text{ and} \\ H_{uvw}^f &= (A_u^f \cup B_v \cup C_w, E \cap K_3^{(2)}(G_{uvw}^f)). \end{aligned}$$

We know that for each $u, v, w \in [k]$ and $f \in I$, (H_{uvw}^f, G_{uvw}^f) satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$, with density at least $1/2$ if $w \leq f(u, v)$ and less than $1/2$ if $w > f(u, v)$. By Theorem 4.1, there are at least $(1/\ell)^{\binom{m}{2}} (1/2)^{\binom{m}{3}} (n/t)^m > 0$ many tuples $(a_u^f)_{u \in [k], f \in I} (b_v)_{v \in [k]} (c_w)_{w \in [k]} \in \prod_{u \in [k], f \in I} A_u^f \times \prod_{v=1}^k B_v \times \prod_{w=1}^k C_w$ such that $a_u^f b_v c_w \in E$ if and only if $w \leq f(u, v)$. Note that for each such tuple, $c_w \in V_{i_w}$ for each $w \in [k]$, $\{a_u^f : u \in [k], f \in I\} \subseteq V_{j_0}$, and $\{b_v : v \in [k]\} \subseteq V_{k_0}$. Thus (1) holds.

To prove (2), we proceed in the same way, but make the following small adjustments in the third paragraph. For each $v \in [k]$, set $B_v = V_{j_0}$, and for each $u \in [k]$ and $f \in I$, set $A_u^f = V_{k_0}$. As before, define $C_w = V_{i_w}$ for each $w \in [k]$. Then for each $u, v, w \in [k]$ and $f \in I$, define $Q_{A_u^f C_w} = P_{k_0, i_w}^{\alpha_w}$, $Q_{B_v C_w} = P_{j_0, i_w}^{\beta_w}$ and $Q_{A_u^f B_v} = P_{j_0 k_0}^{\gamma f(u, v)}$. One then finishes the proof in the same way to obtain (2). \square

Note that in Proposition 6.7, we in fact found many copies of $F(k)$ in $\text{Trip}(H)$. We are now able to prove Theorem 6.5.

PROOF OF THEOREM 6.5: Suppose that \mathcal{H} encodes $U(k)$. Let $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ be as in Proposition 6.6. Let $T = T(\epsilon_1, \epsilon_2)$ and $L = (\epsilon_1, \epsilon_2)$ be as in the definition of $U(k)$ being encoded in \mathcal{H} . Let $N = N(T, L)$ be as in Proposition 6.6. Since $U(k)$ is encoded in \mathcal{H} there is $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, some $t \leq T$, $\ell \leq L$, a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V , and an encoding of $U(k)$ in (H, \mathcal{P}) . By Proposition 6.6, H has k -IP₂, so \mathcal{H} has k -IP₂.

Suppose now that \mathcal{H} encodes $H(k)$. Let $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ be as in Proposition 6.7. Let $T = T(\epsilon_1, \epsilon_2)$ and $L = (\epsilon_1, \epsilon_2)$ be as in the definition of $H(k)$ being encoded in \mathcal{H} . Let $N = N(T, L)$ be as in Proposition 6.6. Since $H(k)$ is encoded in \mathcal{H} there are $H = (V, E) \in \mathcal{H}$ with $|V| \geq N$, some $t \leq T$, $\ell \leq L$, a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition \mathcal{P} of V , and an encoding of $H(k)$ in (H, \mathcal{P}) . By Proposition 6.7, H has k -FOP₂, so \mathcal{H} has k -FOP₂. \square

Theorem 6.5 showed that if \mathcal{H} encodes $H(k)$ (respectively $U(k)$) for all k , then \mathcal{H} has FOP₂ (respectively IP₂). Our next goal is to show that the converse of these implications is also true, i.e. if \mathcal{H} has FOP₂ (respectively IP₂), then it encodes $H(k)$ (respectively $U(k)$) for all k .

We will first prove this for FOP₂. This will require Lemma 6.8 below, which shows we can construct certain 3-graphs from $F(n)$, when n is large.

Lemma 6.8. *For all $m, k \geq 1$, there is N such that the following holds. Suppose $n \geq N$, $U = \{u_1, \dots, u_m\}$, $W = \{w_1, \dots, w_m\}$ are sets $\bigcup_{\alpha \leq k} P^\alpha$ is a partition of $K_2[U, W]$, and $H = (A \cup B \cup C, E)$ is a clean copy of $F(n)$.*

Then there exist disjoint sets $Z_1, \dots, Z_k \subseteq C$, each of size m , along with $\{a_1, \dots, a_m\} \subseteq A$ and $\{b_1, \dots, b_m\} \subseteq B$ such that the following hold, where $Q^\alpha = \{a_i b_j : u_i w_j \in P^\alpha\}$. For all $c \in Z_\beta$ and $ab \in Q^\alpha$, $abc \in E$ if and only if $\beta \leq \alpha$.

Proof. Fix $m, k \geq 1$. Choose $N \gg m, k$. Suppose $U = \{u_1, \dots, u_m\}$ and $W = \{w_1, \dots, w_m\}$ are sets of size m and $\bigcup_{\alpha \leq k} P^\alpha$ is a partition of $K_2[U, W]$.

Assume $n \geq N$ and $H = (A \cup B \cup C, E)$ is a clean copy of $F(n)$. We may assume that $A = \{a_i^f : i \in [n], f : [n]^2 \rightarrow [n]\}$, $B = \{b_i : i \in [n]\}$, and $C = \{c_i : i \in [n]\}$ are such that $a_i^f b_j c_s \in E$ if and only if $s \leq f(i, j)$.

Define a function $g : [n]^2 \rightarrow [n]$, as follows. For each $(s, t) \in [m]^2$, set $g(s, t) = m\alpha$, where α is such that $u_s w_t \in P_{UW}^\alpha$. For all $(s, t) \in [n]^2 \setminus [m]^2$, define $g(s, t) = 1$. Let $X = \{a_1^g, \dots, a_n^g\}$ and $Y = \{b_1, \dots, b_m\}$. For each $1 \leq i \leq k$, define $Z_\alpha = \{c_j : j \in ((\alpha - 1)m, \alpha m]\}$. For each $\alpha \leq k$, let $Q^\alpha = \{a_i^g b_j : u_i w_j \in P_{UW}^\alpha\}$. Now consider some $1 \leq i, j, s \leq m$, and $a_i^g b_j c_s \in K_3[X, Y, Z_1 \cup \dots \cup Z_k]$. By construction there are some $\alpha, \beta \in [k]$ so that $c_s \in Z_\beta$ and $a_i^g b_j \in Q^\alpha$. By assumption on H , $a_i^g b_j c_s \in E$ if and only if $s \leq g(i, j)$. By definition of g , $g(i, j) = m\alpha$, and by definition of Z_β , $(\beta - 1)m < s \leq \beta m$. Thus $s \leq g(i, j)$ if and only if $\beta \leq \alpha$, so $a_i^g b_j c_s \in E$ if and only if $\beta \leq \alpha$. \square

Note that in Lemma 6.8, the partition of $K_2[U, W]$ is completely arbitrary. This will be used in the “(2) \Rightarrow (1)” direction of the following equivalence.

Proposition 6.9. *Suppose \mathcal{H} is a hereditary 3-graph property. Then the following are equivalent.*

- (1) $H(k)$ is encoded in \mathcal{H} for all $k \geq 1$,
- (2) \mathcal{H} has k -FOP₂ for all $k \geq 1$.

Proof. That (1) implies (2) is immediate from Proposition 6.7 and Definition 6.4. We now prove (2) implies (1). Suppose \mathcal{H} has m -FOP₂ for all m . Fix $k \geq 1$. We show $H(k)$ is binary encoded in \mathcal{H} . Fix $\epsilon_1, \rho > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Let $N, T, L \geq 1$ be as in Theorem 2.8 for ϵ_1, ϵ_2 , $t_0 = 2k + k^{2k}$, and $\ell_0 = 1$. Let $n \gg_{\epsilon_1, \epsilon_2, T, L} N$. Fix sets U, W, V_1, \dots, V_k , each of size n , and any partitions $K_2[U, W] = \bigcup_{\alpha \leq k} P_{UW}^\alpha$ such that P_{UW}^α satisfies $\text{disc}_2(\epsilon_2(L), 1/k)$. Such a partition exists by Lemma 3.24.

Let $M \gg n$. Since \mathcal{H} has FOP₂, by Lemma 3.31, it contains a clean copy of $F(M)$, say $H = (X \sqcup Y \sqcup Z, E)$. By Lemma 6.8, there are $X' = \{x_1, \dots, x_n\} \subseteq X$, $Y' = \{y_1, \dots, y_n\} \subseteq Y$, and $Z'_1, \dots, Z'_k \subseteq Z$ such that if $Q_{XY}^\alpha = \{x_i y_j : u_i w_j \in P_{UW}^\alpha\}$, then the following holds: for all $z \in Z_\beta$ and $x_i y_j \in Q_{XY}^\alpha$, $zx_i y_j \in E$ if and only if $\beta \leq \alpha$.

For each $m \in [k]$, choose partitions $K_2[X', Z'_m] = \bigcup_{\alpha \leq k} Q_{XZ'_m}^\alpha$ and $K_2[Y', Z'_m] = \bigcup_{\alpha \leq k} Q_{YZ'_m}^\alpha$ so that for each $\alpha \leq k$, $Q_{XZ'_m}^\alpha$ and $Q_{YZ'_m}^\alpha$ satisfy $\text{disc}_2(\epsilon_2(k), 1/k)$ (such partitions exist by Lemma 3.24). Define a decomposition \mathcal{Q} of $V' = X' \cup Y' \cup \bigcup_{i=1}^k Z'_i$ by setting $\mathcal{Q}_{\text{vert}} = \{X', Y', Z'_1, \dots, Z'_k\}$ and $\mathcal{Q}_{\text{edge}} = \{Q_{XY}^\alpha : \alpha \leq k\} \cup \{Q_{YZ'_m}^\alpha, Q_{XZ'_m}^\alpha : m \in [k], \alpha \leq k\}$. By construction, \mathcal{Q} is a $(2 + k, k, \epsilon_1, \epsilon_2(k))$ -decomposition V' . Let $H' = H[V']$. Clearly there is an encoding of $H(k)$ in (H', \mathcal{Q}) , namely (f, g) , where $f : a_i \mapsto Q_{XY}^i$ and $g : b_j \mapsto Q_{XZ'_j}^1 Q_{XZ'_j}^1$. We have now shown (1) holds, as desired. \square

We will now prove the analogous result for $U(k)$ and IP₂.

Proposition 6.10. *Suppose \mathcal{H} is a hereditary 3-graph property. Then the following are equivalent.*

- (1) $U(k)$ is binary encoded in \mathcal{H} for all $k \geq 1$,
- (2) \mathcal{H} has k -IP₂ for all $k \geq 1$.

Proof. That (1) implies (2) follows immediately from Proposition 6.6 and Definition 6.4. Conversely, assume \mathcal{H} has m -IP₂ for all $m \geq 1$. Fix $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and set $\ell = 2^k$ and $t = 2^k + 2$. Choose N as in Lemma 3.24 for $\epsilon = \delta = \epsilon_2(\ell)$ and ℓ . Suppose $n \geq N$, and B, C , and A_1, \dots, A_ℓ are sets of size n . By Lemma 3.24, there exists a partition $K_2[B, C] = \bigcup_{S \subseteq [k]} P_{BC}^S$ so that each P_{BC}^S has $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Let $H = (V, E)$ be the 3-partite 3-graph with vertex set $A_1 \cup \dots \cup A_\ell \cup B \cup C$, and edge set $\bigcup_{S \subseteq [k]} \bigcup_{i \in S} \bigcup_{a \in A_i} \{abc : bc \in P_{BC}^S\}$.

By Fact 3.30, there is $H' = (V', E') \in \mathcal{H}$ be a clean copy of H . We may assume $V' = X_1 \cup \dots \cup X_\ell \cup Y \cup Z$ where $Y = \{y_b : b \in B\}$, $Z = \{z_c : c \in C\}$, for each $i \in [\ell]$, $X_i = \{x_a : a \in A_i\}$, such that $x_a y_b z_c \in E'$ if and only if $abc \in E$. Use Lemma 3.24 to choose partitions $K_2[X_i, Y] = \bigcup_{\alpha \leq \ell} Q_{X_i Y}^\alpha$ and $K_2[X_i, Z] = \bigcup_{\alpha \leq \ell} P_{X_i Z}^\alpha$ so that each $Q_{X_i Y}^\alpha$, $Q_{X_i Z}^\alpha$ satisfies $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. For each $S \subseteq [k]$, let $Q_{YZ}^S = \{y_b z_c : bc \in P_{BC}^S\}$. Let \mathcal{Q} be the decomposition of V' with $\mathcal{Q}_{\text{vert}} = \{X_1, \dots, X_\ell, Y, Z\}$ and

$$\mathcal{Q}_{\text{edge}} = \{Q_{X_i Y}^\alpha : i \in [k], \alpha \leq \ell\} \cup \{Q_{X_i Z}^\alpha : i \in [k], \alpha \leq \ell\} \cup \{Q_{YZ}^S : S \subseteq [k]\}.$$

By construction, \mathcal{Q} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V' . Clearly $U(k)$ is encoded in (H', \mathcal{Q}) via $f : a_i \mapsto Q_{X_i Y}^1 Q_{X_i Z}^1$ and $g : c_S \mapsto Q_{YZ}^S$. This shows $U(k)$ is encoded in \mathcal{H} , so (1) holds. \square

6.2. Removal in almost stable bipartite graphs. In this section, we prove a removal type result, which roughly says the following. If a bipartite graph $G = (U \cup W, E)$ contains few d -trees with leaves in W , then it is close to a graph in which $\text{rk}(W) \leq d$ (see Section 5.4). We begin with a definition which will help us keep tabs on the locations of the leaves and nodes of a d -tree.

Definition 6.11. Suppose $d \geq 1$, $G = (V, E)$ is a bipartite graph, and $A, B \subseteq V$. An (A, B, d) -tree in G is a d -tree in G with leaves in B and nodes in A .

We now define a notion of closeness tailored to the bipartite setting.

Definition 6.12. Suppose $\delta > 0$ and $G = (U \cup W, E)$, $G' = (U \cup W, E')$ are bipartite graphs. Then G and G' are *uniformly (U, W, δ) -close* if for all $u \in U$, $|N_G(u) \Delta N_{G'}(u)| \leq \delta |W|$.

Definition 6.12 is clearly stronger than the notion of δ -closeness (recall G and G' on the same vertex set are δ -close if $|E(G) \Delta E(G')| \leq \delta |V(G)|^2$). We will need this extra strength in our proof of Theorem 2.49. We can now state our removal lemma.

Theorem 6.13. *For all $d \geq 1$ and $0 < \epsilon < 1/9$, there is $\delta_0(\epsilon, d)$ such that for all $0 < \delta < \delta_0$ there is $\rho = \rho(\epsilon, d, \delta) > 0$ and $N = N(\epsilon, d, \rho)$ such that the following holds.*

Suppose $G = (U \cup W, E)$ is a bipartite graph with $|U|, |W| \geq N$ and the number of (U, W, d) -trees in G is at most $\rho |U|^{2^d-1} |W|^{2^d}$. Then there is $U_0 \subseteq U$ and $W_0 \subseteq W$ such that $|U_0| \leq \delta |U|$, $|W_0| \leq \epsilon |W|$, and a bipartite graph $G' = ((U \setminus U_0) \cup (W \setminus W_0), E')$ and so

that $G[U \setminus U_0, W \setminus W_0]$ is uniformly $(U \setminus U_0, W \setminus W_0, \delta)$ -close to G' , and so that G' contains no $(U \setminus U_0, W \setminus W_0, d)$ -trees.

In the proof of Theorem 2.49, we will apply Theorem 6.13 to a graph $G = (U \cup W, E)$ containing few (U, W, d) -trees and where $|W| \ll |U|$. This imbalance in the size of the parts is another reason we require Theorem 6.13, and cannot use an off-the-shelf removal lemma, such as Lemma 3.7.

We begin with a lemma which shows that if a bipartite graph $G = (U \cup W, E)$ contains few (U, W, d) -trees, then W can be almost partitioned into almost good sets. This can be thought of as a version of Theorem 5.15 with a weaker conclusion, but which has the advantage of more general hypotheses. In particular, Lemma 6.14 applies when there are few (U, W, d) -trees (rather than none, as required in Theorem 5.15).

Lemma 6.14. *For all $d \geq 1$, there is $C = C(d) > 0$ such that for all $0 < \epsilon < 1/9$, there is $\mu_0(\epsilon, d) > 0$, such that for all $0 < \mu < \mu_0$, there are $M = M(\epsilon, d, \mu)$ and $N = N(\epsilon, d, \mu)$ such that the following holds.*

Suppose $G = (U \cup W, E)$ is a bipartite graph with $|U|, |W| \geq N$ and such that the number of (U, W, d) -trees in G is at most $\mu^C |U|^{2^d-1} |W|^{2^d}$. Then there is $m \leq M$, $U' \subseteq U$ of size at most $\mu |U|$, and a partition $W_0 \cup W_1 \cup \dots \cup W_m$ such that $|W_0| \leq \epsilon |W|$, and such that for each $1 \leq i \leq m$, $|W_i| \geq |W|/M$ and W_i is μ -good with respect to all $u \in U \setminus U'$.

Proof. We proceed by induction on d . Suppose first $d = 1$. In this case, set $C = 4$. Given $0 < \epsilon < 1/4$, set $N = M = 1$ and let $\mu_0 = \epsilon$.

Now assume $0 < \mu < \mu_0$, and $G = (U \cup W, E)$ is a bipartite graph containing at most $\mu^C |U| |W|^2$ many $(U, W, 1)$ -trees. Note that for all $u \in U$, if $uw_1 \in E$ and $uw_0 \notin E$, then (u, w_0, w_1) is a $(U, W, 1)$ -tree. Thus, for all $u \in U$, there are at $|N(u) \cap W| |\neg N(u) \cap W|$ many $(U, W, 1)$ -trees in G with $a_{<>} = u$. Consequently there must be a set $U' \subseteq U$ of size at most $\mu^{C/2} |U|$, such that for all $u \in U \setminus U'$, $|N(u) \cap W| |\neg N(u) \cap W| \leq \mu^{C/2} |W|^2$ (since otherwise, G would contain more than $\mu^C |U| |W|^2$ many $(U, W, 1)$ -trees). This means that for all $u \in U \setminus U'$, we must have that $\min\{|N(u) \cap W|, |\neg N(u) \cap W|\} \leq \mu^{C/4} |W| = \mu |W|$. This shows that W itself is μ -good with respect to all $u \in U \setminus U'$. This finishes the case $d = 1$.

Suppose now $d > 1$, and assume that for all $1 \leq d' < d$, we have defined $C(d')$, $\mu_0(d', \epsilon')$, $N(d', \epsilon', \mu)$, and $M(d', \epsilon', \mu)$, for all $0 < \epsilon' < 1/4$ and all $0 < \mu < \mu_0(d', \epsilon')$. Set $C' = C(d-1)$, and define $C(d) = 8C' + 2^d + 2$. Fix $0 < \epsilon < 1/4$. Define $\mu_0(d, \epsilon) = \epsilon^{2^{d+1}} \mu_0(\epsilon, d-1)/4$. Given $0 < \mu < \mu_0$, let $N' = N(d-1, \epsilon^5, \mu^4)$ and $M' = M(d-1, \epsilon^5, \mu^4)$. Finally, we define $M = M(d, \epsilon) := \epsilon^{-100} M'$ and $N = N(d, \epsilon) := \epsilon^{-100} N' M'$.

Suppose $G = (U \cup W, E)$ is a bipartite graph with $|U|, |W| \geq N$, and assume there are at most $\mu^C |U|^{2^d-1} |W|^{2^d}$ many (U, W, d) -trees in G . We define an integer s and a sequence W_0, \dots, W_s and Z_0, \dots, Z_s as follows.

Step 0: Let $W_0 = \emptyset$ and $Z_0 = W$.

Step $i+1$: Suppose we have defined $W_0, \dots, W_i, Z_0, \dots, Z_i$ so that for each $0 \leq i' \leq i$, $Z_{i'} = W \setminus (\bigcup_{u=0}^{i'} W_u)$, and for all $1 \leq i' \leq i$, there are at most $\mu^{4C'} |U|^{2^{d-1}-1} |W_{i'}|^{2^{d-1}}$ many

$(U, W_{i'}, d-1)$ -trees in G , and $|W_{i'}| \geq \epsilon^2 \mu |W|$. Let

$$U_{i+1} = \{u \in U : \min\{|N(u) \cap Z_i|, |\neg N(u) \cap Z_i|\} \geq \mu |Z_i|\}.$$

If either $|Z_i| \leq \epsilon^2 |W|$ or $|U_{i+1}| \leq \mu |U|$, define $s = i$ and end the construction.

Otherwise, for each $u \in U_{i+1}$ set $W_{i,1}(u) = N(u) \cap Z_i$ and $W_{i,0}(u) = \neg N(u) \cap Z_i$. By definition of U_{i+1} and by assumption on the size of Z_i , we know that for all $u \in U_{i+1}$, both $W_{i,1}(u)$ and $W_{i,0}(u)$ have size at least $\mu |Z_i| \geq \mu \epsilon^2 |W|$.

Let U'_{i+1} be the set of $u \in U_{i+1}$ for which there more than $\mu^{4C'} |U|^{2^{d-1}-1} |W_{i,1}(u)|^{2^{d-1}}$ many $(U, W_{i,1}(u), d-1)$ -trees and more than $\mu^{4C'} |U|^{2^{d-1}-1} |W_{i,0}(u)|^{2^{d-1}}$ many $(U, W_{i,0}(u), d-1)$ -tree in G . We claim that $U_{i+1} \setminus U'_{i+1} \neq \emptyset$. Indeed, suppose not. Then we construct many copies of $T(d)$ as follows.

- Choose $u \in U'_{i+1}$ and set $u_{<>} = u$. There are $|U'_{i+1}| = |U_{i+1}| > \mu |U|$ choices.
- Choose $(u_\sigma^1)_{\sigma \in 2^{<d-1}} (w_\rho^1)_{\rho \in 2^{d-1}}$ a $(U, W_{i,1}(u), d-1)$ -tree in G , and for each $\sigma \in 2^{<d-1}$ and $\rho \in 2^{d-1}$, set $u_{1 \wedge \sigma} = u_\sigma^1$ and $w_{1 \wedge \rho} = w_\rho^1$. By assumption, there more than

$$\mu^{4C'} |U|^{2^{d-1}-1} |W_{i,1}(u)|^{2^{d-1}} \geq \mu^{4C'} |U|^{2^{d-1}-1} (\mu \epsilon^2 |W|)^{2^{d-1}}$$

ways to do this, where the inequality is because $u \in U_{i+1}$.

- Choose $(u_\sigma^0)_{\sigma \in 2^{<d-1}} (w_\rho^0)_{\rho \in 2^{d-1}}$ a $(U, W_{i,0}(u), d-1)$ -tree in G , and for each $\sigma \in 2^{<d-1}$ and $\rho \in 2^{d-1}$, set $u_{0 \wedge \sigma} = u_\sigma^0$ and $w_{0 \wedge \rho} = w_\rho^0$. By assumption, there are more than

$$\mu^{4C'} |U|^{2^{d-1}-1} |W_{i,1}(u)|^{2^{d-1}} \geq \mu^{4C'} |U|^{2^{d-1}-1} (\mu \epsilon^2 |W|)^{2^{d-1}}$$

ways to do this, where the inequality is because $u \in U_{i+1}$.

Every such construction produces a distinct tuple $(u_\sigma)_{\sigma \in 2^{<d}} (w_\rho)_{\rho \in 2^d}$ which is a (U, W, d) -tree in G . But this means the number of (U, W, d) -trees in G is at least the following,

$$(\mu |U|) (\mu^{4C'} |U|^{2^{d-1}-1} (\mu \epsilon^2 |W|)^{2^{d-1}})^2 = \epsilon^{2^{d+1}} \mu^{8C'+2^d} |U|^{2^{d-1}} |W|^{2^d} > \mu^C |U|^{2^{d-1}} |W|^{2^d},$$

where the inequality uses that $\mu < \mu_0$ and the definition of C . This is a contradiction, so we must have that $U_{i+1} \setminus U'_{i+1} \neq \emptyset$.

Thus we may choose some $u \in U_{i+1} \setminus U'_{i+1}$. By definition of U'_{i+1} , there is some $W_{i+1} \in \{W_{i,1}(u), W_{i,0}(u)\}$ with the property that there are at most $\mu^{4C'} |U|^{2^{d-1}-1} |W_{i+1}|^{2^{d-1}}$ many $(U, W_{i+1}, d-1)$ -trees in G . Set $Z_{i+1} = W \setminus (\bigcup_{j=1}^{i+1} W_j)$. This ends step $i+1$.

Clearly this construction will end in some $s \leq \epsilon^{-2} \mu^{-1}$ steps (since otherwise $|\bigcup_{i=1}^s W_i| > |W|$). At the end, we will have a partition $W = W_1 \cup \dots \cup W_s \cup Z_s$, where $|Z_s| \leq \epsilon^2 |W|$, and where for each $1 \leq i \leq s$, there are at most $\mu^{4C'} |U|^{2^{<d-1}} |W_i|^{2^{d-1}}$ many $(U, W_i, d-1)$ -trees in G and $|W_i| \geq \mu \epsilon^2 |W| \geq N' = N(\epsilon^2, d-1, \mu^4)$. For each $i \in [s]$, the inductive hypothesis applied to $G[U \cup W_i]$ implies there is some $s_i \leq M' = M(\epsilon^2, d-1, \mu^4)$, a partition $W_i = W_{i,0} \cup W_{i,1} \cup \dots \cup W_{i,s_i}$, and $U_i \subseteq U$ such that $|U_i| \leq \mu^4 |U|$, $|W_{i,0}| \leq \epsilon^2 |W_i|$, and such that for each $j \in [s_i]$, $W_{i,j}$ is μ^4 -good in $G[U \cup W_i]$ with respect to every $u \in U \setminus U_i$. Clearly this implies each $W_{i,j}$ is μ^4 -good with respect to every $u \in U \setminus U_i$ in G as well. Observe that each $|W_{i,j}| \geq \mu \epsilon^2 |W| / M' \geq |W| / M$.

Set $U_0 = \bigcup_{i=1}^s U_i$ and $W_0 = Z_s \cup \bigcup_{i=1}^s W_{i,0}$. Since $s \leq \epsilon^{-2}\mu^{-1}$, $|U_0| \leq \mu^4\epsilon^{-2}\mu^{-1}|U| < \mu^2|U|$. Further,

$$|W_0| \leq \epsilon^2|W| + \epsilon^2 \sum_{i=1}^s |W_i| \leq 2\epsilon^2|W| \leq \epsilon|W|.$$

Let $W'_1, \dots, W_{m'}$ enumerate $\{W_{i,j} : 1 \leq i \leq s, 1 \leq j \leq s_i\}$. Note

$$m' = \sum_{i=1}^s s_i \leq sM' \leq \epsilon^{-2}\mu^{-1}M' \leq M.$$

Thus we have a partition $W = W_0 \cup W'_1 \cup \dots \cup W_{m'}$, where $m' \leq M$, $|W_0| \leq \epsilon|W|$, and such that for each $1 \leq u \leq m'$, W'_u is μ -good to every $u \in U \setminus U_0$ and satisfies $|W'_u| \geq \mu|W|/M$. This finishes the proof. \square

The last remaining ingredient is a bipartite counting lemma (for a proof, see Appendix E of [60]).

Lemma 6.15. *For all $k \geq 1$ and $\epsilon > 0$ there are $\eta > 0$ and N such that the following hold. Suppose $F = (U \cup W, R)$ is a bipartite graph with $|U| + |W| = k$. Assume $G = (V, E)$ is a graph, and $V = \bigcup_{u \in U} V_u \cup \bigcup_{w \in W} V_w$ is a partition of V into parts of size at least N . Suppose that for each $uw \in R$, $|E \cap K_2[V_u, V_w]| \geq (1 - \eta)|V_u||V_w|$, and for each $uw \notin R$, $|E \cap K_2[V_u, V_w]| \leq \eta|V_u||V_w|$. Then there are at least $(1 - \epsilon) \prod_{x \in U \cup W} |V_x|$ many tuples $(v_u)_{u \in U} (v_w)_{w \in W} \in \prod_{u \in U} V_u \times \prod_{w \in W} V_w$ such that $v_u v_w \in E$ if and only if $uw \in R$.*

PROOF OF THEOREM 6.13: Fix $d \geq 1$ and $0 < \epsilon < 1/9$. Let $\eta = \eta(2^d + 2^d - 1, 1/2)$ and $N_2 = N_2(2^d + 2^d - 1, 1/2)$ be from Lemma 6.15. Let $C = C(d)$ and $\mu_0(\epsilon, d)$ be from Lemma 6.14. Set $\delta_0 = \min\{\mu_0(\epsilon, d), \eta\}$. Given $0 < \delta < \delta_0$, let $M = M(\epsilon, \delta, d)$, $N_1 = N_1(\epsilon, \delta, d)$ be from Lemma 6.14. Then define $\rho = \frac{1}{4}(\delta/M2^M)^C$ and let $N = 2^M \mu^{-1} \max\{N_1, N_2\}$.

Suppose $G = (U \cup W, E)$ is a bipartite graph with $|U|, |W| \geq N$ such that the number of (U, W, d) -trees in G is at most $\rho|U|^{2^d-1}|W|^{2^d}$. Since $\rho < \delta^C$, Lemma 6.14 implies there exist $m \leq M$, $U_0 \subseteq U$ of size at most $\delta|U|$, a partition $W_0 \cup W_1 \cup \dots \cup W_m$ such that $|W_0| \leq \epsilon|W|$, and such that for each $1 \leq i \leq m$, $|W_i| \geq |W|/M \geq N_2$ and W_i is δ -good with respect to all $u \in U \setminus U_0$.

For each $1 \leq i \leq m$ and $u \in U \setminus U_0$, let $\xi(u, W_i)$ be such that $|N^{\xi(u, W_i)}(u) \cap W_i| \geq (1 - \delta)|W_i|$. Partition $U \setminus U_0$ into a minimal number of parts $U_1 \cup \dots \cup U_t$ so that if u and u' are in the same part, then $\xi(u, W_j) = \xi(u', W_j)$ for all $j \in [m]$. Clearly $t \leq 2^m$. Then define $I = \{i \in [t] : |U_i| < \mu|U|/2^t\}$, and set $U'_0 = U_0 \cup \bigcup_{i \in I} U_i$. Clearly $|U'_0| \leq 2\delta|U|$. Let U'_1, \dots, U'_t be a reindexing of $\{U_i : i \in [t] \setminus I\}$. For each $i \in [t]$, let $\xi(U'_i, W_j)$ be such that $\xi(u, W_j) = \xi(U'_i, W_j)$ for all $u \in U'_i$.

Set $\mathbf{U} = \{U'_1, \dots, U'_t\}$ and $\mathbf{W} = \{W_1, \dots, W_m\}$, and define the bipartite graph

$$\mathbf{R} = (\mathbf{U} \cup \mathbf{W}, \{\xi(U'_i, W_j) : \xi(U'_i, W_j) = 1\}).$$

Note that each $|W_i| \geq |W|/M \geq N/M \geq N_2$, and each $|U'_i| \geq \delta|U|/2^t \geq \delta N/2^t \geq N_2$. We claim there are no $(\mathbf{U}, \mathbf{W}, d)$ -trees in \mathbf{R} .

Suppose towards a contradiction there exists $(U_\sigma)_{\sigma \in 2^{<d}} (W_\lambda)_{\lambda \in 2^d}$ a $(\mathbf{U}, \mathbf{W}, d)$ -tree in \mathbf{R} . Then for each $\sigma \in 2^{<d}$ and $\lambda \in 2^d$, $|E \cap K_2[U_\sigma, W_\lambda]|/|K_2[U_\sigma, W_\lambda]| \in [0, \delta) \cup (1 - \delta, 1]$. By

Lemma 6.15, the number of (U, W, d) -trees $(u_\sigma) \in 2^{<d}(w_\lambda)_{\lambda \in 2^d} \in \prod_{\sigma \in 2^{<d}} U_\sigma \times \prod_{\lambda \in 2^d} W_\lambda$ is at least

$$\frac{1}{2} \prod_{\sigma \in 2^{<d}} |U_\sigma| \prod_{\lambda \in 2^d} |W_\lambda| \geq \frac{1}{2} (\delta |U| / 2^t)^{2^d-1} (|W|/M)^{2^d} > \rho |U|^{2^d-1} |W|^{2^d},$$

a contradiction. Thus there exist no $(\mathbf{U}, \mathbf{W}, d)$ -trees in \mathbf{R} . Define

$$G' = ((U \setminus U'_0) \cup (W \setminus W'_0), \cup E') \text{ where } E' = \bigcup_{U_i, W_j \in E(\mathbf{R})} K_2[U_i, W_j].$$

We claim that G' contains no $(U \setminus U'_0, W \setminus W'_0, d)$ -trees. Suppose towards a contradiction there exists some $(u_\sigma)_{\sigma \in 2^{<d}}(w_\lambda)_{\lambda \in 2^d}$ which is a $(U \setminus U'_0, W \setminus W'_0, d)$ -tree in G' . By the definition of G' and the fact that there are no $(\mathbf{U}, \mathbf{W}, d)$ -trees in \mathbf{R} , we know there must be some $\lambda_1 \neq \lambda_2$ in 2^d and a $W_i \in \mathbf{W}$ with $w_{\lambda_1}, w_{\lambda_2} \in W_i$. Since $\lambda_1 \neq \lambda_2$, there is some $\tau \in 2^{<d}$ such that $\tau \wedge 0 \leq \lambda_1$ and $\tau \wedge 1 \leq \lambda_2$, or vice versa. Without loss of generality, assume $\tau \wedge 0 \leq \lambda_1$ and $\tau \wedge 1 \leq \lambda_2$. Thus there is $u_\tau \in U \setminus U'_0$ so that $u_\tau w_{\lambda_1} \in E$ and $u_\tau w_{\lambda_2} \notin E$. But there is some $U'_j \in \mathbf{U}$ with $u_\tau \in U'_j$. But by definition, since $w_{\lambda_1}, w_{\lambda_2} \in W_i$, we have that $u_\tau w_{\lambda_1}$ and $u_\tau w_{\lambda_2}$ are both in $E^{\delta(U'_j, W_i)}$, a contradiction. This finishes our verification that G' contains no $(U \setminus U'_0, W \setminus W'_0, d)$ -trees.

It is easy to see that for all $u \in U \setminus U'_0$,

$$|(N_{G'}(u) \Delta N_G(u)) \cap (W \setminus W'_0)| \leq \sum_{W_j \in \mathbf{W}} |(N_{G'}(u) \Delta N_G(u)) \cap W_j| \leq \sum_{W_j \in \mathbf{W}} \delta |W_j| \leq \delta |W \setminus W'_0|.$$

Thus $G[U \setminus U'_0, W \setminus W'_0]$ is uniformly $(U \setminus U'_0, W \setminus W'_0, \delta)$ -close to G' . \square

6.3. A property admits linear $\text{disc}_{2,3}$ -error if and only if it is NFOP_2 . In this subsection we prove our main result about NFOP_2 properties, which is that they are exactly the properties admitting linear $\text{disc}_{2,3}$ -error. The harder direction is Theorem 2.49, which implies that an NFOP_2 property admits linear $\text{disc}_{2,3}$ -error. The proof will apply Theorem 6.13 along with a refinement of Theorem 5.15 to auxiliary graph structures like the ones defined in Section 6.1 (see Definition 6.2).

Before we start the proof of Theorem 2.49, we prove two additional lemmas. The first is a refinement of Theorem 5.15 which has additional equitability requirements.

Theorem 6.16. *Suppose $d \geq 0$. For all $\epsilon > 0$ and non-increasing functions $f : \mathbb{N} \rightarrow (0, 1]$ with $\lim_{n \rightarrow \infty} f(n) = 0$, there is $M \geq 1$ such that the following hold.*

Suppose $G = (U \cup W, E)$ is a bipartite graph, and $W = W_1 \cup \dots \cup W_m$ is a partition satisfying $|W_1| = \dots = |W_m|$, and such that for each $i \in [m]$, G contains no (U, W_i, d) -trees.

Then there are some $m' \leq M$, $n' \leq |W|$, and a set $\Omega \subseteq [m]$, such that $|\bigcup_{u \in \Omega} W_u| \leq \epsilon |W|$, and such that for all $u \in [m] \setminus \Omega$, there is a partition

$$W_u = W_{u,0} \cup W_{u,1} \cup \dots \cup W_{u,m'},$$

so that $|W_{u,0}| \leq \epsilon |W_u|$, and so that for each $1 \leq i \leq m'$, $W_{u,i}$ is $f(m')$ -good in G and $|W_{u,i}| = n'$.

The difference between the above and Theorem 5.15 is that we begin with an equipartition, and end up refining each W_i into the same number of parts, all of which are of the same size.

PROOF OF THEOREM 6.16: Fix $d \geq 0$, $\epsilon > 0$, and a non-increasing function $f : \mathbb{N} \rightarrow (0, 1]$ with $\lim_{n \rightarrow \infty} f(n) = 0$. Define $\psi(x) = \epsilon^4 f(\epsilon^{-4}x)/x^2$. Let M_1 be as in Theorem 5.15 for d , ϵ^8 , and ψ . Then define $M = \epsilon^{-2}(M_1)^2$.

Suppose $G = (U \cup W, E)$ is a bipartite graph, and $W = W_1 \cup \dots \cup W_m$ is a partition so that $|W_1| = \dots = |W_m|$, and so that G contains no (U, W_i, d) -trees for each $i \in [m]$.

By Theorem 5.15, there is some $m_1 \leq M_1$ and a set $\Omega \subseteq [m]$, such that $|\bigcup_{u \in \Omega} W_u| \leq \epsilon^8 |W|$ and so that for each $u \in [m] \setminus \Omega$, there are $s_u \leq m_1$ and a partition

$$W_u = W_{u,0} \cup W_{u,1} \cup \dots \cup W_{u,s_u},$$

so that $|W_{u,0}| \leq \epsilon^8 |W_u|$, and so that for each $1 \leq i \leq s_u$, $W_{u,i}$ is $\psi(m_1)$ -good. By reindexing if necessary, we may assume that for each $u \in [m] \setminus \Omega$, $|W_{u,1}| \geq \dots \geq |W_{u,s_u}|$. Set $K = \lceil \epsilon^2 |W| / m_1 \rceil$, and let $\Omega' = \Omega \cup \{u \in [m] \setminus \Omega : |W_{u,1}| < K\}$.

For each $u \in [m] \setminus \Omega'$, let p_u be maximal so that $|W_{u,p_u}| \geq K$, and for each $1 \leq v \leq p_u$, choose a partition

$$W_{u,v} = W_{u,v,0} \cup W_{u,v,1} \cup \dots \cup W_{u,v,s_{u,v}},$$

with the property that $|W_{u,v,0}| < K$ and for each $1 \leq w \leq s_{u,v}$, $|W_{u,v,w}| = K$. Then set $q_u = \sum_{v \in [p_u]} s_{u,v}$, and fix an enumeration

$$\{W_{u,v,w} : v \in [p_u], w \in [s_{u,v}]\} = \{X_{u,1}, \dots, X_{u,q_u}\} \text{ so that } |X_{u,1}| \geq \dots \geq |X_{u,q_u}|.$$

Then observe that

$$|W_u \setminus \left(\bigcup_{v=1}^{q_u} X_{u,v} \right)| \leq |W_{u,0}| + \left| \bigcup_{v=p_u+1}^{r_u} W_{u,v} \right| + K \leq \epsilon^2 |W_u| + \frac{\epsilon^2 |W_u| r_u}{m_1} + \frac{\epsilon^2 |W_u|}{m_1} \leq 4\epsilon^2 |W_u|.$$

Note that for all $u \in [m] \setminus \Omega'$, $|(\bigcup_{v=1}^{q_u} X_{u,v})| = q_u K \geq (1 - 4\epsilon^2) |W_u| = (1 - 4\epsilon^2) |W_1|$. Therefore, setting $q = \min\{q_u : u \in [m]\}$, we know $|W_1| \geq qK \geq (1 - 4\epsilon^2) |W_1|$. For each $u \in [m] \setminus \Omega'$, define

$$X_{u,0} = W_{u,0} \cup \left(\bigcup_{v=p_u+1}^{r_u} W_{u,v} \right) \cup \left(\bigcup_{v=1}^{p_u} W_{u,v,0} \right) \cup \left(\bigcup_{q < q' \leq q_u} X_{u,q'} \right).$$

Then we note that

$$\begin{aligned} |X_{u,0}| &\leq \epsilon^2 |W_u| + \frac{2(r_u - p_u)\epsilon^2 |W_u|}{m_1} + \frac{2p_u\epsilon^4 |W_u|}{m_1} + (|W_u| - qK) \leq 5\epsilon^2 |W_u| + 4\epsilon^2 |W_1| \\ &= 9\epsilon^2 |W_u| \\ &< \epsilon |W_u|, \end{aligned}$$

where the last inequality is because $\epsilon < 1/4$. For all $u \in [m] \setminus \Omega'$, we now have a new partition of W_u , namely

$$W_u = X_{u,0} \cup X_{u,1} \cup \dots \cup X_{u,q}.$$

where $|X_{u,0}| \leq \epsilon|W_u|$ and for each $1 \leq v \leq q$, $|X_{u,v}| = K$. Further, note that since $qK \leq |W_1|$, $q \leq \epsilon^{-4}m_1 \leq M$.

We now show that for all $u \in [m] \setminus \Omega'$ and $q' \in [q]$, $X_{u,q'}$ is $f(q)$ -good. Fix $u \in [m] \setminus \Omega'$ and $1 \leq q' \leq q$. By construction, there is some $v \in [p_u]$ and $w \in [s_{u,v}]$ so that $X_{u,q'} = W_{u,v,w}$, where $W_{u,v,w}$ has size K , and is contained in $W_{u,v}$, a set which is $\psi(m_1)$ -good. Thus for all $u \in U$ there is some $\delta \in \{0, 1\}$ so that $|N^\delta(u) \cap W_{u,v}| \leq \psi(m_1)|W_{u,v}|$. By definition of ψ and K , we have that

$$\begin{aligned} |N^\delta(u) \cap W_{u,v,w}| &\leq \psi(m_1)|W_{u,v}| \leq \frac{\psi(m_1)|W_u|}{K}|W_{u,v,w}| \leq \psi(m_1)\epsilon^{-4}m_1|W_{u,v,w}| \\ &\leq f(q)|W_{u,v,w}|. \end{aligned}$$

This shows $X_{u,q'}$ is $f(q)$ -good, which finishes the proof. \square

Our second lemma, Lemma 6.17 below, allow us to ensure certain divisibility requirements hold in regular decompositions.

Lemma 6.17. *For all $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $k \geq 1$, there are $\mu_1 > 0$ and $\mu_2 : \mathbb{N} \rightarrow (0, 1]$, such that for all $L, T \geq 1$, there is $N \geq 1$ such that the following holds.*

Suppose $H = (V, E)$ is a 3-graph and $|V| = n \geq N$. Suppose $1 \leq t \leq T$, $1 \leq \ell \leq L$, and \mathcal{P} is an $(\mu_1, \mu_2(\ell), t, \ell)$ decomposition of V which is $(\mu_1, \mu_2(\ell), t, \ell)$ -regular and μ_1 -homogeneous with respect to H . Then there exists \mathcal{P}' is an $(\epsilon_1, \epsilon_2(\ell'), t, \ell')$, decomposition of V , such that $\ell' = k\ell$, and so that \mathcal{P}' is $(\epsilon_1, \epsilon_2(\ell'), t, \ell')$ -regular and ϵ_1 -homogeneous with respect to H .

Proof. Choose μ'_1, μ'_2 as in Lemma 3.28 for ϵ_1 and ϵ_2 . Define $\mu_2 : \mathbb{N} \rightarrow (0, 1]$ by choosing $\mu_2(x) \ll_k \mu'_2(x)$ for all x . Given $T, L \geq 1$, choose $N \gg N_1, T, L, \epsilon_2(L)^{-1}$.

Now suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$. Suppose $1 \leq t \leq T$, $1 \leq \ell \leq L$, and \mathcal{P} is an $(\mu'_1, \mu_2(\ell), t, \ell)$ decomposition of V which is $(\mu'_1, \mu_2(\ell), t, \ell)$ -regular and μ'_1 -homogeneous with respect to H . For each $P_{ij}^\alpha \in \mathcal{P}_{edge}$ satisfying $\text{disc}_2(\mu_2(\ell), 1/\ell)$, apply Lemma 3.24 with $\epsilon = \delta = \mu_2(\ell)$ to obtain a partition $P_{ij}^\alpha = P_{ij}^{\alpha,1} \cup \dots \cup P_{ij}^{\alpha,\ell k}$, so that each $P_{ij}^{\alpha,u}$ satisfies $\text{disc}_2(\mu_2(\ell))$ and has density $(1 \pm 2\mu_2(\ell))\frac{1}{k\ell}$. By our choice of μ_2 , each such $P_{ij}^{\alpha,\beta}$ satisfies $\text{disc}_2(\mu'_2(k\ell), 1/k\ell)$. Let \mathcal{P}' have $\mathcal{P}'_{vert} = \mathcal{P}_{vert}$ and

$$\mathcal{P}'_{edge} = \{P_{ij}^{\alpha,\beta} : P_{ij}^\alpha \text{ satisfies } \text{disc}_2(\mu_2(\ell), 1/\ell), 1 \leq \beta \leq k\} \cup \{P_{ij}^\alpha : P_{ij}^\alpha \text{ fails } \text{disc}_2(\mu_2(\ell), 1/\ell)\}.$$

Then \mathcal{P}' is a $(\mu'_1, \mu'_2(\ell'), t, \ell')$, decomposition of V , where $t' = t$ and $\ell' = \ell k$. It is straightforward to see that \mathcal{P}' is a $(0, 0)$ -approximate refinement of \mathcal{P} (see Definition 3.26). By Lemma 3.28, \mathcal{P}' is a $(t, \ell', \epsilon_1, \epsilon_2(\ell'))$ -decomposition of V which is $(\epsilon_1, \epsilon_2(\ell'))$ -regular and ϵ_1 -homogeneous with respect to H . \square

We are now ready to prove Theorem 2.49.

PROOF OF THEOREM 2.49: Fix $k \geq 1$, $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $\ell_0, t_0 \geq 1$. Choose $0 < \epsilon \ll \epsilon_1$, and let $d = d(k)$ as in part (2) in Theorem 5.12.

Choose $\epsilon_1^* > 0$ and $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ as in Proposition 6.7 for k . Define a function $\psi : \mathbb{N} \rightarrow (0, 1]$ by setting $\psi(x) = \epsilon^{100}/x^3$ for each $x \geq 1$. Then let $M_1 = M_1(d, \epsilon^{100}, f)$

and $N_1 = N_1(d, \epsilon^{100}, f)$ be as in Theorem 6.16, where $f : \mathbb{N} \rightarrow (0, 1]$ is defined by $f(x) = \epsilon^{100} \psi(2\epsilon^{-100}x(x+1))/(3x(x+1))$. Let $M \gg \epsilon^{-100}M_1$, and choose $\epsilon_0 \ll \epsilon/(M!)^3$. Let $\delta_0 = \delta_0(d, \epsilon_0)$ and $N_2 = N_2(d, \epsilon_0)$ be as in Theorem 6.13. Then choose

$$\epsilon'_1 \ll \delta \ll f((M!)^3)/(M!)^3, \epsilon_1^*, \epsilon_1, \delta_0, \epsilon^5, d^{-1}.$$

Define $\epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ by choosing $\epsilon'_2(x) \ll \epsilon'_1 \epsilon_2(x)/xM!$. Set $t'_0 = t_0$ and $\ell'_0 = \ell_0 N_3 \delta^{-1}$.

Now choose $\epsilon''_1 = \epsilon''_1(\epsilon'_1, \epsilon'_2)$ and $\epsilon''_2 = \epsilon''_2(\epsilon'_1, \epsilon'_2)$ as in Lemma 3.25. Let $\mu_1 \ll \epsilon''_1 \ll \epsilon'_1$, and define $\mu_2, \epsilon''_2 : \mathbb{N} \rightarrow (0, 1]$ by choosing $\mu_2(x) \ll \epsilon''_2(x) \ll \epsilon'_2(x)$ for all x . Then choose μ'_1 and μ'_2 be as in Lemma 6.17 for μ_1, μ_2 and $(M!)^3$. Finally, choose T, L , and N_3 as in Theorem 2.23 for t'_0, ℓ'_0, μ'_1 and μ'_2 , and let $N \gg N_1 N_2 N_3, T, M!, (\mu'_1)^{-1}, (\mu'_2(M!L))^{-1}, \delta^{-1}$.

Suppose $H = (V, E)$ is a 3-graph with $|V| \geq N$ which does not have k -FOP₂. Since H has VC₂-dimension less than k (see Fact 2.22), Theorem 2.23 implies there exist $\ell'_0 \leq \ell_1 \leq L$, $t'_0 \leq t \leq T$, and \mathcal{P}_1 a $(t, \ell_1, \mu'_1, \mu'_2(\ell_1))$ -decomposition of V which is $(\mu'_1, \mu'_2(\ell_1))$ -regular and μ'_1 -homogeneous with respect to H . By Lemma 6.17, there is \mathcal{P}_2 a $(t, \ell, \mu_1, \mu_2(\ell))$ -decomposition of V , where $\ell = (M!)^3 \ell_1$, and so that \mathcal{P}_2 is $(\mu_1, \mu_2(\ell))$ -regular and μ_1 -homogeneous with respect to H . By Lemma 3.25, there is \mathcal{P} a $(t, \ell, \epsilon''_1, \epsilon''_2(\ell))$ -decomposition of V , so that \mathcal{P} is $(\epsilon''_1, \epsilon''_2(\ell))$ -regular and ϵ''_1 -homogeneous with respect to H , and so that \mathcal{P} has no disc₂-irregular triads. Say $\mathcal{P}_{vert} = \{V_1, \dots, V_t\}$ and $\mathcal{P}_{edge} = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \in [\ell]\}$.

Given $ijs \in \binom{[t]}{3}$ and $1 \leq \alpha, \beta, \gamma \leq \ell$, set $G_{ijs}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_s, P_{ij}^\alpha \cup P_{js}^\beta \cup P_{is}^\gamma)$, $H_{ijs}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_s, E \cap K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma}))$ and let $d_{ijs}^{\alpha, \beta, \gamma}$ be such that $|E \cap K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma})| = d_{ijs}^{\alpha, \beta, \gamma} |K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma})|$. We will use throughout that since $\epsilon''_2(x) \ll \epsilon'_2(x)$, Corollary 3.6 implies that for all $ijs \in \binom{[t]}{3}$ and $\alpha, \beta, \gamma \in [\ell]$,

$$(5) \quad |K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma})| = (1 \pm \epsilon''_2(\ell))(n/\ell t)^3.$$

We will use \mathcal{P} to construct a different decomposition, of V , which will be called \mathcal{Q} , and which will have the same vertex partition as \mathcal{P} , but different edge partitions.

We begin defining several sets which will help us control the location of the error triads of \mathcal{P} . First, set

$$\begin{aligned} \Gamma_1 &= \{G_{ijs}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P}) : (H_{ijs}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \gamma}) \text{ has } \text{disc}_{2,3}(\epsilon''_1, \epsilon''_2(\ell)) \text{ and } d_{ijs}^{\alpha, \beta, \gamma} \geq 1 - \epsilon''_1\}, \\ \Gamma_0 &= \{G_{ijs}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P}) : (H_{ijs}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \gamma}) \text{ has } \text{disc}_{2,3}(\epsilon''_1, \epsilon''_2(\ell)) \text{ and } d_{ijs}^{\alpha, \beta, \gamma} \leq \epsilon''_1\}, \text{ and} \\ \Gamma_{err} &= \text{Triads}(\mathcal{P}) \setminus (\Gamma_0 \cup \Gamma_1). \end{aligned}$$

By assumption, $\text{Triads}(\mathcal{P}) = \Gamma_{err} \sqcup \Gamma_1 \sqcup \Gamma_0$, and at most $\epsilon''_1 n^3$ triples $xyz \in \binom{V}{3}$ are in an element of Γ_{err} . By (5), this implies $|\Gamma_{err}| \leq \epsilon''_1 n^3 / ((n^3/t^3 \ell^3)(1 - \epsilon''_2(\ell))) \leq \epsilon'_1 t^3 \ell^3$. We now define the set of triples from \mathcal{P}_{vert} which have non-trivial intersection with Γ_{err} .

$$\Sigma_0 = \{V_i V_j V_s : ijs \in \binom{[t]}{3} \text{ and } |\{(\alpha, \beta, \gamma) \in [\ell]^3 : G_{ijk}^{\alpha, \beta, \gamma} \in \Gamma_{err}\}| \geq 2(\epsilon'_1)^{1/2} \ell^3\}.$$

From (5), we have that $|\Sigma_0| \leq (\epsilon'_1)^{1/2} t^3$. We next define the set of pairs $V_i V_j$ contained in a non-trivial number of triples from Σ_0 .

$$\Psi = \{V_i V_j : |\{s \in [t] : V_i V_j V_s \in \Sigma_0\}| \geq (\epsilon'_1)^{1/4} t\}.$$

Since $|\Sigma_0| \leq 2(\epsilon'_1)^{1/2}t^3$, we have that $|\Psi| \leq (\epsilon'_1)^{1/4}t^2$. We now define, for each $V_iV_j \notin \Psi$, an auxiliary edge-colored graph $(U_{ij} \cup W_{ij}, E_{ij}^1, E_{ij}^0, E_{ij}^2)$, where

$$\begin{aligned} W_{ij} &= \{P_{ij}^\alpha : \alpha \leq \ell\} \text{ and } U_{ij} = \{P_{is}^\beta P_{js}^\gamma : s \in [t] \setminus \{i, j\}, \alpha, \beta \in [\ell]\}, \text{ and} \\ E_{ij}^1 &= \{P_{ij}^\alpha (P_{is}^\beta P_{js}^\gamma) \in K_2[W_{ij}, U_{ij}] : G_{ijk}^{\alpha, \beta, \gamma} \in \Gamma_1 \text{ and } V_iV_jV_s \notin \Sigma_0\}, \\ E_{ij}^0 &= \{P_{ij}^\alpha (P_{is}^\beta P_{js}^\gamma) \in K_2[W_{ij}, U_{ij}] : G_{ijk}^{\alpha, \beta, \gamma} \in \Gamma_0 \text{ and } V_iV_jV_s \notin \Sigma_0\}, \\ E_{ij}^2 &= K_2[W_{ij}, U_{ij}] \setminus (E_{ij}^1 \cup E_{ij}^0). \end{aligned}$$

Define $H_{ij} = (U_{ij} \cup W_{ij}, E_{ij}^1)$. Note H_{ij} is a bipartite graph with parts W_{ij}, U_{ij} , where $|W_{ij}| = \ell$ and $|U_{ij}| = (t-2)\ell^2$. Our next step is to prove Claim 6.18 below, which shows that for all $V_iV_j \notin \Psi$, there is a large subset $U_{ij}^* \cup W_{ij}^*$ of $V(H_{ij})$, on which H_{ij} is uniformly δ -close to some H_{ij}^* which contains no d -trees with leaves in W_{ij}^* . The idea behind the proof is as follows. First, by Proposition 6.7, we know there are no encodings of $H(k)$ in (H, \mathcal{P}) . Using Theorem 5.12, we will show that this implies that any (U_{ij}, W_{ij}, d) -tree in H_{ij} must involve some triad $G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_{err}$ where $V_iV_jV_s \notin \Sigma_0$. If $V_iV_j \notin \Psi$, then there are few such triads, so there will be few such (U_{ij}, W_{ij}, d) -trees in H_{ij} . We then apply Theorem 6.16 to obtain the desired H_{ij}^* .

Claim 6.18. *Suppose $V_iV_j \notin \Psi$. Then there are $U_{ij}^{bad} \subseteq U_{ij}$ and $W_{ij}^{bad} \subseteq W_{ij}$ such that the following hold, where $U_{ij}^* = U_{ij} \setminus U_{ij}^{bad}$ and $W_{ij}^* = W_{ij} \setminus W_{ij}^{bad}$.*

- $|W_{ij}^{bad}| \leq \epsilon_0|W_{ij}|$ and $|U_{ij}^{bad}| \leq \delta|U_{ij}|$,
- $H_{ij}[U_{ij}^*, W_{ij}^*]$ is uniformly $(U_{ij}^*, W_{ij}^*, \delta)$ -close to a bipartite graph $H_{ij}^* = (U_{ij}^* \cup W_{ij}^*, E_{ij}^*)$ containing no (U_{ij}^*, W_{ij}^*, d) -trees.

PROOF OF CLAIM 6.18: Fix $V_iV_j \notin \Psi$. We first show there are at most $(\epsilon'_1)^{1/4}|U_{ij}|^{2^{<d}}|W_{ij}|^{2^d}$ many (W_{ij}, U_{ij}, d) -trees in H_{ij} . Define $\Omega_{ij} = \{G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_{err} : V_iV_jV_s \notin \Sigma_0\}$. Note that if V_s satisfies $V_iV_jV_s \notin \Sigma_0$, then by definition of Σ_0 , $|\{(\alpha, \beta, \gamma) \in [\ell]^3 : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_{err}\}| \leq 2(\epsilon'_1)^{1/2}\ell^3$. Therefore, since $V_iV_j \notin \Psi$,

$$|\Omega_{ij}| \leq (\epsilon'_1)^{1/4}t\ell^3 + 2(\epsilon'_1)^{1/2}t\ell^3 \leq 3(\epsilon'_1)^{1/4}t\ell^3.$$

We now make some observations about any (U_{ij}, W_{ij}, d) -tree in H_{ij} . Suppose

$$(P_{is_\sigma}^{\alpha_\sigma} P_{js_\sigma}^{\beta_\sigma})_{\sigma \in 2^{<d}} (P_{ij}^{\gamma_\rho})_{\rho \in 2^d}$$

is a (U_{ij}, W_{ij}, d) -tree in H_{ij} . Note that for all $\sigma \in 2^{<d}$, there is $\rho \in 2^d$ so that $\sigma \wedge 1 \trianglelefteq \rho$, and for this ρ , $(P_{is_\sigma}^{\alpha_\sigma} P_{js_\sigma}^{\beta_\sigma}) P_{ij}^{\gamma_\rho} \in E_{ij}^1$. By definition of E_{ij}^1 , this implies $V_iV_jV_{s_\sigma} \notin \Sigma_0$. Thus for all $\sigma \in 2^{<d}$, $V_iV_jV_{s_\sigma} \notin \Sigma_0$.

By part (1) of Theorem 5.12, there are $\{P_{ij}^{\gamma'_u} : u \leq k\} \subseteq \{P_{ij}^{\gamma_\rho} : \rho \in 2^{d+1}\}$ and $\{P_{is'_u}^{\alpha'_u} P_{js'_u}^{\beta'_u}\} \subseteq \{P_{is_\sigma}^{\alpha_\sigma} P_{js_\sigma}^{\beta_\sigma} : \sigma \in 2^{<d}\}$ so that $P_{is'_u}^{\alpha'_u} P_{js'_u}^{\beta'_u} P_{ij}^{\gamma'_v} \in E_{ij}^1$ if and only if $u \leq v$. Translating, this means that for all $u \leq v$, $G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \in \Gamma_1$ and for all $v < u$, $G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \notin \Gamma_1$ (i.e. $G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \in \Gamma_{err} \cup \Gamma_0$). By above, for all $u \in [k]$, $V_iV_jV_{s'_u} \notin \Sigma_0$, so for all $v < u$, we have that

$G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \in \Omega_{ij} \cup \Gamma_0$. If it were true that $G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \in \Gamma_0$ for all $v < u$, then we would have an encoding of $H(k)$ in (H, \mathcal{P}) via $f : a_u \mapsto P_{ij}^{\gamma'_u}$ and $g : b_v \mapsto P_{is'_v}^{\alpha'_v} P_{js'_v}^{\beta'_v}$. By Proposition 6.7, this would imply H has k -FOP₂, a contradiction. Thus there is some $v < u$ so that $G_{ijs'_u}^{\alpha'_u \beta'_u \gamma'_v} \in \Omega_{ij}$.

The argument above shows that for every (U_{ij}, W_{ij}, d) -tree $(P_{is_\sigma}^{\alpha_\sigma} P_{js_\sigma}^{\beta_\sigma})_{\sigma \in 2^{<d}} (P_{ij}^{\gamma_\rho})_{\rho \in 2^d}$ in H_{ij} , there is some $\sigma \in 2^{<d}$ and $\rho \in 2^d$ such that $P_{ijs_\sigma}^{\alpha_\sigma, \beta_\sigma, \gamma_\rho} \in \Omega_{ij}$. Thus, any (U_{ij}, W_{ij}, d) -tree in H_{ij} will be constructed in the following process.

- (1) Choose some $\sigma \in 2^{<d}$ and $\rho \in 2^d$. There are $(2^d - 1)2^d$ choices.
- (2) Choose an element $G_{ijs}^{\alpha, \beta, \gamma} \in \Omega_{ij}$, and set $P_{is_\sigma}^{\alpha_\sigma} P_{js_\sigma}^{\beta_\sigma} = P_{is}^\alpha P_{js}^\beta$ and $P_{ij}^{\gamma_\rho} = P_{ij}^\gamma$. There are at most $|\Omega_{ij}| \leq 2(\epsilon'_1)^{1/2} t \ell^3$ ways to do this.
- (3) For all $\sigma' \in 2^{<d} \setminus \{\sigma\}$ and $\rho' \in 2^d \setminus \{\rho\}$, choose any $P_{is_{\sigma'}}^{\alpha_{\sigma'}} P_{js_{\sigma'}}^{\beta_{\sigma'}} \in U_{ij}$ and any $P_{ij}^{\gamma_{\rho'}} \in W_{ij}$. Then number of choices is at most

$$|U_{ij}|^{2^d-2} |W_{ij}|^{2^d-1} = ((t-2)\ell^2)^{2^d-2} \ell^{2^d-1}.$$

This yields that the number of (U_{ij}, W_{ij}, d) -trees in H_{ij} is at most

$$(2^d - 1)2^d 2(\epsilon'_1)^{1/2} t \ell^3 ((t-2)\ell^2)^{2^d-2} \ell^{2^d-1} < (\epsilon'_1)^{1/4} |U_{ij}|^{2^d-1} |W_{ij}|^{2^d},$$

where the inequality is due to our choice of ϵ'_1 . Now, since $\epsilon'_1 \ll \delta \ll \delta_0$, Theorem 6.13 implies that there is some $W_{ij}^{bad} \subseteq W_{ij}$ of size at most $\epsilon_0 |W_{ij}|$, and $U_{ij}^{bad} \subseteq U_{ij}$ of size at most $\delta |U_{ij}|$ such that, setting $U_{ij}^* = U_{ij} \setminus U_{ij}^{bad}$ and $W_{ij}^* = W_{ij} \setminus W_{ij}^{bad}$, $H_{ij}[U_{ij}^*, W_{ij}^*]$ is uniformly $(U_{ij}^*, W_{ij}^*, \delta)$ -close to a bipartite graph $H_{ij}^* = (U_{ij}^* \cup W_{ij}^*, E_{ij}^*)$ containing no (U_{ij}^*, W_{ij}^*, d) -trees. This finishes the proof of Claim 6.18. \square

Our next goal is to apply Theorem 6.16. Set $X = \min\{|W_{ij}^*| : V_i V_j \notin \Psi\}$. By Claim 6.18, $X \geq (1 - \epsilon_0)\ell$. For each $V_i V_j \notin \Psi$, let W_{ij}^{**} be any X -element subset of W_{ij}^* . Note that $H_{ij}^*[U_{ij}^*, W_{ij}^{**}]$ contains no $(U_{ij}^*, W_{ij}^{**}, d)$ -trees. Set $W = \bigcup_{V_i V_j \notin \Psi} W_{ij}^{**}$ and $U = \bigcup_{V_i V_j \in \Psi} U_{ij}^*$, and define

$$G^* = (U \cup W, \bigcup_{V_i V_j \notin \Psi} E(H_{ij}^*[U \cup W])).$$

Note G^* comes equipped with an equipartition $\mathcal{S} = \{W_{ij}^{**} : V_i V_j \notin \Psi\}$ of W such that for all $W_{ij}^{**} \in \mathcal{S}$, G^* contains no (U, W_{ij}^{**}, d) -trees. Consequently, by Theorem 5.15, there are $1 \leq m_1 \leq M$, $L_1 \leq X$, and a set $\Omega \subseteq \mathcal{S}$ such that $|\bigcup_{W_{ij}^{**} \in \Omega} W_{ij}^{**}| \leq \epsilon^{100} |W|$, and for all $W_{ij}^{**} \in \mathcal{S} \setminus \Omega$, there is a partition

$$W_{ij}^{**} = B_{ij}^0 \cup B_{ij}^1 \cup \dots \cup B_{ij}^{m_1},$$

so that $|B_{ij}^0| \leq \epsilon^{100} |W_{ij}^{**}|$, and for each $u \in [m_1]$, $|B_{ij}^u| = L_1$ and B_{ij}^u is $f(m_1)$ -good in G^* . Set

$$\Psi' := \Psi \cup \{V_i V_j \in \binom{\mathcal{P}_{vert}}{2} \setminus \Psi : W_{ij}^{**} \in \Omega\}.$$

Note $|\Psi'| \leq (\epsilon'_1)^{1/4} t^2 + \epsilon^{100} t^2 \leq 2\epsilon^{100} t^2$. Set $m = \lceil \epsilon^{-100} m_1(m_1 + 1) \rceil$. By construction, $(1 - \epsilon_0) \frac{\ell}{m_1} \leq L_1 \leq \frac{\ell}{m_1}$, so since $m_1 \leq M$, our choice of ϵ_0 implies $L_1 > \frac{\ell}{m}$. Let r be such that $L_1 = rm + c$ for some $c < m$. Then for all $V_i V_j \notin \Psi'$, and each $u \in [m_1]$, we can choose a partition $B_{ij}^u = B_{ij}^{u,0} \cup B_{ij}^{u,1} \cup \dots \cup B_{ij}^{u,r}$ so that $|B_{ij}^{u,0}| = c < \frac{\ell}{m}$ and for all $1 \leq v \leq r$, $|B_{ij}^{u,v}| = \frac{\ell}{m}$ (recall that ℓ is divisible by $(M!)^3$, so we know that $\ell/m \in \mathbb{N}$). Define

$$R_{ij}^0 = B_{ij}^0 \cup \bigcup_{v=1}^{m_1} B_{ij}^{u,0}.$$

Note $|R_{ij}^0| \leq \epsilon^{100} \ell + \frac{m_1 \ell}{m} < 2\epsilon^{100} \ell \leq \epsilon^{75} \ell$. We claim that for all $W_{ij}^{**} \in \mathcal{S} \setminus \Omega$ and all $u \in [m_1]$, and $1 \leq v \leq r$, $B_{ij}^{u,v}$ is $\psi(m)$ -good in H_{ij} with respect to every $u \in U_{ij}^*$. Fix $W_{ij}^{**} \in \mathcal{S} \setminus \Omega$, $u \in [m_1]$, $1 \leq v \leq r$, and $u \in U_{ij}^*$. We know that B_{ij}^u is $f(m_1)$ -good in G^* , so there is some $\tau \in \{0, 1\}$ so that

$$|N_{G^*}^\tau(u) \cap B_{ij}^u| = |N_{H_{ij}^*}^\tau(u) \cap B_{ij}^u| \leq f(m_1) |B_{ij}^u|.$$

Then since H_{ij}^* is uniformly δ -close to $H_{ij}[U_{ij}^*, W_{ij}^*]$,

$$\begin{aligned} |N_{H_{ij}}^\tau(u) \cap B_{ij}^{u,v}| &\leq f(m_1) |B_{ij}^u| + \delta |W_{ij}^*| = |B_{ij}^{u,v}| \left(\frac{f(m_1) |B_{ij}^u|}{|B_{ij}^{u,v}|} + \frac{\delta |W_{ij}^*|}{|B_{ij}^{u,v}|} \right) \\ &\leq |B_{ij}^{u,v}| (f(m) m + \delta m) \\ &\leq |B_{ij}^{u,v}| (f(m_1) + \delta) (2\epsilon^{-100} m_1(m_1 + 1)) \\ &\leq \psi(m) |B_{ij}^{u,v}|, \end{aligned}$$

where the second inequality is by definition of m , since $|B_{ij}^u| = L_1$, and since $|B_{ij}^{u,v}| = \ell/m$, and the last inequality is by definition of f , since $m \leq 2\epsilon^{-100} m_1(m_1 + 1)$, and our choice of δ . We point out to the reader that it is this step (namely deducing goodness of the $B_{ij}^{u,v}$ in H_{ij} from the goodness of W_{ij}^* in H_{ij}^*) where it is important to have the degree of goodness in Theorem 6.16 be small as a function of the number of parts. Now, choose a reindexing

$$\{R_{ij}^1, \dots, R_{ij}^{m_2}\} = \{B_{ij}^{u,v} : u \in [m_1], v \in [r]\},$$

where $m_2 = m_1 r$. Note $m_1 r \leq \epsilon^{-100} m_1(m_1 + 1) < (M!)^3$, so ℓ is divisible by m_2 .

We have now defined, for all $V_i V_j \notin \Psi'$, a partition

$$W_{ij} = W_{ij}^{bad} \cup R_{ij}^0 \cup R_{ij}^1 \cup \dots \cup R_{ij}^{m_2}.$$

This naturally translates into a new partition of $K_2[V_i, V_j]$. In particular, for each $1 \leq u \leq m_2$, we let

$$\mathbf{R}_{ij}^u = \{xy \in K_2[V_i, V_j] : xy \in P_{ij}^\alpha \text{ for some } P_{ij}^\alpha \in R_{ij}^u\}.$$

Setting $\mathbf{R}_{ij}^0 = K_2[V_i, V_j] \setminus \bigcup_{u=1}^{m_2} \mathbf{R}_{ij}^u$, we have a partition $K_2[V_i, V_j] = \mathbf{R}_{ij}^0 \cup \mathbf{R}_{ij}^1 \cup \dots \cup \mathbf{R}_{ij}^{m_2}$. By construction, for each $u \in [m_2]$, \mathbf{R}_{ij}^u is a union of ℓ/m many $P_{ij}^\alpha \in \mathcal{P}_{edge}$, so by Fact 3.23, \mathbf{R}_{ij}^u has $\text{disc}_2((\ell/m)\epsilon_2'''(\ell), (1/\ell)(\ell/m))$. By our choice of $\epsilon_2'''(\ell)$, we can conclude \mathbf{R}_{ij}^u has

$\text{disc}_2(\epsilon_2''(1/m), 1/m)$. By (5), Claim 6.18, and since $\epsilon_0 \ll \epsilon$, we have the following bound on the size of \mathbf{R}_{ij}^0 .

$$|\mathbf{R}_{ij}^0| \leq (1 + \epsilon_2''(\ell)) \frac{n^2}{t^2 \ell} (|R_{ij}^0| + |W_{ij}^{bad}|) \leq (1 + \epsilon_2''(\ell)) \frac{n^2}{t^2} (\epsilon^{75} + \epsilon_0) \leq \epsilon^{50} \frac{n^2}{t^2}.$$

Given $ijs \in \binom{[t]}{3}$ with $V_i V_j, V_j V_s, V_i V_s \notin \Psi'$, and $(u, v, w) \in [m_2]^3$, we define

$$\mathbf{R}_{ijs}^{uvw} := (V_i \cup V_j \cup V_s, \mathbf{R}_{ij}^u \cup \mathbf{R}_{js}^v \cup \mathbf{R}_{is}^w).$$

By Corollary 3.6, for each such \mathbf{R}_{ijs}^{uvw} ,

$$(6) \quad |K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})| = (1 \pm \epsilon_2'(m)) \frac{n^3}{t^3 m^3}.$$

Our next goal is to show most of these \mathbf{R}_{ijs}^{uvw} have density near 0 or 1 with respect to H . In order to do this, we will need a few more sets to help us avoid the error triads. Set

$$\Sigma_1 = \Sigma_0 \cup \{V_i V_j V_s : ijs \in \binom{[t]}{3} \text{ and } \binom{\{V_i, V_j, V_s\}}{2} \cap \Psi' \neq \emptyset\}.$$

By the bounds given above, $|\Sigma_1| \leq (\epsilon_1')^{1/2} t^3 + 2\epsilon^{100} t^2 \leq 3\epsilon^{100} t^3$. Given $ijs \in \binom{[t]}{3}$, we define

$$U_{ij}(s) := \{P_{is}^\beta P_{js}^\gamma : \beta, \gamma \in [\ell]\}.$$

Note that for all $ij \in \binom{[t]}{2}$, $U_{ij} = \bigcup_{s \in [t]} U_{ij}(s)$. Then define

$$\Sigma_3 = \{V_i V_j V_s \in \binom{\mathcal{P}_{vert}}{3} \setminus \Sigma_1 : \max\{|U_{ij}(s) \cap U_{ij}^{bad}|, |U_{js}(i) \cap U_{js}^{bad}|, |U_{is}(j) \cap U_{is}^{bad}|\} \geq \delta^{1/2} \ell^2\}.$$

Observe that for all $V_i V_j \notin \Psi'$, the number of V_s with $V_i V_j V_s \in \Sigma_3$ is at most $\delta^{1/2} t$. Consequently, $|\Sigma_3| \leq \delta t^3$. We now set $\Sigma = \Sigma_1 \cup \Sigma_3$, and note that $|\Sigma| \leq 3\epsilon^{100} t^3 + \delta t^3 \leq 4\epsilon^{100} t^3$. This will end up being the desired Σ at the end of the proof. We will show that if $V_i V_j V_s \notin \Sigma$, then the density of edges on $K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})$ is close to 0 or 1 for all $u, v, w \in [m_2]$ (see Claim 6.19 below).

First, we need to show a couple things about triples outside Σ . Given $V_i V_j V_s \notin \Sigma$, and $u \in [m_2]$, define

$$U_{ij}^1(s) = \{P_{is}^\beta P_{js}^\gamma \in U_{ij}(s) : |\{P_{ij}^\alpha \in R_{ij}^u : G_{ijs}^{\alpha\beta\gamma} \in \Gamma_1\}| \geq (1 - \psi(m)) |R_{ij}^u|\}, \text{ and} \\ U_{ij}^0(s) = \{P_{is}^\beta P_{js}^\gamma \in U_{ij}(s) : |\{P_{ij}^\alpha \in R_{ij}^u : G_{ijs}^{\alpha\beta\gamma} \notin \Gamma_1\}| \geq (1 - \psi(m)) |R_{ij}^u|\}.$$

By Claim 6.18, we know that $U_{ij}(s) = U_{ij}^{bad} \cup U_{ij}^1(s) \cup U_{ij}^0(s)$. However, we are more interested in the following set, rather than $U_{ij}^0(s)$.

$$U_{ij}^{00}(s) = \{P_{is}^\beta P_{js}^\gamma \in U_{ij}(s) : |\{P_{ij}^\alpha \in R_{ij}^u : (P_{is}^\beta P_{js}^\gamma) P_{ij}^\alpha \in E_{ij}^2\}| \geq (1 - \psi(m)^{1/2}) |R_{ij}^u|\}.$$

Setting $U_{ij}^{err}(s) = U_{ij}(s) \setminus (U_{ij}^1(s) \cup U_{ij}^{00}(s))$, we wish to show that $|U_{ij}^{err}(s)| \leq \epsilon^9 |U_{ij}(s)|$. Observe that

$$|K_2[U_{ij}(s), W_{ij}] \cap E_{ij}^2| \geq (\psi(m)^{1/2} - \psi(m)) |U_{ij}^0(s) \setminus U_{ij}^{00}(s)| |R_{ij}^u| \geq \psi(m) |U_{ij}^0(s) \setminus U_{ij}^{00}(s)| \ell / m,$$

where the last inequality uses the fact that $|R_{ij}^u| = \ell/m$ and $\psi(m) < \epsilon < 1/4$. Because $ijs \notin \Sigma_1$, $|K_2[U_{ij}(s), W_{ij}] \cap E_{ij}^2| \leq 2(\epsilon'_1)^{1/2}\ell^3$. Consequently, this implies

$$|U_{ij}^0(s) \setminus U_{ij}^{00}(s)| \leq 2(\epsilon'_1)^{1/2}\ell^3\psi(m)^{-1}\frac{m}{\ell} = 2(\epsilon'_1)^{1/2}m\psi(m)^{-1}\ell^2 \leq (\epsilon'_1)^{1/4}\ell^2,$$

where the last inequality is because $\epsilon'_1 \ll \psi((M!)^3)/(M!)^3$. Further, since $ijs \notin \Sigma_3$, $|U_{ij}(s) \cap U_{ij}^{bad}| \leq \delta^{1/2}\ell^2$, so

$$\begin{aligned} |U_{ij}^1(s) \cup U_{ij}^{00}(s)| &\geq |U_{ij}(s)| - |U_{ij}^0(s) \setminus U_{ij}^{00}(s)| - |U_{ij}^{bad}(s)| \geq (1 - \delta^{1/2})\ell^2 - (\epsilon'_1)^{1/4}\ell^2 \\ &\geq \ell^2(1 - \epsilon^{50}), \end{aligned}$$

where the last inequality is because $\epsilon'_1 \ll \delta \ll \epsilon$. This shows $|U_{ij}^{err}(s)| \leq \epsilon^{50}|U_{ij}(s)|$. We will use this below.

Claim 6.19. *For all $ijs \in \binom{[t]}{3}$ with $V_iV_jV_s \notin \Sigma$, and all $(u, v, w) \in [m_2]^3$,*

$$\frac{|E \cap K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|}{|K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|} \in [0, \epsilon] \cup (1 - \epsilon, 1].$$

Proof. Fix $ijs \in \binom{[t]}{3}$ with $V_iV_jV_s \notin \Sigma$, and $(u, v, w) \in [m_2]^3$. Suppose towards a contradiction that

$$(7) \quad \frac{|E \cap K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|}{|K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|} \in [\epsilon, 1 - \epsilon].$$

To ease notation, define $A = R_{is}^w$, $B = R_{js}^v$, $C = R_{ij}^u$, and let

$$\mathbf{F}_0 = K_2[A, B] \cap U_{ij}^{00}(s) \text{ and } \mathbf{F}_1 = K_2[A, B] \cap U_{ij}^1(s).$$

Then let $\mathbf{F}_{err} = K_2[A, B] \cap U_{ij}^{err}(s)$. Note $|A| = |B| = |C| = \ell/m$, and our observations above show that $|\mathbf{F}_{err}| \leq \epsilon^{50}|A||B|$. We claim that

$$(8) \quad \min\{|\mathbf{F}_1|, |K_2[A, B] \setminus \mathbf{F}_1|\} \geq \epsilon^3|A||B|.$$

Suppose towards a contradiction that (8) is false. Then either $|\mathbf{F}_1| < \epsilon^3|A||B|$ or $|K_2[A, B] \setminus \mathbf{F}_1| < \epsilon^3|A||B|$. Assume first that $|\mathbf{F}_1| < \epsilon^3|A||B|$. Combining this with the above, we have that

$$|K_2[A, B] \setminus \mathbf{F}_0| \leq (\epsilon^9 + \epsilon^3)|A||B| \leq 2\epsilon^3|A||B|.$$

Therefore, using (5),

$$\sum_{\{P_{is}^\alpha P_{js}^\beta \in K_2[A, B] \setminus \mathbf{F}_0\}} \sum_{P_{ij}^\gamma \in C} |K_3(G_{ijk}^{\alpha, \beta, \gamma})| \leq (1 + \epsilon_2''(\ell)) \frac{2\epsilon^3 n^3}{t^3 \ell^3} |A||B||C| \leq 3\epsilon^3 \frac{n^3}{t^3 m^3}.$$

Note that if $P_{is}^\alpha P_{js}^\beta \in \mathbf{F}_0$, then at least a $(1 - \psi(m)^{1/2})|C|$ many elements in $P_{ij}^\gamma \in C$ are such that $G_{ijs}^{\alpha,\beta,\gamma} \in \Gamma_0$. Consequently,

$$\begin{aligned} \sum_{P_{ij}^\gamma \in C} |E \cap K_3(G_{ijk}^{\alpha,\beta,\gamma})| &\leq \sum_{\{P_{ij}^\gamma: G_{ijs}^{\alpha,\beta,\gamma} \in \Gamma_0\}} \epsilon_1'' |K_3(G_{ijk}^{\alpha,\beta,\gamma})| + \sum_{\{P_{ij}^\gamma: G_{ijs}^{\alpha,\beta,\gamma} \notin \Gamma_0\}} |K_3(G_{ijk}^{\alpha,\beta,\gamma})| \\ &\leq (1 + \epsilon_2(\ell)) \frac{n^3}{t^3 \ell^3} (\epsilon_1'' |C| (1 - \psi(m)^{1/2}) + \psi(m)^{1/2} |C|) \\ &\leq \epsilon^5 \frac{n^3}{t^3 m \ell^2}, \end{aligned}$$

where the last inequality uses that $|C| = \ell/m$ and that $\psi(m), \epsilon_1'(\ell), \epsilon_1'' \ll \epsilon$. So we have that

$$|E \cap K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})| \leq 2\epsilon^3 \frac{n^3}{t^3 m^3} + \epsilon^5 |A||B| \frac{n^3}{t^3 m \ell^2} \leq 3\epsilon^3 \left(\frac{n}{t}\right)^3 \frac{1}{m^3}.$$

By (6), this shows that $|E \cap K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})| < \epsilon |K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|$, contradicting (7). Thus we must have that $|K_2[A, B] \setminus \mathbf{F}_1| < \epsilon^3 K_2[A, B]$. In this case,

$$|\mathbf{F}_0 \cup \mathbf{F}_{err}| \leq \epsilon^3 |A||B| = \epsilon^3 \frac{\ell^2}{m^2}.$$

Consequently, using (5),

$$\sum_{\{P_{is}^\alpha P_{js}^\beta \in \mathbf{F}_0 \cup \mathbf{F}_{err}\}} \sum_{P_{ij}^\gamma \in C} |K_3(G_{ijk}^{\alpha,\beta,\gamma})| \leq (1 + \epsilon_2''(\ell)) \frac{\epsilon^3 n^3}{t^2 m^3} \leq \frac{2\epsilon^3 n^3}{t^3 m^3}.$$

On the other hand, for each $P_{is}^\alpha P_{js}^\beta \in \mathbf{F}_1$, we know that at least $(1 - \psi(m))|C|$ many elements in $P_{ij}^\gamma \in C$ are such that $G_{isj}^{\alpha,\beta,\gamma} \in \Gamma_1$. Consequently,

$$\begin{aligned} \sum_{\{P_{is}^\alpha P_{js}^\beta \in \mathbf{F}_0\}} \sum_{P_{ij}^\gamma \in C} |K_3(G_{ijk}^{\alpha,\beta,\gamma}) \setminus E| &\leq (1 + \epsilon_2''(\ell)) \left(\frac{n}{t\ell}\right)^3 |A||B| (\psi(m)|C| + (1 - \psi(m))\epsilon_1'' |C|) \\ &\leq \epsilon^5 \left(\frac{n}{t}\right)^3 \frac{1}{m^3}, \end{aligned}$$

where the last inequality uses that $|C| = \ell/m$ and that $\psi(m), \epsilon_1'(\ell), \epsilon_1'' \ll \epsilon$. Combining these, we obtain the that

$$|K_3^{(2)}(\mathbf{R}_{ijs}^{uvw}) \setminus E| \leq \frac{2\epsilon^3 n^3}{m^3} + \epsilon^5 \frac{n^3}{t^3 m^3} \leq 3\epsilon^3 \left(\frac{n}{t}\right)^3 \frac{1}{m^3}.$$

By (6), this shows that $|K_3^{(2)}(\mathbf{R}_{ijs}^{uvw}) \setminus E| < |K_3^{(2)}(\mathbf{R}_{ijs}^{uvw})|$, contradicting (7). Thus we must have that $|K_2[A, B] \setminus \mathbf{F}_1| \geq \epsilon^3 |A||B|$, and therefore (8) is true.

Consider the bipartite graph $G_{AB} = (A \cup B, \mathbf{F}_1)$. Since (8) is true, Lemma 5.9 implies one of the following holds.

- (1) There is a set $S \subseteq B$ of size at least $\epsilon^6 |B|/2$ such that for all $b \in S$

$$\min\{|N_{G_{AB}}(b)|, |\neg N_{G_{AB}}(b)|\} \geq \epsilon^6 |A|/2.$$

(2) There is a set $S \subseteq A$ of size at least $\epsilon^6|A|/2$ such that for all $a \in S$

$$\min\{|N_{G_{AB}}(a)|, |\neg N_{G_{AB}}(a)|\} \geq \epsilon^6|B|/2.$$

Let us assume (1) holds (case (2) is similar). So there is some $S \subseteq B$ of size at least $\epsilon^6|B|/2$ such that for all $b \in S$

$$\min\{|N_{G_{AB}}(b)|, |\neg N_{G_{AB}}(b)|\} \geq \epsilon^6|A|/2.$$

Recall we showed $|\mathbf{F}_{err}| \leq \epsilon^{50}|A||B|$, so if we let

$$S' = \{b \in S : |\{a \in A : ab \in \mathbf{F}_{err}\}| \leq \epsilon^{25}|A|\},$$

then $|S'| \geq |S| - \epsilon^{25}|B| \geq \epsilon^6|B|/2 - \epsilon^{25}|B| \geq \epsilon^6|B|/4$, where the inequality is because $\epsilon < 1/4$. For each $b \in S'$, set

$$A_0(b) = \{a \in A : ab \in \mathbf{F}_0\} \text{ and } A_1(b) = \{a \in A : ab \in \mathbf{F}_1\}.$$

By definition of S' , $|A_1(b)| \geq \epsilon^6|A|/2$, and $|A_0(b)| \geq \epsilon^6|A|/2 - \epsilon^{25}|A| \geq \epsilon^6|A|/4$. Now consider the bipartite graphs $G_0(b) = (A_0(b) \cup C, E_0(b))$ and $G_1(b) = (A_1(b) \cup C, E_1(b))$, where

$$E_0(b) = \{ca \in K_2[C, A_0(b)] : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_0 \text{ where } c = P_{ij}^\alpha, a = P_{is}^\beta, \text{ and } b = P_{js}^\gamma\}$$

$$E_1(b) = \{ca \in K_2[C, A_1(b)] : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_1 \text{ where } c = P_{ij}^\alpha, a = P_{is}^\beta, \text{ and } b = P_{js}^\gamma\}.$$

We claim that for all $b \in S'$ and $a \in A_0(b)$, $|N_{G_0(b)}(a)| \geq (1 - \psi(m)^{1/2})|C|$. Say $b = P_{js}^\gamma \in S'$ and $a = P_{is}^\beta \in A_0(b)$. Since $ab \in \mathbf{F}_0$, we know that $P_{js}^\gamma P_{is}^\beta \in W_{ij}^{00}(s)$. By definition, this means that

$$|N_{G_0(b)}(a)| = |\{c = P_{ij}^\alpha \in C : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_0\}| \geq (1 - \psi(m)^{1/2})|C|,$$

as desired. Consequently, $|E_0(b)| \geq (1 - \psi(m)^{1/2})|C||A_0(b)|$. Standard arguments show that this implies there is a set $C_0(b) \subseteq C$ with $|C_0(b)| \geq (1 - \psi(m)^{1/4})|C|$ so that for all $c \in C_0(b)$,

$$|N_{G_0(b)}(c)| \geq (1 - \psi(m)^{1/4})|A_0(b)| \geq (1 - \psi(m)^{1/4})\epsilon^6|A|/2 \geq \epsilon^6|A|/4.$$

A similar argument shows that for all $b \in S'$ and $a \in A_1(b)$, $|N_{G_1(b)}(a)| \geq (1 - \psi(m)^{1/2})|C|$, so $|E_1(b)| \geq (1 - \psi(m)^{1/4})|C||A_1(b)|$. Thus there exists a set $C_1(b) \subseteq C$ with $|C_1(b)| \geq (1 - \psi(m)^{1/4})|C|$ so that for all $c \in C_1(b)$,

$$|N_{G_1(b)}(c)| \geq (1 - \psi(m)^{1/4})|A_1(b)| \geq (1 - \psi(m)^{1/4})\epsilon^6|A|/2 \geq \epsilon^6|A|/4.$$

Combining these, we have that for all $b \in S'$, there is a set $C(b) = C_0(b) \cap C_1(b) \subseteq C$, with size $|C(b)| \geq (1 - 2\psi(m)^{1/4})|C|$, such that for all $c \in C(b)$,

$$\min\{|N_{G_0(b)}(c)|, |N_{G_1(b)}(c)|\} \geq \epsilon^6|A|/4.$$

We claim that for all $b \in S'$ and $c \in C(b)$, $bc \in U_{is}^{bad}$. To this end, fix $b \in S'$ and $c \in C(b)$. Then $b = P_{js}^\beta$ and $c = P_{ij}^\alpha$ for some $\beta, \alpha \in [\ell]$. By above and the definition of H_{is} ,

$$\begin{aligned} \epsilon^6 |A|/4 &\leq \min\{|N_{G_0(b)}(c)|, |N_{G_1(b)}(c)|\} \\ &= \min\{|P_{is}^\gamma \in A : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_0\}|, \{|P_{is}^\gamma \in A : G_{ijs}^{\alpha, \beta, \gamma} \in \Gamma_1\}|\} \\ &\leq \min\{|N_{H_{is}}(P_{js}^\beta P_{ij}^\alpha) \cap A|, |\neg N_{H_{is}}(P_{js}^\beta P_{ij}^\alpha) \cap A|\} \end{aligned}$$

This shows $A = R_{is}^w$ is not $\epsilon^6/4$ -good in H_{is} with respect to bc . Since $V_i V_j V_s \notin \Sigma$, we know that R_{is}^w is $\psi(m)$ -good with respect to every $u \in U_{is} \setminus U_{is}^{bad}$, so since $\psi(m) \ll \epsilon$, this implies that $bc \in U_{is}^{bad}$. Thus we have shown that for all $b \in S'$ and $c \in C(b)$, $bc \in U_{is}^{bad}$. Combining this with the bounds above, we have that

$$|U_{is}^{bad}| \geq |S'|(1 - 2\psi(m)^{1/4})|C| \geq \epsilon^6 |B||C|/4 = \epsilon^6 \ell^2 / 4m^2.$$

However, by Claim 6.18, $|U_{is}^{bad}| \leq \delta \ell^2 < \epsilon^6 \ell^2 / 4m^2$, where the last inequality is since $\delta \ll \epsilon/M$ and $m \leq M^3$. Thus, we have arrived at a contradiction, which finishes the proof of Claim 6.19. \square

We are nearly done. The last main task is to redistribute \mathbf{R}_{ij}^0 for each $V_i V_j \notin \Psi'$. Fix $V_i V_j \notin \Psi'$. Recall that $|W_{ij}| = \ell$, and $|R_{ij}^0| = |W_{ij}| - |B_{ij}^1 \cup \dots \cup B_{ij}^{m_2}| = \ell - \ell m_2 / m$. Since $m_2 \leq M$, and $(M!)^3$ divides ℓ , we know m_2 divides ℓ , say $\ell = m_3 m_2$. Note this implies $|R_{ij}^0| = m_2 m_3 - m_2 \ell / m$. Similarly, since $m \leq (M!)^3$, m also divides ℓ , so we can choose a partition $R_{ij}^0 = S_{ij}^1 \cup \dots \cup S_{ij}^{m_2}$ into sets of size $(m_2 m_3 - m_2 \ell / m) / m_2 = m_3 - \ell / m$. For each $u \in [m_2]$, define

$$\mathbf{S}_{ij}^u = \{xy \in K_2[V_i, V_j] : xy \in P_{ij}^\alpha \text{ for some } P_{ij}^\alpha \in S_{ij}^u\}.$$

By Fact 3.23, \mathbf{S}_{ij}^u has $\text{disc}_2(m\epsilon_2'''(\ell))$. By our choice of ϵ_2'' , this implies \mathbf{S}_{ij}^u has $\text{disc}_2(\epsilon_2'(\ell))$. Now set $\mathbf{P}_{ij}^u = \mathbf{R}_{ij}^u \cup \mathbf{S}_{ij}^u$ for each $u \in [m_2]$. Note each \mathbf{P}_{ij}^u is a union of $m_3 - \ell/m + \ell/m = m_3$ elements of \mathcal{P}_{edge} , so by Fact 3.23, \mathbf{P}_{ij}^u satisfies $\text{disc}_2(m\epsilon_2''(\ell))$ and has size $m_3 \frac{n^2}{t} (1 \pm m_3 \epsilon_2'(\ell))$. Combining this with our choice of ϵ_2'' and the fact that $m_3/\ell = 1/m_2$, we have that each \mathbf{P}_{ij}^u satisfies $\text{disc}_2(\epsilon_2(m_2), 1/m_2)$. By construction, we know that each $|S_{ij}^u| \leq |R_{ij}^0|/m_2 \leq \epsilon^{75} \ell / m_2$. Therefore, $|\mathbf{S}_{ij}^u| \leq \epsilon^{25} \frac{n^2}{t^2 m_2^2} \leq \epsilon^{10} |\mathbf{P}_{ij}^u|$.

Now for each $V_i V_j \in \Psi'$, choose an arbitrary partition $K_2[V_i, V_j] = \bigcup_{\alpha \leq m_2} \mathbf{P}_{ij}^\alpha$. Since $|\Psi'| \leq 3\epsilon^{100} t^2$, and every \mathbf{P}_{ij}^u satisfies $\text{disc}_2(\epsilon_2(m_2), 1/m_2)$, we have a $(t, m_2, \epsilon^{50}, \epsilon_2(m_2))$ -decomposition \mathcal{Q} with $\mathcal{Q}_{vert} = \{V_1, \dots, V_t\}$ and

$$\mathcal{Q}_{edge} = \{\mathbf{P}_{ij}^u : ij \in \binom{[t]}{2}, u \in \{1, \dots, m_2\}\}.$$

We claim that for all $V_i V_j V_s \notin \Sigma$ and $\alpha, \beta, \gamma \in [m_2]$, $\mathbf{P}_{ijs}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_s, \mathbf{P}_{ij}^\alpha \cup \mathbf{P}_{is}^\beta \cup \mathbf{P}_{js}^\gamma)$ is 2ϵ -homogeneous with respect to H .

Fix $V_i V_j V_s \notin \Sigma$, and $\alpha, \beta, \gamma \in [m_2]$. By construction, there are $u, v, w \in [m_2]$ so that

$$\mathbf{P}_{ij}^\alpha = \mathbf{R}_{ij}^u \cup \mathbf{S}_{ij}^u, \mathbf{P}_{is}^\beta = \mathbf{R}_{is}^v \cup \mathbf{S}_{is}^v, \text{ and } \mathbf{P}_{js}^\gamma = \mathbf{R}_{js}^w \cup \mathbf{S}_{js}^w.$$

By Claim 6.19, there is $\tau \in \{0, 1\}$ such that $|E^\tau \cap K_3^{(2)}(\mathbf{R}_{ijs}^{u,v,w})| \leq \epsilon \frac{n^3}{t^3 m_2^3}$. Define

$$\begin{aligned} T_{is} &= (V_i \cup V_j \cup V_s, \mathbf{P}_{ij}^\alpha \cup \mathbf{S}_{is}^v \cup \mathbf{P}_{is}^\gamma) \\ T_{ij} &= (V_i \cup V_j \cup V_s, \mathbf{S}_{ij}^u \cup \mathbf{P}_{is}^\beta \cup \mathbf{P}_{is}), \text{ and} \\ T_{js} &= (V_i \cup V_j \cup V_s, \mathbf{P}_{ij}^\alpha \cup \mathbf{P}_{is}^\beta \cup \mathbf{S}_{js}^w). \end{aligned}$$

By Corollary 3.6, and the bound above on the size of each $\mathbf{S}_{ij}^u, \mathbf{S}_{is}^v, \mathbf{S}_{js}^w$, we have that

$$\max\{|K_3^{(2)}(T_{ij})|, |K_3^{(2)}(T_{ij})|, |K_3^{(2)}(T_{ij})|\} \leq \epsilon^{10} \frac{n^3}{t^3 m_2^3} \leq \epsilon^5 |K_3^{(2)}(\mathbf{P}_{ijs}^{\alpha,\beta,\gamma})|,$$

where the last inequality uses Corollary 3.6. Thus we have that

$$|E^\tau \cap K_3^{(2)}(\mathbf{P}_{ijs}^{\alpha,\beta,\gamma})| \leq \epsilon \frac{n^3}{t^3 m_2^3} + \epsilon^{10} |K_3^{(2)}(\mathbf{P}_{ijs}^{\alpha,\beta,\gamma})| \leq 2\epsilon |K_3^{(2)}(\mathbf{P}_{ijs}^{\alpha,\beta,\gamma})|.$$

Since $\epsilon \ll \epsilon_1$, this implies that $\mathbf{P}_{ijs}^{\alpha,\beta,\gamma}$ is ϵ_1 -homogeneous with respect to H and moreover, by Proposition 3.22, satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(m_2))$ with respect to H . This finishes the proof of Theorem 2.49. \square

We now prove Theorem 2.50, which says a hereditary 3-graph property admits linear $\text{disc}_{2,3}$ -error if and only if it is NFOP_2 .

PROOF OF THEOREM 2.50: Suppose \mathcal{H} is an NFOP_2 , hereditary 3-graph property. That \mathcal{H} admits linear $\text{disc}_{2,3}$ -error follows immediately from Theorem 2.49.

Conversely, suppose that \mathcal{H} has FOP_2 . We show that it requires non-linear $\text{disc}_{2,3}$ -error. Choose $0 < \epsilon'_1 \ll \epsilon_1 \ll 1$, and choose $\epsilon_2, \epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ so that for all $x \in \mathbb{N}$, $0 < \epsilon'_2(x) \ll \epsilon_2(x) \leq 1/4x$. Fix $T, L, N \geq 1$. We show that there is some $H \in \mathcal{H}$ with $|V(H)| \geq N$ so that there exists no $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition \mathcal{P} of $V(H)$ which is $\text{disc}_{2,3}$ -regular with respect to H with linear error.

Choose $m \gg L, T, N, (\epsilon'_1)^{-1}, (\epsilon'_2(L))^{-1}$ and $n \gg m$. By Lemma 3.24, there exist sets $B = \{b_i : i \in [n]\}$ and $C = \{c_i : i \in [n]\}$ and $\gamma_1, \dots, \gamma_m \subseteq K_2[B, C]$ so that for each $i \in [m]$, γ_i has $\text{disc}_2(\epsilon_2(m)/T^2, 1/m)$. Let $A = \{a_i : i \in [n]\}$ be a set of size n .

By Lemma 6.8, there exists $H = (V, E) \in \mathcal{H}$ with $V = A \cup B \cup C$, such that for all $i, j, s \in [n]$, $a_i b_j c_s \in E$ if and only if $b_j c_s \in \gamma_u$ for some $u \geq \lfloor in/\ell \rfloor$. We show that there does not exist $1 \leq \ell \leq L$, $1 \leq t \leq T$ and a $(t, \ell, \epsilon'_1, \epsilon'_2)$ -decomposition of V which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular with respect to H with linear $\text{disc}_{2,3}$ -error.

Suppose towards a contradiction that there exists some $1 \leq \ell \leq L$, $1 \leq t \leq T$ and a $(t, \ell, \epsilon'_1, \epsilon'_2)$ -decomposition \mathcal{P} of V which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular with respect to H , and a set $\Sigma \subseteq \binom{[t]}{3}$ with $|\Sigma| \leq \epsilon'_1 t^3$ so that if $G \in \text{Triads}(\mathcal{P})$ fails $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$ with respect to (H, \mathcal{P}) , then $V(G) \subseteq V_i \cup V_j \cup V_k$ for some $ijk \in \Sigma$. Say $\mathcal{P}_{\text{vert}} = \{V_i : i \in [t]\}$ and $\mathcal{P}_{\text{edge}} = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$.

For each $i \in [T]$, and $X \in \{A, B, C\}$, set $X_i = X \cap V_i$. For each $X \neq Y$, $i \neq j$, and $\alpha \in [\ell]$ we let $P_{X_i Y_j}^\alpha = P_{ij}^\alpha \cap K_2[X_i, Y_j]$.

Given $i \in [t]$ and $X \in \{A, B, C\}$, we say that X_i is *trivial* if $|X_i| < \epsilon_1 |V_i|$, and that a vertex $x \in V$ is trivial if $x \in X_i$ for some trivial X_i . For each $i \in [t]$, at most two

elements of $\{A_i, B_i, C_i\}$ are trivial, and therefore, V_i contains at most $2\epsilon_1|V_i|$ many trivial vertices. Thus, at most $2\epsilon_1(3n)$ vertices are trivial. For each $X \in \{A, B, C\}$, let \mathcal{I}_X be the set of $i \in [T]$ such that X_i is non-trivial. Then for each $X \in \{A, B, C\}$, we have $|X \cap (\bigcup_{i \in \mathcal{I}_X} X_i)| \geq (1 - 12\epsilon_1)|X|$, and consequently

$$|\mathcal{I}_X| \geq (1 - 12\epsilon_1)|X|/(3n/t) = (1 - 12\epsilon_1)t/3.$$

Now define

$$\mathcal{S}_1 = \{X_i Y_j : X \neq Y \in \{A, B, C\} \text{ and either } i = j, \text{ or } i \in \mathcal{I}_X \text{ or } j \in \mathcal{I}_Y\}.$$

By the above, $|\mathcal{S}_1| \leq 3t + 3(12\epsilon_1)t^2/3 \leq 40\epsilon_1 t^2$. Now let

$$\mathcal{S}_2 = \{X_i Y_j : X \neq Y \in \{A, B, C\}, ij \notin \mathcal{S}_1 \text{ and } |\{Z_s : X_i Y_j Z_s \in \Sigma\}| \geq \sqrt{\epsilon'_1} t\}.$$

Since $|\Sigma| \leq \epsilon'_1 t^3$, $|\mathcal{S}_2| \leq \sqrt{\epsilon'_1} t^2$. Now let $\mathcal{S} = \mathcal{S}_1 \cup \mathcal{S}_2$. Then $|\mathcal{S}| \leq 2\sqrt{\epsilon'_1} t^2$ (recall $\epsilon'_1 \ll \epsilon_1 \ll 1$). By above, $|\mathcal{I}_B||\mathcal{I}_C| \geq (1 - 12\epsilon_1)^2 t^2/3$, so there must exist some $B_j C_s \notin \mathcal{S}$ with $B_j \in \mathcal{I}_B$ and $C_s \in \mathcal{I}_C$. Define $\mathcal{I}'_A = \{i \in \mathcal{I}_A : A_i B_j C_s \notin \Sigma\}$, and set $A' = \bigcup_{i \in \mathcal{I}'_A} A_i$. By construction, and since $\epsilon'_1 \ll \epsilon_1$,

$$|A'| \geq (1 - 12\epsilon_1)|A| - \sqrt{\epsilon'_1} t(3n/t) \geq (1 - 13\epsilon_1)|A|.$$

Given $\alpha \in [\ell]$, define

$$I_{B_j C_s}^\alpha = \{u \in [\ell] : |P_{B_j C_s}^\alpha \cap \gamma_u| \geq \epsilon_1 |P_{B_j C_s}^\alpha|/\ell\},$$

and set $R_{B_j C_s}^\alpha = \bigcup_{u \in I_{B_j C_s}^\alpha} (P_{B_j C_s}^\alpha \cap \gamma_u)$. Note

$$|P_{B_j C_s}^\alpha \setminus R_{B_j C_s}^\alpha| \leq \ell(\epsilon_1 |P_{B_j C_s}^\alpha|/\ell) < \epsilon_1 |P_{B_j C_s}^\alpha|.$$

Given $v \in [m]$, let $\gamma_v^+ = \bigcup_{u > v} \gamma_u$ and $\gamma_v^- = \bigcup_{v < u} \gamma_u$. Then define

$$I_{B_j C_s}^\alpha(\epsilon_1) = \{u \in I_{B_j C_s}^\alpha : \min\{|P_{B_j C_s}^\alpha \cap \gamma_u^+|, |P_{B_j C_s}^\alpha \cap \gamma_u^-|\} \geq \epsilon_1 |P_{B_j C_s}^\alpha|\}.$$

Clearly $I_{B_j C_s}^\alpha(\epsilon_1)$ is an interval, say $I_{B_j C_s}^\alpha(\epsilon_1) = [u_1, u_2]$. Define

$$R_{B_j C_s}^\alpha(\epsilon_1) = \bigcup_{u \in I_{B_j C_s}^\alpha(\epsilon_1)} (P_{B_j C_s}^\alpha \cap \gamma_u).$$

This implies that

$$|P_{B_j C_s}^\alpha \setminus R_{B_j C_s}^\alpha(\epsilon_1)| \leq |P_{B_j C_s}^\alpha \setminus R_{B_j C_s}^\alpha| + |R_{B_j C_s}^\alpha \cap [u_2 + 1, n]| + |P_{B_j C_s}^\alpha \cap [1, u_1 - 1]| \leq 3\epsilon_1 |P_{B_j C_s}^\alpha|.$$

Consequently,

$$|\bigcup_{\alpha \in [\ell]} R_{B_j C_s}^\alpha(\epsilon_1)| \geq (1 - 3\epsilon_1)|B_j||C_s|,$$

which implies that $|\bigcup_{\alpha \in [\ell]} I_{B_j C_s}^\alpha(\epsilon_1)| \geq (1 - 3\epsilon_1)|B_j||C_s|/(|B_j||C_s|/\ell) \geq (1 - 3\epsilon_1)\ell$.

Given $u \in [\ell]$, let $u^+ = \lceil (u - 1)n/\ell \rceil, n$, $u^- = [1, \lfloor (u + 1)n/\ell \rfloor]$, and set

$$u^* = [\lfloor (u + 1)n/\ell \rfloor, \lceil (u - 1)n/\ell \rceil].$$

Given $i \in [t]$, we then define

$$I_i = \{u \in [\ell] : |A_i \cap u^*| \geq \epsilon_1 |A_i|\},$$

and we set $I_i(\epsilon) = \{u \in [\ell] : \min\{|A_i \cap u^+|, |A_i \cap u^-|\} \geq \epsilon_1 |A_i|\}$. Clearly $I_i(\epsilon_1)$ is an interval, say $I_i(\epsilon_1) = [v_1, v_2]$. Set $A_i(\epsilon_1) = \bigcup_{u \in I_i(\epsilon_1)} (A_i \cap u^*)$, and observe that

$$|A_i \setminus A_i(\epsilon_1)| \leq |A_i \cap v_1^-| + |A_i \cap v_2^+| \leq 2\epsilon_1 |A_i|.$$

Consequently, $|\bigcup_{i \in \mathcal{I}'_A} A_i(\epsilon_1)| \geq (1 - 2\epsilon_1)|A'|$, which implies that

$$|\bigcup_{i \in \mathcal{I}'_A} I_i(\epsilon_1)| \geq (1 - 2\epsilon_1)|A'|/(n/\ell) \geq (1 - 2\epsilon_1)^2 \ell (1 - 13\epsilon_1) \geq (1 - 20\epsilon_1)\ell.$$

This implies, with the above, that

$$|\left(\bigcup_{i \in \mathcal{I}'_A} I_i(\epsilon_1)\right) \cap \left(\bigcup_{\alpha \in [\ell]} I_{B_j C_s}^\alpha(\epsilon_1)\right)| \geq (1 - 23\epsilon_1)\ell > 0.$$

Choose any $u \in \left(\bigcup_{i \in \mathcal{I}'_A} I_i(\epsilon_1)\right) \cap \left(\bigcup_{\alpha \in [\ell]} I_{B_j C_s}^\alpha(\epsilon_1)\right)$ and define

$$\begin{aligned} G_1 &= ((A_i \cap u^+) \cup B_j \cup C_s, (P_{B_j C_s}^\alpha \cap \gamma_u^-) \cup P_{A_i C_s}^\beta \cup P_{B_j A_i}^\gamma) \\ G_0 &= ((A_i \cap u^-) \cup B_j \cup C_s, (P_{B_j C_s}^\alpha \cap \gamma_u^+) \cup P_{A_i C_s}^\beta \cup P_{B_j A_i}^\gamma) \end{aligned}$$

By construction, $K_3^{(2)}(G_1) \subseteq E$ and $K_3^{(2)}(G_0) \cap E \neq \emptyset$. On the other hand, since $V_i V_j V_s \notin \Sigma$, we must have that $(H_{ijs}^{\alpha, \beta, \gamma}, G_{ijs}^{\alpha, \beta, \gamma})$ satisfies $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$. Let $d \in [0, 1]$ be such that $|E \cap K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma})| = d|K_3^{(2)}(G_{ijs}^{\alpha, \beta, \gamma})|$. Without loss of generality, say $d \leq 1/2$ (the other case is similar). By above, and by definition of $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$,

$$\frac{1}{2}|K_3^{(2)}(G_1)| \leq |K_3^{(2)}(G_1)|(1 - d) = \left| |E \cap K_3^{(2)}(G_1)| - d|K_3^{(2)}(G_1)| \right| \leq \epsilon'_1 \frac{n^3}{\ell^3 t^3}.$$

Thus $|K_3^{(2)}(G_1)| \leq 2\epsilon'_1 \frac{n^3}{\ell^3 t^3}$.

Given $b \in B_j$, let $X_b^- = \{a \in A_i \cap u^- : ab \in P_{ij}^\gamma\}$. Because P_{ij}^γ satisfies $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, Lemma 3.18 implies $P_{A_i B_j}^\gamma[A_i \cap u^-, B_j]$ satisfies $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Therefore, a standard argument shows that there is a set $B'_j \subseteq V_j$ of size at least $(1 - \sqrt{\epsilon_2(\ell)})|B_j|$ such that for all $b \in B'_j$,

$$|X_b^-| = \frac{|A_i \cap u^-|}{\ell} (1 \pm \sqrt{\epsilon_2(\ell)}) \geq \frac{|A_i \cap u^-|}{2\ell} \geq \epsilon_1 |V_i|/2\ell.$$

Similarly, let $Y_b^+ = \{c \in C_s : vw \in P_{B_j C_s}^\alpha \cap \gamma_u^+\}$. Because $P_{B_j C_s}^\alpha$ satisfies $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, Lemma 3.18 implies that $P_{B_j C_s}^\alpha$ satisfies $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Thus, there is a set $B''_j \subseteq B_j$ of size at least $(1 - \sqrt{\epsilon_2(\ell)})|B_j|$ so that for all $v \in B''_j$, $|Y_b^+| = \frac{|C_s|}{\ell} (1 \pm \sqrt{\epsilon_2(\ell)}) \geq \epsilon_1 |V_s|/2\ell$. Combining the above with the fact that B_j is non-trivial, we have

$$|B''_j \cap B'_j| \geq (1 - 2\sqrt{\epsilon_2(\ell)})|B_j| \geq \epsilon_1^2 |V_j|/2.$$

Now given $b \in B'_j \cap B''_j$, since $|X_b^-| \geq \frac{1}{2\ell}|A_i| \geq \frac{\epsilon_1}{2\ell}|V_i|$ and $|Y_b^+| \geq \frac{\epsilon_1^2}{2\ell}|C_s| \geq \frac{\epsilon_1^3}{2\ell}|V_s|$, and since P_{is}^β satisfies $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, we must have that $P_{is}^\beta[X_b^-, Y_b^+] \geq \frac{|X_b^-||Y_b^+|}{2\ell}$. Note that for all $ac \in P_{is}^\beta[X_b, Y_b]$, $bac \in K_3^{(2)}(G_1)$. Thus we have shown

$$|K_3^{(2)}(G_1)| \geq |B'_j \cap B''_j| \frac{\epsilon_1|V_s|}{2\ell} \frac{\epsilon_1|V_i|}{2\ell} \geq \epsilon_1^4 |V_i||V_j||V_s|/4\ell^2.$$

Since $\epsilon'_1 \ll \epsilon_1$, this contradicts that $|K_3^{(2)}(G_1)| \leq 2\epsilon'_1 \frac{n^3}{\ell^3 t^3}$. This finishes the proof. \square

7. BINARY $\text{disc}_{2,3}$ -ERROR

In this section, we switch gears and consider binary error. First, in Section 7.1, we will give several examples of properties which admit binary $\text{disc}_{2,3}$ -error but contain $\overline{U}(k)$ for all k . These examples will be used to prove Proposition 2.58, which says that there exist properties which are far from WNIP but which admit $\text{disc}_{2,3}$ -error. We will also show in Section 7.2, that both \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ have finite VC-dimension. This implies, by Proposition 2.44, that to show \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ require non-binary $\text{disc}_{2,3}$ -error, it suffices to show that they both require non-binary vdisc_3 -error.

The rest of the section will then be concerned with showing that \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ require non-binary vdisc_3 -error. There are two ways one could do this. The most straightforward option is to present separate proofs for each. However, the proofs are so similar that this becomes quite repetitive. The second option is to isolate the important commonalities between the two examples, and then give a unified proof for both at once. Although it is a bit more work, we have chosen the second option, as we think it is of independent interest to know this can be done. Thus, in Section 7.3 we make a general definition, namely that of a $(p, \mu, \tau, \alpha, \rho)$ -special 3-graph (see Definition 7.25). Roughly speaking, this is a 3-partite 3-graph equipped with a metric on each of its parts, and which satisfies certain axioms relating the edge relation and the metric structure. We will show that both $\text{HP}(N)$ and $\text{GS}_p(n)$ are $(p, \mu, \tau, \alpha, \rho)$ -special, for appropriately chosen parameters.

In Section 7.4, we show that any vdisc_3 -regular partition of a sufficiently large $(p, \mu, \tau, \alpha, \rho)$ -special 3-graph requires non-binary vdisc_3 -error (see Theorem 7.28). From this we will deduce Theorems 2.53 and 2.56. We will end the section with a proof of Theorem 2.12.

7.1. Admitting binary $\text{disc}_{2,3}$ -error is distinct from admitting zero vdisc_3 -error.

In this section we give a class of examples admitting binary $\text{disc}_{2,3}$ -error. We will use these examples to show that admitting binary $\text{disc}_{2,3}$ -error is distinct from admitting zero vdisc_3 -error, and that there exist properties far from WNIP which still admit binary $\text{disc}_{2,3}$ -error. The idea behind these examples is that if a property is “essentially binary” (in the sense of Definition 7.1 below), then it admits binary $\text{disc}_{2,3}$ -error.

Definition 7.1. Suppose $G = (V \cup U \cup W, E)$ is a finite 3-partite, 3-graph. We say that G is ℓ -binary if the following holds. There are partitions \mathcal{P}_{UV} of $K_2(U, V)$, \mathcal{P}_{UW} of $K_2(U, W)$, and \mathcal{P}_{VW} of $K_2(V, W)$, such that

- $|\mathcal{P}_{UV}|, |\mathcal{P}_{UW}|, |\mathcal{P}_{VW}| \leq \ell$,

- for all $(\alpha, \beta, \gamma) \in \mathcal{P}_{UV} \times \mathcal{P}_{UW} \times \mathcal{P}_{VW}$, if $G_{\alpha, \beta, \gamma} = (V \cup U \cup W, \alpha \cup \beta \cup \gamma)$, then either $K_3^{(2)}(G_{\alpha, \beta, \gamma}) \subseteq E$ or $K_3^{(2)}(G_{\alpha, \beta, \gamma}) \cap E = \emptyset$.

Here are some examples of ℓ -binary 3-graphs.

- Given n , let $A_n = \{a_1, \dots, a_n\}$, $B_n = \{b_1, \dots, b_n\}$, and $C_n = \{c_1, \dots, c_n\}$. Then $G = (A_n \cup B_n \cup C_n, \{a_i b_j c_k : i < j < k\})$ is 2-binary. Indeed, set

$$\mathcal{P}_{AB} = \{\{a_i b_j : i < j\}, \{a_i b_j : i \geq j\}\},$$

and define \mathcal{P}_{BC} and \mathcal{P}_{AC} analogously.

- For any 3-partite graph $G = (V \cup U \cup W, E)$, the hypergraph with vertex set $V \cup U \cup W$ and edge set $\{(x, y, z) \in VUW : xyz \text{ is a triangle in } G\}$ is 2-binary. Indeed, define

$$\mathcal{P}_{UV} = \{E \cap K_2(U, V), K_2(U, V) \setminus E\},$$

and define \mathcal{P}_{UW} and \mathcal{P}_{VW} analogously.

We now show that ℓ -binary hypergraphs admit $\text{disc}_{2,3}$ -regular decompositions with binary $\text{disc}_{2,3}$ -error.

Proposition 7.2. *For all $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ and $\ell \geq 1$, there is n_0 and T, L, N such that the following holds. If $G = (V \cup U \cup W, E)$ is an ℓ -binary, 3-partite, 3-uniform hypergraph with $|U \cup V \cup W| = n \geq N$, then there is $t_0 \leq t \leq T$ and $\ell' \leq L$ such that G has an $(\epsilon_1, \epsilon_2(\ell'))$ -regular $(t, \ell, \epsilon_2(\ell'), \epsilon_1)$ -decomposition with binary $\text{disc}_{2,3}$ -error.*

Proof. Fix $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ and $\ell \geq 1$. Choose $\epsilon'_1 \ll \mu_1 \ll \epsilon_1$, and $\epsilon'_2, \mu : \mathbb{N} \rightarrow (0, 1]$ with the property that $\epsilon'_2(x) \ll \mu_2(x) \ll \epsilon_2(x)$ for all $x \geq 1$. Choose f and g as in Lemma 3.27 for ϵ'_1 and ϵ'_2 . Then set $L = f(3, \ell, 3, 1)$, $T = f(3, \ell, 3, 1)$, and let $N_1 = N_1(3, \ell, 3, 1)$ from Lemma 3.27. Finally, choose some $N \gg N_1, (\epsilon'_1)^{-1}, \epsilon'_2(L)^{-1}$.

Now suppose $H = (V \cup U \cup W, E)$ is a 3-partite 3-graph which is ℓ -binary, and where $|U \cup W \cup V| = n \geq N$. Choose partitions $K_2[U, V] = \bigcup_{\alpha \leq \ell_{UV}} P_\alpha^{UV}$, $K_2[U, W] = \bigcup_{\alpha \leq \ell_{UW}} P_\alpha^{UW}$, and $K_2[V, W] = \bigcup_{\alpha \leq \ell_{VW}} P_\alpha^{VW}$, with $\ell_{UV}, \ell_{UW}, \ell_{VW} \leq \ell$ witnessing that G is ℓ -binary. Note that each triple $(P_\alpha^{UV}, P_\beta^{UW}, P_\gamma^{VW})$ comes equipped with a $\delta = \delta(\alpha, \beta, \gamma) \in \{0, 1\}$ so that $K_3^{(2)}(U \cup V \cup W, P_\alpha^{UV} \cup P_\beta^{UW} \cup P_\gamma^{VW}) \subseteq E^\delta$.

Let \mathcal{P} be the $(3, \ell)$ -decomposition of V with $\mathcal{P}_{\text{vert}} = \{U, V, W\}$ and $\mathcal{P}_{\text{edge}} = \{P_\alpha^{XY} : XY \in \{UV, UW, VW\}, \alpha \leq \ell_{XY}\}$. Let \mathcal{Q} be the $(3, 1)$ decomposition of V consisting of $\mathcal{Q}_{\text{vert}} = \{U, V, W\}$ and $\mathcal{Q}_{\text{edge}} = \{K_2[U, V], K_2[V, W], K_2[U, W]\}$. Apply Lemma 3.27 to \mathcal{P} and \mathcal{Q} to obtain $\ell' \leq L$, $t \leq T$, and a $(t, \ell', \epsilon'_1, \epsilon'_2(\ell'))$ -decomposition \mathcal{R} of V , which has no disc_2 -irregular triads and which is an $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement of \mathcal{P} . Let $\Sigma \subseteq \binom{\mathcal{R}_{\text{vert}}}{2}$ be as in the definition of an $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement of \mathcal{P} . Let $\mathcal{R}_{\text{vert}} = \{V_1, \dots, V_t\}$ and $\mathcal{R}_{\text{edge}} = \{R_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell'\}$.

Suppose $R_{ij}^\alpha, R_{ik}^\beta, R_{jk}^\gamma \in \mathcal{R}_{\text{edge}}$ are such that $V_i V_j, V_j V_k, V_i V_k \notin \Sigma$. Let $R_{ijk}^{\alpha, \beta, \gamma}$ denote the triad $(V_i \cup V_j \cup V_k, R_{ij}^\alpha \cup R_{ik}^\beta \cup R_{jk}^\gamma)$. By definition of an $(\epsilon'_1, \epsilon'_2(\ell'))$ refinement, there are $X, Y, Z \in \{U, V, W\}$ with $|V_i \setminus X| < \epsilon'_1 |V_i|$, $|V_j \setminus Y| < \epsilon'_1 |V_j|$, and $|V_k \setminus Z| < \epsilon'_1 |V_k|$. If

$X = Y$, $X = Z$, or $Y = Z$, then $|E \cap K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma})| = 0$. By Proposition 3.22, $R_{ijk}^{\alpha,\beta,\gamma}$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell'))$ with respect to H .

Assume now that X, Y, Z are pairwise distinct. Then by definition of an $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement, there are α', β', γ' so that the following hold (possibly after relabeling),

$$|R_{ij}^\alpha \setminus P_{\alpha'}^{UV}| < \epsilon'_1 |R_{ij}^\alpha|, \quad |R_{ik}^\beta \setminus P_{\beta'}^{UW}| < \epsilon'_1 |R_{ik}^\beta|, \quad \text{and} \quad |R_{jk}^\gamma \setminus P_{\gamma'}^{VW}| < \epsilon'_1 |R_{jk}^\gamma|,$$

and further, each of $R_{ij}^\alpha \setminus P_{\alpha'}^{UV}$, $R_{ik}^\beta \setminus P_{\beta'}^{UW}$, and $R_{jk}^\gamma \setminus P_{\gamma'}^{VW}$ have $\text{disc}_2(\epsilon'_2(\ell'))$. By Corollary 3.6, $K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma}) = (1 \pm \mu_2(\ell')) \frac{|V_i||V_j||V_k|}{(\ell')^3}$, and the number of triples in $K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma})$ involving a pair from one of $R_{ij}^\alpha \setminus P_{\alpha'}^{UV}$, $R_{ik}^\beta \setminus P_{\beta'}^{UW}$, or $R_{jk}^\gamma \setminus P_{\gamma'}^{VW}$ is at most

$$3\epsilon'_1(1 + \mu_2(\ell')) \frac{1}{(\ell')^3} |V_i||V_j||V_k| \leq 4\mu_1 |K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma})|.$$

Thus, if $\delta = \delta(\alpha', \beta', \gamma')$, then

$$|E^\delta \cap K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma})| \geq (1 - 4\mu_1) |K_3^{(2)}(R_{ijk}^{\alpha,\beta,\gamma})|.$$

By Proposition 3.22, this shows $R_{ijk}^{\alpha,\beta,\gamma}$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell'))$ with respect to H . By definition, this shows that \mathcal{R} has binary $\text{disc}_{2,3}$ -error with respect to H . \square

As a consequence we can now show that many examples admit binary $\text{disc}_{2,3}$ -error. In particular, we can now prove Proposition 2.58, which shows that there is a property containing $\overline{U}(k)$ for every k and which admits binary $\text{disc}_{2,3}$ -error. Recall that given an infinite 3-graph H , $\text{age}(H)$ denotes the class of all finite 3-graphs which are isomorphic to an induced sub-3-graph of H . It is easy to see that $\text{age}(H)$ is always a hereditary 3-graph property.

Definition 7.3. Define \mathcal{W} to be the 3-graph

$$\mathcal{W} = (\{a_i, b_i : i \in \mathbb{N}\} \cup \{c_S : S \subseteq \mathbb{N}\}, \{(a_i, b_j, c_S) : j \in S\}).$$

Let $\mathcal{H}_{\mathcal{W}} := \text{age}(\mathcal{W})$.

PROOF OF PROPOSITION 2.58: Clearly every element in $\mathcal{H}_{\mathcal{W}}$ is 2-binary. Consequently, by Proposition 7.2, $\mathcal{H}_{\mathcal{W}}$ admits binary $\text{disc}_{2,3}$ -error. On the other hand, $\mathcal{H}_{\mathcal{W}}$ contains $\overline{U}(k) \in \mathcal{W}$ for all k by definition. \square

We can use $\mathcal{H}_{\mathcal{W}}$ as an example to demonstrate several other distinctions among the definitions from the introduction. Recall that in Corollary 2.45, we showed that a property \mathcal{H} admitting binary vdisc_3 -error must be vdisc_3 -homogeneous. For a property \mathcal{H} to be vdisc_3 -homogeneous, by Theorem 2.28, there must be a $k \geq 1$ such that $\overline{U}(k) \notin \text{Trip}(\mathcal{H})$. Consequently, $\mathcal{H}_{\mathcal{W}}$ does not admit binary vdisc_3 -error, but does admit binary $\text{disc}_{2,3}$ -error. Moreover, $\mathcal{H}_{\mathcal{W}}$ has unbounded weak VC-dimension (which implies unbounded VC-dimension) and is not weakly stable (and thus not stable). Thus $\mathcal{H}_{\mathcal{W}}$ shows that admitting binary $\text{disc}_{2,3}$ -error is distinct from admitting binary vdisc_3 -error, from NIP, from WNIP, from stability and from weak stability.

7.2. Proof of Proposition 2.57. In this section, we show that both \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ have bounded VC-dimension, which will allow us to apply Proposition 2.44 to both. We recall some terminology from the study of VC-dimension.

Suppose X is a set. A *set system* is a pair (X, \mathcal{F}) where $\mathcal{F} \subseteq \mathcal{P}(X)$. Given $Y \subseteq X$, define $\mathcal{F} \cap Y := \{Y \cap F : F \in \mathcal{F}\}$. We then say that \mathcal{F} *shatters* Y if $|\mathcal{F} \cap Y| = 2^{|Y|}$.

Definition 7.4. The *VC-dimension* of (X, \mathcal{F}) , $\text{VC}(X, \mathcal{F})$, is the largest size of a shattered subset of X .

Note that given a 3-graph $H = (V, E)$, by definition,

$$\text{VC}(H) = \max \left\{ \text{VC} \left(V, \left\{ N_H(ab) : ab \in \binom{V}{2} \right\} \right), \text{VC} \left(\binom{V}{2}, \{ N_H(a) : a \in V \} \right) \right\}.$$

We will now show that $\mathcal{H}_{\text{GS}_p}$ has bounded VC-dimension. The key ingredient is a fact proved in [60].

Theorem 7.5. *For all $p \geq 3$, the set system $\mathcal{F}_{\text{GS}_p} = (\mathbb{F}_p^n, \{A(p, n) - g : g \in \mathbb{F}_p^n\})$ has VC-dimension 3.*

Corollary 7.6. *For all $p \geq 3$, $\mathcal{H}_{\text{GS}_p}$ has VC-dimension 3.*

Proof. Fix $p \geq 3$. We show $\mathcal{H}_{\text{GS}_p}$ has VC-dimension at most 3. Suppose towards a contradiction there is $G = (V, E) \in \mathcal{H}_{\text{GS}_p}$ with VC-dimension at least 4. By definition of $\mathcal{H}_{\text{GS}_p}$ there is $n \geq 1$ such that G is isomorphic to an induced sub-3-graph of $\text{GS}_p(n)$. Thus, $\text{GS}_p(n)$ also has VC-dimension at least 4. Let $A = \{a_g : g \in \mathbb{F}_p^n\}$, $B = \{b_g : g \in \mathbb{F}_p^n\}$, $C = \{c_g : g \in \mathbb{F}_p^n\}$. Then $\text{GS}_p(n)$ has vertex set $V' = A \cup B \cup C$ and edge set $E' = \{a_g b_{g'} c_{g''} : g + g' + g'' \in A(n, p)\}$. Since $\text{VC}(\text{GS}_p(n)) \geq 4$, by definition, one of the following hold.

- (1) There is $X \subseteq V'$ a set of size 4, and for each $Y \subseteq X$, $e_Y \in \binom{V'}{2}$ such that for all $x \in X$, $\{x\} \cup e_Y \in E'$ holds if and only if $x \in Y$.
- (2) There is $X \subseteq \binom{V'}{2}$ of size 4 and for each $Y \subseteq X$, $v_Y \in A \cup B \cup C$ such that for all $e \in X$, $\{v_Y\} \cup e \in E'$ if and only if $e \in Y$.

Suppose first we are in case (1). Because there exists $e \in \binom{V'}{2}$ with $X = N(e) \cap X$, we must have that either $X \subseteq A$, $X \subseteq B$, or $X \subseteq C$. Without loss of generality, let us assume $X \subseteq A$. Say $X = \{a_x, a_y, a_z, a_w\}$. By definition of $\text{GS}_p(n)$, for any $\emptyset \neq Y \subseteq X$, since $N(e_Y) \cap X \neq \emptyset$, must have that $e_Y = b_{g_Y} c_{h_Y}$ for some $g_Y, h_Y \in \mathbb{F}_p^n$. But then $\{x, y, z, w\}$ is shattered by $\mathcal{F}_{\text{GS}_p}$, contradicting Theorem 7.5.

Suppose now we are in case (2). Because there is some $v \in V$ such that $X \cap N(v) = X$, we must have that either $X \subseteq AB$, $X \subseteq AC$ or $X \subseteq BC$. Without loss of generality, assume $X \subseteq AB$, say $X = \{a_{g_1} b_{h_1}, \dots, a_{g_4} b_{h_4}\}$. Note that for any $\emptyset \neq Y \subseteq X$, since $N(e_Y) \cap X' \neq \emptyset$, we must have that $v_Y = c_{\nu_Y}$ for some $\nu_Y \in \mathbb{F}_p^n$. Again, we see that $\{g_1 + h_1, \dots, g_4 + h_4\}$ is shattered by $\mathcal{F}_{\text{GS}_p}$, contradicting Theorem 7.5.

On the other hand, by Theorem 7.5, there is set $X \subseteq \mathbb{F}_p^n$ of size 4 shattered by $\mathcal{F}_{\text{GS}_p}$. Say $X = \{g_1, \dots, g_4\}$. It is easy to see $\{b_0 c_{g_1}, \dots, b_0 c_{g_4}\}$ is shattered in $(V, \{N_{\text{GS}_p(n)}(a) : a \in A\})$, and thus $\mathcal{H}_{\text{GS}_p(n)}$ has VC-dimension at least 3. \square

We now turn to showing that \mathcal{H}_{HP} has bounded VC-dimension. This will be clear to the model theorist, as the edge relation of \mathcal{H}_{HP} is uniformly definable in Presburger arithmetic, for instance. On the other hand, this fact will also be known to many combinatorialists, as the 3-graphs of \mathcal{H}_{HP} are examples of semialgebraic hypergraphs (see e.g. [4]), which are well known to have bounded VC-dimension. However, we include a proof here for the sake of completeness.

Fact 7.7. \mathcal{H}_{HP} has VC-dimension 1.

Proof. By definition of \mathcal{H}_{HP} , it suffices to show that $\text{VC}(\text{HP}(N)) \leq 1$ for all N . Fix $N \geq 1$, and consider the 3-graph $\text{HP}(N) = (V, E)$ where $V = A \cup B \cup C$, $A = \{a_i : i \in [N]\}$, $B = \{b_i : i \in [N]\}$, $C = \{c_i : i \in [N]\}$, and $E = \{a_i b_j c_k : i + j + k \geq N + 2\}$. Suppose towards a contradiction that $\text{HP}(N)$ has VC-dimension at least 2. This means there is either a set $X \subseteq V$ of size 2 shattered in (V, \mathcal{F}_1) where $\mathcal{F}_1 = \{N_{\text{HP}(N)}(uv) : uv \in \binom{V}{2}\}$, or a there is a set $X \subseteq \binom{V}{2}$ of size 2 shattered $(\binom{V}{2}, \mathcal{F}_2)$, where $\mathcal{F}_2 = \{N_{\text{HP}(N)}(u) : u \in V\}$.

Suppose first there is $X \subseteq \binom{V}{2}$ of size 2 shattered in $(\binom{V}{2}, \mathcal{F}_1)$. Arguing as in the proof of Corollary 7.6, it is not difficult to show that because $\text{HP}(N)$ is 3-partite, $X \subseteq K_2[A, B]$, or $X \subseteq K_2[A, C]$, or $X \subseteq K_2[B, C]$. Since A, B , and C behave symmetrically in $\text{HP}(N)$, let us assume that $X \subseteq K_2[A, B]$. Say $X = \{a_{i_1} b_{j_1}, a_{i_2} b_{j_2}\}$. By relabeling if necessary, assume $i_1 + j_1 \leq i_2 + j_2$. It is not difficult to see that because X is shattered in $\text{HP}(N)$, there is some $c_k \in C$ so that $c_k a_{i_1} b_{j_1} \in E$ but $c_k a_{i_2} b_{j_2} \notin E$. But this means that $i_1 + j_1 + k \geq N + 2$ while $i_2 + j_2 + k < N + 2$, which is not possible, as $i_1 + j_1 \leq i_2 + j_2$.

Suppose now that there is $X \subseteq V$ of size 2 shattered (V, \mathcal{F}_2) . Again, it is not hard to see that since $\text{HP}(N)$ is 3-partite, we must have $X \subseteq A$, $X \subseteq B$, or $X \subseteq C$. As the roles of A, B, C are symmetric, we may assume, say, $X = \{a_{i_1}, a_{i_2}\} \subseteq A$, where $i_1 \leq i_2$. Since X is shattered, there is some $b_j c_k$ so that $b_j c_k a_{i_1} \in E$ but $b_j c_k a_{i_2} \notin E$. But this means $j + k + i_1 \geq N + 2$ while $j + k + i_2 < N + 2$, which is not possible, as $i_1 \leq i_2$. \square

7.3. Special 3-graphs and the proof of Theorems 2.53 and 2.56. In this subsection we define $(p, \mu, \tau, \alpha, \rho)$ -special 3-graphs (see Definition 7.25). The purpose of Definition 7.25 is to isolate common properties our main examples, $\text{GS}_p(n)$ and $\text{HP}(n)$, specifically those which are required to prove both examples require non-binary vdisc_3 -error. The main idea is that both examples come equipped with natural metrics on their set of vertices, which interact nicely with their edge sets. As Definition 7.25 is quite technical, we will begin by examining each of the two examples in turn by way of motivation. We will begin with $\text{GS}_p(n)$, where the picture is somewhat simpler.

7.3.1. Special properties of $\text{GS}_p(n)$. In this section we will go through the main properties of $\text{GS}_p(n)$ used in the proof that it requires non-binary vdisc_3 -error. We first recall the definition of $\text{GS}_p(n)$, and prove some basic facts about it. Given $x \in \mathbb{F}_p^n$ and $i \in [n]$, x_i denotes the i th coordinate of x . Given $1 \leq i \leq n$, we let H_i denote the subspace of \mathbb{F}_p^n with the property that for all $x \in H_i$, $x_j = 0$ for all $1 \leq j \leq i$. By convention, we set $H_0 = \mathbb{F}_p^n$.

Recall that $A(p, n) = \bigcup_{i=1}^n H_i + e^i$, where e^i is the i th standard basis vector in \mathbb{F}_p^n . In other words, a vector $x \in \mathbb{F}_p^n$ is in $A(p, n)$ if the first non-zero coordinate in x is 1. Then $\text{GS}_p(n)$ is defined as follows.

$$\text{GS}_p(n) = (\{a_g, b_g, c_g : g \in \mathbb{F}_p^n\}, \{a_g b_{g'} c_{g''} : g + g' + g'' \in A(p, n)\}).$$

To ease notation, while in Section 7.3.1, we will let $A = \{a_g : g \in \mathbb{F}_p^n\}$, $B = \{b_g : g \in \mathbb{F}_p^n\}$, $C = \{c_g : g \in \mathbb{F}_p^n\}$, $E = E(\text{GS}_p(n))$, and $N = p^n$. Give $x, y \in \mathbb{F}_p^n$ we set

$$\lambda(x, y) = \max\{i \in \{0, \dots, n\} : x \text{ and } y \text{ are in the same coset of } H_i\}.$$

Note that by definition, $x_{\lambda(x,y)+1} \neq y_{\lambda(x,y)+1}$ and for all $1 \leq j \leq \lambda(x, y)$, $y_j = x_j$.

Lemma 7.8. *Given $x, y \in \mathbb{F}_p^n$, set $d(x, y) = 0$ if $x = y$ and set $d(x, y) = p^{-\lambda(x,y)}$ if $x \neq y$. Then d is an ultra metric on \mathbb{F}_p^n , with distances in $[0, 1]$, and for all $x \in \mathbb{F}_p^n$, $B_{p^{-i}}(x)$ is the coset of H_{i+2} containing x .*

Proof. If $x \neq y$, then x and y are in different cosets of H_n , so $d(x, y) \geq p^{-n} > 0$. Given $x \in \mathbb{F}_p^n$, it is clear that $d(x, y) = p^{-i}$ if and only if y is in the same coset of H_i as x , but not the same coset of H_{i+1} . Thus $B_{p^{-i}}(x) = \{y \in \mathbb{F}_p^n : d(x, y) < p^{-i}\}$ is the coset of H_{i+2} containing x . This shows that $d(x, y) = d(y, x)$ for all $x, y \in \mathbb{F}_p^n$ and $d(x, y) = 0$ if and only if $x = y$.

Consider $x, y, z \in \mathbb{F}_p^n$. Suppose $d(x, y) = p^{-i}$, so x, y are in the same coset of H_i , say $H_i + g$, but not the same coset of H_{i+1} , say $x \in H_{i+1} + g + g'$ and $y \in H_{i+1} + g + g''$. If $z \notin H_i + g$, then $d(x, z) = d(y, z) = p^{-i-1} = p d(x, y)$. If $z \in H_i + g$, then it cannot be in both $H_{i+1} + g + g'$ and $H_{i+1} + g + g''$, so $\max\{d(z, x), d(z, y)\} = p^{-i} = d(x, y)$. This shows $d(x, y) \leq \max\{d(x, z), d(y, z)\}$, and thus d is an ultrametric. \square

The fact that d is an ultrametric on \mathbb{F}_p^n makes things particularly easy to analyze in $\text{GS}_p(n)$. For example, for any $r \in (0, 1]$ and $x \in \mathbb{F}_p^n$, $|B_r(x)| = |H_{m+1}| = p^{-m-1}N$, where $m = \max\{i \leq n : r \leq p^{-i}\} = \lceil \log_p(r^{-1}) \rceil$. Thus every ball of radius r contains the same number of points, and $p^{-1}rN \leq |B_r(x)| \leq prN$. It is also very simple to partition balls into smaller ones, as our next claim shows.

Claim 7.9 (Balls can be partitioned). *For any $x \in \mathbb{F}_p^n$ and $0 < r_1 < r_2 \leq 1$, there is a partition of $B_{r_2}(x)$ into at most $pr_2r_1^{-1}$ disjoint open balls of radius r_1 , one of which is centered at x .*

Proof. Let $m_1 = \lceil \log_p(r_1^{-1}) \rceil$, $m_2 = \lceil \log_p(r_2^{-1}) \rceil$. Then $B_{r_2}(x)$ is the coset $H_{m_2+2} + g$ which contains x . We then partition $H_{m_2+2} + g$ into cosets of H_{m_1+2} . All of these cosets are balls of radius r_1 , and one of them must contain x . We can easily compute that the number of balls in this partition is $p^{m_1-m_2} \leq pr_2r_1^{-1}$. \square

Another nice feature of $\text{GS}_p(n)$ is that it is vertex regular (i.e. all of its vertices have the same degree).

Claim 7.10 ($\text{GS}_p(n)$ is vertex regular). *For all $u, v \in A \cup B \cup C$, $|N(v)| = |N(u)| = N^2(\frac{1}{p-1} \pm p^{-n})$.*

Proof. Given $a_g, b_{g'} \in \text{GS}_p(n)$, the neighborhood of $a_g b_{g'}$ is

$$N(a_g b_{g'}) = \{c_{g''} : g'' \in A(p, n) - g - g'\}.$$

Clearly $|N(a_g b_{g'})| = |A(n, p)|$. Observe that

$$|A(p, n)| = \sum_{i=1}^n |H_i| = \sum_{i=1}^n p^{n-i} = N \sum_{i=1}^n p^{-i} = N \left(\frac{1}{p-1} - \frac{p^{-n}}{p-1} \right) \geq \frac{1}{p-1} p^n - \frac{1}{p-1}.$$

Thus $\frac{1}{p-1} p^n - 1 \leq |A(n, p)| = |N(a_g b_{g'})| \leq \frac{1}{p-1} p^n$. Then for any a_g , we compute the size of its neighborhood as $|N(a_g)| = \sum_{g' \in \mathbb{F}_p^n} |N(a_g b_{g'})| = p^n |A(n, p)|$. Clearly any b_g or c_g has the same degree. \square

It is also important to note that we showed in Claim 7.10 that the degree of every vertex has the form αN^2 for some α bounded away from 0 and 1 (e.g. $\alpha = 1/2p$ would work). We now consider some important interactions between the metric and hypergraph structure. Equip each of the sets A, B, C with the natural metrics coming Lemma 7.8 (i.e. for $g, g' \in \mathbb{F}_p^n$, set $d(b_g, b_{g'}) = d(c_g, c_{g'}) = d(a_g, a_{g'}) = d(g, g')$). In our next claim, we use the vertex-regularity in $\text{GS}_p(n)$ to show that given open balls $B_1 \subseteq A$ and $B_2 \subseteq B$, there are many edges and nonedges in $K_3[B_1, B_2, C]$. This can be thought of as a beefed-up version of the fact that for every $a_g b_{g'}$, $N(a_g b_{g'})$ and $\neg N(a_g b_{g'})$ are large (see Claim 7.10).

Claim 7.11. *For all $a_g \in A$, $b_{g'} \in B$, and $r \in (0, 1]$, there are $f_0(r, g, g'), f_1(r, g, g') \in \mathbb{F}_p^n$ such that*

$$K_3[B_r(a_g), B_r(b_{g'}), B_r(c_{f_1(r, g, g')})] \subseteq E \text{ and } K_3[B_r(a_g), B_r(b_{g'}), B_r(c_{f_0(r, g, g')})] \cap E = \emptyset.$$

Proof. Set $m = \lceil \log_p(r^{-1}) \rceil$, so $B_r(a_g) = B_{p^{-m}}(a_g)$ and $B_r(b_{g'}) = B_{p^{-m}}(b_{g'})$. Let $h, h' \in \mathbb{F}_p^n$ be such that in \mathbb{F}_p^n , $B_{p^{-m}}(g) = H_{m+1} + g$ and $B_{p^{-m}}(g') = H_{m+1} + g'$. Define $s = \min\{m, \lambda(g, g')\}$.

Let $f_0(r, g, g') = -\sum_{i=1}^{s+1} (g_i + g'_i) e^i + 2e^{s+1}$ and let $f_1(r, g, g') = -\sum_{i=1}^{s+1} (g_i + g'_i) e^i + e^{s+1}$. Note $H_{s+1} + g + g' + f_0(g, g') = H_{s+1} + 2e^{s+1}$ is disjoint from $A(p, n)$ and $H_{s+1} + g + g' + f_1(g, g') = H_{s+1} + e^{s+1}$ is contained in $A(p, n)$. Thus $K_3[B_r(a_g), B_r(b_{g'}), B_s(c_{f_1(r, g, g')})] \subseteq E$ and $K_3[B_r(a_g), B_r(b_{g'}), B_s(c_{f_0(r, g, g')})] \cap E = \emptyset$, as desired. Note that by construction, $d(f_0(r, g, g'), f_1(r, g, g')) \geq \max\{r, d(g, g')\}$. \square

It will be important for us to know that the $f_0(r, g, g'), f_1(r, g, g'), f_0(r', g'', g'''), f_1(r', g'', g''')$ in Claim 7.11 are far apart for certain choices of g, g', g'', g''', r, r' .

Claim 7.12. *If $g' \neq g''$, then for each $u \in \{0, 1\}$, $d(c_{f_u(r, g, g')}, c_{f_u(r, g, g'')}) = d(b_{g'}, b_{g''})$. Further, if $h \neq h', r' \geq r$, and $d(h, h') \geq \max\{d(g, g'), r'\}$. Then*

$$\min\{d(c_{f_u(r, g, h)}, c_{f_v(r', g', h')}) : u, v \in \{0, 1\}\} \geq d(h, h').$$

Proof. Note that if $g'' \neq g'$, then

$$\lambda(c_{f_0(r, g, g')}, c_{f_0(r, g, g'')}) = \lambda\left(-\sum_{i=1}^{s+1} (g_i + g'_i) e^i + 2e^{s+1}, -\sum_{i=1}^{s'+1} (g_i + g''_i) e^i + 2e^{s'+1}\right),$$

where $s' = \min\{m, \lambda(g, g'')\}$. This shows $d(c_{f_0(g, g')}, c_{f_0(g, g'')}) = d(b_{g'}, b_{g''})$. A similar argument shows that $d(c_{f_1(g, g')}, c_{f_1(g, g'')}) = d(b_{g'}, b_{g''})$.

Suppose now that $g \neq g', h \neq h', r' \geq r$, and assume $d(h, h') \geq \max\{d(g, g'), r'\}$. Set $m' = \lceil \log((r')^{-1}) \rceil$, $s = \min\{m, \lambda(g, h)\}$ and $s' = \min\{m', \lambda(g', h')\}$. By assumption, $\lambda(h, h') \leq \min\{\lambda(g, g'), m'\}$. Given $u \in \{0, 1\}$ let $u \dot{+} 1$ be 2 if $u = 1$ and 1 if $u = 0$. Now fix $(u, v) \in \{0, 1\}^2$ and consider $\lambda(f_u(r, h, g), f_v(r', h', g'))$. This is

$$\begin{aligned} & \lambda\left(-\sum_{i=1}^{s+1} (h_i + g_i)e^i + (u \dot{+} 1)e^{s+1}, -\sum_{i=1}^{s'+1} (g'_i + h'_i)e^i + (v \dot{+} 1)e^{s'+1}\right) \\ &= \lambda\left(-\sum_{i=1}^s 2h_i + (u \dot{+} 1 - h_{s+1} - g_{s+1})e^{s+1}, -\sum_{i=1}^{s'} 2h'_i + (v \dot{+} 1 - h'_{s'+1} - g'_{s'+1})e^{s'+1}\right) \\ &= \lambda(h, h'), \end{aligned}$$

where the last equality is by definition of s and s' , and because $\lambda(h, h') \leq \min\{\lambda(g, g'), m'\}$. Thus $d(c_{f_u(r, h, g)}, c_{f_v(r', h', g')}) \geq d(h, h')$, as desired. \square

We now show that the intersections of certain neighborhoods and non-neighborhoods are contained in open balls.

Claim 7.13. *Given $a_g, b_{g'}, b_{g''} \in V(\text{GS}_p(n))$, $N(a_g, b_{g'}) \cap \neg N(a, b_{g''})$ is contained in an open ball in C ball of radius $d(b_{g'}, b_{g''})$.*

Proof. Given $a \in \mathbb{F}_p^n$ and $\alpha \in \mathbb{F}_p$, define $\tau_i^\alpha(a) := \alpha e^i - a$, and note that

$$A(p, n) - a = \bigcup_{i=1}^n H_i + \tau_i^1(a).$$

Fix $a_g, b_{g'}, b_{g''}$ and consider $C' = N(a_g, b_{g'}) \cap \neg N(a, b_{g''})$. Let $h = g + g'$ and $f = g + g''$. Then let $C' = \{z_w : w \in (A(p, n) - h) \cap (\neg A(p, n) - f)\}$, and set $m = \lambda(h, f)$. Note

$$(A(p, n) - h) \cap (\neg A(p, n) - f) = \left(\bigcup_{i=1}^n H_i + \tau_i^1(h)\right) \cap \left(\bigcup_{i=1}^n H_i + \tau_i^0(f)\right).$$

We prove in [60], that the set above is contained in a single coset K of H_{m-1} (see Lemma C.1 there). Consequently, $C' \subseteq \{c_v : v \in K\}$. We are now done since any coset of H_{m-1} is an open ball of radius $p^{-m} = p^{-\lambda(h, f)} = d(b_{g'}, b_{g''})$ \square

Our last lemma shows that sufficiently small balls around distinct points can be “split apart” using the edge relation in $\text{GS}_p(n)$.

Claim 7.14. *Suppose $a_g, a_{g'} \in A$, and $d(a_g, a_{g'}) = r > 0$. Then for all $b_{g''} \in B$, there is $c_h \in C$ such that $K_3[B_r(c_h), B_r(b_{g''}), B_r(a_g)] \subseteq E$ and $K_3[B_r(c_h), B_r(b_{g''}), B_r(a_{g'})] \cap E = \emptyset$, or vice versa.*

Proof. We begin by defining an element $f(a, b) \in \mathbb{F}_p^n$ for all distinct $a, b \in \mathbb{F}_p$ with the property that for all $a \neq b$ in \mathbb{F}_p ,

$$\{a + f(a, b), b + f(a, b)\} = \{1, u\} \text{ for some } u \notin \{0, 1\}.$$

Given $0 < a < b$, let $f(a, b) = a - 1$. Then set $f(0, p - 1) = f(p - 1, 0) = 2$, for each $b \notin \{0, p - 1\}$, and set $f(0, b) = f(b, 0) = 1$. Finally, set $f(1, b) = f(b, 1) = 0$ for all $b \notin \{0, 1\}$.

Now suppose $a_g, a_{g'} \in A$, and $d(a_g, a_{g'}) = r > 0$ and $b_{g''} \in B$. Let $m = \lambda(g, g')$, so $r = p^{-m}$. Note $g_i = g'_i$ for all $i \leq m$, by definition. Set

$$h = \left(\sum_{i=1}^m -g_i - g''_i \right) + f(g_{m+1} + g''_{m+1}, g'_{m+1} + g''_{m+1})e^{m+1}.$$

Then

$$\begin{aligned} H_{m+1} + g + g'' + h &= H_{m+1} + (f(g_{m+1} + g''_{m+1}, g'_{m+1} + g''_{m+1}) + g_{m+1} + g''_{m+1})e^{d+1} \text{ and} \\ H_{m+1} + g + g'' + h &= H_{m+1} + (f(g_{m+1} + g''_{m+1}, g'_{m+1} + g''_{m+1}) + g'_{m+1} + g''_{m+1})e^{d+1}. \end{aligned}$$

By our definition of f , one of these has the form $H_{m+1} + e^{m+1}$ and one of them has the form $H_{m+1} + ue^{m+1}$ for some $u \notin \{0, 1\}$. Thus one is completely contained in $A(p, n)$, and the other is completely disjoint from $A(p, n)$. Consequently, $K_3[B_r(c_h), B_r(b_{g''}), B_r(a_g)] \subseteq E$ and $K_3[B_r(c_h), B_r(b_{g''}), B_{r'}(a_{g'})] \cap E = \emptyset$, or vice versa. \square

The existence of a metric which satisfies Claims 7.9, 7.10, 7.11, 7.12, 7.13, 7.14 is sufficient to show that $\text{GS}_p(n)$ requires non-binary vdisc_3 -error. In the next subsection, we will show very similar axioms hold in $\text{HP}(N)$, although some weakening will be required.

7.3.2. Special properties of $\text{HP}(N)$. In this subsection we show that $\text{HP}(N)$ satisfies weakened versions of the properties of the prior subsection. We begin by equipping $\text{HP}(N)$ with a metric in the obvious way. Recall that

$$\text{HP}(N) = (\{a_i, b_i, c_i : i \in [N]\}, \{a_i b_j c_k : i + j + k \geq N + 2\}).$$

To ease notation, in Section 7.3.2, we let $A = \{a_i : i \in [N]\}$, $B = \{b_i : i \in [N]\}$, $C = \{c_i : i \in [N]\}$, and $E = E(\text{HP}(N))$.

Lemma 7.15. *For each $X \in \{A, B, C\}$ and $x_i, x_j \in X$, set $d(x_i, x_j) = \frac{|i-j|}{N}$. This makes each of A, B, C into a metric space with distances in $[0, 1]$.*

In an ideal world, we would like to prove all the same facts here for $\text{HP}(N)$ as we showed held for $\text{GS}_p(n)$ in the previous subsection. Indeed, some of the facts will go through largely unchanged. For example, the following is a straightforward analogue of Claim 7.13.

Claim 7.16. *Given $a_i \in A$ and $b_j, b_{j'} \in B$, $N(a_i, b_{j'}) \cap \neg N(a_i, b_j)$ is contained in a closed ball in C ball of radius at most $d(b_{j'}, b_j)$.*

Proof. Without loss of generality, assume $j' < j$. Note that

$$N(a_i, b_j) \cap \neg N(a_i, b_{j'}) = \{c_s : s \in [N + 2 - i - j, N] \cap [0, N + 2 - i - j']\}.$$

If this is empty, let C be the ball of radius 0 centered at c_1 . Otherwise, the above is equal to

$$\{c_s : s \in [N + 2 - i - j, N + 2 - i - j']\},$$

which has size $|j - j'| = d(b_j, b_{j'})N$. This set is clearly contained in a closed ball of radius $|j - j'|/2N$ in C . \square

Other properties from Section 7.3.1 must be tweaked in a more serious way before they will hold in $\text{HP}(N)$. For instance, in $\text{HP}(N)$, a ball $B_r(a_i)$ in A is of the form

$$\{a_j : j \in (\min\{i - rN, 1\}, \max\{i + rN, N\})\}.$$

This could have size about $rN/2$ (e.g. if $i = N$), or it could have size $2rN$ (e.g. if $i = \lfloor N/2 \rfloor$, and $r < 1/2$). Consequently, not all balls with the same radius contain the same number of vertices. However, we can clearly give a range of $\lfloor \frac{r}{2}N \rfloor \leq |B_r(a_i)| \leq 2rN$.

Similarly, not every vertex in $\text{HP}(N)$ has the same degree. For instance, if $i, j \geq N/2$, then $|N(a_i b_j)| = N$, while if $i = j = 1$, then $|N(a_i b_j)| = 1$. However, we can show that most vertices have degree within a certain non-trivial range, and that many (but not all) pairs of vertices are in many edges and non-edges.

Claim 7.17. *Suppose $0 < \tau < 1/4$ and $0 < 3\mu < \tau$. The following hold in $\text{HP}(N)$.*

- (1) *For all $i \in (\mu N, 2\mu N)$ and $j \in (\tau N, (1 - \tau)N)$, $|N(a_i b_j)| \in [\tau N/2, (1 - \tau/2)N]$.*
- (2) *For all $i \in (\mu N, 2\mu N)$, $\min\{|N(a_i)|, |\neg N(a_i)|\} \geq N^2/4$.*
- (3) *For all $j \in (\tau N, (1 - \tau)N)$, $|N(b_j)| \in [\mu\tau N^2/4, (1 - \mu\tau/4)N^2]$.*

Proof. Given $i \in (\mu N, 2\mu N)$ and $j \in (\tau N, (1 - \tau)N)$, note

$$N(a_i b_j) = \{c_k : k \in [N + 2 - i - j, N]\}.$$

Therefore,

$$\tau N/2 < \mu N + \tau N - 2 \leq |N(a_i b_j)| = i + j - 2 < N(1 - \tau + 2\mu) - 2 < N(1 - \tau/2).$$

This shows (1) holds. Further, for all $j \in (\tau N, (1 - \tau)N)$, we have

$$|N(b_j)| \geq |(\mu N, 2\mu N)|\tau N/2 \geq \mu\tau N^2/4,$$

and $|N(b_j)| \leq N^2 - |(\mu N, 2\mu N)|\tau N/2 \leq N^2 - \mu\tau N^2/4$, so (3) holds. For (3), we have that for all $i \in (\mu N, 2\mu N)$,

$$N^2/4 \leq \sum_{j=1}^N \sum_{k=N+2-j-2\mu N} k \leq \sum_{j=1}^N \sum_{k=N+2-j-i} k \leq |N(a_i)| \leq \sum_{j=1}^N \sum_{k=1}^{N-j} k \leq N^2/2$$

as claimed. \square

Claim 7.17 shows that almost all vertices (namely those with index in $(\tau N, (1 - \tau)N)$) have degree αN^2 for some α bounded away from 0 and 1. Further, there are many pairs of vertices with degree βN for some β bounded away from 0 and 1 (namely those where one vertex has index in $(\mu N, 2\mu N)$ and the other has index in $(\tau N, (1 - \tau)N)$).

This lemma, and the observations preceding it, illustrate a general strategy for how to deal with $\text{HP}(N)$ (as compared to $\text{GS}_p(n)$). In particular, we must make the statements more approximate, and also relativize some of the quantifications to certain “special small index sets” (e.g. the vertices with indices in $(\mu N, 2\mu N)$) and “special non-small index sets”

(e.g. the vertices with indices in $(\tau N, (1 - \tau)N)$). To ease notation and emphasize this idea, we define, for $\tau, \mu > 0$,

$$A_{lg}(\tau) = \{a_i : i \in (\tau N, (1 - \tau)N)\} \text{ and} \\ A_{sm}(\mu) = \{a_i : i \in (\mu N, 2\mu N)\}.$$

We define $B_{lg}(\tau)$, $B_{sm}(\mu)$, $C_{lg}(\tau)$, and $B_{sm}(\mu)$ in the same way. We will always use these notions with μ being significantly smaller than τ , in which case $A_{lg}(\tau) \cap A_{sm}(\mu) = \emptyset$. Note that Claim 7.17(1) says that every $ab \in A_{sm}(\mu) \times B_{lg}(\tau)$ has degree βN for some β bounded away from 0 and 1, and Claim 7.17(2) says that every $b \in B_{lg}(\tau) \cup B_{sm}(\mu)$ has degree of αN^2 , for some α bounded away from 0 and 1.

When it comes to partitioning open balls into smaller ones (i.e. proving an analogue of Claim 7.9), things are again more complicated. For instance, unlike in the ultrametric space $\text{GS}_p(n)$, we cannot partition an open ball $B_r(a_i)$ into disjoint open sub-balls. This is because any open ball has the form $\{a_j : j \in I\}$, for some open interval I . Any attempt to partition I into disjoint open intervals will always miss a few points of I , namely the endpoints of the sub-intervals. Problems can also arise due to divisibility issues near the end points of intervals. The following lemma will be used to deal with these issues, and says that we can almost partition an interval into almost evenly sized sub-intervals. Below, d_N denotes the usual metric $d_N(i, j) = |i - j|$ on $[N]$. A ball in this metric space has the form $(x - d, x + d) \cap [N]$ for some $x \in [N]$ and $d \in [N]$.

Lemma 7.18. *For all $0 < r_1 < 1$, there is n_0 so that if $N \geq n_0$, the following holds. Suppose $0 < r_1 < r_2$, and $I = (\alpha, \beta)$ is an open ball of radius $r_2 N$ in $[N]$. Then there is $m \leq 4r_2/r_1$ and a collection \mathcal{P} of m open balls contained in I , each of radius at most $r_1 N$ and at least $r_1 N/3$, such that $|I \setminus (\bigcup \mathcal{P})| \leq 2m$.*

Proof. Assume $N \gg r_1^{-1}, r_2^{-1}$. Set $d_2 = \lceil r_2 N \rceil$ and $d_1 = \lceil r_1 N/3 \rceil$. Let $i \in [N]$ be such that

$$I = (\alpha, \beta) = (i - d_2, i + d_2) \cap [1, N].$$

Note $d_2 + 1 \leq |I| = \beta - \alpha + 1 \leq 2d_2 + 1$. Set $s = \lfloor |\beta - \alpha|/(2d_1) \rfloor$, and for each $j \in [s]$, set $x_j = \alpha + (2j - 1)d_1$. Note that for each $j \in [s]$, $x_j + d_1 < \beta$, so the ball of radius d_1 around x_j is contained in I . However, $I \setminus (\bigcup_{j=1}^s B_{d_1}(x_j))$ could be large if $(x_s + d_1, \beta)$ is large. To make sure this is not an issue, we will combine the last two balls into one ball of a slightly different radius.

By definition of x_{s-1} , $2d_1 < \beta - (x_{s-1} - d_1) \leq 4d_1$. Set $d'_1 = \lfloor (\beta - (x_{s-1} - d_1))/2 \rfloor$, and let $x'_{s-1} = x_{s-1} + d'_1$. Set

$$\mathcal{P} = \{B_{d_1}(x_i) : 1 \leq i \leq s - 1\} \cup \{B_{d'_1}(x'_{s-1})\}.$$

By construction, every element in \mathcal{P} is an open ball of radius at most $r_1 N$ and at least $r_1 N/3$, and

$$|I \setminus (\bigcup \mathcal{P})| \leq |\{\alpha, x_1, \dots, x_{s-2}, x'_{s-1}\} \cup \{\beta, \beta - 1\}| \leq s + 2.$$

By definition $s \leq (2r_2 N + 2)/2d_1 N \leq (2r_2 N + 2)/(2r_1 N/3) \leq 4r_2/r_1$, where the last inequality uses that N is large. \square

We can now give a weak analogue of Claim 7.9.

Claim 7.19. *Suppose $0 < r_1 < \mu^2$, $r_1 < r_2 \leq 1$, N is sufficiently large compared to r_1 , and $a \in A$. Then there is some $m \leq 4r_1/r_2$ and a set \mathcal{P} of m disjoint open balls, each of radius at least $r_1/3$ and at most r_1 , all contained in $B_{r_2}(a)$, such that $|B_{r_2}(a) \setminus (\bigcup \mathcal{P})| \leq 2m$.*

Proof. Note $B_{r_2}(a) = \{a_i : i \in I\}$ for some open ball I in $[N]$. The conclusion is thus an immediate corollary of Lemma 7.18. \square

We recall that in Claim 7.9, there was an additional conclusion beyond what we have shown above, namely that we were also able to choose one of the balls in the cover ahead of time. Due to divisibility issues, we cannot always do this in $\text{HP}(N)$, but it turns out that Claim 7.19 along with Claim 7.20 below will suffice. In particular, Claim 7.20 says that given an open ball $B_r(a)$ in A , whose center is not too close to the endpoints of $[N]$ (where problems arise), we can almost partition A into balls, one of which is $B_r(a)$. To help deal with this requirement, we define the following slightly enlarged versions of our special sets:

$$\begin{aligned} A_{sm}(\mu)^+ &= \{a_i : i \in ((\mu - \mu^2)N, (\mu + \mu^2)N)\} \text{ and} \\ A_{lg}(\tau)^+ &= \{a_i : i \in ((\tau - \mu^2)N, (1 - \tau + \mu^2)N)\}. \end{aligned}$$

We define $B_{sm}(\mu)^+$, $B_{lg}(\mu)^+$ and $C_{sm}(\mu)^+$, $C_{lg}(\mu)^+$ similarly.

Claim 7.20. *Assume $0 < \tau < 1/4$, $0 < \mu < \tau/3$, and $0 < r < \mu^2$. For N sufficiently large, the following holds. For all $a_i \in A_{lg}(\tau)^+ \cup A_{sm}(\mu)^+$, there are $m \leq 8r^{-1}$ and a set \mathcal{P} of m disjoint open balls in A , each of radius at most r and at least $r/3$, such that $B_r(a_i) \in \mathcal{P}$ and $|A \setminus (\bigcup \mathcal{P})| \leq 2m$.*

Proof. Assume $N \gg r^{-1}, \mu^{-1}$. Suppose $a_i \in A_{lg}(\tau)^+ \cup A_{sm}(\mu)^+$ and $0 < r < \mu^2$. By assumption, $i \in ((\tau - \mu^2)N, (1 - \tau + \mu^2)N) \cup ((\mu - \mu^2)N, (2\mu + \mu^2)N)$. Let $d = \lceil rN \rceil$. Note that $B_r(a_i)$ has the form $\{a_i : i \in I\}$ for some open ball $I = (i - d, i + d) \cap [N]$ in $[N]$. Since $a_i \in A_{lg}^+(\tau) \cup A_{sm}^+(\mu)$ and $r < \mu^2$, we know that $i - d > \mu - 2\mu^2$ and $i + d < \tau + 2\mu^2$. Consequently, both $i - d$ and $i + d$ are in $[N]$. Let $I_1 = [1, i - d]$ and $I_2 = (i + d, N]$. These can also be written as open balls in $[N]$, say of radii d_1 and d_2 , respectively. Note that $d_1, d_2 \geq (\mu - 2\mu^2)N/2$.

By Lemma 7.18 there are $m_1 \leq 4\frac{r_1}{N}r^{-1}$ and a collection \mathcal{P}_1 of open balls in $([N], d_N)$, each of radius at least $rN/3$ and at most rN such that each element of \mathcal{P}_1 is contained in $(i + d, N)$ and $|(i + d, N) \setminus (\bigcup \mathcal{P}_1)| \leq 2m_1$. Similarly, there are $m_2 \leq 4r_2r^{-1}$ and a collection \mathcal{P}_2 of open balls in $([N], d_N)$, each of radius at least $rN/3$ and at most rN such that each element of \mathcal{P}_2 is contained in $(1, i - d)$ and $|(1, i - d) \setminus (\bigcup \mathcal{P}_2)| \leq 2m_2$. We then let

$$\mathcal{P} = \{B_r(a_i)\} \cup \mathcal{P}_1 \cup \mathcal{P}_2.$$

Note $|\mathcal{P}| = m_1 + m_2 + 1 \leq 4r^{-1}$. By construction, each element of \mathcal{P} is a ball of radius at least $r/3$ and at most r , and $|A \setminus (\bigcup \mathcal{P})| \leq 2m_1 + 2m_2 + 2 \leq 8r^{-1}$, as desired. \square

We will need one more partition fact, namely that given a ball $B_r(a_i)$ whose center a_i is in $A_{sm}(\mu)$, the approximate subcover obtained in Claim 7.19 will have the property that $B_r(a_i) \cap A_{sm}(\mu)$ is mostly covered by balls whose centers are in $A_{sm}(\mu)^+$.

Claim 7.21. *In Claim 7.19, if \mathcal{P}' are the elements of \mathcal{P} centered at elements in $A_{sm}^+(\mu)$, and \mathcal{P}'' are those centered at $A_{lg}(\tau)^+$, then*

$$|(B_{r_2}(a) \cap A_{sm}(\mu)) \setminus (\bigcup \mathcal{P}')| \leq 2m$$

$$|(B_{r_2}(a) \cap A_{lg}(\tau)) \setminus (\bigcup \mathcal{P}'')| \leq 2m$$

Proof. This is immediate from the definitions of $A_{sm}(\mu)^+$ and $A_{lg}(\tau)^+$. \square

We now turn to the analogues of Claims 7.11 and 7.12. For these we must restrict ourselves to our special sets $A_{sm}(\mu)^+$ and $A_{lg}(\mu)^+$.

Claim 7.22. *Suppose $0 < \tau < 1/4$, $0 < \mu < \tau/3$, and $0 < r < \mu^2$. For all sufficiently large N the following holds. For every $a \in A_{lg}(\tau)^+$ and $b \in B_{sm}(\mu)^+$, there are $f_0(r, a, b), f_1(r, a, b) \in [N]$ such that*

$$K_3[B_{r/2}(c_{f_0(r,a,b)}), B_r(a), B_r(b)] \subseteq E \text{ and } K_3[B_{r/2}(c_{f_1(r,a,b)}), B_r(a), B_r(b)] \cap E = \emptyset,$$

and such that $d(c_{f_0(r,a,b)}, c_{f_1(r,a,b)}) \leq 7r$.

Proof. Suppose $(\tau - \mu^2)N < i < (1 - \tau + \mu^2)N$, $(\mu - \mu^2)N < j < (2\mu + \mu^2)N$, and $0 < r < \mu^2$. Set $d_1 = \lceil rN \rceil$, and define

$$f_0(r, a_i, b_j) = N + 2 - i - j - 3d_1 \text{ and } f_1(r, a_i, b_j) = N + 2 - i - j + 3d_1.$$

Note each of these are in $[N]$ due to the assumptions on i, j , and r . Suppose $a_w b_u c_v \in B_r(a_i) \times B_r(b_j) \times B_{r/2}(c_{f_0(r,a_i,b_j)})$. Then

$$w + u + v < (i + d_1) + (j + d_1) + (N + 2 - i - j - 3d_1) + \lceil d_1/2 \rceil = N + 2 - d_1 + \lceil rN/2 \rceil < N + 2.$$

On the other hand, if $a_w b_u c_v \in B_r(a_i) \times B_r(b_j) \times B_{r/2}(c_{f_1(r,a_i,b_j)})$. Then

$$\begin{aligned} w + u + v &> (i - \lfloor rN \rfloor) + (j - \lfloor rN \rfloor) + (N + 2 - i - j + 3d_1) - \lfloor rN/2 \rfloor \\ &\geq N + d_1 - \lfloor rN/2 \rfloor \\ &> N + 2, \end{aligned}$$

where the last inequality uses the N is large. Note that by construction, $d(c_{f_1(r,a_i,b_j)}, c_{f_0(r,a_i,b_j)}) = 6d_1/N \leq 7r$. \square

As in Section 7.3.1, it is important to know the $f_u(r, a, b)$, $f_v(r', a', b')$ obtained in Claim 7.22 are far for certain choices of a, b, r, a', b', r' .

Claim 7.23. *Assume $0 < \tau < 1/4$, $0 < \mu < \tau/3$, and $0 < r \leq r' < \mu^2$. For any $a, a' \in A_{lg}(\tau)^+$ and $b, b' \in B_{sm}(\mu)^+$ the following hold.*

- (1) *For each $u \in \{0, 1\}$, $d(f_u(r, a, b), f_u(r, a, b')) \geq d(b, b')$ and $d(f_u(r, a', b), f_u(r, a, b)) \geq d(a, a')$.*
- (2) *If $d(a, a') \geq d(b, b') + 6r'$, then*

$$\min\{d(c_{f_u(r,a,b)}, c_{f_v(r',a',b')}) : u, v \in \{0, 1\}\} \geq d(a, a') - d(b, b') - 7r'.$$

(3) If $d(b, b') \geq d(a, a') + 6r'$, then

$$\min\{d(c_{f_u(r, a, b)}, c_{f_v(r', a', b')}) : u, v \in \{0, 1\}\} \geq d(b, b') - d(a, a') - 7r'.$$

Proof. Suppose $a_i, a_{i'} \in A_{lg}(\tau)^+$ and $b_j, b_{j'} \in B_{sm}(\mu)^+$. By definition, for each $u \in \{0, 1\}$,

$$d(f_u(r, a_i, b_j), f_u(r, a_{i'}, b_{j'})) = |i - i'|.$$

Thus $d(c_{f_u(r, a_i, b_j)}, d_{f_u(r, a_{i'}, b_{j'})}) = |i - i'|/N = d(a, a')$. A symmetric argument shows that $d(f_u(r, a, b), f_u(r, a', b')) = d(b, b')$, so (1) holds. For (2), suppose that $d(b_j, b_{j'}) \geq d(a_i, a_{i'}) + 6r'$. Without loss of generality, say $j' > j$. Then for any $(u, v) \in \{0, 1\}^2$,

$$\begin{aligned} d(f_u(r, a, b), f_v(r', a', b')) &= |N + 2 - i - j + (-1)^u 3[rN] - (N + 2 - i' - j' + (-1)^v 3[r'N])| \\ &\geq |j' - j| - |i' - i| - 6[r'N], \end{aligned}$$

where the inequality is using the assumption that $d(b_j, b_{j'}) \geq d(a_i, a_{i'}) + 6r'$, and that $r' \geq r$. Consequently,

$$d(c_{f_u(r, a_i, b_j)}, c_{f_v(r', a_{i'}, b_{j'})}) \geq |j' - j| - |i' - i| - 6[r'N]/N \geq d(b_j, b_{j'}) - d(a_i, a_{i'}) - 7r'.$$

The argument for (3) is similar. \square

We now give the analogue of Claim 7.14. The version here is a bit weaker because the balls are smaller, and we will also require things to be sufficiently spaced out.

Claim 7.24. *Suppose $0 < \tau < 1/4$, $0 < 3\mu < \tau$, $0 < r, r' < \mu^2$ and $r'' \geq \max\{r, r'\}$. Then if N is sufficiently large, the following hold.*

For all $a_i \in A_{lg}^+(\tau)$ and $a_{i'} \in A$ with $d(a, a') = r'' + r' + r$, there is $b \in B_{sm}(\mu)$ and $c \in C$ such that if $r_0 = \min\{r, r', r''\}/3$, then $K_3[B_{r_0/2}(c), B_{r_0/2}(b), B_{r'}(a_{i'})] \subseteq E$ and $K_3[B_{r_0/2}(c), B_{r_0/2}(b), B_r(a_i)] \cap E = \emptyset$, or vice versa.

Proof. Fix $a_i \in A_{lg}^+(\tau)$ and $a_{i'} \in A$ satisfying $d(a, a') = r'' + r' + r$, where $r'' > 2\max\{r', r\}$ and $0 < r, r' < \mu^2$. Set $r_0 = \min\{r, r', r''\}/3$, and choose any $b_j \in B_{sm}(\mu)$ (note this implies $\mu N < j < 2\mu N$).

Suppose first that $i' \in ((\tau - \mu^2)N, N(1 - \tau + \mu^2))$ and $i \leq i'$. Let $k = N + 2 - i' - j + [r'N] + [r_0N]$. Our assumptions imply $i' + j \leq (1 - \tau + \mu)N$ so $k \geq N + 2 - i' - j \geq (\tau - \mu^2 - \mu)N \geq 1$. On the other hand,

$$k \leq N + 2 - (\tau - \mu^2)N - \mu N + 2 + r'N + r_0N \leq N,$$

where the last inequality is because $r', r_0 < \mu^2$ and N is large. Thus $k \in [N]$, so it makes sense to define $c = c_k$. Suppose $b_u c_v \in B_{r_0/2}(b_j) \times B_{r_0/2}(c)$, $a_s \in B_{r'}(a_{i'})$ and $a_t \in B_r(a_i)$. Then using that $i' - i = r + r' + r''$ by assumption, we have

$$\begin{aligned} t + u + v &< (i + rN) + (j + r_0N/2) + (N + 2 - i' - j + [r'N] + [r_0N] + r_0N/2) \\ &= N + 2 + r_0N - (i' - i) + [r'N] + [r_0N] \\ &\leq N + 7 - (r + r' + r'')N - r'N + r_0N \\ &< N + 2, \end{aligned}$$

where the last inequality is because $r'' > 2\max\{r, r'\}$ and N is large. Thus $a_t b_u c_v \notin E$. On the other hand,

$$\begin{aligned} s + u + v &> (i' - r'N) + (j - r_0N/2) + (N + 2 - i' - j + \lceil r'N \rceil + \lceil r_0N \rceil - r_0N/2) \\ &= N + 2 - r_0N + \lceil r'N \rceil + \lceil r_0N \rceil \\ &\geq N + 2. \end{aligned}$$

Thus $a_s b_u c_v \in E$.

If $i' \in ((\tau - \mu^2)N, N(1 - \tau + \mu^2))$ and $i' \leq i$, then proceed as above with the roles of i and i' switched.

Suppose now that $i' < (\tau - \mu^2)N$. In this case, set $k = N + 2 - i - j + \lceil rN \rceil + \lceil r_0N \rceil$. It is straightforward to check that $k \in [N]$ since $i \in ((\tau - \mu^2)N, (1 - \tau + \mu^2)N)$ and $j \in (\mu N, 2\mu N)$. Thus it makes sense to define $c = c_k$. Suppose $b_u c_v \in B_{r_0/2}(b_j) \times B_{r_0/2}(c)$, $a_s \in B_{r'}(a_{i'})$ and $a_t \in B_r(a_i)$. Then

$$\begin{aligned} t + u + v &> (i - rN) + (j - r_0N/2) + (N + 2 - i - j + \lceil rN \rceil + \lceil r_0N \rceil - r_0N/2) \\ &= N + 2 - rN + \lceil rN \rceil - r_0N + \lceil r_0N \rceil \\ &\geq N + 2. \end{aligned}$$

Thus $a_t b_u c_v \in E$. On the other hand

$$\begin{aligned} s + u + v &< (i' + r'N) + (j + r_0N/2) + (N + 2 - i - j + \lceil rN \rceil + \lceil r_0N \rceil + r_0N/2) \\ &= N + 2 - (i - i') + r_0N + \lceil rN \rceil + \lceil r_0N \rceil \\ &\leq N + 2 - \lfloor (r + r' + r'')N \rfloor + \lceil rN \rceil + \lceil r_0N \rceil \\ &\leq N + 2, \end{aligned}$$

where the last inequality is since N is large and $r'' > 2\max\{r, r'\} \geq 6r_0$. Thus $a_s b_u c_v \notin E$.

Finally, consider the case where $i' > N(1 - \tau + \mu^2)$. In this case, set

$$k = N + 2 - i - j - \lceil rN \rceil - \lceil r_0N \rceil.$$

As above, it is straightforward to check that $k \in [N]$. Thus it makes sense to define $c = c_k$. Suppose $b_u c_v \in B_{r_0/2}(b_j) \times B_{r_0/2}(c)$, $a_s \in B_{r'}(a_{i'})$ and $a_t \in B_r(a_i)$. Then

$$\begin{aligned} s + u + v &> (i' - r'N) + (j - r_0N/2) + (N + 2 - i - j - \lceil rN \rceil - \lceil r_0N \rceil - r_0N/2) \\ &= N + 2 + (i' - i) - r'N - \lceil rN \rceil - r_0N - \lceil r_0N \rceil \\ &\geq N + 2 + \lfloor (r + r' + r'')N \rfloor - r'N - \lceil rN \rceil - r_0N - \lceil r_0N \rceil \\ &\geq N + 2, \end{aligned}$$

where the last inequality is because N is large and $r'' > 2\max\{r, r'\} \geq 6r_0$. Thus $a_s b_u c_v \notin E$. On the other hand

$$\begin{aligned} t + u + v &< (i + rN) + (j + r_0N/2) + (N + 2 - i - j - \lceil rN \rceil - \lceil r_0N \rceil + r_0N/2) \\ &= N + 2 + rN - \lceil rN \rceil + r_0N - \lceil r_0N \rceil \\ &\leq N + 2, \end{aligned}$$

thus $a_t b_u c_v \notin E$. This finishes the proof. \square

We will see that the properties described in this section are sufficient for showing that $\text{HP}(N)$ requires non-binary vdisc_3 -error. Further, they are strictly weaker than the properties laid out in Section 7.3.1. For this reason, the axioms in Definition 7.25 will be based on what holds in $\text{HP}(N)$.

7.3.3. Special 3-graphs. In this section we define special 3-graphs, and show that both $\text{HP}(N)$ and $\text{GS}_p(n)$ are special, for appropriately chosen parameters.

Definition 7.25. Suppose $p \geq 2$ and $G = (A \cup B \cup C, E)$ is a 3-partite 3-graph where $|A| = |B| = |C| = N$. We say G is $(p, \mu, \tau, \alpha, \rho)$ -special if there is a metric d on A , B , and C with distances in $[0, 1]$, and distinguished subsets $A_{sm} \subseteq A_{sm}^+$ and $A_{lg} \subseteq A_{lg}^+$ of A , $B_{sm} \subseteq B_{sm}^+$ and $B_{lg} \subseteq B_{lg}^+$ of B , and $C_{sm} \subseteq C_{sm}^+$ and $C_{lg} \subseteq C_{lg}^+$ of C , satisfying the following axioms for all permutations XYZ of ABC , where given $X \in \{A, B, C\}$, $B_r(x) = \{x' \in X : d(x, x') < r\}$.

- (1) (Special Subsets Axiom) There are $r_{sm} \geq \mu/2$, $r_{lg} \geq \tau/2$, and $x_{sm}, x_{lg} \in X$ such that $X_{sm} = B_{r_{sm}}(x_{sm})$, $X_{lg} = B_{r_{lg}}(x_{lg})$, $X_{sm}^+ = B_{r_{sm}+\mu^2}(x_{sm})$, and $X_{lg}^+ = B_{r_{lg}+\mu^2}(x_{lg})$.
- (2) (Size Axiom) For each $r \in (0, 1]$ and $x \in X$, $r|X|/2p \leq |B_r(x)| \leq pr|X|$.
- (3) (Degree Axiom) For all $x \in X_{lg}^+ \cup X_{sm}^+$,

$$\min\{|N(x) \cap K_2[Y, Z]|, |\neg N(x) \cap K_2[Y, Z]|\} \geq \alpha n^2.$$

- (4) (Intersection Axiom) For all $y \in Y$ and $z, z' \in Z$, there is a closed ball of radius at most $d(z, z')$ containing $N(y, z) \cap \neg N(y, z')$.
- (5) (Partition Axiom) For any $\rho < r < \mu^2$ and $x \in X_{lg}^+ \cup X_{sm}^+$, there are $m \leq 4p(r^{-1})$ and a set \mathcal{P} of m disjoint open balls in X of radius at least $r/3$ and at most r , one of which is $B_r(x)$, such that $|X \setminus (\bigcup \mathcal{P})| \leq 2m$.
- (6) (Covering Axiom) Suppose $\rho < r_1 < \mu^2$ and $r_1 < r_2 \leq 1$, and B is a ball of radius r_2 in X . Then there is some $m \leq 2pr_1/r_2$, and a set \mathcal{P} of m disjoint open balls, each of radius at least $r_1/3$ and at most r_1 , all contained in B , such that $|B \setminus (\bigcup \mathcal{P})| \leq 2m$. Moreover, the following hold.
 - (a) If $\mathcal{P}' = \{B_r(x) \in \mathcal{P} : x \in X_{lg}^+\}$, then $|(B \cap X_{lg}) \setminus (\bigcup \mathcal{P}')| \leq 2m$
 - (b) If $\mathcal{P}'' = \{B_r(x) \in \mathcal{P} : x \in X_{sm}^+\}$, then $|(B \cap X_{sm}) \setminus (\bigcup \mathcal{P}'')| \leq 2m$.
- (7) (Splitting Axiom 1) For every $x \in X_{lg}^+$ and $y \in Y_{sm}^+$ and $\rho < r < \mu^2$, there are $f_0(r, x, y)$ and $f_1(r, x, y)$ such that $B_{r/2}(f_u(r, x, y)) \times B_r(x) \times B_r(y) \subseteq E^u$ for each $u \in \{0, 1\}$, and such that $d(f_0(r, x, y), f_1(r, x, y)) \leq 3r$. Further, the following hold.
 - (a) If $y \neq y'$, then for each $u \in \{0, 1\}$, $d(f_u(r, x, y), f_u(r, x, y')) \geq d(y, y')$,
 - (b) If $d(y, y') \geq d(x, x') + 6r'$, then
$$\min\{d(f_u(r, x, y), f_v(r', x', y')) : u, v \in \{0, 1\}\} \geq d(y, y') - d(x, x') - 7r'.$$
 - (c) If $d(x, x') \geq d(y', y) + 6r'$, then
$$\min\{d(f_u(r, x, y), f_v(r', x', y')) : u, v \in \{0, 1\}\} \geq d(x, x') - d(y, y') - 7r'.$$
- (8) (Splitting Axiom 2) For all $x \in X_{lg}^+$ and $x' \in X$ satisfying $d(x, x') = r'' + r' + r$, where $\rho \leq r, r' < \mu^2$ and $r'' > 2 \max\{r', r\}$, and all $y \in Y_{sm}$ there is $z = f(x, x', y) \in Z$

such that if $r_0 = \min\{r, r', r''\}/3$, then $K_3[B_{r_0/2}(y), B_{r_0/2}(z), B_{r'}(x')] \subseteq E$ and $K_3[B_{r_0/2}(y), B_{r_0/2}(z), B_r(x)] \cap E = \emptyset$, or vice versa.

(9) (Disjointness Axiom) The sets X_{sm}^+ and X_{lg}^+ are disjoint.

It seems likely that with a bit more work, one could get rid of Splitting Axiom 2 by deducing it from a version of Splitting Axiom. However, optimizing the list of axioms is not our main goal. We can now easily verify that our main examples are special.

Proposition 7.26. *For all $p \geq 2$, there is n_0 such that for all $\rho > 0$ and all $n \geq n_0$, $\text{GS}_p(n)$ is $(p, 1/p^2, 1/p^2, \frac{1}{2(p-1)}, \rho)$ -special.*

Proof. Equip A , B , and C with the metrics defined in Section 7.3.1. For each $X \in \{A, B, C\}$, set

$$X_{lg} = X_{lg}^+ = \{x_g : g \in H_1 + e_1\} \text{ and } X_{sm} = X_{sm}^+ = \{x_g : g \in H_1 + e_2\}.$$

Clearly these are balls of radius at least p^{-2} (see Claim 7.9), and $X_{sm}^+ \cap X_{lg}^+ = \emptyset$. Thus Axiom (1) holds. Axiom (2) holds by the comments following 7.8. Axioms (3)-(7) hold immediately from Claims 7.9, 7.10, 7.11, 7.12, 7.13, 7.14, and the fact that the roles of A , B and C are symmetric in $\text{GS}_p(n)$. \square

Proposition 7.27. *For all $0 < \tau < 1/4$, $0 < \mu < \tau/3$, $0 < \rho < \mu^3$, and $p \geq 2$, there is n_0 such that for all $N \geq n_0$, $\text{HP}(N)$ is $(p, \mu, 1 - \tau, \mu^2, \rho)$ -special.*

Proof. Choose n_0 sufficiently large for τ , μ , and $r, r', r'' \geq \rho$ in Claims 7.19 and 7.24. Assume $N \geq n_0$ and equip each of A, B, C in $\text{HP}(N)$ the metrics described in Section 7.3.2. Set $x_{lg} = x_{[n/2]}$, $x_{sm} = x_{[3\mu N/2]}$, $r_{sm} = \mu/2$, and $r_{lg} = (1 - \tau)/2$. Then define $X_{sm}, X_{sm}^+, X_{lg}, X_{lg}^+$ to satisfy Axiom of 1 of Definition 7.25. These will each differ from the sets defined as $X_{sm}(\mu), X_{sm}^+(\mu), X_{lg}(\tau), X_{lg}(\tau)^+$ in Section 7.3.2 by at most two points each (note not every interval in $[N]$ is an open ball due to divisibility issues). This difference will not effect the results proved in Section 7.3.2 as N is large. Note Axiom (9) holds by the definition of X_{lg}^+, X_{sm}^+ and the assumptions on τ and μ . Axiom (2) clearly holds by the discussion following Claim 7.17. Then (3)-(7) hold by Claims 7.17, 7.22, 7.23, 7.19, 7.21, 7.16, 7.24, along with the fact that the roles of A , B and C are symmetric in $\text{HP}(N)$. \square

7.4. Proof of Theorems 2.53 and 2.56. In this subsection we give the proofs of Theorems 2.53 and 2.56. We begin with Theorem 7.28 below, which shows that special 3-graphs will not admit vdisc_3 -homogeneous partitions with binary error.

Theorem 7.28. *Suppose $p \geq 2$, $0 < \mu < 1/16$, $0 < \tau < 1 - 4\mu$, and $\alpha > 0$. For all $0 < \epsilon < \alpha\mu/p^{16}$, there is $t_0 \geq 1$, so that for all $T \geq t_0$, the following holds. There is $\rho > 0$ such that for all sufficiently large n , if $G = (A \cup B \cup C, E)$ is a $(p, \mu, \tau, \alpha, \rho)$ -special 3-partite 3-graph with $|A| = |B| = |C| = n$, then G does not admit an ϵ -homogeneous partition with binary error and t parts, for any $t_0 \leq t \leq T$.*

Proof. Fix $p \geq 2$, $0 < \mu < 1/16$, $0 < \tau < 1 - 4\mu$, and $\alpha > 0$. Choose $0 < \epsilon \ll \alpha, \mu, \tau$ and $t_0 \gg \epsilon^{-1}$. Fix $T \geq t_0$. Then choose $0 < \rho \ll T^{-1}p^{-100}\mu\tau$, $n \gg T\rho^{-1}\epsilon^{-1}p^{-1}$ and set $N = 3n$.

Suppose $G = (A \cup B \cup C, E)$ is a 3-partite 3-graph with $|A| = |B| = |C| = n$, and which is $(p, \alpha, \mu, \tau, \rho)$ -special. For ease of notation, set $V = A \cup B \cup C$. Suppose towards a contradiction there are $t_0 \leq t \leq T$ and an equipartition $\mathcal{P} = \{V_1 \cup \dots \cup V_t\}$ of V and $\Sigma \subseteq \binom{[t]}{2}$ so that $|\Sigma| \leq \epsilon t^2$, and for all $ijk \in \binom{[t]}{3}$, if $ij, jk, ik \notin \Sigma$, then $\frac{|E \cap K_3[V_i, V_j, V_k]|}{|K_3[V_i, V_j, V_k]|} \in [0, \epsilon) \cup (1 - \epsilon, 1]$.

Enumerate the sets $A = \{a_i : i \in [n]\}$, $B = \{b_i : i \in [n]\}$, and $C = \{c_i : i \in [n]\}$. For each $X \in \{A, B, C\}$, let $X_{sm}, X_{sm}^+, X_{lg}, X_{lg}^+$ be as in Definition 7.25. Set $\epsilon_3 = \epsilon^{1/8}$. We will use throughout that $t \geq t_0 \gg \epsilon^{-1}$, and that each V_i is very large (since $|V_i| = (3n/t) \pm 1$ and $t \leq T$).

We say that a pair $xy \in \binom{V}{2}$ is a *non- Σ pair* if $xy \in K_2[V_i, V_j]$ for some $V_i V_j \in \binom{[t]}{2} \setminus \Sigma$. On the other hand, we say xy is a *Σ -pair* if $xy \in K_2[V_i, V_j]$ for some $V_i V_j \in \Sigma$. Since $|\Sigma| \leq \epsilon t^2$, the number of non- Σ pairs is at least

$$\binom{|V|}{2} - \sum_{i=1}^t \binom{|V_i|}{2} - \sum_{ij \in \Sigma} |V_i||V_j| \geq \binom{|V|}{2} - |V|^2/t - \epsilon|V|^2 \geq (1 - 2\epsilon) \binom{|V|}{2},$$

there the last inequality uses that $t \geq t_0$, and that $|V|$ is large. For each $1 \leq i \leq t$, set

$$A_i = A^* \cap V_i, \quad B_i = B^* \cap V_i, \quad \text{and} \quad C_i = C^* \cap V_i.$$

Given $i \in [t]$ and $X \in \{A, B, C\}$, we say X_i is *trivial* if $|X_i| < \epsilon_3|V_i|$, and a vertex $x \in V$ is *trivial*, if $x \in X_i$ for some trivial X_i . For each $i \in [t]$, at most two elements of $\{A_i, B_i, C_i\}$ are trivial, and therefore, the number of trivial vertices in V_i is at most $2\epsilon_3|V_i|$. Consequently, at most $4\epsilon_3N$ vertices are trivial. For each $X \in \{A, B, C\}$, let

$$\mathcal{I}_X := \{i \in [t] : X_i \text{ is non-trivial}\}.$$

Since all but at most $4\epsilon_3N$ vertices are non-trivial,

$$(9) \quad \left| \bigcup_{i \in \mathcal{I}_X} X_i \right| \geq (1 - 12\epsilon_3)|X|,$$

and consequently, $|\mathcal{I}_X| \geq (1 - 12\epsilon_3)|X|/(N/t) \geq (1 - 12\epsilon_3)t/3$, for each $X \in \{A, B, C\}$. We now turn to the sets of pairs which are in some sense “error pairs” with respect to the partition $\{V_1, \dots, V_t\}$. In particular, we define

$$\mathcal{S}_1 = \{X_i Y_j : i, j \in [t], XY \in \binom{\{A, B, C\}}{2} \text{ and either } i = j, \text{ or one of } X_i, Y_j \text{ is trivial}\}, \text{ and}$$

$$\mathcal{S}_2 = \{X_i Y_j \notin \mathcal{S}_1 : V_i V_j \in \Sigma\}.$$

Set $\mathcal{S} = \mathcal{S}_1 \cup \mathcal{S}_2$. Note that $|\mathcal{S}_2| \leq 3|\Sigma| \leq 3\epsilon t^2$. Using (9), we have that

$$(10) \quad |\mathcal{S}| \leq 3|\Sigma| + 3t + t \left| \bigcup_{i \notin \mathcal{I}_X} X_i \right|/(N/t) \leq 13\epsilon_3 t^2,$$

where the last inequality uses that $t \gg t_0 \gg \epsilon^{-1}$. We think of the set \mathcal{S} as controlling all potential error triples. Indeed, we will show that any triple $A_i B_j C_k$ avoiding \mathcal{S} will be somewhat homogenous with respect to H . Specifically, we claim that if $A_i B_j C_k$ has $A_i B_j, A_i C_k, B_j C_k \notin \mathcal{S}$, then $A_i B_j C_k$ is $\sqrt{\epsilon}\mu^8$ -homogeneous with respect to H . Suppose

towards a contradiction that $A_i B_j, A_i C_k, B_j C_k \notin \mathcal{S}$, but $|E \cap K_3[A_i, B_j, C_k]|/|A_i||B_j||C_k| \in (\sqrt{\epsilon}\mu^8, 1 - \sqrt{\epsilon}\mu^8)$. Since $A_i B_j, A_i C_k, B_j C_k \notin \Sigma$, by assumption, there is $u \in \{0, 1\}$ such that $|E^u \cap K_3[V_i, V_j, V_k]| \geq (1 - \epsilon)|V_i||V_j||V_k|$. But we also have

$$|K_3[V_i, V_j, V_k] \setminus E^u| \geq \sqrt{\epsilon}|K_3[A_i, B_j, C_k]| \geq \epsilon^{1/2}\mu^8\epsilon_3^3|V_i||V_j||V_k| > \epsilon|V_i||V_j||V_k|,$$

where the last inequality is by our choice of $\epsilon \ll \mu$. This contradicts that $|E^u \cap K_3[V_i, V_j, V_k]| \geq (1 - \epsilon)|V_i||V_j||V_k|$. Thus any triple $A_i B_j C_k$ avoiding \mathcal{S} is $\sqrt{\epsilon}\mu^8$ -homogenous with respect to H .

Eventually, we will contradict (10). We will use the following definition. Given $X \in \{A, B, C\}$ and $i \in [t]$, set

$$Irr(X_i) = \{Y_j Z_k : X_i Y_j, X_i Z_k \notin \mathcal{S}, |E \cap K_3[X_i, Y_j, Z_k]|/|K_3[X_i, Y_j, Z_k]| \in (\epsilon^{1/2}\mu^8, 1 - \epsilon^{1/2}\mu^8)\}.$$

If X_i is non-trivial, then our arguments above imply that if $Y_j Z_k \in Irr(X_i)$, then $Y_j Z_k \in \mathcal{S}$. Consequently, for any non-trivial X_i , $|Irr(X_i)| \leq 13\epsilon_3 t^2$, and thus $|\bigcup_{Y_j Z_k \in Irr(X_i)} Y_j Z_k| \leq 39\epsilon_3 n^2$.

It will be important to know that, based on our assumptions, the pairs in \mathcal{S} are somewhat constrained in their distribution. This is the purpose of our next series of definitions. First, set

$$S_1 = \bigcup_{X_i Y_j \in \mathcal{S}_1} K_2[X_i, Y_j], \text{ and } S_2 = \bigcup_{X_i Y_j \in \mathcal{S}_2} K_2[X_i, Y_j].$$

By (9), we have that

$$\begin{aligned} |S_1| &\leq \sum_{i=1}^t t \binom{V_i}{2} + \sum_{X \in \{A, B, C\}} \sum_{i \in \mathcal{I}_X} |X_i||V \setminus X| \leq t(3n/t)^2 + 2n \sum_{X \in \{A, B, C\}} \sum_{i \in \mathcal{I}_X} |X_i| \\ &\leq 9n^2/t + 6(12\epsilon_3)n^2 \\ &\leq 61\epsilon_3 n^2, \end{aligned}$$

where the last inequality uses that $t \gg \epsilon^{-1}$. Given X_i , set

$$\mathcal{S}_1(X_i) := \{Y_j : X_i Y_j \in \mathcal{S}_1\} \text{ and } S_1(X_i) = \bigcup_{Y_j \in \mathcal{S}_1(X_i)} Y_j.$$

If X_i is non-trivial, then using the above and our bound on the number of non-trivial vertices, we have

$$|S_1(X_i)| \leq |V_i \setminus X_i| + \left| \bigcup_{j \notin \mathcal{I}_Y} Y_j \cup \bigcup_{j \notin \mathcal{I}_Z} Z_j \right| \leq |V_i| + 4\epsilon_3 N \leq 7\epsilon_3(2n) = 14\epsilon_3 n.$$

Because $|\mathcal{S}_2| \leq 3\epsilon t^2$, $|S_2| \leq 9\epsilon n^2$. Similar to the above, given X_i , we set

$$\mathcal{S}_2(X_i) = \{Y_j : X_i Y_j \in \mathcal{S}_2\} \text{ and } S_2(X_i) = \bigcup_{Y_j \in \mathcal{S}_2(X_i)} Y_j.$$

To deduce that $S_2(X_i)$ is small, it is not enough to just know that X_i is non-trivial. This is the motivation of the next definition. Given $X \in \{A, B, C\}$, define

$$Gd(X) := \{x \in X : x \text{ is non-trivial and } |\{y \in V \setminus X : xy \in S_2\}| \leq 3\sqrt{\epsilon}(2n)\}.$$

We say $x \in X$ is *good* if $x \in Gd(X)$. Since $|S_2| \leq 9\epsilon n^2$, we must have that for any $X \in \{A, B, C\}$, $|Gd(X)| \geq (1 - 3\epsilon^{1/2})n$. Given $X \in \{A, B, C\}$ and $i \in [t]$, we say that X_i is *good* if X_i is non-trivial and $|X_i \cap Gd(X)| \geq (1 - (3\epsilon)^{1/3})|X_i|$. Note that for each $X \in \{A, B, C\}$,

$$\left| \bigcup_{X_i \text{ not good}} X_i \right| \leq \left| \bigcup_{i \notin \mathcal{I}_X} X_i \right| + \frac{|X \setminus Gd(X)|}{(3\epsilon)^{1/3}\epsilon_3(3n/t)} \leq 12\epsilon_3|X| + 3\epsilon^{1/2-1/3-1/8}|X| \leq 4\epsilon_3^{1/3}|X|.$$

Thus, most of the elements in X are in a good X_i . We claim that if X_i is good, then $|\mathcal{S}_2(X_i)| \leq 4\epsilon^{5/24}t$. Suppose X_i is good. By definition,

$$\begin{aligned} |\{xy : x \in X, xy \in S_2\}| &\leq (3\sqrt{\epsilon})2n|X_i \cap Gd(X)| + 2n|X_i \setminus Gd(X)| \leq 6\sqrt{\epsilon}n|X_i| + 2n(3\epsilon)^{1/3}|X_i| \\ &\leq 12\epsilon^{1/3}|X_i|n. \end{aligned}$$

Note that if $Y_j \in \mathcal{S}_2(X_i)$, then Y_j is non-trivial so $K_2[X_i, Y_j]$ contains at least $|X_i|\epsilon_3(3n/t)$ many elements of $S_2(X_i)$. Consequently,

$$|\mathcal{S}_2(X_i)| \leq (12\epsilon^{1/3}|X_i|n)/(|X_i|\epsilon_3(3n/t)) = 4\epsilon^{5/24}t.$$

This also shows that if X_i is good, then $|\mathcal{S}_2(X_i)| \leq (3n/t)4\epsilon^{5/24}t = 12\epsilon^{5/24}n$. We have now establishes the required properties about the distribution of \mathcal{S} . We now turn to considering the implications of the axioms from Definition 7.25.

We say that X_i is *mixed-degree* if it is good, and moreover,

$$|\{x \in X_i : \min\{|N(x)|, |\neg N(x)|\} \geq \alpha n^2\}| \geq \epsilon_3^{1/10}|X_i|.$$

Note that by the Degree Axiom, if a set X_i is good and satisfies $|X_i \cap (X_{sm}^+ \cup X_{lg}^+)| \geq \epsilon_3^{1/10}|X_i|$, then X_i is mixed degree. Our next goal is to show that every mixed-degree X_i is mostly contained in an open ball of small radius. Given $\eta \in (0, 1)$, define

$$Mix(\eta, X_i) = \{yz \in YZ : |N_H(yz)|/|X_i| \in (\eta, 1 - \eta)\}.$$

We can already show that if X_i is good, then $Mix(\eta, X_i)$ is not too large for certain values of η .

Claim 7.29. *If X_i is good then $|Mix(\epsilon^{1/4}\mu^4, X_i)| \leq 2\epsilon^{1/4}\mu^4n^2$.*

Proof. Recall that since X_i is non-trivial, for any $X_iY_j, X_iY_j \notin \mathcal{S}$ and $Y_jZ_k \notin Irr(X_i)$, the triple $X_iY_jZ_k$ is $\epsilon^{1/2}\mu^8$ -homogeneous. Thus, for any such Y_j, Z_k , there is some $\eta \in \{0, 1\}$ so that at least $(1 - \epsilon^{1/4}\mu^4)|Y_j||Z_k|$ many elements $yz \in Y_jZ_k$ have $|N_H^\eta(yz) \cap X_i| \geq (1 - \epsilon^{1/4}\mu^4)|X_i|$. Using the size estimates from earlier in the proof, and the fact that X_i is

good, we have

$$\begin{aligned}
|\text{Mix}(\epsilon^{1/4}\mu^4, X_i)| &\leq \epsilon^{1/4}\mu^4 \left| \bigcup_{Y_j Z_k \notin (\mathcal{S}_1(X_i) \cup \mathcal{S}_2(X_i)) \setminus \text{Irr}(X_i)} K_2[Y_j, Z_k] \right| + \\
&\quad + \left| \bigcup_{Y_j Z_k \in \text{Irr}(X_i)} |K_2[Y_j, Z_k]| + n|S_1(X_i)| + n|S_2(X_i)| \right| \\
&\leq \epsilon^{1/4}\mu^4 n^2 + 39\epsilon_3 n^2 + 14\epsilon_3 n^2 + 12\epsilon^{1/3} n^2 \\
&\leq 2\epsilon^{1/4}\mu^4 n^2,
\end{aligned}$$

where the last inequality is because $\epsilon \ll \mu$. \square

Claim 7.29 tells us that, for any mixed degree X_i , there is only a small number of pairs yz with $N(yz) \cap X_i$ and $\neg N(yz) \cap X_i$ both being large. On the other hand, because X_i has mixed degree, its vertices are involved in many edges as well as many non-edges. This suggests that there must be many pairs yz for which $N(yz) \cap X_i$ is most of X_i , as well as many pairs yz for which $\neg N(yz) \cap X_i$ is almost all of X_i . This idea is important for the rest of the proof.

Consider the following sets.

$$\begin{aligned}
\Gamma^0(X_i) &:= \{yz \in K_2[Y, Z] : |(\neg N(yz)) \cap X_i| \geq (1 - 2\epsilon^{1/4})|X_i|\} \text{ and} \\
\Gamma^1(X_i) &:= \{yz \in K_2[Y, Z] : |N(yz) \cap X_i| \geq (1 - 2\epsilon^{1/4})|X_i|\}.
\end{aligned}$$

Note that $\text{Mix}(2\epsilon^{1/4}, X_i) = K_2[Y, Z] \setminus (\Gamma^0(X_i) \cup \Gamma^1(X_i))$. Define

$$\begin{aligned}
Q(X_i) &= \{yzz' \in K_3[Y, Z, Z] : yz \in \Gamma^1(X_i), yz' \in \Gamma^0(X_i)\} \\
&\quad \cup \{yy'z \in K_3[Y, Y, Z] : yz \in \Gamma^1(X_i), y'z \in \Gamma^0(X_i)\}.
\end{aligned}$$

We show in our next claim that if X_i is mixed-degree, then $Q(X_i)$ is nonempty. Later, we will use this, along with the Intersection Axiom, to deduce that every mixed-degree X_i is mostly contained in an open ball of small radius.

Claim 7.30. *If X_i is mixed-degree, then $Q(X_i)$ is nonempty.*

Proof. Fix $X \in \{A, B, C\}$ and suppose X_i is mixed-degree. Let Y, Z be such that $\{X, Y, Z\} = \{A, B, C\}$. Define

$$J_Y = \{Y_j : X_i Y_j \notin \mathcal{S}\} \text{ and } J_Z = \{Z_k : X_i Z_k \notin \mathcal{S}\}.$$

Note that since X_i is non-trivial, $Y_j \in J_Y$ if and only if $j \neq i$, Y_j is non-trivial and $Y_j \notin \mathcal{S}_2(X_i)$. Consequently, since X_i is good,

$$|J_Y| \geq |\mathcal{I}_Y| - 1 - |\mathcal{S}_2(X_i)| \geq |\mathcal{I}_Y| - 1 - 4\epsilon^{5/24}t \geq (1 - 13\epsilon_3)|\mathcal{I}_Y| \geq (1 - 13\epsilon_3)^2 t/3,$$

where the last inequality uses the fact that $|\mathcal{I}_Y| \geq (1 - 12\epsilon_3)t/3$. Similarly $Z_k \in J_Z$ if and only if $k \neq i$, Z_k is non-trivial and $Z_k \notin \mathcal{S}_2(X_i)$. A similar argument also shows

$|J_Z| \geq (1 - 13\epsilon_3)|\mathcal{I}_Z| \geq (1 - 13\epsilon_3)^2 t/3$. Define

$$F_0 = \{Y_j Z_k \in K_2[J_Y, J_Z] : |E \cap K_3[X_i, Y_j, Z_k]|/|X_i||Y_j||Z_k| \leq \epsilon^{1/2}\} \text{ and}$$

$$F_1 = \{Y_j Z_k \in K_2[J_Y, J_Z] : |E \cap K_3[X_i, Y_j, Z_k]|/|X_i||Y_j||Z_k| \geq 1 - \epsilon^{1/2}\}.$$

Note that by the definition of J_Y, J_Z , we have that if $Y_j Z_k \in K_2[J_Y, J_Z] \setminus (F_1 \cup F_0)$, then $Y_j Z_k \in \text{Irr}(X_i)$.

We claim that it suffices to show that either there is some $Y_j Z_k Z_{k'}$ with $Y_j Z_k \in F_0$ and $Y_j Z_{k'} \in F_1$, or there is some $Y_j Y_{j'} Z_k$ with $Y_j Z_k \in F_0$ and $Y_{j'} Z_k \in F_1$. Indeed, suppose there exists some $Y_j Z_k Z_{k'}$ with $Y_j Z_k \in F_0$ and $Y_j Z_{k'} \in F_1$. Since $Y_j Z_k \in F_0$, standard counting arguments imply that there is a set $\Omega_0 \subseteq Y_j$ of size at least $(1 - \epsilon^{1/4})|Y_j|$ such that for all $y \in \Omega_0$,

$$|\{zx \in K_2[Z_k, X_i] : xyz \notin E\}| \geq (1 - \epsilon^{1/4})|X_i||Z_k|.$$

Similarly, since $Y_j Z_{k'} \in F_1$, there is a set $\Omega_1 \subseteq Y_j$ of size at least $(1 - \epsilon^{1/4})|Y_j|$ such that for all $y \in \Omega_1$,

$$|\{zx \in K_2[Z_k, X_i] : xyz \in E\}| \geq (1 - \epsilon^{1/4})|X_i||Z_k|.$$

Choose any $y_\star \in \Omega_1 \cap \Omega_0$ (clearly $\Omega_0 \cap \Omega_1$ is nonempty, as its size is at least $(1 - 2\epsilon^{1/4})|Y_j|$). Consider the bipartite graphs

$$G_0 = (Z_k \cup X_i, N_H(y_\star) \cap K_2[Z_k, X_i]) \text{ and } G_1 = (Z_{k'} \cup X_i, N_H(y_\star) \cap K_2[Z_{k'}, X_i]).$$

By construction, the density of edges in G_0 is at most $\epsilon^{1/4}$, and the density of edges in G_1 is at least $1 - \epsilon^{1/4}$. Thus, there exists a $z \in Z_k$ of degree at most $\epsilon^{1/4}|X_i|$ in G_0 , and there exists a $z' \in Z_{k'}$ with degree at least $(1 - \epsilon^{1/4})|X_i|$ in G_1 . Now

$$\min\{|\neg N_H(y_\star z) \cap X_i|, |N_H(y_\star z') \cap X_i|\} = \min\{|N_{G_1}(z')|, |N_{G_0}(z)|\} \geq (1 - \epsilon^{1/4})|X_i|,$$

so $y_\star z z' \in Q(X_i)$. An symmetric argument shows that if there is some $Y_j Y_{j'} Z_k$ with $Y_j Z_k \in F_0$ and $Y_{j'} Z_k \in F_1$, then there exists some $yy' z_\star \in Q(X_i)$. Thus we just need to show that either there is some $Y_j Y_{j'} Z_k$ with $Y_j Z_k \in F_0$ and $Y_{j'} Z_{k'} \in F_1$, or there is some $Y_j Z_k Z_{k'}$ with $Y_j Z_k \in F_0$ and $Y_{j'} Z_k \in F_1$.

Suppose towards a contradiction that no such $Y_j Y_{j'} Z_k$ or $Y_j Z_k Z_{k'}$ exist. Let \mathcal{G} be the bipartite graph \mathcal{G} with vertex set $J_Y \cup J_Z$ and edge set F_1 . Recall that for all $Y_j Z_k \in (J_Y \times J_Z) \setminus (F_0 \cup F_1)$, we must have $Y_j Z_k \in \text{Irr}(X_i)$. Consequently,

$$|K_2[J_Y, J_Z] \setminus (F_0 \cup F_1)| \leq 13\epsilon_3 t^2 \leq 100\epsilon_3 |J_Y||J_Z|,$$

where the last inequality uses our lower bounds on $|J_Y|, |J_Z|$. Therefore, if we set

$$J'_Y = \{Y_j \in J_Y : |\{Z_k \in J_Z : Y_j Z_k \notin F_0 \cup F_1\}| \leq \sqrt{100\epsilon_3}|J_Z|\} \text{ and}$$

$$J'_Z = \{Z_k \in J_Z : |\{Y_j \in J_Y : Y_j Z_k \notin F_0 \cup F_1\}| \leq \sqrt{100\epsilon_3}|J_Y|\},$$

then $|J'_Y| \geq \sqrt{100\epsilon_3}|J_Y|$ and $|J'_Z| \geq \sqrt{100\epsilon_3}|J_Z|$. Because there does not exist $Y_j, Z_k, Z_{k'}$ with $Y_j Z_k \in F_0$ and $Y_j Z_{k'} \in F_1$, we have that for all $Y_j \in J'_Y$, either $|N_{\mathcal{G}}(Y_j)| \geq (1 - \sqrt{100\epsilon_3})|J_Z|$ or $|N_{\mathcal{G}}(Y_j)| = 0$. Similarly, because there does not exist $Y_j, Y_{j'}, Z_k$ with $Y_j Z_k \in F_0$ and $Y_{j'} Z_k \in F_1$, we have that for all $Z_k \in J'_Z$, either $|N_{\mathcal{G}}(Z_k)| \geq$

$(1 - \sqrt{100\epsilon_3})|J_Z|$ or $|N_{\mathcal{G}}(Z_k)| = 0$. By Lemma 5.9, either $|F_1| \leq 2(100\epsilon_3)^{1/4}|J_Y||J_Z|$ or $|F_1| \geq (1 - 2(100\epsilon_3)^{1/4})|J_Y||J_Z|$.

Suppose first that $|F_1| \leq 2(100\epsilon_3)^{1/4}|J_Y||J_Z|$. Then

$$\begin{aligned}
\left| \bigcup_{x \in X_i} N_H(x) \right| &\leq |X_i| \left| \bigcup_{Y_j Z_k \notin F_0 \cup F_1} K_2[Y_j, Z_k] \right| + \sum_{Y_j Z_k \in F_1} |X_i||Y_j||Z_k| + \sum_{Y_j Z_k \in F_0} |E \cap K_3[X_i, Y_j, Z_k]| \\
&\leq |X_i| 39\epsilon_3 n^2 + |F_1|(3n/t)^2 + \epsilon^{1/2}|J_Y||J_Z|(3n/t)^2 \\
&\leq |X_i| \left(39\epsilon_3 n^2 + 3(100\epsilon_3)^{1/4} t^2 (3n/t)^2 + \epsilon^{1/2} t^2 (3n/t)^2 \right) \\
&= |X_i| n^2 \left(39\epsilon_3 n^2 + 27(100\epsilon_3)^{1/4} + 9\epsilon^{1/2} \right) \\
&\leq \epsilon_3^{1/5} |X_i| n^2,
\end{aligned}$$

where the last inequality is because ϵ is small. This implies that there is a set $X'_i \subseteq X_i$ such that $|X'_i| \geq (1 - \epsilon_3^{1/10})|X_i|$, and such that for all $x \in X'_i$, $|N_H(x)| \leq \epsilon_3^{1/10} n^2 < \alpha n^2$, but this contradicts that X_i is mixed degree.

Suppose on the other hand that $|F_1| \geq (1 - 2(100\epsilon_3)^{1/4})|J_Y||J_Z|$. Note that

$$\begin{aligned}
\left| \bigcup_{Y_j Z_k \notin F_1} K_2[Y_j, Z_k] \right| &\leq \left| \bigcup_{Y_j Z_k \notin F_0 \cup F_1} K_2[Y_j, Z_k] \right| + \left| \bigcup_{Y_j Z_k \in F_0} K_2[Y_j, Z_k] \right| \\
&\leq 39\epsilon_3 n^2 + 2(100\epsilon_3)^{1/4} t^2 \left(\frac{3n}{t} \right)^2 \\
&\leq \epsilon_3^{1/5} n^2 / 2,
\end{aligned}$$

where the last inequality is because ϵ is small. Consequently, we have that

$$\left| \bigcup_{x \in X_i} N_H(x) \right| \geq (1 - 2\epsilon^{1/4}) \sum_{Y_j Z_k \in F_1} |Y_j||Z_k| \geq (1 - \epsilon^{1/2}) n^2 (1 - \epsilon_3^{1/5} / 2) \geq n^2 (1 - \epsilon_3^{1/5}).$$

From this we deduce there is a set $X'_i \subseteq X_i$ of size at least $(1 - \epsilon_3^{1/10})|X_i|$, so that for all $x \in X'_i$, $|N_H(x)| \geq (1 - \epsilon_3^{1/10}) n^2 > (1 - \alpha) n^2$, contradicting that X_i is mixed-degree. This finishes the proof of Claim 7.30. \square

We now deduce that every mixed-degree X_i is mostly contained in an open ball. Given a mixed-degree X_i , define

$$D(X_i) = \{d > 0 : \text{there is } yzz' \in Q(X_i) \text{ with } d(z, z') = d \text{ or } yy'z \in Q(X_i) \text{ with } d(y, y') = d\}.$$

Since $D(X_i)$ is nonempty by Claim 7.30 (and V is finite), we may choose $d_{X_i} := \min D(X_i)$ and define a ball B_{X_i} as follows. If d_{X_i} is witnessed by some $yy'z' \in Q(X_i)$, let $\rho_{X_i} \leq d_{X_i}$ be minimal such that for some x , $N(yz) \cap \neg N(y', z) \subseteq \overline{B_{\rho_{X_i}}(x)}$ (such an x and ρ_{X_i} exist by the Intersection Axiom). If on the other hand, d_{X_i} is witnessed by some $yy'z' \in Q(X_i)$, let $\rho_{X_i} \leq d_{X_i}$ be minimal such that for some x , $N(yz) \cap \neg N(y, z') \subseteq \overline{B_{\rho_{X_i}}(x)}$ (again such an x and ρ_{X_i} exists by the Intersection Axiom). In either case, set $B_{X_i} = \overline{B_{\rho_{X_i}}(x)}$, and note

$|X_i \cap B_{X_i}| \geq (1 - 2\epsilon^{1/4})|X_i|$. Combining this with the Size Axiom and the fact that X_i is non-trivial (since it is good) we have that

$$(1 - 2\epsilon^{1/4})\epsilon_3(3n/t) \leq |B_{X_i}| \leq p\rho_{X_i}n,$$

and thus, $\rho_{X_i} \geq (1 - 2\epsilon^{1/4})t^{-1}p^{-1}\epsilon_3 > 50\rho p^{50}$, where the last inequality is because $t \leq T$ and by our choice of ρ .

We say X_i is *sm-interesting* if it is good and $|X_i \cap X_{sm}| \geq \epsilon_3^{1/10}|X_i|$. Similarly, we say X_i is *lg-interesting* if it is good and $|X_i \cap X_{lg}| \geq \epsilon_3^{1/10}|X_i|$. We will say X_i is simply *interesting* if it is either sm-interesting or lg-interesting. Note that by the Degree Axiom, any interesting X_i is automatically mixed-degree. Therefore, when X_i is sm-interesting, we can apply Claim 7.30 to find B_{X_i} . In this case, we have that $|X_i \cap B_{X_i}| \geq (1 - 2\epsilon^{1/4})|X_i|$ and $|X_i \cap X_{sm}| \geq \epsilon_3^{1/10}|X_i|$. This implies that

$$|X_i \cap B'_{X_i} \cap X_{sm}| \geq (\epsilon_3^{1/10} - 2\epsilon^{1/4})|X_i| \geq \epsilon_3^{1/10}|X_i|/2 > 0.$$

Thus, we may choose some $x_i \in X_i \cap B_{X_i} \cap X_{sm}$, and define $B'_{X_i} = B_{2\rho_{X_i}}(x_i)$.

Similarly, when X_i is lg-interesting, $|X_i \cap B_{X_i}| \geq (1 - 2\epsilon^{1/4})|X_i|$ and $|X_i \cap X_{lg}| \geq \epsilon_3|X_i|$ imply

$$|X_i \cap B'_{X_i} \cap X_{lg}| \geq (\epsilon_3^{1/10} - 2\epsilon^{1/4})|X_i| \geq \epsilon_3^{1/10}|X_i|/2 > 0.$$

Thus, we may choose some $x_i \in X_i \cap B_{X_i} \cap X_{lg}$, and define $B'_{X_i} = B_{2\rho_{X_i}}(x_i)$.

We have now shown that when X_i is sm-interesting (respectively lg-interesting), there is an open ball B'_{X_i} of radius $\rho'_{X_i} := 2\rho_{X_i} \leq 2d_{X_i}$, with center $x_i \in X_{sm}$ (respectively $x_i \in X_{lg}$), and so that $|X_i \cap B'_{X_i}| \geq (1 - 2\epsilon^{1/4})|X_i|$. Our next goal is to show that this ρ'_{X_i} is not too big.

Claim 7.31. *If X_i is interesting, then $\rho'_{X_i} \leq \mu^2/2$.*

Proof. Suppose towards a contradiction there exists an interesting X_i with $\rho'_{X_i} > \mu^2/2$. If X_i is sm-interesting, set $X_* = X_{sm}$ and $Y_* = Y_{lg}$ if X_i is lg-interesting, set $X_* = X_{lg}$ and $Y_* = Y_{sm}$.

By the Covering Axiom, there exists \mathcal{Q}_1 a collection of m_1 open balls of radius, each of radius at most $\mu^2/32$, and at least $\mu^2/96$, each contained in B'_{X_i} , such that $m_1 \leq 2p(\mu^2/32)(\rho'_{X_i})^{-1}$, and such that if \mathcal{Q}'_1 is set of elements of \mathcal{Q}_1 centered at a point in X_*^+ , then $|(B'_{X_i} \cap X_*) \setminus (\bigcup \mathcal{Q}'_1)| \leq 2m_1$. Note that by assumption, $m_1 \leq 2p(\mu^2/32)(\rho'_{X_i})^{-1} \leq 2p$. By the pigeonhole principle, there is $B_{r_1}(x') \in \mathcal{Q}'_1$ such that

$$|B_{r_1}(x') \cap X_* \cap X_i| \geq (\epsilon_3^{1/10}|X_i|/2 - 2m_1)/m_1 \geq \epsilon_3^{1/10}|X_i|/4p,$$

where the last inequality is because $m_1 \leq 2p \ll |X_i|$. We claim

$$|B_{\mu^2/16}(x') \cap X_i| < (1 - 2\epsilon^{1/4})|X_i|.$$

Suppose towards a contradiction $|B_{\mu^2/16}(x') \cap X_i| \geq (1 - 2\epsilon^{1/4})|X_i|$. Choose any $y \in Y_*$. By Splitting Axiom 1, there exist $z_0, z_1 \in Z$ such that $d(z_0, z_1) \leq 3\mu^2/16$ such that $B_{\mu^2/16}(x') \subseteq N_H(z_1y) \cap \neg N_H(z_0y)$. But now $yz_1z_0 \in Q(X_i)$, so $d(z_1, z_0) \in D(X_i)$. But $d(z_0, z_1) < \mu^2/4 < \rho'_{X_i}/2 = \rho_{X_i}$, a contradiction to the definition of ρ_{X_i} .

Thus $|B_{\mu^2/16}(x') \cap X_i| < (1 - 2\epsilon^{1/4})|X_i|$. By the Partition Axiom, there is a set \mathcal{Q}_2 of m_2 open balls, each of radius at most $\mu^2/16$, and at least $\mu^2/48$, one of which is $B_{\mu^2/16}(x')$, such that $m_2 \leq 16 \cdot 4p\mu^{-2}$, and such that $|X \setminus (\bigcup \mathcal{P})| \leq 2m_2$. Since $|B_{\mu^2/16}(x') \cap X_i| < (1 - 2\epsilon^{1/4})|X_i|$, by pigeonhole, there is some $B_{r_2}(x'') \in \mathcal{Q}_2$, distinct from $B_{\mu^2/16}(x')$, such that

$$|X_i \cap B_{r_2}(x'')| \geq (2\epsilon^{1/4}|X_i| - 2m_2)/m_2 \geq \frac{\epsilon^{1/4}\mu^2|X_i|}{16 \cdot 2p} - 2 \geq \frac{\epsilon^{1/4}\mu^2|X_i|}{16 \cdot 4p},$$

where the last inequalities use the upper bound on m_2 , and the fact that X_i is large. By the Covering Axiom, there exists a collection \mathcal{Q}_3 of m_3 disjoint open balls, each of radius at most $r_2/4$ and at least $r_2/12$, each contained in $B_{r_2}(x'')$, such that $m_3 \leq 8p$, and such that $|B_{r_2}(x'') \setminus (\bigcup \mathcal{Q}_3)| \leq 2m_3$. By the pigeonhole principle, there is $B_{r_3}(x''') \in \mathcal{Q}_3$ such that

$$|X_i \cap B_{r_3}(x''')| \geq (\epsilon^{1/4}\mu^2|X_i|/(16 \cdot 4p) - 2m_3)/m_3 \geq \frac{\epsilon^{1/4}\mu^2|X_i|}{16 \cdot 32p^2} - 2 \geq \epsilon^{1/4}\mu^4|X_i|,$$

where the last inequalities use the upper bound on m_3 , the fact that $\epsilon \ll \mu \ll 1$, and the fact that X_i is large. Note that $r_2 > r_3 \geq \rho$.

Now, since $B_{\mu^2/16}(x')$ is disjoint from $B_{r_2}(x'')$ and $B_{r_3}(x''') \subseteq B_{r_2}(x'')$, we have that $d(x''', x') \geq \mu^2/16 + r_3$. Set $r'' = d(x''', x') - r_1 - r_3$, and $r_0 = \min\{r_1, r_3\}/3$. Recall $r_1 \leq \mu^2/32$ and $\mu^2/48 \cdot 12 \leq r_2/12 \leq r_3 \leq r_2/4 \leq \mu^2/16 \cdot 4$. Consequently,

$$d(x''', x') \geq \mu^2/16 + r_3 > 2 \max\{r_1, r_3\}.$$

Therefore, by Splitting Axiom 2, there are is an open ball B_1 in Y and an open ball B_2 in Z , each of radius $r_0/3$, such that $K_3[B_1, B_2, B_{r_1}(x')] \subseteq E$ and $K_3[B_1, B_2, B_{r_3}(x'')] \subseteq \neg E$, or vice versa. Since

$$|B_{r_1}(x') \cap X_i| \geq \epsilon_3^{1/10}|X_i|/4p \text{ and } |B_{r_3}(x'') \cap X_i| \geq \epsilon^{1/4}\mu^4|X_i|,$$

we must have that $K_2[B_1, B_2] \subseteq \text{Mix}(\epsilon^{1/4}\mu^4, X_i)$. Since $r_1 \geq \mu^2/96$ and $r_3 \geq \mu^2/(48 \cdot 12)$, we have $r_0/3 \geq \mu^2/(48 \cdot 36)$, so the Size Axiom implies we have

$$\min\{|B_1|, |B_2|\} \geq \mu^2 n / (48 \cdot 36)(2p) = \mu^2 n / (96 \cdot 36p).$$

Consequently

$$|\text{Mix}(\epsilon^{1/4}\mu^4, X_i)| \geq |K_2[B_1, B_2]| \geq \frac{\mu^4 n^2}{(96 \cdot 36p)^2}.$$

However, since X_i is good, Claim 7.29 implies $|\text{Mix}(\epsilon^{1/4}\mu^4, X_i)| \leq 2\epsilon^{1/4}\mu^4 n^2$. Because we chose $\epsilon \ll \mu$, $2\epsilon^{1/4}\mu^4 n^2 < \frac{\mu^4 n^2}{(96 \cdot 36p)^2}$, so this is a contradiction. This finishes the proof of Claim 7.31. \square

Note that Claim 7.31 tells us that if X_i is sm-interesting, then since the center of B'_{X_i} is in X_{sm} and the radius of B'_{X_i} is less than $\mu^2/2$, $B'_{X_i} \subseteq X_{sm}^+$. Consequently $|B'_{X_i} \cap X_i \cap X_{sm}^+| \geq (1 - 2\epsilon^{1/4})|X_i|$. Similarly if X_i is lg-interesting, then $B'_{X_i} \subseteq X_{lg}^+$, and consequently $|B'_{X_i} \cap X_i \cap X_{lg}^+| \geq (1 - 2\epsilon^{1/4})|X_i|$.

Let K_X^{lg} be the set of lg-interesting X_i , and let K_X^{sm} be the set of sm-interesting X_i . Using the above, we have that

$$\begin{aligned} |X_{sm} \setminus (\bigcup_{X_i \notin K_X^{sm}} X_i)| &\leq |\bigcup_{X_i \text{ not good}} X_i| + \sum_{X_i \text{ good}, X_i \notin K_X^{sm}} |X_i \cap X_{sm}| \leq 2\epsilon_3^{1/3}|X| + 2\epsilon^{1/4}|X| \\ &\leq 3\epsilon_3^{1/3}|X_{sm}|, \end{aligned}$$

where the last inequality is because $\epsilon \ll \mu$, and because, by the Size and Special Sets Axioms, $|X_{sm}| \geq \mu|X|/3p$. Consequently,

$$(11) \quad |\bigcup_{X_i \in K_X^{sm}} X_i| \geq |X_{sm}|(1 - 4\epsilon_3^{1/3}).$$

A similar argument shows

$$(12) \quad |\bigcup_{X_i \in K_X^{lg}} X_i| \geq |X_{lg}|(1 - 4\epsilon_3^{1/3}).$$

Thus, for each $X \in \{A, B, C\}$, most of X_{sm} is in an sm-interesting X_i , and most of X_{lg} is in an lg-interesting X_i .

Claim 7.32. *For each $X \in \{A, B, C\}$ and all interesting X_i , and for each $u \in [3]$, there is a set $\phi_u(X_i)$ of open balls in X of radius at most ρ'_{X_i}/p^{2+u} , and at least $\rho'_{X_i}/3p^{2+u}$ such that the following holds for each $u \in [3]$.*

- (1) For all $B \in \phi_u(X_i)$, $B \subseteq B'_{X_i}$,
- (2) For each $B \in \phi_u(X_i)$, $|X_i \cap B| \geq (1 - 2\epsilon^{1/4})|X_i|/p^{3+u}$,
- (3) For each $u \in [3]$, $|X_i \cap (\bigcup_{B \in \phi_u(X_i)} B)| \geq (1 - 3\epsilon^{1/4})|X_i|$.

Proof. Suppose X_i is interesting. If X_i is lg-interesting, let $X_* = X_{lg}$ and if sm-interesting, let $X_* = X_{sm}$.

By the Covering Axiom, for each $u \in [3]$, there is a set \mathcal{P}_u of $m_u \leq 2p^{3+u}$ disjoint open balls, each of some radius $r_u \leq \rho'_{X_i}/p^{2+u}$, such that $|B'_{X_i} \setminus (\bigcup \mathcal{P}_u)| \leq 2m_u$, and such that each ball in \mathcal{P}_u is contained in B'_{X_i} . For each $u \in [3]$, let

$$\phi_u(X_i) = \{B_{\rho'_{X_i}/p^{2+u}}(x') : B_{\rho'_{X_i}/p^{2+u}}(x') \in \mathcal{P}_u \text{ and } |B_{\rho'_{X_i}/p^{2+u}}(x') \cap X_i| \geq (1 - 2\epsilon^{1/4})|X_i|/m_u\}.$$

Note (1) and (2) hold by definition of $\phi_u(X_i)$. We show (3). By, assumption, $|B'_{X_i} \setminus (\bigcup \mathcal{P}_u)| \leq 2m_u$. Thus,

$$|X_i \cap (\bigcup_{B' \in \phi_u(X_i)} B')| \geq |X_i \cap B'_{X_i}| - 2m_u \geq (1 - 2\epsilon^{1/4})|X_i| - 4p^{3+u} \geq (1 - 3\epsilon^{1/4})|X_i|,$$

where the last inequality is because X_i is large. \square

For ease of notation, given an interesting X_i and $u \in \{1, 2, 3\}$, let $P_u(X_i)$ be the set of centers of the balls in $\phi_u(X_i)$, and let $\Psi_u(X_i) = \bigcup_{B' \in \phi_u(X_i)} B'$. Note that $P_u(X_i) \subseteq X_{sm}^+$ when X_i is sm-interesting, and $P_u(X_i) \subseteq X_{lg}^+$ when X_i is lg-interesting.

Our next claim is crucial to the overall argument. The basic idea is that for certain sm-interesting Y_j , and any lg-interesting X_i , then, given any $B' \in \phi_2(X_i)$ and $B'' \in \phi_2(Y_j)$, there is a ball $B''' = B'''(B', B'')$ in Z , which is not too small, and such that $K_2[B'', B''']$ contains many xz which “split” $Y_j \cap B''$, in the sense that both $N(xz) \cap (Y_j \cap B')$ and $\neg N(xz) \cap (Y_j \cap B'')$ are large. It will also be important to know this produces different B''' as we let B' and B'' vary (see (i) and (ii) below).

Claim 7.33. *Suppose $X \neq Y \in \{A, B, C\}$, Y_j is sm-interesting and X_i is lg-interesting.*

- (1) *Suppose $\rho'_{X_j} \geq \rho'_{Y_i}$. Then for each $x_0 \in P_2(X_i)$, and each $y_0 \in P_2(Y_j)$, there is $f_{X_i Y_j Z}(x_0 y_0) \in Z$ so that for at least half the $x \in B_{\rho'_{X_i}/2p^4}(x_0) \cap X_i$, and all the $z \in B_{\rho'_{X_i}/2p^4}(f_{Y_j X_i Z}(x_0 y_0))$, $xz \in \text{Mix}(2\epsilon^{1/4}/p^5, Y_j)$.*
- (2) *Suppose $\rho'_{Y_j} \geq \rho'_{X_i}$. Then for each $x_0 \in P_2(X_i)$, and each $y_0 \in P_2(Y_j)$, there is $f_{Y_j X_i Z}(y_0 x_0) \in Z$ so that for at least half the $y \in B_{\rho'_{Y_j}/2p^4}(y_0) \cap Y_j$, and all the $z \in B_{\rho'_{Y_j}/2p^4}(f_{X_i Y_j Z}(y_0 x_0))$, $yz \in \text{Mix}(2\epsilon^{1/4}/p^5, X_i)$.*

Moreover, the following hold.

- (i) *If X_i, X_i are lg-interesting, and $\rho'_{X_i} \geq \rho'_{X_{i'}} \geq \rho'_{Y_j}$, then for all $x \in P_2(X_i)$, $x' \in P_2(X_{i'})$ with $d(x, x') \geq 10\rho_{X_i}$, and all $y, y' \in P_2(Y_j)$,*

$$d(f_{X_i Y_j Z}(xy), f_{X_{i'} Y_j Z}(x'y')) \geq 5\rho'_{X_i}.$$

- (ii) *If $Y_j, Y_{j'}$ are sm-interesting, and $\rho'_{Y_j} \geq \rho'_{Y_{j'}} \geq \rho'_{X_i}$, then for all $y \in P_2(Y_j)$, $y' \in P_2(Y_{j'})$ with $d(y, y') \geq 10\rho'_{Y_j}$, and all $x, x' \in P_2(X_i)$,*

$$d(f_{Y_j X_i Z}(yx), f_{Y_{j'} X_i Z}(y'x')) \geq 5\rho'_{Y_j}.$$

Proof. Assume first that $\rho'_{X_i} \geq \rho'_{Y_j}$. Suppose $y_0 \in P_2(Y_j)$ and $x_0 \in P_2(X_i)$. By Claim 7.31, $\rho'_{X_i} \leq \mu^2/2$. Set $r = \rho'_{X_i}/4p^2$ and let $B' = B_r(y_0)$ and $B'' = B_r(x_0)$. Apply Splitting Axiom 1 to obtain $z_0 = f_0(r, x_0, y_0)$ and $z_1 = f_1(r, x_0, y_0)$, so that so that for each $\eta \in \{0, 1\}$, $B_{r/2}(z_\eta) \times B' \times B'' \subseteq E^\eta$ and $d(z_0, z_1) \leq 3r$. Suppose there were $y \in Y$, $z \in B_{r/2}(z_0)$, and $z' \in B_{r/2}(z_0)$ such that $(y, z) \in \Gamma_1(X_i)$ and $(y, z') \in \Gamma_0(X_i)$, or vice versa. Then we would have $yz z' \in Q(X_i)$, contradicting our choice of ρ_{X_i} (since $d(z, z') < 4r \leq \rho_{X_i}$). Thus, for all $y \in Y$, there is $\xi = \xi(y) \in \{0, 1\}$ such that

$$(\{y\} \times B_{r/2}(z_1)) \cup (\{y\} \times B_{r/2}(z_0)) \subseteq \text{Mix}(2\epsilon^{1/4}, X_i) \cup \Gamma_\xi(X_i).$$

Let $B(1) = \{y \in B' \cap Y_j : \xi(y) = 1\}$ and $B(0) = \{y \in B' \cap Y_j : \xi(y) = 0\}$. Then

$$\max\{|B(0)|, |B(1)|\} \geq |B' \cap Y_j|/2.$$

Choose $\xi^* \in \{0, 1\}$ so that $|B(\xi^*) \cap Y_j| \geq |B' \cap Y_j|/2$. Then for all $y \in B(\xi^*) \cap Y_j$, and all $z \in B_{|1-\xi^*|}(z_{|1-\xi^*|})$, one of the following holds.

- $yz \in \text{Mix}(2\epsilon^{1/4}, X_i)$. In this case, by definition, $\min\{|N(yz) \cap X_i|, |\neg N(yz) \cap X_i|\} \geq 2\epsilon^{1/4}|X_i|$,

- $yz \in \Gamma_{\xi^*}(X_i)$. In this case $|N^{\xi^*}(yz)| \geq (1 - 2\epsilon^{1/4})|X_i|$ (by definition of $\Gamma_{\xi^*}(X_i)$). Since $z \in B_{|1-\xi^*|}(z_{|1-\xi^*|})$, we also know $|\neg N(yz) \cap X_i| \geq |B'' \cap X_i| \geq 2\epsilon^{1/4}|X_i|/p^5$, where the last inequality is because $B'' \in \phi_2(X_i)$.

This shows $(B(\xi^*) \cap Y_j) \times B_{|1-\xi^*|}(z_{|1-\xi^*|}) \subseteq \text{Mix}(2\epsilon^{1/4}/p^5, X_i)$. Define $f_{X_i Y_j Z}(y_0 x_0) = z_{|1-\xi^*|}$. This finishes the proof of (1). The proof of (2) is symmetric, with the roles of Y and X switched everywhere.

For (i), suppose X_i and $X_{i'}$ are lg-interesting, and $\rho'_{X_i} \geq \rho'_{X_{i'}} \geq \rho'_{Y_j}$. Suppose $x \in P_2(X_i)$, $x' \in P_2(X_{i'})$ and $y, y' \in P_2(Y_j)$ are such that $d(x, x') \geq 10\rho'_{X_i}$. Since we chose $f_{X_i Y_j Z}(xy)$ and $f_{X_{i'} Y_j Z}(x'y')$ using Splitting Axiom 1 with $r = \rho'_{X_i}/4p^2$ and $r = \rho'_{X_{i'}}/4p^2$, respectively, we have by part (c) of that axiom that, since $d(y, y') \leq \rho'_{Y_j} \leq \rho'_{X_i}$,

$$d(f_{X_i Y_j Z}(xy), f_{X_{i'} Y_j Z}(x'y')) \geq d(x, x') - d(y, y') - 7(\rho'_{X_i}/4p^4) \geq 5\rho'_{X_i}.$$

The proof of (ii) is symmetric, with the roles of X and Y switched everywhere, and with Splitting Axiom part (b) in place of part (c). □

We are now ready to embark on the main argument of the proof. Given $X \in \{A, B, C\}$, let $I_X(0) = K_X^{sm} \cup K_X^{lg}$. Then for $X \neq Y$, define

$$M^Y(X_i) = \{Y_j \in I_Y(0) : X_i Y_j \notin \mathcal{S} \text{ and } \rho'_{Y_j} \leq \rho'_{X_i}\}.$$

We will construct an integer t^* , and for each $1 \leq u \leq t^*$ and a collection of objects w_u , W_u , \mathcal{S}_{W_u} and $L_X(u)$, $I_X(u)$, $J_X(u)$ for each $X \in \{A, B, C\}$ so that the following hold.

- (1) For each $1 \leq u \leq t^*$, $W_u \in I_A(0) \cup I_B(0) \cup I_C(0)$, and $w_u \in P_2(W_u)$,
- (2) For each $1 \leq u \leq t^*$ and $X \in \{A, B, C\}$, $L_X(u) = \{W_1, \dots, W_u\} \cap I_X(0)$, and

$$\begin{aligned} I_X(u) &= \{X_s \in I_X(0) \setminus L_X(u) : X_s \in M^X(W_u) \text{ for all } W_u \in L_Y(u) \text{ with } Y \neq X\}, \\ J_X(u) &= \{X_s \in I_X(u) : \text{for all } X_i \in L_X(u), d_X(x_s, x_i) \geq 10\rho'_{X_i}\}. \end{aligned}$$

- (3) For each $1 \leq u \leq t^*$ and $X \in \{A, B, C\}$, if $W_u \in I_X(0)$ and YZ lists $\{A, B, C\} \setminus \{X\}$ in alphabetical order, the following hold. If W_u is lg-interesting, then

$$\begin{aligned} \mathcal{S}_{W_u} &= \{Y_j Z_k \in K_2[J_Z(u), J_Y(u)] : Y_j \in K_Y^{sm}, \text{ and for some } y \in P_2(Y_j), \\ &\quad \Psi_3(Z_k) \cap B_{\rho'_{W_u}/4}(f_{W_u Y_j Z}(w_u y)) \neq \emptyset\}, \end{aligned}$$

and if W_u is sm-interesting, then

$$\begin{aligned} \mathcal{S}_{W_u} &= \{Y_j Z_k \in K_2[J_Z(u), J_Y(u)] : Y_j \in K_Y^{lg} \text{ and for some } y \in P_2(Y_j), \\ &\quad \Psi_3(Z_k) \cap B_{\rho'_{W_u}/4}(f_{W_u Y_j Z}(w_u y)) \neq \emptyset\}. \end{aligned}$$

- (4) For each $1 \leq u \leq t^*$, $\mathcal{S}_{W_u} \subseteq \mathcal{S}$ and $|\mathcal{S}_{W_u}| \geq (1 - \sqrt{\epsilon})\epsilon_3^{1/4} \rho'_{W_u} t/2$.
- (5) For all $1 \leq u \neq u' \leq t^*$, $\mathcal{S}_{W_u} \cap \mathcal{S}_{W_{u'}} = \emptyset$.
- (6) For some $X \in \{A, B, C\}$, $|\bigcup_{X_v \in L_X(t_0)} B'_{X_v}| \geq \epsilon^{1/16} n$.

Step 0: There is nothing to do, as this case is vacuous.

Step $i+1$: Suppose $i \geq 0$, and for each $1 \leq u \leq i$, you have chosen W_u , \mathcal{S}_{W_u} , and sets $L_U(u)$, $I_U(u)$, $J_U(u)$ for each $U \in \{A, B, C\}$, such that (1)-(4) hold for each $1 \leq u \leq i$, such that (5) holds for each $1 \leq u \neq u' \leq i$, and such that (6) holds for all $1 \leq u < i$ (note the hypotheses say nothing if $i = 0$).

If $|\bigcup_{U_s \in L_U(i)} B'_{U_s}| \geq \epsilon^{1/16}n$ for some $U \in \{A, B, C\}$, set $t^* = i$ and end the construction (clearly this will not happen if $i = 0$). Otherwise, we have that for each $U \in \{A, B, C\}$, $|\bigcup_{U_s \in L_U(i)} B_{U_s}| < \epsilon^{1/16}n$. By the Size Axiom, this implies that for each $U_v \in L_U(i)$, $\rho'_{U_v} \leq 2p\epsilon^{1/16}n$. Consequently, using (12),

$$\begin{aligned} \left| \bigcup_{U_s \in I_U(i)} U_s \right| &\geq \left| \bigcup_{U_s \in I_U(0)} U_s \right| - \left| \bigcup_{U_s \in L_U(i)} B'_{U_s} \right| \geq |U_{lg}^+ \cup U_{sm}^+|(1 - 4\epsilon_3^{1/3}) - \epsilon^{1/16}n \\ (13) \qquad \qquad \qquad &\geq |U_{lg}^+ \cup U_{sm}^+|(1 - 5\epsilon_3^{1/3}). \end{aligned}$$

By our assumptions, for each $U \in \{A, B, C\}$ and each $U_{i_s}, U_{i_v} \in L_U(i)$,

$$B_{10\rho'_{U_{i_s}}}(u_{i_s}) \cap B_{10\rho'_{U_{i_v}}}(u_{i_v}) = \emptyset.$$

Consequently, by the Size Axiom and the definition of $J_U(i)$,

$$\begin{aligned} \left| \bigcup_{U_s \in J_U(i)} U_s \right| &\geq \left| \bigcup_{U_s \in I_U(0)} U_s \right| - \left| \bigcup_{U_{i_s} \in L_U(i)} B_{10\rho'_{U_{i_s}}}(u_{i_s}) \right| \geq |U_{lg}^+ \cup U_{sm}^+|(1 - 5\epsilon_3^{1/3}) - 10p\epsilon^{1/16}n \\ (14) \qquad \qquad \qquad &\geq |U_{lg}^+ \cup U_{sm}^+|(1 - 6\epsilon_3^{1/3}). \end{aligned}$$

For each $U \in \{A, B, C\}$, set $r_{i+1}^U = \max\{\rho'_{U_j} : U_j \in J_U(i)\}$. Then define

$$r_{i+1} = \max\{r_{i+1}^A, r_{i+1}^B, r_{i+1}^C\}.$$

Choose $X \in \{A, B, C\}$ such that $r_{i+1}^X = r_{i+1}$, and any $j_{i+1} \in J_X(i)$ such that $\rho'_{X_{j_{i+1}}} = r_{i+1}$. Then define $W_{i+1} := X_{j_{i+1}}$, $L_X(i+1) = L_X(i) \cup \{W_{i+1}\}$, and for each $U \neq X$, set $L_U(i+1) = L_U(i)$. For each $U \in \{A, B, C\} \setminus \{X\}$, define $I_U(i+1) = M^U(W_{i+1}) \cap I_U(i)$ and $J_U(i+1) = J_U(i) \cap M^U(W_{i+1})$. Then set

$$I_X(i+1) := I_X(i) \setminus \{W_{i+1}\} \text{ and } I_X(i+1) := \{X_s \in I_X(i+1) : d(u_s, x_{j_{i+1}}) \geq 10\rho'_{W_{i+1}}\}.$$

Let YZ list the elements of $\{A, B, C\} \setminus X$ in alphabetical order, and define

$$\begin{aligned} K_Y^{sm}(i+1) &= \{Y_j \in J_Y(i+1) : Y_j \text{ is sm-interesting}\}, \text{ and} \\ K_Y^{lg}(i+1) &= \{Y_j \in J_Y(i+1) : Y_j \text{ is lg-interesting}\}. \end{aligned}$$

Using (11) and (14), and since $J_Y(i+1) = J_Y(i)$,

$$\left| \bigcup_{Y_j \in K_Y^{sm}(i+1)} Y_j \right| \geq \left| \bigcup_{Y_j \in K_Y^{sm}(i+1) \setminus J_Y(i+1)} Y_j \right| \geq (1 - 6\epsilon_3^{1/3})|Y_{sm}|.$$

A similar argument shows that

$$\left| \bigcup_{Y_j \in K_Y^{lg}(i+1)} Y_j \right| \geq (1 - 6\epsilon_3^{1/3})|Y_{lg}|.$$

Consequently, combining this with the Size and Special Sets Axioms, and the fact that $\epsilon \ll \mu, \tau$, we have

$$|K_Y^{sm}(i+1)| \geq (\mu n/2p)(1 - 6\epsilon_3^{1/3})/(3n/t) \geq \mu(1 - 6\epsilon_3^{1/3})t/6p \geq \mu t/7p,$$

and similarly $|K_Y^{lg}(i+1)| \geq \tau t/7p$. By construction, W_{i+1} is either lg-interesting or sm-interesting. If W_{i+1} is lg-interesting, let $Y_* = Y_{sm}$ and $K^* = K_Y^{sm}(i+1)$. If W_{i+1} is sm-interesting, let $Y_* = Y_{lg}$ and $K^* = K_Y^{lg}(i+1)$.

Fix $w_{i+1} \in P_2(W_{i+1})$. Given $Y_j \in K^*$ and $y \in P_2(Y_j)$, consider $B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y))$ as in Claim 7.33. Note that if $Z_k \in J_Z(i+1)$, $B'' \in \phi_3(Z_k)$, and $B'' \cap B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \neq \emptyset$, then $B'' \subseteq B_{r_{i+1}/2}(f_{W_{i+1}Y_jZ}(w_{i+1}y))$ (since by construction, $r_{i+1} = \rho'_{W_{i+1}} \geq \rho'_{Z_k}$). Thus, for all such Z_k , we have by Claim 7.33 that

$$|E \cap K_3[Z_k, W_{i+1}, Y_j]| / |K_3[Z_k, W_{i+1}, Y_j]| \in (\epsilon^{1/4}\mu^4/2, 1 - \epsilon^{1/4}\mu^4/2).$$

Since we assumed $Y_j \in J_Y(i+1) \subseteq M^Y(W_{i+1})$, $W_{i+1}Y_j \notin \mathcal{S}$. Therefore, if $W_{i+1}Z_k \notin \mathcal{S}$, and Z_k is as above, then we would have to have $Y_jZ_k \in Irr(W_{i+1})$. Thus, if $Z_k \in J_Z(i+1)$ and $\Psi_3(Z_k) \cap B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \neq \emptyset$, then $Y_jZ_k \in Irr(W_{i+1})$. Let $\mathcal{S}_{W_{i+1}Y_j}$ be the set of pairs arising in this way for Y_j , i.e.

$$\mathcal{S}_{W_{i+1}Y_j} = \{Y_jZ_k : Z_k \in J_Z(i+1), \text{ and for some } y \in P_2(Y_j), \\ \Psi_3(Z_k) \cap B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \neq \emptyset\}$$

We have shown that for all $Y_j \in K^*$, $\mathcal{S}_{W_{i+1}Y_j} \subseteq Irr(W_{i+1})$. Clearly $|\mathcal{S}_{W_{i+1}Y_j}| \geq s_j$, where we define s_j as follows.

$$s_j := \frac{|(\bigcup_{y \in P_2(Y_j)} B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \cap (\bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k)))|}{|N/t|}.$$

Note that for any $Y_j \in K^*$,

$$\begin{aligned} & \left| \left(\bigcup_{y \in P_2(Y_j)} B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \setminus \left(\bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right) \right) \right| \\ & \geq \left| \bigcup_{y \in P_2(Y_j)} B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \right| - s_j(N/t). \end{aligned}$$

By the Size Lemma, for each $y \in P_2(Y_j)$,

$$|B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y))| \geq r_{i+1}n/8p = \rho'_{W_{i+1}}n/8p.$$

Thus

$$\left| \left(\bigcup_{y \in P_2(Y_j)} B_{r_{i+1}/4}(f_{W_{i+1}Y_jZ}(w_{i+1}y)) \setminus \left(\bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right) \right) \right| \geq \rho'_{W_{i+1}}n/8p - s_j(N/t).$$

Now define

$$\mathcal{G}^* = \{Y_j \in K^* : s_j \geq \rho'_{W_{i+1}}(1 - \sqrt{\epsilon})t/2p^3\}.$$

We claim $|\bigcup_{Y_j \in \mathcal{G}^*} Y_j| \geq \epsilon_3^{1/4}n$. Suppose towards a contradiction this is not the case, i.e. that $|\bigcup_{Y_j \in \mathcal{G}^*} Y_j| < \epsilon_3^{1/4}n$. Note that if $Y_j \in K^* \setminus \mathcal{G}^*$, then by definition of \mathcal{G}^* and the above,

$$\begin{aligned} \left| \left(\bigcup_{y \in P_2(Y_j)} B_{r_{i+1}/4}(f_{W_{i+1}Y_j Z}(w_{i+1}y)) \right) \setminus \left(\bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right) \right| &\geq \rho'_{W_{i+1}}n/8p - \rho'_{W_{i+1}}(1 - \sqrt{\epsilon})N/4p^3 \\ &= \rho'_{W_{i+1}}n \left(\frac{1}{8p} - \frac{3(1 - \sqrt{\epsilon})}{4p^3} \right) \\ &\geq \rho'_{W_{i+1}}n/16p^3. \end{aligned}$$

By assumption $|\bigcup_{Y_j \in \mathcal{G}^*} Y_j| \leq \epsilon_3^{1/4}n$, so

$$(15) \quad \left| \bigcup_{Y_j \in K^* \setminus \mathcal{G}^*} Y_j \right| \geq \left| \bigcup_{Y_j \in K^*} Y_j \right| - \epsilon_3^{1/4}n \geq |Y_*|(1 - 7\epsilon_3^{1/4}),$$

where the last inequality also uses the Size Axiom, the Special Sets Axiom, and the fact that $\epsilon \ll \mu$ (if $Y_* = Y_{sm}$) and $\epsilon \ll \tau$ (if $Y_* = Y_{lg}$). Let K be a maximal $10\rho'_{W_{i+1}}$ -separated subset of $\bigcup_{K^* \setminus \mathcal{G}^*} P_2(Y_j)$. Since

$$\bigcup_{Y_j \in K^* \setminus \mathcal{G}^*} Y_j \subseteq \bigcup_{y \in K} B_{10\rho'_{X_i}}(y),$$

we know that by (15), the Size Axiom, and the Special Sets Axiom,

$$\min\{\mu, \tau\}(1 - 7\epsilon_3^{1/4})n/2p \leq |Y_*|(1 - 7\epsilon_3^{1/4}) \leq |K|10p\rho'_{W_{i+1}}n.$$

Thus $|K| \geq (\rho'_{W_{i+1}})^{-1}(1 - 7\epsilon_3^{1/4})\min\{\mu, \tau\}/20p^2$. By definition, for all $y \neq y' \in K$, $d(y, y') > 10\rho'_{W_{i+1}}$, and therefore, by Claim 7.33(i) or (ii), $d(f_{W_{i+1}Y_j Z}(w_{i+1}y), f_{W_{i+1}Y_{j'} Z}(w_{i+1}y')) \geq 5\rho'_{W_{i+1}}$. Consequently,

$$B_{r_{i+1}/4}(f_{W_{i+1}Y_j Z}(w_{i+1}y)) \cap B_{r_{i+1}/4}(f_{W_{i+1}Y_{j'} Z}(w_{i+1}y')) = \emptyset.$$

Thus

$$\begin{aligned} \left| \bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right| &\leq n - \left| \left(\bigcup_{Y_j \in K^* \setminus \mathcal{G}^*} \bigcup_{y \in K} B_{r_{i+1}/4}(f_{W_{i+1}Y_j Z}(w_{i+1}y)) \right) \setminus \bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right| \\ &\leq n - |K|\rho'_{W_{i+1}}n/16p^3 \\ &\leq n - (\rho'_{W_{i+1}})^{-1}(1 - 7\epsilon_3^{1/4})\min\{\mu, \tau\}(20p^2)^{-1}\rho'_{W_{i+1}}n(16p^3)^{-1} \\ &\leq n(1 - \min\{\mu, \tau\}/240p^6). \end{aligned}$$

However, by the definition of $\Psi_3(Z_k)$ in Claim 7.32, we know that for all $Z_k \in J_Z(i+1)$, $|\Psi(Z_k)| \geq (1 - 3\epsilon^{1/4})|Z_k|$. Consequently by (14),

$$\begin{aligned} \left| \bigcup_{Z_k \in J_Z(i+1)} \Psi_3(Z_k) \right| &\geq (1 - 6\epsilon^{1/3}) \left| \bigcup_{Z_k \in J_Z(i+1)} Z_k \right| \geq (1 - 6\epsilon^{1/3})(1 - \epsilon_3^{1/2})|Z_{lg}^+ \cup Z_{sm}^+| \\ &> (1 - \min\{\mu, \tau\}/240p^6)n, \end{aligned}$$

where the last inequality uses that $\epsilon \ll \mu, \tau$. This is a contradiction, so we must have that $|\bigcup_{Y_j \in \mathcal{G}^*} Y_j| \geq \epsilon_3^{1/4}n$, so $|\mathcal{G}^*| \geq \epsilon_3^{1/4}t/3$. Now define

$$\mathcal{S}_{W_{i+1}} = \bigcup_{Y_j \in \mathcal{G}^*} \mathcal{S}_{W_{i+1}Y_j}$$

Then we have shown that $\mathcal{S}_{W_{i+1}} \subseteq \mathcal{S}$ and

$$|\mathcal{S}_{W_{i+1}}| \geq |\mathcal{G}^*| \rho'_{W_{i+1}} (1 - \sqrt{\epsilon})t/2p^3 \geq \rho'_{W_{i+1}} \epsilon_3^{1/4} (1 - \sqrt{\epsilon})^2 t^2 / 12p^3.$$

This finishes the definition of $\mathcal{S}_{W_{i+1}}$.

We now show that for all $u \leq i$, $\mathcal{S}_{W_u} \cap \mathcal{S}_{W_{i+1}} = \emptyset$. Suppose towards a contradiction, there is some $u \leq i$ with $\mathcal{S}_{W_u} \cap \mathcal{S}_{W_{i+1}} \neq \emptyset$, say $Y_j Z_k \in \mathcal{S}_{W_u} \cap \mathcal{S}_{W_{i+1}}$. By construction, this implies $W_u \in L_X(u)$, say $W_u = X_{j_u}$ for some $j_u \neq j_{i+1}$. We claim that either both $W_u, W_{i+1} \in K_X^{lg}$ or both $W_u, W_{i+1} \in K_X^{sm}$. Suppose $W_u \in K_X^{sm}$ and $W_{i+1} \in K_X^{lg}$. Then $W_u \in K_X^{sm}$ implies $Y_j \in K_Y^{lg}$, while $W_{i+1} \in K_X^{lg}$ implies $Y_j \in K_Y^{sm}$. But now $Y_j \in K_Y^{sm} \cap K_Y^{lg}$. This implies that $|Y_j \cap Y_{sm}^+| > |Y_j|/2$ and $|Y_j \cap Y_{lg}^+| > |Y_j|/2$, which implies $Y_{sm}^+ \cap Y_{lg}^+ \neq \emptyset$, contradicting the Disjointness Axiom. A similar argument shows we cannot have $W_{i+1} \in K_X^{sm}$ and $W_u \in K_X^{lg}$.

Therefore we have that either $W_u, W_{i+1} \in K_X^{lg}$ or $W_u, W_{i+1} \in K_X^{sm}$. By construction, there is $w_u \in P_2(W_u)$ such that one of the following holds.

- (1) Both $W_u, W_{i+1} \in K_X^{lg}$ and $Y_j Z_k \in K_2[K_Y^{sm}(u), J_Z(u)] \cap K_2[K_Y^{sm}(i+1), J_Z(i+1)]$, and

$$\Psi_3(Z_k) \cap B_{r_u/4}(f_{W_u Y_j Z}(w_u y)) \neq \emptyset \text{ some } y \in P_2(Y_j), \text{ and}$$

$$\Psi_3(Z_k) \cap B_{r_{i+1}/4}(f_{W_{i+1} Y_j Z}(w_{i+1} y)) \neq \emptyset \text{ some } y \in P_2(Y_j).$$

- (2) Both $W_u, W_{i+1} \in K_X^{sm}$ and $Y_j Z_k \in K_2[K_Y^{lg}(u), J_Z(u)] \cap K_2[K_Y^{lg}(i+1), J_Z(i+1)]$, and

$$\Psi_3(Z_k) \cap B_{r_u/4}(f_{W_u Y_j Z}(w_u y)) \neq \emptyset \text{ some } y \in P_2(Y_j) \text{ and}$$

$$\Psi_3(Z_k) \cap B_{r_{i+1}/4}(f_{W_{i+1} Y_j Z}(w_{i+1} y)) \neq \emptyset \text{ some } y \in P_2(Y_j).$$

Since $Y_j \in J_Y(i+1)$, we know $\rho'_{Y_j} \leq \min\{\rho'_{W_u}, \rho'_{W_{i+1}}\}$. By construction, the centers of B_{W_u} and $B_{W_{i+1}}$ have distance at least $10\rho'_{W_u}$ from each other. Combining this with the fact that for all $y, y' \in P_2(Y_j)$, $d(y, y') \leq \rho'_{Y_j}$, we have by Claim 7.33 that the following hold.

$$(16) \quad d(f_{Y_j W_u Z}(y w_u), f_{W_{i+1} Y_j Z}(w_{i+1} y')) \geq 5\rho'_{W_u} \geq 5\rho'_{Z_k},$$

But (16) implies that it is not possible to have both $B_{r_u/4}(f_{W_u Y_j Z}(w_u y))$ and $B_{r_{i+1}/4}(f_{W_{i+1} Y_j Z}(w_{i+1} y'))$ intersect B_{Z_k} , a contradiction. This finishes our verification that $\mathcal{S}_{W_u} \cap \mathcal{S}_{W_{i+1}} = \emptyset$, and consequently the $(i+1)$ -st step of our construction.

Clearly this will end at some stage t^* , where we will have that for some $X \in \{A, B, C\}$, $|\bigcup_{X_v \in L_X(t^*)} B'_{X_v}| \geq \epsilon^{1/16} N$. Say $1 \leq i_1 < \dots < i_s \leq t^*$ are such that

$$W_{i_1}, \dots, W_{i_s} \subseteq \{X_i : i \in [t]\},$$

and $|\sum_{j=1}^s B'_{W_{i_j}}| \geq \epsilon^{1/16} N$. By the size axiom, this implies $\sum_{j=1}^s \rho'_{W_{i_j}} \geq \epsilon^{1/16}/4p$. Therefore, we have that

$$|\mathcal{S}| \geq \sum_{j=1}^s |\mathcal{S}_{W_{i_j}}| \geq (1 - 2\sqrt{\epsilon})^2 \epsilon_3^{1/4} t^2 (\sum_{j=1}^s \rho'_{W_{i_j}}) / 4p^3 > \epsilon_3^{3/4} t^2 / 20p^4 > 13\epsilon_3 t^2,$$

where the last inequality uses that $\epsilon \ll p^{-1}$ and $t \geq t_0$. But contradicts (10), which finishes the proof. \square

We obtain as an immediate corollary that \mathcal{H}_{HP} and $\mathcal{H}_{\text{GS}_p}$ require non-binary vdisc_3 -error.

Corollary 7.34. \mathcal{H}_{HP} , as well as $\mathcal{H}_{\text{GS}_p}$ for any prime $p \geq 3$ require non-binary vdisc_3 -error.

Proof. This follows immediately from Lemmas 7.26, 7.27, and Theorem 7.28. \square

We will now use this fact to prove Theorems 2.53 and 2.56, which state that if \mathcal{H} is a hereditary 3-graph property and either $\mathcal{H}_{\text{HP}} \subseteq \text{Trip}(\mathcal{H})$ or $\mathcal{H}_{\text{GS}_p} \subseteq \text{Trip}(\mathcal{H})$, then \mathcal{H} requires non-binary $\text{disc}_{2,3}$ -error.

PROOF OF THEOREMS 2.53 AND 2.56: Suppose towards a contradiction there exists a hereditary 3-graph property \mathcal{H}_2 such that one of (1) or (2) below holds, and such that \mathcal{H}_2 admits binary $\text{disc}_{2,3}$ -error.

- (1) $\mathcal{H}_1 = \mathcal{H}_{\text{HP}} \subseteq \text{Trip}(\mathcal{H}_2)$, or
- (2) $\mathcal{H}_1 = \mathcal{H}_{\text{GS}_p} \subseteq \text{Trip}(\mathcal{H}_2)$.

By Corollary 7.34, there is some $\epsilon_1 > 0$ such that for all T, M , there is $N_{T,M}$ such that for all $n \geq N$, neither $\text{HP}(n)$ nor $\text{GS}_p(n)$ admit an ϵ_1 -homogeneous partition $G_{T,M}$ into at most T parts with binary error.

Let $0 < \epsilon_1 \ll \epsilon_1$, and choose $\epsilon_2 : \mathbb{N} \rightarrow (0, 1)$ as in Proposition 5.1 for ϵ'_1 and $k = 8$. Define $\epsilon'_2 : \mathbb{N} \rightarrow (0, 1]$ by choosing $\epsilon'_2(x) \ll \epsilon'_1 \epsilon_2(x)$ for all x . By assumption, there are T_1, L_1, M_1 such that for all $G' = (V', E') \in \mathcal{H}_2$ with $|V'| \geq M_1$, there is are $1 \leq t \leq T_1$, $1 \leq \ell \leq L_1$ and a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V' which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular with respect to G' with binary $\text{disc}_{2,3}$ -error.

Let $T_2 = T_1(\epsilon'_1)^{-1}$ and choose $M_2 \gg 3M_1 T_1(\epsilon'_1)^{-1}$. Now fix $n \geq M_2, N_{T_2, M_2}$ and let $G = \text{HP}(n) = (A \cup B \cup C, E)$ if (1) holds, and $G = \text{GS}_p(n) = (A \cup B \cup C, E)$ if (2) holds.

By assumption, $G \in \text{Trip}(\mathcal{H}_2)$. By definition this means there is some $H = (V, F) \in \mathcal{H}_2$ and sets

$$A' = \{v_a : a \in A\} \subseteq V', \quad B' = \{v_b : b \in B\} \subseteq V', \quad \text{and} \quad C' = \{v_c : c \in C\} \subseteq V',$$

such that $v_{ag}v_{bg'}v_{cg''} \in F$ if and only if $abc \in E$. Note this implies $|V'| \geq n \geq M_1$. Let $H' = H[A' \cup B' \cup C']$. Then $H' \in \mathcal{H}_2$, and $n \leq |V(H')| \leq 3n$. Since $n \geq M_1$, there are $t_0 \leq t \leq T_1$, $\ell_0 \leq \ell \leq L_1$ and a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition \mathcal{P} of $V(H')$ which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular with respect to H' , and which has binary $\text{disc}_{2,3}$ -error. Say $\mathcal{P}_{\text{vert}} = V'_1 \cup \dots \cup V'_t$ and $\mathcal{P}_{\text{edge}} = \{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Note that for each $1 \leq i \leq t$, $n/t \leq |V_i| \leq 3n/t$. Define

$$\Sigma = \{ij \in \binom{[t]}{2} : \text{some } P_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P}) \text{ fails } \text{disc}_{2,3}(\epsilon_2(\ell), 1/\ell) \text{ with respect to } H'\}.$$

By assumption, $|\Sigma| \leq \epsilon'_1 t^2$. For each $1 \leq i \leq t$, let

$$A_i = \{a \in A : v_a \in V'_i\}, \quad B_i = \{b \in B : v_b \in V'_i\} \text{ and } C_i = \{c \in C : v_c \in V'_i\}.$$

Then $\{A_i, B_i, C_i : i \in [t]\}$ forms a partition of $A \cup B \cup C$. Set $t_0 = 3\lceil \epsilon_1^{-2} \rceil$ and $K = \lfloor 3n/t_0 t \rfloor$. For each $i \in [t]$ and $X \in \{A, B, C\}$, let $X_i = X_{i0} \cup X_{i1} \cup \dots \cup X_{ik_i^X}$ be a partition with the property that $|X_{i0}| < K$ and so that for each $1 \leq s \leq k_i^X$, $|X_{is}| = K$. Let $k_X = \sum_{i=1}^t k_i^X$ and set $X_0 = \bigcup_{i=1}^t X_{i0}$. Note that by definition of t_0 and K , $|X_0| \leq tK \leq 3\epsilon_1^2 n$. Choose $\{X_0^{is} : 1 \leq i \leq t, 1 \leq s \leq k_i^X\}$ any equipartition of X_0 into k parts, and for each $1 \leq i \leq t$ and $1 \leq s \leq k_i^X$, set $X'_{is} = X_{is} \cup X_0^{is}$. Clearly $|X'_{is} \setminus X_{is}| \leq 3\epsilon_2^2 |X_{is}|$.

We now have an partition $\mathcal{Q}_{\text{vert}}$ of $V(G)$ given by

$$\{A'_{is} : 1 \leq i \leq t, 1 \leq s \leq k_i^A\} \cup \{B'_{is} : 1 \leq i \leq t, 1 \leq s \leq k_i^B\} \cup \{C'_{is} : 1 \leq i \leq t, 1 \leq s \leq k_i^C\}.$$

For each $1 \leq \alpha \leq \ell$, $X \neq Y$ in $\{A, B, C\}$, $ij \in \binom{[t]}{2}$, $1 \leq s \leq k_i^X$, and $1 \leq r \leq k_j^Y$, set $Q_{ij, sr, XY}^\alpha = P_{ij}^\alpha \cap K_2[X'_{is}, Y'_{jr}]$. Note that by Lemma 3.18, if P_{ij}^α satisfies $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, then each $Q_{ij, sr, XY}^\alpha$ satisfies $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$.

Then for each $X \in \{A, B, C\}$, $1 \leq i \leq t$, and $sr \in \binom{[k_i^X]}{2}$, choose any partition $K_2[W'_{is}, W'_{ir}] = \bigcup_{\alpha \leq \ell} Q_{is, ir', XX}^\alpha$ so that each $Q_{is, ir', XX}^\alpha$ has $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$ (such a partition exists by Lemma 3.24). We now set

$$\begin{aligned} \mathcal{Q}_{\text{edge}} = & \{Q_{ij, sr, AB}^\alpha : 1 \leq i, j \leq t, 1 \leq s \leq k_i^A, 1 \leq r \leq k_j^B\} \cup \\ & \{Q_{ij, sr, AC}^\alpha : 1 \leq i, j \leq t, 1 \leq s \leq k_i^A, 1 \leq r \leq k_j^C\} \cup \\ & \{Q_{ij, sr, BC}^\alpha : 1 \leq i, j \leq t, 1 \leq s \leq k_i^B, 1 \leq r \leq k_j^C\} \cup \\ & \{Q_{ij, sr, X}^\alpha : X \in \{A, B, C\}, 1 \leq s, r \leq k_i^X\}. \end{aligned}$$

It is straightforward to check that \mathcal{Q} is a $(tk, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of $A \cup B \cup C$. We then define

$$\begin{aligned} \Sigma' = & \{X_{is}Y_{jr} : X \neq Y \in \{A, B, C\}, ij \in \Sigma, s \in [k_i^X], r \in [k_j^Y]\} \cup \\ & \{X_{is}Y_{ir} : X, Y \in \{A, B, C\}, s \in [k_i^X], r \in [k_i^Y]\}. \end{aligned}$$

It is straightforward to check that since $\epsilon'_1 \ll \epsilon_1$ and by our choice of t_0 , we have $|\Sigma'| \leq \epsilon_1 |\mathcal{Q}_{\text{vert}}|^2$. We claim that any triple of sets from $\mathcal{Q}_{\text{vert}}$ avoiding Σ' is ϵ_1 -homogeneous with respect to G . Clearly this holds of any triple of the form $X_{is}X_{js}Y_{pq}$ for $X, Y \in \{A, B, C\}$

or $X_{is}X_{js}X_{pq}$ for some $X \in \{A, B, C\}$. Indeed, any such triple will have edge density 0 in G , since G is 3-partite with parts A, B, C .

So fix some triple $A_{is}B_{jr}C_{pq} \in (\mathcal{Q}_{vert}^3)$ such that $A_{is}B_{jr}, B_{jr}C_{pq}, A_{is}C_{pq} \notin \Sigma'$, and a triad

$$\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma} = (A_{is} \cup B_{jr} \cup C_{pq}, Q_{ij, sr, AB}^\alpha \cup Q_{jp, rq, BC}^\beta \cup Q_{ip, sq, AC}^\gamma).$$

By construction, $ij, ip, jp \notin \Sigma$, so the triad $P_{ijp}^{\alpha,\beta,\gamma} = (V_i' \cup V_j' \cup V_p', P_{ij}^\alpha \cup P_{jp}^\beta \cup P_{ip}^\gamma)$ satisfies $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$ with respect to H' . Let $d \in [0, 1]$ be such that $|E(H') \cap K_3^{(2)}(P_{ijp}^{\alpha,\beta,\gamma})| = d|K_3^{(2)}(P_{ijp}^{\alpha,\beta,\gamma})|$. Now suppose Γ' is a subgraph of $\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma}$.

Consider the graph $R_{ijp}^{\alpha,\beta,\gamma}$ with vertex set $V_i' \cup V_j' \cup V_p'$ and edge set

$$\{xyz \in K_3[V_i', V_j', V_p'] : x = v_a, y = v_b, \text{ and } z = v_c \text{ for some } abc \in E(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})\},$$

and consider the subgraph R' with $V_i' \cup V_j' \cup V_p'$ and edge set

$$\{xyz \in K_3[V_i', V_j', V_p'] : x = v_a, y = v_b, \text{ and } z = v_c \text{ for some } abc \in K_3^{(2)}(\Gamma')\},$$

Both $R_{ijp}^{\alpha,\beta,\gamma}$ and R' are subgraphs of $P_{ijp}^{\alpha,\beta,\gamma}$, so by the definition of $\text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))$,

$$||E(H') \cap K_3^{(2)}(R_{is,jr,pq}^{\alpha,\beta,\gamma})| - d|K_3^{(2)}(R_{is,jr,pq}^{\alpha,\beta,\gamma})|| \leq \epsilon'_1 \frac{1}{\ell^3} |V_i'| |V_j'| |V_p'| \text{ and}$$

$$||E(H') \cap K_3^{(2)}(R')| - d|K_3^{(2)}(R')|| \leq \epsilon'_1 \frac{1}{\ell^3} |V_i'| |V_j'| |V_p'|$$

Since i, j, p are distinct, we have that

$$|K_3^{(2)}(R_{is,jr,pq}^{\alpha,\beta,\gamma})| = |K_3^{(2)}(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})| \text{ and } |K_3^{(2)}(R')| = |K_3^{(2)}(\Gamma')|,$$

and

$$|E(H') \cap K_3^{(2)}(R_{is,jr,pq}^{\alpha,\beta,\gamma})| = |E(G) \cap K_3^{(2)}(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})| \text{ and } |E(H') \cap K_3^{(2)}(R')| = |E(G) \cap K_3^{(2)}(\Gamma')|.$$

Consequently, if we let d' be such that $|E(G) \cap K_3^{(2)}(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})| = d'|K_3^{(2)}(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})|$, then

$$\begin{aligned} |d' - d| &\leq \epsilon'_1 \frac{1}{\ell^3} |V_i'| |V_j'| |V_p'| (|K_3^{(2)}(\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma})|)^{-1} \leq \epsilon'_1 \frac{1}{\ell^3} |V_i'| |V_j'| |V_p'| 3\epsilon_1^{-2} \\ &\leq \epsilon_1^8 \frac{1}{\ell^3} |A_{is}| |B_{jr}| |C_{pq}|, \end{aligned}$$

where the second inequality is from Corollary 3.6 and the definition of \mathcal{Q}_{vert} , and the last inequality is since $\epsilon'_1 \ll \epsilon_1$. We also have from the above that

$$||E(G) \cap K_3^{(2)}(\Gamma')| - d'|K_3^{(2)}(\Gamma')|| \leq \epsilon'_1 \frac{1}{\ell^3} |V_i'| |V_j'| |V_p'| \leq \epsilon_1^8 \frac{1}{\ell^3} |A_{is}| |B_{jr}| |C_{pq}|,$$

where the last inequality is since $\epsilon'_1 \ll \epsilon_1$ and the definition of \mathcal{Q}_{vert} . Combining these, we have that

$$||E(G) \cap K_3^{(2)}(\Gamma')| - d'|K_3^{(2)}(\Gamma')|| \leq 2\epsilon_1^8 \frac{1}{\ell^3} |A_{is}| |B_{jr}| |C_{pq}|,$$

which shows that $\Gamma_{is,jr,pq}^{\alpha,\beta,\gamma}$ satisfies $\text{disc}_{2,3}(2\epsilon_1^8, \epsilon_2(\ell))$ with respect to G . We can conclude By Proposition 5.1, (since G has VC-dimension at most 8) that

$$|E(G) \cap K_3[A_{is}, B_{jr}, C_{pq}]|/|C_{is}||B_{jr}||C_{pq}| \in [0, \epsilon_1^2) \cup (1 - \epsilon_1^2, 1].$$

This shows that any triple form \mathcal{Q}_{vert} avoiding Σ' is ϵ_1^2 -homogeneous with respect to G . As defined \mathcal{Q}_{vert} may not be itself an equipartition, although its parts will differ in size by at most $\epsilon_1^2 n/|\mathcal{Q}_{vert}|$. It is now not too difficult to show that one can redistribute some vertices to make \mathcal{Q}_{vert} into an equipartition without changing the densities within any of the triads by very much. The result will be an ϵ_1 -homogeneous equipartition of G with binary error and at most T_2 parts, a contradiction. \square

We now prove as a Corollary 2.59, which says that a property admitting binary, zero, or linear $\text{disc}_{2,3}$ -error is $\text{disc}_{2,3}$ -homogeneous.

PROOF OF COROLLARY 2.59: Suppose \mathcal{H} admits zero $\text{disc}_{2,3}$ -error. By Corollary 5.6, there is k such that $\overline{H}(k) \notin \text{Trip}(\mathcal{H})$. By Fact 2.22, \mathcal{H} has finite VC₂-dimension, so by Theorem 2.24, it is $\text{disc}_{2,3}$ -homogeneous. Suppose now \mathcal{H} admits binary $\text{disc}_{2,3}$ -error. By Theorem 2.53, there is $k \geq 1$ such that $\text{HP}(k) \notin \text{Trip}(\mathcal{H})$. By Fact 2.22, \mathcal{H} has finite VC₂-dimension, so by Theorem 2.24, it is $\text{disc}_{2,3}$ -homogeneous. Suppose now \mathcal{H} admits linear $\text{disc}_{2,3}$ -error. By Theorem 2.50, there is $k \geq 1$ such that $F(k) \notin \text{Trip}(\mathcal{H})$. By Fact 2.22, \mathcal{H} has finite VC₂-dimension, so by Theorem 2.24, it is $\text{disc}_{2,3}$ -homogeneous. \square

A similar argument will be important in our proof of Theorem 2.12. In particular, we will use the following lemma.

Lemma 7.35. *There are $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$ such that for all T, L and N , there is $n \geq N$ such that for all $t_0 \leq t \leq T_0$, and $\ell_0 \leq \ell \leq T_0$, any $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of $V(\text{HP}(n))$ which is $(\epsilon_1, \epsilon_2(\ell))$ -regular, has a disc_3 -irregular triad.*

Proof. Since \mathcal{H}_{HP} has finite VC-dimension it is $\text{disc}_{2,3}$ -homogeneous by Theorem 2.24. Suppose towards a contradiction that for all $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$ there are T, L and N , such that for all $n \geq N$ there is $t_0 \leq t \leq T_0$, and $\ell_0 \leq \ell \leq L_0$, any $(t, \ell, \epsilon_2, \epsilon_1)$ -decomposition of $V(\text{HP}(n))$ which is (ϵ_1, ϵ_2) -regular and has no disc_3 -irregular triad. By Theorem 2.53, there are $\mu_1, \mu_2, t_0, \ell_0$ such that for all T, L and N , there is $n \geq N$ such that for all $t_0 \leq t \leq T_0$, and $\ell \leq T_0$, any $(t, \ell, \mu_1, \mu_2(\ell))$ -decomposition of $V(\text{HP}(n))$ which is $(\mu_1, \mu_2(\ell))$ -regular, has non-binary $\text{disc}_{2,3}$ -error.

By our assumption, and because \mathcal{H}_{HP} is $\text{disc}_{2,3}$ -homogeneous, there are T, L and N , so that for all $n \geq N$ there is $t_0 \leq t \leq T_0$, and $\ell_0 \leq \ell \leq L_0$, and a $(t, \ell, \mu_1, \mu_2(\ell))$ -decomposition of $V(\text{HP}(n))$ which is $(\mu_1, \mu_2(\ell))$ -regular, with no disc_3 -irregular triads, and which is $\text{disc}_{2,3}$ -homogeneous. Let Σ be the set of pairs $V_i V_j$ where at least $(\epsilon'_1)^{1/2} |V_i| |V_j|$ of the $xy \in K_2[V_i V_j]$ are in a disc_2 -irregular $P \in \mathcal{P}_{edge}$. By assumption, $|\Sigma| \leq (\epsilon'_1)^{1/2} t^2$. If $V_i V_j, V_i V_k, V_j V_k \notin \Sigma$, then by Proposition 5.1, $|E \cap K_3[V_i, V_j, V_k]|/K_3[V_i, V_j, V_k] \in [0, \epsilon_1) \cup (1 - \epsilon_1, 1]$. But now \mathcal{P} is vdisc_3 -homogeneous with binary error, a contradiction. \square

We now prove Theorem 2.12.

PROOF OF THEOREM 2.12: Fix a hereditary 3-graph property \mathcal{H} , and assume it admits binary $\text{disc}_{2,3}$ -error. Fix $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ nonincreasing, and $t_0, \ell_0 \geq 1$. Let $\epsilon'_1 = \epsilon_1^2$. By assumption, there is T, L and N so that if $H \in \mathcal{H}$ has at least N vertices, then there exist $\ell_0 \leq \ell \leq L$, $t_0 \leq t \leq T$, and \mathcal{P} a $(t, \ell, \epsilon'_1, \epsilon_2(\ell))$ -decomposition of $V(H)$ which is $(\epsilon'_1, \epsilon_2(\ell))$ -regular, with binary $\text{disc}_{2,3}$ -error with respect to H . Say \mathcal{P} consists of $\{V_i : i \in [t]\}$ and $\{P_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$. Let $\Sigma \subseteq \binom{[t]}{2}$ be such that every $\text{disc}_{2,3}$ -irregular triad of \mathcal{P} uses a pair $V_i V_j$ with $ij \in \Sigma$, and where $|\Sigma| \leq \epsilon'_1 t^2$.

Let $\ell_1 = \lfloor \ell/2 \rfloor$ and $\ell_2 = \lceil \ell/2 \rceil$. For each $i \in [t]$, let $V_i = V_{i1} \cup V_{i2}$ be an equipartition. For each $i < j$ with $ij \in \Sigma$, use Lemma 3.24 to choose partitions $V_{i1} \times V_j = \bigcup_{\alpha \leq \ell_1} Q_{ij}^\alpha$ and $V_{i2} \times V_j = \bigcup_{\ell_1+1 \leq \alpha \leq \ell} Q_{ij}^\alpha$ so that for each $1 \leq \alpha \leq \ell_1$, Q_{ij}^α has $\text{disc}_2(\epsilon_1(\ell_1), 1/\ell_1)$ in $V_{i1} \times V_j$, and for each $\ell_1 + 1 \leq \alpha \leq \ell$, Q_{ij}^α has $\text{disc}_2(\epsilon_1(\ell_2), 1/\ell_2)$ in $V_{i1} \times V_j$. Clearly each Q_{ij}^α fails $\text{disc}_2(\epsilon_2(\ell), 1/\ell)$. Now define \mathcal{Q} to have vertex partition $\{V_i : i \in [t]\}$ and edge partition

$$\{P_{ij}^\alpha : ij \in \binom{[t]}{2} \setminus \Sigma, \alpha \leq \ell\} \cup \{Q_{ij}^\alpha : ij \in \Sigma, \alpha \leq \ell\}.$$

By construction, any triad from \mathcal{Q} involving a pair $V_i V_j \in \Sigma$ will be disc_2 -irregular, and thus cannot be disc_3 -irregular. Any triad from \mathcal{Q} not involving a pair $V_i V_j$ from Σ will be $\text{disc}_{2,3}$ -regular, and thus cannot be disc_3 -irregular. Thus \mathcal{Q} has no disc_3 -irregular triads.

Conversely, suppose \mathcal{H} has the property that for all $\ell_0, t_0 \geq 1$, $\epsilon_1 > 0$, all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, there are n_0 and T, L such that for all $n \geq n_0$, every $G \in \mathcal{H}$ with at least n_0 vertices has an (ϵ_1, ϵ_2) -regular, $(t, \ell, \epsilon_2, \epsilon_1)$ -decomposition for some $t_0 \leq t \leq T$, and $\ell_0 \leq \ell \leq L$, such that no triad of \mathcal{P} is disc_3 -irregular. By Lemma 7.35, there is k such that $\text{HP}(k) \notin \text{Trip}(\mathcal{H})$, so by Fact 2.22, \mathcal{H} has finite VC_2 -dimension, and so by Theorem 2.24, \mathcal{H} is $\text{disc}_{2,3}$ -homogeneous. Fix $\epsilon_1 > 0$, a non-increasing $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$.

By the above, there exist n_0, L_0 and T_0 so that for all $n \geq n_0$, every $G \in \mathcal{H}$ with at least n_0 vertices, there are $t_0 \leq t \leq T_0$ and $\ell_0 \leq \ell \leq L_0$ and a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of $V(G)$ which is $(\epsilon_1, \epsilon_2(\ell))$ -regular and $\text{disc}_{2,3}$ -homogeneous with respect to G , and such that, moreover, no triad from \mathcal{P} is disc_3 -irregular with respect to G .

Assume $n \geq n_0$ and $H = (V, E) \in \mathcal{H}_n$. By assumption, there exists $t_0 \leq t \leq T_0$, $\ell_0 \leq \ell \leq L_0$ and \mathcal{P} , a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V which is $(\epsilon'_1, \epsilon'_2(\ell))$ -regular and $\text{disc}_{2,3}$ -homogeneous, and such that no triad of \mathcal{P} is disc_3 -irregular with respect to H . Let Σ be the set of pairs $V_i V_j$ where at least $(\epsilon'_1)^{1/2} |V_i| |V_j|$ of the $xy \in K_2[V_i V_j]$ are in a disc_2 -irregular $P \in \mathcal{P}_{\text{edge}}$. By assumption, $|\Sigma| \leq (\epsilon'_1)^{1/2} t^2$.

Set $s = \lceil (\epsilon'_1)^{1/2} \ell \rceil$. Fix $ij \notin \Sigma$. By reindexing if necessary, we may assume that $P_{ij}^{s+1}, \dots, P_{ij}^\ell$ all satisfy $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$. Set $E'_0 = \bigcup_{i=1}^s P_{ij}^\alpha$, and for each $s+1 \leq u \leq \ell$, set $E'_{u-s} = P_{ij}^u$. Note the density of $E'_0 = \frac{s}{\ell}(1 \pm s\epsilon'_2(\ell))$.

Apply Lemma 3.24 to E'_0 with $\epsilon = \delta = \epsilon_2(\ell)$ and $p = 1/(\ell - s)$ to obtain $R_{ij}^1, \dots, R_{ij}^{\ell-s}$, where each R_{ij}^u has $\text{disc}_2(2s\epsilon'_2(\ell)^{1/2}, \frac{s}{\ell(\ell-s)})$. For each $1 \leq m \leq \ell - s$, let $Q_{ij}^m = S_{ij}^m \cup R_{ij}^m$. By Fact 3.23, each Q_{ij}^m has $\text{disc}_2(2s\epsilon'_2(\ell)^{1/2}, \frac{1}{\ell} + \frac{s}{\ell(\ell-s)})$. It is not hard to check that $\frac{1}{\ell} + \frac{s}{\ell(\ell-s)} = \frac{1}{\ell-s}$. Thus, each Q_{ij}^m has $\text{disc}_2(\epsilon_2(k'), \frac{1}{k'})$ (using choice of ϵ'_2), where $k' = \ell - s$. Given $ij \notin \Sigma$, choose any partition $V_i \times V_j = \bigcup_{\alpha \leq k'} Q_{ij}^\alpha$.

Let \mathcal{Q} consist of $\{V_1, \dots, V_t\}$, and $\{Q_{ij}^\alpha : ij \in \binom{[t]}{2}\}$. By construction, this is a $(t, k', \epsilon_1, \epsilon_2(k'))$ -decomposition of V . Suppose $ij, ik, jk \notin \Sigma$, and consider a triad of the form $Q_{ijk}^{\alpha, \beta, \gamma} = (V_i \cup V_j \cup V_k, Q_{ij}^\alpha \cup Q_{ik}^\beta \cup Q_{jk}^\gamma)$, for some $1 \leq \alpha, \beta, \gamma \leq k'$. By construction, there was some triad of the form $P_{ijk}^{\alpha', \beta', \gamma'}$ which had $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$ such that

$$|Q_{ij}^\alpha \setminus P_{ij}^\alpha| < \frac{s}{\ell k'}(1 + 2s\epsilon'_2(\ell))|V_i||V_j| < \epsilon_1^2|Q_{ij}^\alpha|,$$

where the inequality is because $s \leq (\epsilon'_1)^{1/2}\ell + 1$ and $\epsilon'_1 \ll \epsilon_1$. By assumption, $P_{ijk}^{\alpha', \beta', \gamma'}$ was a $\text{disc}_{2,3}$ -regular triad of \mathcal{P} . We show $Q_{ijk}^{\alpha, \beta, \gamma}$ is a $\text{disc}_{2,3}$ -regular triad of \mathcal{Q} . By Proposition 3.22, it suffices to show the density on $Q_{ijk}^{\alpha, \beta, \gamma}$ is in $[0, \epsilon_1^2] \cup (1 - \epsilon_1^2, 1]$. Since $P_{ijk}^{\alpha', \beta', \gamma'}$ has $\text{disc}_2(\epsilon'_2(\ell), 1/\ell)$, it is by assumption, $\text{disc}_{2,3}$ -regular, and thus also $\text{disc}_{2,3}$ -homogeneous. Thus there is $u \in \{0, 1\}$ such that

$$|E^u \cap K_3^{(2)}(P_{ijk}^{\alpha', \beta', \gamma'})| \geq (1 - \epsilon'_1)|K_3^{(2)}(P_{ijk}^{\alpha', \beta', \gamma'})|.$$

Then, using Corollary 3.6,

$$\begin{aligned} |E^u \cap K_3^{(2)}(Q_{ijk}^{\alpha, \beta, \gamma})| &\geq |K_3^{(2)}(Q_{ijk}^{\alpha, \beta, \gamma})| - \epsilon'_1|K_3^{(2)}(P_{ijk}^{\alpha', \beta', \gamma'})| - \frac{9\epsilon_1^2}{(k')^3}|V_i||V_j||V_k| \\ &\geq \frac{|V_i||V_j||V_k|}{(k')^3} \left(1 - \epsilon_1^2 - \epsilon'_1 \frac{(k')^3}{\ell^3} - 9\epsilon_1^2\right) \\ &\geq (1 - \epsilon_1)|K_3^{(2)}(Q_{ijk}^{\alpha, \beta, \gamma})|, \end{aligned}$$

where the inequality is by our choice of $\epsilon'_1 \ll \epsilon_1$ and since $k' \geq (1 - \epsilon_1)\ell$. Thus $Q_{ijk}^{\alpha, \beta, \gamma}$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(k'))$. Thus in \mathcal{Q} all irregular triads use a pair from Σ , so it has binary $\text{disc}_{2,3}$ -error, as desired. \square

APPENDIX A. FACTS ABOUT \sim -CLASSES

This section shows various facts about \sim -classes. We begin with Proposition 2.19, which says that all the decomposition properties of interest in this paper are closed under \sim -classes. This boils down to the following two lemmas.

Lemma A.1. *For all $\epsilon > 0$ and $T \geq 1$, there is $\delta > 0$ so that the following holds. Suppose $H = (V, E)$, and $H' = (V, E')$ are two graphs on the same vertex set V , and assume H and H' are δ -close. Suppose V_1, \dots, V_t is an equipartition of V with $t \leq T$. If $ijk \in \binom{[t]}{3}$ and $V_i V_j V_k$ is has $\text{vdisc}_3(\epsilon)$ in H with density r , then it has $\text{vdisc}_3(3\epsilon)$ in H' with density $r \pm \epsilon$.*

Proof. Fix $\epsilon > 0$ and $T \geq 1$. Define $\delta = \epsilon^2/T^3$ and $N = \epsilon^{-1}T$. Suppose $H = (V, E)$, and $H' = (V, E')$ are two graphs on the same vertex set V of size $n \geq N$, and assume H and H' are δ -close. Suppose V_1, \dots, V_t is an equipartition of V with $t \leq T$. Suppose $ijk \in \binom{[t]}{3}$ and $V_i V_j V_k$ is has $\text{vdisc}_3(\epsilon)$ in H with density r . Then

$$|E' \cap K_3[V_i, V_j, V_k]| = |E \cap K_3[V_i, V_j, V_k]| \pm \delta|V|^3 = r|V_i||V_j||V_k| \pm \delta n^3 = (r \pm \delta t^3)|V_i||V_j||V_k|.$$

By definition of δ and since $t \leq T$, this shows the density of $V_i V_j V_k$ in H' is within ϵ of r . Suppose now that $V'_i \subseteq V_i$, $V'_j \subseteq V_j$, and $V'_k \subseteq V_k$. Then, since H and H' are δ close, and $V_i V_j V_k$ has $\text{vdisc}_3(\epsilon)$ in H ,

$$\begin{aligned} |E' \cap K_3[V'_i, V'_j, V'_k]| &= |E \cap K_3[V'_i, V'_j, V'_k]| \pm \delta |V|^3 = (r \pm \epsilon) |V'_i| |V'_j| |V'_k| \pm \delta n^3 \\ &= (r \pm 2\epsilon) |V_i| |V_j| |V_k|. \end{aligned}$$

Since $V_i V_j V_k$ has density $r \pm \epsilon$, this shows $V_i V_j V_k$ satisfies $\text{vdisc}_3(3\epsilon)$ in H' . \square

Lemma A.2. *For all $\epsilon > 0$, there is $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ such that for all $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfying $\epsilon_2(x) \leq \epsilon_2^*(x)$ for all $x \geq 1$, and all $L, T \geq 1$, there is $\delta > 0$ so that the following holds. Suppose $H = (V, E)$, and $H' = (V, E')$ are two graphs on the same vertex set V of size $n \geq N$, such that H and H' are δ -close. Assume \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . Then for any $G_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ satisfying $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H with density r , then $G_{ijk}^{\alpha, \beta, \gamma}$ satisfies $\text{disc}_{2,3}(3\epsilon_1, \epsilon_2(\ell))$ with respect to H' with density $r \pm 2\epsilon_1$.*

Proof. Fix $\epsilon > 0$. Define $\epsilon_2^* : \mathbb{N} \rightarrow (0, 1]$ so that $\epsilon_2^*(x) \ll \epsilon/x^2$ for all $x \geq 1$. Now suppose $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ satisfies $\epsilon_2(x) \leq \epsilon_2^*(x)$ for all $x \geq 1$, and $L, T \geq 1$. Choose $\delta \ll \epsilon_1^4 \epsilon_2(L)^2 L^{-3} T^{-3}$.

Suppose $H = (V, E)$, and $H' = (V, E')$ are two graphs on the same vertex set V , and assume H and H' are δ -close. Suppose \mathcal{P} is a $(t, \ell, \epsilon_1, \epsilon_2(\ell))$ -decomposition of V . Suppose $G \in \text{Triads}(\mathcal{P})$ has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ in H with density r . Then using Corollary 3.6,

$$|(E \Delta E') \cap K_3(G)| \leq \delta n^3 \leq \epsilon_1^2 |K_3(G)|.$$

Consequently, if r' is the density of G with respect to H' , then $|r' - r| \leq \frac{|(E \Delta E') \cap K_3(G)|}{|K_3(G)|} \leq \epsilon_1^2$. Now suppose $G' \subseteq G$ is a subgraph. Then since G has $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$ with respect to H with density r ,

$$\left| |E \cap K_3^{(2)}(G')| - r |K_3^{(2)}(G')| \right| \leq \epsilon_1 \frac{n^3}{\ell^3 t^3}.$$

Thus

$$\begin{aligned} & \left| |E' \cap K_3^{(2)}(G')| - r' |K_3^{(2)}(G')| \right| \\ & \leq \left| |E' \cap K_3^{(2)}(G')| - |E \cap K_3^{(2)}(G')| \right| + \left| |E \cap K_3^{(2)}(G')| - r' |K_3^{(2)}(G')| \right| \\ & \leq \delta n^3 + \left| |E \cap K_3^{(2)}(G')| - r |K_3^{(2)}(G')| \right| + |r - r'| |K_3^{(2)}(G')| \\ & \leq \delta n^3 + \epsilon_1 \frac{n^3}{\ell^3 t^3} + \epsilon_1^2 |K_3^{(2)}(G')| \\ & \leq 3\epsilon_1 \frac{n^3}{\ell^3 t^3}. \end{aligned}$$

This shows that G has $\text{disc}_{2,3}(3\epsilon_1, \epsilon_2(\ell))$ with respect to H' with density r' . \square

PROOF OF PROOF OF PROPOSITION 2.19: Assume \mathcal{H} and \mathcal{H}' are hereditary 3-graph properties and \mathcal{H} is close to \mathcal{H}' . Suppose \mathcal{H}' satisfies one of the following.

- (a) \mathcal{H}' is vdisc_3 -homogeneous.
- (b) \mathcal{H}' admits binary vdisc_3 -error.
- (c) \mathcal{H}' admits zero vdisc_3 -homogeneous.
- (d) \mathcal{H}' is $\text{disc}_{2,3}$ -homogeneous.
- (e) \mathcal{H}' admits linear vdisc_3 -error.
- (f) \mathcal{H}' admits binary vdisc_3 -error.
- (g) \mathcal{H}' admits zero vdisc_3 -error.

Then we show \mathcal{H} satisfies each of these, respectively. Assume first that (a), (b), or (c) hold. Fix $\epsilon > 0$ and $t_0 \geq 1$. By assumption, there exist $T = T(\epsilon/2, t_0)$, $N = N(\epsilon/2, t_0)$ be such that for every $n \geq N$, the following holds. For every $G \in \mathcal{H}'_n$, there is a decomposition $V(G) = V_1 \cup \dots \cup V_t$ for some $t_0 \leq t \leq T$, such that the following holds (respectively).

- (1) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon t^3/5$ and for all $ijk \notin \Sigma$, $V_i V_j V_k$ is $\epsilon/5$ -homogeneous with respect to G .
- (2) There is $\Gamma \subseteq \binom{[t]}{2}$ such that $|\Gamma| \leq \epsilon t^2/5$ and for all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, (V_i, V_j, V_k) has $\text{vdisc}_3(\epsilon/5)$ with respect to G .
- (3) For all $ijk \in \binom{[t]}{3}$, $V_i V_j V_k$ satisfies $\text{vdisc}_3(\epsilon/5)$ with respect to G .

Let δ be as in Lemma A.1 for $\epsilon/5$ and T . Choose N' sufficiently large so that for every $n \geq N'$ and $G \in \mathcal{H}_n$, there is $G' \in \mathcal{H}'_n$ such that G and G' are δ -close. Now assume $n \geq N'$ and $G = (V, E) \in \mathcal{H}$ with $|V| = n$. Then there is $G' = (V, E') \in \mathcal{H}'$ which is δ -close to G . Further, there are $t_0 \leq t \leq T$ and an equipartition $V = V_1 \cup \dots \cup V_t$ such that the following holds (respectively).

- (1) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon t^3/5$ and for all $ijk \notin \Sigma$, $V_i V_j V_k$ is $\epsilon/5$ -homogeneous with respect to G .
- (2) There is $\Gamma \subseteq \binom{[t]}{2}$ such that $|\Gamma| \leq \epsilon t^2/5$ and for all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, (V_i, V_j, V_k) has $\text{vdisc}_3(\epsilon/5)$ with respect to G .
- (3) For all $ijk \in \binom{[t]}{3}$, $V_i V_j V_k$ satisfies $\text{vdisc}_3(\epsilon/5)$ with respect to G .

By Lemma A.1, the following holds (respectively).

- (1) For all $ijk \notin \Sigma$, $V_i V_j V_k$ is ϵ -homogeneous with respect to G' .
- (2) For all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, (V_i, V_j, V_k) has $\text{vdisc}_3(\epsilon)$ with respect to G' .
- (3) For all $ijk \in \binom{[t]}{3}$, $V_i V_j V_k$ satisfies $\text{vdisc}_3(\epsilon)$ with respect to G' .

Thus we have shown that \mathcal{H} satisfies (a), (b), or (c), respectively.

Assume now that (d), (e), (f), or (g) hold, respectively. Fix $\epsilon_1 > 0$, $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$, and $t_0, \ell_0 \geq 1$. Let ϵ_2^* be as in Lemma A.2 for $\epsilon_1/5$. Let $\epsilon'_2(x) = \min\{\epsilon_2(x), \epsilon_2^*(x)\}$. Given $\ell_0, t_0 \geq 1$, choose N, T, L such that for every $G = (V, E) \in \mathcal{H}'$ on at least N vertices, there are $t_0 \leq t \leq T$ and $\ell_0 \leq \ell \leq L$ and an $(\epsilon_1/5, \epsilon'_2(\ell), t, \ell)$ -decomposition \mathcal{P} of V such that the following hold (respectively).

- (1) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon_1 t^3/5$ and for all $ijk \notin \Sigma$, every triad $P \in \text{Triads}(\mathcal{P})$ is $\epsilon_1/5$ -homogeneous with respect to G .
- (2) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon_1 t^3/5$ and for all $ijk \in \binom{[t]}{3} \setminus \Sigma$, every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon_2(\ell))$ with respect to G .

- (3) There is $\Gamma \subseteq \binom{[t]}{2}$ such that $|\Gamma| \leq \epsilon_1 t^2/5$ and for all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon_2(\ell))$ with respect to G .
- (4) Every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon_2(\ell))$ with respect to G .

Let δ be as in Lemma A.2 for $\epsilon_1/5$, ϵ'_2 , T and L . Let N be such that any $G \in \mathcal{H}$ on at least N vertices is δ -close to some element of \mathcal{H}' . Now assume $n \geq N$ and $G = (V, E) \in \mathcal{H}$ with $|V| = E$. Then there is $G' = (V, E') \in \mathcal{H}'$ which is δ -close to G . By assumption there are $t_0 \leq t \leq T$ and $\ell_0 \leq \ell \leq L$ and a $(t, \ell, \epsilon_1/5, \epsilon'_2(\ell))$ -decomposition \mathcal{P} of V such that the following hold (respectively).

- (1) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon_1 t^3/5$ and for all $ijk \notin \Sigma$, every triad $P_{ijk}^{\alpha, \beta, \gamma} \in \text{Triads}(\mathcal{P})$ is $\epsilon_1/5$ -homogeneous with respect to G' .
- (2) There is $\Sigma \subseteq \binom{[t]}{3}$ such that $|\Sigma| \leq \epsilon_1 t^3/5$ and for all $ijk \in \binom{[t]}{3} \setminus \Sigma$, every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon'_2(\ell))$ with respect to G' .
- (3) There is $\Gamma \subseteq \binom{[t]}{2}$ such that $|\Gamma| \leq \epsilon_1 t^2/5$ and for all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon'_2(\ell))$ with respect to G' .
- (4) Every triad $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon'_2(\ell))$ with respect to G' .

By Lemma A.2, the following hold (respectively).

- (1) For all $ijk \notin \Sigma$, every $P \in \text{Triads}(\mathcal{P})$ is ϵ_1 -homogeneous with respect to G' .
- (2) For all $ijk \in \binom{[t]}{3} \setminus \Sigma$, every $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon'_2(\ell))$ with respect to G' .
- (3) For all $ijk \in \binom{[t]}{3}$ with $ij, jk, ik \notin \Gamma$, every $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon'_2(\ell))$ with respect to G' .
- (4) Every $P \in \text{Triads}(\mathcal{P})$ satisfies $\text{disc}_{2,3}(\epsilon_1/5, \epsilon'_2(\ell))$ with respect to G' .

This shows that \mathcal{H} has (d), (e), (f), or (g), respectively. \square

We now prove Theorem 2.31, whose proof is very similar to that of Theorem 2.41.

PROOF OF THEOREM 2.31: Suppose $\text{Trip}(\mathcal{H})$ contains $\overline{U}(n)$ for all n . Fix a WNIP property \mathcal{H}' . We show \mathcal{H} is far from \mathcal{H}' . By assumption, there is some k so that $U^*(k) \notin \text{Trip}(\mathcal{H}')$. Let $\epsilon = (1/2k)^{2^k+k+1}/|\mathcal{B}(U^*(k))|$ and let $N \gg 2^k \epsilon^{-1}$. For all $n \geq N$, there is $G = (V, E) \in \mathcal{H}$ such that $\text{Trip}(G)$ contains $\overline{U}(n)$ and an induced sub-3-graph. This means there are $\{a_i, c_i : i \in [n]\} \subseteq V$ and $\{b_S : S \subseteq \{c_1, \dots, c_n\}\} \subseteq V$ such that $a_i b_S c_t \in E$ if and only if $c_t \in S$. Set $n_1 = \lfloor n/k \rfloor$ and choose C_1, \dots, C_k to be disjoint subsets of $\{c_1, \dots, c_{kn_1}\}$ of size n_1 . For each $X \subseteq [k]$, let

$$B_X = \{b_S : \text{for each } i \in X, C_i \subseteq S \text{ and for each } i \notin X, C_i \cap S = \emptyset\}.$$

Note $|B_X| \geq 2^{n-kn_1}$, so we can take $B'_X \subseteq B_X$ of size n_1 . We construct many induced subgraphs of G as follows.

- (1) Choose a_i for some $i \in [n]$. There are n choices.
- (2) Choose some $(c_{i_1}, \dots, c_{i_k}) \in C_1 \times \dots \times C_k$. There are n_1^k choices.
- (3) For each $X \subseteq [k]$, choose some $d_X \in B'_X$. There are $n_1^{2^k}$ choices.
- (4) Put $G[\{a_i\} \cup \{c_{i_1}, \dots, c_{i_k}\} \cup \{d_X : X \subseteq [k]\}]$ in S .

By construction, every element of S is isomorphic to some element in $\mathcal{B}(U^*(k))$ and

$$|S| \geq n_1^{k+2^k+1} > \epsilon |\mathcal{B}(U^*(k))| n^{k+2^k+1},$$

where the inequality is by definition of n_1 and ϵ . By the pigeonhole principle, there is some $G(n) \in \mathcal{B}(U^*(k))$ so that S contains at least ϵn^{k+2^k+1} elements isomorphic to $G(n)$. By the pigeonhole principle, there is $G' \in \mathcal{B}(U^*(k))$ so that $G' = G(n)$ for arbitrarily large n . By definition, this implies $G' \in \Delta(\mathcal{H})$. Clearly $G' \notin \Delta(\mathcal{H}')$ (since $U^*(k) \notin \text{Trip}(\mathcal{H}')$), so \mathcal{H} is far from \mathcal{H}' .

Conversely, suppose \mathcal{H} is far from every WNIP property \mathcal{H}' . By Lemma 3.31, \mathcal{H}' is WNIP if and only if, for some $k \geq 1$, \mathcal{H}' contains no clean copies of $U^*(k)$. Thus, since \mathcal{H} is far from every WNIP property, we know that for all $k \geq 1$, there is $\delta > 0$ so that for arbitrarily large n , there is $G(n) \in \mathcal{H}$ and $\Gamma(n) \in \mathcal{B}(U^*(k))$ so that $G(n)$ is not $\delta/|\mathcal{B}(U^*(k))|$ -close to being $\Gamma(n)$ -free. For each $\Gamma \in \mathcal{B}(U^*(k))$, let $c_k(\Gamma)$, $N_k(\Gamma)$ be as in Theorem 3.14 for Γ and $\delta/|\mathcal{B}(U^*(k))|$ -close. Then set $c_k = \min\{c_k(\Gamma) : \Gamma \in \mathcal{B}(U^*(k))\}$ and $N_k = \max\{N_k(\Gamma) : \Gamma \in \mathcal{B}(U^*(k))\}$. Choose any $N > N_k$. Then there is some $n \geq N$, so that that $G(n)$ is not δ -close to being $\Gamma(n)$ -free. Say $V(\Gamma(n)) = \{a, b_i, c_S : i \in [k], S \subseteq [k]\}$ and $ab_i c_S \in E(\Gamma(n))$ if and only if $i \in S$. By Theorem 3.14, G contains at least $c_k n^{k+1+2^k}$ induced sub-3-graphs isomorphic to $\Gamma(n)$. Let S be the set of these induced sub-3-graphs. For each $H \in S$, there is a vertex a_H and sets $D_H = \{d_i^H : i \in [k]\}$, $E_H = \{e_S^H : S \subseteq [k]\}$, such that $a_H d_i^H e_S^H \in E$ if and only if $i \in S$. Since $|S| \geq c_k n^{k+1+2^k}$, there are sets D and E and $X \subseteq S$ of size at least $c_k n/k!(2^k)!$ such that $(D, E) = (D_H, E_H)$ for all $H \in X$. Then we must have that $\{a_H : H \in X\} = |X|$. Let $Y \subseteq \{a_H : H \in X\}$ any set of size k . Then $G[Y \cup D \cup E] \in \mathcal{H}$ and $\text{Trip}(G[Y \cup D \cup E]) \cong \overline{U}(k)$. Thus $\overline{U}(k) \in \text{Trip}(\mathcal{H})$. \square

APPENDIX B. LEMMAS ABOUT COMBINING AND REFINING DECOMPOSITIONS

In this section, we prove the results stated in Section 3.2. We will also prove Theorem 4.1 here, as it uses some of these lemmas.

We begin with the proof of Corollary 3.6.

PROOF OF COROLLARY 3.6: Fix $t \geq 2$, $\epsilon > 0$ and $r \in (0, 1]$. Without loss of generality, assume $\epsilon \leq 1/2$. Choose $\mu \ll_t \gamma \ll_t \epsilon, r$.

Now assume $V = U_1 \cup \dots \cup U_t$ and $G = (V, E)$ is a t -partite graph with vertex partition U_1, \dots, U_t , such that for each $1 \leq i < j \leq t$, $G[U_i, U_j]$ has $\text{disc}_2(\mu, r)$. For each $1 \leq i < j \leq t$, let d_{ij} be such that $|E \cap K_2[U_i, U_j]| = d_{ij}|U_i||U_j|$. By assumption, each $d_{ij} = r \pm \mu$. By Proposition 3.5,

$$\left| K_t(G) - \prod_{ij \in \binom{[t]}{2}} d_{ij} \prod_{i=1}^t |U_i| \right| \leq \gamma \prod_{i=1}^t |U_i|.$$

Let $d = \prod_{ij \in \binom{[t]}{2}} d_{ij}$. Note $d = (1 \pm \mu)^{\binom{t}{2}} r^{\binom{t}{2}} = (1 \pm \epsilon/2)^{\binom{t}{2}}$, where the last equality is since $\mu \ll \epsilon$. Thus

$$|K_t(G)| = (d \pm \gamma) \prod_{i=1}^t |U_i| = d \left(1 \pm \gamma d^{-1}\right) \prod_{i=1}^t |U_i|$$

Since $d = (1 \pm \epsilon/2)^{\binom{t}{2}}$ and $\gamma \ll r, \epsilon$, $\gamma d^{-1} \in (-2\epsilon^2, 2\epsilon^2)$. Thus

$$|K_t(G)| = (1 \pm 2\epsilon^2) d \prod_{i=1}^t |U_i| = (1 \pm 2\epsilon^2) (1 \pm \epsilon/2)^{\binom{t}{2}} \prod_{i=1}^t |U_i|.$$

Since $\epsilon \leq 1/2$ this is equal to $(1 \pm \epsilon) r^{\binom{t}{2}} \prod_{i=1}^t |U_i|$, as desired. \square

We now prove Lemma 3.19.

PROOF OF LEMMA 3.19: Fix $k \geq 1$ and $\delta_3 > 0$. Let $\delta'_3 = \delta_3/32k^3$. Suppose now $d_2, d_3 \in (0, 1]$ and $0 < \delta_2 \leq d_2^3/8$. Set $\gamma = \delta_2/2k^3$ and let $\mu = \mu(3, \gamma, d_2)$ and $N_1 = N_1(3, \gamma, d_2)$ as in Proposition 3.5. Let $\delta'_2 \ll \mu/4k^2, \delta_2/4k^2, d_2\delta_2/(6k^4)$, and choose $N \gg k, N_1$.

Suppose $V = V_1 \cup V_2 \cup V_3$ is a vertex set of size $n \geq N$, and $G = (V_1 \cup V_2 \cup V_3, E)$ is a 3-partite graph. Assume $H = (V, R)$ is a 3-graph, and let $H' = (V, R \cap K_3^{(2)}(G))$. Assume that for each $1 \leq i < j \leq 3$, $G[V_i, V_j]$ has $\text{disc}_2(\delta'_2, d_2)$, and (H', G) has $\text{disc}_{2,3}(\delta'_2, \delta'_3)$ with density d_3 . Suppose that for each $i \in [3]$, $V'_i \subseteq V_i$ satisfies $|V'_i| = m'_i \geq m_i/k$, where $m_i := |V_i|$. Set $G' = G[V'_1, V'_2, V'_3]$ and $H'' = (V'_1 \cup V'_2 \cup V'_3, R \cap K_3^{(2)}(G'))$.

For each $1 \leq i < j \leq 3$, let d_{ij} be the density of $G[V_i, V_j]$. By assumption $d_{ij} = d_2 \pm \delta'_2$. By Lemma 3.18, for each $ij \in \binom{[3]}{2}$, $G'[V_i, V_j]$ has $\text{disc}_2(2k^2\delta'_2, d'_{ij})$ for some $d'_{ij} = d_{ij} \pm k^2\delta'_2$. Since $\delta'_2 \ll \delta_2/4k^2, \mu$, this means $G'[V'_i, V'_j]$ has $\text{disc}_2(\mu, d_2)$. Thus, by Proposition 3.5, we have

$$|K_3^{(2)}(G')| = d_2^3(1 \pm \gamma)m'_1m'_2m'_3.$$

Since (H', G) has $\text{disc}_{2,3}(\delta'_2, \delta'_3)$,

$$||R \cap K_3^{(2)}(G')| - d_3|K_3^{(2)}(G')|| \leq \delta'_3 d_2^3 m_1 m_2 m_3.$$

Therefore if d'_3 is such that $|R \cap K_3^{(2)}(G')| = d'_3 |K_3^{(2)}(G')|$, then

$$\left| d'_3 - d_3 \right| \leq \frac{\delta'_3 d_2^3 m_1 m_2 m_3}{|K_3^{(2)}(G')|} \leq \frac{\delta'_3 d_2^3 m_1 m_2 m_3}{d_2^3 (1 - \gamma) m'_1 m'_2 m'_3} \leq 2\delta'_3,$$

where the last inequality is by assumption, because $\gamma \leq 1/2$, and because each $m'_i \geq m_i/k$. Suppose $G'' \subseteq G'$. Since (H', G) has $\text{disc}_{2,3}(\delta'_2, \delta'_3)$,

$$\left| |R \cap K_3^{(2)}(G'')| - d_3 |K_3^{(2)}(G'')| \right| \leq \delta'_3 d_2^3 m_1 m_2 m_3 \leq \delta_3 d_2^3 m'_1 m'_2 m'_3 / 2,$$

where the equality is by definition of δ'_3 and since each $m_i \geq km_0$. Thus

$$\begin{aligned}
\left| |R \cap K_3^{(2)}(G'')| - d'_3 |K_3^{(2)}(G'')| \right| &\leq \delta_3 d_2^3 m'_1 m'_2 m'_3 / 2 + \left| d_3 |K_3^{(2)}(G'')| - d'_3 |K_3^{(2)}(G'')| \right| \\
&\leq \delta_3 d_2^3 m'_1 m'_2 m'_3 / 2 + 2\delta'_3 |K_3^{(2)}(G'')| \\
&\leq \delta_3 d_2^3 m'_1 m'_2 m'_3 / 2 + 2\delta'_3 d_2^3 (1 + \gamma) m'_1 m'_2 m'_3 \\
&\leq \delta_3 d_2^8 m'_1 m'_2 m'_3,
\end{aligned}$$

where the last inequality is by definition of δ'_2 and γ . This finishes our verification that (H'', G') has $\text{disc}_{2,3}(\delta_3, \delta_2)$. \square

PROOF OF FACT 3.23: Assume E_1, E_2 are disjoint subsets of $K_2[U, V]$. We begin with part (a). Assume $(U \cup V, E_1)$ has $\text{disc}_2(\epsilon_1, d_1)$, $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon_2, d_2)$. We show $(U \cup V, E_1 \cup E_2)$ has $\text{disc}_2(\epsilon_1 + \epsilon_2, d_1 + d_2)$. Suppose $U' \subseteq U$ and $V' \subseteq V$. By assumption, $||E_1 \cap K_2[U', V']| - d_1|U'|||V'||| \leq \epsilon_1|U||V|$, and $||E_2 \cap K_2[U', V']| - d_2|U'|||V'||| \leq \epsilon_2|U||V|$. Then since $E_1 \cap E_2 = \emptyset$,

$$\begin{aligned}
|(E_1 \cup E_2) \cap K_2[U', V']| &= |E_1 \cap K_2[U', V']| + |E_2 \cap K_2[U', V']| - |(E_1 \cap E_2) \cap K_2[U', V']| \\
&= |E_1 \cap K_2[U', V']| + |E_2 \cap K_2[U', V']|.
\end{aligned}$$

By above, this is at most $(d_1 + d_2)|U'|||V'| + \epsilon_1|U||V| + \epsilon_2|U||V|$ and at least $(d_1 + d_2)|U'|||V'| - \epsilon_1|U||V| - \epsilon_2|U||V|$ since $(U \cup V, E_1)$ has $\text{disc}_2(\epsilon_1, d_1)$, $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon_2, d_2)$. Thus $|(E_1 \cup E_2) \cap K_2[U', V']| - (d_1 + d_2)|U'|||V'| \leq (\epsilon_1 + \epsilon_2)|U||V|$, as desired.

We now show (b). Assume $(U \cup V, E_1 \cup E_2)$ has $\text{disc}_2(\epsilon, d)$, $(U \cup V, E_1)$ has $\text{disc}_2(\epsilon_1, d_1)$. We show $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon + \epsilon_2, d - d_1)$. Suppose $U' \subseteq U$ and $V' \subseteq V$. Then

$$\begin{aligned}
&\left| |E_2 \cap K_2[U', V']| - (d - d_1)|U'|||V'| \right| \\
&= \left| |(E_1 \cup E_2) \cap K_2[U', V']| - |E_1 \cap K_2[U', V']| - (d - d_1)|U'|||V'| \right| \\
&\leq \left| |(E_1 \cup E_2) \cap K_2[U', V']| - d|U'|||V'| \right| + \left| |E_1 \cap K_2[U', V']| - d_1|U'|||V'| \right|.
\end{aligned}$$

By assumption, this is at most $\epsilon|U||V| + \epsilon_1|U||V| = (\epsilon + \epsilon_1)|U||V|$. This shows $(U \cup V, E_2)$ has $\text{disc}_2(\epsilon + \epsilon_2, d - d_1)$. \square

Our next goal is to prove Lemma 3.27. This will require two additional tools. The first is a multi-colored version of Szemerédi's regularity lemma (see e.g. [24, 32]).

Theorem B.1 (Multi-colored regularity lemma). *For all $r \geq 1$, $\epsilon > 0$, and $m \geq 1$, there is $M = M(\epsilon, m)$ such that the following holds. Suppose $G = (V, E_1, \dots, E_r)$ where $|V| \geq M$ and E_1, \dots, E_r are disjoint subsets of $\binom{V}{2}$. Then for any equipartition $\mathcal{P} = \{V_1, \dots, V_m\}$ of V , there is $m' \leq M$ and an equipartition, $\mathcal{P}' = \{V'_1, \dots, V'_{m'}\}$ which refines \mathcal{P} , such that all but at most $\epsilon(m')^2$ pairs (V'_i, V'_j) , $(V'_i \cup V'_j, E_u)$ satisfies $\text{disc}_2(\epsilon)$ for all $1 \leq u \leq r$.*

Note that $M = M(\epsilon, m)$ does not depend on the number of colors. In fact, it is the same bound as in the usual regularity lemma. Indeed, it is straightforward to prove Theorem B.1 by following the usual proof of the regularity lemma, with a slightly altered mean square

density. The mean square density increases at each step in an identical way to the usual proof, so the bound ends up identical. We now give a sketch of this proof, as we could not find this explicitly in the literature.

We will follow closely the proof of the usual regularity lemma (see, for example [2]). First, suppose that $r \geq 1$, for each $u \in [r]$, (V, E_u) is a graph on the vertex set V , where $|V| = n$, and $u \neq v$ implies $E_u \cap E_v = \emptyset$. Given $X, Y \subseteq V$, define

$$d_u(X, Y) = \frac{|E_u \cap K_2[X, Y]|}{|X||Y|}.$$

Note that $d_1(X, Y) + \dots + d_r(X, Y) \leq 1$. We then define, for each $u \in [r]$,

$$\text{msd}_u(X, Y) = \frac{|X||Y|}{n^2} d_u^2(X, Y).$$

Here, msd stands for “mean square density.” Note this is just the usual mean square density for the graph (V, E_u) . We then define

$$\text{msd}(X, Y) = \sum_{u=1}^r \text{msd}_u(X, Y) = \frac{|X||Y|}{n^2} \sum_{u=1}^r d_u^2(X, Y).$$

Note that since $0 \leq d_1(X, Y) + \dots + d_r(X, Y) \leq 1$, $\sum_{u=1}^r d_u^2(X, Y) \leq 1$, so $\text{msd}(X, Y) \leq \frac{|X||Y|}{n^2}$. If \mathcal{P}_X and \mathcal{P}_Y are partitions of X and Y , define, for each $u \in [r]$,

$$\text{msd}_u(\mathcal{P}_X, \mathcal{P}_Y) = \sum_{U \in \mathcal{P}_X} \sum_{W \in \mathcal{P}_Y} \text{msd}_u(U, W),$$

and set

$$\text{msd}(\mathcal{P}_X, \mathcal{P}_Y) = \sum_{u=1}^r \text{msd}_u(\mathcal{P}_X, \mathcal{P}_Y).$$

For a partition $\mathcal{P} = \{V_1, \dots, V_k\}$ of V , let $\text{msd}_u(\mathcal{P}) = \text{msd}_u(\mathcal{P}, \mathcal{P})$ and $\text{msd}(\mathcal{P}) = \text{msd}(\mathcal{P}, \mathcal{P})$. Note

$$\text{msd}(\mathcal{P}) = \sum_{i=1}^k \sum_{j=1}^k \frac{|V_i||V_j|}{n^2} \left(\sum_{u=1}^r d_u^2(V_i, V_j) \right) \leq \sum_{i=1}^k \sum_{j=1}^k \frac{|V_i||V_j|}{n^2} = 1$$

Lemma B.2. *For any partitions \mathcal{P}_X of X and \mathcal{P}_Y of Y , $\text{msd}(\mathcal{P}_X, \mathcal{P}_Y) \geq \text{msd}(X, Y)$.*

Proof. It is well known that for each $1 \leq u \leq r$, $\text{msd}_u(\mathcal{P}_X, \mathcal{P}_Y) \geq \text{msd}_u(X, Y)$ (see e.g. [26, Corollary 3.7]). By definition, this implies $\text{msd}(\mathcal{P}_X, \mathcal{P}_Y) \geq \text{msd}(X, Y)$. \square

Lemma B.3. *If $(X \cup Y, E_u \cap K_2[X, Y])$ does not satisfy $\text{disc}_2(\epsilon)$, then there are $X_1 \subseteq X$ and $Y_1 \subseteq Y$, such that*

$$\text{msd}(\{X_1, X \setminus X_1\}, \{Y_1, Y \setminus Y_1\}) > \text{msd}(X, Y) + \epsilon^4 \frac{|X||Y|}{n^2}.$$

Proof. Suppose $(X \cup Y, E_u \cap K_2[X, Y])$ fails $\text{disc}_2(\epsilon)$. Then it is easy to check that (X, Y) is not ϵ -regular, say this is witnessed by $X_1 \subseteq X$ and $Y_1 \subseteq Y$. It is well known this implies that $\text{msd}_u(\{X_1, X \setminus X_1\}, \{Y_1, Y \setminus Y_1\}) > \text{msd}_u(X, Y) + \epsilon^4 \frac{|X||Y|}{n^2}$ (see e.g. [26]). So

$$\begin{aligned} \text{msd}(\{X_1, X \setminus X_1\}, \{Y_1, Y \setminus Y_1\}) &= \sum_{s=1}^r \text{msd}_s(\{X_1, X \setminus X_1\}, \{Y_1, Y \setminus Y_1\}) \\ &> \text{msd}_u(X, Y) + \epsilon^4 \frac{|X||Y|}{n^2} + \sum_{s \in [r] \setminus \{u\}} \text{msd}_s(\{X_1, X \setminus X_1\}, \{Y_1, Y \setminus Y_1\}) \\ &\geq \text{msd}_u(X, Y) + \epsilon^4 \frac{|X||Y|}{n^2} + \sum_{s \in [r] \setminus \{u\}} \text{msd}_s(X, Y) \\ &= \text{msd}(X, Y) + \epsilon^4 \frac{|X||Y|}{n^2}, \end{aligned}$$

where the last inequality is due to Lemma B.3. \square

One can now deduce Theorem B.1 exactly as in usual proofs of the regularity lemma (see e.g. [47], [2]). The second tool used in the proof of Lemma 3.27 is Lemma B.4 below, which is a corollary of Lemma 3.24.

Lemma B.4. *Suppose $\epsilon_1 > 0$, $\ell \geq 1$, and $0 < \epsilon_2 < \epsilon_1/(64\ell')$, where $\ell' := \lceil 32\ell^2\epsilon_1^{-2} \rceil$. There is $m_0 = m_0(\epsilon_1, \epsilon_2, \ell, \delta)$ such that the following holds. Suppose $|U| = |V| = m \geq m_0$, $s \leq \ell$, and $K_2(U, V) = E_0 \cup \dots \cup E_s$ is a partition such that for each $i \in [s]$, $(U \cup V, E_i)$ has $\text{disc}(\epsilon_2)$, and $|E_0| \leq \epsilon_1 m^2$. Then there is a partition*

$$K_2(U, V) = E'_{0,1} \cup \dots \cup E'_{0,k} \cup E'_1 \cup \dots \cup E'_{k'},$$

where $k + k' = \ell'$, such that the following hold.

- (i) For each $1 \leq i \leq k$, and $1 \leq \alpha \leq k'$, $(U \cup V, E'_{0,i})$ and $(U \cup V, E_\alpha)$ have $\text{disc}_2(\epsilon_2^{1/3}, 1/\ell')$.
- (ii) For each $1 \leq \alpha \leq k'$, there is $1 \leq j \leq s$ such that $|E_j| \geq (\sqrt{2/32}\epsilon_1/s)m^2$ and $|E'_\alpha \setminus E_j| < (2\epsilon_2^{1/2}/\ell')m^2$. Moreover, if $k' < \ell'$, then $E'_\alpha \setminus E_j = \emptyset$.
- (iii) If $E'_0 = \bigcup_{j=1}^k E'_{0,j}$, then $|E'_0| \leq 2\epsilon_1 m^2$.

Proof. Fix $\epsilon_1 > 0$, $\ell \geq 1$, and $0 < \epsilon_2 < \epsilon_1/(64\ell')$, where $\ell' := \lceil 32\ell^2\epsilon_1^{-2} \rceil$. Set $\delta = \epsilon_2/2(\ell')^2$, and let $m_0 = m_0(\epsilon_2/2\ell', \ell', \delta/2\ell')$ be as in Lemma 3.24. Suppose $s \leq \ell$, $|U| = |V| = m \geq m_0$, and $K_2(U, V) = E_0 \cup \dots \cup E_s$ is a partition such that for each $1 \leq i \leq s$, $(U \cup V, E_i)$ has $\text{disc}_2(\epsilon_2)$, and such that $|E_0| \leq \epsilon_1 m^2$. For each $i \in \{0, \dots, s\}$, let $\rho_i = |E_i|/m^2$, and define

$$\Sigma = \left\{ i \in [s] : \rho_i \leq \sqrt{\frac{2}{\ell'}} \right\}.$$

Given $i \in [s] \setminus \Sigma$, set $p(i) = 1/\rho_i \ell'$ and $u(i) = \lceil 1/p(i) \rceil$. Observe that for all $i \in [s] \setminus \Sigma$, $\rho_i \geq \sqrt{2/\ell'}$ implies $p(i) < \rho_i/2$, and further, $\epsilon_2 \leq \epsilon_1$ implies $\rho_i \geq 2\epsilon_2$. Therefore, for each $i \in [s] \setminus \Sigma$, Lemma 3.24 implies there is a partition $E_i = R_{i,0} \cup \dots \cup R_{i,u(i)}$ such that

$|R_{i,0}| < p(i)\rho_i(1+\delta)m^2$ and such that for each $1 \leq j \leq u(i)$, $R_{i,j}$ has $\text{disc}_2(\epsilon_2)$ and density $\rho_i p(i)(1 \pm \delta) = \frac{1}{\ell'}(1 \pm \delta)$.

Let $k' = \sum_{i \in [s] \setminus \Sigma} u(i)$. Since $\delta < 1/(\ell')^2$ and

$$m^2 \geq \left| \bigcup_{i=1}^s \bigcup_{j=1}^{u(i)} R_{i,j} \right| \geq (1 - k'\delta) \frac{k'}{\ell'} m^2,$$

we must have that $k' \leq \ell'$. Fix an enumeration

$$\{R_{i,j} : i \in [s] \setminus \Sigma, 1 \leq j \leq u(i)\} = \{\Gamma_1, \dots, \Gamma_{k'}\},$$

and set $\Gamma_0 = E_0 \cup \bigcup_{i \in [s] \setminus \Sigma} R_{i,0} \cup \bigcup_{i \in \Sigma} E_i$. Note

$$\begin{aligned} |\Gamma_0| &\leq \epsilon_1 m^2 + (s - |\Sigma|) \left(\frac{1}{\ell'} (1 + \delta) m^2 \right) + |\Sigma| \sqrt{\frac{2}{\ell'}} m^2 \\ &\leq m^2 \left(\epsilon_1 + \frac{s}{\ell'} (1 + \delta) + |\Sigma| \left(\sqrt{\frac{2}{\ell'}} - \frac{1}{\ell'} (1 + \delta) \right) \right) \\ &\leq 4\epsilon_1 m^2, \end{aligned}$$

where the inequality is because $s \leq \ell \leq \epsilon_1 \ell'$, and $\sqrt{2}|\Sigma|/\sqrt{\ell'} \leq \sqrt{2}s\epsilon_1/\ell \leq 3\epsilon_1$. On other hand, since for each $i \in [k']$, Γ_i has density $1/\ell'(1 \pm \delta)$, we have that

$$(17) \quad |\Gamma_0| = \frac{\ell' - k'}{\ell'} (1 \pm k'\delta) m^2.$$

If $k' = \ell'$, then (17) implies that $|\Gamma_0| \leq k'\delta m^2 < \epsilon_2 m^2 / 2\ell'$. In this case, choose any equipartition of Γ_0 into ℓ' pieces, say $\Gamma_0 = E_1'' \cup \dots \cup E_{\ell'}''$, and for each $\alpha \in [\ell']$, set $E'_\alpha = \Gamma_\alpha \cup E_\alpha''$. Then set $k = 0$, so $k + k' = \ell'$. By construction, for each $\alpha \in [\ell']$, there is $j \in [s]$ such that $|E'_\alpha \setminus E_j| = |E_\alpha''| \leq \epsilon_2 m^2 / 2\ell'$, so (ii) holds. Further, since each E_α has $\text{disc}_2(\epsilon_2, 1/\ell')$ and $|E'_\alpha \Delta E_\alpha| \leq \epsilon_2 m^2 / 2\ell'$, it follows that E'_α has $\text{disc}_2(2\epsilon_2^{1/2})$ and density $1/\ell' \pm 2\sqrt{\epsilon_2}$, so (i) is satisfied. Since (iii) holds trivially, we are done.

If $k' = \ell' - 1$, set $k = 1$, so $k + k' = \ell'$. Define $E_{0,1} = \Gamma_0$, and for each $1 \leq i \leq \ell'$, set $E'_i = \Gamma_i$. Note that by Fact 3.23, $\Gamma_1 \cup \dots \cup \Gamma_{k'}$ has $\text{disc}_2(k'\epsilon_2, k'/\ell')$. Since $E_{0,1}$ is the compliment of this union, it has $\text{disc}_2(k'\epsilon_2, 1 - k'/\ell')$. Consequently, $E_{0,1}$ has $\text{disc}_2(\epsilon_2^{1/2}, 1/\ell')$ (since $k' = \ell' - 1$). Thus (i)-(iii) hold for $E'_{0,1} \cup E'_1 \cup \dots \cup E'_{k'}$.

Finally, assume $k' \leq \ell' - 2$. In this case, set $k = \ell' - k'$ and define $\rho'_0 = |\Gamma_0|/m^2$. Since $|\Gamma_0| \leq 2\epsilon_1 m^2$, $\rho'_0 \leq 2\epsilon_1$. By (17), this implies $(1 - 2\epsilon_1)\ell' \leq k'$. Let $\mu = \epsilon_2^2 / 2\ell'$, $u = k$, and $p = 1/u$. Using that $(1 - 2\epsilon_1)\ell' \leq k'$ and (17), we have that $p < \rho'_0 / 2$. By definition, $2\mu \leq p^{-1}$. Thus we may apply Lemma 3.24 to Γ_0 with μ , p , u , and $\delta' = \delta/(\ell')^2$. From this we obtain a partition $\Gamma_0 = E_{0,1} \cup \dots \cup E_{0,u}$ such that each $E_{0,i}$ has $\text{disc}_2(\mu)$ and density $\rho'_0 \frac{1}{u} (1 \pm \delta') = \frac{1}{\ell'} (1 \pm \delta)$. Thus each $E_{0,i}$ has $\text{disc}_2(\epsilon_2^{1/2}, 1/\ell')$, as desired. Thus (i)-(iii) are satisfied with this $k + k' = \ell'$ and $E_{0,1} \cup \dots \cup E_{0,k} \cup \Gamma_1 \cup \dots \cup \Gamma_{k'}$. \square

We now prove Lemma 3.27.

PROOF OF LEMMA 3.27: Fix $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Choose $\mu_1 \ll \epsilon_1$. Define $\mu_2 : \mathbb{N} \rightarrow (0, 1]$ by setting $\mu_2(x) \ll \mu_1 \epsilon_2 (\mu_1^{-1} x^2) / x^2$ for all $x \geq 1$. Let $M_{Sz}(x, y)$ denote the bound from Theorem B.1 with parameters x and y . Then set $f(x_1, y_1, x_2, y_2) = 32(y_1 y_2)^2 \mu_1^{-1}$, and set $g(x_1, y_1, x_2, y_2) = \mu_1^{-1} M_{Sz}(\mu_1^{-1} x_1 x_2, \mu_2 f(x_1, y_1, x_2, y_2))$. Now fix $\ell, t, r, s \geq 1$. Set $L = g(t, \ell, s, r)$, $T = f(t, \ell, s, r)$, and choose $N \gg T, L, \mu_2(L)^{-1}$.

Suppose $|V| = n \geq N$, \mathcal{P} is a (t, ℓ) -decomposition of V consisting of $\mathcal{P}_{vert} = \{V_i : i \in [t]\}$, $\mathcal{P}_{edge} = \{P_{ij}^\alpha : \alpha \leq \ell\}$, and assume \mathcal{Q} is an (s, r) -decomposition of V consisting of $\mathcal{Q}_{vert} = \{Q_i : i \in [s]\}$, $\mathcal{Q}_{edge} = \{Q_{ij}^\alpha : \alpha \in [r]\}$. For each $\alpha \in [\ell]$ and $\beta \in [r]$, define $P^\alpha = \bigcup_{ij \in \binom{[t]}{2}} P_{ij}^\alpha$ and $Q^\beta = \bigcup_{ij \in \binom{[s]}{2}} Q_{ij}^\beta$, and set $E^{\alpha, \beta} = P^\alpha \cap Q^\beta$.

For each $(i, j) \in [t] \times [s]$, define $X_{ij} = V_i \cap Q_j$, and then let $\mathcal{X} = \{X_{ij} : ij \in [t] \times [s]\}$. Set $m = (\mu_1)^2 n / ts$, and for each X_{ij} choose a partition $X_{ij} = X_{ij}^0 \cup \dots \cup X_{ij}^{s_{ij}}$ with the property that for each $1 \leq u \leq s_{ij}$, $|X_{ij}^u| = m$ and $|X_{ij}^0| < m$. Setting $X^0 = \bigcup_{ij \in [t] \times [s]} X_{ij}^0$, it is easy to see that $|X^0| \leq (\mu_1)^2 n$. Fix an enumeration U_1, \dots, U_τ of the set $\{X_{ij}^u : 1 \leq u \leq s_{ij}\}$, and note we must have $\tau \leq n/m \leq (\mu_1)^{-2} ts$. Let $\ell' = \lceil 32(\ell r)^2 (\mu_1)^{-2} \rceil$, and define $\rho_1 = \mu_1$ and $\rho_2 = \mu_2(\ell')$. Note that $\ell' \leq L$ and $\rho_2 < \rho_1 / (64\ell')$.

Consider the edge-colored graph with vertex set $V' = \bigcup_{i=1}^\tau U_i$, edge colors $F^{\alpha, \beta} := E^{\alpha, \beta} \cap \binom{V'}{2}$, for $\alpha \in [\ell]$ and $\beta \in [r]$. By construction $\mathcal{U} = \{U_1, \dots, U_\tau\}$ is an equipartition of V' . By Theorem B.1, there is $K \leq T$, and $\mathcal{S} = \{S_1, \dots, S_K\}$ an equipartition refining \mathcal{U} such that all but at most $\rho_2 K^2$ pairs from \mathcal{S} satisfy $\text{disc}_2(\rho_2)$ with respect to each of $F^{\alpha, \beta}$. Let Σ_1 be the set of pairs (S_i, S_j) from \mathcal{S} which do not satisfy $\text{disc}_2(\rho_2)$ with respect to all the $F^{\alpha, \beta}$. By assumption, $|\Sigma_1| \leq \rho_2 K^2$.

Our next step is to partition each $K_2[S_i, S_j]$ into equally sized quasirandom graphs.

For each $(S_i, S_j) \notin \Sigma_1$, define I_{ij} to be the set of $F^{\alpha, \beta}$ with density at most $\rho_1 / \ell s$ in (S_i, S_j) , and set $E_{ij}^0 = (\bigcup_{I_{ij}} E^{\alpha, \beta}) \cap K_2[S_i, S_j]$. Note $|E_{ij}^0| \leq \rho_1 |S_i| |S_j|$. By Lemma B.4, there is some $k_{ij} + k'_{ij} = \ell'$ and a partition

$$S_i \times S_j = W_{ij}^{0,1} \cup \dots \cup W_{ij}^{0,k_{ij}} \cup W_{ij}^1 \cup \dots \cup W_{ij}^{k'_{ij}}$$

such that each W_{ij}^u and $W_{ij}^{0,v}$ have $\text{disc}_2(\rho_2^{1/3}, 1/\ell')$, such that for each $1 \leq u \leq k'_{ij}$, there is $(\alpha, \beta) \in [\ell] \times [r]$ with $|W_{ij}^u \setminus E^{\alpha, \beta}| < (2\rho_2^{1/3}/\ell') |S_i| |S_j|$, and such that $W_{ij}^0 := \bigcup_{v=1}^{k_{ij}} W_{ij}^{0,v}$ has size at most $\rho_1 |S_i| |S_j|$. Since $|W_{ij}^0| = (1 \pm k_{ij} \rho_2^{1/3}) m^2 (k_{ij} / \ell')$ and $\rho_2 \ll \rho_1$, this implies $k_{ij} \leq 2\rho_1 \ell'$.

Now let $q = \min\{k'_{ij} : (S_i, S_j) \notin \Sigma_1\}$. Note that by above, $q \geq (1 - 2\rho_1) \ell'$. For each $(S_i, S_j) \notin \Sigma_1$, consider $K_2[S_i, S_j] \setminus (\bigcup_{u=1}^q W_{ij}^u)$. Let

$$\Gamma_{ij} := K_2[S_i, S_j] \setminus \left(\bigcup_{u=1}^q W_{ij}^u \right),$$

and note that by construction, Γ_{ij} is a union of $r_{ij} := k_{ij} + (k'_{ij} - q)$ many graphs, each with $\text{disc}_2(\rho_2^{1/3}, 1/\ell')$. If $r_{ij} = 0$, then let $E_{ij}^u = W_{ij}^u$ for each $u \in [q]$. Otherwise, apply

Lemma 3.24 to obtain a partition

$$\Gamma_{ij} = X_{ij}^1 \cup \dots \cup X_{ij}^q,$$

each with $\text{disc}_2(\rho_2)$ and density $\frac{r_{ij}}{\ell^q}(1 \pm \rho_2)$. Now for each $1 \leq u \leq q$, define $E_{ij}^u = W_{ij}^u \cup X_{ij}^u$. By Fact 3.23, each E_{ij}^u satisfies $\text{disc}_2(2\rho_2^{1/3}, 1/q)$. Note $K_2[S_i, S_j] = \bigcup_{u=1}^q E_{ij}^u$.

Now, for each $(S_i, S_j) \in \Sigma_1$, choose any partition $S_i \times S_j = \bigcup_{u=1}^q E_{ij}^u$ such that E_{ij}^u has $\text{disc}_2(\rho_2, 1/q)$ (such a partition exists by Lemma 3.24).

We now deal with the left over vertices, i.e. those in X^0 . Choose an equipartition $X^0 = X_1^0 \cup \dots \cup X_K^0$. Suppose first there is some X_i^0 with $|X_i^0| < \rho_2^{1/3}n/K$. Then for each $1 \leq i \leq K$, $|X_i^0| \leq \rho_2^{1/3}n/K$. In this case, set $S'_i = S_i \cup X_i^0$ for each $i \in [K]$, and for each $1 \leq i < j \leq K$, choose any partition $K_2[X_i^0, S'_j] \cup K_2[S'_i, X_j^0] = \bigcup_{\alpha=1}^q B_{ij}^\alpha$. Note that for each α , $|B_{ij}^\alpha| \leq \rho_2^{1/3}m(n/K)/q \leq \rho_1|S'_i||S'_j|/q$. For each $ij \in \binom{[K]}{2}$, set $R_{ij}^u = E_{ij}^u \cup B_{ij}^u$. Note that for each $S_i S_j \notin \Sigma_1$, and each $1 \leq u \leq q$, there is some $E^{\alpha, \beta}$ so that

$$|R_{ij}^u \setminus E^{\alpha, \beta}| \leq |B_{ij}^u| + 3\rho_1|E_{ij}^u| \leq 4\rho_1|R_{ij}^u|.$$

In fact, $R_{ij}^u \setminus E^{\alpha, \beta}$ has the form $Y_{ij}^u \cup (W_{ij}^{u'} \setminus E^{\alpha, \beta}) \cup B_{ij}^u$, for some $1 \leq u' \leq k'_{ij}$, and α, β . Thus since Y_{ij}^u has $\text{disc}_2(\rho_2^{1/3})$ and since $|(W_{ij}^{u'} \setminus E^{\alpha, \beta}) \cup B_{ij}^u| < \rho_2^{1/3}|W_{ij}^{u'}|$, it is easy to check that $R_{ij}^u \setminus E^{\alpha, \beta}$ has $\text{disc}_2(\rho_2^{1/6})$.

Suppose now that each X_i^0 has size at least $\rho_2^{1/6}n/K$. Let $b = \lceil \rho_2^{1/6}n/K \rceil$. For each $i \in [K]$, choose any partitions $X_i^0 = X_{i,0}^0 \cup X_{i,1}^0 \cup \dots \cup X_{i,a}^0$ and $S'_i = S_{i,0} \cup S_{i,1} \cup \dots \cup S_{i,c}$ such that for each $1 \leq j \leq a$ and $1 \leq j' \leq c$, $|X_{i,j}^0| = |S_{i,j'}| = b$, and $|X_{i,0}|, |S_{i,0}| < b$. Note $a, c \leq 2\rho_2^{-1/6}$.

Given $1 \leq i < j \leq K$, for each $1 \leq i' \leq c$ and $1 \leq j' \leq a$, use Lemma 3.24 with parameters q and $\delta = \epsilon = \rho_2$ to choose partitions $K_2[X_{i,i'}^0, S_{j,j'}] = D_{ii',jj'}^1 \cup \dots \cup D_{ii',jj'}^q$ so that for each $\alpha \leq q$, $(X_{i,i'}^0 \cup S_{j,j'}, D_{ii',jj'}^\alpha)$ satisfies $\text{disc}_2(\rho_2, 1/q)$. Note this means that each $|D_{ii',jj'}^\alpha| \leq \frac{ac}{q}(1 + \rho_2) < 2\rho_1|S'_i||S'_j|/q$. Then choose an arbitrary partition

$$K_2[X_{i,0}^0, S'_j] = \bigcup_{\alpha \leq \ell'} B_{ij}^\alpha.$$

Note that each $|B_{ij}^\alpha| < \rho_2^{1/6}|S'_i||S'_j|$. Now define, for each $1 \leq u \leq q$,

$$R_{ij}^u = E_{ij}^u \cup \left(\bigcup_{i'=1}^c \bigcup_{j'=1}^a D_{ii',jj'}^u \right) \cup B_{ij}^u.$$

By Fact 3.23, $R_{ij}^u \setminus B_{ij}^u$ has $\text{disc}_2(2\rho_2^{-1/6}\rho_2^{1/3}) = \text{disc}_2(\rho_2^{1/6})$, and size

$$|E_{ij}^u| + ac \frac{b^2}{q} (1 \pm \rho_2^{1/6}) = \frac{1}{q} (1 \pm 5\rho_2^{1/6}).$$

Thus since each B_{ij}^u is very small, we have that $|R_{ij}^u| = \frac{1}{q}(1 \pm 5\rho_2^{1/6})$, and R_{ij}^u has $\text{disc}_2(4\rho_2^{1/6}(q))$. Observe that by construction, for each u , there is $E^{\alpha,\beta}$ so that

$$|R_{ij}^u \setminus E^{\alpha,\beta}| \leq |E_{ij}^u \setminus E^{\alpha,\beta}| + ac \frac{b^2}{q} (1 \pm \rho_2^{1/6}) < 4\rho_1 |R_{ij}^u|.$$

Setting $\mathcal{R}_{\text{vert}} = \{S'_i : i \in [K]\}$ and $\mathcal{R}_{\text{edge}} = \{E_{ij}^u : ij \in \binom{[K]}{2}, u \in [q]\}$, we have that \mathcal{R} is an approximate $(\epsilon_1, \epsilon_2(q))$ -refinement of both \mathcal{P} and \mathcal{Q} . \square

We now prove Lemma 3.28.

PROOF OF LEMMA 3.28: Fix $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$. Choose $\epsilon'_1 \ll \epsilon_1$. Define $\rho, \epsilon'_2, \epsilon''_2 : \mathbb{N} \rightarrow (0, 1]$ by taking $\epsilon''_2(x) \ll \epsilon_1 \epsilon'_1 / x$, $\rho_2(x) \ll \min\{\epsilon_2(x), \epsilon'_2(x)\}$, and $\epsilon'_2(x) \ll \rho_2(x)/x$, for all $x \in \mathbb{N}$. Fix $t, \ell \geq 1$ and $t', \ell' \geq 1$, and choose $N \gg \ell, t(\epsilon'_1)^{-1} \epsilon'_2(\ell)^{-1}, \ell', t'$.

Suppose $H = (V, E)$ is a 3-graph with $|V| = n \geq N$ and \mathcal{Q} is a $(t, \ell, \epsilon'_1, \epsilon'_2(\ell))$ -decomposition of V which is ϵ'_1 -homogeneous with respect to H . Suppose \mathcal{P} is a $(t', \ell', \epsilon'_1, \epsilon'_2(\ell'))$ -decomposition of V which is a $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement of \mathcal{Q} . Say $\mathcal{Q}_{\text{vert}} = \{V_1, \dots, V_t\}$, and $\mathcal{Q}_{\text{edge}} = \{Q_{ij}^\alpha : ij \in \binom{[t]}{2}, \alpha \leq \ell\}$ and $\mathcal{P}_{\text{vert}} = \{W_1, \dots, W_{t'}\}$, and $\mathcal{P}_{\text{edge}} = \{P_{ij}^\alpha : ij \in \binom{[t']}{2}, \alpha \leq \ell'\}$. To ease notation, given $i, j, k \in [t]$ and $\alpha, \beta, \gamma \in [\ell]$, we let $Q_{ijk}^{\alpha,\beta,\gamma}$ denote the triad $(V_i \cup V_j \cup V_k, Q_{ij}^\alpha \cup Q_{ik}^\beta \cup Q_{jk}^\gamma)$. Similarly, given $i, j, k \in [t']$ and $\alpha, \beta, \gamma \in [\ell']$, we let $P_{ijk}^{\alpha,\beta,\gamma}$ denote the triad $(W_i \cup W_j \cup W_k, P_{ij}^\alpha \cup P_{ik}^\beta \cup P_{jk}^\gamma)$. Define

$$\begin{aligned} \Gamma_1 &= \{Q_{ijk}^{\alpha,\beta,\gamma} : Q_{ijk}^{\alpha,\beta,\gamma} \text{ is not } \text{disc}_{2,3}(\epsilon'_1, \epsilon'_2(\ell))\text{-regular and } \epsilon'_1\text{-homogeneous with respect to } H\}, \\ \Gamma_2 &= \{P_{ijk}^{\alpha,\beta,\gamma} : P_{ijk}^{\alpha,\beta,\gamma} \text{ is not } \text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell'))\text{-regular and } \epsilon_1\text{-homogeneous with respect to } H\}. \end{aligned}$$

Setting $\Omega_1 = \bigcup_{Q_{ijk}^{\alpha,\beta,\gamma} \in \Gamma_1} K_2^{(2)}(Q_{ijk}^{\alpha,\beta,\gamma})$, we have by assumption that $|\Omega_1| \leq \epsilon'_1 n^3$. Our goal is to show that $|\Omega_2| \leq \epsilon_1 n^3$, where $\Omega_2 := \bigcup_{P_{ijk}^{\alpha,\beta,\gamma} \in \Gamma_2} K_2^{(2)}(P_{ijk}^{\alpha,\beta,\gamma})$.

Let $\Sigma \subseteq \binom{\mathcal{P}_{\text{vert}}}{2}$ witness that \mathcal{Q} is an $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement of \mathcal{Q} . By the definition of an approximate refinement, for each $V_i V_j \notin \Sigma$, and each $P_{ij}^\alpha \in \mathcal{P}_{\text{edge}}$, there is some $f(P_{ij}^\alpha) \in \mathcal{Q}_{\text{edge}}$ such that $|P_{ij}^\alpha \setminus f(P_{ij}^\alpha)| < \epsilon'_1 |P_{ij}^\alpha|$ and such that $P_{ij}^\alpha \setminus f(P_{ij}^\alpha)$ has $\text{disc}_2(\epsilon'_2(\ell'))$. Similarly, for all $P_{ijk}^{\alpha,\beta,\gamma} \in \text{Triads}(\mathcal{P})$ with $V_i V_j, V_i V_k, V_j V_k \notin \Sigma$, we let $f(P_{ijk}^{\alpha,\beta,\gamma})$ denote the triad $Q_{uvw}^{\alpha',\beta',\gamma'}$ with the property that $f(P_{ij}^\alpha) = Q_{uv}^{\alpha'}$, $f(P_{ik}^\beta) = Q_{uv}^{\beta'}$, and $f(P_{jk}^\gamma) = Q_{vw}^{\gamma'}$. We now define

$$\Sigma' = \Sigma \cup \{P_{ij}^\alpha \in \mathcal{P}_{\text{edge}} : P_{ij}^\alpha \text{ fails } \text{disc}_2(\epsilon'_1, \epsilon'_2(\ell'))\}.$$

Note that we must have $|\Sigma'| \leq \epsilon'_1 (t')^2 + \sqrt{\epsilon'_1} (t')^2 \leq 2\sqrt{\epsilon'_1} (t')^2$, where the first inequality is by definition of a $(\epsilon'_1, \epsilon'_2(\ell'))$ -approximate refinement, and the second is because \mathcal{P} is a $(t', \ell, \epsilon'_1, \epsilon'_2(\ell'))$ -decomposition of V . Now let

$$E^* = \bigcup_{ij \notin \Sigma'} P_{ij}^\alpha \setminus f(P_{ij}^\alpha) \text{ and } V^* = \bigcup_{W_i \in \mathcal{P}_{\text{vert}}} W_i \setminus V_i.$$

By assumption, $|E^*| \leq \epsilon'_2(\ell')|V|^2$ and $|V^*| \leq \epsilon'_1|V|$. Define

$$\Sigma_{\mathcal{Q}} = \{Q_{ij}^\alpha \in \mathcal{Q}_{edge} : |Q_{ij}^\alpha \cap E^*| \geq \sqrt{\epsilon'_1}|Q_{ij}^\alpha| \text{ or } \min\{|V_i \cap V^*|, |V_j \cap V^*|\} \geq \sqrt{\epsilon'_1}|V_j|\}.$$

Clearly, the bounds on E^* and V^* implies $|\Sigma_{\mathcal{Q}}| \leq 3\sqrt{\epsilon'_1}\ell t^2$. Now set

$$\Gamma'_1 = \{Q_{ijs}^{\alpha,\beta,\gamma} : Q_{ij}^\alpha, Q_{is}^\beta, \text{ or } Q_{js}^\gamma \in \Sigma_{\mathcal{Q}}\}.$$

For each $ij \in \binom{[t]}{2}$, set $I_{ij}^\alpha = \{P_{uv}^\beta \in \mathcal{P}_{edge} : f(P_{uv}^\beta) = Q_{ij}^\alpha\}$, and define, for each $ijs \in \binom{[t]}{3}$ and $\alpha, \beta, \gamma \leq \ell$,

$$\mathcal{I}_{ijs}^{\alpha,\beta,\gamma} = \{P_{uvw}^{\rho,\gamma,\tau} \in \text{Triads}(\mathcal{P}) : P_{uv}^\rho \in I_{ij}^\alpha, P_{uw}^\mu \in I_{is}^\beta, P_{vw}^\tau \in I_{js}^\gamma\}.$$

Suppose $Q_{ijs}^{\alpha,\beta,\gamma} \notin \Gamma_1 \cup \Gamma'_1$. We claim that almost all triples contained in $K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})$ are in an ϵ_1 -homogeneous triad of \mathcal{P} . First, we observe that by definition of Γ_1 , there is a $\delta = \delta(ijs, \alpha\beta\gamma) \in \{0, 1\}$ so that

$$|E^\delta \cap K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})| \geq (1 - \epsilon'_1)|K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})|.$$

Next, note that

$$\bigcup_{P_{uvw}^{\rho,\gamma,\tau} \in \mathcal{I}_{ijs}^{\alpha,\beta,\gamma}} K_3^{(2)}(P_{uvw}^{\rho,\gamma,\tau} \cap Q_{ijs}^{\alpha,\beta,\gamma}) \subseteq K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma}).$$

This implies that

$$|\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}| \geq (1 - 3\sqrt{\epsilon'_1}) \frac{n^3}{t^3 \ell^3} / (1 - \epsilon_2(\ell)) \frac{n^3}{(t')^3 (\ell')^3} \geq (1 - 4\sqrt{\epsilon'_1}) (t')^3 (\ell')^3 / t^3 \ell^3.$$

For all $P_{uvw}^{\rho,\gamma,\tau} \in \mathcal{I}_{ijs}^{\alpha,\beta,\gamma}$, each of $P_{uv}^\rho \setminus f(P_{uv}^\rho)$, $P_{uw}^\mu \setminus f(P_{uw}^\mu)$ and $P_{vw}^\tau \setminus f(P_{vw}^\tau)$ satisfy $\text{disc}_2(\epsilon'_2(\ell))$ and have size at most $\epsilon'_1 \frac{n^2}{(t')^2}$. Combining this with the observation above and Corollary 3.6, we have

$$(1 - 3\epsilon'_1)^3 \frac{n^3}{(t')^3 (\ell')^3} |\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}| \leq |K_2^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})| \leq (1 + \epsilon'_1) \frac{n^3}{\ell^3 t^3}.$$

Consequently, using that ϵ'_1 is small, we have that $|\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}| \leq \frac{(t')^3 (\ell')^3}{t^3 \ell^3} (1 - \sqrt{\epsilon'_1})$. On the other hand, since $Q_{ijs}^{\alpha,\beta,\gamma} \notin \Gamma'_1$,

$$|K_2^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma}) \setminus \bigcup_{P_{uvw}^{\rho,\gamma,\tau} \in \mathcal{I}_{ijs}^{\alpha,\beta,\gamma}} K_3^{(2)}(P_{uvw}^{\rho,\gamma,\tau} \cap Q_{ijs}^{\alpha,\beta,\gamma})| \leq 3\sqrt{\epsilon'_1} \frac{n^3}{t^3 \ell^3}.$$

Using the inequalities above and Corollary 3.6, we have

$$\begin{aligned}
\sum_{\text{Triads}(I_{ijs}^{\alpha,\beta,\gamma})} |E^\delta \cap K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau})| &\geq |E^\delta \cap \bigcup_{P_{uvw}^{\rho,\mu,\tau} \in \mathcal{I}_{ijs}^{\alpha,\beta,\gamma}} K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau} \cap Q_{ijs}^{\alpha,\beta,\gamma})| \\
&\geq (1 - 3\sqrt{\epsilon'_1}) |E^\delta \cap K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})| \\
&\geq (1 - 3\sqrt{\epsilon'_1})(1 - \epsilon'_1) |K_3^{(2)}(Q_{ijs}^{\alpha,\beta,\gamma})| \\
&\geq (1 - 3\sqrt{\epsilon'_1})(1 - \epsilon'_1)^2 \frac{n^3}{t^3 \ell^3}.
\end{aligned}$$

This implies

$$\sum_{\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}} \frac{|E^\delta \cap K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau})|}{|K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau})|} \geq \frac{t^3 \ell^3}{(t')^3 (\ell')^3} (1 - 4(\epsilon'_1)^{1/2}) \geq (1 - (\epsilon'_1)^{1/4}) |\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}|.$$

Since each term in the summand above is in $[0, 1]$, and $\epsilon'_1 \ll \epsilon_1$, this implies that there is a set $\mathcal{J}_{ijs}^{\alpha,\beta,\gamma} \subseteq \mathcal{I}_{ijs}^{\alpha,\beta,\gamma}$ such that $|\mathcal{J}_{ijs}^{\alpha,\beta,\gamma}| \geq (1 - \epsilon_1^2) |\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}|$, and for all $P_{uvw}^{\rho,\mu,\tau} \in \mathcal{J}_{ijs}^{\alpha,\beta,\gamma}$

$$\frac{|E^\delta \cap K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau})|}{|K_3^{(2)}(P_{uvw}^{\rho,\mu,\tau})|} \geq (1 - \epsilon_1^2).$$

Thus, the number of triples from $Q_{ijs}^{\alpha,\beta,\gamma}$ which are in an ϵ_1^2 -homogeneous triad of \mathcal{P} is at least

$$\begin{aligned}
|\mathcal{J}_{ijs}^{\alpha,\beta,\gamma}| (1 - \epsilon_1^2) \frac{n^3}{(t')^3 (\ell')^3} &\geq (1 - \epsilon_1^2) |\mathcal{I}_{ijs}^{\alpha,\beta,\gamma}| (1 - \epsilon_1^2) \frac{n^3}{(t')^3 (\ell')^3} \\
&\geq (1 - \epsilon_1^2) (1 - 4\sqrt{\epsilon'_1}) ((t')^3 (\ell')^3 / t^3 \ell^3) (1 - \epsilon_1^2) \frac{n^3}{(t')^3 (\ell')^3} \\
&= (1 - \epsilon_1^2)^2 (1 - 4\sqrt{\epsilon'_1})^2 \frac{n^3}{t^3 \ell^3}.
\end{aligned}$$

We know have that

$$\begin{aligned}
|\Omega_2| &\geq |\text{Triads}(\mathcal{Q}) \setminus (\Gamma_1 \cup \Gamma'_1)| (1 - \epsilon_1^2)^2 (1 - 4\sqrt{\epsilon'_1})^2 \frac{n^3}{t^3 \ell^3} \\
&\geq (1 - 10\sqrt{\epsilon'_1}) t^3 \ell^3 (1 - \epsilon_1^2)^2 (1 - 4\sqrt{\epsilon'_1})^2 \frac{n^3}{t^3 \ell^3} \\
&\geq (1 - \epsilon_1) n^3.
\end{aligned}$$

This finishes the proof. \square

Finally, we prove Theorem 4.1.

PROOF OF THEOREM 4.1: Given $\xi, d_3 > 0$ and $t \geq 1$, let $\xi_0 > 0$ be sufficiently small so that $(1 + \xi_0)^{t^3+t^1} \leq 1 + \xi$. Choose $\delta_3 = \delta_3(t, \xi_0, d_3/2)$ as in Theorem 3.13. Fix $d_2 > 0$, and choose $\delta_2 = \delta_2(t, \xi_0, d_3/2, \delta_3, d_2)$ and $n_0(t, \xi_0, d_3/2, \delta_3, d_2)$ as in Theorem 3.13. Let m_0 be as in Fact 3.4 for δ_2 and d_2 , and let $\delta'_2 = d_2^8 \delta_2 / 4$.

Let $\delta'_3 = \delta'_3(\min\{\delta_3, d_3\xi_0\}, 2)$, $\delta''_2 = \delta''_2(\delta'_2, \min\{\delta_3, d_3\xi_0\}, d_3, 2)$, and $M_0 = M_0(\delta'_2, \min\{\delta_3, d_3\xi_0\}, d_3, 2)$ be from Lemma 3.19. Now set $N = tm_0n_0M_0(\delta''_2)^{-4}$.

Fix $t_a, t_b, t_c \geq 1$ such that $t_a + t_b + t_c = t$, and $F = (U \cup W \cup Z, R_F)$ a 3-partite 3-graph with $U = \{u_1, \dots, u_{t_a}\}$, $W = \{w_1, \dots, w_{t_b}\}$, and $Z = \{z_1, \dots, z_{t_c}\}$. Let V be a set of size $n \geq N$, equipped with a partition

$$V = A_1 \cup \dots \cup A_{t_a} \cup B_1 \cup \dots \cup B_{t_b} \cup C_1 \cup \dots \cup C_{t_c},$$

where each part has size $\frac{n}{t}(1 \pm \delta''_2)$. Let $A = \bigcup_{i=1}^{t_a} A_i$, $B = \bigcup_{i=1}^{t_b} B_i$, and $C = \bigcup_{i=1}^{t_c} C_i$. Suppose $H = (V, R)$ is a 3-graph and $G = (V, E)$ is a t -partite graph with the vertex partition displayed above. Assume H and G satisfy the hypotheses of Theorem 4.1 with $\delta''_2, d_2, d_3, \delta'_3$.

Let $H' = (V, E \setminus ((\binom{A}{2} \cup \binom{B}{2} \cup \binom{C}{2}))$. We define a new graph G' on V as follows. For each $X \in \{A, B, C\}$ and $ij \in \binom{[t_x]}{2}$, define $G'[X_i, X_j]$ to be any bipartite graph on $X_i \cup X_j$ with $\text{disc}_2(\delta'_2, d_2)$ (such a graph exists by Fact 3.4). For $X \neq Y \in \{A, B, C\}$, $i \in [t_x]$ and $j \in [t_y]$, define $G'[X_i, Y_j] = G[X_i, Y_j]$. For each $XYZ \in \{A, B, C\}^3$ and $(i, j, s) \in [t_x] \times [t_y] \times [t_z]$, set $G_{XYZ, ijs} := G'[X_i, Y_j, Z_s]$ and $H_{XYZ, ijs} = (X_i \cup Y_j \cup Z_s, R \cap K_3^{(2)}(G_{XYZ, ijs}))$, and let $d_{XYZ, ijs}$ be such that $|R \cap K_3^{(2)}(G_{XYZ, ijs})| = d_{XYZ, ijs}|K_3^{(2)}(G_{XYZ, ijs})|$. If $XYZ = ABC$, then by assumption, $(H_{XYZ, ijs}, G_{XYZ, ijs})$ satisfies $\text{disc}_{2,3}(\epsilon_1, \epsilon_2(\ell))$. If $XYZ \neq ABC$, then by definition $(H_{XYZ, ijs}, G_{XYZ, ijs})$ has density 0, so by Proposition 3.22, it satisfies $\text{disc}_{2,3}(\delta'_3, \delta''_2)$. Set

$$d = \left(\prod_{u_i v_j w_s \in R_F} d_{ABC, ijs} \right) \left(\prod_{u_i v_j w_s \in R_F} (1 - d_{ABC, ijs}) \right).$$

Let $m = \min\{|A_i|, |B_j|, |C_k| : i \in [t_a], j \in [t_b], k \in [t_c]\}$, and for each $(i, j, k) \in [t_a] \times [t_b] \times [t_c]$, choose any m -element subsets A'_i , B'_j , and C'_k of A_i , B_j , and C_k respectively. Note $m \geq (1 - \delta'_2)\frac{n}{t} \geq \max\{n_0, m_0\}$. Define $V' = \bigcup_{i=1}^{t_a} A'_i \cup \bigcup_{i=1}^{t_b} B'_i \cup \bigcup_{i=1}^{t_c} C'_i$, and set $G'' = G'[V']$ and $H'' = H'[V']$.

Then for each $XYZ \in \{A, B, C\}^3$ and $(i, j, s) \in [t_x] \times [t_y] \times [t_z]$, let $G'_{XYZ, ijs} = G_{XYZ, ijs}[V']$ and $H'_{XYZ, ijs} = H_{XYZ, ijs}[V']$. By Lemma 3.19, each $(G'_{XYZ, ijs}, H'_{XYZ, ijs})$ has $\text{disc}_{2,3}(\delta_3, \delta'_2)$ with density $d_{XYZ, ijs}(1 \pm \xi_0)$.

By Theorem 3.13, there is a set $\Sigma \subseteq \prod_{i=1}^{t_1} A'_i \times \prod_{i=1}^{t_2} B'_i \times \prod_{i=1}^{t_3} C'_i$ such that for all $\bar{a}\bar{b}\bar{c} \in \Sigma$, $x_i y_j z_s \in \binom{\bar{a}\bar{b}\bar{c}}{3}$ if and only if $x_i y_j z_s \in E(H')$, and such that

$$|\Sigma| = d_2^{\binom{t}{2}} d' (1 \pm \xi_0) m^t,$$

where $d' = (\prod_{(u_i, v_j, w_s) \in R} d'_{ABC, ijs}) (\prod_{(u_i, v_j, w_s) \notin R} (1 - d'_{ABC, ijs}))$. Recall that each $d'_{ABC, ijs} = d_{ABC, ijs}(1 \pm \xi_0)$. Consequently, $d' = d(1 \pm \xi_0)^{t^3}$. Thus

$$\begin{aligned} |\Sigma| &= d_2^{\binom{t}{2}} d(1 \pm \xi_0) m^t = d_2^{\binom{t}{2}} d(1 \pm \xi_0)^{t^3+t} \prod_{i=1}^{t_1} |A_i| \prod_{i=1}^{t_2} |B_i| \prod_{i=1}^{t_3} |C_i| \\ &= d_2^{\binom{t}{2}} d(1 \pm \xi) \prod_{i=1}^{t_1} |A_i| \prod_{i=1}^{t_2} |B_i| \prod_{i=1}^{t_3} |C_i|. \end{aligned}$$

By construction, for all $\bar{a}\bar{b}\bar{c} \in \Sigma$, $a_i b_j c_s \in R$ if and only if $u_i w_j z_s \in R_F$, so this finishes the proof. \square

APPENDIX C. RAMSEY FACTS

For convenience, we will use model theoretic language in this section along with the compactness theorem. This is essentially only to cut down on the notation required, and all the proofs can be easily finitized by replacing infinite sets with sufficiently large finite ones.

Let $\mathcal{L} = \{R(x, y, z)\}$ consist of a single ternary relation symbol. Recall that for any hereditary 3-graph property, there is a universal \mathcal{L} -theory $T_{\mathcal{H}}$ such that \mathcal{H} is the class of finite models of $T_{\mathcal{H}}$.

PROOF OF FACT 3.30: Suppose \mathcal{H} has IP_2 . By compactness, there is an infinite 3-graph $\mathcal{M} = (V, E) \models T_{\mathcal{H}}$ with sets $A = \{a_i : i \in \mathbb{N}\} \subseteq V$, $B = \{b_i : i \in \mathbb{N}\} \subseteq V$, and $C = \{c_S : S \subseteq A \times B, S \text{ finite}\} \subseteq V$ so that $a_i, b_j, c_S \in E$ if and only if $(i, j) \in S$. Clearly we may choose $A' \subseteq A$ and $B' \subseteq B$ such that each of A' and B' are infinite and $A' \cap B' = \emptyset$. Suppose $H = (V_1 \cup V_2 \cup V_3, E)$ is a finite 3-partite 3-graph.

Let $f : V_1 \rightarrow A'$, $g : V_2 \rightarrow B'$ be any injections. By construction $f(V_1) \cap g(V_2) = \emptyset$. For each $w \in V_3$, set $S_w = \{c_S \in C : S \cap (f(V_1) \times g(V_2)) = \{f(u)f(v) : uvw \in E\}\}$. There are infinitely many distinct $S \subseteq A \times B$ with the property that $S \cap (f(V_1) \times g(V_2)) = \{f(u)f(v) : uvw \in E\}$, and thus S_w is infinite for all $w \in V_3$. Thus, for all $w \in V_3$, we have that $S'_w = S_w \setminus (f(V_1) \cup g(V_2))$ is also infinite. Since V_3 is finite, we can clearly choose elements $h(w) \in S'_w$ so that $h(V_3) = \{h(w) : w \in V_3\}$ has size V_3 . By construction, $H' := \mathcal{M}[f(V_1) \cup g(V_2) \cup h(V_3)]$ is a clean copy of H . Since $\mathcal{M} \models T_{\mathcal{H}}$ and universal sentences are closed under substructures, $H' \models T_{\mathcal{H}}$, and consequently, $H' \in \mathcal{H}$ is as desired. \square

PROOF OF LEMMA 3.31: We begin with (1). Suppose $H^*(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. By compactness there is a 3-graph $\mathcal{M} = (V, E) \models T_{\mathcal{H}}$, a vertex $c \in M$, and subsets $A = \{a_i : i \in \mathbb{N}\}$, $B = \{b_i : i \in \mathbb{N}\} \subseteq V$ such that $ca_i b_j \in E$ if and only if $i \leq j$. It is easy to see that all elements in A and B must be pairwise distinct, respectively. Given $a_i \in A$, since \mathcal{M} is a 3-graph, for any $j \geq i$, $ca_i b_j \in E$ implies $a_i \neq b_j$. Thus, there can be at most one element in B equal to a_i , and it would have to be in the set $\{b_1, \dots, b_i\}$. Similarly, since c is in an edge with every element of A and B , it is distinct from all of $A \cup B$.

Fix $k \geq 1$. We define sequences (m_k, \dots, m_1) and (I_k, \dots, I_1) as follows.

Start with $m_k = k^3$, and $I_k = \{j < k^3 : b_j \neq a_{k^3}\}$.

Suppose now that $0 \leq i \leq k-1$, and assume we have defined m_k, \dots, m_{k-i} and I_k, \dots, I_{k-i} . Let $m_{k-i-1} = \min I_{k-i}$ and let $I_{k-i-1} = \{j < m_{k-i-1} : b_j \neq a_{k-i-1}\}$. At the end we will have that $ca_{m_u}b_{m_v}$ holds if and only if $u \leq v$, and $X = \{c\} \cup \{a_{m_u}, b_{m_u} : u \in [k]\}$ has size $2k+1$. Thus $\mathcal{M}[X]$ is a clean copy of $H^*(k)$. As above, $\mathcal{M}[X] \in \mathcal{H}$, so we are done.

For part (2), suppose $\overline{H}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. By compactness there is a 3-graph $\mathcal{M} = (V, E) \models T_{\mathcal{H}}$ with subsets $C = \{c_i : i \in \mathbb{N}\}$, $A = \{a_i : i \in \mathbb{N}\}$, and $B = \{b_i : i \in \mathbb{N}\} \subseteq M$ such that $c_s a_i b_j \in E$ if and only if $i \leq j$, and $c_i \neq c_j$ for each $i \neq j$. It is easy to see that all elements in A and B must be pairwise distinct, respectively. Given $a_i \in A$, since \mathcal{M} is a 3-graph, for any $j \geq i$, $c_1 a_i b_j \in E$ implies $a_i \neq b_j$. Thus, there can be at most one element in B equal to a_i , and it would have to be in the set $\{b_1, \dots, b_i\}$. Similarly, since each c_s is in an edge with every element of A and also some edge with every element of B , we automatically know $C \cap A = C \cap B = \emptyset$.

Fix $k \geq 1$. Construct a sequence m_k, \dots, m_1 and I_k, \dots, I_1 exactly as in part (1). Then for all $s \in [k]$, $c_s a_{m_u} b_{m_v}$ holds if and only if $u \leq v$, and $X = \{c_1, \dots, c_k\} \cup \{a_{m_u}, b_{m_u} : u \in [k]\}$ has size $3k$. Thus $\mathcal{M}[X]$ is a clean copy of $\overline{H}(k)$. As above, $\mathcal{M}[X] \in \mathcal{H}$, so we are done.

We now do part (3). Suppose $F(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. By compactness there is a 3-graph $\mathcal{M} = (V, E) \models T_{\mathcal{H}}$ with subsets $C = \{c_i : i \in \mathbb{N}\}$, $A = \{a_i : i \in \mathbb{N}\}$, and $B = \{b_i^f : f : \mathbb{N}^2 \rightarrow \mathbb{N}, i \in \mathbb{N}\} \subseteq M$ such that $a_i b_j^f c_s \in E$ if and only if $s \leq f(i, j)$. It is easy to see this implies that the elements of C and A are pairwise distinct, respectively. Further, since every a_i is in an edge with every c_s (if g is the constant function $g(x, y) = s$, then $a_i b_1^g c_s \in E$), we must have $A \cap C = \emptyset$.

Fix $k \geq 1$ and let $N \gg k$. Let I be the set of functions from $[k]^2 \rightarrow [k]$. Fix any enumeration $I = \{f_u : u \in k^{2k}\}$. For each $f_u \neq f_v \in I$, let $(\alpha_{uv}, \beta_{uv})$ be the least element of $[k]^2$ (in the lexicographic order) such that $f_u(\alpha_{uv}, \beta_{uv}) \neq f_v(\alpha_{uv}, \beta_{uv})$. Given $f_u : [k]^2 \rightarrow [k]$, and $w \in [k^2]$ define $g_{f_u}^w : [N^2] \rightarrow [N]$ as follows. For all $i, j \in [k]$, set $g_{f_u}^w(i, j) = f_u(i, j)$, and for each $u \neq v \in k^{2k}$, and $j \in [k]$, define $g_{f_u}^w(k + 3^u 4^v, j) = g_{f_u}^w(k + 3^v 4^u, j) = f_u(\alpha_{uv}, \beta_{uv})$, and for each $j \neq j' \in [k]$, let $g_{f_u}^w(k + 5^j 7^{j'}, j) = 1$ and $g_{f_u}^w(k + 5^j 7^{j'}, j') = 2$, and then for each $w < w' \leq N$, define $g_{f_u}^w(k + 11^w 13^{w'}, j) = w$ and $g_{f_u}^{w'}(k + 11^w 13^{w'}, j) = w'$. The point of all this is that $|\{e_i^{g_{f_u}^w} : w \in [N], f_u \in I, i \in [k]\}| = |I|k^2 N$. For each $f_u \in I$, and each $j \in [j]$, let $X_{u,j} = \{e_j^{g_{f_u}^w} : w \in [N]\}$. Then $X_{u,j}$ has size N , so it contains a subset $X'_{u,j}$ of size at least $N - 2k$ which is disjoint from $\{c_i, a'_i : i \in [k]\}$.

Choose $(b_1^{f_1}, \dots, b_k^{f_k}) \in X_{1,1} \times \dots \times X_{1,k}$ so that the coordinates are pairwise distinct. This is possible by the size of each $X_{1,j}$.

At step $u+1$, choose $(b_1^{f_{u+1}}, \dots, b_k^{f_{u+1}}) \in X_{u+1,1} \times \dots \times X_{u+1,k}$ which are pairwise distinct, and also distinct from all b_j^v for all $1 \leq v \leq u$ and $j \in [k]$, which is possible by how large the $X_{u+1,j}$ are.

In the end, we have a set $V'' = \{c_i, a'_i, b_i^{f_u} : i \in [k], f_u \in I\}$ such that $a_i b_j^{f_u} c_s \in E$ if and only if $s \leq f_u(i, j)$, and such that the elements of V'' are all pairwise distinct. In other words, $\mathcal{M}[V'']$ is a clean copy of $F(k)$. As before, this tells us $\mathcal{M}[V''] \in \mathcal{H}$, as desired. \square

PROOF OF LEMMA 3.33: Suppose $\overline{U}(k) \in \text{Trip}(\mathcal{H})$ for all $k \geq 1$. By compactness there is a 3-graph $\mathcal{M} = (V, E) \models T_{\mathcal{H}}$ with subsets $C = \{c_i : i \in \mathbb{N}\}$, $A = \{a_i : i \in \mathbb{N}\}$, and $B = \{b_S : S \subseteq \mathbb{N}\}$ such that $c_j a_i b_S \in E$ if and only if $i \in S$. It is easy to see that all elements in A and B must be pairwise distinct, respectively, and the c_i are pairwise distinct by assumption. Given a_i, c_j , and b_S , with $i \in S$, since $c_j a_i b_S \in E$, we know $a_i \neq c_j$, $b_S \neq c_j$. Thus, $(B \setminus \{b_{\emptyset}\}) \cap A = (B \setminus \{b_{\emptyset}\}) \cap C = \emptyset$.

Fix a finite bipartite graph $G = (U \cup V, E')$ and $n \geq 1$. Choose $\{A_v : v \in V\}$ so that each $A_v \subseteq A$ has size $|V|^n$ and so that $v \neq v'$ implies $A_v \cap A_{v'} = \emptyset$. For each $u \in U$, let

$$B_u = \left\{ b_S : \bigcup_{v \in N_G(u)} A_v \subseteq S \text{ and } S \cap \left(\bigcup_{v \in V \setminus N_G(u)} A_v \right) = \emptyset \right\}.$$

Note that B_u is infinite for every $u \in U$. Thus, we may clearly choose a tuple $(b_u)_{u \in U}$ and $(x_v)_{v \in V}$ so that $x_v \in A_v$ for each $v \in V$, $b_u \in B_u$ for each $u \in U$, and so that all the elements of $\{b_u : u \in U\} \cup \{x_v : v \in V\}$ are pairwise distinct. Now choose any c_1, \dots, c_n also disjoint from this set (this is possible as C is infinite). We now have that $c_j b_u x_v \in E$ if and only if $v \in N_G(u)$, i.e. if and only if $uv \in E'$. It is now clear that $\mathcal{M}[\{b_u : u \in U\} \cup \{x_v : v \in V\} \cup \{c_1, \dots, c_n\}] \in \mathcal{H}$ is a clean copy of $n \otimes G$, as desired. \square

APPENDIX D. GENERAL PROPERTIES OF NFOP₂-FORMULAS

This appendix contains some general properties of NFOP₂ formulas. We will use model theoretic notation. First, we observe that formally, all three variables play a distinct role in the definition of FOP₂. For this reason, when we give the definition of FOP₂ in general, we must partition the variables into three parts (rather than the more typical two).

Definition D.1. Suppose \mathcal{L} is a first-order language, \mathcal{M} is an \mathcal{L} -structure, and $\phi(x, y, z)$ is an \mathcal{L} -formula. We say the tripartitioned formula $\phi(x; y; z)$ has k -FOP₂ in \mathcal{M} if there are $b_1, \dots, b_k, c_1, \dots, c_k \in M$, and for each $f : [k]^2 \rightarrow [k]$, $a_1^f, \dots, a_k^f \in M$ such that $\mathcal{M} \models \phi(a_i^f, b_j, c_k)$ if and only if $k \leq f(i, j)$.

It is easy to see that if a formula $\phi(x, y, z)$ has VC₂-dimension at least ℓ , then it has ℓ -FOP₂ (see e.g. Fact 2.22). Let \mathcal{M} be the Fraïssé limit of all 3-graphs. It is well known that $Th(\mathcal{M})$ is simple (this can be shown via an argument similar to Corollary 7.3.14 in [56]). Since the edge relation in this theory has unbounded VC₂-dimension, this shows that simple $\not\models$ NFOP₂. On the other hand, using the machinery of this paper (namely Theorem 2.50 and Section 7.1), it is possible to show that any quantifier-free formula $\phi(x; y; z)$ in the graph language is NFOP₂ in any graph. This implies, for example, that any quantifier-free formula in three variables in the Henson graph is NFOP₂. Consequently, NFOP₂ $\not\models$ simple. We also show in [60] that if a formula has ℓ -FOP₂ then it has VC-dimension at least ℓ .

In our next proposition, we give an equivalent formulation of ℓ -FOP₂, and also show that NFOP₂ is closed under negations.

Proposition D.2. Suppose \mathcal{L} is a first-order language, \mathcal{M} is an \mathcal{L} -structure, and $\phi(x, y, z)$ is an \mathcal{L} -formula.

- (i) If $\phi(x_1; x_2; x_3)$ has ℓ -FOP₂ in \mathcal{M} , then $\neg\phi(x_1; x_2; x_3)$ has $(\ell - 1)$ -FOP₂.

(ii) *The following are equivalent.*

(a) $\phi(x_1; x_2; x_3)$ has ℓ -FOP₂ for all ℓ .

(b) For all ℓ , there are $b_1, \dots, b_\ell, c_1, \dots, c_\ell$ and b_f for $f \in [\ell]^{[\ell]}$ such that $\phi(a^f, b_i, c_k)$ holds if and only if $k \leq f(i)$.

Proof. We begin with (i). Let $y_1, \dots, y_\ell, z_1, \dots, z_\ell$ and $\{x_i^f : i \in [\ell], f : [\ell]^2 \rightarrow [\ell]\}$ be such that $\phi(x_i^f, y_j, z_k)$ holds in \mathcal{M} if and only if $k \leq f(i, j)$. For each $f : [\ell - 1]^2 \rightarrow [\ell - 1]$, let $g_f : [\ell]^2 \rightarrow [\ell]$ be the function defined by $g_f(x, y) = \ell - f(x, y) + 1$ for $x, y \in [\ell - 1]$, and $g_f(x, y) = 1$ otherwise. For each $i \in [\ell - 1]$, let $w_i = z_{\ell-i+1}$, $v_i = x_i$, and for each $f : [\ell - 1]^2 \rightarrow [\ell - 1]$, let $u_i^f = x_i^{g_f}$. Then for each $i, j, k \in [\ell]$, $\phi(u_i^f, v_j, w_k) = \phi(x_i^{g_f}, y_j, z_{\ell-k+2})$ holds if and only if $\ell - k + 2 < g_f(i, j)$, if and only if $\ell - k + 1 < \ell - f(i, j) + 1$, if and only if $f(i, j) < k$. Thus $\neg\phi(u_i^f, v_j, w_k)$ holds if and only if $k \leq f(i, j)$. This shows $\neg\phi(x; y; z)$ has $(\ell - 1)$ -FOP₂.

We now prove (ii). We begin with the direction (a) \Rightarrow (b). Enumerate $[\ell]^{[\ell]} = \{f_1, \dots, f_m\}$. Define $f : [m]^2 \rightarrow [m]$ such that $f(i, j) = f_i(j)$ if $j \leq \ell$ and $f(i, j) = \ell$ if $j > \ell$. Choose $a_1^f, \dots, a_m^f, b_1, \dots, b_m, c_1, \dots, c_m$ so that $\phi(a_i^f, b_j, c_k)$ holds if and only if $k \leq f(i, j)$, i.e. if and only if $k \leq f_i(j)$. For each $t \in [m]$, set $b_{f_t} = b_t^f$. Then for all $j, k \leq \ell$, we have $\phi(a_{f_t}, b_j, c_k)$ if and only if $k \leq f_t(i)$.

We now show (b) \Rightarrow (a). Fix $\ell \geq 1$. Let $b_1, \dots, b_\ell, c_1, \dots, c_\ell$ and a_f for $f \in [\ell]^{[\ell]}$ be as in (b). Fix $f : [\ell]^2 \rightarrow [\ell]$ and for each $j \leq \ell$, define $f_j : [\ell] \rightarrow [\ell]$ so that $f_j(i) = f(i, j)$. For each $i \in [\ell]$, set $a_i^f = a_{f_i}$. Then $\phi(a_i^f, b_j, c_k)$ holds if and only if $k \leq f_j(i)$, i.e., $k \leq f(i, j)$. \square

Proposition D.2 implies that being NFOP₂ is closed under negations. We now show that it is also closed under conjunctions.

Theorem D.3. *For all $k \geq 1$ there is $t \geq 1$ so that the following holds. Suppose \mathcal{L} is a first-order language, \mathcal{M} is an \mathcal{L} -structure, and $\phi_1(x, y, z), \phi_2(x, y, z)$ are \mathcal{L} -formulas. Suppose $\phi_1(x; y; z)$ and $\phi_2(x; y; z)$ have no k -FOP₂ in \mathcal{M} . Then $(\phi_1 \wedge \phi_2)(x; y; z)$ has no t -FOP₂.*

Proof. Suppose towards a contradiction that there exists $k \geq 1$ so that for all $t \geq 1$, there exists an \mathcal{L} -structure $\mathcal{M}(t)$ in which $\phi_1(x; y; z)$ and $\phi_2(x; y; z)$ have no k -FOP₂, but in which $(\psi_1 \wedge \psi_2)(x; z; y)$ has t -FOP₂, where $\psi_1(x, z, y) := \phi_1(x, y, z)$ and $\psi_2(x, z, y) := \phi_2(x, y, z)$.

By compactness, there exists an \mathcal{L} -structure \mathcal{M} in which $\phi_1(x; y; z)$ and $\phi_2(x; y; z)$ have no k -FOP₂, but where $\psi_1 \wedge \psi_2(x; z; y)$ has m -FOP₂ for all $m \geq 1$.

Let $s = s(k)$ be such that the conjunction of two k -stable formulas is s -stable. Let ϵ_1, ϵ_2 be as in Proposition 6.7 for k and choose $S \gg \epsilon_1^{-1}s$. Choose $\epsilon_1'' \ll \epsilon_1' \ll \epsilon_1/S$, and define $\epsilon_2'', \epsilon_2' : \mathbb{N} \rightarrow (0, 1]$ by defining, for each x , $\epsilon_2''(x) \ll \epsilon_2'(x) \ll \epsilon_2(x)/kx$.

Let $\mu_1', \mu_1, \mu_2, \mu_2'$ be as in Lemma 3.28 for ϵ_1', ϵ_2' . Choose $f, g : \mathbb{N}^4 \rightarrow (0, 1]$ as in Lemma 3.27 for μ_1'', μ_2'' , where $\mu_1'' = \min\{\mu_1, \mu_1'\}$ and $\mu_2''(x) = \min\{\mu_2(x), \mu_2'(x)\}$. Define $\epsilon_2''''(x) \ll \epsilon_2'''(x) \ll \mu_2''(x)\epsilon_2''(x)$ for all x .

Let T_1, L_1, N_1 be as in Proposition 2.49 for k, μ_1'' , and μ_2'' , and set $T = f_1(T_1, L_1, T_1, L_1)$ and $L = g_1(T_1, L_1, T_1, L_1)$. Choose $S \gg L$ and $N \gg SLT\mu_2''(LT)^{-1}(\mu_1'')^{-1}N_1N_2$. Define $n = n(k, N)$ from Lemma 6.8.

Now let $U = \{u_1, \dots, u_N\}$, $W = \{w_1, \dots, w_N\}$ be two disjoint sets, each of size N , and let $\bigcup_{\alpha \leq S} P_{UW}^\alpha$ be a partition of $K_2[U, W]$ such that each P_{UW}^α has $\text{disc}_2(\epsilon_2'''(LT), 1/S)$ (such a partition exists by Lemma 3.24).

By assumption and Lemma 3.31, there are disjoint sets $A, B, C \subseteq M$, with

$$A = \{a_i^f : i \in [n], f : [n]^2 \rightarrow [n]\}, B = \{b_i : i \in [n]\}, \text{ and } C = \{c_i : i \in [n]\},$$

so that $\mathcal{M} \models (\psi_1 \wedge \psi_2)(a_i^f, b_j, c_k)$ if and only if $k \leq f(i, j)$. Let

$$E = \{abc \in K_3[A, B, C] : \mathcal{M} \models (\psi_1 \wedge \psi_2)(a, b, c)\},$$

and for each $\alpha \in \{0, 1\}$, let $E_\alpha = \{abc \in K_3[A, B, C] : \mathcal{M} \models \psi_\alpha(a, b, c)\}$. Note that $(A \cup B \cup C, E)$, $(A \cup B \cup C, E_0)$ and $(A \cup B \cup C, E_1)$ are each 3-partite 3-graphs.

By Lemma 6.8, there are subsets $X = \{x_i : i \in [N]\} \subseteq A$, $Y = \{y_i : i \in [N]\} \subseteq B$, and $Z_1, \dots, Z_{S_1} \subseteq C$ each of size N , so that if we define $Q_{XY}^\alpha := \{x_i y_j : u_i w_j \in P_{UW}^\alpha\}$ for each $\alpha \in [S]$, then for all $x_i y_j \in Q_{XY}^\alpha$ and $z \in Z_\beta$, we have $x_i y_j z \in E$ if and only if $\beta \leq \alpha$.

Let $V' = X \cup Y \cup Z_1 \cup \dots \cup Z_S$, and let \mathcal{Q} be the $(S+2, S)$ -decomposition of V' consisting of $\{X, Y, Z_1, \dots, Z_k\}$ and $\{Q_{XY}^\alpha : \alpha, m \leq S\}$. Define $H_1 = (V', E_1[V'])$, $H_2 = (V', E_2[V'])$, and $H_{12} = (V', (E_1 \wedge E_2)[V'])$. Note that by assumption, H_1 and H_2 have no k -FOP₂, and thus both have VC₂-dimension at most k .

By Theorem 2.49, there are $t_1 \leq T_1$, $\ell_1 \leq L_1$ and a $(t_1, \ell_1, \mu_1'', \mu_2''(\ell_1))$ -decomposition \mathcal{P}_1 of V' which is μ_1'' -homogeneous with respect to H_1 and which has linear error, say witnessed by Σ_1 . Similarly, by Theorem 2.49, there are $t_2 \leq T_1$, $\ell_2 \leq L_1$ and an $(\mu_1'', \mu_2''(\ell_2), t_2, \ell_2)$ -decomposition \mathcal{P}_2 of V' which is μ_1'' -homogeneous and which has linear error with respect to H_2 , say witnessed by Σ_2 . Let $\mathcal{P}_4 = \mathcal{P}_1 \wedge \mathcal{P}_2$ be the coarsest decomposition of V refining both \mathcal{P}_1 and \mathcal{P}_2 . Let \mathcal{P}_3 be the decomposition of V with $(\mathcal{P}_3)_{\text{vert}} = \mathcal{Q}_{\text{vert}}$ and $(\mathcal{P}_3)_{\text{edge}} = \{K_2[F, F'] : FF' \in \binom{\mathcal{Q}_{\text{vert}}}{2}\}$.

By Lemma 3.27, there exist some $t_3 \leq f_1(t_1 t_2, \ell_1 \ell_2, S+2, 1)$, $\ell_3 \leq g_1(t_1 t_2, \ell_1 \ell_2, S+2, 1)$, and a $(t_3, \ell_3, \mu_1'', \mu_2''(\ell_3))$ -decomposition \mathcal{R} of V' which is an $(\mu_1'', \mu_2''(\ell_3))$ -approximate refinement of \mathcal{P}_3 and \mathcal{P}_4 . Say this is witnessed by $\Sigma \subseteq \binom{\mathcal{R}_{\text{vert}}}{2}$, as in Definition 3.26.

By Lemma 3.28, \mathcal{R} is $(\epsilon_1'', \epsilon_2''(\ell_3))$ -regular and ϵ_1'' -homogeneous with respect to both E_1 and E_2 . Since $\mathcal{R}_{\text{vert}}$ refines $\mathcal{Q}_{\text{vert}}$, it has the form

$$\mathcal{R}_{\text{vert}} = \{X_i : i \in [t]\} \cup \{Y_i : i \in [t]\} \cup \{Z_{i,j} : i \in [S], j \in [t]\},$$

for some $t \leq t_3$, and where each $X_i \subseteq X$, $Y_i \subseteq Y$ and $Z_{i,j} \subseteq Z_i$. Let $\mathcal{R}_X = \{X_i : i \in [t]\}$, $\mathcal{R}_Y = \{Y_i : i \in [t]\}$, and $\mathcal{R}_Z = \{Z_{i,j} : i \in [S], j \in [t]\}$. For each $i, j \in [t]$ and $i' \in [S]$, we also have the partitions $K_2[X_i, Y_j] = \bigcup_{\alpha \leq \ell_3} R_{X_i Y_j}^\alpha$, $K_2[X_i, Z_{i',j}] = \bigcup_{\alpha \leq \ell_3} R_{X_i Z_{i',j}}^\alpha$ and $K_2[Y_i, Z_{i',j}] = \bigcup_{\alpha \leq \ell_3} R_{Y_i Z_{i',j}}^\alpha$ coming from $\mathcal{R}_{\text{edge}}$. For each $1 \leq i, j \leq t$, set $Q_{X_i Y_j}^\alpha = Q_{XY}^\alpha[X_i, Y_j]$. Note that by Lemma 3.18, each $Q_{X_i Y_j}^\alpha$ satisfies $\text{disc}_2(\epsilon_2'''(S), 1/S)$.

For each $X_i Y_j \notin \Sigma$ and $\tau \in [\ell_3]$, let $f_1(R_{X_i Y_j}^\tau) \in (\mathcal{P}_1)_{\text{edge}}$ and $f_2(R_{X_i Y_j}^\tau) \in (\mathcal{P}_2)_{\text{edge}}$ be such that $|R_{X_i Y_j}^\tau \setminus (f_1(R_{X_i Y_j}^\tau) \cap f_2(R_{X_i Y_j}^\tau))| \leq \mu_1'' |R_{X_i Y_j}^\tau|$, and so that $R_{X_i Y_j}^\tau \setminus (f_1(R_{X_i Y_j}^\tau) \cap f_2(R_{X_i Y_j}^\tau))$ has $\text{disc}_2(\mu_2''(\ell_3))$. Setting

$$C(X_i, Y_j) = \bigcup_{\alpha \in \ell_4} R_{X_i Y_j}^\alpha \setminus f(R_{X_i Y_j}^\alpha),$$

we have, by definition of a $(\mu_1'', \mu_2''(\ell_4))$ -approximate refinement, that for all $X_i Y_j \notin \Sigma$, $|C(X_i, Y_j)| \leq \mu_1'' |X_i| |Y_j|$. Define

$$J(X_i, Y_j) = \{Q_{XY}^\alpha \in \mathcal{Q}_{edge} : |Q_{XY}^\alpha \cap C(X_i, Y_j)| \geq \sqrt{\mu_1''} |Q_{XY}^\alpha|\}.$$

Then $|\{xy \in K_2[X_i, Y_j] : xy \in Q_{XY}^\alpha \text{ for some } Q_{XY}^\alpha \in J(X_i, Y_j)\}| \leq \sqrt{\mu_1''} |X_i| |Y_j|$, and consequently, $|J(X_i, Y_j)| \leq 2\sqrt{\mu_1''} S$. We now define

$$\Gamma = \{G \in \text{Triads}(\mathcal{R}) : \text{for each } u \in \{0, 1\}, (H_u | G, G) \text{ has } \text{disc}_{2,3}(\epsilon_1'', \epsilon_2''(\ell_3)) \text{ and}$$

$$\frac{|E_u \cap K_3^{(2)}(G)|}{|K_3^{(2)}(G)|} \in [0, \epsilon_1''] \cup (1 - \epsilon_1'', 1]\},$$

and set $\Gamma = \{xyz \in K_3[X, Y, Z] : xyz \in K_3^{(2)}(G) \text{ for some } G \in \Gamma\}$. By assumption, $|\Gamma| \geq (1 - \epsilon_1'') |V|^3$. Therefore, setting

$$I = \left\{ xy \in K_2[X, Y] : |\{z \in Z : xyz \in \Gamma\}| \geq (1 - \epsilon_1') |Z|, xy \notin \bigcup_{X_i Y_j \in \Sigma} C(X_i, Y_j) \right\},$$

we must have that $|I| \geq (1 - \epsilon_1') |V|^2$ (here we are using that $\mu_1'' \ll \epsilon_1'' \ll \epsilon_1'$). Therefore, there must be some $X_i \in \mathcal{R}_X$ and $Y_j \in \mathcal{R}_Y$ so that $|I \cap K_2[X_i, Y_j]| \geq (1 - \epsilon_1') |X_i| |Y_j|$. Given $\alpha, \beta, \gamma \leq \ell_3$, $b \in [t]$, and $u \in [S]$, let

$$R_{X_i Y_j Z_{u,b}}^{\alpha, \beta, \gamma} = (X_i \cup Y_j \cup Z_{u,b}, R_{X_i Y_j}^\alpha \cup R_{X_i Z_{u,b}}^\beta \cup R_{Y_j Z_{u,b}}^\gamma).$$

Let $\Gamma' = \Gamma \cap \{R_{X_i Y_j Z_{u,b}}^{\alpha, \beta, \gamma} : \alpha, \beta, \gamma \leq \ell_3, u \in [S], b \in [t]\}$. We now define a bipartite graph $\mathbf{G} = (U \cup W, F)$ where

$$U = \{R_{X_i Y_j}^\alpha : \alpha \leq \ell_3\}, \quad W = \{R_{X_i Z_{u,b}}^\beta R_{Y_j Z_{u,b}}^\gamma : \gamma, \beta \leq \ell_3, b \in [t], u \in [S]\} \text{ and}$$

$$F = \{R_{X_i Y_j}^\alpha (R_{X_i Z_{u,b}}^\beta R_{Y_j Z_{u,b}}^\gamma) \in K_2[U, W] : R_{X_i Y_j Z_{u,b}}^{\alpha, \beta, \gamma} \in \Gamma'\}.$$

By our choice of $X_i Y_j$,

$$\left| \left(\bigcup_{G \in \Gamma'} K_3^{(2)}(G) \right) \cap K_3[X_i, Y_j, Z] \right| \geq (1 - \epsilon_1')^2 |X_i| |Y_j| |Z|.$$

Consequently, using Corollary 3.6,

$$|F| \geq \frac{(1 - \epsilon_1')^2 |X_i| |Y_j| |Z|}{(1 + \epsilon_1') |X_i| |Y_j| |Z| / St \ell_3^3} \geq (1 - \epsilon_1')^3 St \ell_3^3 \geq (1 - 3\epsilon_1') St \ell_3^3.$$

Set

$$\Omega = \{Z_\alpha : \alpha \in [S] \text{ and for some } c \in W, |N_{\mathbf{G}}(c)| \geq (1 - \sqrt{3\epsilon_1'}) \ell_3\}.$$

Then $|F| \geq (1 - 3\epsilon_1') St \ell_3^3$ implies then $|\Omega| \geq (1 - \sqrt{3\epsilon_1'}) S$. Now set

$$\Omega' = \{Z_\alpha \in \Omega : Q_{X_i Y_j}^\alpha \notin J(X_i, Y_j)\}.$$

Then by above, $|\Omega'| \geq (1 - \sqrt{3\epsilon'_1} - 2\sqrt{\mu''_1})S$. Choose $Z_{i_1}, \dots, Z_{i_s} \in \Omega'$ with $i_1 < \dots < i_s$, and for each $1 \leq v \leq s$, choose any $c_v = R_{X_i Z_{i_v}, b_v}^{\beta_v} R_{Y_j Z_{i_v}, b_v}^{\gamma_v} \in Z_{i_v}$ so that $|N_{\mathbf{G}}(c)| \geq (1 - \sqrt{3\epsilon'_1})\ell_3$. Consider

$$O = \bigcap_{v=1}^s N_{\mathbf{G}}(c_v).$$

By construction, $|O| \geq (1 - s\sqrt{3\epsilon'_1})\ell_3$. Since $\epsilon'_1 \ll \epsilon/S$ and $S \gg s$, $|O| \geq (1 - \epsilon_1)\ell_3$. For each $v \in [s]$, consider

$$L_v = \{xy : \text{there is } P \in O \text{ with } xy \in P\} \cap Q_{XY}^{i_v}[X_i, Y_j].$$

By definition of Ω' , and because $\epsilon'_1 \ll \epsilon_1/S$,

$$\begin{aligned} |L_v| &\geq |Q_{XY}^{i_v}[X_i, Y_j]| - |Q_{XY}^{i_v}[X_i, Y_j] \cap C(X_i, Y_j)| \geq |Q_{XY}^{i_v}[X_i, Y_j]| - \sqrt{\epsilon'_1}|X_i||Y_j| \\ &\geq \left(\frac{1}{S} - 2\sqrt{\epsilon'_1}\right)|X_i||Y_j| \\ &\geq (1 - \epsilon_1)|Q_{XY}^{i_v}[X_i, Y_j]|. \end{aligned}$$

Clearly this implies there is some $R_{X_i Y_j}^{\tau_v} \in O$ with $|R_{X_i Y_j}^{\tau_v} \setminus Q_{XY}^{i_v}| \leq \epsilon_1 |R_{X_i Y_j}^{\tau_v}|$. It is not difficult to see now that for each $u \leq v$,

$$|(E_1 \cap E_2) \cap K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})| \geq (1 - \epsilon_1) |K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|,$$

while if $v < u$, then

$$|(E_1 \cap E_2) \cap K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})| < \epsilon_1 |K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|.$$

Since each of these triads $R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u}$ are in Γ , we have that for all $u \leq v$,

$$\min\{|E_1 \cap K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|, |E_2 \cap K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|\} \geq (1 - \mu''_1) |K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|,$$

and for all $v < u$, there $w_{u,v} \in \{0, 1\}$ so that

$$|E_{w_{u,v}} \cap K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})| \leq \mu''_1 |K_3^{(2)}(R_{X_i Y_j Z_{i_u}}^{\tau_v \alpha_u \beta_u})|.$$

By our choice of s , there is a subsequence $1 \leq i'_1 < \dots < i'_k \leq s$ and $w \in \{0, 1\}$ so that so $E_{w_{i'_u}, i'_u} = E_w$ for all $v < u$. Let $f : \{a_1, \dots, a_k\} \rightarrow \mathcal{R}_{edge}$ and $g : \{b_1, \dots, b_k\} \rightarrow \mathcal{R}_{cnr}$ be defined by $f(a_v) = R_{X_i Y_i}^{\tau_{i'_v}}$ and $g(b_u) = R_{X_i Z_{i'_u}}^{\alpha_{i'_u}} R_{Y_j Z_{i'_u}}^{\beta_{i'_u}}$. Then (f, g) is an encoding of $H(k)$ in (H_w, \mathcal{P}) with corresponding partition $(X_i, Y_j, \bigcup_{u=1}^k Z_{i'_u})$. By Proposition 6.7, there are $(c_1, \dots, c_k) \in Z_{i'_1} \times \dots \times Z_{i'_k}$, $\{a_1^f, \dots, a_k^f : f : [k]^2 \rightarrow [k]\} \subseteq X_i$, and $\{b_1, \dots, b_k\} \subseteq Y_j$ such that $a_i^f b_j c_k \in E_w$ if and only if $k \leq f(i, j)$. By definition of E_w , this implies that $\phi_w(x; y; z)$ has k -FOP₂ in \mathcal{M} , a contradiction. \square

Combined with Proposition D.2, this shows that being NFOP₂ is closed under finite boolean combinations.

We end by constructing for all $k \geq 1$, a 3-uniform hypergraph which has k -FOP₂ and VC₂-dimension 1.

Example D.4. Choose $\epsilon_1 > 0$ and $\epsilon_2 : \mathbb{N} \rightarrow (0, 1]$ as in Proposition 6.7 for k , and let N be as in Proposition 6.7 for $\epsilon_1, \epsilon_2, \ell = k$ and $t = k + 2$, and let $n \gg N$. Let $U = \{u_1, \dots, u_n\}$, $W = \{w_1, \dots, w_n\}$ and let $K_2[U, W] = \bigcup_{\alpha \leq N} P_{UW}^\alpha$ be a partition so that each P_{UW}^α has $\text{disc}_2(\epsilon_2(k), 1/k)$ (such a partition exists by Lemma 3.24). Let Z_1, \dots, Z_k be disjoint sets, each of size n , and set $Z = \bigcup_{i=1}^k Z_i$. We now let $V = U \cup W \cup Z$ and define $H = (V, E)$ to be the 3-partite 3-graph where $E = \{uwz : z \in Z_i, uw \in P_{UW}^\alpha \text{ for some } \alpha \leq i\}$. By Proposition 6.7, H has k -FOP₂. We show it has VC₂-dimension at most 1.

Fix any $x_1, x_2, y_1, y_2 \in V$, and let $X = \{x_1, x_2\}$ and $Y = \{y_1, y_2\}$ be two 2-element subsets. We claim that the edge relation in H cannot shatter $X \times Y$. Suppose towards a contradiction this was not the case. Then there are, for each $S \subseteq X \times Y$, some $z_S \in V$ so that $z_S x_i y_j \in E$ if and only if $(i, j) \in S$. Clearly this implies $x_1 \neq x_2$ and $y_1 \neq y_2$. Further, since $z_{X \times Y} x_i y_j$ for each $i, j \in [2]$, and since H is 3-partite, we must have that there is a relabeling $\{A, B, C\} = \{U, W, Z\}$ so that $x_1, x_2 \in A$, $y_1, y_2 \in B$, and $z_S \in C$ for all $S \neq \emptyset$.

Suppose first that $A = Z$. Note that for all $i \in [k]$ and $u \in U$ and $w \in W$, either $Z_i \subseteq N_H(uw)$ or $Z_i \cap N_H(uw) = \emptyset$. Thus we clearly cannot have x_1, x_2 in the same Z_i . Therefore, without loss of generality, we may assume that for some $1 \leq i < j \leq k$, $x_1 \in Z_i$ and $x_2 \in Z_j$. But now there exists no $uw \in K_2[U, W]$ with $x_2 uw \in E$ and $x_1 uw \notin E$, a contradiction. Thus, $A \neq Z$. A symmetric argument shows $B \neq Z$.

Thus $\{A, B\} = \{U, W\}$. Without loss of generality, assume $A = U$ and $B = W$. For each $(i, j) \in [2]^2$, let α_{ij} be such that $x_i y_j \in P_{UW}^{\alpha_{ij}}$. Clearly we must have $\alpha_{11} \neq \alpha_{12}$, since $z_{\{(1,1)\}} x_1 y_1 \in E$ but $z_{\{(1,2)\}} x_1 y_1 \notin E$. Assume $\alpha_{11} < \alpha_{12}$ (the other case is similar). But now there exists no $z \in Z$ with $z x_1 y_2 \in E$, and $z x_1 y_1 \notin E$. Again this is a contradiction, and finishes the proof.

REFERENCES

- [1] Nathanael Ackerman, Cameron Freer, and Rehana Patel, *Stable regularity for relational structures*, arXiv:1712.09305 (2017).
- [2] Noga Alon, Eldar Fischer, Michael Krivelevich, and Mario Szegedy, *Efficient Testing of Large Graphs*, *Combinatorica*, **20** (2000), no. 4, 451–476.
- [3] Noga Alon, Eldar Fischer, and Ilan Newman, *Efficient testing of bipartite graphs for forbidden induced subgraphs*, *SIAM Journal on Computing*, **37** (2007), no. 3, 959–976.
- [4] Noga Alon, János Pach, Rom Pinchasi, and Radoš Radoičić, *Crossing patterns of semi-algebraic sets*, *Journal of Combinatorial Theory*, **111** (2005), no. 2, 310–326.
- [5] Noga Alon, Asaf Shapira, and Uri Stav, *Can a graph have distinct regular partitions?*, *SIAM Journal on Discrete Mathematics*, **23** (2009), no. 1, 278–287.
- [6] Matthias Aschenbrenner, Alf Dolich, Deirdre Haskell, Dugald Macpherson, and Sergei Starchenko, *Vapnik-Chervonenkis density in some theories without the independence property, I*, *Transactions of the American Mathematical Society*, **8** (2016), no. 368, 5889–5949.
- [7] József Balogh, Bela Bollobas, and David Weinreich, *The Speed of Hereditary Properties of Graphs*, *Journal of Combinatorial Theory, Series B*, **79** (2000), no. 2, 131–156.
- [8] József Balogh and Dhruv Mubayi, *Almost all triangle-free triple systems are tripartite*, *Combinatorica*, **32** (2012), no. 2, 143–169.
- [9] Saugata Basu, *Combinatorial complexity in o-minimal geometry*, *Proceedings of the London Mathematical Society*, **100** (2009), no. 2, 405–428.

- [10] David Conlon and Jacob Fox, *Graph removal lemmas*, Surveys in combinatorics 2013, London Math. Soc. Lecture Note Ser., **409** (2013), 1–49.
- [11] Artem Chernikov and Nadja Hempel, *On n -dependent Groups and Fields II*, Forum of Mathematics, Sigma, **9**, E38, 1–51.
- [12] Artem Chernikov, Daniel Palacín, and Kota Takeuchi, *On n -dependence*, Notre Dame Journal of Formal Logic, **60** (2019), no. 2, 195–214.
- [13] Artem Chernikov and Sergei Starchenko, *Regularity lemma for distal structures*, Journal of the European Mathematical Society, **20** (2018), no. 10, 2437–2466.
- [14] ———, *Definable regularity lemmas for NIP hypergraphs*, The Quarterly Journal of Mathematics, **72** (2021), no. 4, 1401–1433.
- [15] Artem Chernikov and Henry Towsner, *Hypergraph regularity and higher arity VC-dimension*, arXiv:2010.00726 (2020).
- [16] Fan Chung, Ron Graham, and R M Wilson, *Quasirandom graphs*, Proceedings of the National Academy of Sciences of the United States of America, **85** (1988), no. 4, 969–970.
- [17] Fan Chung, *Regularity lemmas for hypergraphs and quasirandomness*, Random Structures & Algorithms, **2** (1991), no. 2, 241–252.
- [18] Fan Chung and Ron Graham, *quasirandom hypergraphs*, Random Structures and Algorithms, **1** (1990), no. 1, 105–124.
- [19] Y. Dementieva, P. E. Haxell, B. Nagle, and V. Rödl, *On characterizing hypergraph regularity*, Random Structures & Algorithms, **21** (2002), no. 3–4, 293–335.
- [20] Paul Erdős, Péter Frankl, and Vojtěch Rödl, *The asymptotic number of graphs not containing a fixed subgraph and a problem for hypergraphs having no exponent*, Graphs and Combinatorics, **2** (1986), no. 2, 113–121.
- [21] Jacob Fox and László Miklós Lovász, *A tight bound for Szemerédi’s regularity lemma*, Combinatorica, **37** (2017), no. 5, 911–951.
- [22] Jacob Fox, János Pach, and Andrew Suk, *Erdős-Hajnal Conjecture for Graphs with Bounded VC-Dimension*, Discrete and Computational Geometry, **61** (2019), no. 4, 809–829.
- [23] Péter Frankl and Vojtěch Rödl, *The uniformity lemma for hypergraphs*, Graphs and Combinatorics, **8** (1992), no. 4, 309–312.
- [24] ———, *Extremal problems on set systems*, Random Structures & Algorithms, **20** (2002), no. 2, 131–164.
- [25] Timothy Gowers, *Lower bounds of tower type for Szemerédi’s uniformity lemma*, Geometric and Functional Analysis, **7** (1997), no. 2, 322–337.
- [26] ———, *Quasirandomness, counting and regularity for 3-uniform hypergraphs*, Combinatorics, Probability and Computing, **15** (2006), no. 1–2, 143–184.
- [27] ———, *Hypergraph regularity and the multidimensional Szemerédi theorem*, Annals of Mathematics. Second Series, **166** (2007), no. 3, 897–946.
- [28] Ben Green and Tom Sanders, *Fourier uniformity on subspaces*, arXiv:1510.08739 (2015).
- [29] Nadja Hempel, *On n -dependent groups and fields*, Mathematical Logic Quarterly, **62** (2016), no. 3, 215–224.
- [30] Yoshiharu Kohayakawa, Brendan Nagle, Vojtěch Rödl, and Mathias Schacht, *Weak hypergraph regularity and linear hypergraphs*, Journal of Combinatorial Theory, Series B, **100** (2010), no. 2, 151–160.
- [31] Julie Haviland and Andrew Thomason, *Pseudo-random hypergraphs*, Discrete Mathematics, **75** (1989) no. 1–3, 255–278.
- [32] János Komlós and Miklos Simonovits, *Szemerédi’s regularity lemma and its applications in graph theory*, Combinatorics, Paul Erdős is eighty, **2** (1996), 295–352.
- [33] John Lenz and Dhruv Mubayi, *The poset of hypergraph quasirandomness*, Random Structures & Algorithms, **46** (2012), no. 4, 762–800.
- [34] László Lovász and Balázs Szegedy, *An Irregular Mind, Szemerédi is 70*, Bolyai Society Mathematical Studies (2010), 415–446.

- [35] László Lovász and Balázs Szegedy, *Szemerédi's lemma for the analyst*, Geometric and Functional Analysis, **17** (2007), no. 1, 252–270.
- [36] M. Malliaris and S. Shelah, *Regularity lemmas for stable graphs*, Transactions of the American Mathematical Society, **366** (2014), no. 3, 1551–1585.
- [37] Guy Moshkovitz and Asaf Shapira, *A Short Proof of Gowers' Lower Bound for the Regularity Lemma*, Combinatorica, **36** (2016), 187–194.
- [38] ———, *A Tight Bound for Hypergraph Regularity*, Geometric and Functional Analysis, **29** (2019), 1531–1578.
- [39] Brendan Nagle, Annika Poerschke, Vojtěch Rödl, and Matthias Schacht, *Hypergraph regularity and quasi-randomness*, Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms (2009), 227–235.
- [40] Brendan Nagle, Vojtěch Rödl, *Regularity properties for triple systems*, Random Structures & Algorithms, **23** (2003), no. 3, 264–332.
- [41] Brendan Nagle, Vojtěch Rödl, and Mathias Schacht, *The counting lemma for regular k -uniform hypergraphs*, Random Structures & Algorithms, **28** (2006), no. 2, 113–179.
- [42] Yury Person and Mathias Schacht, *Almost all hypergraphs without Fano planes are bipartite*, SODA '09: Proceedings of the Twentieth Annual ACM-SIAM Symposium on Discrete Algorithms (2009), 217–226.
- [43] Anand Pillay, *Geometric stability theory*, The Clarendon Press, Oxford University Press, New York, vol. 32, 1996.
- [44] Anand Pillay, *Domination and regularity*, The Bulletin of Symbolic Logic, **26** (2020), no. 2, 103–117.
- [45] Maryanthe Malliaris and Anand Pillay, *The stable regularity lemma revisited*, Proceedings of the American Mathematical Society, **144** (2016), no. 4, 1761–1765.
- [46] Vojtěch Rödl and Mathias Schacht, *Generalizations of the removal lemma*, Combinatorica, **29** (2009), no. 4, 467–501.
- [47] Vojtěch Rödl and Mathias Schacht, *Regularity Lemmas for Graphs*, Bolyai Society Mathematical Studies (2010), 287–325.
- [48] Vojtěch Rödl and Mathias Schacht, *Regular partitions of hypergraphs: regularity lemmas*, Combinatorics, Probability and Computing, **16** (2007) no. 6, 833–885.
- [49] Vojtěch Rödl and Jozef Skokan, *Counting subgraphs in quasirandom 4-uniform hypergraphs*, Random Structures & Algorithms, **26** (2005), no. 1–2, 160–203.
- [50] Vojtěch Rödl and Jozef Skokan, *Regularity lemma for k -uniform hypergraphs*, Random Structures & Algorithms, **25** (2004), no. 1, 1–42.
- [51] Saharon Shelah, *Classification Theory and the Number of Non-Isomorphic Models*, 2nd ed., Studies in Logic and The Foundations of Mathematics, vol. 92, Elsevier, 1990.
- [52] Saharon Shelah, *Definable Groups for dependent and 2-dependent theories*, Sarajevo J. Math, **13**(25) (2017), no. 1, 3–25.
- [53] Saharon Shelah, *Strongly dependent theories*, Israel Journal of Mathematics, **205** (2014), no. 1, 1–83.
- [54] Endre Szemerédi, *On sets of integers containing no k elements in arithmetic progression*, Acta Arithmetica, **27** (1975), 199–245.
- [55] Kota Takeuchi, *On 2-order property*, Slides from a talk given at the Asian Logic Conference 2017, Daejeon, Korea, <https://alc15korea.wixsite.com/alckorea/talk-slides> (2017), 30pp.
- [56] Katrin Tent and Martin Ziegler, *A course in model theory*, Cambridge University Press, Cambridge University Press, 2012.
- [57] Caroline Terry, *An improved bound for regular decompositions of 3-uniform hypergraphs of bounded VC₂-dimension*, arXiv: 2204.08537 (2022).
- [58] ———, *VC_ℓ-dimension and the jump to the fastest speed of a hereditary \mathcal{L} -property*, Proceedings of the American Mathematical Society, **146** (2018), 3111–3126.
- [59] Caroline Terry and Julia Wolf, *A note on generalized Green-Sanders examples*, forthcoming.
- [60] ———, *Higher-order generalizations of stability and arithmetic regularity*, arXiv: 2111.01739 (2021).

- [61] Henry Towsner, *Sigma-algebras for quasirandom hypergraphs*, Random Structures and Algorithms, **50** (2016), no. 1, 114–139.

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43210, USA
Email address: `terry.376@osu.edu`

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, CENTRE FOR MATHEMATICAL SCIENCES, WILBERFORCE ROAD, CAMBRIDGE CB3 0WB, UK
Email address: `julia.wolf@dpms.cam.ac.uk`