

Legendre pairs of lengths $\ell \equiv 0 \pmod{5}$

Dursun Bulutoglu* Ilias Kotsireas†
 Christoph Koutschan‡ Jonathan Turner§

November 4, 2021

Abstract

Compression of complementary sequences has been proved to be a valuable tool to discover several new complementary sequences of various kinds in the past decade. Compression based search algorithms encompass a two-stage process, that typically involves the computation of several candidate compression sequences, followed by the computationally expensive decompression phase. In this paper we show how to shorten the former phase in the case of Legendre pairs of lengths $\ell \equiv 0 \pmod{5}$, using a plausible conjecture, supported by overwhelming numerical evidence. This is achieved by significantly decreasing the number of candidate compression sequences by using a sums-of-squares Diophantine equation. We verify our conjecture for all odd values from 3 to 17. In particular, this allows us to exhibit the first known examples of Legendre Pairs of length 85, which has been the smallest open length. We also find Legendre pairs of the open length 87 by assuming a balanced compression. As a consequence, there remains eleven lengths less than 200 for which the question of existence of a Legendre pair remains open.

1 Introduction

Compression of complementary sequences [1] has been proved to be a valuable tool to discover several new complementary sequences of various kinds in the past decade.

*AFIT, Ohio, USA

†Supported by an NSERC grant

‡Supported by the Austrian Science Fund (FWF): F5011-N15.

§AFIT, Ohio, USA

2 Legendre Pairs of lengths $\ell \equiv 0 \pmod{5}$

We consider Legendre Pairs (A, B) of length $\ell \equiv 0 \pmod{5}$ and set $m = \ell/5$. In this case, it is known that there are exact evaluations of $\text{PSD}_A(m)$ and $\text{PSD}_B(m)$, given in [2], that we rewrite here in a slightly different form:

$$\begin{aligned}\text{PSD}_A(m) &= p_2(\mathcal{A}_m) - \frac{1}{2}e_2(\mathcal{A}_m) + \frac{\sqrt{5}}{2}(\text{PAF}_{\mathcal{A}_m}(1) - \text{PAF}_{\mathcal{A}_m}(2)), \\ \text{PSD}_B(m) &= p_2(\mathcal{B}_m) - \frac{1}{2}e_2(\mathcal{B}_m) + \frac{\sqrt{5}}{2}(\text{PAF}_{\mathcal{B}_m}(1) - \text{PAF}_{\mathcal{B}_m}(2)),\end{aligned}\tag{1}$$

where $\mathcal{A}_m = [A_1, A_2, A_3, A_4, A_5]$ and $\mathcal{B}_m = [B_1, B_2, B_3, B_4, B_5]$ are the m -compressions of A and B respectively. In addition, we have the standard relation

$$\text{PSD}_A(m) + \text{PSD}_B(m) = 2\ell + 2.$$

Since $A_1 + \dots + A_5 = B_1 + \dots + B_5 = \pm 1$ (throughout the paper, WLOG, we are going to assume $A_1 + \dots + A_5 = B_1 + \dots + B_5 = 1$), we must have

$$\begin{aligned}p_2(\mathcal{A}_m) - \frac{1}{2}e_2(\mathcal{A}_m) &= \frac{1}{4}(5p_2(\mathcal{A}_m) - 1), \\ p_2(\mathcal{B}_m) - \frac{1}{2}e_2(\mathcal{B}_m) &= \frac{1}{4}(5p_2(\mathcal{B}_m) - 1).\end{aligned}\tag{2}$$

Experimental evidence gathered for $\ell = 15, 25, 35, 45, 55$ indicates that there are Legendre pairs of these orders such that

$$\begin{aligned}\frac{1}{4}(5p_2(\mathcal{A}_m) - 1) &= \ell + 1, \\ \frac{1}{4}(5p_2(\mathcal{B}_m) - 1) &= \ell + 1.\end{aligned}\tag{3}$$

Equations (3) simplify to

$$\begin{aligned}p_2(\mathcal{A}_m) &= 4m + 1, \\ p_2(\mathcal{B}_m) &= 4m + 1.\end{aligned}\tag{4}$$

The following lemma expresses the m th entry of the power spectral density $\text{PSD}_A(m)$ of a sequence A of length $\ell = 5m$ in terms of the values in its PAF sequence.

Lemma 1. Let $m \in \mathbb{Z}^{\geq 1}$, w be the ℓ th primitive root of unity and $w' = w^m$, and A be a sequence of length $\ell = 5m$. Then

$$\begin{aligned}
\text{PSD}_A(m) &= \sum_{j=0}^{\ell-1} \text{PAF}_A(j) w^{mj} = \sum_{j=0}^{\ell-1} \text{PAF}_A(j) (w')^j = \sum_{i=0}^4 \sum_{j=0}^{\frac{\ell}{5}-1} \text{PAF}_A(5j+i) (w')^i \\
&= \sum_{j=0}^{\frac{\ell}{5}-1} \text{PAF}_A(5j) + \sum_{i=1}^2 2 \cos\left(\frac{2\pi i}{5}\right) \sum_{j=0}^{\frac{\ell}{5}-1} \text{PAF}_A(5j+i) \\
&= \sum_{j=0}^{\frac{\ell}{5}-1} \text{PAF}_A(5j) + \sum_{j=0}^{\frac{\ell}{5}-1} \sum_{i=1}^2 \left(\frac{-1 + (-1)^{i+1} \sqrt{5}}{2}\right) \text{PAF}_A(5j+i) \\
&= \sum_{j=0}^{\frac{\ell}{5}-1} \text{PAF}_A(5j) - \sum_{j=0}^{\frac{\ell}{5}-1} \sum_{i=1}^2 \frac{\text{PAF}_A(5j+i)}{2} + \frac{\sqrt{5}}{2} \left(\sum_{j=0}^{\frac{\ell}{5}-1} \sum_{i=1}^2 (-1)^{i+1} \text{PAF}_A(5j+i) \right).
\end{aligned}$$

Proof. The proof follows from the Wiener Khintchin Theorem. ■

The following proposition follows from Lemma 1.

Proposition 1. For a pair of sequences (A, B) of length $\ell = 5m$, the Legendre pair constraint

$$\text{PSD}_A(m) + \text{PSD}_B(m) = 2\ell + 2$$

is satisfied if and only if

$$\sum_{j=0}^{\frac{\ell}{5}-1} (\text{PAF}_A(5j) + \text{PAF}_B(5j)) - \sum_{j=0}^{\frac{\ell}{5}-1} \sum_{i=1}^2 \frac{\text{PAF}_A(5j+i) + \text{PAF}_B(5j+i)}{2} = 2\ell + 2,$$

and

$$\sum_{j=0}^{\frac{\ell}{5}-1} \sum_{i=1}^2 (-1)^{i+1} \frac{\text{PAF}_A(5j+i) + \text{PAF}_B(5j+i)}{2} = 0.$$

The following proposition follows directly from equations (1) and the fact that a Legendre pair (A, B) of length $\ell = 5m$ satisfies $\text{PSD}_A(m) + \text{PSD}_B(m) = 2\ell + 2$.

Proposition 2. *Let (A, B) be a Legendre pair of length $\ell = 5m$. Then there exists an even integer x and $n_1, n_2 \in \mathbb{Z}$ with $n_1 + n_2 = 2\ell + 2$ such that*

$$\begin{aligned}\text{PSD}_A(m) &= n_1 + \frac{\sqrt{5}}{2} \cdot x, \\ \text{PSD}_B(m) &= n_2 - \frac{\sqrt{5}}{2} \cdot x.\end{aligned}$$

Exhaustive searches for $\ell = 15, 25$ reveal that all Legendre pairs of these lengths have $n_1 = n_2 = \ell + 1$, where n_1, n_2 are as in Corollary 2. Non-exhaustive searches for larger odd values of ℓ which are multiples of 5 reveal the same pattern, i.e. the standard relationship $\text{PSD}_A(m) + \text{PSD}_B(m) = 2\ell + 2$ is materialized (often, but not always) in the same very specific manner. These observations give rise to the following conjecture.

Conjecture 1. *For every odd positive integer $\ell \equiv 0 \pmod{5}$, there exist Legendre pairs (A, B) of length $\ell = 5m$, such that*

$$\begin{aligned}\text{PSD}_A(m) &= \ell + 1 + \frac{\sqrt{5}}{2} \cdot x, \\ \text{PSD}_B(m) &= \ell + 1 - \frac{\sqrt{5}}{2} \cdot x\end{aligned}\tag{5}$$

for some even integer $x \geq 0$.

Conjecture 1 says that there exists a Legendre pair (A, B) such that $n_1 = n_2 = \ell + 1$ in Proposition 2, i.e., the constant $2\ell + 2$ can be distributed in a “balanced” manner, in two equal parts, in $\text{PSD}_A(m)$ and $\text{PSD}_B(m)$. Table 1 provides computational evidence for Conjecture 1.

The following proposition provides valid constraints on the m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$, of a Legendre pair (A, B) of length $\ell = 5m$.

Proposition 3. *Let $\ell = 5m$ be an odd integer. Then a pair of m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$, of a Legendre pair (A, B) of length $\ell = 5m$, must satisfy*

$$\text{PAF}_{\mathcal{A}_{(m)}}(1) - \text{PAF}_{\mathcal{A}_{(m)}}(2) = -(\text{PAF}_{\mathcal{B}_{(m)}}(1) - \text{PAF}_{\mathcal{B}_{(m)}}(2)) = x,$$

where x is as in Proposition 2.

Proof. This result follows directly from equations (1) and the fact that a Legendre pair (A, B) of length $\ell = 5m$ satisfies $\text{PSD}_A(m) + \text{PSD}_B(m) = 2\ell + 2$. \blacksquare

m	$\ell = 5m$	x
1	5	2
3	15	{0, 4, 8}
5	25	2
7	35	16
9	45	6
11	55	24
13	65	14
15	75	48
17	85	36

Table 1: Computationally verified cases for Conjecture 1 with their corresponding x values

The following are some observations pertaining to Conjecture 1.

1. For a fixed value of ℓ , the value of x in Conjecture 1 is not necessarily unique.
2. The value of the even integer x is determined by the m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$.
3. There exist Legendre pairs of length $\ell = 5m$ that do not satisfy Conjecture 1.
4. A proof of Conjecture 1 would entail an infinite class of Legendre pairs, of lengths $\ell = 5 \cdot m$, for every odd m .

The following proposition provides valid constraints on the m -compressed sequences $\mathcal{A}_{(m)}$ and $\mathcal{B}_{(m)}$ of a Legendre pair (A, B) of length $\ell = 5m$ satisfying equations (5).

Proposition 4. *A pair of m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$ of a Legendre pair (A, B) of length $\ell = 5m$ satisfying equations (5) must satisfy*

$$p_2(\mathcal{A}_m) = \text{PAF}_{\mathcal{A}_{(m)}}(0) = p_2(\mathcal{B}_m) = \text{PAF}_{\mathcal{B}_{(m)}}(0) = 4m + 1.$$

Proof. This result follows directly from equations (2) and (4). ■

The first two conditions of Proposition 4 can be recast as the following sums-of-squares Diophantine equations

$$\begin{aligned} A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4m + 1, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4m + 1. \end{aligned} \tag{6}$$

In addition, we are only interested in all-odd solutions, as A_i, B_i are odd numbers for $i = 1, \dots, 5$. Proposition 4 drastically reduces the search space by drastically truncating the number of possible m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$ that can give rise to a Legendre pair of order $\ell = 5m$. We illustrate this fact in the next section.

Another consequence of Proposition 4 is that the alphabet of the possible m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$ is also significantly truncated. More specifically, while the full alphabet for candidate m -compressed sequences is

$$\{-m, -(m-2), \dots, -1, +1, \dots, (m-2), m\},$$

using Proposition 4, the full alphabet for candidate m -compressed sequences is

$$\{-7, -5, -3, -1, 1, 3, 5, 7\},$$

for $m = 3, 5, 7, 9, 11, 13, 15, 17, 19$, i.e., Proposition 4 dictates a mild growth on the size of the alphabet for all odd ℓ which are multiples of 5 for $1 < \ell < 100$.

Finally, the linear constraints $A_1 + A_2 + A_3 + A_4 + A_5 = 1$ and $B_1 + B_2 + B_3 + B_4 + B_5 = 1$ are used to rule out some, if not any of the solutions of the sums-of-squares Diophantine equations (6). Hence, this sometimes provides additional reductions in the number of possible candidate m -compressed sequences $(\mathcal{A}_{(m)}, \mathcal{B}_{(m)})$.

3 Computational evidence for Conjecture 1

In this section, we verify Conjecture 1 for Legendre pairs of the seven known lengths $\ell = 15, 25, 35, 45, 55, 65, 75$ and the (subsequent) eighth open length $\ell = 85$. Therefore, we exhibit the first known example of a Legendre pair of length $\ell = 85$, which has been the smallest previously unknown length case for Legendre pairs.

3.1 $\ell = 15, m = 3$

All Legendre pairs of length 15 satisfy:

$$\begin{aligned} \text{PSD}_A(3) &= 15 + 1 + \frac{\sqrt{5}}{2} \cdot x, \\ \text{PSD}_B(3) &= 15 + 1 - \frac{\sqrt{5}}{2} \cdot x. \end{aligned}$$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned} A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 3 + 1 = 13, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 3 + 1 = 13. \end{aligned}$$

Each of these equations has a unique solution (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 1, 3]$.

3.2 $\ell = 25, m = 5$

All Legendre pairs of length 25 satisfy:

$$\begin{aligned}\text{PSD}_A(5) &= 25 + 1 + \frac{\sqrt{5}}{2} \cdot x, \\ \text{PSD}_B(5) &= 25 + 1 - \frac{\sqrt{5}}{2} \cdot x.\end{aligned}$$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 5 + 1 = 21, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 5 + 1 = 21.\end{aligned}$$

Each of these equations has a unique solution (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 3, 3]$.

3.3 $\ell = 35, m = 7$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 7 + 1 = 29, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 7 + 1 = 29.\end{aligned}$$

Each of these equations has only two solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 1, 5]$, $[1, 1, 3, 3, 3]$.

3.4 $\ell = 45, m = 9$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 9 + 1 = 37, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 9 + 1 = 37.\end{aligned}$$

Each of these equations has only two solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 3, 5]$, $[1, 3, 3, 3, 3]$.

3.5 $\ell = 55, m = 11$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 11 + 1 = 45, \\B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 11 + 1 = 45.\end{aligned}$$

Each of these equations has only two solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 3, 3, 5]$, $[3, 3, 3, 3, 3]$. Since the linear equations $A_1 + A_2 + A_3 + A_4 + A_5 = 1$ and $B_1 + B_2 + B_3 + B_4 + B_5 = 1$ must be satisfied, solutions based on $[3, 3, 3, 3, 3]$ are ruled out.

3.6 $\ell = 65, m = 13$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 13 + 1 = 53, \\B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 13 + 1 = 53.\end{aligned}$$

Each of these equations has only three solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 1, 7]$, $[1, 1, 1, 5, 5]$, $[1, 3, 3, 3, 5]$. Since the linear equations $A_1 + A_2 + A_3 + A_4 + A_5 = 1$ and $B_1 + B_2 + B_3 + B_4 + B_5 = 1$ must be satisfied, solutions based on $[1, 1, 1, 1, 7]$ are ruled out.

3.7 $\ell = 75, m = 15$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 15 + 1 = 61, \\B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 15 + 1 = 61.\end{aligned}$$

Each of these equations has only three solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 1, 3, 7]$, $[1, 1, 3, 5, 5]$, $[3, 3, 3, 3, 5]$.

3.8 $\ell = 85, m = 17$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned}A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 17 + 1 = 69, \\B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 17 + 1 = 69.\end{aligned}$$

Each of these equations has only two solutions (up to sign changes and permutations of entries) with all-odd values, namely $[1, 1, 3, 3, 7]$, $[1, 3, 3, 5, 5]$.

We used the subgroup of order 2, $H = \{1, 69\}$ which acts on \mathbb{Z}_{85}^* and yields 16 cosets of size 1 and 34 cosets of size 2. We choose 12 cosets of size 1 and 15 cosets of size 2, to make a block of size $12 \cdot 1 + 15 \cdot 2 = 42$. Therefore, the size of the search space is $\binom{16}{12} \cdot \binom{34}{15} = 1820 \cdot 1,855,967,520 = 3,377,860,886,400$. We found 4 Legendre pairs of length $\ell = 85$, made out of 6 different sequences, this was not an exhaustive search. Their LexRank encoding is given as

$$\begin{aligned} & \{\{12, 1321116338\}, \{42, 1275934280\}\}, \\ & \{\{12, 1843909851\}, \{42, 606586783\}\}, \\ & \{\{42, 1275934280\}, \{9, 1555522731\}\}, \\ & \{\{42, 606586783\}, \{9, 788215097\}\}. \end{aligned}$$

For the first Legendre pair (A, B) of length $\ell = 85$ shown above, i.e. $\{\{12, 1321116338\}, \{42, 1275934280\}\}$, in order to decode the LexRank encoding one can proceed as follows.

- In the space of $\binom{16}{12} = 1820$ 12-subsets of $\{1, \dots, 16\}$, decode

$$\begin{aligned} 12 \text{ as } A_{\text{ones}} &= \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 14, 15\}, \\ 42 \text{ as } B_{\text{ones}} &= \{1, 2, 3, 4, 5, 6, 7, 8, 10, 11, 14, 15\}. \end{aligned}$$

- In the space of $\binom{34}{15} = 1,855,967,520$ 15-subsets of $\{1, \dots, 34\}$, decode

$$\begin{aligned} 1321116338 \text{ as } A_{\text{twos}} &= \{3, 4, 5, 7, 10, 11, 22, 24, 25, 27, 28, 29, 30, 31, 34\}, \\ 1275934280 \text{ as } B_{\text{twos}} &= \{2, 8, 10, 11, 12, 15, 19, 21, 23, 25, 26, 28, 29, 33, 34\}. \end{aligned}$$

- Enumerate the 16 cosets of size 1 in increasing order as

$$O_1 = \{\{5\}, \{10\}, \{15\}, \{20\}, \{25\}, \{30\}, \{35\}, \{40\}, \{45\}, \{50\}, \{55\}, \{60\}, \{65\}, \{70\}, \{75\}, \{80\}\}.$$

- Enumerate the 34 cosets of size 2 in increasing order of their first element as

$$\begin{aligned} O_2 = & \{\{1, 69\}, \{2, 53\}, \{3, 37\}, \{4, 21\}, \{6, 74\}, \{7, 58\}, \{8, 42\}, \{9, 26\}, \{11, 79\}, \\ & \{12, 63\}, \{13, 47\}, \{14, 31\}, \{16, 84\}, \{17, 68\}, \{18, 52\}, \{19, 36\}, \{22, 73\}, \{23, 57\}, \\ & \{24, 41\}, \{27, 78\}, \{28, 62\}, \{29, 46\}, \{32, 83\}, \{33, 67\}, \{34, 51\}, \{38, 72\}, \{39, 56\}, \\ & \{43, 77\}, \{44, 61\}, \{48, 82\}, \{49, 66\}, \{54, 71\}, \{59, 76\}, \{64, 81\}\}. \end{aligned}$$

- Make the block of size 42 of the indices of the positions of the -1 elements in A , by combining 12 elements of O_1 whose indices are given by A_{ones} and 15 elements

of O_2 whose indices are given by A_{twos} . This yields the following A -block of size 42: $\{5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 70, 75, 3, 37, 4, 21, 6, 74, 8, 42, 12, 63, 13, 47, 29, 46, 33, 67, 34, 51, 39, 56, 43, 77, 44, 61, 48, 82, 49, 66, 64, 81\}$.

- Make the block of size 42 of the indices of the positions of the -1 elements in B , by combining 12 elements of O_1 whose indices are given by B_{ones} and 15 elements of O_2 whose indices are given by B_{twos} . This yields the following B -block of size 42: $\{5, 10, 15, 20, 25, 30, 35, 40, 50, 55, 70, 75, 2, 53, 9, 26, 12, 63, 13, 47, 14, 31, 18, 52, 24, 41, 28, 62, 32, 83, 34, 51, 38, 72, 43, 77, 44, 61, 59, 76, 64, 81\}$.
- It can readily be verified that the above A -block and B -block (both of size 42), give a Legendre pair for $\ell = 85$.

For the first Legendre pair (A, B) of length $\ell = 85$ shown above, we have that the 17-compressions of A, B are

$$\mathcal{A}_{(17)} = [1, 3, 3, 1, -7], \mathcal{B}_{(17)} = [3, 1, 1, 3, -7].$$

These possess the properties required by Proposition 4, i.e.,

$$\text{PAF}_{\mathcal{A}_{(17)}}(0) = \text{PAF}_{\mathcal{B}_{(17)}}(0) = 4 \cdot 17 + 1 = 69.$$

As a consequence, we have that the coefficients of $\frac{\sqrt{5}}{2}$ in $\text{PSD}_A(17)$ and $\text{PSD}_B(17)$ cancel out, where

$$\text{PSD}_A(17) = 85 + 1 + \frac{\sqrt{5}}{2} \cdot 36 = 126.2492236,$$

$$\text{PSD}_B(17) = 85 + 1 - \frac{\sqrt{5}}{2} \cdot 36 = 45.75077641.$$

3.9 $\ell = 95, m = 19$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned} A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 19 + 1 = 77, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 19 + 1 = 77. \end{aligned}$$

These equations have only four solutions (up to sign changes and permutations of entries) with all-odd values, namely

$$[1, 1, 1, 5, 7], [1, 1, 5, 5, 5], [1, 3, 3, 3, 7], [3, 3, 3, 5, 5].$$

However, solutions based on $[1, 1, 5, 5, 5]$ are ruled out, as the linear equations $A_1 + A_2 + A_3 + A_4 + A_5 = 1$ and $B_1 + B_2 + B_3 + B_4 + B_5 = 1$ cannot be satisfied.

3.10 $\ell = 115, m = 23$

Proposition 4 implies the following two sums-of-squares Diophantine equations

$$\begin{aligned} A_1^2 + A_2^2 + A_3^2 + A_4^2 + A_5^2 &= 4 \cdot 23 + 1 = 93, \\ B_1^2 + B_2^2 + B_3^2 + B_4^2 + B_5^2 &= 4 \cdot 23 + 1 = 93. \end{aligned}$$

These equations have only three solutions (up to sign changes and permutations of entries) with all-odd values, namely

$$[1, 1, 1, 3, 9], [1, 3, 3, 5, 7], [3, 3, 5, 5, 5].$$

However, solutions based on $[1, 1, 1, 3, 9]$ are ruled out, as the linear equations $A_1 + A_2 + A_3 + A_4 + A_5 = 1$ and $B_1 + B_2 + B_3 + B_4 + B_5 = 1$ cannot be satisfied.

4 Newly discovered Legendre pairs of order $\ell = 87$ with balanced compressions

By a property of compression of Legendre pairs that can be deduced from a theorem in [1], a 3-compression of a Legendre pair of order $\ell = 87$ must contain 14 elements with absolute value equal to three and $2 \cdot 29 - 14 = 44$ elements with absolute value equal to 1. Experimental evidence for other orders that are divisible by 3 indicate that “balanced” configurations of candidate 3-compressions that have exactly half of the total number of elements with absolute value equal to 3, are likely to yield Legendre pairs. Hence, we computed approximately six thousand candidate 3-compressions satisfying the following constraints, where only the last constraint is not necessary.

1. The sequences $\mathcal{A}_3, \mathcal{B}_3 \in \{-3, -1, +1, +3\}^{29}$ contain 14 elements with absolute value equal to 3 and 44 elements with absolute value equal to 1.
2. $\text{PAF}_{\mathcal{A}_3}(s) + \text{PAF}_{\mathcal{B}_3}(s) = (-2) \cdot 3 = -6$.
3. $\text{PSD}_{\mathcal{A}_3}(s) + \text{PSD}_{\mathcal{B}_3}(s) = 2 \cdot 87 + 2 = 176$.
4. $A_1 + \dots + A_{29} = 1$.
5. $B_1 + \dots + B_{29} = 1$.
6. Each of \mathcal{A}_3 and \mathcal{B}_3 contains 7 elements with absolute value equal to 3 and 22 elements with absolute value equal to 1.

Subsequently, we run our C 3-uncompression code for approximately two thousand candidate 3-compressions and discovered the two Legendre pairs of order $\ell = 87$ below.

$$\begin{aligned}
A_{87} &= [-1, -1, -1, -1, 1, -1, -1, -1, 1, 1, -1, 1, -1, 1, 1, -1, -1, 1, 1, -1, -1, 1, -1, 1, 1, \\
&\quad -1, -1, 1, 1, 1, -1, -1, -1, -1, -1, 1, -1, -1, 1, -1, -1, 1, 1, 1, 1, -1, -1, -1, 1, -1, 1, 1, \\
&\quad 1, -1, 1, 1, -1, -1, -1, -1, 1, 1, 1, 1, 1, 1, -1, -1, 1, 1, 1, -1, 1, 1, 1, 1, 1, -1, 1, -1, 1, -1], \\
B_{87} &= [-1, -1, -1, -1, -1, -1, -1, -1, 1, 1, 1, 1, -1, 1, 1, -1, 1, 1, -1, 1, 1, 1, 1, -1, -1, 1, 1, \\
&\quad -1, 1, 1, -1, 1, -1, -1, 1, -1, -1, 1, 1, 1, -1, 1, -1, -1, 1, -1, -1, 1, 1, 1, 1, 1, -1, -1, \\
&\quad -1, 1, 1, 1, -1, -1, 1, -1, 1, -1, -1, 1, -1, -1, -1, 1, -1, -1, 1, 1, 1, -1, 1, -1, 1, -1], \\
A_{87} &= [-1, -1, -1, -1, 1, -1, -1, -1, 1, 1, 1, 1, 1, 1, -1, 1, -1, -1, -1, 1, 1, 1, -1, 1, -1, -1, -1, 1, \\
&\quad 1, -1, -1, 1, 1, 1, -1, -1, -1, 1, -1, -1, 1, -1, 1, -1, -1, 1, 1, 1, 1, -1, -1, 1, 1, 1, -1, 1, 1, \\
&\quad -1, -1, -1, -1, 1, -1, 1, 1, -1, -1, -1, -1, 1, 1, 1, -1, 1, 1, -1, 1, -1, -1, 1, 1, 1, -1], \\
B_{87} &= [-1, 1, -1, 1, 1, -1, -1, 1, 1, 1, 1, 1, 1, -1, 1, -1, 1, 1, 1, -1, -1, 1, -1, 1, -1, 1, -1, \\
&\quad -1, -1, 1, -1, -1, -1, -1, -1, -1, -1, 1, -1, 1, 1, 1, -1, 1, -1, -1, 1, -1, -1, 1, 1, 1, -1, -1, \\
&\quad 1, 1, -1, -1, 1, -1, 1, -1, -1, 1, -1, 1, -1, -1, 1, -1, -1, 1, -1, -1, 1, 1, 1, 1, 1, 1, 1, -1].
\end{aligned}$$

Both of the above Legendre pairs of length $\ell = 87$ 3-compress to

$$\begin{aligned}
A_{29} &= [-3, -1, 1, -1, 1, -3, -3, -1, 1, 1, 1, 1, -1, 1, -3, 1, 1, 1, -1, 3, 3, -1, -1, 1, -1, 1, -1, 3, 1], \\
B_{29} &= [-3, -1, 1, -1, 1, -3, -3, 1, -1, 1, 1, 1, 1, -1, 3, -3, 1, 1, -1, -1, -1, 1, 1, 3, 1, 1, -1, 3, -1].
\end{aligned}$$

The lengths $\ell = 85, 87$ have been the smallest open lengths < 100 for Legendre pairs, see [4], [3].

5 Conclusion

Recently, a Legendre pair of (the previously open) length 77 and of lengths 117, 129, 133, 147 were found recently in [4] and [3] respectively. In this paper, we find Legendre pairs of (the previously open) lengths 85, 87. This reduces the list of integers less than 200 for which the question of existence of Legendre pairs remains open to

$$115, 145, 147, 159, 161, 169, 175, 177, 185, 187, 195.$$

References

- [1] Dragomir Z. Djokovic and Ilias S. Kotsireas. Compression of periodic complementary sequences and applications. *Des. Codes Cryptogr.*, 74(2):365–377, 2015.

- [2] Ilias S. Kotsireas, Christos Koukouvinos, and Jennifer Seberry. Weighing matrices and string sorting. *Ann. Comb.*, 13(3):305–313, 2009.
- [3] Ilias S. Kotsireas and Christoph Koutschan. Legendre pairs of lengths $\ell = 0 \pmod 3$. *Journal of Combin. Designs*, 29(12):870–887, 2021.
- [4] Jonathan S. Turner, Ilias S. Kotsireas, Dursun A. Bulutoglu, and Andrew J. Geyer. A Legendre pair of length 77 using complementary binary matrices with fixed marginals. *Des. Codes Cryptogr.*, 89(6):1321–1333, 2021.