

On the clique behavior of graphs of low degree

Rafael Villarroel-Flores
 Universidad Autónoma del Estado de Hidalgo
 Carretera Pachuca-Tulancingo km. 4.5
 Pachuca 42184 Hgo. México
 email: rafaelv@uaeh.edu.mx
 MSC: 05C76

keywords: iterated clique graphs, convergent graphs
 Partially supported by CONACYT, grant A1-S-45528.

May 20, 2022

Abstract

To any simple graph G , the clique graph operator K associates the graph $K(G)$ which is the intersection graph of the maximal complete subgraphs of G . The iterated clique graphs are defined by $K^0(G) = G$ and $K^n(G) = K(K^{n-1}(G))$ for $n \geq 1$. If there are $m < n$ such that $K^m(G)$ is isomorphic to $K^n(G)$ we say that G is convergent, otherwise, G is divergent. The first example of a divergent graph was shown by Neumann-Lara in the 1970s, and is the graph of the octahedron. In this paper, we prove that among the connected graphs with maximum degree 4, the octahedron is the only one that is divergent.

1 Introduction

All graphs in this paper are finite and simple. Following [5], a *clique* in a graph G is a maximal and complete subgraph of G . The *clique graph* of G , denoted by $K(G)$, is the intersection graph of its cliques. We define the sequence of *iterated clique graphs* by $K^0(G) = G$ and $K^n(G) = K(K^{n-1}(G))$ for $n \geq 1$. We say that the graph G is *convergent* if the sequence of iterated clique graphs has, up to isomorphism, a finite number of graphs. This is equivalent to the existence of $m < n$ such that $K^m(G)$ is isomorphic to $K^n(G)$ and to the fact that the set of orders of the iterated clique graphs is bounded. If G is not convergent we say that G is *divergent*. The graph of

the octahedron was the first known example of a divergent graph, given by Neumann-Lara in [15].

The problem of determining the *behavior* of G under iterated applications of the clique operator is one of the main topics in this theory, as the clique graph operator is considered one of the most complex graph operators ([17]). There are many families of graphs for which criterions have been proved, and in some cases the behavior can even be determined in polynomial time (see for example [7], [8], [11], [9], [10], [13], [14], [6], [4], [1]), however the problem has been found to be undecidable for automatic graphs ([2]).

We denote the maximum degree in G as $\Delta(G)$. The purpose of this paper is to show the following:

Main Theorem. *If G is a connected graph with $\Delta(G) \leq 4$ and G is not the graph of the octahedron, then G is convergent.*

We use standard notation for graph theory, such as $x \sim y$ whenever the vertices x, y are adjacent, $N(x)$ for the set of neighbors of the vertex x , $N[x] = N(x) \cup \{x\}$, and $|A|$ for the cardinality of the set A . Throughout the paper, we usually identify a subset of the vertex set of a graph with the subgraph that it induces. When we say that two sets *meet*, we mean that they have non-empty intersection. A complete subgraph will be called just a *complete*. The difference of the sets A, B is denoted as $A - B$.

2 Preliminaries

We say that a collection of sets is *intersecting* if any two members of the collection has nonempty intersection. The collection has the *Helly property* if any intersecting subcollection has nonempty intersection. A graph G is *Helly* if the collection of the cliques of G has the Helly property.

The following theorem gives a criterion for convergence.

Theorem 2.1. ([3]) *If a graph G is Helly, then G is convergent.*

Not every connected graph with $\Delta(G) \leq 4$ different from the octahedron is Helly. However, we will prove that for any such graph, one has that $K^2(G)$ is Helly. In order to do that, we will have to study the vertices of $K^2(G)$, that is, the cliques of cliques of G . We will pick some terminology from [10], but for the convenience of the reader, we include all the relevant definitions here.

Given $x \in G$, we define the *star* of x , denoted by x^* , as the set of cliques of G that contain x , that is, $x^* = \{q \in K(G) \mid x \in q\}$. This is a complete




Figure 1: Hajós diagram

subgraph of $K(G)$. If $x^* \in K^2(G)$, we say that x is a *normal vertex*. If $Q \in K^2(G)$ is such that $\cap Q = \emptyset$, we say that Q is a *necktie* of G . Thus, vertices of $K^2(G)$ are partitioned in stars and neckties of G , and G is Helly precisely when G has no neckties.

A *triangle* in a graph G is a complete with three vertices. A triangle T in G will be called here an *inner triangle* if T is a clique of G and for any of the edges of T there is a triangle $T' \neq T$ such that $T \cap T'$ is precisely that edge. Given an inner triangle T in G , we define $Q_T = \{q \in K(G) \mid |q \cap T| \geq 2\}$. This is always a complete in $K(G)$. We will prove that if G is a graph with $\Delta(G) \leq 4$, and G is not the octahedron, then Q_T is always a necktie, and also that for any necktie Q there is a triangle T of G , such that $Q = Q_T$. In this case, we will say that T is the *center* of the necktie Q_T , and any other element of the necktie that is not the center, will be called an *ear*.

A graph G is *hereditary Helly* if all its induced subgraphs are Helly. Hence any hereditary Helly graph is Helly. The strategy of the proof of the Main Theorem will be to prove that if G satisfies the hypothesis, then $K^2(G)$ is hereditary Helly. We will use the characterization proved in [16] as reformulated in [12].

Theorem 2.2. (Theorem 2.1 in [16]) *A graph G is hereditary Helly if and only if it is compatible with the diagram in Figure 1, in the sense that whenever the graph with the solid edges is a subgraph of G , then one of the dashed edges is induced.*

3 Neckties in low-degree graphs

In what follows, we will call a connected graph G with $\Delta(G) \leq 4$ and different from the octahedron, a *low degree graph*.

Note that because of the degree condition on G , the cliques of G can have at most five vertices. However, the only connected low degree graph that has a clique of five vertices is precisely K_5 , which is convergent. Hence, for our purposes we may assume that all cliques of G have at most four vertices.

Since the graphs that have no neckties are Helly and hence also convergent, we might as well suppose that G has neckties.

Lemma 3.1. *Let G be any graph, and q_1, q_2 be cliques in G such that if $\{x, y\}$ is any edge between vertices in $q_1 \cup q_2$, with $x \in q_1$ and $y \in q_2$, then one of the vertices x, y is in $q_1 \cap q_2$. Then if c is a complete subgraph such that $c \subseteq q_1 \cup q_2$, then $c \subseteq q_1$ or $c \subseteq q_2$.*

Proof. With the given hypothesis, suppose that there is $x \in c$ such that $x \notin q_1$ and that there is $y \in c$ such that $y \notin q_2$. Then $\{x, y\}$ is an edge with $x \in q_1$ and $y \in q_2$, such that $\{x, y\} \cap (q_1 \cap q_2) = \emptyset$. \square

Note that the hypothesis of Lemma 3.1 is satisfied in the case that $|q_1 - q_2| = 1$. Because in that case, if $\{x\} = q_1 - q_2$, and there was an edge from x to the vertex $y \in q_2 - q_1$, then $q_1 \cup \{y\}$ would be a complete subgraph properly containing q_1 .

Lemma 3.2. *If G is a graph with $\Delta(G) \leq 4$, then the neckties of G can contain only triangles.*

Proof. Let Q be a necktie in G . If $q_1 \in Q$ was an edge $\{x, y\}$, choose $q_2 \in Q$ such that $y \notin q_2$ and $q_3 \in Q$ such that $x \notin q_3$. Then $q_1 \cap q_2 = \{x\}$ and $q_1 \cap q_3 = \{y\}$. Let $z \in q_2 \cap q_3$. Then $\{x, y, z\}$ would be a complete subgraph properly containing q_1 , which is a contradiction.

Now, suppose $q_1 \in Q$ is such that $|q_1| = 4$. If there was $q' \in Q$ such that $|q_1 \cap q'| = 1$, then, since by the previous paragraph, we have $|q'| \geq 3$, we would have that if $x \in q_1 \cap q'$, then $N[x] \supseteq q_1 \cup q'$, and this last set has at least six elements, against our hypothesis. Suppose now that there is $q_2 \in Q$ with $|q_1 \cap q_2| = 3$. Then $|q_2| = 4$. Suppose that $q_1 \cap q_2 = \{x, y, z\}$. Then there is $q_3 \in Q$ such that $x \notin q_3$. If q_3 were disjoint to $q_1 \cap q_2$, then $q_1 \cup q_2$ would be a complete of five vertices. Hence we may assume $y \in q_3$. Then $q_3 \subseteq N[y] = q_1 \cup q_2$, but by Lemma 3.1 this would imply that either $q_3 \subseteq q_1$ or that $q_3 \subseteq q_2$, which is a contradiction.

Hence every clique in Q other than q_1 meets q_1 in exactly two elements, and has exactly three elements. Let $q_2 \in Q$ such that $q_2 \neq q_1$. If there was $q_3 \in Q$ such that $q_3 \cap q_1$ is disjoint to $q_2 \cap q_1$, we would have $q_1 = (q_1 \cap q_2) \cup (q_1 \cap q_3)$, and if $z \in q_2 \cap q_3$, then $z \notin q_1$, but $q_1 \cup \{z\}$ is a complete subgraph, contradicting our choice of q_1 . Hence all cliques in Q meet $q_1 \cap q_2$. Suppose $q_1 = \{x, y, u, v\}$ and $q_2 = \{x, y, w\}$. Let $q_3 \in Q$ such that $y \notin q_3$. Then $x \in q_3$, and if there was $z' \in q_3 - (q_1 \cup q_2)$, we would have $\{y, u, v, w, z'\} \subseteq N(x)$, which is a contradiction. Hence $q_3 \subseteq q_1 \cup q_2$.

By Lemma 3.1, we would have that $q_3 \subseteq q_1$ or $q_3 \subseteq q_2$, which is also a contradiction. \square

Lemma 3.3. *If G is a low-degree graph, for any necktie Q in G there is an inner triangle T of G such that $Q = Q_T$, moreover Q_T has exactly three ears. Conversely, if T is an inner triangle, then Q_T is indeed a necktie.*

Proof. We adapt the proof of Proposition 10 in [10] to this situation. For G a low-degree graph, let Q be a necktie of G . From Lemma 3.2, we know that Q contains only triangles. Let $T \in Q$. We claim that there must be a triangle in Q that meets T in only one vertex. Suppose every other triangle in Q meets T at an edge. Let $T_1 \in Q$, with $T_1 \neq T$ and suppose $T = \{a, b, c\}$ and $T_1 = \{a, b, c'\}$. Now, there must be $T_2 \in Q$ such that $b \notin T_2$. But then $T_2 \cap T = \{a, c\}$ and $T_2 \cap T_1 = \{a, c'\}$, which implies $T_2 \subseteq T \cup T_1$. By Lemma 3.1, this is not possible.

Let then $T = \{a, b, c\} \in Q$, and $T_1 = \{a, b', c'\}$ with $T \cap T_1 = \{a\}$. Choose $T_2 \in Q$ such that $a \notin T_2$. Then $T_2 \cap T$ must consist of exactly one vertex, since if $T_2 \cap T = \{b, c\}$, then T would be properly contained in a complete graph of four vertices. Similarly, $T_2 \cap T_1$ consists of only one vertex. Without loss, assume that $T_2 = \{a', b, c'\}$, with $a' \notin T \cup T_1$. Then $T_3 = \{a, b, c'\}$ is a triangle of G . It must be a clique of G because if, for example, was contained properly in $\{a, b, c', z\}$, we would have that $\{b, b', c, c', z\} \subseteq N(a)$, or if, say, we had $z = c$, then T would be contained in a tetrahedron. Hence T_3 is an inner triangle.

We claim that $T_3 \in Q$. If this were not so, there should be $T_4 \in Q$ such that $T_4 \cap T_3 = \emptyset$, but that meets T , T_1 and T_2 . We must have then that $T_4 \cap T = \{c\}$, $T_4 \cap T_1 = \{b'\}$ and $T_4 \cap T_2 = \{a'\}$. This implies that $T_4 = \{a', b', c\}$, and that the subgraph of G induced by $\{a, a', b, b', c, c'\}$ is an octahedron. This contradicts our assumption on G , and so this proves our claim that $T_3 \in Q$.

We have now that $\{T, T_1, T_2, T_3\} \subseteq Q$. Suppose there was a triangle T_4 of G , with $T_4 \in Q$, and different from T, T_1, T_2, T_3 . We must have $T_4 \cap T_3 \neq \emptyset$, without loss, assume that $a \in T_4 \cap T_3$. Suppose that $T_4 = \{a, r, s\}$. But then $T_4 \subseteq N[a] = \{a, b, c, b', c'\}$. It is immediate to verify that any choice of r, s among b, c, b', c' leads to contradict either that T_4 is different from all of T, T_1, T_2, T_3 , that each of these are cliques, or that $T_4 \cap T_2 \neq \emptyset$. Hence $Q = \{T, T_1, T_2, T_3\} = Q_{T_3}$, where T, T_1, T_2 are the ears of Q_{T_3} .

Now, for the converse, let $T = \{a, b, c\}$ be an inner triangle. Each of the cliques in Q_T contains at least two vertices of T , so Q_T is a complete in $K(G)$. Since T is an internal triangle, there are cliques q_1, q_2, q_3 in G ,

each different from T and such that $\{a, b\} \subseteq q_1 \cap T$, $\{a, c\} \subseteq q_2 \cap T$ and $\{b, c\} \subseteq q_3 \cap T$. Now, since $|T| = 3$, such inclusions must in fact be equalities, which implies q_1, q_2, q_3 are three different cliques. Let $c' \in q_1 - T$, $b' \in q_2 - T$ and $a' \in q_3 - T$. Since T is a clique it follows that c' is not adjacent to c , and similarly, b' is not adjacent to b and a' is not adjacent to a . This implies that $\{a', b', c'\}$ are three different vertices. Now, we have that $\{b, c, b', c'\} \subseteq N(a)$, and given that G is a low degree graph, this inclusion is equality. It follows then that $q_1 = \{a, b, c'\}$, $q_2 = \{a, c, b'\}$ and similarly, $q_3 = \{b, c, a'\}$. We have now that $\{T, q_1, q_2, q_3\} \subseteq Q_T$. Suppose $q \in Q_T$ with $q \neq T$. Without loss, assume that $q \cap T = \{a, b\}$. Then $q \subseteq N[a] \cap N[b] = \{a, b, c, c'\}$. Since $q \neq T$ we have that $c \notin q$ and so $q = \{a, b, c'\} = q_1$. It follows then that $Q_T = \{T, q_1, q_2, q_3\}$. Suppose there was $q \in K(G)$ such that $Q_T \cup \{q\}$ is a complete subgraph of $K(G)$ with $q \notin Q_T$. Then $|q \cap T| = 1$, and without loss, assume $q \cap T = \{a\}$. Then $q \subseteq N[a] = \{a, b, c, b', c'\}$. On the other hand, must have $q \cap q_3 \neq \emptyset$. Since $|q \cap T| = 1$, we have that neither of b, c could be an element of q . It would follow that $a' \in q \cap q_3$. But this contradicts that a is not adjacent to a' . It follows then that Q_T is a clique of $K(G)$, in fact, it is a necktie. \square

4 Forbidden subgraphs

We will apply Theorem 2.2 to $K^2(G)$, but we will first state a series of lemmas to help with the several cases that will arise. Some facts that were proven in [10] in the context of graphs of local girth at least 7 are true also in the case of low degree graphs.

Lemma 4.1. *For any low degree graph G :*

1. *Two stars x^* and y^* are adjacent in $K^2(G)$ if and only if $x \sim y$ in G .*
2. *A star x^* and a necktie Q_T centered at the triangle T are adjacent in $K^2(G)$ if and only if x is adjacent to two vertices of T .*
3. *Two neckties $Q_T, Q_{T'}$ are adjacent in $K^2(G)$ if and only if:*
 - $|T \cap T'| = 2$, or
 - $|T \cap T'| = 1$ and if $v \in T \cap T'$, there is an edge (called a crossbar) joining a vertex in $T - v$ to a vertex in $T' - v$.

Proof. In fact, it is immediate to prove that the first two statements are true in any simple graph. Now, for the third, if $|T \cap T'| = 2$, then $T \in Q_T \cap Q_{T'}$,

and so Q_T is adjacent to $Q_{T'}$ in $K^2(G)$ in this case. In the case that $T \cap T' = \{v\}$ and there is an edge $\{u, u'\}$ with $u \in T - v$ and $u' \in T' - v$, then the triangle $\{v, u, u'\}$ can be extended to a clique q such that $q \in Q_T \cap Q_{T'}$.

Now, suppose T, T' are internal triangles in a low degree graph such that Q_T is adjacent to $Q_{T'}$ in $K^2(G)$, and that $|T \cap T'| \leq 1$. Let $q \in Q_T \cap Q_{T'}$. By Lemma 3.2, we have that $|q| = 3$. Since $|q \cap T| \geq 2$ and $|q \cap T'| \geq 2$, we have that $T \cap T'$ cannot be empty, so that $|T \cap T'| = 1$. Let $T \cap T' = \{v\}$. If $x \in (q \cap T) - v$ and $x' \in (q \cap T') - v$, then $\{x, x'\}$ is a crossbar. \square

Lemma 4.2. *Let G be a low-degree graph. If T and T' are triangles in G , where $T \cap T' = \{x\}$ and T is an inner triangle, then $N(x)$ induces a 4-cycle. Moreover, if T' is also inner, then none of the ears of Q_T is an inner triangle, and none of the vertices of T besides x is a normal vertex.*

Proof. Suppose that $T = \{x, e, f\}$ and $T' = \{x, g, h\}$. Since the triangle T is inner, considering the edge $\{x, e\}$ there must be $y \in G$ and a triangle $T_1 = \{x, e, y\} \neq T$. We have that $y \in N(x) = \{e, f, g, h\}$. Since $y \neq e$ because $|T_1| = 3$ and $y \neq f$ because $T_1 \neq T$, we must have that $y = g$ or $y = h$. Without loss, assume $y = g$, hence $e \sim g$. Now, considering analogously the edge $\{x, f\}$, and given that T is a clique of G , we must have $f \sim h$. Since T is a clique, we have that e is not adjacent to h , and also that f is not adjacent to g . Hence $N(x)$ induces a 4-cycle. Given that the degree of x is 4, we must have that T' is a clique. Since T is inner, then there is a vertex u such that the triangle $\{u, e, f\}$ is different from T . Besides, $u \notin \{g, h\}$ because T is a clique.

In the case that T' is inner, then analogously there is a vertex u' such that $\{u', g, h\}$ is different from T' . We have that $u \neq u'$, otherwise, G would be 4-regular and an octahedron. Let us consider the triangle $T_1 = \{e, g, x\}$, which is an ear in Q_T . If T_1 were inner, there would be a vertex y such that $\{e, g, y\}$ is a triangle different from T_1 . But $y \in N(e) \cap N(g) = \{x\}$, which is not possible. Hence T_1 is not inner, and by symmetry, $\{f, h, x\}$ is not inner either. Now, consider $T_2 = \{e, f, u\}$, and suppose there is a vertex z such that $\{e, u, z\}$ is a triangle different from T_2 . Then $z \in N(e) = \{u, f, x, g\}$. None of these is possible by our assumptions. Hence T_2 is not inner. Finally, neither e^* nor f^* are cliques, since they are properly contained in Q_T . \square

As a corollary to the two previous lemmas, we have that if T_1 and T_2 are internal triangles where $T_1 \cap T_2 \neq \emptyset$, then $Q_{T_1} \sim Q_{T_2}$ in $K^2(G)$. Because if $|T_1 \cap T_2| = 1$, then there are (two) crossbars from T_1 to T_2 . The situation in this case is pictured in Figure 2.




Figure 2: Case $|T_1 \cap T_2| = 1$




Figure 3: Case $|T_1 \cap T_2| = 2$

Lemma 4.3. *Let G be a low-degree graph, and let T, T' be internal triangles such that $T \cap T' = \{x, y\}$. Then $N(x)$ induces a 4-cycle, the only ear of Q_T that is an inner triangle is T' , and the only normal vertices of T are x, y .*

Proof. Suppose that $T = \{x, y, u\}$, and $T' = \{x, y, u'\}$. Considering the edge $\{x, u\}$ of the inner triangle T , let $x' \in G$ such that $T_1 = \{x, u, x'\} \neq T$. Since T is a clique, we must have $x' \neq u'$. Applying the Lemma 4.2 to the inner triangle T' and T_1 , we obtain that $N(x)$ is a 4-cycle, hence $x' \sim u'$. Applying the previous steps to the edge $\{y, u\}$ of the triangle T , we obtain a vertex $y' \in G$ that is a neighbor of u, y, u' so that $N(y)$ is a 4-cycle. Since T is a clique, we have $y' \neq x'$. Now, the triangle T_1 is an ear of Q_T . If T_1 were inner, there would be a vertex z such that $\{z, u, x'\}$ is a triangle different from T_1 . The vertex z is different from y' since G is not an octahedron. Moreover $z \in N(u) = \{x, x', y, y'\}$, but this contradicts our other assumptions. Hence T_1 is not inner. Now, u is not a normal vertex since u^* is contained properly in Q_T . \square

The situation of the previous lemma is shown in Figure 3.

Lemma 4.4. *If G is a low-degree graph, there is no path (induced or not) in $K^2(G)$, that consists of three neckties.*

Proof. Let T_1, T_2 and T_3 be three inner triangles in G such that $Q_{T_1} \sim Q_{T_2} \sim Q_{T_3}$ in $K^2(G)$. From $Q_{T_1} \sim Q_{T_2}$, we have one of the two situations shown




Figure 4: Forbidden diamonds

in either Figure 2 or Figure 3. But in none of those cases it is possible to have an inner triangle T_3 , different from T_1 , and satisfying $T_3 \cap T_2 \neq \emptyset$. \square

Lemma 4.5. *Let G be a low-degree graph, where $a, b \in G$ are normal vertices and T_1, T_2 are inner triangles. Then $a^*, b^*, Q_{T_1}, Q_{T_2}$ do not induce a diamond in $K^2(G)$ of one of the forms shown in Figure 4.*

Proof. Consider the case to the left in Figure 4. Since $Q_{T_1} \sim Q_{T_2}$, we have $|T_1 \cap T_2| \neq \emptyset$. If $|T_1 \cap T_2| = 1$, we have the situation of Figure 2. Since b^* is a neighbor of both Q_{T_1} and Q_{T_2} , and b is a normal vertex, we have that $b \in T_1 \cap T_2$. It is immediate to verify that no neighbor of b is a normal vertex. Now, if $|T_1 \cap T_2| = 2$, then again our assumptions on b imply that $b \in T_1 \cap T_2$, and all neighbors x of b that are normal vertices are such that x^* is a neighbor of both Q_{T_1} and Q_{T_2} .

Now, with respect to the case on the right, since Q_{T_1} is not adjacent to Q_{T_2} in $K^2(G)$, then $T_1 \cap T_2 = \emptyset$. Since b must be a neighbor to two vertices in T_1 , to two vertices in T_2 and to the vertex a , and the degree of b is at most four, we have that $a \in T_1 \cup T_2$. Without loss, assume $a \in T_1$. By a similar argument, we must have $b \in T_2$. Let $T_1 = \{a, u, v\}$ and $T_2 = \{b, w, r\}$. Since a is a neighbor to two vertices in T_2 , without loss we may assume that $a \sim w$. Applying Lemma 4.2 to the inner triangle T_1 and the triangle $T = \{a, b, w\}$, we have that $N(a)$ is a 4-cycle, and without loss, we may assume $u \sim b$ and $v \sim w$. Similarly, applying the same lemma to T_2 and the triangle $T' = \{a, b, u\}$ we have that $u \sim r$. But then T and T' are inner triangles, and Lemma 4.3 applies. We have that T is the only ear in $Q_{T'}$ that is an inner triangle, which contradicts that T_1 is inner. \square

Lemma 4.6. *Let G be a low-degree graph, and let $a, b, x, z \in G$ be normal vertices and T an inner triangle in G forming the subgraph of $K^2(G)$ with the solid edges in Figure 5. Then one of the dashed edges has to be present.*

Proof. Suppose there is a subgraph with the solid edges of the form in Figure 5, but without the dashed edges. Then in G we have that $\{a, x, z\} \subseteq$




Figure 5: Forbidden subgraph




Figure 6: Forbidden subgraph

$N(b)$. Since $b^* \sim Q_T$ in $K^2(G)$, we have that two neighbors of b are in T , and so, at least one of $\{a, x, z\}$ is an element of T . But since $z \not\sim Q_T$, we have that $z \notin T$, and it is not possible for a, b to be both elements of T .

Suppose first that $a \in T$ (and so, $b \notin T$). Let us write $T = \{a, u, v\}$. Now, b is a neighbor of two elements of T , let us say without loss that $b \sim v$. Applying Lemma 4.2, we must then have that $z \sim u$. But this contradicts that $z \not\sim Q_T$. Consider then the case that $a \notin T$ and $x \in T$. Suppose $T = \{x, u, v\}$. If we had that $b \in \{u, v\}$, we could assume without loss that $b = v$. Since a is a neighbor of two elements of T , and $a \not\sim x$ by our hypothesis, then $a \sim u$. Applying Lemma 4.2 to the inner triangle T and the triangle $\{b, a, z\}$, we obtain that $z \sim x$, which again would mean that z is a neighbor of two elements of T . This shows that $b \notin \{u, v\}$. Then $a^* \sim Q_T$ in $K^2(G)$ implies that $a \sim u$ and $a \sim v$ in G . We also have $b^* \sim Q_T$ in $K^2(G)$, without loss, assume $b \sim v$. Then the triangle $\{a, b, v\}$ is inner, and we are in the situation of Figure 4.2, which shows that x is not a normal vertex, contradicting our hypothesis. \square

Lemma 4.7. *Let G be a low-degree graph, and let $a, b, x, z \in G$ be normal vertices and T an inner triangle in G forming the subgraph of $K^2(G)$ with the solid lines in Figure 6. Then one of the dashed lines has to be present.*

Proof. Suppose there is a subgraph with the solid edges of the form in Figure 6, but without the dashed edges. Then in G we have that the normal vertices $\{a, c, b, x\}$ induce a diamond. Since $b^* \sim Q_T$ and $\{a, c, x\} \subseteq N(b)$, at least one of $\{a, c, x\}$ is in the inner triangle T . But since $c^* \not\sim Q_T$, we have that $c \notin T$, and at most one of a, b is in T . Consider first the




Figure 7: Forbidden subgraph

case where $a \in T$, and let $T = \{a, u, v\}$. Since $c^* \not\sim Q_T$ we must have $\{u, v\} \cap \{b, c, x\} = \emptyset$. Applying Lemma 4.2, to the inner triangle T and the triangle $\{a, b, c\}$, we have that c would have to be adjacent to either of u or v , but this contradicts $c^* \not\sim Q_T$. So, let us consider the case $a \notin T$ and $x \in T$. Then $T = \{x, u, v\}$ with $\{u, v\} \cap \{a, b, c\} = \emptyset$. As before, an application of Lemma 4.2 leads to $c \sim u$, or to $c \sim v$ but this is a contradiction. \square

5 Proof of the Main Theorem

Theorem 5.1. *Let G be a low-degree graph. If the graph with the solid edges in Figure 7 appears as a subgraph of $K^2(G)$, then there is necessarily at least one of the dotted edges.*

Proof. Suppose there is a subgraph with the solid edges of the form in Figure 7, but without the dashed edges. We consider several cases:

1. Suppose first that Q_1, Q_2, Q_3 are stars, and set $Q_i = a_i^*$ for $i = 1, 2, 3$. As a subcase, consider when Q_4, Q_5, Q_6 are also stars. Then there is a subgraph of G isomorphic to the one in Figure 7, whose vertices are normal vertices. However, then the triangle $T = \{a_1, a_2, a_3\}$ is inner. Since a_1 is a normal vertex, there must be $q \in K(G)$ such that $q \in a_1^*$ but $q \notin Q_T$. Since $q \subseteq N[a_1] = \{a_2, a_3, a_4, a_6\}$, we must have $q = \{a_1, a_4, a_6\}$, and so a_4 is adjacent to a_6 . Similarly, considering that a_2 is a normal vertex we obtain that $a_4 \sim a_5$, and considering a_3 we obtain that $a_5 \sim a_6$. But then this would imply that G is an octahedron. Hence, this subcase is not possible. Now, consider the subcase when one of Q_4, Q_5, Q_6 is a necktie. Without loss, suppose that Q_4 is a necktie. By Lemma 4.7, neither Q_5 nor Q_6 are stars. Pick then inner triangles T_1, T_2, T_3 such that $Q_4 = Q_{T_3}$, $Q_5 = Q_{T_1}$ and $Q_6 = Q_{T_2}$. We claim that one of $T_1 \cap T_2$, $T_2 \cap T_3$, $T_1 \cap T_3$ must be non-empty. Suppose that $T_1 \cap T_3 = \emptyset$ and $T_2 \cap T_3 = \emptyset$. Since a_1 is a

neighbor of two vertices of T_2 and of two vertices of T_3 , and of a_2 and of a_3 , we must have that $a_2 \in T_3$ and $a_3 \in T_2$. Similarly, considering the neighbors of a_2 , we get that $a_1 \in T_3$ and $a_3 \in T_1$. This means that $T_1 \cap T_2 \neq \emptyset$. This proves our claim, so without loss, assume that $T_1 \cap T_2 \neq \emptyset$. If $|T_1 \cap T_2| = 1$, we are in the situation of Figure 2. In this case, one can verify that the only normal vertex such that its star is a neighbor in $K^2(G)$ of both Q_{T_1} and Q_{T_2} is the vertex in $T_1 \cap T_2$, then $T_1 \cap T_2 = \{a_3\}$. But then none of the neighbors of a_3 is a normal vertex. So we are left with the possibility that $|T_1 \cap T_2| = 2$, that is, the situation of Figure 3. But we obtain that this subcase is not possible either, since, given that a_3 is a normal vertex with $a_3^* \sim Q_{T_1}$ and $a_3^* \sim Q_{T_2}$, it must happen that $a_3 \in T_1 \cap T_2$. But then a_3 cannot have two different neighbors that are normal vertices.

2. Suppose now that Q_1 is a necktie, and Q_2, Q_3 are stars. Considering the graph to the right of Figure 4, we obtain that Q_5 must be a star. Considering the graph to the left of Figure 4, we obtain that Q_6 must also be a star. But by Lemma 4.6, this is not possible.
3. Suppose now that Q_1, Q_2 are neckties, and Q_3 is a star. By Lemma 4.4, Q_5 cannot be a necktie. But by the graph to the left of Figure 4, Q_5 cannot be a star either.
4. The case when Q_1, Q_2, Q_3 are all neckties, is also impossible by Lemma 4.4.

This finishes the proof of the Main Theorem. \square

Corollary 5.2. *If G is a connected, divergent graph, that is different from the octahedron, then G has a vertex with degree at least five.*

References

- [1] M. Baumeister and A. M. Limbach. Clique dynamics of locally cyclic graphs with $\delta \geq 6$, 2021. Preprint at arXiv.
- [2] C. Cedillo and M. A. Pizaña. Clique-convergence is undecidable for automatic graphs. *Journal of Graph Theory*, 96(3):414–437, 2021.
- [3] F. Escalante. Über iterierte Clique-Graphen. *Abh. Math. Sem. Univ. Hamburg*, 39:59–68, 1973.

- [4] M. E. Frías-Armenta, F. Larrión, V. Neumann-Lara, and M. A. Pizaña. Edge contraction and edge removal on iterated clique graphs. *Discrete Appl. Math.*, 161(10-11):1427–1439, 2013.
- [5] F. Harary. *Graph theory*. Addison-Wesley Publishing Co., Reading, Mass.-Menlo Park, Calif.-London, 1969.
- [6] F. Larrión, C. P. de Mello, A. Morgana, V. Neumann-Lara, and M. A. Pizaña. The clique operator on cographs and serial graphs. *Discrete Math.*, 282(1-3):183–191, 2004.
- [7] F. Larrión and V. Neumann-Lara. Clique divergent graphs with unbounded sequence of diameters. *Discrete Math.*, 197/198:491–501, 1999. 16th British Combinatorial Conference (London, 1997).
- [8] F. Larrión and V. Neumann-Lara. Locally C_6 graphs are clique divergent. *Discrete Math.*, 215(1-3):159–170, 2000.
- [9] F. Larrión, V. Neumann-Lara, and M. A. Pizaña. On expansive graphs. *European J. Combin.*, 30(2):372–379, 2009.
- [10] F. Larrión, V. Neumann-Lara, and M. A. Pizaña. Whitney triangulations, local girth and iterated clique graphs. *Discrete Math.*, 258(1-3):123–135, 2002.
- [11] F. Larrión, V. Neumann-Lara, and M. A. Pizaña. Graph relations, clique divergence and surface triangulations. *J. Graph Theory*, 51(2):110–122, 2006.
- [12] F. Larrión and M. A. Pizaña. On hereditary clique-Helly self-clique graphs. *Discrete Appl. Math.*, 156(7):1157–1167, 2008.
- [13] F. Larrión, M. A. Pizaña, and R. Villarreal-Flores. The clique operator on matching and chessboard graphs. *Discrete Math.*, 309(1):85–93, 2009.
- [14] M. Lin, F. J. Souignac, and J. L. Szwarcfiter. The clique operator on circular-arc graphs. *Discrete Appl. Math.*, 158(12):1259–1267, 2010.
- [15] V. Neumann-Lara. On clique-divergent graphs. Problèmes combinatoires et théorie des graphes, Orsay 1976, Colloq. int. CNRS No. 260, 313-315 (1978)., 1978.
- [16] E. Prisner. Hereditary clique-Helly graphs. *J. Combin. Math. Combin. Comput.*, 14:216–220, 1993.

- [17] E. Prisner. *Graph dynamics*, volume 338 of *Pitman Research Notes in Mathematics Series*. Longman, Harlow, 1995.