# *N*-tuple wights noncommutative Orlicz spaces and some geometrical properties \*

Ma Zhenhua<sup>a,\*</sup>, Deng Quancai<sup>a</sup>

<sup>a</sup>Hebei University of Architecture, Zhangjiakou, 075024, P. R. China

# Abstract

This paper studies the *N*-tuple noncommutative Orlicz spaces  $\bigoplus_{j=1}^{n} L_{p,\lambda}^{(\Phi_j)}(\widetilde{\mathcal{M}}, \tau)$ , where  $L^{(\Phi_j)}(\widetilde{\mathcal{M}}, \tau)$  is noncommutative Orlicz spaces and  $\widetilde{\mathcal{M}}$  is the  $\tau$ -measurable operators. Based on the maximum principle, we give the Riesz-Thorin interpolation theorem on  $\bigoplus_{j=1}^{n} L_{p,\lambda}^{(\Phi_j)}(\widetilde{\mathcal{M}}, \tau)$ . As applications, the Clarkson inequality and some geometrical properties such as uniform convexity and unform smooth of noncommutative Orlicz space  $L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau), 0 < s \leq 1$  are given.

Keywords: Noncommutative Orlicz spaces,  $\tau$ -measurable operator, von Neumann algebra, Orlicz function, Riesz-Thorin interpolation 2010 MSC: 46L52, 47L10, 46A80

# 1. Preliminaries

In 1936, for study uniform convexity of  $L^p$  space, Clarkson gave some famous inequalities named Clarkson inequality [3]. In [10], the author used the noncommutative Riesz-Thorin interpolation theorem get the Clarkson inequality on noncommutative  $L^p$  space. The principal objective of this paper is to investigate Riesz-Thorin interpolation theorem on noncommutative Orlicz spaces which yields the Clarkson inequality of noncommutative  $L^p$  space. As applications, some geometrical properties such as uniform convexity and unform smooth of noncommutative Orlicz space  $L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau), 0 < s \leq 1$  are given.

<sup>\*</sup>The research has been supported by Research project of basic scientifific research business expenses of provincial colleges and universities in Hebei Province(2021QNJS11); Innovation and improvement project of academic team of Hebei University of Architecture Mathematics and Applied Mathematics (NO. TD202006); The Major Project of Education Department in Hebei (No. ZD2021039); Nature Science Foundation of Heibei Province under (No. A2019404009; China Postdoctoral Science Foundation (No. 2019M661047);Postdoctoral Foundation of Heibei Province under Grant B2019003016.

<sup>\*</sup>Corresponding author

*Email addresses:* mazhenghua\_1981@163.com (Ma Zhenhua ), dqc\_com@163.com (Deng Quancai)

- The theory of Orlicz spaces associated to a trace was introduced by Muratov [6] and Kunze [12]. Let  $\mathcal{M}$  be a semi-finite von Neumann algebra acting on a Hilbert space  $\mathcal{H}$  with a normal semi-finite faithful trace  $\tau$ . A densely-defined closed linear operator  $A : \mathcal{D}(A) \to \mathcal{H}$  with domain  $\mathcal{D}(A) \subseteq \mathcal{H}$  is called affiliated with  $\mathcal{M}$  if and only if  $U^*AU = A$  for all unitary operators U belonging to the commutant  $\mathcal{M}'$  of
- <sup>15</sup>  $\mathcal{M}$ . Clearly, if  $A \in \mathcal{M}$  then A is affiliated with  $\mathcal{M}$ . If A is a (densely-defined closed) operator affiliated with  $\mathcal{M}$  and A = U|A| the polar decomposition, where  $|A| = (A^*A)^{\frac{1}{2}}$  and U is a partial isometry, then A said to be  $\tau$ -measurable if and only if there exists a number  $\lambda \geq 0$  such that  $\tau(e_{(\lambda,\infty)}(|A|)) < \infty$ , where  $e_{[0,\lambda]}$  is the spectral projection of |A| and  $\tau$  is the trace of normal faithful and semifinite. The
- <sup>20</sup> collection of all  $\tau$ -measurable operators is denoted by  $\mathcal{M}$ . The spectral decomposition implies that a von Neumann algebra  $\mathcal{M}$  is generated by its projections. Recall that an element  $A \in \mathcal{M}_+$  is a linear combination of mutually orthogonal projections if  $A = \sum_{k=1}^{n} \alpha_k e_k$  with  $\alpha_k \in \mathbb{R}_+$  and projection  $e_k \in \mathcal{M}$  such that  $e_k e_j = 0$  whenever  $k \neq j$  [10].
- Next we recall the definition and some basic properties of noncommutative Orlicz spaces.

A function  $\Phi : [0, \infty) \to [0, \infty]$  is called an Orlicz function if and only if  $\Phi(u) = \int_{0}^{|u|} p(t)dt$ , where the right derivative p of  $\Phi$  satisfies p is right-continuous and nondecreasing, p(t) > 0 whenever t > 0 and p(0) = 0 with  $\lim_{t\to\infty} p(t) = \infty$  [11]. Further we say an Orlicz function  $\Phi$  satisfies the  $\Delta_2$ -condition for large t (for small t, or for all t), written often as  $\Phi \in \Delta_2$ , if there exist constants  $t_0 > 0, K > 2$  such that  $\Phi(2t) \leq K\Phi(t)$  for  $|t| \geq t_0$  [7].

If  $A \in \mathcal{M}$  and  $\Phi$  is an Orlicz function, we denote  $\tilde{\rho}_{\Phi}(A) = \tau(\Phi(|A|))$ , hence we can define a corresponding space, which is named the noncommutative Orlicz space, as follows:

$$L^{\Phi}(\widetilde{\mathcal{M}},\tau) = \{A \in \widetilde{\mathcal{M}} : \tau(\Phi(\lambda|A|)) < \infty \text{ for some } \lambda > 0\}.$$

Also we could define the subspace

$$E^{\Phi}(\widetilde{\mathcal{M}},\tau) = \{A \in \widetilde{\mathcal{M}} : \tau(\Phi(\lambda|A|)) < \infty \text{ for any } \lambda > 0\}.$$

We equip these spaces with the Luxemburg norm

$$||A||_{(\Phi)} = \inf\{\lambda > 0 : \tau\left(\Phi\left(\frac{|A|}{\lambda}\right)\right) \le 1\}.$$

In the case of  $\Phi(A) = |A|^p$ ,  $1 \le p < \infty$ ,  $L^{(\Phi)}(\widetilde{\mathcal{M}}, \tau)$  is nothing but the noncommutative space  $L^p(\widetilde{\mathcal{M}}, \tau) = \left\{ A \in \widetilde{\mathcal{M}} : \tau(|A|^p) < \infty \right\}$  [5] and the Luxemburg norm

generated by this function is expressed by the formula

$$||A||_p = (\tau(|A|^p))^{\frac{1}{p}}.$$

One can define another norm on  $L^{\Phi}(\widetilde{\mathcal{M}}, \tau)$  as follows

$$||A||_{\Phi} = \sup\{\tau(|AB|) : B \in L^{\Psi}(\widetilde{\mathcal{M}}, \tau) \text{ and } \tau(\Psi(B)) \le 1\},\$$

where  $\Psi : [0, \infty) \to [0, \infty]$  is defined by  $\Psi(u) = \sup\{uv - \Phi(v) : v \ge 0\}$ . Here we call  $\Psi$  the complementary function of  $\Phi$ . In the following, we use  $L^{(\Phi)}(\widetilde{\mathcal{M}}, \tau)$  and  $L^{\Phi}(\widetilde{\mathcal{M}}, \tau)$  denote the Orlicz which equipped Luxemberg and Orlicz norm respectively.

The same as  $E^{(\Phi)}(\widetilde{\mathcal{M}},\tau)$  and  $E^{\Phi}(\widetilde{\mathcal{M}},\tau)$ .

For more information on the theory of noncommutative Orlicz spaces we refer the reader to [8, 9, 2, 12, 5, 6].

## 2. Riesz-Thorin interpolation theorem of Noncommutative Orlicz spaces

40

35

In this section, we will give the definition of N-tuple noncommutative Orlicz spaces, also give some norm inequalities. For research the Riesz-Thorin interpolation theorem, a equivalent definition of Luxmburg norm must be given. As a corollary, the Clarkson inequality of noncommutive  $L^p$  space could be get. The main ideas and proof ideas in this article are derived from literatures [10] and [7].

Now let  $\mathcal{N} = \mathcal{M} \oplus \mathcal{M} \oplus \cdots \oplus \mathcal{M}$  be the *n*-th von Neumann algebra direct sum of  $\mathcal{M}$  with it self. We know that  $\mathcal{N}$  acts on the direct sum Hilbert space  $\mathcal{H} \oplus \mathcal{H} \oplus \cdots \oplus \mathcal{H}$  coordinatewise:

$$(A_1, A_2, \dots, A_n)(x_1, x_2, \dots, x_n) = \sum_{j=1}^n A_j x_n,$$

<sup>45</sup> where  $A_j \in \mathcal{M}, i = 1, 2, \dots n$ . Then  $\mathcal{N}_+ = \mathcal{M}_+ \oplus \mathcal{M}_+ \oplus \dots \oplus \mathcal{M}_+$ .

Define:  $v : \mathcal{N}_+ \to \mathbb{C}$  by  $v(A_1, A_2, \ldots, A_n) = \sum_{j=1}^n \lambda_j \tau(A_j)$ , where  $\lambda_j \ge 0$  and  $\tau$  is a normal faithful normal faithful normalized trace on  $\mathcal{M}$ , then v is a normal faithful normal faithful normalized trace on  $\mathcal{N}$ . Now we give the following definition:

**Definition 2.1.** Let  $\Phi = (\Phi_1, \Phi_2, \dots, \Phi_n)$  be an n-tuple of N functions  $\Phi_j$ . For each  $p \ge 1, \lambda_j \ge 0$ , and n-tuple of wights  $\lambda = (\lambda_1, \dots, \lambda_n)$  consider the direct sum space, named n-tuple of wights noncommutative Orlicz spaces as follows:

$$\bigoplus_{j=1}^{n} E_{p,\lambda}^{(\Phi_j)} = \{ A = (A_1, A_2, \dots, A_n) : A_j \in E^{(\Phi_j)}(\widetilde{\mathcal{M}}, \tau), 1 \le j \le n \}$$

with norm  $\|\cdot\|_{(\Phi),p,\lambda}$  defined for  $A_j \in E^{(\Phi_j)}(\widetilde{\mathcal{M}},\tau)$ :

50

$$|A||_{(\Phi),p,\lambda} = \begin{cases} \left[\sum_{j=1}^{n} \lambda_{j} ||A_{j}||_{(\Phi_{j})}^{p}\right]^{\frac{1}{p}}, & 1 \le p < \infty, \\ \max_{j} ||A_{j}||_{(\Phi_{j})}, & p = \infty. \end{cases}$$

or the norm  $\|\cdot\|_{\Phi,p,\lambda}$  defined in the same way as before in which  $\|\cdot\|_{(\Phi_j)}$  is replaced by the Orlicz norm  $\|\cdot\|_{\Phi_j}$ , denotes by  $\bigoplus_{j=1}^n E_{p,\lambda}^{\Phi_j}$ . The same way, if  $\Psi_j$  is the complementary

N-function of  $\Phi_j$ , denotes by  $\bigoplus_{j=1}^n E_{q,\lambda}^{(\Psi_j)}$  which equip with  $\|\cdot\|_{(\Psi),q,\lambda}$  and  $\bigoplus_{j=1}^n E_{q,\lambda}^{\Psi_j}$  which equip with  $\|\cdot\|_{\Psi,q,\lambda}$  for the same weights  $\lambda = (\lambda_1, \ldots, \lambda_n)$  and  $q = \frac{p}{p-1}$ . The same way, we also could define  $\bigoplus_{j=1}^n L_{p,\lambda}^{(\Phi_j)}$ ,  $\bigoplus_{j=1}^n L_{q,\lambda}^{\Phi_j}$ ,  $\bigoplus_{j=1}^n L_{q,\lambda}^{(\Psi_j)}$ ,  $\bigoplus_{j=1}^n L_{q,\lambda}^{(\Psi_j)}$ , and the norm as before.

**Remark 1.** By the Theorem 3.4 of [5], if for any  $1 \le j \le n$  with  $\Phi_j \in \Delta_2$ , we have <sup>55</sup>  $\bigoplus_{j=1}^{n} L_{p,\lambda}^{(\Phi_j)} = \bigoplus_{j=1}^{n} E_{p,\lambda}^{(\Phi_j)}$ .

Lemma 2.1. If 
$$A \in \bigoplus_{j=1}^{n} L_{p,\lambda}^{(\Phi_{j})}$$
 and  $B \in \bigoplus_{j=1}^{n} L_{q,\lambda}^{\Psi_{j}}$ , where  $1 \leq p < \infty$ , we have  
(1) If  $||A||_{(\Phi),p,\lambda} \leq 1$ , then we have  $v(\Phi(A)) \leq ||A||_{(\Phi),p,\lambda} \cdot \delta_{1}$ , where  $\delta_{1} = \left(\sum_{j=1}^{n} \lambda_{j}\right)^{\frac{1}{q}}$ .  
(2) If  $||A||_{(\Phi),p,\lambda} > 1$ , then we have  $v(\Phi(A)) > \delta_{2}$ , where  $\delta_{2} = \left[\sum_{j=1}^{n} \lambda_{j}^{p} ||A_{j}||_{(\Phi_{j})}^{p}\right]^{\frac{1}{p}}$ .  
(3)  $v(AB) \leq ||A||_{(\Phi),p,\lambda} \cdot ||B||_{\Psi,q,\lambda}$ .

<sup>60</sup> Proof. (1) If  $||A||_{(\Phi),p,\lambda} \leq 1$ , then from Proposition 3.4 of [2] and classical Hölder inequality, we have

$$\begin{aligned}
\upsilon(\Phi(A)) &= \sum_{j=1}^{n} \lambda_j \tau(\Phi_j(A_j)) \\
&= \sum_{j=1}^{n} \lambda_j^{\frac{1}{q}} \cdot \lambda_j^{\frac{1}{p}} \tau(\Phi_j(A_j)) \\
&\leq \left[ \sum_{j=1}^{n} \lambda_j \left[ \tau(\Phi_j(A_j)) \right)^p \right]^{\frac{1}{p}} \cdot \left( \sum_{j=1}^{n} \lambda_j \right)^{\frac{1}{q}} \\
&\leq \left[ \sum_{j=1}^{n} \lambda_j \|A_j\|_{(\Phi_j)}^p \right]^{\frac{1}{p}} \cdot \delta_1 \\
&= \|A\|_{(\Phi), p, \lambda} \cdot \delta_1.
\end{aligned}$$

(2) If  $||A||_{(\Phi),p,\lambda} > 1$ , then from Proposition 3.4 of [2], we have

$$[\upsilon(\Phi(A))]^{p} = \left[\sum_{j=1}^{n} \lambda_{j} \tau(\Phi_{j}(A_{j}))\right]^{p}$$
$$> \left[\sum_{j=1}^{n} \lambda_{j} \|A_{j}\|_{(\Phi_{j})}\right]^{p}$$
$$\geq \sum_{j=1}^{n} \lambda_{j}^{p} \|A_{j}\|_{(\Phi_{j})}^{p},$$

which means that

$$\upsilon(\Phi(A)) > \left[\sum_{j=1}^n \lambda_j^p \|A_j\|_{(\Phi_j)}^p\right]^{\frac{1}{p}} = \delta_2.$$

(3) From Theorem 3.3 of [2] and classical Hölder inequality, we get that

$$\begin{aligned}
\upsilon(AB) &= \sum_{j=1}^{n} \lambda_j |\tau(A_j B_j)| \\
&\leq \sum_{j=1}^{n} \lambda_j^{\frac{1}{p} + \frac{1}{q}} ||A_j||_{(\Phi)} ||B_j||_{\Psi} \\
&\leq \left( \sum_{j=1}^{n} \lambda_j ||A_j||_{(\Phi)}^p \right)^{\frac{1}{p}} \cdot \left( \sum_{j=1}^{n} \lambda_j ||B_j||_{\Psi}^q \right)^{\frac{1}{q}} \\
&= ||A||_{(\Phi), p, \lambda} \cdot ||B||_{\Psi, q, \lambda},
\end{aligned}$$

Remark 2. If  $\Phi$  is 1-tuple N-function and  $\lambda = 1$ , the Lemma 2.1 just be the Theorem 3.3 and Proposition 3.4 of [2].

**Theorem 2.1.** If  $A \in \bigoplus_{j=1}^{n} E_{p,\lambda}^{(\Phi_j)}$ , then for  $1 \leq p < \infty$ , the weighted norms  $\|\cdot\|_{(\Phi),p,\lambda}$  is given by

$$||A||_{(\Phi),p,\lambda} = \sup \{ v(AB) : ||B||_{\Psi,q,\lambda} \le 1 \},\$$

*Proof.* If  $||B||_{\Psi,q,\lambda} \leq 1$ . One side, by (3) of the Lemma 2.1, we have

$$\upsilon(AB) \le \|A\|_{(\Phi),p,\lambda} \cdot \|B\|_{\Psi,q,\lambda} \le \|A\|_{(\Phi),p,\lambda}$$

The other side, for simplicity, we may take that  $||A||_{(\Phi),p,\lambda} = 1$  and assume that  $A_j \ge 0$ . Let  $\{e_{jn}\}$  be the projection of  $A_j$  and  $0 < \tau(e_{jn}) < \infty$ . By Proposition 3.4 of [2], for any  $\varepsilon > 0$ , we have  $\tau [\Phi_j ((1 + \varepsilon)A_j)] \ge ||(1 + \varepsilon)A_j||_{(\Phi_j)} = 1 + \varepsilon$ .

If we define the operator  $A_{jm} = A_j(e_{j1} + e_{j2} + \dots + e_{jm})(m \le n)$ , where  $A_j = \sum_{k=1}^{n} \alpha_k e_{jk}$  and  $e_{jk} = 0, k > n$ , then  $A_{jm} \uparrow A_j$  as  $m \to \infty$ , there exists an  $m_0$  such that for  $m \ge m_0$  one have

$$\upsilon\left[\frac{1}{\delta_2}\Phi\left((1+\varepsilon)A_m\right)\right] = \sum_{k=1}^n \frac{1}{\delta_2}\lambda_j\Phi_j\left((1+\varepsilon)A_{jm}\right) \ge \left(1+\frac{\varepsilon}{2}\right).$$

If we set

$$B_{jm} = \frac{\delta_2^{-1} p((1+\varepsilon)\lambda_j A_{jm})}{\delta_1 \left(1 + \tau(\Psi_j(\delta_2^{-1} p((1+\varepsilon)\lambda_j A_{jm}))))\right)},$$

then  $B_{jm}$  is bounded operators and  $B_m \in \bigoplus_{j=1}^n E_{q,\lambda}^{\Psi_j}$  for each m. Moreover by definition 1.7 of [5] and 1.9 of [11] we have,  $\|B_{jm}\|_{\Psi_j,q,\lambda} \leq 1$  since  $\|A\|_{(\Phi),p,\lambda} = 1$ . Hence,

$$||B_m||_{\Psi,q,\lambda} = \left(\sum_{j=1}^n \lambda_j ||B_{jm}||_{\Psi_j}^q\right)^{\frac{1}{q}} \le 1.$$

However, one has

$$\sup\{v(AB)\} = \sup\left\{\sum_{j=1}^{n} \lambda_{j}\tau(A_{j}B_{j}) : B_{j} \in E^{\Psi_{j}}, \|B\|_{\Psi,q,\lambda} \leq 1\right\}$$

$$\geq \sup_{m \geq m_{0}} \left\{\sum_{j=1}^{n} \lambda_{j}\tau(A_{j}B_{jm}) : B_{jm} \in E^{\Psi_{j}}, \|B_{m}\|_{\Psi,q,\lambda} \leq 1\right\}$$

$$\geq \frac{1}{1+\varepsilon} \sup_{m \geq m_{0}} \left\{\sum_{j=1}^{n} \tau((1+\varepsilon)\lambda_{j}A_{jm}B_{jm})\right\}$$

$$= \frac{1}{1+\varepsilon} \sup_{m \geq m_{0}} \left\{\sum_{j=1}^{n} \frac{\tau(\Phi_{j}(1+\varepsilon)\lambda_{j}A_{jm}) + \tau(\Psi_{j}(\delta_{2}^{-1}p(1+\varepsilon)\lambda_{j}A_{jm})))}{\delta_{1}\delta_{2}\left(1 + \tau(\Psi_{j}(\delta_{2}^{-1}p(1+\varepsilon)\lambda_{j}A_{jm}))\right)}\right\}$$

$$\geq \frac{1}{1+\varepsilon},$$

since  $\varepsilon > 0$  is arbitrary we get the desired inequality.

**Definition 2.2.** [1] Let  $\Phi_1$  and  $\Phi_2$  be N-functions and define  $\Phi_s$  to be the inverse of  $\Phi_s^{-1}(u) = [\Phi_1^{-1}(u)]^{1-s} [\Phi_2^{-1}(u)]^s$  for  $0 \le s \le 1, u \ge 0$ , where  $\Phi^{-1}$  is the unique inverse of the N-function  $\Phi$ .

**Theorem 2.2.** Let  $\Phi_i = (\Phi_{i1}, \Phi_{i2}, \dots, \Phi_{in}), Q_i = (Q_{i1}, Q_{i2}, \dots, Q_{in}), i = 1, 2$  be ntuples of N-functions and  $0 \leq r_1, r_2, t_1, t_2 \leq \infty, \lambda = (\lambda_1, \dots, \lambda_n)$  be given positive numbers. Next let  $\Phi_s = (\Phi_{s1}, \Phi_{s2}, \dots, \Phi_{sn}), Q_s = (Q_{s1}, Q_{s2}, \dots, Q_{sn})$  be the associated

intermediate N-functions,

$$\frac{1}{r_s} = \frac{1-s}{r_1} + \frac{s}{r_2}, \frac{1}{t_s} = \frac{1-s}{t_1} + \frac{s}{t_2}, 0 \le s \le 1.$$

$$\begin{split} If \ T : \bigoplus_{j=1}^{n} E_{r_{i},\lambda}^{(\Phi_{ij})} &\to \bigoplus_{j=1}^{n} L_{t_{i},\lambda}^{(Q_{ij})} \ is \ a \ bounded \ linear \ operator \ with \ bounds \ K_{1}, K_{2}, \ such \\ that \ \|TA\|_{(Q_{i}),t_{i},\lambda} &\leq K_{i} \|A\|_{(\Phi_{i}),r_{i},\lambda}, A \in \bigoplus_{j=1}^{n} E_{r_{i},\lambda}^{(\Phi_{ij})}, i = 1, 2. \\ Then \ T \ is \ also \ defined \ on \ \bigoplus_{j=1}^{n} E_{r_{s},\lambda}^{(\Phi_{sj})} \ into \ \bigoplus_{j=1}^{n} L_{r_{s},\lambda}^{(\Phi_{sj})} \ for \ all \ 0 \leq s \leq 1 \ and \ one \ have \\ the \ bound \\ \|TA\|_{(Q_{s}),t_{s},\lambda} &\leq K_{1}^{1-s}K_{2}^{s} \|A\|_{(\Phi_{s}),r_{s},\lambda}, \end{split}$$

where  $A \in \bigoplus_{j=1}^{n} E_{r_s,\lambda}^{(\Phi_{sj})}$ .

Proof. Let  $A = (A_1, A_2, \dots, A_n) \in \bigoplus_{j=1}^n E_{r_s,\lambda}^{(\Phi_{sj})}, B = (B_1, B_2, \dots, B_n) \in \bigoplus_{j=1}^n E_{t_s,\lambda}^{\Psi_{tj}}$  with polar decompositions  $A_k = U_k |A_k|, B_k = V_k |B_k|$ . Assume that  $||A||_{(\Phi_i), r_s, \lambda} \leq 1, ||B||_{\Psi_i, t_s, \lambda} \leq 1$ where  $|A_k| = \sum_{j=1}^n \alpha_j e_{kj}, |B_k| = \sum_{j=1}^n \beta_j e'_{kj}$ . Define for  $z = \mathbb{C}$  and  $k = 1, 2, \dots, n$ 

$$A(z) = (A_1(z), A_2(z), \dots, A_n(z))$$

and

$$B(z) = (B_1(z), B_2(z), \dots, B_n(z)),$$

where

$$A_k(z) = U_k \Phi_{sk} \left[ (\Phi_{1k}^{-1})^{1-z} (\Phi_{2k}^{-1})^z \right] (|A_k|),$$
  
$$B_k(z) = V_k \Psi_{sk} \left[ (\Psi_{1k}^{-1})^{1-z} (\Psi_{2k}^{-1})^z \right] (|B_k|).$$

Then,

$$A_{k}(z) = U_{k}\Phi_{sk} \left[ \left( \Phi_{1k}^{-1} \left( \sum_{j=1}^{n} \alpha_{j} e_{kj} \right) \right)^{1-z} \left( \Phi_{2k}^{-1} \left( \sum_{j=1}^{n} \alpha_{j} e_{kj} \right) \right)^{z} \right] \\ = \sum_{j=1}^{n} \Phi_{sk} \left[ \left( \Phi_{1k}^{-1}(\alpha_{j}) \right)^{1-z} \left( \Phi_{2k}^{-1}(\alpha_{j}) \right)^{z} \right] U_{k} e_{kj}$$

Hence,  $z \to A(z)$  is an analytic function on  $\mathbb{C}$  with value in  $\widetilde{\mathcal{M}}$ . The same reduction applies to B.

Now we could define a bounded entire function

$$H(z) = K_1^{z-1} K_2^{-z} \tau(B(z)TA(z)).$$

If z = it for  $t \in \mathbb{R}$ , we have

$$A_{k}(it) = \sum_{j=1}^{n} \Phi_{sk} \left[ \Phi_{sk}^{-1}(\alpha_{j}) \right] U_{k} e_{kj}$$

$$= \sum_{j=1}^{n} \Phi_{sk} \left[ \left( \Phi_{1k}^{-1}(\alpha_{j}) \right)^{1-it} \left( \Phi_{2k}^{-1}(\alpha_{j}) \right)^{it} \right] U_{k} e_{kj}$$

$$= \sum_{j=1}^{n} \Phi_{sk} \left[ \left( \frac{\Phi_{2k}^{-1}(\alpha_{j})}{\Phi_{1k}^{-1}(\alpha_{j})} \right)^{it} \right] U_{k} e_{kj} \cdot \sum_{j=1}^{n} \Phi_{sk} \left[ \Phi_{1k}^{-1}(\alpha_{j}) \right] U_{k} e_{kj}$$

$$= \left[ \Phi_{sk} \left( \frac{\Phi_{2k}^{-1}}{\Phi_{1k}^{-1}} (|A_{k}|) \right) \right]^{it} \cdot \Phi_{sk} \left( \Phi_{1k}^{-1} (|A_{k}|) \right).$$

Hence,

$$|A_k(it)|^2 = A_k(it)^* A_k(it) = \left[\Phi_{sk}\left(\Phi_{1k}^{-1}(|A_k|)\right)\right]^2$$

which means

$$|A_k(it)| = \Phi_{sk} \left( \Phi_{1k}^{-1}(|A_k|) \right).$$

Hence for any  $1 \leq k \leq n$  we have  $\tau(\Phi_{1k}(A_k(it))) = \tau(\Phi_{sk}(A_k))$  which implies that

$$||A_j(it)||_{(\Phi_{1j})} = ||A_j||_{(\Phi_{sj})}$$

and

$$\begin{aligned}
\upsilon(\Phi_1(A(it))) &= \sum_{j=1}^n \lambda_j \tau \left[ \Phi_{1j} \left[ \Phi_{sk} \left( \Phi_{1j}^{-1}(|A_j|) \right) \right] \right] \\
&= \lambda_1 \tau(\Phi_{s1}(|A_1|)) + \lambda_2 \tau(\Phi_{s2}(|A_2|)) + \ldots + \lambda_n \tau(\Phi_{sn}(|A_n|)) \\
&= \upsilon(\Phi_s(|A|)),
\end{aligned}$$

we get that

$$||A(it)||_{(\Phi_1), r_s, \lambda} = ||A||_{(\Phi_s), r_s, \lambda} \le 1.$$

Similar  $||B(it)||_{\Psi_1,t_s,\lambda} = ||B||_{\Psi_s,t_s,\lambda} \leq 1$ . Thus by (3) of the Lemma 2.1 and the assumption on T, we have

$$|\tau(B(it)TA(it))| \le K_1 ||B(it)||_{\Psi_1, t_s, \lambda} ||A(it)||_{(\Phi_1), r_s, \lambda} \le K_1.$$

It then follows that  $|H(it)| \leq 1$  for any  $t \in \mathbb{R}$ . In the same way, we show  $|H(1+it)| \leq 1$ 

1. Therefore, by the maximum principle, for any  $\theta \in \mathbb{C}$ , we get

$$|H(\theta)| = |K_1^{\theta-1} K_2^{-\theta} \tau(B(\theta) T A(\theta))| \le 1.$$

Hence,

$$|\tau(BTA)| \le K_1^{1-\theta} K_2^{\theta}$$

By the Theorem 2.1 we could get that

$$||TA||_{(Q_s),r_s,\lambda} \le K_1^{1-\theta} K_2^{\theta} ||A||_{(\Phi_s),r_s,\lambda}$$

**Theorem 2.3.** Let  $\Phi$  be an N-function and  $\Phi_s$  be the inverse which satisfies that  $\Phi_s^{-1}(u) = [\Phi^{-1}(u)]^{1-s} [\Phi_0^{-1}(u)]^s = [\Phi^{-1}(u)]^{1-s} u^{\frac{s}{2}}$  where  $0 < s \leq 1$  and  $\Phi_0(u) = u^2$ . If  $L^{(\Phi)}(\widetilde{\mathcal{M}}, \tau)$  is the noncommutative Orlicz space, then we have for  $A, B \in L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau)$ :

$$\left(\|A+B\|_{(\Phi_s)}^{\frac{2}{s}}+\|A-B\|_{(\Phi_s)}^{\frac{2}{s}}\right)^{\frac{s}{2}} \le 2^{\frac{s}{2}} \left(\|A\|_{(\Phi_s)}^{\frac{2}{2-s}}+\|B\|_{(\Phi_s)}^{\frac{2}{2-s}}\right)^{\frac{2-s}{2}}$$

*Proof.* Let  $\Phi_1 = (\Phi, \Phi)$  be the 2-vector of N-functions,  $\lambda = (1, 1), 1 \le r_1 \le \infty$  and set

$$\bigoplus_{j=1}^{\bullet} E_{r_1}^{(\Phi)}(\widetilde{\mathcal{M}}, \tau) = \{ (A, B) : A, B \in E^{(\Phi)}(\widetilde{\mathcal{M}}, \tau), \| (A, B) \|_{(\Phi_1), r_1} < \infty \},\$$

where

$$\|(A,B)\|_{(\Phi_1),r_1} = \begin{cases} \left[ \|A\|_{(\Phi)}^{r_1} + \|B\|_{(\Phi)}^{r_1} \right]^{\frac{1}{r_1}}, & 1 \le r_1 < \infty, \\ \max\{\|A\|_{(\Phi)}, \|B\|_{(\Phi)}\}, & r_1 = \infty. \end{cases}$$

Take  $Q_1 = \Phi_1 = (\Phi, \Phi)$  and  $Q_2 = \Phi_2 = (\Phi_0, \Phi_0)$  where  $\Phi_0(u) = u^2$ .

90

Set  $r_1 = 1, r_2 = t_2 = 2$  and  $t_1 = +\infty$ . Define the linear operator  $T : \bigoplus_{j=1}^2 E_{r_i}^{(\Phi_i)} \to C_{r_j}^2$ 

 $\bigoplus_{j=1}^{2} L_{t_i}^{(Q_i)}$  by the equation T(A, B) = (A + B, A - B), we then have

$$\|T(A,B)\|_{(Q_1),t_1} = \max\{\|A+B\|_{(\Phi)}, \|A-B\|_{(\Phi)}\} \\ \leq \|A\|_{(\Phi)} + \|B\|_{(\Phi)} \\ = K_1\|(A,B)\|_{(\Phi_1),r_1}.$$

Hence,  $K_1 = 1$  and since  $\|\cdot\|_{(\Phi_0)} = \|\cdot\|_2$ , we find

$$\begin{aligned} \|T(A,B)\|_{(Q_2),t_2} &= \left[ \|A+B\|_2^2 + \|A-B\|_2^2 \right]^{\frac{1}{2}} \\ &= \sqrt{2} \left[ \|A\|_2^2 + \|B\|_2^2 \right]^{\frac{1}{2}} \\ &= K_2 \|(A,B)\|_{(\Phi_2),r_2}. \end{aligned}$$

Thus  $K_2 = \sqrt{2}$ . Let  $r_s$  and  $t_s$  be given by

$$\frac{1}{r_s} = \frac{1-s}{r_1} + \frac{s}{r_2}, \frac{1}{t_s} = \frac{1-s}{t_1} + \frac{s}{t_2}$$

then we have,  $r_s = \frac{2}{2-s}, t_s = \frac{2}{s}$ .

By the results of Theorem 2.2,

$$||T(A,B)||_{(Q_s),t_s} \le 2^{\frac{s}{s}} ||(A,B)||_{(\Phi_s),r_s}$$

since  $K_1^{1-s}K_2^s = 2^{\frac{s}{2}}$ . Hence, we have

$$\|(A,B)\|_{(Q_s),r_s} = \left[ \|A\|_{(\Phi_s)}^{\frac{2}{2-s}} + \|B\|_{(\Phi_s)}^{\frac{2}{2-s}} \right]^{\frac{2-s}{2}}$$

and

95

$$||T(A,B)||_{(Q_s),t_s} = \left(||A+B||_{(\Phi_s)}^{\frac{2}{s}} + ||A-B||_{(\Phi_s)}^{\frac{2}{s}}\right)^{\frac{s}{2}}$$

which we could get the result.

The following corollary is Clarkson inequality of noncommutative  $L^p$  space and proof the process is completely similar to the P42 of [7].

**Corollary 2.1.** Suppose that  $1 and <math>q = \frac{p}{p-1}$ . Then for  $A, B \in L^p(\widetilde{\mathcal{M}}, \tau)$ , we have

$$\left(\|A+B\|_{p}^{q}+\|A-B\|_{p}^{q}\right)^{\frac{1}{q}} \leq 2^{\frac{1}{q}} \left(\|A\|_{p}^{p}+\|B\|_{p}^{p}\right)^{\frac{1}{p}}, 1$$

and

$$\left(\|A+B\|_{p}^{p}+\|A-B\|_{p}^{p}\right)^{\frac{1}{p}} \leq 2^{\frac{1}{p}} \left(\|A\|_{p}^{q}+\|B\|_{p}^{q}\right)^{\frac{1}{q}}, 2 \leq p \leq \infty.$$

*Proof.* If  $1 , let <math>1 < \alpha < p \le 2$  and  $\Phi(u) = |u|^{\alpha}, \Phi_0(u) = |u|^2, s = \frac{2(p-\alpha)}{p(2-\alpha)}$ . Then  $0 < s \le 1$  and  $\Phi_s^{-1}(u) = |u|^{\frac{1}{p}}$  or  $\Phi_s(u) = |u|^p$ . Hence  $\|\cdot\|_{(\Phi_s)} = \|\cdot\|_{(p)}$  and since  $\lim_{\alpha \downarrow 1} \frac{2}{s} = \frac{p}{p-1} = q$ ;  $\lim_{\alpha \downarrow 1} \frac{2-s}{2} = \frac{1}{p}$  by the Theorem 2.3 we get the first inequality.

100

Similar let  $2 \leq p < \beta < \infty$  and  $\Phi(u) = |u|^{\beta}, \Phi_0(u) = |u|^2, s = \frac{2(\beta-p)}{p(\beta-2)}$ . Then  $0 \leq s \leq 1$  and  $\Phi_s(u) = |u|^p, \lim_{\beta \uparrow \infty} \frac{2}{s} = p; \lim_{\beta \uparrow \infty} \frac{2-s}{2} = \frac{1}{q}$ , by the Theorem 2.3 we get the second inequality.

#### 3. Some geometrical properties

105

This section we contains some geometrical properties of noncommutative Orlicz spaces. These include uniform convexity, uniform smoothness which generalize the results of noncommutative  $L^p$  spaces. All these properties are based on Clarkson inequalities.

**Definition 3.1.** [4] Let X be a Banach space. We define its modulus of convexity by

$$\delta_X(\varepsilon) = \inf\left\{1 - \left\|\frac{x+y}{2}\right\| : x, y \in X, \|x\| = \|y\| = 1, \|x-y\| = \varepsilon\right\}, 0 < \varepsilon < 2$$

and its modulus of smoothness by

$$\rho_X(t) = \sup\left\{\frac{\|x + ty\| + \|x - ty\|}{2} - 1 : x, y \in X, \|x\| = \|y\| = 1\right\}, t > 0.$$

X is said to be uniformly convex if  $\delta_X(\varepsilon) > 0$  for every  $2 \ge \varepsilon > 0$ , and uniformly smooth if  $\lim_{t \to 0} \frac{\rho_X(t)}{t} = 0$ .

**Theorem 3.1.** Let  $\Phi$  be an N-function and  $\Phi_s$  be the inverse which satisfies that  $\Phi_s^{-1}(u) = [\Phi^{-1}(u)]^{1-s} [\Phi_0^{-1}(u)]^s = [\Phi^{-1}(u)]^{1-s} u^{\frac{s}{2}}$  where  $0 < s \leq 1$  and  $\Phi_0(u) = u^2$ , then we have for  $0 < \varepsilon \leq 2$ ,

$$\delta_{L^{(\Phi_s)}}(\varepsilon) \ge 1 - \frac{1}{2} \left[ 2^{\frac{2}{s}} - \varepsilon^{\frac{2}{s}} \right]^{\frac{s}{2}}$$

and

$$\rho_{L(\Phi_s)(t)} \le \left(1 + t^{\frac{2}{2-s}}\right)^{\frac{2-s}{2}} - 1.$$

*Proof.* First, if  $||A - B||_{(\Phi_s)} = \varepsilon$ , then theorem 2.3 implies for  $A, B \in L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau)$ ,

$$\left(\|A+B\|_{(\Phi_s)}^{\frac{2}{s}} + \varepsilon^{\frac{2}{s}}\right)^{\frac{s}{2}} \le 2^{\frac{s}{2}} \cdot 2^{\frac{2-s}{2}} = 2$$

Hence,

$$1 - \frac{1}{2} \|A + B\|_{(\Phi_s)} \ge 1 - \frac{1}{2} \left[ 2^{\frac{2}{s}} - \varepsilon^{\frac{2}{s}} \right]^{\frac{s}{2}}.$$

Taking infimum of  $||A||_{(\Phi_s)} = ||B||_{(\Phi_s)} = 1$  we can get the desired result and  $L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau)$  is uniform convexity if  $0 < \varepsilon \leq 2$ , and reflexive.

Second, if  $||A||_{(\Phi_s)} = ||B||_{(\Phi_s)} = 1$ , then since  $\frac{2}{s} \ge 2$ ,

$$\begin{split} \left[ \frac{1}{2} \left( \|A + tB\|_{(\Phi_s)} + \|A - tB\|_{(\Phi_s)} \right) \right]^{\frac{2}{s}} &\leq \frac{1}{2} \left[ \|A + tB\|_{(\Phi_s)}^{\frac{2}{s}} + \|A - tB\|_{(\Phi_s)}^{\frac{2}{s}} \right] \\ &\leq \frac{1}{2} \left[ 2^{\frac{s}{2}} \left( \|A\|_{(\Phi_s)}^{\frac{2}{2-s}} + \|tB\|_{(\Phi_s)}^{\frac{2}{2-s}} \right)^{\frac{2-s}{2}} \right]^{\frac{2}{s}} \\ &= \frac{1}{2} \left[ 2^{\frac{s}{2}} \left( 1 + t^{\frac{2}{2-s}} \right)^{\frac{2-s}{2}} \right]^{\frac{2}{s}} \\ &= \left( 1 + t^{\frac{2}{2-s}} \right)^{\frac{2-s}{s}} . \end{split}$$

Hence,

$$\frac{1}{2} \left( \|A + tB\|_{(\Phi_s)} + \|A - tB\|_{(\Phi_s)} \right) - 1 \le \left( 1 + t^{\frac{2}{2-s}} \right)^{\frac{2-s}{2}} - 1$$

Taking the supremum on the left we can get the conclusion. Since t > 0, we have that  $L^{(\Phi_s)}(\widetilde{\mathcal{M}}, \tau)$  is uniformly smooth.

From corollary 2.1, we can easily get the following results which appeared on [10]. **Corollary 3.1.** Suppose that 1 and <math>t > 0. Then for  $A, B \in L^p(\widetilde{\mathcal{M}}, \tau)$ , we have (1) If 1 , then

$$\delta_{L^p}(\varepsilon) \ge \frac{\varepsilon^q}{q \cdot 2^q} \quad and \quad \rho_{L^p(t)} \le \frac{t^p}{p}$$

(2) If 2 , then

$$\delta_{L^p}(\varepsilon) \ge \frac{\varepsilon^p}{p \cdot 2^p} \quad and \quad \rho_{L^p(t)} \le \frac{t^q}{q}$$

(3)  $L^p(\widetilde{\mathcal{M}}, \tau)$  is uniformly convex and uniformly smooth. Consequently its reflexive.

## 120 Acknowledgement

We want to express our gratitude to the referee for all his/her careful revision and suggestions which has improved the final version of this work.

# References

## References

<sup>125</sup> [1] C. E. Cleaver, On the extension of Lipschitz-Hlder maps on Orlicz spaces[J]. Studia Mathematica, 1972, 42(3).

- [2] G. Ghadir, Non-commutative Orlicz spaces associated to a modular on  $\tau$ -measurable operators, J. Math. Anal. Appl. 395 (2012), 705-715.
- [3] J. A. Clarkson, Uniformly Convex Spaces[J]. Transactions of the American Mathematical Society, 1936, 40(3):396-414.
- [4] J. Lindenstrauss and L. Tzafriri. Classical Banach spaces. II. Springer-Verlag, Berlin, 1979.
- [5] L. N. Jiang, Z. H. Ma, Closed subspaces and some basic topological properties of noncommutative Orlicz spaces, Proceedings Mathematical Sciences, 127(3) (2017), 525-536.
- [6] M. Muratov, Noncommutative Orlicz spaces, Dokl. Akad. Nauk UzSSR 6 (1978), 11-13.
- [7] M. M. Rao, Z. D. Ren, Theory of Orlicz spaces (New York, Basel, Hong Kong: Marcel Dekker Inc.) (1981).
- [8] M. H. A. Al-Rashed, B. Zegarlinski, Noncommutative Orlicz spaces associated to a state, Studia Math., 180 (2007), 199-209.
  - [9] M. H. A. Al-Rashed, B. Zegarlinski, Noncommutative Orlicz spaces associated to a state II, Linear Algebra and its Applications., 435 (2011), 2999-3013.
  - [10] Q. Xu, Operator spaces and noncommutative  $L_p$  spaces, 2007.
- <sup>145</sup> [11] S. T. Chen, Geometry of Orlicz spaces, in: Dissertations Mathematicae, Warszawa, 1996.
  - [12] W. Kunze, Noncommutative Orlicz spaces and generalized Arens algebras, Math. Nachr. 147 (1990), 123-138.

130

135