

Directed flow-augmentation*

Eun Jung Kim¹, Stefan Kratsch², Marcin Pilipczuk³, and Magnus Wahlström⁴

¹Université Paris-Dauphine, PSL Research University, CNRS, UMR 7243, LAMSADE, 75016, Paris, France.

²Humboldt-Universität zu Berlin, Germany

³University of Warsaw, Warsaw, Poland

⁴Royal Holloway, University of London, TW20 0EX, UK

Abstract

We show a flow-augmentation algorithm in directed graphs: There exists a polynomial-time algorithm that, given a directed graph G , two integers $s, t \in V(G)$, and an integer k , adds (randomly) to G a number of arcs such that for every minimal st -cut Z in G of size at most k , with probability $2^{-\text{poly}(k)}$ the set Z becomes a *minimum* st -cut in the resulting graph.

The directed flow-augmentation tool allows us to prove fixed-parameter tractability of a number of problems parameterized by the cardinality of the deletion set, whose parameterized complexity status was repeatedly posed as open problems:

1. CHAIN SAT, defined by Chitnis, Egri, and Marx [ESA'13, Algorithmica'17] [CEM17],
2. a number of weighted variants of classic directed cut problems, such as WEIGHTED st -CUT, WEIGHTED DIRECTED FEEDBACK VERTEX SET, or WEIGHTED ALMOST 2-SAT.

By proving that CHAIN SAT is FPT, we confirm a conjecture of Chitnis, Egri, and Marx that, for any graph H , if the LIST H -COLORING problem is polynomial-time solvable, then the corresponding vertex-deletion problem is fixed-parameter tractable.



European Research Council
Established by the European Commission



*This research is a part of a project that have received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme Grant Agreement 714704 (M. Pilipczuk). Eun Jung Kim is supported by the grant from French National Research Agency under JCJC program (ASSK: ANR-18-CE40-0025-01).

Contents

1	Introduction	1
2	Overview	4
3	Preliminaries	11
4	Directed Flow-Augmentation	12
4.1	Reachability patterns, leaders, and mincut sequences	13
4.2	Recursion structure and initial steps	15
4.3	Base case: only self-loops in H	16
4.4	Small ℓ case	17
4.5	Large ℓ case	20
4.6	Probability analysis	30
5	Applications: weighted st-cut, weighted DFVS, weighted Chain SAT	31
5.1	Tractable case of WEIGHTED BUNDLED CUT	31
5.2	Applications	33
6	Weighted Almost 2-SAT	34
6.1	Overview and outlook	36
6.2	Min-cut instances	37
6.2.1	Initial reductions	37
6.2.2	Solving arcless instances	39
6.3	General case	43

1 Introduction

The study of graph separation problems has been one of the more vivid areas of parameterized complexity in the recent 10–15 years. The term “graph separation problems” captures a number of classic graph problems where, given a (undirected or directed) graph G with a cut budget k (and possibly some annotations, such as terminal vertices), one aims at obtaining some separation via at most k edge or vertex deletions. For example, the classic st -CUT problem asks to delete at most k edges so that there is no s -to- t path in the resulting graph and the FEEDBACK VERTEX SET asks to remove at most k vertices so that the resulting graph does not contain any cycles (i.e., is a forest in the undirected setting or a DAG in the directed setting). In all these problems, the cardinality of the deletion set is a natural parameter to study.

In 2004, Marx introduced the notion of *important separators* [Mar04, Mar06] that turned out to be the key to fixed-parameter tractability of MULTIWAY CUT and DIRECTED FEEDBACK VERTEX SET [CLL⁺08], among many other examples. In subsequent years, the study of graph separation problems resulted in a rich toolbox of algorithmic techniques, such as *shadow removal* [MR14], treewidth reduction [MOR13], or randomized contractions [CCH⁺16, CLP⁺19, CKL⁺21]. At SODA’21, the current authors added one more item to this list: a flow-augmentation technique in undirected graphs [KKPW21]. As a result of these developments, most open problems regarding parameterized complexity of graph separation problems in undirected graphs has been resolved. Our work [KKPW21] shows that this statement can be formalized in some sense: every problem that can be formalized as a MIN UNSAT problem for some Boolean CSP language Γ that cannot express implication, is either $W[1]$ -hard or FPT by one of the aforementioned techniques.

For directed graphs, the chartered landscape is much less complete. The notion of important separators and related technique of shadow removal generalizes to directed graphs, leading to fixed-parameter tractability of DIRECTED FEEDBACK VERTEX SET [CLL⁺08], DIRECTED MULTIWAY CUT [CHM13], and DIRECTED SUBSET FEEDBACK VERTEX SET [CCHM15]. A number of problems whose undirected counterparts are FPT turned out to be $W[1]$ -hard in the directed setting, including DIRECTED MULTICUT [CHM13, PW18] or DIRECTED ODD CYCLE TRANSVERSAL [LRSZ20].

However, these results are far from satisfactorily charting the parameterized complexity of directed graph separation problems. Arguably, we seem to lack algorithmic techniques. Most notably, the powerful treewidth reduction theorem [MOR13], stating that (in undirected graphs) all separations of size at most k between two fixed terminals live in a part of the graph with treewidth bounded by $2^{O(k)}$, seems not to have any meaningful counterpart in directed graphs. As a result, essentially all known FPT algorithms for graph separation problems in directed graphs rely in some part on important separators, which is a greedy argument bounding the number of cuts between two terminals of bounded size that have inclusion-wise maximal set of vertices reachable from one of the terminal. This “greedy” part of this argument breaks down whenever a problem at hand has some weights or annotations.

On the other hand, despite efforts in the last years, we were not able to prove lower bounds for FPT algorithms for many directed graph separation problems. This suggests that maybe there are still more algorithmic techniques to be explored, leading to more positive tractability results.

In this work, we provide such a technique: we show that flow-augmentation, introduced by current authors in [KKPW21] for undirected graphs, generalizes to directed graphs.

Theorem 1.1. *There exists a randomized polynomial-time algorithm that, given a directed graph G , two vertices $s, t \in V(G)$, and an integer k , outputs a set $A \subseteq V(G) \times V(G)$ such that the following holds: for every minimal st -cut $Z \subseteq E(G)$ of size at most k , with probability $2^{-O(k^4 \log k)}$ Z remains an st -cut in $G + A$ and, furthermore, Z is a minimum st -cut in G .*

Here, a set $Z \subseteq E(G)$ is an st -cut if there is no path from s to t in $G - Z$; it is *minimal* if no proper subset of Z is an st -cut and *minimum* if it is of minimum possible cardinality. By $G + A$ we mean the graph G with all elements of A added as infinity-capacity arcs.

The proof of Theorem 1.1 is sketched in Section 2 and presented in full detail in Section 4. While it borrows a general outline and course of action from its undirected counterpart [KKPW21], at lower level most details are different, as directed graphs pose different challenges. The main differences are discussed in Section 2.

Applications To illustrate the applicability of the directed flow-augmentation, let us first consider the WEIGHTED st -CUT problem. Here, we are given a directed graph G with two terminals $s, t \in V(G)$, a weight function $\omega : E(G) \rightarrow \mathbb{Z}_+$, and two integers $k, W \in \mathbb{Z}_+$, and we ask for an st -cut Z of cardinality at most k and total weight at most W . This problem is known to be NP-hard and FPT when parameterized by $k + W$ [KLM⁺20].

By using directed flow-augmentation, we can ensure that the sought solution Z is actually a *minimum* st -cut (i.e., of minimum cardinality). Then, a solution can easily be found in polynomial-time: take $M := 1 + \sum_{e \in E(G)} \omega(e)$, set the capacity of every edge e as $\omega(e) + M$, and find an st -cut of minimum capacity. By repeating the algorithm $2^{\mathcal{O}(k^4 \log k)}$ times to ensure constant success probability, we obtain the following.

Theorem 1.2. *WEIGHTED st -CUT can be solved in randomized time $2^{\mathcal{O}(k^4 \log k)} n^{\mathcal{O}(1)}$.*

The above approach turns out to be more general. An instance of BUNDLED CUT consists of a directed graph G , vertices $s, t \in V(G)$, a nonnegative integer k , and a family \mathcal{B} of pairwise disjoint subsets of $E(G)$. The elements of \mathcal{B} are henceforth called *bundles*. A *cut* in a BUNDLED CUT instance $\mathcal{I} = (G, s, t, k, \mathcal{B})$ is an st -cut Z with $Z \subseteq \bigcup \mathcal{B}$. The BUNDLED CUT problem asks for an st -cut that intersects at most k bundles, that is, $|\{B \in \mathcal{B} \mid Z \cap B \neq \emptyset\}| \leq k$. One can define a weighted variant of the BUNDLED CUT problem where every bundle $B \in \mathcal{B}$ is equipped with a weight $\omega(B) \in \mathbb{Z}_+$, we are given also a weight bound W , and we ask for an st -cut Z that intersects at most k bundles whose total weight is at most W .

If all bundles are singletons, then BUNDLED CUT just asks for an st -cut of size at most k (with the edges of $E(G) \setminus \bigcup \mathcal{B}$ being undeletable), so it is polynomial-time solvable. It is known that, when restricted to bundles of size 2, BUNDLED CUT is $W[1]$ -hard when parameterized by k [MR09]. To get tractability, we define the following restriction called (WEIGHTED) BUNDLED CUT WITH ORDER where we require the following: for every $B \in \mathcal{B}$ one can order the arcs of B as $e_1^B, \dots, e_{|B|}^B$ such that for every $1 \leq i < j \leq |B|$ there exists a path $P_{i,j}^B$ in G that goes from one endpoint of e_i^B to one endpoint of e_j^B and uses only undeletable arcs (i.e., from $E(G) \setminus \bigcup \mathcal{B}$) and arcs of B .

Using directed flow-augmentation, we show the following.

Theorem 1.3. *WEIGHTED BUNDLED CUT WITH ORDER is randomized FPT when parameterized by k and maximum size of a bundle.*

For an integer $\ell \geq 1$, the ℓ -CHAIN SAT problem is the BUNDLED CUT problem, where every bundle is a path of length at most ℓ . At ESA'13, Chitnis, Egri, and Marx [CEM13, CEM17] defined ℓ -CHAIN SAT and showed that fixed-parameter tractability of ℓ -CHAIN SAT (for every fixed $\ell \geq 1$) is equivalent to the following conjecture.

Conjecture 1.4 (Conjecture 1.1 of [CEM17]). *For every graph H , if LIST H -COLORING problem is polynomial-time solvable, then the vertex-deletion variant (delete at most k vertices from the input graph to obtain a yes-instance to LIST H -COLORING) is fixed-parameter tractable when parameterized by k .*

Clearly, a ℓ -CHAIN SAT instance is a BUNDLED CUT WITH ORDER instance with maximum bundle size at most ℓ . Hence, Theorem 1.3 implies the following.

Corollary 1.5. *ℓ -CHAIN SAT is (randomized) FPT when parameterized by ℓ and k , even in the weighted setting. Consequently, Conjecture 1.4 is confirmed (with randomized FPT algorithms).*

By standard reductions, a DIRECTED FEEDBACK VERTEX SET instance can also be represented as a BUNDLED CUT WITH ORDER instance (with maximum size of a bundle and budget bounded linearly in the parameter of the input instance). Furthermore, in case of a weighted variant¹ the reduction preserves weights. Hence, we obtain the following.

Corollary 1.6. *WEIGHTED DIRECTED FEEDBACK VERTEX SET, parameterized by the cardinality of the deletion set, is (randomized) FPT.*

The question of parameterized complexity of WEIGHTED DIRECTED FEEDBACK VERTEX SET (which we answer affirmatively in Corollary 1.6) has been asked e.g. at Recent Advances in Parameterized Complexity school in December 2017 [Sau17] and in [LRS16].

We also take the applicability of directed flow-augmentation one step further and prove fixed-parameter tractability of WEIGHTED ALMOST 2-SAT. Here, we are given a 2-CNF formula ϕ , where every clause C has weight $\omega(C)$, and positive integers k and W ; the goal is to delete at most k clauses of total weight at most W to get a satisfiable instance.

Theorem 1.7. *WEIGHTED ALMOST 2-SAT parameterized by k is (randomized) FPT.*

The FPT algorithm for (unweighted) ALMOST 2-SAT problem was one of the first applications of important separators and one of the first bridges between graph separation problems and (boolean) constraint satisfaction problems [RO09].

Compared to the previous applications, applying directed flow-augmentation to WEIGHTED ALMOST 2-SAT in Theorem 1.7 is much more technically advanced. By known reductions [KW20], (WEIGHTED) ALMOST 2-SAT is FPT-equivalent to (WEIGHTED) DIGRAPH PAIR CUT. In this problem, we are given a directed graph G , a vertex $s \in V(G)$, and a number of pairs $\mathcal{T} \subseteq \binom{V(G)}{2}$. The goal is to delete at most k edges of G such that for every pair $xy \in \mathcal{T}$, either x or y is not reachable from s . (In the weighted variant, edges have positive integer weights and we also require that the total weight of deleted edges is not larger than a given threshold.)

In the process of solving a (WEIGHTED) DIGRAPH PAIR CUT instance, an algorithm makes some decisions about some pairs $xy \in \mathcal{T}$ whether x or y will be not reachable from s in the final graph. The vertices guessed to be unreachable can be merged into one sink t ; the solution Z needs to be then an st -cut, but may be required to be more than a *minimal* st -cut to satisfy other pairs of \mathcal{T} . On the other hand, in an inclusion-wise minimal solution Z every arc (u, v) serves some purpose: u is reachable from s in $G - Z$, but v is not. This leads to a definition of a *star* st -cut: a set $Z \subseteq E(G)$ is a star st -cut if it is an st -cut and, for every $(u, v) \in Z$, the vertex u is reachable from s in $G - Z$ but v is not. Clearly, every minimal st -cut is a star st -cut, but the reverse implication is far from true.

To apply directed flow-augmentation to WEIGHTED ALMOST 2-SAT, we enhanced the procedure of Theorem 1.1 to star st -cuts: the procedure returns a set $A \subseteq V(G) \times V(G)$ and a maximum st -flow \mathcal{P} in $G + A$ such that for every star st -cut Z of size at most k , with probability $2^{-\mathcal{O}(k^4 \log k)}$ the set Z remains a star st -cut in $G + A$ and, furthermore, the edges of $Z \cap E(\mathcal{P})$ is exactly a minimum st -cut in $G + A$. That

¹In WEIGHTED DIRECTED FEEDBACK VERTEX SET, the input graph is equipped with vertex weights being positive integers and one asks for a solution of cardinality at most k and total weight bounded by a threshold given on input.

is, the arcs of A make Z contain a minimum st -cut and, furthermore, the flow \mathcal{P} avoids edges of Z that are not part of the said minimum st -cut.

Future work We believe the new technique introduced in this work opens new research directions. We would like to point out to two of them.

In the $\text{MIN UNSAT}(\Gamma)$ problem, we are given an instance of a Constraint Satisfaction Problem over language Γ and an integer k to delete at most k clauses to get a satisfiable instance. Here, we restrict only to binary alphabet. Note that if Γ allows unary clauses and equalities, $\text{MIN UNSAT}(\Gamma)$ becomes the question of minimum (undirected) st -cut and when Γ allows unary clauses and implications, $\text{MIN UNSAT}(\Gamma)$ becomes the question of minimum (directed) st -cut.

Recall that the undirected flow-augmentation algorithm [KKPW21] led to full classification of the $\text{MIN UNSAT}(\Gamma)$ for Boolean alphabet into FPT problems and $W[1]$ -hard problems for languages Γ that are unable to express implication (and thus directed cuts). Up to this work, the main roadblock into generalizing this result to all Boolean languages was the ℓ -CHAIN SAT problem (which is easily expressible in the CSP world). Now this roadblock is removed and exploring this classification is a promising research direction.

Moreover, a similar generalization to the “star st -cut” was also present in the undirected case and was pivotal to fixed-parameter tractability of GENERALIZED COUPLED MINCUT problem [KKPW21]. We believe the exemplary case of Theorem 1.7 shows that directed flow-augmentation has potential to be a handy tool in proving fixed-parameter tractability of further Boolean MIN UNSAT CSP problems.

Corollary 1.6 provides an alternative algorithm for the classic DIRECTED FEEDBACK VERTEX SET algorithm, albeit with worse time complexity. We remark that up to now only one approach to fixed-parameter tractability of DIRECTED FEEDBACK VERTEX SET was known, namely the one using important separators [CLL⁺08]. Another newly opened research direction is: have we learned anything new about DIRECTED FEEDBACK VERTEX SET? To be more precise, let us recall two other important open problems about DIRECTED FEEDBACK VERTEX SET: is it solvable in time $2^{\mathcal{O}(k)} n^{\mathcal{O}(1)}$ and does it admit a polynomial kernel?

2 Overview

In this section we sketch the proof of Theorem 1.1.

Note that the actual formal statement in Section 4 is a bit more general: it handles a bit wider family of cuts than only minimal cuts (called later *star st -cuts*) and provides also a handy object called *witnessing flow*. While these extensions are critical for the WEIGHTED ALMOST 2-SAT algorithm in Section 6, already the case of minimal st -cuts encompasses main technical ideas and thus this overview focuses only on this case.

Basic notation Before we start, let us agree on basic nomenclature. All our graphs can be multigraphs. Edges may have capacities 1 or $+\infty$. The input graph has all edge capacities equal to 1. Since we never consider flows of value greater than k , a $+\infty$ -capacity edge is equivalent to $(k + 1)$ copies of an edge of capacity 1. If G is a graph and $A \subseteq V(G) \times V(G)$, then $G + A$ is the graph G with every arc of A added with capacity $+\infty$.

For vertices $s, t \in V(G)$, an st -flow is a collection \mathcal{P} of paths from s to t such that no edge of capacity 1 lies on more than one path of \mathcal{P} . The *value* of the flow is the number of paths. By $\lambda_G(s, t)$ we denote the maximum possible value of an st -flow; note that it may happen that $\lambda_G(s, t) = +\infty$. An st -flow is a *maximum st -flow* or *st -maxflow* if its value is $\lambda_G(s, t)$. By convention, if $\lambda_G(s, t) = +\infty$, then any st -flow that contains a path with all edges of capacity $+\infty$ is considered an st -maxflow.

A set $Z \subseteq E(G)$ is an *st-cut* if it contains no edge of capacity $+\infty$ and there is no path from s to t in $G - Z$. An *st-cut* Z is *minimal* if no proper subset of Z is an *st-cut* and *minimum* (or *st-mincut*) if it has minimum possible cardinality. By Menger’s theorem, if $\lambda_G(s, t) < +\infty$ then the size of every *st-mincut* is exactly $\lambda_G(s, t)$ and there are no *st-cuts* if $\lambda_G(s, t) = +\infty$.

For an *st-cut* Z , the *s-side* of Z is the set of vertices reachable from s in $G - Z$, and the *t-side* of Z is the complement of the *s-side*. Note that this is not symmetric as we do not mandate that t is reachable from all elements of the *t-side* in $G - Z$. If we have two (inclusion-wise) minimal *st-cuts* C_1 and C_2 such that both endpoints of every edge of C_1 lie in the *s-side* of C_2 (and thus both endpoints of every edge of C_2 lie in the *t-side* of C_1), then a vertex is *between* C_1 and C_2 if it is both in the *t-side* of C_1 and *s-side* of C_2 .

One flow-augmentation step If we do not pay particular attention to the exact bound on the probability of a success and we are happy with any $1/f(k)$ bound for a computable function f , it suffices to prove the following:

Theorem 2.1. *There exists a computable function $f : \mathbb{N} \rightarrow \mathbb{Z}_+$ and a polynomial-time algorithm that, given a directed graph G , two vertices $s, t \in V(G)$, and an integer k , outputs a set $A \subseteq V(G) \times V(G)$ such that $\lambda_{G+A}(s, t) > \lambda_G(s, t)$ and for every minimal *st-cut* Z of size at most k that is not a minimum *st-cut*, Z remains an *st-cut* in $G + A$ with probability at least $1/f(k)$.*

Indeed, an algorithm for Theorem 1.1 may first randomly guess the size of the minimal *st-cut* Z it cares about and then repeat the algorithm of Theorem 2.1, adding the output set A to the graph G , as long as $\lambda_G(s, t)$ is smaller than the guessed size of Z . Thus, in the remainder of this section we sketch the proof of Theorem 2.1.

For the proof of Theorem 2.1, we fix one minimal *st-cut* Z of size at most k that is not a minimum *st-cut* and we bound from below the probability that the algorithm never adds an arc from the *s-side* to the *t-side* of Z .

The algorithm for Theorem 2.1 is recursive. We start with computing an *st-maxflow* \mathcal{P} in G ; the goal is to push somehow an extra unit of flow from s to t by adding arcs so that no added arc violates the cut Z . The algorithm tries to isolate smaller and smaller parts of the graph through which it tries to push an extra unit of flow, guessing some properties of the cut Z along the way to guide the recursion. To obtain the desired success probability, we need to very restrictively guess properties of the cut Z .

Natural candidates as separations between distinct areas for recursive calls are minimum *st-cuts*. This was also the case for the undirected counterpart of [KKPW21]: the algorithm there constructed a maximal sequence of noncrossing and pairwise disjoint minimum *st-cuts*, partitioned them into minimal subsegments where the graph in-between was connected, and recursed on them.

In the undirected case, the edges of a minimum *st-cut* C is the entire “interface” between the *s-side* and *t-side* of C . In the directed case, there can be an unbounded number of arcs with tails in the *t-side* of C and heads in the *s-side* of C . This makes the interaction between the *s-side* of C and *t-side* of C much more complicated and harder to grasp in recursive calls.

One of the main ideas of the proof of Theorem 2.1 – and one of the main differences between it and its undirected counterpart [KKPW21] – is a very careful choice of the minimum *st-cuts* we want to separate along. Let us now introduce it in detail.

Reachability patterns, leaders, and mincut sequences An *instance* is a tuple $\mathcal{I} = (G, s, t, k)$ where G is a directed graph, $s, t \in V(G)$, and k is a nonnegative integer. An *instance with a flow* is a pair $(\mathcal{I}, \mathcal{P})$ where $\mathcal{I} = (G, s, t, k)$ is an instance and $\mathcal{P} = \{P_1, P_2, \dots, P_\lambda\}$ is an *st-flow*. In an *instance with a*

maximum flow we additionally require $\lambda = \lambda_G(s, t)$, that is, \mathcal{P} is a maximum st -flow. With an instance with a flow $(\mathcal{I}, \mathcal{P})$ we associate the residual graph $G^{\mathcal{P}}$.

For an instance with a maximum flow $(\mathcal{I} = (G, s, t, k), \mathcal{P} = \{P_1, \dots, P_\lambda\})$, a *reachability pattern* is a directed graph H with vertex set $V(H) = [\lambda]$, that is, each vertex $i \in V(H)$ corresponds to a path $P_i \in \mathcal{P}$, that contains a self-loop at every vertex. The *pattern associated with* $(\mathcal{I}, \mathcal{P})$ is a graph H with $V(H) = [\lambda]$ and $(i, j) \in E(H)$ if and only if there exists $v \in V(P_i) \setminus \{s, t\}$ and $u \in V(P_j) \setminus \{s, t\}$ such that there is a v to u path in $G^{\mathcal{P}}$. Note that a pattern associated with $(\mathcal{I}, \mathcal{P})$ is a reachability pattern if and only if every path P_i has at least one internal vertex.

For a vertex $v \in V(G)$, the set $\text{RReach}(v)$ is the set of vertices reachable from v in $G^{\mathcal{P}}$. For a vertex $v \in V(G)$ and an index $i \in [\lambda]$, the *last vertex on P_i residually reachable from v* , denoted $\text{LastReach}(v, P_i)$, is the last (closest to t) vertex u on P_i that is reachable from v in $G^{\mathcal{P}}$. Note that if u is reachable from v in $G^{\mathcal{P}}$, then all vertices preceding u on P_i are also reachable from v in $G^{\mathcal{P}}$, that is, the set of vertices of P_i reachable from v in $G^{\mathcal{P}}$ form a prefix of P_i . Furthermore, note that if v_1 and v_2 are on P_j and v_1 is earlier than v_2 on P_j , then $\text{LastReach}(v_1, P_i)$ is not later than $\text{LastReach}(v_2, P_i)$ on P_i (they may be equal).

Let C be an st -mincut in \mathcal{I} and let H be a reachability pattern. For $i \in [\lambda]$, the *leader of the path P_i after C* (with respect to the pattern H), denoted $\text{leader}_H(C, i)$, is the first (closest to s) vertex v on P_i such that for every $(i, j) \in E(H)$ the vertex $\text{LastReach}(v, P_j)$ is after C on P_j . A few remarks are in place. First, the notion of the leader is well-defined for any H as the vertex t is always a feasible candidate. Second, since a reachability pattern is required to contain a self-loop at every vertex, for every $i \in [\lambda]$ there is at least one edge $(i, j) \in E(H)$ to consider. Third, as C is oriented from the t -side to the s -side in $G^{\mathcal{P}}$, $\text{leader}_H(C, i)$ is after C on P_i and, furthermore, the entire path in $G^{\mathcal{P}}$ from $\text{leader}_H(C, i)$ to $\text{LastReach}(v, P_j)$ for every $(i, j) \in E(H)$ lies on the t -side of C .

For an st -mincut C and a reachability pattern H , we define the *mincut H -subsequent to C* as follows. If $t \in \text{RReach}(\text{leader}_H(C, i))$ for some $i \in [\lambda]$, the mincut H -subsequent to C is undefined. Otherwise, Lemma 4.2 implies that $\delta^+(\text{RReach}(\text{leader}_H(C, i)))$ is an st -mincut for every $i \in [\lambda]$. Let $X = \bigcup_{i \in [\lambda]} \text{RReach}(\text{leader}_H(C, i))$. By submodularity, $C' := \delta^+(X)$ is also an st -mincut; we proclaim C' to be the mincut H -subsequent to C . Observe that, by definition, for every $i \in [\lambda]$, the leader $\text{leader}_H(C, i)$ lies in the t -side of C and s -side of C' , the set $\text{RReach}(\text{leader}_H(C, i))$ lies in the s -side of C' and for every $(i, j) \in E(H)$ any path in $G^{\mathcal{P}}$ from $\text{leader}_H(C, i)$ to a vertex on P_j in the t -side of C (in particular, to $\text{LastReach}(\text{leader}_H(C, i), P_j)$) lies entirely in the t -side of C and s -side of C' . Furthermore, for every tail of an edge $(u, v) \in C' \cap E(P_j)$, either u is the leader $\text{leader}_H(C, j)$ or there exists $\text{leader}_H(C, i)$ (possibly with $i = j$) such that $u \in \text{RReach}(\text{leader}_H(C, i))$. Note that in the latter case, there is no st -mincut with $\text{leader}_H(C, i)$ in its s -side and u in its t -side because there is a residual path from $\text{leader}_H(C, i)$ to u . This proves the following observation.

Lemma 2.2. *Let $((G, s, t, k), \mathcal{P})$ be an instance with maximum flow, let H be the reachability pattern associated with it, let C be an st -mincut and let C' be an st -mincut whose endpoints are entirely in the t -side of C . Let G' be a graph constructed from G by contracting the s -side of C onto s and t -side of C' onto t and let \mathcal{P}' be an st -flow in G' consisting of the subpaths of paths of \mathcal{P} between the edge of C and the edge of C' . Suppose \mathcal{P}' is an st -maxflow in G' . If C' is H -subsequent to C , then the pattern associated with $((G', s, t, k), \mathcal{P}')$ is still H . Moreover, the st -mincut H -subsequent to C is an st -mincut closest to C with this property.*

For a reachability pattern H , an H -sequence of mincuts is a sequence C_1, C_2, \dots, C_ℓ of mincuts defined as follows. C_1 is the st -mincut closest to s and for $a > 1$ the mincut C_a is the mincut H -subsequent to C_{a-1} , as long as it is defined. Observe that for every $1 \leq a < b \leq \ell$ and $i \in [\lambda]$ we have that the edge of C_a on P_i lies strictly before the edge of C_b on P_i . That is, the s -side of C_a is contained in the s -side of C_b and, furthermore, on every path P_i the s -side of C_a is a strict subset of the s -side of C_b .

Preprocessing and the base case of the recursion The input to a recursive call is an instance with a flow $(\mathcal{I} = (G, s, t, k), \mathcal{P} = \{P_1, \dots, P_\lambda\})$ and the goal is to return a set $A \subseteq V(G) \times V(G)$ such that $\lambda < \lambda_{G+A}(s, t)$ and for every minimal st -cut Z with $\lambda < |Z| \leq k$ the set Z remains an st -cut in $G + A$ with good enough probability. In the initial, root call we set \mathcal{P} to be any st -maxflow.

A recursive call first performs a few preprocessing steps. If $\lambda < \lambda_G(s, t)$, it is safe to return $A = \emptyset$. If $\lambda_G(s, t) = 0$ or $\lambda_G(s, t) \geq k$, then there is no minimal st -cut of size at most k that is not a minimum st -cut. If this is the case, we can return $A = \{(s, t)\}$. Henceforth we assume $0 < \lambda_G(s, t) < k$ and that \mathcal{P} is an st -maxflow.

A relatively standard preprocessing step (that thus we do not describe here) can ensure that both $\delta^+(s)$ and $\delta^-(t)$ (the set of edges with tail in s and the set of edges with head in t , respectively) are disjoint and are both st -mincuts. In particular, this implies that the pattern H associated with \mathcal{I} and \mathcal{P} is a reachability pattern.

In the recursive steps, we will aim at either decreasing $2k - \lambda$ or keeping k and λ intact, but decreasing the number of edges in the pattern H associated with $(\mathcal{I}, \mathcal{P})$. Intuitively, less edges in $E(H)$ means that the graph G is somewhat simpler.

In the base case of the recursion, H consists of λ isolated vertices, with a loop on every vertex of $V(H) = [\lambda]$. In other words, $G^{\mathcal{P}}$ contains no path from a vertex of $V(P_i) \setminus \{s, t\}$ to a vertex of $V(P_j) \setminus \{s, t\}$ for any distinct $i, j \in [\lambda]$, or equivalently $G \setminus \{s, t\}$ contains no path from a vertex of $V(P_i)$ to a vertex of $V(P_j)$. In this case, the paths P_i are somewhat independent in G and it is relatively easy to show the following. (Here, an edge $e \in E(G)$ is *bottleneck* if there exists a minimum st -cut C with $e \in C$.)

Lemma 2.3. *Let $(\mathcal{I}, \mathcal{P})$ be an instance with maximum flow with proper boundaries and let H be its reachability pattern. If $|V(H)| = |E(H)|$, then for every minimal st -cut Z that is not a minimum cut, there exists $i \in [\lambda]$ such that no edge of $Z \cap E(P_i)$ is a bottleneck edge.*

Lemma 2.3 allows the following step: sample $i \in [\lambda]$ at random and return A consisting of duplicates of all bottleneck edges of P_i . Lemm 2.3 ensures that with probability at least $\lambda^{-1} \geq k^{-1}$, a minimal st -cut that is not a minimum st -cut remains an st -cut in $G + A$, while $\lambda_{G+A}(s, t) > \lambda_G(s, t)$ follows from the fact that we duplicated all bottleneck edges on one flow path.

If $|E(H)| > |V(H)|$, we aim at decomposing the graph further and make some recursive calls. We compute the H -sequence of mincuts C_1, C_2, \dots, C_ℓ . Note that we have $C_1 = \delta^+(s)$ and it is easy to see that C_2 is defined, and thus $\ell \geq 2$. We set a threshold ℓ^{big} depending on k (in the actual proof we have ℓ^{big} being polynomially bounded in k) and split into two cases: $\ell \leq \ell^{\text{big}}$ or $\ell > \ell^{\text{big}}$.

Small ℓ case. In this case, we can afford guessing (by random coin flip) for each $i \in [\ell]$ and $e \in C_i$, whether the endpoints of $e = (u, v)$ are in the s -side or t -side of Z .

There are a few simple cases. If for some $e = (u, v)$ we guessed that u is in the s -side of Z and v is in the t -side, then $e \in Z$; we can recurse on $G' := G - e$ and $\mathcal{P}' := \mathcal{P} \setminus \{P\}$ with parameter $k - 1$, where $P \in \mathcal{P}$ is a flow path containing e , and obtain a set A' , and return $A := A' \cup \{(s, u), (v, t)\}$. If for some $e = (u, v)$ we guessed that u is in the t -side of Z and v is in the s -side of Z , then we can return $A = \{(s, v), (u, t)\}$ as $s - v - u - t$ is an augmenting path in $G + A$ with respect to \mathcal{P} , so $\lambda_{G+A}(s, t) > \lambda$. Finally, if for every $e \in C_\ell$, both endpoints of e are in the s -side of Z , then we can recurse on G' being the graph G with the s -side of C_ℓ contracted onto s ; the fact that $C_{\ell+1}$ is undefined implies that in the recursive call the associated pattern is a proper subgraph of H .

In the main case, let A_0 consist of edges (s, v) for any endpoint v of an edge $e \in C_i$, $i \in [\ell]$ that is guessed to be in the s -side of Z and edges (u, t) for any endpoint u of an edge $e \in C_i$, $i \in [\ell]$ that is guessed

to be in the t -side of Z . If the guess is correct, Z remains an st -cut in $G + A_0$. If $\lambda_{G+A_0}(s, t) > \lambda$, then we can return A_0 , so assume otherwise.

Let C be the minimum st -cut in $G + A_0$ that is closest to t . For every endpoint of an edge of C , we guess whether it is in the s -side or t -side of Z . We again have a few simple cases. If for some $e = (u, v) \in C$, u is in the s -side of Z and v is in the t -side of Z , we have $e \in Z$ and we recurse on $G - e$ as before. If a head v of an edge of C is in the s -side of Z , then we return $A := A_0 \cup \{(s, v)\}$; $\lambda_{G+A}(s, t) > \lambda$ follows from the fact that C is the closest to t minimum st -cut in $G + A_0$.

We are left with the most interesting case where C is completely in the t -side of Z . We define $a \in [\ell]$ to be the maximum index such that all endpoints of C_a are in the s -side of Z . This is well-defined as $C_1 = \delta^+(s)$ is in the s -side of Z (as we are not in any of the simple cases). Also we have $a < \ell$ and C_{a+1} is defined (again, we are not in any of the simple cases) and also C lies entirely in the t -side of C_a . Let G' be the graph G with the s -side of C_a contracted onto s and the t -side of C contracted onto t . Let \mathcal{P}' be the flow \mathcal{P} projected onto G' (i.e., we shorten all paths to start from the edges of C_a and end with the edges of C).

The crucial observation is that the pattern associated with G' and \mathcal{P}' is a *proper* subgraph of H . Indeed, by the choice of a (and exclusion of the simple cases), there is at least one tail u of an edge of C_{a+1} that is in the t -side of Z ; this vertex u lies in the t -side of C_a and the s -side of C_{a+1} . Moreover, u is in the t -side of C as the edge (u, t) has been added to A_0 and C is an st -mincut in $G + A_0$. Since C_{a+1} is the mincut H -subsequent to C_a , by Lemma 2.2 the subgraph between C_a and C_{a+1} maintains all reachability (in the residual graph) described by H , but C_{a+1} is chosen to be the closest-to- C_a minimum st -cut that satisfies Lemma 2.2. Since u is in the t -side of C but s -side of C_{a+1} , we can infer that for at least one edge of H , the corresponding reachability is not present in G' and \mathcal{P}' . We recurse on G' and \mathcal{P}' ; this decrease in $|E(H)|$ is a progress in the recursive step.

We remark here that the above progress in the recursion is the main reason to introduce the notions of reachability patterns and sequences of mincuts. It seems to us very delicate; we were not able to reproduce a similar measure of progress with different ways of defining sequences of cuts C_1, \dots, C_ℓ .

Large ℓ case. The progress in the previous case is possible partially due to the fact that ℓ is bounded in k and we could afford guessing the s -side/ t -side assignment of all endpoints of all cuts C_i . If ℓ is large (unbounded in k), we need to resort to a color-coding step to split the long sequence of C_i 's into smaller chunks that are eligible for the small ℓ case.

Let Z_{ts} be the set of edges of paths of \mathcal{P} whose tail is in the t -side of Z but head is in the s -side of Z . On each flow path P_i , we have $|E(P_i) \cap Z| = |E(P_i) \cap Z_{ts}| + 1$ and thus $|Z_{ts}| \leq k - \lambda$. An index $a \in [\ell]$ is *touched* if there is an endpoint of an arc of $Z \cup Z_{ts}$ that is in the t -side of C_a and s -side of C_{a+1} . We have at most $2|Z \cup Z_{ts}| \leq 4k - 2\lambda$ touched indices; see Claim 4.

Recall that the construction of the cuts C_1, C_2, \dots, C_ℓ ensures that, for every $a \in [\ell - 1]$, between C_a and C_{a+1} one can find a path from P_i to P_j in the residual graph for every $(i, j) \in E(H)$. On the other hand, $G - \{s, t\}$ features no path in the residual graph from P_i to P_j if $(i, j) \notin E(H)$. In some sense, it means that the connectivity between paths P_i that is present in the entire graph G is repeatedly realized between each two consecutive cuts C_a .

The fact that we are operating with the residual graph was essential for the small ℓ case. Here, we need to depart from the residual graph and observe the following: for every $(i, j) \in E(H)$ and $1 \leq a \leq \ell - \lambda$, there is a path in G from P_i to P_j that starts between C_a and C_{a+1} , ends between $C_{a+\lambda-1}$ and $C_{a+\lambda}$, and is contained between C_a and $C_{a+\lambda}$.

This has a number of consequences. First, if ℓ^{big} is large enough, then H must be transitive.

Second, for every $a \in [\ell]$ let $L_a \subseteq [\lambda]$ be the set of indices i such that both endpoints of the unique edge of $E(P_i) \cap C_a$ are in the s -side of Z . Observe that if for some $1 \leq a \leq \ell - \lambda$, all indices $a \leq b \leq a + \lambda$ are untouched, then the set $L_{a+\lambda}$ is *downward-closed*: there is no arc $(i, j) \in E(H)$ such that $i \in L_{a+\lambda}$ but $j \notin L_{a+\lambda}$ (see Claim 7).

Third, observe that if $L \subseteq [\lambda]$ is downward-closed, then for every $i \in L$ and $j \notin L$, the graph $G - \{s, t\}$ contains no path from P_i to P_j . Hence, if we denote by $\text{cl}(L_a)$ be the minimal superset of L_a that is downward-closed, we have that $L_b \subseteq \text{cl}(L_a)$ whenever $1 \leq a \leq b \leq \ell$ (see Claim 6).

In particular, if L_a is downward-closed, we have the following.

1. $L_b \subseteq L_a$ for $b \geq a$.
2. For every $i \in L_a$, the entire prefix of P_i up to the edge of C_a is in the s -side of Z , as every path from a vertex v on such prefix to t needs to intersect $C_a \cap \bigcup_{j \in L_a} E(P_j)$ (and Z is a minimal st -cut).
3. Symmetrically for every $i \notin L_a$, the entire suffix of P_i from the edge of C_a is in the t -side of Z .
4. There is no edge $e = (u, v) \in Z$ with u in the t -side of C_a and v in the s -side of C_a . Indeed, if $e = (u, v)$ were such an edge, then by minimality of Z the graph $G - Z$ would need to feature a path from s to v and from u to t ; the first path needs to traverse an edge of $C_a \cap E(P_i)$ for $i \in L_a$ and the second path needs to traverse an edge of $C_a \cap E(P_j)$ for $j \notin L_a$, hence in between we have a path from P_i to P_j in $G - \{s, t\}$.

Let T be the set of touched indices. We say that two indices $a < b$ are *close* if $b - a \leq \lambda$. The set T partitions into *blocks*: maximal sets of indices such that two consecutive indices are close to each other. Let B_1, \dots, B_r be the blocks and for $1 \leq \alpha \leq r$, let a_α and b_α be the minimum and maximum index of the block B_α . We have $b_\alpha - a_\alpha = \mathcal{O}(k\lambda)$ while indices $a_\alpha - \lambda, a_\alpha - \lambda + 1, \dots, a_\alpha - 1$ and $b_\alpha + 1, \dots, b_\alpha + \lambda$ are untouched (we ignore here boundary cases when $a_\alpha \leq \lambda$ or $b_\alpha + \lambda > \ell$ which are easy to adjust to). In particular, $L_{a_\alpha - 1}$ and $L_{b_\alpha + \lambda}$ are downward-closed. Denote $L_\alpha^\leftarrow = L_{a_\alpha - 1}$, $L_\alpha^\rightarrow = L_{b_\alpha + \lambda}$, and $D^\alpha = L_\alpha^\leftarrow \setminus L_\alpha^\rightarrow$.

An important observation from the downward-closedness of L_α^\leftarrow and L_α^\rightarrow is the following: for every edge $e \in Z$, there is a block B_α such that *both* endpoints of e lie between C_{a_α} and $C_{b_\alpha + 1}$ (see Claim 10). Hence, Z partitions into $Z_1 \uplus Z_2 \uplus \dots \uplus Z_r$ where Z_α is the set of edges of Z with both endpoints between C_{a_α} and $C_{b_\alpha + 1}$.

For every $\alpha \in [r]$, define a graph G_α as follows: contract the s -side of $C_{a_\alpha - 1}$ onto s , contract the t -side of $C_{b_\alpha + \lambda}$ onto t , and from the edges incident with s or t leave only the ones on paths P_i for $i \in D_\alpha$. The flow paths P_i for $i \in D_\alpha$ naturally project to a flow \mathcal{P}_α of size $|D^\alpha|$ in G_α . The important observation again from the downward-closedness of L_α^\leftarrow and L_α^\rightarrow is as follows: in G_α , Z_α is a minimal st -cut (Claim 14).

This in particular implies that $D^\alpha \neq \emptyset$ and thus

$$[\lambda] = L_1^\leftarrow \supsetneq L_1^\rightarrow = L_2^\leftarrow \supsetneq L_2^\rightarrow = \dots = L_r^\leftarrow \supsetneq L_r^\rightarrow = \emptyset.$$

We would like to recurse on instances $\mathcal{I}_\alpha = (G_\alpha, \mathcal{P}_\alpha, k_\alpha)$ where $k_\alpha = |Z_\alpha|$, but there is a significant problem: we do not know the blocks B_α . However, following a similar step in the undirected algorithm of [KKPW21], we can deal with it with color-coding.

Sample $\Gamma \subseteq [\ell]$, aiming at follows: for every a that is close to a touched index, we want $a \in \Gamma$ if and only if a is touched. If we put every $a \in [\ell]$ into Γ with probability 0.5 independently of other indices, the success probability is $2^{-\mathcal{O}(k\lambda)}$. Then, the indices of Γ partition into Γ -blocks: one Γ -block is a maximal

subset of Γ in which two consecutive indices are close. If the guess is successful, every block is a Γ -block, but there may be numerous Γ -blocks that are in fact wholly untouched.

We can guess $r \leq 4k - 2\lambda$, sets $L_\alpha^{\leftarrow}, L_\alpha^{\rightarrow}$ as well as sizes $k_\alpha = |Z_\alpha|$ for $\alpha \in [r]$. For every Γ -block, we guess its “type” in $[r]$, aiming that a true block B_α has type α . For a Γ -block B of type $\alpha(B)$, we construct from it an instance $(G_B, s, t, k_{\alpha(B)})$ with flow \mathcal{P}_B in the same way as we constructed $G_{\alpha(B)}$ and $\mathcal{P}_{\alpha(B)}$ for $B_{\alpha(B)}$. If $k_{\alpha(B)} > |D_{\alpha(B)}|$ (i.e., $Z_{\alpha(B)}$ is not a minimum cut in $G_{\alpha(B)}$) we recurse on it, obtaining a set A_B (we denote $A_B = \emptyset$ for cases $k_{\alpha(B)} = |D_{\alpha(B)}|$).

The final set A consists of:

1. for every Γ -block B with minimum index a and maximum index b , the following edges:
 - (a) for every $(u, v) \in A_B$:
 - if $u \neq s$ and $v \neq t$, just the arc (u, v) itself;
 - if $u = s$ and $v \neq t$, all arcs (u', v) where u' ranges over all tails of edges of C_{a-1} on paths P_i for $i \in D_\alpha$;
 - if $u \neq s$ and $v = t$, all arcs (u, v') where v' ranges over all tails of edges of $C_{b+\lambda}$ on paths P_i for $i \in D_\alpha$;
 - if $u = s$ and $v = t$, all arcs (u', v') where u' ranges over all tails of edges of C_{a-1} on paths P_i for $i \in D_\alpha$ and v' ranges over all tails of edges of $C_{b+\lambda}$ on paths P_i for $i \in D_\alpha$;
 - (b) for every $i \in [\lambda] \setminus D_\alpha$ and arc from the tail of the edge of $C_{a-1} \cap E(P_i)$ to the tail of the edge of $C_{b+\lambda} \cap E(P_i)$;
2. for every $1 \leq c \leq \ell$ for which there is no Γ -block B with minimum index a and maximum index b and $a - 1 \leq c \leq b + \lambda$:
 - for every $i \in [\lambda]$, an edge from the tail of the edge of $C_c \cap E(P_i)$ to the tail of the edge of $C_{c+1} \cap E(P_i)$;
3. for every $1 \leq c \leq \ell$ for which there is no Γ -block B with minimum index a and maximum index b and $a - 1 < c < b + \lambda$:
 - for every $1 \leq \alpha \leq r$, for every $i, j \in D_\alpha$, an edge from the tail of the edge of $C_c \cap E(P_i)$ to the tail of the edge of $C_c \cap E(P_j)$.

To see that $\lambda_{G+A}(s, t) > \lambda$, notice first that as Z is not a minimum st -cut, there exists an index α where $k_\alpha = |Z_\alpha| > |D_\alpha|$. We push an extra unit of flow along flow paths $P_i, i \in D_\alpha$ as follows.

- For every Γ -block B of type α (with minimum index a and maximum index b), we push $|D_\alpha| + 1$ units of flow from the tails of edges of C_{a-1} on paths P_i for $i \in D_\alpha$ to the tails of edges of $C_{b+\lambda}$ on paths P_i for $i \in D_\alpha$ using edges of A_B (Point 1a) and inductive assumption from the recursive call.
- For every Γ -block B of type $\alpha' \neq \alpha$, we push $|D_\alpha| + 1$ units of flow along edges added in Point 1b for paths $P_i, i \in D_\alpha$.
- For every $1 \leq c \leq \ell$ for which there is no Γ -block B with minimum index a and maximum index b and $a - 1 \leq c \leq b + \lambda$, we push $|D_\alpha| + 1$ units of flow along edges added in Point 2 for paths $P_i, i \in D_\alpha$.

- We use edges added in Point 3 to reshuffle the flow on paths $P_i, i \in D_\alpha$ between the steps above, if needed.

Let us now briefly analyse the progress in the above recursion. If $r > 1$, then all recursive calls treat strictly smaller value of k . For $r = 1$, we have $D_1 = \lambda$ and possibly $k_\alpha = k$. However, then the recursive calls fall into the “small ℓ case” as the cuts C_c inside recursive call remain an H -sequence of mincuts. This is the desired progress and it finishes the overview of the proof of Theorem 2.1.

3 Preliminaries

All our graphs can be multigraphs. Edges may have capacities 1 or $+\infty$. The input graph has all edge capacities equal to 1. Since we never consider flows of value greater than k , a $+\infty$ -capacity edge is equivalent to $(k + 1)$ copies of an edge of capacity 1. If G is a graph and $A \subseteq V(G) \times V(G)$, then $G + A$ is the graph G with every arc of A added *with capacity* $+\infty$.

For a directed graph G and a set $X \subseteq V(G)$, the set $\delta_G^+(X)$ is the set of arcs with tails in X and heads in $V(G) \setminus X$. Similarly $\delta_G^-(X)$ is the set of arcs with tails in $V(G) \setminus X$ and heads in X . We write $\delta_G^+(v)$ and $\delta_G^-(v)$ for $\delta_G^+(\{v\})$ and $\delta_G^-(\{v\})$. When the graph G under consideration is clear from the context, we omit the subscript G .

Let G be a directed graph and let $s, t \in V(G)$. An st -flow is a collection \mathcal{P} of paths from s to t such that no edge of capacity 1 lies on more than one path of \mathcal{P} . The *value* of the flow is the number of paths. By $\lambda_G(s, t)$ we denote the maximum possible value of an st -flow; note that it may happen that $\lambda_G(s, t) = +\infty$. An st -flow is a *maximum st -flow* or *st -maxflow* if its value is $\lambda_G(s, t)$. By convention, if $\lambda_G(s, t) = +\infty$, then any st -flow that contains a path with all edges of capacity $+\infty$ is considered an st -maxflow.

A set $Z \subseteq E(G)$ is an st -cut if it contains no edge of capacity $+\infty$ and there is no path from s to t in $G - Z$. An st -cut Z is *minimal* if no proper subset of Z is an st -cut and *minimum* (or *st -mincut*) if it has minimum possible cardinality. By Menger’s theorem, if $\lambda_G(s, t) < +\infty$ then the size of every st -mincut is exactly $\lambda_G(s, t)$ and there are no st -cuts if $\lambda_G(s, t) = +\infty$.

An st -cut Z is a *star st -cut* if for every $(u, v) \in Z$, in the graph $G - Z$ there is a path from s to u but there is no path from s to v . Note that every minimal st -cut is a star st -cut, but the implication in the other direction does not hold in general. For a star st -cut Z in G , by $\text{core}_G(Z) \subseteq Z$ we denote the set of arcs $(u, v) \in Z$ such that there exists a path from v to t in $G - Z$. We drop the subscript if the graph G is clear from the context.

We observe the following.

Lemma 3.1. *If Z is a star st -cut in a graph G , then $\text{core}(Z)$ is a minimal st -cut.*

For an st -cut Z , the s -side of Z is the set of vertices reachable from s in $G - Z$, and the t -side of Z is the complement of the s -side. Note that this is not symmetric, we do not mandate that t is reachable from all elements of the t -side in $G - Z$. In a star st -cut Z , all tails of edges of Z are in the s -side of Z and all heads of edges of Z are in the t -side of Z .

We say that a set of arcs $A \subseteq V(G) \times V(G)$ is *compatible* with a star st -cut Z if the following holds: for every $v \in V(G)$, there is a path from s to v in $G - Z$ if and only if there is a path from s to v in $(G + A) - Z$. Equivalently: the s -sides and t -sides of Z are equal in G and $G + A$, or no arc of A has its tail in the s -side of Z in G and its head in the t -side of Z in G .

An immediate yet important observation is as follows.

Lemma 3.2. *If Z is a star st -cut in G and $A \subseteq V(G) \times V(G)$ is compatible with Z , then Z is a star st -cut in $G + A$ as well.*

We remark that albeit in the setting of Lemma 3.2 the cut Z remains a star st -cut in $G + A$, it may happen that $\text{core}_G(Z) \subsetneq \text{core}_{G+A}(Z)$, as the arcs of A may add some new reachability towards t .

Assume Z is a star st -cut in G such that $\text{core}(Z)$ is an st -mincut. An st -maxflow \mathcal{P} is a *witnessing flow* if $E(\mathcal{P}) \cap Z = \text{core}(Z)$, that is, \mathcal{P} contains one edge of $\text{core}(Z)$ on each flow path and no other edge of Z . A witnessing flow may not exist in general, even if $\text{core}(Z)$ is an st -mincut. In contrast, on undirected graphs an arbitrary st -maxflow is a witnessing flow for the counterpart of $\text{core}(Z)$ when the latter is an st -mincut [KKPW21]. However, our flow-augmentation procedure will ensure that not only $\text{core}(Z)$ becomes an st -mincut in the augmented graph, but also a flow is returned that is a witnessing flow in the augmented graph. Formally, for a star st -cut Z in G , $A \subseteq V(G) \times V(G)$, and an st -maxflow $\widehat{\mathcal{P}}$ in $G + A$, we say that $(A, \widehat{\mathcal{P}})$ is *compatible* with Z if A is compatible with Z , $\text{core}_{G+A}(Z)$ is an st -mincut in $G + A$, and $\widehat{\mathcal{P}}$ is a witnessing flow for Z in $G + A$. Our flow-augmenting procedure will return a pair $(A, \widehat{\mathcal{P}})$ that is compatible with a fixed star st -cut Z with good probability.

An edge $e \in E(G)$ is a *bottleneck* edge if there exists an st -mincut that contains e .

For an st -flow \mathcal{P} , the *residual network* $G^{\mathcal{P}}$ is constructed from G by first adding a capacity-1 arc (u, v) for every $P \in \mathcal{P}$ and $(v, u) \in E(P)$ (an edge (u, v) is added multiple times if (v, u) appears on multiple paths of \mathcal{P}) and then deleting all capacity-1 arcs on paths of \mathcal{P} . An *augmenting path* is a path from s to t in $G^{\mathcal{P}}$. Recall that \mathcal{P} is an st -maxflow if and only if there is no augmenting path for \mathcal{P} and G .

Let C be an st -mincut and \mathcal{P} be an st -maxflow. First, note that for every $P \in \mathcal{P}$ there is exactly one edge of $C \cap E(P)$. This edge splits P into a part in the s -side of C and a part in the t -side of C . We say that an edge or a vertex on P is *before* (*after*) C on P if it is in the s -side (t -side, respectively) of C . We infer that every path in G that goes from a vertex in the s -side of C to a vertex in the t -side of C visits an edge of C and, furthermore, no augmenting path goes from a vertex in the s -side of C to a vertex in the t -side of C .

By submodularity, if Z_1 and Z_2 are st -mincuts and X_1 is the s -side of Z_1 and X_2 is the s -side of Z_2 , then both $\delta^+(X_1 \cup X_2)$ and $\delta^+(X_1 \cap X_2)$ are st -mincuts. This implies that there exists a unique st -mincut with inclusion-wise minimal s -side and a unique st -mincut with inclusion-wise maximal s -side. We call them the st -mincut *closest to s* and the st -mincut *closest to t* .

4 Directed Flow-Augmentation

An *instance* is a tuple $\mathcal{I} = (G, s, t, k)$ where G is a directed graph, $s, t \in V(G)$, and k is a nonnegative integer. An *instance with a flow* is a pair $(\mathcal{I}, \mathcal{P})$ where $\mathcal{I} = (G, s, t, k)$ is an instance and $\mathcal{P} = \{P_1, P_2, \dots, P_\lambda\}$ is an st -flow. In an *instance with a maximum flow* we additionally require $\lambda = \lambda_G(s, t)$, that is, \mathcal{P} is a maximum st -flow. With an instance with a flow $(\mathcal{I}, \mathcal{P})$ we associate the residual graph $G^{\mathcal{P}}$.

In this section we prove the following result.

Theorem 4.1. *There exists a polynomial-time algorithm that, given an instance $\mathcal{I} = (G, s, t, k)$, returns a set $A \subseteq V(G) \times V(G)$ and an st -maxflow $\widehat{\mathcal{P}}$ in $G + A$ such that for every star st -cut Z of size at most k , with probability $2^{-\mathcal{O}(k^4 \log k)}$ the pair $(A, \widehat{\mathcal{P}})$ is compatible with Z .*

A few remarks are in place. First, Lemma 3.2 ensures that if A is compatible with Z , then Z remains a star st -cut in $G + A$. Second, we not only guarantee that A is compatible with Z with good probability and that $\text{core}_{G+A}(Z)$ is an st -mincut in $G + A$, but also that $\widehat{\mathcal{P}}$ is a witnessing flow; in particular, that a witnessing flow exists at all. Third, a usage of Theorem 4.1 makes sense even for cuts Z with $\text{core}(Z)$ being an st -mincut for the sake of the witnessing flow: we may need to add some edges to G for a witnessing flow to exist, let alone being returned by the algorithm. Finally, due to the convention that for $\lambda_{G+A}(s, t) = +\infty$,

any flow containing an st -path with all arcs of capacity $+\infty$ is an st -maxflow, the algorithm is allowed to return a pair $(A, \widehat{\mathcal{P}})$ such that $\lambda_{G+A}(s, t) = +\infty$ and $\widehat{\mathcal{P}}$ contains an st -path with all edges of capacity $+\infty$ (but clearly such a pair $(A, \widehat{\mathcal{P}})$ is not compatible with any star st -cut Z of size at most k).

If $\lambda_G(s, t) = 0$, then for every star st -cut Z in G we have $\text{core}(Z) = \emptyset$, so we can return $A = \emptyset$ and $\mathcal{P} = \emptyset$. If $\lambda_G(s, t) > k$, then there is no star st -cut Z of size at most k , so $A = \emptyset$ and \mathcal{P} being any st -maxflow is a valid outcome. Henceforth we assume $0 < \lambda_G(s, t) \leq k$.

We set a threshold $\ell^{\text{big}} := 4k^2 + 3$ for later use. Note that while the algorithm is recursive and the value of k may decrease in the recursive calls, the threshold ℓ^{big} is set once at the beginning, using the initial value of k .

4.1 Reachability patterns, leaders, and mincut sequences

Let $(\mathcal{I} = (G, s, t, k), \mathcal{P} = \{P_1, \dots, P_\lambda\})$ be an instance with a maximum flow.

A *reachability pattern* is a directed graph H with vertex set $V(H) = [\lambda]$, that is, each vertex $i \in V(H)$ corresponds to a path $P_i \in \mathcal{P}$, that contains a self-loop at every vertex. The *pattern associated with* $(\mathcal{I}, \mathcal{P})$ is a graph H with $V(H) = [\lambda]$ and $(i, j) \in E(H)$ if and only if there exists $v \in V(P_i) \setminus \{s, t\}$ and $u \in V(P_j) \setminus \{s, t\}$ such that there is a v to u path in $G^{\mathcal{P}}$. Note that H is a reachability pattern if and only if every path P_i has at least one internal vertex.

For a vertex $v \in V(G)$, the set $\text{RReach}(v)$ is the set of vertices reachable from v in $G^{\mathcal{P}}$. For a vertex $v \in V(G)$ and an index $i \in [\lambda]$, the *last vertex on P_i residually reachable from v* , denoted $\text{LastReach}(v, P_i)$, is the last (closest to t) vertex u on P_i that is reachable from v in $G^{\mathcal{P}}$. Note that if u is reachable from v in $G^{\mathcal{P}}$, then all vertices preceding u on P_i are also reachable from v in $G^{\mathcal{P}}$, that is, the set of vertices of P_i reachable from v in $G^{\mathcal{P}}$ form a prefix of P_i . Furthermore, note that if v_1 and v_2 are on P_j and v_1 is earlier than v_2 on P_j , then $\text{LastReach}(v_1, P_i)$ is not later than $\text{LastReach}(v_2, P_i)$ on P_i (they may be equal).

We have the following simple observation.

Lemma 4.2. *Let \mathcal{P} be an st -maxflow in instance $\mathcal{I} = (G, s, t, k)$ and assume $\lambda_G(s, t) > 0$. Then for every $v \in V(G)$ exactly one of the following options hold:*

- $s \in \text{RReach}(v)$, $t \notin \text{RReach}(v)$, and $\delta_G^+(\text{RReach}(v))$ is an st -mincut;
- $s, t \in \text{RReach}(v)$, v is on the t -side of every st -mincut, and $\delta_G^+(\text{RReach}(v)) = \emptyset$;
- $s, t \notin \text{RReach}(v)$, t is not reachable from v in G , and $\delta_G^+(\text{RReach}(v)) = \emptyset$.

Proof. Note that $s \in \text{RReach}(v)$ if and only if there is a path from v to a vertex on \mathcal{P} in G . The forward implication follows from that any directed path from v to $V(\mathcal{P})$ in $G^{\mathcal{P}}$ contains as a prefix a subpath from v to $V(\mathcal{P})$ which is also a path in G . The backward implication is trivial.

To begin with, consider the case when no vertex on \mathcal{P} is reachable from v in G . From the observation in the previous paragraph, it follows that no vertex on \mathcal{P} is reachable from v in $G^{\mathcal{P}}$ as well. In particular, we have $s, t \notin \text{RReach}(v)$. It remains to note that any edge contained in $\delta_G^+(\text{RReach}(v))$ is on \mathcal{P} since otherwise the head of the edge would have been included in $\text{RReach}(v)$. From $\text{RReach}(v) \cap V(\mathcal{P}) = \emptyset$, we conclude $\delta_G^+(\text{RReach}(v)) = \emptyset$.

Therefore, we turn to the case when some vertex on \mathcal{P} is reachable from v in G , which subsumes $s \in \text{RReach}(v)$. First, if $t \notin \text{RReach}(v)$, observe that the set of vertices on P_i reachable from v in $G^{\mathcal{P}}$ forms a proper prefix of P_i for every $i \in [\lambda]$. Therefore, $\delta_G^+(\text{RReach}(v))$ takes precisely one edge from each $P_i \in \mathcal{P}$. It suffices to show that $C := \delta_G^+(\text{RReach}(v))$ is an st -cut. Indeed, an st -path in $G - C$ contains a path in $G - E(\mathcal{P})$ from a vertex on P_i before $C \cap E(P_i)$ to a vertex on P_j after $C \cap E(P_j)$ for some

$i, j \in [\lambda]$. That is, a vertex on \mathcal{P} after $C \cap E(\mathcal{P})$ is reachable from v in $G^{\mathcal{P}}$, a contradiction. Second, if $t \in \text{RReach}(v)$, then $\text{RReach}(v)$ contains the entire vertices on \mathcal{P} and thus $\delta_G^+(\text{RReach}(v)) = \emptyset$. That v is on the t -side of every st -mincut immediately follows from the condition $t \in \text{RReach}(v)$. \square

Let C be an st -mincut in \mathcal{I} and let H be a reachability pattern. For $i \in [\lambda]$, the *leader of the path P_i after C* (with respect to the pattern H), denoted $\text{leader}_H(C, i)$, is the first (closest to s) vertex v on P_i such that for every $(i, j) \in E(H)$ the vertex $\text{LastReach}(v, P_j)$ is after C on P_j . A few remarks are in place. First, the notion of the leader is well-defined as the vertex t is always a feasible candidate. Second, since a reachability pattern is required to contain a self-loop at every vertex, for every $i \in [\lambda]$ there is at least one edge $(i, j) \in E(H)$ to consider. Third, as C is oriented from the t -side to the s -side in $G^{\mathcal{P}}$, $\text{leader}_H(C, i)$ is after C on P_i and, furthermore, the entire path in $G^{\mathcal{P}}$ from $\text{leader}_H(C, i)$ to $\text{LastReach}(\text{leader}_H(C, i), P_j)$ for every $(i, j) \in E(H)$ lies on the t -side of C .

For an st -mincut C and a reachability pattern H , we define the *mincut H -subsequent to C* as follows. If $t \in \text{RReach}(\text{leader}_H(C, i))$ for some $i \in [\lambda]$, the mincut H -subsequent to C is undefined. Otherwise, Lemma 4.2 implies that $\delta^+(\text{RReach}(\text{leader}_H(C, i)))$ is an st -mincut for every $i \in [\lambda]$. Let $X = \bigcup_{i \in [\lambda]} \text{RReach}(\text{leader}_H(C, i))$. By submodularity, $C' := \delta^+(X)$ is also an st -mincut; we proclaim C' to be the mincut H -subsequent to C . Observe that, by definition, for every $i \in [\lambda]$, the leader $\text{leader}_H(C, i)$ lies in the t -side of C and s -side of C' , the set $\text{RReach}(\text{leader}_H(C, i))$ lies in the s -side of C' and for every $(i, j) \in E(H)$ any path in $G^{\mathcal{P}}$ from $\text{leader}_H(C, i)$ to a vertex on P_j in the t -side of C (in particular, to $\text{LastReach}(\text{leader}_H(C, i), P_j)$) lies entirely in the t -side of C and s -side of C' .

For a reachability pattern H , an *H -sequence of mincuts* is a sequence C_1, C_2, \dots, C_ℓ of mincuts defined as follows. C_1 is the st -mincut closest to s and for $a > 1$ the mincut C_a is the mincut H -subsequent to C_{a-1} , as long as it is defined. Observe that for every $1 \leq a < b \leq \ell$ and $i \in [\lambda]$ we have that the edge of C_a on P_i lies strictly before the edge of C_b on P_i . That is, the s -side of C_a is contained in the s -side of C_b and, furthermore, on every path P_i the s -side of C_a is a strict subset of the s -side of C_b .

We say that the instance \mathcal{I} has *proper boundaries* if both $\delta^+(s)$ and $\delta^-(t)$ are st -mincuts and $\delta^+(s) \cap \delta^-(t) = \emptyset$, that is, there is no arc (s, t) . In other words, \mathcal{I} has proper boundaries if $\delta^+(s)$ is the st -mincut closest to s , $\delta^-(t)$ is the st -mincut closest to t , and these cuts are disjoint. Note that in particular if \mathcal{I} has proper boundaries, then the pattern H associated with it is a reachability pattern. We have the following observations.

Lemma 4.3. *Let $(\mathcal{I}, \mathcal{P})$ be an instance with a maximum flow such that \mathcal{I} has proper boundaries and let H be the associated reachability pattern. Let C_1, C_2, \dots, C_ℓ be the H -sequence of mincuts. Then $C_1 = \delta^+(s)$ and $\ell \geq 2$, that is, C_2 is defined.*

Proof. $C_1 = \delta^+(s)$ by the definition of having proper boundaries. By the definition of H , for every $(i, j) \in E(H)$ there exists a vertex $v_{i,j} \in V(P_i) \setminus \{s, t\}$ and $u_{i,j} \in V(P_j) \setminus \{s, t\}$ such that $u_{i,j}$ is reachable from $v_{i,j}$ in $G^{\mathcal{P}}$. Consequently, $\text{leader}_H(C_1, i) \neq t$ for every $i \in [\lambda]$. Since $\delta^-(t)$ is an st -mincut, $t \notin \text{RReach}(\text{leader}_H(C_1, i))$ for every $i \in [\lambda]$. Thus, C_2 is defined. \square

Lemma 4.4. *Let $(\mathcal{I}, \mathcal{P})$ be an instance with a maximum flow such that \mathcal{I} has proper boundaries and let H be the associated reachability pattern. Let C_1, C_2, \dots, C_ℓ be the H -sequence of mincuts. If $\ell \geq 3$, then H is transitive.*

Proof. Let (i_1, i_2) and (i_2, i_3) be two arcs of H . Our goal is to prove that $(i_1, i_3) \in E(H)$.

By the construction of C_3 , since $(i_1, i_2) \in E(H)$, there exists $v \in V(P_{i_1})$ and $u \in V(P_{i_2})$ such that both v and u are after C_2 and before C_3 on paths P_{i_1} and P_{i_2} , respectively, and u is reachable from v in

$G^{\mathcal{P}}$. Similarly, by the construction of C_2 , since $(i_2, i_3) \in E(H)$, there exists $v' \in V(P_{i_2})$ and $u' \in V(P_{i_3})$ such that both v' and u' are after C_1 but before C_2 on paths P_{i_2} and P_{i_3} , respectively, and u' is reachable from v' in $G^{\mathcal{P}}$. Since u is after C_2 on P_{i_2} and v' is before C_2 on P_{i_2} , v' is before u on P_{i_2} and hence v' is reachable from u in $G^{\mathcal{P}}$. We infer that $u' \in V(P_{i_3}) \setminus \{s, t\}$ is reachable from $v \in V(P_{i_1}) \setminus \{s, t\}$ in $G^{\mathcal{P}}$, hence $(i_1, i_3) \in E(H)$, as desired. \square

4.2 Recursion structure and initial steps

The algorithm of Theorem 4.1 is recursive. The input is an instance with a flow $(\mathcal{I} = (G, s, t, k), \mathcal{P} = \{P_1, \dots, P_\lambda\})$ and an integer κ . The goal is to return a set $A \subseteq V(G) \times V(G)$ and an st -maxflow $\widehat{\mathcal{P}}$ in $G + A$ such that $\lambda_{G+A}(s, t) \geq \kappa$ and for every star st -cut Z with $|Z| \leq k$ and $|\text{core}_G(Z)| \geq \kappa$, $(A, \widehat{\mathcal{P}})$ is compatible with Z (i.e., A is compatible with Z , $\text{core}_{G+A}(Z)$ is an st -mincut in $G + A$, and $\widehat{\mathcal{P}}$ is a witnessing flow for Z in $G + A$) with good probability.

For an input $(\mathcal{I}, \mathcal{P}, \kappa)$, the algorithm may perform some randomized choices and a number of recursive calls. In each case, we will argue that the returned output $(A, \widehat{\mathcal{P}})$ is compatible with Z , usually using the fact that a similar property holds for outputs of recursive subcalls. While performing random choices, we will always say what to aim for (e.g., for a vertex v we may randomly guess if it is on the s -side or t -side of the hypothetical cut Z) and bound the probability of a correct choice. Furthermore, for some recursive calls $(\mathcal{I}', \mathcal{P}', \kappa')$ we will define a star st -cut Z' (depending usually on Z) and prove that if the returned output $(A', \widehat{\mathcal{P}}')$ is compatible with Z' , then the final output $(A, \widehat{\mathcal{P}})$ is compatible with Z .

However, for sake of clarity in the description of the algorithm we will not perform a formal probability analysis, but we postpone it to Section 4.6. Instead, we always informally indicate what is the measure of progress in the recursive subcalls, so that the claim that the success probability is lower bounded by a function of k would be clear (but the exact estimate on the success probability requires some tedious calculations).

A meticulous reader can notice that the flow \mathcal{P} is not used in the definitions above. We will use it to keep track of the progress of the algorithm; in some recursive calls the only progress will be that the flow \mathcal{P} at hand changes structure in some sense.

Let us proceed now to the description of the algorithm.

Given $(\mathcal{I}, \mathcal{P}, \kappa)$, the algorithm first performs a number of preprocessing steps. First, if \mathcal{P} is not an st -maxflow, we compute an st -maxflow \mathcal{P}' and recurse on $(\mathcal{I}, \mathcal{P}', \kappa)$. Thus, we henceforth assume \mathcal{P} is an st -maxflow, $\lambda = \lambda_G(s, t)$.

If $\lambda = 0$, then every star st -cut Z satisfies $\text{core}(Z) = \emptyset$. In this case we return $A = \emptyset$ and $\widehat{\mathcal{P}} = \emptyset$ if $\kappa = 0$ and $A = \{(s, t)\}$ and $\widehat{\mathcal{P}}$ consisting of one flow path along the edge (s, t) if $\kappa > 0$.

If $\lambda > k$, then there is no star st -cut Z with $|Z| \leq k$. Hence, we can return $A = \{(s, t)\}$ and $\widehat{\mathcal{P}}$ consisting of one flow path along (s, t) .

In the remaining (and most interesting) case $0 < \lambda = \lambda_G(s, t) \leq k$, we start by computing C_{\leftarrow} and C_{\rightarrow} , the st -mincuts closest to s and t , respectively. We can also assume $\kappa \geq \lambda$, as we can replace $\kappa := \max(\kappa, \lambda)$.

If $C_{\leftarrow} \neq \delta^+(s)$, we proceed as follows. Let A_{\leftarrow} be the set of tails of edges of C_{\leftarrow} , except for s . We make a random guess: with probability 0.5 we guess that there exists a vertex $v \in A_{\leftarrow}$ that is on the t -side of Z , and with the remaining probability we guess that there is no such vertex. The guess is correct with probability 0.5.

In the first case, we additionally guess one tail $v \in A_{\leftarrow}$ that is on the t -side of Z (there are $|A_{\leftarrow}| \leq \lambda \leq k$ options, so we are correct with probability at least k^{-1}). Let $G' = G + \{(v, t)\}$ and note that $\lambda_{G'}(s, t) > \lambda_G(s, t)$ as C_{\leftarrow} is the st -mincut closest to s . We recurse on $((G', s, t, k), \mathcal{P}, \kappa)$, obtaining a pair

$(A', \widehat{\mathcal{P}}')$ and return $(A' \cup \{(v, t)\}, \widehat{\mathcal{P}}')$. Clearly, if the guess is correct, Z remains a star st -cut in G' and, if furthermore the output $(A', \widehat{\mathcal{P}}')$ is compatible with Z in G' , then $(A' \cup \{(v, t)\}, \widehat{\mathcal{P}}')$ is compatible with Z in G . The probability that we correctly entered this case is at least $(2k)^{-1}$, and in the recursive call the value $\lambda_G(s, t)$ increased.

In the second case, we define G' as G with the whole s -side of C_{\leftarrow} contracted onto s , recurse on G' (with the flow paths of \mathcal{P} shortened to start from the edges of C_{\leftarrow}), obtaining a pair $(A', \widehat{\mathcal{P}}')$, and return $A := A' \cup \{(s, v) \mid v \in A_{\leftarrow}\}$ and $\widehat{\mathcal{P}}$ being the flow $\widehat{\mathcal{P}}'$ with every path potentially prepended with the appropriate edge (s, v) , $v \in A_{\leftarrow}$. If all tails of edges of C_{\leftarrow} are on the s -side of Z , then we did not contract any edge of $\text{core}(Z)$, $\text{core}(Z)$ remains a minimal st -cut in G' and $Z' := Z \cap E(G')$ remains a star st -cut in G' . It is straightforward to observe if furthermore $(A', \widehat{\mathcal{P}}')$ is compatible with Z' in G' , then $(A, \widehat{\mathcal{P}})$ is compatible with Z in G . Furthermore, in the recursive call the st -mincut closest to s in G' is $\delta^+(s)$ (being the image of C_{\leftarrow}).

We perform a symmetric process if $C_{\rightarrow} \neq \delta^-(t)$. That is, let A_{\rightarrow} be the set of heads of edges of C_{\rightarrow} , except for t . If there is a $v \in A_{\rightarrow}$ that is in the s -side of Z , we guess so (with probability 0.5), guess v (with probability at least k^{-1}), recurse on $G' = G + \{(s, v)\}$, obtaining a pair $(A', \widehat{\mathcal{P}}')$ and return $(A' \cup \{(s, v)\}, \widehat{\mathcal{P}}')$. As before, in the recursive call $\lambda_{G'}(s, t) > \lambda_G(s, t)$ as C_{\rightarrow} is the st -mincut closest to t . Otherwise, we guess that this is the case (with probability 0.5) and contract the t -side of C_{\rightarrow} onto t . We observe that if the guess is correct, then no edge of Z is contracted onto t , Z remains a star st -cut in the resulting graph G' . Similarly as before, we recurse on G' , obtaining a pair $(A', \widehat{\mathcal{P}}')$ and return $A := A' \cup \{(v, t) \mid v \in A_{\rightarrow}\}$ and $\widehat{\mathcal{P}}$ constructed from $\widehat{\mathcal{P}}'$ by appending an appropriate arc (v, t) , $v \in A_{\rightarrow}$ to some of the paths. In the recursive call the st -mincut closest to t is $\delta^-(t)$ (being the image of C_{\rightarrow}).

As a result, we either already recursed and returned a pair $(A, \widehat{\mathcal{P}})$ (being a correct guess with probability $\Omega(k^{-1})$) or we are left with the graph where $\delta^+(s)$ and $\delta^-(t)$ are st -mincuts.

As a last preprocessing step, we check if G contains an arc (s, t) . If this is the case, then clearly $(s, t) \in Z$ for any st -cut Z and (s, t) is one of the paths in \mathcal{P} , say P_i . We recurse on $G' := G - \{(s, t)\}$ with $\mathcal{P} \setminus \{P_i\}$, parameter $k - 1$ instead of k , and $\kappa - 1$ instead of κ , obtaining a pair $(A', \widehat{\mathcal{P}}')$. Furthermore, $Z \setminus \{(s, t)\}$ remains a star (s, t) -cut in G' . Thus, we can safely return $(A', \widehat{\mathcal{P}}' \cup \{P_i\})$ in this case.

Hence, we are left with \mathcal{I} being an instance with proper boundaries. We compute the reachability pattern H of $(\mathcal{I}, \mathcal{P})$ and the H -sequence of mincuts C_1, C_2, \dots, C_ℓ . We have $C_1 = \delta^+(s)$ and, by Lemma 4.3, $\ell \geq 2$. We split into three cases, tackled in the next three subsections:

- $|E(H)| = |V(H)|$, that is, H contains only self-loops at every vertex;
- $\ell \leq \ell^{\text{big}}$, that is, the H -sequence C_1, \dots, C_ℓ is short;
- $\ell > \ell^{\text{big}}$, that is, the H -sequence C_1, \dots, C_ℓ is long.

4.3 Base case: only self-loops in H

We now deal with the base case $|E(H)| = |V(H)|$, that is, H consist of λ vertices with self-loops and nothing else.

We have the following observation.

Lemma 4.5. *Let $(\mathcal{I}, \mathcal{P})$ be an instance with maximum flow with proper boundaries and let H be its reachability pattern. Assume $|V(H)| = |E(H)|$. For every $i \in [\lambda]$, $v \in V(P_i) \setminus \{t\}$, and a bottleneck edge e on P_i , if e lies before v on P_i , then any path from s to v in G visits e . Consequently, any set consisting of one bottleneck edge from each path P_i is an st -mincut in G .*

Proof. For the first claim, let Q be a path from s to v in G avoiding e . By the assumption on proper boundaries, $t \notin V(Q)$. If Q contains a vertex of $V(P_j) \setminus \{s, t\}$ for some $j \neq i$, then a minimal subpath of Q from a vertex of $\bigcup_{j \neq i} V(P_j) \setminus \{s, t\}$ to a vertex of $V(P_i) \setminus \{s, t\}$ is a path also in G^P and thus contradicts the assumption that H contains only self-loops.

Let C be an st -mincut witnessing that e is a bottleneck edge. Since e is before v on P_i , the concatenation of Q and a subpath of P_i from v to t is an st -path disjoint with C , a contradiction. This finishes the proof of the first claim.

For the second claim, pick a bottleneck edge $e_i \in E(P_i)$ on every path P_i and let $Y = \{e_i \mid i \in [\lambda]\}$. Assume Y is not an st -mincut. This implies that there exists a path Q in G that starts on a path P_i in a vertex v before e_i , ends on a path P_j in a vertex u after e_j and does not contain any edge of Y nor any internal vertex on paths of \mathcal{P} . In particular, Q does not contain any edge of \mathcal{P} .

Note that if $v \neq s$ and $u \neq t$, then $(i, j) \in E(H)$ by the definition of the graph H . Hence, either $i = j$, or $v = s$, or $u = t$. Note that if $v = s$ or $t = u$ we could have chosen $i = j$ anyway, so it suffices to consider the case $i = j$. Prepend Q with a subpath of P_i from s to v , obtaining a path Q' from s to u avoiding e_i . This contradicts the first claim. \square

Lemma 4.6. *Let $(\mathcal{I}, \mathcal{P})$ be an instance with maximum flow with proper boundaries and let H be its reachability pattern. Assume $|V(H)| = |E(H)|$ and let Z be a star st -cut in G . If $|\text{core}(Z)| > \lambda_G(s, t)$, then there exists $i \in [\lambda]$ such that no edge of $Z \cap E(P_i)$ is a bottleneck edge. Otherwise, if $|\text{core}(Z)| = \lambda_G(s, t)$, then no edge of $Z - \text{core}(Z)$ is a bottleneck edge of G .*

Proof. If there is $i \in [\lambda]$ such that no edge of $Z \cap E(P_i)$ is a bottleneck edge, then $\text{core}(Z)$ is not an st -mincut and we are done. For every $i \in [\lambda]$, arbitrarily pick one bottleneck edge $e_i \in Z \cap E(P_i)$ and denote $Z' = \{e_i \mid i \in [\lambda]\}$. By Lemma 4.5, Z' is an st -mincut. Since Z is a star st -cut, this proves that there was only one choice of e_i for every $i \in [\lambda]$, as desired. \square

We proceed as follows. If $\kappa = \lambda$, with probability 0.5 guess that $\text{core}(Z)$ is an st -mincut in G . In this case, for every $i \in [\lambda]$ proceed as follows. Let $(u_{i,1}, v_{i,1}), \dots, (u_{i,a_i}, v_{i,a_i})$ be the bottleneck edges of P_i , in the order of their appearance on P_i . Denote $v_{i,0} = s, u_{i,a_i+1} = t, A_i = \{(v_{i,b}, u_{i,b+1}) \mid 0 \leq b \leq a_i\}$ and P'_i to be the path consisting of edges $(s, u_{i,1}), (u_{i,1}, v_{i,1}), (v_{i,1}, u_{i,2}), \dots, (u_{i,a_i}, v_{i,a_i}), (v_{i,a_i}, t)$. We return $(A := \bigcup_{i \in [\lambda]} A_i, \widehat{\mathcal{P}} = \{P'_i \mid i \in [\lambda]\})$. By Lemma 4.6, $(A, \widehat{\mathcal{P}})$ is compatible with Z .

If $\kappa > \lambda$, or $\kappa = \lambda$ but (with the remaining probability 0.5) we guessed that $\text{core}(Z)$ is not an st -mincut in G , we randomly guess $i \in [\lambda]$, aiming at $Z \cap E(P_i)$ containing no bottleneck edge. We set A_0 to be the copies of all bottleneck edges of P_i , recurse on $G' := G + A_0$, obtaining a pair $(A', \widehat{\mathcal{P}}')$, and return $(A := A' \cup A_0, \widehat{\mathcal{P}}')$. Clearly, if the guess is correct, A_0 is compatible with Z and Z remains an star st -cut in G' . Furthermore, if $(A', \widehat{\mathcal{P}}')$ is compatible with Z in G' , then $(A, \widehat{\mathcal{P}}')$ is compatible with Z in G . Since we added a copy of every bottleneck edge on one flow path, $\lambda_{G'}(s, t) > \lambda_G(s, t)$. This finishes the description of this case.

4.4 Small ℓ case

In the next case we assume $|E(H)| > |V(H)|$ and $\ell \leq \ell^{\text{big}}$.

If there is no star st -cut Z of size at most k with $|\text{core}(Z)| \geq \kappa$, then any pair $(A, \widehat{\mathcal{P}})$ with $\lambda_{G+A}(s, t) \geq \kappa$ is a valid outcome, so we can assume at least one such star st -cut exists. For the sake of analysis, fix one star st -cut Z with $|Z| \leq k$ and $|\text{core}(Z)| \geq \kappa$.

Let $B = \bigcup_{i=1}^{\ell} V(C_i)$, that is, B is the set of endpoints of the cuts C_1, \dots, C_ℓ . Let B_{\leftarrow} be the set of elements of B on the s -side of Z and B_{\rightarrow} be the set of vertices of B on the t -side of Z .

For every $i \in [\lambda]$, we guess the last vertex v_i of B_{\leftarrow} (there is always one, as s is a candidate) and the first vertex u_i of B_{\rightarrow} on P_i (or guess $u_i = \perp$ meaning that there is no such vertex). There are at most $2^{\ell^{\text{big}}}(2^{\ell^{\text{big}}} + 1)$ options per index i , so in total the success probability is $2^{-\mathcal{O}(k \log k)}$.

We now consider a number of corner cases. We use the first applicable corner case, if possible, so that in subsequent corner cases we can assume that the earlier ones are not applicable.

Corner case 1: $B_{\rightarrow} = \emptyset$, that is, for every $i \in [\lambda]$ we guessed $u_i = \perp$. In other words, all endpoints of edges of $\bigcup_{a=1}^{\ell} C_a$ are on the s -side of Z . In particular, all endpoints of C_{ℓ} are on the s -side of Z and t is not a head of any edge of C_{ℓ} . Let G' be the graph G with the s -side of C_{ℓ} contracted onto s . Then, $\text{core}(Z)$ is a minimal st -cut in G' as well, $Z' := Z \cap E(G')$ is a star st -cut in G' , and the paths P_i , shortened to start from the edge of C_{ℓ} , form a maximum st -flow $\mathcal{P}_{G'}$ in G' . Furthermore, as G has proper boundaries, so has G' (the cut C_{ℓ} becomes $\delta_{G'}^+(s)$ in G'). Let H' be the reachability pattern of G' and $\mathcal{P}_{G'}$. We claim the following.

Claim 1. $E(H') \subsetneq E(H)$.

Proof. Clearly $E(H') \subseteq E(H)$, as any path in the residual graph of G' and $\mathcal{P}_{G'}$ witnessing $(i, j) \in E(H')$ is also a path in $G^{\mathcal{P}}$ and thus witnesses $(i, j) \in E(H)$.

We claim that if $E(H') = E(H)$, then $C_{\ell+1}$ would have been defined. Indeed, from the definition of H' for every $(i, j) \in E(H')$ there exist vertices $v_{i,j} \in V(P_i) \setminus \{t\}$ and $u_{i,j} \in V(P_j) \setminus \{t\}$ that are after C_{ℓ} on their P_i and P_j respectively such that $u_{i,j}$ is reachable from $v_{i,j}$ in the residual graph of G' and $\mathcal{P}_{G'}$ (and hence also in $G^{\mathcal{P}}$). If $E(H') = E(H)$, this witnesses that $\text{leader}_H(C_{\ell}, i) \neq t$ for every $i \in [\lambda]$. Since $\delta_G^-(t)$ is an st -mincut in G and thus t is not reachable from any vertex on $V(P_i) \setminus \{t\}$ in $G^{\mathcal{P}}$ for every $i \in [\lambda]$, we have $t \notin \text{RReach}(\text{leader}_H(C_{\ell}, i))$ for every $i \in [\lambda]$. Therefore, $C_{\ell+1}$ would have been defined, a contradiction. \square

We recurse on G' , $\mathcal{P}_{G'}$, k , and κ obtaining a pair $(A', \widehat{\mathcal{P}}')$. We set $A_0 = \{(s, v) \mid v \in B\}$ and return $(A := A_0 \cup A', \widehat{\mathcal{P}})$, where the every flow path of $\widehat{\mathcal{P}}$ is constructed from a flow path of $\widehat{\mathcal{P}}'$ by potentially prepending it with an appropriate edge of A_0 . Clearly, if $(A', \widehat{\mathcal{P}}')$ is compatible with Z' in G' , then $(A, \widehat{\mathcal{P}})$ is compatible with Z in G .

Corner case 2: for some $i \in [\lambda]$, there is a vertex of B_{\rightarrow} before a vertex of B_{\leftarrow} on P_i . That is, for some $i \in [\lambda]$ the guessed first vertex $u_i \in B_{\rightarrow} \cap V(P_i)$ is before the guessed last vertex $v_i \in B_{\leftarrow} \cap V(P_i)$. Define $A_0 = \{(s, v_i), (u_i, t)\}$ and observe that A_0 is compatible with Z and in $G + A_0$ there is an augmenting path for \mathcal{P} : start with the edge $(s, v_i) \in A_0$, go along reversed P_i to u_i , and finish with the edge $(u_i, t) \in A_0$. Hence, we can just recurse on $G' := G + A_0$, k , and κ , obtaining a pair $(A', \widehat{\mathcal{P}}')$ and return $(A' \cup A_0, \widehat{\mathcal{P}}')$.

Note that if Corner case 2 does not happen, from the guesses we can deduce whole sets B_{\leftarrow} and B_{\rightarrow} : for every $i \in [\lambda]$, all vertices of $B \cap V(P_i)$ up to the (guessed) last vertex of B_{\leftarrow} are in B_{\leftarrow} and the remainder of $B \cap V(P_i)$ is in B_{\rightarrow} . Hence, we can define

$$A_0 = \{(s, v) \mid v \in B_{\leftarrow}\} \cup \{(u, t) \mid u \in B_{\rightarrow}\}.$$

Note that A_0 is compatible with Z (if the guesses are correct).

In $G + A_0$, we define an st -flow $\mathcal{P}' = \{P'_i \mid i \in [\lambda]\}$ as follows: for every $i \in [\lambda]$, the flow path P'_i consists of the arc $(s, v_i) \in A_0$, a subpath of P_i from v_i to u_i , and the arc $(u_i, t) \in A_0$.

Corner case 3: $\lambda_{G+A_0}(s, t) > \lambda_G(s, t)$. In this case we can just recurse on $G' := G + A_0$, \mathcal{P}' , k , and κ , obtaining a pair $(A', \widehat{\mathcal{P}}')$, and return $(A' \cup A_0, \widehat{\mathcal{P}}')$.

If we are not in this corner case, we have that \mathcal{P}' is an st -maxflow in $G + A_0$.

Corner case 4: there is $i \in [\lambda]$ such that P'_i contains exactly one edge e of capacity 1. Then clearly $e \in \text{core}(Z)$. We can recurse on the graph $G' := G + A_0 - \{e\}$, the flow $\mathcal{P}' \setminus \{P'_i\}$, the parameter $k-1$ instead of k , and the parameter $\kappa-1$ instead of κ , obtaining a pair $(A', \widehat{\mathcal{P}}')$, and return $(A := A' \cup A_0, \widehat{\mathcal{P}}' \cup \{P'_i\})$.

Two remarks are in place. First, this corner case covers the case where $Z \cap \bigcup_{a=1}^{\ell} C_a \neq \emptyset$. Second, if we are not in this corner case, then in particular the whole C_1 is in the s -side of Z as $C_1 = \delta^+(s)$.

Main case We are left with the main case, where no of the aforementioned corner cases occur.

Since on every P_i the arcs of C_1, C_2, \dots, C_ℓ appear in this order, by the excluded Corner case 1 at least one head of an arc of C_ℓ is in B_{\rightarrow} . By the excluded Corner case 4, the tail of the said arc of C_ℓ is also in B_{\rightarrow} .

Let $a \in [\ell]$ be maximum such that all endpoints of C_a are in B_{\leftarrow} . This is well-defined as $C_1 = \delta_G^+(s)$ and all endpoints of C_1 are in B_{\leftarrow} as we are not in the Corner case 4. Also, by the assumption of this main case, $a < \ell$, that is, C_{a+1} is defined.

Let C be the st -mincut in $G + A_0$ closest to t . Since A_0 features arcs from s to all endpoints of C_a , C lies entirely in the t -side of C_a . By the definition of a , there is an arc of C_{a+1} whose endpoints lie in B_{\rightarrow} and thus have arcs to t in A_0 . Hence, C contains at least one arc whose endpoints are on the t -side of C_a and on the s -side of C_{a+1} . Furthermore, clearly C is an st -mincut in G as well (but of course not necessarily closest to t).

We randomly guess whether there is a vertex $v \in V(C)$ that is in the s -side of Z . With probability 0.5 we guess that there is such a vertex. Then, with probability 0.5 we guess if there is a head of an edge of C that is in the s -side of Z . If this is the case, we guess such a head $v \in V(C)$ (at most λ options) and otherwise we guess a tail $v \in V(C)$ that is in the s -side of Z (again, at most λ options). In the end, we obtain either an edge $(v, u) \in C$ with v on the s -side of Z and u on the t -side of Z (thus $(v, u) \in Z$) or a head v of an edge of C that is in the s -side of Z . The probability of a correct guess is at least $(4\lambda)^{-1}$.

In the first case, we recurse on $G' := G - \{(v, u)\}$ with flow $\mathcal{P} \setminus \{P_i\}$ where P_i is the flow path containing (v, u) and the parameter $k-1$, obtaining a pair $(A', \widehat{\mathcal{P}}')$. It is straightforward to check that then we can return $(A := A' \cup \{(s, v), (u, t)\}, \widehat{\mathcal{P}} := \widehat{\mathcal{P}}' \cup \{(s, v), (v, u), (u, t)\})$.

In the second case, we recurse on $G' := G + (A_0 \cup \{(s, v)\})$, k , and κ , obtaining a pair $(A', \widehat{\mathcal{P}}')$. Note that $A_0 \cup \{(s, v)\}$ is compatible with Z , if the guess is correct. Hence, we can return $(A_0 \cup \{(s, v)\} \cup A', \widehat{\mathcal{P}}')$. Furthermore, as C is the closest to t mincut of $G + A_0$, we have $\lambda_{G'}(s, t) > \lambda$.

Thus we are left with the most interesting case, guessed with probability 0.5, where all vertices of $V(C)$ are in the t -side of Z . The crucial observation now is the following.

Claim 2. *There exists $i \in [\lambda]$ such that $\text{leader}_H(C_a, i)$ lies in the t -side of C .*

Proof. By the definition of a , there exists $e \in C_{a+1}$ with an endpoint in B_{\rightarrow} . By the excluded Corner case 4, both endpoints of e are in B_{\rightarrow} ; let v be the tail of e . Clearly v is on the t -side of C because $(v, t) \in A_0$.

By the definition of C_{a+1} , there exists $i \in [\lambda]$ such that $v \in \text{RReach}(\text{leader}_H(C_a, i))$. Again by the definition of C_{a+1} , the whole $\text{RReach}(\text{leader}_H(C_a, i))$ is on the s -side of C_{a+1} , in particular $\text{leader}_H(C_a, i)$ is. Since C is an st -mincut in G as well, while v is reachable from $\text{leader}_H(C_a, i)$ in $G^{\mathcal{P}}$ and v is on the t -side of C , $\text{leader}_H(C_a, i)$ is also on the t -side of C . This finishes the proof. \square

Let G' be constructed from G by contracting first the s -side of C_a onto s and then the t -side of C onto t . Note that both C_a and C are st -mincuts in G and on every path P_i the arc of C_a appears strictly before the

arc of C . Hence, G' has proper boundaries and the paths of P_i naturally shorten to a maximum st -flow $\mathcal{P}_{G'}$ in G' . We recurse on G' , $\mathcal{P}_{G'}$, and k , obtaining a pair $(A', \widehat{\mathcal{P}}')$. We return $A = A_0 \cup A' \cup \{(v, t) \mid v \in V(C)\}$ and a flow $\widehat{\mathcal{P}}$ constructed from flow paths $\widehat{\mathcal{P}}'$ by potentially prepending them with an edge $(s, v) \in A_0$ for v being a head of an edge of C_a or adding at the end an edge $(v, t) \in A$ for v being a tail of an edge of C .

Consider a star st -cut Z in G . Assume that all vertices of B_{\leftarrow} are in the s -side of Z and vertices of B_{\rightarrow} and all endpoints of C are in the t -side of Z . Then, $\text{core}(Z)$ is a minimal st -cut in G' and $Z' := Z \cap E(G')$ is a star st -cut in G' . Furthermore, if $(A', \widehat{\mathcal{P}}')$ is compatible with Z' in G' , then $(A, \widehat{\mathcal{P}})$ is compatible with Z in G , as desired.

Let H' be the reachability pattern of G' and $\mathcal{P}_{G'}$. To bound the success probability, the following observation is crucial.

Claim 3. $E(H') \subsetneq E(H)$.

Proof. Clearly, any path witnessing $(i, j) \in E(H')$ in the residual network of G' and $\mathcal{P}_{G'}$ is also a path in $G^{\mathcal{P}}$ so it witnesses $(i, j) \in E(H)$. We need to exhibit an element of $E(H) \setminus E(H')$.

Let $i \in [\lambda]$ be the index asserted by Claim 2. Since $v := \text{leader}_H(C_a, i)$ lies in the t -side of C , while the head of the edge of $C_a \cap E(P_i)$ lies in the s -side of C , the predecessor v' of v on P_i lies in the t -side of C_a . There is a reason why v' is before $\text{leader}_H(C_a, i)$ on P_i : there exists $(i, j) \in E(H)$ such that $\text{LastReach}(v', P_j)$ is before C_a on P_j . Since v is in the t -side of C , we have that for every $u \in V(P_i) \cap V(G') \setminus \{s, t\}$ it holds that $\text{LastReach}(u, P_j)$ is before C_a on P_j . Consequently, $(i, j) \notin E(H')$, as desired. \square

Hence, in the recursive call $|E(H')|$ decreases, which is the progress giving the desired bound on the success probability. (For formal proof of the success probability lower bound, see Section 4.6.)

4.5 Large ℓ case

In the remaining case we have $|E(H)| > |V(H)|$ and $\ell > \ell^{\text{big}}$. By Lemma 4.4, H is transitive. Also, $|E(H)| > |V(H)|$ implies that $\lambda \geq 2$.

In the argumentation, we will frequently use the predicate of being in the s -side or t -side of a cut C_a . For brevity, we extend this notion to all indices $a \in \mathbb{Z}$, not only $a \in [\ell]$. That is, every vertex and edge of G is in the t -side of C_a for $a \leq 0$ and in the s -side of C_a for $a > \ell$ while neither a vertex nor an edge of G is in the s -side of C_a for $a \leq 0$ and in the t -side of C_a for $a > \ell$.

Fix a star st -cut Z of size at most k such that $|\text{core}(Z)| \geq \kappa$. Let Z_{ts} be the set of those edges $(v, u) \in \bigcup_{i=1}^{\lambda} E(P_i)$ such that v is on the t -side of Z while u is on the s -side of Z . Note that for every $i \in [\lambda]$ we have $|Z_{ts} \cap E(P_i)| \leq |Z \cap E(P_i)| - 1$ and hence $|Z_{ts}| \leq k - \lambda$.

We say that an index $a \in \mathbb{Z}$ is *touched* if there exists $e \in Z \cup Z_{ts}$ with an endpoint that is in the t -side of C_a and in the s -side of C_{a+1} . Note that a touched index a satisfies $0 \leq a \leq \ell$.

We have the following observation.

Claim 4. *There are at most $2|Z \cup Z_{ts}| \leq 4k - 2\lambda$ touched indices.*

Proof. Every endpoint v of an edge $e \in Z \cup Z_{ts}$ gives raise to at most one index a being touched, namely the maximum $a \in \mathbb{Z}$ such that v is in the t -side of C_a . Hence, there are at most $2|Z \cup Z_{ts}| \leq 4k - 2\lambda$ touched indices as desired. \square

A set $L \subseteq V(H)$ is *downward-closed* if there is no arc $(i, j) \in E(H)$ with $i \in L$ but $j \notin L$. For a set $L \subseteq V(H)$, let $\text{cl}(L)$ be the minimal superset of L that is downward-closed; since H is transitive and contains a loop at every vertex, we have

$$\text{cl}(L) = \{i \in [\lambda] \mid \exists j \in L (j, i) \in E(H)\}.$$

For $a \in [\ell]$, let $L_a \subseteq [\lambda]$ be the set of those indices for which both endpoints of the unique edge of $C_a \cap E(P_i)$ lie in the s -side of Z . We also define $L_a = [\lambda]$ for $a \leq 0$ and $L_a = \emptyset$ for $a > \ell$.

We have established a few properties of the objects defined above.

Claim 5. *Let $L \subseteq [\lambda]$ be downward closed. Then G admits no path*

$$\text{from } \bigcup_{i \in L} V(P_i) \setminus \{s, t\} \text{ to } \bigcup_{i \in [\lambda] \setminus L} V(P_i) \setminus \{s, t\}.$$

Proof. Assume the contrary, let Q be a shortest such path. By minimality, no internal vertex of G lies in $\bigcup_{i \in [\lambda]} V(P_i) \setminus \{s, t\}$. Since G has proper boundaries, neither s nor t lies on Q . Hence, Q is a path in $G^{\mathcal{P}}$, contradicting the definition of H and the assumption that L is downward-closed. \square

Claim 6. *Let $a \leq b$ be two integers. Then $L_b \subseteq \text{cl}(L_a)$. In particular, if L_a is downward-closed, then $L_b \subseteq L_a$.*

Proof. The claim is obvious if $a < 1$ (as then $L_a = [\lambda]$) or $b > \ell$ (as then $L_b = \emptyset$), so assume $1 \leq a \leq b \leq \ell$.

For every $i \in [\lambda] \setminus \text{cl}(L_a)$, pick an endpoint v_i of the unique edge of $C_a \cap E(P_i)$ that is in the t -side of Z and let X_i be the set of vertices of $V(P_i) \setminus \{t\}$ that lie after v_i on P_i . By Claim 5 applied to $L := \text{cl}(L_a)$, for every vertex $v \in \bigcup_{i \in [\lambda] \setminus \text{cl}(L_a)} X_i$, any path from s to v of G passes through $\{v_i \mid i \in [\lambda] \setminus \text{cl}(L_a)\}$. We infer that such v is in the t -side of Z . Hence, $L_b \subseteq \text{cl}(L_a)$ as desired. \square

Claim 7. *Assume $a < b$ are two integers such that $b - a \geq \lambda - 1$ and for every $a \leq c < b$, c is untouched. Then, $L_b = \text{cl}(L_a)$. In particular, L_b is downward-closed.*

Proof. Clearly, $L_b \subseteq \text{cl}(L_a)$ by Claim 6. Also, since every c with $a \leq c < b$ is untouched, for every $i \in L_a$ no edge of P_i between the edge of C_a and the edge of C_b is in Z , so $L_a \subseteq L_b$.

If $a < 1$, then $L_a = [\lambda]$ and we have $L_b = [\lambda]$ as well. If $b > \ell$, then $L_b = \emptyset$ so from $L_a \subseteq L_b$ it follows that $L_a = \emptyset$ as well.

Hence, we can assume $1 \leq a < b \leq \ell$. Since $L_a \subseteq L_b \subseteq \text{cl}(L_a)$, it suffices to show that if for some $1 \leq c < \ell$ we have that c is untouched and L_c is not downward-closed, then L_{c+1} is a strict superset of L_c .

Clearly, as c is untouched $L_c \subseteq L_{c+1}$. Since L_c is not downward-closed, there is $(i, j) \in E(H)$ with $i \in L_c$ but $j \notin L_c$. By the construction of C_{c+1} , there exists $v \in V(P_i)$ and $u \in V(P_j)$, both in the t -side of C_c and s -side of C_{c+1} and a path Q in $G^{\mathcal{P}}$ from v to u . Pick i, j , and Q as above so that the length of Q is minimum possible.

Since in $G^{\mathcal{P}}$ one cannot cross neither C_c nor C_{c+1} from the s -side to the t -side, all vertices of Q are in the t -side of C_c and s -side of C_{c+1} . By the minimality of Q , no internal vertex of Q lies on any path P_ι , $\iota \in [\lambda]$. Hence, Q is a path in G as well. Since c is untouched, no edge of Q lies in Z . Also, no edge of P_i nor P_j between C_c and C_{c+1} (including C_{c+1}) lies in Z . We infer that $j \in L_{c+1}$, as desired. This finishes the proof of the claim. \square

Claim 8. *Assume $a \in [\ell]$ is such that L_a is downward-closed. Let $i \in [\lambda]$ and e be the unique edge of $C_a \cap E(P_i)$.*

- If $i \notin L_a$, then the following holds. The entire suffix of P_i from the head of e to t is in the t -side of Z . In particular, no edge of the said suffix is in Z .
- If $i \in L_a$, then the following holds. The entire prefix of P_i from s to the tail of e is in the s -side of $\text{core}(Z)$. In particular, no edge of the said prefix is in $\text{core}(Z)$.

Proof. We start with the first claim. Let $v \in V(P_i)$ be in the s -side of Z and in t -side of C_a . By the definition of the s -side, there is a path Q from s to v in $G - Z$.

Since v is in the t -side of C_a , there is an edge of C_a on Q . Assume $j \in [\lambda]$ is such that the edge of $E(P_j) \cap C_a$ lies on Q . Since Q is a path in $G - Z$ starting in s , it is entirely in the s -side of Z . By the definition of L_a , we have $j \in L_a$. However, as $v \in V(P_i)$ and $i \notin L_a$, the subpath of Q from the edge of $C_a \cap E(P_j)$ to v contradicts Claim 5.

We now move to the second claim. By contradiction, assume there is a vertex on P_i that is in the t -side of $\text{core}(Z)$ and s -side of C_a . Let v be the earliest (on P_i) such vertex. Since v is in the t -side of $\text{core}(Z)$, $v \neq s$. By the choice of v , the predecessor of v on P_i is in the s -side of Z , that is, the edge on P_i with the head v is in $\text{core}(Z)$. By the minimality of $\text{core}(Z)$, there is a path Q from v to t in $G - \text{core}(Z)$.

Since v is in the s -side of C_a , there is an edge of C_a on Q . Assume $j \in [\lambda]$ is such that the edge of $E(P_j) \cap C_a$ lies on Q . Since Q is a path in $G - \text{core}(Z)$ ending in t , it is entirely in the t -side of $\text{core}(Z)$. By the definition of L_a , we have $j \notin L_a$. However, as $v \in V(P_i)$ and $i \in L_a$, the subpath of Q from v to the edge of $E(P_j) \cap C_a$ contradicts Claim 5. \square

Claim 9. Let $a \in [\ell]$ be such that for every $a - (\lambda - 1) \leq c < a$ the index c is untouched. Then, for every $i \in L_a$ there exists in $G - Z$ a path from s to the tail of the unique edge of $C_a \cap E(P_i)$ whose all vertices lie in the s -side of C_a .

Proof. If $a - (\lambda - 1) \leq 1$ (i.e., $a \leq \lambda$), then there is no endpoint of the edge of Z in the s -side of C_a (note that if 0 is touched, then so is 1) and hence the prefix of P_i from s to C_a satisfies the desired properties. We henceforth assume then that $b := a - (\lambda - 1) > 1$.

Let $e = (u, v)$ be the unique edge of $C_a \cap E(P_i)$. Since $i \in L_a$, both u and v are in the s -side of Z . Let Q be a path from s to u in $G - Z$.

Since Q ends in u , Q contains an edge of C_b . Let e' be the first edge of C_b on Q and let $j \in [\lambda]$ be such that $e' \in E(P_j) \cap C_b$. Since Q is in the s -side of Z , we have $j \in L_b$.

For $b < c \leq a$, let L'_c be the family of indices $i' \in [\lambda]$ such that there is a path from the head of e' to the tail of the unique edge of $C_c \cap E(P_{i'})$ with all vertices in the t -side of C_b and s -side of C_c . We also define $L'_b = \{j\}$. Since for every $b \leq c < a$, c is untouched, we have that any path witnessing $i' \in L'_c$ is contained completely in the s -side of Z and hence $L'_c \subseteq L_c$. Also, $L'_c \subseteq L'_{c'}$ for any $b \leq c \leq c' \leq a$.

Note that it suffices to prove that $i \in L'_c$ for some $b \leq c \leq a$, as then the desired path can be formed by concatenating the prefix of Q up to e' , the path witnessing $i \in L'_c$, and the part of P_i from C_c to u . As $b = a - (\lambda - 1)$, it suffices to show that if $i \notin L'_c$ for some $b \leq c < a$, then L'_{c+1} is a proper superset of L'_c .

Fix $b \leq c < a$ such that $i \notin L'_c$. As $j \in L'_c$ and $e' \in E(P_j)$, there is a subpath of Q , between e' and u , that starts in a vertex of some $P_{j'}$, $j' \in L'_c$, ends in a vertex of some $P_{i'}$, $i' \notin L'_c$, and contains no internal vertices on paths of \mathcal{P} . This path witnesses that $(j', i') \in E(H)$ and thus there is a path R in $G^{\mathcal{P}}$ from a vertex of $P_{j'}$ to a vertex of $P_{i'}$ that lies completely in the t -side of C_c and s -side of C_{c+1} . A path R contains a subpath R' with no internal vertices on \mathcal{P} that starts on a path $P_{j''}$ for some $j'' \in L'_c$ and ends on a path $P_{i''}$ for some $i'' \notin L'_c$. This path is present in G . Consequently, by using a subpath of $P_{j''}$ from C_c to the start or R' , the path R' , and a subpath of $P_{i''}$ from the end of R' to C_{c+1} we witness that $i'' \in L'_{c+1}$, as desired. \square

Claims 7, 8, and 9 motivate the following definition. An index $a \in [\ell]$ is a *milestone* if for every $a - (\lambda - 1) \leq c \leq a$, c is untouched. Claim 7 implies for a milestone a , the set L_a is downward-closed.

An arc (v, u) is a *long backarc* if there exists a milestone a with v in the t -side of C_a and u in the s -side of C_a . Not all assertions of the following observations are used later, but we present them anyway as they give good intuition about why the long backarcs are essentially irrelevant for the problem.

Claim 10. *Let \mathbf{L} be the family of long backarcs. Then,*

- \mathcal{P} is (still) a maximum st -flow in $G - \mathbf{L}$;
- $G - \mathbf{L}$ is (still) properly bounded;
- the reachability pattern of $(G - \mathbf{L}, \mathcal{P})$ is (still) H ;
- C_1, C_2, \dots, C_ℓ is (still) an H -sequence of mincuts in $(G - \mathbf{L}, H)$;
- $\text{core}(Z)$ is disjoint with \mathbf{L} and is (still) a minimal st -cut in $G - \mathbf{L}$;
- $Z' \subseteq Z$, consisting of those arcs of $Z \setminus \mathbf{L}$ whose tail is reachable from s in $G - (Z \cup \mathbf{L})$, contains $\text{core}(Z)$ and is a star st -cut in $G - \mathbf{L}$.

Proof. The first two properties are immediate as \mathbf{L} does not contain any edge of \mathcal{P} . For the third and fourth properties, recall that $\ell > \ell^{\text{big}} > 2$ and notice that when defining C_a for $1 < a \leq \ell$, all paths in $G^{\mathcal{P}}$ that witness where $\text{leader}_H(C_{a-1}, i)$ is for $i \in [\lambda]$ are contained in the t -side of C_{a-1} and in the s -side of C_a . Hence, these paths do not use any edge of \mathbf{L} .

For the penultimate property, clearly $Z \setminus \mathbf{L}$ is an st -cut in $G - \mathbf{L}$. Let $e \in \text{core}(Z)$. Since $\text{core}(Z)$ is a minimal st -cut, there exists a path Q from s to t whose only intersection with $\text{core}(Z)$ is e . Pick such a path Q that minimizes the number of edges of \mathbf{L} . To prove the last property, it suffices to show that there is no long backarc on Q (which in particular implies that $e \notin \mathbf{L}$).

To this end, we use Claim 8. Assume that Q contains at least one edge of \mathbf{L} and let $f = (v, u)$ be the last such edge. Let a be the milestone index witnessing that $f \in \mathbf{L}$. Since v is in the t -side of C_a and u is in the s -side of C_a , Q contains an edge e_1 of C_a before f and an edge e_2 of C_a after f . Let $i_1, i_2 \in [\lambda]$ be such that $e_1 \in E(P_{i_1})$ and $e_2 \in E(P_{i_2})$.

If $e_1 \notin L_a$, then e is not later on Q than e_1 and, by Claim 8, we can substitute the suffix of Q from the head of e_1 to t with the suffix of P_{i_1} from the head of e_1 to t , contradicting the minimality of Q . If $e_2 \in L_a$, then e_2 is before e on Q and, by Claim 8, we can substitute the prefix of Q from s to the tail of e_2 with the prefix of P_{i_2} from s to the tail of e_2 , again contradicting the minimality of Q . Hence, $e_1 \in L_a$ but $e_2 \notin L_a$. However, as e_1 is before e_2 on Q , the subpath of Q from e_1 to e_2 contradicts Claim 5.

For the final property, note that the penultimate property implies $\text{core}(Z) \subseteq Z'$ and thus Z' is an st -cut. Since every arc of $Z \setminus Z'$ has its tail not reachable from s in $G - (Z \cup \mathbf{L})$, the sets of vertices reachable from s in $G - (Z \cup \mathbf{L})$ and $G - (Z' \cup \mathbf{L})$ are equal. Hence, for every $e \in Z'$, the tail of e is reachable from s and the head of e is not reachable from s in $G - (Z' \cup \mathbf{L})$. This finishes the proof that Z' is a star st -cut. \square

We remark here that it is possible that $\mathbf{L} \cap Z \neq \emptyset$ or $Z' \neq Z \setminus \mathbf{L}$; Claim 10 only asserts that the properties of $\text{core}(Z)$ remain unchanged upon deletion of \mathbf{L} .

We say that two touched indices a and b are *close* if $|b - a| \leq \lambda$. Let $0 \leq a_1 < a_2 < \dots < a_r \leq \ell$ be the touched indices; by Claim 4 we have $r \leq 4k - 2\lambda$. A *block* is a maximal subsequence $a_\alpha, a_{\alpha+1}, \dots, a_\beta$ such that a_i and a_{i+1} are close for every $\alpha \leq i < \beta$. Note that we have $a_\beta - a_\alpha \leq (\beta - \alpha)\lambda \leq (r - 1)\lambda$.

Naturally, the sequence a_1, \dots, a_r partitions into a number of blocks. An index $a \in \mathbb{Z}$ is *interesting* if $a_\alpha - \lambda \leq a \leq a_\beta + \lambda$ for some block a_α, \dots, a_β . Note that this condition is equivalent to $|a - a_\alpha| \leq \lambda$ for some $\alpha \in [r]$. We infer that the number of interesting indices is bounded by $r \cdot (2\lambda + 1) \leq 12k\lambda$.

We randomly sample a subset $\Gamma \subseteq \{0, 1, \dots, \ell\}$ in the following skewed way: every $a \in \{0, 1, \dots, \ell\}$ belongs to Γ with probability k^{-1} , independently of the other indices. We aim at the following: for every interesting index a , $a \in \Gamma$ if and only if a is touched. Note that no integer outside the set $\{0, \dots, \ell\}$ is touched, there are at most $4k - 2\lambda$ touched indices and at most $12k\lambda$ interesting indices. Thus we are successful with probability at least

$$k^{-(4k-2\lambda)} \cdot (1 - k^{-1})^{12k\lambda} = 2^{-O(k \log k)}.$$

A Γ -*milestone* is an index $a \in [\ell]$ such that for every $a - (\lambda - 1) \leq c \leq a$, $c \notin \Gamma$. Note that if the guess is correct, every Γ -milestone is a milestone.

A Γ -*block* is a maximal sequence $b_1 < b_2 < \dots < b_\zeta$ of elements of Γ such that b_i is close to b_{i+1} for $1 \leq i < \zeta$. We perform the following cleaning procedure: for every Γ -block $b_1 < b_2 < \dots < b_\zeta$ such that $\zeta > 4k - 2\lambda$ or $b_\zeta - b_1 > (4k - 2\lambda - 1)\lambda$, we delete all elements of the said block from Γ . If we are successful, then even after the cleaning procedure every block is a Γ -block, but there may be numerous Γ -blocks that are not blocks. Furthermore, if we are successful and $(b_1 < \dots < b_\zeta)$ is a Γ -block, then both $b_1 - 1$ (if $b_1 > 1$) and $b_\zeta + \lambda$ (if $b_\zeta + \lambda \leq \ell$) are Γ -milestones (and thus in particular actual milestones).

Let $\mathbf{B}^1, \mathbf{B}^2, \dots, \mathbf{B}^\xi$ be the sampled Γ -blocks. For a Γ -block $\mathbf{B}^\alpha = (b_1 < b_2 < \dots < b_\zeta)$, we define $b_{\zeta-}^\alpha = b_1$, $b_{\zeta+}^\alpha = b_\zeta$, and let Z_0^α be the set of edges of Z with both endpoints on the t -side of C_{b_1} and s -side of $C_{b_\zeta+\lambda}$.

We henceforth assume that the guess of Γ is a correct one. We have the following observations.

Claim 11. *For every $e \in Z \setminus \mathbf{L}$ there exists a Γ -block \mathbf{B}^α that is an actual block such that both endpoints of e are in the t -side of $C_{b_{\zeta-}^\alpha}$ and in the s -side of $C_{b_{\zeta+}^\alpha+1}$.*

Proof. Let a be maximum such that the head of e is in the t -side of C_a . Such a exists and $a \geq 1$ as $a = 1$ is one of the candidates. Clearly, a is touched. Let \mathbf{B}^α be the Γ -block (and an actual block) where a lies. It suffices to prove that the tail of e lies in the t -side of $C_{b_{\zeta-}^\alpha}$ and in the s -side of $C_{b_{\zeta+}^\alpha+1}$.

By assumption, e is not a long backarc. If $b_{\zeta-}^\alpha > 1$ then clearly also e cannot have a tail in the s -side of $C_{b_{\zeta-}^\alpha-1}$ as the head of e is in the t -side of $C_{b_{\zeta-}^\alpha}$. If $b_{\zeta+}^\alpha + \lambda \leq \ell$, then $b_{\zeta+}^\alpha + \lambda$ is a milestone and hence e does not have its tail on the t -side of $C_{b_{\zeta+}^\alpha+\lambda}$. We infer that actually both endpoints of e are on the t -side of $C_{b_{\zeta-}^\alpha-1}$ and on the s -side of $C_{b_{\zeta+}^\alpha+\lambda}$. By the definition of being touched, this implies that both endpoints of e are in the t -side of $C_{b_{\zeta-}^\alpha}$ and in the s -side of $C_{b_{\zeta+}^\alpha+1}$. \square

Claim 11 asserts that $(Z_0^\alpha)_{\alpha \in [\xi]}$ is a partition of $Z - \mathbf{L}$.

Claim 12. *For every \mathbf{B}^α , the following holds.*

- $L_{b_{\zeta-}^\alpha-1} = L_{b_{\zeta+}^\alpha+\lambda}$ if \mathbf{B}^α is not an actual block or $\text{core}(Z) \cap Z_0^\alpha = \emptyset$
- $L_{b_{\zeta-}^\alpha-1} \supsetneq L_{b_{\zeta+}^\alpha+\lambda}$ if \mathbf{B}^α is an actual block and $\text{core}(Z) \cap Z_0^\alpha \neq \emptyset$.

Proof. Claim 7 implies that both $L' := L_{b_{\zeta-}^\alpha-1}$ and $L'' := L_{b_{\zeta+}^\alpha+\lambda}$ are downward-closed (recall that $L_a = [\lambda]$ for $a \leq 0$ and $L_a = \emptyset$ for $a > \ell$). Then, Claim 6 implies $L'' \subseteq L'$.

Assume first $L' \neq L''$ and pick $i \in L' \setminus L''$. Consider the path P_i . By Claim 8, there is no edge of $\text{core}(Z)$ on P_i neither before $C_{b_{\zeta-}^\alpha-1}$ nor after $C_{b_{\zeta+}^\alpha+\lambda}$. As $\text{core}(Z)$ is a minimal st -cut, we infer that there

is an edge of $\text{core}(Z)$ between $C_{b_{\leftarrow}^{\alpha}-1}$ and $C_{b_{\rightarrow}^{\alpha}+\lambda}$. By Claim 11, both endpoints of the said edge is between $C_{b_{\leftarrow}^{\alpha}}$ and $C_{b_{\rightarrow}^{\alpha}+1}$. Hence, Z_0^{α} contains an edge of $\text{core}(Z)$.

Assume now $L' = L''$ and, by contradiction, let e be any edge of $\text{core}(Z) \cap Z_0^{\alpha}$. That is, both endpoints of e are in the t -side of $C_{b_{\leftarrow}^{\alpha}}$ and in the s -side of $C_{b_{\rightarrow}^{\alpha}+1}$.

By the definition of $\text{core}(Z)$, there exists an st -path Q whose only intersection with Z is the edge e . Let $f = (v, u)$ be the first edge of $C_{b_{\leftarrow}^{\alpha}}$ on Q , let $i \in [\lambda]$ be such that $f \in E(P_i)$, and let Q_1 be the prefix of Q from s to v . Since Q_1 is contained in the s -side of $C_{b_{\leftarrow}^{\alpha}}$, while e is in the t -side of $C_{b_{\leftarrow}^{\alpha}}$, Q_1 is a path in $G - \text{core}(Z)$ as well. Since $b_{\leftarrow}^{\alpha} - 1$ is untouched, $f \notin \text{core}(Z)$ and thus u is in the s -side of $\text{core}(Z)$. This witnesses that $i \in L_{b_{\leftarrow}^{\alpha}}$ and hence $L' \neq \emptyset$. Since $L' = L''$, it holds that $L_{b_{\rightarrow}^{\alpha}+\lambda} \neq \emptyset$. In particular, $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$ and thus $C_{b_{\rightarrow}^{\alpha}+\lambda}$ is defined.

Recall that Q contains e . Since e has its head in the s -side of $C_{b_{\rightarrow}^{\alpha}+1}$, Q visits an edge of $C_{b_{\rightarrow}^{\alpha}+\lambda}$ after e . Let f' be the last edge of $C_{b_{\rightarrow}^{\alpha}+\lambda}$ on Q and let Q_2 be the suffix of Q starting from the head of f' . Clearly, f' is after e on Q so e does not lie on Q_2 . Also, Q_2 is contained completely on the t -side of $C_{b_{\rightarrow}^{\alpha}+\lambda}$ by the choice of f' . Hence, Q_2 is also a path in $G - \text{core}(Z)$. Consequently, as $\text{core}(Z)$ is an st -cut, f' lies on a path P_j for some $j \notin L_{b_{\rightarrow}^{\alpha}+\lambda} = L''$. However, then the subpath Q_3 of Q from the head of f to the tail of f' contradicts Claim 5. This is the desired contradiction. \square

Let $\alpha_1 < \alpha_2 < \dots < \alpha_{\eta}$ be the indices of Γ -blocks that are actual blocks. We randomly guess the integer $1 \leq \eta \leq 4k - 2\lambda$. We sample a sequence $\mathcal{L} = ([\lambda] = L_Z^0 \supseteq L_Z^1 \supseteq \dots \supseteq L_Z^{\eta} = \emptyset)$ of downward-closed sets, aiming at $L_Z^{\iota-1} = L_{b_{\leftarrow}^{\alpha_{\iota}}-1}$ and $L_Z^{\iota} = L_{b_{\rightarrow}^{\alpha_{\iota}}+\lambda}$ for every $1 \leq \iota \leq \eta$. We define $J \subseteq [\eta]$ to be the set of those indices $\iota \in [\eta]$ such that $L_Z^{\iota-1} \subsetneq L_Z^{\iota}$.

For every $1 \leq \alpha \leq \xi - 1$, we sample a set $L^{\alpha} \in \mathcal{L}$, and further denote $L^0 = [\lambda]$ and $L^{\xi} = \emptyset$. We aim that for every $1 \leq \iota \leq \eta$ actually $L^{\alpha_{\iota}-1} = L_Z^{\iota-1}$ and $L^{\alpha_{\iota}} = L_Z^{\iota}$.

In other words, for every $1 \leq \alpha \leq \xi - 1$, the sampled set L^{α} is a guess for the downward-closed set $L_{b_{\rightarrow}^{\alpha}+\lambda}$ (and thus $L^{\alpha-1}$ is also a guess for the downward-closed set $L_{b_{\leftarrow}^{\alpha}-1}$). We aim at being correct in this guess around (just before and just after) Γ -blocks that are actual blocks.

If this is the case, then consider a single Γ -block \mathbf{B}^{α} . If there is an index $1 \leq \iota \leq \eta$ such that $L^{\alpha-1} = L_Z^{\iota-1}$ and $L^{\alpha} = L_Z^{\iota}$, then it is possible that \mathbf{B}^{α} is an actual block and $\alpha = \alpha_{\iota}$. There may be multiple such blocks α for a single value of ι and we will treat them in the same way, considering them candidates for $\mathbf{B}^{\alpha_{\iota}}$. On the other hand, if for an index α no such index ι exists, we know that \mathbf{B}^{α} is not an actual block and we can add some edges to the returned set A that simply bypass \mathbf{B}^{α} .

To implement the above intuition, we need a few notions. We say that an index $1 \leq \alpha \leq \xi$ is *good* if there exists $1 \leq \iota \leq \eta$ such that $L^{\alpha-1} = L_Z^{\iota-1}$, and $L^{\alpha} = L_Z^{\iota}$. A good index α is *excellent* if additionally this ι belongs to J , that is, $L^{\alpha-1} \neq L^{\alpha}$. Note that all indices α_{ι} for $1 \leq \iota \leq \eta$ are good and α_{ι} is excellent if and only if $L_Z^{\iota-1} \neq L_Z^{\iota}$, which is equivalent to $\text{core}(Z) \cap Z_0^{\alpha_{\iota}} \neq \emptyset$ by Claim 12. Furthermore, if α is excellent, then the corresponding index $\iota \in J$ is uniquely defined and we denote it $\iota(\alpha)$.

For every excellent $1 \leq \alpha \leq \xi$, we define an instance $\mathcal{I}^{\alpha} = (G^{\alpha}, s, t, k^{\alpha})$ with a maximum flow \mathcal{P}^{α} as follows. Let $D^{\alpha} = L^{\alpha-1} \setminus L^{\alpha} = L_Z^{\iota(\alpha)-1} \setminus L_Z^{\iota(\alpha)}$ (recall that for excellent α we have $L^{\alpha-1} \supsetneq L^{\alpha}$). Define first a graph G_0^{α} that consists of:

- a subgraph of G induced by all vertices $v \in V(G)$ which are on the t -side of $C_{b_{\leftarrow}^{\alpha}-1}$ and on the s -side of $C_{b_{\rightarrow}^{\alpha}+\lambda}$, except for the vertices s and t ;
- vertices s and t (but so far no arcs incident with them)
- if $b_{\leftarrow}^{\alpha} > 1$, then for every $i \in D^{\alpha}$ an edge (s, v_i) where v_i is the head of the edge of $E(P_i) \cap C_{b_{\leftarrow}^{\alpha}-1}$;

- if $b_{\leftarrow}^{\alpha} \leq 1$, then all edges of the form (s, v_i) that lie on paths $\{P_i \mid i \in D^{\alpha}\}$;
- if $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$, then for every $i \in D^{\alpha}$ an edge (u_i, t) where u_i is the tail of the edge of $E(P_i) \cap C_{b_{\rightarrow}^{\alpha} + \lambda}$;
- if $b_{\rightarrow}^{\alpha} + \lambda > \ell$, then all edges of the form (u_i, t) that lie on paths $\{P_i \mid i \in D^{\alpha}\}$

The flow \mathcal{P}^{α} is of size $\lambda^{\alpha} := |D^{\alpha}|$ and consists of, for every $i \in D^{\alpha}$, the arc (s, v_i) , the subpath of P_i between v_i and u_i , and the arc (u_i, t) . The graph G^{α} is the subgraph of G_0^{α} induced by all vertices that are at the same time reachable from s and from which one can reach t in G_0^{α} . Note that the flow \mathcal{P}^{α} is present in G^{α} .

We define $Z^{\alpha} \subseteq Z_0^{\alpha}$ as the set of those $(u, v) \in Z$ such that either:

- $u, v \in V(G^{\alpha}) \setminus \{s, t\}$,
- $b_{\leftarrow}^{\alpha} \leq 1$ and $u = s, v = v_i$ for some $i \in D^{\alpha}$, or
- $b_{\rightarrow}^{\alpha} + \lambda > \ell$ and $u = u_i, v = t$ for some $i \in D^{\alpha}$.

We remark that $Z_0^{\alpha} \setminus Z^{\alpha}$ may be nonempty. For example, Z_0^{α} may contain an edge e on a path P_i for some $i \notin D^{\alpha}$ that has both endpoints between $C_{b_{\leftarrow}^{\alpha}}$ and $C_{b_{\rightarrow}^{\alpha}}$ and is an edge of $Z \setminus \text{core}(Z)$. This edge is not in G^{α} and hence also not in Z^{α} .

For every $\iota \in J$ we guess integers $0 \leq \kappa_{\leftarrow}^{\iota} \leq k_{\leftarrow}^{\iota} \leq k$, aiming at $\kappa_{\leftarrow}^{\iota} = |Z^{\alpha_{\iota}} \cap \text{core}(Z)|$ and $k_{\leftarrow}^{\iota} = |Z^{\alpha_{\iota}}|$. We mandate that for every $\iota \in J$ it holds that $|L_{\leftarrow}^{\iota-1} \setminus L_{\leftarrow}^{\iota}| \leq \kappa_{\leftarrow}^{\iota} \leq k_{\leftarrow}^{\iota}$ and that $\sum_{\iota \in J} k_{\leftarrow}^{\iota} \leq k$. If these conditions are not satisfied, we just return either $(A = \emptyset, \mathcal{P})$ if $\kappa = \lambda$ or $(A = \{(s, t)\}, \{(s, t)\})$ if $\kappa > \lambda$ from this recursive call and terminate. Furthermore, for every excellent $1 \leq \alpha \leq \xi$, we denote $\kappa^{\alpha} = \kappa_{\leftarrow}^{\iota(\alpha)}$ and $k^{\alpha} = k_{\leftarrow}^{\iota(\alpha)}$ for brevity. This defines the parameter k^{α} in the instance \mathcal{I}^{α} for an excellent α and a parameter κ^{α} we will pass down in the recursion.

Let us lower bound the probability that all guesses are as we aim for. The probability of guessing η correctly is $\Omega(k^{-1})$. Guessing the sequence \mathcal{L} boils down to guessing, for every $i \in [\lambda]$, the maximum index $0 \leq \iota \leq \eta$ for which $i \in L_{\leftarrow}^{\iota}$. Hence, the success probability here is at least $(\eta + 1)^{-\lambda} = 2^{-\mathcal{O}(k \log k)}$. The integers $\{\kappa_{\leftarrow}^{\iota} \mid \iota \in J\}$ and $\{k_{\leftarrow}^{\iota} \mid \iota \in J\}$ are guessed correctly with probability $2^{-\mathcal{O}(k \log k)}$ and similarly correct sets L^{α} for $\alpha \in \bigcup_{\iota=1}^{\eta} \{\alpha_{\iota} - 1, \alpha_{\iota}\}$ are guessed with probability $2^{-\mathcal{O}(k \log k)}$. Hence, overall success probability is $2^{-\mathcal{O}(k \log k)}$.

We observe the following.

Claim 13. *For every excellent $1 \leq \alpha \leq \xi$, \mathcal{P}^{α} is a maximum st -flow in G^{α} and G^{α} has proper boundaries. Furthermore, one of the following options hold:*

- either $D^{\alpha} \neq [\lambda]$ and thus $|\mathcal{P}^{\alpha}| = |D^{\alpha}| < \lambda$ and $k^{\alpha} \leq k - (\lambda - |D^{\alpha}|) < k$, or
- $D^{\alpha} = [\lambda]$, H is the reachability pattern of $(G^{\alpha}, \mathcal{P}^{\alpha})$, and $\delta_{G^{\alpha}}^{+}(s), C_{b_{\leftarrow}^{\alpha}}, C_{b_{\leftarrow}^{\alpha}+1}, \dots, C_{b_{\rightarrow}^{\alpha}+\lambda-1}, \delta_{G^{\alpha}}^{-}(t)$ is the H -sequence of mincuts in $(G^{\alpha}, \mathcal{P}^{\alpha})$ (without the first term if $b_{\leftarrow}^{\alpha} = 1$, without the last term if $b_{\rightarrow}^{\alpha} + \lambda > \ell$, and without all terms C_c for which $c > \ell$).

Proof. The claim that \mathcal{P}^{α} is a maximum st -flow in G^{α} and that G^{α} is properly bounded follows directly from how we chose edges incident with s and t in G^{α} .

Consider first the simpler case $D^{\alpha} \neq [\lambda]$. The fact that $|\mathcal{P}^{\alpha}| = |D^{\alpha}| < \lambda$ is immediate. Since we mandate $k_{\leftarrow}^{\iota} \geq |L_{\leftarrow}^{\iota-1} \setminus L_{\leftarrow}^{\iota}|$ and $\sum_{\iota=1}^{\eta} k_{\leftarrow}^{\iota} \leq k$, we have $k^{\alpha} \leq k - (\lambda - |D^{\alpha}|)$.

Consider then the case $D^{\alpha} = [\lambda]$. Observe that G_0^{α} is isomorphic to the graph G with the following modifications: if $b_{\leftarrow}^{\alpha} > 1$, then contract the s -side of $C_{b_{\leftarrow}^{\alpha}-1}$ onto s and if $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$, contract the t -side of

$C_{b_{\leftarrow}^{\alpha} + \lambda}$ onto t . Furthermore, \mathcal{P}^{α} can be then defined as the projection of \mathcal{P} shortened to the part in the t -side of $C_{b_{\leftarrow}^{\alpha} - 1}$ and in the s -side of $C_{b_{\leftarrow}^{\alpha} + \lambda}$.

To see that the reachability pattern of $(G^{\alpha}, \mathcal{P}^{\alpha})$ is still H , first note that every path in the residual graph of G^{α} and \mathcal{P}^{α} that does not use s nor t is also a path in $G^{\mathcal{P}}$, and hence the residual graph of G^{α} and \mathcal{P}^{α} is a subgraph of H . In the second direction, since the entire part of G in the t -side of $C_{b_{\leftarrow}^{\alpha}}$ and s -side of $C_{b_{\leftarrow}^{\alpha} + 1}$ lies in G_0^{α} (recall $\lambda \geq 2$), and $C_{b_{\leftarrow}^{\alpha} + 1}$ is the H -subsequent mincut to $C_{b_{\leftarrow}^{\alpha}}$, for every $(i, j) \in E(H)$ there is a path from $V(P_i) \setminus \{s, t\}$ to $V(P_j) \setminus \{s, t\}$ in $G^{\mathcal{P}}$ that lies entirely in the t -side of $C_{b_{\leftarrow}^{\alpha}}$ and in the s -side of $C_{b_{\leftarrow}^{\alpha} + 1}$. By the definition of G^{α} , this path is also present in G^{α} . Hence, it is also present in the residual graph of G^{α} and \mathcal{P}^{α} .

To see that $\delta_{G^{\alpha}}^{+}(s), C_{b_{\leftarrow}^{\alpha}}, C_{b_{\leftarrow}^{\alpha} + 1}, \dots, C_{b_{\leftarrow}^{\alpha} + \lambda - 1}, \delta_{G^{\alpha}}^{-}(t)$ is the H -sequence of mincuts in $(G^{\alpha}, \mathcal{P}^{\alpha})$, recall that $\delta_{G^{\alpha}}^{+}(s)$ is the projection of $C_{b_{\leftarrow}^{\alpha} - 1}$ if $b_{\leftarrow}^{\alpha} > 1$ and $\delta_{G^{\alpha}}^{-}(t)$ is the projection of $C_{b_{\leftarrow}^{\alpha} + \lambda}$ if $b_{\leftarrow}^{\alpha} + \lambda \leq \ell$. \square

Claim 14. *Let $1 \leq \alpha \leq \xi$ be such that \mathbf{B}^{α} is an actual block and α is excellent. Then, for every $v \in V(G^{\alpha})$, it holds that:*

- *there is a path from s to v in $G^{\alpha} - Z^{\alpha}$ if and only if v is in the s -side of Z ;*
- *there is a path from v to t in $G^{\alpha} - Z^{\alpha}$ if and only if there is a path from v to t in $G - Z$.*

Proof. We first prove the \Rightarrow implication for the first claim. It is immediate for $v = s$, so assume $v \neq s$. Let Q be a path from s to v in $G^{\alpha} - Z^{\alpha}$ and let $(s, v_i), i \in D^{\alpha}$, be the first edge of this path. By Claim 9, there exists a path Q_i from s to v_i in $G - Z$ that visits only vertices in the s -side of $C_{b_{\leftarrow}^{\alpha}}$ (except for the endpoint). The concatenation of Q_i and Q witnesses that v is in the s -side of Z .

The proof for the \Rightarrow implication of the second claim is very similar. It is immediate for $v = t$, so assume $v \neq t$. Let Q be a path from v to t in $G^{\alpha} - Z^{\alpha}$ and let $(u_i, t), i \in D^{\alpha}$, be the least edge of this path. By Claim 8, the suffix of the path P_i from $C_{b_{\leftarrow}^{\alpha} + \lambda}$ to t is disjoint with Z and in the t -side of Z . Hence, the concatenation of Q and this suffix is a path from v to t in $G - Z$.

We now move to the \Leftarrow direction for the first claim. Let Q be a path from s to v in $G - Z$, witnessing that v is in the s -side of Z . Note that $v \neq t$. By the definition of G^{α} , there is a path R from v to t in G^{α} ; let (u_j, t) be the last edge of this path.

Since v is in the t -side of $C_{b_{\leftarrow}^{\alpha} - 1}$, Q intersects $C_{b_{\leftarrow}^{\alpha} - 1}$; let e_1 be the last edge of $C_{b_{\leftarrow}^{\alpha} - 1}$ on Q and let $i_1 \in [\lambda]$ be such that e_1 lies on P_{i_1} . Since both endpoints of e_1 are in the s -side of Z , $i_1 \in L^{\alpha - 1}$.

We observe that no vertex w on Q can lie on a path P_i for $i \in L^{\alpha}$. If w lies on both Q and P_i for $i \in L^{\alpha}$, then the subpath of Q from w to v and the subpath of R from v to u_j contradicts Claim 5 as $j \in D^{\alpha}$ so $j \notin L^{\alpha}$ and L^{α} is downward-closed. This in particular implies that $i_1 \in D^{\alpha}$.

In particular, Q contains no edge of $C_{b_{\leftarrow}^{\alpha} + \lambda}$, as all edges of $C_{b_{\leftarrow}^{\alpha} + \lambda}$ that are in the s -side of Z lie on paths $P_i, i \in L^{\alpha}$. Consequently, the subpath Q' of Q from the head of e_1 to v is completely contained in the t -side of $C_{b_{\leftarrow}^{\alpha} - 1}$ and the s -side of $C_{b_{\leftarrow}^{\alpha} + \lambda}$. Concatenating Q' with R witnesses that the whole Q' is in fact present in G^{α} . Hence, Q' , prepended with (s, v_{i_1}) is a path from s to v in $G^{\alpha} - Z^{\alpha}$, as desired.

We are left with the \Leftarrow direction for the second claim, which is similar to the previous argumentation. Let Q be a path from v to t in $G - Z$; note that $v \neq s$. By the definition of G^{α} , there is a path R from s to v in G^{α} ; let (s, v_j) be the first edge of this path.

Since v is in the s -side of $C_{b_{\leftarrow}^{\alpha} + \lambda}$, Q intersects $C_{b_{\leftarrow}^{\alpha} + \lambda}$; let e_2 be the first edge of $C_{b_{\leftarrow}^{\alpha} + \lambda}$ on Q and let $i_2 \in [\lambda]$ be such that e_2 lies in P_{i_2} . We have $i_2 \notin L^{\alpha}$.

We observe that no vertex w on Q can lie on a path P_i for $i \notin L^{\alpha - 1}$. If w lies on both Q and P_i for $i \notin L^{\alpha - 1}$, then the subpath of R from v_j to v and the subpath of Q from w to v contradicts Claim 5 as $j \in D^{\alpha}$ so $j \in L^{\alpha - 1}$ and $L^{\alpha - 1}$ is downward-closed. This in particular implies $i_2 \in D^{\alpha}$.

In particular, Q contains no edge of $C_{b_{\leftarrow}^{\alpha}-1}$, as all edges of $C_{b_{\leftarrow}^{\alpha}-1}$ that are in the t -side of Z lie on paths P_i , $i \notin L^{\alpha-1}$. Consequently, the subpath Q' of Q from v to the tail of e_2 is completely contained in the t -side of $C_{b_{\leftarrow}^{\alpha}-1}$ and the s -side of $C_{b_{\rightarrow}^{\alpha}+\lambda}$. Concatenating R with Q' witnesses that the whole Q' is in fact present in G^{α} . Hence, Q' with the edge (u_{i_1}, t) is a path from v to t in $G^{\alpha} - Z^{\alpha}$, as desired. \square

Claim 15. *Let $1 \leq \alpha \leq \xi$ be such that \mathbf{B}^{α} is an actual block and α is excellent. Then Z^{α} is a star st -cut in G^{α} . Furthermore, $\text{core}(Z^{\alpha}) = \text{core}(Z) \cap Z^{\alpha}$.*

Proof. This is an easy corollary of Claim 14.

Since Z is an st -cut, there is no vertex of G that has both a path from s to v and a path from v to t in $G - Z$. Claim 14 implies that no vertex of G^{α} admits both a path from s to v and from v to t in $G^{\alpha} - Z^{\alpha}$. Thus, Z^{α} is an st -cut. The first point of Claim 14 implies that from Z being a star st -cut in G follows that for every $(u, v) \in Z^{\alpha}$, u is in the s -side of Z^{α} in G^{α} and v is in the t -side of Z^{α} in G^{α} . Finally, an arc $(u, v) \in Z^{\alpha}$ we have that $(u, v) \in \text{core}(Z^{\alpha})$ if and only if there is path from v to t in $G^{\alpha} - Z^{\alpha}$, which by Claim 14 is equivalent to an existence of a path from v to t in $G - Z$, which is equivalent to $(u, v) \in \text{core}(Z)$. \square

Claim 15 allows the following recursive step. For every excellent $1 \leq \alpha \leq \xi$, recurse on \mathcal{I}^{α} , κ^{α} , and \mathcal{P}^{α} , obtaining a pair $(A^{\alpha}, \widehat{\mathcal{P}}^{\alpha})$. By Claim 13, all recursive calls either consider smaller value of k or consider properly bounded instance with the same k , λ , and H , but fall into the $\ell \leq \ell^{\text{big}}$ case.

We aim at $(A^{\alpha}, \widehat{\mathcal{P}}^{\alpha})$ being compatible with Z^{α} for all excellent α where \mathbf{B}^{α} is an actual block. In this case, for any such α , we have $\kappa^{\alpha} \leq \lambda_{G^{\alpha}+A^{\alpha}}(s, t) = |\text{core}_{G^{\alpha}+A^{\alpha}}(Z^{\alpha})| \leq |Z^{\alpha}| = k^{\alpha}$. For every $\iota \in J$, we randomly guess an integer $\kappa_Z^{\iota} \leq \lambda_Z^{\iota} \leq k_Z^{\iota}$, aiming at $\lambda_Z^{\iota} = \lambda_{G^{\alpha}+A^{\alpha\iota}}(s, t)$. The guess is correct with probability $2^{-\mathcal{O}(k \log k)}$. We say that an excellent index α is *superb* if $\lambda_{G^{\alpha}+A^{\alpha}}(s, t) = \lambda_Z^{\iota(\alpha)}$. Note that if our guess is correct, for every excellent α , if \mathbf{B}^{α} is an actual block, then α is superb.

Return a set A consisting of:

1. for every superb $1 \leq \alpha \leq \xi$, the following edges:
 - (a) all edges of A^{α} with the following replacement:
 - if $b_{\leftarrow}^{\alpha} > 1$, replace every arc $(s, v) \in A^{\alpha}$ with arcs (u, v) where u ranges over all tails of edges of $C_{b_{\leftarrow}^{\alpha}-1}$ on paths P_i for $i \in D^{\alpha}$;
 - if $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$, replace every arc $(v, t) \in A^{\alpha}$ with arcs (v, u) where u ranges over all tails of edges of $C_{b_{\rightarrow}^{\alpha}+\lambda}$ on paths P_i for $i \in D^{\alpha}$;
 - (b) for every $i \in [\lambda] \setminus D^{\alpha}$ an arc from the tail of the edge of $C_{b_{\leftarrow}^{\alpha}-1} \cap E(P_i)$ (or s if $b_{\leftarrow}^{\alpha} = 1$) to the tail of the edge of $C_{b_{\rightarrow}^{\alpha}+\lambda} \cap E(P_i)$ (or t if $b_{\rightarrow}^{\alpha} + \lambda > \ell$);
2. for every good but not superb $1 \leq \alpha \leq \xi$, the following edges:
 - for every $i \in [\lambda]$, an edge from the tail of the edge of $C_{b_{\leftarrow}^{\alpha}-1} \cap E(P_i)$ (or s if $b_{\leftarrow}^{\alpha} \leq 1$) to the tail of the edge of $C_{b_{\rightarrow}^{\alpha}+\lambda} \cap E(P_i)$ (or t if $b_{\rightarrow}^{\alpha} + \lambda > \ell$).
3. for every $1 \leq a \leq \ell$ for which there is no good α such that the block $\mathbf{B}^{\alpha} = (b_{\leftarrow}^{\alpha} < \dots < b_{\rightarrow}^{\alpha})$ satisfies $b_{\leftarrow}^{\alpha} \leq a \leq b_{\rightarrow}^{\alpha} + 1$:
 - for every $i \in [\lambda]$, an edge from the tail of the edge of $C_a \cap E(P_i)$ to the tail of the edge of $C_{a+1} \cap E(P_i)$ (or t if $a = \ell$);

4. for every Γ -milestone a , for every $\iota \in J$, for every $i, j \in L_Z^{\iota-1} \setminus L_Z^\iota$, an edge from the tail of the edge of $C_a \cap E(P_i)$ to the tail of the edge of $C_a \cap E(P_j)$.

Return a flow $\widehat{\mathcal{P}}$ of value $\sum_{\iota \in J} \lambda_Z^\iota$ constructed as follows. Intuitively, for every $\iota \in [\eta]$, we push λ_Z^ι units of flow along the flow paths $P_i, i \in L_Z^{\iota-1} \setminus L_Z^\iota$. We go from s to t .

- For every superb $1 \leq \alpha \leq \xi$, denote $\iota := \iota(\alpha)$ and:
 - push $\kappa^\alpha = \kappa_Z^\iota$ units of flow from $C_{\max(b_{\leftarrow}^\alpha - 1, 1)}$ to $C_{\min(b_{\rightarrow}^\alpha + \lambda, \ell)}$ using the flow $\widehat{\mathcal{P}}^\alpha$ of value λ_Z^ι in $G^\alpha + A^\alpha$ (and arcs added in Point 1a), using the arcs added in Point 4 at milestones $b_{\leftarrow}^\alpha - 1$ (if $b_{\leftarrow}^\alpha > 1$) and $b_{\rightarrow}^\alpha + \lambda$ (if $b_{\rightarrow}^\alpha + \lambda \leq \ell$) to reshuffle the flow between flow paths $P_i, i \in L_Z^{\iota-1} \setminus L_Z^\iota$ in $C_{b_{\leftarrow}^\alpha - 1}$ and $C_{b_{\rightarrow}^\alpha + \lambda}$, respectively;
 - for every $\iota' \in J \setminus \{\iota\}$ push $\lambda_Z^{\iota'}$ units of flow from $C_{\max(b_{\leftarrow}^\alpha - 1, 1)}$ to $C_{\min(b_{\rightarrow}^\alpha + \lambda, \ell)}$ using edges added in Point 1b.
- For every good but not superb $1 \leq \alpha \leq \xi$, for every $\iota \in J$ push λ_Z^ι units of flow along the corresponding edges from $C_{\max(b_{\leftarrow}^\alpha - 1, 1)}$ to $C_{\min(b_{\rightarrow}^\alpha + \lambda, \ell)}$ added in Point 2.
- For every $1 \leq a \leq \ell$ that such that there is no good α with $b_{\leftarrow}^\alpha - 1 \leq a \leq b_{\rightarrow}^\alpha + \lambda$, for every $\iota \in J$ push λ_Z^ι units of flow along the corresponding edges added for a and $i \in L_Z^{\iota-1} \setminus L_Z^\iota$ in Point 3.

Recall that we work under the assumption that all random guesses in this recursive call were correct for the cut Z and, furthermore, for every $\iota \in J$, the pair $(A^{\alpha_\iota}, \widehat{\mathcal{P}}^{\alpha_\iota})$ is compatible with Z^{α_ι} in G^{α_ι} . The following two observations show correctness of the recursion above.

Claim 16. *Let*

$$Z' := \bigcup_{\iota \in J} \text{core}_{G^{\alpha_\iota} + A^{\alpha_\iota}}(Z^{\alpha_\iota}).$$

Then Z' is an st -cut in $G + A$.

Proof. By contradiction, let Q be an st -path in $G + A - Z'$.

For every $\iota \in J$, if $b_{\rightarrow}^{\alpha_\iota} + \lambda \leq \ell$, let D_ι be the set of all tails of edges of $C_{b_{\rightarrow}^{\alpha_\iota} + \lambda}$. Note that D_ι forms a vertex separator from s to t in $G + A$, that is, any path from s to t in $G + A$ needs to contain a vertex of D_ι .

Assume that for some $\iota \in J$ for which D_ι is defined (i.e., $b_{\rightarrow}^{\alpha_\iota} + \lambda \leq \ell$) the path Q contains a vertex $v \in D_\iota$ that is a tail of an edge of $C_{b_{\rightarrow}^{\alpha_\iota} + \lambda}$ on path P_i where $i \notin L_Z^\iota$. Let ι, v , and i be such that v is the first such vertex on Q . For brevity, denote $\alpha = \alpha_\iota$.

As L_Z^ι is downward-closed, Claim 5 implies that no vertex on Q before v (except for s) lies on a path $P_{i'}$ for $i' \in L_Z^\iota$. In particular, by the choice of v , there is no vertex of D_ι before v on Q .

If $b_{\leftarrow}^\alpha \leq 1$, then the prefix of Q until v is present in $G^\alpha + A^\alpha - Z^\alpha$, a contradiction to the assumption that A^α is compatible with Z^α . Otherwise, let D be the set of tails of edges of $C_{b_{\leftarrow}^\alpha - 1}$ and observe that Q needs to pass a vertex of D before v . Let w be the last such vertex and $j \in [\lambda]$ be such that w is a tail of an edge of $C_{b_{\leftarrow}^\alpha - 1} \cap E(P_j)$. We have $w \in L_Z^{\iota-1}$: if $\iota = 1$, then $L_Z^{\iota-1} = [\lambda]$ and the statement is immediate and, otherwise, it follows from the choice of v . Since $L_Z^{\iota-1}$ is downward-closed, Claim 5 implies that $i \in L_Z^{\iota-1}$. Hence, $i, j \in L_Z^{\iota-1} \setminus L_Z^\iota = D^\alpha$.

We infer that the subpath of Q from w to v projects to a path from s to t in $G^\alpha + A^\alpha - Z^\alpha$, a contradiction to the assumption that A^α is compatible with Z^α . Hence, such a vertex v does not exist.

As $L_Z^\eta = \emptyset$, this is only possible if $\eta \in J$ and $b_{\rightarrow}^{\alpha_\eta} + \lambda > \ell$. For brevity, denote $\alpha = \alpha_\eta$. Since $\ell > \ell^{\text{big}}$, we have $b_{\leftarrow}^\alpha > 1$. Let D be the set of tails of edges of $C_{b_{\leftarrow}^\alpha - 1}$ and observe that Q needs to pass a vertex of

D. Let v be the last such vertex and assume v is a tail of an edge of $C_{b_{\leftarrow}^{\alpha}-1} \cap E(P_i)$ for some $i \in [\lambda]$. We have $i \in L_Z^{\eta-1}$, as otherwise the index ι would be defined. Hence, the subpath of Q from v to t projects to a path from s to t in $G^{\alpha} + A^{\alpha} - Z^{\alpha}$, a contradiction to the assumption that A^{α} is compatible with Z^{α} . This finishes the proof of the claim. \square

Claim 17. $(A, \widehat{\mathcal{P}})$ is compatible with Z in G .

Proof. We first check if A is compatible with Z in G , that is, if no edge of A goes from the s -side of Z to the t -side of Z .

For edges added in Point 1a, consider two cases. If \mathbf{B}^{α} is an actual block, it suffices to recall Claim 14: a vertex $v \in V(G^{\alpha})$ is in the s -side of Z^{α} if and only if it is in the s -side of Z in G and, furthermore for $i \in D^{\alpha}$ the endpoints of the edge of $E(P_i) \cap C_{b_{\leftarrow}^{\alpha}-1}$ are in the s -side of Z (if $b_{\leftarrow}^{\alpha} > 1$) and the endpoint of the edge of $E(P_i) \cap C_{b_{\rightarrow}^{\alpha}+\lambda}$ (if $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$) are in the t -side of Z . If \mathbf{B}^{α} is not an actual block, we observe that all vertices of $V(G^{\alpha}) \setminus \{s, t\}$ lie either entirely in the s -side of Z or entirely in the t -side of Z due to the fact that we kept in G^{α} only vertices reachable from s and from which one can reach t in G_0^{α} .

For edges added in Point 1b, we first observe that for every $i \in [\lambda] \setminus D^{\alpha}$ it holds that $i \in L_{b_{\leftarrow}^{\alpha}-1}$ if and only if $i \in L_{b_{\rightarrow}^{\alpha}+\lambda}$: if \mathbf{B}^{α} is an actual block it follows from the assumption that $L_{b_{\leftarrow}^{\alpha}-1} = L^{\alpha-1}$ and $L_{b_{\rightarrow}^{\alpha}+\lambda} = L^{\alpha}$ and otherwise we have $L_{b_{\leftarrow}^{\alpha}-1} = L_{b_{\rightarrow}^{\alpha}+\lambda}$. Hence, as both $b_{\leftarrow}^{\alpha} - 1$ (if $b_{\leftarrow}^{\alpha} > 1$) and $b_{\rightarrow}^{\alpha} + \lambda$ (if $b_{\rightarrow}^{\alpha} + \lambda \leq \ell$) are milestones, we have that any edge added in Point 1b lie on the same side of Z .

For edges added in Point 2, note that we have $L_{b_{\leftarrow}^{\alpha}-1} = L^{\alpha-1} = L^{\alpha} = L^{b_{\rightarrow}^{\alpha}+\lambda}$, so an edge added for $i \in L^{\alpha}$ connects two vertices in the s -side of Z and for $i \notin L^{\alpha}$ connects two vertices in the t -side of Z .

For edges added in Point 3, note that any such a is untouched and hence for every $i \in [\lambda]$, the entire subpath of P_i from the edge of C_a to the edge of C_{a+1} (or t if $a = \ell$) lies in the same side of Z . Hence, again for any edge added in Point 3, both its endpoints are in the same side of Z .

Finally, for edges added in Point 4, the claim follows from the fact that every Γ -milestone a is a milestone, and hence a is untouched and $L_a \in \{L_Z^0, \dots, L_Z^{\eta}\}$. Thus, both endpoints of an edge added in Point 4 lies in the same side of Z .

This concludes the proof that A is compatible with Z in G . In particular, Z is a star st -cut in $G + A$.

Claim 16 asserts that $\widehat{\mathcal{P}}$ is an st -maxflow in $G + A$ and, furthermore, Z contains an st -cut Z' of size $|\widehat{\mathcal{P}}|$. Hence, as Z is a star st -cut, $Z' = \text{core}_{G+A}(Z)$.

It remains to check that $\widehat{\mathcal{P}}$ is a witnessing flow for Z in $G + A$. Note that the only edges of $\widehat{\mathcal{P}}$ that are not in A are edges inside subinstances G^{α} from recursive calls for superb α . If \mathbf{B}^{α} is an actual block, then $\widehat{\mathcal{P}}$ does not use any edge of $Z \setminus Z'$ in G^{α} as $\widehat{\mathcal{P}}^{\alpha}$ is a witnessing flow for Z^{α} in $G^{\alpha} + A^{\alpha}$. If \mathbf{B}^{α} is not an actual block, there is no edge of Z in G^{α} . This finishes the proof of the lemma. \square

4.6 Probability analysis

Fix a star st -cut Z of $|Z| \leq k$. We want to lower bound the probability that the recursive algorithm finds a pair $(A, \widehat{\mathcal{P}})$ compatible with Z . We say that the algorithm is *correct* at some random step if it correctly guesses the properties of Z .

In the preprocessing phase, the algorithm either directly returns a desired pair (without any random choice), recurses on an instance with smaller value of $2k - \lambda$ (being correct with probability $\Omega(k^{-1})$), or recurses on an instance with the same value of k and λ and, being properly boundaried (with probability $\Omega(1)$).

In the base case $|E(H)| = |V(H)|$, we either return a correct outcome or recurse on an instance with larger λ (making correct choices with probability $\Omega(\lambda^{-1}) = \Omega(k^{-1})$).

In the case of $\ell \leq \ell^{\text{big}}$, the algorithm makes correct guesses about sets B_{\leftarrow} and B_{\rightarrow} with probability at least $2^{-\mathcal{O}(k \log k)}$. If $\lambda_{G+A_0}(s, t) > \lambda_G(s, t)$ at this point, the algorithm recurses on an instance with smaller value of $2k - \lambda$. If a vertex of $V(C)$ is in the s -side of Z , where C is the st -mincut of $G + A_0$ closest to t , the algorithm makes correct guesses with probability $\Omega(k^{-1})$ and again recurses on an instance with a smaller value of $2k - \lambda$. If $V(C)$ is contained in the t -side of Z , the algorithm guess so with probability $\Omega(1)$ and recurses on an instance with the same k and λ , but smaller $|E(H)|$. Note that $|E(H)|$ can decrease (while keeping k and λ the same) less than k^2 times.

In the case of $\ell > \ell^{\text{big}}$, recall the algorithm makes correct guesses with probability $2^{-\mathcal{O}(k \log k)}$. Recall also that we are interested that in every Γ -block \mathbf{B}^α that is an actual block and α is superb, the returned pair $(A^\alpha, \widehat{\mathcal{P}}^\alpha)$ is compatible with Z^α in G^α . By Claim 13, there is either one such recursive call (if $L_Z^{\iota-1} \setminus L_Z^\iota = [\lambda]$ for some $1 \leq \iota \leq \eta$) with the same k, λ, H , but falling into case $\ell \leq \ell^{\text{big}}$, or a number of such calls with values $2k^\alpha - |\mathcal{P}^\alpha|$ summing up to at most $2k - |\mathcal{P}|$.

Hence, the depth of the recursion is $\mathcal{O}(k^3)$, as $2k - \lambda$ can decrease only $2k$ times and, between these decreases, $|E(H)|$ can decrease less than k^2 times. Furthermore, in the entire recursion tree we care only about being correct $\mathcal{O}(k^3)$ recursive calls, as even if in the last case there are multiple subcalls $(G^\alpha, \mathcal{P}^\alpha)$ with $Z^\alpha \neq \emptyset$, the values of $2k^\alpha - |\mathcal{P}^\alpha|$ for these calls sum up to at most $2k - |\mathcal{P}|$. At each recursive call, we make correct guesses with probability $2^{-\mathcal{O}(k \log k)}$. Hence, the overall success probability is $2^{-\mathcal{O}(k^4 \log k)}$, as desired.

5 Applications: weighted st -cut, weighted DFVS, weighted Chain SAT

5.1 Tractable case of WEIGHTED BUNDLED CUT

An instance of BUNDLED CUT consists of a directed graph G , vertices $s, t \in V(G)$, a nonnegative integer k , and a family \mathcal{B} of pairwise disjoint subsets of $E(G)$. An element $B \in \mathcal{B}$ is called a *bundle*. An edge that is part of a bundle is *deletable*, otherwise it is *undeletable*. A *cut* in a BUNDLED CUT instance $\mathcal{I} = (G, s, t, k, \mathcal{B})$ is an st -cut Z that does not contain any undeletable edge, that is, $Z \subseteq \bigcup \mathcal{B}$. A cut Z *touches* a bundle $B \in \mathcal{B}$ if $Z \cap B \neq \emptyset$. The *cost* of a cut Z is the number of bundles it touches. A cut Z is a *solution* if its cost is at most k . The BUNDLED CUT problem asks if there exists a solution to the input instance.

An instance of WEIGHTED BUNDLED CUT consists of a BUNDLED CUT instance $\mathcal{I} = (G, s, t, k, \mathcal{B})$ and additionally a weight function $\omega : \mathcal{B} \rightarrow \mathbb{Z}_+$ and an integer $W \in \mathbb{Z}_+$. The *weight* of a cut Z in \mathcal{I} is the total weight of all bundles it touches. A solution Z to \mathcal{I} is a solution to the WEIGHTED BUNDLED CUT instance (\mathcal{I}, ω, W) if additionally the weight of Z is at most W . The WEIGHTED BUNDLED CUT problem asks if there is a solution to the input instance. Note that any BUNDLED CUT instance can be treated as a WEIGHTED BUNDLED CUT instance by setting ω uniformly equal 1 and $W = k$.

When parameterized by k , it is well known that BUNDLED CUT is W[1]-hard even if all bundles are of size 2 [MR09]. To get tractability, we define the following restriction called *Weighted Bundled Cut with Order* where we require the following: in the input instance $((G, s, t, k, \mathcal{B}), \omega, W)$, for every $B \in \mathcal{B}$ one can order the arcs of B as $e_1^B, \dots, e_{|B|}^B$ such that for every $1 \leq i < j \leq |B|$ there exists a path $P_{i,j}^B$ in G that goes from one endpoint of e_i^B to one endpoint of e_j^B and uses only undeletable arcs and arcs of B .

In this subsection we prove the following theorem.

Theorem 5.1. *WEIGHTED BUNDLED CUT WITH ORDER is randomized FPT when parameterized by k and maximum size of a bundle.*

The main workhorses behind the proof of Theorem 5.1 are the flow-augmentation routine and the following lemma.

Lemma 5.2. *Assume we are given a WEIGHTED BUNDLED CUT WITH ORDER instance $\mathcal{I} = ((G, s, t, k, \mathcal{B}), \omega, W)$. Then one can in time FPT with parameters $\lambda_G(s, t)$, maximum size of a bundle, and k check if there is a solution to \mathcal{I} that is an st -mincut, where for $\lambda_G(s, t)$ we treat every deletable edge as of capacity 1 and every undeletable edge as of capacity $+\infty$.*

Proof of Theorem 5.1 using Lemma 5.2. Let b be the maximum size of a bundle. We will describe a randomized algorithm that runs in time FPT in b and k and, if given a yes-instance, answers yes with probability $2^{-\mathcal{O}((bk)^4 \log(bk))}$, and never answers yes to a no-instance. Repeating the algorithm $2^{\mathcal{O}((bk)^4 \log(bk))}$ times gives the desired algorithm.

Consider a solution Z to the input WEIGHTED BUNDLED CUT WITH ORDER instance $\mathcal{I} = ((G, s, t, k, \mathcal{B}), \omega, W)$. Without loss of generality, we can assume that Z is a minimal st -cut. Note that $|Z| \leq bk$. Apply flow-augmentation (Theorem 4.1) to G and parameter bk , obtaining a pair $(A, \widehat{\mathcal{P}})$, and add all arcs of A to G as undeletable arcs. Note that adding arcs A to G does not break the requirements for a WEIGHTED BUNDLED CUT WITH ORDER instance. With probability $2^{-\mathcal{O}((bk)^4 \log(bk))}$ the set Z is still a solution in the final instance and is an st -mincut. Run the algorithm of Lemma 5.2, returning its answer. \square

Thus, we are left with proving Lemma 5.2.

Proof of Lemma 5.2. For every $B \in \mathcal{B}$, fix an order $B = (e_1^B, \dots, e_{|B|}^B)$ as in the definition of a WEIGHTED BUNDLED CUT WITH ORDER instance. Let b be the maximum size of a bundle.

Compute a maximum st -flow $\mathcal{P} = \{P_1, P_2, \dots, P_\lambda\}$ where $\lambda := \lambda_G(s, t)$. Fix a hypothetical solution Z ; note that Z contains exactly one deletable edge on every path P_i ; denote by f_i the unique edge of $E(P_i) \cap Z$. Branch, guessing the following information about Z . First, guess the number $\kappa \leq k$ of bundles touched by Z . Let those bundles be $F_1, F_2, \dots, F_\kappa \in \mathcal{B}$. For every $i \in [\lambda]$, guess the indices $\alpha(i) \in [\kappa]$ and $\beta(i) \in [b]$ such that $f_i = e_{\beta(i)}^{F_{\alpha(i)}}$. There are $2^{\mathcal{O}(\lambda \log(kb))}$ options. We perform a simple sanity check we consider only options where the mapping $i \mapsto (\alpha(i), \beta(i))$ is injective.

We also guess a partition of \mathcal{B} into sets $\mathcal{B}_1, \dots, \mathcal{B}_\kappa$, aiming at $F_j \in \mathcal{B}_j$. By standard objects for derandomizing color-coding, this can be turned into a branch into $2^{\mathcal{O}(\kappa)} \log |\mathcal{B}|$ options.

For every $1 \leq j \leq \kappa$ and every $B \in \mathcal{B}_j$ we make a sanity check: we expect that, for every $i \in [\lambda]$ such that $\alpha(i) = j$ the edge $e_{\beta(i)}^B$ actually lies on P_i . If this is not the case, we remove B from \mathcal{B}_j and \mathcal{B} (making its edges undeletable). Clearly, F_j remains in \mathcal{B}_j in the branch where the guesses are correct.

We now make the following (main) filtering step. Iterate over all $1 \leq j \leq \kappa$ and indices $1 \leq i_1, i_2 \leq \lambda$ such that $\alpha(i_1) = \alpha(i_2) = j$ but $\beta(i_1) < \beta(i_2)$. Consider $B \in \mathcal{B}_j$ such that there exists another $B' \in \mathcal{B}_j$ such that $e_{\beta(i_1)}^{B'}$ is before $e_{\beta(i_1)}^B$ on P_{i_1} but $e_{\beta(i_2)}^{B'}$ is after $e_{\beta(i_2)}^B$ on P_{i_2} . Assume $B = F_j$. Then $B' \cap Z = \emptyset$. By the property of the WEIGHTED BUNDLED CUT WITH ORDER instance, G contains a path Q from an endpoint of $e_{\beta(i_1)}^{B'}$ to an endpoint of $e_{\beta(i_2)}^{B'}$ that uses only undeletable edges and edges of B' . Hence, Q is disjoint with Z . However, as $e_{\beta(i_1)}^B \in Z$ lies after $e_{\beta(i_1)}^{B'}$ on P_{i_1} while $e_{\beta(i_2)}^B \in Z$ lies before $e_{\beta(i_2)}^{B'}$ on P_{i_2} , Q goes from the s -side of Z to the t -side of Z , a contradiction. Hence, $B \neq F_j$ and we can remove such B from \mathcal{B}_j and \mathcal{B} .

After we perform the filtering step exhaustively, for every $1 \leq j \leq \kappa$ we can order \mathcal{B}_j as $B_{j,1}, \dots, B_{j,|\mathcal{B}_j|}$ such that for every $1 \leq \xi < \zeta \leq |\mathcal{B}_j|$ and for every $i \in [\lambda]$, if $\alpha(i) = j$, then $e_{\beta(i)}^{B_{j,\xi}}$ is before $e_{\beta(i)}^{B_{j,\zeta}}$ on P_i . Let a_j^Z be such that $F_j = B_{j,a_j^Z}$.

In the end, we also check if still $\lambda = \lambda_G(s, t)$. If this is not the case, we terminate the current branch.

An edge $e \in E(G)$ is *vulnerable* if $e = e_{\beta(i)}^B$ for some $i \in [\lambda]$ and $B \in \mathcal{B}_{\alpha(i)}$. Note that all edges of Z are vulnerable, all vulnerable edges are deletable, and vulnerable edges appear only on paths of \mathcal{P} .

We now construct an auxiliary weighted directed graph H as follows. Start with H consisting of two vertices s and t . For every $1 \leq j \leq \kappa$, add a path P_j^H from s to t with $|\mathcal{B}_j|$ edges; denote the a -th edge as $e_{j,a}$ and set its weight as $\omega(e_{j,a}) = \omega(B_{j,a})$. Furthermore, for every $1 \leq i_1, i_2 \leq \lambda$, denote $j_1 = \alpha(i_1)$, $j_2 = \alpha(i_2)$, for every $1 \leq a_1 \leq |\mathcal{B}_{j_1}|$ and $1 \leq a_2 \leq |\mathcal{B}_{j_2}|$, for every endpoint u_1 of $e_{\beta(i_1)}^{B_{j_1, a_1}}$, for every endpoint u_2 of $e_{\beta(i_2)}^{B_{j_2, a_2}}$, if G contains a path from u_1 to u_2 consisting only of nonvulnerable edges, then add to H an edge of weight $+\infty$ from the corresponding endpoint of e_{j_1, a_1} (i.e., tail if and only if u_1 is a tail of $e_{\beta(i_1)}^{B_{j_1, a_1}}$) to the corresponding endpoint of e_{j_2, a_2} (i.e., tail if and only if u_2 is a tail of $e_{\beta(i_2)}^{B_{j_2, a_2}}$).

Observe that $Z' := \{e_{j, a_j^Z} \mid 1 \leq j \leq \kappa\}$ is an st -cut in H . Indeed, if H would contain an arc (v, u) with v before e_{j, a_j^Z} and u after $e_{j', a_{j'}^Z}$ for some $j, j' \in [\kappa]$, then this arc was added to H because of some path between the corresponding endpoints in G and such a path would lead from the s -side to t -side of Z . Also, in the other direction, observe that if $Y' = \{e_{j, a_j} \mid 1 \leq j \leq \kappa\}$ is an st -mincut in H , then

$$Y = \left\{ e_{\beta(i)}^{B_{\alpha(i), \alpha(i)}} \mid i \in [\lambda] \right\}$$

is a solution to \mathcal{I} of the same weight.

Hence, it suffices to find in H an st -cut of cardinality κ and minimum possible weight. Since $\kappa \leq \lambda_H(s, t)$, this can be done in polynomial time. \square

5.2 Applications

In all three discussed problems, we are given both a bound k on the cardinality of the solution and a bound W on the total weight of the solution. The parameter is k .

In WEIGHTED CUT, the solution is an st -cut. In WEIGHTED DFAS, the solution is a feedback arc set in a directed graph. Note that by standard reduction, this is equivalent to WEIGHTED DFVS. In WEIGHTED b -CHAIN SAT, the problem is in fact a WEIGHTED BUNDLED CUT where every bundle is a path of length at most b .

Note that WEIGHTED CUT is WEIGHTED BUNDLED CUT with bundles of size 1 and no undeletable edges while WEIGHTED b -CHAIN SAT is WEIGHTED BUNDLED CUT with bundles of size at most b . Furthermore, in both problems the assumption for WEIGHTED BUNDLED CUT WITH ORDER is satisfied (in the latter case, order the edges along the path). We infer the following.

Theorem 5.3. *WEIGHTED CUT is randomized FPT when parameterized by k .*

Theorem 5.4. *WEIGHTED b -CHAIN SAT is randomized FPT when parameterized by k and b .*

By standard approach, WEIGHTED DFAS can be solved using a subroutine for WEIGHTED SKEW MULTICUT. Here, we are given a directed graph G , a tuple $(s_i, t_i)_{i=1}^b$ of terminal pairs, a weight function $\omega : E(G) \rightarrow \mathbb{Z}_+$, and integers k, W . The goal is to find a set $Z \subseteq E(G)$ of cardinality at most k , weight at most W , and such that there is no path from s_i to t_j in $G - Z$ for any $1 \leq i \leq j \leq b$. We observe the following reduction.

Lemma 5.5. *Given a WEIGHTED SKEW MULTICUT instance $\mathcal{I} = (G, (s_i, t_i)_{i=1}^b, \omega, k, W)$, one can in polynomial time construct an equivalent WEIGHTED BUNDLED CUT WITH ORDER instance $\mathcal{I}' = (G', \mathcal{B}, \omega, k, W)$ with the same k and W and bundles of size b each.*

Proof. To construct the graph G' , we start with b disjoint copies G^1, \dots, G^b of the graph G . By v^i, e^i , etc., we denote the copy of vertex v or edge e in the copy G^i . We set $B_e = \{e^i \mid i \in [b]\}$ for $e \in E(G)$ and $\mathcal{B} = \{B_e \mid e \in E(G)\}$, that is, all b copies of one edge of G form a bundle. We set weights of bundles as $\omega'(B_e) = \omega(e)$. There will be no more bundles, so all arcs introduced later to G' are undeletable.

For every $1 \leq i < j \leq b$ and $v \in V(G)$, we add to G' an arc (v^i, v^j) . These arcs make the instance satisfy the “order” property with the natural order (e^1, e^2, \dots, e^b) for the bundle B_e . Furthermore, we add to G' vertices s and t and arcs (s, s_i^i) and (t_i^i, t) for every $i \in [b]$. This finishes the description of the instance $\mathcal{I}' = (G', \mathcal{B}, \omega', k, W)$.

It is straightforward to observe that if Z is a solution to the instance \mathcal{I} , then $\bigcup_{e \in Z} B_e$ is a solution to \mathcal{I}' of the same weight and touching $|Z|$ bundles. In the other direction, note that if Z' is a solution to \mathcal{I}' then $Z = \{e \in E(G) \mid Z' \cap B_e \neq \emptyset\}$ is a solution to \mathcal{I} . \square

We deduce the following.

Theorem 5.6. *WEIGHTED DFAS and WEIGHTED DFVS are randomized FPT when parameterized by k .*

6 Weighted Almost 2-SAT

In this section, we present the algorithm for WEIGHTED ALMOST 2-SAT. Let us recall the problem definition. The input consists of a 2-CNF formula ϕ , viewed as a set of 2-clauses; a weight function $\omega: \phi \rightarrow \mathbb{Z}_+$; and integers $k, W \in \mathbb{Z}_+$. The objective is to find a clause deletion set $Z \subseteq \phi$ such that $\phi - Z$ is satisfiable, $|Z| \leq k$, and $\omega(Z) = \sum_{C \in Z} \omega(C) \leq W$. We refer to a clause set $Z \subseteq \phi$ such that $\phi - Z$ is satisfiable as a *solution*. Our goal is thus to find such a solution Z , with $|Z| \leq k$ and $\omega(Z) \leq W$, in time FPT parameterized by k . We show the following.

Theorem 6.1. *WEIGHTED ALMOST 2-SAT is randomized FPT parameterized by k , with a running time of $2^{k^{O(1)}} n^{O(1)}$.*

We also show a corollary for valued CSPs; see end of section.

As in the ALMOST 2-SAT-algorithm of Kratsch and Wahlström [KW20], we solve the problem by reduction to the auxiliary problem DIGRAPH PAIR CUT. Let us define the weighted version of this. An instance $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ of WEIGHTED DIGRAPH PAIR CUT consists of a digraph G with special vertices s and t , a weight function $\omega: E(G) \rightarrow \mathbb{Z}_+$, a set $\mathcal{T} \subseteq \binom{V(G)}{2}$ of pairs of vertices of G , and integers $k, W \in \mathbb{Z}_+$. Refer to a set of edges $Z \subseteq E(G)$ as a *solution* if Z is an st -cut and for every pair $p \in \mathcal{T}$, at most one endpoint of p is reachable from s in $G - Z$. The goal is then to find a solution $Z \subseteq E(G)$ such that $|Z| \leq k$ and $\omega(Z) \leq W$. Note that the addition of a sink vertex t differs from the original formulation of DIGRAPH PAIR CUT [KW20]; we add such a vertex t in order to make use of the flow-augmentation technique.

Let us note that WEIGHTED ALMOST 2-SAT FPT-reduces to WEIGHTED DIGRAPH PAIR CUT.

Lemma 6.2. *Let $\mathcal{I} = (\phi, \omega, k, W)$ be an instance of WEIGHTED ALMOST 2-SAT. In time $2^{O(k)}$ we can produce a list of instances $\mathcal{I}' = (G, s, t, \omega', \mathcal{T}, k', W')$ of WEIGHTED DIGRAPH PAIR CUT such that \mathcal{I} is a yes-instance if and only if at least one instance \mathcal{I}' is a yes-instance, where $k' = O(k)$ for each produced instance.*

Proof. We use the reduction of Kratsch and Wahlström [KW20], slightly simplified, and verify that it preserves weights. In time $2^{O(k)}$, we can find a solution $Z_0 \subseteq \phi$ to \mathcal{I} with $|Z_0| \leq k$ or conclude that no such solution exists [KW20]. In the latter case, we report that \mathcal{I} is a no-instance. If Z_0 exists and $\omega(Z_0) \leq W$, we may simply accept \mathcal{I} . Hence, we assume that Z_0 exists and $\omega(Z_0) > W$.

Perform the following operation. Pick any satisfying assignment f of $\phi - Z_0$ and for every variable v such that $f(v)$ is true, negate v in the entire formula ϕ , that is, replace all literals v with $\neg v$ and all literals $\neg v$ with v . In this manner, we do not change the space of solutions to the input instance \mathcal{I} , while the assignment that sets false to every variable satisfies $\phi - Z_0$. In other words, $\phi - Z_0$ does not contain a clause with two positive literals. Furthermore, we remove from Z_0 every clause that contains at least one negative literal; clearly $\phi - Z_0$ is still satisfied by the all-false assignment. Hence, all clauses of Z_0 have only positive literals, while every clause of $\phi - Z_0$ has at least one negative literal.

Let X be the set of variables occurring in clauses of Z_0 . Branch over every assignment $f: X \rightarrow \{0, 1\}$; there are $2^{|X|} \leq 4^k$ such assignments.

In every branch, perform the following. Let Z_1 be the set of all clauses in ϕ that have every literal false under f and Z_2 the set of all clauses in ϕ that have at least one true literal under f ; observe that $Z_0 \subseteq Z_1 \cup Z_2$. If $|Z_1| > k$ or $\omega(Z_1) > W$ we abort the branch; otherwise proceed with $\phi \leftarrow \phi - (Z_1 \cup Z_2)$. Next, initialize a digraph G_0 over vertex set $V(\phi)$ (i.e., we produce one vertex v for every variable v in ϕ), with a directed edge (u, v) if and only if the clause $C = (\neg u \vee v)$ occurs in ϕ . Create a weight function ω' where $\omega'((u, v)) = \omega(C)$ for every such edge (u, v) . For every clause $C = (\neg u \vee \neg v)$ in ϕ , create a pair of fresh vertices u' and v' in G_0 , add edges (u, u') and (v, v') , and add the pair $\{u', v'\}$ to \mathcal{T} . Let $\omega'((u, u')) = \omega'((v, v')) = \omega(C)$. Finally contract the vertices of $f^{-1}(1)$ in G_0 into a new vertex s , and contract the vertices of $f^{-1}(0)$ in G_0 into a new vertex t . We note that $f^{-1}(1) \neq \emptyset$, as otherwise $Z_1 \supseteq Z_0$ and $\omega(Z_1) > W$, and the branch is aborted. Finally, if no vertex t was created, we simply add an isolated vertex t . Let G be the resulting graph, and output the instance $\mathcal{I}' = (G, s, t, \omega', \mathcal{T}, k - |Z_1|, W - \omega(Z_1))$.

We now show correctness. First assume that there is a (minimal) solution Z to \mathcal{I} with $|Z| \leq k$ and $\omega(Z) \leq W$. Let g be a satisfying assignment to $\phi - Z$ and consider the branch where the assignment f to X matches g . Since f and g match on X , every clause in Z_1 is false in g and every clause in Z_2 is true in g , i.e., $(Z_1 \cup Z_2) \cap Z = Z_1$. The other clauses $C \in Z$ are of two types, clauses $(\neg u \vee v)$ with one negation and clauses $(\neg u \vee \neg v)$ with two negations, as $C \in Z \setminus (Z_1 \cup Z_2) \subseteq Z \setminus Z_0$. Define an edge set Z' consisting of every arc (u, v) such that $(\neg u \vee v) \in Z$, and of one arc (u, u') for every pair $\{u, v\} \in \mathcal{T}$ such that $(\neg u \vee \neg v) \in Z$, disambiguated arbitrarily between (u, u') and (v, v') . Note that Z' contains precisely $|Z \setminus Z_1| \leq k - |Z_1|$ edges and $\omega(Z') = \omega(Z) - \omega(Z_1)$. We show that Z' is a solution to \mathcal{I}' . Let $S \subseteq V(G)$ be the vertices reachable from s in $G - Z'$. Then $g(v) = 1$ for every $v \in S$ where v corresponds to a variable in ϕ (i.e., omitting vertices v' added in the pair creation step). This follows by induction on the distance of v from s in $G - Z'$. As base case, consider an arc (s, u) in $G - Z'$. Then there is a clause $C = (\neg u_0 \vee u)$ in $\phi - Z$ where $u_0 \in X$ and $f(u_0) = 1$; hence $g(u_0) = g(u) = 1$. Otherwise, inductively consider a vertex $v \in S$ and let $P = su_1 \dots u_\ell v$ be a shortest sv -path in $G - Z'$. Then $g(u_\ell) = 1$ and $\phi - Z$ contains a clause $(\neg u_\ell \vee v)$; hence $g(v) = 1$. It follows that Z' is an st -cut, since either t is isolated or there is at least one vertex $v \in X$ with $f(v) = g(v) = 0$ contracted into t during construction; hence $t \notin S$. Finally, for every clause $C = (\neg u \vee \neg v) \in \phi - Z$ at most one of its variables is in S , or else g would falsify it; hence the corresponding pair $\{u', v'\}$ is not contained in S . For all other pairs $\{u', v'\}$, the corresponding clause $C = (\neg u \vee \neg v)$ is in Z and, thus, Z' contains (u, u') or (v, v') , which implies $\{u', v'\} \subseteq S$. Thus Z' is a solution to \mathcal{I}' .

In the other direction, assume that the instance \mathcal{I}' created in some branch f has a solution Z' , i.e., no pair in \mathcal{T} is reachable from s in $G - Z'$, and both $|Z'| \leq k - |Z_1|$ and $\omega'(Z') \leq W - \omega(Z_1)$, where Z_1 is the set of clauses false under f . Let $X_1 = f^{-1}(1)$, let S be the set of vertices reachable from s in $G - Z'$, and let g be the assignment to ϕ where $g(v) = 1$ if $v \in S \cup X_1$ and $g(v) = 0$ otherwise. Note that g matches f on X . Consider a clause $C \in \phi$ that is false under g . Then either $C \in Z_1$, but those are exactly $|Z_1|$ many and their weight is $\omega(Z_1)$; or, either $C = (\neg u \vee v)$ where $\{u, v\} \cap (S \cup X_1) = \{u\}$, or $C = (\neg u \vee \neg v)$

where $u, v \in S \cup X_1$. In the former case, there is an arc (s, v) or (u, v) in $E(G)$ which must belong to Z' ; in the latter case we have a pair $\{u', v'\} \in \mathcal{T}$ and since Z' is a solution at least one of the arcs into u' and v' is cut by Z' . In both cases, the clause of ϕ that is false under g is mapped to a distinct arc in G of the same weight; all other false clauses are those in Z_1 with total weight $\omega(Z_1)$. Hence the clauses of ϕ false under g form a solution to \mathcal{I} of cardinality $|Z| = |Z'| + |Z_1| \leq k$ and weight $\omega(Z) = \omega'(Z') + \omega(Z_1) \leq W$. \square

In the sequel, we therefore focus on WEIGHTED DIGRAPH PAIR CUT. Furthermore, although the problem was formally defined as taking a weight threshold W , we will solve the problem in FPT time by finding a min-weight solution Z among all solutions Z with $|Z| \leq k$. Clearly, this is sufficient.

6.1 Overview and outlook

Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an instance of WEIGHTED DIGRAPH PAIR CUT and let Z be a minimal solution to \mathcal{I} . Then Z must be a star st -cut; i.e., for every $(u, v) \in Z$, u is reachable from s in $G - Z$ but v is not. Therefore, our algorithm is based around the flow augmentation for star st -cuts (Theorem 4.1). However, compared to previous applications in this paper, the process to get an FPT algorithm is more involved.

In rough overview, our algorithm has three phases. First, we use standard branching methods to set the stage for an application of Theorem 4.1. After this phase, and after a successful call to Theorem 4.1, we have an augmented graph $G + A$ and an st -maxflow \mathcal{P} in $G + A$ such that (1) the st -mincut closest to s in $G + A$ is a solution, although it may have large weight; and (2) (A, \mathcal{P}) is compatible with Z , meaning that Z is a star st -cut in $G + A$ such that $\text{core}_{G+A}(Z)$ is an st -mincut and Z intersects every path of \mathcal{P} precisely once. This phase is summarized in Lemma 6.13 in Section 6.3.

Second, given a pair (A, \mathcal{P}) compatible with Z , we wish to compute further augmentations so that Z becomes an st -mincut rather than a star cut. This is handled in Section 6.3. We are not able to compute such an augmentation “non-destructively”; however, by carefully analysing how the pairs \mathcal{T} interact with the maxflow \mathcal{P} , we are able to reduce the instance \mathcal{I} , with parameter k and optimal solution Z of cardinality k , to an augmented instance \mathcal{I}' with parameter $k' \leq 2k$ and optimal solution $Z' = Z \cup Z^+$ of cardinality k' .

Finally, we assume that we have an instance where the solution Z is an st -mincut. That is, the flow \mathcal{P} consists of k edge-disjoint st -paths, and Z cuts each path of \mathcal{P} once and makes no other cuts. Even this restricted case turns out to be non-trivial; the solution is given in the next subsection.

In fact, due to the extra expressive power given by edge weights, the remaining problem in the min-cut situation is equivalent to the following question: Let $G = (V, E)$ be a digraph with vertices $s, t \in V$ and edge weights $\omega: E \rightarrow \mathbb{Z}_+$, and let ϕ be a 2-CNF formula on variable set $V \setminus \{s, t\}$. Let $k = \lambda_G(s, t)$. We then wish to find an st -mincut Z of smallest possible weight, such that Z is consistent with ϕ in the following sense: Define an assignment f such that $f(v) = 1$ if v is reachable from s in $G - Z$, and $f(v) = 0$ otherwise. Then f is required to be a satisfying assignment to ϕ . Note that ϕ is no longer required to be a “digraph pair cut-formula”, i.e., all three clause types $(u \vee v)$, $(\neg u \vee v)$ and $(\neg u \vee \neg v)$ can occur in ϕ .

Due to basic manipulations, this problem is also equivalent to the following high-domain Valued CSP. The input consists of k variables x_i , with domains of size $O(n)$, with a language supporting arbitrary weighted unary constraints and crisp binary constraints of the types $((x_i \geq a) \vee (x_j \geq b))$, $((x_i < a) \vee (x_j \geq b))$ and $((x_i < a) \vee (x_j < b))$, for $i, j \in [k]$ and a, b being values in the corresponding domains. The task is then to find a min-weight solution that satisfies all crisp constraints.

Our results thus imply that both of these problems have randomized FPT algorithms parameterized by k .

6.2 Min-cut instances

In this section, we solve the problem when Z is an st -mincut. More precisely, let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an instance of WEIGHTED DIGRAPH PAIR CUT. Say that \mathcal{I} is a *min-cut DPC instance* if $\lambda_G(s, t) = k$, where $\lambda_G(s, t)$ is computed with respect to the *unweighted* graph G , ignoring the edge weights ω . Hence every solution Z with $|Z| \leq k$ must be an st -mincut in G .

6.2.1 Initial reductions

Let $P = sv_1 \dots v_n t$ be an st -path in G . A *twin path* of P is an st -path $P' = sv'_n \dots v'_1 t$ where for every min-weight solution Z with $|Z| \leq k$ and every internal vertex v of P , v lies in the s -side of Z if and only if v' lies in the t -side of Z . Note that, if we write $v'' = v$, then the relation is symmetric, and we say that P and P' is a *twin path pair*. We show that, given an instance \mathcal{I} and a maxflow \mathcal{P} , we can introduce a twin path for every path in \mathcal{P} .

Lemma 6.3. *Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be a min-cut DPC instance and let $\mathcal{P} = \{P_1, \dots, P_k\}$ be an st -maxflow. In polynomial time, we can compute a min-cut DPC instance $\mathcal{I}' = (G', s, t, \omega', \mathcal{T}', 2k, W + W_0)$ for some $W_0 > 0$, and an st -maxflow $\mathcal{P}' = (P_1, P'_1, \dots, P_k, P'_k)$ in \mathcal{I}' , such that P_i and P'_i is a twin path pair for every $i \in [k]$, and the cost of an optimal solution in \mathcal{I}' is precisely W_0 higher than in \mathcal{I} .*

Proof. To begin with, let Z_0 be the st -mincut closest to s . If Z_0 is not a solution, i.e., if some pair $\{u, v\} \in \mathcal{T}$ is contained in the s -side of Z_0 , then \mathcal{I} has no solution of cardinality at most k and we may reject the instance. Indeed, by choice of Z_0 , every vertex reachable from s in $G - Z_0$ is also reachable from s in $G - Z$ for any st -mincut Z . Otherwise, if Z_0 is a solution, set $W' = \max(W, \omega(Z_0)) + 1$. The value W' will serve as a prohibitively large weight, guaranteed to be larger than the weight of an optimal solution.

Let $P = sv_1 \dots v_r t$ be an st -path and let $P' = sv'_r \dots v'_1 t$ be an induced path where v'_i is a fresh vertex for every $i \in [r]$. For every $i \in [r]$, add the pair $\{v_i, v'_i\}$ to \mathcal{T} . Set $\omega(s, v'_r) = rW'$, $\omega(v'_{i+1}, v'_i) = iW'$ for every $i \in [r-1]$, and $\omega(v'_1, t) = 0$. Add rW' to $\omega(s, v_1)$, add $(r-i)W'$ to $\omega(v_i, v_{i+1})$ for every $i \in [r-1]$, and keep $\omega(v_r, t)$ unchanged. Let ω' be the resulting weights. We claim that P and P' is a twin path pair; observe that the resulting instance \mathcal{I}' is still a min-cut instance.

On the one hand, let Z be a solution to \mathcal{I} with $|Z| \leq k$, and let $Z \cap E(P) = \{(v_i, v_{i+1})\}$ (interpreting v_0 as s and v_{r+1} as t). Let $Z' = Z + (v'_{i+1}, v'_i)$. Then Z' is a solution to \mathcal{I}' , and $\omega'(Z') = \omega(Z) + rW'$. Furthermore, v_i is in the s -side of Z' if and only if v'_i is not. We show that these are the only solutions of weight at most $W + rW'$.

For this, let Z' be a solution to \mathcal{I}' with $|Z'| \leq k + 1$, and first assume that there is an $i \in [r]$ such that both v_i and v'_i are in the s -side of Z' . But then the pair $\{v_i, v'_i\}$ is contained in the s -side of Z' , contradicting that Z' is a solution. On the other hand, assume that v_i and v'_i are both in the t -side. Then P is cut at the latest on edge (v_{i-1}, v_i) , and P' is cut at the latest on edge (v'_{i+1}, v'_i) . These two edges contribute weight $(r+1)W' > W + rW'$ to $\omega'(Z')$; cutting earlier edges on either path would add even more weight. By the previous paragraph, such a solution cannot be min-weight. Hence P and P' is a twin path pair.

Apply the above procedure for every path $P_i \in \mathcal{P}$, and add the resulting paths P'_i to \mathcal{P} . Return the resulting instance, with updated weights ω' and parameter $k' = 2k$, the resulting flow \mathcal{P}' , and the value $W_0 = O(nkW')$ produced as the sum of the additive terms rW' . \square

Next, we observe that the twin paths allow us to reduce \mathcal{I} to a very special form where G consists of k internally vertex-disjoint induced st -paths and nothing more.

Lemma 6.4. *Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be a min-cut DPC instance and let \mathcal{P} be an st -maxflow with paths partitioned into twin path pairs. In polynomial time, we can reduce \mathcal{I} to an equivalent min-cut DPC instance $\mathcal{I}' = (G', s, t, \omega, \mathcal{T}', k, W)$ and an st -maxflow \mathcal{P}' in G' such that every path $P \in \mathcal{P}'$ is an induced path, the paths in \mathcal{P}' are pairwise internally vertex-disjoint, and G' contains no arcs or vertices except those of \mathcal{P}' .*

Proof. First, we compute the *projection* of \mathcal{I} onto the vertices $V(\mathcal{P})$. Let $G' = G - E(\mathcal{P})$, and note that $Z \cap E(G') = \emptyset$ for every solution Z with $|Z| \leq k$ since \mathcal{I} is a min-cut instance. For this, we compute the set of all arcs (u, v) and pairs $\{u, v\}$ over $V(\mathcal{P})$ that are implied by \mathcal{T} and G' . Specifically, for every $u, v \in V(\mathcal{P})$ such that G' contains a path from u to v , add the arc (u, v) to a new arc set A ; and for every $u, v \in V(\mathcal{P})$ such that there exists a pair $\{x, y\} \in \mathcal{T}$ such that u reaches x in G' and v reaches y in G' , add the pair $\{u, v\}$ to \mathcal{T} . After this process, delete all vertices of $V(G) \setminus V(\mathcal{P})$ and restrict \mathcal{T} to the set of pairs $p \in \mathcal{T}$ contained in $V(\mathcal{P})$.

We observe that this preserves the set of solutions Z with $|Z| \leq k$. In the first case, consider the instance with graph $G + A$ and the extended set of pairs \mathcal{T} , with $\omega(e) = W + 1$ for every $e \in A$. On the one hand, any solution Z to this instance of weight at most W is also a solution to \mathcal{I} . On the other hand, let Z be a solution to \mathcal{I} with $|Z| = k$. Then as noted $Z \cap E(G') = \emptyset$, hence every path Q in G' is uncut by Z . Therefore Z cuts no arc $e \in A$ (i.e., if $e = (u, v)$ and u is in the s -side of Z , then so is v). Similarly, let $\{u, v\}$ be a pair that was added to \mathcal{T} . Let $\{x, y\} \in \mathcal{T}$ be a pair in the input from which $\{u, v\}$ was created. If u and v are both in the s -side of Z , then there are paths in G' from u, v to x, y that are unbroken in Z , hence $\{x, y\}$ is contained in the s -side of Z , contradicting that Z is a solution. Hence Z satisfies all added arcs and pairs.

Next, let $G^* = (G + A)[V(\mathcal{P})]$. Clearly, taking an induced subgraph does not eliminate any solutions; we show that it also does not introduce any new unwanted solutions. Let Z be a solution to the new instance. First, we note that for every vertex $v \in V(\mathcal{P})$, v is in the s -side of Z in $G + A$ if and only if v is in the s -side of Z in G^* . On the one hand, any sv -path in $G^* - Z$ also exists in $G + A - Z$. On the other hand, let Q be an sv -path in $G + A - Z$. If $V(Q) \subseteq V(\mathcal{P})$ then Q exists in $G^* - Z$. Otherwise, let Q' be a subpath of Q with endpoints in $V(\mathcal{P})$ and internal vertices disjoint from $V(\mathcal{P})$. Assume Q' goes from p to q . But then $(p, q) \in A$, hence replacing Q' by the arc (p, q) in Q yields a shorter sv -path, unbroken by Z . Hence we can generate an sv -path Q^* in $G^* - Z$, showing equivalence. In particular, this shows that t is not in the s -side in $G + A - Z$, i.e., Z is an st -cut. Next, assume that there is a pair $\{x, y\} \in \mathcal{T}$ contained in the s -side of Z in G . Let Q_x and Q_y be paths in $G - Z$ from s to u and v , respectively. Let u be the last vertex of $Q_x \cap V(\mathcal{P})$ and v the last vertex of $Q_y \cap V(\mathcal{P})$. Then \mathcal{T} contains a pair $\{u, v\}$, but by the above both u and v are in the s -side of Z in G^* , a contradiction. Hence we showed the equivalence of taking the projection on vertex set $V(\mathcal{P})$.

Next, we expand G^* so that the paths of \mathcal{P} are pairwise internally vertex-disjoint. For this, let $\mathcal{P} = \{P_1, \dots, P_k\}$. For every $i \in [k]$ and every $v \in V(P_i)$, create a fresh vertex $v^{(i)}$, and for every pair $i, j \in [k]$ such that $v \in V(P_i) \cap V(P_j)$, we add arcs $(v^{(i)}, v^{(j)})$ and $(v^{(j)}, v^{(i)})$. Furthermore, for every arc (u, v) not on a flow path, add arcs $(u^{(i)}, v^{(j)})$ for all copies of u and v , and similarly create all pairs $\{u^{(i)}, v^{(j)}\}$ for pairs $\{u, v\} \in \mathcal{T}$. Clearly, this preserves all solutions of cardinality at most k .

Finally, for every arc (u, v) that is not contained in a flow, add the pair $\{u, v'\}$, where v' is the negation of v found on the corresponding twin path. It is easy to see that this is equivalent; the arc (u, v) would contradict a proposed solution Z if and only if u is in the s -side and v in the t -side, which is equivalent to both u and v' being in the s -side.

Since every step in the above sequence preserves all solutions of cardinality at most k , we are done. \square

We refer to the resulting instances as *arcless* (even though technically speaking, they still contain arcs

in the paths).

6.2.2 Solving arcless instances

We now assume that \mathcal{I} is an arcless instance. From this point on we forget the path pairings, and treat \mathcal{P} as an unordered collection of st -paths, $\mathcal{P} = \{P_1, \dots, P_k\}$ (where k is twice its original value). For two vertices $u, v \in V(P_i)$, we write $u \leq v$ if u lies between s and v on P_i , and $u < v$ if additionally $u \neq v$. We extend the notation to pairs: For two pairs p, p' intersecting distinct paths P_i and P_j , write $p \leq p'$ if $p \cap V(P_i) \leq p' \cap V(P_i)$ and $p \cap V(P_j) \leq p' \cap V(P_j)$.

Define an undirected graph H on $[k]$ where $\{i, j\} \in E(H)$ if and only if there exists a pair with endpoints on P_i and P_j . Furthermore, for all $\{i, j\} \in E(H)$ let $\mathcal{T}_{i,j} \subseteq \mathcal{T}$ be the set of all pairs $p \in \mathcal{T}$ intersecting P_i and P_j , where p furthermore is *minimal*, i.e., if there is a pair $p' \in \mathcal{T}$ intersecting P_i and P_j , with $p' \leq p$, then $p = p'$.

Lemma 6.5. *If H is bipartite, then the instance is polynomial-time solvable.*

Proof. Compute a bipartition $[k] = A \cup B$ of H . Compute a weighted maxflow instance as follows. Let $W^+ = \sum_{e \in E(G)} \omega(e) + 1$ and $\varepsilon = 1/W^+$. Replicate paths $P_i, i \in A$ as-is, letting $\omega'(e) = 1 + \varepsilon\omega(e)$ for every arc $e \in E(P_i)$. For every $j \in B$, reverse the path $P_j = sv_1 \dots v_n t$ into a path $sv_n \dots v_1 t$, letting $\omega'((u, v)) = 1 + \varepsilon\omega((v, u))$ for every arc $(u, v) \in E(P_j)$. Finally, for every pair $\{u, v\} \in \mathcal{T}$, where u lies on a path in A and v on a path in B , introduce the arc (u, v) with weight $k + 1$. Let G' be the resulting graph. We claim that \mathcal{I} has a solution Z with cardinality at most k and weight $W, 0 \leq W < W^+$, if and only if (G', s, t, ω') has an st -cut of weight at most $k + \varepsilon W < k + 1$.

On the one hand, let Z be a solution with $|Z| \leq k$ and $\omega(Z) = W$. Since \mathcal{I} is a min-cut instance, $|Z| = k$ and Z contains precisely one edge per path. Let Z' be the corresponding edge set in G' , i.e., Z' contains the edge $Z \cap E(P_i)$ for every $i \in A$ as well as the reversal of $Z \cap E(P_j)$ for every $j \in B$; thus, $\omega'(Z') = k + \varepsilon W$. We show that Z' is an st -cut. Assume to the contrary that $G' - Z'$ contains an st -path Q . Let $(u, v) \in E(Q)$ be an edge such that u lies before Z' on its path and v lies after Z' on its path. But then $\{u, v\} \in \mathcal{T}$, and by design both u and v are in the s -side of Z in G , contradicting that Z is a solution. Hence we also conclude that the weight of an st -mincut in G' is strictly less than $k + 1$, since we assume that \mathcal{I} has a solution of cardinality k . On the other hand, let Z' be a min-weight st -cut in G' , of weight $k + \varepsilon W, W < W^+$. (Note $\omega(Z') \geq k$ since the paths in G' form a flow of weight at least k .) Then Z' is also an st -cut of minimum cardinality, containing precisely one edge per path P_i . Let Z be the corresponding edge set in \mathcal{I} , and assume there is a pair $\{u, v\} \in \mathcal{T}$ such that both u and v are in the s -side of Z . Assume that u lies on a path $P_i, i \in A$ and v on a path $P_j, j \in B$. Then $(u, v) \in E(G')$. But by construction u is in the s -side of Z' and v is not, contradicting that Z' is an st -cut. \square

Finally, we show how to reduce the non-bipartite case to bipartite with some additional branching. We show that when H is not bipartite, we can inspect an odd cycle in H and with $f(k)$ guesses simplify the instance such that $|E(H)|$ strictly decreases.

To support the discussion, we define a contraction-and-cleanup procedure. Recall that all vertices of G lie on paths in \mathcal{P} .

Definition 6.6 (Contraction and cleanup). Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an arcless instance, with maxflow \mathcal{P} . Let $v \in V(G)$ be a vertex, $v \notin \{s, t\}$, and let $P \in \mathcal{P}$ be the path with $v \in V(P)$. To *contract v into s* (respectively *into t*) corresponds to the following two-step procedure. In the first step, we contract v and every predecessor u of v on P into s (respectively, contract v and every successor u of v on P into t).

For any pair $p \in \mathcal{T}$, and vertex $v \in p$ contracted into s (respectively into t) in the process, retain the pair $p \in \mathcal{T}$ with v replaced by s (respectively by t). We then apply a *cleanup* procedure, repeating the following steps to exhaustion, always applying the earliest step that applies.

1. If any pair has both endpoints in s , reject the instance.
2. If any pair has an endpoint in t , discard it.
3. If any pair has an endpoint in s , contract its other endpoint into t .

Note that after cleanup, no pair intersects s or t .

Lemma 6.7. *Let $v \in V(G)$, $v \notin \{s, t\}$, and let $P \in \mathcal{P}$ with $v \in V(P)$. Contracting a vertex v into t never causes an instance to be rejected, and contracting v into s causes rejection if and only if there is a pair $p \in \mathcal{T}$ with both members contained between s and v (inclusively) on P .*

Proof. Initially, before contractions, no pair intersects s or t , and no step of the cleanup procedure contracts any vertex into s . Hence the only case when an instance \mathcal{I} is rejected, due to a pair $p \in \mathcal{T}$, is if both endpoints of p were contracted into s in one step, i.e., v is contracted into s and this caused both endpoints of p to be contracted into s . \square

We show correctness of the contract-and-cleanup procedure.

Lemma 6.8. *Let \mathcal{I}' be the result of applying a contraction of a vertex v into s (respectively into t) in an arcless instance \mathcal{I} and cleaning up. Then \mathcal{I}' is an arcless instance, and any solution Z' to \mathcal{I}' is a solution to \mathcal{I} of the same cost. Furthermore, if Z is a solution to \mathcal{I} where v is in the s -side (respectively in the t -side), then Z remains a solution of the same cost in \mathcal{I}' .*

Proof. It is clear that the process maintains arcless instances. Assume that a contraction and cleanup on \mathcal{I} yields an instance $\mathcal{I}' = (G', s, t, \omega, \mathcal{T}', k, W)$ without terminating and let Z' be a solution to \mathcal{I}' . By design and by the arcless form, Z' is an st -mincut in G as well, and every vertex in $V(G')$ is in the s -side (resp. t -side) of Z' in G if and only if it is in the s -side (resp. t -side) of Z' in G' . Assume towards a contradiction that there is a pair $p \in \mathcal{T}$ with both endpoints in the s -side of Z' in G . Since $p \notin \mathcal{T}'$, p must have been discarded in cleanup. But then at least one endpoint of p must have been contracted into s , which, since \mathcal{I} was not rejected, implies that the other endpoint would have been contracted into t , contradicting that p is contained in the s -side. Thus Z' is a solution to \mathcal{I} . The cost of Z' is unchanged.

Finally, let Z be a solution to \mathcal{I} , and assume that v was contracted into x and v is in the x -side of Z , for $x \in \{s, t\}$. We argue that every vertex u contracted into s (respectively into t) this way was in the s -side (respectively t -side) of Z . Let $P \in \mathcal{P}$ be the path containing v , and first consider $x = t$. Then the only contraction is of the suffix of v on P into t , and clearly this entire suffix is in the t -side of Z . Otherwise $x = s$, and the contraction is of v into s on P possibly followed by some number of contractions of vertices u_i into t . As above, the prefix of P is entirely in the s -side of Z . Furthermore, consider some vertex u_i contracted into t in the cleanup phase. Then there must exist a pair $\{y, u_i\} \in \mathcal{T}$ such that y precedes v on P , as no other cascaded contractions are performed. But then y is in the s -side of Z , and since Z is a solution u_i must be in the t -side. It now follows that Z remains a solution in \mathcal{I}' : Since every edge $(u, u') \in Z$ has u in the s -side and u' in the t -side, no edge of Z was deleted by a contraction; thus Z is an st -mincut in G' . Hence the t -side of Z in G' is contained in the t -side of Z in G , implying that for every pair $p \in \mathcal{T}'$ there is a vertex $u \in p$ in the t -side of Z . Thus Z is a solution in \mathcal{I}' , and clearly the cost of Z is unchanged. \square

As a further pre-cleaning, assume that there is a pair $p = \{u, v\} \in \mathcal{T}$ where u and v lie on the same path $P \in \mathcal{P}$, with $u \leq v$ on P . Then we may contract v into t and clean up, as above. Hence we assume that all pairs hit two distinct paths in \mathcal{P} .

Now, we observe some beneficial contraction operations that will yield instances where $|E(H)|$ has been reduced.

Lemma 6.9. *Let $\{i, j\} \in E(H)$ and let $\mathcal{T}_{i,j}$ be the minimal pairs on paths P_i and P_j . Let $s < v_1^i < \dots < v_r^i < t$ be the endpoints on P_i of pairs $p \in \mathcal{T}_{i,j}$, in order from s to t . Performing any of the following operations and cleaning up the result will eliminate all pairs between P_i and P_j in the instance.*

1. To contract v_1^i into t .
2. To contract v_a^i into s and v_{a+1}^i into t for any $a \in [r - 1]$.
3. To contract v_r^i into s .
4. To contract both v_a^i and v_a^j into t , where $\{v_a^i, v_a^j\} \in \mathcal{T}_{i,j}$.

Proof. For the first three, consider a pair $\{v_a^i, v_a^j\} \in \mathcal{T}_{i,j}$ and assume that the cleanup terminates without rejecting the instance. Then v_a^i is contracted either into s or into t . In the former case, v_a^j is contracted into t in cleanup; in both cases, the pair contains at least one endpoint in t and is dropped. For the last case, consider a pair $\{v_b^i, v_b^j\} \in \mathcal{T}_{i,j}$. If $a \leq b$, then v_b^i is contracted into t ; otherwise, since $\mathcal{T}_{i,j}$ is an anti-chain we have $v_b^j \leq v_a^j$ and v_b^j is contracted into t . For any other pair $p = \{u, v\} \in \mathcal{T}$ occurring on paths P_i and P_j , there is a pair $p' \in \mathcal{T}_{i,j}$ dominating p , and one vertex of p' is contracted into t . Since $p' \leq p$, the same holds for p and all pairs on paths P_i and P_j are eliminated. Finally, note that throughout the cleanup process, the only case where a vertex u is contracted into s is if the contraction was of a vertex v into s and $u \leq v$ (in particular, u and v lie on the same path). Given that no pair initially intersects s , that no pair has both endpoints on the same path, and that $\mathcal{T}_{i,j}$ contains all minimal pairs, it follows that neither of these four contraction operations can result in a terminated instance. \square

Now assume H is not bipartite and let $i_1 \dots i_d i_1$ be an odd cycle in H , i.e., d is odd and for every $j \in [d]$ we have $\{i_j, i_{j+1}\} \in E(H)$, where $i_{d+1} = i_1$. Let $L = \{p_1, \dots, p_{d-1}\}$ be a set of minimal pairs, constructed as follows.

1. Start with an arbitrary pair $p_1 = \{u_1, v_1\} \in \mathcal{T}_{i_1, i_2}$ where u_1 lies on path i_1 and v_1 on path i_2 .
2. For every $j = 2, \dots, d - 1$, select a pair $p_j = \{u_j, v_j\} \in \mathcal{T}_{i_j, i_{j+1}}$, where u_j lies on path i_j , such that no pair of $\mathcal{T}_{i_j, i_{j+1}}$ lies between v_{j-1} and u_j . Note that this may imply $v_{j-1} = u_j$.

We refer to such a set L as an H -path from i_1 to i_d , and say that L induces a *connection* between vertices u_1 on path i_1 and v_{d-1} on path i_d .

Lemma 6.10. *Let L be an H -path inducing a connection between u and v . Then we may either perform one of $O(k)$ beneficial contractions as in Lemma 6.9, or Z hits P_{i_1} before u if and only if it hits P_{i_d} before v .*

Proof. Let $L = \{p_1, \dots, p_{d-1}\}$ as above and assume that for some $j \in [d - 1]$ the solution Z separates v_j from u_{j+1} . Then this implies one of the first three cases of Lemma 6.9 with respect to pairs between i_{j+1} and i_{j+2} . Similarly, if for some $j \in [d - 1]$ the pair $p_j \in L$ is contained in the t -side, then this matches the fourth case of Lemma 6.9. This gives at most $2d = O(k)$ possible beneficial contractions, each of which after cleanup will lead to the removal of an edge of H .

Otherwise, assume that no such event occurs, and consider the case that u_1 is in the s -side of Z . Then v_1 must be in the t -side, and by assumption so is u_2 ; then, for p_2 not to be contained in the t -side v_2 must be in the s -side; and so on, in an alternating process. Since d is odd, we conclude that vertex v_{d-1} on path i_d must be in the s -side. The case that u_1 is in the t -side similarly implies that v_{d-1} is in the t -side. \square

We now have all the tools we need to break the cycle.

Lemma 6.11. *Let $i_1 \dots i_d i_1$ be an odd cycle in H . Then we can branch into $O(k)$ directions, such that in each branch the parameter k is unchanged and $|E(H)|$ has decreased.*

Proof. We assemble a set F of connections between paths i_1 and i_d corresponding to H -paths as above. We add one member $f \in F$ for every pair $p \in \mathcal{T}_{i_1, i_d}$, where f and p intersect P_{i_1} in the same vertex. If there is a choice in constructing the H -path, i.e., in moving from a vertex v_a to a vertex u_{a+1} , select u_{a+1} to be in the s -side of v_a . Order the pairs \mathcal{T}_{i_1, i_d} as $p_1 < \dots < p_m$ according to their order of occurrence on path i_1 , and similarly order $F = \{f_1, \dots, f_r\}$ in increasing order according to their occurrence on path i_1 .

Claim 18. *The pairs of F are in non-decreasing order, i.e., if $a < b$ then $f_a \leq f_b$ on both paths i_1 and i_d .*

Proof. Let $1 \leq a < b \leq r$, and let L_a respectively L_b be the H -connections yielding f_a respectively f_b . Since each set $\mathcal{T}_{a,b}$ contains only the minimal pairs, any pair that starts later on path P_a arrives earlier on P_b . Then, moving from a vertex v_j to a vertex u_{j+1} in the construction of an H -path, a search that starts at a later vertex v'_j arrives at a vertex u'_{j+1} with $u_{j+1} \leq u'_{j+1}$, with equality possible. Thus on any odd index $q \in [d]$ we have $f_a \leq f_b$ on path i_q , and on an even index q we have $f_a \geq f_b$ on path i_q . In particular, we have $f_a \leq f_b$ on paths i_1 and i_d . \square

We establish some shorthand terminology for the rest of the process. For a connection $f \in F$, produced by an H -path L , to *branch on f* refers to applying Lemma 6.10, either branching into one of $O(k)$ beneficial contractions or concluding that either both endpoints of f are in the s -side of Z or both are on the t -side of Z . For a pair $p \in \mathcal{T}$, we say that p is *double hit by Z* if both endpoints of p are on the t -side of Z . If we guess this to be the case, then by Lemma 6.9 we may apply a contraction procedure which will remove an edge from $E(H)$. Finally, assume that $p, p' \in \mathcal{T}_{a,b}$ are adjacent on P_a , i.e., no pair in $\mathcal{T}_{a,b}$ connects between p and p' on P_a , and assume that Z cuts P_a between the two pairs. In this case, we may again apply Lemma 6.9 to remove an edge of $E(H)$.

We now describe the search for a branching. First assume that there is a connection $f \in F$ and a pair of adjacent pairs $p_a, p_{a+1} \in \mathcal{T}_{i_1, i_d}$ such that p_a and p_{a+1} *cross* f , i.e., on path i_1 we have a relative order $p_a \leq f \leq p_{a+1}$ and on path i_d we have a relative order $p_{a+1} \leq f \leq p_a$. Then we have a branching as follows. If Z cuts i_1 or i_d between p_a and p_{a+1} , then by Lemma 6.9 we remove an edge from H . Next, we branch on f , either applying Lemma 6.9 or concluding that either both endpoints of f lie in the s -side of Z or both endpoints of f lie in the t -side of Z . But in the former remaining case the pairs p_a and p_{a+1} are violated by Z , and in the latter case p_a and p_{a+1} are double hit by Z . In all cases, Lemma 6.9 applies and we have a reduction of $E(H)$.

Now assume that no such configuration occurs in F and \mathcal{T}_{i_1, i_d} . Introduce artificial members $f_0 = \{s\}$ and $f_{r+1} = \{t\}$ to F ; clearly, they are consistent with the ordering. Now let f_{a+1} be the earliest member of F such that there exists a pair $p \in \mathcal{T}_{i_1, i_d}$ with $p < f_{a+1}$. Note that f_{a+1} exists with $a \geq 0$ since $f_0 < p < f_{r+1}$ for every $p \in \mathcal{T}_{i_1, i_d}$. Consider f_a . First, assume that there exists a pair $p' \in \mathcal{T}_{i_1, i_d}$ with $f_a \leq p'$. In this case, we branch on f_a and on f_{a+1} , as well as on the case that Z cuts path i_1 between f_a and f_{a+1} . This gives $O(k)$ applications of Lemma 6.9. In the remaining case, Z hits i_1 either before f_a or after f_{a+1} , and by Lemma 6.10 Z hits i_d in the same way. But if Z hits both paths after f_{a+1} , then p is

contained in the s -side since $p < f_{a+1}$, which contradicts Z being a solution. Finally, then, Z must hit both paths i_1 and i_d before f_a , implying that p' is double hit. In all cases, Lemma 6.9 applies.

Next, assume that there are pairs $p, p' \in \mathcal{T}_{i_1, i_d}$ such that $p < f_a \leq p'$ on i_1 . Since we excluded the case that $f_a \leq p'$ overall, we must have $p' < f_a$ on i_d for any such p' . Now we simply pick p as the last pair on i_1 meeting the condition, so that $p = p_b$ and $p' = p_{b+1}$. Since we also excluded $p < f_a$, we conclude $f_a \leq p$ on i_d . Then p and p' are adjacent pairs crossing f_a , hitting a case we already considered.

We now merely have two mop-up cases remaining. If $p < f_a$ on i_1 for every $p \in \mathcal{T}_{i_1, i_d}$, we branch into two cases. Either Z hits path i_1 between p_m and t , or before p_m . In the former case Lemma 6.9 applies. In the latter case, we further branch on f_a , leaving the case that both endpoints of f_a lie in t . But then, since $p_m < f_a$ is excluded, we have $f_a \leq p_m$ on path i_d , implying that Z double hits p_m .

Finally, $f_a \leq p$ for every $p \in \mathcal{T}_{i_1, i_d}$. Branch on f_a and f_{a+1} as well as on the Z hitting i_1 between f_a and f_{a+1} . As above, it is excluded that f_{a+1} is contained in the s -side of Z , leaving the case that f_a is on the t -side of Z . But then Z hits the gap between s and the first member of \mathcal{T}_{i_1, i_d} on path i_1 , and in all cases Lemma 6.9 applies. \square

We summarize the result of this section.

Lemma 6.12. *Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an instance of WEIGHTED DIGRAPH PAIR CUT where $\lambda_G(s, t) = k$. There is a polynomial-time procedure that with probability at least $1/2^{k^{O(1)}}$ produces a min-cost solution to \mathcal{I} (and in the failure cases either rejects the instance or produces solutions of weight higher than optimum).*

Proof. By Lemma 6.3 feeding into Lemma 6.4 we can reduce \mathcal{I} to an arcless instance \mathcal{I}' with parameter $k' = 2k$. Let H be the graph describing intersections of pairs in \mathcal{T} with paths in \mathcal{I}' . If H is bipartite, then we apply Lemma 6.5 and solve the problem in polynomial time. Otherwise we let C be an odd cycle in H and apply Lemma 6.11, giving us a collection of $O(k)$ branches; we guess a branch with success probability $1/O(k)$. Repeat this until H is bipartite, which happens after at most $O(k^2)$ repetitions. In total our probability of consistently selecting the right branch is at least $1/2^{O(k^2 \log k)}$, and since the branches consist of contractions a false choice never gives false positives or solutions with a too-low weight. \square

6.3 General case

We now consider the general case of WEIGHTED DIGRAPH PAIR CUT, where the solution is a star st -cut which is not an st -mincut. Let the input be an instance $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$, where $\lambda_G(s, t) < k$. We begin with a bootstrapping step, leading up to the application of Theorem 4.1.

Lemma 6.13. *Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an instance of WEIGHTED DIGRAPH PAIR CUT and let Z be a min-cost solution with $|Z| \leq k$. There is a polynomial-time procedure that produces a pair (A, \mathcal{P}) where $A \subseteq V^2$ and \mathcal{P} is an st -maxflow in $G + A$, such that the following holds. Let $G' = G + A$, and let ω' be ω updated so that $\omega'(e) = W + 1$ for every $e \in A$. Let*

$$\mathcal{I}' = (G', s, t, \omega', \mathcal{T}, k, W)$$

be the resulting instance. Then with probability at least $1/f(k)$ for some $f(k) = 2^{k^{O(1)}}$, the following holds: the st -mincut closest to s in G' is a solution, Z is a min-cost solution in \mathcal{I}' , and \mathcal{P} is a witnessing flow for Z .

Proof. Throughout this proof, as in the majority of the paper, all mincuts and maxflows are unweighted, i.e., we only concern ourselves with cardinality, not weight. Let Z_0 be the st -mincut closest to s in G , with $Z_0 = \emptyset$ if t is isolated. Recall that Z_0 is well-defined due to submodularity of cuts, and can be found in

polynomial time. As long as there is a pair $p \in \mathcal{T}$ with both endpoints reachable from s in $G - Z_0$, select a random endpoint $u \in p$, add (u, t) to A , and recompute Z_0 . Since Z_0 was the unique st -mincut closest to s , this procedure increases $\lambda_G(s, t)$ every time it is applied, hence it can only occur at most k times. If $\lambda_G(s, t) > k$, abort the computation and return $A = \{(s, t)\}$ and an arbitrary maxflow in $G + A$. Next, apply Theorem 4.1 with instance $(G + A, s, t, k)$, and let the result be (A', \mathcal{P}) , where A' is the augmentation and \mathcal{P} is an st -maxflow in $G + A + A'$. Add the edges of A' to A . If the st -mincut closest to s in $G + A$ is not a solution, repeat the branching process above until this is the case, then reapply Theorem 4.1. If we ever reach a stage where $\lambda_{G+A}(s, t) > k$, abort as above, otherwise return (A, \mathcal{P}) .

We argue correctness. Since every modification that is made makes the instance harder, i.e., solutions are lost but not added, it suffices to show some probability that Z survives as a solution. For the initial branching, this is clear. If a pair p is entirely on the s -side of Z_0 , then Z separates at least one of its members from s , hence our guess of vertex $u \in p$ is correct with probability at least $1/2$. Subject to this, Z remains a star st -cut if the edge (u, t) is added. Furthermore, at the end of the process we have $\lambda_G(s, t) \leq |Z| \leq k$. Next, assume, with probability $2^{-\mathcal{O}(k^4 \log k)}$, that the application of Theorem 4.1 returns a pair (A', \mathcal{P}) compatible with Z . Then Z remains a star st -cut in $G + A$ and \mathcal{P} is a witnessing flow for Z in $G + A$ by definition. Thus, with some probability $1/f(k)$, Z remains a star st -cut throughout the process of branching and augmentation and the process terminates in a state where all the conditions of the lemma are fulfilled. \square

We henceforth proceed with the output of Lemma 6.13, i.e., we assume that the st -mincut Z_0 closest to s in G is a solution, although $\omega(Z_0) > W$, and we carry with us a witnessing flow \mathcal{P} for the optimal solution Z . Furthermore, write $Z = Z_{s,t} \cup Z'$ where $Z_{s,t} = \text{core}_G(Z)$, and let $k = |Z|$, $k' = |Z'|$. We will make progress by decreasing k' (at the cost of increasing k). If $k' = 0$, we invoke the algorithm from the previous subsection. Hence Z is a star st -cut, $Z_{s,t}$ is an st -mincut, and \mathcal{P} is a witnessing flow for Z .

Let $v \in V(G) \setminus V(\mathcal{P})$. The *projection* of v onto P_i , for a path $P_i \in \mathcal{P}$, is the earliest vertex $u \in V(P_i)$ (i.e., closest to s) such that there is a path Q from u to v in G , where Q intersects $V(\mathcal{P})$ only in u . Note that such a projection may not exist, in case all paths from P_i to v pass through other paths in \mathcal{P} . When the projection u exists, we refer to such a path Q as the *attachment path* from u to v .

Lemma 6.14. *Let $P \in \mathcal{P}$, and let v_1, \dots, v_p be a sequence of vertices in $V \setminus V(\mathcal{P})$ such that the projection u_i of v_i onto P exists for every $i \in [p]$ and such that the projections u_i are all distinct. For $i \in [p]$, let Q_i be an attachment path from u_i to v_i . Then for all $i \neq j$, $i, j \in [p]$, Q_i and Q_j are vertex-disjoint.*

Proof. Assume that $i \neq j$, $i, j \in [p]$, and u_i is before u_j on P . Let $w \in V(Q_i) \cap V(Q_j)$. Then there exists a walk Q' from u_i to w to v_j , disjoint from $V(\mathcal{P})$ except at u_i , implying the existence of a path Q'' with the same property. This contradicts the assumption that u_j is the projection of v_j onto P . \square

Let us make a quick observation. Let v be a vertex in the s -side of $Z_{s,t}$. We say that $P_i \in \mathcal{P}$ *activates* v if the projection u of v onto P_i exists, and u is in the s -side of Z . We show that an activating path always exists.

Lemma 6.15. *Let v be in the s -side of $Z_{s,t}$. Then there exists a path $P_i \in \mathcal{P}$ such that the projection u of v onto P_i exists and u is in the s -side of Z .*

Proof. If v lies on a path P_i , then $u = v$, and since \mathcal{P} is a witnessing flow, the s -side of Z on P_i equals the s -side of $Z_{s,t}$. Otherwise, let Q be a path from s to v in $G - Z_{s,t}$ and let u' be the last vertex of Q that lies in $V(\mathcal{P})$. Let $P_i \in \mathcal{P}$ be a path containing u' . (If the choice of P_i is ambiguous, select P_i arbitrarily among the options. In particular, note that we may have $u' = s$ in which case any path $P \in \mathcal{P}$ is a valid choice.)

Let u be the projection of v onto P_i , which exists since $u = u'$ is an option. Then Q proves that u' lies in the s -side of $Z_{s,t}$, and since u precedes u' on P_i and \mathcal{P} is a witnessing flow, u lies in the s -side of Z . \square

Let $\{u, v\}$ be a pair. We say that $\{u, v\}$ is *active* in Z if u and v are in the s -side of $Z_{s,t}$. Then by the above lemma, there exist activating paths $P_i, P_j \in \mathcal{P}$, where we may have $P_i = P_j$, so that the projections of u onto P_i and of v onto P_j exist and both are in the s -side of Z . We note that this implies the presence of arcs from the set Z' .

Lemma 6.16. *If $k' > 0$, then there exists an active pair $\{u, v\} \in \mathcal{T}$. Furthermore, let $\{u, v\} \in \mathcal{T}$ be an active pair, where $P_i \in \mathcal{P}$ activates u and $P_j \in \mathcal{P}$ activates v . Let x be the projection of u onto P_i and y the projection of v onto P_j . Then Z' hits either every attachment path from x to u or every attachment path from y to v .*

Proof. In the first case, assume there is no active pair. Then for every pair $\{u, v\} \in \mathcal{T}$, at least one vertex is in the t -side of $Z_{s,t}$. Thus $Z_{s,t}$ is a solution, and $Z = Z_{s,t}$ by minimality, so $k' = 0$. Otherwise, let $\{u, v\} \in \mathcal{T}$ be an active pair. Since Z is a solution, there exists a vertex $z \in \{u, v\}$ that is in the t -side of Z ; w.l.o.g. $z = u$. Since u is in the s -side of $Z_{s,t}$, by Lemma 6.15 the activating path P_i exists, and the projection x of u onto P_i is in the s -side of Z . Let Q_i be an attachment path from x to u . Then Q_i leads from the s -side to the t -side of Z , and is edge-disjoint from $Z_{s,t}$; hence Q_i is hit by Z' . The case $z = v$ is symmetric. \square

Now, let $P_i, P_j \in \mathcal{P}$ be a pair of paths such that there exists a pair $\{u, v\} \in \mathcal{T}$ which is active in Z , where P_i and P_j activate u and v , respectively. Define

$$\mathcal{T}_{i,j} = \{(u, v) \mid \{u, v\} \in \mathcal{T}, u \text{ projects onto } P_i, v \text{ projects onto } P_j\}.$$

Note that $\mathcal{T}_{i,j}$ contains ordered pairs. For a pair $p = (u, v) \in \mathcal{T}_{i,j}$, let $\pi(u)$ denote the projection of u onto P_i , let $\pi(v)$ denote the projection of v onto P_j , and let $\pi(p) = (\pi(u), \pi(v))$. Define

$$S_{i,j} = \{\pi(p) \mid p \in \mathcal{T}_{i,j}\}.$$

We define a partial order on $S_{i,j}$. For $p_1, p_2 \in S_{i,j}$, with $p_1 = (u_1, v_1)$ and $p_2 = (u_2, v_2)$, we write $p_1 \leq p_2$ if u_1 precedes u_2 on P_i and v_1 precedes v_2 on P_j . Similarly, for pairs $p_1, p_2 \in \mathcal{T}_{i,j}$, we write $p_1 \leq p_2$ if $\pi(p_1) \leq \pi(p_2)$. In the latter case, we say that p_1 *dominates* p_2 . Note that \leq in $\mathcal{T}_{i,j}$ is merely a preorder, as there can be non-trivial equivalence classes of pairs with identical projections. Write $p_1 < p_2$ if $p_1 \leq p_2$ and $\pi(p_1) \neq \pi(p_2)$, and say that p_1 *strictly dominates* p_2 if $p_1 < p_2$.

Lemma 6.17. *Let $p_1 = (u_1, v_1), p_2 = (u_2, v_2)$ be two pairs in $\mathcal{T}_{i,j}$ such that $p_1 \leq p_2$. If p_2 is active in Z , P_i activates u_2 , and P_j activates v_2 , then p_1 is active in Z , P_i activates u_1 , and P_j activates v_1 .*

Proof. If P_i activates u_2 and P_j activates v_2 , then by assumption $\pi(u_2)$ and $\pi(v_2)$ are in the s -side of $Z_{s,t}$. If $p_1 \leq p_2$, then $\pi(u_1)$ precedes $\pi(u_2)$ and $\pi(v_1)$ precedes $\pi(v_2)$. Since \mathcal{P} is a witnessing flow for Z , $\pi(u_1)$ and $\pi(v_1)$ are in the s -side of $Z_{s,t}$ as well. Since the attachment paths from $\pi(u_1)$ to u_1 and from $\pi(v_1)$ to v_1 are disjoint from $Z_{s,t}$, p_1 is active and is activated by the paths P_i and P_j , as required. \square

Let $S'_{i,j} \subseteq S_{i,j}$ be all pairs minimal with respect to domination, i.e., those pairs $p_1 \in S_{i,j}$ such that if $p_2 \leq p_1$ for some $p_2 \in S_{i,j}$ then $p_1 = p_2$. Let $\mathcal{T}'_{i,j}$ contain one representative pair for every $q \in S'_{i,j}$; i.e., for every $q \in S_{i,j}$ there is a pair $p := \pi^{-1}(q)$ in $\mathcal{T}'_{i,j}$ such that $\pi(p) = q$, where p is chosen arbitrarily in case of ambiguity. By Lemma 6.17, if there is an active pair in Z activated by P_i and P_j , then we will find such a pair among $\mathcal{T}'_{i,j}$.

Furthermore, note that the members of $S'_{i,j}$ are pairwise vertex-disjoint, i.e., if $(x, y), (x', y') \in S'_{i,j}$ are distinct then $x \neq x'$ and $y \neq y'$. Order $S'_{i,j}$ as $S'_{i,j} = \{q_1, \dots, q_m\}$, where $q_a = (u_a, v_a)$ for every $a \in [m]$, so that for any $1 \leq a < b \leq m$, u_a precedes u_b on P_i . Then, since the pairs form an antichain, v_b precedes v_a on P_j .

Finally, for $a \in [m]$, let $\pi^{-1}(q_a) = p_a = (x_a, y_a)$. We observe that the sequence of vertices (x_a) is almost monotone in Z .

Lemma 6.18. *Let $a \in [m]$ be an index such that pair p_a is activated by P_i and P_j . Then x_b is in the s -side of Z for all but at most k' indices $b < a$, and x_b is in the t -side of Z for all but at most k' indices $b > a$.*

Proof. Since x_a is activated by u_a , for every $b \leq a$ we have that u_b is in the s -side of Z . Therefore, for any index $b < a$ such that x_b is in the t -side of Z , at least one edge of Z' is needed to cut the attachment path from u_b to x_b . Since attachment paths for distinct projections on P_i are pairwise disjoint, this will exhaust the budget k' if there are too many such indices. Dually, consider an index $b > a$. Since y_a is activated by P_j from attachment vertex v_a , every such vertex v_b is in the s -side of Z . Therefore, similarly, for any index $b > a$ such that x_b is in the s -side of Z , the attachment path from v_b to y_b must be cut by Z' , and again all such attachment paths are disjoint. \square

Lemma 6.19. *Let $a \in [m]$ be an index such that p_a is activated by P_i and P_j . If a exists, then we can sample an index set $I \subseteq [m]$ such that with probability $\Omega(1/k)$ the following hold: $a \in I$; for every $b \in I, b < a$, the vertex x_b is in the s -side of Z ; and for every $b \in I, b > a$, the vertex x_b is in the t -side of Z .*

Proof. Flip a biased coin, sampling every index $b \in [m]$ independently with probability $1/k$. Then success is conditioned on sampling $a \in I$ and not sampling $b \in I$ for every “exceptional” index b as in Lemma 6.18. The former has probability $1/k$, the latter probability at least $(1 - 1/k)^{2k} = \Omega(1)$. \square

Now assume we have sampled (P_i, P_j) and the index sequence I as above, such that the sequence of vertices $(x_b)_{b \in I}$ is monotone with respect to Z , and for one unknown index $a \in I$, Z' separates x_a from u_a . Then we “move” this cut from Z' into $Z_{s,t}$ by (1) adding a new st -path $sx_{i_1} \dots x_{i_r}t$ to G , where $I = (i_1, \dots, i_r)$ in increasing order, and (2) reapplying Theorem 4.1.

Indeed, let $Z^+ = Z \cup \{(x_b, x_a)\}$, where (x_b, x_a) is the newly created edge into x_a , and consider an attachment path Q from u_a to x_a . Let e be the last arc of Z' on Q . Then s reaches the tail of e in $G - Z^+$, since Z is a star st -cut; the head of e reaches x_a by assumption; and x_a reaches t in $G - Z^+$ using the new path. Thus e has moved from the auxiliary part Z' to the core in the new graph. Let us formalize this progress.

Lemma 6.20. *Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be an instance of WEIGHTED DIGRAPH PAIR CUT, let Z be an optimal solution to \mathcal{I} , and let \mathcal{P} be a witnessing flow for Z . There is a polynomial-time procedure that, with probability at least $1/f(k)$ for some $f(k) = 2^{k^{O(1)}}$, produces an output instance $\mathcal{I}' = (G', s, t, \omega', \mathcal{T}, k + 1, 2W)$ and an st -maxflow \mathcal{P}' in G' such that the following holds. There is an optimal solution Z^+ to \mathcal{I}' such that $|Z^+| = |Z| + 1$ and $\omega(Z^+) = \omega(Z) + W$, \mathcal{P}' is a witnessing flow for Z^+ , and $|Z^+| - \lambda_{G'}(s, t) < |Z| - \lambda_G(s, t)$. Furthermore, if \mathcal{I} is a no-instance (i.e., $\omega(Z) > W$), then every output instance is a no-instance.*

Proof. We assume that $k' = |Z| - \lambda_G(s, t) > 0$, as otherwise we can invoke the algorithm of the previous section. Given this, the procedure follows the scheme above. Sample an ordered pair of paths $(P_i, P_j) \in \mathcal{P}^2$. Compute the set $S_{i,j}$ of projections of pairs $p \in \mathcal{T}$ onto (P_i, P_j) , and abort if $S_{i,j} = \emptyset$. Otherwise let $S'_{i,j} \subseteq S_{i,j}$ be the domination-minimal subset of $S_{i,j}$, and compute a set $\mathcal{T}'_{i,j}$ of representatives for $S'_{i,j}$, disambiguating according to some fixed global ordering on \mathcal{T} . Let $\mathcal{T}'_{i,j} = (x_1, \dots, x_m)$ in order of

appearance of $\pi(x_a)$ from s to t on P_i . Let $I \subseteq [m]$ be a subset where every index $a \in [m]$ is sampled independently with probability $1/k$. Let (x'_1, \dots, x'_r) be the subsequence of (x_1, \dots, x_m) indexed by I . Add to G a new path $sx'_1 \dots x'_r t$, let every new edge have weight W , and let $\mathcal{I}' = (G', s, t, \omega', \mathcal{T}, k+1, 2W)$ be the new instance. Finally, apply Lemma 6.13 to \mathcal{I}' and output the resulting instance.

We show correctness. On the one hand, it is easy to see that no false positives are produced. The procedure adds a new st -path of weight W , hence any solution to the output of cardinality at most $k+1$ and weight at most $2W$ must contain a solution to \mathcal{I} of cardinality at most k and weight at most W . On the other hand, let Z be an optimal solution to \mathcal{I} . By Lemma 6.16 there exists at least one active pair $p \in \mathcal{T}$ and a pair of paths $(P_i, P_j) \in \mathcal{P}^2$ that activates p . Let $\mathcal{T}' = \bigcup_{(i,j) \in [\lambda]^2} \mathcal{T}'_{i,j}$, let p be a lowest-index member of \mathcal{T}' according to the assumed ordering, and let (i, j) be such that $p \in \mathcal{T}'_{i,j}$. We claim that $p \in \mathcal{T}'_{j,i}$ as well. Indeed, let $p = (x, y)$, and let u be the projection of x onto P_i and v be the projection of y onto P_j , where P_i activates x and P_j activates y . Then the definitions are symmetrical, and $(u, v) \in S_{i,j}$ and $(v, u) \in S_{j,i}$ both hold. Furthermore, if there existed a pair $q = (v', u') \in S_{j,i}$ that dominates (v, u) , then $(u', v') \in S_{i,j}$ would exist and would dominate (u, v) in $S_{i,j}$. Hence $(v, u) \in S'_{i,j}$. Finally, clearly $p \in \mathcal{T}'_{j,i}$ will be chosen in the disambiguation by choice of p in the ordering.

Now, by Lemma 6.16, either Z' hits every attachment path from u to x , or from v to y . Assume the former by exploiting symmetry, and assume that we sampled the pair of paths (P_i, P_j) . Then by the above reasoning, $p \in \mathcal{T}'_{i,j}$. By Lemma 6.18 we may compute a sequence of vertices (x'_1, \dots, x'_r) such that the sequence contains x , say $x = x'_a$, and x'_b is in the s -side of Z for $b \in [r]$ if and only if $b < a$. Reaching this state has a success probability of $1/k^{\mathcal{O}(1)}$. Let $P' = sx'_1 \dots x'_r t$ be the path newly added to G' and let $Z^+ = Z + (x'_{a-1}, x'_a)$, interpreting $x'_0 = s$. Then $|Z^+| = |Z| + 1$ and $\omega(Z^+) = \omega(Z) + W$, and Z^+ is a solution to the resulting instance \mathcal{I}' . In fact, Z^+ must be an optimal solution, since the cost of any solution increases by at least W from \mathcal{I} to \mathcal{I}' . Furthermore, Z' hits every path from u to $x = x'_a$. As argued previously, for any attachment path Q from u to x , the last edge of $E(Q) \cap Z'$ before x is in $\text{core}_{G'}(Z^+)$, and so is (x'_{a-1}, x'_a) . Therefore $|Z^+| - |\text{core}_{G'}(Z^+)| < |Z| - |\text{core}_G(Z)|$, and this fact is invariant under adding further augmentation edges A' . Now the conditions of Lemma 6.13 apply, and with some probability $1/f(k)$ produces an output (A', \mathcal{P}') where \mathcal{P}' is a witnessing flow for Z^+ in $G' + A'$. Thus we are done. \square

We can now finish the proof.

Proof of Theorem 6.1. Let $\mathcal{I} = (G, s, t, \omega, \mathcal{T}, k, W)$ be the input instance and let Z be an optimal solution to \mathcal{I} , i.e., $\omega(Z)$ is minimum among all solutions with $|Z| \leq k$. We can verify that such a solution exists by an algorithm for unweighted ALMOST 2-SAT. Further, we assume inductively that we have verified that all solutions Z' with cardinality $|Z'| < k$ have weight $\omega(Z') > W$. By Lemma 6.13 we can produce an augmented instance \mathcal{I}' such that, with some probability $1/f(k)$, Z remains an optimal solution for \mathcal{I}' and we have a witnessing flow \mathcal{P} for Z in \mathcal{I}' . Let the current augmented graph be G' , let $\lambda = \lambda_{G'}(s, t)$, and let $k' = k - \lambda$. If $k' = 0$, use the algorithm of Lemma 6.12. Otherwise, we repeat the application of Lemma 6.20 until we have an instance where $\lambda_{G'}(s, t) = k$ and call Lemma 6.12. In total, since the number of edges of the solution not captured by the st -flow decreases each time, we need $\mathcal{O}(k)$ repetitions, giving a total success probability of $1/2^{k^{\mathcal{O}(1)}}$. In total, the value $|Z|$ increases by one per iteration, and the max weight is doubled each iteration, meaning that at termination we have a parameter of $|Z^+| \leq 2k$ and a weight threshold $W' \leq 2^k W$, i.e., the parameter is $\mathcal{O}(k)$ and the weight coding length has increased by only a polynomial factor in k . Thus the total success probability is $1/2^{k^{\mathcal{O}(1)}}$ and the running time of each attempt remains $n^{\mathcal{O}(1)}$. Repeating the procedure some $2^{k^{\mathcal{O}(1)}}$ times gives a randomized algorithm with FPT running time and constant success probability. At a small cost to running time, the success probability can be boosted arbitrarily close to 1 by further repetitions the usual way. \square

We also show a corollary, for a problem that can be translated into WEIGHTED ALMOST 2-SAT. A *valued CSP* (valued constraint satisfaction problem) is an optimization problem defined by a *valued constraint language* defining the types of constraints that are allowed. For every fixed language, the resulting valued CSP is either in P or NP-hard [TZ16, KKR17].

We show that WEIGHTED ALMOST 2-SAT implies that a natural (NP-hard) high-domain valued CSP is FPT parameterized by the number of variables.

Corollary 6.21. *Consider the constraint language consisting of arbitrary unary valued constraints and crisp binary constraints of the form $((x_i \geq a) \vee (x_j \geq b))$, $((x_i \geq a) \vee (x_j < b))$ and $((x_i < a) \vee (x_j < b))$. The corresponding valued CSP on k variables with domain size n can be solved in randomized time $2^{k^{O(1)}} n^{O(1)}$.*

Proof. Let \mathcal{I} be an instance of the valued CSP with variables $X = \{x_1, \dots, x_k\}$. For $i \in [k]$, let D_i be the domain of x_i ; w.l.o.g., we use $D_i = [n_i]$ for some $n_i \in \mathbb{N}$. Let $\omega(x_i = a)$ for $i \in [k]$, $a \in D_i$ denote the total weight of the assignment $x_i = a$ over all unary valued constraints on x_i , or $\omega(x_i = a) = 0$ if no such constraints exist. We create a WEIGHTED ALMOST 2-SAT instance with parameter k and a mixture of crisp and weighted 2-clauses as follows (where a crisp 2-clause C is implemented by adding $k + 1$ copies of C to the formula). Let s and t be Boolean variables forced to take the values 1 and 0, respectively (e.g., create $k + 1$ copies of clauses $(s \vee t)$, $(s \vee \neg t)$ and $(\neg s \vee \neg t)$). For every $i \in [k]$ and $a \in D_i$, create a Boolean variable $v_{i,a}$. Create $k + 1$ copies of the clause $(\neg s \vee v_{i,1})$, and for every $a \in D_i$ create the clause $C = (\neg v_{i,a} \vee v_{i,a+1})$ (if $a < n_i$) respectively $C = (\neg v_{i,a} \vee t)$ (if $a = n_i$), at weight $\omega'(C) = \omega(x_i = a)$. For the remaining constraints, which are all binary clauses, we create crisp 2-clauses, by translating $(x_i \geq a)$ into the literal $v_{i,a}$ and $(x_i < a)$ into the literal $\neg v_{i,a}$. Assign weights $\omega'(C)$ arbitrarily to crisp 2-clauses C . We output the resulting formula ϕ with weights ω' and parameter k .

Correctness of the translation should be clear. For every $i \in [k]$, the weighted clauses on variables $v_{i,a}$ form a chain where at least one clause must be broken, hence by the budget constraint only one clause can be broken per chain. In such an assignment, every chain is monotone, i.e., there is a value $a \in D_i$ such that $v_{i,b} = 1$ if and only if $b \leq a$, where $C = (\neg v_{i,a} \vee v')$ is the broken clause of the chain. The translation of binary constraints of \mathcal{I} to 2-clauses over variables $v_{j,b}$ is now easy to verify. Since WEIGHTED ALMOST 2-SAT is FPT parameterized by k , so is the above valued CSP. \square

References

- [CCH⁺16] Rajesh Chitnis, Marek Cygan, MohammadTaghi Hajiaghayi, Marcin Pilipczuk, and Michal Pilipczuk. Designing FPT algorithms for cut problems using randomized contractions. *SIAM J. Comput.*, 45(4):1171–1229, 2016.
- [CCHM15] Rajesh Hemant Chitnis, Marek Cygan, Mohammad Taghi Hajiaghayi, and Dániel Marx. Directed subset feedback vertex set is fixed-parameter tractable. *ACM Trans. Algorithms*, 11(4):28:1–28:28, 2015.
- [CEM13] Rajesh Hemant Chitnis, László Egri, and Dániel Marx. List h-coloring a graph by removing few vertices. In Hans L. Bodlaender and Giuseppe F. Italiano, editors, *Algorithms - ESA 2013 - 21st Annual European Symposium, Sophia Antipolis, France, September 2-4, 2013. Proceedings*, volume 8125 of *Lecture Notes in Computer Science*, pages 313–324. Springer, 2013.
- [CEM17] Rajesh Chitnis, László Egri, and Dániel Marx. List H-coloring a graph by removing few vertices. *Algorithmica*, 78(1):110–146, 2017.
- [CHM13] Rajesh Hemant Chitnis, MohammadTaghi Hajiaghayi, and Dániel Marx. Fixed-parameter tractability of directed multiway cut parameterized by the size of the cutset. *SIAM J. Comput.*, 42(4):1674–1696, 2013.
- [CKL⁺21] Marek Cygan, Pawel Komosa, Daniel Lokshtanov, Marcin Pilipczuk, Michal Pilipczuk, Saket Saurabh, and Magnus Wahlström. Randomized contractions meet lean decompositions. *ACM Trans. Algorithms*, 17(1):6:1–6:30, 2021.

- [CLL⁺08] Jianer Chen, Yang Liu, Songjian Lu, Barry O’Sullivan, and Igor Razgon. A fixed-parameter algorithm for the directed feedback vertex set problem. *J. ACM*, 55(5):21:1–21:19, 2008.
- [CLP⁺19] Marek Cygan, Daniel Lokshtanov, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. Minimum bisection is fixed-parameter tractable. *SIAM J. Comput.*, 48(2):417–450, 2019.
- [KKPW21] Eun Jung Kim, Stefan Kratsch, Marcin Pilipczuk, and Magnus Wahlström. Solving hard cut problems via flow-augmentation. In Dániel Marx, editor, *Proceedings of the 2021 ACM-SIAM Symposium on Discrete Algorithms, SODA 2021, Virtual Conference, January 10 - 13, 2021*, pages 149–168. SIAM, 2021.
- [KKR17] Vladimir Kolmogorov, Andrei A. Krokhnin, and Michal Rolinek. The complexity of general-valued CSPs. *SIAM J. Comput.*, 46(3):1087–1110, 2017.
- [KLM⁺20] Stefan Kratsch, Shaohua Li, Dániel Marx, Marcin Pilipczuk, and Magnus Wahlström. Multi-budgeted directed cuts. *Algorithmica*, 82(8):2135–2155, 2020.
- [KW20] Stefan Kratsch and Magnus Wahlström. Representative sets and irrelevant vertices: New tools for kernelization. *J. ACM*, 67(3):16:1–16:50, 2020.
- [LRS16] Daniel Lokshtanov, M. S. Ramanujan, and Saket Saurabh. A linear time parameterized algorithm for directed feedback vertex set. *CoRR*, abs/1609.04347, 2016.
- [LRSZ20] Daniel Lokshtanov, M. S. Ramanujan, Saket Saurabh, and Meirav Zehavi. Parameterized complexity and approximability of directed odd cycle transversal. In Shuchi Chawla, editor, *Proceedings of the 2020 ACM-SIAM Symposium on Discrete Algorithms, SODA 2020, Salt Lake City, UT, USA, January 5-8, 2020*, pages 2181–2200. SIAM, 2020.
- [Mar04] Dániel Marx. Parameterized graph separation problems. In Rodney G. Downey, Michael R. Fellows, and Frank K. H. A. Dehne, editors, *Parameterized and Exact Computation, First International Workshop, IWPEC 2004, Bergen, Norway, September 14-17, 2004, Proceedings*, volume 3162 of *Lecture Notes in Computer Science*, pages 71–82. Springer, 2004.
- [Mar06] Dániel Marx. Parameterized graph separation problems. *Theor. Comput. Sci.*, 351(3):394–406, 2006.
- [MOR13] Dániel Marx, Barry O’Sullivan, and Igor Razgon. Finding small separators in linear time via treewidth reduction. *ACM Trans. Algorithms*, 9(4):30:1–30:35, 2013.
- [MR09] Dániel Marx and Igor Razgon. Constant ratio fixed-parameter approximation of the edge multicut problem. *Inf. Process. Lett.*, 109(20):1161–1166, 2009.
- [MR14] Dániel Marx and Igor Razgon. Fixed-parameter tractability of multicut parameterized by the size of the cutset. *SIAM J. Comput.*, 43(2):355–388, 2014.
- [PW18] Marcin Pilipczuk and Magnus Wahlström. Directed multicut is $W[1]$ -hard, even for four terminal pairs. *ACM Trans. Comput. Theory*, 10(3):13:1–13:18, 2018.
- [RO09] Igor Razgon and Barry O’Sullivan. Almost 2-SAT is fixed-parameter tractable. *J. Comput. Syst. Sci.*, 75(8):435–450, 2009.
- [Sau17] Saket Saurabh. What’s next? future directions in parameterized complexity, 2017. Recent Advances in Parameterized Complexity school, Tel Aviv, December 2017.
- [TZ16] Johan Thapper and Stanislav Zivny. The complexity of finite-valued CSPs. *J. ACM*, 63(4):37:1–37:33, 2016.