

Multiple $SO(5)$ isovector pairing and seniority $Sp(2\Omega)$ multi- j algebras with isospin

V.K.B. Kota¹, R. Sahu²

¹Physical Research Laboratory, Ahmedabad 380 009, India

²National Institute of Science and Technology, Palur Hills, Berhampur-761008, Odisha, India

Received Day Month Year (Insert date of submission)

Abstract.

With nucleons occupying several shell model j orbits, the isovector pair creation operator A_μ^1 (creates a two particle state with angular momentum $J = 0$ and isospin $T = 1$) is no longer unique. Choosing it to be a sum of single- j isovector pair creation operators each with a phase, there will be multiple pair $SO(5)$ algebras with isospin; with r number of j orbits, there will be 2^{r-1} $SO(5)$ algebras each with a corresponding complementary $Sp(2\Omega)$ algebra [$2\Omega = \sum_j (2j+1)$] that gives seniority and reduced isospin quantum numbers. Three applications of multiple $SO(5)$ algebras are presented demonstrating the usefulness of considering $SO(5)$ pairing algebras with general sign factors.

KEY WORDS: pairing, multiple algebras, multi- j , $SO(5)$, $Sp(2\Omega)$.

1 Introduction

With identical nucleons (protons or neutrons) in a single- j shell, the pair creation and annihilation operators (S_+ and S_- respectively) and the number operator (\hat{n}) generate remarkably the quasi-spin $SU(2)$ algebra. On the other hand, the $Sp(2j+1)$ subalgebra of the Spectrum Generating Algebra (SGA) $U(2j+1)$ is ‘complementary’ to the quasi-spin- $SU(2)$ algebra and the seniority quantum number v that labels the states w.r.t. $Sp(2j+1)$ algebra, i.e. irreducible representation (irreps), corresponds to the quasi-spin quantum number and similarly, particle number m that labels the irreps of $U(2j+1)$ corresponds to the z -component of quasi-spin. More importantly, this solves the pairing Hamiltonian $H_p = -S_+S_-$ and allows one to extract m dependence of many particle matrix elements of a given operator; see [1] for full details and applications.

All the single- j shell results extend to the multi- j shell situation i.e. for identical nucleons occupying several- j orbits, with $(2j+1)$ replaced by $2\Omega = \sum_j (2j+1)$.

Multiple pairing algebras with isospin

In this situation, v is called generalized seniority. Now, a new result is that with r number of j -orbits there will be 2^{r-1} number of quasi-spin $SU(2)$ and the corresponding $Sp(2\Omega)$ algebras ($S_+ = \sum_j \alpha_j S_+(j)$; $\alpha_j = \pm 1$). These multiple quasi-spin algebras (one for each α_j choice) play an important role in deciding selection rules for electric and magnetic multipole operators. See [2] for details regarding these multiple pairing algebras and [1, 3–5] for the goodness of multi- j seniority in certain nuclei. Also, multi- j seniority provides a framework for shell model theory of the interacting boson model [6, 7].

With isospin (T) degree of freedom, the algebra changes to the more complex $SO(5)$ algebra. With m -nucleons in a single- j orbit the SGA is $U(2(2j+1))$ with the 2 coming from isospin. An isospin conserving subalgebra chain then is $U(2(2j+1)) \supset [U(2j+1) \supset Sp(2j+1)] \otimes SU_T(2)$. Particle number m labels $U(2(2j+1))$, (m, T) label $U(2j+1)$ and T labels $SU_T(2)$ irreps. It is recognized in very early years of shell model that the isovector pair creation and annihilation operators, isospin and the number operator generate a $SO(5)$ algebra, which is 'complementary' to the above $Sp(2j+1)$ algebra. The irreps of $SO(5)$ contain two labels and they can be written in terms of the seniority v and reduced isotopic spin t quantum numbers with (v, t) uniquely labeling the $Sp(2j+1)$ irreps. Another important result is that the isovector pairing Hamiltonian is simply related to the quadratic Casimir invariants of $SO(5)$ and $Sp(2j+1)$. However, an unsatisfactory aspect of the $SO(5)$ algebra of the shell model is that it does not contain isoscalar pair operators in its algebra. For the first papers on single- j shell pairing $Sp(2j+1)$ algebra and the corresponding $SO(5)$ algebra see [8–13]. Similarly, for technical work on these algebras (for example deriving analytical formulas for the Wigner coefficients of $SO(5)$) see [12, 14–18] and for recent applications see [19–23] and references therein. Although many of the single- j shell results extend to the multi- j shell systems, for the multi- j shell situation a crucial aspect is that there will be multiple $SO(5)$ algebras as the isovector pair creation operator here is no longer unique. The purpose of this paper is to introduce and analyze these multiple $SO(5)$ isovector pairing algebras with isospin.

Before proceeding further, let us add that the general mathematical theory describing complementarity between identical nucleon number non-conserving quasi-spin $SU(2)$ and the number conserving $Sp(2\Omega)$, complementarity between $SO(5)$ and $Sp(2\Omega)$ with isospin and similar complementarity between many other algebras (for example proton-neutron pairing $SO(8)$ algebra with LST coupling and the corresponding number conserving algebras [24, 25]) including those for boson systems (see for example [26, 27]) is due to Neergard based on Howe's general duality theorem [28, 29]; first proof of complementarity is due to Helmers [10] and later work is due to Rowe et al. [30]. Now we will give a preview.

In Section 2, multiple pairing $SO(5)$ and $Sp(2\Omega)$ algebras for the multi- j situation are introduced. Section 3 gives formulas for constructing many-particle

matrix elements of the pairing Hamiltonian generating multiple $SO(5)$ algebras. In Section 4 presented are some applications. Finally, Section 5 gives conclusions.

2 Multiple multi- j shell $SO(5)$ and $Sp(2\Omega)$ algebras with isospin

Consider the angular momentum zero coupled isovector pair creation operator $A_\mu^1(j)$ for nucleons in a single- j shell, $A_\mu^1(j) = \sqrt{(2j+1)/2} (a_{j\frac{1}{2}}^\dagger a_{j\frac{1}{2}}^\dagger r)_{0,\mu}^{0,1}$. Now, with nucleons in (j_1, j_2, \dots, j_r) orbits, pair creation operator can be taken as a linear combination of the single- j shell pair creation operators but with different phases giving the generalized isovector pairing operator to be,

$$\mathcal{A}_\mu^1(\beta) = \sum_{p=1}^r \beta_{j_p} A_\mu^1(j_p); \quad \{\beta\} = \{\beta_{j_1}, \beta_{j_2}, \dots, \beta_{j_r}\} = \{\pm 1, \pm 1, \dots\}. \quad (1)$$

Strikingly, the ten operators $\mathcal{A}_\mu^1(\beta)$, $[\mathcal{A}_\mu^1(\beta)]^\dagger$, T_μ^1 and $Q_0 = [\hat{n} - 2\Omega]/2$ (equivalently \hat{n}) form a pairing $SO^{(\beta)}(5)$ algebra for each $\{\beta\}$ set when $\beta_{j_p} = \pm 1$ as in Eq. (1). Note that $2\Omega = \sum_j (2j+1)$, isospin generators $T_\mu^1 = \sum_j \sqrt{(2j+1)/2} (a_{j\frac{1}{2}}^\dagger \tilde{a}_{j\frac{1}{2}})_{\mu}^{0,1}$ and the number operator $\hat{n} = \sum_j \sqrt{2(2j+1)} (a_{j\frac{1}{2}}^\dagger \tilde{a}_{j\frac{1}{2}})_{0,0}^{0,0}$. Without loss of generality we can choose $\beta_{j_1} = +1$ and then the remaining β_{j_p} will be ± 1 . Thus, there will be 2^{r-1} $SO(5)$ algebras. Then, with two j orbits we have two $SO(5)$ algebras $SO^{(+,+)}(5)$ and $SO^{(+,-)}(5)$, with three we have four $SO(5)$ algebras $SO^{(+,+,+)}(5)$, $SO^{(+,+, -)}(5)$, $SO^{(+,-,+)}(5)$ and $SO^{(+,-,-)}(5)$, with four j orbits there will be eight $SO(5)$ algebras and so on. Significantly, the isovector pairing Hamiltonians $H_p(\beta) = -G \sum_\mu \mathcal{A}_\mu^1(\beta) [\mathcal{A}_\mu^1(\beta)]^\dagger$ with G the pairing strength, is simply related to $\mathcal{C}_2(SO^{(\beta)}(5))$, the quadratic Casimir invariant of $SO^{(\beta)}(5)$; $\mathcal{C}_2(SO^{(\beta)}(5)) = 2 \sum_\mu \mathcal{A}_\mu^1(\beta) [\mathcal{A}_\mu^1(\beta)]^\dagger + T^2 + Q_0(Q_0 - 3)$. Further, $SO(5)$ irreps are labeled by (ω_1, ω_2) with ω_1 and ω_2 both integers or half integers and $\omega_1 \geq \omega_2 \geq 0$ [31]. Then, the eigenvalues of $\mathcal{C}_2(SO^{(\beta)}(5))$ are $\langle \mathcal{C}_2(SO^{(\beta)}(5)) \rangle^{(\omega_1, \omega_2)} = \omega_1(\omega_1 + 3) + \omega_2(\omega_2 + 1)$. Expressions for Casimir invariants are given by Racah very early [32]. We will now turn to the complementary $Sp^{(\beta)}(2\Omega)$ algebras.

Consider one-body operators $u_{m_k, m_t}^{k,t}(j_1, j_2)$ defined in terms of the single particle creation and annihilation operators in (jt) space, $u_{m_k, m_t}^{k,t}(j_1, j_2) = (a_{j_1\frac{1}{2}}^\dagger \tilde{a}_{j_2\frac{1}{2}})_{m_k, m_t}^{k,t}$ where $\tilde{a}_{j-m, \frac{1}{2}-m_t} = (-1)^{j-m+\frac{1}{2}-m_t} a_{jm, \frac{1}{2}m_t}$. Now, it is easy to prove that the operators $u^{k,t}(j_1, j_2)$ generate the $U(4\Omega)$ SGA. Moreover, we have the subalgebra $U(4\Omega) \supset [U(2\Omega) \supset Sp(2\Omega)] \otimes SU_T(2)$ with $u_{m_k, 0}^{k,0}(j_1, j_2)$ operators generating $U(2\Omega)$ and $SU_T(2)$ generating isospin. Following the results in [2, 11, 15], it is easy to recognize the generators of $Sp(2\Omega)$

Multiple pairing algebras with isospin

and they are,

$$\begin{aligned} & u_{\mu,0}^{k,0}(j, j) ; k = \text{odd} \\ & V_{\mu,0}^{k,0}(j_1, j_2) = u_{\mu,0}^{k,0}(j_1, j_2) + X(j_1, j_2, k) u_{\mu,0}^{k,0}(j_2, j_1) ; j_1 > j_2 . \end{aligned} \quad (2)$$

Now, the most important result is that for every $SO^{(\beta)}(5)$, there will be a complementary $Sp^{(\beta)}(2\Omega)$ algebra with generators given by Eq. (2) provided

$$X(j_1, j_2, k) = (-1)^{j_1+j_2+k} \beta_{j_1} \beta_{j_2} . \quad (3)$$

Proof for the complementarity is given first by Helmers [10]. Thus, the multiple $SO^{(\beta)}(5)$ algebras that have number non-conserving generators and $Sp^{(\beta)}(2\Omega)$ algebras with only number conserving generators are complementary provided Eq. (3) is satisfied along with Eqs. (1) and (2).

Turning to the irreps, all m nucleon states transform as the antisymmetric irrep $\{1^m\}$ of $U(4\Omega)$ and the irreps of $U(2\Omega)$ will be two columned irreps $\{2^{m_1} 1^{m_2}\}$ in Young tableaux notation with $2m_1 + m_2 = m$ and $T = m_2/2$. Similarly, the $Sp(2\Omega)$ irreps are two columned denoted by $\langle 2^{v_1} 1^{v_2} \rangle$ giving $v = 2v_1 + v_2$ the seniority quantum number and $t = v_2/2$ the reduced isospin. Group theory allows us to obtain $(m, T) \rightarrow (v, t)$ reductions or $(v, t) \rightarrow T$ for a given m [31]. More importantly, it can be shown that (ω_1, ω_2) are equivalent to (v, t) giving $\omega_1 = \Omega - (v/2)$ and $\omega_2 = t$. With all these, the eigenvalues of $H_p^{(\beta)}$ are [11],

$$\left\langle H_p^{(\beta)} \right\rangle^{m,T;v,t} = -\frac{G}{4} \left[(m-v)(2\Omega+3 - \frac{m+v}{2}) - 2T(T+1) + 2t(t+1) \right] . \quad (4)$$

Note that $SO^{(\beta)}(5) \supset [SO(3) \supset SO(2)] \otimes U(1)$ with $SO(3)$ generating T , $SO(2)$ generating M_T (T_z quantum number $(N-Z)/2$) and $U(1)$ generating particle number or $H_1 = (m - 2\Omega)/2$. Then, the eigenstates of $H_p^{(\beta)}$ are

$$\left| \Psi_{H_p^{(\beta)}} \right\rangle \Rightarrow \left| \left(\Omega - \frac{v}{2}, t \right), H_1 = \frac{m - 2\Omega}{2}, T, M_T = \frac{N-Z}{2} \right\rangle . \quad (5)$$

and the labels do not depend on (β) . However, explicit structure of the wavefunctions do depend on (β) ; see Section 4. Thus, they will effect various selection rules and matrix elements of certain transition operators (see Section 4). Finally, with $SO(5)$ algebra it is possible to factorize (m, T) dependence of various matrix elements [1, 15, 21]. In order to enumerate the irrep labels in Eq. (5), used is $U(4\Omega) \supset [U(2\Omega) \supset Sp(2\Omega)] \otimes SU_T(2)$ reductions starting with the $\{1^m\}$ irrep of $U(4\Omega)$. All the rules for these are known [24, 31, 33].

3 Construction of many-particle matrix for pairing Hamiltonian generating multiple $SO(5)$ algebras

In order to probe the role of multiple pair $SO^{(\beta)}(5)$ algebras with isospin, we need to obtain the eigenstates of the pairing Hamiltonian \mathcal{H}_p as a func-

Table 1. Basis states for the $m = 6$ system with $T = 0$ considered in Section 4. Here, $\Omega_1 = 6$ and $\Omega_2 = 5$. See text for details.

#	$ (v_1, t_1)m_1, T_1 : (v_2, t_2)m_2, T_2 ; T = 0\rangle$
1	$ (6, 0), 6, 0 : (0, 0)0, 0 ; 0\rangle$
2	$ (4, 1), 6, 0 : (0, 0)0, 0 ; 0\rangle$
3	$ (2, 0), 6, 0 : (0, 0)0, 0 ; 0\rangle$
4	$ (4, 0), 4, 0 : (2, 0)2, 0 ; 0\rangle$
5	$ (4, 1), 4, 1 : (0, 0)2, 1 ; 0\rangle$
6	$ (4, 1), 4, 1 : (2, 1)2, 1 ; 0\rangle$
7	$ (2, 1), 4, 0 : (2, 0)2, 0 ; 0\rangle$
8	$ (2, 0), 4, 1 : (0, 0)2, 1 ; 0\rangle$
9	$ (2, 0), 4, 1 : (2, 1)2, 1 ; 0\rangle$
10	$ (2, 1), 4, 1 : (0, 0)2, 1 ; 0\rangle$
11	$ (2, 1), 4, 1 : (2, 1)2, 1 ; 0\rangle$
12	$ (0, 0), 4, 0 : (2, 0)2, 0 ; 0\rangle$
13	$ (0, 0), 2, 1 : (4, 1)4, 1 ; 0\rangle$
14	$ (2, 0), 2, 0 : (4, 0)4, 0 ; 0\rangle$
15	$ (2, 1), 2, 1 : (4, 1)4, 1 ; 0\rangle$
16	$ (2, 0), 2, 0 : (0, 0)4, 0 ; 0\rangle$
17	$ (2, 0), 2, 0 : (2, 1)4, 0 ; 0\rangle$
18	$ (2, 1), 2, 1 : (2, 0)4, 1 ; 0\rangle$
19	$ (2, 1), 2, 1 : (2, 1)4, 1 ; 0\rangle$
20	$ (0, 0), 2, 1 : (0, 0)4, 1 ; 0\rangle$
21	$ (0, 0), 2, 1 : (2, 1)4, 1 ; 0\rangle$
22	$ (0, 0), 0, 0 : (6, 0)6, 0 ; 0\rangle$
23	$ (0, 0), 0, 0 : (4, 1)6, 0 ; 0\rangle$
24	$ (0, 0), 0, 0 : (2, 0)6, 0 ; 0\rangle$

tion of $\{\beta\}$'s. A convenient basis for constructing the \mathcal{H}_p matrix is the product basis defined by the single- j shell $SO(5)$ basis. We will illustrate this using two j -orbits say j_1 and j_2 . Hereafter, we call the corresponding spaces a and b respectively (or 1 and 2). Then, the basis states are, $\Psi_{ab}(T M_T) = |(\omega_1^a \omega_2^a) H^a T^a, (\omega_1^b \omega_2^b) H^b T^b ; T M_T\rangle$ or equivalently $|(v_1, t_1)m_1 T_1, (v_2, t_2)m_2 T_2 ; T M_T\rangle$. Given m number of nucleons, with m_1 in number in the first orbit and m_2 in the second orbit, $m = m_1 + m_2$. Note that $\Omega_1 = j_1 + \frac{1}{2}$, $\Omega_2 = j_2 + \frac{1}{2}$, $H^a = \frac{m_1}{2} - \Omega_1$ and $H^b = \frac{m_2}{2} - \Omega_2$. Similarly T^a and T^b are the isospins in the two spaces respectively. Now, a general pairing Hamiltonian [with $\mathcal{A}_\mu^1(\alpha) = A_\mu^1(j_1) + \alpha A_\mu^1(j_2)$] is,

$$\mathcal{H}_p(\xi, \alpha) = \frac{(1-\xi)}{m} \hat{n}_2 - \frac{\xi}{m^2} \left\{ 4 \sum_{\mu} \mathcal{A}_\mu^1(\alpha) [\mathcal{A}_\mu^1(\alpha)]^\dagger \right\}. \quad (6)$$

Here, \hat{n}_2 is the number operator for the second orbit and ξ and α are parameters changing from 0 to 1 and +1 to -1 respectively. Note that for $\xi = 1$ and $\alpha = +1$ we have a $SO^{(+)}(5)$ algebra in the total two-orbit space and similarly for $\xi = 1$

Multiple pairing algebras with isospin

and $\alpha = -1$ the $SO^{(-)}(5)$ algebra. Diagonal matrix elements of \mathcal{H}_p in our basis follow easily from Eq. (4); note that \hat{n}_2 gives m_2 and the other part giving diagonal matrix elements is $\sum_{\mu} [\{A_{\mu}^1(j_1)[A_{\mu}^1(j_1)]^{\dagger}\} + \alpha^2 \{A_{\mu}^1(j_2)[A_{\mu}^1(j_2)]^{\dagger}\}]$. The off-diagonal matrix elements involve $SO(5) \supset SO(3) \otimes U(1)$ reduced Wigner coefficients and using Eq. (14) of [15] will give,

$$\begin{aligned}
& \langle (\omega_1^a \omega_2^a) H_f^a T_f^a, (\omega_1^b \omega_2^b) H_f^b T_f^b; T M_T | \mathcal{H}_p(\zeta, \alpha) \\
& | (\omega_1^a \omega_2^a) H_i^a T_i^a, (\omega_1^b \omega_2^b) H_i^b T_i^b; T M_T \rangle = - \left(\frac{4\xi}{m^2} \right) (\alpha) \\
& \times [(\omega_1^a(\omega_1^a + 3) + \omega_2^a(\omega_2^a + 1))^{1/2} [(\omega_1^b(\omega_1^b + 3) + \omega_2^b(\omega_2^b + 1))^{1/2}] \\
& \times (-1)^{T_f^a + T_f^b + T + 1} \sqrt{(2T_f^a + 1)(2T_f^b + 1)} \begin{Bmatrix} T & T_f^b & T_f^a \\ 1 & T_i^a & T_i^b \end{Bmatrix} \quad (7) \\
& \times \left\langle (\omega_1^a \omega_2^a) H_i^a T_i^a \quad (11) 1, 1 \quad || \quad (\omega_1^a \omega_2^a) H_f^a T_f^a \right\rangle \\
& \times \left\langle (\omega_1^b \omega_2^b) H_i^b T_i^b \quad (11) -1, 1 \quad || \quad (\omega_1^b \omega_2^b) H_f^b T_f^b \right\rangle
\end{aligned}$$

for $H_f^a = H_i^a + 1$ and $H_f^b = H_i^b - 1$; $H_i^a = \frac{m_1}{2} - \Omega_1$, $H_i^b = \frac{m_2}{2} - \Omega_2$. The $\langle \dots || \dots \rangle$ factors above are the $SO(5) \supset SO(3) \otimes U(1)$ reduced Wigner coefficients. For the $m = 6$ system considered in the next Section, the needed Wigner coefficients follow from Tables III in [15] and Table A.1 in [16]. It is important to mention that for simplicity, in Eq. (7) we are not showing the additional label that is required as discussed in [15, 16]. This label is called β in [16]. Finally, Eq. (7) can be extended to three or more orbits by using isospin T couplings. Thus, \mathcal{H}_p construction is possible with multiple $SO^{(\beta)}(5)$ algebras provided all the needed Wigner coefficients in Eq. (7) are known.

4 Applications of multiple $SO(5)/Sp(2\Omega)$ algebras

Electromagnetic transition operators T^{EL} and T^{ML} are one-body operators and their $SO(5) \supset [SO_T(3) \supset SO(2)] \otimes U(1)$ tensorial structure is $T_{H_1, T, M_T}^{(\omega_1, \omega_2)}$ with $(\omega_1 \omega_2) = (11) \oplus (10) \oplus (00)$. More importantly, Eqs. (2) and (3) show that it is possible for T^{EL} and T^{ML} to be generators of $Sp^{(\beta)}(2\Omega)$ giving selection rules under certain conditions. We have the following results: (i) isovector parts of T^{EL} and T^{ML} will not be $Sp^{(\beta)}(2\Omega)$ scalars as the generators of this algebras are only isoscalar operators; (ii) the isoscalar part of T^{ML} with L odd (they preserve parity) or even can be $Sp^{(\beta)}(2\Omega)$ scalars provided $\beta_{j(\ell)} = (-1)^\ell$ for the $j(\ell)$ orbits; (iii) the T^{EL} with L even or odd will not be generators of any $Sp^{(\beta)}(2\Omega)$ as the $X(j_f, j_i, L)$ (see Eqs. (2) and (3)) given by the isoscalar part will not lead to a formula for β_{j_i} real. With the phase choice $\beta_{j(\ell)} = (-1)^\ell$, the selection rule from the generators that they will not change (v, t) or $(\omega_1 \omega_2)$ irreps can be used in experimental tests of this phase choice. Besides this, EL and ML transitions can change seniority only by units of 2, i.e. transition for $v \rightarrow v, v \pm 2$ states are only allowed. In addition, the (m, T) dependence of say quadrupole moments and $B(E2)$'s can be written down using $SO(5)$ algebra.




Figure 1. Energy spectra of the pairing Hamiltonian \mathcal{H}_p in Eq. (6) as a function of ξ and α . Note that the energies (E) are unitless. See text for details.

In the second application, let us consider a two level system with first level having $\Omega_1 = 6$ with $-ve$ parity and the second level having $\Omega_2 = 5$ with $+ve$ parity. This is appropriate for nuclei in $A=56-80$ region so that the $(1p_{3/2}, 0f_{5/2}, 1p_{1/2})$ orbits with degenerate single particle levels give the $\Omega_1 = 6$ orbit (we will call it orbit #1 or a) and $0g_{9/2}$ gives the $\Omega_2 = 5$ orbit (we will call it orbit #2 or b). In our numerical calculations we use the system with six nucleons in the above two orbits. Then, the number of $+ve$ parity basis states for $m = 6$ and $T = 0$ will be 24 as shown in Table 1. Using these basis states, the matrix for \mathcal{H}_p defined by Eq. (6) is constructed following the formulation in Section 3. Diagonalization of $\mathcal{H}_p(\xi = 1, \alpha = \pm 1)$ will give eigenvalues that must be same as those given by Eq. (4) with $(m = 6, T = 0)$ and $(v, t) = (6, 0), (4, 1)$ and $(2, 0)$. The eigenvalues are $0, -44/m^2$ and $-84/m^2$ respectively with degeneracies 13, 9 and 2 respectively. It is easy to see that the wavefunctions are of the form $|(v_1, t_1)(v_2, t_2)(v, t)\gamma, m = 6, T = 0\rangle$ where γ are additional labels. Therefore, a sum of $\mathcal{C}_2(SO^{(a)}(5))$ and $\mathcal{C}_2(SO^{(b)}(5))$ will remove some of the degeneracies in the spectrum without changing the eigenvectors. By adding a

Multiple pairing algebras with isospin

term $-(\xi/m^2)[\mathcal{C}_2(SO^{(a)}(5)) + \mathcal{C}_2(SO^{(b)}(5))]$ to $\mathcal{H}_p(\xi, \alpha)$ we have calculated the eigenvalues for the $(m = 6, T = 0)$ system and shown in Fig. 1 are the energies of the 24 states as a function of ξ for nine α values. For example, wavefunctions for the lowest two degenerate states are,

$$\begin{aligned} |\Psi_1^{m=6, T=0}\rangle &= \sqrt{\frac{13}{35}} |4(0, 0)0; 2(20)0\rangle - \alpha \sqrt{\frac{16}{35}} |2(0, 0)1; 2(20)1\rangle + \sqrt{\frac{6}{35}} |0(0, 0)0; 6(20)0\rangle, \\ |\Psi_2^{m=6, T=0}\rangle &= \sqrt{\frac{11}{42}} |6(2, 0)0; 0(00)0\rangle - \alpha \sqrt{\frac{20}{42}} |4(2, 0)1; 2(00)1\rangle + \sqrt{\frac{11}{42}} |2(2, 0)0; 4(00)0\rangle. \end{aligned} \quad (8)$$

Here the notation used is $|m_1(v_1, t_1)T_1; m_2(v_2, t_2)T_2\rangle$. Eq. (8) shows the role of α , i.e. the two $SO(5)$ algebras. As seen from Fig. 1, clearly by changing (ξ, α) it is possible to study order-chaos-order transitions. Detailed analysis of this including all T 's will be reported elsewhere.

Two-particle transfer strengths form the third application. As an example let us consider removal of a isovector pair from the lowest two states [these are Ψ_1 and Ψ_2 in Eq. (8)] of the $(m = 6, T = 0)$ system generating the states of $(m = 4, T = 1)$ system. To study the transfer strengths, we have diagonalized $\mathcal{H}_p(\xi = 1, \alpha = \pm 1)$ in $(m = 4, T = 1)$ space and the basis states here are 14 in number. Then, the eigenstates belong to $(v, t) = (21), (20)$ and (41) irreps in the 4 nucleon space. There are three, two and nine states respectively with these irreps and the corresponding eigenvalues are $-44/m^2, -40/m^2$ and 0 respectively. The transition operator for example can be chosen to be $P = [A_\mu^1(j_i)]^\dagger$ or it can be $[\mathcal{A}_\mu^1(\alpha)]^\dagger$ with $\alpha = +1$ or -1 . These will not change $(v_1 t_1)$ and $(v_2 t_2)$ of the states. From Eq. (8) it is easy to see that the transfer is allowed to the two states with $(v, t) = (2, 0)$ and these are

$$\begin{aligned} |\Phi_1^{m=4, T=1}\rangle &= \sqrt{\frac{1}{2}} |4(2, 0)1; 0(00)0\rangle + \alpha \sqrt{\frac{1}{2}} |2(2, 0)0; 2(00)1\rangle, \\ |\Phi_2^{m=4, T=1}\rangle &= \sqrt{\frac{2}{5}} |0(0, 0)0; 4(20)1\rangle + \alpha \sqrt{\frac{3}{5}} |2(0, 0)1; 2(20)0\rangle. \end{aligned} \quad (9)$$

With the α dependence in both the six and four particle states [see Eqs. (8) and (9)], clearly, the two-particle transfer strengths depend on α . In practice we need to add a term in the Hamiltonian that mixes the states Ψ_1 and Ψ_2 and similarly Φ_1 and Φ_2 . This is being investigated and explicit formulas for the transfer strengths will be reported elsewhere.

5 Conclusions

Extending the previous results [2] on multiple $SU(2)$ pairing algebras for identical nucleons occupying several j -orbits, in this paper it is shown that there are multiple $SO(5)$ pairing algebras for nucleons (with isospin) occupying several j -orbits. Further, a method to analyze the results, based on the algebra in [15], due to multiple $SO(5)$ (or the equivalent $Sp(2\Omega)$) algebras is described. Finally, in three applications are briefly discussed. More detailed investigations of multiple $SO(5)$ algebras and their applications will be reported elsewhere. Present

work complements the corresponding investigations without isospin in [2] and on multiple $SU(3)$ algebras in [34, 35].

Acknowledgements

R. Sahu is thankful to SERB of DST (Government of India) for financial support.

References

- [1] I. Talmi(1993) "Simple Models of Complex Nuclei: The Shell Model and the Interacting Boson Model". Harwood, New York.
- [2] V.K.B. Kota (2017) Multiple multi-orbit fermionic and bosonic pairing and rotational $SU(3)$ algebras. *Bulg. J. Phys.* **44** 454-465.
- [3] B. Maheswari, A.K. Jain (2016) Odd tensor electric transitions in high-spin Sn-isomers and generalized seniority. *Phys. Lett.* **B753** 122-125.
- [4] B. Maheswari, A.K. Jain, B. Singh (2016) Asymmetric behavior of the $B(E2; 0^+ \rightarrow 2^+)$ values in $^{104-130}\text{Sn}$ and generalized seniority. *Nucl. Phys.* **A952** 62-69.
- [5] B. Maheswari, D. Choudhury, A.K. Jain (2021) Generalized seniority isomers in and around $Z=82$ closed shell: A survey of Hg, Pb and Po isotopes. *Nucl. Phys.* **A1014** 122262/1-12.
- [6] F. Iachello, A. Arima (1987) "The Interacting Boson Model" Cambridge University Press, Cambridge.
- [7] F. Iachello, I. Talmi (1987) Shell-model foundations of the interacting boson model. *Rev. Mod. Phys.* **59**, 339-361.
- [8] B.H. Flowers (1952) Studies in jj-coupling I. Classification of nuclear and atomic states. *Proc. Roy. Soc. (London)* **A212** 248-263.
- [9] A. Kerman (1961) Pairing forces and nuclear collective motion. *Ann. Phys. (N.Y.)* **12** 300-329.
- [10] K. Helmers (1961) Symplectic invariants and Flower's classification of shell model states. *Nucl. Phys.* **23** 594-611.
- [11] B.H. Flowers, S. Szpikowski (1964) A generalized quasi-spin formalism. *Proc. Phys. Soc. (London)* **84** 193-199.
- [12] K.T. Hecht (1965) Some simple R_5 Wigner coefficients and their application. *Nucl. Phys.* **63** 177-213.
- [13] J.C. Parikh (1965) The role of isospin in pair correlations for configurations of the type $(j)^N$. *Nucl. Phys.* **63** 214-232.
- [14] J.N. Ginocchio (1965) Generalized quasi-spin in neutron-proton systems. *Nucl. Phys.* **74** 321-347.
- [15] K.T. Hecht (1965) Five-dimensional quasi-spin. Exact solutions of a pairing hamiltonian in the J-T scheme. *Phys. Rev.* **139** B794-B817.
- [16] R.P. Hemenger, K.T. Hecht (1970) Five-dimensional quasi-spin. Towards a complete classification of the isospin characteristics of shell model states in the seniority scheme. *Nucl. Phys.* **A145** 468-496.
- [17] K.T. Hecht, J.P. Elliott (1985) Coherent-state theory for the proton-neutron quasi-spin group. *Nucl. Phys.* **A438** 29-40.

Multiple pairing algebras with isospin

- [18] K.T. Hecht (1989) Wigner coefficients for the proton-neutron quasi-spin group: An application of vector coherent state technique. *Nucl. Phys.* **A493** 29-60.
- [19] J. Engel, K. Langanke, P. Vogel (1996) Pairing and isospin symmetry in proton-rich nuclei. *Phys. Lett.* **B389** 211-216.
- [20] D. Stefanik, F. Simkovic, K. Muto, A. Faessler (2013) Two-neutrino double- β decay Fermi transition and two-nucleon interaction. *Phys. Rev. C* **88** 025503/1-7.
- [21] J.P. Elliott, J.A. Evans, G.L. Long, V.S. Lac (1994) Shell-model matrix elements in a neutron-proton quasi-spin formalism for several shells. *J. Phys. A: Math. Gen.* **27** 4465-4471.
- [22] J.A. Evans, J.P. Elliott, V.S. Lac, G.L. Long (1995) An IBM-3 hamiltonian from a multi-j-shell model. *Nucl. Phys.* **A593** 85-94.
- [23] J.P. Draayer, C.S. Han, K.J. Weeks, K.T. Hecht (1981) Band crossing and the pre-alignment B(E2) anomaly in ^{126}Ba . *Nucl. Phys.* **A365** 127-141.
- [24] V.K.B. Kota, J.A. Castilho Alcarás (2006) Classification of states in $SO(8)$ proton-neutron pairing model. *Nucl. Phys.* **A764** 181-204.
- [25] D. J. Rowe, M. J. Carvalho, J. Repka (2012) Dual pairing of symmetry and dynamical groups in physics. *Rev. Mod. Phys.* **84** 711-757.
- [26] V.K.B. Kota (2000) $O(36)$ Symmetry Limit of IBM-4 with good s , d and sd Boson Spin-Isospin Wigner's $SU(4) \sim O(6)$ Symmetries for $N \approx Z$ Odd-Odd Nuclei. *Ann. Phys. (N.Y.)* **280** 1-34.
- [27] S. Lerma H, B. Errea, J. Dukelsky, S. Pittel, P. Van Isacker (2006) Exactly solvable models of proton and neutron interacting bosons. *Phys. Rev. C* **74** 024314/1-7.
- [28] K. Neergard (2019) Proof by characters of the orthogonal orthogonal duality and relations of Casimir invariants. *J. Math. Phys.* **60** 081705/1-14.
- [29] K. Neergard (2020) Fock space dualities. *J. Math. Phys.* **61** 081702/1-14; arXiv:2006.08047v4 [math-ph] (2021); article in this volume.
- [30] D.J. Rowe, J. Repka, M. J. Carvalho (2011) Simple unified proofs of four duality theorems. *J. Math. Phys.* **52**, 013507/1-24.
- [31] B.G. Wybourne (1970) "Symmetry Principles and Atomic Spectroscopy". Wiley, New York.
- [32] G. Racah (1952) Nuclear levels and Casimir operator. L. Farkas memorial volume, edited by A. Farkas and E.P. Wigner, Research Council of Israel, Jerusalem, pp. 294-304; <http://cds.cern.ch/record/104181>.
- [33] D.E. Littlewood (1950) "The Theory of Group Characters and Matrix Representations of Groups". 2nd edn, Oxford University Press, Oxford.
- [34] V.K.B. Kota, R. Sahu, P.C. Srivastava (2019) Shell model analysis of multiple $SU(3)$ algebras in nuclei. *Bulg. J. Phys.* **46** 313-324; R. Sahu, V.K.B. Kota, P.C. Srivastava (2020) Quadrupole properties of the eight $SU(3)$ algebras in ($sdgi$) space. *Eur. Phys. J. Special Topics* **229** 2389-2403.
- [35] V.K.B. Kota, R. Sahu (2021) Multiple $SU(3)$ algebras in interacting boson model and shell model: Results for (β, γ) bands and scissors 1^+ band. *Nucl. Phys.* **A1016**, 122313/1-27.